



Office de la Propriété

Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2725600 A1 2010/01/07

(21) **2 725 600**

(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2009/06/23
(87) Date publication PCT/PCT Publication Date: 2010/01/07
(85) Entrée phase nationale/National Entry: 2010/11/23
(86) N° demande PCT/PCT Application No.: US 2009/048194
(87) N° publication PCT/PCT Publication No.: 2010/002622
(30) Priorité/Priority: 2008/07/01 (US12/165,743)

(51) Cl.Int./Int.Cl. *C07C 43/13* (2006.01)

(71) **Demandeur/Applicant:**
E. I. DU PONT DE NEMOURS AND COMPANY, US

(72) **Inventeur/Inventor:**
MCALAIN, STEPHAN J., US

(74) **Agent:** TORYS LLP

(54) Titre : ALCOOLS FLUORES
(54) Title: FLUORINATED ALCOHOLS

(57) **Abrégé/Abstract:**

A compound comprising Formula 5: $R_fOCFHC_2O(CH_2CH_2O)_vH$, wherein R_f is $C_cF(2c+1)$; c is 2 to about 6; and v is 2 to about 4; and a process for its preparation comprising contacting a compound of Formula 6: $R_f-O-CF=CF_2$, wherein R_f is $C_cF(2c+1)$, and c is 2 to about 6, with a compound of Formula 7 $HO-(CH_2CH_2O)_vH$, wherein v is 2 to about 4.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(10) International Publication Number
WO 2010/002622 A1(43) International Publication Date
7 January 2010 (07.01.2010)(51) International Patent Classification:
C07C 43/13 (2006.01)(21) International Application Number:
PCT/US2009/048194(22) International Filing Date:
23 June 2009 (23.06.2009)(25) Filing Language:
English(26) Publication Language:
English(30) Priority Data:
12/165,743 1 July 2008 (01.07.2008) US(71) Applicant (for all designated States except US): **E. I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US).**

(72) Inventor; and

(75) Inventor/Applicant (for US only): **MCLAINE, Stephan, J. [US/US]; 4405 Emerson Road, Wilmington, DE 19802 (US).**(74) Agents: **MAYER, Nancy, S. et al.; E. I. du Pont de Nemours and Company, Legal Patent Records Center, 4417 Lancaster Pike, Wilmington, DE 19805 (US).**(81) Designated States (unless otherwise indicated, for every kind of national protection available): **AE, AG, AL, AM,**

AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))



WO 2010/002622 A1

(54) Title: **FLUORINATED ALCOHOLS**(57) Abstract: A compound comprising Formula 5: $R_fOCFHC_2O(CH_2CH_2O)_v-H$, wherein R_f is $C_cF(2c+1)$; c is 2 to about 6; and v is 2 to about 4; and a process for its preparation comprising contacting a compound of Formula 6: $R_f-O-CF=CF_2$, wherein R_f is $C_cF(2c+1)$, and c is 2 to about 6, with a compound of Formula 7 $HO-(CH_2CH_2O)_v-H$, wherein v is 2 to about 4.

TITLE**FLUORINATED ALCOHOLS**BACKGROUND OF THE INVENTION

Historically, many fluoroalkyl surfactants were based on the 5 perfluoroalkylethanols, $F(CF_2CF_2)_qCH_2CH_2OH$, the so-called “Telomer B alcohols”, where q was typically about 2 to 10. The Telomer B alcohols and their preparation are described by Kirchner et al. in US Patent 5,202,506. Other fluoroalkyl surfactants based on Telomer B alcohols have included “twin-tailed” anionic surfactants such as

10 $F(CF_2CF_2)_q(CH_2CH_2)OCOCH_2CH(SO_3Na)COO(CH_2CH_2)(CF_2CF_2)_qF$, where q is as defined above, prepared by firstly reacting two moles of one or more perfluoroalkylethanols with one of maleic anhydride and, secondly, reacting the diester product with sodium hydrogen sulfite solution, as described, for instance, by Yoshino et al. in “Surfactants having polyfluoroalkyl chains. II. Syntheses of 15 anionic surfactants having two polyfluoroalkyl chains including a trifluoromethyl group at each tail and their flocculation-redispersion ability for dispersed magnetite particles in water”, Journal of Fluorine Chemistry (1995), 70(2), 187-91. Yoshino et al. reported examples wherein q was 2, 3, and 4 for use in supercritical carbon dioxide. Yoshino et al. report twin-tailed surfactants 20 wherein both end groups are limited to perfluoroalkyl groups.

European Patent 1,522,536 discloses the preparation of fluorine-containing alcohols useful as intermediates of formula $R^1CY^1HCY^2Y^3OQ^1OH$ or $HOQ^2OCZ^1Z^2CZ^3HR^2CZ^4HCZ^5Z^6OQ^2OH$ wherein R^1 and R^2 are fluorine-containing groups, Y^1 to Y^3 and Z^1 to Z^6 are fluorine atoms, and Q^1 and Q^2 are 25 alkylene groups.

Nagai et al. in US Patent Application 2008/0093582, describe twin-tailed surfactants of the structure $R_f-(CH_2)_{n1}-(X^1)_{p1}-CH(SO_3M)(X^2)_{q1}-R_h$ wherein R_f is a fluoroalkyl group that may contain an ether bond, X^1 and X^2 are the same or different divalent linking groups; M is H, an alkali metal, half an 30 alkaline earth metal, or ammonium; R_h is an alkyl group; $n1$ is an integer of 1 to 10; and $p1$ and $q1$ are each 0 or 1.

One common route to perfluoroalkylethanols used to make such surfactants is a multi-step process using tetrafluoroethylene. Tetrafluoroethylene is a hazardous and expensive intermediate with limited availability. It is 35 desirable to provide fluorinated surfactants that use less or no tetrafluoroethylene

in their preparation. It is also desirable to provide new and improved fluorinated surfactants in which the perfluoroalkyl group of the prior art is replaced by partially fluorinated terminal groups that require less tetrafluoroethylene and show increased fluorine efficiency. By "fluorine efficiency" is meant the ability to use 5 a minimum amount of fluorochemical to obtain a desired surface effect or surfactant properties, when applied to substrates, or to obtain better performance

using the same level of fluorine. A polymer having high fluorine efficiency generates the same or greater level of surface effect using a lower amount of fluorine than a comparative polymer. The present invention provides intermediates used to prepare such improved fluorinated surfactants.

5

SUMMARY OF THE INVENTION

The present invention further comprises compound of Formula 5



wherein

R_f is $C_cF_{(2c+1)}$;

10 c is 2 to about 6; and v is 2 to about 4.

The present invention further comprises a process for the preparation of a compound of Formula 5



wherein

15 R_f is $C_cF_{(2c+1)}$;

c is 2 to about 6; and v is 2 to about 4,

comprising contacting a compound of Formula 6



wherein R_f is $C_cF_{(2c+1)}$, and c is 2 to about 6, with a compound of Formula 7





wherein v is 2 to about 4.

DETAILED DESCRIPTION

Herein trademarks are shown in upper case.

Surfactants of interest have the structure of Formulae 1A, 1B, or 1C.




wherein

R is H or a linear or branched alkyl group $C_bH_{(2b+1)}$ - wherein b is from 1 to about 18, preferably from about 6 to about 18;

each R_f is independently $C_cF_{(2c+1)}$ having c of from about 2 to about 6, preferably from 2 to 4, and more preferably 4;

5 X is a linear or branched difunctional alkyl sulfonate group



wherein e is from 2 or 3, preferably 3; and M is a monovalent cation which is hydrogen, ammonium, alkali metal, or alkaline earth metal, and is preferably Na;

10 R_a is selected from the group consisting of structure (i) through (vi) wherein R_f is as defined above, and g is 1 to about 4, preferably from 1 to 3, and more preferably 2:


- (i) $R_f(CH_2CF_2)_d-(C_gH_{2g})$ - wherein d is 1 to about 3, preferably from 1 to 2, and more preferably 1;
- (ii) $R_fOCF_2CF_2-(C_gH_{2g})$ -;
- 15 (iii) $R_fOCFHCFC_2O(CH_2CH_2O)_v-(C_gH_{2g})$ - wherein v is 1 to about 4, preferably from 1 to 2, and more preferably 2;
- (iv) $R_fOCFHCFC_2O(C_wH_{2w})$ - wherein w is from about 3 to about 12, preferably from 4 to 6, and more preferably 4;
- (v) $R_fOCF(CF_3)CONH-(C_gH_{2g})$ -; or
- 20 (vi) $R_f(CH_2)_h[(CF_2CF_2)_i(CH_2CH_2)_j]_k$ wherein h is 1 to about 6, preferably from 1 to 3, and more preferably 1; and i, j, and k are each independently 1, 2, or 3, or a mixture thereof, preferably 1 or 2, and more preferably 1; provided that the total number of carbon atoms in group (vi) is from about 8 to about 22. The preferred R_a groups are (i), (ii), (iii), and (iv).

25 Preferred embodiments of Formula 1A, 1B and 1C are those wherein R_a is group (i) $R_f(CH_2CF_2)_d-(C_gH_{2g})$ -, (ii) $R_fOCF_2CF_2-(C_gH_{2g})$ -; (iii) $R_fOCFHCFC_2O(CH_2CH_2O)_v-(C_gH_{2g})$ -; or (iv) $R_fOCFHCFC_2O(C_wH_{2w})$ -; when c is 3 or 4; and X is $CH_2CH(SO_3M)$, $CH_2CH(CH_2SO_3M)$, $CH(CH_3)CH(SO_3M)$, $CH_2CH(SO_3M)CH_2$, or $CH_2CH(SO_3M)CH_2CH_2$. More specifically preferred embodiments of Formula 1A, 1B and 1C are when d is 1, g is 1 or 2, and R_f is C_3F_7 or C_4F_9 . Also specifically preferred are compounds wherein R_f is C_3F_7 or C_4F_9 and X is $C_3H_5(SO_3Na)$ or $CH_2CH(SO_3Na)$. The compound of Formula 1B wherein R_a is $C_4F_9CH_2CF_2CH_2CH_2$ or $C_3F_7CH_2CF_2CH_2CH_2$ and R_f is $(CF_2)_6F$ is also preferred. A preferred embodiment of Formula 1A ($R_aOCO-X-$



COOR_a) is C₄F₉CH₂CF₂CH₂CH₂OC(O)C₃H₅(SO₃Na)C(O)O-CH₂CH₂CF₂-CH₂C₄F₉. A preferred embodiment of Formula 1B is C₄F₉CH₂CF₂CH₂CH₂-OC(O)CH₂CH(SO₃Na)C(O)O-CH₂CH₂(CF₂)₆F. A preferred embodiment of Formula 1C is C₄F₉CH₂CF₂CH₂-CH₂OC(O)CH₂CH(SO₃Na)C(O)O(CH₂)₆H.

5 The surfactants of Formulae 1A, 1B, and 1C economize on the use of tetrafluoroethylene in their preparation and provide comparable or improved properties, versus prior art surfactants derived from Telomer B alcohols.

10 The surfactants of Formulae 1A, 1B, and 1C are prepared via the unsaturated intermediates of Formulae 2A, 2B, and 2C according to the following Reaction Scheme A:



The unsaturated intermediates used in the preparation of Formula 1A, 1B and 1C are compounds of Formula 2A, 2B and 2C:



wherein

R_a is the group (i) $R_f(CH_2CF_2)_d-(C_gH_{2g})-$; (ii) $R_fOCF_2CF_2-(C_gH_{2g})-$; (iii) $R_fOCFHCFC_2O(CH_2CH_2O)_v-(C_gH_{2g})-$; (iv) $R_fOCFHCFC_2O(C_wH_{2w})-$; (v) $R_fOCF(CF_3)CONH-(C_gH_{2g})-$; or (vi) $R_f(CH_2)_h[(CF_2CF_2)_i(CH_2CH_2)_j]_k-$

10 each R_f is independently $C_cF_{(2c+1)}$;

c is 2 to about 6, preferably from 2 to 4, more preferably 4;

d is 1 to about 3, preferably from 1 to 2, more preferably 1;

g is 1 to 4, preferably from 1 to 3, more preferably 2;

v is 1 to about 4, preferably from 2 to 3, more preferably 2;

15 w is from about 3 to about 12, preferably from 4 to 6, more preferably 4;

h is 1 to about 6, preferably from 1 to 3, more preferably 2;

i, j , and k are each independently 1, 2, or 3, or a mixture thereof, preferably 1 or 2, more preferably 1;

20 provided that the total number of carbon atoms in group (vi) is from about 8 to about 22;

Y is a linear or branched diradical having olefinic unsaturation of the formula $-C_eH_{(2e-2)}-$


wherein


e is 2 or 3, preferably 2; R is H or a linear or branched alkyl group

25 $C_bH_{(2b+1)}-$; and b is from 1 to about 18, preferably from 6 to 18.

The surfactants of Formula 1A are prepared by reacting two moles of fluoroalcohols of formula R_a-OH wherein R_a is defined as above with one mole of an unsaturated linear or branched dibasic acid of the structure $C_eH_{(2e-2)}(COOH)_2$ or its anhydride of Formula D

30





5 wherein e is 2 or 3 to form the unsaturated diester of Formula 2A as an intermediate. Methods for carboxylic acid esterification are conventional as discussed by Jain and Masse in "Carboxylic acid esters: synthesis from carboxylic acids and derivatives" in *Science of Synthesis* (2006), 20b, 711-723. An acid catalyst or dehydrating agent is preferred when reacting the free acid groups with 10 alcohols. An example of an acid catalyst is p-toluenesulfonic acid in toluene, and an example of a dehydrating agent is dicyclohexylcarbodiimide in methylene chloride. Preferred unsaturated dibasic acids and corresponding anhydrides are maleic, itaconic (methylenesuccinic acid), citraconic (methylmaleic acid), *trans*-glutaconic ($\text{HOOCCH}_2\text{CH}=\text{CHCOOH}$), and *trans-beta*-hydromuconic 15 ($\text{HOOCCH}_2\text{CH}=\text{CHCH}_2\text{COOH}$) acids and anhydrides. The unsaturated diester of Formula 2A is then reacted with aqueous sodium hydrogen sulfite to form the sulfonic acid. Sulfonation techniques are described by Roberts in "Sulfonation Technology for Anionic Surfactant Manufacture", *Organic Process Research & Development* 1998, 2, 194-202, and by Sekiguchi et al. in US Patent 4,545,939. 20 Alternatively, the olefinic precursors described above can be converted to the sulfonates of Formulae 1A, 1B, and 1C by the addition of sulfur trioxide to the double bond. The free sulfonic acid can be used as the surfactant, or the sulfonic acid can be converted to the ammonium salt, the alkali metal salt, or an alkaline earth metal salt, and preferably to the sodium salt. Those skilled in the art will 25 recognize other sulfonation methods, such as those described by Roberts and Sekiguchi (above) are applicable and are included in the present invention.

Addition of the sulfonate group across the double bond of Formulae 1A, 1B, and 1C to make the surfactants of Formulae 2A, 2B, and 2C results in the formation of stereo-isomers and regio-isomers. For the purposes of the present 30 invention, all the isomers are equivalent and all are included in the definitions of Formulae 2A, 2B, and 2C.

The surfactants of Formulae 1B and 1C are prepared by reacting one mole of a fluoroalcohol of formula $\text{R}_a\text{-OH}$ with one mole of an unsaturated linear or branched dibasic acid anhydride of the structure of Formula D at a lower 35 temperature (between about 50 - 85°C). The esterification is then continued at a higher temperature (between about 100 - 120°C) with a mole of a fluoroalcohol,

preferably of formula $R_fCH_2CH_2-OH$, to produce Formula 2B, or a mole of alcohol of formula $R-OH$, to produce Formula 2C. Any of a variety of conventional fluorinated alcohols are suitable for use at this point of the process. This is followed by conversion to the sulfonates. The anhydride is preferred in

5 the preparation of surfactants of Formulae 1B and 1C. The opening of the anhydride ring by the first esterification is a faster reaction than the second esterification of the intermediate acid ester. As indicated above, in the second esterification acid catalysts or dehydrating agents are used. Use of the dibasic acid thus tends to give mixtures of products. The sequence of use of the two

10 alcohols in the esterifications can be reversed.

The surfactants of Formula 1A can also be prepared using two additions of the same alcohol and the two-temperature procedure described for Formulae 1B and 1C, followed by conversion to the sulfonates. However, this two-step procedure is not preferred.


15 Mixtures of surfactants of compositions of Formulae 1A, 1B, and 1C can be prepared by using two moles of a mixture of two or more of R_a-OH , $R_f-CH_2CH_2-OH$, and $R-OH$ alcohols. The two moles of a mixture of two or more alcohols are reacted with one mole of an unsaturated linear or branched dibasic acid of the structure $C_eH_{(2e-2)}(COOH)_2$ or its anhydride (Formula D) as

20 described above for the preparation of surfactants of Formula 1A, followed by conversion to the sulfonates. Such surfactant mixtures can be used as is or separated into the component fractions. Such separations are not preferred.

Alcohols containing the R_a group (i) of $R_f(CH_2CF_2)_d-CH_2CH_2$ include the fluorinated telomer alcohols of formula (V):




wherein R_f is a linear or branched perfluoroalkyl group having 2 to 6 carbon atoms, and subscripts q and r are each independently integers of 1 to 3. These telomer alcohols are available by synthesis according to Scheme 1 wherein R_f , q and r are as defined for Formula (V).



Scheme 1

The telomerization of vinylidene fluoride with linear or branched perfluoroalkyl iodides produces compounds of the structure $R_f(CH_2CF_2)_q\text{I}$, wherein, q is 1 or more and R_f is a C_2 to C_6 perfluoroalkyl group. For example, see Balague, et al, "Synthesis of fluorinated telomers, Part 1, Telomerization of vinylidene fluoride with perfluoroalkyl iodides", J. Fluorine Chem. (1995), 70(2), 215-23. The specific telomer iodides are isolated by fractional distillation. The telomer iodides are treated with ethylene by procedures described in US Patent 3,979,469 to provide the telomer ethylene iodides (VI of Scheme 1) wherein r is 1 to 3 or more. The telomer ethylene iodides (VI of Scheme 1) are treated with oleum and hydrolyzed to provide the corresponding telomer alcohols (V of Scheme 1) according to procedures disclosed in WO 95/11877. Alternatively, the telomer ethylene iodides (VI of Scheme 1) can be treated with N -methyl formamide followed by ethyl alcohol/acid hydrolysis.

The R_a group of Formula (ii), $R_f\text{OCF}_2\text{CF}_2(C_g\text{H}_{2g})-$, is obtained by preparing fluoroalcohols of the formula $R_f\text{OCF}_2\text{CF}_2\text{-CH}_2\text{CH}_2\text{OH}$ which are available by the following series of reactions wherein R_f is a linear or branched C_2 to C_6 perfluoroalkyl optionally interrupted by one to three oxygen atoms and q is an integer of 1 to 3:



Scheme 2


The perfluoroalkyl ether iodides of formula (V of Scheme 2) above can be made by the procedure described in US Patent 5,481,028 in Example 8, which discloses the preparation of compounds of formula (V of Scheme 2) from perfluoro-n-propyl vinyl ether. The perfluoroalkyl ether iodide (V of Scheme 2) is 5 reacted with an excess of ethylene at an elevated temperature and pressure. While the addition of ethylene can be carried out thermally, the use of a suitable catalyst is preferred. Preferably the catalyst is a peroxide catalyst such as benzoyl peroxide, isobutyryl peroxide, propionyl peroxide, or acetyl peroxide. More preferably the peroxide catalyst is benzoyl peroxide. The temperature of the 10 reaction is not limited, but a temperature in the range of 110°C to 130°C is preferred. The reaction time can vary with the catalyst and reaction conditions, but 24 hours is usually adequate. The product is purified by any means that separates unreacted starting material from the final product, but distillation is preferred. Satisfactory yields up to 80% of theory have been obtained using about 15 2.7 mols of ethylene per mole of perfluoroalkyl ether iodide, a temperature of 110°C and autogenous pressure, a reaction time of 24 hours, and purifying the product by distillation.

The perfluoroalkylether ethylene iodides (VI of Scheme 2) are treated with oleum and hydrolyzed to provide the corresponding alcohols (VII of Scheme 2) 20 according to procedures disclosed in WO 95/11877 (Elf Atochem S.A.). Alternatively, the perfluoroalkylether ethyl iodides can be treated with N-methyl formamide followed by ethyl alcohol/acid hydrolysis. A temperature of about 130° to 160°C is preferred.

The R_a group of Formula (iii), R_fOCFHCF₂O(CH₂CH₂O)_v-(C_gH_{2g})-, is 25 prepared by reacting a fluorinated vinyl ether with a polyethylene glycol. Typically the vinyl ether is slowly added to the glycol in a molar ratio of from about 1:1 to about 3:1, preferably at about 2:1. The reaction is conducted in the presence of sodium hydride, which is a catalyst that is basic enough to generate equilibrium amounts of the alkoxide anion from the glycol. Other suitable base 30 catalysts include potassium hydride, sodium amide, lithium amide, potassium tert-butoxide, and potassium hydroxide. The reaction is conducted under an inert atmosphere such as nitrogen gas. Suitable solvents include dimethylformamide, dimethylacetamide, acetonitrile, tetrahydrofuran, and dioxane. Preferred is dimethylformamide. Cooling is employed to maintain the reaction temperature at 35 from about 0°C to about 30°C. The reaction is usually conducted for 1 to about 18 hours. The solvent is then removed using conventional techniques; such as evaporation in vacuum on a rotary evaporator, or in cases where the product is

water insoluble and the solvent is water soluble, addition of the mixture to an excess of water followed by separation of the layers.

The reaction of perfluoropropyl vinyl ether with polyethylene glycol does not always go to completion. The average degree of conversion of the 5 polyethylene glycol hydroxyl groups can be determined by ^1H NMR spectroscopy. Typically mixtures of unreacted polyethylene glycol, the product of fluorinated vinyl ether adding to one end of polyethylene glycol (for example, structure B below), and the product of fluorinated vinyl ether adding to both ends of the polyethylene glycol (for example, structure A below) can be obtained. The 10 relative amounts of the components of the mixture are affected by the ratio of the reactants, the reaction conditions, and the way in which the product is isolated. High ratios of the vinyl ether to glycol and long reaction times tend to favor Structure A, shown below. Lower ratios of vinyl ether to glycol and shorter reaction times give increased amounts of Structure B, shown below, and 15 unreacted polyethylene glycol. It is sometimes possible to use the differences in solubility between Structures A, B, and the starting glycol to do selective solvent extraction of mixtures to obtain samples that are highly enriched in Structures A or B. The alcohol of Structure B is a required composition for the group R_a (iii).




Polyethylene glycols suitable for the use are commercially available from Sigma-Aldrich, Milwaukee, WI. The fluorinated vinyl ether used in the above reaction is made by various methods. These methods include making fluorinated vinyl ethers by reacting a 2-alkoxypropionyl fluoride in a stationary bed of a 25 metal carbonate, a tubular reactor filled with dried metal carbonate and equipped with a screw blade running through the tube, or a fluidized bed of metal carbonate. US Patent Application 2007/0004938 describes a process to produce fluorinated vinyl ether by reacting a 2-alkoxypropionyl fluoride with a metal carbonate under anhydrous conditions in a stirred bed reactor at a temperature 30 above the decarboxylation temperature of an intermediate carboxylate to produce fluorinated vinyl ether. Examples of fluorinated vinyl ethers suitable for use include $\text{CF}_3\text{-O-CF=CF}_2$, $\text{CF}_3\text{CF}_2\text{-O-CF=CF}_2$, $\text{CF}_3\text{CF}_2\text{CF}_2\text{-O-CF=CF}_2$, and $\text{CF}_3\text{CF}_2\text{CF}_2\text{CF}_2\text{-O-CF=CF}_2$, each of which are available from E. I. du Pont de Nemours and Company, Wilmington, DE.

35 The R_a group of Formula (iv) $\text{R}_f\text{OCFHC}\text{F}_2\text{O}(\text{C}_w\text{H}_{2w})$ - wherein w is from about 3 to about 12, is prepared by the reaction of a

perfluorohydrocarbonvinyl ether with a diol in the presence of an alkali metal compound. Preferred ethers include those of formula R_f -O-CF=CF₂ wherein R_f is a perfluoroalkyl of two to six carbons. The diol is used at about 1 to about 15 mols per mol of ether, preferably from about 1 to 5 about 5 mols per mol of ether. Suitable alkali metal compounds include an alkali metal, alkali earth metal, alkali hydroxide, alkali hydride, or an alkali amide. Preferred are alkali metals such as Na, K or Cs, or alkali hydrides such as NaH or KH. The reaction is conducted at a temperature of from about 40°C to about 120°C. The reaction can be conducted in an optional solvent, such as ether or nitrile.

The R_a group of Formula (v), R_f OCF(CF₃)CONH-CH₂CH₂-⁻, is prepared by making a fluorinated alcohol of Formula 4:



wherein

20 R_f is a straight or branched perfluoroalkyl group having from about 2 to about 6 carbon atoms, or a mixture thereof;

X^3 is oxygen or X^1 ;

each X^1 is independently an organic divalent linking group having from about 1 to about 20 carbon atoms, optionally containing an oxygen, nitrogen, or sulfur, or a combination thereof;

25 G is F or CF₃; A is an amide; j' is zero or positive integer; X² is an organic linking group; h' is zero or one; B is H; and E is hydroxyl.

The compound of Formula 4 is prepared by reaction between a perfluorinated ester (prepared according to reported methods in US Patent 6,054,615 and US Patent 6,376,705 with a triamine or diamine alcohol with or without solvent. The conditions of this reaction are dependent on structure of the ester. The reaction of alpha, alpha-difluorosubstituted ester with diamine is reacted at a temperature of from about 5°C to about 35°C. Suitable solvents for this reaction include tetrahydrofuran,

methyl isobutyl ketone, acetone, CHCl_3 , CH_2Cl_2 , or ether. The reaction of ester without alpha-fluorine substitution with diamine is conducted at a temperature of from about 90°C to about 160°C, preferably at between about 100°C to about 140°C. Preferably no solvent is employed for this reaction, but suitable solvents 5 include chlorobenzene, dimethylformamide, or 2-methoxyethyl ether.

The compound of Formula 4 is also prepared by reaction between a perfluorinated acyl fluoride with a diamine alcohol or amine alcohol. This reaction is conducted at a temperature of from about -30°C to about 40°C, preferably at between about 5°C to about 25°C. Suitable solvents for this reaction 10 include tetrahydrofuran, methyl isobutyl ketone, acetone, CHCl_3 , CH_2Cl_2 , or 2-methoxyethyl ether, diethyl ether.



The R_a group of Formula (vi) above, $R_f(\text{CH}_2)_h[(\text{CF}_2\text{CF}_2)_i(\text{CH}_2\text{CH}_2)_j]_k$, is obtained by preparation of fluoroalcohols of the formula $R_f(\text{CH}_2)_h[(\text{CF}_2\text{CF}_2)_i(\text{CH}_2\text{CH}_2)_j]_k\text{OH}$, wherein R_f is a C_2 to C_6 perfluoroalkyl, subscript h is 1 to 15 about 6, and subscripts i , j , and k are each independently 1, 2, 3, or a mixture thereof. These alcohols are prepared from oligomeric iodides ($\text{C}_n\text{F}_{2n+1}\text{C}_2\text{H}_4\text{I}$, $\text{C}_n\text{F}_{2n+1}\text{CH}_2\text{I}$ or $\text{C}_n\text{F}_{2n+1}\text{I}$) wherein subscript n is an integer from 1 to about 6, using an oleum treatment and hydrolysis. It has been found, for example, that reacting with oleum (15% SO_3) at about 60°C for about 1.5 hours, followed by 20 hydrolysis using an iced dilute K_2SO_3 solution, and then followed by heating to about 100°C for about 30 minutes gives satisfactory results. But other reaction conditions can also be used. After being cooled to ambient room temperature, a solid is precipitated, isolated and purified. For example, the liquid is then 25 decanted and the solid is dissolved in ether and washed with water saturated with NaCl , dried over anhydrous Na_2SO_4 , and concentrated and dried under vacuum. Other conventional purification procedures can be employed.

Alternatively, the alcohols of formula $R_f(\text{CH}_2)_h[(\text{CF}_2\text{CF}_2)_i(\text{CH}_2\text{CH}_2)_j]_k\text{OH}$ as defined above can be prepared by heating the oligomeric iodides $R_f(\text{CH}_2)_h[(\text{CF}_2\text{CF}_2)_i(\text{CH}_2\text{CH}_2)_j]_k\text{I}$ wherein R_f , and subscripts h , i , j , and 30 k are as defined above for the corresponding alcohol, with N -methylformamide to about 150°C and holding for about 19 hours. The reaction mixture is washed with water to give a residue. A mixture of this residue with ethanol and concentrated hydrochloric acid is gently refluxed (at about 85°C bath temperature) for about 2.5 hours. The reaction mixture is washed with water, diluted with 35 dichloromethane, and dried over sodium sulfate. The dichloromethane solution is concentrated and distilled at reduced pressure to give the alcohol. Optionally

N,N-dimethylformamide can be used instead of N-methylformamide. Other conventional purification procedures can also be employed.

The iodides of formula $R_f(CH_2)_h[(CF_2CF_2)_i(CH_2CH_2)_j]_kI$ are preferably prepared by oligomerization of $C_nF_{2n+1}C_2H_4I$, $C_nF_{2n+1}CH_2I$ or $C_nF_{2n+1}I$ wherein n is 1 to about 6 using a mixture of ethylene and tetrafluoroethylene. The reaction can be conducted at any temperature from room temperature to about 150°C with a suitable radical initiator. Preferably the reaction is conducted at a temperature of from about 40° to about 100°C with an initiator which has about a 10 hour half-life in that range. The feed ratio of the starting materials in the gas phase, that is the moles of $C_nF_{2n+1}C_2H_4I$, $C_nF_{2n+1}CH_2I$ or $C_nF_{2n+1}I$ wherein n is 1 to about 6, versus the combined moles of ethylene and tetrafluoroethylene, can be used to control conversion of the reaction. This mole ratio is from about 1:3 to about 20:1, preferably from about 1:2 to 10:1, more preferably from about 1:2 to about 5:1. The mole ratio of ethylene to tetrafluoroethylene is from about 1:10 to about 10:1, preferably from about 3:7 to about 7:3, and more preferably from about 4:6 to about 6:4.

The unsaturated intermediates used in the preparation of the surfactants of Formulae 1A, 1B, and 1C are formed prior to the addition of the sulfonic acid group. The unsaturated intermediates have the structure of Formulae 2A, 2B, and 2C:




wherein R_a is the group (i) $R_f(CH_2CF_2)_d-(C_gH_{2g})-$; (ii) $R_fOCF_2CF_2-(C_gH_{2g})-$; (iii) $R_fOCFHCF_2O(CH_2CH_2O)_v-(C_gH_{2g})-$; (iv) $R_fOCFHCF_2O(C_wH_{2w})-$; (v) $R_fOCF(CF_3)CONH-(C_gH_{2g})-$; or (vi) $R_f(CH_2)_h[(CF_2CF_2)_i(CH_2CH_2)_j]_k$

each R_f is independently $C_cF_{(2c+1)}$;
 c is 2 to about 6, preferably from 2 to 4, more preferably 4;
 30 d is 1 to about 3, preferably from 1 to 2, more preferably 1;
 g is 1 to 4, preferably from 1 to 3, more preferably 2;
 v is 1 to about 4, preferably from 2 to 3, more preferably 2;
 w is from about 3 to about 12, preferably from 4 to 6, more preferably 4;

h is 1 to about 6, preferably from 1 to 3, more preferably 2;
 i, j, and k are each independently 1, 2, or 3, or a mixture thereof,
 preferably 1 or 2, more preferably 1;
 provided that the total number of carbon atoms in group (vi) is from about
 5 8 to about 22;

Y is a linear or branched diradical having olefinic unsaturation of the
 formula



wherein e is 2 or 3, preferably 2; R is H or a linear or branched alkyl group
 10 C_bH_(2b+1)-; and b is from 1 to about 18, preferably from 6 to 18.

Compounds of Formulae 2A, 2B, and 2C are prepared as discussed above
 for Formulae 1A, 1B, and 1C except that the sulfonation step is omitted.
 Compounds of Formula 2A, 2B, and 2C are also monomers that can be
 polymerized alone or in admixture with other monomers to confer soil and water
 15 repellency to the resulting polymers and to surfaces to which the resulting
 polymers are applied.

Preferred compounds of Formula 2A, 2B, and 2C are those wherein R_a is
 R_f(CH₂CF₂)_d-(C_gH_{2g})-; R_fOCF₂CF₂-(C_gH_{2g})-; R_fOCFHCF₂O-
 (CH₂CH₂O)_v-(C_gH_{2g})-; or R_fOCFHCF₂O(C_wH_{2w}O)-(C_gH_{2g})-. Also
 20 preferred are those compounds of Formula 2A, 2B and 2C wherein c is 3 or 4, or
 wherein Y is CH=CH, CH₂C(=CH₂), C(CH₃)=CH₂, CH=CHCH₂, or
 CH₂CH=CHCH₂. More preferred are those compounds wherein R_a is
 R_f(CH₂CF₂)_d-(C_gH_{2g})-; R_fOCF₂CF₂-(C_gH_{2g})-; R_fOCFHCF₂O-
 (CH₂CH₂O)_v-(C_gH_{2g})-; or R_fOCFHCF₂O(C_wH_{2w}O)-(C_gH_{2g})-; d is 1, g is 1,
 25 R_f is C₃F₇ or C₄F₉, and Y is CH=CH, CH₂C(=CH₂), or C(CH₃)=CH₂. Also
 preferred are compounds of Formula 2B wherein R_a is C₄F₉CH₂CF₂CH₂CH₂ or
 C₃F₇CH₂CF₂CH₂CH₂ and R_f is (CF₂)₆F.

The compounds of Formula 2A, 2B and 2C are useful as intermediates to
 prepare partially fluorinated sulfonated surfactants, in particular those of Formula
 30 1A, 1B and 1C as previously defined.

A method of altering the surface behavior of a liquid comprises adding to
 the liquid a compound of Formulae 1A, 1B, and 1C, as defined above, in a wide
 variety of applications. The surfactants of Formula 1A, 1B, and 1C are typically
 used by simply blending with or adding to water, aqueous solutions, and aqueous
 35 emulsions. The surfactants of Formulae

1A, 1B, and 1C typically lower surface and interfacial tensions and provide low critical micelle concentrations. Examples of surface behavior alteration include improvements in the properties of wetting, penetration, spreading, leveling, flowing, emulsifying, stabilization of dispersions in liquids, repellency, releasing, 5 lubricating, etching, and bonding.

Examples of such applications where low surface tension is required include coating compositions and aqueous and non-aqueous cleaning products, each for glass, wood, metal, brick, concrete, cement, natural and synthetic stone, tile, synthetic flooring, laminates, paper, textile materials, linoleum and other 10 plastics, resins, natural and synthetic rubbers, fibers and fabrics, and paints; polymers; and waxes, finishes, leveling and gloss agents for floors, furniture, shoes, inks, and automotive care. Wetting agent applications include wetting agents for compositions containing herbicides, fungicides, weed killers, hormone growth regulators, parasiticides, insecticides, germicides, bactericides, 15 nematocides, microbiocides, defoliants or fertilizers, therapeutic agents, antimicrobials, fluorochemical blood substitutes, textile treatment baths, and fiber spin finishes. Applications in personal care products include shampoos, conditioners, creme rinses, cosmetic products for the skin (such as therapeutic or protective creams and lotions, oil and water repellent cosmetic powders, 20 deodorants and antiperspirants), nail polish, lipstick, and toothpaste. Further applications include fabric care products (such as stain pretreatments and/or stain removers for clothing, carpets and upholstery), and laundry detergents. Other applications include rinse-aids (for car washes and in automatic dishwashers), for 25 oil well treatments (including drilling muds and additives to improve tertiary oil well recovery), extreme pressure lubricants, lubricating cutting oil to improve penetration times, writing inks, printing inks, photography developer solutions, emulsions for fighting forest fires, dry chemical fire extinguishing agents, aerosol-type fire extinguishers, thickening agents to form gels for solidifying or encapsulating medical waste, photoresists, developers, cleaning solutions, etching 30 compositions, developers, polishers, and resist inks in the manufacturing, processing, and handling of semiconductors and electronics. The surfactants of Formulae 1A, 1B, and 1C can be incorporated into products that function as antifogging agents for glass surfaces and photography films, and as antistatic agents for magnetic tapes, phonograph records, floppy disks, disk drives, rubber 35 compositions, PVC, polyester film, and photography films, and as surface treatments for optical elements (such as glass, plastic, or ceramics). Other applications are in emulsifying agents, foaming agents, release agents, repellency

agents, flow modifiers, film evaporation inhibitors, wetting agents, penetrating agents, cleaners, grinding agents, electroplating agents, corrosion inhibitors, soldering agents, dispersion aids, microbial agents, pulping aids, rinsing aids, polishing agents, drying agents, antistatic agents, antiblocking agents, bonding agents, and oil field chemicals.

The compounds of Formulae 1A, 1B, and 1C are also useful as foam control agents in polyurethane foams, spray-on oven cleaners, foamed kitchen and bathroom cleansers and disinfectants, aerosol shaving foams, and in textile treatment baths. The surfactants of Formulae 1A, 1B, and 1C are useful as emulsifying agents for polymerization, particularly of fluoromonomers, as latex stabilizers, as mold release agents for silicones, photoemulsion stabilizers, inorganic particles, and pigments. Such fluorosurfactants are also useful for supercritical carbon dioxide emulsions and dispersion of nanoparticles or pigments in water.

A low concentration of less than about 0.1%, preferably less than about 0.01% by weight of a compound of Formulae 1A, 1B, or 1C in the liquid is effective. Consequently, the surfactants of Formulae 1A, 1B, and 1C are useful in a wide variety of end use applications.

The present invention comprises compounds of Formula 5



wherein

R_f is $C_cF_{(2c+1)}$;

c is 2 to about 6, preferably from 2 to 4, more preferably 4; and

v is 2 to about 4, preferably from 2 to 3, more preferably 2.

Preferred compounds of Formula 5 are those wherein c is 3 or 4, g is 2, and v is 2 or 3. The compounds of Formula 5 are useful as intermediates in making partially fluorinated sulfonated surfactants. In particular, Formula 5 compounds are useful in making surfactants of Formula 1A, 1B and 1C as previously defined.

30 The present invention further comprises a process for the preparation of a compound of Formula 5



wherein

R_f is $C_c F_{(2c+1)}$; c is 2 to about 6; and v is 2 to about 4,

comprising contacting a compound of Formula 6



wherein R_f is $C_cF_{(2c+1)}$, and c is 2 to about 6,

with a compound of Formula 7



wherein v is 2 to about 4.

In the process of the present invention the compound of Formula 5 is prepared by the reaction of a perfluorohydrocarbonvinyl ether with a diol in the presence of an alkali metal compound. Preferred ethers include those of 10 formula $R_f-O-CF=CF_2$ wherein R_f is a perfluoroalkyl of one to six carbons. Preferred diols include diethylene glycol. The diol is used at about 1 to about 15 mols per mol of ether, preferably from about 1 to about 5 mols per mol of ether. Suitable alkali metal compounds include an alkali metal, alkali earth metal, alkali hydroxide, alkali hydride, or an alkali amide. Preferred are 15 alkali metals such as Na, K or Cs or alkali hydrides such as NaH or KH. The reaction is conducted at a temperature of from about ambient temperature to about 120°C, preferably from about 40°C to about 120°C. The reaction can be conducted in an optional solvent, such as ether or nitrile. The process is useful to prepare alcohols of Formula 5 which are used to make derivatives, 20 such as surfactants.

The surfactants of Formula 1A, 1B, and 1C, in many cases, require less tetrafluoroethylene in their preparation when compared to conventional fluorosurfactants made from Telomer B alcohols. While tetrafluoroethylene may be used in the R_f portion of the R_a-OH alcohol precursors when R_a is (ii) 25 $R_fOCF_2CF_2-(C_gH_{2g})-$; (iii) $R_fOCFHCFC_2O(CH_2CH_2O)_v-(C_gH_{2g})-$; (iv) $R_fOCFHCFC_2O(C_wH_{2w})-$; or (v) $R_fOCF(CF_3)CONH-(C_gH_{2g})$; tetrafluoroethylene is not otherwise used in the preparation of the compounds of Formula 1A, 1B or 1C or of Formula 2A, 2B or 2C. For the surfactants of Formulac 1A, 1B, and 1C, some of the fluorine is replaced with other atoms or 30 monomers, compared to the typical perfluoroalkyl groups of 1 to 20 carbons in surfactants made from traditional Telomer B alcohols. So less tetrafluoroethylene is used in the preparation of compounds of Formula 1A, 1B and 1C or of Formula 2A, 2B and 2C containing the R_a (i) and (vi) groups.

The monomer moieties replacing tetrafluoroethylene in most cases also 35 contain a lower proportion of fluorine. Consequently, in many cases the

surfactants of Formulae 1A, 1B, and 1C are more fluorine efficient than many conventional surfactants. By "fluorine efficiency" is meant the ability to use a minimum amount of fluorochemical to obtain a desired surface effect or surfactant properties, when applied to substrates, or to obtain better performance 5 using the same level of fluorine.

MATERIALS AND TEST METHODS

The following materials and test methods were used in the Examples herein.

All common organic and inorganic compounds were obtained from 10 Sigma-Aldrich (Milwaukee WI) and used without purification. These included maleic anhydride, sodium hydrogen sulfite, toluene, hexane, p-toluene sulfonic acid, itaconic anhydride, citraconic anhydride, *trans*-glutaconic acid, *trans-beta*-hydromuconic acid, and other routine compounds employed in the Examples.

SIMULSOL SL8: octyl/decyl polyglucoside is available from Kreglinger 15 Europe, Antwerp, Belgium.

TRITON X100: *p-tert*-octylphenoxy polyethyl alcohol is available from Sigma-Aldrich, Saint Louis, MO.

DOWANOL DB: 1-butoxy-2-ethoxyethane is available from Dow Chemical Company, Midland, MI.

20 SOLKANE 365 MFC is 1,1,1,3,3-pentafluorobutane is available from Solvay Fluorides, Thorofare NJ.

The following fluorinated chemicals are available from E. I. du Pont de Nemours and Company, Wilmington DE: Perfluoro-2-methyl-3-oxahexanoyl 25 fluoride; Perfluorobutyl iodide; Vinylidene fluoride; Perfluoropropylvinyl ether; and Perfluoroethylmethylethyl iodide.

The following fluorinated chemicals were prepared as indicated below:

30 $C_4F_9CH_2CF_2I$ and $C_4F_9(CH_2CF_2)_2I$ were prepared by reacting perfluorobutyl iodide and vinylidene fluoride as described by Balague, et al, "Synthesis of Fluorinated Telomers, Part 1, Telomerization of Vinylidene Fluoride with Perfluoroalkyl Iodides", J. Fluorine Chem. (1995), 70(2), 215-23. The specific telomer iodides are isolated by fractional distillation.

$C_3F_7OCF_2CF_2I$ was prepared by reacting perfluoropropyl vinyl ether with iodine chloride and hydrofluoric acid with boron trifluoride as a catalyst as described by Viacheslav et al. in US 5,481,028.

Test Method 1 - Measurement of the Critical Micelle Concentration (CMC) and the Surface Tension beyond CMC.

Surface tensions of aqueous surfactant solutions were measured at various weight percents in mN/m using a Kruss K11 Tensiometer (from Kruss USA, 5 Charlotte, NC). Compounds having the lowest surface tension have the highest effectiveness.

The critical micelle concentration (CMC) is defined as the concentration at which increased concentrations of surfactant essentially no longer reduce the surface tension. To determine CMC, the surface tension is measured as a function 10 of surfactant concentration. Surface tension is then plotted (abscissa) vs. log concentration (ordinate). The resulting curve has a nearly horizontal portion at concentrations higher than the CMC and has a negative steep slope at concentrations less than the CMC. The CMC is the concentration at the intersection of the extrapolated steep slope and the extrapolated near horizontal 15 line. The Surface Tension beyond CMC is the value in the flat portion of the curve. The CMC should be as low as possible to provide the lowest cost for effective performance.

Test Method 2 - Spreading on Cyclohexane

Test Method 2 is adapted from Stern et al. in WO1997046283A1, wherein 20 surfactants were applied to the surface of n-heptane to provide a screening evaluation for Advance Fire Fighting Foams (AFFF). Cyclohexane was used in Test Method 2 to replace the n-heptane used by Stern et al. Test Method 2 measures the ability of the surfactant solution to spread across the surface of a less 25 dense flammable liquid. When the surfactant solution spreads across the surface (“excellent” rating), a barrier is established between the flammable liquid and the air. If the surfactant solution does not spread completely across the surface (“good” or “fair” rating depending on the extent of the partial spreading) the barrier between air and flammable liquid is incomplete. If the surfactant solution sinks into the flammable liquid (“poor” rating), no barrier between air and 30 flammable liquid is established.

A surfactant solution was prepared by combining fluorosurfactant (0.9 g/L of active ingredient), hydrocarbon surfactant (either SIMULSOL SL8 or TRITON X100; 2.4 g/L of active ingredient), butyl carbitol (DOWANOL DB; 4.2 g/L of active ingredient), and mixed thoroughly. A Petri dish (11.5 cm diameter) was 35 filled about half-way with 75 mL of cyclohexane. After the surface of the cyclohexane had completely calmed (about 1 minute), 100 microliters of the

solution of fluorosurfactant, hydrocarbon surfactant, butyl carbitol, and water was deposited dropwise with a micropipette beginning at the center of the Petri dish and outwardly along a radial line to the outer edge of the Petri dish. The timer was started. In poor performing formulations, the surfactant solution "sinks" 5 immediately" below the cyclohexane. In fair performing formulations, the surfactant solution merely "floats" on the surface of the cyclohexane without sinking. In good performing formulations, the surfactant spreads across the surface of the cyclohexane. The time was noted when the extent of the spreading of the surfactant solution across the surface of the cyclohexane ceased and the 10 extent of the surface coverage (<100%) at that point was recorded. In excellent performing formulations, the surfactant solution rapidly spreads across the entire surface of the cyclohexane. In excellent performing formulations, the time was noted when extent of the spreading of the surfactant solution first covered the entire surface (100%).

15

EXAMPLES

Example 1

A mixture of ethanolamine (13 g, 28 mmol) and ether (30 mL) was cooled to 15°C. Perfluoro-2-methyl-3-oxahexanoyl fluoride (33 g in ether 50 mL) was added dropwise to keep the reaction temperature below 25°C. After the addition, 20 the reaction mixture was stirred at room temperature for one hour. The solid was removed by filtration and the filtrate was washed with hydrochloric acid (0.5N, 30 mL), water (2 times 30 mL), sodium hydrogen carbonate solution (0.5N, 20 mL), water (30 mL), and sodium chloride solution (saturated, 20 mL). It was then concentrated and dried in vacuum over night at room temperature to give a white 25 solid 35 g, yield 95%. The product was analyzed using ^1H NMR and the structure confirmed as is N-(perfluoro-2-methyl-3-oxahexanoyl)-2-aminoethanol, $\text{C}_3\text{F}_7\text{OCF}(\text{CF}_3)\text{CONHCH}_2\text{CH}_2\text{OH}$.

A mixture of maleic anhydride (0.60 g, 6.1 mmol), $\text{C}_3\text{F}_7\text{OCF}(\text{CF}_3)\text{CONHCH}_2\text{CH}_2\text{OH}$ (4.5 g, 12 mmol, prepared as described 30 above), *p*-toluenesulfonic acid monohydrate (0.12 g) and toluene (50 mL) was stirred continuously and heated to reflux under nitrogen. The temperature was maintained at 111°C for approximately 22 h until 90% of water was removed azeotropically with the aid of a Dean-Stark trap. A liquid chromatography/mass spectrum (LC/MS) was taken to show the completion to the diester. The solution 35 was separated and extracted with two washings of 5% sodium bicarbonate solution. The combined organic extracts were dried over anhydrous magnesium

sulfate ($MgSO_4$) and then toluene was removed by rotary evaporation. The yellow oil (3.12 g, 61.9% yield, 90% purity) was analyzed by 1H NMR and LC/MS to confirm the structure as $C_3F_7OCF(CF_3)C(O)NHCH_2CH_2OC(O)CH=CHC(O)OCH_2CH_2NHC(O)-CF(CF_3)OC_3F_7$.

5 Example 2

Ethylene (25 g) was introduced to an autoclave charged with $C_4F_9CH_2CF_2I$ (217 g) and d-(+)-limonene (1 g), and the reactor heated at 240°C for 12 hours. The product was isolated by vacuum distillation to provide $C_4F_9CH_2CF_2CH_2CH_2I$. Fuming sulfuric acid (70mL) was added slowly to 50 g of $C_4F_9CH_2CF_2CH_2CH_2I$ and mixture was stirred at 60 °C for 1.5 hours. The reaction was quenched with ice-cold 1.5 wt% Na_2SO_3 aqueous solution and heated at 95 °C for 0.5 hours. The bottom layer was separated and washed with 10 wt% aqueous sodium acetate and distilled to provide $C_4F_9CH_2CF_2CH_2CH_2OH$ bp 54-57 °C at 2 mmHg (267 Pa).

15 The esterification procedure of Example 1 was used to make Di(1H,1H,2H,2H,4H,4H-perfluorooctyl) maleate (7.76 g, 95% yield, 95% purity) by the reaction of maleic anhydride (1.07 g, 11 mmol). $C_4F_9CH_2CF_2CH_2CH_2OH$ (7.13 g, 22 mmol, prepared as described above) and *p*-toluenesulfonic acid monohydrate (0.21 g, 1.1 mmol) in 50 mL of toluene at 20 111°C for 40 h. The pale yellow product was analyzed by 1H NMR and LC/MS to confirm the structure as $C_4F_9CH_2CF_2CH_2CH_2OC(O)-CH=CH-C(O)OCH_2CH_2CF_2CH_2C_4F_9$.

Example 3

Ethylene (56 g) was introduced to an autoclave charged with $C_4F_9(CH_2CF_2)_2I$ (714 g) and d-(+)-limonene (3.2 g), and the reactor heated at 240 °C for 12 hours. The product was isolated by vacuum distillation to provide $C_4F_9(CH_2CF_2)_2CH_2CH_2I$. A mixture of $C_4F_9(CH_2CF_2)_2CH_2CH_2I$ (10 g, 0.02 mol) and N-methylformamide (8.9 mL, 0.15 mol) was heated to 150 °C for 26 hours. The mixture was cooled to 100 °C, followed by the addition of water to 30 separate the crude ester. Ethyl alcohol (3 mL) and p-toluene sulfonic acid (0.09 g) were added and the mixture stirred at 70 °C for 0.25 hours. Ethyl formate and ethyl alcohol were removed by distillation to give a crude product. The crude product was dissolved in ether, washed with 10 wt % aqueous sodium sulfite, water and brine, in turn, and dried over magnesium sulfate. Distillation provided 35 the product $C_4F_9(CH_2CF_2)_2CH_2CH_2OH$ (6.5 g, 83 % yield): bp 94-95 °C at 2 mm Hg (266 Pa).

Maleic anhydride (0.65 g, 6.7 mmol), C₄F₉CH₂CF₂CH₂CF₂CH₂OH (4.37 g, 1.333*10⁻² mol, prepared as described above), *p*-toluenesulfonic acid monohydrate (0.13 g, 0.67 mmol) and toluene (50mL) were mixed together and heated to reflux at 110°C for 48 h. The work-up was carried out as in Example 1.

5 The resulting pale yellow liquid (2.90 g, 51.4% yield, >99% purity) was analyzed by ¹H NMR and LC/MS to confirm the structure as

C₄F₉CH₂CF₂CH₂CF₂CH₂CH₂O-C(O)CH=CH-

C(O)OCH₂CH₂CF₂CH₂CF₂CH₂C₄F₉.

Example 4

10 C₃F₇OCF₂CF₂I (100 g, 0.24 mol) and benzoyl peroxide (3 g) were charged to a pressure vessel under nitrogen. A series of three vacuum/nitrogen gas sequences was then executed at -50 °C and ethylene (18 g, 0.64 mol) was introduced. The vessel was heated for 24 hour at 110 °C. The autoclave was cooled to 0 °C and opened after degassing. Then the product was collected in a

15 bottle. The product was distilled giving 80 g of C₃F₇OCF₂CF₂CH₂CH₂I in 80% yield. The boiling point was 56~60°C at 25 mm Hg (3.3 kPa).

20 A mixture of C₃F₇OCF₂CF₂CH₂CH₂I (300 g, 0.68 mol, prepared as described above) and N-methyl-formamide (300 mL), was heated to 150 °C for 26 h. Then the reaction was cooled to 100 °C, followed by the addition of water to separate the crude ester. Ethyl alcohol (77 mL) and *p*-toluene sulfonic acid (2.59 g) were added to the crude ester, and the reaction was stirred at 70 °C for 15 minutes. Then ethyl formate and ethyl alcohol were distilled out to give a crude product. The crude product was dissolved in ether, washed with aqueous sodium sulfite, water, and brine in turn, then dried over magnesium sulfate. The product 25 was then distilled to give 199 g of C₃F₇OCF₂CF₂CH₂CH₂OH in 85 % yield. The boiling point was 71~73 °C at 40 mm Hg (5333 Pa).

30 A procedure as Example 1 was conducted. Maleic anhydride (0.66 g, 6.8 mmol), C₃F₇OCF₂CF₂CH₂CH₂OH (4.46 g, 14 mmol, prepared as described above), *p*-toluenesulfonic acid monohydrate (0.13 g, 0.68 mmol) and toluene (50mL) were mixed together and refluxed for 50 h at 112°C. The pale yellow crude product (4.12 g, 82.4%, >99% purity) was analyzed by ¹H NMR and LC/MS to confirm the structure as C₃F₇OCF₂CF₂-

CH₂CH₂OC(O)CH=CHC(O)OCH₂CH₂NHC(O)CF(CF₃)OC₃F₇.

Example 5

35 In a dry box, a 500 mL Pyrex bottle was charged with diethylene glycol (99%, Aldrich Chemical Company) (175 mL, 1.84 mole) and 80 mL of

anhydrous tetrahydrofuran (Aldrich Sure/SealTM). NaH (3.90 g, 0.163 mole) was added slowly with magnetic stirring until the completion of hydrogen evolution. The capped bottle was removed from the drybox, and the solution was transferred to a 400 mL metal shaker tube in a nitrogen filled glovebag. The shaker tube was 5 cooled to an internal temperature of -18 °C, shaking was started, and perfluoropropylvinyl ether (PPVE, 41 g 0.145 mole) was added from a metal cylinder. The mixture was allowed to warm to room temperature and was shaken for 20 h. The reaction mixture was combined with a duplicate reaction run in a separate 400 mL shaker tube. The combined reaction mixtures were added to 600 10 mL of water, and this mixture was extracted with 3 x 200 mL of diethyl ether in a separatory funnel. The ether extracts were dried over MgSO₄, filtered, and concentrated *in vacuo* on a rotary evaporator to give a liquid (119.0 g) ¹H NMR in CD₃OD, and analysis by gas chromatography both showed a small amount of diethylene glycol. This material was dissolved in 150 mL of diethyl ether and 15 extracted with water (3 x 150 mL) in a separatory funnel. The ether layer was dried over MgSO₄, filtered, and concentrated *in vacuo* on a rotary evaporator at high vacuum to give a liquid (99.1 g) ¹H NMR (C₆D₆, ppm downfield of TMS) shows 97 mole% desired mono-PPVE adduct: 1.77 (broad s, OH), 3.08-3.12 (m, OCH₂CH₂OCH₂CH₂OH), 3.42 (t, OCH₂CH₂OCH₂CH₂OH), 3.61 (t, 20 OCH₂CH₂OCH₂CH₂OH), 5.496 (doublet of triplets, ²J_{H-F} = 53 Hz, ³J_{H-F} = 3 Hz OCF₂CHFOC₃F₇), and 3 mole% of the bis PPVE adduct: 5.470 (doublet of triplets, ²J_{H-F} = 53 Hz, ³J_{H-F} = 3 Hz, C₃F₇OCHFCF₂OCH₂CH₂OCH₂CH₂OCF₂CHFOC₃F₇) The other peaks for the bis PPVE adduct overlap with the mono PPVE adduct.

25 A mixture of maleic anhydride (0.59 g, 6.1 mmol), C₃F₇OCHFCF₂OCH₂CH₂OCH₂CH₂OH (4.5 g, 12 mmol, prepared as above, *p*-toluenesulfonic acid monohydrate (0.12 g, 0.61 mmol) and toluene (50mL) were stirred continuously together and heated to reflux at 114°C for a period of 25 h. The reaction was confirmed to be completed through LC/MS and the removal of 30 water. The work-up as in Example 1 was carried out to produce a pale yellow liquid (4.48 g, 90.0% yield, 87% purity. ¹H NMR and LC/MS were used to confirm the complete conversion to the diester and the structure as C₃F₇OCFHCF₂OCH₂CH₂-OCH₂CH₂OC(O)CH=CHC(O)OCH₂CH₂O-CH₂CH₂OCF₂CFHOC₃F₇.

35 Example 6

A one-gallon reactor was charged with perfluoroethyl iodide (850 g). After cool evacuation, ethylene and tetrafluoroethylene in a ratio of 27:73 were

added until pressure reached 60 psig (414 kPa). The reaction was then heated to 70°C. More ethylene and tetrahydrofuran in a 27:73 ratio were added until pressure reached 160 psig (1.205 MPa). A lauroyl peroxide solution (4g lauroyl peroxide in 150 g perfluoroethyl ethyl iodide) was added at a 1 mL/min. rate for 1 5 hour. Gas feed ratio was adjusted to 1:1 of ethylene and tetrafluoroethylene and the pressure was kept at 160 psig (1.205 MPa). After about 67g of ethylene was added, both ethylene and tetrafluoroethylene feeds were stopped. The reaction was heated at 70°C for another 8 hours. The volatiles were removed by vacuum distillation at room temperature. A solid of oligomer ethylene-tetrafluoroethylene 10 iodides $C_2F_5(CH_2)_2[(CF_2CF_2)(CH_2CH_2)]_k-I$ (773g) wherein k is a mixture of 2 and 3 in about a 2:1 ratio was obtained.

An oligomer iodide mixture, prepared as described above (46.5 g) without separation of the iodides was mixed with N-methylformamide (NMF, 273 mL) and heated to 150°C for 19 h. The reaction mixture was washed with water (4 X 15 500 mL) to give a residue. A mixture of this residue, ethanol (200 mL), and concentrated hydrochloric acid (1 mL) was gently refluxed (85°C bath temperature) for 24 h. The reaction mixture was poured into water (300 mL). The solid was washed with water (2 X 75 mL) and dried under vacuum (2 torr, 267 Pa) to give a solid, 24.5 g. About 2g of product was sublimed. The total 20 yield of oligomer alcohols $C_2H_5(CH_2)_h[(CF_2CF_2)(CH_2CH_2)]_k-OH$ wherein k is a mixture of 2 and 3 in about a 2:1 ratio was 26.5 g.

A mixture of maleic anhydride (1.74 g, 18 mmol) and $C_2F_5(CH_2)_2[(CF_2CF_2)(CH_2CH_2)]_k-OH$ (6.26 g) were stirred continuously together and heated to 70°C. The reaction was carried out neat for 45 h and a gas 25 chromatogram (GC) was taken at several intervals to notice the disappearance of reactants and the introduction of the half acid/ester.

$C_2F_5CH_2CH_2[(CF_2CF_2)(CH_2CH_2)]_k-OC(O)CH=CHC(O)OH$. The half acid/ester (4.75 g, 9.7 mmol), $C_2F_5CH_2CH_2[(CF_2CF_2)(CH_2CH_2)]_k-OH$ (3.80 g, 9.7 mmol), and *p*-toluenesulfonic acid monohydrate (0.12 g, 0.97 mmol) were heated to 30 reflux in toluene (50 mL) at 114°C for 19 h. The product was isolated using extraction with CH_3CN (3 x 100 mL), concentration, extraction with tetrahydrofuran, concentration, and drying was to produce the yellow/orange solid product (7.46 g, 93.3% yield, 97%), which was analyzed by 1H NMR and LC/MS to confirm the structure as

35 $C_2F_5CH_2CH_2[(CF_2CF_2)(CH_2CH_2)]_kOOC(O)CH=CHC(O)O[(CH_2CH_2)-(CF_2CF_2)]_k-CH_2CH_2C_2F_5$ wherein k is a mixture of 2 and 3.

Example 7

A mixture of ethanolamine (13 g, 28 mmol) and ether (30 mL) was cooled to 15°C. Perfluoro-2-methyl-3-oxahexanoyl fluoride (33 g in ether 50 mL) was added dropwise to keep the reaction temperature below 25°C. After the addition, 5 the reaction mixture was stirred at room temperature for one hour. The solid was removed by filtration and the filtrate was washed with hydrochloric acid (0.5N, 30 mL), water (2 times 30 mL), sodium hydrogen carbonate solution (0.5N, 20 mL), water (30 mL), and sodium chloride solution (saturated, 20 mL). It was then 10 concentrated and dried in vacuum over night at room temperature to give a white solid 35 g, yield 95%. Analysis by ^1H NMR and F NMR showed the product was N-(perfluoro-2-methyl-3-oxahexanoyl)-2-aminoethanol, $\text{C}_3\text{F}_7\text{OCF}(\text{CF}_3)\text{CONHCH}_2\text{CH}_2\text{OH}$.

Itaconic anhydride (0.67 g, 6.0 mmol), $\text{C}_3\text{F}_7\text{OCF}(\text{CF}_3)\text{CONH-CH}_2\text{CH}_2\text{OH}$ (4.44 g, 12 mmol, prepared as described above), *p*-toluenesulfonic acid monohydrate (0.11g, 0.60 mmol), and toluene (50mL) were stirred 15 continuously and heated to reflux at 111°C for a period of 25 h. The toluene was decanted off to leave a yellow, viscous solid. The product was firstly air-dried and then placed in a vacuum oven for 2 h. The product (3.62g, 72.4%, 65% 20 purity) was analyzed by ^1H NMR and LC/MS to confirm complete conversion and the structure as $\text{C}_3\text{F}_7\text{OCF-}(\text{CF}_3)\text{C}(\text{O})\text{NHCH}_2\text{CH}_2\text{OC}(\text{O})\text{CH}_2\text{C}(\text{=CH}_2)\text{C}(\text{O})\text{OCH}_2\text{CH}_2\text{-NHC}(\text{O})\text{CF}(\text{CF}_3)\text{OC}_3\text{F}_7$.

Example 8

Ethylene (25 g) was introduced to an autoclave charged with 25 $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{I}$ (217 g) and d-(+)-limonene (1 g), and the reactor heated at 240 °C for 12 hours. The product was isolated by vacuum distillation to provide $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{I}$. Fuming sulfuric acid (70 mL) was added slowly to 50 g of $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{I}$ and mixture was stirred at 60°C for 1.5 hours. The 30 reaction was quenched with ice-cold 1.5 wt% Na_2SO_3 aqueous solution and heated at 95°C for 0.5 hours. The bottom layer was separated and washed with 10 wt% aqueous sodium acetate and distilled to provide $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{OH}$ bp 54-57 °C at 2 mmHg (267 Pa).

Itaconic anhydride (0.75 g, 6.7 mmol), $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{OH}$ (4.37 g, 13 mmol, prepared as described above), *p*-toluenesulfonic acid monohydrate 35 (0.13 g, 0.67 mmol) and toluene (50 mL) were refluxed for a period of 19 h at a temperature of 113°C. The resulting pale yellow liquid (4.53 g, 90.6% yield, 72%

purity) was analyzed by ^1H NMR and LC/MS to confirm the structure as $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{OC(O)CH}_2\text{C(=CH}_2\text{)C(O)OCH}_2\text{CH}_2\text{CF}_2\text{CH}_2\text{C}_4\text{F}_9$.

Example 9

A mixture of ethanolamine (13 g, 28 mmol) and ether (30 mL) was cooled to 15°C. Perfluoro-2-methyl-3-oxahexanoyl fluoride (33 g in ether 50 mL) was added dropwise to keep the reaction temperature below 25°C. After the addition, the reaction mixture was stirred at room temperature for one hour. The solid was removed by filtration and the filtrate was washed with hydrochloric acid (0.5N, 30 mL), water (2 times 30 mL), sodium hydrogen carbonate solution (0.5N, 20 mL), water (30 mL), and sodium chloride solution (saturated, 20 mL). It was then concentrated and dried in vacuum over night at room temperature to give a white solid 35 g, yield 95%. Analysis by ^1H NMR and F NMR showed the product was N-(perfluoro-2-methyl-3-oxahexanoyl)-2-aminoethanol, $\text{C}_3\text{F}_7\text{OCF(CF}_3\text{)CONHCH}_2\text{CH}_2\text{OH}$.

Citraconic anhydride (0.67 g, 6.0 mmol), $\text{C}_3\text{F}_7\text{OCF(CF}_3\text{)CONHCH}_2\text{CH}_2\text{OH}$ (4.44 g, 12 mmol, prepared as described above), *p*-toluenesulfonic acid monohydrate (0.11 g, 0.60 mmol) and toluene (50 mL) were added together and heated to reflux at 111°C, with continual stirring, for 40 h. There were two noticeable solid materials present within the toluene solution. The pinkish solid was removed and the white solid was vacuum filtered. Both materials were analyzed by LC/MS, which confirmed that the pinkish solid was the product and the white solid was unreacted alcohol (1.22g). The product (2.98 g, 59.6% yield, 65% purity) was analyzed by ^1H NMR and LC/MS to confirm the structure as $\text{C}_3\text{F}_7\text{OCF(CF}_3\text{)C(O)NHCH}_2\text{CH}_2\text{OC(O)-C(CH}_3\text{)=CH}_2\text{C(O)OCH}_2\text{CH}_2\text{NHC(O)CF(CF}_3\text{)OC}_3\text{F}_7$.

Example 10

Ethylene (25 g) was introduced to an autoclave charged with $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{I}$ (217 g) and d-(+)-limonene (1 g), and the reactor heated at 240 °C for 12 hours. The product was isolated by vacuum distillation to provide $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{I}$. Fuming sulfuric acid (70mL) was added slowly to 50 g of $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{I}$ and mixture was stirred at 60 °C for 1.5 hours. The reaction was quenched with ice-cold 1.5 wt% Na_2SO_3 aqueous solution and heated at 95 °C for 0.5 hours. The bottom layer was separated and washed with 10 wt% aqueous sodium acetate and distilled to provide $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{OH}$: bp 54-57 °C at 2 mmHg (267 Pa).

Citraconic anhydride (0.75 g, 6.7 mmol), $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{OH}$ (4.37 g, 13.3 mmol, prepared as described above), *p*-toluenesulfonic acid monohydrate (0.13 g), and toluene (50 mL) were refluxed for about 46 h at 112°C, after which only the diester was observed in the LC/MS analysis. The 5 work-up was carried out as in example 1 to give a pale yellow liquid (2.98 g, 59.6% yield, >99% purity) which was analyzed by ^1H NMR and LC/MS to confirm the diester structure as

$\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{OC(O)C(CH}_3\text{)=CH}_2\text{C(O)OCH}_2\text{CH}_2\text{CF}_2\text{CH}_2\text{C}_4\text{F}_9$.

Example 11

10 Ethylene (25 g) was introduced to an autoclave charged with $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{I}$ (217 g) and d-(+)-limonene (1 g), and the reactor heated at 240 °C for 12 hours. The product was isolated by vacuum distillation to provide $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{I}$. Fuming sulfuric acid (70mL) was added slowly to 50 g of $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{I}$ and mixture was stirred at 60 °C for 1.5 hours. The 15 reaction was quenched with ice-cold 1.5 wt% Na_2SO_3 aqueous solution and heated at 95 °C for 0.5 hours. The bottom layer was separated and washed with 10 wt% aqueous sodium acetate and distilled to provide

$\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{OH}$: bp 54-57 °C at 2 mmHg (267 Pa).

20 Trans-glutaconic acid (0.87 g, 6.7 mmol), $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{OH}$ (4.37 g, 13 mmol, prepared as described above), *p*-toluenesulfonic acid monohydrate (0.13g, 0.67 mmol) and toluene (50mL) were stirred continuously together and heated to reflux at 111°C for 24 h. The work-up procedure was carried out as in Example 1. The resulting white solid (2.52 g, 50.4% yield, 80% 25 purity) was dried in a vacuum oven and analyzed by ^1H NMR and LC/MS to confirm the structure as


$\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{OC(O)CH=CHCH}_2\text{C(O)OCH}_2\text{CH}_2\text{CF}_2\text{CH}_2\text{C}_4\text{F}_9$.

Example 12

30 Ethylene (56 g) was introduced to an autoclave charged with $\text{C}_4\text{F}_9(\text{CH}_2\text{CF}_2)_2\text{I}$ (714 g) and d-(+)-limonene (3.2 g), and the reactor heated at 240 °C for 12 hours. The product was isolated by vacuum distillation to provide $\text{C}_4\text{F}_9(\text{CH}_2\text{CF}_2)_2\text{CH}_2\text{CH}_2\text{I}$. A mixture of $\text{C}_4\text{F}_9(\text{CH}_2\text{CF}_2)_2\text{CH}_2\text{CH}_2\text{I}$ (10 g, 0.02 mol) and N-methylformamide (8.9 mL, 0.15 mol) was heated to 150 °C for 26 hours. The mixture was cooled to 100 °C, followed by the addition of water to separate the crude ester. Ethyl alcohol (3 mL) and *p*-toluene sulfonic acid (0.09 35 g) were added and the mixture stirred at 70 °C for 0.25 hours. Ethyl formate and ethyl alcohol were removed by distillation to give a crude product. The crude

product was dissolved in ether, washed with 10 % by weight aqueous sodium sulfate, water and brine, in turn, and dried over magnesium sulfate. Distillation provided the product $C_4F_9(CH_2CF_2)_2CH_2CH_2OH$ (6.5 g, 83 % yield): bp 94-95 °C at 2 mm Hg (266 Pa).

5 Trans-glutaconic acid (0.75 g, 5.8 mmol), $C_4F_9CH_2CF_2CH_2CF_2CH_2CH_2OH$ (4.54 g, 12 mmol, prepared as described above), *p*-toluenesulfonic acid monohydrate (0.11g, 0.58 mmol) and toluene (50mL) were stirred continuously together and heated to reflux at 111°C for a period of 16 h. The progress was monitored by LC/MS and the removal of water 10 azeotropically. The orange/yellow solid was filtered and washed with 5% sodium bicarbonate solution (50mL). The filtrate was separated and the organic layer was washed with 5% sodium bicarbonate solution (50mL), and then with deionized water (50mL). The combined organic extracts were dried over anhydrous $MgSO_4$ and the toluene was then concentrated (140.30mmHg, 67°C). The orange solid 15 (4.14 g, 81.5% yield, 85% purity) was dried in a vacuum oven and analyzed by 1H NMR and LC/MS to confirm the structure as



Example 13

20 Ethylene (25 g) was introduced to an autoclave charged with $C_4F_9CH_2CF_2I$ (217 g) and d-(+)-limonene (1 g), and the reactor heated at 240 °C for 12 hours. The product was isolated by vacuum distillation to provide $C_4F_9CH_2CF_2CH_2CH_2I$. Fuming sulfuric acid (70mL) was added slowly to 50 g of $C_4F_9CH_2CF_2CH_2CH_2I$ and mixture was stirred at 60 °C for 1.5 hours. The 25 reaction was quenched with ice-cold 1.5 wt% Na_2SO_3 aqueous solution and heated at 95 °C for 0.5 hours. The bottom layer was separated and washed with 10 wt% aqueous sodium acetate and distilled to provide $C_4F_9CH_2CF_2CH_2CH_2OH$: bp 54-57 °C at 2 mmHg (267 Pa).

30 A melt reaction was undertaken by reacting maleic anhydride (2.00 g, 20 mmol) with $C_4F_9CH_2CF_2CH_2CH_2OH$ (6.69 g, 20 mmol, prepared as described above). The reaction was sustained at 70°C for a period of 34 h, during which aliquots were taken for GC analysis. The white solid half acid/ester (8.02 g, 92.3% yield, >98% purity) was analyzed by 1H NMR and LC/MS to confirm the structure as $C_4F_9CH_2CF_2CH_2CH_2OC(O)CH=CHC(O)OH$.

Example 14

The maleate, prepared as described in Example 13, (6.68 g, 17 mmol), *p*-toluenesulfonic acid monohydrate (0.30 g, 1.7 mmol) and hexyl alcohol (1.60 g, 17 mmol) were mixed together along with toluene (50 mL). The mixture was stirred continuously together and heated to reflux at 114°C for 19 h. The work-up procedure as in example 1 was conducted to produce a clear liquid (7.14 g, 89.3% yield, 98% purity), that was analyzed by ¹H NMR and LC/MS to confirm the structure as C₄F₉CH₂CF₂CH₂CH₂OC(O)CH=CHC(O)O-(CH₂)₆H.


Example 15

The maleate, prepared as described in Example 13, (4.41 g, 10 mmol), *p*-toluenesulfonic acid monohydrate (0.20 g, 1.0 mmol) and C₆F₁₃CH₂CH₂OH (3.77 g, 10 mmol) were added together along with toluene (50 mL). The contents were refluxed for 19 h at 114°C and the work-up procedure was carried out as in Example 1. The pale yellow liquid (5.78 g, 72.3% yield, 90% purity) was analyzed by ¹H NMR and LC/MS to confirm the formation of the mixed diester and the structure as



Example 16

The maleate, prepared as described in Example 1, (2.62 g, 3.2 mmol) and isopropyl alcohol (IPA, 31 g) were added together at 50°C until the mixture was dissolved; about 10 minutes. Aqueous sodium bisulfite (0.17 g, 1.6 mmol) was dissolved in deionized water (8 mL) and added dropwise to the isopropyl alcohol solution, which was then heated to reflux (86°C) for 26 h. The isopropyl alcohol and water were removed by rotary evaporation followed by drying in a vacuum oven at 50°C to generate a viscous yellow liquid (1.70 g, 57.6% yield, 75% purity), which was confirmed to be the diester sulfonate by ¹H NMR and LC/MS analyses to confirm the structure as



The product was evaluated for CMC and surface tension beyond the CMC by Test Method 1; the results are shown in Table 2.

Example 17

The maleate, prepared as described in Example 2, (7.74 g, 11 mmol) and isopropyl alcohol (31 g) were stirred continuously together. The temperature was raised to 61°C and then a solution of sodium bisulfite (1.09 g, 11 mmol), dissolved in deionized water (53 mL), was added dropwise. The mixture was

heated to reflux at an elevated temperature of 82°C for 24 h. The solution was concentrated to remove the isopropyl alcohol/water solution. The remaining pale yellow liquid was dried overnight in an oven to produce a white solid (6.96 g, 78.8% yield, 98% purity) and was then analyzed by ¹H NMR and LC/MS to 5 confirm the structure as C₄F₉CH₂CF₂CH₂CH₂OC(O)-CH₂CH(SO₃Na)C(O)O-CH₂CH₂CF₂CH₂C₄F₉. The product was evaluated for CMC and surface tension beyond the CMC by Test Method 1, with results shown in Table 2, and spreading on cyclohexane by Test Method 2, with results shown in Table 3.

Example 18

10 The maleate, prepared as described in Example 3, (2.88 g, 3.3 mmol) and isopropyl alcohol (31 g) were stirred continuously at 82°C, with the addition of aqueous sodium bisulfite (1.54 g, 15 mmol), dissolved in deionised water (20 mL), for 28 h. The white solid (2.58 g, 80.1% yield, >95% purity) was collected by concentrating the isopropyl alcohol/water solution and then dried in a vacuum 15 oven overnight. The product was analyzed by ¹H NMR and LC/MS to confirm the structure as C₄F₉CH₂CF₂CH₂CF₂CH₂CH₂OC(O)CH₂CH(SO₃Na)C(O)O-CH₂CH₂CF₂CH₂CF₂CH₂C₄F₉. The product was evaluated for CMC and surface tension beyond the CMC by Test Method 1, with results shown in Table 2, and spreading on cyclohexane by Test Method 2, with results shown in Table 3.

20 Example 19

25 The maleate, prepared as described in Example 4, (4.10 g, 5.5 mmol), isopropyl alcohol (31g) and aqueous sodium bisulfite (0.28 g, 2.8 mmol) dissolved in deionized water (14 mL) were stirred continuously for 18 h at a temperature of 82°C. The white solid (3.36g, 71.9% yield, >95% purity) was collected by rotary evaporating the isopropyl alcohol/water solution and then the product was dried in a vacuum oven. The product was analyzed by ¹H NMR and LC/MS to confirm conversion to the diester sulfonate and the structure as C₃F₇OCF₂CF₂CH₂CH₂OC(O)CH₂CH(SO₃Na)C(O)O-CH₂CH₂CF₂CF₂OC₃F₇. The product was evaluated for CMC and surface tension beyond the CMC by Test Method 1, with results shown in Table 2, and spreading on cyclohexane by Test Method 2, with results shown in Table 3.

Example 20

30 The maleate, prepared as described in Example 5, (1.49 g, 1.8 mmol), isopropyl alcohol (31 g) and aqueous sodium bisulfite (0.29 g, 2.8 mmol) dissolved in deionised water (14 mL) were mixed together and refluxed for 27 h at 82°C. The isopropyl alcohol was concentrated and the white solid (1.46 g, 87.1%

yield, >97% purity) obtained was dried in a vacuum oven and analyzed by ^1H NMR and LC/MS to confirm the structure as $\text{C}_3\text{F}_7\text{OCFHCF}_2\text{OCH}_2\text{CH}_2\text{OCH}_2\text{CH}_2\text{OC(O)CH}_2\text{CH(SO}_3\text{Na)\text{C(O)O-CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2\text{OCF}_2\text{CFHOC}_3\text{F}_7}$. The product was evaluated for CMC and 5 surface tension beyond the CMC by Test Method 1, with results shown in Table 2, and spreading on cyclohexane by Test Method 2, with results shown in Table 3.

Example 21

10 The maleate, prepared as described in Example 6, (7.54 g, 8.7 mmol) and isopropyl alcohol (31 g) were heated to 50°C until the solid had dissolved in solution. A solution of aqueous sodium bisulfite (0.91 g, 8.7 mmol) dissolved in deionized water (43 mL) was transferred to the mixture and the contents were refluxed at 82°C for 20 h. The isopropyl alcohol/water solution was removed by rotary evaporation to attain the orange/brown solid (7.22 g, 87.3% yield, 92% purity). The product was analyzed by ^1H NMR and LC/MS to confirm the structure 15 as $\text{C}_2\text{F}_5\text{CH}_2\text{CH}_2[(\text{CF}_2\text{CF}_2)(\text{CH}_2\text{CH}_2)]_k\text{OOC(O)CH=CHC(O)O-[(CH}_2\text{CH}_2)(\text{CF}_2\text{CF}_2)]_k\text{CH}_2\text{CH}_2\text{C}_2\text{F}_5$, wherein k is a mixture of 2 and 3 in a 2:1 ratio. The product was evaluated for CMC and surface tension beyond the CMC by Test Method 1; the results are shown in Table 2.

Example 22

20 The itaconate, prepared as described in Example 7, (3.60 g, 4.3 mmol) and isopropyl alcohol (31 g) were stirred continuously together. An aqueous solution 5 of sodium bisulfite (0.45 g, 4.3 mmol) dissolved in deionized water (21 mL) was added slowly to the solution and the temperature was raised to 82°C for 23 h. The isopropyl alcohol/water was concentrated to leave the yellow gel-like product 25 (3.73 g, 92.1% yield, 75% purity), which was placed in a vacuum oven overnight analyzed by ^1H NMR and LC/MS to confirm the structure as $\text{C}_3\text{F}_7\text{OCF(CF}_3\text{)C(O)NHCH}_2\text{CH}_2\text{OC(O)C}_3\text{H}_5(\text{SO}_3\text{Na)\text{C(O)O-CH}_2\text{CH}_2\text{NHC(O)CF(CF}_3\text{)OC}_3\text{F}_7}$. The product was evaluated for CMC and 30 surface tension beyond the CMC by Test Method 1, with results shown in Table 2, and spreading on cyclohexane by Test Method 2, with results shown in Table 3.


Example 23

35 The itaconate, prepared as described in Example 8, (2.00 g, 2.7 mmol), isopropyl alcohol (31 g) and aqueous sodium bisulfite (0.28 g, 2.7 mmol) dissolved in deionized water (14 mL) were refluxed for 22 h at 82°C. The white solid (precipitate) was filtered off and washed with deionized water (50 mL) to remove unreacted NaHSO_3 . The white dry solid (2.11 g, 91.5% yield, >95%

purity) was analyzed by ^1H NMR and LC/MS to confirm the structure as $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{OC(O)C}_3\text{H}_5(\text{SO}_3\text{Na})\text{C(O)OCH}_2\text{CH}_2\text{CF}_2\text{CH}_2\text{C}_4\text{F}_9$. The product was evaluated for CMC and surface tension beyond the CMC by Test Method 1, with results shown in Table 2, and spreading on cyclohexane by Test 5 Method 2, with results shown in Table 3.

Example 24

The citraconate, prepared as described in Example 9, (2.96 g, 3.5 mmol), and isopropyl alcohol (31 g) were stirred continuously together and heated to reflux. A solution of aqueous sodium bisulfite (0.37 g, 3.5 mmol) dissolved in 10 deionized water (18 mL) was added dropwise to the mixture. The solution was maintained at 82°C for 23 h. The solution was concentrated and two noticeable layers were observed. The small top layer was yellow in colour and the bottom was white. Each layer was analyzed by ^1H NMR, which confirmed that the top layer was likely to be impurities. The product was tested in isopropyl alcohol and 15 also in water, and the alcohol was also similarly tested. The results indicated that the product was soluble in water but insoluble in isopropyl alcohol, and the opposite was true for the alcohol. Therefore, if the impurity layer contained some alcohol this would be removed by filtration when water was added. If any of the starting acid remained this would not affect the surface tension results. The 20 bottom layer (2.62 g, 78.8% yield, 85% purity) was analyzed by ^1H NMR and LC/MS to confirm the structure as



The product was evaluated for CMC and surface tension beyond the CMC 25 by Test Method 1; the results are shown in Table 2.

Example 25

The citraconate, prepared as described in Example 10, (2.70 g, 3.6 mmol) and isopropyl alcohol (31 g) were mixed together at 50°C until dissolved; about 10 minutes. Aqueous sodium bisulfite (1.54 g, 14.8 mmol) was dissolved in 30 deionized water (15 mL) and added dropwise to the isopropyl alcohol solution, which was then heated to about 82°C for about 22 h. The isopropyl alcohol and water were removed by rotary evaporation followed by drying in a vacuum oven at 50°C to give an off-white solid (1.56 g, 50.8% yield, Purity: >99%), which was analyzed by ^1H NMR and LC/MS to confirm the formation of the diester 35 sulfonate and the structure as $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{OC(O)C}_3\text{H}_5(\text{SO}_3\text{Na})\text{C(O)OCH}_2\text{CH}_2\text{CF}_2\text{CH}_2\text{C}_4\text{F}_9$. The product was evaluated for CMC and surface


tension beyond the CMC by Test Method 1, with results shown in Table 2, and spreading on cyclohexane by Test Method 2, with results shown in Table 3.

Example 26

5 The trans-glutaconate, prepared as described in Example 11, (2.52 g, 3.4 mmol) was added to isopropyl alcohol (31 g) and heated to 60°C. At this point a solution of sodium bisulfite (0.31 g, 3.0 mmol) dissolved in deionized water (15 mL) was added dropwise, and the temperature was raised to 82°C for 22 h. The pale yellow solid (2.26 g, 78.8% yield, 80% purity) was analyzed by ^1H NMR and LC/MS to confirm the structure as
10 $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{OC(O)C}_3\text{H}_5(\text{SO}_3\text{Na})\text{C(O)OCH}_2\text{CH}_2\text{CF}_2\text{CH}_2\text{-C}_4\text{F}_9$.
The product was evaluated for CMC and surface tension beyond the CMC by Test Method 1; the results are shown in Table 2.

Example 27

15 The trans-glutaconate, prepared as described in Example 12, (4.08 g, 4.6 mmol) was added to isopropyl alcohol (31 g) and heated to 50°C. A solution of sodium bisulfite (0.31 g, 3.0 mmol) dissolved in deionized water (15 mL) was added dropwise to the solution and the mixture was heated to 82°C for 23 h. The yellow solid (3.94 g, 86.3% yield, 90% purity) was collected by rotary
20 evaporating the isopropyl alcohol/water solution and analyzed by ^1H NMR and LC/MS to confirm the structure as



The product was evaluated for CMC and surface tension beyond the CMC by Test Method 1; the results are shown in Table 2.

25 Example 28

30 The maleate, prepared as described in Example 13, (4.20 g, 9.4 mmol) and isopropyl alcohol (31 g) were heated to approximately 50°C to allow for the solid to dissolve in solution. Aqueous sodium bisulfite (0.99 g, 9.4 mmol) dissolved in deionized water (47 mL) was transferred to the solution and the contents were refluxed for 22 h at 82°C. The isopropyl alcohol/water solution was rotary
evaporated to leave the white solid (4.14 g, 82.8% yield, 90% purity) that was
35 analyzed by ^1H NMR and LC/MS to confirm the structure as
 $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{OC(O)CH}_2\text{CH}(\text{SO}_3\text{Na})\text{C(O)OH}$. The product was evaluated for CMC and surface tension beyond the CMC by Test Method 1; the results are shown in Table 2.

Example 29

The mixed diester, prepared as described in Example 14, (7.10 g, 13.9 mmol) and isopropyl alcohol (32 g) were stirred continuously together and heated to 50°C until the two liquids became miscible. Aqueous sodium bisulfite (1.45 g, 5 13.9 mmol) dissolved in deionized water (70 mL) was transferred to the mixture and the contents were refluxed for 22 h at 82°C. The isopropyl alcohol/water solution was evaporated off and the white gel product (6.26 g, 73.3% yield, 98% purity) was dried in vacuum oven for 2 h. The product was analyzed by ^1H NMR and LC/MS to confirm the structure as

10 $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{OC(O)CH}_2\text{CH}(\text{SO}_3\text{Na})\text{C(O)O(CH}_2)_6\text{H}$. The product was evaluated for CMC and surface tension beyond the CMC by Test Method 1, with results shown in Table 2, and spreading on cyclohexane by Test Method 2, with results shown in Table 3.

Example 30

15 The mixed diester, prepared as in Example 15 (5.78 g, 7.5 mmol) and isopropyl alcohol (31 g) were added together and heated to 60°C for 10 minutes. A solution of sodium bisulfite (0.78 g, 7.5 mmol) dissolved in deionized water (37 mL) was added to the solution and the mixture was heated to reflux at 82°C for a period of 20 h. The isopropyl alcohol/water solution was rotary evaporated off to 20 leave a colorless gel product (4.26 g, 65% yield, 98% purity) that was analyzed by ^1H NMR and LC/MS to confirm the structure as

$\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{OC(O)CH}_2\text{CH}(\text{SO}_3\text{Na})\text{C(O)OCH}_2\text{CH}_2(\text{CF}_2)_6\text{F}$. The product was evaluated for CMC and surface tension beyond the CMC by Test Method 1; the results are shown in Table 2.

25 Comparative Example A

Maleic anhydride (0.63 g, 6.5 mmol), 1H,1H,2H,2H-perfluoro-1-octanol (4.74 g, 13 mmol), *p*-toluenesulfonic acid monohydrate (*p*-TsOH) (0.19 g, 1.0 mmol) and toluene (50mL) were added to a flask and heated to reflux for 96 hours at 111°C. The solution was separated and extracted with two washings of 5% 30 sodium bicarbonate (50mL each). The combined organic extracts were dried over anhydrous magnesium sulfate, and concentrated to remove the toluene at 140.30 mmHg (18.7 kPa) and 67°C). The structure of the resulting liquid product di(1H,1H,2H,2H-perfluoroctyl) maleate (4.88 g, 93.4% yield, >80% purity) was confirmed by ^1H NMR and LC/MS.

35 Di(1H,1H,2H,2H-perfluoroctyl) maleate (4.70 g, 5.8 mmol, prepared as described above) was added to isopropyl alcohol (isopropyl alcohol, 31 g) and

heated to 50°C for a period of 10 min. with continual stirring. A solution of sodium bisulfite (0.61 g, 5.8 mmol) dissolved in deionized water (10mL) was added dropwise to the solution. The mixture was refluxed for 22 h at 82°C. The progress was checked by LC/MS and a further addition of aqueous sodium 5 bisulfite (0.61 g, 5.9 mmol) was added. The mixture was refluxed for a further 70.3 h. The isopropyl alcohol/water solution was removed by rotary evaporation to produce a white solid. (2.70 g, 52.2% yield, 99% purity). The product 10 composition was confirmed by ^1H NMR and LC/MS as the sodium salt of di(1H,1H,2H,2H-perfluoroctyl) maleate-2-sulfosuccinate. The product was evaluated for CMC and surface tension beyond the CMC by Test Method 1; the results are shown in Table 1.

Comparative Example B

1H,1H,2H,2H-perfluoroctanol (8.02 g, 22 mmol), 15 dicyclohexylcarbodiimide (DCC) (4.27 g, 21 mmol) and dichloromethane CH_2Cl_2 , 35mL) were added to a flask, equipped with a nitrogen inlet, overhead stirrer and two stoppers. The solution was cooled to 0°C and the citraconic acid (1.28 g, 9.8 mmol) dissolved in tetrahydrofuran (15mL) was added dropwise. The solution was stirred for 10 min. and then the ice bath was removed to allow 20 the solution to warm to room temperature. The mixture was left to stir overnight. The resulting mixture was filtered to remove the traces of 1,3-dicyclohexylurea that was produced as a by-product and then washed with excess tetrahydrofuran (50mL). The tetrahydrofuran and CH_2Cl_2 were concentrated at 378.14 mmHg (kPa) and 46°C and the product was dried in a vacuum oven for 3 hours. The product was analyzed through ^1H NMR and LC/MS, which indicated the 25 conversion to monoester. A similar procedure was carried out again but with the addition of another mole of alcohol. The alcohol (6.30 g, 13 mmol), DCC (2.63 g, 13 mmol) and CH_2Cl_2 (35 mL) were added to the flask and cooled to 0°C. The monoester, that was produced previously, was re-dissolved in tetrahydrofuran 30 (15mL) and added dropwise to the solution. The work-up method was carried out and the resulting product was a pale yellow liquid (6.46 g, 80.0% yield, 75% purity). The product was analyzed by ^1H NMR and LC/MS to confirm the structure as di(1H,1H,2H,2H-perfluoroctyl) citraconate.

35 Di(1H,1H,2H,2H-perfluoroctyl) citraconate (4.99 g, 6.1 mmol, prepared as described above) and isopropyl alcohol (32 g) were transferred to a flask and heated to 50°C for 10 min. A solution of aqueous sodium bisulfite (1.53 g, 15 mmol) dissolved in deionized water was added to the solution and heated to reflux (82°C) for 22 h. The white solid was dried in an oven overnight (2.98 g, 53.0%

yield, 95% purity). The product composition was confirmed by ^1H NMR and LC/MS as the sodium salt of di(1H,1H,2H,2H-perfluorooctyl) citraconate-2-sulfosuccinate. The product was evaluated for CMC and surface tension beyond the CMC by Test Method 1, with results shown in Table 2, and spreading on cyclohexane by Test Method 2, with results in Table 3.

Comparative Example C

Maleic anhydride (17.2 g, 176 mmol), 1H,1H,2H,2H,-perfluorohexanol (93.1 g, 353 mmol), *p*-toluenesulfonyl hydroxide (*p*-TsOH) (3.4 g, 17.6 mmol) and toluene (500mL) were heated to reflux for 8 h. An additional amount of *p*-TsOH (3.4 g, 17.6 mmol) was added after 4 h of reflux. The solution was stirred overnight at room temperature. The solution was diluted with ethyl acetate (500mL) and washed three times with brine (250mL each). The combined extracts were washed with a further washing of ethyl acetate (300mL). The combined organics were dried over anhydrous MgSO_4 and concentrated to yield a colorless oil (85.8 g, 80% yield, 98% purity). The structure of the product was confirmed by ^1H NMR and LC/MS as di(1H,1H,2H,2H-perfluorohexyl) maleate.

Di(1H,1H,2H,2H-perfluorohexyl) maleate (1.5 g, 2.5 mmol, prepared as described above) was added to isopropyl alcohol (32 g) and heated for a period of 10 min. until the two liquids became miscible. A solution of sodium bisulfite (1.5 g, 14 mmol) dissolved in deionized water (15 mL) was transferred to the flask and the contents were heated to reflux at 82°C for 22 h. The white solid product resulted after the removal of isopropyl alcohol/water solution (0.98 g, 55.8% yield, 99% purity). The product composition was confirmed by ^1H NMR and LC/MS as the sodium salt of di(1H,1H,2H,2H-perfluorohexyl) maleate-2-sulfosuccinate. The product was evaluated for CMC and surface tension beyond the CMC by Test Method 1, with results shown in Table 2, and spreading on cyclohexane by Test Method 2, with results shown in Table 3.

Comparative Example D

Ethylene (25 g) was introduced to an autoclave charged with $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{I}$ (217 g) and d-(+)-limonene (1 g), and the reactor heated at 240 °C for 12 hours. The product was isolated by vacuum distillation to provide $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{I}$. Fuming sulfuric acid (70mL) was added slowly to 50 g of $\text{C}_4\text{F}_9\text{CH}_2\text{CF}_2\text{CH}_2\text{CH}_2\text{I}$ and mixture was stirred at 60 °C for 1.5 hours. The reaction was quenched with ice-cold 1.5 wt% Na_2SO_3 aqueous solution and heated at 95 °C for 0.5 hours. The bottom layer was separated, washed with 10

wt% aqueous sodium acetate, distilled to provide $C_4F_9CH_2CF_2CH_2CH_2OH$: bp 54-57 °C at 2 mmHg (267 Pa).

5 Trans- β -hydromuconic acid (0.94 g, 6.5 mmol), *p*-toluenesulfonic acid monohydrate (0.12 g, 0.65 mmol), $C_4F_9CH_2CF_2CH_2CH_2OH$ (4.29 g, 13 mmol) and toluene were added together and the contents were heated to reflux at 111°C for 25 h. The work-up as in example 1 was conducted. The white solid (3.82 g, 76.4% yield, 95% purity) analyzed by 1H NMR and LC/MS to confirm the structure as $C_4F_9CH_2CF_2CH_2CH_2OC(O)CH_2-CH=CHCH_2C(O)OCH_2CH_2CF_2CH_2-C_4F_9$.

10 The trans- β -hydromuconate, prepared as described above, (3.80 g, 5.0 mmol) was added to isopropyl alcohol (31 g) and heated to 60°C. A solution of aqueous sodium bisulfite (0.52 g, 5.0 mmol) was dissolved in deionized water and transferred to the mixture. The temperature was raised to 82°C and maintained for 22 h. The white precipitate was collected by vacuum filtration and the filtrate was concentrated to remove the isopropyl alcohol/water solution. The white solid 15 (3.88 g, 89.9% yield, 98% purity) was analyzed by 1H NMR and LC/MS to confirm the structure as

$C_4F_9CH_2CF_2CH_2CH_2OC(O)CH_2CH(SO_3Na)CH_2CH_2C(O)O-$
 $CH_2CH_2CF_2CH_2C_4F_9$. The product was evaluated for CMC and surface tension beyond the CMC by Test Method 1; the results are shown in Table 2.

20 Comparative Example E

25 $C_3F_7OCF_2CF_2I$ (100 g, 0.24 mol) and benzoyl peroxide (3 g) were charged to a pressure vessel under nitrogen. A series of three vacuum/nitrogen gas sequences was then executed at -50 °C and ethylene (18 g, 0.64 mol) was introduced. The vessel was heated for 24 hour at 110 °C. The autoclave was cooled to 0 °C and opened after degassing. Then the product was collected in a bottle. The product was distilled giving 80 g of $C_3F_7OCF_2CF_2CH_2CH_2I$ in 80% yield. The boiling point was 56~60°C at 25 mm Hg (3.3 kPa).

30 A mixture of $C_3F_7OCF_2CF_2CH_2CH_2I$ (300 g, 0.68 mol, prepared as described above) and N-methyl-formamide (300 mL), was heated to 150 °C for 26 h. Then the reaction was cooled to 100 °C, followed by the addition of water to separate the crude ester. Ethyl alcohol (77 mL) and *p*-toluene sulfonic acid (2.59 g) were added to the crude ester, and the reaction was stirred at 70 °C for 15 minutes. Then ethyl formate and ethyl alcohol were distilled out to give a crude product. The crude product was dissolved in ether, washed with aqueous sodium sulfite, water, and brine in turn, then dried over magnesium sulfate. The product 35

was then distilled to give 199 g of $C_3F_7OCF_2CF_2CH_2CH_2OH$ in 85 % yield. The boiling point was 71-73°C at 40 mm Hg (5.3 kPa).

Trans- β -hydromuconic acid (0.94 g, 6.5 mmol), $C_3F_7OCF_2CF_2CH_2CH_2OH$ (4.30 g, 13 mmol, prepared as described above), *p*-toluenesulfonic acid monohydrate H (0.13 g, 0.65 mmol) and toluene (50mL) were stirred continuously together and heated to reflux (111°C for 25 h). The work-up was carried out to produce a pale yellow liquid (3.90 g, 78.0% yield, 99% purity), which was analyzed by 1H NMR and LC/MS to confirm the structure as $C_3F_7OCF_2CF_2CH_2CH_2OC(O)CH_2CH=CHCH_2C(O)O-CH_2CH_2CF_2CF_2OC_3F_7$.

The trans- β -hydromuconate, prepared as described above, (3.88 g, 5.1 mmol) was stirred continuously with isopropyl alcohol (31 g) for a period of 10mins at an elevated temperature of 65°C. A solution of sodium bisulfite (0.29 g, 2.8 mmol) dissolved in deionized water (14 mL) was added dropwise to the mixture. The temperature was raised to 82°C and maintained for a period of 22 h. The solution was concentrated to remove the isopropyl alcohol, and the resulting liquid was left in a vacuum oven overnight. The white solid (3.72 g, 84.4% yield, 87% purity) obtained was analyzed by 1H NMR and LC/MS to confirm the structure as $C_3F_7OCF_2CF_2CH_2CH_2OC(O)CH_2CH(SO_3Na)CH_2CH_2C(O)O-CH_2CH_2CF_2CF_2OC_3F_7$. The product was evaluated for CMC and surface tension beyond the CMC by Test Method 1; the results are shown in Table 2.

Table 1 - Comparative Examples and Surface Tension Measurements

Comparative Example	R _f	X	CMC (wt%)	Surface Tension beyond CMC (mN/m)
Comp. Ex. A	C ₆ F ₁₃	-CH ₂ -CH-(SO ₃ M)-	0.024	13.8
Comp. Ex. B	C ₆ F ₁₃	-CH ₂ -CH(CH ₂ -SO ₃ M)-	0.068	16.3
Comp. Ex. C	C ₄ F ₉	-CH ₂ CH-(SO ₃ M)-	0.26	17.1
Comp. Ex. D	C ₄ F ₉ CH ₂ CF ₂ CH ₂ -CH ₂ -	-CH ₂ CH(SO ₃ M)-CH ₂ CH ₂ -	0.33	18.0
Comp. Ex. E	C ₃ F ₇ OCF ₂ CF ₂ CH ₂ -CH ₂ -	-CH ₂ CH(SO ₃ M)-CH ₂ CH ₂ -	0.88	20.8

Table 2 - Formulae 1A, 1B, and 1C and Surface Tension Measurements

Ex.	R _a	X	Critical Micelle Concn. (wt%)	Surface Tension Beyond CMC (mN/m)
16	C ₃ F ₇ OCF(CF ₃)CONH-CH ₂ CH ₂ -	-CH ₂ CH(SO ₃ M)-	0.016	22.0
17	C ₄ F ₉ CH ₂ CF ₂ CH ₂ CH ₂ -	-CH ₂ CH(SO ₃ M)-	0.051	18.9
18	C ₄ F ₉ CH ₂ CF ₂ CH ₂ CF ₂ -CH ₂ CH ₂ -	-CH ₂ CH(SO ₃ M)-	0.014	20.7
19	C ₃ F ₇ OCF ₂ CF ₂ CH ₂ CH ₂ -	-CH ₂ CH(SO ₃ M)-	0.034	17.1
20	C ₃ F ₇ OCFHCF ₂ O-CH ₂ CH ₂ OCH ₂ CH ₂ -	-CH ₂ CH(SO ₃ M)-	0.039	17.8
21	C ₂ H ₅ CH ₂ CH ₂ [(CF ₂ C _F ₂) _i -(CH ₂ CH ₂) _j] _k	-CH ₂ CH(SO ₃ M)-	0.028	21.1
22	C ₃ F ₇ OCF(CF ₃)CONH-CH ₂ CH ₂ -	-CH ₂ CH(CH ₂ SO ₃ M)-	0.083	18.2
23	C ₄ F ₉ CH ₂ CF ₂ CH ₂ CH ₂ -	-CH ₂ CH(CH ₂ SO ₃ M)-	0.0095	19.4
24	C ₃ F ₇ OCF(CF ₃)CONH-CH ₂ CH ₂ -	-CH(CH ₃)CH(SO ₃ M)-	0.095	25.6

25	C ₄ F ₉ CH ₂ CF ₂ CH ₂ CH 2-	- CH(CH ₃)CH(SO ₃ M)-	0.019	16.8
26	C ₄ F ₉ CH ₂ CF ₂ CH ₂ CH 2-	-CH ₂ CH(SO ₃ M)CH ₂ -	0.042	18.0
27	C ₄ F ₉ CH ₂ CF ₂ CH ₂ CF ₂ -CH ₂ CH ₂ -	-CH ₂ CH(SO ₃ M)CH ₂ -	0.030	18.2
R_a / R				
28*	C ₄ F ₉ CH ₂ CF ₂ CH ₂ CH 2- / -H	-CH ₂ CH(SO ₃ M)-	0.064	19.7
29	C ₄ F ₉ CH ₂ CF ₂ CH ₂ CH 2- / -(CH ₂) ₆ H	-CH ₂ CH(SO ₃ M)-	0.038	16.0
R_a / R_f				
30	C ₄ F ₉ CH ₂ CF ₂ CH ₂ CH 2- / -(CF ₂) ₆ F	-CH ₂ CH(SO ₃ M)-	0.023	20.8

* Example 28 was measured at pH 3.0. Since the R_a is H, the performance of the compound is sensitive to pH.

Table 2 shows that the surfactants of the invention gave low critical micelle concentrations (less than 0.1 weight percent) and low surface tension levels beyond CMC (less than 20 mN/m in water). Table 1 provides data for Comparative Examples. Comparative Example C, having an R_f of C₄F₉ contained a similar fluorine level to the Examples of the invention, but had a far higher CMC value, thus indicating superior performance by the Examples of the invention. Comparative Examples A and B each contained an R_f of C₆F₁₃, which was a higher fluorine level than the Examples of the invention. The Examples of the invention had CMC values similar to Comparative Examples A and B despite the lower level of fluorine. Thus the Examples of the invention had a higher level of fluorine efficiency in providing comparable performance with less fluorine present. Beyond the CMC all of the examples demonstrated comparable surface tension.


Table 3 - Spreading on Cyclohexane

Ex. #	Hydrocarbon surfactant Trials (I) and (II)	Spreading on cyclohexane (extent and time)	Performance Category
17	I) SIMULSOL SL8 II) TRITON X100	30% in 30 s Floats without spreading	Good Fair
18	I) SIMULSOL SL8 II) TRITON X100	50% in 30 s 100% in 30 seconds	Good Excellent
19	I) SIMULSOL SL8 II) TRITON X100	Sink immediately 70% in 40 s	Poor Good
20	I) SIMULSOL SL8 II) TRITON X100	100% in 6 seconds 50% in 30 s	Excellent Good
22	I) SIMULSOL SL8 II) TRITON X100	50% in 10 s Floats without spreading	Good Fair
23	I) SIMULSOL SL8 II) TRITON X100	20% in 20 s 100% in 25 s	Good Excellent
25	I) SIMULSOL SL8 II) TRITON X100	10% in 20 s Sink immediately	Good Poor
29	I) SIMULSOL SL8 II) TRITON X100	100% in 3 s 100% in 3 s	Excellent Excellent
B	I) SIMULSOL SL8 II) TRITON X100	Sink immediately Sink immediately	Poor Poor
C	I) SIMULSOL SL8 II) TRITON X100	Sink immediately Sink immediately	Poor Poor

Table 3 shows that the surfactants of the present invention, when combined with hydrocarbon surfactant SIMULSOL SL8 or TRITON X100 in an aqueous formulation, spread more quickly and more completely across cyclohexane than either Comparative Example B or C, which both sank. Spreading across cyclohexane is predictive of an effective fire fighting foam. Table 3 shows that low critical micelle concentrations and low Surface Tension levels beyond CMC are necessary but not sufficient criteria for an effective fire fighting foam.

CLAIMSWhat is claimed is:

1. A compound comprising Formula 5



5 wherein


R_f is $C_cF_{(2c+1)}$;

c is 2 to 6; and v is 2 to 4.

2. The compound of claim 1 wherein c is 3 or 4.

3. The compound of claim 1 wherein v is 2 or 3.

10 4. A process for the preparation of a compound of Formula 5



wherein


R_f is $C_cF_{(2c+1)}$;

c is 2 to 6; and v is 2 to 4,

15 comprising contacting a compound of Formula 6



wherein R_f is $C_cF_{(2c+1)}$, and c is 2 to 6, with a compound of Formula 7



wherein v is 2 to 4.

20 5. The process of claim 4 wherein c is 3 or 4.

6. The process of claim 5 wherein v is 2.

7. The process of claim 4 wherein the mol ratio of $R_f-O-CF=CF_2$ to $HO-(CH_2CH_2O)_v-H$ during the contacting is 1 to 15.

25 8. The process of claim 4 wherein the contacting is conducted in the presence of an alkali metal, alkali earth metal, alkali hydroxide, alkali hydride, or alkali amide.

9. The process of claim 8 wherein the alkali metal is Na, K, or Cs, and the alkali hydride is NaH or KH.