



(86) Date de dépôt PCT/PCT Filing Date: 1998/09/10
(87) Date publication PCT/PCT Publication Date: 1999/03/18
(45) Date de délivrance/Issue Date: 2005/02/08
(85) Entrée phase nationale/National Entry: 2000/03/09
(86) N° demande PCT/PCT Application No.: JP 1998/004087
(87) N° publication PCT/PCT Publication No.: 1999/013059
(30) Priorité/Priority: 1997/09/10 (9/245154) JP

(51) Cl.Int.⁷/Int.Cl.⁷ C12N 9/10, C12N 15/54, C12N 1/19,
C12P 19/00

(72) Inventeurs/Inventors:
YANAI, KOJI, JP;
NAKANE, AKITAKA, JP;
KONO, TOSHIAKI, JP

(73) Propriétaire/Owner:
MEIJI SEIKA KAISHA, LTD., JP

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : β -FRUCTOFURANOSIDASE ET GENE CORRESPONDANT
(54) Title: β -FRUCTOFURANOSIDASE AND ITS GENE

(57) **Abrégé/Abstract:**

A novel β -fructofuranosidase and its gene are disclosed. A polypeptide comprising the amino acid sequence of SEQ ID No. 1 or No. 3 is an enzyme having β -fructofuranosidase activity and high transferase activity, and is capable of efficiently producing fructooligosaccharides.



ABSTRACT

A novel β -fructofuranosidase and its gene are disclosed. A polypeptide comprising the amino acid sequence of SEQ ID No. 1 or No. 3 is an enzyme having β -fructofuranosidase activity and high transferase activity, and is capable of efficiently producing fructooligosaccharides.

β -FRUCTOFURANOSIDASE AND ITS GENEBackground of the InventionField of the Invention

5 The present invention relates to a β -fructofuranosidase having a fructose transferase activity, which is useful for the industrial production of fructooligosaccharides, and its gene and use.

Background Art

10 The molecular structure of a fructooligosaccharide is the same as that of sucrose, except that the fructose half of a fructooligosaccharide is coupled with another one to three fructose molecules at positions C1 and C2 via a β -bond. Fructooligosaccharides are indigestible sugars known for their
15 physiological advantages, such as the facilitation of Bifidobacterial growth in the intestines, metabolic stimulation for cholesterol and other lipids, and little cariogenicity.

 Fructooligosaccharides are found in plants, such as asparagus, onion, Jerusalem-artichoke and honey. They are also
20 synthesized from sucrose by the newly industrialized mass production technique using fructosyltransfer reaction which is catalyzed by a β -fructofuranosidase derived from a microorganism.

 The molecular structure of 1-kestose and nystose, which
25 make up component of industrially produced fructooligosaccharide mixtures of today, are the same as that of sucrose except that their fructose half is coupled with one and two molecules of fructose, respectively. It has been found recently that their high-purity crystals exhibit new desirable characteristics both
30 in physical properties and food processing purpose while maintaining the general physiological advantages of fructooligosaccharides (Japanese Patent Application No. 222923/1995, Japanese Laid-Open Publication No. 31160/1994). In this sense, they are fructooligosaccharide preparations having
35 new features.

 In consideration of the above, some of the inventors have already proposed an industrial process for producing crystal

1-ketose from sucrose (Japanese Patent Application No. 64682/1996, Japanese Patent Application No. 77534/1996, and Japanese Patent Application No. 77539/1996). According to this process, a β -fructofuranosidase harboring fructosyltransferase activity is first allowed to act on sucrose to produce 1-kestose; the resultant 1-kestose is fractionated to a purity of 80% or higher by chromatographic separation; then, using this fraction as a crystallizing sample, crystal 1-kestose is obtained at a purity of 95% or higher. The β -fructofuranosidase harboring fructosyltransferase activity used in this process should be able to produce 1-kestose from sucrose at a high yield while minimizing the byproduct nystose, which inhibits the reactions in the above steps of chromatographic separation and crystallization. In the enzyme derived from Aspergillus niger, which is currently used for the industrial production of fructooligosaccharides mixtures, the 1-kestose yield from sucrose is approximately 44%, while 7% is turned to nystose (Japanese Patent Application No. 64682/1996). These figures suggest that the enzyme has room for improvement in view of the industrial production of crystal 1-ketose.

As a next step, some of the inventors have successfully screened new enzymes having more favorable characteristics from Penicillium roqueforti and Scopulariopsis brevicaulis. These enzymes were able to turn 47% and 55% of sucrose into 1-kestose, respectively, and 7% and 4% to nystose (Japanese Patent Application No. 77534/1996, and Japanese Patent Application No. 77539/1996). These enzymes are inferior in productivity and stability to the enzyme derived from Aspergillus niger, and have room for improvement in view of the industrial production of crystal 1-ketose.

Thus, some of the inventors had paid attention to the procedure of genetic engineering as a process for improving the productivity of the enzyme, isolated the gene encoding β -fructofuranosidase from Penicillium roqueforti and Scopulariopsis brevicaulis, respectively, and conducted the structure analysis (PCT/JP97/00757). As a result, the translation regions encoding 565 amino acids and 574 amino acids as a mature protein were respectively deduced in the β -

fructofuranosidase genes from Penicillium roqueforti and Scopulariopsis brevicaulis and their expression products were shown to have β -fructofuranosidase activity, like the β -fructofuranosidase gene from Aspergillus niger (L.M. Boddy et al., Curr. Genet., 24, 60-66 (1993)).

Summary of the Invention

The inventors have now found that the addition of 38 and 39 amino acids to the C-terminal of the β -fructofuranosidase genes from Penicillium roqueforti and Scopulariopsis brevicaulis, which were previously found by some of the inventors, improves its activity.

Thus, an object of the present invention is to provide a novel β -fructofuranosidase and its gene.

The novel β -fructofuranosidase according to the present invention is a polypeptide comprising the amino acid sequence of SEQ ID No. 1 or 3 or a homologue thereof.

Furthermore, the gene according to the present invention is a DNA encoding the above polypeptide.

The amino acid sequence of SEQ ID No. 1 or 3 according to the present invention is constructed by adding 38 and 39 amino acids to the C-terminals of the β -fructofuranosidase genes from Penicillium roqueforti and Scopulariopsis brevicaulis, which were previously found by some of the inventors as described above.

It has been found that an intron actually exists at the region of the β -fructofuranosidase gene, which was presumed to encode the C-terminal amino acids by some of the present inventors and that the β -fructofuranosidase genes further encode 38 and 39 amino acids of the C-terminal. Surprisingly, the β -fructofuranosidase activity was remarkably improved by adding these amino acids to the C-terminal, as compared with the protein to which these sequences are not added.

Brief Description of the Drawing

Figures 1A, B, C and D show the construction of expression vector pYPEN02 in which a gene encoding the enzyme protein consisting of the amino acid sequence of SEQ ID No. 1 is introduced,

and expression vector pYPEN01 in which a gene encoding the enzyme protein consisting of the amino acid sequence from 1 to 565 of amino acid sequence of SEQ ID No. 1 is introduced.

Figures 2A and B show the construction of expression vector pYSCOP02 in which a gene encoding the enzyme protein consisting of the amino acid sequence of SEQ ID No. 3 is introduced, and expression vector pYSCOP01 in which a gene encoding the enzyme protein consisting of the amino acid sequence from 1 to 574 of amino acid sequence of SEQ ID No. 3 is introduced.

10

Detailed Description of the Invention

β -fructofuranosidase

The polypeptide according to the present invention comprises the amino acid sequence of SEQ ID No. 1 or 3. This polypeptide having the amino acid sequence of SEQ ID No. 1 or 3 has enzymatic activity as β -fructofuranosidase. The polypeptide according to the present invention involves a homologue of the amino acid sequence of SEQ ID No. 1 or 3 as shown in the sequence listing. The term "homologue" refers to an amino acid sequence in which one or more amino acids (for example, one to several amino acids) are inserted, substituted or deleted in, or added to either or both of the terminals of, the amino acid sequence of SEQ ID Nos. 1 and 3 while retaining β -fructofuranosidase activity. Such a homologue can be selected and produced by those skilled in the art without undue experiments by referring to the sequence of SEQ ID No. 1 or 3.

The β -fructofuranosidase having the amino acid sequence of SEQ ID Nos. 1 and 3 according to the present invention has high fructosyltransferase activity and efficiently produces fructooligosaccharides. Specifically, when a sucrose solution at a concentration of 30 wt% or more is used as a substrate for reaction, the fructosyltransferase activity of β -fructofuranosidase having the amino acid sequence of SEQ ID No. 1 is at least 4 times higher, and the fructosyltransferase activity of β -fructofuranosidase having the amino acid sequence of SEQ ID No. 3 is at least 7 times higher than hydrolytic activity. Furthermore, 50% or more of sucrose is converted to

fructooligosaccharides in both cases.

β -fructofuranosidase gene

The novel gene encoding β -fructofuranosidase according to the present invention comprises a DNA sequence encoding the amino acid sequence of SEQ ID Nos. 1 and 3 or a homologue thereof.

Generally, a nucleotide sequence which encodes the amino acid sequence of a given protein can be easily determined from the reference chart known as "codon table". A variety of nucleotide sequence are available from those encoding the amino acid sequence of SEQ ID No. 1 or 3. Therefore, the term "a nucleotide sequence encoding the amino acid sequence of SEQ ID No. 1 or 3" refers to the meaning including the nucleotide sequence of SEQ ID No. 2 or 4, as well as nucleotide sequences which consist of the same codons as above allowing for degeneracy and encode the amino acid sequence of SEQ ID No. 1 or 3.

A preferred embodiment of the present invention provides, as a preferred example of the novel gene according to the present invention, a DNA fragment comprising the nucleotide sequence of SEQ ID No. 2 or 4.

As described above, the present invention encompasses a homologue of the amino acid sequence of SEQ ID No. 1 or 3. Therefore, the DNA fragment according to the present invention involves a nucleotide sequence which encodes such a homologue.

As the nucleotide sequence of the DNA fragment according to the present invention is determined, the DNA fragment may be obtained according to the procedure for the synthesis of a nucleic acid.

This sequence can also be obtained from Penicillium roqueforti or Scopulariopsis brevicaulis, preferably Penicillium roqueforti IAM7254 or Scopulariopsis brevicaulis IFO4843, according to the procedure of genetic engineering.

Expression of β -fructofuranosidase Gene

The β -fructofuranosidase according to the present invention can be produced in a host cell which has been transformed by a DNA fragment encoding the enzyme. More specifically, a DNA fragment encoding the β -fructofuranosidase according to the present invention is introduced in a host cell in the form of

a DNA molecule which is replicatable in the host cell and can express the above gene, particularly an expression vector, in order to transform the host cell. Then, the obtained transformant is cultivated.

5 Therefore, the present invention provides a DNA molecule which comprises a gene encoding the β -fructofuranosidase according to the present invention, particularly an expression vector. This DNA molecule is obtained by introducing a DNA
10 fragment encoding the β -fructofuranosidase according to the present invention in a vector molecule. According to a preferred embodiment of the present invention, the vector is a plasmid.

 The DNA molecule according to the present invention may be prepared by the standard technique of genetic engineering.

 The vector applicable in the present invention can be
15 selected as appropriate from viruses, plasmids, cosmid vectors, etc., considering the type of the host cell used. For example, a bacteriophage in the λ phage group or a plasmid in the pBR or pUC group may be used for *E. coli* host cells, a plasmid in the pUB group for *Bacillus subtilis*, and a vector in the YE_p or YC_p
20 group for yeast.

 It is preferable that the plasmid contain a selectable marker to ensure the selection of the obtained transformance, such as a drug-resistance marker or marker gene complementing an auxotrophic mutation. Preferred example of marker genes
25 include ampicillin-resistance gene, kanamycin-resistance gene, and tetracycline-resistance gene for bacterium host cells; N-(5'-phosphoribosyl)-anthranilate isomerase gene (TRP1), orotidine-5'-phosphate decarboxylase gene (URA3), and β -isopropylmalate dehydrogenase gene (LEU2) for yeast; and
30 hygromycin-resistance gene (hph), bialophos-resistance gene (bar), and nitrate reductase gene (niaD) for mold.

 It is also preferable that the DNA molecule for use as an expression vector according to the present invention contain nucleotide sequence necessary for the expression of the β -
35 fructofuranosidase gene, including transcription and translation control signals, such as a promoter, a transcription initiation signal, a ribosome binding site, a translation

termination signal, and a transcription termination signal.

Examples of preferred promoters include, in addition to the promoter on the inserted fragment which is able to function in the host, promoters such as those of lactose operon (lac), and tryptophan operon (trp) for E. coli; promoters such as those of alcohol dehydrogenase gene (ADH), acid phosphatase gene (PHO), galactose regulated gene (GAL), and glyceraldehyde-3-phosphate dehydrogenase gene (GPD) for yeast; and promoters such as those of α -amylase gene (amy) and cellobiohydrolase I gene (CBHI) for mold.

When the host cell is Bacillus subtilis, yeast or mold, it is also advantageous to use a secretion vector to allow it to extracellularly secrete the produced recombinant β -fructofuranosidase. Any host cell with an established host-vector system may be used, preferably yeast, mold, etc. It is preferable also to use the mold fungus having no β -fructofuranosidase activity described in PCT/JP97/00757.

A novel recombinant enzyme produced by the transformant described above is obtained by the following procedure: first, the host cell described above is cultivated under suitable conditions to obtain the supernatant or cell bodies from the resultant culture, using a known technique such as centrifugation; cell bodies should be further suspended in a suitable buffer solution, then homogenized by freeze-and-thaw, ultrasonic treatment, or mortar, followed by centrifugation or filtration to separate a cell body extract containing the novel recombinant enzyme.

The enzyme can be purified by combining the standard techniques for separation and purification. Examples of such as techniques include processes such as heat treatment, which rely on the difference in thermal resistance; processes such as salt sedimentation and solvent sedimentation, which rely on the difference in solubility; processes such as dialysis, ultrafiltration and gel filtration, and SDS-polyacrylamide gel electrophoresis, which rely on the difference in molecular weight; processes such as ion exchange chromatography, which rely on the difference in electric charge; processes such as affinity

chromatography, which rely on specific affinity; processes such as hydrophobic chromatography and reversed-phase partition chromatography, which rely on the difference in hydrophobicity; and processes such as isoelectric focusing, which rely on the difference in isoelectric point.

Production of fructooligosaccharides using the β -fructofuranosidase

The present invention further provides a process for producing fructooligosaccharide using the recombinant host or recombinant β -fructofuranosidase described above.

In the process for producing fructooligosaccharides according to the present invention, the recombinant host or recombinant β -fructofuranosidase described above is brought into contact with sucrose.

The mode and conditions where the recombinant host or recombinant β -fructofuranosidase according to the present invention comes in contact with sucrose are not limited in any way provided that the novel recombinant enzyme is able to act on sucrose. A preferred embodiment for contact in solution is as follows: The sucrose concentration may be selected as appropriate in the range where sucrose can be dissolved. However, considering the conditions such as the specific activity of the enzyme and reaction temperature, the concentration should generally fall in the range of 5% to 80%, preferably 30% to 70%. The temperature and pH for the reaction of sucrose by the enzyme should preferably be optimized for the characteristics of the novel recombinant enzyme. Therefore, the reasonable conditions are about 30°C to 80°C, pH 4 to 10, preferably 40°C to 70°C, pH 5 to 7.

The degree of purification of the novel recombinant enzyme may be selected as appropriate. The enzyme may be used either as unpurified in the form of supernatant from a transformant culture or cell body homogenate, as purified after processed in various purification steps, or as isolated after processed by various purification means.

Furthermore, the enzyme may be brought into contact with sucrose as fixed on a carrier using the standard technique.

The fructooligosaccharides thus produced are purified from the resulting solution according to known procedures. For example, the solution may be heated to inactivate the enzyme, decolorized using activated carbon, then desalted using ion exchange resin.

Examples

Example 1: Determination of translation region of β -fructofuranosidase gene from Penicillium roqueforti IAM7254

A DNA fragment of about 2 kbp containing the β -fructofuranosidase gene from Aspergillus niger was amplified by PCR using a chromosomal DNA prepared from Aspergillus niger ATCC20611 according to the standard procedure as a template and synthetic DNAs of SEQ ID Nos. 5 and 6 as primers. This DNA fragment was fractionated by agarose gel electrophoresis, extracted according to the standard procedure, purified, and then dissolved in sterilized water to 0.1 $\mu\text{g}/\mu\text{l}$ to prepare a DNA sample for probe.

In the next step, a chromosomal DNA from Penicillium roqueforti IAM7254 was prepared, about 20 μg of the chromosomal DNA was digested completely with EcoRI, followed by agarose gel electrophoresis to recover about 4 kbp DNA fragments.

The recovered DNA fragments of about 4 kbp (about 0.5 μg) were ligated with 1 μg of $\lambda\text{gt}10$ vector which had been digested with EcoRI and treated with phosphatase, packaged using an in vitro packaging kit, GIGAPACK^{*} II Gold (Stratagene L.L.C.), and then introduced in E. coli NM514, to prepare a library.

A probe was prepared from DNA sample for probe above described. As a result of plaque hybridization using ECL^{*} Direct DNA/RNA Labelling & Detection System (Amersham International), 4 clones turned out positive in about 25,000 plaques. These positive clones were purified by a second screening to prepare phage DNA, which was then analyzed using restriction enzymes. The result showed that all the clones had an identical EcoRI fragment of about 4 kbp.

The EcoRI fragments of about 4 kbp were subdivided into a small fragment to select the desired DNA region using restriction enzymes, then subcloned to plasmid vector pUC118 or pUC119. The plasmid DNA was obtained from the subclone according

^{*}Trade-mark

10

to the standard procedure and sequenced using ALFred DNA Sequencer (Pharmacia) as shown in SEQ ID No.7.

5 The sequence consisting 50 bases from 1695 to 1744 in this sequence was identified as an intron because it showed a typical intron structure of filamentous fungi. As a result, the sequence of SEQ ID No. 2 as a sequence encoding protein was obtained by deleting the intron from the sequence of SEQ ID No. 7. The encoded amino acid sequence was shown in SEQ ID No. 1.

10 Example 2: Expression of β -fructofuranosidase gene from *Penicillium roqueforti* IAM7254 in *Saccharomyces cerevisiae*

Plasmid pYPEN01 and pYPEN02 for expressing the β -fructofuranosidase gene from *Penicillium roqueforti* were prepared as follows (Figure 1A, B, C and D).

15 pYPR2831 (H. Horiuchi et al., Agric. Biol. Chem., 54, 1771-1779, 1990) was digested with EcoRI and SalI, and then its terminals were blunted with T4 DNA polymerase. The obtained fragment was ligated with BamHI linker (5'-CGGATCCG-3'), digested with BamHI, followed by self-ligation to obtain vector pY2831 for expression in yeast.

20 Next, single-stranded DNA was prepared from the plasmid pPRS01 obtained by inserting an about 4 kbp EcoRI DNA fragment containing the β -fructofuranosidase gene prepared in Example 1 into plasmid pUC118. Using the single-stranded DNA as a template and a synthetic DNA of SEQ ID No. 8 as a primer, the translated region of the β -fructofuranosidase gene was subjected to site-specific mutagenesis to disrupt the BamHI site without changing the encoded amino acid sequence (pPRS02).

30 A part of the translated region of the β -fructofuranosidase gene was prepared as an about 1.8 kbp BamHI fragment by PCR using plasmid pPRS02 as a template and synthetic DNAs of SEQ ID Nos. 9 and 10 as primers, and inserted into the BamHI site of plasmid pY2831 to prepare pYPEN01. Thus, plasmid pYPEN01 is designed to produce an enzyme protein comprising an amino acid sequence from 1 to 565 in the amino acid sequence of SEQ ID No. 1, which is a mature β -fructofuranosidase following secretion signal sequence.

35 Further, a DNA fragment containing the translated region
*Trade-mark

of the β -fructofuranosidase gene was prepared as an about 1.8 kbp BamHI fragment by PCR using plasmid pPRS02 as a template and synthetic DNAs of SEQ ID Nos. 9 and 11 as primers, and inserted into the BamHI site of plasmid pUC118 to prepare plasmid pPRS03.

5 A single-stranded DNA was prepared from plasmid pPRS03. As a result of site-specific mutagenesis using this as a template and a synthetic DNA of SEQ ID No. 12 as a primer, the intron sequence was removed (pPRS04). The translated region of the β -fructofuranosidase gene was prepared as an about 1.8 kbp BamHI

10 fragment from plasmid pPRS04, and inserted into the BamHI site of plasmid pY2831 to prepare plasmid pYPEN02. Thus, plasmid pYPEN02 is designed to produce an enzyme protein comprising an amino acid sequence of SEQ ID No. 1, which is a mature β -fructofuranosidase following secretion signal sequence.

15 Plasmids pYPEN01 and pYPEN02 were introduced into Saccharomyces cerevisiae MS-161 (Suc⁻, ura3, trp1) by the lithium-acetate method (Ito, H. et al., J. Bacteriol., 153, 163-168, 1983) to obtain transformants. The transformants were cultivated in an SD-Ura medium (0.67% yeast nitrogen base

20 (Difco), 2% glucose and 50 μ g/ml uracil) at 30°C overnight. The culture was seeded in a production medium (0.67% yeast nitrogen base (Difco), 2% glucose, 2% casamino acid and 50 μ g/ml uracil) at a final concentration of 1% and cultivated at 30°C for 2 days. The culture supernatant was analyzed for β -fructofuranosidase

25 activity, in units, i.e., the quantity of free glucose (μ mol) released in 1 minute in 10 wt% sucrose solution, pH 5.5, at 40°C for 60 minutes. As a result, the transformant with plasmid pYPEN01 exhibited 4×10^{-4} units/ml or less of activity while the transformant with plasmid pYPEN02 exhibited 0.38 units/ml of

30 activity.

Example 3: Determination of the translated region of β -fructofuranosidase gene from Scopulariopsis brevicaulis IFO4843

The chromosomal DNA was prepared from Scopulariopsis brevicaulis IFO4843. About 20 μ g of a chromosomal DNA sample was

35 completely digested with EcoRI, and electrophoresed through an agarose gel to recover an about 10 kbp DNA fragment.

The recovered DNA fragment of about 10 kbp (about 0.5 µg) were ligated with 1 µg of λDASHII vector digested with HindIII and EcoRI, and packaged using an in vitro packaging kit, GIGAPACK II Gold (Stratagene L.L.C.), then introduced in E. coli XL1-Blue MRA (P2), to prepare a library.

As a result of plaque hybridization using ECL Direct DNA/RNA Labelling & Detection System (Amersham International), with the about 2 kbp DNA fragment used in Example 1 as a probe, 3 clones turned out positive in about 15,000 plaques. These positive clones were purified by a second screening to prepare phage DNA, which was then analyzed using restriction enzymes. The result showed that all the clones had an identical EcoRI fragment of about 10 kbp.

These EcoRI fragments of about 10 kbp were subdivided into a small fragment to select the desired DNA region using restriction enzymes, then subcloned to plasmid vector pUC118 or pUC119. The plasmid DNA was obtained from the subclone according to the standard procedure and sequenced using ALFred DNA Sequencer (Pharmacia) as shown in SEQ ID No.13.

The sequence comprising 55 bases from 1722 to 1776 in this sequence was identified as an intron because it showed a typical intron structure of filamentous fungi. As a result, the sequence of SEQ ID No. 4 as a sequence encoding protein was obtained by deleting the intron from the sequence of SEQ ID No. 13. The encoded amino acid sequence was shown SEQ ID No. 3.

Example 4: Expression of β-fructofuranosidase gene from Scopulariopsis brevicaulis IFO4843 in Saccharomyces cerevisiae

Plasmid pYSCOP01 and pYSCOP02 for expressing the β-fructofuranosidase gene from Scopulariopsis brevicaulis were prepared as follows (Figure 2A and B).

A part of the translated region of the β-fructofuranosidase gene was prepared as an about 1.8 kbp BamHI fragment by PCR using about 10 kbp EcoRI DNA fragment prepared in Example 3 containing the β-fructofuranosidase gene as a template and synthetic DNAs of SEQ ID Nos. 14 and 15 as primers, and inserted into the BamHI site of plasmid pY2831 to prepare pYSCOP01. Thus, plasmid pYSCOP01 is designed to produce an enzyme protein comprising an

amino acid sequence from 1 to 574 in the amino acid sequence of SEQ ID No. 3, which is a mature β -fructofuranosidase following secretion signal sequence.

Next, a DNA fragment containing the translated region of the β -fructofuranosidase gene was prepared as an about 1.9 kbp BamHI fragment by PCR using an about 10 kbp EcoRI fragment containing the β -fructofuranosidase gene as a template and synthetic DNAs of SEQ ID Nos. 14 and 16 as primers, and inserted into the BamHI site of plasmid pUC118 to prepare plasmid pSCB01.

A single-stranded DNA was prepared from plasmid pSCB01. As a result of site-specific mutagenesis using this as a template and the synthetic DNA of SEQ ID No. 17 as a primer, the intron sequence was removed (pSCB02). The translated region of the β -fructofuranosidase gene was prepared as an about 1.9 kbp BamHI fragment from plasmid pSCB02, and inserted into the BamHI site of plasmid pY2831 to prepare plasmid pYSCOP02. Thus, plasmid pYSCOP02 is designed to produce an enzyme protein comprising an amino acid sequence of SEQ ID No. 3, which is a mature β -fructofuranosidase following secretion signal sequence.

Plasmids pYSCOP01 and pYSCOP02 were introduced into Saccharomyces cerevisiae MS-161 (Suc⁻, ura3, trp1) by the lithium-acetate method to obtain transformants. The transformants were cultivated in an SD-Ura medium at 30°C overnight. The culture was seeded a production medium at a final concentration of 1% and cultivated at 30°C for 2 days. The culture supernatant was analyzed for β -fructofuranosidase activity in the same manner as described in Example 2. As a result, the transformant with plasmid pYSCOP01 exhibited 4×10^{-4} units/ml or less of activity, while the transformant with plasmid pYSCOP02 exhibited 6.5×10^{-3} units/ml of activity.

14

SEQUENCE LISTING

<110> MEIJI SEIKA KAISHA, LTD.

<120> Beta-fructofuranosidase and its gene

<130> 116875-475

<140>

<141> 1998-9-10

<150> JP/245154/1997

10 <151> 1997-9-10

<160> 17

<170> PatentIn* Ver. 2.0

<210> 1

<211> 603

<212> PRT

<213> Penicillium roqueforti IAM7254

<220>

<221> mat peptide

<222> (1)...(603)

20 <400> 1

Val	Asp	Phe	His	Thr	Pro	Ile	Asp	Tyr	Asn	Ser	Ala	Pro	Pro	Asn	Leu
1				5					10					15	

Ser	Thr	Leu	Ala	Asn	Ala	Ser	Leu	Phe	Lys	Thr	Trp	Arg	Pro	Arg	Ala
		20						25					30		

His	Leu	Leu	Pro	Pro	Ser	Gly	Asn	Ile	Gly	Asp	Pro	Cys	Gly	His	Tyr
	35					40						45			

Thr	Asp	Pro	Lys	Thr	Gly	Leu	Phe	His	Val	Gly	Trp	Leu	Tyr	Ser	Gly
	50				55					60					

Ile	Ser	Gly	Ala	Thr	Thr	Asp	Asp	Leu	Val	Thr	Tyr	Lys	Asp	Leu	Asn
65				70					75					80	

Pro	Asp	Gly	Ala	Pro	Ser	Ile	Val	Ala	Gly	Gly	Lys	Asn	Asp	Pro	Leu
		85						90						95	

Ser	Val	Phe	Asp	Gly	Ser	Val	Ile	Pro	Ser	Gly	Ile	Asp	Gly	Met	Pro
		100					105						110		

*Trade-mark

	Thr	Leu	Leu	Tyr	Thr	Ser	Val	Ser	Tyr	Leu	Pro	Ile	His	Trp	Ser	Ile	
			115					120					125				
	Pro	Tyr	Thr	Arg	Gly	Ser	Glu	Thr	Gln	Ser	Leu	Ala	Val	Ser	Tyr	Asp	
		130					135					140					
10	Gly	Gly	His	Asn	Phe	Thr	Lys	Leu	Asn	Gln	Gly	Pro	Val	Ile	Pro	Thr	
		145				150					155					160	
	Pro	Pro	Phe	Ala	Leu	Asn	Val	Thr	Ala	Phe	Arg	Asp	Pro	Tyr	Val	Phe	
					165					170					175		
	Gln	Ser	Pro	Ile	Leu	Asp	Lys	Ser	Val	Asn	Ser	Thr	Gln	Gly	Thr	Trp	
				180					185					190			
	Tyr	Val	Ala	Ile	Ser	Gly	Gly	Val	His	Gly	Val	Gly	Pro	Cys	Gln	Phe	
20			195					200					205				
	Leu	Tyr	Arg	Gln	Asn	Asp	Ala	Asp	Phe	Gln	Tyr	Trp	Glu	Tyr	Leu	Gly	
		210					215					220					
	Gln	Trp	Trp	Lys	Glu	Pro	Leu	Asn	Thr	Thr	Trp	Gly	Lys	Gly	Asp	Trp	
		225				230					235					240	
	Ala	Gly	Gly	Trp	Gly	Phe	Asn	Phe	Glu	Val	Gly	Asn	Val	Phe	Ser	Leu	
					245					250					255		
30	Asn	Ala	Glu	Gly	Tyr	Ser	Glu	Asp	Gly	Glu	Ile	Phe	Ile	Thr	Leu	Gly	
				260					265					270			
	Ala	Glu	Gly	Ser	Gly	Leu	Pro	Ile	Val	Pro	Gln	Val	Ser	Ser	Ile	Arg	
			275					280					285				
	Asp	Met	Leu	Trp	Val	Thr	Gly	Asn	Val	Thr	Asn	Asp	Gly	Ser	Val	Thr	
		290					295					300					
40	Phe	Lys	Pro	Thr	Met	Ala	Gly	Val	Leu	Asp	Trp	Gly	Val	Ser	Ala	Tyr	
		305				310					315					320	
	Ala	Ala	Ala	Gly	Lys	Ile	Leu	Pro	Ala	Ser	Ser	Gln	Ala	Ser	Thr	Lys	
					325					330					335		
	Ser	Gly	Ala	Pro	Asp	Arg	Phe	Ile	Ser	Tyr	Val	Trp	Leu	Thr	Gly	Asp	
				340					345					350			
	Leu	Phe	Glu	Gln	Val	Lys	Gly	Phe	Pro	Thr	Ala	Gln	Gln	Asn	Trp	Thr	
			355				360						365				
50	Gly	Ala	Leu	Leu	Leu	Pro	Arg	Glu	Leu	Asn	Val	Arg	Thr	Ile	Ser	Asn	
		370					375					380					
	Val	Val	Asp	Asn	Glu	Leu	Ser	Arg	Glu	Ser	Leu	Thr	Ser	Trp	Arg	Val	
						390					395					400	
	Ala	Arg	Glu	Asp	Ser	Gly	Gln	Ile	Asp	Leu	Glu	Thr	Met	Gly	Ile	Ser	
					405					410					415		
60	Ile	Ser	Arg	Glu	Thr	Tyr	Ser	Ala	Leu	Thr	Ser	Gly	Ser	Ser	Phe	Val	
				420					425						430		

Glu Ser Gly Lys Thr Leu Ser Asn Ala Gly Ala Val Pro Phe Asn Thr
 435 440 445
 Ser Pro Ser Ser Lys Phe Phe Val Leu Thr Ala Asn Ile Ser Phe Pro
 450 455 460
 10 Thr Ser Ala Arg Asp Ser Gly Ile Gln Ala Gly Phe Gln Val Leu Ser
 465 470 475 480
 Ser Ser Leu Glu Ser Thr Thr Ile Tyr Tyr Gln Phe Ser Asn Glu Ser
 485 490 495
 Ile Ile Val Asp Arg Ser Asn Thr Ser Ala Ala Ala Arg Thr Thr Ala
 500 505 510
 Gly Ile Leu Ser Asp Asn Glu Ala Gly Arg Leu Arg Leu Phe Asp Val
 515 520 525
 20 Leu Arg Asn Gly Lys Glu Gln Val Glu Thr Leu Glu Leu Thr Ile Val
 530 535 540
 Val Asp Asn Ser Val Leu Glu Val Tyr Ala Asn Gly Arg Phe Ala Leu
 545 550 555 560
 Gly Thr Trp Ala Arg Ser Trp Tyr Ala Asn Ser Thr Lys Ile Asn Phe
 565 570 575
 30 Phe His Asn Gly Val Gly Glu Ala Thr Phe Glu Asp Val Thr Val Phe
 580 585 590
 Glu Gly Leu Tyr Asp Ala Trp Pro Gln Arg Lys
 595 600

<210> 2

<211> 1809

<212> DNA

<213> Penicillium roqueforti IAM7254

40 <220>

<221> mat peptide

<222> (1)...(1809)

<400> 2

gttgatttcc ataccgccgat tgactataac tcggctccgc caaacctttc taccctggca 60
 aacgcattctc ttttcaagac atggagaccc agagcccatc ttctccctcc atctgggaac 120
 ataggcgacc cgtgcgggca ctataccgat cccaagactg gtctcttcca cgtggggttg 180
 ctttacagtg ggatttcggg agcgacaacc gacgatctcg ttacctataa agacctcaat 240
 cccgatggag ccccgatcaat tgttgcagga ggaaagaacg accctctttc tgtcttcgat 300

ggctcgggtca ttccaagcgg tatagacggc atgccaactc ttctgtatac ctctgtatca 360
 tacctcccaa tccactgggtc catcccctac acccggggaa gcgagacaca atccttggcc 420
 gtttcctatg acggtgggtca caacttcacc aagctcaacc aaggggccgt gatccctacg 480
 cctccgtttg ctctcaatgt caccgctttc cgtgaccctt acgttttcca aagcccaatt 540
 ctggacaaat ctgtcaatag taccgaagga acatgggtatg tcgccatatac tggcgggtgtc 600
 cacgggtgtcg gaccttgtca gttcctctac cgtcagaacg acgcagattt tcaatattgg 660
 gaatatctcg ggcaatgggtg gaaggagccc ctttaatacca cttggggaaa gggtgactgg 720
 gccggggggtt ggggcttcaa ctttgagggtt ggcaacgtct ttagtctgaa tgcagagggg 780
 10 tatagtgaag acggcgagat attcataacc ctcggtgctg agggttcggg acttcccatac 840
 gttcctcaag tctcctctat tcgcgatatg ctgtgggtga ccggcaatgt cacaatgac 900
 ggctctgtca ctttcaagcc aaccatggcg ggtgtgcttg actggggcgt gtcggcatat 960
 gctgctgcag gcaagatctt gccggccagc tctcaggcat ccacaaagag cggtgcccc 1020
 gatcggttca tttcctatgt ctggctcact ggagatctat tcgagcaagt gaaaggattc 1080
 cctaccgctc aacaaaactg gaccggggcc ctcttactgc cgcgagagct gaatgtccgc 1140
 actatctcta acgtgggtga taacgaactt tcgcgtgagt ccttgacatac gtggcgcggtg 1200
 gccgcgaag actctgggtca gatcgacctt gaaacaatgg gaatctcaat ttccagggag 1260
 acttacagcg ctctcacatac cggctcatct tttgtcgagt ctggtaaaac gttgtcgaat 1320
 gctggagcag tgcccttcaa tacctcacc tcaagcaagt tcttcgtgct gacagcaaata 1380
 20 atatctttcc cgacctctgc ccgtgactct ggcattccagg ctggtttcca ggttttatcc 1440
 tctagtcttg agtctacaac tatctactac caattctcca acgagtccat catcgtcgac 1500
 cgcagcaaca cgagtgtctg gccgagaaca actgctggga tcctcagtga taacgaggcg 1560
 ggacgtctgc gcctcttcga cgtgttgoga aatggaaaag aacagggttg aactttggag 1620
 ctactatcg tgggtggataa tagtgtactg gaagtatatg ccaatggacg ctttgctcta 1680
 ggcacttggg ctcggtcttg gtacgccaac tcgactaaaa ttaacttctt ccataacggc 1740
 gtgggagaag cgacattcga agatgtgacg gtctttgaag gactgtatga tgcctggcca 1800
 caaaggaag 1809

<210> 3

30 <211> 613

<212> PRT

<213> Scopulariopsis brevicaulis IFO4843

<220>

<221> mat peptide

<222> (1)...(613)

<400> 3

	Gln	Pro	Thr	Ser	Leu	Ser	Ile	Asp	Asn	Ser	Thr	Tyr	Pro	Ser	Ile	Asp	
	1				5					10					15		
10	Tyr	Asn	Ser	Ala	Pro	Pro	Asn	Leu	Ser	Thr	Leu	Ala	Asn	Asn	Ser	Leu	
				20					25					30			
	Phe	Glu	Thr	Trp	Arg	Pro	Arg	Ala	His	Val	Leu	Pro	Pro	Gln	Asn	Gln	
			35					40					45				
	Ile	Gly	Asp	Pro	Cys	Met	His	Tyr	Thr	Asp	Pro	Glu	Thr	Gly	Ile	Phe	
		50					55					60					
20	His	Val	Gly	Trp	Leu	Tyr	Asn	Gly	Asn	Gly	Ala	Ser	Gly	Ala	Thr	Thr	
	65					70				75						80	
	Glu	Asp	Leu	Val	Thr	Tyr	Gln	Asp	Leu	Asn	Pro	Asp	Gly	Ala	Gln	Met	
					85					90					95		
	Ile	Leu	Pro	Gly	Gly	Val	Asn	Asp	Pro	Ile	Ala	Val	Phe	Asp	Gly	Ala	
				100					105					110			
	Val	Ile	Pro	Ser	Gly	Ile	Asp	Gly	Lys	Pro	Thr	Met	Met	Tyr	Thr	Ser	
30			115					120					125				
	Val	Ser	Tyr	Met	Pro	Ile	Ser	Trp	Ser	Ile	Ala	Tyr	Thr	Arg	Gly	Ser	
		130					135					140					
	Glu	Thr	His	Ser	Leu	Ala	Val	Ser	Ser	Asp	Gly	Gly	Lys	Asn	Phe	Thr	
	145					150					155					160	
	Lys	Leu	Val	Gln	Gly	Pro	Val	Ile	Pro	Ser	Pro	Pro	Phe	Gly	Ala	Asn	
					165					170					175		
40	Val	Thr	Ser	Trp	Arg	Asp	Pro	Phe	Leu	Phe	Gln	Asn	Pro	Gln	Phe	Asp	
				180					185					190			
	Ser	Leu	Leu	Glu	Ser	Glu	Asn	Gly	Thr	Trp	Tyr	Thr	Val	Ile	Ser	Gly	
			195					200					205				
	Gly	Ile	His	Gly	Asp	Gly	Pro	Ser	Ala	Phe	Leu	Tyr	Arg	Gln	His	Asp	
		210					215					220					
50	Pro	Asp	Phe	Gln	Tyr	Trp	Glu	Tyr	Leu	Gly	Pro	Trp	Trp	Asn	Glu	Glu	
	225					230					235				240		
	Gly	Asn	Ser	Thr	Trp	Gly	Ser	Gly	Asp	Trp	Ala	Gly	Arg	Trp	Gly	Tyr	
					245					250					255		
	Asn	Phe	Glu	Val	Ile	Asn	Ile	Val	Gly	Leu	Asp	Asp	Asp	Gly	Tyr	Asn	
				260					265					270			

	Pro	Asp	Gly	Glu	Ile	Phe	Ala	Thr	Val	Gly	Thr	Glu	Trp	Ser	Phe	Asp
			275					280					285			
	Pro	Ile	Lys	Pro	Gln	Ala	Ser	Asp	Asn	Arg	Glu	Met	Leu	Trp	Ala	Ala
		290					295					300				
10	Gly	Asn	Met	Thr	Leu	Glu	Asp	Gly	Asp	Ile	Lys	Phe	Thr	Pro	Ser	Met
	305					310					315					320
	Ala	Gly	Tyr	Leu	Asp	Trp	Gly	Leu	Ser	Ala	Tyr	Ala	Ala	Ala	Gly	Lys
					325					330					335	
	Glu	Leu	Pro	Ala	Ser	Ser	Lys	Pro	Ser	Gln	Lys	Ser	Gly	Ala	Pro	Asp
				340					345					350		
	Arg	Phe	Val	Ser	Tyr	Leu	Trp	Leu	Thr	Gly	Asp	Tyr	Phe	Glu	Gly	His
			355					360					365			
20	Asp	Phe	Pro	Thr	Pro	Gln	Gln	Asn	Trp	Thr	Gly	Ser	Leu	Leu	Leu	Pro
		370					375					380				
	Arg	Glu	Leu	Ser	Val	Gly	Thr	Ile	Pro	Asn	Val	Val	Asp	Asn	Glu	Leu
	385					390					395					400
	Ala	Arg	Glu	Thr	Gly	Ser	Trp	Arg	Val	Gly	Thr	Asn	Asp	Thr	Gly	Val
					405					410					415	
30	Leu	Glu	Leu	Val	Thr	Leu	Lys	Gln	Glu	Ile	Ala	Arg	Glu	Thr	Leu	Ala
				420					425					430		
	Glu	Met	Thr	Ser	Gly	Asn	Ser	Phe	Thr	Glu	Ala	Ser	Arg	Asn	Val	Ser
			435					440					445			
	Ser	Pro	Gly	Ser	Thr	Ala	Phe	Gln	Gln	Ser	Leu	Asp	Ser	Lys	Phe	Phe
		450					455					460				
40	Val	Leu	Thr	Ala	Ser	Leu	Ser	Phe	Pro	Ser	Ser	Ala	Arg	Asp	Ser	Asp
	465					470					475					480
	Leu	Lys	Ala	Gly	Phe	Glu	Ile	Leu	Ser	Ser	Glu	Phe	Glu	Ser	Thr	Thr
					485					490					495	
	Val	Tyr	Tyr	Gln	Phe	Ser	Asn	Glu	Ser	Ile	Ile	Ile	Asp	Arg	Ser	Asn
				500					505					510		
	Ser	Ser	Ala	Ala	Ala	Leu	Thr	Thr	Asp	Gly	Ile	Asp	Thr	Arg	Asn	Glu
			515					520					525			
50	Phe	Gly	Lys	Met	Arg	Leu	Phe	Asp	Val	Val	Glu	Gly	Asp	Gln	Glu	Arg
		530					535					540				
	Ile	Glu	Thr	Leu	Asp	Leu	Thr	Ile	Val	Val	Asp	Asn	Ser	Ile	Val	Glu
	545					550					555					560
	Val	His	Ala	Asn	Gly	Arg	Phe	Ala	Leu	Ser	Thr	Trp	Val	Arg	Ser	Trp
				565						570					575	
60	Tyr	Glu	Ser	Ser	Lys	Asp	Ile	Lys	Phe	Phe	His	Asp	Gly	Asp	Ser	Thr
				580					585					590		

Val Gln Phe Ser Asn Ile Thr Val Tyr Glu Gly Leu Phe Asp Ala Trp
 595 600 605

Pro Glu Arg Ala Arg
 610

<210> 4

10 <211> 1839

<212> DNA

<213> Scopulariopsis brevicaulis IF04843

<220>

<221> mat peptide

<222> (1)...(1839)

<400> 4

caacctacgt ctctgtcaat cgacaattcc acgtatcctt ctatcgacta caactccgcc 60
 cctccaaacc tctcgactct tgccaacaac agcctcttcg agacatggag gccgagggca 120
 cacgtccttc cgccccagaa ccagatcggc gatccgtgta tgcactacac cgacccccgag 180
 20 acaggaatct tccacgtcgg ctggctgtac aacggcaatg gcgcttccgg cgccacgacc 240
 gaggatctcg tcacctatca ggatctcaac cccgacggag cgcagatgat ccttccgggt 300
 ggtgtgaatg accccattgc tgtctttgac ggcgcgggta ttcccagtgg cattgatggg 360
 aaaccaccca tgatgtatac ctcggtgtca tacatgcccc tctcctggag catcgcttac 420
 accaggggaa gcgagaccca ctctctcgca gtgtcgtccg acggcggtaa gaacttcacc 480
 aagctgggtgc agggccccgt cattccttcg cctcccttcg gcgccaacgt gaccagctgg 540
 cgtgaccctt tctgtttcca aaacccccag ttcgactctc tctcgaag cgagaacggc 600
 acgtggtaca ccgttatctc tgggtggcatc cacggtgacg gcccctccgc gttcctctac 660
 cgtcagcacg accccgactt ccagtactgg gagtaccttg gaccgtgggtg gaacgaggaa 720
 gggaactcga cctggggcag cggtgactgg gctggccgggt ggggctacaa cttcgaggtc 780
 30 atcaacattg tcggtcttga cgatgatggc tacaaccccg acggtgaaat ctttgccacg 840
 gtaggtaccg aatggtcggt tgaccccatc aaaccgcagg cctcggacaa caggagatg 900
 ctctgggccg cgggcaacat gactctcgag gacggcgata tcaagttcac gccaaagcatg 960
 gcgggctacc tcgactgggg tctatcggcg tatgccgccg ctggcaagga gctgcccgtc 1020
 tcttcaaagc cttcgcagaa gagcgggtgc cggaccgggt tcgtgtcgta cctgtggctc 1080
 accggtgact acttcgaggg ccacgacttc cccaccccg cgcagaattg gaccggctcg 1140

	cttttgcttc	cgcgtagagct	gagcgtcggg	acgattccca	acgttggtcga	caacgagctt	1200
	gctcgcgaga	cgggctcttg	gagggttggc	accaacgaca	ctggcgtgct	tgagctggtc	1260
	actctgaagc	aggagattgc	tcgcgagacg	ctggctgaaa	tgaccagcgg	caactccttc	1320
	accgaggcga	gcaggaatgt	cagctcgccc	ggatctaccg	ccttccagca	gtccctggat	1380
	tccaagttct	tcgtcctgac	cgctcgctc	tccttccctt	cgtcggctcg	cgactccgac	1440
	ctcaaggctg	gtttcgagat	cctgtcgctc	gagtttgagt	cgaccacggt	ctactaccag	1500
	ttttccaacg	agtccatcat	cattgaccgg	agcaactcga	gtgctgccgc	cttgactacc	1560
10	gatggaatcg	acacccgcaa	cgagtttggc	aagatgcgcc	tgtttgatgt	tgtcgagggg	1620
	gaccaggagc	gtatcgagac	gctcgatctc	actattgtgg	ttgataactc	gatcgttgag	1680
	gttcatgcca	acgggcgatt	cgctctgagc	acttggggtt	gttcgtggta	cgagtcgtcc	1740
	aaggacatca	agttcttcca	cgatggcgac	agcacgggtt	agttctcgaa	catcacccgc	1800
	tacgaggggac	tgtttgacgc	ctggccggag	cgggccagg			1839

	<210> 5	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
20	<400> 5	
	caatgaagct caccactacc	20
	<210> 6	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<400> 6	
	atccccggtca atttctctcc	20
30	<210> 7	
	<211> 1859	

22

<212> DNA

<213> *Penicillium roqueforti* IAM7254

<220>

<221> CDS

<222> (1)...(1694)

<221> intron

<222> (1695)...(1744)

<221> CDS

<222> (1745)...(1859)

10 <400> 1

gttgatttcc ataccccgat tgactataac tcggctccgc caaacctttc taccctggca 60

aacgcatctc ttttcaagac atggagaccc agagcccatc ttctccctcc atctgggaac 120

ataggcgacc cgtgcgggca ctataccgat cccaagactg gtctcttcca cgtgggttgg 180

ctttacagtg ggatttcggg agcgacaacc gacgatctcg ttacctataa agacctcaat 240

cccgatggag ccccgatcaat tggtgcagga ggaaagaacg accctctttc tgtcttcgat 300

ggctcgggtca ttccaagcgg tatagacggc atgccaaactc ttctgtatac ctctgtatca 360

tacctcccaa tccactggtc catcccctac acccggggaa gcgagacaca atccttggcc 420

gtttcctatg acggtgggtca caacttcacc aagctcaacc aagggcccggt gatccctacg 480

cctccgtttg ctctcaatgt caccgctttc cgtgaccctt acgttttcca aagcccaatt 540

20 ctggacaaat ctgtcaatag tacccaagga acatggtatg tcgccatatac tggcgggtgtc 600

cacggtgtcg gaccttgtca gttcctctac cgtcagaacg acgcagattt tcaatattgg 660

gaatatctcg ggcaatgggtg gaaggagccc cttaatacca cttggggaaa gggtgactgg 720

gccggggggtt ggggcttcaa ctttgagggtt ggcaacgtct ttagtctgaa tgcagagggg 780

tatagtgaag acggcgagat attcataacc ctcggtgctg agggttcggg acttcccatc 840

gttcctcaag tctcctctat tcgcatatg ctgtgggtga ccggcaatgt cacaaatgac 900

ggctctgtca ctttcaagcc aaccatggcg ggtgtgcttg actggggcgt gtcggcatat 960

gctgctgcag gcaagatctt gccggccagc tctcaggcat ccacaaagag cggtgccccc 1020

gatcggttca tttcctatgt ctggctcact ggagatctat tcgagcaagt gaaaggattc 1080

cctaccgctc aacaaaactg gaccggggcc ctcttactgc cgcgagagct gaatgtccgc 1140

30 actatctcta acgtgggtgga taacgaactt tcgcgtgagt ccttgacatc gtggcgcggtg 1200

gcccgcgaag actctgggtca gatcgacctt gaaacaatgg gaatctcaat ttccagggag 1260

acttacagcg ctctcacatc cggctcatct tttgtcgagt ctggtaaaac gttgtcgaat 1320
 gctggagcag tgcccttcaa tacctcaccc tcaagcaagt tcttcgtgct gacagcaaatt 1380
 atatctttcc cgacctctgc ccgtgactct ggcattccagg ctggttttcca ggttttatcc 1440
 tctagtcttg agtctacaac tatctactac caattctcca acgagtccat catcgtcgac 1500
 cgcagcaaca cgagtgtgctc ggcgagaaca actgctggga tcctcagtga taacgaggcg 1560
 ggacgtctgc gcctcttcga cgtgttgcca aatggaaaag aacagggtga aactttggag 1620
 ctactatcg tggtaggataa tagtgtactg gaagtatatg ccaatggacg ctttgctcta 1680
 ggcacttggg ctcggttaagt ctctcttggt tatggaagat tggtaaaaaa ctaaccgcat 1740
 10 gaaggtcttg gtacgccaac tcgactaaaa ttaacttctt ccataacggc gtgggagaag 1800
 cgacattcga agatgtgacg gtctttgaag gactgtatga tgcttgcca caaaggaag 1859

<210> 8

<211> 22

<212> DNA

<213> Artificial Sequence

<400> 8

caactgctgg catcctcagt ga

22

20 <210> 9

<211> 29

<212> DNA

<213> Artificial Sequence

<400> 9

gcggatccat gaagctatca aatgcaatc

29

<210> 10

<211> 26

<212> DNA

30 <213> Artificial Sequence

24

<400> 10

gcggatcctt accgagccca agtgcc

26

<210> 11

<211> 27

<212> DNA

<213> Artificial Sequence

<400> 11

gcggatcctc acttcctttg tggccag

27

10

<210> 12

<211> 30

<212> DNA

<213> Artificial Sequence

<400> 12

gttggcgtac caagaccgag cccaagtgcc

30

<210> 13

<211> 1894

20 <212> DNA

<213> Scopulariopsis brevicaulis IFO4843

<220>

<221> CDS

<222> (1)...(1721)

<221> intron

<222> (1722)...(1776)

<221> CDS

<222> (1777)...(1894)

<400> 13

30 caacctacgt ctctgtcaat cgacaattcc acgtatcctt ctatcgacta caactccgcc 60

cctccaaacc tctcgactct tgccaacaac agcctcttcg agacatggag gccgagggca 120

cacgtccttc cgccccagaa ccagatcggc gatccgtgta tgcactacac cgacccccgag 180
 acaggaatct tccacgtcgg ctggctgtac aacggcaatg gcgcttccgg cgccacgacc 240
 gaggatctcg tcacctatca ggatctcaac cccgacggag cgcagatgat ccttccgggt 300
 ggtgtgaatg accccattgc tgtctttgac ggcgcgggta ttcccagtgg cattgatggg 360
 aaaccaccca tgatgtatac ctcggtgtca tacatgcca tctcctggag catcgcttac 420
 accaggggaa gcgagacca ctctctcgca gtgtcgtccg acggcggtaa gaacttcacc 480
 aagctgggtgc agggccccgt cattccttcg cctcccttcg gcgccaacgt gaccagctgg 540
 cgtgacccct tcctgttcca aaacccccag ttcgactctc tcctcgaaag cgagaacggc 600
 10 acgtggtaca ccgttatctc tgggtggcatc cacggtgacg gccctccgc gttcctctac 660
 cgtcagcacg accccgactt ccagtactgg gagtaccttg gaccgtggtg gaacgaggaa 720
 gggaactcga cctggggcag cggtgactgg gctggccggg ggggctacaa cttcgaggtc 780
 atcaacattg tcggtcttga cgatgatggc tacaaccccg acggtgaaat ctttgccacg 840
 gtaggtaccg aatggtcgtt tgaccccatc aaaccgcagg cctcggacaa caggagatg 900
 ctctgggccg cgggcaacat gactctcgag gacggcgata tcaagttcac gccaagcatg 960
 gcgggctacc tcgactgggg tctatcggcg tatgcccggc ctggcaagga gctgcccgt 1020
 tcttcaaagc cttcgcagaa gagcgggtgcg ccggaccggg tcgtgtcgta cctgtggctc 1080
 accggtgact acttcgaggg ccacgacttc cccaccccgc agcagaattg gaccggctcg 1140
 cttttgcttc cgcgtgagct gagcgtcggg acgattcca acgttgtcga caacgagctt 1200
 20 gctcgcgaga cgggctcttg gagggttggc accaacgaca ctggcgtgct tgagctggtc 1260
 actctgaagc aggagattgc tcgagagacg ctggctgaaa tgaccagcgg caactccttc 1320
 accgaggcga gcaggaatgt cagctcgccc ggatctaccg ccttcacgca gtccctggat 1380
 tccaagttct tcgtcctgac cgctcgtctc tccttcctt cgtcggctcg cgactccgac 1440
 ctcaaggctg gtttcgagat cctgtcgtcc gagtttgagt cgaccacggg ctactaccag 1500
 ttttccaacg agtccatcat cattgaccgg agcaactcga gtgctgccgc cttgactacc 1560
 gatggaatcg acacccgcaa cgagtttggc aagatgcgcc tgtttgatgt tgtcgagggt 1620
 gaccaggagc gtatcgagac gctcgatctc actattgtgg ttgataactc gatcgttgag 1680
 gttcatgcca acgggcgatt cgctctgagc acttgggttc ggtaagtggg acgccgacca 1740
 cgcttcttaa tttctccaat actaacatta tcacagttcg tggtagagt cgtccaagga 1800
 30 catcaagttc ttccacgatg gcgacagcac gggttcagttc tcgaacatca ccgtctacga 1860
 gggactgttt gacgcctggc cggagcgggc cagg 1894

26

<210> 14

<211> 26

<212> DNA

<213> Artificial Sequence

<400> 14

gcggatccat gaaactctca actggt

26

<210> 15

10 <211> 26

<212> DNA

<213> Artificial Sequence

<400> 15

gcggatcctt accgaaccga agtgct

26

<210> 16

<211> 25

<212> DNA

<213> Artificial Sequence

20 <400> 16

gcggatcctt acctggcccg ctccg

25

<210> 17

<211> 33

<212> DNA

<213> Artificial Sequence

<400> 17

cgactcgtac cacgaacgaa cccaagtgt cag

33

CLAIMS:

1. A polypeptide comprising the amino acid sequence of SEQ ID No. 1.
2. A DNA encoding the polypeptide according to
5 claim 1.
3. The DNA according to claim 2 comprising the nucleotide sequence of SEQ ID No. 2.
4. A polypeptide comprising the amino acid sequence of SEQ ID No. 3.
- 10 5. A DNA encoding the polypeptide according to claim 4.
6. The DNA according to claim 5 comprising the nucleotide sequence of SEQ ID No. 4.
7. A vector comprising the DNA according to
15 claim 2, 3, 5 or 6.
8. An isolated host cell transformed by the vector according to claim 7.
9. The vector according to claim 7, which is an expression vector comprising, in addition to the DNA,
20 nucleotide sequences necessary to express the DNA.
10. An isolated host cell transformed by the vector according to claim 9.
11. A process for producing β -fructofuranosidase comprising the steps of:
25 cultivating the host cell according to claim 10,
and

collecting the β -fructofuranosidase from the host cell, a culture thereof or both the host cell and the culture.

12. A process for producing fructooligosaccharides
5 comprising the step of bringing sucrose into contact with the host cell according to claim 10 or a β -fructofuranosidase obtained by the process according to claim 11.

FETHERSTONHAUGH & CO.
OTTAWA, CANADA

PATENT AGENTS

1 / 6




FIG. 1A




FIG. 1B




FIG. 1C




FIG. 1D




FIG. 2 A




FIG. 2B