




(86) **Date de dépôt PCT/PCT Filing Date:** 2008/07/25
(87) **Date publication PCT/PCT Publication Date:** 2009/01/29
(45) **Date de délivrance/Issue Date:** 2016/07/05
(85) **Entrée phase nationale/National Entry:** 2010/01/22
(86) **N° demande PCT/PCT Application No.:** US 2008/071256
(87) **N° publication PCT/PCT Publication No.:** 2009/015368
(30) **Priorités/Priorities:** 2007/07/25 (US60/951,906);
2007/07/25 (US60/951,901); 2008/02/15 (US61/029,196)

(51) **Cl.Int./Int.Cl.** **C07D 313/00** (2006.01),
A61K 31/365 (2006.01), **A61P 35/00** (2006.01),
A61P 35/02 (2006.01), **A61P 35/04** (2006.01),
C07D 401/12 (2006.01), **C07D 405/10** (2006.01),
C07D 413/10 (2006.01), **C07D 493/04** (2006.01),
C07F 9/655 (2006.01)

(72) **Inventeurs/Inventors:**
AGOULNIK, SERGEI, US;
DECOSTA, BRUCE, US;
DU, HONG, US;
JIANG, YIMIN, US;
LI, XIANG-YI, US;
...

(73) **Propriétaire/Owner:**

(54) **Titre : INHIBITEURS MULTIKINASE A UTILISER DANS LE TRAITEMENT DU CANCER**
(54) **Title: MULTIKINASE INHIBITORS FOR USE IN THE TREATMENT OF CANCER**



(57) **Abrégé/Abstract:**

The present invention provides compounds, pharmaceutical compositions and methods for the treatment of specific cancers. Such compositions may generally comprise a compound of formula (I):
..... wherein R₁-R₃ are as defined herein, or pharmaceutically acceptable salts or esters thereof; and a pharmaceutically acceptable carrier.



(72) **Inventeurs(suite)/Inventors(continued):** NOMOTO, KENICHI, US; WANG, JOHN (YUAN), US; ZHANG, HUIMING, US

(73) **Propriétaires(suite)/Owners(continued):**EISAI R & D MANAGEMENT CO., LTD., JP

(74) **Agent:** BORDEN LADNER GERVAIS LLP

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 January 2009 (29.01.2009)

PCT

(10) International Publication Number
WO 2009/015368 A2

(51) International Patent Classification:

A61K 31/365 (2006.01) A61P 35/02 (2006.01)
C07D 313/00 (2006.01) A61P 35/04 (2006.01)
A61P 35/00 (2006.01)

Kenichi [JP/US]; 72 Washington Street, Belmont, MA 02478 (US). WANG, John (Yuan) [US/US]; 344 River Road, Andover, MA 01810 (US). ZHANG, Huiming [CN/US]; 2 Dorset Circle, Andover, MA 01810 (US).

(21) International Application Number:

PCT/US2008/071256

(74) Agents: HANLEY, Elizabeth, A. et al.; Lahive & Cockfield, LLP, One Post Office Square, Boston, MA 02109-2127 (US).

(22) International Filing Date: 25 July 2008 (25.07.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/951,901 25 July 2007 (25.07.2007) US
60/951,906 25 July 2007 (25.07.2007) US
61/029,196 15 February 2008 (15.02.2008) US

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(71) Applicant (for all designated States except US): EISAI R & D MANAGEMENT CO., LTD. [JP/JP]; 6-10, Koshikawa 4-chome, Tokyo, Bunkyo-ku (JP).

(72) Inventors; and


(75) Inventors/Applicants (for US only): AGOULNIK, Sergei [US/US]; 91 Marion Street, Wilmington, MA 01887 (US). DECOSTA, Bruce [GB/US]; 99 Cluff Crossing Road, Apt. C-8, Salem, NH 03079 (US). DU, Hong [CN/US]; 43 Upland Street, North Andover, MA 01845 (US). JIANG, Yimin [US/US]; 16 Devonshire Lane, Londonderry, NH 03053 (US). LI, Xiang-Yi [US/US]; 229 High Plain Road, Andover, MA 01810 (US). NOMOTO,

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

(54) Title: MULTIKINASE INHIBITORS FOR USE IN THE TREATMENT OF CANCER



(I)

(57) Abstract:


The present invention provides compounds, pharmaceutical compositions and methods for treatment of specific cancers. Such compositions may generally comprise a compound of formula

(I):..... wherein R1-R3 are as defined herein, or pharmaceutically acceptable salts or esters thereof; and a pharmaceutically acceptable carrier.

WO 2009/015368 A2

MULTIKINASE INHIBITORS FOR USE IN THE TREATMENT OF CANCER**BACKGROUND OF THE INVENTION**

F152 (LL-Z1640-2), a zearalenone-like macrolide, was first isolated from shake flask fermentation, crude extracts of which inhibited the ciliated protozoan *Tetrahymena pyriformis* (see, McGahren *et al. J. Org. Chem.* **1978**, *43*, 2339). Although initial biological



F152

studies using this natural product failed to yield any particularly interesting activities, the possibility of preparing additional derivatives and/or further exploring their biological activity was undertaken. For example, F152 and certain isomers thereof inhibit the phosphorylating enzyme Map/Erk kinase (MEK) and may be useful for the treatment of certain MEK-related cancers and other diseases characterized by the formation of neoangiogenesis (see, *e.g.*, GB 323 845). Derivatives of F152 have also been shown to have activity as tyrosine kinase inhibitors, which are useful, for example, for the treatment of cancer and inflammatory disorders (see, *e.g.*, EP 606 044; WO 00/38674; JP 8-40893; WO 96/13259; 5,728,726; 5,674,892; 5,795,910). Often, however, F152 and derivatives thereof are obtained by fermentation techniques and modifications to the natural product and thus were limited in the number and types of derivatives that could be prepared and evaluated for biological activity. Additionally, although F152 and certain derivatives thereof have demonstrated potent *in vitro* activities, these compounds may be biologically unstable (for example, they are susceptible to enone isomerization in mouse and human plasma), thereby limiting the development of these compounds as therapeutics for the treatment of humans or other animals.

Recently, F152 analogues have been shown to be inhibitors of NF- κ B activation, and MEK1 (See, *e.g.*, U.S. Application Serial No.: 10/507,067 and U.S. Application Publication No.: US 2004/0224936). The compounds were also reported to inhibit AP-1 activation and some protein kinases (for example, MEKK, PDGFr, VEGFr). Based on these mechanisms of action, it was suggested that the compounds inhibit the production of various pro-inflammatory and/or immunologic cytokines such as TNF α , IL-1, IL-6, IL-8, IL-2 etc, and also inhibit the production of various pro-inflammatory molecules under the regulation of NF- κ B pathway such as prostaglandins produced from COX-2, ICAM-1 and MMP-1 and 3 etc. Also, it was reported that the compounds have the ability to inhibit cell proliferation under the regulation of AP-1 pathway through the inhibition of MEK1. In addition, it was reported that the compounds have ability to inhibit angiogenesis, likely based on the inhibitory activity on VEGFr kinase and weak inhibitory activity on PDGFr kinase.

SUMMARY OF THE INVENTION

The present invention is based, at least in part, on the discovery that the compounds of formula (I) or (II) have unique multikinase inhibition profiles. Accordingly, in some aspects, the present invention provides compounds having unique multikinase inhibition profiles. In some aspects, the present invention provides compounds useful as SrcTK/MEK inhibitors. Such compounds may be useful against specific cancers selected based upon the unique multikinase inhibition profile of the compounds. Generally, chemotherapeutic drugs often lack of specificity and have unwanted side effects associated therewith. Accordingly, it is an object of the present invention to provide cancer-specific agents for targeted cancer therapy and targeted methods for treating and/or preventing cancer which may circumvent adverse side effects often associated with chemotherapeutics.


Accordingly, in some aspects, the present invention provides methods for treating a SrcTK/MEK associated cancer in a subject in need thereof. The methods

generally include administering to the subject a composition comprising a SrcTK/MEK inhibitor in an amount effective to treat the SrcTK/MEK associated cancer. In some aspects, the present invention also relates to the use of a SrcTK/MEK inhibitor in the manufacture of a medicament for the treatment of a SrcTK/MEK associated cancer.

In some embodiments, the SrcTK/MEK inhibitor inhibits the activity of at least one member of the Src tyrosine kinase family by at least about 75% at a concentration of about 0.1 μ M. The members of the Src tyrosine kinase family can be any member of the Src family including cSrc, Fyn, Lyn, Lck, Yes, Fgr, Hck and Blk. In some embodiments, the SrcTK/MEK inhibitor inhibits the activity of at least five members of

the Src tyrosine kinase family by at least about 50% at a concentration of about 0.1 μ M. In some embodiments, the SrcTK/MEK inhibitor inhibits the activity of MEK1 by at least about 50% at a concentration of about 0.1 μ M.


In some embodiments, the SrcTK/MEK inhibitor is a compound of formula (I):



wherein:

R_1 is H

R_2 is a moiety selected from the group consisting of H and trifluoromethylcarbonyl; or R_1 and R_2 are taken together with the core structure to form a heterocyclodiyl moiety of formula (a):




R_3 is a moiety selected from the group consisting of $-OR_a$ and $-NR_bR_c$;

R_a is a C_{1-4} alkyl group optionally substituted with an imidazolyl;

R_b is a moiety selected from the group consisting of a C_{1-4} alkyl group and a C_{1-4} alkoxy group, wherein R_b is substituted with 0, 1 or 2 groups each independently selected from the group consisting of $-OCH_3$, $-C(O)OH$, $-C(O)NR'R''$, $-NH(C_{1-3} \text{ alkyl})$, $-NH(CH_2CH_2O)_nCH_3$, wherein n is 2-4, piperazinyl, N-methylpiperazinyl, piperidinyl, N-methylpiperidinyl, N-morpholinyl, imidazolyl, pyrrolidinyl, $-OPO_3H_2$ and hydroxyl; wherein the $-NH(C_{1-3} \text{ alkyl})$ group is substituted with 0, 1 or 2 hydroxyl moieties and wherein R' and R'' are each independently selected from $-H$ or $-CH_3$; and

R_c is H, or R_b and R_c are taken together with the nitrogen to which they are attached to form a heterocyclyl moiety selected from the group consisting of

formula (b) and formula (c):




or a pharmaceutically acceptable salt or ester thereof.

In some embodiments, the SrcTK/MEK associated cancer is chronic myeloid leukemia or colorectal cancer. In some embodiments, the SrcTK/MEK inhibitor also inhibits the activity of Bcr-Abl, and the SrcTK/MEK associated cancer is leukemia. In some embodiments, the SrcTK/MEK inhibitor also inhibits the activity of TrkB. In some embodiments, the SrcTK/MEK inhibitor also inhibits the activity of TrkB at least to the extent described further herein, *e.g.*, by at least about 75% at a concentration of about 0.1 μ M.

In some aspects, the present invention is directed to methods for treating a TrkB/MEK associated cancer in a subject in need thereof. The methods generally include administering to the subject a composition comprising a TrkB/MEK inhibitor in an amount effective to treat the TrkB/MEK associated cancer. In some aspects, the present invention also relates to the use of a TrkB/MEK inhibitor in the manufacture of a medicament for the treatment of a TrkB/MEK associated cancer.

In some embodiments, the TrkB/MEK inhibitor inhibits the activity of TrkB by at least about 75% at a concentration of about 0.1 μ M. In some embodiments, the TrkB/MEK inhibitor inhibits the activity of MEK1 by at least about 50% at a concentration of about 0.1 μ M.


In some embodiments, the TrkB/MEK inhibitor is a compound of formula (I):



wherein:

R_1 is H

R_2 is a moiety selected from the group consisting of H and trifluoromethylcarbonyl; or R_1 and R_2 are taken together with the core structure to form a heterocyclodiyl moiety of formula (a):




R_3 is a moiety selected from the group consisting of $-OR_a$ and $-NR_bR_c$;

R_a is a C_{1-4} alkyl group optionally substituted with an imidazolyl;


R_b is a moiety selected from the group consisting of a C_{1-4} alkyl group and a C_{1-4} alkoxy group, wherein R_b is substituted with 0, 1 or 2 groups each independently selected from the group consisting of $-OCH_3$, $-C(O)OH$, $-C(O)NR'R''$, $-NH(C_{1-3} \text{ alkyl})$, $-NH(CH_2CH_2O)_nCH_3$, wherein n is 2-4, piperazinyl, N-methylpiperazinyl, piperidinyl, N-methylpiperidinyl, N-morpholinyl, imidazolyl, pyrrolidinyl, $-OPO_3H_2$ and hydroxyl; wherein the $-NH(C_{1-3} \text{ alkyl})$ group is substituted with 0, 1 or 2 hydroxyl moieties and wherein R' and R'' are each independently selected from $-H$ or $-CH_3$; and

R_c is H, or R_b and R_c are taken together with the nitrogen to which they are attached to form a heterocyclyl moiety selected from the group consisting of formula (b) and formula (c):



or a pharmaceutically acceptable salt or ester thereof.

In one embodiment, compound (I) is a compound of formula (II):



wherein

R_3 is $-NHR_b$, and R_b is C_1 - C_3 alkyl substituted with 0, 1, or 2 hydroxyl moieties; or a pharmaceutically acceptable salt or ester thereof.

Reference herein to compounds of formula (I) or (II) (or compositions comprising the same) includes pharmaceutically acceptable salts and esters of the compounds (or the compositions comprising the same).

In some embodiments, the TrkB/MEK associated cancer is pancreatic cancer, neural cancer, glioma, neuroblastoma or retinoblastoma. In some embodiments, the TrkB/MEK inhibitor also inhibits the activity of at least one member of the Src tyrosine kinase family. In some embodiments, the TrkB/MEK inhibitor also inhibits the activity of at least one member of the Src tyrosine kinase family at least to the extent described further herein, *e.g.*, by at least about 75% at a concentration of about $0.1\mu M$.

In some aspects, the present invention provides methods for treating at least one cancer selected from the group consisting of chronic myeloid leukemia, pancreatic cancer, neural cancer, glioma, neuroblastoma, retinoblastoma, ovarian cancer, thyroid cancer, colorectal cancer and melanoma, in a subject in need thereof. The methods generally include administering to the subject a composition comprising a compound of formula (I), (II), or a pharmaceutically acceptable salt or ester thereof, in an amount effective for treating the cancer. The compound of formula (I) or (II) can be any of the compounds of formula (I) or (II) as described herein.


In some aspects, the present invention also relates to the use of a compound of formula (I) or (II) in the manufacture of a medicament for the treatment of chronic myeloid leukemia, pancreatic cancer, neural cancer, glioma, neuroblastoma, retinoblastoma, ovarian cancer, thyroid cancer, colorectal cancer or melanoma. In some aspects, the present invention provides methods for inhibition of a metastatic process in


a subject in need thereof. The methods generally include administering to the subject a composition comprising a compound of formula (I) or (II) or a pharmaceutically acceptable salt or ester thereof, in an amount effective for inhibiting the metastatic process. The compound of formula (I) or (II) can be any of the compounds of formula (I) or (II) as described herein.

In other aspects, the present invention provides methods for treating a B-RAF mutated cancer in a subject in need thereof. The methods generally include administering to the subject a composition comprising at least one compound of formula (I) or (II) or a pharmaceutically acceptable salt or ester thereof; in an amount effective for treating the B-RAF mutated cancer. The compound of formula (I) or (II) can be any of the compounds of formula (I) or (II) as described herein. In some embodiments, the B-RAF mutated cancer is a B-RAF V600E mutated cancer. In some embodiments, the B-RAF mutated cancer is ovarian cancer, thyroid cancer, colorectal cancer or melanoma.

In other aspects, the present invention provides methods for treating a FLT3 mutated cancer in a subject in need thereof. For example, the method can comprise administration of a compound of formula (I) or (II) or a pharmaceutically acceptable salt or ester thereof (or a composition comprising same) to the subject in an amount effective for treating the FLT3 mutated cancer. In certain embodiments of the method, the FLT3 mutated cancer carries a mutation of residue D835 and/or a mutation of residue I836, for example, a D835Y mutation. In some embodiments, the FLT3 mutated cancer is acute myeloid leukemia (AML).


In other aspects, the present invention provides methods for treating a central nervous system tumor in a subject in need thereof. The methods generally include administering to the subject a composition comprising a compound of formula (I) or (II) or a pharmaceutically acceptable salt or ester thereof; an amount effective for treating the tumor, such that at least 20% of the compound of formula (I) or (II) penetrates the blood brain barrier. The compound of formula (I) or (II) can be any of the compounds of formula (I) or (II) as described herein. In some embodiments, at least about 65% of the compound of formula (I) or (II) penetrates the blood brain barrier, based upon a ratio of amount of a compound of formula (I) or (II) in the brain to amount of a compound of formula (I) or (II) in the plasma. In some embodiments, the central nervous system tumor is a brain tumor, a glioma or a neuroblastoma.

In some embodiments,  in formula (I) represents a double bond. In

some embodiments,  represents a triple bond. In some embodiments, R_3 is -

NR_bR_c , R_c is H


and R_b is an unsubstituted C_{1-4} alkyl. In some embodiments, R_c is H and R_b is methyl or ethyl. For example, compounds of formula (I) include the following compounds:



























and pharmaceutically acceptable salts and esters thereof.


In some embodiments, any of the methods of the present invention include intravenous administration of the compositions described herein. In some embodiments, any of the methods of the present invention include administration of a dosage between about 0.10 mg/kg to about 25 mg/kg of body weight of the compositions described herein.

In some aspects, the present invention is directed to the following compounds:




 (Compound 014)	 (Compound 054)
 (Compound 015)	 (Compound 065)
 (Compound 016)	 (Compound 076)
 (Compound 018)	 (Compound 114)
 (Compound 019)	 (Compound 122)
 (Compound 022)	 (Compound 127)

 (Compound 024)	 (Compound 137)
 (Compound 025)	 (Compound 144)
 (Compound 034)	 (Compound 155)
 (Compound 041)	 (Compound 156)
 (Compound 045)	 (Compound 157)
 (Compound 046)	 (Compound 158)



and pharmaceutically acceptable salts and esters thereof.

With regard to compounds of formula (II), in some embodiments, R₃ is an unsubstituted C₁₋₃ alkylamino. In some embodiments, R₃ is a group selected from the group consisting of methylamino and ethylamino. In some embodiments, R₃ is a group selected from the group consisting of 2-hydroxyethylamino and 2,3-dihydroxypropylamino. For example, compounds of formula (II) include, but are not limited to, the following compounds:



and pharmaceutically acceptable salts and esters thereof.

In some embodiments, any of the methods of the present invention include administration of a second chemotherapeutic drug.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts a schematic representation of the RAS-MAPK signaling pathway.

Figure 2A describes results of exemplary compounds of formula (I) or (II) in biochemical and cell based MEK1 assays.

Figure 2B describes exemplary kinase inhibition results of a compound .

Figure 3 describes results of exemplary compounds of formula (I) or (II) in cell growth inhibition assays in a variety of cancer cell lines.

Figure 4 includes graphs depicting results from *in vivo* Antitumor Activity of exemplary compounds of formula (I) or (II) in Colo-205 Human Colon Cancer Xenografts.

Figure 5 includes graphs depicting results from *in vivo* Antitumor Activity of exemplary compounds of formula (I) or (II) in BxPC-3 Human Pancreatic Cancer Xenografts.

Figure 6 includes graphs depicting results from *in vivo* Antitumor Activity of exemplary compounds of formula (I) or (II) in LOX Human Melanoma Xenografts.

Figure 7 describes effects of a compound on B-cell driven hematological malignant cancer cell growth.

Figure 8 describes effects of a compound on the anticancer activity of CPT-11 (SN-38) in PANC-1 Pancreatic Cancer Cells.

Figure 9 describes blood brain barrier (BBB) experiments of a compound.

Figure 10 describes effects of a compound on the phosphorylation of Flt-3 receptor in MV-4-11 AML cells.

Figure 11 is a graph which depicts inhibition of MV-4-11 cell proliferation by a compound.

DETAILED DESCRIPTION OF THE INVENTION

As discussed in WO 05/023792 and WO 03/076424, certain analogs of F152 have been reported to have inhibitory activity against MEK1, have demonstrated increased stability and are potent inhibitors of NF- κ B activation, AP-1 activation and protein kinases (for example, MEKK, MEK1, PDGFr, VEGFr). The present invention is based, at least in part, on the discovery that certain F152 macrolide analogs also have unique multikinase inhibition profiles. That is, compounds of formula (I) or (II) are effective as inhibitors of multiple, specific kinases as discussed in further detail herein. Certain multikinases have previously been shown to be effective against cancer. It has recently been discovered, for example, that imatinib mesylate (STI571, GLEEVEC), an inhibitor of the Bcr-Abl kinase, has been successful in the treatment of chronic myeloid leukemia. This success is attributed, at least in part, to the multikinase activity of imatinib mesylate. Accordingly, without wishing to be bound to any particular theory, it

is believed that compounds with these unique multikinase inhibition profiles, *e.g.*, F152 macrolide analogs of formula (I) or (II), have a broader therapeutic spectrum than selective MEK1 inhibitors. For example, it is believed that compounds having this unique multikinase inhibition profile can be used in the treatment and/or prevention of specific cancers.

Definitions

In order to more clearly and concisely describe the subject matter of the claims, the following definitions are intended to provide guidance as to the meaning of specific terms used herein.

It is to be noted that the singular forms “a,” “an,” and “the” as used herein include “at least one” and “one or more” unless stated otherwise. Thus, for example, reference to “a pharmacologically acceptable carrier” includes mixtures of two or more carriers as well as a single carrier, and the like.

As used herein, the term “SrcTK/MEK inhibitor” refers to a single compound that inhibits both MEK1 and at least one member of the Src tyrosine kinase family. Exemplary members of the Src tyrosine kinase family include cSrc, Fyn, Lyn, Lck, Yes, Fgr, Hck and Blk. In some embodiments, the term “SrcTK/MEK inhibitor” refers to a single compound that inhibits both MEK1 and from 1 to 5 members of the Src tyrosine kinase family, *e.g.*, 1, 2, 3, 4, or 5 members of the Src tyrosine kinase family.

As used herein, the term “TrkB/MEK inhibitor” refers to a single compound that inhibits both MEK1 and TrkB.

When used herein, a specific protein kinase is referred to by a specific name, *e.g.*, Src, TrkB, and other kinases mentioned specifically herein, but may have a number of synonyms known in the art. For example, Src protein kinase may also be referred to as ASV, c-Src, p60-Src, pp60c-src, Proto-oncogene tyrosine-protein kinase Src, SRC1, AW259666, Neuronal proto-oncogene tyrosine-protein kinase, fc54g04, wu:fc54g0 and/or SDR. Additionally, TrkB can also be referred to as BDNF/NT-3 growth factors receptor precursor, GP145-TrkB, Neurotrophic tyrosine kinase receptor type 2, Trk-B, TrkB tyrosine kinase, AI848316, C030027L06Rik, GP145-TrkB/GP95-TrkB and/or Tkrb. Yes may also be referred to as c-Yes, HsT441, p61-Yes, Proto-oncogene tyrosine-protein kinase Yes, Yes, AI323763, p61-Yes, Tyrosine-protein kinase Yes, MGC94936 and/or p60c-yes. Lck may also be referred to as LSK, Lymphocyte cell-specific protein-

tyrosine kinase, p56-LCK, Proto-oncogene tyrosine-protein kinase LCK, T cell-specific protein-tyrosine kinase, Hck-3 and/or Lsk-t. Fyn may also be referred to as MGC45350, p59-Fyn, Protooncogene Syn, Proto-oncogene tyrosine-protein kinase Fyn, SLK, SYN, AI448320, AW552119, MGC115870, MGC132140, MGC52878, Src Kinase p59, c-fyn and/or zgc:86720. Lyn may also be referred to as FLJ26625, JTK8, Tyrosine-protein kinase Lyn, AA407514, Hck-2, Lyncein and/or TBC1 domain family member 1. It is well within the capability of the skilled artisan to determine whether a name is a synonym for the protein kinases recited herein using, *e.g.*, searchable protein databases.

When used herein, a specific protein kinase includes not only the naturally occurring kinase, *e.g.*, Src, TrkB, and other kinases mentioned specifically herein, but also modified kinases. Accordingly, in some embodiments, the compounds of formula (I) or (II) inhibit kinases which share at least 90% sequence identity with naturally occurring counterparts. In other embodiments, the compounds of formula (I) or (II) inhibit kinases which share at least 95% sequence identity with naturally occurring counterparts. In still other embodiments, the compounds of formula (I) or (II) inhibit kinases which share at least 97% sequence identity with naturally occurring counterparts. In yet other embodiments, the compounds of formula (I) or (II) inhibit kinases which share at least 99% sequence identity with naturally occurring counterparts. Kinases which share sequence identity with naturally occurring counterparts can be naturally occurring or synthetic. The term "sequence identity" generally refers to the degree to which two polynucleotide or polypeptide sequences are identical on a residue-by-residue basis over a particular region of comparison. The term "percentage of sequence identity" is calculated by comparing two optimally aligned sequences over that region of comparison, determining the number of positions at which the identical nucleic acid base (*e.g.*, A, T, C, G, U, or I, in the case of nucleic acids) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the region of comparison (*i.e.*, the window size), and multiplying the result by 100 to yield the percentage of sequence identity. In some embodiments, the window size is at least 20 nucleotide positions, for example, at least 25-50 nucleotides. In other embodiments, the window size is greater than 50 nucleotides. In some embodiments, the window size is the entire naturally occurring protein kinase. In other embodiments, the window size is the sequence which corresponds at least to the ATP binding site on the naturally occurring protein kinase.

As used herein, the term “SrcTK/MEK associated cancer” refers to cancers that are associated with or otherwise affected by the activity or function of MEK1 and at least one member of the Src tyrosine kinase family. In one aspect, the activity or function of MEK1 is elevated, and the activity or function of at least one member of the Src tyrosine kinase family is elevated in a SrcTK/MEK associated cancer. Exemplary SrcTK/MEK associated cancers are discussed in more detail herein, and may include chronic myeloid leukemia (CML) and colorectal cancer.

As used herein, the term “TrkB/MEK associated cancer” refers to cancers that are associated with or otherwise affected by the activity or function of MEK1 and TrkB. In one aspect, the activity or function of MEK1 is elevated, and the activity or function of TrkB is elevated in a TrkB/MEK associated cancer. Exemplary TrkB/MEK associated cancers are also discussed in more detail herein, and may include pancreatic cancer, neural cancer, glioma, neuroblastoma and retinoblastoma.

In some embodiments, cancers that are “associated with or otherwise affected by the activity or function” of one or more protein kinases are cancers which are attributable to dysregulation of protein kinase activity. In other embodiments, cancers that are “associated with or otherwise affected by the activity or function” of one or more protein kinases are cancers which are attributable to dysregulation of an upstream or downstream event in a protein kinase pathway. In other embodiments, cancers that are “associated with or otherwise affected by the activity or function” of one or more protein kinases are cancers which are attributable to mutations in genes which regulate protein kinase activity. In some embodiments, cancers that are “associated with or otherwise affected by the activity or function” of one or more protein kinases are cancers which are attributable to increased or constitutive activation of one or more signal molecules. In some embodiments, cancers that are “associated with or otherwise affected by the activity or function” of one or more protein kinases are cancers which are attributable to the increased or constitutive ability of the protein kinase to phosphorylate proteins. In still other embodiments, cancers that are “associated with or otherwise affected by the activity or function” of one or more protein kinases are cancers which are attributable to alteration of binding of adenosine triphosphate (ATP) to the enzyme's binding site such that kinase activity is increased. In some embodiments, cancers that are “associated with or otherwise affected by the activity or function” of one or more

protein kinases are cancers which are attributable to ligand binding to and/or dimerization of the protein kinase.

Cancers designated “SrcTK/MEK associated,” “TrkB/MEK associated,” “associated with B-RAF mutations” and/or “associated with FLT3 mutations” are cancers which have been chosen because of their apparent association with certain protein kinases or specific protein mutations, as described in more detail herein. Once a cancer (e.g., pancreatic cancer) is determined to be associated with MEK1, Src, TrkB, or any of the other kinases or protein mutations described herein, (e.g., in the literature or through experimentation), the skilled artisan would understand that treatment of the specific cancer is indicative of treatment of a cancer of the present invention (e.g., a TrkB/MEK associated cancer).

Compounds of formula (I) or (II) inhibit a number of protein kinases as described herein. As used herein, the term “inhibit” when used in relation to a protein kinase, refers to the ability of a compound of formula (I) or (II) to at least partially reduce the activity of the protein kinase. Without wishing to be bound by any particular theory, it is believed that an inhibitor of a protein kinase may either compete directly with ATP for the binding pocket of a specific kinase or otherwise interact with the kinase such that the binding pocket is no longer available to the ATP. The amount to which a compound will inhibit a protein kinase is ascertainable by a number of assays known in the art. For example, a compound may be diluted in an acceptable buffer (e.g., MOPS, EDTA, Brij™-35, Glycerol, NaCl, BSA, HEPES, Triton™ X-100, or TRIS), and exposed to the protein kinase in the presence of a specific amount of [γ - ^{33}P -ATP] and a substrate peptide. Following incubation and quenching, an aliquot will be taken to determine the amount of ^{33}P incorporated. For example, ionizing radiation will be measured with a scintillation counter, to determine the amount of γ - ^{33}P incorporated into the substrate which is associated with the kinase activity. This value would be compared to a suitable control, e.g., the same process without the inhibitor present. It will be appreciated that ability of a compound to inhibit kinase activity can be assessed or measured by any suitable method. For example, antibodies which bind phospho-peptide or phospho-protein substrates can be used to detect such phosphorylation products, which are indicative of kinase activity. Binding of antibody to substrate can be detected by any suitable method, such as labeled second antibody. In some embodiments, the ability of a compound to inhibit transphosphorylation of receptor tyrosine kinase

subunits or to inhibit autophosphorylation of receptor tyrosine kinase can be assessed or measured. In some embodiments, the inhibitors of the present invention inhibit at least about 30% of the activity of the specified protein kinase at an inhibitor concentration of about 0.1 μ M. In some embodiments, the inhibitors of the present invention inhibit at least about 45% of the activity of the specified protein kinase at an inhibitor concentration of about 0.1 μ M. In some embodiments, the inhibitors of the present invention inhibit at least about 50% of the activity of the specified protein kinase at an inhibitor concentration of about 0.1 μ M. In some embodiments, the inhibitors of the present invention inhibit at least about 55% of the activity of the specified protein kinase at an inhibitor concentration of about 0.1 μ M. In some embodiments, the inhibitors of the present invention inhibit at least about 40% of the activity of the specified protein kinase at an inhibitor concentration of about 1.0 μ M. In some embodiments, the inhibitors of the present invention inhibit at least about 50%, of the activity of the specified protein kinase at an inhibitor concentration of about 1.0 μ M. In some embodiments, the inhibitors of the present invention inhibit at least about 75%, 80%, 85%, or 90% of the activity of the specified protein kinase at an inhibitor concentration of about 1.0 μ M. In some embodiments, the inhibitors of the present invention inhibit about 95% of the activity of the specified protein kinase at an inhibitor concentration of about 1.0 μ M. In some embodiments, the assay to determine amount of inhibition is performed in the presence of ATP. In some embodiments, the assay to determine amount of inhibition is performed in the presence of 10 μ M ATP. In some embodiments, the assay to determine amount of inhibition is performed in the presence of $[\gamma\text{-}^{33}\text{P}\text{-ATP}]$. In some embodiments, the assay to determine amount of inhibition is performed in the presence of an amount of $[\gamma\text{-}^{33}\text{P}\text{-ATP}]$ which affords a specific activity of about 500 cpm/pmol. In some embodiments, the assay to determine amount of inhibition is performed in the presence of trace amounts of $[\gamma\text{-}^{33}\text{P}\text{-ATP}]$.

As used herein, the term “inhibition of a metastatic process” refers to the ability of a compound of formula (I) or (II) to inhibit not only the growth of a single tumor, but also the spread of tumors from a primary site to distant organs. Without wishing to be bound by any particular theory, it is believed that metastasis is a common cause of failure of cancer treatment. In one embodiment, inhibition of a metastatic process includes inhibiting the ability of the tumor cells to separate from its neighboring cells. In one embodiment, inhibition of a metastatic process includes inhibiting the ability of

the tumor cells to pass through the extracellular matrix. In one embodiment, inhibition of a metastatic process includes inhibiting the ability of the tumor cells to secrete enzymes that break down the matrix. In one embodiment, inhibition of a metastatic process includes inhibiting the ability of the tumor cells to pass through the endothelial lining of the blood vessels. In one embodiment, inhibition of a metastatic process includes inhibiting the ability of the tumor cells to pass through the lymphatic system. In one embodiment, inhibition of a metastatic process includes inhibiting the ability of the tumor cells to survive in the circulatory system. In one embodiment, inhibition of a metastatic process includes inhibiting the ability of the tumor cells to extravasate into the surrounding tissue of the organ. It is to be understood that, when used in reference to a metastatic process, the term “inhibition” may refer to complete halting of a metastatic process in a subject or to a slowing of the metastatic process in a subject. Without wishing to be bound by any particular theory, it is believed that the ability of a compound of formula (I) or (II) to inhibit the metastatic process is due, at least in part, to the ability of the compound to inhibit a Src tyrosine kinase.

“Treatment”, or “treating” as used herein, is defined as the application or administration of a therapeutic agent (*e.g.*, specific multikinase inhibitors of the invention) to a patient, or to an isolated tissue or cell line from a patient. The patient generally has a disease or disorder, a symptom of disease or disorder or a predisposition toward a disease or disorder. The purpose of treatment is generally to cure, heal, alleviate, relieve, remedy, ameliorate, or improve such disease, disorder, symptoms or predisposition. “Treated,” as used herein, refers to the disease or disorder being cured, healed, alleviated, relieved, remedied, ameliorated, or improved. For example, methods of treatment of the instant invention provide for administration of an inhibitor as described herein, such that the progression of a specific cancer or a specific class of cancer (*e.g.*, a SrcTK/MEK associated cancer) is slowed or stopped. Methods of treatment of the instant invention further include the administration of an inhibitor, such that a specific cancer is cured.

The term “effective dose” or “effective dosage” is defined as an amount sufficient to achieve a desired effect. The term “desired effect” refers generally to any result that is anticipated by the skilled artisan when a compound or composition of formula (I) or (II) is administered to a subject. In some embodiments, the desired effect is a complete remission of the disease or disorder. In other embodiments, the desired

effect is a partial treatment of a disease or disorder. In still other embodiments, the desired effect is a full or partial treatment of the symptoms of a disease or disorder. For example, in some embodiments, the desired effect is the shrinkage of a solid tumor. In another exemplary embodiment, the desired effect is the elimination of a solid tumor. The term “therapeutically effective dose” is defined as an amount sufficient to cure or at least partially arrest the disease and its complications in a subject already suffering from the disease.

The term “subject,” as used herein, refers to animals such as mammals, including, but not limited to, humans, primates, cows, sheep, goats, horses, pigs, dogs, cats, rabbits, guinea pigs, rats, mice or other bovine, ovine, equine, canine, feline, rodent or murine species. In some embodiments, the subject is a human.

Numerous values and ranges are recited in connection with various embodiments of the present invention, *e.g.*, amount of a compound of formula (I) or (II) present in a composition. It is to be understood that all values and ranges which fall between the values and ranges listed are intended to be encompassed by the present invention unless explicitly stated otherwise. The term “about” as used herein in association with parameters, ranges and amounts, means that the parameter or amount is within $\pm 1\%$ of the stated parameter or amount.

The term “stable”, as used herein, generally refers to compounds which possess stability sufficient to allow manufacture and which maintain the integrity of the compound for a sufficient period of time to be detected. In some embodiments, stable compounds of formula (I) or (II) are those which maintain their integrity for a sufficient period of time to be useful for the purposes detailed herein. For example, a prodrug may be considered “stable” even though it is eventually metabolized in the body.


As used herein, “alkyl” groups include saturated hydrocarbons having one or more carbon atoms, including straight-chain alkyl groups (*e.g.*, methyl, ethyl, propyl, butyl, pentyl, hexyl, etc.), cyclic alkyl groups (or “cycloalkyl” or “alicyclic” or “carbocyclic” groups) (*e.g.*, cyclopropyl, cyclopentyl, cyclohexyl, etc.), branched-chain alkyl groups (isopropyl, *tert*-butyl, *sec*-butyl, isobutyl, etc.), and alkyl-substituted alkyl groups (*e.g.*, alkyl-substituted cycloalkyl groups and cycloalkyl-substituted alkyl groups). In certain embodiments, a straight-chain or branched-chain alkyl group may have 8 or fewer carbon atoms in its backbone, *e.g.*, C₁-C₈ for straight-chain or C₃-C₈ for branched-chain. In certain embodiments, a straight-chain or branched-chain alkyl group


may have 6 or fewer carbon atoms in its backbone, *e.g.*, C₁-C₆ for straight-chain or C₃-C₆ for branched-chain. In still other embodiments, an alkyl group includes about 1 to 4 carbons. In other embodiments, an alkyl group includes about 1 to 3 carbons. In yet other embodiments, an alkyl group includes about 1 or 2 carbons. The term “lower alkyl” refers to alkyl groups having from 1 to 6 carbons in the chain, and to cycloalkyl groups having from 3 to 6 carbons in the ring structure. The term “C₁-C₆” as in “C₁-C₆ alkyl” means alkyl groups containing 1 to 6 carbon atoms. The terms “alkenyl” and “alkynyl” refer to unsaturated aliphatic groups analogous to alkyls, but which contain at least one double or triple carbon-carbon bond respectively.

The term “alkoxy” as used herein means an alkyl group having an oxygen atom attached thereto. In some embodiments, alkoxy groups include groups having 1 to about 8 carbon atoms. In other embodiments, alkoxy groups include groups having 1 to about 6 carbon atoms. In still other embodiments, alkoxy groups include groups having fewer than about 4 carbon atoms. Examples of alkoxy groups include, but are not limited to, methoxy, ethoxy, isopropoxy, propoxy, butoxy, and pentoxy groups. The alkoxy groups can be straight-chain or branched.

The term “heterocyclic group” includes closed ring structures analogous to carbocyclic groups in which one or more of the carbon atoms in the ring is an element other than carbon, for example, nitrogen, sulfur, or oxygen. Heterocyclic groups may be saturated or unsaturated. Additionally, heterocyclic groups (such as pyrrolyl, pyridyl, isoquinolyl, quinolyl, purinyl, and furyl) may have aromatic character, in which case they may be referred to as “heteroaryl” or “heteroaromatic” groups. Exemplary imidazolyl, *e.g.*,

heterocyclic groups include, but are not limited to





The term “amine” or “amino,” as used herein, refers to an unsubstituted or substituted moiety of the formula $-NR^aR^b$, in which R^a and R^b are each independently hydrogen, alkyl, aryl, or heterocyclyl, or R^a and R^b , taken together with the nitrogen atom to which they are attached, form a cyclic moiety having from 3 to 8 atoms in the ring. Thus, the term amino includes cyclic amino moieties such as piperidinyl or pyrrolidinyl groups, unless otherwise stated.

Regarding connectivity, an “arylalkyl” group, for example, is an alkyl group substituted with an aryl group (*e.g.*, phenylmethyl (*i.e.*, benzyl)). An “alkylaryl” moiety is an aryl group substituted with an alkyl group (*e.g.*, *p*-methylphenyl (*i.e.*, *p*-tolyl)). Thus, the term imidazolyl-alkyl refers to an alkyl group substituted with an imidazolyl moiety.

When specified, the chemical moieties of the compounds of formula (I) or (II), including those groups discussed above, may be “substituted or unsubstituted.” In some embodiments, the term “substituted” means that the moiety has substituents placed on the moiety other than hydrogen (*i.e.*, in most cases, replacing a hydrogen), which allow the molecule to perform its intended function. It will be understood that “substitution” or “substituted with” includes the implicit proviso that such substitution is in accordance with the permitted valence of the substituted atom and the substituent, and that the substitution results in a stable compound, *e.g.*, which does not spontaneously undergo transformation such as by rearrangement, cyclization, elimination, etc. As used herein,

the term “substituted” is meant to include all permissible substituents of organic compounds. In a broad aspect, permissible substituents include acyclic and cyclic, branched and unbranched, carbocyclic and heterocyclic, aromatic and nonaromatic substituents of organic compounds. The compounds of formula (I) or (II) may have one or more substitutions, as described herein.

As used in the description and drawings herein, an optional double/triple bond is represented by two solid lines together with a third dashed line, and refers to a covalent linkage between two carbon atoms which can be either a double bond or a triple bond. For example, the structure:



can represent either butene or butyne.

When compounded chemical names, *e.g.*, “alkylaryl,” “aryloxy,” and the like, are used herein, they are understood to have a specific connectivity to the core of the chemical structure. The moiety listed farthest to the right (*e.g.*, aryl in “alkylaryl”), is the moiety which is directly connected to the core. That is, for example, in a structure V-CH₂CH₂CH₃, when the variable “V” is an alkylaryl, the structure is understood to be alkyl-aryl-CH₂CH₂CH₃.

As used herein, the term “compound” is intended to mean a substance made up of molecules that further consist of atoms. A compound may be any natural or non-natural material, for example, peptide or polypeptide sequences, organic or inorganic molecules or compositions, nucleic acid molecules, carbohydrates, lipids or combinations thereof. A compound generally refers to a chemical entity, whether in the solid, liquid or gaseous phase, and whether in a crude mixture or purified and isolated. Compounds encompass the chemical compound itself as well as, where applicable: amorphous and crystalline forms of the compound, including polymorphic forms, said forms in mixture or in isolation; free acid and free base forms of the compound; isomers of the compound, including geometric isomers, optical isomers, and tautomeric isomers, said optical isomers to include enantiomers and diastereomers, chiral isomers and non-chiral isomers, said optical isomers to include isolated optical isomers or mixtures of optical isomers including racemic and non-racemic mixtures; said geometric isomers to include transoid and cisoid forms, where an isomer may be in isolated form or in admixture with one or more other isomers; isotopes of the compound, including

deuterium- and tritium-containing compounds, and including compounds containing radioisotopes, including therapeutically- and diagnostically-effective radioisotopes; multimeric forms of the compound, including dimeric, trimeric, etc. forms; salts of the compound, including acid addition salts and base addition salts, including organic counterions and inorganic counterions, and including zwitterionic forms, where if a compound is associated with two or more counterions, the two or more counterions may be the same or different; and solvates of the compound, including hemisolvates, monosolvates, disolvates, etc., including organic solvates and inorganic solvates, said inorganic solvates including hydrates; where if a compound is associated with two or more solvent molecules, the two or more solvent molecules may be the same or different.

In some embodiments, the term “compound” explicitly refers to small molecules. As used herein, the term “small molecule,” as used herein, refers to a compound that is not itself the product of gene transcription or translation (*e.g.*, protein, RNA, or DNA) and preferably has a low molecular weight, *e.g.*, organic molecules having a molecular weight of less than about 2,000 daltons (Da). In some embodiments, small molecules have a molecular weight of less than about 1,500 Da. In other embodiments, small molecules have a molecular weight of less than about 1,000 Da. In still other embodiments, small molecules have a molecular weight of less than about 750 Da. In yet other embodiments, small molecules have a molecular weight of less than about 500 Da.

The phrase, “pharmaceutically acceptable prodrugs” as used herein, refers to derivatives of a compound, usually with significantly reduced pharmacological activity, which contain an additional moiety, which are susceptible to removal *in vivo* yielding the parent molecule as the pharmacologically active species. An example of a pro-drug is an ester, which is cleaved *in vivo* to yield a compound of interest. Pro-drugs of a variety of compounds, and materials and methods for derivatizing the parent compounds to create the pro-drugs, are known and may be adapted to the present invention.


As used herein the term “ester” refers to a group having the formula R'–COOR, where one of R' or R is, for example, an alkyl group, a haloalkyl group or an aromatic group, and the other of R' or R is the active moiety. A “pharmaceutically acceptable ester” refers to an ester which acts as a prodrug, *e.g.*, an ester which is subject to at least partial removal (*e.g.*, by hydrolysis or other cleavage) from the core structure *in vivo*.

The term “protecting group” as used herein (*e.g.*, with regard to the synthesis of compounds of formula (I) or (II)), refers to a particular functional moiety, *e.g.*, O, S, or N, being temporarily blocked so that a reaction can be carried out selectively at another reactive site in a multifunctional compound. In preferred embodiments, a protecting group reacts selectively in good yield to give a protected substrate that is stable to the projected reactions; the protecting group must be selectively removed in good yield by readily available, preferably nontoxic reagents that do not attack the other functional groups; the protecting group forms an easily separable derivative (more preferably without the generation of new stereogenic centers); and the protecting group has a minimum of additional functionality to avoid further sites of reaction.

The term “amine” or “amino,” as used herein, refers to an unsubstituted or substituted moiety of the formula $-NR^aR^b$, in which R^a and R^b are each independently, for example, hydrogen, alkyl, aryl, or heterocyclyl, or R^a and R^b , taken together with the nitrogen atom to which they are attached, form a cyclic moiety having from 3 to 8 atoms in the ring.

Compounds of Formula (I) and (II)


The invention includes compounds of formula (I):



wherein:

R_1 is H

R_2 is a moiety selected from the group consisting of H and trifluoromethylcarbonyl; or R_1 and R_2 are taken together with the core structure to form a heterocyclidiyl moiety of formula (a):




R_3 is a moiety selected from the group consisting of $-OR_a$ and $-NR_bR_c$;


R_a is a C_{1-4} alkyl group optionally substituted with an imidazolyl;


R_b is a moiety selected from the group consisting of a C_{1-4} alkyl group and a C_{1-4} alkoxy group, wherein R_b is substituted with 0, 1 or 2 groups each independently selected from the group consisting of $-OCH_3$, $-C(O)OH$, $-C(O)NR'R''$, $-NH(C_{1-3} \text{ alkyl})$, $-NH(CH_2CH_2O)_nCH_3$, wherein n is 2-4, piperazinyl, N-methylpiperazinyl, piperidiny, N-methylpiperidiny, N-morpholinyl, imidazolyl, pyrrolidinyl, $-OPO_3H_2$ and hydroxyl; wherein the $-NH(C_{1-3} \text{ alkyl})$ group is substituted with 0, 1 or 2 hydroxyl moieties and wherein R' and R'' are each independently selected from $-H$ or $-CH_3$; and

R_c is H, or R_b and R_c are taken together with the nitrogen to which they are attached to form a heterocyclyl moiety selected from the group consisting of formula (b) and formula (c):



or a pharmaceutically acceptable salt or ester thereof.


In some embodiments,  represents a double bond. In some


embodiments,  represents a triple bond. In some embodiments, R_2 is H. In some embodiments, R_3 is


$-NR_bR_c$, where R_b is an unsubstituted C_{1-4} alkyl. In some embodiments, R_b is methyl. In some embodiments, R_b is ethyl. In some embodiments,


R_3 is $-NR_bR_c$, where R_b is a C_{1-4} alkyl, substituted with 1 or 2 $-OH$ groups. In some embodiments, R_b is 2-hydroxyethyl. In some embodiments, R_b is 2,3-dihydroxypropyl. In some embodiments, R_3 is $-OR_a$, where R_a is a C_{2-3} alkyl substituted with an imidazolyl group.


Exemplary imidazolyl groups include, but are not limited to,







imidazolyl group is . Exemplary morpholinyl groups include, but are not limited to,







group is . Exemplary piperidinyl groups include, but are not limited to,






piperidinyl group is . Exemplary N-methylpiperidinyl groups include, but are


not limited to, ,  and . In some embodiments, the

N-methylpiperidinyl group is . Exemplary pyrrolidinyl groups include, but are


not limited to, ,  and . In some embodiments, the

pyrrolidinyl group is . Exemplary piperazinyl groups include, but are not









limited to, , and . In some embodiments, the piperazinyl group is









. Exemplary













N-methylpiperazinyl groups include, but are not limited to,




In other embodiments, compounds of formula (I) or (II) include the compounds listed hereinbelow:

 (Compound 013)	 (Compound 054)
 (Compound 014)	 (Compound 065)
 (Compound 015)	 (Compound 069)
 (Compound 016)	 (Compound 076)


 (Compound 018)	 (Compound 091)
 (Compound 019)	 (Compound 092)
 (Compound 022)	 (Compound 106)
 (Compound 024)	 (Compound 114)

 (Compound 025)	 (Compound 122)
 (Compound 029)	 (Compound 127)
 (Compound 034)	 (Compound 137)
 (Compound 041)	 (Compound 144)
 (Compound 045)	 (Compound 155)
 (Compound 046)	 (Compound 156)



and pharmaceutically acceptable salts or esters thereof.


In some embodiments, the compound is at least one compound selected from the group consisting of:

























and pharmaceutically acceptable salts or esters thereof.


In at least one embodiment, the compound of formula (I) or (II) is not compound 029, compound 069, compound 091 or compound 106.

In some aspects, the present invention is directed to novel compounds of formula (I). All novel compounds of formula (I) described herein are included in the invention as compounds. In some embodiments, the present invention is directed to one or more of the compounds listed hereinbelow:




 (Compound 014)	 (Compound 054)
 (Compound 015)	 (Compound 065)
 (Compound 016)	 (Compound 076)
 (Compound 018)	 (Compound 114)
 (Compound 019)	 (Compound 122)

 (Compound 022)	 (Compound 127)
 (Compound 024)	 (Compound 137)
 (Compound 025)	 (Compound 144)
 (Compound 034)	 (Compound 155)
 (Compound 041)	 (Compound 156)
 (Compound 045)	 (Compound 157)



and pharmaceutically acceptable salts or esters thereof.

In some embodiments, the compounds of formula (I) include compounds of formula (II):




wherein

R_3 is $-NHR_b$, and R_b is C_{1-3} alkyl substituted with 0, 1, or 2 hydroxyl moieties; or a pharmaceutically acceptable salt or ester thereof.

In some embodiments, R_3 is an unsubstituted C_{1-3} alkyl-amino. In some embodiments, R_3 is methylamino. In other embodiments, R_3 is ethylamino. In some embodiments, R_3 is a C_{1-3} alkyl-amino substituted with one hydroxyl moiety. In some embodiments, R_3 is a C_{1-3} alkyl-amino substituted with two hydroxyl moieties. The hydroxyl moieties can be on any of the carbons in the C_{1-3} alkyl chain. Additionally, more than one hydroxyl moiety can be on a single carbon of the C_{1-3} alkyl chain. In some embodiments, there is a hydroxyl moiety on the 2-carbon of the alkyl chain. In

some embodiments, R_3 is hydroxyethylamino, *e.g.*, 2-hydroxyethylamino. In other embodiments, R_3 is dihydroxypropylamino, *e.g.*, 2,3-dihydroxypropylamino. In some embodiments, the C_{1-3} alkyl is an acyclic C_{1-3} alkyl chain.


In some embodiments, the compound is at least one compound selected from the group consisting of:



and pharmaceutically acceptable salts or esters thereof.

In at least one embodiment, the compound of formula (I) or (II) is not compound **029**, compound **069**, compound **091** or compound **106**.

In some aspects, the present invention is directed to novel compounds of formula (I). All novel compounds of formula (I) described herein are included in the invention as compounds. In some embodiments, the present invention is directed to one or more of the compounds listed hereinbelow:



and pharmaceutically acceptable salts or estersprodrugs thereof.

In at least one embodiment, the compound of formula (I) or (II) is not compound **029**, compound **091** or compound **106**. In at least one embodiment, the compound of formula (I) or (II) is not a specific compound listed in US Application Publication No. 2004/0224936.

The compounds of formula (I) or (II) may include one or more asymmetric centers, and thus can exist in various isomeric forms, *e.g.*, stereoisomers and/or

diastereomers. Thus, compounds and pharmaceutical compositions of formula (I) or (II) may be in the form of an individual enantiomer, diastereomer or geometric isomer, or may be in the form of a mixture of stereoisomers. In certain embodiments, the compounds of formula (I) or (II) are enantiopure compounds. In certain other embodiments, mixtures of stereoisomers or diastereomers are provided.

Compounds of formula (I) or (II) may also have one or more double bonds that can exist as either the Z or E isomer, unless otherwise indicated. The invention additionally encompasses the compounds as individual isomers substantially free of other isomers and alternatively, as mixtures of various isomers, *e.g.*, racemic mixtures of stereoisomers.

The compounds of formula (I) or (II) may further exist as one or a combination of crystalline forms, *e.g.*, polymorphs, solvates or hydrates of compounds of formula (I) or (II). Various crystalline forms may be identified and/or prepared using different solvents, or different mixtures of solvents for recrystallization; by performing crystallizations at different temperatures; or by using various modes of cooling, ranging from very fast to very slow cooling during crystallizations. Different crystalline forms may also be obtained by heating or melting the compound followed by gradual or fast cooling. The presence of polymorphs may be determined by solid probe NMR spectroscopy, IR spectroscopy, differential scanning calorimetry, powder X-ray diffractogram and/or other techniques.

Synthetic Methodology

Guidance for preparing compounds useful for practicing the present invention can be found, for example, in WO 05/023792 on pages 32-38 and WO 03/076424 on pages 28-36, as well as in the Examples presented therein. These references in combination with the information contained herein and the additional body of knowledge with regard to macrolide chemistry provides a person of skill in the art with guidance on synthetic strategies, protecting groups, and other materials and methods useful for the synthesis of the compounds of formula (I) or (II). For example, the above references provide background information on preparing compounds similar to the compounds described herein or relevant intermediates, as well as information on formulation, uses, and administration of such compounds.

Specific guidance and examples are also provided herein relating to various exemplary compounds and intermediates thereof. Accordingly, compounds of formula (I) or (II) and their preparation can be understood further by the examples herein, which illustrate some of the processes by which these compounds are prepared or used. It will be appreciated, however, that these examples do not limit the invention. Variations of the invention, now known or further developed, are considered to fall within the scope of the present invention as described herein and as hereinafter claimed.

According to the present invention, any available techniques can be used to make or prepare the inventive compounds or compositions including them. For example, a variety of solution phase synthetic methods such as those discussed in detail below may be used. Alternatively or additionally, the inventive compounds may be prepared using any of a variety combinatorial techniques, parallel synthesis and/or solid phase synthetic methods known in the art.

Starting materials and reagents useful in synthesizing the compounds of formula (I) or (II) are either commonly available from commercial suppliers or can be prepared by methods known to a person of ordinary skill in the art. The starting materials, intermediates, and compounds of formula (I) or (II) may be isolated and purified using conventional techniques, including filtration, distillation, crystallization, chromatography, and the like. They may be characterized using conventional methods, including physical constants and spectral data.

Exemplary syntheses of the compounds of formula (I) or (II) are provided in the Exemplification. It will be appreciated that the methods as described herein can be applied to each of the compounds as disclosed herein and equivalents thereof. Additionally, the reagents and starting materials are well known to those skilled in the art. Although the schemes described in the above-cited references depict certain exemplary compounds, it will be appreciated that the use of alternate starting materials will yield other analogs of formula (I) or (II).

Unless mentioned specifically, reaction mixtures were cooled to room temperature or below then quenched, when necessary, with either water or a saturated aqueous solution of ammonium chloride. Desired products were extracted by partitioning between water and a suitable water-immiscible solvent (*e.g.*, ethyl acetate, dichloromethane, diethyl ether). The desired product containing extracts were washed appropriately with water followed by a saturated solution of brine. On occasions where

the product containing extract was deemed to contain residual oxidants, the extract was washed with a 10% solution of sodium sulphite in saturated aqueous sodium bicarbonate solution, prior to the aforementioned washing procedure. On occasions where the product containing extract was deemed to contain residual acids, the extract was washed with saturated aqueous sodium bicarbonate solution, prior to the aforementioned washing procedure (except in those cases where the desired product itself had acidic character). On occasions where the product containing extract was deemed to contain residual bases, the extract was washed with 10% aqueous citric acid solution, prior to the aforementioned washing procedure (except in those cases where the desired product itself had basic character). Post washing, the desired product containing extracts were dried over anhydrous magnesium sulphate, and then filtered. The crude products were then isolated by removal of solvent(s) by rotary evaporation under reduced pressure, at an appropriate temperature (generally less than 45°C).

On occasions where triphenylphosphine oxide was a major byproduct of the reaction, the reaction mixture was added directly to a large volume of well-stirred hexane. The resultant precipitate of triphenylphosphine oxide was removed by filtration and the filtrate processed in the usual manner.

Unless mentioned specifically, chromatographic purification was used. Chromatographic purification refers to flash column chromatography on silica, using a single solvent or mixed solvent as eluent. Suitably purified desired product containing elutes were combined and concentrated under reduced pressure at an appropriate temperature (generally less than 45°C) to constant mass. If solid, final products were submitted for biological testing in their solid form. If freeze-drying was desired or necessary, final compounds were generally dissolved in 50% aqueous acetonitrile or other appropriate solvent or mixture of solvents, filtered and transferred to vials, then freeze-dried under high vacuum before submission for biological testing.

Pharmaceutical Compositions

In another aspect of the present invention, pharmaceutical compositions are provided, which comprise any one of the compounds described herein (or a pharmaceutically acceptable salt or ester thereof), and optionally comprise a pharmaceutically acceptable carrier. In certain embodiments, these compositions optionally further comprise one or more additional therapeutic agents. Alternatively, a

compound of this invention may be administered to a patient in need thereof in combination with the administration of one or more other therapeutic agents. For example, additional therapeutic agents for conjoint administration or inclusion in a pharmaceutical composition with a compound of this invention may be an anticancer agent approved for the treatment of cancer, as discussed in more detail herein, or it may be any one of a number of agents undergoing approval in the Food and Drug Administration that ultimately obtain approval for the treatment of cancer. It will also be appreciated that certain of the compounds of present invention can exist in free form for treatment, or where appropriate, as a pharmaceutically acceptable salt or ester thereof.

As used herein, the term “pharmaceutically acceptable salt” refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts of amines, carboxylic acids, and other types of compounds, are well known in the art. For example, S.M. Berge, *et al.* describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 66: 1-19 (1977). The salts can be prepared *in situ* during the final isolation and purification of the compounds of formula (I) or (II), or separately by reacting a free base or free acid function with a suitable reagent, as described generally below. For example, a free base function can be reacted with a suitable acid. Furthermore, where the compounds of formula (I) or (II) carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may, include metal salts such as alkali metal salts, *e.g.* sodium or potassium salts; and alkaline earth metal salts, *e.g.* calcium or magnesium salts. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, harnisulfate, heptanoate,

hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, lower alkyl sulfonate and aryl sulfonate.

Additionally, as used herein, the term “pharmaceutically acceptable ester” refers to esters that hydrolyze *in vivo* and include those that break down readily in the human body to leave the parent compound or a salt thereof. Suitable ester groups include, for example, those derived from pharmaceutically acceptable aliphatic carboxylic acids, particularly alkanoic, alkenoic, cycloalkanoic and alkanedioic acids, in which each alkyl or alkenyl moiety advantageously has not more than 6 carbon atoms. Examples of particular esters include formates, acetates, propionates, butyrates, acrylates and ethylsuccinates.

As described above, the pharmaceutical compositions comprising compounds of formula (I) or (II) may additionally comprise a pharmaceutically acceptable carrier, which, as used herein, may include any and all solvents, diluents, or other liquid vehicle, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like, as suited to the particular dosage form desired. Remington's Pharmaceutical Sciences, Sixteenth Edition, E. W. Martin (Mack Publishing Co., Easton, Pa., 1980) discloses various carriers used in formulating pharmaceutical compositions and known techniques for the preparation thereof. Except insofar as any conventional carrier medium is incompatible with the compounds of formula (I) or (II), such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutical composition, its use is contemplated to be within the scope of this invention. Some examples of materials which can serve as pharmaceutically acceptable carriers include, but are not limited to, sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose

and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatine; talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil; safflower oil, sesame oil; olive oil; corn oil and soybean oil; glycols; such as propylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogenfree water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.

Compositions comprising compounds of formula (I) or (II) may be formulated to have any concentration desired. In some embodiments, the composition is formulated such that it comprises at least a therapeutically effective amount. In some embodiments, the composition is formulated such that it comprises an amount that would not cause one or more unwanted side effects. In certain embodiments, compositions are formulated so that the compound is present at a concentration of between about 1 mg/mL and about 20 mg/mL; between about 1 mg/mL and about 15 mg/mL; between about 1 mg/mL and about 10 mg/mL; between about 2 mg/mL and about 9 mg/mL; between about 3 mg/mL and about 8 mg/mL; between about 4 mg/mL and about 7 mg/mL; between about 4 mg/mL and about 6 mg/mL. In certain embodiments, compositions are formulated such that the compound is present at a concentration of about 5 mg/mL.

Methods of Treatment

As discussed above, compounds of formula (I) or (II) exhibit a unique kinase inhibition profile. In some embodiments, based at least in part on this unique kinase inhibition profile, the compounds of formula (I) or (II) are useful for treating and/or preventing a variety of cancers, as discussed in further detail herein. Exemplary cancers include solid tumors, leukemias, lymphomas and myelomas such as multiple myeloma, neural cancer (e.g., glioma, neuroblastoma or retinoblastoma), leukemia (e.g., chronic myeloid leukemia, acute myeloid leukemia and/or acute lymphoblastic leukemia (ALL)), melanoma, pancreatic cancer, colorectal, thyroid and/or breast cancer. Accordingly, in some aspects, the present invention is directed to methods for the treatment and/or

prevention of a cancer in a subject. For example, in some embodiments, the present invention is directed to methods for the treatment a SrcTK/MEK associated cancer in a subject. In other embodiments, the present invention is directed to methods for the treatment a TrkB/MEK associated cancer in a subject.

The methods of the present invention generally include administering an effective amount of a composition comprising a selected multikinase inhibitor to the subject, such that the cancer is treated. The selected multikinase inhibitor is chosen such that it has a unique multikinase inhibition profile, which targets or is otherwise associated with the cancer. For example, in some embodiments, the methods for treating a SrcTK/MEK associated cancer include administering a SrcTK/MEK inhibitor or another compound of formula (I) or (II) to a subject in need thereof. In other embodiments, the methods for treating a TrkB/MEK associated cancer include administering a TrkB/MEK inhibitor or another compound of formula (I) or (II) to a subject in need thereof.

Compounds of formula (I) or (II) have a unique multikinase inhibitory profile. For example, in some embodiments, compounds of formula (I) or (II) have inhibitory activity against a number of protein kinases including the Src tyrosine kinase family, MEK1, MEKK1, TrkB, BCR-ABL, FLT-3 and KDR kinases. Such compounds include, for example compounds **106** and **091**, synthesis and certain biological properties of which are characterized in the Examples and Figures herein. This unique kinase inhibition profile suggests that compounds of formula (I) or (II) may have a broader therapeutic spectrum than selective MEK1 inhibitors.

In some embodiments, compounds of formula (I) or (II) include those which exhibit activity as inhibitors of MEK1; inhibitors of members of the Src family (*e.g.*, Src, Lyn, Fyn, Lck, Yes); inhibitors of the NF- κ B pathway through MEKK1 inhibition; inhibitors of TrkB; inhibitors of Bcr-Abl; inhibitors of FLT-3; and/or inhibitors of KDR.

In some embodiments, compounds of formula (I) or (II) include those which exhibit activity as inhibitors of MEK1. Accordingly, in some embodiments, an inhibitor of the present invention inhibits the activity of MEK1 by at least about 50% at a concentration of about 0.1 μ M. In some embodiments, an inhibitor of the present invention inhibits the activity of MEK1 by at least about 51% at a concentration of about 0.1 μ M. In some embodiments, an inhibitor of the present invention inhibits the activity of MEK1 by at least about 52% at a concentration of about 0.1 μ M. In some

embodiments, an inhibitor of the present invention inhibits the activity of MEK1 by at least about 53%, 54%, 55%, 56%, 57%, 58%, 59% or even 60% at a concentration of about 0.1 μ M. In further embodiments, an inhibitor of the present invention inhibits the activity of MEK1 by at least about 85% at a concentration of about 1.0 μ M. In still further embodiments, an inhibitor of the present invention inhibits the activity of MEK1 by at least about 86% at a concentration of about 1.0 μ M. In other embodiments, an inhibitor of the present invention inhibits the activity of MEK1 by at least about 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, or even 95% at a concentration of about 1.0 μ M.

In some embodiments, the compounds of formula (I) or (II) inhibit MEK1. For example, in some embodiments, compounds of formula (I) or (II) exhibit IC_{50} values less than 10 μ M. In certain other embodiments, compounds of formula (I) or (II) exhibit IC_{50} values less than 7.5 μ M. In certain embodiments, compounds of formula (I) or (II) exhibit IC_{50} values less than 5 μ M. In certain other embodiments, compounds of formula (I) or (II) exhibit IC_{50} values less than 2.5 μ M, less than 1 μ M, less than 0.75 μ M, less than 0.5 μ M, less than 0.25 μ M, less than 0.1 μ M, less than 75 nM or less than 50 nM. All ranges and values between the listed values are meant to be encompassed by the present invention.

The RAS-MAPK signaling pathway is also viewed as an important pathway for anticancer therapies, based upon its central role in regulating the growth and survival of cells from a broad spectrum of human tumors (Figure 1). MEK1 is downstream of RAS and RAF proteins, which are often mutated and abnormally active in cancer. Tumors with elevated MEK signaling based on B-RAF mutations are thus attractive targets for using the compositions and methods of the present invention.

It is believed that the Ras/Raf/MEK/ERK signal transduction pathway regulates cell proliferation in diverse types of cells. Mutations in this pathway are often observed in transformed cell lines and frequently linked with human cancer. Davies *et al.* (*Nature* **417**, 949-954, 2002), for example, identified B-RAF (an isoform of Raf) somatic missense mutations in 67% of malignant melanomas and 12% of colorectal cancers. All mutations were within the kinase domain with a single substitution, such as V600E (V599E), which accounts for 90% of these mutations. Mutated B-RAF proteins have elevated kinase activity, leading to constitutive activation of MEK, which then triggers ERK phosphorylation, and activates downstream pathway. It is believed that B-RAF

mutated cancers, including melanoma, are attractive therapeutic targets for MEK inhibitors. The frequency of B-RAF mutations in melanoma and colorectal cancers, and the relative lack of effective therapies for advanced stages of these diseases suggest that inhibition of B-RAF activity may be an important new strategy in these cancer types. Moreover, in certain embodiments, B-RAF mutation can be used as molecular marker for “patient enrichment strategies” (*i.e.*, for targeting B-RAF mutation carrying patients that will benefit from compositions and methods of the invention).

It was discovered by the present inventors that that compounds of formula (I) or (II) are potent inhibitors of B-RAF mutated cancer cell growth, and thus may selectively target B-RAF mutated cancers. As used herein, the term “B-RAF mutated cancer” refers to cancers that are associated with a particular point mutation of B-RAF. Accordingly, in some embodiments, the present invention provides compositions and methods for the treatment of B-RAF mutated cancers. Compounds of formula (I) or (II) include compounds which exhibit an antiproliferative activity against B-RAF mutated cancers; exhibit an antiproliferative effect on suitable cell lines maintained *in vitro*, or in animal studies using a scientifically acceptable model; and/or exhibit a favorable therapeutic profile (*e.g.*, safety, efficacy, and stability).

In certain embodiments, compounds of formula (I) or (II) are useful for treating and/or preventing tumors with elevated MEK signaling, including, but not limited to, those with B-RAF mutations. A number of proteins have received attention as targets for elevated MEK signaling (see Table 1 below, modified from Table 1 in Nature Reviews Cancer: 4:937-947, 2004). Mutations in these proteins could lead to activation of MEK-ERK pathway. Specifically, ovarian cancer, thyroid cancer, colorectal cancer and melanoma show high frequencies of B-RAF mutations.

Table 1

Tumor type	Pathway mutations in patient tumors
Colon	KRAS (45%), B-RAF (12%)
Pancreatic	KRAS (90%)
Ovarian	B-RAF (30%)
Melanoma	NRAS(15%), B-RAF (67%)
Non-small-cell lung	KRAS (35%)
Papillary thyroid	HRAS, KRAS and NRAS (60%); B-RAF (30-70%)
ALL, AML	NRAS (30%)

Accordingly, in certain embodiments, the methods of the present invention include methods for treating and/or preventing tumors with elevated MEK signaling, including, but not limited to, those with B-RAF mutations. In certain embodiments, compounds of formula (I) or (II) are useful for treating and/or preventing melanoma, colorectal cancer, ovarian cancer and/or thyroid cancer. Accordingly, in certain embodiments, the methods of the present invention include methods for treating and/or preventing a B-RAF mutated cancer. In certain embodiments, compounds of formula (I) or (II) are useful for treating and/or preventing cancers carrying a B-RAF mutation (e.g., a V600E mutation). In certain embodiments, the methods of the present invention include methods for treating and/or preventing a B-RAF mutated cancer carrying a V600E mutation.

In some embodiments, compounds of formula (I) or (II) include those which exhibit activity as inhibitors of a number of members of the Src tyrosine kinase family. Accordingly, in some embodiments, an inhibitor of the present invention inhibits the activity of at least five members of the Src tyrosine kinase family by at least about 40% at a concentration of about 0.1 μ M. In other embodiments, an inhibitor of the present invention inhibits the activity of at least five members of the Src tyrosine kinase family by at least about 45% at a concentration of about 0.1 μ M. In still other embodiments, an inhibitor of the present invention inhibits the activity of at least five members of the Src tyrosine kinase family by at least about 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, or even 55% at a concentration of about 0.1 μ M. In still other embodiments, an

inhibitor of the present invention inhibits the activity of at least five members of the Src tyrosine kinase family by at least about 85% at a concentration of about 1.0 μ M. In other embodiments, an inhibitor of the present invention inhibits the activity of at least five members of the Src tyrosine kinase family by at least about 90% at a concentration of about 1.0 μ M. In still other embodiments, an inhibitor of the present invention inhibits the activity of at least five members of the Src tyrosine kinase family by at least about 91%, 92%, 93%, 94%, or even 95% at a concentration of about 1.0 μ M. Exemplary members of the Src tyrosine kinase family include, but are not limited to cSrc, Fyn, Lyn, Lck and Yes.

In some embodiments, an inhibitor of the present invention inhibits the activity of at least four members of the Src tyrosine kinase family by at least about 40% at a concentration of about 0.1 μ M. In other embodiments, an inhibitor of the present invention inhibits the activity of at least four members of the Src tyrosine kinase family by at least about 45% at a concentration of about 0.1 μ M. In still other embodiments, an inhibitor of the present invention inhibits the activity of at least four members of the Src tyrosine kinase family by at least about 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, or even 55% at a concentration of about 0.1 μ M. In still other embodiments, an inhibitor of the present invention inhibits the activity of at least four members of the Src tyrosine kinase family by at least about 90% at a concentration of about 1.0 μ M. In still other embodiments, an inhibitor of the present invention inhibits the activity of at least four members of the Src tyrosine kinase family by at least about 91%, 92%, 93%, 94%, 95%, 96%, or even 97% at a concentration of about 1.0 μ M. In some embodiments, the four members of the Src tyrosine kinase family which are inhibited by the inhibitors, *e.g.*, SrcTK/MEK inhibitors, of the present invention are cSrc, Fyn, Lyn and Lck.

In some embodiments, an inhibitor of the present invention inhibits the activity of at least three members of the Src tyrosine kinase family by at least about 55% at a concentration of about 0.1 μ M. In other embodiments, an inhibitor of the present invention inhibits the activity of at least three members of the Src tyrosine kinase family by at least about 60% at a concentration of about 0.1 μ M. In still other embodiments, an inhibitor of the present invention inhibits the activity of at least three members of the Src tyrosine kinase family by at least about 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68% or even 69% at a concentration of about 0.1 μ M. In still other embodiments, an inhibitor of the present invention inhibits the activity of at least three members of the Src tyrosine

kinase family by at least about 93% at a concentration of about 1.0 μ M. In still other embodiments, an inhibitor of the present invention inhibits the activity of at least three members of the Src tyrosine kinase family by at least about 94%, 95%, 96%, 97% or even 98% at a concentration of about 1.0 μ M. In some embodiments, the three members of the Src tyrosine kinase family which are inhibited by the inhibitors, *e.g.*, SrcTK/MEK inhibitors, of the present invention are selected from cSrc, Fyn, Lyn and Lck. For example, in some embodiments, the three members of the Src tyrosine kinase family which are inhibited by the compounds and/or inhibitors of the present invention are Fyn, Lyn and Lck. For example, in other embodiments, the three members of the Src tyrosine kinase family which are inhibited by the inhibitors of the present invention are cSrc, Fyn and Lck.

In some embodiments, an inhibitor of the present invention inhibits the activity of at least two members of the Src tyrosine kinase family by at least about 65% at a concentration of about 0.1 μ M. In other embodiments, an inhibitor of the present invention inhibits the activity of at least two members of the Src tyrosine kinase family by at least about 70% at a concentration of about 0.1 μ M. In still other embodiments, an inhibitor of the present invention inhibits the activity of at least two members of the Src tyrosine kinase family by at least about 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or even 79% at a concentration of about 0.1 μ M. In still other embodiments, an inhibitor of the present invention inhibits the activity of at least two members of the Src tyrosine kinase family by at least about 95% at a concentration of about 1.0 μ M. In still other embodiments, an inhibitor of the present invention inhibits the activity of at least two members of the Src tyrosine kinase family by at least about 96%, 97%, 98% or even 99% at a concentration of about 1.0 μ M. In some embodiments, the two members of the Src tyrosine kinase family which are inhibited by the compounds and/or inhibitors, *e.g.*, SrcTK/MEK inhibitors, of the present invention are Fyn and Lck.

In some embodiments, an inhibitor of the present invention inhibits the activity of at least one member of the Src tyrosine kinase family by at least about 70% at a concentration of about 0.1 μ M. In other embodiments, an inhibitor of the present invention inhibits the activity of at least one member of the Src tyrosine kinase family by at least about 75% at a concentration of about 0.1 μ M. In still other embodiments, an inhibitor of the present invention inhibits the activity of at least one member of the Src tyrosine kinase family by at least about 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%,

84% or even 85% at a concentration of about 0.1 μ M. In still other embodiments, an inhibitor of the present invention inhibits the activity of at least one member of the Src tyrosine kinase family by at least about 97% at a concentration of about 1.0 μ M. In still other embodiments, an inhibitor of the present invention inhibits the activity of at least one member of the Src tyrosine kinase family by at least about 98%, 99% or even 100% at a concentration of about 1.0 μ M. In some embodiments, the one member of the Src tyrosine kinase family which is inhibited by the compounds and/or inhibitors, *e.g.*, SrcTK/MEK inhibitors, of the present invention is selected from Fyn and Lck.

The skilled artisan would understand that any or all of the above-indicated Src tyrosine kinase profiles can be accurate with regard to one single compound. That is, a single inhibitor of the present invention at a concentration of about 0.1 μ M can inhibit five members of the Src tyrosine kinase family by at least about 45%, four members by at least about 50%, three members by at least about 60%, two members by at least about 70% and one member by at least about 75%. It is also understood that, in the foregoing scenario, the one member of the Src tyrosine kinase family which is inhibited by at least 75% may also be one of the five members which are inhibited by at least 45%. Thus, in an exemplary case, a compound of formula (I) or (II) may have the following profile at a concentration of about 0.1 μ M: 50% inhibition of cSrc, 31% inhibition of Fyn, 42% inhibition of Lyn, 73% inhibition of Lck and 56% inhibition of Yes.

Without wishing to be bound by any particular theory, it is also believed that members of the Src tyrosine kinase family may also be useful in inhibition of the metastatic process. For example, Src has been associated with the establishment of metastasis through cell motility and invasion. Accordingly, in some embodiments, the methods of the present invention include methods for inhibition of a metastatic process in a subject.

Recently, it has also been shown that Lyn, a Src tyrosine kinase family member, plays an important role in the mechanism of imatinib mesylate resistance. Without wishing to be bound to any particular theory, it is also believed that certain members of the Src tyrosine kinase family are associated with chronic myeloid leukemia and/or colorectal cancer. Accordingly, in some embodiments, the present invention is directed to methods for treating chronic myeloid leukemia and/or colorectal cancer. Moreover, certain members of the Src tyrosine kinase family have been associated with imatinib mesylate resistance, *e.g.*, in chronic myeloid leukemia. Accordingly, the present

invention is directed to methods for treating cancers which are resistant to imatinib and pharmaceutically acceptable salts and esters thereof, *e.g.*, imatinib mesylate. In some embodiments, the present invention is directed to imatinib mesylate resistant chronic myeloid leukemia.

As used herein, the term “resistance” refers to the occasion where a subject becomes less responsive to a therapeutic agent over time. Accordingly, in some embodiments, resistance to a therapeutic agent refers to a subject’s complete non-responsiveness to the agent (*e.g.*, where rate of growth of a tumor is not inhibited) as compared to the absence of administration of the agent. In some embodiments, resistance to a therapeutic agent refers to a subject’s partial non-responsiveness to the agent (*e.g.*, where rate of growth of a tumor is inhibited only to a very low degree) as compared to the absence of administration of the agent. The quality of being resistant to a therapeutic agent is a highly variable one, with different tumors exhibiting different levels of “resistance” to a given therapeutic agent, under different conditions. In other embodiments, resistance to a therapeutic agent refers to a subject’s complete or partial non-responsiveness to the agent as compared to a previous administration of the agent. In still other embodiments, resistance to a therapeutic agent refers to a subject who is being administered the agent, and desires additional therapy.

In some embodiments, compounds of formula (I) or (II) include those which exhibit activity as inhibitors of TrkB. Accordingly, in some embodiments, an inhibitor of the present invention inhibits the activity of TrkB by at least about 65% at a concentration of about 0.1 μ M. In some embodiments, an inhibitor of the present invention inhibits the activity of TrkB by at least about 70% at a concentration of about 0.1 μ M. In some embodiments, an inhibitor of the present invention inhibits the activity of TrkB by at least about 75% at a concentration of about 0.1 μ M. In some embodiments, an inhibitor of the present invention inhibits the activity of TrkB by at least about 76%, 78%, 80%, 82%, 84%, 86% or even 88% at a concentration of about 0.1 μ M. In further embodiments, an inhibitor of the present invention inhibits the activity of TrkB by at least about 85% at a concentration of about 1.0 μ M. In still further embodiments, an inhibitor of the present invention inhibits the activity of TrkB by at least about 90% at a concentration of about 1.0 μ M. In other embodiments, a compound or inhibitor of the present invention inhibits the activity of TrkB by at least

about 91%, 92%, 93%, 94%, 95%, 96%, 97%, or even 98% at a concentration of about 1.0 μ M.

Without wishing to be bound by any particular theory, it is believed that TrkB, the nerve derived growth factor (NDGF) receptor, may play important roles in proliferation and survival of neural cancers such as glioma, retinoblastoma and neuroblastoma. Moreover, it is also believed that TrkB activity may be associated with pancreatic cancer. For example, signal transduction through TrkB may cause severe pain in patients with pancreatic cancer. Accordingly, in some embodiments, the methods of the present invention include methods for treating and/or preventing glioma, retinoblastoma and neuroblastoma. In other embodiments, the methods of the present invention include methods for treating and/or preventing pancreatic cancer.

In certain embodiments, compounds of formula (I) or (II) or active metabolites thereof exhibit high penetration through the blood brain barrier (BBB). When used with regard to the amount which crosses the BBB, the term “a compound of formula (I) or (II)” is intended to include active metabolites thereof. Accordingly, in some embodiments, the methods of the present invention include methods for treating a central nervous system tumor in a subject in need thereof. The method generally includes administration of a composition as described herein (including a compound of formula (I) or (II)) such that at least a portion of the compound of formula (I) or (II) penetrates the blood brain barrier, as demonstrated in a suitable animal. In some embodiments, at least about 20% of the compound of formula (I) or (II) penetrates the blood brain barrier, based upon a ratio of amount of a compound of formula (I) or (II) in the brain to amount of a compound of formula (I) or (II) in the plasma, as demonstrated in a suitable animal. In some embodiments, at least about 40% of the compound of formula (I) or (II) penetrates the blood brain barrier, based upon a ratio of amount of the compound in the brain to amount of a compound of formula (I) or (II) in the plasma, as demonstrated in a suitable animal. In other embodiments, at least about 50% of the compound of formula (I) or (II) penetrates the blood brain barrier, as demonstrated in a suitable animal. In still other embodiments, at least about 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or even 100% of the compound of formula (I) or (II) penetrates the blood brain barrier, as demonstrated in a suitable animal. Thus, whether a compound of formula (I) or (II) penetrates the blood brain barrier of a subject (e.g., a human) can be determined in a suitable animal (e.g., a rodent, such as a mouse). It is to be noted that

the percentages are calculated based upon the area under the concentration-time curve for a given time period (AUC_{0-t}) in the brain versus the plasma. Accordingly, the percentages represent a ratio of concentrations. That is, if (AUC_{0-24h}) for a compound is 20ng/mL in the brain and 80ng/mL in the plasma, then the percentage of the compound that penetrates the blood brain barrier is 20% (20ng/ml in the brain divided by the total concentration of (20ng/mL + 80ng/mL)). In some embodiments, the percentages are calculated based upon the area under the concentration-time curve for the time period from $t=0$ (time of dosing) to the last quantifiable concentration point, *i.e.*, (AUC_{0-last}). Exemplary central nervous system tumors include, but are not limited to, brain tumors, glioma and neuroblastoma.

In some embodiments, the inhibitors of the present invention inhibit Bcr-Abl and/or FLT-3. Bcr-Abl and FLT-3 are known targets for leukemia treatment. Accordingly, in certain embodiments, the methods of the present invention include methods for treating and/or preventing leukemia. In some embodiments, an inhibitor of the present invention inhibits the activity of Bcr-Abl by at least about 40% at a concentration of about 0.1 μ M. In some embodiments, an inhibitor of the present invention inhibits the activity of Bcr-Abl by at least about 45% at a concentration of about 0.1 μ M. In some embodiments, an inhibitor of the present invention inhibits the activity of Bcr-Abl by at least about 46%, 47%, 48%, 49%, 50%, 52% or even 54% at a concentration of about 0.1 μ M. In further embodiments, an inhibitor of the present invention inhibits the activity of Bcr-Abl by at least about 75% at a concentration of about 1.0 μ M. In still further embodiments, an inhibitor of the present invention inhibits the activity of Bcr-Abl by at least about 80% at a concentration of about 1.0 μ M. In other embodiments, an inhibitor of the present invention inhibits the activity of Bcr-Abl by at least about 81%, 82%, 83%, 84%, 85%, 86%, 88%, or even 90% at a concentration of about 1.0 μ M. In some embodiments, an inhibitor of the present invention inhibits the activity of FLT-3 by at least about 45% at a concentration of about 0.1 μ M. In some embodiments, an inhibitor of the present invention inhibits the activity of FLT-3 by at least about 50% at a concentration of about 0.1 μ M. In some embodiments, an inhibitor of the present invention inhibits the activity of FLT-3 by at least about 52%, 54%, 56%, 58%, 60%, 65% or even 70% at a concentration of about 0.1 μ M. In further embodiments, an inhibitor of the present invention inhibits the activity of FLT-3 by at least about 85% at a concentration of about 1.0 μ M. In still further embodiments, an

inhibitor of the present invention inhibits the activity of FLT-3 by at least about 90% at a concentration of about 1.0 μ M. In other embodiments, an inhibitor of the present invention inhibits the activity of FLT-3 by at least about 91%, 92%, 93%, 94%, 95% or even 96% at a concentration of about 1.0 μ M.

Compounds and compositions of formula (I) or (II) are useful for treating a *FLT3* mutated cancer in a subject. As used herein, FLT3 mutated cancers are cancers associated with activation mutations in *FLT3* which activate the kinase function of the encoded protein.

The FMS-like tyrosine kinase 3 gene (*FLT3* or *FLT-3*) encodes a member of the type III platelet-derived growth factor family of receptor tyrosine kinases. *FLT3* is expressed in many hematological malignancies, and its signaling cascade has been implicated in multiple tumorigenic pathways. For example, *FLT3* mutations can lead to activation of the STAT5 and Ras/MAPK pathways. (Parcells *et al.*, *Stem Cells*, 24(5); 1174 -1184 (2006))

Activation mutations of *FLT3* have been reported in leukemia patients. For example, the two most frequent types of *FLT3* mutations associated with adult cytogenetically normal acute myeloid leukemia (CN-AML) are: (1) internal tandem duplications (ITD) in the juxtamembrane domain (present in approximately 30% of CN-AML patients); and (2) mutations in the second tyrosine kinase domain (TKD) (present in approximately 7% of CN-AML patients). Common TKD mutations include nucleotide substitutions, deletions, or insertions affecting amino acid residues D835 and/or I836 in the activation loop (A-loop) of the TKD. The D835Y substitution constitutes approximately 50% of TKD mutations reported. Y842C, K663Q, and V592A mutations have also been observed. FLT3 proteins with these ITD or TKD mutations exhibit ligand-independent FLT3 dimerization and constitutive activation through autophosphorylation. (Whitman *et al.*, *Blood*, 111(3):1552-1559 (2008); see also, Parcells *et al.*, *Stem Cells*, 24(5); 1174 -1184 (2006)) Activation mutations of *FLT3* have been associated with an inferior prognosis. (Stone, *Blood*, 104I: 915-916 (2004))

In some embodiments, the compounds of formula (I) or (II) can inhibit FLT3 mutated cancer cell growth. In certain embodiments, the compounds can exhibit antiproliferative activity against FLT3 mutated cancers; an antiproliferative effect on suitable cell lines maintained *in vitro* or in animal models (e.g., comprising an activation mutation of *FLT3*); and/or a favorable therapeutic profile (e.g., safety, efficacy, and

stability).

Accordingly, in some embodiments, the invention provides a method of treating a FLT3 mutated cancer in a subject in need thereof, comprising administering a therapeutically effective amount of a compound or composition of formula (I) or (II) to said subject. In certain embodiments of the method, a compound of formula (I) or (II) or a composition comprising same is used for treating cancers carrying a FLT3 mutation. In certain embodiments of the method, the FLT3 mutated cancer carries an internal tandem duplication (ITD) in the juxtamembrane domain. In certain embodiments of the method, the FLT3 mutated cancer carries a mutation in the second tyrosine kinase domain, for example, in the activation loop. In certain embodiments of the method, the FLT3 mutated cancer carries a mutation of residue D835 and/or a mutation of residue I836, for example, a D835Y mutation. In certain embodiments of the method, the FLT3 mutated cancer carries a Y842C, K663Q or V592A mutation. Moreover, in certain embodiments, FLT3 mutation can be used as molecular marker for “patient enrichment strategies” in order to identify patients for treatment.

As noted above, activation mutations of *FLT3* have been reported in leukemia patients, and the invention provides a method of treating leukemia in a subject, comprising administering a therapeutically effective amount of a compound or composition of formula (I) or (II) to the subject. In one embodiment of the method the leukemia is acute myeloid leukemia (AML).

Without wishing to be bound by any particular theory, it is believed that the unique multikinase inhibition profiles of the compounds of formula (I) or (II) indicates that such compounds may be useful in targeting hematological malignancies as well as solid tumors. (See, *e.g.*, Warmuth, M et al. *Ann Hematol.* 78(2):49-64 (1999) and Harder, KW et al. *Immunity.* 15(4):603-15 (2001).) In certain embodiments, compounds of formula (I) or (II) exhibit anticancer activity with evidence of apoptosis in B-cell non-Hodgkin’s lymphoma, possibly by inhibition of MEKK1, an upstream molecule for NF- κ B activation, which leads to chemoresistance to apoptosis. Accordingly, in certain embodiments, the methods of the present invention include methods for treating and/or preventing hematologic malignant cancer cell growth such as multiple myeloma and B-cell lymphomas.

In still other embodiments, the compounds of formula (I) or (II) inhibit the growth of tumor cell lines *in vitro*. For example, in some embodiments, compounds of

formula (I) or (II) exhibit IC_{50} values less than 10 μ M. In certain other embodiments, compounds of formula (I) or (II) exhibit IC_{50} values less than 7.5 μ M. In certain embodiments, compounds of formula (I) or (II) exhibit IC_{50} values less than 5 μ M. In certain other embodiments, compounds of formula (I) or (II) exhibit IC_{50} values less than 2.5 μ M, less than 1 μ M, less than 0.75 μ M, less than 0.5 μ M, less than 0.25 μ M, less than 0.1 μ M, less than 75 nM or less than 50 nM. All ranges and values between the listed values are meant to be encompassed by the present invention.

In some embodiments, compounds of formula (I) or (II) cause tumor regression *in vivo*. In certain exemplary embodiments, compounds of formula (I) or (II) cause tumor regression *in vivo* in suitable mouse tumor xenograft models. As demonstrated in the Examples herein, certain compounds of formula (I) or (II) have strong *in vivo* efficacy in multiple xenografts by *i.v.* dosing. In certain embodiments, compounds of formula (I) or (II) cause reduction of tumor size to below 70% of the size at the start of compound administration in a suitable cancer cell xenograft model. In certain embodiments, compounds of formula (I) or (II) cause reduction of tumor size to below 65% of the size at the start of compound administration in a suitable cancer cell xenograft model. In certain embodiments, compounds of formula (I) or (II) cause reduction of tumor size to below 60% below 55% or below 50% of the size at the start of compound administration in a suitable cancer cell xenograft model. All values and ranges between the listed values are intended to be encompassed by the present invention.

In certain embodiments, compounds of formula (I) or (II) cause inhibition of tumor growth *in vivo*. For example, in some embodiments, compounds of formula (I) or (II) cause significant inhibition of tumor growth in suitable cancer cell xenograft models. In certain embodiments, compounds of formula (I) or (II) cause inhibition of tumor growth in treated animals by $> 50\%$ compared to that of control animals (*i.e.*, “treated” tumor size $< 50\%$ “control” tumor size; or T/C value $< 50\%$) in suitable cancer cell xenograft models. In certain embodiments, compounds of formula (I) or (II) have T/C values $< 70\%$. In certain embodiments, compounds of formula (I) or (II) have T/C values $< 65\%$, $< 60\%$, or $< 55\%$. All values and ranges between the listed values are intended to be encompassed by the present invention.

Exemplary compounds, including F152 (natural product), were tested for their growth inhibitory activity in MDA-MB-435 and other cancer cell lines. Several

compounds were shown to have remarkable cancer cell growth inhibitory activities (*e.g.*, with nM IC₅₀ values). Based on biological characterization in various relevant assays (See Examples and Figures), it is believed that certain compounds of formula (I) or (II) exhibit potent cell growth inhibition especially toward B-RAF mutated cells; possess a multikinase inhibitory profile, including MEK1, MEKK1, Src kinase family tyrosine kinase, TrkB, FLT-3, and Bcr-Abl; have potent *in vivo* antitumor activity in multiple tumor xenografts including “major tumor types” without significant body weight loss; have projecting efficacy in Q4D and/or Q7D *iv* dosing in subjects; have signs of better efficacy than certain MEK1 inhibitors known in the art in combination therapy; be able to penetrate the blood-brain barrier, which renders the compounds potentially useful for treating/preventing brain tumors; be potentially useful for targeted therapy based on possible biomarkers for specific cancers.

In some embodiments, compounds of formula (I) or (II) are natural product-based multi-kinase inhibitors that are highly effective against cancer-relevant kinases including the Src family (Src, Lyn, Fyn, Lck, Yes), NF-κB pathway through MEKK1 inhibition, TrkB, Bcr-Abl, FLT-3 and KDR, as well as B-RAF mutated cancer cells. This unique profile may offer the compounds of formula (I) or (II) a competitive advantage in the clinic, for example low levels of cytotoxicity against non-dividing cells, and therefore potentially good therapeutic margin.

In certain embodiments, compounds of formula (I) or (II) exhibit synergy in combination therapy with another anticancer agent. In certain exemplary embodiments, compounds of formula (I) or (II) exhibit synergy in combination therapy with SN-38 (active metabolite of CPT-11). In some embodiments, compounds of formula (I) or (II) are more effective in tumor cell killing when used in combination with another anticancer agent (*e.g.*, SN-38, an active metabolite of CPT-11— see, *e.g.*, Example 9).

In certain embodiments, compounds of formula (I) or (II) exhibit inhibitory activity against NF-κB. As such, they are useful for treating and/or preventing cancers related to NF-κB activity. In certain embodiments, compounds of formula (I) or (II) are useful for treating and/or preventing hematologic malignancies such as multiple myeloma and/or B cell lymphomas. In some embodiments, compounds of formula (I) or (II) are beneficial for cancer patients showing resistance in conventional chemotherapy *via* NF-κB activation.

According to the present invention, the inventive compounds may be assayed in any of the available assays known in the art for identifying compounds having, for example, MEK1 inhibitory activity, protein kinase inhibitory activity, NF- κ B activation inhibitory activity and cancer cell growth inhibitory activity. For example, the assay may be cellular or non-cellular, *in vivo* or *in vitro*, high- or low-throughput format, etc.

Dosages and Modes of Administration

It will be appreciated that the compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for the treatment of cancer. Thus, the expression “effective amount” as used herein, refers to a sufficient amount of agent to inhibit the growth of tumor cells, or refers to a sufficient amount to reduce the effects of cancer. The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the diseases, the particular anticancer agent, its mode of administration, and the like. The compounds of formula (I) or (II) are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. The expression “dosage unit form” as used herein refers to a physically discrete unit of therapeutic agent appropriate for the patient to be treated. It will be understood, however, that the total daily usage of the compounds and compositions of formula (I) or (II) will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient or organism will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts (see, for example, Goodman and Gilman’s, “The Pharmacological Basis of Therapeutics”, Tenth Edition, A. Gilman, J. Hardman and L. Limbird, eds., McGraw-Hill Press, 155-173, 2001).

In certain embodiments, the compounds or compositions are administered systemically. As used herein, “systemic administration” refers to any means by which a compound of formula (I) or (II) can be made systemically available. For example,

systemic administration encompasses enteral (*e.g.*, oral and rectal) and parenteral methods of administration. In certain embodiments, systemic administration encompasses intravenous administration, intraperitoneal administration, intramuscular administration, intracoronary administration, intraarterial administration (*e.g.*, into a carotid artery), intradermal administration, subcutaneous administration, transdermal delivery, intratracheal administration, subcutaneous administration, intraarticular administration, intraventricular administration, inhalation (*e.g.*, aerosol), intracerebral, nasal, nasal, oral, intraocular, pulmonary administration, impregnation of a catheter, by suppository and direct injection into a tissue, or systemically absorbed topical or mucosal administration. Mucosal administration includes administration to the respiratory tissue, *e.g.*, by inhalation, nasal drops, ocular drop, etc.; anal or vaginal routes of administration, *e.g.*, by suppositories; and the like. In certain exemplary embodiments, the compositions of formula (I) or (II) are administered intravenously.

It will be appreciated that the inventive compound may be administered systemically in dosage forms, formulations or *e.g.* suitable delivery devices or implants containing conventional, non-toxic pharmaceutically acceptable carriers and adjuvants such that the compound effectiveness is optimized. For example, the inventive compound may be formulated together with appropriate excipients into a pharmaceutical composition, which, upon administration of the composition to the subject, systemically releases the active substance in a controlled manner. Alternatively, or additionally, compound dosage form designs may be optimized so as to increase the compound effectiveness upon administration. The above strategies (*i.e.*, dosage form design and rate control of drug input), when used alone or in combination, can result in a significant increase in compound effectiveness and are considered part of the invention.

Furthermore, after formulation with an appropriate pharmaceutically acceptable carrier in a desired dosage, the pharmaceutical compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, creams or drops), buccally, as an oral or nasal spray, or the like. In certain embodiments, the compounds of formula (I) or (II) may be administered at dosage levels of about 0.001 mg/kg to about 50 mg/kg, from about 0.01 mg/kg to about 25 mg/kg, or from about 0.1 mg/kg to about 10 mg/kg of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect. It will also be appreciated that dosages smaller than 0.001 mg/kg or

greater than 50 mg/kg (for example 50-100 mg/kg) can be administered to a subject. In certain embodiments, compounds are administered orally or parenterally. In certain embodiments, administration is conducted by intermittent dosing one to three times a week.

In certain embodiments, the compounds of formula (I) or (II) may be administered at dosage levels of about 1-500 mg/m²; *e.g.*, about 5-400 mg/m²; about 10-250 mg/m² one or more times a week to obtain the desired therapeutic effect. Specific ranges and values which fall between the listed ranges are also contemplated by the present invention.

In certain embodiments, the compounds of formula (I) or (II) may be administered intravenously one to three times a week. In certain embodiments, the compounds of formula (I) or (II) may be administered intravenously once a week. In certain embodiments, the compounds of formula (I) or (II) may be administered intravenously twice a week.

Administration of the compounds of formula (I) or (II) can be continuous over a specified period of time, can be intermittent or can be in a bolus. For example, in one embodiment, the compounds of formula (I) or (II) are formulated for intravenous use and can be administered to patients as a slow injection, *e.g.*, over about 30 minutes to about 3 hours. In another embodiment, the compounds of formula (I) or (II) are formulated for intravenous use and can be administered to patients as a bolus injection, also referred to as IV (intravenous) push. In still another embodiment, the compounds of formula (I) or (II) are formulated for intravenous use and can be administered to patients in any number of discreet injections that are separated by time periods which may or may not be equivalent.

Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral

compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

In order to prolong the effect of a drug, it is often desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension or crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution that, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle. Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include (poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions, which are compatible with body tissues.

Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository

wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar--agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

The active compounds can also be in microencapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active compound may be admixed with at least one

inert diluent such as sucrose, lactose and starch. Such dosage forms may also comprise, as in normal practice, additional substances other than inert diluents, *e.g.*, tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions, which can be used, include polymeric substances and waxes.

The present invention encompasses pharmaceutically acceptable topical formulations of inventive compounds. The term "pharmaceutically acceptable topical formulation", as used herein, means any formulation which is pharmaceutically acceptable for intradermal administration of a compound of formula (I) or (II) by application of the formulation to the epidermis. In certain embodiments of the invention, the topical formulation comprises a carrier system. Pharmaceutically effective carriers include, but are not limited to, solvents (*e.g.*, alcohols, poly alcohols, water), creams, lotions, ointments, oils, plasters, liposomes, powders, emulsions, microemulsions, and buffered solutions (*e.g.*, hypotonic or buffered saline) or any other carrier known in the art for topically administering pharmaceuticals. A more complete listing of art-known carriers is provided by reference texts that are standard in the art, for example, Remington's Pharmaceutical Sciences, 16th Edition, 1980 and 17th Edition, 1985, both published by Mack Publishing Company, Easton, Pa. In certain other embodiments, the topical formulations of the invention may comprise excipients. Any pharmaceutically acceptable excipient known in the art may be used to prepare the inventive pharmaceutically acceptable topical formulations. Examples of excipients that can be included in the topical formulations of the invention include, but are not limited to, preservatives, antioxidants, moisturizers, emollients, buffering agents, solubilizing agents, other penetration agents, skin protectants, surfactants, and propellants, and/or additional therapeutic agents used in combination to the inventive compound. Suitable preservatives include, but are not limited to, alcohols, quaternary amines, organic acids, parabens, and phenols. Suitable antioxidants include, but are not limited to, ascorbic acid and its esters, sodium bisulfite, butylated hydroxytoluene, butylated hydroxyanisole, tocopherols, and chelating agents like EDTA and citric acid. Suitable moisturizers

include, but are not limited to, glycerine, sorbitol, polyethylene glycols, urea, and propylene glycol. Suitable buffering agents for use with the invention include, but are not limited to, citric, hydrochloric, and lactic acid buffers. Suitable solubilizing agents include, but are not limited to, quaternary ammonium chlorides, cyclodextrins, benzyl benzoate, lecithin, and polysorbates. Suitable skin protectants that can be used in the topical formulations of the invention include, but are not limited to, vitamin E oil, allantoin, dimethicone, glycerin, petrolatum, and zinc oxide.

In certain embodiments, the pharmaceutically acceptable topical formulations of the invention comprise at least a compound of formula (I) or (II) and a penetration enhancing agent. The choice of topical formulation will depend on several factors, including the condition to be treated, the physicochemical characteristics of the inventive compound and other excipients present, their stability in the formulation, available manufacturing equipment, and costs constraints. As used herein the term "penetration enhancing agent" means an agent capable of transporting a pharmacologically active compound through the stratum corneum and into the epidermis or dermis, preferably, with little or no systemic absorption. A wide variety of compounds have been evaluated as to their effectiveness in enhancing the rate of penetration of drugs through the skin. See, for example, *Percutaneous Penetration Enhancers*, Maibach H. I. and Smith H. E. (eds.), CRC Press, Inc., Boca Raton, Fla. (1995), which surveys the use and testing of various skin penetration enhancers, and Buyuktimkin *et al.*, *Chemical Means of Transdermal Drug Permeation Enhancement in Transdermal and Topical Drug Delivery Systems*, Gosh T. K., Pfister W. R., Yum S. I. (Eds.), Interpharm Press Inc., Buffalo Grove, Ill. (1997). In certain exemplary embodiments, penetration agents for use with the invention include, but are not limited to, triglycerides (*e.g.*, soybean oil), aloe compositions (*e.g.*, aloe-vera gel), ethyl alcohol, isopropyl alcohol, octolyphenylpolyethylene glycol, oleic acid, polyethylene glycol 400, propylene glycol, N-decylmethylsulfoxide, fatty acid esters (*e.g.*, isopropyl myristate, methyl laurate, glycerol monooleate, and propylene glycol monooleate) and N-methyl pyrrolidone.

In certain embodiments, the compositions may be in the form of ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. In certain exemplary embodiments, formulations of the compositions according to the invention are creams, which may further contain saturated or unsaturated fatty acids such as stearic

acid, palmitic acid, oleic acid, palmito-oleic acid, cetyl or oleyl alcohols, stearic acid being particularly preferred. Creams of the invention may also contain a non-ionic surfactant, for example, polyoxy-40-stearate. In certain embodiments, the active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, eardrops, and eye drops are also contemplated as being within the scope of this invention. Additionally, the present invention contemplates the use of transdermal patches, which have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms are made by dissolving or dispensing the compound in the proper medium. As discussed above, penetration enhancing agents can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

In certain exemplary embodiments, pharmaceutically acceptable topical formulations of the invention are contained in a patch that is applied adjacent to the area of skin to be treated. As used herein a "patch" comprises at least a topical formulation and a covering layer, such that, the patch can be placed over an area of skin. Any patch known in the art can be used in conjunction with the compositions and methods of the present invention.

In certain exemplary embodiments, the inventive compounds may be used as coating for stents. Guidance for using compounds of formula (I) or (II) in this capacity can be found, for example, in WO 05/023792.

Coadministration

It will also be appreciated that the compounds and pharmaceutical compositions of formula (I) or (II) can be formulated and employed in combination therapies, that is, the compounds and pharmaceutical compositions can be formulated with or administered concurrently with, prior to, or subsequent to, one or more other desired therapeutics or medical procedures. The particular combination of therapies (therapeutics or procedures) to employ in a combination regimen will take into account compatibility of the desired therapeutics and/or procedures and the desired therapeutic effect to be achieved. It will also be appreciated that the therapies employed may achieve a desired effect for the same disorder (for example, an inventive compound may

be administered concurrently with another anticancer agent), or they may achieve different effects (*e.g.*, control of any adverse effects).

For example, other therapies or anticancer agents that may be used in combination with the inventive compounds of formula (I) or (II) include surgery, radiotherapy (in but a few examples, γ -radiation, neutron beam radiotherapy, electron beam radiotherapy, proton therapy, brachytherapy, and systemic radioactive isotopes, to name a few), endocrine therapy, biologic response modifiers (interferons, interleukins, and tumor necrosis factor (TNF) to name a few), hyperthermia and cryotherapy, agents to attenuate any adverse effects (*e.g.*, antiemetics), and other approved chemotherapeutic drugs, including, but not limited to, alkylating drugs (mechlorethamine, chlorambucil, Cyclophosphamide, Melphalan, Ifosfamide), antimetabolites (Methotrexate), purine antagonists and pyrimidine antagonists (6-Mercaptopurine, 5-Fluorouracil, Cytarabine, Gemcitabine), spindle poisons (Vinblastine, Vincristine, Vinorelbine, Paclitaxel), podophyllotoxins (Etoposide, Irinotecan, Topotecan), antibiotics (Doxorubicin, Bleomycin, Mitomycin), nitrosoureas (Carmustine, Lomustine), inorganic ions (Cisplatin, Carboplatin), enzymes (Asparaginase), and hormones (Tamoxifen, Leuprolide, Flutamide, and Megestrol), to name a few. Other anticancer agents that may be used in combination with compounds or compositions of formula (I) or (II), include a therapeutic antibody or antibody fragment, such as a human, humanized, chimeric, and/or single-chain antibody or antibody fragment (*e.g.*, Fv, Fab, Fab', F(ab')₂, dAb) or the like (*e.g.*, Trastuzumab, Bevacizumab, Cetuximab, Rituximab). For a more comprehensive discussion of updated cancer therapies see, The Merck Manual, Seventeenth Ed. 1999. See also the National Cancer Institute (NCI) website (www.nci.nih.gov) and the Food and Drug Administration (FDA) website for a list of the FDA approved oncology drugs (www.fda.gov/cder/cancer/druglistframe – See Appendix A).

In certain embodiments, the pharmaceutical compositions comprising compounds of formula (I) or (II) further comprise one or more additional therapeutically active ingredients (*e.g.*, chemotherapeutic and/or palliative). For purposes of the invention, the term "*Palliative*" refers to treatment that is focused on the relief of symptoms of a disease and/or side effects of a therapeutic regimen, but is not curative. For example, palliative treatment encompasses painkillers, anti-nausea medications and anti-sickness drugs. In addition, chemotherapy, radiotherapy and surgery can all be used

palliatively (that is, to reduce symptoms without going for cure; *e.g.*, for shrinking tumors and reducing pressure, bleeding, pain and other symptoms of cancer).

Treatment Kits

In some aspects, the present invention provides a kit for carrying out the methods in accordance with the present invention. In general, the pharmaceutical pack or kit comprises one or more of the ingredients of the pharmaceutical compositions of formula (I) or (II). For example, the kit may include one or more containers filled (completely or partially) with a compound or composition of formula (I) or (II). Such kits are suited for, *e.g.*, the delivery of solid oral forms such as tablets or capsules. In some embodiments, the kit includes a number of unit dosages, and may also include a card having the dosages oriented in the order of their intended use. If desired, a memory aid can be provided, for example in the form of numbers, letters, or other markings or with a calendar insert, designating the days in the treatment schedule in which the dosages can be administered. Alternatively, placebo dosages, or calcium dietary supplements, either in a form similar to or distinct from the dosages of the pharmaceutical compositions, can be included to provide a kit in which a dosage is taken every day. In some embodiments, the kit includes instructions or directions for the use of the other components of the kit. In some embodiments, the kit comprises a composition comprising a compound of formula (I) or (II) and a second therapeutic agent, with instructions on coadministration of the two agents. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceutical products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

EXEMPLIFICATION

The compounds of this invention and their preparation can be understood further by the examples that illustrate some of the processes by which these compounds are prepared or used. It will be appreciated, however, that these examples do not limit the invention. Variations of the invention, now known or further developed, are considered to fall within the scope of the present invention as described herein and as hereinafter claimed.


The schemes listed herein are merely illustrative of some methods by which the compounds of this invention can be synthesized, and it is to be understood that various modifications to these schemes can be made.

Example 1: Synthesis of Compounds


General Reaction Procedures:

Unless mentioned specifically, reaction mixtures were stirred using a magnetically driven stirrer bar. An inert atmosphere generally refers to either dry argon or dry nitrogen. Reactions were monitored either by thin layer chromatography, by proton nuclear magnetic resonance (NMR) or by high-pressure liquid chromatography (HPLC), of a suitable sample of the reaction mixture.


Common abbreviations include m-CPBA: meta-chloroperbenzoic acid; DDQ: 2,3-Dichloro-5,6-dicyano-1,4-benzoquinone; DEAD Diethyl azodicarboxylate; DIBAL-H: Diisobutyl aluminum hydride; DMAP: N,N-Dimethylaminopyridine; DMF: N,N-Dimethylformamide; HMPA: Hexamethylphosphoramide; LDA: Lithium diisopropyl amide; LiHMDS: Lithium bis (trimethylsilyl) amide; PCC: Pyridinium chlorochromate; TBAF: Tetrabutylammonium fluoride; THF: Tetrahydrofuran; CH₂Cl₂ or DCM: methylene chloride; MPM: para-methoxy benzyl; Boc: tert-butyl oxy carbamate; TfOH: trifluoromethane sulfonic acid; Tf₂O: trifluoromethane sulfonic acid anhydride; TFA: trifluoroacetic acid; and TFAA: trifluoroacetic acid anhydride..

Synthesis of a common starting material**Compound 003:**

To a solution of trihydroxy-benzoic acid (120 g) in 350 mL of acetone, 500 mL of TFA (trifluoro acetic acid) was added at 40°C under stirring. After 1h at 40°C, 300 mL of TFAA (trifluoro acetic anhydride) was added. The mixture was heated for 3 days. The mixture was distilled under house vacuum at 50°C to remove solvents. The crude product was then diluted with 4L of CH₂Cl₂, washed with water, sat. NaHCO₃, dried and concentrated to give 85 g of semi pure solid **001**. The solid was crystallized in EtOH (1g/ 2mL) to give 20g of pure crystal. The mother liquor was then purified by silica gel with CH₂Cl₂ to 5% MeOH/CH₂Cl₂ to give 55g of additional product **001**.



To a solution of **001** (50g, 238 mmol) in 156 mL of pyridine, Tf₂O (100mL, 595 mmol, 2.5eq.) was added at 0°C in 3h. Then it was warmed to room temperature and stirred for 2h. The reaction mixture was diluted with water. The mixture was filtered, and the solid on the filter was washed with water, dried under vacuum to give 100g of solid **002**.




Ditriflate **002** (45.35g), BocNH₂ (17.22g), Pd₂(dba)₃ (4.38g) and *Pt*-Bu₃ (4.38g) were mixed in 150 mL of toluene. Tri-ethylamine (26.92 mL) was added to this mixture

and the reaction was heated under inert atmosphere at 80°C for 4h. The crude reaction mixture was cooled and filtered through a pad of celite. The filtrates were concentrated and purified on silica gel with Hex/EtOAc, 9:1, 4:1 to give 28.3 g of desired product, compound **003**.

Synthesis of common intermediates

Intermediates **009** and **010** were synthesized as illustrated in Scheme 1.



Scheme 1

Compound 004:

Compound **003** (18.6g, 42.1 mmol) in DMPU (42 ml) was cooled to 0°C. A solution of lithium hexamethyldisilylamide in THF (1.0 M, 47 ml, 47 mmol) was added slowly using a syringe pump maintaining internal temperature below 5°C. The resulting dark red-brown solution was stirred at 0°C for an additional 1 hr 20 min. TBSO-ethyl iodide (16.0g, 55.9 mmol) was added via canula over 10 min, while maintaining internal temperature below 3°C. The reaction mixture was stirred at 0°C for an

additional 10 min and subsequently stirred at room temperature over night. The reaction was quenched with saturated sodium bicarbonate (30 ml) and the solvent was removed under reduced pressure. The residue was purified by flash chromatography eluting with 5% EtOAc/Hexanes to obtain 18.5g (74%) pure desired product, compound **004**. ^1H -NMR (400 MHz, C_6D_6) δ : 7.20 (d, $J = 2.0$ Hz, 1H), 3.59 (t, $J = 5.2$ Hz, 2H), 3.41 (d, $J = 5.2$ Hz, 2H), 1.29 (s, 9H), 1.18 (s, 6H), 0.82 (s, 9H), -0.07 (s, 6H); MS m/e : ($M+23$) = 622.

Compound 006:

To **005** (13.0 g, 33.4 mmol) and **004** (18.6 g, 31.0 mmol) were added N-methylpyrrolidinone (30 mL), dicyclohexylmethylamine (8 ml, 37.4 mmol) and $\text{Pd}_2(\text{DBA})_3$ (8.51g, 9.29 mmol). The mixture was heated at 100°C for 2.5 hr and then at 110°C for 10 hr. The solvents were then removed under reduced pressure with heating. The residue was purified by flash chromatography eluting with a 10-17% gradient solvent of EtOAc/Hexanes to yield 24.7g (90%) of the compound **006**. ^1H -NMR (400 MHz, C_6D_6) δ : 8.10 (d, $J = 16.0$ Hz, 1H), 7.45 (d, $J = 2.0$ Hz, 1H), 7.30 (d, $J = 8.8$ Hz, 2H), 7.05 (d, $J = 2.0$ Hz, 1H), 6.92 (d, $J = 8.8$ Hz, 2H), 6.57-6.49 (m, 1H), 5.62-5.48 (m, 2H), 4.6 (d, 1H), 4.48 (m, 1H), 4.38-4.32 (m, 1H), 4.22 (d, 1H), 4.17-4.13 (m, 1H), 3.68 (m, 2H), 3.58 (m, 2H), 3.49-3.45 (m, 1H), 3.34 (s, 3H), 2.85-2.77 (m, 1H), 2.73-2.67 (m, 1H), 2.63-2.56 (m, 1H), 1.44 (s, 3H), 1.37 (s, 9H), 1.24 (s, 6H), 1.20 (s, 3H), 1.13 (d, $J = 6.0$ Hz, 3H), 1.03 (d, $J = 6.8$ Hz, 3H), 0.88 (s, 9H), -0.02 (s, 6H); MS m/e : ($M+23$) = 862.

Compound 007:

To **006** (12.0 g, 14.3 mmol) in THF (1.48 L) was added potassium hexamethyldisilylamide in toluene (0.5M, 30 mL, 6.0 mmol) slowly at room temperature via a syringe pump down wall over a period of 1.5 hr. The reaction was quenched with aqueous ammonium chloride solution (800 ml). The layers were separated and the aqueous layer was back extracted three times with MTBE (3x400mL) and three times with ethyl acetate (3x400mL). The combined extracts were dried with sodium sulfate and concentrated. The residue was purified by flash chromatography eluting with 5% then 10% EtOAc/Hexanes to give 5.99 g (54%) of compound **007** and 2.32 g (19%) of recovered starting material **006**. ^1H -NMR of **007** (400 MHz, C_6D_6) δ : 7.29 (d, $J = 2.4$

Hz, 1H), 7.19 (d, $J = 15.6$ Hz, 1H), 7.14 (d, $J = 2.4$ Hz, 1H), 6.93 (d, $J = 8.4$ Hz, 1H), 6.78 (d, $J = 8.4$ Hz, 1H), 5.93-5.86 (m, 1H), 5.67 (dd, $J = 8.8$ and 9.2 Hz, 1H), 5.46 (dd, $J = 10.4$ and 8.8 Hz, 1H), 5.22 (dd, $J = 9.6$ and 10.4 Hz, 1H), 4.88-4.82 (m, 1H), 4.42-4.36 (m, 1H), 4.25-4.19 (m, 2H), 3.80-3.76 (m, 1H), 3.72-3.60 (m, 4H), 3.29 (s, 3H), 3.11-3.02 (m, 1H), 2.83-2.74 (m, 1H), 2.64-2.57 (m, 1H), 1.42 (s, 3H), 1.35 (s, 9H), 1.23 (s, 3H), 1.04 (d, $J = 6.4$ Hz, 3H), 0.90 (s, 9H), 0.77 (d, $J = 7.2$ Hz, 3H), 0.01 (s, 6H); MS m/e : (M+23) = 804.

Compound 008:

To **007** (5.94 g, 7.60 mmol) in THF cooled at 0°C (ice/water bath) were added DBU (4.9 mL, 33 mmol) then methoxymethylchloride (2.3 mL, 30.3 mmol) in portions down wall. After stirring for 3hr, the reaction mixture was washed three times with aqueous ammonium chloride solution, water and brine. The aqueous layers were back extracted three times with MTBE. The combined extracts were dried with sodium sulfate and concentrated to get 6.27g (99%) of a white foam crude product which was clean and directly used for the next desilylation reaction. A small amount of crude product was purified by prep-TLC (eluting with 30% EtOAc/Hexanes) to verify the desired product, **008**. $^1\text{H-NMR}$ (400 MHz, C_6D_6) δ : 7.32 (d, $J = 8.8$ Hz, 1H), 7.16 (br s, Hz, 1H), 7.06 (s, 1H), 7.86 (d, $J = 8.4$ Hz, 2H), 6.82 (d, $J = 18.0$ Hz, 1H), 6.49-6.41 (m, 1H), 5.52-5.46 (m, 2H), 5.33 (dd, $J = 10.4$ and 10 Hz, 1H), 4.98 (dd, $J = 8.8$ and 8.8 Hz, 1H), 4.82 (s, 2H), 4.74 (d, $J = 10.8$ Hz, 1H), 4.58 (d, $J = 11.2$ Hz, 1H), 4.37 (dd, $J = 6.4$ and 6.4 Hz, 1H), 4.17 (dd, $J = 9.6$ and 7.2 Hz, 1H), 3.75-3.63 (m, 4H), 3.26 (s, 3H), 3.23-3.16 (m, 1H), 3.09 (s, 3H), 2.67-2.59 (m, 2H), 1.35 (s, 9H), 1.30 (d, $J = 6.8$ Hz, 3H), 1.27 (s, 3H), 1.11 (s, 3H), 0.95 (d, $J = 6.4$ Hz, 3H), 0.90 (s, 9H), 0.01 (s, 3H), 0.00 (s, 3H); MS m/e : (M+23) = 848.


Compound 009:

Compound **008** (6.27g) was dissolved in THF (230ml) in 1 L round bottom flask and cooled to 0°C. To this was added buffered TBAF (1.0 M TBAF and 0.5M imidazole/HCl in THF, 15.2ml) via a syringe pump over 26 min. After completion of addition, the mixture was stirred at room temperature for 12 hr. The reaction mixture was washed with saturated aqueous bicarbonate solution, water and brine sequentially. Aqueous phases were extracted with MTBE. The combined organic extracts were dried

with sodium sulfate and purified by chromatography to afford 5.08g (94%) of **009** as a colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 7.34-6.85 (m, 6H), 6.78 (d, $J = 16.0$ Hz, 1H), 6.43 (m, 1H), 5.52-5.46 (m, 2H), 5.33 (dd, $J = 10.4$, and 10.0 Hz, 1H), 4.96 (t, $J = 10.0$ Hz, 1H), 4.75 (s, 2H), 5.25 (d, $J = 11.2$ Hz, 1H), 4.57 (d, $J = 11.2$ Hz, 1H), 4.37 (dd, $J = 6.4$, and 6.4 Hz, 1H), 4.20-4.15 (m, 1H), 3.56-3.40 (m, 4H), 3.25 (s, 3H), 3.22-3.16 (m, 1H), 3.07 (s, 3H), 2.66-2.50 (m, 2H), 1.30 (d, $J = 7.2$ Hz, 3H), 1.29 (s, 9H), 1.27 (s, 3H), 1.11 (s, 3H), 0.94 (d, $J = 6.8$ Hz, 3H); MS, m/e ($m+23$)=734. 113mg (2%) of side product, Compound **023**, was also isolated and was used to synthesize Compound **024**.


Compound 010:

Into a round bottom flask were added **009** (500 mg, 0.702 mmol) and methylene chloride (14.0 mL) and the mixture was cooled to 0°C . Triethylamine (0.29 mL, 2.10 mmol) was added, then methanesulfonyl chloride (0.082 mL, 1.00 mmol) was added slowly. After 2hrs, TLC and MS check indicated the reaction was complete, clean and desired product was observed. The reaction mixture was poured into a saturated sodium bicarbonate solution, and extracted with DCM three times. The extracts were combined and dried over sodium sulfate, concentrated and then azeotroped with toluene. The colorless oil crude product, **010**, was reasonably clean and generally directly subjected to the nucleophiles in next step reaction without purification. Yield 555mg (100%). ^1H NMR (400 MHz, CDCl_3) δ 7.38-6.90 (m, 6H), 6.83 (d, $J = 16.4$ Hz, 1H), 6.53 (m, 1H), 5.57-5.50 (m, 2H), 5.38 (dd, $J = 10.0$, and 10.4 Hz, 1H), 5.00 (t, $J = 8.8$ Hz, 1H), 4.88 (AB, 2H), 4.78 (d, $J = 11.2$ Hz, 1H), 4.61 (d, $J = 11.2$ Hz, 1H), 4.42 (dd, $J = 6.8$, and 6.4 Hz, 1H), 4.24-4.20 (m, 1H), 4.02 (t, $J = 5.6$ Hz, 2H), 3.51 (t, $J = 5.6$ Hz, 2H), 3.30 (s, 3H), 3.25-3.21 (m, 1H), 3.12 (s, 3H), 2.70-2.53 (m, 2H), 2.20 (s, 3H), 1.38-1.33 (s and d overlapped, 12H), 1.33 (s, 3H), 1.16 (s, 3H), 0.98 (d, $J = 6.8$ Hz, 3H); MS, m/e ($M+23=812$).


Synthesis of Selected Compounds**Compound 011:**

To **010** (554 mg, 0.702 mmol) prepared as described above was added DMF (5.44 mL) and imidazole sodium derivative (1.26 g, 14.0 mmol). The mixture was stirred over night at room temperature. A check via TLC indicated that the reaction was complete. The solvent was removed under reduced pressure and the residue was partitioned between dichloromethane (DCM) and aqueous bicarbonate. The mixture was extracted with DCM (3x100ml) and the combined organic extracts were dried over sodium sulfate and concentrated. The residue was purified by flash chromatography eluting with 10-50% gradient of EtOAc/Hexanes and 0-7% MeOH/DCM to obtain 531mg (100%) of **011** as a white foam.


Compound **011** was then subjected to the sequential treatment of DDQ, Dess-Martin oxidation, to give compound **012** and TFA/water, and TfOH/water, similar to the synthesis of compound **091**, below, to obtain analog **013** as a white powder. Yield: 41mg (55%). ¹H NMR (400 MHz, 10% CD₃OD/CDCl₃) δ: 1.07 (d, 3H, J=8 Hz), 1.29 (d, 3H, J=8 Hz), 1.97 (m, 1H), 2.06 (m, 1H), 3.42 (m, 3H, J = 4 Hz), 3.85 (br s, 2H), 3.93 (m, 1H), 4.05 (t, 2H, 8 Hz), 4.39 (d, 1H J = 3 Hz), 4.78 (m, 1H, J = 3 Hz), 5.82 (m, 1H, 3 Hz), 5.89 (d, 1H, J = 3 Hz), 5.92 (d, 1H, J = 3 Hz), 6.02 (dd, 1H, J = 12 Hz), 6.13 (d, 1H, J = 12 Hz), 6.73 (d, 1H, J = 16 Hz), 6.88 (d, 1H, J = 16 Hz), 7.38 (s, 1H). MS m/e: (m+1) 456 (5%), (m+23) 478 (100%).

Compound 014:

Compound **014** was synthesized from compound **010** following the same procedures as the synthesis of compound **013**, except N-methylpiperidine was used in place of sodium imidazole. Yield 18mg (30%). ^1H NMR (400 MHz, CD_3OD) δ : 1.15 (d, 2H, $J = 6$ Hz), 1.36 (d, 2H, $J = 6$ Hz), 1.45 (t, 2H, $J = 6$ Hz), 1.97 (s, 1H), 2.08 (m, 1H), 2.70 (t, 2H, $J = 6$ Hz), 2.79 (s, 3H), 3.30 (m, 2H), 3.48 (m, 2H), 4.05 (m, 1H), 4.48 (d, 1H, $J = 4$ Hz), 5.95 (m, 2H), 6.12 (dd, 2H, $J = 8$ Hz), 6.32 (d, 1H, $J = 8$ Hz), 6.88 (d, 1H, $J = 16$ Hz). MS m/e : ($m+1$) 488 (100%).

Compound 015:


Compound **015** was synthesized from compound **010** following the same procedures as the synthesis of compound **013**, except morpholine was used in place of sodium imidazole. The yield was a white powder, 80 mg (76%). ^1H NMR (400 MHz, C_6D_6) δ : 0.78 (d, 3H, $J=8$ Hz), 0.93 (t, 3H, $J=8$ Hz), 1.00 (d, 3H, $J=8$ Hz), 1.94 (m, 6H), 2.13 (m, 2H), 2.61 (q, 2H, $J = 4$ Hz), 3.46 (t, 4H, $J = 4$ Hz), 3.94 (m, 1H), 4.30 (d, 1H, $J = 3$ Hz), 4.66 (m, 2H), 5.41 (t, 1H, $J = 8$ Hz), 5.55 (d, 1H, $J = 12$ Hz), 5.97 (m, 1H), 6.02 (d, 1H, $J = 4$ Hz), 6.22 (d, 1H, $J = 3$ Hz), 6.94 (d, 1H, $J = 16$ Hz). MS m/e : ($m+1$) 475 (100%), ($m+23$) 497 (9%).

Compound 016:

Compound **016** was synthesized from compound **010** following the same procedures as the synthesis of compound **013**, except pyrrolidine was used in place of


sodium imidazole. Yield 14.6mg (47%). ^1H NMR (400 MHz, CD_3OD) δ : 1.14 (d, 3H, $J = 3$ Hz), 1.27 (s, 2H), 1.34 (d, 3H, $J = 3$ Hz), 1.85, (m, 4H, $J = 3$ Hz), 2.08 (m, 2H), 2.72 (m, 4H), 2.79 (t, 2H, $J = 8$ Hz), 3.31 (t, 2H, $J = 8$ Hz), 3.46 (m, 1H), 4.04 (m, 1H), 4.48 (d, 1H, 3 Hz), 5.95 (m, 1H) 5.97 (d, 1H, $J = 4$ Hz), 6.09 (d, 1H, $J = 12$ Hz), 6.14 (d, 1H, $J = 4$ Hz), 6.30 (d, 1H, $J = 12$ Hz), 6.87 (d, 1H, $J = 16$ Hz). MS m/e : (m^{+1}) 459 (100).

Compounds 018 and 019:




Compound **017** was prepared from compound **010** following the same procedures as the synthesis of compound **013**, except 2,2-dimethyl-1,3-dioxolane-4-methanamine was used in place of sodium imidazole. **017** was subjected to sequential treatment with DDQ, PCC and TFA to obtain compound **018**. Yield 32mg. MS m/e : (m^{+1}) 579.

Compound **018** was then treated with TfOH, in the same manner as the synthesis of compound **013** to yield compound **019** (9.2mg, 23%). ^1H -NMR (400 MHz, CD_3OD) δ : 1.16 (d, 3H, $J = 6.73$ Hz), 1.36 (d, 3H, $J = 6$ Hz), 2.09 (m, 2H), 2.65 (s, 1H), 3.07 (m, 1H), 3.23 (m, 2H), 3.30 (m, 2H), 3.40 (m, 4H), 3.54 (m, 1H), 3.9 (m, 2H), 4.04 (m, 2H), 4.48 (d, 1H, $J = 2.4$ Hz), 5.58 (m, 1H), 6.06 (d, 1H, $J = 2.4$ Hz), 6.12 (m, 2H), 6.19 (d, 1H, $J = 2.4$ Hz), 6.32 (d, 2H, $J = 11.5$ Hz), 6.9 (d, 1H, $J = 16$ Hz). MS, m/e : ($m+23$) 479 (100%).

Compound 022:


Compound **020** was prepared from compound **010** following the same procedures as the synthesis of **017**, except 2-TBSO-ethanamine was used 2,2-dimethyl-1,3-dioxolane-4-methanamine. Compound **020** was then subjected to sequential treatments with Boc₂O, DDQ and PCC to offer **021** which was subsequently treated with TFA and TfOH in the same manner as the procedure for the synthesis of **019**. After reverse phase HPLC purification, compound **022** (5.2mg) was obtained as a TFA salt form. ¹H-NMR (400 MHz, CD₃OD) δ: 1.18 (d, 3H, J = 6.75 Hz), 1.41 (d, 3H, J = 6.15 Hz), 2.1 (m, 2H), 2.65 (s, 1H), 3.15 (t, 2H, J = 5.25 Hz), 3.22 (t, 2H, J = 6.75 Hz), 3.50 (m, 2H), 3.56 (m, 2H), 3.63 (m, 2H), 3.77 (m, 2H), 3.91 (s, 1H), 4.02 (m, 1H), 4.49 (t, 1H, J = 2.4 Hz), 4.93 (m, 1H), 6.01 (m, 1H), 6.13 (m, 2H), 6.23 (s, 1H), 6.32 (d, 1H, J = 11.5 Hz), 6.92 (d, 1H, J = 15.85 Hz). MS, m/e (m+1) 445 (100).

Compound 024:

Compound **023** (113mg) was obtained as a side product during the synthesis of **009**. Compound **023** was then subjected to the same process as the synthesis of compound **013** to obtain analog compound **024**. Yield 16.8mg (30% in 3 steps). ¹H-NMR (400 MHz, CDCl₃) δ: 7.18 (d, J = 2.4 Hz, 1H), 7.05 (d, J = 2.4 Hz, 1H), 6.96 (d, J = 15.2 Hz, 1H), 6.34 (d, J = 11.2 Hz, 1H), 6.17 (d, J = 10.4 Hz, 1H), 6.10 (m, 1H), 5.05-


4.94 (m, 1H), 4.50-4.46 (m, 3H), 4.10-4.05 (m, 3H), 3.56-3.49 (m, 1H), 2.16-2.10 (m, 1H), 1.40 (d, $J=6.4\text{Hz}$, 3H), 1.17 (d, $J=6.4\text{Hz}$, 3H); MS, m/e ($M+23$)=454, ($M-1$)=430.

Compound 025




To compound **010** (220mg, 0.28mmol) was added ammonia solution in methanol (2M, 100ml) and concentrated ammonia aqueous solution (22ml). The mixture was stirred at room temperature over night. The reaction mixture was poured into saturated bicarbonate solution (500ml) and extracted four times with DCM and once with ethyl acetate. The combined extracts were concentrated and the residue was purified by chromatography to obtain 100mg of urea compound (60%).


The urea compound was then subjected to the same process as in the synthesis of compound **013**, except PCC oxidation was applied in place of Dess-Martin oxidation. Compound **025** was obtained in 91% yield (49 mg). $^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 1.14 (d, 3H, $J = 8\text{ Hz}$), 1.38 (d, 3H, $J = 6\text{ Hz}$), 2.07 (m, 2H), 2.17 (m, 2H), 2.82 (s, 2H), 3.18 (s, 1H), 3.60 (m, 1H), 4.02 (m, 2H), 4.07 (m, 2H), 4.52 (m, 2H), 4.95 (m, 1H), 5.33 (s, 1H), 6.12 (m, 2H), 6.25 (d, 1H, $J = 12\text{ Hz}$), 6.89 (s, 1H), 6.93 (d, 1H, $J = 3\text{ Hz}$), 7.25 (d, 1H, $J = 3\text{ Hz}$). MS, m/e ($m+23$) 453 (100%).

Compound 029:

Compound **029** was synthesized from **008**. Compound **008** was subjected to the sequential treatment of DDQ, Dess-Martin oxidation, to give **027**. Similar to the synthesis of **013**, **027** was then treated with TFA/water and TfOH/water, to obtain analog **029** as a white powder. Yield: 40.1mg (45%). ¹H-NMR (400 MHz, CDCl₃) δ: 1.16 (d, 3H, J = 8 Hz), 1.20 (d, 1H, J = 4 Hz), 1.12 (s, 1H), 1.25 (t, 8H, J = 8 Hz), 1.39 (d, 3H, J = 8 Hz), 2.04 (s, 8H), 3.31 (t, 2H, J = 4 Hz), 3.54 (m, 1H), 3.82 (t, 2H, 4 Hz), 3.91 (m, 1H), 4.04 (m, 1H), 4.48 (s, 1H), 4.86 (m, 1H), 5.30 (s, 1H), 5.94 (m 1H), 6.03 (d, 1H, J = 4 Hz), 6.09 (d, 1H, J = 4 Hz), 6.16 (q, 2H, J = 8 Hz), 6.83 (d, 1H, J = 16 Hz). MS m/e: (m+23) 428 (100%).

Compound 034:

Compound **009** (310.0 mg, 0.0004355 mol), silver (I) oxide (202 mg, 0.000871 mol), and methyl iodide (542.2 μL , 0.008710 mol) were mixed in anhydrous acetonitrile (1.581 mL, 0.03027 mol) and the mixture was heated by microwaves in a sealed tube for 5 min intervals. Reaction was complete as shown by TLC (5% MeOH/DCM). The reaction mixture was filtered through celite 545 and the residue was purified by chromatography: 12g Redi-Sep pack Si-Gel column using 0-5% acetone/ DCM, resulted in incomplete separation, it was necessary to repurify on 2mm Prep Plate X2 using 5%acetone/DCM. This process was repeated to remove all of the side product, from our desired product. EtOAc/Hexane co-solvent helped remove all impurities from the desired product, **030**. Yield: 160mg (51%). Then, following the same procedures as the synthesis of **013**, compound **034** was obtained. Yield: 25mg (47%). $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 1.17 (d, 3H, $J = 8$ Hz), 1.26 (t, 1H, $J = 8$ Hz), 1.40 (d, 3H, $J = 8$ Hz), 2.05 (s, 1H), 2.14 (m, 2H), 2.62 (d, 1H, $J = 3$ Hz) 3.31 (t, 2H, $J = 3$ Hz), 3.38 (s, 3H), 3.58 (t, 3H, $J = 4$ Hz), 3.86 (d, 1H, $J = 4$ Hz), 4.01 (s, 1H), 4.12 (q, 1H, $J = 4$ Hz), 4.49 (m, 1H) 4.53 (br s, 1H), 4.88 (m 1H), 5.94 (m, 1H), 6.03 (d, 1H, $J = 4$ Hz), 6.09 (d, 1H, $J = 4$ Hz), 6.17 (m 2H), 6.84 (d, 1H, $J = 16$ Hz). MS m/e : ($m+1$) 420 (30%), ($m+23$) 442 (100%).

Compound 041:

009 (26.9 mg, 0.0378 mmol) and triphenylphosphine (16.8 mg, 0.0642 mmol) were dissolved in toluene (0.40 mL, 3.78 mmol). Methyl iodide (3.06 μ L, 0.049 mmol) and diethyl azodicarboxylate (6.54 μ L, 0.0416 mmol) was added at the same time in one shot. Reaction mixture was stirred for 1.5hrs and monitored by TLC at 15min intervals. Reaction was allowed to stir over night. TLC (50% EtOAc/hexane) the next morning showed a 1:1 mixture of product to starting material, this was confirmed by MS. Additional triphenylphosphine (16.8 mg, 0.0642 mmol), followed by methyl iodide (3.06 μ L, 0.0491 mmol) and diethyl azodicarboxylate (6.54 μ L, 0.0416 mmol) were added simultaneously. The reaction was monitored at 10min intervals, and still showed substantial amount of starting material. Additional triphenylphosphine (16.8 mg, 0.0642 mmol), followed by methyl iodide (3.06 μ L, 0.0491 mmol) and diethyl azodicarboxylate (6.54 μ L, 0.0416 mmol) were added simultaneously. The reaction was monitored at 10min intervals by TLC, and showed that it went to completion. Aqueous workup was performed and extracted with MTBE. The organic extracts were dried over sodium sulfate and concentrated. The residue was loaded onto a 4g Redi-sep column, and eluted


with 10% Ethanol/Hexanes. Compound **035** was isolated as a clear oil confirmed by TLC, MS and NMR. Yield: 31mg (100%)

An amino-polyether fragment was added to **035** to produce compound **036** using the same procedures as used in the synthesis of compound **013**. The product structure was verified by NMR, MS, and TLC. Yield: 150.2mg (61%).

Into an oven dried round bottom flask was added the **036** (174.5 mg, 0.0002036 mol), followed by tetrahydrofuran (1.651 mL, 20.36 mmol) and di-tert-butylidicarbonate (222.2 mg, 1.018 mmol) and 4-dimethylaminopyridine (4.97 mg, 0.0407 mmol) under an atmosphere of nitrogen. The reaction was stirred for 1 hour, and determined to be complete by TLC and MS. The reaction mixture was diluted with water and extracted with MTBE, dried over sodium sulfate and concentrated. The residue, compound **037**, was purified on red-sep column eluting with 0-5% MeOH/DCM. Yield: 162.3mg (83%).

The same procedures as used in the synthesis of compound **013** were performed with compound **037** in order to give the final polyether product **041**. Yield: 43.2mg (80%). ¹H-NMR (400 MHz, CDCl₃) δ : 1.15 (d, 3H, J = 7 Hz), 1.37 (d, 3H, J = 7 Hz), 2.05 (s, 1H), 2.07 (m, 2H), 3.05 (t, 2H, J = 3 Hz), 3.08 (t, 2H, J = 3 Hz), 3.36 (s, 3H), 3.38 (m, 2H), 3.54 (m, 2H), 3.64 (m, 8H), 3.69 (m, 2H), 4.03 (s, 1H), 4.18 (q, 1H, J = 7.1 Hz), 4.47 (d, 1H, J = 3 Hz), 4.84 (m, 1H), 5.40 (s, 1H), 5.99 (m, 1H), 6.00 (d, 1H, J = 3 Hz), 6.12 (m, 4H), 6.79 (d, 1H, J = 16 Hz). MS m/e: (m+1) 551 (100%), (m+23) 573 (10%).

Compound 042




To a 250 ml round bottom flask were added dichloromethane (4 ml) and oxalyl chloride solution in dichloromethane (2.0M, 2.1ml). The solution was cooled to -78°C and DMSO was added. After stirring for 5 min, **009** in dichloromethane (0.05M, 55 ml,

2.75 mmol) was added. After stirring additional 30 min, triethylamine (2.3ml) was added and the reaction mixture was allowed to warm up to 0°C over 45 min. The mixture was poured into saturated aqueous sodium bicarbonate solution. The layers were separated and the aqueous layer was back extracted three times using dichloromethane. The combined organic extracts were dried with sodium sulfate and concentrated.

The above crude aldehyde product (1.95g) was mixed with t-butyl alcohol (52 ml) in a 200 ml round bottom flask. To this were added 2-methyl-2-butene via a syringe and needle, sodium chlorite (2.5 g), potassium hydrogen phosphate (1.8 g) and water (25ml). The reaction mixture was stirred at room temperature for 1 hour. Aqueous sodium thiosulfate solution (150ml) and ethyl acetate (150ml) were added and the mixture was stirred for 1 hour. The layers were separated, and the aqueous layer was cooled to 0°C, acidified to a pH of about 3 using 1N HCl, and back extracted three times using ethyl acetate. The combined organic extracts were washed three times with distilled water, dried with sodium sulfate and concentrated to obtain 1.8g (90%) over 2 steps. Four analogs were synthesized from **042** as shown in the scheme above.


Compound **045**:



Compound **042** (316 mg), dimethylammonium-OOBt salt (370mg), HBTU (672mg), triethyl amine (0.43ml) and dichloromethane (3 ml) were added to a 10 ml round bottom flask and the mixture was stirred at room temperature over night. The reaction mixture was poured into a saturated aqueous sodium bicarbonate solution and extracted three times using dichloromethane. The combined extracts were dried with sodium sulfate and purified by Biotage chromatography to obtain 320 mg (98%) of **043**. Compound **043** was then subjected to the same procedures as the synthesis of compound **013** to obtain compound **045**. ¹H-NMR (400 MHz, CD₃OD) δ: 6.88 (d, *J* = 15.2 Hz,


1H), 6.32 (d, $J=11.6$ Hz, 1H), 6.19 (d, $J=2.4$ Hz, 1H), 6.12 (dd, $J=11.6$ and 10.0 Hz, 1H), 6.02-5.95 (m, 1H), 5.96 (d, $J=2.4$ Hz, 1H), 4.48 (d, $J=2.0$ Hz, 1H), 4.06-4.03 (m, 1H), 3.99 (s, 2H), 3.50-3.43 (m, 1H), 3.07 (s, 3H), 2.97 (s, 3H), 2.16-2.02 (m, 2H), 1.36 (d, $J=6.4$ Hz, 3H), 1.15 (d, $J=7.2$ Hz, 3H); MS m/e : ($M+23$) = 469.

Compound 046:




The synthesis of compound **046** was the same as the synthesis **045** except methylammonium-OOBt salt was used in place of dimethylammonium-OOBt. MS m/e : ($M+23$) = 455.

Compound 047:



The synthesis of compound **047** was the same as the synthesis of **045** except ammonium-OOBt salt was used in place of dimethylammonium-OOBt and pyridiniumchlorochromate (PCC) was used in place of Dess-Martin reagent. $^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 6.89 (d, $J=15.2$ Hz, 1H), 6.31 (d, $J=11.2$ Hz, 1H), 6.16 (d, $J=2.4$ Hz, 1H), 6.12 (dd, $J=11.2$ and 9.6 Hz, 1H), 6.02-5.94 (m, 1H), 5.92 (d, $J=2.4$ Hz, 1H), 4.48 (d, $J=2.0$ Hz, 1H), 4.06-4.03 (m, 1H), 3.77 (s, 2H), 3.50-3.43 (m, 1H), 2.16-2.02 (m, 2H), 1.35 (d, $J=6.0$ Hz, 3H), 1.15 (d, $J=7.2$ Hz, 3H); MS m/e : ($M+23$) = 441.


Compound 048:



Compound **042** was subjected to the same procedure as for the synthesis of compound **013** to yield 8.5mg (58%) of compound **048**. $^1\text{H-NMR}$ (400 MHz, CD_3OD)

δ : 1.16 (d, 3H, $J = 6.75$ Hz), 1.36 (d, 3H, $J = 5.86$ Hz), 2.0 (s, 2H), 2.10 (m, 2H), 3.3 (m, 1H), 3.47 (m, 1H), 3.64 (s, 1H), 3.87 (s, 2H), 4.05 (m, 1H), 4.09 (d, 1H, $J = 7$ Hz), 4.12 (m, 1H), 4.48 (d, 1H, $J = 2.40$ Hz), 5.91 (d, 1H, $J = 2.40$ Hz), 5.97 (m, 1H), 6.10 (m, 1H), 6.12 (m, 1H), 6.16 (d, 1H, $J = 3$ Hz), 6.32 (d, 1H, $J = 12$ Hz), 6.89 (d, 1H, $J = 15.5$ Hz). MS m/e : (m-1) 418 (100%).

Compound 054:




Compound **009** (200 mg, 0.00028 mol) was dissolved in methylene chloride (1.44 mL, 0.0225 mol). The diallyldiisopropylphosphoramidite (172 mg, 0.000702 mol), and pyridinium trifluoroacetate (136 mg, 0.000702 mol) was added. The reaction mixture was stirred at 20-25°C. The reaction was cooled to about 0-5°C and 30% H_2O_2 in water (3:7, H_2O_2 :water, 0.14 mL) was added. After 15 min TLC (MTBE) showed disappearance of intermediate. Into the reaction was added water (1.98 mL, 0.1101 mol) and the product was extracted three times with MTBE. The organic phase

was dried and then concentrated on rotavap to a crude colorless film. The product, compound **049**, was purified on a 4g redi-sep column eluting with 0-10% acetone/DCM. Yield: 242mg (99%). DDQ Deprotection of **049** was the same as in the synthesis of compound **013** affording 201mg (97%) of compound **050**.

Into a round bottom flask under an atmosphere of nitrogen, compound **050** (100.5 mg, 0.134 mmol) was dissolved in tetrahydrofuran (4 mL) was added, followed by the addition of tetrakis(triphenylphosphine) palladium(0) (31 mg, 0.027 mmol), silane, phenyl- (99.0 μ L, 0.000802 mol) and triphenylphosphine (42 mg, 0.00016 mol). The mixture was stirred for 60 minutes, and the reaction was monitored at 15 minute intervals until TLC (1:1 Maigic) showed that the reaction was complete. Solvent was removed under vacuum. The residue was dissolved in 2:3:1 solvent system and loaded onto DEAE resin in a 1" diameter column. The resin was flushed with the 2:3:1 solvent system to remove all catalyst and non-polar materials. After the filtrate turned clear, 0.04 M ammonium acetate was used to continue washing the resin and elute the product, compound **051**. The concentration of ammonium acetate was increased to 0.1M to drive off the remaining product from the resin. Fractions containing the product were collected and concentrated to a residue; the flask was placed under high vacuum and heated to 40°C to remove all traces of ammonium acetate. Yield: 94.3mg (100%).

The remainder of the procedure for the synthesis of compound **054** was similar to the analog compound **013**, except that the workup and purification for 054 was accomplished with the DEAE Resin as shown above for compound **051**. ^1H -NMR, ^{31}P -NMR, and MS verify structure. Product **054** submitted in free form. Yield 7.0mg (30%); ^1H -NMR (400 MHz, CD_3OD) δ : 1.15 (d, 3H, $J = 6.75$ Hz), 1.36 (d, 3H, $J = 5.9$ Hz), 2.10 (m, 2H), 3.3 (m, 2H), 3.45 (m, 1H), 4.01 (q, 2H, $J = 6$ Hz), 4.06 (m, 2H), 4.48 (d, 1H, $J = 3$ Hz), 5.93 (d, 1H, $J = 3$ Hz), 5.98 (m, 1H),), 6.12 (m, 1H), 6.18 (d, 1H, $J = 3$ Hz), 6.32 (s, 1H, $J = 12$ Hz), 6.87 (s, 1H, $J = 16$ Hz). ^{31}P NMR (400 MHz, CD_3OD) δ : 2.60 (s, 1P). MS m/e : (m^{-1}) 484


Compound 065:

Compound **055** was synthesized in a manner similar to the synthesis of compound **006**, shown above. To a solution of **055** (5.3 g, 7.95 mmol) and 2,6-lutidine (2.3 ml, 19.87 mmol) was added TBSOTf (2.19 ml, 9.54 mmol) dropwise at room temperature. The reaction mixture was quenched with aqueous sodium bicarbonate and the aqueous was extracted with ethyl acetate. The organics were combined and dried over Na₂SO₄. The dried solution was filtered and concentrated *in vacuo*. Purification of the residue by flash column chromatography on SiO₂ eluted with 5-10% ethyl acetate in hexane gave 5.26 g of desired product, compound **056**.

To **056** (5.26 g, 6.73 mmol) in methanol (100 ml) was added K₂CO₃ (1.86 g) at room temperature. The reaction mixture was stirred for 2 hours and concentrated in vacuo. Purification of the residue by flash column chromatography on SiO₂ eluted with 10% ethyl acetate in hexane gave 2.48 g of desired product **057**.

To **057** (2.48 g, 3.56 mmol) and pyridine (0.86 ml) in methylene chloride (20 ml) was added Tf₂O (0.72 ml) dropwise at 0°C. The reaction mixture was allowed to warm to room temperature and stirred for 2 hours. The reaction was quenched with aqueous NaHCO₃. The aqueous layer was extracted with ethyl acetate and dried with MgSO₄. The dried solution was filtered and concentrated in vacuo. Purification of the residue by

flash column chromatography on SiO₂ eluted with 5-10% ethyl acetate in hexane gave 2.77 g of desired product **058**.



Compound **058** (1.74 g, 2.1 mmol), compound **059** (0.62 g) and (rac)-BINAP (196.1 mg) were weighed into a 100 ml round bottom reaction flask and azeotropically dried with toluene. Pd(OAc)₂ (47.1 mg) and Cs₂CO₃ (821.1 mg) were weighed in dry box to the reaction flask. The flask was removed from dry box and added de-gassed toluene (20 ml). The reaction mixture was heated to 80°C for 15 hours and concentrated in vacuo. Purification of the residue by flash column chromatography on SiO₂ eluted with 5-10% ethyl acetate in hexane gave 1.64 g of desired product **060**.

The buffered TBAF was prepared by dissolving imidazole HCl (659.0 mg) in TBAF (1.0 M in THF, 12.6 ml). The solution of **060** (1.67 g) in buffered TBAF was heated to 60°C and stirred for 60 hours. The reaction was quenched by aqueous NH₄Cl. The aqueous was extracted with ether. The organics was combined and dried with

Na₂SO₄. The dried solution was filtered and concentrated in vacuo. Purification of the residue by flash column chromatography on SiO₂ eluted with 10-30% ethyl acetate in hexane gave 0.80 g of desired product **061**.

To the solution of t-BuOK(1.31 ml, 1.0 M in THF) in THF (9 ml) was added compound **061** (710 mg, 0.87 mmol) in THF (3 ml) slowly at 0°C and then rinsed with 2 ml of THF. After the addition of starting material was completed, TBSCl (138.7 mg) in THF (2ml) was added at 0°C. The reaction was quenched in 20 min and the aqueous was extracted with AcOEt. The organics was combined and dried over Na₂SO₄. The dried solution was filtered and concentrated in vacuo. Purification of the residue by flash column chromatography on SiO₂ eluted with 10% ethyl acetate in hexane gave 401.3 mg of desired product, compound **062**.


To **062** (401.3mg, 0.46 mmol) in methylene chloride and water (10 ml, 4:1 mixture) was added DDQ (128.0 mg) and stirred at room temperature for 2 hours. The reaction was quenched with aqueous Na₂SO₃ and NaHCO₃. The aqueous material was extracted with AcOEt. The organics were combined and dried over Na₂SO₄. The dried solution was filtered and concentrated in vacuo. Purification of the residue by flash column chromatography on SiO₂ eluted with 10% ethyl acetate in hexane gave 335.4 mg of desired product **063**.

To **063** (335.4 mg, 0.45 mmol) and pyridine (54.4 µl) in methylene chloride (8 ml) was added Dess-Martin periodinane (259.2 mg) at room temperature. The reaction mixture was stirred for 1 hour and quenched with aqueous NaHCO₃. The aqueous material was extracted with AcOEt. The organics were combined and dried over Na₂SO₄. The dried solution was filtered and concentrated in vacuo. Purification of the residue by flash column chromatography on SiO₂ eluted with 10% ethyl acetate in hexane gave 232.5 mg of desired product **064**.

To **064** (47.1 mg, 0.063 mmol) in toluene/water (1 ml/0.035 ml) was added CF₃SO₃H (16.8 µl) at 0°C. The reaction mixture was stirred for 1.5 hours and warmed to room temperature. Additional TfOH (16.8 µl) was added and the reaction was quenched in 5 minutes with saturated aqueous NaHCO₃. The aqueous was extracted with AcOEt. The organics was combined and dried over Na₂SO₄. The dried solution was filtered and concentrated in vacuo. Purification of the residue by flash column chromatography on SiO₂ eluted with 10% methanol in methylene chloride gave 7.4 mg of desired product **065**. ¹HNMR (CD₃OD): 6.92, (d, *J* = 15.6 Hz, 1H), 6.36, (s, 1H), 6.35 (s, 1H), 6.34 (d, *J*

= 11.6 Hz, 1H), 6.32 (d, J = 15.2 Hz), 6.13 (dd, J = 11.6, 10.0 Hz, 1H), 6.04-5.96 (m, 1H), 4.96-4.86 (m, 1H), 4.48 (d, J = 2.0 Hz, 1H). 4.08-4.02 (m, 1H), 3.95-3.87 (m, 2H), 3.86-3.79 (m, 1H), 3.64-3.53 (m, 2H), 3.52-3.43 (m, 1H), 2.18-2.02 (m, 2H), 1.38 (d, J = 6.4 Hz, 3H), 1.16 (d, J = 7.2 Hz, 3H).

Compound 069:



Into a round bottom flask was added **057** (2.331 g, 0.003345 mol), triphenylphosphine (1.14 g, 0.00435 mol) and tetrahydrofuran (13.6 mL, 0.167 mol). The mixture was cooled to 0°C under a nitrogen atmosphere. Imidazole ethanol fragment (0.450 g, 0.00401 mol) was added, followed by addition on diethyl azodicarboxylate (0.685 mL, 0.00435 mol). The reaction was allowed to warm slowly to room temperature while stirring. A TLC test revealed an intense product spot together with some unreacted starting material. The reaction was cooled to 0°C and an additional 0.2eq of the amine, triphenylphosphine, and DEAD were added. The ice bath


was removed and reaction was allowed to warm to room temperature. After 30min, TLC showed the reaction was complete and all starting material was consumed. All solvent was removed under high vacuum. The residue was dissolved in MTBE, the mixture was cooled in an ice bath and some of the triphenyl phosphate was filtered off. The residue was purified on 40g column, using 0-5% MeOH/DCM. Product fractions containing compound **066** were combined and concentrated down to a residue. Yield: 3.65g(100%).

The TBS protecting group was removed using buffered TBAF with imidazole hydrochloride, for a total compound **067** yield of ~2.0g (90%).

Compound **068** was produced from **067** using the same procedure as used in the synthesis of compound **007**. Combining all batches, the total yield was 467mg (50%).

The intermediate compound **068** was deprotected with DDQ, then oxidized with Dess-martin reagent and deprotected with TfOH to give the product, compound **069** (100mg). ¹H-NMR (400 MHz, CDCl₃) δ 1.19 (d, J=7.2Hz, 3H), 1.43 (d, J=6.0Hz, 3H), 2.10-2.25 (m, 2H), 3.58-3.65 (m, 1H), 4.00-4.03 (m, 1H), 4.22 (t, J=5.2Hz, 2H), 4.34 (t, J=5.2Hz, 2H), 4.50 (d, J=2.4, 1H), 4.88-4.95 (m, 1H), 5.96-6.03 (m, 1H), 6.14 (dd, J=9.6, 11.2Hz, 1H), 6.22 (d, J=11.6, 1H), 6.33 (d, J=2.4Hz, 1H), 6.37 (d, J=2.8Hz), 6.86 (d, J=15.6Hz, 1H), 7.03 (s, 1H), 7.08 (s, 1H), 12.17 (s, 1H).


Compound 076:



Compound **058** as synthesized above (459.7 mg, 0.55 mmol), compound **070** (104.6 mg) and (rac)-BINAP (51.8 mg) were weighed into a 25 ml round bottom reaction flask and azeotropically dried with toluene. Pd(OAc)₂ (12.4 mg) and Cs₂CO₃ (216.8 mg) were weighed in dry box to the reaction flask. The flask was removed from dry box and added de-gassed toluene (3 ml). The reaction mixture was heated to 80°C for 60 hours and concentrated in vacuo. Purification of the residue by flash column chromatography on SiO₂ eluted with 5-10% ethyl acetate in hexane gave 247.5 mg of desired product, compound **071**.

Buffered TBAF was prepared by dissolving imidazole HCl (108.3 mg) in TBAF (1.0 M in THF, 2.1 ml). The solution of **071** (247.5 mg, 0.3 mmol) in buffered TBAF was heated to 60°C and stirred for 36 hours. The reaction was quenched by aqueous NH₄Cl. The aqueous material was extracted with ether. The organics were combined and dried with Na₂SO₄. The dried solution was filtered and concentrated in vacuo. Purification of the residue by flash column chromatography on SiO₂ eluted with 10-20% ethyl acetate in hexane gave 116.6 mg of desired product, compound **072**.

To a solution of t-BuOK (0.24 ml, 1.0 M in THF) in THF (2 ml) was added **072** (116.6 mg, 0.16 mmol) in THF (1 ml) slowly at 0°C and rinsed with 2 ml of THF. After the addition of starting material was completed, TBSCl (25.6 mg) in THF (2 ml) was added at 0°C. The reaction was quenched in 20 min and the aqueous material was extracted with AcOEt. The organics were combined and dried over Na₂SO₄. The dried solution was filtered and concentrated in vacuo. Purification of the residue by pTLC developed with 30% ethyl acetate in hexane gave 40.0 mg of desired product, compound **073**.




To compound **073** (40.0 mg, 0.051 mmol) in methylene chloride and water (2 ml, 4:1 mixture) was added DDQ (14.2 mg) and stirred at room temperature overnight. The reaction was quenched with aqueous Na_2SO_3 and NaHCO_3 . The aqueous was extracted with AcOEt. The organics was combined and dried over Na_2SO_4 . The dried solution was filtered and concentrated in vacuo. Purification of the residue by pTLC developed with 20% ethyl acetate in hexane gave 25.0 mg of desired product, compound **074**.

To compound **074** (25.0 mg, 0.038 mmol) and pyridine (6.1 μl) in methylene chloride (2 ml) was added Dess-Martin periodinane (31.0 mg) at room temperature. The reaction mixture was stirred for 5 h and quenched with aqueous NaHCO_3 . The aqueous was extracted with AcOEt. The organics was combined and dried over Na_2SO_4 . The dried solution was filtered and concentrated in vacuo. Purification of the residue by pTLC developed with 30% ethyl acetate in hexane gave 24.0 mg of desired product, compound **075**.

To compound **075** (24.0 mg, 0.037 mmol) in toluene/water (1 ml/0.035 ml) was added $\text{CF}_3\text{SO}_3\text{H}$ (19.4 μl) at 0°C . The reaction mixture was stirred for 10 min and warmed to room temperature. The reaction was quenched in 1 h with saturated aqueous NaHCO_3 . The aqueous material was extracted with AcOEt. The organics were combined and dried over Na_2SO_4 . The dried solution was filtered and concentrated *in vacuo*. Purification of the residue by pTLC developed with 10% methanol in methylene chloride gave 2.4 mg of desired product, compound **076**. ^1H NMR (CDCl_3): 6.86 (d, $J = 15.2$ Hz, 1H), 6.29 (d, $J = 2.8$ Hz, 1H), 6.26-6.12 (m, 3H), 5.99-5.90 (m, 1H), 4.94-4.84 (m, 1H), 4.56-4.46 (m, 2H), 4.06-3.98 (m, 1H), 3.84 (d, $J = 5.6$ Hz, 1H), 3.65-3.54 (m,

1H), 2.50-2.42 (m, 2H), 2.24-2.08 (m, 2H), 1.42 (d, J = 6.0 Hz, 3H), 1.26 (t, J = 7.2 Hz, 1H), 1.18 (d, J = 6.8 Hz, 3H), 0.82-0.76 (m, 2H), 0.56-0.51 (m, 2H).


Compound 081:




DEAD was added to a solution of compound **077a**, Ph_3P and 2-TMS-ethanol in toluene at 0°C . The mixture was stirred at room temperature for 1 hour before it was quenched with aqueous NaHCO_3 . The organic phase was dried (Na_2SO_4), concentrated and purified by silica gel chromatography to afford **077b**.

MOM-Cl (2.8 mL, 37 mmol) was added to a solution of **077b** (2.1 g, 7.4 mmol) and DBU (7.1 mL, 48 mmol) in 10 mL of CH_2Cl_2 at 0°C . The mixture was stirred at room temperature for 15 minutes before the organic phase was washed with NaHCO_3 , concentrated and purified by silica gel chromatography to furnish compound **078** (1.8 g, 5.5 mmol) from a silica gel plug in 74% yield.

2.5 M $n\text{-BuLi}$ in hexanes was introduced drop wise to the stirring solution of diisopropylamine in of THF at -5°C . The solution was stirred at 0°C for 30 minutes before it was cooled to -78°C . Compound **078** in THF was added to the cold LDA slowly so that the internal temperature was kept below -78°C . The mixture was stirred at -78°C for 45 minutes, and then $(\text{PhSe})_2$ in THF was added slowly so that the internal temperature was kept below -60°C . The reaction mixture was stirred for 40 minutes before it was quenched with aqueous NH_4Cl at -78°C . EtOAc was added to the mixture at room temperature. After separation, the organic layer was dried (Na_2SO_4) and concentrated. 5% EtOAc in toluene was used to elute **079** from a silica gel column.



2.5 N NaOH was added to **079** in EtOH. The mixture was refluxed for 12 hours before it was acidified and extracted with EtOAc. Organic phase was dried (Na₂SO₄) and concentrated to afford solid **080**. DEAD was added to a solution of **080**, Ph₃P and 2-TMS-ethanol in toluene at 0°C. The mixture was stirred at room temperature for 1 hour before it was quenched with aqueous NaHCO₃. The organic phase was dried (Na₂SO₄), concentrated and purified by silica gel chromatography to afford compound **081**.

Compound 087:

LiHMDS in THF was introduced drop wise into a cold (-78°C) solution of compounds **081** and **082** in 10:1 THF-HMPA. The internal temperature was kept below -70°C during the addition. The mixture was stirred at -78°C for about 30 minutes before it was quenched with aqueous NH_4Cl and diluted with EtOAc. Organic phase was dried (Na_2SO_4), concentrated and purified by silica gel chromatography to afford **083** (1.9 g, 1.6 mmol) in 82% yield.


m-CPBA (1.0 g, 4.2 mmol) was added in three portions into a cold (0°C) solution of **083** in CH_2Cl_2 . The mixture was stirred at 0°C for 1 hour before the addition of Et_3N . After the reaction mixture was stirred at room temperature for 1 hour, it was quenched with aqueous $\text{Na}_2\text{S}_2\text{O}_3$ and diluted with aqueous NaHCO_3 . The organic phase was dried (Na_2SO_4), concentrated and purified by silica gel chromatography to afford compound **084**.

In a solution of TBAF buffered with 0.33 mole equivalent of imidazole•HCl was introduced to the solution of **084** in THF. The mixture was stirred at 50°C for 12 hours before it was diluted with Et_2O and washed with aqueous NH_4Cl . Organic phase was

dried (Na_2SO_4) and concentrated to furnish crude **085**, which was dissolved in CH_2Cl_2 and was added drop wise to a refluxing mixture of 2-chloro-1-methylpyridinium iodide and $n\text{-Bu}_3\text{N}$ in of CH_2Cl_2 . The mixture was refluxed for 2 hours before it was diluted with Et_2O and washed with 0.05 N HCl , H_2O and NaHCO_3 . Organic phase was dried (Na_2SO_4), concentrated and purified by silica gel chromatography to afford compound **086**.

TBAF in THF (7.0 mL, 7.0 mmol) was introduced to the solution of **086** (0.87 g, 1.3 mmol) in 7 mL of THF. The mixture was stirred at room temperature for 4 hours before it was diluted with Et_2O and washed with brine. Organic phase was dried (Na_2SO_4), concentrated and purified by silica gel chromatography to afford compound **087** (0.43 g, 0.78 mmol) in 58% yield over 3 steps.

Compound 091:



Tf_2O (0.42 ML, 2.5 mmole) was added to a solution of **087** (0.95 g, 1.7 mmole) and Et_3N (0.58 ML, 4.2 mmole) in 20 ML of CH_2Cl_2 at 0°C . The mixture was stirred for 10 min before the addition of aqueous NaHCO_3 . The aqueous layer was extracted

twice with CH_2Cl_2 . The organic phase was concentrated and passed through a short plug of silica gel (20% EtOAc/Hex).

The triflate thus obtained was added $\text{Pd}(\text{OAc})_2$ (19 mg, 0.08 mmole), BINAP (64 mg, 0.10 mmole) and Cs_2CO_3 (0.66 g, 2.0 mmole) in a dry box. Benzophenone imine (0.32 mL, 1.9 mmole) and 30 mL of toluene was added under nitrogen before the mixture was heated at 90°C for 14 h. The mixture was then diluted with EtOAc and brine. The organic layer was dried (Na_2SO_4) and concentrated.

The resultant crude material was dissolved in 8 mL of MeOH and 5 mL of THF before the addition of NaOAc (0.56 g, 6.8 mmole) and $\text{NH}_2\text{OH}\cdot\text{HCl}$ (0.24 g, 3.4 mmole) at room temperature. After 50 min, EtOAc and brine was added. The organic layer was dried (Na_2SO_4), concentrated and purified with silica gel (30% EtOAc/Hex) to produce crystalline compound **088** (0.88 g, 1.6 mmole).


LiHMDS (1N in THF, 8.0 mmole) was added slowly to a solution of **088** (0.88 g, 1.6 mmole) in 16 mL of THF at -55 to -50°C . The mixture was stirred at -45°C for 5 min before the addition of BOC_2O (0.38 mL, 1.8 mmole). After the mixture was stirred at -40°C for 30 min, MeI (0.60 mL, 9.6 mmole) was added. After 10 minutes, the mixture was warmed up to room temperature for 2 hours. The mixture was recooled to -35°C and 72 mL of 1N NaOH and 48 mL of EtOH were added. After it was heated at 45°C for 12 hours, the mixture was diluted with 100 mL of water and 150 mL of CH_2Cl_2 . The aqueous layer was extracted twice with 50 mL of CH_2Cl_2 . The organic layer was concentrated and purified by silica gel chromatography (30% EtOAc/Hex) to furnish colorless gel **089** (0.58 g, 1.0 mmole).

A suspension of compound **089** (0.40 g, 0.71 mmole), PCC (0.46 g, 2.1 mmole), 4A molecular sieves (0.50 g), and celite (0.50 g) in 8 mL of CH_2Cl_2 was stirred at room temperature for 2.5 h before the addition of Et_3N (0.29 mL, 2.1 mmole). After 5 min, 30 mL of Et_2O was added and the mixture was filtered. The filtrate was concentrated and passed through a short silica gel plug (75% EtOAc/Hex) to provide colorless crystalline **090** (0.35 g, 0.63 mmole).


TFA (5% water, 6 mL) was added slowly to the solution of compound **090** (0.35 g, 0.63 mmole) in CH_2Cl_2 at -35°C . The mixture was stirred at -20°C for 1 hour before the addition of saturated aq. NaHCO_3 (PH ~ 8) and CH_2Cl_2 . The aqueous layer was extracted twice with CH_2Cl_2 . The organic layer was dried (Na_2SO_4), concentrated and purified by silica gel chromatography (75% EtOAc/Hex) to afford compound **091**

(124 mg, 0.33 mmole) in 25% overall yield over 8 steps. NMR: $^1\text{H-NMR}$ δ (Benzene- d_6) 0.80 (d, $J = 6.8$ Hz, 3H, C4- CH_3), 1.02 (d, $J = 6.0$ Hz, 3H, C3- CH_3), 2.07 (d, $J = 5.1$ Hz, 3H, NCH_3), 2.11 (m, 2H, 10- CH_2), 2.26 (d, $J = 11.3$ Hz, 1H, C9-OH), 3.06 (m, 1H, NH), 3.44 (m, 1H, 4-CH), 3.82 (d, $J = 5.1$ Hz, 1H, C8-OH), 3.89 (ddt, $J_1 = 11.3$ Hz, $J_2 = 2.8$ Hz, $J_3 = 7.0$ Hz, 1H, 9-CH), 4.22 (dd, $J_1 = 2.6$ Hz, $J_2 = 5.1$ Hz, 1H, 8-CH), 4.67 (dq, $J_1 = 6.0$ Hz, $J_2 = 10.4$ Hz, 1H, 3-CH), 5.40 (dd, $J_1 = 9.6$ Hz, $J_2 = 11.3$ Hz, 1H, 5-CH), 5.48 (d, $J = 11.3$ Hz, 1H, 6-CH), 5.85 (d, $J = 2.3$ Hz, 1H, aromatic CH), 6.00 (dt, $J_1 = 15.2$ Hz, $J_2 = 7.5$ Hz, 1H, 11-CH), 6.13 (d, $J = 2.3$ Hz, 1H, aromatic CH), 6.98 (d, $J = 15.4$ Hz, 1H, 12-CH), 13.03 (s, 1H, phenol). MS: 398 (M^+Na , 100%).


Compound 092:



To a solution of compound **091** (16 mg) in CH_3CN was added DMAP (20 mg). The solution was then stirred at 60°C for 3 hours and then at room temperature for 12 hours. The crude mixture was purified on Silica gel and desired product **092** (13 mg) was eluted with 60% EtOAc/Hexanes. $^1\text{HNMR}$ (Benzene- d_6): 6.79 (2H, dd, $J_1=16.0$ Hz, $J_2=5.95$ Hz); 6.09 (1H, d, $J=2.3$ Hz); 5.92 (1H, dd, $J_1=16.0$ Hz, $J_2=1.4$ Hz); 5.82 (1H, m); 5.81 (1H, d, $J=2.3$ Hz); 4.98 (1H, m); 4.42 (1H, m); 3.97 (1H, m); 3.80 (1H, d, $J=5.7$ Hz); 3.04 (1H, m); 2.37 (2H, m); 2.04 (3H, d, $J=5.7$ Hz); 2.04 (1H, m); 1.67 (1H, m); 0.86 (3H, d, $J=7.1$ Hz); 0.56 (3H, d, $J=7.1$ Hz). Note that the coupling constants of the 2 enone olefin hydrogens changed from 11 Hz in **091** to 16 Hz in **092**.



yield).




To a solution of $(\text{COCl})_2$ (25 mL, 295 mmol, 2 eq.) in CH_2Cl_2 (870 mL), DMSO (41.85 mL, 590 mmol, 4 eq.) was slowly added at -78°C . After stirring for 30 min at -78°C , a solution of **096** (75 g, 147.4 mmol) in CH_2Cl_2 (160 mL) was added over 45 minutes. After the mixture was stirred at -78°C for 45 min, Et_3N (82.2 mL, 590 mmol, 4 eq.) was added at the same temperature. After the reaction is further stirred for 30 min, it was warmed to 0°C for 1.5 h. The reaction was quenched with 750 mL of saturated NH_4Cl and extracted three times with EtOAc. The extracts were dried and concentrated. The crude product was re-suspended with 2.5 L of 1:1 solution of EtOAc/Hexanes, washed three times with water and once with brine, and then dried and concentrated. The crude product **097** was used directly for next step.

To a suspension of $\text{Ph}_3\text{PCH}_2\text{Br}$ (115.8 mL, 324.3 mmol, 2.2 eq.) in a mixture of THF (870 mL) and DMSO (433.6 mL), *n*-BuLi (184.3 mL of 1.6 M solution, 294.8 mmol, 2 eq.) was added at 0°C . After stirred for 1h, a solution of **097** (74.7 g in 175 mL of THF, 147.4 mmol) was added at 0°C . After 30 min, it was warmed to room temperature. After 2 hours, it was quenched with 1.1 L of saturated NH_4Cl and extracted three times with EtOAc. The extracts were washed with water and brine and then dried and concentrated. The crude product was purified on silica gel with 5-10% EtOAc/Hexanes to give 66.5g of **098** as an oil (89% yield).

To a mixture of **098** (2.5 g, 4.95 mmol) and triflate **003** (2.7g, 6.4 mmol, 1.3 eq.),

$\text{Pd}_2(\text{dba})_3$ (1.36 g, 1.48 mmol, 0.3 eq.) was added in the dry box. After the mixture was moved out of dry box, it was suspended in 8.3 mL of dioxane and N-methyl N-dicyclohexane amine (2.1 mL, 9.9 mmol, 2eq.) was added. The reaction was heated at 100°C for 12 hours with vigorous stirring. After cooled, 6 g of celite was added and diluted with EtOAc. The mixture was filtered through a celite plug and rinsed with EtOAc. The filtrates were concentrated. The crude product was purified on silica gel with 10-20% EtOAc/Hexanes to give 3g of pure **099** (76% yield).




To a solution of **099** (46.3 g, 58.16 mmol) in DMPU (291 mL), LiHMDS (116 mL of 1M solution in THF, 116.3 mmol, 2 eq.) was added at 0°C. After the mixture was stirred for 40 minutes at the same temperature, EtI (27.9 mL, 349 mmol, 6 eq.) was added. After 5 minutes, the mixture was warmed to room temperature. After the mixture was stirred for 3 hours, it was quenched with 1L of Sat. NH_4Cl at 0°C. The mixture was extracted with MTBE/Hexanes (1:1). The extracts were washed with brine, and then dried and concentrated. The crude product was purified on silica gel with 15-20% EtOAc/Hexanes to give 40 g of desired product, **100** (84% yield).

To a solution of **100** (48 g, 58.2 mmol) in 230 mL, a solution of TBAF (407 mL of 1M solution, 407 mmol, 7 eq.) and imidazole.HCl (21.3 g, 203.9 mmol, 3.5 eq.) was added. The reaction was heated at 60°C for 72 hours. After the mixture was cooled to room temperature, it was quenched with saturated NH_4Cl and was extracted three times

with ether. The organic layers were washed with water and brine, and then dried and concentrated. The crude product was purified on silica gel with 20-30% EtOAc/Hexanes to give 31.4 g (76% yield) of **101**.

To a solution of **101** (20.3 g, 28.6 mmol) in 3L of THF, 0.5M KHMDS solution (60 mL, 30 mmol, 1.05 eq.) was added slowly by syringe pump over 120 minutes. After the mixture was stirred for 5 minutes, it was quenched with 1.5L of sat. NH_4Cl . The mixture was extracted three times with ether. The extracts were washed with brine, and then dried and concentrated. The crude product was purified on silica gel with 10-20%-50% EtOAc/Hexanes to give 14.2 g of **102** (76% yield).




To a solution of **102** (19 g, 29.15 mmol) in DMF (194 mL), imidazole (4 g, 58.3 mmol, 2 eq.) and TBSCl (5.27 g, 35 mmol, 1.2 eq.) were added. After the mixture was stirred overnight, it was quenched with a saturated solution of NaHCO_3 and water. The mixture was extracted with EtOAc. The organic layer was washed with water and brine, and then dried and concentrated. The crude product was purified on silica gel column to give 22 g (99% yield) of **103**.

To a solution of **103** (22 g, 28.7 mmol) in a mixture of CH_2Cl_2 (230 mL) and

H₂O (57.4 mL), DDQ (9.78g, 43 mmol, 1.5 eq.) was added at 0°C. After the mixture was stirred for 2 hours, it was quenched with 1L of a 1:1 mixture of aqueous saturated NaHCO₃ and 10% aqueous Na₂S₂O₃. The mixture was extracted three times, each with of ether. The extracts were washed with brine, and then dried and concentrated. The crude product was purified on silica gel to give 18.1 g of **104**.

To a solution of **104** (18g, 27.9 mmol) in 279 mL of CH₂Cl₂, dried 4Å molecular sieves (18g) and PCC (18 g) were added. After the mixture was stirred for 90 min, it was quenched with Et₃N (19.45 mL). The reaction mixture was filtered through a plug of celite and the plug was rinsed with 75% EtOAc in hexanes (900 mL). The filtrates were concentrated. The crude product was purified on silica gel column with 10-15-20% EtOAc/Hexanes to give 14.6 g (81%) of **105**.


In a 2L flask, **105** (8.5g, 13.2 mmol) was dissolved in CH₂Cl₂ (82.5 mL) and the mixture was cooled to 0°C. A solution of 5% H₂O/TFA (4.1/78.1 mL) was added slowly (~30min) and the mixture was stirred at 0°C for 14.5 hours. The reaction was monitored by TLC. The reaction mixture was diluted with CH₂Cl₂ at 0°C. The reaction was then quenched with a solution of NaHCO₃ in water (~1.2eq compared to TFA). The reaction was cooled to room temperature and extracted three times with CH₂Cl₂, dried with Na₂SO₄. The solution was then filtered and concentrated. Chromatography on Si-Gel, 50-60-75% EtOAc/hexane gave **106**: 4.8g, 93% yield. NMR: ¹H-NMR δ (Benzene-d₆) 0.65 (t, *J* = 7.2 Hz, 3H, NCH₂CH₃), 0.81 (d, *J* = 7.0 Hz, 3H, C4-CH₃), 1.03 (d, *J* = 6.0 Hz, 3H, C3-CH₃), 2.12 (m, 2H, 10-CH₂), 2.35 (d, *J* = 10.9 Hz, 1H, C9-OH), 2.52 (dt, *J*₁ = 5.3 Hz, *J*₂ = 7.2 Hz, 2H, NCH₂CH₃), 3.22 (t, *J* = 5.1 Hz, 1H, NH), 3.44 (m, 1H, 4-CH), 3.89 (m, 2H, 9-CH and C8-OH), 4.24 (dd, *J*₁ = 2.4 Hz, *J*₂ = 4.8 Hz, 1H, 8-CH), 4.67 (dq, *J*₁ = 6.0 Hz, *J*₂ = 10.4 Hz, 1H, 3-CH), 5.40 (dd, *J*₁ = 9.7 Hz, *J*₂ = 11.4 Hz, 1H, 5-CH), 5.49 (d, *J* = 11.5 Hz, 1H, 6-CH), 5.84 (d, *J* = 2.3 Hz, 1H, aromatic CH), 6.01 (m, 1H, 11-CH), 6.17 (d, *J* = 2.5 Hz, 1H, aromatic CH), 6.97 (d, *J* = 15.2 Hz, 1H, 12-CH), 13.05 (s, 1H, phenol).

Compound 114:

To a solution of **003** (2.914 g, 6.60 mmol) in 1, 3-Dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (20.0 mL, 0.166 mol) at 0 to -5 °C was added lithium hexamethyldisilazide (8.0 mL, 1.0M in THF). After the reaction mixture was stirred at cold for 30 minutes, to the solution was added 3-Iodo-1-Propene (0.74 mL, 7.9 mmol). The resulting solution was warmed up to room temperature and stirred for 4 hours. The reaction mixture was diluted with ether, added 10.0 ml of 1.0N HCl, extracted with ether. The combined ether solution was washed with saturated bicarbonate (30 mL), and brine (20 mL), and dried with MgSO₄. Chromatography purification with 10, 15% EtOAc/Hexane gave compound **107** (2.4 g, 75%) as white solid.

To a solution of compound **107** in *t*-BuOH:THF:water (100.0 mL, *t*-BuOH :THF:water =10:3:1) at room temperature was added N-Methylmorpholine N-oxide (0.79 g, 6.7 mmol) and Osmium tetroxide (1.70 g, 0.167 mmol). After the reaction mixture was stirred at room temperature for two hours, reaction progressed slowly. To the reaction mixture was added additional osmium tetroxide (4.71 g, 0.463 mmol). The reaction mixture was stirred at room temperature over night. The reaction mixture was diluted with EtOAc (250 mL), and was added sodium thiosulfate (10%, 150 mL). The reaction mixture was extracted the aqueous with EtOAc (2x250ml), dried and concentrated. Evaporation of the solvent gave compound **108** (1.8g, 70%).

To a solution of **108** in acetone (20 mL) was added 2,2-Dimethoxypropane (2.0 mL, 0.016 mol) and PTS (110 mg). The reaction mixture was stirred at room temperature for 30 minutes. The reaction mixture was then diluted with EtOAc (50 mL), washed with 5% NaOH (40 mL). The aqueous was extracted with EtOAc (3x50 mL). The combined organic solvent was concentrated. Chromatography purification (20- 30% EtOAc/hexane) gave compound **109** (1.63g, 84%) as oil.




To a 10 ml flask was added compound **005** (0.9624 g, 0.002464 mol) and **109** (1.45 g, 0.00261 mol), the mixture was dried under high vacuum. To the mixture was added Tris(dibenzylideneacetone)dipalladium(0) (0.677 g, 0.000739 mol), and N-Methylpyrrolidinone (2.50 mL, 0.0259 mol) and N-cyclohexyl-N-methylcyclohexanamine (1.06 mL, 0.00493 mol). The reaction mixture was stirred at 80°C over night. The reaction was cooled to room temperature, and was diluted with TBME. The organic solution was washed with 1N HCl, brine, and was concentrated under reduced pressure. Chromatography purification (20% EtOAc/Hexane) gave compound **110** (1.73g, 88%).

To a solution of 1.0M potassium tert-Butoxide (3.2 mL, 3.2mmol) in THF (10.0 mL) in a 50 mL round bottomed flask at 0°C was added a solution of compound **110** (1.72g, 2.16 mmol) in THF (5.0 mL) via syringe pump. The reaction mixture was stirred for 10 minutes after the addition of compound **110** and was quenched with tert-

butyldimethylsilyl chloride (490 mg, 0.0032 mol) in THF (5.0 mL) at 0°C. The reaction mixture was diluted with water and was extracted with EtOAc. Chromatography purification (20% EtOAc/hexane) gave compound **111** (0.64g, 35%).

To a solution of **111** (0.65 g, 0.00076 mol) in DCM (8.0 mL) at 0°C was added water (2.0 mL) and 2,3-Dichloro-5, 6-dicyanohydroquinone (0.245 g, 0.00106 mol). The reaction mixture was stirred at room temperature for 3 hours. After checking TLC (30% EtOAc/hexane), the reaction was quenched with sodium thiosulfate in saturated bicarbonate. The reaction mixture was extracted with TBME, dried and concentrated. Chromatography purification (20% - 40% EtOAc/hexane) gave compound **112** (0.557g, 100%).




To a solution of compound **112** (111.2 mg, 0.0001519 mol) in DCM (1.0 mL) was added Pyridine (18 μ L, 0.00023 mol) and Dess-Martin periodinane (0.077 g, 0.00018 mol) at 0 °C. The reaction mixture was stirred for 30 minutes at 0°C, still some starting material was detected by TLC. The reaction mixture was then allowed to warm to room temperature and stirred for 30 minutes. The reaction mixture was then diluted with 30% EtOAc/hexane, the resulting solution was filtered through a silicon plug, and washed with 30% EtOAc/hexane. Chromatography purification (30% EtOAc/hexane) gave compound **113** (103.8 mg, 94%).

To a solution of **113** (101.5 mg, 0.0001390 mol) in DCM (1.0 mL) was added water (20 micro liter) and Trifluoromethanesulfonic acid (61.5 μ L, 0.000695 mol) at 0°C. The reaction mixture was stirred for 20 minutes at 0°C, and was allowed to warm to room temperature and was stirred for another 30 minutes. The reaction mixture was quenched with saturated sodium bicarbonate, and was extracted with DCM. Chromatography purification (5%MeOH/DCM) gave compound **114** (42.0 mg, 69.4%).

^1H NMR (CD_3OD): 6.86 (*d*, $J = 15.6$ Hz, 1H), 6.30 (*d*, $J = 11.6$ Hz, 1H), 6.10-6.18 (*m*, 2H), 5.93-6.00 (*m*, 2H), 4.47 (*d*, $J = 2.0$ Hz, 1H), 4.04 (*t*, $J = 2.8$ Hz, 1H), 3.78 (*m*, 1H), 3.55 (*m*, 2H), 3.43 (*m*, 1H), 3.30 (*m*, 3H), 3.08 (*m*, 1H), 2.08 (*m*, 2H), 1.35 (*d*, $J = 5.6$ Hz, 3H), 1.14 (*d*, $J = 7.2$ Hz, 3H). Mass 457.95 observed.

Compound 122:




Triethylsilane (510 μL , 0.0032 mol) and Trifluoroacetic Acid (75 μL , 0.97 mmol) were added to a solution of **003** (176 mg, 0.399 mmol) in DCM (3.0 mL) at 0°C . The reaction mixture was stirred at 0°C for 1 hour, still a lot starting material left after checking TLC (30% EtOAc/Hex). The reaction mixture was warmed up to room temperature and stirred at ambient temperature over night. There was not much progress by checking TLC. To the reaction mixture was added additional TFA (1.0 mL) at 0°C . The reaction mixture was stirred at room temperature for 30 minutes. There was no starting material by TLC. The reaction mixture was poured into saturate bicarbonate, extracted with EtOAc (3x20 mL). The combined organic layer was dried and filtered. The mixture was concentrated and purified on preparative TLC (30%EtOAc/hexane) gave compound **115** (0.107g, 79%).

Sodium triacetoxyborohydride (0.57 g, 0.0027 mol) was added to a solution of *N-tert*-butoxycarbonyl piperidine-1-carboxylate (448.1 mg, 0.002101 mol), compound **115** (348.0 mg, 0.001020 mol), and acetic acid (1300 μL , 0.023 mol) in THF (10.0 mL) at room temperature. The reaction mixture was stirred at room temperature for 5 hours.

The reaction mixture was then concentrated without aqueous workup. Purification using preparative TLC (35% EtOAc/hexane) gave compound **116** (0.4164g, 76%).

Di-tert-Butyldicarbonate (0.0251 g, 0.000115 mol) was added to a solution of compound **116** (31.0 mg, 0.0000576 mol) in DCM (1.0 mL). To the reaction mixture was added 4-Dimethylaminopyridine (0.00703 g, 0.0000576 mol) and triethylamine (8.02 μ l, 0.0000576 mol). The reaction mixture was then stirred at ambient temperature over night. Preparative TLC purification (30% EtOAc/hexane) gave compound **117** (16.4 mg, 45%).




Compounds **117** (0.3916 g, 0.0006132 mol) and **005** (0.2422 g, 0.0006202 mol) were combined in a reaction flask and was dried under high vacuum.

Tris(dibenzylideneacetone) dipalladium(0) (0.17 g, 0.00019 mol) was added to the above mixture in a drybox. N-Methylpyrrolidinone (2.0 mL, 0.021 mol) and Cyclohexanamine, N-cyclohexyl-N-methyl- (268 μ l, 0.00125 mol) was added to the reaction flask. The reaction mixture was stirred at 80°C in an oil bath for 20 hours. The mixture was diluted with MTBE, and washed with 1.0 M hydrochloric acid. The aqueous was extracted with MTBE and was concentrated. Chromatography purification (20-40%EtOAc/hexane) gave compound **118** (0.373 g, 69%).

Compound **118** (0.37 g, 0.00042 mol) in THF (7.2 mL) was added to a solution of potassium tert-Butoxide in tetrahydrofuran (0.716 mL, 1.00M) in tetrahydrofuran (7.20 mL, 0.0888 mol) at 0°C over 5 minutes. No starting material was observed (TLC,

30% EtOAc/hexane) after the reaction mixture was stirred for 10 minutes. The reaction was quenched with tert-Butyldimethylsilyl chloride (0.108 g, 0.000716 mol) in THF (1.0 mL) at 0°C and was stirred for 10 minutes. Chromatography purification (20%EtOAc/hexane) gave compound **119** (0.182g, 46%).


Water (2.0 mL) and Dichlorodicyanoquinone (57.3 mg, 0.000252 mol) was added to a solution of compound **119** (181.6 mg, 0.0001942 mol) in DCM (8.0 mL) at 0°C. The reaction mixture was then warmed to ambient temperature and was stirred for 3 hours. The reaction was quenched with 1:1 10% sodium thiosulfate/saturated bicarbonate after no starting material was detected (TLC, 30%EtOAc/hexane). The aqueous was extracted with TBME. Chromatography purification (20%-40% EtOAc/hexane) gave compound **120** (0.1461g, 92%).



To a solution of compound **120** (146.1 mg, 0.179 mmol) in DCM (1.0 mL) was added Pyridine and Dess-Martin periodinane (0.099 g, 0.23 mmol) at 0°C. The reaction mixture was stirred for 30 minutes at 0°C, still some starting material was detected by TLC. The reaction mixture was then allowed to warm to room temperature and stirred at ambient temperature until all the starting material was consumed. The reaction mixture was then diluted with 30% EtOAc/hexane, the resulting solution was filtered through a silicon plug, and washed with 30% EtOAc/hexane. Chromatography purification (30% EtOAc/hexane) gave compound **121** (107.4 mg, 74%).

To a solution of compound **121** (107.4 mg, 0.0001321 mol) in Methylene chloride (1.50 mL, 0.0234 mol) was added Water (22.4 μ L, 0.00124 mol) and

Trifluoromethanesulfonic acid (55.0 μ l, 0.000622 mol) at 0°C. The reaction mixture was stirred at 0°C for 90 minutes. Still some starting material was not consumed (TLC, 10%NH₄OH/EtOH). The reaction mixture was stirred for another 30 minutes at room temperature and was diluted with 20 ml TBME. To the reaction mixture was added saturated sodium bicarbonate, extracted with EtOAc (3x20 mL), the aqueous was further extracted with EtOAc (3x20 mL). The combined organic solvent was dried with sodium sulfate, and filtered. The crude material was loaded onto preparative TLC, and was eluted with 10% NH₄OH/EtOH (Rf 0.1). The desired band was collected, and was rinsed with 10% NH₄OH/EtOH to give compound **122** (46.8 mg, 77%). ¹HNMR (CD₃OD): 6.85 (*d*, *J* = 15.2 Hz, 1H), 6.29 (*d*, *J* = 12.0 Hz, 1H), 6.08-6.11 (*m*, 2H), 5.91-5.98 (*m*, 2H), 4.47 (*d*, *J* = 2.4 Hz, 1H), 4.04 (*m*, 1H), 3.46 (*m*, 1H), 3.02 (*d*, *J* = 12.0 Hz, 2H), 2.97 (*d*, *J* = 6.4 Hz, 2H), 2.56 (*m*, 2H), 2.09 (*m*, 2H), 1.71 (*m*, 4H), 1.34 (*d*, *J* = 6.0 Hz, 3H), 1.22, (*m*, 2H), 1.14 (*d*, *J* = 6.8 Hz, 3H).

Compound 127:

A solution of acetic acid/water (4.0 mL, Acetic acid:water = 4:1) at 0°C was added to compound **123** (prepared in the same manner as compound **064**) (195.2 mg, 0.0002617 mol). The reaction mixture was kept at 0°C for 30 minutes and then was warmed up to ambient temperature. After being stirred over night at ambient temperature, the reaction mixture was diluted with sodium bicarbonate, and was extracted with EtOAc. Preparative TLC (75% EtOAc/Hexane) gave compound **124** (0.1686 g, 91%).


To a solution of compound **124** (168.6 mg, 0.0002388 mol) in Toluene (3.0 mL, 0.028 mol) and DCM (1.0 mL) was added Lead tetraacetate (0.127 g, 0.000287 mol). The reaction mixture was stirred for 20 to 30 minutes at ambient temperature, and was

filtered through celite. The celite was washed with DCM. Preparative TLC (50% EtOAc/hexane) gave compound **125** (106.1 mg, 66%).

To a solution of compound **125** (106.0 mg, 0.0001573 mol) in tetrahydrofuran (0.50 mL, 0.0062 mol), tert-Butyl alcohol (1.50 mL, 0.0157 mol), 2-Methylbut-2-ene (0.50 mL, 0.0047 mol) was added a solution of sodium chlorite (88.9 mg, 0.000786 mol) and potassium phosphate monobasic (100 mg, 0.0007 mol) in 0.6 ml of water at ambient temperature. The reaction mixture was stirred at ambient temperature for 2 hours, and was concentrated. The residue was diluted with water, acidified to pH 2 with dilute HCl. The aqueous was extracted with EtOAc. Preparative TLC gave compound **126** (66.1 mg, 61%).

To a solution of compound **126** (32.5 mg, 0.0471 mmol) in methylene chloride (1.0 mL, 0.016 mol) and water (1.7 μ l, 0.094 mmol) at 0°C was added trifluoromethanesulfonic acid (17 μ l, 0.00019 mol). The reaction mixture was stirred at 0°C for 60 minutes and at ambient temperature for 60 minutes. The reaction mixture was quenched with saturated sodium bicarbonate. The aqueous phase was extracted with DCM (2X20mL). No desired product appeared in a TLC of the DCM solution. The aqueous phase was then acidified to pH 3, and was extracted with EtOAc (3x10 mL). Preparative TLC gave compound **127** (10.2 mg, 50%).


Compound 137:



To a solution of compound **098** (1.0g, 1.98mmol) in MeOH/CH₂Cl₂ (5ml/5ml) at -78°C was bubbled O₃. After 7min, N₂ was bubbled through for 10min and 2 pipettes of Me₂S were added. The reaction mixture was warmed up to room temperature and solvents were removed by evaporation. Purification by silica gel chromatography gave compound **128** (735mg, 73%).


To a reaction flask containing CBr₄ (969mg, 2.90mmol) and PPh₃ (15.2g 5.80mmol) in 10ml of CH₂Cl₂ was added a solution of compound **128** (734mg, 1.45mmol) and Et₃N (202μl, 1.45mmol) in 5ml of CH₂Cl₂ at 0°C. The ice bath was removed and the reaction mixture was stirred at room temperature for 1 hour. After titration with Hexane, purification by silica gel chromatography gave compound **129** (557g, 58%).

To a solution of compound **129** (557mg, 0.84mmol) in THF (5ml) at -78°C was added n-BuLi (840μl, 2.5M in Hexane). After 1hr the reaction was quenched with saturated aqueous NH₄Cl. The aqueous layer was extracted with MTBE. Purification by silica gel chromatography gave compound **130** (360mg, 85%).



To a solution of compound **130** (358mg, 0.71mmol) in THF (5ml) at room temperature was added TBAF (1.1ml, 1.0M in THF). The reaction mixture was heated at 50°C for 4 hours. The mixture was then diluted with MTBE, washed with water and brine. Purification by silica gel chromatography gave compound **131** (277mg, 100%).

To a solution of compound **131** (263.8mg, 0.678mmol) and compound **132** (637.4mg, 1.356mmol) in benzene (1.8ml) and Et₂NH (1.2ml) at room temperature was added Pd(PPh₃)₄ (39.2mg, 0.034mmol) and CuI (12.9mg, 0.068mmol). The reaction mixture was stirred for 5 hours. The mixture was then diluted with MTBE, washed with saturated NH₄Cl, saturated NaHCO₃ and brine. Purification by silica gel chromatography gave compound **133** (251.5mg, 52%).



To a solution of t-BuOK (565μl, 1.0M in THF) in THF (13.0ml) at 0°C was added the solution of compound **133** (200.0mg, 0.28mmol) in THF (8.0ml) at rate of 0.1ml/min. After 1 hour, TBSCl (169mg, 0.56mmol) was added. After 5 minutes, the reaction mixture was quenched with saturated NaHCO₃, diluted with water and extracted with MTBE. Purification by silica gel chromatography gave compound **134** (175.4mg, 82%).


To a solution of compound **134** (200.0mg, 0.28mmol) in DCM 4.0ml) and water (1.0ml) at room temperature was added DDQ (96.2mg, 0.42mmol). After 2 hours, the reaction was quenched with 10% Na₂S₂O₃ in a saturated NaHCO₃ aqueous solution and

extracted with MTBE. Purification by silica gel chromatography gave compound **135** (133.1mg, 74%).

To a solution of compound **135** (133.0mg, 0.207mmol) in DCM (3.0ml) at room temperature was added pyridine (42.0μl, 0.518mmol) and Dess-Martin reagent (175.0mg, 0.414mmol). After 1 hour, the reaction was quenched with 10% Na₂S₂O₃ in saturated NaHCO₃ aqueous solution and extracted with DCM. Purification by silica gel chromatography gave compound **136** (114.8mg, 86%).

To a solution of compound **136** (113.0mg, 0.176mmol) in DCM (2.0ml) at 0°C was added water (19.0μl, 1.056mmol) and TfOH (46.0μl, 0.528mmol). The reaction was allowed to warm up to room temperature. After 30 minutes, the reaction was quenched with saturated NaHCO₃ and extracted with DCM. Purification by silica gel chromatography gave compound **137** (46.4mg, 68%). ¹H NMR (400 MHz, CDCl₃) δ 1.20 (d, J=6.8Hz, 3H), 1.25 (t, J=6.8, 3H), 1.38 (d, J=6.0, 3H), 2.47-2.48 (m, 2H), 2.85 (d, J=11.2Hz), 3.70-3.21 (m, 2H), 3.41-3.45 (m, 1H), 3.95 (d, J=5.2, 1H), 4.00-4.40 (m, 1H), 4.27-4.33 (m, 1H), 4.54-4.56 (m, 1H), 4.93-5.00 (m, 1H), 6.04 (d, J=2.4, 1H), 6.23-6.32 (m, 3H), 12.10 (s, 1H).

Compound 144:




To a solution of compound **130** (475mg, 0.944mmol) and compound **138** (558mg, 1.227mmol) in benzene (2.2ml) and Et₂NH (1.5ml) at room temperature was added Pd(PPh₃)₄ (55mg, 0.0472mmol) and CuI (18mg, 0.0944mmol). The reaction mixture was stirred for 3 hours. The reaction mixture was then diluted with MTBE,

washed with saturated NH_4Cl , saturated NaHCO_3 and brine. Purification by silica gel chromatography gave compound **139** (485mg, 56%).

To a solution of compound **139** (485mg, 0.60mmol) in THF (5.0ml) was added buffered TBAF (3.6ml, 0.5N buffered with imidazole/HCl) at room temperature. The reaction mixture was heated at 50°C for 60 hours. The reaction mixture was then diluted with MTBE and washed with water and brine. Purification by silica gel chromatography gave compound **140** (280.8mg, 67%).

To a solution of t-BuOK (720 μl , 1.0M in THF) in THF (13.5ml) at 0°C was added the solution of compound **140** (250mg, 0.36mmol) in THF (13.5ml). After 2 hours, TBSCl (217mg, 1.44mmol) was added. After 5 minutes, the reaction mixture was quenched with saturated NaHCO_3 , diluted with water and extracted with MTBE. Purification by silica gel chromatography gave compound **141** (155mg, 52%).




To a solution of compound **141** (153mg, 0.204mmol) in DCM 4.0ml) and water (1.0ml) at room temperature was added DDQ (69.5mg, 0.306mmol). After 2.5 hours, the reaction was quenched with saturated NaHCO_3 and extracted with MTBE. Purification by silica gel chromatography gave compound **142** (124.0mg, 96%).

To a solution of compound **142** (124.0mg, 0.197mmol) in DCM (5.0ml) at room temperature was added pyridine (40.0 μl , 0.492mmol) and Dess-Martin reagent (167.0mg, 0.394mmol). After 1hr, the reaction was quenched with 10% $\text{Na}_2\text{S}_2\text{O}_3$ in saturated NaHCO_3 aqueous solution and extracted with DCM. Purification by silica gel chromatography gave compound **143** (122.8mg, 99%).

To a solution of compound **143** (54.5mg, 0.087mmol) in DCM (2.0ml) at 0°C was added water (9.4μl, 0.522mmol) and TfOH (23.0μl, 0.261mmol). The reaction was allowed to warm up to room temperature. After 30 minutes, it was quenched with saturated NaHCO₃ and extracted with DCM. Purification by silica gel chromatography gave compound **144** (30.0mg, 92%). ¹H NMR (400 MHz, CDCl₃) δ 1.20 (d, J=6.8Hz, 3H), 1.38 (d, J=6.0Hz, 3H), 2.47-2.48 (m, 2H), 2.85 (d, J=5.2Hz, 3H), 3.40-3.50 (m, 1H), 3.94 (d, J=5.2), 4.12-4.19 (m, 1H), 4.28-4.33 (m, 1H), 4.54-4.56 (m, 1H), 4.93-5.00 (m, 1H), 6.05 (d, J=2.4Hz, 1H), 6.30 (s, 1H), 6.33 (m, 2H), 12.11 (s, 1H).


Compound 155:



To a solution of **005** (2.46g, 4.87mmol) in MeOH (40ml) at 0°C was added p-toluenesulfonic acid (1.02g, 5.36mmol). The reaction was warmed up to room temperature and stirred for 18 hours. The reaction mixture was then quenched with saturated NaHCO₃ and extracted with MTBE. Purification by silica gel chromatography gave compound **145** (1.42g, 64%).

To a solution of compound **145** (1.53g, 4.37mmol) in DCM at 0°C was added Et₃N (1.83ml, 13.11mmol) and TBSOTf (2.10ml, 9.18mmol). The reaction was warmed up to room temperature and stirred for 35 minutes. The reaction mixture was then quenched with saturated NaHCO₃ and extracted with DCM. Purification by silica gel chromatography gave compound **146** (1.18g, 41%).


To a solution of compound **146** (1.09g, 1.88mmol) in DCM (10.0ml) at room temperature was added pyridine (303μl, 3.76mmol) and Dess-Martin reagent (1.20g, 2.82mmol). After 5 hours, the reaction was quenched with 10% Na₂S₂O₃ in saturated NaHCO₃ aqueous solution and extracted with MTBE. Purification by silica gel chromatography gave compound **147** as crude product.



To a solution of compound **147** in MeOH (15.0ml) at 0°C was added NaBH₄ (142mg, 3.76mmol). After 10 minutes, the reaction was quenched with saturated NH₄Cl and extracted with MTBE. Purification by silica gel chromatography gave compound **148** (714mg, 66% from 1006-216B).

To a solution of compound **148** (710mg, 1.23mmol) in THF (10ml) at room temperature was added TBAF (3.7ml, 1.0M in THF). The reaction mixture was heated at 50°C for 17 hours. The reaction mixture was then diluted with MTBE, washed with saturated NaHCO₃ and brine. Purification by silica gel chromatography gave compound **149** (419mg, 97%).


To a solution of compound **149** (400mg, 1.14mmol) in acetone (10ml) at room temperature was added 2,2-dimethoxypropane (419μl, 3.42mmol) and PTSA (100mg). After 20min, it was diluted with MTBE, washed with saturated NaHCO₃ and brine. Purification by silica gel chromatography gave compound **150** (450mg, 95%).



To a solution of compound **150** (384mg, 0.98mmol) and compound **138** (582mg, 1.27mmol) in NMP (2.0ml) at room temperature was added $\text{Pd}_2(\text{DBA})_3$ (90.0mg, 0.098mmol) and Cy_2NMe (272 μl , 1.27mmol). The reaction mixture was heated at 80°C for 24 hours. The reaction mixture was then diluted with MTBE, filtered through celite, washed with 1.0N aqueous HCl and brine. Purification by silica gel chromatography gave compound **151** (441.3mg, 65%).

To a solution of t-BuOK (884 μl , 1.0M in THF) in THF (15.0ml) at 0°C was added the solution of compound **151** (410mg, 0.59mmol) in THF (5.0ml). After 30 minutes, TBSCl (178mg, 1.18mmol) was added. After 10 minutes, the reaction mixture was quenched with saturated NaHCO_3 , diluted with water and extracted with EtOAc. Purification by silica gel chromatography gave compound **152** (170.1mg, 34%).


To a solution of compound **152** (170mg, 0.226mmol) in DCM 4.0ml) and water (1.0ml) at room temperature was added DDQ (77.0mg, 0.339mmol). After 2.5 hours, the reaction was quenched with 10% $\text{Na}_2\text{S}_2\text{O}_3$ in saturated NaHCO_3 and extracted with MTBE. Purification by silica gel chromatography gave compound **153** (122.6mg, 86%).



To a solution of compound **153** (61.0mg, 0.097mmol) in DCM (2.5ml) at room temperature was added pyridine (19.6 μl , 0.243mmol) and Dess-Martin reagent (82.0mg, 0.194mmol). After 1 hour, the reaction was quenched with 10% $\text{Na}_2\text{S}_2\text{O}_3$ in saturated NaHCO_3 aqueous solution and extracted with MTBE. Purification by silica gel chromatography gave compound **154** (46.1mg, 76%).

To a solution of compound **154** (41.8mg, 0.066mmol) in DCM (2.0ml) at 0°C was added water (7.1 μl , 0.396mmol) and TfOH (17.6 μl , 0.198mmol). The reaction was allowed to warm up to room temperature. After 30 minutes, it was quenched with saturated NaHCO_3 and extracted with DCM. Purification by silica gel chromatography gave compound **155** (11.1mg, 45%). ^1H NMR (400 MHz, CDCl_3) δ 0.64 (d, $J=6.80$, 3H), 0.93 (d, $J=6.0$, 3H), 2.06(d, $J=3.6\text{Hz}$, 3H), 2.37-2.44 (m, 1H), 2.54-2.64 (m, 1H), 2.79 (br, 1H), 3.05-3.22 (m, 2H), 3.81-3.91 (m, 1H), 4.18-4.33 (m, 1H), 4.60-4.71 (m, 1H), 5.33 (t, $J=11.6\text{Hz}$, 1H), 5.86 (d, $J=2.8\text{Hz}$, 1H), 5.91-6.03 (m, 1H), 6.04(d, $J=11.6\text{Hz}$, 1H), 6.11 (d, $J=2.0\text{Hz}$, 1H), 7.00 (d, $J=15.6\text{Hz}$, 1H), 12.86 (s, 1H).


Compound 156:




Compound **106** (1 wt) was dissolved in TFA (3 vols, 15 eq.) and stirred at room temperature. The progress of the reaction was monitored by TLC. After 48 h stirring,

the reaction mixture was diluted with DCM (20 vols), washed with *sat. aq.* NaHCO₃, and brine. The organic layer was concentrated and purified by flash chromatography to give **156** as white solid (48.3% yield) (NMR data below). Compound **106** was also recovered (9.1% recovery rate). ¹HNMR (C₆D₆): δ = 6.83 (d, 1H), 6.12 (s, 1H), 5.81 (dt, 1H), 5.70 (s, 1H), 5.55 (d, 1H), 5.40 (dd, 1H), 5.12 (s, 1H), 4.61 (m, 1H), 4.19 (s, 1H), 3.86 (d, 1H), 3.26 (q, 1H), 3.15 (br s, 1H), 2.48 (q, 2H), 2.30-2.35 (m, 1H), 2.12 (q, 1H), 0.99 (d, 3H), 0.75 (d, 3H), 0.60 (t, 3H), -2.00 (s, 1H)


Compound 157:




Compound **106** was dissolved in 1,4-dioxane (11 vols) and ⁿBu₃P (1.2 eq.) was slowly added under N₂ atmosphere. The reaction mixture was stirred at room temperature, and the progress of the reaction was followed by TLC and HPLC. After 23 h stirring at r.t., the reaction mixture was diluted with TBME (100 vols), washed with water (2 x 10 vols) and brine (10 vols). The aqueous layers were back extracted with TBME (50 vols). The combined organic layer was dried over Na₂SO₄ and concentrated. The residue was purified by flash chromatography to give **157** as white solid (58.5%). MS (ES⁺): M+Na⁺ = 412.1, M+H⁺ = 390.1; MS (ES⁻): M-H⁺ = 388.0; ¹HNMR (C₆D₆): δ = 6.78-6.84 (m, 2H), 6.13 (d, 1H), 5.95 (d, 1H), 5.86 (m, 1H), 5.83 (d, 1H), 4.97-5.02 (m, 1H), 4.47 (br s, 1H), 4.02 (br s, 1H), 3.93 (br s, 1H), 3.26 (t, 1H), 2.64 (br s, 1H), 2.52 (m, 2H), 2.39-2.43 (m, 1H), 2.04-2.11 (m, 1H), 1.68 (m, 1H), 0.88 (d, 3H), 0.64 (t, 3H), 0.58 (d, 3H), -2.38 (s, 1H).

Compound 158:


The starting material (**159**) (100.0mg, 0.13mmol) was dissolved in 1.0ml of methanol at 0°C. p-Toluenesulfonic acid (49.4mg, 0.28mmol) was added. The reaction mixture was stirred at 0°C for 25min, then at room temperature over night. It was quenched with sat. aqueous sodium bicarbonate solution and extracted with ethyl acetate. After purification on silica gel column, **160** was obtained in 84% yield. The chiral starting material (**159**) can be obtained by separation of a racemic mixture of **094** (see preparation of **106**) and carrying through the steps to achieve **103**.




The two stereoisomers of **094** (at position 7) were separated by chromatography on silica gel using a 12:1 mixture of hexane / ethyl acetate as eluent. Identification of the stereochemistry was made by NMR spectroscopy. The less polar isomer was found to have the “R” stereochemistry at position 7 while the more polar has the “S” stereochemistry. The “R” isomer, **094-(7R)**, was used to produce **159** with the same synthetic route that was used to produce **103** from **094** (see example compound **106** for reference).




Compound **160** (46.0mg, 0.063mmol) was dissolved in 2.0ml of dichloromethane at 0°C. Pyridine (51.0μl, 0.63mmol) and phosgene (167.0μl, 0.32mmol) were added, respectively. The reaction mixture was stirred at 0°C for 15min, it was quenched with sat. sodium bicarbonate solution and extracted with methyl t-butyl ether. After purification on silica gel prep plate, **161** was obtained in quantitative yield.



Compound **161** (47.4mg, 0.063mmol) was dissolved in the mixture of 2.0ml dichloromethane and 0.5ml water. 2,3-Dichloro-5,6-dicyano-1,4-benzoquinone (31.5mg, 0.14mmol) was added. After stirring at room temperature for 2h, the reaction mixture was quenched with 10% sodium thiosulfate in sat. sodium bicarbonate solution and extracted with methyl t-butyl ether. After purification on silica gel prep plate, **162** was obtained in 69% yield.



Compound **162** (27.6mg, 0.044mmol) was dissolved in dichloromethane. Pyridine (8.9μl, 0.11mmol) and Dess-Martin reagent (37.0mg, 0.088mmol) were added, respectively. The reaction mixture was stirred at room temperature for 1.5h. It was quenched with 10% sodium thiosulfate in sat. sodium bicarbonate aqueous solution, and extracted with methyl t-butyl ether. After purification on silica gel column, **163** was obtained in 100% yield.



Compound **163** (27.7mg, 0.044mmol) was dissolved in 1.0ml dichloromethane. Triflic acid (11.6μl, 0.13mmol) and water (4.8μl, 0.26mmol) were added at 0°C. After stirring at room temperature for 1h, the reaction mixture was quenched with sat. sodium bicarbonate solution and extracted with dichloromethane. After purification on silica gel column, **158** was obtained in 89% yield. ¹H NMR (CDCl₃) δ 12.10 (s, 1H), 7.08 (dd, J=15.2, 2.4 Hz, 1H), 6.11 (m, 2H), 6.03 (dd, J=13.2, 2.4 Hz, 2H), 5.53 (m, 1H), 5.14 (d, J=8.4 Hz, 1H), 5.04 (m, 1H), 4.94 (m, 1H), 4.11 (s, 1H), 3.92 (m, 1H), 3.19 (dd, J=21.6, 7.2 Hz, 2H), 2.86 (m, 1H), 2.75 (m, 1H), 1.45 (d, J=6.0 Hz, 3H), 1.26 (t, J=7.2, 3H), 1.14 (d, J=6.8 Hz, 3H).

Examples 2-12 Biological Assays

A set of thirty compounds of formula (I) or (II) was randomly selected, and each compound was tested for its growth inhibitory activity in MDA-MB-435 and HT-29 cancer cell lines. Several compounds were found to have remarkable cancer cell growth inhibitory activity (*e.g.*, nM IC₅₀ values). Two compounds (herein referred to as Compounds **091** and **106**) were selected for further investigation to further evaluate and characterize biological properties of the compounds of this class (*i.e.*, F152 analogs). It is to be understood for the biological assays, *e.g.*, the cell based assays, that results given may be exemplary of more than one set of data. In certain cases, *e.g.*, in certain *in vitro* assays, a mean and standard deviation is representative of the acquired data. In other cases, *e.g.*, in certain *in vivo* assays, a single representative data set is chosen from several substantially similar data sets (*e.g.*, data sets taken from the same assay using various dosages, etc.). The representative data set is often chosen based upon the assay conditions most similar to assay conditions used in clinical studies for human dosages. The skilled artisan would understand that different assay conditions, *e.g.*, different dosages or dosing schedules, may give different results.

Example 2

A biochemical MEK1 assay was carried out to investigate whether compounds of formula (I) or (II), *e.g.*, compounds **029**, **091**, **106** and **114** are MEK1 inhibitors. Biochemical MEK1 inhibition of the compounds was assessed using an ELISA assay. All assays were performed in a 96 well format. A plate coated with ERK2, a substrate protein for MEK1 was used for biochemical MEK1 assay. In a final volume of 40 μ L, MEK1 (25 ng) was incubated with 50 mmol/L HEPES, 20 mmol/L $MgCl_2$, 4 mmol/L $MnCl_2$, and 0.5 mmol/L Na_3VO_4 in the absence or presence of the test compounds. The reaction was initiated by the addition of 1 μ mol/L ATP. After incubation for 40 minutes at 30°C, the reaction mixture was discarded and the plate was then washed 4 times with 0.05% Tween20/PBS. Each well was incubated with 100 μ L of 1:4000 diluted phospho-ERK1/2 monoclonal antibody (Cell Signaling, MA) for 1.5 hours at room temperature. The antibody was then discarded and the plate was washed 6 times with 0.05% Tween20/PBS. Finally, 100 μ L of 1:2000 diluted of HRP conjugated anti-mouse IgG antibody (Cell Signaling, MA) was added. After incubation for 90 minutes, this antibody was discarded and the plate was washed 8 times with 0.05% Tween20/PBS. Phosphorylated ERK2 was then visualized by peroxidase reaction, and read at 450 nm. Figure 2A is a table depicting results from MEK1 assays.

The experiments confirm that compounds **091** and **106** are potent MEK1 inhibitors with low nM activity (See Figure 2A). Compound **091** showed more potent MEK1 inhibitory activity at a low concentration of ATP, suggesting modes of binding to MEK1 are ATP competitive.

Example 3

A study was carried out to investigate the specificity of compounds to various cancer relevant kinases *in vitro*. Compounds **091** and **106** were tested at the concentrations of 0.1 μ M and 1 μ M against a panel of 95 cancer relevant kinases including: Abl(h), Abl (T315I)(h), ALK(h), AMPK(r), Arg(m), Aurora-A(h), Axl(h), Blk(m), Bmx(h), BTK(h), CaMKII(r), CaMKIV(h), CDK1/cyclinB(h), CDK2/cyclinA(h), CDK2/cyclinE(h), CDK3/cyclinE(h), CDK5/p35(h), CDK6/cyclinD3(h), CDK7/cyclinH/MAT1(h), CHK1(h), CHK2(h), CK1 δ (h), CK2(h), c-RAF(h), CSK(h), cSRC(h), EGFR(h), EphB2(h), EphB4(h), Fes(h), FGFR3(h), Flt3(h), Fms(h), Fyn(h), GSK3 α (h), GSK3 β (h), IGF-1R(h),

IKK α (h), IKK β (h), IR(h), JNK1 α 1(h), JNK2 α 2(h), JNK3 (h), Lck(h), Lyn(h), MAPK1(h), MAPK2(h), MAPKAP-K2(h), MEK1(h), Met(h), MKK4(m), MKK6(h), MKK7 β (h), MSK1(h), MST2(h), NEK2(h), p70S6K(h), PAK2(h), PAR-1B α (h), PDGFR α (h), PDGFR β (h), PDK1(h), PKA(h), PKB α (h), PKB β (h), PKB γ (h), PKC α (h), PKC β II(h), PKC γ (h), PKC δ (h), PKC ϵ (h), PKC η (h), PKC ι (h), PKC μ (h), PKC θ (h), PKC ζ (h), PKD2(h), PRAK(h), PRK2(h), ROCK-II(r), ROCK-II(h), Ros(h), Rsk1(h), Rsk2(h), Rsk3(h), SAPK2a(h), SAPK2b(h), SAPK3(h), SAPK4(h), SGK(h), Syk(h), Tie2(h), TrkB(h), Yes(h) & ZAP-70(h) (Figure 2B).

In a final reaction volume of 25 μ L, each human enzyme was incubated with appropriate buffer. The reaction was initiated by the addition of the MgATP mix with [γ ^{33} P-ATP]. After incubation for 40 minutes at room temperature, the reaction was stopped by the addition of 5 μ L of a phosphoric acid solution. All assays were performed at an ATP concentration of 10 μ M.

10 μ L of the reaction was spotted onto a P30 filtermat and washed three times for 5 minutes in a phosphoric acid solution. ^{33} P transferred to a substrate peptide was then determined by scintillation counting. The results were expressed as a percentage of that in control incubations. The kinases (1) with >50% inhibition at the 0.1 μ M concentration of compound, and (2) with concentration-response of inhibitions between 0.1 μ M and 1 μ M were considered as target kinases of the compounds.

Besides strong MEK1 inhibition, the compounds also inhibited members of the Src tyrosine kinase family (Lyn, Fyn, Lck, Yes, c-Src), FLT-3 and TrkB kinase activities at 0.1 μ M concentration. Moreover, it appears that the compounds are highly selective for tyrosine kinases, as indicated by their lack of inhibition of a multitude of serine/threonine kinases.

Furthermore, inhibitory activities of the compounds were confirmed against six of these kinases with a full range of concentrations. Three cytoplasmic Src family kinases were studied: c-Src, Fyn and Lyn. We also studied cytoplasmic Abl kinase and two receptor tyrosine kinases: Flt3, and TrkB. A standard ATP concentration of 10 μ M was used in all reactions. All experiments were conducted with duplicate measurements. As seen in Figure 2B, six kinases showed IC₅₀ below 214 nM for Compound **091**. TrkB and members of the Src family were the most sensitive enzymes to Compound **091** inhibition.

Example 4

***In vitro* anticancer activity of exemplary compounds.** Compounds **091** and **106**, were extensively tested in solid cancer cell lines (shown in Figure 3 for 18 cell lines for compound **091** and 20 cell lines for compound **106**). All cell lines were introduced into 96-well plates and grown in the absence or continuous presence of 0.3–10000 nM of either compound **091** or compound **106** for 96 hours. Cell growth was assessed using a CellTiter-Glo[®] Luminescent Cell Viability Assay (Promega) or a methylene blue assay. IC₅₀ values were determined as the concentration of a substance which inhibits cell growth by 50% compared to untreated cell populations.

Cell lines which carried a B-RAF V600E mutation were very sensitive to Compounds **091** and **106** in the low-nM or sub-μM concentration range, as shown in Figure 3.

Example 5

***In vivo* Antitumor Activity in Colo205 Human Colon Cancer Xenografts** (Figure 4). A study was carried out to investigate anticancer activity of compounds **091** and **106** in human Colo205 BRAF mutated colon cancer xenografts. The compounds were evaluated against subcutaneous-implanted Colo205 human colon cancer xenografts in female athymic mice.

The human colon cancer cell line, Colo205 which has B-RAF (V600E) mutation was transplanted subcutaneously into female nude mice. The experiment consisted of 6 mice per group on the first day of treatment. The compound was administered intravenously for two weekly rounds of five daily injections (q1d x 5) with a two-day rest period between the rounds. Compound **091** was administered in a vehicle volume of 0.1 mL/10 g of body weight.

The general health of the mice was monitored daily. Tumor dimensions and body weights were recorded twice a week starting on the first day of treatment. Tumor weights were calculated using the equation $(l \times w^2)/2$, where l and w refer to the larger and smaller dimensions collected at each measurement. When the mean tumor volume reached 140 mm³, vehicle (cremophor:ethanol:5%glucose; 1:1:8) or the test compound (30 mg/kg) was administered to mice (6/group).

As can be seen from Figure 4, compound **091** generated an approximate 70% tumor regression in COLO205 BRAF mutated colon cancer xenografts. In a similar

study with a different dosing schedule, compound **106** caused 70% tumor regression from the first day of treatment in COLO205 BRAF mutated colon cancer xenografts.

Example 6

***In vivo* Antitumor Activity in BxPC-3 Human Pancreatic Cancer Xenografts** (Figure 5). A study was carried out to investigate anticancer activity of compound **091** in human BxPC-3 pancreatic cancer xenografts. The compound was evaluated against subcutaneous-implanted BxPC-3 human pancreatic cancer xenografts in female athymic mice.

The human pancreatic cancer cell line, BxPC-3-T1 (no mutations on B-RAF and Ras genes) was transplanted subcutaneously into female nude mice. The experiment consisted of 8 mice per group on the first day of treatment. The compound was administered intravenously for two weekly rounds of five daily injections (q1d x 5) with a two-day rest period between the rounds. The compound was administered in a vehicle volume of 0.1 mL/10 g of body weight.

The general health of the mice was monitored daily. Tumor dimensions and body weights were recorded twice a week starting on the first day of treatment. Tumor weights were calculated using the equation $(l \times w^2)/2$, where l and w refer to the larger and smaller dimensions collected at each measurement. When the mean tumor volume reached 130 mm³, vehicle (cremophor:ethanol: 5%glucose; 1:1:8) or the test compound (15, 30 mg/kg) was administered to mice (8/group).

As can be seen in Figure 5, 30 mg/kg of compound **091** generated 50% tumor regression.

Example 7

***In vivo* Antitumor Activity in LOX Human Melanoma Xenografts** (Figure 6). A study was carried out to investigate anticancer activity of compounds **091** and **106** in human LOX BRAF mutated melanoma xenografts. The compounds were evaluated against subcutaneous-implanted LOX human melanoma xenografts in female athymic mice.

The human melanoma cell line, LOX which has B-RAF (V600E) mutation was transplanted subcutaneously into female nude mice. The experiment consisted of 8 mice per group on the first day of treatment. Compounds were independently administered

intravenously for two weekly rounds of five daily injections (q1d x 5) with a two-day rest period between the rounds. Compounds were administered in a vehicle volume of 0.1 mL/10 g of body weight.

The general health of the mice was monitored daily. Tumor dimensions and body weights were recorded twice a week starting on the first day of treatment. Tumor weights were calculated using the equation $(l \times w^2)/2$, where l and w refer to the larger and smaller dimensions collected at each measurement. When the mean tumor volume reached 128 mm³, vehicle (cremophor:ethanol:5%glucose; 1:1:8) or one of the test compounds (10, 20, 30, or 40 mg/kg) was administered to mice (8/group).

As can be seen in Figure 6, 10 mg/kg Compound **091** generated 100% tumor regression for the first three weeks. Compound **091** also generated 100% tumor regression at doses of 20, 30, 40 mg/kg in all mice. Mice given **091** were tumor free at the end of the study. Moreover, administration of compound **106** at three dosages, 10, 20, and 40 mg/kg/dose, caused statistically significant anticancer activity. 10 mg/kg of **106** caused 5 out of 8 mice to be tumor free, and 20 mg/kg or 40 mg/kg both caused 8 out of 8 mice to be tumor free at the end of study.

Example 8

Inhibition of B-cell Driven Hematological Malignant Cancer Cell Growth / Non-Hodgkin's Lymphoma (Figure 7). A study was carried out to investigate anticancer activity of compounds **091** and **106** in a panel of human hematologic cancer cell lines *in vitro*. Antiproliferative effects of compounds **091** and **106** were evaluated in 7 human cancer cell lines representing several different cancer types of hematologic malignancy. Cultured human cancer cells were placed into 96-well plates and grown in the absence or continuous presence of 0.3 – 1000 nM test compounds for 96 h. Cell growth was assessed using the MTT assay. IC₅₀ values were determined as the concentration of a substance which inhibits cell growth by 50% compared to untreated cell populations.

As indicated in Figure 7, two cell lines derived from B-cell non-Hodgkin's lymphoma, were sensitive to Compound **091** and cell lines that were derived from leukemia, or B-cells such non-Hodgkin's lymphoma were sensitive to compound **106**.

Example 9

Enhancement of Anticancer Activity of CPT-11 (SN-38) in PANC-1

Pancreatic Cancer Cells (Figure 8). SN-38, active metabolite of CPT-11 is well known to increase NF- κ B activity, which leads to chemo-resistance. To determine whether compounds **091** or **106** in combination with SN-38 would enhance the anticancer activity of chemotherapy due to MEKK1 inhibition, PANC-1 human pancreatic cancer cells were examined.

It is noted that 1 μ M of compound **091** or compound **106** alone did not affect PANC-1 cancer cell growth in separate experiments (data not shown). Antiproliferative effects of SN-38 were evaluated in PANC-1 pancreatic cancer cells in combination with compound **091** or **106**. Cultured human cancer cells were placed into 96-well plates and grown in the absence or continuous presence of 0.03 – 1000 nM SN-38 for 96 h with or without 1 μ M of test compound. Cell growth was assessed using CellTiter-Glo[®] Luminescent Cell Viability Assay (Promega). IC₅₀ values were determined as the concentration of a substance which inhibits cell growth by 50% compared to untreated cell populations.

IC₅₀ values for cell growth inhibition by SN-38 alone was 82.5 nM. 1 μ M of both compound **091** and compound **106** enhanced cytotoxicity of SN-38 in combination setting (e.g., with IC₅₀ value of 16.2 nM). Both compounds **091** and **106** enhance the anticancer activity of CPT-11 (SN-38) in this cell line in two independent experiments. These results suggest that compounds **091** and **106** block chemotherapy-induced NF- κ B activation, leading to enhancement of chemo-sensitivity, although these experiments are still ongoing. Compounds **091** and **106** may enhance anticancer activity in combination setting.

Example 10

Compound 091 Penetrates Blood Brain Barrier (BBB) (Figure 9). Penetration of Compound **091** into the brain tissue was notably high, as measured by the brain/plasma AUC_{0-t} ratio of 3.170. These data suggest that Compound **091** may be useful in treating CNS tumors including glioma *via* the inhibition of a combination of kinases as shown herein, *e.g.*, the inhibition of MEK and TrkB and/or other kinases.

Example 11

Inhibition of Phosphorylation of Flt-3 Receptor Tyrosine Kinase (FLT3)

(Figure 10). A study was carried out to investigate the ability of compound **106** to inhibit phosphorylation of Flt-3 receptor in a FLT3-mutated (AML) cell line *in vitro*. MV-4-11 human acute myeloid leukemia (AML) cells (Catalog # CRL-9591, ATCC, Rockville, MD), which carry an activation mutation of Flt-3, were plated into a 25 mm³ flasks and incubated for 1 day. Then, compound **106** at concentrations of 0.1 nmol/L-300 nmol/L (or culture medium) were added to each flask and cultured for 24 h. The control dish received cultured medium alone. Cells lysates were prepared using ice cold lysis buffer.

In order to detect phospho-Flt-3 protein, MV-4-11 cell lysate proteins were subjected to SDS-PAGE under reducing conditions and immunoblotted with an antibody against phospho-Flt-3 antibody (Catalog # 3464, Cell Signaling Technology, Danvers, MA). Western blots were probed with secondary IRDye antibodies (LI-COR, Inc., Lincoln, NE). Blotted proteins were visualized using the Odyssey infrared imaging system (LI-COR, Inc., Lincoln, NE). Anti-actin antibody (Catalog # sc-8432, Santa Cruz Biotechnologies, Inc., Santa Cruz, CA) was used as a loading control.

MV-4-11 cells showed constitutive phosphorylation of Flt-3 receptor in the absence of compound **106**. The protein level of phospho-Flt-3 was decreased by compound **106** in the MV-4-11 cell line in a concentration dependent manner. 10nmol/L compound **106** completely inhibited constitutive phosphorylation of Flt-3 receptor, showing that compound **106** is a potent inhibitor of Flt-3 receptor tyrosine kinase in whole cells.

Example 12

Inhibition of Cell Proliferation of a FLT3-mutated cell line *in vitro* (Figure

11). A study was carried out to investigate the anticancer activity of compound **106** in a FLT-3 mutated (AML) cell line *in vitro*. The antiproliferative effects of compound **106** were determined in a cell proliferation assay using MV-4-11 AML cells.


MV-4-11 human acute myeloid leukemia (AML) cells, which carry an activation mutation of Flt-3, were plated into 96-well plates and grown in the absence or continuous presence of 0.03 – 1,000 nmol/L compound **106** for 96 h. Cell growth was assessed using CellTiter-Glo[®] Luminescent Cell Viability Assay (Promega, Madison,

WI). Luminescence was read on the EnVision 2102 Multilabel Counter (Perkin-Elmer/Wallac, Waltham, MA). The software, Graphpad Prism (ver.4, San Diego, CA) was used for determination of IC₅₀ values. The raw value for the untreated control represents an average from 4 wells.

MV-4-11 cells which carry an activation mutation of Flt-3 receptor tyrosine kinase were sensitive to compound **106** in the low-nmol/L concentration range. Compound **106** showed antiproliferative activity against FLT-3 mutated human cancer cells *in vitro*, with an IC₅₀ value in the low-nmol/L range.

CLAIMS:

1. Use of a SrcTK/MEK inhibitor for treating a SrcTK/MEK associated cancer in a subject in need thereof, wherein the SrcTK/MEK inhibitor is a compound of formula (II):




(II)

wherein

R_3 is $-NHR_b$, and R_b is C_1 - C_3 alkyl substituted with 0, 1, or 2 hydroxyl moieties; or a pharmaceutically acceptable salt or ester thereof.

2. The use of claim 1, wherein the SrcTK/MEK inhibitor inhibits the activity of at least one member of the Src tyrosine kinase family by at least about 75% at a concentration of about $0.1\mu\text{M}$.
3. The use of claim 1 or 2, wherein the SrcTK/MEK inhibitor inhibits the activity of at least five members of the Src tyrosine kinase family by at least about 50% at a concentration of about $0.1\mu\text{M}$.
4. The use of claim 3, wherein the members of the Src tyrosine kinase family include cSrc, Fyn, Lyn, Lck and Yes.


5. The use of any one of claims 1-4, wherein the SrcTK/MEK inhibitor inhibits the activity of MEK1 by at least about 50% at a concentration of about 0.1 μ M.
6. The use of any one of claims 1-5, wherein the SrcTK/MEK associated cancer is a cancer selected from the group consisting of chronic myeloid leukemia and colorectal cancer.
7. The use of any one of claims 1-6, wherein the cancer is resistant to imatinib mesylate.
8. The use of any one of claims 1-7, wherein the SrcTK/MEK inhibitor also inhibits the activity of Bcl-Abl, and the SrcTK/MEK associated cancer is leukemia.
9. The use of any one of claims 1-8, wherein the SrcTK/MEK inhibitor also inhibits the activity of TrkB.
10. Use of a TrkB/MEK inhibitor for treating a TrkB/MEK associated cancer in a subject in need thereof, wherein the TrkB/MEK inhibitor is a compound of formula (II):



wherein

R_3 is $-NHR_b$, and R_b is C_1 - C_3 alkyl substituted with 0, 1, or 2 hydroxyl moieties; or a pharmaceutically acceptable salt or ester thereof.

11. The use of claim 10, wherein the TrkB/MEK inhibitor inhibits the activity of TrkB by at least about 75% at a concentration of about 0.1 μ M.
12. The use of claim 10 or 11, wherein the TrkB/MEK inhibitor inhibits the activity of MEK1 by at least about 55% at a concentration of about 0.1 μ M.
13. The use of any one of claims 10-12, wherein the TrkB/MEK associated cancer is a cancer selected from the group consisting of pancreatic cancer, neural cancer, glioma, neuroblastoma and retinoblastoma.
14. The use of any one of claims 10-13, wherein the TrkB/MEK inhibitor also inhibits the activity of at least one member of the Src tyrosine kinase family.
15. Use for inhibition of a metastatic process in a subject in need thereof of a compound of formula (II):




wherein

R_3 is $-NHR_b$, and R_b is C_1 - C_3 alkyl substituted with 0, 1, or 2 hydroxyl moieties;

or a pharmaceutically acceptable salt or ester thereof.

16. Use for treating a B-RAF mutated cancer in a subject in need thereof

of a compound of formula (II):




(II)

wherein

R₃ is -NHR_b, and R_b is C₁-C₃ alkyl substituted with 0, 1, or 2 hydroxyl moieties;

or a pharmaceutically acceptable salt or ester thereof.

17. The use of claim 16, wherein the B-RAF mutated cancer is a B-RAF V600E mutated cancer.
18. The use of claim 16 or 17, wherein the B-RAF mutated cancer is a cancer selected from the group consisting of ovarian cancer, thyroid cancer, colorectal cancer and melanoma.
19. Use for treating a FLT3 mutated cancer in a subject in need thereof of at least one compound of formula (II):




(II)

wherein

R_3 is $-NHR_b$, and R_b is C_1 - C_3 alkyl substituted with 0, 1, or 2 hydroxyl moieties;
or a pharmaceutically acceptable salt or ester thereof.

20. The use of claim 19, wherein the FLT3 mutated cancer comprises an internal tandem duplication.
21. The use of claim 19 or 20, wherein the FLT3 mutated cancer comprises a D835Y, Y842C, K663Q or V592A mutation.
22. The use of any one of claims 19-21, wherein the FLT3 mutated cancer is a hematologic cancer.
23. The use of any one of claims 19-22, wherein the FLT3 mutated cancer is leukemia.
24. The use of claim 23, wherein the FLT3 mutated cancer is acute myeloid leukemia.
25. Use for treating a central nervous system tumor in a subject in need thereof of a compound of formula (II):



wherein


R_3 is $-NHR_b$, and R_b is C_1 - C_3 alkyl substituted with 0, 1, or 2 hydroxyl moieties;

or a pharmaceutically acceptable salt or ester thereof,

wherein the compound of formula (II) is capable of penetrating the blood brain barrier.


26. The use of claim 25, wherein at least about 20% of the compound of formula (II) is capable of penetrating the blood brain barrier, based upon a ratio of the amount of the compound of formula (II) in the brain to the amount of the compound of formula (II) in the plasma in an animal.
27. The use of claim 25, wherein at least about 50% of the compound of formula (II) is capable of penetrating the blood brain barrier, based upon a ratio of amount of a compound of formula (II) in the brain to amount of a compound of formula (II) in the plasma in an animal.
28. The use of any one of claims 25-27, wherein the central nervous system tumor is a tumor selected from the group consisting of a brain tumor, a glioma and a neuroblastoma.

29. The use of any one of claims 1, 10, 15, 16, 19 and 25, wherein the compound is at least one compound selected from the group consisting of:




and pharmaceutically acceptable salts and esters thereof.

30. The use of claim 29, wherein the compound is:




or a pharmaceutically acceptable salt thereof.

31. The use of claim 29, wherein the compound is:













32. The use of any one of claims 1, 10, 15, 16, 19 and 25, wherein the compound is at least one compound selected from the group consisting of:





















and pharmaceutically acceptable salts and esters thereof.

33. The use of any one of claims 1, 10, 15, 16, 19, 25, 30 and 31, wherein the use further comprises a second chemotherapeutic drug.
34. The use of any one of claims 1, 10, 15, 16, 19, 25, 30 and 31, wherein the compound of formula (II) is for intravenous administration.
35. The use of any one of claims 1, 10, 15, 16, 19, 25, 30 and 31, wherein the compound of formula (II) is for administration at a dosage between about 0.10 mg/kg to about 25 mg/kg of body weight.

36. A compound selected from the group consisting of:


 (Compound 013)	 (Compound 048)
 (Compound 014)	 (Compound 054)
 (Compound 015)	 (Compound 065)
 (Compound 016)	 (Compound 076)
 (Compound 018)	 (Compound 114)

 (Compound 019)	 (Compound 122)
 (Compound 022)	 (Compound 127)
 (Compound 024)	 (Compound 137)
 (Compound 025)	 (Compound 144)
 (Compound 034)	 (Compound 155)

 (Compound 041)	 (Compound 156)
 (Compound 045)	 (Compound 157)
 (Compound 046)	 (Compound 158)
 (Compound 047)	 (Compound 114)

and pharmaceutically acceptable salts and esters thereof.


37. The compound of claim 36, which is



(Compound **013**),

or a pharmaceutically acceptable salt or ester thereof.


38. The compound of claim 36, which is



(Compound **014**),

or a pharmaceutically acceptable salt or ester thereof.


39. The compound of claim 36, which is



(Compound **015**),

or a pharmaceutically acceptable salt or ester thereof.


40. The compound of claim 36, which is



(Compound **016**),

or a pharmaceutically acceptable salt or ester thereof.


41. The compound of claim 36, which is



(Compound **018**),

or a pharmaceutically acceptable salt or ester thereof.


42. The compound of claim 36, which is



(Compound **019**),

or a pharmaceutically acceptable salt or ester thereof.


43. The compound of claim 36, which is



(Compound **022**),

or a pharmaceutically acceptable salt or ester thereof.


44. The compound of claim 36, which is



(Compound **024**),

or a pharmaceutically acceptable salt or ester thereof.


45. The compound of claim 36, which is



(Compound **025**),

or a pharmaceutically acceptable salt or ester thereof.


46. The compound of claim 36, which is



(Compound **034**),

or a pharmaceutically acceptable salt or ester thereof.


47. The compound of claim 36, which is



(Compound **041**),

or a pharmaceutically acceptable salt or ester thereof.


48. The compound of claim 36, which is



(Compound **045**),

or a pharmaceutically acceptable salt or ester thereof.


49. The compound of claim 36, which is



(Compound **046**),

or a pharmaceutically acceptable salt or ester thereof.


50. The compound of claim 36, which is



(Compound **047**),

or a pharmaceutically acceptable salt or ester thereof.


51. The compound of claim 36, which is



(Compound **048**),

or a pharmaceutically acceptable salt or ester thereof.


52. The compound of claim 36, which is



(Compound **054**),

or a pharmaceutically acceptable salt or ester thereof.


53. The compound of claim 36, which is



(Compound **065**),

or a pharmaceutically acceptable salt or ester thereof.


54. The compound of claim 36, which is



(Compound **076**),


or a pharmaceutically acceptable salt or ester thereof.

55. The compound of claim 36, which is

(Compound **114**),


or a pharmaceutically acceptable salt or ester thereof.

56. The compound of claim 36, which is

(Compound **122**),


or a pharmaceutically acceptable salt or ester thereof.

57. The compound of claim 36, which is

(Compound **127**),

or a pharmaceutically acceptable salt or ester thereof.


58. The compound of claim 36, which is



(Compound **137**),

or a pharmaceutically acceptable salt or ester thereof.


59. The compound of claim 36, which is



(Compound **144**),

or a pharmaceutically acceptable salt or ester thereof.


60. The compound of claim 36, which is



(Compound **155**),

or a pharmaceutically acceptable salt or ester thereof.


61. The compound of claim 36, which is



(Compound **156**),

or a pharmaceutically acceptable salt or ester thereof.


62. The compound of claim 36, which is



(Compound **157**),

or a pharmaceutically acceptable salt or ester thereof.


63. The compound of claim 36, which is



(Compound **158**),

or a pharmaceutically acceptable salt or ester thereof.

64. The compound of claim 36, which is




(Compound **114**),

or a pharmaceutically acceptable salt or ester thereof.

1/12

RAS-MAPK Signaling Pathway

*Fig. 1*

2/12

Biochemical MEK1 Assays

<i>Compound</i>	<i>Biochemical MEK1 inhibition (IC₅₀; nM)</i>	
	<i>10 μM ATP</i>	<i>100 μM ATP</i>
091	10	70

<i>Compound</i>	<i>IC₅₀(nM)</i>	<i>Compound</i>	<i>IC₅₀(nM)</i>	
Compound 013	120	Compound 054	>330	
Compound 014	471 946	Compound 065	187	
Compound 015	91 101	Compound 069	37 9.5	
Compound 016	356 640	Compound 076	657	
Compound 018	-	Compound 091	34 37 22 10.9 24 2.8	24.4 51 33 113 73 71
Compound 019	658	Compound 092	27 4.8	14.8 715
Compound 022	846	Compound 106	107 13 172	45 71
Compound 024	187	Compound 114	56 101	
Compound 025	484	Compound 122	356	
Compound 029	91 198	Compound 127	>330	
Compound 034	172	Compound 137	182	
Compound 041	756	Compound 144	75 163	104 117
Compound 045	318	Compound 155	433 421	
Compound 046	657	Compound 156	79	
Compound 047	99 310	Compound 157	240 421	
Compound 048	366	Compound 158	145	

Fig. 2A

3/12
Kinase inhibition

Kinase	Class	Importance in cancer	Compound 091 IC50 (nM)
Abl	TK	Leukemia	214
c-Src	Src TK family	Colon, Pancreatic	91
Flt3	Receptor TK	Leukemia	91
Fyn	Src TK family	Leukemia	47
Lyn	Src TK family	Leukemia	35
TrkB	Receptor TK	Prostate, Pancreatic Glioma	26
MEK1	MAPKK	Proliferation	10
MEKK1	MAPKKK	Antiapoptosis MM/Lymphoma	46*
KDR	Receptor TK	Angiogenesis	500*
* Result from separate study			

<i>Percent Kinase Activity Remaining</i>				
Kinase	Compound 091		Compound 106	
	0.1 μ M	1 μ M	0.1 μ M	1 μ M
Abl	69	22	51	15
cSRC	47	5	31	1
Flt3	48	8	33	6
Fyn	25	2	24	0
Lck	38	3	27	0
Lyn	15	2	45	2
MEK1	30	4	-	-
TrkB	15	4	24	4
Yes	46	7	54	8

Fig. 2B


4/12

Inhibition of B-RAF Mutated Cancer Cell Growth

Cell line	Description	K-ras mutation	B-RAF mutation	Mean IC ₅₀ , nmol/L	
				Cpd 106	Cpd 091
AU-565	breast cancer		WT	>10000	>10000
MCF-7			WT	>10000	>10000
MDA-MB-231		G13D	WT	156	119
MDA-MB-435			V600E	71	50
DU4475			V600E	52	30
SK-MEL-2	melanoma	Q61R (N-ras)	WT	135	113
SK-MEL-3			V600E	24	19
SK-MEL-24			V600E	66	53
SK-MEL-31			V600E	225	209
HCT-116	colon cancer	G13D	WT	423	257
DLD-1		G13D	WT	649	466
LoVo		G13D	WT	132	90
SW-620		G12V	WT	282	207
COLO205		WT	V600E	43	27
HT-29		WT	V600E	84	65
Mia PaCa-2	pancreatic cancer	G12C	WT	734	582
PANC-1		G12D	WT	>10000	>10000
BxPC-3		WT	WT	—	412
MES-SA	sarcoma		WT	7490	5780
MES-SA/Dx5-Rx1	sarcoma (P-glycoprotein overex.)		WT	1590	908
IMR-90	Non-dividing fibroblast			—	>10000
“WT” = wild type; “—” = not tested					

Fig. 3

5/12

Antitumor Activity in Colo-205 Colon Xenografts*Fig. 4*

6/12

*In vivo Antitumor Activity in BxPC-3 Human
Pancreatic Cancer Xenografts*





Fig. 5

7/12

Antitumor Activity in LOX Xenografts*Fig. 6*

8/12

***Inhibition of B-cell Driven Hematological Malignant
Cancer Cell Growth***

Cell line	Type	Compound 091 IC ₅₀ (μM)	Compound 106 IC ₅₀ (μM)
HL-60	Acute Promyelocytic Leukemia	0.05 ± 0.001	0.06 ± 0.004
THP-1	Acute Monocytic Leukemia	0.09 ± 0.012	0.12 ± 0.008
K562	Chronic Myelogenous Leukemia	1.43 ± 0.20	1.80 ± 0.09
U266	MM	1.00 ± 0.58	0.95 ± 0.22
RPMI- 8226	MM	>4.83 ± 2.65	>7.05 ± 2.95
RL	Non-Hodgkin's Lymphoma (B-lymphoblast)	1.62 ± 0.34	1.65 ± 0.37
NALM-6	Non-Hodgkin's Lymphoma (Pre-B lymphoma)	0.65 ± 0.20	0.62 ± 0.11

Fig. 7

9/12

***Enhancement of Anticancer Activity of CPT-11
(SN-38) in PANC-1 Pancreatic Cancer Cells***




Fig. 8

10/12


Penetration of Blood Brain Barrier

Penetration of compound 091 into the brain tissue,
as measured by the brain/plasma AUC_{0-t} ratio, is as follows:

Species	Compound	Dose	Brain penetration ratio (AUC _{0-t} ,brain/AUC _{0-t} ,plasma)
Mouse (n=10)	091	20 mg/kg, <i>i.v.</i>	3.170 ± 1.289

Fig. 9

11/12

*Fig. 10*

12/12

*Inhibition of MV-4-11 cell proliferation
by compound 106*




Fig. 11

