

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
1 July 2004 (01.07.2004)

PCT

(10) International Publication Number
WO 2004/054566 A1

(51) International Patent Classification⁷: **A61K 31/197**,
A61P 1/00, 25/18, 29/00, 43/00, 5/00

Research and Development, Ann Arbor Laboratories, 2800
Plymouth Road, Ann Arbor, MI 48105 (US).

(21) International Application Number:
PCT/IB2003/005712

(74) Agents: **LUMB, J., Trevor** et al.; c/o David J. Wood,
Pfizer Global Research and Development, UK Patent
Department, Ramsgate Road, Sandwich, Kent CT13 9NJ
(GB).

(22) International Filing Date: 3 December 2003 (03.12.2003)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,
MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT,
RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(26) Publication Language: English

(30) Priority Data:
60/433,491 13 December 2002 (13.12.2002) US
60/487,740 16 July 2003 (16.07.2003) US

(71) Applicant (*for all designated States except US*):
WARNER-LAMBERT COMPANY LLC [US/US];
201 Tabor Road, Morris Plains, NJ 07950 (US).

(84) Designated States (*regional*): ARIPO patent (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): **DOOLEY, David,**
James [US/US]; Pfizer Global Research and Develop-
ment, Ann Arbor Laboratories, 2800 Plymouth Road, Ann
Arbor, MI 48105 (US). **TAYLOR, Charles, Price, Jr.**
[US/US]; Pfizer Global Research and Development, Ann
Arbor Laboratories, 2800 Plymouth Road, Ann Arbor, MI
48105 (US). **THORPE, Andrew, John** [GB/US]; Pfizer
Global Research and Development, Ann Arbor Laborato-
ries, 2800 Plymouth Road, Ann Arbor, MI 48105 (US).
WUSTROW, David, Juergen [US/US]; Pfizer Global

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: PREGABALIN DERIVATIVES FOR THE TREATMENT OF FIBROMYALGIA AND OTHER DISORDERS



(57) Abstract: This invention relates to a method of treating a disorder selected from OCD, agoraphobia, agoraphobia without history of panic disorder, specific phobia, social phobia, PTSD, restless legs syndrome, premenstrual dysphoric disorder, hot flashes, and fibromyalgia by administering a compound of the formula (1), or a pharmaceutically acceptable salt thereof, wherein: a) R¹ is hydrogen, straight or branched alkyl of from 1 to 6 carbon atoms or phenyl; and b) R² is straight or branched alkyl of from 4 to 8 carbon atoms, straight or branched alkenyl of from 2 to 8 carbon atoms, cycloalkyl of from 3 to 7 carbon atoms, alkoxy of from 1

to 6 carbon atoms, -alkylcycloalkyl, -alkylalkoxy, -alkyl OH, -alkylphenyl, -alkylphenoxy, or -substituted phenyl. The invention also relates to a method of treating the above disorders by administering the compound (3S, 5R)-3-Aminomethyl-5-methyl-octanoic acid.

PREGABALIN DERIVATIVES FOR THE TREATMENT OF FIBROMYALGIA AND OTHER DISORDERS

BACKGROUND OF THE INVENTION

5 This invention relates to the use of certain alpha2delta ligand for the treatment of fibromyalgia and other central nervous system disorders. Fibromyalgia (FM) is a chronic syndrome characterized mainly by widespread pain, unrefreshing sleep, disturbed mood, and fatigue. The main symptoms fibromyalgia include pain, sleep, mood disturbances and fatigue. Syndromes commonly associated with fibromyalgia include irritable bowel syndrome, and
10 migraine headaches, among others. Success of treating fibromyalgia with a single pharmacological agent has been characterized as modest and results of clinical trials have been characterized as disappointing. It is believed that based on current understanding of the mechanisms and pathways involved in fibromyalgia, multiple agents will be required, aimed at the major symptoms of pain, disturbed
15 sleep, mood disturbances, and fatigue. Fibromyalgia patients are often sensitive to side effects of medications, a characteristic perhaps related to the pathophysiology of this disorder (Barkhuizen A, Rational and Targeted pharmacologic treatment of fibromyalgia. Rheum Dis Clin N Am 2002; 28: 261-290; Leventhal LJ. Management of fibromyalgia. Ann Intern Med 1999;131:850-8).

20 While fibromyalgia is a complex disorder with multiple facets, this complexity can be well assessed (Yunus MB, A comprehensive medical evaluation of patients with fibromyalgia syndrome, Rheum Dis N Am 2002; 28:201-217). The diagnosis of FM is usually based on the 1990 recommendations of the American College of Rheumatology classification criteria (Bennett RM, The rational management of fibromyalgia patients. Rheum Dis Clin N Am 2002;
25 28: 181-199; Wolfe F, Smythe HA, Yunus MB, Bennett RM, Bombardier C, Goldenberg DL, et al. The American College of Rheumatology 1990 criteria for the classification of fibromyalgia: Report of the Multicenter Criteria Committee. Arthritis Rheum 1990; 33:160-72). Evaluation, management, and
30 pharmacological treatment of fibromyalgia have been reviewed (Barkhuizen A,

-2-

Rational and Targeted pharmacologic treatment of fibromyalgia. *Rheum Dis Clin N Am* 2002; Buskila D, Fibromyalgia, chronic fatigue syndrome and myofascial pain syndrome. *Current opinions in Rheumatology* 2001; 13: 117-127; Leventhal LJ. Management of fibromyalgia. *Ann Intern Med* 1999;131:850-8; Bennett RM, 5 The rational management of fibromyalgia patients. *Rheum Dis Clin N Am* 2002; 28: 181-199; Yunus MB, A comprehensive medical evaluation of patients with fibromyalgia syndrome, *Rheum Dis N Am* 2002; 28:201-217).

Restless Legs Syndrome (RLS) has been characterized by the minimal diagnostic criteria: (a) desire to move the extremities, often associated with 10 paresthesias/dysesthesias; (b) motor restlessness; (c) worsening of symptoms at rest with at least temporary relief by activity, and (d) worsening of symptoms in the evening or night. Other features commonly seen in RLS include sleep disturbance, periodic limb movements in sleep and similar involuntary movements while awake, a normal neurological examination in the idiopathic form, a 15 tendency for the symptoms to be worse in middle to older age, and, in some cases, a family history suggestive of an autosomal dominant mode of inheritance. See Walters AS; Toward a better definition of the restless legs syndrome; The International Restless Legs Syndrome Study Group; *Mov Disord.* (1995) 10(5): 634-42. See also, Bhatia M and Bhowmik D; Restless legs syndrome in 20 maintenance haemodialysis patients; *Nephrol Dial Transplant* (2003) 18: 217. Restless legs syndrome can be a primary disorder, or a secondary disorder associated with, for example, renal insufficiency, heredity, pregnancy, poliomyelitis, infectious diseases, avitaminosis, different types of anemia, diabetes, and certain drugs (e.g. prochloroperazine, lithium, and mianserin).

25 Gabapentin, pregabalin and other alpha2delta ligands including 4H-[1,2,4]oxadiazol-5-one, C-[1-(1H-Tetrazol-5-ylmethyl)-cycloheptyl]-methylamine, (3S,4S)-(1-Aminomethyl-3,4-dimethyl-cyclopentyl)-acetic acid, (1 α ,3 α ,5 α)-(3-amino-methyl-bicyclo[3.2.0]hept-3-yl)-acetic acid , and (3S,5R)-3-aminomethyl-5-methyl-heptanoic acid , and pharmaceutically acceptable salts and 30 solvates thereof, are referred to in United States Patent 4,024,175; U.S. Patent 4,087,544; U.S. Patent 6,306,910; WO9921824, WO0190052, WO0128978,

-3-


EP0641330, WO9817627, and WO0076958,. The foregoing patents and applications are incorporated herein by reference in their entirety.

The compounds employed in the methods of the present invention are mono- and disubstituted 3-propyl gamma-aminobutyric acids. United States
5 Patent Applications 10/009,938 filed December 10, 2001, and 10/324,929 filed December 20, 2002 refer to the compounds employed in the methods of the present invention discussed below and disclose various utilities for these. The entire contents of applications 10/009,938 and 10/324,929 are hereby incorporated herein by reference.

10

SUMMARY OF THE INVENTION

This invention relates to a method for treating a disorder in a mammal, including a human, comprising administering to said mammal a therapeutically effective amount of a compound of formula 1



15

or a pharmaceutically acceptable salt thereof, wherein:

R¹ is hydrogen, straight or branched alkyl of from 1 to 6 carbon atoms or phenyl;

R² is straight or branched alkyl of from 4 to 8 carbon atoms, straight or branched alkenyl of from 2 to 8 carbon atoms, cycloalkyl of from 3 to 7 carbon atoms, alkoxy of from 1 to 6 carbon atoms, -alkylcycloalkyl,, -alkylalkoxy, -alkyl OH,

20

-alkylphenyl, -alkylphenoxy, and -substituted phenyl, and wherein said disorder is selected from OCD, phobias, PTSD, restless legs syndrome, premenstrual dysphoric disorder, hot flashes, and fibromyalgia. This method is hereinafter also referred to as the "inventive method".

25

The invention also relates to a method for treating a disorder in a mammal, including a human, comprising administering to said mammal a therapeutically

-4-

effective amount of the compound (3S, 5R)-3-Aminomethyl-5-methyl-octanoic acid or a pharmaceutically acceptable salt thereof wherein said disorders is selected from OCD, phobias, PTSD, restless legs syndrome, premenstrual dysphoric disorder, hot flashes, and fibromyalgia. The compound (3S, 5R)-3-Aminomethyl-5-methyl-octanoic acid is hereinafter also referred to as "Compound A".

The invention also relates to a method for treating fibromyalgia and a concomitant disorder in a mammal, including a human, comprising administering to said mammal a therapeutically effective amount of a compound of the formula 1 as described above wherein said concomitant disorder is selected from migraine headaches, temporomandibular joint dysfunction, dysautonomia, endocrine dysfunction, dizziness, cold intolerance, chemical sensitivity, sicca symptoms, cognitive dysfunction, generalized anxiety disorder, premenstrual dysphoric disorder, irritable bowel syndrome, functional abdominal pain, neuropathic pain, somatoform disorders, OCD, phobias, and PTSD.

The invention also relates to a method for treating fibromyalgia and a concomitant disorder in a mammal, including a human, comprising administering to said mammal a therapeutically effective amount of Compound A or a pharmaceutically acceptable salt thereof wherein said concomitant disorder is selected from the concomitant disorders described above.

The invention also relates to a method of increasing slow wave sleep in a human subject being treated with an active pharmaceutical agent that decreases slow wave sleep comprising administering to a human subject in need of such treatment:

(a) a compound of the formula 1, as described above, or a pharmaceutically acceptable salt thereof; and

(b) a human growth hormone or human growth hormone secretagogue, or a pharmaceutically acceptable salt thereof;

wherein the amounts of the active agents "a" and "b" are chosen so as to render the combination effective in increasing slow wave sleep.

The invention also relates to a method of increasing slow wave sleep in a human subject being treated with an active pharmaceutical agent that decreases

-5-

slow wave sleep comprising administering to a human subject in need of such treatment:

(a) Compound A; and

(b) a human growth hormone or human growth hormone secretagogue, or a pharmaceutically acceptable salt thereof;

wherein the amounts of the active agents "a" and "b" are chosen so as to render the combination effective in increasing slow wave sleep.

The invention also relates to a method of increasing slow wave sleep in a human subject comprising administering to a human subject in need of such treatment:

(a) a compound of the formula 1, as described above, or a pharmaceutically acceptable salt thereof; and

(b) a human growth hormone or human growth hormone secretagogue, or a pharmaceutically acceptable salt thereof;

wherein the amounts of the active agents "a" and "b" are chosen so as to render the combination effective in increasing slow wave sleep.

The invention also relates to a method of increasing slow wave sleep in a human subject comprising administering to a human subject in need of such treatment:

(a) Compound A or a pharmaceutically acceptable salt thereof; and

(b) a human growth hormone or human growth hormone secretagogue, or a pharmaceutically acceptable salt thereof;

wherein the amounts of the active agents "a" and "b" are chosen so as to render the combination effective in increasing slow wave sleep.

A more specific embodiment of this invention relates to any one of the above method for increasing slow wave sleep wherein the human growth hormone secretagogue that is employed is 2-amino-N-[2-(3a-benzyl-2-methyl-3-oxo-2,3,3a,4,6,7-hexahydro-pyrazole[4,3-c]pyridin-5-yl)-1-benzyloxymethyl-2-oxo-ethyl]-2-methyl-propionamide.

This invention also relates to any one of the above methods for increasing slow wave sleep in a human subject being treated with an active pharmaceutical agent that decreases slow wave sleep, such as morphine or another opioid analgesic agent, comprising administering to such human subject an amount of a compound of

-6-

the formula **1**, as defined above, compound A, or a pharmaceutically acceptable salt thereof, that is effective in increasing slow wave sleep.

In one embodiment, the invention relates to any one of the methods described above wherein the disorder being treated is a phobia selected from agoraphobia, agoraphobia without history of panic disorder, specific phobia, and social phobia.

In another embodiment, the invention relates to any one of the methods described above wherein the compound administered is Compound A or a pharmaceutically acceptable salt thereof, and wherein the disorder being treated is OCD, PTSD, or a phobia.

In another embodiment, the invention relates to any one of the methods described above wherein the compound administered is Compound A or a pharmaceutically acceptable salt thereof, and wherein the disorder being treated is a phobia selected from agoraphobia and specific phobias.

In another embodiment, the invention relates to any one of the methods described above wherein the compound administered is Compound A or a pharmaceutically acceptable salt thereof, and the disorder being treated is fibromyalgia.

In another embodiment, the invention relates to any one of the methods described above wherein the compound administered is selected from 3-Aminomethyl-5-methylheptanoic acid; (3S,5R)-3-Aminomethyl-5-methylheptanoic acid; 3-Aminomethyl-5-methyl-octanoic acid; 3-Aminomethyl-4,5-dimethyl-hexanoic acid; (3S,4S)-3-Aminomethyl-4,5-dimethyl-hexanoic acid; (3R,4R)-3-Aminomethyl-4,5-dimethyl-hexanoic acid MP; 3-Aminomethyl-4-isopropyl-hexanoic acid; 3-Aminomethyl-4-isopropyl-heptanoic acid; (3S,5S)-3-Aminomethyl-6-fluoro-5-methyl-hexanoic acid; (3S,5S)-3-Aminomethyl-7-fluoro-5-methyl-heptanoic acid; (3S,5S)-3-Aminomethyl-7,7,7-trifluoro-5-methyl-heptanoic acid; (3S,5R)-3-Aminomethyl-8,8,8-trifluoro-5-methyl-octanoic acid; (3S,5S)-3-Aminomethyl-5,6-dimethyl-heptanoic acid; (3R,4R,5R)-3-Aminomethyl-4,5-dimethyl-heptanoic acid; and (3R,4R,5R)-3-Aminomethyl-4,5-dimethyl-octanoic acid or a pharmaceutically acceptable salt thereof, and wherein said disorder is selected from OCD, agoraphobia, agoraphobia without history of panic disorder, specific phobia, social phobia, PTSD, and fibromyalgia. The

-7-

compounds listed immediately above in the foregoing method are hereinafter also referred to as "the Group A compounds".

5 In another embodiment, the invention relates to any one of the methods described above wherein the disorder being treated is OCD or PTSD and the compound that is administered is a Group A compound or a pharmaceutically acceptable salt thereof.

10 In another embodiment, the invention relates to any one of the methods described above wherein the disorder being treated is fibromyalgia and the compound or salt that is administered is a Group A compound or a pharmaceutically acceptable salt thereof.

In another embodiment, the invention relates to any one of the methods described above wherein the disorder being treated is fibromyalgia, and wherein the compound administered is Compound A or a pharmaceutically acceptable salt thereof.

15 Another embodiment of the invention relates to any one of the above methods for treating fibromyalgia and a concomitant disorder wherein said concomitant disorder is generalized anxiety disorder, dysphoric dysthemia, irritable bowel syndrome, functional abdominal pain, neuropathic pain, a somatoform disorder, or migraine headache.

20 This invention also relates to a method of treating a disorder or condition selected from acute pain, chronic pain, pain resulting from soft tissue and peripheral damage such as acute trauma; complex regional pain syndrome also referred to as reflex sympathetic dystrophy; postherpetic neuralgia, occipital neuralgia, trigeminal neuralgia, segmental or intercostal neuralgia and other
25 neuralgias; pain associated with osteoarthritis and rheumatoid arthritis; musculo-skeletal pain such as pain associated with strains, sprains and trauma such as broken bones; spinal pain, central nervous system pain such as pain due to spinal cord or brain stem damage; lower back pain, sciatica, dental pain, myofascial pain syndromes, episiotomy pain, gout pain, and pain resulting from burns; deep and
30 visceral pain, such as heart pain; muscle pain, eye pain, inflammatory pain, orofacial pain, for example, odontalgia; abdominal pain, and gynecological pain, for example, dysmenorrhoea, labour pain and pain associated with endometriosis; somatogenic pain; pain associated with nerve and root damage, such as pain

-8-

associated with peripheral nerve disorders, for example, nerve entrapment, brachial plexus avulsions, and peripheral neuropathies ; pain associated with limb amputation, tic douloureux, neuroma, or vasculitis; diabetic neuropathy, chemotherapy-induced-neuropathy, acute herpetic and postherpetic neuralgia; atypical facial pain, nerve root damage, neuropathic lower back pain, HIV related neuropathic pain, cancer related neuropathic pain, diabetes related neuropathic pain and arachnoiditis, trigeminal neuralgia, occipital neuralgia, segmental or intercostal neuralgia, HIV related neuralgias and AIDS related neuralgias and other neuralgias; allodynia, hyperalgesia, burn pain, idiopathic pain, pain caused by chemotherapy; occipital neuralgia, psychogenic pain, brachial plexus avulsion, pain associated with restless legs syndrome; pain associated with gallstones; pain caused by chronic alcoholism or hypothyroidism or uremia or vitamin deficiencies; neuropathic and non-neuropathic pain associated with carcinoma, often referred to as cancer pain, phantom limb pain, functional abdominal pain, headache, including migraine with aura, migraine without aura and other vascular headaches, acute or chronic tension headache, sinus headache and cluster headache; temperomandibular pain and maxillary sinus pain; pain resulting from ankylosing spondylitis and gout; pain caused by increased bladder contractions; pain associated with gastrointestinal (GI) disorders, disorders caused by *helicobacter pylori* and diseases of the GI tract such as gastritis, proctitis, gastroduodenal ulcers, peptic ulcers, dyspepsia, disorders associated with the neuronal control of viscera, ulcerative colitis, chronic pancreatitis, Crohn's disease and emesis; post operative pain, scar pain, and chronic non-neuropathic pain such as pain associated with HIV, anthralgia and myalgia, vasculitis and fibromyalgia in a mammal, comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof.

This invention also relates to a method of treating a disorder or condition selected from the group consisting of mood disorders, such as depression, or more particularly, depressive disorders, for example, single episodic or recurrent major depressive disorder, severe unipolar recurrent major depressive episodes, dysthymic disorder, depressive neurosis and neurotic depression, melancholic depression including anorexia, weight loss, insomnia, early morning waking or

psychomotor retardation, atypical depression (or reactive depression) including increased appetite, hypersomnia, psychomotor agitation or irritability; treatment resistant depression; seasonal affective disorder and pediatric depression; premenstrual syndrome, premenstrual dysphoric disorder, hot flashes, bipolar disorders or manic depression, for example, bipolar I disorder, bipolar II disorder and cyclothymic disorder; seasonal affective disorder, conduct disorder and disruptive behavior disorder; stress related somatic disorders and anxiety disorders, such as childhood anxiety disorder, panic disorder with or without agoraphobia, phobia including agoraphobia without history of panic disorder and specific phobias (*e.g.*, specific animal phobias), social anxiety disorder, social phobia, obsessive-compulsive disorder(OCD), autism and associated disorders including pervasive developmental delay, mood disorders associated with psychotic disorders such as acute mania and depression associated with bipolar disorder, mood disorders associated with schizophrenia, behavioral disturbances associated with mental retardation, autistic disorder, conduct disorder and disruptive behavior disorder, borderline personality disorder, psychotic episodes of anxiety, and anxiety associated with psychosis; stress disorders including post-traumatic stress disorder (PTSD) and acute stress disorder, and generalized anxiety disorder in a mammal, comprising administering to a mammal in need of such treatment a therapeutically effective amount of therapeutically effective amount of a compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof.

It will be appreciated that for the treatment of depression or anxiety, a compound employed in the methods of the present invention may be used in conjunction with other antidepressant or anti-anxiety agents. Suitable classes of antidepressant agents include norepinephrine reuptake inhibitors, selective serotonin reuptake inhibitors (SSRIs), monoamine oxidase inhibitors (MAOIs), reversible inhibitors of monoamine oxidase (RIMAs), serotonin and noradrenaline reuptake inhibitors (SNRIs), corticotropin releasing factor (CRF) antagonists, α -adrenoreceptor antagonists and atypical antidepressants. Suitable norepinephrine reuptake inhibitors include tertiary amine tricyclics and secondary amine tricyclics. Suitable examples of tertiary amine tricyclics include amitriptyline,

-10-

clomipramine, doxepin, imipramine and trimipramine, and pharmaceutically acceptable salts thereof. Suitable examples of secondary amine tricyclics include amoxapine, desipramine, maprotiline, nortriptyline and protriptyline, and pharmaceutically acceptable salts thereof. Suitable selective serotonin reuptake inhibitors include fluoxetine, fluvoxamine, paroxetine and sertraline, and pharmaceutically acceptable salts thereof. Suitable monoamine oxidase inhibitors include isocarboxazid, phenelzine, tranylcypromine and selegiline, and pharmaceutically acceptable salts thereof. Suitable reversible inhibitors of monoamine oxidase include moclobemide, and pharmaceutically acceptable salts thereof. Suitable serotonin and noradrenaline reuptake inhibitors of use in the present invention include venlafaxine, and pharmaceutically acceptable salts thereof. Suitable CRF antagonists include those compounds described in International Patent Application Nos. WO 94/13643, WO 94/13644, WO 94/13661, WO 94/13676 and WO 94/13677. Suitable atypical anti-depressants include bupropion, lithium, nefazodone, trazodone and viloxazine, and pharmaceutically acceptable salts thereof. Suitable classes of anti-anxiety agents include benzodiazepines and 5-HT_{1A} agonists or antagonists, especially 5-HT_{1A} partial agonists, and corticotropin releasing factor (CRF) antagonists. Suitable benzodiazepines include alprazolam, chlordiazepoxide, clonazepam, chlorazepate, diazepam, halazepam, lorazepam, oxazepam, and prazepam, and pharmaceutically acceptable salts thereof. Suitable 5-HT_{1A} receptor agonists or antagonists include, in particular, the 5-HT_{1A} receptor partial agonists buspirone, flesinoxan, gepirone and ipsapirone, and pharmaceutically acceptable salts thereof.

This invention also relates to a method of treating a disorder or condition selected from the group consisting of sleep disorders such as insomnia (*e.g.*, primary insomnia including psychophysiological and idiopathic insomnia, secondary insomnia including insomnia secondary to restless legs syndrome, insomnia related to peri- and/or postmenopause, Parkinson's disease or another chronic disorder, and transient insomnia), somnambulism, sleep deprivation, REM sleep disorders, sleep apnea, hypersomnia, parasomnias, sleep-wake cycle disorders, jet lag, narcolepsy, sleep disorders associated with shift work or irregular work schedules, deficient sleep quality due to a decrease in slow wave sleep caused by medications or other sources, and other sleep disorders in a mammal, in a mammal, comprising

-11-

administering to a mammal in need of such treatment a therapeutically effective amount of therapeutically effective amount of a compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof.

5 This invention also relates to a method of increasing slow wave sleep and increasing growth hormone secretion in a human subject in a mammal, comprising administering to a human subject in need of such treatment a therapeutically effective amount of therapeutically effective amount of a compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof.

10 This invention also relates to a method of treating a disorder or condition selected from the group consisting of respiratory diseases, particularly those associated with excess mucus secretion, such as chronic obstructive airways disease, bronchopneumonia, chronic bronchitis, cystic fibrosis, adult respiratory distress syndrome, and bronchospasm; cough, whooping cough, angiotensin converting enzyme (ACE) induced cough, pulmonary tuberculosis, allergies such
15 as eczema and rhinitis; contact dermatitis, atopic dermatitis, urticaria, and other eczematoid dermatitis; itching, hemodialysis associated itching; inflammatory diseases such as inflammatory bowel disease, psoriasis, osteoarthritis, cartilage damage (*e.g.*, cartilage damage resulting from physical activity or osteoarthritis), rheumatoid arthritis, psoriatic arthritis, asthma, pruritis and sunburn; and
20 hypersensitivity disorders such as poison ivy in a mammal, including a human, comprising administering to a mammal in need of such treatment a therapeutically effective amount of therapeutically effective amount of a compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof.

25 Other more specific embodiments of this invention include any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of any two or more comorbid disorders or conditions selected from those disorders and conditions referred to in any of the above methods.

30 Another more specific embodiment of this invention relates to any of the above methods for treating fibromyalgia wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of fibromyalgia and concomitant generalized anxiety disorder.

-12-

Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of major depressive disorder and concomitant irritable bowel syndrome.

5 Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of major depressive disorder and concomitant functional abdominal pain.

10 Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of major depressive disorder and concomitant neuropathic pain.

15 Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of fibromyalgia and concomitant premenstrual dysphoric disorder.

20 Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of major depressive disorder and concomitant dysthymia.

Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of major depressive disorder and concomitant fibromyalgia.

25 Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of dysthymia and concomitant fibromyalgia.

30 Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of major depressive disorder and a concomitant somatoform disorder selected from somatization disorder, conversion disorder, body dysmorphic disorder,

-13-

hypochondriasis, somatoform pain disorder, undifferentiated somatoform disorder and somatoform disorder not otherwise specified. See Diagnostic and Statistical manual of Mental Disorders, Fourth Edition (DSM-IV), American Psychiatric Association, Washington, D.C., May 1194, pp. 435-436.

5 Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of fibromyalgia and concomitant irritable bowel syndrome.

10 Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of fibromyalgia and concomitant functional abdominal pain.

15 Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of fibromyalgia and concomitant neuropathic pain.

20 Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of generalized anxiety disorder and concomitant premenstrual dysphoric disorder.

25 Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of generalized anxiety disorder and concomitant dysthymia.

 Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of generalized anxiety disorder and concomitant fibromyalgia.

30 Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of generalized anxiety disorder and a concomitant somatoform disorder

-14-

selected from somatization disorder, conversion disorder, hypochondriasis, somatoform pain disorder (or simply "pain disorder"), body dysmorphic disorder, undifferentiated somatoform disorder, and somatoform disorder not otherwise specified. See Diagnostic and Statistical manual of Mental Disorders, Fourth Edition
5 (DSM-IV), American Psychiatric Association, Washington, D.C., May 1194, pp. 435-436.

Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the
10 treatment of fibromyalgia and a concomitant somatoform disorder selected from somatization disorder, conversion disorder, hypochondriasis, somatoform pain disorder (or simply "pain disorder"), body dysmorphic disorder, undifferentiated somatoform disorder, and somatoform disorder not otherwise specified. See Diagnostic and Statistical manual of Mental Disorders, Fourth Edition (DSM-IV),
15 American Psychiatric Association, Washington, D.C., May 1194, pp. 435-436.

Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of major depressive disorder accompanied by one or more somatic
20 symptoms selected from loss of appetite, sleep disturbances (*e.g.*, insomnia, interrupted sleep, early morning awakening, tired awakening), loss of libido, restlessness, fatigue, constipation, dyspepsia, heart palpitations, aches and pains (*e.g.*, headache, neck pain, back pain, limb pain, joint pain, abdominal pain), dizziness, nausea, heartburn, nervousness, tremors, burning and tingling sensations, morning
25 stiffness, abdominal symptoms (*e.g.*, abdominal pain, abdominal distention, gurgling, diarrhea), and the symptoms associated with generalized anxiety disorder (*e.g.*, excessive anxiety and worry (apprehensive expectation), occurring more days than not for at least six months, about a number of events and activities, difficulty controlling the worry, etc.) See Diagnostic and Statistical manual of Mental
30 Disorders, Fourth Edition (DSM-IV), American Psychiatric Association, Washington, D.C., May 1194, pp. 435-436 and 445-469. This document is incorporated herein by reference in its entirety.

-15-

Another more specific embodiment of this invention relates to any of the above methods wherein the formula **1**, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of major depressive disorder accompanied by one or more somatic symptoms selected from fatigue, headache, neck pain, back pain, limb pain, joint pain, abdominal pain, abdominal distention, gurgling, diarrhea nervousness, and the symptoms associated with generalized anxiety disorder (*e.g.*, excessive anxiety and worry (apprehensive expectation), occurring more days than not for at least six months, about a number of events and activities, difficulty controlling the worry, etc. See Diagnostic and Statistical manual of Mental Disorders, Fourth Edition (DSM-IV), American Psychiatric Association, Washington, D.C., May 1194, pp. 435-436 and 445-469.

Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula **1**, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of generalized anxiety disorder accompanied by one or more somatic symptoms selected from loss of appetite, sleep disturbances (*e.g.*, insomnia, interrupted sleep, early morning awakening, tired awakening), loss of libido, restlessness, fatigue, constipation, dyspepsia, heart palpitations, aches and pains (*e.g.*, headache, neck pain, back pain, limb pain, joint pain, abdominal pain), dizziness, nausea, heartburn, nervousness, tremors, burning and tingling sensations, morning stiffness, abdominal symptoms (*e.g.*, abdominal pain, abdominal distention, gurgling, diarrhea), and the symptoms associated with major depressive disorder (*e.g.*, sadness, tearfulness, loss of interest, fearfulness, helplessness, hopelessness, fatigue, low self esteem, obsessive ruminations, suicidal thoughts, impaired memory and concentration, loss of motivation, paralysis of will, reduced appetite, increased appetite).

Another more specific embodiment of this invention relates to any of the above methods wherein the compound of formula **1**, Compound A, or a pharmaceutically acceptable salt thereof, is administered to a human for the treatment of generalized anxiety disorder accompanied by one or more somatic symptoms selected from fatigue, headache, neck pain, back pain, limb pain, joint pain, abdominal pain, abdominal distention, gurgling, diarrhea nervousness, and the symptoms associated with major depressive disorder (*e.g.*, sadness, tearfulness, loss

-16-

of interest, fearfulness, helplessness, hopelessness, low self esteem, obsessive ruminations, suicidal thoughts, fatigue, impaired memory and concentration, loss of motivation, paralysis of will, reduced appetite, increased appetite).

5 This invention also relates to a method of treating a disorder or condition selected from the group consisting of sleep disorders such as insomnia (*e.g.*, primary insomnia including psychophysiological and idiopathic insomnia, secondary insomnia including insomnia secondary to restless legs syndrome, Parkinson's disease or another chronic disorder, and transient insomnia), somnambulism, sleep deprivation, REM sleep disorders, sleep apnea, hypersomnia, parasomnias, sleep-
10 wake cycle disorders, jet lag, narcolepsy, sleep disorders associated with shift work or irregular work schedules, deficient sleep quality due to a decrease in slow wave sleep caused by medications or other sources, and other sleep disorders in a mammal, comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of the formula 1, Compound A, or
15 a pharmaceutically acceptable salt thereof.

This invention also relates to a method of increasing slow wave sleep in a human subject comprising administering to a human subject in need of such treatment a therapeutically effective amount of a compound of the formula 1 Compound A, or a pharmaceutically acceptable salt thereof.

20 This invention also relates to a method of increasing growth hormone secretion in a human subject comprising administering to a human subject in need of such treatment a therapeutically effective amount of a compound of the formula 1, compound A, or a pharmaceutically acceptable salt thereof.

25 This invention also relates to a method of treating irritable bowel syndrome in a mammal, preferably a human, comprising administering to a human subject in need of such treatment a therapeutically effective amount of a compound of the formula 1, Compound A, or a pharmaceutically acceptable salt thereof.

30 Another more specific embodiment of this invention relates to any of the above methods wherein the disorder is restless legs syndrome, and wherein the restless legs syndrome is a secondary syndrome associated with another disorder or condition included but not limited to renal insufficiency, heredity, pregnancy,

-17-

poliomyelitis, infectious diseases, avitaminosis, different types of anemia, diabetes. certain drugs (e.g. prochloroperazine, lithium, and mianserin.

Another more specific embodiment of this invention relates to any of the above methods wherein the disorder is restless legs syndrome, and wherein the restless legs syndrome is a secondary syndrome associated with certain drugs including but not limited to prochloroperazine, lithium, and mianserin.

Another more specific embodiment of this invention relates to method for treating a disorder in a mammal, including a human, comprising administering a therapeutically effective amount of a compound or a pharmaceutically acceptable salt thereof to said mammal wherein said compound is Compound A and wherein said disorder is selected from restless legs syndrome, premenstrual dysphoric disorder, and hot flashes.

Another more specific embodiment of this invention relates to method for treating a disorder in a mammal, including a human, comprising administering a therapeutically effective amount of a compound or a pharmaceutically acceptable salt thereof to said mammal wherein said compound is Compound A and wherein said disorder is selected from restless legs syndrome, premenstrual dysphoric disorder, and hot flashes.

Another more specific embodiment of this invention relates to method for treating restless legs syndrome in a mammal, including a human, comprising administering to said mammal a therapeutically effective amount of a compound or a pharmaceutically acceptable salt thereof wherein said compound is compound A.

Another more specific embodiment of this invention relates to a method for treating a disorder in a mammal, including a human, comprising administering to said mammal a therapeutically effective amount of a compound or a pharmaceutically acceptable salt thereof to a mammal wherein said compound is a compound selected from Group A compounds and wherein said disorder is selected from restless legs syndrome, premenstrual dysphoric disorder, and hot flashes.

This invention also relates to a method of treating a disorder or condition selected from the group consisting of movement disorders such as primary movement disorders, akinesias, dyskinesias (*e.g.*, familial paroxysmal dyskinesia,

-18-

tardive dyskinesia, tremor, chorea, myoclonus, tics and other dyskinesias) spasticities including muscle spasticity, periodic limb movement disorder, hypotonia with paralysis, Tourette's syndrome, Scott syndrome, palsys (*e.g.*, Bell's palsy, cerebral palsy, birth palsy, brachial palsy, wasting palsy, ischemic palsy, progressive bulbar palsy and other palsys), akinetic-rigid syndrome; extra-pyramidal movement disorders such as medication-induced movement disorders, for example, neuroleptic-induced Parkinsonism, neuroleptic malignant syndrome, neuroleptic-induced acute dystonia, neuroleptic-induced acute akathisia, neuroleptic-induced tardive dyskinesia and medication-induced postural tremour; restless legs syndrome and movement disorders associated with Parkinson's disease or Huntington's disease in a mammal, comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of formula 1, Compound A, or a pharmaceutically acceptable salt thereof.

Particular embodiments of the invention relate to any of the above methods employing a compound of formula 1 wherein the compound or salt that is employed in such method is one wherein R¹ is hydrogen, and R² is straight or branched alkyl of from 4 to 8 carbon atoms.

Other embodiments of the invention relate to any of the above methods wherein the compound or salt that is employed in the method is selected from the following compounds and their pharmaceutically acceptable salts:

3-Aminomethyl-5-methyl-nonanoic acid;
3-Aminomethyl-5-methyl-decanoic acid;
3-Aminomethyl-5-methyl-undecanoic acid;
3-Aminomethyl-5-methyl-dodecanoic acid;
3-Aminomethyl-5-methyl-tridecanoic acid;
3-Aminomethyl-5-cyclopropyl-hexanoic acid;
3-Aminomethyl-5-cyclobutyl-hexanoic acid;
3-Aminomethyl-5-trifluoromethyl-hexanoic acid;
3-Aminomethyl-5-(2-chlorophenyl)-hexanoic acid;
3-Aminomethyl-5-(3-chlorophenyl)-hexanoic acid;
3-Aminomethyl-5-(4-chlorophenyl)-hexanoic acid;
3-Aminomethyl-5-(2-methoxyphenyl)-hexanoic acid;

-19-

- 3-Aminomethyl-5-(3-methoxyphenyl)-hexanoic acid;
3-Aminomethyl-5-(4-methoxyphenyl)-hexanoic acid;
3-Aminomethyl-5-(phenylmethyl)-hexanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-octanoic acid;
5 (3S,5R)-3-Aminomethyl-5-methyl-nonanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-decanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-undecanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-dodecanoic acid;
(3S,5R)-3-Aminomethyl-5,9-dimethyl-decanoic acid;
10 (3S,5R)-3-Aminomethyl-5-methyl-heptanoic acid;
(3S,5R)-3-Aminomethyl-5,7-dimethyl-octanoic acid;
(3S,5R)-3-Aminomethyl-5,10-dimethyl-undecanoic acid;
(3S,5R)-3-Aminomethyl-6-cyclopropyl-5-methyl-hexanoic acid;
(3S,5R)-3-Aminomethyl-6-cyclobutyl-5-methyl-hexanoic acid;
15 (3S,5R)-3-Aminomethyl-6-cyclopentyl-5-methyl-hexanoic acid;
(3S,5R)-3-Aminomethyl-6-cyclohexyl-5-methyl-hexanoic acid;
(3S,5R)-3-Aminomethyl-7-cyclopropyl-5-methyl-heptanoic acid;
(3S,5R)-3-Aminomethyl-7-cyclobutyl-5-methyl-heptanoic acid;
(3S,5R)-3-Aminomethyl-7-cyclopentyl-5-methyl-heptanoic acid;
20 (3S,5R)-3-Aminomethyl-7-cyclohexyl-5-methyl-heptanoic acid;
(3S,5R)-3-Aminomethyl-8-cyclopropyl-5-methyl-octanoic acid;
(3S,5R)-3-Aminomethyl-8-cyclobutyl-5-methyl-octanoic acid;
(3S,5R)-3-Aminomethyl-8-cyclopentyl-5-methyl-octanoic acid;
(3S,5R)-3-Aminomethyl-8-cyclohexyl-5-methyl-octanoic acid;
25 (3S,5R)-3-Aminomethyl-8-fluoro-5-methyl-octanoic acid;
(3S,5R)-3-Aminomethyl-9-fluoro-5-methyl-nonanoic acid;
(3S,5S)-3-Aminomethyl-5-methoxy-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-ethoxy-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-propoxy-hexanoic acid;
30 (3S,5S)-3-Aminomethyl-5-isopropoxy-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-*tert*-butoxy-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-fluoromethoxy-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-(2-fluoro-ethoxy)-hexanoic acid;

-20-

- (3S,5S)-3-Aminomethyl-5-(3,3,3-trifluoro-propoxy)-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-phenoxy-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-(4-chloro-phenoxy)-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-(3-chloro-phenoxy)-hexanoic acid;
5 (3S,5S)-3-Aminomethyl-5-(2-chloro-phenoxy)-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-(4-fluoro-phenoxy)-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-(3-fluoro-phenoxy)-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-(2-fluoro-phenoxy)-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-(4-methoxy-phenoxy)-hexanoic acid;
10 (3S,5S)-3-Aminomethyl-5-(3-methoxy-phenoxy)-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-(2-methoxy-phenoxy)-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-(4-nitro-phenoxy)-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-(3-nitro-phenoxy)-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-(2-nitro-phenoxy)-hexanoic acid;
15 (3S,5S)-3-Aminomethyl-5-methyl-6-propoxy-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-isopropoxy-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-*tert*-butoxy-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-fluoromethoxy-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(2-fluoro-ethoxy)-5-methyl-
20 hexanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl-6-(3,3,3-trifluoro-propoxy)-
hexanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl-6-phenoxy-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(4-chloro-phenoxy)-5-methyl-
25 hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(3-chloro-phenoxy)-5-methyl-
hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(2-chloro-phenoxy)-5-methyl-
hexanoic acid;
30 (3S,5S)-3-Aminomethyl-6-(4-fluoro-phenoxy)-5-methyl-
hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(3-fluoro-phenoxy)-5-methyl-
hexanoic acid;

-21-

- (3S,5S)-3-Aminomethyl-6-(2-fluoro-phenoxy)-5-methyl-hexanoic acid;
- (3S,5S)-3-Aminomethyl-6-(4-methoxy-phenoxy)-5-methyl-hexanoic acid;
- 5 (3S,5S)-3-Aminomethyl-6-(3-methoxy-phenoxy)-5-methyl-hexanoic acid;
- (3S,5S)-3-Aminomethyl-6-(2-methoxy-phenoxy)-5-methyl-hexanoic acid;
- (3S,5S)-3-Aminomethyl-5-methyl 6-(4-trifluoromethyl-phenoxy)-hexanoic acid;
- 10 (3S,5S)-3-Aminomethyl-5-methyl 6-(3-trifluoromethyl-phenoxy)-hexanoic acid;
- (3S,5S)-3-Aminomethyl-5-methyl 6-(2-trifluoromethyl-phenoxy)-hexanoic acid;
- 15 (3S,5S)-3-Aminomethyl-5-methyl 6-(4-nitro-phenoxy)-hexanoic acid;
- (3S,5S)-3-Aminomethyl-5-methyl 6-(3-nitro-phenoxy)-hexanoic acid;
- (3S,5S)-3-Aminomethyl-5-methyl 6-(2-nitro-phenoxy)-hexanoic acid;
- 20 (3S,5S)-3-Aminomethyl-6-benzyloxy-5-methyl-hexanoic acid;
- (3S,5S)-3-Aminomethyl-7-hydroxy-5-methyl-heptanoic acid;
- (3S,5S)-3-Aminomethyl-7-methoxy-5-methyl-heptanoic acid;
- (3S,5S)-3-Aminomethyl-5-methyl-7-propoxy-heptanoic acid;
- 25 (3S,5S)-3-Aminomethyl-7-isopropoxy-5-methyl-heptanoic acid;
- (3S,5S)-3-Aminomethyl-7-*tert*-butoxy-5-methyl-heptanoic acid;
- (3S,5S)-3-Aminomethyl-7-fluoromethoxy-5-methyl-heptanoic acid;
- (3S,5S)-3-Aminomethyl-7-(2-fluoro-ethoxy)-5-methyl-heptanoic acid;
- 30 (3S,5S)-3-Aminomethyl-5-methyl-7-(3,3,3-trifluoro-propoxy)-heptanoic acid;
- (3S,5S)-3-Aminomethyl-7-benzyloxy-5-methyl-heptanoic acid;

-22-

- (3S,5S)-3-Aminomethyl-5-methyl-7-phenoxy-heptanoic acid;
(3S,5S)-3-Aminomethyl-7-(4-chloro-phenoxy)-5-methyl-heptanoic acid;
(3S,5S)-3-Aminomethyl-7-(3-chloro-phenoxy)-5-methyl-heptanoic acid;
5 (3S,5S)-3-Aminomethyl-7-(2-chloro-phenoxy)-5-methyl-heptanoic acid;
(3S,5S)-3-Aminomethyl-7-(4-fluoro-phenoxy)-5-methyl-heptanoic acid;
10 (3S,5S)-3-Aminomethyl-7-(3-fluoro-phenoxy)-5-methyl-heptanoic acid;
(3S,5S)-3-Aminomethyl-7-(2-fluoro-phenoxy)-5-methyl-heptanoic acid;
(3S,5S)-3-Aminomethyl-7-(4-methoxy-phenoxy)-5-methyl-heptanoic acid;
15 (3S,5S)-3-Aminomethyl-7-(3-methoxy-phenoxy)-5-methyl-heptanoic acid;
(3S,5S)-3-Aminomethyl-7-(2-methoxy-phenoxy)-5-methyl-heptanoic acid;
20 (3S,5S)-3-Aminomethyl-5-methyl-7-(4-trifluoromethyl-phenoxy)-heptanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl-7-(3-trifluoromethyl-phenoxy)-heptanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl-7-(2-trifluoromethyl-phenoxy)-heptanoic acid;
25 (3S,5S)-3-Aminomethyl-5-methyl-7-(4-nitro-phenoxy)-heptanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl-7-(3-nitro-phenoxy)-heptanoic acid;
30 (3S,5S)-3-Aminomethyl-5-methyl-7-(2-nitro-phenoxy)-heptanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl-6-phenyl-hexanoic acid;

-23-

- (3S,5S)-3-Aminomethyl-6-(4-chloro-phenyl)-5-methyl-hexanoic acid;
- (3S,5S)-3-Aminomethyl-6-(3-chloro-phenyl)-5-methyl-hexanoic acid;
- 5 (3S,5S)-3-Aminomethyl-6-(2-chloro-phenyl)-5-methyl-hexanoic acid;
- (3S,5S)-3-Aminomethyl-6-(4-methoxy-phenyl)-5-methyl-hexanoic acid;
- (3S,5S)-3-Aminomethyl-6-(3-methoxy-phenyl)-5-methyl-hexanoic acid;
- 10 (3S,5S)-3-Aminomethyl-6-(2-methoxy-phenyl)-5-methyl-hexanoic acid;
- (3S,5S)-3-Aminomethyl-6-(3-fluoro-phenyl)-5-methyl-hexanoic acid;
- 15 (3S,5S)-3-Aminomethyl-6-(2-fluoro-phenyl)-5-methyl-hexanoic acid;
- (3S,5R)-3-Aminomethyl-5-methyl-7-phenyl-heptanoic acid;
- (3S,5R)-3-Aminomethyl-7-(4-chloro-phenyl)-5-methyl-heptanoic acid;
- 20 (3S,5R)-3-Aminomethyl-7-(3-chloro-phenyl)-5-methyl-heptanoic acid;
- (3S,5R)-3-Aminomethyl-7-(2-chloro-phenyl)-5-methyl-heptanoic acid;
- (3S,5R)-3-Aminomethyl-7-(4-methoxy-phenyl)-5-methyl-heptanoic acid;
- 25 (3S,5R)-3-Aminomethyl-7-(3-methoxy-phenyl)-5-methyl-heptanoic acid;
- (3S,5R)-3-Aminomethyl-7-(2-methoxy-phenyl)-5-methyl-heptanoic acid;
- 30 (3S,5R)-3-Aminomethyl-7-(4-fluoro-phenyl)-5-methyl-heptanoic acid;
- (3S,5R)-3-Aminomethyl-7-(3-fluoro-phenyl)-5-methyl-heptanoic acid;

(3S,5R)-3-Aminomethyl-7-(2-fluoro-phenyl)-5-methyl-heptanoic acid;

- 5 (3S,5S)-3-Aminomethyl-5-methyl-hept-6-enoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-oct-7-enoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-non-8-enoic acid;
(E)-(3S,5S)-3-Aminomethyl-5-methyl-oct-6-enoic acid;
(Z)-(3S,5S)-3-Aminomethyl-5-methyl-oct-6-enoic acid;
(Z)-(3S,5S)-3-Aminomethyl-5-methyl-non-6-enoic acid;
(E)-(3S,5S)-3-Aminomethyl-5-methyl-non-6-enoic acid;
10 (E)-(3S,5R)-3-Aminomethyl-5-methyl-non-7-enoic acid;
(Z)-(3S,5R)-3-Aminomethyl-5-methyl-non-7-enoic acid;
(Z)-(3S,5R)-3-Aminomethyl-5-methyl-dec-7-enoic acid;
(E)-(3S,5R)-3-Aminomethyl-5-methyl-undec-7-enoic acid;
(3S,5S)-3-Aminomethyl-5,6,6-trimethyl-heptanoic acid;
15 (3S,5S)-3-Aminomethyl-5-cyclopropyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-cyclobutyl-hexanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-8-phenyl-octanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl-6-phenyl-hexanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-7-phenyl-heptanoic acid;
20 3-Aminomethyl-5-methylheptanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-heptanoic acid;
3-Aminomethyl-5-methyl-octanoic acid;
3-Aminomethyl-4,5-dimethyl-hexanoic acid;
(3S,4S)-3-Aminomethyl-4,5-dimethyl-hexanoic acid;
25 (3R,4R)-3-Aminomethyl-4,5-dimethyl-hexanoic acid;
3-Aminomethyl-4-isopropyl-hexanoic acid;
3-Aminomethyl-4-isopropyl-heptanoic acid;
(3S,5S)-3-Aminomethyl-6-fluoro-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-7-fluoro-5-methyl-heptanoic acid;
30 (3S,5S)-3-Aminomethyl-7,7,7-trifluoro-5-methyl-heptanoic acid;
(3S,5R)-3-Aminomethyl-8,8,8-trifluoro-5-methyl-octanoic acid;
(3S,5S)-3-Aminomethyl-5,6-dimethyl-heptanoic acid;
(3R,4R,5R)-3-Aminomethyl-4,5-dimethyl-heptanoic acid; and

-25-

(3R,4R,5R)-3-Aminomethyl-4,5-dimethyl-octanoic acid.

Especially useful as an agent for the treatment of fibromyalgia, fibromyalgia and one or more concomitant disorders; and/or obsessive compulsive disorder (OCD), post traumatic stress syndrome (PTSD), agoraphobia, agoraphobia without history of panic disorder, specific phobia, social phobia, restless legs syndrome, premenstrual dysphoric disorder, and/or hotflashes is (3S,5R)-3-Aminomethyl-5-methyl-octanoic acid .

The term "lower alkyl" is a straight or branched alkyl group of from 1 to 4 carbons.

The term "alkyl" is a straight or branched hydrocarbon radical of from 1 to 6 carbon atoms including but not limited to methyl, ethyl, propyl, n-propyl, isopropyl, butyl, 2-butyl, tert-butyl, pentyl, except as where otherwise stated.

The phenyl groups of compounds of the formula 1 may be unsubstituted or substituted by from 1 to 3 substituents selected from hydroxy, carboxy, carboalkoxy, halogen, CF₃, nitro, alkyl, and alkoxy. Preferred substituents are halogens.

Since amino acids are amphoteric, pharmacologically compatible salts when R is hydrogen can be salts of appropriate inorganic or organic acids, for example, hydrochloric, sulphuric, phosphoric, acetic, oxalic, lactic, citric, malic, salicylic, malonic, maleic, succinic, methanesulfonic acid, and ascorbic. Starting from corresponding hydroxides or carbonates, salts with alkali metals or alkaline earth metals, for example, sodium, potassium, magnesium, or calcium are formed. Salts with quaternary ammonium ions can also be prepared with, for example, the tetramethyl-ammonium ion. The carboxyl group of the amino acids can be esterified by known means.

Certain of the compounds employed in the methods of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms, including hydrated forms, are equivalent to unsolvated forms and are intended to be encompassed within the scope of the present invention.

Certain of the compounds employed in the methods of the present invention possess one or more chiral centers and each center may exist in the R(D)

-26-

or S(L) configuration. The present invention includes all enantiomeric and epimeric forms as well as the appropriate mixtures thereof.

One benefit of using the compounds of this invention to treat fibromyalgia is that they are not addictive. In these methods, the compounds can be combined with other agents including antidepressant and/or anti-anxiety agents.


DETAILED DESCRIPTION OF THE INVENTION


Compounds of the formula 1 can be prepared as described below and in United States Patent Applications 10/009,938, which was filed December 10, 2001, and 10/324,929, which was filed December 20, 2002. In the schemes and discussion that follow, R¹ and R² are defined as above.

The following examples are illustrative of the instant invention; they are not intended to limit the scope.

General synthetic schemes


Method 1





- 5
- a) LiAlH_4 ;
 - b) pyridinium dichormate;
 - c) triethylphosphonoacetate, NaH ;
 - d) Nitromethane DBU;
 - e) i. H_2 Pd/C; ii. HCl ; iii ion exchange chromatography.

Method 2



10

X = OEt or chiral oxazolidine auxiliary.


- a) Triethylphosphonoacetate, NaH ;

-28-

- b) i. NaOH, ii. Pivaloyl chloride, Et₃N, XH;
 c) R¹MgBr, CuBr₂ DMS;
 d) NaHMDS, BrCH₂CO₂tBu;
 e) R = tBu i. LiOH, H₂O₂; ii. BH₃, iii. TsCl, Et₃N, iv. NaN₃, DMSO;
 5 f) R = Et i. LiOH, H₂O₂; ii. BH₃, iii. PTSA, THF; iv HBr EtOH,
 v. NaN₃ DMSO;
 g) i. H₂ Pd/C; ii. HCl, iii. Ion exchange chromatography.

Specific Examples

Synthesis of Example 1: **3-Aminomethyl-5-methylheptanoic acid**



- a) PDC, CH₂Cl₂;
 b) NaH, triethylphosphonoacetate;
 15 c) DBU, CH₃NO₂;
 d) H₂, 10% Pd/C;
 e) 6N HCl, reflux, ion exchange resin (Dowex 50WX8, strongly acidic).

-29-

3-Methyl-1-pentanal 11

To a stirred suspension of pyridinium dichromate (112.17 g, 298.1 mmol) in dichloromethane 500 mL was added 3-methyl-1-pentanol 10 (15 g, 146.79 mmol). After stirring for 2.5 hours, ether 400 mL was added, and stirring was continued for another 5 minutes. The filtrate from the mixture was concentrated to a small volume and applied to a column of Florisil. The compound was eluted with petroleum ether, and further chromatographed on silica gel column using 10% ether in petroleum ether as eluent gave 11 (6.5 g, 44%).
¹H-NMR (CDCl₃) δ 9.72, (d, -CHO), 2.38 (dd, 1H, -CH₂CHO), 2.19 (dd, 1H, -CH₂CHO), 1.95 (m, 1H, C₂H₅(CH₃)CHCH₂-), 1.4-1.0 (m), 0.9-0.8 (m).

Ethyl 5-methyl-2-heptenoate 12

Sodium hydride (60% dispersion, 2.4 g, 65 mmol) was washed with hexane and suspended in dimethoxyethane 60 mL. While cooling in ice water bath triethyl phosphonoacetate was slowly added, calcd. 5 minutes. The reaction was stirred for 15 minutes at 0°C and a solution of 3-methyl-1-pentanal 11 (6.5 g, 65 mmol) in imethoxyethane 20 mL was added. After refluxing overnight, it was concentrated, water and hexane were added, the organic phase was separated, and the aqueous portion discarded. The solution was washed twice with brine and dried on magnesium sulfate. The solvent was evaporated to give 12 (6.75 g, 61%).
¹H-NMR (CDCl₃) δ 6.89 (m, 1H, -CH₂CH:CHCOOEt), 5.77 (d, 1H, -CH₂CH:CHCOOEt), 4.16 (q, 2H, -COOCH₂CH₃), 2.15 and 1.98 (1H each and a multiplet, -CH₂CH:CHCOOEt), 1.48 (m, 1H, C₂H₅(CH₃)CHCH₂-), 1.30-1.10 (m), and 0.83.

Ethyl 5-methyl-3-nitromethylheptanoate 13

Ethyl 5-methyl-2-heptanoate 12 (6.75 g, 39.70 mmol), DBU (6.0 g, 39.7 mmol), nitromethane (21.97 g, 359.9 mmol) in acetonitrile 80 mL was stirred at room temperature under nitrogen atmosphere overnight. The mixture was concentrated to an oil. A solution of the oil in ether was washed with 1N HCl, brine and dried. It was evaporated to give a light oil which was chromatographed on silica gel, eluting with 5% to 10% ether in Pet. ether to give 13 (3.6 g, 42%).

-30-

$^1\text{H-NMR}$ (CDCl_3) δ 4.49-4.39 (m), 4.12-4.07 (m), 3.61 (m), 2.36 (m), 1.36-1.18 (m), 0.86-0.79.

3-Aminomethyl-5-methylheptanoic acid (Example 1)

Ethyl 5-methyl-3-nitromethylheptanoate 13 (3.6 g) was hydrogenated in ethanol in the presence of 20% Pd/C and evaporated to give 14. Six normal hydrochloric acid 30 mL was added and refluxed overnight. The solvent was evaporated at reduced pressure, and the residue was azeotroped with toluene. Aqueous solution of the residue was applied to Dowex 50WX 8-100 ion exchange resin that had been washed to neutral pH with HPLC grade water. The column was eluted with water until eluent was neutral pH, and then with 0.5N. NH_4OH solution to give fractions containing 3-aminomethyl-5-methylheptanoic acid. The fractions were combined and further chromatographed on a C_{18} column. The compound was eluted with 40% water in methanol and crystallized from methanol-ether to give Example 1 630 mg. $^1\text{H-NMR}$ (CD_3OD) δ 2.83 (m, 1H), 2.75 (m, 1H), 2.35 (m, 1H), 2.15 (m, 1H), 1.95 (1H, bs), 1.38 (1H, m), 1.3-1.15 (m, 2H), 1.14-0.95 (m, 2H). 0.80 (m, 2 CH_3). MS found molecular ion at (M+1) 174 and other ions at 156, 139, and 102. Anal. Calcd. for $\text{C}_9\text{H}_{19}\text{NO}_2$: C, 62.39; H 11.05; N 8.08. Found C, 62.00; H, 10.83; N, 7.98.


In a similar way the following examples can be prepared.

- 20 3-Aminomethyl-5-methyl-heptanoic acid;
- 3-Aminomethyl-5-methyl-octanoic acid;
- 3-Aminomethyl-5-methyl-nonanoic acid;
- 3-Aminomethyl-5-methyl-decanoic acid;
- 3-Aminomethyl-5-methyl-undecanoic acid;
- 25 3-Aminomethyl-5-methyl-dodecanoic acid;
- 3-Aminomethyl-5-methyl-tridecanoic acid;
- 3-Aminomethyl-5-cyclopropyl-hexanoic acid;
- 3-Aminomethyl-5-cyclobutyl-hexanoic acid;
- 3-Aminomethyl-5-trifluoromethyl-hexanoic acid;
- 30 3-Aminomethyl-5-(2-chlorophenyl)-hexanoic acid;


-31-

- 3-Aminomethyl-5-(3-chlorophenyl)-hexanoic acid;
 3-Aminomethyl-5-(4-chlorophenyl)-hexanoic acid;
 3-Aminomethyl-5-(2-methoxyphenyl)-hexanoic acid;
 3-Aminomethyl-5-(3-methoxyphenyl)-hexanoic acid;
 5 3-Aminomethyl-5-(4-methoxyphenyl)-hexanoic acid; and
 3-Aminomethyl-5-(phenylmethyl)-hexanoic acid.

Synthesis of Example 2: (3R,4S)3-Aminomethyl-4,5-dimethyl-hexanoic acid



-32-



Example 2

Reagents and Conditions:

- (R) -(-)-4-phenyl-2-oxazolidinone, $(\text{CH}_3)_3\text{CCOCl}$, Et_3N , LiCl , THF, -20 to 23°C ;
- MeMgCl , CuBrSMe_2 , THF, -35°C ;
- NaHMDS , $\text{BrCH}_2\text{CO}_2\text{tBu}$, THF, -78°C to -40°C ;
- LiOH , H_2O_2 , THF, H_2O , 25°C ;
- BH_3SMe_2 , THF, 0 to 25°C ;
- pTsCl , pyridine, 25°C ;
- NaN_3 , DMSO, 60°C ;
- Raney nickel, MeOH , H_2 ; i) 3M HCl , reflux, ion exchange resin (Dowex 50WX8, strongly acidic).

[R-(E)]3-(4-Methyl-pent-2-enoyl)-4-phenyl-oxazolidin-2-one 16

Trimethylacetyl chloride (7.8 g, 0.065 mol) was added to acid **14** (6.9 g, 0.06 mol) and triethylamine (18 g, 0.187 mol) in THF (200 mL) at -20°C. After 1 hour, lithium chloride (2.35 g, 0.55 mol) and (*R*)-(-)-4-phenyl-2-oxazolidinone (8.15 g, 0.05 mol) were added and the thick suspension warmed to room temperature. After 20 hours, the suspension was filtered and the filtrate concentrated. The resultant solid was recrystallized from hexane/ethyl acetate (5:1) to give the oxazolidinone **16** as a white solid (8.83 g, 68%). ¹H NMR

-33-

(CDCl₃) δ 7.35 (m, 5H), 7.18 (dd, 1H, J = 15.4 and 1.2 Hz), 7.02 (dd, 1H, J = 15.4 and 6.8 Hz), 5.45 (dd, 1H, J = 8.8 and 3.9 Hz), 4.68 (t, 1H, J = 8.8 Hz), 4.22 (dd, 1H, J = 8.8 and 3.9 Hz), 2.50 (m, 1H), 1.04 (d, 1H, J = 1.4 Hz), 1.02 (d, 1H, J = 1.4 Hz). MS, m/z (relative intensity): 260 [M+H, 100%].

5 **(3R,3R*)3-(3,4-Dimethyl-pentanoyl)-4-phenyl-oxazolidin-2-one 17**

To copper(I) bromide-dimethyl sulphide complex in THF (45 mL) at -20°C was added methylmagnesium chloride (as a 3 M solution in THF). After 20 minutes, the oxazolidinone **16** (3.69 g, 0.014 mol) in THF (20 mL) was added dropwise over 10 minutes. After 2.5 hours, the reaction was quenched through the addition of a saturated aqueous solution of ammonium chloride. The resultant two layers were separated and the aqueous phase extracted with ether. The combined organic phases were washed with 1 M hydrochloric acid, then with 5% aqueous ammonium hydroxide. The organic phases were dried (MgSO₄) and concentrated to give the oxazolidinone **17** as a white solid (3.39 g, 88%). ¹H NMR (CDCl₃) δ 7.30 (m, 1H), 5.40 (dd, 1H, J = 8.8 and 3.7 Hz), 4.63 (t, 1H, J = 8.8 Hz), 4.21 (dd, 1H, J = 8.8 and 3.7 Hz), 2.85 (dd, 1H, J = 16.1 and 5.6 Hz), 2.8 (dd, 1H, J = 16.1 and 8.5 Hz), 1.90 (m, 1H), 1.56 (m, 2H), 0.83 (d, 3H, J = 6.8 Hz), 0.78 (d, 3H, J = 6.8 Hz), 0.75 (d, 3H, J = 6.8 Hz). MS, m/z (relative intensity): 276 [M+H, 100%].

20 **[3R-(3R*(R*),4S*)]-4,5-Dimethyl-3-(2-oxo-4-phenyl-oxazolidine-3-carbonyl)-hexanoic acid tert-butyl ester 18**

Sodium bis(trimethylsilyl)amide (14.4 mL, 0.014 mol of a 1 M solution in THF) was added to a solution of the oxazolidinone **17** (3.37 g, 0.012 mol) in THF (35 mL) at -78°C. After 35 minutes, *tert*-butyl bromoacetate (3.5 g, 0.018 mol) was added and the solution immediately warmed to -40°C. After 3 hours, the reaction was quenched through the addition of a saturated aqueous solution of ammonium chloride. The resultant two layers were separated and the aqueous phase extracted with ether. The combined organic phases were dried (MgSO₄) and concentrated. Flash chromatography (9:1 to 5:1 hexane/ethyl acetate gradient) gave the ester **18** (3.81 g, 82%) as a white solid. ¹H NMR (CDCl₃) δ 7.35 (m,

-34-

5H), 5.37 (dd, 1H, $J = 8.4$ and 3.1 Hz), 4.67 (t, 1H, $J = 8.7$ Hz), 4.41 (dt, 1H, $J = 12.0$ and 3.5 Hz), 4.25 (dd, 1H, $J = 8.68$ and 3.1 Hz), 2.65 (dd, 1H, $J = 16.9$ and 12.0 Hz), 2.25 (dd, 1H, $J = 16.9$ and 3.5 Hz), 1.6 (m, 1H), 1.45 (m, 1H), 1.23 (s, 9H), 1.02 (d, 1H, $J = 6.5$ Hz), 0.93 (d, 1H, $J = 6.7$ Hz), 0.80 (d, 1H, $J = 7.0$ Hz). MS, m/z (relative intensity): 429 [M-H+CH₃CN, 100%], 388 [M-H, 20%].

(3R,4S)-2-(1,2-Dimethyl-propyl)-succinic acid 4-tert-butyl ester 19

To the oxazolidinone **18** (3.62 g, 9.3 mmol) in THF (54 mL)/water (15 mL) was added a premixed solution of lithium hydroxide (20 mL of a 0.8 M aqueous solution, 0.016 mol)/H₂O₂ (5.76 mL of a 30% aqueous solution). After 10 7 hours, the solution was diluted with water and sodium bisulfite added (~10 g). After stirring for a further 0.5 hours, the two layers were separated and the aqueous phase extracted with ether. The aqueous phase was then rendered acidic (pH 2) with 1 M hydrochloric acid and extracted with ether. The combined 15 organic phases were dried (MgSO₄) and concentrated. Flash chromatography (5:1 hexane/ethyl acetate) gave the acid **19** (2.1 g, 95%) as a colorless oil. ¹H NMR (CDCl₃) δ 3.0 (m, 1H), 2.55 (dd, 1H, $J = 16.6$ and 11.2 Hz), 2.27 (dd, 1H, $J = 16.6$ and 3.4 Hz), 1.70 (m, 1H), 1.53 (m, 1H), 1.45 (m, 1H), 1.43 (s, 9H), 0.95 (d, 1H, $J = 6.8$ Hz), 0.90 (d, 1H, $J = 6.6$ Hz), 0.83 (d, 1H, $J = 6.8$ Hz). MS, 20 m/z (relative intensity): 243 [M-H, 100%].

(3R,4S)-3-Hydroxymethyl-4,5-dimethyl-hexanoic acid tert-butyl ester 20

Borane-methyl sulfide complex (16 mL, 0.032 mol of a 2 M solution in THF) was added to a stirred solution of the acid **19** (1.96 g, 8 mmol) in THF (20 mL) at 0°C. After 20 hours, methanol was added until effervescence ceased 25 and the solution concentrated. Flash chromatography (5:1 hexane/ethyl acetate gradient) gave the alcohol **20** (1.29 g, 70%) as a colorless oil. ¹H NMR (CDCl₃) δ 3.62 (m, 1H), 2.32 (m, 1H), 2.14 (m, 1H), 1.6 (m, 1H), 1.45 (s, 9H), 1.35 (m, 1H), 0.93 (d, 1H, $J = 6.8$ Hz), 0.86 (d, 1H, $J = 6.8$ Hz), 0.77 (d, 1H, $J = 6.9$ Hz). MS, m/z (relative intensity): 175 [M-tBu, 100%].

-35-

(3R,4S)-4,5-Dimethyl-3-(toluene-4-sulfonyloxymethyl)-hexanoic acid tert-butyl ester 21

p-Toluenesulfonyl chloride (847 mg, 4.4 mmol) was added to a stirred solution of the alcohol **6** (850 mg, 3.7 mmol), DMAP (10 mg, 0.08 mmol) and triethylamine (1.23 mL, 8.88 mmol) in CH₂Cl₂ (20 mL) at 0°C and the solution warmed to room temperature. After 15 hours, the solution was washed with 1N hydrochloric acid then with brine. The combined organic phases were dried (MgSO₄) and concentrated. Flash chromatography (100 to 92% hexane/ethyl acetate gradient) gave the tosylate **7** (1.22 g, 86%) as a thick gum. ¹H NMR (CDCl₃) δ 7.80 (d, 2H, *J* = 8.2 Hz), 7.25 (d, 2H, *J* = 8.2 Hz), 3.92 (m, 1H), 2.38 (s, 3H), 2.20 (m, 2H), 1.95 (m, 1H), 1.40 (m, 1H), 1.32 (s, 9H), 1.27 (m, 1H), 0.78 (d, 1H, *J* = 6.6 Hz), 0.73 (d, 1H, *J* = 6.6 Hz), 0.63 (d, 1H, *J* = 7.1 Hz). MS, *m/z* (relative intensity): 311 [85%], 198 [100%], 157 [95%].

(3R,4S)-3-Azidomethyl-4,5-dimethyl-hexanoic acid tert-butyl ester 22

A solution of the tosylate **21** (1.19 g, 3.1 mmol) and sodium azide (402 mg, 6.2 mmol) in DMSO (15 mL) was warmed to 60°C for 2.5 hours. Water (100 mL) was added and the solution extracted with ether. The combined organic phases were dried (MgSO₄) and concentrated. Flash chromatography (9:1 hexane/ethyl acetate) gave the azide **22** (628 mg, 80%) as a colorless oil. ¹H NMR (CDCl₃) δ 3.4 (dd, 1H, *J* = 12.21 and 6.11 Hz), 3.3 (dd, 1H, *J* = 21.11 and 6.59 Hz), 2.30 (dd, 1H, *J* = 15.14 and 3.66 Hz), 2.25 (m, 1H), 2.05 (dd, 1H, *J* = 15.14 and 9.04 Hz), 1.55 (m, 1H), 1.45 (s, 9H), 1.35 (m, 1H), 0.95 (d, 1H, *J* = 6.59 Hz), 0.90 (d, 1H, *J* = 6.83 Hz), 0.80 (d, 1H, *J* = 7.08 Hz). MS (*m/z*): (relative intensity): 228 [M-N₂, 35%], 172 [M-N₂-tBu, 100%].

(3R,4S)-3-Aminomethyl-4,5-dimethyl-hexanoic acid tert-butyl ester 23 and [4R-[4R*(S*)]]-4-(1,2-Dimethyl-propyl)-pyrrolidin-2-one 24

The azide **8** (640 mg, 2.5 mmol) and Raney nickel (1 g) in methanol (50 mL) were shaken under an atmosphere of hydrogen for 4 hours. The solution was filtered and the filtrate concentrated to give a mixture of the amine **23** and lactam **24** which was used without further purification in the next step.

-36-

(3R,4S)-3-Aminomethyl-4,5-dimethyl-hexanoic acid (Example 2)

A solution of the amine **23** and lactam **24** (500 mg) in 3 M hydrochloric acid were heated to reflux for 9 hours, then stirred at room temperature for 15 hours. The solution was concentrated and the resultant solid subjected to a sequential purification which involved ion exchange chromatography (Dowex 50WX8, strongly acidic), oxalate salt formation then further purification by ion exchange chromatography (Dowex 50WX8, strongly acidic) to give the Example 2 (343 mg) as a white solid. ¹H NMR (D₂O) δ 2.87 (m, 2H), 2.22 (dd, 1H, *J* = 15.4 and 3.4 Hz), 2.12 (m, 1H), 1.93 (dd, 1H, *J* = 15.4 and 9.5 Hz), 1.38 (m, 1H), 1.12 (m, 1H), 0.77 (d, 1H, *J* = 6.6 Hz), 0.74 (d, 1H, *J* = 6.6 Hz), 0.70 (d, 1H, *J* = 6.8 Hz). MS, *m/z* (relative intensity): 174 [M+H, 100%].

In a similar way, the following examples can be prepared:

3-Aminomethyl-4,5-dimethyl-hexanoic acid;

(3R,4S)-3-Aminomethyl-4,5-dimethyl-hexanoic acid MP;

(3S,4S)-3-Aminomethyl-4,5-dimethyl-hexanoic acid;

(3R,4R)-3-Aminomethyl-4,5-dimethyl-hexanoic acid MP;

3-Aminomethyl-4-isopropyl-hexanoic acid;

3-Aminomethyl-4-isopropyl-heptanoic acid;


3-Aminomethyl-4-isopropyl-octanoic acid;

3-Aminomethyl-4-isopropyl-nonanoic acid;

3-Aminomethyl-4-isopropyl-decanoic acid; and

3-Aminomethyl-4-phenyl-5-methyl-hexanoic acid.

-37-

Method 3

where
 R₃ = OMe or H
 R₄ = Me, Et
 n = 0 to 2

5

A compound of structure 30 could be prepared from a compound of structure 29 by treatment with an aqueous acid such as hydrochloric acid and alike at a temperature between room temperature and reflux. As an alternative, a compound of structure 30 can be prepared from a compound of structure 32 by treatment with trifluoroacetic acid in a solvent such as CH₂Cl₂ or EtOAc and alike. Compound 32 could be prepared by base mediate hydrolysis of a Boc

-38-

protected lactam such as compound 31 which itself could be prepared from a compound of structure 29 by treatment with di-tert-butyl dicarbonate in a solvent such as THF and alike. The treatment of the Boc-lactam 31 with aqueous sodium hydroxide for example would give rise to the acid 32.

5 A compound of structure 29 could be prepared from compound of structure 28 ($n = 0$) by treatment with sodium or lithium metal in ammonia. Preferably, the reaction is carried out with sodium metal in ammonia. Alternatively, a compound of structure 29 could be prepared from compound of structure 28 ($n = 1$ or 2) by treatment with ceric ammonium nitrate in a mixture of
10 acetonitrile and water. Other methods known in the literature for the removal of substituted alkoxy benzyl groups from nitrogen are described in Green, *Protective Groups in Organic Synthesis*, Wiley, 2 ed, 1991 and could be utilized.

 A compound of structure 28 could be prepared from a compound of structure 27 (where LG is a suitable leaving group such as a halide or an alkyl
15 sulphonate, preferably an iodide would be used) by carbon-carbon bond forming reactions known in the art. Several methods exist in the literature for the coupling of organohalides or organoalkyl sulphonates with organometallic reagents in the presence of various metal salts as summarized in *Comprehensive Organic Synthesis*, volume 3:413 which could be utilized. For example, a compound of
20 structure 28 could be prepared from a compound of structure 27 (where LG is iodide) by treatment with a suitable secondary halide (chloride or iodide) in the presence of magnesium metal, iodine and copper bromide dimethylsulphide in a solvent such as tetrahydrofuran and alike. Alternatively the method according to El Marini, *Synthesis*, 1992:1104 could be used. Hence, a compound of
25 structure 28 could be prepared from a compound of structure 27 (where LG is iodide) by treatment with suitable methyl-substituted secondary halide such as an iodide in the presence of magnesium, iodine and lithium tetrachlorocuprate in a solvent such as tetrahydrofuran and alike.

 A compound of structure 27 incorporates a suitable leaving group, which
30 would undergo nucleophilic substitution with suitable nucleophile. Examples of such leaving groups include halides such as chloride, bromide, or iodide, and sulphonic esters such as mesylate, tosylate, triflate, nosylate, and alike.


-39-

A compound of structure 27 (where LG = iodide) could be prepared from a compound of structure 26 through treatment with iodine, triphenylphosphine, and imidazole in a solvent such as toluene and alike.


A compound of structure 26 could be prepared from compound of structure 25 by treatment with a metal borohydride, such as sodium borohydride in a solvent such as tetrahydrofuran or DME and alike.

Compound 25 could be prepared in a similar fashion to the procedures of Zoretic et al, *J. Org. Chem.*, 1980;45:810-814 or Nielsen et al *J. Med. Chem.*, 1990;33:71-77 using an appropriate benzylamine, such as but not limited to benzylamine, 4-methoxybenzylamine or 2,4-dimethoxybenzylamine.

As an alternative approach, a compound of structure 26 could be treated with sodium metal and ammonia to give 4-hydroxymethyl-pyrrolidinone which could be iodinated affording 4-iodomethyl-pyrrolidinone. 4-iodomethyl-pyrrolidinone could then be coupled with organometallic reagents according to the above procedures avoiding protection of the lactam nitrogen as below.



Analogous to the above methods a lactam of structure 33 (see Nielsen et. al., *J. Med. Chem.*, 1990;33:71-77 for general method of preparation) could be employed thus establishing fixed stereochemistry at C3 of the final amino acids.



Compounds which could be prepared in this manner include:

-40-

- 3-Aminomethyl-5-methyl-6-phenyl-hexanoic acid;
3-Aminomethyl-6-(4-chloro-phenyl)-5-methyl-hexanoic acid;
3-Aminomethyl-6-(3-chloro-phenyl)-5-methyl-hexanoic acid;
3-Aminomethyl-6-(2-chloro-phenyl)-5-methyl-hexanoic acid;
5 3-Aminomethyl-6-(4-fluoro-phenyl)-5-methyl-hexanoic acid;
3-Aminomethyl-6-(3-fluoro-phenyl)-5-methyl-hexanoic acid;
3-Aminomethyl-6-(2-fluoro-phenyl)-5-methyl-hexanoic acid;
3-Aminomethyl-5-methyl-7-phenyl-heptanoic acid;
3-Aminomethyl-7-(4-chloro-phenyl)-5-methyl-heptanoic acid;
10 3-Aminomethyl-7-(3-chloro-phenyl)-5-methyl-heptanoic acid;
3-Aminomethyl-7-(2-chloro-phenyl)-5-methyl-heptanoic acid;
3-Aminomethyl-7-(4-fluoro-phenyl)-5-methyl-heptanoic acid;
3-Aminomethyl-7-(3-fluoro-phenyl)-5-methyl-heptanoic acid;
3-Aminomethyl-7-(2-fluoro-phenyl)-5-methyl-heptanoic acid;
15 (3S)-3-Aminomethyl-6-cyclopropyl-5-methyl-hexanoic acid;
(3S)-3-Aminomethyl-6-cyclobutyl-5-methyl-hexanoic acid;
(3S)-3-Aminomethyl-6-cyclopentyl-5-methyl-hexanoic acid;
(3S)-3-Aminomethyl-6-cyclohexyl-5-methyl-hexanoic acid;
(3S)-3-Aminomethyl-7-cyclopropyl-5-methyl-heptanoic acid;
20 (3S)-3-Aminomethyl-7-cyclobutyl-5-methyl-heptanoic acid;
(3S)-3-Aminomethyl-7-cyclopentyl-5-methyl-heptanoic acid;
(3S)-3-Aminomethyl-7-cyclohexyl-5-methyl-heptanoic acid;
(3S)-3-Aminomethyl-8-cyclopropyl-5-methyl-octanoic acid;
(3S)-3-Aminomethyl-8-cyclobutyl-5-methyl-octanoic acid;
25 (3S)-3-Aminomethyl-8-cyclopentyl-5-methyl-octanoic acid;
(3S)-3-Aminomethyl-8-cyclohexyl-5-methyl-octanoic acid;
(3S)-3-Aminomethyl-5-methyl-heptanoic acid;
(3S)-3-Aminomethyl-5-methyl-octanoic acid;
(3S)-3-Aminomethyl-5-methyl-nonanoic acid;
30 (3S)-3-Aminomethyl-5-methyl-decanoic acid;
(3S)-3-Aminomethyl-5-methyl-undecanoic acid;
(3S)-3-Aminomethyl-5,7-dimethyl-octanoic acid;
(3S)-3-Aminomethyl-5,8-dimethyl-nonanoic acid;


-41-

- (3S)-3-Aminomethyl-5,9-dimethyl-decanoic acid;
(3S)-3-Aminomethyl-5,6-dimethyl-heptanoic acid;
(3S)-3-Aminomethyl-5,6,6-trimethyl-heptanoic acid;
(3S)-3-Aminomethyl-5-cyclopropyl-hexanoic acid;
5 (3S)-3-Aminomethyl-6-fluoro-5-methyl-hexanoic acid;
(3S)-3-Aminomethyl-7-fluoro-5-methyl-heptanoic acid;
(3S)-3-Aminomethyl-8-fluoro-5-methyl-octanoic acid;
(3S)-3-Aminomethyl-7,7,7-trifluoro-5-methyl-heptanoic acid;
(3S)-3-Aminomethyl-8,8,8-trifluoro-5-methyl-octanoic acid;
10 (3S)-3-Aminomethyl-5-methyl-hept-6-enoic acid;
(3S)-3-Aminomethyl-5-methyl-oct-7-enoic acid;
(3S)-3-Aminomethyl-5-methyl-non-8-enoic acid;
(E)-(3S)-3-Aminomethyl-5-methyl-oct-6-enoic acid;
(Z)-(3S)-3-Aminomethyl-5-methyl-oct-6-enoic acid;
15 (E)-(3S)-3-Aminomethyl-5-methyl-non-6-enoic acid;
(Z)-(3S)-3-Aminomethyl-5-methyl-non-6-enoic acid;
(E)-(3S)-3-Aminomethyl-5-methyl-non-7-enoic acid;
(Z)-(3S)-3-Aminomethyl-5-methyl-non-7-enoic acid;
(E)-(3S)-3-Aminomethyl-5-methyl-dec-7-enoic acid;
20 (Z)-(3S)-3-Aminomethyl-5-methyl-dec-7-enoic acid;
3-Aminomethyl-6-cyclopropyl-5-methyl-hexanoic acid;
3-Aminomethyl-6-cyclobutyl-5-methyl-hexanoic acid;
3-Aminomethyl-6-cyclopentyl-5-methyl-hexanoic acid;
3-Aminomethyl-6-cyclohexyl-5-methyl-hexanoic acid;
25 3-Aminomethyl-7-cyclopropyl-5-methyl-heptanoic acid;
3-Aminomethyl-7-cyclobutyl-5-methyl-heptanoic acid;
3-Aminomethyl-7-cyclopentyl-5-methyl-heptanoic acid;
3-Aminomethyl-7-cyclohexyl-5-methyl-heptanoic acid;
3-Aminomethyl-8-cyclopropyl-5-methyl-octanoic acid;
30 3-Aminomethyl-8-cyclobutyl-5-methyl-octanoic acid;
3-Aminomethyl-8-cyclopentyl-5-methyl-octanoic acid;
3-Aminomethyl-8-cyclohexyl-5-methyl-octanoic acid;
3-Aminomethyl-5-methyl-heptanoic acid;

-42-

- 3-Aminomethyl-5-methyl-octanoic acid;
3-Aminomethyl-5-methyl-nonanoic acid;
3-Aminomethyl-5-methyl-decanoic acid;
3-Aminomethyl-5-methyl-undecanoic acid;
5 3-Aminomethyl-5,7-dimethyl-octanoic acid;
3-Aminomethyl-5,8-dimethyl-nonanoic acid;
3-Aminomethyl-5,9-dimethyl-decanoic acid;
3-Aminomethyl-5,6-dimethyl-heptanoic acid;
3-Aminomethyl-5,6,6-trimethyl-heptanoic acid;
10 3-Aminomethyl-5-cyclopropyl-hexanoic acid;
3-Aminomethyl-6-fluoro-5-methyl-hexanoic acid;
3-Aminomethyl-7-fluoro-5-methyl-heptanoic acid;
3-Aminomethyl-8-fluoro-5-methyl-octanoic acid;
3-Aminomethyl-7,7,7-trifluoro-5-methyl-heptanoic acid;
15 3-Aminomethyl-8,8,8-trifluoro-5-methyl-octanoic acid;
3-Aminomethyl-5-methyl-hept-6-enoic acid;
3-Aminomethyl-5-methyl-oct-7-enoic acid;
3-Aminomethyl-5-methyl-non-8-enoic acid;
(E)-3-Aminomethyl-5-methyl-oct-6-enoic acid;
20 (Z)-3-Aminomethyl-5-methyl-oct-6-enoic acid;
(E)-3-Aminomethyl-5-methyl-non-6-enoic acid;
(Z)-3-Aminomethyl-5-methyl-non-6-enoic acid;
(E)-3-Aminomethyl-5-methyl-non-7-enoic acid;
(Z)-3-Aminomethyl-5-methyl-non-7-enoic acid;
25 (E)-3-Aminomethyl-5-methyl-dec-7-enoic acid; and
(Z)-3-Aminomethyl-5-methyl-dec-7-enoic acid.

-43-

Method 4

5 A compound of structure 40 could be prepared from compound of structure 39 through treatment with diethylaminosulphur trifluoride in a solvent such as methylene chloride at a temperature between -78°C and room temperature. Other methods for the fluorination of alcohols are known and could be utilized as exemplified in Wilkinson, *Chem. Rev.* 1992;92:505-519. Compounds of structure 40 can be converted to the requisite γ -amino acid as described in method 3 above.

10 A compound of structure 39 could be prepared from compound of structure 38 through treatment with osmium tetroxide and sodium periodate in a

-44-

solvent such as THF and water and reduction of the resultant intermediate with sodium borohydride in a solvent such as ethanol.

Compounds of structures **38** and **34** could be prepared from compound of structure **33** according to the principles described in method 3.


5 An alternative procedure for the synthesis of alcohol **39** (n = 0) involves the treatment of a compound of structure **36** with a metal borohydride, such as sodium borohydride in a solvent such as tetrahydrofuran or DME and alike to give a compound of structure **37**, the fluorination of which could be achieved in a similar manner to the preparation of a compound of structure **40**. A compound of structure **36** could be prepared from compound of structure **35** through treatment with sodium or lithium chloride in aqueous DMSO at a temperature between room temperature and reflux. Preferably the reaction is carried out using sodium chloride in aqueous DMSO at reflux. A compound of structure **35** could be prepared from compound of structure **34** through treatment with a suitable methyl malonic acid diester, such as dimethyl methylmalonate and alike with sodium hydride in a solvent such as DMSO or THF and alike. Preferably the reaction is carried out by adding NaH to a solution of dimethyl methylmalonate in DMSO followed by the addition of the lactam **34** (where LG is preferably iodide or as defined in method 3) pre-dissolved in DMSO.

20 Compounds **39** and **37** can be converted to the free amino acids bearing a hydroxyl group by the methods described above.

The following compounds could be prepared in this manner:

(3S)-3-Aminomethyl-6-fluoro-5-methyl-hexanoic acid;
(3S)-3-Aminomethyl-6-fluoro-5-methyl-hexanoic acid;
25 (3S)-3-Aminomethyl-7-fluoro-5-methyl-heptanoic acid;
(3S)-3-Aminomethyl-8-fluoro-5-methyl-octanoic acid;
(3S)-3-Aminomethyl-9-fluoro-5-methyl-nonanoic acid;
(3S)-3-Aminomethyl-7-hydroxy-5-methyl-heptanoic acid; and
(3S)-3-Aminomethyl-6-hydroxy-5-methyl-hexanoic acid.

-45-

Method 5

A compound of structure **41** could be prepared from compound of
 structure **39** through treatment with a suitable alkyl iodide (or alkyl sulphonate),
 such as methyl iodide and alike, and a base such as n-butyl lithium or sodium
 5 hydride and alike, in a solvent such as DMSO or THF and alike. Preferably the
 reaction is carried out by adding NaH to a solution of the alcohol in DMSO
 followed by the addition of the alkyl iodide and heating of the reaction mixture at
 a temperature between room temperature and reflux. The conversion of
 10 compounds of structure **41** to the γ -amino acids has been described above.


Alternatively, compounds of structure **41** could be derived from
 compounds of structure **42** (where LG = iodide, bromide or an sulphonic acid
 ester, as exemplified in method 3) by treatment of an appropriate alkoxy anion in a
 solvent such as DMSO or THF and alike. A compound of structure **42** would also
 15 serve as a substrate for carbon-carbon bond forming procedures as outlined in
 method 3.

Compounds which could be prepared in this manner include:

-46-

- (3S)-3-Aminomethyl-7-hydroxy-5-methyl-heptanoic acid;
(3S)-3-Aminomethyl-7-methoxy-5-methyl-heptanoic acid;
(3S)-3-Aminomethyl-7-ethoxy-5-methyl-heptanoic acid;
(3S)-3-Aminomethyl-5-methyl-7-propoxy-heptanoic acid;
5 (3S)-3-Aminomethyl-7-fluoromethoxy-5-methyl-heptanoic acid;
(3S)-3-Aminomethyl-7-(2-fluoro-ethoxy)-5-methyl-heptanoic acid;
(3S)-3-Aminomethyl-5-methyl-7-(3,3,3-trifluoro-propoxy)-heptanoic acid;
(3S)-3-Aminomethyl-6-hydroxy-5-methyl-hexanoic acid;
(3S)-3-Aminomethyl-6-methoxy-5-methyl-hexanoic acid;
10 (3S)-3-Aminomethyl-6-ethoxy-5-methyl-hexanoic acid;
(3S)-3-Aminomethyl-5-methyl-6-propoxy-hexanoic acid;
(3S)-3-Aminomethyl-6-fluoromethoxy-5-methyl-hexanoic acid;
(3S)-3-Aminomethyl-6-(2-fluoro-ethoxy)-5-methyl-hexanoic acid; and
(3S)-3-Aminomethyl-5-methyl-6-(3,3,3-trifluoro-propoxy)-hexanoic acid.

-47-

Method 6

Compounds of structure **53** could be prepared from a compound of structure **45** as shown above and by the general procedures described in Hoekstra et. al., *Organic Process Research and Development*, 1997;1:26-38.

Compounds of structure **45** can be prepared from compounds of structure **44** by treatment with a solution of chromium trioxide in water/sulfuric acid. Alternative methods of cleaving the olefin in **44** could be utilized as detailed in Hudlicky, *Oxidations in Organic Chemistry*, ACS Monograph 186, ACS 1990:77.

-48-

Compounds of structure **44** (where R₂ = alkyl, branched alkyl, cycloalkyl, alkyl-cycloalkyl) could be prepared from (S)-citronellyl bromide by carbon-carbon bond forming reactions known in the art and as described in method 3. The substitution of the halide in (S)-citronellyl bromide with alkoxy anions could also be used to provide compounds of structure **44** where R = alkoxy or phenoxy ethers (and appropriate substitutions thereof as according to Formula 1). Alternatively (S)-citronellol could be utilized to afford compounds of structure **44** by treatment of (S)-citronellol with a base such as sodium hydride, and treatment of the resultant alkoxide with an appropriate alkyl halide to afford ethers. In another method (S)-citronellyl bromide (or an appropriate sulphonic ester such as, but not limited to, methanesulfonic acid (S)-3,7-dimethyl-oct-6-enyl ester) could be reduced with an appropriate metal borohydride or with an aluminum hydride species, such as LAH, to provide (R)-2,6-dimethyl-oct-2-ene.

To one skilled in the art it will be appreciated that rational choice of either *R*- or *S*-citronellol or *R*- or *S*-citronellyl bromide would give rise to the requisite isomer at C5 of the final amino acid.

Compounds which could be prepared in this manner include:

(3S,5S)-3-Aminomethyl-7-methoxy-5-methyl-heptanoic acid;

(3S,5S)-3-Aminomethyl-5-methyl-7-propoxy-heptanoic acid;

(3S,5S)-3-Aminomethyl-7-isopropoxy-5-methyl-heptanoic acid;

(3S,5S)-3-Aminomethyl-7-*tert*-butoxy-5-methyl-heptanoic acid;

(3S,5S)-3-Aminomethyl-7-fluoromethoxy-5-methyl-heptanoic acid;

(3S,5S)-3-Aminomethyl-7-(2-fluoro-ethoxy)-5-methyl-heptanoic acid;

(3S,5S)-3-Aminomethyl-5-methyl-7-(3,3,3-trifluoro-propoxy)-heptanoic

acid;

(3S,5S)-3-Aminomethyl-7-benzyloxy-5-methyl-heptanoic acid;

(3S,5S)-3-Aminomethyl-5-methyl-7-phenoxy-heptanoic acid;

(3S,5S)-3-Aminomethyl-7-(4-chloro-phenoxy)-5-methyl-heptanoic acid;

(3S,5S)-3-Aminomethyl-7-(3-chloro-phenoxy)-5-methyl-heptanoic acid;

(3S,5S)-3-Aminomethyl-7-(2-chloro-phenoxy)-5-methyl-heptanoic acid;

(3S,5S)-3-Aminomethyl-7-(4-fluoro-phenoxy)-5-methyl-heptanoic acid;


(3S,5S)-3-Aminomethyl-7-(3-fluoro-phenoxy)-5-methyl-heptanoic acid;

-49-

- (3S,5S)-3-Aminomethyl-7-(2-fluoro-phenoxy)-5-methyl-heptanoic acid;
(3S,5S)-3-Aminomethyl-7-(4-methoxy-phenoxy)-5-methyl-heptanoic acid;
(3S,5S)-3-Aminomethyl-7-(3-methoxy-phenoxy)-5-methyl-heptanoic acid;
5 (3S,5S)-3-Aminomethyl-7-(2-methoxy-phenoxy)-5-methyl-heptanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl-7-(4-trifluoromethyl-phenoxy)-heptanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl-7-(3-trifluoromethyl-phenoxy)-
10 heptanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl-7-(2-trifluoromethyl-phenoxy)-heptanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl-7-(4-nitro-phenoxy)-heptanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl-7-(3-nitro-phenoxy)-heptanoic acid;
15 (3S,5S)-3-Aminomethyl-5-methyl-7-(2-nitro-phenoxy)-heptanoic acid;
(3S,5R)-3-Aminomethyl-7-cyclopropyl-5-methyl-heptanoic acid;
(3S,5R)-3-Aminomethyl-7-cyclobutyl-5-methyl-heptanoic acid;
(3S,5R)-3-Aminomethyl-7-cyclopentyl-5-methyl-heptanoic acid;
(3S,5R)-3-Aminomethyl-7-cyclohexyl-5-methyl-heptanoic acid;
20 (3S,5R)-3-Aminomethyl-8-cyclopropyl-5-methyl-octanoic acid;
(3S,5R)-3-Aminomethyl-8-cyclobutyl-5-methyl-octanoic acid;
(3S,5R)-3-Aminomethyl-8-cyclopentyl-5-methyl-octanoic acid;
(3S,5R)-3-Aminomethyl-8-cyclohexyl-5-methyl-octanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-heptanoic acid;
25 (3S,5R)-3-Aminomethyl-5-methyl-octanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-nonanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-decanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-undecanoic acid;
(3S,5R)-3-Aminomethyl-5,9-dimethyl-decanoic acid;
30 (3S,5S)-3-Aminomethyl-7-fluoro-5-methyl-heptanoic acid;
(3S,5R)-3-Aminomethyl-8-fluoro-5-methyl-octanoic acid;
(3S,5R)-3-Aminomethyl-8,8,8-trifluoro-5-methyl-octanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-7-phenyl-heptanoic acid;

-50-

- (3S,5R)-3-Aminomethyl-7-(4-chloro-phenyl)-5-methyl-heptanoic acid;
 (3S,5R)-3-Aminomethyl-7-(3-chloro-phenyl)-5-methyl-heptanoic acid;
 (3S,5R)-3-Aminomethyl-7-(2-chloro-phenyl)-5-methyl-heptanoic acid;
 (3S,5R)-3-Aminomethyl-7-(4-methoxy-phenyl)-5-methyl-heptanoic acid;
 5 (3S,5R)-3-Aminomethyl-7-(3-methoxy-phenyl)-5-methyl-heptanoic acid;
 (3S,5R)-3-Aminomethyl-7-(2-methoxy-phenyl)-5-methyl-heptanoic acid;
 (3S,5R)-3-Aminomethyl-7-(4-fluoro-phenyl)-5-methyl-heptanoic acid;
 (3S,5R)-3-Aminomethyl-7-(3-fluoro-phenyl)-5-methyl-heptanoic acid;
 (3S,5R)-3-Aminomethyl-7-(2-fluoro-phenyl)-5-methyl-heptanoic acid; and
 10 (3S,5R)-3-Aminomethyl-5,10-dimethyl-undecanoic acid.

Method 7

A compound of structure 58 can be prepared from a compound of structure 57 by treatment with borontrifluoride diethyletherate and triethylsilane in a solvent such as CH₂Cl₂. Alternatively the method described in Meyers, *J. Org. Chem.*, 1993;58:36-42, could be utilized thus treating a compound of structure 57 with sodium cyanoborohydride in a solvent such as THF/methanol with 3% HCl in methanol.

-51-

A compound of structure **57** can be prepared from a compound of structure **56** by treatment with dimethylamine in a solvent such as DMF and alike according to the procedure of Koot, *Tetrahedron Lett.*, 1992;33:7969-7972.

5 A compound of structure **56** can be prepared from a compound of structure **54** by treatment of a suitable primary halide **55** (iodide, bromide, or chloride) under standard transmetallation conditions with tBuLi and treatment of the resultant organometallic reagent with suitable copper salt, such as but not limited to, copper bromide or copper iodide. The resultant organo-cuprate is added to lactam (see Koot et al, *J. Org. Chem.*, 1992;57:1059-1061 for the preparation of
10 the chiral lactam **54**) in a solvent such as THF and alike. The procedure of Koot, *Tetrahedron Lett.*, 1992;33:7969-7972 exemplifies this method.

To one skilled in the art it will be appreciated that rational choice of either *R*- or *S*-primary halides **55** would give rise to the requisite isomer at C5 of the final amino acid.

15 Compounds which could be prepared in this manner include:
(3*S*,5*S*)-3-Aminomethyl-5-methoxy-hexanoic acid;
(3*S*,5*S*)-3-Aminomethyl-5-ethoxy-hexanoic acid;
(3*S*,5*S*)-3-Aminomethyl-5-propoxy-hexanoic acid;
(3*S*,5*S*)-3-Aminomethyl-5-isopropoxy-hexanoic acid;
20 (3*S*,5*S*)-3-Aminomethyl-5-*tert*-butoxy-hexanoic acid;
(3*S*,5*S*)-3-Aminomethyl-5-fluoromethoxy-hexanoic acid;
(3*S*,5*S*)-3-Aminomethyl-5-(2-fluoro-ethoxy)-hexanoic acid;
(3*S*,5*S*)-3-Aminomethyl-5-(3,3,3-trifluoro-propoxy)-hexanoic acid;
(3*S*,5*S*)-3-Aminomethyl-5-phenoxy-hexanoic acid;
25 (3*S*,5*S*)-3-Aminomethyl-5-(4-chloro-phenoxy)-hexanoic acid;
(3*S*,5*S*)-3-Aminomethyl-5-(3-chloro-phenoxy)-hexanoic acid;
(3*S*,5*S*)-3-Aminomethyl-5-(2-chloro-phenoxy)-hexanoic acid;
(3*S*,5*S*)-3-Aminomethyl-5-(4-fluoro-phenoxy)-hexanoic acid;
(3*S*,5*S*)-3-Aminomethyl-5-(3-fluoro-phenoxy)-hexanoic acid;
30 (3*S*,5*S*)-3-Aminomethyl-5-(2-fluoro-phenoxy)-hexanoic acid;
(3*S*,5*S*)-3-Aminomethyl-5-(4-methoxy-phenoxy)-hexanoic acid;
(3*S*,5*S*)-3-Aminomethyl-5-(3-methoxy-phenoxy)-hexanoic acid;
(3*S*,5*S*)-3-Aminomethyl-5-(2-methoxy-phenoxy)-hexanoic acid;

-52-


- (3S,5S)-3-Aminomethyl-5-(4-nitro-phenoxy)-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-(3-nitro-phenoxy)-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-(2-nitro-phenoxy)-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl-6-propoxy-hexanoic acid;
5 (3S,5S)-3-Aminomethyl-6-isopropoxy-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-*tert*-butoxy-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-fluoromethoxy-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(2-fluoro-ethoxy)-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl-6-(3,3,3-trifluoro-propoxy)-hexanoic
10 acid;
(3S,5S)-3-Aminomethyl-5-methyl-6-phenoxy-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(4-chloro-phenoxy)-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(3-chloro-phenoxy)-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(2-chloro-phenoxy)-5-methyl-hexanoic acid;
15 (3S,5S)-3-Aminomethyl-6-(4-fluoro-phenoxy)-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(3-fluoro-phenoxy)-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(2-fluoro-phenoxy)-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(4-methoxy-phenoxy)-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(3-methoxy-phenoxy)-5-methyl-hexanoic acid;
20 (3S,5S)-3-Aminomethyl-6-(2-methoxy-phenoxy)-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl 6-(4-trifluoromethyl-phenoxy)-hexanoic
acid;
(3S,5S)-3-Aminomethyl-5-methyl 6-(3-trifluoromethyl-phenoxy)-hexanoic
acid;
25 (3S,5S)-3-Aminomethyl-5-methyl 6-(2-trifluoromethyl-phenoxy)-hexanoic
acid;
(3S,5S)-3-Aminomethyl-5-methyl 6-(4-nitro-phenoxy)-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl 6-(3-nitro-phenoxy)-hexanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl 6-(2-nitro-phenoxy)-hexanoic acid;
30 (3S,5S)-3-Aminomethyl-6-benzyloxy-5-methyl-hexanoic acid;
(3S,5R)-3-Aminomethyl-6-cyclopropyl-5-methyl-hexanoic acid;
(3S,5R)-3-Aminomethyl-6-cyclobutyl-5-methyl-hexanoic acid;
(3S,5R)-3-Aminomethyl-6-cyclopentyl-5-methyl-hexanoic acid;

-53-

- (3S,5R)-3-Aminomethyl-6-cyclohexyl-5-methyl-hexanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-heptanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-octanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-nonanoic acid;
5 (3S,5R)-3-Aminomethyl-5-methyl-decanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-undecanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-dodecanoic acid;
(3S,5R)-3-Aminomethyl-5,7-dimethyl-octanoic acid;
(3S,5R)-3-Aminomethyl-5,9-dimethyl-decanoic acid;
10 (3S,5R)-3-Aminomethyl-5,10-dimethyl-undecanoic acid;
(3S,5S)-3-Aminomethyl-5,6-dimethyl-heptanoic acid;
(3S,5S)-3-Aminomethyl-5,6,6-trimethyl-heptanoic acid;
(3S,5S)-3-Aminomethyl-5-cyclopropyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-fluoro-5-methyl-hexanoic acid;
15 (3S,5S)-3-Aminomethyl-7-fluoro-5-methyl-heptanoic acid;
(3S,5R)-3-Aminomethyl-8-fluoro-5-methyl-octanoic acid;
(3S,5S)-3-Aminomethyl-7,7,7-trifluoro-5-methyl-heptanoic acid;
(3S,5R)-3-Aminomethyl-8,8,8-trifluoro-5-methyl-octanoic acid;
(3S,5S)-3-Aminomethyl-5-methyl-6-phenyl-hexanoic acid;
20 (3S,5S)-3-Aminomethyl-6-(4-chloro-phenyl)-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(3-chloro-phenyl)-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(2-chloro-phenyl)-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(4-methoxy-phenyl)-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(3-methoxy-phenyl)-5-methyl-hexanoic acid;
25 (3S,5S)-3-Aminomethyl-6-(2-methoxy-phenyl)-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(3-fluoro-phenyl)-5-methyl-hexanoic acid;
(3S,5S)-3-Aminomethyl-6-(2-fluoro-phenyl)-5-methyl-hexanoic acid;
(3S,5R)-3-Aminomethyl-5-methyl-7-phenyl-heptanoic acid;
(3S,5R)-3-Aminomethyl-7-(4-chloro-phenyl)-5-methyl-heptanoic acid;
30 (3S,5R)-3-Aminomethyl-7-(3-chloro-phenyl)-5-methyl-heptanoic acid;
(3S,5R)-3-Aminomethyl-7-(2-chloro-phenyl)-5-methyl-heptanoic acid;
(3S,5R)-3-Aminomethyl-7-(4-methoxy-phenyl)-5-methyl-heptanoic acid;
(3S,5R)-3-Aminomethyl-7-(3-methoxy-phenyl)-5-methyl-heptanoic acid;

-54-

- (3S,5R)-3-Aminomethyl-7-(2-methoxy-phenyl)-5-methyl-heptanoic acid;
 (3S,5R)-3-Aminomethyl-7-(4-fluoro-phenyl)-5-methyl-heptanoic acid;
 (3S,5R)-3-Aminomethyl-7-(3-fluoro-phenyl)-5-methyl-heptanoic acid;
 (3S,5R)-3-Aminomethyl-7-(2-fluoro-phenyl)-5-methyl-heptanoic acid;
 5 (3S,5S)-3-Aminomethyl-5-methyl-hept-6-enoic acid;
 (3S,5R)-3-Aminomethyl-5-methyl-oct-7-enoic acid;
 (3S,5R)-3-Aminomethyl-5-methyl-non-8-enoic acid;
 (E)-(3S,5S)-3-Aminomethyl-5-methyl-oct-6-enoic acid;
 (Z)-(3S,5S)-3-Aminomethyl-5-methyl-oct-6-enoic acid;
 10 (Z)-(3S,5S)-3-Aminomethyl-5-methyl-non-6-enoic acid;
 (E)-(3S,5S)-3-Aminomethyl-5-methyl-non-6-enoic acid;
 (E)-(3S,5R)-3-Aminomethyl-5-methyl-non-7-enoic acid;
 (Z)-(3S,5R)-3-Aminomethyl-5-methyl-non-7-enoic acid;
 (Z)-(3S,5R)-3-Aminomethyl-5-methyl-dec-7-enoic acid; and
 15 (E)-(3S,5R)-3-Aminomethyl-5-methyl-undec-7-enoic acid.

Method 8

A compound of structure **60** can be prepared from a compound of structure **59** through treatment with an appropriately substituted phenol (including phenol itself) under conditions described by Mitsunobu, *Synthesis*, 1981:1.

A compound of structure **59** could be prepared from compound of structure **39** by treatment with sodium or lithium metal and alike in ammonia. Preferably, the reaction is carried out with sodium metal in ammonia.

-55-

The direct hydrolysis of compound **60** would give rise to the desired amino acid or the approach via hydrolysis of the Boc protected lactam could be utilized.

Compounds which could be prepared in this manner include:

- (3S)-3-Aminomethyl-5-methyl-7-phenoxy-heptanoic acid;
- 5 (3S)-3-Aminomethyl-7-(4-chloro-phenoxy)-5-methyl-heptanoic acid;
- (3S)-3-Aminomethyl-7-(3-chloro-phenoxy)-5-methyl-heptanoic acid;
- (3S)-3-Aminomethyl-7-(2-chloro-phenoxy)-5-methyl-heptanoic acid;
- (3S)-3-Aminomethyl-7-(4-fluoro-phenoxy)-5-methyl-heptanoic acid;
- (3S)-3-Aminomethyl-7-(3-fluoro-phenoxy)-5-methyl-heptanoic acid;
- 10 (3S)-3-Aminomethyl-7-(2-fluoro-phenoxy)-5-methyl-heptanoic acid;
- (3S)-3-Aminomethyl-7-(4-methoxy-phenoxy)-5-methyl-heptanoic acid;
- (3S)-3-Aminomethyl-7-(3-methoxy -phenoxy)-5-methyl-heptanoic acid;
- (3S)-3-Aminomethyl-7-(2-methoxy -phenoxy)-5-methyl-heptanoic acid;
- (3S)-3-Aminomethyl-5-methyl-7-(4-trifluoromethyl-phenoxy)-heptanoic
- 15 acid;
- (3S)-3-Aminomethyl-5-methyl-7-(3-trifluoromethyl-phenoxy)-heptanoic
- acid;
- (3S)-3-Aminomethyl-5-methyl-7-(2-trifluoromethyl-phenoxy)-heptanoic
- acid;
- 20 (3S)-3-Aminomethyl-5-methyl-7-(4-nitro-phenoxy)-heptanoic acid;
- (3S)-3-Aminomethyl-5-methyl-7-(3-nitro-phenoxy)-heptanoic acid;
- (3S)-3-Aminomethyl-5-methyl-7-(2-nitro-phenoxy)-heptanoic acid;
- (3S)-3-Aminomethyl-6-(3-chloro-phenoxy)-5-methyl-hexanoic acid;
- (3S)-3-Aminomethyl-6-(2-chloro-phenoxy)-5-methyl-hexanoic acid;
- 25 (3S)-3-Aminomethyl-6-(4-fluoro-phenoxy)-5-methyl-hexanoic acid;
- (3S)-3-Aminomethyl-6-(3-fluoro-phenoxy)-5-methyl-hexanoic acid;
- (3S)-3-Aminomethyl-6-(2-fluoro-phenoxy)-5-methyl-hexanoic acid;
- (3S)-3-Aminomethyl-6-(4-methoxy-phenoxy)-5-methyl-hexanoic acid;
- (3S)-3-Aminomethyl-6-(3-methoxy-phenoxy)-5-methyl-hexanoic acid;
- 30 (3S)-3-Aminomethyl-6-(2-methoxy-phenoxy)-5-methyl-hexanoic acid;
- (3S)-3-Aminomethyl-5-methyl 6-(4-trifluoromethyl-phenoxy)-hexanoic
- acid;

-56-

(3S)-3-Aminomethyl-5-methyl 6-(3-trifluoromethyl-phenoxy)-hexanoic acid;

(3S)-3-Aminomethyl-5-methyl 6-(2-trifluoromethyl-phenoxy)-hexanoic acid;

5 (3S)-3-Aminomethyl-5-methyl 6-(4-nitro-phenoxy)-hexanoic acid;

(3S)-3-Aminomethyl-5-methyl 6-(3-nitro-phenoxy)-hexanoic acid;


(3S)-3-Aminomethyl-5-methyl 6-(2-nitro-phenoxy)-hexanoic acid;

(3S)-3-Aminomethyl-5-methyl-6-phenoxy-hexanoic acid; and

(3S)-3-Aminomethyl-6-(4-chloro-phenoxy)-5-methyl-hexanoic acid.

10 Method 9

Synthesis of C-4 substituted analogs



A compound of structure **64** could be prepared from compound of structure **63** by treatment of **63** with hydrogen at 50 psi in the presence of a catalyst such as Raney nickel in the presence of a base such as triethyl amine in an organic solvent for example methanol. The resulting product is then treated with an aqueous acid such as 6N HCl at a temperature between room temperature and reflux. The resulting mixture could be subjected to ion exchange chromatography to isolate the product **64**.

20 A compound of structure **63** can be prepared from a compound of structure **62B** by treatment with an appropriate base, such as but not limited too sodium hydride, n-butyl lithium and alike, and an alkylating reagent such as t-butylbromoacetate or benzylbromoacetate in a solvent such as DMSO or THF an alike. Preferably, the reaction is carried out by treating a solution of a compound

-57-


of structure **62B** in THF with sodium hydride and alkylation of the resultant anion with t-butylbromoacetate.

A compound of structure **62B** can be prepared from a compound of structure **62A** by treatment with sodium chloride in a solvent such as aqueous DMSO at a temperature between 50°C and reflux.

A compound of structure **62A** can be prepared from a compound of structure **61** by treatment with an appropriate alkylmetalhalide such as an alkyllithium reagent or an organomagnesium halide in a solvent such as THF or ether in the presence of a copper salt, such as but not limited to copper iodide, copper bromide dimethylsulphide. Alternatively, the reaction may be carried out by the treatment of the nitrile in a solvent such as ether at, or below, room temperature with an alkylmagnesium chloride.

A compound such as **61** can be prepared according to known literature procedures between the condensation of isobutyraldehyde and methylcyanoacetate.

Method 10: C-4 Substitution



-58-

Doubly branched 3-substituted GABA analogs **72** can be prepared in two steps from the azide **71** through hydrogenation of the azide **71** in the presence of a noble metal catalyst such as 5% palladium on carbon and hydrolysis of the resulting lactam with a strong acid such as 6 N HCl at reflux. The final product **72** can then be isolated using ion exchange chromatography.

Compound **71** can be prepared in two steps by treatment of a lactone such as **70** with HBr in a solvent such as ethanol at a temperature such as 0°C and reacting the resulting bromide with sodium azide in a solvent such as dimethyl sulfoxide at a temperature between 10°C and 80°C.

Lactone **70** can be prepared in two steps by oxidation of a compound such as **69** with an oxidant such as sodium periodate in the presence of a catalytic amount of ruthenium trichloride in a solvent such as acetonitrile at a temperature between 0°C and 100°C and treatment of the resulting compound with potassium carbonate in methanol followed at a temperature between 25°C and 70°C and then treatment with an acid such as p-toluene sulfonic acid in a solvent such as THF at reflux or an aqueous acid such as HCl in water at ambient temperature.

A compound such as **69** can be prepared by a by reduction of a compound such as **68** with a hydride reducing agent such as lithium aluminum hydride in a solvent such as ether or THF and reaction of the resulting alcohol with an acylating agent such as acetic anhydride in the presence of a base such as triethyl amine or pyridine or the like.


Compounds of structure **68** can be prepared by reaction of a compound such as **67** with hydrogen at approximately 50 psi in the presence of a noble metal catalyst such as 5% palladium on carbon in a solvent such as ethanol. A compound of the formula **67** can be prepared by reaction of a compound of structure **66** with a solution of ethanol saturated with hydrogen bromide gas. A compound such as **66** can be prepared from a compound such as **65** by treatment of a compound such as one with a strong base such as lithium diisopropyl amine in a solvent such as THF at a temperature such as -78°C and reaction of the resulting anion with a compound such as benzyl bromide or benzyl iodide. Compounds of the structure **66** (R = H or lower alkyl) can be prepared in optical form from methods known in the literature (Davies, *J. Org. Chem.*,

-59-

1999;64(23):8501-8508; Koch J. *Org. Chem.*, 1993;58(10):2725-37; Afonso, *Tetrahedron*, 1993;49(20):4283-92; Bertus, *Tetrahedron*, Asymmetry 1999;10(7):1369-1380; Yamamoto, *J. Am. Chem. Soc.*, 1992;114(20):7652-60).

Specific Examples

5 Example 3: Synthesis of 3-Aminomethyl-5-methyl-octanoic acid



1-Benzyl-4-hydroxymethyl-pyrrolidine-2-one 74

Sodium borohydride (8.0 g, 0.211 mol) was added to a solution of methyl-1-benzyl-5-oxo-3-pyrrolidene-2-carboxylate **73** (See Zoretic et al, *J. Org. Chem.*, 1980;45:810-814 for general method of synthesis) (32.0 g, 0.137 mol) in 1,2-dimethoxyethane (600 mL) and refluxed for 19 hours. The reaction was cooled to room temperature and 200 mL of water was added. The reaction was quenched with 1 M citric acid and concentrated under reduced pressure. The residue was extracted with dichloromethane, dried over magnesium sulfate, and evaporated to dryness to give 17.47 g, 62% of the alcohol **74** as clear oil.

^1H NMR (CDCl_3) δ 7.30 (m, 5H), 4.38 (d, 1H, $J = 14.7$), 4.46 (d, 1H, $J = 14.7$), 3.56 (m, 2H), 3.36 (m, 1H), 3.10 (m, 1H), 2.52 (m, 2H), 2.26 (m, 1H). MS, m/z (relative intensity): 207 [$M+2H$, 66%]. IR (KBr) 3345, 2946, 2866, 1651, 1445, 1025, 737, and 698 cm^{-1} .

20 1-Benzyl-4-iodomethyl-pyrrolidin-2-one 75

To alcohol lactam **74** (11.18 g, 0.056 mol) in 210 mL toluene was added in turn, triphenylphosphine (20.0 g, 0.076 mol), imidazole (10.8 g, 0.159 mol), and

-60-

iodine (19.0 g, 0.075 mol). After stirring the suspension for 1.5 hours, the supernatant was poured into another flask. The sticky yellow residue was washed twice with ether and the solutions were combined. The solvent was evaporated and the residue was chromatographed on silica, eluting with 1:1 acetone/hexane to give 7.92 g, 46% of the iodolactam **75** as yellow oil. ^1H NMR (CDCl_3) δ 7.25 (m, 5H), 4.38 (d, 1H, $J = 14.6$), 4.46 (d, 1H, $J = 14.6$), 3.38 (dd, 1H, $J = 7.8$ and 2.2), 3.20 (dd, 1H, $J = 5.6$ and 4.4), 3.12 (dd, 1H, $J = 7.3$ and 2.4), 2.96 (dd, 1H, $J = 5.8$ and 4.4), 2.60 (m, 2H), 2.22 (dd, 1H, $J = 10.5$ and 9.7). MS, m/z (relative intensity): 224 [$\text{M}-\text{H}-\text{Bn}$, 94%], 317 [$\text{M}+2\text{H}$, 64%]. IR 3027, 2917, 1688, 1438, 1267, and 701 cm^{-1} .

1-Benzyl-4-(2-methyl-pentyl)-pyrrolidin-2-one **76**

To a suspension of magnesium turnings (0.50 g, 0.021 mol) in 15 mL of dry THF under nitrogen, was added an iodine crystal and 2-bromopentane (2.88 g, 0.019 mol). After an exothermic reaction which was periodically cooled in an ice bath, the reaction was stirred at room temperature for 2 hours. Eight milliliters of Li_2CuCl_4 (made from 84 mg LiCl and 134 mg CuCl_2 in 10 mL of dry THF) was added at 0°C followed by dropwise addition of 1-Benzyl-4-iodomethyl-pyrrolidine-2-one **75** in 15 mL dry THF, and the resulting suspension was let stir at 0°C for 3 hours. Stirring was continued at room temperature for 1 hour before quenching with a saturated solution of ammonium chloride. Water was added to dissolve the precipitate formed, and the solution was then extracted with ether and dried over magnesium sulfate. The solvent was evaporated under vacuum and the residue chromatographed on silica eluting with 1:1 acetone/hexane to give 1.13 g, 69% of the 1-benzyl-4-(2-methyl-pentyl)-pyrrolidin-2-one **76**. ^1H NMR (CDCl_3) δ 7.30 (m, 5H), 4.44 (m, 2H), 3.32 (m, 1H), 2.86 (m, 1H), 2.56 (m, 1H), 2.40 (m, 1H), 2.10 (m, 1H), 1.30 (m, 6H), 1.10 (m, 1H), 0.90 (m, 6H). MS, m/z (relative intensity): 261 [$\text{M}+2\text{H}$, 100%], 301 [$\text{M}-\text{H}+\text{CH}_3\text{CN}$, 82%], 260 [$\text{M}+\text{H}$, 72%].

4-(2-Methyl-pentyl)-pyrrolidin-2-one **77**

A 250 mL 3-neck flask equipped with a dry ice condenser was chilled to -78°C . Ammonia (80 mL) was condensed into the flask and 1-benzyl-4-(2-methyl-

-61-


pentyl)-pyrrolidin-2-one **76** (1.67 g, 0.006 mol) in 15 mL THF was added. Freshly cut sodium beads were added until a deep blue color persisted. The cooling bath was removed and the reaction stirred at reflux (-33°C) for 1 hour. The reaction was quenched with ammonium chloride and the excess ammonia was allowed to evaporate. The resulting residue was diluted with water, extracted with dichloromethane, and dried over magnesium sulfate. Evaporation of the solvent followed by chromatography on silica eluting with 1:1 acetone/hexane gave 0.94 g, 86% of the 4-(2-Methyl-pentyl)-pyrrolidin-2-one **77**. ¹H NMR (CDCl₃) δ 6.25 (br, 1H), 3.44 (m, 1H), 2.95 (m, 1H), 2.54 (m, 1H), 2.40 (m, 1H), 1.98 (m, 1H), 1.30 (m, 6H), 0.80 (m, 6H). MS, *m/z* (relative intensity): 212 [M+2H+CH₃CN, 100%], 171 [M+2H, 72%], 170 [M+1H, 65%].

3-Aminomethyl-5-methyl-octanoic acid (Example 3)

The 4-(2-methyl-pentyl)-pyrrolidin-2-one **77** (0.94 g, 0.007 mol) was dissolved in 70 mL of 6N HCl and refluxed for 20 hours. The solution was evaporated under vacuum and an aqueous solution of the residue was applied to Dowex 50WX 8-100 (strongly acidic) ion exchange resin that had been washed with HPLC grade water. The column was eluted, first with water until the eluent was at constant pH, and then with 5% ammonium hydroxide solution. The ammonium hydroxide fractions were evaporated and azeotroped with toluene. The white solid was washed with acetone filtered and dried in a vacuum oven for 24 hours to give the amino acid 0.61 g, 59%. ¹H NMR (CD₃OD) δ 3.00 (m, 1H), 2.85 (m, 1H), 2.48 (m, 1H), 2.30 (m, 1H), 2.14 (brm, 1H), 1.60 (brm, 1H), 1.38 (m, 4H), 1.18 (m, 2H), 0.60 (m, 6H). MS, *m/z* (relative intensity): 188 [M+H, 100%].

-62-

Example 4: Synthesis of 3-Aminomethyl-5,7-dimethyl-octanoic acid



1-(4-Methoxy-benzyl)-5-oxo-pyrrolidine-3-carboxylic acid methyl ester 79

5 To 4-methoxybenzylamine (42 g, 0.306 mol) in methanol (40 mL) at 0°C was added the dimethyl itaconate (48 g, 0.306 mol) in methanol (13 mL). The solution was stirred at room temperature for 4 days. 1N HCl was added to the solution followed by ether. The two layers were separated and the aqueous phase

-63-

extracted with ether. The combined organic phases were dried (MgSO₄). Upon filtration of the drying agent the desired material **79** precipitated from solution that was collected and dried under vacuum. 23.26 g, 29%. MS, *m/z* (relative intensity): 264 [M+H, 100%]. Anal. Calcd for C₁₄H₁₇N₁O₄: C, 63.87; H, 6.51; N, 5.32.

5 Found: C, 63.96; H, 6.55; N, 5.29.

4-Hydroxymethyl-1-(4-methoxy-benzyl)-pyrrolidine-2-one **80**

NaBH₄ (15 g, 0.081 mol) was added in portions to ester **79** in ethanol (600 mL) at room temperature. After 4.5 hours water (~200 mL) was carefully added to the reaction and the solution stirred at room temperature overnight. The resultant solid was removed by filtration and the filtrate concentrated to give alcohol **80** as an oil. 15.33 g, 81%. MS, *m/z* (relative intensity): 235 [M+H, 100%].

10

4-Iodomethyl-1-(4-methoxy-benzyl)-pyrrolidin-2-one **81**

To alcohol **80** (12.9 g, 0.055 mol) in PhMe was added triphenylphosphine (20 g, 0.077 mol), imidazole (10.8 g, 0.16 mol), and iodine (19 g, 0.075 mol). The suspension was stirred at room temperature 5 hours. A saturated aqueous solution of sodium thiosulphate was added and the two layers separated. The aqueous phase was extracted with ether and the combined organic phases washed with brine, dried (MgSO₄) and concentrated. Flash chromatography (6:1 to 4:1 toluene/acetone) of the residue gave iodide **81** as an oil. 11.9g, 63%. MS, *m/z* (relative intensity): 346 [M+H, 100%].

15

20

4-(2,4-Dimethyl-pentyl)-1-(4-methoxy-benzyl)-pyrrolidin-2-one **82**

A procedure similar to the preparation of 1-benzyl-4-(2-methyl-pentyl)-pyrrolidin-2-one **76** was utilized to give 4-(2,4-dimethyl-pentyl)-1-(4-methoxy-benzyl)-pyrrolidin-2-one as an oil. 1.22g, 29%. MS, *m/z* (relative intensity): 304 [M+H, 100%].

25

4-(2,4-Dimethyl-pentyl)-pyrrolidin-2-one **83**


-64-

To the lactam (1.17 g, 3.86 mmol) in MeCN (20 mL) at 0°C was added ceric ammonium nitrate (4.2 g, 7.7 mmol) in H₂O (10 mL). After 50 minutes a further portion of ceric ammonium nitrate (2.1 g, 3.86 mmol) was added, and after 1 hour the mixture was absorbed onto silica and flash chromatographed to give an oil. MS, *m/z* (relative intensity): 183 [M+H, 100%].

3-Aminomethyl-5,7-dimethyl-octanoic acid (Example 4)

A procedure similar to the preparation of 3-aminomethyl-5-methyl-octanoic acid (Example 3) was utilized to give the amino acid as a solid. MS, *m/z* (relative intensity): 202 [M+H, 100%].

Example 5: Synthesis of (S)-3-Aminomethyl-5-methyl-octanoic acid



(S)-4-Hydroxymethyl-1-((S)-1-phenylethyl)-pyrrolidin-2-one **84**

To the ester **33** (49 g, 0.198 mol) in EtOH (600 mL) was added sodium borohydride (22 g, 0.595 mol). After 7 hours, 1 M citric acid was carefully added and, after effervescence had ceased, water was added to fully quench the reaction. The ethanol was removed under reduced pressure and ethyl acetate added. The

-65-

resultant two layers were separated, the aqueous phase was extracted with EtOAc, and the combined organic phases dried (MgSO₄) and concentrated to give a heavy oil. MS, *m/z* (relative intensity): [M+H, 100%].

(S)-4-Iodomethyl-1-((S)-1-phenyl-ethyl)-pyrrolidin-2-one 85

5 A procedure similar to the iodination of compound **80** was utilized giving iodide **85** as an oil. 35.2 g, 56%. Anal. Calcd for C₁₃H₁₆I₁N₁O₁: C, 47.43; H, 4.90; N, 4.25. Found: C, 47.41; H, 4.83; N, 4.17.

4-(2-Methyl-pentyl)-1-((S)-1-phenyl-ethyl)-pyrrolidin-2-one 86

10 A procedure similar to the preparation of 1-benzyl-4-(2-methyl-pentyl)-pyrrolidin-2-one **76** was utilized to give 2.71 g, 81.0% of **86** as an oil. MS, *m/z* (relative intensity): 274 [M+1H, 100%], 315 [M+H+CH₃CN, 65%].

(S)-4-(2-Methyl-pentyl)-pyrrolidin-2-one 87


15 A procedure similar to the preparation of 4-(2-methyl-pentyl)-pyrrolidin-2-one **77** was used to give 1.14 g, 72.8% of **87** as an oil. MS, *m/z* (relative intensity): 170 [M+1H, 10%], 211 [M+1H+CH₃CN, 90%].

Example 5: (S)-3-Aminomethyl-5-methyl-octanoic acid

20 A procedure similar to the preparation of 3-aminomethyl-5-methyl-octanoic acid (Example 3) was used to give the amino acid (example 5) 0.88 g, 74.3%. ¹H NMR (CD₃OD) δ 2.95 (m, 1H), 2.80 (m, 1H), 2.40 (m, 1H), 2.25 (m, 1H), 2.05 (brm, 1H), 1.50 (brm, 1H), 1.30 (m, 4H), 1.10 (m, 2H), 0.90 (m, 6H). MS, *m/z* (relative intensity): 188 [M+1H, 100%], 186 [M-1H, 100%], 229 [M+1H+CH₃CN, 30%].

-66-

Example 6: Synthesis of (S)-3-Aminomethyl-7-methoxy-5-methyl-heptanoic acid



(S)-4-(2-Methyl-pent-4-enyl)-1-((S)-1-phenyl-ethyl)-pyrrolidin-2-one **88**

A procedure similar to the preparation of 1-benzyl-4-(2-methyl-pentyl)-pyrrolidin-2-one **76** was followed giving the adduct **88** as an oil. 6 g, 74%. MS, m/z (relative intensity): 272 [$\text{M}+\text{H}$, 100%].

(S)-4-(4-Hydroxy-2-methyl-butyl)-1-((S)-1-phenyl-ethyl)-pyrrolidin-2-one **89**

OsO_4 (2 mL of a 4% wt solution in $t\text{-BuOH}$) was added to the alkene **88** (5.8 g, 0.021 mol) in $\text{THF}/\text{H}_2\text{O}$ (3:1, 100 mL). After 1 hour, sodium periodate (11.4 g, 0.053 mol) was added. After 2 hours, the suspension was filtered and the solids washed with dichloromethane. The filtrate was concentrated and the residue azeotroped with toluene. The residue was dissolved in ethanol and sodium borohydride (2.5 g) added. The suspension was stirred at room temperature overnight. 1N citric acid was added and the mixture diluted with ether. The resultant two layers were separated and the aqueous phase was extracted with ether and the combined organic dried (MgSO_4) and concentrated. Flash

-67-

chromatography (1:1 hexane/EtOAc) of the residue gave an oil. 4.2 g, 73%. MS, m/z (relative intensity): 276 [M+H, 100%].

(S)-4-(4-Methoxy-2-methyl-butyl)-1-((S)-1-phenyl-ethyl)-pyrrolidin-2-one 90

To alcohol **89** (2 g, 7.66 mmol) in DMSO (60 mL) at room temperature was added NaH (368 mg, 60% in oil). After 30 minutes the methyl iodide (1.08 g, 7.66 mmol) was added and the solution stirred at room temperature overnight, upon which the reaction was diluted with water (500 mL). The solution was extracted with ether, and the combined organic extracts were dried (MgSO₄) and concentrated. Flash chromatography (90% to 50% hexane/acetone) of the residue gave the product **90** as an oil (1.1 g, 52%). MS m/z 290 (M+H, 100%).

(S)-4-(4-Methoxy-2-methyl-butyl)-pyrrolidin-2-one 91


A procedure similar to the synthesis of 4-(2-methyl-pentyl)-pyrrolidin-2-one **77** was utilized giving lactam **91** as an oil. MS m/z 186 (M+H, 100%).

Example 6: (S)-3-Aminomethyl-7-methoxy-5-methyl-heptanoic acid

A procedure similar to the synthesis of example 3 was followed. The resultant amino acid isolated from ion-exchange chromatography was recrystallized from methanol/ethyl acetate to give the example 6 as a white solid. MS m/z 204 (M+H, 100%). Anal. Calcd for C₁₀H₂₁N₁O₃: C, 59.09; H, 10.41; N, 6.89. Found: C, 58.71; H, 10.21; N, 6.67.

-68-

Example 7: Synthesis of (S)-3-Aminomethyl-6-fluoro-5-methyl-hexanoic acid

2-Methyl-2-[(S)-5-oxo-1-((S)-1-phenyl-ethyl)-pyrrolidin-3-ylmethyl]-malonic acid dimethyl ester **92**

To dimethyl methylmalonate (1.06 g, 7.29 mmol) in DMSO (7 mL) at room temperature was added NaH (291 mg of a 60% dispersion in oil). After the effervescence had ceased the lactam **85** (2 g, 7.29 mol) in DMSO (5 mL) was added. After 1 hour water was added and the aqueous solution extracted with ether. The combined organic extracts were dried (MgSO₄) and concentrated. Flash chromatography (1:1 hexane/acetone) of the residue gave the product as an oil (1.7 g, 81%). MS *m/z* 348 (M+H, 100%).

2-Methyl-3-[(S)-5-oxo-1-((S)-1-phenyl-ethyl)-pyrrolidin-3-yl]-propionic acid methyl ester **93**

The ester **92** (483 mg, 1.4 mmol), NaCl (104 mg, 1.8 mmol), water (105 μ L) and DMSO (5 mL) were heated to reflux for 2 hours. The solution was cooled to room temperature water was added and the aqueous solution extracted

-69-

with ether. The combined organic extracts were dried (MgSO_4) and concentrated. Flash chromatography (80% to 66% hexane/acetone) of the residue gave the product as an oil (160 mg, 40%). MS m/z 290 (M+H, 100%).

(S)-4-(3-Hydroxy-2-methyl-propyl)-1-((S)-1-phenyl-ethyl)-pyrrolidin-2-one
37

To the ester **93** (4.82 g, 0.017 mol) in EtOH (100 mL) was added NaBH_4 (3.7 g, 0.10 mol) and the mixture heated to reflux for 2.5 hours. The solution was cooled to 0°C and 1 M citric acid carefully added followed by water. The solution was concentrated to half volume added and extracted with ether. The combined organic extracts were dried (MgSO_4) and concentrated. Flash chromatography (1:1 hexane/acetone) of the residue gave the product as an oil (2.6 g, 59%). MS m/z 262 (M+H, 100%).

(S)-4-(3-Fluoro-2-methyl-propyl)-1-((S)-1-phenyl-ethyl)-pyrrolidin-2-one 94

To DAST (1 g, 6.2 mmol) in CH_2Cl_2 (20 mL) at -78°C was added the alcohol **37** in CH_2Cl_2 (10 mL). After 1 hour at -78°C the solution was warmed to room temperature. After 7 hours the solution was carefully quenched with a saturated aqueous solution of sodium bicarbonate and the two layers separated. The organic phase was dried (MgSO_4) and concentrated. Flash chromatography (90% to 66% hexane/acetone) of the residue gave the product as an oil (600 mg, 37%). MS m/z 264 (M+H, 100%).

(S)-4-(3-Fluoro-2-methyl-propyl)-pyrrolidin-2-one 95

A procedure similar to the preparation of 4-(2-methyl-pentyl)-pyrrolidin-2-one **77** was utilized affording the lactam as an oil (242 mg, 68%). MS m/z 159 (M, 100%).


Example 7 (S)-3-Aminomethyl-6-fluoro-5-methyl-hexanoic acid

A procedure similar to the synthesis of example 3 was followed. The resultant amino acid isolated from ion-exchange chromatography was recrystallized from methanol/ethyl acetate to give example 7 as a white solid. MS

-70-

m/z 177 (M, 100%). Anal. Calcd for $C_8H_{16}F_1N_1O_2 \cdot 0.02 H_2O$: C, 54.11; H, 9.10; N, 7.89. Found: C, 53.75; H, 9.24; N, 7.72.

Example 8: Synthesis of (S)-3-Aminomethyl-6-methoxy-5-methyl-hexanoic acid



Example 8

(S)-4-(3-Methoxy-2-methyl-propyl)-1-((S)-1-phenyl-ethyl)-pyrrolidin-2-one
96

A procedure similar to the synthesis of (S)-4-(4-methoxy-2-methyl-butyl)-1-((S)-1-phenyl-ethyl)-pyrrolidin-2-one **90** was utilized giving ether **96** as an oil (90 mg, 37%). MS m/z 276 (M+H, 100%).

(S)-4-(3-Methoxy-2-methyl-propyl)-pyrrolidin-2-one 97

A procedure similar to the synthesis of 4-(2-methyl-pentyl)-pyrrolidin-2-one **77** was utilized giving **97** as an oil (760 mg, 93%). MS m/z 171 (M+H, 100%).


Example 8 (S)-3-Aminomethyl-6-methoxy-5-methyl-hexanoic acid

A procedure similar to the synthesis of example 3 was followed. The resultant amino acid isolated from ion-exchange chromatography was recrystallized from methanol/ethyl acetate to give Example 8 as a white solid. MS m/z 190 (M+H, 100%). Anal. Calcd for $C_9H_{19}N_1O_3$: C, 57.12; H, 10.12; N,

-71-

7.40. Found: C, 57.04; H, 10.37; N, 7.30. A second batch precipitated from the mother liquors (1:5 ratio of C5 isomers by ^1H NMR). MS m/z 190 ($\text{M}+\text{H}$, 100%).

Example 9: Synthesis of (3S,5R)-3-Aminomethyl-5-methyl-octanoic acid hydrochloride



5

(R)-2,6-Dimethyl-non-2-ene 98

To (S)-citronellyl bromide (50 g, 0.228 mol) in THF (800 mL) at 0°C was added LiCl (4.3 g) followed by CuCl_2 (6.8 g). After 30 minutes methylmagnesium chloride (152 mL of a 3 M solution in THF, Aldrich) was added and the solution warmed to room temperature. After 10 hours the solution was cooled to 0°C and a saturated aqueous solution of ammonium chloride carefully added. The resultant two layers were separated and the aqueous phase extracted with ether. The combined organic phases were dried (MgSO_4) and

10

-72-

concentrated to give an oil. 32.6 g; 93%. Used without further purification. ^{13}C NMR (100 MHz; CDCl_3) 131.13, 125.28, 39.50, 37.35, 32.35, 25.92, 25.77, 20.31, 19.74, 17.81, 14.60.

(R)-4-Methyl-heptanoic acid 99

5 To alkene **98** (20 g, 0.13 mol) in acetone (433 mL) was added a solution of CrO_3 (39 g, 0.39 mol) in H_2SO_4 (33 mL)/ H_2O (146 mL) over 50 minutes. After 6 hours a further amount of CrO_3 (26 g, 0.26 mol) in H_2SO_4 (22 mL)/ H_2O (100 mL) was added. After 12 hours the solution was diluted with brine and the solution extracted with ether. The combined organic phases were dried (MgSO_4)
10 and concentrated. Flash chromatography (gradient of 6:1 to 2:1 hexane/EtOAc) gave the product **99** as an oil. 12.1 g; 65%. MS, m/z (relative intensity): 143 [M-H, 100%].

(4R,5S)-4-Methyl-3-((R)-4-methyl-heptanoyl)-5-phenyl-oxazolidin-2-one 100

15 To the acid **99** (19 g, 0.132 mol) and triethylamine (49.9 g, 0.494 mol) in THF (500 mL) at 0°C was added trimethylacetylchloride (20 g, 0.17 mol). After 1 hour LiCl (7.1 g, 0.17 mol) was added followed by the oxazolidinone (30 g, 0.17 mol). The mixture was warmed to room temperature and after 16 hours the filtrate was removed by filtration and the solution concentrated under reduced pressure. Flash chromatography (7:1 hexane/EtOAc) gave the product **100** as an
20 oil. 31.5 g; 79%. $[\alpha]_{\text{D}} = 5.5$ (c 1 in CHCl_3). MS, m/z (relative intensity): 304 [M+H, 100%].

(3S,5R)-5-Methyl-3-((4R,5S)-4-methyl-2-oxo-5-phenyl-oxazolidine-3-carbonyl)-octanoic acid tert-butyl ester 101

25 To oxazolidinone **100** (12.1 g, 0.04 mol) in THF (200 mL) at -50°C was added NaHMDS (48 mL of a 1 M solution in THF). After 30 t-butylbromoacetate (15.6 g, 0.08 mol) was added. The solution was stirred for 4 hours at -50°C and then warmed to room temperature. After 16 hours a saturated aqueous solution of ammonium chloride was added and the two layers separated. The aqueous phase was extracted with ether and the combined organic phases dried (MgSO_4) and

-73-

concentrated. Flash chromatography (9:1 hexane/EtOAc) gave the product **101** as a white solid 12 g; 72%. $[\alpha]_D = 30.2$ (c 1 in CHCl_3). ^{13}C NMR (100 MHz; CDCl_3) 176.47, 171.24, 152.72, 133.63, 128.87, 125.86, 80.85, 78.88, 55.34, 39.98, 38.77, 38.15, 37.58, 30.60, 28.23, 20.38, 20.13, 14.50, 14.28.

5 **(S)-2-((R)-2-Methyl-pentyl)-succinic acid 4-tert-butyl ester 102**

To ester **101** (10.8 g, 0.025 mol) in H_2O (73 mL) and THF (244 mL) at 0°C was added a premixed solution of LiOH (51.2 mL of a 0.8 M solution) and H_2O_2 (14.6 mL of a 30% solution). After 4 hours a further 12.8 mL LiOH (0.8 M solution) and 3.65 mL of H_2O_2 (30% solution) was added. After 30 minutes
10 sodium bisulfite (7 g), sodium sulfite (13 g), and water (60 mL) was added followed by hexane (100 mL) and ether (100 mL). The two layers were separated and the aqueous layer extracted with ether. The combined organic phases were concentrated to an oil that was dissolved in heptane (300 mL). The resultant solid was filtered off and the filtrate dried (MgSO_4) and concentrated to afford an oil
15 (6 g, 93%) which was used without further purification. MS, m/z (relative intensity): 257 $[\text{M}+\text{H}, 100\%]$.

(3S,5R)-3-Hydroxymethyl-5-methyl-octanoic acid tert-butyl ester 103

To acid **102** (3.68 g, 0.014 mol) in THF (100 mL) at 0°C was added $\text{BH}_3\cdot\text{Me}_2$ (36 mL of a 2 M solution in THF, Aldrich) upon which the solution was
20 warmed to room temperature. After 15 hours ice was **carefully** added (in order to control the effervescence) to the solution followed by brine. The solution was extracted with ether and the combined organic phases dried (MgSO_4) and concentrated under reduced pressure. Flash chromatography (4:1 hexane/EtOAc) gave alcohol **103** as an oil (2.0 g, 59%). ^{13}C NMR (100 MHz; CDCl_3) 173.56,
25 80.85, 65.91, 39.74, 39.20, 38.90, 35.65, 29.99, 28.31, 20.18, 19.99, 14.56.

-74-

(3S,5R)-5-Methyl-3-(toluene-4-sulfonyloxymethyl)-octanoic acid tert-butyl ester 104

To alcohol **103** (1.98 g, 8.1 mmol) in CH₂Cl₂ (40 mL) at room temperature was added triethylamine (2.4 g, 0.024 mol), DMAP (20 mg) and tosyl chloride (2.3 g, 0.012 mol). After 14 hours 1N HCl was added and the two layers separated. The aqueous phase was extracted with ether and the combined organic phases dried (MgSO₄) and concentrated. Flash chromatography (95% hexane/EtOAc) gave tosylate **104** as an oil (2.94 g, 91%). ¹³C NMR (100 MHz; CDCl₃) 171.60, 144.92, 133.07, 130.02, 128.12, 80.80, 72.15, 39.73, 38.09, 37.89, 32.67, 29.71, 28.22, 21.83, 20.10, 19.54, 14.49.

(3S,5R)-3-Azidomethyl-5-methyl-octanoic acid tert-butyl ester 105

Tosylate **104** (2.92 g, 7.3 mmol) and sodium azide (1.43 g, 0.02 mol) were warmed to ~50°C in DMSO (30 mL). After 2 hours the solution was cooled to room temperature and diluted with water. The solution was extracted with ether and the combined organic phases dried (MgSO₄) and concentrated to give an oil 1.54 g, 79%. Further purification by flash chromatography (95% hexane/EtOAc) gave an oil. [α]_D = -8.3 (c 1 in CHCl₃). ¹³C NMR (100 MHz; CDCl₃) 172.01, 80.73, 54.89, 39.73, 39.46, 39.00, 33.40, 29.85, 28.30, 20.15, 19.82, 14.52.

(S)-4-((R)-2-Methyl-pentyl)-pyrrolidin-2-one 107 and (3S,5R)-3-aminomethyl-5-methyl-octanoic acid tert-butyl ester 106

Azide **105** was treated with 5% Pd/C and shaken under an atmosphere of hydrogen for 20 hours where upon a further 200 mg of 5% Pd/C added. After 6 hours the filtrate was concentrated to afford an oil which by ¹H NMR was found to be a mixture of primary amine **106** and lactam **107** (1.75 g) which was used without further purification.

Example 9 (3S,5R)-3-Aminomethyl-5-methyl-octanoic acid hydrochloride


The mixture of the amine **106** and the lactam **107** (1.74 g) was treated with 3N HCl (40 mL) and the solution warmed to 50°C for 4 hours then cooled to room temperature. After 12 hours the solution was concentrated and the residue

-75-

recrystallized from ethyl acetate to give the amino acid as a white solid 605 mg. MS, m/z (relative intensity): 188 [M+H, 100%]. Anal. Calcd for $C_{10}H_{21}N_1O_2 \cdot H_1Cl_1$ C, 53.68; H, 9.91; N, 6.26. Found: C, 53.83; H, 10.12; N, 6.07.

-76-

Example 10: Synthesis of (3S,5R)-3-Aminomethyl-5-methyl-heptanoic acid



Methanesulfonic acid (S)-3,7-dimethyl-oct-6-enyl ester 108

To S-(-)-citronellol (42.8 g, 0.274 mol) and triethylamine (91 mL, 0.657 mol) in CH₂Cl₂ (800 mL) at 0°C was added methanesulphonyl chloride (26 mL, 0.329 mol) in CH₂Cl₂ (200 mL). After 2 hours at 0°C the solution was washed with 1N HCl then brine. The organic phase was dried (MgSO₄) and concentrated to afford an oil (60.5 g, 94%) which was used without further

-77-

purification. ^1H NMR (400 MHz; CDCl_3) 5.05 (1H, m), 4.2 (2H, m), 2.95 (3H, s), 1.98 (2H, m), 1.75 (1H, m), 1.6 (3H, s), 1.5 (4H, m), 1.35 (2H, m), 1.2 (1H, m), 0.91 (3H, d, $J = 6.5$ Hz).

(R)-2,6-Dimethyl-oct-2-ene 109

5 To alkene **108** (60 g, 0.256 mol) in THF (1 L) at 0°C was added lithium aluminum hydride (3.8 g, 0.128 mol). After 7 hours, a further 3.8 g of lithium aluminum hydride was added and the solution warmed to room temperature. After 18 hours, a further 3.8 g of lithium aluminum hydride was added. After a further 21 hours, the reaction was carefully quenched with 1N citric acid and the solution
10 diluted further with brine. The resultant two phases were separated and the organic phase was dried (MgSO_4) and concentrated to afford an oil which was used without further purification. MS, m/z (relative intensity): 139 [M-H, 100%].

(R)-4-Methyl-hexanoic acid 110

A procedure similar to the synthesis of (R)-4-methyl-heptanoic acid **99**
15 was utilized giving the acid as an oil (9.3 g, 56%). MS, m/z (relative intensity): 129 [M-H, 100%].

(4R, 5S)-4-Methyl-3-((R)-4-methyl-hexanoyl)-5-phenyl-oxazolidin-2-one 111

A procedure similar to the synthesis of (4R,5S)-4-methyl-3-((R)-4-methyl-heptanoyl)-5-phenyl-oxazolidin-2-one **100** was utilized giving oxazolidinone **111**
20 as an oil (35.7 g, 95%). MS, m/z (relative intensity): 290 [M+H, 100%].

(3S,5R)-5-Methyl-3-[1-((4R,5S)-4-methyl-2-oxo-5-phenyl-oxazolidin-3-yl)-methanoyl]-heptanoic acid tert-butyl ester 112

A procedure similar to the preparation of (3S,5R)-5-methyl-3-((4R,5S)-4-methyl-2-oxo-5-phenyl-oxazolidine-3-carbonyl)-octanoic acid tert-butyl ester **101**
25 was followed giving **112** as an oil (7.48 g; 31%).

(S)-2-((R)-2-Methyl-butyl)-succinic acid 4-tert-butyl ester 113

To ester **112** (7.26 g, 0.018 mol) in H_2O (53 mL) and THF (176 mL) at 0°C was added a premixed solution of LiOH (37 mL of a 0.8 M solution) and

-78-

H₂O₂ (10.57 mL of a 30% solution) and the solution warmed to room temperature. After 2 hours sodium bisulfite (7 g), sodium sulfite (13 g), and water (60 mL) was added and the two layers were separated and the aqueous layer extracted with ether. The combined organic phases were concentrated to an oil that was dissolved in heptane (200 mL). The resultant solid was filtered off and the filtrate dried (MgSO₄) and concentrated to afford an oil (4.4 g) that was used without further purification.

(3S,5R)-3-Hydroxymethyl-5-methyl-heptanoic acid tert-butyl ester 114

A procedure similar to the preparation of (3S,5R)-3-hydroxymethyl-5-methyl-octanoic acid tert-butyl ester **103** was utilized giving alcohol **114** as an oil (2.68 g, 69%). MS, *m/z* (relative intensity): 216 [89%], 174 [M-(CH₃)₃C, 100%].

(3S,5R)-5-Methyl-3-(toluene-4-sulfonyloxymethyl)-heptanoic acid tert-butyl ester 115

To **114** alcohol (2.53 g, 0.011 mmol) in CH₂Cl₂ (140 mL) at 0°C was added pyridine (2.6 g, 0.033 mol), DMAP (100 mg), and tosyl chloride (3.15 g, 0.016 mol) and the solution warmed to room temperature for 3.5 hours whereupon more DMAP and TsCl (3.15 g) were added. After 14 hours 1N HCl was added and the two layers separated. The organic phase was washed with brine then or dried (MgSO₄) and concentrated. Flash chromatography (95% to 86% hexane/EtOAc) gave tosylate **115** as an oil (1.53 g, 36%). ¹³C NMR (100 MHz; CDCl₃) 130.03, 128.12, 72.18, 37.89, 37.71, 32.67, 31.49, 29.88, 28.22, 21.83, 19.07, 11.37.

(3S,5R)-3-Azidomethyl-5-methyl-heptanoic acid tert-butyl ester 116

A procedure similar to the preparation of (3S,5R)-3-azidomethyl-5-methyl-octanoic acid tert-butyl ester **105** was utilized giving an oil 0.956 g, 97%. MS, *m/z* (relative intensity): 228 [M-N₂, 80%].

-79-

(S)-4-((R)-2-Methyl-butyl)-pyrrolidin-2-one 118 and (3S,5R)-3-Aminomethyl-5-methyl-heptanoic acid tert-butyl ester 117


5 Azide **116** (689 mg) was treated with 20% Pd/C (90 mg) in THF (20 mL) and shaken under an atmosphere of hydrogen for 36 hours. The catalyst was removed by filtration and the resultant oil used without further purification.

Example 10 (3S,5R)-3-Aminomethyl-5-methyl-heptanoic acid

10 The mixture of amine **117** and lactam **118** was treated with 6N HCl and the solution warmed to 50°C for 17 hours then cooled to room temperature and concentrated. The resultant oil was subjected to ion-exchange chromatography (Dowex, strongly acidic resin) using 5% ammonium hydroxide to give a cream solid which was recrystallized from methanol/ethyl acetate to give (3S, 5R)-3-aminomethyl-5-methyl-heptanoic acid, example 10. MS, *m/z* (relative intensity): 174 [M+H, 100%]. Anal. Calcd for C₁₉H₁₉N₁O₂. C, 62.39; H, 11.05; N, 8.08. Found: C, 62.23; H, 11.33; N, 7.89.

-80-

Example 11: Synthesis of (3S,5S)-3-Aminomethyl-5-methyl-octanoic acid



Example 11

(S)-2,6-Dimethyl-non-2-ene 119

5 CuCl₂ (5.36 g, 39.7 mmol) and LiCl (3.36, 80.0 mmol) were stirred together in dry THF (40 mL) for 15 minutes. The resulting solution was added to methylmagnesium chloride, 3.0 M in THF (168 mL) at 0°C under nitrogen atmosphere and stirred at that temperature for 15 minutes. To the reaction suspension was added slowly (R)-(-)-Citronellyl bromide (55.16 g, 251.8 mmol) in THF (100 mL), and stirred at 0°C for 2.5 hours. It was warmed to room temperature and stirring was continued for an additional 1 hour. The mixture was cooled to 0°C and quenched with saturated ammonium chloride solution. The suspension was then extracted into ether, washed with water, and dried over MgSO₄. The solution was concentrated under reduced pressure to afford 36.3 g;

10

-81-

94% of (S)-2,6-Dimethyl-non-2-ene as an oil. MS, m/z (relative intensity): 153 [M-1H, 100%], 194 [M-1H+CH₃CN, 45%].

(S)-4-Methyl-heptanoic acid 120

To the (S)-2,6-Dimethyl-non-2-ene **119** (39.0 g, 253.2 mmol) in acetone
5 (1L) at 0°C was added Jones reagent (2.7 M, 600 mL) dropwise over 1.5 hours
and let stir at room temperature for 18 hours. The reaction mixture was poured
into a saturated solution of Na₂SO₄ and extracted into ether. It was washed with
brine and concentrated in vacuo. The oily residue was dissolved in methanol
(70 mL) and 1 M NaOH (700 mL) and then stirred for 30 minutes. The aqueous
10 solution was washed with CH₂Cl₂, acidified with 10% HCl and extracted into
CH₂Cl₂. The solution was dried over MgSO₄ and concentrated to dryness to give
24.22 g; 66% of (S)-4-Methyl-heptanoic acid as an oil. MS, m/z (relative
intensity): 143 [M-1H, 100%].

(4R,5S)-4-Methyl-3-((S)-4-methyl-heptanoyl)-5-phenyl-oxazolidin-2-one 121

15 A procedure similar to the preparation of (4R,5S)-4-methyl-3-((R)-4-
methyl-heptanoyl)-5-phenyl-oxazolidin-2-one **100** was utilized giving (4R,5S)-4-
methyl-3-((S)-4-methyl-heptanoyl)-5-phenyl-oxazolidin-2-one **121** 6.2 g ; 80.0%,
as an oil. MS, m/z (relative intensity): 304 [M+1H, 90%], 355 [M+1H+CH₃CN,
60%].

20 **(3S,5S)-5-Methyl-3-((4R,5S)-4-methyl-2-oxo-5-phenyl-oxazolidine-3-
carbonyl)-octanoic acid tert-butyl ester 122**

n-BuLi, 1.6 M in Hexane (18.0 mL, 30.1 mmol) was added dropwise to a
solution of diisopropylamine (4.6 mL, 32.6 mmol) in dry THF (50 mL) under
nitrogen at -5°C keeping the temperature below 0°C during addition. The mixture
25 was let stir at -5°C for 20 minutes and then cooled to -78°C. **121** (7.6 g,
25.1 mmol) in dry THF (12 mL) was added to the LDA solution and stirred at
-78°C for 30 minutes. *t*-Butylbromo acetate (4.8 mL, 32.6 mmol) as added to the
reaction and stirring at -78°C was continued for 2 hours. It was let warm to room
temperature before stirring for an additional 18 hours. The reaction was quenched

-82-

with a saturated solution NaH_2PO_4 , extracted into ethylacetate, and dried over MgSO_4 . The solution was concentrated to give a solid residue which was dissolved in hot hexane. The hexane solution was allowed to cool to room temperature before cooling further in an ice bath. The resulting precipitate was collected and allowed to air dry to give **122** as a fluffy white solid. 4.3 g; 41%. MS, m/z (relative intensity): 362 [$\text{M}-\text{C}(\text{CH}_3)_3+1\text{H}$, 100%], 418 [$\text{M}+1\text{H}$, 20%].

(S)-2-((S)-2-Methyl-pentyl)-succinic acid 4-tert-butyl ester and (3S,5S)-3-Hydroxymethyl-5-methyl-octanoic acid tert-butyl ester 123

To the ester **122** in a mixture of THF (203.0 mL) and water (61.0 mL) at 0°C was added a premixed solution of 30% H_2O_2 (12.2 mL) and LiOH (0.8 M, 42.7 mL). The resulting solution was stirred at 0°C for 4 hours. To the reaction was added sodium bisulfite (7 g), sodium sulfite (13 g), and water (60 mL). A 1:1 mixture of ether/hexane (200 mL) was then added and the organic phase was separated. The aqueous phase was extracted with ether and the combined organic extract was dried over MgSO_4 and concentrated in vacuo. The residue was dissolved in heptane and let stir for 5 minutes. The resulting precipitate was filtered and the filtrate was concentrated to dryness to give as an oil.

(3S,5S)-3-Hydroxymethyl-5-methyl-octanoic acid tert-butyl ester 123

A procedure similar to the preparation of (3S,5R)-3-hydroxymethyl-5-methyl-octanoic acid tert-butyl ester **103** was followed giving **123** as an oil. 4.0 g; 76.0%. MS, m/z (relative intensity): 230 [$\text{M}-\text{C}(\text{CH}_3)_3+1\text{H}+\text{CH}_3\text{CN}$, 100%], 189 [$\text{M}-\text{C}(\text{CH}_3)_3+1\text{H}$, 70%].

(3S,5S)-5-Methyl-3-(toluene-4-sulfonyloxymethyl)-octanoic acid tert-butyl ester 124

A procedure similar to the preparation of (3S,5R)-5-methyl-3-(toluene-4-sulfonyloxymethyl)-octanoic acid tert-butyl ester **104** was followed giving 6.9 g of **124**. MS, m/z (relative intensity): 343 [$\text{M}-\text{C}(\text{CH}_3)_3+1\text{H}$, 70%], 384 [$\text{M}-\text{C}(\text{CH}_3)_3+1\text{H}+\text{CH}_3\text{CN}$, 100%].

-83-

(3S,5S)-3-Azidomethyl-5-methyl-heptanoic acid tert-butyl ester 125

A procedure similar to the preparation of (3S,5R)-3-azidomethyl-5-methyl-octanoic acid tert-butyl ester **105** was followed giving 2.9 g; 66% of **125** as an oil. MS, m/z (relative intensity): 212 [M-C(CH₃)₃ -1H, 45%].

5 **(3S,5S)-3-Aminomethyl-5-methyl-octanoic acid tert-butyl ester 126**


A mixture of **125** (2.8 g, 10.4 mmol) and 10% Pd/C (1.0 g) in methanol (50.0 mL) was hydrogenated at 41 PSI for 96 hours. The solution was filtered to give 1.7 g of crude **126** which was used in the next step without further purification. MS, m/z (relative intensity): 244 [M +1H, 100%], 285
10 [M+1H+CH₃CN, 25%].

Example 11 (3S,5S)-3-Aminomethyl-5-methyl-octanoic acid

A procedure similar to the preparation of example 10 (3S,5R)-3-aminomethyl-5-methyl-heptanoic acid was followed giving example 11. 380 mg; 29.0%. ¹H NMR (CD₃OD) δ 2.90 (dd, J = 3.9, 8.8 Hz, 1H), 2.80 (dd, J = 7.6, 5.1
15 Hz, 1H), 2.40 (dd, J = 3.2, 12.51 Hz, 1H), 2.20 (dd, J = 8.8, 6.8 Hz, 1H), 2.05 (m, 1H), 1.55 (m, 1H), 1.30 (m, 3H), 1.10 (m, 2H), 0.85 (m, 6H); MS, m/z (relative intensity): 187 [M+1H, 100%], 211 [M+1H+CH₃CN, 30%].

-84-

Example 12: Synthesis of (3S,5S)-3-Aminomethyl-5-methyl-heptanoic acid

(S)-2,6-Dimethyl-oct-2-ene **127**

(R)-(-)-Citronellyl bromide (49.1 g, 224.2 mmol) was dropwise added to a solution of LAH 1.0 M in THF (336 mL, 336 mmol) at 0°C over a 45-minute period. Stirring was continued for an additional 4 hours at 0°C. The reaction was slowly quenched with a saturated solution of ammonium chloride followed by the addition of ether (100 mL). The resulting white slurry was filtered and the filtrate was dried over MgSO₄. The solution was concentrated under reduced pressure to afford 26.2 g; 83% of **127** as an oil. MS, *m/z* (relative intensity): 180 [M-1H+CH₃CN, 100%], 139 [M-1H, 90%].

-85-

(S)-4-Methyl-hexanoic acid 128

A procedure similar to that used to prepare compound **120** was used giving 15.9 g of **128** as an oil. MS, *m/z* (relative intensity): 129 [M-1H, 100%], 170 [M-1H+CH₃CN, 70%].

5 **(4R,5S)-4-Methyl-3-((S)-4-methyl-hexanoyl)-5-phenyl-oxazolidin-2-one 129**

A procedure similar to that used to prepare (4R,5S)-4-Methyl-3-((S)-4-methyl-heptanoyl)-5-phenyl-oxazolidin-2-one **121** was used giving 35.0 g of crude (4R,5S)-4-methyl-3-((S)-4-methyl-hexanoyl)-5-phenyl-oxazolidin-2-one **129** as an oil. It was used in the next step without further purification. MS, *m/z* (relative intensity): 290 [M+1H, 100%], 331 [M+1H+CH₃CN, 20%].

10

(3S,5S)-5-Methyl-3-((4R,5S)-4-methyl-2-oxo-5-phenyl-oxazolidine-3-carbonyl)-heptanoic acid tert-butyl ester 130

A procedure similar to that used to prepare (3S,5S)-5-methyl-3-((4R,5S)-4-methyl-2-oxo-5-phenyl-oxazolidine-3-carbonyl)-octanoic acid tert-butyl ester **122** was used to give 4.6.0 g, 25.4% of **130** as a white solid. MS, *m/z* (relative intensity): 348 [M-C(CH₃)₃+1H, 100%], 443 [M-1H+CH₃CN, 100%], 402 [M-1H, 55%], 404 [M+1H, 45%].

15

(3S,5S)-3-Hydroxymethyl-5-methyl-heptanoic acid tert-butyl ester 131

A procedure similar to that used to prepare (3S,5S)-3-Hydroxymethyl-5-methyl-octanoic acid tert-butyl ester **123** was giving 1.2 g, 52.1% of **131** as an oil. MS, *m/z* (relative intensity): 175 [M-C(CH₃)₃+1H, 100%], 173 [M-C(CH₃)₃-1H, 100%], 216 [M-C(CH₃)₃+1H +CH₃CN, 95%].

20

(3S,5S)-5-Methyl-3-(toluene-4-sulfonyloxymethyl)-heptanoic acid tert-butyl ester 132

A procedure similar to the preparation of (3S,5R)-5-methyl-3-(toluene-4-sulfonyloxymethyl)-octanoic acid tert-butyl ester **104** was followed giving 2.1 g of **132** as an oil. The product was used in the next step without further

25

-86-

purification. MS, m/z (relative intensity): 329 [M-C(CH₃)₃+1H, 85%], 370 [M-C(CH₃)₃+1H +CH₃CN, 65%].

(3S,5S)-3-Azidomethyl-5-methyl-heptanoic acid tert-butyl ester 133

5 A procedure similar to the preparation of (3S,5R)-3-azidomethyl-5-methyl-octanoic acid tert-butyl ester **105** was followed giving 0.76 g, 54.0% of **133** as an oil. MS, m/z (relative intensity): 198 [M-C(CH₃)₃-1H, 100%]

(3S,5S)-3-Aminomethyl-5-methyl-heptanoic acid tert-butyl ester 134


10 A procedure similar to that used for (3S,5S)-3-aminomethyl-5-methyl-octanoic acid tert-butyl ester **126** was used giving 0.62 g of **134** as an oil. The product was used in the next step without further purification. MS, m/z (relative intensity): 230 [M+1H, 100%], 271 [M+1H +CH₃CN, 45%].

Example 12 (3S,5S)-3-Aminomethyl-5-methyl-heptanoic acid

15 A procedure similar to that used for Example 11 was used giving (3S,5S)-3-aminomethyl-5-methyl-heptanoic acid (0.3 g, 65.1%) as a white solid. ¹H NMR (CD₃OD) δ 2.80-3.00 (m, 2H), 2.40 (m, 1H), 2.20 (dd, J = 8.2, 7.1 Hz, 1H), 2.05 (m, 1H), 1.30-1.50 (m, 3H), 1.00-1.20 (m, 2H), 0.9 (m, 6H); MS, m/z (relative intensity): 187 [M+1H, 100%], 211 [M+1H+CH₃CN, 30%]. MS, m/z (relative intensity): 174 [M+1H, 100%], 172 [M-1H, 100%], 215 [M+1H +CH₃CN, 20%].

-87-

Example 13: Synthesis of (3*S*,5*R*)-3-Aminomethyl-5-methyl-nonanoic acid hydrochloride



(R)-4-Methyl-octanoic acid **136**

5 Lithium chloride (0.39 g, 9.12 mmol) and copper (I) chloride (0.61 g, 4.56 mmol) were combined in 45 ml THF at ambient temperature and stirred 15 minutes, then cooled to 0°C at which time ethylmagnesium bromide (1 M solution in THF, 45 mL, 45 mmol) was added. (S)-citronellyl bromide (5.0 g, 22.8 mmol) was added dropwise and the solution was allowed to warm slowly to ambient temperature with stirring overnight. The reaction was quenched by cautious addition of sat. NH₄Cl (aq), and stirred with Et₂O and sat. NH₄Cl (aq) for 30 minutes. The phases were separated and the organic phase dried (MgSO₄) and concentrated. The crude product was used without purification.

15 To a solution of alkene **135** (3.8 g, 22.8 mmol) in 50 mL acetone at 0°C was added Jones' reagent (2.7 M in H₂SO₄ (aq), 40 mL, 108 mmol) and the solution was allowed to warm slowly to ambient temperature with stirring overnight. The mixture was partitioned between Et₂O and H₂O, the phases were

-88-

separated, and the organic phase washed with brine, dried (MgSO₄), and concentrated. The residue was purified by flash chromatography (8:1 hexanes:EtOAc) to afford 2.14 g (59%) of acid **136** as a colorless oil: LRMS: *m/z* 156.9 (M⁺); ¹H NMR (CDCl₃): δ 2.33 (m, 2H), 1.66 (m, 1H), 1.43 (m, 2H), 1.23 (m, 5H), 1.10 (m, 1H), 0.86 (m, 6H). Jones' reagent was prepared as a 2.7M solution by combining 26.7g CrO₃, 23 mL H₂SO₄, and diluting to 100 mL with H₂O.

(4R, 5S)-4-Methyl-3-((R)-4-methyl-octanoyl)-5-phenyl-oxazolidin-2-one 137

To acid **136** (2.14 g, 13.5 mmol) in 25 mL CH₂Cl₂ at 0°C was added 3 drops DMF, followed by oxalyl chloride (1.42 mL, 16.2 mmol) resulting in vigorous gas evolution. The solution was warmed directly to ambient temperature, stirred 30 minutes, and concentrated. Meanwhile, to a solution of the oxazolidinone (2.64 g, 14.9 mmol) in 40 mL THF at -78°C was added *n*-butyllithium (1.6 M soln in hexanes, 9.3 mL, 14.9 mmol) dropwise. The mixture was stirred for 10 minutes at which time the acid chloride in 10 mL THF was added dropwise. The reaction was stirred 30 minutes at -78°C, then warmed directly to ambient temperature and quenched with sat. NH₄Cl. The mixture was partitioned between Et₂O and sat. NH₄Cl (aq), the phases were separated, and the organic phase dried (MgSO₄), and concentrated to furnish 3.2 g of oxazolidinone **137** as a colorless oil. LRMS: *m/z* 318.2 (M⁺); ¹H NMR (CDCl₃): δ 7.34 (m, 5H), 5.64 (d, *J* = 7.3 Hz, 1H), 4.73 (quint, *J* = 6.8 Hz, 1H), 2.96 (m, 1H), 2.86 (m, 1H), 1.66 (m, 1H), 1.47 (m, 2H), 1.26 (m, 5H), 1.13 (m, 1H), 0.88 (m, 9H). The crude product was used without purification.

(3S,5R)-5-Methyl-3-((4R,5S)-4-methyl-2-oxo-5-phenyl-oxazolidine-3-carbonyl)-nonanoic acid tert-butyl ester 138

To a solution of diisopropylamine (1.8 mL, 12.6 mmol) in 30 mL THF at -78°C was added *n*-butyllithium (1.6 M soln in hexanes, 7.6 mL, 12.1 mmol), and the mixture stirred 10 minutes at which time oxazolidinone **137** (3.2 g, 10.1 mmol) in 10 mL THF was added dropwise. The solution was stirred for

-89-

30 minutes, *t*-butyl bromoacetate (1.8 mL, 12.1 mmol) was added quickly dropwise at -50°C, and the mixture was allowed to warm slowly to 10°C over 3 hours. The mixture was partitioned between Et₂O and sat. NH₄Cl (aq), the phases were separated, and the organic phase dried (MgSO₄), and concentrated.

5 The residue was purified by flash chromatography (16:1 to 8:1 hexanes:EtOAc) to provide 2.65 g (61%) of ester **138** as a colorless crystalline solid, mp = 84-86°C.

[α]_D²³ +17.1 (c = 1.00, CHCl₃); ¹H NMR (CDCl₃): δ 7.34 (m, 5H), 5.62 (d, *J* = 7.3 Hz, 1H), 4.73 (quint, *J* = 6.8 Hz, 1H), 4.29 (m, 1H), 2.67 (dd, *J* = 9.8, 16.4 Hz, 1H), 2.40 (dd, *J* = 5.1, 16.4 Hz, 1H), 1.69 (m, 1H), 1.38 (s, 9H), 1.28 (m, 10 7H), 1.08 (m, 1H), 0.88 (m, 9H); ¹³C NMR (CDCl₃) δ 176.45, 171.22, 152.71, 133.64, 128.86, 125.86, 80.83, 78.87, 55.33, 40.02, 38.21, 37.59, 36.31, 30.86, 29.29, 28.22, 23.14, 20.41, 14.36, 14.26. Anal. Calcd for C₂₅H₃₇NO₅: C, 69.58; H, 8.64; N, 3.25. Found: C, 69.37; H, 8.68; N, 3.05.

(S)-2-((R)-2-Methyl-hexyl)-succinic acid 4-tert-butyl ester 139

15 To a solution of ester **138** (2.65 g, 6.14 mmol) in 20 mL THF at 0°C was added a precooled (0°C) solution of LiOH monohydrate (1.0 g, 23.8 mmol) and hydrogen peroxide (30 wt% aqueous soln, 5.0 mL) in 10 mL H₂O. The mixture was stirred vigorously for 90 minutes, then warmed to ambient temperature and stirred 90 minutes. The reaction was quenched at 0°C by addition of 100 mL 10%
20 NaHSO₃ (aq), then extracted with Et₂O. The phases were separated, and the organic phase washed with brine, dried (MgSO₄), and concentrated. The crude acid **139** was used without purification.

(3S,5R)-3-Hydroxymethyl-5-methyl-nonanoic acid tert-butyl ester 140

25 To a solution of the crude acid **139** (6.14 mmol) in 30 mL THF at 0°C was added borane-dimethyl sulfide complex (2.0 M soln in THF, 4.6 mL, 9.2 mmol), and the mixture was allowed to warm slowly to ambient temperature overnight. Additional BH₃-DMS was added until the acid was completely consumed (ca. 5 mL). The reaction was quenched by addition of MeOH, then partitioned between Et₂O and sat. NaHCO₃ (aq). The phases were separated, and the organic

-90-

phase washed with brine, dried (MgSO₄), and concentrated to provide alcohol

140. LRMS: m/z 226.1; ¹H NMR (CDCl₃): δ 3.63 (dd, J = 11.0, 4.2 Hz, 1H), 3.42 (dd, J = 11.0, 6.8 Hz, 1H), 2.30 (dd, J = 14.9, 7.6 Hz, 1H), 2.20 (dd, J = 14.9, 5.6 Hz, 1H), 2.03 (m, 2H), 1.42 (s, 9H), 1.24 (m, 6H), 1.02 (m, 2H), 0.85 (m, 6H).

5 The crude product was used without purification.

(3S,5R)-5-Methyl-3-(toluene-4-sulfonyloxymethyl)-nonanoic acid tert-butyl ester 141

To alcohol **140** (6.14 mmol) in 30 mL CH₂Cl₂ at 0°C was added DMAP (0.1 g), *p*-toluenesulfonyl chloride (1.37 g, 7.2 mmol), and then triethylamine (1.8 mL, 13 mmol) was added quickly dropwise. The mixture was warmed immediately to ambient temperature following addition and stirred overnight, and did not proceed to completion. The mixture was partitioned between Et₂O and 1N HCl (aq), the phases were separated, and the organic phase washed with sat. NaHCO₃ (aq), dried (MgSO₄), and concentrated to provide tosylate **141**. The product was used without further purification.

(3S,5R)-3-Azidomethyl-5-methyl-nonanoic acid tert-butyl ester 142

A procedure similar to the preparation of (3S,5R)-3-azidomethyl-5-methyl-octanoic acid tert-butyl ester **105** was followed giving azide **142** as a colorless oil. LRMS: m/z 200.1; ¹H NMR (CDCl₃): δ 3.31 (dd, J = 12.2, 4.2 Hz, 1H), 3.19 (dd, J = 12.2, 5.9 Hz, 1H), 2.22 (m, 1H), 2.10 (m, 1H), 1.39 (s, 9H), 1.21 (m, 8H), 1.00 (m, 2H), 0.81 (m, 6H).


Example 13 (3S,5R)-3-Aminomethyl-5-methyl-nonanoic acid hydrochloride

The azide **142** (1.0 g) was hydrogenated in the presence of 20% Pd/C, EtOH, at 45 psi of H₂ for 15 hours to provide the crude amino ester **143** which was concentrated and used without purification. To the amino ester **143** was added 6 mL 6N HCl (aq) and the mixture was heated to reflux 90 minutes, cooled, and concentrated. Recrystallization from EtOAc:hexanes provided 0.38 g (45% from azide) of (3S,5R)-3-aminomethyl-5-methyl-nonanoic acid hydrochloride as a colorless crystalline solid (HCl salt), and a second crop of 82 mg (10% from

-91-

azide) was also obtained. mp = 146-156°C. LRMS: m/z 200.1 (M⁺); ¹H NMR (CDCl₃): δ 2.87 (dd, J = 13.2, 5.4 Hz, 1H), 2.79 (dd, J = 13.2, 7.3 Hz, 1H), 2.29 (d, J = 6.8 Hz, 2H), 2.08 (m, 1H), 1.31 (m, 1H), 1.09 (m, 7H), 0.92 (m, 1H), 0.68 (m, 6H). Anal. Calcd for C₁₁H₂₄NO₂Cl: C, 55.57; H, 10.17; N, 5.89. Found: C, 55.69; H, 10.10; N, 5.86.

Example 14: Synthesis of (3S, 5S)-3-Aminomethyl-5-methyl-nonanoic acid



The (S)-acid **145** was prepared from (R)-citronellyl bromide according to the procedure outlined above for (R)-4-methyl-octanoic acid **136**. The yield was comparable and the ¹H NMR spectrum was identical to that of the (R)-acid enantiomer. LRMS: m/z 158.9 (M⁺).

Oxazolidinone **146** was prepared from acid **145** as described above for (4R, 5S)-4-methyl-3-((R)-4-methyl-octanoyl)-5-phenyl-oxazolidin-2-one **137**. LRMS: m/z 290.1 (M-27); ¹H NMR (CDCl₃): δ 7.38 (m, 3H), 7.28 (m, 2H), 5.64 (d, J = 7.1 Hz, 1H), 4.74 (quint, J = 6.8 Hz, 1H), 2.92 (m, 2H), 1.71 (m, 1H), 1.42 (m, 7H), 1.18 (m, 1H), 0.88 (m, 9H).

-92-

t-Butyl ester **147** was prepared from oxazolidinone **146** as described above for compound **138**. LRMS: m/z 348.1 (M-83).


Alcohol **149** was prepared from the *t*-butyl ester **147** as described above for (3S,5R)-3-hydroxymethyl-5-methyl-nonanoic acid tert-butyl ester **140**. LRMS: m/z 156.9 (M-100); ^1H NMR (CDCl_3): δ 3.60 (dd, $J = 11.0, 4.6$ Hz, 1H), 3.45 (dd, $J = 11.0, 6.8$ Hz, 1H), 2.24 (m, 2H), 2.04 (m, 2H), 1.42 (s, 9H), 1.17-1.38 (m, 7H), 1.11 (m, 1H), 0.84 (m, 6H).

Example 14: (3S, 5S)-3-Aminomethyl-5-methyl-nonanoic acid

(3S, 5S)-3-Aminomethyl-5-methyl-nonanoic acid was obtained from **149** as described above for (3S,5R)-3-aminomethyl-5-methyl-nonanoic acid hydrochloride. The crude HCl salt thus obtained was purified by ion exchange chromatography on Dowex 50WX8 50-100 mesh, H-Form resin, using 10% NH_4OH as eluant to provide the free base. The waxy solid was washed twice with Et_2O and dried to furnish an amorphous white solid, mp 144-146°C. LRMS: m/z 172.0 (M-28); ^1H NMR (CDCl_3): δ 2.76 (d, $J = 5.9$ Hz, 2H), 2.14 (m, 1H), 1.96 (m, 2H), 1.25 (m, 1H), 1.12 (m, 6H), 0.96 (m, 2H), 0.66 (m, 6H).

-93-

Example 15: Synthesis of (3S,5R)-3-Aminomethyl-5-methyl-decanoic acid



Example 15

(R)-2,6-Dimethylundec-2-ene **153**

A procedure similar to the preparation of (S)-2,6-dimethyl-non-2-ene **119** was used giving **153** as a colorless oil (20.16 g, 98%). ^1H NMR (400 MHz, CDCl_3) δ 5.10-5.06 (m, 1H), 2.10-1.89 (m, 2H), 1.66 (s, 3H), 1.58 (s, 3H), 1.34-1.23 (m, 4H), 1.15-1.06 (m, 2H), 0.88-0.81 (m, 11H).

(R)-4-methylnonanoic acid **154**

(R)-2,6-Dimethylundec-2-ene **153** (10.03 g, 55.03 mmol) was dissolved in acetone (270 mL) and cooled to 0°C . Jones reagent ($\text{CrO}_3/\text{H}_2\text{SO}_4$) (2.7 M, 120 mL) was added dropwise, and the reaction allowed to warm to room temperature over 18 hours. The reaction was poured on to water/ Na_2SO_4 (200 mL), and the aqueous layer extracted with ethyl acetate (4×100 mL). The combined organics were dried over MgSO_4 , filtered and rotovapped to give an oil.

The crude oil was dissolved in CH_2Cl_2 (400 mL) and cooled to -78°C . Ozone was

-94-

bubbled into reaction until blue to remove traces of the impurity (6E)(3S)-3,7-dimethylocta-1,6-diene. Dimethylsulfide (5 mL) was added, and the reaction stirred at room temperature for 2 hours. The solvent was removed, and the crude material chromatographed on silica eluting with 20% EtOAc/hex to give oil. The oil was dissolved in ether (100 mL) and extracted with 10% NaOH (2 × 25 mL). The aqueous layers were combined and extracted with ether (50 mL). The aqueous layer was cooled to 0°C and acidified with HCl. The acidic layer was extracted with EtOAc (3 × 100 mL), and the combined extracts dried over MgSO₄, filtered and rotovapped to give **154** as an oil (6.86 g, 54%). ¹H NMR (400 MHz, CDCl₃) δ 2.40-2.25 (m, 4H), 1.70-1.62 (m, 2H), 1.47-1.11 (m, 8H), 0.87-0.84 (m, 6H); [α]_D = -11.4 (c1 in CHCl₃).

(4R,5S)-4-Methyl-3-((R)-4-methyl-nonanoyl)-5-phenyl-oxazolidin-2-one 155

Compound **154** (6.504 g, 37.76 mmol) was dissolved in THF (95 mL) and cooled to 0°C. Triethylamine (19.74 mL, 141.6 mmol) was added dropwise, followed by dropwise addition of trimethylacetyl chloride (6.98 mL, 56.64 mmol). The thick white suspension was stirred at 0°C for 90 minutes. LiCl (1.86 g, 41.54 mmol), (4R)-4-methyl-5-phenyl-1,3-oxazolidin-2-one (6.824 g, 38.51 mmol), and THF (70 mL) were added, and the reaction warmed to room temperature overnight. The solvent was evaporated. The solids were taken up in EtOAc, filtered off, and washed generously with EtOAc. The filtrate was washed with water (2 × 50 mL), and brine. The organics were dried over MgSO₄, filtered, and rotovapped. The crude material was chromatographed on silica eluting with 10% EtOAc/hexanes to give **155** as an oil (10.974 g, 88%). ¹H NMR (400 MHz, CDCl₃) δ 7.44-7.35 (m, 3H), 7.31-7.26 (m, 2H), 5.66 (d, *J* = 7.33 Hz, 1H), 4.76 (quin, *J* = 7.03 Hz, 1H), 3.04-2.96 (m, 1H), 2.93-2.86 (m, 1H), 1.74-1.66 (m, 1H), 1.52-1.47 (m, 1H), 1.46-1.36 (m, 2H), 1.27-1.16 (m, 2H), 0.92-0.87 (m, 8H); [α]_D = +34.1 (c1 in CHCl₃).

-95-

(3S,5R)-5-Methyl-3-((4R,5S)-4-methyl-2-oxo-5-phenyl-oxazolidine-3-carbonyl)-decanoic acid tert-butyl ester 156

A procedure similar to the preparation of (3S,5S)-5-methyl-3-((4R,5S)-4-methyl-2-oxo-5-phenyl-oxazolidine-3-carbonyl)-octanoic acid tert-butyl ester **122** was followed giving (3S,5R)-5-methyl-3-((4R,5S)-4-methyl-2-oxo-5-phenyl-oxazolidine-3-carbonyl)-decanoic acid tert-butyl ester **156** as an oil (0.668g, 90%). ¹H NMR (400 MHz, CDCl₃) δ 7.41-7.28 (m, 5H), 5.63 (d, *J* = 7.33 Hz, 1H), 4.74 (quin, *J* = 6.84 Hz, 1H), 4.33-4.26 (m, 1H), 2.68 (dd, *J* = 16.4, 9.77 Hz, 1H), 2.41 (dd, *J* = 16.6, 4.88 Hz, 1H), 1.68 (quin, *J* = 6.6 Hz, 1H), 1.50-1.32 (m, 10H), 1.28-1.21 (m, 1H), 1.15-1.08 (m, 1H), 0.90-0.86 (m, 9H); MS (APCI) *m/z* 348 (M⁺-97, 100%); [α]_D = +18.8 (c1 in CHCl₃).

(S)-2-((R)-2-Methyl-heptyl)-succinic acid 4-tert-butyl ester 157

Compound **156** (5.608 g, 12.59 mmol) was dissolved in THF/H₂O (60 mL/14 mL) and cooled to 0°C. LiOH (1N, 18.89 mL) and H₂O₂ (35%, 4.45 mL, 50.4 mmol) were combined, and then added to the reaction dropwise keeping T < 5°C. the reaction was stirred at 0°C for 4 hours, and quenched with Na₂SO₃ (6.3 g) and NaHSO₃ (3.4 g) in 50 mL H₂O added dropwise. The reaction was stirred for 15 minutes, and the layers separated. The aqueous layer was extracted with EtOAc (3 × 100 mL), and the combined extracts dried over MgSO₄, filtered, and rotovapped to give an oil. The crude material was dissolved in EtOAc (10 mL) and added dropwise to heptane (250 mL). The suspension was stirred for 20 minutes, and the solids filtered and washed with heptane. The filtrate was washed with 60°C H₂O (100 mL), dried over MgSO₄, filtered, and rotovapped to give **157** as an oil (3.52 g). the material was used directly in the next step.

(3S,5R)-3-Hydroxymethyl-5-methyl-decanoic acid tert-butyl ester 158

Compound **157** (3.52 g, 12.3 mmol) was dissolved in anhydrous THF (123 mL) and cooled to 0°C. Borane dimethylsulfide complex (10 M, 3.69 mL) was added dropwise, and the reaction then warmed to room temperature and

-96-

stirred for 1 hour. The reaction was cooled to 0°C, and quenched with MeOH (20 mL) added dropwise. The reaction was stirred for 18 hours, and the solvent rotovapped off. The crude material was chromatographed on silica eluting with 20% EtOAc/hexanes to give **158** (2.28 g, 68%) as an oil. ¹H NMR (400 MHz, CDCl₃) δ 3.65-3.59 (m, 1H), 3.43 (dd, *J* = 11.1, 6.96 Hz, 1H), 2.31 (dd, *J* = 14.9, 7.57 Hz, 1H), 2.21 (dd, *J* = 15.1, 5.62 Hz, 1H), 2.06-2.02 (m, 1H), 1.43 (s, 9H), 1.40-1.25 (m, 4H), 1.07-1.13 (m, 1H), 1.03-0.96 (m, 1H), 0.86-0.84 (m, 6H); MS (APCI) *m/z* 216 (*M*⁺-56, 100%).

(3S,5R)-5-Methyl-3-(toluene-4-sulfonyloxymethyl)-decanoic acid tert-butyl ester 159

Compound **158** (2.27 g, 8.33 mmol) was dissolved in CH₂Cl₂ (30 mL) and cooled to 0°C. Tosyl chloride (1.91 g, 10.0 mmol) and catalytic DMAP were added, followed by dropwise addition of triethylamine (2.55 mL, 18.33 mmol). The reaction was then stirred at 0°C for 18 hours. The solvent was rotovapped off (removed under reduced pressure), and the crude material washed with EtOAc and filtered. The solids were washed with EtOAc, and the filtrate washed with 0.5N HCl (20 mL), brine (30 mL), dried over MgSO₄, filtered and rotovapped. The oil was chromatographed on silica eluting with a 5% EtOAc/hexanes gradient to 10% EtOAc/hexanes to give **159** (3.399 g, 96%) as an oil. ¹H NMR (400 MHz, CDCl₃) δ 7.75 (d, *J* = 8.30 Hz, 2H), 7.31 (d, *J* = 8.30 Hz, 2H), 3.99 (dd, *J* = 9.65, 3.54 Hz, 1H), 3.89 (dd, *J* = 9.52, 5.37 Hz, 1H), 2.42 (s, 3H), 2.28 (dd, *J* = 14.7, 6.23 Hz, 1H), 2.19-2.14 (m, 1H), 2.10 (dd, *J* = 14.9, 6.35 Hz, 1H), 1.38 (s, 9H), 1.31-1.17 (m, 3H), 1.08-0.81 (m, 2H), 0.79-0.76 (m, 6H); [α]_D = -10.1 (c1 in CHCl₃).

(3S,5R)-3-Azidomethyl-5-methyl-decanoic acid tert-butyl ester 160

Compound **159** (3.01 g, 7.05 mmol), sodium azide (1.26 g, 19.40 mmol) and DMSO (12 mL) were combined and heated to 60°C for 3 hours. EtOAc (100 mL) was added to the reaction and filtered. The solids were washed with EtOAc (20 mL), and the filtrate evaporated. The crude material was

-97-

chromatographed on silica eluting with 5% EtOAc/hexanes to give **160** as an oil (1.86 g, 89%).

(3S,5R)-3-Aminomethyl-5-methyl-decanoic acid tert-butyl ester 161


5 A solution of compound **160** (1.86 g, 6.25 mmol) in THF (50 mL) was shaken over 5% Pd/C under hydrogen and pressure for 8 hours with three purges of hydrogen. The catalyst was filtered off and the filtrate evaporated. The crude material was chromatographed on silica eluting with methanol to give **161** as an oil (1.21 g, 71%). ¹H NMR (400 MHz, CDCl₃) δ 2.70 (dd, *J* = 12.9, 4.40 Hz, 1H), 2.54 (dd, *J* = 12.7, 6.59 Hz, 1H), 2.26 (dd, *J* = 14.5, 6.96, 1H), 2.12 (dd, *J* = 14.5, 6.47 Hz, 1H), 1.91 (m, 1H), 1.91 (m, 1H), 1.43 (s, 12H), 1.39-1.25 (m, 4H), 1.14-1.07 (m, 1H), 1.03-0.97 (m, 1H), 0.86-0.82 (m, 6H).

Example 15 (3S,5R)-3-Aminomethyl-5-methyl-decanoic acid

15 Compound **161** (1.20 g, 4.44 mmol) was heated to 50°C in 3N HCl (30 mL) for 4 hours. The solvent was evaporated, and the oil washed with toluene, and evaporated. The crude material was passed through an ion exchange column (Dowex 50WX8-100, strongly acidic) eluting with water, then 0.5N NH₄OH. Isolate (3S,5R)-3-aminomethyl-5-methyl-decanoic acid as a white solid (0.725 g, 75%): mp = 174-175°C; ¹H NMR (400 MHz, CDCl₃) δ 2.83 (dd, *J* = 12.69, 4.88 Hz, 1H), 2.70 (dd, *J* = 13.1, 7.45 Hz, 1H), 2.08 (d, *J* = 6.59 Hz, 2H), 1.98 (m, 1H), 1.28-1.20 (m, 1H), 1.19-1.09 (m, 2H), 0.99-0.91 (m, 2H), 0.66 (m, 6H); MS (APCI) *m/z* 215 (M⁺, 10%), 174 (M⁺-41, 100%); [α]_D = -5.7 (c1.025 in H₂O).

-98-

Example 16: Synthesis of (3S,5S)-3-Aminomethyl-5-methyl-decanoic acid

(S)-2,6-Dimethyl-undec-2-ene **162**

nPropylmagnesium chloride/ether solution (2.0 M, 228 mL) was cooled to
 5 -20°C under a N₂ atmosphere. LiCl (3.87 g, 91.25 mmol), CuCl₂ (6.13 g,
 45.63 mmol), and distilled THF (456 mL) were combined and stirred for
 30 minutes. The Li₂CuCl₄ solution was added via cannula to the Grignard
 reagent, and the resulting solution stirred for 30 minutes at -20°C. R-(-)-
 Citronellyl bromide (50 g, 228.1 mmol) was dissolved in THF (60 mL) and added
 10 dropwise to the Grignard solution. The reaction was stirred at 0°C for 1 hour. The
 reaction was cooled to -40°C and quenched with NH₄Cl (sat'd, 200 mL) added
 dropwise. The layers were separated and the aqueous layer extracted with ether
 (3 × 100 mL). The combined organics were dried over MgSO₄, filtered, and
 rotovapped to give an oil. The crude material was chromatographed on silica
 15 eluting with hexanes to give **162** as a colorless oil (9.15 g, 22%). ¹H NMR (400

-99-

MHz, CDCl₃) δ 5.10-5.06 (m, 1H), 2.10-1.89 (m, 2H), 1.66 (s, 3H), 1.58 (s, 3H), 1.34-1.23 (m, 4H), 1.15-1.06 (m, 2H), 0.88-0.81 (m, 11H).

(S)-4-Methylnonanoic acid 163

Compound **162** (7.97 g, 43.7 mmol) was dissolved in acetone (214 mL) and cooled to 0°C. Jones reagent (CrO₃/H₂SO₄) (2.7 M, 95 mL) was added dropwise, and the reaction allowed to warm to room temperature over 18 hours. The reaction was poured on to water/Na₂SO₄ (200 mL), and the aqueous layer extracted with ethyl acetate (4 × 100 mL). The combined organics were dried over MgSO₄, filtered, and rotovapped to give an oil. The crude oil was chromatographed on silica eluting with hexanes to give **163** as an oil (5.56 g, 74%). ¹H NMR (400 MHz, CDCl₃) δ 2.40-2.25 (m, 4H), 1.70-1.62 (m, 2H), 1.47-1.11 (m, 8H), 0.87-0.84 (m, 6H); MS APCI *m/z* 170.9 (M⁺-1, 100%).

(4R,5S)-4-Methyl-3-((S)-4-methyl-nonanoyl)-5-phenyl-oxazolidin-2-one 164

A procedure similar to that used to prepare compound **155** was used except that (S)-4-methylnonanoic acid **163** (5.56 g, 32.27 mmol) was used as a reactant to give **164** as an oil (10.70 g 100%). ¹H NMR (400 MHz, CDCl₃) δ 7.42-7.34 (m, 3H), 7.28 (d, *J* = 6.59 Hz, 2H), 5.64 (d, *J* = 7.33 Hz, 1H), 4.74 (quin, *J* = 6.78 Hz, 1H), 2.94-2.85 (m, 2H), 1.73-1.67 (m, 1H), 1.47-1.43 (m, 1H), 1.39-1.22 (m, 7H), 0.90-0.84 (m, 8H).

(3S,5S)-5-Methyl-3-((4R,5S)-4-methyl-2-oxo-5-phenyl-oxazolidine-3-carbonyl)-decanoic acid tert-butyl ester 165

A procedure similar to that used to prepare compound **156** was used to give **165** as a solid (4.25 g, 61%). MS (APCI) *m/z* 446 (M⁺+1, 10%), 390 (M⁺-55, 100%, -tBu).

(S)-2-((S)-2-Methyl-heptyl)-succinic acid 4-tert-butyl ester 166

A procedure similar to that used for compound **157** was used except that ester **165** (8.42 g, 18.89 mmol) was used as a reactant to give **166** as an oil

-100-

(5.81 g). The material was used directly in the next step. MS (APCI) m/z 285 ($M-1$, 100%).

(3S,5S)-3-Hydroxymethyl-5-methyl-decanoic acid tert-butyl ester 167

5 A procedure similar to that used to prepare compound **158** was used except that (S)-2-((S)-2-methyl-heptyl)-succinic acid 4-tert-butyl ester **166** (5.78 g, 20.18 mmol) was used as a reactant to give **167** as an oil (4.18 g, 76%). ^1H NMR (400 MHz, CDCl_3) δ 3.64-3.58 (m, 1H), 3.84-3.42 (m, 1H), 2.28-2.20 (m, 1H), 2.09-2.02 (m, 1H), 1.43 (s, 9H), 1.26-1.18 (m, 8H), 1.11-1.04 (m, 2H), 0.87-0.83 (m, 6H); MS (APCI) m/z 217 (M^+ -55, 50%, -tBu).

10 **(3S,5S)-5-Methyl-3-(toluene-4-sulfonyloxymethyl)-decanoic acid tert-butyl ester 168**

A procedure similar to that used to prepare compound **159** was used except that (3S,5S)-3-Hydroxymethyl-5-methyl-decanoic acid tert-butyl ester **167** (4.164 g, 15.29 mmol) was used as a reactant to give **168** as an oil (4.17 g, 64%).
15 ^1H NMR (400 MHz, CDCl_3) δ 7.75 (d, $J = 8.30$ Hz, 2H), 7.31 (d, $J = 8.30$ Hz, 2H), 3.97 (dd, $J = 9.52$, 4.15 Hz, 1H), 3.90 (dd, $J = 9.52$, 5.13 Hz, 1H), 2.42 (s, 3H), 2.28, 2.19-2.13 (m, 2H), 1.37 (s, 9H), 1.27-1.01 (m, 11H), 0.85 (t, $J = 7.08$ Hz, 3H), 0.76 (d, $J = 6.35$ Hz, 3H).

(3S,5S)-3-Azidomethyl-5-methyl-decanoic acid tert-butyl ester 169

20 A procedure similar to that used to prepare compound **160** was used except (3S,5S)-5-methyl-3-(toluene-4-sulfonyloxymethyl)-decanoic acid tert-butyl ester **168** (4.155 g, 9.74 mmol) was used as a reactant to give **169** as an oil (2.77 g, 96%). MS (APCI) m/z 270 (M^+ -27, 30%, $-\text{N}_2$), 214 (M^+ -87, 100%, -tBu, $-\text{N}_2$).

(3S,5S)-3-Aminomethyl-5-methyl-decanoic acid tert-butyl ester 170


25 A procedure similar to that used to prepare compound **161** was used except that (3S,5S)-3-Azidomethyl-5-methyl-decanoic acid tert-butyl ester **169** (2.50 g, 8.405 mmol) was used as a reactant to give **170** as an oil (1.648 g, 72%). MS (APCI) m/z 272 (M^+ +1, 100%).

-101-

Example 14 (3S,5S)-3-Aminomethyl-5-methyl-decanoic acid

A procedure similar to that used for Example 15 was used except tert-butyl (3S,5S)-3-(aminomethyl)-5-methyldecanoate **170** (1.6 g, 6.00 mmol) was used as a reactant to give Example **16** as a white solid (72%). MS (APCI) m/z 272 ($M^+ + 1$, 100%). mp = 174-175 °C; ^1H NMR (400 MHz, CD_3OD) δ 2.91 (dd, $J = 12.9$, 3.91 Hz, 1H), 2.83 (dd, $J = 12.7$, 7.57 Hz, 1H), 2.43 (dd, $J = 15.6$, 3.17 Hz, 1H), 2.19 (dd, $J = 15.6$, 8.80 Hz, 1H), 2.08-2.04 (m, 1H), 1.53 (m, 1H), 1.38-1.27 (m, 7H), 1.78-1.03 (m, 2H), 0.90-0.86 (m, 6H), 0.66 (m, 6H); MS (APCI) m/z 216 ($M^+ + 1$, 100%), 214 (M^- , 100%); $[\alpha]_D = +21.4$ (c1 in MeOH).

Example 17: Synthesis of (3R,4R)-3-Aminomethyl-4,5-dimethyl-hexanoic acid



-102-

(S)-2-Benzyl-3-methyl-butan-1-ol 172Ref. *JACS* 1997;119:6510. Amide 171.**Large scale procedure for the synthesis of acetic acid (S)-2-benzyl-3-methyl-butyl ester 173 from 171**

5 A of n-butyl lithium (10 M in hexane, 100 mL, 1000 mmol, 3.9 equiv.) was added to a solution of diisopropylamine (108.9 g, 150.9 mL, 1.076 mol, 4.20 equiv.) in THF (600 mL), at -78°C. The resulting solution was stirred for 10 minutes and warmed to 0°C, and hold at the temperature for 10 minutes. Borane-ammonia complex (31.65 g, 1.025 mmol, and 4.0 equiv) was added in one
10 portion, and the suspension was stirred at 0°C for 15 minutes, and at 23°C for 15 minutes, and then cooled to 0°C. A solution of amide **171** (86 g, 256.41 mmol, 1 equiv.) in THF was added to the cold hydride via a cannula over 3 minutes. The reaction was stirred at 23°C for overnight, then cooled to 0°C. Excess hydride was quenched by the slow addition of 3N HCl (700 mL). The reaction mixture was
15 diluted with more aqueous HCl (3N, 200 mL), and brine and then extracted with ether (4 × 15 mL). The ether solution was concentrated to a small volume, and 200 mL 2N NaOH was added, and stirred at 23°C for 2.5 hours. More ether was added and the layers were separated. The aqueous layer was saturated with salt and extracted with ether (3 × 200 mL). The combined organic was washed with
20 brine and dried on sodium sulfate. The residue was flash chromatographed (Pet. ether-25% ether -TEA) to give alcohol **172**, 50 g. NMR (CDCl₃) δ 7.35-7.16 (m, 5H, C₆H₅), 3.55 (app. t, 2H, -CH₂OH), 2.71 (dd, 1H, ArCH₂CH-), 2.52 (dd, 1H, ArCH₂CH), 1.87 (m, 1H, CHCH(Me)), 1.67 (m, 1H, CH(Me)₂), 0.98 (d, 3H, CH₃) and 0.96 (d, 3H, CH₃).

25 A sample 3.3 g was saved for characterization and the rest was immediately acetylated (triethylamine 50 mL, DMAP 4.6 g, acetic acid anhydride 32 mL) overnight at room temperature. Work up followed by chromatography on silica gel eluted with pet ether and then 10% ether in pet ether gave 62 g of **173**. NMR (CDCl₃) δ 7.30-7.14 (m, 5H, C₆H₅), 3.98 (m, 2H, -CH₂OAc), 2.71 (dd,

-103-

1H, ArCH₂CH-), 2.51 (dd, 1H, ArCH₂CH), 1.99 (s, 3H, CH₃C=O), 1.82 (m, 1H, CHCH(Me) and CH(Me)₂), 0.97 (d, 3H, CH₃) and 0.95 (d, 3H, CH₃).

(S)-Acetoxymethyl-4-methyl-pentanoic acid 174 and (S)-4-Isopropyl-dihydro-furan-2-one 175

5 Acetate **173** (15 g, 68.18 mmol) was dissolved in CH₃CN (150 mL), carbon tetrachloride (150 mL) and HPLC grade water (300 mL) and stirred. Sodium periodate (262.50 g, 1220 mmol) was added followed by ruthenium chloride (650 mg, 3.136 mmol). After overnight stirring it was diluted with ether and water, and filtered through a pad of Celite. The organic portion was separated
10 and the aqueous phase was further extracted with ether. After drying on magnesium sulfate the solvent was evaporated. Potassium carbonate (42 g) was added to the residue and refluxed overnight in methanol (250 mL) and cooled to room temperature. After evaporation, water was added to dissolve the solid, and conc. HCl was added to bring the pH to 2. Chloroform was added and extracted
15 overnight. The organic phase was separated, and aqueous was further extracted with chloroform. The combined organic extracts were dried, evaporated, and the product was purified on a silica gel column and the compound was eluted with 20% ether in methylene chloride. Fractions were monitored by tlc, and spots were detected with I₂/KI solution. Fractions were combined to give 4.6 g of lactone
20 **175**. NMR (CDCl₃) δ 4.38 (dd, 1H, CH_aH_bO), 3.93 (app. t, 1H, CH_aH_bO), 2.54 (dd, 1H, CH_cH_d C=O), 2.23 (m, 2H, CHCH(Me) and CH_cH_d C=O), 1.60 (m, 1H, CH(Me)₂), 0.92 (d, 3H, CH₃) and 0.85 (d, 3H, CH₃).

(3R,4R)-3-Benzyl-4-isopropyl-dihydro-furan-2-one 176

25 Lithium bis(trimethylsilyl)amide (1.0 M solution in THF, 92 mL, 92 mmol) was added in 3-5 minutes to a solution of (S)-β-(2-propyl)-γ-butyrolactone **175** (11.68 g, 91.25 mmol) in dry THF 100 mL at -78°C under argon atmosphere. It was stirred for 1 h and a solution of benzyl iodide (21.87 g, 100.37 mmol) in dry THF was added rapidly. Stirring was continued for 1.5 hours and quenched at -78°C by the addition of a solution of brine followed by ethyl
30 acetate. The organic phase was separated and the aqueous was further extracted

-104-

with ether. Chromatography on silica gel first eluted with 5% methylene chloride in pet ether, and finally with 10% ether in pet ether gave desired compound 11.6 g, 58%. NMR (CDCl₃) δ 7.19 (m, 5H, C₆H₅), 4.02 (app. t, 1H, CH_aH_bO), 3.87 (dd, 1H, CH_aH_bO), 2.98 (d, 2H, ArCH₂), 2.57 (q, 1H, BnCHC=O), 2.05 (m, 1H, CHCH(Me)₂), 1.55 (m, 1H, CH(Me)₂), 0.81 (d, 3H, CH₃) and 0.72 (d, 3H, CH₃).

(2R,3R)-2-Benzyl-3-bromomethyl-4-methyl-pentanoic acid ethyl ester 177

Lactone **176** (6.5 g, 29.8 mmol) was dissolved in abs. ethanol (80 mL) and cooled in ice bath. Anhydrous HBr was bubbled through the solution for 1 hour and stirred at room temperature overnight while maintaining reaction under dry atmosphere. It was poured onto ice cooled mixture of pet ether and brine. The organic phase was separated, and the aqueous was further extracted with pet ether. The combined organic solution was washed repeatedly with cold water and dried. Solvent was removed *in vacuo* to give crude compound 7.0 g. NMR (CDCl₃) δ 7.27 (m, 5H, C₆H₅), 4.02 (m, 2H, CH₃CH₂O), 3.70 (dd, 1H, CH_aH_bBr), 3.55 (dd, 1H, CH_aH_bBr), 2.97 (m, 2H, ArCH₂), 2.83 (q, 1H, BnCHC=O), 2.11 (m, 1H, CHCH(Me)₂), 1.97 (m, 1H, CH(Me)₂), 1.10 (t, 3H, CH₃CH₂O), 0.96 (d, 3H, CH₃) and 0.93 (d, 3H, CH₃).

(2R,3R)-2-Benzyl-3,4-dimethyl-pentanoic acid ethyl ester 178

Bromoester **177** (7.25 g, about 80% pure), in ethanol (100 mL) containing triethylamine (3.2 mL) was hydrogenated overnight in the presence of 20% Pd/C (1.0 g). It was filtered through a pad of Celite, and the cake was washed with ethanol. Solvent was evaporated, and the residue was taken up in ether, whereupon solid (Et₃N.HCl) separated. The solid was removed by filtration. The filtrate was concentrated, and the procedure was repeated to eliminate all hydrochloride salt. Product was chromatographed on a silica gel column which was eluted with pet ether to give the desired debrominated compound 3.35 g. NMR (CDCl₃) δ 7.21 (m, 5H, C₆H₅), 3.95 (m, 2H, CH₃CH₂O), 2.85 (m, 2H, ArCH₂), 2.64 (q, 1H, BnCHC=O), 1.85 (m, 1H, CHCH(Me)₂), 1.62 (m, 1H,

-105-

$CH(Me)_2$), 1.05 (t, 3H, CH_3CH_2O), 0.95 (d, 3H, CH_3) 0.84 (d, 3H, CH_3) and 0.82 (d, 3H, CH_3). MS gave 290 ($M + CH_3CN$), 249 ($M + 1$), and others at 203. Further elution with ether gave lactone (2.25 g) that was carried over from previous step.

5 **Acetic acid (2R,3R)-2-benzyl-3,4-dimethyl-pentyl-ester 179**

Ethyl ester **178** (3.20 g, 12.85 mmol) was dissolved in anhydrous ether and cooled in ice bath under inert atmosphere. Lithium aluminum hydride (500 mg, 13.15 mmol) was added, and the suspension was stirred at room temperature overnight. Excess LAH was destroyed by careful addition of ethyl acetate while
10 the reaction was stirred in ice bath. Saturated sodium sulfate was added cautiously to coagulate the alumina that separated at room temperature as white precipitate. The reaction mixture was diluted with methylene chloride, and anhydrous sodium sulfate was added to dry the mixture. After filtration the solution was concentrated to give an oil 3.0 g.

15 The material (3.0 g) was dissolved in dichloromethane (30 mL) and triethylamine (2.5 mL), DMAP (200 mg), and acetic anhydride (1.5 mL) were added. It was stirred at room temperature for 3 hours, and diluted with ether. The ether solution was washed with water, 1N HCl, saturated sodium bicarbonate, brine and dried. The solution was concentrated in vacuo to give the acetoxy
20 compound **179** 3.16 g. NMR ($CDCl_3$) δ 7.19 (m, 5H, C_6H_5), 4.03 (m, 2H, CH_3CH_2O), 2.69 (m, 2H, $ArCH_2$), 2.09 (m, 1H, $BnCHCH_2O$), 2.02 (s, 3H, $CH_3C=O$), 1.68 (m, 1H, $CH_3CHCH(Me)_2$), 1.23 (m, 1H, $CH(Me)_2$), 0.87 (d, 3H, CH_3), 0.84 (d, 3H, CH_3) and 0.81 (d, 3H, CH_3).

(R)-4-((R)-1,2-Dimethyl-propyl)-dihydro-furan-2-one 180

25 To a solution of aromatic compound **179** (5.0 g, 20.16 mmol) in HPLC grade acetonitrile (60 mL), carbon tetrachloride (60 mL), and water (120 mL) was added sodium periodate (86.24 g, 403.32 mmol, 20 equiv.), followed by $RuCl_3$ (414 mg, 10 mol %). The mixture was stirred vigorously overnight at room temperature, and diluted with methylene chloride (400 mL). The mixture was
30 filtered through a pad of Celite to remove the solid precipitate. The organic

-106-

portion was separated, and the aqueous was further extracted with methylene chloride. After the combined organic portions concentrated, the residue was dissolved in ether and applied to a column of Florisil. The compound was eluted with 3% methanol in ether, evaporated to a paste that was dissolved in methanol (100 mL). Potassium carbonate (8.0 g) was added, and the mixture was refluxed for 6 hours. The solvent was evaporated, and the solid residue was dissolved in water. The pH was adjusted to 2 by the careful addition of concentrated HCl while being cooled in ice water bath and stirred. Chloroform (200 mL) was added to the solution and stirred as such overnight at room temperature. The organic phase was separated, and the aqueous portion was further extracted with chloroform. After drying, the solvent was evaporated to give the lactone **180** 5.0 g. NMR (CDCl₃) δ 4.36 (app. t, 1H, CH_aH_bO), 3.85 (app. t, 1H, CH_aH_bO), 2.46 (m, 2H, CH_cH_d C=O), 2.13 (m, 2H, CHCH₂C=O), 1.60 (m, 1H, CH(Me)₂), 1.35 (m, 1H, CH₃CHCH(Me)₂), 0.86 (d, 3H, CH₃) and 0.72 (t, 3H, CH₃).

(3R,4R)-3-Bromomethyl-4,5-dimethyl-hexanoic acid ethyl ester 181

Lactone **180** (5.0 g) was dissolved in absolute ethanol (25 mL) and flushed with argon. While being cooled in ice water bath, anhydrous HBr gas was bubbled through the mixture for 45 minutes and allowed to stand at room temperature overnight. The mixture was poured into ice-salt water and hexane. The organic phase was separated, and the aqueous was further extracted with hexane. The combined organic extract was dried and evaporated. Flash chromatography with 10% ether in pet ether on a silica gel column gave the bromoester **181** 3.54 g. NMR (CDCl₃) δ 4.14 (q, 2H, CH₃H₂O), 3.60 (dd, 1H, CH_aH_bBr), 3.41 (dd, 1H, CH_cH_d Br), 2.54 (dd, 1H, CH_aH_bC=O), 2.44 (dd, 1H, CH_aH_bC=O), 2.22 (m, 1H, O=CCH₂CHCH₂Br), 1.67 (m, 1H, CHCH₃CH(Me)₂), 1.37 (m, 1H, CH(Me)₂), 1.26 (t, 3H, CH₃CH₂O), 0.94 (d, 3H, CHCH₃CH(Me)₂), 0.81 (d, 3H, ((CH₃)₂)CHCH₃CH) and 0.79 (d, 3H, ((CH₃)₂)CHCH₃CH).

(3R,4R)-3-Azidomethyl-4,5-dimethyl-hexanoic acid ethyl ester 182 and


Example 17 (3R,4R)-3-Aminomethyl-4,5-dimethyl-hexanoic acid

-107-

Bromoester **181** (3.54 g, 13.34 mmol), sodium azide (1.04 g, 16.13 mmol) in anhydrous DMF (8.0 mL) was stirred at room temperature overnight. Water (16 mL) and hexane were added, the organic portion was separated, and the aqueous portion was further extracted with hexane. It was dried and evaporated to give azido ester 3.0 g. NMR (CDCl₃) δ 4.14 (q, 2H, CH₃H₂O), 3.48 (dd, 1H, CH_aH_bN₃), 3.21 (dd, 1H, CH_cH_b N₃), 2.34 (m 2H, CH_aH_bC=O), 2.20 (m, 1H, O=CCH₂CHCH₂ N₃), 1.60 (m, 1H, CHCH₃CH(Me)₂). Compound was submitted for hydrogenation (HPL, 66480 $\times$ 100). The hydrogenated crude was dissolved in 6N HCl and refluxed overnight. The solvent was evaporated in vacuo the residue was azeotroped with toluene. The crude was further purified by loading onto an ion exchange column chromatography (Dowex 50Wb $\times$ 8-100), washed to neutral eluent with HPLC grade water followed by elution of compound with 0.5N NH₄OH solution. Crystallization of product from methanol gave 720 mg. NMR (CD₃OD) δ 3.04 (dd, 1H, CH_aH_bNH₂), 2.82 (dd, 1H, CH_cH_b NH₂), 2.52 (dd, 1H, CH_aH_bC=O), 2.40 (dd, 1H, CH_cH_bC=O), 2.07 (m, 1H, O=CCH₂CHCH₂NH₂), 1.67 (m, 1H, CHCH₃CH(Me)₂), 1.35 (m, 1H, CH(Me)₂), 0.97 (d, 3H, CHCH₃CH(Me)₂), 0.88 (d, 3H, ((CH₃)₂)CHCH₃CH) and 0.83 (d, 3H, ((CH₃)₂)CHCH₃CH). [α]_D -5.3 (c, MeOH, 1.9 mg/mL). Anal. Calcd for C₉H₁₉NO₂: C 62.39, H 11.05, N 8.08. Found C 62.01, H 11.35, N 7.88. MS showed ions at 215 (M + CH₃CN), 197 (M + Na⁺), 174 (M + H⁺). Analysis of derivative by reverse phase HPLC, Hypersil BDS C₁₈ 5 micron and mobile phase 50/50 CH₃CN-water containing 0.1%TFA gave 99.93% purity at retention time of 8.21 minutes.

-108-

Examples 18-20: Synthesis of 3-Aminomethyl-4-isopropyl-heptanoic acid



2-Cyano-4-methyl-2-pentenoic acid methyl ester 61

5 A solution of isobutyraldehyde (30.0 g, 416 mmol), methyl-cyano-acetate (20.6 g, 208 mmol), ammonium hydroxide (3.2 g, 41.6 mmol) and acetic acid (5.0 g, 83.2 mmol) in 500 mL of toluene is warmed to reflux under a Dean-Stark

-109-

trap for 12 hours. The mixture is cooled to room temperature and extracted with saturated NaHSO_3 (3×100 mL), saturated NaHCO_3 (3×100 mL), and 100 mL of brine. The organic layer is dried over Na_2SO_4 , and the solvent is evaporated. The remaining oil is distilled under high vacuum (0.5 mm Hg, B.P. = 115-120°C) to give 28.8 g of 2-cyano-4-methyl-2-pentenoic acid methyl ester **61** as an oil (90% yield).

2-Cyano-3-isopropyl-hexanoic acid methyl ester 183

A 2.0 M solution of propyl magnesium chloride in Et_2O (9.8 mL, 19.6 mmol) is added to a solution of 2-cyano-4-methyl-2-pentenoic acid (3.0 g, 19.6 mmol) in 50 mL of THF which is cooled in an IPA/dry ice bath to -40°C under argon. The solution is stirred for 4 hours, and the reaction is quenched by addition of 50 mL of saturated KH_2PO_4 . The THF is evaporated, and the remaining oil is chromatographed under medium pressure over silica gel with 50% CH_2Cl_2 /hexane. Yield = 1.9 g (50%) of 2-cyano-3-isopropyl-hexanoic acid methyl ester as an oil.

2-Cyano-2-(1-isopropyl-butyl)-succinic acid 4-*tert*-butyl ester 1-methyl ester 184

A solution of 2-cyano-3-isopropyl-hexanoic acid methyl ester (1.9 g, 9.6 mmol) in 10 mL of THF is added to a slurry of NaH (washed with hexane, 0.23 g, 9.6 mmol) in 20 mL of THF which is cooled in an ice water bath under argon. The solution is stirred for 10 minutes, and *t*-butyl bromoacetate (2.1 g, 10.6 mmol) is added. The solution is warmed to room temperature. After 12 hours, the reaction is quenched by addition of 50 mL of saturated KH_2PO_4 and the THF is evaporated. The organic products are extracted into Et_2O (3×50 mL), and the combined organic layers are dried over MgSO_4 . The solvent is evaporated, and the remaining oil is chromatographed under medium pressure over silica gel in 25% hexane/ CH_2Cl_2 . Yield of 2-cyano-2-(1-isopropyl-butyl)-succinic acid 4-*tert*-butyl ester 1-methyl ester = 1.3 g (42%) as an oil.

-110-

3-Cyano-4-isopropyl-heptanoic acid t-butyl ester 185

A mixture of 2-cyano-2-(1-isopropyl-butyl)-succinic acid 4-*tert*-butyl ester 1-methyl ester (1.3 g, 4.2 mmol), NaCl (0.25 g, 4.2 mmol), and H₂O (0.15 g, 8.3 mmol) in 25 mL of DMSO is warmed to 130°C for 12 hours. The mixture is cooled to room temperature and diluted with 100 mL of brine. The organic products are extracted into Et₂O (3 × 50 mL). The organic layers are combined and washed with 50 mL of H₂O and 50 mL of brine. Drying over Na₂SO₄ and evaporation of the solvent gives 0.8 g (75% yield) of 3-cyano-4-isopropyl-heptanoic acid t-butyl ester as an oil.

4-(1-Isopropyl-butyl)-2-pyrrolidone 186

3-Cyano-4-isopropyl-heptanoic acid t-butyl ester (0.8 g, 3.2 mmol) is reduced under 50 psi of H₂ in MeOH containing TEA and Ra Ni. When the theoretical amount of H₂ is taken up, the catalyst is removed by filtration, and the solvent is evaporated to give 0.6 g (100% yield) of 4-(1-isopropyl-butyl)-2-pyrrolidone as an oil.

Example 18: 3-Aminomethyl-4-isopropyl-heptanoic acid

4-(1-Isopropyl-butyl)-2-pyrrolidone (0.6 g, 2.3 mmol) is warmed to reflux in 50 mL of 6.0 M HCl for 12 hours. The solution is cooled to room temperature and filtered through Celite. The filtrate is evaporated, and the solid remaining is recrystallized from MeOH/EtOAc. Yield 0.035 g (6% yield) of 3-aminomethyl-4-isopropyl-heptanoic acid as an HCl salt, mp 160-170°C. ¹H NMR (CD₃OD) δ 0.9 (m, 9H), 1.30 (m, 5H), 1.78 (m, 1H), 2.30 (m, 2H), 2.45 (m, 1H), 2.95 (m, 2H). MS (APCI, CH₃CN, H₂O) 201 (M⁺, 100%).


Example 19: 3-Aminomethyl-4-isopropyl-octanoic acid

Prepared according to the procedure of Example 18. Yield = 0.13 g (15%) of 3-aminomethyl-4-isopropyl-octanoic acid. mp = 160-170°C. ¹H NMR (CD₃OD) δ 0.9 (m, 9H), 1.30 (m, 7H), 1.78 (m, 1H), 2.30 (m, 1H), 2.45 (m, 2H), 2.95 (m, 2H). MS (APCI, CH₃CN, H₂O) 198 (M-17, 100%), 216 (M⁺, 50%).

-111-

Example 20: 3-Aminomethyl-4-isopropyl-hexanoic acid

Prepared according to the procedure of Example 18. Yield = 0.11 g (42%) of 3-aminomethyl-4-isopropyl-hexanoic acid. mp = 170-180°C. ¹H NMR (CD₃OD) δ 0.9 (m, 9H), 1.18 (m, 1H), 1.39 (m, 3H), 1.78 (m, 1H), 2.30 (m, 1H), 2.45 (m, 1H), 2.95 (m, 2H). MS (APCI, CH₃CN, H₂O) 188 (M⁺, 100%).

Example 21

(i) MeO₂CCH=PPh₃, THF, 40°C; (ii) MeNO₂, DBU; (iii) Raney Nickel, H₂, MeOH; (iv) Pd-C, MeOH, H₂; (v) 6N HCl

Synthesis of the unsaturated ester 188

(S)-(-)-citronellal **187** (2.0 mL, 11.03 mmol) was stirred at 40°C in dry tetrahydrofuran (30 mL) with methyl triphenylphosphoranylidene acetate (3.69 g, 11.03 mmol). After 8 hours the mixture was cooled to room temperature and stirred overnight. The solvent was removed in vacuo and the residue stirred with n-pentane (50 mL). After 1 hour the solid was removed by filtration and the solvent removed in vacuo to give an oil which was purified by flash chromatography (silica, ethyl acetate:heptane 1:9) to give 2.05 g (88%) of **188** as a clear oil. ¹H NMR (400 MHz) (CDCl₃) δ 0.90 (3H, d, J = 6 Hz); 1.12-1.40 (2H, m); 1.60 (3H, s); 1.62 (1H, m); 1.68 (3H, s); 2.01 (3H, m); 2.21 (1H, m);

-112-

3.73 (3H, s); 5.08 (1H, m); 5.82 (1H, d, $J = 16$ Hz); 6.94 (1H, m).
MS (CI⁺) (m/z): 211 (MH⁺, 75%), 179 (78%), 151 (100%).
IR (thin film) (cm⁻¹) v: 1271, 1436, 1728, 2917.

Synthesis of the nitroester 189

5 The ester **188** (2.02 g, 9.6 mmol) was dissolved in nitromethane (25 mL) with 1,8-diazabicyclo[5,4,0]undec-7-ene (1.44 mL, 9.6 mmol) and stirred at room temperature. After 23 hours the mixture was diluted with diethyl ether (150 mL) and washed with water (50 mL) and then 2N HCl (50 mL). The organic phase was collected, dried (MgSO₄), and the solvent removed in vacuo. The residue was
10 purified by flash chromatography (silica, ethyl acetate:heptane 3:7) to give 2.26 g (87%) of **189** as a clear oil. Note that this and all subsequent compounds are equimolar mixtures of 2 diastereoisomers. ¹H NMR (400 MHz) (CDCl₃) δ 0.90 (2 $\times$ 3H, each d, $J = 6$ Hz); 1.09-1.58 (10H, m); 1.602 (6H, s); 1.685 (6H, s); 1.94 (4H, m); 2.42 (4H, m); 2.66 (2H, m); 3.70 (6H, s); 4.42 (4H, m);
15 5.07 (2H, m).
MS (CI⁺) (m/z): 272 (MH⁺, 90%), 240 (100%), 151 (100%).
IR (thin film) (cm⁻¹) v: 1554, 1739, 2918.

Synthesis of the lactam 191

20 The nitro ester **189** (2.09 g, 7.7 mmol) was dissolved in methanol (75 mL) and shaken over Raney Nickel (catalytic, prewashed with water and then methanol) under an atmosphere of hydrogen gas (39 psi) at 35°C. After 17 hours the mixture was filtered through Celite. The solvent was removed in vacuo to give an oil. ¹H NMR showed there had been partial reduction of the double bond so this was carried on without further purification. A sample of this partial reduced
25 product (440 mg, 2.1 mmol) was dissolved in methanol (40 mL) and shaken over 5% Pd-C under an atmosphere of hydrogen gas. After 18 hours the catalyst was removed by filtration through Celite to obtain 442 mg (99% from partial reduced material) as a clear oil which did not need purification. Note that this and all subsequent compounds are equimolar mixtures of 2 diastereoisomers. ¹H NMR
30 (400 MHz) (CDCl₃) δ : 0.88 (18H, m); 1.04-1.58 (20H, m); 1.96 (2H, m);

-113-

2.40 (2H, m); 2.58 (2H, m); 2.98 (2H, m); (3.45 (2H, m), 5.82 (2H, br s).

MS (CI⁺) (m/z): 212 (MH⁺, 100%).

Synthesis of Example 21

5 The lactam **191** (428 mg, 2.0 mmol) was heated to reflux in 6N HCl (20 mL). After 5 hours the mixture was cooled to room temperature and washed with dichloromethane (2 × 10 mL). The aqueous phase was collected and the solvent removed in vacuo. The residue was dissolved in water (10 mL) and freeze-dried to give 382 mg (71%) of Example 34 as a white solid. Note that this compound is an equimolar mixture of 2 diastereoisomers. ¹H NMR (400 MHz) (d₆-DMSO) δ 0.82 (18H, m); 0.95-1.55 (20H, m); 2.05-2.45 (6H, m); 2.75 (4H, m); 7.98 (6H, br s).

10 MS (CI⁺) (m/z): 230 ([MH-HCl]⁺, 90%), 212 (100%).

Microanalysis: Calculated for C₁₃H₂₈NO₂Cl:

C 58.74; H 10.62; N 5.27.

15 Found: C 58.46; H 10.50; N 5.33.

To one skilled in the art, the use of (R)-(+)-citronellal would afford compounds of opposite C5-stereochemistry to Example 21.

The compounds of the present invention can be prepared and administered in a wide variety of oral and parenteral dosage forms. Thus, the compounds of the present invention can be administered by injection, that is, intravenously, 20 intramuscularly, intracutaneously, subcutaneously, intraduodenally, or intraperitoneally. Also, the compounds of the present invention can be administered by inhalation, for example, intranasally. Additionally, the compounds of the present invention can be administered transdermally. It will be 25 obvious to those skilled in the art that the following dosage forms may comprise as the active component, either a compound of formula **1** or a corresponding pharmaceutically acceptable salt of a compound of formula **1**.

For preparing pharmaceutical compositions from the compounds of the present invention, pharmaceutically acceptable carriers can be either solid or

-114-

liquid. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier can be one or more substances which may also act as diluents, flavoring agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material.

5 In powders, the carrier is a finely divided solid which is in a mixture with the finely divided active component.

 In tablets, the active component is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

10 The powders and tablets preferably contain from five or ten to about seventy percent of the active compound. Suitable carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the
15 formulation of the active compound with encapsulating material as a carrier providing a capsule in which the active component, with or without other carriers, is surrounded by a carrier, which is thus in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid dosage forms suitable for oral administration.

20 For preparing suppositories, a low melting wax, such as a mixture of fatty acid glycerides or cocoa butter, is first melted, and the active component is dispersed homogeneously therein, as by stirring. The molten homogenous mixture is then poured into convenient sized molds, allowed to cool, and thereby to solidify.

25 Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water propylene glycol solutions. For parenteral injection liquid preparations can be formulated in solution in aqueous polyethylene glycol solution.

 Aqueous solutions suitable for oral use can be prepared by dissolving the
30 active component in water and adding suitable colorants, flavors, stabilizing and thickening agents as desired.

 Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or

-115-

synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-known suspending agents.

Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for oral administration.

5 Such liquid forms include solutions, suspensions, and emulsions. These preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

10 The pharmaceutical preparation is preferably in unit dosage form. In such form the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the
15 appropriate number of any of these in packaged form.

The quantity of active component in a unit dose preparation may be varied or adjusted from 0.1 mg to 1 g according to the particular application and the potency of the active component. In medical use the drug may be administered three times daily as, for example, capsules of 100 or 300 mg. The composition
20 can, if desired, also contain other compatible therapeutic agents.

In therapeutic use, the compounds utilized in the pharmaceutical method of this invention are administered at the initial dosage of about 0.01 mg to about 100 mg/kg daily. A daily dose range of about 0.01 mg to about 100 mg/kg is preferred. The dosages, however, may be varied depending upon the requirements
25 of the patient, the severity of the condition being treated, and the compound being employed. Determination of the proper dosage for a particular situation is within the skill of the art. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached.
30 For convenience, the total daily dosage may be divided and administered in portions during the day, if desired.

-116-

Illustrative examples of compounds made in accordance with the present invention are tested as described in EXAMPLE 22.

EXAMPLE 22

5 **RAT MODEL OF FOOTPAD TACTILE ALLODYNIA FROM
PRIOR INJECTION OF ACID INTO THE GASTROCNEMIUS MUSCLE**

Patients with fibromyalgia syndrome (FMS) typically demonstrate widespread, chronic musculoskeletal pain, which is often accompanied by tactile allodynia (pain in response to a relatively light tactile stimulus that is normally not painful). A rat model of persistent mechanical allodynia has been developed that is consistent with the muscle tenderness found in these patients. Multiple injections of acidified saline into the gastrocnemius muscle in rats produce a long-lasting allodynia (conveniently measured at the footpad) that is thought to be centrally mediated (Sluka K, Kalra A, Moore S. Unilateral intramuscular injections of acidic saline produce a bilateral, long-lasting hyperalgesia. Muscle Nerve 2001;24:37-46; Sluka K, Rohlwing J, Bussey R, et al. Chronic muscle pain induced by repeated acid injection is reversed by spinally administered mu- and delta-, but not kappa-, opioid receptor agonists. J Pharmacol Exp Ther 2002;302:1146-50). This model was utilized to evaluate a compound of the invention for its ability to inhibit allodynia.

20 Allodynia is induced as described by Sluka, et al. (Sluka K, Kalra A, Moore S. Unilateral intramuscular injections of acidic saline produce a bilateral, long-lasting hyperalgesia. Muscle Nerve 2001;24:37-46) with minor modifications. On Day 0, male Sprague-Dawley rats (~200 g body weight) in their dark cycle are placed in suspended wire-bottom cages and allowed to acclimate for 0.5 hours. The baseline paw withdrawal threshold is determined on the right hind paw by Von Frey monofilament hairs (bending forces of 2.0, 3.6, 5.5, 8.5, 15.1, and 28.8 g) using the Dixon Up-Down method (Dixon W. Efficient analysis of experimental observations. Ann Rev Pharmacol Toxicol 1980;20:441-62). Von Frey hairs are applied to the plantar surface for up to 6 seconds, and a flinching of the paw during that time frame is considered a positive response. After assessment, the right gastrocnemius muscle is shaved, swabbed with

-117-

alcohol, and injected with 0.1 mL of 0.9% NaCl solution acidified to pH 4 with HCl. The injection is repeated on Day 5. Animals are manipulated with a dynamic plantar aesthesiometer (Ugo Basile, Comerio-Varese, Italy) on Days 6, 7, and 8 to facilitate induction of the allodynia. To screen the rats for the development of allodynia, the 15.1 g Von Frey hair is applied to the ipsilateral paw on Day 11. Positive responders from that test are included in the compound evaluation study. On Day 12 (the day of peak allodynia), animals are assigned into treatment groups, and then their ipsilateral paw withdrawal thresholds are determined to establish the allodynia (reduction in paw withdrawal threshold) compared to baseline values. Rats are then orally dosed with 10 mL/kg vehicle (0.5% hydroxypropyl-methylcellulose/0.2% Tween 80) or an indicated dose of a compound of the invention. Paw withdrawal thresholds are reassessed by Von Frey hairs in blinded fashion 2 hours after dosing for the dose-response study, and 2, 5, 8, and 24 hours after dosing in the time course experiment. The inhibition of allodynia is determined for each animal by dividing the increase in paw withdrawal threshold after treatment by the difference between baseline and pretreatment paw withdrawal values. This fraction is then converted to percent inhibition by multiplying by 100.

In this manner, one or more compounds described above is evaluated for its ability to dose-dependently attenuate allodynia, and minimum effective doses of are determined). To determine the time course of inhibition, allodynia is monitored at various time points after a minimum effective dose of a compound of the invention.


In this manner, it is evaluated whether administration of one or more compounds described above reduces tactile allodynia to the footpad caused by prior injection of acidic saline. The time period for sustained efficacy is also determined. The results allow evaluation of a compound of the invention for treating the allodynia associated with fibromyalgia syndrome.

-118-

CLAIMS

What is claimed is:

1. A method for treating a disorder in a mammal, including a human, comprising administering to said mammal a therapeutically effective amount of a compound of the formula 1



or a pharmaceutically acceptable salt thereof, wherein:

- 10 R^1 is hydrogen, straight or branched alkyl of from 1 to 6 carbon atoms or phenyl; and

- 15 R^2 is straight or branched alkyl of from 4 to 8 carbon atoms, straight or branched alkenyl of from 2 to 8 carbon atoms, cycloalkyl of from 3 to 7 carbon atoms, alkoxy of from 1 to 6 carbon atoms, -alkylcycloalkyl, -alkylalkoxy, -alkyl OH, -alkylphenyl, -alkylphenoxy, or -substituted phenyl; and wherein said disorder is selected from OCD, phobias, PTSD, restless legs syndrome, premenstrual dysphoric disorder, hot flashes, and fibromyalgia.

- 20 2. A method for treating a disorder in a mammal, including a human, comprising administering to said mammal a therapeutically effective amount of the compound (3S, 5R)-3-Aminomethyl-5-methyl-octanoic acid or a pharmaceutically acceptable salt thereof wherein said disorders is selected from OCD, phobias, PTSD, restless legs syndrome, premenstrual dysphoric disorder, hot flashes, and fibromyalgia.
- 25 3. The method according to claim 2 wherein said disorder is selected from restless legs syndrome, premenstrual dysphoric disorder, hot flashes, and fibromyalgia.


-119-

4. The method according to claim 2 wherein said disorder is fibromyalgia.

5. The method according to claim 1 or 2 wherein said disorder is selected from restless legs syndrome, premenstrual dysphoric disorder, hot flashes, and fibromyalgia.

6. The method according to claim 1 or 2 wherein said disorder is selected from OCD, PTSD, and phobia; and wherein said phobia is selected from agoraphobia and specific phobias.

7. A method for treating fibromyalgia and a concomitant disorder in a mammal, including a human, comprising administering to said mammal a therapeutically effective amount of a compound of the formula 1



1

or a pharmaceutically acceptable salt thereof, wherein:

R^1 is hydrogen, straight or branched alkyl of from 1 to 6 carbon atoms or phenyl; and

R^2 is straight or branched alkyl of from 4 to 8 carbon atoms, straight or branched alkenyl of from 2 to 8 carbon atoms, cycloalkyl of from 3 to 7 carbon atoms, alkoxy of from 1 to 6 carbon atoms, -alkylcycloalkyl, -alkylalkoxy, -alkyl OH, -alkylphenyl, -alkylphenoxy, or -substituted phenyl; and wherein said concomitant disorder is selected from migraine headaches, temporomandibular joint dysfunction, dysautonomia, endocrine dysfunction, dizziness, cold intolerance, chemical sensitivity, sicca symptoms, cognitive dysfunction, generalized anxiety disorder, premenstrual dysphoric disorder, irritable bowel syndrome, functional abdominal pain, neuropathic pain, somatoform disorders, OCD, phobias, and PTSD.

-120-

8. A method for treating fibromyalgia and a concomitant disorder in a mammal, including a human, comprising administering to said mammal a therapeutically effective amount of the compound (3S, 5R)-3-Aminomethyl-5-methyl-octanoic acid or a pharmaceutically acceptable salt thereof wherein said
5 concomitant disorder is selected from migraine headaches, temporomandibular joint dysfunction, dysautonomia, endocrine dysfunction, dizziness, cold intolerance, chemical sensitivity, sicca symptoms, cognitive dysfunction, generalized anxiety disorder, premenstrual dysphoric disorder, irritable bowel syndrome, functional abdominal pain, neuropathic pain, somatoform disorders,
10 OCD, phobias, and PTSD.

9. The method according to claim 7 or 8 wherein the compound administered is selected from

(3S,5R)-3-Aminomethyl-5-methyl-nonanoic acid,
15 (3S,5R)-3-Aminomethyl-5-methyl-decanoic acid,
(3S,5R)-3-Aminomethyl-5-methyl-undecanoic acid, and
(3S,5R)-3-Aminomethyl-5-methyl-dodecanoic acid;
or a pharmaceutically acceptable salt thereof.

20 10. The method according to claim 8 wherein said concomitant disorder is generalized anxiety disorder, premenstrual dysphoric disorder, a somatoform disorder, irritable bowel syndrome, functional abdominal pain, neuropathic pain, or migraine headache.

25 11. A method of increasing slow wave sleep in a human subject being treated with an active pharmaceutical agent that decreases slow wave sleep comprising administering to a human subject in need of such treatment:

(a) a compound of the formula 1, as defined in claim 1, or a pharmaceutically acceptable salt thereof; and

30 (b) a human growth hormone or human growth hormone secretagogue, or a pharmaceutically acceptable salt thereof;

wherein the amounts of the active agents "a" and "b" are chosen so as to render the combination effective in increasing slow wave sleep.

-121-

12. A method of increasing slow wave sleep in a human subject being treated with an active pharmaceutical agent that decreases slow wave sleep comprising administering to a human subject in need of such treatment:

(a) the compound (3S, 5R)-3-Aminomethyl-5-methyl-octanoic acid or a pharmaceutically acceptable salt thereof; and

(b) a human growth hormone or human growth hormone secretagogue, or a pharmaceutically acceptable salt thereof;

wherein the amounts of the active agents "a" and "b" are chosen so as to render the combination effective in increasing slow wave sleep.

13. A method of increasing slow wave sleep in a human subject comprising administering to a human subject in need of such treatment:

(a) a compound of the formula **1**, as defined in claim 1, or a pharmaceutically acceptable salt thereof; and

(b) a human growth hormone or human growth hormone secretagogue, or a pharmaceutically acceptable salt thereof;

wherein the amounts of the active agents "a" and "b" are chosen so as to render the combination effective in increasing slow wave sleep.

14. A method of increasing slow wave sleep in a human subject comprising administering to a human subject in need of such treatment:

(a) the compound (3S, 5R)-3-Aminomethyl-5-methyl-octanoic acid or a pharmaceutically acceptable salt thereof; and

(b) a human growth hormone or human growth hormone secretagogue, or a pharmaceutically acceptable salt thereof;

wherein the amounts of the active agents "a" and "b" are chosen so as to render the combination effective in increasing slow wave sleep.

15. The method of claim 1, 7, 11, or 13 where in said compound is selected from:

(3S,5R)-3-Aminomethyl-5-methyl-nonanoic acid,

(3S,5R)-3-Aminomethyl-5-methyl-decanoic acid,

-122-

(3S,5R)-3-Aminomethyl-5-methyl-undecanoic acid, and
(3S,5R)-3-Aminomethyl-5-methyl-dodecanoic acid;
or a pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

Intern: Application No
PCT/IB 03/05712

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/197 A61P1/00 A61P25/18 A61P29/00 A61P43/00
A61P5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data, EMBASE, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SELA I: "Pregabalin (Pfizer)" CURRENT OPINION IN INVESTIGATIONAL DRUGS, CURRENT DRUGS, LONDON, GB, vol. 2, no. 6, June 2001 (2001-06), pages 828-834, XP009020712 ISSN: 0967-8298 the whole document	1-15
Y	GUYSMER E K ET AL: "AN APPROACH TO MANAGING FIBROMYALGIA" MEDICINE TODAY, XX, JP, vol. 3, no. 12, 1 December 2002 (2002-12-01), pages 58-59, XP009027465 the whole document	1-15

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

° Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

22 March 2004

Date of mailing of the international search report

01/04/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Trifilieff-Riolo, S

INTERNATIONAL SEARCH REPORT

Intern: Application No

PCT/IB 03/05712

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	AMERINGEN VAN M ET AL: "DRUGS IN DEVELOPMENT FOR SOCIAL ANXIETY DISORDER: MORE TO SOCIAL ANXIETY THAN MEETS THE SSRI" EXPERT OPINION ON INVESTIGATIONAL DRUGS, ASHLEY PUBLICATIONS LTD., LONDON, GB, vol. 9, no. 10, 2000, pages 2215-2231, XP009027468 ISSN: 1354-3784 the whole document ---	1-15
Y	MELTZER ET AL.: "PREGABALIN ENHANCES NON-RAPID EYE MOVEMENT SLEEP AND EEG SLOW-WAVE ACTIVITY IN RATS" SOCIETY FOR NEUROSCIENCE ABSTRACTS, vol. 27, no. 2, 2001, page 2245 XP001180051 abstract -----	1-15

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IB 03/05712

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 1-15 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.