

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2016356037 B2**

(54) Title
Metconazole-containing pesticide composition for preventing and treating fusarium head blight and application thereof

(51) International Patent Classification(s)
A01N 43/653 (2006.01) **A01P 3/00** (2006.01)
A01N 43/16 (2006.01)

(21) Application No: **2016356037** (22) Date of Filing: **2016.05.11**

(87) WIPO No: **WO17/084249**

(30) Priority Data

(31) Number	(32) Date	(33) Country
201510801884.5	2015.11.19	CN

(43) Publication Date: **2017.05.26**

(44) Accepted Journal Date: **2018.06.14**

(71) Applicant(s)
Nanjing Agricultural University

(72) Inventor(s)
Zhou, Mingguo;Duan, Yabing;Wang, Jianxin

(74) Agent / Attorney
Shelston IP Pty Ltd., Level 21, 60 Margaret Street, Sydney, NSW, 2000, AU

(56) Related Art
CN 102057914 A

(12) 按照专利合作条约所公布的国际申请

(19) 世界知识产权组织
国际局



(43) 国际公布日
2017年5月26日 (26.05.2017)

WIPO | PCT

(10) 国际公布号
WO 2017/084249 A1

- (51) 国际专利分类号:
A01N 43/653 (2006.01) A01P 3/00 (2006.01)
A01N 43/16 (2006.01)
- (21) 国际申请号: PCT/CN2016/081655
- (22) 国际申请日: 2016年5月11日 (11.05.2016)
- (25) 申请语言: 中文
- (26) 公布语言: 中文
- (30) 优先权:
201510801884.5 2015年11月19日 (19.11.2015) CN
- (71) 申请人: 南京农业大学 (NANJING AGRICULTURAL UNIVERSITY) [CN/CN]; 中国江苏省南京市卫岗1号, Jiangsu 210095 (CN)。
- (72) 发明人: 周明国 (ZHOU, Mingguo); 中国江苏省南京市卫岗1号, Jiangsu 210095 (CN)。 段亚冰 (DUAN, Yabing); 中国江苏省南京市卫岗1号, Jiangsu 210095 (CN)。 王建新 (WANG, Jianxin); 中国江苏省南京市卫岗1号, Jiangsu 210095 (CN)。
- (74) 代理人: 南京苏高专利商标事务所 (普通合伙) (NANJING SUGAO PATENT AND TRADEMARK FIRM (ORDINARY PARTNERSHIP)); 中国江苏省南京市中山东路198号龙台国际大厦1912室, Jiangsu 210005 (CN)。
- (81) 指定国 (除另有指明, 要求每一种可提供的国家保护): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW。
- (84) 指定国 (除另有指明, 要求每一种可提供的地区保护): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), 欧亚 (AM, AZ, BY, KG, KZ, RU, TJ, TM), 欧洲 (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG)。

根据细则 4.17 的声明:

— 发明人资格(细则 4.17(iv))

本国际公布:


- 包括国际检索报告(条约第 21 条(3))。
- 包括经修改的权利要求(条约第 19 条(1))。

(54) Title: METCONAZOLE-CONTAINING PESTICIDE COMPOSITION FOR PREVENTING AND TREATING FUSARIUM HEAD BLIGHT AND APPLICATION THEREOF

(54) 发明名称: 一种用于防治麦类赤霉病的含叶菌唑农药组合物及其用途

(57) Abstract: The present invention relates to a pesticide composition for preventing and treating fusarium head blight comprising, for each mu of a farmland, 1 to 150 parts by weight of validamycin and 1 to 20 parts by weight of metconazole. The pesticide composition of the present invention has a significant synergistic effect on preventing and treating fusarium head blight, can significantly improve the head blight preventing and treating effect and the head blight preventing and controlling capability of mankind; the pesticide composition significantly reduces the pollution level of DON toxin in wheat, reduces the dosage, the costs, environment pollution and pesticide residues, retards the drug resistance of wheat pathogenic fungus to metconazole to prevent and treat fusarium head blight; moreover, the pesticide composition prolongs the duration of action of validamycin, reduces the application frequency, and can also treat powdery mildew, rust disease, leaf blight, and sheath blight of Triticeae crops, thus the use costs of the pesticide are reduced; and due to a great reduction in the dosage of a chemical pesticide, the pesticide composition is environment-friendly.

(57) 摘要: 本发明涉及一种用于防治麦类赤霉病的农药组合物, 每亩用含有 1~150 重量份井冈霉素与 1~20 重量份叶菌唑。本发明的农药组合物用于防治麦类赤霉病具有明显的增效作用, 可显著提高防治赤霉病的效果, 提高人类防控麦类作物赤霉病的能力; 显著降低谷物 DON 毒素的污染水平, 减少用药量、降低成本、减少环境污染和农药残留, 延缓小麦病原真菌对叶菌唑的抗药性, 防治麦类赤霉病; 延长井冈霉素的持效期、减少用药次数、兼治麦类白粉病、锈病、叶枯病和纹枯病, 减少农药使用的用工成本的优点, 大幅度降低了化学农药的使用剂量, 对环境友好。



WO 2017/084249 A1

20 Apr 2018

2016356037

METCONAZOLE-CONTAINING PESTICIDE COMPOSITION FOR PREVENTING AND CONTROLLING FUSARIUM HEAD BLIGHT AND APPLICATION THEREOF

5 **FIELD OF THE INVENTION**

The invention belongs to the technical field of pesticides. More specifically, the present invention relates to a pesticide composition for preventing and controlling fusarium head blight, and further relates to an agricultural fungicide for preventing and controlling fusarium head blight.

0 **BACKGROUND OF THE INVENTION**

Any discussion of the prior art throughout the specification should in no way be considered as an admission that such prior art is widely known or forms part of common general knowledge in the field.

5 Wheat and barley are the most important cultivated grain crops in the world as well as one of the main food crops in China. Grain crops are often confronted with the threat of fungal diseases such as head blight, powdery mildew, sheath blight, rust and leaf blight. These diseases may occur in worldwide grain growing regions, severely affecting the yield and quality of agricultural grain products. Especially, fusarium head blight caused by fusaria not only can result in devastating yield loss, but also can produce toxins, such as deoxynivalenol (DON) and
20 zearalenon in infected grains, consequently, the quality of food, such as fermented food and brewed liquors, is severely affected, and the health of the human being and animals are harmed.

The deoxynivalenol toxin includes DON, 3A DON and 15A DON toxins, which have acute and chronic toxicity to human being and animals. As the DON toxin has stable chemical properties and won't be decomposed when heated, the DON toxin widely exists in head blight
25 fungus-infected barley and wheat, processed food thereof and animal products fed with grains infected by head blight fungi. It has been known that the main symptoms of human and animal poisoning caused by the DON toxin include nausea, emesis, dizziness, drowsiness, headache, numb hands and feet, general weakness and decrease in immunity, the fluctuation of breath, pulse, body temperature and blood pressure, bleeding, miscarriage and even death

can be seen in severe cases, and the DON toxin can cause human Kaschin-Beck disease (KBD) and has obvious embryotoxicity and teratogenic and carcinogenic effects. Since the DON toxin has severe harm to human health. In the joint conference on food additives and

contaminants held in Geneva in October, 1973, the United Nations world food and agriculture organization (FAO) and the world health organization (WHO) decided to include fusarium toxins into the fifth topic among sixteen topics given priority in research, and many countries have also always considered the prevention and control of head blight as an important research topic.

Moreover, many international organizations and national health organizations have laid down corresponding laws and regulations to strictly limit DON toxin content in food. FAO provides that the content of DON in food must be less than 1mg/kg, and WHO provides that the content of the DON toxin in food and feed cannot respectively exceed the standards of 1mg/kg and 5mg/kg. The European Union food safety standard requires that the content of the DON toxin cannot exceed 0.75mg/kg in edible kernels and cannot exceed 0.5mg/kg in bread and other foods. The national standard established by the Ministry of Health in China provides that the content of the DON toxin in edible wheat flour and corn cannot exceed 1mg/kg.

Fusarium head blight is an ascomycete disease, including bud rot, seedling blight and ear rot/ear blight, caused by a variety of fusaria (*Fusarium* spp.). It includes nearly twenty varieties of pathogens, such as fusarium graminearum (*F. graminearum*), fusarium asiaticum (*F. asiaticum*), fusarium culmorum (*F. culmorum*), fusarium avenaceum (*F. avenaceum*), fusarium moniliforme (*F. moniliforme*) and fusarium nivale (*F. nivale*). Fusarium head blight in most of the regions in China is caused by the mixed population of fusarium graminearum (*F. graminearum*) and fusarium asiaticum (*F. asiaticum*), fusarium asiaticum is mainly in the southern wheat region, and fusarium graminearum is mainly in the northern wheat region.

As no grain crop varieties or gene resources which are immune or highly resistant to fusaria have been found as yet, using fungicides has become the only effective fusarium head blight prevention and control technology. For almost a century, people have been carrying out a great deal of work to research and develop fungicides for triticeae fungal diseases, and have developed inorganic fungicides, organosulfurous fungicides and aromatic hydrocarbon protective fungicides in succession, especially selective fungicides developed after 1960s, such as benzimidazole fungicides, ergosterol biosynthesis inhibitors, cytochrome b inhibitors and succinodehydrogenase inhibitors, so that the epidemic harm of a lot of important triticeae crop diseases, such as grain crop smut, powdery mildew, rust, leaf blight and eyespot, has been effectively controlled. However, as the deoxynivalenol (DON) toxin produced by fusaria

(*Fusarium* spp.) has the function of a pathogenic factor, the resistance to a variety of fungicides is enhanced, and as a result, the prevention and control effect of the majority of fungicides on fusarium head blight is not very ideal. Therefore, how to effectively control grain crop head blight is a current significant social demand in guaranteeing grain safety and food safety.

The DON toxin contamination level of diseased grain depends on the quantity of pathogenic fungi infecting the grain and the toxin biosynthesis capability of thalli. Aimed at the situation that the prevention and control effect of today's domestic and foreign main fungicides for preventing and controlling fusarium head blight is not ideal and that DON toxin contamination often exceeds edible safety standards, the present invention studies and discovers two types of β -tubulin (β_1 - and β_2 -) receptors for benzimidazole fungicides existing in fusaria and a genetic negative regulation mechanism for fungicide resistance and fungicide sensitivity for the first time, and discovers that the 240th amino acid phenylalanine composing the β_2 -tubulin receptor is a main cause of fungicide sensitivity decrease and that the point mutation of the 167th amino acid will cause carbendazim to lose the prevention and control effect and greatly enhance the DON toxin synthesis capability, mycelial colonization speed and pathogenicity of thalli, decreasing the prevention control effect of fungicides; in addition, the study also discovers an ergosterol biosynthesis inhibitor (EBI) receptor of fusaria, i.e. cytochrome P₄₅₀ monooxygenase, or fungicide sensitivity or fungicide resistance to EBI can be decreased by the overexpression or point mutation of encoding genes. Therefore, although some fungicides show very high activity in the lab, high dosages are needed when the fungicides are applied in fields, for example, 40g to 50g (600-750g a.i./hm²) of benzimidazole fungicide with above active ingredients or 15g to 20g (225-300g a.i./hm²) of EBI with above active ingredients need to be used per mu to obtain about 70 percent of fusarium head blight control effect.

As these selective fungicides with single acting mechanisms are used in a large amount for a long term, fungicide resistance will appear in pathogenic fungus populations in the nature, and the effect in use will decrease year by year. After nearly thirty years of fungicide resistance monitoring, the inventor discovers that fusaria which have developed fungicide resistance to benzimidazole fungicides, such as carbendazim, have formed a dominant population in Eastern China, and the commonly used benzimidazole fungicides, such as carbendazim and thiophanate, have almost lost the value in preventing and controlling

2016356037 20 Apr 2018

5 fusarium head blight. Furthermore, because the capability of fungicide-resistant pathogenic fungi in producing DON toxin is not less than five times the capability of sensitive strains, the harm of fusarium head blight and the risk of food safety in China are increasingly exacerbated. In order to reduce yield loss, farmers often double the use of fungicides, such as carbendazim, and as a result, fungicide resistance, pesticide residue, environment pollution and food safety problem are further exacerbated.

Therefore, the inventor finally accomplishes the present invention through a great deal of experimental research and analysis on the basis of summarizing the prior art.

SUMMARY OF THE INVENTION

0 [Technical Problems to Be Solved]

It is an object of the present invention to overcome or ameliorate at least one of the disadvantages of the prior art, or to provide a useful alternative.

The objective of a preferred embodiment of the present invention is to provide a pesticide composition for preventing and controlling fusarium head blight.

5 The other objective of a preferred embodiment of the present invention is to provide an agricultural fungicide for preventing and controlling fusarium head blight and decreasing grain DON toxin contamination.

[Technical Solution]

The present invention is implemented through the following technical solution.

20 According to a first aspect the present invention provides an application of a pesticide composition in preventing and controlling fusarium head blight, wherein the pesticide composition comprises jinggangmycin and metconazole according to a weight ratio of (1-150):(1-20).

25 According to a second aspect the present invention provides an application of a pesticide composition in inhibiting deoxynivalenol biosynthesis, wherein the pesticide composition comprises jinggangmycin and metconazole according to a weight ratio of (1-150):(1-20).

30 Unless the context clearly requires otherwise, throughout the description and the claims, the words “comprise”, “comprising”, and the like are to be construed in an inclusive sense as opposed to an exclusive or exhaustive sense; that is to say, in the sense of “including, but not limited to”.

2016356037 20 Apr 2018

The present invention relates to a pesticide composition for preventing and controlling fusarium head blight.

The pesticide composition comprises jinggangmycin and metconazole according to a weight ratio of (1-150):(1-20).

5 According to one preferred embodiment of the present invention, the pesticide composition comprises jinggangmycin and metconazole according to a weight ratio of (30-100):(3-18).

According to another preferred embodiment of the present invention, the pesticide composition comprises jinggangmycin and metconazole according to a weight ratio of (50-65):(6-16).

0 The present invention further relates to an agricultural fungicide for preventing and controlling fusarium head blight. The agricultural fungicide contains 0.1 to 90.0 percent by

weight of pesticide composition, and the balance is a carrier and/or an adjuvant acceptable in pesticides.

According to one preferred embodiment of the present invention, the agricultural fungicide contains 10 to 75.0 percent by weight of pesticide composition, and the balance is the carrier and/or the adjuvant acceptable in pesticides.

According to another preferred embodiment of the present invention, the agricultural fungicide contains 26 to 58.0 percent by weight of pesticide composition, and the balance is the carrier and/or the adjuvant acceptable in pesticides.

According to another preferred embodiment of the present invention, the agricultural fungicide is a suspending agent, an emulsion in water, a microemulsion, a wettable powder or a water-dispersible granule.

According to another preferred embodiment of the present invention, the carrier is one or more of water, attapulgite, kaolin and light calcium carbonate.

According to another preferred embodiment of the present invention, the adjuvant is chosen from ethanol, methanol, ethylene glycol, propylene glycol, NNO-1, NNO-7, xanthan gum, polyethylene glycol, glycerol, nekal, sodium dodecyl sulfate, sodium dodecyl benzene sulfonate, ammonium sulfate, alkylphenol ethoxylates, alkylphenol polyoxyethylene ether phosphate, polyoxyethylene fatty acid, benzoic acid, sodium lignin sulfonate, carboxymethylcellulose and polyvinyl alcohol.


According to another preferred embodiment of the present invention, the fusarium head blight is wheat scab and barley scab, including bud rot, seedling blight, ear rot or ear blight of triticeae crops caused by fusarium pathogenic fungi (*Fusarium* spp.) of carbendazim-resistant fusaria.

The present invention is described in more detail below.

The present invention relates to a pesticide composition for preventing and controlling fusarium head blight.


The pesticide composition comprises jinggangmycin and metconazole according to a weight ratio of (1-150):(1-20).

The chemical structural formula of metconazole used in the present invention is as follows:



Metconazole is a triazole ergosterol biosynthesis inhibitor, which comprises cis-isomers (1RS, 5SR) with high antifungal activity and trans-isomers with low activity, and metconazole used by the present invention is a mixture of cis-isomers and trans-isomers. Metconazole has high inhibitory activity for the growth of fusarium hyphae, and, mainly by inhibiting ergosterol biosynthesis, destroys the cell membrane permeability and membrane structure of fungi, strongly inhibiting the growth of the fungal hyphae. However, not only do metconazole acceptors Cyt.P₄₅₀ in different plant pathogenic fungus cells have structural diversity, but also a variety of regulatory and antagonistic physiological mechanisms exist, so that different fungi show different sensitivities. At present, this fungicide is mainly internationally used for preventing and controlling rust (*Puccinia* spp.), leaf blight or glume blight (*Septoria* spp.), net blotch (*Pyrenophora teres*) and scald (*Rhynchosporium secalis*) of grain crops, but only has a medium prevention and control effect on triticeae powdery mildew (*Erysiphe* spp.) and head blight (*Fusarium* spp.).

The chemical structural formula of jinggangmycin or validamycin used by the present invention is as follows:



Jinggangmycin is a secondary metabolite of actinomycetes, and contains six types of similar amino glucan glycoside derivatives A, B, C, D, E, F. A large number of domestic and foreign studies have shown that the component A is the main active component of jinggangmycin. Therefore, jinggangmycin in the present invention is jinggangmycin A.

Jinggangmycin shows interference specific to the development of hyphae at the tips of rhizoctonia fungi (*Rhizoctonia* spp.) among basidiomycetes in vitro. Therefore, since jinggangmycin was discovered in 1970s, jinggangmycin has always served as an agricultural antibiotic for specifically preventing and controlling plant diseases caused by rhizoctonia, or

has been processed into mixtures along with pesticides for preventing and controlling other plant pests to control rhizoctonia (*Rhizoctonia* spp.) diseases. These rhizoctonia diseases mainly include rice sheath blight, wheat sheath blight and sheath blight or damping-off of other crops. In recent years, it has also been discovered that jinggangmycin can also be used for preventing and controlling false smut caused by hard-to-culture imperfect fungi (*Ustilagoideae virens*).

As validamycin or jinggangmycin only shows specific antifungal activity on rhizoctonia in vitro and does not have antifungal activity on fusaria of head blight, people have always studied the selective mechanism of jinggangmycin with rhizoctonia as an object for decades, discovering that validamycin/jinggangmycin interferes with the inositol and trehalose metabolism of rhizoctonia, as a result, the structure of the cell wall is destroyed, rhizoctonia is prevented from infecting plants, and thereby jinggangmycin has a good protective effect. For decades, this antibiotic has always been limited to use for preventing and controlling rhizoctonia diseases of crops, especially rice sheath blight.

As everyone knows, fungicide-resistant pathogenic fungi can be easily screened out from specific fungicides with single effects, so the specific fungicides will lose application value shortly after being applied widely. The inventor has studied the risk of the fungicide resistance of rhizoctonia to selective jinggangmycin since 1980s, discovering that what is different from any other selective antibiotics and chemical fungicides with extremely high resistance risk is that jinggangmycin-resistant rhizoctonia cannot be screened out from jinggangmycin under both lab and field conditions. It can be inferred therefrom that besides interfering with known inositol and trehalose biosynthesis, jinggangmycin may also have other special disease-resistant effect mechanisms.

On the basis of studying the DON toxin biosynthesis pathway and regulatory mechanism of fusarium, the inventor carried out the screening of a large number of compounds for inhibiting DON toxin biosynthesis. In the screening process, it was surprisingly discovered that with certain treating dosage, jinggangmycin can strongly inhibit biochemical reaction in the early pathway of DON toxin biosynthesis of wheat scab fungi. Therefore, providing a technique for preventing and controlling fusarium head blight with a jinggangmycin and metconazole composition to reduce DON contamination has become an important objective of the present invention.

The inventor carries out the inhibition of the synthesis of the pathogenic factor DON of

5 fusaria by jinggangmycin/validamycin and the screening of synergistic formulas of mixtures of jinggangmycin/validamycin and other various fungicides for preventing and controlling fusarium head blight in fields for the first time in the world, and creatively discovered that the jinggangmycin and metconazole composition has the advantages of outstanding synergism, DON toxin contamination reduction and great metconazole dosage reduction in preventing and controlling fusarium head blight.

10 In the pesticide composition of the present invention, if the amount of jinggangmycin is within a range from 1 part by weight to 150 parts by weight while the amount of metconazole is less than 1 part by weight, then the prevention and control effect on head blight will be decreased; if the amount of metconazole is greater than 20 parts by weight, then the synergistic effect will be decreased, and the pesticide composition has an inhibitory effect on the growth of wheat. Therefore, it is reasonable that the amount of metconazole is 1 to 20 parts by weight.

15 Likewise, if the amount of metconazole is within a range from 1 part by weight to 20 parts by weight while the amount of jinggangmycin is less than 1 part by weight, then the synergistic effect of jinggangmycin in preventing and controlling head blight and decreasing the toxin contamination level in cooperation with metconazole will be lost; if the amount of jinggangmycin is greater than 150 parts by weight, then cost will be increased, and the synergistic effect of jinggangmycin in preventing and controlling head blight and reducing
20 toxin contamination in cooperation with metconazole will be decreased; and therefore, it is appropriate that the amount of jinggangmycin is 1 to 150 parts by weight.

Preferably, the pesticide composition comprises jinggangmycin and metconazole according to a weight ratio of (30-100):(3-18).

25 More preferably, the pesticide composition comprises jinggangmycin and metconazole according to a weight ratio of (50-65):(6-16).

Both jinggangmycin and metconazole which are used by the present invention are products which are sold in the market at present, such as 3%, 5%A, 20%, 40%A and 60%A jinggangmycin powders and aqueous solutions of Zhejiang Qianjiang Biochemical Co., Ltd. and metconazole produced by Jiangsu Gengyun Chemical Co., Ltd.

30 The present invention further relates to an agricultural fungicide preparation for preventing and controlling fusarium head blight. The agricultural fungicide contains 0.1 to

90.0 percent by weight of pesticide composition, and the balance is a carrier and/or an adjuvant acceptable in pesticides.

In the present invention, it should be understood that the pesticide carrier or adjuvant is an auxiliary substance which is used to improve the physical and chemical properties of the agent, enhance efficacy and facilitate transportation and storage when the pesticide preparation is processed or used, such as a wetting agent, an emulsifier, a dispersing agent, a binder, a stabilizer or a synergist. According to the present invention, any chemical substance which has these properties and does not have any adverse influence on the agricultural fungicide of the present invention can be used in the present invention, and they are all within the protection scope of the present invention.

The carrier used in the present invention, for example, is one or more of water, attapulgite, kaolin and light calcium carbonate. Besides, the present invention can also use other carriers which can meet the requirement of the present invention and are commonly used in the art.

For example, the adjuvant used by the present invention is chosen from ethanol, methanol, ethylene glycol, propylene glycol, NNO-1 (chemical name: naphthalene sulfonate formaldehyde condensate sodium salt), NNO-7 (chemical name: naphthalene sulfonate formaldehyde condensate calcium salt), xanthan gum, polyethylene glycol, glycerol, nekal, sodium dodecyl sulfate, sodium dodecyl benzene sulfonate, ammonium sulfate, alkylphenol ethoxylates, alkylphenol polyoxyethylene ether phosphate, polyoxyethylene fatty acid, benzoic acid, sodium lignin sulfonate, carboxymethylcellulose and polyvinyl alcohol. The adjuvant used by the present invention is a product which is sold widely in the market at present, such as dispersing agents sold with the product names "NNO-1" and "NNO-7" by Xinyi Feihuang Chemical Co., Ltd., butylnaphthalenesulfonic acid sodium salt emulsifying, dispersing and permeating agent sold with the product name "nekal" by Hubei Prosperity Galaxy Chemical Co.,Ltd., alkylphenol ethoxylates sold by Shanghai Honesty Fine Chemical Co., Ltd., alkylphenol polyoxyethylene ether phosphate sold by Guangzhou XiLu Chemical Co., Ltd., polyoxyethylene fatty acid sold by Shandong Liangshan Hongtai Epoxidized Soybean Oil Co., Ltd. and carboxymethylcellulose sold by Jiangxi Pingxiang Mashan Additives Factory. Besides, the present invention can also use other adjuvants which can meet the requirement of the present invention and are commonly used in the art.

In the present invention, in the agricultural fungicide, if the amount of the pesticide

composition is less than 0.1 percent, then the disease prevention effect will be decreased or the synergistic effect will be lost; if the amount of the pesticide composition is greater than 90.0 percent, then the synergistic effect will be lost or the disease prevention effect will be decreased ; and therefore, it is reasonable that the amount of the pesticide composition is 0.1 to 90.0 percent.

Preferably, the agricultural fungicide contains 10 to 75 percent by weight of pesticide composition, and the balance is the carrier and/or the adjuvant acceptable in pesticides.

More preferably, the agricultural fungicide contains 26 to 58 percent by weight of pesticide composition, and the balance is the carrier and/or the adjuvant acceptable in pesticides.

In practical application, according to methods known by those skilled in the art, the fungicidal composition of the present invention and the carrier and/or the adjuvant can be prepared into agent forms adopted commonly in agriculture, such as a suspending agent, an emulsion in water, a microemulsion, a wettable powder or a water-dispersible granule.

While the suspending agent is prepared, the applicable carrier or adjuvant, for example, can be a dispersing agent chosen from alkyl naphthalene sulfonate, polycarboxylate and lignosulfonate, a wetting agent chosen from alkylphenol polyoxyethylene ether phosphate, phenethyl phenol polyoxyethylene ether phosphate, alkyl sulfate, alkyl sulfonate and naphthalene sulfonate, a thickening agent chosen from xanthan gum, aluminium-magnesium silicate and bentonite, a preservative chosen from benzoic acid and sodium benzoate, an organosilicon defoaming agent and an antifreezing agent chosen from glycerol, urea, ethylene glycol and propylene glycol.

While the emulsion in water is prepared, the applicable carrier or adjuvant, for example, can be an emulsifier chosen from pesticide emulsifier **700**, pesticide emulsifier **2201**, **Span-60** and emulsifier **T-60**, a solvent chosen from xylene, methylbenzene and cyclohexanone, a stabilizer chosen from triphenyl phosphite and epichlorohydrin, an antifreezing agent chosen from ethylene glycol, propylene glycol, glycerol and urea, a thickening agent chosen from aluminium-magnesium silicate, bentonite and xanthan gum or a preservative chosen from benzoic acid and sodium benzoate.

While the microemulsion is prepared, the applicable carrier or adjuvant, for example, can be an emulsifier chosen from calcium dodecyl benzene sulfonate (pesticide emulsifier **500**),

2016356037 20 Apr 2018

pesticide emulsifier **700**, pesticide emulsifier **2201**, Span-**60**, Tween-**80** and **TX-10**, a cosolvent chosen from methanol, isopropanol, n-butyl alcohol and ethyl alcohol, a solvent chosen from dimethylbenzene, methylbenzene, cyclohexanone and N-methylpyrrolidone or a stabilizer chosen from triphenyl phosphite and epichlorohydrin.

5 While the wettable powder is prepared, the applicable carrier or adjuvant, for example, can be a dispersing agent chosen from polycarboxylate, lignosulfonate and alkyl naphthalene sulfonate, a wetting agent chosen from alkyl sulfonate, alkyl sulfate and naphthalene sulfonate or a filler chosen from light calcium carbonate, talcum powder, diatomite, kaolin and attapulgite.

0 While the water-dispersible granule is prepared, the applicable carrier or adjuvant, for example, can be a dispersing agent chosen from polycarboxylate, lignosulfonate and alkyl naphthalene sulfonate, a wetting agent chosen from alkyl sulfate, polyoxyethylene alcohol, alkyl sulfonate and naphthalene sulfonate, a disintegrating agent chosen from citric acid, ammonium sulfate, glucose, urea and sodium bicarbonate, a binder chosen from corn starch, 5 microcrystalline celluloses and diatomite or a filler chosen from light calcium carbonate, sepiolite, diatomite, kaolin, attapulgite and talcum powder.

His research results are respectively elaborated below.

I. Screening of Active Compounds for Inhibiting Fusarium Growth

10 The present invention adopts a conventional method of fungicide bioassay, jinggangmycin and metconazole are respectively prepared into 2mg/mL of mother solution by using sterile water and methanol, a control agent, raw carbendazim, is dissolved into 0.1M/L of hydrochloric acid solution, phenamacril is dissolved in methanol, and thereby 2mg/mL of mother solution is prepared. When potato dextrose agar media (PDA) are cooled to the temperature of about 45°C, jinggangmycin and metconazole are respectively added in to designed concentrations (see Table 25 1), and are then poured into culture dishes to produce plates treated by the different agents, the process is repeated each time after three dishes are treated, hyphal clumps of wild sensitive strains (sensitive strains for short) and carbendazim-resistant and phenamacril-resistant strains (fungicide-resistant strains for short) of common fusarium graminearum and fusarium asiaticum causing fusarium head blight are inoculated, and are cultured under the temperature 30 of 25°C for four days, the diameters of the fungal colonies are measured by a crossing method, effective medium dosages (ED₅₀ values) when different treatments inhibit the growth of the

2016356037 20 Apr 2018

pathogenic fungi by 50 percent are calculated, and antifungal activities are compared.

The experimental result indicates that no matter whether jinggangmycin is used alone or mixed with metconazole, jinggangmycin almost does not have inhibitory activity on the growth of the sensitive strains and fungicide-resistant strains of the two types of fusaria causing head blight in vitro, and only when the concentration of jinggangmycin is as high as 50µg/mL can jinggangmycin have 6.5 to 7.8 percent of inhibitory effect on the growth of the two types of fusaria. However, metconazole has a similar strong inhibitory effect on the hyphal growth of the sensitive strains and fungicide-resistant strains of the two types of fusaria, and the inhibitory effect of 0.125µg/mL of metconazole for treatment on hyphal growth can exceed 50 percent. Jinggangmycin does not have a synergistic effect for metconazole in inhibiting hyphal growth in vitro (Table 1).

According to growth inhibition rates of the wild sensitive strains and carbendazim-resistant and phenamacril-resistant strains of fusarium asiaticum and fusarium graminearum treated by different doses of metconazole, an effective medium dosage (ED₅₀) of metconazole for inhibiting the growth of the different fungicide-sensitive strains is calculated, a result shows that the sensitivities of the sensitive strains, the carbendazim-resistant strains and the phenamacril-resistant strains to metconazole are similar, ED₅₀ is 0.11 to 0.12µg/mL, and an experimental result is listed in Table 2.

With ED₅₀ of the agents as parameters, the activities of the different fungicides in inhibiting the growth of the fusaria are compared, and it is discovered that the activity of metconazole is about four times the activity (ED₅₀ is 0.45µg/mL for both types of fusaria) of carbendazim on the sensitive strains and about 1.5 times the activity (ED₅₀ is 0.165µg/mL for both types of fusaria) of phenamacril on the sensitive strains. The result indicates that metconazole has strong activity in inhibiting the growth of the wild sensitive strains and the carbendazim-resistant and phenamacril-resistant fusaria, helping to decrease the contamination level of DON toxin infecting grains and prevent and control fungicide-resistant diseases.

Table 1: Influence of Jinggangmycin and Metconazole on Growth of Sensitive Strains of Two types of Fusaria In Vitro

Strains		Average Fungal Colony Diameter (mm)		Growth Inhibition Rate (%)	
Agents	Concentration (µg/mL)	<i>F.g</i> *	<i>F.a</i> *	<i>F.g</i>	<i>F.a</i>
Jinggangmycin A	0.1	76.3	76.3	/	/
	1.0	76.2	76.3	/	/
	10.0	75.4	75.6	/	/
	50.0	70.3	69.4	6.5	7.8
Metconazole	0.0625	52.1	52.2	30.7	30.7
	0.125	31.2	31.2	58.5	58.6
	0.25	18.3	18.4	75.7	75.6
	0.5	8.2	8.1	89.1	89.2
	1.0	3.0	3.1	96.0	95.9
Jinggangmycin A + Metconazole	0.1+0.125	32.3	32.2	57.0	57.2
	1.0+0.125	32.2	32.3	57.2	57.1
	10.0+0.125	31.4	31.3	58.2	58.4
	50.0+0.125	30.1	30.1	60.0	60.0
	0.1+0.25	19.3	19.2	74.3	74.5
	1.0+0.25	18.7	18.6	75.1	75.3
	10.0+0.25	18.7	18.8	75.1	75.0
	50.0+0.25	16.0	15.6	78.7	79.3

Comparison	0	75.2	75.3	/	/
------------	----------	-------------	-------------	---	---

**F.g* and *F.a* are abbreviations of *Fusarium graminearum* and *Fusarium asiaticum*, similarly hereinafter.

Table 2: Growth Inhibition Activity of Metconazole on Sensitive Strains and Carbendazim-resistant

and Phenamacril-resistant Strains of Two Types of Fusaria

Strains	<i>F.g</i> sensitive strains	<i>F.g</i> carbendazi m- resistant strains	<i>F.g</i> phenamacr il-resistant strains	<i>F.a</i> sensitive strains	<i>F.a</i> carbendazi m- resistant strains	<i>F.a</i> phenamacril- resistant strains
ED ₅₀ (µg/mL)	0.117	0.113	0.115	0.114	0.112	0.116

II. Inhibitory Activity of Jिंगgangmycin on Toxin Biosynthesis Capability of Fusaria

As the fungicide sensitivities of fusarium graminearum and fusarium asiaticum to jिंगgangmycin and metconazole are the same, the inventor chooses fusarium asiaticum with higher toxin synthesis capability (the weight of synthesized DON per unit fungus amount, µgDON/g by dry weight of hyphae) as a material to further study toxin synthesis. Carbendazim-resistant fusarium asiaticum causing fusarium head blight is inoculated into 3 percent of sterile mung bean soup, the solution is shaken for culture under the temperature of 25°C and 12/24 hours of scattered light for ten days, and conidia are collected centrifugally. The conidia are inoculated into potato sucrose (PS) culture solutions containing different doses of jिंगgangmycin according to final 10⁶/mL and shaken to be cultured under the temperature of 25°C and 12/24 hours of scattered light, cultures are filtered after seven and fourteen days, toxin contents in the culture solutions are assayed respectively and the dry weights of hyphae are measured, and toxin synthesis capabilities (the amount of toxin produced per unit weight of hyphae) are analyzed.

2016356037 20 Apr 2018

Toxin Determination Method: The culture filtrate is equal-volume extracted with ethyl acetate for two times, the extracts are combined, and then depressurized, distilled and dried, the combined extract is dissolved by 1mL of acetonitrile and transferred into a new centrifuge tube, and after being distilled and dried again, the solution is stored under -20°C for later assay. During assay, 100µL of TMS derivatization reagent (TMSI:TMCS=100:1) is added, 1mL of ultrapure water is added after 10 minutes of uniform mixing, supernate is extracted and added into a GC sample bottle after shaking for layering, and a gas chromatograph (GC-ECD) with an electron capture detector is used to carry out toxin content assay. With a DON reagent of Sigma as a standard sample, a standard curve is created, and DON contents, including DON, 3ADON and 15ADON, in the culture solutions are calculated. Meanwhile, the filtered hyphae are dried to constant weights under 80°C, and the dry weights of the hyphae are measured. In addition, after seven days of shaking for culture, the hyphae are taken out, and the expression levels of the key gene *Tri5* for toxin synthesis are assayed.

It is discovered from an experimental result (Table 3) that the hyphal growths of the head blight fungi increase as culture time extends, but, when shaken to be cultured in the media treated by the different doses of jinggangmycin, hyphal growths are not notably changed in comparison with that of the blank control. It is indicated that jinggangmycin does not have the inhibitory effect on the growth of the head blight fungi cultured in liquid, which is the same as a linear growth rate determination result on the PDA plates. However, it is discovered for the first time that the amount (µg DON/g by dry weight of hyphae) of DON toxin synthesized per unit hyphal weight is remarkably decreased as the treating dosage of jinggangmycin is increased. Moreover, the inhibitory effect of jinggangmycin on DON synthesis decreases as culture time extends, and especially, the decreasing amplitude of low-concentration treatment is greater. It is indicated that jinggangmycin may be degraded as experimental time extends, and as a result, the inhibitory effect on toxin biosynthesis may be decreased.

According to toxin synthesis gene expression level analysis after seven days of treatment, it is creatively discovered that although jinggangmycin does not have adverse influence on the growth and hyphal morphology of fusaria in vitro, a very low treating dose of jinggangmycin can strongly inhibit the expression of the key gene *tri5* for DON toxin synthesis, weakening the toxin biosynthesis capability of thalli and reducing DON biosynthesis, and an experimental result is listed in Table 4.

Table 3: Effect of Jinggangmycin in Inhibiting DON Toxin Synthesis Capability of

Fusaria

Treating Dosage of Jinggaangmycin ($\mu\text{g/ml}$)	Toxin Production Capability of Hyphae ($\mu\text{gDON/g}$ by dry weight of hyphae)		DON Biosynthesis Inhibition Rate (%)	
	7 Days	14 Days	7 Days	14 Days
0	88.0	52.32	/	/
1	62.82	51.43	28.61	1.70
10	56.76	48.20	35.50	7.87
100	48.23	36.68	45.19	29.89
1000	30.35	22.52	65.51	56.96
10000	21.23	12.85	75.88	75.44

Table 4: Influence of Jinggaangmycin on Gene Expression of DON Synthesis Gene *Tri5* of Fusaria

Treating Dosage of Jinggaangmycin ($\mu\text{g/ml}$)	Relative Expression Level of <i>Tri5</i> Gene	Rate of Inhibition on Relative Expression Level of <i>Tri5</i> Gene (%)
0	1	/
1	0.41	59
10	0.34	66
100	0.16	84

1000	0.13	87
10000	0.10	90

III. Synergistic Effect of Metconazole for Jinglyngmycin in Inhibiting DON Toxin Biosynthesis Capability of Fusaria

Metconazole treatment can destroy the cell membrane permeability of the head blight fungi, inhibiting the growth of hyphae. Metconazole and jinglyngmycin composition treatment can enhance the absorption and utilization of jinglyngmycin by the pathogenic fungi. While the research content II determines the inhibition of jinglyngmycin on DON toxin biosynthesis of the fusaria, inhibitory effects of jinglyngmycin on the DON toxin biosynthesis capability of fusarium asiaticum under the existence of 0.1µg/mL of metconazole are determined after seven and fourteen days of culture, and the synergistic effect of metconazole for jinglyngmycin in inhibiting DON toxin biosynthesis is analyzed. The DON assay method is the same as that of the research content II.

It has been known from the research contents I and II that 0.1µg/ml of metconazole for treatment alone has an inhibition rate approximating 50 percent on the hyphal growth of the head blight fungi. It can be seen from the result in Table 5 that 0.1µg/ml of metconazole does not have significant inhibitory effects on the toxin biosynthesis capability of the fusaria in comparison with the toxin biosynthesis capability of the blank control after seven and fourteen days of treatment, which indicates that metconazole only has hyphal growth inhibition activity, and has no inhibitory effect on toxin synthesis capability. Nevertheless, when 0.1µg/ml of metconazole exists in each treatment concentration, the inhibitory effect of jinglyngmycin on DON toxin biosynthesis capability is greatly enhanced. Moreover, as culture time extends, the decreasing speed of the inhibitory effect of jinglyngmycin on toxin synthesis capability is remarkably lower than that of control treatment without metconazole, and in particular, the synergistic effect for low-concentration jinglyngmycin treatment and the prolonging of action time are more obvious. Based on the results of the simultaneous experiments in Table 3, synergistic effects for jinglyngmycin in inhibiting the toxin biosynthesis of the fusaria after seven and fourteen days of treatment under the existence of 0.10µg/mL of metconazole are calculated, and results are listed in Table 5. These results indicate: (1) jinglyngmycin has a strong effect in decreasing the DON toxin biosynthesis capability of the fusaria, while metconazole does not have this effect; (2) metconazole has an

obvious synergistic effect for jinggangmycin in inhibiting the toxin biosynthesis of the fusaria, and as the treating dosage of jinggangmycin is reduced, the synergistic effect is enhanced; (3) metconazole can prolong the time of the inhibitory effect of jinggangmycin on the toxin synthesis of the fusaria, and the synergistic effect is enhanced as the time of treatment extends.

Table 5: Synergistic Effect of 0.1µg/mL of Metconazole for Jinggangmycin in Inhibiting DON Toxin Synthesis of Wheat Strab Fungi

Treating Dosage of Jinggangmycin+Metconazole (µg/ml)	Toxin Production Capability (µgDON/g by dry weight of hyphae)		DON Biosynthesis Inhibition Rate (%)		Synergistic coefficient of 0.1µg/ml of Metconazole for Jinggangmycin in Inhibiting DON Synthesis *	
	7 Days	14 Days	7 Days	14 Days	7 Days	14 Days
0+0	88.00	52.32	/	/	/	/
0+0.1	87.42	52.50	0.66	-0.34	/	/
1+0.1	47.20	45.15	46.36	13.70	162.0	805.9
10 +0.1	26.06	27.21	70.39	47.99	198.3	609.8
100 +0.1	15.13	10.08	82.81	80.73	180.4	270.1
1000 +0.1	8.65	5.82	90.17	88.88	137.6	156.0
10000+0.1	5.85	3.12	93.35	94.04	123.0	124.7

* Synergistic Coefficient Calculation Method: When 0.1µg/ml of metconazole exists, an inhibition rate of jinggangmycin on toxin synthesis is divided by a toxin synthesis inhibition rate (Table 3) of a corresponding dose of jinggangmycin for treatment alone, and the result is multiplied by 100.

IV. Effect and Synergistic Effect of Jिंगgangmycin and Metconazole Composition Suspending Agent in Treating Seeds to Prevent and Control Head Blight and Other Diseases at Wheat Seedling Stage

According to weight ratios of jinggangmycin to metconazole which are 1:1.5, 10:1.5, 100:1.5, 1:1, 10:1, 100:1, 2:1, 20:1, 200:1, 5:1, 50:1 and 500:1, composition suspending agent samples, the contents of which are 3% (1:1.5, 1:1, 2:1), 5% (10:1.5, 10:1, 5:1), 10% (50:1, 100:1, 20:1) and 20% (100:1.5, 200:1, 500:1), are prepared respectively. In a sample preparation method, raw metconazole and raw jinggangmycin are mixed according to the above-mentioned weight ratios, added into a solution containing 40 percent by weight of water, 8 percent by weight of ethylene glycol and propylene glycol mixed solvent (volume ratio: 1:1), 1.0 percent by weight of NNO-1 and 1.5 percent by weight of NNO-7 dispersing agents (produced by Xinyi Feihuang Chemical Co., Ltd.), 0.5 percent by weight of xanthan gum binder and 1 percent by weight of polyethylene glycol and, finally, supplemented with water to 100 percent, and the granule diameter of agents milled to 90 percent by a sand mill is less than or equal to 5 μ m.

According to dosages of the single agents in a mixed agent for treatment per 100kg of seeds, single dosages of jinggangmycin and metconazole in corresponding dosages are designed (see Table 6), and before seeding, the samples are respectively mixed with the seeds according to 5L of liquor per 100kg of seeds to treat the wheat seeds infected by wheat scab fungi after being diluted by water. Twenty five seeds are sowed per pot with the diameter of 20cm, each treatment is repeated for ten pots, and the seeds are put into a greenhouse for culture. Ten days after emergence, the rate of emergence and the rate of dead seedlings are checked, and prevention and control effects on bud rot and seedling blight caused by the head blight fungi are calculated. Meanwhile, ten seedlings which are uniform in growth are kept, furthermore, a bran medium containing rhizoctonia cerealis is inoculated in substrate, powdery mildew fungus and leaf rust fungus spores are inoculated on leaves, one hundred wheat seedlings per treatment are investigated fourteen days after inoculation, and incidences of sheath blight, powdery mildew and rust and prevention and control effects are calculated. Meanwhile, according to prevention effects of the single agents on bud rot and seedling blight caused by the head blight fungi, theoretical prevention effect ($E=X+(100-X)Y/100$, wherein E is the theoretical prevention effect, X is the prevention effect of the single jinggangmycin agent, and Y is the prevention effect of the single metconazole agent) and synergistic

coefficient (actual prevention effect of the composition in application÷theoretical prevention effect×100) of the composition are calculated by the Abbott (1925) method, and results can be seen in Table 6.

5 Table 6: Effect and Synergistic Effect of Jingga mycin and Metconazole Composition Suspending Agent in Treating Seeds to Prevent and Control Wheat Diseases

Serial Number	Treating Dosage Per 100kg of Seeds (g ai. agent)		Incidence Prevention Effect %				Prevention and Control of Head Blight	
	Jingga mycin	Metconazole	Head Blight	Sheath Blight	Powdery Mildew	Rust	Theoretical Prevention Effect %	Synergistic Coefficient *
	10	15	100	98.5	90.5	98.5	60.5	165.3
	100	15	99.5	99.5	90.0	98.5	66.03	150.7
	1000	15	100	100	91.5	99.5	68.60	145.8
	10	10	95.0	97.0	82.5	94.5	58.5	162.4
	100	10	96.5	99.0	82.0	95.5	64.31	150.1
	1000	10	99.5	100	82.5	97.5	67.0	148.5
	10	5	87.5	80.5	74.0	86.0	46.0	175.0
	100	5	89.5	86.0	75.5	88.0	53.56	167.1
	1000	5	94.5	95.5	74.5	89.5	57.07	165.6
0	10	2	32.0	60.5	60.5	70.0	15.0	213.3

1	100	2	51.5	75.0	62.0	83.5	26.9	191.4
2	1000	2	58.5	81.5	64.0	83.0	32.43	180.4
3	10	0	0	10.5	0	0	/	/
4	100	0	14.0	26.0	0	0	/	/
5	1000	0	20.5	56.0	6.5	4.5	/	/
6	0	15	60.5	35.0	75.0	85.5	/	/
7	0	10	58.5	19.5	68.5	82.0	/	/
8	0	5	46.0	14.5	61.0	71.0	/	/
9	0	2	15.0	11.0	40.0	54.5	/	/
0	Blank Incidence %	Control	16.5	21.5	65.0	24.0	/	/

* Abbott believed that if the ratio of the actual prevention effect to theoretical prevention effect of a composition is greater than 1 (or a synergistic coefficient is greater than 100), then the composition has a synergistic effect (similarly hereinafter) (Abbott, 1925).

5 The above-mentioned experimental result indicates that when being mixed with the wheat seeds to treat the wheat seeds infected by head blight, the jinggangmycin and metconazole composition not only can effectively prevent and control bud rot and seedling blight caused

by the head blight fungi, but also has an excellent synergistic effect in preventing and controlling seedling blight caused by the head blight fungi within the ranges of the experimental dosages for treatment (10-1000g ai of jinggangmycin and 2-15g ai of metconazole/100kg of seeds), the synergistic coefficient is far greater than 100, and the synergistic effect is extremely outstanding. The single jinggangmycin agent has a certain prevention effect on sheath blight, but is almost ineffective to powdery mildew and rust; and except having a good prevention effect on rust at the seedling stage, the single metconazole agent only has a medium prevention effect on head blight, sheath blight and powdery mildew at the seedling stage, but can show an obvious synergistic effect in preventing and controlling the above-mentioned diseases when mixed with jinggangmycin.

V. Synergistic Effect and Toxin-reducing Effect of Jinggangmycin and Metconazole Composition Suspending Agent in Preventing and Controlling Wheat Scab

According to weight ratios of jinggangmycin to metconazole which are 1:50, 1:20, 1:3, 1:2, 1:1.5, 1:1, 2:1, 5:1, 5:1.5, 50:3, 10:1, 10:1.5, 50:2.5, 50:1.5 and 50:1, experimental composition suspending agent samples, the contents of which are 20% (1:50, 1:20, 1:3, 1:2, 1:1.5, 1:1, 2:1, 5:1, 5:1.5) and 40% (10:1, 10:1.5, 50:3, 50:2.5, 50:1.5, 50:1) by weight, are prepared respectively. In a sample preparation method, after raw metconazole and raw jinggangmycin are mixed according to the above-mentioned weight ratios, each mixture is added into a solution containing 5 percent by weight of ethylene glycol and propylene glycol mixed solvent (volume ratio: 1:1), 2 percent by weight of glycerol, 1.0 percent by weight of NNO-1, 1.5 percent by weight of NNO-7 dispersing agent, 3 percent by weight of sodium dodecyl sulfate and a small amount of water, finally, the solution is supplemented with water to 100 percent, and the granule diameter of agents milled to 90 percent by a sand mill is less than or equal to 5 μ m.

Different dosages of the jinggangmycin and metconazole composition used per mu are designed, and meanwhile, corresponding single-agent treatments are designed according to dosages of the single agents in each mixed agent in use. In April, 2013, experimental screening for preventing and controlling head blight in fields was carried out in Jiangsu White Horse Lake Farm, the wheat variety was Huaimai 22, each agent sample was mixed with water and sprayed for treatment at the initial stage of wheat blooming, five days later the agent sample is sprayed for the second time, and the amount of sprayed water per mu was 50kg. Each treatment was repeated for three plots, and the area of each plot was 50 square

2016356037 20 Apr 2018

5 meters. Meanwhile, 80g of carbendazim wettable powder which was 50 percent was used as a control agent for treatment per mu. According to a corresponding method specified by the industrial standard of Guidelines for the Field Efficacy Trials of Fungicides issued by the Ministry of Agriculture, the occurrence of wheat scab was investigated at the stage of milky ripeness, and according to the actual effect of each treatment in preventing and controlling head blight, the synergistic effects of the composition were calculated. According to the Abbott (ditto, 1925) method, theoretical prevention effect ($E=X+(100-X)Y/100$, wherein E is the theoretical prevention effect, X is the prevention effect of the single jinggangmycin agent, and Y is the prevention effect of the single metconazole agent) and synergistic coefficient (actual prevention effect of the composition in application÷theoretical prevention effect×100) of the composition were calculated.

5 Toxin Determination Method: Two hundred wheat ears were sampled every five treated points at the stage of wax ripeness, and were threshed indoors, and after drying, 30g of kernels were sampled randomly and ground. According to Goswami and Kistler methods, 5g of flour was put into a centrifuge tube, 20mL of extract of acetonitrile and water (84:16) was added into the flour and uniformly mixed by a vortex mixer, the solution was then shaken on a shaker for 24 hours, and was centrifuged at 5000rpm for 10 minutes, and 2mL of supernate was blow-dried by nitrogen in an Eppendorf centrifuge tube and preserved under -20°C. During assay, 100µL of TMS derivatization reagent (TMSI:TMCS=100:1) was added, 1mL of ultrapure water 10 was added after 10 minutes of uniform mixing, supernate was extracted and added into a GC sample bottle after shaking for layering, and a gas chromatograph (GC-ECD) with an electron capture detector was used to carry out toxin content assay.

25 Kernel-inflecting Fungus Amount Determination Method: 2g of flour for toxin assay was put into a 50mL centrifuge tube, added with 20 percent of CTAB pathogenic fungus DNA extraction buffer, then added with 50µL of protease K and 30µL of RNA enzyme and mixed sufficiently, the solution was centrifuged at 10000rpm for 10 minutes after being incubated under 65°C for 3 hours, 20mL of supernate was completely transferred into a 50mL centrifuge tube, then added with phenol, chloroform and isoamyl alcohol (25:24:1) of the same volume, shaken violently and centrifuged at 10000rpm for 5 minutes, 10mL of supernate was added with 30 3molL⁻¹ of cold sodium acetate accounting for one tenth of the volume and absolute ethyl alcohol two times the volume, precipitation was carried out under -20°C for 24 hours, centrifugation was carried out at 10000rpm for 10 minutes, precipitate was eluted two times by

20mL of 70 percent ethyl alcohol, the precipitate was dissolved into 500 μ L of TE solution after being dried, and the DNA template was stored under -20°C. Primers were designed according to the key gene *Tri5* for DON synthesis, DNA in a sample is amplified by adopting real time quantitative PCR, moreover, the *Tri5* DNA content (μ gDNA/g of wheat) in each gram of wheat sample was calculated, and the inhibitory effect of each treatment of the composition on the toxin synthesis capability of thalli was calculated. The result can be seen in Table 7.

Table 7: Synergistic Effect of Jinggangmycin and Metconazole Composition Suspending Agent in Preventing and Controlling Wheat Scab and Effect of Jinggangmycin and Metconazole Composition Suspending Agent in Reducing DON Toxin Contamination

Serial Number	Treating Dosage (g agent/mu)		Disease Prevention Effect %			DON Synthesis Capability (μ g of toxin/g of DNA)	Grain Toxin Content (μ g of toxin/g of kernels)	DON Contamination Reduction Rate %
	Jinggangmycin	Metconazole	Actual Prevention Effect	Theoretical Prevention Effect	Synergistic Coefficient			
1	0.5	25	98.0	85.2	115.0	21.40	0.26	97.5
2	1	20	93.3	75.7	123.2	20.11	0.34	96.8
3	5	5	80.3	48.42	154.4	15.29	0.82	92.3
4	5	10	90.1	58.34	157.9	14.89	0.28	97.4
5	5	15	96.2	73.71	130.5	14.85	0.21	98.0
6	10	3	66.0	41.30	159.8	8.80	1.51	85.8

7	10	5	80.6	50.55	159.4	8.28	0.43	95.9
8	10	10	94.2	60.06	156.8	7.93	0.32	97.0
9	10	15	98.1	74.80	131.1	8.26	0.27	97.5
10	50	3	75.2	49.65	151.5	6.01	1.10	89.6
11	50	5	88.5	57.59	153.7	5.92	0.37	96.5
12	50	10	94.3	65.75	143.4	5.73	0.20	98.1
13	50	15	98.2	78.39	125.3	5.84	0.18	98.3
14	100	3	80.8	53.69	150.5	3.76	0.65	93.9
15	100	5	93.7	60.99	1.536	3.35	0.21	98.0
16	100	10	94.3	68.5	137.7	3.55	0.23	97.8
17	100	15	97.2	80.13	121.3	3.28	0.20	98.1
18	150	3	87.9	59.31	148.2	3.03	0.31	97.1
19	0.5	/	0	/	/	21.52	9.4	11.3
20	1	/	0	/	/	20.15	8.9	16.0
21	5	/	10.6	/	/	15.05	8.6	18.9
22	10	/	14.3	/	/	8.46	6.6	37.7
23	50	/	26.5	/	/	5.89	4.7	55.7
24	100	/	32.4	/	/	3.42	4.5	57.5
25	150	/	40.6	/	/	2.88	4.0	62.3
26	/	3	31.5	/	/	20.16	5.8	45.3

27	/	5	42.3	/	/	19.80	4.9	53.8
28	/	10	53.4	/	/	19.96	4.2	60.4
29	/	15	70.6	/	/	19.54	3.6	66.0
30	/	20	75.7	/	/	19.20	1.9	82.1
31	/	25	85.2	/	/	19.38	0.42	96.0
32	Control Agent: Carbendazim 40g a.i.		67.5			28.94	4.4	58.5
33	Blank Control Disease Index		47.2	/	/	21.51	10.6	/

The field screening and experiment results of the synergistic effect of the composition for preventing and controlling fusarium head blight in the present invention indicate that when the jinggangmycin and metconazole composition is sprayed from the initial stage of wheat blooming to the filling stage and the mixed agent of 1g to 150g of active jinggangmycin ingredient and 2g to 20g of active metconazole ingredient is used per mu, not only does the jinggangmycin and metconazole composition have an outstanding synergistic effect (the synergistic coefficient is greater than 100) in preventing and controlling head blight, but also, after being applied, the jinggangmycin and metconazole composition has a remarkable synergistic effect in decreasing the DON contamination level of grains, the DON content can be reduced by 85 percent or above, and the DON contamination level can be controlled at a safe level not higher than 1mg of DON/kg of grain. Meanwhile, the result of quantitative pathogenic fungus analysis for the inflected kernels also indicates that jinggangmycin has the effect of inhibiting the toxin synthesis capability of thalli while metconazole does not have such an effect. Although carbendazim has 67.5 percent of prevention effect on head blight when used for treatment alone, the toxin contamination level of grains is still 4.4mg/kg of grains, and is only decreased by 58.5 percent in comparison with the blank control.

VI. Synergistic Effect and Toxin-reducing Effect of Jinggangmycin and Metconazole Composition Suspending Agent in Preventing and Controlling Wheat Scab

								Contro lling Head blight			
1	1	15	92.2	90.5	85.3	97.8	65.8	140	0.20	98.4	44.2
2	3	10	91.9	87.2	80.1	90.4	61.3	150	0.26	97.9	44.6
3	5	5	77.4	78.6	75.6	84.3	51.3	151	0.87	93.1	44.3
4	5	10	80.7	85.2	79.6	87.4	63.5	127	0.32	97.5	44.4
5	5	15	92.5	87.9	83.8	95.3	67.7	137	0.28	97.8	44.3
6	10	3	73.2	74.5	74.2	75.5	47.6	154	1.14	91.0	43.8
7	10	10	85.9	86.9	80.5	84.8	66.4	129	0.41	96.7	44.7
8	10	15	91.3	94.8	84.9	90.2	70.3	130	0.29	97.7	45.6
9	50	3	81.5	82.8	68.5	83.4	55.4	147	0.68	94.6	44.8
10	50	10	90.6	86.4	79.4	88.6	71.4	127	0.26	97.9	44.7
11	50	15	94.6	96.1	83.6	98.2	83.1	114	0.22	98.3	45.2
12	100	5	82.6	84.4	80.8	85.6	65.0	127	0.35	97.2	45.2
13	100	10	91.8	90.2	80.1	87.8	73.8	124	0.28	97.8	45.7
14	100	/	32.2	8.3	7.5	4.9	32.2	/	3.36	73.3	43.2
15	50	/	26.1	7.8	4.5	3.8	26.1	/	3.80	69.8	42.8
16	10	/	13.3	7.8	0	0	13.3	/	4.98	60.5	42.9
17	5	/	5.6	0	0	0	5.6	/	7.92	37.1	42.6

18	3	/	0	0	0	0	0	0	8.98	28.7	42.0
19	1	/	0	0	0	0	0	/	11.05	12.3	39.6
20	/	15	65.8	76.6	78.7	83.8	65.8	/	1.46	88.4	43.4
21	/	10	61.3	71.1	74.3	78.4	61.3	/	2.53	79.9	42.8
22	/	5	48.4	52.1	70.2	66.7	48.4	/	3.60	71.4	42.4
23	/	3	39.6	38.2	48.6	41.9	39.6	/	4.21	66.6	40.4
24	Carbendazim 40g a.i.		38.6	48.5	52.6	61.2	/	/	5.67	55.0	41.6
25	Control Disease Index		28.1	5.2	18.5	16.9	/	/	12.6	/	38.9

The field application results of the above-mentioned mixed agents indicate that when 1g to 100g of jinggangmycin as an active ingredient and 3g to 15g of metconazole as an active ingredient are used per mu to prevent and control wheat scab, the jinggangmycin and metconazole composition suspending agent not only has an excellent prevention and control effect on wheat scab, leaf blight, powdery mildew and rust, which is far better than that of commonly used carbendazim (the assay showed that pathogenic fungi which showed resistance to carbendazim in pathogenic fungus colonies in the experimental field accounted for 37.7 percent), moreover, the jinggangmycin and metconazole composition suspending agent has extremely remarkable synergistic and prevention and control effects on head blight caused by fungicide-resistant head blight fungi, and the result was similar to the field experiment result of White Horse Lake Farm in the same year. The difference between the effect of 3g to 50g of jinggangmycin and 3g to 15g of metconazole used per mu and the prevention effect of higher dosages is not great, DON toxin contamination can be controlled within a safe range, moreover, the dosage is less, the cost is low, the environment can be protected, and benefits are better. It is particularly worth noting that the thousand kernel weights of grains treated by all fungicides are obviously higher than that of the control, especially the jinggangmycin and metconazole composition for treatment increases thousand

kernel weight by not less than 13.9 percent, and the yield-increasing effect is more prominent.

VII. Synergistic Effect of Jिंगgangmycin and Metconazole Composition Wettable Powder in Preventing and Controlling Wheat Scab and Toxin Contamination-reducing Effect of Jिंगgangmycin and Metconazole Composition Wettable Powder

5 According to the good result of the jinggangmycin and metconazole composition
suspending agent in the field prevention and control of head blight and the treatment of other
diseases, the present invention designs weight proportions which can help to reduce the
dosage of the pesticide and cost, and prepares composition wettable powders with different
contents of jinggangmycin and metconazole. Preparation method: After mixing according to a
10 weight ratio of raw metconazole to raw jinggangmycin, the mixture is added with 30 percent
by weight of attapulgit carrier, 1.0 percent by weight of NNO-1 dispersing agent, 3 percent
by weight of nekal wetting agent and 2 percent by weight of sodium dodecyl benzene
sulfonate surfactant and, finally, added with light calcium carbonate as a filler to 100 percent,
and the mixture is milled by a sand mill and screened by a 300-mesh screen. Jिंगgangmycin
15 and metconazole composition wettable powder samples, the contents of which are 20% (1:3,
1:2, 1:1.5, 1:1) and 60% (5:1, 10:3, 10:1), are prepared respectively.

Meanwhile, according to the dosages of the single agents in the composition used per mu,
the single-agent treatments of jinggangmycin and metconazole and a common polyketone
wetable powder control agent are designed. In 2014, each composition wettable powder
20 sample was mixed with water and sprayed for treatment at the initial stage of wheat blooming,
the composition wettable powder sample was sprayed for the second time 5 days later, and the
amount of sprayed water per mu was 50kg. Each treatment was repeated three times, and the
area of each plot was 50 square meters. According to a corresponding method specified by
Guidelines for the Field Efficacy Trials of Fungicides, the occurrences of barley scab,
25 powdery mildew, rust and leaf blight were investigated at the stage of milky ripeness, and
disease indexes and prevention and control effects were calculated. Meanwhile, according to
the Abbott (1925) method of the research content V, the synergistic effects of the mixed
agents in preventing and controlling barley scab were calculated. One hundred and fifty wheat
ears were sampled randomly every five treated points at the stage of wax ripeness, taken and
30 threshed indoors and assayed according to the method of the research content V, and toxin
contents of the kernels were calculated. The treating dosages, prevention effects, synergistic
effects and influences of the agents on toxin content can be seen in Table 9.

Table 9: Effect of Jिंगgangmycin and Metconazole Composition Wettable Powder in Preventing and Controlling Barley Scab

Experimental Number	Treatment Dosage (g ai. agent/mu)		Disease Index Prevention Effect %				Prevention and Control of Head Blight		Grain Toxin Content (mg of DON/Kg of grain)	Toxin Reduction Percentage (%)
	Jिंगgangmycin	Metconazole	Head Blight	Leaf Blight	Powdery Mildew	Rust	Theoretical Prevention Effect	Synergistic Coefficient		
	3	10	80.4	81.2	80.5	85.1	61.6	132		
5	15	84.0	87.3	84.6	96.4	67.4	125	0.83	90.3	
5	10	83.1	86.2	80.8	88.0	63.3	131	0.85	90.0	
10	15	90.3	93.6	86.2	91.0	70.2	129	0.66	92.3	
5	5	78.0	80.4	78.0	81.3	49.2	159	0.91	89.3	
10	10	85.8	86.7	81.6	85.4	66.4	129	0.78	90.9	
50	10	88.6	85.4	82.4	86.5	71.7	124	0.49	94.3	
10	3	75.2	72.4	70.2	73.1	41.0	183	0.80	90.6	
100	10	88.6	93.2	82.1	90.8	74.1	120	0.46	94.6	
	3	/	0	0	0	0	/	/	6.67	21.8
0	5	/	4.5	0	0	0	/	/	5.28	38.1

1	10	/	12.6	0	0	0	/	/	4.62	45.8
2	50	/	26.4	0	0	0	/	/	3.86	54.7
3	100	/	32.5	6.9	3.4	0	/	/	3.49	59.1
4	/	3	32.5	56.2	52.6	63.0	/	/	3.86	54.7
5	/	5	46.8	70.4	65.3	68.2	/	/	3.70	56.6
6	/	10	61.6	73.7	71.8	76.1	/	/	3.09	63.8
7	/	15	65.9	80.0	82.2	88.2	/	/	3.26	61.8
8	Polyketone 30g a.i.		67.3	62.6	74.1	73.3	/	/	5.86	31.3
9	Control Disease Index		20.5	25.3	14.4	8.6	/	/	8.53	/

The field application of the composition indicates that the single-agent treatment of the single metconazole agent has a good prevention effect on barley scab, leaf blight, powdery mildew and rust while, except having a low prevention effect on head blight, the single jinggangmycin agent almost has no effect on other diseases. Nevertheless, when the composition containing 3g to 50g ai of jinggangmycin and 3g to 15g ai of metconazole is used, the prevention effect on head blight, leaf blight, powdery mildew and rust is greatly enhanced, the prevention and control effect reaches not less than 80 to 90 percent, better than the effect of the common control agent polyketone, and the synergistic effect is outstanding.

				t	Milde w		on Head Blight %		kernels)
1	3	15	90.6	93.6	90.1	100	66.7	136	0.24
2	5	10	89.2	87.4	80.3	97.6	64.3	139	0.23
3	5	5	84.2	78.3	74.2	87.7	54.1	156	0.45
4	10	5	88.8	83.6	76.6	89.1	57.2	155	0.38
5	50	3	85.2	75.4	72.8	74.3	51.3	166	0.40
6	100	3	90.2	85.7	71.8	87.5	54.4	166	0.31
7	100	/	31.4	8.5	7.3	5.1	31.4	/	2.89
8	50	/	26.7	7.9	4.4	3.6	26.7	/	3.24
9	10	/	12.8	0	4.5	0	12.8	/	3.86
10	5	/	6.5	0	0	0	6.5	/	4.20
11	3	/	0	0	0	0	0	/	4.77
12	/	15	66.7	76.8	73.7	82.6	66.7	/	2.01
13	/	10	61.8	71.7	70.5	77.4	61.8	/	2.24
14	/	5	50.9	53.4	63.5	69.2	40.9	/	2.38
15	/	3	33.6	32.4	33.6	40.4	33.6	/	2.94
16	Polyketone 57g a.i.		69.4	72.1	75.3	74.6	/	/	3.16
17	Blank	45.6	18.4	20.6	24.8	/	/	/	5.84

Control Disease Index									
-----------------------------	--	--	--	--	--	--	--	--	--

The above-mentioned research results indicate that when the single jinggangmycin agent and the single metconazole agent in are mixed in a tank on spot and then sprayed, the effect of the treatment using 3g to 100g of active jinggangmycin ingredient per mu and the effect of the treatment using 3g to 15g of active metconazole ingredient per mu in preventing and controlling head blight are equivalent to the effect of a prepared mixed agent applied on barley and wheat, moreover, the synergistic effect on wheat scab, powdery mildew, rust and leaf blight is also obvious, and the prevention and control effect is better than that of polyketone. In particular, the synergistic effect on wheat scab is prominent, the results are similar to the results of the other examples, which is reflected in that within the experimental dosage rang, as the dosage of metconazole is reduced, the synergistic effect is enhanced. 3g to 50g of active jinggangmycin ingredient used per mu and 3g to 15g of active metconazole ingredient used per mu can both achieve an ideal effect in preventing and controlling head blight, controlling leaf blight, powdery mildew and rust and reducing DON toxin contamination.

In a word, fusarium head blight described in the present invention is wheat scab and barley scab, including bud rot, seedling blight, ear rot or ear blight of triticeae crops caused by fusarium pathogenic fungi (*Fusarium* spp.) of carbendazim-resistant fusaria.

The above-mentioned research results indicate that the composition containing 1g to 150g of active jinggangmycin ingredient and 1g to 20g of active metconazole ingredient used per mu has a remarkable synergistic effect in treating (coating or mixing) wheat seeds to prevent and control fusarium head blight before sowing, and moreover, the composition can also control powdery mildew, rust and sheath blight of triticeae; and when mixed with water and sprayed at the stages of wheat heading, blooming and filling, the composition not only has a synergistic effect for the prevention and control of barley scab and wheat scab, but also has the effect of decreasing the DON toxin contamination level, increasing thousand kernel weight and controlling powdery mildew, rust and leaf blight.

[Beneficial Effect]

The beneficial effect of the present invention is that compared with conventional

fungicides, the fungicide composition of the present invention has the following advantages: (1) compared with the application of the single agents, the composition has an obvious synergistic effect when used for preventing and controlling fusarium head blight, and can remarkably enhance the effect of preventing and controlling head blight, increasing the ability of the human being in preventing and controlling head blight of triticeae crops; (2) the contamination level of DON toxin in grain can be remarkably decreased; (3) dosage, cost, environmental pollution and pesticide residue can be reduced; (4) the pressure of fungicide choice is decreased, the resistance of pathogenic fungi of wheat to metconazole can be retarded, and fusarium head blight which generates resistance to carbendazim can be prevented and controlled; (5) the persistent period of jinggangmycin is prolonged, the frequency of application is decreased, the powdery mildew, rust, leaf blight and sheath blight of triticeae can also be controlled, and the labor cost of pesticide application is reduced; (6) jinggangmycin is a pollution-free microbial pesticide with high selectivity, and by being mixed with metconazole, greatly reduces the dosage of the chemical pesticide, and the composition is environment-friendly.

DETAILED DESCRIPTION OF THE INVENTION

The present invention will be better understood through the following embodiments.

Embodiment 1: Experiment on 42 Percent of Jinggangmycin and Metconazole Composition Suspending Agent for Preventing and Controlling Wheat Diseases

Time and Place of Experiment: April, 2015, Jiangsu Xinyang Farm

Experimental Material: wheat, variety: Huaimai 33

Experimental Method:

Sample Preparation Method: Raw metconazole and raw jinggangmycin were mixed according to weight ratios of 1:20, 1:6, 5:16, 5:2 and 6:1, and according to the method of the research V described in the specification of the present application, 42 percent of experimental composition suspending agent samples with different ratios were obtained by processing. 25mL of composition preparation, 50mL of composition preparation and 100mL of composition preparation were used per mu. Each composition preparation was mixed with water and sprayed at the initial stage of wheat blooming, and was sprayed for the second time 5 days later, and the amount of sprayed water per mu was 50kg. Meanwhile, a corresponding dose of single wettable powder and spraying clean water were used as controls, carbendazim

served as a control agent, each treatment was repeated two times, and the area of each plot was 100 square meters.

According to a corresponding method specified by Guidelines for the Field Efficacy Trials of Fungicides, the occurrences of wheat scab, powdery mildew, rust and leaf blight were investigated at the stage of milky ripeness, disease indexes, prevention and control effects, synergistic effects on head blight and effects in controlling toxin contamination were calculated, and results are listed in Table 11.

Table 11: Effect of 42 Percent of Jinggangmycin and Metconazole Composition Suspending Agent in Preventing and Controlling Wheat Diseases

Serial Number	Treating Dosage (g ai. agent/mu)		Disease Index Prevention Effect %				Theoretical Prevention Effect of Mixed Agent on Head Blight %	Synergistic Coefficient	Grain Toxin Content (mg DON /Kg grain)	Rate of Decrease in Toxin Content in Grains (%)
	Jinggangmycin	Metconazole	Head Blight	Leaf Blight	Powdery Mildew	Rust				
1	1	20	89.8	87.4	92.6	100	75.1	120	0.28	95.7
2	0.5	10	88.4	85.1	82.3	85.4	70.1	126	0.36	94.4
3	3	18	90.1	84.6	87.2	91.5	74.9	120	0.30	95.4
4	1.5	9	85.6	80.3	80.5	88.4	67.2	127	0.41	93.7
5	5	16	92.3	81.0	85.6	90.3	74.6	124	0.26	96.0
6	2.5	8	88.2	79.6	78.7	85.8	65.1	135	0.44	93.2

7	15	6	89.1	80.2	73.4	82.8	65.8	135	0.43	93.3
8	7.5	3	64.2	65.6	64.4	75.9	40.4	162	0.78	87.9
9	36	6	88.3	83.6	74.3	88.2	69.1	128	0.48	92.6
10	18	3	73.8	73.5	56.9	72.6	49.0	151	0.62	90.4
11	36	/	31.6	5.8	8.5	7.6	/	/	2.61	59.6
12	18	/	28.9	5.0	7.8	7.2	/	/	2.86	55.7
13	15	/	24.3	4.6	7.6	7.2	/	/	2.94	54.5
14	7.5	/	16.9	0	0	0	/	/	3.42	47.1
15	5	/	5.2	0	0	0	/	/	3.86	40.2
16	2.5	/	0	0	0	0	/	/	3.92	39.3
17	3	/	0	0	0	0	/	/	3.96	38.7
18	1.5	/	0	0	0	0	/	/	4.82	25.4
19	1	/	0	0	0	0	/	/	5.80	10.2
20	0.5	/	0	0	0	0	/	/	5.86	9.3
21	/	20	75.1	80.5	80.1	91.2	/	/	1.11	82.8
22	/	18	74.9	80.2	74.6	88.4	/	/	0.99	84.7
23	/	16	73.2	79.6	71.4	85.6	/	/	1.08	83.3
24	/	10	70.1	75.3	63.4	82.1	/	/	1.34	79.3
25	/	9	67.2	75.0	60.8	82.4	/	/	1.56	75.9
26	/	8	65.1	74.3	58.6	79.4	/	/	1.61	75.1

27	/	6	54.8	70.2	51.8	72.6	/	/	1.89	70.7
28	/	3	28.3	28.1	26.4	30.2	/	/	3.22	50.2
29	Carbendazim 50g a.i.		49.04	/	/	/	/	/	4.35	32.7
30	Blank Control Disease Index		35.2	14.3	22.7	20.1	/	/	6.46	/

The above-mentioned field application results indicate that 0.5g to 36g of jinggangmycin and 3g to 20g of metconazole have a good wheat disease prevention and control effect when used. The synergistic coefficient for the prevention and control of head blight is 120 to 162, so the synergistic effect is prominent. However, when 3g to 15g of jinggangmycin and 3g to 16g of metconazole are applied per mu, the toxin contamination level is remarkably decreased, the synergistic effect is extremely obvious, the effect in preventing and controlling head blight is not less than 85 percent, and furthermore, powdery mildew, rust and leaf blight of wheat can also be effectively controlled. When 18g to 36g of jinggangmycin and 16g to 20g of metconazole are applied per mu, the disease prevention and control effect is not increased notably. These results also indicate that when 0.5g to 36g of jinggangmycin and 3g to 20g of metconazole are adopted per mu, the DON toxin content in grains is reduced by not less than 90 percent in comparison with that in the control, and in comparison with the toxin reduction effect of jinggangmycin or metconazole used for treatment alone, the toxin reduction effect is also extremely outstanding.

Embodiment 2: Effect of 64 Percent of Jinggangmycin and Metconazole Composition Wettable Powder in Preventing and Controlling Wheat Diseases

Time and Place of Experiment: April, 2015, Jiangsu Yanjiang Institute of Agricultural Sciences (Rugao)

Experimental Material: wheat cultivar: Yangmai 4

Experimental Method: Raw metconazole and raw jinggangmycin were mixed according to a weight ratio of 1:1, and 64 percent of experimental wettable powder sample was prepared according to the method of the research VII described in the specification of the present application. The composition wettable powder containing 4g to 20g of the active

jinggangmycin ingredient and the active metconazole ingredient was respectively used per mu. The composition wettable powder was mixed with water and sprayed on Yangmai 4 at the initial stage of wheat blooming, and was sprayed for the second time 5 days later, and the amount of sprayed water per mu was 45kg. A dose of 20 percent of single-agent wettable powder used in mixture and spraying clean water were used as controls, and 60g of 50 percent of polyketone wettable powder per mu was adopted as a control agent. Each treatment was repeated three times, and the area of each plot was 50 square meters. According to a corresponding method specified by Guidelines for the Field Efficacy Trials of Fungicides, the occurrences of wheat scab, powdery mildew, rust and leaf blight were investigated at the stage of milky ripeness, and disease indexes and prevention and control effects were calculated. The result can be seen in Table 12.

Table 12: Effect of 64 Percent of Jinggangmycin and Metconazole Composition Wettable Powder in Preventing and Controlling Wheat Disease

Serial Number	Preparation Dosage (g/mu)	Active Ingredient Dosage (g ai. agent/mu)		Disease Index Prevention Effect %				Theoretical Prevention Effect on Head Blight %	Synergistic Coefficient	Thousand Kernel Weight (g)
		Jinggangmycin	Metconazole	Head Blight	Leaf Blight	Powdery Mildew	Rust			
1	12.5	4	4	85.2	79.4	78.8	75.3	52.0	164	42.8
2	25	8	8	96.3	85.2	87.3	84.4	71.9	134	43.2
3	50	16	16	96.6	91.8	99.2	98.6	78.5	123	43.6
4	62.5	20	20	97.5	92.4	100	100	85.2	114	43.6

5	100	20	/	32.0	8.6	12.6	6.8	/	/	39.2
6	80	16	/	25.4	8.8	11.9	6.5	/	/	41.1
7	40	8	/	14.7	5.8	0	0	/	/	39.8
8	20	4	/	0	0	0	0	/	/	39.5
9	100	/	20	78.2	72.8	83.2	84.4	/	/	42.8
10	80	/	16	71.2	70.6	75.4	77.3	/	/	41.8
11	40	/	8	67.1	68.3	69.2	70.6	/	/	42.0
12	20	/	4	52.0	50.8	60.6	63.8	/	/	39.2
13	60	Polyketone 30		54.6	65.2	75.4	76.8	/	/	39.3
14	Blank Control Disease Index			35.6	14.8	25.1	20.6	/	/	37.2

The above-mentioned field application results indicate that the compositions in which the effective contents of jinggangmycin and metconazole according to the weight ratio of 1:1 are 4g to 20g each has a good prevention and control effect on wheat scab, leaf blight, powdery mildew and rust when used per mu, but the effect is better when 4g to 8g of active jinggangmycin ingredient and 4g to 8g of active metconazole ingredient are used. The composition for treatment can notably increase thousand kernel weight, and the yield-increasing effect is obvious.

Embodiment 3: Effect of 20 Percent of Jinggangmycin and Metconazole Composition Wettable Powder in Preventing and Controlling Wheat Diseases and Reducing Toxin Contamination

Time and Place of Experiment: April, 2015, Jiangsu Yanjiang Institute of Agricultural Sciences (Rugao), arranging the experiment in the same plots in embodiment 2.

Experimental Material: wheat cultivar: Yangmai 4

Experimental Method: After raw metconazole and raw jinggangmycin were mixed according to a weight ratio of 1:1, the mixture was added with 10 percent by weight of kaolin

1	30	3.0	3.0	76.2	70.4	68.8	73.3	39.1	195	41.2
2	75	7.5	7.5	93.8	83.3	80.4	83.8	75.1	125	42.6
3	100	10	10	95.4	86.6	90.1	94.2	83.5	114	44.1
5	50	10	/	32.4	8.6	12.6	6.8	/	/	39.6
6	37.5	7.5	/	24.2	8.8	11.9	6.5	/	/	39.0
7	15	3	/	0	5.8	0	0	/	/	37.9
8	50	/	10	75.6	72.8	83.2	84.4	/	/	43.2
9	37.5	/	7.5	67.1	70.6	75.4	77.3	/	/	42.6
10	15	/	3	39.1	68.3	69.2	70.6	/	/	39.8
11	60	Polyketone 30		54.6	65.2	75.4	76.8	/	/	39.3
12	Blank Control Disease Index			35.6	14.8	25.1	20.6	/	/	37.2

The above-mentioned experimental results indicate that the jinggangmycin and metconazole composition has an outstanding synergistic effect for the prevention and control of head blight and an outstanding yield-increasing effect. Such effects are related to a used dosage, but have no significant relationship with the content and adjuvant of the processed preparation.

Embodiment 4: Effect of 30 Percent of Jinggangmycin and Metconazole Composition Water-dispersible Granule in Preventing and Controlling Wheat Diseases and Reducing Toxin Contamination

Time and Place of Experiment: April, 2015, Jiangsu White Horse Lake Farm

Experimental Material: wheat cultivar: Huamai 4

Experimental Method: 30 percent of composition water-dispersible granule was obtained by processing according to a weight ratio of 5:1 of jinggangmycin to metconazole. A processing method was as follows: 39.1 parts by weight of 64 percent of jinggangmycin (25 parts of active ingredient), 5.2 parts by weight of 96.2 percent of raw metconazole (5 parts of

active ingredient), 30 parts by weight of ammonium sulfate, 17.2 parts by weight of light calcium carbonate, 2.5 parts by weight of alkylphenol ethoxylates, 2.5 parts by weight of sodium dodecyl benzene sulfonate and 3.5 parts by weight of polyoxyethylene fatty acid were ground, screened and pelletized.

5 200g, 100g, 50g and 30g of composition preparations were used per mu, mixed with water and sprayed at the initial stage of wheat blooming and sprayed for the second time 5 days later, and the amount of sprayed water per mu was 50kg. 100g of 20 percent of single wetttable powder per mu and spraying clean water were used as controls, each treatment was repeated three times, and the area of each plot was 50 square meters. According to a
10 corresponding method specified by Guidelines for the Field Efficacy Trials of Fungicides, the occurrences of wheat scab, powdery mildew, rust and leaf blight were investigated at the stage of milky ripeness, and disease indexes and prevention and control effects were calculated. Experimental results are listed in Table 14.

15 Table 14: Effect of 60 Percent of Jिंगgangmycin and Metconazole Composition Water-dispersible Granule in Preventing and Controlling Wheat Diseases

Serial Number	Preparation Dosage (g/mu)	Active Ingredient Dosage (g ai. agent/mu)		Disease Index Prevention Effect %			
		Jिंगgangmycin	Metconazole	Head Blight	Leaf Blight	Powdery Mildew	Rust
1	200	100	20	89.2	84.4	96.8	95.2
2	100	50	10	88.3	85.2	91.4	92.1
3	50	25	5	88.6	84.8	86.7.2	83.3
4	30	15	3	85.1	68.2	72.9	70.1
5	100	20	0	40.2	7.8	10.6	6.5

6	100	0	20	83.4	84.6	98.2	100
7	Blank Control Disease Index			25.9	23.4	29.7	24.0

The above-mentioned field application effect indicates that when sprayed according to 18g to 120g of active ingredients of the composition water-dispersible granule with 5 parts of jinggangmycin and 1 part of metconazole, the composition water-dispersible granule has a good prevention and control effect on wheat scab, powdery mildew, rust and leaf blight, and in particular, when 18g to 60g of active ingredients of the composition is sprayed per mu, the cost performance is best.

Embodiment 5: Effect of 32 Percent of Jinggangmycin and Metconazole Composition Emulsion in Water in Preventing and Controlling Wheat Diseases

Time and Place of Experiment: April, 2015, Jiangsu White Horse Lake Farm

Experimental Material: wheat cultivar: Huamai 4

Experimental Method: A preparation method for 32 percent of composition emulsion in water was as follows: raw metconazole was prepared into a suspending agent, and the suspending agent was added with an ethyleneoxide-propyleneoxide block polymer non-ionic emulsifier and stirred into the uniform oil phase; and in addition, a jinggangmycin aqueous solution, an ethylene glycol antifreezing agent and 1 percent of benzoic acid were mixed into the aqueous phase. Under high-speed stirring, the aqueous phase was gradually added into the oil phase, so that 32 percent of oil-in-water emulsion in water in which the ratio of jinggangmycin to metconazole was 1:3 was prepared.

100mL, 50mL, 25mL and 12.5mL of composition emulsions in water were used per mu, each composition emulsion in water was mixed with water and sprayed at the initial stage of wheat blooming and sprayed for the second time 5 days later, and the amount of sprayed water per mu was 50kg. 100g of 20 percent of single wettable powder per mu and spraying clean water were used as controls, each treatment was repeated three times, and the area of each plot was 50 square meters. According to a corresponding method specified by Guidelines for the Field Efficacy Trials of Fungicides, the occurrences of wheat scab, powdery mildew, rust and leaf blight were investigated at the stage of milky ripeness, and disease indexes and prevention and control effects were calculated. Experimental results are listed in Table 15.

Table 15: Effect of 32 Percent of Jिंगgangmycin and Metconazole Composition Emulsion in water in Preventing and Controlling Wheat Diseases

Serial Number	Preparation Dosage (mL/mu)	Active Ingredient Dosage (g ai. agent/mu)		Disease Index Prevention Effect %			
		Jिंगgangmycin	Metconazole	Head Blight	Leaf Blight	Powdery Mildew	Rust
1	100	8	24	94.6	88.4	99.8	97.2
2	50	4	12	91.3	87.5	95.6	93.7
3	25	2	6	82.6	80.8	81.7	80.2
4	12.5	1	3	60.2	61.4	62.3	62.2
5	100	20	0	34.2	7.8	10.6	6.5
6	100	0	20	82.4	80.6	78.2	85.0
7	Blank Control Disease Index			35.9	23.4	29.7	24.6

The above-mentioned field application results indicate that when 8g to 32g of active ingredients of the jिंगgangmycin and metconazole composition is used per mu, the prevention and control effect on wheat scab, powdery mildew, rust and leaf blight is good, and when 4g of active ingredients of the composition is used per mu, the effect in preventing and controlling these wheat diseases is also outstanding.

Embodiment 6: Effect of 45 Percent of Jिंगgangmycin and Metconazole Wettable Powder in Preventing and Controlling Wheat Diseases and Reducing Toxin Contamination

Time and Place of Experiment: April, 2015, Jiangsu Yanjiang Institute of Agricultural Sciences (Rugao)

Experimental Material: wheat cultivar: Yangmai 4

Experimental Method: 45 percent of experimental wettable powder sample in which the weight ratio of jinggangmycin to metconazole was 2:1 was prepared according to the method of the research VII described in the specification of the present application, and Jiangsu Yanjiang Institute of Agricultural Sciences was commissioned to carry out a field experiment. 5 50g, 40g, 30g and 20g of composition wettable powder preparations were used per mu respectively. Each composition wettable powder preparation was mixed with water and sprayed at the initial stage of wheat blooming, and was sprayed for the second time 5 days later, and the amount of sprayed water per mu was 50kg. 20 percent of single jinggangmycin and metconazole wettable powder was adopted, doses of single agents in the used mixed 10 agent and 40g of active carbendazim ingredient were adopted as control agents for treatment, spraying clean water was adopted as a blank control, each treatment is repeated three times, and the area of each plot was 50 square meters. According to a corresponding method specified by Guidelines for the Field Efficacy Trials of Fungicides, the occurrences of wheat scab and powdery mildew were investigated at the stage of milky ripeness, and disease 15 indexes and prevention and control effects were calculated.

During harvest, two hundred ears were sampled in total every five treated points, and after being threshed, the samples were sent to Nanjing Agricultural University for toxin content assay. The toxin assay method is the same as that mentioned above. Experimental results are listed in Table 16.

20

25 Table 16: Effect of 45 Percent of Jinggangmycin and Metconazole Composition Wettable Powder in Preventing and Controlling Wheat Diseases

Seri al Nu	Prepar ation Dosag	Active Ingredient Dosage	Disease Index Prevention Effect %	Synergistic Effect	Grain Toxin Content mg	Toxin Redu ction
------------------	--------------------------	--------------------------------	---	--------------------	------------------------------	------------------------

mb er	e (g/mu)	(g ai. agent/mu)								of DON/kg of grain	Perce ntage %
		Jing gan gmy cin	Metc onaz ole	Head Blight	Powd ery Milde w	Theor etical Preve ntion Effect on Head Blight	Theor etical Preve ntion Effect on Powd ery Milde w	Syne rgisti c Coef ficie nt on Head Blig ht	Synerg istic Coeffi cient on Powde ry Milde w		
1	50	15	7.5	94.6	90.8	70.8	70.8	134	128	0.22	95.1
2	40	12	6	89.3	85.6	57.2	63.2	156	135	0.34	92.5
3	30	9	4.5	82.6	84.7	54.3	51.1	152	166	0.66	85.4
4	20	6	3	70.2	66.3	33.9	46.1	207	144	1.28	71.7
5	75	15	/	34.2	8.6	/	/	/	/	2.86	36.7
6	60	12	/	10.2	6.6	/	/	/	/	3.82	15.5
7	45	9	/	9.4	3.0	/	/	/	/	4.02	11.1
8	30	6	/	7.4	1.8	/	/	/	/	4.14	8.4
9	37.5	/	7.5	55.6	68.8	/	/	/	/	2.01	55.5
10	30	/	6	52.3	60.6	/	/	/	/	2.17	52.0
11	22.5	/	4.5	49.6	58.6	/	/	/	/	2.06	54.4
12	15	/	3	28.6	45.1	/	/	/	/	3.12	31.0

13	80g of 50% of Carbendazim Wettable Powder	64.6	62.8	/	/	/	/	2.95	34.7
14	Blank Control Disease Index	21.8	15.2	/	/	/	/	4.52	/

It can be seen from the experimental results of the above-mentioned field application embodiments 1-6 that based on the discovery that jinggangmycin can inhibit the pathogenic factor DON toxin biosynthesis of the head blight fungi, the jinggangmycin and metconazole composition has an outstanding synergistic effect for the prevention and control of wheat scab when adopted to treat wheat seeds and be sprayed in fields. Moreover, the prevention and control effect on powdery mildew, rust, leaf blight and sheath blight of triticeae crops is also ideal. In addition, after the inventor adopted some of the combined jinggangmycin and metconazole compositions to carry out experimental demonstration in Jiangsu Yanjiang Institute of Agricultural Sciences, Jiangsu Xinyang Farm and White Horse Lake Farm for three years, the jinggangmycin and metconazole compositions all showed an excellent prevention effect on barley scab and wheat scab, and could also effectively control powdery mildew, leaf blight and rust. The present invention not only solves the major problem confronting China at present that the wheat scab fungi are hard to prevent and control due to carbendazim resistance, but also can overcome the food safety problem caused by wheat contaminated by DON toxin. Compared with conventional fungicides in the prior art, the jinggangmycin and metconazole composition of the present invention can greatly reduce the contamination of grains by the DON toxin of fusaria, and can also greatly reduce chemical pesticide dosage and application cost because of the synergistic effect, and thereby the jinggangmycin and metconazole composition has a good application prospect in reducing environmental pollution, controlling pesticide-resistant fungal diseases of triticeae and guaranteeing food safety.

CLAIMS

1. An application of a pesticide composition in preventing and controlling fusarium head blight, wherein the pesticide composition comprises jinggangmycin and metconazole according to a weight ratio of (1-150):(1-20).

2. The application according to claim 1, wherein fusarium head blight is wheat scab and barley scab, including bud rot, seedling blight, ear rot or ear blight of triticeae crops caused by fusarium pathogenic fungi (*Fusarium* spp.).

3. The application of claim 1 or claim 2 wherein the fusarium head bright is caused mainly by *Fusarium graminearum*.

4. An application of a pesticide composition in inhibiting deoxynivalenol biosynthesis, wherein the pesticide composition comprises jinggangmycin and metconazole according to a weight ratio of (1-150):(1-20).

5. The application according to any one of claims 1 to 4, wherein the pesticide composition comprises jinggangmycin and metconazole according to a weight ratio of (30-100):(3-18).

6. The application according to any one of claims 1 to 4, wherein the pesticide composition comprises jinggangmycin and metconazole according to a weight ratio of (50-65):(6-16).

7. The application according to any one of claims 1 to 4, wherein the pesticide composition comprises a carrier and/or an adjuvant acceptable in pesticides, and a mixture which is composed of jinggangmycin and metconazole according to a weight ratio of 1-150: 1-20 accounts for 0.1 to 90.0 percent of the total weight of the pesticide composition.

8. The application according to claim 7, wherein the mixture which is composed of jinggangmycin and metconazole according to a weight ratio of 1-150: 1-20 accounts for 10 to 75.0 percent of the total weight of the pesticide composition.

9. The application according to claim 7, wherein the mixture which is composed of jinggangmycin and metconazole according to a weight ratio of 1-150: 1-20 accounts for 26.0 to 58.0 percent of the total weight of the pesticide composition.

10. The application according to any one of claims 7-9, wherein the pesticide composition is a suspending agent, an emulsion in water, a microemulsion, a wettable powder or a water-dispersible granule.

11. The application according to any one of claims 7-9, wherein the carrier is one or more of water, attapulgite, kaolin and light calcium carbonate.

12. The application according to any one of claims 7-9, wherein the adjuvant is chosen from ethanol, methanol, ethylene glycol, propylene glycol, NNO-1, NNO-7, xanthan gum, polyethylene glycol, glycerol, nekal, sodium dodecyl sulfate, sodium dodecyl benzene sulfonate, ammonium sulfate, alkylphenol ethoxylates, alkylphenol polyoxyethylene ether phosphate, polyoxyethylene fatty acid, benzoic acid, sodium lignin sulfonate, carboxymethylcellulose and polyvinyl alcohol.

2016356037 20 Apr 2018