
(12) UK Patent Application (19) GB (11) 2 054 635 A

(21) Application No **8020493**

(22) Date of filing
23 Jun 1980

(30) Priority data

(31) **54/082185**

(32) **29 Jun 1979**

(33) **Japan (JP)**

(43) Application published
18 Feb 1981

(51) **INT CL³ C11D 1/29**
(C11D 1/29 1/38)

(52) Domestic classification
C5D 6A10 6A5B 6A5C
6A5D1 6A5D2 6A5E
6B10A 6B11C 6B12B1
6B12B2 6B12B3 6B12E
6B12F2 6B12G2A
6B12K2 6B12L 6B12N1
6B12NX 6C6 6C8

(56) Documents cited
GB 1500966

(58) Field of search
C5D

(71) Applicant
Kao Soap Co Ltd
1 Nihonbashi-Kayabacho
1 chome
Chuo-ku
Tokyo
Japan

(72) Inventors
Moriyasu Murata
Akira Suzuki

(74) Agents
Withers & Rogers
4 Dyer's Buildings
Holborn
London EC1N 2JT

(54) **Detergent composition**

(57) A detergent composition for cleaning and improving the feel of clothes comprising from 0.5 to 20% by weight of cationic surfactant for imparting a fabric softening effect to clothes and from 0.2 to 4 parts by weight, per part by weight of said cationic surfactant of an alkyl or alkenyl ether sulfate having oxypropylene and/or oxybutylene groups in the molecule.

U
K
P
A
T
E
N
T
O
F
F
I
C
E

SPECIFICATION

Detergent composition

5 The present invention relates to a detergent composition. More particularly, the present invention relates to a detergent composition which possesses an excellent recontamination-preventing effect. 5

Household treatments for cleaning and improving the feel of clothes have heretofore been carried out by a two-step process in which, in the first step, dirt and stains are removed by a detergent and, in the second step, the clothes are dipped in a clean water bath containing in solution a feel improving agent such as a fabric softening agent or an antistatic agent. Recently, washing and treating agent compositions that make possible simultaneous performance of the washing step and the feel improving step have been developed. From the viewpoint of shortening the washing time and from the economical viewpoint, the demand for such products has been increasing. However, because the commercially available products contain a cationic surfactant as the feel improving agent, for example, a soft finishing agent, there arises the problem that previously removed dirt and staining agents may become re-attached to the fabric in the washing bath, i.e. recontamination of the fabric may occur. 10 15

We have studied the prevention of recontamination using a detergent containing a cationic surfactant as the feel improving agent. As a result, we have discovered that if a cationic surfactant is used in combination with a specific anionic surfactant and a complex of the anionic surfactant and the cationic surfactant is formed, a high recontamination preventing effect can be obtained without reduction in the desired feel-improving effect. 20

More specifically, in accordance with the present invention, there is provided a detergent composition comprising a cationic surfactant and, as an anionic surfactant, an alkyl or alkenyl ether sulfuric acid ester salt containing a specific oxyalkylene group, which is represented by the following general formula (1): 25

$$R-O-A-SO_3M \quad (1)$$
 30

wherein R stands for a linear or branched, primary or secondary alkyl or alkenyl group having 8 to 22 carbon atoms, A stands for a polyoxyalkylene group containing oxypropylene and/or oxybutylene groups, in which the average number of the total polyoxyalkylene groups is from 0.5 to 30, and M stands for an alkali metal, an alkaline earth metal or an alkanolamine having 2 or 3 carbon atoms. 35

It is critical that the specific alkyl or alkenyl sulfuric acid ester salt having specific oxyalkylene groups, which is represented by the general formula (1), should be contained in an amount of 0.2 to 4 parts by weight, preferably 1 to 3 parts by weight, per one part by weight of the cationic surfactant. The cationic surfactant is incorporated in an amount of 0.5 to 20% by weight, preferably 1 to 5% by weight, based on the total weight of the detergent composition. 40

The oxyalkylene group (A) of the alkyl or alkenyl ether sulfuric acid ester salt of formula (1), includes oxypropylene and/or oxybutylene groups. The conventionally used oxyethylene groups can optionally also be included in the formula (1) material. The ratio of the number of moles of the oxypropylene groups and the oxybutylene groups and the ratio of the number of moles of the oxyethylene groups optionally included therein are not particularly critical. 45

Concerning the oxypropylene and oxybutylene groups used as the specific oxyalkylene groups in the oxyalkylene chain (A), oxypropylene groups are preferred, and combinations of oxyethylene and oxypropylene groups are especially preferred. The compounds of general formula (1) are industrially obtained in the form of a mixture in which the total number of oxyalkylene groups differs from molecule to molecule. The average number of the total oxyalkylene groups is 0.5 to 30, preferably 1 to 10. When oxyethylene groups are added, the ratio of oxyethylene groups to oxypropylene and/or oxybutylene groups is preferably in the range of from 1/4 to 4/1 and the average total oxyalkylene group number is preferably from 1 to 10. 50

The specific alkyl or alkenyl ether sulfuric acid ester salt containing specific oxyalkylene groups, which is represented by the general formula (1), is obtained by condensing at least one higher alcohol with at least one alkylene oxide, esterifying the condensate with sulfuric acid and neutralizing the ester with an alkali. Either an alkali catalyst or an acid catalyst can be used as the catalyst for the alkylene oxide condensation. Both natural higher alcohols and synthetic higher alcohols can be used as the starting higher alcohol. Primary and secondary synthetic higher alcohols can be used, but an oxo process synthetic higher alcohol (having an iso ratio of 20 to 80%) is most preferred, and a straight linear higher alcohol (higher alcohol derived from coconut oil or beef tallow) is second most preferred. More specifically, in the case of a liquid detergent composition, an oxo process synthetic higher alcohol having an average carbon number of 10 to 14 and an iso ratio of at least 30% is preferred. In the case of a powder detergent composition, a higher alcohol having an average carbon number of 12 to 18 is 65

preferred.

As preferred examples of the counter ion M in the general formula (1), there can be mentioned sodium, potassium, monoethanolamine, diethanolamine, triethanolamine and magnesium.

- 5 As preferred examples of the specific alkyl or alkenyl sulfuric acid ester salt having the specific oxyalkylene groups, which is represented by the general formula (1), there can be mentioned compounds represented by the following general formulae (2) to (7): 5

- 10 $R_1O(PO)_{m_1}SO_3M_1$ (2) 10
 10 $R_1O(PO)_{m_2}(EO)_{m_3}SO_3M_1$ (3) 10
 $R_1O(BO)_{m_2}(EO)_{m_3}SO_3M_1$ (4)
 $R_1O(PO,EO)_{m_1}SO_3M_1$ (5)
 $R_1O(BO)_{m_1}SO_3M_1$ (6) and
 15 $R_1O(PO,EO,BO)_{m_1}SO_3M_1$ (7) 15

- wherein R_1O stands for an alkyloxy and/or alkenyloxy residue of a coconut oil-derived higher alcohol, a beef tallow-derived higher alcohol, an oxo process synthetic higher alcohol (having an iso ratio of 20 to 80%) or a synthetic secondary higher alcohol, PO stands for an oxypropylene group, BO stands for an oxybutylene group, EO stands for an oxyethylene group, m_1 is a number of from 1 to 10, the sum of m_2 and m_3 is in the range of from 1 to 10, the ratio of m_2/m_3 is in the range of from 4/1 to 1/4, (PO,EO) and (PO,EO,BO) stand for randomly arranged cumulative oxyalkylene group assemblies, and M_1 stands for sodium, potassium, monoethanolamine, diethanolamine, triethanolamine or magnesium. 20

- The cationic surfactants that can be used in the present invention are not particularly critical. 25
 25 For example, the following cationic surfactants can be used in the present invention. 25
 (a) Di-(long-chain alkyl) quaternary ammonium salts represented by the following general formula:



- wherein R_2 and R_3 stand for an alkyl group having 10 to 26 carbon atoms, preferably 14 to 20 carbon atoms, R_4 and R_5 stand for an alkyl group having 1 to 5 carbon atoms, preferably 1 or 2 carbon atoms, and X stands for a halogen atom or a methyl sulfate or ethyl sulfate group (the same definitions will apply hereinafter). 35

- (b) Mono-(long-chain alkyl) quaternary ammonium salts represented by the following general formula: 40



- (c) Di-(long-chain alkyl) polyoxyethylene quaternary ammonium salts represented by the following general formula: 45




- wherein n is a number of 1 to 20, preferably 1 to 10 (the same definition will apply hereinafter), 55 or 55




- wherein m is a number of 1 to 20, preferably 1 to 10 (the same definition will apply hereinafter). 65
 65 (d) Mono-(long-chain alkyl) polyoxyethylene quaternary ammonium salts represented by the 65

following general formula:




(e) Bis-(hydroxyalkyl) quaternary ammonium salts represented by the following general formula:

10




(f) Quaternary ammonium salts having an amide or ester linkage, such as reaction products of compounds represented by the following general formula:

20



25 wherein p is a number of 1 to 5, preferably 2 or 3 (the same definition will apply hereinafter), with compounds represented by the following general formula:



and quaternary ammonium salts represented by the following general formula:



or



(g) Cationic polyamide compounds prepared by reacting 1 mole of diethylene triamine or dipropylene triamine with about 2 moles of a fatty acid having 12 to 24 carbon atoms to obtain a condensate having an acid value smaller than 10, adding about 1 to about 2 moles of epichlorohydrin to the thus-obtained condensate, subjecting the adduct to ring-opening polymerization in the presence of an alkaline agent and neutralizing the formed polymer with a monobasic acid in an amount of 0.3 to 1.5 moles per mole of said amine.


55 (h) Di-quaternary salts represented by the following general formula: 55



or




30 wherein q is a number of 2 to 8 and B stands for the group



35


or $-CH=CH-$ (the same definitions will apply hereinafter).

(i) Poly(N,N-dimethyl-3,5-methylenepiperidinium chlorides) having an average molecular weight of 1000 to 500000, which are represented by the following general formula:



wherein r stands for the total number of the monomer units.

(j) Products obtained by quaternizing a vinylpyrrolidonedimethylaminoethyl methacrylate copolymer with dimethyl sulfate, which have a molecular weight of 1000 to 500000 and are represented by the following general formula:




wherein s and t each stand for the total number of the monomer units.


(k) Products obtained by adding trimethylamine to an adduct of epichlorohydrin to hydroxyethyl cellulose to effect quaternization, which have a molecular weight of 200000 to 1000000 and

65


are represented by the following general formula:



(l) Aminoethyl acrylate phosphate/acrylate copolymers having a molecular weight of 1000 to 500000, which are represented by the following general formula:



(m) Products obtained by adding trimethylamine to an adduct of epichlorohydrin to starch to effect quaternization, which are represented by the following general formula:



Among the foregoing cationic compounds, quaternary ammonium salts (a), (b), (c) and (d) are preferred. More specifically, as preferred cationic compounds, there can be mentioned di-(beef tallow-alkyl)dimethyl ammonium chloride, mono-(beef tallow-alkyl)trimethyl ammonium chloride, di-(beef tallow-alkyl)dipolyoxyethylene ammonium chloride and mono-(beef tallow-alkyl)monomethyldipolyoxyethylene ammonium chloride.

A powdery or liquid detergent composition for clothes comprising the above-mentioned critical ingredients, according to the present invention, can additionally contain one or more of various anionic surface active agents and amphoteric surface active agents (1) to (9) described below, in amounts of 1 to 50% by weight, preferably 5 to 40% by weight. As the counter ion or the anionic surface active agents, there can be mentioned, for example, ions of alkali metals such as sodium and potassium, ions of alkaline earth metals such as calcium and magnesium, an ammonium ion, and alkanolamines containing 1 to 3 alkanol groups having 2 or 3 carbon atoms, such as monoethanolamine, diethanolamine, triethanolamine and triisopropanolamine.

(1) Linear or branched alkylbenzene sulfonate salts containing an alkyl group having 10 to 16 carbon atoms on the average.

(2) Alkyl- or alkenyl-ethoxy sulfate salts containing a linear or branched alkyl or alkenyl group having 10 to 20 carbon atoms on the average and having 0.5 to 8 moles of added ethylene oxide units on the average in one molecule.

(3) Alkyl or alkenyl sulfate salts containing an alkyl or alkenyl group having 10 to 20 carbon atoms on the average.


(4) Olefin sulfonate salts containing 10 to 20 carbon atoms on the average in one molecule.

(5) Alkane sulfonate salts containing 10 to 20 carbon atoms on the average in one molecule.

(6) Saturated or unsaturated fatty acid salts having 10 to 24 carbon atoms on the average in one molecule.

(7) Alkyl or alkenyl ether carboxylate salts containing an alkyl or alkenyl group having 10 to 20 carbon atoms on the average and having 0.5 to 8 moles of added ethylene oxide units, propylene oxide units, butylene oxide units, mixed ethylene oxide and propylene oxide units (molar ratio of 0.1/9.9 to 9.9/0.1) or mixed ethylene oxide and butylene oxide units (molar ratio of from 0.1/9.9 to 9.9/0.1) on the average in one molecule.

(8) α -Sulfo-fatty acid salts or esters represented by the following general formula:



wherein Y stands for an alkyl group having 1 to 3 alkyl groups or a counter ion as described above with respect to the anionic surface active agents, Z stands for a counter ion as described above with respect to the anionic surface active agents, and R₇ stands for an alkyl or alkenyl group having 10 to 20 carbon atoms.

(9) Amphoteric surface active agents represented by the following general formula:



wherein R_8 stands for an alkyl or alkenyl group having 10 to 20 carbon atoms, R_9 and R_{10} each stand for an alkyl group having 1 to 4 carbon atoms, p' is an integer of from 1 to 3, and X stands for a group $-COO^\ominus$ or $-SO_3^\ominus$.

In the present invention, a nonionic surface active agent can be incorporated as an optional ingredient. Ordinary nonionic surface active agents customarily used for detergent compositions can be used without any particular limitation. For example, the following nonionic surface active agents can be mentioned.

(A) Polyoxyethylene alkyl or alkenyl ethers containing an alkyl or alkenyl group having 10 to 20 carbon atoms on the average and having 1 to 20 moles of added ethylene oxide units.

(B) Polyoxyethylene alkylphenyl ethers containing an alkyl group having 6 to 12 carbon atoms on the average and having 1 to 20 moles of added ethylene oxide units.

(C) Polyoxypropylene alkyl or alkenyl ethers containing an alkyl or alkenyl group having 10 to 20 carbon atoms on the average and having 1 to 20 moles of added propylene oxide units.

(D) Polyoxybutylene alkyl or alkenyl ethers containing an alkyl or alkenyl group having 10 to 20 carbon atoms on the average and having 1 to 20 moles of added butylene oxide units.

(E) Nonionic surfactants containing an alkyl or alkenyl group having 10 to 20 carbon atoms on the average and 1 to 30 moles of added ethylene oxide and propylene oxide units or added ethylene oxide and butylene oxide units (the molar ratio of the ethylene oxide units to the propylene oxide or butylene oxide units is in the range of from 0.1/9.9 to 9.9/0.1).

(F) Higher fatty acid alkanolamides and alkylene oxide adducts thereof, which are represented by the following general formula:



wherein R_{11} stands for an alkyl or alkenyl group having 10 to 20 carbon atoms, R_{12} stands for H or CH_3 , n' is an integer of from 1 to 3, and m' is an integer of from 0 to 3.

(G) Sucrose fatty acid esters consisting of sucrose and a fatty acid having 10 to 20 carbon atoms on the average.

(H) Fatty acid glycerin monoesters consisting of glycerin and a fatty acid having 10 to 20 carbon atoms on the average.

(I) Alkylamine oxides represented by the following general formula:



wherein R_{13} stands for an alkyl or alkenyl group having 10 to 20 carbon atoms, and R_{14} and R_{15} each stand for an alkyl group having 1 to 3 carbon atoms.

In addition to the above-mentioned critical ingredients, the detergent composition of the present invention may further comprise up to 50% by weight of one or more alkali metal salts as builder salts. As such builder salts, there can be mentioned, for example, condensed phosphoric acid salts such as tripolyphosphoric acid salts, pyrophosphoric acid salts and metaphosphoric acid salts, aminopolyacetic acid salts, nitrilotriacetic acid salts, ethylenediamine-tetraacetic acid salts and diethylene-triamine-pentaacetic acid salts, hydroxycarboxylic acid salts such as citric acid salts, malic acid salts and glycollic acid salts, and polymeric electrolytes such as salts of alkali hydrolysis products of polyacrylic acid and vinyl acetate-maleic anhydride copolymers.


One or more alkali metal salts, such as alkali metal silicates, carbonates and sulfates, can be incorporated as an alkaline agent or an inorganic electrolyte, in amounts of 1 to 50% by weight,

preferably 5 to 30% by weight, based on the total weight of the composition. Moreover, alkanolamines represented by triethanolamine, diethanolamine, monoethanolamine and triisopropanolamine can be incorporated as organic alkaline agents.


5 Still further, one or more recontamination preventing agents such as polyethylene glycol, polyvinyl alcohol, polyvinylpyrrolidone and carboxymethyl cellulose can be incorporated in amounts of 0.1 to 5% by weight based on the composition. 5

10 Furthermore, bleaching agents such as sodium percarbonate, sodium perborate, sodium sulfate and sodium chloride-hydrogen peroxide adducts, whitening agents such as commercially available fluorescent dyes, and other additives such as perfumes, enzymes and bluing agents can be incorporated in the composition of the present invention according to need. 10

15 Moreover, talc, finely divided silica, clay, calcium silicate (for example, Microcell manufactured by Johns-Manvill Co.) and zeolites having a divalent metal ion exchange capacity, which are described below, can be incorporated as a water-insoluble substance, according to need. As the zeolites, there can be mentioned, for example, crystalline and/or amorphous aluminosilicates represented by the following formula (u): 15







20 wherein M stands for sodium and/or potassium, and x, y and w are the mole numbers of the respective components, which satisfy the requirements of $0.7 \leq x \leq 1.2$ and $1.6 \leq y \leq 2.8$, and w being an optional positive number inclusive of 0, and crystalline and/or amorphous zeolites represented by the following formula (v): 20



25 wherein M stands for Na or K, and X, Y, Z and W are mole numbers of the respective components, which satisfy the requirements of $0.20 \leq X \leq 1.10$, $0.20 \leq Y \leq 4.00$ and $0.001 \leq Z \leq 0.80$, and W being an optional positive number inclusive of 0. 25

30 Fluorescent dyes, for example, those represented by the following structural formulae (w), (x) and (y), can be incorporated: 30



55 Furthermore, lower alcohols such as ethanol and isopropanol, glycols such as ethylene glycol and propylene glycol, urea, benzene-sulfonic acid salts, p-toluene-sulfonic acid salts, xylene-sulfonic acid salts, benzoic acid salts and salicylic acid salts can be incorporated as a viscosity 55

65

reducing agent, and polyvinyl acetate, vinyl acetate-styrene copolymers and polystyrene can be incorporated as an opacifying agent.

Thus, a fabric-softening detergent having an excellent recontamination preventing effect can be provided according to the present invention.

5 The present invention will now be further described in detail with reference to the following illustrative Examples that by no means limit the scope of the invention. In these Examples, all references to "%" mean percent by weight. 5

In the Examples, the recontamination preventing test and softness test were carried out according to the following methods and the test results were evaluated according to the standards described below. Recontamination preventing test. 10

In 1l of service water was dissolved 2.5 g of the detergent composition, and 0.20 g of carbon black was added to the solution and was uniformly dispersed therein by applying ultrasonic vibrations for 10 minutes. The thus-formed test bath was transferred into a washing tank of a Terg-O-Meter. Clean white test cloth specimens (10 cm × 10 cm) were immersed in the bath and stirred at 25°C for 15 minutes in the Terg-O-Meter. The bath was discharged and 1l of clear service water maintained at 25°C was added in its place and the specimens were stirred in the clear water for 5 minutes to effect rinsing. Then, the water was removed from the specimens by a centrifugal separator and the specimens were air-dried. 15

In the Examples, "wool", "acryl" and "cotton" indicate the following woven fabrics. 20
 "wool": commercially available crimped white sweater
 "acryl": acryl jersey knit wear
 "cotton": white cloth formed by boiling and washing in water a shirting as defined by the Japanese Industrial Standard (JIS L-0206). 20

The whiteness of the tested specimen was examined and the recontamination preventing effect was evaluated as follows: 25

○: whiteness was the same as that of the original specimen before the test
 X: the tested specimen was more black than the original specimen before the test
 Softness test

Five sheets of acryl jersey (30 cm × 60 cm) or one wool sweater (the same as used in the recontamination preventing test) were hand-washed with 5l of an aqueous solution containing 0.25% of the detergent, which was maintained at 30°C. After air drying, the feel of the acryl jersey or wool sweater was examined by five experts and the softness was evaluated according to the following rating criteria. 30

A: softer than the specimen washed with the standard detergent
 B: as soft as the specimen washed with the standard detergent
 C: hard finish touch 35

Example 1

Light-duty liquid detergent compositions comprising the following ingredients were prepared and subjected to the recontamination preventing test. The obtained results are shown in Table 1. 40

Ingredients Incorporated	Amounts (%)	
Secondary alcohol ethoxylate (average carbon number = 11, average added ethylene oxide mole number = 7)		45
Coconut fatty acid diethanolamide	20	
Beef tallow-alkyl-trimethyl ammonium chloride	2	
Recontamination preventing agent (polyoxyalkylene alkyl or alkenyl ether sulfuric acid ester salt or conventional recontamination preventing agent or surface active agent) (see Table 1)	1.5	50
Ethanol	3.5	55
Water	5	
	68	

*: In Table 1, PO stands for propylene oxide group, BO stands for propylene oxide group and EO stands for ethylene oxide group (the same definitions will apply hereinafter). 60

Table 1

Recontamination Preventing Agent	Recontamination Preventing Effect	
5		5
<i>Present Invention</i>		
$C_8H_{17}O(PO)_1SO_3Na$	○	
$C_{12}H_{25}O(PO)_3SO_3Na$	○	
R-beef tallow- $O(PO)_8SO_3Na$	○	
10 $R_1oxoO(PO)_{25}SO_3 \cdot$ triethanolamine	○	10
$R_2oxoO(PO)_4SO_3 \cdot$ triethanolamine	○	
$R_3oxoO(PO)_{10}SO_3 \cdot Na$	○	
$R_4oxoO(PO)_{30}SO_3 \cdot Na$	○	
R-coconut- $O(PO)_3SO_3 \cdot Na$	○	
15 $R_1oxoO(PO)_8SO_3 \cdot Mg_{1/2}$	○	15
$R_1oxoO(PO)_8SO_3 \cdot Ca_{1/2}$	○	
$C_8H_{17}O(BO)_1SO_3Na$	○	
$C_{12}H_{25}O(BO)_3SO_3Na$	○	
$R_1oxoO(BO)_3SO_3 \cdot$ triethanolamine	○	
20 $C_{10}H_{23}O(PO)_1(EO)_1SO_3Na$	○	20
$C_{12}H_{23}O(PO)_2(EO)_1SO_3Na$	○	
$R_1oxoO(PO)_2(EO)_1SO_3Na$	○	
$R_1oxoO(PO)_3(EO)_3SO_3Na$	○	
$R_1oxoO(PO)_3(EO)_8SO_3Na$	○	
25 $R_5oxoO(PO)_4(EO)_8SO_3Na$	○	25
$R_1oxoO(EO)_3(PO)_2SO_3Na$	○	
$R_1oxoO(BO)_1(EO)_2SO_3Na$	○	
<i>Comparison</i>		
30 not added	X	30
$C_{12}H_{25}O(EO)_3SO_3Na$	X	
polyethylene glycol (molecular weight = 6,000)	X	
ethylene oxide-propylene oxide		
35 block copolymer (molecular weight = 2,000)	X	35
$R_1oxoO(EO)_9H$	X	
lauryl betaine	X	
sodium linear-alkyl-benzene-sulfonate		
40 (average carbon number = 13)	X	40
$R_1oxoOSO_3Na$	X	
sodium α -olefin-sulfonate		
(average carbon number = 17)	X	
sodium coconut fatty acid	X	
45 sodium sulfosuccinate	X	45
carboxymethyl cellulose	X	
sodium polyacrylate	X	
50 In the Table, "R-beef tallow-O" stands for a natural beef tallow-derived higher alcohol group, "R-coconut-O" stands for a natural coconut oil-derived higher alcohol group and "RoxoO" stands for an oxo process synthetic higher alcohol group, in which R_1oxo is characterized by an average carbon number of 12 and an iso ratio of 35%, R_2oxo is characterized by an average carbon number of 14 and an iso ratio of 36%, R_3oxo is characterized by an average carbon number of 12 and an iso ratio of 50%, R_4oxo is characterized by an average carbon number of 14 and an iso ratio of 21% and R_5oxo is characterized by an average carbon number of 17 and an iso ratio of 71%.		50
55		55
<i>Example 2</i>		
60 A light-duty liquid detergent composition comprising the following ingredients was prepared and subjected to the recontamination preventing test. The obtained results are shown in Table 2.		60

Ingredients Incorporated	Amounts (%)	
Secondary alcohol ethoxylate (same as used in Example 1)	20	
5 Cationic surfactant (see Table 2)	0 or 2	5
R,oxoO(PO) ₃ (EO) ₁ SO ₃ Na	5	
Ethanol	10	
Water	balance	

10 Table 2

Cationic Surfactant	Recontamination Preventing Effect	
15 Comparison		
Not added	X	
Present Invention		
20 R-beef tallow $\begin{matrix} \text{CH}_3 \\ \\ \text{N}^+ \\ \\ \text{CH}_3 \end{matrix} \cdot \text{Cl}^-$	$\begin{matrix} \text{CH}_3 \\ \\ \text{N}^+ \\ \\ (\text{CH}_2\text{CH}_2\text{O})_8\text{H} \end{matrix} \text{Cl}^-$	20
25 $\begin{matrix} \text{C}_{18}\text{H}_{37} \\ \\ \text{N}^+ \\ \\ \text{CH}_3 \end{matrix} \cdot \text{Cl}^-$	$\begin{matrix} (\text{C}_2\text{H}_4\text{O})_m \text{H} \\ \\ \text{N}^+ \\ \\ (\text{C}_2\text{H}_4\text{O})_n \text{H} \end{matrix} \cdot \text{Cl}^-$ (m+n=8)	25
30 $\begin{matrix} \text{C}_{16}\text{H}_{33}\text{CH}(\text{OH})\text{CH}_2 \\ \\ \text{N}^+ \\ \\ \text{CH}_3 \end{matrix} \cdot \text{Cl}^-$	$\begin{matrix} \text{CH}_3 \\ \\ \text{N}^+ \\ \\ \text{CH}_3 \end{matrix} \cdot \text{Cl}^-$	30
35 $\begin{matrix} \text{C}_{16}\text{H}_{33}\text{CH}(\text{OH})\text{CH}_2 \\ \\ \text{N}^+ \\ \\ \text{CH}_3 \end{matrix} \cdot \text{Cl}^-$	$\begin{matrix} \text{CH}_3 & \text{CH}_3 \\ & \\ \text{N}^+ \\ \\ (\text{CH}_2)_2 \end{matrix} - *$	35
40 * $\text{NHCOC}_{16}\text{H}_{33} \cdot \text{Cl}^-$	$\begin{matrix} \text{CH}_3 & \text{CH}_3 \\ & \\ \text{N}^+ \\ \\ (\text{CH}_2)_2 \end{matrix} - *$	40
45 R-beef tallow- $\text{CONHC}_2\text{H}_4 \cdot \text{N}-\text{C}_2\text{H}_4\text{N}^+ *$	$\begin{matrix} \text{CH}_3 & \text{CH}_3 \\ & \\ \text{N}^+ \\ \\ (\text{CH}_2)_2 \end{matrix} - *$	45
* $(\text{CH}_3)_3 \cdot \text{Cl}^-$	$\begin{matrix} \text{COOR} & \text{beef tallow} \\ \\ \text{N}^+ \\ \\ (\text{CH}_2)_2 \end{matrix} - *$	45
50 (R-beef tallow $\text{CONHCH}_2)_2 \text{NC}_2\text{H}_4\text{N}^+ *$	$\begin{matrix} \text{CH}_3 \\ \\ \text{N}^+ \\ \\ (\text{CH}_2)_2 \end{matrix} - *$	50
* $(\text{CH}_2\text{NHOCR-beef tallow})_2 \text{Cl}^-$	$\begin{matrix} \text{CH}_3 \\ \\ \text{N}^+ \\ \\ (\text{CH}_2)_2 \end{matrix} - *$	50
55 (R-beef tallow- $\text{COOC}_2\text{H}_4)_2 \text{NC}_2\text{H}_4\text{N}^+ *$	$\begin{matrix} \text{CH}_3 \\ \\ \text{N}^+ \\ \\ (\text{CH}_2)_2 \end{matrix} - *$	55
* $(\text{C}_2\text{H}_4\text{OCOR beef tallow})_2 \text{Cl}^-$	$\begin{matrix} \text{CH}_3 \\ \\ \text{N}^+ \\ \\ (\text{CH}_2)_2 \end{matrix} - *$	55

Table 2 continued

		Recontamination Preventing Effect	
5	Cationic Surfactant		5
	cationic surfactant (g)	○	
10	$\left[\begin{array}{c} \text{R-beef tallow} \quad \text{CH}_3 \\ \quad \quad \quad \quad \quad \\ \quad \quad \quad \quad \quad \text{N} \cdot \text{CH}_2\text{CHCH}_2\text{N} \quad \text{CH}_3 \\ \quad \quad \quad \quad \quad / \quad \quad \quad \quad \quad \backslash \\ \text{R-beef tallow} \end{array} \right] \cdot \text{CH}_2\text{CHCH}_2\text{N} \quad \text{CH}_3 \quad *$	○	10
15	$\left. \begin{array}{l} * \text{ R-beef tallow} \\ \text{R-beef tallow} \end{array} \right\}^{2+} \quad 2 \text{Cl}^-$		15
20	cationic surfactant (i) cationic surfactant (j) cationic surfactant (k) cationic surfactant (l) cationic surfactant (m)	○ ○ ○ ○ ○	20
25	Cationic surfactant (g):		25
	Cationic polyamide compound was prepared by reacting 1 mole of diethylene triamine with 2 moles of hydrogenated beef tallow fatty acid to form a condensate (having an acid value of 4.2), adding 1.5 moles of epichlorohydrin to the obtained condensate, subjecting the adduct to ring-opening polymerization in the presence of 0.2 mole of sodium hydroxide and neutralizing the resulting polymer with glycolic acid in an amount of 1.0 mole per mole of the above-mentioned amine.		30
30	Cationic surfactant (i):		
	Merquat 100 manufactured by Merck Co.		
35	Cationic surfactant (j):		35
	Gafquat manufactured by GAF Co.		
	Cationic surfactant (k):		
	Polymer JR manufactured by Union Carbide Corporation Cationic surfactant (l):		
	Cartex L manufactured by National Starch Co. Cationic surfactant (m):		
	Excell manufactured by Nichiden Kagaku K.K.		
40			40
	<i>Example 3</i>		
	A powdery detergent composition comprising the following ingredients was prepared and tested. The results shown in Table 3 were obtained.		
45	<i>Ingredients Incorporated</i>	<i>Amounts (%)</i>	45
	Sodium linear-alkyl-benzene-sulfonate (average carbon number = 12)	20	
	Completely hardened beef tallow fatty acid sodium salt	1	
50	Sodium tripolyphosphate	8.7	50
	Sodium pyrophosphate	4.7	
	Sodium orthophosphate	5.8	
	Sodium silicate	10	
	Sodium carbonate	5	
55	Polyethylene glycol (molecular weight = 6,000)	1	55
	Carboxymethyl cellulose	1	
	Zeolite of type 4A	5	
	Water	10	
60	R-O-A-SO ₃ M (see Table 3)	2.5	60
	Cationic surfactant (see Table 3)	2	
	Glauber salt	balance	

Table 3

Compound of General Formula (1) of	Cationic Surfactant	Recontamination Preventing Effect	Softening Effect	
5 R-O-A-SO ₃ M				5
R: average carbon number of 14.5 and iso ratio of 36%	di-(beef tallow-alkyl) dimethyl ammonium chloride	○	A	
10 A: (PO) ₁ (EO) ₂ M: sodium not added	di-(beef tallow-alkyl) dimethyl ammonium chloride	X	standard	10
15 R: average carbon number of 18 and iso ratio of 0%	di-(beef tallow-alkyl) dimethyl ammonium chloride	○	A	15
A: (PO,EO) ₅ M: sodium not added	di-(beef tallow-alkyl) dimethyl ammonium chloride	X	standard	20
20				
25 CLAIMS				25
1. A detergent composition for washing and softening clothes comprising: from 0.5 to 20% by weight of cationic surfactant having fabric softening properties; from 0.2 to 4 parts by weight, per part by weight of the cationic surfactant, of at least one anionic surfactant having the formula				
30 R-O-(A) _m -SO ₃ M				30
wherein R is alkyl or alkenyl having from 8 to 22 carbon atoms; A is oxypropylene, oxbutylene or oxyethylene, and m has an average value from 0.5 to 30 with the proviso that at least 20% of the A units are oxypropylene and/or oxybutylene and M is an alkali metal, alkaline earth metal or alkanolamine having 2 or 3 carbon atoms.				
35 2. A detergent composition according to claim 1 wherein the anionic surfactant is selected from compounds having the formulae (2) to (7):				35
40 R ₁ O-(-PO-) _{m1} SO ₃ M ₁ (2)				40
R ₁ O-(-PO) _{m2} (EO) _{m3} SO ₃ M ₁ (3)				
R ₁ O-(-BO) _{m2} (EO) _{m3} SO ₃ M ₁ (4)				
R ₁ O-(-PO,EO) _{m1} SO ₃ M ₁ (5)				
R ₁ O-(-BO) _{m1} SO ₃ M ₁ (6)				
45 R ₁ O-(-PO,EO,BO) _{m1} SO ₃ M ₁ (7)				45
wherein R ₁ O is an alkyloxy and/or alkenyloxy residue of a coconut oil-derived higher alcohol, a beef tallow-derived higher alcohol, an oxo process synthetic higher alcohol having an iso ratio of 20 to 80% or a synthetic secondary high alcohol, PO is oxypropylene, BO is oxybutylene, EO is oxyethylene, m ₁ is an integer from 1 to 10, the sum of m ₂ and m ₃ is in the range of from 1 to 10, the ratio of m ₂ /m ₃ is in the range of from 4/1 to 1/4, the PO, EO and BO units in (PO,EO) and (PO,EO,BO) are randomly arranged and M ₁ is sodium, potassium, monoethanolamine, diethanolamine, triethanolamine or magnesium.				
50 3. A detergent composition according to claim 2 wherein R ₁ O is a residue of an oxo process synthetic higher alcohol having an iso ratio of 20 to 80%.				50
55 4. A detergent composition according to claim 2 wherein said anionic surfactant has the formula (2):				55
60 R ₁ O(PO) _{m1} SO ₃ M ₁ (2)				60
5. A detergent composition according to claim 2 wherein said anionic surfactant has the formula (3):				
R ₁ O(PO) _{m2} (EO) _{m3} SO ₃ M ₁ (3)				

6. A detergent composition according to any preceding claim wherein the cationic surfactant is selected from (a) di-(long-chain alkyl) quaternary ammonium salts, (b) mono-(long-chain alkyl) quaternary ammonium salts, (c) di-(long-chain alkyl) polyoxyethylene quaternary ammonium salts and (d) mono-(long-chain alkyl) polyoxyethylene quaternary ammonium salts.
- 5 7. A detergent composition according to claim 6 wherein the amount of cationic surfactant is 1 to 5% by weight. 5
8. A detergent composition according to claim 2 further containing up to 50% by weight of an anionic synthetic water-soluble organic surfactant effective for washing clothes and different from said anionic surfactant of formula (1), and/or an amphoteric synthetic, water-soluble organic surfactant effective for washing clothes, and/or a non-ionic synthetic, water-soluble organic surfactant effective for washing clothes or mixture thereof; and/or up to 50% by weight of water-soluble builder salts for detergents, water-soluble polymeric polyelectrolytes for detergents or mixture thereof; and/or up to 50% by weight of water-soluble inorganic alkaline agent for detergents, water-soluble organic alkaline agent for detergents or mixture thereof; and
- 10 15 water. 10 15
9. A detergent composition according to claim 8 containing from 5 to 40% by weight of said synthetic water-soluble organic surfactant; from 5 to 30% by weight of alkali metal salts selected from the group consisting of alkali silicates, alkali metal carbonates and alkali metal sulfates.