



US008987457B2

(12) **United States Patent**
Burger et al.(10) **Patent No.:** **US 8,987,457 B2**
(45) **Date of Patent:** **Mar. 24, 2015**(54) **RING-SUBSTITUTED N-PYRIDINYL AMIDES AS KINASE INHIBITORS**(71) Applicant: **Novartis AG**, Basel (CH)(72) Inventors: **Matthew Burger**, Albany, CA (US); **Joseph E. Drumm, III**, Walnut Creek, CA (US); **Gisele Nishiguchi**, Albany, CA (US); **Alice Rico**, Castro Valley, CA (US); **Robert Lowell Simmons**, San Francisco, CA (US); **Benjamin Taft**, Oakland, CA (US); **Huw Tanner**, Alameda, CA (US)(73) Assignee: **Novartis AG**, Basel (CH)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **13/897,927**(22) Filed: **May 20, 2013**(65) **Prior Publication Data**

US 2013/0336965 A1 Dec. 19, 2013

Related U.S. Application Data

(60) Provisional application No. 61/695,533, filed on Aug. 31, 2012, provisional application No. 61/649,645, filed on May 21, 2012.

(51) **Int. Cl.**

C07D 211/68 (2006.01)
C07D 213/44 (2006.01)
C07D 417/00 (2006.01)
A61K 31/445 (2006.01)
A61K 31/41 (2006.01)
C07D 213/81 (2006.01)
C07D 403/12 (2006.01)
C07D 401/14 (2006.01)
C07D 401/12 (2006.01)
C07D 403/14 (2006.01)
C07D 407/14 (2006.01)
C07D 409/14 (2006.01)
C07D 417/12 (2006.01)
C07D 417/14 (2006.01)
A61K 31/4439 (2006.01)
A61K 31/444 (2006.01)
A61K 31/4545 (2006.01)
A61K 31/506 (2006.01)
A61K 31/675 (2006.01)
A61K 45/06 (2006.01)
C07D 405/14 (2006.01)
C07F 9/6558 (2006.01)

(52) **U.S. Cl.**

CPC **C07D 213/81** (2013.01); **C07D 403/12** (2013.01); **C07D 401/14** (2013.01); **C07D 401/12** (2013.01); **C07D 403/14** (2013.01); **C07D 407/14** (2013.01); **C07D 409/14** (2013.01); **C07D 417/12** (2013.01); **C07D 417/14** (2013.01); **A61K 31/4439** (2013.01); **A61K 31/444** (2013.01); **A61K 31/4545**

(2013.01); **A61K 31/506** (2013.01); **A61K 31/675** (2013.01); **A61K 45/06** (2013.01); **C07D 405/14** (2013.01); **C07F 9/65583** (2013.01)

USPC **546/193**; 546/262; 546/270.7; 514/318; 514/382

(58) **Field of Classification Search**

USPC 546/193, 262, 270.7; 514/318, 382
 See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

8,168,794 B2 * 5/2012 Burger et al. 546/270.7
 8,329,732 B2 * 12/2012 Burger et al. 514/332
 2005/0245530 A1 11/2005 Borzilleri et al.

(Continued)

FOREIGN PATENT DOCUMENTS

CA 2649043 11/2007
 WO WO 01/55155 8/2001

(Continued)

OTHER PUBLICATIONS

Johnson et al., "Relationships Between Drug Activity in NC1 Preclinical in vitro and in vivo Models and Early Clinical Trials," *British Journal of Cancer* 84:1424-1431, 2001.

(Continued)


Primary Examiner — Paul Zarek

Assistant Examiner — George W Kosturko

(74) Attorney, Agent, or Firm — Stephen Johnson

(57) **ABSTRACT**

The present invention provides a compound of formula (A):



as described herein, and pharmaceutically acceptable salts, enantiomers, rotamers, tautomers, or racemates thereof. Also provided are methods of treating a disease or condition mediated by PIM kinase using the compounds of Formula I, and pharmaceutical compositions comprising such compounds.

9 Claims, No Drawings

(56)

References Cited

U.S. PATENT DOCUMENTS

2006/0004197 A1 1/2006 Thrash
 2009/0286766 A1 11/2009 Sugasawa et al.
 2010/0056576 A1* 3/2010 Burger et al. 514/332
 2010/0216839 A1* 8/2010 Burger et al. 514/318
 2010/0311980 A1 12/2010 Rao et al.
 2012/0208815 A1 8/2012 Burger et al.

FOREIGN PATENT DOCUMENTS

WO WO 02/057261 7/2002
 WO WO 02/076986 10/2002
 WO WO 03/008365 1/2003
 WO WO 2005/033097 4/2005
 WO WO 2005/042488 5/2005
 WO WO 2005/056547 6/2005
 WO WO 2006/105081 10/2006
 WO WO 2007/019344 2/2007
 WO WO 2007/044724 4/2007
 WO WO 2007/058942 5/2007
 WO WO 2007/123269 11/2007
 WO WO 2008/022164 2/2008
 WO WO 2008/054701 5/2008
 WO WO 2008/054702 5/2008
 WO WO 2008/054749 5/2008
 WO WO 2008/106692 9/2008
 WO WO 2009/014637 1/2009
 WO WO 2009/103739 8/2009

WO WO 2009/109541 9/2009
 WO WO 2009/109576 9/2009
 WO WO 2009/110844 9/2009
 WO WO 2009/111309 9/2009
 WO WO 2009/111337 9/2009
 WO WO 2009/118475 10/2009
 WO WO 2010/026124 3/2010
 WO WO 2011/016234 8/2010
 WO WO 2012/004217 1/2012
 WO WO 2012/100135 7/2012
 WO 2012/120415 A1 9/2012

OTHER PUBLICATIONS

Merkel, "PIM 1 Kinase as a Target for Cancer Therapy" *Expert Opin. Investig. Drug* [Early Online], 1-12, 2012.
 Trisha Gura "Cancer Models: Systems for Identifying New Drugs are Often Faculty" *Science* 278(5340):1041-1042, Nov. 1, 1997.
 Simone, *Oncology: Introduction*, Cecil Textbook of Medicine, 20th Edition, 1996, 1:1004-1010.
 Dorwald, *Side Reactions in Organic Synthesis*, 2005 Wiley: VCH Weinheim Preface, pp. 1-15 & Chapter 8, pp. 279-308.
 Garham Atwell et al., "Potential Antitumor Agents, 55, 6-Phenylphenanthride-4-Carboxamides: A New Class of DNA-Intercalating Antitumor Agents" *J. Med. Chem.* 31:774-779, 1988.
 Second Annual Anti-Inflammatories: Small Molecule Approaches, "Newer Kinase Inflammation Targets (BTK and Beyond)" Apr. 12-13, 2011, San Diego, CA.

* cited by examiner

1

RING-SUBSTITUTED N-PYRIDINYL AMIDES AS KINASE INHIBITORS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. provisional application Ser. Nos. 61/649,645, filed 21 May 2012, and 61/695,533, filed 31 Aug. 2012, the contents of which are incorporated herein by reference in their entirety.

FIELD OF THE INVENTION

The present invention relates to new compounds and their tautomers and pharmaceutically acceptable salts, esters, metabolites or prodrugs thereof, compositions of the new compounds together with pharmaceutically acceptable carriers, and uses of the new compounds, either alone or in combination with at least one additional therapeutic agent, in the prophylaxis or treatment of cancer and other cellular proliferation disorders.

BACKGROUND

Infection with the Moloney retrovirus and genome integration in the host cell genome results in development of lymphomas in mice. Provirus Integration of Moloney Kinase (PIM-Kinase) was identified as one of the frequent proto-oncogenes capable of being transcriptionally activated by this retrovirus integration event (Cuypers H T et al., "Murine leukemia virus-induced T-cell lymphomagenesis: integration of proviruses in a distinct chromosomal region," *Cell* 37(1): 141-50 (1984); Selten G, et al., "Proviral activation of the putative oncogene Pim-1 in MuLV induced T-cell lymphomas" *EMBO J* 4(7):1793-8 (1985)), thus establishing a correlation between over-expression of this kinase and its oncogenic potential. Sequence homology analysis demonstrated that there are three highly homologous Pim-Kinases (Pim1, 2 & 3), Pim1 being the proto-oncogene originally identified by retrovirus integration. Furthermore, transgenic mice over-expressing Pim1 or Pim2 show increased incidence of T-cell lymphomas (Breuer M et al., "Very high frequency of lymphoma induction by a chemical carcinogen in pim-1 transgenic mice" *Nature* 340(6228):61-3 (1989)), while over-expression in conjunction with c-myc is associated with incidence of B-cell lymphomas (Verbeek S et al., "Mice bearing the E mu-myc and E mu-pim-1 transgenes develop pre-B-cell leukemia prenatally" *Mol Cell Biol* 11(2):1176-9 (1991)). Thus, these animal models establish a strong correlation between Pim over-expression and oncogenesis in hematopoietic malignancies.

In addition to these animal models, Pim over-expression has been reported in many human malignancies. Pim1, 2 & 3 over-expression is frequently observed in hematopoietic malignancies (Amson R et al., "The human protooncogene product p33pim is expressed during fetal hematopoiesis and in diverse leukemias," *PNAS USA* 86(22):8857-61 (1989); Cohen A M et al., "Increased expression of the hPim-2 gene in human chronic lymphocytic leukemia and non-Hodgkin lymphoma," *Leuk Lymph* 45(5):951-5 (2004); Huttmann A et al., "Gene expression signatures separate B-cell chronic lymphocytic leukaemia prognostic subgroups defined by ZAP-70 and CD38 expression status," *Leukemia* 20:1774-1782 (2006)) and in prostate cancer (Dhanasekaran S M, et al., "Delineation of prognostic biomarkers in prostate cancer," *Nature* 412(6849):822-6 (2001); Cibull T L, et al., "Overexpression of Pim-1 during progression of prostatic adenocar-

2

cinoma," *J Clin Pathol* 59(3):285-8 (2006)), while over-expression of Pim3 is frequently observed in hepatocellular carcinoma (Fujii C, et al., "Aberrant expression of serine/threonine kinase Pim-3 in hepatocellular carcinoma development and its role in the proliferation of human hepatoma cell lines," *Int J Cancer* 114:209-218 (2005)) and pancreatic cancer (Li Y Y et al., "Pim-3, a proto-oncogene with serine/threonine kinase activity, is aberrantly expressed in human pancreatic cancer and phosphorylates bad to block bad-mediated apoptosis in human pancreatic cancer cell lines," *Cancer Res* 66(13):6741-7 (2006)).

Pim1, 2 & 3 are Serine/Threonine kinases that normally function in survival and proliferation of hematopoietic cells in response to growth factors and cytokines. Cytokines signaling through the Jak/Stat pathway leads to activation of transcription of the Pim genes and synthesis of the proteins. No further post-translational modifications are required for the Kinase Pim activity. Thus, signaling downstream is primarily controlled at the transcriptional/translational and protein turnover level. Substrates for Pim kinases include regulators of apoptosis such as the Bcl-2 family member BAD (Aho T et al., "Pim-1 kinase promotes inactivation of the pro-apoptotic Bad protein by phosphorylating it on the Ser112 gatekeeper site," *FEBS Letters* 571: 43-49 (2004)), cell cycle regulators such as p21^{WAF1/CIP1} (Wang Z, et al., "Phosphorylation of the cell cycle inhibitor p21Cip1/WAF1 by Pim-1 kinase," *Biochem Biophys Acta* 1593:45-55 (2002)), CDC25A (1999), C-TAK (Bachmann M et al., "The Oncogenic Serine/Threonine Kinase Pim-1 Phosphorylates and Inhibits the Activity of Cdc25C-associated Kinase 1 (C-TAK1). A novel role for Pim-1 at the G2/M cell cycle checkpoint," *J Biol Chem* 279:48319-48328 (2004)) and NuMA (Bhattacharya N, et al., "Pim-1 associates with protein complexes necessary for mitosis," *Chromosoma* 111(2): 80-95 (2002)) and the protein synthesis regulator 4EBP1 (Hammerman P S et al., "Pim and Akt oncogenes are independent regulators of hematopoietic cell growth and survival," *Blood* 105(11):4477-83 (2005)). The effects of Pim(s) in these regulators are consistent with a role in protection from apoptosis and promotion of cell proliferation and growth. Thus, over-expression of Pim(s) in cancer is thought to play a role in promoting survival and proliferation of cancer cells and, therefore, their inhibitions should be an effective way of treating cancers in which they are over-expressed. In fact several reports indicate that knocking down expression of Pim(s) with siRNA results in inhibition of proliferation and cell death (Dai J M, et al., "Antisense oligodeoxynucleotides targeting the serine/threonine kinase Pim-2 inhibited proliferation of DU-145 cells," *Acta Pharmacol Sin* 26(3):364-8 (2005); Fujii et al. 2005; Li et al. 2006).

Furthermore, mutational activation of several well known oncogenes in hematopoietic malignancies is thought to exert its effects at least in part through Pim(s). For example, targeted down-regulation of Pim expression impairs survival of hematopoietic cells transformed by Flt3 and BCR/ABL (Adam et al. 2006). Thus, inhibitors to Pim1, 2 and 3 would be useful in the treatment of these malignancies.

In addition to a potential role in cancer treatment and myeloproliferative diseases, such inhibitor could be useful to control expansion of immune cells in other pathologic condition such as autoimmune diseases, allergic reactions and in organ transplantation rejection syndromes. This notion is supported by the findings that differentiation of Th1 Helper T-cells by IL-12 and IFN- α results in induction of expression of both Pim1 and Pim2 (Aho T et al., "Expression of human Pim family genes is selectively up-regulated by cytokines promoting T helper type 1, but not T helper type 2, cell


differentiation," *Immunology* 116: 82-88 (2005)). Moreover, Pim(s) expression is inhibited in both cell types by the immunosuppressive TGF- β (Aho et al. 2005). These results suggest that Pim kinases are involved in the early differentiation process of Helper T-cells, which coordinate the immunological responses in autoimmune diseases, allergic reaction and tissue transplant rejection. Recent reports demonstrate that Pim kinase inhibitors show activity in animal models of inflammation and autoimmune diseases. See J E Robinson "Targeting the Pim Kinase Pathway for Treatment of Autoimmune and Inflammatory Diseases," for the Second Annual Conference on Anti-Inflammatories: Small Molecule Approaches," San Diego, Calif. (Conf. April 2011; Abstract published earlier on-line).

A continuing need exists for compounds that inhibit the proliferation of capillaries, inhibit the growth of tumors, treat cancer, modulate cell cycle arrest, and/or inhibit molecules such as Pim1, Pim2 and Pim3, and pharmaceutical formulations and medicaments that contain such compounds. A need also exists for methods of administering such compounds, pharmaceutical formulations, and medicaments to patients or subjects in need thereof. The present invention addresses such needs.

Earlier patent applications have described compounds that inhibit Pims and function as anticancer therapeutics, see, e.g., WO2012/004217, WO2010/026124, WO 2008/106692 and WO2011/124580, and as treatment for inflammatory conditions such as Crohn's disease, inflammatory bowel disease, rheumatoid arthritis, and chronic inflammatory diseases, see e.g., WO 2008/022164. The present invention provides compounds that inhibit activity of one or more Pims, preferably two or more Pims, more preferably Pim1, Pim2 and Pim3, at nanomolar levels (e.g., IC₅₀ under 100 nM or under 50 nM) and exhibit distinctive characteristics that may provide improved therapeutic effects and pharmacokinetic properties, such as reduced drug-drug interactions associated with inhibition of cytochrome oxidases, relative to compounds previously disclosed. Compounds of the invention contain novel substitution combinations on one or more rings that provide these distinctive properties and are suitable for treating Pim-related conditions such as those described herein.

SUMMARY OF THE INVENTION

The invention provides compounds of Formula A, having three substituents on a cyclohexyl or piperidinyl ring that is attached to an N-pyridinyl amide moiety of Formula (A):



wherein:

groups attached to the ring containing Q that are depicted inside the ring are all syn to each other, and all groups attached to that ring that are depicted outside the ring are syn to one another;

Q is C or N;

R^q is H when Q is C, and R^q is absent when Q is N;

R¹ is selected from H, C₁₋₄ alkyl, C₃₋₆ cycloalkyl, C₄₋₆ heterocyclyl, —(CR'₂)₁₋₃OR' and —OR',

where each R' is independently H or C₁₋₄ alkyl,

and each alkyl, cycloalkyl and heterocyclyl is optionally substituted with up to two groups selected from halo, CN, NH₂, hydroxy, oxo, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, and C₁₋₄ haloalkoxy;

R^{1b}, R^{1c}, and R^{1d} are each independently selected from H, halo, OR', R', —(CH₂)₁₋₂OR', and CONR'₂;

one of R^{2a} and R^{2b} is H, and the other of R^{2a} and R^{2b} is selected from CN, halo, azido, amino, —OR, —O(CH₂)₁₋₃OR, —NRC(O)R, —NRC(O)OR, —NHSO₂R, —SO₂R, —OSO₂R, —SR, —S(O)R, —OP(O)R₂, and 1-pyridonyl or 1-triazolyl optionally substituted with up to two groups selected from halo, C₁₋₄ alkyl, hydroxy-substituted C₁₋₄ alkyl, C₂₋₄ alkenyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, and C₅₋₆ heteroaryl;

where each R is independently C₅₋₆ heteroaryl or C₁₋₄ alkyl

optionally substituted with up to three groups selected from cyano, halo, hydroxy, carboxy, C₁₋₄ alkylsulfonyl, and C₁₋₄ alkoxy; or R^{2a} and R^{2b} taken together may form a dialkyl ketal or 5-6 membered cyclic ketal, =O or =N—OR", where R" is H or C₁₋₄ alkyl;

ring A is selected from pyridinyl, pyrimidinyl, pyrazinyl, and thiazolyl, each having N positioned as shown in Formula (I); and

Ring A is optionally substituted with 1 or 2 groups selected from halo, CN, NH₂, hydroxy, C₁₋₄ alkyl, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, and C₁₋₄ haloalkoxy;

or a pharmaceutically acceptable salt thereof.

In some embodiments of the compounds of Formula A, R^{1b} is H and R¹ is selected from H, C₁₋₄ alkyl, C₃₋₆ cycloalkyl, C₄₋₆ heterocyclyl, —(CR'₂)₁₋₃OR' and —OR', where each R' is independently H or C₁₋₄ alkyl, and each alkyl, cycloalkyl and heterocyclyl is optionally substituted with up to two groups selected from halo, CN, NH₂, hydroxy, oxo, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, and C₁₋₄ haloalkoxy;

one of R^{2a} and R^{2b} is H, and the other of R^{2a} and R^{2b} is selected from CN, halo, azido, amino, —OR, —O(CH₂)₁₋₃OR, —NRC(O)R, —NRC(O)OR, —NHSO₂R, —SO₂R, —OSO₂R, —SR, —S(O)R, —OP(O)R₂, and triazolyl optionally substituted with up to two groups selected from halo, C₁₋₄ alkyl, hydroxy-substituted C₁₋₄ alkyl, C₁₋₄ alkoxy, and C₅₋₆ heteroaryl;

where each R is independently C₁₋₄ alkyl optionally substituted with up to three groups selected from cyano, halo, hydroxy, carboxy, C₁₋₄ alkylsulfonyl, and C₁₋₄ alkoxy; or R^{2a} and R^{2b} taken together form a dialkyl ketal or 5-6 membered cyclic ketal, =O or =N—OR", where R" is H or C₁₋₄ alkyl;

Ring A is selected from pyridinyl, pyrimidinyl, pyrazinyl, and thiazolyl, each having N positioned as shown in Formula (I); and

Ring A is optionally substituted with 1 or 2 groups selected from halo, CN, NH₂, hydroxy, C₁₋₄ alkyl, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, and C₁₋₄ haloalkoxy;

or a pharmaceutically acceptable salt thereof.

Additional embodiments of these compounds and pharmaceutical compositions and uses for these compounds and compositions are described below.

These compounds are inhibitors of Pim kinases as further discussed herein. These compounds and their pharmaceuti-

cally acceptable salts, and pharmaceutical compositions containing these compounds and salts are useful for therapeutic methods such as treatment of cancers and autoimmune disorders that are caused by or exacerbated by excessive levels of Pim kinase activity.

DETAILED DESCRIPTION OF EMBODIMENTS OF THE INVENTION

“PIM inhibitor” is used herein to refer to a compound that exhibits an IC_{50} with respect to PIM Kinase activity of no more than about 100 μ M and more typically not more than about 50 μ M, as measured in the PIM depletion assays described herein below for at least one of Pim1, Pim2 and Pim3. Preferred compounds have an IC_{50} below about 1 micromolar on at least one Pim, and generally have an IC_{50} below 100 nM on each of Pim1, Pim2 and Pim3.

The phrase “alkyl” refers to hydrocarbon groups that do not contain heteroatoms, i.e., they consist of carbon atoms and hydrogen atoms. Thus the phrase includes straight chain alkyl groups such as methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl and the like. The phrase also includes branched chain isomers of straight chain alkyl groups, including but not limited to, the following which are provided by way of example: $-\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}(\text{CH}_2\text{CH}_3)_2$, $-\text{C}(\text{CH}_3)_3$, $-\text{C}(\text{CH}_2\text{CH}_3)_3$, $-\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, $-\text{CH}(\text{CH}_3)_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, and others. Thus the term “alkyl” includes primary alkyl groups, secondary alkyl groups, and tertiary alkyl groups. Typical alkyl groups include straight and branched chain alkyl groups having 1 to 12 carbon atoms, preferably 1-6 carbon atoms. The term “lower alkyl” or “loweralkyl” and similar terms refer to alkyl groups containing up to 6 carbon atoms.

The term “alkenyl” refers to alkyl groups as defined above, wherein there is at least one carbon-carbon double bond, i.e., wherein two adjacent carbon atoms are attached by a double bond. The term “alkynyl” refers to alkyl groups wherein two adjacent carbon atoms are attached by a triple bond. Typical alkenyl and alkynyl groups contain 2-12 carbon atoms, preferably 2-6 carbon atoms. Lower alkenyl or lower alkynyl refers to groups having up to 6 carbon atoms. An alkenyl or alkynyl group may contain more than one unsaturated bond, and may include both double and triple bonds, but of course their bonding is consistent with well-known valence limitations.

The term “alkoxy” refers to $-\text{OR}$, wherein R is alkyl.

As used herein, the term “halogen” or “halo” refers to chloro, bromo, fluoro and iodo groups. Typical halo substituents are F and/or Cl. “Haloalkyl” refers to an alkyl radical substituted with one or more halogen atoms. The term “haloalkyl” thus includes monohalo alkyl, dihalo alkyl, trihalo alkyl, perhaloalkyl, and the like.

“Amino” refers herein to the group $-\text{NH}_2$. The term “alkylamino” refers herein to the group $-\text{NRR}'$ where R and R' are each independently selected from hydrogen or a lower alkyl, provided $-\text{NRR}'$ is not $-\text{NH}_2$. The term “aryl amino” refers herein to the group $-\text{NRR}'$ where R is aryl and R' is hydrogen, a lower alkyl, or an aryl. The term “aralkyl amino” refers herein to the group $-\text{NRR}'$ where R is a lower aralkyl and R'

is hydrogen, a loweralkyl, an aryl, or a loweraralkyl. The term cyano refers to the group $-\text{CN}$. The term nitro refers to the group $-\text{NO}_2$.

The term “alkoxyalkyl” refers to the group $-\text{alk}_1-\text{O}-\text{alk}_2$ where alk_1 is an alkyl or alkenyl linking group, and alk_2 is alkyl or alkenyl. The term “loweralkoxyalkyl” refers to an alkoxyalkyl where alk_1 is loweralkyl or loweralkenyl, and alk_2 is loweralkyl or loweralkenyl. The term “aryl-oxyalkyl” refers to the group $-\text{alkyl}-\text{O}-\text{aryl}$, where $-\text{alkyl}-$ is a C_{1-12} straight or branched chain alkyl linking group, preferably C_{1-6} . The term “aralkoxyalkyl” refers to the group $-\text{alkylenyl}-\text{O}-\text{aralkyl}$, where aralkyl is preferably a loweraralkyl.

The term “aminocarbonyl” refers herein to the group $-\text{C}(\text{O})-\text{NH}_2$. “Substituted aminocarbonyl” refers herein to the group $-\text{C}(\text{O})-\text{NRR}'$ where R is loweralkyl and R' is hydrogen or a loweralkyl. In some embodiments, R and R', together with the N atom attached to them may be taken together to form a “heterocycloalkylcarbonyl” group. The term “arylaminocarbonyl” refers herein to the group $-\text{C}(\text{O})-\text{NRR}'$ where R is an aryl and R' is hydrogen, loweralkyl or aryl. “aralkylaminocarbonyl” refers herein to the group $-\text{C}(\text{O})-\text{NRR}'$ where R is loweraralkyl and R' is hydrogen, loweralkyl, aryl, or loweraralkyl.

“Aminosulfonyl” refers herein to the group $-\text{S}(\text{O})_2-\text{NH}_2$. “Substituted aminosulfonyl” refers herein to the group $-\text{S}(\text{O})_2-\text{NRR}'$ where R is loweralkyl and R' is hydrogen or a loweralkyl. The term “aralkylaminosulfonyl” refers herein to the group $-\text{aryl}-\text{S}(\text{O})_2-\text{NH}-\text{aralkyl}$, where the aralkyl is loweraralkyl.

“Carbonyl” refers to the divalent group $-\text{C}(\text{O})-$. “Carboxy” refers to $-\text{C}(=\text{O})-\text{OH}$. “Alkoxycarbonyl” refers to ester $-\text{C}(=\text{O})-\text{OR}$ wherein R is optionally substituted lower alkyl. “Loweralkoxycarbonyl” refers to ester $-\text{C}(=\text{O})-\text{OR}$ wherein R is optionally substituted lower loweralkyl. “Cycloalkyloxycarbonyl” refers to $-\text{C}(=\text{O})-\text{OR}$ wherein R is optionally substituted C3-C8 cycloalkyl.

“Cycloalkyl” refers to a mono- or polycyclic, carbocyclic alkyl substituent. Carbocycloalkyl groups are cycloalkyl groups in which all ring atoms are carbon. Typical cycloalkyl substituents have from 3 to 8 backbone (i.e., ring) atoms. When used in connection with cycloalkyl substituents, the term “polycyclic” refers herein to fused and non-fused alkyl cyclic structures. The term “partially unsaturated cycloalkyl”, “partially saturated cycloalkyl”, and “cycloalkenyl” all refer to a cycloalkyl group wherein there is at least one point of unsaturation, i.e., wherein to adjacent ring atoms are connected by a double bond or a triple bond. Such rings typically contain 1-2 double bonds for 5-6 membered rings, and 1-2 double bonds or one triple bond for 7-8 membered rings. Illustrative examples include cyclohexenyl, cyclooctenyl, cyclopropenyl, cyclobutenyl, cyclohexadienyl, and the like.

The term “heterocycloalkyl” refers herein to cycloalkyl substituents that have from 1 to 5, and more typically from 1 to 4 heteroatoms as ring members in place of carbon atoms. Preferably, heterocycloalkyl or “heterocyclyl” groups contain one or two heteroatoms as ring members, typically only one heteroatom for 3-5 membered rings and 1-2 heteroatoms for 6-8 membered rings. Suitable heteroatoms employed in heterocyclic groups of the present invention are nitrogen, oxygen, and sulfur. Representative heterocycloalkyl moieties include, for example, pyrrolidinyl, tetrahydrofuranyl, oxirane, oxetane, oxepane, thiirane, thietane, azetidine, morpholino, piperazinyl, piperidinyl and the like.


The terms “substituted heterocycle”, “heterocyclic group” or “heterocycle” as used herein refers to any 3- or 4-membered ring containing a heteroatom selected from nitrogen, oxygen, and sulfur or a 5- or 6-membered ring containing

7


from one to three heteroatoms, preferably 1-2 heteroatoms, selected from the group consisting of nitrogen, oxygen, or sulfur; wherein the 5-membered ring has 0-2 double bonds and the 6-membered ring has 0-3 double bonds; wherein the nitrogen and sulfur atom may be optionally oxidized; wherein the nitrogen and sulfur heteroatoms may be optionally quar-

ternized; and including any bicyclic group in which any of the above heterocyclic rings is fused to a benzene ring or another 5- or 6-membered heterocyclic ring as described herein. Preferred heterocycles include, for example: diazapinyl, pyrrolinyl, pyrrolidinyl, pyrazolinyl, pyrazolidinyl, imidazolinyl, imidazolidinyl, piperidinyl, piperazinyl, N-methyl piperazinyl, azetidiny, N-methylazetidiny, oxazolidinyl, isoazolidinyl, morpholinyl, thiazolidinyl, isothiazolidinyl, and oxiranyl. The heterocyclic groups may be attached at various positions as will be apparent to those having skill in the organic and medicinal chemistry arts in conjunction with the disclosure herein.

The term 'syn to one another' as used herein means the groups being described are all on the same face of the ring they are attached to, e.g., the groups are either all above the plane of the ring, or they are all below the plane of the ring. Compounds of Formula I can thus be depicted as (Ia) or (Ib) when Q is C:



(Ia)



(Ib)

Heterocyclic moieties can be unsubstituted or they can be substituted with one or more substituents independently selected from hydroxy, halo, oxo (C=O), alkylimino (RN=, wherein R is a loweralkyl or loweralkoxy group), amino, alkylamino, dialkylamino, acylaminoalkyl, alkoxy, thioalkoxy, lower alkoxyalkoxy, loweralkyl, cycloalkyl or haloalkyl. Typically, substituted heterocyclic groups will have up to four substituent groups. The term "cyclic ether" as used herein refers to a 3-7 membered ring containing one oxygen atom (O) as a ring member. Where the cyclic ether is "optionally substituted" it can be substituted at any carbon atom with a group suitable as a substituent for a heterocyclic

8

group, typically up to three substituents selected from lower alkyl, lower alkoxy, halo, hydroxy, —C(O)-lower alkyl, and —C(O)-lower alkoxy. In preferred embodiments, halo, hydroxy and lower alkoxy are not attached to the carbon atoms of the ring that are bonded directly to the oxygen atom in the cyclic ether ring. Specific examples include oxirane, oxetane (e.g., 3-oxetane), tetrahydrofuran (including 2-tetrahydrofuranyl and 3-tetrahydrofuranyl), tetrahydropyran (e.g., 4-tetrahydropyran), and oxepane.

"Aryl" refers to monocyclic and polycyclic aromatic groups having from 5 to 14 backbone carbon or hetero atoms, and includes both carbocyclic aryl groups and heteroaromatic aryl groups. Carbocyclic aryl groups are aryl groups in which all ring atoms in the aromatic ring are carbon, typically including phenyl and naphthyl. Exemplary aryl moieties employed as substituents in compounds of the present invention include phenyl, pyridyl, pyrimidinyl, thiazolyl, indolyl, imidazolyl, oxadiazolyl, tetrazolyl, pyrazinyl, triazolyl, thiophenyl, furanyl, quinolinyl, purinyl, naphthyl, benzothiazolyl, benzopyridyl, and benzimidazolyl, and the like. When used in connection with aryl substituents, the term "polycyclic aryl" refers herein to fused and non-fused cyclic structures in which at least one cyclic structure is aromatic, such as, for example, benzodioxolo (which has a heterocyclic structure fused to a phenyl group, naphthyl, and the like. Where "aryl" is used, the group is preferably a carbocyclic group; the term "heteroaryl" is used for aryl groups when ones containing one or more heteroatoms are preferred.

The term "heteroaryl" refers herein to aryl groups having from 1 to 4 heteroatoms as ring atoms in an aromatic ring with the remainder of the ring atoms being carbon atoms, in a 5-14 atom aromatic ring system that can be monocyclic or polycyclic. Monocyclic heteroaryl rings are typically 5-6 atoms in size. Exemplary heteroaryl moieties employed as substituents in compounds of the present invention include pyridyl, pyrimidinyl, thiazolyl, indolyl, imidazolyl, oxadiazolyl, tetrazolyl, pyrazinyl, triazolyl, thiophenyl, furanyl, quinolinyl, purinyl, benzothiazolyl, benzopyridyl, and benzimidazolyl, and the like.

"Alkyl" or "arylalkyl" refers to an aryl group connected to a structure through an alkylene linking group, e.g., a structure such as —(CH₂)₁₋₄—Ar, where Ar represents an aryl group. "Lower alkyl" or similar terms indicate that the alkyl linking group has up to 6 carbon atoms.

"Optionally substituted" or "substituted" refers to the replacement of one or more hydrogen atoms with a monovalent or divalent radical. Alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl groups described herein may be substituted or unsubstituted. Suitable substitution groups include, for example, hydroxy, nitro, amino, imino, cyano, halo, thio, sulfonyl, thioamido, amidino, imidino, oxo, oxamidino, methoxamidino, imidino, guanidino, sulfonamido, carboxyl, formyl, loweralkyl, haloloweralkyl, loweralkylamino, haloloweralkylamino, loweralkoxy, haloloweralkoxy, loweralkoxyalkyl, alkylcarbonyl, aminocarbonyl, arylcarbonyl, aralkylcarbonyl, heteroarylcarbonyl, heteroarylcarbonyl, alkylthio, aminoalkyl, cyanoalkyl, aryl and the like, provided that oxo, imidino or other divalent substitution groups are not placed on aryl or heteroaryl rings due to the well known valence limitations of such rings.

The substitution group can itself be substituted where valence permits, i.e., where the substitution group contains at least one CH, NH or OH having a hydrogen atom that can be replaced. The group substituted onto the substitution group can be carboxyl, halo (on carbon only); nitro, amino, cyano, hydroxy, loweralkyl, loweralkoxy, C(O)R, —OC(O)R, —OC

(O)OR, —NRCOR, —CONR₂, —NRCOOR, —C(S)NR₂, —NRC(S)R, —OC(O)NR₂, —SR, —SO₃H, —SO₂R or C3-8 cycloalkyl or 3-8 membered heterocycloalkyl, where each R is independently selected from hydrogen, lower haloalkyl, lower alkoxyalkyl, and loweralkyl, and where two R on the same atom or on directly connected atoms can be linked together to form a 5-6 membered heterocyclic ring.

When a substituted substituent includes a straight chain group, the substitution can occur either within the chain (e.g., 2-hydroxypropyl, 2-aminobutyl, and the like) or at the chain terminus (e.g., 2-hydroxyethyl, 3-cyanopropyl, and the like). Substituted substituents can be straight chain, branched or cyclic arrangements of covalently bonded carbon or heteroatoms.

It is understood that the above definitions are not intended to include impermissible substitution patterns (e.g., methyl substituted with five fluoro groups or a halogen atom substituted with another halogen atom). Such impermissible substitution patterns are well known to the skilled artisan.

“Syn” as used herein has its ordinary meaning, and is used in connection with Formula I to indicate that the specified groups are attached to sp³ hybridized (tetrahedral) carbon centers and extend out from one face of the cyclohexyl or piperidinyl ring, i.e., those groups all project toward the ‘alpha’ face of the ring, or they all project toward the ‘beta’ face of the ring. This is thus used as a convenient way to define the relative orientations of two or more groups, without limiting the compounds to a specific chiral configuration. This reflects the fact that the compounds of the invention have such groups in a specific relative orientation, but are not limited to either enantiomer of that specific relative orientation. Accordingly, unless described as optically active, such compounds may be racemic, but also include each of the two enantiomers having the specified relative stereochemistry. In some embodiments, the compounds of the invention are optically active form as further described herein, and in preferred embodiments of the invention, the compounds are obtained and used in optically active form. Preferably, the enantiomer having greater potency as an inhibitor of at least two of Pim1, Pim2 and Pim3 is selected.

It will also be apparent to those skilled in the art that the compounds of the invention, as well as the pharmaceutically acceptable salts, esters, metabolites and prodrugs of any of them, may be subject to tautomerization and may therefore exist in various tautomeric forms wherein a proton of one atom of a molecule shifts to another atom and the chemical bonds between the atoms of the molecules are consequently rearranged. See, e.g., March, *Advanced Organic Chemistry: Reactions, Mechanisms and Structures*, Fourth Edition, John Wiley & Sons, pages 69-74 (1992). As used herein, the term “tautomer” refers to the compounds produced by the proton shift, and it should be understood that all tautomeric forms, insofar as they may exist, are included within the invention.

The compounds of the invention comprise one or more asymmetrically substituted carbon atoms. Such asymmetrically substituted carbon atoms can result in the compounds of the invention existing in enantiomers, diastereomers, and other stereoisomeric forms that may be defined, in terms of absolute stereochemistry, such as in (R)- or (S)-forms. The compounds of the invention are sometimes depicted herein as single enantiomers, and are intended to encompass the specific configuration depicted and the enantiomer of that specific configuration (the mirror image isomer of the depicted configuration), unless otherwise specified. The depicted structures herein describe the relative stereochemistry of the compounds where two or more chiral centers, but the invention is not limited to the depicted enantiomer’s absolute stereochemistry unless otherwise stated.

The invention includes both enantiomers, each of which will exhibit Pim inhibition, even though one enantiomer will be more potent than the other. In some instances, compounds of the invention have been synthesized in racemic form and separated into individual isomers by chiral chromatography or similar conventional methods, and the analytical data about the two enantiomers do not provide definitive information about absolute stereochemical configuration. In such cases, the absolute stereochemistry of the most active enantiomer has been identified based on correlation with similar compounds of known absolute stereochemistry, rather than by a definitive physical method such as X-ray crystallography. Therefore, in certain embodiments, the preferred enantiomer of a compound described herein is the specific isomer depicted or its opposite enantiomer, whichever has the lower IC₅₀ for Pim kinase inhibition using the assay methods described herein, i.e., the enantiomer that is more potent as a Pim inhibitor for at least two of Pim1, Pim2, and Pim3.

The terms “S” and “R” configuration, as used herein, are as defined by the IUPAC 1974 RECOMMENDATIONS FOR SECTION E, FUNDAMENTAL STEREOCHEMISTRY, *Pure Appl. Chem.* 45:13-30 (1976). The terms α and β are employed for ring positions of cyclic compounds. The α -side of the reference plane is that side on which the preferred substituent lies at the lower numbered position. Those substituents lying on the opposite side of the reference plane are assigned β descriptor. It should be noted that this usage differs from that for cyclic stereoparents, in which “ α ” means “below the plane” and denotes absolute configuration. The terms α and β configuration, as used herein, are as defined by the CHEMICAL ABSTRACTS INDEX GUIDE-APPENDIX IV (1987) paragraph 203.

As used herein, the term “pharmaceutically acceptable salts” refers to the nontoxic acid or base addition salts of the compounds of Formula I or II, wherein the compound acquires a positive or negative charge as a result of adding or removing a proton; the salt then includes a counterion of opposite charge from the compound itself, and the counterion is preferably one suitable for pharmaceutical administration under the conditions where the compound would be used. These salts can be prepared in situ during the final isolation and purification of the compounds of Formula I or II, or by separately reacting the base or acid functions with a suitable organic or inorganic acid or base, respectively. Representative salts include but are not limited to the following: acetate, adipate, alginate, citrate, aspartate, benzoate, benzene-sulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, cyclopentanepropionate, dodecylsulfate, ethanesulfonate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate and undecanoate.


Also, a basic nitrogen-containing group in compounds of the invention can be quaternized with such agents as lower-alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides, and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl, and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides, and others. Water or oil-soluble or dispersible products are thereby obtained. These quaternized ammonium salts when paired with a pharmaceutically acceptable anion can also serve as pharmaceutically acceptable salts.

or a pharmaceutically acceptable salt thereof.


13

In some embodiments of the compounds of Formula (A), R^{1c} or R^{1d} or both R^{1c} and R^{1d} represent halo, preferably F. Typically, one of R^1 and R^{1b} , but not both, represents H. In some embodiments, R^1 is H and R^{1b} is halo or CONR'_2 , such as $-\text{CONHR}'$ where R' is C_{1-4} alkyl. In other embodiments, R^{1b} is H, in which case R^1 is preferably selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{4-6} heterocyclyl, $-(\text{CR}'_2)_{1-3}-\text{OR}'$ and $-\text{OR}'$, where each R' is independently H or C_{1-4} alkyl, and each alkyl, cycloalkyl and heterocyclyl is optionally substituted with up to two groups selected from halo, CN, NH_2 , hydroxy, and C_{1-4} alkoxy.

In preferred embodiments of the compounds of Formula (A), Q is CH. In many embodiments of these compounds, Ring A is a pyridine or a thiazole ring. Typically, these compounds are of this formula:




where Z^3 is H or NH_2 ,
or of this formula



wherein Z^3 is H or NH_2 ,
and R^1 , R^{1b} , R^{1c} , R^{1d} , R^{2a} and R^{2b} are as defined above for any of the embodiments of compounds of Formula (A).


Specific embodiments of these compounds include Formula A2 and A3:



14

-continued

(A4)




where each Z^2 is H or amino, Y is halo or H, and R^1 , R^{1b} , R^{1c} and R^{2a} are as defined above for any of the embodiments of compounds of Formula (A). In some embodiments, for example, R^{1c} is F, and R^{1b} is H.

In some of these embodiments, R^{2a} is preferably methoxy. In other embodiments wherein R^{2b} is H, R^{2a} is $-\text{OMe}$ or $-\text{O}(\text{CH}_2)_2-\text{X}$, wherein X is $-\text{OMe}$, COOH , CN or $-\text{SO}_2\text{Me}$, or R^{2a} is 1-triazolyl (e.g., 1,2,3-triazolyl) that is optionally substituted with up to two groups selected from halo, C_{1-4} alkyl, hydroxy-substituted C_{1-4} alkyl, C_{2-4} alkenyl, and C_{1-4} alkoxy, or $-\text{SO}_2\text{Me}$.

In some embodiments of these compounds of Formula (A), A1, A2, A3 or A4, R^{2a} is $-\text{OMe}$ or $-\text{O}(\text{CH}_2)_2-\text{X}$, wherein X is $-\text{OMe}$, COOH , CN or $-\text{SO}_2\text{Me}$; preferably R^{2a} is $-\text{OMe}$ or $-\text{OCH}_2\text{CH}_2\text{CN}$. R^1 in some embodiments of these compounds can be selected from 2-hydroxy-2-propyl, methoxymethyl, ethoxymethyl, methoxy, ethoxy, isopropoxy, cyclopropyl, cyclobutyl, 1-hydroxycyclobutyl, cyclopentyl, tetrahydropyranyl, 4-f, 4-hydroxy-4-tetrahydropyranyl, 4-tetrahydropyranyloxy, and 4-tetrahydropyranyl.

In certain embodiments, the compounds of Formula (A) have the Formula (I):



R^1 is selected from H, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{4-6} heterocyclyl, $-(\text{CR}'_2)_{1-3}-\text{OR}'$ and $-\text{OR}'$,

where each R' is independently H or C_{1-4} alkyl, and each alkyl, cycloalkyl and heterocyclyl is optionally substituted with up to two groups selected from halo, CN, NH_2 , hydroxy, oxo, C_{1-4} haloalkyl, C_{1-4} alkoxy, and C_{1-4} haloalkoxy;

one of R^{2a} and R^{2b} is H, and the other of R^{2a} and R^{2b} is selected from CN, halo, azido, amino, $-\text{OR}$, $-\text{O}(\text{CH}_2)_{1-3}\text{OR}$, $-\text{NRC}(\text{O})\text{R}$, $-\text{NRC}(\text{O})\text{OR}$, $-\text{NHSO}_2\text{R}$, $-\text{SO}_2\text{R}$, $-\text{OSO}_2\text{R}$, $-\text{SR}$, $-\text{S}(\text{O})\text{R}$, $-\text{OP}(\text{O})\text{R}_2$, and triazolyl optionally substituted with up to two groups selected from halo, C_{1-4} alkyl, hydroxy-substituted C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, and C_{5-6} heteroaryl;

15

where each R is independently C₁₋₄ alkyl optionally substituted with up to three groups selected from cyano, halo, hydroxy, carboxy, C₁₋₄ alkylsulfonyl, and C₁₋₄ alkoxy;

or R^{2a} and R^{2b} taken together form a dialkyl ketal or 5-6 membered cyclic ketal, =O or =N—OR", where R" is H or C₁₋₄ alkyl;


ring A is selected from pyridinyl, pyrimidinyl, pyrazinyl, and thiazolyl, each having N positioned as shown in Formula (I); and

Ring A is optionally substituted with 1 or 2 groups selected from halo, CN, NH₂, hydroxy, C₁₋₄ alkyl, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, and C₁₋₄ haloalkoxy;

or a pharmaceutically acceptable salt thereof.

The following enumerated embodiments represent additional aspects and variations of the invention:

1. A compound of Formula (A)



wherein:

groups attached to the ring containing Q that are depicted inside the ring are all syn to each other, and all groups attached to that ring that are depicted outside the ring are syn to one another;

Q is C or N;

R^{2c} is H when Q is C, and R^{2c} is absent when Q is N;

R¹ is selected from H, C₁₋₄ alkyl, C₃₋₆ cycloalkyl, C₄₋₆ heterocyclyl, —(CR'₂)₁₋₃—OR' and —OR',

where each R' is independently H or C₁₋₄ alkyl,

and each alkyl, cycloalkyl and heterocyclyl is optionally substituted with up to two groups selected from halo, CN, NH₂, hydroxy, oxo, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, and C₁₋₄ haloalkoxy;

R^{1b}, R^{1c}, and R^{1d} are each independently selected from H, halo, OR', R', —(CH₂)₁₋₃OR', and CONR'₂;

one of R^{2a} and R^{2b} is H,

and the other of R^{2a} and R^{2b} is selected from CN, halo, azido, amino, —OR, —O(CH₂)₁₋₃OR, —NRC(O)R, —NRC(O)OR, —NHSO₂R, —SO₂R, —OSO₂R, —SR, —S(O)R, —OP(O)R₂, and 1-pyridonyl or 1-triazolyl optionally substituted with up to two groups selected from halo, C₁₋₄ alkyl, hydroxy-substituted C₁₋₄ alkyl, C₂₋₄ alkenyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, and C₅₋₆ heteroaryl;

where each R is independently C₅₋₆ heteroaryl or C₁₋₄ alkyl optionally substituted with up to three groups selected from cyano, halo, hydroxy, carboxy, C₁₋₄ alkylsulfonyl, and C₁₋₄ alkoxy;

or R^{2a} and R^{2b} taken together may form a dialkyl ketal or 5-6 membered cyclic ketal, =O or =N—OR", where R" is H or C₁₋₄ alkyl;

ring A is selected from pyridinyl, pyrimidinyl, pyrazinyl, and thiazolyl, each having N positioned as shown in Formula (I); and

16

Ring A is optionally substituted with 1 or 2 groups selected from halo, CN, NH₂, hydroxy, C₁₋₄ alkyl, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, and C₁₋₄ haloalkoxy;

or a pharmaceutically acceptable salt thereof

2. The compound of Formula (A) according to embodiment 1, wherein one but not both, of R^{1b} and R¹ represents H.

3. The compound of embodiment 1, wherein R^{1d} is F.

4. The compound of embodiment 1 or 2, wherein R^{1c} is F.

5. The compound of any of embodiments 1-3, wherein R^{1b} is H.

6. The compound of any of embodiments 1-3, wherein R^{1b} is H or CONR'₂.


7. The compound of any of embodiments 1-6, wherein R^{2b} is H.

8. The compound of any of embodiments 1-7, wherein R^{2a} is —OMe, —SO₂Me, —NHCOOMe, or —O(CH₂)₂—X, wherein X is —OMe, COOH, CN or —SO₂Me, or R^{2a} is 1-triazolyl (e.g., 1,2,3-triazolyl) or 1-pyridonyl that is optionally substituted with up to two groups selected from halo, C₁₋₄ alkyl, hydroxy-substituted C₁₋₄ alkyl, C₂₋₄ alkenyl, C₃₋₆ cycloalkyl, and C₁₋₄ alkoxy, or —SO₂Me.

9. The compound of any of embodiments 1-8, wherein R¹ is selected from 2-hydroxy-2-propyl, methoxymethyl, ethoxymethyl, methoxy, ethoxy, isopropoxy, cyclopropyl, cyclobutyl, 1-hydroxycyclobutyl, cyclopentyl, tetrahydropyranyl, 4-F, 4-hydroxy-4-tetrahydropyranyl, 4-tetrahydropyranyloxy, and 4-tetrahydropyranyl.

10. The compound of any of embodiments 1-6, wherein R^{2b} is OMe.

11. The compound of embodiment 1, which is a compound of Formula (I):



wherein:

groups attached to the ring containing Q that are depicted inside the ring are all syn to each other, and all groups attached to that ring that are depicted outside the ring are syn to one another;

Q is C or N;

R^{2c} is H when Q is C, and R^{2c} is absent when Q is N;

R¹ is selected from H, C₁₋₄ alkyl, C₃₋₆ cycloalkyl, C₄₋₆ heterocyclyl, —(CR'₂)₁₋₃—OR' and —OR',

where each R' is independently H or C₁₋₄ alkyl,

and each alkyl, cycloalkyl and heterocyclyl is optionally substituted with up to two groups selected from halo, CN, NH₂, hydroxy, oxo, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, and C₁₋₄ haloalkoxy;

one of R^{2a} and R^{2b} is H,

and the other of R^{2a} and R^{2b} is selected from CN, halo, azido, amino, —OR, —O(CH₂)₁₋₃OR, —NRC(O)R, —NRC(O)OR, —NHSO₂R, —SO₂R, —OSO₂R, —SR, —S(O)R, —OP(O)R₂, and N-pyridonyl or 1-triazolyl optionally sub-

17

stituted with up to two groups selected from halo, C₁₋₄ alkyl, hydroxy-substituted C₁₋₄ alkyl, C₂₋₄ alkenyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, and C₅₋₆ heteroaryl;

where each R is independently C₅₋₆ heteroaryl or C₁₋₄ alkyl optionally substituted with up to three groups selected from cyano, halo, hydroxy, carboxy, C₁₋₄ alkylsulfonyl, and C₁₋₄ alkoxy;

or R^{2a} and R^{2b} taken together may form a dialkyl ketal or 5-6 membered cyclic ketal, =O or =N—OR", where R" is H or C₁₋₄ alkyl;

Ring A is selected from pyridinyl, pyrimidinyl, pyrazinyl, and thiazolyl, each having N positioned as shown in Formula (I); and

Ring A is optionally substituted with 1 or 2 groups selected from halo, CN, NH₂, hydroxy, C₁₋₄ alkyl, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, and C₁₋₄ haloalkoxy;

or a pharmaceutically acceptable salt thereof

12. The compound of embodiment 11, wherein R^{2a} is H.

13. The compound of embodiment 11, wherein R^{2b} is H.

14. The compound of embodiment 11 or 13, wherein R^{2a} is —NHCOOMe.

15. The compound of embodiment 11 or 13, wherein R^{2a} is —O(CH₂)₂—CN.

16. The compound of embodiment 11 or 13, wherein R^{2a} is —O(CH₂)₂—SO₂Me.

17. The compound of embodiment 11 or 13, wherein R^{2a} is —OMe.

18. The compound of embodiment 11 or 13, wherein R^{2a} is —SO₂Me.

19. The compound of and embodiment 11 or 12, wherein R^{2b} is —OMe.

20. The compound of embodiment 11, wherein R^{2a} and R^{2b} taken together form =O or =N—OR", where R" is H or C₁₋₄ alkyl.

21. The compound of any of embodiments 11-20, wherein R¹ is selected from H, methyl, ethyl, isopropyl, 2-hydroxy-2-propyl, methoxymethyl, ethoxymethyl, methoxy, ethoxy, isopropoxy, cyclopropyl, cyclobutyl, 1-hydroxycyclobutyl, cyclopentyl, tetrahydrofuranyl, 4-fluoro-4-tetrahydrofuranyl, tetrahydrothiopyran, and 4-tetrahydrothiopyran-1,1-dioxide.

22. The compound of embodiment 21, wherein R¹ is selected from H, Me, tetrahydropyran, methoxymethyl, and ethoxymethyl.

23. The compound of any one of the preceding embodiments, wherein Ring A is pyridine, and is optionally substituted with F or amino.

24. The compound of any one of embodiments 11-21, wherein Ring A is a thiazolyl, optionally substituted with amino.

25. The compound of any of embodiments 11-24, wherein R^{2a} is selected from the group consisting of —OR, —OCH₂CH₂OR, —OCH₂CH₂CN, —OCH₂CH₂COOH, —OCH₂CH₂SO₂R, —CN, —NHC(O)OR, —NHC(O)R, N₃, NH₂, F, —NHSO₂R, —SO₂R, —SR, —S(O)R, unsubstituted triazole, and triazole substituted with Me, ethyl, cyclopropyl, hydroxymethyl, C₂₋₄ alkenyl, or thienyl;

wherein each R is methyl, ethyl or isopropyl.

26. The compound of any one of embodiments 11-24, wherein R^{2b} is selected from the group consisting of —OR, —OCH₂CH₂OR, —OCH₂CH₂CN, —OCH₂CH₂COOH, —OCH₂CH₂SO₂R, —CN, —NHC(O)OR, —NHC(O)R, N₃, NH₂, F, —NHSO₂R, —SO₂R, —SR, —S(O)R, unsubstituted triazole, and triazole substituted with Me, ethyl, cyclopropyl, hydroxymethyl, C₂₋₄ alkenyl, or thienyl;

wherein each R is methyl, ethyl or isopropyl.

27. The compound of embodiment 11, wherein R^{2a} is —SO₂Me, —OCH₂CH₂CN, 1,2,4-triazol-1-yl, —OCH₂CH₂COOH, —OCH₂CH₂SO₂R, —CN, or —NHC(O)OMe.

18

28. The compound of embodiment 12, wherein R^{2b} is —OMe, —CN, —OCH₂CH₂CN, 1,2,4-triazol-1-yl, —OCH₂CH₂COOH, —OCH₂CH₂SO₂R, —CN, or —NHC(O)OMe.

29. The compound of any of the preceding embodiments, wherein Q is C.



30. The compound of any of embodiments 1-28, wherein Q is N.

31. The compound of any of the preceding embodiments, which is optically active and has a lower IC₅₀ than its opposite enantiomer on at least one Pim kinase.

32. The compound of embodiment 31, wherein the at least one Pim kinase is Pim2 kinase.


33. The compound of any one of embodiments 11-32, which is an optically active compound of Formula IIa or IIb:

or




wherein Y is H or F, and Z is H or NH₂.

34. The compound of any one of embodiments 11-32, which is an optically active compound of Formula IIIa or IIIb:



19

-continued



wherein Y is H or F, and Z is H or NH₂.

35. The compound of embodiment 33, which is a compound of Formula IIa.

36. The compound of embodiment 33, which is a compound of Formula IIb.

37. The compound of embodiment 34, which is a compound of Formula IIa.

38. The compound of embodiment 34, which is a compound of Formula IIb.

39. A compound selected from the group consisting of the compounds in Table 1 and Table 2, and the pharmaceutically acceptable salts thereof

40. A pharmaceutical composition comprising a compound of any of embodiments 1-39, admixed with at least one pharmaceutically acceptable excipient.

41. The pharmaceutical composition of embodiment 40 which further comprises an additional agent for treatment of cancer.

42. The pharmaceutical composition of embodiment 41, wherein the additional therapeutic agent is selected from MEK inhibitors, irinotecan, topotecan, gemcitabine, 5-fluorouracil, cytarabine, daunorubicin, PI3 Kinase inhibitors, mTOR inhibitors, DNA synthesis inhibitors, leucovorin, carboplatin, cisplatin, taxanes, tezacitabine, cyclophosphamide, vinca alkaloids, imatinib, anthracyclines, rituximab, lenalidomide, bortezomib and trastuzumab

43. A compound of any of embodiments 1-39 for use in the treatment of a condition that responds to inhibitors of Protein Integration of Moloney Kinase (PIM Kinase) activity.

44. The compound according to embodiment 43, wherein the condition is a cancer.

45. The compound according to embodiment 44, wherein the cancer is selected from carcinoma of the lungs, pancreas, thyroid, ovaries, bladder, breast, prostate or colon, melanoma, myeloid leukemia, multiple myeloma, erythro leukemia, villous colon adenoma, and osteosarcoma.

46. The compound of embodiment 45, wherein the condition is an autoimmune disorder.

47. A method of treating a disease or condition mediated by PIM kinase, comprising administering to a subject in need thereof a therapeutically effective amount of a compound according to any one of embodiments 1-39, or a pharmaceutically acceptable salt thereof

48. The method of embodiment 47, wherein the disease is selected from carcinoma of the lungs, pancreas, thyroid, ovaries, bladder, breast, prostate or colon, melanoma, myeloid leukemia, multiple myeloma, erythro leukemia, villous colon adenoma, and osteosarcoma; or the disease is an autoimmune disorder.

49. The method of embodiment 48, wherein the disease is an autoimmune disorder.

20


50. The method of embodiment 49, wherein the autoimmune disorder is selected from Crohn's disease, inflammatory bowel disease, rheumatoid arthritis, and chronic inflammatory diseases.

5 In some embodiments, at least one substituent for Ar is selected from F, Cl, NH₂, Me, Et, OMe, OEt, OCF₃, OCHF₂, OCH₂CF₃, CN, CF₃, SMe, SOMe, SO₂Me, —COOMe, —C(O)Me, —C(Me)₂-OH, MeOCH₂—, HOCH₂—, hydroxyethyl, hydroxyethoxy, methoxyethyl, methoxyethoxy, oxetanyl (e.g., 3-oxetanyl), isopropoxy, tetrahydropyranyloxy (e.g., 4-tetrahydropyranyloxy), cyclopropyl, and CN. At least one substituent for Ar is preferably selected from Me, F, NH₂, OMe, MeOCH₂—, HOCH₂—, hydroxyethyl, hydroxyethoxy, methoxyethyl, methoxyethoxy, and CN.


15 These compounds may be used in racemic form, or the individual enantiomers may be used, or mixtures of the enantiomers may be used. Each enantiomer can be used, and preferably the compound to be used is the enantiomer that has greater activity as a Pim inhibitor.

20 The cyclohexyl or piperidine ring in these compounds has three non-hydrogen substituents, not counting its attachment to the pyridinyl ring in Formula I. The invention provides novel combinations of substituents and their relative stereochemical orientation on the cyclohexyl or piperidine ring, to provide advantageous biological activities. Advantages provided by preferred compounds include reduced drug-drug interactions, due to reduction of time-dependent Cyp inhibition or pharmacokinetic superiority based on improved clearance and metabolic properties.

30 In one preferred embodiment of the claimed invention, the compound is of formula IIc or IId,



(IIc)





(IId)

65 wherein Y is H or F, and Z² is H or NH₂; or. In these compounds, R^{2a} is preferably —OR, —O(CH₂)₁₋₃OR, —SO₂R, or —NRC(O)OR, where each R is independently C₁₋₄ alkyl optionally substituted with cyano, halo, hydroxy, carboxy, C₁₋₄ alkylsulfonyl, or C₁₋₄ alkoxy; and R^{2b} is H. In

21

some such embodiments, R^{2a} is preferably —OR, —O(CH₂)₁₋₃OR, or —NRC(O)OR, where each R is independently C₁₋₄ alkyl optionally substituted with cyano, halo, hydroxy, carboxy, C₁₋₄ alkylsulfonyl, or C₁₋₄ alkoxy; and R^{2b} is H. In some preferred embodiments, R^{2a} is OMe. In these compounds, R^1 can be H, Me, 4-tetrahydropyranyl, or 2-hydroxy-2-propyl. In specific embodiments, R^{2a} is —OMe, —OCH₂CH₂CN, —OCH₂CH₂OMe, or —OCH₂CH₂SO₂Me or —NHCOOMe, or —N(Me)COOMe. Preferably when R^{2b} is H, R^1 is not H.

In another embodiment, the compound is of Formula IIe or IIf:



wherein Y is H or F, and Z^2 is H or NH₂. In these compounds, R^{2a} is preferably —OR, —O(CH₂)₁₋₃OR, —SO₂R, or —NRC(O)OR, where each R is independently C₁₋₄ alkyl optionally substituted with cyano, halo, hydroxy, carboxy, C₁₋₄ alkylsulfonyl, or C₁₋₄ alkoxy; and R^{2b} is H. In some such embodiments, R^{2a} is —OR, —O(CH₂)₁₋₃OR, or —NRC(O)OR, where each R is independently C₁₋₄ alkyl optionally substituted with cyano, halo, hydroxy, carboxy, C₁₋₄ alkylsulfonyl, or C₁₋₄ alkoxy; and R^{2b} is H. In some preferred embodiments, R^{2a} is OMe. In these compounds, R^1 can be H, Me, 4-tetrahydropyranyl, or 2-hydroxy-2-propyl. In specific embodiments, R^{2a} is —OMe, —OCH₂CH₂CN, —OCH₂CH₂OMe, or —OCH₂CH₂SO₂Me or —NHCOOMe, or —N(Me)COOMe. Preferably when R^{2b} is H, R^1 is not H.

Each of the species in Tables 1 and 2 are preferred embodiments of the invention.

For purposes of the present invention, a therapeutically effective dose will generally be a total daily dose administered to a host in single or divided doses may be in amounts, for example, of from 0.001 to 1000 mg/kg body weight daily, typically 0.01 to 10 mg/kg per day, and more preferred from 0.1 to 30 mg/kg body weight daily. Generally, daily dosage amounts of 1 to 4000 mg, or from 5 to 3000, or from 10 to

22

2000 mg, or from 100 to 2000 mg are anticipated for human subjects. Dosage unit compositions may contain such amounts of submultiples thereof to make up the daily dose.

The compounds of the present invention may be administered orally, parenterally, sublingually, by aerosolization or inhalation spray, rectally, or topically in dosage unit formulations containing conventional nontoxic pharmaceutically acceptable carriers, adjuvants, and vehicles as desired. Topical administration may also involve the use of transdermal administration such as transdermal patches or ionophoresis devices. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection, or infusion techniques.

Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-propanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or di-glycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

Suppositories for rectal administration of the drug can be prepared by mixing the drug with a suitable nonirritating excipient such as cocoa butter and polyethylene glycols, which are solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum and release the drug.

Solid dosage forms for oral administration may include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound may be admixed with at least one inert diluent such as sucrose lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

Liquid dosage forms for oral administration may include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions may also comprise adjuvants, such as wetting agents, emulsifying and suspending agents, cyclodextrins, and sweetening, flavoring, and perfuming agents.

The compounds of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multilamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the phospholipids and phosphatidylcholines (lecithins), both natural and synthetic. Methods to form liposomes are known in the art. See, for example, Prescott, Ed., *Methods in Cell Biology*, Volume XIV, Academic Press, New York, N.Y., p. 33 et seq. (1976).

While the compounds of the invention can be administered as the sole active pharmaceutical agent, they can also be used in combination with one or more other agents used in the treatment of cancer. The compounds of the present invention are also useful in combination with known therapeutic agents and anti-cancer agents, and combinations of the presently disclosed compounds with other anti-cancer or chemothera-

peutic agents are within the scope of the invention. Examples of such agents can be found in *Cancer Principles and Practice of Oncology*, V. T. Devita and S. Hellman (editors), 6th edition (Feb. 15, 2001), Lippincott Williams & Wilkins Publishers. A person of ordinary skill in the art would be able to discern which combinations of agents would be useful based on the particular characteristics of the drugs and the cancer involved. Such anti-cancer agents include, but are not limited to, the following: MEK inhibitors, estrogen receptor modulators, androgen receptor modulators, retinoid receptor modulators, cytotoxic/cytostatic agents, antiproliferative agents, prenyl-protein transferase inhibitors, HMG-CoA reductase inhibitors and other angiogenesis inhibitors, inhibitors of cell proliferation and survival signaling, apoptosis inducing agents and agents that interfere with cell cycle checkpoints. The compounds of the invention are also useful when co-administered with radiation therapy.

Therefore, in one embodiment of the invention, the compounds of the invention are also used in combination with known therapeutic or anticancer agents including, for example, estrogen receptor modulators, androgen receptor modulators, retinoid receptor modulators, cytotoxic agents, antiproliferative agents, prenyl-protein transferase inhibitors, HMG-CoA reductase inhibitors, HIV protease inhibitors, reverse transcriptase inhibitors, and other angiogenesis inhibitors.

In certain presently preferred embodiments of the invention, representative therapeutic agents useful in combination with the compounds of the invention for the treatment of cancer include, for example, MEK inhibitors, irinotecan, topotecan, gemcitabine, 5-fluorouracil, cytarabine, daunorubicin, PI3 Kinase inhibitors, mTOR inhibitors, DNA synthesis inhibitors, leucovorin carboplatin, cisplatin, taxanes, tezacitabine, cyclophosphamide, vinca alkaloids, imatinib (Gleevec), anthracyclines, rituximab, trastuzumab, Revlimid, Velcade, dexamethasone, daunorubicin, cytarabine, clofarabine, Mylotarg, lenalidomide, bortezomib, as well as other cancer chemotherapeutic agents including targeted therapeutics.

The above compounds to be employed in combination with the compounds of the invention will be used in therapeutic amounts as indicated in the *Physicians' Desk Reference (PDR)* 47th Edition (1993), which is incorporated herein by reference, or such therapeutically useful amounts as would be known to one of ordinary skill in the art, or provided in prescribing materials such as a drug label for the additional therapeutic agent.

The compounds of the invention and the other anticancer agents can be administered at the recommended maximum clinical dosage or at lower doses. Dosage levels of the active

compounds in the compositions of the invention may be varied so as to obtain a desired therapeutic response depending on the route of administration, severity of the disease and the response of the patient. The combination can be administered as separate compositions or as a single dosage form containing both agents. When administered as a combination, the therapeutic agents can be formulated as separate compositions, which are given at the same time or different times, or the therapeutic agents, can be given as a single composition.


In one embodiment, the invention provides a method of inhibiting Pim1, Pim2 or Pim3 in a human or animal subject. The method includes administering an effective amount of a compound, or a pharmaceutically acceptable salt thereof, of any of the embodiments of compounds of Formula I or II to a subject in need thereof.

The present invention will be understood more readily by reference to the following examples, which are provided by way of illustration and are not intended to be limiting of the present invention.

Synthetic Methods


The compounds of the invention can be obtained through procedures known to those skilled in the art. As shown in Scheme 1, 5-alkyl, 4-hydroxy, 3-aminopiperidines can be prepared and modified to yield 5-alkyl, 4-substituted, 3-aminopiperidinyl pyridine amides VI as follows. Reaction of Garner's aldehyde with (R)-4-benzyl-3-propionyloxazolidin-2-one followed by TBS protection of the resulting alcohol affords compound I. Reduction of the oxazolidinone followed by introduction of the azide group yields intermediate II. Deprotection under acidic conditions reveals the corresponding amino alcohol, which upon protection with the Boc group followed by mesylation of the primary alcohol yields intermediate III. Reduction of the azide affords formation of the piperidine which is subsequently reacted with 4-chloro-3-nitropyridine and following nitro reduction pyridyl aniline IV is obtained. Aniline IV can be coupled with heterocyclic acids, which after silyl group deprotection (Va) and modification of the hydroxyl (activation as mesylate, displacement and potentially further modification of the displaced group) and Boc deprotection can afford target amides VI. Alternatively, aniline IV can be Boc protected, silyl deprotected (Vb), modified at the hydroxyl position and after aniline Boc deprotection (Vc), amide coupling and Boc deprotection, 5-alkyl, 4-substituted, 3-aminopiperidinyl pyridine amides VI can be obtained.

Scheme 1




25

26




In Scheme 2, synthetic methods to prepare tetrasubstituted aminocyclohexylpyridyl amides X are depicted. Methyl cyclohexanone can be converted via the monotriflate to the corresponding cyclohexenoneboronate ester which can undergo palladium mediated carbon bond formation with 4-chloro, 3-nitro pyridine to yield nitropyridine substituted cyclohexenone VII. Ketone reduction followed by dehydration yields a cyclohexadiene which upon epoxidation (via bromohydrin formation and HBr elimination), azide epoxide opening, azide reduction and amine Boc protection yields cyclohexenyl Boc amino alcohol nitro pyridyl compound VIII. Nitro pyridyl VIII can be converted to the trans protected amino hydroxy aniline IXa by alcohol protection and alkene and nitro reduction. Alternatively, the alcohol moiety of nitropyridyl VIII can be inverted via a mesylation, cyclization, Boc protection and hydrogenation sequence to provide the all cis substituted cyclohexyl pyridyl aniline IXb, where the cis hydroxy is protected in the form of a cyclic carbamate. As described above in Scheme 1 for the preparation of substituted piperidine compounds of the invention, upon amide coupling of the cyclohexyl pyridyl anilines IXa or IXb to heterocyclic acids and subsequent hydroxyl and amine deprotection, substituted cyclohexyl compounds of the invention Xa and Xb can be prepared.

Scheme 2




27

-continued




28


-continued



29
-continued






30
-continued




In Scheme 3, additional synthetic methods to prepare tetrasubstituted aminocyclohexylpyridyl amides X are depicted in which the cyclohexyl systems are modified prior to coupling of the aniline to heterocyclic acids. For example, aniline IXa can be Boc protected, acetyl deprotected, modified at the hydroxyl position and after a net aniline Boc deprotection, amide coupling and Boc deprotection, substituted aminocyclohexylpyridyl amides Xa can be obtained. In an alternative manner, the cyclohexenyl Boc amino alcohol nitro pyridyl compound VIII can be modified at the hydroxyl position and after alkene and nitro reduction, amide coupling and Boc deprotection, substituted aminocyclohexylpyridyl amides Xc can be obtained. Additionally, the hydroxyl group of VIII can be inverted via mesylation, intramolecular cyclization, Boc protection and cyclic carbamate opening to yield an alcohol that after processing as described above can yield substituted aminocyclohexylpyridyl amides Xd. For the sequences depicted in Schemes 1-3, the aliphatic amine is primarily protected as a Boc derivative. As one skilled in the art would expect, alternative protecting groups and subsequent deprotection conditions for the amine and hydroxy moieties can be utilized.

Scheme 3



31

-continued




A representative route to a heterocyclic acid that can be incorporated into compounds VI, Xa-Xd of the invention is depicted in Scheme 4. Lithiation of 5-methoxy, 1,3-difluorobenzene and reaction with 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane yields a boronate ester which upon

32

Suzuki coupling with the bromofluoro picolinate ester and subsequent ester hydrolysis yields the fluoro picolinic acid XI.

Scheme 4



EXAMPLES

Referring to the examples that follow, compounds of the preferred embodiments were synthesized using the methods described herein, or other methods, which are known in the art.

The compounds and/or intermediates were characterized by high performance liquid chromatography (HPLC) using a Waters Millenium chromatography system with a 2695 Separation Module (Milford, Mass.). The analytical columns were reversed phase Phenomenex Luna C18-5 μ , 4.6 $\times$ 50 mm, from Alltech (Deerfield, Ill.). A gradient elution was used (flow 2.5 mL/min), typically starting with 5% acetonitrile/95% water and progressing to 100% acetonitrile over a period of 10 minutes. All solvents contained 0.1% trifluoroacetic acid (TFA). Compounds were detected by ultraviolet light (UV) absorption at either 220 or 254 nm. HPLC solvents were from Burdick and Jackson (Muskegan, Mich.), or Fisher Scientific (Pittsburgh, Pa.).

In some instances, purity was assessed by thin layer chromatography (TLC) using glass or plastic backed silica gel plates, such as, for example, Baker-Flex Silica Gel 1B2-F flexible sheets. TLC results were readily detected visually under ultraviolet light, or by employing well-known iodine vapor and other various staining techniques.

Mass spectrometric analysis was performed on one of three LCMS instruments: a Waters System (Alliance HT HPLC

33

and a Micromass ZQ mass spectrometer; Column: Eclipse XDB-C18, 2.1×50 mm; gradient: 5-95% (or 35-95%, or 65-95% or 95-95%) acetonitrile in water with 0.05% TFA over a 4 min period; flow rate 0.8 mL/min; molecular weight range 200-1500; cone Voltage 20 V; column temperature 40° C.), another Waters System (ACQUITY UPLC system and a ZQ 2000 system; Column: ACQUITY UPLC HSS-C18, 1.8 um, 2.1×50 mm; gradient: 5-95% (or 35-95%, or 65-95% or 95-95%) acetonitrile in water with 0.05% TFA over a 1.3 min period; flow rate 1.2 mL/min; molecular weight range 150-850; cone Voltage 20 V; column temperature 50° C.) or a Hewlett Packard System (Series 1100 HPLC; Column: Eclipse XDB-C18, 2.1×50 mm; gradient: 5-95% acetonitrile in water with 0.05% TFA over a 4 min period; flow rate 0.8 mL/min; molecular weight range 150-850; cone Voltage 50 V; column temperature 30° C.). All masses were reported as those of the protonated parent ions.

Nuclear magnetic resonance (NMR) analysis was performed on some of the compounds with a Varian 400 MHz NMR (Palo Alto, Calif.). The spectral reference was either TMS or the known chemical shift of the solvent.

Preparative separations are carried out using a Flash 40 chromatography system and KP-Sil, 60A (Biotage, Charlottesville, Va.), or by flash column chromatography using silica gel (230-400 mesh) packing material on ISCO or Analogix purification systems, or by HPLC using a Waters 2767 Sample Manager, C-18 reversed phase column, 30×50 mm, flow 75 mL/min. Typical solvents employed for the Flash 40 Biotage, ISCO or Analogix system for silica gel column chromatography are dichloromethane, methanol, ethyl acetate, hexane, n-heptanes, acetone, aqueous ammonia (or ammonium hydroxide), and triethyl amine. Typical solvents employed for the reverse phase HPLC are varying concentrations of acetonitrile and water with 0.1% trifluoroacetic acid.

Chiral separations of enantiomeric mixtures were performed by the following analytical and preparative general methods:

Chiral SFC-Analytical Method: Chiral compounds were separated on a Waters Supercritical Fluid Chromatography (SFC). The separation used a Chiralpak AD (AS, OD, OJ, IC or IA) 4.6×100 mm column at 40 C temperature at a flow rate of 5 mL/min using an isocratic method. The mobile phase was 15% MeOH (or EtOH or IPA or with 0.1% Diethyl amine): 85% CO₂. The detection wavelength was 220 nm (or 250 nm or Diode Array)

Chiral SFC-Purification Method: Chiral compounds were separated on a Waters Supercritical Fluid Chromatography (SFC). The separation used a Chiralpak AD (AS, OD, OJ, IC or IA) 21×250 mm column at 40 C temperature at a flow rate of 100 mL/min using an isocratic method. The mobile phase was 15% MeOH (or EtOH or IPA or with 0.1% Diethyl amine): 85% CO₂. The detection wavelength was 220 nm (or 250 nm or Diode Array).

Chiral HPLC-Analytical Method: Chiral compounds were separated on a Waters 2695 HPLC system. The separation used a Chiralpak AD (AS, OD, OJ, IC or IA) 4.6×100 mm column at room temperature at a flow rate of 1 mL/min using an isocratic method. The mobile phase was 15% EtOH (or IPA or with 0.1% Diethyl amine): 85% Heptane. The detection wavelength was 220 nm (or 250 nm or Diode Array).

Chiral HPLC-Purification Method: Chiral compounds were separated on a Waters 2767 HPLC system. The separation used a Chiralpak AD (AS, OD, OJ, IC or IA) 21×250 mm column at room temperature at a flow rate of 20 (or 10-15) mL/min using an isocratic method. The mobile phase was

34

15% EtOH (or IPA or with 0.1% Diethyl amine): 85% Heptane. The detection wavelength was 220 nm (or 250 nm or Diode Array).

It should be understood that the organic compounds according to the preferred embodiments may exhibit the phenomenon of tautomerism. As the chemical structures within this specification can only represent one of the possible tautomeric forms, it should be understood that the preferred embodiments encompasses any tautomeric form of the drawn structure.

It is understood that the invention is not limited to the embodiments set forth herein for illustration, but embraces all such forms thereof as come within the scope of the above disclosure.

The examples below as well as throughout the application, the following abbreviations have the following meanings. If not defined, the terms have their generally accepted meanings.


ABBREVIATIONS

BINAP	2,2'-Bis(diphenylphosphino)-1,1'-binaphthalene
Boc ₂ O	di-tert-butyl dicarbonate
Boc-OSu	N-(tert-Butoxycarbonyloxy)succinimide
Cbz-OSu	N-(Benzyloxycarbonyloxy)succinimide
DAST	(diethylamino)sulfurtrifluoride
DBU	1,8-Diazabicyclo[5.4.0]undec-7-ene
DCM	Dichloromethane
DIAD	diisopropylazodicarboxylate
DIEA	diisopropylethylamine
DMA	Dimethylacetamide
DMAP	4-dimethylaminopyridine
DME	1,2-dimethoxyethane
DMF	N,N-dimethylformamide
DPPF	1,1'-bis(diphenylphosphino)ferrocene
EDC	1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
EtOAc	ethyl acetate
EtOH	Ethanol
HOAT	Hydroxyazabenzotriazole
K ₂ CO ₃	Potassium carbonate
KSAc	Potassium thioacetate
Lawesson's reagent	2,4-bis(4-methoxyphenyl)-1,3,2,4-dithiadiphosphetane-2,4-dithione
LiOH	Lithium hydroxide
MCPBA	Meta-chloroperoxybenzoic acid
MeCN	Acetonitrile
methylDAST	(dimethylamino)sulfurtrifluoride
MgSO ₄	Magnesium sulfate
MeOH	Methanol
MsCl	Methane sulfonyl chloride
Na ₂ CO ₃	sodium carbonate
NaCl	Sodium chloride
NaHCO ₃	sodium bicarbonate
NaHMDS	Sodium bis(trimethylsilyl)amide
NBS	N-bromosuccinimide
NMP	N-methyl-2-pyrrolidone
oxone	Potassium peroxymonosulfate
p-TSA	para-toluene sulfonic acid
Pd ₂ (dba) ₃	Tris(dibenzylideneacetone)dipalladium(0)
Pd(PPh ₃) ₄	Tetrakis(triphenylphosphine)palladium(0)
Pd(dppf)Cl ₂	Dichloro-(1,2-bis(diphenylphosphino)ethane)-
DCM	Palladium(II)—dichloromethane adduct
RT or rt	room temperature
TBAF	Tetrabutyl ammonium fluoride
TBDMSCl	tert-butyl dimethylsilylchloride
TEA	Triethylamine
THF	tetrahydrofuran

35


EXAMPLES

Synthesis of 6-bromo-5-fluoropicolinic acid



To 2-bromo-3-fluoro-6-methylpyridine (1.0 equiv.) in H₂O (30 mL) was added potassium permanganate (1.0 equiv.). The solution was heated at 100° C. for 5 hours at which time more potassium permanganate (1.0 equiv.) was added. After heating for an additional 48 hours the material was filtered through celite (4 cm×2 inches) and rinsed with H₂O (150 mL). The combined aqueous was acidified with 1N HCl to pH=4, extracted with ethyl acetate (200 mL), washed with NaCl(sat.), dried over MgSO₄, filtered and concentrated to yield 6-bromo-5-fluoropicolinic acid (17%) as a white solid. LCMS (m/z): 221.9 (MH⁺); LC Rt=2.05 min.


Synthesis of methyl 6-bromo-5-fluoropicolinate



To a solution of 6-bromo-5-fluoropicolinic acid (1.0 equiv.) in methanol (0.2 M) was added H₂SO₄ (4.2 equiv.) and the reaction was stirred at room temperature for two hours. Upon completion of the reaction as monitored by LC/MS, the reaction was diluted with ethyl acetate and quenched slowly with saturated aqueous NaHCO₃. The reaction was poured into a separatory funnel and extracted with ethyl acetate. The organic phase was dried with magnesium sulfate, filtered, and concentrated in vacuo to provide methyl 6-bromo-5-fluoropicolinate as a white solid (>99%). LC/MS=233.9/235.9 (M+H), Rt=0.69 min.

Method 1

Synthesis of methyl 6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinate




36

To a solution of methyl 6-bromo-5-fluoropicolinate (1.0 equiv.) in THF and water (10:1, 0.1 M) was added 2,6-difluoro-4-methoxyphenylboronic acid (2.5 equiv.) and potassium fluoride (3.3 equiv.). The reaction was degassed with nitrogen, then Pd₂(dba)₃ (0.25 equiv.) and tri-tert-butylphosphine (0.5 equiv.) were added and the reaction was heated to 80° C. for one hour. LC/MS analysis indicated complete conversion of the starting material to product. The reaction was cooled to room temperature, then concentrated in vacuo and fused to silica gel. The crude product was purified by ISCO flash chromatography eluting with ethyl acetate and hexanes (0% to 30% ethyl acetate) to provide methyl 6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinate as a white solid in 85% yield. LC/MS=298.0 (M+H), Rt=0.89 min.

Method 2


Synthesis of 6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinic acid



To a solution of methyl 6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinate (1.0 equiv.) in THF/MeOH (2:1, 0.09 M) was added LiOH (1.5 equiv.) and the reaction was stirred at room temperature for 1 hour. The solution was quenched with 1N HCl, extracted with ethyl acetate, washed with brine, dried with sodium sulfate, filtered and concentrated to give 6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinic acid in 84% yield. LC/MS=284.1 (M+H), Rt=0.76 min.

Method 3

Synthesis of 2-(2,6-difluoro-4-methylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborane




To a solution of 1,3-difluoro-5-methylbenzene (1.0 eq) in dry THF (0.2 M) under an atmosphere of N₂ at -78° C. was

37


added n-butyllithium (1 eq, 1.6 M in hexanes) slowly keeping the internal temperature below -65°C . The reaction was stirred for 2 hrs at -78°C ., followed by the addition of 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.15 eq). The reaction was allowed to warm to room temperature. Upon completion, the reaction was quenched with NaHCO_3 (sat.) and extracted with EtOAc. The organics were washed with brine, dried over Na_2SO_4 , filtered and concentrated to yield 2-(2,6-difluoro-4-methylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane as a white solid in 92%. ^1H NMR (400 MHz, CDCl_3) δ ppm 6.67 (dd, $J=9.39, 0.78$ Hz, 2H), 2.34 (s, 3H), 1.38 (s, 12H).

Synthesis of methyl
6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinate



Method 1 was followed using methyl 6-bromo-5-fluoropicolinate (1.0 equiv.) and 2-(2,6-difluoro-4-methylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.75 equiv.) to give methyl 6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinate as a solid in 85% yield. LC/MS=282.0 (M+H), $R_t=0.87$ min.

Synthesis of
6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinic
acid




To a solution of methyl 6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinate (1.0 eq) in THF (0.1 M) was added LiOH (5.5 eq, 2 M) and allowed to stir at room temperature for 4 hrs. The volatiles were removed in vacuo, and the residual aqueous was acidified with 2 M HCl to pH 4. The precipitate was filtered and dried to yield 6-(2,6-difluoro-4-methylphenyl)-

38


5-fluoropicolinic acid as a light yellow solid in 73.5%. LCMS (m/z): 268.0 (MH^+), $R_t=0.76$ min.

Synthesis of (2-(3,5-difluorophenyl)propan-2-yloxy)
triisopropylsilane



To a solution of 1-(3,5-difluorophenyl)ethanone (1.0 equiv) in THF (0.2 M) at 0°C . was added methylmagnesium bromide (1.0 M in THF, 1.15 equiv). After stirring for 4 hours the reaction was quenched by addition of NH_4Cl (sat.), diluted with EtOAc, washed with NaCl (sat.), dried over MgSO_4 , filtered, concentrated and purified by ISCO SiO_2 chromatography to yield 2-(3,5-difluorophenyl)propan-2-ol. To a solution of 2-(3,5-difluorophenyl)propan-2-ol in CH_2Cl_2 (0.1 M) at 0°C . was added 2.6 lutidine (6 equiv.) and then triisopropylsilyl trifluoromethanesulfonate (3.0 equiv.). After stirring for 3 hours at 0°C . and six hours at rt the solution was partitioned between EtOAc and NaHCO_3 (sat.), separated, washed with NaCl (sat.), dried over MgSO_4 , filtered, concentrated and purified by ISCO SiO_2 chromatography to yield (2-(3,5-difluorophenyl)propan-2-yloxy)triisopropylsilane. (400 MHz, CDCl_3) δ ppm 1.05-1.08 (m, 21H) 1.57 (s, 6H) 6.63 (s, 1H) 7.00 (dd, $J=9.39, 2.35$ Hz, 2H).

Synthesis of (2-(3,5-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)propan-2-yloxy)
triisopropylsilane




To a solution of (2-(3,5-difluorophenyl)propan-2-yloxy)triisopropylsilane (1.0 eq) in dry THF (0.2 M) under an atmosphere of N_2 at -78°C . was added n-butyllithium (1 eq, 1.6 M in hexanes) slowly keeping the internal temperature below -65°C . The reaction was stirred for 2 hrs at -78°C ., followed by the addition of 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.15 eq). The reaction was allowed to warm to room temperature. Upon completion, the reaction was quenched with NaHCO_3 (sat.) and extracted with EtOAc. The organics were washed with brine, dried over Na_2SO_4 , filtered and concentrated to yield (2-(3,5-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)propan-2-yloxy)triisopropylsilane.

39


ethyl-1,3,2-dioxaborolan-2-yl)phenyl)propan-2-yloxy)triisopropylsilane in 99%. ^1H NMR (400 MHz, CDCl_3) δ ppm 1.03-1.08 (m, 21H) 1.24 (s, 12H) 1.38 (s, 3H) 1.57 (s, 3H) 6.92-7.03 (m, 2H).

Synthesis of
tert-butyl(3,5-difluorophenoxy)dimethylsilane



To a solution of 3,5-difluorophenol (1.0 equiv.) and imidazole (2.2 equiv.) in DMF (0.8 M) at 0°C . was added TBDMSCl (1.1 equiv.). The ice bath was removed and after stirring for 3 hours the solution was diluted with EtOAc, washed with water, brine, dried over MgSO_4 , filtered, concentrated and purified by SiO_2 chromatography to yield tert-butyl(3,5-difluorophenoxy)dimethylsilane in 73% yield. ^1H NMR (400 MHz, CDCl_3) δ ppm 0.23 (s, 6H) 0.99 (s, 9H) 6.33-6.40 (m, 2H) 6.44 (tt, 1H).

Synthesis of tert-butyl(3,5-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy)dimethylsilane




To a solution of tert-butyl(3,5-difluorophenoxy)dimethylsilane (1.0 eq) in dry THF (0.2 M) under an atmosphere of N_2 at -78°C . was added n-butyllithium (1 eq, 1.6 M in hexanes) slowly keeping the internal temperature below -65°C . The reaction was stirred for 1 hr at -78°C ., followed by the addition of 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (2.1 eq). The reaction was allowed to warm to room temperature. Upon completion, the reaction was quenched with NaHCO_3 (sat.) and extracted with EtOAc. The organics were washed with brine, dried over Na_2SO_4 , filtered and concentrated to yield tert-butyl(3,5-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy)dimethylsilane in 99% yield. ^1H NMR (400 MHz, CDCl_3) δ ppm 1.03-1.08 (m, 21H) 1.24 (s, 12H) 1.38 (s, 3H) 1.57 (s, 3H) 6.92-7.03 (m, 2H).

40


ramethyl-1,3,2-dioxaborolan-2-yl)phenoxy)dimethylsilane in 91% yield. ^1H NMR (400 MHz, CDCl_3) δ ppm 0.21 (s, 6H) 0.97 (s, 9H) 1.37 (s, 12H) 6.33 (d, $J=9.39$ Hz, 2H).

Synthesis of methyl 6-(2,6-difluoro-4-hydroxyphenyl)-5-fluoropicolinate



Method 1 was followed using methyl 6-bromo-5-fluoropicolinate (1.0 equiv.) and tert-butyl(3,5-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy)dimethylsilane (1.75 equiv.) to give methyl 6-(2,6-difluoro-4-hydroxyphenyl)-5-fluoropicolinate in 65% yield. The reaction was heated for an additional 30 minutes at 100°C . in the microwave to drive to completion the deprotection of the TBDMS group. LC/MS=283.9 (M+H), $R_t=0.69$ min.


Synthesis of methyl 6-(4-(2-(tert-butyl dimethylsilyloxy)ethoxy)-2,6-difluorophenyl)-5-fluoropicolinate



To a solution of methyl 6-(2,6-difluoro-4-hydroxyphenyl)-5-fluoropicolinate (1.0 equiv.) and potassium carbonate (4.0 equiv.) in DMF (0.4 M) was added (2-bromoethoxy)(tert-butyl)dimethylsilane (2 equiv.). After stirring for 72 hours at rt the heterogeneous solution was diluted with water, extracted with EtOAc, dried over MgSO_4 , filtered, concentrated and purified by ISCO SiO_2 chromatography to yield methyl 6-(4-(2-(tert-butyl dimethylsilyloxy)ethoxy)-2,6-difluorophenyl)-5-fluoropicolinate in 74% yield. LC/MS=442.1 (M+H), $R_t=1.22$ min.


41

Synthesis of 6-(4-(2-(tert-butyldimethylsilyloxy)ethoxy)-2,6-difluorophenyl)-5-fluoropicolinic acid




Method 2 was followed using methyl 6-(4-(2-(tert-butyldimethylsilyloxy)ethoxy)-2,6-difluorophenyl)-5-fluoropicolinate to give 6-(4-(2-(tert-butyldimethylsilyloxy)ethoxy)-2,6-difluorophenyl)-5-fluoropicolinic acid in 94% yield. LC/MS=428.1 (M+H), R_f =1.13 min.

Synthesis of
1,3-difluoro-5-(2-methoxyethoxy)benzene



To a solution of 3,5-difluorophenol (1.0 equiv.), 2-methoxyethanol (3.0 equiv.) and triphenylphosphine (3.0 equiv.) in THF (0.1 M) was added DIAD (3.0 equiv.). After stirring at rt for 18 hours, the volatiles were removed in vacuo and the residue was purified by SiO_2 chromatography to yield 1,3-difluoro-5-(2-methoxyethoxy)benzene in 95% yield. ^1H NMR (400 MHz, CDCl_3) δ ppm 6.41-6.47 m (3H), 4.08 (t, 2H), 3.74 (t, 2H), 3.45 (s, 3H).


Synthesis of 2-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane



42


To a solution of 1,3-difluoro-5-(2-methoxyethoxy)benzene (1.0 eq) in dry THF (0.2 M) under an atmosphere of N_2 at -78°C . was added n-butyllithium (1 eq, 1.6 M in hexanes) slowly keeping the internal temperature below -65°C . The reaction was stirred for 1 hr at -78°C ., followed by the addition of 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (2.1 eq). The reaction was allowed to warm to room temperature. Upon completion, the reaction was quenched with NaHCO_3 (sat.) and extracted with EtOAc. The organics were washed with brine, dried over Na_2SO_4 , filtered and concentrated to yield 2-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane. ^1H NMR (400 MHz, CDCl_3) ppm 6.42 (d, 2H), 4.10 (m, 2H), 3.74 (m, 2H), 3.44 (s, 3H), 1.37 (s, 12H).

Synthesis of methyl 6-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-5-fluoropicolinate



Method 1 was followed using methyl 6-bromo-5-fluoropicolinate (1.0 equiv.) and 2-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.75 equiv.) at 80°C . for 1 hour to give methyl 6-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-5-fluoropicolinate in 95% yield. LC/MS=341.9 (M+H), R_f =0.89 min.


Synthesis of 6-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-5-fluoropicolinic acid



Method 2 was followed using methyl 6-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-5-fluoropicolinate to give 6-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-5-fluoropicolinic acid in 98% yield. LC/MS=327.9 (M+H), R_f =0.71 min.


43

Synthesis of methyl 6-(2,6-difluoro-4-(2-hydroxypropan-2-yl)phenyl)-5-fluoropicolinate




Method 1 was followed using methyl 6-bromo-5-fluoropicolinate (1.0 equiv.) and (2-(3,5-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)propan-2-yloxy)triisopropylsilane (1.6 equiv.) at 100° C. for 30 min in the microwave to give methyl 6-(2,6-difluoro-4-(2-hydroxypropan-2-yl)phenyl)-5-fluoropicolinate in 90% yield. LC/MS=325.9 (MH⁺), R_t=0.81 min. ¹H NMR (400 MHz, <cdcl3>) δ ppm 1.59 (s, 6H), 4.00 (s, 3H), 7.15 (d, J=9.00 Hz, 2H), 7.62-7.68 (m, 1H), 8.23-8.29 (m, 1H).

Synthesis of 6-(2,6-difluoro-4-(2-hydroxypropan-2-yl)phenyl)-5-fluoropicolinic acid



Method 2 was followed using methyl 6-(2,6-difluoro-4-(2-hydroxypropan-2-yl)phenyl)-5-fluoropicolinate to give 6-(2,6-difluoro-4-(2-hydroxypropan-2-yl)phenyl)-5-fluoropicolinic acid in 94% yield. LC/MS=312.0 (MH⁺), R_t=0.69 min.

Synthesis of 4-(3,5-difluorophenyl)tetrahydro-2H-pyran-4-ol




To a solution of 1-bromo-3,5-difluorobenzene (1.6 equiv.) in THF (0.26 M) under Ar was added Mg turnings (1.6

44


equiv.). A reflux condenser was attached and the solution was submerged in a 90° C. oil bath and refluxed for two hours. The dihydro-2H-pyran-4(3H)-one (1.0 equiv.) was added in THF via syringe. The solution was left stirring at rt under Ar for 5 hrs. The reaction solution was quenched by addition of NH₄Cl_(sat) and the solution was extracted with EtOAc, washed with NaCl_(sat), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography (0-100% EtOAc/n-heptanes gradient) to yield 4-(3,5-difluorophenyl)tetrahydro-2H-pyran-4-ol in 71% yield. ¹H NMR (400 MHz, CHLOROFORM-d) δ ppm 1.59-1.68 (m, 3H), 2.07-2.19 (m, 2H), 3.87-3.93 (m, 4H), 6.72 (tt, J=8.75, 2.20 Hz, 1H), 6.97-7.06 (m, 2H).

Synthesis of 4-(3,5-difluorophenyl)-3,6-dihydro-2H-pyran



4-(3,5-difluorophenyl)tetrahydro-2H-pyran-4-ol (1.0 equiv.) was dissolved in DCM (0.2 M) and cooled to 0° C. TEA (2.8 equiv.) was added to the solution, followed by MsCl (1.3 equiv.). The reaction was stirred at rt for 2 hrs. The solution was cooled to 0° C. and DBU (3.0 equiv.) was added. The reaction was stirred at rt for 18 hrs. The solution was concentrated and the residue was purified by SiO₂ chromatography (0-100% EtOAc in Heptanes) to afford 4-(3,5-difluorophenyl)-3,6-dihydro-2H-pyran in 38% yield. ¹H NMR (400 MHz, <cdcl3>) δ ppm 2.42-2.49 (m, 2H), 3.93 (t, J=5.48 Hz, 2H), 4.32 (q, J=2.74 Hz, 2H), 6.16-6.22 (m, 1H), 6.70 (tt, J=8.80, 2.35 Hz, 1H), 6.85-6.94 (m, 2H).

Synthesis of 4-(3,5-difluorophenyl)tetrahydro-2H-pyran




To a solution of 4-(3,5-difluorophenyl)-3,6-dihydro-2H-pyran (1.0 equiv.) in methanol (0.2 M) was added 10% Pd/C (0.05 equiv.). The reaction was placed under an atmosphere of hydrogen and stirred for 18 hours. Upon completion, the solution was filtered over a pad of Celite, the pad was washed with DCM, the filtrate was concentrated in vacuo to give 4-(3,5-difluorophenyl)tetrahydro-2H-pyran in 71% yield. ¹H

45


NMR (400 MHz, CDCl_3) δ ppm 1.76 (br. s., 4H), 2.75 (br. s., 1H), 3.50 (br. s., 2H), 4.08 (d, $J=9.78$ Hz, 2H), 6.56-6.94 (m, 3H).

Synthesis of 2-(2,6-difluoro-4-(tetrahydro-2H-pyran-4-yl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane



Method 3 was followed using 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (2.2 equiv.), butyllithium (1.1 equiv.) and 4-(3,5-difluorophenyl)tetrahydro-2H-pyran (1.0 equiv.) to give 2-(2,6-difluoro-4-(tetrahydro-2H-pyran-4-yl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane in 100% yield. ^1H NMR (400 MHz, CDCl_3) δ ppm 1.16-1.19 (m, 12H), 1.65-1.74 (m, 4H), 2.60-2.75 (m, 1H), 3.37-3.51 (m, 2H), 4.01 (dt, $J=11.54$, 3.42 Hz, 2H), 6.67 (d, $J=8.22$ Hz, 2H).


Synthesis of methyl 6-(2,6-difluoro-4-(tetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinate



Method 1 was followed using methyl 6-bromo-5-fluoropicolinate (1.0 equiv.) and 2-(2,6-difluoro-4-(tetrahydro-2H-pyran-4-yl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3.0 equiv.) at 100°C for 20 min in microwave to give methyl 6-(2,6-difluoro-4-(tetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinate in 59% yield. LC/MS=352.2 (MH^+), $R_t=0.92$ min.


46

Synthesis of 6-(2,6-difluoro-4-(tetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinic acid



Method 2 was followed using methyl 6-(2,6-difluoro-4-(tetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinate to give 6-(2,6-difluoro-4-(tetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinic acid in 71% yield. LC/MS=338.1 (MH^+), $R_t=0.80$ min.

Synthesis of 4-(3,5-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)tetrahydro-2H-pyran-4-ol




Method 3 was followed using 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (2.5 equiv.), butyllithium (2.4 equiv.) and 4-(3,5-difluorophenyl)tetrahydro-2H-pyran-4-ol (1.0 equiv.) to give 4-(3,5-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)tetrahydro-2H-pyran-4-ol in 97% yield. ^1H NMR (400 MHz, CDCl_3) δ ppm 1.32-1.42

47


(m, 12H), 1.56-1.65 (m, 2H), 2.11 (d, J=3.13 Hz, 2H), 3.86-3.92 (m, 4H), 6.99 (d, J=9.00 Hz, 2H).

Synthesis of methyl 6-(2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinate



Method 1 was followed using methyl 6-bromo-5-fluoropicolinate (1.0 equiv.) and 4-(3,5-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)tetrahydro-2H-pyran-4-ol (1.8 equiv.) at to give methyl 6-(2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinate in 72% yield. LC/MS=368.0 (MH⁺), R_t=0.75 min.

Synthesis of 6-(2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinic acid



Method 2 was followed using methyl 6-(2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinate to give 6-(2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-

48

ran-4-yl)phenyl)-5-fluoropicolinic acid in 69% yield. LC/MS=354.0 (MH⁺), R_t=0.64 min.


Synthesis of methyl 6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinate

10

15

20

25



To a solution of methyl 6-(2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinate (1.0 equiv.) in CH₂Cl₂ (0.04 M) at -78° C. under Ar was added methylDAST (2.0 equiv.). After addition, the solution was stirred under Ar at -78° C. for 10 minutes and then the bath was removed. The reaction was allowed to warm up to rt and quenched by addition of NaHCO_{3(sat.)}. The solution was diluted with EtOAc, washed with NaHCO_{3(sat.)}, NaCl_(sat.), dried over MgSO₄, filtered, concentrated, purified by ISCO SiO₂ chromatography (0-100 EtOAc/n-heptanes) to yield methyl 6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinate in 100% yield. LC/MS=370.0 (MH⁺), R_t=0.94 min.


Synthesis of 6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinic acid

45

50

55

60




Method 2 was followed using methyl 6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinate to give 6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-

49


yl)phenyl)-5-fluoropicolinic acid in 95% yield.
LC/MS=355.9 (MH⁺), R_f=0.81 min.

Synthesis of 1-(3,5-difluorophenyl)cyclobutanol



To a solution of 1-bromo-3,5-difluorobenzene (1.0 equiv.) in THF (0.26 M) under Ar was added Mg turnings (1.6 equiv.). A reflux condenser was attached and the solution was submerged in a 90° C. oil bath and refluxed for two hours. The cyclobutanone (1.0 equiv.) was added in THF via syringe. The solution was left stirring at rt under Ar for 5 hrs. The reaction solution was quenched by addition of NH₄Cl_(sat) and the solution was extracted with EtOAc, washed with NaCl_(sat), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography (0-100% EtOAc/n-heptanes gradient) to yield 1-(3,5-difluorophenyl)cyclobutanol in 54% yield. ¹H NMR (400 MHz, CHLOROFORM-d) δ ppm 1.69-1.83 (m, 1H), 2.03-2.13 (m, 1H), 2.31-2.43 (m, 2H), 2.45-2.56 (m, 2H), 6.71 (tt, J=8.80, 2.35 Hz, 1H), 6.98-7.07 (m, 2H).


Synthesis of 1-(3,5-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)cyclobutanol



Method 3 was followed using 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (2.5 equiv.), butyllithium (2.4 equiv.) and 1-(3,5-difluorophenyl)cyclobutanol (1.0 equiv.) to give 1-(3,5-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)cyclobutanol in 100% yield. ¹H NMR (400 MHz, <cdcl3>) δ ppm 1.23-1.25 (m, 12H), 1.69-1.82 (m, 1H), 2.05-2.12 (m, 1H), 2.37 (br. s., 2H), 2.47 (br. s., 2H), 7.00 (d, J=8.80 Hz, 2H).


50

Synthesis of methyl 6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinate




Method 1 was followed using methyl 6-bromo-5-fluoropicolinate (1.0 equiv.) and 1-(3,5-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)cyclobutanol (1.6 equiv.) at 100° C. for 30 min in microwave to give methyl 6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinate in 71% yield. LC/MS=338.0 (MH⁺), R_f=0.85 min.

Synthesis of 6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinic acid



Method 2 was followed using methyl 6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinate to give 6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinic acid in 90% yield. LC/MS=323.9 (MH⁺), R_f=0.74 min.


Synthesis of 4-(3,5-difluorophenoxy)tetrahydro-2H-pyran



51


To a solution of 3,5-difluorophenol (1.0 equiv.), tetrahydro-2H-pyran-4-ol (1.2 equiv.), and triphenylphosphine (2.0 equiv.) in THF (0.33 M) at 0° C. was added DIAD (2.0 equiv.) dropwise. The reaction mixture was stirred at rt overnight. The mixture was concentrated and purified by flash chromatography over silica gel (heptanes:ethyl acetate gradient) to give 4-(3,5-difluorophenoxy)tetrahydro-2H-pyran in 90% yield. ¹H NMR (400 MHz, <cdcl3>) δ ppm 1.72-1.84 (m, 2H), 1.96-2.09 (m, 2H), 3.59 (ddd, J=11.64, 8.31, 3.52 Hz, 2H), 3.90-4.04 (m, 2H), 4.44 (tt, J=7.78, 3.77 Hz, 1H), 6.32-6.53 (m, 3H).

Synthesis of 2-(2,6-difluoro-4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane



Method 3 was followed using 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.5 equiv.), butyllithium (1.3 equiv.) and 4-(3,5-difluorophenoxy)tetrahydro-2H-pyran (1.0 equiv.) to give 2-(2,6-difluoro-4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane in 33% yield. ¹H NMR (400 MHz, <cdcl3>) δ ppm 1.21-1.34 (m, 12H), 1.78 (dtd, J=12.72, 8.31, 8.31 Hz, 2H), 1.93-2.09 (m, 2H), 3.59 (ddd, J=11.64, 8.31, 3.13 Hz, 2H), 3.89-4.01 (m, 2H), 4.48 (tt, J=7.78, 3.77 Hz, 1H), 6.40 (d, J=9.39 Hz, 2H).


Synthesis of methyl 6-(2,6-difluoro-4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-5-fluoropicolinate



52


Method 1 was followed using methyl 6-bromo-5-fluoropicolinate (1.0 equiv.) and 2-(2,6-difluoro-4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.5 equiv.) at 100° C. for 30 min in microwave to give methyl 6-(2,6-difluoro-4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-5-fluoropicolinate in 77% yield. LC/MS=368.0 (MH⁺), R_t=0.95 min.

Synthesis of 6-(2,6-difluoro-4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-5-fluoropicolinic acid



Method 2 was followed using methyl 6-(2,6-difluoro-4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-5-fluoropicolinate to give 6-(2,6-difluoro-4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-5-fluoropicolinic acid in 100% yield. LC/MS=353.9 (MH⁺), R_f=0.82 min.


Synthesis of methyl 6-(4-(ethoxymethyl)-2,6-difluorophenyl)-5-fluoropicolinate



To a solution of methyl 6-(2,6-difluoro-4-(hydroxymethyl)phenyl)-5-fluoropicolinate (1.0 equiv.) in DMF (0.20 M) (colorless) at 0° C. was added sodium hydride (1.2 equiv.) and the reaction was stirred at 0° C. for 2 min. Ethyl iodide (1.2 equiv.) was added and the reaction was allowed to warm to room temperature. After 1 h, additional 1.0 equiv. of NaH was added and stirred for 15 ml. Reaction was quenched by the addition of sat. Ammonium chloride. The aqueous was acidified with conc HCl to pH3 and extracted with ethyl acetate three times. The organics were combined, dried with MgSO₄, filtered and concentrated. The crude mixture was used as is. LC/MS=326.0 (MH⁺), R_f=0.94 min.


53

Synthesis of 6-(4-(ethoxymethyl)-2,6-difluorophenyl)-5-fluoropicolinic acid




Method 2 was followed using methyl 6-(4-(ethoxymethyl)-2,6-difluorophenyl)-5-fluoropicolinate to give 6-(4-(ethoxymethyl)-2,6-difluorophenyl)-5-fluoropicolinic acid. LC/MS=311.9 (MH⁺), R_t=0.82 min.

Synthesis of 1,3-difluoro-5-isopropoxybenzene



To a solution of 3,5-difluorophenol (1.0 equiv.) in DMF (0.26 M) was added potassium carbonate (2.2 equiv.) followed by 2-iodopropane (1.1 equiv.) and the reaction was stirred overnight at room temperature. The reaction was poured into a separatory funnel and diluted with a 3:1 (v/v) solution of EtOAc:heptanes. The organic phase was washed with water, then sat'd NaHCO₃. The remaining organic phase was dried over MgSO₄, filtered and concentrated in vacuo to provide 1,3-difluoro-5-isopropoxybenzene in 88% yield. ¹H NMR (400 MHz, <cdcl3>) δ ppm 1.33 (d, J=6.26 Hz, 6H), 4.48 (dt, J=11.93, 6.16 Hz, 1H), 6.31-6.47 (m, 3H).

Synthesis of 2-(2,6-difluoro-4-isopropoxyphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane




Method 3 was followed using 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (2.2 equiv.), butyllithium (1.2

54


equiv.) and 1,3-difluoro-5-isopropoxybenzene (1.0 equiv.) to give 2-(2,6-difluoro-4-isopropoxyphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane in 99% yield. ¹H NMR (400 MHz, <cdcl3>) δ ppm 1.24 (s, 12H), 1.31-1.33 (m, 6H), 4.43-4.56 (m, 1H), 6.31-6.44 (m, 2H).

Synthesis of methyl 6-(2,6-difluoro-4-isopropoxyphenyl)-5-fluoropicolinate




Method 1 was followed using methyl 6-bromo-5-fluoropicolinate (0.8 equiv.) and 2-(2,6-difluoro-4-isopropoxyphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.0 equiv.) at 70° C. for 1 hour to give methyl 6-(2,6-difluoro-4-isopropoxyphenyl)-5-fluoropicolinate. LC/MS=325.9 (MH⁺), R_t=1.04 min.

Synthesis of 6-(2,6-difluoro-4-isopropoxyphenyl)-5-fluoropicolinic acid



Method 2 was followed using methyl 6-(2,6-difluoro-4-isopropoxyphenyl)-5-fluoropicolinate to give 6-(2,6-difluoro-4-isopropoxyphenyl)-5-fluoropicolinic acid. LC/MS=311.9 (MH⁺), R_t=0.92 min.


Synthesis of 3-(3,5-difluorophenyl)oxetane



55


3,5-difluorophenylboronic acid (2.0 equiv.), (1R,2R)-2-aminocyclohexanol (0.06 equiv.), NaHMDS (2.0 equiv.), and nickel(II) iodide (0.06 equiv.) were dissolved in 2-propanol (0.35 M). The mixture was degassed with N₂, stirred at rt for 10 min and then a solution of 3-iodooxetane (1.0 equiv.) in 2-Propanol (0.70 M) was added. The mixture was sealed and heated at 80° C. in the microwave for 20 min. The mixture was filtered through celite, eluting with EtOH and concentrated. The crude residue was purified by ISCO SiO₂ chromatography eluting with 0-100% EtOAc in Heptanes to afford 3-(3,5-difluorophenyl)oxetane in 63% yield. ¹H NMR (400 MHz, <cdcl₃>) δ 6.88-6.96 (m, 2H), 6.72 (tt, J=2.20, 8.95 Hz, 1H), 5.08 (dd, J=6.26, 8.22 Hz, 2H), 4.71 (t, J=6.26 Hz, 2H), 4.14-4.24 (m, 1H).

Synthesis of 2-(2,6-difluoro-4-(oxetan-3-yl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane



Method 3 was followed using 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.3 equiv.), butyllithium (1.1 equiv.) and 3-(3,5-difluorophenyl)oxetane (1.0 equiv.) to give 2-(2,6-difluoro-4-(oxetan-3-yl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane. ¹H NMR (400 MHz, <cdcl₃>) δ ppm 6.90 (d, J=8.22 Hz, 2H), 5.07 (dd, J=6.06, 8.41 Hz, 2H), 4.70 (t, J=6.26 Hz, 2H), 4.13-4.23 (m, 1H), 1.39 (s, 12H).


Synthesis of methyl 6-(2,6-difluoro-4-(oxetan-3-yl)phenyl)-5-fluoropicolinate



56


Method 1 was followed using methyl 6-bromo-5-fluoropicolinate (1.2 equiv.) and 2-(2,6-difluoro-4-(oxetan-3-yl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.0 equiv.) at 80° C. for 15 min in microwave to give methyl 6-(2,6-difluoro-4-(oxetan-3-yl)phenyl)-5-fluoropicolinate in 47% yield. LC/MS=324.0 (MH⁺), Rt=0.75 min.

Synthesis of 6-(2,6-difluoro-4-(oxetan-3-yl)phenyl)-5-fluoropicolinic acid



Method 2 was followed using methyl 6-(2,6-difluoro-4-(oxetan-3-yl)phenyl)-5-fluoropicolinate to give 6-(2,6-difluoro-4-(oxetan-3-yl)phenyl)-5-fluoropicolinic acid in 71% yield. LC/MS=309.9 (MH⁺), Rt=0.69 min.

Synthesis of methyl 2',6,6'-trifluoro-4'-(trifluoromethylsulfonyloxy)biphenyl-3-carboxylate




To a solution of methyl 2',6,6'-trifluoro-4'-hydroxybiphenyl-3-carboxylate (1.0 equiv.) in DCM (0.35 M) at 0° C. was added pyridine (1.5 equiv.) and allowed to stir for 5 mins, followed by the addition of Triflic Anhydride (1.1 equiv.). The reaction was allowed to stir warming to RT. The reaction was quenched with NaHCO₃(sat), extracted in DCM and the organics were washed with water and brine. The organics were dried over Na₂SO₄, filtered, and concentrated to yield

57


methyl 2',6,6'-trifluoro-4'-(trifluoromethylsulfonyloxy)bi-phenyl-3-carboxylate in 81% yield.

Synthesis of methyl 6-(4-(3,6-dihydro-2H-thiopyran-4-yl)-2,6-difluorophenyl)-5-fluoropicolinate



To a degassed solution of methyl 6-(2,6-difluoro-4-(trifluoromethylsulfonyloxy)phenyl)-5-fluoropicolinate (1.0 equiv.) and 3,6-dihydro-2H-thiopyran-4-ylboronic acid (1.5 equiv.) in DME/2M Na₂CO₃ (3/1, 0.10 M) was added PdCl₂(dppf).CH₂Cl₂ adduct (0.10 equiv.). The reaction was heated to 90° C. in an oil bath for 15 min. The reaction mixture was partitioned with water and EtOAc; the organics were dried over MgSO₄, filtered, and concentrated. The crude was purified via ISCO SiO₂ chromatography. Pure fractions were combined and concentrated to yield methyl 6-(4-(3,6-dihydro-2H-thiopyran-4-yl)-2,6-difluorophenyl)-5-fluoropicolinate in 60% yield. LC/MS=366.1 (M+H), Rt=1.00 min.

Synthesis of methyl 6-(4-(1,1-dioxido-3,6-dihydro-2H-thiopyran-4-yl)-2,6-difluorophenyl)-5-fluoropicolinate




To a solution of methyl 6-(4-(3,6-dihydro-2H-thiopyran-4-yl)-2,6-difluorophenyl)-5-fluoropicolinate (1.0 equiv.) in DCM (0.10 M) at rt was added oxone (6.0 equiv.) in one portion. The resulting mixture was stirred at RT overnight, and then refluxed at 40° C. for 4 hrs. 10.0 equiv. of oxone were

58


added and the reaction was allowed to stir at 40° C. over the weekend. The reaction mixture was then diluted with DCM and washed with water the aqueous layer was then separated and extracted with DCM. The combined organic were then dried over MgSO₄ and concentrated in vacuo to yield methyl 6-(4-(1,1-dioxido-3,6-dihydro-2H-thiopyran-4-yl)-2,6-difluorophenyl)-5-fluoropicolinate in 100% yield. LC/MS=398.0 (M+H), Rt=0.76 min.

Synthesis of 6-(4-(1,1-dioxido-3,6-dihydro-2H-thiopyran-4-yl)-2,6-difluorophenyl)-5-fluoropicolinic acid



Method 2 was followed using methyl 6-(4-(1,1-dioxido-3,6-dihydro-2H-thiopyran-4-yl)-2,6-difluorophenyl)-5-fluoropicolinate to give 6-(4-(1,1-dioxido-3,6-dihydro-2H-thiopyran-4-yl)-2,6-difluorophenyl)-5-fluoropicolinic acid in 74% yield. LC/MS=384.0 (M+H), Rt=0.64 min.

Synthesis of 6-(4-(1,1-dioxido-3,6-dihydro-2H-thiopyran-4-yl)-2,6-difluorophenyl)-5-fluoropicolinic acid




To a degassed solution of 6-(4-(1,1-dioxido-3,6-dihydro-2H-thiopyran-4-yl)-2,6-difluorophenyl)-5-fluoropicolinic acid (1.0 equiv.) in EtOH (0.10 M) was added Pd/C (0.1 equiv.). The mixture was stirred at rt under H₂ for 16 hrs. Add Pd/C (0.1 equiv.) and the reaction was stirred for additional 16 hrs. The reaction was filtered and concentrated to yield 6-(4-

59


(1,1-dioxidotetrahydro-2H-thiopyran-4-yl)-2,6-difluorophenyl)-5-fluoropicolinic acid in 100% yield. LC/MS=386.0 (M+H), Rt=0.64 min.

Synthesis of methyl 6-(2,6-difluoro-4-(prop-1-en-2-yl)phenyl)-5-fluoropicolinate



To a degassed solution of methyl 6-(2,6-difluoro-4-(trifluoromethylsulfonyloxy)phenyl)-5-fluoropicolinate (1.0 equiv.) in DME/2M Na₂CO₃ (3/1, 0.09 M) was added 4,4,5,5-tetramethyl-2-(prop-1-en-2-yl)-1,3,2-dioxaborolane (1.5 equiv.) and PdCl₂(dppf)-CH₂Cl₂Adduct (0.1 equiv.). The reaction was heated to 90° C. in an oil bath for 15 min. The mixture was cooled to rt and partitioned between water and ethyl acetate. The organic phase was dried with sodium sulfate, filtered and concentrated. The crude material was purified via silica gel column chromatography (Analogix, eluting with 0-100% ethyl acetate). The pure fractions were concentrated to yield methyl 6-(2,6-difluoro-4-(prop-1-en-2-yl)phenyl)-5-fluoropicolinate. LC/MS=308.2 (M+H), Rt=0.99 min. ¹H NMR (400 MHz, <cdcl3>) δ ppm 2.15 (s, 3H), 4.01 (s, 3H), 5.23 (s, 1H), 5.47 (s, 1H), 7.11 (d, J=9.39 Hz, 2H), 7.65 (t, J=8.41 Hz, 1H), 8.26 (dd, J=8.61, 3.91 Hz, 1H).

Synthesis of methyl 6-(2,6-difluoro-4-isopropylphenyl)-5-fluoropicolinate




To a degassed solution of methyl 6-(2,6-difluoro-4-(prop-1-en-2-yl)phenyl)-5-fluoropicolinate (1.0 equiv.) in MeOH (0.09 M) was added Pd/C (0.1 equiv.) and the reaction was stirred at rt under an atmosphere of hydrogen. After overnight stirring, filtered through a pad of Celite and washed with Methanol. The filtrate was concentrated and dried under

60


vacuo to give methyl 6-(2,6-difluoro-4-isopropylphenyl)-5-fluoropicolinate. LC/MS=310.0 (M+H), Rt=1.00 min.

Synthesis of 6-(2,6-difluoro-4-isopropylphenyl)-5-fluoropicolinic acid




Method 2 was followed using methyl 6-(2,6-difluoro-4-isopropylphenyl)-5-fluoropicolinate to give 6-(2,6-difluoro-4-isopropylphenyl)-5-fluoropicolinic acid in 100% yield. LC/MS=296.2 (M+H), Rt=0.89 min.

Synthesis of ethyl 2-(2,6-difluorophenyl)thiazole-4-carboxylate



A solution of 2,6-difluorobenzothioamide (1.0 eq) and ethylbromopyruvate (1.0 eq.) in ethanol (1.0 M) was heated in the microwave at 130° C. for 30 minutes. Upon removal of volatiles in vacuo, ethyl acetate was added and the solution was washed with Na₂CO₃(sat.), with NaCl(sat.), was dried over MgSO₄, filtered and concentrated yielding ethyl 2-(2,6-difluorophenyl)thiazole-4-carboxylate (84%). LCMS (m/z): 270.1 (MH⁺); LC R_t=3.79 min.


Synthesis of 2-(2,6-difluorophenyl)thiazole-4-carboxylic acid



61


To a solution of ethyl 2-(2,6-difluorophenyl)thiazole-4-carboxylate (1.0 eq.) in 2:1 THF/MeOH (0.17 M) was added 1.0 M LiOH (2.0 eq.). After standing for 16 hours, 1.0 M HCl (2.0 eq.) was added and the THF/MeOH was removed in vacuo. The resulting solid was filtered, rinsed with H₂O and dried, yielding 2-(2,6-difluorophenyl)thiazole-4-carboxylic acid (88%) as a crusty solid. LC/MS (m/z): 251.1 (MH⁺); LC R_t=2.68 min.

Synthesis of ethyl 2-amino-2-cyanoacetate




To a solution of ethyl 2-cyano-2-(hydroxyimino)acetate (1 eq) in ethanol (1.4 M) was added PtO₂ (0.05 eq) and the solution was put under an H₂ atmosphere (4 bar) in a steel bomb and was stirred overnight. The reaction was filtered through a pad of celite, rinsing with CH₂Cl₂ and upon removal of volatiles in vacuo ethyl 2-amino-2-cyanoacetate was obtained in 89% yield. LC/MS (m/z): 129.0 (MH⁺), R_t: 0.25 min.

Synthesis of ethyl 2-cyano-2-(2,6-difluorobenzamido)acetate



To a solution of ethyl 2-amino-2-cyanoacetate (1 eq) in 6 mL of dichloromethane was added pyridine (1.5 eq) and 2,6-difluorobenzoyl chloride (1 eq) at 0° C. The reaction mixture was stirred at room temperature for 3 hours. The mixture was diluted with ethyl acetate, washed with brine, then dried over anhydrous MgSO₄, filtered, and concentrated in vacuo. The crude residue was purified by flash chromatography (EtOAc:hexanes=1:1) to give the titled compound (84%). LC/MS (m/z): 269.1 (MH⁺), R_t: 0.69 min.


Synthesis of ethyl 5-amino-2-(2,6-difluorophenyl)thiazole-4-carboxylate



62


To a solution of the ethyl 2-cyano-2-(2,6-difluorobenzamido)acetate (1 eq) in pyridine (0.1 M) was added Lawesson's reagent (1.5 eq.). The mixture was stirred at reflux under Ar for 18 hours. Solvents were removed under reduced pressure. The crude residue was purified by flash chromatography (EtOAc:hexanes=1:1) to give the ethyl 5-amino-2-(2,6-difluorophenyl)thiazole-4-carboxylate in 25% yield. LC/MS (m/z): 284.9 (MH⁺), R_t: 0.76 min.

Synthesis of 5-((tert-butoxycarbonyl)amino)-2-(2,6-difluorophenyl)thiazole-4-carboxylic acid




To a solution of the ethyl 5-amino-2-(2,6-difluorophenyl)thiazole-4-carboxylate (1 eq) in CH₂Cl₂ (0.1 M) was added Boc₂O (1.2 eq.) and DMAP (0.05 eq.). Upon stirring for 1 hour the volatiles were removed in vacuo, THF (0.1 M) and 2.0 M LiOH_(aq.) (5 equiv) were added and the solution was stirred at 55° C. for 2 days. The volatiles were removed in vacuo and the remaining aqueous solution was adjusted to pH 5 by addition of 2 M HCl. The resulting solid was filtered, rinsed with H₂O and pumped on to yield 5-((tert-butoxycarbonyl)amino)-2-(2,6-difluorophenyl)thiazole-4-carboxylic acid 25% yield. LC/MS (m/z): 357.1 (MH⁺), R_t: 0.97 min.

Synthesis of Methyl 3-amino-5-fluoropicolinate




To a steel bomb reactor, 2-bromo-5-fluoropyridin-3-amine (1.0 equiv.), triethylamine (1.6 equiv.), Pd(BINAP)Cl₂ (0.0015 equiv.) and anhydrous methanol (0.4 M solution) were added. After degassed by nitrogen stream for 15 min, the steel bomb reactor was closed and filled with CO gas up to 60 psi. The reactor was then heated to 100° C. After 3 h, more Pd catalyst (0.0015 equiv.) was added and the reaction mixture was re-heated to the same temperature for 3 h. After cooling down to room temperature, a brown precipitate was filtered off and the filtrate was extracted with EtOAc, which was washed with water and brine, dried over anhydrous sodium sulfate, and filtered. After removing volatile materials, the crude yellow product was obtained and used for the next step without further purification (40%). LC/MS (m/z): 271.2 (MH⁺); LC R_t=3.56 min.


63

Synthesis of Methyl
3-amino-6-bromo-5-fluoropicolinate

To a solution of methyl 3-amino-5-fluoropicolinate (1.0 equiv.) in acetonitrile (0.3 M solution) was added NBS (1.1 equiv.) for 2 minutes at room temperature. After quenched with water, the reaction mixture was extracted with EtOAc. The crude product was purified by silica column chromatography (20% to 50% EtOAc in hexanes) to give methyl 3-amino-6-bromo-5-fluoropicolinate (41%). LCMS (m/z): 249.1 (MH⁺); LC R_t=2.80 min.

Synthesis of methyl
3-amino-6-(2,6-difluorophenyl)-5-fluoropicolinate


Method 1 was followed using methyl 3-amino-6-bromo-5-fluoropicolinate (1.0 equiv.) and 2,6-difluorophenylboronic acid (1.3 equiv.) and Pd(dppf)Cl₂-DCM (0.05 equiv.) to give methyl 3-amino-6-(2,6-difluorophenyl)-5-fluoropicolinate in 94% yield. LCMS (m/z): 283.0 (MH⁺), R_t=0.76 min.

Synthesis of
3-amino-6-(2,6-difluorophenyl)-5-fluoropicolinic
acid

Method 2 was followed using methyl 3-amino-6-(2,6-difluorophenyl)-5-fluoropicolinate (1.0 equiv.) and LiOH (1.0


64

equiv.) to give 3-amino-6-(2,6-difluorophenyl)-5-fluoropicolinic acid in 79% yield. LCMS (m/z): 269.0 (MH⁺), R_t=0.79 min.

Synthesis of
2-(2,6-difluorophenyl)pyrimidine-4-carboxylic acid

To a solution of 2-chloropyrimidine-4-carboxylic acid (1.0 equiv.) in DME and 2M Na₂CO₃ (3:1, 0.25 M) was added 2,6-difluorophenylboronic acid (1.3 equiv.) and Pd(dppf)Cl₂-DCM (0.05 equiv.) in a microwave vial. The vial was heated in the microwave at 120° C. for 30 minutes. The mixture was diluted with ethyl acetate and 1N NaOH was added. The organic phase was separated and extracted three more times with 1N NaOH and once with 6N NaOH. The combined aqueous phases were filtered and acidified to pH 1 by the addition of concentrated HCl and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, filtered, and concentrated to give 2-(2,6-difluorophenyl)pyrimidine-4-carboxylic acid in 81%. LCMS (m/z): 237.0 (MH⁺), R_t=0.54 min.

Synthesis of 5-amino-2-(2,6-difluorophenyl)pyrimidine-4-carboxylic acid




A 2.68 M NaOEt in EtOH solution (3 eq) was added to an ice-bath cooled mixture of 2,6-difluorobenzimidamide hydrochloride (2 eq) in EtOH (0.1 M). The resulting mixture was allowed to warm to rt and stirred under N₂ for 30 min. To the reaction mixture was added drop wise a solution of mucobromic acid (1 eq) in EtOH and the reaction was heated in a 50° C. oil bath for 2.5 hr. After cooling to rt the reaction mixture was concentrated in vacuo. H₂O and 1.0 N NaOH were added and the aqueous mixture was washed with EtOAc. The aqueous phase was acidified to pH=4 with 1.0 N HCl then extracted with EtOAc. Combined organic extracts were washed once with brine, then dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo to give 5-bromo-2-(2,6-difluorophenyl)pyrimidine-4-carboxylic acid. The crude product was used for the next step without further purification. LC/MS (m/z): 316.9 (MH⁺). LC: R_t: 2.426 min.

65


CuSO₄ (0.1 eq) was added to a mixture of 5-bromo-2-(2,6-difluorophenyl)pyrimidine-4-carboxylic acid (1 eq) and 28% aqueous ammonium hydroxide solution in a microwave reaction vessel. The reaction mixture was heated in a microwave reactor at 110° C. for 25 min. The reaction vessel was cooled in dry ice for 30 min then unsealed and concentrated in vacuo. To the resulting solids was added 1.0 N HCl and the mixture was extracted with EtOAc. Combined organic extracts were washed once with brine, then dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo to give 5-amino-2-(2,6-difluorophenyl)pyrimidine-4-carboxylic acid. The crude product was used for the next step without further purification. LCMS (m/z): 252.0 (MH⁺), R_t=2.0 min.

Synthesis of
2-(2,6-difluorophenyl)-3-fluoro-6-methylpyridine



To a solution of 2-bromo-3-fluoro-6-methylpyridine (1.0 equiv.) in THF and Water (10:1, 0.2 M) was added 2,6-difluorophenylboronic acid (2.0 equiv.) and potassium fluoride (3.3 equiv.). The reaction was degassed for 10 minutes, then Pd₂(dba)₃ (0.05 equiv.) was added, followed by tri-*t*-butylphosphine (0.1 equiv.). The reaction was stirred to 60° C. for 1 hour at which point, all starting material was consumed as indicated by LC/MS. The reaction was allowed to cool to room temperature, partitioned with ethyl acetate and water, the organic phase was dried with sodium sulfate, filtered, and concentrated. The crude material was diluted in EtOH to 0.1 M, and 0.5 equiv. of NaBH₄ was added to reduce the dba. The reaction was stirred for one hour at room temperature, then quenched with water and concentrated under vacuo to remove the ethanol. The product was extracted in ether, washed with brine, the organics were dried over sodium sulfate, filtered, and concentrated. The crude material was loaded on silica gel and purified via column chromatography (ISCO) eluting with hexanes and ethyl acetate (0%-10% ethyl acetate). The pure fractions were combined, and concentrated to yield 2-(2,6-difluorophenyl)-3-fluoro-6-methylpyridine as a light yellow oil in 86% yield. LC/MS=224.0 (M+H), R_t=0.84 min.


Synthesis of 6-(2,6-difluorophenyl)-5-fluoropicolinic acid



66


To a solution of 2-(2,6-difluorophenyl)-3-fluoro-6-methylpyridine (1.0 equiv.) in water (0.05 M) was added KMnO₄ (2.0 equiv.) and the reaction was heated to reflux overnight. Another 2.0 equiv. of KMnO₄ were added and stirred at reflux for another 8 hours. The solution was cooled to room temperature, filtered through Celite and washed with water. The filtrate was acidified with 6N HCl to pH=3, the white precipitate was filtered. The filtrate was further acidified to pH=1 and filtered again. The filtrate was extracted with ethyl acetate until no more product was in the aqueous layer. The organic phase was washed with brine and dried over magnesium sulfate, filtered, and concentrated. The residue was dissolved in ethyl acetate, washed with 1N NaOH, the aqueous layer was acidified to pH=1 and the white crystals were filtered. The combined products yielded 6-(2,6-difluorophenyl)-5-fluoropicolinic acid in 32% yield as a white solid. LC/MS=254.0 (M+H), R_t=0.71 min.

Synthesis of methyl 3-amino-6-(2-fluoro-5-isopropylcabamoyl)phenyl-picolinate



A solution of methyl 3-amino-6-bromopicolinate (1.0 equiv.), N-isopropyl 3-borono-4-fluorobenzamide (1.1 equiv.), and Pd(dppf)Cl₂-DCM (0.15 equiv.) in DME/2M Na₂CO₃ (3:1), at a concentration of 0.5 M, was stirred at 120° C. for 1.5 hours. The reaction was filtered and washed with EtOAc. The organic was partitioned with H₂O (25 mL), washed with NaCl_(sat.) (25 mL), dried over MgSO₄, and the volatiles were removed in vacuo. The residue was diluted in EtOAc and passed through a silica gel plug and the volatiles were removed in vacuo yielding methyl 3-amino-6-(2-fluoro-5-isopropylcabamoyl)phenyl-picolinate (60%). LCMS (m/z): 332.2 (MH⁺); LC R_t=2.9 min.


Synthesis of 3-amino-6-(2-fluoro-5-isopropylcabamoyl)phenyl-picolinic acid



67


To a solution of methyl 3-amino-6-(2-fluoro-5-isopropylcabamoyl)phenylpicolinate (1.0 equiv) in THF (0.5 M), was added 1M LiOH (4.0 equiv). After stirring for 4 hours at 60° C., 1 N HCl (4.0 equiv.) was added and the THF was removed in vacuo. The resulting solid was filtered and rinsed with cold H₂O (3×20 mL) to yield 3-amino-6-(2-fluoro-5-isopropylcabamoyl)phenylpicolinic acid (98%). LCMS (m/z): 318.1 (MH⁺); LC R_t=2.4 min.

Synthesis of (R)-tert-butyl 4-((1R,2R)-3-((R)-4-benzyl-2-oxooxazolidin-3-yl)-1-hydroxy-2-methyl-3-oxopropyl)-2,2-dimethyloxazolidine-3-carboxylate



To a solution of (R)-4-benzyl-3-propionyloxazolidin-2-one (1.0 equiv.) in DCM (0.13 M) was added TiCl₄ (1.0 equiv.) at -40° C. The mixture was stirred at -40° C. for 10 min (yellow suspension), then DIPEA (2.5 equiv.) was added (dark red solution) and stirred at 0° C. for 20 min. (R)-tert-butyl 4-formyl-2,2-dimethyloxazolidine-3-carboxylate (1.0 equiv.) in DCM (0.5 M) was then added dropwise and the resulting mixture was stirred for 1.5 hours. The reaction was quenched by the addition of aqueous ammonium chloride and the mixture was extracted with ethyl acetate. The organic phase was separated, washed with brine, dried with magnesium sulfate, filtered, and concentrated. The residue was purified via column chromatography eluting with ethyl acetate and hexanes (1:4) to give (R)-tert-butyl 4-((1R,2R)-3-((R)-4-benzyl-2-oxooxazolidin-3-yl)-1-hydroxy-2-methyl-3-oxopropyl)-2,2-dimethyloxazolidine-3-carboxylate as the major product (5:2) in 58% yield. LC/MS=363.3 (M+H-Boc), R_t=1.09 min.


Synthesis of (R)-tert-butyl 4-((1R,2R)-3-((R)-4-benzyl-2-oxooxazolidin-3-yl)-1-(tert-butyldimethylsilyloxy)-2-methyl-3-oxopropyl)-2,2-dimethyloxazolidine-3-carboxylate



68


To a solution of (R)-tert-butyl 4-((1R,2R)-3-((R)-4-benzyl-2-oxooxazolidin-3-yl)-1-hydroxy-2-methyl-3-oxopropyl)-2,2-dimethyloxazolidine-3-carboxylate (1.0 equiv.) and lutidine (1.8 equiv.) in DCM (0.1 M) was added TBSOTf (1.4 equiv.) at -40° C. The reaction mixture was stirred at -40° C. for 2 hours. The solution was diluted with ethyl acetate and washed with sat. NaHCO₃, sat. NaCl, dried with magnesium sulfate, filtered, and concentrated. The residue was purified by silica gel column chromatography eluting with ethyl acetate and hexanes (1:4) to give (R)-tert-butyl 4-((1R,2R)-3-((R)-4-benzyl-2-oxooxazolidin-3-yl)-1-(tert-butyldimethylsilyloxy)-2-methyl-3-oxopropyl)-2,2-dimethyloxazolidine-3-carboxylate as the major product (5:2) in 83% yield. LC/MS=577.3 (M+H), R_t=1.33 min (Frac 65%-95% method).

Synthesis of (R)-tert-butyl 4-((1R,2S)-1-(tert-butyldimethylsilyloxy)-3-hydroxy-2-methylpropyl)-2,2-dimethyloxazolidine-3-carboxylate



To a solution of (R)-tert-butyl 4-((1R,2R)-3-((R)-4-benzyl-2-oxooxazolidin-3-yl)-1-(tert-butyldimethylsilyloxy)-2-methyl-3-oxopropyl)-2,2-dimethyloxazolidine-3-carboxylate (1.0 equiv.) and ethanol (3.0 equiv.) in THF (0.09 M) was added LiBH₄ (3.0 equiv.) at -30° C. The reaction mixture was allowed to warm up to 0° C. and stirred at that temperature for 3 hours. The solution was then diluted with diethyl ether and 1N NaOH was added. The resulting mixture was extracted with ethyl acetate, the organic layer was separated, washed with sat. NaCl, dried over magnesium sulfate, filtered, and concentrated. The residue was purified via silica gel column chromatography eluting with ethyl acetate and hexanes (1:4) to give (R)-tert-butyl 4-((1R,2S)-1-(tert-butyldimethylsilyloxy)-3-hydroxy-2-methylpropyl)-2,2-dimethyloxazolidine-3-carboxylate as the major product (5:2 ratio) in 71% yield. LC/MS=304.3 (M+H-Boc), R_t=0.95 min (Frac 65%-95% method).

Synthesis of (R)-tert-butyl 4-((1R,2S)-3-azido-1-(tert-butyldimethylsilyloxy)-2-methylpropyl)-2,2-dimethyloxazolidine-3-carboxylate




To a solution of (R)-tert-butyl 4-((1R,2S)-1-(tert-butyldimethylsilyloxy)-3-hydroxy-2-methylpropyl)-2,2-dimethyloxazolidine-3-carboxylate (1.0 equiv.), DIAD (2.0 equiv.), and PPh₃ (2.0 equiv.) in THF (0.18 M) was added DPPA (2.0 equiv., 1M solution in THF). The reaction mixture was stirred at room temperature overnight. Upon removal of the volatiles under vacuo, the residue was purified by silica

69


gel column chromatography eluting with ethyl acetate and hexanes (1:6) to give (R)-tert-butyl 4-((1R,2S)-3-azido-1-(tert-butyldimethylsilyloxy)-2-methylpropyl)-2,2-dimethylloxazolidine-3-carboxylate as the major product (5:2) in 86% yield. LC/MS=329.3 (M+H-Boc), Rt=1.40 min (Frac 65%-95% method).

Synthesis of tert-butyl (2R,3R,4S)-5-azido-3-(tert-butyldimethylsilyloxy)-1-hydroxy-4-methylpentan-2-ylcarbamate



To a solution of (R)-tert-butyl 4-((1R,2S)-3-azido-1-(tert-butyldimethylsilyloxy)-2-methylpropyl)-2,2-dimethylloxazolidine-3-carboxylate (1.0 equiv.) in EtOH (0.1 M) was added PPTS (1.3 equiv.) and the mixture was refluxed for 2 days. The volatiles were removed under vacuo, the residue was dissolved in DCM (0.1 M) and DIEA (1.5 equiv.) and Boc₂O (1.0 equiv.) were added to the reaction mixture. The solution was stirred for 3 hours at room temperature. The solvents were removed under reduced pressure and the residue was diluted with ethyl acetate, washed with water, aqueous NaHSO₄, aqueous NaHCO₃, sat. NaCl, the organic phase was dried with magnesium sulfate, filtered, and concentrated. The residue was purified via silica gel column chromatography eluting with ethyl acetate and hexanes (1:3) to give tert-butyl (2R,3R,4S)-5-azido-3-(tert-butyldimethylsilyloxy)-1-hydroxy-4-methylpentan-2-ylcarbamate as the major isomer (5:2) in 70% yield. LC/MS=289.3 (M+H-Boc), Rt=0.76 min (Frac 65%-95% method).


Synthesis of (2R,3R,4S)-5-azido-2-(tert-butoxycarbonylamino)-3-(tert-butyldimethylsilyloxy)-4-methylpentyl methanesulfonate



To a solution of tert-butyl (2R,3R,4S)-5-azido-3-(tert-butyldimethylsilyloxy)-1-hydroxy-4-methylpentan-2-ylcarbamate (1.0 equiv.) in pyridine (0.2 M) was added MsCl (1.3 equiv.) followed by DMAP (catalytic amount) at 0° C. The mixture was stirred at that temperature for 1 hour. The solution was diluted with ether and ethyl acetate (4:1), washed with aq. NaHSO₄, sat. NaHCO₃, brine, dried over magnesium sulfate, filtered, and concentrated. The residue was purified by silica gel column chromatography eluting with ethyl acetate and hexanes (1:3) to give (2R,3R,4S)-5-azido-2-(tert-butoxycarbonylamino)-3-(tert-butyldimethylsilyloxy)-4-methylpentyl methanesulfonate as the major isomer (5:2) in 90% yield. LC/MS=367.3 (M+H-Boc), Rt=0.81 min (Frac 65%-95% method).


70

Synthesis of tert-butyl (3R,4R,5S)-4-(tert-butyldimethylsilyloxy)-5-methylpiperidin-3-ylcarbamate




A solution of (2R,3R,4S)-5-azido-2-(tert-butoxycarbonylamino)-3-(tert-butyldimethylsilyloxy)-4-methylpentyl methanesulfonate in MeOH (0.09 M) was degassed with nitrogen for 20 min. DIEA (2.5 equiv.) was added, followed by 10% Pd/C (0.1 equiv.). The reaction mixture was stirred under a hydrogen balloon for 2 hours. The solution was filtered and the filtrate was concentrated under vacuo to afford tert-butyl (3R,4R,5S)-4-(tert-butyldimethylsilyloxy)-5-methylpiperidin-3-ylcarbamate as the major isomer (5:2) in >99% yield. LC/MS=345.2 (M+H-Boc), Rt=0.95 and 0.99 min.

Synthesis of tert-butyl (3R,4R,5S)-4-(tert-butyldimethylsilyloxy)-5-methyl-1-(3-nitropyridin-4-yl)piperidin-3-ylcarbamate



To a solution of tert-butyl (3R,4R,5S)-4-(tert-butyldimethylsilyloxy)-5-methylpiperidin-3-ylcarbamate (1.0 equiv.) in i-PrOH (0.09 M) was added DIEA (2.5 equiv.) and 4-chloro-3-nitropyridine (1.5 equiv.). The reaction mixture was stirred at 60° C. for 2 hours. The volatiles were removed under vacuo, the residue was diluted with ethyl acetate and washed with sat. NaCl. The organic phase was dried with magnesium sulfate, filtered, and concentrated. The crude material was purified by silica gel column chromatography eluting with ethyl acetate and hexanes (1:2) to give tert-butyl (3R,4R,5S)-4-(tert-butyldimethylsilyloxy)-5-methyl-1-(3-nitropyridin-4-yl)piperidin-3-ylcarbamate in 76% yield. LC/MS=467.3 (M+H), Rt=1.09 min.


Synthesis of tert-butyl (3R,4R,5S)-1-(3-aminopyridin-4-yl)-4-(tert-butyldimethylsilyloxy)-5-methylpiperidin-3-ylcarbamate



71


A solution of tert-butyl (3R,4R,5S)-4-(tert-butyldimethylsilyloxy)-5-methyl-1-(3-nitropyridin-4-yl)piperidin-3-ylcarbamate (1.0 equiv.) in MeOH (0.05 M) was degassed with nitrogen for 20 min. 10% Pd/C (0.2 equiv.) was added to the mixture and the solution was stirred under a hydrogen balloon for 3 hours. The reaction was filtered and the filtrate was concentrated under reduced pressure to give tert-butyl (3R,4R,5S)-1-(3-aminopyridin-4-yl)-4-(tert-butyldimethylsilyloxy)-5-methylpiperidin-3-ylcarbamate as the desired product in 94% yield. LC/MS=437.4 (M+H), Rt=1.08 min. ¹H-NMR (300 MHz, CDCl₃): δ 8.01 (s, 1H), 7.95 (d, J=6.0 Hz, 1H), 6.76 (d, J=6.0 Hz, 1H), 4.44 (br s, 1H), 3.74 (br s, 2H), 3.59-3.55 (m, 1H), 3.25-3.13 (m, 2H), 2.47-2.35 (m, 2H), 1.89 (br s, 2H), 1.44 (s, 9H), 1.04 (d, J=6.0, 3H), 0.92 (s, 9H), 0.13 (d, J=9.0, 6H).

Synthesis of 5-methyl-3-oxocyclohex-1-enyltrifluoromethanesulfonate



To a solution of 5-methylcyclohexane-1,3-dione (1.0 equiv.) in DCM (0.5 M) was added Na₂CO₃ (1.1 equiv.) and cooled to 0° C. Added Tf₂O (1.0 equiv.) in DCM (5.0 M) dropwise over 1 hr at 0° C. under a nitrogen atmosphere. Upon addition, the reaction was stirred for 1 hr at room temperature (dark red solution). The solution was filtered and the filtrate was quenched by careful addition of saturated NaHCO₃ with vigorous stirring until pH=7. The solution was transferred to a separatory funnel and the layers were separated. The organic layer was washed with brine, dried with Na₂SO₄, filtered, concentrated under vacuo and dried under high vacuum for 15 min to yield 5-methyl-3-oxocyclohex-1-enyl trifluoromethanesulfonate as light yellow oil in 78% yield. The triflate decomposes upon storage and should be used immediately for the next reaction. LC/MS=259.1/300.1 (M+H and M+CH₃CN); Rt=0.86 min, LC=3.84 min. ¹H-NMR (400 MHz, CDCl₃) δ ppm: 6.05 (s, 1H), 2.70 (dd, J=17.2, 4.3, 1H), 2.53 (dd, J=16.6, 3.7, 1H), 2.48-2.31 (m, 2H), 2.16 (dd, J=16.4, 11.7, 1H), 1.16 (d, J=5.9, 3H).

Synthesis of 5-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)cyclohex-2-enone




To a solution of 5-methyl-3-oxocyclohex-1-enyl trifluoromethanesulfonate (1.0 equiv.) in degassed dioxane (0.7 M) was added bis(pinacolato)diboron (2.0 equiv.), KOAc (3.0

72


equiv.), and Pd(dppf)Cl₂-DCM (0.03 equiv.). The reaction was heated to 80° C. for 10 h then cooled to room temperature and filtered through a coarse frit glass funnel. The cake was rinsed with more dioxane and the filtrate solution was used for the next step without further purification. LC/MS=155.1 (M+H of boronic acid); Rt=0.41 min, LC=1.37 min.

Synthesis of 5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enone



To a solution of 5-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)cyclohex-2-enone (1.0 equiv.) in degassed dioxane (0.5 M) and 2M Na₂CO₃ (2 equiv.) was added 4-chloro-3-nitropyridine (1.3 equiv.) and Pd(dppf)Cl₂-DCM (0.05 equiv.). The reaction was placed under a reflux condenser and heated in an oil bath to 110° C. for 1 h. Cooled to room temperature, filtered through a pad of Celite, washed the pad with ethyl acetate and concentrated the filtrate under vacuo. The residue was further pumped at 80° C. on a rotary evaporator for one hour to remove boronate by-products (M+H=101) via sublimation. The residue was partitioned between brine and ethyl acetate, and the layers were separated, the aqueous phase was further extracted with ethyl acetate (4×), the organics were combined, dried over sodium sulfate, filtered, and concentrated. The crude was purified via silica gel chromatography loading in DCM and eluting with 2-50% ethyl acetate and hexanes. The pure fractions were concentrated in vacuo to yield an orange oil. The oil was placed under high vacuum (~500 mtorr) with seed crystals overnight to yield an orange solid. The solid was further purified via trituration in hexanes to yield 5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enone (48% 2 steps). LC/MS=233.2 (M+H); Rt=0.69 min, LC=2.70 min. ¹H-NMR (400 MHz, CDCl₃) δ ppm: 9.31 (s, 1H), 8.88 (d, J=5.1, 1H), 7.30 (d, J=5.1, 1H), 6.00 (d, J=2.4, 1H), 2.62 (dd, J=16.4, 3.5, 1H), 2.53-2.34 (m, 3H), 2.23 (dd, J=16.1, 11.7, 1H), 1.16 (d, J=6.3, 3H).


Synthesis of cis-(+/-)-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enol



73


To a solution of 5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enone (1.0 equiv.) in EtOH (0.3 M) was added $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$ (1.2 equiv.). The reaction was cooled to 0°C ., then NaBH_4 (1.2 equiv.) was added in portions. Stirred for 1 h at 0°C ., then quenched by adding water, concentrated to remove the EtOH, added EtOAc, extracted the organics, washed with brine, then dried with Na_2SO_4 , filtered and concentrated to yield cis-(+/-)-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enol (94%). LC/MS=235.2 (M+H), LC=2.62 min.

Synthesis of (+/-)-4-(5-methylcyclohexa-1,3-dienyl)-3-nitropyridine



To a solution of (+/-)-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enol (1.0 equiv.) in dioxane (0.1 M) was added p-TSA (1.0 equiv.), and the reaction was stirred at 100°C . for 3 h. The solution was cooled to room temperature, then passed through a pad of neutral alumina eluting with EtOAc to yield (+/-)-4-(5-methylcyclohexa-1,3-dienyl)-3-nitropyridine as a yellow oil in 68% yield. LC/MS=217.1 (M+H), LC=3.908 min.

Synthesis of (+/-)-6-bromo-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enol




To a solution of 4-(5-methylcyclohexa-1,3-dienyl)-3-nitropyridine (1.0 equiv.) in THF and water (1:1, 0.13 M) was added NBS (1.5 equiv.) and the reaction was stirred at room temperature for 30 min. Upon completion, ethyl acetate and water were added to the reaction, the organic phase was dried with brine, then sodium sulfate, filtered, and concentrated. The crude material was purified via silica gel column chromatography eluting with ethyl acetate and hexanes (1:1) to

74


give (+/-)-6-bromo-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enol as a yellow oil in 80% yield. LC/MS=315.0/313.0 (M+H), LC=2.966 min.

Synthesis of (+/-)-2-azido-6-methyl-4-(3-nitropyridin-4-yl)cyclohex-3-enol



To a solution of (+/-)-6-bromo-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enol (1.0 equiv.) in THF (0.1 M) was added potassium tert-butoxide (1.5 equiv.). The reaction turned from orange to black almost immediately. By TLC, the formation of product is clean in 30 min. Quenched by adding saturated ammonium chloride and ethyl acetate. The organic phase was dried with brine, then sodium sulfate, filtered, and concentrated. The crude product was dissolved in ethanol and water (3:1, 0.1 M), and ammonium chloride (2.0 equiv) and sodium azide (2.0 equiv.) were added. The dark orange reaction was stirred at room temperature overnight. The conversion to product is clean as indicated by LC/MS. The reaction was concentrated to remove the ethanol, ethyl acetate and water were added, and the organic phase was dried with sodium sulfate, filtered, and concentrated. The crude material was purified via silica gel column chromatography eluting with ethyl acetate and hexanes (1:1) to give (+/-)-2-azido-6-methyl-4-(3-nitropyridin-4-yl)cyclohex-3-enol in 55% yield. LC/MS=276.0 (M+H), LC=2.803 min.

Synthesis of (+/-)-tert-butyl 6-hydroxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate




To a solution of (+/-)-2-azido-6-methyl-4-(3-nitropyridin-4-yl)cyclohex-3-enol (1.0 equiv.) in pyridine and ammonium hydroxide (8:1, 0.08 M) was added trimethylphosphine (3.0 equiv.) and the brown solution was stirred at room temperature for 2 h. Upon completion, EtOH was added and the solution was concentrated in vacuo. More ethanol was added and the reaction was concentrated again. Dioxane and sat. NaHCO_3 (1:1, 0.08 M) were added to the crude, followed by Boc_2O (1.0 equiv.). Stirred the reaction mixture at room tem-

75


perature for 2 h, then added water and ethyl acetate. The organic phase was dried with MgSO_4 , and concentrated. The crude product was purified via silica gel column chromatography eluting with ethyl acetate and hexanes (1:1) to afford (+/-)-tert-butyl 6-hydroxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate (59%). LC/MS=350.1 (M+H), Rt: 0.76 min.

Synthesis of (+/-)-2-(tert-butoxycarbonylamino)-6-methyl-4-(3-nitropyridin-4-yl)cyclohex-3-enyl acetate



To a solution of (+/-)-tert-butyl 6-hydroxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate (1.0 equiv.) in pyridine (0.1 M) was added Ac_2O (2.0 equiv.) and the reaction was stirred at room temperature overnight. Upon completion, the reaction was concentrated to dryness, then worked-up with ethyl acetate and water. The organic phase was dried with brine, then sodium sulfate, filtered, and concentrated to give (+/-)-2-(tert-butoxycarbonylamino)-6-methyl-4-(3-nitropyridin-4-yl)cyclohex-3-enyl acetate in 94% yield. LC/MS=392.2 (M+H), Rt=0.94 min.

Synthesis of (1S,2S,4S,6R)-4-(3-aminopyridin-4-yl)-2-((tert-butoxycarbonyl)amino)-6-methylcyclohexyl acetate and (1R,2R,4R,6S)-4-(3-aminopyridin-4-yl)-2-((tert-butoxycarbonyl)amino)-6-methylcyclohexyl acetate




To a degassed solution of (+/-)-2-(tert-butoxycarbonylamino)-6-methyl-4-(3-nitropyridin-4-yl)cyclohex-3-enyl acetate (1.0 equiv.) in MeOH and EtOAc (1:1, 0.1 M) was added 10% Pd/C (0.1 equiv.) and the reaction was stirred at room temperature under a hydrogen balloon for 3 days. Upon completion, the solution was filtered through a pad of Celite, the pad was washed with ethyl acetate and the filtrate was concentrated. The crude material contained about 10% of the undesired isomer. The crude was dissolved in ethyl acetate (~20%) and hexanes and heated until all dissolved. The solu-

76


tion was allowed to sit at room temperature for 2 days. The precipitate was then collected to give (+/-)-4-(3-aminopyridin-4-yl)-2-(tert-butoxycarbonylamino)-6-methylcyclohexyl acetate as the pure product in 59% yield. LC/MS=364.3 (M+H), Rt=0.63 min. The racemic material was resolved using an AD-H chiral column (20% i-PrOH/80% n-heptanes, 20 mL/min flow rate) to (1S,2S,4S,6R)-4-(3-aminopyridin-4-yl)-2-((tert-butoxycarbonyl)amino)-6-methylcyclohexyl acetate (peak#1, R_t =3.76 min on AD-H chiral analytical column, 20% i-PrOH/80% n-heptanes, 1 mL/min) and (1R,2R,4R,6S)-4-(3-aminopyridin-4-yl)-2-((tert-butoxycarbonyl)amino)-6-methylcyclohexyl acetate (peak#2, R_t =6.79 min on AD-H chiral analytical column, 20% i-PrOH/80% n-heptanes, 1 mL/min).

Synthesis of 2-(tert-butoxycarbonylamino)-6-methyl-4-(3-nitropyridin-4-yl)cyclohex-3-enyl methanesulfonate



To a solution of tert-butyl 6-hydroxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate (1.0 equiv.) in DCM (0.09 M) was added triethylamine (1.5 equiv.) and the reaction was cooled to 0° C. MsCl (1.2 equiv.) was added to the reaction and stirred for 3 h. Another 1.0 equiv. of MsCl was added to the reaction and stirred for another 2 h. Worked up the reaction by adding water, the organic phase was dried with brine, sodium sulfate, and concentrated. The crude product was purified via silica gel column chromatography eluting with ethyl acetate and hexanes (1:1) to afford 2-(tert-butoxycarbonylamino)-6-methyl-4-(3-nitropyridin-4-yl)cyclohex-3-enyl methanesulfonate as a white foam in 65% yield. LC/MS=428.2 (M+H), LC: 3.542 min.


Synthesis of (+/-)-tert-butyl 7-methyl-5-(3-nitropyridin-4-yl)-2-oxo-3a,6,7,7a-tetrahydrobenzo[d]oxazole-3(2H)-carboxylate



77


A solution of (+/-)-2-(tert-butoxycarbonylamino)-6-methyl-4-(3-nitropyridin-4-yl)cyclohex-3-enyl methanesulfonate (1.0 equiv.) in pyridine (0.2 M) was heated in the microwave at 110° C. for 10 min. The orange reaction was then concentrated under vacuo, the crude was dissolved in ethyl acetate and water, the organic phase was dried with sodium sulfate and concentrated under vacuo. The crude material was dissolved in DCM (0.2 M), triethylamine (1.8 equiv.) was added, followed by Boc₂O (1.2 equiv.). The reaction was stirred for 40 min, then concentrated to dryness. The crude material was purified via silica gel column chromatography eluting with hexane and ethyl acetate (1:1) to afford (+/-)-tert-butyl 7-methyl-5-(3-nitropyridin-4-yl)-2-oxo-3a,6,7,7a-tetrahydrobenzo[d]oxazole-3(2H)-carboxylate as a white foam in 66% yield. LC/MS=376.0 (M+H), LC: 3.424 min.

Synthesis of (+/-)-tert-butyl 5-(3-aminopyridin-4-yl)-7-methyl-2-oxohexahydrobenzo[d]oxazole-3(2H)-carboxylate



To a degassed solution of (+/-)-tert-butyl 7-methyl-5-(3-nitropyridin-4-yl)-2-oxo-3a,6,7,7a-tetrahydrobenzo[d]oxazole-3(2H)-carboxylate (1.0 equiv.) in MeOH and EtOAc (1:1, 0.1 M) was added 10% Pd/C (0.1 equiv.). The reaction was stirred under a hydrogen balloon overnight. Upon completion, the solution was filtered through a pad of Celite and the pad was washed with ethyl acetate. The filtrate was concentrated under vacuo to give (+/-)-tert-butyl 5-(3-aminopyridin-4-yl)-7-methyl-2-oxohexahydrobenzo[d]oxazole-3(2H)-carboxylate as the desired product as a yellow foam in 93% yield. LC/MS=348.1 (M+H), Rt=055 min.


Synthesis of tert-butyl (1R,2R,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-2-hydroxy-3-methylcyclohexylcarbamate



78


To a solution of (1R,2R,4R,6S)-4-(3-aminopyridin-4-yl)-2-(tert-butoxycarbonylamino)-6-methylcyclohexyl acetate (1.0 equiv.) in DMF/Ethanol (1/5, 0.05 M) was added 6-(2,6-difluorophenyl)-5-fluoropicolinic acid (1.3 equiv.), azadipic acid (1.3 equiv.) and EDC (1.3 equiv.). The mixture was stirred at rt for 6 hrs. The solution was diluted with EtOAc, washed with H₂O, 1N NaOH, NaCl (sat.), dried over MgSO₄, filtered and concentrated to yield crude protected amide. The material was dissolved in EtOH (0.45 M), Cs₂CO₃ (1.0 equiv.) was added and the solution was submerged in a 60° C. oil bath and stirred for 90 mins. The volatiles were removed in vacuo; the residue was partitioned between with EtOAc and H₂O. The organic layer was washed with NaCl (sat.), dried over MgSO₄, filtered, concentrated and pumped on to yield tert-butyl (1R,2R,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-2-hydroxy-3-methylcyclohexylcarbamate in 99% yield. LC/MS (m/z)=557.3 (MH⁺), R_t=0.80 min.

Synthesis of (1R,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl methanesulfonate



To a solution of tert-butyl (1R,2R,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-2-hydroxy-3-methylcyclohexylcarbamate (1.0 equiv.) in CH₂Cl₂ was added TEA (4.0 equiv.) and MsCl (2.0 equiv.). The capped solution was stirred for 5 minutes and then the homogeneous solution was left standing at rt for 16 hrs. The volatiles were removed in vacuo, the residue was dissolved in DMSO, purified by RP HPLC and the product fractions were lyophilized directly. The Boc protected product was treated with 25% TFA/CH₂Cl₂ for 20 minutes at which time the volatiles were removed in vacuo and the residue was dissolved in DMSO and purified by RP-HPLC. The product fractions were lyophilized directly to yield (1R,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl methanesulfonate in 31% yield. LC/MS (m/z)=535.2 (MH⁺), R_t=0.61 min.


Synthesis of tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-2-hydroxy-3-methylcyclohexylcarbamate



79

To a solution of (3aR,5R,7S,7aS)-tert-butyl 5-(3-aminopyridin-4-yl)-7-methyl-2-oxohexahydrobenzo[d]oxazole-3 (2H)-carboxylate (1.0 equiv.) in DMF (0.2 M) was added 6-(2,6-difluorophenyl)-5-fluoropicolinic acid (1.3 equiv.), aza-HOBt (1.3 equiv.) and EDC (1.3 equiv.). The mixture was stirred at rt for 16 hrs. The solution was diluted with EtOAc, washed with H₂O, 1N NaOH, NaCl_(sat.), dried over MgSO₄, filtered and concentrated to yield crude protected amide. The material was dissolved in EtOH (0.45 M), Cs₂CO₃ (1.0 equiv.) was added and the solution was submerged in a 60° C. oil bath and stirred for 90 mins. The volatiles were removed in vacuo; the residue was partitioned between with EtOAc and H₂O. The organic layer was washed with NaCl_(sat.), dried over MgSO₄, filtered, concentrated and pumped on to yield tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-2-hydroxy-3-methylcyclohexylcarbamate in 100% yield. LC/MS (m/z)=557.3 (MH⁺), R_f=0.83 min.


Synthesis of (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl methanesulfonate



To a solution of tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-2-hydroxy-3-methylcyclohexylcarbamate (1.0 equiv.) in pyridine (0.12 M) was added MsCl (7.0 equiv.). The capped solution was stirred for 5 minutes and then the homogeneous solution was left standing at rt for 16 hrs. The volatiles were removed in vacuo and the residue was partitioned between EtOAc and H₂O. The organic layer was washed with H₂O, 10% CuSO₄, H₂O, Na₂CO_{3(sat.)}, NaCl_(sat.), dried over MgSO₄, filtered, concentrated and purified by ISCO chromatography. The Boc protected product was treated with 25% TFA/CH₂Cl₂ for 20 minutes at which time the volatiles were removed in vacuo and the residue was dissolved in DMSO and purified by RP-HPLC. The product fractions were lyophilized directly to yield (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl methanesulfonate in 45% yield. LC/MS (m/z)=535.2 (MH⁺), R_f=0.60 min.


80

Synthesis of (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl dimethylphosphinate



To a solution of tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-2-hydroxy-3-methylcyclohexylcarbamate (1.0 equiv.) in pyridine (0.05 M) was added phosphonic chloride (5.0 equiv.). The capped homogeneous solution was left stirring at rt for 1 hr. The volatiles were removed in vacuo and the residue was partitioned between EtOAc and H₂O. The organic layer was washed with H₂O, 10% CuSO₄, Na₂CO_{3(sat.)}, NaCl_(sat.), dried over MgSO₄, filtered, concentrated to yield crude Boc protected product. The Boc group was removed by treating with 25% TFA/CH₂Cl₂ for 30 minutes at which time the volatiles were removed in vacuo and the residue was dissolved in DMSO and purified by RP-HPLC. The product fractions were lyophilized directly to yield (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl dimethylphosphinate in 43% yield. LC/MS (m/z)=533.3 (MH⁺), R_f=0.59 min.

Synthesis of (1R,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl dimethylphosphinate




To a solution of tert-butyl (1R,2R,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-2-hydroxy-3-methylcyclohexylcarbamate (1.0 equiv.) in pyridine (0.05 M) was added phosphonic chloride (5.0 equiv.). The capped homogeneous solution was left stirring at rt for 1 hr. The volatiles were removed in vacuo and the residue was partitioned between EtOAc and H₂O. The organic layer was washed with H₂O, 10% CuSO₄, Na₂CO_{3(sat.)}, NaCl_(sat.), dried over MgSO₄, filtered, concentrated to yield crude Boc protected product. The Boc group was removed by treating

81


with 25% TFA/CH₂Cl₂ for 30 minutes at which time the volatiles were removed in vacuo and the residue was dissolved in DMSO and purified by RP-HPLC. The product fractions were lyophilized directly to yield (1R,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl dimethylphosphinate in 43% yield. LC/MS (m/z)=533.3 (MH⁺), R_t=0.60 min.

Synthesis of S-(1S,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl ethanethioate



To a solution of (1R,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl methanesulfonate (1.0 equiv.) in DMF (0.19 M) was added KSAc (3.0 equiv.). The capped solution was left stirring at rt for 20 hrs. The solution was partitioned between EtOAc and H₂O. The organic layer was washed with H₂O, Na₂CO₃(sat.), NaCl(sat.), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography to yield S-(1S,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl ethanethioate in 35% yield. LC/MS (m/z)=615.2 (MH⁺), R_t=0.95 min.

Synthesis of tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(methylthio)cyclohexylcarbamate




To a solution of S-(1S,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl ethanethioate (1.0 equiv.) in MeOH (0.07 M) was added K₂CO₃ (3.0 equiv.). The heterogeneous solution was capped and left stirring at rt for 1 hr. Methyl iodide (1.5 equiv.) was added and stirred at rt for 10

82


min. The volatiles were removed in vacuo and the residue was partitioned between EtOAc and H₂O. The organic layer was washed with NaCl(sat.), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography to yield tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(methylthio)cyclohexylcarbamate in 78% yield. LC/MS (m/z)=587.2 (MH⁺), R_t=1.00 min.

Synthesis of N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylthio)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide




To a solution of HCl (30.0 equiv.) in dioxane was added tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(methylthio)cyclohexylcarbamate (1.0 equiv.). The solution was capped and left standing at rt for 1 hr. The volatiles were removed in vacuo, dissolved in DMSO and purified by reverse phase HPLC to yield N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylthio)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 94% yield. LC/MS (m/z)=487.2 (MH⁺), R_t=0.65 min. ¹H NMR (400 MHz, <dmsO>) δ ppm 10.44 (s, 1H), 8.58 (d, J=4.0, 1H), 8.47 (d, J=4.0, 1H), 8.34 (dd, J=8.0, 4.0, 1H), 8.20 (dd, J=8.0, 8.0, 1H), 8.20 (dd, J=16.0, 4.0, 1H), 7.67-7.74 (m, 1H), 7.36 (dd, J=8.0, 8.0, 2H), 7.26 (d, J=4.0, 1H), 2.85-2.95 (m, 2H), 2.18 (s, 3H), 1.88-1.98 (m, 1H), 1.74-1.84 (m, 1H), 1.48-1.56 (m, 1H), 1.38-1.48 (m, 1H), 1.18-1.28 (m, 1H), 1.02 (d, J=8.0, 3H).

Synthesis of tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-((R)-methylsulfinyl)cyclohexylcarbamate and tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(methylsulfonyl)cyclohexylcarbamate




83

-continued



To a solution of tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(methylthio)cyclohexylcarbamate (1.0 equiv.) in CH_2Cl_2 (0.10 M) in a 0°C . bath was added mCPBA (1.2 equiv.). The capped solution was left stirring at rt for 1 hr. cyclohexene (10.0 equiv.) was added to quench any remaining mCPBA and after stirring for 5 minutes the solution was directly loaded onto ISCO SiO_2 column and purified to yield tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-((R)-methylsulfinyl)cyclohexylcarbamate in 31% yield, LC/MS=603.2 (MH⁺), Rt=0.66 min; and tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(methylsulfonyl)cyclohexylcarbamate in 37% yield. LC/MS (m/z)=619.2 (MH⁺), R_f =0.88 min.

Synthesis of N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-((R)-methylsulfinyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide




To a solution of tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-((R)-methylsulfinyl)cyclohexylcarbamate in DCM (0.10 M) was added TFA (30.0 equiv.). The solution was capped and left standing at rt for 1 hr. The volatiles were removed in vacuo; the residue was dissolved in DMSO and purified by reverse phase HPLC. The product fractions were lyophilized directly to yield N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-((R)-methylsulfinyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 54% yield. LC/MS (m/z)=503.2 (MH⁺), Rt=0.57 min. ^1H NMR (400 MHz, <dms>) δ ppm 10.48 (s, 1H), 8.58 (d, J=4.0, 1H), 8.50 (d, J=4.0, 1H), 8.34 (dd, J=8.0, 4.0, 1H), 8.20 (dd, J=8.0, 8.0, 1H), 8.20 (dd, J=16.0, 4.0, 1H), 7.67-7.74 (m, 1H), 7.42 (d, J=4.0, 1H), 7.36 (dd, J=8.0, 8.0, 2H), 3.40-3.42 (m, 1H),

84


3.06-3.20 (m, 1H), 2.92 (s, 3H), 2.06-2.20 (m, 1H), 1.95-2.04 (m, 2H), 1.70-1.80 (m, 1H), 1.56-1.70 (m, 1H), 0.86 (d, J=8.0, 3H).

Synthesis of N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide



To a solution of tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(methylsulfonyl)cyclohexylcarbamate in DCM (0.02 M) was added TFA (30.0 equiv.). The solution was capped and left standing at rt for 1 hr. The volatiles were removed in vacuo; the residue was dissolved in DMSO and purified by reverse phase HPLC. The product fractions were lyophilized directly to yield N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 73% yield. LC/MS (m/z)=519.2 (MH⁺), R_f =0.58 min. ^1H NMR (400 MHz, <dms>) δ ppm 10.52 (s, 1H), 8.51-8.54 (m, 2H), 8.34 (dd, J=8.0, 4.0, 1H), 8.20 (dd, J=8.0, 8.0, 1H), 8.15 (dd, J=16.0, 4.0, 1H), 7.67-7.74 (m, 1H), 7.36 (dd, J=8.0, 8.0, 2H), 7.28 (d, J=4.0, 1H), 3.77-3.79 (m, 1H), 3.18 (s, 3H), 3.02-3.20 (m, 1H), 1.94-2.40 (m, 5H), 1.57-1.62 (m, 1H), 1.24 (d, J=8.0, 3H).

Synthesis of tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-2-(2-methoxyethylthio)-3-methylcyclohexylcarbamate




To a solution of S-(1S,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl ethanethioate (1.0 equiv.) in MeOH (0.04 M) was added K_2CO_3 (3.0 equiv.). The heterogeneous solution was capped and left stirring at rt for 1 hr. 1-bromo-2-methoxyethane (7.0 equiv.) was added and stirred at rt for 6 hrs. Quench the reaction with diisopropylamine and

85


the volatiles were removed in vacuo and the residue was partitioned between EtOAc and H₂O. The organic layer was washed with NaCl_(sat.), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography to yield tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-2-(2-methoxyethylthio)-3-methylcyclohexylcarbamate in 63% yield. LC/MS (m/z)=631.2 (MH⁺), R_t=0.98 min.

Synthesis of N-(4-((1R,3R,4S,5S)-3-amino-4-((2-methoxyethyl)sulfonyl)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide



To a solution of tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-2-(2-methoxyethylthio)-3-methylcyclohexylcarbamate (1.0 equiv.) in THF (0.04 M) in a 0° C. bath was added oxone (2.0 equiv.) as a solution in H₂O. The solution was left stirring at rt for 5 hrs. The solution was diluted with EtOAc, washed with H₂O, NaCl_(sat.), dried over MgSO₄, filtered, concentrated to yield Boc protected product. The material was treated with 25% TFA/CH₂Cl₂ for 30 minutes, at which time the volatiles were removed in vacuo, the residue was dissolved in DMSO and purified by reverse phase HPLC. The product fractions were lyophilized directly to yield N-(4-((1R,3R,4S,5S)-3-amino-4-((2-methoxyethyl)sulfonyl)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 76% yield. LC/MS (m/z)=563.3 (MH⁺), R_t=0.62 min. ¹H NMR (400 MHz, <dms>) δ ppm 10.52 (s, 1H), 8.53 (s, 1H), 8.50 (d, J=4.0, 1H), 8.35 (dd, J=8.0, 4.0, 1H), 8.20 (t, J=8.0, 1H), 7.80 (bs, 2H), 7.70 (quintet, J=8.0, 1H), 7.36 (t, J=8.0, 2H), 7.31 (d, J=4.0, 1H), 3.82-3.88 (m, 2H), 3.73-3.78 (m, 2H), 3.55-3.61 (m, 1H), 3.42-3.49 (m, 1H), 3.31 (s, 3H), 3.06-3.16 (m, 1H), 1.92-2.18 (m, 4H), 1.54-1.64 (m, 1H), 1.22 (d, J=4.0, 3H).


Synthesis of tert-butyl (1R,2S,3S,5R)-2-(2-(tert-butyldimethylsilyloxy)ethylthio)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexylcarbamate



86

To a solution of S-(1S,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl ethanethioate (1.0 equiv.) in MeOH (0.04 M) was added K₂CO₃ (5.0 equiv.). The heterogeneous solution was capped and left stirring at rt for 1 hr. (2-bromoethoxy)(tert-butyl)dimethylsilane (7.0 equiv.) was added and stirred at rt for 6 hrs. Quench the reaction with diisopropylamine and the volatiles were removed in vacuo and the residue was partitioned between EtOAc and H₂O. The organic layer was washed with NaCl_(sat.), dried over MgSO₄, filtered, concentrated. The residue was dissolved in DMF (0.05 M) and imidazole (20.0 equiv.) and TBDMSCl (7.0 equiv.) were added. After stirring at rt for 1 hr, the solution was partitioned between EtOAc and H₂O. The organic layer was washed with NaCl_(sat.), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography to yield tert-butyl (1R,2S,3S,5R)-2-(2-(tert-butyldimethylsilyloxy)ethylthio)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexylcarbamate in 57% yield. LC/MS (m/z)=731.4 (MH⁺), R_t=1.27 min.

Synthesis of N-(4-((1R,3R,4S,5S)-3-amino-4-(2-hydroxyethylsulfonyl)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide




To a solution of tert-butyl (1R,2S,3S,5R)-2-(2-(tert-butyldimethylsilyloxy)ethylthio)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexylcarbamate (1.0 equiv.) in THF (0.04 M) in a 0° C. bath was added oxone (2.0 equiv.) as a solution in H₂O. The solution was left stirring at rt for 48 hrs. The solution was diluted with EtOAc, washed with H₂O, Na₂CO_{3(sat.)}, NaCl_(sat.), dried over MgSO₄, filtered, concentrated to yield Boc protected product. The material was treated with 25% TFA/CH₂Cl₂ for 30 minutes, at which time the volatiles were removed in vacuo, the residue was dissolved in DMSO and purified by reverse phase HPLC. The product fractions were lyophilized directly to N-(4-((1R,3R,4S,5S)-3-amino-4-(2-hydroxyethylsulfonyl)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 44% yield. LC/MS (m/z)=549.6 (MH⁺), R_t=0.58 min. ¹H NMR (400 MHz, <dms>) δ ppm 10.51 (s, 1H), 8.53 (s, 1H), 8.51 (d, J=4.0, 1H), 8.33 (dd, J=8.0, 4.0, 1H), 8.20 (t, J=8.0, 1H), 8.09 (broad doublet, J=4.0, 2H), 7.65-7.75 (m, 1H), 7.36 (t, J=8.0, 2H), 7.30 (d, J=8.0, 1H), 3.84-4.02 (m, 3H), 3.58-3.68 (m,

87


1H), 3.43-3.53 (m, 1H), 3.28-3.36 (m, 1H), 3.04-3.14 (m, 1H), 1.92-2.18 (m, 4H), 1.56-1.63 (m, 1H), 1.24 (d, J=8.0, 3H).

Synthesis of S-(1R,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl ethanethioate



To a solution of (1S,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl methanesulfonate (1.0 equiv.) in DMF (0.25 M) was added KSAc (6.0 equiv.). The capped solution was left stirring at rt for 20 hrs. The solution was partitioned between EtOAc and H₂O. The organic layer was washed with H₂O, Na₂CO_{3(sat.)}, NaCl_(sat.), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography to yield S-(1R,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl ethanethioate in 50% yield. LC/MS (m/z)=615.2 (MH⁺), R_t=0.96 min.

Synthesis of tert-butyl (1R,2R,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(methylthio)cyclohexylcarbamate




To a solution of S-(1R,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl ethanethioate (1.0 equiv.) in MeOH (0.07 M) was added K₂CO₃ (3.0 equiv.). The heterogeneous solution was capped and left stirring at rt for 1 hr. Methyl iodide (1.5 equiv.) was added and stirred at rt for 10 min. The volatiles were removed in vacuo and the residue was partitioned between EtOAc and H₂O. The organic layer was washed with NaCl_(sat.), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography to yield tert-butyl (1R,2R,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-

88


fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(methylthio)cyclohexylcarbamate in 72% yield. LC/MS (m/z)=587.2 (MH⁺), R_t=0.96 min.

5 Synthesis of N-(4-((1R,3R,4R,5S)-3-amino-5-methyl-4-(methylthio)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide



To a solution of HCl (30.0 equiv.) in dioxane was added to tert-butyl (1R,2R,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(methylthio)cyclohexylcarbamate (1.0 equiv.). The solution was capped and left standing at rt for 1 hr. The volatiles were removed in vacuo, dissolved in DMSO and purified by reverse phase HPLC to yield N-(4-((1R,3R,4R,5S)-3-amino-5-methyl-4-(methylthio)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 43% yield. LC/MS (m/z)=487.2 (MH⁺), R_t=0.65 min.


40 Synthesis of N-(4-((1R,3R,4R,5S)-3-amino-5-methyl-4-((S)-methylsulfinyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide and N-(4-((1R,3R,4R,5S)-3-amino-5-methyl-4-((R)-methylsulfinyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide



89

To a solution of tert-butyl ((1R,2R,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(methylthio)cyclohexylcarbamate (1.0 equiv.) in THF (0.05 M) in a 0° C. bath was added the oxone (1.0 equiv.) as a solution in H₂O. The solution was left stirring at 0° C. for 5 mins. The solution was diluted with EtOAc, washed with H₂O, NaCl_(sat.), dried over MgSO₄, filtered, concentrated to yield Boc protected product. The material was treated with 25% TFA/CH₂Cl₂ for 30 minutes, at which time the volatiles were removed in vacuo, the residue was dissolved in DMSO and purified by reverse phase HPLC. The product fractions were lyophilized directly to yield N-(4-((1R,3R,4R,5S)-3-amino-5-methyl-4-((S)-methylsulfinyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 33% yield, LC/MS=503.2 (MH⁺), Rt=0.58 min; and N-(4-((1R,3R,4R,5S)-3-amino-5-methyl-4-((R)-methylsulfinyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 37% yield. LC/MS (m/z)=503.2 (MH⁺), R_f=0.60 min.

Synthesis of N-(4-((1R,3R,4R,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide




To a solution of tert-butyl ((1R,2R,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(methylthio)cyclohexylcarbamate (1.0 equiv.) in THF (0.04 M) in a 0° C. bath was added the oxone (2.0 equiv.) as a solution in H₂O. The solution was left stirring at rt for 5 hrs. The solution was diluted with EtOAc, washed with H₂O, NaCl_(sat.), dried over MgSO₄, filtered, concentrated to yield Boc protected product. The material was treated with 25% TFA/CH₂Cl₂ for 30 minutes, at which time the volatiles were removed in vacuo, the residue was dissolved in DMSO and purified by reverse phase HPLC. The product fractions were lyophilized directly to yield N-(4-((1R,3R,4R,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 68% yield. LC/MS (m/z)=519.2 (MH⁺), R_f=0.60 min. ¹H NMR (400 MHz, <dmso>) δ ppm 10.44 (s, 1H), 8.60 (d, J=8.0, 1H), 8.48 (d, J=4.0, 1H), 8.34 (dd, J=8.0, 4.0, 1H), 8.20 (dd, J=8.0, 8.0, 1H), 8.00 (dd, J=16.0, 4.0, 1H), 7.67-7.74 (m, 1H), 7.34-7.38

90


(m, 3H), 3.14 (s, 3H), 3.02-3.12 (m, 2H), 2.18-2.24 (m, 1H), 1.84-1.96 (m, 3H), 1.62-1.72 (m, 1H), 1.38-1.48 (m, 1H), 1.18 (d, J=4.0, 3H).

Synthesis of (+/-)-tert-butyl ((1R,2R,3S,5R)-5-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-2-hydroxy-3-methylcyclohexyl)carbamate



A solution of (+/-)-(1R,2R,4R,6S)-4-(3-aminopyridin-4-yl)-2-(tert-butoxycarbonylamino)-6-methylcyclohexyl acetate (1.0 equiv.) and Boc₂O (2.1 equiv.) in dioxane (0.34 M) was submerged in an 120° C. oil bath, fitted with a condenser and left stirring under Ar for 6 hrs. The reaction was cooled to rt and the volatiles were removed in vacuo. The residue was dissolved in EtOH (0.34 M), K₂CO₃ (10.0 equiv.) was added, a refluxing head was attached and the heterogeneous solution was submerged in an 50° C. oil bath and left stirring for 24 hrs. The reaction was cooled to rt. The volatiles were removed in vacuo and the residue was partitioned between EtOAc and H₂O. The organic layer was washed with Na₂CO₃(sat.), NaCl_(sat.), dried over MgSO₄, filtered, and concentrated. The residue was dissolved in CH₂Cl₂/Heptane and left standing. The solid that form was sonicated, filtered, rinsed with CH₂Cl₂ and pumped on to yield (+/-)-tert-butyl ((1R,2R,3S,5R)-5-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-2-hydroxy-3-methylcyclohexyl)carbamate in 85% yield. LC/MS (m/z)=422.3 (MH⁺), R_f=0.65 min.

Synthesis of (+/-)-(1R,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(tert-butoxycarbonylamino)pyridin-4-yl)-6-methylcyclohexyl methanesulfonate




To a solution of (+/-)-tert-butyl ((1R,2R,3S,5R)-5-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-2-hydroxy-3-methylcyclohexyl)carbamate (1.0 equiv.) in pyridine (0.17 M) was added MsCl (5.0 equiv.). The capped solution was stirred for 5 minutes and then the homogeneous solution was left standing at rt for 16 hrs. The volatiles were removed in vacuo and the residue was partitioned between EtOAc and

91


H₂O. The organic layer was washed with H₂O, 10% CuSO₄, H₂O, Na₂CO₃(*sat.*), NaCl(*sat.*), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography to yield (+/-)-(1R,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(tert-butoxycarbonylamino)pyridin-4-yl)-6-methylcyclohexyl methanesulfonate in 59% yield. LC/MS (m/z)=500.3 (MH⁺), R_t=0.74 min.

Synthesis of (+/-)S-((1S,2R,4R,6S)-2-((tert-butoxycarbonylamino)-4-(3-((tert-butoxycarbonylamino)pyridin-4-yl)-6-methylcyclohexyl)ethanethioate



To a solution of (+/-)-(1R,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(tert-butoxycarbonylamino)pyridin-4-yl)-6-methylcyclohexyl methanesulfonate (1.0 equiv.) in DMF (0.25 M) was added potassium thioacetate (6.0 equiv.). The mixture was stirred for 6 hours in a 60° C. bath under Ar. Upon cooling and the residue was partitioned between EtOAc and H₂O. The organic layer was washed with H₂O (3x), Na₂CO₃(*sat.*), NaCl(*sat.*), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography to yield (+/-)S-((1S,2R,4R,6S)-2-((tert-butoxycarbonylamino)-4-(3-((tert-butoxycarbonylamino)pyridin-4-yl)-6-methylcyclohexyl)ethanethioate in 87% yield. LC/MS (m/z)=480.3 (MH⁺), R_t=0.82 min.

Synthesis of (+/-)tert-butyl ((1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-3-methyl-2-(methylsulfonyl)cyclohexyl)carbamate




To a solution of (+/-)S-((1S,2R,4R,6S)-2-((tert-butoxycarbonylamino)-4-(3-((tert-butoxycarbonylamino)pyridin-4-yl)-6-methylcyclohexyl)ethanethioate (1.0 equiv.) in MeOH (0.09 M) was added potassium carbonate (3.0 equiv.). The mixture was stirred for 15 minutes at which time methyl

92

iodide (1.1 eq.) was added and the solution was stirred at rt for 15 minutes. The volatiles were removed in vacuo and the residue was partitioned between EtOAc and H₂O. The organic layer was washed with H₂O, NaCl(*sat.*), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography to yield the methyl sulfide product in 99% yield. LC/MS (m/z)=452.3 (MH⁺), R_t=0.87 min. To a solution of methyl sulfide (1.0 eq.) in THF (0.05 M) at rt was added an aqueous solution of oxone (2.2 eq.) dropwise over 10 minutes. After stirring at rt for 1 hour the solution was partitioned between EtOAc and H₂O. The organic layer was washed with H₂O, NaCl(*sat.*), dried over MgSO₄, filtered, concentrated to yield the bis Boc protected methyl sulfone product in 95% yield. LC/MS (m/z)=484.2 (MH⁺), R_t=0.77 min. The bis boc protected cyclohexyl sulfone (1.0 equiv) was treated with 4M HCl in dioxane for 3 hours to removed both Boc groups. Upon removal of the volatiles in vacuo, the residue was suspended in 1:1 dioxane/Na₂CO₃(*sat.*) and N-(tert-Butoxycarbonyloxy)succinimide (1.2 eq.) was added. After stirring for 1 hour, additional N-(tert-Butoxycarbonyloxy)succinimide (1.2 eq.) was added. After stirring for an additional 2 hours the solution was extracted with CH₂Cl₂, the combined organic layers were washed with H₂O, NaCl(*sat.*), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography to yield the (+/-)tert-butyl ((1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-3-methyl-2-(methylsulfonyl)cyclohexyl)carbamate in 56% yield. LC/MS (m/z)=384.3 (MH⁺), R_t=0.57 min. Chiral purification was completed via SFC (20% EtOH/80% n-heptanes, 20 mL/min, OJ column) to isolate the pure enantiomers. The second peak correlated with tert-butyl ((1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-3-methyl-2-(methylsulfonyl)cyclohexyl)carbamate.

Synthesis of tert-butyl (1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-3-methyl-2-(1H-1,2,4-triazol-1-yl)cyclohexylcarbamate and tert-butyl (1S,2R,3R,5S)-5-(3-aminopyridin-4-yl)-3-methyl-2-(1H-1,2,4-triazol-1-yl)cyclohexylcarbamate




A solution of (+/-)-(1R,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(tert-butoxycarbonylamino)pyridin-4-yl)-6-methylcyclohexyl methanesulfonate (1.0 equiv.), 1H-1,2,4-triazole (3.0 equiv.) and Cs₂CO₃ (3.0 equiv.) in DMF (0.15 M) was stirred at 80° C. for 5 hrs. The volatiles were removed in vacuo and the residue was partitioned between EtOAc and H₂O. The organic layer was washed with Na₂CO₃(*sat.*), NaCl(*sat.*), dried over MgSO₄, filtered, concentrated and purified by RP HPLC, followed by free basing by partitioning

93


between an equal volume of EtOAc and Na₂CO₃, separating, washing with NaCl_(sat.), drying over MgSO₄, filtering, concentrating. Purification was completed via SFC (20% MeOH, 100 mL/min, AD column) to yield tert-butyl (1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-3-methyl-2-(1H-1,2,4-triazol-1-yl)cyclohexylcarbamate (19% yield, 99% ee) and tert-butyl (1S,2R,3R,5S)-5-(3-aminopyridin-4-yl)-3-methyl-2-(1H-1,2,4-triazol-1-yl)cyclohexylcarbamate (19% yield, 99% ee). LC/MS (m/z)=373.3 (MH⁺), R_t=0.54 min.

Synthesis of tert-butyl tert-butoxycarbonyl(4-((3R,4R,5S)-3-((tert-butoxycarbonyl)amino)-4-((tert-butyl)dimethylsilyloxy)-5-methylpiperidin-1-yl)pyridin-3-yl)carbamate



To a solution of tert-butyl (3R,4R,5S)-1-(3-aminopyridin-4-yl)-4-(tert-butyl)dimethylsilyloxy)-5-methylpiperidin-3-ylcarbamate (1.0 equiv.) in CH₂Cl₂ (0.50 M) at RT was added Boc₂O (6.0 equiv.), followed by DMAP (2.0 equiv.). The resulting mixture was stirred at RT for 16 hrs. The reaction mixture was then diluted with EtOAc and water. The organic layer was washed with Brine, dried over MgSO₄, concentrated and purified by flash column chromatography to yield tert-butyl tert-butoxycarbonyl(4-((3R,4R,5S)-3-((tert-butoxycarbonyl)amino)-4-((tert-butyl)dimethylsilyloxy)-5-methylpiperidin-1-yl)pyridin-3-yl)carbamate in 57% yield. LC/MS (m/z)=637.3 (MH⁺), R_t=1.17 min.

Synthesis of tert-butyl tert-butoxycarbonyl(4-((3R,4R,5S)-3-((tert-butoxycarbonyl)amino)-4-hydroxy-5-methylpiperidin-1-yl)pyridin-3-yl)carbamate




To a solution of tert-butyl tert-butoxycarbonyl(4-((3R,4R,5S)-3-((tert-butoxycarbonyl)amino)-4-((tert-butyl)dimethylsilyloxy)-5-methylpiperidin-1-yl)pyridin-3-yl)carbamate (1.0 equiv.) in THF (0.20 M) at RT was added TBAF (1.0 equiv.). The resulting mixture was stirred at rt for 4 hrs. The reaction mixture was then diluted with EtOAc and water. The organic layer was washed with Brine, dried over MgSO₄,

94


concentrated and purified by flash column chromatography to yield tert-butyl tert-butoxycarbonyl(4-((3R,4R,5S)-3-((tert-butoxycarbonyl)amino)-4-hydroxy-5-methylpiperidin-1-yl)pyridin-3-yl)carbamate in 87% yield. LC/MS (m/z)=523.4 (MH⁺), R_t=0.72 min. ¹H NMR (400 MHz, <cdcl3>) δ ppm 1.03 (d, J=6.65 Hz, 3H), 1.34-1.51 (m, 54H), 1.72-1.87 (m, 1H), 2.05 (s, 1H), 2.46-2.57 (m, 1H), 2.69 (t, J=11.35 Hz, 1H), 2.78-2.94 (m, 1H), 3.00-3.14 (m, 1H), 3.45 (d, J=12.52 Hz, 1H), 3.53-3.76 (m, 1H), 4.63 (d, J=6.26 Hz, 1H), 6.88 (d, J=5.48 Hz, 3H), 8.13 (s, 3H), 8.26-8.36 (m, 3H).

Synthesis of (3R,4R,5S)-1-(3-(bis(tert-butoxycarbonyl)amino)pyridin-4-yl)-3-((tert-butoxycarbonyl)amino)-5-methylpiperidin-4-yl methanesulfonate



To a solution of tert-butyl tert-butoxycarbonyl(4-((3R,4R,5S)-3-((tert-butoxycarbonyl)amino)-4-hydroxy-5-methylpiperidin-1-yl)pyridin-3-yl)carbamate (1.0 equiv.) in DCM (0.20 M) was added TEA (1.7 equiv.), followed by MsCl (1.3 equiv.). The capped solution was stirred at rt for 90 mins. The reaction mixture was quenched with NaHCO_{3(sat.)}, and extracted with EtOAc. The organic layer was washed with NaCl_(sat.), dried over MgSO₄, filtered, concentrated to yield (3R,4R,5S)-1-(3-(bis(tert-butoxycarbonyl)amino)pyridin-4-yl)-3-((tert-butoxycarbonyl)amino)-5-methylpiperidin-4-yl methanesulfonate in 99% yield. LC/MS (m/z)=601.3 (MH⁺), R_t=0.83 min.

Synthesis of tert-butyl (3R,4S,5S)-1-(3-aminopyridin-4-yl)-5-methyl-4-(1H-1,2,4-triazol-1-yl)piperidin-3-ylcarbamate




A solution of (3R,4R,5S)-3-(tert-butoxycarbonylamino)-1-(3-(tert-butoxycarbonylamino)pyridin-4-yl)-5-methylpiperidin-4-yl methanesulfonate (1.0 equiv.), 1H-1,2,4-triazole (3.0 equiv.) and Cs₂CO₃ (3.0 equiv.) in DMF (0.17 M) was stirred at 90° C. for 3 hrs. The mixture was diluted with DMF,

95

filtered and purified by RP HPLC, followed by free basing by partitioning between an equal volume of EtOAc and Na₂CO₃, separating, washing with NaCl_(sat.), drying over MgSO₄, filtering, concentrating to yield tert-butyl (3R,4S,5S)-1-(3-aminopyridin-4-yl)-5-methyl-4-(1H-1,2,4-triazol-1-yl)piperidin-3-ylcarbamate in 8% yield. LC/MS (m/z)=374.3 (MH⁺), R_t=0.53 min.

Synthesis of tert-butyl ((1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-3-methyl-2-(2-oxopyridin-1(2H)-yl)cyclohexyl)carbamate, tert-butyl ((1S,2R,3R,5S)-5-(3-aminopyridin-4-yl)-3-methyl-2-(2-oxopyridin-1(2H)-yl)cyclohexyl)carbamate, tert-butyl ((1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-3-methyl-2-(pyridin-2-yloxy)cyclohexyl)carbamate and tert-butyl ((1S,2R,3R,5S)-5-(3-aminopyridin-4-yl)-3-methyl-2-(pyridin-2-yloxy)cyclohexyl)carbamate




To a solution of (1R,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(tert-butoxycarbonylamino)pyridin-4-yl)-6-methylcyclohexyl methanesulfonate (1.0 equiv.) Cs₂CO₃ (3.0 equiv.) in DMF (0.15 M) was added pyridin-2-ol (1.0 equiv.) 55 The mixture was stirred at 70° C. for 5 hrs. The reaction was worked up with EtOAc and Brine. The organic layer was concentrated and was treated with 4N HCl (30.0 equiv.) in Dioxane for 2 hrs at which time the volatiles were removed in vacuo. The residue was dissolved in THF (0.15 M) and tert-butyl 2,5-dioxopyrrolidin-1-yl carbonate (1.5 equiv.) was added, followed by DIEA (3.0 equiv.). After stirring at rt for 3 hrs, the reaction was quenched with sat. NaHCO₃ and extracted with EtOAc. The organic layer was washed with Brine, dried over Na₂SO₄ and concentrated. The crude was purified by prep HPLC to yield two major peaks. The fractions of the first product peak was combined and neutralized

96

with sat. NaHCO₃ and extracted with EtOAc. The organic layer was washed with Brine, dried over Na₂SO₄ and concentrated. Purification was completed via SFC (20% (MeOH with 10% DEA), 100 mL/min, AD column) to yield tert-butyl ((1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-3-methyl-2-(2-oxopyridin-1(2H)-yl)cyclohexyl)carbamate (8% yield, 99% ee) and tert-butyl ((1S,2R,3R,5S)-5-(3-aminopyridin-4-yl)-3-methyl-2-(2-oxopyridin-1(2H)-yl)cyclohexyl)carbamate (9% yield, 99% ee). LC/MS (m/z)=399.2 (MH⁺), R_t=0.60 min. ¹H NMR (400 MHz, <dms>) δ ppm 0.72 (s, 3H), 1.30 (s, 9H), 1.63-1.81 (m, 2H), 1.89-2.00 (m, 2H), 2.02-2.18 (m, 2H), 3.04-3.13 (m, 1H), 4.03-4.12 (m, 1H), 4.91-5.02 (m, 1H), 5.04-5.13 (m, 2H), 6.17-6.26 (m, 1H), 6.31-6.41 (m, 1H), 6.98-7.07 (m, 1H), 7.32-7.41 (m, 2H), 7.68-7.74 (m, 1H), 7.74-7.82 (m, 1H), 7.85-7.90 (m, 1H). The fractions of the second product peak was combined and neutralized with sat. NaHCO₃ and extracted with EtOAc. The organic layer was washed with Brine, dried over Na₂SO₄ and concentrated. Purification was completed via SFC (20% (MeOH with 10% DEA), 100 mL/min, AD column) to yield tert-butyl ((1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-3-methyl-2-(pyridin-2-yloxy)cyclohexyl)carbamate (14% yield, 99% ee) and tert-butyl ((1S,2R,3R,5S)-5-(3-aminopyridin-4-yl)-3-methyl-2-(pyridin-2-yloxy)cyclohexyl)carbamate (15% yield, 99% ee). LC/MS (m/z)=399.2 (MH⁺), R_t=0.65 min. ¹H NMR (400 MHz, <cdcl3>) δ ppm 0.95 (d, J=6.65 Hz, 3H), 1.35-1.78 (m, 12H), 1.97-2.08 (m, 2H), 2.76 (t, J=11.93 Hz, 1H), 3.72 (br. s., 2H), 3.88 (d, J=4.70 Hz, 1H), 5.43 (d, J=7.43 Hz, 1H), 5.59 (br. s., 1H), 6.79-6.94 (m, 2H), 7.08 (d, J=5.09 Hz, 1H), 7.56-7.65 (m, 1H), 7.98-8.17 (m, 3H).

Synthesis of tert-butyl ((1R,2S,3S,5R)-5-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-2-azido-3-methylcyclohexyl)carbamate and tert-butyl ((1S,2R,3R,5S)-5-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-2-azido-3-methylcyclohexyl)carbamate




To a solution of (+/-)-(1R,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(tert-butoxycarbonylamino)pyridin-4-yl)-6-methylcyclohexyl methanesulfonate (1.0 equiv.) in DMF (0.13 M) was added NaN₃ (1.0 equiv.) 55 The solution was submerged in an 80° C. oil bath and left stirring under Ar for 16 hrs. The solution was cooled to rt and left stirring under Ar overnight. The solution was partitioned between EtOAc and H₂O. The organic layer was washed with Na₂CO₃(sat.), NaCl_(sat.), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography. Purification was completed via SFC (15% IPA, 100 mL/min, IA column) to yield tert-butyl ((1R,2S,3S,5R)-5-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-2-azido-3-methylcyclohexyl)carbamate (21% yield, 99% ee) and tert-butyl ((1S,2R,3R,5S)-5-(3-

97


((tert-butoxycarbonyl)amino)pyridin-4-yl)-2-azido-3-methylcyclohexyl)carbamate (22% yield, 99% ee). LC/MS (m/z)=447.3 (MH^+), R_f =0.86 min.

Synthesis of tert-butyl (1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-2-azido-3-methylcyclohexylcarbamate



A solution of 4 M HCl in dioxane (30.0 equiv.) was added to tert-butyl ((1R,2S,3S,5R)-5-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-2-azido-3-methylcyclohexyl)carbamate (1.0 equiv.). The solution started to go homogeneous for a few minutes, but then a ppt formed and the solution went very thick. After sitting at rt for 1 hour, the volatiles were removed in vacuo and the solid was pumped on for 5 minutes on the high vac. To the residue was added CH_2Cl_2 (0.15 M), TEA (5.0 equiv.) and Boc_2O (1.0 equiv.). The solution was left stirring at rt for 1 hr. The volatiles were removed in vacuo and the residue was partitioned between EtOAc and H_2O . The organic layer was washed with Na_2CO_3 (sat.), $NaCl$ (sat.), dried over $MgSO_4$, filtered, concentrated and purified by ISCO SiO_2 chromatography to yield tert-butyl (1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-2-azido-3-methylcyclohexylcarbamate in 57% yield. LC/MS (m/z)=347.3 (MH^+), R_f =0.70 min. 1H NMR (400 MHz, $CHLOROFORM-d$) δ ppm 1.13 (d, J =6.65 Hz, 3H), 1.43-1.58 (m, 11H), 1.79 (d, J =12.52 Hz, 1H), 1.95 (d, J =6.26 Hz, 1H), 2.60 (br. s., 1H), 3.61 (br. s., 2H), 3.77-3.91 (m, 2H), 4.78 (d, J =7.43 Hz, 1H), 6.96 (d, J =4.70 Hz, 1H), 7.97-8.07 (m, 2H).

Synthesis of tert-butyl (1S,2R,3R,5S)-5-(3-aminopyridin-4-yl)-2-azido-3-methylcyclohexylcarbamate




A solution of 4 M HCl in dioxane (30.0 equiv.) was added to tert-butyl ((1S,2R,3R,5S)-5-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-2-azido-3-methylcyclohexyl)carbamate (1.0 equiv.). The solution started to go homogeneous for a few minutes, but then a ppt formed and the solution went very thick. After sitting at rt for 1 hour, the volatiles were removed in vacuo and the solid was pumped on for 5 minutes on the

98


high vac. To the residue was added CH_2Cl_2 (0.15 M), TEA (5.0 equiv.) and Boc_2O (1.0 equiv.). The solution was left stirring at rt for 1 hr. The volatiles were removed in vacuo and the residue was partitioned between EtOAc and H_2O . The organic layer was washed with Na_2CO_3 (sat.), $NaCl$ (sat.), dried over $MgSO_4$, filtered, concentrated to yield tert-butyl (1S,2R,3R,5S)-5-(3-aminopyridin-4-yl)-2-azido-3-methylcyclohexylcarbamate in 98% yield. LC/MS (m/z)=347.3 (MH^+), R_f =0.71 min.

Synthesis of tert-butyl (1S,2R,3R,5S)-2-azido-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexylcarbamate



EDC (2.0 equiv.) and HOAt (2.0 equiv.) was added to a solution of tert-butyl (1S,2R,3R,5S)-5-(3-aminopyridin-4-yl)-2-azido-3-methylcyclohexyl carbamate (1.0 equiv.) and 6-(2,6-difluorophenyl)-5-fluoropicolinic acid (1.5 equiv.) in DMF (0.08 M). The mixture was stirred at ambient temperature overnight. The reaction mixture was diluted with water and extracted with ethyl acetate. The combined extracts were washed sequentially with 1M aqueous sodium hydroxide and brine, dried over sodium sulfate, filtered, concentrated and purified by ISCO SiO_2 chromatography to yield tert-butyl (1S,2R,3R,5S)-2-azido-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexyl carbamate in 31% yield. LC/MS (m/z)=582.3 (MH^+), R_f =1.00 min.

Synthesis of tert-butyl (1R,2S,3S,5R)-2-azido-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexylcarbamate




EDC (2.0 equiv.) and HOAt (2.0 equiv.) was added to a solution of tert-butyl (1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-2-azido-3-methylcyclohexylcarbamate (1.0 equiv.) and 6-(2,6-difluorophenyl)-5-fluoropicolinic acid (1.5 equiv.) in DMF (0.08 M). The mixture was stirred at ambient temperature overnight. The reaction mixture was diluted with water

99

and extracted with ethyl acetate. The combined extracts were washed sequentially with 1M aqueous sodium hydroxide and brine, dried over sodium sulfate, filtered, concentrated and purified by ISCO SiO₂ chromatography to yield tert-butyl (1R,2S,3S,5R)-2-amino-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexylcarbamate 59% yield. LC/MS (m/z)=582.3 (MH⁺), R_t=0.97 min.


Synthesis of tert-butyl (1R,2S,3S,5R)-2-amino-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexylcarbamate



Degass a solution of tert-butyl (1R,2S,3S,5R)-2-amino-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexylcarbamate (1.0 equiv.) in MeOH/EtOAc (3/1, 0.04 M). To this solution was added Pd/C (0.2 equiv.) and purge with Ar and H₂. The mixture was stirred under H₂ for 16 hrs. Filter the mixture over celite and wash the cake with MeOH. Concentrate the filtrate to yield tert-butyl (1R,2S,3S,5R)-2-amino-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexylcarbamate in 100% yield. LC/MS (m/z)=556.3 (MH⁺), R_t=0.73 min.

Method 4

Synthesis of ethyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate




To a solution of tert-butyl (1R,2S,3S,5R)-2-amino-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexylcarbamate (1.0 equiv.) in CH₂Cl₂ (0.03 M) at 0° C. was added DIEA (3.0 equiv.) and then ETHYL CHLOROFORMATE (1.0 equiv.). The homogeneous solution was left standing at 0° C. at for 4 hrs. Neutralize the

100


reaction with sat. NaHCO₃ solution. The solution was partitioned between EtOAc and H₂O. The organic layer was washed with Na₂CO₃(sat.), NaCl(sat.), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography. The Boc protected product was treated with 25% TFA/CH₂Cl₂ for 20 minutes at which time the volatiles were removed in vacuo and the residue was dissolved in DMSO and purified by RP-HPLC. The product fractions were lyophilized directly to yield ethyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate in 14% yield. LC/MS (m/z)=528.2 (MH⁺), R_t=0.65 min.

Synthesis of isopropyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate



Method 4 was followed using tert-butyl (1R,2S,3S,5R)-2-amino-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexyl carbamate and isopropyl carbonochloridate to give isopropyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methyl cyclohexylcarbamate in 6% yield. LC/MS (m/z)=542.3 (MH⁺), R_t=0.68 min.


Synthesis of N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-propionamidocyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide



Method 4 was followed using tert-butyl (1R,2S,3S,5R)-2-amino-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexyl carbamate, TEA and propionyl chloride to give N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-propionamidocyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 8% yield. LC/MS (m/z)=512.1 (MH⁺), R_t=0.62 min.


101

Synthesis of N-(4-(((1R,3R,4S,5S)-3-amino-4-isobutyramido-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide




Method 4 was followed using tert-butyl (1R,2S,3S,5R)-2-amino-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexyl carbamate, TEA and isobutyryl chloride to give N-(4-(((1R,3R,4S,5S)-3-amino-4-isobutyramido-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 16% yield. LC/MS (m/z)=526.3 (MH⁺), R_t=0.66 min.

Synthesis of N-(4-(((1R,3R,4S,5S)-3-amino-4-(2-methoxyacetamido)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide



Method 4 was followed using tert-butyl (1R,2S,3S,5R)-2-amino-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexyl carbamate, TEA and 2-methoxyacetyl chloride to give N-(4-(((1R,3R,4S,5S)-3-amino-4-(2-methoxyacetamido)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 24% yield. LC/MS (m/z)=528.2 (MH⁺), R_t=0.62 min.

Synthesis of N-(4-(((1R,3R,4R,5S)-4-acetamido-3-amino-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide




102


Method 4 was followed using tert-butyl ((1R,2R,3S,5R)-2-amino-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexyl)carbamate and acetic anhydride to give N-(4-(((1R,3R,4R,5S)-4-acetamido-3-amino-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 13% yield. LC/MS (m/z)=498.3 (MH⁺), R_t=0.58 min.

Method 5

Synthesis of N-(4-(((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonamido)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide and N-(4-(((1S,3S,4R,5R)-3-amino-5-methyl-4-(methylsulfonamido)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide



Single Enantiomer





Single Enantiomer

To a solution of (+/-)-tert-butyl (1R,2S,3S,5R)-2-amino-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexyl carbamate (1.0 equiv.) in CH₂Cl₂ (0.03 M) at 0° C. was added DIEA (3.0 equiv.) and then METHANESULFONYL CHLORIDE (1.5 equiv.). The homogeneous solution was left standing at 0° C. at for 2 hrs. Neutralize the reaction with sat. NaHCO₃ solution. The solution was partitioned between EtOAc and H₂O. The organic layer was washed with Na₂CO₃(sat.), NaCl(sat.), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography. Purification was completed via SFC (MeOH, 100 mL/min, OD column) to yield tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(methylsulfonamido)cyclohexylcarbamate (14% yield, 99% ee) and tert-butyl (1S,2R,3R,5S)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(methylsulfonamido)cyclohexylcarbamate (13% yield, 99% ee). LC/MS (m/z)=634.3 (MH⁺), R_t=0.86

103

min. Each Boc protected enantiomer was treated respectively with 25% TFA/CH₂Cl₂ for 20 minutes at which time the volatiles were removed in vacuo and the residue was dissolved in DMSO and purified by RP-HPLC to yield N-4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonylamido)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 94% yield; N-4-((1S,3S,4R,5R)-3-amino-5-methyl-4-(methylsulfonylamido)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 97% yield. LC/MS (m/z)=534.2 (MH⁺), R_t=0.59 min.



Synthesis of N-4-((1R,3R,4S,5S)-4-acetamido-3-amino-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide and N-4-((1S,3S,4R,5R)-4-acetamido-3-amino-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide



Method 5 was followed using (+/-)-tert-butyl (1R,2S,3S,5R)-2-amino-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexyl carbamate and acetic anhydride to give N-4-((1R,3R,4S,5S)-4-acetamido-3-amino-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 20% yield; N-4-((1S,3S,4R,5R)-4-acetamido-3-amino-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 25% yield. LC/MS (m/z)=498.2 (MH⁺), R_t=0.59 min.


104

Synthesis of methyl (1R,2S,4S,6R)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate and methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate



Method 5 was followed using (+/-)-tert-butyl (1R,2S,3S,5R)-2-amino-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexyl carbamate and methyl carbonochloridate to give methyl (1R,2S,4S,6R)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate in 9% yield; methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate in 10% yield. LC/MS (m/z)=514.2 (MH⁺), R_t=0.62 min.


Synthesis of (+/-)-(1S,2R,6S)-2-((tert-butoxycarbonyl)amino)-6-methyl-4-(3-nitropyridin-4-yl)cyclohex-3-en-1-yl methanesulfonate



105


To a solution of (+/-)-tert-butyl ((1R,5S,6S)-6-hydroxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-en-1-yl)carbamate (1.0 equiv.) in pyridine (0.20 M) was added MsCl (5.0 equiv.). The capped solution was stirred for 5 minutes and then the homogeneous solution was left standing at rt for 16 hrs. The volatiles were removed in vacuo and the residue was partitioned between EtOAc and H₂O. The organic layer was washed with 10% CuSO₄, H₂O, Na₂CO_{3(sat.)}, NaCl_(sat.), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography to yield (+/-)-(1S,2R,6S)-2-((tert-butoxycarbonyl)amino)-6-methyl-4-(3-nitropyridin-4-yl)cyclohex-3-en-1-yl methanesulfonate in 46% yield. LC/MS (m/z)=428.2 (MH⁺), R_t=0.89 min.

Synthesis of (+/-)-(1S,2R,4R,6S)-4-(3-aminopyridin-4-yl)-2-((tert-butoxycarbonyl)amino)-6-methylcyclohexyl methanesulfonate



Degass a solution of (+/-)-(1S,2R,6S)-2-((tert-butoxycarbonyl)amino)-6-methyl-4-(3-nitropyridin-4-yl)cyclohex-3-en-1-yl methanesulfonate (1.0 equiv.) in Ethanol (0.20 M). To this solution was added Pd/C (0.2 equiv.) and purge with Ar and H₂. The mixture was stirred under H₂ for 16 hrs. Filter the mixture over celite and wash the cake with MeOH. Concentrate the filtrate to yield (+/-)-(1S,2R,4R,6S)-4-(3-aminopyridin-4-yl)-2-((tert-butoxycarbonyl)amino)-6-methylcyclohexyl methanesulfonate in 49% yield. LC/MS (m/z)=400.3 (MH⁺), R_t=0.62 min.

Synthesis of (+/-)-tert-butyl ((1R,2R,3S,5R)-5-(3-aminopyridin-4-yl)-2-azido-3-methylcyclohexyl)carbamate




To a solution of (+/-)-(1S,2R,4R,6S)-4-(3-aminopyridin-4-yl)-2-((tert-butoxycarbonyl)amino)-6-methylcyclohexyl methanesulfonate (1.0 equiv.) in DMF (0.20 M) was added NaN₃ (7.0 equiv.). The solution was submerged in a 70° C. oil bath and left stirring under Ar for 4 hrs. The solution was cooled to rt and left stirring under Ar overnight. The solution was partitioned between EtOAc and H₂O. The organic layer

106


was washed with Na₂CO_{3(sat.)}, NaCl_(sat.), dried over MgSO₄, filtered, concentrated to yield (+/-)-tert-butyl ((1R,2R,3S,5R)-5-(3-aminopyridin-4-yl)-2-azido-3-methylcyclohexyl)carbamate in 87% yield. LC/MS (m/z)=347.3 (MH⁺), R_t=0.68 min.

Synthesis of (+/-)-tert-butyl ((1R,2R,3S,5R)-2-amino-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexyl)carbamate




EDC (2.0 equiv.) and HOAt (2.0 equiv.) was added to a solution of (+/-)-tert-butyl ((1R,2R,3S,5R)-5-(3-aminopyridin-4-yl)-2-azido-3-methylcyclohexyl)carbamate (1.0 equiv.) and 6-(2,6-difluorophenyl)-5-fluoropicolinic acid (1.5 equiv.) in DMF (0.20 M). The mixture was stirred at ambient temperature overnight. The reaction mixture was diluted with water and extracted with ethyl acetate. The combined extracts were washed sequentially with 1M aqueous sodium hydroxide and brine, dried over sodium sulfate, filtered, concentrated and purified by ISCO SiO₂ chromatography. To a degassed a solution of the azide (1.0 equiv.) in 2-propanol (0.10 M) was added Pd/C (0.2 equiv.). The mixture was stirred under H₂ for 48 hrs. Filter the mixture over celite and wash the cake with MeOH. Concentrate the filtrate to yield (+/-)-tert-butyl ((1R,2R,3S,5R)-2-amino-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexyl)carbamate in 35% yield. LC/MS (m/z)=556.3 (MH⁺), R_t=0.64 min.

Synthesis of N-(4-((1R,3R,4R,5S)-3-amino-5-methyl-4-(methylsulfonamido)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide and N-(4-((1S,3S,4S,5R)-3-amino-5-methyl-4-(methylsulfonamido)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide




Single Enantiomer

107



Method 5 was followed using (+/-)-tert-butyl (1R,2R,3S,5R)-2-amino-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexylcarbamate and methanesulfonyl chloride to give N-(4-((1R,3R,4R,5S)-3-amino-5-methyl-4-(methylsulfonamido)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 13% yield; N-(4-((1S,3S,4S,5R)-3-amino-5-methyl-4-(methylsulfonamido)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 14% yield. LC/MS (m/z)=534.2 (MH⁺), R_t=0.58 min.

Synthesis of tert-butyl (1R,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-oxocyclohexylcarbamate




To a solution of tert-butyl (1R,2R,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-2-hydroxy-3-methylcyclohexylcarbamate (1.0 equiv.) in DCM (0.10 M) was added Dess-MartinPeriodinane (1.2 equiv.). The flask was capped and the homogeneous solution was left stirring at rt for 3 hrs. The solution was partitioned between EtOAc and 1:1 10% Na₂S₂O₃/NaHCO₃ (sat.). The organic layer was washed with NaCl(sat.), dried over MgSO₄, filtered and concentrated, and purified by ISCO SiO₂ chromatography to yield tert-butyl (1R,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-oxocyclohexylcarbamate

108

nyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-oxocyclohexylcarbamate in 83% yield. LC/MS (m/z)=555.4 (MH⁺), R_t=0.87 min.

5 Synthesis of N-(4-((1R,3R,5S,E)-3-amino-4-(methoxyimino)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide and N-(4-((1R,3R,5S,Z)-3-amino-4-(methoxyimino)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide




A solution of methoxylamine-HCl (1.0 equiv.) and tert-butyl (1R,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-oxocyclohexylcarbamate (1.0 equiv.) in EtOH/pyridine (1/1, 0.01 M) was capped and left standing at rt for 16 hrs. The volatiles were removed in vacuo and the residue was partitioned between EtOAc and Na₂CO₃ (sat.). The organic layer was washed with NaCl(sat.), dried over MgSO₄, filtered, concentrated. The Boc group was removed with 25% TFA/CH₂Cl₂. After 45 minutes, the volatiles were removed in vacuo and the residue was pumped on for 5 minutes, dissolved in CH₂Cl₂ and neutralized with TEA. The volatiles were removed in vacuo and after pumping the residue was dissolved in DMSO and purified by RP-HPLC to yield N-(4-((1R,3R,5S,E)-3-amino-4-(methoxyimino)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 16% yield. LC/MS (m/z)=484.2 (MH⁺), R_t=0.64 min. ¹H NMR (400 MHz, DMSO-d₆) δ ppm 10.49 (s, 1H), 8.57 (s, 1H), 8.47 (d, J=4.0, 1H), 8.35 (dd, J=8.0, 4.0, 1H), 8.25 (broad doublet, J=4.0, 2H), 8.20 (t, J=8.0, 1H), 7.67-7.74 (m, 1H), 7.42 (d, J=8.0, 1H), 7.36 (t, J=8.0, 2H), 4.04-4.08 (m, 1H), 3.79 (s, 3H), 3.23-3.29 (m, 1H), 2.39-2.45 (m, 1H), 2.11 (d, J=8.0, 1H), 2.10 (d, J=8.0, 1H), 1.90 (q, J=12, 1H), 1.40 (q, J=12, 1H), 1.01 (d, J=4.0, 3H); and N-(4-((1R,3R,5S,Z)-3-amino-4-(methoxyimino)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 27% yield. LC/MS (m/z)=484.2 (MH⁺), R_t=0.66 min. ¹H NMR (400 MHz, DMSO-d₆) δ ppm 10.48 (s, 1H), 8.57 (s, 1H), 8.47 (d, J=4.0, 1H), 8.35 (dd, J=8.0, 4.0, 1H), 8.20 (t, J=8.0, 1H), 8.08 (broad singlet, 2H), 7.67-7.74 (m, 1H), 7.42 (d, J=8.0, 1H), 7.36 (t,

109


J=8.0, 2H), 3.88-3.92 (m, 1H), 3.80 (s, 3H), 3.22-3.28 (m, 1H), 2.51-2.58 (m, 1H), 2.25 (d, J=12.0, 1H), 1.86 (d, J=12.0, 1H), 1.70 (q, J=12, 1H), 1.62 (q, J=12, 1H), 1.34 (d, J=4.0, 3H).

Synthesis of N-(4-((1R,3R,5S,Z)-3-amino-4-(hydroxyimino)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide



A solution of hydroxylamine-HCl (4.0 equiv.) and tert-butyl (1R,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-oxocyclohexylcarbamate (1.0 equiv.) in EtOH/pyridine (1/1, 0.01 M) was capped and left standing at rt for 16 hrs. The volatiles were removed in vacuo and the residue was partitioned between EtOAc and Na₂CO₃(*sat.*). The organic layer was washed with NaCl(*sat.*), dried over MgSO₄, filtered, concentrated. The Boc group was removed with 25% TFA/CH₂Cl₂. After 45 minutes, the volatiles were removed in vacuo and after pumping the residue was dissolved in DMSO and purified by RP-HPLC to yield N-(4-((1R,3R,5S,Z)-3-amino-4-(hydroxyimino)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 14% yield. LC/MS (m/z)=470.3 (MH⁺), R_f=0.60 min. ¹H NMR (300 MHz, DMSO-d₆) δ ppm 10.94 (s, 1H), 10.49 (s, 1H), 8.60 (s, 1H), 8.48 (d, J=4.0, 1H), 8.35 (dd, J=8.0, 4.0, 1H), 8.20 (t, J=8.0, 1H), 8.02 (broad doublet, J=4.0, 2H), 7.67-7.74 (m, 1H), 7.42 (d, J=4.0, 1H), 7.36 (t, J=8.0, 2H), 4.24 (m, 1H), 3.82-3.86 (m, 1H), 3.21-3.27 (m, 1H), 2.50-2.55 (m, 1H), 2.24 (d, J=12.0, 1H), 1.86 (d, J=16.0, 1H), 1.68 (q, J=12.0, 1H), 1.59 (q, J=12.0, 1H), 1.40 (d, J=8.0, 3H).

Synthesis of N-(4-((1R,3R,5S)-3-amino-5-methyl-4-oxocyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide




Tert-butyl (1R,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-oxocyclohexylcarbamate was treated with 25% TFA/CH₂Cl₂ for 30 minutes. The volatiles were removed in vacuo, the residue was dissolved in DMSO and purified by reverse phase HPLC

110

to yield N-(4-((1R,3R,5S)-3-amino-5-methyl-4-oxocyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 98% yield. LC/MS (m/z)=455.1 (MH⁺), R_f=0.57 min. ¹H NMR (300 MHz, DMSO-d₆) δ ppm 10.55 (s, 1H), 8.55 (s, 1H), 8.47 (d, J=4.0, 1H), 8.37 (dd, J=8.0, 4.0, 1H), 8.21 (t, J=8.0, 1H), 8.16 (broad doublet, J=4.0, 2H), 7.67-7.74 (m, 1H), 7.40 (d, J=8.0, 1H), 7.36 (t, J=8.0, 2H), 4.20-4.26 (m, 1H), 3.50-3.70 (m, 2H), 2.76-2.82 (m, 1H), 2.49-2.54 (m, 1H), 2.32-2.36 (m, 1H), 2.16-2.18 (m, 1H), 1.91 (q, J=12, 1H), 1.65 (q, J=12, 1H), 0.97 (d, J=8.0, 3H).


Synthesis of (+/-)-tert-butyl ((1R,5S,6R)-6-methoxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-en-1-yl)carbamate



(+/-)-Tert-butyl (1R,5S,6R)-6-hydroxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate (1.0 equiv.) was suspended in iodomethane (100.0 equiv.). Silver oxide (6.0 equiv.) was added to the mixture and the reaction vessel was wrapped in foil (kept dark) and allowed to stir 45° C. for 10 hrs. The reaction was diluted with THF and filtered through a pad of celite. The celite cake was further washed with MeOH. The organics were concentrated and the crude was taken up in DCM, washed with NaHCO₃(*aq.*), dried over Na₂SO₄, filtered and concentrated. The crude was loaded onto silica gel and purified via ISCO to yield (+/-)-tert-butyl ((1R,5S,6R)-6-methoxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-en-1-yl)carbamate in 35% yield. LC/MS (m/z)=364.1 (MH⁺), R_f=0.89 min.

Method 6


Synthesis of tert-butyl ((1S,2S,3R,5S)-5-(3-aminopyridin-4-yl)-2-methoxy-3-methylcyclohexyl)carbamate and tert-butyl ((1R,2R,3S,5R)-5-(3-aminopyridin-4-yl)-2-methoxy-3-methylcyclohexyl)carbamate



111


To a solution of (+/-)-tert-butyl ((1R,5S,6R)-6-methoxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-en-1-yl)carbamate (1.0 equiv.) in degassed EtOH (0.10 M) was added Pd/C (0.1 equiv.). The mixture was purged with H₂, and allowed to stir under an atm of H₂ overnight at RT. The reaction was filtered through a pad of celite and the cake was washed with MeOH. The organics were concentrated and purified by ISCO SiO₂ chromatography. Purification was completed via SFC (30% MeOH, 100 mL/min, AD column) to yield tert-butyl ((1S,2S,3R,5S)-5-(3-aminopyridin-4-yl)-2-methoxy-3-methylcyclohexyl)carbamate (15% yield, 99% ee) and tert-butyl ((1R,2R,3S,5R)-5-(3-aminopyridin-4-yl)-2-methoxy-3-methylcyclohexyl)carbamate (12% yield, 99% ee). LC/MS (m/z)=336.3 (MH⁺), R_f=0.58 min.

Synthesis of (+/-)-tert-butyl ((1R,5S,6S)-6-hydroxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-en-1-yl)carbamate



To a solution of (+/-)-(3aR,7S,7aS)-tert-butyl 7-methyl-5-(3-nitropyridin-4-yl)-2-oxo-3a,6,7,7a-tetrahydrobenzo[d]oxazole-3(2H)-carboxylate (1.0 equiv.) in THF (0.20 M) was added 2M LiOH (3.0 equiv.) was added. The mixture was stirred overnight 20 hrs at 22° C. The mixture was diluted with EtOAc and NaHCO_{3(aq.)}. The layers were separated and the aqueous was extracted with EtOAc. The combined organics were washed with brine, dried over Na₂SO₄, filtered, and concentrated. The golden foam was purified by ISCO chromatography to afford (+/-)-tert-butyl ((1R,5S,6S)-6-hydroxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-en-1-yl)carbamate in 83% yield. LC/MS (m/z)=350.2 (MH⁺), R_f=0.82 min.


Synthesis of (+/-)-tert-butyl (1R,5S,6S)-6-(2-cyanoethoxy)-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate



112


A mixture of (+/-)-Tert-butyl (1R,5S,6S)-6-hydroxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate (1.0 equiv.), acrylonitrile (30.0 equiv.) and CESIUM CARBONATE (1.2 equiv.) in t-BuOH (0.57 M) was stirred at 35° C. for 3 hrs. The reaction was cooled to room temperature, followed by the addition of NaHCO_{3(aq.)} and water. The mixture was extracted with EtOAc and the combined organics were dried over MgSO₄, filtered, and concentrated. The sample was purified by ISCO SiO₂ chromatography to yield (+/-)-tert-butyl (1R,5S,6S)-6-(2-cyanoethoxy)-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate in 94% yield. LC/MS (m/z)=403.3 (MH⁺), R_f=0.92 min. ¹H NMR (400 MHz, CHLOROFORM-d) δ ppm 1.13 (d, J=6.46 Hz, 3H), 1.46 (s, 9H), 1.99-2.18 (m, 2H), 2.20-2.36 (m, 1H), 2.65 (t, J=6.06 Hz, 2H), 3.68 (br. s., 1H), 3.88 (t, J=5.99 Hz, 2H), 4.51 (br. s., 1H), 4.99 (d, J=9.15 Hz, 1H), 5.39 (br. s., 1H), 7.25 (d, J=4.99 Hz, 1H), 8.73 (d, J=4.94 Hz, 1H), 9.10 (s, 1H).

Synthesis of tert-butyl ((1S,2R,3R,5S)-5-(3-aminopyridin-4-yl)-2-(2-cyanoethoxy)-3-methylcyclohexyl)carbamate and tert-butyl ((1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-2-(2-cyanoethoxy)-3-methylcyclohexyl)carbamate



Method 6 was followed using (+/-)-tert-butyl (1R,5S,6S)-6-(2-cyanoethoxy)-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate with SFC (15% EtOH, 100 mL/min, OJ column) to yield tert-butyl ((1S,2R,3R,5S)-5-(3-aminopyridin-4-yl)-2-(2-cyanoethoxy)-3-methylcyclohexyl)carbamate (31% yield, 99% ee) and tert-butyl ((1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-2-(2-cyanoethoxy)-3-methylcyclohexyl)carbamate (26% yield, 99% ee). LC/MS (m/z)=375.3 (MH⁺), R_f=0.65 min.


Synthesis of (+/-)-tert-butyl ((1R,5S,6S)-6-methoxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-en-1-yl)carbamate



113

To a solution of (+/-)-tert-butyl ((1R,5S,6S)-6-hydroxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-en-1-yl)carbamate (1.0 equiv.) in MeI (100.0 equiv.) was added Ag₂O (5.5 equiv.). A reflux condenser was attached and the heterogeneous solution under Ar was submerged in a 50° C. bath and the reaction was gently refluxing for 6 hrs. The solids were filtered, rinsed with CH₂Cl₂. The volatiles were removed in vacuo, the residue was partitioned between CH₂Cl₂ and NaHCO₃(sat.). The organic layer was dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography to yield (+/-)-tert-butyl ((1R,5S,6S)-6-hydroxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-en-1-yl)carbamate in 59% yield. LC/MS (m/z)=364.5 (MH⁺), R_f=1.02 min. ¹H NMR (400 MHz, <cdcl3>) δ ppm 1.11 (d, J=6.65 Hz, 3H), 1.46 (s, 9H), 1.95-2.13 (m, 2H), 2.18-2.28 (m, 1H), 3.47 (d, J=3.52 Hz, 1H), 3.57 (s, 3H), 4.45 (d, J=7.83 Hz, 1H), 5.01 (d, J=9.39 Hz, 1H), 5.44 (br. s., 1H), 7.24 (s, 1H), 8.71 (d, J=5.09 Hz, 1H), 9.08 (s, 1H).


Synthesis of tert-butyl ((1S,2R,3R,5S)-5-(3-aminopyridin-4-yl)-2-methoxy-3-methylcyclohexyl)carbamate and tert-butyl ((1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-2-methoxy-3-methylcyclohexyl)carbamate



To a degassed solution of (+/-)-tert-butyl ((1R,5S,6S)-6-hydroxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-en-1-yl)carbamate (1.0 equiv.) in i-PrOH (0.07 M) was added Pd/C (0.1 equiv.). The solution was degassed and purged to H₂ and left stirring under a balloon of H₂ at rt for 16 hrs. The solution was degassed and purged to Ar, diluted with CH₂Cl₂, filtered through a pad of celite, concentrated and purified by ISCO SiO₂ chromatography. Purification was completed via SFC (20% MeOH, 100 mL/min, AD column) to yield tert-butyl ((1S,2R,3R,5S)-5-(3-aminopyridin-4-yl)-2-methoxy-3-methylcyclohexyl)carbamate (42% yield, 99% ee) and tert-butyl ((1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-2-methoxy-3-methylcyclohexyl)carbamate (39% yield, 99% ee). LC/MS (m/z)=336.2 (MH⁺), R_f=0.67 min. ¹H NMR (400 MHz, <cdcl3>) δ ppm 1.08 (d, J=7.04 Hz, 3H), 1.43-1.49 (m, 10H), 1.52-1.64 (m, 2H), 1.70-1.81 (m, 2H), 2.52-2.64 (m, 1H), 3.39 (br. s., 1H), 3.52-3.57 (m, 3H), 3.62 (br. s., 2H), 3.66-3.75 (m, 1H), 4.75-4.87 (m, 1H), 6.98 (d, J=5.09 Hz, 1H), 7.95-8.05 (m, 2H).


114

Synthesis of (+/-)-tert-butyl (1R,5S,6S)-6-ethoxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate



(+/-)-Tert-butyl (1R,5S,6S)-6-hydroxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate (1.0 equiv.) was suspended in iodoethane (100.0 equiv.). Silver oxide (6.0 equiv.) was added to the mixture and the reaction vessel was wrapped in foil (kept dark) and allowed to stir 55° C. for 10 hrs. The reaction was diluted with THF and filtered through a pad of celite. The celite cake was further washed with MeOH. The organics were concentrated and the crude was taken up in DCM, washed with NaHCO₃ (aq), dried over Na₂SO₄, filtered and concentrated. The crude was loaded onto silica gel and purified via ISCO to yield (+/-)-tert-butyl (1R,5S,6S)-6-ethoxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate in 31% yield. LC/MS (m/z)=378.1 (MH⁺), R_f=0.99 min.

Synthesis of tert-butyl ((1S,2R,3R,5S)-5-(3-aminopyridin-4-yl)-2-ethoxy-3-methylcyclohexyl)carbamate and tert-butyl ((1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-2-ethoxy-3-methylcyclohexyl)carbamate




Method 6 was followed using (+/-)-tert-butyl (1R,5S,6S)-6-ethoxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate with Chiral HPLC (Heptane/EtOH=90/10, 20 mL/min, IC column) to yield tert-butyl ((1S,2R,3R,5S)-5-(3-aminopyridin-4-yl)-2-ethoxy-3-methylcyclohexyl)carbamate (33% yield, 99% ee) and tert-butyl ((1R,2S,3S,5R)-5-(3-

115


aminopyridin-4-yl)-2-ethoxy-3-methylcyclohexyl)
 carbamate (28% yield, 99% ee). LC/MS (m/z)=350.2 (MH^+),
 R_f =0.72 min.

Synthesis of (+/-)-methyl 3-((1R,2R,6S)-2-(tert-
 butoxycarbonylamino)-6-methyl-4-(3-nitropyridin-4-
 yl)cyclohex-3-enyloxy)propanoate



A mixture of (+/-)-tert-butyl (1R,5S,6R)-6-hydroxy-5-
 methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate
 (1.0 equiv.), methyl acrylate (30.0 equiv.) and CESIUM
 CARBONATE (1.2 equiv.) in t-BuOH (0.38 M) was stirred at
 35° C. for 16 hrs. The reaction was cooled to room tempera-
 ture, followed by the addition of $NaHCO_3(aq.)$ and water. The
 mixture was extracted with EtOAc and the combined organics
 were dried over $MgSO_4$, filtered, and concentrated. The
 sample was purified by ISCO chromatography to yield (+/-)-
 methyl 3-((1R,2R,6S)-2-(tert-butoxycarbonylamino)-6-me-
 thyl-4-(3-nitropyridin-4-yl)cyclohex-3-enyloxy)propanoate
 in 48% yield. LC/MS (m/z)=436.1 (MH^+), R_f =0.91 min.

Synthesis of (+/-)-methyl 3-((1R,2R,4R,6S)-4-(3-
 aminopyridin-4-yl)-2-(tert-butoxycarbonylamino)-6-
 methylcyclohexyloxy)propanoate




To a solution of (+/-)-methyl 3-((1R,2R,6S)-2-(tert-bu-
 toxycarbonylamino)-6-methyl-4-(3-nitropyridin-4-yl)cyclo-
 hex-3-enyloxy)propanoate (1.0 equiv.) in degassed EtOH

116

(0.07 M) was added Pd/C (0.3 equiv.). The mixture was
 purged with H_2 , and allowed to stir under H_2 overnight at RT.
 The reaction was filtered through a pad of celite and the cake
 was washed with MeOH. The organics were concentrated and
 purified by ISCO SiO_2 chromatography to yield (+/-)-methyl
 3-((1R,2R,4R,6S)-4-(3-aminopyridin-4-yl)-2-(tert-butoxy-
 carbonylamino)-6-methylcyclohexyloxy)propanoate in
 100% yield. LC/MS (m/z)=408.2 (MH^+), R_f =0.62 min.


Synthesis of methyl 3-((1R,2R,4R,6S)-2-(tert-bu-
 toxycarbonylamino)-4-(3-(6-(2,6-difluorophenyl)-5-
 fluoropicolinamido)pyridin-4-yl)-6-methylcyclo-
 hexyloxy)propanoate and methyl 3-(((1S,2S,4S,6R)-
 2-((tert-butoxycarbonyl)amino)-4-(3-(6-(2,6-
 difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-
 6-methylcyclohexyl)oxy)propanoate



EDC (2.0 equiv.) and HOAt (2.0 equiv.) was added to a
 solution of (+/-)-methyl 3-((1R,2R,4R,6S)-4-(3-aminopyri-
 din-4-yl)-2-(tert-butoxycarbonylamino)-6-methylcyclo-
 hexyloxy)propanoate (1.0 equiv.) and 6-(2,6-difluorophe-
 nyl)-5-fluoropicolinic acid (1.5 equiv.) in DMF (0.08 M). The
 mixture was stirred at ambient temperature overnight. The
 reaction mixture was diluted with water and extracted with
 ethyl acetate. The combined extracts were washed sequen-
 tially with 1M aqueous sodium hydroxide and brine, dried
 over sodium sulfate, filtered, concentrated and purified by
 ISCO SiO_2 chromatography. Purification was completed via
 SFC (20% IPA, 20 mL/min, AD column) to yield methyl
 3-((1R,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(6-(2,
 6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-
 methylcyclohexyloxy)propanoate (22% yield, 99% ee) and
 methyl 3-((1S,2S,4S,6R)-2-(tert-butoxycarbonyl)amino)-4-
 (3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-
 yl)-6-methylcyclohexyl)oxy)propanoate (21% yield, 99%
 ee). LC/MS (m/z)=643.4 (MH^+), R_f =0.92 min.


117

Synthesis of 3-((1R,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyloxy)propanoic acid



To a solution of methyl 3-((1R,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyloxy)propanoate (1.0 equiv.) was added 4 M HCl (40.0 equiv.) in dioxane. After stirring at rt overnight, the mixture was concentrated and dissolved in MeOH (0.05 M). LiOH (20.0 equiv.) was added. After stirred 10 min at rt, the mixture was concentrated, neutralized with HCl to pH 7 and extracted with EtOAc/t-Butanol (1/1). The organic layer was wash with brine, dried over MgSO₄, filtered, and concentrated. The sample was purified by RP HPLC to yield 3-((1R,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyloxy)propanoic acid in 34% yield. LC/MS (m/z)=529.3 (MH⁺), R_t=0.63 min.

Synthesis of (+/-)-tert-butyl (1R,5S,6S)-5-methyl-6-(2-(methylsulfonyl)ethoxy)-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate




A mixture of (+/-)-Tert-butyl (1R,5S,6S)-6-hydroxy-5-methyl-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate (1.0 equiv.), methylsulfonyl ethene (30.0 equiv.) and CESIUM CARBONATE (1.2 equiv.) in t-BuOH (0.22 M) was stirred at 22° C. for 5 hrs. The reaction was cooled to room temperature, followed by the addition of NaHCO₃(aq.) and water. The mixture was extracted with EtOAc and the combined organics were dried over MgSO₄, filtered, and concentrated. The sample was purified by ISCO chromatography to yield (+/-)-tert-butyl (1R,5S,6S)-5-methyl-6-(2-(methylsulfonyl)ethoxy)-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate in 84% yield. LC/MS (m/z)=456.2 (MH⁺), R_t=0.87 min. ¹H NMR (400 MHz, CHLOROFORM-d) δ ppm 1.10 (d, J=6.60 Hz, 3H), 1.47 (s, 9H), 1.92-2.16 (m, 2H),

118


2.17-2.32 (m, 1H), 2.93-3.00 (m, 1H), 3.09 (s, 3H), 3.18 (d, J=14.87 Hz, 1H), 3.38-3.52 (m, 1H), 3.63 (d, J=2.40 Hz, 1H), 3.95-4.06 (m, 1H), 4.21 (td, J=9.84, 2.42 Hz, 1H), 4.56 (d, J=7.58 Hz, 1H), 5.58 (br. s., 1H), 5.66 (d, J=9.44 Hz, 1H), 5.72 (d, J=4.99 Hz, 1H), 8.73 (d, J=4.99 Hz, 1H), 9.05 (s, 1H).

Synthesis of (+/-)-tert-butyl (1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-3-methyl-2-(2-(methylsulfonyl)ethoxy)cyclohexylcarbamate




To a solution of (+/-)-tert-butyl (1R,5S,6S)-5-methyl-6-(2-(methylsulfonyl)ethoxy)-3-(3-nitropyridin-4-yl)cyclohex-2-enylcarbamate (1.0 equiv.) in degassed EtOH (0.17 M) was added Pd/C (0.3 equiv.). The mixture was purged with H₂, and allowed to stir under H₂ overnight at RT. The reaction was filtered through a pad of celite and the cake was washed with MeOH. The organics were concentrated and purified by ISCO SiO₂ chromatography to yield (+/-)-tert-butyl (1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-3-methyl-2-(2-(methylsulfonyl)ethoxy)cyclohexylcarbamate in 55% yield. LC/MS (m/z)=428.2 (MH⁺), R_t=0.59 min. ¹H NMR (400 MHz, CHLOROFORM-d) δ ppm 1.07 (d, J=6.85 Hz, 3H), 1.46 (s, 9H), 1.72-1.88 (m, 2H), 2.62 (tt, J=12.23, 3.30 Hz, 1H), 3.08 (s, 3H), 3.21 (d, J=14.62 Hz, 1H), 3.35-3.47 (m, 1H), 3.58 (br. s., 1H), 3.64 (br. s., 2H), 3.70-3.86 (m, 1H), 3.94-4.10 (m, 1H), 4.10-4.22 (m, 1H), 5.43 (d, J=9.00 Hz, 1H), 6.89 (d, J=5.04 Hz, 1H), 7.98 (d, J=4.99 Hz, 1H), 8.03 (s, 1H).

Synthesis of tert-butyl (1S,2R,3R,5S)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(2-(methylsulfonyl)ethoxy)cyclohexylcarbamate and tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(2-(methylsulfonyl)ethoxy)cyclohexylcarbamate




119



EDC (2.0 equiv.) and HOAt (2.0 equiv.) were added to a solution of (+/-)-tert-butyl (1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-3-methyl-2-(2-(methylsulfonyl)ethoxy)cyclohexylcarbamate (1.0 equiv.) and 6-(2,6-difluorophenyl)-5-fluoropicolinic acid (1.5 equiv.) in DMF (0.08 M). The mixture was stirred at ambient temperature overnight. The reaction mixture was diluted with water and extracted with ethyl acetate. The combined extracts were washed sequentially with 1M aqueous sodium hydroxide and brine, dried over sodium sulfate, filtered, concentrated and purified by ISCO SiO₂ chromatography. Purification was completed via SFC (50% MeOH, 100 mL/min, IC column) to yield tert-butyl (1S,2R,3R,5S)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(2-(methylsulfonyl)ethoxy)cyclohexylcarbamate (48% yield, 99% ee) and tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(2-(methylsulfonyl)ethoxy)cyclohexylcarbamate (48% yield, 99% ee). LC/MS (m/z)=663.2 (MH⁺), R_f=0.88 min. ¹H NMR (400 MHz, ³CDCl₃) δ ppm 0.95 (d, J=6.75 Hz, 3H), 1.45 (s, 8H), 1.50-1.84 (m, 5H), 2.80-2.95 (m, 1H), 3.08 (s, 3H), 3.14-3.29 (m, 1H), 3.32-3.45 (m, 1H), 3.56 (br. s., 1H), 3.63-3.77 (m, 1H), 3.95-4.08 (m, 1H), 4.12 (q, J=7.12 Hz, 1H), 5.36 (d, J=8.56 Hz, 1H), 7.04-7.18 (m, 3H), 7.50 (tt, J=8.47, 6.35 Hz, 1H), 7.77 (t, J=8.56 Hz, 1H), 8.33-8.46 (m, 2H), 9.26 (s, 1H), 9.81 (s, 1H).

Synthesis of (+/-)-tert-butyl (1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-2-cyano-3-methylcyclohexylcarbamate




To a solution of (+/-)-(1R,2R,4R,6S)-2-(tert-butoxycarbonylamino)-4-(3-(tert-butoxycarbonylamino)pyridin-4-yl)-6-methylcyclohexyl methanesulfonate (1.0 equiv.) in DMF

120

(0.20 M) was added NaCN (5.0 equiv.). The solution was submerged in an 85° C. oil bath and left stirring under Ar for 16 hrs. The solution was cooled to rt and left stirring under Ar overnight. The solution was partitioned between EtOAc and H₂O. The organic layer was washed with Na₂CO₃(sat.), NaCl(sat.), dried over MgSO₄, filtered, concentrated. To a solution of the bis-Boc product (10 equiv.) in DCM (0.20 M) was added TFA (62.0 equiv.). The mixture was stirred at ambient temperature for 40 min and concentrated and neutralized with saturated aqueous sodium bicarbonate. Dioxane (0.15 M) and Boc₂O (4.0 equiv.) were added. The reaction mixture was stirred vigorously at ambient temperature for 16 hrs. Volatiles were removed in vacuo. The aqueous phase was extracted with 10:1 DCM:MeOH. The combined extracts were dried over sodium sulfate, filtered, and concentrated. The crude material was purified by ISCO chromatography over silica gel to give (+/-)-tert-butyl (1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-2-cyano-3-methylcyclohexylcarbamate in 20% yield. LC/MS (m/z)=331.2 (MH⁺), R_f=0.62 min.

Synthesis of tert-butyl (1R,2S,3S,5R)-2-cyano-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexylcarbamate and tert-butyl (1S,2R,3R,5S)-2-cyano-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexylcarbamate




EDC (1.1 equiv.) and HOAt (1.1 equiv.) were added to a solution of (+/-)-tert-butyl (1R,2S,3S,5R)-5-(3-aminopyridin-4-yl)-2-cyano-3-methylcyclohexylcarbamate (1.0 equiv.) and 6-(2,6-difluorophenyl)-5-fluoropicolinic acid (1.5 equiv.) in DMF (0.11 M). The mixture was stirred at ambient temperature overnight. The reaction mixture was diluted with water and extracted with ethyl acetate. The combined extracts were washed sequentially with 1M aqueous sodium hydroxide and brine, dried over sodium sulfate, filtered, concentrated and purified by ISCO SiO₂ chromatography. Purification was completed via SFC (15% IPA, 100 mL/min, IA column) to yield tert-butyl (1R,2S,3S,5R)-2-

121


cyano-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexylcarbamate (25% yield, 99% ee) and tert-butyl (1S,2R,3R,5S)-2-cyano-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexylcarbamate (27% yield, 99% ee). LC/MS (m/z)=566.2 (MH⁺), R_t=0.90 min.

Synthesis of tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(1H-1,2,3-triazol-1-yl)cyclohexylcarbamate



A solution of tert-butyl (1R,2S,3S,5R)-2-azido-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methylcyclohexylcarbamate (1.0 equiv.) in vinyl acetate (0.06 M) to give a suspension was heated in microwave at 160° C. for 1 hr. The reaction was concentrated to yield tert-butyl (1R,2S,3S,5R)-5-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-3-methyl-2-(1H-1,2,3-triazol-1-yl)cyclohexylcarbamate in 50% yield. LC/MS (m/z)=608.3 (MH⁺), R_t=0.89 min.


Synthesis of (+/-)-tert-butyl ((1R,2S,3S,5R)-5-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-2-amino-3-methylcyclohexyl)carbamate



To a degassed a solution of (+/-)-tert-butyl ((1R,2S,3S,5R)-5-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-2-azido-3-methylcyclohexyl)carbamate (1.0 equiv.) in ethanol (0.10 M) was added Pd/C (0.2 equiv.). The mixture was stirred under H₂ for 4 hrs. Filter the mixture over celite and wash the cake with MeOH. Concentrate the filtrate to yield (+/-)-tert-butyl ((1R,2S,3S,5R)-5-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-2-amino-3-methylcyclohexyl)carbamate in 88% yield. LC/MS (m/z)=421.3 (MH⁺), R_t=0.58 min.


122

Synthesis of (+/-)-tert-butyl ((1R,2S,3S,5R)-5-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-3-methyl-2-(methylamino)cyclohexyl)carbamate



To a solution of (+/-)-tert-butyl ((1R,2S,3S,5R)-5-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-2-amino-3-methylcyclohexyl)carbamate (1.0 equiv.) in MeOH (0.10 M) was added benzaldehyde (1.3 equiv.). After 3 hrs, sodium cyanotrihydroborate (2.5 equiv.) was added and the mixture was stirred at rt for 16 hrs. The reaction mixture was quenched by the addition of water, and volatiles were removed in vacuo. The mixture was extracted with ethyl acetate. The combined organic phases were dried with sodium sulfate, filtered and concentrated. The residue was dissolved in MeOH (0.10 M) and paraformaldehyde (5.0 equiv.) was added. After 16 hrs, sodium cyanotrihydroborate (5.0 equiv.) was added and the mixture was left stirred at rt for 16 hrs. The reaction was quenched by the addition of water, and volatiles were removed in vacuo. The mixture was extracted with DCM. The combined organic phases were dried with sodium sulfate, filtered, concentrated and purified by ISCO Chromatography. The product was dissolved in MeOH (0.10 M) and treated with Pd(OH)₂ (0.50 equiv.) under H₂ for 5 hrs at RT. The reaction was filtered through a pad of celite and the cake was washed with MeOH. The organics were concentrated to yield (+/-)-tert-butyl ((1R,2S,3S,5R)-5-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-3-methyl-2-(methylamino)cyclohexyl)carbamate in 75% yield. LC/MS (m/z)=435.2 (MH⁺), R_t=0.64 min.


Synthesis of (+/-)-methyl ((1S,2R,4R,6S)-4-(3-aminopyridin-4-yl)-2-((tert-butoxycarbonyl)amino)-6-methylcyclohexyl)(methyl)carbamate



123

To a solution of (+/-)-tert-butyl ((1R,2S,3S,5R)-5-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-3-methyl-2-(methylamino)cyclohexyl)carbamate (1.0 equiv.) in DCM (0.05 M) at 0° C. was added DIEA (3.0 equiv.) and then methyl chloroformate (1.5 equiv.). The homogeneous solution was left standing at 0° C. at for 4 hrs. The reaction was quenched partitioned between NaHCO₃ solution and EtOAc. The organic layer was washed with Brine, dried over Na₂SO₄, concentrated and purified by ISCO chromatography. The product was treated with 4 M HCl in dioxane (30.0 equiv.) at rt for 1 hour. The volatiles were removed in vacuo and the solid was pumped on for 5 minutes on the high vac. To the residue was added CH₂Cl₂ (0.15 M), DIEA (5.0 equiv.) and tert-butyl 2,5-dioxopyrrolidin-1-yl carbonate (1.6 equiv.). The solution was left stirring at rt for 1 hr. The volatiles were removed in vacuo and the residue was partitioned between EtOAc and H₂O. The organic layer was washed with Na₂CO_{3(sat.)}, NaCl_(sat.), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography to yield (+/-)-methyl ((1S,2R,4R,6S)-4-(3-aminopyridin-4-yl)-2-((tert-butoxycarbonyl)amino)-6-methylcyclohexyl)(methyl)carbamate in 20% yield. LC/MS (m/z)=393.2 (MH⁺), R_t=0.60 min.

Synthesis of methyl ((1S,2R,4R,6S)-2-((tert-butoxycarbonyl)amino)-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl)(methyl)carbamate and methyl ((1R,2S,4S,6R)-2-((tert-butoxycarbonyl)amino)-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl)(methyl)carbamate




EDC (1.1 equiv.) and HOAt (1.1 equiv.) were added to a solution of (+/-)-methyl ((1S,2R,4R,6S)-4-(3-aminopyridin-4-yl)-2-((tert-butoxycarbonyl)amino)-6-methylcyclohexyl)(methyl)carbamate (1.0 equiv.) and 6-(2,6-difluorophenyl)-5-fluoropicolinic acid (1.5 equiv.) in DMF (0.05 M). The mixture was stirred at ambient temperature overnight. The

124


reaction mixture was diluted with water and extracted with ethyl acetate. The combined extracts were washed sequentially with 1M aqueous sodium hydroxide and brine, dried over sodium sulfate, filtered, concentrated and purified by ISCO SiO₂ chromatography. Purification was completed via SFC (40% EtOH, 100 mL/min, IC column) to yield methyl ((1S,2R,4R,6S)-2-((tert-butoxycarbonyl)amino)-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl)(methyl)carbamate (15% yield, 99% ee) and methyl ((1R,2S,4S,6R)-2-((tert-butoxycarbonyl)amino)-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl)(methyl)carbamate (15% yield, 99% ee). LC/MS (m/z)=628.3 (MH⁺), R_t=0.89 min.

Synthesis of tert-butyl (3R,4S,5S)-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methyl-4-(1H-1,2,3-triazol-1-yl)piperidin-3-ylcarbamate



A solution of tert-butyl (3R,4S,5S)-4-azido-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-3-ylcarbamate (1.0 equiv.) in vinyl acetate (0.06 M) to give a suspension was heated at 110° C. for 88 hrs. The reaction was concentrated and purified by ISCO chromatography to yield tert-butyl (3R,4S,5S)-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methyl-4-(1H-1,2,3-triazol-1-yl)piperidin-3-ylcarbamate in 48% yield. LC/MS (m/z)=609.3 (MH⁺), R_t=0.83 min.


Synthesis of tert-butyl (3R,4S,5S)-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methyl-4-(4-(thiophen-3-yl)-1H-1,2,3-triazol-1-yl)piperidin-3-ylcarbamate



125


In a high pressure vial was added tert-butyl (3R,4S,5S)-4-azido-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-3-ylcarbamate (1.0 equiv.), copper in charcoal (0.4 equiv.), 3-ethynyl thiophene (5.0 equiv.) and triethylamine (1.0 equiv.), in dioxane (0.09 M) to give a black suspension. The pressure tube was sealed and the mixture was stirred with heating to 100° C. overnight. The reaction was cooled to RT, filtered through celite, concentrated to yield tert-butyl (3R,4S,5S)-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methyl-4-(4-(thiophen-3-yl)-1H-1,2,3-triazol-1-yl)piperidin-3-ylcarbamate in 48% yield. LC/MS (m/z)=691.2 (MH⁺), R_t=0.98 min.

Synthesis of tert-butyl (3R,4S,5S)-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methyl-4-(4-methyl-1H-1,2,3-triazol-1-yl)piperidin-3-ylcarbamate



In a high pressure vial was added tert-butyl (3R,4S,5S)-4-azido-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-3-ylcarbamate (1.0 equiv.), copper in charcoal (0.2 equiv.), prop-1-yne (10.0 equiv.) and triethylamine (1.5 equiv.), in dioxane (0.15 M) to give a black suspension. The pressure tube was sealed and the mixture was stirred with heating to 60° C. for 48 hrs. The reaction was cooled to RT, filtered through celite, concentrated to yield tert-butyl (3R,4S,5S)-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methyl-4-(4-methyl-1H-1,2,3-triazol-1-yl)piperidin-3-ylcarbamate in 95% yield. LC/MS (m/z)=623.2 (MH⁺), R_t=0.87 min.


Synthesis of tert-butyl (3R,4S,5S)-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-4-(4-(methoxymethyl)-1H-1,2,3-triazol-1-yl)-5-methylpiperidin-3-ylcarbamate



126


In a high pressure vial was added tert-butyl (3R,4S,5S)-4-azido-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-3-ylcarbamate (1.0 equiv.), copper in charcoal (0.2 equiv.), 3-methoxyprop-1-yne (1.5 equiv.) and triethylamine (1.5 equiv.), in dioxane (0.20 M) to give a black suspension. The pressure tube was sealed and the mixture was stirred with heating to 70° C. for 16 hrs. The reaction was cooled to RT, filtered through celite, concentrated to yield tert-butyl (3R,4S,5S)-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-4-(4-(methoxymethyl)-1H-1,2,3-triazol-1-yl)-5-methylpiperidin-3-ylcarbamate in 95% yield. LC/MS (m/z)=653.2 (MH⁺), R_t=0.86 min.

Synthesis of tert-butyl ((3S,4S,5R)-1-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-4-cyano-5-methylpiperidin-3-yl)carbamate



To a solution of (3R,4R,5S)-1-(3-(bis(tert-butoxycarbonyl)amino)pyridin-4-yl)-3-(tert-butoxycarbonylamino)-5-methylpiperidin-4-ylmethanesulfonate (1.0 equiv.) in DMF (0.10 M) was added NaCN (5.0 equiv.). The mixture was stirred at 80° C. for 6 hrs and partitioned between EtOAc and H₂O. The organic layer was washed NaCl_(sat.), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography to yield tert-butyl ((3S,4S,5R)-1-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-4-cyano-5-methylpiperidin-3-yl)carbamate in 5% yield. LC/MS (m/z)=432.2 (MH⁺), R_t=0.73 min.

Synthesis of tert-butyl (3S,4S,5R)-1-(3-aminopyridin-4-yl)-4-cyano-5-methylpiperidin-3-ylcarbamate




Tert-butyl ((3S,4S,5R)-1-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-4-cyano-5-methylpiperidin-3-yl)carbamate (1.0 equiv.) was treated with 4 M HCl in dioxane (30.0 equiv.) at rt for 1 hour. The volatiles were removed in vacuo and the solid was pumped on for 5 minutes on the high vac. To the residue was added CH₂Cl₂ (0.05 M), DIEA (5.0 equiv.) and Boc-OSu (1.6 equiv.). The solution was left stirring at rt for 1 hr. The volatiles were removed in vacuo and the residue was

127


partitioned between EtOAc and H₂O. The organic layer was washed with Na₂CO₃(sat.), NaCl(sat.), dried over MgSO₄, filtered, concentrated to yield tert-butyl (3S,4S,5R)-1-(3-aminopyridin-4-yl)-4-cyano-5-methylpiperidin-3-ylcarbamate in 100% yield. LC/MS (m/z)=332.1 (MH⁺), Rt=0.59 min.

Synthesis of tert-butyl (4-((3R,4S,5S)-4-azido-3-((tert-butoxycarbonyl)amino)-5-methylpiperidin-1-yl)pyridin-3-yl)(tert-butoxycarbonyl)carbamate



To a solution of (3R,4R,5S)-1-(3-(bis(tert-butoxycarbonyl)amino)pyridin-4-yl)-3-((tert-butoxycarbonyl)amino)-5-methylpiperidin-4-ylmethanesulfonate (1.0 equiv.) in DMF (0.13 M) was added NaN₃ (5.0 equiv.). The solution was submerged in an 80° C. oil bath and left stirring under Ar for 24 hrs. The solution was cooled to rt and left stirring under Ar overnight. The solution was partitioned between EtOAc and H₂O. The organic layer was washed with Na₂CO₃(sat.), NaCl(sat.), dried over MgSO₄, filtered, concentrated to yield tert-butyl (4-((3R,4S,5S)-4-azido-3-((tert-butoxycarbonyl)amino)-5-methylpiperidin-1-yl)pyridin-3-yl)(tert-butoxycarbonyl)carbamate in 60% yield. LC/MS (m/z)=548.4 (MH⁺), R_f=0.94 min. ¹H NMR (400 MHz, <cdcl3>) δ ppm 0.97-1.13 (m, 3H), 1.33-1.52 (m, 30H), 2.03-2.19 (m, 1H), 2.66-2.87 (m, 2H), 3.16 (dd, J=12.72, 2.15 Hz, 1H), 3.22-3.32 (m, 1H), 3.81-4.01 (m, 2H), 4.78 (d, J=9.00 Hz, 1H), 6.76-6.88 (m, 1H), 8.05-8.18 (m, 1H), 8.26-8.37 (m, 1H).

Synthesis of tert-butyl ((3R,4S,5S)-1-(3-aminopyridin-4-yl)-4-azido-5-methylpiperidin-3-yl)carbamate




A solution of 4 M HCl in dioxane (30.0 equiv.) was added to tert-butyl (4-(3R,4S,5S)-4-azido-3-((tert-butoxycarbonyl)amino)-5-methylpiperidin-1-yl)pyridin-3-yl)(tert-butoxycarbonyl)carbamate (1.0 equiv.). The solution started to go homogeneous for a few minutes, but then a ppt formed and the solution went very thick. After sitting at rt for 1 hour, the volatiles were removed in vacuo and the solid was pumped on for 5 minutes on the high vac. To the residue was added CH₂Cl₂ (0.11 M), TEA (5.0 equiv.) and Boc₂O (1.0 equiv.). The solution was left stirring at rt for 1 hr. The volatiles were

128


removed in vacuo and the residue was partitioned between EtOAc and H₂O. The organic layer was washed with Na₂CO₃(sat.), NaCl(sat.), dried over MgSO₄, filtered, concentrated and purified by ISCO SiO₂ chromatography to yield tert-butyl ((3R,4S,5S)-1-(3-aminopyridin-4-yl)-4-azido-5-methylpiperidin-3-yl)carbamate in 33% yield. LC/MS (m/z)=348.2 (MH⁺), R_f=0.70 min. ¹H NMR (400 MHz, CHLOROFORM-d) δ ppm 1.02-1.18 (m, 3H), 1.36-1.54 (m, 10H), 2.19 (qd, J=6.91, 3.91 Hz, 1H), 2.57 (q, J=10.96 Hz, 2H), 2.96 (d, J=9.00 Hz, 1H), 3.20 (dd, J=11.15, 3.72 Hz, 1H), 3.55-3.73 (m, 2H), 3.90 (br. s., 1H), 4.01 (br. s., 1H), 4.81 (d, J=8.61 Hz, 1H), 6.72-6.83 (m, 1H), 7.96 (d, J=5.09 Hz, 1H), 8.02 (s, 1H).

Synthesis of tert-butyl ((3R,4S,5S)-4-amino-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-3-yl)carbamate



EDC (1.5 equiv.) and HOAt (1.5 equiv.) were added to a solution of tert-butyl ((3R,4S,5S)-1-(3-aminopyridin-4-yl)-4-azido-5-methylpiperidin-3-yl)carbamate (1.0 equiv.) and 6-(2,6-difluorophenyl)-5-fluoropicolinic acid (1.3 equiv.) in DMF (0.20 M). The mixture was stirred at ambient temperature overnight. The reaction mixture was diluted with water and extracted with ethyl acetate. The combined extracts were washed sequentially with 1M aqueous sodium hydroxide and brine, dried over sodium sulfate, filtered, concentrated and purified by ISCO SiO₂ chromatography. To a degassed solution of the azide (1.0 equiv.) in 2-propanol (0.10 M) was added Pd/C (0.2 equiv.). The mixture was stirred under H₂ for 48 hrs. Filter the mixture over celite and wash the cake with MeOH. Concentrate the filtrate to yield tert-butyl ((3R,4S,5S)-4-amino-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-3-yl)carbamate in 58% yield. LC/MS (m/z)=557.1 (MH⁺), R_f=0.69 min.


Synthesis of N-(4-((3R,4S,5S)-4-acetamido-3-amino-5-methylpiperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide



129


Method 4 was followed using tert-butyl ((3R,4S,5S)-4-amino-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-3-yl)carbamate and acetic anhydride to give N-(4-((3R,4S,5S)-4-acetamido-3-amino-5-methylpiperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 40% yield. LC/MS (m/z)=499.1 (MH⁺), R_t=0.58 min.

Synthesis of N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonamido)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide



Method 4 was followed using tert-butyl ((3R,4S,5S)-4-amino-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-3-yl)carbamate and methanesulfonyl chloride to give N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonamido)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 28% yield. LC/MS (m/z)=535.0 (MH⁺), R_t=0.58 min.

Synthesis of di-tert-butyl 4-((3R,4S,5S)-3-((tert-butoxycarbonyl)amino)-5-methyl-4-(methylamino)piperidin-1-yl)pyridin-3-yl)iminodibutylcarbonate




To a solution of di-tert-butyl 4-((3R,4S,5S)-4-azido-3-((tert-butoxycarbonyl)amino)-5-methylpiperidin-1-yl)pyridin-3-yliminodibutylcarbonate (1.0 equiv.) in DCM (0.14 M) at rt was added PMe₃ (2.0 equiv.). After stirring at rt for 2 hr, PARA-FORMALDEHYDE (5.0 equiv.) was added and the mixture was stirred at rt for another 2.5 hrs. The reaction was added MeOH (0.14 M), cooled to 0° C. and added NaBH₄ (5.0 equiv.). After 30 min at rt, the reaction was quenched with sat. NaHCO₃ and extract with EtOAc to yield di-tert-butyl 4-((3R,4S,5S)-3-((tert-butoxycarbonyl)amino)-5-methyl-4-

130


(methylamino)piperidin-1-yl)pyridin-3-yliminodibutylcarbonate in 85% yield. LC/MS (m/z)=536.3 (MH⁺), R_t=0.61 min.

Synthesis of methyl ((3R,4S,5S)-1-(3-aminopyridin-4-yl)-3-((tert-butoxycarbonyl)amino)-5-methylpiperidin-4-yl)(methyl)carbamate



To a solution of di-tert-butyl 4-((3R,4S,5S)-3-((tert-butoxycarbonyl)amino)-5-methyl-4-(methylamino)piperidin-1-yl)pyridin-3-yliminodibutylcarbonate (1.0 equiv.) in DCM (0.10 M) was added DIEA (3.0 equiv.) the reaction mixture was then cooled to 0° C. To this solution was added methyl chloroformate (1.2 equiv.). The resulting mixture was at RT for 50 min. The reaction mixture was quenched with NaHCO₃ and diluted with EtOAc. The aqueous layer was separated and extracted with EtOAc, the combined organics were then dried over MgSO₄ and concentrated in vacuo. 4 M HCl (43.0 equiv.) in dioxane was added to the residue. After 1 hr, the volatile was removed in vacuo. To the solution of the residue in DCM (0.10 M) at 0° C. was added DIEA (3.0 equiv.) and BocOSu (1.0 equiv.). After 60 min at rt, the reaction mixture was quenched with NaHCO₃ and diluted with EtOAc. The aqueous layer was separated and extracted with EtOAc, the combined organics were then dried over MgSO₄ and concentrated in vacuo to yield a yellow residue, which was purified by ISCO SiO₂ chromatography to yield methyl ((3R,4S,5S)-1-(3-aminopyridin-4-yl)-3-((tert-butoxycarbonyl)amino)-5-methylpiperidin-4-yl)(methyl)carbamate in 44% yield. LC/MS (m/z)=394.2 (MH⁺), R_t=0.59 min. ¹H NMR (400 MHz, <cdcl3>) δ ppm 1.06 (d, J=7.04 Hz, 3H), 1.40-1.52 (m, 10H), 2.28-2.42 (m, 1H), 2.89 (d, J=4.70 Hz, 2H), 3.08 (dd, J=11.93 Hz, 4.50 Hz, 1H), 3.15 (s, 3H), 3.47 (dd, J=11.15, 4.11 Hz, 1H), 3.67-3.79 (m, 5H), 4.13-4.23 (m, 1H), 4.56-4.77 (m, 1H), 6.80 (d, J=5.09 Hz, 1H), 7.97 (d, J=5.48 Hz, 1H), 8.04 (s, 1H).


Synthesis of tert-butyl ((3R,4S,5S)-1-(3-aminopyridin-4-yl)-5-methyl-4-(N-methylacetamido)piperidin-3-yl)carbamate



131

To a solution of di-tert-butyl 4-((3R,4S,5S)-3-(tert-butoxycarbonylamino)-5-methyl-4-(methylamino)piperidin-1-yl)pyridin-3-yliminodicarbonate (1.0 equiv.) in DCM (0.10 M) was added DIEA (3.0 equiv.) the reaction mixture was then cooled to 0° C. To this solution was added acetic anhydride (1.2 equiv.). The resulting mixture was at RT for 50 min. The reaction mixture was quenched with NaHCO₃ and diluted with EtOAc. The aqueous layer was separated and extracted with EtOAc, the combined organics were then dried over MgSO₄ and concentrated in vacuo. 4 M HCl (43.0 equiv.) in Dioxane was added to the residue. After 1 hr, the volatile was removed in vacuo. To the solution of the residue in DCM (0.10 M) at 0° C. was added DIEA (3.0 equiv.) and BocOSu (1.0 equiv.). After 60 min at rt, the reaction mixture was quenched with NaHCO₃ and diluted with EtOAc. The aqueous layer was separated and extracted with EtOAc, the combined organics were then dried over MgSO₄ and concentrated in vacuo to yield a yellow residue, which was purified by SiO₂ chromatography to yield tert-butyl ((3R,4S,5S)-1-(3-aminopyridin-4-yl)-5-methyl-4-(N-methylacetamido)piperidin-3-yl)carbamate in 60% yield. LC/MS (m/z)=378.2 (MH⁺), R_t=0.50 min. ¹HNMR (400 MHz, <cdcl3>) δ ppm 1.06 (d, J=7.83 Hz, 3H), 1.39-1.50 (m, 9H), 2.20 (br. s., 3H), 2.34-2.48 (m, 1H), 2.84-3.28 (m, 6H), 3.77 (d, J=18.00 Hz, 2H), 4.19-4.62 (m, 1H), 6.82 (d, J=5.09 Hz, 1H), 7.97 (br. s., 1H), 8.05 (br. s., 1H).


Synthesis of tert-butyl tert-butoxycarbonyl(4-((3R,4S,5S)-3-((tert-butoxycarbonyl)amino)-4-(4-(hydroxymethyl)-1H-1,2,3-triazol-1-yl)-5-methylpiperidin-1-yl)pyridin-3-yl)carbamate



In a round-bottom flask was added tert-butyl 4-((3R,4S,5S)-4-azido-3-((tert-butoxycarbonyl)amino)-5-methylpiperidin-1-yl)pyridin-3-yl)tert-butoxycarbonyl carbamate (1.0 equiv.), copper in charcoal (0.02 equiv.) and propargyl alcohol (1.0 equiv.) in t-BuOH/H₂O (0.15 M) to give a blue solution. And sodium ascorbate (0.1 equiv.) was added. The mixture was stirred at rt for 16 hrs. The reaction was diluted with H₂O and cooled to 0° C., then filtered, and the precipitate was collected to yield tert-butyl tert-butoxycarbonyl(4-((3R,4S,5S)-3-((tert-butoxycarbonyl)amino)-4-(4-(hydroxymethyl)-1H-1,2,3-triazol-1-yl)-5-methylpiperidin-1-yl)pyridin-3-yl)carbamate in 85% yield. LC/MS (m/z)=604.3 (MH⁺), R_t=0.69 min.


132

Synthesis of tert-butyl (3R,4S,5S)-1-(3-aminopyridin-4-yl)-4-(4-(hydroxymethyl)-1H-1,2,3-triazol-1-yl)-5-methylpiperidin-3-ylcarbamate



To di-tert-butyl 4-((3R,4S,5S)-3-(tert-butoxycarbonylamino)-4-(4-(hydroxymethyl)-1H-1,2,3-triazol-1-yl)-5-methylpiperidin-1-yl)pyridin-3-yliminodicarbonate (1.0 equiv.) was added 4 M HCl (30.0 equiv.) in dioxane. After 1 hr, the volatile was removed in vacuo. To the solution of the residue in DCM (0.10 M) at 0° C. was added DIEA (30.0 equiv.) and BocOSu (1.0 equiv.). After 4 hrs at rt, the reaction mixture was quenched with NaHCO₃ and diluted with EtOAc. The aqueous layer was separated and extracted with EtOAc, the combined organics were then dried over MgSO₄ and concentrated in vacuo to yield a yellow residue, which was purified by ISCO SiO₂ chromatography to yield tert-butyl (3R,4S,5S)-1-(3-aminopyridin-4-yl)-4-(4-(hydroxymethyl)-1H-1,2,3-triazol-1-yl)-5-methylpiperidin-3-ylcarbamate in 57% yield. LC/MS (m/z)=404.3 (MH⁺), R_t=0.47 min.

Synthesis of N-(4-((3R,4S,5S)-3-amino-4-(4-(hydroxymethyl)-1H-1,2,3-triazol-1-yl)-5-methylpiperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide




To a round-bottomed flask was added tert-butyl (3R,4S,5S)-1-(3-aminopyridin-4-yl)-4-(4-(hydroxymethyl)-1H-1,2,3-triazol-1-yl)-5-methylpiperidin-3-ylcarbamate (1.0 equiv.) and BSA (2.5 equiv.) in Acetonitrile/DMF (4/1, 0.05 M) to give an orange suspension. The mixture was stirred at room temperature for 1 hr at which time all of the solids had dissolved into solution. The mixture was taken to dryness. The crude in DMF (0.10 M) was added EDCI (2.4 equiv.), HOAT (2.4 equiv.) and 6-(2,6-difluorophenyl)-5-fluoropicolinic

133


acid (2.4 equiv.). The mixture was stirred at rt for 16 hrs. The reaction was taken to dryness and dissolved in EtOAc and cooled to 0° C. to precipitate the urea byproduct. The mixture was filtered and the organics were washed with H₂O, dried over MgSO₄, filtered and concentrated. To a solution of the residue in EtOH (0.01 M) was added potassium carbonate (10.0 equiv.). The mixture was stirred at room temperature for 2 hrs. The reaction was diluted with DCM and quenched with NaHCO₃(aq). The organics were separated and the aqueous solution was extracted with DCM. The combined organics were dried over MgSO₄, filtered, and concentrated. The crude alcohol was then dissolved in 25% TFA/DCM. The solution was stirred at room temp for 1 hr. The volatiles were removed on the rotovap. The crude product was taken up in DMSO, filtered, purified via prep-HPLC to yield N-(4-((3R,4S,5S)-3-amino-4-(4-(hydroxymethyl)-1H-1,2,3-triazol-1-yl)-5-methylpiperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide in 69% yield. LC/MS (m/z)=539.3 (MH⁺), R_t=0.57 min.

Synthesis of (+/-)-tert-butyl ((1R,3S,5R)-5-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-3-methyl-2-oxocyclohexyl)carbamate



To a solution of (+/-)-tert-butyl ((1R,2R,3S,5R)-5-(3-((tert-butoxycarbonyl)amino)pyridin-4-yl)-2-hydroxy-3-methylcyclohexyl)carbamate (1.0 equiv.) DCM (0.10 M) was added Dess-Martin Periodinane (1.2 equiv.). The mixture was stirred at room temperature for 18 hrs. The volatiles were removed under reduced pressure. The residue was partitioned between EtOAc and 10% sodium thiosulfate/sat. sodium bicarbonate aqueous solution. The organic was washed with brine, dried over sodium sulfate, filtered, concentrated and purified by ISCO chromatography to afford (+/-)-tert-butyl ((1R,3S,5R)-5-(3-acetamidopyridin-4-yl)-3-methyl-2-oxocyclohexyl)carbamate in 65% yield. LC/MS (m/z)=420.2 (MH⁺), R_t=0.73 min.

Synthesis of (+/-)-benzyl ((1R,3S,5R)-5-(3-((benzoxycarbonyl)amino)pyridin-4-yl)-3-methyl-2-oxocyclohexyl)carbamate




(+/-)-Tert-butyl ((1R,3S,5R)-5-(3-acetamidopyridin-4-yl)-3-methyl-2-oxocyclohexyl)carbamate (1.0 equiv.) was

134


dissolved in 4.0 M HCl (50.0 equiv.) in p-dioxane. After stirring at room temperature for 16 hrs, the mixture was concentrated. The crude was dissolved in DCM (0.20 M) and CBZ-OSu (5.0 equiv.) was added followed by DIEA (8.0 equiv.). The reaction was stirred at room temperature for 20 hrs. The solution was diluted with EtOAc and washed with water, sat. sodium bicarbonate, and brine and dried over sodium sulfate, filtered and concentrated. The crude residue was purified by ISCO chromatography to afford (+/-)-benzyl ((1R,3S,5R)-5-(3-((benzoxycarbonyl)amino)pyridin-4-yl)-3-methyl-2-oxocyclohexyl)carbamate in 50% yield. LC/MS (m/z)=488.2 (MH⁺), R_t=0.79 min.

Synthesis of (+/-)-benzyl ((6R,8R,10S)-8-(3-((benzoxycarbonyl)amino)pyridin-4-yl)-10-methyl-1,4-dioxaspiro[4.5]decan-6-yl)carbamate



(+/-)-Benzyl ((1R,3S,5R)-5-(3-((benzoxycarbonyl)amino)pyridin-4-yl)-3-methyl-2-oxocyclohexyl)carbamate (1.0 equiv.) was dissolved in dry THF (0.20 M) under nitrogen and ETHYLENE GLYCOL (8.0 equiv.) was added followed by BF₃.OEt₂ (1.4 equiv.). The solution was microwave vial and heated at 100° C. for 60 mins. The solution was diluted with EtOAc and washed with sat. sodium bicarbonate, water, brine, dried over sodium sulfate and concentrated. The crude material was purified by ISCO chromatography to afford (+/-)-benzyl ((6R,8R,10S)-8-(3-((benzoxycarbonyl)amino)pyridin-4-yl)-10-methyl-1,4-dioxaspiro[4.5]decan-6-yl)carbamate in 100% yield. LC/MS (m/z)=532.2 (MH⁺), R_t=0.81 min.

Synthesis of benzyl (6S,8S,10R)-8-(3-aminopyridin-4-yl)-10-methyl-1,4-dioxaspiro[4.5]decan-6-ylcarbamate and benzyl ((6R,8R,10S)-8-(3-aminopyridin-4-yl)-10-methyl-1,4-dioxaspiro[4.5]decan-6-yl)carbamate



Single Enantiomer

Single Enantiomer

135

(+/-)-Benzyl ((6R,8R,10S)-8-(3-((benzoxycarbonyl)amino)pyridin-4-yl)-10-methyl-1,4-dioxaspiro[4.5]decan-6-yl)carbamate (1.0 equiv.) was dissolved in MeOH (0.10 M) and degassed with Argon to vacuum. Pd/C (0.05 equiv.) was added and the mixture was stirred under a H₂ balloon for 20 hrs. The mixture was filtered and concentrated. The crude and CBZ-OSu (0.99 equiv.) were dissolved in DCM (0.20 M). The reaction was stirred at room temperature for 2 hrs. The solution was diluted with EtOAc and washed with water, dried over sodium sulfate, filtered and concentrated. The crude residue was purified by ISCO chromatography. Purification was completed via SFC (40% EtOH, 15 mL/min, OJ column) to yield benzyl (6S,8S,10R)-8-(3-aminopyridin-4-yl)-10-methyl-1,4-dioxaspiro[4.5]decan-6-ylcarbamate (28% yield, 99% ee) and benzyl (6R,8R,10S)-8-(3-aminopyridin-4-yl)-10-methyl-1,4-dioxaspiro[4.5]decan-6-ylcarbamate (13% yield, 99% ee). LC/MS (m/z)=398.2 (MH⁺), R_f=0.64 min.

Method 7

A solution of N-Boc protected amine was treated with excess 4M HCl/dioxane for 14 hours or with 25% TFA/

136

CH₂Cl₂ for 2 hours. Upon removal of the volatiles in vacuo, the material was purified by RP HPLC yielding after lyophilization the amide product as the TFA salt. Alternatively, the HPLC fractions could be added to EtOAc and solid Na₂CO₃, separated and washed with NaCl_(sat.). Upon drying over MgSO₄, filtering and removing the volatiles in vacuo the free base was obtained. Upon dissolving in MeCN/H₂O, adding 1 eq. of 1 N HCl and lyophilizing, the HCl salt of the amide product was obtained.

If an OBn or NCbz group was present, it was deprotected by treatment with 10% Pd/C (0.2 equiv.) under an atmosphere of hydrogen in ethyl acetate and methanol (1:2). Upon completion, the reaction was filtered through Celite, washed with methanol, and the filtrate was concentrated in vacuo.

If a CO₂Me group was present, it could be converted to the corresponding CO₂H following Method 2.

Following the procedures of Method 7, the following compounds were prepared:

TABLE 1

Ex #	Structure	LC/MS (MH+ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name	
1		Chiral	508.2	0.60	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
2		Chiral	528.2	0.62	N-(4-((1R,3R,4S,5S)-3-amino-4-(2-methoxyacetamido)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 1-continued

Ex #	Structure	LC/MS (MH+ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
3		Chiral	526.3 0.67	N-(4-((1R,3R,4S,5S)-3-amino-4-isobutyramido-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
4		Chiral	591.2 0.72	N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(4-(thiophen-3-yl)-1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
5		Chiral	509.2 0.57	N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
6		Chiral	470.3 0.61	N-(4-((1R,3R,5S,Z)-3-amino-4-(hydroxyimino)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 1-continued

Ex #	Structure		LC/MS (MH+ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
7		Chiral	523.1	0.60	N-((3R,4S,5S)-3-amino-5-methyl-4-(4-methyl-1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
8		Chiral	528.2	0.65	ethyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate
9		Chiral	542.3	0.68	isopropyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate
10		Chiral	512.1	0.62	N-((1R,3R,4S,5S)-3-amino-5-methyl-4-propionamidocyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 1-continued

Ex #	Structure		LC/MS (MH+ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
11		Chiral	455.1	0.57	N-(4-((1R,3R,5S)-3-amino-5-methyl-4-oxocyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
12		Chiral	484.2	0.63	N-(4-((1R,3R,5S,E)-3-amino-4-(methoxyimino)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
13		Chiral	484.2	0.64	N-(4-((1R,3R,5S,Z)-3-amino-4-(methoxyimino)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
14		Chiral	553.2	0.60	N-(4-((3R,4S,5S)-3-amino-4-(4-(methoxymethyl)-1H-1,2,3-triazol-1-yl)-5-methylpiperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 1-continued

Ex #	Structure		LC/MS	LC/MS	Chemical Name
			(MH+ on UPLC)	(Rf on UPLC)	
15		Chiral	549.2	0.58	N-(4-((1R,3R,4S,5S)-3-amino-4-(2-hydroxyethylsulfonyl)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
16		Chiral	534.2	0.57	N-(4-((1S,3S,4S,5R)-3-amino-5-methyl-4-(methylsulfonylamido)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
17		Chiral	534.2	0.57	N-(4-((1R,3R,4R,5S)-3-amino-5-methyl-4-(methylsulfonylamido)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
18		Chiral	457.0	0.57	N-(4-((3R,4S,5S)-3,4-diamino-5-methylpiperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 1-continued

Ex #	Structure		LC/MS	LC/MS	Chemical Name
			(MH+ on UPLC)	(Rf on UPLC)	
19		Chiral	535.0	0.58	N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonamido)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
20		Chiral	563.3	0.61	N-(4-((1R,3R,4S,5S)-3-amino-4-(2-methoxyethyl)sulfonyl)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
21		Chiral	499.1	0.58	N-(4-((3R,4S,5S)-4-acetamido-3-amino-5-methylpiperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
22		Chiral	514.2	0.61	methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate

TABLE 1-continued

Ex #	Structure		LC/MS	LC/MS	Chemical Name
			(MH+ on UPLC)	(Rf on UPLC)	
23		Chiral	514.3	0.61	methyl (1R,2S,4S,6R)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate
24		Chiral	498.2	0.59	N-(4-((1S,3S,4R,5R)-4-acetamido-3-amino-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
25		Chiral	498.3	0.59	N-(4-((1R,3R,4S,5S)-4-acetamido-3-amino-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
26		Chiral	534.2	0.58	N-(4-((1S,3S,4R,5R)-3-amino-5-methyl-4-(methylsulfonamido)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 1-continued

Ex #	Structure		LC/MS	LC/MS	Chemical Name
			(MH+ on UPLC)	(Rf on UPLC)	
27		Chiral	534.2	0.58	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonamido)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
28		Chiral	482.1	0.64	N-(4-((1S,3S,4R,5R)-3-amino-4-azido-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
29		Chiral	482.2	0.63	N-(4-((1R,3R,4S,5S)-3-amino-4-azido-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
30		Chiral	498.3	0.57	N-(4-((1R,3R,4R,5S)-4-acetamido-3-amino-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 1-continued

Ex #	Structure		LC/MS	LC/MS	Chemical Name
			(MH+ on UPLC)	(Rf on UPLC)	
31		Chiral	503.2	0.59	N-(4-((1R,3R,4R,5S)-3-amino-5-methyl-4-((R)-methylsulfinyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
32		Chiral	503.2	0.57	N-(4-((1R,3R,4R,5S)-3-amino-5-methyl-4-((S)-methylsulfinyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
33		Chiral	487.2	0.64	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylthio)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
34		Chiral	519.2	0.60	N-(4-((1R,3R,4R,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 1-continued

Ex #	Structure		LC/MS	LC/MS	Chemical Name
			(MH+ on UPLC)	(Rf on UPLC)	
35		Chiral	487.2	0.65	N-(4-((1R,3R,4R,5S)-3-amino-5-methyl-4-(methylthio)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
36		Chiral	503.2	0.57	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-((R)-methylsulfinyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
37		Chiral	519.2	0.58	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
38		Chiral	533.3	0.60	(1R,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl dimethylphosphinate

TABLE 1-continued

Ex #	Structure		LC/MS (MH+ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
39		Chiral	533.3	0.59	(1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl dimethylphosphinate
40		Chiral	535.2	0.60	(1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl methanesulfonate
41		Chiral	535.2	0.61	(1R,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl methanesulfonate

Method 8

A homogeneous solution of 1 eq each of amine, carboxylic acid, HOAT and EDC in DMF, at a concentration of 0.5 M, was left standing for 24 hours at which time water and ethyl acetate were added. The organic phase was dried with sodium sulfate and purified via silica gel column chromatography eluting with ethyl acetate and hexanes to give the desired protected amide product. Alternatively the crude reaction mixture was directly purified by HPLC. Upon lyophilization, the TFA salt of the protected amide product was obtained. Alternatively, the HPLC fractions could be added to EtOAc and solid Na₂CO₃, separated and washed with NaCl_(sat.). Upon drying over MgSO₄, filtering and removing the volatiles in vacuo, the protected amide product was obtained as a free base. Alternatively, the crude reaction mixture was used for the deprotection step without further purification.

If an N-Boc protected amine was present, it was removed by treating with excess 4M HCl/dioxane for 14 hours or by

treating with 25% TFA/CH₂Cl₂ for 2 hours. Upon removal of the volatiles in vacuo, the material was purified by RP HPLC yielding after lyophilization the amide product as the TFA salt. Alternatively, the HPLC fractions could be added to EtOAc and solid Na₂CO₃, separated and washed with NaCl_(sat.). Upon drying over MgSO₄, filtering and removing the volatiles in vacuo the free base was obtained. Upon dissolving in MeCN/H₂O, adding 1 eq. of 1 N HCl and lyophilizing, the HCl salt of the amide product was obtained.

If an N-Boc, OAc group were present, prior to Boc deprotection, the acetate group could be cleaved by treating with K₂CO₃ (2.0 equiv.) in ethanol at a concentration of 0.1 M for 24 hours.

If a TBDMS ether was present, it was deprotected prior to Boc removal by treating with 6N HCl, THF, methanol (1:2:1) at room temperature for 12 h. After removal of volatiles in vacuo, the Boc amino group was deprotected as described above. Alternatively, the TBDMS ether and Boc group could

157

be both deprotected with 6N HCl, THF, methanol (1:2:1) if left at rt for 24 hours, or heated at 60° C. for 3 hours.

If a OBn or Cbz protecting group was present, it was deprotected by treatment with 10% Pd/C (0.2 equiv.) under an atmosphere of hydrogen in ethyl acetate and methanol (1:2). Upon completion, the reaction was filtered through Celite,

158

washed with methanol, and the filtrate was concentrated in vacuo.

If a CO₂Me group was present, it could be converted to the corresponding CO₂H following Method 2.

Following the procedures of Method 8, the following compounds were prepared:

TABLE 2

Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
42		Chiral	499.2	0.64	N-((6R,8R,10S)-6-amino-10-methyl-1,4-dioxaspiro[4.5]decan-8-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
43		Chiral	499.2	0.64	N-((6S,8S,10R)-6-amino-10-methyl-1,4-dioxaspiro[4.5]decan-8-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
44		Chiral	529.3	0.63	3-((1R,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyloxy)propanoic acid

TABLE 2-continued

Ex#	Structure		LC/MS (MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
45		Chiral	529.3	0.62	3-((1S,2S,4S,6R)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyloxy)propanoic acid
46		Chiral	531.3	0.59	methyl (1R,2S,4S,6R)-2-amino-4-(3-(5-amino-2-(2,6-difluorophenyl)thiazole-4-carboxamido)pyridin-4-yl)-6-methylcyclohexyl(methyl) carbamate
47		Chiral	531.3	0.58	methyl (1S,2R,4R,6S)-2-amino-4-(3-(5-amino-2-(2,6-difluorophenyl)thiazole-4-carboxamido)pyridin-4-yl)-6-methylcyclohexyl(methyl) carbamate
48		Chiral	508.3	0.59	N-(4-((1S,3S,4R,5R)-3-amino-5-methyl-4-(1H-1,2,4-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 2-continued





Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
49		Chiral	508.3	0.59	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,4-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
50		Chiral	459.2	0.58	N-(4-((1R,3R,4S,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide
51		Chiral	454.3	0.54	N-(4-((1R,3R,4S,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)pyrimidine-4-carboxamide
52		Chiral	542.2	0.68	methyl (1R,2S,4S,6R)-2-amino-4-(3-(6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl(methyl) carbamate

TABLE 2-continued





Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
53		Chiral	542.2	0.67	methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl(methyl) carbamate
54		Chiral	470.0	0.57	5-amino-N-(4-((3S,4S,5R)-3-amino-4-cyano-5-methylpiperidin-1-yl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide
55		Chiral	481.1	0.66	N-(4-((3S,4S,5R)-3-amino-4-cyano-5-methylpiperidin-1-yl)pyridin-3-yl)-6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamide
56		Chiral	509.2	0.57	N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,4-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 2-continued

Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
57		Chiral	467.0	0.60	N-(4-((3S,4S,5R)-3-amino-4-cyano-5-methylpiperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
58		Chiral	543.2	0.67	methyl (3R,4S,5S)-3-amino-1-(3-(6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-4-yl(methyl)carbamate
59		Chiral	528.1	0.64	methyl (1R,2S,4S,6R)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl(methyl)carbamate
60		Chiral	528.2	0.63	methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl(methyl)carbamate

TABLE 2-continued





Ex#	Structure		LC/MS (MH+ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
61		Chiral	529.2	0.63	methyl (3R,4S,5S)-3-amino-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-4-yl(methyl)carbamate
62		Chiral	485.1	0.69	N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamide
63		Chiral	513.2	0.61	5-amino-N-(4-((1S,3S,4R,5R)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide
64		Chiral	513.2	0.61	5-amino-N-(4-((1R,3R,4S,5S)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide

TABLE 2-continued

Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
65		Chiral	513.2	0.53	N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(N-methylacetamido)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
66		Chiral	539.3	0.57	N-(4-((3R,4S,5S)-3-amino-4-(4-(hydroxymethyl)-1H-1,2,3-triazol-1-yl)-5-methylpiperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
67		Chiral	466.1	0.61	N-(4-((1S,3S,4R,5R)-3-amino-4-cyano-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
68		Chiral	466.1	0.61	N-(4-((1R,3R,4S,5S)-3-amino-4-cyano-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 2-continued

Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
69		Chiral	474.1	0.58	5-amino-N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide
70		Chiral	469.1	0.54	5-amino-N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)pyrimidine-4-carboxamide
71		Chiral	488.2	0.68	5-amino-N-(4-((1R,3R,4S,5S)-3-amino-4-ethoxy-5-methylcyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide
72		Chiral	518.1	0.60	methyl ((3R,4S,5S)-3-amino-1-(2-(5-amino-2-(2,6-difluorophenyl)thiazole-4-carboxamido)phenyl)-5-methylpiperidin-4-yl)carbamate

TABLE 2-continued





Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
73		Chiral	577.2	0.65	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamide
74		Chiral	577.2	0.65	N-(4-((1S,3S,4R,5R)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamide
75		Chiral	471.1	0.66	N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
76		Chiral	471.1	0.67	N-(4-((1S,3S,4S,5R)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 2-continued





Ex#	Structure		LC/MS (MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
77		Chiral	563.3	0.61	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
78		Chiral	563.3	0.61	N-(4-((1S,3S,4R,5R)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
79		Chiral	469.3	0.53	5-amino-N-(4-((1R,3R,4S,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)pyrimidine-4-carboxamide
80		Chiral	529.1	0.69	methyl ((3R,4S,5S)-3-amino-1-(3-(6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-4-yl)carbamate

TABLE 2-continued

Ex#	Structure	Chiral	LC/MS (MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
81		Chiral	515.1	0.64	methyl ((3R,4S,5S)-3-amino-1-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-4-yl)carbamate
82		Chiral	474.2	0.59	5-amino-N-(4-((1R,3R,4S,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide
83		Chiral	524.3	0.67	N-(4-((1R,3R,4S,5S)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamide
84		Chiral	524.3	0.67	N-(4-((1S,3S,4R,5R)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamide

TABLE 2-continued





Ex#	Structure		LC/MS (MH+ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
85		Chiral	510.3	0.62	N-(4-((1R,3R,4S,5S)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
86		Chiral	510.3	0.62	N-(4-((1S,3S,4R,5R)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
87		Chiral	483.1	0.62	N-(4-((3R,4S,5S)-3-amino-4-azido-5-methylpiperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
88		Chiral	499.2	0.73	N-(4-((1R,3R,4S,5S)-3-amino-4-ethoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 2-continued





Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
89		Chiral	499.2	0.73	N-(4-((1S,3S,4R,5R)-3-amino-4-ethoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamide
90		Chiral	485.2	0.68	N-(4-((1S,3S,4R,5R)-3-amino-4-ethoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
91		Chiral	485.2	0.68	N-(4-((1R,3R,4S,5S)-3-amino-4-ethoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamide
92		Chiral	471.1	0.64	N-(4-((1R,3R,4S,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 2-continued




Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
93		Chiral	545.3	0.64	N-(4-((1R,3R,4S,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-5-fluoropicolinamide
94		Chiral	603.3	0.63	N-(4-((1R,3R,4S,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(4-(1,1-dioxidotetrahydro-2H-thiopyran-4-yl)-2,6-difluorophenyl)-5-fluoropicolinamide
95		Chiral	513.2	0.79	N-(4-((1R,3R,4S,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-isopropylphenyl)-5-fluoropicolinamide

TABLE 2-continued




Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
96		Chiral	573.3	0.76	N-(4-((1R,3R,4S,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide
97		Chiral	529.3	0.68	N-(4-((1R,3R,4S,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(2-hydroxypropan-2-yl)phenyl)-5-fluoropicolinamide
98		Chiral	541.3	0.75	N-(4-((1R,3R,4S,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinamide

TABLE 2-continued




Ex#	Structure		LC/MS (MH+ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
99		Chiral	555.3	0.66	N-(4-((1R,3R,4S,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(tetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide
100		Chiral	529.3	0.80	N-(4-((1R,3R,4S,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(4-(ethoxymethyl)-2,6-difluorophenyl)-5-fluoropicolinamide
101		Chiral	529.3	0.86	N-(4-((1R,3R,4S,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-isopropoxyphenyl)-5-fluoropicolinamide

TABLE 2-continued





Ex#	Structure		LC/MS (MH+ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
102		Chiral	485.2	0.73	N-(4-((1R,3R,4S,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamide
103		Chiral	485.4	0.68	N-(4-((1S,3S,4R,5R)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamide
104			502.3	0.50	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)pyrimidine-4-carboxamide
105			389.4	0.62	methyl ((3R,4S,5S)-3-amino-1-(3-(6-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-4-yl)carbamate

TABLE 2-continued

Ex#	Structure	LC/MS (MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
106		588.4	0.63	methyl ((1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl)carbamate
107		619.4	0.65	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-5-fluoropicolinamide
108		589.4	0.62	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinamide

TABLE 2-continued




Ex#	Structure	LC/MS		Chemical Name
		(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
109		583.4	0.58	3-amino-N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-6-(2-fluoro-5-(isopropylcarbamoyl)phenyl)picolinamide
110		621.4	0.66	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide
111		574.4	0.59	3-amino-N-(4-((1R,3R,4S,5S)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-6-(2-fluoro-5-(isopropylcarbamoyl)phenyl)picolinamide

TABLE 2-continued

Ex#	Structure	LC/MS		Chemical Name
		(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
112		549.3	0.63	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinamide
113		534.2	0.69	N-(4-((1S,3S,4R,5R)-3-amino-5-methyl-4-(pyridin-2-yloxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
114		534.3	0.61	3-amino-N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
115		522.2	0.57	5-amino-N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide

TABLE 2-continued




Ex#	Structure	LC/MS (MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
116	 Chemical structure of N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-6-(4-(ethoxymethyl)2,6-difluorophenyl)-5-fluoropicolinamide AND Enantiomer. The structure shows a cyclohexane ring with a methylsulfonyl group, an amino group, and a methyl group, connected to a pyridine ring, which is further connected to a picolinamide moiety. The picolinamide moiety is substituted with a 4-(ethoxymethyl)2,6-difluorophenyl group.	577.3	0.66	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-6-(4-(ethoxymethyl)2,6-difluorophenyl)-5-fluoropicolinamide
117	 Chemical structure of N-(4-((1R,3R,4S,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide. The structure shows a cyclohexane ring with a methoxy group, an amino group, and a methyl group, connected to a pyridine ring, which is further connected to a picolinamide moiety. The picolinamide moiety is substituted with a 2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl group.	571.4	0.59	N-(4-((1R,3R,4S,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide
118	 Chemical structure of N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide. The structure shows a cyclohexane ring with a 2-(methylsulfonyl)ethoxy group, an amino group, and a methyl group, connected to a pyridine ring, which is further connected to a picolinamide moiety. The picolinamide moiety is substituted with a 2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl group.	663.5	0.58	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide

TABLE 2-continued

Ex#	Structure	LC/MS (MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
119		593.3	0.62	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-5-fluoropicolinamide
120		507.2	0.55	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide
121		619.4	0.56	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide
122		533.4	0.63	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamide

TABLE 2-continued

Ex#	Structure		LC/MS (MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
123		Chiral	534.4	0.69	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(pyridin-2-yloxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
124		Chiral	545.3	0.67	N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-5-fluoropicolinamide
125		Chiral	621.3	0.72	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(4-(ethoxymethyl)-2,6-difluorophenyl)-5-fluoropicolinamide
126		Chiral	535.4	0.61	3-amino-N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2-fluoro-5-(isopropylcarbamoyl)phenyl)picolinamide

TABLE 2-continued

Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
127		Chiral	573.3	0.70	N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide
128		Chiral	627.4	0.59	3-amino-N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2-fluoro-5-(isopropylcarbamoyl)phenyl)picolinamide
129		Chiral	665.4	0.68	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide

TABLE 2-continued

Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
130		Chiral	637.4	0.68	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-5-fluoropicolinamide
131		Chiral	497.2	0.62	N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide
132		Chiral	583.3	0.68	N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-5-fluoropicolinamide

TABLE 2-continued

Ex#	Structure		LC/MS (MH+ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
133		Chiral	551.1	0.61	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide
134			459.3	0.69	N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide
135		Chiral	566.2	0.63	5-amino-N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide
136		Chiral	529.3	0.74	N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(4-(ethoxymethyl)-2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 2-continued




Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
137		Chiral	571.4	0.67	N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-5-fluoropicolinamide
138		Chiral	541.3	0.64	N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinamide
139		Chiral	633.4	0.63	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinamide

TABLE 2-continued




Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
140		Chiral	663.4	0.66	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-5-fluoropicolinamide
141		Chiral	621.4	0.81	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-isopropoxyphenyl)-5-fluoropicolinamide
142		Chiral	529.3	0.82	N-(4-((1R,3R,4R,5S)-3-amino-5-methoxy-4-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-isopropoxyphenyl)-5-fluoropicolinamide

TABLE 2-continued

Ex#	Structure		LC/MS (MH+ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
143		Chiral	501.1	0.70	N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinamide
144		Chiral	593.3	0.68	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinamide
145		Chiral	486.1	0.69	3-amino-N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
146		Chiral	578.2	0.66	3-amino-N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 2-continued

Ex#	Structure		LC/MS		Chemical Name
			(MH+ on UPLC)	LC/MS (Rf on UPLC)	
147		Chiral	567.3	0.72	N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluoro-4-isopropoxyphenyl)-5-fluoropicolinamide
148		Chiral	573.3	0.72	methyl (3R,4S,5S)-3-amino-1-(3-(6-(2,6-difluoro-4-isopropoxyphenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-4-ylcarbamate
149		Chiral	572.3	0.74	methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-isopropoxyphenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate

TABLE 2-continued

Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
150		Chiral	534.4	0.60	N-(4-((1S,3S,4R,5R)-3-amino-5-methyl-4-(2-oxopyridin-1(2H)-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
151		Chiral	611.4	0.67	N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide
152		Chiral	534.4	0.60	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-oxopyridin-1(2H)-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 2-continued





Ex#	Structure		LC/MS (MH+ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
153		Chiral	612.4	0.70	N-(4-((1R,3R,4S,5S)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide
154		Chiral	511.2	0.58	5-amino-N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide
155		Chiral	566.3	0.68	N-(4-((1S,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(4-(ethoxymethyl)-2,6-difluorophenyl)-5-fluoropicolinamide
156		Chiral	512.2	0.56	5-amino-N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide

TABLE 2-continued




Ex#	Structure		LC/MS		Chemical Name
			(MH+ on UPLC)	LC/MS (Rf on UPLC)	
157		Chiral	567.3	0.67	N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(4-(ethoxymethyl)-2,6-difluorophenyl)-5-fluoropicolinamide
158		Chiral	585.4	0.62	methyl (3R,4S,5S)-3-amino-1-(3-(6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-4-ylcarbamate
159		Chiral	615.4	0.64	methyl (3R,4S,5S)-3-amino-1-(3-(6-(2,6-difluoro-4-(tetrahydro-2H-pyran-4-yloxy)phenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-4-ylcarbamate

TABLE 2-continued




Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
160		Chiral	609.4	0.59	N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide
161		Chiral	523.3	0.65	N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamide
162		Chiral	571.3	0.58	methyl (3R,4S,5S)-3-amino-1-(3-(6-(2,6-difluoro-4-(oxetan-3-yl)phenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-4-ylcarbamate

TABLE 2-continued




Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
163		Chiral	566.3	0.61	N-(4-((1R,3R,4S,5S)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(oxetan-3-yl)phenyl)-5-fluoropicolinamide
164		Chiral	565.3	0.48	N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluoro-4-(oxetan-3-yl)phenyl)-5-fluoropicolinamide
165		Chiral	540.1	0.67	N-(4-((1R,3R,4S,5S)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinamide

TABLE 2-continued

Ex#	Structure		LC/MS (MH+ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
166		Chiral	525.1	0.66	3-amino-N-(4-((1R,3R,4S,5S)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
167		Chiral	584.4	0.63	methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate
168		Chiral	582.3	0.64	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-5-fluoropicolinamide
169		Chiral	496.2	0.57	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide

TABLE 2-continued




Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	(Rf on UPLC)	
170		Chiral	614.4	0.66	methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-(tetrahydro-2H-pyran-4-yloxy)phenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate
171		Chiral	608.4	0.65	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(tetrahydro-2H-pyran-4-yloxy)phenyl)-5-fluoropicolinamide
172		Chiral	578.4	0.62	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinamide

TABLE 2-continued

Ex#	Structure	LC/MS (MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
173		545.3	0.64	methyl ((3R,4S,5S)-3-amino-1-(3-(6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-4-yl)carbamate
174		530.3	0.62	methyl ((3R,4S,5S)-3-amino-1-(3-(3-amino-6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-4-yl)carbamate
175		529.3	0.65	methyl (1S,2R,4R,6S)-2-amino-4-(3-(3-amino-6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate
176		523.3	0.63	3-amino-N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 2-continued




Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
177		Chiral	615.3	0.59	methyl (3R,4S,5S)-3-amino-1-(3-(6-(2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-4-ylcarbamate
178		Chiral	614.3	0.59	methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate
179		Chiral	608.3	0.58	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide

TABLE 2-continued

Ex#	Structure	LC/MS (MH+ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
180		528.2	0.66	methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate
181		522.2	0.65	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamide
182		570.3	0.61	methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-(oxetan-3-yl)phenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate
183		610.4	0.70	N-(4-((1R,3R,4S,5S)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-5-fluoropicolinamide

TABLE 2-continued




Ex#	Structure	LC/MS (MH+ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
184		580.4	0.66	N-(4-((1R,3R,4S,5S)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinamide
185		533.3	0.66	N-(4-((1R,3R,4S,5S)-3-amino-4-(4-cyano-1H-1,2,3-triazol-1-yl)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
186		548.4	0.67	N-(4-((1R,3R,4S,5S)-3-amino-4-(4-cyclopropyl-1H-1,2,3-triazol-1-yl)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide

TABLE 2-continued




Ex#	Structure		LC/MS		Chemical Name
			(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
187		Chiral	548.3	0.69	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(4-(prop-1-en-2-yl)-1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
188		Chiral	566.4	0.62	N-(4-((1R,3R,4S,5S)-3-amino-4-(4-(2-hydroxypropan-2-yl)-1H-1,2,3-triazol-1-yl)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
189		Chiral	515.3	0.62	2-((1R,2S,4S,6R)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyloxy)acetic acid

TABLE 2-continued





Ex#	Structure		LC/MS (MH+ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
190		Chiral	515.3	0.62	2-((1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyloxy)acetic acid
191		Chiral	544.3	0.66	methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate
192		Chiral	538.3	0.65	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinamide
193		Chiral	539.3	0.64	N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinamide

TABLE 2-continued

Ex#	Structure	Chiral	LC/MS		Chemical Name
			(MH+ on UPLC)	LC/MS (Rf on UPLC)	
194		Chiral	524.3	0.62	3-amino-N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide
195			610.4	0.67	N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide
196			572.4	0.59	3-amino-N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2-fluoro-5-(isopropylcarbamoyl)phenyl)picolinamide

TABLE 2-continued




Ex#	Structure	LC/MS		Chemical Name
		(MH ⁺ on UPLC)	LC/MS (Rf on UPLC)	
197		573.4	0.58	3-amino-N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(2-fluoro-5-(isopropylcarbamoyl)phenyl)picolinamide
198		579.3	0.60	methyl (3R,4S,5S)-3-amino-1-(3-(3-amino-6-(2-fluoro-5-(isopropylcarbamoyl)phenyl)picolinamido)pyridin-4-yl)-5-methylpiperidin-4-ylcarbamate
199		617.4	0.68	methyl (3R,4S,5S)-3-amino-1-(3-(6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-4-ylcarbamate

TABLE 2-continued





Ex#	Structure	LC/MS		Chemical Name	
		(MH+ on UPLC)	LC/MS (Rf on UPLC)		
200		578.4	0.60	methyl (1S,2R,4R,6S)-2-amino-4-(3-(3-amino-6-(2-fluoro-5-(isopropylcarbamoyl)phenyl)picolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate	
201		616.4	0.68	methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate	
202		Chiral	572.2	0.67	methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(4-(ethoxymethyl)-2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate

TABLE 2-continued

Ex#	Structure		LC/MS (MH+ on UPLC)	LC/MS (Rf on UPLC)	Chemical Name
203		Chiral	573.2	0.66	methyl (3R,4S,5S)-3-amino-1-(3-(6-(4-(ethoxymethyl)-2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-4-ylcarbamate

In addition to LC/MS and LC characterization, representative compounds were analyzed by ¹H-NMR. The following

data in Table 3 are typical spectra of the compounds of the invention.

TABLE 3

Ex #	¹ H-NMR data
1	¹ H NMR (400 MHz, Methanol-d) δ ppm 0.65 (d, J = 6.70 Hz, 1 H) 1.81 (d, J = 13.11 Hz, 1 H) 2.00-2.20 (m, 1 H) 2.22-2.37 (m, 1 H) 2.61-2.79 (m, 1 H) 3.34-3.45 (m, 1 H) 3.89 (dt, J = 12.76, 4.30 Hz, 1 H) 4.86 (s, 5 H) 5.00 (br. s., 1 H) 7.22 (t, J = 8.31 Hz, 1 H) 7.64 (tt, J = 8.48, 6.46 Hz, 1 H) 7.83-7.92 (m, 1 H) 8.04 (t, J = 8.73 Hz, 1 H) 8.08 (s, 1 H) 8.42 (dd, J = 8.63, 3.94 Hz, 1 H) 8.60 (br. s., 1 H) 8.90 (br. s., 1 H)
4	¹ H NMR (400 MHz, METHANOL-d ₄) δ ppm 0.48 (d, J = 6.90 Hz, 3 H) 2.40-2.55 (m, 1 H) 2.64 (s, 1 H) 3.49-3.61 (m, 1 H) 3.67-3.79 (m, 1 H) 3.90 (dd, J = 11.40, 4.30 Hz, 1 H) 3.97-4.08 (m, 1 H) 4.14-4.28 (m, 1 H) 4.96 (t, J = 4.38 Hz, 1 H) 7.26 (t, J = 8.39 Hz, 2 H) 7.49-7.57 (m, 3 H) 7.61-7.72 (m, 1 H) 7.82 (dd, J = 2.64, 1.52 Hz, 1 H) 8.05 (t, J = 8.71 Hz, 1 H) 8.33 (s, 1 H) 8.39 (dd, J = 6.58, 0.90 Hz, 1 H) 8.46 (dd, J = 8.68, 3.94 Hz, 1 H) 9.15 (s, 1 H)
5	¹ H NMR (400 MHz, METHANOL-d ₄) δ ppm 0.41 (d, J = 6.85 Hz, 3 H) 2.50 (td, J = 11.57, 4.55 Hz, 1 H) 3.59 (t, J = 12.94 Hz, 1 H) 3.81 (dd, J = 13.60, 3.81 Hz, 1 H) 3.94-4.15 (m, 2 H) 4.18-4.33 (m, 1 H) 5.11 (t, J = 4.16 Hz, 1 H) 7.29 (t, J = 8.49 Hz, 2 H) 7.62 (d, J = 6.75 Hz, 1 H) 7.70 (tt, J = 8.49, 6.50 Hz, 1 H) 7.90 (d, J = 0.93 Hz, 1 H) 8.07 (t, J = 8.68 Hz, 1 H) 8.17 (d, J = 0.68 Hz, 1 H) 8.42 (d, J = 6.65 Hz, 1 H) 8.48 (dd, J = 8.66, 3.96 Hz, 1 H) 9.13 (s, 1 H)
6	¹ H NMR in DMSO-d ₆ : δ 10.94 (s, 1H), 10.49 (s, 1H), 8.60 (s, 1H), 8.48 (d, J = 4.0, 1H), 8.35 (dd, J = 8.0, 4.0, 1H), 8.20 (t, J = 8.0, 1H), 8.02 (broad doublet, J = 4.0, 2H), 7.67-7.74 (m, 1H), 7.42 (d, J = 4.0, 1H), 7.36 (t, J = 8.0, 2H), 4.24 (m, 1H), 3.82-3.86 (m, 1H), 3.21-3.27 (m, 1H), 2.50-2.55 (m, 1H), 2.24 (d, J = 12.0, 1H), 1.86 (d, J = 16.0, 1H), 1.68 (q, J = 12.0, 1H), 1.59 (q, J = 12.0, 1H), 1.40 (d, J = 8.0, 3H).
7	¹ H NMR (400 MHz, METHANOL-d ₄) δ ppm 0.42 (d, J = 6.85 Hz, 3 H) 2.37 (s, 3 H) 2.46 (dd, J = 11.71, 7.02 Hz, 1 H) 3.55 (t, J = 12.84 Hz, 1 H) 3.77 (dd, J = 13.62, 3.99 Hz, 1 H) 3.89-4.09 (m, 2 H) 4.12-4.30 (m, 1 H) 4.93 (t, J = 4.40 Hz, 1 H) 7.27 (t, J = 8.41 Hz, 2 H) 7.56 (d, J = 6.75 Hz, 1 H) 7.68 (tt, J = 8.49, 6.48 Hz, 1 H) 7.83 (d, J = 0.59 Hz, 1 H) 8.06 (t, J = 8.71 Hz, 1 H) 8.36-8.42 (m, 1 H) 8.47 (dd, J = 8.68, 3.94 Hz, 1 H) 9.14 (s, 1 H)
11	¹ H NMR in DMSO-d ₆ : δ 10.55 (s, 1H), 8.55 (s, 1H), 8.47 (d, J = 4.0, 1H), 8.37 (dd, J = 8.0, 4.0, 1H), 8.21 (t, J = 8.0, 1H), 8.16 (broad doublet, J = 4.0, 2H), 7.67-7.74 (m, 1H), 7.40 (d, J = 8.0, 1H), 7.36 (t, J = 8.0, 2H), 4.20-4.26 (m, 1H), 3.50-3.70 (m, 2H), 2.76-2.82 (m, 1H), 2.49-2.54 (m, 1H), 2.32-2.36 (m, 1H), 2.16-2.18 (m, 1 H), 1.91 (q, J = 12, 1H), 1.65 (q, J = 12, 1H), 0.97 (d, J = 8.0, 3H).
12	¹ H NMR in DMSO-d ₆ : δ 10.49 (s, 1H), 8.57 (s, 1H), 8.47 (d, J = 4.0, 1H), 8.35 (dd, J = 8.0, 4.0, 1H), 8.25 (broad doublet, J = 4.0, 2H), 8.20 (t, J = 8.0, 1H), 7.67-7.74 (m, 1H), 7.42 (d, J = 8.0, 1H), 7.36 (t, J = 8.0, 2H), 4.04-4.08 (m, 1H), 3.79 (s, 3H), 3.23-3.29 (m, 1H), 2.39-2.45 (m, 1H), 2.11 (d, J = 8.0, 1H), 2.10 (d, J = 8.0, 1H), 1.90 (q, J = 12, 1H), 1.40 (q, J = 12, 1H), 1.01 (d, J = 4.0, 3H).
13	¹ H NMR in DMSO-d ₆ : δ 10.48 (s, 1H), 8.57 (s, 1H), 8.47 (d, J = 4.0, 1H), 8.35 (dd, J = 8.0, 4.0, 1H), 8.20 (t, J = 8.0, 1H), 8.08 (broad singlet, 2H), 7.67-7.74 (m, 1H), 7.42 (d, J = 8.0, 1H), 7.36 (t, J = 8.0, 2H), 3.88-3.92 (m, 1H), 3.80 (s, 3H), 3.22-3.28 (m, 1H), 2.51-2.58 (m, 1H), 2.25 (d, J = 12.0, 1H), 1.86 (d, J = 12.0, 1H), 1.70 (q, J = 12, 1H), 1.62 (q, J = 12, 1H), 1.34 (d, J = 4.0, 3H).

TABLE 3-continued

Ex # ¹H-NMR data

- 14 ¹H NMR (400 MHz, METHANOL-d₄) δ ppm 0.43 (d, J = 6.85 Hz, 3 H)
2.36-2.53 (m, 1 H) 3.43 (s, 3 H) 3.46-3.58 (m, 1 H) 3.69 (dd, J = 13.57, 3.79 Hz, 1 H) 3.88 (dd, J = 12.30, 4.52 Hz, 1 H) 4.02 (dt, J = 11.63, 4.68 Hz, 1 H)
4.12-4.26 (m, 1 H) 4.60 (s, 2 H) 4.95 (t, J = 4.38 Hz, 2 H) 7.28 (t, J = 8.39 Hz, 2 H) 7.54 (d, J = 6.55 Hz, 1 H) 7.69 (tt, J = 8.50, 6.47 Hz, 1 H) 8.00-8.12 (m, 2 H) 8.39 (dd, J = 6.55, 0.88 Hz, 1 H) 8.47 (dd, J = 8.66, 3.96 Hz, 1 H) 9.18 (s, 1 H)
- 15 ¹H NMR in DMSO-d₆: δ 10.51 (s, 1H), 8.53 (s, 1H), 8.51 (d, J = 4.0, 1H), 8.33 (dd, J = 8.0, 4.0, 1H), 8.20 (t, J = 8.0, 1H), 8.09 (broad doublet, J = 4.0, 2H), 7.65-7.75 (m, 1H), 7.36 (t, J = 8.0, 2H), 7.30 (d, J = 8.0, 1H), 3.84-4.02 (m, 3H), 3.58-3.68 (m, 1H), 3.43-3.53 (m, 1H), 3.28-3.36 (m, 1H), 3.04-3.14 (m, 1H), 1.92-2.18 (m, 4H), 1.56-1.63 (m, 1H), 1.24 (d, J = 8.0, 3H).
- 30 (¹H NMR, DMSO-d₆) δ 10.48 (s, 1H), 8.57 (s, 1H), 8.48 (d, J = 8.0, 1H), 8.33 (dd, J = 8.0, 4.0, 1H), 8.20 (t, J = 8.0, 1H), 7.80 (bs, 2H), 7.71 (quintet, J = 8.0, 2H), 7.43 (d, J = 4.0, 1H), 7.37 (t, J = 8.0, 2H), 3.43-3.50 (m, 2H), 2.88-2.96 (m, 1H), 2.04 (d, J = 8.0, 1H), 1.90 (s, 3H), 1.82 (d, J = 12.0, 1H), 1.64 (q, J = 12.0, 1H), 1.46-1.55 (m, 1H), 1.32 (q, J = 12.0, 1H), 0.85 (d, J = 4.0, 3H).
- 33 ¹H NMR, 400 MHz DMSO-d₆, δ 10.44 (s, 1H), 8.58 (d, J = 4.0, 1H), 8.47 (d, J = 4.0, 1H), 8.34 (dd, J = 8.0, 4.0, 1H), 8.20 (dd, J = 8.0, 8.0, 1H), 8.20 (dd, J = 16.0, 4.0, 1H), 7.67-7.74 (m, 1H), 7.36 (dd, J = 8.0, 8.0, 2H), 7.26 (d, J = 4.0, 1H), 2.85-2.95 (m, 2H), 2.18 (s, 3H), 1.88-1.98 (m, 1H), 1.74-1.84 (m, 1H), 1.48-1.56 (m, 1H), 1.38-1.48 (m, 1H), 1.18-1.28 (m, 1H), 1.02 (d, J = 8.0, 3H)
- 34 ¹H NMR, 400 MHz DMSO-d₆, δ 10.44 (s, 1H), 8.60 (d, J = 8.0, 1H), 8.48 (d, J = 4.0, 1H), 8.34 (dd, J = 8.0, 4.0, 1H), 8.20 (dd, J = 8.0, 8.0, 1H), 8.00 (dd, J = 16.0, 4.0, 1H), 7.67-7.74 (m, 1H), 7.34-7.38 (m, 3H), 3.14 (s, 3H), 3.02-3.12 (m, 2H), 2.18-2.24 (m, 1H), 1.84-1.96 (m, 3H), 1.62-1.72 (m, 1H), 1.38-1.48 (m, 1H), 1.18 (d, J = 4.0, 3H).
- 36 ¹H NMR, 400 MHz DMSO-d₆, δ 10.48 (s, 1H), 8.58 (d, J = 4.0, 1H), 8.50 (d, J = 4.0, 1H), 8.34 (dd, J = 8.0, 4.0, 1H), 8.20 (dd, J = 8.0, 8.0, 1H), 8.20 (dd, J = 16.0, 4.0, 1H), 7.67-7.74 (m, 1H), 7.42 (d, J = 4.0, 1H), 7.36 (dd, J = 8.0, 8.0, 2H), 3.40-3.42 (m, 1H), 3.06-3.20 (m, 1H), 2.92 (s, 3H), 2.06-2.20 (m, 1H), 1.95-2.04 (m, 2H), 1.70-1.80 (m, 1H), 1.56-1.70 (m, 1H), 0.86 (d, J = 8.0, 3H).
- 37 ¹H NMR, 400 MHz DMSO-d₆, δ 10.52 (s, 1H), 8.51-8.54 (m, 2H), 8.34 (dd, J = 8.0, 4.0, 1H), 8.20 (dd, J = 8.0, 8.0, 1H), 8.15 (dd, J = 16.0, 4.0, 1H), 7.67-7.74 (m, 1H), 7.36 (dd, J = 8.0, 8.0, 2H), 7.28 (d, J = 4.0, 1H), 3.77-3.79 (m, 1H), 3.18 (s, 3H), 3.02-3.20 (m, 1H), 1.94-2.40 (m, 4H), 1.57-1.62 (m, 1H), 1.24 (d, J = 8.0, 3H).
- 56 ¹H NMR, (400, DMSO-d₆) δ 10.45 (s, 1H), 8.92 (s, 1H), 8.56 (s, 1H), 8.43 (d, J = 4.0, 1H), 8.40 (dd, J = 8.0, 4.0, 1H), 8.24 (t, J = 8.0, 1H), 8.17-8.24 (m, 2H), 8.17 (s, 1H), 7.70-7.80 (m, 1H), 7.35-7.43 (m, 3H), 4.780-4.82 (m, 1H), 3.96 (t, J = 12.0, 1H), 3.60-3.90 (m, 3H), 3.50 (t, J = 12.0, 1H), 2.25-2.35 (m, 1H), 0.31 (d, J = 8.0, 3H).
- 64 ¹H NMR, (400 MHz, METHANOL-d₄) δ ppm 1.19 (d, J = 6.90 Hz, 3 H) 1.42 (d, J = 12.76 Hz, 1 H) 1.63-1.77 (m, 2 H) 1.84-2.04 (m, 3 H) 2.71-2.91 (m, 2 H) 3.13-3.24 (m, 2 H) 3.40-3.52 (m, 1 H) 3.73 (s, 1 H) 3.76-3.87 (m, 1 H) 3.93 (td, J = 8.68, 5.14 Hz, 1 H) 7.14 (t, J = 8.75 Hz, 2 H) 7.47 (tt, J = 8.44, 6.25 Hz, 1 H) 7.68 (d, J = 5.62 Hz, 1 H) 8.46 (d, J = 5.62 Hz, 1 H) 9.18 (s, 1 H)
- 66 ¹H NMR, (400 MHz, METHANOL-d₄) δ ppm 0.43 (d, J = 6.80 Hz, 3 H) 2.37-2.60 (m, 1 H) 3.60 (t, J = 12.94 Hz, 1 H) 3.83 (dd, J = 13.64, 4.11 Hz, 1 H) 3.95-4.14 (m, 2 H) 4.17-4.32 (m, 1 H) 4.75 (s, 2 H) 5.03 (t, J = 4.01 Hz, 1 H) 7.27 (t, J = 8.46 Hz, 2 H) 7.59 (d, J = 6.85 Hz, 1 H) 7.62-7.75 (m, 1 H) 7.97-8.12 (m, 2 H) 8.40 (dd, J = 6.80, 0.93 Hz, 1 H) 8.47 (dd, J = 8.66, 3.91 Hz, 1 H) 9.12 (d, J = 0.83 Hz, 1 H).
- 74 ¹H NMR, (400 MHz, DMSO-d₆) δ ppm 0.96 (d, J = 6.85 Hz, 3 H) 1.16-1.36 (m, 1 H) 1.46 (d, J = 12.81 Hz, 1 H) 1.50-1.78 (m, 3 H) 2.41 (s, 3 H) 2.89 (t, J = 12.06 Hz, 1 H) 3.07 (s, 3 H) 3.24 (br. s., 2 H) 3.86 (dt, J = 10.20, 5.12 Hz, 2 H) 3.99 (td, J = 9.29, 4.30 Hz, 1 H) 7.17 (d, J = 9.29 Hz, 2 H) 7.22 (d, J = 5.23 Hz, 1 H) 7.86 (br. s., 3 H) 8.08-8.20 (m, 1 H) 8.28 (dd, J = 8.66, 4.06 Hz, 1 H) 8.43 (d, J = 5.18 Hz, 1 H) 8.58 (s, 1 H) 10.38 (s, 1 H).
- 77 ¹H NMR, (400 MHz, DMSO-d₆) δ ppm 1.00 (d, J = 6.80 Hz, 3 H) 1.42-1.81 (m, 3 H) 2.92 (t, J = 12.32 Hz, 1 H) 3.10 (s, 2 H) 3.28 (br. s., 1 H) 3.33-3.44 (m, 1 H) 3.54 (br. s., 1 H) 3.59-3.72 (m, 1 H) 3.89 (dt, J = 10.17, 5.09 Hz, 1 H) 4.02 (td, J = 9.26, 4.23 Hz, 1 H) 7.23 (d, J = 5.23 Hz, 1 H) 7.36 (t, J = 8.31 Hz, 2 H) 7.63-7.77 (m, 1 H) 7.89 (br. s., 2 H) 8.13-8.27 (m, 1 H) 8.33 (dd, J = 8.68, 4.08 Hz, 1 H) 8.46 (d, J = 5.13 Hz, 1 H) 8.60 (s, 1 H) 10.43 (s, 1 H).
- 86 ¹H NMR, (400 MHz, METHANOL-d₄) δ ppm 1.13 (d, J = 6.80 Hz, 3 H) 1.46-1.86 (m, 3 H) 1.88-2.06 (m, 2 H) 2.72-2.93 (m, 2 H) 3.08-3.21 (m, 1 H) 3.34-3.42 (m, 1 H) 3.70 (s, 1 H) 3.76-3.88 (m, 1 H) 3.93 (td, J = 8.69, 5.01 Hz, 1 H) 7.22 (t, J = 8.27 Hz, 2 H) 7.55-7.77 (m, 2 H) 8.04 (t, J = 8.71 Hz, 1 H) 8.42 (dd, J = 8.66, 3.96 Hz, 1 H) 8.55 (d, J = 5.62 Hz, 1 H) 9.03 (s, 1 H)
- 90 ¹H NMR, (400 MHz, <cd3od>) δ ppm 8.95 (s, 1 H), 8.50 (d, J = 5.48 Hz, 1 H), 8.39 (dd, J = 8.61, 3.91 Hz, 1 H), 8.00 (t, J = 8.80 Hz, 1 H), 7.60 (d, J = 5.48 Hz, 1 H), 7.05 (d, J = 9.39 Hz, 2 H), 3.72 (quin, J = 7.24 Hz, 2 H), 3.57 (br. s., 1 H), 3.08 (t, J = 11.74 Hz, 1 H), 2.47 (s, 3 H), 1.84-2.00 (m, 2 H), 1.72 (d, J = 5.48 Hz, 1 H), 1.54-1.65 (m, 2 H), 1.28 (t, J = 7.04 Hz, 3 H), 1.06 (d, J = 6.65 Hz, 3 H)
- 91 ¹H NMR, (400 MHz, <cd3od>) δ ppm 8.96 (s, 1 H), 8.50 (d, J = 5.48 Hz, 1 H), 8.41 (dd, J = 8.61, 3.91 Hz, 1 H), 8.03 (t, J = 8.80 Hz, 1 H), 7.58-7.68 (m, 2 H), 7.21 (t, J = 8.41 Hz, 2 H), 3.72 (quin, J = 7.24 Hz, 2 H), 3.57 (br. s., 1

TABLE 3-continued

Ex # ¹H-NMR data

- H), 3.03-3.15 (m, 1 H), 1.84-2.00 (m, 2 H), 1.72 (br. s., 1 H),
1.51-1.66 (m, 2 H), 1.28 (t, J = 7.04 Hz, 3 H), 1.06 (d, J = 6.65 Hz, 3 H)
- 92 ¹H NMR, (300 MHz, <cd3od>) δ ppm 8.93-9.05 (m, 1 H), 8.46-8.57 (m, 1 H),
8.35-8.46 (m, 1 H), 7.97-8.09 (m, 1 H), 7.55-7.73 (m, 2 H), 7.15-7.30 (m, 2 H),
3.58 (s, 3 H), 3.42-3.51 (m, 1 H), 3.00-3.18 (m, 1 H), 1.83-2.00 (m, 2 H),
1.45-1.81 (m, 3 H), 1.07 (d, J = 6.74 Hz, 3 H)
- 94 ¹H NMR, (400 MHz, METHANOL-d4) δ ppm 8.82 (s, 1 H), 8.48 (d, J = 5.48 Hz,
1 H), 8.39 (dd, J = 8.61, 3.91 Hz, 1 H), 8.01 (t, J = 8.61 Hz, 1 H), 7.54 (d, J = 5.09 Hz,
1 H), 7.19 (d, J = 9.39 Hz, 2 H), 3.58 (s, 3 H), 3.46 (s, 1 H), 3.00-3.21 (m, 5 H),
2.23-2.40 (m, 4 H), 1.85-1.95 (m, 2 H), 1.75 (br. s., 1 H), 1.50-1.67 (m,
2 H), 1.08 (d, J = 6.65 Hz, 3 H)
- 95 ¹H NMR, (400 MHz, METHANOL-d4) δ ppm 8.92 (s, 1 H), 8.47 (d, J = 5.48 Hz,
1 H), 8.36 (dd, J = 8.63, 3.94 Hz, 1 H), 7.98 (t, J = 8.71 Hz, 1 H), 7.56 (d, J = 5.48 Hz,
1 H), 7.08 (d, J = 9.44 Hz, 2 H), 3.56 (s, 3 H), 3.44 (s, 1 H), 2.95-3.10 (m, 2 H),
1.81-1.97 (m, 2 H), 1.71 (d, J = 5.97 Hz, 1 H), 1.46-1.66 (m, 2 H) 1.30 (d,
J = 6.94 Hz, 6 H), 1.05 (d, J = 6.80 Hz, 3 H)
- 96 ¹H NMR, (400 MHz, METHANOL-d4) δ ppm 8.92 (s, 1 H), 8.49 (d, J = 5.53 Hz,
1 H), 8.39 (dd, J = 8.68, 3.94 Hz, 1 H), 8.01 (t, J = 8.73 Hz, 1 H), 7.60 (d, J = 5.58 Hz,
1 H), 7.30 (d, J = 9.24 Hz, 2 H), 3.90-4.01 (m, 2 H), 3.84 (td, J = 11.91, 1.61 Hz,
2 H), 3.44 (s, 1 H), 2.99-3.15 (m, 1 H), 2.30 (td, J = 13.14, 5.45 Hz, 1 H),
2.20 (td, J = 13.14, 5.40 Hz, 1 H), 1.81-1.98 (m, 4 H), 1.66-1.79 (m, 1 H),
1.44-1.65 (m, 2 H), 1.05 (d, J = 6.85 Hz, 3 H)
- 97 ¹H NMR, (400 MHz, METHANOL-d4) δ ppm 8.93 (s, 1 H), 8.46 (d, J = 5.43 Hz,
1 H), 8.37 (dd, J = 8.63, 3.94 Hz, 1 H), 7.99 (t, J = 8.73 Hz, 1 H), 7.55 (d, J = 5.48 Hz,
1 H), 7.29 (d, J = 9.68 Hz, 2 H), 3.56 (s, 3 H), 3.42-3.45 (m, 1 H),
3.05 (ddd, J = 15.75, 11.59, 3.91 Hz, 1 H), 1.80-1.98 (m, 2 H), 1.66-1.78 (m, 1 H),
1.45-1.65 (m, 8 H), 1.05 (d, J = 6.85 Hz, 3 H)
- 98 ¹H NMR, (300 MHz, <cd3od>) δ ppm 8.98 (s, 1 H), 8.49 (d, J = 5.57 Hz, 1 H),
8.40 (dd, J = 8.64, 3.96 Hz, 1 H), 8.02 (t, J = 8.64 Hz, 1 H), 7.60 (d, J = 5.57 Hz, 1 H),
7.35 (d, J = 9.67 Hz, 2 H), 3.58 (s, 3 H), 3.46 (s, 1 H), 3.00-3.15 (m, 1 H),
2.35-2.62 (m, 4 H), 2.02-2.21 (m, 1 H), 1.78-1.99 (m, 3 H), 1.45-1.68 (m, 1 H),
1.07 (d, J = 6.74 Hz, 3 H)
- 99 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.97 (d, J = 6.80 Hz, 3 H) 1.36-1.52 (m,
1 H) 1.63-1.73 (m, 1 H) 1.75-1.91 (m, 7 H) 2.83-3.02 (m, 1 H) 3.25 (s, 2 H)
3.51-3.54 (m, 3 H) 3.56-3.63 (m, 1 H) 4.06 (dt, J = 11.21, 2.95 Hz, 4 H)
7.12 (d, J = 9.19 Hz, 2 H) 7.38 (d, J = 5.33 Hz, 1 H) 7.97 (t, J = 8.73 Hz, 1 H)
8.29-8.41 (m, 2 H) 8.95 (s, 1 H)
- 100 ¹H NMR, (300 MHz, <cd3od>) δ ppm 8.97 (s, 1 H), 8.50 (d, J = 5.57 Hz, 1 H),
8.40 (dd, J = 8.64, 3.96 Hz, 1 H), 8.02 (t, J = 8.79 Hz, 1 H), 7.62 (d, J = 5.57 Hz, 1 H),
7.19 (d, J = 9.08 Hz, 2 H), 4.61 (s, 2 H), 3.63 (q, J = 7.03 Hz, 2 H), 3.58 (s, 3 H),
3.43-3.50 (m, 1 H), 3.01-3.16 (m, 1 H), 1.83-1.99 (m, 2 H),
1.68-1.82 (m, 1 H), 1.48-1.67 (m, 2 H), 1.28 (t, J = 7.03 Hz, 3 H), 1.07 (d, J = 6.74 Hz, 3 H)
- 101 ¹H NMR, (300 MHz, <cd3od>) δ ppm 8.97-9.13 (m, 1 H), 8.48-8.60 (m, 1 H),
8.33-8.45 (m, 1 H), 7.94-8.08 (m, 1 H), 7.60-7.72 (m, 1 H), 6.71-6.87 (m, 2 H),
4.92-5.04 (m, 1 H), 4.65-4.79 (m, 1 H), 3.47-3.53 (m, 1 H) 3.60 (s, 3 H),
3.02-3.20 (m, 1 H), 1.86-2.06 (m, 2 H), 1.71-1.85 (m, 1 H), 1.48-1.70 (m, 2 H),
1.39 (d, J = 6.15 Hz, 6 H), 1.10 (d, J = 6.74 Hz, 3 H)
- 104 ¹H NMR, (400 MHz, <cd3od>) δ ppm 9.26 (d, J = 5.1 Hz, 1 H), 8.74 (s, 1 H),
8.55 (d, J = 5.5 Hz, 1 H), 8.21 (d, J = 5.1 Hz, 1 H), 7.51-7.70 (m, 2 H),
7.10-7.27 (m, 2 H), 3.80 (br. s., 1 H), 3.60-3.74 (m, 1 H), 3.20-3.28 (m, 1 H),
3.16 (s, 3 H), 2.32 (quin, J = 12.7 Hz, 2 H), 2.04-2.24 (m, 2 H), 1.65-1.84 (m, 1 H),
1.37 (d, J = 7.0 Hz, 3 H)
- 107 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.36 (d, J = 7.04 Hz, 3 H) 1.75 (dtd,
J = 12.98, 8.61, 8.61, 3.99 Hz, 3 H) 2.01-2.23 (m, 4 H) 2.24-2.39 (m, 2 H)
3.16 (s, 3 H) 3.19-3.28 (m, 1 H) 3.54-3.73 (m, 3 H) 3.79 (br. s., 1 H)
3.89-4.04 (m, 2 H) 4.64-4.75 (m, 1 H) 6.83 (d, J = 9.98 Hz, 2 H) 7.61 (d, J = 5.53 Hz,
1 H) 7.96 (t, J = 8.68 Hz, 1 H) 8.34 (dd, J = 8.61, 3.91 Hz, 1 H) 8.53 (d, J = 5.48 Hz,
1 H) 8.72 (s, 1 H)
- 108 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.36 (d, J = 6.99 Hz, 3 H) 1.75 (d, J = 13.50 Hz,
1 H) 1.80-1.91 (m, 1 H) 2.03-2.23 (m, 3 H) 2.23-2.37 (m, 2 H)
2.38-2.62 (m, 4 H) 3.17 (s, 3 H) 3.61-3.73 (m, 1 H) 3.78 (br. s., 1 H) 7.34 (d, J = 9.59 Hz,
2 H) 7.59 (d, J = 5.48 Hz, 1 H) 8.01 (t, J = 8.66 Hz, 1 H) 8.38 (dd, J = 8.61,
3.91 Hz, 1 H) 8.52 (d, J = 5.43 Hz, 1 H) 8.73 (s, 1 H)
- 109 ¹H NMR (400 MHz, <dms_o-d₆>) δ ppm 1.15 (dd, J = 6.46, 1.76 Hz, 7 H) 1.22 (d,
J = 6.65 Hz, 4 H) 1.60 (d, J = 12.91 Hz, 1 H) 1.86-2.06 (m, 2 H) 2.08-2.35 (m, 3 H)
4.10 (dd, J = 13.89, 6.85 Hz, 4 H) 7.26-7.47 (m, 4 H) 7.77 (dd, J = 8.61, 2.35 Hz,
1 H) 7.88 (ddd, J = 8.31, 4.60, 2.35 Hz, 1 H) 8.15 (d, J = 3.13 Hz, 3 H)
8.35 (d, J = 7.83 Hz, 1 H) 8.40-8.54 (m, 2 H) 8.76 (s, 1 H) 10.36 (s, 1 H)
- 110 ¹H NMR (400 MHz, <dms_o-d₆>) δ ppm 1.23 (d, J = 6.65 Hz, 5 H) 1.59 (br. s., 2 H)
1.78-2.37 (m, 14 H) 3.00-3.13 (m, 2 H) 3.51-3.73 (m, 6 H) 3.77 (br. s., 2 H)
3.89 (dd, J = 11.35, 5.09 Hz, 4 H) 7.29 (d, J = 5.48 Hz, 2 H) 7.39-7.55 (m, 4 H)
8.04-8.24 (m, 7 H) 8.32 (dd, J = 8.61, 4.30 Hz, 2 H) 8.44-8.57 (m, 3 H)
10.48-10.57 (m, 2 H)
- 111 ¹H NMR (400 MHz, <dms_o-d₆>) δ ppm 0.95 (d, J = 6.65 Hz, 2 H) 1.15 (d, J = 6.26 Hz,
4 H) 1.48 (br. s., 3 H) 2.75-2.86 (m, 1 H) 3.02 (br. s., 1 H) 3.64-3.81 (m,
2 H) 4.00-4.18 (m, 1 H) 7.24-7.50 (m, 2 H) 7.78 (dd, J = 8.61, 2.35 Hz, 1 H)
7.83-8.01 (m, 3 H) 8.35 (d, J = 7.43 Hz, 1 H) 8.39-8.49 (m, 1 H) 8.88 (s, 1 H)
10.34 (s, 1 H)

TABLE 3-continued

Ex # ¹H-NMR data

- 112 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.36 (d, J = 7.04 Hz, 3 H) 1.71-1.83 (m, 1 H) 2.05-2.25 (m, 2 H) 2.25-2.41 (m, 2 H) 3.13-3.21 (m, 3 H) 3.23-3.29 (m, 1 H) 3.63-3.73 (m, 1 H) 3.80-3.86 (m, 1 H) 3.89 (s, 3 H) 6.80 (d, J = 9.78 Hz, 2 H) 7.59-7.71 (m, 1 H) 7.97 (s, 1 H) 8.31-8.40 (m, 1 H) 8.51-8.59 (m, 1 H) 8.83 (s, 1 H)
- 114 ¹H NMR (400 MHz, <cd3od>) δ ppm 1.32 (d, J = 6.65 Hz, 3 H) 1.69-1.84 (m, 1 H) 2.04-2.19 (m, 2 H) 2.20-2.45 (m, 2 H) 3.13-3.22 (m, 3 H) 3.24-3.29 (m, 1 H) 3.57-3.73 (m, 1 H) 3.89 (br. s., 1 H) 7.05-7.19 (m, 3 H) 7.47-7.59 (m, 1 H) 7.76 (d, J = 5.48 Hz, 1 H) 8.45-8.59 (m, 1 H) 9.16 (s, 1 H)
- 115 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.26 (d, J = 6.65 Hz, 3 H) 1.54-1.67 (m, 1 H) 1.89-2.02 (m, 2 H) 2.08 (q, J = 12.52 Hz, 1 H) 2.94-3.15 (m, 1 H) 3.16-3.24 (m, 3 H) 3.65 (br. s., 1 H) 3.81 (br. s., 1 H) 7.10-7.34 (m, 3 H) 7.44-7.68 (m, 3 H) 8.15 (d, J = 3.91 Hz, 3 H) 8.46 (d, J = 5.48 Hz, 1 H) 8.62 (s, 1 H) 9.59 (s, 1 H)
- 116 ¹H NMR (400 MHz, <dms_o-d₆>) δ ppm 1.03-1.31 (m, 6 H) 1.57 (d, J = 12.91 Hz, 1 H) 1.81-2.08 (m, 3 H) 2.09-2.20 (m, 1 H) 2.92-3.11 (m, 1 H) 3.17 (s, 3 H) 3.53 (q, J = 7.04 Hz, 2 H) 3.60 (d, J = 5.87 Hz, 1 H) 3.79 (br. s., 1 H) 4.48-4.67 (m, 2 H) 7.15-7.37 (m, 3 H) 8.01-8.22 (m, 4 H) 8.31 (dd, J = 8.61, 3.91 Hz, 1 H) 8.44-8.59 (m, 2 H) 10.32-10.60 (m, 1 H)
- 117 ¹H NMR, (400 MHz, <cd3od>) δ ppm 9.03 (s, 1 H), 8.45 (d, J = 5.1 Hz, 1 H), 8.40 (dd, J = 8.6, 3.9 Hz, 1 H), 8.02 (t, J = 8.6 Hz, 1 H), 7.53 (d, J = 5.5 Hz, 1 H), 7.35 (d, J = 10.2 Hz, 2 H), 3.91-4.03 (m, 2 H), 3.79-3.90 (m, 2 H), 3.53 (s, 3 H), 3.05-3.21 (m, 2 H), 2.93 (t, J = 10.2 Hz, 1 H), 2.09-2.26 (m, 3 H), 1.93 (dd, J = 13.5, 2.5 Hz, 1 H), 1.78 (q, J = 12.1 Hz, 1 H), 1.67 (d, J = 13.7 Hz, 2 H), 1.21-1.51 (m, 2 H), 1.07 (d, J = 6.3 Hz, 3 H)
- 118 ¹H NMR, (400 MHz, <cd3od>) δ ppm 8.95 (s, 1 H), 8.51 (d, J = 5.5 Hz, 1 H), 8.40 (dd, J = 8.6, 3.9 Hz, 1 H), 8.02 (t, J = 8.6 Hz, 1 H), 7.63 (d, J = 5.5 Hz, 1 H), 7.35 (d, J = 9.8 Hz, 2 H), 4.26 (dt, J = 10.7, 5.0 Hz, 1 H), 4.00-4.12 (m, 1 H), 3.91-4.00 (m, 2 H), 3.79-3.90 (m, 2 H), 3.69 (s, 1 H), 3.55 (t, J = 5.3 Hz, 2 H), 3.35-3.46 (m, 1 H), 3.03-3.20 (m, 4 H), 2.16 (td, J = 13.0, 4.9 Hz, 2 H), 1.88-2.01 (m, 2 H), 1.78 (d, J = 6.3 Hz, 1 H), 1.49-1.72 (m, 4 H), 1.11 (d, J = 6.7 Hz, 3 H)
- 119 ¹H NMR, (400 MHz, <cd3od>) δ ppm 8.76 (s, 1 H), 8.52 (d, J = 5.5 Hz, 1 H), 8.34 (dd, J = 8.6, 3.9 Hz, 1 H), 7.97 (t, J = 8.8 Hz, 1 H), 7.61 (d, J = 5.5 Hz, 1 H), 6.82 (d, J = 10.2 Hz, 2 H), 4.21 (dd, J = 5.1, 3.5 Hz, 2 H), 3.73-3.87 (m, 3 H), 3.63-3.72 (m, 1 H), 3.43 (s, 3 H), 3.19-3.28 (m, 1 H), 3.17 (s, 3 H), 2.32 (qd, J = 12.7, 6.1 Hz, 2 H), 2.05-2.24 (m, 2 H), 1.75 (d, J = 13.3 Hz, 1 H), 1.36 (d, J = 7.0 Hz, 3 H)
- 120 ¹H NMR, (300 MHz, <cd3od>) δ ppm 8.77 (s, 1 H), 8.62 (s, 1 H), 8.54 (d, J = 5.5 Hz, 1 H), 7.55-7.69 (m, 2 H), 7.23 (t, J = 8.6 Hz, 2 H), 3.81 (br. s., 1 H), 3.67-3.78 (m, 1 H), 3.17 (s, 3 H), 2.20-2.46 (m, 3 H), 2.08-2.19 (m, 1 H), 1.78 (d, J = 13.3 Hz, 1 H), 1.40 (d, J = 7.0 Hz, 3 H)
- 121 ¹H NMR, (400 MHz, <cd3od>) δ ppm 8.72 (s, 1 H), 8.53 (d, J = 5.5 Hz, 1 H), 8.38 (dd, J = 8.6, 3.9 Hz, 1 H), 8.00 (t, J = 8.6 Hz, 1 H), 7.62 (d, J = 5.5 Hz, 1 H), 7.34 (d, J = 9.8 Hz, 2 H), 3.82-4.01 (m, 4 H), 3.79 (br. s., 1 H), 3.63-3.73 (m, 1 H), 3.20-3.28 (m, 1 H), 3.17 (s, 3 H), 2.31 (qd, J = 12.7, 8.4 Hz, 2 H), 2.06-2.24 (m, 4 H), 1.75 (d, J = 13.3 Hz, 1 H), 1.65 (d, J = 13.3 Hz, 2 H), 1.36 (d, J = 7.0 Hz, 3 H)
- 122 ¹H NMR, (400 MHz, <cd3od>) δ ppm 8.77 (s, 1 H), 8.54 (d, J = 5.5 Hz, 1 H), 8.36 (dd, J = 8.8, 4.1 Hz, 1 H), 7.99 (t, J = 8.6 Hz, 1 H), 7.63 (d, J = 5.5 Hz, 1 H), 7.03 (d, J = 9.4 Hz, 2 H), 3.80 (br. s., 1 H), 3.62-3.73 (m, 1 H), 3.20-3.29 (m, 1 H), 3.17 (s, 3 H), 2.46 (s, 3 H), 2.25-2.40 (m, 2 H), 2.06-2.24 (m, 2 H), 1.68-1.82 (m, 1 H), 1.36 (d, J = 7.0 Hz, 3 H)
- 123 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.90 (d, J = 6.65 Hz, 3 H) 1.72-1.81 (m, 2 H) 1.88-2.07 (m, 2 H) 2.07-2.21 (m, 1 H) 3.24-3.28 (m, 1 H) 3.53-3.63 (m, 1 H) 4.87-4.97 (m, 1 H) 5.59-5.69 (m, 1 H) 6.94-7.08 (m, 2 H) 7.23 (s, 2 H) 7.58-7.70 (m, 1 H) 7.72-7.84 (m, 2 H) 8.01-8.09 (m, 1 H) 8.15-8.22 (m, 1 H) 8.39-8.48 (m, 1 H) 8.55-8.62 (m, 1 H) 9.08 (s, 1 H)
- 124 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.08 (d, J = 6.26 Hz, 3 H) 1.43 (q, J = 12.52 Hz, 1 H) 1.59-1.75 (m, 1 H) 1.81 (q, J = 12.39 Hz, 1 H) 1.94 (dd, J = 13.30, 2.74 Hz, 1 H) 2.15-2.30 (m, 1 H) 2.96 (t, J = 9.98 Hz, 1 H) 3.08-3.22 (m, 2 H) 3.44 (s, 3 H) 3.53 (s, 3 H) 3.73-3.83 (m, 2 H) 4.11-4.28 (m, 2 H) 6.83 (d, J = 10.17 Hz, 2 H) 7.69 (d, J = 5.48 Hz, 1 H) 8.00 (t, J = 8.80 Hz, 1 H) 8.38 (dd, J = 8.61, 3.52 Hz, 1 H) 8.51 (d, J = 5.48 Hz, 1 H) 9.20 (s, 1 H)
- 125 ¹H NMR (400 MHz, <dms_o-d₆>) δ ppm 1.03 (d, J = 6.65 Hz, 3 H) 1.33 (q, J = 12.52 Hz, 1 H) 1.45-1.68 (m, 2 H) 1.70-1.88 (m, 2 H) 2.86-3.04 (m, 1 H) 3.04-3.15 (m, 3 H) 3.22-3.43 (m, 3 H) 3.58 (br. s., 2 H) 3.61-3.72 (m, 2 H) 3.89 (dt, J = 10.08, 4.94 Hz, 6 H) 4.02 (td, J = 9.20, 4.30 Hz, 5 H) 7.15-7.34 (m, 3 H) 7.45-7.69 (m, 3 H) 7.95 (br. s., 3 H) 8.42 (d, J = 5.48 Hz, 1 H) 8.78 (s, 1 H) 9.48 (s, 1 H)
- 126 ¹H NMR (400 MHz, <dms_o-d₆>) δ ppm 0.93 (d, J = 6.26 Hz, 7 H) 1.15 (d, J = 6.65 Hz, 14 H) 1.32-1.46 (m, 2 H) 1.49-1.70 (m, 5 H) 1.72-1.85 (m, 2 H) 2.09 (d, J = 10.17 Hz, 2 H) 2.88 (t, J = 9.98 Hz, 2 H) 2.98-3.14 (m, 5 H) 3.30 (s, 7 H) 4.00-4.17 (m, 2 H) 7.31-7.41 (m, 4 H) 7.45 (d, J = 5.09 Hz, 2 H) 7.79 (dd, J = 9.00, 1.96 Hz, 2 H) 7.85-8.00 (m, 8 H) 8.33 (d, J = 7.83 Hz, 2 H) 8.46 (d, J = 5.09 Hz, 4 H) 8.89 (s, 2 H) 10.36 (s, 2 H)
- 127 ¹H NMR (400 MHz, <dms_o-d₆>) δ ppm 0.95 (d, J = 6.26 Hz, 3 H) 1.19-1.39 (m, 1 H) 1.47-1.67 (m, 2 H) 1.74 (d, J = 11.35 Hz, 1 H) 1.87 (t, J = 11.93 Hz, 2 H)

TABLE 3-continued

Ex # ¹H-NMR data

- 2.00 (br. s., 1 H) 2.11-2.37 (m, 2 H) 2.82-3.13 (m, 3 H) 3.37 (s, 3 H) 3.69 (t, J = 11.35 Hz, 2 H) 3.89 (dd, J = 11.35, 5.09 Hz, 2 H) 7.37 (d, J = 5.48 Hz, 1 H) 7.44 (d, J = 9.39 Hz, 2 H) 7.91 (br. s., 3 H) 8.15-8.25 (m, 1 H) 8.33 (dd, J = 8.80, 4.11 Hz, 1 H) 8.44 (d, J = 5.09 Hz, 1 H) 8.62-8.70 (m, 1 H) 10.39 (s, 1 H)
- 128 ¹H NMR (400 MHz, <dms_o-d₆>) δ ppm 0.96 (d, J = 6.65 Hz, 3 H) 1.16 (d, J = 6.65 Hz, 6 H) 1.29-1.64 (m, 3 H) 1.70-1.88 (m, 2 H) 2.97-3.14 (m, 4 H) 3.30-3.44 (m, 2 H) 3.53 (br. s., 1 H) 3.57-3.69 (m, 1 H) 3.88 (dt, J = 10.17, 5.09 Hz, 1 H) 4.00 (td, J = 9.19, 4.30 Hz, 1 H) 4.11 (dq, J = 13.60, 6.68 Hz, 1 H) 7.25-7.47 (m, 3 H) 7.78 (dd, J = 8.80, 2.15 Hz, 1 H) 7.84-8.01 (m, 4 H) 8.28-8.50 (m, 3 H) 8.90 (s, 1 H) 10.35 (s, 1 H)
- 129 ¹H NMR (400 MHz, <dms_o-d₆>) δ ppm 0.98 (d, J = 7.04 Hz, 3 H) 1.30 (q, J = 12.52 Hz, 1 H) 1.47 (d, J = 12.52 Hz, 1 H) 1.53-1.79 (m, 3 H) 1.87 (t, J = 11.93 Hz, 2 H) 2.10-2.38 (m, 2 H) 2.91 (t, J = 12.33 Hz, 1 H) 3.09 (s, 3 H) 3.26 (br. s., 1 H) 3.32-3.44 (m, 1 H) 3.52 (br. s., 1 H) 3.58-3.77 (m, 3 H) 3.79-3.95 (m, 3 H) 4.01 (td, J = 9.19, 4.30 Hz, 1 H) 7.23 (d, J = 5.09 Hz, 1 H) 7.38-7.53 (m, 2 H) 7.87 (br. s., 3 H) 8.19 (t, J = 8.80 Hz, 1 H) 8.32 (dd, J = 8.61, 3.91 Hz, 1 H) 8.45 (d, J = 5.09 Hz, 1 H) 8.53-8.63 (m, 1 H) 10.37-10.49 (m, 1 H)
- 130 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.13 (d, J = 6.65 Hz, 3 H) 1.46-1.72 (m, 2 H) 1.79 (d, J = 5.87 Hz, 1 H) 1.88-2.05 (m, 2 H) 3.10 (s, 4 H) 3.35-3.47 (m, 4 H) 3.50-3.63 (m, 2 H) 3.71 (s, 1 H) 3.76-3.81 (m, 2 H) 4.05 (ddd, J = 10.86, 6.95, 4.11 Hz, 1 H) 4.17-4.33 (m, 3 H) 6.83 (d, J = 10.17 Hz, 2 H) 7.71 (d, J = 5.48 Hz, 1 H) 7.99 (t, J = 8.61 Hz, 1 H) 8.37 (dd, J = 8.80, 3.72 Hz, 1 H) 8.54 (d, J = 5.87 Hz, 1 H) 9.09 (s, 1 H)
- 131 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.53 (d, J = 7.04 Hz, 3 H) 2.57-2.82 (m, 1 H) 3.56-3.72 (m, 1 H) 3.83 (dd, J = 13.30, 3.91 Hz, 1 H) 4.04 (d, J = 10.17 Hz, 1 H) 4.12-4.36 (m, 2 H) 5.20 (t, J = 3.91 Hz, 1 H) 7.16-7.36 (m, 2 H) 7.50-7.71 (m, 2 H) 7.89 (d, J = 0.78 Hz, 1 H) 8.12 (d, J = 0.78 Hz, 1 H) 8.34-8.48 (m, 1 H) 8.73 (s, 1 H) 9.08 (s, 1 H)
- 132 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.34 (d, J = 6.65 Hz, 3 H) 2.22-2.53 (m, 1 H) 3.36 (s, 3 H) 3.39-3.56 (m, 1 H) 3.60-3.76 (m, 3 H) 3.85 (d, J = 6.65 Hz, 1 H) 3.93-4.03 (m, 1 H) 4.10-4.25 (m, 3 H) 4.94 (br. s., 1 H) 6.80 (d, J = 10.17 Hz, 2 H) 7.46 (d, J = 7.04 Hz, 1 H) 7.81 (s, 1 H) 7.92 (t, J = 8.61 Hz, 1 H) 8.00 (d, J = 0.78 Hz, 1 H) 8.26-8.45 (m, 2 H) 9.08 (s, 1 H)
- 133 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.19 (d, J = 6.65 Hz, 3 H) 1.56-1.77 (m, 2 H) 1.84-2.05 (m, 3 H) 3.11 (s, 3 H) 3.17-3.27 (m, 1 H) 3.47 (ddd, J = 11.64, 4.40, 2.74 Hz, 1 H) 3.53-3.63 (m, 2 H) 3.73 (s, 1 H) 4.06 (ddd, J = 10.76, 6.46, 4.70 Hz, 1 H) 4.23-4.37 (m, 1 H) 7.24 (t, J = 8.61 Hz, 2 H) 7.62 (tt, J = 8.61, 6.26 Hz, 1 H) 7.73 (d, J = 5.87 Hz, 1 H) 8.55 (d, J = 5.48 Hz, 1 H) 8.66 (s, 1 H) 9.02 (s, 1 H)
- 134 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.15 (d, J = 6.26 Hz, 3 H) 1.43-1.58 (m, 1 H) 1.69-1.92 (m, 2 H) 1.93-2.08 (m, 1 H) 2.20-2.35 (m, 1 H) 3.00 (t, J = 10.17 Hz, 1 H) 3.17-3.29 (m, 2 H) 3.54 (s, 3 H) 7.25 (t, J = 8.61 Hz, 2 H) 7.62 (tt, J = 8.56, 6.31 Hz, 1 H) 7.76 (d, J = 5.87 Hz, 1 H) 8.55 (d, J = 5.48 Hz, 1 H) 8.67 (s, 1 H) 9.11 (s, 1 H)
- 135 ¹H NMR (400 MHz, <dms_o-d₆>) δ ppm 1.03 (d, J = 6.65 Hz, 3 H) 1.33 (q, J = 12.52 Hz, 1 H) 1.45-1.68 (m, 2 H) 1.70-1.88 (m, 2 H) 2.86-3.04 (m, 1 H) 3.04-3.15 (m, 3 H) 3.22-3.43 (m, 3 H) 3.58 (br. s., 2 H) 3.61-3.72 (m, 2 H) 3.89 (dt, J = 10.08, 4.94 Hz, 6 H) 4.02 (td, J = 9.20, 4.30 Hz, 5 H) 7.15-7.34 (m, 3 H) 7.45-7.69 (m, 3 H) 7.95 (br. s., 3 H) 8.42 (d, J = 5.48 Hz, 1 H) 8.78 (s, 1 H) 9.48 (s, 1 H)
- 136 ¹H NMR (400 MHz, <dms_o>) d ppm 0.83-1.01 (m, 3 H) 1.09-1.22 (m, 3 H) 1.30 (q, J = 12.52 Hz, 1 H) 1.45-1.68 (m, 2 H) 1.74 (d, J = 10.96 Hz, 1 H) 2.02 (d, J = 9.78 Hz, 1 H) 2.82-2.95 (m, 1 H) 2.95-3.12 (m, 2 H) 3.37 (s, 3 H) 3.54 (q, J = 7.04 Hz, 2 H) 4.57 (s, 2 H) 7.09-7.32 (m, 2 H) 7.43 (d, J = 5.09 Hz, 1 H) 7.99 (br. s., 3 H) 8.10-8.23 (m, 1 H) 8.32 (dd, J = 8.61, 4.30 Hz, 1 H) 8.47 (d, J = 5.09 Hz, 1 H) 8.71 (s, 1 H) 10.41 (s, 1 H)
- 137 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.09 (d, J = 6.46 Hz, 3 H) 1.36-1.50 (m, 1 H) 1.63-1.87 (m, 4 H) 1.95 (dd, J = 13.55, 2.59 Hz, 1 H) 2.05-2.14 (m, 2 H) 2.17-2.29 (m, 1 H) 2.91-3.01 (m, 1 H) 3.10-3.23 (m, 2 H) 3.54 (s, 3 H) 3.64 (ddd, J = 11.73, 8.74, 3.01 Hz, 2 H) 3.93-4.02 (m, 2 H) 4.66-4.75 (m, 1 H) 6.85 (d, J = 10.12 Hz, 2 H) 7.64 (d, J = 5.53 Hz, 1 H) 8.00 (t, J = 8.71 Hz, 1 H) 8.38 (dd, J = 8.63, 3.89 Hz, 1 H) 8.50 (d, J = 5.48 Hz, 1 H) 9.12 (s, 1 H)
- 138 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.09 (d, J = 6.50 Hz, 3 H) 1.36-1.51 (m, 1 H) 1.64-2.00 (m, 4 H) 2.04-2.27 (m, 2 H) 2.38-2.50 (m, 2 H) 2.51-2.63 (m, 2 H) 2.85-3.03 (m, 2 H) 3.11-3.24 (m, 2 H) 3.54 (s, 3 H) 7.36 (d, J = 9.49 Hz, 2 H) 7.64 (d, J = 5.58 Hz, 1 H) 8.04 (t, J = 8.71 Hz, 1 H) 8.42 (dd, J = 8.66, 3.91 Hz, 1 H) 8.50 (d, J = 5.58 Hz, 1 H) 9.15 (s, 1 H)
- 139 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.12 (d, J = 6.85 Hz, 3 H) 1.52-1.71 (m, 2 H) 1.72-1.98 (m, 4 H) 2.05-2.18 (m, 1 H) 2.35-2.60 (m, 4 H) 3.06-3.17 (m, 4 H) 3.55 (t, J = 5.31 Hz, 2 H) 3.68 (s, 1 H) 3.97-4.11 (m, 1 H) 4.19-4.34 (m, 1 H) 7.35 (d, J = 9.44 Hz, 2 H) 7.58 (br. s., 1 H) 8.02 (t, J = 8.71 Hz, 1 H) 8.40 (dd, J = 8.73, 3.94 Hz, 1 H) 8.49 (d, J = 5.38 Hz, 1 H) 8.91 (br. s., 1 H)
- 140 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.14 (d, J = 6.94 Hz, 3 H) 1.55-1.83 (m, 5 H) 1.87-2.01 (m, 2 H) 2.03-2.18 (m, 2 H) 3.12 (s, 3 H) 3.14 (dd, J = 3.28, 1.66 Hz, 1 H) 3.52-3.73 (m, 5 H) 3.91-4.11 (m, 3 H) 4.22-4.35 (m, 1 H) 4.72 (tt, J = 8.08, 4.13 Hz, 1 H) 6.86 (d, J = 10.08 Hz, 2 H) 7.64 (d, J = 5.53 Hz, 1 H) 7.99 (t, J = 8.71 Hz, 1 H) 8.37 (dd, J = 8.63, 3.94 Hz, 1 H) 8.52 (d, J = 5.53 Hz, 1 H) 8.95 (s, 1 H)

TABLE 3-continued

Ex # ¹H-NMR data

- 141 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.08 (d, J = 6.50 Hz, 3 H) 1.34-1.49 (m, 7 H) 1.63-1.76 (m, 1 H) 1.81-1.99 (m, 2 H) 2.18-2.28 (m, 1 H) 2.99 (t, J = 10.00 Hz, 1 H) 3.12-3.25 (m, 2 H) 3.54 (s, 3 H) 4.72 (dt, J = 12.07, 6.02 Hz, 1 H) 6.76 (d, J = 10.22 Hz, 2 H) 7.75 (d, J = 5.62 Hz, 1 H) 7.99 (t, J = 8.73 Hz, 1 H) 8.38 (dd, J = 8.63, 3.89 Hz, 1 H) 8.53 (d, J = 5.48 Hz, 1 H) 9.26 (s, 1 H)
- 142 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.12 (d, J = 6.85 Hz, 3 H) 1.38 (d, J = 6.02 Hz, 6 H) 1.56-1.64 (m, 1 H) 1.65-1.73 (m, 1 H) 1.76-1.86 (m, 1 H) 1.92-2.01 (m, 2 H) 3.11 (s, 3 H) 3.13-3.22 (m, 1 H) 3.37-3.45 (m, 1 H) 3.47-3.64 (m, 2 H) 3.74 (s, 1 H) 4.06 (ddd, J = 10.69, 7.31, 3.72 Hz, 1 H) 4.27 (ddd, J = 10.45, 6.39, 3.94 Hz, 1 H) 4.72 (dt, J = 12.07, 6.02 Hz, 1 H) 6.78 (d, J = 10.17 Hz, 2 H) 7.74 (d, J = 5.72 Hz, 1 H) 7.99 (t, J = 8.71 Hz, 1 H) 8.37 (dd, J = 8.61, 3.91 Hz, 1 H) 8.55 (d, J = 5.62 Hz, 1 H) 9.13 (s, 1 H)
- 143 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.08 (d, J = 6.26 Hz, 3 H) 1.34-1.53 (m, 1 H) 1.60-1.75 (m, 1 H) 1.79-2.00 (m, 2 H) 2.17-2.29 (m, 1 H) 2.98 (s, 1 H) 3.09-3.24 (m, 2 H) 3.48-3.58 (m, 3 H) 3.91 (s, 3 H) 6.80 (d, J = 9.78 Hz, 2 H) 7.77 (s, 1 H) 8.00 (s, 1 H) 8.31-8.44 (m, 1 H) 8.47-8.60 (m, 1 H) 9.28 (s, 1 H)
- 144 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.13 (d, J = 7.04 Hz, 3 H) 1.51-1.72 (m, 2 H) 1.74-1.87 (m, 1 H) 1.89-1.98 (m, 2 H) 3.11 (s, 4 H) 3.35-3.43 (m, 1 H) 3.51-3.60 (m, 2 H) 3.66-3.74 (m, 1 H) 3.86-3.95 (m, 3 H) 4.00-4.11 (m, 1 H) 4.22-4.32 (m, 1 H) 6.75-6.87 (m, 2 H) 7.64-7.72 (m, 1 H) 7.93-8.03 (m, 1 H) 8.33-8.42 (m, 1 H) 8.48-8.57 (m, 1 H) 9.00-9.08 (m, 1 H)
- 145 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.03-1.11 (m, 3 H) 1.34-1.49 (m, 1 H) 1.63 (ddd, J = 12.42, 6.16, 3.33 Hz, 1 H) 1.81 (q, J = 12.13 Hz, 1 H) 1.92 (dd, J = 13.30, 2.74 Hz, 1 H) 2.14-2.27 (m, 1 H) 2.89-2.99 (m, 1 H) 3.05-3.19 (m, 2 H) 3.53 (s, 3 H) 7.10-7.18 (m, 3 H) 7.49-7.59 (m, 1 H) 7.70 (d, J = 5.48 Hz, 1 H) 8.47 (d, J = 5.48 Hz, 1 H) 9.35 (s, 1 H)
- 146 ¹H NMR (400 MHz, <dms_o-d₆>) δ ppm 0.96 (d, J = 6.65 Hz, 3 H) 1.26 (d, J = 12.52 Hz, 1 H) 1.46 (br. s., 1 H) 1.54-1.80 (m, 3 H) 2.90 (br. s., 1 H) 3.07 (s, 3 H) 3.15-3.27 (m, 1 H) 3.38 (br. s., 1 H) 3.55 (br. s., 1 H) 3.64 (d, J = 5.48 Hz, 1 H) 3.87-3.91 (m, 1 H) 4.00 (d, J = 3.91 Hz, 1 H) 7.20-7.29 (m, 4 H) 7.30-7.44 (m, 1 H) 7.58 (s, 1 H) 7.95 (br. s., 2 H) 8.43 (d, J = 5.09 Hz, 1 H) 8.78 (s, 1 H) 10.12 (s, 1 H)
- 147 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.44 (d, J = 6.85 Hz, 3 H) 1.40 (dd, J = 6.02, 3.03 Hz, 6 H) 2.54 (ddd, J = 11.57, 4.62, 2.20 Hz, 1 H) 3.52-3.62 (m, 1 H) 3.63-3.72 (m, 1 H) 3.88 (dd, J = 11.64, 4.16 Hz, 1 H) 4.03-4.11 (m, 1 H) 4.14-4.23 (m, 1 H) 4.78 (dt, J = 12.06, 6.05 Hz, 1 H) 5.09 (t, J = 4.35 Hz, 1 H) 6.84 (d, J = 10.42 Hz, 2 H) 7.52 (d, J = 6.50 Hz, 1 H) 7.89 (d, J = 1.03 Hz, 1 H) 8.00 (t, J = 8.71 Hz, 1 H) 8.09 (d, J = 1.03 Hz, 1 H) 8.34-8.46 (m, 2 H) 9.18 (s, 1 H)
- 148 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.65 (d, J = 6.85 Hz, 3 H) 1.40 (dd, J = 6.02, 2.93 Hz, 6 H) 2.17-2.27 (m, 1 H) 2.83 (t, J = 12.98 Hz, 1 H) 3.22-3.29 (m, 1 H) 3.47-3.62 (m, 2 H) 3.64-3.73 (m, 4 H) 4.06-4.15 (m, 1 H) 4.76 (dt, J = 12.07, 6.02 Hz, 1 H) 6.83 (d, J = 10.51 Hz, 2 H) 7.25 (d, J = 9.68 Hz, 1 H) 7.51 (d, J = 6.60 Hz, 1 H) 8.01 (t, J = 8.66 Hz, 1 H) 8.37-8.44 (m, 2 H) 9.33 (s, 1 H)
- 149 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.94 (d, J = 6.75 Hz, 3 H) 1.38 (d, J = 6.02 Hz, 6 H) 1.43-1.57 (m, 1 H) 1.71 (d, J = 13.60 Hz, 1 H) 1.78-2.02 (m, 3 H) 3.10-3.21 (m, 1 H) 3.41 (dt, J = 12.41, 3.58 Hz, 1 H) 3.72 (s, 3 H) 4.08-4.16 (m, 1 H) 4.72 (dt, J = 12.09, 6.06 Hz, 1 H) 6.78 (d, J = 10.22 Hz, 2 H) 7.21 (d, J = 9.73 Hz, 1 H) 7.82 (d, J = 5.62 Hz, 1 H) 7.98 (t, J = 8.71 Hz, 1 H) 8.37 (dd, J = 8.61, 3.91 Hz, 1 H) 8.54 (d, J = 5.58 Hz, 1 H) 9.08 (s, 1 H)
- 150 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.91-1.01 (m, 3 H) 1.73-1.86 (m, 1 H) 1.90-2.01 (m, 1 H) 2.12-2.31 (m, 2 H) 2.34-2.46 (m, 1 H) 3.53-3.66 (m, 1 H) 3.77-3.88 (m, 1 H) 5.15-5.25 (m, 1 H) 6.03-6.13 (m, 1 H) 6.52-6.61 (m, 1 H) 7.10-7.24 (m, 2 H) 7.42-7.50 (m, 1 H) 7.53-7.66 (m, 1 H) 7.70-7.77 (m, 1 H) 7.77-7.85 (m, 1 H) 7.99-8.05 (m, 1 H) 8.36-8.43 (m, 1 H) 8.51-8.58 (m, 1 H) 8.76-8.82 (m, 1 H)
- 151 ¹H NMR (400 MHz, <dms_o-d₆>) δ ppm -0.10-0.12 (m, 3 H) 1.64 (br. s., 2 H) 1.85-2.12 (m, 3 H) 3.43 (t, J = 11.15 Hz, 4 H) 3.52-3.73 (m, 5 H) 4.68 (br. s., 1 H) 7.07-7.26 (m, 3 H) 7.61 (s, 1 H) 7.85 (s, 3 H) 7.92-8.00 (m, 1 H) 8.05-8.18 (m, 2 H) 8.59 (br. s., 1 H) 10.16 (br. s., 1 H)
- 152 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.95 (d, J = 7.04 Hz, 3 H) 1.72-1.87 (m, 1 H) 1.96 (d, J = 14.09 Hz, 1 H) 2.11-2.46 (m, 3 H) 3.60 (br. s., 1 H) 3.84 (br. s., 1 H) 5.20 (d, J = 5.48 Hz, 1 H) 6.09 (t, J = 6.46 Hz, 1 H) 6.57 (d, J = 9.00 Hz, 1 H) 7.16 (t, J = 8.61 Hz, 2 H) 7.46 (t, J = 7.83 Hz, 1 H) 7.59 (quin, J = 7.43 Hz, 1 H) 7.74 (d, J = 7.04 Hz, 1 H) 7.86 (br. s., 1 H) 8.03 (t, J = 8.80 Hz, 1 H) 8.40 (dd, J = 8.41, 3.72 Hz, 1 H) 8.55 (d, J = 5.09 Hz, 1 H) 8.83 (d, J = 9.00 Hz, 1 H)
- 153 ¹H NMR (400 MHz, <dms_o-d₆>) δ ppm 0.96 (d, J = 6.65 Hz, 3 H) 1.25-1.40 (m, 1 H) 1.40-1.52 (m, 1 H) 1.56-1.76 (m, 3 H) 1.77-1.93 (m, 2 H) 2.11-2.36 (m, 2 H) 2.74-2.99 (m, 3 H) 3.24 (br. s., 1 H) 3.83-3.93 (m, 2 H) 7.24 (d, J = 5.09 Hz, 1 H) 7.36-7.50 (m, 2 H) 7.86 (br. s., 3 H) 8.18 (t, J = 9.00 Hz, 1 H) 8.30 (dd, J = 8.80, 4.11 Hz, 1 H) 8.45 (d, J = 5.09 Hz, 1 H) 8.56 (s, 1 H) 10.41 (s, 1 H)
- 154 ¹H NMR (400 MHz, <dms_o-d₆>) δ ppm 0.52 (d, J = 6.65 Hz, 3 H) 1.64 (d, J = 13.30 Hz, 1 H) 1.79-2.02 (m, 2 H) 2.16-2.41 (m, 2 H) 3.20 (t, J = 12.13 Hz, 1 H) 3.81 (br. s., 1 H) 4.91 (t, J = 3.72 Hz, 1 H) 7.26 (t, J = 8.41 Hz, 2 H) 7.47-7.68 (m, 4 H) 7.80-7.87 (m, 1 H) 8.00 (br. s., 3 H) 8.13 (s, 1 H) 8.51 (d, J = 5.48 Hz, 1 H) 8.72 (s, 1 H) 9.60 (s, 1 H)

TABLE 3-continued

Ex # ¹H-NMR data

- 155 ¹H NMR (400 MHz, <dms-*d*₆>) δ ppm 0.48 (d, J = 6.65 Hz, 3 H) 1.07-1.24 (m, 3 H) 1.60 (d, J = 13.30 Hz, 1 H) 1.78-1.97 (m, 2 H) 2.10-2.23 (m, 1H) 2.33 (q, J = 12.52 Hz, 1 H) 3.15 (t, J = 12.33 Hz, 1 H) 3.52 (q, J = 7.04 Hz, 2 H) 4.49-4.62 (m, 2 H) 4.88 (t, J = 3.72 Hz, 1 H) 7.27 (d, J = 8.61 Hz, 2H) 7.52 (d, J = 5.48 Hz, 1 H) 7.79-7.87 (m, 1 H) 8.00 (br. s., 3 H) 8.17 (t, J = 8.80 Hz, 1 H) 8.31 (dd, J = 8.61, 3.91 Hz, 1 H) 8.55 (d, J = 5.09 Hz, 1 H), 8.58 (s, 1 H) 10.52 (s, 1 H)
- 156 ¹H NMR (400 MHz, <dms-*d*₆>) δ ppm 0.01 (d, J = 7.04 Hz, 3 H) 2.94-3.13 (m, 2 H) 3.21 (d, J = 10.56 Hz, 2 H) 3.31-3.44 (m, 1 H) 3.45-3.63 (m, 2H) 4.71 (t, J = 3.91 Hz, 1 H) 6.89 (t, J = 8.80 Hz, 2 H) 7.06 (d, J = 6.65 Hz, 1 H) 7.12-7.23 (m, 1 H) 7.31 (br. s., 2 H) 7.48 (s, 1 H) 7.80 (s, 2 H) 8.00 (d, J = 6.26 Hz, 1 H) 8.60 (s, 1 H) 9.04 (br. s., 1 H)
- 157 ¹H NMR (400 MHz, <dms-*d*₆>) δ ppm 0.01 (d, J = 7.04 Hz, 3 H) 0.85-0.99 (m, 3 H) 2.01-2.15 (m, 1 H) 3.07-3.24 (m, 1 H) 3.30 (q, J = 7.04 Hz, 3H) 3.43 (d, J = 10.17 Hz, 2 H) 3.52-3.70 (m, 4 H) 4.26-4.40 (m, 2 H) 4.71 (br. s., 1 H) 6.95-7.09 (m, 2 H) 7.17 (d, J = 6.65 Hz, 1 H) 7.62 (s, 1 H), 7.88 (s, 1 H) 7.91-8.03 (m, 3 H) 8.07-8.20 (m, 2 H) 8.66 (s, 1 H) 10.17 (br. s., 1 H)
- 158 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.63 (d, J = 6.85 Hz, 3 H) 1.89 (dt, J = 11.38, 8.98 Hz, 1 H) 2.08-2.22 (m, 2 H) 2.40-2.53 (m, 2 H) 2.54-2.65 (m, 2 H) 2.78 (t, J = 12.76 Hz, 1 H) 3.08-3.27 (m, 2 H) 3.34-3.43 (m, 2 H) 3.46-3.59 (m, 2 H) 3.71 (s, 3 H) 4.02 (br. s., 1 H) 7.38-7.47 (m, 3 H) 8.06 (t, J = 8.68 Hz, 1 H) 8.39 (d, J = 6.46 Hz, 1 H) 8.46 (dd, J = 8.66, 3.96 Hz, 1 H) 9.39 (s, 1 H)
- 159 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.65 (d, J = 6.80 Hz, 3 H) 1.74-1.89 (m, 2 H) 2.06-2.25 (m, 3 H) 2.66 (s, 1 H) 2.76 (t, J = 12.18 Hz, 1 H) 3.12-3.22 (m, 1 H) 3.25-3.28 (m, 1 H) 3.36 (dd, J = 3.86, 2.20 Hz, 3 H) 3.45-3.58 (m, 4 H) 3.61-3.70 (m, 3 H) 3.72 (s, 3 H) 3.93-4.04 (m, 2 H) 4.04-4.10 (m, 1 H) 6.88-6.99 (m, 2 H) 7.42 (d, J = 6.31 Hz, 1 H) 8.03 (t, J = 8.68 Hz, 1 H) 8.33-8.47 (m, 2 H) 9.44 (s, 1 H)
- 160 ¹H NMR, (400 MHz, <cd3od>) δ ppm 9.16 (s, 1 H), 8.48 (dd, J = 8.6, 3.9 Hz, 1 H), 8.41 (d, J = 6.3 Hz, 1 H), 8.02-8.13 (m, 1 H), 7.93 (s, 1 H), 7.57 (d, J = 6.7 Hz, 1 H), 7.44 (d, J = 9.8 Hz, 2 H), 5.08 (t, J = 4.1 Hz, 1 H), 4.20-4.32 (m, 1 H), 4.06-4.16 (m, 1 H), 3.95-4.05 (m, 3 H), 3.87-3.95 (m, 2 H), 3.76-3.86 (m, 1 H), 3.56-3.69 (m, 1 H), 2.46-2.60 (m, 1 H), 2.16-2.32 (m, 2 H), 1.73 (d, J = 13.7 Hz, 2 H), 0.43 (d, J = 7.0 Hz, 3 H)
- 161 ¹H NMR, (400 MHz, <cd3od>) δ ppm 9.15 (s, 1 H), 8.46 (dd, J = 8.6, 3.9 Hz, 1 H), 8.41 (d, J = 6.7 Hz, 1 H), 8.12 (s, 1 H), 8.05 (t, J = 8.8 Hz, 1 H), 7.92 (s, 1 H), 7.58 (d, J = 7.0 Hz, 1 H), 7.11 (d, J = 9.8 Hz, 2 H), 5.08 (t, J = 4.3 Hz, 1 H), 4.19-4.31 (m, 1 H), 4.04-4.15 (m, 1 H), 3.99 (dd, J = 11.9, 4.1 Hz, 1 H), 3.80 (dd, J = 13.5, 4.1 Hz, 1 H), 3.60 (t, J = 12.9 Hz, 1 H), 2.45-2.60 (m, 4 H), 0.44 (d, J = 7.0 Hz, 3 H)
- 162 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.58 (d, J = 6.70 Hz, 3 H) 2.19 (br. s., 1 H) 2.79 (t, J = 13.01 Hz, 1 H) 3.20-3.28 (m, 1 H) 3.45 (d, J = 11.25 Hz, 1 H) 3.52-3.65 (m, 2 H) 3.71 (s, 3 H) 4.02 (br. s., 1 H) 4.43 (quin, J = 7.25 Hz, 1 H) 4.79 (t, J = 6.31 Hz, 1 H) 5.15-5.22 (m, 2 H) 7.39 (d, J = 9.63 Hz, 2 H) 7.48 (d, J = 6.46 Hz, 1 H) 8.07 (t, J = 8.71 Hz, 1 H) 8.40 (d, J = 6.26 Hz, 1 H) 8.47 (dd, J = 8.66, 3.86 Hz, 1 H) 9.39 (s, 1 H)
- 163 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.87-0.99 (m, 1 H) 1.14 (d, J = 6.70 Hz, 3 H) 1.26-1.38 (m, 1 H) 1.66 (d, J = 8.75 Hz, 2 H) 1.80 (br. s., 1 H) 1.89-1.98 (m, 2 H) 2.78-2.89 (m, 1 H) 3.12 (d, J = 11.74 Hz, 1 H) 3.68 (br. s., 1 H) 3.83 (br. s., 1 H) 3.88-3.97 (m, 1 H) 4.35-4.42 (m, 1 H) 4.75 (t, J = 6.26 Hz, 2 H) 5.15 (t, J = 7.29 Hz, 2 H) 7.31 (d, J = 9.54 Hz, 2 H) 7.56 (d, J = 4.99 Hz, 1 H) 8.02 (t, J = 8.80 Hz, 1 H) 8.40 (d, J = 5.33 Hz, 1 H) 8.50 (d, J = 5.04 Hz, 1 H) 8.82 (s, 1 H)
- 164 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.46 (d, J = 6.70 Hz, 3 H) 2.39 (d, J = 5.62 Hz, 1 H) 2.85 (t, J = 12.84 Hz, 1 H) 3.59 (br. s., 1 H) 3.80-3.90 (m, 1 H) 3.98-4.14 (m, 4 H) 4.25 (d, J = 12.03 Hz, 1 H) 5.08-5.17 (m, 1 H) 5.32 (d, J = 11.88 Hz, 1 H) 5.46 (br. s., 1 H) 6.88 (d, J = 9.05 Hz, 1 H) 7.55 (d, J = 6.70 Hz, 1 H) 7.69 (d, J = 10.12 Hz, 1 H) 8.10 (t, J = 8.61 Hz, 1 H) 8.33 (d, J = 6.46 Hz, 1 H) 8.39 (dd, J = 8.58, 3.45 Hz, 1 H) 8.97 (s, 1 H) 9.19 (s, 1 H) 9.55 (s, 1 H)
- 165 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.12 (d, J = 6.65 Hz, 3 H) 1.54-1.70 (m, 2 H) 1.72-1.84 (m, 1 H) 1.87-2.00 (m, 2 H) 2.71-2.90 (m, 2 H) 3.01-3.17 (m, 1 H) 3.64-3.71 (m, 1 H) 3.76-3.97 (m, 5 H) 6.74-6.83 (m, 2 H) 7.54-7.63 (m, 1 H) 7.92-8.02 (m, 1 H) 8.31-8.39 (m, 1 H) 8.46-8.52 (m, 1 H) 8.89-8.96 (m, 1 H)
- 166 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.02-1.18 (m, 3 H) 1.50-1.75 (m, 3 H) 1.87-2.03 (m, 2 H) 2.70-2.90 (m, 2 H) 3.02-3.18 (m, 1 H) 3.34-3.50 (m, 1 H) 3.66 (br. s., 1 H) 3.76-3.96 (m, 2 H) 7.08-7.18 (m, 3 H) 7.47-7.68 (m, 2 H) 8.46 (d, J = 5.48 Hz, 1 H) 9.20 (s, 1 H)
- 167 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.94 (d, J = 6.70 Hz, 3 H) 1.24-1.36 (m, 1 H) 1.41-1.58 (m, 1 H) 1.62-1.99 (m, 5 H) 2.11 (br. s., 1 H) 2.37-2.62 (m, 4 H) 3.03-3.17 (m, 1 H) 3.73 (s, 3 H) 4.09 (br. s., 1 H) 7.17 (d, J = 9.10 Hz, 1 H) 7.36 (d, J = 9.54 Hz, 2 H) 7.68 (d, J = 5.33 Hz, 1 H) 8.02 (t, J = 8.68 Hz, 1 H) 8.40 (dd, J = 8.63, 3.74 Hz, 1 H) 8.49 (d, J = 5.28 Hz, 1 H) 8.91 (s, 1 H)
- 168 ¹H NMR, (400 MHz, <cd3od>) δ ppm 8.85 (s, 1 H), 8.58 (d, J = 5.48 Hz, 1 H), 8.38 (dd, J = 8.61, 3.52 Hz, 1 H), 8.07 (s, 1 H), 7.99 (t, J = 8.80 Hz, 1 H), 7.90 (s, 1 H), 7.82 (d, J = 5.09 Hz, 1 H), 6.84 (d, J = 10.17 Hz, 2 H), 4.98 (br. s., 1 H), 4.22 (br. s., 2 H), 3.89 (d, J = 12.52 Hz, 1 H), 3.78 (br. s., 2 H), 3.44 (s, 3 H), 2.64-2.74 (m, 2 H), 2.29 (br. s., 1 H), 2.02-2.18 (m, 2 H), 1.81 (d, J = 13.69 Hz, 1 H), 0.67 (d, J = 6.65 Hz, 3 H)

TABLE 3-continued

Ex # ¹H-NMR data

- 169 ¹H NMR, (400 MHz, <cd3od>) δ ppm 8.84 (s, 1 H), 8.65 (s, 1 H), 8.60 (d, J = 5.09 Hz, 1 H), 8.08 (s, 1 H), 7.90 (s, 1 H), 7.85 (d, J = 5.48 Hz, 1 H), 7.63 (t, J = 7.83 Hz, 1 H), 7.25 (t, J = 8.80 Hz, 2 H), 5.00 (br. s., 1 H), 3.95 (d, J = 12.52 Hz, 1 H), 2.63-2.76 (m, 1 H), 2.38 (br. s., 1 H), 2.06-2.21 (m, 2 H), 1.84 (d, J = 12.91 Hz, 1 H), 0.70 (d, J = 6.65 Hz, 3 H)
- 170 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.94 (d, J = 6.60 Hz, 3 H) 1.50 (q, J = 13.17 Hz, 1 H) 1.65-1.84 (m, 4 H) 1.85-2.01 (m, 2 H) 2.09 (d, J = 11.59 Hz, 2 H) 3.04-3.18 (m, 2 H) 3.35-3.51 (m, 2 H) 3.64 (t, J = 9.24 Hz, 2 H) 3.73 (s, 3 H) 3.93-4.02 (m, 2 H) 4.09 (br. s., 1 H) 4.71 (br. s., 1 H) 6.86 (d, J = 10.37 Hz, 2 H) 7.18 (d, J = 10.07 Hz, 1 H) 7.70 (d, J = 5.43 Hz, 1 H) 7.99 (t, J = 8.61 Hz, 1 H) 8.36 (dd, J = 8.73, 3.94 Hz, 1 H) 8.50 (d, J = 5.33 Hz, 1 H) 8.93 (s, 1 H)
- 171 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.67 (d, J = 6.70 Hz, 3 H) 1.68-1.87 (m, 3 H) 2.00-2.20 (m, 4 H) 2.30 (br. s., 1 H) 2.63-2.77 (m, 1 H) 3.63 (t, J = 8.95 Hz, 2 H) 3.84-4.03 (m, 3 H) 4.66-4.76 (m, 1 H) 4.98 (br. s., 1 H) 6.86 (d, J = 10.17 Hz, 2 H) 7.82 (d, J = 5.18 Hz, 1 H) 7.89 (s, 1 H) 7.99 (t, J = 8.66 Hz, 1 H) 8.07 (s, 1 H) 8.37 (dd, J = 8.66, 3.86 Hz, 1 H) 8.58 (d, J = 5.28 Hz, 1 H) 8.80 (s, 1 H)
- 172 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.65 (d, J = 6.60 Hz, 3 H) 1.73-1.92 (m, 2 H) 2.00-2.17 (m, 3 H) 2.27 (br. s., 1 H) 2.37-2.60 (m, 4 H) 2.63-2.76 (m, 1 H) 3.86 (br. s., 1 H) 4.96 (br. s., 2 H) 7.35 (d, J = 9.63 Hz, 2 H) 7.78 (d, J = 5.28 Hz, 1 H) 7.89 (s, 1 H) 7.99-8.08 (m, 2 H) 8.37-8.43 (m, 1 H) 8.56 (d, J = 5.43 Hz, 1 H) 8.79 (s, 1 H)
- 174 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.50 (d, J = 7.04 Hz, 3 H) 1.84-1.96 (m, 1 H) 2.54-2.66 (m, 1 H) 3.04-3.15 (m, 1 H) 3.25-3.45 (m, 3 H) 3.61 (s, 3 H) 3.76-3.83 (m, 1 H) 7.01-7.16 (m, 3 H) 7.32-7.41 (m, 1 H) 7.46-7.58 (m, 1 H) 8.22-8.32 (m, 1 H) 9.33-9.42 (m, 1 H)
- 175 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.87 (d, J = 6.65 Hz, 3 H) 1.39-1.53 (m, 1 H) 1.62-1.99 (m, 4 H) 3.04-3.16 (m, 1 H) 3.32-3.37 (m, 1 H) 3.70 (s, 3 H) 4.03-4.11 (m, 1 H) 7.08-7.21 (m, 4 H) 7.49-7.60 (m, 1 H) 7.80 (d, J = 5.48 Hz, 1 H) 8.48 (m, 1 H) 9.29 (s, 1 H)
- 176 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.52 (d, J = 6.65 Hz, 3 H) 1.65-1.75 (m, 1 H) 1.90-2.16 (m, 3 H) 2.59-2.73 (m, 1 H) 3.26-3.35 (m, 1 H) 3.71-3.81 (m, 1 H) 4.91-4.97 (m, 1 H) 7.02-7.13 (m, 3 H) 7.40-7.51 (m, 1 H) 7.79 (s, 2 H) 8.00 (s, 1 H) 8.45-8.52 (m, 1 H) 9.12 (s, 1 H)
- 177 ¹H NMR, (400 MHz, <cd3od>) δ ppm 9.46 (s, 1 H), 8.48 (dd, J = 8.6, 3.9 Hz, 1 H), 8.42 (d, J = 6.3 Hz, 1 H), 8.08 (t, J = 8.8 Hz, 1 H), 7.39-7.53 (m, 3 H), 3.86-4.09 (m, 5 H), 3.72 (s, 3 H), 3.57 (dd, J = 8.4, 3.7 Hz, 2 H), 3.37-3.47 (m, 1 H), 3.19-3.28 (m, 1 H), 2.77 (t, J = 12.9 Hz, 1 H), 2.10-2.36 (m, 3 H), 1.64-1.86 (m, 2 H), 0.56 (d, J = 6.7 Hz, 3 H)
- 178 ¹H NMR, (400 MHz, <cd3od>) δ ppm 8.99 (s, 1 H), 8.54 (d, J = 5.5 Hz, 1 H), 8.42 (dd, J = 9.0, 3.9 Hz, 1 H), 8.04 (t, J = 8.8 Hz, 1 H), 7.75 (d, J = 5.5 Hz, 1 H), 7.39 (d, J = 9.8 Hz, 2 H), 7.19 (d, J = 9.8 Hz, 1 H), 4.12 (d, J = 9.0 Hz, 1 H), 3.84-4.04 (m, 4 H), 3.74 (s, 3 H), 3.39 (d, J = 3.5 Hz, 1 H), 3.14 (t, J = 12.1 Hz, 1 H), 2.12-2.28 (m, 2 H), 1.86-2.03 (m, 2 H), 1.64-1.85 (m, 4 H), 1.51 (q, J = 12.8 Hz, 1 H), 0.94 (d, J = 6.7 Hz, 3 H)
- 179 ¹H NMR, (400 MHz, <cd3od>) δ ppm 8.81 (s, 1 H), 8.60 (d, J = 5.5 Hz, 1 H), 8.42 (dd, J = 8.6, 3.9 Hz, 1 H), 8.08 (s, 1 H), 8.04 (t, J = 8.6 Hz, 1 H), 7.91 (s, 1 H), 7.84 (d, J = 5.5 Hz, 1 H), 7.38 (d, J = 9.8 Hz, 2 H), 5.00 (t, J = 3.9 Hz, 1 H), 3.81-4.04 (m, 4 H), 3.37 (br. s., 1 H), 2.64-2.80 (m, 1 H), 2.26-2.39 (m, 1 H), 2.02-2.25 (m, 4 H), 1.82 (d, J = 13.3 Hz, 1 H), 1.68 (d, J = 13.3 Hz, 2 H), 0.67 (d, J = 6.7 Hz, 3 H)
- 180 ¹H NMR, (400 MHz, <cd3od>) δ ppm 9.03 (s, 1 H), 8.55 (d, J = 5.5 Hz, 1 H), 8.41 (dd, J = 8.6, 3.9 Hz, 1 H), 8.02 (t, J = 8.6 Hz, 1 H), 7.80 (d, J = 5.9 Hz, 1 H), 7.20 (d, J = 9.8 Hz, 1 H), 7.07 (d, J = 9.4 Hz, 2 H), 4.12 (d, J = 9.4 Hz, 1 H), 3.74 (s, 3 H), 3.36-3.47 (m, 1 H), 3.09-3.23 (m, 1 H), 2.49 (s, 3 H), 1.65-2.04 (m, 4 H), 1.52 (q, J = 13.2 Hz, 1 H), 0.95 (d, J = 7.0 Hz, 3 H)
- 181 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.68 (d, J = 6.70 Hz, 3 H) 1.76-1.88 (m, 1 H) 2.02-2.20 (m, 2 H) 2.23-2.38 (m, 1 H) 2.49 (s, 3 H) 2.64-2.80 (m, 1 H) 3.35-3.39 (m, 1 H) 3.90 (dt, J = 12.72, 4.43 Hz, 1 H) 4.97-5.03 (m, 1 H) 7.07 (d, J = 9.44 Hz, 2 H) 7.84 (d, J = 5.48 Hz, 1 H) 7.91 (d, J = 0.98 Hz, 1 H) 8.02 (t, J = 8.71 Hz, 1 H) 8.09 (d, J = 1.03 Hz, 1 H) 8.41 (dd, J = 8.61, 3.96 Hz, 1 H) 8.60 (d, J = 5.48 Hz, 1 H) 8.84 (s, 1 H)
- 182 ¹H NMR, (400 MHz, <cd3od>) δ ppm 0.94 (d, J = 6.75 Hz, 3 H) 1.51 (q, J = 12.78 Hz, 1 H) 1.71 (d, J = 13.64 Hz, 1 H) 1.80 (q, J = 12.41 Hz, 1 H) 1.88-2.02 (m, 2 H) 3.10-3.20 (m, 1 H) 3.41 (dt, J = 12.47, 3.55 Hz, 1 H) 3.74 (s, 3 H) 4.08-4.14 (m, 1 H) 4.36-4.45 (m, 1 H) 4.77 (td, J = 6.28, 1.47 Hz, 2 H) 5.16 (dd, J = 8.22, 6.21 Hz, 2 H) 7.20 (d, J = 9.73 Hz, 1 H) 7.33 (d, J = 9.10 Hz, 2 H) 7.78 (d, J = 5.58 Hz, 1 H) 8.04 (t, J = 8.71 Hz, 1 H) 8.42 (dd, J = 8.68, 3.94 Hz, 1 H) 8.55 (d, J = 5.53 Hz, 1 H) 8.98 (s, 1 H)
- 183 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.13 (d, J = 6.85 Hz, 3 H) 1.53-1.85 (m, 5 H) 1.85-2.00 (m, 2 H) 2.00-2.15 (m, 2 H) 2.70-2.94 (m, 2 H) 3.03-3.16 (m, 2 H) 3.56-3.70 (m, 3 H) 3.75-3.86 (m, 1 H) 3.87-4.05 (m, 3 H) 4.70 (dt, J = 8.03, 4.22 Hz, 1 H) 6.84 (d, J = 10.12 Hz, 2 H) 7.57 (d, J = 5.48 Hz, 1 H) 7.97 (t, J = 8.71 Hz, 1 H) 8.35 (dd, J = 8.63, 3.94 Hz, 1 H) 8.49 (d, J = 5.48 Hz, 1 H) 8.86 (s, 1 H)
- 184 ¹H NMR, (400 MHz, <cd3od>) δ ppm 1.13 (d, J = 6.85 Hz, 3 H) 1.58-1.73 (m, 2 H) 1.73-2.02 (m, 4 H) 2.05-2.23 (m, 1 H) 2.33-2.63 (m, 4 H) 2.72-2.92 (m, 2 H) 3.02-3.18 (m, 1 H) 3.69 (s, 1 H) 3.82 (dt, J = 9.06, 5.42 Hz, 1 H)

TABLE 3-continued

Ex #	¹ H-NMR data
	3.94 (td, J = 8.63, 5.09 Hz, 1 H) 7.36 (d, J = 9.49 Hz, 2 H) 7.59 (d, J = 5.48 Hz, 1 H) 8.03 (t, J = 8.71 Hz, 1 H) 8.40 (dd, J = 8.63, 3.94 Hz, 1 H) 8.51 (d, J = 5.43 Hz, 1 H) 8.92 (s, 1 H)
185	¹ H NMR, (400 MHz, <cd3od>) δ ppm 0.69 (d, J = 6.75 Hz, 3 H) 1.84 (d, J = 13.69 Hz, 1 H) 1.90-2.10 (m, 1 H) 2.16 (d, J = 12.47 Hz, 1 H) 2.32 (br. s., 1 H) 2.50-2.72 (m, 1 H) 3.94 (dt, J = 12.89, 4.56 Hz, 1 H) 5.04 (t, J = 4.16 Hz, 1 H) 7.22 (t, J = 8.29 Hz, 2 H) 7.58-7.70 (m, 1 H) 7.75 (d, J = 5.43 Hz, 1 H) 8.04 (t, J = 8.71 Hz, 1 H) 8.42 (dd, J = 8.66, 3.96 Hz, 1 H) 8.57 (d, J = 5.38 Hz, 1 H) 8.75 (s, 1 H) 8.81 (s, 1 H)
186	¹ H NMR, (400 MHz, <cd3od>) δ ppm 0.67 (d, J = 6.70 Hz, 3 H) 0.76-0.91 (m, 2 H) 0.98-1.08 (m, 2 H) 1.72-1.84 (m, 1 H) 1.96-2.17 (m, 3 H) 2.17-2.36 (m, 1 H) 2.58-2.78 (m, 1 H) 3.84 (dt, J = 12.67, 4.45 Hz, 1 H) 7.22 (t, J = 8.27 Hz, 2 H) 7.64 (tt, J = 8.49, 6.47 Hz, 1 H) 7.77 (s, 1 H) 7.82 (d, J = 5.58 Hz, 1 H) 8.04 (t, J = 8.73 Hz, 1 H) 8.42 (dd, J = 8.66, 3.96 Hz, 1 H) 8.59 (d, J = 5.38 Hz, 1 H) 8.86 (s, 1 H)
187	¹ H NMR, (400 MHz, <cd3od>) δ ppm 0.69 (d, J = 6.65 Hz, 3 H) 1.80 (d, J = 13.30 Hz, 1 H) 2.00-2.14 (m, 2 H) 2.17 (s, 3 H) 2.20-2.36 (m, 1 H) 2.60-2.80 (m, 1 H) 3.87 (dt, J = 12.63, 4.32 Hz, 1 H) 4.91 (d, J = 6.36 Hz, 1 H) 5.19 (s, 1 H) 5.75 (s, 1 H) 7.22 (t, J = 8.29 Hz, 2 H) 7.63 (tt, J = 8.48, 6.49 Hz, 1 H) 7.84 (br. s., 1 H) 8.03 (t, J = 8.71 Hz, 1 H) 8.09 (d, J = 1.86 Hz, 1 H) 8.41 (dd, J = 8.66, 3.91 Hz, 1 H) 8.54-8.63 (m, 1 H) 8.76-8.91 (m, 1 H)
188	¹ H NMR, (400 MHz, <cd3od>) δ ppm 0.66 (d, J = 6.75 Hz, 3 H) 1.63 (s, 6 H) 1.71-1.82 (m, 1 H) 1.90-2.18 (m, 2 H) 2.24 (d, J = 12.57 Hz, 1 H) 2.61-2.79 (m, 1 H) 3.85 (dt, J = 12.64, 4.34 Hz, 1 H) 7.21 (t, J = 8.29 Hz, 2 H) 7.57-7.70 (m, 1 H) 7.81 (d, J = 5.48 Hz, 1 H) 7.87 (s, 1 H) 8.02 (t, J = 8.71 Hz, 1 H) 8.41 (dd, J = 8.63, 3.99 Hz, 1 H) 8.57 (d, J = 5.58 Hz, 1 H) 8.82 (s, 1 H)
189	¹ H NMR, (400 MHz, <cd3od>) δ ppm 0.97 (s, 3 H) 1.45-1.60 (m, 2 H) 1.60-1.71 (m, 1 H) 1.80-1.90 (m, 1 H) 1.91-2.03 (m, 1 H) 2.97-3.11 (m, 1 H) 3.26-3.31 (m, 1 H) 3.60-3.67 (m, 1 H) 4.08-4.17 (m, 1 H) 4.35-4.45 (m, 1 H) 7.06-7.17 (m, 2 H) 7.48-7.65 (m, 2 H) 7.90-7.98 (m, 1 H) 8.28-8.36 (m, 1 H) 8.41-8.47 (m, 1 H) 8.91-8.97 (m, 1 H)
190	¹ H NMR, (400 MHz, <cd3od>) δ ppm 1.02-1.10 (m, 3 H) 1.52-1.67 (m, 2 H) 1.68-1.80 (m, 1 H) 1.88-1.96 (m, 1 H) 1.97-2.10 (m, 1 H) 3.04-3.16 (m, 1 H) 3.35-3.41 (m, 1 H) 3.67-3.73 (m, 1 H) 4.16-4.25 (m, 1 H) 4.43-4.52 (m, 1 H) 7.14-7.24 (m, 2 H) 7.56-7.69 (m, 2 H) 7.97-8.05 (m, 1 H) 8.36-8.43 (m, 1 H) 8.47-8.55 (m, 1 H) 8.94-9.00 (m, 1 H)
191	¹ H NMR, (400 MHz, <cd3od>) δ ppm 0.88-0.97 (m, 3 H) 1.42-1.55 (m, 1 H) 1.63-1.80 (m, 2 H) 1.85-1.98 (m, 2 H) 3.03-3.16 (m, 1 H) 3.36-3.41 (m, 1 H) 3.66-3.74 (m, 3 H) 3.85-3.92 (m, 3 H) 4.04-4.12 (m, 1 H) 6.78 (s, 1 H) 7.64-7.72 (m, 1 H) 7.92-8.01 (m, 1 H) 8.31-8.39 (m, 1 H) 8.45-8.51 (m, 1 H) 8.86-8.94 (m, 1 H)
192	¹ H NMR, (400 MHz, <cd3od>) δ ppm 0.64 (d, J = 6.65 Hz, 3 H) 1.73-1.85 (m, 1 H) 2.00-2.16 (m, 2 H) 2.21-2.34 (m, 1 H) 2.61-2.76 (m, 1 H) 3.34-3.41 (m, 1 H) 3.88 (s, 4 H) 4.95-5.02 (m, 1 H) 6.74-6.84 (m, 2 H) 7.79-7.85 (m, 1 H) 7.85-7.91 (m, 1 H) 7.93-8.01 (m, 1 H) 8.02-8.09 (m, 1 H) 8.31-8.40 (m, 1 H) 8.53-8.61 (m, 1 H) 8.81-8.89 (m, 1 H)
193	¹ H NMR, (400 MHz, <cd3od>) δ ppm 0.43 (d, J = 6.65 Hz, 3 H) 2.44-2.58 (m, 1 H) 3.57-3.67 (m, 1 H) 3.76-3.84 (m, 1 H) 3.93 (s, 3 H) 3.97-4.13 (m, 2 H) 4.19-4.29 (m, 1 H) 5.06-5.12 (m, 1 H) 6.79-6.89 (m, 2 H) 7.52-7.60 (m, 1 H) 7.84-7.91 (m, 1 H) 7.95-8.04 (m, 1 H) 8.07-8.14 (m, 1 H) 8.34-8.44 (m, 2 H) 9.07-9.13 (m, 1 H)
194	¹ H NMR, (400 MHz, <cd3od>) δ ppm -0.02-0.04 (m, 3 H) 1.94-2.07 (m, 1 H) 3.01-3.11 (m, 2 H) 3.16-3.24 (m, 1 H) 3.35-3.44 (m, 1 H) 3.50-3.59 (m, 1 H) 3.72-3.82 (m, 1 H) 6.78 (s, 1 H) 7.12-7.29 (m, 2 H) 7.49-7.52 (m, 1 H) 7.70-7.74 (m, 1 H) 7.97-8.04 (m, 1 H) 8.92-8.97 (m, 1 H)
198	¹ H NMR (400 MHz, <dms _o -d ₆ >) δ ppm 0.59 (d, J = 6.65 Hz, 3 H) 1.16 (dd, J = 8.41, 6.85 Hz, 6 H) 2.06 (d, J = 3.91 Hz, 1 H) 3.07-3.33 (m, 5 H) 3.56 (s, 3 H) 3.86 (d, J = 9.78 Hz, 1 H) 4.18 (dq, J = 13.60, 6.68 Hz, 1 H) 7.19 (br. s., 1 H) 7.28-7.52 (m, 4 H) 7.78 (dd, J = 8.61, 1.96 Hz, 1 H) 7.86-8.06 (m, 4 H) 8.28-8.52 (m, 3 H) 9.16 (s, 1 H) 10.24 (s, 1 H)
199	¹ H NMR (400 MHz, <dms _o -d ₆ >) δ ppm 0.45 (d, J = 7.04 Hz, 3 H) 1.74-2.02 (m, 3 H) 2.07-2.41 (m, 2 H) 2.84-2.99 (m, 1 H) 3.08-3.22 (m, 1 H) 3.30 (d, J = 14.48 Hz, 2 H) 3.48 (d, J = 9.39 Hz, 1 H) 3.56 (s, 3 H) 3.69 (t, J = 11.15 Hz, 2 H) 3.79-4.00 (m, 3 H) 7.28 (d, J = 6.26 Hz, 1 H) 7.36 (d, J = 10.17 Hz, 1 H) 7.47 (d, J = 9.39 Hz, 2 H) 8.03 (br. s., 3 H) 8.23 (t, J = 8.80 Hz, 1 H) 8.29-8.46 (m, 2 H) 9.00 (br. s., 1 H) 10.36 (s, 1 H)
200	¹ H NMR (400 MHz, <dms _o -d ₆ >) δ ppm 0.75 (d, J = 6.65 Hz, 3 H) 1.06-1.21 (m, 6 H) 1.42 (d, J = 12.91 Hz, 1 H) 1.51-1.71 (m, 2 H) 1.77 (d, J = 12.13 Hz, 2 H) 2.99 (br. s., 1 H) 3.32 (br. s., 1 H) 3.52-3.68 (m, 3 H) 3.85-3.99 (m, 1 H) 4.09 (dq, J = 13.79, 6.75 Hz, 1 H) 7.31-7.44 (m, 3 H) 7.69-7.85 (m, 5 H) 7.88 (ddd, J = 8.31, 4.60, 2.35 Hz, 1 H) 8.32 (d, J = 7.43 Hz, 1 H) 8.40 (dd, J = 7.83, 1.96 Hz, 1 H) 8.49 (d, J = 5.09 Hz, 1 H) 8.88 (s, 1 H) 10.36 (s, 1 H)
201	¹ H NMR (400 MHz, <dms _o -d ₆ >) δ ppm 0.76 (d, J = 6.65 Hz, 3 H) 1.29-1.44 (m, 1 H) 1.44-1.80 (m, 4 H) 1.80-1.95 (m, 2 H) 2.18 (td, J = 13.21, 5.28 Hz, 1 H) 2.28 (td, J = 13.01, 5.67 Hz, 1 H) 2.79-2.93 (m, 1 H) 3.20 (br. s., 1 H) 3.53-3.63 (m, 3 H) 3.68 (t, J = 11.35 Hz, 2 H) 3.79-3.98 (m, 4 H) 7.33-7.53 (m, 3 H) 7.67-7.90 (m, 4 H) 8.11-8.23 (m, 1 H) 8.25-8.35 (m, 1 H) 8.48 (d, J = 5.48 Hz, 1 H) 8.54-8.63 (m, 1 H) 10.37-10.47 (m, 1 H)

TABLE 3-continued

Ex # ¹ H-NMR data	
202	¹ H NMR (400 MHz, <cdcl ₃ >) δ ppm 0.87 (br. s., 3 H) 1.27 (t, J = 6.65 Hz, 3 H) 1.45-1.68 (m, 2 H) 1.80 (br. s., 1 H) 2.05 (br. s., 1 H) 2.24 (br. s., 1 H) 2.99 (br. s., 1 H) 3.41-3.68 (m, 6 H) 4.23 (br. s., 1 H) 4.51-4.66 (m, 2 H) 6.79-7.19 (m, 3 H) 7.70-8.11 (m, 2 H) 8.20-8.64 (m, 3 H) 9.40-9.64 (m, 1 H) 10.17-10.40 (m, 1 H)
203	¹ H NMR (400 MHz, <cdcl ₃ >) δ ppm 0.42-0.71 (m, 3 H) 1.28 (br. s., 3 H) 2.06 (br. s., 1 H) 2.88 (br. s., 1 H) 3.25 (br. s., 1 H) 3.47-3.83 (m, 8 H) 4.10 (br. s., 1 H) 4.62 (br. s., 2 H) 6.83-7.06 (m, 1 H) 7.18 (br. s., 2 H) 7.48 (br. s., 1 H) 7.75 (br. s., 1 H) 8.34 (br. s., 2 H) 9.45-9.68 (m, 1 H) 10.05 (br. s., 1 H)

Pim1, Pim2, Pim3 AlphaScreen Assays

Pim 1, Pim 2 & Pim 3 AlphaScreen assays using high ATP (11-125× ATP Km) were used to determine the biochemical activity of the inhibitors. The activity of Pim 1, Pim 2, & Pim 3 is measured using a homogeneous bead based system quantifying the amount of phosphorylated peptide substrate resulting from kinase-catalyzed phosphoryl transfer to a peptide substrate. Compounds to be tested are dissolved in 100% DMSO and directly distributed to a white 384-well plate at 0.25 µl per well. To start the reaction, 5 µl of 100 nM Bad peptide (Biotin-AGAGRSRHSYPAGT-OH (SEQ ID NO:1)) and ATP (concentrations described below) in assay buffer (50 mM Hepes, pH=7.5, 5 mM MgCl₂, 0.05% BSA, 0.01% Tween-20, 1 mM DTT) is added to each well. This is followed by the addition of 5 µl/well of Pim 1, Pim 2 or Pim 3 kinase in assay buffer (concentrations described below). Final assay concentrations (described below) are in 2.5% DMSO. The reactions are performed for ~2 hours, then stopped by the addition of 10 µl of 0.75 µg/ml anti-phospho Ser/Thr antibody (Cell Signaling), 10 µg/ml Protein A AlphaScreen beads (Perkin Elmer), and 10 µg/ml streptavidin coated AlphaScreen beads in stop/detection buffer (50 mM EDTA, 95 mM Tris, pH=7.5, 0.01% Tween-20). The stopped reactions are incubated overnight in the dark. The phosphorylated peptide is detected via an oxygen anion initiated chemiluminescence/fluorescence cascade using the Envision plate reader (Perkin Elmer).

AlphaScreen Assay Conditions

Enzyme source	Enzyme conc. (nM)	b-BAD peptide conc. (nM)	ATP conc. (uM)	ATP Km (app) (uM)
Pim 1 (INV)	0.0025	50	2800	246
Pim 2 (INV)	0.01	50	500	4
Pim 3 (NVS)	0.005	50	2500	50

Compounds of the foregoing examples were tested by the Pim 1, Pim 2 & Pim 3 AlphaScreen assays and found to exhibit an IC₅₀ values as shown in Table 4 below. IC₅₀, the half maximal inhibitory concentration, represents the concentration of test compound that is required for 50% inhibition of its target in vitro.

Cell Proliferation Assay

KMS11 (human myeloma cell line), were cultured in IMDM supplemented with 10% FBS, sodium pyruvate and antibiotics. Cells were plated in the same medium at a density of 2000 cells per well into 96 well tissue culture plates, with outside wells vacant, on the day of assay.

Test compounds supplied in DMSO were diluted into DMSO at 500 times the desired final concentrations before

dilution into culture media to 2 times final concentrations. Equal volumes of 2× compounds were added to the cells in 96 well plates and incubated at 37° C. for 3 days.

After 3 days plates were equilibrated to room temperature and equal volume of CellTiter-Glow Reagent (Promega) was added to the culture wells. The plates were agitated briefly and luminescent signal was measured with luminometer. The percent inhibition of the signal seen in cells treated with DMSO alone vs. cells treated with control compound was calculated and used to determine EC₅₀ values (i.e., the concentration of a test compound that is required to obtain 50% of the maximum effect in the cells) for tested compounds, as shown in Table 4.

Using the procedures of the Pim1, Pim2, Pim3 AlphaScreen Assays the IC₅₀ concentrations of compounds of the previous examples were determined as shown in the Table 4.

Using the procedures of Cell Proliferation Assay, the EC₅₀ concentrations of compounds of the examples were determined in KMS11 cells as shown in Table 4.

TABLE 4

Ex#	Pim1 IC50 µM	Pim2 IC50 µM	Pim3 IC50 µM	KMS11 EC50 µM
1	0.00006	0.00253	0.00252	0.094
2	0.00036	0.03007	0.01572	1.008
3	0.00014	0.04942	0.03053	1.130
4	0.00006	0.00428	0.00154	0.148
5	0.00003	0.00124	0.00064	0.033
6	0.00024	0.00681	0.01100	3.090
7	0.00004	0.00139	0.00086	0.040
8	0.00035	0.02050	0.01423	0.499
9	0.00040	0.05937	0.01778	0.670
10	0.00057	0.03407	0.02088	1.833
11	0.00038	0.03331	0.01956	4.179
12	0.00021	0.01607	0.01572	0.716
13	0.00009	0.00482	0.00509	0.507
14	0.00004	0.00172	0.00102	0.045
15	0.00007	0.00356	0.00207	0.146
16	0.17209	21.7	8.3	>10
17	0.00256	0.38852	0.15521	8.260
18	0.00005	0.00086	0.00099	0.267
19	0.00017	0.00259	0.00321	0.287
20	0.00014	0.00668	0.00678	0.476
21	0.00005	0.00766	0.00092	0.756
22	0.00037	0.01388	0.01165	0.530
23	0.02746	1.4	1.0	>10
24	0.06819	2.7	1.7	>10
25	0.00063	0.02504	0.01580	2.867
26	0.14276	7.1	4.8	>10
27	0.00212	0.04198	0.05333	1.751
28	0.00680	0.23750	0.15437	4.874
29	0.00020	0.00944	0.00888	0.768
30	0.00072	0.07280	0.04426	9.686
31	0.00021	0.01045	0.01240	0.633
32	0.00297	0.10611	0.06386	7.958
33	0.00002	0.00068	0.00061	0.072
34	0.00076	0.04724	0.03991	3.753
35	0.00014	0.00794	0.00524	0.365
36	0.00059	0.02943	0.02876	0.979

269

TABLE 4-continued

Ex#	Pim1 IC50 μ M	Pim2 IC50 μ M	Pim3 IC50 μ M	KMS11 EC50 μ M
37	0.00009	0.00300	0.00503	0.178
38	0.00056	0.04551	0.04371	5.949
39	0.00039	0.03092	0.02182	2.989
40	0.00017	0.01241	0.00875	0.964
41		0.02032		5.980
42	0.00004	0.00196	0.00100	0.063
43	0.00586	0.32741	0.11823	4.577
44	0.00338	0.28911	0.22295	>10
45	0.64280	>25	>25	>10
46	0.00020	0.02139	0.00304	0.348
47	0.17919	15.8	2.7	>10
48	0.01063	0.42676	0.41111	9.416
49	0.00003	0.00120	0.00116	0.031
50	0.00024	0.01707	0.00535	0.470
51	0.00177	0.06911	0.03579	0.970
52	0.00850	0.72623	0.59094	4.386
53	0.00013	0.01432	0.01054	0.363
54	0.00009	0.00468	0.00174	0.891
55	0.00013	0.01118	0.00629	0.553
56	0.00002	0.00151	0.00062	0.036
57	0.00017	0.00647	0.00571	0.462
58	0.00002	0.00151	0.00049	0.168
59	0.01678	0.86628	0.58283	7.309
60	0.00022	0.01356	0.01080	0.202
61	0.00003	0.00107	0.00049	0.041
62	0.00003	0.00290	0.00128	0.106
63	0.00630	0.36441	0.09475	4.486
64	0.00001	0.00100	0.00022	0.044
65	0.04056	2.8	1.1	>10
66	0.00003	0.00118	0.00073	0.224
67	0.00148	0.07360	0.07771	3.205
68	0.00028	0.01196	0.01485	0.847
69	0.00004	0.00117	0.00040	0.114
70	0.00075	0.01070	0.00819	0.144
71	0.00004	0.00170	0.00054	0.156
72	0.00003	0.00083	0.00043	0.065
73	0.00002	0.00137	0.00091	0.069
74	0.04346	2.48310	1.50872	7.843
75	0.00008	0.00337	0.00308	0.147
76	0.13046	6.6	4.3	>10
77	0.00004	0.00144	0.00162	0.080
78	0.02538	1.1	1.0	>10
79	0.00028	0.00257	0.00536	0.132
80	0.00003	0.00082	0.00079	0.050
81	0.00004	0.00089	0.00074	0.036
82	0.00003	0.00062	0.00032	0.048
83	0.00001	0.00063	0.00064	0.037
84	0.00133	0.06413	0.04181	3.201
85	0.00003	0.00113	0.00096	0.057
86	0.00308	0.09956	0.06764	5.155
87	0.00003	0.00074	0.00068	0.166
88	0.00003	0.00237	0.00124	0.093
89	0.01317	1.47124	0.58270	3.990
90	0.03105	1.9	1.0	>10
91	0.00005	0.00226	0.00264	0.128
92	0.00010	0.00471	0.00369	0.175
93	0.00004	0.00368	0.00188	0.154
94	0.00003	0.00235	0.00279	0.356
95	0.00002	0.00138	0.00143	0.289
96	0.00004	0.00242	0.00507	0.186
97	0.00004	0.00133	0.00220	0.051
98	0.00004	0.00172	0.00158	0.074
99	0.00001	0.00108	0.00125	0.051
100	0.00006	0.00330	0.00373	0.165
101	0.00002	0.00301	0.00145	0.212
102	0.00003	0.00218	0.00210	0.454
103	0.00212	0.13033	0.10610	3.451
104	0.00535	0.13520	0.19792	>10
105	0.00006	0.00374	0.00161	0.117
106	0.00031	0.02440	0.01673	0.792
107	0.00005	0.00769	0.00542	0.364
108	0.00008	0.00234	0.00581	0.155
109	0.00040	0.00161	0.01117	0.616
110	0.00009	0.01038	0.01791	0.825
111	0.00008	0.00074	0.00141	0.169
112	0.00008	0.00737	0.00415	0.258
113	0.00250	0.12003	0.11490	1.826

270

TABLE 4-continued

Ex#	Pim1 IC50 μ M	Pim2 IC50 μ M	Pim3 IC50 μ M	KMS11 EC50 μ M
114			0.00188	0.053
115	0.00007	0.00248	0.00144	0.315
116	0.00012	0.00578	0.01210	0.323
117	0.00003	0.00153	0.00310	0.026
118	0.00002	0.00124	0.00174	2.455
119	0.00007	0.00907	0.00525	0.316
120	0.00140	0.14659	0.06752	6.060
121	0.00005	0.00593	0.01043	0.625
122	0.00005	0.00385	0.00384	0.321
123	0.00003	0.00153	0.00153	0.054
124	0.00005	0.00403	0.00202	0.040
125	0.00004	0.00179	0.00239	0.046
126	0.00070	0.00256	0.00778	0.267
127	0.00005	0.00198	0.00451	0.076
128	0.00024	0.00121	0.00411	2.099
129	0.00005	0.00246	0.00544	0.107
130	0.00003	0.00254	0.00170	0.109
131	0.00005	0.00255	0.00149	0.135
132	0.00004	0.00145	0.00108	0.016
133	0.00057	0.02866	0.01050	2.304
134	0.00054	0.02207	0.01019	0.468
135	0.00002	0.00078	0.00026	0.193
136	0.00004	0.00147	0.00249	0.038
137	0.00002	0.00103	0.00093	0.027
138	0.00003	0.00122	0.00258	0.020
139	0.00002	0.00124	0.00155	0.083
140	0.00002	0.00269	0.00134	0.100
141	0.00002	0.00117	0.00124	0.047
142	0.00002	0.00079	0.00087	0.078
143	0.00005	0.00302	0.00217	0.118
144	0.00002	0.00186	0.00123	0.056
145	0.00004	0.00090	0.00089	0.057
146	0.00002	0.00047	0.00039	0.018
147	0.00003	0.00169	0.00103	0.089
148	0.00002	0.00079	0.00055	0.091
149	0.00008	0.00888	0.00797	0.231
150	0.38044	>25	>25	>10
151	0.00004	0.00204	0.00218	0.113
152	0.05163	6.8	4.6	>10
153	0.00001	0.00074	0.00145	0.026
154	0.00005	0.00104	0.00074	0.257
155	0.00006	0.00220	0.00481	0.114
156	0.00004	0.00077	0.00080	0.178
157	0.00004	0.00120	0.00150	0.057
158	0.00004	0.00081	0.00132	0.040
159	0.00002	0.00109	0.00075	0.046
160	0.00005	0.00190	0.00222	0.082
161	0.00002	0.00087	0.00076	0.068
162		0.00168		0.059
163		0.00183		0.053
164		0.27024		>10
165	0.00002	0.00113	0.00095	0.066
166	0.00002	0.00040	0.00036	0.015
167	0.00018	0.00607	0.01257	0.240
168	0.00004	0.00531	0.00195	0.099
169	0.00079	0.06566	0.02227	2.363
170	0.00012	0.01384	0.00986	0.352
171	0.00002	0.01018	0.00251	0.398
172	0.00004	0.00169	0.00272	0.086
173	0.00005	0.00267	0.00132	0.136
174	0.00002	0.00065	0.00041	0.014
175	0.00009	0.00215	0.00252	0.091
176	0.00002	0.00077	0.00084	0.056
177	0.00004	0.00189	0.00296	0.069
178	0.00016	0.01494	0.02525	2.190
179	0.00003	0.00319	0.00516	0.266
180	0.00013	0.00747	0.00668	0.206
181	0.00003	0.00210	0.00175	0.108
182	0.00030	0.03743	0.02985	1.045
183	0.00001	0.00308	0.00075	0.058
184	0.00002	0.00204	0.00090	0.032
185		0.01333		0.417
186	0.00005	0.00405	0.00287	0.291

271

TABLE 4-continued

Ex#	Pim1 IC50 μ M	Pim2 IC50 μ M	Pim3 IC50 μ M	KMS11 EC50 μ M
187	0.00004	0.00279	0.00203	0.163
188		0.00527		0.403
189	0.03192	4.4	1.3	>10
190	0.00087	0.04726	0.04313	>10
191	0.00019	0.02718	0.00971	0.747
192	0.00004	0.00442	0.00210	0.194
193	0.00003	0.00179	0.00062	0.092
194	0.00005	0.00154	0.00117	0.039
195	0.00010	0.01374	0.01423	0.528
196	0.00031	0.00260	0.00690	0.969
197	0.00007	0.00515	0.00209	0.400
198	0.00028	0.00218	0.00305	0.265
199	0.00005	0.00281	0.00356	0.098
200	0.00170	0.00519	0.01829	2.415
201	0.00017	0.02347	0.02474	0.424
202	0.00038	0.01614	0.01967	0.358
203	0.00005	0.00167	0.00152	0.059

Compound structures in the tables marked as "Chiral" were prepared and tested in optically active form, having the absolute stereochemistry as shown; other compounds were prepared and tested in racemic form, and the depicted structure represents the relative stereochemistry at each chiral center.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 1

<210> SEQ ID NO 1

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: artificial sequence

<220> FEATURE:


<223> OTHER INFORMATION: Bad Peptide - biotinylated

<400> SEQUENCE: 1

Ala Gly Ala Gly Arg Ser Arg His Ser Ser Tyr Pro Ala Gly Thr
1 5 10 15

The invention claimed is:

1. A compound of Formula (A)



wherein:

groups attached to the ring containing Q that are depicted inside the ring are all syn to each other, and all groups attached to that ring that are depicted outside the ring are syn to one another;

Q is C;

272

R^q is H;

R¹ is selected from C₁₋₄ alkyl, C₃₋₆ cycloalkyl, and C₄₋₆ heterocyclyl, each alkyl, cycloalkyl and heterocyclyl is optionally substituted with up to two groups selected from halo, CN, NH₂, hydroxy, oxo, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, and C₁₋₄ haloalkoxy;

R^{1c} is halo, and R^{1d} is halo and R^{1b} is selected from H, halo, OR', R', —(CH₂)₁₋₂OR', and CONR'₂, where each R' is independently H or C₁₋₄ alkyl;

R^{2a} is —OMe, —SO₂Me, —NHCOOMe, or —O(CH₂)₂—X, wherein X is —OMe, COOH, CN or —SO₂Me, or R^{2a} is 1,2,3-triazolyl or 1-pyridonyl that is optionally substituted with up to two groups selected from halo, C₁₋₄ alkyl, hydroxy-substituted C₁₋₄ alkyl, C₂₋₄ alkenyl, C₃₋₆ cycloalkyl, and C₁₋₄ alkoxy, or —SO₂Me and R^{2b} is H;

or R^{2a} and R^{2b} taken together may form a dialkyl ketal or 5-6 membered cyclic ketal, =O or =N—OR", where R" is H or C₁₋₄ alkyl;

ring A is selected from pyridinyl, pyrimidinyl, pyrazinyl, and thiazolyl, each having N positioned as shown in Formula (I); and


Ring A is optionally substituted with 1 or 2 groups selected from halo, CN, NH₂, hydroxy, C₁₋₄ alkyl, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, and C₁₋₄ haloalkoxy;

or a pharmaceutically acceptable salt thereof.

2. The compound of claim 1, wherein R^{1d} is F.

3. The compound of claim 1, wherein R^{1c} is F.

4. The compound of claim 1, which is a compound of Formula (I):



N-4-(((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-(((3R,4S,5S)-3-amino-5-methyl-4-(4-(thiophen-3-yl)-1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-(((1R,3R,5S,Z)-3-amino-4-(hydroxyimino)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-(((1R,3R,5S)-3-amino-5-methyl-4-oxocyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-(((1R,3R,5S,E)-3-amino-4-(methoxyimino)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-(((1R,3R,5S,Z)-3-amino-4-(methoxyimino)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-(((3R,4S,5S)-3-amino-4-(4-(methoxymethyl)-1H-1,2,3-triazol-1-yl)-5-methylpiperidin-1-yl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-(((1R,3R,4S,5S)-3-amino-4-(2-hydroxyethylsulfonyl)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-(((1S,3S,4S,5R)-3-amino-5-methyl-4-(methylsulfonamido)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-(((1R,3R,4R,5 S)-3-amino-5-methyl-4-(methylsulfonamido)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-(((1R,3R,4S,5 S)-3-amino-4-((2-methoxyethyl)sulfonyl)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-(((1S,3S,4R,5R)-3-amino-5-methyl-4-(methylsulfonamido)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-(((1R,3R,4S,5S)-3-amino-5-methyl-4-(methylsulfonamido)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-(((1S,3S,4R,5R)-3-amino-4-azido-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-(((1R,3R,4S,5 S)-3-amino-4-azido-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-(((1R,3R,4R,5 S)-3-amino-5-methyl-4-((S)-methylsulfinyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-(((1R,3R,4R,5 S)-3-amino-5-methyl-4-((S)-methylsulfinyl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-(((1R,3R,4R,5S)-3-amino-5-

[illegible]

N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2-fluoro-5-(isopropylcarbamoyl)phenyl)picolinamide, N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide, N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-5-fluoropicolinamide, N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide, N-(4-((3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-5-fluoropicolinamide, N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide, N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide, 5-amino-N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide, N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(4-(ethoxymethyl)-2,6-difluorophenyl)-5-fluoropicolinamide, N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-5-fluoropicolinamide, N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinamide, N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinamide, N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(tetrahydro-2H-pyran-4-yl)oxy)phenyl)-5-fluoropicolinamide, N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-isopropoxyphenyl)-5-fluoropicolinamide, N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-isopropoxyphenyl)-5-fluoropicolinamide, N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinamide, N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinamide, 3-amino-N-(4-((1R,3R,4R,5S)-3-amino-4-methoxy-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, 3-amino-N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-(methylsulfonyl)ethoxy)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-isopropoxyphenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate, N-(4-((1S,3S,4R,5R)-3-amino-5-methyl-4-(2-oxopyridin-1(2H)-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-(4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(2-oxopyridin-1(2H)-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropi-

279

colinamide, 5-amino-N-4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide, N-4-((1S,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(4-(ethoxymethyl)-2,6-difluorophenyl)-5-fluoropicolinamide, N-4-((1R,3R,4S,5S)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(oxetan-3-yl)phenyl)-5-fluoropicolinamide, N-4-((1R,3R,4S,5S)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinamide, 3-amino-N-4-((1R,3R,4S,5S)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide N-4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(2-methoxyethoxy)phenyl)-5-fluoropicolinamide, N-4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-2-(2,6-difluorophenyl)thiazole-4-carboxamide, N-4-((1R,3R,4S,5 S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(tetrahydro-2H-pyran-4-yloxy)phenyl)-5-fluoropicolinamide, N-4-((1R,3R,4S,5 S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinamide, 3-amino-N-4-((1R,3R,4S,5 S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide, N-4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamide, N-4-((1R,3R,4S,5S)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-((tetrahydro-2H-pyran-4-yl)oxy)phenyl)-5-fluoropicolinamide, N-4-((1R,3R,4S,5 S)-3-amino-4-(2-cyanoethoxy)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinamide, N-4-((1R,3R,4S,5 S)-3-amino-4-(4-cyano-1H-1,2,3-triazol-1-yl)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-((1R,3R,4S,5 S)-3-amino-4-(4-cyclopropyl-1H-1,2,3-triazol-1-yl)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-((1R,3R,4S,5 S)-3-amino-5-methyl-4-(4-(prop-1-en-2-yl)-1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-((1R,3R,4S,5S)-3-amino-4-(4-(2-hydroxypropan-2-yl)-1H-1,2,3-triazol-1-yl)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, 2-((1R,2S,4S,6R)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyloxy)acetic acid, 2-((1 S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyloxy)acetic acid, N-4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinamide N-4-((1R,3R,

280

4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamide, 3-amino-N-4-((1R,3R,4S,5S)-3-amino-5-methyl-4-(1H-1,2,3-triazol-1-yl)cyclohexyl)pyridin-3-yl)-6-(2-fluoro-5-(isopropylcarbonyl)phenyl)picolinamide, N-4-((1R,3R,4S,5 S)-3-amino-4-(2-methoxyacetamido)-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-((1R,3R,4S,5S)-3-amino-4-isobutyramido-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, ethyl (1 S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate, isopropyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate, N-4-((1R,3R,4S,5S)-3-amino-5-methyl-4-propionamidocyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate, methyl (1R,2S,4S,6R)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate, N-4-((1S,3S,4R,5R)-4-acetamido-3-amino-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-((1R,3R,4S,5S)-4-acetamido-3-amino-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, N-4-((1R,3R,4R,5 S)-4-acetamido-3-amino-5-methylcyclohexyl)pyridin-3-yl)-6-(2,6-difluorophenyl)-5-fluoropicolinamide, methyl (1R,2S,4S,6R)-2-amino-4-(3-(5-amino-2-(2,6-difluorophenyl)thiazole-4-carboxamido)pyridin-4-yl)-6-methylcyclohexyl (methyl)carbamate, methyl (1S,2R,4R,6S)-2-amino-4-(3-(5-amino-2-(2,6-difluorophenyl)thiazole-4-carboxamido)pyridin-4-yl)-6-methylcyclohexyl (methyl)carbamate, methyl (1R,2S,4S,6R)-2-amino-4-(3-(6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl (methyl)carbamate, methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl (methyl)carbamate, methyl (3R,4S,5S)-3-amino-1-(3-(6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamido)pyridin-4-yl)-5-methylpiperidin-4-yl(methyl)carbamate, methyl (1R,2S,4S,6R)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl(methyl)carbamate, methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexyl(methyl)carbamate, methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-isopropoxyphenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate, methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-(1-hydroxycyclobutyl)phenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate, methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-(tetrahydro-2H-pyran-4-yloxy)phenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate, methyl (1S,2R,4R,6S)-2-

281

amino-4-(3-(3-amino-6-(2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate, methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-(4-hydroxytetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate, methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-methylphenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate, methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-(oxetan-3-yl)phenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate, methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(2,6-difluoro-4-methoxyphenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate, methyl (1S,2R,4R,6S)-2-amino-4-(3-(3-amino-6-(2-fluoro-5-(isopropylcarbamoyl)phenyl)picolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate, methyl (1S,2R,4R,

282

6S)-2-amino-4-(3-(6-(2,6-difluoro-4-(4-fluorotetrahydro-2H-pyran-4-yl)phenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate, methyl (1S,2R,4R,6S)-2-amino-4-(3-(6-(4-(ethoxymethyl)-2,6-difluorophenyl)-5-fluoropicolinamido)pyridin-4-yl)-6-methylcyclohexylcarbamate, and the pharmaceutically acceptable salts thereof.

8. A pharmaceutical composition comprising a compound of claim **1**, admixed with at least one pharmaceutically acceptable excipient.

9. The pharmaceutical composition of claim **8** which further comprises an additional therapeutic agent selected from MEK inhibitors, irinotecan, topotecan, gemcitabine, 5-fluorouracil, cytarabine, daunorubicin, PI3 Kinase inhibitors, mTOR inhibitors, DNA synthesis inhibitors, leucovorin, carboplatin, cisplatin, taxanes, tezacitabine, cyclophosphamide, vinca alkaloids, imatinib, anthracyclines, rituximab, lenalidomide, bortezomib and trastuzumab.

* * * * *