



(51) International Patent Classification:
C12N 15/82 (2006.01)

(21) International Application Number:
PCT/EP2010/064095

(22) International Filing Date:
24 September 2010 (24.09.2010)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09171385.9 25 September 2009 (25.09.2009) EP
09171364.4 25 September 2009 (25.09.2009) EP
61/252,220 16 October 2009 (16.10.2009) US
61/252,215 16 October 2009 (16.10.2009) US
61/260,942 13 November 2009 (13.11.2009) US
09175989.4 13 November 2009 (13.11.2009) EP

(71) Applicants (for all designated States except US): **BASF PLANT SCIENCE COMPANY GMBH** [DE/DE]; 67056 Ludwigshafen (DE). **CROP FUNCTIONAL GENOMICS CENTER** [KR/KR]; San 56-1, Shillim 9-Dong, Gawanak-Gu, Seoul 151-921 (KR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CHOI, Yang Do** [KR/KR]; San-56-1, Shilim 9-Dong, Seoul 151-921 (KR). **CHOI, Doil** [KR/KR]; 2-908, Samick APT, Dogok-dong, Gangnam-gu, Seoul 137-857 (KR). **REUZEAU, Christophe** [FR/FR]; rue de Cimetière, 24350 La Chapelle Gonaguet (FR). **SONG, Ji-Young** [KR/KR]; Jijok 919, Daejeon 305-770 (KR). **PARK, Youn-II** [KR/KR]; Jijok-dong 919, Youseong-gu, Daejeon 305-770 (KR).

(74) Agent: **MISTRY, Meeta**; Basf Se, Gvx - C 006, 67056 Ludwigshafen (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: PLANTS HAVING ENHANCED YIELD-RELATED TRAITS AND A METHOD FOR MAKING THE SAME

(57) Abstract: The present invention relates generally to the field of molecular biology and concerns a method for enhancing yield-related traits in plants by modulating expression in a plant of a nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide. The present invention also concerns plants having modulated expression of a nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, which plants have enhanced yield-related traits relative to corresponding wild type plants or other control plants. The invention also provides constructs useful in the methods of the invention.



Plants having enhanced yield-related traits and a method for making the same

The present invention relates generally to the field of molecular biology and concerns a method for enhancing yield-related traits in plants by modulating expression in a plant of a nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide. The present invention also concerns plants having modulated expression of a nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, which plants have enhanced yield-related traits relative to corresponding wild type plants or other control plants. The invention also provides constructs useful in the methods of the invention.

The ever-increasing world population and the dwindling supply of arable land available for agriculture fuels research towards increasing the efficiency of agriculture. Conventional means for crop and horticultural improvements utilise selective breeding techniques to identify plants having desirable characteristics. However, such selective breeding techniques have several drawbacks, namely that these techniques are typically labour intensive and result in plants that often contain heterogeneous genetic components that may not always result in the desirable trait being passed on from parent plants. Advances in molecular biology have allowed mankind to modify the germplasm of animals and plants. Genetic engineering of plants entails the isolation and manipulation of genetic material (typically in the form of DNA or RNA) and the subsequent introduction of that genetic material into a plant. Such technology has the capacity to deliver crops or plants having various improved economic, agronomic or horticultural traits.

A trait of particular economic interest is increased yield. Yield is normally defined as the measurable produce of economic value from a crop. This may be defined in terms of quantity and/or quality. Yield is directly dependent on several factors, for example, the number and size of the organs, plant architecture (for example, the number of branches), seed production, leaf senescence and more. Root development, nutrient uptake, stress tolerance and early vigour may also be important factors in determining yield. Optimizing the abovementioned factors may therefore contribute to increasing crop yield.

Seed yield is a particularly important trait, since the seeds of many plants are important for human and animal nutrition. Crops such as corn, rice, wheat, canola and soybean account for over half the total human caloric intake, whether through direct consumption of the seeds themselves or through consumption of meat products raised on processed seeds. They are also a source of sugars, oils and many kinds of metabolites used in industrial processes. Seeds contain an embryo (the source of new shoots and roots) and an endosperm (the source of nutrients for embryo growth during germination and during early growth of seedlings). The development of a seed involves many genes, and requires the transfer of metabolites from the roots, leaves and stems into the growing seed. The

endosperm, in particular, assimilates the metabolic precursors of carbohydrates, oils and proteins and synthesizes them into storage macromolecules to fill out the grain.

Another important trait for many crops is early vigour. Improving early vigour is an important objective of modern rice breeding programs in both temperate and tropical rice cultivars. Long roots are important for proper soil anchorage in water-seeded rice. Where rice is sown directly into flooded fields, and where plants must emerge rapidly through water, longer shoots are associated with vigour. Where drill-seeding is practiced, longer mesocotyls and coleoptiles are important for good seedling emergence. The ability to engineer early vigour into plants would be of great importance in agriculture. For example, poor early vigour has been a limitation to the introduction of maize (*Zea mays* L.) hybrids based on Corn Belt germplasm in the European Atlantic.

A further important trait is that of improved abiotic stress tolerance. Abiotic stress is a primary cause of crop loss worldwide, reducing average yields for most major crop plants by more than 50% (Wang et al., *Planta* 218, 1-14, 2003). Abiotic stresses may be caused by drought, salinity, extremes of temperature, chemical toxicity and oxidative stress. The ability to improve plant tolerance to abiotic stress would be of great economic advantage to farmers worldwide and would allow for the cultivation of crops during adverse conditions and in territories where cultivation of crops may not otherwise be possible.

Crop yield may therefore be increased by optimising one of the above-mentioned factors.

Depending on the end use, the modification of certain yield traits may be favoured over others. For example for applications such as forage or wood production, or bio-fuel resource, an increase in the vegetative parts of a plant may be desirable, and for applications such as flour, starch or oil production, an increase in seed parameters may be particularly desirable. Even amongst the seed parameters, some may be favoured over others, depending on the application. Various mechanisms may contribute to increasing seed yield, whether that is in the form of increased seed size or increased seed number.

One approach to increasing yield (seed yield and/or biomass) in plants may be through modification of the inherent growth mechanisms of a plant, such as the cell cycle or various signalling pathways involved in plant growth or in defence mechanisms.

It has now been found that various yield-related traits may be improved in plants by modulating expression in a plant of a nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, in a plant.

Background

1. SGT1 polypeptides

SGT1 is known as a suppressor allele of *skp1* mutant.

Chung et al., 2006 report that SGT1 plays a crucial role in developmental processes. SGT1 has unique domains necessary for protein functions: tetratricopeptide repeat domain (TPR), CHORD and SGT1 motif (CS) and SGT1-specific motif (SGS motif). The TPR domain has been known to mediate protein–protein interactions among multicomplex proteins functioning as chaperone, cell cycle, transcription, or protein transport complexes. For example, the TPR domain of SGT1 was shown to bind to heat-shock protein 70 (HSP70). However, the CS domain of SGT1 is similar to the one in the human p23 protein, which is known to interact with HSP90 and participate in the folding of different regulatory proteins.

2. CLC-pKG polypeptides

In both prokaryotic and eukaryotic organism, anion channels/transporters appear as key players in the control of metabolism, in the maintenance of electrochemical gradients and in signalling pathways leading to adaptation to abiotic and biotic environmental stresses. In plants, they contribute to various physiological functions such as control of stomatal movements regulating gas exchanges in leaves, plant–pathogen interaction, root xylem loading, compartmentation of metabolites and coupling with proton gradients (reviewed in De Angeli et al. 2007 *Phil. Trans. R. Soc. B* 2009 364, 195-201). Anion channel activities and associated regulation mechanisms have been characterized primarily using electrophysiological techniques. They were reported in all plant membranes including the plasma membrane, tonoplast, endoplasmic reticulum, mitochondria and chloroplasts, plasma membrane channels being by far the best characterized compared to those located on other membranes. In model plants such as rice and *Arabidopsis* there are up to seven genes encoding CLCs which are spread in over two distinct subfamilies (Marmagne et al. 2007, *Journal of Experimental Botany*, Vol. 58, No. 12, pp. 3385–3393). One of such subfamilies comprising the AtCLCe and AtCLCf proteins is close to the prokaryotic CLC, altogether belonging to the Chloride Channel Prokaryotic Group while the other class is closer to the eukaryotic CLCs. The *Arabidopsis thaliana* AtCLCf protein is reported to have similar subcellular distribution and presumably similar function as those CLCs of *Synechocystis* CLC, considered to represent the ancestor precursor of plant chloroplast.

Nitrate and malate represent the majority of anions in a plant cell. Nitrate is a nutrient but can act as a signalling molecule as well. Plants have a sophisticated nitrate uptake system involving both low- and high-affinity transporters, nitrate is next either transported through the xylem to enter into the cellular metabolism or is stored locally. Cells assimilate nitrate via the nitrate reductase pathway or store it in the tonoplast, and a dynamic balance exists between cytosolic and vacuolar nitrate levels, regulated by uptake of extracellular nitrate, storage in the vacuole and anabolism. The discovery of the chloride channel (CLC) family

allowed unravelling the mechanism of proton/nitrate exchange between tonoplasts and cytosol. Determination of subcellular localization, expression patterns, and characterization of knockout mutant phenotypes, gave insight in the physiological role of CLC proteins. Phenotypic analyses showed that *clca-1* and *clca-2* mutant plants have a reduced nitrate compared to that of wild-type in root and shoot tissues. Also *clcc* and *clce* mutants showed lower nitrate levels compared to control plants. An overview of the art is provided by De Angeli et al., *Phil. Trans. R. Soc. B* 364, 195-201, 2009; and the references cited therein. However, still little is known about the precise role of CLC proteins, and the effect of overexpressing CLC genes on plant phenotypes.

3. HD-hydrolase-like polypeptides

The HD domain comprises a sequence of two small amino acids, followed by two hydrophobic amino acids, a histidine and an aspartic acid, again a hydrophobic amino acid, a small amino acid and a charged amino acid. Because of its weak sequence conservation, it was only recently discovered (Aravind and Koonin, *Trends in Biochemical Sciences*, 23: 469-472, 1998). The domain is reportedly present in metal-dependent phosphohydrolases, including nucleic acid polymerases and helicases (Aravind and Koonin, 1998). Although the HD-domain comprising proteins are predicted to exhibit phosphohydrolase activity and appears to be involved in nucleic acid metabolism or in signal transduction (Galperin et al., *J. Mol. Microbiol Biotechnol.* 1, 303-305, 1999), the precise biological role of the HD domain remains to be elucidated.

Summary

1. SGT1 polypeptides

Surprisingly, it has now been found that modulating expression of a nucleic acid encoding an SGT1 polypeptide gives plants having enhanced yield-related traits, in particular increased seed yield relative to control plants.

According to one embodiment of the present invention, there is provided a method for improving yield-related traits in plants relative to control plants, comprising modulating expression in a plant of a nucleic acid encoding an SGT1 polypeptide.

2. CLC-pKG polypeptides

Surprisingly, it has now been found that modulating expression of a nucleic acid encoding a CLC-pKG polypeptide gives plants having enhanced yield-related traits relative to control plants.

According one embodiment, there is provided a method for enhancing or improving yield-related traits in plants relative to control plants, comprising modulating expression in a plant of a nucleic acid encoding a CLC-pKG polypeptide.

3. HD-hydrolase-like polypeptides

Surprisingly, it has now been found that modulating expression of a nucleic acid encoding a HD-hydrolase-like polypeptide gives plants having enhanced yield-related traits, in particular increased yield relative to control plants.

According one embodiment, there is provided a method for improving yield-related traits in plants relative to control plants, comprising modulating expression in a plant of a nucleic acid encoding a HD-hydrolase-like polypeptide.

Definitions

The following definitions will be used throughout the present specification.

Polypeptide(s)/Protein(s)

The terms “polypeptide” and “protein” are used interchangeably herein and refer to amino acids in a polymeric form of any length, linked together by peptide bonds.

Polynucleotide(s)/Nucleic acid(s)/Nucleic acid sequence(s)/nucleotide sequence(s)

The terms “polynucleotide(s)”, “nucleic acid sequence(s)”, “nucleotide sequence(s)”, “nucleic acid(s)”, “nucleic acid molecule” are used interchangeably herein and refer to nucleotides, either ribonucleotides or deoxyribonucleotides or a combination of both, in a polymeric unbranched form of any length.

Homologue(s)

“Homologues” of a protein encompass peptides, oligopeptides, polypeptides, proteins and enzymes having amino acid substitutions, deletions and/or insertions relative to the unmodified protein in question and having similar biological and functional activity as the unmodified protein from which they are derived.

A deletion refers to removal of one or more amino acids from a protein.

An insertion refers to one or more amino acid residues being introduced into a predetermined site in a protein. Insertions may comprise N-terminal and/or C-terminal fusions as well as intra-sequence insertions of single or multiple amino acids. Generally, insertions within the amino acid sequence will be smaller than N- or C-terminal fusions, of the order of about 1 to 10 residues. Examples of N- or C-terminal fusion proteins or peptides include the binding domain or activation domain of a transcriptional activator as used in the yeast two-hybrid system, phage coat proteins, (histidine)-6-tag, glutathione S-transferase-tag, protein A, maltose-binding protein, dihydrofolate reductase, Tag•100 epitope, c-myc epitope, FLAG®-epitope, lacZ, CMP (calmodulin-binding peptide), HA epitope, protein C epitope and VSV epitope.

A substitution refers to replacement of amino acids of the protein with other amino acids having similar properties (such as similar hydrophobicity, hydrophilicity, antigenicity, propensity to form or break α -helical structures or β -sheet structures). Amino acid substitutions are typically of single residues, but may be clustered depending upon functional constraints placed upon the polypeptide and may range from 1 to 10 amino acids; insertions will usually be of the order of about 1 to 10 amino acid residues. The amino acid substitutions are preferably conservative amino acid substitutions. Conservative substitution tables are well known in the art (see for example Creighton (1984) *Proteins*. W.H. Freeman and Company (Eds) and Table 1 below).

Table 1: Examples of conserved amino acid substitutions

Residue	Conservative Substitutions	Residue	Conservative Substitutions
Ala	Ser	Leu	Ile; Val
Arg	Lys	Lys	Arg; Gln
Asn	Gln; His	Met	Leu; Ile
Asp	Glu	Phe	Met; Leu; Tyr
Gln	Asn	Ser	Thr; Gly
Cys	Ser	Thr	Ser; Val
Glu	Asp	Trp	Tyr
Gly	Pro	Tyr	Trp; Phe
His	Asn; Gln	Val	Ile; Leu
Ile	Leu, Val		

Amino acid substitutions, deletions and/or insertions may readily be made using peptide synthetic techniques well known in the art, such as solid phase peptide synthesis and the like, or by recombinant DNA manipulation. Methods for the manipulation of DNA sequences to produce substitution, insertion or deletion variants of a protein are well known in the art. For example, techniques for making substitution mutations at predetermined sites in DNA are well known to those skilled in the art and include M13 mutagenesis, T7-Gen in vitro mutagenesis (USB, Cleveland, OH), QuickChange Site Directed mutagenesis (Stratagene, San Diego, CA), PCR-mediated site-directed mutagenesis or other site-directed mutagenesis protocols.

Derivatives

“Derivatives” include peptides, oligopeptides, polypeptides which may, compared to the amino acid sequence of the naturally-occurring form of the protein, such as the protein of interest, comprise substitutions of amino acids with non-naturally occurring amino acid residues, or additions of non-naturally occurring amino acid residues. “Derivatives” of a protein also encompass peptides, oligopeptides, polypeptides which comprise naturally occurring altered (glycosylated, acylated, prenylated, phosphorylated, myristoylated, sulphated etc.) or non-naturally altered amino acid residues compared to the amino acid sequence of a naturally-occurring form of the polypeptide. A derivative may also comprise

one or more non-amino acid substituents or additions compared to the amino acid sequence from which it is derived, for example a reporter molecule or other ligand, covalently or non-covalently bound to the amino acid sequence, such as a reporter molecule which is bound to facilitate its detection, and non-naturally occurring amino acid residues relative to the amino acid sequence of a naturally-occurring protein. Furthermore, "derivatives" also include fusions of the naturally-occurring form of the protein with tagging peptides such as FLAG, HIS6 or thioredoxin (for a review of tagging peptides, see Terpe, Appl. Microbiol. Biotechnol. 60, 523-533, 2003).

Orthologue(s)/Parologue(s)

Orthologues and paralogues encompass evolutionary concepts used to describe the ancestral relationships of genes. Paralogues are genes within the same species that have originated through duplication of an ancestral gene; orthologues are genes from different organisms that have originated through speciation, and are also derived from a common ancestral gene.

Domain, Motif/Consensus sequence/Signature

The term "domain" refers to a set of amino acids conserved at specific positions along an alignment of sequences of evolutionarily related proteins. While amino acids at other positions can vary between homologues, amino acids that are highly conserved at specific positions indicate amino acids that are likely essential in the structure, stability or function of a protein. Identified by their high degree of conservation in aligned sequences of a family of protein homologues, they can be used as identifiers to determine if any polypeptide in question belongs to a previously identified polypeptide family.

The term "motif" or "consensus sequence" or "signature" refers to a short conserved region in the sequence of evolutionarily related proteins. Motifs are frequently highly conserved parts of domains, but may also include only part of the domain, or be located outside of conserved domain (if all of the amino acids of the motif fall outside of a defined domain).

Specialist databases exist for the identification of domains, for example, SMART (Schultz et al. (1998) Proc. Natl. Acad. Sci. USA 95, 5857-5864; Letunic et al. (2002) Nucleic Acids Res 30, 242-244), InterPro (Mulder et al., (2003) Nucl. Acids. Res. 31, 315-318), Prosite (Bucher and Bairoch (1994), A generalized profile syntax for biomolecular sequences motifs and its function in automatic sequence interpretation. (In) ISMB-94; Proceedings 2nd International Conference on Intelligent Systems for Molecular Biology. Altman R., Brutlag D., Karp P., Lathrop R., Searls D., Eds., pp53-61, AAAI Press, Menlo Park; Hulo et al., Nucl. Acids. Res. 32:D134-D137, (2004)), or Pfam (Bateman et al., Nucleic Acids Research 30(1): 276-280 (2002)). A set of tools for in silico analysis of protein sequences is available on the ExPASy proteomics server (Swiss Institute of Bioinformatics (Gasteiger et al., ExPASy: the proteomics server for in-depth protein knowledge and analysis, Nucleic Acids

Res. 31:3784-3788(2003)). Domains or motifs may also be identified using routine techniques, such as by sequence alignment.

Methods for the alignment of sequences for comparison are well known in the art, such methods include GAP, BESTFIT, BLAST, FASTA and TFASTA. GAP uses the algorithm of Needleman and Wunsch ((1970) J Mol Biol 48: 443-453) to find the global (i.e. spanning the complete sequences) alignment of two sequences that maximizes the number of matches and minimizes the number of gaps. The BLAST algorithm (Altschul et al. (1990) J Mol Biol 215: 403-10) calculates percent sequence identity and performs a statistical analysis of the similarity between the two sequences. The software for performing BLAST analysis is publicly available through the National Centre for Biotechnology Information (NCBI). Homologues may readily be identified using, for example, the ClustalW multiple sequence alignment algorithm (version 1.83), with the default pairwise alignment parameters, and a scoring method in percentage. Global percentages of similarity and identity may also be determined using one of the methods available in the MatGAT software package (Campanella et al., BMC Bioinformatics. 2003 Jul 10;4:29. MatGAT: an application that generates similarity/identity matrices using protein or DNA sequences.). Minor manual editing may be performed to optimise alignment between conserved motifs, as would be apparent to a person skilled in the art. Furthermore, instead of using full-length sequences for the identification of homologues, specific domains may also be used. The sequence identity values may be determined over the entire nucleic acid or amino acid sequence or over selected domains or conserved motif(s), using the programs mentioned above using the default parameters. For local alignments, the Smith-Waterman algorithm is particularly useful (Smith TF, Waterman MS (1981) J. Mol. Biol 147(1);195-7).

Reciprocal BLAST

Typically, this involves a first BLAST involving BLASTing a query sequence (for example using any of the sequences listed in Table A of the Examples section) against any sequence database, such as the publicly available NCBI database. BLASTN or TBLASTX (using standard default values) are generally used when starting from a nucleotide sequence, and BLASTP or TBLASTN (using standard default values) when starting from a protein sequence. The BLAST results may optionally be filtered. The full-length sequences of either the filtered results or non-filtered results are then BLASTed back (second BLAST) against sequences from the organism from which the query sequence is derived. The results of the first and second BLASTs are then compared. A paralogue is identified if a high-ranking hit from the first blast is from the same species as from which the query sequence is derived, a BLAST back then ideally results in the query sequence amongst the highest hits; an orthologue is identified if a high-ranking hit in the first BLAST is not from the same species as from which the query sequence is derived, and preferably results upon BLAST back in the query sequence being among the highest hits.

High-ranking hits are those having a low E-value. The lower the E-value, the more significant the score (or in other words the lower the chance that the hit was found by chance). Computation of the E-value is well known in the art. In addition to E-values, comparisons are also scored by percentage identity. Percentage identity refers to the number of identical nucleotides (or amino acids) between the two compared nucleic acid (or polypeptide) sequences over a particular length. In the case of large families, ClustalW may be used, followed by a neighbour joining tree, to help visualize clustering of related genes and to identify orthologues and paralogues.

Hybridisation

The term "hybridisation" as defined herein is a process wherein substantially homologous complementary nucleotide sequences anneal to each other. The hybridisation process can occur entirely in solution, i.e. both complementary nucleic acids are in solution. The hybridisation process can also occur with one of the complementary nucleic acids immobilised to a matrix such as magnetic beads, Sepharose beads or any other resin. The hybridisation process can furthermore occur with one of the complementary nucleic acids immobilised to a solid support such as a nitro-cellulose or nylon membrane or immobilised by e.g. photolithography to, for example, a siliceous glass support (the latter known as nucleic acid arrays or microarrays or as nucleic acid chips). In order to allow hybridisation to occur, the nucleic acid molecules are generally thermally or chemically denatured to melt a double strand into two single strands and/or to remove hairpins or other secondary structures from single stranded nucleic acids.

The term "stringency" refers to the conditions under which a hybridisation takes place. The stringency of hybridisation is influenced by conditions such as temperature, salt concentration, ionic strength and hybridisation buffer composition. Generally, low stringency conditions are selected to be about 30°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. Medium stringency conditions are when the temperature is 20°C below T_m , and high stringency conditions are when the temperature is 10°C below T_m . High stringency hybridisation conditions are typically used for isolating hybridising sequences that have high sequence similarity to the target nucleic acid sequence. However, nucleic acids may deviate in sequence and still encode a substantially identical polypeptide, due to the degeneracy of the genetic code. Therefore medium stringency hybridisation conditions may sometimes be needed to identify such nucleic acid molecules.

The T_m is the temperature under defined ionic strength and pH, at which 50% of the target sequence hybridises to a perfectly matched probe. The T_m is dependent upon the solution conditions and the base composition and length of the probe. For example, longer sequences hybridise specifically at higher temperatures. The maximum rate of hybridisation is obtained from about 16°C up to 32°C below T_m . The presence of monovalent cations in the hybridisation solution reduce the electrostatic repulsion between

the two nucleic acid strands thereby promoting hybrid formation; this effect is visible for sodium concentrations of up to 0.4M (for higher concentrations, this effect may be ignored). Formamide reduces the melting temperature of DNA-DNA and DNA-RNA duplexes with 0.6 to 0.7°C for each percent formamide, and addition of 50% formamide allows hybridisation to be performed at 30 to 45°C, though the rate of hybridisation will be lowered. Base pair mismatches reduce the hybridisation rate and the thermal stability of the duplexes. On average and for large probes, the T_m decreases about 1°C per % base mismatch. The T_m may be calculated using the following equations, depending on the types of hybrids:

1) DNA-DNA hybrids (Meinkoth and Wahl, Anal. Biochem., 138: 267-284, 1984):

$$T_m = 81.5^\circ\text{C} + 16.6 \times \log_{10}[\text{Na}^+]^a + 0.41 \times \%[\text{G/C}^b] - 500 \times [\text{L}^c]^{-1} - 0.61 \times \% \text{ formamide}$$

2) DNA-RNA or RNA-RNA hybrids:

$$T_m = 79.8^\circ\text{C} + 18.5 (\log_{10}[\text{Na}^+]^a) + 0.58 (\% \text{G/C}^b) + 11.8 (\% \text{G/C}^b)^2 - 820/\text{L}^c$$

3) oligo-DNA or oligo-RNA^d hybrids:

$$\text{For } <20 \text{ nucleotides: } T_m = 2 (I_n)$$

$$\text{For } 20\text{--}35 \text{ nucleotides: } T_m = 22 + 1.46 (I_n)$$

^a or for other monovalent cation, but only accurate in the 0.01–0.4 M range.

^b only accurate for %GC in the 30% to 75% range.

^c L = length of duplex in base pairs.

^d oligo, oligonucleotide; I_n = effective length of primer = $2 \times (\text{no. of G/C}) + (\text{no. of A/T})$.

Non-specific binding may be controlled using any one of a number of known techniques such as, for example, blocking the membrane with protein containing solutions, additions of heterologous RNA, DNA, and SDS to the hybridisation buffer, and treatment with Rnase. For non-homologous probes, a series of hybridizations may be performed by varying one of (i) progressively lowering the annealing temperature (for example from 68°C to 42°C) or (ii) progressively lowering the formamide concentration (for example from 50% to 0%). The skilled artisan is aware of various parameters which may be altered during hybridisation and which will either maintain or change the stringency conditions.

Besides the hybridisation conditions, specificity of hybridisation typically also depends on the function of post-hybridisation washes. To remove background resulting from non-specific hybridisation, samples are washed with dilute salt solutions. Critical factors of such washes include the ionic strength and temperature of the final wash solution: the lower the salt concentration and the higher the wash temperature, the higher the stringency of the wash. Wash conditions are typically performed at or below hybridisation stringency. A positive hybridisation gives a signal that is at least twice of that of the background. Generally, suitable stringent conditions for nucleic acid hybridisation assays or gene amplification detection procedures are as set forth above. More or less stringent conditions may also be selected. The skilled artisan is aware of various parameters which may be altered during washing and which will either maintain or change the stringency conditions.

For example, typical high stringency hybridisation conditions for DNA hybrids longer than 50 nucleotides encompass hybridisation at 65°C in 1x SSC or at 42°C in 1x SSC and 50% formamide, followed by washing at 65°C in 0.3x SSC. Examples of medium stringency hybridisation conditions for DNA hybrids longer than 50 nucleotides encompass hybridisation at 50°C in 4x SSC or at 40°C in 6x SSC and 50% formamide, followed by washing at 50°C in 2x SSC. The length of the hybrid is the anticipated length for the hybridising nucleic acid. When nucleic acids of known sequence are hybridised, the hybrid length may be determined by aligning the sequences and identifying the conserved regions described herein. 1×SSC is 0.15M NaCl and 15mM sodium citrate; the hybridisation solution and wash solutions may additionally include 5x Denhardt's reagent, 0.5-1.0% SDS, 100 µg/ml denatured, fragmented salmon sperm DNA, 0.5% sodium pyrophosphate.

For the purposes of defining the level of stringency, reference can be made to Sambrook et al. (2001) *Molecular Cloning: a laboratory manual*, 3rd Edition, Cold Spring Harbor Laboratory Press, CSH, New York or to *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989 and yearly updates).

Splice variant

The term “splice variant” as used herein encompasses variants of a nucleic acid sequence in which selected introns and/or exons have been excised, replaced, displaced or added, or in which introns have been shortened or lengthened. Such variants will be ones in which the biological activity of the protein is substantially retained; this may be achieved by selectively retaining functional segments of the protein. Such splice variants may be found in nature or may be manmade. Methods for predicting and isolating such splice variants are well known in the art (see for example Foissac and Schiex (2005) *BMC Bioinformatics* 6: 25).

Allelic variant

Alleles or allelic variants are alternative forms of a given gene, located at the same chromosomal position. Allelic variants encompass Single Nucleotide Polymorphisms (SNPs), as well as Small Insertion/Deletion Polymorphisms (INDELs). The size of INDELs is usually less than 100 bp. SNPs and INDELs form the largest set of sequence variants in naturally occurring polymorphic strains of most organisms.

Endogenous gene

Reference herein to an “endogenous” gene not only refers to the gene in question as found in a plant in its natural form (i.e., without there being any human intervention), but also refers to that same gene (or a substantially homologous nucleic acid/gene) in an isolated form subsequently (re)introduced into a plant (a transgene). For example, a transgenic plant containing such a transgene may encounter a substantial reduction of the transgene expression and/or substantial reduction of expression of the endogenous gene. The

isolated gene may be isolated from an organism or may be manmade, for example by chemical synthesis.

Gene shuffling/Directed evolution

Gene shuffling or directed evolution consists of iterations of DNA shuffling followed by appropriate screening and/or selection to generate variants of nucleic acids or portions thereof encoding proteins having a modified biological activity (Castle et al., (2004) Science 304(5674): 1151-4; US patents 5,811,238 and 6,395,547).

Construct

Additional regulatory elements may include transcriptional as well as translational enhancers. Those skilled in the art will be aware of terminator and enhancer sequences that may be suitable for use in performing the invention. An intron sequence may also be added to the 5' untranslated region (UTR) or in the coding sequence to increase the amount of the mature message that accumulates in the cytosol, as described in the definitions section. Other control sequences (besides promoter, enhancer, silencer, intron sequences, 3'UTR and/or 5'UTR regions) may be protein and/or RNA stabilizing elements. Such sequences would be known or may readily be obtained by a person skilled in the art.

The genetic constructs of the invention may further include an origin of replication sequence that is required for maintenance and/or replication in a specific cell type. One example is when a genetic construct is required to be maintained in a bacterial cell as an episomal genetic element (e.g. plasmid or cosmid molecule). Preferred origins of replication include, but are not limited to, the f1-ori and colE1.

For the detection of the successful transfer of the nucleic acid sequences as used in the methods of the invention and/or selection of transgenic plants comprising these nucleic acids, it is advantageous to use marker genes (or reporter genes). Therefore, the genetic construct may optionally comprise a selectable marker gene. Selectable markers are described in more detail in the "definitions" section herein. The marker genes may be removed or excised from the transgenic cell once they are no longer needed. Techniques for marker removal are known in the art, useful techniques are described above in the definitions section.

Regulatory element/Control sequence/Promoter

The terms "regulatory element", "control sequence" and "promoter" are all used interchangeably herein and are to be taken in a broad context to refer to regulatory nucleic acid sequences capable of effecting expression of the sequences to which they are ligated. The term "promoter" typically refers to a nucleic acid control sequence located upstream from the transcriptional start of a gene and which is involved in recognising and binding of RNA polymerase and other proteins, thereby directing transcription of an operably linked nucleic acid. Encompassed by the aforementioned terms are transcriptional regulatory

sequences derived from a classical eukaryotic genomic gene (including the TATA box which is required for accurate transcription initiation, with or without a CCAAT box sequence) and additional regulatory elements (i.e. upstream activating sequences, enhancers and silencers) which alter gene expression in response to developmental and/or external stimuli, or in a tissue-specific manner. Also included within the term is a transcriptional regulatory sequence of a classical prokaryotic gene, in which case it may include a -35 box sequence and/or -10 box transcriptional regulatory sequences. The term "regulatory element" also encompasses a synthetic fusion molecule or derivative that confers, activates or enhances expression of a nucleic acid molecule in a cell, tissue or organ.

A "plant promoter" comprises regulatory elements, which mediate the expression of a coding sequence segment in plant cells. Accordingly, a plant promoter need not be of plant origin, but may originate from viruses or micro-organisms, for example from viruses which attack plant cells. The "plant promoter" can also originate from a plant cell, e.g. from the plant which is transformed with the nucleic acid sequence to be expressed in the inventive process and described herein. This also applies to other "plant" regulatory signals, such as "plant" terminators. The promoters upstream of the nucleotide sequences useful in the methods of the present invention can be modified by one or more nucleotide substitution(s), insertion(s) and/or deletion(s) without interfering with the functionality or activity of either the promoters, the open reading frame (ORF) or the 3'-regulatory region such as terminators or other 3' regulatory regions which are located away from the ORF. It is furthermore possible that the activity of the promoters is increased by modification of their sequence, or that they are replaced completely by more active promoters, even promoters from heterologous organisms. For expression in plants, the nucleic acid molecule must, as described above, be linked operably to or comprise a suitable promoter which expresses the gene at the right point in time and with the required spatial expression pattern.

For the identification of functionally equivalent promoters, the promoter strength and/or expression pattern of a candidate promoter may be analysed for example by operably linking the promoter to a reporter gene and assaying the expression level and pattern of the reporter gene in various tissues of the plant. Suitable well-known reporter genes include for example beta-glucuronidase or beta-galactosidase. The promoter activity is assayed by measuring the enzymatic activity of the beta-glucuronidase or beta-galactosidase. The promoter strength and/or expression pattern may then be compared to that of a reference promoter (such as the one used in the methods of the present invention). Alternatively, promoter strength may be assayed by quantifying mRNA levels or by comparing mRNA levels of the nucleic acid used in the methods of the present invention, with mRNA levels of housekeeping genes such as 18S rRNA, using methods known in the art, such as Northern blotting with densitometric analysis of autoradiograms, quantitative real-time PCR or RT-PCR (Heid et al., 1996 *Genome Methods* 6: 986-994). Generally by "weak promoter" is intended a promoter that drives expression of a coding sequence at a low level. By "low

level” is intended at levels of about 1/10,000 transcripts to about 1/100,000 transcripts, to about 1/500,000 transcripts per cell. Conversely, a “strong promoter” drives expression of a coding sequence at high level, or at about 1/10 transcripts to about 1/100 transcripts to about 1/1000 transcripts per cell. Generally, by “medium strength promoter” is intended a promoter that drives expression of a coding sequence at a lower level than a strong promoter, in particular at a level that is in all instances below that obtained when under the control of a 35S CaMV promoter.

Operably linked

The term “operably linked” as used herein refers to a functional linkage between the promoter sequence and the gene of interest, such that the promoter sequence is able to initiate transcription of the gene of interest.

Constitutive promoter

A “constitutive promoter” refers to a promoter that is transcriptionally active during most, but not necessarily all, phases of growth and development and under most environmental conditions, in at least one cell, tissue or organ. Table 2a below gives examples of constitutive promoters.

Table 2a: Examples of constitutive promoters

Gene Source	Reference
Actin	McElroy et al, Plant Cell, 2: 163-171, 1990
HMGP	WO 2004/070039
CAMV 35S	Odell et al, Nature, 313: 810-812, 1985
CaMV 19S	Nilsson et al., Physiol. Plant. 100:456-462, 1997
GOS2	de Pater et al, Plant J Nov;2(6):837-44, 1992, WO 2004/065596
Ubiquitin	Christensen et al, Plant Mol. Biol. 18: 675-689, 1992
Rice cyclophilin	Buchholz et al, Plant Mol Biol. 25(5): 837-43, 1994
Maize H3 histone	Lepetit et al, Mol. Gen. Genet. 231:276-285, 1992
Alfalfa H3 histone	Wu et al. Plant Mol. Biol. 11:641-649, 1988
Actin 2	An et al, Plant J. 10(1); 107-121, 1996
34S FMV	Sanger et al., Plant. Mol. Biol., 14, 1990: 433-443
Rubisco small subunit	US 4,962,028
OCS	Leisner (1988) Proc Natl Acad Sci USA 85(5): 2553
SAD1	Jain et al., Crop Science, 39 (6), 1999: 1696
SAD2	Jain et al., Crop Science, 39 (6), 1999: 1696
nos	Shaw et al. (1984) Nucleic Acids Res. 12(20):7831-7846
V-ATPase	WO 01/14572
Super promoter	WO 95/14098
G-box proteins	WO 94/12015

Ubiquitous promoter

A ubiquitous promoter is active in substantially all tissues or cells of an organism.

Developmentally-regulated promoter

A developmentally-regulated promoter is active during certain developmental stages or in parts of the plant that undergo developmental changes.

Inducible promoter

An inducible promoter has induced or increased transcription initiation in response to a chemical (for a review see Gatz 1997, Annu. Rev. Plant Physiol. Plant Mol. Biol., 48:89-108), environmental or physical stimulus, or may be “stress-inducible”, i.e. activated when a plant is exposed to various stress conditions, or a “pathogen-inducible” i.e. activated when a plant is exposed to exposure to various pathogens.

Organ-specific/Tissue-specific promoter

An organ-specific or tissue-specific promoter is one that is capable of preferentially initiating transcription in certain organs or tissues, such as the leaves, roots, seed tissue etc. For example, a “root-specific promoter” is a promoter that is transcriptionally active predominantly in plant roots, substantially to the exclusion of any other parts of a plant, whilst still allowing for any leaky expression in these other plant parts. Promoters able to initiate transcription in certain cells only are referred to herein as “cell-specific”.

Examples of root-specific promoters are listed in Table 2b below:

Table 2b: Examples of root-specific promoters

Gene Source	Reference
RCc3	Plant Mol Biol. 1995 Jan;27(2):237-48
Arabidopsis PHT1	Koyama et al. J Biosci Bioeng. 2005 Jan;99(1):38-42.; Mudge et al. (2002, Plant J. 31:341)
Medicago phosphate transporter	Xiao et al., 2006, Plant Biol (Stuttg). 2006 Jul;8(4):439-49
Arabidopsis Pyk10	Nitz et al. (2001) Plant Sci 161(2): 337-346
root-expressible genes	Tingey et al., EMBO J. 6: 1, 1987.
tobacco auxin-inducible gene	Van der Zaal et al., Plant Mol. Biol. 16, 983, 1991.
β -tubulin	Oppenheimer, et al., Gene 63: 87, 1988.
tobacco root-specific genes	Conkling, et al., Plant Physiol. 93: 1203, 1990.
B. napus G1-3b gene	United States Patent No. 5, 401, 836
SbPRP1	Suzuki et al., Plant Mol. Biol. 21: 109-119, 1993.
LRX1	Baumberger et al. 2001, Genes & Dev. 15:1128
BTG-26 Brassica napus	US 20050044585
LeAMT1 (tomato)	Lauter et al. (1996, PNAS 3:8139)
The LeNRT1-1 (tomato)	Lauter et al. (1996, PNAS 3:8139)
class I patatin gene (potato)	Liu et al., Plant Mol. Biol. 17 (6): 1139-1154

KDC1 (<i>Daucus carota</i>)	Downey et al. (2000, J. Biol. Chem. 275:39420)
TobRB7 gene	W Song (1997) PhD Thesis, North Carolina State University, Raleigh, NC USA
OsRAB5a (rice)	Wang et al. 2002, Plant Sci. 163:273
ALF5 (<i>Arabidopsis</i>)	Diener et al. (2001, Plant Cell 13:1625)
NRT2;1Np (<i>N. plumbaginifolia</i>)	Quesada et al. (1997, Plant Mol. Biol. 34:265)

A seed-specific promoter is transcriptionally active predominantly in seed tissue, but not necessarily exclusively in seed tissue (in cases of leaky expression). The seed-specific promoter may be active during seed development and/or during germination. The seed specific promoter may be endosperm/aleurone/embryo specific. Examples of seed-specific promoters (endosperm/aleurone/embryo specific) are shown in Table 2c to Table 2f below. Further examples of seed-specific promoters are given in Qing Qu and Takaiwa (Plant Biotechnol. J. 2, 113-125, 2004), which disclosure is incorporated by reference herein as if fully set forth.

Table 2c: Examples of seed-specific promoters

Gene source	Reference
seed-specific genes	Simon et al., Plant Mol. Biol. 5: 191, 1985;
	Scofield et al., J. Biol. Chem. 262: 12202, 1987.;
	Baszczynski et al., Plant Mol. Biol. 14: 633, 1990.
Brazil Nut albumin	Pearson et al., Plant Mol. Biol. 18: 235-245, 1992.
legumin	Ellis et al., Plant Mol. Biol. 10: 203-214, 1988.
glutelin (rice)	Takaiwa et al., Mol. Gen. Genet. 208: 15-22, 1986;
	Takaiwa et al., FEBS Letts. 221: 43-47, 1987.
zein	Matzke et al Plant Mol Biol, 14(3):323-32 1990
napA	Stalberg et al, Planta 199: 515-519, 1996.
wheat LMW and HMW glutenin-1	Mol Gen Genet 216:81-90, 1989; NAR 17:461-2, 1989
wheat SPA	Albani et al, Plant Cell, 9: 171-184, 1997
wheat α , β , γ -gliadins	EMBO J. 3:1409-15, 1984
barley ltr1 promoter	Diaz et al. (1995) Mol Gen Genet 248(5):592-8
barley B1, C, D, hordein	Theor Appl Gen 98:1253-62, 1999; Plant J 4:343-55, 1993; Mol Gen Genet 250:750-60, 1996
barley DOF	Mena et al, The Plant Journal, 116(1): 53-62, 1998
blz2	EP99106056.7
synthetic promoter	Vicente-Carbajosa et al., Plant J. 13: 629-640, 1998.
rice prolamin NRP33	Wu et al, Plant Cell Physiology 39(8) 885-889, 1998
rice α -globulin Glb-1	Wu et al, Plant Cell Physiology 39(8) 885-889, 1998
rice OSH1	Sato et al, Proc. Natl. Acad. Sci. USA, 93: 8117-8122, 1996

rice α -globulin REB/OHP-1	Nakase et al. Plant Mol. Biol. 33: 513-522, 1997
rice ADP-glucose pyrophosphorylase	Trans Res 6:157-68, 1997
maize ESR gene family	Plant J 12:235-46, 1997
sorghum α -kafirin	DeRose et al., Plant Mol. Biol 32:1029-35, 1996
KNOX	Postma-Haarsma et al, Plant Mol. Biol. 39:257-71, 1999
rice oleosin	Wu et al, J. Biochem. 123:386, 1998
sunflower oleosin	Cummins et al., Plant Mol. Biol. 19: 873-876, 1992
PRO0117, putative rice 40S ribosomal protein	WO 2004/070039
PRO0136, rice alanine aminotransferase	unpublished
PRO0147, trypsin inhibitor ITR1 (barley)	unpublished
PRO0151, rice WSI18	WO 2004/070039
PRO0175, rice RAB21	WO 2004/070039
PRO005	WO 2004/070039
PRO0095	WO 2004/070039
α -amylase (Amy32b)	Lanahan et al, Plant Cell 4:203-211, 1992; Skriver et al, Proc Natl Acad Sci USA 88:7266-7270, 1991
cathepsin β -like gene	Cejudo et al, Plant Mol Biol 20:849-856, 1992
Barley Ltp2	Kalla et al., Plant J. 6:849-60, 1994
Chi26	Leah et al., Plant J. 4:579-89, 1994
Maize B-Peru	Selinger et al., Genetics 149:1125-38, 1998

Table 2d: examples of endosperm-specific promoters

Gene source	Reference
glutelin (rice)	Takaiwa et al. (1986) Mol Gen Genet 208:15-22; Takaiwa et al. (1987) FEBS Letts. 221:43-47
zein	Matzke et al., (1990) Plant Mol Biol 14(3): 323-32
wheat LMW and HMW glutenin-1	Colot et al. (1989) Mol Gen Genet 216:81-90, Anderson et al. (1989) NAR 17:461-2
wheat SPA	Albani et al. (1997) Plant Cell 9:171-184
wheat gliadins	Rafalski et al. (1984) EMBO 3:1409-15
barley ltr1 promoter	Diaz et al. (1995) Mol Gen Genet 248(5):592-8
barley B1, C, D, hordein	Cho et al. (1999) Theor Appl Genet 98:1253-62; Muller et al. (1993) Plant J 4:343-55; Sorenson et al. (1996) Mol Gen Genet 250:750-60
barley DOF	Mena et al, (1998) Plant J 116(1): 53-62
blz2	Onate et al. (1999) J Biol Chem 274(14):9175-82
synthetic promoter	Vicente-Carbajosa et al. (1998) Plant J 13:629-640

rice prolamin NRP33	Wu et al, (1998) Plant Cell Physiol 39(8) 885-889
rice globulin Glb-1	Wu et al. (1998) Plant Cell Physiol 39(8) 885-889
rice globulin REB/OHP-1	Nakase et al. (1997) Plant Molec Biol 33: 513-522
rice ADP-glucose pyrophosphorylase	Russell et al. (1997) Trans Res 6:157-68
maize ESR gene family	Opsahl-Ferstad et al. (1997) Plant J 12:235-46
sorghum kafirin	DeRose et al. (1996) Plant Mol Biol 32:1029-35

Table 2e: Examples of embryo specific promoters:

Gene source	Reference
rice OSH1	Sato et al, Proc. Natl. Acad. Sci. USA, 93: 8117-8122, 1996
KNOX	Postma-Haarsma et al, Plant Mol. Biol. 39:257-71, 1999
PRO0151	WO 2004/070039
PRO0175	WO 2004/070039
PRO005	WO 2004/070039
PRO0095	WO 2004/070039

Table 2f: Examples of aleurone-specific promoters:

Gene source	Reference
α -amylase (Amy32b)	Lanahan et al, Plant Cell 4:203-211, 1992; Skriver et al, Proc Natl Acad Sci USA 88:7266-7270, 1991
cathepsin β -like gene	Cejudo et al, Plant Mol Biol 20:849-856, 1992
Barley Ltp2	Kalla et al., Plant J. 6:849-60, 1994
Chi26	Leah et al., Plant J. 4:579-89, 1994
Maize B-Peru	Selinger et al., Genetics 149:1125-38,1998

A green tissue-specific promoter as defined herein is a promoter that is transcriptionally active predominantly in green tissue, substantially to the exclusion of any other parts of a plant, whilst still allowing for any leaky expression in these other plant parts.

Examples of green tissue-specific promoters which may be used to perform the methods of the invention are shown in Table 2g below.

Table 2g: Examples of green tissue-specific promoters

Gene	Expression	Reference
Maize Orthophosphate dikinase	Leaf specific	Fukavama et al., Plant Physiol. 2001 Nov;127(3):1136-46
Maize Phosphoenolpyruvate carboxylase	Leaf specific	Kausch et al., Plant Mol Biol. 2001 Jan;45(1):1-15
Rice Phosphoenolpyruvate carboxylase	Leaf specific	Lin et al., 2004 DNA Seq. 2004 Aug;15(4):269-76
Rice small subunit Rubisco	Leaf specific	Nomura et al., Plant Mol Biol.

		2000 Sep;44(1):99-106
rice beta expansin EXBP9	Shoot specific	WO 2004/070039
Pigeonpea small subunit Rubisco	Leaf specific	Panguluri et al., Indian J Exp Biol. 2005 Apr;43(4):369-72
Pea RBCS3A	Leaf specific	

Another example of a tissue-specific promoter is a meristem-specific promoter, which is transcriptionally active predominantly in meristematic tissue, substantially to the exclusion of any other parts of a plant, whilst still allowing for any leaky expression in these other plant parts. Examples of green meristem-specific promoters which may be used to perform the methods of the invention are shown in Table 2h below.

Table 2h: Examples of meristem-specific promoters

Gene source	Expression pattern	Reference
rice OSH1	Shoot apical meristem, from embryo globular stage to seedling stage	Sato et al. (1996) Proc. Natl. Acad. Sci. USA, 93: 8117-8122
Rice metallothionein	Meristem specific	BAD87835.1
WAK1 & WAK 2	Shoot and root apical meristems, and in expanding leaves and sepals	Wagner & Kohorn (2001) Plant Cell 13(2): 303-318

Terminator

The term "terminator" encompasses a control sequence which is a DNA sequence at the end of a transcriptional unit which signals 3' processing and polyadenylation of a primary transcript and termination of transcription. The terminator can be derived from the natural gene, from a variety of other plant genes, or from T-DNA. The terminator to be added may be derived from, for example, the nopaline synthase or octopine synthase genes, or alternatively from another plant gene, or less preferably from any other eukaryotic gene.

Selectable marker (gene)/Reporter gene

"Selectable marker", "selectable marker gene" or "reporter gene" includes any gene that confers a phenotype on a cell in which it is expressed to facilitate the identification and/or selection of cells that are transfected or transformed with a nucleic acid construct of the invention. These marker genes enable the identification of a successful transfer of the nucleic acid molecules via a series of different principles. Suitable markers may be selected from markers that confer antibiotic or herbicide resistance, that introduce a new metabolic trait or that allow visual selection. Examples of selectable marker genes include genes conferring resistance to antibiotics (such as nptII that phosphorylates neomycin and kanamycin, or hpt, phosphorylating hygromycin, or genes conferring resistance to, for example, bleomycin, streptomycin, tetracyclin, chloramphenicol, ampicillin, gentamycin,

geneticin (G418), spectinomycin or blasticidin), to herbicides (for example bar which provides resistance to Basta®; aroA or gox providing resistance against glyphosate, or the genes conferring resistance to, for example, imidazolinone, phosphinothricin or sulfonylurea), or genes that provide a metabolic trait (such as manA that allows plants to use mannose as sole carbon source or xylose isomerase for the utilisation of xylose, or antinutritive markers such as the resistance to 2-deoxyglucose). Expression of visual marker genes results in the formation of colour (for example β -glucuronidase, GUS or β -galactosidase with its coloured substrates, for example X-Gal), luminescence (such as the luciferin/luciferase system) or fluorescence (Green Fluorescent Protein, GFP, and derivatives thereof). This list represents only a small number of possible markers. The skilled worker is familiar with such markers. Different markers are preferred, depending on the organism and the selection method.

It is known that upon stable or transient integration of nucleic acids into plant cells, only a minority of the cells takes up the foreign DNA and, if desired, integrates it into its genome, depending on the expression vector used and the transfection technique used. To identify and select these integrants, a gene coding for a selectable marker (such as the ones described above) is usually introduced into the host cells together with the gene of interest. These markers can for example be used in mutants in which these genes are not functional by, for example, deletion by conventional methods. Furthermore, nucleic acid molecules encoding a selectable marker can be introduced into a host cell on the same vector that comprises the sequence encoding the polypeptides of the invention or used in the methods of the invention, or else in a separate vector. Cells which have been stably transfected with the introduced nucleic acid can be identified for example by selection (for example, cells which have integrated the selectable marker survive whereas the other cells die).

Since the marker genes, particularly genes for resistance to antibiotics and herbicides, are no longer required or are undesired in the transgenic host cell once the nucleic acids have been introduced successfully, the process according to the invention for introducing the nucleic acids advantageously employs techniques which enable the removal or excision of these marker genes. One such a method is what is known as co-transformation. The co-transformation method employs two vectors simultaneously for the transformation, one vector bearing the nucleic acid according to the invention and a second bearing the marker gene(s). A large proportion of transformants receives or, in the case of plants, comprises (up to 40% or more of the transformants), both vectors. In case of transformation with *Agrobacteria*, the transformants usually receive only a part of the vector, i.e. the sequence flanked by the T-DNA, which usually represents the expression cassette. The marker genes can subsequently be removed from the transformed plant by performing crosses. In another method, marker genes integrated into a transposon are used for the transformation together with desired nucleic acid (known as the Ac/Ds technology). The transformants can be crossed with a transposase source or the transformants are transformed with a nucleic acid construct conferring expression of a transposase, transiently or stable. In some cases

(approx. 10%), the transposon jumps out of the genome of the host cell once transformation has taken place successfully and is lost. In a further number of cases, the transposon jumps to a different location. In these cases the marker gene must be eliminated by performing crosses. In microbiology, techniques were developed which make possible, or facilitate, the detection of such events. A further advantageous method relies on what is known as recombination systems; whose advantage is that elimination by crossing can be dispensed with. The best-known system of this type is what is known as the Cre/lox system. Cre1 is a recombinase that removes the sequences located between the loxP sequences. If the marker gene is integrated between the loxP sequences, it is removed once transformation has taken place successfully, by expression of the recombinase. Further recombination systems are the HIN/HIX, FLP/FRT and REP/STB system (Tribble et al., J. Biol. Chem., 275, 2000: 22255-22267; Velmurugan et al., J. Cell Biol., 149, 2000: 553-566). A site-specific integration into the plant genome of the nucleic acid sequences according to the invention is possible. Naturally, these methods can also be applied to microorganisms such as yeast, fungi or bacteria.

Transgenic/Transgene/Recombinant

For the purposes of the invention, "transgenic", "transgene" or "recombinant" means with regard to, for example, a nucleic acid sequence, an expression cassette, gene construct or a vector comprising the nucleic acid sequence or an organism transformed with the nucleic acid sequences, expression cassettes or vectors according to the invention, all those constructions brought about by recombinant methods in which either

- (a) the nucleic acid sequences encoding proteins useful in the methods of the invention, or
- (b) genetic control sequence(s) which is operably linked with the nucleic acid sequence according to the invention, for example a promoter, or
- (c) a) and b)

are not located in their natural genetic environment or have been modified by recombinant methods, it being possible for the modification to take the form of, for example, a substitution, addition, deletion, inversion or insertion of one or more nucleotide residues. The natural genetic environment is understood as meaning the natural genomic or chromosomal locus in the original plant or the presence in a genomic library. In the case of a genomic library, the natural genetic environment of the nucleic acid sequence is preferably retained, at least in part. The environment flanks the nucleic acid sequence at least on one side and has a sequence length of at least 50 bp, preferably at least 500 bp, especially preferably at least 1000 bp, most preferably at least 5000 bp. A naturally occurring expression cassette – for example the naturally occurring combination of the natural promoter of the nucleic acid sequences with the corresponding nucleic acid sequence encoding a polypeptide useful in the methods of the present invention, as defined above – becomes a transgenic expression cassette when this expression cassette is modified by non-natural, synthetic ("artificial") methods such as, for example, mutagenic treatment. Suitable methods are described, for example, in US 5,565,350 or WO 00/15815.

A transgenic plant for the purposes of the invention is thus understood as meaning, as above, that the nucleic acids used in the method of the invention are not present in, or originating from, the genome of said plant, or are present in the genome of said plant but not at their natural locus in the genome of said plant, it being possible for the nucleic acids to be expressed homologously or heterologously. However, as mentioned, transgenic also means that, while the nucleic acids according to the invention or used in the inventive method are at their natural position in the genome of a plant, the sequence has been modified with regard to the natural sequence, and/or that the regulatory sequences of the natural sequences have been modified. Transgenic is preferably understood as meaning the expression of the nucleic acids according to the invention at an unnatural locus in the genome, i.e. homologous or, preferably, heterologous expression of the nucleic acids takes place. Preferred transgenic plants are mentioned herein.

It shall further be noted that in the context of the present invention, the term “isolated nucleic acid” or “isolated polypeptide” may in some instances be considered as a synonym for a “recombinant nucleic acid” or a “recombinant polypeptide”, respectively and refers to a nucleic acid or polypeptide that is not located in its natural genetic environment and/or that has been modified by recombinant methods.

Modulation

The term “modulation” means in relation to expression or gene expression, a process in which the expression level is changed by said gene expression in comparison to the control plant, the expression level may be increased or decreased. The original, unmodulated expression may be of any kind of expression of a structural RNA (rRNA, tRNA) or mRNA with subsequent translation. For the purposes of this invention, the original unmodulated expression may also be absence of any expression. The term “modulating the activity” shall mean any change of the expression of the inventive nucleic acid sequences or encoded proteins, which leads to increased yield and/or increased growth of the plants. The expression can increase from zero (absence of, or immeasurable expression) to a certain amount, or can decrease from a certain amount to immeasurable small amounts or zero.

Expression

The term “expression” or “gene expression” means the transcription of a specific gene or specific genes or specific genetic construct. The term “expression” or “gene expression” in particular means the transcription of a gene or genes or genetic construct into structural RNA (rRNA, tRNA) or mRNA with or without subsequent translation of the latter into a protein. The process includes transcription of DNA and processing of the resulting mRNA product.

Increased expression/overexpression

The term “increased expression” or “overexpression” as used herein means any form of expression that is additional to the original wild-type expression level. For the purposes of this invention, the original wild-type expression level might also be zero, i.e. absence of expression or immeasurable expression.

Methods for increasing expression of genes or gene products are well documented in the art and include, for example, overexpression driven by appropriate promoters, the use of transcription enhancers or translation enhancers. Isolated nucleic acids which serve as promoter or enhancer elements may be introduced in an appropriate position (typically upstream) of a non-heterologous form of a polynucleotide so as to upregulate expression of a nucleic acid encoding the polypeptide of interest. For example, endogenous promoters may be altered in vivo by mutation, deletion, and/or substitution (see, Kmiec, US 5,565,350; Zarling et al., WO9322443), or isolated promoters may be introduced into a plant cell in the proper orientation and distance from a gene of the present invention so as to control the expression of the gene.

If polypeptide expression is desired, it is generally desirable to include a polyadenylation region at the 3'-end of a polynucleotide coding region. The polyadenylation region can be derived from the natural gene, from a variety of other plant genes, or from T-DNA. The 3' end sequence to be added may be derived from, for example, the nopaline synthase or octopine synthase genes, or alternatively from another plant gene, or less preferably from any other eukaryotic gene.

An intron sequence may also be added to the 5' untranslated region (UTR) or the coding sequence of the partial coding sequence to increase the amount of the mature message that accumulates in the cytosol. Inclusion of a spliceable intron in the transcription unit in both plant and animal expression constructs has been shown to increase gene expression at both the mRNA and protein levels up to 1000-fold (Buchman and Berg (1988) Mol. Cell biol. 8: 4395-4405; Callis et al. (1987) Genes Dev 1:1183-1200). Such intron enhancement of gene expression is typically greatest when placed near the 5' end of the transcription unit. Use of the maize introns Adh1-S intron 1, 2, and 6, the Bronze-1 intron are known in the art. For general information see: The Maize Handbook, Chapter 116, Freeling and Walbot, Eds., Springer, N.Y. (1994).

Decreased expression

Reference herein to “decreased expression” or “reduction or substantial elimination” of expression is taken to mean a decrease in endogenous gene expression and/or polypeptide levels and/or polypeptide activity relative to control plants. The reduction or substantial elimination is in increasing order of preference at least 10%, 20%, 30%, 40% or 50%, 60%, 70%, 80%, 85%, 90%, or 95%, 96%, 97%, 98%, 99% or more reduced compared to that of control plants.

For the reduction or substantial elimination of expression an endogenous gene in a plant, a sufficient length of substantially contiguous nucleotides of a nucleic acid sequence is required. In order to perform gene silencing, this may be as little as 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10 or fewer nucleotides, alternatively this may be as much as the entire gene (including the 5' and/or 3' UTR, either in part or in whole). The stretch of substantially contiguous nucleotides may be derived from the nucleic acid encoding the protein of interest (target gene), or from any nucleic acid capable of encoding an orthologue, paralogue or homologue of the protein of interest. Preferably, the stretch of substantially contiguous nucleotides is capable of forming hydrogen bonds with the target gene (either sense or antisense strand), more preferably, the stretch of substantially contiguous nucleotides has, in increasing order of preference, 50%, 60%, 70%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, 100% sequence identity to the target gene (either sense or antisense strand). A nucleic acid sequence encoding a (functional) polypeptide is not a requirement for the various methods discussed herein for the reduction or substantial elimination of expression of an endogenous gene.

This reduction or substantial elimination of expression may be achieved using routine tools and techniques. A preferred method for the reduction or substantial elimination of endogenous gene expression is by introducing and expressing in a plant a genetic construct into which the nucleic acid (in this case a stretch of substantially contiguous nucleotides derived from the gene of interest, or from any nucleic acid capable of encoding an orthologue, paralogue or homologue of any one of the protein of interest) is cloned as an inverted repeat (in part or completely), separated by a spacer (non-coding DNA).

In such a preferred method, expression of the endogenous gene is reduced or substantially eliminated through RNA-mediated silencing using an inverted repeat of a nucleic acid or a part thereof (in this case a stretch of substantially contiguous nucleotides derived from the gene of interest, or from any nucleic acid capable of encoding an orthologue, paralogue or homologue of the protein of interest), preferably capable of forming a hairpin structure. The inverted repeat is cloned in an expression vector comprising control sequences. A non-coding DNA nucleic acid sequence (a spacer, for example a matrix attachment region fragment (MAR), an intron, a polylinker, etc.) is located between the two inverted nucleic acids forming the inverted repeat. After transcription of the inverted repeat, a chimeric RNA with a self-complementary structure is formed (partial or complete). This double-stranded RNA structure is referred to as the hairpin RNA (hpRNA). The hpRNA is processed by the plant into siRNAs that are incorporated into an RNA-induced silencing complex (RISC). The RISC further cleaves the mRNA transcripts, thereby substantially reducing the number of mRNA transcripts to be translated into polypeptides. For further general details see for example, Grierson et al. (1998) WO 98/53083; Waterhouse et al. (1999) WO 99/53050).

Performance of the methods of the invention does not rely on introducing and expressing in a plant a genetic construct into which the nucleic acid is cloned as an inverted repeat, but

any one or more of several well-known "gene silencing" methods may be used to achieve the same effects.

One such method for the reduction of endogenous gene expression is RNA-mediated silencing of gene expression (downregulation). Silencing in this case is triggered in a plant by a double stranded RNA sequence (dsRNA) that is substantially similar to the target endogenous gene. This dsRNA is further processed by the plant into about 20 to about 26 nucleotides called short interfering RNAs (siRNAs). The siRNAs are incorporated into an RNA-induced silencing complex (RISC) that cleaves the mRNA transcript of the endogenous target gene, thereby substantially reducing the number of mRNA transcripts to be translated into a polypeptide. Preferably, the double stranded RNA sequence corresponds to a target gene.

Another example of an RNA silencing method involves the introduction of nucleic acid sequences or parts thereof (in this case a stretch of substantially contiguous nucleotides derived from the gene of interest, or from any nucleic acid capable of encoding an orthologue, paralogue or homologue of the protein of interest) in a sense orientation into a plant. "Sense orientation" refers to a DNA sequence that is homologous to an mRNA transcript thereof. Introduced into a plant would therefore be at least one copy of the nucleic acid sequence. The additional nucleic acid sequence will reduce expression of the endogenous gene, giving rise to a phenomenon known as co-suppression. The reduction of gene expression will be more pronounced if several additional copies of a nucleic acid sequence are introduced into the plant, as there is a positive correlation between high transcript levels and the triggering of co-suppression.

Another example of an RNA silencing method involves the use of antisense nucleic acid sequences. An "antisense" nucleic acid sequence comprises a nucleotide sequence that is complementary to a "sense" nucleic acid sequence encoding a protein, i.e. complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA transcript sequence. The antisense nucleic acid sequence is preferably complementary to the endogenous gene to be silenced. The complementarity may be located in the "coding region" and/or in the "non-coding region" of a gene. The term "coding region" refers to a region of the nucleotide sequence comprising codons that are translated into amino acid residues. The term "non-coding region" refers to 5' and 3' sequences that flank the coding region that are transcribed but not translated into amino acids (also referred to as 5' and 3' untranslated regions).

Antisense nucleic acid sequences can be designed according to the rules of Watson and Crick base pairing. The antisense nucleic acid sequence may be complementary to the entire nucleic acid sequence (in this case a stretch of substantially contiguous nucleotides derived from the gene of interest, or from any nucleic acid capable of encoding an orthologue, paralogue or homologue of the protein of interest), but may also be an

oligonucleotide that is antisense to only a part of the nucleic acid sequence (including the mRNA 5' and 3' UTR). For example, the antisense oligonucleotide sequence may be complementary to the region surrounding the translation start site of an mRNA transcript encoding a polypeptide. The length of a suitable antisense oligonucleotide sequence is known in the art and may start from about 50, 45, 40, 35, 30, 25, 20, 15 or 10 nucleotides in length or less. An antisense nucleic acid sequence according to the invention may be constructed using chemical synthesis and enzymatic ligation reactions using methods known in the art. For example, an antisense nucleic acid sequence (e.g., an antisense oligonucleotide sequence) may be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acid sequences, e.g., phosphorothioate derivatives and acridine substituted nucleotides may be used. Examples of modified nucleotides that may be used to generate the antisense nucleic acid sequences are well known in the art. Known nucleotide modifications include methylation, cyclization and 'caps' and substitution of one or more of the naturally occurring nucleotides with an analogue such as inosine. Other modifications of nucleotides are well known in the art.

The antisense nucleic acid sequence can be produced biologically using an expression vector into which a nucleic acid sequence has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest). Preferably, production of antisense nucleic acid sequences in plants occurs by means of a stably integrated nucleic acid construct comprising a promoter, an operably linked antisense oligonucleotide, and a terminator.

The nucleic acid molecules used for silencing in the methods of the invention (whether introduced into a plant or generated in situ) hybridize with or bind to mRNA transcripts and/or genomic DNA encoding a polypeptide to thereby inhibit expression of the protein, e.g., by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid sequence which binds to DNA duplexes, through specific interactions in the major groove of the double helix. Antisense nucleic acid sequences may be introduced into a plant by transformation or direct injection at a specific tissue site. Alternatively, antisense nucleic acid sequences can be modified to target selected cells and then administered systemically. For example, for systemic administration, antisense nucleic acid sequences can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, e.g., by linking the antisense nucleic acid sequence to peptides or antibodies which bind to cell surface receptors or antigens. The antisense nucleic acid sequences can also be delivered to cells using the vectors described herein.

According to a further aspect, the antisense nucleic acid sequence is an α -anomeric nucleic acid sequence. An α -anomeric nucleic acid sequence forms specific double-stranded

hybrids with complementary RNA in which, contrary to the usual b-units, the strands run parallel to each other (Gaultier et al. (1987) Nucl Ac Res 15: 6625-6641). The antisense nucleic acid sequence may also comprise a 2'-o-methylribonucleotide (Inoue et al. (1987) Nucl Ac Res 15, 6131-6148) or a chimeric RNA-DNA analogue (Inoue et al. (1987) FEBS Lett. 215, 327-330).

The reduction or substantial elimination of endogenous gene expression may also be performed using ribozymes. Ribozymes are catalytic RNA molecules with ribonuclease activity that are capable of cleaving a single-stranded nucleic acid sequence, such as an mRNA, to which they have a complementary region. Thus, ribozymes (e.g., hammerhead ribozymes (described in Haselhoff and Gerlach (1988) Nature 334, 585-591) can be used to catalytically cleave mRNA transcripts encoding a polypeptide, thereby substantially reducing the number of mRNA transcripts to be translated into a polypeptide. A ribozyme having specificity for a nucleic acid sequence can be designed (see for example: Cech et al. U.S. Patent No. 4,987,071; and Cech et al. U.S. Patent No. 5,116,742). Alternatively, mRNA transcripts corresponding to a nucleic acid sequence can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules (Bartel and Szostak (1993) Science 261, 1411-1418). The use of ribozymes for gene silencing in plants is known in the art (e.g., Atkins et al. (1994) WO 94/00012; Lenne et al. (1995) WO 95/03404; Lutziger et al. (2000) WO 00/00619; Prinsen et al. (1997) WO 97/13865 and Scott et al. (1997) WO 97/38116).

Gene silencing may also be achieved by insertion mutagenesis (for example, T-DNA insertion or transposon insertion) or by strategies as described by, among others, Angell and Baulcombe ((1999) Plant J 20(3): 357-62), (Amplicon VIGS WO 98/36083), or Baulcombe (WO 99/15682).

Gene silencing may also occur if there is a mutation on an endogenous gene and/or a mutation on an isolated gene/nucleic acid subsequently introduced into a plant. The reduction or substantial elimination may be caused by a non-functional polypeptide. For example, the polypeptide may bind to various interacting proteins; one or more mutation(s) and/or truncation(s) may therefore provide for a polypeptide that is still able to bind interacting proteins (such as receptor proteins) but that cannot exhibit its normal function (such as signalling ligand).

A further approach to gene silencing is by targeting nucleic acid sequences complementary to the regulatory region of the gene (e.g., the promoter and/or enhancers) to form triple helical structures that prevent transcription of the gene in target cells. See Helene, C., Anticancer Drug Res. 6, 569-84, 1991; Helene et al., Ann. N.Y. Acad. Sci. 660, 27-36 1992; and Maher, L.J. Bioassays 14, 807-15, 1992.

Other methods, such as the use of antibodies directed to an endogenous polypeptide for inhibiting its function in planta, or interference in the signalling pathway in which a polypeptide is involved, will be well known to the skilled man. In particular, it can be envisaged that manmade molecules may be useful for inhibiting the biological function of a target polypeptide, or for interfering with the signalling pathway in which the target polypeptide is involved.

Alternatively, a screening program may be set up to identify in a plant population natural variants of a gene, which variants encode polypeptides with reduced activity. Such natural variants may also be used for example, to perform homologous recombination.

Artificial and/or natural microRNAs (miRNAs) may be used to knock out gene expression and/or mRNA translation. Endogenous miRNAs are single stranded small RNAs of typically 19-24 nucleotides long. They function primarily to regulate gene expression and/or mRNA translation. Most plant microRNAs (miRNAs) have perfect or near-perfect complementarity with their target sequences. However, there are natural targets with up to five mismatches. They are processed from longer non-coding RNAs with characteristic fold-back structures by double-strand specific RNases of the Dicer family. Upon processing, they are incorporated in the RNA-induced silencing complex (RISC) by binding to its main component, an Argonaute protein. MiRNAs serve as the specificity components of RISC, since they base-pair to target nucleic acids, mostly mRNAs, in the cytoplasm. Subsequent regulatory events include target mRNA cleavage and destruction and/or translational inhibition. Effects of miRNA overexpression are thus often reflected in decreased mRNA levels of target genes.

Artificial microRNAs (amiRNAs), which are typically 21 nucleotides in length, can be genetically engineered specifically to negatively regulate gene expression of single or multiple genes of interest. Determinants of plant microRNA target selection are well known in the art. Empirical parameters for target recognition have been defined and can be used to aid in the design of specific amiRNAs, (Schwab et al., *Dev. Cell* 8, 517–527, 2005). Convenient tools for design and generation of amiRNAs and their precursors are also available to the public (Schwab et al., *Plant Cell* 18, 1121-1133, 2006).

For optimal performance, the gene silencing techniques used for reducing expression in a plant of an endogenous gene requires the use of nucleic acid sequences from monocotyledonous plants for transformation of monocotyledonous plants, and from dicotyledonous plants for transformation of dicotyledonous plants. Preferably, a nucleic acid sequence from any given plant species is introduced into that same species. For example, a nucleic acid sequence from rice is transformed into a rice plant. However, it is not an absolute requirement that the nucleic acid sequence to be introduced originates from the same plant species as the plant in which it will be introduced. It is sufficient that there is

substantial homology between the endogenous target gene and the nucleic acid to be introduced.

Described above are examples of various methods for the reduction or substantial elimination of expression in a plant of an endogenous gene. A person skilled in the art would readily be able to adapt the aforementioned methods for silencing so as to achieve reduction of expression of an endogenous gene in a whole plant or in parts thereof through the use of an appropriate promoter, for example.

Transformation

The term "introduction" or "transformation" as referred to herein encompasses the transfer of an exogenous polynucleotide into a host cell, irrespective of the method used for transfer. Plant tissue capable of subsequent clonal propagation, whether by organogenesis or embryogenesis, may be transformed with a genetic construct of the present invention and a whole plant regenerated there from. The particular tissue chosen will vary depending on the clonal propagation systems available for, and best suited to, the particular species being transformed. Exemplary tissue targets include leaf disks, pollen, embryos, cotyledons, hypocotyls, megagametophytes, callus tissue, existing meristematic tissue (e.g., apical meristem, axillary buds, and root meristems), and induced meristem tissue (e.g., cotyledon meristem and hypocotyl meristem). The polynucleotide may be transiently or stably introduced into a host cell and may be maintained non-integrated, for example, as a plasmid. Alternatively, it may be integrated into the host genome. The resulting transformed plant cell may then be used to regenerate a transformed plant in a manner known to persons skilled in the art.

The transfer of foreign genes into the genome of a plant is called transformation. Transformation of plant species is now a fairly routine technique. Advantageously, any of several transformation methods may be used to introduce the gene of interest into a suitable ancestor cell. The methods described for the transformation and regeneration of plants from plant tissues or plant cells may be utilized for transient or for stable transformation. Transformation methods include the use of liposomes, electroporation, chemicals that increase free DNA uptake, injection of the DNA directly into the plant, particle gun bombardment, transformation using viruses or pollen and microprojection. Methods may be selected from the calcium/polyethylene glycol method for protoplasts (Krens, F.A. et al., (1982) *Nature* 296, 72-74; Negrutiu I et al. (1987) *Plant Mol Biol* 8: 363-373); electroporation of protoplasts (Shillito R.D. et al. (1985) *Bio/Technol* 3, 1099-1102); microinjection into plant material (Crossway A et al., (1986) *Mol. Gen Genet* 202: 179-185); DNA or RNA-coated particle bombardment (Klein TM et al., (1987) *Nature* 327: 70) infection with (non-integrative) viruses and the like. Transgenic plants, including transgenic crop plants, are preferably produced via *Agrobacterium*-mediated transformation. An advantageous transformation method is the transformation in planta. To this end, it is possible, for example, to allow the *agrobacteria* to act on plant seeds or to inoculate the

plant meristem with agrobacteria. It has proved particularly expedient in accordance with the invention to allow a suspension of transformed agrobacteria to act on the intact plant or at least on the flower primordia. The plant is subsequently grown on until the seeds of the treated plant are obtained (Clough and Bent, *Plant J.* (1998) 16, 735–743). Methods for Agrobacterium-mediated transformation of rice include well known methods for rice transformation, such as those described in any of the following: European patent application EP 1198985 A1, Aldemita and Hodges (*Planta* 199: 612-617, 1996); Chan et al. (*Plant Mol Biol* 22 (3): 491-506, 1993), Hiei et al. (*Plant J* 6 (2): 271-282, 1994), which disclosures are incorporated by reference herein as if fully set forth. In the case of corn transformation, the preferred method is as described in either Ishida et al. (*Nat. Biotechnol* 14(6): 745-50, 1996) or Frame et al. (*Plant Physiol* 129(1): 13-22, 2002), which disclosures are incorporated by reference herein as if fully set forth. Said methods are further described by way of example in B. Jené et al., *Techniques for Gene Transfer*, in: *Transgenic Plants*, Vol. 1, Engineering and Utilization, eds. S.D. Kung and R. Wu, Academic Press (1993) 128-143 and in Potrykus *Annu. Rev. Plant Physiol. Plant Molec. Biol.* 42 (1991) 205-225). The nucleic acids or the construct to be expressed is preferably cloned into a vector, which is suitable for transforming *Agrobacterium tumefaciens*, for example pBin19 (Bevan et al., *Nucl. Acids Res.* 12 (1984) 8711). Agrobacteria transformed by such a vector can then be used in known manner for the transformation of plants, such as plants used as a model, like *Arabidopsis* (*Arabidopsis thaliana* is within the scope of the present invention not considered as a crop plant), or crop plants such as, by way of example, tobacco plants, for example by immersing bruised leaves or chopped leaves in an agrobacterial solution and then culturing them in suitable media. The transformation of plants by means of *Agrobacterium tumefaciens* is described, for example, by Höfgen and Willmitzer in *Nucl. Acid Res.* (1988) 16, 9877 or is known inter alia from F.F. White, *Vectors for Gene Transfer in Higher Plants*; in *Transgenic Plants*, Vol. 1, Engineering and Utilization, eds. S.D. Kung and R. Wu, Academic Press, 1993, pp. 15-38.

In addition to the transformation of somatic cells, which then have to be regenerated into intact plants, it is also possible to transform the cells of plant meristems and in particular those cells which develop into gametes. In this case, the transformed gametes follow the natural plant development, giving rise to transgenic plants. Thus, for example, seeds of *Arabidopsis* are treated with agrobacteria and seeds are obtained from the developing plants of which a certain proportion is transformed and thus transgenic [Feldman, KA and Marks MD (1987). *Mol Gen Genet* 208:1-9; Feldmann K (1992). In: C Koncz, N-H Chua and J Shell, eds, *Methods in Arabidopsis Research*. World Scientific, Singapore, pp. 274-289]. Alternative methods are based on the repeated removal of the inflorescences and incubation of the excision site in the center of the rosette with transformed agrobacteria, whereby transformed seeds can likewise be obtained at a later point in time (Chang (1994). *Plant J.* 5: 551-558; Katavic (1994). *Mol Gen Genet*, 245: 363-370). However, an especially effective method is the vacuum infiltration method with its modifications such as the “floral dip” method. In the case of vacuum infiltration of *Arabidopsis*, intact plants under reduced

pressure are treated with an agrobacterial suspension [Bechthold, N (1993). C R Acad Sci Paris Life Sci, 316: 1194-1199], while in the case of the "floral dip" method the developing floral tissue is incubated briefly with a surfactant-treated agrobacterial suspension [Clough, SJ and Bent AF (1998) The Plant J. 16, 735-743]. A certain proportion of transgenic seeds are harvested in both cases, and these seeds can be distinguished from non-transgenic seeds by growing under the above-described selective conditions. In addition the stable transformation of plastids is of advantages because plastids are inherited maternally in most crops reducing or eliminating the risk of transgene flow through pollen. The transformation of the chloroplast genome is generally achieved by a process which has been schematically displayed in Klaus et al., 2004 [Nature Biotechnology 22 (2), 225-229]. Briefly the sequences to be transformed are cloned together with a selectable marker gene between flanking sequences homologous to the chloroplast genome. These homologous flanking sequences direct site specific integration into the plastome. Plastidal transformation has been described for many different plant species and an overview is given in Bock (2001) Transgenic plastids in basic research and plant biotechnology. J Mol Biol. 2001 Sep 21; 312 (3):425-38 or Maliga, P (2003) Progress towards commercialization of plastid transformation technology. Trends Biotechnol. 21, 20-28. Further biotechnological progress has recently been reported in form of marker free plastid transformants, which can be produced by a transient co-integrated marker gene (Klaus et al., 2004, Nature Biotechnology 22(2), 225-229).

The genetically modified plant cells can be regenerated via all methods with which the skilled worker is familiar. Suitable methods can be found in the abovementioned publications by S.D. Kung and R. Wu, Potrykus or Höfgen and Willmitzer.

Generally after transformation, plant cells or cell groupings are selected for the presence of one or more markers which are encoded by plant-expressible genes co-transferred with the gene of interest, following which the transformed material is regenerated into a whole plant. To select transformed plants, the plant material obtained in the transformation is, as a rule, subjected to selective conditions so that transformed plants can be distinguished from untransformed plants. For example, the seeds obtained in the above-described manner can be planted and, after an initial growing period, subjected to a suitable selection by spraying. A further possibility consists in growing the seeds, if appropriate after sterilization, on agar plates using a suitable selection agent so that only the transformed seeds can grow into plants. Alternatively, the transformed plants are screened for the presence of a selectable marker such as the ones described above.

Following DNA transfer and regeneration, putatively transformed plants may also be evaluated, for instance using Southern analysis, for the presence of the gene of interest, copy number and/or genomic organisation. Alternatively or additionally, expression levels of the newly introduced DNA may be monitored using Northern and/or Western analysis, both techniques being well known to persons having ordinary skill in the art.

The generated transformed plants may be propagated by a variety of means, such as by clonal propagation or classical breeding techniques. For example, a first generation (or T1) transformed plant may be selfed and homozygous second-generation (or T2) transformants selected, and the T2 plants may then further be propagated through classical breeding techniques. The generated transformed organisms may take a variety of forms. For example, they may be chimeras of transformed cells and non-transformed cells; clonal transformants (e.g., all cells transformed to contain the expression cassette); grafts of transformed and untransformed tissues (e.g., in plants, a transformed rootstock grafted to an untransformed scion).

T-DNA activation tagging

T-DNA activation tagging (Hayashi et al. Science (1992) 1350-1353), involves insertion of T-DNA, usually containing a promoter (may also be a translation enhancer or an intron), in the genomic region of the gene of interest or 10 kb up- or downstream of the coding region of a gene in a configuration such that the promoter directs expression of the targeted gene. Typically, regulation of expression of the targeted gene by its natural promoter is disrupted and the gene falls under the control of the newly introduced promoter. The promoter is typically embedded in a T-DNA. This T-DNA is randomly inserted into the plant genome, for example, through *Agrobacterium* infection and leads to modified expression of genes near the inserted T-DNA. The resulting transgenic plants show dominant phenotypes due to modified expression of genes close to the introduced promoter.

TILLING

The term "TILLING" is an abbreviation of "Targeted Induced Local Lesions In Genomes" and refers to a mutagenesis technology useful to generate and/or identify nucleic acids encoding proteins with modified expression and/or activity. TILLING also allows selection of plants carrying such mutant variants. These mutant variants may exhibit modified expression, either in strength or in location or in timing (if the mutations affect the promoter for example). These mutant variants may exhibit higher activity than that exhibited by the gene in its natural form. TILLING combines high-density mutagenesis with high-throughput screening methods. The steps typically followed in TILLING are: (a) EMS mutagenesis (Redei GP and Koncz C (1992) In *Methods in Arabidopsis Research*, Koncz C, Chua NH, Schell J, eds. Singapore, World Scientific Publishing Co, pp. 16–82; Feldmann et al., (1994) In Meyerowitz EM, Somerville CR, eds, *Arabidopsis*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, pp 137-172; Lightner J and Caspar T (1998) In J Martinez-Zapater, J Salinas, eds, *Methods on Molecular Biology*, Vol. 82. Humana Press, Totowa, NJ, pp 91-104); (b) DNA preparation and pooling of individuals; (c) PCR amplification of a region of interest; (d) denaturation and annealing to allow formation of heteroduplexes; (e) DHPLC, where the presence of a heteroduplex in a pool is detected as an extra peak in the chromatogram; (f) identification of the mutant individual; and (g) sequencing of the mutant

PCR product. Methods for TILLING are well known in the art (McCallum et al., (2000) Nat Biotechnol 18: 455-457; reviewed by Stemple (2004) Nat Rev Genet 5(2): 145-50).

Homologous recombination

Homologous recombination allows introduction in a genome of a selected nucleic acid at a defined selected position. Homologous recombination is a standard technology used routinely in biological sciences for lower organisms such as yeast or the moss *Physcomitrella*. Methods for performing homologous recombination in plants have been described not only for model plants (Offringa et al. (1990) EMBO J 9(10): 3077-84) but also for crop plants, for example rice (Terada et al. (2002) Nat Biotech 20(10): 1030-4; Iida and Terada (2004) Curr Opin Biotech 15(2): 132-8), and approaches exist that are generally applicable regardless of the target organism (Miller et al, Nature Biotechnol. 25, 778-785, 2007).

Yield related Traits

Yield related traits are traits or features which are related to plant yield. Yield-related traits may comprise one or more of the following non-limitative list of features: early flowering time, yield, biomass, seed yield, early vigour, greenness index, increased growth rate, improved agronomic traits, such as e.g. improved Water Use Efficiency (WUE), improved Nitrogen Use Efficiency (NUE), etc..

Yield

The term "yield" in general means a measurable produce of economic value, typically related to a specified crop, to an area, and to a period of time. Individual plant parts directly contribute to yield based on their number, size and/or weight, or the actual yield is the yield per square meter for a crop and year, which is determined by dividing total production (includes both harvested and appraised production) by planted square meters.

The terms "yield" of a plant and "plant yield" are used interchangeably herein and are meant to refer to vegetative biomass such as root and/or shoot biomass, to reproductive organs, and/or to propagules such as seeds of that plant.

Taking corn as an example, male inflorescences (tassels) and female inflorescences (ears). The female inflorescence produces pairs of spikelets on the surface of a central axis (cob). Each of the female spikelets encloses two fertile florets, one of whose will usually mature into a maize kernel once fertilized. Hence a yield increase in maize may be manifested as one or more of the following: increase in the number of plants established per square meter, an increase in the number of ears per plant, an increase in the number of rows, number of kernels per row, kernel weight, thousand kernel weight, ear length/diameter, increase in the seed filling rate, which is the number of filled florets (i.e. florets containing seed) divided by the total number of florets and multiplied by 100), among others.

Inflorescences in rice plants are called panicles. The panicle bears spikelets. The spikelet is the basic unit of the panicles and consists of a pedicel and a floret. The floret is born on the pedicel. A floret includes a flower that is covered by two protective glumes: a larger glume (the lemma) and a shorter glume (the palea). Hence, taking rice as an example, a yield increase may manifest itself as an increase in one or more of the following: number of plants per square meter, number of panicles per plant, panicle length, number of spikelets per panicle, number of flowers (or florets) per panicle, increase in the seed filling rate which is the number of filled florets (i.e. florets containing seeds divided by the total number of florets and multiplied by 100), increase in thousand kernel weight, among others. In rice, submergence tolerance may also result in increased yield.

Early flowering time

Plants having an “early flowering time” as used herein are plants which start to flower earlier than control plants. Hence this term refers to plants that show an earlier start of flowering. Flowering time of plants can be assessed by counting the number of days (“time to flower”) between sowing and the emergence of a first inflorescence. The “flowering time” of a plant can for instance be determined using the method as described in WO 2007/093444.

Early vigour

“Early vigour” refers to active healthy well-balanced growth especially during early stages of plant growth, and may result from increased plant fitness due to, for example, the plants being better adapted to their environment (i.e. optimizing the use of energy resources and partitioning between shoot and root). Plants having early vigour also show increased seedling survival and a better establishment of the crop, which often results in highly uniform fields (with the crop growing in uniform manner, i.e. with the majority of plants reaching the various stages of development at substantially the same time), and often better and higher yield. Therefore, early vigour may be determined by measuring various factors, such as thousand kernel weight, percentage germination, percentage emergence, seedling growth, seedling height, root length, root and shoot biomass and many more.

Increased growth rate

The increased growth rate may be specific to one or more parts of a plant (including seeds), or may be throughout substantially the whole plant. Plants having an increased growth rate may have a shorter life cycle. The life cycle of a plant may be taken to mean the time needed to grow from a dry mature seed up to the stage where the plant has produced dry mature seeds, similar to the starting material. This life cycle may be influenced by factors such as speed of germination, early vigour, growth rate, greenness index, flowering time and speed of seed maturation. The increase in growth rate may take place at one or more stages in the life cycle of a plant or during substantially the whole plant life cycle. Increased growth rate during the early stages in the life cycle of a plant may reflect enhanced vigour. The increase in growth rate may alter the harvest cycle of a plant allowing plants to be sown later and/or harvested sooner than would otherwise be possible (a similar effect may be

obtained with earlier flowering time). If the growth rate is sufficiently increased, it may allow for the further sowing of seeds of the same plant species (for example sowing and harvesting of rice plants followed by sowing and harvesting of further rice plants all within one conventional growing period). Similarly, if the growth rate is sufficiently increased, it may allow for the further sowing of seeds of different plants species (for example the sowing and harvesting of corn plants followed by, for example, the sowing and optional harvesting of soybean, potato or any other suitable plant). Harvesting additional times from the same rootstock in the case of some crop plants may also be possible. Altering the harvest cycle of a plant may lead to an increase in annual biomass production per square meter (due to an increase in the number of times (say in a year) that any particular plant may be grown and harvested). An increase in growth rate may also allow for the cultivation of transgenic plants in a wider geographical area than their wild-type counterparts, since the territorial limitations for growing a crop are often determined by adverse environmental conditions either at the time of planting (early season) or at the time of harvesting (late season). Such adverse conditions may be avoided if the harvest cycle is shortened. The growth rate may be determined by deriving various parameters from growth curves, such parameters may be: T-Mid (the time taken for plants to reach 50% of their maximal size) and T-90 (time taken for plants to reach 90% of their maximal size), amongst others.

Stress resistance

An increase in yield and/or growth rate occurs whether the plant is under non-stress conditions or whether the plant is exposed to various stresses compared to control plants. Plants typically respond to exposure to stress by growing more slowly. In conditions of severe stress, the plant may even stop growing altogether. Mild stress on the other hand is defined herein as being any stress to which a plant is exposed which does not result in the plant ceasing to grow altogether without the capacity to resume growth. Mild stress in the sense of the invention leads to a reduction in the growth of the stressed plants of less than 40%, 35%, 30% or 25%, more preferably less than 20% or 15% in comparison to the control plant under non-stress conditions. Due to advances in agricultural practices (irrigation, fertilization, pesticide treatments) severe stresses are not often encountered in cultivated crop plants. As a consequence, the compromised growth induced by mild stress is often an undesirable feature for agriculture. "Mild stresses" are the everyday biotic and/or abiotic (environmental) stresses to which a plant is exposed. Abiotic stresses may be due to drought or excess water, anaerobic stress, salt stress, chemical toxicity, oxidative stress and hot, cold or freezing temperatures.

"Biotic stresses" are typically those stresses caused by pathogens, such as bacteria, viruses, fungi, nematodes and insects.

The "abiotic stress" may be an osmotic stress caused by a water stress, e.g. due to drought, salt stress, or freezing stress. Abiotic stress may also be an oxidative stress or a cold stress. "Freezing stress" is intended to refer to stress due to freezing temperatures, i.e.

temperatures at which available water molecules freeze and turn into ice. "Cold stress", also called "chilling stress", is intended to refer to cold temperatures, e.g. temperatures below 10°, or preferably below 5°C, but at which water molecules do not freeze. As reported in Wang et al. (Planta (2003) 218: 1-14), abiotic stress leads to a series of morphological, physiological, biochemical and molecular changes that adversely affect plant growth and productivity. Drought, salinity, extreme temperatures and oxidative stress are known to be interconnected and may induce growth and cellular damage through similar mechanisms. Rabbani et al. (Plant Physiol (2003) 133: 1755-1767) describes a particularly high degree of "cross talk" between drought stress and high-salinity stress. For example, drought and/or salinisation are manifested primarily as osmotic stress, resulting in the disruption of homeostasis and ion distribution in the cell. Oxidative stress, which frequently accompanies high or low temperature, salinity or drought stress, may cause denaturing of functional and structural proteins. As a consequence, these diverse environmental stresses often activate similar cell signalling pathways and cellular responses, such as the production of stress proteins, up-regulation of anti-oxidants, accumulation of compatible solutes and growth arrest. The term "non-stress" conditions as used herein are those environmental conditions that allow optimal growth of plants. Persons skilled in the art are aware of normal soil conditions and climatic conditions for a given location. Plants with optimal growth conditions, (grown under non-stress conditions) typically yield in increasing order of preference at least 97%, 95%, 92%, 90%, 87%, 85%, 83%, 80%, 77% or 75% of the average production of such plant in a given environment. Average production may be calculated on harvest and/or season basis. Persons skilled in the art are aware of average yield productions of a crop.

In particular, the methods of the present invention may be performed under non-stress conditions. In an example, the methods of the present invention may be performed under non-stress conditions such as mild drought to give plants having increased yield relative to control plants.

In another embodiment, the methods of the present invention may be performed under stress conditions.

In an example, the methods of the present invention may be performed under stress conditions such as drought to give plants having increased yield relative to control plants.

In another example, the methods of the present invention may be performed under stress conditions such as nutrient deficiency to give plants having increased yield relative to control plants.

Nutrient deficiency may result from a lack of nutrients such as nitrogen, phosphates and other phosphorous-containing compounds, potassium, calcium, magnesium, manganese, iron and boron, amongst others.

In yet another example, the methods of the present invention may be performed under stress conditions such as salt stress to give plants having increased yield relative to control plants. The term salt stress is not restricted to common salt (NaCl), but may be any one or more of: NaCl, KCl, LiCl, MgCl₂, CaCl₂, amongst others.

In yet another example, the methods of the present invention may be performed under stress conditions such as cold stress or freezing stress to give plants having increased yield relative to control plants.

Increase/Improve/Enhance

The terms “increase”, “improve” or “enhance” are interchangeable and shall mean in the sense of the application at least a 3%, 4%, 5%, 6%, 7%, 8%, 9% or 10%, preferably at least 15% or 20%, more preferably 25%, 30%, 35% or 40% more yield and/or growth in comparison to control plants as defined herein.

Seed yield

Increased seed yield may manifest itself as one or more of the following:

- (a) an increase in seed biomass (total seed weight) which may be on an individual seed basis and/or per plant and/or per square meter;
- (b) increased number of flowers per plant;
- (c) increased number of seeds;
- (d) increased seed filling rate (which is expressed as the ratio between the number of filled florets divided by the total number of florets);
- (e) increased harvest index, which is expressed as a ratio of the yield of harvestable parts, such as seeds, divided by the biomass of aboveground plant parts; and
- (f) increased thousand kernel weight (TKW), which is extrapolated from the number of seeds counted and their total weight. An increased TKW may result from an increased seed size and/or seed weight, and may also result from an increase in embryo and/or endosperm size.

An increase in seed yield may also be manifested as an increase in seed size and/or seed volume. Furthermore, an increase in seed yield may also manifest itself as an increase in seed area and/or seed length and/or seed width and/or seed perimeter.

Greenness Index

The “greenness index” as used herein is calculated from digital images of plants. For each pixel belonging to the plant object on the image, the ratio of the green value versus the red value (in the RGB model for encoding color) is calculated. The greenness index is expressed as the percentage of pixels for which the green-to-red ratio exceeds a given threshold. Under normal growth conditions, under salt stress growth conditions, and under reduced nutrient availability growth conditions, the greenness index of plants is measured in

the last imaging before flowering. In contrast, under drought stress growth conditions, the greenness index of plants is measured in the first imaging after drought.

Biomass

The term “biomass” as used herein is intended to refer to the total weight of a plant. Within the definition of biomass, a distinction may be made between the biomass of one or more parts of a plant, which may include any one or more of the following:

- aboveground parts such as but not limited to shoot biomass, seed biomass, leaf biomass, etc. ;
- aboveground harvestable parts such as but not limited to shoot biomass, seed biomass, leaf biomass, etc. ;
- parts below ground, such as but not limited to root biomass, etc.;
- harvestable parts below ground, such as but not limited to root biomass, etc.;
- vegetative biomass such as root biomass, shoot biomass, etc., ;
- reproductive organs; and
- propagules such as seed.

Marker assisted breeding

Such breeding programmes sometimes require introduction of allelic variation by mutagenic treatment of the plants, using for example EMS mutagenesis; alternatively, the programme may start with a collection of allelic variants of so called “natural” origin caused unintentionally. Identification of allelic variants then takes place, for example, by PCR. This is followed by a step for selection of superior allelic variants of the sequence in question and which give increased yield. Selection is typically carried out by monitoring growth performance of plants containing different allelic variants of the sequence in question. Growth performance may be monitored in a greenhouse or in the field. Further optional steps include crossing plants in which the superior allelic variant was identified with another plant. This could be used, for example, to make a combination of interesting phenotypic features.

Use as probes in (gene mapping)

Use of nucleic acids encoding the protein of interest for genetically and physically mapping the genes requires only a nucleic acid sequence of at least 15 nucleotides in length. These nucleic acids may be used as restriction fragment length polymorphism (RFLP) markers. Southern blots (Sambrook J, Fritsch EF and Maniatis T (1989) Molecular Cloning, A Laboratory Manual) of restriction-digested plant genomic DNA may be probed with the nucleic acids encoding the protein of interest. The resulting banding patterns may then be subjected to genetic analyses using computer programs such as MapMaker (Lander et al. (1987) Genomics 1: 174-181) in order to construct a genetic map. In addition, the nucleic acids may be used to probe Southern blots containing restriction endonuclease-treated genomic DNAs of a set of individuals representing parent and progeny of a defined genetic cross. Segregation of the DNA polymorphisms is noted and used to calculate the position of

the nucleic acid encoding the protein of interest in the genetic map previously obtained using this population (Botstein et al. (1980) *Am. J. Hum. Genet.* 32:314-331).

The production and use of plant gene-derived probes for use in genetic mapping is described in Bernatzky and Tanksley (1986) *Plant Mol. Biol. Reporter* 4: 37-41. Numerous publications describe genetic mapping of specific cDNA clones using the methodology outlined above or variations thereof. For example, F2 intercross populations, backcross populations, randomly mated populations, near isogenic lines, and other sets of individuals may be used for mapping. Such methodologies are well known to those skilled in the art.

The nucleic acid probes may also be used for physical mapping (i.e., placement of sequences on physical maps; see Hoheisel et al. In: *Non-mammalian Genomic Analysis: A Practical Guide*, Academic press 1996, pp. 319-346, and references cited therein).

In another embodiment, the nucleic acid probes may be used in direct fluorescence in situ hybridisation (FISH) mapping (Trask (1991) *Trends Genet.* 7:149-154). Although current methods of FISH mapping favour use of large clones (several kb to several hundred kb; see Laan et al. (1995) *Genome Res.* 5:13-20), improvements in sensitivity may allow performance of FISH mapping using shorter probes.

A variety of nucleic acid amplification-based methods for genetic and physical mapping may be carried out using the nucleic acids. Examples include allele-specific amplification (Kazazian (1989) *J. Lab. Clin. Med* 11:95-96), polymorphism of PCR-amplified fragments (CAPS; Sheffield et al. (1993) *Genomics* 16:325-332), allele-specific ligation (Landegren et al. (1988) *Science* 241:1077-1080), nucleotide extension reactions (Sokolov (1990) *Nucleic Acid Res.* 18:3671), Radiation Hybrid Mapping (Walter et al. (1997) *Nat. Genet.* 7:22-28) and Happy Mapping (Dear and Cook (1989) *Nucleic Acid Res.* 17:6795-6807). For these methods, the sequence of a nucleic acid is used to design and produce primer pairs for use in the amplification reaction or in primer extension reactions. The design of such primers is well known to those skilled in the art. In methods employing PCR-based genetic mapping, it may be necessary to identify DNA sequence differences between the parents of the mapping cross in the region corresponding to the instant nucleic acid sequence. This, however, is generally not necessary for mapping methods.

Plant

The term "plant" as used herein encompasses whole plants, ancestors and progeny of the plants and plant parts, including seeds, shoots, stems, leaves, roots (including tubers), flowers, and tissues and organs, wherein each of the aforementioned comprise the gene/nucleic acid of interest. The term "plant" also encompasses plant cells, suspension cultures, callus tissue, embryos, meristematic regions, gametophytes, sporophytes, pollen and microspores, again wherein each of the aforementioned comprises the gene/nucleic acid of interest.

Plants that are particularly useful in the methods of the invention include all plants which belong to the superfamily Viridiplantae, in particular monocotyledonous and dicotyledonous plants including fodder or forage legumes, ornamental plants, food crops, trees or shrubs selected from the list comprising *Acer* spp., *Actinidia* spp., *Abelmoschus* spp., *Agave sisalana*, *Agropyron* spp., *Agrostis stolonifera*, *Allium* spp., *Amaranthus* spp., *Ammophila arenaria*, *Ananas comosus*, *Annona* spp., *Apium graveolens*, *Arachis* spp., *Artocarpus* spp., *Asparagus officinalis*, *Avena* spp. (e.g. *Avena sativa*, *Avena fatua*, *Avena byzantina*, *Avena fatua* var. *sativa*, *Avena hybrida*), *Averrhoa carambola*, *Bambusa* sp., *Benincasa hispida*, *Bertholletia excelsa*, *Beta vulgaris*, *Brassica* spp. (e.g. *Brassica napus*, *Brassica rapa* ssp. [canola, oilseed rape, turnip rape]), *Cadaba farinosa*, *Camellia sinensis*, *Canna indica*, *Cannabis sativa*, *Capsicum* spp., *Carex elata*, *Carica papaya*, *Carissa macrocarpa*, *Carya* spp., *Carthamus tinctorius*, *Castanea* spp., *Ceiba pentandra*, *Cichorium endivia*, *Cinnamomum* spp., *Citrullus lanatus*, *Citrus* spp., *Cocos* spp., *Coffea* spp., *Colocasia esculenta*, *Cola* spp., *Corchorus* sp., *Coriandrum sativum*, *Corylus* spp., *Crataegus* spp., *Crocus sativus*, *Cucurbita* spp., *Cucumis* spp., *Cynara* spp., *Daucus carota*, *Desmodium* spp., *Dimocarpus longan*, *Dioscorea* spp., *Diospyros* spp., *Echinochloa* spp., *Elaeis* (e.g. *Elaeis guineensis*, *Elaeis oleifera*), *Eleusine coracana*, *Eragrostis tef*, *Erianthus* sp., *Eriobotrya japonica*, *Eucalyptus* sp., *Eugenia uniflora*, *Fagopyrum* spp., *Fagus* spp., *Festuca arundinacea*, *Ficus carica*, *Fortunella* spp., *Fragaria* spp., *Ginkgo biloba*, *Glycine* spp. (e.g. *Glycine max*, *Soja hispida* or *Soja max*), *Gossypium hirsutum*, *Helianthus* spp. (e.g. *Helianthus annuus*), *Hemerocallis fulva*, *Hibiscus* spp., *Hordeum* spp. (e.g. *Hordeum vulgare*), *Ipomoea batatas*, *Juglans* spp., *Lactuca sativa*, *Lathyrus* spp., *Lens culinaris*, *Linum usitatissimum*, *Litchi chinensis*, *Lotus* spp., *Luffa acutangula*, *Lupinus* spp., *Luzula sylvatica*, *Lycopersicon* spp. (e.g. *Lycopersicon esculentum*, *Lycopersicon lycopersicum*, *Lycopersicon pyriforme*), *Macrotyloma* spp., *Malus* spp., *Malpighia emarginata*, *Mammea americana*, *Mangifera indica*, *Manihot* spp., *Manilkara zapota*, *Medicago sativa*, *Melilotus* spp., *Mentha* spp., *Miscanthus sinensis*, *Momordica* spp., *Morus nigra*, *Musa* spp., *Nicotiana* spp., *Olea* spp., *Opuntia* spp., *Ornithopus* spp., *Oryza* spp. (e.g. *Oryza sativa*, *Oryza latifolia*), *Panicum miliaceum*, *Panicum virgatum*, *Passiflora edulis*, *Pastinaca sativa*, *Pennisetum* sp., *Persea* spp., *Petroselinum crispum*, *Phalaris arundinacea*, *Phaseolus* spp., *Phleum pratense*, *Phoenix* spp., *Phragmites australis*, *Physalis* spp., *Pinus* spp., *Pistacia vera*, *Pisum* spp., *Poa* spp., *Populus* spp., *Prosopis* spp., *Prunus* spp., *Psidium* spp., *Punica granatum*, *Pyrus communis*, *Quercus* spp., *Raphanus sativus*, *Rheum rhabarbarum*, *Ribes* spp., *Ricinus communis*, *Rubus* spp., *Saccharum* spp., *Salix* sp., *Sambucus* spp., *Secale cereale*, *Sesamum* spp., *Sinapis* sp., *Solanum* spp. (e.g. *Solanum tuberosum*, *Solanum integrifolium* or *Solanum lycopersicum*), *Sorghum bicolor*, *Spinacia* spp., *Syzygium* spp., *Tagetes* spp., *Tamarindus indica*, *Theobroma cacao*, *Trifolium* spp., *Tripsacum dactyloides*, *Triticosecale rimpau*, *Triticum* spp. (e.g. *Triticum aestivum*, *Triticum durum*, *Triticum turgidum*, *Triticum hybernum*, *Triticum macha*, *Triticum sativum*, *Triticum monococcum* or *Triticum vulgare*), *Tropaeolum minus*, *Tropaeolum majus*, *Vaccinium* spp.,

Vicia spp., Vigna spp., Viola odorata, Vitis spp., Zea mays, Zizania palustris, Ziziphus spp., amongst others.

Control plant(s)

The choice of suitable control plants is a routine part of an experimental setup and may include corresponding wild type plants or corresponding plants without the gene of interest. The control plant is typically of the same plant species or even of the same variety as the plant to be assessed. The control plant may also be a nullizygote of the plant to be assessed. Nullizygotes are individuals missing the transgene by segregation. A "control plant" as used herein refers not only to whole plants, but also to plant parts, including seeds and seed parts.

Detailed description of the invention

Surprisingly, it has now been found that modulating expression in a plant of a nucleic acid encoding an SGT1 polypeptide gives plants having enhanced yield-related traits relative to control plants. According to a first embodiment, the present invention provides a method for enhancing yield-related traits in plants relative to control plants, comprising modulating expression in a plant of a nucleic acid encoding an SGT1 polypeptide and optionally selecting for plants having enhanced yield-related traits.

The invention also provides hitherto unknown SGT1-encoding nucleic acids and SGT1 polypeptides useful for conferring enhanced yield-related traits in plants relative to control plants.

Furthermore, it has now surprisingly been found that modulating expression in a plant of a nucleic acid encoding a CLC-pKG polypeptide gives plants having enhanced yield-related traits relative to control plants. According to a first embodiment, the present invention provides a method for enhancing yield-related traits in plants relative to control plants, comprising modulating expression in a plant of a nucleic acid encoding a CLC-pKG polypeptide and optionally selecting for plants having enhanced yield-related traits.

Furthermore, it has now surprisingly been found that modulating expression in a plant of a nucleic acid encoding a HD-hydrolase-like polypeptide gives plants having enhanced yield-related traits relative to control plants. According to a first embodiment, the present invention provides a method for enhancing yield-related traits in plants relative to control plants, comprising modulating expression in a plant of a nucleic acid encoding a HD-hydrolase-like polypeptide and optionally selecting for plants having enhanced yield-related traits.

A preferred method for modulating (preferably, increasing) expression of a nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like

polypeptide, is by introducing and expressing in a plant a nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide respectively.

In one embodiment a “protein useful in the methods of the invention” is taken to mean an SGT1 polypeptide as defined herein. Any reference hereinafter to a “nucleic acid useful in the methods of the invention” is taken to mean a nucleic acid capable of encoding such an SGT1 polypeptide. The nucleic acid to be introduced into a plant (and therefore useful in performing the methods of the invention) is any nucleic acid encoding the type of protein which will now be described, hereinafter also named “SGT1 nucleic acid” or “SGT1 gene”.

An “SGT1 polypeptide” as defined herein refers to any polypeptide comprising the following in any order:

1. at least one tetratricopeptide (TPR) repeat;
2. at least one CS domain; and
3. an SGS domain.

The TPR is a structural motif involved in the mediation of protein-protein interactions and the assembly of multiprotein complexes. PFam Accession number PF00515 represents TPRs. See also Interpro Accession number IPR001440 for further information on TPRs. SGT1 polypeptides useful in the methods of the invention comprise at least one TPR, preferably two TPRs and most preferably three TPRs.

The first TPR is a Type 1 TPR (TPR1) and is represented by the sequence given below or by a sequence having in increasing order of preference at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity to the sequence of TPR1.

TPR1: SKAYLRKGLACMKLEEYQTAKAALETGASLAPGE (SEQ ID NO: 37)

The second TPR is a Type 2 TPR (TPR2) and is represented by the sequence given below or by a sequence having in increasing order of preference at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity to the sequence of TPR2a.

TPR2a: ASDLETRAKEAFIDDFELAVDLYTQAISLSPKN (SEQ ID NO: 38)

The third TPR is also a Type 2 TPR (TPR2) is represented by the sequence given below or by a sequence having in increasing order of preference at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%,

90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity to the sequence of TPR2b.

TPR2b: PELFADRAQANIKLNYFTEAVVDANKAIELDPYM (SEQ ID NO: 39)

PFam Accession number PF04969 represents CS domains. See also Interpro Accession number IPR017447 for further information on CS domains. The CS domain is represented by the sequence below or by a sequence having in increasing order of preference at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity to the sequence below.

CS Domain: YRHEFYQKPEEVVVTIFAKGIPAKNVVDFGEQILSVSIDLPGGETYSFQPRLF GKITPAKCRYEVMSTKIEIRLAK (SEQ ID NO: 40)

PFam Accession number PF05002 represents SGS domains. See also Interpro Accession number IPR007699 for further information on SGS domains. The SGS domain is represented by the sequence below or by a sequence having in increasing order of preference at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity to the sequence below.

SGS Domain: NVDWDKLEAQVKKEEKDEKLDGDAALNKFFRDIYKDADEDTRRAMMKSF VESNGTVLSTNWKVKVGTKKVEGSPDGMELKKW (SEQ ID NO: 41)

According to a preferred feature of the present invention, SGT1 polypeptides useful in the methods of the invention may comprise any one or more of Motifs 1 to 8 as defined below:

Motif 1: DLETRAKEAFIDDFELAVDLYTQAI, or a sequence having in increasing order of preference at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity to the sequence of Motif 1 (SEQ ID NO: 42).

Motif 2: ADRAQANIKL, or a sequence having in increasing order of preference at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity to the sequence of Motif 2 (SEQ ID NO: 43).

Motif 3: LEEYQTAKAALETGAS, or a sequence having in increasing order of preference at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity to the sequence of Motif 3 (SEQ ID NO: 44).

Motif 4: KYRHEFYQKP EEEVV TIFAKGIP, or a sequence having in increasing order of preference at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity to the sequence of Motif 4 (SEQ ID NO: 45).

Motif 5: QPRLFGKITPAKCRYEVMSTKIEIRLAKAE, or a sequence having in increasing order of preference at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity to the sequence of Motif 5 (SEQ ID NO: 46).

Motif 6: EAQVKKEEKDEKLDGDA, or a sequence having in increasing order of preference at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity to the sequence of Motif 6 (SEQ ID NO: 47).

Motif 7: ALNKFFRDIYKDADEDTRAMMKSFVESNGTVLSTNWKKVGTKKVEG, or a sequence having in increasing order of preference at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity to the sequence of Motif 7 (SEQ ID NO: 48).

Motif 8: SPPDGMELKKWEI, or a sequence having in increasing order of preference at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more sequence identity to the sequence of Motif 8 (SEQ ID NO: 49).

Preferably, the SGT1 polypeptide comprises in increasing order of preference, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7 or all 8 of the motifs as defined above.

Additionally, SGT1 polypeptides typically have in increasing order of preference at least 25%, 26%, 27%, 28%, 29%, 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% overall sequence identity to the amino acid represented by SEQ ID NO: 2.

The overall sequence identity is determined using a global alignment algorithm, such as the Needleman Wunsch algorithm in the program GAP (GCG Wisconsin Package, Accelrys), preferably with default parameters and preferably with sequences of mature proteins (i.e. without taking into account secretion signals or transit peptides). Compared to overall

sequence identity, the sequence identity will generally be higher when only conserved domains or motifs are considered.

Preferably, the polypeptide sequence which when used in the construction of a phylogenetic tree, such as the one depicted in Figure 2, clusters with the group of SGT1 polypeptides comprising the amino acid sequence represented by SEQ ID NO: 2 rather than with any other group.

In another embodiment a "protein useful in the methods of the invention" is taken to mean a CLC-pKG polypeptide as defined herein. Any reference hereinafter to a "nucleic acid useful in the methods of the invention" is taken to mean a nucleic acid capable of encoding such a CLC-pKG polypeptide. The nucleic acid to be introduced into a plant (and therefore useful in performing the methods of the invention) is any nucleic acid encoding the type of protein which will now be described, hereafter also named "CLC-pKG nucleic acid" or "CLC-pKG gene".

A "CLC-pKG polypeptide" as defined herein refers to any polypeptide comprising a Voltage CLC domain (Pfam entry PF00654) and C-terminally thereof a CBS domain (Pfam entry PF00571; Interpro entry IPR000644) and optionally a USP domain (Pfam entry PF00582; Interpro entry IPR006016). CLC-pKG polypeptides are part of a family of voltage-gated channels that is highly conserved over various kingdoms of life and are structurally unrelated to the other known voltage-gated channels. They are found in organisms ranging from bacteria to yeasts and plants, and also in animals. CLC-pKG polypeptides typically have 10 or 12 transmembrane (TM) domains, occasionally less than 10 TM domains or more than 12, up to 18 or 20 TM domains may occur. The CLC-pKG polypeptide represented by the Arabidopsis AtCLCe and AtCLCf proteins show up to 12 membrane-spanning domains Marmagne et al. 2007. CBS (cystathionine-beta-synthase) domains are small intracellular modules, mostly found in two or four copies within a protein (two copies for CLC-pKG proteins), and are present in several different proteins in all kingdoms of life. Tandem pairs of CBS domains, also called Baeteman domains, can act as binding domains for adenosine derivatives and may regulate the activity of attached enzymatic or other domains. Typically CBS domains in CLC-pKG polypeptides are proposed to play a role as a sensor for the energy status of the cell. USP (Universal Stress Protein) domains are conserved protein regions present in proteins whose expression in nature is typically enhanced when the cell is exposed to stress agents. It was first characterized in the universal stress protein UspA (Nystrom T, Neidhardt FC. Mol. Microbiol. 11 537-44 1994), which is proposed to have a a general "stress endurance" activity in E. coli.

Preferably, the CLC-pKG polypeptide useful in the methods of the present inventions comprises a domain having in increasing order of preference at least 25%, 26%, 27%, 28%, 29%, 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%,

61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100 % sequence identity to:

- (i) a CLC in SEQ ID NO: 54 as represented by the sequence located between amino acids 77 and 422 of SEQ ID NO: 54;
- (ii) a CBS domain in SEQ ID NO: 54 as represented by the sequence located between amino acids 455 to 510 or 516 to 571 of SEQ ID NO: 54; and optionally
- (iii) a USP domain in SEQ ID NO: 54 as represented by the sequence located between amino acids 597-731 of SEQ ID NO: 54.

Preferably, the CLC-pKG polypeptide useful in the methods of the present inventions comprises one or more of motifs 9 or 10:

- (i) Motif 9: P(T/S)(V/I)(H/Q)(I/V)GA(A/S/G)LA;
- (ii) Motif 10: (P/S)GG(I/L/V)(F/L)XP(A/S/T)LX(L/M)G(S/A/T)XLG

Wherein up to one, two, three, four, five, six, seven amino acids maybe substituted by any amino acid preferably by a semiconservative amino acid more preferably by a conservative amino acid.

Additionally or alternatively, the CLC-pKG polypeptide useful in the methods of the invention or a homologue thereof has in increasing order of preference at least 25%, 26%, 27%, 28%, 29%, 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% overall sequence identity to the amino acid represented by any one of the polypeptides of Table A2, preferably by SEQ ID NO: 54, provided that the polypeptide comprises the conserved domains as outlined above. The overall sequence identity is determined using a global alignment algorithm, such as the Needleman Wunsch algorithm in the program GAP (GCG Wisconsin Package, Accelrys), preferably with default parameters and preferably with sequences of mature proteins (i.e. without taking into account secretion signals or transit peptides). Compared to overall sequence identity, the sequence identity will generally be higher when only conserved domains or motifs are considered.

In a preferred embodiment the CLC-pKG nucleic acid and/or polypeptide useful in the methods of the invention is of natural origin, more of plant origin, most preferably of prokaryotic origin. Preferred plant CLC-pKG polypeptides are AtCIC-e (At4g35440) and AtCIC-f (At1g55620) as described by Marmagne et al 2007 and having the locus number or Spinach CLCf as described by Teardo et al. 2005 FEBS Letters 579 (2005) 4991–4996. As described by Teardo et al. 2005, AtCIC-f and AtCIC-e are close related in sequence structure to cyanobacterial CIC channels (See Table 1 of Teardo et al. 2005). in particular the similarity with the the *Synechocystis* sp. PCC 6803 protein, sl1864, the sequence

similarity is of 20% with AtCLC-e and of 21% with AtCLC-f. Preferred prokaryotic CLC-pKG polypeptides are the polypeptides of *Synechocystis* species, more preferably of *Synechocystis* sp. PCC 6803, most preferably polypeptide sll1864 of *Synechocystis* sp. PCC 6803, herein provided under SEQ ID NO: 54.

Alternatively or additionally, the CLC-pKG polypeptide sequence useful in the methods of the invention when used in the construction of a phylogenetic tree, such as the one depicted in Figure 1 of Marmagne et al. 2007 (herein incorporated by reference), clusters with the group of CLC polypeptides of bacterial origin, preferably with SCLC (P74477) from *Synechocystis* sp.; or with the group of plant origin comprising AtCLCe and AtCLCf than with any other anion group. Figure 1 of Marmagne et al. 2007 is given in Figure 6 herein.

Furthermore, CLC-like polypeptides (at least in their native form) typically have nitrate/H⁺ exchanging activity. Tools and techniques for measuring anion transporting activity are well known in the art, for example by complementation of yeast mutants (Marmagne et al., J. Exp. Bot. 58, 3385-3393, 2007) or complementation of *Arabidopsis* mutants (De Angeli, 2009). Further details are provided in the Example section.

Selectivity of CLC chloride channels for different anions vary amongst classes of CLC proteins. Thus AtCLCa of *Arabidopsis thaliana*, closer to CLC proteins of eukaryotic origin (see Figure 6) was shown to have a higher selectivity for NO₃⁻ anions than for Chloride, Cl⁻. A further preferred CLC-pKG polypeptide useful in the methods of the invention has a higher selectivity for Chloride than for Nitrate anions.

Plant CLC polypeptides are localized to membranes, with different classes of CLC localized to different membranes, for example to the plasma, the tonoplast, the chloroplast or the golgi membranes. A yet further preferred CLC-pKG polypeptide useful in the methods of the invention localizes to the Golgi or to the Thylakoid membranes when expressed in a plant cell.

In addition or alternatively, CLC-pKG polypeptides, when expressed in rice according to the methods of the present invention as outlined in the Examples section herein give plants having increased yield related traits in comparison to control plants, in particular an increase in any one or more of seed yield, harvest index, height of the plant, leaf biomass when grown under non- stress conditions.

In yet another embodiment a "protein useful in the methods of the invention" is taken to mean a HD-hydrolase-like polypeptide as defined herein. Any reference hereinafter to a "nucleic acid useful in the methods of the invention" is taken to mean a nucleic acid capable of encoding such a HD-hydrolase-like polypeptide. The nucleic acid to be introduced into a plant (and therefore useful in performing the methods of the invention) is any nucleic acid

encoding the type of protein which will now be described, hereafter also named "HD-hydrolase-like nucleic acid" or "HD-hydrolase-like gene".

A "HD-hydrolase-like polypeptide" as defined herein refers to a protein comprising the CRISPR-associated protein Crm2 domain (InterPro accession IPR013407). Preferably the HD-hydrolase-like protein comprises one or more of the following motifs:

Motif 11 (SEQ ID NO: 148): (A/G/S/Q)(R/Y/F/L)(L/F/N)H(D/S)(P/S)(A/P/L)

Preferably, motif 11 is (G/A)LLHDPA

Motif 12 (SEQ ID NO: 149): P(A/Q)(D/E/H)(T/Y/S)(R/I)(I/F/A/L)P(D/N)

Preferably, motif 12 is PA(D/E)TR(I/F/A/L)P(D/N)

Motif 13 (SEQ ID NO: 150): (T/S)(L/I/F)(G/S/N)P(V/I)Q(E/S/Q/D/R)(L/F/Y)I

Preferably, motif 13 is (T/S)(L/I/F)(G/S)P(V/I)Q(E/S)(L/F/Y)I

Motif 14 (SEQ ID NO: 151): D(L/F)(W/Y/F)(A/W/M/T)(G/S)S(H/Y/W/F/M)(L/I/M)L(A/S/H)(R/Y/D)(L/F).

Preferably, motif 14 is D(L/F)(W/Y/F)(A/W/M/T)GS(H/Y/W)(L/I/M)L(A/S/H)(R/Y)(L/F)

Motif 15 (SEQ ID NO: 152): (L/I/V)(F/Y/T)PXL

Wherein X can be any amino acid but preferably one of Q, R, E, N, S, C or D

Motif 16 (SEQ ID NO: 153): (L/M/F/I)PN(R/V/T)(F/I/L)

Motif 17 (SEQ ID NO: 154): (M/L/G/F)DGD(X)(M/I)

Wherein X can be any amino acid but preferably one of R, Y, N, H, G, S or D

Preferably, motif 7 is GDGD(G/S/D)M

Motif 18 (SEQ ID NO: 155): (S/T/G/A)P(A/G/S/T)(R/V/L/T/I)H(L/S/Q/N/V/A)(A/L/T/G/F)(I/L/V)(S/N)(G/E/R/L)

Motif 19 (SEQ ID NO: 156): G(R/K)(V/L)(L/I/V)Y(A/S)GGDD(L/V)(M/F/L)(A/T)

Motif 20 (SEQ ID NO: 157): (T/S)(A/M/I/L)S(M/A/F/C/L)G(A/V/I/L)(V/T/A)I(A/V/G)(H/D)(H/Y/K)(K/Q/S)

Additionally or alternatively, the HD-hydrolase-like polypeptide comprises one or more of the following motifs:

Motif 21 (SEQ ID NO: 158): IKASRK(F/I/M)(L/R/K)DFWAGS(Y/W)(L/I)LHYLSA

Motif 22 (SEQ ID NO: 159): L(A/L)TAGFPN

Motif 23 (SEQ ID NO: 160): GLFDG(S/T/R/I)E(M/K/Q)LNA(L/T)E(L/I/T)(T/V/L)KR

Motif 24 (SEQ ID NO: 161): KRMGP(A/S)TH(V/S/N)(G/A)L(N/S)RALLDFS

Motif 25 (SEQ ID NO: 162): Y(S/A)GGDDV

Motif 26 (SEQ ID NO: 163): LFT(M/L)G(F/K/S/Q/D)(G/C/L/A/N)A(T/S)(M/I)S(L/F)G(I/V/L)(V/T)I(V/A)(D/H)(K/H)S(V/I)PL(P/A)(V/I)ALE(A/S/N)LW(E/Q)AE

More preferably, the HD-hydrolase-like polypeptide comprises in increasing order of preference, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, or all 16 motifs.

Additionally or alternatively, the homologue of a HD-hydrolase-like protein has in increasing order of preference at least 25%, 26%, 27%, 28%, 29%, 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% overall sequence identity to the amino acid represented by SEQ ID NO: 119, provided that the homologous protein comprises any one or more of the conserved motifs as outlined above. The overall sequence identity is determined using a global alignment algorithm, such as the Needleman Wunsch algorithm in the program GAP (GCG Wisconsin Package, Accelrys), preferably with default parameters and preferably with sequences of mature proteins (i.e. without taking into account secretion signals or transit peptides). Compared to overall sequence identity, the sequence identity will generally be higher when only conserved domains or motifs are considered. Preferably the motifs in a HD-hydrolase-like polypeptide have, in increasing order of preference, at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to any one or more of the motifs represented by SEQ ID NO: 148 to SEQ ID NO: 163 (Motifs 11 to 26).

Preferably, the polypeptide sequence which when used in the construction of a phylogenetic tree, such as the one depicted in Figure 10, clusters with the group of HD-hydrolase-like polypeptides comprising the amino acid sequence represented by SEQ ID NO: 119 rather than with any other group.

HD-hydrolase-like polypeptides, when expressed in rice according to the methods of the present invention as outlined in Examples 6 and 7, give plants having increased yield related traits, such as increased seed yield.

The terms “domain”, “signature” and “motif” are defined in the “definitions” section herein.

Concerning SGT1 polypeptides, the present invention is illustrated by transforming plants with the nucleic acid sequence represented by SEQ ID NO: 1, encoding the polypeptide sequence of SEQ ID NO: 2. However, performance of the invention is not restricted to these sequences; the methods of the invention may advantageously be performed using any SGT1-encoding nucleic acid or SGT1 polypeptide as defined herein.

Examples of nucleic acids encoding SGT1 polypeptides are given in Table A1 of the Examples section herein. Such nucleic acids are useful in performing the methods of the invention. The amino acid sequences given in Table A1 of the Examples section are example sequences of orthologues and paralogues of the SGT1 polypeptide represented by SEQ ID NO: 2, the terms “orthologues” and “paralogues” being as defined herein. Further orthologues and paralogues may readily be identified by performing a so-called

reciprocal blast search as described in the definitions section; where the query sequence is SEQ ID NO: 1 or SEQ ID NO: 2, the second BLAST (back-BLAST) would be against Capsicum sequences.

Concerning CLC-pKG polypeptides, the present invention is illustrated by transforming plants with the nucleic acid sequence represented by SEQ ID NO: 53, encoding the polypeptide sequence of SEQ ID NO: 54. However, performance of the invention is not restricted to these sequences; the methods of the invention may advantageously be performed using any CLC-pKG-encoding nucleic acid or CLC-pKG polypeptide as defined herein.

Examples of nucleic acids encoding CLC-pKG polypeptides are given in Table A2 of the Examples section herein. Such nucleic acids are useful in performing the methods of the invention. The amino acid sequences given in Table A2 of the Examples section are example sequences of orthologues and paralogues of the CLC-pKG polypeptide represented by SEQ ID NO: 54, the terms "orthologues" and "paralogues" being as defined herein. Further orthologues and paralogues may readily be identified by performing a so-called reciprocal blast search as described in the definitions section; where the query sequence is SEQ ID NO: 53 or SEQ ID NO: 54, the second BLAST (back-BLAST) would be against rice sequences.

Concerning HD-hydrolase-like polypeptides, the present invention is illustrated by transforming plants with the nucleic acid sequence represented by SEQ ID NO: 118, encoding the polypeptide sequence of SEQ ID NO: 119. However, performance of the invention is not restricted to these sequences; the methods of the invention may advantageously be performed using any HD-hydrolase-like-encoding nucleic acid or HD-hydrolase-like polypeptide as defined herein.

Examples of nucleic acids encoding HD-hydrolase-like polypeptides are given in Table A3 of the Examples section herein. Such nucleic acids are useful in performing the methods of the invention. The amino acid sequences given in Table A3 of the Examples section are example sequences of orthologues and paralogues of the HD-hydrolase-like polypeptide represented by SEQ ID NO: 119, the terms "orthologues" and "paralogues" being as defined herein. Further orthologues and paralogues may readily be identified by performing a so-called reciprocal blast search as described in the definitions section; where the query sequence is SEQ ID NO: 118 or SEQ ID NO: 119, the second BLAST (back-BLAST) would be against Synechocystis sequences.

Nucleic acid variants may also be useful in practising the methods of the invention. Examples of such variants include nucleic acids encoding homologues and derivatives of any one of the amino acid sequences given in Table A1 to A3 of the Examples section, the terms "homologue" and "derivative" being as defined herein. Also useful in the methods of

the invention are nucleic acids encoding homologues and derivatives of orthologues or paralogues of any one of the amino acid sequences given in Table A1 to A3 of the Examples section. Homologues and derivatives useful in the methods of the present invention have substantially the same biological and functional activity as the unmodified protein from which they are derived. Further variants useful in practising the methods of the invention are variants in which codon usage is optimised or in which miRNA target sites are removed.

Further nucleic acid variants useful in practising the methods of the invention include portions of nucleic acids encoding SGT1 polypeptides, or CLC-pKG polypeptides, or HD-hydrolase-like polypeptides, nucleic acids hybridising to nucleic acids encoding SGT1 polypeptides, or CLC-pKG polypeptides, or HD-hydrolase-like polypeptides, splice variants of nucleic acids encoding SGT1 polypeptides, or CLC-pKG polypeptides, or HD-hydrolase-like polypeptides, allelic variants of nucleic acids encoding SGT1 polypeptides and variants of nucleic acids encoding SGT1 polypeptides, or CLC-pKG polypeptides, or HD-hydrolase-like polypeptides, obtained by gene shuffling. The terms hybridising sequence, splice variant, allelic variant and gene shuffling are as described herein.

Nucleic acids encoding SGT1 polypeptides, or CLC-pKG polypeptides, or HD-hydrolase-like polypeptides, need not be full-length nucleic acids, since performance of the methods of the invention does not rely on the use of full-length nucleic acid sequences. According to the present invention, there is provided a method for enhancing yield-related traits in plants, comprising introducing and expressing in a plant a portion of any one of the nucleic acid sequences given in Table A1 to A3 of the Examples section, or a portion of a nucleic acid encoding an orthologue, paralogue or homologue of any of the amino acid sequences given in Table A1 to A3 of the Examples section.

A portion of a nucleic acid may be prepared, for example, by making one or more deletions to the nucleic acid. The portions may be used in isolated form or they may be fused to other coding (or non-coding) sequences in order to, for example, produce a protein that combines several activities. When fused to other coding sequences, the resultant polypeptide produced upon translation may be bigger than that predicted for the protein portion.

Concerning SGT1 polypeptides, portions useful in the methods of the invention, encode a SGT1 polypeptide as defined herein, and have substantially the same biological activity as the amino acid sequences given in Table A1 of the Examples section. Preferably, the portion is a portion of any one of the nucleic acids given in Table A1 of the Examples section, or is a portion of a nucleic acid encoding an orthologue or paralogue of any one of the amino acid sequences given in Table A1 of the Examples section. Preferably the portion is at least 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000 consecutive nucleotides in length, the consecutive nucleotides being of any one of the nucleic acid sequences given in Table A1 of the Examples section, or of a nucleic acid encoding an

orthologue or paralogue of any one of the amino acid sequences given in Table A1 of the Examples section. Most preferably the portion is a portion of the nucleic acid of SEQ ID NO: 1. Preferably, the portion encodes a fragment of an amino acid sequence which, when used in the construction of a phylogenetic tree, clusters with the group of SGT1 polypeptides comprising the amino acid sequence represented by SEQ ID NO: 2 rather than with any other.

Concerning CLC-pKG polypeptides, portions useful in the methods of the invention, encode a CLC-pKG polypeptide as defined herein, and have substantially the same biological activity as the amino acid sequences given in Table A2 of the Examples section. Preferably, the portion is a portion of any one of the nucleic acids given in Table A2 of the Examples section, or is a portion of a nucleic acid encoding an orthologue or paralogue of any one of the amino acid sequences given in Table A2 of the Examples section. Preferably the portion is at least 100, 200, 300, 400, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000 consecutive nucleotides in length, the consecutive nucleotides being of any one of the nucleic acid sequences given in Table A2 of the Examples section, or of a nucleic acid encoding an orthologue or paralogue of any one of the amino acid sequences given in Table A2 of the Examples section. Most preferably the portion is a portion of the nucleic acid of SEQ ID NO: 53. Preferably, the portion encodes a fragment of an amino acid sequence which, when used in the construction of a phylogenetic tree, such as the one depicted in Figure 1 of Marmagne et al. 2007 (herein incorporated by reference), clusters with the group of CLC polypeptides of bacterial origin, preferably with SCLC (P74477) from *Synechocystis* sp.; or with the group of plant origin comprising AtCLCe and AtCLCf than with any other anion group.

Concerning HD-hydrolase-like polypeptides, portions useful in the methods of the invention, encode a HD-hydrolase-like polypeptide as defined herein, and have substantially the same biological activity as the amino acid sequences given in Table A3 of the Examples section. Preferably, the portion is a portion of any one of the nucleic acids given in Table A3 of the Examples section, or is a portion of a nucleic acid encoding an orthologue or paralogue of any one of the amino acid sequences given in Table A3 of the Examples section. Preferably the portion is at least 1000, 1100, 1200, 1300, 1400, 1500, 1600, 1700, 1800, 1900, 2000, 2100, 2200, 2300, 2400, 2500, 2600, 2700, 2800, 2900, 3000, 3100, 3200, 3300, 3400, 3500, 3600, 3700 consecutive nucleotides in length, the consecutive nucleotides being of any one of the nucleic acid sequences given in Table A3 of the Examples section, or of a nucleic acid encoding an orthologue or paralogue of any one of the amino acid sequences given in Table A3 of the Examples section. Most preferably the portion is a portion of the nucleic acid of SEQ ID NO: 118. Preferably, the portion encodes a fragment of an amino acid sequence which, when used in the construction of a phylogenetic tree, such as the one depicted in Figure 10, clusters with the group of HD-hydrolase-like polypeptides comprising the amino acid sequence represented by SEQ ID

NO: 119 rather than with any other group and/or comprises any one or more of the motifs 11 to 26 defined above.

Another nucleic acid variant useful in the methods of the invention is a nucleic acid capable of hybridising, under reduced stringency conditions, preferably under stringent conditions, with a nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, as defined herein, or with a portion as defined herein.

According to the present invention, there is provided a method for enhancing yield-related traits in plants, comprising introducing and expressing in a plant a nucleic acid capable of hybridizing to any one of the nucleic acids given in Table A1 to A3 of the Examples section, or comprising introducing and expressing in a plant a nucleic acid capable of hybridising to a nucleic acid encoding an orthologue, paralogue or homologue of any of the nucleic acid sequences given in Table A1 to A3 of the Examples section.

Concerning SGT1 polypeptides, hybridising sequences useful in the methods of the invention encode an SGT1 polypeptide as defined herein, having substantially the same biological activity as the amino acid sequences given in Table A1 of the Examples section. Preferably, the hybridising sequence is capable of hybridising to the complement of any one of the nucleic acids given in Table A1 of the Examples section, or to a portion of any of these sequences, a portion being as defined above, or the hybridising sequence is capable of hybridising to the complement of a nucleic acid encoding an orthologue or paralogue of any one of the amino acid sequences given in Table A1 of the Examples section. Most preferably, the hybridising sequence is capable of hybridising to the complement of a nucleic acid as represented by SEQ ID NO: 1 or to a portion thereof.

Preferably, the hybridising sequence encodes a polypeptide with an amino acid sequence which, when substantially full-length and used in the construction of a phylogenetic tree, clusters with the group of SGT1 polypeptides comprising the amino acid sequence represented by SEQ ID NO: 2 rather than with any other.

Concerning CLC-pKG polypeptides, hybridising sequences useful in the methods of the invention encode a CLC-pKG polypeptide as defined herein, having substantially the same biological activity as the amino acid sequences given in Table A2 of the Examples section. Preferably, the hybridising sequence is capable of hybridising to the complement of any one of the nucleic acids given in Table A2 of the Examples section, or to a portion of any of these sequences, a portion being as defined above, or the hybridising sequence is capable of hybridising to the complement of a nucleic acid encoding an orthologue or paralogue of any one of the amino acid sequences given in Table A2 of the Examples section. Most preferably, the hybridising sequence is capable of hybridising to the complement of a nucleic acid as represented by SEQ ID NO: 53 or to a portion thereof.

Preferably, the hybridising sequence encodes a polypeptide with an amino acid sequence which, when full-length and used in the construction of a phylogenetic tree, such as the one depicted in Figure 1 of Marmagne et al. 2007 (herein incorporated by reference), clusters with the group of CLC polypeptides of bacterial origin, preferably with SCLC (P74477) from *Synechocystis* sp.; or with the group of plant origin comprising AtCLCe and AtCLCf than with any other anion group.

Concerning HD-hydrolase-like polypeptides, hybridising sequences useful in the methods of the invention encode a HD-hydrolase-like polypeptide as defined herein, having substantially the same biological activity as the amino acid sequences given in Table A3 of the Examples section. Preferably, the hybridising sequence is capable of hybridising to the complement of any one of the nucleic acids given in Table A3 of the Examples section, or to a portion of any of these sequences, a portion being as defined above, or the hybridising sequence is capable of hybridising to the complement of a nucleic acid encoding an orthologue or paralogue of any one of the amino acid sequences given in Table A3 of the Examples section. Most preferably, the hybridising sequence is capable of hybridising to the complement of a nucleic acid as represented by SEQ ID NO: 118 or to a portion thereof.

Preferably, the hybridising sequence encodes a polypeptide with an amino acid sequence which, when full-length and used in the construction of a phylogenetic tree, such as the one depicted in Figure 10, clusters with the group of HD-hydrolase-like polypeptides comprising the amino acid sequence represented by SEQ ID NO: 119 rather than with any other group and/or comprises any one or more of the motifs 11 to 26 defined above.

Another nucleic acid variant useful in the methods of the invention is a splice variant encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, as defined hereinabove, a splice variant being as defined herein.

According to the present invention, there is provided a method for enhancing yield-related traits in plants, comprising introducing and expressing in a plant a splice variant of any one of the nucleic acid sequences given in Table A1 to A3 of the Examples section, or a splice variant of a nucleic acid encoding an orthologue, paralogue or homologue of any of the amino acid sequences given in Table A1 to A3 of the Examples section.

Concerning SGT1 polypeptides, preferred splice variants are splice variants of a nucleic acid represented by SEQ ID NO: 1, or a splice variant of a nucleic acid encoding an orthologue or paralogue of SEQ ID NO: 2. Preferably, the amino acid sequence encoded by the splice variant, when used in the construction of a phylogenetic tree, clusters with the group of SGT1 polypeptides comprising the amino acid sequence represented by SEQ ID NO: 2 rather than with any other group.

Concerning CLC-pKG polypeptides, preferred splice variants are splice variants of a nucleic acid represented by SEQ ID NO: 53, or a splice variant of a nucleic acid encoding an orthologue or paralogue of SEQ ID NO: 54. Preferably, the amino acid sequence encoded by the splice variant, when used in the construction of a phylogenetic tree, such as the one depicted in Figure 1 of Marmagne et al. 2007 (herein incorporated by reference), clusters with the group of CLC polypeptides of bacterial origin, preferably with SCLC (P74477) from *Synechocystis* sp.; or with the group of plant origin comprising AtCLCe and AtCLCf than with any other anion group.

Concerning HD-hydrolase-like polypeptides, preferred splice variants are splice variants of a nucleic acid represented by SEQ ID NO: 118, or a splice variant of a nucleic acid encoding an orthologue or paralogue of SEQ ID NO: 119. Preferably, the amino acid sequence encoded by the splice variant, when used in the construction of a phylogenetic tree, such as the one depicted in Figure 10, clusters with the group of HD-hydrolase-like polypeptides comprising the amino acid sequence represented by SEQ ID NO: 119 rather than with any other group and/or comprises any one or more of the motifs 11 to 26 defined above.

Another nucleic acid variant useful in performing the methods of the invention is an allelic variant of a nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, as defined hereinabove, an allelic variant being as defined herein.

According to the present invention, there is provided a method for enhancing yield-related traits in plants, comprising introducing and expressing in a plant an allelic variant of any one of the nucleic acids given in Table A1 to A3 of the Examples section, or comprising introducing and expressing in a plant an allelic variant of a nucleic acid encoding an orthologue, paralogue or homologue of any of the amino acid sequences given in Table A1 to A3 of the Examples section.

Concerning SGT1 polypeptides, the polypeptides encoded by allelic variants useful in the methods of the present invention have substantially the same biological activity as the SGT1 polypeptide of SEQ ID NO: 2 and any of the amino acids depicted in Table A1 of the Examples section. Allelic variants exist in nature, and encompassed within the methods of the present invention is the use of these natural alleles. Preferably, the allelic variant is an allelic variant of SEQ ID NO: 1 or an allelic variant of a nucleic acid encoding an orthologue or paralogue of SEQ ID NO: 2. Preferably, the amino acid sequence encoded by the allelic variant, when used in the construction of a phylogenetic tree, clusters with the SGT1 polypeptides comprising the amino acid sequence represented by SEQ ID NO: 2 rather than with any other group.

Concerning CLC-pKG polypeptides, the polypeptides encoded by allelic variants useful in the methods of the present invention have substantially the same biological activity as the CLC-pKG polypeptide of SEQ ID NO: 54 and any of the amino acids depicted in Table A2 of the Examples section. Allelic variants exist in nature, and encompassed within the methods of the present invention is the use of these natural alleles. Preferably, the allelic variant is an allelic variant of SEQ ID NO: 53 or an allelic variant of a nucleic acid encoding an orthologue or paralogue of SEQ ID NO: 54. Preferably, the amino acid sequence encoded by the allelic variant, when used in the construction of a phylogenetic tree, such as the one depicted in Figure 1 of Marmagne et al. 2007 (herein incorporated by reference), clusters with the group of CLC polypeptides of bacterial origin, preferably with SCLC (P74477) from *Synechocystis* sp.; or with the group of plant origin comprising AtCLCe and AtCLCf than with any other anion group.

Concerning HD-hydrolase-like polypeptides, the polypeptides encoded by allelic variants useful in the methods of the present invention have substantially the same biological activity as the HD-hydrolase-like polypeptide of SEQ ID NO: 119 and any of the amino acids depicted in Table A3 of the Examples section. Allelic variants exist in nature, and encompassed within the methods of the present invention is the use of these natural alleles. Preferably, the allelic variant is an allelic variant of SEQ ID NO: 118 or an allelic variant of a nucleic acid encoding an orthologue or paralogue of SEQ ID NO: 119. Preferably, the amino acid sequence encoded by the allelic variant, when used in the construction of a phylogenetic tree, such as the one depicted in Figure 10, clusters with the group of HD-hydrolase-like polypeptides comprising the amino acid sequence represented by SEQ ID NO: 119 rather than with any other group and/or comprises any one or more of the motifs 11 to 26 defined above.

Gene shuffling or directed evolution may also be used to generate variants of nucleic acids encoding SGT1 polypeptides, or CLC-pKG polypeptides, or HD-hydrolase-like polypeptides, as defined above; the term "gene shuffling" being as defined herein.

According to the present invention, there is provided a method for enhancing yield-related traits in plants, comprising introducing and expressing in a plant a variant of any one of the nucleic acid sequences given in Table A1 to A3 of the Examples section, or comprising introducing and expressing in a plant a variant of a nucleic acid encoding an orthologue, paralogue or homologue of any of the amino acid sequences given in Table A1 to A3 of the Examples section, which variant nucleic acid is obtained by gene shuffling.

Concerning SGT1 polypeptides, preferably, the amino acid sequence encoded by the variant nucleic acid obtained by gene shuffling, when used in the construction of a phylogenetic tree, clusters with the group of SGT1 polypeptides comprising the amino acid sequence represented by SEQ ID NO: 2 rather than with any other group.

Concerning CLC-pKG polypeptides, preferably, the amino acid sequence encoded by the variant nucleic acid obtained by gene shuffling, when used in the construction of a phylogenetic tree such as the one depicted in Figure 1 of Marmagne et al. 2007 (herein incorporated by reference), clusters with the group of CLC polypeptides of bacterial origin, preferably with SCLC (P74477) from *Synechocystis* sp.; or with the group of plant origin comprising AtCLCe and AtCLCf than with any other anion group.

Concerning HD-hydrolase-like polypeptides, preferably, the amino acid sequence encoded by the variant nucleic acid obtained by gene shuffling, when used in the construction of a phylogenetic tree, such as the one depicted in Figure 10, clusters with the group of HD-hydrolase-like polypeptides comprising the amino acid sequence represented by SEQ ID NO: 119 rather than with any other group and/or comprises any one or more of the motifs 11 to 26 defined above.

Furthermore, nucleic acid variants may also be obtained by site-directed mutagenesis. Several methods are available to achieve site-directed mutagenesis, the most common being PCR based methods (Current Protocols in Molecular Biology. Wiley Eds.).

Nucleic acids encoding SGT1 polypeptides may be derived from any natural or artificial source. The nucleic acid may be modified from its native form in composition and/or genomic environment through deliberate human manipulation. Preferably the SGT1 polypeptide-encoding nucleic acid is from a plant, further preferably from pepper, most preferably the nucleic acid is from *Capsicum annuum*.

Nucleic acids encoding CLC-pKG polypeptides may be derived from any natural or artificial source. The nucleic acid may be modified from its native form in composition and/or genomic environment through deliberate human manipulation. Preferably the CLC-pKG polypeptide-encoding nucleic acid is from a cyanobacterial origin, further preferably from *Synechocystis* species, most preferably from *Synechocystis* sp. PCC 6803.

Nucleic acids encoding HD-hydrolase-like polypeptides may be derived from any natural or artificial source. The nucleic acid may be modified from its native form in composition and/or genomic environment through deliberate human manipulation. Preferably the HD-hydrolase-like polypeptide-encoding nucleic acid is from a plant or a bacterium, further preferably from a cyanophyte, more preferably from the family Chroococcaceae, most preferably the nucleic acid is from *Synechocystis* sp.

Performance of the methods of the invention gives plants having enhanced yield-related traits. In particular, performance of the methods of the invention gives plants having increased yield, especially increased seed yield relative to control plants. The terms "yield" and "seed yield" are described in more detail in the "definitions" section herein.

Reference herein to enhanced yield-related traits is taken to mean an increase early vigour and/or in biomass (weight) of one or more parts of a plant, which may include aboveground (harvestable) parts and/or (harvestable) parts below ground. In particular, such harvestable parts are seeds, and performance of the methods of the invention results in plants having increased seed yield relative to the seed yield of control plants.

The present invention provides a method for increasing yield-related traits, especially seed yield in plants, relative to control plants, which method comprises modulating expression in a plant of a nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, as defined herein.

Since the transgenic plants according to the present invention have increased yield and/or increased yield-related traits, it is likely that these plants exhibit an increased growth rate (during at least part of their life cycle), relative to the growth rate of control plants at a corresponding stage in their life cycle.

According to a preferred feature of the present invention, performance of the methods of the invention gives plants having an increased growth rate relative to control plants. Therefore, according to the present invention, there is provided a method for increasing the growth rate of plants, which method comprises modulating expression in a plant of a nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, as defined herein.

Performance of the methods of the invention gives plants grown under non-stress conditions or under mild drought conditions increased yield relative to control plants grown under comparable conditions. Therefore, according to the present invention, there is provided a method for increasing yield in plants grown under non-stress conditions or under mild drought conditions, which method comprises modulating expression in a plant of a nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide.

Performance of the methods of the invention gives plants grown under conditions of nutrient deficiency, particularly under conditions of nitrogen deficiency, increased yield relative to control plants grown under comparable conditions. Therefore, according to the present invention, there is provided a method for increasing yield in plants grown under conditions of nutrient deficiency, which method comprises modulating expression in a plant of a nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide.

Performance of the methods of the invention gives plants grown under conditions of salt stress, increased yield relative to control plants grown under comparable conditions. Therefore, according to the present invention, there is provided a method for increasing

yield in plants grown under conditions of salt stress, which method comprises modulating expression in a plant of a nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide.

The invention also provides genetic constructs and vectors to facilitate introduction and/or expression in plants of nucleic acids encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide. The gene constructs may be inserted into vectors, which may be commercially available, suitable for transforming into plants and suitable for expression of the gene of interest in the transformed cells. The invention also provides use of a gene construct as defined herein in the methods of the invention.

More specifically, the present invention provides a construct comprising:

- (a) a nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, as defined above;
- (b) one or more control sequences capable of driving expression of the nucleic acid sequence of (a); and optionally
- (c) a transcription termination sequence.

Preferably, the nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, is as defined above. The term “control sequence” and “termination sequence” are as defined herein.

Plants are transformed with a vector comprising any of the nucleic acids described above. The skilled artisan is well aware of the genetic elements that must be present on the vector in order to successfully transform, select and propagate host cells containing the sequence of interest. The sequence of interest is operably linked to one or more control sequences (at least to a promoter).

Concerning SGT1 polypeptides, advantageously, any type of promoter, whether natural or synthetic, may be used to drive expression of the nucleic acid sequence, but preferably the promoter is of plant origin. A root-specific promoter is particularly useful in the methods. The root-specific promoter is preferably an RCc3 promoter (Plant Mol Biol. 1995 Jan;27(2):237-48), more preferably the RCc3 promoter is from rice, further preferably the RCc3 promoter is represented by a nucleic acid sequence substantially similar to SEQ ID NO: 50, most preferably the promoter is as represented by SEQ ID NO: 50. Examples of other root-specific promoters which may also be used to perform the methods of the invention are shown in Table 2b in the “Definitions” section above.

Concerning CLC-pKG polypeptides, advantageously, any type of promoter, whether natural or synthetic, may be used to drive expression of the nucleic acid sequence, but preferably the promoter is of plant origin. A root promoter is particularly useful in the methods. Preferably the root promoter is ubiquitous within the root. Also useful in the methods of the

invention is constitutive promoter. Preferably the constitutive promoter is a ubiquitous constitutive promoter of medium strength. See the "Definitions" section herein for definitions of the various promoter types.

Concerning HD-hydrolase-like polypeptides, advantageously, any type of promoter, whether natural or synthetic, may be used to drive expression of the nucleic acid sequence, but preferably the promoter is of plant origin. A constitutive promoter is particularly useful in the methods. Preferably the constitutive promoter is a ubiquitous constitutive promoter of medium strength. See the "Definitions" section herein for definitions of the various promoter types.

Concerning SGT1 polypeptides, it should be clear that the applicability of the present invention is not restricted to the SGT1 polypeptide-encoding nucleic acid represented by SEQ ID NO: 1, nor is the applicability of the invention restricted to expression of a SGT1 polypeptide-encoding nucleic acid when driven by a root-specific promoter.

Optionally, one or more terminator sequences may be used in the construct introduced into a plant. Preferably, the construct comprises an expression cassette comprising an RCc3 promoter, substantially similar to SEQ ID NO: 50, and the nucleic acid encoding an SGT1 polypeptide.

According to a preferred feature of the invention, the modulated expression is increased expression. Methods for increasing expression of nucleic acids or genes, or gene products, are well documented in the art and examples are provided in the definitions section.

Concerning CLC-pKG polypeptides, it should be clear that the applicability of the present invention is not restricted to the CLC-pKG polypeptide-encoding nucleic acid represented by SEQ ID NO: 53, nor is the applicability of the invention restricted to expression of a CLC-pKG polypeptide-encoding nucleic acid when driven by root-specific a promoter, or when driven by a constitutive promoter.

The constitutive promoter is preferably a medium strength promoter, more preferably selected from a plant derived promoter, such as a GOS2 promoter, more preferably is the promoter GOS2 promoter from rice.

According to another preferred feature of the invention, the nucleic acid encoding a CLC-pKG polypeptide is operably linked to a root-specific promoter. The root-specific promoter is preferably an RCc3 promoter (Plant Mol Biol. 1995 Jan;27(2):237-48), more preferably the RCc3 promoter is from rice, further preferably the RCc3 promoter is represented by a nucleic acid sequence substantially similar to SEQ ID NO: 115, most preferably the promoter is as represented by SEQ ID NO: 115. Examples of other root-specific promoters

which may also be used to perform the methods of the invention are shown in Table 2b in the "Definitions" section above.

According to a preferred feature of the invention, the modulated expression is increased expression. Methods for increasing expression of nucleic acids or genes, or gene products, are well documented in the art and examples are provided in the definitions section.

Concerning HD-hydrolase-like polypeptides, it should be clear that the applicability of the present invention is not restricted to the HD-hydrolase-like polypeptide-encoding nucleic acid represented by SEQ ID NO: 118, nor is the applicability of the invention restricted to expression of a HD-hydrolase-like polypeptide-encoding nucleic acid when driven by a constitutive promoter.

The constitutive promoter is preferably a medium strength promoter of plant origin, such as a GOS2 promoter, more preferably the promoter is the GOS2 promoter from rice. Further preferably the constitutive promoter is represented by a nucleic acid sequence substantially similar to SEQ ID NO: 164, most preferably the constitutive promoter is as represented by SEQ ID NO: 164. See the "Definitions" section herein for further examples of constitutive promoters.

Optionally, one or more terminator sequences may be used in the construct introduced into a plant. Preferably, the construct comprises an expression cassette comprising a GOS2 promoter, substantially similar to SEQ ID NO: 164, and the nucleic acid encoding the HD-hydrolase-like polypeptide.

According to a preferred feature of the invention, the modulated expression is increased expression. Methods for increasing expression of nucleic acids or genes, or gene products, are well documented in the art and examples are provided in the definitions section.

As mentioned above, a preferred method for modulating expression of a nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, is by introducing and expressing in a plant a nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide respectively; however the effects of performing the method, i.e. enhancing yield-related traits may also be achieved using other well known techniques, including but not limited to T-DNA activation tagging, TILLING, homologous recombination. A description of these techniques is provided in the definitions section.

The invention also provides a method for the production of transgenic plants having enhanced yield-related traits relative to control plants, comprising introduction and expression in a plant of any nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, as defined hereinabove.

More specifically, the present invention provides a method for the production of transgenic plants having enhanced yield-related traits, particularly increased (seed) yield, which method comprises:

- (i) introducing and expressing in a plant or plant cell nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide; and
- (ii) cultivating the plant cell under conditions promoting plant growth and development.

The nucleic acid of (i) may be any of the nucleic acids capable of encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, as defined herein.

The nucleic acid may be introduced directly into a plant cell or into the plant itself (including introduction into a tissue, organ or any other part of a plant). According to a preferred feature of the present invention, the nucleic acid is preferably introduced into a plant by transformation. The term "transformation" is described in more detail in the "definitions" section herein.

The present invention clearly extends to any plant cell or plant produced by any of the methods described herein, and to all plant parts and propagules thereof. The present invention encompasses plants or parts thereof (including seeds) obtainable by the methods according to the present invention. The plants or parts thereof comprise a nucleic acid transgene encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, as defined above. The present invention extends further to encompass the progeny of a primary transformed or transfected cell, tissue, organ or whole plant that has been produced by any of the aforementioned methods, the only requirement being that progeny exhibit the same genotypic and/or phenotypic characteristic(s) as those produced by the parent in the methods according to the invention.

The invention also includes host cells containing an isolated nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, as defined hereinabove. Preferred host cells according to the invention are plant cells. Host plants for the nucleic acids or the vector used in the method according to the invention, the expression cassette or construct or vector are, in principle, advantageously all plants, which are capable of synthesizing the polypeptides used in the inventive method.

The methods of the invention are advantageously applicable to any plant. Plants that are particularly useful in the methods of the invention include all plants which belong to the superfamily Viridiplantae, in particular monocotyledonous and dicotyledonous plants including fodder or forage legumes, ornamental plants, food crops, trees or shrubs. According to a preferred embodiment of the present invention, the plant is a crop plant.

Examples of crop plants include soybean, sunflower, canola, alfalfa, rapeseed, linseed, cotton, tomato, potato and tobacco. Further preferably, the plant is a monocotyledonous plant. Examples of monocotyledonous plants include sugarcane. More preferably the plant is a cereal. Examples of cereals include rice, maize, wheat, barley, millet, rye, triticale, sorghum, emmer, spelt, secale, einkorn, teff, milo and oats.

The invention also extends to harvestable parts of a plant such as, but not limited to seeds, leaves, fruits, flowers, stems, roots, rhizomes, tubers and bulbs, which harvestable parts comprise a recombinant nucleic acid encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide. The invention furthermore relates to products derived, preferably directly derived, from a harvestable part of such a plant, such as dry pellets or powders, oil, fat and fatty acids, starch or proteins.

The present invention also encompasses use of nucleic acids encoding SGT1 polypeptides, or CLC-pKG polypeptides, or HD-hydrolase-like polypeptides, as described herein and use of these SGT1 polypeptides, or CLC-pKG polypeptides, or HD-hydrolase-like polypeptides, in enhancing any of the aforementioned yield-related traits in plants. For example, nucleic acids encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, described herein, or the SGT1 polypeptides, or CLC-pKG polypeptides, or HD-hydrolase-like polypeptides, themselves, may find use in breeding programmes in which a DNA marker is identified which may be genetically linked to gene encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide,. The nucleic acids/genes, or the SGT1 polypeptides, or CLC-pKG polypeptides, or HD-hydrolase-like polypeptides, may be used to define a molecular marker. This DNA or protein marker may then be used in breeding programmes to select plants having enhanced yield-related traits as defined hereinabove in the methods of the invention. Furthermore, allelic variants of nucleic acid/gene encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, may find use in marker-assisted breeding programmes. Nucleic acids encoding an SGT1 polypeptide, or a CLC-pKG polypeptide, or a HD-hydrolase-like polypeptide, may also be used as probes for genetically and physically mapping the genes that they are a part of, and as markers for traits linked to those genes. Such information may be useful in plant breeding in order to develop lines with desired phenotypes.

Items

1. SGT1 polypeptides

In one aspect, the invention is characterised by any of the following items

1. A method for enhancing yield-related traits in plants relative to control plants, comprising modulating expression in a plant of a nucleic acid encoding an SGT1 polypeptide, wherein said SGT1 polypeptide comprises the following in any order:
 - (i) at least one tetratricopeptide (TPR) repeat;
 - (ii) at least one CS domain; and

- (iii) an SGS domain.
2. Method according to item 1, wherein said modulated expression is effected by introducing and expressing in a plant a nucleic acid encoding an SGT1 polypeptide.
 3. Method according to item 1 or 2, wherein said nucleic acid encoding an SGT1 polypeptide encodes any one of the proteins listed in Table A1 or is a portion of such a nucleic acid, or a nucleic acid capable of hybridising with such a nucleic acid.
 4. Method according to any one of items 1 to 3, wherein said nucleic acid sequence encodes an orthologue or paralogue of any of the proteins given in Table A1.
 5. Method according to any preceding item, wherein said enhanced yield-related traits comprise increased seed yield relative to control plants.
 6. Method according to any one of items 1 to 5, wherein said enhanced yield-related traits are obtained under non-stress conditions.
 7. Method according to any one of items 2 to 6, wherein said nucleic acid is operably linked to a root-specific promoter, preferably to an RCc3 promoter, most preferably to an RCc3 promoter from rice.
 8. Method according to any one of items 1 to 7, wherein said nucleic acid encoding a SGT1 polypeptide is of plant origin, preferably from *Capsicum annuum* a dicotyledonous plant, further preferably from the family Solanaceae, more preferably from the genus *Capsicum*, most preferably from *Capsicum annuum*.
 9. Plant or part thereof, including seeds, obtainable by a method according to any one of items 1 to 8, wherein said plant or part thereof comprises a recombinant nucleic acid encoding an SGT1 polypeptide.
 10. Construct comprising:
 - (i) nucleic acid encoding an SGT1 polypeptide as defined in item 1;
 - (ii) one or more control sequences capable of driving expression of the nucleic acid sequence of (a); and optionally
 - (iii) a transcription termination sequence.
 11. Construct according to item 10, wherein one of said control sequences is a root-specific promoter, preferably an RCc3 promoter, most preferably an RCc3 promoter from rice.

12. Use of a construct according to item 10 or 11 in a method for making plants having increased yield, particularly increased seed yield relative to control plants.
13. Plant, plant part or plant cell transformed with a construct according to item 10 or 11.
14. Method for the production of a transgenic plant having increased yield, particularly increased seed yield relative to control plants, comprising:
 - (i) introducing and expressing in a plant a nucleic acid encoding an SGT1 polypeptide as defined in item 1; and
 - (ii) cultivating the plant cell under conditions promoting plant growth and development.
15. Transgenic plant having increased yield, particularly increased seed yield, relative to control plants, resulting from modulated expression of a nucleic acid encoding an SGT1 polypeptide as defined in item 1, or a transgenic plant cell derived from said transgenic plant.
16. Transgenic plant according to item 9, 13 or 15, or a transgenic plant cell derived thereof, wherein said plant is a crop plant or a monocot or a cereal, such as rice, maize, wheat, barley, millet, rye, triticale, sorghum emmer, spelt, secale, einkorn, teff, milo and oats.
17. Harvestable parts of a plant according to item 16, wherein said harvestable parts are preferably shoot biomass and/or seeds.
18. Products derived from a plant according to item 16 and/or from harvestable parts of a plant according to item 17.
19. Use of a nucleic acid encoding an SGT1 polypeptide in increasing yield, particularly in increasing seed yield relative to control plants.
20. Use of a nucleic acid encoding an SGT1 polypeptide as a molecular marker for plants having increased yield, particularly increased seed yield.

2. CLC-pKG polypeptides

In another aspect, the invention is characterised by any of the following items

1. A method for enhancing yield-related traits in plants relative to control plants, comprising modulating expression in a plant of a nucleic acid encoding a CLC-pKG polypeptide, wherein said CLC-pKG polypeptide comprises a Voltage_CLC domain

(Pfam entry PF00654) and a CBS domain (Pfam entry PF00571) and optionally a USP domain (PF00582).

2. Method according to item 1, wherein said CLC-pKG polypeptide comprises a domain having in increasing order of preference at least 25%, 26%, 27%, 28%, 29%, 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100 % sequence identity to:
 - (i) a CLC in SEQ ID NO: 54 as represented by the sequence located between amino acids 77 and 422 of SEQ ID NO: 54;
 - (ii) a CBS domain in SEQ ID NO: 54 as represented by the sequence located between amino acids 455 to 510 or 516 to 571 SEQ ID NO: 54; and optionally
 - (iii) a USP domain in SEQ ID NO: 54 as represented by the sequence located between amino acids 597-731 of SEQ ID NO: 54.
3. Method according to item 1 or 2, wherein said modulated expression is effected by introducing and expressing in a plant a genetic construct comprising a nucleic acid encoding a CLC-pKG polypeptide.
4. Method according to any one of items 1 to 3, wherein said nucleic acid encoding a CLC-pKG polypeptide encodes any one of the proteins listed in Table A2 or is a portion of such a nucleic acid, or a nucleic acid capable of hybridising with such a nucleic acid.
5. Method according to any one of items 1 to 4, wherein said nucleic acid encodes an orthologue or paralogue of any of the proteins given in Table A2.
6. Method according to any preceding item, wherein said enhanced yield-related traits comprise increased yield, preferably increased biomass and/or increased seed yield relative to control plants.
7. Method according to any one of items 1 to 6, wherein said enhanced yield-related traits are obtained under non-stress conditions or under conditions of drought stress, or salt stress.
8. Method according to any one of items 3 to 7, wherein said nucleic acid is operably linked to a constitutive promoter, preferably to a RCc3 promoter, most preferably to a RCc3 promoter from rice.

9. Method according to any one of items 1 to 8, wherein said nucleic acid encoding a CLC-pKG polypeptide is of cyanobacterial origin, further preferably from *Synechosystis* species, more preferably from *Synechocystis* sp. PCC 6803.
10. Plant or part thereof, including seeds, obtainable by a method according to any one of items 1 to 9, wherein said plant or part thereof comprises a recombinant nucleic acid encoding a CLC-pKG polypeptide.
11. Construct comprising:
 - (i) nucleic acid encoding a CLC-pKG polypeptide as defined in items 1 or 2;
 - (ii) one or more control sequences capable of driving expression of the nucleic acid sequence of (a); and optionally
 - (iii) a transcription termination sequence.
12. Construct according to item 11, wherein one of said control sequences is a constitutive promoter, preferably a RCc3 promoter, most preferably a RCc3 promoter from rice.
13. Use of a construct according to item 11 or 12 in a method for making plants having increased yield, particularly increased biomass and/or increased seed yield relative to control plants.
14. Plant, plant part or plant cell transformed with a construct according to item 11 or 12.
15. Method for the production of a transgenic plant having increased yield, particularly increased biomass and/or increased seed yield relative to control plants, comprising:
 - (i) introducing and expressing in a plant a nucleic acid encoding a CLC-pKG polypeptide as defined in item 1 or 2; and
 - (ii) cultivating the plant cell under conditions promoting plant growth and development.
16. Transgenic plant having increased yield, particularly increased biomass and/or increased seed yield, relative to control plants, resulting from modulated expression of a recombinant nucleic acid encoding a CLC-pKG polypeptide as defined in item 1 or 2, or a transgenic plant cell derived from said transgenic plant.
17. Transgenic plant according to item 10, 14 or 16, or a transgenic plant cell derived thereof, wherein said plant is a crop plant or a monocot or a cereal, such as rice, maize, wheat, barley, millet, rye, triticale, sorghum emmer, spelt, secale, einkorn, teff, milo and oats.
18. Harvestable parts of a plant according to item 17, wherein said harvestable parts are preferably shoot biomass and/or seeds.

19. Products derived from a plant according to item 17 and/or from harvestable parts of a plant according to item 18.
20. Use of a genetic construct comprising a nucleic acid encoding a CLC-pKG polypeptide in increasing yield, particularly in increasing seed yield and/or shoot biomass in plants, relative to control plants.

3. HD-hydrolase-like polypeptides

In yet another aspect, the invention is characterised by any of the following items

1. A method for enhancing yield-related traits in plants relative to control plants, comprising modulating expression in a plant of a nucleic acid encoding a HD-hydrolase-like polypeptide, wherein said HD-hydrolase-like polypeptide comprises a CRISPR-associated protein Crm2 domain.
2. Method according to item 1, wherein said HD-hydrolase-like polypeptide comprises one or more of the motifs 11 to 26.
3. Method according to item 1 or 2, wherein said modulated expression is effected by introducing and expressing in a plant a nucleic acid encoding a HD-hydrolase-like polypeptide.
4. Method according to any one of items 1 to 3, wherein said nucleic acid encoding a HD-hydrolase-like polypeptide encodes any one of the proteins listed in Table A3 or is a portion of such a nucleic acid, or a nucleic acid capable of hybridising with such a nucleic acid.
5. Method according to any one of items 1 to 4, wherein said nucleic acid sequence encodes an orthologue or paralogue of any of the proteins given in Table A3.
6. Method according to any preceding item, wherein said enhanced yield-related traits comprise increased yield, preferably increased biomass and/or increased seed yield relative to control plants.
7. Method according to any one of items 1 to 6, wherein said enhanced yield-related traits are obtained under non-stress conditions.
8. Method according to any one of items 3 to 7, wherein said nucleic acid is operably linked to a constitutive promoter, preferably to a GOS2 promoter, most preferably to a GOS2 promoter from rice.

9. Method according to any one of items 1 to 8, wherein said nucleic acid encoding a HD-hydrolase-like polypeptide is of plant origin, preferably from a cyanophyte, further preferably from the family Chroococcaceae, more preferably from the genus *Synechocystis*.
10. Plant or part thereof, including seeds, obtainable by a method according to any one of items 1 to 9, wherein said plant or part thereof comprises a recombinant nucleic acid encoding a HD-hydrolase-like polypeptide.
11. Construct comprising:
 - (i) nucleic acid encoding a HD-hydrolase-like polypeptide as defined in items 1 or 2;
 - (ii) one or more control sequences capable of driving expression of the nucleic acid sequence of (a); and optionally
 - (iii) a transcription termination sequence.
12. Construct according to item 11, wherein one of said control sequences is a constitutive promoter, preferably a GOS2 promoter, most preferably a GOS2 promoter from rice.
13. Use of a construct according to item 11 or 12 in a method for making plants having increased yield, particularly increased biomass and/or increased seed yield relative to control plants.
14. Plant, plant part or plant cell transformed with a construct according to item 11 or 12.
15. Method for the production of a transgenic plant having increased yield, particularly increased biomass and/or increased seed yield relative to control plants, comprising:
 - (i) introducing and expressing in a plant a nucleic acid encoding a HD-hydrolase-like polypeptide as defined in item 1 or 2; and
 - (ii) cultivating the plant cell under conditions promoting plant growth and development.
16. Transgenic plant having increased yield, particularly increased biomass and/or increased seed yield, relative to control plants, resulting from modulated expression of a nucleic acid encoding a HD-hydrolase-like polypeptide as defined in item 1 or 2, or a transgenic plant cell derived from said transgenic plant.
17. Transgenic plant according to item 10, 14 or 16, or a transgenic plant cell derived thereof, wherein said plant is a crop plant or a monocot or a cereal, such as rice, maize, wheat, barley, millet, rye, triticale, sorghum emmer, spelt, secale, einkorn, teff, milo and oats.

18. Harvestable parts of a plant according to item 17, wherein said harvestable parts are preferably shoot biomass and/or seeds.
19. Products derived from a plant according to item 17 and/or from harvestable parts of a plant according to item 18.
20. Use of a nucleic acid encoding a HD-hydrolase-like polypeptide in increasing yield, particularly in increasing seed yield and/or shoot biomass in plants, relative to control plants.

Description of figures

The present invention will now be described with reference to the following figures in which: Figure 1 represents a multiple alignment of various SGT1 polypeptides. Alignment of polypeptide sequences was performed using the ClustalW 2.0 algorithm of progressive alignment (Thompson et al. (1997) *Nucleic Acids Res* 25:4876-4882; Chenna et al. (2003). *Nucleic Acids Res* 31:3497-3500) with standard setting (slow alignment, similarity matrix: Gonnet, gap opening penalty 10, gap extension penalty: 0.2). Minor manual editing was done to further optimise the alignment. The SGT1 polypeptides are aligned in Figure 2.

Figure 2 shows a further alignment highlighting the conserved domains common to SGT1 polypeptides.

Figure 3 shows a phylogenetic tree of SGT1 polypeptides constructed using a neighbour-joining clustering algorithm as provided in the AlignX programme from the Vector NTI (Invitrogen).

Figure 4 represents the binary vector used for increased expression in *Oryza sativa* of an SGT1-encoding nucleic acid under the control of a rice RCc3 promoter (pRCc3).

Figure 5 represents a multiple alignment of various CLC-pKG polypeptides. The asterisks indicate identical amino acids among the various protein sequences, colons represent highly conserved amino acid substitutions, and the dots represent less conserved amino acid substitution; on other positions there is no sequence conservation. These alignments can be used for defining further motifs, when using conserved amino acids.

Figure 6 shows phylogenetic tree of CLC-pKG polypeptides.

Figure 7 represents the binary vector used for increased expression in *Oryza sativa* of a CLC-PKG-encoding nucleic acid under the control of a rice RCc3 promoter (pRCc3).

Figure 8 represents the domain structure of SEQ ID NO: 119 with conserved motifs 11 to 26, underlined and numbered. The CRISPR-associated protein Crm2 (IPR013407) domain is shown in bold.

Figure 9 represents a multiple alignment of various HD-hydrolase-like polypeptides. The asterisks indicate identical amino acids among the various protein sequences, colons represent highly conserved amino acid substitutions, and the dots represent less conserved amino acid substitution; on other positions there is no sequence conservation. These alignments can be used for defining further motifs, when using conserved amino acids.

Figure 10 shows phylogenetic tree of HD-hydrolase-like polypeptides. The identifiers are the accession numbers as provided for the proteins in the sequence listing.

Figure 11 represents the binary vector used for increased expression in *Oryza sativa* of a HD-hydrolase-like-encoding nucleic acid under the control of a rice GOS2 promoter (pGOS2).

Examples

The present invention will now be described with reference to the following examples, which are by way of illustration alone. The following examples are not intended to completely define or otherwise limit the scope of the invention.

DNA manipulation: unless otherwise stated, recombinant DNA techniques are performed according to standard protocols described in (Sambrook (2001) *Molecular Cloning: a laboratory manual*, 3rd Edition Cold Spring Harbor Laboratory Press, CSH, New York) or in Volumes 1 and 2 of Ausubel et al. (1994), *Current Protocols in Molecular Biology*, Current Protocols. Standard materials and methods for plant molecular work are described in *Plant Molecular Biology Labfax* (1993) by R.D.D. Croy, published by BIOS Scientific Publications Ltd (UK) and Blackwell Scientific Publications (UK).

Example 1: Identification of sequences related to the nucleic acid sequence used in the methods of the invention

1. SGT1 polypeptides

Sequences (full length cDNA, ESTs or genomic) related to SEQ ID NO: 1 and SEQ ID NO: 2 were identified amongst those maintained in the Entrez Nucleotides database at the National Center for Biotechnology Information (NCBI) using database sequence search tools, such as the Basic Local Alignment Tool (BLAST) (Altschul et al. (1990) *J. Mol. Biol.* 215:403-410; and Altschul et al. (1997) *Nucleic Acids Res.* 25:3389-3402). The program is used to find regions of local similarity between sequences by comparing nucleic acid or polypeptide sequences to sequence databases and by calculating the statistical significance of matches. For example, the polypeptide encoded by the nucleic acid of SEQ ID NO: 1 was used for the TBLASTN algorithm, with default settings and the filter to ignore low complexity sequences set off. The output of the analysis was viewed by pairwise comparison, and ranked according to the probability score (E-value), where the score reflect the probability that a particular alignment occurs by chance (the lower the E-value, the more significant the hit). In addition to E-values, comparisons were also scored by percentage identity. Percentage identity refers to the number of identical nucleotides (or amino acids) between the two compared nucleic acid (or polypeptide) sequences over a particular length. In some instances, the default parameters may be adjusted to modify the stringency of the search. For example the E-value may be increased to show less stringent matches. This way, short nearly exact matches may be identified.

Table A1 provides a list of nucleic acid sequences related to SEQ ID NO: 1 and SEQ ID NO: 2.

Table A1: Examples of SGT1 nucleic acids and polypeptides:

Name	Nucleic acid SEQ ID NO:	Polypeptide SEQ ID NO:
SGT1 from Pepper	1	2
A.thaliana_AT4G23570.1#1	3	4
A.thaliana_AT4G23570.2#1	5	6
A.thaliana_AT4G11260.1#1	7	8
A.thaliana_AT4G23570.3#1	9	10
B.napus_BN06MC26625_51284379@26524#1	11	12
G.max_GM06MC38560_sab91h08@37598#1	13	14
H.vulgare_c62589592hv270303@6491#1	15	16
M.truncatula_CT025844_7.4#1	17	18
O.sativa_LOC_Os01g43540.1#1	19	20
P.trichocarpa_scaff_IV.839#1	21	22
P.trichocarpa_scaff_44.273#1	23	24
S.lycopersicum_TC192025#1	25	26
S.lycopersicum_NP9243669#1	27	28
S.lycopersicum_AW398985#1	29	30
T.aestivum_TC280790#1	31	32
Z.mays_ZM07MC27211_BFb0187N13@27130#1	33	34
Z.mays_ZM07MC23831_BFb0038K08@23765#1	35	36

Sequences have been tentatively assembled and publicly disclosed by research institutions, such as The Institute for Genomic Research (TIGR; beginning with TA). The Eukaryotic Gene Orthologs (EGO) database may be used to identify such related sequences, either by keyword search or by using the BLAST algorithm with the nucleic acid sequence or polypeptide sequence of interest. Special nucleic acid sequence databases have been created for particular organisms, such as by the Joint Genome Institute.

2. CLC-pKG polypeptides

Sequences (full length cDNA, ESTs or genomic) related to SEQ ID NO: 53 and SEQ ID NO: 54 were identified amongst those maintained in the Entrez Nucleotides database at the National Center for Biotechnology Information (NCBI) using database sequence search tools, such as the Basic Local Alignment Tool (BLAST) (Altschul et al. (1990) J. Mol. Biol. 215:403-410; and Altschul et al. (1997) Nucleic Acids Res. 25:3389-3402). The program is used to find regions of local similarity between sequences by comparing nucleic acid or polypeptide sequences to sequence databases and by calculating the statistical significance of matches. For example, the polypeptide encoded by the nucleic acid of SEQ ID NO: 53 was used for the TBLASTN algorithm, with default settings and the filter to ignore

low complexity sequences set off. The output of the analysis was viewed by pairwise comparison, and ranked according to the probability score (E-value), where the score reflect the probability that a particular alignment occurs by chance (the lower the E-value, the more significant the hit). In addition to E-values, comparisons were also scored by percentage identity. Percentage identity refers to the number of identical nucleotides (or amino acids) between the two compared nucleic acid (or polypeptide) sequences over a particular length. In some instances, the default parameters may be adjusted to modify the stringency of the search. For example the E-value may be increased to show less stringent matches. This way, short nearly exact matches may be identified.

Table A2 provides a list of nucleic acid sequences and polypeptides related to SEQ ID NO: 53 and SEQ ID NO: 54.

Table A2: Examples of CLC-pKG polypeptides:

Organism	Nucleic acid name	Nucleic acid SEQ ID NO
Synechocystis sp. PCC 6803]	gi 16329170:2216476-2219175	53
Cyanothece sp. PCC 7424]	gi 218437013:4004295-4006922	55
Cyanothece sp. PCC 7822]	gi 196258358:38011-40635	57
Cyanothece sp. CCY0110]	gi 126659182:36041-38674	59
Cyanothece sp. ATCC 51142]	gi 172034917:71619-74267	61
Cyanothece sp. PCC 8801]	gi 218244892:3580780-3583419	63
Crocospaera watsonii WH 8501]	gi 67923302:15782-18430	65
Microcystis aeruginosa PCC 7806]	gi 159026870:19707-22334	67
Microcystis aeruginosa NIES-843]	gi 166362741:3545508-3548135	69
Synechococcus sp. PCC 7002]	gi 170076636:589624-592260	71
Nodularia spumigena CCY9414]	gi 119510275:28473-31136	73
Microcoleus chthonoplastes PCC 7420]	gi 254413420:221158-223773	75
Nostoc punctiforme PCC 73102]	gi 186680550:3804594-3807185	77
Nostoc punctiforme PCC 73102]	gi 186680550:2232857-2235448	79
Anabaena variabilis ATCC 29413]	gi 75906225:2674416-2677004	81
Nostoc sp. PCC 7120]	gi 17227497:5827615-5830230	83
'Nostoc azollae' 0708]	gi 225522173:3082-5673	85
Nostoc punctiforme PCC 73102]	gi 186680550:2232857-2235448	87
Anabaena variabilis ATCC 29413]	gi 75906225:2674416-2677004	89
Nostoc sp. PCC 7120]	gi 17227497:5827615-5830230	91
'Nostoc azollae' 0708]	gi 225522173:3082-5673	93

Nostoc punctiforme PCC 73102]	gi 186680550:1989068-1991719	95
'Nostoc azollae' 0708]	gi 225516157:49369-51948	97
Lyngbya sp. PCC 8106]	gi 119486739:136813-139515	99
Cyanothece sp. PCC 7425]	gi 220905643:4270175-4272766	101
Gloeobacter violaceus PCC 7421]	gi 37519569:4380220-4382796	103
Arthrospira maxima CS-328]	gi 209525652:71572-74244	105
Acaryochloris marina MBIC11017]	gi 158333233:913406-916021	107
Cyanothece sp. PCC 7425]	gi 220905643:2873784-2876477	109
Synechococcus elongatus PCC 7942]	gi 81298811:1653255-1655834	111

Organism	Polypeptide name	Polypeptide SEQ ID NO
Synechocystis sp. PCC 6803]	gi 16331172 ref NP_441900	54
Cyanothece sp. PCC 7424]	gi 218440518 ref YP_002378847	56
Cyanothece sp. PCC 7822]	gi 196258397 ref ZP_03156931	58
Cyanothece sp. CCY0110]	gi 126659209 ref ZP_01730347	60
Cyanothece sp. ATCC 51142]	gi 172034995 ref YP_001801496	62
Cyanothece sp. PCC 8801]	gi 218248189 ref YP_002373560	64
Crocospaera watsonii WH 8501]	gi 67923321 ref ZP_00516804	66
Microcystis aeruginosa PCC 7806]	gi 159026894 emb CAO89145	68
Microcystis aeruginosa NIES-843]	gi 166366612 ref YP_001658885	70
Synechococcus sp. PCC 7002]	gi 170077188 ref YP_001733826	72
Nodularia spumigena CCY9414]	gi 119510301 ref ZP_01629437	74
Microcoleus chthonoplastes PCC 7420]	gi 254413597 ref ZP_05027367	76
Nostoc punctiforme PCC 73102]	gi 186683293 ref YP_001866489	78
Nostoc punctiforme PCC 73102]	gi 186682226 ref YP_001865422	80
Anabaena variabilis ATCC 29413]	gi 75908381 ref YP_322677	82
Nostoc sp. PCC 7120]	gi 17232383 ref NP_488931	84
'Nostoc azollae' 0708]	gi 225522175 ref ZP_03768989	86
Nostoc punctiforme PCC 73102]	gi 186682226 ref YP_001865422	88
Anabaena variabilis ATCC 29413]	gi 75908381 ref YP_322677	90
Nostoc sp. PCC 7120]	gi 17232383 ref NP_488931	92
'Nostoc azollae' 0708]	gi 225522175 ref ZP_03768989	94
Nostoc punctiforme PCC 73102]	gi 186682049 ref YP_001865245	96

'Nostoc azollae' 0708]	gi 225516217 ref ZP_03763190	98
Lyngbya sp. PCC 8106]	gi 119486852 ref ZP_01620827	100
Cyanothece sp. PCC 7425]	gi 220909579 ref YP_002484890	102
Gloeobacter violaceus PCC 7421]	gi 37523751 ref NP_927128	104
Arthrospira maxima CS-328]	gi 209525724 ref ZP_03274261	106
Acaryochloris marina MBIC11017]	gi 158334130 ref YP_001515302	108
Cyanothece sp. PCC 7425]	gi 220908306 ref YP_002483617	110
Synechococcus elongatus PCC 7942]	gi 81300397 ref YP_400605	112

Sequences have been tentatively assembled and publicly disclosed by research institutions, such as The Institute for Genomic Research (TIGR; beginning with TA). The Eukaryotic Gene Orthologs (EGO) database may be used to identify such related sequences, either by keyword search or by using the BLAST algorithm with the nucleic acid sequence or polypeptide sequence of interest. Special nucleic acid sequence databases have been created for particular organisms, such as by the Joint Genome Institute. Furthermore, access to proprietary databases, has allowed the identification of novel nucleic acid and polypeptide sequences.

3. HD-hydrolase-like polypeptides

Sequences (full length cDNA, ESTs or genomic) related to SEQ ID NO: 118 and SEQ ID NO: 119 were identified amongst those maintained in the Entrez Nucleotides database at the National Center for Biotechnology Information (NCBI) using database sequence search tools, such as the Basic Local Alignment Tool (BLAST) (Altschul et al. (1990) J. Mol. Biol. 215:403-410; and Altschul et al. (1997) Nucleic Acids Res. 25:3389-3402). The program is used to find regions of local similarity between sequences by comparing nucleic acid or polypeptide sequences to sequence databases and by calculating the statistical significance of matches. For example, the polypeptide encoded by the nucleic acid of SEQ ID NO: 118 was used for the TBLASTN algorithm, with default settings and the filter to ignore low complexity sequences set off. The output of the analysis was viewed by pairwise comparison, and ranked according to the probability score (E-value), where the score reflect the probability that a particular alignment occurs by chance (the lower the E-value, the more significant the hit). In addition to E-values, comparisons were also scored by percentage identity. Percentage identity refers to the number of identical nucleotides (or amino acids) between the two compared nucleic acid (or polypeptide) sequences over a particular length. In some instances, the default parameters may be adjusted to modify the stringency of the search. For example the E-value may be increased to show less stringent matches. This way, short nearly exact matches may be identified.

Table A3 provides a list of nucleic acid sequences related to SEQ ID NO: 118 and SEQ ID NO: 119.

Table A3: Examples of HD-hydrolase-like nucleic acids and polypeptides:

Name	Source	Nucleic acid SEQ ID NO:	Polypeptide SEQ ID NO:
NP_942378.1	Synechocystis sp.	118	119
YP_001736234.1	Synechococcus sp.	120	121
YP_477086.1	Synechococcus sp.	122	123
ZP_01732553.1	Cyanothece sp.	124	125
YP_001801768.1	Cyanothece sp.	126	127
YP_475929.1	Synechococcus sp.	128	129
ZP_03274910.1	Arthrospira maxima	130	131
YP_001002214.1	Halorhodospira halophila	132	133
ZP_05334193.1	Thermoanaerobacter italicus	134	135
ZP_05378933.1	Thermoanaerobacter mathranii subsp. Mathranii	136	137
NP_624155.1	Thermoanaerobacter tengcongensis	138	139
ZP_05091540.1	Carboxydibrachium pacificum	140	141
YP_001616323.1	Sorangium cellulosum	142	143
ZP_05335917.1	Thermoanaerobacterium thermosaccharolyticum	144	145
ZP_01125571.1	Nitrococcus mobilis	146	147

Sequences have been tentatively assembled and publicly disclosed by research institutions, such as The Institute for Genomic Research (TIGR; beginning with TA). The Eukaryotic Gene Orthologs (EGO) database may be used to identify such related sequences, either by keyword search or by using the BLAST algorithm with the nucleic acid sequence or polypeptide sequence of interest. Special nucleic acid sequence databases have been created for particular organisms, such as by the Joint Genome Institute. Furthermore, access to proprietary databases, has allowed the identification of novel nucleic acid and polypeptide sequences.

Example 2: Alignment of sequences related to the polypeptide sequences used in the methods of the invention

1. SGT1 polypeptides

Alignment of polypeptide sequences was performed using the ClustalW 2.0 algorithm of progressive alignment (Thompson et al. (1997) Nucleic Acids Res 25:4876-4882; Chenna et al. (2003). Nucleic Acids Res 31:3497-3500) with standard setting (slow alignment, similarity matrix: Gonnet, gap opening penalty 10, gap extension penalty: 0.2). Minor

manual editing was done to further optimise the alignment. The SGT1 polypeptides are aligned in Figures 1 and 2.

A phylogenetic tree of SGT1 polypeptides (Figure 3) was constructed using a neighbour-joining clustering algorithm as provided in the AlignX programme from the Vector NTI (Invitrogen).

2. CLC-pKG polypeptides

Alignment of polypeptide sequences was performed using the ClustalW 1.8 algorithm of progressive alignment (Thompson et al. (1997) *Nucleic Acids Res* 25:4876-4882; Chenna et al. (2003). *Nucleic Acids Res* 31:3497-3500) with standard setting (slow alignment, similarity matrix: Blosum 62, gap opening penalty 10, gap extension penalty: 0.2). Minor manual editing was done to further optimise the alignment. The CLC-pKG polypeptides are aligned in Figure 5.

A phylogenetic tree of CLC-pKG polypeptides (Figure 6) was taken from Marmagne et al. 2007.

3. HD-hydrolase-like polypeptides

Alignment of polypeptide sequences was performed using the ClustalW 1.81 algorithm of progressive alignment (Thompson et al. (1997) *Nucleic Acids Res* 25:4876-4882; Chenna et al. (2003). *Nucleic Acids Res* 31:3497-3500) with standard setting (slow alignment, similarity matrix: Gonnet, gap opening penalty 10, gap extension penalty: 0.2). Minor manual editing was done to further optimise the alignment.

A phylogenetic tree of HD-hydrolase-like polypeptides (Figure 10) was constructed using a neighbour-joining clustering algorithm as provided in the AlignX program of Vector NTI (Invitrogen).

Example 3: Calculation of global percentage identity between polypeptide sequences useful in performing the methods of the invention

1. SGT1 polypeptides

Global percentages of similarity and identity between full length polypeptide sequences useful in performing the methods of the invention were determined using one of the methods available in the art, the MatGAT (Matrix Global Alignment Tool) software (BMC Bioinformatics. 2003 4:29. MatGAT: an application that generates similarity/identity matrices using protein or DNA sequences. Campanella JJ, Bitincka L, Smalley J; software hosted by Ledion Bitincka). MatGAT software generates similarity/identity matrices for DNA or protein sequences without needing pre-alignment of the data. The program performs a series of pair-wise alignments using the Myers and Miller global alignment algorithm (with a gap opening penalty of 12, and a gap extension penalty of 2), calculates similarity and identity

using for example Blosum 62 (for polypeptides), and then places the results in a distance matrix.

Parameters used in the comparison are: Scoring matrix: Blosum62, First Gap: 12, Extending Gap: 2.

A MATGAT table for local alignment of a specific domain, or data on % identity/similarity between specific domains may also be performed.

2. CLC-pKG polypeptides

Global percentages of similarity and identity between full length polypeptide sequences useful in performing the methods of the invention were determined using one of the methods available in the art, the MatGAT (Matrix Global Alignment Tool) software (BMC Bioinformatics. 2003 4:29. MatGAT: an application that generates similarity/identity matrices using protein or DNA sequences. Campanella JJ, Bitincka L, Smalley J; software hosted by Ledion Bitincka). MatGAT software generates similarity/identity matrices for DNA or protein sequences without needing pre-alignment of the data. The program performs a series of pair-wise alignments using the Myers and Miller global alignment algorithm (with a gap opening penalty of 12, and a gap extension penalty of 2), calculates similarity and identity using for example Blosum 62 (for polypeptides), and then places the results in a distance matrix. Parameters typically used in the comparisons are: Scoring matrix: Blosum62, First Gap: 12, Extending Gap: 2.

3. HD-hydrolase-like polypeptides

Global percentages of similarity and identity between full length polypeptide sequences useful in performing the methods of the invention were determined using one of the methods available in the art, the MatGAT (Matrix Global Alignment Tool) software (BMC Bioinformatics. 2003 4:29. MatGAT: an application that generates similarity/identity matrices using protein or DNA sequences. Campanella JJ, Bitincka L, Smalley J; software hosted by Ledion Bitincka). MatGAT software generates similarity/identity matrices for DNA or protein sequences without needing pre-alignment of the data. The program performs a series of pair-wise alignments using the Myers and Miller global alignment algorithm (with a gap opening penalty of 12, and a gap extension penalty of 2), calculates similarity and identity using for example Blosum 62 (for polypeptides), and then places the results in a distance matrix. Sequence similarity is shown in the bottom half of the dividing line and sequence identity is shown in the top half of the diagonal dividing line.

Parameters used in the comparison were: Scoring matrix: Blosum62, First Gap: 12, Extending Gap: 2.

Results of the software analysis are shown in Table B1 for the global similarity and identity over the full length of the polypeptide sequences. The sequence identity (in %) between

the HD-hydrolase-like polypeptide sequences useful in performing the methods of the invention can be as low as 17 % compared to SEQ ID NO: 118 (NP_942378.1).

Table B1: MatGAT results for global similarity and identity over the full length of the polypeptide sequences.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1. NP_942378.1		68.1	48.3	51.5	47.2	48.3	27.2	19.6	23.6	23.5	21.5	21.5	17.2	20.5	20.4
2. YP_001736234.1	80.1		46.4	50.7	46.7	49.3	27.9	19.7	25.5	24.5	21.0	20.9	19.9	22.6	22.8
3. ZP_01732553.1	61.8	61.0		44.7	45.5	44.0	25.7	15.4	19.4	18.9	21.2	21.3	22.2	19.0	18.9
4. YP_477086.1	66.7	66.2	59.3		78.1	47.5	28.2	21.2	26.0	24.0	21.2	21.2	18.1	21.5	21.0
5. YP_475929.1	62.6	60.9	61.2	83.8		44.2	27.9	20.8	24.3	23.6	20.1	20.0	17.5	20.6	20.3
6. YP_001801768.1	63.5	64.4	58.2	65.1	59.6		26.7	19.5	23.4	22.4	22.9	21.9	18.7	20.9	20.4
7. ZP_03274910.1	44.6	46.3	40.4	47.1	44.8	47.9		18.7	21.8	19.9	19.3	19.4	17.7	20.5	20.1
8. YP_001616323.1	33.7	34.1	28.4	34.3	32.7	35.1	33.9		24.1	23.4	18.3	18.3	15.4	20.1	20.2
9. YP_001002214.1	42.2	41.8	36.0	41.7	39.4	40.3	37.5	34.6		56.1	25.4	25.5	20.2	26.3	26.6
10. ZP_01125571.1	38.8	40.5	35.4	40.8	40.3	39.1	36.2	35.9	66.8		24.9	24.8	19.6	26.7	27.2
11. ZP_05334193.1	39.4	39.8	39.8	41.2	40.4	41.4	35.6	34.3	42.9	45.5		99.4	29.2	26.3	26.7
12. ZP_05378933.1	39.1	39.7	40.4	40.9	40.5	41.8	35.6	34.2	43.0	45.6	99.8		29.3	26.4	26.7
13. ZP_05335917.1	33.8	36.4	38.7	35.5	35.3	35.2	32.5	28.8	35.1	37.6	47.8	47.8		22.3	22.6
14. NP_624155.1	40.2	40.3	37.2	41.1	41.4	39.4	37.8	33.5	43.3	44.6	48.4	48.4	40.6		97.6
15. ZP_05091540.1	40.2	40.8	36.9	40.1	41.1	39.3	37.6	33.5	43.5	44.4	49.1	49.4	40.4	98.6	


Example 4: Identification of domains comprised in polypeptide sequences useful in performing the methods of the invention

1. SGT1 polypeptides

The Integrated Resource of Protein Families, Domains and Sites (InterPro) database is an integrated interface for the commonly used signature databases for text- and sequence-based searches. The InterPro database combines these databases, which use different methodologies and varying degrees of biological information about well-characterized proteins to derive protein signatures. Collaborating databases include SWISS-PROT, PROSITE, TrEMBL, PRINTS, ProDom and Pfam, Smart and TIGRFAMs. Pfam is a large collection of multiple sequence alignments and hidden Markov models covering many common protein domains and families. Pfam is hosted at the Sanger Institute server in the United Kingdom. Interpro is hosted at the European Bioinformatics Institute in the United Kingdom.

The results of the InterPro scan of the polypeptide sequence as represented by SEQ ID NO: 2 are presented in Table C1.

Table C1: InterPro scan results (major accession numbers) of the polypeptide sequence as represented by SEQ ID NO: 2.



sanger

institute

HOME

SEARCH

BROWSE

FTP


HELP

Sequence search results

Show

the detailed description of this results page.

We found 6 Pfam-A matches to your search sequence (3 significant and 3 insignificant) . You did not choose to search for Pfam-B matches on your sequence.



Show

the search options and sequence that you submitted.

Return
to the search form to look for Pfam domains on a new sequence.

Significant Pfam-A Matches

show

or

hide

all alignments.

Pfam-A	Description	Entry type	Sequence		HMM		Bits score	E-value	Alignment mode
			Start	End	From	To			
IPR_1	Tetratricopeptide repeat	Repeat	70	103	1	34	21.0	7.3e-05	fs
#HMM	*->eeayniShaylkigkydeAieayekAeldFnk--*								
#MATCH	++y ++G++kl+-y+ A +te +l P								
#SEQ	SKATLRPLAENLEFYTAFAALITGAPLAGE 103								
CS	CS domain	Domain	172	248	1	33	94.3	4.5e-27	fs
#HMM	*--ayrnyvsihvticplgwygxtkthwVekprilkevshgpgwagyllGeplegldpccfawksicshkvtlthk--*								
#MATCH	++++G ++G++G++G++ +++++V++++ lqvsl +pgy + G + Llg+L p ++ + ++ k+el L N								
#SEQ	VHEFTNPKFETVTLFARLI---FANWYVDFEGHLYVSHLPQ6-ETTFPSP-1SGRTYAKCKEYTNST-KTLIRLAK 248								
SGS	SGS domain	Domain	207	368	1	91	186.1	6.9e-53	fs
#HMM	*--kKkSDkltaveKrikwKkchisodEklcGcAAlmniFKk1YedpddakPAdwTStEEnGTvLSTNKKdVgkthVetkPpccapKew--*								
#MATCH	+GKk+e ++KkEk dEkldc+ARLn++F++-Y+O+D+G++RAdKSE+ERnGTvLSTNKK Vg kKVe++lp+GdS+K+w								
#SEQ	RYVSDKLA-----QVNNKK---DEPLDGDALNKKFPVLYKDAEDTFANWESFVESHGTVLSINWVVGTYVVEGSPYDGHLPFW 368								

2. CLC-pKG polypeptides

The Integrated Resource of Protein Families, Domains and Sites (InterPro) database is an integrated interface for the commonly used signature databases for text- and sequence-based searches. The InterPro database combines these databases, which use different

methodologies and varying degrees of biological information about well-characterized proteins to derive protein signatures. Collaborating databases include SWISS-PROT, PROSITE, TrEMBL, PRINTS, ProDom and Pfam, Smart and TIGRFAMs. Pfam is a large collection of multiple sequence alignments and hidden Markov models covering many common protein domains and families. Pfam is hosted at the Sanger Institute server in the United Kingdom. Interpro is hosted at the European Bioinformatics Institute in the United Kingdom.

The results of the pfam scan of the polypeptide sequence as represented by SEQ ID NO: 54 are presented in Table C2.

Table C2: pfam scan results (major accession numbers) of the polypeptide sequence as represented by SEQ ID NO: 54.

Database	Accession number	Accession name	Amino acid coordinates on SEQ ID NO 54
Pfam	PF00571	Voltage Gate Chloride Channel	77-422
Pfam	PF00571	CBS domain	455-510
Pfam	PF00571	CBS domain	516-571
Pfam	PF00582	Universal Stress Protein Family	597-731

3. HD-hydrolase-like polypeptides

The Integrated Resource of Protein Families, Domains and Sites (InterPro) database is an integrated interface for the commonly used signature databases for text- and sequence-based searches. The InterPro database combines these databases, which use different methodologies and varying degrees of biological information about well-characterized proteins to derive protein signatures. Collaborating databases include SWISS-PROT, PROSITE, TrEMBL, PRINTS, ProDom and Pfam, Smart and TIGRFAMs. Pfam is a large collection of multiple sequence alignments and hidden Markov models covering many common protein domains and families. Pfam is hosted at the Sanger Institute server in the United Kingdom. Interpro is hosted at the European Bioinformatics Institute in the United Kingdom.

The results of the InterPro scan of the polypeptide sequence as represented by SEQ ID NO: 119 are presented in Table C3.

Table C3: InterPro scan results (major accession numbers) of the polypeptide sequence as represented by SEQ ID NO: 119.

Database	Accession number	Accession name	Amino acid coordinates on SEQ ID NO 119
InterPro	IPR013407	CRISPR-associated protein Crm2	
HMMTigr	TIGR02577	cas_TM1794_Crm2: CRISPR-associated pr	[221-842] 8.7e-13

Example 5: Topology prediction of the polypeptide sequences useful in performing the methods of the invention

1. SGT1 polypeptides

TargetP 1.1 predicts the subcellular location of eukaryotic proteins. The location assignment is based on the predicted presence of any of the N-terminal pre-sequences: chloroplast transit peptide (cTP), mitochondrial targeting peptide (mTP) or secretory pathway signal peptide (SP). Scores on which the final prediction is based are not really probabilities, and they do not necessarily add to one. However, the location with the highest score is the most likely according to TargetP, and the relationship between the scores (the reliability class) may be an indication of how certain the prediction is. The reliability class (RC) ranges from 1 to 5, where 1 indicates the strongest prediction. TargetP is maintained at the server of the Technical University of Denmark.

For the sequences predicted to contain an N-terminal pre-sequence a potential cleavage site can also be predicted.

A number of parameters are selected, such as organism group (non-plant or plant), cutoff sets (none, predefined set of cutoffs, or user-specified set of cutoffs), and the calculation of prediction of cleavage sites (yes or no).

Many other algorithms can be used to perform such analyses, including:

- ChloroP 1.1 hosted on the server of the Technical University of Denmark;
- Protein Prowler Subcellular Localisation Predictor version 1.2 hosted on the server of the Institute for Molecular Bioscience, University of Queensland, Brisbane, Australia;
- PENCE Proteome Analyst PA-GOSUB 2.5 hosted on the server of the University of Alberta, Edmonton, Alberta, Canada;
- TMHMM, hosted on the server of the Technical University of Denmark
- PSORT (URL: psort.org)
- PLOC (Park and Kanehisa, Bioinformatics, 19, 1656-1663, 2003).

2. CLC-pKG polypeptides

TargetP 1.1 predicts the subcellular location of eukaryotic proteins. The location assignment is based on the predicted presence of any of the N-terminal pre-sequences: chloroplast transit peptide (cTP), mitochondrial targeting peptide (mTP) or secretory pathway signal peptide (SP). Scores on which the final prediction is based are not really probabilities, and they do not necessarily add to one. However, the location with the highest score is the most likely according to TargetP, and the relationship between the scores (the reliability class) may be an indication of how certain the prediction is. The reliability class (RC) ranges from

1 to 5, where 1 indicates the strongest prediction. TargetP is maintained at the server of the Technical University of Denmark.

For the sequences predicted to contain an N-terminal presequence a potential cleavage site can also be predicted.

In addition or alternatively, many other algorithms can be used to perform such analyses, including:

- ChloroP 1.1 hosted on the server of the Technical University of Denmark;
- Protein Prowler Subcellular Localisation Predictor version 1.2 hosted on the server of the Institute for Molecular Bioscience, University of Queensland, Brisbane, Australia;
- PENCE Proteome Analyst PA-GOSUB 2.5 hosted on the server of the University of Alberta, Edmonton, Alberta, Canada;
- TMHMM, hosted on the server of the Technical University of Denmark
- PSORT (URL: psort.org)
- PLOC (Park and Kanehisa, Bioinformatics, 19, 1656-1663, 2003).

3. HD-hydrolase-like polypeptides

TargetP 1.1 predicts the subcellular location of eukaryotic proteins. The location assignment is based on the predicted presence of any of the N-terminal pre-sequences: chloroplast transit peptide (cTP), mitochondrial targeting peptide (mTP) or secretory pathway signal peptide (SP). Scores on which the final prediction is based are not really probabilities, and they do not necessarily add to one. However, the location with the highest score is the most likely according to TargetP, and the relationship between the scores (the reliability class) may be an indication of how certain the prediction is. The reliability class (RC) ranges from 1 to 5, where 1 indicates the strongest prediction. TargetP is maintained at the server of the Technical University of Denmark.

For the sequences predicted to contain an N-terminal presequence a potential cleavage site can also be predicted.

A number of parameters were selected, such as organism group (non-plant or plant), cutoff sets (none, predefined set of cutoffs, or user-specified set of cutoffs), and the calculation of prediction of cleavage sites (yes or no).

The results of TargetP 1.1 analysis of the polypeptide sequence as represented by SEQ ID NO: 119 are presented Table D1. The "plant" organism group has been selected, no cutoffs defined, and the predicted length of the transit peptide requested. The subcellular localization of the polypeptide sequence as represented by SEQ ID NO: 119 may be the cytoplasm or nucleus, no transit peptide is predicted.

Table D1: TargetP 1.1 analysis of the polypeptide sequence as represented by SEQ ID NO: 119. Abbreviations: Len, Length; cTP, Chloroplastic transit peptide; mTP, Mitochondrial transit peptide, SP, Secretory pathway signal peptide, other, Other subcellular targeting, Loc, Predicted Location; RC, Reliability class; TPlen, Predicted transit peptide length.

Name	Len	cTP	mTP	SP	other	Loc	RC	TPlen

SEQ ID NO: 119	979	0.103	0.321	0.050	0.554	_	4	-

cutoff	0.000	0.000	0.000	0.000				

Many other algorithms can be used to perform such analyses, including:

- ChloroP 1.1 hosted on the server of the Technical University of Denmark;
- Protein Prowler Subcellular Localisation Predictor version 1.2 hosted on the server of the Institute for Molecular Bioscience, University of Queensland, Brisbane, Australia;
- PENCE Proteome Analyst PA-GOSUB 2.5 hosted on the server of the University of Alberta, Edmonton, Alberta, Canada;
- TMHMM, hosted on the server of the Technical University of Denmark
- PSORT (URL: psort.org)
- PLOC (Park and Kanehisa, Bioinformatics, 19, 1656-1663, 2003).

Example 6: Assay related to the polypeptide sequences useful in performing the methods of the invention

1. CLC-pKG polypeptides

Marmagne et al 2007 described a method functionally complements the gef-1 yeast mutant. In brief, suspensions of the indicated strains W303 (WT), RGY86 (gef-1 mutant,) and each transformant (gef-1:CLC-of interest) are set up at DO1 (53108 cells). Serial dilutions (d1–d16) are spotted onto the appropriate media. For the low-iron phenotype, a 4-fold dilution of each suspension may be spotted onto YPEG media supplemented with 0.6 mM ferrozine, pH 5.8 (A, B) and grown for 3 d at 30 °C. For the pH-dependent phenotype, an 8-fold dilution is spotted onto YPD (C), SD 6 0.1 mM CuSO₄, pH 7 (D, E) and SGE 6 CuSO₄ 0.1 mM, pH7 (F, G) media and is grown for 3 d at 30 °C.

In addition or alternatively the patch clamp based method described by De Angeli (Nature 442, 939-942, 2006) may be used to measure currents across the tonoplast. The patch clamp technique is well known in the art for studying ion channel activity in cells, see for example Hamill et al., Pflügers Archiv (European Journal of Physiology) 391, 85-100, 1981.

Example 7: Cloning of the nucleic acid sequence used in the methods of the invention

1. SGT1 polypeptides

The nucleic acid sequence was amplified by PCR using as template a Capsicum annuum cDNA library (in pCMV Sport 6.0; Invitrogen, Paisley, UK). PCR was performed using Hifi

Taq DNA polymerase in standard conditions, using 200 ng of template in a 50 µl PCR mix. The primers used were prm14133 (SEQ ID NO: 51; sense, start codon in bold): 5'-aaaaa gcaggctcacaatggagaatgggaaaagagac-3' and prm14134 (SEQ ID NO: 52; reverse, complementary): 5'-agaaagctgggttggtttaactagttccaccg-3', which include the AttB sites for Gateway recombination. The amplified PCR fragment was purified also using standard methods. The first step of the Gateway procedure, the BP reaction, was then performed, during which the PCR fragment recombined in vivo with the pDONR201 plasmid to produce, according to the Gateway terminology, an "entry clone", pSGT1. Plasmid pDONR201 was purchased from Invitrogen, as part of the Gateway® technology.

The entry clone comprising SEQ ID NO: 1 was then used in an LR reaction with a destination vector used for *Oryza sativa* transformation. This vector contained as functional elements within the T-DNA borders: a plant selectable marker; a screenable marker expression cassette; and a Gateway cassette intended for LR in vivo recombination with the nucleic acid sequence of interest already cloned in the entry clone. A rice RCc3 promoter (SEQ ID NO: 50) for root-specific expression was located upstream of this Gateway cassette.

After the LR recombination step, the resulting expression vector pRCc3::SGT1 (Figure 4) was transformed into *Agrobacterium* strain LBA4044 according to methods well known in the art.

2. CLC-pKG polypeptides

The nucleic acid sequence was amplified by PCR using as template a custom-made *Synechocystis* DNA. PCR was performed using Hifi Taq DNA polymerase in standard conditions, using 200 ng of template in a 50 µl PCR mix. The primers used were: 5'-ggggacaagttgtacaaaaaagcaggcttaacaatggcttggttccctttt-3' (SEQ ID NO: 116) and 5'-ggggaccactttgtacaagaaagctgggtttactcgtctagaccgaaatctg-3' (SEQ ID NO: 117), which include the AttB sites for Gateway recombination. The amplified PCR fragment was purified also using standard methods. The first step of the Gateway procedure, the BP reaction, was then performed, during which the PCR fragment recombined in vivo with the pDONR201 plasmid to produce, according to the Gateway terminology, an "entry clone", pCLC-pKG. Plasmid pDONR201 was purchased from Invitrogen, as part of the Gateway® technology.

The entry clone comprising the longest Open Reading Frame of SEQ ID NO: 53 was then used in an LR reaction with a destination vector used for *Oryza sativa* transformation. This vector contained as functional elements within the T-DNA borders: a plant selectable marker; a screenable marker expression cassette; and a Gateway cassette intended for LR in vivo recombination with the nucleic acid sequence of interest already cloned in the entry clone. A rice RCc3 promoter (SEQ ID NO: 115) for root specific expression was located upstream of this Gateway cassette.

After the LR recombination step, the resulting expression vector pGOS2::CLC-pKG (Figure 7) was transformed into *Agrobacterium* strain LBA4044 according to methods well known in the art.

3. HD-hydrolase-like polypeptides

The nucleic acid sequence was amplified by PCR using as template a custom-made *Synechocystis* cDNA library (in pCMV Sport 6.0; Invitrogen, Paisley, UK). PCR was performed using Hifi Taq DNA polymerase in standard conditions, using 200 ng of template in a 50 µl PCR mix. The primers used were prm14240 (SEQ ID NO: 165; sense, start codon in bold): 5'-ggggacaagttgtacaaaaagcaggcttaacaatgtctgaagttactggcaag-3' and prm14239 (SEQ ID NO: 166; reverse, complementary): 5'-ggggaccactttgtacaagaaagctgggttaattgggaaatttaagtgaacg-3', which include the AttB sites for Gateway recombination. The amplified PCR fragment was purified also using standard methods. The first step of the Gateway procedure, the BP reaction, was then performed, during which the PCR fragment recombined in vivo with the pDONR201 plasmid to produce, according to the Gateway terminology, an "entry clone", pHD-hydrolase-like. Plasmid pDONR201 was purchased from Invitrogen, as part of the Gateway® technology.

The entry clone comprising SEQ ID NO: 118 was then used in an LR reaction with a destination vector used for *Oryza sativa* transformation. This vector contained as functional elements within the T-DNA borders: a plant selectable marker; a screenable marker expression cassette; and a Gateway cassette intended for LR in vivo recombination with the nucleic acid sequence of interest already cloned in the entry clone. A rice GOS2 promoter (SEQ ID NO: 164) for constitutive specific expression was located upstream of this Gateway cassette.

After the LR recombination step, the resulting expression vector pGOS2::HD-hydrolase-like (Figure 11) was transformed into *Agrobacterium* strain LBA4044 according to methods well known in the art.

Example 8: Plant transformation

Rice transformation

The *Agrobacterium* containing the expression vector was used to transform *Oryza sativa* plants. Mature dry seeds of the rice japonica cultivar Nipponbare were dehusked. Sterilization was carried out by incubating for one minute in 70% ethanol, followed by 30 minutes in 0.2% HgCl₂, followed by a 6 times 15 minutes wash with sterile distilled water. The sterile seeds were then germinated on a medium containing 2,4-D (callus induction medium). After incubation in the dark for four weeks, embryogenic, scutellum-derived calli were excised and propagated on the same medium. After two weeks, the calli were multiplied or propagated by subculture on the same medium for another 2 weeks.

Embryogenic callus pieces were sub-cultured on fresh medium 3 days before co-cultivation (to boost cell division activity).

Agrobacterium strain LBA4404 containing the expression vector was used for co-cultivation. *Agrobacterium* was inoculated on AB medium with the appropriate antibiotics and cultured for 3 days at 28°C. The bacteria were then collected and suspended in liquid co-cultivation medium to a density (OD₆₀₀) of about 1. The suspension was then transferred to a Petri dish and the calli immersed in the suspension for 15 minutes. The callus tissues were then blotted dry on a filter paper and transferred to solidified, co-cultivation medium and incubated for 3 days in the dark at 25°C. Co-cultivated calli were grown on 2,4-D-containing medium for 4 weeks in the dark at 28°C in the presence of a selection agent. During this period, rapidly growing resistant callus islands developed. After transfer of this material to a regeneration medium and incubation in the light, the embryogenic potential was released and shoots developed in the next four to five weeks. Shoots were excised from the calli and incubated for 2 to 3 weeks on an auxin-containing medium from which they were transferred to soil. Hardened shoots were grown under high humidity and short days in a greenhouse.

Approximately 35 independent T0 rice transformants were generated for one construct. The primary transformants were transferred from a tissue culture chamber to a greenhouse. After a quantitative PCR analysis to verify copy number of the T-DNA insert, only single copy transgenic plants that exhibit tolerance to the selection agent were kept for harvest of T1 seed. Seeds were then harvested three to five months after transplanting. The method yielded single locus transformants at a rate of over 50 % (Aldemita and Hodges 1996, Chan et al. 1993, Hiei et al. 1994).

Example 9: Transformation of other crops

Corn transformation

Transformation of maize (*Zea mays*) is performed with a modification of the method described by Ishida et al. (1996) *Nature Biotech* 14(6): 745-50. Transformation is genotype-dependent in corn and only specific genotypes are amenable to transformation and regeneration. The inbred line A188 (University of Minnesota) or hybrids with A188 as a parent are good sources of donor material for transformation, but other genotypes can be used successfully as well. Ears are harvested from corn plant approximately 11 days after pollination (DAP) when the length of the immature embryo is about 1 to 1.2 mm. Immature embryos are cocultivated with *Agrobacterium tumefaciens* containing the expression vector, and transgenic plants are recovered through organogenesis. Excised embryos are grown on callus induction medium, then maize regeneration medium, containing the selection agent (for example imidazolinone but various selection markers can be used). The Petri plates are incubated in the light at 25 °C for 2-3 weeks, or until shoots develop. The green shoots are transferred from each embryo to maize rooting medium and incubated at 25 °C for 2-3 weeks, until roots develop. The rooted shoots are transplanted to soil in the

greenhouse. T1 seeds are produced from plants that exhibit tolerance to the selection agent and that contain a single copy of the T-DNA insert.

Wheat transformation

Transformation of wheat is performed with the method described by Ishida et al. (1996) *Nature Biotech* 14(6): 745-50. The cultivar Bobwhite (available from CIMMYT, Mexico) is commonly used in transformation. Immature embryos are co-cultivated with *Agrobacterium tumefaciens* containing the expression vector, and transgenic plants are recovered through organogenesis. After incubation with *Agrobacterium*, the embryos are grown in vitro on callus induction medium, then regeneration medium, containing the selection agent (for example imidazolinone but various selection markers can be used). The Petri plates are incubated in the light at 25 °C for 2-3 weeks, or until shoots develop. The green shoots are transferred from each embryo to rooting medium and incubated at 25 °C for 2-3 weeks, until roots develop. The rooted shoots are transplanted to soil in the greenhouse. T1 seeds are produced from plants that exhibit tolerance to the selection agent and that contain a single copy of the T-DNA insert.

Soybean transformation

Soybean is transformed according to a modification of the method described in the Texas A&M patent US 5,164,310. Several commercial soybean varieties are amenable to transformation by this method. The cultivar Jack (available from the Illinois Seed foundation) is commonly used for transformation. Soybean seeds are sterilised for in vitro sowing. The hypocotyl, the radicle and one cotyledon are excised from seven-day old young seedlings. The epicotyl and the remaining cotyledon are further grown to develop axillary nodes. These axillary nodes are excised and incubated with *Agrobacterium tumefaciens* containing the expression vector. After the cocultivation treatment, the explants are washed and transferred to selection media. Regenerated shoots are excised and placed on a shoot elongation medium. Shoots no longer than 1 cm are placed on rooting medium until roots develop. The rooted shoots are transplanted to soil in the greenhouse. T1 seeds are produced from plants that exhibit tolerance to the selection agent and that contain a single copy of the T-DNA insert.

Rapeseed/canola transformation

Cotyledonary petioles and hypocotyls of 5-6 day old young seedling are used as explants for tissue culture and transformed according to Babic et al. (1998, *Plant Cell Rep* 17: 183-188). The commercial cultivar Westar (Agriculture Canada) is the standard variety used for transformation, but other varieties can also be used. Canola seeds are surface-sterilized for in vitro sowing. The cotyledon petiole explants with the cotyledon attached are excised from the in vitro seedlings, and inoculated with *Agrobacterium* (containing the expression vector) by dipping the cut end of the petiole explant into the bacterial suspension. The explants are then cultured for 2 days on MSBAP-3 medium containing 3 mg/l BAP, 3 % sucrose, 0.7 % Phytagar at 23 °C, 16 hr light. After two days of co-cultivation with *Agrobacterium*, the

petiole explants are transferred to MSBAP-3 medium containing 3 mg/l BAP, cefotaxime, carbenicillin, or timentin (300 mg/l) for 7 days, and then cultured on MSBAP-3 medium with cefotaxime, carbenicillin, or timentin and selection agent until shoot regeneration. When the shoots are 5 – 10 mm in length, they are cut and transferred to shoot elongation medium (MSBAP-0.5, containing 0.5 mg/l BAP). Shoots of about 2 cm in length are transferred to the rooting medium (MS0) for root induction. The rooted shoots are transplanted to soil in the greenhouse. T1 seeds are produced from plants that exhibit tolerance to the selection agent and that contain a single copy of the T-DNA insert.

Alfalfa transformation

A regenerating clone of alfalfa (*Medicago sativa*) is transformed using the method of (McKersie et al., 1999 Plant Physiol 119: 839–847). Regeneration and transformation of alfalfa is genotype dependent and therefore a regenerating plant is required. Methods to obtain regenerating plants have been described. For example, these can be selected from the cultivar Rangelander (Agriculture Canada) or any other commercial alfalfa variety as described by Brown DCW and A Atanassov (1985. Plant Cell Tissue Organ Culture 4: 111-112). Alternatively, the RA3 variety (University of Wisconsin) has been selected for use in tissue culture (Walker et al., 1978 Am J Bot 65:654-659). Petiole explants are cocultivated with an overnight culture of *Agrobacterium tumefaciens* C58C1 pMP90 (McKersie et al., 1999 Plant Physiol 119: 839–847) or LBA4404 containing the expression vector. The explants are cocultivated for 3 d in the dark on SH induction medium containing 288 mg/ L Pro, 53 mg/ L thioproline, 4.35 g/ L K₂SO₄, and 100 µm acetosyringinone. The explants are washed in half-strength Murashige-Skoog medium (Murashige and Skoog, 1962) and plated on the same SH induction medium without acetosyringinone but with a suitable selection agent and suitable antibiotic to inhibit *Agrobacterium* growth. After several weeks, somatic embryos are transferred to BOi2Y development medium containing no growth regulators, no antibiotics, and 50 g/ L sucrose. Somatic embryos are subsequently germinated on half-strength Murashige-Skoog medium. Rooted seedlings were transplanted into pots and grown in a greenhouse. T1 seeds are produced from plants that exhibit tolerance to the selection agent and that contain a single copy of the T-DNA insert.

Cotton transformation

Cotton is transformed using *Agrobacterium tumefaciens* according to the method described in US 5,159,135. Cotton seeds are surface sterilised in 3% sodium hypochlorite solution during 20 minutes and washed in distilled water with 500 µg/ml cefotaxime. The seeds are then transferred to SH-medium with 50µg/ml benomyl for germination. Hypocotyls of 4 to 6 days old seedlings are removed, cut into 0.5 cm pieces and are placed on 0.8% agar. An *Agrobacterium* suspension (approx. 10⁸ cells per ml, diluted from an overnight culture transformed with the gene of interest and suitable selection markers) is used for inoculation of the hypocotyl explants. After 3 days at room temperature and lighting, the tissues are transferred to a solid medium (1.6 g/l Gelrite) with Murashige and Skoog salts with B5 vitamins (Gamborg et al., Exp. Cell Res. 50:151-158 (1968)), 0.1 mg/l 2,4-D, 0.1 mg/l 6-

furfurylaminopurine and 750 µg/ml MgCL₂, and with 50 to 100 µg/ml cefotaxime and 400-500 µg/ml carbenicillin to kill residual bacteria. Individual cell lines are isolated after two to three months (with subcultures every four to six weeks) and are further cultivated on selective medium for tissue amplification (30°C, 16 hr photoperiod). Transformed tissues are subsequently further cultivated on non-selective medium during 2 to 3 months to give rise to somatic embryos. Healthy looking embryos of at least 4 mm length are transferred to tubes with SH medium in fine vermiculite, supplemented with 0.1 mg/l indole acetic acid, 6 furfurylaminopurine and gibberellic acid. The embryos are cultivated at 30°C with a photoperiod of 16 hrs, and plantlets at the 2 to 3 leaf stage are transferred to pots with vermiculite and nutrients. The plants are hardened and subsequently moved to the greenhouse for further cultivation.

Example 10: Phenotypic evaluation procedure

10.1 Evaluation setup

Approximately 35 independent T₀ rice transformants were generated. The primary transformants were transferred from a tissue culture chamber to a greenhouse for growing and harvest of T₁ seed. Six events, of which the T₁ progeny segregated 3:1 for presence/absence of the transgene, were retained. For each of these events, approximately 10 T₁ seedlings containing the transgene (hetero- and homo-zygotes) and approximately 10 T₁ seedlings lacking the transgene (nullizygotes) were selected by monitoring visual marker expression. The transgenic plants and the corresponding nullizygotes were grown side-by-side at random positions. Greenhouse conditions were of shorts days (12 hours light), 28°C in the light and 22°C in the dark, and a relative humidity of 70%. Plants grown under non-stress conditions were watered at regular intervals to ensure that water and nutrients were not limiting and to satisfy plant needs to complete growth and development.

Four T₁ events were further evaluated in the T₂ generation following the same evaluation procedure as for the T₁ generation but with more individuals per event. From the stage of sowing until the stage of maturity the plants were passed several times through a digital imaging cabinet. At each time point digital images (2048x1536 pixels, 16 million colours) were taken of each plant from at least 6 different angles.

10.2 Statistical analysis: F test

A two factor ANOVA (analysis of variants) was used as a statistical model for the overall evaluation of plant phenotypic characteristics. An F test was carried out on all the parameters measured of all the plants of all the events transformed with the gene of the present invention. The F test was carried out to check for an effect of the gene over all the transformation events and to verify for an overall effect of the gene, also known as a global gene effect. The threshold for significance for a true global gene effect was set at a 5% probability level for the F test. A significant F test value points to a gene effect, meaning

that it is not only the mere presence or position of the gene that is causing the differences in phenotype.

Where two experiments with overlapping events were carried out, a combined analysis was performed. This is useful to check consistency of the effects over the two experiments, and if this is the case, to accumulate evidence from both experiments in order to increase confidence in the conclusion. The method used was a mixed-model approach that takes into account the multilevel structure of the data (i.e. experiment - event - segregants). P values were obtained by comparing likelihood ratio test to chi square distributions.

10.3 Parameters measured

Biomass-related parameter measurement

From the stage of sowing until the stage of maturity the plants were passed several times through a digital imaging cabinet. At each time point digital images (2048x1536 pixels, 16 million colours) were taken of each plant from at least 6 different angles.

The plant aboveground area (or leafy biomass) was determined by counting the total number of pixels on the digital images from aboveground plant parts discriminated from the background. This value was averaged for the pictures taken on the same time point from the different angles and was converted to a physical surface value expressed in square mm by calibration. Experiments show that the aboveground plant area measured this way correlates with the biomass of plant parts above ground. The above ground area is the area measured at the time point at which the plant had reached its maximal leafy biomass. The early vigour is the plant (seedling) aboveground area three weeks post-germination. Increase in root biomass is expressed as an increase in total root biomass (measured as maximum biomass of roots observed during the lifespan of a plant); or as an increase in the root/shoot index (measured as the ratio between root mass and shoot mass in the period of active growth of root and shoot).

Early vigour was determined by counting the total number of pixels from aboveground plant parts discriminated from the background. This value was averaged for the pictures taken on the same time point from different angles and was converted to a physical surface value expressed in square mm by calibration. The results described below are for plants three weeks post-germination.

Seed-related parameter measurements

The mature primary panicles were harvested, counted, bagged, barcode-labelled and then dried for three days in an oven at 37°C. The panicles were then threshed and all the seeds were collected and counted. The filled husks were separated from the empty ones using an air-blowing device. The empty husks were discarded and the remaining fraction was counted again. The filled husks were weighed on an analytical balance. The number of filled seeds was determined by counting the number of filled husks that remained after the

separation step. The total seed yield was measured by weighing all filled husks harvested from a plant. Total seed number per plant was measured by counting the number of husks harvested from a plant. Thousand Kernel Weight (TKW) is extrapolated from the number of filled seeds counted and their total weight. The Harvest Index (HI) in the present invention is defined as the ratio between the total seed yield and the above ground area (mm²), multiplied by a factor 10⁶. The total number of flowers per panicle as defined in the present invention is the ratio between the total number of seeds and the number of mature primary panicles. The seed fill rate as defined in the present invention is the proportion (expressed as a %) of the number of filled seeds over the total number of seeds (or florets).

Examples 11: Results of the phenotypic evaluation of the transgenic plants

1. SGT1 polypeptides

The results of the evaluation of transgenic rice plants in the T2 generation and expressing a nucleic acid comprising the longest Open Reading Frame in SEQ ID NO: 1 under non-stress conditions are presented below.

Parameter	% Overall
Total weight seeds	10.5
Fill rate	10.6
Harvest index	10.0
Number filled seeds	11.3
Number of Flowers per panicle	6.6

For each parameter, the percentage overall difference between transgenic and control plants (corresponding nullizygotes) is shown if value reaches $p < 0.05$ and above the 5% threshold.

2. CLC-pKG polypeptides

The results of the evaluation of transgenic rice plants in the T1 generation and expressing a nucleic acid comprising the Open Reading Frame in SEQ ID NO: 53 encoding the polypeptide of SEQ ID NO: 54 under non-stress conditions are presented below. See previous Examples for details on the generations of the transgenic plants.

The results of the evaluation of transgenic rice plants under non-stress conditions are presented below. An increase of at least 5 % was observed for aboveground biomass (AreaMax), plant height (HeightMax), emergence vigour (early vigour; EmerVigor), root biomass (RootMax), total seed yield (totalwgseeds), harvest index (harvestindex), Proportion of thick roots compared to thin roots (RootThickMax), Gravity center of the canopy of the plants (GravityYMax). The last two parameters are indicative of a change in the architecture of the root system and of the canopy of the plant, respectively.

Table E1: Data summary for transgenic rice plants; for each yield-related trait, the overall percent increase is shown for the T1 generation, for each parameter the p-value is <0.05.

Yield related trait	% increase in the transgenic compared to the control plants
AreaMax	10.6
EmerVigor	6.0
RootMax	11.1
totalwgseeds	17.4
harvestindex	9.3
HeightMax	5.7
GravityYMax	7.3
RootThickMax	7.8

3. HD-hydrolase-like polypeptides

The results of the evaluation of transgenic T1 rice plants and expressing a nucleic acid encoding the polypeptide of SEQ ID NO: 119 under non-stress conditions are presented below in Table E2.

Table E2: Data summary for transgenic rice plants; for each parameter, the overall percent increase is shown for the confirmation (T1 generation), for each parameter the p-value is <0.05.

Parameter	Overall increase
totalwgseeds	15.4
fillrate	12.7
harvestindex	14.1
flowerperpan	9.2
Height	7.0

Claims

1. A method for enhancing yield-related traits in plants relative to control plants, comprising modulating expression in a plant of a nucleic acid encoding an SGT1 polypeptide, wherein said SGT1 polypeptide comprises the following in any order:
 - (i) at least one tetratricopeptide (TPR) repeat;
 - (ii) at least one CS domain; and
 - (iii) an SGS domain.
2. Method according to claim 1, wherein said modulated expression is effected by introducing and expressing in a plant a nucleic acid encoding an SGT1 polypeptide.
3. Method according to claim 1 or 2, wherein said nucleic acid encoding an SGT1 polypeptide encodes any one of the proteins listed in Table A1 or is a portion of such a nucleic acid, or a nucleic acid capable of hybridising with such a nucleic acid.
4. Method according to any one of claims 1 to 3, wherein said nucleic acid sequence encodes an orthologue or paralogue of any of the proteins given in Table A1.
5. Method according to any preceding claim, wherein said enhanced yield-related traits comprise increased seed yield relative to control plants.
6. Method according to any one of claims 1 to 5, wherein said enhanced yield-related traits are obtained under non-stress conditions.
7. Method according to any one of claims 2 to 6, wherein said nucleic acid is operably linked to a root-specific promoter, preferably to an RCc3 promoter, most preferably to an RCc3 promoter from rice.
8. Method according to any one of claims 1 to 7, wherein said nucleic acid encoding a SGT1 polypeptide is of plant origin, preferably from *Capsicum annuum* a dicotyledonous plant, further preferably from the family Solanaceae, more preferably from the genus *Capsicum*, most preferably from *Capsicum annuum*.
9. Plant or part thereof, including seeds, obtainable by a method according to any one of claims 1 to 8, wherein said plant or part thereof comprises a recombinant nucleic acid encoding an SGT1 polypeptide.
10. Construct comprising:
 - (i) nucleic acid encoding an SGT1 polypeptide as defined in claim 1;
 - (ii) one or more control sequences capable of driving expression of the nucleic acid sequence of (a); and optionally

- (iii) a transcription termination sequence.
11. Construct according to claim 10, wherein one of said control sequences is a root-specific promoter, preferably an RCc3 promoter, most preferably an RCc3 promoter from rice.
 12. Use of a construct according to claim 10 or 11 in a method for making plants having increased yield, particularly increased seed yield relative to control plants.
 13. Plant, plant part or plant cell transformed with a construct according to claim 10 or 11.
 14. Method for the production of a transgenic plant having increased yield, particularly increased seed yield relative to control plants, comprising:
 - (i) introducing and expressing in a plant a nucleic acid encoding an SGT1 polypeptide as defined in claim 1; and
 - (ii) cultivating the plant cell under conditions promoting plant growth and development.
 15. Transgenic plant having increased yield, particularly increased seed yield, relative to control plants, resulting from modulated expression of a nucleic acid encoding an SGT1 polypeptide as defined in claim 1, or a transgenic plant cell derived from said transgenic plant.
 16. Transgenic plant according to claim 9, 13 or 15, or a transgenic plant cell derived thereof, wherein said plant is a crop plant or a monocot or a cereal, such as rice, maize, wheat, barley, millet, rye, triticale, sorghum emmer, spelt, secale, einkorn, teff, milo and oats.
 17. Harvestable parts of a plant according to claim 16, wherein said harvestable parts are preferably shoot biomass and/or seeds.
 18. Products derived from a plant according to claim 16 and/or from harvestable parts of a plant according to claim 17.
 19. Use of a nucleic acid encoding an SGT1 polypeptide in increasing yield, particularly in increasing seed yield relative to control plants.
 20. Use of a nucleic acid encoding an SGT1 polypeptide as a molecular marker for plants having increased yield, particularly increased seed yield.
 21. A method for enhancing yield-related traits in plants relative to control plants, comprising modulating expression in a plant of a nucleic acid encoding a CLC-pKG

polypeptide, wherein said CLC-pKG polypeptide comprises a Voltage_CLC domain (Pfam entry PF00654) and a CBS domain (Pfam entry PF00571) and optionally a USP domain (PF00582).

22. Method according to claim 21, wherein said CLC-pKG polypeptide comprises a domain having in increasing order of preference at least 25%, 26%, 27%, 28%, 29%, 30%, 31%, 32%, 33%, 34%, 35%, 36%, 37%, 38%, 39%, 40%, 41%, 42%, 43%, 44%, 45%, 46%, 47%, 48%, 49%, 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100 % sequence identity to:
- (i) a CLC in SEQ ID NO: 54 as represented by the sequence located between amino acids 77 and 422 of SEQ ID NO: 54;
 - (ii) a CBS domain in SEQ ID NO: 54 as represented by the sequence located between amino acids 455 to 510 or 516 to 571 SEQ ID NO: 54; and optionally
 - (iii) a USP domain in SEQ ID NO: 54 as represented by the sequence located between amino acids 597-731 of SEQ ID NO: 54.
23. Method according to claim 21 or 22, wherein said modulated expression is effected by introducing and expressing in a plant a genetic construct comprising a nucleic acid encoding a CLC-pKG polypeptide.
24. Method according to any one of claims 21 to 23, wherein said nucleic acid encoding a CLC-pKG polypeptide encodes any one of the proteins listed in Table A2 or is a portion of such a nucleic acid, or a nucleic acid capable of hybridising with such a nucleic acid.
25. Method according to any one of claims 21 to 24, wherein said nucleic acid encodes an orthologue or paralogue of any of the proteins given in Table A2.
26. Method according to any one of claims 21 to 25, wherein said enhanced yield-related traits comprise increased yield, preferably increased biomass and/or increased seed yield relative to control plants.
27. Method according to any one of claims 21 to 26, wherein said enhanced yield-related traits are obtained under non-stress conditions or under conditions of drought stress, or salt stress.
28. Method according to any one of claims 23 to 27, wherein said nucleic acid is operably linked to a constitutive promoter, preferably to a RCc3 promoter, most preferably to a RCc3 promoter from rice.

29. Method according to any one of claims 21 to 28, wherein said nucleic acid encoding a CLC-pKG polypeptide is of cyanobacterial origin, further preferably from *Synechosystis* species, more preferably from *Synechocystis* sp. PCC 6803.
30. Plant or part thereof, including seeds, obtainable by a method according to any one of claims 21 to 29, wherein said plant or part thereof comprises a recombinant nucleic acid encoding a CLC-pKG polypeptide.
31. Construct comprising:
 - (i) nucleic acid encoding a CLC-pKG polypeptide as defined in claims 21 or 22;
 - (ii) one or more control sequences capable of driving expression of the nucleic acid sequence of (a); and optionally
 - (iii) a transcription termination sequence.
32. Construct according to claim 31, wherein one of said control sequences is a constitutive promoter, preferably a RCc3 promoter, most preferably a RCc3 promoter from rice.
33. Use of a construct according to claim 31 or 32 in a method for making plants having increased yield, particularly increased biomass and/or increased seed yield relative to control plants.
34. Plant, plant part or plant cell transformed with a construct according to claim 31 or 32.
35. Method for the production of a transgenic plant having increased yield, particularly increased biomass and/or increased seed yield relative to control plants, comprising:
 - (i) introducing and expressing in a plant a nucleic acid encoding a CLC-pKG polypeptide as defined in claim 21 or 22; and
 - (ii) cultivating the plant cell under conditions promoting plant growth and development.
36. Transgenic plant having increased yield, particularly increased biomass and/or increased seed yield, relative to control plants, resulting from modulated expression of a recombinant nucleic acid encoding a CLC-pKG polypeptide as defined in claim 21 or 22, or a transgenic plant cell derived from said transgenic plant.
37. Transgenic plant according to claim 30, 34 or 36, or a transgenic plant cell derived thereof, wherein said plant is a crop plant or a monocot or a cereal, such as rice, maize, wheat, barley, millet, rye, triticale, sorghum emmer, spelt, secale, einkorn, teff, milo and oats.

38. Harvestable parts of a plant according to claim 37, wherein said harvestable parts are preferably shoot biomass and/or seeds.
39. Products derived from a plant according to claim 37 and/or from harvestable parts of a plant according to claim 38.
40. Use of a genetic construct comprising a nucleic acid encoding a CLC-pKG polypeptide in increasing yield, particularly in increasing seed yield and/or shoot biomass in plants, relative to control plants.
41. A method for enhancing yield-related traits in plants relative to control plants, comprising modulating expression in a plant of a nucleic acid encoding a HD-hydrolase-like polypeptide, wherein said HD-hydrolase-like polypeptide comprises a CRISPR-associated protein Crm2 domain.
42. Method according to claim 41, wherein said HD-hydrolase-like polypeptide comprises one or more of the motifs 11 to 26.
43. Method according to claim 41 or 42, wherein said modulated expression is effected by introducing and expressing in a plant a nucleic acid encoding a HD-hydrolase-like polypeptide.
44. Method according to any one of claims 41 to 43, wherein said nucleic acid encoding a HD-hydrolase-like polypeptide encodes any one of the proteins listed in Table A3 or is a portion of such a nucleic acid, or a nucleic acid capable of hybridising with such a nucleic acid.
45. Method according to any one of claims 41 to 44, wherein said nucleic acid sequence encodes an orthologue or paralogue of any of the proteins given in Table A3.
46. Method according to any one of claims 41 to 45, wherein said enhanced yield-related traits comprise increased yield, preferably increased biomass and/or increased seed yield relative to control plants.
47. Method according to any one of claims 41 to 46, wherein said enhanced yield-related traits are obtained under non-stress conditions.
48. Method according to any one of claims 43 to 47, wherein said nucleic acid is operably linked to a constitutive promoter, preferably to a GOS2 promoter, most preferably to a GOS2 promoter from rice.

49. Method according to any one of claims 41 to 48, wherein said nucleic acid encoding a HD-hydrolase-like polypeptide is of plant origin, preferably from a cyanophyte, further preferably from the family Chroococcaceae, more preferably from the genus *Synechocystis*.
50. Plant or part thereof, including seeds, obtainable by a method according to any one of claims 41 to 49, wherein said plant or part thereof comprises a recombinant nucleic acid encoding a HD-hydrolase-like polypeptide.
51. Construct comprising:
 - (i) nucleic acid encoding a HD-hydrolase-like polypeptide as defined in claims 41 or 42;
 - (ii) one or more control sequences capable of driving expression of the nucleic acid sequence of (a); and optionally
 - (iii) a transcription termination sequence.
52. Construct according to claim 51, wherein one of said control sequences is a constitutive promoter, preferably a GOS2 promoter, most preferably a GOS2 promoter from rice.
53. Use of a construct according to claim 51 or 52 in a method for making plants having increased yield, particularly increased biomass and/or increased seed yield relative to control plants.
54. Plant, plant part or plant cell transformed with a construct according to claim 51 or 52.
55. Method for the production of a transgenic plant having increased yield, particularly increased biomass and/or increased seed yield relative to control plants, comprising:
 - (i) introducing and expressing in a plant a nucleic acid encoding a HD-hydrolase-like polypeptide as defined in claim 41 or 42; and
 - (ii) cultivating the plant cell under conditions promoting plant growth and development.
56. Transgenic plant having increased yield, particularly increased biomass and/or increased seed yield, relative to control plants, resulting from modulated expression of a nucleic acid encoding a HD-hydrolase-like polypeptide as defined in claim 41 or 42, or a transgenic plant cell derived from said transgenic plant.
57. Transgenic plant according to claim 50, 54 or 56, or a transgenic plant cell derived thereof, wherein said plant is a crop plant or a monocot or a cereal, such as rice, maize, wheat, barley, millet, rye, triticale, sorghum emmer, spelt, secale, einkorn, teff, milo and oats.

58. Harvestable parts of a plant according to claim 57, wherein said harvestable parts are preferably shoot biomass and/or seeds.
59. Products derived from a plant according to claim 57 and/or from harvestable parts of a plant according to claim 58.
60. Use of a nucleic acid encoding a HD-hydrolase-like polypeptide in increasing yield, particularly in increasing seed yield and/or shoot biomass in plants, relative to control plants.

1/31

CLUSTAL W (1.81) multiple sequence alignment

```

A.thaliana_AT4G23570.1#1      ----MAKELADKAKEAFVDDDDFDVAVDLYSKAIDLDPNCAEFFADRAQAY
A.thaliana_AT4G23570.2#1      ----MAKELADKAKEAFVDDDDFDVAVDLYSKAIDLDPNCAEFFADRAQAY
A.thaliana_AT4G23570.3#1      ----MAKELADKAKEAFVDDDDFDVAVDLYSKAIDLDPNCAEFFADRAQAY
A.thaliana_AT4G11260.1#1      ----MAKELAEKAKEAFVDDDDFDVAVDLYSKAIDLDPNCAEFFADRAQAN
B.napus_BN06MC26625_51284379@2 ----MAKELAEKAKEAFVDDDDFDVAVDLYSKAIDLDPSCASFFADRAQAN
Z.mays_ZM07MC27211_BFb0187N13@ ---MAASDLESKAKEAFVDDDDFELAAELYTQAI DAGPATADLYADRAQAH
Z.mays_ZM07MC23831_BFb0038K08@ ---MAASDLESKAKEAFVDDDDFELATELYSQAI DAGPATADLYADRAQAH
O.sativa_LOC_Os01g43540.1#1   MATAAASDLESKAKAAFVDDDDFELAAELYTQAI EASPATAELYADRAQAH
T.aestivum_TC280790#1         -----
M.truncatula_CT025844_7.4#1    ----MAQELENKAKEAFVDDDDFTLAVDFYSQAI EIDPTNANLFADRAQSH
G.max_GM06MC38560_sab91h08@375 ----MASDLEAKAKEAFVEDHFELAVDLLSQAI HLEPNKAEFYADRAQAN
TraitMillCDS_25598_CDS6534_69_ ----MASDLETRAKEAFIDHDFELAVDLYTQAI SLSPKNPELFADRAQAN
SGT1_from_Pepper_KOGENT2_     ----MASDLETRAKEAFIDHDFELAVDLYTQAI SLSPKNPELFADRAQAN
S.lycopersicum_TC192025#1      ----MASDLETRAKEAFIDHDFELAVDLYTQAI TLSPKNPELFADRAQAN
S.lycopersicum_NP9243669#1     ----MASDLETRAKEAFIDHDFELAVDLYTQAI AMTPKNAELFADRAQAN
P.trichocarpa_scaff_IV.839#1   ----MADLEKKAKEAFIDHDFELAVDLYTQAI ALNPTNPDLADRAQAN
P.trichocarpa_scaff_44.273#1   ----MADLERKAKEAFIDHDFELAVDLYTQAI ALNPTNPDLFADRAQAN
S.lycopersicum_AW398985#1      ----MASDLVTKAKEAFIDHDFELAVDLYSQAI AITPNNADLFADRAQAN
H.vulgare_c62589592hv270303@64 ----MADEAKAKGNAAFSAGRFEAAAGHFSDAI ALAPANHVLYSNRSAAL

A.thaliana_AT4G23570.1#1      IKLESFT-EAVADANKAIELDPSLT KAYLRKGTACMKLEEYRTAKTALEK
A.thaliana_AT4G23570.2#1      IKLESFT-EAVADANKAIELDPSLT KAYLRKGTACMKLEEYRTAKTALEK
A.thaliana_AT4G23570.3#1      IKLESFTAEAVADANKAIELDPSLT KAYLRKGTACMKLEEYRTAKTALEK
A.thaliana_AT4G11260.1#1      IKIDNFT-EAVVDANKAIELEPTLAKAYLRKGTACMKLEEYSTAKAALEK
B.napus_BN06MC26625_51284379@2 IKLLNFT-EAVVDANKAIELEPTLAKAYLRKGTACMKLEEYATAKAALEK
Z.mays_ZM07MC27211_BFb0187N13@ IKLGNYT-EAVADANKAIGLDPTMHKAYYRKGAAIC KLEEYQTAKAALEL
Z.mays_ZM07MC23831_BFb0038K08@ IKLGNYT-EAVADANKAIELDPMMHKAYYRKGAAIC KLEEYQTAKAALEL
O.sativa_LOC_Os01g43540.1#1   IKLGNYT-EAVADANKAIELDPSMHKAYLRKGAACIR LEEYQTAKAALEL
T.aestivum_TC280790#1         -----
M.truncatula_CT025844_7.4#1    IKLNFT-EAVSDANKAIQLNPNSKAYLRKGTACIN LEEYHTAKVALEK
G.max_GM06MC38560_sab91h08@375 IKLLNFT-EAVADANKAIELNPSP LKAYLRKGTACMKLEEYETAKAALEV
TraitMillCDS_25598_CDS6534_69_ IKLLNFT-EAVVDANKAIELDPYMSKAYLRKGLACMK LEEYQTAKAALET
SGT1_from_Pepper_KOGENT2_     IKLNYFT-EAVVDANKAIELDPYMSKAYLRKGLACMK LEEYQTAKAALET
S.lycopersicum_TC192025#1      IKLNYFT-EAVVDANKAIELDPTMSKAYLRKGLACMK LEEYQTAKAALEI
S.lycopersicum_NP9243669#1     IKLNYFT-EAVVDANKAIELDPSMSKAYLRKGLACMK LEEYQTAKAALQT
P.trichocarpa_scaff_IV.839#1   IKLNSLT-GMLN-----
P.trichocarpa_scaff_44.273#1   IKLLNFT-EAVADASRAIALDASLAKAYLRKGIACMK LEEYQTAKAALEV
S.lycopersicum_AW398985#1      IKLRKFT-----
H.vulgare_c62589592hv270303@64 ASIHRYS-DALADAECTVELKPDWAKGY SRLGAHLGLGDAASAAAAYEK

A.thaliana_AT4G23570.1#1      GASITPSESKFKKLIDECNFLITEEEKDLVQVPV PSTLPSSVTAPPVSELD
A.thaliana_AT4G23570.2#1      GASITPSESKFKKLIDECNFLITEEEKDLVQVPV PSTLPSSVTAPPVSELD
A.thaliana_AT4G23570.3#1      GASITPSESKFKKLIDECNFLITEEEKDLVQVPV PSTLPSSVTAPPVSELD
A.thaliana_AT4G11260.1#1      GASVAPNEPKFKKMIDECNLLIAEEEKDLVQVPM PPSSTTPLATEAD
B.napus_BN06MC26625_51284379@2 GASVAPNESKFEKMIDECNLLIAEEEKDLVQVPM PPSSTTPLAIAAD
Z.mays_ZM07MC27211_BFb0187N13@ GSSYAPGDSRFTRLLEKCECIAEESSQAPAKNVEA ----PVAATVEDK
Z.mays_ZM07MC23831_BFb0038K08@ GSSYASGDSRFARLLKECDERIAEESSQAPVKNVEA ----TVAATIEDK
O.sativa_LOC_Os01g43540.1#1   GYSFASGDSRFTRLMKECDERIAEELSEVPVKKAEDGAAAPSVASFVEEK
T.aestivum_TC280790#1         -----
M.truncatula_CT025844_7.4#1    GASFAPDDSRFTNLIQQCRFIAAEESESLTSLPPN GPKSSVASVDDTH
G.max_GM06MC38560_sab91h08@375 GASLSPDNSRFATLIKECDKLIAEESYTPPIIEEKT TTEDATPKDVQQQD
TraitMillCDS_25598_CDS6534_69_ GASLAPGESRFTKLMKECEDSIAEEAGELPNISLEK TSANAVASSASELS
SGT1_from_Pepper_KOGENT2_     GASLAPGESRFTKLMKECEDSIAEEAGELPNISLEK TSANAVASSASELS
S.lycopersicum_TC192025#1      GASLAPGELRFSKLMKECEDHIAEEAGELPNLSVDK TSASVVAHPASELP
S.lycopersicum_NP9243669#1     GASLAPAESRFTKLIKECDERIAEEAGELPNQSVDK TSGNVVTAPP-ESL
P.trichocarpa_scaff_IV.839#1   -----
P.trichocarpa_scaff_44.273#1   GASLAPGESRFTNLIKECEDCIAEETGGSKNHAADAP VNTVSIEDVEPED
S.lycopersicum_AW398985#1      -----
H.vulgare_c62589592hv270303@64 GLALDPSNEGLKAGLADAKAAAAPRRPPSGGADAIG QMFQGPPELWTKI

```

FIGURE 1

2/31

A.thaliana_AT4G23570.1#1
 A.thaliana_AT4G23570.2#1
 A.thaliana_AT4G23570.3#1
 A.thaliana_AT4G11260.1#1
 B.napus_BN06MC26625_51284379@2
 Z.mays_ZM07MC27211_BFb0187N13@
 Z.mays_ZM07MC23831_BFb0038K08@
 O.sativa_LOC_Os01g43540.1#1
 T.aestivum_TC280790#1
 M.truncatula_CT025844_7.4#1
 G.max_GM06MC38560_sab91h08@375
 TraitMillCDS_25598_CDS6534_69_
 SGT1_from_Pepper_KOGENT2_
 S.lycopersicum_TC192025#1
 S.lycopersicum_NP9243669#1
 P.trichocarpa_scaff_IV.839#1
 P.trichocarpa_scaff_44.273#1
 S.lycopersicum_AW398985#1
 H.vulgare_c62589592hv270303@64

A.thaliana_AT4G23570.1#1
 A.thaliana_AT4G23570.2#1
 A.thaliana_AT4G23570.3#1
 A.thaliana_AT4G11260.1#1
 B.napus_BN06MC26625_51284379@2
 Z.mays_ZM07MC27211_BFb0187N13@
 Z.mays_ZM07MC23831_BFb0038K08@
 O.sativa_LOC_Os01g43540.1#1
 T.aestivum_TC280790#1
 M.truncatula_CT025844_7.4#1
 G.max_GM06MC38560_sab91h08@375
 TraitMillCDS_25598_CDS6534_69_
 SGT1_from_Pepper_KOGENT2_
 S.lycopersicum_TC192025#1
 S.lycopersicum_NP9243669#1
 P.trichocarpa_scaff_IV.839#1
 P.trichocarpa_scaff_44.273#1
 S.lycopersicum_AW398985#1
 H.vulgare_c62589592hv270303@64

A.thaliana_AT4G23570.1#1
 A.thaliana_AT4G23570.2#1
 A.thaliana_AT4G23570.3#1
 A.thaliana_AT4G11260.1#1
 B.napus_BN06MC26625_51284379@2
 Z.mays_ZM07MC27211_BFb0187N13@
 Z.mays_ZM07MC23831_BFb0038K08@
 O.sativa_LOC_Os01g43540.1#1
 T.aestivum_TC280790#1
 M.truncatula_CT025844_7.4#1
 G.max_GM06MC38560_sab91h08@375
 TraitMillCDS_25598_CDS6534_69_
 SGT1_from_Pepper_KOGENT2_
 S.lycopersicum_TC192025#1
 S.lycopersicum_NP9243669#1
 P.trichocarpa_scaff_IV.839#1
 P.trichocarpa_scaff_44.273#1
 S.lycopersicum_AW398985#1
 H.vulgare_c62589592hv270303@64

VTPT-----
 VTPT-----
 VTPT-----
 APPVPIPA-----
 APPVPSPA-----
 EDVANMDN-----
 EDFTNMEN-----
 DDAANMDN-----
 QDGANMEN-----
 MCDKSDETSK-----
 DLLEKPTVT-----
 DNVAIAPRDP-----
 DNVAIAPRDP-----
 DNVAIAPEGA-----
 DNVAVAPKDA-----

 TSSQAPMVI-----

 ASDPATRAYLDQPDFMQMLRDVQRDPSSLNMFSLSDPRMMQVLSLMLNIKI

-----AKYRHEYYQKPEEVVTVFAKGIP
 -----AKYRHEYYQKPEEVVTVFAKGIP
 -----AKYRHEYYQKPEEVVTVFAKGIP
 -----APAKPMFRHEFYQKPEEAVTIFAKKVP
 -----PPAKPMFRHEFYQKPEEVVTVFAKGIP
 -----TPPVVEPPSKPKYRHDYNSATEVVLTIFAKGVP
 -----TPPVEIEPPSKPKYRHDYNSATEVVLTIFAKGVP
 -----TPPMVEVK--PKYRHDYNSATEVVLTIFAKGVP
 -----AQPTVEVPSPKPKYRHDYNTPTPEVVLTIFAKGVP
 -----EPQRDSPASQTNVAVAPVRPKYRHEYYQKPEEVVTVFAKGIP
 -----VTKPKYRHEFYQKHDQLVVTIFAKKIP
 -----QSTVN----LSHQESAAPKPKYRHEFYQKPEEVVTVFAKGIP
 -----QSTVN----LSHQESAAPKPKYRHEFYQKPEEVVTVFAKGIP
 -----QPTVN----QSHQESAAPKPKYRHEFYQKPEEVVTVFAKGIP
 -----QPSVN----LSYQGSAAAPKPKYRHEFYQKPEEVVTVFAKGIP

 -----PSKPKYRHEFYQKPEEVVVSIFAKGVQ

 QTPQSDSDFPQSSSPSQPPPQQQRQQPETKAREMEPEPQPEPMEVSDDEKE

KQNVNIDFGEQILSVVIEVPGEDAYYLQPRLFGKIIPDKCKYEVLTSTKIE
 KQNVNIDFGEQILSVVIEVPGEDAYYLQPRLFGKIIPDKCKYEVLTSTKIE
 KQNVNIDFGEQILSVVIEVPGEDAYYLQPRLFGKIIPDKCKYEVLTSTKIE
 KENVTVFGEQILSVVIDVAGEEAYHLQPRLFGKIIPDKCKYEVLTSTKIE
 KQNVNVEFGDQILSVVIDVAGEEAYHFQPRLFGKIIPDKCKYEVLTSTKIE
 ADSVVIDFGEQILSVVIEVPGEEPYPHFQPRLFSKIIPDKCKYQVLSTKVE
 ADSVVIDFGEQILSVVIEVPGEEPYPHFQPRLFSKIIPDKCKYQVLSTKVE
 AENVVVDGEQILSVVIEVPGEEPYPHFQPRLFSKIIPDKCKYQVLSTKVE
 ADSVVVDGEQILSVVIEVPGEEPYPHFQPRLFSKIIPDKCKYQVLSTKVE
 AENVVVDGEQILSVVIDVPGQDAYHYQPRLFGKIIPDKCKYVVLSTKIE
 KESITVDGEQILSVSINVPGEDTYAFQPRLFGKIIPSNQCYEVLTSTKIE
 AKNVVVDGEQILSVSIDLPGGETYSFQPRLFGKITPAKCRYEVMSTKIE
 AKNVVVDGEQILSVSIDLPGGETYSFQPRLFGKITPAKCRYEVMSTKIE
 AKNVVVDGEQILSVSIDVPGEEYYSFQPRLFGKITPAKCRYEVMSTKIE
 AKNVVVDGEQILSVSIDVPGEEYYSFQPRLFGKITPAKCRYEVMSTKIE

 ASWISVDGEQILSVRIEVPGEDGYHFQPRLFGKIIPDKCKYNILSTKVE
 -----GNLFVNMVPLHSR-----
 KKERKAAALKEKEAGNASYKKKDFETAIQHYTKAMELDDDEDISYLTNRRA

FIGURE 1 (continued)

FIGURE 1 (continued)

4/31

```

A.thaliana_AT4G23570.1#1 -----KFFREIYQNADEDMRR-----A
A.thaliana_AT4G23570.2#1 -----KFFREIYQNADEDMRR-----A
A.thaliana_AT4G23570.3#1 -----KFFREIYQNADEDMRR-----A
A.thaliana_AT4G11260.1#1 -----KFFSDIYSSADEDMRR-----A
B.napus_BN06MC26625_51284379@2 -----KFFSDIYQSADEDMRR-----A
Z.mays_ZM07MC27211_BFb0187N13@ -----KFFRDIYKDADEDMRR-----A
Z.mays_ZM07MC23831_BFb0038K08@ -----KFFRDIYKDADEDMRR-----A
O.sativa_LOC_Os01g43540.1#1 -----KFFRDIYSDADEDMRR-----A
T.aestivum_TC280790#1 -----KFFREIYSDADEDMRR-----A
M.truncatula_CT025844_7.4#1 -----KLFRDIYQNADEDMRR-----A
G.max_GM06MC38560_sab91h08@375 -----KFFREIYQDADEDTRR-----A
TraitMillCDS_25598_CDS6534_69_ -----KFFRDIYKDADEDTRR-----A
SGT1_from_Pepper_KOGEN2_ -----KFFRDIYKDADEDTRR-----A
S.lycopersicum_TC192025#1 -----KFFRDIYQDADEDTRR-----A
S.lycopersicum_NP9243669#1 -----KFFRDIYKDADEDTRR-----A
P.trichocarpa_scaff_IV.839#1 -----
P.trichocarpa_scaff_44.273#1 -----KFFREIYQDADEDTRR-----A
S.lycopersicum_AW398985#1 -----
H.vulgare_c62589592hv270303@64 LKDAEKCIELDPTFSKGYTRKGAVQFFMKEYEKAMETYQAGLKLDPNNQE

```

```

A.thaliana_AT4G23570.1#1 MSKSFVESNGTVLSTNWQEVGTKTIESTPPDGMELKKWEI-----
A.thaliana_AT4G23570.2#1 MSKSFVESNGTVLSTNWQEVGTKTIESTPPDGMELKKWEI-----
A.thaliana_AT4G23570.3#1 MSKSFVESNGTVLSTNWQEVGTKTIESTPPDGMELKKWEI-----
A.thaliana_AT4G11260.1#1 MNKSFAESNGTVLSTNWKEVGTKKVESTPPDGMELKKWEY-----
B.napus_BN06MC26625_51284379@2 MNKSFAESNGTVLSTNWKEVGTKKVESTPPDGMELKKWEY-----
Z.mays_ZM07MC27211_BFb0187N13@ MDKSFRESNGTVLSTNWKDVGSKTVEASPPDGMELKKWEI-----
Z.mays_ZM07MC23831_BFb0038K08@ MMKSFVESNGTVLSTNWKDVGAKKVEGSPPDGMELKKWEY-----
O.sativa_LOC_Os01g43540.1#1 MMKSFVESNGTVLSTNWKDVGSKKVEGSPPDGMELKKWEY-----
T.aestivum_TC280790#1 MMKSFVESNGTVLSTNWKDVGKKTVEGSPPDGMELKKWEY-----
M.truncatula_CT025844_7.4#1 MSKSFLESNGTVLSTDWKEVGSKKVEGSPPEGMEVKKWEY-----
G.max_GM06MC38560_sab91h08@375 MKKSFVESNGTVLSTNWKEVGSKKVEGSPDGMELKKWEY-----
TraitMillCDS_25598_CDS6534_69_ MMKSFVESNGTVLSTNWKKVGTKKVEGSPPDGMELKKWEI-----
SGT1_from_Pepper_KOGEN2_ MMKSFVESNGTVLSTNWKKVGTKKVEGSPPDGMELKKWEI-----
S.lycopersicum_TC192025#1 MMKSFVESNGTVLSTNWKEVGTKKVEGSPPDGMELKKWEI-----
S.lycopersicum_NP9243669#1 MMKSFVESNGTVLSTNWKEVGAKKVEGSPPDGMELKKWEI-----
P.trichocarpa_scaff_IV.839#1 -----
P.trichocarpa_scaff_44.273#1 MKKSFVESNGTVLSTNWKEVGTKKVEGSPPDGMEMRKWEY-----
S.lycopersicum_AW398985#1 -----
H.vulgare_c62589592hv270303@64 LLDGIRRCVQQINKANRGDISQEDLQEQSKAMQDPEIQNILTDPIMRQV

```

```

A.thaliana_AT4G23570.1#1 -----
A.thaliana_AT4G23570.2#1 -----
A.thaliana_AT4G23570.3#1 -----
A.thaliana_AT4G11260.1#1 -----
B.napus_BN06MC26625_51284379@2 -----
Z.mays_ZM07MC27211_BFb0187N13@ -----
Z.mays_ZM07MC23831_BFb0038K08@ -----
O.sativa_LOC_Os01g43540.1#1 -----
T.aestivum_TC280790#1 -----
M.truncatula_CT025844_7.4#1 -----
G.max_GM06MC38560_sab91h08@375 -----
TraitMillCDS_25598_CDS6534_69_ -----
SGT1_from_Pepper_KOGEN2_ -----
S.lycopersicum_TC192025#1 -----
S.lycopersicum_NP9243669#1 -----
P.trichocarpa_scaff_IV.839#1 -----
P.trichocarpa_scaff_44.273#1 -----
S.lycopersicum_AW398985#1 -----
H.vulgare_c62589592hv270303@64 LMDFQENPRAAQDHLKDPGVAQKIQLINAGIVQTR

```

FIGURE 1 (continued)




FIGURE 2

6/31




FIGURE 3

7/31




FIGURE 4

8/31

CLUSTAL W (1.81) multiple sequence alignment

```

gi|218440518|ref|YP_002378847.      --MNIYLPKRLRGWFKARRFGRSSVDTRYAMLEACLIGILSAVAALVLK
gi|196258397|ref|ZP_03156931.1      --MNLPPFLKRLHGWFKARRFGRSSVDTRYAMLESCLIGILSAIAALLLK
gi|126659209|ref|ZP_01730347.1      -----MPPRFYQWLKSRKLAFNATDTRYALVEASLIGVLSAIAALILK
gi|172034995|ref|YP_001801496.      --MQPHLMPQRLYQWFKSRNFALNATDTRYALVEASLIGVLSAIAALILK
gi|67923321|ref|ZP_00516804.1|      --MGPHLIFQRFYRWLKSRLRTVDATDTRYALAEACLIGILSAIAALILK
gi|218248189|ref|YP_002373560.      --MKPHLIVKRLYQWVKSSHLGVNATDSRYAFVEACLLGLLSGIAALILK
gi|159026894|emb|CAO89145.1|_e      --MGLFARYQNLQWFKSQHFGRSATDSRYALLVACLIGILSALAAILK
gi|166366612|ref|YP_001658885.      --MGLFARHQNLQWFKSQHFGRSATDSRYALLVACLIGILSALAAILK
gi|16331172|ref|NP_441900.1|_c      --MAWFPPWQRWQWLRSRHYGRTAIDSRYAIAEACVIGFVSALAALVIK
gi|170077188|ref|YP_001733826.      -----MRNWWIFRSLGQHSNNNYLLIEACLIGVISGLAALFLK
gi|158334130|ref|YP_001515302.      -----MKLISAFQPLRHLTQSSRFATIEACLIGLISGLAVALQ
gi|119510301|ref|ZP_01629437.1      -----MAISGLTQRFRLWLQPTRGLAIAEACIIGLVAALSAVFLK
gi|186682049|ref|YP_001865245.      -----MNQRFRTWWQPRRGLAIAEACVIGLVAALSAVFLK
gi|75908381|ref|YP_322677.1|_C      -----MYFRSWLQPRRGLAIAEACVIGLVAALSAVFLK
gi|17232383|ref|NP_488931.1|_h      -----MYFRSWLQPRRGLAIAEACVIGLVAALSAVFLK
gi|225516217|ref|ZP_03763190.1      -----MSKARGRLPLPFWSMGYANAEASMIGIVAALSAVLLK
gi|254413597|ref|ZP_05027367.1      -----MINPIRQWLLPKRRLAIAEACLIGVVSALSAVLLQ
gi|186683293|ref|YP_001866489.      -----MLARGKILWLRLSRQLVQPRRLAFVEACLIGLVSGLAVALLG
gi|225522175|ref|ZP_03768989.1      -----MLARSKIFWLRLSRQLVKPRRLAFVEACVIGLVSGLAVALLG
gi|186682226|ref|YP_001865422.      -----MLARGKILWLRLSRQLLRPRRLAFVEACIIGLVSGLAVALLG
gi|220909579|ref|YP_002484890.      -----MGSLAALRRLQQLLRPRRVAVLEACFIGLVSALAAGGMA
gi|37523751|ref|NP_927128.1|_c      -----MQSWLQNFRLQLLRPQRLATLEACLIGLVAGLAAVILK
gi|119486852|ref|ZP_01620827.1      MLRISIPEQLSKRFKIIATQILQPSQRRLAILEACMIGLVSGLAFFCLR
gi|209525724|ref|ZP_03274261.1      ----MYWPSLTNQFQSLAKSIIG--GSQRLAILEAGLIGLISGIAAFFLR
gi|220908306|ref|YP_002483617.      -----MALMKGTIWLTSQSLQRLLRPKSVAILEACLIGLVSALAVALLK
gi|81300397|ref|YP_400605.1|_C      -----MFRLRFPPRFKAQQAQWVLEACAIGLVSGASAFLLK
                                     :  :*.:. :*  :

```

```

gi|218440518|ref|YP_002378847.      QGVGWLGGWRLS--LTRTFGALWILPLGGILGLLAGWVIEILSPAAAGG
gi|196258397|ref|ZP_03156931.1      QGVGWLGGWRVH--ITTKLGAVWLLPIGGLLLGLLAGQLIQVISPAAGG
gi|126659209|ref|ZP_01730347.1      QGVGWLGGWRVA--LANQFGAMWVPLPLGGILGLYLSGLIETFSPTATGG
gi|172034995|ref|YP_001801496.      QGVGWLGGWRVA--LANEFGGIVVPLPLGGILGLYLAGWIIETLSPTATGG
gi|67923321|ref|ZP_00516804.1|      QGVGWLGGKRVA--LANQFGAIWVPLPLGGILGLSMAGWLIETLSPTATGG
gi|218248189|ref|YP_002373560.      EGVGWLGGWRVS--LANQLGAIWVPLPLGGFLGLSMAGWLIETLSPTAAGG
gi|159026894|emb|CAO89145.1|_e      LGIGWLGGWRVH--LVANSSPFLVPLPLGGFLLGYSAGWIVEHFSPAAAGG
gi|166366612|ref|YP_001658885.      LGIGWLGGWRVH--LVANSSPFLVPLPLGGFLLGYSAGWIVEHFSPAAAGG
gi|16331172|ref|NP_441900.1|_c      NGVNWLGSSRLS--LADTYGAIWVLPFCFGLVFGGLAGALIEQFSRPAGGG
gi|170077188|ref|YP_001733826.      QGIGFVGGRYVI--LSRQYGAWLILPLFGVVCWGLAGYLLETFAPEAKGG
gi|158334130|ref|YP_001515302.      DGIGWLGGRLIS--LANQFHSPLILPLIGVIGGLIAGLLIETIAPDAKGS
gi|119510301|ref|ZP_01629437.1      FGSGFLGTWRVQ--TSQIVPALLVLPTIGLSFGFIAGSLVNRFAPEASGS
gi|186682049|ref|YP_001865245.      VGSGWLGTWRVH--TTHLFPAPWLVLPLVIGLVLFAGWLVDRLAPEAYGS
gi|75908381|ref|YP_322677.1|_C      VCSGLLGAWRVH--SSHVLPAWVVLPIIGLGFGLYLAGLMVQRLAPEAAGS
gi|17232383|ref|NP_488931.1|_h      VCSGLLGAWRVH--SSHILPAWVVLPIIGLSFGYLAGLMVQRLAPEAAGS
gi|225516217|ref|ZP_03763190.1      QGSGCLGTWRVH--STQFVPAWLILPLIGMSFGFLAGWLQRLAPEAAGS
gi|254413597|ref|ZP_05027367.1      HSIGWLGSWRVQ--TSEQFPFPLVLPFAFGLVLGALSGFLVEYLAPDASGS
gi|186683293|ref|YP_001866489.      QAVDWAGAWRVH--LSYHWSAYLVLPGIGLAGGLLAGWLVEHFAPASGS
gi|225522175|ref|ZP_03768989.1      QAVDWAGALVR--VSYHCPAYLVLPISIGLVGGILAGWLVECFAPASGS
gi|186682226|ref|YP_001865422.      QAVDWAGAWRVH--LSYIWPAYLVLPGIGLVGGILAGWLVERFAPEASGS
gi|220909579|ref|YP_002484890.      WGVGWMGTWRVN--LAEQWPDWLVLPLFGAIGALLSGALVLEAPEAAGS
gi|37523751|ref|NP_927128.1|_c      QGADLLSHWRLD--AAGSFG--WWVLPLLGLLGGVTAGWLVERFAPEASGS
gi|119486852|ref|ZP_01620827.1      VGAGWLGSWRIYGALNSPLPVWVFLPSVGLIGGFLTGFVERFAPETTGS
gi|209525724|ref|ZP_03274261.1      TGAGWLGSWRVQSAH--FSAWLFLPILGVVGGLLSGFLVERIAPEAAGS
gi|220908306|ref|YP_002483617.      QAIAAMESWTAA--IGADLPPLLILPVIGLLGGLLSGWLIEWAAPEATGS
gi|81300397|ref|YP_400605.1|_C      LGATGVQDLRDR-----PGLPLGLQLLLPPILGAIAGWLVRFAPEAEGS
                                     .      .      :* : :      :      :*.

```

FIGURE 5

GIPQVKAALARFPVPLSLRVALAKLIGAILVLGAGLTLGRRAPT VHIGAA
GIPQVKAALARFPVPLSLRVAIAKLLGATLVLGAGLTLGRRAPT VHIGAA
GITQVKAALARYPVPLSLRVAVVKLIGITILVLGGGLTLGRRAPT VHIGAA
GITQVKAALARYPVPLSLRVAIVKLIGITILVLGGGLTLGRRAPT VHIGAA
GMTQVKAALARYPVPLSLRVAVVKLIGITILVLGGGLTLGRRAPT VHIGAA
GITQVKAALARYPVPLSLRVAVVKLIGITILVLGGGLTLGRRAPT VHIGAA
GIPQVKAALAKYPLPLSWRVAVVKMIGAILILGGGLTLGRRAPT VHIGAA
GIPQVKAALAKYPLPLSWRVAVVKMIGAILILGGGLTLGRRAPT VHIGAA
GIPQVKAALARYPVPLDWRVAVVKLLGTVLVLGGGMTLGRRAPT VHIGAA
GVPQVKVVL AQFPPLPNWKVALAKSIGTILVLGGGLTLGRRGPSVHIGAA
GIPQVKAAL AQFPPIALNWQVALVKLFSGITALTGLPLGRQGPTVQIGAA
GIPQVKATLANVPITLSWRVAVVKLSAIIATIGSGMALGRQGPTVHVAG
GIPQVKASLANVPVNLSSWRVAVVKLSAIIATIGSGMTLGRQGPTVQVAG
GIPQVKATLANIPMKLSWRVAVIKLLSAIIALGSGITLGRQGPTVQVAG
GIPQVKATLANIPMKLSWRVAIVKLLSAIIALGSGTLGRQGPTVQVAG
GIPQVKASLANIPIKLSWRVEGIKLLSTIIISLGGITLGRQGPTVQIGAS
GIPQIKAALGGYPIALDLRVALVKLVSSILATAAGLTLGRQGPTVHIGAA
GMSEVKAVLARVPMPLNLRIALVKLIGATLVLGSGMPLGREGPTVQIGAA
GMSEVKAVLARVPMPLNLRIALVKLISATLVLGSGMPLGREGPTVQIGAA
GMSEVKAVLARVPMPLNLRIALVKLVSAITLVLGSGMPLGREGPTVQIGAA
GMSQVKAVLARVPMPLNLRIALVKLISAVLALGSGLALGREGPTVQVAGAA
GIPQVKAVLARVPMALDLRVALVKLVGTMVTLGSGFPLGREGPTIQIGAA
GVPQVKAALSGMPI SLDFRVAISKLLGTMFTMGSGTLGRQGPTVQIGAA
GIPQVKAALGGIKTSLDLRVAVVKLISTMITMASGFTLGRQGPTVQIGAA
GIPQVKAALGYASIALDLRVAVVKLVSTTLALGAGLALGRQGPTVQIGAA
GISQVQAALSGSRIALNRVAIVKIIASTMLVVGSGPLGRQGPTVQVGAS
* : : : : * : * : : * : : : : * : : : : * : : : *

gi|218440518|ref|YP_002378847.
gi|196258397|ref|ZP_03156931.1
gi|126659209|ref|ZP_01730347.1
gi|172034995|ref|YP_001801496.
gi|67923321|ref|ZP_00516804.1
gi|218248189|ref|YP_002373560.
gi|159026894|emb|CA089145.1|_e
gi|166366612|ref|YP_001658885.
gi|16331172|ref|NP_441900.1|_c
gi|170077188|ref|YP_001733826.
gi|158334130|ref|YP_001515302.
gi|119510301|ref|ZP_01629437.1
gi|186682049|ref|YP_001865245.
gi|75908381|ref|YP_322677.1|_C
gi|17232383|ref|NP_488931.1|_h
gi|225516217|ref|ZP_03763190.1
gi|254413597|ref|ZP_05027367.1
gi|186683293|ref|YP_001866489.
gi|225522175|ref|ZP_03768989.1
gi|186682226|ref|YP_001865422.
gi|220909579|ref|YP_002484890.
gi|37523751|ref|NP_927128.1|_c
gi|119486852|ref|ZP_01620827.1
gi|209525724|ref|ZP_03274261.1
gi|220908306|ref|YP_002483617.
gi|81300397|ref|YP_400605.1|_c

LAAQLSQWLPTSPEHRRQMI AAGAAAGLAAGFGTPIAGVLFVIEELMRDV
LAAQLSQWVPTSPEHRRQMI AAGAAAGLAAGFGTPIAGVLFVVEELMRDV
LAAQLSNWVPTSPEHRRQMI AAGAAAGLAAGFTTPIAGVLFVIEELMRDV
LAAQLSSWVPTSPEHRRQMI AAGAAAGLAAGFTTPIAGVLFVIEELMRDV
LAAQLSSWVPTSPEHRRQMI AAGAAAGLAAGFTTPIAGVLFVIEELMRDV
LAAQLSSWVPTSPEHRRQMI AAGAAAGLAAGFGTPIAGVLFVVEELMRDV
LAAQLSVWLPTS PDHRRQMI AAGAAAGLAAGFTTPIAGVLFVIEELMRDV
LAAQLSVWLPTS PDHRRQMI AAGAAAGLAAGFTTPIAGVLFVIEELMRDV
LAAQLSRWFPTSPEHRRQMI AAGAAAGLAAGFTTPIAGVLFVVEELMRDV
LAAQLSQWLPTT PNHRRQMI AAGAAAGLAAGFNTPIAGVLFVVEELMRDM
LAAQLSYWFPTAPNHRRQMI AAGAGAGLAAGFAAPIAGVLFVIEELLHDL
LAAGMSRWVPTS PDHRRQMI AAGAGAGLAAAFNAPIAGVLFVIEELLQDL
LAAGMSRWVPTS PDHRRQMI AAGAGAGLAAAFNAPIAGVLFVIEELLQDL
LAAGMSRLVPTS PDHRRQMI AAGAGAGLAAAFNAPIAGVLFVIEELLQDL
LAAGMSRLVPTS PDHRRQMI AAGAGAGLAAAFNAPIAGVLFVIEELLQDL
LAAGMSRTFPTSPEHRRQMI AAGAGAGLSAAFNAPIAGVLFVIEELLQDL
LAAQFSRWVPTS PSHRRQLI AAGSAAGLAAGFNTPIAGILFVVEELLHDF
LANQLSNWAPTSPEHRRQLI AAGAGAGLAAAFNAPIAGVLFVVEELLQDV
LANQFSNWAPTSPEHRRQLI AAGAGAGLAAAFNAPIAGVLFVVEELLQDV
LANQLSNWVPTSPEHRRQLI AAGAGAGLAAAFNAPIAGVLFVVEELLQDV
LAAQLSHWLPTSPEHRRQLI AAGAGAGLAAAFNAPIAGVLFVVEELLQDV
LANQLSRWVPTS PAYRRQLVAAGAGAGLAAAFDAP IAGVIFVVEQLLQNV
LAAWIGRWVTPSPNYSRQI IACGAAAGLAAGFNAPIAGVLFVVEDLLHDI
LAAWVSRWVTPSPNYRQQLI ACGAAAGLAAGFNAPIAGVLFVVEDLLHDI
LAGQFSRWLATSPAYQRQLI AAGAAAGLAAGFNAPIAGVLFVIEELLQDL
LAGQMSQWFPTS PDYRRQLI ASGAAAGLAAGFNAPLAGVMLVLEQLLHDV
* * * * *

FIGURE 5 (continued)

FIGURE 5 (continued)

FIGURE 5 (continued)

12/31

gi|218440518|ref|YP_002378847.
 gi|196258397|ref|ZP_03156931.1
 gi|126659209|ref|ZP_01730347.1
 gi|172034995|ref|YP_001801496.
 gi|67923321|ref|ZP_00516804.1|
 gi|218248189|ref|YP_002373560.
 gi|159026894|emb|CA089145.1|_e
 gi|166366612|ref|YP_001658885.
 gi|16331172|ref|NP_441900.1|_c
 gi|170077188|ref|YP_001733826.
 gi|158334130|ref|YP_001515302.
 gi|119510301|ref|ZP_01629437.1
 gi|186682049|ref|YP_001865245.
 gi|75908381|ref|YP_322677.1|_C
 gi|17232383|ref|NP_488931.1|_h
 gi|225516217|ref|ZP_03763190.1
 gi|254413597|ref|ZP_05027367.1
 gi|186683293|ref|YP_001866489.
 gi|225522175|ref|ZP_03768989.1
 gi|186682226|ref|YP_001865422.
 gi|220909579|ref|YP_002484890.
 gi|37523751|ref|NP_927128.1|_c
 gi|119486852|ref|ZP_01620827.1
 gi|209525724|ref|ZP_03274261.1
 gi|220908306|ref|YP_002483617.
 gi|81300397|ref|YP_400605.1|_C

TAVVRAPVTAIVIVFELRANFNIVLPLMITCAVSYLVAENVFSRSIYEHL
 TAVVRAPVTAIVIVFELKANFNIVLPLMLTCAVSYLVAENVFSRSIYEHL
 TAVVRVPVTAIVFVFELNTDFNIVLPLMVTCAVSYIVAESVSRGSLYEHL
 TAVVRVPVTAIVFVFELNTDFNIVLPLMVTCAVSYIVAESVSRGSLYEHL
 TAVVRVPVTAIVFVFELNTNFNIVLPLMVTCAVSYIVAESVSKGSLYEHL
 SAVVGVPVTAIVIVFELNSNFNIVLPLMIVSSVAYVADSFVRGSLYERM
 TGVVRVPVTAIVIVFELHRNFNIVLPLMLTCAVSYITAESIIRPGSLYEHL
 TGVVRVPVTAIVIVFELHRNFNIVLPLMLTCAVSYITAESIHPGSLYQHL
 TSVVRVPVTAIIIVFELTGNFNVVLPMLVACATSYLVAESLFPRSLYDHL
 TGVARVPVTAIVIVFELTDFNVLVPLMLASVTALIVAESIFKGSIIYEYL
 GAVSRVPVTGIVIIIFEITQDFNLVLPMLISSVVAYLTAEQINQSSIIDLL
 SAVSKVPITAIVIVFEMTDFNVLVPLMIVSVAAYLVSDKVMPGSLYDKL
 SAVSKVPITAIVIVFEMTDFNVLVPLMIVSVAAYLVADKVVPGSLYEKL
 SAVSKVPITAIVIIIFEMTDFNVLVPLMIVSVTAYLVADKVVPGSLYEKL
 SAVSKVPITAIVIVFEMTDFNVLVPLMIVSVTAYLVADKVVPGSLYEKL
 SAVSKVPITAIVIVFEMTDFNVLVPLMIVSVISYLVADQLVPGSLYDKL
 SAVSKVPITAIVIIIFEITADFNVLVPLMITCVVSYLIADQLSKGSLYQRL
 SAVSKVPITAVVIVFEMTDFNVLVPLMIASVVAYLVAEKIDHRSLYDLL
 SAVSKVPITAVVIVFEMTDFNVLVPLMIASVVAYLVAEKIDHRSLYDLL
 SAVSKVPITAVVIVFEMTDFNVLVPLMIVSVIAYLVAEKIDPRSLYDLL
 AATAKAPVTAIVIVFELTHDFNVLVPLMIASVIAYLVSERLFPFSFYDRI
 CAVTRVPITAVIIIFEITNDFNVLVPLMVVAVASLVAERLNEGSIIDQL
 CAVSRTPITAVVIVFEITRDFNVLVQLMICSVVAYFVADKIDKHSLYDRL
 CAVSRTPVITAVVIVFEITNDFNVLVPLMICSVVSYLVAEKIEKDSLYDRL
 SAVTRGPITAIVIVFELTMDFDVVLPLMIGSVVAYLVAEKVSKGSIIYTHL
 SAVTQGPITAIVIVFEMTRDFNAVLPMLVASITAYGIASLARSPIKAAAV
 .. *:*****: *: ** *: . : . * :

gi|218440518|ref|YP_002378847.
 gi|196258397|ref|ZP_03156931.1
 gi|126659209|ref|ZP_01730347.1
 gi|172034995|ref|YP_001801496.
 gi|67923321|ref|ZP_00516804.1|
 gi|218248189|ref|YP_002373560.
 gi|159026894|emb|CA089145.1|_e
 gi|166366612|ref|YP_001658885.
 gi|16331172|ref|NP_441900.1|_c
 gi|170077188|ref|YP_001733826.
 gi|158334130|ref|YP_001515302.
 gi|119510301|ref|ZP_01629437.1
 gi|186682049|ref|YP_001865245.
 gi|75908381|ref|YP_322677.1|_C
 gi|17232383|ref|NP_488931.1|_h
 gi|225516217|ref|ZP_03763190.1
 gi|254413597|ref|ZP_05027367.1
 gi|186683293|ref|YP_001866489.
 gi|225522175|ref|ZP_03768989.1
 gi|186682226|ref|YP_001865422.
 gi|220909579|ref|YP_002484890.
 gi|37523751|ref|NP_927128.1|_c
 gi|119486852|ref|ZP_01620827.1
 gi|209525724|ref|ZP_03274261.1
 gi|220908306|ref|YP_002483617.
 gi|81300397|ref|YP_400605.1|_C

LDASGIHLT-EEIPVNDFLSKLKASDVMQSQVETLDSYLSLEAVLQAMSI
 LDASGIHLS-EEVPVNDFLSKLKAADVMQSQVESLESHTLTDKVLQAMSI
 LHARGIYIN-EAATGQDFLSKLTASQVMESQVETLSSDILTDEVLQAMSN
 LHARGIYIN-EAATGQDFLSKLTASQVMESHVETLSSDILTDEVLQAMSN
 LKARGIYIN-EGATGQDFLSKLTASEVMQSQVETLSSDFTLDDVLQAMSA
 LKARGIYLD-NNPVVQDFLSQLTAADVMQSQVETLPSDILTLDQVIQAMSL
 LSASGIILN-EETPANDVLAHLSAMDVMQSQVEILAADLPGEVVKIMSR
 LSASGIILN-EETPSNDVLAHLSAMDVMQSQVEILPADLPGEVVKIMSR
 LETKGIFLA-EEKPDHDFLADIRAGQVMKTEVESLEQSLTAAQVLPIMSN
 LEASGIHLE-EEKNPQ-VLTDLTAAQVMQPEVETLESHLNKDLVPILSE
 LQQQGIQLQNASPSAQRMLDALTAEDIMQRQVETLPSDLSLDAARKIFSR
 LQLKGITITKAVSPEG-ILTKLTANDVMQHRVETLDAADMTLEEAMQVFSR
 LELKGITITLKQVPMEG-ALTKLTAKDVMQERVETLDAEMSLEEAMQSFAR
 LLLNGITLTKQMSVEG-ILSQMTAKDVMQQRVETLDAADMTLEEAKQAFAS
 LLLNGITLTKQMSVEG-ILSQMTAKDVMQQRVETLDAEITLEEAKQAFAS
 LLLN--GIKKDATREG-VLTQLTAQDVMQERVETLETQMSVDEVIQAFSR
 LEWGYTSPTQKVDKG-SLVGLTAEDLMQRQVETLSLHMTLDDALQMVS
 LEWKGIHITKEPST-EGLLAQLSALDVMQRRVETLSSQMSTDEAVQAFSH
 LEWKGIHITKEPSR-EVLLAQLSAVDVMQRRLETLSQMSTDEAVQAFSH
 LEWKGIHITKEPGT-EGILAQISAADVMQRRVETLSSQMSTDEAVQAFSD
 LELNGIYLNELTTV-NTSLADMTAAQVMQRQVETLRSMDMPLEEAVQAFSR
 LAWNGIRLGEEAVSDEKLLAQLAAGDVMQTRLETLESTLPLSEVSQAFSR
 LKLNIGIELKEDQ-LDREALSKLLARDVMQRQVETLSSQLPLLQVRQEF
 LALSGIELEPDQNAADRVALNMLHARDVMQRQVETLEDQLSLEQVRSFASQ
 LELRGIQLDQKFAGDQ-QLQGLLAADIMQRRVETLPADMTIAAAIQFFEQ
 VDALTPLNS-----SLGLTAAQVMASPVETLEASLPLTEVIQQFNR
 : : * ::* : * * :

FIGURE 5 (continued)

FIGURE 5 (continued)

14/31

gi|218440518|ref|YP_002378847.
 gi|196258397|ref|ZP_03156931.1
 gi|126659209|ref|ZP_01730347.1
 gi|172034995|ref|YP_001801496.
 gi|67923321|ref|ZP_00516804.1|
 gi|218248189|ref|YP_002373560.
 gi|159026894|emb|CA089145.1|_e
 gi|166366612|ref|YP_001658885.
 gi|16331172|ref|NP_441900.1|_c
 gi|170077188|ref|YP_001733826.
 gi|158334130|ref|YP_001515302.
 gi|119510301|ref|ZP_01629437.1
 gi|186682049|ref|YP_001865245.
 gi|75908381|ref|YP_322677.1|_C
 gi|17232383|ref|NP_488931.1|_h
 gi|225516217|ref|ZP_03763190.1
 gi|254413597|ref|ZP_05027367.1
 gi|186683293|ref|YP_001866489.
 gi|225522175|ref|ZP_03768989.1
 gi|186682226|ref|YP_001865422.
 gi|220909579|ref|YP_002484890.
 gi|37523751|ref|NP_927128.1|_c
 gi|119486852|ref|ZP_01620827.1
 gi|209525724|ref|ZP_03274261.1
 gi|220908306|ref|YP_002483617.
 gi|81300397|ref|YP_400605.1|_C

gi|218440518|ref|YP_002378847.
 gi|196258397|ref|ZP_03156931.1
 gi|126659209|ref|ZP_01730347.1
 gi|172034995|ref|YP_001801496.
 gi|67923321|ref|ZP_00516804.1|
 gi|218248189|ref|YP_002373560.
 gi|159026894|emb|CA089145.1|_e
 gi|166366612|ref|YP_001658885.
 gi|16331172|ref|NP_441900.1|_c
 gi|170077188|ref|YP_001733826.
 gi|158334130|ref|YP_001515302.
 gi|119510301|ref|ZP_01629437.1
 gi|186682049|ref|YP_001865245.
 gi|75908381|ref|YP_322677.1|_C
 gi|17232383|ref|NP_488931.1|_h
 gi|225516217|ref|ZP_03763190.1
 gi|254413597|ref|ZP_05027367.1
 gi|186683293|ref|YP_001866489.
 gi|225522175|ref|ZP_03768989.1
 gi|186682226|ref|YP_001865422.
 gi|220909579|ref|YP_002484890.
 gi|37523751|ref|NP_927128.1|_c
 gi|119486852|ref|ZP_01620827.1
 gi|209525724|ref|ZP_03274261.1
 gi|220908306|ref|YP_002483617.
 gi|81300397|ref|YP_400605.1|_C

KQLGGG--TMQGSKPAPSYVYQTRSPALGNRILLPLSDGEMAGTLLQI
 KQLGG---GTQGTKPPPSYVVYQTRSPALGNRILLPLSDPEMAATLFQI
 NQLQGDDTTINEVVPDPSYIAYQTRAPLAGDGRILLPLVNPDTATSLFKI
 NQLQGDDTTINEVVPDPSYIAYQTRAPLAGEGRILLPLVNPDTATSLFKI
 NQLQGDDTTINEVVPDPSYIAYQTRAPLVGVGRILLPLTNINTATSLFKI
 NQLEGND SKLNEAIPESYIAYQTRAPATGGGRILLPLADPKTATALFQI
 DQLGGVCQLPQPT--TPSYVVYQTRSPAVGIGRILLPIANPDTATALFKI
 DQLGGVCQLPQPS--TPSYVVYQTRSPAVGIGRILLPIANPDTATALFKI
 SQLGG----QIIHPPHPAYSVYQTRSPVLGEGRILLPIRSDSALVLFQI
 QELLG--KSHHAVQRTPSYCVYQTHSDNYKRKLWVAIANPNTAEALITF
 DHISGQ--KQQDNQAFAPSYGVYLTRGPATGEGRILLPLANPDTAEPLLVK
 EHLNYG--NVTPESSQPEPSYVVYQTRSPNIGRILLVPVANPETAAILLQM
 DHLNCG--NGTPKLQPEPSYIVYQTRSPNIGRILLVPVANPETAAILLQM
 DRLNCE--NPNPGPQPEPSYVVYQTRSPSTGRGRILLVPVANPETAATLLKM
 DRLNCE--NPNPGPQPEPSYVVYQTRSPNTGRGRILLVPVANPETAATLLKM
 DHLNCE--NGVSGPQPEPSYVVYQTRSPNIGRILLVTVANPDTAETLLQM
 DQLDSK--TTQTRSQSEPSYVVYQTRSPETGRGRILLVPPIANPQTAPILVKL
 ERLSGN--SQQIERKSAPSYVVYQTRAPATGKGRILLVPLSHPQTAETLLEM
 ERLSGN--SEQMERKLAPSYVVYQSRAPATGKGRILLVPLSHPQTAETLLEM
 ERLNGN--TQQIESKSATSYVIYQTRAPATGKGRILLVPLSHPQTAETLLEM
 SELSGE--RQQLAPRVQPSYLVYQTRCTASGQGRILLPLANPATADRLLQM
 DLLSGD--TQAVGPRPDPSYVVYQTRAPALGGGRILLPLANPDTAPALLEF
 GILSGE--DTQVGLHPEPSYVVYQTRAPQVGGGRMLVSLHNPKTAPSLLLEI
 GYISG--SHQVGHHATPSYVVYQTRGPMVGHGRILLVALNNPPTAPALLHI
 DRLSGE--ASPLKVKPAPSYVYQTSSTTAQGRILLVPLSNPQTTDALLRL
 ELLSGQ--DIAATPIAPSYVAYETRSPATGQGRILLVLLANPRTAQSLLQL
 : .: * * .: . .: : : . : * . .

AAAIARHRNYEIECLQVIKVPRHQFPSEARIDIHHSRKLQMQRMERLARHL
 AATIAHRHRHYEIECLQVIKPRHQFPSEARVEVHHSRKLQMQRMERLGRHL
 SSAIAQQNQEIIECLQVITVPKHSDPAQTPVITQHSRQLLQRTERLGRHY
 SAAIAQQNQEIIECLQVITVPKHSDPAQTPVITQHSRQLLQRTERLGRQY
 SAAIAEQNQEIIECLQVIKPKHSDPAQTTPVITQQSRQLLQRAERLGRQH
 GAAIARQQNDELECLQVIKVPKHKKPAQTPVKSSQHSRHLHRMERLGRRQ
 AAAVARERNYEIDCLYVITVPRLSPPAEVKVDTREGRKLLHRLERLARQQ
 AAAIASERNYEIDCLYVITVPRLSPPAEVKVDTREGRKLLHRLERLARQQ
 AAAIAKKQNYEIDCLQVVKVSKAQSPSVQRVQWQRQRLMQKLERIARHQ
 AAAIAKEKNAMVYCLNVVQVAATYPLTDAHMNTYPERHLMKLERLGETL
 AVAMAQALNYEIECLHIIIRLPNRNLPATPVQTVKGRLLQKAIRYGRH
 AAAIARDRHYEIECVQIILVSRHNSPSETSVRTAKSRLLRQAEVLAKKW
 AAAIARDRHYEIDCVQVMLIPRHSSPSETQVRTAKSRLLRQAEVLAKKW
 AAAIARDRHYEIECVQVMLVSRHSSPSETTVRTAKSRLLRQAEVLAKKW
 AAAIARDRHYEIECVQVMLVSRHSSPSETTVRTAKSRLLRQAEVLAKKW
 AATIAARDRHYEIECLQIILSRHSSPSETQVNTTKSRLLRQAKALAKKW
 AAAIARDRHYEIECLQVILVSRSTPAETPVTTTKSRLLRQAEKWRD
 AVAIAKDRNYEIECLQVIIVPSGRIPSETPVQISKSLQLLQRAILLGENS
 AVAIAKDRNYEIESLQVIIVPSSRIPSETPVQISKSLQLLQRAILLGENS
 AGAIAKANNYEIECLQVIIVPSNRIPSETPVQITKSLELLQKAILLGEKW
 GLAIAREFQLEIECLQVIIVPRHSDPSTTPVTRTTLNRRLLQRMERLARPD
 ALAIARARNLEIECLHVIAPRTSAPSSTPVAVAPAVNLVRQAVRAGRAV
 AAAIARERNYEIECLNVITIPRHRSPSETPVRLTKSRLLMRQAEQMGRAW
 AAAIAAERNYEIECLTIITARNLSPAETPVSTIKCRLLRQAEERIAKSW
 AAAIAQQHHYDLECLHIIIPRHRSPSETSVDTITASRQMEHAVAGGQTW
 AAGLASLRNYEIECLHLIPVSREADPSQTAVSTLGARRMLRQAEERLGRW
 . : * . : . : : . : : .: : : ..

FIGURE 5 (continued)

15/31

gi|218440518|ref|YP_002378847.
 gi|196258397|ref|ZP_03156931.1
 gi|126659209|ref|ZP_01730347.1
 gi|172034995|ref|YP_001801496.
 gi|67923321|ref|ZP_00516804.1|
 gi|218248189|ref|YP_002373560.
 gi|159026894|emb|CA089145.1|_e
 gi|166366612|ref|YP_001658885.
 gi|16331172|ref|NP_441900.1|_c
 gi|170077188|ref|YP_001733826.
 gi|158334130|ref|YP_001515302.
 gi|119510301|ref|ZP_01629437.1
 gi|186682049|ref|YP_001865245.
 gi|75908381|ref|YP_322677.1|_C
 gi|17232383|ref|NP_488931.1|_h
 gi|225516217|ref|ZP_03763190.1
 gi|254413597|ref|ZP_05027367.1
 gi|186683293|ref|YP_001866489.
 gi|225522175|ref|ZP_03768989.1
 gi|186682226|ref|YP_001865422.
 gi|220909579|ref|YP_002484890.
 gi|37523751|ref|NP_927128.1|_c
 gi|119486852|ref|ZP_01620827.1
 gi|209525724|ref|ZP_03274261.1
 gi|220908306|ref|YP_002483617.
 gi|81300397|ref|YP_400605.1|_C

NVPVHTQIRIAQDQAQSIETIQRHINLMMSWKGGGDTQGAIFGNVTD
 NIAVHTQVRVAQDQAESIIETIQRHISLMMSWKGISDSQGAIFGSVAD
 HIPVHTQIRVAHDRAQAAILDTIRERHINLMVLEWTDHHTPGTIFDVID
 HLPVHTQIRIAHDRAQAAILDTIRERHIDLMVLQWTDHNSPGTIFDVID
 QISVHTQIRVAHDRAQAAILDTIRERHINLIVLEWNPDKKTPGSVFGPVID
 KLPVHTQVRVAHDIAQGILDTIRERHINLMMEWNGETSTPGAIFGHVVD
 NISVHTQISVAQDIAEGILATIRERHSNLLIMGWTGERSTTGAIFGFLVD
 HISVHTQISVAQDIADGILATIRERHSNLLIMGWTGEKSTTGAIFGFSVD
 RVLFTHEIKLAYSITDTILDTIQTRHSDLLILEWQGEMPIGGQIFGQITD
 GISIHTQVILAYDVAGAILQEISSKDQ-GLILGWQGDHRHPPGQLFGSVME
 QVSVHTQIIAAQDVAQAILEVEQTEHINLLMGWQGGQFFSGGQIGNPTVR
 KIPLHTQIRVAHDPAEAIETIHERHINLILMGWKGETSTPGRIFGNVVD
 KIPLHTQIRVTHNVAGAILETINEQHIDLIILMGWKNTSTPGRVFSRVVD
 RIPLHTQIRVAHDPAHAILETIKDRHIDLIILMGWKGSTSTPGRIFGNVVD
 RIPLHTQIRVAHDPAHAILETIKDRHIDLIILMGWKGSTSTPGRIFGNVVD
 HIPIHTQIRVAHDIAQAILETINERHIDLIILMGWKNTSTPGRIFGNVVD
 NIPVHTQIRVSQDITQGILDTIKERHINLMMMGWKGSTSTPGRIFGGVVD
 RIPVHTQIRVAHNVAGAIETVKERHIDLVLMGWKGSTSTPGRVFSRVVD
 RIPVHTQIRVAHNVAGAIETVKERHIDLVLMGWKGSTSTPGRVFSRVVD
 RIPVHTQIRVTHNVAGAIETVKDRHIDLVLMGWKGTSTPGRVFSRVVD
 SVPLHTQIRVAQDVAGAILETISDRHIDVLMGWEGESLTPGSIFGSVVD
 GVPVHTQVRVAHDIAQAVLETVKERRIGLVLMGWRGVNTSPERIFSNTVD
 KIPLHTQVRVTNDVAQAILETTKERHIDLIILMGWQKGSSTDRVFGNVVD
 RVPIHTQIRVSHDVSGAIETVQERNIDLTILGWQGERSATNRVFGTVVD
 QVPVHTQIRVAQDIAMAILETIRDRHIDLVLMGWEGVTLTPGRIFGTVD
 QIPVHTQVRVCHDLAAILETISDRHINDLVLMGWGGGASLAQRLFREGID
 : .**:: . : :: . . : * :

gi|218440518|ref|YP_002378847.
 gi|196258397|ref|ZP_03156931.1
 gi|126659209|ref|ZP_01730347.1
 gi|172034995|ref|YP_001801496.
 gi|67923321|ref|ZP_00516804.1|
 gi|218248189|ref|YP_002373560.
 gi|159026894|emb|CA089145.1|_e
 gi|166366612|ref|YP_001658885.
 gi|16331172|ref|NP_441900.1|_c
 gi|170077188|ref|YP_001733826.
 gi|158334130|ref|YP_001515302.
 gi|119510301|ref|ZP_01629437.1
 gi|186682049|ref|YP_001865245.
 gi|75908381|ref|YP_322677.1|_C
 gi|17232383|ref|NP_488931.1|_h
 gi|225516217|ref|ZP_03763190.1
 gi|254413597|ref|ZP_05027367.1
 gi|186683293|ref|YP_001866489.
 gi|225522175|ref|ZP_03768989.1
 gi|186682226|ref|YP_001865422.
 gi|220909579|ref|YP_002484890.
 gi|37523751|ref|NP_927128.1|_c
 gi|119486852|ref|ZP_01620827.1
 gi|209525724|ref|ZP_03274261.1
 gi|220908306|ref|YP_002483617.
 gi|81300397|ref|YP_400605.1|_C

TLIHKAPCDMLVLKLGASPTAYPHN-----LEHNATWLVP
 TLIHKAPCDMLVLKLGASANAYPLN-----LEHNATWLVP
 ILIRKATCELILVLKLGQKEAYPQI-----LDKDTTWLIP
 VLIRKATCELVLVLKLGQKEAYPQT-----LDKDATWLLP
 ILIRKATCELVIVVLKLGQKEAYPQT-----LDKDTTWLIP
 LLIRKAPCELVLVLKLGNGREPSYDPS-----LNRDATWLIP
 TLIAQAPCETILVLKLG-TKDCFPND-----PNRERIWLIP
 TLIAEAPCETILVLKLG-TKDCFPND-----LNRERIWLIP
 RLIDQAPCSLLMVKQGNTNDHAYPRY-----LSPMAHWLMP
 YLVKKVPCDVILVKTSHAHPLQFKA-----QTGKIRALVP
 TILQQAQSVLVLIRPATHEFTG-----KNQRWLIP
 TIIRQATCDVMLVLKLGNTPHSLLP--IPHS-----PFPTPKFNRWLVT
 NIIRQATCEVVLVLKLGKTQHKPNHSYPQTA-----LQTQHSFNRWLVP
 TIIRQATCDVVLVLKLGTS-----PIPNPPFNRWLVP
 TIIRQATCDVVLVLKLGSSPFPIPN---TQY-----PISNPQFNRWLVP
 TIIRQAACDVVLVLKFGNIS-----DSNHFNRLVP
 AVIRQAPCDVMVLKLGEMMHPPYIP-----PFFQRWLIP
 TIIRQAGCDVILAKLDDKRS-----FDRWLLP
 TIIRQAGCDVILAKLDDKRS-----FDRWLLP
 TIIRQAGCDVILAKLDDKRA-----FDRWLLP
 TIVHQAECDVVLVLKLAQENVR-----FDRWLVP
 TIIRQAPCEVLLVLKLG-GDAI-----PRRWLVP
 MIIRACECEMVLVWKPMVDPPFGSKRK-LR-----LHSLLGWQRWLVP
 TIIRQVPGEVVLVLKVGKNVNIIFSDPSDQR-----SLQFPVWRWLVP
 TIVREASCNVVLVVRPGKQLILPLAKAADLSGQSALTLLMRLFTLNRWLVP
 QILQQAPCRVLLVLKPGAFLAKAPLLQGDR-----PTLPLRHWLIP
 :: :: : *

FIGURE 5 (continued)

16/31

gi|218440518|ref|YP_002378847.
 gi|196258397|ref|ZP_03156931.1
 gi|126659209|ref|ZP_01730347.1
 gi|172034995|ref|YP_001801496.
 gi|67923321|ref|ZP_00516804.1|
 gi|218248189|ref|YP_002373560.
 gi|159026894|emb|CA089145.1|_e
 gi|166366612|ref|YP_001658885.
 gi|16331172|ref|NP_441900.1|_c
 gi|170077188|ref|YP_001733826.
 gi|158334130|ref|YP_001515302.
 gi|119510301|ref|ZP_01629437.1
 gi|186682049|ref|YP_001865245.
 gi|75908381|ref|YP_322677.1|_C
 gi|17232383|ref|NP_488931.1|_h
 gi|225516217|ref|ZP_03763190.1
 gi|254413597|ref|ZP_05027367.1
 gi|186683293|ref|YP_001866489.
 gi|225522175|ref|ZP_03768989.1
 gi|186682226|ref|YP_001865422.
 gi|220909579|ref|YP_002484890.
 gi|37523751|ref|NP_927128.1|_c
 gi|119486852|ref|ZP_01620827.1
 gi|209525724|ref|ZP_03274261.1
 gi|220908306|ref|YP_002483617.
 gi|81300397|ref|YP_400605.1|_C

MAGGPNAQRAIELLPSLTTLTYANPN----APELWLCKVYPP--TELEPNS
 MAGGPNAQRAIELLPSLTRLYSNPD----APQLWLCKVYPP--SEIEPDS
 IAGGPNAKRAMELLPGLTLYTRPR----SPIIWLCQVFPP--HKSTPNY
 IAGGPNAKRAMELLPGLIKLYTRPR----SPIIWLCQVFSP--HKATPNY
 IAGGPNAKRAMELLPGLTQLYTRPR----SPIIWLCQVFSP--HKPTPDY
 IAGGPNAQRALELLPGLTGLYDRPR----SPIIWLCQVFPP--AGALPDY
 TAGGPNAQRALELLPSLLSLSESSD----YPKLWLCKIYSP--TDLLPDL
 TAGGPNAQRALELLPSLLSLSESSD----HPELWLCKIYSP--TELLPDL
 VAGGPNIKMLNLLPALFSLYPEEN----NPQLLISKVYLPQRSQRYDPF
 IAGGPNAERAQEFIPALLHLSPEVVRYSLPSQICLCQVYQP--QQRQHF
 VSGGPNVQQAIIQLLPALVPIISKSPK-----LKLQVYFP--SSDLPDQ
 MAGGPNVRAIKLLPALATLGNDP-----YIRLTQVFKP--SQLKPDMT
 MAGGPNSPLAIKLLPALITLGNDP-----QIRLTQVFN--SELKPDMS
 MAGGPNARIAIKLLPALVTLSDDP-----QIRLTRVFKP--WEFRPDMT
 MAGGPNARIAIKLLPALVTLSDDP-----QIRLTRVFKP--WEFRPDMT
 MAGGPNVTKAVKLLPALVTLGNNT-----QIHLTQVFKP--SELKPDIT
 IAGGPNSQRALEFLPALIPLSSKPP-----DIRLCQVFP--EDTIMDTT
 MAGGPNSSQAIKLLPALSSSLSTS-----PQIKLCQVFP--TNSILD
 IAGGPNSSQAIKLLPALSSSLSTS-----SQINLCQVFP--NNSIPD
 MAGGPNSSQAIKLLPALASLSKS-----PQINLCQIFQP--TKSLPD
 IAGGPNVQRAMQLLPGLTRLSPA-----PQVRLCQVFP--EATLLD
 IGGGPNAQEAIIKILPALLRGPGK-----SEIACVQVFP--DNPAHD
 IRDDAKKSVAVQLLPALTHLSHQ-----AEIRLLKVMNK--EVSUSE
 LRTGGPPPAAMGILPGLARLSLR-----PDIRLLTVVKN--ALSEAE
 VAGGPNAQYALTLLPALVALSQHP-----QIRLCQVFP--SEPYHD
 LSAQVPQTGLAEPLEQILQLITDPD-----ISLCQIDL--RQTVS

: : : : :

gi|218440518|ref|YP_002378847.
 gi|196258397|ref|ZP_03156931.1
 gi|126659209|ref|ZP_01730347.1
 gi|172034995|ref|YP_001801496.
 gi|67923321|ref|ZP_00516804.1|
 gi|218248189|ref|YP_002373560.
 gi|159026894|emb|CA089145.1|_e
 gi|166366612|ref|YP_001658885.
 gi|16331172|ref|NP_441900.1|_c
 gi|170077188|ref|YP_001733826.
 gi|158334130|ref|YP_001515302.
 gi|119510301|ref|ZP_01629437.1
 gi|186682049|ref|YP_001865245.
 gi|75908381|ref|YP_322677.1|_C
 gi|17232383|ref|NP_488931.1|_h
 gi|225516217|ref|ZP_03763190.1
 gi|254413597|ref|ZP_05027367.1
 gi|186683293|ref|YP_001866489.
 gi|225522175|ref|ZP_03768989.1
 gi|186682226|ref|YP_001865422.
 gi|220909579|ref|YP_002484890.
 gi|37523751|ref|NP_927128.1|_c
 gi|119486852|ref|ZP_01620827.1
 gi|209525724|ref|ZP_03274261.1
 gi|220908306|ref|YP_002483617.
 gi|81300397|ref|YP_400605.1|_C

QALEEAAQTLTEKLNKSIIP-ISIRSHSVADALIHQAQAEVCDLIILGAS
 QGLEEAAQILRDKNKPIFP-ISIRSNVSVDALIQARAEECDVIVLGAS
 QALEIEAKRLKEELERPVP-LPVRQSVDALIHQAQAEVCDVIVLGAS
 QSLEVAAKRLKEELERPVP-LPVRQSVDALIHQAQAEVCDVIVLGAS
 QSLEAKAQQKKEQLERPVP-LPIRSQSVTDALIVHLAESESCDVIVLGAS
 QVLEQTAQAVKDAVARPVP-LPIRSQSVDAVHLLAAETCDVIVLGAS
 SILEVLQASLQKAIAPVVP-LPIRSASPAEAINLVESEDCSLVLLGAS
 STLEVLQASLQKAIAPVVP-LPIRSASPAEAVINLVESEDCSLVLLGAS
 YDLKNLAERWSEQLQRPPI-IPVCSTSIADALSDLAEMRECAIVLGAS
 PLLVKAKRGLDPLVEIPI-PIRSSHIADSLIRLIKEETYDLVILGAS
 SILEHYSQFLKQACPDASVKRTQICQNVADAIVELARLQKTDIIVIGAS
 VLEQAIRILMRRRLSSNVVAIPVQADSVAEGVINLVKTEGYDVVVLGAS
 ISEQAIRQLMRRRLSSNVVAIPVQADSVAEGVINLVKTEGYDVVVLGAS
 VLEQAIRQLMRRRLSSNVVAIPVQADSVAEGVINLVKTEGYDVVVLGAS
 VLEQAIRQLMRRRLSSNVVAIPVQADSVAEGVINLVKTEGYDVVVLGAS
 VLEESTHQLMRHRLQSTVVAAPVQANSVCEGVINLVQTERYDVVVLGAS
 AIEEASDFLKKQTEG--RIRRIPIRASEVSDAVVYIANNLCDVIVLGAS
 TTLLDKSVHFLQRRVSGKVATPVRANSVSDAVLKCAELNSDVIVLGAS
 TTLLDKSVNLFQCRVSGKVATPVRANSVSDAVLQCAEQDKSDVIVLGAS
 TTLLDKSVHFLQRRVSGKVATPVRANSVSEAVLECAQDQNSDVIVLGAS
 TQALEQGMALLKPHLNGALSSLSLSKSI AEAI IELAHAEQDDVILVGAS
 TGLLDADAESLGRSLGYPVRALPLFSKSAQAIVQCTSEGHFDGVIIGAS
 KRAWEHTSQELSSLLNSKVRITAVTSEFVPDAVIDFAYQEHCDVIVLGAS
 MAKLDHLVEELKSPINGEVVATAVCARSVVDVLLDIANHDYCDVIVLGAS
 TTLLQDADFLNQQLQAPVITLTLCADSVADAIVDLVEKDQCDVIVLGAS
 ANRLQQIQRRLNRLRQPIQADLICARSVTAVLLDVMQSRGIEAVMLSMT

. : : : : :

FIGURE 5 (continued)

FIGURE 5 (continued)

18/31




FIGURE 6

19/31




FIGURE 7

20/31

MSEVYWQAKIWGLLHDPALKALHNNSSGRGGEGAWQSLGCMQNWVSPKSSTEEKDGE LSN
---1---

SWLDHIGLSDLIASASDRGAIHYIGTPIDYDGKGLELSHLLSGAKLPLKLANHSEIIGQ

GNRRTYLEQKELELIQQMPAEFLHGTDKAQECFWWLWRCLPEAVAEQFGPESLLMPAET
---2

RLPDASIWSHGSM TAALAGALAGYDTNIEDIAKGARNTPKSQAYLAVFTFSPIQELIKA
2--- ----3----

SRKIKDFWAGSWILHYLSAKVCWELAQQYGPDCFLYPSLYGQPLIDHWLLEKYGEQQGS
-----4----- --5--

--11-----
QWVPQPGDRQLLTAGFPNVIMLVLPKEKVSAA MQSAKNHLLNAWRNIAHLVFAELQQRH
--6--

---12---
WQKQLNPTDPTWKHWLDAQWQTYWSAMAIGAEGKTLKSVG IPTQTPE SKAQ RDQWVREQ

NEAFSASLFAQEELDFIEKSTDLFIQQRNRNYAGSLNVGSWWADIFGQTRFTLSAVKNA

RNWKIPTAFGPRSTISGLGPVVHPQAPSHQRDWWTEGDTQKYWQRQAGLFDGSEQLNAT
-----13--

ETVKRGLEKVL PALLGRGAKELKTYYPDLTVGIAGYLKTQPDGLPIFEQACSAISQKIL

GDRHKVADSVTQDWGIPWVEENMEKKYHPRL LNSGWLVEELEDLDQTELPSYRQALQTE

IEKFYPQNNPTSWYVLAAGDGDGMSEWLKGTKMRTYGDYFPQVLTVPEDLQPTFQPFSE
---7--

QPKRMGPATHNALSRALLDFSNQLVPYLTEQRYAGR LIYSGGDDVLAYTNLWEWDQWLW
-----8----- -----9-----

-----14----- ---15--
DIRQA FRGDRDEHREFDSTGHYWH SKTAHKNLPQRPLFTMGDGATISFGITIAHHSVPL
-----10-----

-----16-----
AIALEHLWEAEEEEAKEHEYGEGEDKKSKDAVQVRVIYGN GNVL TATSKFEVFKTWKDLL
---16-----

DIETIDASTYETAATVLEQHPIPVREAIMPWVNV LVERRDALDKDQQSTLRSRLACFLI

QLWQTTSQKNWEKEAKNWLKVAAFMKRNR YIKFPN

FIGURE 8

21/31

CLUSTAL W (1.81) multiple sequence alignment

```

gi|121997427|ref|YP_001002214.-----MTEATLMDIDLARNKLLARLHD
gi|88810314|ref|ZP_01125571.1|-----MTDRLWQAKLHARLHD
gi|255254649|ref|ZP_05334193.1|-----MSKKEELFENKLKAYFHD
gi|255338062|ref|ZP_05378933.1|-----MSKKEELFENKLKAYFHD
gi|20808984|ref|NP_624155.1|_h-----MEKVWLLKLKAFLLHD
gi|254478158|ref|ZP_05091540.1|-----MEKVWLLKLKAFLLHD
gi|38505758|ref|NP_942378.1|_h-----MSEVYWQAKIWGLLHD
gi|170079601|ref|YP_001736234.-----MAGKAYWQAKIWGLLHD
gi|126661502|ref|ZP_01732553.1|-----
gi|172035267|ref|YP_001801768.-----MSHLYHPYWQAKIWALLHD
gi|86608324|ref|YP_477086.1|_C-----MSYYWQAKLWGLLHD
gi|86607166|ref|YP_475929.1|_C-----
gi|255256456|ref|ZP_05335917.1|-----
gi|209526382|ref|ZP_03274910.1|-----MDNFLGSLNHS
gi|162453956|ref|YP_001616323.-----MSERDRDRDRNLAVAHVPVHGRTGDDEIEHHGTEADTRFWQQLQLLHD

gi|121997427|ref|YP_001002214.-----PAEKALVLLRDPAGHEGGTVRTLIEEILG-----
gi|88810314|ref|ZP_01125571.1|-----PAEKALVLLRDPAGPEGGTSRVLHERLFP-----
gi|255254649|ref|ZP_05334193.1|-----SPDKPFILLTGENHEKRAKEISK-----
gi|255338062|ref|ZP_05378933.1|-----SPDKPFILLTGENHEKRAKEISK-----
gi|20808984|ref|NP_624155.1|_h-----PPHKHWIISMDNEECIKKFQLQEKGRCHEILIKSKVI-----
gi|254478158|ref|ZP_05091540.1|-----PPHKHWIISMDNEECIKKFQLQEKGRCHEILIKSKVI-----
gi|38505758|ref|NP_942378.1|_h-----PALKALHNNSGRGEGAWQSLGCMQNWVSPKSSTEEK-----
gi|170079601|ref|YP_001736234.-----PALKALYGNPGRGEEGLWKKLACMEGWASPKSSEEKQ-----
gi|126661502|ref|ZP_01732553.1|-----
gi|172035267|ref|YP_001801768.-----PALKPLSDRYGFAREGQWQLLRMCQGCKSPKDNKKVLG-----
gi|86608324|ref|YP_477086.1|_C-----PALKALHDRTGRSKEGLWTRLAAMQGWSSPKA-----
gi|86607166|ref|YP_475929.1|_C-----
gi|255256456|ref|ZP_05335917.1|-----
gi|209526382|ref|ZP_03274910.1|-----SLCNIVFLRKLKALVG-----
gi|162453956|ref|YP_001616323.-----PPGKAFFLRQGAGGHKAVAADLFQATAGVPLKYVRPGPDWAASGADRPVS

gi|121997427|ref|YP_001002214.-----EVPQ-AAWAPVRKADRWASAADRPQFPRERDGRFARWSQVRFAEQP
gi|88810314|ref|ZP_01125571.1|-----QGMAGDLRATVRKADWWASAADRPQFPRDGKEGYPYARWSQVNFAEQP
gi|255254649|ref|ZP_05334193.1|-----IDIPYHKTIASDVLASFMEYYIPKDASKNKNLQVVFEEQP
gi|255338062|ref|ZP_05378933.1|-----IDIPYHKTIASDVLASSMEYYIPKDASKNKNLQVVFEEQP
gi|20808984|ref|NP_624155.1|_h-----EPLLGEDLTETEEEEKLIEKADTQAYPVNRIPLPPVAVKIRGEDVIFDDMF
gi|254478158|ref|ZP_05091540.1|-----EPLFGEDLTETEEEEKLIKADTQAYPVNRIPLPPVAVKIRGEDVIFDDMF
gi|38505758|ref|NP_942378.1|_h-----DGELSNSWLDHIGLSDLIASASDRGAIHYIG----TPIDYDGK----G
gi|170079601|ref|YP_001736234.-----ASELSGTWLDHVGLCDLISSASDRGIIHYIG----TAVDYGAE----G
gi|126661502|ref|ZP_01732553.1|-----
gi|172035267|ref|YP_001801768.-----DCVLQGNWLNHVGLCDLIVASASDRSTIGRLDPQYSAVTYQQQT----G
gi|86608324|ref|YP_477086.1|_C-----EGSSPIGDADLIASASDRAAIGHLPVAIDYQVEDRGQGLGSG
gi|86607166|ref|YP_475929.1|_C-----MGDADLIVASASDRAAIGHLPVAIDYQVADCSQSLGPG
gi|255256456|ref|ZP_05335917.1|-----
gi|209526382|ref|ZP_03274910.1|-----DRHLCSLDCLKTSEGQAALFWWEQNQHLEAIASSSDRV
gi|162453956|ref|YP_001616323.-----SPPRPAHVSVDWVKNPILTHPLASGTAAIDLGRDLPASREQVRALAGDV

```

FIGURE 9

22/31

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

VLRHPLTGEKIDLGEHGKL--REIELDTLKAGSLEHFRHLIHRDEAGSV
 VPIHPLTGKAFDLRQQGGL--RETRIEAVKDQSQAHFKRLK--AAGGDD
 EFVHPLAANRYNNFVIEKE--VFKKAVDEAVDELSKLD-----FKD
 EFVHPLAANRYNNFVIEKE--VFKKAVDEAVDELSKLD-----FKD
 NKKIYYDILAETDYETENK--RFLELVEKLAKNYNLSR-----IKE
 NKKIYYDILAETDYETENK--RFLELVEKLAKNYNLSR-----IKE
 LELSHLLSGAKLPLKLANHSEIIGQGNRRTYLEQKELELIQMPAEFLHG
 LELSHLLSGEKLPLRLQNHAEILGQSSRKTYLEQTEADLIDQMPAEIRDD

 LQIRHLLSGEPQNLKVN-----NNWQENLITKRKKLEAKEAEFLDKISQ
 LEMSHLLSGRKHFWQLANE--QHQAALNAHSRADYLMEREENAI PDWVRQE
 LEMSHLLSGRKRFWQLPEE--EHRALRQASSRAEYLLAQEARVVPDWIWQE

 NLEPGSSTERQTKERI IKHPISAQQQTIQTPDNREIAFP-----QWVQQ
 GRTLRELLERRVSQEDELDDLPEEEERERFLSERAEQVESELAALPRWHD

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

SWRTLLALWRFGEPEPATSGDATGLGELWRYLPADTRIPDHTIWQHLDLSS
 PRMALTWRFGEPELDTADDTARLGELWRFPLADTRIPDHSIWDHLDLVS
 SYKKFVYLWRYLKELLKKHTPSEYRKYWDLAPADTRFPNHTIFEHLKLAS
 SYKKFVYLWRYLKELLKKHTPSEYRKYWDLAPADTRFPNHTIFEHLKLAS
 EEKKEK---AKILLFLWRFYQEIFHWMRIHPADTRAPNHSIYDHLVQTS
 EEKKEK---AKILLFLWRFYQEIFHWMRIHPADTRAPNHSIYDHLVQTS
 TDKAQECFWWLWRLCPQAVAEQFGPE--SLLMPAETRLPDASIWSHGSMATA
 PEAAQACFWWLWRLCPQAVAEFGAA--SFLMPAETRLPDGSIWSHASMTA
 -----CLPEATCQLFDDASLLMPAETRLPDSSHWSHASLTA
 WEDPKKVFALWRCYAEVLEKE--EALIHLLPAETRLPDGSLWSHVSMTS
 TD-PQKVFWFWRCFPQAICQEFGDESLLLLPAETRLPDGSLWSHASVTA
 QEDIQKVFWFWRCFPQALAREFGDESLLLLPAETRLPDGSLWSHVSMTS
 -----MMEEKIYESFAENSS
 EQNPQNVFNWLWRMMPELRRQQ--DSDALLDPQHYILPCPIHSYRATVS
 DARLQQACLQLWRRLPPEEPPPGVAEVVWRHQPADSRAPDHSIWDHLDLVS
 : .. :

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

ALTGAFAGDGQ-----GRCALLNVTGLGPVQELIAAARTTS
 AFAGAFVADAN-----GECALLNISIGPVQEFIAAARTTS
 AVNAYFYNELNL-----NNMTLFIFTIGPVQEFYIVQARKTQ
 AVNAYFYNELNL-----NNMTLFIFTIGPVQEFYIVQARKTQ
 TLVSALP-----KPAFLFLTIGPVQSFIATARKTS
 TLVSALP-----KPAFLFLTIGPVQSFIATARKTS
 ALAGALAGYDTNIEDIAK--GARNTPKSQAYLAVFTFSPQELIKASRKIK
 ALAGALAGYDTTELDDIPK--GGQNTPKSHPYLAVFTFSPQELIKASRKMK
 AMAGALSGYDLTTEDLTQNGEANKPLSHPYLASFTFSPQELIKASRKMR
 ALAGGLAGYYKQAEENYPIKNQKFNDYSRPLYVNFTFSPQELIKASRKMR
 ALAGCLAGYDE-----PKFARPLVSFTFNPIQELIKASRKMR
 ALAGCLAGYDE-----PKFTRPFLASLTFNPIQELIKASRKMR
 NVS-----LFLFTIGPVQQFISQARKTQ
 AIAGAVDFDERG-----IVTQYPYLLLTFTFSPQDFIKASRKFL
 SLSFLSGRRREG-----PVMPWLLAFSLGPVQRFIAQSRTST
 . : * : * * : *

FIGURE 9 (continued)

DLWAGSHLLARLAWAMRVVCERLGPDAVLFPPQLRGVPPQVDLWLLLEQGL-
DLWAGSHLLARLSWEAMRVVCERLGPEALFPPRLRGI PQVDLWLLLRDCG-
DLYWGSYILSYLTWVAIEKVEIAYGPDSI IFPELKEQPLCDFWIAKLFN-
DLYWGSYILSYLTWVAIEKVEIAYGPDSI IFPELKEQPLCDFWIAKLFN-
DLWAGSYMLSYFIWKIMKFVVERYGADVIVYPNLLGQPLVDLWLSSEVFG
DLWAGSYMLSYFIWKIMKFVVERYGADVIVYPNLLGQPLVDLWLSSEVFG
DFWAGSWILHYLSAKVCWELAHQQYGPDCFLYPSLYGQPLIDHWLLLEKYGE
DFWAGSWILHYLSAKVCWRLLAKYGGADCFLYPSLYGQPLIDHWLLQKHS-
DFWAGSWILHYLSAKVSWTLANIYGPDCFIYPNLYQQPLIDHWLLQKYSK
DFWAGSWILHYLSAKVSYAIANEYGPD TLLYPCLYQQPLIDNWLLLEEYS-
DFWAGSWLLHYLSARISWRLLAEKYGPDCLLYPSLYGQPLIDYWLREKYP-
DFWAGSWLLHYLSARLCWRLLAEKYGPDCLLYPSLYGQPLIDAWLLEKYP-
DLFMGSFLLSYLTFIGMEEVIDRYGPKSIIYPNLSAQPLMEWHLLKRCNIN
DFWAGSYMLLHYLSAQLCWFAETYGVDAVITPSLWGQEI IDALLLKTYP I
DLWTSSMLLDLVLVHAMVPFVERYGPEAIVY PDLRANPRADVLWLEQANG
* : * : * : . . . * : * : . . . : :

```
gi|121997427|ref|YP_001002214.
gi|88810314|ref|ZP_01125571.1|
gi|255254649|ref|ZP_05334193.1|
gi|255338062|ref|ZP_05378933.1|
gi|20808984|ref|NP_624155.1|_h
gi|254478158|ref|ZP_05091540.1|
gi|38505758|ref|NP_942378.1|_h
gi|170079601|ref|YP_001736234.
gi|126661502|ref|ZP_01732553.1|
gi|172035267|ref|YP_001801768.
gi|86608324|ref|YP_477086.1|_C
gi|86607166|ref|YP_475929.1|_C
gi|255256456|ref|ZP_05335917.1|
gi|209526382|ref|ZP_03274910.1|
gi|162453956|ref|YP_001616323.
```

```

-----APELFAKEPWNERTVTDANPLFSAALPNRFLAVVPAD-----
-----LDRNLFGAQNWMDPTDGNPLFSAALPNRFLALVPAS-----
-----NSATTTKDLKTPTLPNRFFAILPTK-----
-----NSATTTKDLKTPTLPNRFFAILPTK-----
NSLTHLSQKEFDEWFKEWNNIPRSKGLEEKLTIANMPNRFLAIAPIN---
NLLTHLSQKEFDEWFKEWNNIPRSKGLEEKLSIANMPNRFLAIVPIN---
-----QGFSQWVPQPGDRQLLTAGFPNVIMLVLP-----
-----EFEDWIQPPDDRQLLTAGFPNVIMLVLP-----
-----FGFNQWIKQPSQRSLLTAGFPNVLVLVLP-----
-----DFKEWIEPHKPDQLLTAGFPNVIIMILPNNHKS-----
-----EFEEWIPQPSNALLTAGFPNVLLLLLLP-----
-----PFAEWIPQPSANTLLTAGFPNVILLLLP-----
-----RKSSISSFIDQPTIPNRFVALIPES-----
FR---ESFQELSPDGTDLPLTRFNERLSTTLATAGFPNTITILVGSS---
ADG-----AAPPAHAARYQDVLPEQLNPCSYAGIFPNTFIALVPLGGER
          : ** : :

```

```
gi|121997427|ref|YP_001002214.
gi|88810314|ref|ZP_01125571.1|
gi|255254649|ref|ZP_05334193.1|
gi|255338062|ref|ZP_05378933.1|
gi|20808984|ref|NP_624155.1|_h
gi|254478158|ref|ZP_05091540.1|
gi|38505758|ref|NP_942378.1|_h
gi|170079601|ref|YP_001736234.
gi|126661502|ref|ZP_01732553.1|
gi|172035267|ref|YP_001801768.
gi|86608324|ref|YP_477086.1|_C
gi|86607166|ref|YP_475929.1|_C
gi|255256456|ref|ZP_05335917.1|
gi|209526382|ref|ZP_03274910.1|
gi|162453956|ref|YP_001616323.
```

```

-----QAEAIGR-AIEAHLHEWVAETAEGVSDLLRAVNLHTDP-----
-----AAEAIAQ-DITEQLRQWTVGRTEAALQRVLQEAGVEDRR-----
-----NVEEIRLLNLKKTVKDEYIKIGEYVFSHLVKSND-----
-----NVEEIRLLNLKKTVKDEYIKIGEYVFSHLVKSND-----
-----EGEALKLGEECEKNFKEQLKSLAKKTVKRLESILNLEGD-----
-----EGEALKLGKECKNFKEQLKSLAKKTVKRLESILNLEGD-----
-----KEKVSAAAMQSAKNHLLNAWRNIAHLVFAELQQR-HWQKQ-----
-----EAQVAAAMQSAKNFLLDAREIAKEVLAEALQGDHRHWQTN-----
-----KDKVETAMQTAKSTLISEWLNISKLVFSELEER-YWMKS-----
QQDLKDNPIYAATQLAKRTLKEEWKKLGDQSLFLQDSKQWQN-----
-----EERVEAAMQFAQQTlKEEWKQLAQRVLDHLQADRRWRLRQ-----
-----EERLQAAMQFAQQTlKEEWRHLAERALEHLQLERRWLVQ-----
-----EESKI1KLAHKMEDIVRKKWKEMVQD1VNLNFKFLDKIRLDK-----
-----ETAQQLGETLCEKLRKSEWSAIAQNI1REGIPDHPIPELRQGIRDR-----
-----YLTLKLDLGEAREAVLGRWRLOADLARYFLEOAAKRPEAMSDS-----

```

FIGURE 9 (continued)

24/31

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

-----TTPAYQQAREQLAGFPEVYWSAVGWDEIATDANGEPEAD-----
 -----DLHAHRQLREQLQGFPEVHWAAVPYSALTADGERKVRTG-----
 -----EQKKLLLRQLSNFPDVYWVALPLEN-----
 -----EQKKLLLRQLSNFPDVYWVALPLEN-----
 -----LEKIQLQIENQLLNIFYQVYWAVMPWFEEKESP-----
 -----LEKIQLQIENQLLNIFYQVYWAVMPWFEEKESP-----
 -----LNPTDPTWKHWLDAQWQTYWSAMAIGAEGKTLKSVG-----
 -----LSDTDPSPWKGWLDQAQWQTYWSAVAIGDKNSSFKSVG-----
 -----LSFDHKTWKGWLSQWQTYWTAVPIGVDDKKQEDQEGQLEKIALT
 -----INPNTWDKWLKCQWQTYTAFPIGDPESDLTCSLRVG-----
 -----LHPEDPTWQGWLEHQWQHYWAAVPLGSPAHALKDTG-----
 -----LRPEDPTWQGWLEHQWQFYWAAVPLGSPQQLKDTA-----
 -----KDEDCNAVIKKQTLDFPEIYWVAMPFKKDG-----
 TIAFLQNPQNHKRIEKILEEFSQSGGHDEQNRHDLNWKRSCEWWRG--
 -----ERKAFEATWARQHEDVLFTSWSAAAWPSIERVNDPESLAIR----

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

-----RLAEALRPFFYEPQAEAPGFLSS-----
 -----ALAEALRPFFYPGDPIEPGFLGS-----
 -----GDVNRPDWEIQLDKIKDYFN-----
 -----GDVNRPDWEIQLDKIKDYFN-----
 -----KEVLDDYKEIIGETELYKTI-----
 -----KEVLDDYKEIIGETELYKTI-----
 -----IPTQTPEKAKQDQWVREQNEAFSA-----SLF
 -----IPTQTSKKADLEKWIEQQNKAFNG-----KLF
 SSEIYRENEEIKDSEIEKDRSWVAIQNQAAYGLNE-----NTALF
 -----KNETEEQKKDAFEAWCHQQNQLTNPKESLFET-----NEEF
 -----ILRPPQVSIDADPWVSAQNRTTHHLEA-----KQKLF
 -----ILRS-QESIERDPWVSAQNRTYHLKG-----KRQLF
 -----RYITEEDFKDFFEEVQ-----
 -----LWNAQIDQWTQPYWTAVPLGDPNTSLMIEKNDDQLSEDWI
 -----EALVAQRPAQALPREDAEALAAWR-----ERHTT

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

-EAWRVLQRPPIEVIDPEAGTPATFYRPNPGVLYPALYELSERVMGGVKAT
 -RAWRVLSRPLELDG-----AATFYQPNPGVLYPVYDLVDRVLVAAKSV
 -SDEVEEIKDLLRYIK----DKGEYGPNIIGNIYGLLYSFMEKMMGTRKGI
 -SDEVEEIKDLLRYIK----DKGEYGPNIIGNIYGLLYSFMEKMMGTRKGI
 -SEFIESVPKIWSKNK----EKDEEYRDFSIAYSLLLELLEKLLGARKSI
 -SEFIESVPKIWSKNK----EKDEEYRDFSIAYSLLLELLEKLLGARKSM
 AQEELDFIEKSTDLEFIQQRNRNYAGSLNVGSWWADIFGQTRFTLSAVKNA
 TDTERDFFEKATAKFTQNRNRNYAGSLNAGSWWADICGQTRLALNTVKNA
 LEKELTFLQKAAKLRQDQNK-YPFNTNIGSWWAAIFDQTRLALTAVKNG
 LKAIFKLTDIETEKEPKTKYSKQPNLNVGSWWAYVFDQLRTSLNAVKNA
 GDEEELFLRVSFWDPERQQRQYKAGLNVGSWWAAVEQSRRLQAACKQA
 ADPELKFRLRISFWDPERQQRQYKAGLNVGSWWAAVEATRRLQAACKQA
 -----KENVGLIYHLGYSALEKSMGIRKNL
 KKQEKVAQTRVINPPPTTAEKMLYTTLLNGTWWGSIQSRCCQAIQAIKNT
 WIPGETFARYAEARYVYARTNRTVHQCERGFDPYLLHHALVSRHGLRKAE

: . : *

FIGURE 9 (continued)

25/31

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

RP-FAAQHEGGYRCSLTGEVEWLTHDRSHLDLPPGQREGHDT-----
 RA-FDALEQHGRCYCSLTAEAEWLTDNPAHLSLPPGQRDQADT-----
 RN-FNQYEEIGRKCSICGEHNVIIYRCTREEDEKIEKGKESY-----
 RN-FNQYEEIGRKCSICGEHNVIIYRCTREEDEKIEKGKESY-----
 RE-FEQLQKGRKCSLCGEFEVLPLDWEKLSKEKGLVKEKE-----
 RE-FEHLQKGRKCSLCGEFEVLPLDWEKLSKEKGLVKEKE-----
 RN-WKIPTAFGPRSTISGLGPVVHPQAPSHQRDWVTEGDTQK-----
 RN-WKLPTAFGTRSTISGLGPVVHPQNGQHRDWLTEGEAQH-----
 RD-WQIPTAFSTRSTISGLGSVVHPS-----DDWIEEGKTKA-----
 RT-WELPTAFTVRSSLSGIGSAVHPIVNEEKPDRISEKEINE-----
 RD-WQLPVAFGPRSTVSGLGPVVHPS---PRQGWITEGESRR-----
 RD-WQLPVTFGPRSTVSGLGPVVHPS---PRNRWISSEGESRR-----
 RN-FSQINEYGKKCNICGTKEGVKAG-MGSLQVGKYISEKE-----
 RT-WAIPISPGVRSTLSGQLSALHPQFRYSGNFREGYGLPIESMRLF---
 AQGAIAEESGEKCTLCGLRQALGAGDAGASVDAQRETARAFWRRFDRNS
 :.: . :

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

-----LWSLVAKRHPSWARKGEHLGALAALKRL-----WPQRFRCR
 -----LWSRLGRKRPTWVRKGEHLGALATLKLRL-----WPTLFCE
 -----KIKLLREQNAI IKKSDDKSITYKYLAQG-----EGLCSRC
 -----KIKLLREQNAI IKKSDDKSITYKYLAQG-----EGLCSRC
 -----QLCGVCLAKRLFPKVMKEELNLSEEMKF-----PSTSEMA
 -----QLCGVCLAKRLFPKVMKEELSLSEEMKF-----PSTSEMA
 -----YWQRQAGLFDGSEQLNATETVKRGLEKV-----LPALLGR
 -----YWQRDAGLFDGIEQLNATETVKRGLEKV-----LPKLLDR
 -----LWKRQAGLFDGIEQLNATETVKRGLHLV-----LPKLLLEL
 -----FWSEKFGFLFDGTEKLNATEIVKRVLHKL-----LPEILNY
 -----LWQRQAGLFDGREQLNATETLKRGLYKV-----LPELFRE
 -----LWQRQAGLFDGREQLNATETVKRALDKV-----LPELFPT
 -----ALCIPCFVKRALDKYLGDKVDGKFIDYT-----FPSTAEAM
 --WRLIAEVYPGLFDGSEMLNALELTKRMAWVYGGVGESLGIDFSNLENE
 DDGAERLCAVCTMKRVLVRAGVATDERGARRRVGLTAAWAGPATPLDDVC

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

ELAGTLGQEQTFRFVVSTHTMALATS-----LARAEEPNPAGG
 ELKDTLN-MSFSRFVVSTHTMALATT-----LDKLG--DAPLAI
 LTKRAAEIYFKDMFGKNNIEESFPST-----AEIALLDIINNNSD
 LTKRAAEIYFKDMFGKNNIEESFPST-----AEIALLDIINNNSD
 TIGEKRRADNKEFIAEFKELFGEFK-----QKLLLPNTVS---
 TIGEKRRADNKEFIAEFKELFGEFK-----QKLLLPNTVS---
 G-AKELKTYYPDLTVGIAGYLKTQPD-----GLPIFEQACSAISQKI
 D-PTLKYYPDLTVGVAGYLKTQSEKNG---EALNQYQDACRAIRTKI
 E-ADKIQMSYPDLTAGMAGYLKNGTS-----EDLEHFEKTSKAVINEY
 E-EQEISLYYPDLTAGVAGYLKNALLNNQ---SEIDYYKRSCHKIYKEI
 LDEDQIAASYPDLTAGVAGYLKTQGR-----REREHFEQACQAVEARY
 LDEDEVAASYPDLTAGVAGYLKTHGR-----RELEHFEQACRAVEAIH
 ASSNFKRKALKDAEVQFNAYIDDIKK-----IVGENV
 KIDYSQLIRFPNLSSIASARFAYQDLQDQDNHGKLSQYWQKLQINIKQF
 DRDELRVFPFSTATIAAQGYLCAVATRR--ELQAQVAEVVRCCEDARLPR

FIGURE 9 (continued)

26/31

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

LSAEQAEAIRNADRVALPRRLAQR-----LHHLEHGEL-----LARLP
 DTELRNEIEQRGervalPRRLADK-----LREHPNREL-----IERIPA
 NDLTNIKRYKEVFKKWCGDNFDY-----ELLYEENLN-----ENYFKT
 NDLTNIKRYKEVFKKWCGDNFDY-----ELLYEENLN-----ENYFKT
 -----VPKLKDNQLFEIDG-----QWLMRESYR-----EEYIKS
 -----VPKLKDNQLFEIDG-----QWLMRESYR-----EEYIKS
 LGDRHKVADSVTQDWGIPWVEEN-----MEKKYHPRLLN-----SGWLVE
 TADHHEIANFVRQEWGIPWIDETH-DPLFENLPHPRLLN-----AGWLVE
 LAEHNKIPDEITDKWGIPWADTN-----PTIKYHPRLLN-----AGWLVE
 QPLLD--NDNLSQSWGIPWINNNKELKLSNTIHHPRLLN-----AGWLVE
 P-EVRGVLREMRGKWGIPWADRQ-----NLRYHPRLLN-----AGWLVE
 P-GMREVIREMRGKWGIPWADQQ-----NLRYHPRLLN-----AGWLVE
 LRQIQVKPLNKIEGDFNNVNIEG-----EWFEEENLC-----TENIKK
 GKTSQEYQDFLAQTRGRPFHIPKTDICINRQNLPGRYNGVMFSSKWLA
 TSFVRSLSLAEAAQRADGAGLRFLEYEAEELTVFPEVLAAAEERAPEGGG

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

WVE-----QQRESDDGDETAARQ
 WIE-----AQR--DEGKAEAAIS
 YDFN-----SKRLGKLKELLKSID
 YDFK-----SKRLGKLKELLKSID
 EHTE-----NFKVEEFQKMSQKIT
 EHTE-----NFKVEEFQKMSQKIT
 ELED-----LDQTELPYRQALQTE
 ELEN-----LDKEAQKVYRQDLQNE
 DLE-----KESTAENRLKLQKL
 DVEISHDSNI-----EYREKLEELSTKRRTTIQV
 DLEL-----DEETQRSYRQDIETL
 DLEA-----DEATQKRCRQEIEAT
 QFG-----IEIDDKIEDLRKSL
 DMGLE-----KEQRPELRKIVDLTH
 REQGDGAAQRGGLRDRDGKRVPRSKVEALRRAVESLRRAAKELDRPAGSA

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

AIRKVTGTEAEAYYALLLMDGDRMGAWLSGGDGTTPYRAAFHP-----
 RLAHALGYKPETYYALLLMDGDYMGAWLSGDERFTIPYRAAFHP-----
 KRIKDRLNKKKYYAVIKLDGDNMGKWLAGE--LAPYMLEMYHS-----
 KRIKDRLNKKKYYAVIKLDGDNMGKWLAGE--LAPYMLEMYHS-----
 KLEKHRVNPSRYYAILQMDGDHMGKWLKGE--NNPPIMETLHP-----
 KLEKHRVNPSRYYAILQMDGDHMGKWLKGE--NNPPIMETLHP-----
 IEKFYPQNNPTSWSYVLAAGDGDGMSEWLKGTkmrtyGDYFPQVL-----
 IQKFYPQNNPASWYAIAAGDGDGMSEWLKGTkMEAYKEYFPKAL-----
 VKDYYPNNPTDCYIIAAGDGDGMSEWLKGGKLSYGHYTPENL-----
 INKYYPNNPTNWWYILVAGDGDSMRDWLRGSKLKNYGEYLPDELREKI--
 LGRYYPNVNPASWYVLAAGDGDGMGKWLARGEKMSYYDYLQPLQRLANG
 LSRYYPSINPASWYVVAAGDGDGMGQWLSGGKMKPYDYLPQPLQRLADG
 KQITDKVGEPNPYAVVLFDDGDNMGEWLAGKLLHNFEHIYGSDLWN----
 SECFGDSSPTDWWVMILGDGDGMGQYVSGKKLHPYQEYLTETHPETFID
 RRAGDRIPAPGSQVALIALDGDRISSQILLGEKIGARWRDVIHPE-----

: *** : : *

FIGURE 9 (continued)

27/31

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

-----DLADAVDTRFGDHTVLQAYLDTPRAVSPARHLAISGA
 -----KIQAGLQARFQDNSPLDDYLNAQRAVSPGRHLAISGA
 -----KLQGFLPEDFKEKIK-----DKKRLMTPAVHSLISEA
 -----KLQGFLPEDFKEKIK-----DKKRLMTPAVHSLISEA
 -----KVGEHIKQHAKDNLSS--ILCKKHPTTPSLHQTLSRK
 -----KVGEHIKQHAKDNLSS--ILCKKHPTTPSLHQTLSRK
 -----TVPEDLQPTFQPFSE-----QPKRMGPATHNALSRA
 -----NVPNDLKKSFDKFS-----EAKRMGPATHNALSRA
 -----NVEGDIAEAFNEFLT-----VPKRMGPSTHNSALSRA
 -----INDSNDEIPKEYKEPLQNFLE-----VRKRMGPSTHSALSRA
 QTEAIGEVFGKDKEELEQVKDAFQKFVKN-----SRKRMGPATHSALSRA
 QRDAIGRVLDS--EELKQVKDAFSEFVKA-----SRKRMGPATHSALSRA
 -----KLPNKVKEELKTSIKN-----KLLAPTIHSLISLS
 RDRYPDLDENQYEQKKRQFIKAFRDDKHS--LLKTRKRMGPATHVGLNRA
 -----AVATMESNEVTRAARWPELLGRKRLMGPSTHAFVNRV
 : * * :.

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

LNDFSTTLARRVVEECHHGRVLYAGGDDLMLPTGDLLSAMRELRLAYS
 LNDFSTVIAREVVEREHIGRVLYAGGDDLMLFAVSDLLSAMRRLRLAYS
 LKNYSKYVKEIVEETRAGKVIYSGGDDVFAIVNLNLYLLDVMVKLRAAF
 LKNYSKYVKEIVEETRAGKVIYSGGDDVFAIVNLNLYLLDVMVKLRAAF
 ISTFALQEVRRIVEETHYGLVYAGGDDVLALLPVEEVLECAVELQNAFK
 ISTFALQEVRRIVEETHYGLVYAGGDDVLALLPVEEVLECAVELQNAFK
 LLDFSNQLVPYLTEQRYAGRLIYSGGDDVLAYTNLWEWDQWLWDIRQAFR
 LLDFSNQLVPYLTEQRYAGRLIYSGGDDVLAYTNLWEWDQWLWDIRQAFR
 LLDFSNQLVPYLTEQRYAGRLIYSGGDDVLAYTNLWEWDQWLWDIRQAFR
 LLDFSNQLVPYLTEQRYAGRLIYSGGDDVLAYTNLWEWDQWLWDIRQCFR
 LLDFSNQLVPYLTEQRYAGRLIYSGGDDVLAYTNLWEWDQWLWDIRQCFR
 LLDFSNQLVPYLTEQRYAGRLIYSGGDDVLAYTNLWEWDQWLWDIRQCFR
 LRNYSLEFVKKIVEDDHLGKLIYSGGDDVLAFVNLKDLFEVMMRRLRAAFS
 LLDFSNRLVPFITEKRFVCGRVVYSGGDDVMTVLPLEDLPDYLLSLRAAWC
 LAEFAHTIVPWWVEREFSGRLIYAGGDDVLAIAPAGEALDLCARLAQLYS
 : :: . :.* *:::*****::: : :

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

GTATSG-----
 GLAPEE-----
 GHLIVN-----
 GHLIVN-----
 EVLSS-----
 EVLSS-----
 GDRD-----E
 GDKD-----D
 GDKDPLGFNNDG-----TFNLE
 GQED-----P
 GGED-----D
 GGED-----D
 GHKIE-----
 GGDDPYQPQAG-----DPD
 AAWVLDTPGEGPWAWRAKTWTSSTSIRASERFVVVAPSETGEPLPWPIP

FIGURE 9 (continued)

28/31

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

-----EAEEDAWCRNGFIYRQDRLYLTMGSAQATASMGAVIAHHTPLT
 -----GEAAHEMRFGNGFARRRGRLLYLTMGENATASMGAVIAHHQAPLG
 -----NEIIPDFTIDAGFVERKEEIDVMMGNKATASMGVIVIAHYKEDLK
 -----NKIIPDFTIDAGFVERKEEIDVMMGNKATASMGVIVIAHYKEDLK
 -----EASMSAGIVIVHHKYPLY
 -----EASMSAGIVIVHHKYPLY
 HREFDSTGHYWHSTAHKN--LPQRPLFTMGDGATISFGITIAHHSVPLA
 CGEFNNQGHYQWQADRFR-----PLFTMGKNATISFGIVIAHHSVPLA
 KSEFKNEGNYWQWNQENKPNHLAERPLFTMGQATISFGIVIAHHSVPLA
 YNEFKNEGQWQWKNPKPKNIAQRPLFTMGKCATISFGIVIAHHSIPLA
 WREFENLGDYQWQKGRPPEGLSRRPLFTLGSATISFGVIVIAHHSVPLA
 WREFANEGDYWQWQKQALPAGLSRRPLFTLGSASISFGVIVIAHHSVPLA
 -----NNEIKVDWSNNTGFVEKDGKLLLTMGKNATASCGVIVIAHYKTPLK
 LKFQSQGGYWRPTSDTNDQGLPDRSLFTMGFGATMSLGIVIVDKSVPLP
 EASRRGPIPAGDAAAASAAESAERLLPMMGAGQTLASGIAIGHYKTSLG
 : * * . * . . *

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

AALAEELRQAEQRAKNE-----
 AVLRLALRGAEKTAKAYNR-----
 NVIASVDEVEEEYAKKVE-----
 NVIASVDEVEEEYAKKVE-----
 LALKEVQLAQKKAKDERQ-----
 LALKEVQLAQKKAKDERQ-----
 IALEHLWEAEEEEAKEHEYGEDED-----
 IALENLWEAEEGAKDHEYEDFALEGKER-----
 IALENLWEAEEEEAKEHYINENN-----
 IALESLWEAEEEEAKEHYTQKDNK-----
 IALENLWQAEAAEAKGHLYAHGCP-----
 IALENLWEAEEQGAKEQAYAQSS-----
 MVMDKVRETEKKAKSNTE-----
 VALEALWEAESDRAKELSGGLMSIDLRLGLEWRSLSLMRFVLSEIFRL
 GLVKAAWDERDRVKGEDEERDCAKEDNEE-----RIRVKGDDEERDRATR
 : ..

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

---GGRDAFALSVLKRSGGALRHVGRWAPAAGIDEMSALRDFAE---LH
 ---GSRDAFSLTIKIRAGGALELTLPWQMG--EDVIGALRTLSAA---LQ
 ---G-KDAFAIKLILHSGENYIARAKWNYDDARNKEGTIGLLKDI---NS
 ---G-KDAFAIKLILHSGENYIARAKWNYDDARNKEGTIGLLKDI---NS
 ---YNRNAPCLKFIKSGS-----ALKECGGKWALMDFLREL---IE
 ---YNRNAPCLKFIKSGS-----ALKECGGKWALMDFLRGL---IE
 --KSKSDAVQVRVIYGNNGNLTATSKFEVFKTWKDLDDIET-----
 --KKKKDAVQLRVIYGNNGNLTATCKFDTFRAWQQLLEITG-----
 --KTYKDAVQVRVIYGNNGNLTATSKFEVFEQWKLLRINE-----
 --DTYKDAVQVRVLYGNNGTLKSTSKFDIFNQWKQLLKLD-----
 ----KDSVQVRVLFANGNQLVSTSKFDTFDKWKDLDFDWKLMG---QQ
 ----KDSAQVRVLFANGNQLVSTCKFDTFDKWKDLDFDWKLLG---QK
 ----KDSFAISLLKRSGEKIIVSKWKYGDMDVLEILKELSYLF---RR
 SSVPNKNGLCFRVLYSSGNTLEALIKGELYAGWRDWITTEHHQALSSLLY
 EDLAVKNAFISIRRYTRGGEKARLRLSWGRCAAPGKALAGYLRVKAVIEG
 : : . . *

FIGURE 9 (continued)

29/31

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

ANPEASRRAAYNVAGWLADLPEPDTLPPEDGVAGYLEAVLHYQFCRQGLE
 KNAGASRRAAYNVRGWLRLDLPKPVAVGGEAPFRRLLGQMLHYQFQRQKLE
 FFKEDKLSISFVGKLLAALERLDVDSLPPQGIFISELKRNKRSLNENLSK
 FFKEDKLSISFVGKLLAALERLDVDSLPPQGIFISELKRNKRSLNENLSK
 HFKKGEFSSTFPYQFFEVVDRLYDDENKEQIIGILRNELKRIFLRQSGNR
 HFKKGEFSSTFPYQFFEVVDRLYDDENKEQVIGILRNELKRIFLRQSGNR
 -IDASTYETAATVLEQHPIPVREAIMPWVNVLVERRDALDKDQQS--TLR
 -LEASTFETAATLLEQHPIPTSVAIAPWVSFAVVERRSNLNEEQKI--GLQ
 -LEASLFEQAATLWTQHPIPIQEAIKPWTVAFCSSRETLDNDIRK--SFQ
 -LEPSLFEQAATIWEQHPIPDKSAIEPWTKAFCSSRD-IGEELQQ--DYQ
 QDQSSLFEQAQLWEQHPAPSPEAVPAWATLFCSSRDLFKQESEQKEHFQ
 QDQSSLFEHAAQLWEQHPAPSLEAVPAWARLFCSSRDLFHGESEQTDRFQ
 NQSGVRISRRIVYTLAEFAKLKDKCGNYIAETGIINTEIKRVVLRAVDT
 RLAEELPRHADLTPSSYLIAQAARAIARRDDKDLVQENIDKLLIWLRYW
 FKTGALPGRLPYKLREVEGAVLAAWLVEAKALPGGLTSREARRELVKRVE

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

QQGQPIHARRLGSLASVSPTGDELSSAAKIRDRLSHLVGVAEFLARETRR
 KEGAPIHAERLAKLCPVENP-----AKAADFIENMLSVAEFLARETRS
 DEKEKVERDLYILLYSLYKE-----VKYDNFISLLTILAFNLNRGGEK
 DEKEKVERDLYILLYSLYKE-----VKYDNFISLLTILAFNLNRGGEK
 DILEKYASEILLPKFDEIVK-----ETPDRVIDFANMLIARSIAME
 DILEKYASEILLPKFDEIVK-----ETPDRVIDFANMLIARSIAME
 SRLACFLIQLWQTTSQKN-----WEKEAKNWLKAAAFMKRNRYI
 KYLAHFLQELWNTTNADD-----WDKESKNWLKAAAFNLRTKRV
 TALSKFIEDLWNTTSEEQ-----LNQDVQSWFKLTAFLVLRNRKI
 KILTSFIQGLWITTPQNN-----LDKEVKNWKLAAAFVKNRNRKI
 NKLVEFIQHFVQHSPLAQPP-----EWDLEIKNWFKLAAAFVLRNRTV
 TKLVEFIQHFVQHSPLAQPG-----AWDLAIKNWFKLAAAFVLRNRTV
 VSGNIGKEEKKNISKEISDVLKN--LYHKYDDIDNFKLLEISAFIGKE
 EDWARSVHVQWLIDREKNPDTPKPIG--IDLKDLNLIREFSAFWLDKMQQ
 PFARGLFQAQCAAEVEGAFFIWWQGLLSALRAHERRWRAPDAVPPGATPV

gi|121997427|ref|YP_001002214.
 gi|88810314|ref|ZP_01125571.1|
 gi|255254649|ref|ZP_05334193.1
 gi|255338062|ref|ZP_05378933.1
 gi|20808984|ref|NP_624155.1|_h
 gi|254478158|ref|ZP_05091540.1
 gi|38505758|ref|NP_942378.1|_h
 gi|170079601|ref|YP_001736234.
 gi|126661502|ref|ZP_01732553.1
 gi|172035267|ref|YP_001801768.
 gi|86608324|ref|YP_477086.1|_C
 gi|86607166|ref|YP_475929.1|_C
 gi|255256456|ref|ZP_05335917.1
 gi|209526382|ref|ZP_03274910.1
 gi|162453956|ref|YP_001616323.

 LDT-----

 HSCNE-----
 ARI-----
 KFPN-----

 KLGGKA-----
 SLNQE-----
 KLGRNGR-----
 TLGRNGR-----
 GE-----
 RQQWHRQDQEA-----
 DDGAPSRGLGSEQGLAVCRFLGALARGEEDAQ

FIGURE 9 (continued)

30/31




FIGURE 10

31/31




FIGURE 11

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2010/064095

A. CLASSIFICATION OF SUBJECT MATTER
INV. C12N15/82

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CAMP WIM VAN: "Yield enhancement genes: seeds for growth" CURRENT OPINION IN BIOTECHNOLOGY, vol. 16, no. 2, April 2005 (2005-04), pages 147-153, XP002613153 ISSN: 0958-1669 page 151	1-20
X	----- WANG YALING ET AL: "OsRAR1 and OsSGT1 physically interact and function in rice basal disease resistance" MOLECULAR PLANT-MICROBE INTERACTIONS, vol. 21, no. 3, March 2008 (2008-03), pages 294-303, XP002613154 ISSN: 0894-0282 abstract; figure 4 page 297 - page 298 ----- -/-	10,13



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier document but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
 "&" document member of the same patent family

Date of the actual completion of the international search

9 December 2010

Date of mailing of the international search report

17/02/2011

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Krüger, Julia

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2010/064095

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHUNG EUNSOOK ET AL: "Suppression of pepper SGT1 and SKP1 causes severe retardation of plant growth and compromises basal resistance" PHYSIOLOGIA PLANTARUM, vol. 126, no. 4, April 2006 (2006-04), pages 605-617, XP002613155 ISSN: 0031-9317 abstract; figure 7 page 611 - page 614	10,13
X	----- WO 2005/113777 A1 (MAX PLANCK GESELLSCHAFT [DE]; ROMEIS TINA [DE]; WITTE CLAUS-PETER [DE]) 1 December 2005 (2005-12-01) page 90 - page 91; figures 14, 20 page 100 - page 101	10,13
X	----- NOEL LAURENT D ET AL: "Interaction between SGT1 and Cytosolic/Nuclear HSC70 chaperones regulates Arabidopsis immune responses" PLANT CELL, vol. 19, no. 12, December 2007 (2007-12), pages 4061-4076, XP002613156 ISSN: 1040-4651 page 4062, column 2	10,13
X	----- GRAY WILLIAM M ET AL: "Arabidopsis SGT1b is required for SCFTIR1-mediated auxin response." PLANT CELL, vol. 15, no. 6, June 2003 (2003-06), pages 1310-1319, XP002613157 ISSN: 1040-4651 page 1315, column 1	10,13
X	----- AZEVEDO CRISTINA ET AL: "Role of SGT1 in resistance protein accumulation in plant immunity" EMBO (EUROPEAN MOLECULAR BIOLOGY ORGANIZATION) JOURNAL, vol. 25, no. 9, May 2006 (2006-05), pages 2007-2016, XP002613158 ISSN: 0261-4189 page 2009, column 2, paragraph 2 - page 2010, column 1, paragraph 1; figure 4 ----- -/--	10,13

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2010/064095

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>CHOE S ET AL: "Overexpression of DWARF4 in the brassinosteroid biosynthetic pathway results in increased vegetative growth and seed yield in Arabidopsis" PLANT JOURNAL, BLACKWELL SCIENTIFIC PUBLICATIONS, OXFORD, GB, vol. 26, no. 6, 1 June 2001 (2001-06-01), pages 573-582, XP002370275 ISSN: 0960-7412 DOI: DOI:10.1046/J.1365-313X.2001.01055.X -----</p>	1-20

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2010/064095

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2005113777	A1	01-12-2005	NONE

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP2010/064095

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.b of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application and necessary to the claimed invention, the international search was carried out on the basis of:
 - a. (means)

☐

 on paper
 - ☒

 in electronic form
 - b. (time)

☒

 in the international application as filed
 - ☐

 together with the international application in electronic form
 - ☐

 subsequently to this Authority for the purpose of search
2.

☐

 In addition, in the case that more than one version or copy of a sequence listing and/or table relating thereto has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments:

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2010/064095

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

see additional sheet(s)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-20

Methods for enhancing yield related traits in plants relative to control plants, comprising modulation of the expression of a SGT1 polypeptide, constructs, plants and products in relation thereto.

2. claims: 21-40

Methods for enhancing yield related traits in plants relative to control plants, comprising modulation of the expression of a CLC-pKG polypeptide, constructs, plants and products in relation thereto.

3. claims: 41-60

Methods for enhancing yield related traits in plants relative to control plants, comprising modulation of the expression of a HD-hydrolase-like polypeptide, constructs, plants and products in relation thereto.
