

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 January 2002 (17.01.2002)

PCT

(10) International Publication Number
WO 02/04607 A1

(51) International Patent Classification⁷: **C12N 5/08**, 5/24,
C07K 16/00, G01N 33/53, A61P 35/00, C12N 15/07,
G01N 33/577 // A61K 39/395, A61P 35/00

(21) International Application Number: PCT/GB01/03059

(22) International Filing Date: 9 July 2001 (09.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0016824.5 7 July 2000 (07.07.2000) GB
0017139.7 12 July 2000 (12.07.2000) GB
60/270,588 23 February 2001 (23.02.2001) US

(71) Applicant (for all designated States except US): **CEN-
FOLD HOLDINGS S.A.** [PA/PA]; Panama City (PA).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **KARPAS, Abraham**
[GB/GB]; Cambridge Serologicals Limited, MRC Centre,
Hills Road, Cambridge CB2 2QH (GB).

(74) Agents: **MASCHIO, Antonio** et al.; D Young & Co, 21
New Fetter Lane, London EC4A 1DA (GB).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA,
ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv)) for US only

Published:

— with international search report
— before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: HUMAN MYELOMA CELL LINE

(57) Abstract: There is provided a human myeloma cell line for use in a method for producing a human monoclonal antibody.



WO 02/04607 A1

HUMAN MYELOMA CELL LINE

The present invention relates to a human myeloma cell line. In particular it relates to the use of a human myeloma cell line in a method for making a human hybridoma capable of producing human monoclonal antibodies.

BACKGROUND

The technique for providing murine monoclonal antibodies was devised in 1975 by Köhler & Milstein (1,2). In this method, antibody-producing spleen cells from an immunised mouse are fused with cells from a mouse myeloma to produce hybrid cells known as hybridomas. After fusion, the cells are selected using drugs that kill the myeloma parental cell, whereas the unfused parental spleen cells have a limited life-span and soon die, so that only hybrid cells survive. These hybridomas have the capacity to proliferate indefinitely and to secrete antibody specific for the antigen used to immunise the spleen cell donor. Those hybridomas producing antibody of the desired specificity are cloned, resulting in a hybridoma cell line producing a homogenous population of antibody molecules (i.e. monoclonal antibodies).

Much research effort has been directed to the adaptation of this technology to produce human monoclonal antibodies. Attempts to use the mouse myelomas to derive human monoclonal antibodies were unsuccessful because the hetero-specific hybrids quickly lose the relevant human chromosomes. Attempts have also been made to use known antibody-secreting human myeloma cell lines (such as the IgE λ secreting U266 human myeloma cell line, and the RPMI-8226 cell line which contains a heterogeneous mixture of lymphoblastoid and plasmablast cells (3)) for the production of human monoclonal antibodies, but they have all failed, in spite of promising early reports.

Due to the lack of success in obtaining human antibody-producing hybridomas, various alternative cell lines have been considered. For example, the Epstein-Barr virus (EBV) has been used to immortalise antibody-producing human peripheral blood B-lymphocytes. However, such cell lines have a number of drawbacks when compared with the murine monoclonal antibody-producing hybridomas. For example, they tend to be unstable with

respect to growth, they are low producers of antibodies, and EBV does not preferentially immortalise lymphoblasts engaged in antibody responses (4).

5 Strategies have also been developed to produce monoclonal antibodies with human character which bypass the need for an antibody-producing human cell line. For example, useful mouse monoclonal antibodies have been "humanised" by linking rodent variable regions and human constant regions (chimeric antibodies) (13). This somewhat reduces the human anti-mouse immunogenicity of the antibody but residual immunogenicity is retained by virtue of the foreign V-region framework. Moreover, the antigen-binding
10 specificity is essentially that of the murine donor. CDR-grafting and framework manipulation (EP 0239400) has improved and refined antibody manipulation to the point where it is possible to produce humanised murine antibodies which are acceptable for therapeutic use in humans. However, the antibody specificity remains determined by the murine binding site donor. Moreover, their production involves complex manipulation of
15 sequences and is laborious.

Phage display technology has been developed for the *in vitro* generation of human monoclonal antigen-binding fragments, whose V genes can be recovered and engineered into antibody genes to produce monoclonal antibodies (14). However, production of
20 human monoclonal antibodies by this route is a complex multi-step procedure, and the binding specificity of the antibodies, since it is engineered *in vitro* as a result of combination of randomly-generated sequences, is not subjected to *in vivo* affinity maturation as is the case with natural antibodies.

25 A transgenic mouse strain which contains human instead of mouse immunoglobulin genes has been developed (15). This mouse strain contains human genes and produces human antibodies; however because the diversity created is selected not in a human but the mouse host, and the antibodies undergo affinity maturation in the mouse, the antibodies are not truly human.

30

From an analytical perspective, none of the known methods is capable of dissecting man's immune response to specific antigen challenges. Direct analysis of genes from heavy and light chains following polymerase chain reaction (PCR) amplification has become the method of choice to characterise the type of antibodies expressed by people in certain

conditions. This method has the disadvantage that the information of the heavy-light chains pairing is lost.

5 The murine hybridoma procedure for producing monoclonal antibodies has the added advantage (over the above-mentioned non hybridoma based procedures) that it can preferentially immortalise antigen activated B cells, thus producing a long-lasting supply of monoclonal antibodies.

10 There is thus a need for a method for producing human monoclonal antibodies which has all the advantages of the conventional murine monoclonal antibody production method.

SUMMARY OF THE INVENTION

15 The present invention provides a human myeloma cell line which is capable of acting in an analogous manner to the mouse myeloma used in conventional methods for making murine monoclonal antibodies. Therefore, for the first time, the present invention enables the reliable production of truly human hybridomas and monoclonal antibodies.

20 The present invention thus concerns a human myeloma cell line which is capable of being fused with a human lymphocyte using a standard technique, such as a polyethylene glycol (PEG) fusion protocol, use of inactivated Sendai virus (2) or electrofusion methods (17), to produce hybrid cells. The present inventor has also developed a method for producing a myeloma cell line with a considerable growth advantage (i.e. a much shorter doubling time than a primary plasma cell culture from a myeloma patient).

25

30 The human myeloma cell line of the present invention can be used as a general antibody producing cell. For example, previously characterised antibodies can be produced by transfection of the myeloma cell line with known antibody genes. The myeloma cell line is suited for "physiological" antibody production because the post-translational modifications of proteins produced by the myeloma cells are likely to be identical to those produced by human antibody-producing cells.

In a preferred embodiment, however, the myeloma cell line of the present invention is used in a method for producing a human hybridoma which is capable of producing human monoclonal antibodies. Further aspects of the invention, therefore, relate to: a method for producing a hybridoma, which comprises the step of fusing a human myeloma cell line
5 with a human lymphocyte; a human hybridoma cell line produced by such a method; and a human monoclonal antibody produced by such a hybridoma.

The hybridoma-production method the present invention can be used to immortalise human antibodies that are provoked by the natural body reaction in pathological
10 conditions. This enables an immune response to be monitored over time, and also provides a source of antibodies which are more likely to be useful in therapy than antibodies prepared by other methods. Analysis of the specificity of such antibodies can be used to provide information on the disease-specific antigens which provoked the antibody *in vivo*. This information can, in turn be used to devise antigen-based
15 therapeutic strategies, such as improved vaccines.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1: Quantitation of IgG secretion by two hybridomas. 707/164(o) and
20 707/100(●). Each hybridoma was seeded at a concentration of 3×10^5 cells/ml in RPMI-1640 growth medium supplemented with 10% fetal bovine serum. Samples were collected at the indicated intervals for IgG quantitation by immunofixation and for cell counts. After four days the cell count of the 707/164 hybridoma reached 8×10^5 cells/ml while that of the 707/100 hybridoma was 6×10^5 cells/ml. Thereafter the number of
25 viable cells decreased and by the 10th day nearly all the cells were dead when the level of IgG in the medium was at its highest, namely 210 µg/ml of IgG for both cultures.

Figure 2: Analysis of immunoglobulin secreted by the myeloma Karpas 707 cell line and hybridomas. The cells were grown at a concentration of 2×10^6 cells/ml in L-methionine, L-cystine deficient medium which was supplemented with 10% dialysed fetal
30 bovine serum and with [³⁵S] methionine and cystein at 250µCi/ml (Amersham International). The cells were incubated for 8 hours at 37°C in a CO₂ incubator. After incubation the cell suspensions were centrifuged at 1000g for 5 minutes. The supernatant

was analysed by sodium dodecyl sulfate polyacrylamide gel electrophoresis after total reduction. The identity of radiolabelled secreted Ig was confirmed by immunofixation.

Panel:

5

1 Small quantity of κ light chain produced by the myeloma cells

2 IgG that react with the gp41 HIV-1 produced by the hybrid with the 164 cells.

10 3,4 IgG producing hybridoma formed by fusing the myeloma cells with fresh white blood cells.

5, 8, 9, 16 IgG producing hybridoma formed with tonsil cells.

15 6,7 Two hybridomas with tonsil cells that secrete the κ light chain only.

10-12 IgM secreted by hybridoma formed with fresh tonsil cells. Band 12 suggests that this hybridoma secretes both light chains.

20 13 This hybridoma appears to secrete only the two light chains since it was negative for the heavy chains G and M.

14 IgG with the two light chains produced by a hybrid between the myeloma and tonsil cells.

25

15. Hybridoma with tonsil cells that secrete high levels of light chain only.

Figure 3a Karyotype of the chromosomes of the Karpas 707 cell line which are near tetraploid. Note that chromosomes 2, 14 and 22 which contain the immunoglobulin genes
30 are normal and diploid.

83,XX,-X,-X,del(1)(p11p36)x4,+del(1)(q11q44),-2,-2,der(3)del(3)(p21p25)inv(3)
(q23q25),+del(3)(q13q21),der(5)t(1;5)(p11;q31),der(5)t(2;5)(q21;q31),
del(5)(q13q33)x2,+add(5)(q31),-6,-6,add(7)(p25),del(8)(q22q24),-9,

del(11)(q23q25),-11,del(12)(p12p13),-13,-13,-14,-14,add(15)(p11)x2,
 der(16)t(1;16)(p11;p11),add(16)(p11)x2,-16,der(18)t(1;18)(q23;p11),
 add(18)(p11),add(18)(q23),-18,add(19)(q13)x2,+der(19)t(1;19)(q11;p13)
 +der(19)t(dup(1)(q23q44);19)(q11;p13),-21,-22,-22,+mar1x2,+mar2

5

b. Karyotype of the hybridoma Karpas 707/164 which is near hexaploid containing the normal pairs of chromosomes which are readily obvious in the case of chromosomes No. 1.

10 125,XXX,-X,-X,-X,del(1)(p11p36)x4,+del(1)(q11q44),-2,-2,add(3)(p25),
 der(3)del(3)(p21p25)inv(3)(q23q25),add(4)(q31),-4,der(5)t(1;5)(p11;q31),
 der(5)t(2;5)(q21;q31),del(5)(q13q33)x2,+add(5)(q31),del(6)(q21q23)
 -6,-6,add(7)(p25)x2,del(8)(q22q24),-9,del(11)(q23q25),-11,
 del(12)(p12p13)x2,-13,-13,-14,-14,add(15)(p11)x2,der(16)t(1;16)(p11;p11),
 15 add(16)(p11)x2,-16,der(18)t(1;18)(q23;p11),add(18)(p11),add(18)(q23),
 -18,add(19)(q13)x2,+der(19)t(1;19)(q11;p13),+der(19)t(dup(1)(q23q44);19)(q11;p13),
 -21,-21,-22,-22,+mar1x2,+mar2

Figure 4: Electron micrograph of the myeloma (a) and hybridoma (b). In contrast to
 20 the cytoplasm of the myeloma cells which does not appear to contain detectable levels of
 rough endoplasmic reticulum (RER), the cytoplasm of the hybridoma is taken up by
 parallel and concentric RER resembling plasma cells. These morphological observations
 can be correlated with the low level of κ light chain produced by the myeloma and high
 levels of IgG which is secreted by a hybridoma. The cells were fixed in 3%
 25 glutaraldehyde and the ultra-thin sections stained in uranyl acetate and lead citrate.

DETAILED DESCRIPTION OF INVENTION

In a first aspect, the present invention relates to a human myeloma cell line which is
 30 suitable for use in a method for producing a human monoclonal antibody-secreting
 hybridoma.

The myeloma cell line may be developed from primary plasma cell cultures from patients with myeloma, for example from the bone marrow or peripheral blood of such patients.

In order to develop a homogenous cell line, the cells from the patients are cultured *in vitro* and amplified.

5 It is normal for newly developed human myeloma cell lines to have a relatively long doubling time, of the order of 70 hours. However, the present inventor has surprisingly found that it is possible to produce a much faster growing cell line by culturing the cells *in vitro*. For example, it is possible to reduce the doubling time to approximately 35 hours or less.

10 The myeloma cell line of the present invention may have a doubling time which is less than 0.75, preferable less than 0.5 times the doubling time of the cell line before *in vitro* culture treatment. The myeloma cell line of the present invention may have a doubling time of less than 50 hours, preferably less than 40 hours, more preferably about 30 hours.

15 In order to use the standard polyethylene glycol (PEG) fusion protocol, the myeloma cell line of the present invention may be resistant to PEG. The present inventor found that 99% of human myeloma cells were killed after first treatment with PEG for lymphocyte fusion. Myeloma cells may be made PEG-resistant by treating cells with multiples cycles of PEG. The present inventor has found that in the region of twenty cycles of treatment
20 produces a cell line in which only about 25% of cells are killed by PEG treatment. Selection of a PEG-resistant myeloma cell results in a cell line which can be fused with human lymphocytes to produce a human/human hybridoma in accordance with the present invention.

25 In the standard murine monoclonal antibody-secreting hybridoma production method, fused cells are selected using hypoxanthin, aminopterin, thymidin (HAT) medium. Murine myeloma cells are HAT-sensitive because they lack the enzyme hypoxanthine:guanosine phosphoribosyl transferase (HGPRT), whereas hybridoma cells are HAT-resistant, the HGPRT gene being provided by the spleen cell fusion partner. In
30 order to use the same selection procedure, the myeloma cell line of the invention should be sensitive to HAT. HAT sensitivity may be selected by growth in 8-azaguanine and/or 6-thioguanine or other selective agents as known in the art. As used herein, the term "HAT-sensitive" means that, in a hybridoma fusion procedure, treatment with HAT

medium results in preferential killing of unfused myelomas such that hybridoma cells may be isolated.

5 It is convenient for the myeloma cell line to be Ouabain resistant, to facilitate the fusion and selection procedure with cell lines or with fresh cells isolated from tissues (such as bone marrow, tonsil, lymph node, spleen and malignant tissue). Cells may be rendered Ouabain resistant by continued culture in increasing Ouabain concentrations. The resulting cell lines are HAT-sensitive and Ouabain resistant.

10 It has been found that the myeloma cells having a growth advantage have an increased chromosome number compared to myeloma cells with longer doubling times. The myeloma cells of the present invention may have an increase in the number of chromosomes (anuploid), or be near tetraploid, and may have a number of gross chromosomal abnormalities. However, one or more (preferably all) of chromosomes 2,
15 14 and 22 may be normal and/or diploid. Chromosomal analysis may be performed using standard techniques such as the G-banding method.

The present inventor has produced and characterised a human myeloma cell line which has a short doubling time, is PEG resistant, Ouabain resistant and HAT sensitive (see the
20 Examples). This cell line is known as "Karpas 707", and a sample has been deposited at the ECACC, Salisbury, Wiltshire, SP4 0JG, under Deposit No. 00071108.

The second aspect of the invention relates to a method for producing a myeloma cell line.
25 In addition to steps selecting for: a growth advantage, HAT-sensitivity, Ouabain-resistance and/or PEG sensitivity (as described above), the method may also comprise the step of introducing into the cells a marker capable of conferring sensitivity or resistance to an agent other than HAT, Ouabain and PEG. Such a marker or marker(s) may be introduced into the cell by, for example, genetic engineering methods known in the art.
30 Representative markers: enzymes producing a detectable product, such as β -galactosidase; proteins which confer antibiotic resistance or susceptibility; and genes whose presence is fatal to the cell under certain conditions ("suicide genes"), such as herpes simplex virus thymidine kinase (HSVTK) which causes cell death in the presence of gancyclovir.

The third aspect of the invention relates to a method for producing a human hybridoma cell line. The hybridoma may be produced by a method analogous to the conventional method for producing murine hybridomas (7). For example, the myeloma cell line of the first aspect of the present invention may be fused with a human lymphocyte using, for example, a standard PEG fusion protocol, inactivated Sendai virus or electrofusion methods.

Hybrid cells may be selected from unfused cells by any of the means known in the art. For example, hybrid cells may be selected and/or isolated by cell sorting (such as by fluorescence-activated cell sorting (FACS)), using magnetic beads or using cytotoxicity. The conventional murine method for selecting hybrid cells involves growing the cell population post-fusion in a medium containing a chemical which will selectively kill myeloma cells which are not fused with lymphocytes (such as HAT). The hybrid cells may then optionally be cloned to produce a homogenous hybridoma cell line. General methods for hybridoma production are known in the art. For example, the following procedure may be used:

A single-cell suspension of human lymphocytes from an immunised individual is formed in serum free RPMI 1640, supplemented with 2 mM L-glutamine, 1 mM sodium pyruvate, 100 units/ml penicillin, and 100 µg/ml streptomycin (RPMI). The cell suspension is filtered through sterile 70-mesh Nitex cell strainer, and washed twice by centrifuging at 200 g for 5 minutes and resuspending the pellet in 20 ml serum free RPMI.

25 2×10^8 lymphocytes are combined with 4×10^7 myeloma cells (kept in log phase in RPMI with 11% foetal bovine serum (FBS) for three days prior to fusion), centrifuged and the supernatant is aspirated. The cell pellet is dislodged and 2 ml of 37°C PEG 1500 (50% in 75 mM HEPES, pH 8.0) is added while stirring over the course of one minute, followed by the addition of 14 ml of serum free RPMI over seven minutes. Additional RPMI can be added and the cells are centrifuged at 200 g for 10 minutes. After discarding the supernatant, the pellet is resuspended in 200 ml RPMI containing 15% FBS, 100 mM sodium hypoxanthine, 0.4 mM aminopterin, 16 mM thymidine

(HAT), 25 units/ml IL-6 and 1.5×10^6 lymphocytes/ml. The suspension is dispensed into ten 96-well flat bottom tissue culture plates at 200 μ l/well. Cells are fed on days 2, 4, and 6 days post-fusion by aspirating 100 μ l from each well with an 18 G needle, and adding 100 μ l/well plating medium with or without 10 U/ml IL-6 and lacking lymphocytes.

When cell growth reaches 60-80% confluence, culture supernatants are taken from each well and screened for reactivity to the desired antigen by ELISA. ELISAs may be performed as follows. Immulon 4 plates are coated at 4°C with 50 μ l/well with 100ng/well of antigen in 50 mM carbonate buffer, pH 9.6. Plates are washed with PBS with 0.05%, Tween 20 and blocked 30 minutes at 37°C with 0.5% Fish Skin Gelatin. Plates are washed as described above and 50 μ l culture supernatant is added. After incubation at 37°C for 30 minutes, 50 μ l of horseradish peroxidase conjugated goat anti-human IgG(fc) [diluted 1:10,000 in PBST] is added. Plates are incubated at 37°C for 30 minutes, washed with PBST and 100 μ l of substrate, consisting of 1 mg/ml TMB and 0.15ml/ml 30% H₂O₂ in 100 mM Citrate, pH 4.5, is added. The colour reaction is stopped with the addition of 50 ml of 15% H₂SO₄. A₄₅₀ is read on a plate reader.

The term lymphocyte is intended to include any antibody-producing cell. For example, the lymphocyte may be a lymphoblast, such as an EBV-transformed B cell. The lymphocyte may also be an *in vitro* cultured cell. Alternatively, the lymphocyte may be freshly isolated from a human subject. For example, the lymphocytes can be obtained from organs such as tonsils as well as lymph nodes, spleen, blood or bone marrow.

A human subject may be a healthy individual, in particular an individual who has been immunised with a particular antigen, or who has had (and recovered from) a particular disease. The human subject may alternatively be an individual who *has* a particular disease. A lymphocyte from the human subject may, therefore, be capable of producing an antibody of interest.

Myelomas according to the invention may also be fused with non-antibody-producing lymphocytes, or other white blood cells, which produce lymphokines. Especially preferred are tumour-infiltrating cells which produce anti-tumor antibodies or

lymphokines of therapeutic utility. The hybridomas thus obtained are capable of reliably producing anti-tumor antibodies or lymphokines in large quantities.

One advantage with the hybridoma-production method of the present invention, as well as
5 the conventional method for producing mouse monoclonal antibody-secreting hybridomas, is that the myeloma cell line preferentially fuses with antigen-activated lymphocytes.

The present invention also provides a hybridoma cell line, and a monoclonal antibody
10 produced by such a cell line.

Preferably the hybridoma of the present invention produces a high yield of antibodies. For example, the hybridoma may secrete more than 100 µg/ml antibody, preferably at least 200µg/ml antibody, more preferably at least 300 µg/ml antibody.

15

The hybridoma of the present invention may have extensive rough endoplasmic reticulum (RER). In contrast to the cytoplasm of the myeloma cells (which do not appear to contain detectable levels of rough endoplasmic reticulum), the present inventor has found that the cytoplasm of a hybridoma cell of the present invention is taken up by parallel and
20 concentric RER resembling plasma cells. The term "extensive" is used herein to indicate that the levels of RER in the hybridoma are comparable to those found in plasma cells.

The monoclonal antibody may be of any isotype: for example IgG, IgM, IgD, IgA or IgE. It may comprise light chains, heavy chains or both.

25

Preferably the antibody is useful in the prevention or treatment of a disease. For example, if a disease-specific antibody-producing cell is isolated from one patient, the cell could be immortalised by the method of the present invention, enabling the antibody to be produced in large quantities. The antibody may be given to a patient in a passive
30 immunisation protocol to confer resistance to the disease. Alternatively, the antibody may be given to a patient suffering from the disease to boost immune responses and help clear the infection. It is possible that the derived hybridomas with various white blood cells will give rise to a range of therapeutic humoral factors.

Human monoclonal antibodies produced according to the method of the present invention may be employed in substantially any process which involves ligand-polypeptide binding, including *in vivo* therapeutic and prophylactic applications, *in vitro* and *in vivo* diagnostic applications, *in vitro* assay and reagent applications, and the like. For example, antibody
5 molecules may be used in antibody based assay techniques, such as ELISA techniques, according to methods known to those skilled in the art.

As alluded to above, the antibodies produced according to the invention are of use in diagnostic, prophylactic and therapeutic procedures. For example, antibodies produced
10 according to the invention are of use diagnostically in agglutination, ELISA, immunoperoxidase, Western blot and *in situ* protein detection by standard immunohistochemical procedures. For use in these applications, the antibodies may be labelled in accordance with techniques known to the art. In addition, such antibodies may be used preparatively in affinity chromatography procedures, when complexed to a
15 chromatographic support, such as a resin. All such techniques are well known to one of skill in the art.

Therapeutic and prophylactic uses of antibodies prepared according to the invention involve the administration of thereof to a recipient or patient.

20 Substantially pure antibodies of at least 90 to 95% homogeneity are preferred for administration to a patient, and 98 to 99% or more homogeneity is most preferred for pharmaceutical uses. Once purified, partially or to homogeneity as desired, the selected antibodies may be used diagnostically or therapeutically (including extracorporeally) or in
25 developing and performing assay procedures, immunofluorescent stainings and the like (Lefkovite and Pernis, (1979 and 1981) Immunological Methods, Volumes I and II, Academic Press, NY).

The antibodies thereof of the present invention will typically find use in preventing,
30 suppressing or treating inflammatory states, allergic hypersensitivity, cancer, bacterial or viral infection, and autoimmune disorders (which include, but are not limited to, Type I diabetes, multiple sclerosis, rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease and myasthenia gravis).

In the instant application, the term "prevention" involves administration of the protective composition prior to the induction of the disease. "Suppression" refers to administration of the composition after an inductive event, but prior to the clinical appearance of the disease. "Treatment" involves administration of the protective composition after disease symptoms become manifest.

Animal model systems which can be used to screen the effectiveness of the antibodies in protecting against or treating the disease are available. Methods for the testing of systemic lupus erythematosus (SLE) in susceptible mice are known in the art (Knight *et al.* (1978) *J. Exp. Med.*, **147**: 1653; Reinersten *et al.* (1978) *New Eng. J. Med.*, **299**: 515). Myasthenia Gravis (MG) is tested in SJL/J female mice by inducing the disease with soluble AchR protein from another species (Lindstrom *et al.* (1988) *Adv. Immunol.*, **42**: 233). Arthritis is induced in a susceptible strain of mice by injection of Type II collagen (Stuart *et al.* (1984) *Ann. Rev. Immunol.*, **42**: 233). A model by which adjuvant arthritis is induced in susceptible rats by injection of mycobacterial heat shock protein has been described (Van Eden *et al.* (1988) *Nature*, **331**: 171). Thyroiditis is induced in mice by administration of thyroglobulin as described (Maron *et al.* (1980) *J. Exp. Med.*, **152**: 1115). Insulin dependent diabetes mellitus (IDDM) occurs naturally or can be induced in certain strains of mice such as those described by Kanasawa *et al.* (1984) *Diabetologia*, **27**: 113. EAE in mouse and rat serves as a model for MS in human. In this model, the demyelinating disease is induced by administration of myelin basic protein (see Paterson (1986) *Textbook of Immunopathology*, Mischer *et al.*, eds., Grune and Stratton, New York, pp. 179-213; McFarlin *et al.* (1973) *Science*, **179**: 478; and Satoh *et al.* (1987) *J. Immunol.*, **138**: 179).

The antibodies of the present invention may also be used in combination with other antibodies, particularly human monoclonal antibodies (MAbs) reactive with other markers on human cells responsible for the diseases. For example, suitable T-cell markers can include those grouped into the so-called "Clusters of Differentiation," as named by the First International Leukocyte Differentiation Workshop (Bernhard *et al.* (1984) *Leukocyte Typing*, Springer Verlag, NY).

Generally, the present antibodies will be utilised in purified form together with pharmacologically appropriate carriers. Typically, these carriers include aqueous or

alcoholic/aqueous solutions, emulsions or suspensions, any including saline and/or buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride and lactated Ringer's. Suitable physiologically-acceptable adjuvants, if necessary to keep a polypeptide complex in suspension, may be chosen from
5 thickeners such as carboxymethylcellulose, polyvinylpyrrolidone, gelatin and alginates.

Intravenous vehicles include fluid and nutrient replenishers and electrolyte replenishers, such as those based on Ringer's dextrose. Preservatives and other additives, such as antimicrobials, antioxidants, chelating agents and inert gases, may also be present (Mack
10 (1982) *Remington's Pharmaceutical Sciences*, 16th Edition).

The antibodies of the present invention may be used as separately administered compositions or in conjunction with other agents. These can include various immunotherapeutic drugs, such as cyclosporine, methotrexate, adriamycin or cisplatin, and immunotoxins. Pharmaceutical compositions can include "cocktails" of various
15 cytotoxic or other agents in conjunction with the selected antibodies, receptors or binding proteins thereof of the present invention, or even combinations of selected polypeptides according to the present invention having different specificities, such as polypeptides selected using different target ligands, whether or not they are pooled prior to
20 administration.

The route of administration of pharmaceutical compositions according to the invention may be any of those commonly known to those of ordinary skill in the art. For therapy, including without limitation immunotherapy, the antibodies of the invention can be
25 administered to any patient in accordance with standard techniques. The administration can be by any appropriate mode, including parenterally, intravenously, intramuscularly, intraperitoneally, transdermally, *via* the pulmonary route, or also, appropriately, by direct infusion with a catheter. The dosage and frequency of administration will depend on the age, sex and condition of the patient, concurrent administration of other drugs,
30 counterindications and other parameters to be taken into account by the clinician.

The antibodies of this invention can be lyophilised for storage and reconstituted in a suitable carrier prior to use. This technique has been shown to be effective with conventional immunoglobulins and art-known lyophilisation and reconstitution

techniques can be employed. It will be appreciated by those skilled in the art that lyophilisation and reconstitution can lead to varying degrees of antibody activity loss (e.g. with conventional immunoglobulins, IgM antibodies tend to have greater activity loss than IgG antibodies) and that use levels may have to be adjusted upward to compensate.

5

The compositions containing the present antibodies or a cocktail thereof can be administered for prophylactic and/or therapeutic treatments. In certain therapeutic applications, an adequate amount to accomplish at least partial inhibition, suppression, modulation, killing, or some other measurable parameter, of a population of selected cells is defined as a "therapeutically-effective dose". Amounts needed to achieve this dosage will depend upon the severity of the disease and the general state of the patient's own immune system, but generally range from 0.005 to 5.0 mg of antibody *per* kilogram of body weight, with doses of 0.05 to 2.0 mg/kg/dose being more commonly used. For prophylactic applications, compositions containing the present selected polypeptides or cocktails thereof may also be administered in similar or slightly lower dosages.

10
15

A composition containing an antibody according to the present invention may be utilised in prophylactic and therapeutic settings to aid in the alteration, inactivation, killing or removal of a select target cell population in a mammal. In addition, the antibodies described herein may be used extracorporeally or *in vitro* selectively to kill, deplete or otherwise effectively remove a target cell population from a heterogeneous collection of cells. Blood from a mammal may be combined extracorporeally with the selected antibodies, cell-surface receptors or binding proteins thereof whereby the undesired cells are killed or otherwise removed from the blood for return to the mammal in accordance with standard techniques.

20
25

The capacity to produce unlimited quantities of a particular antibody from an antibody-producing cell in a patient can also be used to gain valuable information about the immune response. For example, if the patient has a disease, it is possible to deduce the disease-specific antigens targeted by the immune system. This information can be used to design better drugs and vaccines to treat and/or prevent the disease.

30

Almost every disease produces an antibody-mediated immune response, and so is a candidate for prevention/treatment by an antibody-mediated therapy. The procedure of the

present invention will be invaluable to immortalise human antibodies provoked by the natural defensive reaction of pathological conditions. Of particular interest, however, is the use of the method of the present invention to immortalise tumour-infiltrating lymphocytes. This may produce useful anti-cancer antibodies and also uncover tumour
5 specific antigens. Such antigens could in turn be used as anti-cancer vaccines.

By analysing antibodies produced at different times, it is also possible to map the course of an immune response to antigenic challenge. This may provide valuable information on the optimum timing for a particular treatment strategy or vaccination protocol.

10

In a further aspect of the present invention there is provided the use of a human myeloma cell line according to the first aspect to produce an antibody, by transfection of the cell line with antibody-producing genes. The post-translational modifications effected by the human myeloma are identical to those formed in human antibodies. Thus, for
15 biotechnological production, the human myeloma line is preferred to produce antibodies derived not only from human hybridomas but also by transfection of already known antibody genes.

A human hybridoma cell line of the present invention may also be used to select a mutant
20 that will be also HAT-sensitive in the same way that the murine Sp2/0-Ag14 was developed to produce a hybridoma (16) following fusion with another lymphocyte. In particular, a hybridoma produced using the myeloma cell line of the present invention which produces low/undetectable levels of antibody, but has high levels of RER, may be selected for fusion with a second lymphocyte capable of producing an antibody of
25 interest. In this way, a hybridoma may be produced which is capable of secreting high titres of the desired antibody.

The following examples serve to illustrate the present invention, but should not be construed as a limitation thereof. The invention particularly relates to the specific
30 embodiments described in these examples.

EXAMPLES

Example 1 – Production of an improved human myeloma cell line

The present inventor has been trying to develop new human myeloma cell lines for over twenty years. The present inventor has previously described a myeloma cell line which was developed from the bone marrow and peripheral blood of a 53 year old male patient
5 diagnosed as having multiple myeloma (5,6). Since this cell line now secretes only κ light chain it has a potential practical advantage over myeloma cell lines that also produce the heavy chain.

Selecting for myeloma cells having a growth advantage

10 The doubling time of this myeloma cell line after isolation was over 70 hours, but following continuous *in vitro* culture over many months a faster growing cell emerged with a doubling time of approximately 30-40 hours, with an increase tendency to stick to the surfaces of glass or plastic when in stationary culture. The adherent cells can be detached easily by forceful pipetting in a similar way to the mouse myeloma cells used to
15 generate hybridomas.

Selection for HAT-sensitivity and Ouabain-resistance

The conventional method for making murine hybridomas relies on the fact that the unfused murine myeloma cells are sensitive to hypoxanthin, aminopterin, thymidin
20 (HAT). In order to select for human myeloma cells which are (HAT)-sensitive, they were grown in 8-azaguanine. Azaguanine-resistant cells that grew at a concentration of 30 μ g/ml were further subjected to 30 μ g/ml of 2-Amino-6-mercaptopurine (6-Thioguanine). The first twenty clones that grew out were tested for their sensitivity to HAT medium. The fastest growing clone which did not seem to produce any resistance
25 was selected for further work. In order to be able to fuse the myeloma cells also with antibody producing cell lines it was decided to select also for an Ouabain resistant subline. Growing the myeloma cells with an increasing concentration of Ouabain eventually led to the emergence of a HAT-sensitive Ouabain-resistant subline.

30 Example 2 – Production of a human hybridoma

In order to optimise the conditions and screening for hybridoma formation the present inventor decided to develop (for the initial fusions) an antibody-producing human B-cell lymphocyte with known specificity. Therefore the peripheral blood lymphocytes from healthy HIV-1 infected individual were infected with EBV. From the EBV-immortalised

B-cells, cells that produced monoclonal antibodies to HIV-1 gp41 were selected and cloned, and referred to as Cell line Karpas 164. The present inventor then attempted unsuccessfully to fuse the myeloma cells with the EBV-immortalised human lymphocytes using polyethylene glycol (PEG) of different molecular weights.

5

In order to try to investigate the reasons for the repeated failure to develop hybrids the myeloma cells alone were treated with PEG using the same procedure as that used for cell fusion. Thereafter, the cells were seeded in growth medium, but without HAT or Ouabain, and examined for viability. Examination of the PEG-treated cells revealed that
10 the PEG was highly toxic, killing over 99% of the cells.

Development of a PEG-resistant sub-line

In view of the above findings, the present inventor decided to try and develop a PEG-resistant subline. This was achieved by treating about 10^7 myeloma cells with PEG
15 (1500MW) following the protocol used for murine hybridoma formation. The few viable cells which eventually grew out were treated again with PEG. He repeated such cycles of treatment more than twenty times over a period of about 18 months before a subline of the myeloma emerged where only about one quarter of the cells were killed by the PEG treatment. However, in the process HAT-resistant cells reappeared. The cells were
20 therefore treated once more with 8-azaguanine and cloned. A clone was isolated which did not revert when grown in HAT medium. This cell line is known as the myeloma Karpas 707 cell line, a sample of which has been deposited as Deposit No. ECACC 00071108.

25 *Fusion with EBV-immortalised lymphocytes*

The HAT-sensitive, PEG-resistant cells were then fused with the EBV immortalised human B-lymphocytes 164 cells producing the monoclonal antibodies to gp41 using a standard PEG fusion protocol (7). The cell suspension was seeded in multi-well plastic trays. Following growth in the presence of HAT and Ouabain, clones resembling the
30 myeloma cell line grew out. The tissue culture fluid gave strong staining of the membranes of acetone fixed HIV-1 infected Karpas 45 T-cells in the Karpas AIDS cell test for anti-viral antibodies (8).

Characterisation of the human monoclonal antibody produced by the hybridoma

Further studies using several polypeptides of the gp41 of HIV-1 enabled us to establish that our monoclonal antibody reacted against the peptide LAVERYLKDQQLGIWG but it failed to react against four other peptides of the gp41 of HIV-1 (results not shown). The epitope that this monoclonal antibody recognises seems not to be present in the African NDK strain of HIV-1 (9), since the antibody fails to react with NDK infected T-cells but reacts not only with the Cambridge isolate of HIV-1 (10) but also with the early French LAV isolate (11). A database search confirms that both the French (LAV) and British (CAM1) isolates of HIV-1 share the above amino acid sequence, whereas the African isolate has a lysine to arginine mutation.

Fusion with fresh human lymphocytes

After the successful fusion with the 164 cell line the present inventor decided to establish whether their myeloma cells would also form antibody-producing hybridomas with fresh human lymphocytes. They fused the myeloma cells with unfractionated white blood cells (WBC) from the peripheral blood of an adult and with tonsil cells prepared from surgically removed tonsils from two children. The cell suspensions obtained after the PEG fusion were seeded in 24 and 96 wells plastic tissue culture plates and grown in RPMI-1640 medium supplemented with 20% foetal bovine serum and HAT for the first 12 days. Thereafter, growth medium alone was used and colony formation was observed during the following 2-3 weeks in approximately 25% of the wells when the myeloma cells were fused with tonsil cells and in about 10% of the wells when the myeloma cells were fused with blood WBC. Tissue culture fluid collected from wells with active cell growth was assayed by ELISA for the presence of IgG. In this method, protein-A was used on the plastic wells to capture the IgG monoclonal antibodies produced by the newly formed hybridomas. The culture fluid from over 100 of the wells contained IgG.

The present inventor has, to date, developed over 20 independent Ig secreting hybrids from fusion with peripheral blood lymphocytes and over 100 antibody producing hybrids from fusion with tonsil cells.

Example 3 – Characterisation of the human myeloma cell line and hybridomas*Morphological analysis of the hybridomas*

The growth pattern of many of the cells that grew out was variable. When in stationary culture, myeloma cells grow in a uniform pattern of adherent cells forming an irregular mosaic which covered the plastic surface with continuous release of cells into the culture fluid. When grown in roller flasks, most cells remain in suspension as clusters of variable size. The growth of some hybrids resembled the parent myeloma cells but many others acquired a distinct morphology and growth forms. In stationary culture, some hybridomas adhered to the plastic surfaces and displayed numerous thick but short spindles, rectangular and triangular shapes. Other grew mainly in suspension as single cells or clusters.

Ultrastructural examination of the cultured myeloma revealed that in contrast to the cells which contained large amounts of rough endoplasmic reticulum (RER) during the first year in culture (5), after long-term *in vitro* growth the cells have lost most of the RER. However, examination of two hybridomas, 707/164 and 707/105 (formed with blood B-cells) revealed that in both cases the hybrids resembled plasma cells (Figure 4). A large part of the cytoplasm was taken up by parallel and concentric RER while the degree of dilatation of the cisternae varied from cell to cell. The contents of the RER appeared as fine granular material which is probably immunoglobulin. A large number of mitochondria could be seen in most sections, while sections through the Golgi area revealed highly developed apparatus. Since the ultrastructural examination of EBV-immortalised 164 B-cells revealed only a small number of short and thin strands of RER it will not be unreasonable to suggest that the fused lymphoblast somehow activates the latent myeloma RER for the production of Ig.

Chromosomal analysis of the myeloma cell line

To confirm that the antibody producing cells were indeed a hybrid of myeloma and human lymphocyte, cell cultures were prepared for chromosome analysis using the G-banding method described by Czepulkowski et al. (12). Early on, the cultured myeloma cells (like the fresh patient cells) were found to be hypodiploid with 45 chromosomes containing several chromosomal abnormalities (5). However, the myeloma cells which had the selective growth advantage (after multiple cycles of *in vitro* culture) and which had increased PEG-resistance were revealed to be near tetraploid containing numerous

and unusual high number of gross chromosomal abnormalities. In fact 13 of the 23 chromosomes were abnormal. Karyotypic analysis revealed 5 copies of chromosome 1 all of which were also shorter than usual, and 3 marker chromosomes. It is interesting to note that the chromosomes 2, 14 and 22 (which contain the immunoglobulin genes) are normal and diploid.

In order to establish whether the repeated cycles of PEG treatment induced these chromosomal karyotype abnormalities, the chromosomes of cells which were frozen prior to the PEG treatment were also analysed. The analysis revealed that all these changes were present before the PEG treatment. Hence it appears that, during the continuous *in vitro* culture over many months and treatment with 8-azaguanine, 6-thioguanine, HAT and Ouabain, a near tetraploid cell line emerged which had a growth advantage over the hypodiploid myeloma cells.

15 *Chromosomal analysis of the hybridoma cell lines*

Karyotype analysis of the hybridoma with the 164 cells revealed that the cells are near hexaploid, containing the near tetraploid chromosome of the myeloma cell and the normal chromosome from the lymphoblast.

20 Karyotype analysis of the EBV-transformed lymphoblasts (Karpas 164 line) that produce monoclonal antibody to gp41 of HIV-1 shows that about 10% of the cells display a similar translocation, t(1;16) (p11;q10) to that detected also in the myeloma cell line. This may be due to prolonged *in vitro* culture.

25 *Analysis of IgG production*

As to the quantity of IgG production, a comparative assay for the secretion of antibodies to the gp41 of HIV-1 by the EBV infected lymphoblasts and the hybrid (707/164) revealed that the hybridoma produced at least eight times more antibodies than the lymphoblasts following culture of identical number of cells (10^6 /ml) under the same conditions: Specifically, the hybrid 707/164 secreted 470 μ g/ml of IgG while the 164 cells secreted less than the assay detection level, which is 70 μ g/ml after four days in culture. Since both cell cultures have the same genetic information for the antibody synthesis it is clear that the hybridoma cells of the present invention are far more effective in antibody production.

Tests for the secretion of immunoglobulin confirm that the myeloma cells produced only a small quantity of light chain. The fusions with the tonsil cells and blood lymphocytes produced clones that secreted IgG as well as IgM. One of the hybrids which was formed
5 with tonsil cells produces only light chains ($\lambda + \kappa$). Such diversity of Ig production suggests that the myeloma cell line is capable of forming hybridomas with most if not all the range of antibody producing cells. In fact the present inventor has found that the hybridomas that are formed with the myeloma cell line described herein are far more effective in antibody production than murine hybridomas (specifically, the hybridoma that
10 secreted antibodies against gp41 of HIV-1 produced 470 $\mu\text{g/ml}$ yield of IgG, and the hybridomas that were formed following fusion with blood lymphocytes produced 210 $\mu\text{g/ml}$ and 150 $\mu\text{g/ml}$ of IgG). It correlates with extensive rough endoplasmic reticulum (RER) which is present in the human hybridoma as compared to the sparse RER of the murine hybridoma.

15

Various modifications and variations of the described methods and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed
20 should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in chemistry or biology or related fields are intended to be covered by the present invention. All publications mentioned in the above specification are herein incorporated by reference.

REFERENCES

1. Cotton, R.G.H. and Milstein, C. *Nature* **244**, 42-43 (1973).
2. Köhler, G. and Milstein, C. *Nature* **256**, 495-497 (1975).
- 5 3. Nilsson, K. *Int. J. Cancer* **7**, 380-396 (1971).
4. Roder, J.C., Cole, S.P.C. and Kozbor, D. *Meth Enzym.* **121**, 140-167 (1989).
5. Karpas, A., Fischer, P., Swirsky, D. *Science* **216**, 997-999 (1982).
6. Karpas, A., Fischer, P., Swirsky, D. *Lancet* **i**, 931-933 (1982).
7. Galfre, G. & Milstein, C. *Meth. Enzymology*, **7B**, 3-46 (1981)
- 10 8. Karpas, A., Gilson, W., Bevan, P.C., Oates, J.K. *Lancet* **ii**, 695-697 (1985).
9. Ellrodt, A. *et al.* *Lancet* **i**, 1383-1385 (1984)
10. Karpas, A. *Mole. Biol. Med.* **1**, 457-459 (1983).
11. Barré-Sinoussi, F. *et al.* *Science* **220**, 868-871 (1983).
12. Czepulkowski, B.H., Bhatt, B., Rooney, D.E. in Human cytogenetics: A practical
15 approach. **Vol 2**, IRL Press, Oxford University Press (1992).
13. Winter, G. and Milstein, C. *Nature* **349**, 293-299 (1991).
14. Winter, G., Griffiths, A.D., Hawkins, R.E. and Hoogenboom, H.R. *Annu. Rev. Immunol.* **12**, 433-455 (1994).
15. Bruggemann, M. and Neuberger, M.S. *Immunol. Today* **17**, 391-397 (1996).
- 20 16. Shulman *et al.* *Nature* **276**, 269-270 (1978).
17. Panova & Gustafsson, (1995) *Hybridoma* **14**,3, 265-269

CLAIMS

1. A human myeloma cell line for use in a method for producing a human monoclonal antibody, which has been selected for growth advantage by multiple cycles of continuous *in vitro* culture.
2. A human myeloma cell line for use in a method for producing a human monoclonal antibody, which possesses one or more properties selected from the group consisting of
 - a) PEG-resistance;
 - b) HAT-sensitivity; and
 - c) Ouabain resistance.
3. A human myeloma cell line according to claim 2, which possesses each of the properties a, b and c.
4. A myeloma cell line according to any preceding claim which has an increased chromosome number and or one or more chromosomal abnormalities.
5. A myeloma cell line according to claim 4 which is near tetraploid.
6. A myeloma cell line according to claim 4 or 5, wherein one or more of chromosomes 2, 14 and 22 are diploid and free from chromosomal abnormalities.
7. A myeloma cell line according to any preceding claim, substantially as described herein, and known herein as Karpas 707, which is the same as, or derivable from, the sample deposited in Deposit No. ECACC 00071108.
8. A method for producing a human myeloma cell line, comprising the steps of:
 - a) isolating a primary human myeloma cell from a myeloma patient;
 - b) culturing and multiplying the cell *in vitro* by multiple cycles of *in vitro* culture so as to select for a growth advantage over primary myeloma cells;
 - c) exposing the cells to an agent which enables selection for one or more HAT-sensitive cells;

d) optionally, exposing the cells to Ouabain and selecting one or more cells which are Ouabain-resistant; and

e) exposing the cells to polyethylene glycol (PEG), and selecting one or more PEG-resistant cells;

5 wherein the above steps c-e may be performed individually or sequentially in any order.

9. A method according to claim 8, which includes the step

 f) introducing into the cells a marker capable of conferring sensitivity or
10 resistance to an agent other than HAT, Ouabain and PEG
 wherein step f) can occur at any point in the method after step a).

10. A method for production of a human hybridoma, which comprises the step of
 fusion of a human myeloma cell line according to any of claims 1 to 7 with a human
15 white blood cell.

11. A method according to claim 10, comprising the steps of:

 a) incubating a myeloma cell line according to any one of claims 1 to 7 with a
 human white blood cell, in the presence of polyethylene glycol (PEG), inactivated Sendai
20 virus, or under electrofusion conditions;
 b) selecting for hybrid cells.

12. A method according to claim 11, wherein step b) comprises

 i) continuing the incubation in the presence of HAT with or without Ouabain;
25 ii) selecting HAT-resistant, optionally Ouabain-resistant cells; and
 iii) culturing the HAT-resistant, optionally Ouabain-resistant cells and selecting
 individual hybridoma clones.

13. A method according to any one of claims 10 to 12, wherein the white blood cell is
30 an antibody-producing lymphocyte, a lymphoblast such as an EBV-immortalised
 lymphocyte, or a lymphokine-producing white blood cell.

14. A method according to any one of claims 10 to 13, wherein the white blood cell is
 freshly isolated from a human subject.

15. A method according to any one of claims 10 to 12, wherein the human white blood cell is part of a population of lymphocytes showing varying degrees of antigen activation, and the human myeloma cell line preferentially fuses with antigen activated lymphocytes from the population.
16. A human hybridoma cell line produced by a method according to any of claims 10 to 15.
17. A human hybridoma cell line according to claim 16, having extensive rough endoplasmic reticulum.
18. A human monoclonal antibody produced by a hybridoma cell line according to claim 16 or 17.
19. A human monoclonal antibody according to claim 18, for use in a method for treating or preventing a disease.
20. A human monoclonal antibody according to claim 19, for use in conferring humoral immunity to a human subject.
21. A method of producing an antibody, which comprises the following steps:
- (i) isolation of a lymphocyte from a human subject;
 - (ii) production of a human hybridoma by using the lymphocyte in a method according to any of claims 10 to 15; and
 - (iii) isolation of an antibody secreted by the hybridoma.
22. A method according to claim 21, wherein the antibody is capable of treating and/or preventing a disease.
23. A method according to claim 22, wherein the lymphocyte is a tumour-infiltrating lymphocyte, and the antibody is useful in the treatment or prevention of cancer.

24. A method for identifying a disease-specific antigen which comprises the following steps:
- (i) production of a disease-specific antibody by a method according to claim 22 or 23; and
 - 5 (ii) deduction of the antigen which is specifically recognised by the antibody.
25. A method of monitoring an immune response in a human which comprises the following steps:
- (i) isolation of lymphocytes from a human subject at varying stages of the
10 immune response;
 - (ii) production of a human hybridomas by using the lymphocytes in a method according to any of claims 10-15;
 - (iii) isolation of an antibody secreted by each hybridoma.
- 15 26. A method for producing a human antibody which comprises the following steps:
- (i) transfection of a human myeloma cell line according to any of claim 1 to 7 with one or more antibody encoding gene(s);
 - (ii) production of the antibody by the transfected myeloma cell line.
- 20 27. A method for producing a humoral factor, which comprises the step of fusing a cell from a myeloma cell line according to any of claims 1 to 7 with a white blood cell which is capable of producing the humoral factor.
28. The use of a human hybridoma cell line according to claim 16, in a method for
25 selecting a mutant that will lead to a hybrid hybridoma.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 01/03059

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N5/08 C12N5/24 C07K16/00 G01N33/53 A61P35/00
 C12N15/07 G01N33/577 //A61K39/395, A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, MEDLINE, LIFESCIENCES, EMBASE, CHEM ABS Data, BIOSIS, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GUO YONGJUN ET AL.: "Establishment of human myeloma cell line KM-2R and its preliminary application to human-human hybridoma research." CHIN MED SCI, vol. 6, no. 2, June 1991 (1991-06), pages 92-95, XP001021594 page 93	1-6, 8-14, 21-24, 26,27
X	HISAYOSHI O'OKA ET AL.: "Establishment of Stable Cell Lines producing anti-Pseudomonas aeruginosa monoclonal antibodies and their protective effect for the infection in mice" MICROBIOL. IMMUNOL, vol. 36, no. 12, 1992, pages 1305-1316, XP001025835 page 1306	2,3, 10-14, 21,22, 24,27

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *Z* document member of the same patent family

Date of the actual completion of the international search

17 October 2001

Date of mailing of the international search report

05/11/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Wagner, R

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 01/03059

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HULETTE C M ET AL: "PRODUCTION OF A HUMAN MONOCLONAL ANTIBODY TO HLA BY HUMAN-HUMAN HYBRIDOMA TECHNOLOGY A PRELIMINARY REPORT" AMERICAN JOURNAL OF PATHOLOGY, PHILADELPHIA, PA, US, vol. 121, no. 1, October 1985 (1985-10), pages 10-14, XP000971774 ISSN: 0002-9440 page 11	2,3,10, 14,21, 24,27
X	---- "HUMAN MYELOMA CELL LINE CARRYING A PHILADELPHIA CHROMOSOME" SCIENCE, AMERICAN ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE,, US, vol. 216, no. 4549, 28 May 1992 (1992-05-28), pages 997-999, XP001018528 ISSN: 0036-8075 cited in the application abstract	7
X	---- KARPAS A ET AL: "HUMAN PLASMACYTOMA WITH AN UNUSUAL KARYOTYPE GROWING IN VITRO AND PRODUCING LIGHT-CHAIN IMMUNOGLOBULIN" LANCET THE, LANCET LIMITED. LONDON, GB, 24 April 1982 (1982-04-24), pages 931-935, XP001018518 ISSN: 0140-6736 cited in the application abstract	7
P,X	---- KARPAS ET AL: "A human myeloma cell line suitable for the generation of human monoclonal antibodies" PNAS, vol. 98, no. 4, 13 February 2001 (2001-02-13), pages 1799-1804, XP002179323 page 1799 -page 1804	2-14, 21-27
A	---- US 4 720 459 A (WINKELHAKE JEFFREY L) 19 January 1988 (1988-01-19) the whole document	2-14, 21-27
A	---- US 4 833 077 A (ABE TSUTOMU ET AL) 23 May 1989 (1989-05-23) the whole document	2-14, 21-27
	---- -/--	

INTERNATIONAL SEARCH REPORT

Int nal Application No
PCT/GB 01/03059

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>OLSSON L ET AL: "HUMAN-HUMAN HYBRIDOMAS PRODUCING MONOCLONAL ANTIBODIES OF PREDEFINED ANTIGENIC SPECIFICITY" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, NATIONAL ACADEMY OF SCIENCE. WASHINGTON, US, vol. 77, no. 9, September 1980 (1980-09), pages 5429-5431, XP001021693 ISSN: 0027-8424 the whole document</p>	2-14, 21-27
A	<p>COLE S P C ET AL: "HUMAN MONOCLONAL ANTIBODIES" MOLECULAR AND CELLULAR BIOCHEMISTRY, NORWELL, MA, US, vol. 62, no. 2, June 1984 (1984-06), pages 109-120, XP001021694 ISSN: 0300-8177 the whole document</p>	2-14, 21-27

INTERNATIONAL SEARCH REPORT

Int. Application No

PCT/GB 01/03059

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4720459	A	19-01-1988	NONE
US 4833077	A	23-05-1989	JP 59175896 A 04-10-1984
			JP 1756958 C 23-04-1993
			JP 4039999 B 01-07-1992
			JP 58216125 A 15-12-1983
			DE 3379055 D1 02-03-1989
			EP 0096839 A1 28-12-1983