

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 723 948 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
26.03.2003 Bulletin 2003/13

(51) Int Cl.⁷: **C06D 7/00**

(21) Application number: **95300595.6**

(22) Date of filing: **31.01.1995**

(54) Capsaicinoid lachrymator

"Capsaicinoid" enthaltendes Tränengas

Gaz lacrymogène à base de "capsaicinoid"

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IE IT LI NL PT SE

(56) References cited:

US-A- 3 900 560 **US-A- 5 217 708**
US-A- 5 307 960

(30) Priority: **24.01.1995 US 373834**

- P.A. SANDERS 'Principles of Aerosol Technology' 1970, VAN NOSTRAND REINHOLD COMPANY, NEW YORK, USA * page 4 - page 7 *
* page 70 *
- R.C. WEAST 'Handbook of Chemistry and Physics, 54th Edition' 1973, CHEMICAL RUBBER COMPANY PRESS, CLEVELAND, OHIO, USA * page C-231 *
- Section Ch, Week 9011, Derwent Publications Ltd., London, GB; Class B07, AN 90-079285 & JP-A-2 032 190 (OSAKA AEROSOL KOGYO KK)
1 February 1990

(43) Date of publication of application:
31.07.1996 Bulletin 1996/31

(73) Proprietor: **DEFENSE TECHNOLOGY CORPORATION OF AMERICA;**
(a Delaware corporation)
City of Yulee, State of Florida 32097 (US)

(72) Inventor: **DuBay, David K.**
Casper, Wyoming 82604 (US)

(74) Representative: **Nachshen, Neil Jacob et al**
D Young & Co
21 New Fetter Lane
London EC4A 1DA (GB)

EP 0 723 948 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description**Field of the Invention**

[0001] The present invention relates to a liquid composition useful as a lachrymator. More particularly, the invention concerns a lachrymator comprising capsaicinoids and a nonflammable water-based carrier.

Background of the Invention

[0002] The prior art provides various compositions which constitute strong irritants to the eyes, mouth and nose of a human and other animals. Such compositions are generally classified as lachrymators. Lachrymators are better known, and commonly referred to as "tear gas" or "riot gas." Lachrymators serve a significant role in society because they allow the police, military and other authorities to control unruly or disruptive persons, including persons under the influence of alcohol or other mind-altering drugs, without having to resort to physical means that may inflict long-term or permanent bodily harm or damage.

[0003] Upon application of a lachrymator, the recipient of the lachrymator is overcome by eye, nose and mouth irritation and rendered harmless. More particularly, upon application, the recipient is temporarily disabled with intense burning eye pain, blepharospasm, acute bronchitis and respiratory irritation. The prior art provides various means for dispensing or applying lachrymators, such as, for example, pressurized canisters, hand grenades, and munition cartridges.

[0004] The need for an aerosol irritant-type spray or lachrymator, for enforcement and defensive purposes, is historic as well as present day evident. Social unrest, demonstrations and rioting during the 1960's caused the extensive use of military type tear gas agents, known typically as the chemicals CN (chloroacetophen-one) and CS (ortho-chlorobenzalmalononitrile). These tear gases are actually irritants causing pain and discomfort to the lacrimal glands (tear ducts) and the upper respiratory system. Delivery systems for these agents were comprised mainly of grenades and projectiles, commonly designed as pyrotechnic (burning) type devices. In Wortley, Jr. et al U.S. Patent 3,192,105, there is disclosed a method of combining CN with a colloidal silica to produce a solid phase lachrymator. As disclosed in the '105 patent, this solid phase lachrymator is well suited for use in munitions which disperse the lachrymator by an explosion or similar means.

[0005] In 1966, Smith and Wesson, through acquisition of General Ordance Equipment Company, introduced CHEMICAL MACE® lachrymator, a liquid based CN chemical formula which was filled into pressurized aerosol spray containers. CHEMICAL MACE® lachrymator has been the forerunner of all such products since its introduction in 1966. Its use greatly lessens the amount of physical force which might be necessarily ap-

plied by a law enforcement officer, corrections officer or security officer when attempting to detain a suspect or disperse a crowd of unruly persons. Lachrymators are generally classified as "nonlethal weapons."

5 [0006] The original CHEMICAL MACE® lachrymator formula consisted of trichlorotrifluoroethanes (CFC's 111, 113) and cosmetic kerosene as the carrier agent/solvent, blended with the active ingredient (CN), and pressurized with carbon dioxide as the propellant. This formula was used extensively until about 1990, but beginning about 1990 there was increasing pressure on the chemical industry to phase out ozone depleting substances (i.e., chlorofluorocarbons--CFC's) under the provisions of the Clean Air Act.

10 [0007] The CFC's, which serve as a nonflammable carrier, provide a significant advantage for the lachrymator solution. Specifically, lachrymator solutions that employ a CFC carrier are generally less likely to ignite. More particularly, when a lachrymator is dispensed, there is always the possibility that an open flame or other source of ignition may be present that could ignite the carrier of the mixture resulting in serious bodily harm to both the user and the intended recipient of the lachrymator. A nonflammable carrier is less likely to ignite when exposed to such sources of ignition and is thus a preferred carrier for use in lachrymators.

15 [0008] However, governmental actions have been taken worldwide to reduce and eliminate the use of CFC's because of the damage caused to the ozone layer. It is likely that CFC's will be totally phased out before the year 2000.

20 [0009] Manufacturers of CFC substances have been relentless in their search for acceptable substitutes. However, to this date, manufacturers have only been able to provide solvent replacements that do not totally eliminate the problem. The most recent substitute offered is a halogenated fluorohydrocarbon, HCFC-141b (often referred to as a soft CFC). However, soft CFC's still have an adverse effect on the ozone layer and their pressure has continued for developing carrier systems for lachrymators which do not employ CFC's or HCFC's.

25 [0010] Capsicum (also known as cayenne pepper) and its chemical equivalents such as capsaicin ($C_{18}H_{27}NO_2$) have also been utilized to produce lachrymators for many years. Such lachrymators are commonly utilized in the liquid phase and are dispensed from pressurized canisters or bottles. Prior art capsicum containing lachrymators have comprised a mixture of capsicum, soybean oil and an alcohol type solvent. Capsicum containing lachrymators are preferred by some users. More particularly, some users believe that capsicum containing lachrymators are more effective than other prior art lachrymators. Additionally, some users prefer capsicum because it is a "natural" material as compared to the man-made chemicals CS and CN.

30 [0011] Unfortunately, because the prior art capsicum containing lachrymators also contain carriers which are primarily alcohol, they are considered unacceptable by

some users because they may present too much of a fire hazard. Additionally, some persons consider the prior art capsicum containing lachrymators to be unacceptable because they believe the carrier which is primarily alcohol may have an adverse impact upon tests which are utilized to determine the blood-alcohol level of a person exposed to the lachrymator. Thus, there is a need for a carrier system which does not adversely impact upon the blood-alcohol level of a recipient, is suitable for use with capsicum, does not contain CFC's and is nonflammable.

[0012] U.S. Patent 5,217,708 describes a capsicum lachrymator solution and canister for storing and applying such solutions. The solutions comprise capsicum (generally about 0.1% to about 0.8% by weight), about 7% to about 23% by weight of a glycol, about 31% to about 39% by weight of ethyl alcohol, and from about 46% to about 54% by weight of water.

Summary of the Invention

[0013] The present invention provides a new and useful capsicum containing liquid lachrymator solution which comprises capsaicinoids and a nonflammable carrier, said solution comprising from 0.1% to 0.5% by weight of capsaicinoids, from 7% to 23% by weight of a glycol, from 20% to 30% by weight of ethanol and from 55% to 65% by weight of water.

Description of the Invention

[0014] The lachrymator of the present invention includes a nonflammable carrier that is generally acceptable for use around or in the vicinity of potential sources of ignition such as, for example, lit cigarettes or cigars, candles and stoves. Furthermore, the nonflammable carrier does not contain any CFC's. Also, the lachrymator tends not to adversely impact upon the blood-alcohol level of a recipient. Additionally, the lachrymator is suitable for use in conventional pressurized aerosol spray canisters or containers and thus it does not have to be thermally vaporized by a heat source in order to perform properly. Further, the lachrymator is substantially water-based yet it is very stable for it does not freeze through a normal range of temperatures and it does not separate over periods of time. The lachrymator is also stable in that it does not spoil or lose its effectiveness over periods of time and there are no adverse interactions or reactions observed as between the various components of the lachrymator. Further, the lachrymator can be disposed of in a conventional manner.

[0015] A lachrymator solution made in accordance with the principles of the present invention comprises capsaicinoids and a nonflammable carrier. The nonflammable carrier comprises water, ethyl alcohol and glycol. The glycol may comprise ethylene glycol or propylene glycol or mixtures thereof. In a preferred embodiment, the glycol is propylene glycol. As used in this

specification and claims the following terms used in the specification are defined as follows: "capsicum" includes capsicum and chemical equivalents such as capsaicin; "lachrymator" is intended to be afforded a broad

5 interpretation including any irritant-type product that may be utilized for defense purposes; and "nonflammable" means having a flash point in excess of about 100°F (38°C) or an ignition point above 500°F (260°C).
[0016] The lachrymator of the present invention is 10 produced utilizing a mixture of capsaicinoids and a nonflammable carrier. More particularly, the lachrymator solutions of the present invention comprise from 0.1% to 0.5% by weight of capsaicinoids, from 7% to 23% by weight of a glycol, from 20% to 30% by weight of ethanol, 15 and from 56% to 65% by weight of water. In other embodiments, the lachrymator solutions of the present invention comprise from 0.1% or 0.15% up to about 0.25% or 0.3% or 0.4% by weight of capsaicinoids, from about 10% or 12% to about 16% or 20% by weight of a glycol, 20 from about 26% or 27% up to 29% or 30% by weight of ethanol and from 56% up to 60% by weight of water. In a more preferred embodiment, the solutions useful as lachrymators comprise about 0.18% to about 0.22% by weight of capsaicinoids, about 12% to about 16% by weight of propylene glycol, about 27% to about 29% by weight of ethanol and about 57% to about 59% by weight of water.

[0017] Common sources of capsaicinoids useful in 25 this invention include peppers such as cayenne peppers or chili peppers. The process usually involves drying and grinding peppers into a fine powder. Oleoresin capsicum present in the powder generally is extracted through the use of an organic solvent such as hexane. This mixture of powder and solvent is referred to as *Miscella*. Once the oleoresin extraction is completed, the 30 solvent is removed by evaporation or distillation. After the solvent is removed from the *Miscella*, the remaining oily substance is raw oleoresin capsicum, the strength of which is dependent upon the grade of the peppers. 35 That were utilized. The raw oleoresin capsicum generally is diluted to various strengths according to customer specifications utilizing a glycol and water if the diluted product is to be water-soluble. Emulsifiers and surfactants may be included in the diluted mixture to improve 40 the stability of the mixture (generally a suspension). These diluted products are generally referred to in the industry as oleoresin capsicum regardless of the concentration or strength of the raw oleoresin capsicum contained in the mixtures.

[0018] Oleoresin capsicum consists of three major 45 components that cause the pungency or hotness. These components are known as capsaicinoids, and the three major capsaicinoids in oleoresin capsicum are capsaicin, dihydrocapsaicin and nordihydrocapsaicin. Minor 50 capsaicinoids present in oleoresin capsicum include homocapsaicin and homodihydrocapsaicin. The capsaicinoid content of a given solution or mixture is the determining factor of how hot a product will be. For example,

the higher the capsaicinoid concentration, the hotter the product, and vice versa. Capsaicinoid concentration is the determination of the active components in oleoresin capsicum. Thus, a measurement of the amount of capsaicinoids in oleoresin capsicum provides an accurate level of pungency. The amounts and identification of the capsaicinoids in oleoresin capsicum can be detected utilizing high performance liquid chromatography (HPLC).

[0019] One example of a capsicum and propylene glycol mixture suitable for use in the present invention and which is commercially available is an oleoresin capsicum sold by Kalsec, Inc. of Kalamazoo, Michigan under the designation "Oleoresin Capsicum, W/S, decolorized 2% MC. This particular propylene glycol/water-based oleoresin capsicum solution contains about 5% oleoresin capsicum containing about 2.0% by weight of capsaicinoids.

[0020] An example of a suitable source of ethyl alcohol for use with the present invention is a denatured ethyl alcohol sold under the trade designation SDA-40-B by Midwest Grain Products, Inc. of Weston, Missouri 64098. SDA-40-B denatured ethyl alcohol contains about 99.9% by weight ethyl alcohol and about 0.1% by weight tertiary butyl alcohol and BITREX (denatonium benzoate).

[0021] An example of a suitable source of propylene glycol is a food grade propylene glycol USP available under the product code 70531 from the Dow Chemical Company of Midland, Michigan 48674.

[0022] When preparing the lachrymators of the invention, the indicated components are mixed together in a vessel using a stirrer for a minimum of about 10 to 15 minutes. In one embodiment, the oleoresin capsicum is first mixed with the ethanol followed by addition of the glycol and water. When preparing large quantities of the formulations as set forth in the following examples, the components can be mixed in a proportion shown for a period of about one hour in 55-gallon drums utilizing mixers. Thorough mixing is required in order to eliminate any separation of the components of the solution which would require additional mixing of the solution at a later time. Upon completion of the mixing step, the solution should appear clear and homogenous when viewed with a flashlight.

[0023] The following examples illustrate the lachrymator solutions of the present invention. Unless otherwise indicated in the following examples and elsewhere in the specification and claims, all parts and percentages are by weight, temperatures are in degrees Centigrade and pressures are at or near atmospheric.

Example 1

[0024] Denatured ethyl alcohol (114 parts) is charged to a mixing vessel followed by the addition of 42 parts of the Kalsec oleoresin capsicum solution described above (containing about 2.0% of capsaicinoids) which

is first thoroughly mixed to restore homogeneity before it is added to the ethyl alcohol. After the oleoresin capsicum is thoroughly mixed with the ethanol, 19.5 parts of propylene glycol USP grade are added to the mixing vessel followed by 209.6 parts of distilled water. The contents of the vessel are mixed for an additional 15 minutes. The mixture is the desired product which is analyzed and found to contain about 0.20% by weight of capsaicinoids, 14% by weight of propylene glycol, 28% by weight of ethanol and 58% by weight of water.

[0025] The in following Examples 2-5, the amount of the Kalsec oleoresin capsicum solution used in the examples is an amount sufficient to provide the indicated amount of capsaicinoids.

Example 2

[0026]

	%/wt
Capsaicinoids	0.3
Propylene glycol	14
Ethanol	26
Water	59.7

Example 3

[0027]

Capsaicinoids	0.22
Propylene glycol	16
Ethanol	26
Water	57.78

Example 4

[0028]

Capsaicinoids	0.4
Propylene glycol	14
Ethanol	28
Water	57.6

Example 5

[0029]

Capsaicinoids	0.15
Propylene glycol	12
Ethanol	27.85
Water	60

[0030] The lachrymator solutions made in accordance with the present invention may be utilized in any number of conventional pressurized aerosol spray con-

tainers or canisters. In order to be properly utilized in a pressurized canister, the nonflammable carrier should be a liquid or a gas at ambient temperature (i.e., about 70°F (21°C) and at ambient pressure. Preferably, the carrier has a boiling point of less than about 225°F (about 107°C) in order to ensure sufficient dispersion or distribution of the capsaicum once the solution is dispensed from the pressurized canister. Additionally, the nonflammable carrier does not solidify or freeze at low temperatures as low as -20°F. Further, the amounts of propylene glycol, ethanol and water present in the solution must be such that the components of the lachrymator solution do not separate out or come out of solution over extended periods of time such as three days, preferably about three to four weeks.

[0031] The lachrymator solution of the present invention offers several advantages over the lachrymators claimed in U.S. Patent 5,217,708. For example, these solutions of the present invention are characterized as having a higher flash point and a lower vapor pressure. For example, a lachrymator solution prepared in accordance with the procedure of Example XV in U.S. Patent 5,217,708 is found to have a flash point of about 80°F (25°C). A lachrymator solution prepared in accordance with the procedure of Example 1 of the present application is found to have a flash point of 89°F (31.6°C). Because of increased concern over flammability, this increase in flash point is a significant improvement. The reduction in vapor pressure exhibited by the solutions of the present invention result in more stable products and products which are able to withstand higher temperature extremes.

[0032] Another advantage of the solutions of the present invention when compared to the solutions described and claimed in U.S. Patent 5,217,708 is the presence of larger sized droplets when the product is aerosolized. The ballistic droplets that are produced from an aerosol lachrymator are comprised proportionately of the product ingredients. As the liquids and the droplets begin to evaporate, the droplets become smaller, thus leaving smaller particles. The smaller the aerodynamic particle size, the further into the lungs the particle may travel when inhaled, and since lachrymators are strong irritants, smaller particle sizes are not beneficial since they could yield undesirable effects. The relative amounts of ethanol and water in an aerosolized droplet, therefore, have an affect on the relative size of the droplet and particle size. Because ethanol evaporates more readily than water, those droplets containing more ethanol become smaller and leave smaller particles. Therefore, since the solutions of the present invention contain relatively lesser amounts of ethanol than the solutions described in the '708 patent, the droplets containing more water and less ethanol will not reduce in size as much as the droplets formed from the solution of the '708 patent, and the particles are larger in size.

Claims

1. A solution for use as a lachrymator comprising capsaicinoids and a nonflammable carrier, said solution comprising from 0.1% to 0.5% by weight of capsaicinoids, from 7% to 23% by weight of a glycol, from 20% to 30% by weight of ethanol and from 56% to 65% by weight of water.
- 5 2. A solution according to claim 1 wherein said glycol comprises propylene glycol.
- 10 3. A solution according to claim 1 or claim 2 comprising from 0.1% to about 0.4% by weight of capsaicinoids.
- 15 4. A solution according to claim 3 comprising from about 0.15% to about 0.25% by weight of capsaicinoids.
- 20 5. A solution according to any one of claims 2 to 4 comprising from about 10% to about 20% by weight of propylene glycol.
- 25 6. A solution according to any one of the preceding claims comprising from about 26% to 30% by weight of ethanol.
- 30 7. A solution according to any one of the preceding claims comprising from 56% to about 60% by weight of water.
- 35 8. A solution according to any one of the preceding claims wherein the capsaicinoids comprise capsaicin, dihydrocapsaicin and nordihydrocapsaicin.
- 40 9. A lachrymator solution for use in a pressurized aerosol container that does not have a source of heat for vaporizing said solution, said solution comprising capsaicinoids and a nonflammable carrier suitable for use in a lachrymator, said carrier being liquid at ambient temperature and pressure and having a boiling point of less than about 110°C, and said solution comprises from 0.1% to 0.4% by weight of capsaicinoids, from 7% to 23% by weight of propylene glycol, from 20% to 30% by weight of ethanol and from 56% to 65% by weight of water.
- 45 10. A lachrymator solution according to claim 9 comprising from about 0.15% to about 0.25% by weight of capsaicinoids.
- 50 11. A lachrymator solution according to claim 9 or claim 10 comprising from about 10% to about 20% by weight of propylene glycol.
- 55 12. A lachrymator solution according to any one of claims 9 to 11 comprising from about 26% to 30%

by weight of ethanol.

13. A lachrymator solution according to any one of claims 9 to 12 comprising from 56% to about 60% by weight of water. 5

14. A lachrymator solution according to any one of claims 9 to 14 wherein the capsaicinoids comprise capsaicin, dihydrocapsaicin and nordihydrocapsaicin. 10

15. A lachrymator solution for use in a pressurized aerosol container wherein said solution comprises capsaicinoids and a nonflammable carrier suitable for use in the lachrymator, and said solution comprises from 0.15% to 0.25% by weight of capsaicinoids, from 10% to 20% by weight of propylene glycol, from 26% to 30% by weight of ethanol and from 56% to 60% by weight of water. 15

16. A lachrymator solution according to claim 15 wherein the capsaicinoids comprise capsaicin, dihydrocapsaicin and nordihydrocapsaicin. 20

Patentansprüche

1. Lösung für die Verwendung als ein Tränengas mit Capsaicinoiden und einem nichtbrennbaren Träger, wobei diese Lösung 0,1 bis 0,5 Gew.% Capsaicinoide, 7 bis 23 Gew.% Glycol, 20 bis 30 Gew.% Ethanol und 56 bis 65 Gew.% Wasser umfasst.
2. Lösung nach Anspruch 1, worin das Glycol Propylenglycol umfasst.
3. Lösung nach Anspruch 1 oder 2, die 0,1 bis etwa 0,4 Gew.% Capsaicinoide umfasst.
4. Lösung nach Anspruch 3, die etwa 0,15 bis 0,25 Gew.% Capsaicinoide umfasst.
5. Lösung nach einem der Ansprüche 2 bis 4, die etwa 10 bis 20 Gew.% Propylenglycol umfasst.
6. Lösung nach einem der vorausgehenden Ansprüche, die etwa 26 bis 30 Gew.% Ethanol umfasst.
7. Lösung nach einem der vorausgehenden Ansprüche, die 56 bis etwa 60 Gew.% Wasser umfasst.
8. Lösung nach einem der vorausgehenden Ansprüche, worin die Capsaicinoide Capsaicin, Dihydrocapsaicin und Nordihydrocapsaicin umfassen.
9. Tränengaslösung für die Verwendung in einem unter Druck stehenden Aerosolbehälter, der keine Heizquelle zum Verdampfen der Lösung hat, wobei

5 diese Lösung Capsaicinoide und einen nichtbrennbaren Träger umfasst, der für die Verwendung in einem Tränengas geeignet ist, wobei dieser Träger bei Umgebungstemperatur und -druck flüssig ist und einen Siedepunkt geringer als etwa 110 °C hat und die Lösung 0,1 bis 0,4 Gew.% Capsaicinoide, 7 bis 23 Gew.% Propylenglycol, 20 bis 30 Gew.% Ethanol und 56 bis 65 Gew.% Wasser umfasst.

10 10. Tränengaslösung nach Anspruch 9, die etwa 0,15 bis 0,25 Gew.% Capsaicinoide umfasst.

11. Tränengaslösung nach Anspruch 9 oder 10, die etwa 10 bis 20 Gew.% Propylenglycol umfasst. 15

12. Tränengaslösung nach einem der Ansprüche 9 bis 11, die etwa 26 bis 30 Gew.% Ethanol umfasst.

13. Tränengaslösung nach einem der Ansprüche 9 bis 20 12, die 56 bis etwa 60 Gew.% Wasser umfasst.

14. Tränengaslösung nach einem der Ansprüche 9 bis 25 13, worin die Capsaicinoide Capsaicin, Dihydrocapsaicin und Nordihydrocapsaicin umfassen.

15. Tränengaslösung für die Verwendung in einem unter Druck stehenden Aerosolbehälter, worin diese 30 Lösung Capsaicinoide und einen nichtbrennbaren Träger umfasst, der für die Verwendung in dem Tränengas geeignet ist und wobei diese Lösung 0,15 bis 0,25 Gew.% Capsaicinoide, 10 bis 20 Gew.% Propylenglycol, 26 bis 30 Gew.% Ethanol und 56 bis 60 Gew.% Wasser umfasst.

35 16. Tränengaslösung nach Anspruch 15, worin die Capsaicinoide Capsaicin, Dihydrocapsaicin und Nordihydrocapsaicin umfassen.

Revendications

1. Solution à utiliser en tant que gaz lacrymogène comprenant des capsaïcinoïdes et un véhicule ininflammable, ladite solution comprenant de 0,1 % à 45 0,5 % en poids de capsaïcinoïdes, de 7 % à 23 % en poids d'un glycol, de 20 % à 30 % en poids d'éthanol et de 56 % à 65 % en poids d'eau.
2. Solution selon la revendication 1, dans laquelle ledit glycol est constitué de propylèneglycol.
3. Solution selon la revendication 1 ou la revendication 2, comprenant de 0,1 % à environ 0,4 % en poids de capsaïcinoïdes. 55
4. Solution selon la revendication 3, comprenant environ 0,15 % à environ 0,25 % en poids de capsaïcinoïdes.

5. Solution selon l'une quelconque des revendications 2 à 4, comprenant environ 10 % à environ 20 % en poids de propylèneglycol. 5

6. Solution selon l'une quelconque des revendications précédentes, comprenant environ 26 % à 30 % en poids d'éthanol. 10

7. Solution selon l'une quelconque des revendications précédentes, comprenant de 56 % à environ 60 % en poids d'eau. 15

8. Solution selon l'une quelconque des revendications précédentes, dans laquelle les capsaïcinoïdes comprennent de la capsaïcine, de la dihydrocapsaïcine et de la nordihydrocapsaïcine. 20

9. Solution de gaz lacrymogène à utiliser dans un conteneur pressurisé pour aérosol qui n'a pas de source de chaleur pour vaporiser ladite solution, ladite solution comprenant des capsaïcinoïdes et un véhicule ininflammable approprié à une utilisation dans un gaz lacrymogène, ledit véhicule étant liquide à la température et à la pression ambiantes et ayant un point d'ébullition inférieur à environ 110°C, et ladite solution comprenant de 0,1 % à 0,4 % en poids de capsaïcinoïdes, de 7 % à 23 % en poids de propylèneglycol, de 20 % à 30 % en poids d'éthanol et de 56 % à 65 % en poids d'eau. 25

10. Solution de gaz lacrymogène selon la revendication 9, comprenant environ 0,15 % à environ 0,25 % en poids de capsaïcinoïdes. 30

11. Solution de gaz lacrymogène selon la revendication 9 ou la revendication 10, comprenant environ 10 % à environ 20 % en poids de propylèneglycol. 35

12. Solution de gaz lacrymogène selon l'une quelconque des revendications 9 à 11, comprenant environ 26 % à 30 % en poids d'éthanol. 40

13. Solution de gaz lacrymogène selon l'une quelconque des revendications 9 à 12, comprenant de 56 % à environ 60 % en poids d'eau. 45

14. Solution de gaz lacrymogène selon l'une quelconque des revendications 9 à 14, dans laquelle les capsaïcinoïdes comprennent de la capsaïcine, de la dihydrocapsaïcine et de la nordihydrocapsaïcine. 50

15. Solution de gaz lacrymogène à utiliser dans un conteneur pressurisé pour aérosol, ladite solution comprenant des capsaïcinoïdes et un véhicule ininflammable approprié à une utilisation dans le gaz lacrymogène, et ladite solution comprenant de 0,15 % à 0,25 % en poids de capsaïcinoïdes, de 10 % à 20 % en poids de propylèneglycol, de 26 % à 30 % en poids d'éthanol et de 56 % à 60 % en poids d'eau. 55

16. Solution de gaz lacrymogène selon la revendication 15, dans laquelle les capsaïcinoïdes comprennent de la capsaïcine, de la dihydrocapsaïcine et de la nordihydrocapsaïcine. 60