

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

07 December 2017 (07.12.2017)



(10) International Publication Number

WO 2017/207368 A1

(51) International Patent Classification:

A01P 3/00 (2006.01)

A01N 43/56 (2006.01)

A01N 43/653 (2006.01)

(21) International Application Number:

PCT/EP2017/062454

(22) International Filing Date:

23 May 2017 (23.05.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

16172693.0

02 June 2016 (02.06.2016)

EP

(71) Applicant: BASF AGRO B.V. [NL/NL]; Groningsingel 1, 6835 Arnhem (NL).

(72) Inventor: GEWEHR, Markus; Friedrich-Ebert-Strasse 18, 69502 Hemsbach (DE).

(74) Agent: BASF IP ASSOCIATION; BASF SE, G-FLP - C006, 67056 Ludwigshafen (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: FUNGICIDAL COMPOSITIONS

(57) Abstract: The present invention relates to compositions comprising, 1) as component I a compound selected from the following compounds I-1 to I-3, and 2) as component II a compound II-1 or II-2 and 3) as component III a compound selected from compounds III-1 to III-53. The invention furthermore relates to the use of the inventive compositions as pesticides in particular for controlling phytopathogenic fungi as detailed herein and preparations or compositions comprising them. The invention furthermore also relates to seed comprising the compositions. The invention furthermore also relates to methods for controlling pests, in particular phytopathogenic fungi as detailed herein, wherein the fungi or the materials, plants, the soil or seed to be protected from fungal attack are treated with an effective amount of a compositions according to the invention. The invention furthermore also relates to processes for preparing the compositions according to the invention.



WO 2017/207368 A1

FUNGICIDAL COMPOSITIONS

Description

The present invention relates to compositions comprising,

1) as component I a compound selected from the following compounds I-1 to I-3:

5 compound I-1: 1-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-cyclopropyl-2-(1,2,4-triazol-1-yl)ethanol;

compound I-2: 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-3-methyl-1-(1,2,4-triazol-1-yl)butan-2-ol;

10 compound I-3: 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1,2,4-triazol-1-yl)propan-2-ol;

or the N-oxides or agriculturally acceptable salts of each of the compounds; and

2) as component II the

compound II-1 pydiflumetofen, or

15 compound II-2 N-[(5-chloro-2-isopropyl-phenyl)methyl]-N-cyclopropyl-5-fluoro-1,3-dimethylpyrazole-4-carboxamide;

and

3) as component III, a compound selected from the following compounds III-1 to III-53 from the groups A) to H) and K):

A) Respiration inhibitors

- 20 - Inhibitors of complex III at Q_o site: azoxystrobin (III-1), fluoxastrobin (III-2), picoxystrobin (III-3), pyraclostrobin (III-4), trifloxystrobin (III-5), (1-[3-chloro-2-[[1-(4-chlorophenyl)pyrazol-3-yl]oxymethyl]phenyl]-4-methyl-tetrazol-5-one) (III-6), 1-[3-bromo-2-[[1-(4-chlorophenyl)pyrazol-3-yl]oxymethyl]phenyl]-4-methyl-tetrazol-5-one (III-7), 1-[2-[[1-(4-chlorophenyl)pyrazol-3-yl]oxymethyl]-3-methyl-phenyl]-4-methyl-tetrazol-5-one (III-8), 1-[2-[[1-(4-chlorophenyl)pyrazol-3-yl]oxymethyl]-3-fluoro-phenyl]-4-methyl-tetrazol-5-one (III-9), 1-[2-[[1-(2,4-dichlorophenyl)pyrazol-3-yl]oxymethyl]-3-fluoro-phenyl]-4-methyl-tetrazol-5-one (III-10), 1-[3-cyclopropyl-2-[[2-methyl-4-(1-methylpyrazol-3-yl)phenoxy]methyl]phenyl]-4-methyl-tetrazol-5-one (III-11), 1-[3-(difluoromethoxy)-2-[[2-methyl-4-(1-methylpyrazol-3-yl)phenoxy]methyl]phenyl]-4-methyl-tetrazol-5-one (III-12), 1-methyl-4-[3-methyl-2-[[2-methyl-4-(1-methylpyrazol-3-yl)phenoxy]methyl]phenyl]tetrazol-5-one (III-13), (Z,2E)-5-[1-(2,4-dichlorophenyl)pyrazol-3-yl]-oxy-2-methoxyimino-N,3-dimethyl-pent-3-enamide (III-14), (Z,2E)-5-[1-(4-chlorophenyl)pyrazol-3-yl]oxy-2-methoxyimino-N,3-dimethyl-pent-3-enamide (III-15);
- inhibitors of complex II: benzovindiflupyr (III-16), bixafen (III-17), boscalid (III-18), fluxapyroxad (III-19), isopyrazam (III-20);

- 35 - other respiration inhibitors: ametocetradin (III-21);

B) Sterol biosynthesis inhibitors (SBI fungicides)

- C14 demethylase inhibitors: triazoles: difenoconazole (III-22), epoxiconazole (III-23), fluquinconazole (III-24), metconazole (III-25), propiconazole (III-26), prothioconazole (III-27), tebuconazole (III-28); imidazoles: prochloraz (III-29);
- Delta14-reductase inhibitors: fenpropimorph (III-30);
- 5 C) Nucleic acid synthesis inhibitors
 - phenylamides or acyl amino acid fungicides: benalaxyl (III-31), metalaxyl (III-32);
- D) Inhibitors of cell division and cytoskeleton
 - tubulin inhibitors: carbendazim (III-33), thiophanate-methyl (III-34);
 - other cell division inhibitors: metrafenone (III-35);
- 10 E) Inhibitors of amino acid and protein synthesis
 - methionine synthesis inhibitors: cyprodinil (III-36), pyrimethanil (III-37);
- F) Signal transduction inhibitors
 - MAP / histidine kinase inhibitors: fluoroimide (III-38), iprodione (III-39), vinclozolin (III-40), fludioxonil (III-41);
- 15 G) Lipid and membrane synthesis inhibitors
 - phospholipid biosynthesis and cell wall deposition: dimethomorph (III-42), mandipropamid (III-43), iprovalicarb (III-44);
 - inhibitors of oxysterol binding protein: oxathiapiprolin (III-45), 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}phenyl
- 20 methanesulfonate (III-46), 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl) 1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate (III-47);
- H) Inhibitors with Multi Site Action
 - thio- and dithiocarbamates: mancozeb (III-48), metiram (III-49);
 - organochlorine compounds: chlorothalonil (III-50);
- 25 - guanidines and others: dithianon (III-51); and
- K) Unknown mode of action

picarbutrazox (III-52), 3-(4,4-difluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline (III-53);

or the N-oxides or agriculturally acceptable salts of each of the compounds.
- 30 The invention furthermore relates to the use of the inventive compositions as pesticides in particular for controlling phytopathogenic fungi as detailed herein and preparations or compositions comprising them. The invention furthermore also relates to seed comprising the compositions. The invention furthermore also relates to methods for controlling pests, in particular phytopathogenic fungi as detailed herein, wherein the fungi or the materials, plants,
- 35 the soil or seed to be protected from fungal attack are treated with an effective amount of a composition according to the invention. The invention furthermore also relates to processes for preparing the compositions according to the invention.


With a view to reducing the application rates and broadening the activity spectrum of the known compounds, it was an object of the present invention to provide compositions which, at a reduced total amount of active compounds applied, show improved activity against important pests, in particular phytopathogenic fungi, in particular for certain indications. It was a further object to provide for compositions that are useful for the control of specific pathogens in specific important crops that are often susceptible to the attack of pathogens.

Accordingly we have found the compositions and uses defined at the outset and in the following description.


The compounds I (component I), its preparation and use in crop protection are described in WO 2013/007767, which also disclose certain compositions with other active compounds. Owing to the basic character of their nitrogen atoms, the compounds I-1, I-2 and I-3 are capable of forming salts or adducts with inorganic or organic acids or with metal ions, in particular salts with inorganic acids.

The structures are as follows:


I-1: 1-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-cyclopropyl-2-(1,2,4-triazol-1-yl)ethanol



I-2: 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-3-methyl-1-(1,2,4-triazol-1-yl)butan-2-ol



I-3: 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1,2,4-triazol-1-yl)propan-2-ol



In each case, the respective N-oxide and the agriculturally acceptable salts of the compounds as further defined below are encompassed.

Compounds I-1 to I-3 comprise chiral centers and they are generally obtained in the form of racemates. The R- and S-enantiomers of the compounds contained as component I in the compositions according to the invention can be separated and isolated in pure form with methods known by the skilled person, e.g. by using chiral HPLC. Suitable for use in the compositions are both the enantiomers and compositions thereof. Furthermore, said components I can be present in different crystal modifications, which may differ in biological activity.

In particular, in each case, a racemic composition of the respective components I, namely of compound I-1, I-2 or I-3, respectively, is present. Furthermore, any other proportions of the (R)-

enantiomer and the (S)-enantiomer of the respective compound I-1, I-2 or I-3, respectively, may be present according to the present invention. For example, the (R)-enantiomer of compound I-3 is (R)-2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1,2,4-triazol-1-yl)propan-2-ol; the S-enantiomer of I-3 is (S)-2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1,2,4-triazol-1-yl)propan-2-ol. This applies to the other compounds accordingly. According to one specific embodiment, the respective compound I-1, I-2 or I-3, respectively, is provided and used as (R)-enantiomer with an enantiomeric excess (e.e.) of at least 40%, for example, at least 50%, 60%, 70% or 80%, preferably at least 90%, more preferably at least 95%, yet more preferably at least 98% and most preferably at least 99%. According to a further specific embodiment, the respective compound I-1, I-2 or I-3, respectively, is provided and used as (S)-enantiomer with an enantiomeric excess (e.e.) of at least 40%, for example, at least 50%, 60%, 70% or 80%, preferably at least 90%, more preferably at least 95%, yet more preferably at least 98% and most preferably at least 99%. This applies to every composition detailed herein.

According to one embodiment of the invention, component I is compound I-1.

According to a further embodiment of the invention, component I is compound I-2.


According to a further embodiment of the invention, component I is compound I-3.

Component II is compound II-1 or compound II-2.

According to one embodiment of the invention, component II is compound II-1.


According to a further embodiment of the invention, component II is compound II-2.

Compound II-1 is pydiflumetofen, i.e. 3-(difluoromethyl)-N-methoxy-1-methyl-N-[1-methyl-2-(2,4,6-trichlorophenyl)ethyl]pyrazole-4-carboxamide



The preparation, the pesticidal action and possible uses of compound II-1 can be found in WO2010/063700, WO2013/167651, WO2013/127764, WO2014/206855, WO2014/023628, WO2014/016279 and WO 2015/124543.

Compound II-1 is N-[(5-chloro-2-isopropyl-phenyl)methyl]-N-cyclopropyl-5-fluoro-1,3-dimethylpyrazole-4-carboxamide, having the following structure:



The preparation, the pesticidal action and possible uses for compound II-2 can be found in WO2012/143127, WO2014/060502, WO2014/060521 and WO2014/076007

Component III is a compound selected from the compounds III-1 to III-53.

- 5 The compounds having common names can be found for example in the e-Pesticide Manual V5.2 (ISBN 978 1 901396 85 0) (2008-2011) and other publications (cf.: <http://www.alanwood.net/pesticides/>). They are mainly commercially available. Compounds III-6 to III-13, their preparation and uses are e.g. known from WO 2013/162072, WO 2013/162072, WO 2013/162072, WO 2013/162072, WO 2013/162072, WO 2013/162077, WO 2013/162077, 10 WO 2014/051161, WO 2014/051161, WO 2014/051165 and WO 2014/084223. Compounds III-14 and III-15, their preparation and uses are e.g. known from WO 13/092224. Compounds III-46 and III-47, their preparation and uses are e.g. known from WO 2008/013622. Compounds III-53 its preparation and uses are e.g. known from WO 2005/070917 and WO 2011/77514.

- 15 According to one embodiment of the invention, component III is selected from compounds III-6 to III-13.

According to a further embodiment of the invention, component III is selected from compounds III-14 and III-15.

According to a further embodiment of the invention, component III is selected from compounds III-46 and III-47.

- 20 According to a further embodiment of the invention, component III is selected from compounds III-1 to III-5, III-16 to III-45 and III-48 to III-52.

- According to one particular embodiment, component III is III-1. According to a further particular embodiment, component III is III-2. According to still a further embodiment, component III is III-3. According to still a further embodiment, component III is III-4. According to still a further 25 embodiment, component III is III-5. According to still a further embodiment, component III is III-6. According to still a further embodiment, component III is III-7. According to still a further embodiment, component III is III-8. According to still a further embodiment, component III is III-9. According to still a further embodiment, component III is III-10. According to still a further embodiment, component III is III-11. According to still a further embodiment, component III is III- 30 12. According to still a further embodiment, component III is III-13. According to still a further embodiment, component III is III-14. According to still a further embodiment, component III is III-15. According to still a further embodiment, component III is III-16. According to still a further embodiment, component III is III-17. According to still a further embodiment, component III is III-18. According to still a further embodiment, component III is III-19. According to still a further

embodiment, component III is III-20. According to still a further embodiment, component III is III-21. According to still a further embodiment, component III is III-22. According to still a further embodiment, component III is III-23. According to still a further embodiment, component III is III-24. According to still a further embodiment, component III is III-25. According to still a further embodiment, component III is III-26. According to still a further embodiment, component III is III-27. According to still a further embodiment, component III is III-28. According to still a further embodiment, component III is III-29. According to still a further embodiment, component III is III-30. According to a further particular embodiment, component III is III-31. According to a further embodiment, component III is III-32. According to still a further embodiment, component III is III-33. According to still a further embodiment, component III is III-34. According to still a further embodiment, component III is III-35. According to still a further embodiment, component III is III-36. According to one specific embodiment, component III is III-37. According to a further embodiment, component III is III-38. According to still a further embodiment, component III is III-39. According to one specific embodiment, component III is III-40. According to a further embodiment, component III is III-41. According to still a further embodiment, component III is III-42. According to still a further embodiment, component III is III-43. According to still a further embodiment, component III is III-44. According to still a further embodiment, component III is III-45. According to still a further embodiment, component III is III-46. According to still a further embodiment, component III is III-47. According to still a further embodiment, component III is III-48. According to still a further embodiment, component III is III-49. According to still a further embodiment, component III is III-50. According to still a further embodiment, component III is III-51. According to still a further embodiment, component III is III-52. According to still a further embodiment, component III is III-53.

Examples of inorganic acids are hydrohalic acids, such as hydrogen fluoride, hydrogen chloride, hydrogen bromide and hydrogen iodide, carbonic acid, sulfuric acid, phosphoric acid and nitric acid.

Suitable organic acids are, for example, formic acid and alkanoic acids, such as acetic acid, trifluoroacetic acid, trichloroacetic acid and propionic acid, and also glycolic acid, thiocyanic acid, lactic acid, succinic acid, citric acid, benzoic acid and other arylcarboxylic acids, cinnamic acid, oxalic acid, alkylsulfonic acids (sulfonic acids having straight-chain or branched alkyl radicals of 1 to 20 carbon atoms), arylsulfonic acids or aryldisulfonic acids (aromatic radicals, such as phenyl and naphthyl, which carry one or two sulfonic acid groups), alkylphosphonic acids (phosphonic acids having straight-chain or branched alkyl radicals with 1 to 20 carbon atoms), arylphosphonic acids or aryldiphosphonic acids (aromatic radicals, such as phenyl and naphthyl, which carry one or two phosphoric acid radicals), where the alkyl or aryl radicals may carry further substituents, for example p-toluenesulfonic acid, salicylic acid, p-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid etc. Suitable metal ions are in particular the ions of the elements of the second main group, in particular calcium and magnesium, of the third and fourth main group, in particular aluminum, tin and lead, and also of the elements of transition groups one to eight, in particular chromium, manganese, iron, cobalt, nickel, copper, zinc, and others. Particular preference is given to the metal ions of the elements of transition

groups of the fourth period. The metals can be present in the various valencies that they can assume.

In the following, the inventive compositions and their preferred uses are further described. In each case, according to the present invention, the use of the composition for controlling a particular phytopathogenic fungus is also meant to encompass the respective method for controlling the particular phytopathogenic fungi, wherein the fungi or the materials, plants, the soil or seed to be protected from fungal attack are treated with an effective amount of a composition as defined in that particular context.

According to a specific embodiment thereof, exactly three active compounds as defined are present in these ternary compositions. The composition may, of course, contain any kind of additive or the like as detailed herein in order to provide a formulation suitable for use in agriculture.

In the inventive three-component-compositions, the weight ratio of component I to the 1st further active compound (component II) depends on the properties of the active compounds in question and may particularly be 1000:1 to 1:1000, specifically 500:1 to 1:500. Preferably, it is in the range of from 1:100 to 100:1, preferably in the range of from 1:50 to 50:1 and in particular in the range of from 1:20 to 20:1. It may be preferable for the weight ratio to be in the region of from 1:10 to 10:1, preferably from 1:3 to 3:1, in particular from 1:2 to 2:1. The weight ratio of component I to the 2nd further active compound (component III) may particularly be 1000:1 to 1:1000, specifically 500:1 to 1:500. It is preferably in the range of from 1:100 to 100:1, preferably in the range of from 1:50 to 50:1 and in particular in the range of from 1:20 to 20:1. It may be preferable for the weight ratio to be in the region of from 1:10 to 10:1, preferably from 1:3 to 3:1, in particular from 1:2 to 2:1. The weight ratio of 1st further active compound (component II) to 2nd further active compound (component III) is preferably in the range of from 1:100 to 100:1, frequently in the range of from 1:50 to 50:1, preferably in the range of from 1:20 to 20:1, and in particular in the range of from 1:10 to 10:1. It may be preferable for the weight to be in the range of from 1:3 to 3:1, in particular from 1:2 to 2:1.

Three-component compositions of the invention are compiled in Table T1, wherein each row corresponds to one embodiment of the compositions according to the invention, i.e. one specific individualized composition. According to one specific aspect, these are ternary compositions, wherein each one only contains these three components as active compounds. Furthermore, also every combination of the compositions individualized in this table represent embodiments of the present invention.

Table T1: Three-component compositions T1-1 to T1-318 comprising one component I, component II and one component III. According to the respective specific embodiments, these compositions contain the respective components I, II and III as only pesticidally active ingredients:

I = component I selected from compounds I-1, I-2 and I-3;

II = component II is compound II-1 or compound II-2;

III = component III selected from III-1 to III-53 as defined above

composition	I	II	III
T1-1	I-1	II-1	III-1
T1-2	I-1	II-1	III-2
T1-3	I-1	II-1	III-3
T1-4	I-1	II-1	III-4
T1-5	I-1	II-1	III-5
T1-6	I-1	II-1	III-6
T1-7	I-1	II-1	III-7
T1-8	I-1	II-1	III-8
T1-9	I-1	II-1	III-9
T1-10	I-1	II-1	III-10
T1-11	I-1	II-1	III-11
T1-12	I-1	II-1	III-12
T1-13	I-1	II-1	III-13
T1-14	I-1	II-1	III-14
T1-15	I-1	II-1	III-15
T1-16	I-1	II-1	III-16
T1-17	I-1	II-1	III-17
T1-18	I-1	II-1	III-18
T1-19	I-1	II-1	III-19
T1-20	I-1	II-1	III-20
T1-21	I-1	II-1	III-21
T1-22	I-1	II-1	III-22
T1-23	I-1	II-1	III-23
T1-24	I-1	II-1	III-24
T1-25	I-1	II-1	III-25
T1-26	I-1	II-1	III-26
T1-27	I-1	II-1	III-27
T1-28	I-1	II-1	III-28
T1-29	I-1	II-1	III-29
T1-30	I-1	II-1	III-30
T1-31	I-1	II-1	III-31
T1-32	I-1	II-1	III-32
T1-33	I-1	II-1	III-33
T1-34	I-1	II-1	III-34
T1-35	I-1	II-1	III-35
T1-36	I-1	II-1	III-36
T1-37	I-1	II-1	III-37
T1-38	I-1	II-1	III-38

composition	I	II	III
T1-39	I-1	II-1	III-39
T1-40	I-1	II-1	III-40
T1-41	I-1	II-1	III-41
T1-42	I-1	II-1	III-42
T1-43	I-1	II-1	III-43
T1-44	I-1	II-1	III-44
T1-45	I-1	II-1	III-45
T1-46	I-1	II-1	III-46
T1-47	I-1	II-1	III-47
T1-48	I-1	II-1	III-48
T1-49	I-1	II-1	III-49
T1-50	I-1	II-1	III-50
T1-51	I-1	II-1	III-51
T1-52	I-1	II-1	III-52
T1-53	I-1	II-1	III-53
T1-54	I-1	II-2	III-1
T1-55	I-1	II-2	III-2
T1-56	I-1	II-2	III-3
T1-57	I-1	II-2	III-4
T1-58	I-1	II-2	III-5
T1-59	I-1	II-2	III-6
T1-60	I-1	II-2	III-7
T1-61	I-1	II-2	III-8
T1-62	I-1	II-2	III-9
T1-63	I-1	II-2	III-10
T1-64	I-1	II-2	III-11
T1-65	I-1	II-2	III-12
T1-66	I-1	II-2	III-13
T1-67	I-1	II-2	III-14
T1-68	I-1	II-2	III-15
T1-69	I-1	II-2	III-16
T1-70	I-1	II-2	III-17
T1-71	I-1	II-2	III-18
T1-72	I-1	II-2	III-19
T1-73	I-1	II-2	III-20
T1-74	I-1	II-2	III-21
T1-75	I-1	II-2	III-22
T1-76	I-1	II-2	III-23

composition	I	II	III
T1-77	I-1	II-2	III-24
T1-78	I-1	II-2	III-25
T1-79	I-1	II-2	III-26
T1-80	I-1	II-2	III-27
T1-81	I-1	II-2	III-28
T1-82	I-1	II-2	III-29
T1-83	I-1	II-2	III-30
T1-84	I-1	II-2	III-31
T1-85	I-1	II-2	III-32
T1-86	I-1	II-2	III-33
T1-87	I-1	II-2	III-34
T1-88	I-1	II-2	III-35
T1-89	I-1	II-2	III-36
T1-90	I-1	II-2	III-37
T1-91	I-1	II-2	III-38
T1-92	I-1	II-2	III-39
T1-93	I-1	II-2	III-40
T1-94	I-1	II-2	III-41
T1-95	I-1	II-2	III-42
T1-96	I-1	II-2	III-43
T1-97	I-1	II-2	III-44
T1-98	I-1	II-2	III-45
T1-99	I-1	II-2	III-46
T1-100	I-1	II-2	III-47
T1-101	I-1	II-2	III-48
T1-102	I-1	II-2	III-49
T1-103	I-1	II-2	III-50
T1-104	I-1	II-2	III-51
T1-105	I-1	II-2	III-52
T1-106	I-1	II-2	III-53
T1-107	I-2	II-1	III-1
T1-108	I-2	II-1	III-2
T1-109	I-2	II-1	III-3
T1-110	I-2	II-1	III-4
T1-111	I-2	II-1	III-5
T1-112	I-2	II-1	III-6
T1-113	I-2	II-1	III-7
T1-114	I-2	II-1	III-8

compo- sition	I	II	III
T1-115	I-2	II-1	III-9
T1-116	I-2	II-1	III-10
T1-117	I-2	II-1	III-11
T1-118	I-2	II-1	III-12
T1-119	I-2	II-1	III-13
T1-120	I-2	II-1	III-14
T1-121	I-2	II-1	III-15
T1-122	I-2	II-1	III-16
T1-123	I-2	II-1	III-17
T1-124	I-2	II-1	III-18
T1-125	I-2	II-1	III-19
T1-126	I-2	II-1	III-20
T1-127	I-2	II-1	III-21
T1-128	I-2	II-1	III-22
T1-129	I-2	II-1	III-23
T1-130	I-2	II-1	III-24
T1-131	I-2	II-1	III-25
T1-132	I-2	II-1	III-26
T1-133	I-2	II-1	III-27
T1-134	I-2	II-1	III-28
T1-135	I-2	II-1	III-29
T1-136	I-2	II-1	III-30
T1-137	I-2	II-1	III-31
T1-138	I-2	II-1	III-32
T1-139	I-2	II-1	III-33
T1-140	I-2	II-1	III-34
T1-141	I-2	II-1	III-35
T1-142	I-2	II-1	III-36
T1-143	I-2	II-1	III-37
T1-144	I-2	II-1	III-38
T1-145	I-2	II-1	III-39
T1-146	I-2	II-1	III-40
T1-147	I-2	II-1	III-41
T1-148	I-2	II-1	III-42
T1-149	I-2	II-1	III-43
T1-150	I-2	II-1	III-44
T1-151	I-2	II-1	III-45
T1-152	I-2	II-1	III-46
T1-153	I-2	II-1	III-47
T1-154	I-2	II-1	III-48

compo- sition	I	II	III
T1-155	I-2	II-1	III-49
T1-156	I-2	II-1	III-50
T1-157	I-2	II-1	III-51
T1-158	I-2	II-1	III-52
T1-159	I-2	II-1	III-53
T1-160	I-2	II-2	III-1
T1-161	I-2	II-2	III-2
T1-162	I-2	II-2	III-3
T1-163	I-2	II-2	III-4
T1-164	I-2	II-2	III-5
T1-165	I-2	II-2	III-6
T1-166	I-2	II-2	III-7
T1-167	I-2	II-2	III-8
T1-168	I-2	II-2	III-9
T1-169	I-2	II-2	III-10
T1-170	I-2	II-2	III-11
T1-171	I-2	II-2	III-12
T1-172	I-2	II-2	III-13
T1-173	I-2	II-2	III-14
T1-174	I-2	II-2	III-15
T1-175	I-2	II-2	III-16
T1-176	I-2	II-2	III-17
T1-177	I-2	II-2	III-18
T1-178	I-2	II-2	III-19
T1-179	I-2	II-2	III-20
T1-180	I-2	II-2	III-21
T1-181	I-2	II-2	III-22
T1-182	I-2	II-2	III-23
T1-183	I-2	II-2	III-24
T1-184	I-2	II-2	III-25
T1-185	I-2	II-2	III-26
T1-186	I-2	II-2	III-27
T1-187	I-2	II-2	III-28
T1-188	I-2	II-2	III-29
T1-189	I-2	II-2	III-30
T1-190	I-2	II-2	III-31
T1-191	I-2	II-2	III-32
T1-192	I-2	II-2	III-33
T1-193	I-2	II-2	III-34
T1-194	I-2	II-2	III-35

compo- sition	I	II	III
T1-195	I-2	II-2	III-36
T1-196	I-2	II-2	III-37
T1-197	I-2	II-2	III-38
T1-198	I-2	II-2	III-39
T1-199	I-2	II-2	III-40
T1-200	I-2	II-2	III-41
T1-201	I-2	II-2	III-42
T1-202	I-2	II-2	III-43
T1-203	I-2	II-2	III-44
T1-204	I-2	II-2	III-45
T1-205	I-2	II-2	III-46
T1-206	I-2	II-2	III-47
T1-207	I-2	II-2	III-48
T1-208	I-2	II-2	III-49
T1-209	I-2	II-2	III-50
T1-210	I-2	II-2	III-51
T1-211	I-2	II-2	III-52
T1-212	I-2	II-2	III-53
T1-213	I-3	II-1	III-1
T1-214	I-3	II-1	III-2
T1-215	I-3	II-1	III-3
T1-216	I-3	II-1	III-4
T1-217	I-3	II-1	III-5
T1-218	I-3	II-1	III-6
T1-219	I-3	II-1	III-7
T1-220	I-3	II-1	III-8
T1-221	I-3	II-1	III-9
T1-222	I-3	II-1	III-10
T1-223	I-3	II-1	III-11
T1-224	I-3	II-1	III-12
T1-225	I-3	II-1	III-13
T1-226	I-3	II-1	III-14
T1-227	I-3	II-1	III-15
T1-228	I-3	II-1	III-16
T1-229	I-3	II-1	III-17
T1-230	I-3	II-1	III-18
T1-231	I-3	II-1	III-19
T1-232	I-3	II-1	III-20
T1-233	I-3	II-1	III-21
T1-234	I-3	II-1	III-22

composition	I	II	III
T1-235	I-3	II-1	III-23
T1-236	I-3	II-1	III-24
T1-237	I-3	II-1	III-25
T1-238	I-3	II-1	III-26
T1-239	I-3	II-1	III-27
T1-240	I-3	II-1	III-28
T1-241	I-3	II-1	III-29
T1-242	I-3	II-1	III-30
T1-243	I-3	II-1	III-31
T1-244	I-3	II-1	III-32
T1-245	I-3	II-1	III-33
T1-246	I-3	II-1	III-34
T1-247	I-3	II-1	III-35
T1-248	I-3	II-1	III-36
T1-249	I-3	II-1	III-37
T1-250	I-3	II-1	III-38
T1-251	I-3	II-1	III-39
T1-252	I-3	II-1	III-40
T1-253	I-3	II-1	III-41
T1-254	I-3	II-1	III-42
T1-255	I-3	II-1	III-43
T1-256	I-3	II-1	III-44
T1-257	I-3	II-1	III-45
T1-258	I-3	II-1	III-46
T1-259	I-3	II-1	III-47
T1-260	I-3	II-1	III-48
T1-261	I-3	II-1	III-49
T1-262	I-3	II-1	III-50

composition	I	II	III
T1-263	I-3	II-1	III-51
T1-264	I-3	II-1	III-52
T1-265	I-3	II-1	III-53
T1-266	I-3	II-2	III-1
T1-267	I-3	II-2	III-2
T1-268	I-3	II-2	III-3
T1-269	I-3	II-2	III-4
T1-270	I-3	II-2	III-5
T1-271	I-3	II-2	III-6
T1-272	I-3	II-2	III-7
T1-273	I-3	II-2	III-8
T1-274	I-3	II-2	III-9
T1-275	I-3	II-2	III-10
T1-276	I-3	II-2	III-11
T1-277	I-3	II-2	III-12
T1-278	I-3	II-2	III-13
T1-279	I-3	II-2	III-14
T1-280	I-3	II-2	III-15
T1-281	I-3	II-2	III-16
T1-282	I-3	II-2	III-17
T1-283	I-3	II-2	III-18
T1-284	I-3	II-2	III-19
T1-285	I-3	II-2	III-20
T1-286	I-3	II-2	III-21
T1-287	I-3	II-2	III-22
T1-288	I-3	II-2	III-23
T1-289	I-3	II-2	III-24
T1-290	I-3	II-2	III-25

composition	I	II	III
T1-291	I-3	II-2	III-26
T1-292	I-3	II-2	III-27
T1-293	I-3	II-2	III-28
T1-294	I-3	II-2	III-29
T1-295	I-3	II-2	III-30
T1-296	I-3	II-2	III-31
T1-297	I-3	II-2	III-32
T1-298	I-3	II-2	III-33
T1-299	I-3	II-2	III-34
T1-300	I-3	II-2	III-35
T1-301	I-3	II-2	III-36
T1-302	I-3	II-2	III-37
T1-303	I-3	II-2	III-38
T1-304	I-3	II-2	III-39
T1-305	I-3	II-2	III-40
T1-306	I-3	II-2	III-41
T1-307	I-3	II-2	III-42
T1-308	I-3	II-2	III-43
T1-309	I-3	II-2	III-44
T1-310	I-3	II-2	III-45
T1-311	I-3	II-2	III-46
T1-312	I-3	II-2	III-47
T1-313	I-3	II-2	III-48
T1-314	I-3	II-2	III-49
T1-315	I-3	II-2	III-50
T1-316	I-3	II-2	III-51
T1-317	I-3	II-2	III-52
T1-318	I-3	II-2	III-53

One particular embodiment relates to the compositions T1-213 to T1-318. One more specific embodiment relates to the compositions T1-213 to T1-265, in particular selected from T-213, T-216, T-218, T-219, T-220, T-221, T-222, T-223, T-224, T-225, T-226, T-227, T-237, T-269, T-231 and T-234. Another more specific embodiment relates to compositions T1-266 to T1-318, in particular selected from T-237, T-269, T-231, T-234, T-267, T-270, T-271, T-272, T-273, T-274, T-275, T-276, T-277, T-278, T-279, T-280, T-284, T-292 and T-318. One further particular embodiment relates to a composition selected from T-213, T-216, T-218, T-219, T-220, T-221, T-222, T-223, T-224, T-225, T-226, T-227, T-237, T-269, T-231, T-234, T-267, T-270, T-271, T-272, T-273, T-274, T-275, T-276, T-277, T-278, T-279, T-280, T-284, T-292 and T-318.

- 10 One embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 1000:1 to 1:1000.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 500:1 to 1:500.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 100:1 to 1:100.

- 5 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 1:50 to 50:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 1:20 to 20:1.

- 10 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 1:10 to 10:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 1:3 to 3:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 1:2 to 2:1.

- 15 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 1000:1 to 1:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 100:1 to 1:1.

- 20 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 50:1 to 1:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 20:1 to 1:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 10:1 to 1:1.

- 25 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 4:1 to 1:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 2:1 to 1:1.

- 30 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 1:1 to 1:1000.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 1:1 to 1:100.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 1:1 to 1:50.

- 35 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 1:1 to 1:20.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 1:1 to 1:10.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 1:1 to 1:4.

- 5 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component II is from 1:1 to 1:2.

One embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 1000:1 to 1:1000.

- 10 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 500:1 to 1:500.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 100:1 to 1:100.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 1:50 to 50:1.

- 15 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 1:20 to 20:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 1:10 to 10:1.

- 20 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 1:3 to 3:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 1:2 to 2:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 1000:1 to 1:1.

- 25 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 100:1 to 1:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 50:1 to 1:1.

- 30 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 20:1 to 1:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 10:1 to 1:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 4:1 to 1:1.

- 35 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 2:1 to 1:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 1:1 to 1:1000.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 1:1 to 1:100.

- 5 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 1:1 to 1:50.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 1:1 to 1:20.

- 10 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 1:1 to 1:10.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 1:1 to 1:4.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component I to component III is from 1:1 to 1:2.

- 15 One embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 1000:1 to 1:1000.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 500:1 to 1:500.

- 20 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 100:1 to 1:100.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 1:50 to 50:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 1:20 to 20:1.

- 25 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 1:10 to 10:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 1:3 to 3:1.

- 30 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 1:2 to 2:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 1000:1 to 1:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 100:1 to 1:1.

- 35 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 50:1 to 1:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 20:1 to 1:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 10:1 to 1:1.

- 5 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 4:1 to 1:1.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 2:1 to 1:1.

- 10 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 1:1 to 1:1000.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 1:1 to 1:100.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 1:1 to 1:50.

- 15 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 1:1 to 1:20.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 1:1 to 1:10.

- 20 One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 1:1 to 1:4.

One further embodiment of the invention relates to any one of compositions T1-1 to T1-318 wherein the weight ratio of component II to component III is from 1:1 to 1:2.

- 25 The mixtures and compositions according to the invention are suitable as fungicides. They are distinguished by an outstanding effectiveness against a broad spectrum of phytopathogenic fungi, including soil-borne fungi, which derive especially from the classes of the Plasmodiophoromycetes, Peronosporomycetes (syn. Oomycetes), Chytridiomycetes, Zygomycetes, Ascomycetes, Basidiomycetes and Deuteromycetes (syn. Fungi imperfecti). Some are systemically effective and they can be used in crop protection as foliar fungicides, fungicides for seed dressing and soil fungicides. Moreover, they are suitable for controlling
30 harmful fungi, which inter alia occur in wood or roots of plants.

- The mixtures and compositions according to the invention are particularly important in the control of a multitude of phytopathogenic fungi on various cultivated plants, such as cereals, e. g. wheat, rye, barley, triticale, oats or rice; beet, e. g. sugar beet or fodder beet; fruits, such as pomes, stone fruits or soft fruits, e. g. apples, pears, plums, peaches, almonds, cherries,
35 strawberries, raspberries, blackberries or gooseberries; leguminous plants, such as lentils, peas, alfalfa or soybeans; oil plants, such as rape, mustard, olives, sunflowers, coconut, cocoa beans, castor oil plants, oil palms, ground nuts or soybeans; cucurbits, such as squashes, cucumber or melons; fiber plants, such as cotton, flax, hemp or jute; citrus fruit, such as oranges, lemons, grapefruits or mandarins; vegetables, such as spinach, lettuce, asparagus,

cabbages, carrots, onions, tomatoes, potatoes, cucurbits or paprika; lauraceous plants, such as avocados, cinnamon or camphor; energy and raw material plants, such as corn, soybean, rape, sugar cane or oil palm; corn; tobacco; nuts; coffee; tea; bananas; vines (table grapes and grape juice grape vines); hop; turf; natural rubber plants or ornamental and forestry plants, such as flowers, shrubs, broad-leaved trees or evergreens, e. g. conifers; and on the plant propagation material, such as seeds, and the crop material of these plants.

Preferably the inventive mixtures and compositions are used for controlling a multitude of fungi on field crops, such as potatoes sugar beets, tobacco, wheat, rye, barley, oats, rice, corn, cotton, soybeans, rape, legumes, sunflowers, coffee or sugar cane; fruits; vines; ornamentals; or vegetables, such as cucumbers, tomatoes, beans or squashes.

The term "plant propagation material" is to be understood to denote all the generative parts of the plant such as seeds and vegetative plant material such as cuttings and tubers (e. g. potatoes), which can be used for the multiplication of the plant. This includes seeds, roots, fruits, tubers, bulbs, rhizomes, shoots, sprouts and other parts of plants, including seedlings and young plants, which are to be transplanted after germination or after emergence from soil. These young plants may also be protected before transplantation by a total or partial treatment by immersion or pouring.

Preferably, treatment of plant propagation materials with the inventive combination of component I, component II and component III, and compositions thereof, respectively, is used for controlling a multitude of fungi on cereals, such as wheat, rye, barley and oats; rice, corn, cotton and soybeans.

The term "cultivated plants" is to be understood as including plants which have been modified by breeding, mutagenesis or genetic engineering including but not limiting to agricultural biotech products on the market or in development (cf. <http://cera-gmc.org/>, see GM crop database therein). Genetically modified plants are plants, which genetic material has been so modified by the use of recombinant DNA techniques that under natural circumstances cannot readily be obtained by cross breeding, mutations or natural recombination. Typically, one or more genes have been integrated into the genetic material of a genetically modified plant in order to improve certain properties of the plant. Such genetic modifications also include but are not limited to targeted post-translational modification of protein(s), oligo- or polypeptides e. g. by glycosylation or polymer additions such as prenylated, acetylated or farnesylated moieties or PEG moieties.

Plants that have been modified by breeding, mutagenesis or genetic engineering, e. g. have been rendered tolerant to applications of specific classes of herbicides, such as hydroxyphenylpyruvate dioxygenase (HPPD) inhibitors; acetolactate synthase (ALS) inhibitors, such as sulfonyl ureas (see e. g. US 6,222,100, WO 01/82685, WO 00/26390, WO 97/41218, WO 98/02526, WO 98/02527, WO 04/106529, WO 05/20673, WO 03/14357, WO 03/13225, WO 03/14356, WO 04/16073) or imidazolinones (see e. g. US 6,222,100, WO 01/82685, WO 00/26390, WO 97/41218, WO 98/002526, WO 98/02527, WO 04/106529, WO 05/20673, WO 03/014357, WO 03/13225, WO 03/14356, WO 04/16073); enolpyruvylshikimate-3-phosphate synthase (EPSPS) inhibitors, such as glyphosate (see e. g. WO 92/00377); glutamine synthetase (GS) inhibitors, such as glufosinate (see e. g. EP-A 242 236, EP-A 242 246) or oxynil herbicides (see

e. g. US 5,559,024) as a result of conventional methods of breeding or genetic engineering. Several cultivated plants have been rendered tolerant to herbicides by conventional methods of breeding (mutagenesis), e. g. Clearfield® summer rape (Canola, BASF SE, Germany) being tolerant to imidazolinones, e. g. imazamox. Genetic engineering methods have been used to render cultivated plants such as soybean, cotton, corn, beets and rape, tolerant to herbicides such as glyphosate and glufosinate, some of which are commercially available under the trade names RoundupReady® (glyphosate-tolerant, Monsanto, U.S.A.) and LibertyLink® (glufosinate-tolerant, Bayer CropScience, Germany).

Furthermore, plants are also covered that are by the use of recombinant DNA techniques capable to synthesize one or more insecticidal proteins, especially those known from the bacterial genus *Bacillus*, particularly from *Bacillus thuringiensis*, such as δ -endotoxins, e. g. CryIA(b), CryIA(c), CryIF, CryIF(a2), CryIIA(b), CryIIIA, CryIIIB(b1) or Cry9c; vegetative insecticidal proteins (VIP), e. g. VIP1, VIP2, VIP3 or VIP3A; insecticidal proteins of bacteria colonizing nematodes, e. g. *Photorhabdus* spp. or *Xenorhabdus* spp.; toxins produced by animals, such as scorpion toxins, arachnid toxins, wasp toxins, or other insect-specific neurotoxins; toxins produced by fungi, such Streptomyces toxins, plant lectins, such as pea or barley lectins; agglutinins; proteinase inhibitors, such as trypsin inhibitors, serine protease inhibitors, patatin, cystatin or papain inhibitors; ribosome-inactivating proteins (RIP), such as ricin, maize-RIP, abrin, luffin, saporin or bryodin; steroid metabolism enzymes, such as 3-hydroxysteroid oxidase, ecdysteroid-IDP-glycosyl-transferase, cholesterol oxidases, ecdysone inhibitors or HMG-CoA-reductase; ion channel blockers, such as blockers of sodium or calcium channels; juvenile hormone esterase; diuretic hormone receptors (helicokinin receptors); stilbene synthase, bibenzyl synthase, chitinases or glucanases. In the context of the present invention these insecticidal proteins or toxins are to be understood expressly also as pre-toxins, hybrid proteins, truncated or otherwise modified proteins. Hybrid proteins are characterized by a new combination of protein domains, (see, e. g. WO 02/015701). Further examples of such toxins or genetically modified plants capable of synthesizing such toxins are disclosed, e. g., in EP-A 374 753, WO 93/007278, WO 95/34656, EP-A 427 529, EP-A 451 878, WO 03/18810 and WO 03/52073. The methods for producing such genetically modified plants are generally known to the person skilled in the art and are described, e. g. in the publications mentioned above. These insecticidal proteins contained in the genetically modified plants impart to the plants producing these proteins tolerance to harmful pests from all taxonomic groups of arthropods, especially to beetles (Coleoptera), two-winged insects (Diptera), and moths (Lepidoptera) and to nematodes (Nematoda). Genetically modified plants capable to synthesize one or more insecticidal proteins are, e. g., described in the publications mentioned above, and some of which are commercially available such as YieldGard® (corn cultivars producing the Cry1Ab toxin), YieldGard® Plus (corn cultivars producing Cry1Ab and Cry3Bb1 toxins), Starlink® (corn cultivars producing the Cry9c toxin), Herculex® RW (corn cultivars producing Cry34Ab1, Cry35Ab1 and the enzyme Phosphinothricin-*N*-Acetyltransferase [PAT]); NuCOTN® 33B (cotton cultivars producing the Cry1Ac toxin), Bollgard® I (cotton cultivars producing the Cry1Ac toxin), Bollgard® II (cotton cultivars producing Cry1Ac and Cry2Ab2 toxins); VIPCOT® (cotton cultivars producing a VIP-toxin); NewLeaf® (potato cultivars producing the Cry3A toxin); Bt-Xtra®, NatureGard®, KnockOut®, BiteGard®, Protecta®, Bt11 (e. g. Agrisure® CB) and Bt176 from

Syngenta Seeds SAS, France, (corn cultivars producing the Cry1Ab toxin and PAT enzyme), MIR604 from Syngenta Seeds SAS, France (corn cultivars producing a modified version of the Cry3A toxin, c.f. WO 03/018810), MON 863 from Monsanto Europe S.A., Belgium (corn cultivars producing the Cry3Bb1 toxin), IPC 531 from Monsanto Europe S.A., Belgium (cotton cultivars producing a modified version of the Cry1Ac toxin) and 1507 from Pioneer Overseas Corporation, Belgium (corn cultivars producing the Cry1F toxin and PAT enzyme).

Furthermore, plants are also covered that are by the use of recombinant DNA techniques capable to synthesize one or more proteins to increase the resistance or tolerance of those plants to bacterial, viral or fungal pathogens. Examples of such proteins are the so-called "pathogenesis-related proteins" (PR proteins, see, e. g. EP-A 392 225), plant disease resistance genes (e. g. potato cultivars, which express resistance genes acting against *Phytophthora infestans* derived from the mexican wild potato *Solanum bulbocastanum*) or T4-lysozym (e. g. potato cultivars capable of synthesizing these proteins with increased resistance against bacteria such as *Erwinia amylovora*). The methods for producing such genetically modified plants are generally known to the person skilled in the art and are described, e. g. in the publications mentioned above.

Furthermore, plants are also covered that are by the use of recombinant DNA techniques capable to synthesize one or more proteins to increase the productivity (e. g. bio mass production, grain yield, starch content, oil content or protein content), tolerance to drought, salinity or other growth-limiting environmental factors or tolerance to pests and fungal, bacterial or viral pathogens of those plants.

Furthermore, plants are also covered that contain by the use of recombinant DNA techniques a modified amount of substances of content or new substances of content, specifically to improve human or animal nutrition, e. g. oil crops that produce health-promoting long-chain omega-3 fatty acids or unsaturated omega-9 fatty acids (e. g. Nexera® rape, DOW Agro Sciences, Canada).

Furthermore, plants are also covered that contain by the use of recombinant DNA techniques a modified amount of substances of content or new substances of content, specifically to improve raw material production, e. g. potatoes that produce increased amounts of amylopectin (e. g. Amflora® potato, BASF SE, Germany).

The inventive mixtures and compositions are particularly suitable for controlling the following plant diseases:

Albugo spp. (white rust) on ornamentals, vegetables (e. g. *A. candida*) and sunflowers (e. g. *A. tragopogonis*); *Alternaria* spp. (*Alternaria* leaf spot) on vegetables, rape (*A. brassicola* or *brassicae*), sugar beets (*A. tenuis*), fruits, rice, soybeans, potatoes (e. g. *A. solani* or *A. alternata*), tomatoes (e. g. *A. solani* or *A. alternata*) and wheat; *Aphanomyces* spp. on sugar beets and vegetables; *Ascochyta* spp. on cereals and vegetables, e. g. *A. tritici* (anthracnose) on wheat and *A. hordei* on barley; *Bipolaris* and *Drechslera* spp. (teleomorph: *Cochliobolus* spp.), e. g. Southern leaf blight (*D. maydis*) or Northern leaf blight (*B. zeicola*) on corn, e. g. spot blotch (*B. sorokiniana*) on cereals and e. g. *B. oryzae* on rice and turfs; *Blumeria* (formerly *Erysiphe*) *graminis* (powdery mildew) on cereals (e. g. on wheat or barley); *Botrytis cinerea*

- (teleomorph: *Botryotinia fuckeliana*: grey mold) on fruits and berries (e. g. strawberries), vegetables (e. g. lettuce, carrots, celery and cabbages), rape, flowers, vines, forestry plants and wheat; *Bremia lactucae* (downy mildew) on lettuce; *Ceratocystis* (syn. *Ophiostoma*) spp. (rot or wilt) on broad-leaved trees and evergreens, e. g. *C. ulmi* (Dutch elm disease) on elms;
- 5 *Cercospora* spp. (Cercospora leaf spots) on corn (e. g. Gray leaf spot: *C. zeae-maydis*), rice, sugar beets (e. g. *C. beticola*), sugar cane, vegetables, coffee, soybeans (e. g. *C. sojae* or *C. kikuchii*) and rice; *Cladosporium* spp. on tomatoes (e. g. *C. fulvum*: leaf mold) and cereals, e. g. *C. herbarum* (black ear) on wheat; *Claviceps purpurea* (ergot) on cereals; *Cochliobolus* (anamorph: *Helminthosporium* of *Bipolaris*) spp. (leaf spots) on corn (*C. carbonum*), cereals
- 10 (e. g. *C. sativus*, anamorph: *B. sorokiniana*) and rice (e. g. *C. miyabeanus*, anamorph: *H. oryzae*); *Colletotrichum* (teleomorph: *Glomerella*) spp. (anthracnose) on cotton (e. g. *C. gossypii*), corn (e. g. *C. graminicola*: Anthracnose stalk rot), soft fruits, potatoes (e. g. *C. coccodes*: black dot), beans (e. g. *C. lindemuthianum*) and soybeans (e. g. *C. truncatum* or *C. gloeosporioides*); *Corticium* spp., e. g. *C. sasakii* (sheath blight) on rice; *Corynespora cassiicola*
- 15 (leaf spots) on soybeans and ornamentals; *Cycloconium* spp., e. g. *C. oleaginum* on olive trees; *Cylindrocarpon* spp. (e. g. fruit tree canker or young vine decline, teleomorph: *Nectria* or *Neonectria* spp.) on fruit trees, vines (e. g. *C. liriodendri*, teleomorph: *Neonectria liriodendri*: Black Foot Disease) and ornamentals; *Dematophora* (teleomorph: *Rosellinia*) *necatrix* (root and stem rot) on soybeans; *Diaporthe* spp., e. g. *D. phaseolorum* (damping off) on soybeans;
- 20 *Drechslera* (syn. *Helminthosporium*, teleomorph: *Pyrenophora*) spp. on corn, cereals, such as barley (e. g. *D. teres*, net blotch) and wheat (e. g. *D. tritici-repentis*: tan spot), rice and turf; Esca (dieback, apoplexy) on vines, caused by *Formitiporia* (syn. *Phellinus*) *punctata*, *F. mediterranea*, *Phaeomoniella chlamydospora* (earlier *Phaeoacremonium chlamydosporum*), *Phaeoacremonium aleophilum* and/or *Botryosphaeria obtusa*; *Elsinoe* spp. on pome fruits (*E. pyri*), soft fruits (*E. veneta*: anthracnose) and vines (*E. ampelina*: anthracnose); *Entyloma*
- 25 *oryzae* (leaf smut) on rice; *Epicoccum* spp. (black mold) on wheat; *Erysiphe* spp. (powdery mildew) on sugar beets (*E. betae*), vegetables (e. g. *E. pisi*), such as cucurbits (e. g. *E. cichoracearum*), cabbages, rape (e. g. *E. cruciferarum*); *Eutypa lata* (*Eutypa* canker or dieback, anamorph: *Cytosporina lata*, syn. *Libertella blepharis*) on fruit trees, vines and ornamental
- 30 woods; *Exserohilum* (syn. *Helminthosporium*) spp. on corn (e. g. *E. turcicum*); *Fusarium* (teleomorph: *Gibberella*) spp. (wilt, root or stem rot) on various plants, such as *F. graminearum* or *F. culmorum* (root rot, scab or head blight) on cereals (e. g. wheat or barley), *F. oxysporum* on tomatoes, *F. solani* (f. sp. *glycines* now syn. *F. virguliforme*) and *F. tucumaniae* and *F. brasiliense* each causing sudden death syndrome on soybeans, and *F. verticillioides* on corn;
- 35 *Gaeumannomyces graminis* (take-all) on cereals (e. g. wheat or barley) and corn; *Gibberella* spp. on cereals (e. g. *G. zeae*) and rice (e. g. *G. fujikuroi*: Bakanae disease); *Glomerella cingulata* on vines, pome fruits and other plants and *G. gossypii* on cotton; Grainstaining complex on rice; *Guignardia bidwellii* (black rot) on vines; *Gymnosporangium* spp. on rosaceous plants and junipers, e. g. *G. sabinae* (rust) on pears; *Helminthosporium* spp. (syn. *Drechslera*,
- 40 teleomorph: *Cochliobolus*) on corn, cereals and rice; *Hemileia* spp., e. g. *H. vastatrix* (coffee leaf rust) on coffee; *Isariopsis clavispora* (syn. *Cladosporium vitis*) on vines; *Macrophomina phaseolina* (syn. *phaseoli*) (root and stem rot) on soybeans and cotton; *Microdochium* (syn. *Fusarium*) *nivale* (pink snow mold) on cereals (e. g. wheat or barley); *Microsphaera diffusa*

(powdery mildew) on soybeans; *Monilinia* spp., e. g. *M. laxa*, *M. fructicola* and *M. fructigena* (bloom and twig blight, brown rot) on stone fruits and other rosaceous plants; *Mycosphaerella* spp. on cereals, bananas, soft fruits and ground nuts, such as e. g. *M. graminicola* (anamorph: *Septoria tritici*, Septoria blotch) on wheat or *M. fijiensis* (black Sigatoka disease) on bananas;

5 *Peronospora* spp. (downy mildew) on cabbage (e. g. *P. brassicae*), rape (e. g. *P. parasitica*), onions (e. g. *P. destructor*), tobacco (*P. tabacina*) and soybeans (e. g. *P. manshurica*); *Phakopsora pachyrhizi* and *P. meibomia* (soybean rust) on soybeans; *Phialophora* spp. e. g. on vines (e. g. *P. tracheiphila* and *P. tetraspora*) and soybeans (e. g. *P. gregata*: stem rot); *Phoma lingam* (root and stem rot) on rape and cabbage and *P. betae* (root rot, leaf spot and

10 damping-off) on sugar beets; *Phomopsis* spp. on sunflowers, vines (e. g. *P. viticola*: can and leaf spot) and soybeans (e. g. stem rot: *P. phaseoli*, teleomorph: *Diaporthe phaseolorum*); *Physoderma maydis* (brown spots) on corn; *Phytophthora* spp. (wilt, root, leaf, fruit and stem root) on various plants, such as paprika and cucurbits (e. g. *P. capsici*), soybeans (e. g. *P. megasperma*, syn. *P. sojae*), potatoes and tomatoes (e. g. *P. infestans*: late blight) and broad-

15 leaved trees (e. g. *P. ramorum*: sudden oak death); *Plasmodiophora brassicae* (club root) on cabbage, rape, radish and other plants; *Plasmopara* spp., e. g. *P. viticola* (grapevine downy mildew) on vines and *P. halstedii* on sunflowers; *Podosphaera* spp. (powdery mildew) on rosaceous plants, hop, pome and soft fruits, e. g. *P. leucotricha* on apples; *Polymyxa* spp., e. g. on cereals, such as barley and wheat (*P. graminis*) and sugar beets (*P. betae*) and thereby

20 transmitted viral diseases; *Pseudocercospora herpotrichoides* (eyespot, teleomorph: *Tapesia yallundae*) on cereals, e. g. wheat or barley; *Pseudoperonospora* (downy mildew) on various plants, e. g. *P. cubensis* on cucurbits or *P. humili* on hop; *Pseudopezizicola tracheiphila* (red fire disease or 'rotbrenner', anamorph: *Phialophora*) on vines; *Puccinia* spp. (rusts) on various plants, e. g. *P. triticina* (brown or leaf rust), *P. striiformis* (stripe or yellow rust), *P. hordei* (dwarf

25 rust), *P. graminis* (stem or black rust) or *P. recondita* (brown or leaf rust) on cereals, such as e. g. wheat, barley or rye, *P. kuehnii* (orange rust) on sugar cane and *P. asparagi* on asparagus; *Pyrenophora* (anamorph: *Drechslera*) *tritici-repentis* (tan spot) on wheat or *P. teres* (net blotch) on barley; *Pyricularia* spp., e. g. *P. oryzae* (teleomorph: *Magnaporthe grisea*, rice blast) on rice and *P. grisea* on turf and cereals; *Pythium* spp. (damping-off) on turf, rice, corn, wheat, cotton,

30 rape, sunflowers, soybeans, sugar beets, vegetables and various other plants (e. g. *P. ultimum* or *P. aphanidermatum*); *Ramularia* spp., e. g. *R. collo-cygni* (Ramularia leaf spots, Physiological leaf spots) on barley and *R. beticola* on sugar beets; *Rhizoctonia* spp. on cotton, rice, potatoes, turf, corn, rape, potatoes, sugar beets, vegetables and various other plants, e. g. *R. solani* (root and stem rot) on soybeans, *R. solani* (sheath blight) on rice or *R. cerealis* (Rhizoctonia spring

35 blight) on wheat or barley; *Rhizopus stolonifer* (black mold, soft rot) on strawberries, carrots, cabbage, vines and tomatoes; *Rhynchosporium secalis* (scald) on barley, rye and triticale; *Sarocladium oryzae* and *S. attenuatum* (sheath rot) on rice; *Sclerotinia* spp. (stem rot or white mold) on vegetables and field crops, such as rape, sunflowers (e. g. *S. sclerotiorum*) and soybeans (e. g. *S. rolfsii* or *S. sclerotiorum*); *Septoria* spp. on various plants, e. g. *S. glycines* (brown spot) on soybeans, *S. tritici* (Septoria blotch) on wheat and *S.* (syn. *Stagonospora*)

40 *nodorum* (Stagonospora blotch) on cereals; *Uncinula* (syn. *Erysiphe*) *necator* (powdery mildew, anamorph: *Oidium tuckeri*) on vines; *Setosphaeria* spp. (leaf blight) on corn (e. g. *S. turcicum*, syn. *Helminthosporium turcicum*) and turf; *Sphacelotheca* spp. (smut) on corn, (e. g. *S. reiliana*:

head smut), sorghum und sugar cane; *Sphaerotheca fuliginea* (powdery mildew) on cucurbits; *Spongospora subterranea* (powdery scab) on potatoes and thereby transmitted viral diseases; *Stagonospora* spp. on cereals, e. g. *S. nodorum* (Stagonospora blotch, teleomorph: *Leptosphaeria* [syn. *Phaeosphaeria*] *nodorum*) on wheat; *Synchytrium endobioticum* on potatoes (potato wart disease); *Taphrina* spp., e. g. *T. deformans* (leaf curl disease) on peaches and *T. pruni* (plum pocket) on plums; *Thielaviopsis* spp. (black root rot) on tobacco, pome fruits, vegetables, soybeans and cotton, e. g. *T. basicola* (syn. *Chalara elegans*); *Tilletia* spp. (common bunt or stinking smut) on cereals, such as e. g. *T. tritici* (syn. *T. caries*, wheat bunt) and *T. controversa* (dwarf bunt) on wheat; *Typhula incarnata* (grey snow mold) on barley or wheat; *Urocystis* spp., e. g. *U. occulta* (stem smut) on rye; *Uromyces* spp. (rust) on vegetables, such as beans (e. g. *U. appendiculatus*, syn. *U. phaseoli*) and sugar beets (e. g. *U. betae*); *Ustilago* spp. (loose smut) on cereals (e. g. *U. nuda* and *U. avenae*), corn (e. g. *U. maydis*: corn smut) and sugar cane; *Venturia* spp. (scab) on apples (e. g. *V. inaequalis*) and pears; and *Verticillium* spp. (wilt) on various plants, such as fruits and ornamentals, vines, soft fruits, vegetables and field crops, e. g. *V. dahliae* on strawberries, rape, potatoes and tomatoes.

The mixtures and compositions of the invention are also suitable for controlling harmful fungi in the protection of stored products or harvest and in the protection of materials.

The term "protection of materials" is to be understood to denote the protection of technical and non-living materials, such as adhesives, glues, wood, paper and paperboard, textiles, leather, paint dispersions, plastics, cooling lubricants, fiber or fabrics, against the infestation and destruction by harmful microorganisms, such as fungi and bacteria. As to the protection of wood and other materials, the particular attention is paid to the following harmful fungi: Ascomycetes such as *Ophiostoma* spp., *Ceratocystis* spp., *Aureobasidium pullulans*, *Sclerophoma* spp., *Chaetomium* spp., *Humicola* spp., *Petriella* spp., *Trichurus* spp.; Basidiomycetes such as *Coniophora* spp., *Coriolus* spp., *Gloeophyllum* spp., *Lentinus* spp., *Pleurotus* spp., *Poria* spp., *Serpula* spp. and *Tyromyces* spp.; Deuteromycetes such as *Aspergillus* spp., *Cladosporium* spp., *Penicillium* spp., *Trichoderma* spp., *Alternaria* spp., *Paecilomyces* spp. and Zygomycetes such as *Mucor* spp., and in addition in the protection of stored products and harvest the following yeast fungi are worthy of note: *Candida* spp. and *Saccharomyces cerevisiae*.

The mixtures and compositions or any group of mixtures or compositions of the present invention are particularly suitable for controlling phytopathogenic fungi in particular crops, specifically in cereals or soybeans, wherein the cereals are in particular selected from wheat, rye, barley, triticale and oats.

According to one embodiment, the present invention relates to the use of any one of the inventive mixtures or compositions as detailed herein for controlling phytopathogenic fungi in wheat.

According to one embodiment, the present invention relates to the use of any one of the inventive mixtures or compositions as detailed herein for controlling phytopathogenic fungi in rye.

According to a further embodiment, the present invention relates to the use of any one of the inventive mixtures or compositions as detailed herein for controlling phytopathogenic fungi in barley.

5 According to a further embodiment, the present invention relates to the use of any one of the inventive mixtures or compositions as detailed herein for controlling phytopathogenic fungi in triticale.

According to a further embodiment, the present invention relates to the use of any one of the inventive mixtures or compositions as detailed herein for controlling phytopathogenic fungi in/on soybeans.

10 According to a further embodiment, the present invention relates to the use of any one of the inventive mixtures or compositions as detailed herein for controlling phytopathogenic fungi in oats.

More specifically, according to a further embodiment, the present invention relates to the use of the inventive mixtures or compositions for controlling a disease selected from:

15 *M. graminicola* (anamorph: *Septoria tritici*, *Septoria blotch*) on wheat;

Puccinia spp. (rusts) on various plants, e. g. *P. triticina* (brown or leaf rust), *P. striiformis* (stripe or yellow rust), *P. hordei* (dwarf rust), *P. graminis* (stem or black rust) or *P. recondita* (brown or leaf rust);

20 *Pyrenophora* (anamorph: *Drechslera*) *tritici-repentis* (tan spot) on wheat or *P. teres* (net blotch) on barley;

Erysiphe spp. (powdery mildew), e.g. *Erysiphe graminis tritici* on wheat or *Erysiphe graminis hordei* on barley;

Ramularia spp., e. g. *R. collo-cygni* (*Ramularia* leaf spots, *Physiological* leaf spots) on barley;

Rhynchosporium secalis (scald) on barley, rye and triticale;

25 *Phaeosphaeria nodorum* (glume blotch) on wheat;

Microdochium nivale (stem blight and head blight) on wheat;

Pseudocercospora herpotrichoides (eyespot, teleomorph: *Tapesia yallundae*) on cereals, e. g. wheat or barley;

Microdochium (syn. *Fusarium*) *nivale* (pink snow mold) on cereals (e. g. wheat or barley);

30 *Fusarium* (teleomorph: *Gibberella*) spp. (wilt, root or stem rot) e.g. *F. graminearum* or *F. culmorum* (root rot, scab or head blight) on cereals (e. g. wheat or barley);

Gaeumannomyces graminis (take-all) on cereals (e. g. wheat or barley);

Phakopsora pachyrhizi (Asian soybean rust) on soybeans;

P. meibomia (soybean rust) on soybeans; *Cercospora sojina* (Frogeye leaf spot) on soybean;

35 *Cercospora kikuchii* on soybean;

Corynespora cassiicola (target/leaf spots) on soybeans;

Glomerella glycines (Anthracnose) on soybeans; and
Sclerotinia sclerotiorum (White Mould) on soybeans.

According to one particular embodiment, the present invention relates to the use of any one of the compositions T1-213 to T1-265 for controlling phytopathogenic fungi in cereals.

- 5 According to one particular embodiment, the present invention relates to the use of any one of the compositions T1-266 to T1-318 for controlling phytopathogenic fungi in cereals.

According to one particular embodiment, the present invention relates to the use of any one of the compositions T1-213 to T1-265 for controlling phytopathogenic fungi in soybeans.

- 10 According to one particular embodiment, the present invention relates to the use of any one of the compositions T1-266 to T1-318 for controlling phytopathogenic fungi in soybeans.

The mixtures and compositions according to the invention can also be used in the field of protecting stored products or harvest against attack of fungi and microorganisms. According to the present invention, the term "stored products" is understood to denote natural substances of plant or animal origin and their processed forms, which have been taken from the natural life cycle and for which long-term protection is desired. Stored products of crop plant origin, such as plants or parts thereof, for example stalks, leaves, tubers, seeds, fruits or grains, can be protected in the freshly harvested state or in processed form, such as pre-dried, moistened, comminuted, ground, pressed or roasted, which process is also known as post-harvest treatment. Also falling under the definition of stored products is timber, whether in the form of crude timber, such as construction timber, electricity pylons and barriers, or in the form of finished articles, such as furniture or objects made from wood. Stored products of animal origin are hides, leather, furs, hairs and the like. The combinations according to the present invention can prevent disadvantageous effects such as decay, discoloration or mold. Preferably "stored products" is understood to denote natural substances of plant origin and their processed forms, more preferably fruits and their processed forms, such as pomes, stone fruits, soft fruits and citrus fruits and their processed forms.

Compositions of the invention may also be suitable for efficiently controlling arthropodal pests such as arachnids, myriapedes and insects as well as nematodes.

- 30 The mixtures and compositions of the invention may be used for improving the health of a plant. The invention also relates to a method for improving plant health by treating a plant, its propagation material and/or the locus where the plant is growing or is to grow with an effective amount of the inventive compositions or the components thereof, respectively.

The term "plant health" is to be understood to denote a condition of the plant and/or its products which is determined by several indicators alone or in combination with each other such as yield (e. g. increased biomass and/or increased content of valuable ingredients), plant vigor (e. g. improved plant growth and/or greener leaves ("greening effect")), quality (e. g. improved content or composition of certain ingredients) and tolerance to abiotic and/or biotic stress. The above identified indicators for the health condition of a plant may be interdependent or may result from each other.

The compounds of the inventive compositions or mixtures can be present in different crystal modifications whose biological activity may differ. Compositions and mixtures containing the same are likewise subject matter of the present invention.

The inventive compositions are employed as such or in form of agrochemical compositions by treating the fungi or the plants, plant propagation materials, such as seeds, soil, surfaces, materials or rooms to be protected from fungal attack with a fungicidally effective amount of the active substances. The application can be carried out both before and after the infection of the plants, plant propagation materials, such as seeds, soil, surfaces, materials or rooms by the fungi.

Plant propagation materials may be treated with the mixtures and compositions of the invention prophylactically either at or before planting or transplanting.

The invention also relates to agrochemical compositions comprising an auxiliary and a component I, a component II and a component III according to the invention.

An agrochemical composition comprises a fungicidally effective amount of a component I, a component II and a component III. The term "effective amount" denotes an amount of the composition or of the components I and II and III, which is sufficient for controlling harmful fungi on cultivated plants or in the protection of materials and which does not result in a substantial damage to the treated plants. Such an amount can vary in a broad range and is dependent on various factors, such as the fungal species to be controlled, the treated cultivated plant or material, the climatic conditions and the specific component I and II and III used.

The components I, components II and components III, their N-oxides and salts can be converted into customary types of agrochemical compositions, e. g. solutions, emulsions, suspensions, dusts, powders, pastes, granules, pressings, capsules, and mixtures thereof. Examples for composition types are suspensions (e. g. SC, OD, FS), emulsifiable concentrates (e. g. EC), emulsions (e. g. EW, EO, ES, ME), capsules (e. g. CS, ZC), pastes, pastilles, wettable powders or dusts (e. g. WP, SP, WS, DP, DS), pressings (e. g. BR, TB, DT), granules (e. g. WG, SG, GR, FG, GG, MG), insecticidal articles (e. g. LN), as well as gel formulations for the treatment of plant propagation materials such as seeds (e. g. GF). These and further compositions types are defined in the "Catalogue of pesticide formulation types and international coding system", Technical Monograph No. 2, 6th Ed. May 2008, CropLife International.

The compositions are prepared in a known manner, such as described by Mollet and Grubemann, Formulation technology, Wiley VCH, Weinheim, 2001; or Knowles, New developments in crop protection product formulation, Agrow Reports DS243, T&F Informa, London, 2005.

Suitable auxiliaries are solvents, liquid carriers, solid carriers or fillers, surfactants, dispersants, emulsifiers, wetters, adjuvants, solubilizers, penetration enhancers, protective colloids, adhesion agents, thickeners, humectants, repellents, attractants, feeding stimulants, compatibilizers, bactericides, anti-freezing agents, anti-foaming agents, colorants, tackifiers and binders.

Suitable solvents and liquid carriers are water and organic solvents, such as mineral oil fractions of medium to high boiling point, e. g. kerosene, diesel oil; oils of vegetable or animal origin; aliphatic, cyclic and aromatic hydrocarbons, e. g. toluene, paraffin, tetrahydronaphthalene, alkylated naphthalenes; alcohols, e. g. ethanol, propanol, butanol, benzyl alcohol, cyclohexanol; glycols; DMSO; ketones, e. g. cyclohexanone; esters, e. g. lactates, carbonates, fatty acid esters, gamma-butyrolactone; fatty acids; phosphonates; amines; amides, e. g. N-methylpyrrolidone, fatty acid dimethylamides; and mixtures thereof.

Suitable solid carriers or fillers are mineral earths, e. g. silicates, silica gels, talc, kaolins, limestone, lime, chalk, clays, dolomite, diatomaceous earth, bentonite, calcium sulfate, magnesium sulfate, magnesium oxide; polysaccharides, e. g. cellulose, starch; fertilizers, e. g. ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas; products of vegetable origin, e. g. cereal meal, tree bark meal, wood meal, nutshell meal, and mixtures thereof.

Suitable surfactants are surface-active compounds, such as anionic, cationic, nonionic and amphoteric surfactants, block polymers, polyelectrolytes, and mixtures thereof. Such surfactants can be used as emulsifier, dispersant, solubilizer, wetter, penetration enhancer, protective colloid, or adjuvant. Examples of surfactants are listed in McCutcheon's, Vol.1: Emulsifiers & Detergents, McCutcheon's Directories, Glen Rock, USA, 2008 (International Ed. or North American Ed.).

Suitable anionic surfactants are alkali, alkaline earth or ammonium salts of sulfonates, sulfates, phosphates, carboxylates, and mixtures thereof. Examples of sulfonates are alkylarylsulfonates, diphenylsulfonates, alpha-olefin sulfonates, lignin sulfonates, sulfonates of fatty acids and oils, sulfonates of ethoxylated alkylphenols, sulfonates of alkoxyated arylphenols, sulfonates of condensed naphthalenes, sulfonates of dodecyl- and tridecylbenzenes, sulfonates of naphthalenes and alkylnaphthalenes, sulfosuccinates or sulfosuccinamates. Examples of sulfates are sulfates of fatty acids and oils, of ethoxylated alkylphenols, of alcohols, of ethoxylated alcohols, or of fatty acid esters. Examples of phosphates are phosphate esters. Examples of carboxylates are alkyl carboxylates, and carboxylated alcohol or alkylphenol ethoxylates.

Suitable nonionic surfactants are alkoxyates, N-substituted fatty acid amides, amine oxides, esters, sugar-based surfactants, polymeric surfactants, and mixtures thereof. Examples of alkoxyates are compounds such as alcohols, alkylphenols, amines, amides, arylphenols, fatty acids or fatty acid esters which have been alkoxyated with 1 to 50 equivalents. Ethylene oxide and/or propylene oxide may be employed for the alkoxyation, preferably ethylene oxide. Examples of N-substituted fatty acid amides are fatty acid glucamides or fatty acid alkanolamides. Examples of esters are fatty acid esters, glycerol esters or monoglycerides. Examples of sugar-based surfactants are sorbitans, ethoxylated sorbitans, sucrose and glucose esters or alkylpolyglucosides. Examples of polymeric surfactants are home- or copolymers of vinylpyrrolidone, vinylalcohols, or vinylacetate.

Suitable cationic surfactants are quaternary surfactants, for example quaternary ammonium compounds with one or two hydrophobic groups, or salts of long-chain primary amines. Suitable amphoteric surfactants are alkylbetains and imidazolines. Suitable block polymers are block

polymers of the A-B or A-B-A type comprising blocks of polyethylene oxide and polypropylene oxide, or of the A-B-C type comprising alkanol, polyethylene oxide and polypropylene oxide. Suitable polyelectrolytes are polyacids or polybases. Examples of polyacids are alkali salts of polyacrylic acid or polyacid comb polymers. Examples of polybases are polyvinyl amines or polyethylene amines.

Suitable adjuvants are compounds, which have a negligible or even no pesticidal activity themselves, and which improve the biological performance of the component used on the target. Examples are surfactants, mineral or vegetable oils, and other auxiliaries. Further examples are listed by Knowles, Adjuvants and additives, Agrow Reports DS256, T&F Informa UK, 2006, chapter 5.

Suitable thickeners are polysaccharides (e. g. xanthan gum, carboxymethyl cellulose), inorganic clays (organically modified or unmodified), polycarboxylates, and silicates.

Suitable bactericides are bronopol and isothiazolinone derivatives such as alkylisothiazolinones and benzisothiazolinones.

Suitable anti-freezing agents are ethylene glycol, propylene glycol, urea and glycerin.

Suitable anti-foaming agents are silicones, long chain alcohols, and salts of fatty acids.

Suitable colorants (e. g. in red, blue, or green) are pigments of low water solubility and water-soluble dyes. Examples are inorganic colorants (e. g. iron oxide, titan oxide, iron hexacyanoferrate) and organic colorants (e. g. alizarin-, azo- and phthalocyanine colorants).

Suitable tackifiers or binders are polyvinyl pyrrolidones, polyvinyl acetates, polyvinyl alcohols, polyacrylates, biological or synthetic waxes, and cellulose ethers.

Examples for composition types and their preparation are (wherein active substances denote component I, component II and component III):

i) Water-soluble concentrates (SL, LS)

10-60 wt% active substances and 5-15 wt% wetting agent (e. g. alcohol alkoxyates) are dissolved in water and/or in a water-soluble solvent (e. g. alcohols) ad 100 wt%. The active substance dissolves upon dilution with water.

ii) Dispersible concentrates (DC)

5-25 wt% active substances and 1-10 wt% dispersant (e. g. polyvinyl pyrrolidone) are dissolved in organic solvent (e. g. cyclohexanone) ad 100 wt%. Dilution with water gives a dispersion.

iii) Emulsifiable concentrates (EC)

15-70 wt% active substances and 5-10 wt% emulsifiers (e. g. calcium dodecylbenzenesulfonate and castor oil ethoxylate) are dissolved in water-insoluble organic solvent (e. g. aromatic hydrocarbon) ad 100 wt%. Dilution with water gives an emulsion.

iv) Emulsions (EW, EO, ES)

5-40 wt% active substances and 1-10 wt% emulsifiers (e. g. calcium dodecylbenzenesulfonate and castor oil ethoxylate) are dissolved in 20-40 wt% water-insoluble organic solvent (e. g.

aromatic hydrocarbon). This mixture is introduced into water ad 100 wt% by means of an emulsifying machine and made into a homogeneous emulsion. Dilution with water gives an emulsion.

v) Suspensions (SC, OD, FS)

- 5 In an agitated ball mill, 20-60 wt% active substances are comminuted with addition of 2-10 wt% dispersants and wetting agents (e. g. sodium lignosulfonate and alcohol ethoxylate), 0.1-2 wt% thickener (e. g. xanthan gum) and ad water ad 100 wt% to give a fine active substance suspension. Dilution with water gives a stable suspension of the active substance. For FS type composition up to 40 wt% binder (e. g. polyvinyl alcohol) is added.

10 vi) Water-dispersible granules and water-soluble granules (WG, SG)

- 50-80 wt% active substances are ground finely with addition of dispersants and wetting agents (e. g. sodium lignosulfonate and alcohol ethoxylate) ad 100 wt% and prepared as water-dispersible or water-soluble granules by means of technical appliances (e. g. extrusion, spray tower, fluidized bed). Dilution with water gives a stable dispersion or solution of the active
15 substance.

vii) Water-dispersible powders and water-soluble powders (WP, SP, WS)

- 50-80 wt% active substances are ground in a rotor-stator mill with addition of 1-5 wt% dispersants (e. g. sodium lignosulfonate), 1-3 wt% wetting agents (e. g. alcohol ethoxylate) and solid carrier (e. g. silica gel) ad 100 wt%. Dilution with water gives a stable dispersion or solution
20 of the active substance.

viii) Gel (GW, GF)

- In an agitated ball mill, 5-25 wt% active substances are comminuted with addition of 3-10 wt% dispersants (e. g. sodium lignosulfonate), 1-5 wt% thickener (e. g. carboxymethyl cellulose) and water ad 100 wt% to give a fine suspension of the active substance. Dilution with water gives a
25 stable suspension of the active substance.

ix) Microemulsion (ME)

- 5-20 wt% active substances are added to 5-30 wt% organic solvent blend (e. g. fatty acid dimethylamide and cyclohexanone), 10-25 wt% surfactant blend (e. g. alcohol ethoxylate and arylphenol ethoxylate), and water ad 100 wt%. This mixture is stirred for 1 h to produce
30 spontaneously a thermodynamically stable microemulsion.

x) Microcapsules (CS)

- An oil phase comprising 5-50 wt% active substances, 0-40 wt% water insoluble organic solvent (e. g. aromatic hydrocarbon), 2-15 wt% acrylic monomers (e. g. methylmethacrylate, methacrylic acid and a di- or triacrylate) are dispersed into an aqueous solution of a protective colloid (e. g.
35 polyvinyl alcohol). Radical polymerization initiated by a radical initiator results in the formation of poly(meth)acrylate microcapsules. Alternatively, an oil phase comprising 5-50 wt% of active substances, 0-40 wt% water insoluble organic solvent (e. g. aromatic hydrocarbon), and an isocyanate monomer (e. g. diphenylmethene-4,4'-diisocyanatae) are dispersed into an aqueous solution of a protective colloid (e. g. polyvinyl alcohol). The addition of a polyamine (e. g.

hexamethylenediamine) results in the formation of polyurea microcapsules. The monomers amount to 1-10 wt%. The wt% relate to the total CS composition.

xi) Dustable powders (DP, DS)

1-10 wt% active substances are ground finely and mixed intimately with solid carrier (e. g. finely divided kaolin) ad 100 wt%.

xii) Granules (GR, FG)

0.5-30 wt% active substances are ground finely and associated with solid carrier (e. g. silicate) ad 100 wt%. Granulation is achieved by extrusion, spray-drying or fluidized bed.

xiii) Ultra-low volume liquids (UL)

1-50 wt% active substances are dissolved in organic solvent (e. g. aromatic hydrocarbon) ad 100 wt%.

The compositions types i) to xiii) may optionally comprise further auxiliaries, such as 0.1-1 wt% bactericides, 5-15 wt% anti-freezing agents, 0.1-1 wt% anti-foaming agents, and 0.1-1 wt% colorants.

The agrochemical compositions generally comprise between 0.01 and 95%, preferably between 0.1 and 90%, and in particular between 0.5 and 75%, by weight of active substances. The active substances are employed in a purity of from 90% to 100%, preferably from 95% to 100% (according to NMR spectrum).

For the purposes of treatment of plant propagation materials, particularly seeds, solutions for seed treatment (LS), Suspoemulsions (SE), flowable concentrates (FS), powders for dry treatment (DS), water-dispersible powders for slurry treatment (WS), water-soluble powders (SS), emulsions (ES), emulsifiable concentrates (EC), and gels (GF) are usually employed. The compositions in question give, after two-to-tenfold dilution, active substance concentrations of from 0.01 to 60% by weight, preferably from 0.1 to 40%, in the ready-to-use preparations.

Application can be carried out before or during sowing. Methods for applying or treating with component I, component II and component III and compositions thereof, respectively, onto plant propagation material, especially seeds include dressing, coating, pelleting, dusting, and soaking as well as in-furrow application methods. Preferably, component I, component II and component III or the compositions thereof, respectively, are applied on to the plant propagation material by a method such that germination is not induced, e. g. by seed dressing, pelleting, coating and dusting.

When employed in plant protection, the amounts of active substances applied are, depending on the kind of effect desired, from 0.001 to 2 kg per ha, preferably from 0.005 to 2 kg per ha, more preferably from 0.05 to 0.9 kg per ha, and in particular from 0.1 to 0.75 kg per ha.

In treatment of plant propagation materials such as seeds, e. g. by dusting, coating or drenching seed, amounts of active substance of from 0.1 to 1000 g, preferably from 1 to 1000 g, more preferably from 1 to 100 g and most preferably from 5 to 100 g, per 100 kilogram of plant propagation material (preferably seed) are generally required.

When used in the protection of materials or stored products, the amount of active substance

applied depends on the kind of application area and on the desired effect. Amounts customarily applied in the protection of materials are 0.001 g to 2 kg, preferably 0.005 g to 1 kg, of active substance per cubic meter of treated material.

Various types of oils, wetters, adjuvants, fertilizer, or micronutrients, and further pesticides (e. g. herbicides, insecticides, fungicides, growth regulators, safeners) may be added to the active substances or the compositions comprising them as premix or, if appropriate not until immediately prior to use (tank mix). These agents can be admixed with the compositions according to the invention in a weight ratio of 1:100 to 100:1, preferably 1:10 to 10:1.

The user applies the composition according to the invention usually from a predosage device, a knapsack sprayer, a spray tank, a spray plane, or an irrigation system. Usually, the agrochemical composition is made up with water, buffer, and/or further auxiliaries to the desired application concentration and the ready-to-use spray liquor or the agrochemical composition according to the invention is thus obtained. Usually, 20 to 2000 liters, preferably 50 to 400 liters, of the ready-to-use spray liquor are applied per hectare of agricultural useful area.

According to one embodiment, individual components of the composition according to the invention such as parts of a kit or parts of a binary or ternary mixture may be mixed by the user himself in a spray tank or any other kind of vessel used for applications (e. g. seed treater drums, seed pelleting machinery, knapsack sprayer) and further auxiliaries may be added, if appropriate.

In the mixtures and compositions, the compound ratios are advantageously chosen so as to produce a synergistic effect. The term "synergistic effect" is understood to refer in particular to that defined by Colby's formula (Colby, S. R., "Calculating synergistic and antagonistic responses of herbicide combinations", *Weeds*, 15, pp. 20-22, 1967). The term "synergistic effect" is also understood to refer to that defined by application of the Tammes method, (Tammes, P. M. L., "Isoboles, a graphic representation of synergism in pesticides", *Netherl. J. Plant Pathol.* 70, 1964).

In the mixtures and compositions according to the invention, the weight ratio of component I and component II generally depends from the properties of the active substances used, usually it is in the range of from 1:100 to 100:1, regularly in the range of from 1:50 to 50:1, preferably in the range of from 1:20 to 20:1, more preferably in the range of from 1:10 to 10:1, even more preferably in the range of from 1:4 to 4:1 and in particular in the range of from 1:2 to 2:1.

According to further embodiments of the mixtures and compositions according to the invention, the weight ratio of component I versus component II usually is in the range of from 1000:1 to 1:1, often in the range of from 100:1 to 1:1, regularly in the range of from 50:1 to 1:1, preferably in the range of from 20:1 to 1:1, more preferably in the range of from 10:1 to 1:1, even more preferably in the range of from 4:1 to 1:1 and in particular in the range of from 2:1 to 1:1.

According to further embodiments of the mixtures and compositions according to the invention, the weight ratio of component I versus component II usually is in the range of from 1:1 to 1000, often in the range of from 1:1 to 1:100, regularly in the range of from 1:1 to 1:50, preferably in the range of from 1:1 to 1:20, more preferably in the range of from 1:1 to 1:10, even more

preferably in the range of from 1:1 to 1:4 and in particular in the range of from 1:1 to 1:2.

In the mixtures and compositions, the compound ratios are advantageously chosen so as to produce a synergistic effect.

The active compounds, separately or jointly, are prepared as a stock solution comprising 25 mg of active compound which is made up to 10 ml using a mixture of acetone and/or DMSO and the emulsifier Uniperol® EL (wetting agent having an emulsifying and dispersing action based on ethoxylated alkylphenols) in a ratio by volume of solvent/emulsifier of 99:1. The mixture is then made up to 100 ml with water. This stock solution is diluted with the solvent/emulsifier/water mixture described to give the concentration of active compound stated below.

- 10 The visually determined percentages of infected leaf areas are converted into efficacies in % of the untreated control. The efficacy (E) is calculated as follows using Abbot's formula:

$$E = (1 - \alpha/\beta) \cdot 100$$

α corresponds to the fungicidal infection of the treated plants in % and

β corresponds to the fungicidal infection of the untreated (control) plants in %

- 15 An efficacy of 0 means that the infection level of the treated plants corresponds to that of the untreated control plants; an efficacy of 100 means that the treated plants were not infected.

The expected efficacies of active compound combinations were determined using Colby's formula (Colby, S.R. "Calculating synergistic and antagonistic responses of herbicide combinations", Weeds, 15, pp. 20-22, 1967) and compared with the observed efficacies.

- 20 Colby's formula: $E = x + y - x \cdot y/100$

E expected efficacy, expressed in % of the untreated control, when using the mixture of the active compounds A and B at the concentrations a and b

x efficacy, expressed in % of the untreated control, when using the active compound A at the concentration a

- 25 y efficacy, expressed in % of the untreated control, when using the active compound B at the concentration b.

Claims

1. Compositions comprising,

1) as component I a compound selected from the following compounds I-1 to I-3:

compound I-1 1-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-cyclopropyl-2-(1,2,4-triazol-1-yl)ethanol;

compound I-2 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-3-methyl-1-(1,2,4-triazol-1-yl)butan-2-ol; and

compound I-3 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1,2,4-triazol-1-yl)propan-2-ol;

or the N-oxides or agriculturally acceptable salts of each of the compounds;

and

2) as component II

compound II-1 pydiflumetofen, or

compound II-2 N-[(5-chloro-2-isopropyl-phenyl)methyl]-N-cyclopropyl-5-fluoro-1,3-dimethyl-pyrazole-4-carboxamide;

and

3) as component III, a compound selected from the following compounds III-1 to III-53 from the groups A) to H) and K):

A) Respiration inhibitors

- Inhibitors of complex III at Q_o site: azoxystrobin (III-1), fluoxastrobin (III-2), picoxystrobin (III-3), pyraclostrobin (III-4), trifloxystrobin (III-5), (1-[3-chloro-2-[[1-(4-chlorophenyl)pyrazol-3-yl]oxymethyl]phenyl]-4-methyl-tetrazol-5-one) (III-6), 1-[3-bromo-2-[[1-(4-chlorophenyl)pyrazol-3-yl]oxymethyl]phenyl]-4-methyl-tetrazol-5-one (III-7), 1-[2-[[1-(4-chlorophenyl)pyrazol-3-yl]oxymethyl]-3-methyl-phenyl]-4-methyl-tetrazol-5-one (III-8), 1-[2-[[1-(4-chlorophenyl)pyrazol-3-yl]oxymethyl]-3-fluoro-phenyl]-4-methyl-tetrazol-5-one (III-9), 1-[2-[[1-(2,4-dichlorophenyl)pyrazol-3-yl]oxymethyl]-3-fluoro-phenyl]-4-methyl-tetrazol-5-one (III-10), 1-[3-cyclopropyl-2-[[2-methyl-4-(1-methylpyrazol-3-yl)phenoxy]methyl]phenyl]-4-methyl-tetrazol-5-one (III-11), 1-[3-(difluoromethoxy)-2-[[2-methyl-4-(1-methylpyrazol-3-yl)phenoxy]methyl]phenyl]-4-methyl-tetrazol-5-one (III-12), 1-methyl-4-[3-methyl-2-[[2-methyl-4-(1-methylpyrazol-3-yl)phenoxy]methyl]phenyl]tetrazol-5-one (III-13), (Z,2E)-5-[1-(2,4-dichlorophenyl)pyrazol-3-yl]-oxy-2-methoxyimino-N,3-dimethyl-pent-3-enamide (III-14), (Z,2E)-5-[1-(4-chlorophenyl)pyrazol-3-yl]oxy-2-methoxyimino-N,3-dimethyl-pent-3-enamide (III-15);

- inhibitors of complex II: benzovindiflupyr (III-16), bixafen (III-17), boscalid (III-18), fluxapyroxad (III-19), isopyrazam (III-20);

- other respiration inhibitors: ametoctradin (III-21);

- B) Sterol biosynthesis inhibitors (SBI fungicides)
- C14 demethylase inhibitors: triazoles: difenoconazole (III-22), epoxiconazole (III-23), fluquinconazole (III-24), metconazole (III-25), propiconazole (III-26), prothioconazole (III-27), tebuconazole (III-28); imidazoles: prochloraz (III-29);
- 5 - Delta14-reductase inhibitors: fenpropimorph (III-30);
- C) Nucleic acid synthesis inhibitors
- phenylamides or acyl amino acid fungicides: benalaxyl (III-31), metalaxyl (III-32);
- D) Inhibitors of cell division and cytoskeleton
- 10 - tubulin inhibitors: carbendazim (III-33), thiophanate-methyl (III-34);
- other cell division inhibitors: metrafenone (III-35);
- E) Inhibitors of amino acid and protein synthesis
- methionine synthesis inhibitors: cyprodinil (III-36), pyrimethanil (III-37);
- F) Signal transduction inhibitors
- 15 - MAP / histidine kinase inhibitors: fluoroimide (III-38), iprodione (III-39), vinclozolin (III-40), fludioxonil (III-41);
- G) Lipid and membrane synthesis inhibitors
- phospholipid biosynthesis and cell wall deposition: dimethomorph (III-42), mandipropamid (III-43), iprovalicarb (III-44);
- 20 - inhibitors of oxysterol binding protein: oxathiapiprolin (III-45), 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}phenyl methanesulfonate (III-46), 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate (III-47);
- 25 H) Inhibitors with Multi Site Action
- thio- and dithiocarbamates: mancozeb (III-48), metiram (III-49);
 - organochlorine compounds: chlorothalonil (III-50);
 - guanidines and others: dithianon (III-51); and
- K) Unknown mode of action
- 30 picarbutrazox (III-52), 3-(4,4-difluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline (III-53);
- or the N-oxides or agriculturally acceptable salts of each of the compounds.
2. The compositions according to claim 1, wherein component II is compound II-1.
 3. The compositions according to claim 1, wherein component II is compound II-2.
 - 35 4. The compositions according to claim 1, wherein component III is selected from compounds III-1, III-2, III-4, III-5, III-6, III-7, III-8, III-9, III-10, III-11, III-12, III-13, III-14 and III-15.

5. The compositions according to any one of claims 1 to 4, wherein component I is compound I-1.
6. The compositions according to any one of claims 1 to 4, wherein component I is compound I-2.
- 5 7. The compositions according to any one of claims 1 to 4, wherein component I is compound I-3.
8. The compositions according to any one of claims 1 to 7, wherein the weight ratio of component I to component II is from 1000:1 to 1:1000.
9. The compositions according to any one of claims 1 to 8, wherein the weight ratio of
10 component I to component III is from 1000:1 to 1:1000.
10. The compositions according to any one of claims 1 to 9, wherein the weight ratio of component II to component III is from 1000:1 to 1:1000.
11. The compositions according to any one of claims 1 to 10, further comprising a solvent and/or at least one agrochemical auxiliary.
- 15 12. A use of a composition as defined in any one of claims 1 to 11 for combating phytopathogenic fungi.
13. The use of claim 12 for combating phytopathogenic fungi in cereals.
14. The use of claim 12 for combating phytopathogenic fungi in soybeans.
15. A method for combating phytopathogenic harmful fungi, comprising treating the fungi or
20 the materials, plants, the soil or seeds to be protected against fungal attack with an effective amount of a composition as defined in any of the claims 1 to 11.
16. Seed, coated with the components I, II and III of the compositions as defined in any one of claims 1 to 11, in an amount of from 0.1 to 10 kg active substances per 100 kg of seed.
17. Plant propagation material, coated with the components I, II and III of the compositions as
25 defined in any one of claims 1 to 11.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2017/062454

A. CLASSIFICATION OF SUBJECT MATTER
INV. A01P3/00 A01N43/653 A01N43/56
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2015/124543 A1 (BAYER CROPSOURCE AG [DE]) 27 August 2015 (2015-08-27) cited in the application the whole document	1-17
Y	WO 2013/007767 A1 (BASF SE [DE]; DIETZ JOCHEN [DE]; RIGGS RICHARD [DE]; BOUDET NADEGE [DE]) 17 January 2013 (2013-01-17) cited in the application the whole document	1-17
Y,P	WO 2016/174042 A1 (BASF AGRO B.V.) 3 November 2016 (2016-11-03) the whole document	1-17
Y	WO 2015/049168 A1 (SYNGENTA PARTICIPATIONS AG) 9 April 2015 (2015-04-09) the whole document	1-17
	-/--	



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

21 July 2017

Date of mailing of the international search report

02/08/2017

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Breimaier, Waltraud

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2017/062454

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2014/016279 A1 (SYNGENTA PARTICIPATIONS AG) 30 January 2014 (2014-01-30) the whole document -----	1-17
Y	WO 2015/197393 A1 (BASF SE) 30 December 2015 (2015-12-30) the whole document -----	1-17
Y	WO 2014/095994 A1 (BASF SE) 26 June 2014 (2014-06-26) the whole document -----	1-17

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2017/062454

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
WO 2015124543	A1	27-08-2015	CN	106163281 A		23-11-2016
			EP	3107392 A1		28-12-2016
			US	2017055531 A1		02-03-2017
			WO	2015124543 A1		27-08-2015

WO 2013007767	A1	17-01-2013	AR	087194 A1		26-02-2014
			AU	2012282501 A1		30-01-2014
			BR 112014000625	A2		13-09-2016
			CA	2840286 A1		17-01-2013
			CL	2014000067 A1		11-07-2014
			CN	103649057 A		19-03-2014
			CN	105152899 A		16-12-2015
			CO	6890103 A2		10-03-2014
			CR	20130673 A		05-03-2014
			DK	2731935 T3		06-06-2016
			EA	201400125 A1		30-06-2014
			EP	2731935 A1		21-05-2014
			ES	2570187 T3		17-05-2016
			HU	E027461 T2		28-09-2016
			JP	5789340 B2		07-10-2015
			JP	2014520832 A		25-08-2014
			KR	20140022483 A		24-02-2014
			MA	35344 B1		01-08-2014
			NZ	619937 A		24-12-2015
			PE	08262014 A1		09-07-2014
			PL	2731935 T3		30-09-2016
			US	2014155262 A1		05-06-2014
			US	2017081296 A1		23-03-2017
			UY	34203 A		28-02-2013
			WO	2013007767 A1		17-01-2013

WO 2016174042	A1	03-11-2016	NONE			

WO 2015049168	A1	09-04-2015	AR	099642 A1		10-08-2016
			CN	105611836 A		25-05-2016
			EA	201600306 A1		30-09-2016
			EP	3051945 A1		10-08-2016
			JP	2016531883 A		13-10-2016
			US	2016235064 A1		18-08-2016
			WO	2015049168 A1		09-04-2015

WO 2014016279	A1	30-01-2014	AR	091905 A1		11-03-2015
			AU	2013295098 A1		05-02-2015
			CA	2879911 A1		30-01-2014
			CN	104486949 A		01-04-2015
			CN	106614610 A		10-05-2017
			CN	106614611 A		10-05-2017
			CN	106614612 A		10-05-2017
			CN	106614620 A		10-05-2017
			CO	7180221 A2		09-02-2015
			EA	201500173 A1		30-07-2015
			EP	2877025 A1		03-06-2015
			JP	2015522635 A		06-08-2015
			KR	20150038337 A		08-04-2015
			TW	201417711 A		16-05-2014
			US	2015181877 A1		02-07-2015
			US	2017156325 A1		08-06-2017
			US	2017156326 A1		08-06-2017

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2017/062454

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		UY 34950 A	28-02-2014
		WO 2014016279 A1	30-01-2014

WO 2015197393 A1	30-12-2015	AU 2015281187 A1	12-01-2017
		CA 2951177 A1	30-12-2015
		CN 106793776 A	31-05-2017
		EP 3160229 A1	03-05-2017
		US 2017127676 A1	11-05-2017
		WO 2015197393 A1	30-12-2015

WO 2014095994 A1	26-06-2014	AR 094242 A1	22-07-2015
		CA 2894264 A1	26-06-2014
		CN 105050406 A	11-11-2015
		EA 201500653 A1	30-12-2015
		EP 2934147 A1	28-10-2015
		JP 2016501899 A	21-01-2016
		TW 201429398 A	01-08-2014
		US 2015344445 A1	03-12-2015
		UY 35237 A	30-06-2014
		WO 2014095994 A1	26-06-2014
