

(19)



(11)

EP 2 894 167 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

15.07.2015 Bulletin 2015/29

(51) Int Cl.:

C07K 16/22 <small>(2006.01)</small>	A61P 35/00 <small>(2006.01)</small>
A61K 39/395 <small>(2006.01)</small>	C12N 15/13 <small>(2006.01)</small>
C12N 15/63 <small>(2006.01)</small>	A61K 47/48 <small>(2006.01)</small>
A61P 37/00 <small>(2006.01)</small>	

(21) Application number: **14197237.2**

(22) Date of filing: **17.06.2010**

(84) Designated Contracting States:

AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK SM TR

Designated Extension States:

RS

- **Dubridge, Robert B.**
Belmont, CA 94002 (US)
- **Akamatsu, Yoshiko**
Palo Alto, CA 94306 (US)
- **Powers, David B.**
Fairfax, CA 94930 (US)

(30) Priority: **17.06.2009 US 218005 P**

(74) Representative: **Roques, Sarah Elizabeth**

J A Kemp
14 South Square
Gray's Inn
London WC1R 5JJ (GB)

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:

10725590.3 / 2 443 150

(71) Applicant: **AbbVie Biotherapeutics Inc.**

Redwood City, CA 94063 (US)

Remarks:

- This application was filed on 10-12-2014 as a divisional application to the application mentioned under INID code 62.
- Claims filed after the date of filing of the application/ after the date of receipt of the divisional application (Rule 68(4) EPC).

(72) Inventors:

- **Harding, Fiona**
Mountain View, CA 94041 (US)

(54) **Anti-VEGF Antibodies and their uses**

(57) The present disclosure relates to antibodies directed to vascular endothelial growth factor ("VEGF") and uses of such antibodies, for example to treat diseases associated with the activity and/or overproduction of VEGF.

EP 2 894 167 A1

Description**1. CROSS-REFERENCE TO RELATED APPLICATIONS AND SEQUENCE LISTING**

5 [0001] This application claims the benefit under 35 U.S.C. § 119(e) of U.S. provisional application no. 61/218,005, filed June 17, 2009, the contents of which are incorporated herein by reference in their entirety.

[0002] The instant application contains a Sequence Listing which has been submitted via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on June 15, 2010, is named 381493PC.txt and is 141,482 bytes in size.

2. FIELD OF THE INVENTION

10 [0003] The present invention relates to anti-VEGF antibodies, pharmaceutical compositions comprising anti-VEGF antibodies, and therapeutic uses of such antibodies.

3. BACKGROUND

15 [0004] Angiogenesis has emerged as attractive therapeutic target due to its implication in a variety of pathological conditions, including tumor growth, proliferative retinopathies, age-related macular degeneration, rheumatoid arthritis (RA), and psoriasis (Folkman et al., 1992, J. Biol. Chem. 267:10931-10934). The first indication of specific molecular angiogenic factors was based on the observation of the strong neovascular response induced by transplanted tumors. It is now known that angiogenesis is essential for the growth of most primary tumors and their subsequent metastasis. Numerous molecules have since been associated with the positive regulation of angiogenesis, including transforming growth factor (TGF)- α , TGF- β , hepatocyte growth factor (HGF), tumor necrosis factor- α , angiogenin, interleukin (IL)-8, and vascular endothelial growth factor (VEGF, also referred to as VEGFA or vascular permeability factor (VPF)) (Ferrara et al., 2003, Nature Medicine 9:669-676).

20 [0005] The VEGF proteins are important signaling proteins involved in both normal embryonic vasculogenesis (the de novo formation of the embryonic circulatory system) and abnormal angiogenesis (the growth of blood vessels from pre-existing vasculature) (Ferrara et al., 1996, Nature 380:439-442; Dvorak et al., 1995, Am. J. Pathol. 146:1029-1039). VEGF is associated with solid tumors and hematologic malignancies, interocular neovascular syndromes, inflammation and brain edema, and pathology of the female reproductive tract (Ferrara et al., 2003, Nature Medicine 9:669-676). VEGF mRNA is over-expressed in many human tumors, including those of the lung, breast, gastrointestinal tract, kidney, pancreas, and ovary (Berkman et al., 1993, J. Clin. Invest. 91:153-159). Increases in VEGF in the aqueous and vitreous humor of the eyes have been associated with various retinopathies (Aiello et al., 1994, N. Engl. J. Med. 331:1480-1487). Age-related macular degeneration (AMD), a major cause of vision loss in the elderly is due to neovascularization and vascular leakage. The localization of VEGF in the choroidal neovascular membranes in patients affected by AMD has been shown (Lopez et al., 1996, Invest. Ophthalmol. Vis. Sci. 37:855-868).

30 [0006] The VEGF gene family includes the prototypical member VEGFA, as well as VEGFB, VEGFC, VEGFD, and placental growth factor (PLGF). The human VEGFA gene is organized as eight exons separated by seven introns. At least six different isoforms of VEGF exist, VEGF₁₂₁, VEGF₁₄₅, VEGF₁₆₂, VEGF₁₆₅, VEGF_{165b}, VEGF₁₈₃, VEGF₁₈₉, and VEGF₂₀₆, where the subscripts refer to the number of amino acids remaining after signal cleavage. Native VEGF is a 45kDa homodimeric heparin-binding glycoprotein (Ferrara et al., 2003, Nature Medicine 9:669-676). VEGF (specifically VEGFA) binds to two related receptor tyrosine kinases, VEGFR-1 (also referred to as Flt-1) and VEGFR-2 (also referred to as Flk-1 or kinase domain region (KDR) or CD309). Each receptor has seven extracellular and one transmembrane region. VEGF also binds to the neuropilins NRP1 (also referred to as vascular endothelial cell growth factor 165 receptor (VEGF165R) or CD304) and NRP2 also referred to as vascular endothelial cell growth factor 165 receptor 2 (VEGF165R2)).

35 [0007] Given its central role in regulating angiogenesis, VEGF provides an attractive target for therapeutic intervention. Indeed, a variety of therapeutic strategies aimed at blocking VEGF or its receptor signaling system are currently being developed for the treatment of neoplastic diseases. The anti-VEGF antibody bevacizumab, also referred to as rhuMab VEGF or Avastin®, is a recombinant humanized anti-VEGF monoclonal antibody created and marketed by Genentech (Presta et al., 1997, Cancer Res. 57:4593-4599). In order to construct bevacizumab the complementarity-determining regions (CDRs) of the murine anti-VEGF monoclonal antibody A.4.6.1 were grafted onto human frameworks and an IgG constant region. Additional mutations outside the CDRs were then introduced into the molecule to improve binding, affording an antibody in which ~93% of the amino acid sequence is derived from human IgG₁ and ~7% of the sequence is derived from the murine antibody A.4.6.1. Bevacizumab has a molecular mass of about 149,000 Daltons and is glycosylated.

40 [0008] Ranibizumab is an affinity matured Fab fragment derived from bevacizumab. Ranibizumab has a higher

affinity for VEGF and also is smaller in size, allowing it to better penetrate the retina, and thus treat the ocular neovascularization associated with AMD (Lien and Lowman, In: Chernajovsky, 2008, Therapeutic Antibodies. Handbook of Experimental Pharmacology 181, Springer-Verlag, Berlin Heidelberg 131-150). Ranibizumab was developed and is marketed by Genentech under the trade name Lucentis®.

5 **[0009]** Treatment of cancer patients with a regimen that includes Avastin® can result in side effects including hypertension, proteinuria, thromboembolic events, bleeding and cardiac toxicity (Blowers & Hall, 2009, Br. J. Nurs. 18(6):351-6, 358). Also, despite being a humanized antibody, bevacizumab can elicit an immune response when administered to humans. Such an immune response may result in an immune complex-mediated clearance of the antibodies or fragments from the circulation, and make repeated administration unsuitable for therapy, thereby reducing the therapeutic benefit
10 to the patient and limiting the re-administration of the antibody.

[0010] Accordingly, there is a need to provide improved anti-VEGF antibodies or fragments that overcome one or more of these problems, for example, by generating variants with higher affinity than bevacizumab that can be administered at reduced dosages, or variants with reduced immunogenicity and other side-effects as compared to bevacizumab.

15 **[0011]** Citation or identification of any reference in Section 3 or in any other section of this application shall not be construed as an admission that such reference is available as prior art to the present disclosure.

4. SUMMARY

20 **[0012]** The present disclosure relates to variants of the anti-VEGF antibody bevacizumab with reduced immunogenicity and/or improved affinity towards VEGF as compared to bevacizumab or ranibizumab. Bevacizumab has three heavy chain CDRs, referred to herein (in amino- to carboxy-terminal order) as CDR-H1, CDR-H2, and CDR-H3, and three light chain CDRs, referred to herein (in amino- to carboxy-terminal order) as CDR-L1, CDR-L2, and CDR-L3. The sequences of the bevacizumab CDRs are shown in Figures 1A and 1B, and their numbering is set forth in Table 1 (for heavy chain CDRs) and Table 2 (for light chain CDRs). A related antibody, ranibizumab, was generated by affinity maturation of
25 bevacizumab. Ranibizumab has identical CDR-L1, CDR-L2, CDR-L3 and CDR-H2 sequences to bevacizumab, but varies in its CDR-H1 and CDR-H3 sequences from those of bevacizumab. The heavy and light chain sequences of ranibizumab are shown in Figure 1C, and the CDRs are set forth in Figure 1D.

[0013] The antibodies of the disclosure generally have at least one amino acid substitution in at least one heavy chain CDR as compared to bevacizumab and ranibizumab.

30 **[0014]** In certain aspects, the anti-VEGF antibodies include at least one substitution as compared to bevacizumab or ranibizumab selected from N31F in CDR-H1; K64S in CDR-H2; K64Q in CDR-H2; Y53F in CDR-H2; H97E in CDR-H3; H97D in CDR-H3; H97P in CDR-H3; Y98F in CDR-H3; Y99E in CDR-H3; Y99D in CDR-H3; S100aG in CDR-H3, and T51A in CDR-L2. In other aspects, the anti-VEGF antibodies include at least one substitution selected from Tables 8 and 9. Additional mutations that can be incorporated into the improved affinity variant antibodies can be candidate
35 deimmunizing substitutions, such as those described in Table 6, as well as other mutations, e.g., substitutions, that do not destroy the ability of the antibodies to bind to VEGF, including but not limited to the mutations described in Tables 10 and 11, or known mutations, such as the mutations described in Tables 12-1 to 12-9 and 13. Yet further mutations that can be incorporated include but are not limited to the mutations described in Tables 14-16.

40 **[0015]** In specific embodiments, the anti-VEGF antibodies of the disclosure include a combination of substitutions selected from Table 7, and optionally one or more additional mutations, e.g., candidate deimmunizing substitutions, such as those described in Table 6, as well as other mutations, e.g., substitutions, that do not destroy the ability of the antibodies to bind to VEGF, including but not limited to the mutations described in Tables 10 and 11, or known mutations, such as the mutations described in Tables 12-1 to 12-9 and 13. Yet further mutations that can be incorporated into the
45 anti-VEGF antibodies of the disclosure include but are not limited to the mutations described in Tables 14-16.

[0016] In other embodiments, the anti-VEGF antibodies of the disclosure include one or more of the following CDR substitutions: K64S (CDR-H2), K64Q (CDR-H2), Y53F and K64Q (CDR-H2), H97E and Y98F (CDR-H3), or T51A (CDR-L2). The anti-VEGF antibodies can also optionally include one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9, or 13-16.

50 **[0017]** Further CDR substitutions can include N31F (CDR-H1), H97E (CDR-H3), H97D (CDR-H3), H97P (CDR-H3), Y99E (CDR-H3), Y99D (CDR-H3), S100aG (CDR-H3) wherein position 3 in CDR-H3 optionally is not tyrosine, T28P, N31F, N31G and N31M (CDR-H1), H97A, H97Q, H97S, H97T, S100aD, S100aE, and S100Av (CDR-H3), T30W, T30R or T30Q (CDR-H1), Y53F, T58F, A61G, A61K, A61R, A61H, A61Y, K64G, K64E, R65L, R65T, R65A, R65E, and R65D (CDR-H2), and Y98F and Y100eF (CDR-H3). The CDRs optionally contain one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9 and 13.

55 **[0018]** Yet further substitutions can include heavy chain CDR substitutions including a combination of substitutions selected from: (a) N31F in CDR-H1, H97D in CDR-H3, Y99D in CDR-H3, and S100aG in CDR-H3; (b) N31F in CDR-H1, H97P in CDR-H3, Y99D in CDR-H3, and S100aG in CDR-H3; (c) N31F in CDR-H1, H97P in CDR-H3, and Y99E in CDR-H3; (d) N31F in CDR-H1, H97E in CDR-H3, and Y99E in CDR-H3; (e) N31F in CDR-H1, H97D in CDR-H3, and

Y99E in CDR-H3; (f) N31F in CDR-H1, H97E in CDR-H3, Y99D in CDR-H3, and S100aG in CDR-H3; (g) N31F in CDR-H1, Y99D in CDR-H3, and S100aG in CDR-H3; (h) N31F in CDR-H1, H97P in CDR-H3, and Y99D in CDR-H3; (i) N31F in CDR-H1, H97D in CDR-H3, and S100aG in CDR-H3; (j) N31F in CDR-H1 and S100aG in CDR-H3; or (k) N31F in CDR-H1, H97P in CDR-H3, and S100aG in CDR-H3. Further optional substitutions can include one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9 and 13.

[0019] Still further heavy chain substitutions can include at least one substitution selected from A61F in CDR-H2, A61E in CDR-H2, A61D in CDR-H2, D62L in CDR-H2, D62G in CDR-H2, D62Q in CDR-H2, D62T in CDR-H2, D62K in CDR-H2, D62R in CDR-H2, D62E in CDR-H2, D62H in CDR-H2, K64S in CDR-H2, K64V in CDR-H2, K64Q in CDR-H2, R65V in CDR-H2, R65F in CDR-H2, R65H in CDR-H2, R65N in CDR-H2, R65S in CDR-H2, R65Q in CDR-H2, R65K in CDR-H2, R65I in CDR-H2, and Y98H in CDR-H3. Optionally, one or more additional mutations or combinations of mutations can be included as selected from one or more of Tables 7, 8, 9, 10, 11, 12-1 to 12-9 and 13.

[0020] In certain aspects, the antibodies of the disclosure have VH and VL sequences having at least 80% sequence identity (and in certain embodiments, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99% sequence identity) to the VH and VL sequences of bevacizumab or ranibizumab, and include at least one amino acid substitution in at least one CDR as compared to bevacizumab or ranibizumab. In other aspects, the antibodies of the disclosure have VH and VL sequences having at least 80% sequence identity (and in certain embodiments, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99% sequence identity) to the VH and VL sequences of bevacizumab or ranibizumab, and include at least one amino acid substitution in at least one framework region as compared to bevacizumab or ranibizumab. In specific embodiments, the percentage sequence identity for the heavy chain and the light chain compared to the VH and VL sequences of bevacizumab or ranibizumab is independently selected from at least 80%, at least 85%, at least 90%, at least 95% sequence identity, or at least 99% sequence identity. In certain aspects, the antibodies of the disclosure have VH and/or VL sequences having at least 95%, at least 98% or at least 99% sequence identity to the VH and/or VL sequences of bevacizumab or ranibizumab.

[0021] In certain aspects, the antibodies of the disclosure have up to 17 amino acid substitutions in their CDRs as compared to bevacizumab or ranibizumab. Variant antibodies with 17 amino acid substitutions that maintain their target binding capability have been generated by Bostrom et al., 2009, Science 323:1610-14.

[0022] In specific embodiments, an anti-VEGF antibody of the disclosure has, independently:

- up to one, up to two, up to three, up to four, up to five, up to six, up to seven, up to eight, up to nine or up to ten CDR-H1 substitutions as compared to the corresponding CDR of bevacizumab or of ranibizumab;
- up to one, up to two, up to three, up to four, up to five, up to six, up to seven, up to eight, up to nine, up to ten, up to eleven, up to twelve, up to thirteen, up to fourteen, up to fifteen, up to sixteen or up to seventeen CDR-H2 substitutions as compared to the corresponding CDR of bevacizumab or of ranibizumab;
- up to one, up to two, up to three, up to four, up to five, up to six, up to seven, up to eight, up to nine, up to ten, up to eleven, up to twelve, up to thirteen or up to fourteen CDR-H3 substitutions as compared to the corresponding CDR of bevacizumab or of ranibizumab;
- up to one, up to two, up to three, up to four, up to five, up to six, up to seven, up to eight, up to nine, up to ten or up to eleven CDR-L1 substitutions as compared to the corresponding CDR of bevacizumab or of ranibizumab;
- up to one, up to two, up to three, up to four, up to five, up to six or up to seven CDR-L2 substitutions as compared to the corresponding CDR of bevacizumab or of ranibizumab; and
- up to one, up to two, up to three, up to four, up to five, up to six, up to seven, up to eight or up to nine CDR-L3 substitutions as compared to the corresponding CDR of bevacizumab or of ranibizumab.

[0023] The present disclosure further provides pharmaceutical compositions comprising modified anti-VEGF antibodies. In some aspects, the pharmaceutical compositions have increased affinity to VEGF and/or reduced immunogenicity as compared to bevacizumab or ranibizumab.

[0024] Nucleic acids comprising nucleotide sequences encoding the anti-VEGF antibodies of the disclosure are provided herein, as are vectors comprising the nucleic acids. Additionally, prokaryotic and eukaryotic host cells transformed with a vector comprising a nucleotide sequence encoding an anti-VEGF antibody are provided herein, as well as eukaryotic (such as mammalian) host cells engineered to express the nucleotide sequences. Methods of producing anti-VEGF antibodies by culturing host cells are also provided.

[0025] The anti-VEGF antibodies of the disclosure are useful in the treatment of cancers (e.g., colon carcinoma, rectal carcinoma, non-small cell lung cancer, and breast cancer), retinal diseases (e.g., age-related macular degeneration

("AMD"), and immune disorders (e.g., rheumatoid arthritis).

[0026] In certain aspects, the anti-VEGF antibodies of the disclosure can be used in reduced dosages as compared to bevacizumab or ranibizumab, e.g., at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80% or at least 90% lower dosages.

5 **[0027]** It should be noted that the indefinite articles "a" and "an" and the definite article "the" are used in the present application, as is common in patent applications, to mean one or more unless the context clearly dictates otherwise. Further, the term "or" is used in the present application, as is common in patent applications, to mean the disjunctive "or" or the conjunctive "and."

10 **[0028]** All publications mentioned in this specification are herein incorporated by reference. Any discussion of documents, acts, materials, devices, articles or the like that has been included in this specification is solely for the purpose of providing a context for the present disclosure. It is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present disclosure as it existed anywhere before the priority date of this application.

15 **[0029]** The features and advantages of the disclosure will become further apparent from the following detailed description of embodiments thereof.

5. BRIEF DESCRIPTION OF THE TABLES AND FIGURES

20 **[0030]**

Table 1 shows the numbering of the amino acids in the heavy chain CDRs of bevacizumab. CDRs 1-3 are disclosed as SEQ ID NOS:3-5, respectively.

25 **Table 2** shows the numbering of the amino acids in the light chain CDRs of bevacizumab. CDRs 1-3 are disclosed as SEQ ID NOS:6-8, respectively.

Table 3 shows bevacizumab VL peptides that were tested for immunogenicity.

30 **Table 4** shows bevacizumab VH peptides that were tested for immunogenicity.

Table 5 shows identified CD4⁺ T cell epitope regions in bevacizumab. CDR regions are underlined.

35 **Table 6** shows candidate mutations in CDR-H2 and CDR-H3 for lowering immunogenicity of bevacizumab. The numbering of the amino acids in Table 6 corresponds to Kabat numbering in the bevacizumab heavy chain.

40 **Table 7** shows heavy chain CDR amino acid substitutions in bevacizumab resulting improved K_D as analyzed by surface plasmon resonance. Δk_{on} refers to fold improvement in k_{on} (mutant/WT). Δk_{off} refers to fold improvement in k_{off} (WT/mutant). ΔK_D refers to the improvement in the K_D in the mutant relative to wild type. The numbering of the amino acids in Table 7 corresponds to Kabat numbering in the bevacizumab heavy chain.

Table 8 shows mutations in the bevacizumab heavy chain CDRs that preliminary binding studies indicate increase the affinity towards VEGF (data not shown). The numbering of the amino acids in Table 8 corresponds to Kabat numbering in the bevacizumab heavy chain.

45 **Table 9** shows mutations in the bevacizumab heavy chain CDRs that preliminary studies indicate increase the affinity towards VEGF (data not shown). The numbering of the amino acids in Table 9 corresponds to Kabat numbering in the bevacizumab heavy chain.

50 **Table 10** shows mutations in the bevacizumab heavy chain CDRs that do not impact binding and can be incorporated into the antibodies of the disclosure. The numbering of the amino acids in Table 10 corresponds to Kabat numbering in the bevacizumab heavy chain.

55 **Table 11** shows mutations in the bevacizumab light chain CDRs that do not impact binding and can be incorporated into the antibodies of the disclosure. The numbering of the amino acids in Table 11 corresponds to Kabat numbering in the bevacizumab light chain.

Tables 12-1 to 12-9 show known mutations in bevacizumab heavy chain CDRs that can be incorporated into the antibodies of the disclosure. Each row in Tables 12-1 to 12-9 includes a distinct known variant. For each variant,

the known CDR sequences are shaded. The sequence identifiers for each variant identified in Tables 12-1 to 12-9 are set forth in Tables 20-1 to 20-9, respectively. The CDR-H1 column provides a partial sequence of CDR-H1. The final asparagine of CDR-H1 is not shown. This partial sequence corresponds to SEQ ID NO:411. Although known mutations in CDR-H1 are shown in the context of this partial sequence, it is noted that the mutations exist in the context of the full length CDR.

Table 13 shows known mutations in bevacizumab light chain CDRs that can be incorporated into the antibodies of the disclosure. Each row in Table 13 includes a distinct known variant. For each variant, the known CDR sequences are shaded. The sequence identifiers for each variant identified in Table 13 is set forth in Table 20-10.

Table 14 shows bevacizumab CDR2 VH peptides that were tested for immunogenicity, wherein residues unchanged from SEQ ID NO:62 are indicated by a blank box. CD4+ T cell assay results are also provided.

Table 15 shows bevacizumab CDR3 VH peptides that were tested for immunogenicity, wherein residues unchanged from SEQ ID NO:74 are indicated by a blank box. CD4+ T cell assay results are also provided.

Table 16 shows bevacizumab CDR2 VL peptides that were tested for immunogenicity, wherein residues unchanged from SEQ ID NO:25 are indicated by a blank box. CD4+ T cell assay results are also provided.

Table 17 shows selected epitope modifications for the three CD4+ T cell epitopes in bevacizumab.

Table 18 shows single variable region mutants and their associated mean fluorescence intensity (MFI) score.

Table 19 shows combined variable region mutants and their associated EC₅₀.

Tables 20-1 to 20-10 show the SEQ ID NOS, where known, corresponding to the CDRs of the bevacizumab variants listed in Tables 12-1 to 12-9 and Table 13, respectively. N/A indicates an unknown CDR sequence.

Figures 1A-1D. **Figure 1A** shows the amino acid sequences of the bevacizumab heavy and light chain variable regions, SEQ ID NO:1 and SEQ ID NO:2, respectively, with CDR regions in bold, underlined text. **Figure 1B** shows the CDR sequences and corresponding sequence identifiers of bevacizumab. **Figure 1C** shows the amino acid sequences of the ranibizumab heavy and light chains, SEQ ID NO:9 and SEQ ID NO:10, respectively, with CDR regions in bold, underlined text. **Figure 1D** shows the CDR sequences and corresponding sequence identifiers of ranibizumab.

Figures 2A-2B show bevacizumab VL peptide responses. Figure 2A shows percent of donor responses to each VL peptide with a stimulation index of 2.95 or greater. N = 99 donors. Figure 2B shows the average stimulation index for all 99 donors for each peptide plus or minus standard error.

Figures 3A-3B show bevacizumab VH peptide responses. Figure 3A shows percent of donor responses to each VH peptide with a stimulation index of 2.95 or greater. N = 99 donors. Figure 3B shows the average stimulation index for all 99 donors for each peptide plus or minus standard error.

Figures 4A-4C show CD4+ T cell responses to mutant bevacizumab epitope peptides. Average responses to the unmodified parent epitope sequences are indicated with open marks. Large circles indicate selected changes referred to in **Table 17**. **Figure 4A** is directed to VH CDR2 peptides; **Figure 4B** is directed to VH CDR3 peptides; and **Figure 4C** is directed to VL CDR2 peptides.

6. DETAILED DESCRIPTION

6.1 ANTI-VEGF ANTIBODIES

[0031] The present disclosure provides anti-VEGF antibodies. Unless indicated otherwise, the term "antibody" (Ab) refers to an immunoglobulin molecule that specifically binds to, or is immunologically reactive with, a particular antigen, and includes polyclonal, monoclonal, genetically engineered and otherwise modified forms of antibodies, including but not limited to chimeric antibodies, humanized antibodies, heteroconjugate antibodies (e.g., bispecific antibodies, diabodies, triabodies, and tetrabodies), and antigen binding fragments of antibodies, including e.g., Fab', F(ab')₂, Fab, Fv, rIgG, and scFv fragments. Moreover, unless otherwise indicated, the term "monoclonal antibody" (mAb) is meant to

include both intact molecules, as well as, antibody fragments (such as, for example, Fab and F(ab')₂ fragments) which are capable of specifically binding to a protein. Fab and F(ab')₂ fragments lack the Fc fragment of intact antibody, clear more rapidly from the circulation of the animal, and may have less non-specific tissue binding than an intact antibody (Wahl et al., 1983, J. Nucl. Med. 24:316).

5 **[0032]** The term "scFv" refers to a single chain Fv antibody in which the variable domains of the heavy chain and the light chain from a traditional antibody have been joined to form one chain.

[0033] References to "VH" refer to the variable region of an immunoglobulin heavy chain of an antibody, including the heavy chain of an Fv, scFv, or Fab. References to "VL" refer to the variable region of an immunoglobulin light chain, including the light chain of an Fv, scFv, dsFv or Fab. Antibodies (Abs) and immunoglobulins (Igs) are glycoproteins having the same structural characteristics. While antibodies exhibit binding specificity to a specific target, immunoglobulins include both antibodies and other antibody-like molecules which lack target specificity. Native antibodies and immunoglobulins are usually heterotetrameric glycoproteins of about 150,000 Daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each heavy chain has at the amino terminus a variable domain (VH) followed by a number of constant domains. Each light chain has a variable domain at the amino terminus (VL) and a constant domain at the carboxy terminus.

15 **[0034]** The anti-VEGF antibodies of the disclosure bind to human VEGF and inhibit VEGF receptor activity in a cell.

[0035] The anti-VEGF antibodies of the disclosure contain complementarity determining regions (CDRs) that are related in sequence to the CDRs of the antibody bevacizumab (also known as Avastin®) and/or ranibizumab (also known as Lucentis®).

20 **[0036]** CDRs are also known as hypervariable regions both in the light chain and the heavy chain variable domains. The more highly conserved portions of variable domains are called the framework (FR). As is known in the art, the amino acid position/boundary delineating a hypervariable region of an antibody can vary, depending on the context and the various definitions known in the art. Some positions within a variable domain may be viewed as hybrid hypervariable positions in that these positions can be deemed to be within a hypervariable region under one set of criteria while being deemed to be outside a hypervariable region under a different set of criteria. One or more of these positions can also be found in extended hypervariable regions. The disclosure provides antibodies comprising modifications in these hybrid hypervariable positions. The variable domains of native heavy and light chains each comprise four FR regions, largely by adopting a β-sheet configuration, connected by three CDRs, which form loops connecting, and in some cases forming part of, the β-sheet structure. The CDRs in each chain are held together in close proximity by the FR regions in the order FR1-CDR1-FR2-CDR2-FR3-CDR3-FR4 and, with the CDRs from the other chain, contribute to the formation of the target binding site of antibodies (see Kabat et al., Sequences of Proteins of Immunological Interest (National Institute of Health, Bethesda, Md. 1987). As used herein, numbering of immunoglobulin amino acid residues is done according to the immunoglobulin amino acid residue numbering system of Kabat *et al.*, unless otherwise indicated.

30 **[0037]** The sequences of the heavy and light chain variable regions of bevacizumab are represented by SEQ ID NO:1 and SEQ ID NO:2, respectively. The sequences of the heavy and light chain variable regions are also depicted in Figure 1A. The sequences of the CDRs of bevacizumab, and their corresponding identifiers, are presented in Figure 1B. Any nucleotide sequences encoding SEQ ID NO:1 or SEQ ID NO:2 can be used in the compositions and methods of the present disclosure.

35 **[0038]** The sequences of the heavy and light chains of ranibizumab are represented by SEQ ID NO:9 and SEQ ID NO:10, respectively. The sequences of the heavy and light chains are also depicted in Figure 1C. The sequences of the CDRs of ranibizumab, and their corresponding identifiers, are presented in Figure 1D. Any nucleotide sequences encoding SEQ ID NO:9 or SEQ ID NO:10 can be used in the compositions and methods of the present disclosure.

40 **[0039]** The present disclosure further provides anti-VEGF antibody fragments comprising CDR sequences that are related to the CDR sequences of bevacizumab and ranibizumab. The term "antibody fragment" refers to a portion of a full-length antibody, generally the target binding or variable region. Examples of antibody fragments include Fab, Fab', F(ab')₂ and Fv fragments. An "Fv" fragment is the minimum antibody fragment which contains a complete target recognition and binding site. This region consists of a dimer of one heavy and one light chain variable domain in a tight, non-covalent association (VH-VL dimer). It is in this configuration that the three CDRs of each variable domain interact to define a target binding site on the surface of the VH-VL dimer. Often, the six CDRs confer target binding specificity to the antibody. However, in some instances even a single variable domain (or half of an Fv comprising only three CDRs specific for a target) can have the ability to recognize and bind target. "Single-chain Fv" or "scFv" antibody fragments comprise the VH and VL domains of an antibody in a single polypeptide chain. Generally, the Fv polypeptide further comprises a polypeptide linker between the VH and VL domain which enables the scFv to form the desired structure for target binding. "Single domain antibodies" are composed of a single VH or VL domains which exhibit sufficient affinity to the target. In a specific embodiment, the single domain antibody is a camelid antibody (see, e.g., Riechmann, 1999, Journal of Immunological Methods 231:25-38).

55 **[0040]** The Fab fragment contains the constant domain of the light chain and the first constant domain (CH₁) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxyl terminus of the

heavy chain CH₁ domain including one or more cysteines from the antibody hinge region. F(ab') fragments are produced by cleavage of the disulfide bond at the hinge cysteines of the F(ab')₂ pepsin digestion product. Additional chemical couplings of antibody fragments are known to those of ordinary skill in the art.

5 **[0041]** In certain embodiments, the anti-VEGF antibodies of the disclosure are monoclonal antibodies. The term "monoclonal antibody" as used herein is not limited to antibodies produced through hybridoma technology. The term "monoclonal antibody" refers to an antibody that is derived from a single clone, including any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced. Monoclonal antibodies useful in connection with the present disclosure can be prepared using a wide variety of techniques known in the art including the use of hybridoma, recombinant, and phage display technologies, or a combination thereof. The anti-VEGF antibodies of the disclosure include chimeric, primatized, humanized, or human antibodies.

10 **[0042]** The anti-VEGF antibodies of the disclosure can be chimeric antibodies. The term "chimeric" antibody as used herein refers to an antibody having variable sequences derived from a non-human immunoglobulin, such as rat or mouse antibody, and human immunoglobulin constant regions, typically chosen from a human immunoglobulin template. Methods for producing chimeric antibodies are known in the art. See, e.g., Morrison, 1985, *Science* 229(4719):1202-7; Oi et al., 1986, *BioTechniques* 4:214-221; Gillies et al., 1985, *J. Immunol. Methods* 125:191-202; U.S. Pat. Nos. 5,807,715; 4,816,567; and 4,816,397, which are incorporated herein by reference in their entireties.

15 **[0043]** The anti-VEGF antibodies of the disclosure can be humanized. "Humanized" forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other target-binding subdomains of antibodies) which contain minimal sequences derived from non-human immunoglobulin. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin sequence. The humanized antibody can also comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin consensus sequence. Methods of antibody humanization are known in the art. See, e.g., Riechmann et al., 1988, *Nature* 332:323-7; U.S. Patent Nos. 5,530,101; 5,585,089; 5,693,761; 5,693,762; and 6,180,370 to Queen et al.; EP239400; PCT publication WO 91/09967; U.S. Patent No. 5,225,539; EP592106; EP519596; Padlan, 1991, *Mol. Immunol.*, 28:489-498; Studnicka et al., 1994, *Prot. Eng.* 7:805-814; Roguska et al., 1994, *Proc. Natl. Acad. Sci.* 91:969-973; and U.S. Patent No. 5,565,332, all of which are hereby incorporated by reference in their entireties.

20 **[0044]** The anti-VEGF antibodies of the disclosure can be human antibodies. Completely "human" anti-VEGF antibodies can be desirable for therapeutic treatment of human patients. As used herein, "human antibodies" include antibodies having the amino acid sequence of a human immunoglobulin and include antibodies isolated from human immunoglobulin libraries or from animals transgenic for one or more human immunoglobulin and that do not express endogenous immunoglobulins. Human antibodies can be made by a variety of methods known in the art including phage display methods using antibody libraries derived from human immunoglobulin sequences. See U.S. Patent Nos. 35 4,444,887 and 4,716,111; and PCT publications WO 98/46645; WO 98/50433; WO 98/24893; WO 98/16654; WO 96/34096; WO 96/33735; and WO 91/10741, each of which is incorporated herein by reference in its entirety. Human antibodies can also be produced using transgenic mice which are incapable of expressing functional endogenous immunoglobulins, but which can express human immunoglobulin genes. See, e.g., PCT publications WO 98/24893; WO 92/01047; WO 96/34096; WO 96/33735; U.S. Patent Nos. 5,413,923; 5,625,126; 5,633,425; 5,569,825; 5,661,016; 40 5,545,806; 5,814,318; 5,885,793; 5,916,771; and 5,939,598, which are incorporated by reference herein in their entireties. In addition, companies such as Medarex (Princeton, NJ), Astellas Pharma (Deerfield, IL), Amgen (Thousand Oaks, CA) and Regeneron (Tarrytown, NY) can be engaged to provide human antibodies directed against a selected antigen using technology similar to that described above. Completely human antibodies that recognize a selected epitope can be generated using a technique referred to as "guided selection." In this approach a selected non-human monoclonal antibody, e.g., a mouse antibody, is used to guide the selection of a completely human antibody recognizing the same epitope (Jespersen et al., 1988, *Biotechnology* 12:899-903).

45 **[0045]** The anti-VEGF antibodies of the disclosure can be primatized. The term "primatized antibody" refers to an antibody comprising monkey variable regions and human constant regions. Methods for producing primatized antibodies are known in the art. See e.g., U.S. Patent Nos. 5,658,570; 5,681,722; and 5,693,780, which are incorporated herein by reference in their entireties.

50 **[0046]** The anti-VEGF antibodies of the disclosure can be bispecific antibodies. Bispecific antibodies are monoclonal, often human or humanized, antibodies that have binding specificities for at least two different antigens. In the present disclosure, one of the binding specificities can be directed towards VEGF, the other can be for any other antigen, e.g., for a cell-surface protein, receptor, receptor subunit, tissue-specific antigen, virally derived protein, virally encoded envelope protein, bacterially derived protein, or bacterial surface protein, etc. In a specific embodiment, an antibody of the disclosure is a bispecific antibody with binding specificities for both VEGF and CD3.

55 **[0047]** The anti-VEGF antibodies of the disclosure include derivatized antibodies. For example, but not by way of limitation, derivatized antibodies are typically modified by glycosylation, acetylation, pegylation, phosphorylation, ami-

dation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein (see Section 6.6 for a discussion of antibody conjugates), etc. Any of numerous chemical modifications can be carried out by known techniques, including, but not limited to, specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, etc. Additionally, the derivative can contain one or more non-natural amino acids, e.g., using ambrx technology (see, e.g., Wolfson, 2006, Chem. Biol. 13(10):1011-2).

[0048] In yet another embodiment of the disclosure, the anti-VEGF antibodies or fragments thereof can be antibodies or antibody fragments whose sequence has been modified to alter at least one constant region-mediated biological effector function relative to the corresponding wild type sequence.

[0049] For example, in some embodiments, an anti-VEGF antibody of the disclosure can be modified to reduce at least one constant region-mediated biological effector function relative to an unmodified antibody, e.g., reduced binding to the Fc receptor (Fc γ R). Fc γ R binding can be reduced by mutating the immunoglobulin constant region segment of the antibody at particular regions necessary for Fc γ R interactions (see e.g., Canfield and Morrison, 1991, J. Exp. Med. 173:1483-1491; and Lund et al., 1991, J. Immunol. 147:2657-2662). Reduction in Fc γ R binding ability of the antibody can also reduce other effector functions which rely on Fc γ R interactions, such as opsonization, phagocytosis and antigen-dependent cellular cytotoxicity ("ADCC").

[0050] In other embodiments, an anti-VEGF antibody of the disclosure can be modified to acquire or improve at least one constant region-mediated biological effector function relative to an unmodified antibody, e.g., to enhance Fc γ R interactions (see, e.g., US 2006/0134709). For example, an anti-VEGF antibody of the disclosure can have a constant region that binds Fc γ RIIA, Fc γ RIIB and/or Fc γ RIIIA with greater affinity than the corresponding wild type constant region.

[0051] Thus, antibodies of the disclosure can have alterations in biological activity that result in increased or decreased opsonization, phagocytosis, or ADCC. Such alterations are known in the art. For example, modifications in antibodies that reduce ADCC activity are described in U.S. Patent No. 5,834,597. An exemplary ADCC lowering variant corresponds to "mutant 3" shown in Figure 4 of U.S. Patent No. 5,834,597, in which residue 236 is deleted and residues 234, 235 and 237 (using EU numbering) are substituted with alanines.

[0052] In some embodiments, the anti-VEGF antibodies of the disclosure have low levels of or lack fucose. Antibodies lacking fucose have been correlated with enhanced ADCC (activity, especially at low doses of antibody. See Shields et al., 2002, J. Biol. Chem. 277:26733-26740; Shinkawa et al., 2003, J. Biol. Chem. 278:3466-73. Methods of preparing fucose-less antibodies include growth in rat myeloma YB2/0 cells (ATCC CRL 1662). YB2/0 cells express low levels of FUT8 mRNA, which encodes α -1,6-fucosyltransferase, an enzyme necessary for fucosylation of polypeptides.

[0053] In yet another aspect, the anti-VEGF antibodies or fragments thereof can be antibodies or antibody fragments that have been modified to increase or reduce their binding affinities to the fetal Fc receptor, FcRn, for example by mutating the immunoglobulin constant region segment at particular regions involved in FcRn interactions (see, e.g., WO 2005/123780). In particular embodiments, an anti-VEGF antibody of the IgG class is mutated such that at least one of amino acid residues 250, 314, and 428 of the heavy chain constant region is substituted alone, or in any combinations thereof, such as at positions 250 and 428, or at positions 250 and 314, or at positions 314 and 428, or at positions 250, 314, and 428, with positions 250 and 428 a specific combination. For position 250, the substituting amino acid residue can be any amino acid residue other than threonine, including, but not limited to, alanine, cysteine, aspartic acid, glutamic acid, phenylalanine, glycine, histidine, isoleucine, lysine, leucine, methionine, asparagine, proline, glutamine, arginine, serine, valine, tryptophan, or tyrosine. For position 314, the substituting amino acid residue can be any amino acid residue other than leucine, including, but not limited to, alanine, cysteine, aspartic acid, glutamic acid, phenylalanine, glycine, histidine, isoleucine, lysine, methionine, asparagine, proline, glutamine, arginine, serine, threonine, valine, tryptophan, or tyrosine. For position 428, the substituting amino acid residues can be any amino acid residue other than methionine, including, but not limited to, alanine, cysteine, aspartic acid, glutamic acid, phenylalanine, glycine, histidine, isoleucine, lysine, leucine, asparagine, proline, glutamine, arginine, serine, threonine, valine, tryptophan, or tyrosine. Specific combinations of suitable amino acid substitutions are identified in Table 1 of U.S. Patent No. 7,217,797, which table is incorporated by reference herein in its entirety. Such mutations increase the antibody's binding to FcRn, which protects the antibody from degradation and increases its half-life.

[0054] In yet other aspects, an anti-VEGF antibody has one or more amino acids inserted into one or more of its hypervariable regions, for example as described in Jung and Plückthun, 1997, Protein Engineering 10(9):959-966; Yazaki et al., 2004, Protein Eng Des. Sel. 17(5):481-9. Epub 2004 Aug 17; and US 2007/0280931.

[0055] In various embodiments, the anti-VEGF antibodies or fragments thereof can be antibodies or antibody fragments that have been modified for increased expression in heterologous hosts. In certain embodiments, the anti-VEGF antibodies or fragments thereof can be antibodies or antibody fragments that have been modified for increased expression in and/or secretion from heterologous host cells. In some embodiments, the anti-VEGF antibodies or fragments thereof are modified for increased expression in bacteria, such as *E. coli*. In other embodiments, the anti-VEGF antibodies or fragments thereof are modified for increased expression in yeast (Kieke et al., 1999, Proc. Nat'l Acad. Sci. USA 96:5651-5656). In still other embodiments, the anti-VEGF antibodies or fragments thereof are modified for increased expression in insect cells. In additional embodiments, the anti-VEGF antibodies or fragments thereof are modified for

increased expression in mammalian cells, such as CHO cells.

[0056] In certain embodiments, the anti-VEGF antibodies or fragments thereof can be antibodies or antibody fragments that have been modified to increase stability of the antibodies during production. In some embodiments, the antibodies or fragments thereof can be modified to replace one or more amino acids such as asparagine or glutamine that are susceptible to nonenzymatic deamidation with amino acids that do not undergo deamidation (Huang et al., 2005, Anal. Chem. 77:1432-1439). In other embodiments, the antibodies or fragments thereof can be modified to replace one or more amino acids that is susceptible to oxidation, such as methionine, cysteine or tryptophan, with an amino acid that does not readily undergo oxidation. In still other embodiments, the antibodies or fragments thereof can be modified to replace one or more amino acids that is susceptible to cyclization, such as asparagine or glutamic acid, with an amino acid that does not readily undergo cyclization.

6.2 NUCLEIC ACIDS AND EXPRESSION SYSTEMS

[0057] The present disclosure encompasses nucleic acid molecules and host cells encoding the anti-VEGF antibodies of the disclosure.

[0058] An anti-VEGF antibody of the disclosure can be prepared by recombinant expression of immunoglobulin light and heavy chain genes in a host cell. To express an antibody recombinantly, a host cell is transfected with one or more recombinant expression vectors carrying DNA fragments encoding the immunoglobulin light and heavy chains of the antibody such that the light and heavy chains are expressed in the host cell and, optionally, secreted into the medium in which the host cells are cultured, from which medium the antibodies can be recovered. Standard recombinant DNA methodologies are used to obtain antibody heavy and light chain genes, incorporate these genes into recombinant expression vectors and introduce the vectors into host cells, such as those described in Molecular Cloning; A Laboratory Manual, Second Edition (Sambrook, Fritsch and Maniatis (eds), Cold Spring Harbor, N. Y., 1989), Current Protocols in Molecular Biology (Ausubel, F.M. et al., eds., Greene Publishing Associates, 1989) and in U.S. Patent No. 4,816,397.

[0059] In one embodiment, the anti-VEGF antibodies are similar to bevacizumab or ranibizumab but for changes in one or more CDRs. In another embodiment, the anti-VEGF antibodies are similar to bevacizumab or ranibizumab but for changes in one or more framework regions. In yet another embodiment, the anti-VEGF antibodies are similar to bevacizumab or ranibizumab but for changes in one or more CDRs and in one or more framework regions. Such antibodies are referred to herein collectively as having "bevacizumab-related" or "ranibizumab-related" sequences and are sometimes referenced simply as anti-VEGF antibodies of the disclosure. To generate nucleic acids encoding anti-VEGF antibodies, DNA fragments encoding the light and heavy chain variable regions are first obtained. These DNAs can be obtained by amplification and modification of germline DNA or cDNA encoding light and heavy chain variable sequences, for example using the polymerase chain reaction (PCR). Germline DNA sequences for human heavy and light chain variable region genes are known in the art (see, e.g., the "VBASE" human germline sequence database; see also Kabat, E. A. et al., 1991, Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242; Tomlinson et al., 1992, J. Mol. Biol. 22T:116-198; and Cox et al., 1994, Eur. J. Immunol. 24:827-836; the contents of each of which are incorporated herein by reference). A DNA fragment encoding the heavy or light chain variable region of bevacizumab or ranibizumab can be synthesized and used as a template for mutagenesis to generate a variant as described herein using routine mutagenesis techniques; alternatively, a DNA fragment encoding the variant can be directly synthesized.

[0060] Once DNA fragments encoding anti-VEGF VH and VL segments are obtained, these DNA fragments can be further manipulated by standard recombinant DNA techniques, for example to convert the variable region genes to full-length antibody chain genes, to Fab fragment genes or to a scFv gene. In these manipulations, a VL- or VH-encoding DNA fragment is operatively linked to another DNA fragment encoding another protein, such as an antibody constant region or a flexible linker. The term "operatively linked," as used in this context, is intended to mean that the two DNA fragments are joined such that the amino acid sequences encoded by the two DNA fragments remain in-frame.

[0061] The isolated DNA encoding the VH region can be converted to a full-length heavy chain gene by operatively linking the VH-encoding DNA to another DNA molecule encoding heavy chain constant regions (CH₁, CH₂, CH₃ and, optionally, CH₄). The sequences of human heavy chain constant region genes are known in the art (see, e.g., Kabat, E.A. et al., 1991, Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242) and DNA fragments encompassing these regions can be obtained by standard PCR amplification. The heavy chain constant region can be an IgG₁, IgG₂, IgG₃, IgG₄, IgA, IgE, IgM or IgD constant region, but in certain embodiments is an IgG₁ or IgG₄ constant region. For a Fab fragment heavy chain gene, the VH-encoding DNA can be operatively linked to another DNA molecule encoding only the heavy chain CH₁ constant region.

[0062] The isolated DNA encoding the VL region can be converted to a full-length light chain gene (as well as a Fab light chain gene) by operatively linking the VL-encoding DNA to another DNA molecule encoding the light chain constant region, CL. The sequences of human light chain constant region genes are known in the art (see, e.g., Kabat, E. A. et al., 1991, Sequences of Proteins of Immunological Interest, Fifth Edition (U.S. Department of Health and Human Services,

NIH Publication No. 91-3242)) and DNA fragments encompassing these regions can be obtained by standard PCR amplification. The light chain constant region can be a kappa or lambda constant region, but in certain embodiments is a kappa constant region. To create a scFv gene, the VH-and VL-encoding DNA fragments are operatively linked to another fragment encoding a flexible linker, *e.g.*, encoding the amino acid sequence (Gly₄~Ser)₃, such that the VH and VL sequences can be expressed as a contiguous single-chain protein, with the VL and VH regions joined by the flexible linker (see, *e.g.*, Bird et al., 1988, *Science* 242:423-426; Huston et al., 1988, *Proc. Natl. Acad. Sci. USA* 85:5879-5883; McCafferty et al., 1990, *Nature* 348:552-554).

[0063] To express the anti-VEGF antibodies of the disclosure, DNAs encoding partial or full-length light and heavy chains, obtained as described above, are inserted into expression vectors such that the genes are operatively linked to transcriptional and translational control sequences. In this context, the term "operatively linked" is intended to mean that an antibody gene is ligated into a vector such that transcriptional and translational control sequences within the vector serve their intended function of regulating the transcription and translation of the antibody gene. The expression vector and expression control sequences are chosen to be compatible with the expression host cell used. The antibody light chain gene and the antibody heavy chain gene can be inserted into separate vectors or, more typically, both genes are inserted into the same expression vector.

[0064] The antibody genes are inserted into the expression vector by standard methods (*e.g.*, ligation of complementary restriction sites on the antibody gene fragment and vector, or blunt end ligation if no restriction sites are present). Prior to insertion of the light or heavy chain sequences of the anti-VEGF antibodies of the disclosure, the expression vector can already carry antibody constant region sequences. For example, one approach to converting the anti-VEGF VH and VL sequences to full-length antibody genes is to insert them into expression vectors already encoding heavy chain constant and light chain constant regions, respectively, such that the VH segment is operatively linked to the CH segment(s) within the vector and the VL segment is operatively linked to the CL segment within the vector. Additionally or alternatively, the recombinant expression vector can encode a signal peptide that facilitates secretion of the antibody chain from a host cell. The antibody chain gene can be cloned into the vector such that the signal peptide is linked in-frame to the amino terminus of the antibody chain gene. The signal peptide can be an immunoglobulin signal peptide or a heterologous signal peptide (*i.e.*, a signal peptide from a non-immunoglobulin protein).

[0065] In addition to the antibody chain genes, the recombinant expression vectors of the disclosure carry regulatory sequences that control the expression of the antibody chain genes in a host cell. The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (*e.g.*, polyadenylation signals) that control the transcription or translation of the antibody chain genes. Such regulatory sequences are described, for example, in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185 (Academic Press, San Diego, CA, 1990). It will be appreciated by those skilled in the art that the design of the expression vector, including the selection of regulatory sequences may depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, *etc.* Suitable regulatory sequences for mammalian host cell expression include viral elements that direct high levels of protein expression in mammalian cells, such as promoters and/or enhancers derived from cytomegalovirus (CMV) (such as the CMV promoter/enhancer), Simian Virus 40 (SV40) (such as the SV40 promoter/enhancer), adenovirus, (*e.g.*, the adenovirus major late promoter (AdMLP)) and polyoma. For further description of viral regulatory elements, and sequences thereof, see *e.g.*, U.S. Patent No. 5,168,062 by Stinski, U.S. Patent No. 4,510,245 by Bell et al., and U.S. Patent No. 4,968,615 by Schaffner et al.,

[0066] In addition to the antibody chain genes and regulatory sequences, the recombinant expression vectors of the disclosure can carry additional sequences, such as sequences that regulate replication of the vector in host cells (*e.g.*, origins of replication) and selectable marker genes. The selectable marker gene facilitates selection of host cells into which the vector has been introduced (see *e.g.*, U.S. Patents Nos. 4,399,216, 4,634,665 and 5,179,017, all by Axel *et al.*). For example, typically the selectable marker gene confers resistance to drugs, such as G418, puromycin, blasticidin, hygromycin or methotrexate, on a host cell into which the vector has been introduced. Suitable selectable marker genes include the dihydrofolate reductase (DHFR) gene (for use in DHFR⁻ host cells with methotrexate selection/amplification) and the neo gene (for G418 selection). For expression of the light and heavy chains, the expression vector(s) encoding the heavy and light chains is transfected into a host cell by standard techniques. The various forms of the term "transfection" are intended to encompass a wide variety of techniques commonly used for the introduction of exogenous DNA into a prokaryotic or eukaryotic host cell, *e.g.*, electroporation, lipofection, calcium-phosphate precipitation, DEAE-dextran transfection and the like.

[0067] It is possible to express the antibodies of the disclosure in either prokaryotic or eukaryotic host cells. In certain embodiments, expression of antibodies is performed in eukaryotic cells, *e.g.*, mammalian host cells, for optimal secretion of a properly folded and immunologically active antibody. Exemplary mammalian host cells for expressing the recombinant antibodies of the disclosure include Chinese Hamster Ovary (CHO cells) (including DHFR⁻ CHO cells, described in Urlaub and Chasin, 1980, *Proc. Natl. Acad. Sci. USA* 77:4216-4220, used with a DHFR selectable marker, *e.g.*, as described in Kaufman and Sharp, 1982, *Mol. Biol.* 159:601-621), NS0 myeloma cells, COS cells, 293 cells and SP2/0 cells. When recombinant expression vectors encoding antibody genes are introduced into mammalian host cells, the

antibodies are produced by culturing the host cells for a period of time sufficient to allow for expression of the antibody in the host cells or secretion of the antibody into the culture medium in which the host cells are grown. Antibodies can be recovered from the culture medium using standard protein purification methods. Host cells can also be used to produce portions of intact antibodies, such as Fab fragments or scFv molecules. It is understood that variations on the

5 above procedure are within the scope of the present disclosure. For example, it can be desirable to transfect a host cell with DNA encoding either the light chain or the heavy chain (but not both) of an anti-VEGF antibody of this disclosure.

[0068] Recombinant DNA technology can also be used to remove some or all of the DNA encoding either or both of the light and heavy chains that is not necessary for binding to VEGF. The molecules expressed from such truncated DNA molecules are also encompassed by the antibodies of the disclosure.

10 **[0069]** In addition, bifunctional antibodies can be produced in which one heavy and one light chain are an antibody of the disclosure and the other heavy and light chain are specific for an antigen other than VEGF by crosslinking an antibody of the disclosure to a second antibody by standard chemical crosslinking methods. Bifunctional antibodies can also be made by expressing a nucleic acid engineered to encode a bifunctional antibody.

15 **[0070]** In certain embodiments, dual specific antibodies, *i.e.*, antibodies that bind VEGF and an unrelated antigen using the same binding site, can be produced by mutating amino acid residues in the light chain and/or heavy chain CDRs. In various embodiments, dual specific antibodies that bind two antigens, such as HER2 and VEGF, can be produced by mutating amino acid residues in the periphery of the antigen binding site (Bostrom et al., 2009, Science 323:1610-1614). Dual functional antibodies can be made by expressing a nucleic acid engineered to encode a dual specific antibody.

20 **[0071]** For recombinant expression of an anti-VEGF antibody of the disclosure, the host cell can be co-transfected with two expression vectors of the disclosure, the first vector encoding a heavy chain derived polypeptide and the second vector encoding a light chain derived polypeptide. Typically, the two vectors each contain a separate selectable marker. Alternatively, a single vector can be used which encodes both heavy and light chain polypeptides.

25 **[0072]** Once a nucleic acid encoding one or more portions of an anti-VEGF antibody is generated, further alterations or mutations can be introduced into the coding sequence, for example to generate nucleic acids encoding antibodies with different CDR sequences, antibodies with reduced affinity to the Fc receptor, or antibodies of different subclasses.

30 **[0073]** The anti-VEGF antibodies of the disclosure can also be produced by chemical synthesis (*e.g.*, by the methods described in Solid Phase Peptide Synthesis, 2nd ed., 1984 The Pierce Chemical Co., Rockford, IL). Variant antibodies can also be generated using a cell-free platform (*see, e.g.*, Chu et al., 2001, Biochemia No. 2 (Roche Molecular Biologicals)).

35 **[0074]** Once an anti-VEGF antibody of the disclosure has been produced by recombinant expression, it can be purified by any method known in the art for purification of an immunoglobulin molecule, for example, by chromatography (*e.g.*, ion exchange, affinity, particularly by affinity for VEGF after Protein A or Protein G selection, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins. Further, the anti-VEGF antibodies of the present disclosure or fragments thereof can be fused to heterologous polypeptide sequences described herein or otherwise known in the art to facilitate purification.

40 **[0075]** Once isolated, an anti-VEGF antibody can, if desired, be further purified, *e.g.*, by high performance liquid chromatography (*See, e.g.*, Fisher, Laboratory Techniques In Biochemistry And Molecular Biology (Work and Burdon, eds., Elsevier, 1980), or by gel filtration chromatography on a Superdex™ 75 column (Pharmacia Biotech AB, Uppsala, Sweden).

6.3 BIOLOGICAL ACTIVITIES OF ANTI-VEGF ANTIBODIES

45 **[0076]** In certain embodiments, the anti-VEGF antibodies of the disclosure have certain biological activities, such as competing with bevacizumab or ranibizumab for binding to VEGF or neutralizing VEGF activity.

[0077] Accordingly, in certain embodiments, anti-VEGF antibodies of the disclosure compete with bevacizumab or ranibizumab for binding to VEGF. The ability to compete for binding to VEGF can be tested using a competition assay. In one example of a competition assay, VEGF is adhered onto a solid surface, *e.g.*, a microwell plate, by contacting the plate with a solution of VEGF (*e.g.*, at a concentration of 1 µg/mL in PBS over night at 4°C). The plate is washed (*e.g.*, 0.1% Tween 20 in PBS) and blocked (*e.g.*, in Superblock, Thermo Scientific, Rockford, IL). A mixture of sub-saturating amount of biotinylated bevacizumab (80 ng/mL) or an equivalent amount of biotinylated ranibizumab and unlabeled bevacizumab (or ranibizumab as the case may be) (the "reference" antibody) or competing anti-VEGF antibody (the "test" antibody) in serial dilution (*e.g.*, at a concentration of 2.8 µg/mL, 8.3 µg/mL, or 25 µg/mL) in ELISA buffer (*e.g.*, 1% BSA and 0.1% Tween 20 in PBS) is added to wells and plates are incubated for 1 hour with gentle shaking. 50
55 The plate is washed, 1 µg/mL HRP-conjugated Streptavidin diluted in ELISA buffer is added to each well and the plates incubated for 1 hour. Plates are washed and bound antibodies were detected by addition of substrate (*e.g.*, TMB, Biofx Laboratories Inc., Owings Mills, MD). The reaction is terminated by addition of stop buffer (*e.g.*, Bio FX Stop Reagents, Biofx Laboratories Inc., Owings Mills, MD) and the absorbance is measured at 650 nm using microplate reader (*e.g.*,

VERSAmx, Molecular Devices, Sunnyvale, CA). Variations on this competition assay can also be used to test competition between an anti-VEGF antibody of the disclosure and bevacizumab or ranibizumab. For example, in certain aspects, the anti-VEGF antibody is used as a reference antibody and bevacizumab or ranibizumab is used as a test antibody. Additionally, instead of soluble VEGF, membrane-bound VEGF expressed on the surfaces of cell (for example mammalian cells) in culture can be used. Other formats for competition assays are known in the art and can be employed.

[0078] In various embodiments, an anti-VEGF antibody of the disclosure reduces the binding of labeled bevacizumab or ranibizumab by at least 30%, by at least 40%, by at least 50%, by at least 60%, by at least 70%, by at least 80%, by at least 90%, by at least 95%, by at least 99% or by a percentage ranging between any of the foregoing values (e.g., an anti-VEGF antibody of the disclosure reduces the binding of labeled bevacizumab or ranibizumab by 50% to 70%) when the anti-VEGF antibody is used at a concentration of 0.08 $\mu\text{g/mL}$, 0.4 $\mu\text{g/mL}$, 2 $\mu\text{g/mL}$, 10 $\mu\text{g/mL}$, 50 $\mu\text{g/mL}$, 100 $\mu\text{g/mL}$ or at a concentration ranging between any of the foregoing values (e.g., at a concentration ranging from 2 $\mu\text{g/mL}$ to 10 $\mu\text{g/mL}$).

[0079] In other embodiments, bevacizumab or ranibizumab reduces the binding of a labeled anti-VEGF antibody of the disclosure by at least 40%, by at least 50%, by at least 60%, by at least 70%, by at least 80%, by at least 90%, or by a percentage ranging between any of the foregoing values (e.g., bevacizumab or ranibizumab reduces the binding of a labeled an anti-VEGF antibody of the disclosure by 50% to 70%) when bevacizumab or ranibizumab is used at a concentration of 0.4 $\mu\text{g/mL}$, 2 $\mu\text{g/mL}$, 10 $\mu\text{g/mL}$, 50 $\mu\text{g/mL}$, 250 $\mu\text{g/mL}$ or at a concentration ranging between any of the foregoing values (e.g., at a concentration ranging from 2 $\mu\text{g/mL}$ to 10 $\mu\text{g/mL}$).

[0080] In other aspects, an anti-VEGF antibody of the disclosure inhibits (or neutralizes) VEGF activity in a range of *in vitro* assays, such as cell proliferation or cell migration. For example, in one embodiment, the VEGF activity assayed is induction of endothelial cell ("EC") proliferation (see, e.g., protocol of Qin et al., 2006, J. Biol. Chem. 281:32550-32558). In another embodiment, the VEGF activity assayed is induction of EC migration (see, e.g., the *in vitro* scratch assay protocol described of Liang et al., 2007, Nat. Protoc. 2:329-333). In a specific embodiment, an anti-VEGF antibody is tested for the ability to reverse proliferation and cell migration stimulated by VEGF and delocalization of tight junction proteins induced by VEGF₁₆₅ in immortalized bovine retinal endothelial cells (Deissler et al., 2008, British Journal of Ophthalmology 92:839-843). In yet another embodiment, the neutralization of VEGF activity is assayed using a reporter assay (see, e.g., Yohno et al., 2003, Biological & Pharmaceutical Bulletin 26(4):417-20 and U.S. Patent No. 6,787,323).

[0081] Other formats for VEGF neutralization assays are known in the art and can be employed.

[0082] In various embodiments, an anti-VEGF antibody of the disclosure neutralizes VEGF by at least 30%, by at least 40%, by at least 50%, by at least 60%, by at least 70%, by at least 80%, by at least 90%, or by a percentage ranging between any of the foregoing values (e.g., an anti-VEGF antibody of the disclosure neutralizes VEGF activity by 50% to 70%) when the anti-VEGF antibody is used at a concentration of 2 ng/mL, 5 ng/mL, 10 ng/mL, 20 ng/mL, 0.1 $\mu\text{g/mL}$, 0.2 $\mu\text{g/mL}$, 1 $\mu\text{g/mL}$, 2 $\mu\text{g/mL}$, 5 $\mu\text{g/mL}$, 10 $\mu\text{g/mL}$, 20 $\mu\text{g/mL}$, or at a concentration ranging between any of the foregoing values (e.g., at a concentration ranging from 1 $\mu\text{g/mL}$ to 5 $\mu\text{g/mL}$).

[0083] In some embodiments, an anti-VEGF antibody of the disclosure is at least 0.7-fold as effective, 0.8-fold as effective, at least 0.9-fold as effective, at least 1-fold as effective, at least 1.1-fold as effective, at least 1.25-fold as effective, at least 1.5-fold as effective, at least 2-fold as effective, at least 5-fold as effective, at least 10-fold as effective, at least 20-fold as effective, at least 50-fold as effective, at least 100-fold as effective, at least 200-fold as effective, at least 500-fold as effective, at least 1000-fold as effective as bevacizumab or ranibizumab at neutralizing VEGF, or having an effectiveness at neutralizing VEGF relative to bevacizumab or ranibizumab ranging between any pair of the foregoing values (e.g., 0.9-fold to 5-fold as effective as bevacizumab or ranibizumab or 2-fold to 50-fold as effective as bevacizumab or ranibizumab in neutralizing VEGF).

6.4 KINETIC PROPERTIES OF ANTI-VEGF ANTIBODIES

[0084] In certain embodiments, the anti-VEGF antibodies of the disclosure have a high binding affinity for VEGF. In specific embodiments, the anti-VEGF antibodies of the present disclosure have specific association rate constants (k_{on} or k_{a} values), dissociation rate constants (k_{off} or k_{d} values), affinity constants (K_{A} values), dissociation constants (K_{D} values) and/or IC_{50} values. In various embodiments, binding constants for the interaction of the anti-VEGF antibodies with VEGF receptor can be determined using surface plasmon resonance, e.g., according to the method disclosed in Karlsson et al., 1991, J. Immunol. Methods 145:229-240. In certain aspects, such values are selected from the following embodiments.

[0085] In a specific embodiment, an anti-VEGF antibody of the disclosure binds to VEGF with a k_{on} of at least $10^4 \text{ M}^{-1}\text{s}^{-1}$, at least $5 \times 10^4 \text{ M}^{-1}\text{s}^{-1}$, at least $10^5 \text{ M}^{-1}\text{s}^{-1}$, at least $5 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$, at least $10^6 \text{ M}^{-1}\text{s}^{-1}$, at least $5 \times 10^6 \text{ M}^{-1}\text{s}^{-1}$, at least $10^7 \text{ M}^{-1}\text{s}^{-1}$, at least $5 \times 10^7 \text{ M}^{-1}\text{s}^{-1}$, at least $10^8 \text{ M}^{-1}\text{s}^{-1}$, at least $5 \times 10^8 \text{ M}^{-1}\text{s}^{-1}$, at least $10^9 \text{ M}^{-1}\text{s}^{-1}$ or with a k_{on} of any range between any pair of the foregoing values (e.g., 5×10^5 to $5 \times 10^6 \text{ M}^{-1}\text{s}^{-1}$ or 10^7 to $10^8 \text{ M}^{-1}\text{s}^{-1}$).

[0086] In another embodiment, an anti-VEGF antibody of the disclosure binds to VEGF with a k_{off} rate of 10^{-3} s^{-1} or less, $5 \times 10^{-4} \text{ s}^{-1}$ or less, 10^{-4} s^{-1} or less, $5 \times 10^{-5} \text{ s}^{-1}$ or less, 10^{-5} s^{-1} or less, $5 \times 10^{-6} \text{ s}^{-1}$ or less, 10^{-6} s^{-1} or less, $5 \times$

10^{-7} s^{-1} or less, 10^{-7} s^{-1} or less, $5 \times 10^{-8} \text{ s}^{-1}$ or less, 10^{-8} s^{-1} or less, or with a k_{off} rate of any range between any pair of the foregoing values (e.g., 5×10^{-4} to 10^{-6} s^{-1} , or 10^{-3} to $5 \times 10^{-5} \text{ s}^{-1}$).

[0087] In another embodiment, an anti-VEGF antibody of the disclosure binds to VEGF with a K_A ($k_{\text{on}}/k_{\text{off}}$) of at least at least 10^8 M^{-1} , at least $5 \times 10^9 \text{ M}^{-1}$, at least 10^{10} M^{-1} , at least $5 \times 10^{10} \text{ M}^{-1}$, 10^{11} M^{-1} , at least $5 \times 10^{11} \text{ M}^{-1}$, at least 10^{12} M^{-1} , at least $5 \times 10^{12} \text{ M}^{-1}$, at least 10^{13} M^{-1} , at least $5 \times 10^{13} \text{ M}^{-1}$, at least 10^{14} M^{-1} , at least $5 \times 10^{14} \text{ M}^{-1}$, at least 10^{15} M^{-1} or with a K_A of any range between any pair of the foregoing values (e.g., from $5 \times 10^9 \text{ M}^{-1}$ to 10^{11} M^{-1} , or from 10^{11} M^{-1} to $5 \times 10^{14} \text{ M}^{-1}$).

[0088] In other embodiments, an anti-VEGF antibody of the disclosure binds to VEGF with a K_D ($k_{\text{off}}/k_{\text{on}}$) of 10^{-8} M or less, $5 \times 10^{-9} \text{ M}$ or less, 10^{-9} M or less, $5 \times 10^{-10} \text{ M}$ or less, 10^{-10} M or less, $5 \times 10^{-11} \text{ M}$ or less, 10^{-11} M or less, $5 \times 10^{-12} \text{ M}$ or less, 10^{-12} M or less, $5 \times 10^{-13} \text{ M}$ or less, 10^{-13} M or less, $5 \times 10^{-14} \text{ M}$ or less, 10^{-14} M or less, $5 \times 10^{-15} \text{ M}$ or less, 10^{-15} M or less, or with a K_D of any range between any pair of the foregoing values (e.g., 5×10^{-9} to $5 \times 10^{-12} \text{ M}$, or from $5 \times 10^{-11} \text{ M}$ to $5 \times 10^{-13} \text{ M}$).

[0089] In specific embodiments, the K_D ($k_{\text{off}}/k_{\text{on}}$) value is determined by assays well known in the art or described herein, e.g., ELISA, isothermal titration calorimetry (ITC), fluorescent polarization assay or any other biosensors such as BIAcore.

[0090] In some embodiments, an anti-VEGF antibody of the disclosure binds to VEGF and inhibits the binding of VEGF to a VEGF receptor (Flt-1 or Flk-1) at an IC_{50} value of less than $5 \times 10^7 \text{ nM}$, less than 10^7 nM , less than $5 \times 10^6 \text{ nM}$, less than 10^6 nM , less than $5 \times 10^5 \text{ nM}$, less than 10^5 nM , less than $5 \times 10^4 \text{ nM}$, less than 10^4 nM , less than $5 \times 10^3 \text{ nM}$, less than 10^3 nM , less than $5 \times 10^2 \text{ nM}$, less than 100 nM , less than 90 nM , less than 80 nM , less than 70 nM , less than 65 nM , less than 60 nM , less than 50 nM , less than 40 nM , less than 30 nM , less than 25 nM , less than 20 nM , less than 15 nM , less than 12 nM , less than 10 nM , less than 5 nM , less than 1 nM , less than $5 \times 10^{-1} \text{ nM}$, less than 10^{-1} nM , less than $5 \times 10^{-2} \text{ nM}$, less than 10^{-2} nM , less than $5 \times 10^{-3} \text{ nM}$, less than 10^{-3} nM , less than $5 \times 10^{-4} \text{ nM}$, or less than 10^{-4} nM , or with an IC_{50} of any range between any pair of the foregoing values (e.g., 5×10^7 to 50 nM , or 15 nM to $5 \times 10^{-3} \text{ nM}$). IC_{50} can be measured according to methods well known in the art or described herein, e.g., ELISA.

[0091] In other embodiments, an anti-VEGF antibody of the disclosure binds to VEGF and neutralizes the activity of VEGF in a bioassay (e.g., EC proliferation or migration) at an IC_{50} value of less than $5 \times 10^7 \text{ nM}$, less than 10^7 nM , less than $5 \times 10^6 \text{ nM}$, less than 10^6 nM , less than $5 \times 10^5 \text{ nM}$, less than 10^5 nM , less than $5 \times 10^4 \text{ nM}$, less than 10^4 nM , less than $5 \times 10^3 \text{ nM}$, less than 10^3 nM , less than $5 \times 10^2 \text{ nM}$, less than 100 nM , less than 90 nM , less than 80 nM , less than 70 nM , less than 65 nM , less than 60 nM , less than 50 nM , less than 40 nM , less than 30 nM , less than 25 nM , less than 20 nM , less than 15 nM , less than 12 nM , less than 10 nM , less than 5 nM , less than 1 nM , less than $5 \times 10^{-1} \text{ nM}$, less than 10^{-1} nM , less than $5 \times 10^{-2} \text{ nM}$, less than 10^{-2} nM , less than $5 \times 10^{-3} \text{ nM}$, less than 10^{-3} nM , less than $5 \times 10^{-4} \text{ nM}$, or less than 10^{-4} nM , or with an IC_{50} of any range between any pair of the foregoing values (e.g., 5×10^7 to 50 nM , or 15 nM to $5 \times 10^{-3} \text{ nM}$). An exemplary neutralization assay that can be used to measure the IC_{50} of an anti-VEGF antibody is described in Section 6.3 below.

[0092] In certain embodiments, an anti-VEGF antibody binds to VEGF and inhibits the binding of VEGF to Flt-1, Flk-1 or both, or inhibits VEGF activity in a VEGF neutralization assay, at an IC_{50} value of between approximately 1 pM and approximately $1 \text{ }\mu\text{M}$. In specific embodiments, an anti-VEGF antibody binds to VEGF and inhibits the binding of VEGF to Flt-1, Flk-1 or both, or inhibits VEGF activity in a VEGF neutralization assay, at an IC_{50} value of between 10 pM and 100 nM , between 100 pM and 10 nM , between 200 pM and 5 nM , between 300 pM and 4 nM , between 500 pM and 3 nM , between 750 pM and 2 nM , between 1 nM and 20 nM , between 500 pM and 40 nM , between 50 pM and 50 nM , between 250 pM and 100 nM , and between 100 nM and $1 \text{ }\mu\text{M}$, or with an IC_{50} of any range between any pair of the foregoing values (e.g., 10 pM to 50 nM , or 750 pM to 2 nM).

[0093] In certain aspects of the foregoing embodiments, the IC_{50} is measured in the presence of VEGF at a concentration of $0.001 \text{ }\mu\text{M}$, $0.005 \text{ }\mu\text{M}$, $0.01 \text{ }\mu\text{M}$, $0.05 \text{ }\mu\text{M}$, $0.1 \text{ }\mu\text{M}$, $0.5 \text{ }\mu\text{M}$, $1 \text{ }\mu\text{M}$, $10 \text{ }\mu\text{M}$, $20 \text{ }\mu\text{M}$, $30 \text{ }\mu\text{M}$, $40 \text{ }\mu\text{M}$, $50 \text{ }\mu\text{M}$, $60 \text{ }\mu\text{M}$, $70 \text{ }\mu\text{M}$, $80 \text{ }\mu\text{M}$, $90 \text{ }\mu\text{M}$, $100 \text{ }\mu\text{M}$, $200 \text{ }\mu\text{M}$, $300 \text{ }\mu\text{M}$, $400 \text{ }\mu\text{M}$, $500 \text{ }\mu\text{M}$, $600 \text{ }\mu\text{M}$, $700 \text{ }\mu\text{M}$, $800 \text{ }\mu\text{M}$, $900 \text{ }\mu\text{M}$, $1000 \text{ }\mu\text{M}$ or at a concentration of any range between any pair of the foregoing values (e.g., 0.01 to $50 \text{ }\mu\text{M}$, or $10 \text{ }\mu\text{M}$ to $100 \text{ }\mu\text{M}$).

[0094] In certain embodiments, the kinetic properties of an antibody of the disclosure are comparable to, or improved relative to, bevacizumab or of ranibizumab in a comparable assay. For example, in certain embodiments, an anti-VEGF antibody of the disclosure binds to VEGF with a k_{on} rate ranging from approximately $0.5x$ to $1000x$ of the k_{on} of bevacizumab or of ranibizumab, for example a k_{on} of $0.5x$ of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of $0.75x$ of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of $0.9x$ of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of $1x$ of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of $1.1x$ of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of $1.2x$ of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of $1.3x$ of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of $1.4x$ of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of $1.5x$ of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of $1.75x$ of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of $2x$ of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of $2.25x$ of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of $2.5x$ of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of $3x$ of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of $4x$ of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of $5x$ of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of $7.5x$ of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of $10x$ of the k_{on} of

bevacizumab or of ranibizumab, a k_{on} of 15x of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of 20x of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of 50x of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of 75x of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of 100x of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of 150x of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of 200x of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of 200x of the k_{on} of bevacizumab or of ranibizumab or a k_{on} ranging between any pair of the foregoing values, e.g., a k_{on} of 2x-75x of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of 5x-100x of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of 0.5x-1000x of the k_{on} of bevacizumab or of ranibizumab, a k_{on} of 0.75x-200x of the k_{on} of bevacizumab or of ranibizumab, etc.

[0095] In certain embodiments, an anti-VEGF antibody of the disclosure binds to VEGF with a k_{off} rate ranging from 0.001x to 3x of the k_{off} of bevacizumab or of ranibizumab, for example a k_{off} of 0.002x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 0.005x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 0.0075x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 0.01x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 0.025x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 0.05x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 0.075x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 0.1x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 0.25x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 0.5x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 0.75x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 0.9x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 1x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 1.1x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 1.25x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 1.5x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 1.75x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 4x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 3x of the k_{off} of bevacizumab or of ranibizumab, or a k_{off} ranging between any pair of the foregoing values, e.g., a k_{off} of 0.01x to 1.1x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 0.05x-1.25x of the k_{off} of bevacizumab or of ranibizumab, a k_{off} of 0.1x-0.9x of the k_{off} of bevacizumab or of ranibizumab, etc.

[0096] In other embodiments, an anti-VEGF antibody of the disclosure binds to VEGF with a K_A (k_{on}/k_{off}) ranging from 0.25x to 1000x of the K_A of bevacizumab or of ranibizumab, for example a K_A of 0.5x of the K_A of bevacizumab or of ranibizumab, a K_A of 0.75x of the K_A of bevacizumab or of ranibizumab, a K_A of 1x of the K_A of bevacizumab or of ranibizumab, a K_A of 2x of the K_A of bevacizumab or of ranibizumab, a K_A of 4x of the K_A of bevacizumab or of ranibizumab, a K_A of 10x of the K_A of bevacizumab or of ranibizumab, a K_A of 15x of the K_A of bevacizumab or of ranibizumab, a K_A of 20x of the K_A of bevacizumab or of ranibizumab, a K_A of 30x of the K_A of bevacizumab or of ranibizumab, a K_A of 40x of the K_A of bevacizumab or of ranibizumab, a K_A of 50x of the K_A of bevacizumab or of ranibizumab, a K_A of 100x of the K_A of bevacizumab or of ranibizumab, a K_A of 250x of the K_A of bevacizumab or of ranibizumab, a K_A of 500x of the K_A of bevacizumab or of ranibizumab, a K_A of 750x of the K_A of bevacizumab or of ranibizumab, a K_A of 1000x of the K_A of bevacizumab or of ranibizumab or a K_A ranging between any pair of the foregoing values, e.g., a K_A of 0.75x-10 5x of the K_A of bevacizumab or of ranibizumab, a K_A of 1x-100x of the K_A of bevacizumab or of ranibizumab, a K_A of 10x-20x of the K_A of bevacizumab or of ranibizumab, a K_A of 4x-50x of the K_A of bevacizumab or of ranibizumab, a K_A of 2x-20x of the K_A of bevacizumab or of ranibizumab, or any value or range that can be calculated from the k_{on} and k_{off} rates disclosed herein.

[0097] In other embodiments, an anti-VEGF antibody of the disclosure binds to VEGF a K_D (k_{off}/k_{on}) ranging from ranging from 0.001 x to 10x of the K_D of bevacizumab or of ranibizumab, for example a K_D of 0.001x of the K_D of bevacizumab or of ranibizumab, a K_D of 0.005x of the K_D of bevacizumab or of ranibizumab, a K_D of 0.01x of the K_D of bevacizumab or of ranibizumab, a K_D of 0.05x of the K_D of bevacizumab or of ranibizumab, a K_D of 0.075x of the K_D of bevacizumab or of ranibizumab, a K_D of 0.1x of the K_D of bevacizumab or of ranibizumab, a K_D of 0.2x of the K_D of bevacizumab or of ranibizumab, a K_D of 0.3x of the K_D of bevacizumab or of ranibizumab, a K_D of 0.4x of the K_D of bevacizumab or of ranibizumab, a K_D of 0.5x of the K_D of bevacizumab or of ranibizumab, a K_D of 0.6x of the K_D of bevacizumab or of ranibizumab, a K_D of 0.7x of the K_D of bevacizumab or of ranibizumab, a K_D of 0.8x of the K_D of bevacizumab or of ranibizumab, a K_D of 0.9x of the K_D of bevacizumab or of ranibizumab, a K_D of 1x of the K_D of bevacizumab or of ranibizumab, a K_D of 1.5x of the K_D of bevacizumab or of ranibizumab, a K_D of 2x of the K_D of bevacizumab or of ranibizumab, a K_D of 4x of the K_D of bevacizumab or of ranibizumab, a K_D of 7.5x of the K_D of bevacizumab or of ranibizumab, a K_D of 10x of the K_D of bevacizumab or of ranibizumab or a K_D ranging between any pair of the foregoing values, e.g., a K_D of 0.01x-2x of the K_D of bevacizumab or of ranibizumab, a KD of 0.1x-1.5x of the K_D of bevacizumab or of ranibizumab, a K_D of 0.7x-4x of the K_D of bevacizumab or of ranibizumab, a K_D of 0.2x-2x of the K_D of bevacizumab or of ranibizumab or any value or range that can be calculated from the k_{on} and k_{off} rates disclosed herein.

[0098] In some embodiments, an anti-VEGF antibody of the disclosure binds to VEGF and inhibits the binding of VEGF to Flt-1, Flk-1 or both, or neutralize the activity of VEGF at an IC_{50} value ranging from 0.001x to 10x of the IC_{50} of bevacizumab or of ranibizumab, for example an IC_{50} of 0.001x of the IC_{50} of bevacizumab or of ranibizumab, an IC_{50} of 0.005x of the IC_{50} of bevacizumab or of ranibizumab, an IC_{50} of 0.01x of the IC_{50} of bevacizumab or of ranibizumab, an IC_{50} of 0.05x of the IC_{50} of bevacizumab or of ranibizumab, an IC_{50} of 0.075x of the IC_{50} of bevacizumab or of ranibizumab, an IC_{50} of 0.1x of the IC_{50} of bevacizumab or of ranibizumab, an IC_{50} of 0.2x of the IC_{50} of bevacizumab

or of ranibizumab, an IC₅₀ of 0.3x of the IC₅₀ of bevacizumab or of ranibizumab, an IC₅₀ of 0.4x of the IC₅₀ of bevacizumab or of ranibizumab, an IC₅₀ of 0.5x of the IC₅₀ of bevacizumab or of ranibizumab, an IC₅₀ of 0.6x of the IC₅₀ of bevacizumab or of ranibizumab, an IC₅₀ of 0.7x of the IC₅₀ of bevacizumab or of ranibizumab, an IC₅₀ of 0.8x of the IC₅₀ of bevacizumab or of ranibizumab, an IC₅₀ of 0.9x of the IC₅₀ of bevacizumab or of ranibizumab, an IC₅₀ of 1x of the IC₅₀ of bevacizumab or of ranibizumab, an IC₅₀ of 1.5x of the IC₅₀ of bevacizumab or of ranibizumab, an IC₅₀ of 2x of the IC₅₀ of bevacizumab or of ranibizumab, an IC₅₀ of 4x of the IC₅₀ of bevacizumab or of ranibizumab, an IC₅₀ of 7.5x of the IC₅₀ of bevacizumab or of ranibizumab, an IC₅₀ of 10x of the IC₅₀ of bevacizumab or of ranibizumab or an IC₅₀ ranging between any pair of the foregoing values, e.g., an IC₅₀ of 0.01x-2x of the IC₅₀ of bevacizumab or of ranibizumab, an IC₅₀ of 0.1x-1.5x of the IC₅₀ of bevacizumab or of ranibizumab, an IC₅₀ of 0.7x-4x of the IC₅₀ of bevacizumab or of ranibizumab, an IC₅₀ of 0.2x-2x of the IC₅₀ of bevacizumab or of ranibizumab. In certain embodiments, a single CDR substitution can result in the foregoing differences in IC₅₀ as compared to bevacizumab or ranibizumab, whereas an anti-VEGF antibody of the disclosure can comprise such substitution and up to 16 additional CDR substitutions as compared to bevacizumab or ranibizumab.

6.5 REDUCED IMMUNOGENICITY OF ANTI-VEGF ANTIBODIES

[0099] In certain aspects, the present disclosure provides anti-VEGF antibodies having reduced immunogenicity as compared to bevacizumab or ranibizumab. The present disclosure provides anti-VEGF antibodies having single or multiple amino acid substitutions in their CDRs and/or framework regions as compared to the CDRs of bevacizumab, wherein at least one substitution reduces the immunogenicity of the antibody as compared to bevacizumab or ranibizumab. In certain embodiments, the reduced immunogenicity results from one or more amino acid substitutions that result in eliminating or mitigating one or more T cell epitopes.

[0100] In certain aspects, the anti-VEGF antibodies of the disclosure having reduced immunogenicity have comparable or improved biological activity as compared to bevacizumab or ranibizumab, e.g., affinity towards VEGF or neutralization of VEGF activity. Such properties can be tested, for example, by the methods described in Section 6.3 above.

[0101] In certain embodiments, the immunogenicity of an anti-VEGF antibody of the disclosure is reduced relative to bevacizumab or ranibizumab antibody. Such antibodies generally have variant sequences relative to the heavy and/or light chain variable region in regions corresponding to SEQ ID NO:25, SEQ ID NO:62 and/or SEQ ID NO:74. The antibodies will generally have one, two or three amino acid substitutions in one, two or all three sequences corresponding to SEQ ID NO:25, SEQ ID NO:62, and SEQ ID NO:74, although up to four or five substitutions in one, two or all three regions are contemplated herein.

[0102] As used in the present disclosure, a variant with "reduced immunogenicity" refers to an anti-VEGF antibody with a variant sequence in a region corresponding to SEQ ID NO:25, SEQ ID NO:62, and/or SEQ ID NO:74 that elicits a reduced proliferative response in peripheral blood mononuclear cells as compared to a peptide of SEQ ID NO:25, SEQ ID NO:62, or SEQ ID NO:74, respectively. An exemplary proliferation assay that can be used to evaluate the proliferative response is set forth in Section 7 below. The reduced proliferative response can be reflected in terms of the percentage of responders, the stimulation index, or both.

[0103] In other embodiments, as compared to a peptide having the sequence of SEQ ID NO:25, SEQ ID NO:62, or SEQ ID NO:74, the variant sequence results in at least 25% fewer responders, in at least 30% fewer responders, in at least 35% fewer responders, in at least 40% fewer responders, in at least 45% fewer responders, in at least 50% fewer responders, in at least 60% fewer responders, in at least 65% fewer responders, in at least 70% fewer responders, in at least 75% fewer responders, in at least 80% fewer responders, in at least 85% fewer responders, in at least 90% fewer responders, in at least 95% fewer responders, 100% fewer responders, or a reduction in responders in a range between any of the foregoing values, e.g., 25%-75% fewer responders, 50%-90% fewer responders, 60%-100% fewer responders, 70%-90% fewer responders, or the like.

[0104] In other embodiments, the variant sequence results in a stimulation index that is at least 5% less, at least 10% less, at least 15% less, at least 20% less, at least 25% less, at least 30% less, at least 35% less, or at least 40% less than the stimulation index elicited by a peptide of SEQ ID NO:25, SEQ ID NO:62, or SEQ ID NO:74, respectively, or results in a stimulation index reduced by a range between any of the foregoing values as compared to a peptide of SEQ ID NO:25, SEQ ID NO:62, or SEQ ID NO:74, e.g., 5%-20% less, 10%-30% less, 25%-35% less, 30%-40% less, or the like.

[0105] Exemplary embodiments of candidate anti-VEGF antibodies with reduced immunogenicity as compared to bevacizumab or ranibizumab comprise one or more of the CDR substitutions or combinations of substitutions set forth in Table 6. Optionally, anti-VEGF antibodies with reduced immunogenicity as compared to bevacizumab or ranibizumab comprise one or more additional substitutions, such as the CDR mutations in any of Tables 7-13, singly or in combination.

[0106] Yet further exemplary embodiments of candidate anti-VEGF antibodies with reduced immunogenicity as compared to bevacizumab or ranibizumab comprise one or more of the CDR substitutions or combinations of substitutions set forth in Tables 14-16. Some preferred embodiments of anti-VEGF antibodies with reduced immunogenicity as compared to bevacizumab or ranibizumab are provided in Table 19.

6.6 ANTIBODY CONJUGATES

[0107] The anti-VEGF antibodies of the disclosure include antibody conjugates that are modified, *e.g.*, by the covalent attachment of any type of molecule to the antibody, such that covalent attachment does not interfere with binding to VEGF.

5 **[0108]** In certain aspects, an anti-VEGF antibody of the disclosure can be conjugated to an effector moiety or a label. The term "effector moiety" as used herein includes, for example, antineoplastic agents, drugs, toxins, biologically active proteins, for example enzymes, other antibody or antibody fragments, synthetic or naturally occurring polymers, nucleic acids (*e.g.*, DNA and RNA), radionuclides, particularly radioiodide, radioisotopes, chelated metals, nanoparticles and reporter groups such as fluorescent compounds or compounds which can be detected by NMR or ESR spectroscopy.

10 **[0109]** In one example, anti-VEGF antibodies can be conjugated to an effector moiety, such as a cytotoxic agent, a radionuclide or drug moiety to modify a given biological response. The effector moiety can be a protein or polypeptide, such as, for example and without limitation, a toxin (such as abrin, ricin A, *Pseudomonas* exotoxin, or *Diphtheria* toxin), a signaling molecule (such as α -interferon, β -interferon, nerve growth factor, platelet derived growth factor or tissue plasminogen activator), a thrombotic agent or an anti-angiogenic agent (*e.g.*, angiostatin or endostatin) or a biological response modifier such as a cytokine or growth factor (*e.g.*, interleukin-1 (IL-1), interleukin-2 (IL-2), interleukin-6 (IL-6), granulocyte macrophage colony stimulating factor (GM-CSF), granulocyte colony stimulating factor (G-CSF), or nerve growth factor (NGF)).

15 **[0110]** In another example the effector moieties can be cytotoxins or cytotoxic agents. Examples of cytotoxins and cytotoxic agents include taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof.

20 **[0111]** Effector moieties also include, but are not limited to, antimetabolites (*e.g.* methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (*e.g.*, mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C5 and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (*e.g.*, daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (*e.g.*, dactinomycin (formerly actinomycin), bleomycin, mithramycin, anthramycin (AMC), calicheamicins or duocarmycins), and anti-mitotic agents (*e.g.*, vincristine and vinblastine).

25 **[0112]** Other effector moieties can include radionuclides such as, but not limited to, ^{111}In and ^{90}Y , Lu^{177} , Bi^{213} , Ca^{252} , Ir^{192} and Tungsten^{188} / Rhenium^{188} and drugs such as, but not limited to, alkylphosphocholines, topoisomerase I inhibitors, taxoids and suramin.

30 **[0113]** Techniques for conjugating such effector moieties to antibodies are well known in the art (see, *e.g.*, Hellstrom et al., *Controlled Drug Delivery*, 2nd Ed., at pp. 623-53 (Robinson et al., eds., 1987)); Thorpe et al., 1982, *Immunol. Rev.* 62:119-58 and Dubowchik et al., 1999, *Pharmacology and Therapeutics* 83:67-123).

35 **[0114]** In one example, the anti-VEGF antibody or fragment thereof is fused via a covalent bond (*e.g.*, a peptide bond), through the antibody's N-terminus or C-terminus or internally, to an amino acid sequence of another protein (or portion thereof; for example at least a 10, 20 or 50 amino acid portion of the protein). The antibody, or fragment thereof, can be linked to the other protein at the N-terminus of the constant domain of the antibody. Recombinant DNA procedures can be used to create such fusions, for example as described in WO 86/01533 and EP0392745. In another example the effector molecule can increase half-life *in vivo*, and/or enhance the delivery of an antibody across an epithelial barrier to the immune system. Examples of suitable effector molecules of this type include polymers, albumin, albumin binding proteins or albumin binding compounds such as those described in WO 2005/117984.

40 **[0115]** In certain aspects, an anti-VEGF antibody is conjugated to a small molecule toxin. In certain exemplary embodiments, an anti-VEGF antibody of the disclosure is conjugated to a dolastatin or a dolostatin peptidic analogs or derivatives, *e.g.*, an auristatin (U.S. Pat. Nos. 5,635,483 and 5,780,588). The dolastatin or auristatin drug moiety may be attached to the antibody through its N (amino) terminus, C (carboxyl) terminus or internally (WO 02/088172). Exemplary auristatin embodiments include the N-terminus linked monomethylauristatin drug moieties DE and DF, as disclosed in U.S. Patent No. 7,498,298, which is hereby incorporated by reference in its entirety (disclosing, *e.g.*, linkers and methods of preparing monomethylvaline compounds such as MMAE and MMAF conjugated to linkers).

45 **[0116]** In other exemplary embodiments, small molecule toxins include but are not limited to calicheamicin, maytansine (U.S. Pat. No. 5,208,020), trichothene, and CC1065. In one embodiment of the disclosure, the antibody is conjugated to one or more maytansine molecules (*e.g.*, about 1 to about 10 maytansine molecules per antibody molecule). Maytansine may, for example, be converted to May-SS-Me which may be reduced to May-SH3 and reacted with an antibody (Chari et al., 1992, *Cancer Research* 52: 127-131) to generate a maytansinoid-antibody or maytansinoid-Fc fusion conjugate. Structural analogues of calicheamicin that can also be used include but are not limited to γ_1^1 , γ_3^1 , N-acetyl- γ_1^1 , PSAG, and θ_1^1 , (Hinman et al., 1993, *Cancer Research* 53:3336-3342; Lode et al., 1998, *Cancer Research* 58:2925-2928; U.S. Patent No. 5,714,586; U.S. Patent No. 5,712,374; U.S. Patent No. 5,264,586; U.S. Patent No. 5,773,001).

50 **[0117]** Antibodies of the disclosure can also be conjugated to liposomes for targeted delivery (See, *e.g.*, Park et al.,

1997, Adv. Pharmacol. 40:399-435; Marty & Schwendener, 2004, Methods in Molecular Medicine 109:389-401).

[0118] In one example antibodies of the present disclosure can be attached to poly(ethyleneglycol) (PEG) moieties. In one particular example the antibody is an antibody fragment and the PEG moieties can be attached through any available amino acid side-chain or terminal amino acid functional group located in the antibody fragment, for example any free amino, imino, thiol, hydroxyl or carboxyl group. Such amino acids can occur naturally in the antibody fragment or can be engineered into the fragment using recombinant DNA methods. See for example U.S. Patent No. 5,219,996. Multiple sites can be used to attach two or more PEG molecules. PEG moieties can be covalently linked through a thiol group of at least one cysteine residue located in the antibody fragment. Where a thiol group is used as the point of attachment, appropriately activated effector moieties, for example thiol selective derivatives such as maleimides and cysteine derivatives, can be used.

[0119] In a specific example, an anti-VEGF antibody conjugate is a modified Fab' fragment which is PEGylated, *i.e.*, has PEG (poly(ethyleneglycol)) covalently attached thereto, *e.g.*, according to the method disclosed in EP0948544. See also Poly(ethyleneglycol) Chemistry, Biotechnical and Biomedical Applications, (J. Milton Harris (ed.), Plenum Press, New York, 1992); Poly(ethyleneglycol) Chemistry and Biological Applications, (J. Milton Harris and S. Zalipsky, eds., American Chemical Society, Washington D.C., 1997); and Bioconjugation Protein Coupling Techniques for the Biomedical Sciences, (M. Aslam and A. Dent, eds., Grove Publishers, New York, 1998); and Chapman, 2002, Advanced Drug Delivery Reviews 54:531-545. PEG can be attached to a cysteine in the hinge region. In one example, a PEG-modified Fab' fragment has a maleimide group covalently linked to a single thiol group in a modified hinge region. A lysine residue can be covalently linked to the maleimide group and to each of the amine groups on the lysine residue can be attached a methoxypoly(ethyleneglycol) polymer having a molecular weight of approximately 20,000 Da. The total molecular weight of the PEG attached to the Fab' fragment can therefore be approximately 40,000 Da.

[0120] The word "label" when used herein refers to a detectable compound or composition which can be conjugated directly or indirectly to an anti-VEGF antibody of the disclosure. The label can itself be detectable (*e.g.*, radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, can catalyze chemical alteration of a substrate compound or composition which is detectable. Useful fluorescent moieties include, but are not limited to, fluorescein, fluorescein isothiocyanate, rhodamine, 5-dimethylamine-1-naphthalenesulfonyl chloride, phycoerythrin and the like. Useful enzymatic labels include, but are not limited to, alkaline phosphatase, horseradish peroxidase, glucose oxidase and the like.

[0121] Additional anti-VEGF antibody conjugates that are useful for, *inter alia*, diagnostic purposes, are described in Section 6.7 below.

6.7 DIAGNOSTIC USES OF ANTI-VEGF ANTIBODIES

[0122] The anti-VEGF antibodies of the disclosure, including those antibodies that have been modified, *e.g.*, by biotinylation, horseradish peroxidase, or any other detectable moiety (including those described in Section 6.6), can be advantageously used for diagnostic purposes.

[0123] In particular, the anti-VEGF antibodies can be used, for example, but not limited to, to purify or detect VEGF, including both *in vitro* and *in vivo* diagnostic methods. For example, the antibodies have use in immunoassays for qualitatively and quantitatively measuring levels of VEGF in biological samples. See, *e.g.*, Harlow et al., Antibodies: A Laboratory Manual, Second Edition (Cold Spring Harbor Laboratory Press, 1988), which is incorporated by reference herein in its entirety.

[0124] The present disclosure further encompasses antibodies or fragments thereof conjugated to a diagnostic agent. The antibodies can be used diagnostically, for example, to detect expression of a target of interest in specific cells, tissues, or serum; or to monitor the development or progression of an immunologic response as part of a clinical testing procedure to, *e.g.*, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, radioactive materials, positron emitting metals using various positron emission tomographies, and nonradioactive paramagnetic metal ions. The detectable substance can be coupled or conjugated either directly to the antibody (or fragment thereof) or indirectly, through an intermediate (such as, for example, a linker known in the art) using techniques known in the art. Examples of enzymatic labels include luciferases (*e.g.*, firefly luciferase and bacterial luciferase; U.S. Patent No. 4,737,456), luciferin, 2,3-dihydrophthalazin-ediones, malate dehydrogenase, urease, peroxidase such as horseradish peroxidase (HRPO), alkaline phosphatase, β -galactosidase, acetylcholinesterase, glucoamylase, lysozyme, saccharide oxidases (*e.g.*, glucose oxidase, galactose oxidase, and glucose-6-phosphate dehydrogenase), heterocyclic oxidases (such as uricase and xanthine oxidase), lactoperoxidase, microperoxidase, and the like. Examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin; and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{111}In or ^{99}Tc .

[0125] The disclosure provides for the detection of expression of VEGF, comprising contacting a biological sample (cells, tissue, or body fluid of an individual) using one or more anti-VEGF antibodies of the disclosure (optionally conjugated to detectable moiety), and detecting whether or not the sample is positive for VEGF expression, or whether the sample has altered (e.g., reduced or increased) expression as compared to a control sample.

[0126] Diseases that can be diagnosed using the present methods include, but are not limited to, the diseases described herein. In certain embodiments, the tissue or body fluid is peripheral blood, peripheral blood leukocytes, biopsy tissues such as lung or skin biopsies, and tissue.

6.8 THERAPEUTIC METHODS USING ANTI-VEGF ANTIBODIES

6.8.1 Clinical Benefits

[0127] The anti-VEGF antibodies of the disclosure can be used to treat various neoplasms or non-neoplastic conditions characterized by pathological angiogenesis.

[0128] The antibodies of the disclosure are useful in the treatment of tumors in which angiogenesis plays an important role in tumor growth, including cancers and benign tumors. Examples of cancer to be treated herein include, but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia. More particular examples of such cancers include squamous cell cancer, lung cancer (including small-cell lung cancer, non-small cell lung cancer, adenocarcinoma of the lung, and squamous carcinoma of the lung), cancer of the peritoneum, hepatocellular cancer, gastric or stomach cancer (including gastrointestinal cancer), pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer, colon cancer, colorectal cancer, endometrial or uterine carcinoma, salivary gland carcinoma, kidney or renal cancer, liver cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma and various types of head and neck cancer, as well as B-cell lymphoma (including low grade/follicular non-Hodgkin's lymphoma (NHL); small lymphocytic (SL) NHL; intermediate grade/follicular NHL; intermediate grade diffuse NHL; high grade immunoblastic NHL; high grade lymphoblastic NHL; high grade small non-cleaved cell NHL; bulky disease NHL; mantle cell lymphoma; AIDS-related lymphoma; and Waldenstrom's Macroglobulinemia); chronic lymphocytic leukemia (CLL); acute lymphoblastic leukemia (ALL); Hairy cell leukemia; chronic myeloblastic leukemia; and post-transplant lymphoproliferative disorder (PTLD), as well as abnormal vascular proliferation associated with phakomatoses, edema (such as that associated with brain tumors), and Meigs' syndrome. More particularly, cancers that are amenable to treatment by the antibodies of the disclosure include breast cancer, colorectal cancer, rectal cancer, non-small cell lung cancer, non-Hodgkins lymphoma (NHL), renal cell cancer, prostate cancer, liver cancer, pancreatic cancer, soft-tissue sarcoma, kaposi's sarcoma, carcinoid carcinoma, head and neck cancer, melanoma, ovarian cancer, mesothelioma, and multiple myeloma. In some embodiments, the anti-VEGF antibodies of the disclosure are used to treat colorectal cancer in a human patient.

[0129] The present disclosure encompasses anti-angiogenic therapy, a cancer treatment strategy aimed at inhibiting the development of tumor blood vessels required for providing nutrients to support tumor growth. Because angiogenesis is involved in both primary tumor growth and metastasis, the antiangiogenic treatment provided by the disclosure is capable of inhibiting the neoplastic growth of tumor at the primary site as well as preventing metastasis of tumors at the secondary sites.

[0130] Non-neoplastic conditions that are amenable to treatment with the antibodies of the disclosure include retinal diseases (e.g., age-related macular degeneration) and immune and inflammatory diseases (e.g., rheumatoid arthritis, psoriasis, atherosclerosis, diabetic and other proliferative retinopathies including retinopathy of prematurity, retrolental fibroplasia, neovascular glaucoma, age-related macular degeneration, thyroid hyperplasias (including Grave's disease), corneal and other tissue transplantation, chronic inflammation, lung inflammation, nephrotic syndrome, preeclampsia, ascites, pericardial effusion (such as that associated with pericarditis), and pleural effusion).

[0131] Accordingly, the present disclosure provides methods of treating any of the foregoing diseases in a patient in need thereof, comprising: administering to the patient an anti-VEGF antibody of the disclosure. Optionally, said administration is repeated, e.g., after one day, two days, three days, five days, one week, two weeks, three weeks, one month, five weeks, six weeks, seven weeks, eight weeks, two months or three months. The repeated administration can be at the same dose or at a different dose. The administration can be repeated once, twice, three times, four times, five times, six times, seven times, eight times, nine times, ten times, or more. For example, according to certain dosage regimens a patient receives anti-VEGF therapy for a prolonged period of time, e.g., 6 months, 1 year or more. The amount of anti-VEGF antibody administered to the patient is in certain embodiments a therapeutically effective amount. As used herein, a "therapeutically effective" amount of VEGF antibody can be administered as a single dose or over the course of a therapeutic regimen, e.g., over the course of a week, two weeks, three weeks, one month, three months, six months, one year, or longer. Exemplary therapeutic regimens are described in Section 6.11 below.

[0132] According to the present disclosure, treatment of a disease encompasses the treatment of patients already diagnosed as having any form of the disease at any clinical stage or manifestation; the delay of the onset or evolution

or aggravation or deterioration of the symptoms or signs of the disease; and/or preventing and/or reducing the severity of the disease.

[0133] A "subject" or "patient" to whom the anti-VEGF antibody of the disclosure is administered is preferably a mammal such as a non-primate (e.g., cow, pig, horse, cat, dog, rat, *etc.*) or a primate (e.g., monkey or human). In certain embodiments, the subject or patient is a human. In certain aspects, the human is a pediatric patient. In other aspects, the human is an adult patient.

6.9 PHARMACEUTICAL COMPOSITIONS AND ROUTES OF ADMINISTRATION

[0134] Compositions comprising an anti-VEGF antibody of the disclosure and, optionally one or more additional therapeutic agents, such as the combination therapeutic agents described in Section 6.10 below, are provided herein. The compositions will usually be supplied as part of a sterile, pharmaceutical composition that will normally include a pharmaceutically acceptable carrier. This composition can be in any suitable form (depending upon the desired method of administering it to a patient).

[0135] The anti-VEGF antibodies of the disclosure can be administered to a patient by a variety of routes such as orally, transdermally, subcutaneously, intranasally, intravenously, intramuscularly, intraocularly, topically, intrathecally and intracerebroventricularly. The most suitable route for administration in any given case will depend on the particular antibody, the subject, and the nature and severity of the disease and the physical condition of the subject.

[0136] For treatment of indications other than retinal diseases, the effective dose of an anti-VEGF antibody of the disclosure can range from about 0.1 to about 75 mg/kg per single (e.g., bolus) administration, multiple administrations or continuous administration, or to achieve a serum concentration of 0.01-5000 $\mu\text{g}/\text{mL}$ serum concentration per single (e.g., bolus) administration, multiple administrations or continuous administration, or any effective range or value therein depending on the condition being treated, the route of administration and the age, weight and condition of the subject. In certain embodiments, e.g. for the treatment of cancer, each dose can range from about 0.1 mg to about 50 mg per kilogram of body weight, for example from about 3 mg to about 25 mg per kilogram body weight. The antibody can be formulated as an aqueous solution and administered by subcutaneous injection.

[0137] For treatment of retinal diseases (e.g., age-related macular degeneration (AMD)), the dosage suitably results in aqueous humor concentration of the anti-VEGF antibody the injected eye of 1-50 $\mu\text{g}/\text{mL}$. For treatment of AMD, each dose can be from 0.1 mg to about 1 mg, for example from about 0.25 to about 0.5 mg. The antibody can be formulated as an aqueous solution and administered by intravitreal injection.

[0138] Pharmaceutical compositions can be conveniently presented in unit dose forms containing a predetermined amount of an anti-VEGF antibody of the disclosure per dose. Such a unit can contain for example but without limitation 0.1 mg to 5 g, for example 1 mg to 1 g, or 10 to 50 mg. Pharmaceutically acceptable carriers for use in the disclosure can take a wide variety of forms depending, e.g., on the condition to be treated or route of administration.

[0139] Therapeutic formulations of the anti-VEGF antibodies of the disclosure can be prepared for storage as lyophilized formulations or aqueous solutions by mixing the antibody having the desired degree of purity with optional pharmaceutically-acceptable carriers, excipients or stabilizers typically employed in the art (all of which are referred to herein as "carriers"), *i.e.*, buffering agents, stabilizing agents, preservatives, isotoniifiers, non-ionic detergents, antioxidants, and other miscellaneous additives. See, Remington's Pharmaceutical Sciences, 16th edition (Osol, ed. 1980). Such additives must be nontoxic to the recipients at the dosages and concentrations employed.

[0140] Buffering agents help to maintain the pH in the range which approximates physiological conditions. They can be present at concentration ranging from about 2 mM to about 50 mM. Suitable buffering agents for use with the present disclosure include both organic and inorganic acids and salts thereof such as citrate buffers (e.g., monosodium citrate-disodium citrate mixture, citric acid-trisodium citrate mixture, citric acid-monosodium citrate mixture, *etc.*), succinate buffers (e.g., succinic acid-monosodium succinate mixture, succinic acid-sodium hydroxide mixture, succinic acid-disodium succinate mixture, *etc.*), tartrate buffers (e.g., tartaric acid-sodium tartrate mixture, tartaric acid-potassium tartrate mixture, tartaric acid-sodium hydroxide mixture, *etc.*), fumarate buffers (e.g., fumaric acid-monosodium fumarate mixture, fumaric acid-disodium fumarate mixture, monosodium fumarate-disodium fumarate mixture, *etc.*), gluconate buffers (e.g., gluconic acid-sodium glyconate mixture, gluconic acid-sodium hydroxide mixture, gluconic acid-potassium glyconate mixture, *etc.*), oxalate buffer (e.g., oxalic acid-sodium oxalate mixture, oxalic acid-sodium hydroxide mixture, oxalic acid-potassium oxalate mixture, *etc.*), lactate buffers (e.g., lactic acid-sodium lactate mixture, lactic acid-sodium hydroxide mixture, lactic acid-potassium lactate mixture, *etc.*) and acetate buffers (e.g., acetic acid-sodium acetate mixture, acetic acid-sodium hydroxide mixture, *etc.*). Additionally, phosphate buffers, histidine buffers and trimethylamine salts such as Tris can be used.

[0141] Preservatives can be added to retard microbial growth, and can be added in amounts ranging from 0.2%-1% (w/v). Suitable preservatives for use with the present disclosure include phenol, benzyl alcohol, meta-cresol, methyl paraben, propyl paraben, octadecyldimethylbenzyl ammonium chloride, benzalconium halides (e.g., chloride, bromide, and iodide), hexamethonium chloride, and alkyl parabens such as methyl or propyl paraben, catechol, resorcinol, cy-

clohexanol, and 3-pentanol. Isotonicifiers sometimes known as "stabilizers" can be added to ensure isotonicity of liquid compositions of the present disclosure and include polyhydric sugar alcohols, for example trihydric or higher sugar alcohols, such as glycerin, erythritol, arabitol, xylitol, sorbitol and mannitol. Stabilizers refer to a broad category of excipients which can range in function from a bulking agent to an additive which solubilizes the therapeutic agent or helps to prevent denaturation or adherence to the container wall. Typical stabilizers can be polyhydric sugar alcohols (enumerated above); amino acids such as arginine, lysine, glycine, glutamine, asparagine, histidine, alanine, ornithine, L-leucine, 2-phenylalanine, glutamic acid, threonine, *etc.*, organic sugars or sugar alcohols, such as lactose, trehalose, stachyose, mannitol, sorbitol, xylitol, ribitol, myoinositol, galactitol, glycerol and the like, including cyclitols such as inositol; polyethylene glycol; amino acid polymers; sulfur containing reducing agents, such as urea, glutathione, thiocetic acid, sodium thioglycolate, thioglycerol, α -monothioglycerol and sodium thio sulfate; low molecular weight polypeptides (*e.g.*, peptides of 10 residues or fewer); proteins such as human serum albumin, bovine serum albumin, gelatin or immunoglobulins; hydrophilic polymers, such as polyvinylpyrrolidone monosaccharides, such as xylose, mannose, fructose, glucose; disaccharides such as lactose, maltose, sucrose and trisaccharides such as raffinose; and polysaccharides such as dextran. Stabilizers can be present in the range from 0.1 to 10,000 weights per part of weight active protein.

[0142] Non-ionic surfactants or detergents (also known as "wetting agents") can be added to help solubilize the therapeutic agent as well as to protect the therapeutic protein against agitation-induced aggregation, which also permits the formulation to be exposed to shear surface stressed without causing denaturation of the protein. Suitable non-ionic surfactants include polysorbates (20, 80, *etc.*), polyoxamers (184, 188 *etc.*), Pluronic polyols, polyoxyethylene sorbitan monoethers (TWEEN®-20, TWEEN®-80, *etc.*). Non-ionic surfactants can be present in a range of about 0.05 mg/mL to about 1.0 mg/mL, for example about 0.07 mg/mL to about 0.2 mg/mL.

[0143] Additional miscellaneous excipients include bulking agents (*e.g.*, starch), chelating agents (*e.g.*, EDTA), antioxidants (*e.g.*, ascorbic acid, methionine, vitamin E), and cosolvents.

[0144] The formulation herein can also contain a combination therapeutic agent in addition to the anti-VEGF antibody of the disclosure. Examples of suitable combination therapeutic agents are provided in Section 6.10 below.

[0145] The dosing schedule for subcutaneous administration can vary from once every six months to daily depending on a number of clinical factors, including the type of disease, severity of disease, and the patient's sensitivity to the anti-VEGF antibody.

[0146] The dosage of an anti-VEGF antibody of the disclosure to be administered will vary according to the particular antibody, the type of disease (*e.g.*, cancer, inflammatory, *etc.*), the subject, and the severity of the disease, the physical condition of the subject, the therapeutic regimen (*e.g.*, whether a combination therapeutic agent is used), and the selected route of administration; the appropriate dosage can be readily determined by a person skilled in the art.

[0147] It will be recognized by one of skill in the art that the optimal quantity and spacing of individual dosages of an anti-VEGF antibody of the disclosure will be determined by the nature and extent of the condition being treated, the form, route and site of administration, and the age and condition of the particular subject being treated, and that a physician will ultimately determine appropriate dosages to be used. This dosage can be repeated as often as appropriate. If side effects develop the amount and/or frequency of the dosage can be altered or reduced, in accordance with normal clinical practice.

6.10 COMBINATION THERAPY

[0148] Described below are combinatorial methods in which the anti-VEGF antibodies of the disclosure can be utilized. The combinatorial methods of the disclosure involve the administration of at least two agents to a patient, the first of which is an anti-VEGF antibody of the disclosure, and the second of which is a combination therapeutic agent. The anti-VEGF antibody and the combination therapeutic agent can be administered simultaneously, sequentially or separately.

[0149] The combinatorial therapy methods of the present disclosure can result in a greater than additive effect, providing therapeutic benefits where neither the anti-VEGF antibody or combination therapeutic agent administered in an amount that is alone therapeutically effective.

[0150] In the present methods, the anti-VEGF antibody of the disclosure and the combination therapeutic agent can be administered concurrently, either simultaneously or successively. As used herein, the anti-VEGF antibody of the disclosure and the combination therapeutic agent are said to be administered successively if they are administered to the patient on the same day, for example during the same patient visit. Successive administration can occur 1, 2, 3, 4, 5, 6, 7 or 8 hours apart. In contrast, the anti-VEGF antibody of the disclosure and the combination therapeutic agent are said to be administered separately if they are administered to the patient on the different days, for example, the anti-VEGF antibody of the disclosure and the combination therapeutic agent can be administered at a 1-day, 2-day or 3-day, one-week, 2-week or monthly intervals. In the methods of the present disclosure, administration of the anti-VEGF antibody of the disclosure can precede or follow administration of the combination therapeutic agent.

[0151] As a non-limiting example, the anti-VEGF antibody of the disclosure and combination therapeutic agent can be administered concurrently for a period of time, followed by a second period of time in which the administration of the

anti-VEGF antibody of the disclosure and the combination therapeutic agent is alternated.

[0152] Because of the potentially synergistic effects of administering an anti-VEGF antibody of the disclosure and a combination therapeutic agent, such agents can be administered in amounts that, if one or both of the agents is administered alone, is/are not therapeutically effective.

[0153] In certain aspects, the combination therapeutic agent is an anti-angiogenic agent, an anti-rheumatic drug, an anti-inflammatory agent, a chemotherapeutic agent, a radiotherapeutic, an immunosuppressive agent, or a cytotoxic drug.

[0154] It is contemplated that when used to treat various diseases, the anti-VEGF antibodies of the disclosure can be combined with other therapeutic agents suitable for the same or similar diseases. When used for treating cancer, antibodies of the present disclosure may be used in combination with conventional cancer therapies, such as surgery, radiotherapy, chemotherapy or combinations thereof.

[0155] In some other aspects, other therapeutic agents useful for combination tumor therapy with the antibody of the disclosure include antagonists, e.g., antibodies, of other factors that are involved in tumor growth, such as EGFR, ErbB2 (also known as Her2), ErbB3, ErbB4, or TNF- α .

[0156] Sometimes, for treatment of cancers and immune diseases, it may be beneficial to also administer one or more cytokines to the patient. In a preferred embodiment, the VEGF antibody is co-administered with a growth inhibitory agent.

[0157] Suitable dosages for the growth inhibitory agent are those presently used and may be lowered due to the combined action (synergy) of the growth inhibitory agent and anti-VEGF antibody.

[0158] For treatment of cancers, immune diseases and retinal diseases, anti-inflammatory agents can suitably be used in combination with the anti-VEGF antibodies of the disclosure. Anti-inflammatory agents include, but are not limited to, acetaminophen, diphenhydramine, meperidine, dexamethasone, pentasa, mesalazine, asacol, codeine phosphate, benorylate, fenbufen, naprosyn, diclofenac, etodolac and indomethacin, aspirin and ibuprofen.

[0159] For treatment of cancers, chemotherapeutic agents can suitably be used in combination with the anti-VEGF antibodies of the disclosure. Chemotherapeutic agents include, but are not limited to, radioactive molecules, toxins, also referred to as cytotoxins or cytotoxic agents, which includes any agent that is detrimental to the viability of cells, agents, and liposomes or other vesicles containing chemotherapeutic compounds. Examples of suitable chemotherapeutic agents include but are not limited to 1-dehydrotestosterone, 5-fluorouracil decarbazine, 6-mercaptopurine, 6-thioguanine, actinomycin D, adriamycin, aldesleukin, an anti- $\alpha 5\beta 1$ integrin antibody, alkylating agents, allopurinol sodium, altretamine, amifostine, anastrozole, anthramycin (AMC)), anti-mitotic agents, cis-dichlorodiamine platinum (II) (DDP) cisplatin, diamino dichloro platinum, anthracyclines, antibiotics, antimetabolites, asparaginase, BCG live (intravesical), betamethasone sodium phosphate and betamethasone acetate, bicalutamide, bleomycin sulfate, busulfan, calcium leucovorin, calicheamicin, capecitabine, carboplatin, lomustine (CCNU), carmustine (BSNU), Chlorambucil, Cisplatin, Cladribine, Colchicin, conjugated estrogens, Cyclophosphamide, Cyclothosphamide, Cytarabine, Cytarabine, cytochalasin B, Cytosan, Dacarbazine, Dactinomycin, dactinomycin (formerly actinomycin), daunirubicin HCL, daunorubicin citrate, denileukin difitox, Dexrazoxane, Dibromomannitol, dihydroxy anthracin dione, Docetaxel, dolasetron mesylate, doxorubicin HCL, dronabinol, *E. coli* L-asparaginase, eoloxicimab, emetine, epoetin- α , Erwinia L-asparaginase, esterified estrogens, estradiol, estramustine phosphate sodium, ethidium bromide, ethinyl estradiol, etidronate, etoposide citrororum factor, etoposide phosphate, filgrastim, floxuridine, fluconazole, fludarabine phosphate, fluorouracil, flutamide, folinic acid, gemcitabine HCL, glucocorticoids, goserelin acetate, gramicidin D, granisetron HCL, hydroxyurea, idarubicin HCL, ifosfamide, interferon α -2b, irinotecan HCL, letrozole, leucovorin calcium, leuprolide acetate, levamisole HCL, lidocaine, lomustine, maytansinoid, mechlorethamine HCL, medroxyprogesterone acetate, megestrol acetate, melphalan HCL, mercaptopurine, mesna, methotrexate, methyltestosterone, mithramycin, mitomycin C, mitotane, mitoxantrone, nilutamide, octreotide acetate, ondansetron HCL, paclitaxel, pamidronate disodium, pentostatin, pilocarpine HCL, plimycin, polifeprosan 20 with carmustine implant, porfimer sodium, procaine, procarbazine HCL, propranolol, rituximab, sargramostim, streptozotocin, tamoxifen, taxol, teniposide, tenoposide, testolactone, tetracaine, thioepa chlorambucil, thioguanine, thiotepa, topotecan HCL, toremifene citrate, trastuzumab, tretinoin, valrubicin, vinblastine sulfate, vincristine sulfate, and vinorelbine tartrate.

[0160] Any anti-angiogenic agent can be used in conjunction with the anti-VEGF antibodies of the disclosure, including those listed by Carmeliet and Jain, 2000, Nature 407:249-257. In certain embodiments, the anti-angiogenic agent is another VEGF antagonist or a VEGF receptor antagonist such as VEGF variants, soluble VEGF receptor fragments, aptamers capable of blocking VEGF or VEGFR, neutralizing anti-VEGFR antibodies, low molecule weight inhibitors of VEGFR tyrosine kinases and any combinations thereof. Alternatively, or in addition, two or more anti-VEGF antibodies may be co-administered to the patient.

[0161] In certain embodiments, hormone therapy can be used in conjunction with anti-VEGF antibodies of the disclosure. In some embodiments, the hormone therapy includes one or more agents that inhibit estrogen and/or progesterone from promoting cancer cell growth, e.g., a selective estrogen-receptor modulator such as tamoxifen, an aromatase inhibitor such as anastrozole (Arimidex®) or letrozole (Femara), an aromatase inactivator such as exemestane (Aromasin®), or an agent that inhibits estrogen production such as goserelin (Zoladex). In other embodiments, the hormone therapy is one or more agents that inhibit production of hormones from the ovaries.

[0162] In some aspects, an anti-VEGF antibody can be used in conjunction with a small molecule protein tyrosine kinase (PTK) inhibitor. In some embodiments, the PTK inhibitor is specific for a VEGF receptor tyrosine kinase. In other embodiments, the PTK inhibitor binds to more than one of the VEGF receptor family of tyrosine kinases (e.g., VEGFR-1, VEGFR-2). In other embodiments, protein tyrosine kinase inhibitors useful in the compositions and methods of the invention include PTK inhibitors that do not bind selectively to the VEGF family of receptor tyrosine kinases, but also bind to the tyrosine kinase domains of other families of proteins such as HER2, HER3, HER4, PDGFR, and/or Raf.

[0163] In some embodiments, the tyrosine kinase is a receptor tyrosine kinase, i.e., is an intra-cellular domain of a larger protein that has an extra-cellular ligand binding domain and is activated by the binding of one or more ligands. In certain embodiments, the protein tyrosine kinase is a non-receptor tyrosine kinase. PTK inhibitors for use in the methods of the present disclosure include, but are not limited to, gefitinib (ZD-1839, Iressa®), erlotinib (OSI-1774, Tarceva™), canertinib (CI-1033), vandetanib (ZD6474, Zactima®), tyrphostin AG-825 (CAS 149092-50-2), lapatinib (GW-572016), sorafenib (BAY43-9006), AG-494 (CAS 133550-35-3), RG-13022 (CAS 149286-90-8), RG-14620 (CAS 136831-49-7), BIBW 2992 (Tovok), tyrphostin 9 (CAS 136831-49-7), tyrphostin 23 (CAS 118409-57-7), tyrphostin 25 (CAS 118409-58-8), tyrphostin 46 (CAS 122520-85-8), tyrphostin 47 (CAS 122520-86-9), tyrphostin 53 (CAS 122520-90-5), butein (1-(2,4-dihydroxyphenyl)-3-(3,4-dihydroxyphenyl)-2-propen-1-one 2',3,4,4'-Tetrahydroxychalcone; CAS 487-52-5), curcumin ((E,E)-1,7-bis(4-Hydroxy-3-methoxyphenyl)-1,6-heptadiene-3,5-dione; CAS 458-37-7), N4-(1-Benzyl-1H-indazol-5-yl)-N6,N6-dimethyl-pyrido-[3,4-d]-pyrimidine-4,6-diamine (202272-68-2), AG-1478, AG-879, Cyclopropanecarboxylic acid-(3-(6-(3-trifluoromethyl-phenylamino)-pyrimidin-4-ylamino)-phenyl)-amide (CAS 879127-07-8), N8-(3-Chloro-4-fluorophenyl)-N2-(1-methylpiperidin-4-yl)-pyrimido[5,4-d]pyrimidine-2,8-diamine, 2HCl (CAS 196612-93-8), 4-(4-Benzoyloxylanilino)-6,7-dimethoxyquinazoline (CAS 179248-61-4), N-(4-(3-Chloro-4-fluorophenyl)amino)pyrido[3,4-d]pyrimidin-6-yl)2-butynamide (CAS 881001-19-0), EKB-569, HKI-272, and HKI-357.

[0164] In a specific embodiment, an anti-VEGF antibody of the disclosure is used in combination with intravenous 5-fluorouracil-based chemotherapy. This combination is suitable for, *inter alia*, first- or second-line treatment of patients with metastatic carcinoma of the colon or rectum.

[0165] In another specific embodiment, an anti-VEGF antibody of the disclosure is used in combination with carboplatin and paclitaxel. This combination is suitable for, *inter alia*, first-line treatment of patients with unresectable, locally advanced, recurrent or metastatic non-squamous, non-small cell lung cancer.

[0166] In yet another specific embodiment, an anti-VEGF antibody of the disclosure is used in combination with paclitaxel. This combination is suitable for, *inter alia*, treatment of patients who have not received chemotherapy for metastatic HER2-negative breast cancer.

[0167] For treatment of retinal diseases, the anti-VEGF antibodies of the disclosure can be used in combination with E10030, an anti-platelet-derived growth factor (PDGF) pegylated aptamer; with ARC1905, a pegylated aptamer targeting the C5 component of the complement cascade; and volociximab, a monoclonal antibody targeting the $\alpha 5\beta 1$ integrin transmembrane receptor; photodynamic therapy with Visudyne® (PDT); or Macugen®, an aptamer (pegaptanib sodium).

6.11 THERAPEUTIC REGIMENS

[0168] The present disclosure provides therapeutic regimens involving the administration of the anti-VEGF antibodies of the disclosure. The therapeutic regimen will vary depending on the patient's age, weight, and disease condition. The therapeutic regimen can continue for 2 weeks to indefinitely. In specific embodiments, the therapeutic regimen is continued for 2 weeks to 6 months, from 3 months to 5 years, from 6 months to 1 or 2 years, from 8 months to 18 months, or the like. The therapeutic regimen can be a non-variable dose regimen or a multiple-variable dose regimen.

[0169] For the dosage exemplary regimens described below, the anti-VEGF antibody can be administered as a sterile, preservative-free solution for subcutaneous administration.

[0170] For treatment of metastatic colorectal cancer, an anti-VEGF antibody of the disclosure is administered intravenously at a dose of 0.5-15 mg/kg every 2 weeks with bolus-IFL (irinotecan, 5-fluorouracil and leucovorin regimen). In specific embodiments, the dose is 1-4 mg/kg, 2-6 mg/kg, 0.5-3 mg/kg, 1-10 mg/kg, 3-4.8 mg/kg or 1-4.5 mg/kg every two weeks with bolus-IFL.

[0171] In another embodiment for treatment of metastatic colorectal cancer, an anti-VEGF antibody of the disclosure is administered intravenously at a dose of 1-30 mg/kg every 2 weeks with FOLFOX4 (oxaliplatin, leucovorin, and fluorouracil regimen). In specific embodiments, the dose is 2-9 mg/kg, 3-12 mg/kg, 1-7.5 mg/kg, 2-20 mg/kg, 6-9.75 mg/kg or 4-9.5 mg/kg every two weeks with FOLFOX4.

[0172] For treatment of non-squamous non-small cell lung cancer, an anti-VEGF antibody of the disclosure is administered intravenously at a dose of 2-40 mg/kg every three weeks with carboplatin/paclitaxel. In specific embodiments, the dose is 5-14 mg/kg, 4-20 mg/kg, 10-17.5 mg/kg, 7-14 mg/kg, 10-30 mg/kg or 3-30 mg/kg every three weeks with carboplatin/paclitaxel.

[0173] For treatment of metastatic breast cancer, an anti-VEGF antibody of the disclosure is administered intravenously at a dose of 0.5-20 mg/kg every two weeks with paclitaxel. In specific embodiments, the dose is 1-4 mg/kg, 2-6 mg/kg,

0.5-3 mg/kg, 1-10 mg/kg, 3-4.8 mg/kg or 1-4.5 mg/kg every two weeks with paclitaxel.

[0174] For treatment of metastatic breast cancer, an anti-VEGF antibody of the disclosure is administered intravenously at a dose of 0.5-20 mg/kg every two weeks as monotherapy. In specific embodiments, the dose is 1-4 mg/kg, 2-6 mg/kg, 0.5-3 mg/kg, 1-10 mg/kg, 3-4.8 mg/kg or 1-4.5 mg/kg every two weeks as monotherapy.

[0175] For treatment of age-related macular degeneration (AMD), an anti-VEGF antibody of the disclosure is administered at a dose of 0.1-1 mg by intravitreal injection once a month (approximately 28 days). In specific embodiments, the dose is 0.1-0.4 mg, 0.2-0.6 mg, 0.1-0.25 mg, 0.25-0.5 mg, 0.25-0.75 mg, or 0.3-0.45 mg by intravitreal injection once a month (approximately 28 days). In a specific embodiment, a patient treated with an anti-VEGF antibody of the disclosure has wet AMD. In another specific embodiment, a patient has dry AMD.

6.12 DIAGNOSTIC AND PHARMACEUTICAL KITS

[0176] Encompassed by the present disclosure are pharmaceutical kits containing the anti-VEGF antibodies (including antibody conjugates) of the disclosure. The pharmaceutical kit is a package comprising the anti-VEGF antibody of the disclosure (*e.g.*, either in lyophilized form or as an aqueous solution) and one or more of the following:

- A combination therapeutic agent, for example as described in Section 6.10 above;
- A device for administering the anti-VEGF antibody, for example a pen, needle and/or syringe; and
- Pharmaceutical grade water or buffer to resuspend the antibody if the antibody is in lyophilized form.

[0177] In certain aspects, each unit dose of the anti-VEGF antibody is packaged separately, and a kit can contain one or more unit doses (*e.g.*, two unit doses, three unit doses, four unit doses, five unit doses, eight unit doses, ten unit doses, or more). In a specific embodiment, the one or more unit doses are each housed in a syringe or pen.

[0178] Diagnostic kits containing the anti-VEGF antibodies (including antibody conjugates) of the disclosure are also encompassed herein. The diagnostic kit is a package comprising the anti-VEGF antibody of the disclosure (*e.g.*, either in lyophilized form or as an aqueous solution) and one or more reagents useful for performing a diagnostic assay. Where the anti-VEGF antibody is labeled with an enzyme, the kit can include substrates and cofactors required by the enzyme (*e.g.*, a substrate precursor which provides the detectable chromophore or fluorophore). In addition, other additives can be included, such as stabilizers, buffers (*e.g.*, a block buffer or lysis buffer), and the like. In certain embodiments, the anti-VEGF antibody included in a diagnostic kit is immobilized on a solid surface, or a solid surface (*e.g.*, a slide) on which the antibody can be immobilized is included in the kit. The relative amounts of the various reagents can be varied widely to provide for concentrations in solution of the reagents which substantially optimize the sensitivity of the assay. In a specific embodiment, the antibody and one or more reagents can be provided (individually or combined) as dry powders, usually lyophilized, including excipients which on dissolution will provide a reagent solution having the appropriate concentration.

7. EXAMPLE 1: IDENTIFICATION OF T-CELL EPITOPES OF BEVACIZUMAB

7.1 MATERIALS & METHODS

7.1.1 Peptides

[0179] Peptides were synthesized using a multi-pin format by Mimotopes (Adelaide, Australia). The sequences of the bevacizumab light and heavy chain V regions were synthesized as 15-mer peptides overlapping by 12 amino acids (Tables 3 and 4) for a total of 69 peptides. Peptides arrived lyophilized and were re-suspended in DMSO (Sigma-Aldrich) at approximately 1-2 mg/mL. Stock peptides were kept frozen at -20°C.

7.1.2 Human Peripheral Blood Mononuclear Cells

[0180] Community donor buffy coat products were purchased from the Stanford Blood Center, Palo Alto, CA. Buffy coat material was diluted 1:1 v:v with DPBS containing no calcium or magnesium. Diluted buffy coat material (25-35 mls) was underlaid in 50 ml conical centrifuge tubes (Sarstedt or Costar) with 12.5 mls of FicollPaque-PLUS (GE Healthcare). The samples were centrifuged at 900 g for 30 minutes at room temperature. Peripheral blood mononuclear cells (PBMC) were collected from the interface. DPBS was added to bring the final volume to 50 mls and the cells were centrifuged at 350 g for 5 minutes. Pelleted cells were resuspended in DPBS and counted.

7.1.3 Dendritic cells

[0181] For isolation of dendritic cells, T75 culture flasks (Costar) were seeded with 10^8 freshly isolated PBMC in a total volume of 30 mls AIM V media (Invitrogen). Excess PBMC were frozen at -80°C in 90% fetal calf serum (FCS), 10% DMSO at 5×10^7 cells/mL. T75 flasks were incubated at 37°C in 5% CO_2 for 2 hours. Nonadherent cells were removed, and the adherent monolayer was washed with DPBS. To differentiate dendritic cells from monocytes, 30 mls of AIM V media containing 800 units/mL of GM-CSF (R and D Systems) and 500 units/mL IL-4 (R and D Systems) was added. Flasks were incubated for 5 days. On day 5 IL-1 α (Endogen) and TNF α (Endogen) were added to 50 pg/mL and 0.2 ng/mL. Flasks were incubated two more days. On day 7, dendritic cells were collected by the addition of 3 mls of 100 mM EDTA containing 0.5 to 1.0 mg Mitomycin C (Sigma-Aldrich) for a final concentration of 10 mM EDTA and 16.5 to 33 $\mu\text{g}/\text{mL}$ Mitomycin C. Flasks were incubated an additional hour at 37°C and 5% CO_2 . Dendritic cells were collected, and washed in AIM V media 2-3 times.

7.1.4 Cell culture

[0182] On day 7, previously frozen autologous PBMC were thawed quickly in a 37°C water bath. Cells were immediately diluted into DPBS or AIM V media and centrifuged at 350g for 5 minutes. CD4^+ cells were enriched by negative selection using magnetic beads (Easy-Sep CD4^+ kit, Stem Cell Technologies). Autologous CD4^+ T cells and dendritic cells were cocultured at 2×10^5 CD4^+ T cells per 2×10^4 dendritic cells per well in 96 well round bottomed plates (Costar 9077). Peptides were added at approximately 5 $\mu\text{g}/\text{mL}$. Control wells contained the DMSO (Sigma) vehicle alone at 0.25% v:v. Positive control wells contained DMSO at 0.25% and tetanus toxoid (List Biologicals or CalBioChem) at 1 $\mu\text{g}/\text{mL}$. Cultures were incubated for 5 days. On day 5, 0.25 μCi per well of tritiated thymidine (Amersham or GE Healthcare) was added. Cultures were harvested on day 6 to filtermats using a Packard Filtermate Cell harvester. Scintillation counting was performed using a Wallac MicroBeta 1450 scintillation counter (Perkin Elmer).

7.1.5 Data Analysis

[0183] Average background CPM values were calculated by averaging individual results from 6 to 12 replicates. The CPM values of the four positive control wells were averaged. Replicate or triplicate wells for each peptide were averaged. Stimulation index values for the positive control and the peptide wells were calculated by dividing the average experimental CPM values by the average control values. In order to be included in the dataset, a stimulation index of greater than 3.0 in the tetanus toxoid positive control wells was required. A response was noted for any peptide resulting in a stimulation index of 2.95 or greater. Peptides were tested using peripheral blood samples from a group of 99 donors. Responses to all peptides were compiled. For each peptide tested, the percentage of the donor set that responded with a stimulation index of 2.95 or greater was calculated. In addition, the average stimulation index for all donors was also calculated.

7.2 RESULTS

7.2.1 Identification Of CD4^+ T Cell Epitopes In The Bevacizumab VH And VL Regions

[0184] CD4^+ T cell epitope peptides were identified by an analysis of the percent responses to the peptides within the set of 99 donors. The average percent response and standard deviation were calculated for all peptides tested describing the bevacizumab heavy chain and light chain V regions. A response rate greater than or equal to the average background response plus three standard deviations was considered a potential CD4^+ T cell epitope. For the bevacizumab light chain V region, 32 peptides were tested (Table 3) which resulted in an average background percent response of $2.1 \pm 2.7\%$. Three standard deviations above background was determined to be 10.2%. One peptide at position 13 (Q40-T54) displayed this level of response in the bevacizumab light chain peptide dataset, with a response rate of 15.2% (Figure 2A). For the bevacizumab heavy chain V region, 37 peptides were tested (Table 4). The average background percent response was $2.8 \pm 3.1\%$. Three standard deviations above background was 12.1%. One peptide within the bevacizumab heavy chain dataset, #18 (N52-R56), achieved a percent response of 16.2% (Figure 3A). A second peptide at position #30 in the heavy chain dataset achieved a response rate of 9.1%, and was considered an epitope due to an increased stimulation index (see below).

[0185] The average stimulation index was calculated for all peptides in the dataset. Light chain peptide 13 had a high average stimulation index of 1.82 ± 0.24 s.e.m. (Figure 2B). Heavy chain peptide #18 had an average stimulation index value of 2.16 ± 0.35 s.e.m. (Figure 3B). The peptide at position #30 returned an average stimulation index of $1.45 + 0.18$ s.e.m. (Figure 3B) due to an elevated average stimulation index and an above average response rate. The peptide at position #30 was included when determining CD4^+ T cell epitope content of this antibody V region. All of these stimulation index values are significantly higher than the average stimulation index for all peptides in the two datasets

(1.14 ± 0.07 for all 69 heavy chain and light chain peptides).

[0186] These data indicate that there are three CD4+ T cell epitope regions in the bevacizumab V regions (Table 5). In the VL region, an epitope is found at peptide position 13 that encompasses framework 2 and two amino acids from CDR2. The sequence contains a murine back-mutation (V46) inserted into the sequence during humanization. In Table 5, the CDR-derived amino acids are underlined. In the heavy chain, two epitope peptide regions were identified. The stronger epitope at position #18 encompasses all of CDR2. The second epitope peptide region contains both framework 3 and CDR3 amino acids.

7.2.2 Reduced Immunogenicity Variants of Bevacizumab Variant Antibodies

[0187] Bevacizumab was subjected to mutational analysis (see Example 2 below). Based on antigen-binding studies performed in conjunction with the mutational analysis, a set of candidate amino acid substitutions within the CDR-H2 and CDR-H3 region were identified that did not significantly reduce the affinity of the antibody to VEGF (Table 6). These amino acid substitutions were tested singly and in combination to identify variants of bevacizumab with reduced immunogenicity as compared to the wild type antibody.

8. EXAMPLE 2: IDENTIFICATION OF VARIANTS OF BEVACIZUMAB WITH INCREASED AFFINITY TO VEGF

[0188] The bevacizumab antibody was subjected to comprehensive mutational analysis to identify mutants that had increased affinity to VEGF as compared to bevacizumab. The increased affinity of candidate high affinity mutants to VEGF as compared to bevacizumab was analyzed by BIAcore to confirm their binding characteristics.

8.1 MATERIALS & METHODS

8.1.1 BIAcore

[0189] Fifteen variant bevacizumab VH region constructs were cloned along with the unmodified VL region into a human IgG₁-containing plasmid, expressed in 293T/17 cell lines by transient transfection, and antibodies purified by Protein A or Protein G affinity. The affinity of the antibodies for VEGF (R&D systems, Minneapolis, MN) was determined by using a BIAcore 2000 and 3000 surface plasmon resonance system (BIAcore, GE Healthcare, Piscataway, NJ). Polyclonal goat anti-human Fc antibody (Jackson ImmunoResearch) was first immobilized to the biosensor surface using standard BIAcore amine coupling reagents (N-ethyl-N'-dimethylamino-propylcarbodiimide, EDC; N-hydroxysuccinimide, NHS; and ethanolamine HCl, pH 8.5), followed by the capture of anti-VEGF antibodies (bevacizumab and bevacizumab variants) on parallel surfaces at a low flow rate of 5 µL/min. RL was kept low to minimize avidity due to the dimeric nature of VEGF. No capture of the antibody was made on the reference surface to serve as a negative control. Subsequently, VEGF was injected to all flow cells at a flow rate of 50 µL/min for two minutes to monitor association followed by a 25-minute flow of HBS-P running buffer (10 mM HEPES, 150 mM sodium chloride, 0.005% P-20, pH 7.4) to monitor the dissociation phase. At each cycle, VEGF, in 6 different concentrations of VEGF ranging between 0 nM and 512nM and at four-fold increments, was injected over the surface. The surface was regenerated with 1.5% H₃PO₄ at a flow rate of 100 µL/min in two brief pulses at the end of each cycle. Binding data were fit to the 1:1 Langmuir model to extract binding constants from the BIAevaluate software. Double referencing was applied in each analysis to eliminate background responses from the reference surface and buffer only control. All the binding kinetics data were analyzed at least three separate determinations.

8.2 RESULTS

[0190] Results are displayed as absolute numbers and as fold improvement over wild-type. Almost all the variants listed have improved association (k_{on}) and dissociation (k_{off}) rates when compared to bevacizumab or wild-type (Table 7). The final affinity values for the variants were in the 0.1 nM range and reach as low as 0.08 nM for the variant corresponding to SEQ ID NO:82. These values contrast to bevacizumab which has a measured affinity in these experiments of 1.9 nM.

[0191] Tables 8 and 9 show additional heavy chain variants that preliminary binding studies show have a greater affinity to VEGF than bevacizumab (data not shown). Table 10 shows heavy chain variants that preliminary studies indicate have an affinity to VEGF similar to that of bevacizumab (data not shown). Table 11 shows light chain variants that preliminary studies indicate have an affinity to VEGF similar to that of bevacizumab (data not shown).

9. EXAMPLE 3: SELECTION OF DEIMMUNIZED VARIANT PEPTIDES

[0192] Variant peptides corresponding to the immunogenic regions of bevacizumab (see Example 1) were generated (Tables 14-16). The variant peptides were selected on the basis of comprehensive mutational analysis described in Example 2, in which CDR modifications were identified that did not substantially reduce the binding affinity of bevacizumab to VEGF.

[0193] A total of 77 peptides were synthesized and tested based on the antigen-binding studies, including two syntheses of each of the parent 15-mer peptides. A total of 93 donors were tested with the parent and variant peptides utilizing the method described in Section 7.1 and the results are shown in Figures 4A-4C. In particular, Figures 4A-4C show CD4+ T cell responses to mutant bevacizumab epitope peptides. Average responses to the unmodified parent epitope sequences are indicated with open marks. Large circles indicate selected peptides referred to in Table 17 (see below). Figure 4A shows VH CDR2 peptides; Figure 4B shows VH CDR3 peptides; and Figure 4C shows VL CDR2 peptides. Immunogenicity data for selected peptides are shown in Table 17.

[0194] For the heavy chain variable region CDR2 peptides, the average percent response to the parent peptides in this study was 5.38% and 6.45%. Three mutant peptides demonstrated a reduced overall response rate and average stimulation index as compared to the parent peptides.

[0195] The parent peptide response rates for the heavy chain variable region CDR3 epitope peptides in this study were 7.53% and 6.45%. A single mutant peptide sequence was found that demonstrated reduced overall responses as compared to the parent peptide.

[0196] Finally, the light chain variable region CDR2 peptide parent response rates in this study were 25.8% and 15%. One mutant peptide was identified that demonstrated a reduced overall immunogenicity as compared to the parent peptide.

[0197] To demonstrate that the deimmunizing mutations maintained affinity to VEGF as compared to bevacizumab, flow cytometry was used to compare the binding properties of variant antibodies incorporating mutations in the modified epitope peptides (either as single or double amino-acid modifications and bevacizumab). Several deimmunizing mutations had comparable or increased affinity to VEGF as compared to bevacizumab.

[0198] In one study, transiently transfected 293c18 cells expressing surface-bound forms of the bevacizumab variants were stained with Alexa647-conjugated rHuVEGF (Invitrogen Cat #PHG0143) at 3 nM and goat-anti-human-kappa-RPE (Southern Biotech Cat#2063-09) at a 1:400 dilution. Data were gathered by way of flow cytometry using a DakoCytomation CyAn ADP flow cytometer and was analyzed using Treestar's FloJo analysis program. The mean fluorescence intensities (MFI) measured in this work are set forth in Table 18.

[0199] In another study, the EC_{50} of bevacizumab and variant antibody binding to VEGF was measured. Antibody titration plots were generated using bevacizumab and its variants with Alexas647-conjugated rHu VEGF as described above, with the VEGF serially diluted two-fold from 5 μ M to 0.01 μ M. The EC_{50} values are shown in Table 19.

10. SPECIFIC EMBODIMENTS, CITATION OF REFERENCES

[0200] All publications, patents, patent applications and other documents cited in this application are hereby incorporated by reference in their entireties for all purposes to the same extent as if each individual publication, patent, patent application or other document were individually indicated to be incorporated by reference for all purposes. While various specific embodiments have been illustrated and described, it will be appreciated that various changes can be made without departing from the spirit and scope of the invention(s).

PARAGRAPHS OF THE INVENTION:**[0201]**

1. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein CDR-H2 includes the substitution K64S, and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9, or 13-16, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab.

2. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein CDR-H2 includes the substitution K64Q, and optionally one or more additional mutations or combinations of mutations selected from one or more of

Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9, or 13-16, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab.

5 3. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein CDR-H2 includes the substitutions Y53F and K64Q, and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9, or 13-16, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab.

10 4. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein CDR-H3 includes the substitutions H97E and Y98F, and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9, or 13-16, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab.

15 5. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein CDR-L2 includes the substitution T51A and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9, or 13-16, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab, and wherein CDR-L2 in said VEGF antibody does not consist of a CDR-L2 sequence set forth in Table 13.

20 6. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises six complementarity determining regions ("CDRs") having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein said CDR-H1 includes the substitution N31F, and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9 and 13, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab, and wherein CDR-H1 in said VEGF antibody does not consist of a CDR-H1 sequence set forth in Tables 12-1 to 12-9.

25 7. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises six CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein said CDR-H3 includes the substitution H97E, and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9 and 13, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab, and wherein CDR-H3 in said VEGF antibody does not consist of a CDR-H3 sequence set forth in Tables 12-1 to 12-9.

30 8. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises six CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein said CDR-H3 includes the substitution H97D, and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9 and 13, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab, and wherein CDR-H3 in said VEGF antibody does not consist of a CDR-H3 sequence set forth in Tables 12-1 to 12-9.

35 9. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises six CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein said CDR-H3 includes the substitution H97P, and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9 and 13, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab, and wherein CDR-H3 in said VEGF antibody does not consist of a CDR-H3 sequence set forth in Tables 12-1 to 12-9.

10. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises six CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein said CDR-H3 includes the substitution Y99E, and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9 and 13, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab, and wherein CDR-H3 in said VEGF antibody does not consist of a CDR-H3 sequence set forth in Tables 12-1 to 12-9.

11. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises six CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein said CDR-H3 includes the substitution Y99D, and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9 and 13, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab, and wherein CDR-H3 in said VEGF antibody does not consist of a CDR-H3 sequence set forth in Tables 12-1 to 12-9.

12. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises six CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein said CDR-H3 includes the substitution S100aG, and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9 and 13, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab, and wherein CDR-H3 in said VEGF antibody does not consist of a CDR-H3 sequence set forth in Tables 12-1 to 12-9.

13. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein said CDR-H3 includes the substitution S100aG, and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9 and 13, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab, and wherein position 3 in CDR-H3 of said VEGF antibody is not tyrosine.

14. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein CDR-H1 includes at least one substitution selected from T28P, N31F, N31G and N31M, and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9 and 13, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab, and wherein CDR-H1 in said VEGF antibody does not consist of a CDR-H1 sequence set forth in Tables 12-1 to 12-9.

15. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein CDR-H3 includes at least one substitution selected from H97A, H97Q, H97S, H97T, S100aD, S100aE, and S100aV, and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9 and 13, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab, and wherein CDR-H3 in said VEGF antibody does not consist of a CDR-H3 sequence set forth in Tables 12-1 to 12-9.

16. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein CDR-H1 includes the substitution T30W, T30R or T30Q, and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9 and 13, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab, and wherein CDR-H1 in said VEGF antibody does not consist of a CDR-H1 sequence set forth in Tables 12-1 to 12-9.

17. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein CDR-H2 includes at least one substitution selected from Y53F, T58F, A61G, A61K, A61R, A61H, A61Y, K64G, K64E, R65L, R65T, R65A, R65E, and R65D, and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9 and 13, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab, and wherein CDR-H2 in said VEGF antibody does not consist of a CDR-H2 sequence set forth in Tables 12-1 to 12-9.

18. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein CDR-H3 includes at least one substitution selected from Y98F and Y100eF, and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9 and 13, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab, and wherein CDR-H3 in said VEGF antibody does not consist of a CDR-H3 sequence set forth in Tables 12-1 to 12-9.

19. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises CDRs having amino acid sequences corresponding SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein the heavy chain CDRs include a combination of substitutions selected from:

- (a) N31F in CDR-H1, H97D in CDR-H3, Y99D in CDR-H3, and S100aG in CDR-H3;
- (b) N31F in CDR-H1, H97P in CDR-H3, Y99D in CDR-H3, and S100aG in CDR-H3;
- (c) N31F in CDR-H1, H97P in CDR-H3, and Y99E in CDR-H3;
- (d) N31F in CDR-H1, H97E in CDR-H3, and Y99E in CDR-H3;
- (e) N31F in CDR-H1, H97D in CDR-H3, and Y99E in CDR-H3;
- (f) N31F in CDR-H1, H97E in CDR-H3, Y99D in CDR-H3, and S100aG in CDR-H3;
- (g) N31F in CDR-H1, Y99D in CDR-H3, and S100aG in CDR-H3;
- (h) N31F in CDR-H1, H97P in CDR-H3, and Y99D in CDR-H3;
- (i) N31F in CDR-H1, H97D in CDR-H3, and S100aG in CDR-H3;
- (j) N31F in CDR-H1 and S100aG in CDR-H3;
- (k) N31F in CDR-H1, H97P in CDR-H3, and S100aG in CDR-H3;

wherein said antibody or binding fragment optionally comprises one or more additional mutations or combinations of mutations selected from one or more of Tables 6, 7, 8, 9, 10, 11, 12-1 to 12-9 and 13, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab.

20. The anti-VEGF antibody or anti-VEGF binding fragment of paragraph 19, in which, other than the combination of substitutions selected from (a) to (k), the heavy chain CDRs do not comprise additional mutations as compared to the heavy chain CDR sequences of the antibody bevacizumab or as compared to the heavy chain CDR sequences of the antibody ranibizumab.

21. The anti-VEGF antibody or anti-VEGF binding fragment of any one of paragraphs 1 to 19, wherein the six CDRs altogether have up to 16, up to 15, up to 14, up to 13, up to 12, up to 11, up to 10, up to 9, up to 8, up to 7, up to 6, up to 5, or up to 4 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab.

22. The anti-VEGF antibody or anti-VEGF binding fragment of any one of paragraphs 1 to 21, wherein any individual CDR has no more than three amino acid substitutions as compared to the corresponding CDR sequence of the antibody bevacizumab or the antibody ranibizumab.

23. The anti-VEGF antibody or anti-VEGF binding fragment of anti-VEGF antibody or anti-VEGF binding fragment of any one of paragraphs 1 to 21, wherein any individual CDR has no more than two amino acid substitutions as compared to the corresponding CDR sequence of the antibody bevacizumab or the antibody ranibizumab.

EP 2 894 167 A1

24. The anti-VEGF antibody or anti-VEGF binding fragment of any one of paragraphs 1 to 23 which is a monoclonal antibody or anti-VEGF binding fragment of a monoclonal antibody, respectively.
- 5 25. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of paragraphs 1 to 24 which is a human or humanized antibody, or anti-VEGF binding fragment of a humanized or humanized antibody, respectively.
26. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of paragraphs 1 to 25 which is an IgG.
- 10 27. The anti-VEGF antibody or an anti-VEGF binding fragment of paragraph 26 which is an IgG₁.
28. The anti-VEGF antibody or an anti-VEGF binding fragment of paragraph 26 which is an IgG₂.
29. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of paragraphs 1 to 28 which includes one or more mutations in the Fc region that increase ADCC activity.
- 15 30. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of paragraphs 1 to 29 which is non-fucosylated.
31. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of paragraphs 1 to 30 which includes one or more mutations in the Fc region that increase binding to Fc γ R.
- 20 32. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of paragraphs 1 to 30 which includes one or more mutations in the Fc region that increase binding to FcRn.
- 25 33. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of paragraphs 1 to 27 which includes one or more mutations in the Fc region that decrease ADCC activity.
34. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of paragraphs 1 to 34 which has, other than said one or more mutations, a V_H sequence corresponding to SEQ ID NO:1 and a V_L sequence corresponding to SEQ ID NO:2.
- 30 35. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of paragraphs 1 to 34 which has, other than said one or more mutations, a heavy chain sequence corresponding to SEQ ID NO:9 and a light chain sequence corresponding to SEQ ID NO:10.
- 35 36. The anti-VEGF antibody or anti-VEGF binding fragment of any one of paragraphs 1 to 34 which has an affinity that is 1.5- to 50-fold greater than the affinity of an antibody having a V_H sequence corresponding to SEQ ID NO:1 and a V_L sequence corresponding to SEQ ID NO:2.
- 40 37. The anti-VEGF antibody or anti-VEGF binding fragment of any one of paragraphs 1 to 34 which has an affinity that is 1.5- to 50-fold greater than the affinity of an antibody having a heavy chain sequence corresponding to SEQ ID NO:9 and a light chain sequence corresponding to SEQ ID NO:10.
- 45 38. The anti-VEGF antibody or anti-VEGF binding fragment of paragraph 36 which has a 2- to 30-fold greater than the affinity of an antibody having a V_H sequence corresponding to SEQ ID NO:1 and a V_L sequence corresponding to SEQ ID NO:2.
- 50 39. The anti-VEGF antibody or anti-VEGF binding fragment of paragraph 37 which has a 2- to 30-fold greater than the affinity of an antibody having a heavy chain sequence corresponding to SEQ ID NO:9 and a light chain sequence corresponding to SEQ ID NO:10.
40. The anti-VEGF antibody or anti-VEGF binding fragment of any one of paragraphs 36 to 39, wherein the affinity is a measure of K_D as analyzed by a biosensor.
- 55 41. The anti-VEGF antibody or anti-VEGF binding fragment of any one of paragraphs 1 to 40 which is purified.
42. The anti-VEGF antibody or anti-VEGF binding fragment of paragraph 41 which is purified to at least 85%, at least 90%, at least 95% or at least 98% homogeneity.

43. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises six complementarity determining regions ("CDRs") having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein the heavy chain includes at least one substitution selected from A61F in CDR-H2, A61E in CDR-H2, A61D in CDR-H2, D62L in CDR-H2, D62G in CDR-H2, D62Q in CDR-H2, D62T in CDR-H2, D62K in CDR-H2, D62R in CDR-H2, D62E in CDR-H2, D62H in CDR-H2, K64S in CDR-H2, K64V in CDR-H2, K64Q in CDR-H2, R65V in CDR-H2, R65F in CDR-H2, R65H in CDR-H2, R65N in CDR-H2, R65S in CDR-H2, R65Q in CDR-H2, R65K in CDR-H2, R65I in CDR-H2, and Y98H in CDR-H3, and optionally one or more additional mutations or combinations of mutations selected from one or more of Tables 7, 8, 9, 10, 11, 12-1 to 12-9 and 13, wherein the six CDRs altogether have up to 17 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab.

44. The anti-VEGF antibody or an anti-VEGF binding fragment of paragraph 43 wherein wherein the six CDRs altogether have up to 16, up to 15, up to 14, up to 13, up to 12, up to 11, up to 10, up to 9, up to 8, up to 7, up to 6, up to 5, or up to 4 amino acid substitutions as compared to CDR sequences of the antibody bevacizumab or the antibody ranibizumab.

45. The anti-VEGF antibody or an anti-VEGF binding fragment of paragraph 43 or paragraph 44, wherein any individual CDR has no more than three amino acid substitutions as compared to the corresponding CDR sequence of the antibody bevacizumab.

46. The anti-VEGF antibody or an anti-VEGF binding fragment of paragraph 43 or paragraph 44, wherein any individual CDR has no more than two amino acid substitutions as compared to the corresponding CDR sequence of the antibody bevacizumab.

47. The anti-VEGF antibody or anti-VEGF binding fragment of any one of paragraphs 43 to 46 which is a monoclonal antibody or an anti-VEGF binding fragment of a monoclonal antibody, respectively.

48. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of paragraphs 43 to 47 which is a human or humanized antibody, or an anti-VEGF binding fragment of a humanized or humanized antibody, respectively.

49. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of paragraphs 43 to 48 which is an IgG.

50. The anti-VEGF antibody or an anti-VEGF binding fragment of paragraph 49 which is an IgG₁.

51. The anti-VEGF antibody or an anti-VEGF binding fragment of paragraph 49 which is an IgG₂.

52. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of paragraphs 43 to 51 which includes one or mutations in the Fc region that increase ADCC activity.

53. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of paragraphs 43 to 52 which is non-fucosylated.

54. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of paragraphs 43 to 53 which includes one or more mutations in the Fc region that increase binding to Fc γ R.

55. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of paragraphs 43 to 53 which includes one or more mutations in the Fc region that increase binding to FcRn.

56. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of paragraphs 43 to 50 which includes one or mutations in the Fc region that decrease ADCC activity.

57. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of paragraphs 43 to 56 which has, other than said one or more mutations, a V_H sequence corresponding to SEQ ID NO:1 and a V_L sequence corresponding to SEQ ID NO:2.

58. The anti-VEGF antibody or anti-VEGF binding fragment of any one of paragraphs 43 to 57 which is purified.

59. The anti-VEGF antibody or anti-VEGF binding fragment of paragraph 58 which is purified to at least 85%, at least 90%, at least 95% or at least 98% homogeneity.
- 5 60. An antibody-drug conjugate comprising an anti-VEGF antibody or anti-VEGF binding fragment according to any one of paragraphs 1 to 59.
61. A pharmaceutical composition comprising an anti-VEGF antibody or anti-VEGF binding fragment according to any one of paragraphs 1 to 59 or an antibody-drug conjugate according to paragraph 60, and a pharmaceutically acceptable carrier.
- 10 62. A nucleic acid comprising a nucleotide sequence encoding anti-VEGF antibody or anti-VEGF binding fragment of any one of paragraphs 1 to 36 and 43 to 57.
63. A vector comprising the nucleic acid of paragraph 62.
- 15 64. A prokaryotic host cell transformed with a vector according to paragraph 63.
65. A eukaryotic host cell transformed with a vector according to paragraph 63.
- 20 66. A eukaryotic host cell engineered to express the nucleotide sequence of paragraph 62.
67. The eukaryotic host cell of paragraph 66 which is a mammalian host cell.
- 25 68. A method of producing anti-VEGF antibody or anti-VEGF binding fragment, comprising: (a) culturing the eukaryotic host cell of paragraph 66 or paragraph 67 and (b) recovering the anti-VEGF antibody or anti-VEGF binding fragment antibody.
69. A method of treating cancer, comprising administering to a human patient in need thereof a therapeutically effective amount of anti-VEGF antibody or anti-VEGF binding fragment according to any one of paragraphs 1 to 59, an antibody-drug conjugate according to paragraph 60, or a pharmaceutical composition according to paragraph 61.
- 30 70. The method of paragraph 69, wherein the cancer is metastatic carcinoma of the colon, metastatic carcinoma of the rectum, non-squamous non-small cell lung cancer, or metastatic HER2-negative breast cancer.
- 35 71. The method of paragraph 70, wherein the cancer is non-squamous non-small cell lung cancer which is unresectable, locally advanced, recurrent or metastatic.
72. A method of treating age-related macular degeneration, comprising administering to a human patient in need thereof a therapeutically effective amount of anti-VEGF antibody or anti-VEGF binding fragment according to any one of paragraphs 1 to 59, an antibody-drug conjugate according to paragraph 60, or a pharmaceutical composition according to paragraph 61.
- 40 73. A method of treating an immune disorder, comprising administering to a human patient in need thereof a therapeutically effective amount of anti-VEGF antibody or anti-VEGF binding fragment according to any one of paragraphs 1 to 59, an antibody-drug conjugate according to paragraph 60, or a pharmaceutical composition according to paragraph 61.
- 45 74. The method of paragraph 73, wherein the immune disorder is rheumatoid arthritis or Grave's disease.
- 50
- 55

EP 2 894 167 A1

SEQUENCE LISTING

<110> AbbVie Biotherapeutics, Inc.

5

<120> ANTI-VEGF ANTIBODIES AND THEIR USES

<130> P101835EP52

<140>

<141>

10

<150> US 61/218,005

<151> 2009-06-17

<160> 412

15

<170> PatentIn version 3.5

<210> 1

<211> 123

<212> PRT

20

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

25

<400> 1

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

30

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
20 25 30

35

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

40

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
50 55 60

45

Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
65 70 75 80

50

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Tyr Pro His Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
100 105 110

55

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 2

<211> 108

EP 2 894 167 A1

<212> PRT
<213> Artificial Sequence

<220>

5

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 2

10

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr
20 25 30

15

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile
35 40 45

20

Tyr Phe Thr Ser Ser Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

25

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Thr Val Pro Trp
85 90 95

30

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
100 105

<210> 3

35

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

40

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 3

45

Gly Tyr Thr Phe Thr Asn Tyr Gly Met Asn
1 5 10

<210> 4

50

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

55

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 4

EP 2 894 167 A1

Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15

5 Arg

10 <210> 5
<211> 14
<212> PRT
<213> Artificial Sequence

15 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

20 <400> 5
Tyr Pro His Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
1 5 10

25 <210> 6
<211> 11
<212> PRT
<213> Artificial Sequence

30 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

35 <400> 6
Ser Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn
1 5 10

40 <210> 7
<211> 7
<212> PRT
<213> Artificial Sequence

45 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

50 <400> 7
Phe Thr Ser Ser Leu His Ser
1 5

55 <210> 8
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

EP 2 894 167 A1

<400> 8
 Gln Gln Tyr Ser Thr Val Pro Trp Thr
 1 5

5

<210> 9
 <211> 231
 <212> PRT
 <213> Artificial Sequence

10

<220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic polypeptide"

15

<400> 9
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

20

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Asp Phe Thr His Tyr
 20 25 30

25

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
 50 55 60

30

Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
 65 70 75 80

35

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Lys Tyr Pro Tyr Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val
 100 105 110

40

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly
 115 120 125

45

Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly
 130 135 140

50

Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val
 145 150 155 160

Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe
 165 170 175

55

Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val
 180 185 190

EP 2 894 167 A1

Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val
 195 200 205

5 Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys
 210 215 220

10 Ser Cys Asp Lys Thr His Leu
 225 230

<210> 10
 <211> 214
 <212> PRT
 15 <213> Artificial Sequence

<220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 20 polypeptide"

<400> 10
 Asp Ile Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15

25 Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr
 20 25 30

30 Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile
 35 40 45

35 Tyr Phe Thr Ser Ser Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80

40 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Thr Val Pro Trp
 85 90 95

45 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
 100 105 110

Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
 115 120 125

50 Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
 130 135 140

55 Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
 145 150 155 160

EP 2 894 167 A1

Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
165 170 175

5 Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
180 185 190

10 Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
195 200 205

Phe Asn Arg Gly Glu Cys
210

15 <210> 11
<211> 10
<212> PRT
<213> Artificial Sequence

20 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

25 <400> 11
Gly Tyr Asp Phe Thr His Tyr Gly Met Asn
1 5 10

30 <210> 12
<211> 14
<212> PRT
<213> Artificial Sequence

35 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

40 <400> 12
Tyr Pro Tyr Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val
1 5 10

45 <210> 13
<211> 15
<212> PRT
<213> Artificial Sequence

50 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

55 <400> 13
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
1 5 10 15

<210> 14

EP 2 894 167 A1

<211> 15
<212> PRT
<213> Artificial Sequence

5
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10
<400> 14
Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly Asp Arg
1 5 10 15

15
<210> 15
<211> 15
<212> PRT
<213> Artificial Sequence

20
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25
<400> 15
Ser Pro Ser Ser Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile
1 5 10 15

30
<210> 16
<211> 15
<212> PRT
<213> Artificial Sequence

35
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40
<400> 16
Ser Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser
1 5 10 15

45
<210> 17
<211> 15
<212> PRT
<213> Artificial Sequence

50
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

55
<400> 17
Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln
1 5 10 15

<210> 18
<211> 15
<212> PRT
<213> Artificial Sequence

EP 2 894 167 A1

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
5
<400> 18
Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser
1 5 10 15
10
<210> 19
<211> 15
<212> PRT
<213> Artificial Sequence
15
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
20
<400> 19
Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr Leu
1 5 10 15
25
<210> 20
<211> 15
<212> PRT
<213> Artificial Sequence
30
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
35
<400> 20
Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn Trp Tyr
1 5 10 15
40
<210> 21
<211> 15
<212> PRT
<213> Artificial Sequence
45
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
50
<400> 21
Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys
1 5 10 15
55
<210> 22
<211> 15
<212> PRT
<213> Artificial Sequence
55
<220>
<221> source

EP 2 894 167 A1

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 22

5 Asp Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys
1 5 10 15

<210> 23

10 <211> 15

<212> PRT

<213> Artificial Sequence

<220>

<221> source

15 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 23

20 Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys
1 5 10 15

<210> 24

<211> 15

25 <212> PRT

<213> Artificial Sequence

<220>

<221> source

30 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 24

35 Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile
1 5 10 15

<210> 25

<211> 15

40 <212> PRT

<213> Artificial Sequence

<220>

<221> source

45 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 25

50 Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile Tyr Phe Thr
1 5 10 15

<210> 26

55 <211> 15

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

EP 2 894 167 A1

<400> 26
Pro Gly Lys Ala Pro Lys Val Leu Ile Tyr Phe Thr Ser Ser Leu
1 5 10 15

5

<210> 27
<211> 15
<212> PRT
<213> Artificial Sequence

10

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

15

<400> 27
Ala Pro Lys Val Leu Ile Tyr Phe Thr Ser Ser Leu His Ser Gly
1 5 10 15

20

<210> 28
<211> 15
<212> PRT
<213> Artificial Sequence

25

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

30

<400> 28
Val Leu Ile Tyr Phe Thr Ser Ser Leu His Ser Gly Val Pro Ser
1 5 10 15

35

<210> 29
<211> 15
<212> PRT
<213> Artificial Sequence

40

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

45

<400> 29
Tyr Phe Thr Ser Ser Leu His Ser Gly Val Pro Ser Arg Phe Ser
1 5 10 15

50

<210> 30
<211> 15
<212> PRT
<213> Artificial Sequence

55

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 30
Ser Ser Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Gly
1 5 10 15

<210> 31
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 5
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 10
 <400> 31
 His Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr
 1 5 10 15
 15
 <210> 32
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 20
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 25
 <400> 32
 Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
 1 5 10 15
 30
 <210> 33
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 35
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 40
 <400> 33
 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile
 1 5 10 15
 45
 <210> 34
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 50
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 55
 <400> 34
 Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu
 1 5 10 15
 <210> 35

EP 2 894 167 A1

<211> 15
<212> PRT
<213> Artificial Sequence

5
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10
<400> 35
Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu
1 5 10 15

15
<210> 36
<211> 15
<212> PRT
<213> Artificial Sequence

20
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25
<400> 36
Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala
1 5 10 15

30
<210> 37
<211> 15
<212> PRT
<213> Artificial Sequence

35
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40
<400> 37
Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr
1 5 10 15

45
<210> 38
<211> 15
<212> PRT
<213> Artificial Sequence

50
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

55
<400> 38
Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
1 5 10 15

<210> 39
<211> 15
<212> PRT
<213> Artificial Sequence

EP 2 894 167 A1

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
5
<400> 39
Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Thr
1 5 10 15
10
<210> 40
<211> 15
<212> PRT
<213> Artificial Sequence
15
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
20
<400> 40
Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Thr Val Pro Trp
1 5 10 15
25
<210> 41
<211> 15
<212> PRT
<213> Artificial Sequence
30
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
35
<400> 41
Thr Tyr Tyr Cys Gln Gln Tyr Ser Thr Val Pro Trp Thr Phe Gly
1 5 10 15
40
<210> 42
<211> 15
<212> PRT
<213> Artificial Sequence
45
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
50
<400> 42
Cys Gln Gln Tyr Ser Thr Val Pro Trp Thr Phe Gly Gln Gly Thr
1 5 10 15
55
<210> 43
<211> 15
<212> PRT
<213> Artificial Sequence
<220>
<221> source

EP 2 894 167 A1

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 43

5 Tyr Ser Thr Val Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu
1 5 10 15

<210> 44

10 <211> 15

<212> PRT

<213> Artificial Sequence

<220>

<221> source

15 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 44

20 Val Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
1 5 10 15

<210> 45

<211> 15

<212> PRT

25 <213> Artificial Sequence

<220>

<221> source

30 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 45

35 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
1 5 10 15

<210> 46

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<221> source

40 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 46

45 Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu
1 5 10 15

<210> 47

50 <211> 15

<212> PRT

<213> Artificial Sequence

<220>

<221> source

55 <223> /note="Description of Artificial Sequence: Synthetic peptide"

EP 2 894 167 A1

5
5
10
15
20
25
30
35
40
45
50
55

<400> 47
Ser Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser
1 5 10 15

<210> 48
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 48
Gly Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala
1 5 10 15

<210> 49
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 49
Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr
1 5 10 15

<210> 50
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 50
Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr
1 5 10 15

<210> 51
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 51
Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Asn Tyr Gly
1 5 10 15

<210> 52
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 5
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 10
 <400> 52
 Cys Ala Ala Ser Gly Tyr Thr Phe Thr Asn Tyr Gly Met Asn Trp
 1 5 10 15
 15
 <210> 53
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 20
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 25
 <400> 53
 Ser Gly Tyr Thr Phe Thr Asn Tyr Gly Met Asn Trp Val Arg Gln
 1 5 10 15
 30
 <210> 54
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 35
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 40
 <400> 54
 Thr Phe Thr Asn Tyr Gly Met Asn Trp Val Arg Gln Ala Pro Gly
 1 5 10 15
 45
 <210> 55
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 50
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 55
 <400> 55
 Asn Tyr Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 1 5 10 15
 <210> 56

EP 2 894 167 A1

<211> 15
<212> PRT
<213> Artificial Sequence

5
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10
<400> 56
Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
1 5 10 15

15
<210> 57
<211> 15
<212> PRT
<213> Artificial Sequence

20
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25
<400> 57
Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Gly Trp Ile
1 5 10 15

30
<210> 58
<211> 15
<212> PRT
<213> Artificial Sequence

35
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40
<400> 58
Ala Pro Gly Lys Gly Leu Glu Trp Val Gly Trp Ile Asn Thr Tyr
1 5 10 15

45
<210> 59
<211> 15
<212> PRT
<213> Artificial Sequence

50
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

55
<400> 59
Lys Gly Leu Glu Trp Val Gly Trp Ile Asn Thr Tyr Thr Gly Glu
1 5 10 15

<210> 60
<211> 15
<212> PRT
<213> Artificial Sequence

EP 2 894 167 A1

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
5
<400> 60
Glu Trp Val Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr
1 5 10 15
10
<210> 61
<211> 15
<212> PRT
<213> Artificial Sequence
15
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
20
<400> 61
Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp
1 5 10 15
25
<210> 62
<211> 15
<212> PRT
<213> Artificial Sequence
30
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
35
<400> 62
Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Arg
1 5 10 15
40
<210> 63
<211> 15
<212> PRT
<213> Artificial Sequence
45
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
50
<400> 63
Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Arg Arg Phe Thr
1 5 10 15
55
<210> 64
<211> 15
<212> PRT
<213> Artificial Sequence
<220>
<221> source

EP 2 894 167 A1

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 64

5 Pro Thr Tyr Ala Ala Asp Phe Lys Arg Arg Phe Thr Phe Ser Leu
1 5 10 15

<210> 65

10 <211> 15

<212> PRT

<213> Artificial Sequence

<220>

<221> source

15 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 65

20 Ala Ala Asp Phe Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser
1 5 10 15

<210> 66

<211> 15

25 <212> PRT

<213> Artificial Sequence

<220>

<221> source

30 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 66

35 Phe Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr
1 5 10 15

<210> 67

<211> 15

40 <212> PRT

<213> Artificial Sequence

<220>

<221> source

45 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 67

50 Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr Leu
1 5 10 15

<210> 68

55 <211> 15

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

EP 2 894 167 A1

5 <400> 68
Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr Leu Gln Met Asn
1 5 10 15

10 <210> 69
<211> 15
<212> PRT
<213> Artificial Sequence

15 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

20 <400> 69
Asp Thr Ser Lys Ser Thr Ala Tyr Leu Gln Met Asn Ser Leu Arg
1 5 10 15

25 <210> 70
<211> 15
<212> PRT
<213> Artificial Sequence

30 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

35 <400> 70
Lys Ser Thr Ala Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
1 5 10 15

40 <210> 71
<211> 15
<212> PRT
<213> Artificial Sequence

45 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

50 <400> 71
Ala Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
1 5 10 15

55 <210> 72
<211> 15
<212> PRT
<213> Artificial Sequence

60 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

65 <400> 72
Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
1 5 10 15

<210> 73
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 5
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"
 10
 <400> 73
 Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Lys Tyr
 1 5 10 15
 15
 <210> 74
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 20
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"
 25
 <400> 74
 Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Lys Tyr Pro His Tyr
 1 5 10 15
 30
 <210> 75
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 35
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"
 40
 <400> 75
 Thr Ala Val Tyr Tyr Cys Ala Lys Tyr Pro His Tyr Tyr Gly Ser
 1 5 10 15
 45
 <210> 76
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 50
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"
 55
 <400> 76
 Tyr Tyr Cys Ala Lys Tyr Pro His Tyr Tyr Gly Ser Ser His Trp
 1 5 10 15
 <210> 77

EP 2 894 167 A1

<211> 15
<212> PRT
<213> Artificial Sequence

5 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10 <400> 77
Ala Lys Tyr Pro His Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp
1 5 10 15

15 <210> 78
<211> 15
<212> PRT
<213> Artificial Sequence

20 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25 <400> 78
Pro His Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val Trp Gly
1 5 10 15

30 <210> 79
<211> 15
<212> PRT
<213> Artificial Sequence

35 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40 <400> 79
Tyr Gly Ser Ser His Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr
1 5 10 15

45 <210> 80
<211> 15
<212> PRT
<213> Artificial Sequence

50 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

55 <400> 80
Ser His Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Leu Val Thr
1 5 10 15

<210> 81
<211> 15
<212> PRT
<213> Artificial Sequence

EP 2 894 167 A1

<220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"

5

<400> 81
 Tyr Phe Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 1 5 10 15

10

<210> 82
 <211> 123
 <212> PRT
 <213> Artificial Sequence

15

<220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic polypeptide"

20

<400> 82
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

25

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Phe Tyr
 20 25 30

30

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

35

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
 50 55 60

40

Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

45

Ala Lys Tyr Pro Asp Tyr Asp Gly Gly Ser His Trp Tyr Phe Asp Val
 100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

50

<210> 83
 <211> 123
 <212> PRT
 <213> Artificial Sequence

55

<220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic polypeptide"

EP 2 894 167 A1

<400> 83

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

5

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Phe Tyr
20 25 30

10

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

15

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
50 55 60

Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
65 70 75 80

20

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

25

Ala Lys Tyr Pro Pro Tyr Asp Gly Gly Ser His Trp Tyr Phe Asp Val
100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

30

<210> 84

<211> 123

<212> PRT

<213> Artificial Sequence

35

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

40

<400> 84

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

45

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Phe Tyr
20 25 30

50

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
50 55 60

55

Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
65 70 75 80

EP 2 894 167 A1

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

5 Ala Lys Tyr Pro Pro Tyr Glu Gly Ser Ser His Trp Tyr Phe Asp Val
100 105 110

10 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 85

<211> 123

<212> PRT

15 <213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic polypeptide"

20

<400> 85

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

25

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Phe Tyr
20 25 30

30

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

35

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
50 55 60

Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
65 70 75 80

40

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

45

Ala Lys Tyr Pro Glu Tyr Glu Gly Ser Ser His Trp Tyr Phe Asp Val
100 105 110

50

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 86

<211> 123

<212> PRT

55 <213> Artificial Sequence

<220>

EP 2 894 167 A1

<221> source
 <223> /note="Description of Artificial Sequence: Synthetic polypeptide"

5 <400> 86
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Phe Tyr
 10 20 25 30
 Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 15 35 40 45
 Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
 50 55 60
 Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
 20 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 25 85 90 95
 Ala Lys Tyr Pro Asp Tyr Glu Gly Ser Ser His Trp Tyr Phe Asp Val
 100 105 110
 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

35 <210> 87
 <211> 123
 <212> PRT
 <213> Artificial Sequence

40 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic polypeptide"

45 <400> 87
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Phe Tyr
 50 20 25 30
 Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
 55 50 55 60

EP 2 894 167 A1

Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
65 70 75 80

5 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

10 Ala Lys Tyr Pro Glu Tyr Asp Gly Gly Ser His Trp Tyr Phe Asp Val
100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

15 <210> 88
<211> 123
<212> PRT
<213> Artificial Sequence

20 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
polypeptide"

25 <400> 88
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

30 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Phe Tyr
20 25 30

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
50 55 60

40 Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
65 70 75 80

45 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Tyr Pro His Tyr Asp Gly Gly Ser His Trp Tyr Phe Asp Val
100 105 110

50 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

55 <210> 89
<211> 123
<212> PRT

EP 2 894 167 A1

<213> Artificial Sequence

<220>

<221> source

5 <223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 89

10 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Phe Tyr
20 25 30

15 Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

20 Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
50 55 60

25 Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

30 Ala Lys Tyr Pro Pro Tyr Asp Gly Ser Ser His Trp Tyr Phe Asp Val
100 105 110

35 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 90

<211> 123

40 <212> PRT

<213> Artificial Sequence

<220>

<221> source

45 <223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 90

50 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Phe Tyr
20 25 30

55 Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

EP 2 894 167 A1

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
50 55 60

5 Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
65 70 75 80

10 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Tyr Pro Asp Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
100 105 110

15 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

20 <210> 91
<211> 123
<212> PRT
<213> Artificial Sequence

25 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
polypeptide"

30 <400> 91
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

35 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Phe Tyr
20 25 30

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

40 Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
50 55 60

45 Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
65 70 75 80

50 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Tyr Pro His Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
100 105 110

55 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

EP 2 894 167 A1

<210> 92
 <211> 123
 <212> PRT
 <213> Artificial Sequence

5

<220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic polypeptide"

10

<400> 92
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
 20 25 30

20

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
 50 55 60

25

Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
 65 70 75 80

30

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Lys Tyr Pro Pro Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
 100 105 110

35

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

40

<210> 93
 <211> 123
 <212> PRT
 <213> Artificial Sequence

45

<220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic polypeptide"

50

<400> 93
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
 20 25 30

55

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val

EP 2 894 167 A1

35 40 45

5 Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
50 55 60

10 Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
65 70 75 80

15 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

20 Ala Lys Tyr Pro Glu Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
100 105 110

25 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 94
<211> 123
<212> PRT
<213> Artificial Sequence

30 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
polypeptide"

35 <400> 94
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

40 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Phe Tyr
20 25 30

45 Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

50 Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
50 55 60

55 Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
65 70 75 80

60 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

65 Ala Lys Tyr Pro His Tyr Tyr Gly Gly Ser His Trp Tyr Phe Asp Val
100 105 110

70 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser

EP 2 894 167 A1

115

120

5

<210> 95
 <211> 123
 <212> PRT
 <213> Artificial Sequence

10

<220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic polypeptide"

15

<400> 95
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

20

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
 20 25 30

25

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

30

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
 50 55 60

35

Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
 65 70 75 80

40

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

45

Ala Lys Tyr Pro Asp Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
 100 105 110

50

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

55

<210> 96
 <211> 123
 <212> PRT
 <213> Artificial Sequence

<220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic polypeptide"

<400> 96
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr Phe Tyr
 20 25 30

EP 2 894 167 A1

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

5 Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe
50 55 60

10 Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

15 Ala Lys Tyr Pro Pro Tyr Tyr Gly Gly Ser His Trp Tyr Phe Asp Val
100 105 110

20 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 97
<211> 15
25 <212> PRT
<213> Artificial Sequence

<220>
<221> source
30 <223> /note="Description of Artificial Sequence: Synthetic
peptide"

<400> 97
35 Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Arg
1 5 10 15

<210> 98
<211> 15
40 <212> PRT
<213> Artificial Sequence

<220>
<221> source
45 <223> /note="Description of Artificial Sequence: Synthetic
peptide"

<400> 98
Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Phe Asp Phe Lys Arg
1 5 10 15

50 <210> 99
<211> 15
<212> PRT
<213> Artificial Sequence

55 <220>
<221> source

EP 2 894 167 A1

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 99

5 Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Glu Asp Phe Lys Arg
1 5 10 15

<210> 100

<211> 15

10 <212> PRT

<213> Artificial Sequence

<220>

<221> source

15 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 100

20 Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Asp Asp Phe Lys Arg
1 5 10 15

<210> 101

<211> 15

25 <212> PRT

<213> Artificial Sequence

<220>

<221> source

30 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 101

35 Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Gly Phe Lys Arg
1 5 10 15

<210> 102

<211> 15

40 <212> PRT

<213> Artificial Sequence

<220>

<221> source

45 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 102

50 Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Leu Phe Lys Arg
1 5 10 15

<210> 103

<211> 15

55 <212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

EP 2 894 167 A1

5
<400> 103
Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Gln Phe Lys Arg
1 5 10 15

10
<210> 104
<211> 15
<212> PRT
<213> Artificial Sequence

15
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

20
<400> 104
Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Thr Phe Lys Arg
1 5 10 15

25
<210> 105
<211> 15
<212> PRT
<213> Artificial Sequence

30
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

35
<400> 105
Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Lys Phe Lys Arg
1 5 10 15

40
<210> 106
<211> 15
<212> PRT
<213> Artificial Sequence

45
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

50
<400> 106
Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Arg Phe Lys Arg
1 5 10 15

55
<210> 107
<211> 15
<212> PRT
<213> Artificial Sequence

55
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

55
<400> 107
Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Glu Phe Lys Arg
1 5 10 15

5 <210> 108
 <211> 15
 <212> PRT
 <213> Artificial Sequence

<220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"

10 <400> 108
 Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala His Phe Lys Arg
 1 5 10 15

15 <210> 109
 <211> 15
 <212> PRT
 <213> Artificial Sequence

20 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"

25 <400> 109
 Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Ser Arg
 1 5 10 15

30 <210> 110
 <211> 15
 <212> PRT
 <213> Artificial Sequence

35 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"

40 <400> 110
 Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Val Arg
 1 5 10 15

45 <210> 111
 <211> 15
 <212> PRT
 <213> Artificial Sequence

50 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"

55 <400> 111
 Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Gln Arg
 1 5 10 15

<210> 112

EP 2 894 167 A1

<211> 15
<212> PRT
<213> Artificial Sequence

5
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10
<400> 112
Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Val
1 5 10 15

15
<210> 113
<211> 15
<212> PRT
<213> Artificial Sequence

20
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25
<400> 113
Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Phe
1 5 10 15

30
<210> 114
<211> 15
<212> PRT
<213> Artificial Sequence

35
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40
<400> 114
Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys His
1 5 10 15

45
<210> 115
<211> 15
<212> PRT
<213> Artificial Sequence

50
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

55
<400> 115
Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Asn
1 5 10 15

<210> 116
<211> 15
<212> PRT
<213> Artificial Sequence

EP 2 894 167 A1

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
5
<400> 116
Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Ser
1 5 10 15
10
<210> 117
<211> 15
<212> PRT
<213> Artificial Sequence
15
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
20
<400> 117
Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Gln
1 5 10 15
25
<210> 118
<211> 15
<212> PRT
<213> Artificial Sequence
30
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
35
<400> 118
Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Lys
1 5 10 15
40
<210> 119
<211> 15
<212> PRT
<213> Artificial Sequence
45
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
50
<400> 119
Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Ile
1 5 10 15
55
<210> 120
<211> 15
<212> PRT
<213> Artificial Sequence
<220>
<221> source

EP 2 894 167 A1

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 120

5 Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Phe Asp Phe Lys Arg
1 5 10 15

<210> 121

10 <211> 15

<212> PRT

<213> Artificial Sequence

<220>

<221> source

15 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 121

20 Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Glu Asp Phe Lys Arg
1 5 10 15

<210> 122

<211> 15

25 <212> PRT

<213> Artificial Sequence

<220>

<221> source

30 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 122

35 Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Asp Asp Phe Lys Arg
1 5 10 15

<210> 123

<211> 15

40 <212> PRT

<213> Artificial Sequence

<220>

<221> source

45 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 123

50 Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Gly Phe Lys Arg
1 5 10 15

<210> 124

55 <211> 15

<212> PRT

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

EP 2 894 167 A1

5
5
10
15
20
25
30
35
40
45
50
55

<400> 124
Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Leu Phe Lys Arg
1 5 10 15

<210> 125
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 125
Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Gln Phe Lys Arg
1 5 10 15

<210> 126
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 126
Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Thr Phe Lys Arg
1 5 10 15

<210> 127
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 127
Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Lys Phe Lys Arg
1 5 10 15

<210> 128
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 128
Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Arg Phe Lys Arg
1 5 10 15

<210> 129
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 5
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 10
 <400> 129
 Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Glu Phe Lys Arg
 1 5 10 15
 15
 <210> 130
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 20
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 25
 <400> 130
 Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala His Phe Lys Arg
 1 5 10 15
 30
 <210> 131
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 35
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 40
 <400> 131
 Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Ser Arg
 1 5 10 15
 45
 <210> 132
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 50
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 55
 <400> 132
 Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Val Arg
 1 5 10 15
 <210> 133

EP 2 894 167 A1

<211> 15
<212> PRT
<213> Artificial Sequence

5
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10
<400> 133
Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Gln Arg
1 5 10 15

15
<210> 134
<211> 15
<212> PRT
<213> Artificial Sequence

20
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25
<400> 134
Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Val
1 5 10 15

30
<210> 135
<211> 15
<212> PRT
<213> Artificial Sequence

35
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40
<400> 135
Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Phe
1 5 10 15

45
<210> 136
<211> 15
<212> PRT
<213> Artificial Sequence

50
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

55
<400> 136
Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys His
1 5 10 15

<210> 137
<211> 15
<212> PRT
<213> Artificial Sequence

EP 2 894 167 A1

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
5
<400> 137
Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Asn
1 5 10 15
10
<210> 138
<211> 15
<212> PRT
<213> Artificial Sequence
15
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
20
<400> 138
Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Ser
1 5 10 15
25
<210> 139
<211> 15
<212> PRT
<213> Artificial Sequence
30
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
35
<400> 139
Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Gln
1 5 10 15
40
<210> 140
<211> 15
<212> PRT
<213> Artificial Sequence
45
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
50
<400> 140
Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Lys
1 5 10 15
55
<210> 141
<211> 15
<212> PRT
<213> Artificial Sequence
<220>
<221> source

EP 2 894 167 A1

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 141

5 Asn Thr Phe Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Ile
1 5 10 15

<210> 142

10 <211> 15

<212> PRT

<213> Artificial Sequence

<220>

<221> source

15 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 142

20 Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Lys Tyr Pro Asp Tyr
1 5 10 15

<210> 143

25 <211> 15

<212> PRT

<213> Artificial Sequence

<220>

<221> source

30 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 143

Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Lys Tyr Pro Glu Tyr
1 5 10 15

<210> 144

35 <211> 15

<212> PRT

<213> Artificial Sequence

<220>

<221> source

40 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 144

45 Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Lys Tyr Pro Pro Tyr
1 5 10 15

<210> 145

50 <211> 15

<212> PRT

<213> Artificial Sequence

<220>

<221> source

55 <223> /note="Description of Artificial Sequence: Synthetic peptide"

EP 2 894 167 A1

5
<400> 145
Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Lys Tyr Pro His Phe
1 5 10 15

10
<210> 146
<211> 15
<212> PRT
<213> Artificial Sequence

15
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

20
<400> 146
Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Lys Tyr Pro Asp Phe
1 5 10 15

25
<210> 147
<211> 15
<212> PRT
<213> Artificial Sequence

30
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

35
<400> 147
Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Lys Tyr Pro Glu Phe
1 5 10 15

40
<210> 148
<211> 15
<212> PRT
<213> Artificial Sequence

45
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

50
<400> 148
Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Lys Tyr Pro Pro Phe
1 5 10 15

55
<210> 149
<211> 15
<212> PRT
<213> Artificial Sequence

60
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

65
<400> 149
Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Phe Thr
1 5 10 15

<210> 150
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 5
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"
 10
 <400> 150
 Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile Tyr Asp Thr
 1 5 10 15
 15
 <210> 151
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 20
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"
 25
 <400> 151
 Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile Tyr Ser Thr
 1 5 10 15
 30
 <210> 152
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 35
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"
 40
 <400> 152
 Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile Tyr Gly Thr
 1 5 10 15
 45
 <210> 153
 <211> 15
 <212> PRT
 <213> Artificial Sequence
 50
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"
 55
 <400> 153
 Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile Tyr Ala Thr
 1 5 10 15
 <210> 154

EP 2 894 167 A1

<211> 15
<212> PRT
<213> Artificial Sequence

5
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10
<400> 154
Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile Tyr Phe Ala
1 5 10 15

15
<210> 155
<211> 15
<212> PRT
<213> Artificial Sequence

20
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25
<400> 155
Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile Tyr Asp Ala
1 5 10 15

30
<210> 156
<211> 15
<212> PRT
<213> Artificial Sequence

35
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40
<400> 156
Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile Tyr Ser Ala
1 5 10 15

45
<210> 157
<211> 15
<212> PRT
<213> Artificial Sequence

50
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

55
<400> 157
Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile Tyr Gly Ala
1 5 10 15

<210> 158
<211> 15
<212> PRT
<213> Artificial Sequence

EP 2 894 167 A1

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
5
<400> 158
Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile Tyr Ala Ala
1 5 10 15
10
<210> 159
<211> 15
<212> PRT
<213> Artificial Sequence
15
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
20
<400> 159
Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Asp Thr
1 5 10 15
25
<210> 160
<211> 15
<212> PRT
<213> Artificial Sequence
30
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
35
<400> 160
Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Ser Thr
1 5 10 15
40
<210> 161
<211> 15
<212> PRT
<213> Artificial Sequence
45
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
50
<400> 161
Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Gly Thr
1 5 10 15
55
<210> 162
<211> 15
<212> PRT
<213> Artificial Sequence
55
<220>
<221> source

EP 2 894 167 A1

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 162

5 Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Ala Thr
1 5 10 15

<210> 163

10 <211> 15

<212> PRT

<213> Artificial Sequence

<220>

<221> source

15 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 163

20 Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Phe Ala
1 5 10 15

<210> 164

<211> 15

25 <212> PRT

<213> Artificial Sequence

<220>

<221> source

30 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 164

Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Asp Ala
1 5 10 15

<210> 165

<211> 15

35 <212> PRT

<213> Artificial Sequence

<220>

<221> source

40 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 165

45 Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Ser Ala
1 5 10 15

<210> 166

50 <211> 15

<212> PRT

<213> Artificial Sequence

<220>

<221> source

55 <223> /note="Description of Artificial Sequence: Synthetic peptide"

EP 2 894 167 A1

5 <400> 166
Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Gly Ala
1 5 10 15

5 <210> 167
<211> 15
<212> PRT
<213> Artificial Sequence

10 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

15 <400> 167
Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Ala Ala
1 5 10 15

20 <210> 168
<211> 9
<212> PRT
<213> Artificial Sequence

25 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

30 <400> 168
Gly Tyr Asp Phe Thr Asn Tyr Gly Met
1 5

35 <210> 169
<211> 9
<212> PRT
<213> Artificial Sequence

40 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

45 <400> 169
Gly Tyr Thr Phe Thr Asn Tyr Gly Ile
1 5

50 <210> 170
<211> 9
<212> PRT
<213> Artificial Sequence

55 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

55 <400> 170
Gly Tyr Thr Phe Thr His Tyr Gly Met
1 5

<210> 171
 <211> 9
 <212> PRT
 <213> Artificial Sequence
 5
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 10
 <400> 171
 Ala Tyr Thr Phe Thr Asn Tyr Gly Met
 1 5
 15
 <210> 172
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 20
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 25
 <400> 172
 Gly Phe Thr Phe Ser Ser Tyr Ala Met Ser
 1 5 10
 30
 <210> 173
 <211> 9
 <212> PRT
 <213> Artificial Sequence
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 35
 <400> 173
 Gly Tyr Asp Phe Thr His Tyr Gly Ile
 1 5
 40
 <210> 174
 <211> 9
 <212> PRT
 <213> Artificial Sequence
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 50
 <400> 174
 Gly Tyr Thr Phe Thr Asn Tyr Gly Ala
 1 5
 55
 <210> 175

EP 2 894 167 A1

<211> 9
<212> PRT
<213> Artificial Sequence

5
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10
<400> 175
Gly Tyr Asp Phe Gly His Tyr Gly Ile
1 5

15
<210> 176
<211> 9
<212> PRT
<213> Artificial Sequence

20
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25
<400> 176
Gly Tyr Glu Phe Gln His Tyr Gly Ile
1 5

30
<210> 177
<211> 9
<212> PRT
<213> Artificial Sequence

35
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40
<400> 177
Gly Tyr Asp Phe Ser His Tyr Gly Ile
1 5

45
<210> 178
<211> 9
<212> PRT
<213> Artificial Sequence

50
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

55
<400> 178
Gly Tyr Glu Phe Ser His Tyr Gly Ile
1 5

<210> 179
<211> 9
<212> PRT
<213> Artificial Sequence

EP 2 894 167 A1

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
5
<400> 179
Gly Tyr Glu Phe Thr His Tyr Gly Ile
1 5
10
<210> 180
<211> 9
<212> PRT
<213> Artificial Sequence
15
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
20
<400> 180
Gly Tyr Thr Phe Thr His Tyr Gly Ile
1 5
25
<210> 181
<211> 9
<212> PRT
<213> Artificial Sequence
30
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
35
<400> 181
Gly Tyr Asp Phe Thr Asn Tyr Gly Ile
1 5
40
<210> 182
<211> 10
<212> PRT
<213> Artificial Sequence
45
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
50
<400> 182
Gly Tyr Thr Phe Thr Asn Tyr Gly Met Ala
1 5 10
55
<210> 183
<211> 9
<212> PRT
<213> Artificial Sequence
60
<220>
<221> source

EP 2 894 167 A1

<223> /note="Description of Artificial Sequence: Synthetic peptide"

5 <400> 183
Gly Tyr Thr Phe Thr Asn Tyr Ala Met
1 5

10 <210> 184
<211> 9
<212> PRT
<213> Artificial Sequence

15 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

20 <400> 184
Gly Tyr Thr Phe Thr Ala Tyr Gly Met
1 5

25 <210> 185
<211> 9
<212> PRT
<213> Artificial Sequence

30 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 185
Gly Tyr Thr Phe Ala Asn Tyr Gly Met
1 5

35 <210> 186
<211> 9
<212> PRT
<213> Artificial Sequence

40 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

45 <400> 186
Gly Tyr Thr Ala Thr Asn Tyr Gly Met
1 5

50 <210> 187
<211> 9
<212> PRT
<213> Artificial Sequence

55 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 187
Gly Tyr Ala Phe Thr Asn Tyr Gly Met
1 5

5

<210> 188
<211> 9
<212> PRT
<213> Artificial Sequence

10

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

15

<400> 188
Gly Ala Thr Phe Thr Asn Tyr Gly Met
1 5

20

<210> 189
<211> 9
<212> PRT
<213> Artificial Sequence

25

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

30

<400> 189
Gly Tyr Thr Phe Thr Asn Ala Gly Met
1 5

35

<210> 190
<211> 17
<212> PRT
<213> Artificial Sequence

40

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

45

<400> 190
Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Asp Glu Phe Thr
1 5 10 15

50

Arg

55

<210> 191
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

EP 2 894 167 A1

<400> 191
Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Val Asp Glu Phe Lys
1 5 10 15

5

Arg

10

<210> 192
<211> 17
<212> PRT
<213> Artificial Sequence

15

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

20

<400> 192
Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Glu Glu Phe Lys
1 5 10 15

25

Arg

30

<210> 193
<211> 17
<212> PRT
<213> Artificial Sequence

35

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40

<400> 193
Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Pro Glu Phe Lys
1 5 10 15

Arg

45

<210> 194
<211> 17
<212> PRT
<213> Artificial Sequence

50

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

55

<400> 194
Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Gln Glu Phe Lys
1 5 10 15

Arg

5 <210> 195
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
10 <221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

<400> 195
15 Trp Ile Asn Thr Tyr Thr Gly Glu Thr Thr Tyr Ala His Asp Phe Lys
1 5 10 15

Arg

20

<210> 196
<211> 17
<212> PRT
25 <213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"
30

<400> 196
Trp Ile Asn Thr Tyr Asn Gly Glu Thr Thr Tyr Ala Pro Asp Phe Lys
1 5 10 15

Arg

35

<210> 197
<211> 17
40 <212> PRT
<213> Artificial Sequence

<220>
<221> source
45 <223> /note="Description of Artificial Sequence: Synthetic
peptide"

<400> 197
Trp Ile Asn Thr Tyr Asn Gly Glu Pro Thr Tyr Ala Pro Asp Phe Lys
50 1 5 10 15

Arg

55 <210> 198
<211> 17

EP 2 894 167 A1

<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 198
Trp Ile Asn Thr Tyr Thr Gly Glu Ser Thr Tyr Ala Gln Asp Phe Lys
1 5 10 15

Arg

<210> 199
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 199
Trp Ile Asn Thr Tyr Thr Asn Asn Pro Thr Tyr Ala Gln Asp Phe Lys
1 5 10 15

Arg

<210> 200
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 200
Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Asp Asp Phe Lys
1 5 10 15

Glu

<210> 201
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

peptide"

<400> 201

5 Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Glu Glu Phe Thr
1 5 10 15

Arg

10

<210> 202

<211> 17

<212> PRT

<213> Artificial Sequence

15

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

20

<400> 202

Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala His Glu Phe Thr
1 5 10 15

25

Arg

30

<210> 203

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<221> source

35 <223> /note="Description of Artificial Sequence: Synthetic peptide"

40

<400> 203

Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Pro Asp Phe Thr
1 5 10 15

Arg

45

<210> 204

<211> 17

<212> PRT

<213> Artificial Sequence

50

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

55

<400> 204

Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala His Glu Phe Lys
1 5 10 15

Arg

5 <210> 205
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
10 <221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

<400> 205
15 Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Pro Glu Phe Lys
1 5 10 15

Arg

20

<210> 206
<211> 17
<212> PRT
25 <213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"
30

<400> 206
Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Gln Glu Phe Lys
1 5 10 15

Arg

35

<210> 207
<211> 17
<212> PRT
40 <213> Artificial Sequence

<220>
<221> source
45 <223> /note="Description of Artificial Sequence: Synthetic
peptide"

<400> 207
50 Trp Ile Asn Thr Tyr Thr Gly Glu Thr Thr Tyr Ala His Asp Phe Lys
1 5 10 15

Arg

55

<210> 208

EP 2 894 167 A1

<211> 17
<212> PRT
<213> Artificial Sequence

5

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10

<400> 208
Trp Ile Asn Thr Tyr Asn Gly Glu Pro Thr Tyr Ala Pro Asp Phe Lys
1 5 10 15

Arg

15

<210> 209
<211> 17
<212> PRT
<213> Artificial Sequence

20

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25

<400> 209
Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Pro Asp Phe Lys
1 5 10 15

30

Arg

<210> 210
<211> 17
<212> PRT
<213> Artificial Sequence

35

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40

<400> 210
Trp Ile Asn Thr Tyr Thr Gly Glu Ser Thr Tyr Ala Gln Asp Phe Lys
1 5 10 15

45

Arg

50

<210> 211
<211> 17
<212> PRT
<213> Artificial Sequence

55

<220>
<221> source

EP 2 894 167 A1

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 211

5 Trp Ile Asn Thr Asn Asn Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15

Arg

10

<210> 212

<211> 17

<212> PRT

15

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

20

<220>

<221> MOD_RES

<222> (5)..(5)

25

<223> Any amino acid

<400> 212

Trp Ile Asn Thr Xaa Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15

30

Arg

<210> 213

35

<211> 19

<212> PRT

<213> Artificial Sequence

<220>

<221> source

40

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 213

45

Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Lys Phe
1 5 10 15

Lys Asp Arg

50

<210> 214

<211> 17

<212> PRT

<213> Artificial Sequence

55

<220>

<221> source

EP 2 894 167 A1

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 214

5 Val Ile Ser Gly Asp Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

10

<210> 215

<211> 17

<212> PRT

15

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

20

<400> 215

Trp Ile Asn Thr Trp Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15

25

Arg

<210> 216

30

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<221> source

35

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 216

Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15

40

Ala

<210> 217

45

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<221> source

50

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 217

Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Ala
1 5 10 15

55

Arg

5 <210> 218
<211> 17
<212> PRT
<213> Artificial Sequence

10 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

15 <400> 218
Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Ala Lys
1 5 10 15

Arg

20 <210> 219
<211> 17
<212> PRT
<213> Artificial Sequence

25 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

30 <400> 219
Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Ala Phe Lys
1 5 10 15

35

Arg

40 <210> 220
<211> 17
<212> PRT
<213> Artificial Sequence

45 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

50 <400> 220
Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15

Arg

55

EP 2 894 167 A1

<210> 221
<211> 17
<212> PRT
<213> Artificial Sequence

5

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10

<400> 221
Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15

15

Arg

<210> 222
<211> 17
<212> PRT
<213> Artificial Sequence

20

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25

<400> 222
Trp Ile Asn Thr Tyr Thr Gly Glu Pro Ala Tyr Ala Ala Asp Phe Lys
1 5 10 15

30

Arg

<210> 223
<211> 17
<212> PRT
<213> Artificial Sequence

35

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40

<400> 223
Trp Ile Asn Thr Tyr Thr Gly Glu Ala Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15

45

Arg

50

<210> 224
<211> 17
<212> PRT
<213> Artificial Sequence

55

<220>

EP 2 894 167 A1

<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

5 <400> 224
Trp Ile Asn Thr Tyr Thr Gly Ala Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15

10 Arg

<210> 225
<211> 17
15 <212> PRT
<213> Artificial Sequence

<220>
<221> source
20 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 225
Trp Ile Asn Thr Trp Thr Gly Gln Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15

25 Arg

<210> 226
30 <211> 17
<212> PRT
<213> Artificial Sequence

<220>
35 <221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 226
40 Trp Ile Asn Thr Trp Thr Gly Thr Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15

Arg

45 <210> 227
<211> 17
<212> PRT
50 <213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

55 <400> 227
Trp Ile Asn Thr Tyr Thr Ala Glu Pro Thr Tyr Ala Ala Asp Phe Lys

EP 2 894 167 A1

1 5 10 15

Arg

5

<210> 228
<211> 17
<212> PRT
<213> Artificial Sequence

10

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

15

<400> 228
Trp Ile Asn Thr Tyr Ala Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15

20

Arg

25

<210> 229
<211> 17
<212> PRT
<213> Artificial Sequence

30

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

35

<400> 229
Trp Ile Asn Thr Trp Asp Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15

40

Arg

45

<210> 230
<211> 17
<212> PRT
<213> Artificial Sequence

50

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

55

<400> 230
Trp Ile Asn Thr Ala Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15

Arg

EP 2 894 167 A1

5
<210> 231
<211> 17
<212> PRT
<213> Artificial Sequence

10
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

15
<400> 231
Trp Ile Asn Thr Trp Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15

Arg

20
<210> 232
<211> 17
<212> PRT
<213> Artificial Sequence

25
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

30
<400> 232
Trp Ile Asn Ala Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15

Arg

35
<210> 233
<211> 17
<212> PRT
<213> Artificial Sequence

40
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

45
<400> 233
Trp Ile Ala Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15

Arg

50
<210> 234
<211> 17
<212> PRT
<213> Artificial Sequence

EP 2 894 167 A1

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
5
<400> 234
Trp Ala Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15
10 Arg
<210> 235
<211> 17
<212> PRT
<213> Artificial Sequence
<220>
<221> source
20 <223> /note="Description of Artificial Sequence: Synthetic peptide"
<400> 235
Ala Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys
1 5 10 15
25 Arg
<210> 236
<211> 17
<212> PRT
<213> Artificial Sequence
30
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
35
<400> 236
Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Ala Ala Ala Asp Phe Lys
1 5 10 15
40 Arg
<210> 237
<211> 14
<212> PRT
50 <213> Artificial Sequence
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
55
<400> 237

EP 2 894 167 A1

Tyr Pro Lys Tyr Tyr Gly Lys Ser His Trp Tyr Phe Asp Val
1 5 10

5 <210> 238
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
10 <221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 238
15 Tyr Pro Tyr Tyr Tyr Gly Arg Ser His Trp Tyr Phe Asp Val
1 5 10

<210> 239
20 <211> 17
<212> PRT
<213> Artificial Sequence

<220>
<221> source
25 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 239
30 Tyr Pro His Tyr Tyr Val Asn Glu Arg Lys Ser His Trp Tyr Phe Asp
1 5 10 15

Val

35 <210> 240
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
40 <221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 240
45 Tyr Pro Tyr Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val
1 5 10

50 <210> 241
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
55 <221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

EP 2 894 167 A1

5
10
15
20
25
30
35
40
45
50
55

<400> 241
Tyr Pro His Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val
1 5 10

<210> 242
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 242
Tyr Pro His Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Leu
1 5 10

<210> 243
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 243
Tyr Pro His Tyr Tyr Gly Arg Ser Gln Trp Tyr Phe Asp Val
1 5 10

<210> 244
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 244
Tyr Pro His Tyr Tyr Gly Gly Cys His Trp Tyr Phe Asp Val
1 5 10

<210> 245
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 245
Tyr Pro His Tyr Tyr Gly Thr Thr His Trp Tyr Phe Asp Val
1 5 10

<210> 246
 <211> 14
 <212> PRT
 <213> Artificial Sequence
 5
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"
 10
 <400> 246
 Asp Pro His Tyr Tyr Gly Ser Tyr His Trp Tyr Phe Asp Val
 1 5 10
 15
 <210> 247
 <211> 14
 <212> PRT
 <213> Artificial Sequence
 20
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"
 25
 <400> 247
 Tyr Pro His Tyr Tyr Gly Ser Tyr His Trp Tyr Phe Asp Val
 1 5 10
 30
 <210> 248
 <211> 14
 <212> PRT
 <213> Artificial Sequence
 35
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"
 40
 <400> 248
 Arg Pro His Tyr Tyr Gly Ala Ser His Trp Tyr Phe Asp Val
 1 5 10
 45
 <210> 249
 <211> 14
 <212> PRT
 <213> Artificial Sequence
 50
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"
 55
 <400> 249
 Tyr Pro His Tyr Tyr Gly Ala Ser His Trp Tyr Phe Asp Val
 1 5 10
 <210> 250

EP 2 894 167 A1

<211> 18
<212> PRT
<213> Artificial Sequence

5
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10
<400> 250
Tyr Pro His Tyr Tyr Ser Gly Glu Arg Glu Ser Ser His Trp Tyr Phe
1 5 10 15

Asp Val

15

<210> 251
<211> 17
<212> PRT
<213> Artificial Sequence

20
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25
<400> 251
Tyr Pro His Tyr Tyr Ala Arg Glu Gly Gly Ser His Trp Tyr Phe Asp
1 5 10 15

30
Val

35
<210> 252
<211> 17
<212> PRT
<213> Artificial Sequence

40
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

45
<400> 252
Tyr Pro His Tyr Tyr Ser Val Glu Gly Gly Ser His Trp Tyr Phe Asp
1 5 10 15

Val

50
<210> 253
<211> 14
<212> PRT
<213> Artificial Sequence

55
<220>
<221> source

EP 2 894 167 A1

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 253

5 Tyr Pro His Tyr Tyr Gly Gly Ser His Trp Tyr Phe Asp Val
1 5 10

<210> 254

10 <211> 17

<212> PRT

<213> Artificial Sequence

<220>

<221> source

15 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 254

20 Tyr Pro His Tyr Tyr Gln Ser Glu Gly Arg Ser His Trp Tyr Phe Asp
1 5 10 15

Val

25

<210> 255

<211> 18

<212> PRT

<213> Artificial Sequence

30

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

35

<400> 255

Tyr Pro His Tyr Tyr His Thr Arg Gly Gly Ser Ser His Trp Tyr Phe
1 5 10 15

Asp Val

40

<210> 256

<211> 17

45

<212> PRT

<213> Artificial Sequence

<220>

<221> source

50 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 256

55 Tyr Pro His Tyr Tyr Leu Thr Asp His Lys Ser His Trp Tyr Phe Asp
1 5 10 15

Val

EP 2 894 167 A1

5
<210> 257
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10
<400> 257
Tyr Pro His Tyr Tyr Arg Gly Asp Arg Lys Ser His Trp Tyr Phe Asp
1 5 10 15

15
Val

<210> 258
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25
<400> 258
Tyr Pro His Tyr Tyr Glu Lys Gln Arg Lys Ser His Trp Tyr Phe Asp
1 5 10 15

30
Val

<210> 259
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40
<400> 259
Tyr Pro His Tyr Tyr Ser Asp Glu Lys Lys Ser His Trp Tyr Phe Asp
1 5 10 15

45
Val

<210> 260
<211> 18
<212> PRT
<213> Artificial Sequence

50
Val

55
<210> 260
<211> 18
<212> PRT
<213> Artificial Sequence

EP 2 894 167 A1

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

5

<400> 260
Tyr Pro His Tyr Tyr Leu Lys Asp Gly Lys Lys Ser His Trp Tyr Phe
1 5 10 15

10

Asp Val

<210> 261
<211> 17
<212> PRT
<213> Artificial Sequence

15

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

20

<400> 261
Tyr Pro His Tyr Tyr Lys Glu Asp Lys Lys Ser His Trp Tyr Phe Asp
1 5 10 15

25

Val

30

<210> 262
<211> 17
<212> PRT
<213> Artificial Sequence

35

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40

<400> 262
Tyr Pro His Tyr Tyr Leu Arg Asp Lys Lys Ser His Trp Tyr Phe Asp
1 5 10 15

45

Val

<210> 263
<211> 17
<212> PRT
<213> Artificial Sequence

50

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

55

EP 2 894 167 A1

<400> 263

Tyr Pro His Tyr Tyr Leu Ser Asp Lys Lys Ser His Trp Tyr Phe Asp
1 5 10 15

5

Val

<210> 264

<211> 17

<212> PRT

<213> Artificial Sequence

10

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

15

<400> 264

Tyr Pro His Tyr Tyr Arg Arg Asp Lys Lys Ser His Trp Tyr Phe Asp
1 5 10 15

20

Val

25

<210> 265

<211> 17

<212> PRT

<213> Artificial Sequence

30

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

35

<400> 265

Tyr Pro His Tyr Tyr Ser His Gln Lys Arg Ser His Trp Tyr Phe Asp
1 5 10 15

40

Val

<210> 266

<211> 17

<212> PRT

<213> Artificial Sequence

45

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

50

<400> 266

Tyr Pro His Tyr Tyr Thr Tyr Asp Lys Ser Ser His Trp Tyr Phe Asp
1 5 10 15

55

Val

EP 2 894 167 A1

<210> 267
<211> 18
<212> PRT
<213> Artificial Sequence

5

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10

<400> 267
Tyr Pro His Tyr Tyr Arg Gly Gln Arg Lys Ser Ser His Trp Tyr Phe
1 5 10 15

15

Asp Val

<210> 268
<211> 18
<212> PRT
<213> Artificial Sequence

20

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25

<400> 268
Tyr Pro His Tyr Tyr Glu Lys Glu Arg Lys Ser Ser His Trp Tyr Phe
1 5 10 15

30

Asp Val

35

<210> 269
<211> 17
<212> PRT
<213> Artificial Sequence

40

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

45

<400> 269
Tyr Pro His Tyr Tyr Gln Asp Glu Lys Ser Ser His Trp Tyr Phe Asp
1 5 10 15

50

Val

55

<210> 270
<211> 18
<212> PRT

EP 2 894 167 A1

<213> Artificial Sequence

<220>

<221> source

5 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 270

Tyr Pro His Tyr Tyr Glu Arg Asp Gly Lys Ser Ser His Trp Tyr Phe
1 5 10 15

10

Asp Val

15

<210> 271

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<221> source

20 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 271

Tyr Pro His Tyr Tyr Arg Asn Glu Lys Ser Ser His Trp Tyr Phe Asp
1 5 10 15

25

Val

30

<210> 272

<211> 18

<212> PRT

35 <213> Artificial Sequence

<220>

<221> source

40 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 272

Tyr Pro His Tyr Tyr Gln Arg Asn Gly Lys Ser Ser His Trp Tyr Phe
1 5 10 15

40

45

Asp Val

<210> 273

50 <211> 17

<212> PRT

<213> Artificial Sequence

<220>

<221> source

55 <223> /note="Description of Artificial Sequence: Synthetic peptide"

EP 2 894 167 A1

<400> 273

Tyr Pro His Tyr Tyr Arg Thr Glu Lys Ser Ser His Trp Tyr Phe Asp
1 5 10 15

5

Val

10

<210> 274
<211> 17
<212> PRT
<213> Artificial Sequence

15

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

20

<400> 274
Tyr Pro His Tyr Tyr Val Asn Asp Lys Thr Ser His Trp Tyr Phe Asp
1 5 10 15

25

Val

30

<210> 275
<211> 18
<212> PRT
<213> Artificial Sequence

35

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

40

<400> 275
Tyr Pro His Tyr Tyr Leu Lys Asp Gly Met Ser Ser His Trp Tyr Phe
1 5 10 15

Asp Val

45

<210> 276
<211> 14
<212> PRT
<213> Artificial Sequence

50

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic
peptide"

55

<400> 276
Tyr Pro His Tyr Tyr Gly Asn Ser His Trp Tyr Phe Asp Val
1 5 10

EP 2 894 167 A1

5
<210> 277
<211> 17
<212> PRT
<213> Artificial Sequence

10
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

15
<400> 277
Tyr Pro His Tyr Tyr Pro Ser Pro Arg Gly Ser His Trp Tyr Phe Asp
1 5 10 15
Val

20
<210> 278
<211> 18
<212> PRT
<213> Artificial Sequence

25
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

30
<400> 278
Tyr Pro His Tyr Tyr Gln Asn Glu Gly Pro Ser Ser His Trp Tyr Phe
1 5 10 15
Asp Val

35
<210> 279
<211> 14
<212> PRT
<213> Artificial Sequence

40
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

45
<400> 279
Tyr Pro Tyr Tyr Tyr Gly Arg Ser His Trp Tyr Phe Asp Val
1 5 10

50
<210> 280
<211> 14
<212> PRT
<213> Artificial Sequence

55
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

EP 2 894 167 A1

<400> 280
Tyr Pro His Tyr Tyr Gly Arg Ser His Trp Tyr Phe Asp Val
1 5 10

5

<210> 281
<211> 17
<212> PRT
<213> Artificial Sequence

10

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

15

<400> 281
Tyr Pro His Tyr Tyr Arg Asp Glu Arg Glu Ser His Trp Tyr Phe Asp
1 5 10 15

Val

20

<210> 282
<211> 17
<212> PRT
<213> Artificial Sequence

25

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

30

<400> 282
Tyr Pro His Tyr Tyr Ser His Glu Arg Lys Ser His Trp Tyr Phe Asp
1 5 10 15

35

Val

<210> 283
<211> 17
<212> PRT
<213> Artificial Sequence

40

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

45

<400> 283
Tyr Pro His Tyr Tyr Leu Lys Asp Arg Lys Ser His Trp Tyr Phe Asp
1 5 10 15

50

Val

55

<210> 284

EP 2 894 167 A1

<211> 17
<212> PRT
<213> Artificial Sequence

5
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10
<400> 284
Tyr Pro His Tyr Tyr Leu Asn Glu Arg Lys Ser His Trp Tyr Phe Asp
1 5 10 15

Val

15

<210> 285
<211> 17
<212> PRT
<213> Artificial Sequence

20
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25
<400> 285
Tyr Pro His Tyr Tyr Ser Asn Glu Arg Lys Ser His Trp Tyr Phe Asp
1 5 10 15

30
Val

35
<210> 286
<211> 17
<212> PRT
<213> Artificial Sequence

40
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

45
<400> 286
Tyr Pro His Tyr Tyr Val Asn Glu Arg Lys Ser His Trp Tyr Phe Asp
1 5 10 15

Val

50

<210> 287
<211> 17
<212> PRT
<213> Artificial Sequence

55
<220>
<221> source

EP 2 894 167 A1

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 287

5 Tyr Pro His Tyr Tyr Val Thr Asp Arg Lys Ser His Trp Tyr Phe Asp
1 5 10 15

Val

10

<210> 288

<211> 17

<212> PRT

15

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

20

<400> 288

Tyr Pro His Tyr Tyr Gly Asn His Arg Ser Ser His Trp Tyr Phe Asp
1 5 10 15

25

Val

<210> 289

30

<211> 18

<212> PRT

<213> Artificial Sequence

<220>

<221> source

35

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 289

40

Tyr Pro His Tyr Tyr Leu Lys Asp Gly Arg Ser Ser His Trp Tyr Phe
1 5 10 15

Asp Val

45

<210> 290

<211> 18

<212> PRT

<213> Artificial Sequence

50

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

55

<400> 290

Tyr Pro His Tyr Tyr Leu Lys Asp Arg Arg Ser Ser His Trp Tyr Phe
1 5 10 15

Asp Val

5 <210> 291
 <211> 17
 <212> PRT
 <213> Artificial Sequence

10 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"

15 <400> 291
 Tyr Pro His Tyr Tyr Gln Arg Asp Arg Ser Ser His Trp Tyr Phe Asp
 1 5 10 15

Val

20

25 <210> 292
 <211> 17
 <212> PRT
 <213> Artificial Sequence

30 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"

35 <400> 292
 Tyr Pro His Tyr Tyr Arg Asp Glu Arg Ser Ser His Trp Tyr Phe Asp
 1 5 10 15

Val

40 <210> 293
 <211> 18
 <212> PRT
 <213> Artificial Sequence

45 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"

50 <400> 293
 Tyr Pro His Tyr Tyr Glu Arg Asp Gly Arg Ser Ser His Trp Tyr Phe
 1 5 10 15

Asp Val

55

EP 2 894 167 A1

<210> 294
<211> 18
<212> PRT
<213> Artificial Sequence

5

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10

<400> 294
Tyr Pro His Tyr Tyr Leu Arg Asp Gly Arg Ser Ser His Trp Tyr Phe
1 5 10 15

15

Asp Val

<210> 295
<211> 17
<212> PRT
<213> Artificial Sequence

20

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25

<400> 295
Tyr Pro His Tyr Tyr Arg Asn Glu Arg Ser Ser His Trp Tyr Phe Asp
1 5 10 15

30

Val

<210> 296
<211> 17
<212> PRT
<213> Artificial Sequence

35

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40

<400> 296
Tyr Pro His Tyr Tyr Ser His Glu Arg Val Ser His Trp Tyr Phe Asp
1 5 10 15

45

Val

50

<210> 297
<211> 18
<212> PRT
<213> Artificial Sequence

55

<220>
<221> source

EP 2 894 167 A1

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 297

5 Tyr Pro His Tyr Tyr Gln Lys Gln Ser Lys Ser Ser His Trp Tyr Phe
1 5 10 15

Asp Val

10

<210> 298

<211> 14

<212> PRT

15

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

20

<400> 298

Tyr Pro Tyr Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val
1 5 10

25

<210> 299

<211> 14

<212> PRT

<213> Artificial Sequence

30

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

35

<400> 299

Tyr Pro His Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val
1 5 10

40

<210> 300

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<221> source

45

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 300

50

Tyr Pro His Tyr Tyr Val Glu Glu Thr Glu Ser His Trp Tyr Phe Asp
1 5 10 15

Val

55

<210> 301

<211> 17

EP 2 894 167 A1

<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 301
Tyr Pro His Tyr Tyr Val Gly Glu Thr Ser Ser His Trp Tyr Phe Asp
1 5 10 15

Val

<210> 302
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<220>
<221> MOD_RES
<222> (3)..(3)
<223> Any amino acid

<220>
<221> MOD_RES
<222> (7)..(7)
<223> Any amino acid

<400> 302
Tyr Pro Xaa Tyr Tyr Gly Xaa Ser His Trp Tyr Phe Asp Val
1 5 10

<210> 303
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 303
Tyr Pro His Tyr Tyr Leu Ala Asp Arg Ser Ser His Trp Tyr Phe Asp
1 5 10 15

Val

<210> 304

EP 2 894 167 A1

<211> 17
<212> PRT
<213> Artificial Sequence

5
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10
<400> 304
Tyr Pro His Tyr Tyr Val Gly Glu Gln Ser Ser His Trp Tyr Phe Asp
1 5 10 15

Val

15

<210> 305
<211> 17
<212> PRT
<213> Artificial Sequence

20
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25
<400> 305
Tyr Pro His Tyr Tyr Val Gly Glu Thr Ser Ser His Trp Tyr Phe Asp
1 5 10 15

30
Val

35
<210> 306
<211> 17
<212> PRT
<213> Artificial Sequence

40
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

45
<400> 306
Tyr Pro His Tyr Tyr Leu Lys Asp Lys Ser Ser His Trp Tyr Phe Asp
1 5 10 15

Val

50
<210> 307
<211> 17
<212> PRT
<213> Artificial Sequence

55
<220>
<221> source

EP 2 894 167 A1

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 307

5 Tyr Pro His Tyr Tyr Leu Lys Asp Arg Ser Ser His Trp Tyr Phe Asp
1 5 10 15

Val

10

<210> 308

<211> 17

<212> PRT

15 <213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

20

<400> 308

25 Tyr Pro His Tyr Tyr Ala Lys Glu Arg Ser Ser His Trp Tyr Phe Asp
1 5 10 15

25

Val

<210> 309

30

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

35

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 309

40 Tyr Pro His Tyr Tyr Leu Gly Ser Ser His Trp Tyr Phe Asp Val
1 5 10 15

<210> 310

45

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<221> source

50

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 310

55 Tyr Pro His Tyr Tyr Lys Asn Asp Lys Ser Ser His Trp Tyr Phe Asp
1 5 10 15

55

Val

EP 2 894 167 A1

5
<210> 311
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10
<400> 311
Tyr Pro His Tyr Tyr Leu Asn Asp Lys Ser Ser His Trp Tyr Phe Asp
1 5 10 15

15
Val

<210> 312
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25
<400> 312
Tyr Pro His Tyr Tyr Val Asn Glu Arg Ser Ser His Trp Tyr Phe Asp
1 5 10 15

30
Val

35
<210> 313
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40
<400> 313
Tyr Pro His Tyr Tyr Ala Arg Asp Arg Ser Ser His Trp Tyr Phe Asp
1 5 10 15

45
Val

50
<210> 314
<211> 17
<212> PRT
<213> Artificial Sequence

55

EP 2 894 167 A1

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
5
<400> 314
Tyr Pro His Tyr Tyr Tyr Arg Asp Arg Ser Ser His Trp Tyr Phe Asp
1 5 10 15
10 Val

<210> 315
15 <211> 14
<212> PRT
<213> Artificial Sequence

<220>
20 <221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 315
25 Tyr Pro Tyr Tyr Asn Gly Ser Ser His Trp Tyr Phe Asp Val
1 5 10

<210> 316
<211> 14
<212> PRT
30 <213> Artificial Sequence

<220>
<221> source
35 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 316
Pro Pro Tyr Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
1 5 10
40

<210> 317
<211> 14
<212> PRT
45 <213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

50 <400> 317
Tyr Pro Tyr Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
1 5 10

55 <210> 318
<211> 13
<212> PRT

EP 2 894 167 A1

<213> Artificial Sequence

<220>
<221> source
5 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 318
Tyr His Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
1 5 10

10

<210> 319
<211> 14
<212> PRT
15 <213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
20

<400> 319
Tyr Tyr His Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
1 5 10

25

<210> 320
<211> 14
<212> PRT
<213> Artificial Sequence

30

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

35

<400> 320
Thr Pro His Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
1 5 10

40

<210> 321
<211> 14
<212> PRT
<213> Artificial Sequence

45

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

50

<400> 321
Tyr Pro Tyr Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
1 5 10

55

<210> 322
<211> 14
<212> PRT
<213> Artificial Sequence

<220>

EP 2 894 167 A1

<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

5 <400> 322
Tyr Pro His Tyr Tyr Gly Arg Ser His Trp Tyr Phe Asp Val
1 5 10

10 <210> 323
<211> 14
<212> PRT
<213> Artificial Sequence

15 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

20 <400> 323
Tyr Pro His Tyr Tyr Gly Arg Ser His Trp Tyr Phe Asp Val
1 5 10

25 <210> 324
<211> 17
<212> PRT
<213> Artificial Sequence

30 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

35 <400> 324
Tyr Pro His Tyr Tyr Val Asn Glu Arg Lys Ser His Trp Tyr Phe Asp
1 5 10 15

Val

40 <210> 325
<211> 14
<212> PRT
<213> Artificial Sequence

45 <220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

50 <400> 325
Tyr Pro Tyr Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val
1 5 10

55 <210> 326
<211> 14
<212> PRT
<213> Artificial Sequence

EP 2 894 167 A1

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
5
<400> 326
Tyr Pro Tyr Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val
1 5 10
10
<210> 327
<211> 14
<212> PRT
<213> Artificial Sequence
15
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
<400> 327
20 Tyr Pro His Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Ala
1 5 10
25
<210> 328
<211> 14
<212> PRT
<213> Artificial Sequence
30
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
<400> 328
Tyr Pro His Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Tyr
1 5 10
35
<210> 329
<211> 14
<212> PRT
<213> Artificial Sequence
40
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
45
<400> 329
Tyr Pro His Tyr Tyr Gly Ser Ser His Trp Tyr Phe Ala Val
1 5 10
50
<210> 330
<211> 14
<212> PRT
<213> Artificial Sequence
55
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic

peptide"

<400> 330

Tyr Pro His Tyr Tyr Gly Ser Ser His Trp Tyr Ala Asp Val
 1 5 10

5

<210> 331

<211> 14

<212> PRT

<213> Artificial Sequence

10

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

15

<400> 331

Tyr Pro His Tyr Tyr Gly Ser Ser His Trp Ala Phe Asp Val
 1 5 10

20

<210> 332

<211> 14

<212> PRT

<213> Artificial Sequence

25

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

30

<400> 332

Tyr Pro His Tyr Tyr Gly Ser Ser His Ala Tyr Phe Asp Val
 1 5 10

35

<210> 333

<211> 14

<212> PRT

<213> Artificial Sequence

40

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

45

<400> 333

Tyr Pro His Tyr Tyr Gly Ser Ser Ala Trp Tyr Phe Asp Val
 1 5 10

50

<210> 334

<211> 14

<212> PRT

<213> Artificial Sequence

55

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 334

EP 2 894 167 A1

Tyr Pro His Tyr Tyr Gly Ser Ala His Trp Tyr Phe Asp Val
1 5 10

5

<210> 335
<211> 14
<212> PRT
<213> Artificial Sequence

10

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

15

<400> 335
Tyr Pro Tyr Tyr Arg Gln Lys Gly His Trp Tyr Phe Asp Val
1 5 10

20

<210> 336
<211> 14
<212> PRT
<213> Artificial Sequence

25

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

30

<400> 336
Tyr Pro Tyr Tyr Arg Gln Arg Gly His Trp Tyr Phe Asp Val
1 5 10

35

<210> 337
<211> 14
<212> PRT
<213> Artificial Sequence

40

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

45

<400> 337
Tyr Pro Tyr Tyr Arg Asp Ala Ser His Trp Tyr Phe Asp Val
1 5 10

50

<210> 338
<211> 14
<212> PRT
<213> Artificial Sequence

55

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 338
Tyr Pro Tyr Tyr Arg Asn Ala Ser His Trp Tyr Phe Asp Val
1 5 10

<210> 339
 <211> 14
 <212> PRT
 <213> Artificial Sequence
 5
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"
 10
 <400> 339
 Tyr Pro Tyr Tyr Tyr Gly Ala Ser His Trp Tyr Phe Asp Val
 1 5 10
 15
 <210> 340
 <211> 14
 <212> PRT
 <213> Artificial Sequence
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"
 20
 <400> 340
 Tyr Pro His Tyr Tyr Gly Ala Ser His Trp Tyr Phe Asp Val
 1 5 10
 25
 <210> 341
 <211> 14
 <212> PRT
 <213> Artificial Sequence
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"
 30
 <400> 341
 Tyr Pro Tyr Tyr Arg Gly Glu Ser His Trp Tyr Phe Asp Val
 1 5 10
 35
 <210> 342
 <211> 14
 <212> PRT
 <213> Artificial Sequence
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"
 40
 <400> 342
 Tyr Pro Tyr Tyr Arg Lys Gly Ser His Trp Tyr Phe Asp Val
 1 5 10
 45
 <210> 343
 <211> 14
 50
 55

EP 2 894 167 A1

<212> PRT
<213> Artificial Sequence

5
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10
<400> 343
Tyr Pro Tyr Tyr Arg Ser Gly Ser His Trp Tyr Phe Asp Val
1 5 10

15
<210> 344
<211> 14
<212> PRT
<213> Artificial Sequence

20
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25
<400> 344
Tyr Pro Tyr Tyr Lys Gly Gly Ser His Trp Tyr Phe Asp Val
1 5 10

30
<210> 345
<211> 14
<212> PRT
<213> Artificial Sequence

35
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40
<400> 345
Tyr Pro Tyr Tyr Ile Ala Lys Ser His Trp Tyr Phe Asp Val
1 5 10

45
<210> 346
<211> 14
<212> PRT
<213> Artificial Sequence

50
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

55
<400> 346
Tyr Pro Tyr Tyr Ile Asn Lys Ser His Trp Tyr Phe Asp Val
1 5 10

<210> 347
<211> 14
<212> PRT
<213> Artificial Sequence

EP 2 894 167 A1

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
5
<400> 347
Tyr Pro Tyr Tyr Arg Asp Asn Ser His Trp Tyr Phe Asp Val
1 5 10
10
<210> 348
<211> 14
<212> PRT
<213> Artificial Sequence
15
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
<400> 348
20 Tyr Pro Tyr Tyr Arg Gln Asn Ser His Trp Tyr Phe Asp Val
1 5 10
25
<210> 349
<211> 14
<212> PRT
<213> Artificial Sequence
30
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
<400> 349
Tyr Pro Tyr Tyr Tyr Asn Gln Ser His Trp Tyr Phe Asp Val
1 5 10
35
40
<210> 350
<211> 14
<212> PRT
<213> Artificial Sequence
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
45
<400> 350
Tyr Pro Tyr Tyr Ile Glu Arg Ser His Trp Tyr Phe Asp Val
1 5 10
50
<210> 351
<211> 14
<212> PRT
<213> Artificial Sequence
55
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic

peptide"

<400> 351

Tyr Pro Tyr Tyr Thr Asn Arg Ser His Trp Tyr Phe Asp Val
 1 5 10

5

<210> 352

<211> 14

<212> PRT

<213> Artificial Sequence

10

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

15

<400> 352

Tyr Pro Tyr Tyr Thr Thr Arg Ser His Trp Tyr Phe Asp Val
 1 5 10

20

<210> 353

<211> 14

<212> PRT

<213> Artificial Sequence

25

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

30

<400> 353

Tyr Pro Tyr Tyr Thr Gly Arg Ser His Trp Tyr Phe Asp Val
 1 5 10

35

<210> 354

<211> 14

<212> PRT

<213> Artificial Sequence

40

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

45

<400> 354

Tyr Pro Tyr Tyr Lys Asn Thr Ser His Trp Tyr Phe Asp Val
 1 5 10

50

<210> 355

<211> 14

<212> PRT

<213> Artificial Sequence

55

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 355

EP 2 894 167 A1

Tyr Pro Tyr Tyr Arg Asn Thr Ser His Trp Tyr Phe Asp Val
1 5 10

5

<210> 356
<211> 14
<212> PRT
<213> Artificial Sequence

10

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

15

<400> 356
Tyr Pro Tyr Tyr Thr Asn Thr Ser His Trp Tyr Phe Asp Val
1 5 10

20

<210> 357
<211> 14
<212> PRT
<213> Artificial Sequence

25

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

30

<400> 357
Tyr Pro Tyr Tyr Arg Gly Thr Ser His Trp Tyr Phe Asp Val
1 5 10

35

<210> 358
<211> 14
<212> PRT
<213> Artificial Sequence

40

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

45

<400> 358
Tyr Pro Tyr Tyr Trp Gly Thr Ser His Trp Tyr Phe Asp Val
1 5 10

50

<210> 359
<211> 14
<212> PRT
<213> Artificial Sequence

55

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 359
Tyr Pro Tyr Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val
1 5 10

EP 2 894 167 A1

5
<210> 360
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10
<400> 360
Tyr Pro His Tyr Tyr Ala Ser Ser His Trp Tyr Phe Asp Val
1 5 10

15
<210> 361
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

20
<400> 361
Tyr Pro Tyr Tyr Lys Glu Ser Ser His Trp Tyr Phe Asp Val
1 5 10

25
<210> 362
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

30
<400> 362
Tyr Pro Tyr Tyr Arg Asn Ser Ser His Trp Tyr Phe Asp Val
1 5 10

35
<210> 363
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40
<400> 363
Tyr Pro Tyr Tyr Arg Gln Ser Ser His Trp Tyr Phe Asp Val
1 5 10

45
<210> 364
<211> 14

50
<210> 364
<211> 14

55

EP 2 894 167 A1

<212> PRT
<213> Artificial Sequence

5
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10
<400> 364
Tyr Pro His Tyr Ala Gly Ser Ser His Trp Tyr Phe Asp Val
1 5 10

15
<210> 365
<211> 14
<212> PRT
<213> Artificial Sequence

20
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25
<400> 365
Tyr Pro Tyr Tyr Glu Gly Ser Ser His Trp Tyr Phe Asp Val
1 5 10

30
<210> 366
<211> 14
<212> PRT
<213> Artificial Sequence

35
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40
<400> 366
Tyr Pro Tyr Tyr Thr Gly Ser Ser His Trp Tyr Phe Asp Val
1 5 10

45
<210> 367
<211> 14
<212> PRT
<213> Artificial Sequence

50
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

55
<400> 367
Tyr Pro His Ala Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
1 5 10

<210> 368
<211> 14
<212> PRT
<213> Artificial Sequence

EP 2 894 167 A1

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
5
<400> 368
Tyr Pro Ala Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
1 5 10
10
<210> 369
<211> 14
<212> PRT
<213> Artificial Sequence
15
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
<400> 369
20 Tyr Pro Tyr Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
1 5 10
25
<210> 370
<211> 14
<212> PRT
<213> Artificial Sequence
30
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
<400> 370
Ala Ala His Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
1 5 10
35
<210> 371
<211> 14
<212> PRT
<213> Artificial Sequence
40
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
45
<400> 371
Ala Pro His Tyr Tyr Gly Ser Ser His Trp Tyr Phe Asp Val
1 5 10
50
<210> 372
<211> 11
<212> PRT
<213> Artificial Sequence
55
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic

peptide"

<400> 372

Ala Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn
1 5 10

5

<210> 373

<211> 11

<212> PRT

<213> Artificial Sequence

10

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

15

<400> 373

Ser Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn
1 5 10

20

<210> 374

<211> 11

<212> PRT

<213> Artificial Sequence

25

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

30

<400> 374

Ser Ala Ala Gln Asp Ile Ser Asn Tyr Leu Asn
1 5 10

35

<210> 375

<211> 11

<212> PRT

<213> Artificial Sequence

40

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

45

<400> 375

Ser Ala Ser Ala Asp Ile Ser Asn Tyr Leu Asn
1 5 10

50

<210> 376

<211> 11

<212> PRT

<213> Artificial Sequence

55

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 376

EP 2 894 167 A1

Ser Ala Ser Gln Ala Ile Ser Asn Tyr Leu Asn
1 5 10

5

<210> 377
<211> 11
<212> PRT
<213> Artificial Sequence

10

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

15

<400> 377
Ser Ala Ser Gln Asp Ala Ser Asn Tyr Leu Asn
1 5 10

20

<210> 378
<211> 11
<212> PRT
<213> Artificial Sequence

25

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

30

<400> 378
Ser Ala Ser Gln Asp Ile Ala Asn Tyr Leu Asn
1 5 10

35

<210> 379
<211> 11
<212> PRT
<213> Artificial Sequence

40

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

45

<400> 379
Ser Ala Ser Gln Asp Ile Ser Ala Tyr Leu Asn
1 5 10

50

<210> 380
<211> 11
<212> PRT
<213> Artificial Sequence

55

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 380
Ser Ala Ser Gln Asp Ile Ser Asn Ala Leu Asn
1 5 10

5
<210> 381
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10
<400> 381
Ser Ala Ser Gln Asp Ile Ser Asn Tyr Ala Asn
1 5 10

15
<210> 382
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

20
<400> 382
Ser Ala Ser Gln Asp Ile Ser Asn Tyr Leu Ala
1 5 10

25
<210> 383
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

30
<400> 383
Arg Ala Asn Glu Gln Leu Ser Asn Tyr Leu Asn
1 5 10

35
<210> 384
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40
<400> 384
Arg Ala Asn Glu Gln Ile Ser Asn Tyr Leu Asn
1 5 10

45
<210> 385
<211> 11

50
<210> 385
<211> 11

55
<210> 385
<211> 11

<212> PRT
 <213> Artificial Sequence

 <220>
 5 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"

 <400> 385
 10 Arg Ala Ser Gln Ser Ile Ser Asn Tyr Leu Ala
 1 5 10

 <210> 386
 <211> 7
 15 <212> PRT
 <213> Artificial Sequence

 <220>
 <221> source
 20 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"

 <400> 386
 Ala Thr Ser Ser Leu His Ser
 1 5
 25

 <210> 387
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 30

 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 35

 <400> 387
 Phe Ala Ser Ser Leu His Ser
 1 5

 40

 <210> 388
 <211> 7
 <212> PRT
 <213> Artificial Sequence

 45

 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"

 <400> 388
 50 Phe Thr Ala Ser Leu His Ser
 1 5

 <210> 389
 <211> 7
 55 <212> PRT
 <213> Artificial Sequence

EP 2 894 167 A1

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
5
<400> 389
Phe Thr Ser Ala Leu His Ser
1 5

10
<210> 390
<211> 7
<212> PRT
<213> Artificial Sequence

15
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
20
<400> 390
Phe Thr Ser Ser Ala His Ser
1 5

25
<210> 391
<211> 7
<212> PRT
<213> Artificial Sequence

30
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
35
<400> 391
Phe Thr Ser Ser Leu Ala Ser
1 5

40
<210> 392
<211> 7
<212> PRT
<213> Artificial Sequence

45
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"
50
<400> 392
Phe Thr Ser Ser Leu His Ala
1 5

55
<210> 393
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic

peptide"

<400> 393

Ala Ala Ser Ser Leu Glu Ser
1 5

5

<210> 394

<211> 9

<212> PRT

<213> Artificial Sequence

10

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic
peptide"

15

<400> 394

Gln Gln Tyr Ser Thr Val Phe Trp Thr
1 5

20

<210> 395

<211> 9

<212> PRT

<213> Artificial Sequence

25

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic
peptide"

30

<400> 395

Gln Gln Tyr Asn Ser Thr Pro Trp Thr
1 5

35

<210> 396

<211> 9

<212> PRT

<213> Artificial Sequence

40

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic
peptide"

45

<400> 396

Gln Gln Tyr Tyr Ser Thr Pro Trp Thr
1 5

50

<210> 397

<211> 9

<212> PRT

<213> Artificial Sequence

55

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic
peptide"

<400> 397

EP 2 894 167 A1

Gln Gln Ser Tyr Ser Thr Pro Trp Thr
1 5

5

<210> 398
<211> 9
<212> PRT
<213> Artificial Sequence

10

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

15

<400> 398
Gln Gln Tyr Ser Ala Val Pro Trp Thr
1 5

20

<210> 399
<211> 9
<212> PRT
<213> Artificial Sequence

25

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

30

<400> 399
Gln Gln Tyr Ser Asn Val Pro Trp Thr
1 5

35

<210> 400
<211> 9
<212> PRT
<213> Artificial Sequence

40

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

45

<400> 400
Gln Gln Tyr Ser Ser Val Pro Trp Thr
1 5

50

<210> 401
<211> 9
<212> PRT
<213> Artificial Sequence

55

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 401
Ala Gln Tyr Ser Thr Val Pro Trp Thr
1 5

5
<210> 402
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

10
<400> 402
Gln Ala Tyr Ser Thr Val Pro Trp Thr
1 5

15
<210> 403
<211> 9
<212> PRT
<213> Artificial Sequence

20
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

25
<400> 403
Gln Gln Ala Ser Thr Val Pro Trp Thr
1 5

30
<210> 404
<211> 9
<212> PRT
<213> Artificial Sequence

35
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

40
<400> 404
Gln Gln Tyr Ala Thr Val Pro Trp Thr
1 5

45
<210> 405
<211> 9
<212> PRT
<213> Artificial Sequence

50
<220>
<221> source
<223> /note="Description of Artificial Sequence: Synthetic peptide"

55
<400> 405
Gln Gln Tyr Ser Ala Val Pro Trp Thr
1 5

<210> 406
<211> 9

<212> PRT
 <213> Artificial Sequence

 <220>
 5 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"

 <400> 406
 10 Gln Gln Tyr Ser Thr Ala Pro Trp Thr
 1 5

 <210> 407
 <211> 9
 15 <212> PRT
 <213> Artificial Sequence

 <220>
 <221> source
 20 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"

 <400> 407
 Gln Gln Tyr Ser Thr Val Ala Trp Thr
 1 5
 25

 <210> 408
 <211> 9
 <212> PRT
 30 <213> Artificial Sequence

 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"
 35
 <400> 408
 Gln Gln Tyr Ser Thr Val Pro Ala Thr
 1 5
 40

 <210> 409
 <211> 9
 <212> PRT
 <213> Artificial Sequence
 45

 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic
 peptide"

 <400> 409
 50 Gln Gln Tyr Ser Thr Val Pro Trp Ala
 1 5

 <210> 410
 <211> 9
 55 <212> PRT
 <213> Artificial Sequence

5
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"

<400> 410
 Gln Gln Tyr Asn Ser Leu Pro Trp Thr
 1 5

10
 <210> 411
 <211> 9
 <212> PRT
 <213> Artificial Sequence

15
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"

20
 <400> 411
 Gly Tyr Thr Phe Thr Asn Tyr Gly Met
 1 5

25
 <210> 412
 <211> 15
 <212> PRT
 <213> Artificial Sequence

30
 <220>
 <221> source
 <223> /note="Description of Artificial Sequence: Synthetic peptide"

35
 <400> 412
 Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
 1 5 10 15

40 **Claims**

1. An anti-VEGF antibody or an anti-VEGF binding fragment of an antibody which comprises six CDRs that have one or more amino acid substitutions or combinations of amino acid substitutions compared to an anti-VEGF antibody or an anti-VEGF binding fragment that has CDRs having amino acid sequences corresponding to SEQ ID NO:3 (CDR-H1), SEQ ID NO:4 (CDR-H2), SEQ ID NO:5 (CDR-H3), SEQ ID NO:6 (CDR-L1), SEQ ID NO:7 (CDR-L2) and SEQ ID NO:8 (CDR-L3), wherein said one or more amino acid substitution or combination of amino acid substitutions are selected from:

- 45
 50
 55
 (i) the CDR-H2 substitution K64S;
 (ii) the CDR-H2 substitution K64Q;
 (iii) the CDR-H2 substitutions Y53F and K64Q;
 (iv) the CDR-H3 substitution H97E;
 (v) the CDR-H3 substitution H97P;
 (vi) the CDR-H3 substitutions H97E and Y98F; and
 (vii) the CDR-L2 substitution T51A.

2. The anti-VEGF antibody or anti-VEGF binding fragment of claim 1 which is a monoclonal antibody or anti-VEGF binding fragment of a monoclonal antibody, respectively.

3. The anti-VEGF antibody or anti-VEGF binding fragment of claim 1 which is a human or humanized antibody, or anti-VEGF binding fragment of a human or humanized antibody, respectively.
- 5 4. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of claims 1 to 3 which is an IgG, optionally which is an IgG₁ or an IgG₂.
- 10 5. The anti-VEGF antibody or an anti-VEGF binding fragment of claim 1 which includes one or more mutations in the Fc regions that increase ADCC activity, or which includes one or more mutations in the Fc regions that decrease ADCC activity.
- 15 6. The anti-VEGF antibody or an anti-VEGF binding fragment of claim 1 which is non-fucosylated; or which includes one or more mutations in the Fc region that increase binding to Fc γ R; or which includes one or more mutations in the Fc region that increase binding to FcRn.
- 20 7. The anti-VEGF antibody or an anti-VEGF binding fragment of claim 1 which has other than said one or more substitutions, a V_H sequence corresponding to SEQ ID NO:1 and a V_L sequence corresponding to SEQ ID NO:2, or which has other than said one or more substitutions, a V_H sequence corresponding to SEQ ID NO:9 and a V_L sequence corresponding to SEQ ID NO:10.
- 25 8. The anti-VEGF antibody or an anti-VEGF binding fragment of claim 1 which has an affinity that is 1.5- to 50-fold greater than the affinity of an antibody having a V_H sequence corresponding to SEQ ID NO:1 and a V_L sequence corresponding to SEQ ID NO:2, or which has an affinity that is 1.5- to 50-fold greater than the affinity of an antibody having a V_H sequence corresponding to SEQ ID NO:9 and a V_L sequence corresponding to SEQ ID NO:10, optionally wherein said affinity is a measure of K_D as analysed by a biosensor.
- 30 9. The anti-VEGF antibody or an anti-VEGF binding fragment of claim 8 which has an affinity that is 2- to 30-fold greater than the affinity of an antibody having a V_H sequence corresponding to SEQ ID NO:1 and a V_L sequence corresponding to SEQ ID NO:2, or which has an affinity that is 2- to 30-fold greater than the affinity of an antibody having a V_H sequence corresponding to SEQ ID NO:9 and a V_L sequence corresponding to SEQ ID NO:10.
- 35 10. The anti-VEGF antibody or an anti-VEGF binding fragment of any one of the preceding claims which is purified, optionally purified to at least 85%, at least 90%, at least 95%, or at least 98% homogeneity.
- 40 11. An antibody-drug conjugate comprising an anti-VEGF antibody or anti-VEGF binding fragment according to anyone of claims 1 to 10.
- 45 12. A pharmaceutical composition comprising an anti-VEGF antibody or anti-VEGF binding fragment according to anyone of claims 1 to 10 or an antibody-drug conjugate according to claim 11, and a pharmaceutically acceptable carrier.
- 50 13. An anti-VEGF antibody or anti-VEGF binding fragment according to any one of claims 1 to 10, an antibody-drug conjugate according to claim 11, or a pharmaceutical composition according to claim 12, for use in treating cancer in a human patient, optionally wherein the cancer is metastatic carcinoma of the colon, metastatic carcinoma of the rectum, nonsquamous non-small cell lung cancer, or metastatic HER2-negative breast cancer, further optionally wherein the non-squamous non-small cell lung cancer is unresectable, locally advanced, recurrent or metastatic.
- 55 14. An anti-VEGF antibody or anti-VEGF binding fragment according to any one of claims 1 to 10, an antibody-drug conjugate according to claim 11, or a pharmaceutical composition according to claim 12, for use in treating age-related macular degeneration in a human patient.
15. An anti-VEGF antibody or anti-VEGF binding fragment according to any one of claims 1 to 10, an antibody-drug conjugate according to claim 11, or a pharmaceutical composition according to claim 12, for use in treating an immune disorder in a human patient, optionally wherein the immune disorder is rheumatoid arthritis or Grave's disease.

Antibody Chain	CDR No.	Residue	Position in CDR	Kabat No.
Heavy	1	G	1	26
		Y	2	27
		T	3	28
		F	4	29
		T	5	30
		N	6	31
		Y	7	32
		G	8	33
		M	9	34
		N	10	35
Heavy	2	W	1	50
		I	2	51
		N	3	52
		T	4	52a
		Y	5	53
		T	6	54
		G	7	55
		E	8	56
		P	9	57
		T	10	58
		Y	11	59
		A	12	60
		A	13	61
		D	14	62
		F	15	63
		K	16	64
		R	17	65
Heavy	3	Y	1	95
		P	2	96
		H	3	97
		Y	4	98
		Y	5	99
		G	6	100
		S	7	100a
		S	8	100b
		H	9	100c
		W	10	100d
		Y	11	100e
		F	12	100f
		D	13	101
		V	14	102

TABLE 1

Antibody Chain	CDR No.	Residue	Position in CDR	Kabat No.
Light	1	S	1	24
		A	2	25
		S	3	26
		Q	4	27
		D	5	28
		I	6	29
		S	7	30
		N	8	31
		Y	9	32
		L	10	33
		N	11	34
Light	2	F	1	50
		T	2	51
		S	3	52
		S	4	53
		L	5	54
		H	6	55
		S	7	56
Light	3	Q	1	89
		Q	2	90
		Y	3	91
		S	4	92
		T	5	93
		V	6	94
		P	7	95
		W	8	96
		T	9	97

TABLE 2

SEQ ID NO:	VL Peptide
13.	DIQMTQSPSSLSASV
14.	MTQSPSSLSASVGDR
15.	SPSSLSASVGDRVTI
16.	SLSASVGDRVTTITCS
17.	ASVGDRVTTITCSASQ
18.	GDRVTTITCSASQDIS
19.	VTITCSASQDISNYL
20.	TCSASQDISNYLNWY
21.	ASQDISNYLNWYQQK
22.	DISNYLNWYQQKPGK
23.	NYLNWYQQKPGKAPK
24.	NWYQQKPGKAPKVLII
25.	QQKPGKAPKVLIIYFT
26.	PGKAPKVLIIYFTSSL
27.	APKVLIIYFTSSLHSG
28.	VLIYFTSSLHSGVPS
29.	YFTSSLHSGVPSRFS
30.	SSLHSGVPSRFSGSG
31.	HSGVPSRFSGSGSGT
32.	VPSRFSGSGSGTDFT
33.	RFSGSGSGTDFTLTI
34.	GSGSGTDFTLTISSL
35.	SGTDFTLTISSLQPE
36.	DFTLTISSLQPEDFA
37.	LTISSLQPEDFATYY
38.	SSLQPEDFATYYCQQ
39.	QPEDFATYYCQQYST
40.	DFATYYCQQYSTVPW
41.	TYYCQQYSTVPWTFG
42.	CQQYSTVPWTFGQGT
43.	YSTVPWTFGQGTKVE
44.	VPWTFGQGTKVEIKR

TABLE 3

SEQ ID NO:	VH Peptide
45.	EVQLVESGGGLVQPG
46.	LVESGGGLVQPGGSL
47.	SGGGLVQPGGSLRLS
48.	GLVQPGGSLRLSCAA
49.	QPGGSLRLSCAASGY
50.	GSLRLSCAASGYTFT
51.	RLSCAASGYTFTNYG
52.	CAASGYTFTNYGMNW
53.	SGYTFTNYGMNWVRQ
54.	TFTNYGMNWVRQAPG
55.	NYGMNWVRQAPGKGL
56.	MNWVRQAPGKGLEWV
57.	VRQAPGKGLEWVGWI
58.	APGKGLEWVGWINTY
59.	KGLEWVGWINTYTGE
60.	EWVGWINTYTGEPTY
61.	GWINTYTGEPTYAAD
62.	NTYTGEPTYAADFKR
63.	TGEPTYAADFKRRFT
64.	PTYAADFKRRFTFSL
65.	AADFKRRFTFSLDTS
66.	FKRRFTFSLDTSKST
67.	RFTFSLDTSKSTAYL
68.	FSLDTSKSTAYLQMN
69.	DTSKSTAYLQMNSLR
70.	KSTAYLQMNSLRAED
71.	AYLQMNSLRAEDTAV
72.	QMNSLRAEDTAVYYC
73.	SLRAEDTAVYYCAKY
74.	AEDTAVYYCAKYPHY
75.	TAVYYCAKYPHYYS
76.	YYCAKYPHYYSSSH
77.	AKYPHYYSSSHWFYD
78.	PHYYSSSHWFYDVG
79.	YGSSSHWFYDVGQGT
80.	SHWFYDVGQGTLV
81.	YFDVVGQGTLVTVSS

TABLE 4

SEQ ID. NO.	Sequence	n	%	SI	s.e.m.
25	QQKPGKAPKVL <u>I</u> YFT	15	15.15	1.82	0.24
62	NTYTGEPTYAAD <u>F</u> KR	16	16.16	2.16	0.35
74	AEDTAVYYCAK <u>Y</u> PHY	9	9.09	1.45	0.18

TABLE 5

WT	position	Candidate deimmunizing substitutions
CDR-H2		
A	61	F, E, D
D	62	G, L, Q, T, K, R, E, H
K	64	S, V, Q
R	65	V, F, H, N, S, Q, K, I
CDR-H3		
Y	98	H

TABLE 6

SEQ ID NO:	28	31	97	99	100a	k_{on} (1/Ms)	k_{off} (1/s)	K_D (M)	Δk_{on}	Δk_{off}	ΔK_D
82		F	D	D	G	2.08E+06	1.72E-04	8.27E-11	11.6	2.0	23.8
83		F	P	D	G	1.88E+06	1.94E-04	1.06E-10	10.4	1.8	18.6
84		F	P	E		1.60E+06	1.82E-04	1.28E-10	8.9	1.9	15.3
85		F	E	E		1.25E+06	1.75E-04	1.40E-10	6.9	2.0	14.1
86		F	D	E		1.33E+06	2.38E-04	1.79E-10	7.4	1.5	11.0
87		F	E	D	G	8.24E+05	1.60E-04	2.04E-10	4.6	2.2	9.6
88		F		D	G	9.68E+05	2.73E-04	2.82E-10	5.4	1.3	7.0
89		F	P	D		9.04E+05	2.49E-04	2.82E-10	5.0	1.4	7.0
90		F	D		G	6.45E+05	1.97E-04	3.11E-10	3.6	1.8	6.3
91		F				2.11E+05	1.21E-04	5.75E-10	1.2	2.9	3.4
92			P			3.67E+05	2.56E-04	7.08E-10	2.0	1.4	2.8
93			E			2.92E+05	2.25E-04	7.61E-10	1.6	1.6	2.6
94		F			G	2.58E+05	2.20E-04	8.57E-10	1.4	1.6	2.3
95			D			2.62E+05	3.96E-04	1.54E-09	1.5	0.9	1.3
96		F	P		G	2.25E+05	3.79E-04	1.75E-09	1.3	0.9	1.1
WT	T	N	H	Y	S	1.80E+05	3.52E-04	1.97E-09	1.0	1.0	1.0

TABLE 7

WT	Position	Increased affinity heavy chain substitutions
CDR-H1		
T	28	P
N	31	F, G, M
CDR-H3		
H	97	A, D, E, P, Q, S, T
Y	99	D, E
S	100a	D, E, G, V

TABLE 8

WT	Position	Increased affinity heavy chain substitutions
CDR-H1		
T	30	W, R, Q
CDR-H2		
Y	53	F
T	58	F
A	61	G, K, R, H, Y
K	64	G, E
R	65	L, T, A, E, D
CDR-H3		
Y	98	F
Y	100e	F

TABLE 9

WT	Position	Neutral heavy chain substitutions
CDR-H1		
T	30	I, M, V
N	31	Y, S, A, Q
CDR-H2		
A	61	F, E, D
D	62	G, L, Q, T, K, R, E, H
K	64	S, V, Q
R	65	V, F, H, N, S, Q, K, I
CDR-H3		
Y	98	H

TABLE 10

WT	Position	Neutral light chain substitutions
CDR-L1		
S	24	R, W, G
S	26	V
S	30	G
N	31	L, S
CDR-L2		
L	54	M
H	55	R
S	56	G, R
CDR-L3		
T	93	D, E, S

TABLE 11

V#	CDR-H1										CDR-H2										CDR-H3																					
	G	Y	I	F	T	N	Y	G	M	W	I	N	T	Y	T	G	E	P	T	Y	A	A	D	F	K	R	Y	P	H	Y	G	S	S	H	W	Y	F	D	V			
114																																										
115																																										
116																																										
117																																										
118																																										
119	A																																									
120																																										
121	F				S	S	A	MS	V						S	G	D	G	G	S	T	Y	Y	D	S	V	C															
122																																										
123																																										
124																																										
125																																										
126																																										
127																																										
128																																										
129																																										
130																																										
131																																										
132																																										
133																																										
134																																										

TABLE 12-6

V#	CDR-L1											CDR-L2					CDR-L3										
	S	A	S	Q	D	I	S	N	Y	L	N	F	T	S	S	L	H	S	Q	Q	Y	S	T	V	P	W	T
206																											
207																					NS		-				
208																					YS		-				
209																					SY		-				
210																							A				
211																							N				
212																							S				
213	A																										
214		A																									
215			A																								
216				A																							
217					A																						
218						A																					
219							A																				
220								A																			
221									A																		
222										A																	
223											A																
224												A															
225													A														
226														A													
227															A												
228																A											
229																	A										
230																		A									
231																			A								
232																				A							
233																					A						
234																						A					
235																							A				
236																								A			
237																									A		
238																										A	
239																											A
240	R		N	E	Q	L																					
241	R		N	E	Q																						
242	R				S						A	A	A				E					N	S	L			

TABLE 13

SEQ ID NO:	N	I	Y	I	G	E	P	I	Y	A	A	D	F	K	R	# responders*	% responders	Average SI
62																5	5.38	1.67
97			F													9	9.68	1.59
98											F					6	6.45	1.51
99											E					5	5.38	1.82
100											D					7	7.53	1.39
101												G				6	6.45	1.45
102												L				8	8.60	1.44
103												Q				10	10.75	1.46
104												T				4	4.30	1.40
105												K				5	5.38	1.50
106												R				4	4.30	1.32
107												E				5	5.38	1.32
108												H				4	4.30	1.21
109													S			2	2.15	1.22
110													V			4	4.30	1.19
111													Q			2	2.15	1.13
112														V		10	10.75	1.59
113														F		13	13.98	1.82
114														H		9	9.68	1.55
115														N		7	7.53	1.41
116														S		4	4.30	1.24
117														Q		7	7.53	1.33
118														K		4	4.30	1.27
119														I		9	9.68	2.06
* n= 93																		

TABLE 14

SEQ ID NO:	N	I	Y	I	G	E	P	I	Y	A	A	D	F	K	R	# responders*	% responders	Average SI
120			F								F					7	7.53	1.68
121			F								E					8	8.60	1.51
122			F								D					4	4.30	1.34
123			F									G				3	3.23	1.19
124			F									L				6	6.45	1.29
125			F									Q				6	6.45	1.34
126			F									T				6	6.45	1.34
127			F									K				9	9.68	1.48
128			F									R				8	8.60	1.67
129			F									E				10	10.75	1.61
130			F									H				8	8.60	1.64
131			F										S			4	4.30	1.21
132			F										V			3	3.23	1.12
133			F										Q			2	2.15	1.15
134			F											V		6	6.45	1.41
135			F											F		12	12.90	1.65
136			F											H		11	11.83	1.70
137			F											N		9	9.68	1.80
138			F											S		11	11.83	1.80
139			F											Q		7	7.53	1.34
140			F											K		4	4.30	1.17
141			F											I		5	5.38	1.36
62																6	6.45	1.33
* n= 93																		

TABLE 14 (continued)

SEQ ID NO:	A	E	D	T	A	V	Y	Y	C	A	K	Y	P	H	Y	# responders*	% responders	Average SI
74																7	7.53	1.45
142														D		11	11.83	1.62
143														E		6	6.45	1.43
144														P		10	10.75	1.97
145															F	4	4.30	1.26
146														D	F	5	5.38	1.22
147														E	F	1	1.08	1.09
148														P	F	5	5.38	1.22
74																6	6.45	1.34
* n= 93																		

TABLE 15

SEQ ID NO:	Q	Q	K	P	G	K	A	P	K	V	L	I	Y	F	I	# responders*	% responders	Average SI
25																24	25.81	3.69
149										L						18	19.35	2.05
150														D		6	6.45	1.42
151														S		6	6.45	1.43
152														G		2	2.15	1.27
153														A		2	2.15	1.17
154															A	2	2.15	1.16
155														D	A	7	7.53	1.87
156														S	A	8	8.60	2.34
157														G	A	8	8.60	1.63
158														A	A	11	11.83	1.77
159										L				D		12	12.90	1.49
160										L				S		3	3.23	1.25
161										L				G		4	4.30	1.27
162										L				A		5	5.38	1.28
163										L					A	23	24.73	2.49
164										L				D	A	9	9.68	2.23
165										L				S	A	10	10.75	1.39
166										L				G	A	6	6.45	1.52
167										L				A	A	5	5.38	1.27
25																14	15.05	2.30
* n= 93																		

TABLE 16

SEQ ID NO:				
VH	CDR2 16%/2.16 SI	# Responders*	%	SI
62	NTYTGEPTYAADFKR	5	5.38	1.67
109	NTYTGEPTYAADFSR	2	2.15	1.22
111	NTYTGEPTYAADFQR	2	2.15	1.13
133	NTFTGEPTYAADFQR	2	2.15	1.15
62	NTYTGEPTYAADFKR	6	6.45	1.33
VH	CDR3 9%/1.45 SI			
74	AEDTAVYYCAKYPHY	7	7.53	1.45
143	AEDTAVYYCAKYPEF	1	1.08	1.09
74	AEDTAVYYCAKYPHY	6	6.45	1.34
VL	CDR2 15%/1.45 SI			
25	QQKPGKAPKVLIIYFT	24	25.81	3.69
154	QQKPGKAPKVLIIYFA	2	2.15	1.16
25	QQKPGKAPKVLIIYFT	14	15.05	2.30
	* n= 93			

TABLE 17

Single Variable Region Mutants	MFI
WT	194.78
VH K64Q	230.4
VH K64S	168.08
VH Y53F.K64Q	217.4
VH H97E.Y98F	310.32
VL T51A	295.93

TABLE 18

Combined Variable Region Mutants	EC₅₀
WT	152.4
VH K64Q/VL T51A	174.2
VH K64S/VL T51A	263.2
VH Y53F.K64Q/VL T51A	104
VH H97E.Y98F/VL T51A	26.22

TABLE 19

Variant No.	SEQ ID NOS		
	CDR-H1	CDR-H2	CDR-H3
1	n/a	4	237
2	n/a	190	238
3	n/a	191	238
4	n/a	192	238
5	n/a	193	238
6	n/a	194	238
7	n/a	195	238
8	n/a	196	238
9	n/a	197	238
10	n/a	198	238
11	n/a	199	238
12	n/a	4	238
13	n/a	4	239
14	n/a	4	240
15	n/a	4	241
16	n/a	200	n/a
17	n/a	201	n/a
18	n/a	202	n/a
19	n/a	203	5
20	n/a	204	5
21	n/a	205	n/a
22	n/a	206	n/a
23	n/a	207	n/a
24	n/a	208	n/a
25	n/a	209	n/a
26	n/a	210	n/a
27	n/a	211	n/a
28	n/a	212	n/a

TABLE 20-1

Variant No.	SEQ ID NOS		
	CDR-H1	CDR-H2	CDR-H3
29	n/a	n/a	242
30	n/a	n/a	243
31	n/a	n/a	244
32	n/a	n/a	245
33	n/a	n/a	246
34	n/a	n/a	247
35	n/a	n/a	248
36	n/a	n/a	249
37	n/a	n/a	250
38	n/a	n/a	251
39	n/a	n/a	252
40	n/a	n/a	253
41	n/a	n/a	254
42	n/a	n/a	255
43	n/a	n/a	256
44	n/a	n/a	257
45	n/a	n/a	258
46	n/a	n/a	259
47	n/a	n/a	260
48	n/a	n/a	261
49	n/a	n/a	262
50	n/a	n/a	263

TABLE 20-2

Variant No.	SEQ ID NOS		
	CDR-H1	CDR-H2	CDR-H3
51	n/a	n/a	264
52	n/a	n/a	265
53	n/a	n/a	266
54	n/a	n/a	267
55	n/a	n/a	268
56	n/a	n/a	269
57	n/a	n/a	270
58	n/a	n/a	271
59	n/a	n/a	272
60	n/a	n/a	273
61	n/a	n/a	274
62	n/a	n/a	275
63	n/a	n/a	276
64	n/a	n/a	277
65	n/a	n/a	278
66	n/a	n/a	279
67	n/a	n/a	280
68	n/a	n/a	281
69	n/a	n/a	282
70	n/a	n/a	283

TABLE 20-3

Variant No.	SEQ ID NOS		
	CDR-H1	CDR-H2	CDR-H3
71	n/a	n/a	284
72	n/a	n/a	285
73	n/a	n/a	286
74	n/a	n/a	287
75	n/a	n/a	288
76	n/a	n/a	289
77	n/a	n/a	290
78	n/a	n/a	291
79	n/a	n/a	292
80	n/a	n/a	293
81	n/a	n/a	294
82	n/a	n/a	295
83	n/a	n/a	296
84	n/a	n/a	297
85	n/a	n/a	298
86	n/a	n/a	299
87	n/a	n/a	300
88	n/a	n/a	301
89	n/a	n/a	302
90	n/a	n/a	303
91	n/a	n/a	304
92	n/a	n/a	305
93	n/a	n/a	306

TABLE 20-4

Variant No.	SEQ ID NOS		
	CDR-H1	CDR-H2	CDR-H3
94	n/a	n/a	307
95	n/a	n/a	308
96	n/a	n/a	309
97	n/a	n/a	310
98	n/a	n/a	311
99	n/a	n/a	312
100	n/a	n/a	313
101	n/a	n/a	314
102	n/a	n/a	315
103	n/a	n/a	316
104	n/a	n/a	317
105	n/a	n/a	318
106	n/a	n/a	319
107	n/a	n/a	320
108	411	4	321
109	168	4	322
110	411	4	323
111	168	4	324
112	169	4	325
113	411	4	326

TABLE 20-5

Variant No.	SEQ ID NOS		
	CDR-H1	CDR-H2	CDR-H3
114	411	4	321
115	170	213	n/a
116	169	4	5
117	168	4	n/a
118	169	n/a	n/a
119	171	4	5
120	411	4	327
121	172	214	328
122	411	4	329
123	411	4	330
124	411	4	331
125	411	4	332
126	411	4	333
127	411	4	334
128	169	4	335
129	169	4	336
130	169	4	337
131	169	4	338
132	169	4	339
133	411	4	340
134	169	4	341

TABLE 20-6

Variant No.	SEQ ID NOS		
	CDR-H1	CDR-H2	CDR-H3
135	169	4	342
136	169	4	343
137	169	4	344
138	169	4	345
139	169	4	346
140	169	4	347
141	169	4	348
142	169	4	349
143	169	4	350
144	169	4	351
145	169	4	352
146	169	4	353
147	169	4	354
148	169	4	355
149	169	4	356
150	169	4	357
151	169	4	358
152	173	215	359
153	173	4	359
154	169	4	359
155	169	4	241
156	411	4	360
157	169	4	361

TABLE 20-7

Variant No.	SEQ ID NOS		
	CDR-H1	CDR-H2	CDR-H3
158	169	4	362
159	169	4	363
160	411	4	364
161	169	4	365
162	169	4	366
163	411	4	367
164	411	4	368
165	169	4	369
166	411	4	370
167	411	4	371
168	411	216	5
169	411	217	5
170	411	218	5
171	411	219	5
172	411	220	5
173	411	221	5
174	411	222	5
175	411	223	5
176	411	224	5
177	169	225	5
178	169	226	5
179	411	227	5
180	411	228	5
181	169	229	5

TABLE 20-8

Variant No.	SEQ ID NOS		
	CDR-H1	CDR-H2	CDR-H3
182	411	230	5
183	169	231	5
184	411	232	5
185	411	233	5
186	411	234	5
187	411	235	5
188	174	4	5
189	175	4	5
190	176	4	5
191	177	4	5
192	178	4	5
193	173	4	5
194	179	4	5
195	180	4	5
196	181	4	5
197	182	4	5
198	183	4	5
199	184	4	5
200	185	4	5
201	186	4	5
202	187	4	5
203	188	4	5
204	189	4	5
205	411	236	5

TABLE 20-9

Variant #	SEQ ID NOS		
	CDR-L1	CDR-L2	CDR-L3
206	6	7	394
207	n/a	n/a	395
208	n/a	n/a	396
209	n/a	n/a	397
210	n/a	n/a	398
211	n/a	n/a	399
212	n/a	n/a	400
213	372	7	8
214	373	7	8
215	374	7	8
216	375	7	8
217	376	7	8
218	377	7	8
219	378	7	8
220	379	7	8
221	380	7	8
222	381	7	8
223	382	7	8
224	6	386	8
225	6	387	8
226	6	388	8
227	6	389	8
228	6	390	8
229	6	391	8
230	6	392	8
231	6	7	401
232	6	7	402
233	6	7	403
234	6	7	404
235	6	7	405
236	6	7	406
237	6	7	407
238	6	7	408
239	6	7	409
240	383	7	8
241	384	7	8
242	385	393	410

TABLE 20-10

Bevacizumab (Avastin®) VH (SEQ ID NO:1)

EVQLVESGGGLVQPGGSLRLSCAASGYTFTNYGMNWVRQAPGKGLEWVGWINTYTGEPT
YAADFKRRFTFSLDTSKSTAYLQMNSLRAEDTAVYYCAKYPHYYGSSHWFYFDVWGQGL
 VTVSS

Bevacizumab (Avastin®) VL (SEQ ID NO:2)

DIQMTQSPSSLSASVGDRVTITCSASQDISNYLNWYQQKPGKAPKVLIIFTSSLHSGVP
 SRFSGSGSGTDFTLTISLQPEDFATYYCQQYSTVPWTFGQGTKVEIKR

FIGURE 1A

Antibody Chain	CDR No.	Sequence	SEQ ID NO.
Heavy	1	GYTFTNYGMN	3
Heavy	2	WINTYTGEPTYAADFKR	4
Heavy	3	YPHYYGSSHWFYFDV	5
Light	1	SASQDISNYLN	6
Light	2	FTSSLHS	7
Light	3	QQYSTVPWT	8

FIGURE 1B

Ranibizumab (Lucentis®) VH (SEQ ID NO:9)

EVQLVESGGGLVQPGGSLRLSCAAS**GYDFTHYGMN**WVRQAPGKGLEWVG**WINTYTGEPT**
YAADFKRRFTFSLDTSKSTAYLQMNSLRAEDTAVYYCAK**YPYYGTSHWYFDV**WGQGTL
 VTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFP
 AVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHL

Ranibizumab (Lucentis®) VL (SEQ ID NO:10)

DIQLTQSPSSLSASVGDRTITTC**SASQDISNYLN**WYQQKPGKAPKVLII**FTSSLHS**GVP
 SRFSGSGSGTDFTLTITSSLPEDFATYYC**QOYSTVPWT**FGQGTKVEIKRTVAAPSVFIF
 PPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSYSLSS
 TLTLISKADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC

FIGURE 1C

Antibody Chain	CDR No.	Sequence	SEQ ID NO.
Heavy	1	GYDFTHYGMN	11
Heavy	2	WINTYTGEPTYAADFKR	4
Heavy	3	YPYYGTSHWYFDV	12
Light	1	SASQDISNYLN	6
Light	2	FTSSLHS	7
Light	3	QOYSTVPWT	8

FIGURE 1D




FIGURE 2A-2B




FIGURE 3A-3B

FIGURE 4A




FIGURE 4B





FIGURE 4C





EUROPEAN SEARCH REPORT

Application Number
EP 14 19 7237

5

10

15

20

25

30

35

40

45

50

55

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
A	WO 03/099999 A2 (ABMAXIS INC [US]; LUO PEIZHI [US]; HSIEH MARK [US]; ZHONG PINGYU [US];) 4 December 2003 (2003-12-04) * page 142, line 7; figures 9A and 9B, 14B, 16A; example 1; sequences 1,2 *	1-15	INV. C07K16/22 A61P35/00 A61K39/395 C12N15/13 C12N15/63 A61K47/48 A61P37/00
X	US 2002/032315 A1 (BACA MANUEL [AU] ET AL) 14 March 2002 (2002-03-14) * paragraphs [0208], [0304]; claim 9; table 15 *	1-15	
X	WO 2005/054273 A2 (ABMAXIS INC [US]; ZHONG PINGYU [US]; LUO PEIZHI [US]; WANG KEVIN C [US]) 16 June 2005 (2005-06-16) * claim 19 *	1-15	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (IPC)
			C07K
Place of search		Date of completion of the search	Examiner
The Hague		10 April 2015	Vadot, Pierre
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			

2
EPO FORM 1503 03.02 (P04/C01)

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 14 19 7237

5

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

10-04-2015

10

15

20

25

30

35

40

45

50

55

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 03099999 A2	04-12-2003	AU 2003248548 A1	12-12-2003
		CA 2485732 A1	04-12-2003
		CN 1672160 A	21-09-2005
		EP 1514216 A2	16-03-2005
		JP 2005526518 A	08-09-2005
		SG 135053 A1	28-09-2007
		WO 03099999 A2	04-12-2003

US 2002032315 A1	14-03-2002	US 2002032315 A1	14-03-2002
		US 2003190317 A1	09-10-2003
		US 2007196374 A1	23-08-2007
		US 2008226629 A1	18-09-2008
		US 2011052575 A1	03-03-2011

WO 2005054273 A2	16-06-2005	AU 2004295339 A1	16-06-2005
		CA 2547016 A1	16-06-2005
		CN 1946422 A	11-04-2007
		CN 101602806 A	16-12-2009
		CN 101899113 A	01-12-2010
		EP 1699484 A2	13-09-2006
		JP 2008504215 A	14-02-2008
		KR 20070036018 A	02-04-2007
		SG 146696 A1	30-10-2008
		US 2004133357 A1	08-07-2004
		WO 2005054273 A2	16-06-2005

EPO FORM P0158

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 61218005 A [0001]
- US 5807715 A [0042]
- US 4816567 A [0042]
- US 4816397 A [0042] [0058]
- US 5530101 A [0043]
- US 5585089 A [0043]
- US 5693761 A [0043]
- US 5693762 A [0043]
- US 6180370 A, Queen [0043]
- EP 239400 A [0043]
- WO 9109967 A [0043]
- US 5225539 A [0043]
- EP 592106 A [0043]
- EP 519596 A [0043]
- US 5565332 A [0043]
- US 4444887 A [0044]
- US 4716111 A [0044]
- WO 9846645 A [0044]
- WO 9850433 A [0044]
- WO 9824893 A [0044]
- WO 9816654 A [0044]
- WO 9634096 A [0044]
- WO 9633735 A [0044]
- WO 9110741 A [0044]
- WO 9201047 A [0044]
- US 5413923 A [0044]
- US 5625126 A [0044]
- US 5633425 A [0044]
- US 5569825 A [0044]
- US 5661016 A [0044]
- US 5545806 A [0044]
- US 5814318 A [0044]
- US 5885793 A [0044]
- US 5916771 A [0044]
- US 5939598 A [0044]
- US 5658570 A [0045]
- US 5681722 A [0045]
- US 5693780 A [0045]
- US 20060134709 A [0050]
- US 5834597 A [0051]
- WO 2005123780 A [0053]
- US 7217797 B [0053]
- US 20070280931 A [0054]
- US 5168062 A, Stinski [0065]
- US 4510245 A, Bell [0065]
- US 4968615 A, Schaffner [0065]
- US 4399216 A [0066]
- US 4634665 A [0066]
- US 5179017 A [0066]
- US 6787323 B [0080]
- WO 8601533 A [0114]
- EP 0392745 A [0114]
- WO 2005117984 A [0114]
- US 5635483 A [0115]
- US 5780588 A [0115]
- WO 02088172 A [0115]
- US 7498298 B [0115]
- US 5208020 A [0116]
- US 5714586 A [0116]
- US 5712374 A [0116]
- US 5264586 A [0116]
- US 5773001 A [0116]
- US 5219996 A [0118]
- EP 0948544 A [0119]
- US 4737456 A [0124]

Non-patent literature cited in the description

- FOLKMAN et al. *J. Biol. Chem.*, 1992, vol. 267, 10931-10934 [0004]
- FERRARA et al. *Nature Medicine*, 2003, vol. 9, 669-676 [0004] [0005] [0006]
- FERRARA et al. *Nature*, 1996, vol. 380, 439-442 [0005]
- DVORAK et al. *Am. J. Pathol.*, 1995, vol. 146, 1029-1039 [0005]
- BERKMAN et al. *J. Clin. Invest.*, 1993, vol. 91, 153-159 [0005]
- AIELLO et al. *N. Engl. J. Med.*, 1994, vol. 331, 1480-1487 [0005]
- LOPEZ et al. *Invest. Ophthalmo. Vis. Sci.*, 1996, vol. 37, 855-868 [0005]
- PRESTA et al. *Cancer Res.*, 1997, vol. 57, 4593-4599 [0007]
- Therapeutic Antibodies. LIEN ; LOWMAN. Handbook of Experimental Pharmacology. Springer-Verlag, 2008, vol. 181, 131-150 [0008]
- BLOWERS ; HALL. *Br. J. Nurs.*, 2009, vol. 18 (6), 351-6, 358 [0009]
- BOSTROM et al. *Science*, 2009, vol. 323, 1610-14 [0021]
- WAHL et al. *J. Nucl. Med.*, 1983, vol. 24, 316 [0031]

- **KABAT et al.** Sequences of Proteins of Immunological Interest. National Institute of Health, 1987 [0036]
- **RIECHMANN.** *Journal of Immunological Methods*, 1999, vol. 231, 25-38 [0039]
- **MORRISON.** *Science*, 1985, vol. 229 (4719), 1202-7 [0042]
- **Ol et al.** *BioTechniques*, 1986, vol. 4, 214-221 [0042]
- **GILLIES et al.** *J. Immunol. Methods*, 1985, vol. 125, 191-202 [0042]
- **RIECHMANN et al.** *Nature*, 1988, vol. 332, 323-7 [0043]
- **PADLAN.** *Mol. Immunol.*, 1991, vol. 28, 489-498 [0043]
- **STUDNICKA et al.** *Prot. Eng.*, 1994, vol. 7, 805-814 [0043]
- **ROGUSKA et al.** *Proc. Natl. Acad. Sci.*, 1994, vol. 91, 969-973 [0043]
- **JESPERs et al.** *Biotechnology*, 1988, vol. 12, 899-903 [0044]
- **WOLFSON.** *Chem. Biol.*, 2006, vol. 13 (10), 1011-2 [0047]
- **CANFIELD ; MORRISON.** *J. Exp. Med.*, 1991, vol. 173, 1483-1491 [0049]
- **LUND et al.** *J. Immunol.*, 1991, vol. 147, 2657-2662 [0049]
- **SHIELDS et al.** *J. Biol. Chem.*, 2002, vol. 277, 26733-26740 [0052]
- **SHINKAWA et al.** *J. Biol. Chem.*, 2003, vol. 278, 3466-73 [0052]
- **JUNG ; PLÜCKTHUN.** *Protein Engineering*, 1997, vol. 10 (9), 959-966 [0054]
- **YAZAKI et al.** *Protein Eng Des. Sel.*, 17 August 2004, vol. 17 (5), 481-9 [0054]
- **KIEKE et al.** *Proc. Nat'l Acad. Sci. USA*, 1999, vol. 96, 5651-5656 [0055]
- **HUANG et al.** *Anal. Chem.*, 2005, vol. 77, 1432-1439 [0056]
- *Molecular Cloning; A Laboratory Manual.* Cold Spring Harbor, 1989 [0058]
- *Current Protocols in Molecular Biology.* Greene Publishing Associates, 1989 [0058]
- **KABAT, E. A. et al.** Sequences of Proteins of Immunological Interest. NIH Publication No. 91-3242, 1991 [0059] [0062]
- **TOMLINSON et al.** *J. Mol. Biol.*, 1992, vol. 22T, 116-198 [0059]
- **COX et al.** *Eur. J. Immunol.*, 1994, vol. 24, 827-836 [0059]
- **BIRD et al.** *Science*, 1988, vol. 242, 423-426 [0062]
- **HUSTON et al.** *Proc. Natl. Acad. Sci. USA*, 1988, vol. 85, 5879-5883 [0062]
- **MCCAFFERTY et al.** *Nature*, 1990, vol. 348, 552-554 [0062]
- **GOEDEL.** *Gene Expression Technology: Methods in Enzymology.* Academic Press, 1990, vol. 185 [0065]
- **URLAUB ; CHASIN.** *Proc. Natl. Acad. Sci. USA*, 1980, vol. 77, 4216-4220 [0067]
- **KAUFMAN ; SHARP.** *Mol. Biol.*, 1982, vol. 159, 601-621 [0067]
- **BOSTROM et al.** *Science*, 2009, vol. 323, 1610-1614 [0070]
- *Solid Phase Peptide Synthesis.* The Pierce Chemical Co, 1984 [0073]
- **CHU et al.** *Biochemia No. 2 (Roche Molecular Biologicals)*, 2001 [0073]
- **FISHER.** *Laboratory Techniques In Biochemistry And Molecular Biology.* Elsevier, 1980 [0075]
- **QIN et al.** *J. Biol. Chem.*, 2006, vol. 281, 32550-32558 [0080]
- **LIANG et al.** *Nat. Protoc.*, 2007, vol. 2, 329-333 [0080]
- **DEISSLER et al.** *British Journal of Ophthalmology*, 2008, vol. 92, 839-843 [0080]
- **YOHNO et al.** *Biological & Pharmaceutical Bulletin*, 2003, vol. 26 (4), 417-20 [0080]
- **KARLSSON et al.** *J. Immunol. Methods*, 1991, vol. 145, 229-240 [0084]
- **HELLSTROM et al.** *Controlled Drug Delivery.* 1987, 623-53 [0113]
- **THORPE et al.** *Immunol. Rev.*, 1982, vol. 62, 119-58 [0113]
- **DUBOWCHIK et al.** *Pharmacology and Therapeutics*, 1999, vol. 83, 67-123 [0113]
- **CHARI et al.** *Cancer Research*, 1992, vol. 52, 127-131 [0116]
- **HINMAN et al.** *Cancer Research*, 1993, vol. 53, 3336-3342 [0116]
- **LODE et al.** *Cancer Research*, 1998, vol. 58, 2925-2928 [0116]
- **PARK et al.** *Adv. Pharmacol.*, 1997, vol. 40, 399-435 [0117]
- **MARTY ; SCHWENDENER.** *Methods in Molecular Medicine*, 2004, vol. 109, 389-401 [0117]
- *Poly(ethyleneglycol) Chemistry, Biotechnical and Biomedical Applications.* Plenum Press, 1992 [0119]
- *Poly(ethyleneglycol) Chemistry and Biological Applications.* American Chemical Society, 1997 [0119]
- *Bioconjugation Protein Coupling Techniques for the Biomedical Sciences.* Grove Publishers, 1998 [0119]
- **CHAPMAN.** *Advanced Drug Delivery Reviews*, 2002, vol. 54, 531-545 [0119]
- **HARLOW et al.** *Antibodies: A Laboratory Manual.* Cold Spring Harbor Laboratory Press, 1988 [0123]
- *Remington's Pharmaceutical Sciences.* 1980 [0139]
- **CARMELIET ; JAIN.** *Nature*, 2000, vol. 407, 249-257 [0160]

摘要

本发明涉及针对血管内皮生长因子“VEGF”的抗体和所述抗体例如用于治疗与 VEGF 活性和/或 VEGF 过度产生相关的疾病的用途。