

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
21 May 2004 (21.05.2004)

PCT

(10) International Publication Number
WO 2004/042076 A2

(51) International Patent Classification⁷: **C12Q 1/44**,
1/68, A61K 38/17, 48/00, A61P 25/00, 9/00, 7/00, 29/00,
1/00, 5/00

MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT,
RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(21) International Application Number:
PCT/EP2003/011879

(22) International Filing Date: 25 October 2003 (25.10.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
02024994.2 8 November 2002 (08.11.2002) EP

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— *as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii)) for the following designations AE,
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH,
CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES,
FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH,
PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN,
TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO
patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG,
ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU,
TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT,
RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)*

Published:

— *without international search report and to be republished
upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(71) Applicant (*for all designated States except US*): **BAYER
HEALTHCARE AG** [DE/DE]; 51368 Leverkusen (DE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **GOLZ, Stefan**
[DE/DE]; Bückmannsmühle 46, 45326 Essen (DE).
BRÜGGEMEIER, Ulf [DE/DE]; Leysiefen 20, 42799
Leichlingen (DE). **GEERTS, Andreas** [DE/DE]; Schuck-
ertstr. 29, 42113 Wuppertal (DE).

(74) Common Representative: **BAYER HEALTHCARE
AG**; Law and Patents, Patents and Licensing, 51368
Leverkusen (DE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,

(54) Title: DIAGNOSTICS AND THERAPEUTICS FOR DISEASES ASSOCIATED WITH HUMAN PHOSPHODIESTERASE
4A (PDE4A)

(57) Abstract: Diagnostics and Therapeutics for Diseases Associated with Human Phosphodiesterase 4A (PDE4A) Abstract of Dis-
closure The invention provides a human PDE4A which is associated with the disorders of the peripheral and central nervous system,
cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease. The in-
vention also provides assays for the identification of compounds useful in the treatment or prevention of disorders of the peripheral
and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and en-
docrinological disease. The invention also features compounds which bind to and/or activate or inhibit the activity of PDE4A as well
as pharmaceutical compositions comprising such compounds.



WO 2004/042076 A2

Diagnostics and Therapeutics for Diseases Associated with Human Phosphodiesterase 4A (PDE4A)

5 **Technical field of the invention**

The present invention is in the field of molecular biology, more particularly, the present invention relates to nucleic acid sequences and amino acid sequences of a human PDE4A and its regulation for the treatment of disorders of the peripheral and
10 central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in mammals.

Background of the invention

15 PDE4A is a member of the enzyme family of phosphodiesterases (PDEs) [[Livi GP et al., 1990], [Lamontagne S et al., 2001], [Sullivan M et al., 1998], WO0179526, US 5977305]. PDEs catalyze the hydrolyzation of 3', 5' cyclic nucleotides. That results in the formation of the respective nucleoside 5' monophosphates. The cyclic nucleotides cAMP and cGMP serve as crucial second messengers in a number of cellular
20 signaling pathways. The PDEs as well as the guanylyl and adenylyl cyclases, which synthesize the cyclic nucleotides, are important cellular components to regulate the concentration of cyclic nucleotides and, thus, to regulate the signal transduction pathways. Because of their central role in regulating second messenger levels PDEs have been considered chemotherapeutic targets and have been worked on
25 extensively.

Several families of PDEs have been identified. The nomenclature system includes first a number that indicates the PDE family. To date, eleven families (PDE 1-11) are known which are classified by: (i) primary structure; (ii) substrate preference; (iii)
30 response to different modulators; (iv) sensitivity to specific inhibitors; and (v) modes of regulation [Loughney and Ferguson, (1996)]. The number indicating the family is

- 2 -

followed by a capital letter, indicating a distinct gene, and the capital letter followed by a second number, indicating a specific splice variant or a specific transcript that utilizes a unique transcription initiation site.

5 PDEs show of the following structural features:

All mammalian PDEs identified to date possess a highly conserved region of 270-300 amino acids in the carboxy terminal half of the protein [Charbonneau, et al. (1986)]. Here, the catalytic site for cAMP and/or cGMP hydrolysis and two putative zinc binding sites as well as family specific determinants are located [Beavo, (1995); Francis, et al. (1994)]. The amino terminal regions of the various PDEs are highly variable and include other family specific determinants and diverse regulatory motifs such as: (i) calmodulin binding sites (PDE1); (ii) non-catalytic cyclic GMP binding sites (PDE2, PDE5, PDE6); (iii) membrane targeting sites (PDE4); (iv) hydrophobic membrane association sites (PDE3); and (v) phosphorylation sites for either the calmodulin-dependent kinase II (PDE1), the cAMP-dependent kinase (PDE1, PDE3, PDE4), or the cGMP dependent kinase (PDE5) [Beavo, (1995); Manganiello, et al. (1995); Conti, et al. (1995)].

Members of the PDE1 family are calcium-calmodulin dependent. The group is composed of at least three genes with several splicing variants [Kakkar, R. et al. (1999)]; PDE1A and PDE1B preferentially hydrolyze cGMP while PDE1C is dual-specific, it exhibits a high affinity for both cAMP and cGMP. In vitro experiments show regulation of some PDE1 species by phosphorylation, which decreases the affinity of the enzyme for calmodulin [Kakkar, (1999)]. PDE1s have been shown to be expressed in lung, heart and brain.

The PDE2 family is characterized as being specifically stimulated by cGMP [Loughney and Ferguson, supra]. PDE2 species have been found in cerebellum, neocortex, heart, kidney, lung, pulmonary artery, and skeletal muscle [Sadhu, K. et al. (1999)]. Only one gene has been identified, PDE2A. The respective PDE2A protein is specifically inhibited by erythro-9-(2-hydroxy-3-nonyl)adenine (EHNA).

- 3 -

Two genes have been identified in the PDE3 family, PDE3A and PDE3B, both having high affinity for both cAMP and cGMP, although the $V_{\max}$ for cGMP hydrolysis is low enough that cGMP functions as a competitive inhibitor for cAMP hydrolysis. Enzymes in the PDE3 family are specifically inhibited by cGMP. PDE3 enzymes are specifically inhibited by milrinone and enoximone [Loughney and Ferguson, supra].

PDE4s are specific for cAMP hydrolysis. The family is comprised of four genes, PDE4A, PDE4B, PDE4C, and PDE4D. The genes give rise to multiple splice variants and are expressed in airway smooth muscle, the vascular endothelium, and all inflammatory cells. The enzymes can be activated by cAMP-dependent phosphorylation. Members of this family are specifically inhibited by the anti-depressant drug rolipram.

PDE5 is highly selective for cGMP [Turko, I.V. et al. (1998)]. Members of PDE5 family bind cGMP at non-catalytic sites [McAllister-Lucas, L.M. (1995)]. cGMP binding at non-catalytic sites has been suggested to be important for phosphorylation by cGMP-dependent protein kinase. PDE5 is highly expressed in vascular smooth muscle, platelets, lung, and kidney. Only one gene, PDE5A, has been identified.

PDE6s, the photoreceptor enzymes specifically hydrolyze cGMP [Loughney and Ferguson, supra]. PDE6s possess 2 regulatory high affinity cGMP binding sites. Genes include PDE6A and PDE6B (the protein products of which dimerize and bind two copies of a smaller γ inhibitory subunit to form rod PDE), in addition to PDE6C which associates with three smaller proteins to form cone PDE.

The PDE7 family effects cAMP hydrolysis but, in contrast to the PDE4 family, is not inhibited by rolipram [Loughney and Ferguson, supra]. Only one gene, PDE7A, has been identified. PDE7A gives rise to multiple splice variants. PDE7 mRNA can be found in several tissues but PDE7 protein expression appears to be restricted [Han, P.

- 4 -

et al. (1997); Perry, M.J. and G.A. Higgs (1998)]. Not much is known about the physiological function of PDE7.

5 The PDE8 family is closely related to the the PDE4 family. PDE8s have been shown to hydrolyze both cAMP and cGMP and are insensitive to inhibitors specific for PDEs 1-5. PDE8s are found in thyroid gland, testis, eye, liver, skeletal muscle, heart, kidney, ovary, and brain.

10 The PDE9 family preferentially hydrolyzes cAMP and is not sensitive to inhibition by rolipram, a PDE4-specific inhibitor, or isobutyl methyl xanthine (IBMX), a non-specific PDE inhibitor. PDE9 expression has been demonstrated in kidney, liver, lung, brain, spleen, and small intestine. Depending on nomenclature used, PDE9 is also referred to as PDE8, but is distinct from PDE8 mentioned above. To date, two genes have been identified in the PDE9 family.

15 PDE10 family members hydrolyze both cAMP and cGMP. PDE10s show expression in brain, thyroid and testis. [Soderling, S.H. et al. (1999); Fujishige, K. et al. (1999); Loughney, K. et al (1999)]

20 Members of the recently identified PDE11 family are also dualspecific. Interestingly, PDE11 splice variants exhibit different regulatory sequences in the N-terminal region. This suggests the possibility of differential regulation of PDE11s [Hetman JM, Robas N, Baxendale R, Fidock M, Phillips SC, Soderling SH, Beavo JA (2000)].

25 Increased PDE activity and decreased levels of cyclic nucleotides have been shown to be associated with many diseases. Furthermore, specific and non-specific inhibitors of several PDE protein families have been shown to be effective in treating such disorders. For example, the PDE4-specific inhibitor rolipram, mentioned above as an anti-depressant, inhibits lipopolysaccharide-induced expression of TNF- α , and
30 has been effective in treating multiple sclerosis in an animal model. Other PDE4-specific inhibitors are being investigated for use as anti-inflammatory therapeutics,

- 5 -

and efficacy in attenuating the late asthmatic response to allergen challenge has been demonstrated [Harbinson, et al. (1997)]. Inhibitors specific for the PDE3 family have been approved for treatment of congestive heart failure. PDE5 inhibitors such as Sildenafil are in use for treatment of penile erectile dysfunction [Terrett, N. et al. (1996)]. PDE5-inhibitors are under investigation as agents for cardiovascular therapy [Perry, M.J. and G.A. Higgs (1998)].

PDEs cyclic nucleotide levels have been suggested to influence proliferation of different cell types [Conti et al. (1995)]. For example, growth of the prostatic carcinoma cell lines DU145 and LNCaP was inhibited by cAMP derivatives and PDE inhibitors [Bang, Y.J. et al. (1994)]. Furthermore, PDEs have been implemented to additional cancers.

Non-specific inhibitors, such as theophylline and pentoxifylline, are currently used in the treatment of respiratory and vascular disorders, respectively.

In summary, cAMP and cGMP play a central role in intracellular second messenger signaling. Furthermore, the value as pharmaceutical targets has been proven for several PDEs. Selective inhibitors have been developed as therapeutic agents for diseases such as cancer, heart failure, depression and sexual dysfunction. Thus, the identification of further disease implications of PDE species and their splice variants may lead to the development of specific inhibitors or modulators, or suggest new utilities for known compounds affecting PDEs. That in turn will provide additional pharmacological approaches to treat diseases and conditions in which alterations in cyclic nucleotide pathways are involved. This diseases may include, but are not limited to, infections such as bacterial, fungal, protozoan, and viral infections, particularly those caused by HIV viruses, cancers, allergies including asthma, cardiovascular diseases including acute heart failure, hypotension, hypertension, angina pectoris, myocardial infarction, hematological diseases, genito-urinary diseases including urinary incontinence and benign prostate hyperplasia, osteo-

- 6 -

porosis, and peripheral and central nervous system disorders including pain, Alzheimer's disease and Parkinson's disease.

TaqMan-Technology / expression profiling

5

TaqMan is a recently developed technique, in which the release of a fluorescent reporter dye from a hybridisation probe in real-time during a polymerase chain reaction (PCR) is proportional to the accumulation of the PCR product. Quantification is based on the early, linear part of the reaction, and by determining the threshold cycle (CT), at which fluorescence above background is first detected.

10

Gene expression technologies may be useful in several areas of drug discovery and development, such as target identification, lead optimization, and identification of mechanisms of action. The TaqMan technology can be used to compare differences between expression profiles of normal tissue and diseased tissue. Expression profiling has been used in identifying genes, which are up- or downregulated in a variety of diseases. An interesting application of expression profiling is temporal monitoring of changes in gene expression during disease progression and drug treatment or in patients versus healthy individuals. The premise in this approach is that changes in pattern of gene expression in response to physiological or environmental stimuli (e.g., drugs) may serve as indirect clues about disease-causing genes or drug targets. Moreover, the effects of drugs with established efficacy on global gene expression patterns may provide a guidepost, or a genetic signature, against which a new drug candidate can be compared.

15

20

25

PDE4A

The nucleotide sequence of PDE4A is accessible in public databases by the accession number P27815 and is given in SEQ ID NO: 1. The amino acid sequence of PDE4A is depicted in SEQ ID NO: 2.

30

- 7 -

PDE4A is described as a phosphodiesterase [Livi GP et al., 1990]. The phosphodiesterase PDE4A is published in [Sullivan M et al., 1998], WO0179526 and US 5977305. The expression of PDE4A in medulla and brain was previously described [Lamontagne S et al., 2001]. PDE4A shows the highest homology to other phosphodiesterases as shown in example 1.

Summary of the invention

The invention relates to novel disease associations of PDE4A polypeptides and polynucleotides. The invention also relates to novel methods of screening for therapeutic agents for the treatment of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal. The invention also relates to pharmaceutical compositions for the treatment of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising a PDE4A polypeptide, a PDE4A polynucleotide, or regulators of PDE4A or modulators of PDE4A activity. The invention further comprises methods of diagnosing disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal.

Brief Description of the Drawings

Fig. 1 shows the nucleotide sequence of a PDE4A polynucleotide (SEQ ID NO: 1).
Fig. 2 shows the amino acid sequence of a PDE4A polypeptide (SEQ ID NO: 2).
Fig. 3 shows the nucleotide sequence of a primer useful for the invention (SEQ ID NO: 3).
Fig. 4 shows the nucleotide sequence of a primer useful for the invention (SEQ ID NO: 4).

- 8 -

Fig. 5 shows a nucleotide sequence useful as a probe to detect proteins of the invention (SEQ ID NO: 5).

Detailed description of the invention

5

Definition of terms

An "oligonucleotide" is a stretch of nucleotide residues which has a sufficient number of bases to be used as an oligomer, amplimer or probe in a polymerase chain reaction (PCR). Oligonucleotides are prepared from genomic or cDNA sequence and are used to amplify, reveal, or confirm the presence of a similar DNA or RNA in a particular cell or tissue. Oligonucleotides or oligomers comprise portions of a DNA sequence having at least about 10 nucleotides and as many as about 35 nucleotides, preferably about 25 nucleotides.

15

"Probes" may be derived from naturally occurring or recombinant single- or double-stranded nucleic acids or may be chemically synthesized. They are useful in detecting the presence of identical or similar sequences. Such probes may be labeled with reporter molecules using nick translation, Klenow fill-in reaction, PCR or other methods well known in the art. Nucleic acid probes may be used in southern, northern or in situ hybridizations to determine whether DNA or RNA encoding a certain protein is present in a cell type, tissue, or organ.

20

A "fragment of a polynucleotide" is a nucleic acid that comprises all or any part of a given nucleotide molecule, the fragment having fewer nucleotides than about 6 kb, preferably fewer than about 1 kb.

25

"Reporter molecules" are radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents which associate with a particular nucleotide or amino acid sequence, thereby establishing the presence of a certain sequence, or allowing for the quantification of a certain sequence.

30

“Chimeric” molecules may be constructed by introducing all or part of the nucleotide sequence of this invention into a vector containing additional nucleic acid sequence which might be expected to change any one or several of the following PDE4A characteristics: cellular location, distribution, ligand-binding affinities, interchain
5 affinities, degradation/turnover rate, signaling, etc.

“Active”, with respect to a PDE4A polypeptide, refers to those forms, fragments, or domains of a PDE4A polypeptide which retain the biological and/or antigenic
10 activity of a PDE4A polypeptide.

“Naturally occurring PDE4A polypeptide” refers to a polypeptide produced by cells which have not been genetically engineered and specifically contemplates various polypeptides arising from post-translational modifications of the polypeptide including but not limited to acetylation, carboxylation, glycosylation, phos-
15 phorylation, lipidation and acylation.

“Derivative” refers to polypeptides which have been chemically modified by techniques such as ubiquitination, labeling (see above), pegylation (derivatization with polyethylene glycol), and chemical insertion or substitution of amino acids such
20 as ornithine which do not normally occur in human proteins.

“Conservative amino acid substitutions” result from replacing one amino acid with another having similar structural and/or chemical properties, such as the replacement
25 of a leucine with an isoleucine or valine, an aspartate with a glutamate, or a threonine with a serine.

“Insertions” or “deletions” are typically in the range of about 1 to 5 amino acids. The variation allowed may be experimentally determined by producing the peptide synthetically while systematically making insertions, deletions, or substitutions of
30 nucleotides in the sequence using recombinant DNA techniques.

- 10 -

5 A "signal sequence" or "leader sequence" can be used, when desired, to direct the polypeptide through a membrane of a cell. Such a sequence may be naturally present on the polypeptides of the present invention or provided from heterologous sources by recombinant DNA techniques.

10 An "oligopeptide" is a short stretch of amino acid residues and may be expressed from an oligonucleotide. Oligopeptides comprise a stretch of amino acid residues of at least 3, 5, 10 amino acids and at most 10, 15, 25 amino acids, typically of at least 9 to 13 amino acids, and of sufficient length to display biological and/or antigenic activity.

15 "Inhibitor" is any substance which retards or prevents a chemical or physiological reaction or response. Common inhibitors include but are not limited to antisense molecules, antibodies, and antagonists.

20 "Standard expression" is a quantitative or qualitative measurement for comparison. It is based on a statistically appropriate number of normal samples and is created to use as a basis of comparison when performing diagnostic assays, running clinical trials, or following patient treatment profiles.

25 "Animal" as used herein may be defined to include human, domestic (e.g., cats, dogs, etc.), agricultural (e.g., cows, horses, sheep, etc.) or test species (e.g., mouse, rat, rabbit, etc.).

A "PDE4A polynucleotide", within the meaning of the invention, shall be understood as being a nucleic acid molecule selected from a group consisting of

30 (i) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2,

- 11 -

- (ii) nucleic acid molecules comprising the sequence of SEQ ID NO: 1,
- (iii) nucleic acid molecules having the sequence of SEQ ID NO: 1,
- 5 (iv) nucleic acid molecules the complementary strand of which hybridizes under stringent conditions to a nucleic acid molecule of (i), (ii), or (iii); and
- (v) nucleic acid molecules the sequence of which differs from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code;

10

wherein the polypeptide encoded by said nucleic acid molecule has PDE4A activity.

A "PDE4A polypeptide", within the meaning of the invention, shall be understood as being a polypeptide selected from a group consisting of

15

- (i) polypeptides having the sequence of SEQ ID NO: 2,
- (ii) polypeptides comprising the sequence of SEQ ID NO: 2,
- 20 (iii) polypeptides encoded by PDE4A polynucleotides; and
- (iv) polypeptides which show at least 99%, 98%, 95%, 90%, or 80% homology with a polypeptide of (i), (ii), or (iii);

25

wherein said polypeptide has PDE4A activity.

The nucleotide sequences encoding a PDE4A (or their complement) have numerous applications in techniques known to those skilled in the art of molecular biology. These techniques include use as hybridization probes, use in the construction of
30 oligomers for PCR, use for chromosome and gene mapping, use in the recombinant production of PDE4A, and use in generation of antisense DNA or RNA, their

- 12 -

chemical analogs and the like. Uses of nucleotides encoding a PDE4A disclosed herein are exemplary of known techniques and are not intended to limit their use in any technique known to a person of ordinary skill in the art. Furthermore, the nucleotide sequences disclosed herein may be used in molecular biology techniques that have not yet been developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, e.g., the triplet genetic code, specific base pair interactions, etc.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of PDE4A - encoding nucleotide sequences may be produced. Some of these will only bear minimal homology to the nucleotide sequence of the known and naturally occurring PDE4A. The invention has specifically contemplated each and every possible variation of nucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the nucleotide sequence of naturally occurring PDE4A, and all such variations are to be considered as being specifically disclosed.

Although the nucleotide sequences which encode a PDE4A, its derivatives or its variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring PDE4A polynucleotide under stringent conditions, it may be advantageous to produce nucleotide sequences encoding PDE4A polypeptides or its derivatives possessing a substantially different codon usage. Codons can be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic expression host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding a PDE4A polypeptide and/or its derivatives without altering the encoded amino acid sequence include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

- 13 -

Nucleotide sequences encoding a PDE4A polypeptide may be joined to a variety of other nucleotide sequences by means of well established recombinant DNA techniques. Useful nucleotide sequences for joining to PDE4A polynucleotides include an assortment of cloning vectors such as plasmids, cosmids, lambda phage derivatives, phagemids, and the like. Vectors of interest include expression vectors, replication vectors, probe generation vectors, sequencing vectors, etc. In general, vectors of interest may contain an origin of replication functional in at least one organism, convenient restriction endonuclease sensitive sites, and selectable markers for one or more host cell systems.

Another aspect of the subject invention is to provide for PDE4A-specific hybridization probes capable of hybridizing with naturally occurring nucleotide sequences encoding PDE4A. Such probes may also be used for the detection of similar PDE encoding sequences and should preferably show at least 40% nucleotide identity to PDE4A polynucleotides. The hybridization probes of the subject invention may be derived from the nucleotide sequence presented as SEQ ID NO: 1 or from genomic sequences including promoter, enhancers or introns of the native gene. Hybridization probes may be labelled by a variety of reporter molecules using techniques well known in the art.

It will be recognized that many deletional or mutational analogs of PDE4A polynucleotides will be effective hybridization probes for PDE4A polynucleotides. Accordingly, the invention relates to nucleic acid sequences that hybridize with such PDE4A encoding nucleic acid sequences under stringent conditions.

“Stringent conditions” refers to conditions that allow for the hybridization of substantially related nucleic acid sequences. For instance, such conditions will generally allow hybridization of sequence with at least about 85% sequence identity, preferably with at least about 90% sequence identity, more preferably with at least about 95% sequence identity. Hybridization conditions and probes can be adjusted in well-characterized ways to achieve selective hybridization of human-derived

- 14 -

probes. Stringent conditions, within the meaning of the invention are 65°C in a buffer containing 1 mM EDTA, 0.5 M NaHPO₄ (pH 7.2), 7 % (w/v) SDS.

5 Nucleic acid molecules that will hybridize to PDE4A polynucleotides under stringent conditions can be identified functionally. Without limitation, examples of the uses for hybridization probes include: histochemical uses such as identifying tissues that express PDE4A; measuring mRNA levels, for instance to identify a sample's tissue type or to identify cells that express abnormal levels of PDE4A; and detecting polymorphisms of PDE4A.

10

PCR provides additional uses for oligonucleotides based upon the nucleotide sequence which encodes PDE4A. Such probes used in PCR may be of recombinant origin, chemically synthesized, or a mixture of both. Oligomers may comprise discrete nucleotide sequences employed under optimized conditions for identification
15 of PDE4A in specific tissues or diagnostic use. The same two oligomers, a nested set of oligomers, or even a degenerate pool of oligomers may be employed under less stringent conditions for identification of closely related DNAs or RNAs.

20

Rules for designing polymerase chain reaction (PCR) primers are now established, as reviewed by PCR Protocols. Degenerate primers, i.e., preparations of primers that are heterogeneous at given sequence locations, can be designed to amplify nucleic acid sequences that are highly homologous to, but not identical with PDE4A. Strategies are now available that allow for only one of the primers to be required to specifically hybridize with a known sequence. For example, appropriate nucleic acid
25 primers can be ligated to the nucleic acid sought to be amplified to provide the hybridization partner for one of the primers. In this way, only one of the primers need be based on the sequence of the nucleic acid sought to be amplified.

30

PCR methods for amplifying nucleic acid will utilize at least two primers. One of these primers will be capable of hybridizing to a first strand of the nucleic acid to be amplified and of priming enzyme-driven nucleic acid synthesis in a first direction.

- 15 -

The other will be capable of hybridizing the reciprocal sequence of the first strand (if the sequence to be amplified is single stranded, this sequence will initially be hypothetical, but will be synthesized in the first amplification cycle) and of priming nucleic acid synthesis from that strand in the direction opposite the first direction and
5 towards the site of hybridization for the first primer. Conditions for conducting such amplifications, particularly under preferred stringent hybridization conditions, are well known.

Other means of producing specific hybridization probes for PDE4A include the
10 cloning of nucleic acid sequences encoding PDE4A or PDE4A derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerase as T7 or SP6 RNA polymerase and the appropriate reporter molecules.

15 It is possible to produce a DNA sequence, or portions thereof, entirely by synthetic chemistry. After synthesis, the nucleic acid sequence can be inserted into any of the many available DNA vectors and their respective host cells using techniques which are well known in the art. Moreover, synthetic chemistry may be used to introduce
20 mutations into the nucleotide sequence. Alternately, a portion of sequence in which a mutation is desired can be synthesized and recombined with longer portion of an existing genomic or recombinant sequence.

PDE4A polynucleotides may be used to produce a purified oligo-or polypeptide
25 using well known methods of recombinant DNA technology. The oligopeptide may be expressed in a variety of host cells, either prokaryotic or eukaryotic. Host cells may be from the same species from which the nucleotide sequence was derived or from a different species. Advantages of producing an oligonucleotide by recombinant DNA technology include obtaining adequate amounts of the protein for
30 purification and the availability of simplified purification procedures.

Quantitative determinations of nucleic acids

An important step in the molecular genetic analysis of human disease is often the enumeration of the copy number of a nucleic acid or the relative expression of a gene in particular tissues.

Several different approaches are currently available to make quantitative determinations of nucleic acids. Chromosome-based techniques, such as comparative genomic hybridization (CGH) and fluorescent in situ hybridization (FISH) facilitate efforts to cytogenetically localize genomic regions that are altered in tumor cells. Regions of genomic alteration can be narrowed further using loss of heterozygosity analysis (LOH), in which disease DNA is analyzed and compared with normal DNA for the loss of a heterozygous polymorphic marker. The first experiments used restriction fragment length polymorphisms (RFLPs) [Johnson, (1989)], or hypervariable minisatellite DNA [Barnes, 2000]. In recent years LOH has been performed primarily using PCR amplification of microsatellite markers and electrophoresis of the radio labelled [Jeffreys, (1985)] or fluorescently labelled PCR products [Weber, (1990)] and compared between paired normal and disease DNAs.

A number of other methods have also been developed to quantify nucleic acids [Gergen, (1992)]. More recently, PCR and RT-PCR methods have been developed which are capable of measuring the amount of a nucleic acid in a sample. One approach, for example, measures PCR product quantity in the log phase of the reaction before the formation of reaction products plateaus [Thomas, (1980)].

A gene sequence contained in all samples at relatively constant quantity is typically utilized for sample amplification efficiency normalization. This approach, however, suffers from several drawbacks. The method requires that each sample has equal input amounts of the nucleic acid and that the amplification efficiency between samples is identical until the time of analysis. Furthermore, it is difficult using the conventional methods of PCR quantitation such as gel electrophoresis or plate

- 17 -

capture hybridization to determine that all samples are in fact analyzed during the log phase of the reaction as required by the method.

Another method called quantitative competitive (QC)-PCR, as the name implies, relies on the inclusion of an internal control competitor in each reaction [Piatak, (1993), BioTechniques]. The efficiency of each reaction is normalized to the internal competitor. A known amount of internal competitor is typically added to each sample. The unknown target PCR product is compared with the known competitor PCR product to obtain relative quantitation. A difficulty with this general approach lies in developing an internal control that amplifies with the same efficiency than the target molecule.

5' Fluorogenic Nuclease Assays

Fluorogenic nuclease assays are a real time quantitation method that uses a probe to monitor formation of amplification product. The basis for this method of monitoring the formation of amplification product is to measure continuously PCR product accumulation using a dual-labelled fluorogenic oligonucleotide probe, an approach frequently referred to in the literature simply as the "TaqMan method" [Piatak, (1993), Science; Heid, (1996); Gibson, (1996); Holland, (1991)].

The probe used in such assays is typically a short (about 20-25 bases) oligonucleotide that is labeled with two different fluorescent dyes. The 5' terminus of the probe is attached to a reporter dye and the 3' terminus is attached to a quenching dye, although the dyes could be attached at other locations on the probe as well. The probe is designed to have at least substantial sequence complementarity with the probe binding site. Upstream and downstream PCR primers which bind to flanking regions of the locus are added to the reaction mixture. When the probe is intact, energy transfer between the two fluorophors occurs and the quencher quenches emission from the reporter. During the extension phase of PCR, the probe is cleaved by the 5' nuclease activity of a nucleic acid polymerase such as Taq polymerase,

- 18 -

thereby releasing the reporter from the oligonucleotide-quencher and resulting in an increase of reporter emission intensity which can be measured by an appropriate detector.

5 One detector which is specifically adapted for measuring fluorescence emissions such as those created during a fluorogenic assay is the ABI 7700 or 4700 HT manufactured by Applied Biosystems, Inc. in Foster City, Calif. The ABI 7700 uses fiber optics connected with each well in a 96-or 384 well PCR tube arrangement. The instrument includes a laser for exciting the labels and is capable of measuring
10 the fluorescence spectra intensity from each tube with continuous monitoring during PCR amplification. Each tube is re-examined every 8.5 seconds.

Computer software provided with the instrument is capable of recording the fluorescence intensity of reporter and quencher over the course of the amplification.
15 The recorded values will then be used to calculate the increase in normalized reporter emission intensity on a continuous basis. The increase in emission intensity is plotted versus time, i.e., the number of amplification cycles, to produce a continuous measure of amplification. To quantify the locus in each amplification reaction, the amplification plot is examined at a point during the log phase of product
20 accumulation. This is accomplished by assigning a fluorescence threshold intensity above background and determining the point at which each amplification plot crosses the threshold (defined as the threshold cycle number or Ct). Differences in threshold cycle number are used to quantify the relative amount of PCR target contained within each tube. Assuming that each reaction functions at 100% PCR efficiency, a
25 difference of one Ct represents a two-fold difference in the amount of starting template. The fluorescence value can be used in conjunction with a standard curve to determine the amount of amplification product present.

Non-Probe-Based Detection Methods

A variety of options are available for measuring the amplification products as they are formed. One method utilizes labels, such as dyes, which only bind to double stranded DNA. In this type of approach, amplification product (which is double stranded) binds dye molecules in solution to form a complex. With the appropriate dyes, it is possible to distinguish between dye molecules free in solution and dye molecules bound to amplification product. For example, certain dyes fluoresce only when bound to amplification product. Examples of dyes which can be used in methods of this general type include, but are not limited to, Syber Green.TM. and Pico Green from Molecular Probes, Inc. of Eugene, Oreg., ethidium bromide, propidium iodide, chromomycin, acridine orange, Hoechst 33258, Toto-1, Yoyo-1, DAPI (4',6-diamidino-2-phenylindole hydrochloride).

Another real time detection technique measures alteration in energy fluorescence energy transfer between fluorophors conjugated with PCR primers [Livak, (1995)].

Probe-Based Detection Methods

These detection methods involve some alteration to the structure or conformation of a probe hybridized to the locus between the amplification primer pair. In some instances, the alteration is caused by the template-dependent extension catalyzed by a nucleic acid polymerase during the amplification process. The alteration generates a detectable signal which is an indirect measure of the amount of amplification product formed.

For example, some methods involve the degradation or digestion of the probe during the extension reaction. These methods are a consequence of the 5'-3' nuclease activity associated with some nucleic acid polymerases. Polymerases having this activity cleave mononucleotides or small oligonucleotides from an oligonucleotide probe annealed to its complementary sequence located within the locus.

- 20 -

The 3' end of the upstream primer provides the initial binding site for the nucleic acid polymerase. As the polymerase catalyzes extension of the upstream primer and encounters the bound probe, the nucleic acid polymerase displaces a portion of the 5' end of the probe and through its nuclease activity cleaves mononucleotides or oligonucleotides from the probe.

The upstream primer and the probe can be designed such that they anneal to the complementary strand in close proximity to one another. In fact, the 3' end of the upstream primer and the 5' end of the probe may abut one another. In this situation, extension of the upstream primer is not necessary in order for the nucleic acid polymerase to begin cleaving the probe. In the case in which intervening nucleotides separate the upstream primer and the probe, extension of the primer is necessary before the nucleic acid polymerase encounters the 5' end of the probe. Once contact occurs and polymerization continues, the 5'-3' exonuclease activity of the nucleic acid polymerase begins cleaving mononucleotides or oligonucleotides from the 5' end of the probe. Digestion of the probe continues until the remaining portion of the probe dissociates from the complementary strand.

In solution, the two end sections can hybridize with each other to form a hairpin loop. In this conformation, the reporter and quencher dye are in sufficiently close proximity that fluorescence from the reporter dye is effectively quenched by the quencher dye. Hybridized probe, in contrast, results in a linearized conformation in which the extent of quenching is decreased. Thus, by monitoring emission changes for the two dyes, it is possible to indirectly monitor the formation of amplification product.

Probes

The labeled probe is selected so that its sequence is substantially complementary to a segment of the test locus or a reference locus. As indicated above, the nucleic acid

site to which the probe binds should be located between the primer binding sites for the upstream and downstream amplification primers.

Primers

5

The primers used in the amplification are selected so as to be capable of hybridizing to sequences at flanking regions of the locus being amplified. The primers are chosen to have at least substantial complementarity with the different strands of the nucleic acid being amplified. When a probe is utilized to detect the formation of amplification products, the primers are selected in such that they flank the probe, i.e. are located upstream and downstream of the probe.

10

The primer must have sufficient length so that it is capable of priming the synthesis of extension products in the presence of an agent for polymerization. The length and composition of the primer depends on many parameters, including, for example, the temperature at which the annealing reaction is conducted, proximity of the probe binding site to that of the primer, relative concentrations of the primer and probe and the particular nucleic acid composition of the probe. Typically the primer includes 15-30 nucleotides. However, the length of the primer may be more or less depending on the complexity of the primer binding site and the factors listed above.

15

20

Labels for Probes and Primers

The labels used for labeling the probes or primers of the current invention and which can provide the signal corresponding to the quantity of amplification product can take a variety of forms. As indicated above with regard to the 5' fluorogenic nuclease method, a fluorescent signal is one signal which can be measured. However, measurements may also be made, for example, by monitoring radioactivity, colorimetry, absorption, magnetic parameters, or enzymatic activity. Thus, labels which can be employed include, but are not limited to, fluorophores, chromophores,

25

30

- 22 -

radioactive isotopes, electron dense reagents, enzymes, and ligands having specific binding partners (e.g., biotin-avidin).

Monitoring changes in fluorescence is a particularly useful way to monitor the accumulation of amplification products. A number of labels useful for attachment to probes or primers are commercially available including fluorescein and various fluorescein derivatives such as FAM, HEX, TET and JOE (all which are available from Applied Biosystems, Foster City, Calif.); lucifer yellow, and coumarin derivatives.

Labels may be attached to the probe or primer using a variety of techniques and can be attached at the 5' end, and/or the 3' end and/or at an internal nucleotide. The label can also be attached to spacer arms of various sizes which are attached to the probe or primer. These spacer arms are useful for obtaining a desired distance between multiple labels attached to the probe or primer.

In some instances, a single label may be utilized; whereas, in other instances, such as with the 5' fluorogenic nuclease assays for example, two or more labels are attached to the probe. In cases wherein the probe includes multiple labels, it is generally advisable to maintain spacing between the labels which is sufficient to permit separation of the labels during digestion of the probe through the 5'-3' nuclease activity of the nucleic acid polymerase.

Patients Exhibiting Symptoms of Disease

A number of diseases are associated with changes in the copy number of a certain gene. For patients having symptoms of a disease, the real-time PCR method can be used to determine if the patient has copy number alterations which are known to be linked with diseases that are associated with the symptoms the patient has.

PDE4A expression*PDE4A fusion proteins*

5 Fusion proteins are useful for generating antibodies against PDE4A polypeptides and for use in various assay systems. For example, fusion proteins can be used to identify proteins which interact with portions of PDE4A polypeptides. Protein affinity chromatography or library-based assays for protein-protein interactions, such as the yeast two-hybrid or phage display systems, can be used for this purpose. Such
10 methods are well known in the art and also can be used as drug screens.

A PDE4A fusion protein comprises two polypeptide segments fused together by means of a peptide bond. The first polypeptide segment can comprise at least 54, 75, 100, 125, 139, 150, 175, 200, 225, 250, 275, 300, 325 or 350 contiguous amino acids
15 of SEQ ID NO: 2 or of a biologically active variant, such as those described above. The first polypeptide segment also can comprise full-length PDE4A.

The second polypeptide segment can be a full-length protein or a protein fragment. Proteins commonly used in fusion protein construction include, but are not limited to
20 β -galactosidase, β -glucuronidase, green fluorescent protein (GFP), autofluorescent proteins, including blue fluorescent protein (BFP), glutathione-S-transferase (GST), luciferase, horseradish peroxidase (HRP), and chloramphenicol acetyltransferase (CAT). Additionally, epitope tags are used in fusion protein constructions, including histidine (His) tags, FLAG tags, influenza hemagglutinin (HA) tags, Myc tags, VSV-
25 G tags, and thioredoxin (Trx) tags. Other fusion constructions can include maltose binding protein (MBP), S-tag, Lex a DNA binding domain (DBD) fusions, GAL4 DNA binding domain fusions, and herpes simplex virus (HSV) BP16 protein fusions. A fusion protein also can be engineered to contain a cleavage site located adjacent to the PDE4A.

Preparation of Polynucleotides

A naturally occurring PDE4A polynucleotide can be isolated free of other cellular components such as membrane components, proteins, and lipids. Polynucleotides
5 can be made by a cell and isolated using standard nucleic acid purification techniques, or synthesized using an amplification technique, such as the polymerase chain reaction (PCR), or by using an automatic synthesizer. Methods for isolating polynucleotides are routine and are known in the art. Any such technique for obtaining a polynucleotide can be used to obtain isolated PDE4A polynucleotides.
10 For example, restriction enzymes and probes can be used to isolate polynucleotide fragments which comprise PDE4A nucleotide sequences. Isolated polynucleotides are in preparations which are free or at least 70, 80, or 90% free of other molecules.

PDE4A cDNA molecules can be made with standard molecular biology techniques,
15 using PDE4A mRNA as a template. PDE4A cDNA molecules can thereafter be replicated using molecular biology techniques known in the art. An amplification technique, such as PCR, can be used to obtain additional copies of polynucleotides of the invention, using either human genomic DNA or cDNA as a template.

20 Alternatively, synthetic chemistry techniques can be used to synthesize PDE4A polynucleotides. The degeneracy of the genetic code allows alternate nucleotide sequences to be synthesized which will encode PDE4A having, for example, an amino acid sequence shown in SEQ ID NO: 2 or a biologically active variant thereof.

Extending Polynucleotides

Various PCR-based methods can be used to extend nucleic acid sequences encoding human PDE4A, for example to detect upstream sequences of PDE4A gene such as promoters and regulatory elements. For example, restriction-site PCR uses universal
30 primers to retrieve unknown sequence adjacent to a known locus. Genomic DNA is first amplified in the presence of a primer to a linker sequence and a primer specific

to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

5

Inverse PCR also can be used to amplify or extend sequences using divergent primers based on a known region. Primers can be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, Minn.), to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68-72°C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

10

Another method which can be used is capture PCR, which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA. In this method, multiple restriction enzyme digestions and ligations also can be used to place an engineered double-stranded sequence into an unknown fragment of the DNA molecule before performing PCR.

20

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Randomly-primed libraries are preferable, in that they will contain more sequences which contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries can be useful for extension of sequence into 5' non-transcribed regulatory regions.

25

Commercially available capillary electrophoresis systems can be used to analyze the size or confirm the nucleotide sequence of PCR or sequencing products. For example, capillary sequencing can employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser

30

- 26 -

activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity can be converted to electrical signal using appropriate equipment and software (e.g., GENOTYPER and Sequence NAVIGATOR, Perkin Elmer), and the entire process from loading of samples to computer analysis and electronic data display can be computer controlled. Capillary electrophoresis is especially preferable for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

Obtaining Polypeptides

PDE4A can be obtained, for example, by purification from human cells, by expression of PDE4A polynucleotides, or by direct chemical synthesis.

Protein Purification

PDE4A can be purified from any human cell which expresses the enzyme, including those which have been transfected with expression constructs which express PDE4A. A purified PDE4A is separated from other compounds which normally associate with PDE4A in the cell, such as certain proteins, carbohydrates, or lipids, using methods well-known in the art. Such methods include, but are not limited to, size exclusion chromatography, ammonium sulfate fractionation, ion exchange chromatography, affinity chromatography, and preparative gel electrophoresis.

Expression of PDE4A Polynucleotides

To express PDE4A, PDE4A polynucleotides can be inserted into an expression vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art can be used to construct expression vectors containing sequences encoding PDE4A and appropriate transcriptional and translational control elements. These

- 27 -

methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination.

5 A variety of expression vector/host systems can be utilized to contain and express sequences encoding PDE4A. These include, but are not limited to, microorganisms, such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors, insect cell systems infected with virus expression vectors (*e.g.*, baculovirus), plant cell systems transformed with virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (*e.g.*, Ti or pBR322 plasmids), or animal cell systems.

15 The control elements or regulatory sequences are those non-translated regions of the vector - enhancers, promoters, 5' and 3' untranslated regions -- which interact with host cellular proteins to carry out transcription and translation. Such elements can vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, can be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the BLUESCRIPT phagemid (Stratagene, LaJolla, Calif.) or pSPORT1 plasmid (Life Technologies) and the like can be used. The baculovirus polyhedrin promoter can be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (*e.g.*, heat shock, RUBISCO, and storage protein genes) or from plant viruses (*e.g.*, viral promoters or leader sequences) can be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of a nucleotide sequence encoding PDE4A, vectors based on SV40 or EBV can be used with an appropriate selectable marker.

Bacterial and Yeast Expression Systems

In bacterial systems, a number of expression vectors can be selected. For example, when a large quantity of PDE4A is needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified can be used. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene). In a BLUESCRIPT vector, a sequence encoding PDE4A can be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced. pIN vectors or pGEX vectors (Promega, Madison, Wis.) also can be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems can be designed to include heparin, thrombin, or factor Xa protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

Plant and Insect Expression Systems

If plant expression vectors are used, the expression of sequences encoding PDE4A can be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV can be used alone or in combination with the omega leader sequence from TMV. Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters can be used. These constructs can be introduced into plant cells by direct DNA transformation or by pathogen-mediated transfection.

An insect system also can be used to express PDE4A. For example, in one such system *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia*

- 29 -

larvae. Sequences encoding PDE4A can be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of PDE4A will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses can then be used to infect *S. frugiperda* cells or *Trichoplusia* larvae in which PDE4A can be expressed.

Mammalian Expression Systems

A number of viral-based expression systems can be used to express PDE4A in mammalian host cells. For example, if an adenovirus is used as an expression vector, sequences encoding PDE4A can be ligated into an adenovirus transcription/-translation complex comprising the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome can be used to obtain a viable virus which is capable of expressing PDE4A in infected host cells [Engelhard, 1994)]. If desired, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, can be used to increase expression in mammalian host cells.

Human artificial chromosomes (HACs) also can be used to deliver larger fragments of DNA than can be contained and expressed in a plasmid. HACs of 6M to 10M are constructed and delivered to cells via conventional delivery methods (*e.g.*, liposomes, polycationic amino polymers, or vesicles). Specific initiation signals also can be used to achieve more efficient translation of sequences encoding PDE4A. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding PDE4A, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals (including the ATG initiation codon) should be provided. The initiation codon should be in the correct reading frame to ensure translation of the entire insert.

Exogenous translational elements and initiation codons can be of various origins, both natural and synthetic.

Host Cells

5

A host cell strain can be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed PDE4A in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the polypeptide also can be used to facilitate correct insertion, folding and/or function. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC; 10801 University Boulevard, Manassas, VA 20110-2209) and can be chosen to ensure the correct modification and processing of the foreign protein.

10
15

20

25

30

Stable expression is preferred for long-term, high-yield production of recombinant proteins. For example, cell lines which stably express PDE4A can be transformed using expression vectors which can contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells can be allowed to grow for 1-2 days in an enriched medium before they are switched to a selective medium. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced PDE4A sequences. Resistant clones of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type. Any number of selection systems can be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase [Logan, (1984)] and adenine phosphoribosyltransferase [Wigler, (1977)] genes which can be employed in *tk* or *aprf* cells, respectively. Also, antimetabolite, antibiotic, or herbicide resistance

- 31 -

can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate [Lowy, (1980)], *npt* confers resistance to the aminoglycosides, neomycin and G-418 [Wigler, (1980)], and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively [Colbere-Garapin, 1981].

5 Additional selectable genes have been described. For example, *trpB* allows cells to utilize indole in place of tryptophan, or *hisD*, which allows cells to utilize histinol in place of histidine. Visible markers such as anthocyanins, β -glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, can be used to identify transformants and to quantify the amount of transient or stable protein expression

10 attributable to a specific vector system

Detecting Polypeptide Expression

Although the presence of marker gene expression suggests that a PDE4A polynucleotide is also present, its presence and expression may need to be confirmed.

15 For example, if a sequence encoding PDE4A is inserted within a marker gene sequence, transformed cells containing sequences which encode PDE4A can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding PDE4A under the control of a single

20 promoter. Expression of the marker gene in response to induction or selection usually indicates expression of PDE4A polynucleotide.

Alternatively, host cells which contain a PDE4A polynucleotide and which express PDE4A can be identified by a variety of procedures known to those of skill in the art.

25 These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip-based technologies for the detection and/or quantification of nucleic acid or protein. For example, the presence of a polynucleotide sequence encoding PDE4A can be detected by DNA-DNA or DNA-RNA

30 hybridization or amplification using probes or fragments or fragments of polynucleotides encoding PDE4A. Nucleic acid amplification-based assays involve the

- 32 -

use of oligonucleotides selected from sequences encoding PDE4A to detect transformants which contain a PDE4A polynucleotide.

5 A variety of protocols for detecting and measuring the expression of PDE4A, using either polyclonal or monoclonal antibodies specific for the polypeptide, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radio-immunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay using monoclonal antibodies reactive to two non-interfering epitopes on PDE4A can be used, or a competitive binding assay can be
10 employed.

A wide variety of labels and conjugation techniques are known by those skilled in the art and can be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to
15 polynucleotides encoding PDE4A include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, sequences encoding PDE4A can be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and can be used to synthesize RNA probes *in vitro* by addition of labeled nucleotides and an appropriate
20 RNA polymerase such as T7, T3, or SP6. These procedures can be conducted using a variety of commercially available kits (Amersham Pharmacia Biotech, Promega, and US Biochemical). Suitable reporter molecules or labels which can be used for ease of detection include radionuclides, enzymes, and fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles,
25 and the like.

Expression and Purification of Polypeptides

Host cells transformed with PDE4A polynucleotides can be cultured under
30 conditions suitable for the expression and recovery of the protein from cell culture. The polypeptide produced by a transformed cell can be secreted or contained

- 33 -

intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing PDE4A polynucleotides can be designed to contain signal sequences which direct secretion of soluble PDE4A through a prokaryotic or eukaryotic cell membrane or which direct the membrane insertion of membrane-bound PDE4A.

As discussed above, other constructions can be used to join a sequence encoding PDE4A to a nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). Inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, CA) between the purification domain and PDE4A also can be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing PDE4A and 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification by IMAC (immobilized metal ion affinity chromatography) Maddox, (1983)], while the enterokinase cleavage site provides a means for purifying PDE4A from the fusion protein [Porath, (1992)].

Chemical Synthesis

Sequences encoding PDE4A can be synthesized, in whole or in part, using chemical methods well known in the art. Alternatively, PDE4A itself can be produced using chemical methods to synthesize its amino acid sequence, such as by direct peptide synthesis using solid-phase techniques. Protein synthesis can either be performed using manual techniques or by automation. Automated synthesis can be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer).

- 34 -

Optionally, fragments of PDE4A can be separately synthesized and combined using chemical methods to produce a full-length molecule.

5 The newly synthesized peptide can be substantially purified by preparative high performance liquid chromatography. The composition of a synthetic PDE4A can be confirmed by amino acid analysis or sequencing. Additionally, any portion of the amino acid sequence of PDE4A can be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins to produce a variant polypeptide or a fusion protein.

10

Production of Altered Polypeptides

As will be understood by those of skill in the art, it may be advantageous to produce PDE4A polynucleotides possessing non-naturally occurring codons. For example, 15 codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce an RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

20

The nucleotide sequences referred to herein can be engineered using methods generally known in the art to alter PDE4A polynucleotides for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the polypeptide or mRNA product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides 25 can be used to engineer the nucleotide sequences. For example, site-directed mutagenesis can be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, introduce mutations, and so forth.

Antibodies

Any type of antibody known in the art can be generated to bind specifically to an epitope of PDE4A.

5

“Antibody” as used herein includes intact immunoglobulin molecules, as well as fragments thereof, such as Fab, F(ab')₂, and Fv, which are capable of binding an epitope of PDE4A. Typically, at least 6, 8, 10, or 12 contiguous amino acids are required to form an epitope. However, epitopes which involve non-contiguous amino acids may require more, *e.g.*, at least 15, 25, or 50 amino acid. An antibody which specifically binds to an epitope of PDE4A can be used therapeutically, as well as in immunochemical assays, such as Western blots, ELISAs, radioimmunoassays, immunohistochemical assays, immunoprecipitations, or other immunochemical assays known in the art. Various immunoassays can be used to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays are well known in the art. Such immunoassays typically involve the measurement of complex formation between an immunogen and an antibody which specifically binds to the PDE4A immunogen.

10

15

20

Typically, an antibody which specifically binds to PDE4A provides a detection signal at least 5-, 10-, or 20-fold higher than a detection signal provided with other proteins when used in an immunochemical assay. Preferably, antibodies which specifically bind to PDE4A do not detect other proteins in immunochemical assays and can immunoprecipitate PDE4A from solution.

25

30

PDE4A can be used to immunize a mammal, such as a mouse, rat, rabbit, guinea pig, monkey, or human, to produce polyclonal antibodies. If desired, PDE4A can be conjugated to a carrier protein, such as bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin. Depending on the host species, various adjuvants can be used to increase the immunological response. Such adjuvants include, but are not limited to, Freund's adjuvant, mineral gels (*e.g.*, aluminum hydroxide), and surface

- 36 -

active substances (e.g., lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol). Among adjuvants used in humans, BCG (*bacilli Calmette-Guerin*) and *Corynebacterium parvum* are especially useful.

5

Monoclonal antibodies which specifically bind to PDE4A can be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These techniques include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique [Roberge, (1995)].

10

In addition, techniques developed for the production of "chimeric antibodies", the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. Monoclonal and other antibodies also can be "humanized" to prevent a patient from mounting an immune response against the antibody when it is used therapeutically. Such antibodies may be sufficiently similar in sequence to human antibodies to be used directly in therapy or may require alteration of a few key residues. Sequence differences between rodent antibodies and human sequences can be minimized by replacing residues which differ from those in the human sequences by site directed mutagenesis of individual residues or by grating of entire complementarity determining regions. Antibodies which specifically bind to PDE4A can contain antigen binding sites which are either partially or fully humanized, as disclosed in U.S. 5,565,332.

15

20

25

Alternatively, techniques described for the production of single chain antibodies can be adapted using methods known in the art to produce single chain antibodies which specifically bind to PDE4A. Antibodies with related specificity, but of distinct idiotypic composition, can be generated by chain shuffling from random combinatorial immunoglobulin libraries. Single-chain antibodies also can be constructed using a DNA amplification method, such as PCR, using hybridoma cDNA as a template.

30

- 37 -

Single-chain antibodies can be mono- or bispecific, and can be bivalent or tetravalent. Construction of tetravalent, bispecific single-chain antibodies is taught. A nucleotide sequence encoding a single-chain antibody can be constructed using manual or automated nucleotide synthesis, cloned into an expression construct using standard recombinant DNA methods, and introduced into a cell to express the coding sequence, as described below. Alternatively, single-chain antibodies can be produced directly using, for example, filamentous phage technology.

Antibodies which specifically bind to PDE4A also can be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents. Other types of antibodies can be constructed and used therapeutically in methods of the invention. For example, chimeric antibodies can be constructed as disclosed in WO 93/03151. Binding proteins which are derived from immunoglobulins and which are multivalent and multispecific, such as the "diabodies" described in WO 94/13804, also can be prepared.

Antibodies according to the invention can be purified by methods well known in the art. For example, antibodies can be affinity purified by passage over a column to which PDE4A is bound. The bound antibodies can then be eluted from the column using a buffer with a high salt concentration.

Antisense Oligonucleotides

Antisense oligonucleotides are nucleotide sequences which are complementary to a specific DNA or RNA sequence. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form complexes and block either transcription or translation. Preferably, an antisense oligonucleotide is at least 11 nucleotides in length, but can be at least 12, 15, 20, 25, 30, 35, 40, 45, or 50 or more nucleotides long. Longer sequences also can be used. Antisense

oligonucleotide molecules can be provided in a DNA construct and introduced into a cell as described above to decrease the level of PDE4A gene products in the cell.

Antisense oligonucleotides can be deoxyribonucleotides, ribonucleotides, or a combination of both. Oligonucleotides can be synthesized manually or by an automated synthesizer, by covalently linking the 5' end of one nucleotide with the 3' end of another nucleotide with non-phosphodiester internucleotide linkages such as alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, alkylphosphonates, phosphoramidates, phosphate esters, carbamates, acetamidate, carboxymethyl esters, carbonates, and phosphate triesters.

Modifications of PDE4A gene expression can be obtained by designing antisense oligonucleotides which will form duplexes to the control, 5', or regulatory regions of the PDE4A gene. Oligonucleotides derived from the transcription initiation site, *e.g.*, between positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or chaperons. Therapeutic advances using triplex DNA have been described in the literature [Nicholls, (1993)]. An antisense oligonucleotide also can be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Precise complementarity is not required for successful complex formation between an antisense oligonucleotide and the complementary sequence of a PDE4A polynucleotide. Antisense oligonucleotides which comprise, for example, 2, 3, 4, or 5 or more stretches of contiguous nucleotides which are precisely complementary to a PDE4A polynucleotide, each separated by a stretch of contiguous nucleotides which are not complementary to adjacent PDE4A nucleotides, can provide sufficient targeting specificity for PDE4A mRNA. Preferably, each stretch of complementary contiguous nucleotides is at least 4, 5, 6, 7, or 8 or more nucleotides in length. Non-complementary intervening sequences are preferably 1, 2, 3, or 4 nucleotides in

- 39 -

length. One skilled in the art can easily use the calculated melting point of an antisense-sense pair to determine the degree of mismatching which will be tolerated between a particular antisense oligonucleotide and a particular PDE4A polynucleotide sequence. Antisense oligonucleotides can be modified without affecting their ability to hybridize to a PDE4A polynucleotide. These modifications can be internal or at one or both ends of the antisense molecule. For example, internucleoside phosphate linkages can be modified by adding cholesteryl or diamine moieties with varying numbers of carbon residues between the amino groups and terminal ribose. Modified bases and/or sugars, such as arabinose instead of ribose, or a 3', 5'-substituted oligonucleotide in which the 3' hydroxyl group or the 5' phosphate group are substituted, also can be employed in a modified antisense oligonucleotide. These modified oligonucleotides can be prepared by methods well known in the art.

Ribozymes

Ribozymes are RNA molecules with catalytic activity [Uhlmann, (1987)]. Ribozymes can be used to inhibit gene function by cleaving an RNA sequence, as is known in the art. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Examples include engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of specific nucleotide sequences. The coding sequence of a PDE4A polynucleotide can be used to generate ribozymes which will specifically bind to mRNA transcribed from a PDE4A polynucleotide. Methods of designing and constructing ribozymes which can cleave other RNA molecules in trans in a highly sequence specific manner have been developed and described in the art. For example, the cleavage activity of ribozymes can be targeted to specific RNAs by engineering a discrete "hybridization" region into the ribozyme. The hybridization region contains a sequence complementary to the target RNA and thus specifically hybridizes with the target RNA.

- 40 -

Specific ribozyme cleavage sites within a PDE4A RNA target can be identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target RNA
5 containing the cleavage site can be evaluated for secondary structural features which may render the target inoperable. Suitability of candidate PDE4A RNA targets also can be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays. The nucleotide sequences shown in SEQ ID NO: 1 and its complement provide sources of suitable
10 hybridization region sequences. Longer complementary sequences can be used to increase the affinity of the hybridization sequence for the target. The hybridizing and cleavage regions of the ribozyme can be integrally related such that upon hybridizing to the target RNA through the complementary regions, the catalytic region of the ribozyme can cleave the target.

15 Ribozymes can be introduced into cells as part of a DNA construct. Mechanical methods, such as microinjection, liposome-mediated transfection, electroporation, or calcium phosphate precipitation, can be used to introduce a ribozyme-containing DNA construct into cells in which it is desired to decrease PDE4A expression.
20 Alternatively, if it is desired that the cells stably retain the DNA construct, the construct can be supplied on a plasmid and maintained as a separate element or integrated into the genome of the cells, as is known in the art. A ribozyme-encoding DNA construct can include transcriptional regulatory elements, such as a promoter element, an enhancer or UAS element, and a transcriptional terminator signal, for
25 controlling transcription of ribozymes in the cells (U.S. 5,641,673). Ribozymes also can be engineered to provide an additional level of regulation, so that destruction of mRNA occurs only when both a ribozyme and a target gene are induced in the cells.

Screening / Screening Assays

Regulators

5 Regulators as used herein, refer to compounds that affect the activity of PDE4A in vivo and/or in vitro. Regulators can be agonists and antagonists of PDE4A polypeptide and can be compounds that exert their effect on the PDE4A activity via the enzymatic activity, expression, post-translational modifications or by other means. Agonists of PDE4A are molecules which, when bound to PDE4A, increase or
10 prolong the activity of PDE4A. Agonists of PDE4A include proteins, nucleic acids, carbohydrates, small molecules, or any other molecule which activate PDE4A. Antagonists of PDE4A are molecules which, when bound to PDE4A, decrease the amount or the duration of the activity of PDE4A. Antagonists include proteins, nucleic acids, carbohydrates, antibodies, small molecules, or any other molecule
15 which decrease the activity of PDE4A.

The term "modulate", as it appears herein, refers to a change in the activity of PDE4A polypeptide. For example, modulation may cause an increase or a decrease in enzymatic activity, binding characteristics, or any other biological, functional, or
20 immunological properties of PDE4A.

As used herein, the terms "specific binding" or "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, or an antagonist. The interaction is dependent upon the presence of a particular structure of the protein recognized by the binding molecule (i.e., the antigenic determinant or epitope). For
25 example, if an antibody is specific for epitope "A" the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

30

- 42 -

The invention provides methods (also referred to herein as “screening assays”) for identifying compounds which can be used for the treatment of hematological and cardiovascular diseases, disorders of the peripheral and central nervous system, COPD, asthma, genito-urological disorders and inflammation diseases. The methods
5 entail the identification of candidate or test compounds or agents (e.g., peptides, peptidomimetics, small molecules or other molecules) which bind to PDE4A and/or have a stimulatory or inhibitory effect on the biological activity of PDE4A or its expression and then determining which of these compounds have an effect on symptoms or diseases regarding the hematological and cardiovascular diseases,
10 disorders of the peripheral and central nervous system, COPD, asthma, genito-urological disorders and inflammation diseases in an *in vivo* assay.

Candidate or test compounds or agents which bind to PDE4A and/or have a stimulatory or inhibitory effect on the activity or the expression of PDE4A are
15 identified either in assays that employ cells which express PDE4A (cell-based assays) or in assays with isolated PDE4A (cell-free assays). The various assays can employ a variety of variants of PDE4A (e.g., full-length PDE4A, a biologically active fragment of PDE4A, or a fusion protein which includes all or a portion of PDE4A). Moreover, PDE4A can be derived from any suitable mammalian species
20 (e.g., human PDE4A, rat PDE4A or murine PDE4A). The assay can be a binding assay entailing direct or indirect measurement of the binding of a test compound or a known PDE4A ligand to PDE4A. The assay can also be an activity assay entailing direct or indirect measurement of the activity of PDE4A. The assay can also be an expression assay entailing direct or indirect measurement of the expression of
25 PDE4A mRNA or PDE4A protein. The various screening assays are combined with an *in vivo* assay entailing measuring the effect of the test compound on the symptoms of hematological and cardiovascular diseases, disorders of the peripheral and central nervous system, COPD, asthma, genito-urological disorders and inflammation diseases.

30

- 43 -

The present invention includes biochemical, cell free assays that allow the identification of inhibitors and agonists of PDEs suitable as lead structures for pharmacological drug development. Such assays involve contacting a form of PDE4A (e.g., full-length PDE4A, a biologically active fragment of PDE4A, or a fusion protein comprising all or a portion of PDE4A) with a test compound and determining the ability of the test compound to act as an antagonist (preferably) or an agonist of the enzymatic activity of PDE4A. In one embodiment, the assay includes monitoring the PDE activity of PDE4A by measuring the conversion of either cAMP or cGMP to its nucleoside monophosphate after contacting PDE4A with a test compound.

For example, cAMP and cGMP levels can be measured by the use of the tritium containing compounds $^3\text{HcAMP}$ and $^3\text{HcGMP}$ as described in [Hansen, R.S., and Beavo, J.A., PNAS USA1982;79: 2788-92]. To screen a compound pool comprised of a large number of compounds, the microtiter plate-based scintillation proximity assay (SPA) as described in [Bardelle, C. et al. (1999) Anal. Biochem. 275: 148-155] can be applied.

Alternatively, the phosphodiesterase activity of the recombinant protein can be assayed using a commercially available SPA kit (Amersham Pharmacia). The PDE enzyme hydrolyzes cyclic nucleotides, e.g. cAMP and cGMP to their linear counterparts. The SPA assay utilizes the tritiated cyclic nucleotides [^3H]cAMP or [^3H]cGMP, and is based upon the selective interaction of the tritiated non cyclic product with the SPA beads whereas the cyclic substrates are not effectively binding. Radiolabelled product bound to the scintillation beads generates light that can be analyzed in a scintillation counter.

In one embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of PDE4A. Such assays can employ full-length PDE4A, a biologically active fragment of PDE4A, or a fusion protein which includes all or a portion of PDE4A. As described in greater detail

below, the test compound can be obtained by any suitable means, e.g., from conventional compound libraries.

Determining the ability of the test compound to modulate the activity of PDE4A can be accomplished, for example, by determining the ability of PDE4A to bind to or interact with a target molecule. The target molecule can be a molecule with which PDE4A binds or interacts with in nature. The target molecule can be a component of a signal transduction pathway which facilitates transduction of an extracellular signal. The target PDE4A molecule can be, for example, a second intracellular protein which has catalytic activity or a protein which facilitates the association of downstream signaling molecules with PDE4A.

Determining the ability of PDE4A to bind to or interact with a target molecule can be accomplished by one of the methods described above for determining direct binding. In one embodiment, determining the ability of a polypeptide of the invention to bind to or interact with a target molecule can be accomplished by determining the activity of the target molecule. For example, the activity of the target molecule can be determined by detecting induction of a cellular second messenger of the target (e.g., intracellular Ca^{2+} , diacylglycerol, IP_3 , etc.), detecting catalytic/enzymatic activity of the target on an appropriate substrate, detecting the induction of a reporter gene (e.g., a regulatory element that is responsive to a polypeptide of the invention operably linked to a nucleic acid encoding a detectable marker, e.g., luciferase), or detecting a cellular response.

In various embodiments of the above assay methods of the present invention, it may be desirable to immobilize PDE4A (or a PDE4A target molecule) to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to PDE4A, or interaction of PDE4A with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtitre plates, test tubes, and

- 45 -

micro-centrifuge tubes. In one embodiment, a fusion protein can be provided which adds a domain that allows one or both of the proteins to be bound to a matrix. For example, glutathione-S-transferase (GST) fusion proteins or glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical; St. Louis, Mo.) or glutathione derivatized microtitre plates, which are then combined with the test compound or the test compound and either the non-adsorbed target protein or PDE4A, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtitre plate wells are washed to remove any unbound components and complex formation is measured either directly or indirectly, for example, as described above. Alternatively, the complexes can be dissociated from the matrix, and the level of binding or activity of PDE4A can be determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either PDE4A or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated polypeptide of the invention or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals; Rockford, Ill.), and immobilized in the wells of streptavidin-coated plates (Pierce Chemical). Alternatively, antibodies reactive with PDE4A or target molecules but which do not interfere with binding of the polypeptide of the invention to its target molecule can be derivatized to the wells of the plate, and unbound target or polypeptide of the invention trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with PDE4A or target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with PDE4A or target molecule.

- 46 -

Another technique for drug screening which may be used provides for high throughput screening of compounds having suitable binding affinity to the protein of interest as described in published PCT application WO84/03564. In this method, large numbers of different small test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with PDE4A, or fragments thereof, and washed. Bound PDE4A is then detected by methods well known in the art. Purified PDE4A can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding PDE4A specifically compete with a test compound for binding PDE4A. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PDE4A.

The screening assay can also involve monitoring the expression of PDE4A. For example, regulators of expression of PDE4A can be identified in a method in which a cell is contacted with a candidate compound and the expression of PDE4A protein or mRNA in the cell is determined. The level of expression of PDE4A protein or mRNA in the presence of the candidate compound is compared to the level of expression of PDE4A protein or mRNA in the absence of the candidate compound. The candidate compound can then be identified as a regulator of expression of PDE4A based on this comparison. For example, when expression of PDE4A protein or mRNA is greater (statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of PDE4A protein or mRNA expression. Alternatively, when expression of PDE4A protein or mRNA is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as

an inhibitor of PDE4A protein or mRNA expression. The level of PDE4A protein or mRNA expression in the cells can be determined by methods described below.

Binding Assays

5

For binding assays, the test compound is preferably a small molecule which binds to and occupies the active site of PDE4A polypeptide, thereby making the ligand binding site inaccessible to substrate such that normal biological activity is prevented. Examples of such small molecules include, but are not limited to, small peptides or peptide-like molecules. Potential ligands which bind to a polypeptide of the invention include, but are not limited to, the natural ligands of known PDE4A PDEs and analogues or derivatives thereof.

In binding assays, either the test compound or the PDE4A polypeptide can comprise a detectable label, such as a fluorescent, radioisotopic, chemiluminescent, or enzymatic label, such as horseradish peroxidase, alkaline phosphatase, or luciferase. Detection of a test compound which is bound to PDE4A polypeptide can then be accomplished, for example, by direct counting of radioemmission, by scintillation counting, or by determining conversion of an appropriate substrate to a detectable product. Alternatively, binding of a test compound to a PDE4A polypeptide can be determined without labeling either of the interactants. For example, a microphysiometer can be used to detect binding of a test compound with a PDE4A polypeptide. A microphysiometer (*e.g.*, Cytosensor™) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between a test compound and PDE4A [Haseloff, (1988)].

Determining the ability of a test compound to bind to PDE4A also can be accomplished using a technology such as real-time Bimolecular Interaction Analysis (BIA) [McConnell, (1992); Sjolander, (1991)]. BIA is a technology for studying biospecific interactions in real time, without labeling any of the interactants (*e.g.*,

30

BIAcoreTM). Changes in the optical phenomenon surface plasmon resonance (SPR) can be used as an indication of real-time reactions between biological molecules.

5 In yet another aspect of the invention, a PDE4A-like polypeptide can be used as a "bait protein" in a two-hybrid assay or three-hybrid assay [Szabo, (1995); U.S. 5,283,317), to identify other proteins which bind to or interact with PDE4A and modulate its activity.

10 The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. For example, in one construct, polynucleotide encoding PDE4A can be fused to a polynucleotide encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL-4). In the other construct a DNA sequence that encodes an unidentified protein ("prey" or "sample") can be fused to a
15 polynucleotide that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact *in vivo* to form an protein-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (*e.g.*, LacZ), which is operably linked to a transcriptional regulatory
20 site responsive to the transcription factor. Expression of the reporter gene can be detected, and cell colonies containing the functional transcription factor can be isolated and used to obtain the DNA sequence encoding the protein which interacts with PDE4A.

25 It may be desirable to immobilize either the PDE4A (or polynucleotide) or the test compound to facilitate separation of the bound form from unbound forms of one or both of the interactants, as well as to accommodate automation of the assay. Thus, either the PDE4A-like polypeptide (or polynucleotide) or the test compound can be bound to a solid support. Suitable solid supports include, but are not limited to, glass
30 or plastic slides, tissue culture plates, microtiter wells, tubes, silicon chips, or particles such as beads (including, but not limited to, latex, polystyrene, or glass

beads). Any method known in the art can be used to attach PDE4A-like polypeptide (or polynucleotide) or test compound to a solid support, including use of covalent and non-covalent linkages, passive absorption, or pairs of binding moieties attached respectively to the polypeptide (or polynucleotide) or test compound and the solid support. Test compounds are preferably bound to the solid support in an array, so that the location of individual test compounds can be tracked. Binding of a test compound to PDE4A (or a polynucleotide encoding for PDE4A) can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and microcentrifuge tubes.

In one embodiment, PDE4A is a fusion protein comprising a domain that allows binding of PDE4A to a solid support. For example, glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, Mo.) or glutathione derivatized microtiter plates, which are then combined with the test compound or the test compound and the non-adsorbed PDE4A; the mixture is then incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components. Binding of the interactants can be determined either directly or indirectly, as described above. Alternatively, the complexes can be dissociated from the solid support before binding is determined.

Other techniques for immobilizing proteins or polynucleotides on a solid support also can be used in the screening assays of the invention. For example, either PDE4A (or a polynucleotide encoding PDE4A) or a test compound can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated PDE4A (or a polynucleotide encoding biotinylated PDE4A) or test compounds can be prepared from biotin-NHS (N-hydroxysuccinimide) using techniques well known in the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, Ill.) and immobilized in the wells of streptavidin-coated plates (Pierce Chemical). Alternatively, antibodies which specifically bind to PDE4A, polynucleotide, or a test compound, but which do not interfere with a

- 50 -

desired binding site, such as the active site of PDE4A, can be derivatized to the wells of the plate. Unbound target or protein can be trapped in the wells by antibody conjugation.

5 Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies which specifically bind to PDE4A polypeptide or test compound, enzyme-linked assays which rely on detecting an activity of PDE4A polypeptide, and SDS gel electrophoresis under non-reducing conditions.

10

Screening for test compounds which bind to a PDE4A polypeptide or polynucleotide also can be carried out in an intact cell. Any cell which comprises a PDE4A polypeptide or polynucleotide can be used in a cell-based assay system. A PDE4A polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Binding of the test compound to PDE4A or a polynucleotide encoding PDE4A is determined as described above.

15

Functional Assays

20 Test compounds can be tested for the ability to increase or decrease PDE4A activity of a PDE4A polypeptide. The PDE4A activity can be measured, for example, using methods described in the specific examples, below. PDE4A activity can be measured after contacting either a purified PDE4A or an intact cell with a test compound. A test compound which decreases PDE4A activity by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for decreasing PDE4A activity. A test compound which increases PDE4A activity by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for increasing PDE4A activity.

25

Gene Expression

In another embodiment, test compounds which increase or decrease PDE4A gene expression are identified. As used herein, the term "correlates with expression of a polynucleotide" indicates that the detection of the presence of nucleic acids, the same or related to a nucleic acid sequence encoding PDE4A, by northern analysis or realtime PCR is indicative of the presence of nucleic acids encoding PDE4A in a sample, and thereby correlates with expression of the transcript from the polynucleotide encoding PDE4A. The term "microarray", as used herein, refers to an array of distinct polynucleotides or oligonucleotides arrayed on a substrate, such as paper, nylon or any other type of membrane, filter, chip, glass slide, or any other suitable solid support. A PDE4A polynucleotide is contacted with a test compound, and the expression of an RNA or polypeptide product of PDE4A polynucleotide is determined. The level of expression of appropriate mRNA or polypeptide in the presence of the test compound is compared to the level of expression of mRNA or polypeptide in the absence of the test compound. The test compound can then be identified as a regulator of expression based on this comparison. For example, when expression of mRNA or polypeptide is greater in the presence of the test compound than in its absence, the test compound is identified as a stimulator or enhancer of the mRNA or polypeptide expression. Alternatively, when expression of the mRNA or polypeptide is less in the presence of the test compound than in its absence, the test compound is identified as an inhibitor of the mRNA or polypeptide expression.

The level of PDE4A mRNA or polypeptide expression in the cells can be determined by methods well known in the art for detecting mRNA or polypeptide. Either qualitative or quantitative methods can be used. The presence of polypeptide products of PDE4A polynucleotide can be determined, for example, using a variety of techniques known in the art, including immunochemical methods such as radioimmunoassay, Western blotting, and immunohistochemistry. Alternatively, polypeptide synthesis can be determined *in vivo*, in a cell culture, or in an *in vitro* translation system by detecting incorporation of labelled amino acids into PDE4A.

Such screening can be carried out either in a cell-free assay system or in an intact cell. Any cell which expresses PDE4A polynucleotide can be used in a cell-based assay system. The PDE4A polynucleotide can be naturally occurring in the cell or
5 can be introduced using techniques such as those described above. Either a primary culture or an established cell line can be used.

Test Compounds

10 Suitable test compounds for use in the screening assays of the invention can be obtained from any suitable source, e.g., conventional compound libraries. The test compounds can also be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries;
15 spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the "one-bead one-compound" library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of
20 compounds [Lam, (1997)]. Examples of methods for the synthesis of molecular libraries can be found in the art. Libraries of compounds may be presented in solution or on beads, bacteria, spores, plasmids or phage.

Modeling of Regulators

25 Computer modeling and searching technologies permit identification of compounds, or the improvement of already identified compounds, that can modulate PDE4A expression or activity. Having identified such a compound or composition, the active sites or regions are identified. Such sites might typically be the enzymatic active site, regulator binding sites, or ligand binding sites. The active site can be identified using
30 methods known in the art including, for example, from the amino acid sequences of peptides, from the nucleotide sequences of nucleic acids, or from study of complexes

- 53 -

of the relevant compound or composition with its natural ligand. In the latter case, chemical or X-ray crystallographic methods can be used to find the active site by finding where on the factor the complexed ligand is found.

5 Next, the three dimensional geometric structure of the active site is determined. This can be done by known methods, including X-ray crystallography, which can determine a complete molecular structure. On the other hand, solid or liquid phase NMR can be used to determine certain intramolecular distances. Any other
10 experimental method of structure determination can be used to obtain partial or complete geometric structures. The geometric structures may be measured with a complexed ligand, natural or artificial, which may increase the accuracy of the active site structure determined.

15 If an incomplete or insufficiently accurate structure is determined, the methods of computer based numerical modeling can be used to complete the structure or improve its accuracy. Any recognized modeling method may be used, including parameterized models specific to particular biopolymers such as proteins or nucleic acids, molecular dynamics models based on computing molecular motions, statistical
20 mechanics models based on thermal ensembles, or combined models. For most types of models, standard molecular force fields, representing the forces between constituent atoms and groups, are necessary, and can be selected from force fields known in physical chemistry. The incomplete or less accurate experimental structures can serve as constraints on the complete and more accurate structures computed by these modeling methods.

25 Finally, having determined the structure of the active site, either experimentally, by modeling, or by a combination, candidate modulating compounds can be identified by searching databases containing compounds along with information on their molecular structure. Such a search seeks compounds having structures that match the
30 determined active site structure and that interact with the groups defining the active

site. Such a search can be manual, but is preferably computer assisted. These compounds found from this search are potential PDE4A modulating compounds.

5 Alternatively, these methods can be used to identify improved modulating compounds from an already known modulating compound or ligand. The composition of the known compound can be modified and the structural effects of modification can be determined using the experimental and computer modeling methods described above applied to the new composition. The altered structure is then compared to the active site structure of the compound to determine if an improved fit or interaction results. In this manner systematic variations in composition, such as by varying side groups, can be quickly evaluated to obtain modified modulating compounds or ligands of improved specificity or activity.

15 **Therapeutic Indications and Methods**

It was found by the present applicant that PDE4A is expressed in various human tissues.

20 *Neurology*

CNS disorders include disorders of the central nervous system as well as disorders of the peripheral nervous system.

25 CNS disorders include, but are not limited to brain injuries, cerebrovascular diseases and their consequences, Parkinson's disease, corticobasal degeneration, motor neuron disease, dementia, including ALS, multiple sclerosis, traumatic brain injury, stroke, post-stroke, post-traumatic brain injury, and small-vessel cerebrovascular disease. Dementias, such as Alzheimer's disease, vascular dementia, dementia with Lewy bodies, frontotemporal dementia and Parkinsonism linked to chromosome 17, frontotemporal dementias, including Pick's disease, progressive nuclear palsy, 30 corticobasal degeneration, Huntington's disease, thalamic degeneration, Creutzfeld-

Jakob dementia, HIV dementia, schizophrenia with dementia, and Korsakoff's psychosis, within the meaning of the definition are also considered to be CNS disorders.

- 5 Similarly, cognitive-related disorders, such as mild cognitive impairment, age-associated memory impairment, age-related cognitive decline, vascular cognitive impairment, attention deficit disorders, attention deficit hyperactivity disorders, and memory disturbances in children with learning disabilities are also considered to be CNS disorders.

10

- Pain, within the meaning of this definition, is also considered to be a CNS disorder. Pain can be associated with CNS disorders, such as multiple sclerosis, spinal cord injury, sciatica, failed back surgery syndrome, traumatic brain injury, epilepsy, Parkinson's disease, post-stroke, and vascular lesions in the brain and spinal cord (e.g., infarct, hemorrhage, vascular malformation). Non-central neuropathic pain includes that associated with post mastectomy pain, phantom feeling, reflex sympathetic dystrophy (RSD), trigeminal neuralgia, radioculopathy, post-surgical pain, HIV/AIDS related pain, cancer pain, metabolic neuropathies (e.g., diabetic neuropathy, vasculitic neuropathy secondary to connective tissue disease),
15 paraneoplastic polyneuropathy associated, for example, with carcinoma of lung, or leukemia, or lymphoma, or carcinoma of prostate, colon or stomach, trigeminal neuralgia, cranial neuralgias, and post-herpetic neuralgia. Pain associated with peripheral nerve damage, central pain (i.e. due to cerebral ischemia) and various chronic pain i.e., lumbago, back pain (low back pain), inflammatory and/or
20 rheumatic pain. Headache pain (for example, migraine with aura, migraine without aura, and other migraine disorders), episodic and chronic tension-type headache, tension-type like headache, cluster headache, and chronic paroxysmal hemicrania are also CNS disorders.

- 25 Visceral pain such as pancreatitis, intestinal cystitis, dysmenorrhea, irritable Bowel syndrome, Crohn's disease, biliary colic, ureteral colic, myocardial infarction and

pain syndromes of the pelvic cavity, e.g., vulvodynia, orchialgia, urethral syndrome and protatodynia are also CNS disorders.

Also considered to be a disorder of the nervous system are acute pain, for example postoperative pain, and pain after trauma.

The human PDE4A is highly expressed in the following brain tissues: brain, alzheimer brain, cerebral cortex, alzheimer cerebral cortex, frontal lobe, alzheimer brain frontal lobe, occipital lobe, parietal lobe, temporal lobe, precentral gyrus, substantia nigra, cerebral peduncles, hippocampus, thalamus, spinal cord. The expression in brain tissues and in particular the differential expression between diseased tissue alzheimer brain and healthy tissue brain demonstrates that the human PDE4A or mRNA can be utilized to diagnose nervous system diseases. Additionally the activity of the human PDE4A can be modulated to treat nervous system diseases.

15

Cardiovascular Disorders

Heart failure is defined as a pathophysiological state in which an abnormality of cardiac function is responsible for the failure of the heart to pump blood at a rate commensurate with the requirement of the metabolizing tissue. It includes all forms of pumping failures such as high-output and low-output, acute and chronic, right-sided or left-sided, systolic or diastolic, independent of the underlying cause.

20

Myocardial infarction (MI) is generally caused by an abrupt decrease in coronary blood flow that follows a thrombotic occlusion of a coronary artery previously narrowed by arteriosclerosis. MI prophylaxis (primary and secondary prevention) is included as well as the acute treatment of MI and the prevention of complications.

25

Ischemic diseases are conditions in which the coronary flow is restricted resulting in a perfusion which is inadequate to meet the myocardial requirement for oxygen. This group of diseases includes stable angina, unstable angina and asymptomatic

30

ischemia.

Arrhythmias include all forms of atrial and ventricular tachyarrhythmias, atrial tachycardia, atrial flutter, atrial fibrillation, atrio-ventricular reentrant tachycardia, 5 preexcitation syndrome, ventricular tachycardia, ventricular flutter, ventricular fibrillation, as well as bradycardic forms of arrhythmias.

Hypertensive vascular diseases include primary as well as all kinds of secondary arterial hypertension, renal, endocrine, neurogenic, others. The genes may be used as 10 drug targets for the treatment of hypertension as well as for the prevention of all complications arising from cardiovascular diseases.

Peripheral vascular diseases are defined as vascular diseases in which arterial and/or venous flow is reduced resulting in an imbalance between blood supply and tissue 15 oxygen demand. It includes chronic peripheral arterial occlusive disease (PAOD), acute arterial thrombosis and embolism, inflammatory vascular disorders, Raynaud's phenomenon and venous disorders.

Atherosclerosis is a cardiovascular disease in which the vessel wall is remodeled, 20 compromising the lumen of the vessel. The atherosclerotic remodeling process involves accumulation of cells, both smooth muscle cells and monocyte/macrophage inflammatory cells, in the intima of the vessel wall. These cells take up lipid, likely from the circulation, to form a mature atherosclerotic lesion. Although the formation of these lesions is a chronic process, occurring over decades of an adult human life, 25 the majority of the morbidity associated with atherosclerosis occurs when a lesion ruptures, releasing thrombogenic debris that rapidly occludes the artery. When such an acute event occurs in the coronary artery, myocardial infarction can ensue, and in the worst case, can result in death.

30 The formation of the atherosclerotic lesion can be considered to occur in five overlapping stages such as migration, lipid accumulation, recruitment of inflam-

matory cells, proliferation of vascular smooth muscle cells, and extracellular matrix deposition. Each of these processes can be shown to occur in man and in animal models of atherosclerosis, but the relative contribution of each to the pathology and clinical significance of the lesion is unclear.

5

Thus, a need exists for therapeutic methods and agents to treat cardiovascular pathologies, such as atherosclerosis and other conditions related to coronary artery disease.

10 Cardiovascular diseases include but are not limited to disorders of the heart and the vascular system like congestive heart failure, myocardial infarction, ischemic diseases of the heart, all kinds of atrial and ventricular arrhythmias, hypertensive vascular diseases, peripheral vascular diseases, and atherosclerosis.

15 To high or to low levels of fats in the bloodstream, especially cholesterol, can cause long-term problems. The risk to develop atherosclerosis and coronary artery or carotid artery disease (and thus the risk of having a heart attack or stroke) increases with the total cholesterol level increasing. Nevertheless, extremely low cholesterol levels may not be healthy. Examples of disorders of lipid metabolism are
20 hyperlipidemia (abnormally high levels of fats (cholesterol, triglycerides, or both) in the blood, may be caused by family history of hyperlipidemia), obesity, a high-fat diet, lack of exercise, moderate to high alcohol consumption, cigarette smoking, poorly controlled diabetes, and an underactive thyroid gland), hereditary hyperlipidemias (type I hyperlipoproteinemia (familial hyperchylomicronemia), type II
25 hyperlipoproteinemia (familial hypercholesterolemia), type III hyperlipoproteinemia, type IV hyperlipoproteinemia, or type V hyperlipoproteinemia), hypolipoproteinemia, lipidoses (caused by abnormalities in the enzymes that metabolize fats), Gaucher's disease, Niemann-Pick disease, Fabry's disease, Wolman's disease, cerebrotendinous xanthomatosis, sitosterolemia, Refsum's disease, or Tay-Sachs
30 disease.

Kidney disorders may lead to hypertension or hypotension. Examples for kidney problems possibly leading to hypertension are renal artery stenosis, pyelonephritis, glomerulonephritis, kidney tumors, polycystic kidney disease, injury to the kidney, or radiation therapy affecting the kidney. Excessive urination may lead to hypotension.

5

The human PDE4A is highly expressed in the following cardiovascular related tissues: heart, pericardium, coronary artery smooth muscle primary cells, HUVEC cells. Expression in the above mentioned tissues demonstrates that the human PDE4A or mRNA can be utilized to diagnose of cardiovascular diseases. Additionally the activity of the human PDE4A can be modulated to treat cardiovascular diseases.

10

Hematological Disorders

15 Hematological disorders comprise diseases of the blood and all its constituents as well as diseases of organs and tissues involved in the generation or degradation of all the constituents of the blood. They include but are not limited to 1) Anemias, 2) Myeloproliferative Disorders, 3) Hemorrhagic Disorders, 4) Leukopenia, 5) Eosinophilic Disorders, 6) Leukemias, 7) Lymphomas, 8) Plasma Cell Dyscrasias, 9) Disorders of the Spleen in the course of hematological disorders. Disorders according to 1) include, but are not limited to anemias due to defective or deficient hem synthesis, deficient erythropoiesis. Disorders according to 2) include, but are not limited to polycythemia vera, tumor-associated erythrocytosis, myelofibrosis, thrombocythemia. Disorders according to 3) include, but are not limited to vasculitis, thrombocytopenia, heparin-induced thrombocytopenia, thrombotic thrombocytopenic purpura, hemolytic-uremic syndrome, hereditary and acquired disorders of platelet function, hereditary coagulation disorders. Disorders according to 4) include, but are not limited to neutropenia, lymphocytopenia. Disorders according to 5) include, but are not limited to hypereosinophilia, idiopathic hypereosinophilic syndrome. Disorders according to 6) include, but are not limited to acute myeloic leukemia, acute lymphoblastic leukemia, chronic myelocytic leukemia, chronic lymphocytic

20

25

30

- 60 -

leukemia, myelodysplastic syndrome. Disorders according to 7) include, but are not limited to Hodgkin's disease, non-Hodgkin's lymphoma, Burkitt's lymphoma, mycosis fungoides cutaneous T-cell lymphoma. Disorders according to 8) include, but are not limited to multiple myeloma, macroglobulinemia, heavy chain diseases.

5 In extension of the preceding idiopathic thrombocytopenic purpura, iron deficiency anemia, megaloblastic anemia (vitamin B12 deficiency), aplastic anemia, thalassemia, malignant lymphoma bone marrow invasion, malignant lymphoma skin invasion, hemolytic uremic syndrome, giant platelet disease are considered to be hematological diseases too.

10

The human PDE4A is highly expressed in the following tissues of the hematological system: leukocytes (peripheral blood), thymus, spleen, spleen liver cirrhosis. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue spleen liver cirrhosis and healthy tissue spleen demonstrates
15 that the human PDE4A or mRNA can be utilized to diagnose of hematological diseases. Additionally the activity of the human PDE4A can be modulated to treat hematological disorders.

Gastrointestinal and Liver Diseases

20

Gastrointestinal diseases comprise primary or secondary, acute or chronic diseases of the organs of the gastrointestinal tract which may be acquired or inherited, benign or malignant or metaplastic, and which may affect the organs of the gastrointestinal tract or the body as a whole. They comprise but are not limited to 1) disorders of the
25 esophagus like achalasia, vigorous achalasia, dysphagia, cricopharyngeal incoordination, pre-esophageal dysphagia, diffuse esophageal spasm, globus sensation, Barrett's metaplasia, gastroesophageal reflux, 2) disorders of the stomach and duodenum like functional dyspepsia, inflammation of the gastric mucosa, gastritis, stress gastritis, chronic erosive gastritis, atrophy of gastric glands,
30 metaplasia of gastric tissues, gastric ulcers, duodenal ulcers, neoplasms of the stomach, 3) disorders of the pancreas like acute or chronic pancreatitis, insufficiency

of the exocrine or endocrine tissues of the pancreas like steatorrhea, diabetes, neoplasms of the exocrine or endocrine pancreas like 3.1) multiple endocrine neoplasia syndrome, ductal adenocarcinoma, cystadenocarcinoma, islet cell tumors, insulinoma, gastrinoma, carcinoid tumors, glucagonoma, Zollinger-Ellison syndrome, Vipoma syndrome, malabsorption syndrome, 4) disorders of the bowel like chronic inflammatory diseases of the bowel, Crohn's disease, ileus, diarrhea and constipation, colonic inertia, megacolon, malabsorption syndrome, ulcerative colitis, 4.1) functional bowel disorders like irritable bowel syndrome, 4.2) neoplasms of the bowel like familial polyposis, adenocarcinoma, primary malignant lymphoma, carcinoid tumors, Kaposi's sarcoma, polyps, cancer of the colon and rectum.

Liver diseases comprise primary or secondary, acute or chronic diseases or injury of the liver which may be acquired or inherited, benign or malignant, and which may affect the liver or the body as a whole. They comprise but are not limited to disorders of the bilirubin metabolism, jaundice, syndromes of Gilbert's, Crigler-Najjar, Dubin-Johnson and Rotor; intrahepatic cholestasis, hepatomegaly, portal hypertension, ascites, Budd-Chiari syndrome, portal-systemic encephalopathy, fatty liver, steatosis, Reye's syndrome, liver diseases due to alcohol, alcoholic hepatitis or cirrhosis, fibrosis and cirrhosis, fibrosis and cirrhosis of the liver due to inborn errors of metabolism or exogenous substances, storage diseases, syndromes of Gaucher's, Zellweger's, Wilson's - disease, acute or chronic hepatitis, viral hepatitis and its variants, inflammatory conditions of the liver due to viruses, bacteria, fungi, protozoa, helminths; drug induced disorders of the liver, chronic liver diseases like primary sclerosing cholangitis, α_1 -antitrypsin-deficiency, primary biliary cirrhosis, postoperative liver disorders like postoperative intrahepatic cholestasis, hepatic granulomas, vascular liver disorders associated with systemic disease, benign or malignant neoplasms of the liver, disturbance of liver metabolism in the new-born or prematurely born.

The human PDE4A is highly expressed in the following tissues of the gastroenterological system: stomach, colon, colon tumor, ileum chronic inflammation,

- 62 -

rectum, liver, liver liver cirrhosis, HEP G2 cells. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue colon tumor and healthy tissue colon demonstrates that the human PDE4A or mRNA can be utilized to diagnose of gastroenterological disorders. Additionally the activity of the human PDE4A can be modulated to treat gastroenterological disorders.

Endocrine System and Hormones

The endocrine system consists of a group of organs whose main function is to produce and secrete hormones directly into the bloodstream. The major organs of the endocrine system are the hypothalamus, the pituitary gland, thyroid gland, the parathyroid glands, the islets of the pancreas, the adrenal glands, the testes, and the ovaries.

The hypothalamus secretes several hormones that stimulate the pituitary: Some trigger the release of pituitary hormones; others suppress the release of pituitary hormones.

The pituitary gland coordinates many functions of the other endocrine glands, but some pituitary hormones have direct effects.

The insulin-secreting cells of the pancreas respond to glucose and fatty acids. Parathyroid cells respond to calcium and phosphate. The adrenal medulla (part of the adrenal gland) responds to direct stimulation by the parasympathetic nervous system.

When endocrine glands malfunction, hormone levels in the blood can become abnormally high or low, disrupting body functions. Many disorders are caused by malfunction of the endocrine system or hormones. Examples of such disorders are presented in the following.

- 63 -

Diabetes mellitus is a disorder in which blood levels of glucose are abnormally high because the body doesn't release or use insulin adequately.

5 People with type I diabetes mellitus (insulin-dependent diabetes) produce little or no insulin at all. In type I diabetes more than 90 percent of the insulin-producing cells (beta cells) of the pancreas are permanently destroyed. The resulting insulin deficiency is severe, and to survive, a person with type I diabetes must regularly inject insulin.

10 In type II diabetes mellitus (non-insulin-dependent diabetes) the body develops resistance to insulin effects, resulting in a relative insulin deficiency.

15 The pancreas has two major functions: to secrete fluid containing digestive enzymes into the duodenum and to secrete the hormones insulin and glucagon. Chronic pancreatitis is a long-standing inflammation of the pancreas. Eventually, the insulin-secreting cells of the pancreas may be destroyed, gradually leading to diabetes. An insulinoma is a rare type of pancreatic tumor that secretes insulin. The symptoms of an insulinoma result from low blood glucose levels. A gastrinoma is a pancreatic tumor that produces excessive levels of the hormone gastrin, which stimulates the
20 stomach to secrete acid and enzymes, causing peptic ulcers. The excess gastrin secreted by the gastrinoma causes symptoms, called the Zollinger-Ellison syndrome. A glucagonoma is a tumor that produces the hormone glucagon, which raises the level of glucose in the blood and produces a distinctive rash.

25 Diabetes insipidus is a disorder in which insufficient levels of antidiuretic hormone cause excessive thirst (polydipsia) and excessive production of very dilute urine (polyuria). Diabetes insipidus results from the decreased production of antidiuretic hormone (vasopressin).

30 The body has two adrenal glands. The medulla of the adrenal glands secretes hormones such as adrenaline (epinephrine) that affect blood pressure, heart rate,

- 64 -

sweating, and other activities also regulated by the sympathetic nervous system. The cortex secretes many different hormones, including corticosteroids (cortisone-like hormones), androgens (male hormones), and mineralocorticoids, which control blood pressure and the levels of salt and potassium in the body.

5

A diseases characterized by underactive adrenal glands is Addison's disease (adrenocortical insufficiency).

10 Several disorders are characterized by overactive Adrenal Glands. The causes can be changes in the adrenal glands themselves or overstimulation by the pituitary gland. Examples of these diseases are listed in the following.

15 Overproduction of androgenic steroids (testosterone and similar hormones, leads to virilization), overproduction of corticosteroids (causes could be tumors of the pituitary or the adrenal gland, results in Cushing's syndrome), Nelson's syndrome (developed by people who have both adrenal glands removed, characterized by an enlargement of the pituitary gland), Overproduction of aldosterone (hyperaldosteronism), Conn's syndrome (hyperaldosterism caused by a tumor), pheochromocytoma (a tumor that originating from the adrenal gland's chromaffin cells, causing overproduction of catecholamines),
20

The thyroid is a small gland located under the Adam's apple. It secretes thyroid hormones, which control the metabolic rate. The thyroid gland traps iodine and processes it into thyroid hormones. The euthyroid sick syndrome is characterized by
25 lack of conversion of the T4 form of thyroid hormone to the T3 form. Hyperthyroidism (overactive thyroid gland, production of too much hormone) may have several causes. Thyroiditis (an inflammation of the thyroid gland), typically leads to a phase of hyperthyroidism. The inflammation may damage the thyroid gland, so that in later stages the disease is characterized by transient or permanent
30 underactivity (hypothyroidism). Toxic thyroid nodules (adenomas) often produce thyroid hormone in large quantities. Toxic multinodular goiter (Plummer's disease) is

- 65 -

a disorder in which there are many nodules. Graves' disease (toxic diffuse goiter) is believed to be caused by an antibody that stimulates the thyroid to produce too much thyroid hormone. In toxic nodular goiter, one or more nodules in the thyroid produce too much thyroid hormone and aren't under the control of thyroid-stimulating hormone. Secondary hyperthyroidism may (rarely) be caused by a pituitary tumor that secretes too much thyroid-stimulating hormone, by resistance of the pituitary to thyroid hormone, which results in the pituitary gland secreting too much thyroid-stimulating hormone, or by a hydatidiform mole in women. Thyroid storm is a sudden extreme overactivity of the thyroid gland is a life-threatening emergency requiring prompt treatment.

Hypothyroidism is a condition in which the thyroid gland is underactive and produces too little thyroid hormone. Very severe hypothyroidism is called myxedema. In Hashimoto's thyroiditis (autoimmune thyroiditis) the thyroid gland is often enlarged, and hypothyroidism results because the gland's functioning areas are gradually destroyed. Rarer causes of hypothyroidism include some inherited disorders which are caused by abnormalities of the enzymes in thyroid cells. In other rare disorders, either the hypothalamus or the pituitary gland fails to secrete enough of the hormone needed to stimulate normal thyroid function.

Other examples of Thyroiditis are silent lymphocytic thyroiditis, Hashimoto's thyroiditis, or subacute granulomatous thyroiditis.

Thyroid cancer is any one of four main types of malignancy of the thyroid: papillary, follicular, anaplastic, or medullary.

The pituitary is a pea-sized gland that sits in a bony structure (sella turcica) at the base of the brain. The sella turcica protects the pituitary but allows very little room for expansion. If the pituitary enlarges, it tends to push upward, often pressing on the areas of the brain that carry signals from the eyes, possibly resulting in headaches or impaired vision. The pituitary gland has two distinct parts: the anterior (front) and the

- 66 -

posterior (back) lobes. The anterior lobe produces (secretes) hormones that ultimately control the function of the thyroid gland, adrenal glands, and reproductive organs (ovaries and testes); milk production (lactation) in the breasts; and overall body growth. It also produces hormones that cause the skin to darken and that inhibit pain
5 sensations. The posterior lobe produces hormones that regulate water balance, stimulate the let-down of milk from the breasts in lactating women, and stimulate contractions of the uterus.

Examples for disorders of the pituitary gland are Empty Sella Syndrome; hypo-
10 pituitarism (an underactive pituitary gland); acromegaly, which is excessive growth caused by oversecretion of growth hormone, which is almost always caused by a benign pituitary tumor (adenoma); galactorrhea, which is the production of breast milk in men or in women who aren't breastfeeding, in both sexes, the most common cause of galactorrhea is a prolactin-producing tumor (prolactinoma) in the pituitary
15 gland.

The human PDE4A is highly expressed in the following tissues of the endocrinological system: thyroid, thyroid tumor, pancreas, pancreas liver cirrhosis. The expression in the above mentioned tissues and in particular the differential expression
20 between diseased tissue thyroid tumor and healthy tissue thyroid demonstrates that the human PDE4A or mRNA can be utilized to diagnose of endocrinological disorders. Additionally the activity of the human PDE4A can be modulated to treat endocrinological disorders.

25 *Inflammatory Diseases*

Inflammatory diseases comprise diseases triggered by cellular or non-cellular mediators of the immune system or tissues causing the inflammation of body tissues and subsequently producing an acute or chronic inflammatory condition. Examples
30 for such inflammatory diseases are hypersensitivity reactions of type I – IV, for example but not limited to hypersensitivity diseases of the lung including asthma,

- 67 -

atopic diseases, allergic rhinitis or conjunctivitis, angioedema of the lids, hereditary angioedema, antireceptor hypersensitivity reactions and autoimmune diseases, Hashimoto's thyroiditis, systemic lupus erythematosus, Goodpasture's syndrome, pemphigus, myasthenia gravis, Grave's and Raynaud's disease, type B insulin-resistant diabetes, rheumatoid arthritis, psoriasis, Crohn's disease, scleroderma, mixed connective tissue disease, polymyositis, sarcoidosis, glomerulonephritis, acute or chronic host versus graft reactions.

The human PDE4A is highly expressed in the following tissues of the immune system and tissues responsive to components of the immune system as well as in the following tissues responsive to mediators of inflammation: pancreas liver cirrhosis, ileum chronic inflammation, liver liver cirrhosis, leukocytes (peripheral blood), spleen liver cirrhosis. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue pancreas liver cirrhosis and healthy tissue pancreas demonstrates that the human PDE4A or mRNA can be utilized to diagnose of inflammatory diseases. Additionally the activity of the human PDE4A can be modulated to treat inflammatory diseases.

Applications

The present invention provides for both prophylactic and therapeutic methods for disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease.

The regulatory method of the invention involves contacting a cell with an agent that modulates one or more of the activities of PDE4A. An agent that modulates activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring cognate ligand of the polypeptide, a peptide, a peptidomimetic, or any small molecule. In one embodiment, the agent stimulates one or more of the biological activities of PDE4A. Examples of such stimulatory agents include the

active PDE4A and nucleic acid molecules encoding a portion of PDE4A. In another embodiment, the agent inhibits one or more of the biological activities of PDE4A. Examples of such inhibitory agents include antisense nucleic acid molecules and antibodies. These regulatory methods can be performed in vitro (e.g., by culturing
5 the cell with the agent) or, alternatively, in vivo (e.g. by administering the agent to a subject). As such, the present invention provides methods of treating an individual afflicted with a disease or disorder characterized by unwanted expression or activity of PDE4A or a protein in the PDE4A signaling pathway. In one embodiment, the method involves administering an agent like any agent identified or being
10 identifiable by a screening assay as described herein, or combination of such agents that modulate say upregulate or downregulate the expression or activity of PDE4A or of any protein in the PDE4A signaling pathway. In another embodiment, the method involves administering a regulator of PDE4A as therapy to compensate for reduced or undesirably low expression or activity of PDE4A or a protein in the PDE4A
15 signaling pathway.

Stimulation of activity or expression of PDE4A is desirable in situations in which enzymatic activity or expression is abnormally low and in which increased activity is likely to have a beneficial effect. Conversely, inhibition of enzymatic activity or
20 expression of PDE4A is desirable in situations in which activity or expression of PDE4A is abnormally high and in which decreasing its activity is likely to have a beneficial effect.

This invention is further illustrated by the following examples which should not be
25 construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are hereby incorporated by reference.

Pharmaceutical Compositions

30 This invention further pertains to novel agents identified by the above-described screening assays and uses thereof for treatments as described herein.

- 69 -

The nucleic acid molecules, polypeptides, and antibodies (also referred to herein as “active compounds”) of the invention can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein the language “pharmaceutically acceptable carrier” is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

The invention includes pharmaceutical compositions comprising a regulator of PDE4A expression or activity (and/or a regulator of the activity or expression of a protein in the PDE4A signaling pathway) as well as methods for preparing such compositions by combining one or more such regulators and a pharmaceutically acceptable carrier. Also within the invention are pharmaceutical compositions comprising a regulator identified using the screening assays of the invention packaged with instructions for use. For regulators that are antagonists of PDE4A activity or which reduce PDE4A expression, the instructions would specify use of the pharmaceutical composition for treatment of hematological and cardiovascular diseases, disorders of the peripheral and central nervous system, COPD, asthma, genito-urological disorders and inflammation diseases. For regulators that are agonists of PDE4A activity or increase PDE4A expression, the instructions would specify use of the pharmaceutical composition for treatment of hematological and cardiovascular diseases, disorders of the peripheral and central nervous system, COPD, asthma, genito-urological disorders and inflammation diseases.

An inhibitor of PDE4A may be produced using methods which are generally known in the art. In particular, purified PDE4A may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind PDE4A. Antibodies to PDE4A may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies like those which inhibit dimer formation are especially preferred for therapeutic use.

In another embodiment of the invention, the polynucleotides encoding PDE4A, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding PDE4A may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding PDE4A. Thus, complementary molecules or fragments may be used to modulate PDE4A activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding PDE4A.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors which will express nucleic acid sequence complementary to the polynucleotides of the gene encoding PDE4A. These techniques are described, for example, in [Scott and Smith (1990)].

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

- 71 -

An additional embodiment of the invention relates to the administration of a pharmaceutical composition containing PDE4A in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of PDE4A, antibodies to PDE4A, and mimetics, agonists, antagonists, or inhibitors of PDE4A. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs or hormones.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EMTM (BASF, Parsippany, N.J.) or phosphate

buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, a pharmaceutically acceptable polyol like glycerol, propylene glycol, liquid polyethylene glycol, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin. Sterile injectable solutions can be prepared by incorporating the active compound (e.g., a polypeptide or antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be

- 73 -

prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed.

Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from a pressurized container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release

- 74 -

formulation, including implants and microencapsulated delivery systems. Bio-degradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration. For pharmaceutical compositions which include an antagonist of PDE4A activity, a compound which reduces expression of PDE4A, or a compound which reduces expression or activity of a protein in the PDE4A signaling pathway or any combination thereof, the instructions for administration will specify use of the composition for hematological and cardiovascular diseases, disorders of the peripheral and central nervous system, COPD, asthma, genito-urological disorders and inflammation diseases. For pharmaceutical compositions which include an agonist of PDE4A activity, a compound which increases expression of PDE4A, or a compound which increases expression or

- 75 -

activity of a protein in the PDE4A signaling pathway or any combination thereof, the instructions for administration will specify use of the composition for hematological and cardiovascular diseases, disorders of the peripheral and central nervous system, COPD, asthma, genito-urological disorders and inflammation diseases.

5

Diagnostics

In another embodiment, antibodies which specifically bind PDE4A may be used for the diagnosis of disorders characterized by the expression of PDE4A, or in assays to monitor patients being treated with PDE4A or agonists, antagonists, and inhibitors of PDE4A. Antibodies useful for diagnostic purposes may be prepared in the same manner as those described above for therapeutics. Diagnostic assays for PDE4A include methods which utilize the antibody and a label to detect PDE4A in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent joining with a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

20

25

A variety of protocols for measuring PDE4A, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of PDE4A expression. Normal or standard values for PDE4A expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to PDE4A under conditions suitable for complex formation. The amount of standard complex formation may be quantified by various methods, preferably by photometric means. Quantities of PDE4A expressed in subject samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

30

In another embodiment of the invention, the polynucleotides encoding PDE4A may be used for diagnostic purposes. The polynucleotides which may be used include

- 76 -

oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantitate gene expression in biopsied tissues in which expression of PDE4A may be correlated with disease. The diagnostic assay may be used to distinguish between absence, presence, and excess
5 expression of PDE4A, and to monitor regulation of PDE4A levels during therapeutic intervention.

Polynucleotide sequences encoding PDE4A may be used for the diagnosis of disorders of the peripheral and central nervous system, cardiovascular diseases,
10 hematological diseases, inflammation, gastroenterological diseases and endocrinological disease associated with expression of PDE4A. The polynucleotide sequences encoding PDE4A may be used in Southern, Northern, or dot-blot analysis, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and ELISA assays; and in microarrays utilizing fluids or tissues from patient biopsies to
15 detect altered PDE4A expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding PDE4A may be useful in assays that detect the presence of associated disorders, particularly those mentioned
20 above. The nucleotide sequences encoding PDE4A may be labelled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the patient sample is significantly altered
25 from that of a comparable control sample, the nucleotide sequences have hybridized with nucleotide sequences in the sample, and the presence of altered levels of nucleotide sequences encoding PDE4A in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or in
30 monitoring the treatment of an individual patient.

- 77 -

In order to provide a basis for the diagnosis of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease associated with expression of PDE4A, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding PDE4A, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained from normal samples may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Determination of a Therapeutically Effective Dose

The determination of a therapeutically effective dose is well within the capability of those skilled in the art. A therapeutically effective dose refers to that amount of active ingredient which increases or decreases PDE4A activity relative to PDE4A activity which occurs in the absence of the therapeutically effective dose. For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays or in animal models, usually mice, rabbits, dogs, or pigs. The animal model also can be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

Therapeutic efficacy and toxicity, e.g., ED₅₀ (the dose therapeutically effective in 50% of the population) and LD₅₀ (the dose lethal to 50% of the population), can be determined by standard pharmaceutical procedures in cell cultures or experimental animals. The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the ratio, LD₅₀/ED₅₀. Pharmaceutical compositions which

exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage varies
5 within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration. The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active ingredient or to maintain the desired effect. Factors which can be taken into account include the
10 severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions can be administered every 3 to 4 days, every week, or once every two weeks depending on the half-life and clearance rate of the particular formulation.

15 Normal dosage amounts can vary from 0.1 micrograms to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different
20 formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc. If the reagent is a single-chain antibody, polynucleotides encoding the antibody can be constructed and introduced into a cell either *ex vivo* or *in vivo* using well-established techniques including, but not limited to, transferrin-polycation-
25 mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated cellular fusion, intracellular transportation of DNA-coated latex beads, protoplast fusion, viral infection, electroporation, "gene gun", and DEAE- or calcium phosphate-mediated transfection.

30 If the expression product is mRNA, the reagent is preferably an antisense oligonucleotide or a ribozyme. Polynucleotides which express antisense oligonucleotides

- 79 -

or ribozymes can be introduced into cells by a variety of methods, as described above. Preferably, a reagent reduces expression of PDE4A gene or the activity of PDE4A by at least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the reagent. The effectiveness of the mechanism
5 chosen to decrease the level of expression of PDE4A gene or the activity of PDE4A can be assessed using methods well known in the art, such as hybridization of nucleotide probes to PDE4A-specific mRNA, quantitative RT-PCR, immunologic detection of PDE4A, or measurement of PDE4A activity.

10 In any of the embodiments described above, any of the pharmaceutical compositions of the invention can be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy can be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents can act syner-
15 gistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects. Any of the therapeutic methods described above can be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows,
20 horses, rabbits, monkeys, and most preferably, humans.

Nucleic acid molecules of the invention are those nucleic acid molecules which are contained in a group of nucleic acid molecules consisting of (i) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of SEQ ID
25 NO: 2, (ii) nucleic acid molecules comprising the sequence of SEQ ID NO: 1, (iii) nucleic acid molecules having the sequence of SEQ ID NO: 1, (iv) nucleic acid molecules the complementary strand of which hybridizes under stringent conditions to a nucleic acid molecule of (i), (ii), or (iii); and (v) nucleic acid molecules the sequence of which differs from the sequence of a nucleic acid molecule of (iii) due to
30 the degeneracy of the genetic code, wherein the polypeptide encoded by said nucleic acid molecule has PDE4A activity.

- 80 -

Polypeptides of the invention are those polypeptides which are contained in a group of polypeptides consisting of (i) polypeptides having the sequence of SEQ ID NO: 2, (ii) polypeptides comprising the sequence of SEQ ID NO: 2, (iii) polypeptides encoded by nucleic acid molecules of the invention and (iv) polypeptides which show at least 99%, 98%, 95%, 90%, or 80% homology with a polypeptide of (i), (ii), or (iii), wherein said purified polypeptide has PDE4A activity.

An object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising the steps of (i) contacting a test compound with a PDE4A polypeptide, (ii) detect binding of said test compound to said PDE4A polypeptide. E.g., compounds that bind to the PDE4A polypeptide are identified potential therapeutic agents for such a disease.

Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising the steps of (i) determining the activity of a PDE4A polypeptide at a certain concentration of a test compound or in the absence of said test compound, (ii) determining the activity of said polypeptide at a different concentration of said test compound. E.g., compounds that lead to a difference in the activity of the PDE4A polypeptide in (i) and (ii) are identified potential therapeutic agents for such a disease.

Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological

- 81 -

diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising the steps of (i) determining the activity of a PDE4A polypeptide at a certain concentration of a test compound, (ii) determining the activity of a PDE4A polypeptide at the presence of a compound known to be a regulator of a PDE4A polypeptide. E.g., compounds that show similar effects on the activity of the PDE4A polypeptide in (i) as compared to compounds used in (ii) are identified potential therapeutic agents for such a disease.

Other objects of the invention are methods of the above, wherein the step of contacting is in or at the surface of a cell.

Other objects of the invention are methods of the above, wherein the cell is in vitro.

Other objects of the invention are methods of the above, wherein the step of contacting is in a cell-free system.

Other objects of the invention are methods of the above, wherein the polypeptide is coupled to a detectable label.

Other objects of the invention are methods of the above, wherein the compound is coupled to a detectable label.

Other objects of the invention are methods of the above, wherein the test compound displaces a ligand which is first bound to the polypeptide.

Other objects of the invention are methods of the above, wherein the polypeptide is attached to a solid support.

Other objects of the invention are methods of the above, wherein the compound is attached to a solid support.

- 82 -

Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in
5 a mammal comprising the steps of (i) contacting a test compound with a PDE4A polynucleotide, (ii) detect binding of said test compound to said PDE4A polynucleotide. Compounds that, e.g., bind to the PDE4A polynucleotide are potential therapeutic agents for the treatment of such diseases.

10 Another object of the invention is the method of the above, wherein the nucleic acid molecule is RNA.

Another object of the invention is a method of the above, wherein the contacting step is in or at the surface of a cell.

15 Another object of the invention is a method of the above, wherein the contacting step is in a cell-free system.

Another object of the invention is a method of the above, wherein the polynucleotide
20 is coupled to a detectable label.

Another object of the invention is a method of the above, wherein the test compound is coupled to a detectable label.

25 Another object of the invention is a method of diagnosing a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising the steps of (i) determining the amount of a PDE4A polynucleotide in a sample taken
30 from said mammal, (ii) determining the amount of PDE4A polynucleotide in healthy and/or diseased mammal. A disease is diagnosed, e.g., if there is a substantial

similarity in the amount of PDE4A polynucleotide in said test mammal as compared to a diseased mammal.

5 Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising a therapeutic agent which binds to a PDE4A polypeptide.

10 Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising a therapeutic agent which regulates the activity of a PDE4A
15 polypeptide.

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases,
20 inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising a therapeutic agent which regulates the activity of a PDE4A polypeptide, wherein said therapeutic agent is (i) a small molecule, (ii) an RNA molecule, (iii) an antisense oligonucleotide, (iv) a polypeptide, (v) an antibody, or (vi) a ribozyme.

25

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal
30 comprising a PDE4A polynucleotide.

- 84 -

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising a PDE4A polypeptide.

Another object of the invention is the use of regulators of a PDE4A for the preparation of a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal.

Another object of the invention is a method for the preparation of a pharmaceutical composition useful for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising the steps of (i) identifying a regulator of PDE4A, (ii) determining whether said regulator ameliorates the symptoms of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal; and (iii) combining of said regulator with an acceptable pharmaceutical carrier.

Another object of the invention is the use of a regulator of PDE4A for the regulation of PDE4A activity in a mammal having a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease.

30

- 85 -

The examples below are provided to illustrate the subject invention. These examples are provided by way of illustration and are not included for the purpose of limiting the invention.

5 **Examples**

Example 1: Search for homologous sequences in public sequence data bases

10 The degree of homology can readily be calculated by known methods. Preferred methods to determine homology are designed to give the largest match between the sequences tested. Methods to determine homology are codified in publicly available computer programs such as BestFit, BLASTP, BLASTN, and FASTA. The BLAST programs are publicly available from NCBI and other sources in the internet.

15 For PDE4A the following hits to known sequences were identified by using the BLAST algorithm [Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ; Nucleic Acids Res 1997 Sep 1; 25(17): 3389-402] and the following set of parameters: matrix = BLOSUM62 and low complexity filter. The following databases were searched: NCBI (non-redundant database) and DERWENT patent
20 database (Geneseq).

The following hits were found:

25 >ref|NM_006202.1| Homo sapiens phosphodiesterase 4A, cAMP-specific (phosphodiesterase E2 dunce homolog, Drosophila) (PDE4A), mRNA, Length = 4255, Score = 8181 bits (4255), Expect = 0.0, Identities = 4255/4255 (100%)

30 >gb|AF069491.1|HSPDE4A5 Homo sapiens cAMP specific phosphodiesterase 4A variant pde46 (PDE4A) gene, exon 15 and 15a and alternative splice products, complete cds, Length = 3908, Score = 5280 bits (2746), Expect = 0.0, Identities = 2746/2746 (100%)

- 86 -

>gb|AC011461.5| Homo sapiens chromosome 19 clone CTC-429L19, complete sequence, Length = 100680, Score = 5238 bits (2724), Expect = 0.0, Identities = 2745/2748 (99%), Gaps = 3/2748 (0%)

5 >gb|AR083673.1|AR083673 Sequence 64 from patent US 5977305, Length = 3705, Score = 5236 bits (2723), Expect = 0.0, Identities = 2745/2751 (99%), Gaps = 4/2751 (0%)

10 >emb|AX409528.1| Sequence 2175 from Patent WO0229103, Length = 3705, Score = 5236 bits (2723), Expect = 0.0, Identities = 2745/2751 (99%), Gaps = 4/2751 (0%)

15 >emb|AX375039.1| Sequence 42 from Patent WO0210436, Length = 3705, Score = 5236 bits (2723), Expect = 0.0, Identities = 2745/2751 (99%), Gaps = 4/2751 (0%)

20 >emb|AX331581.1|AX331581 Sequence 2090 from Patent WO0194629, Length = 3705, Score = 5236 bits (2723), Expect = 0.0, Identities = 2745/2751 (99%), Gaps = 4/2751 (0%)

>gb|L20965.1|HUMPDEA Human phosphodiesterase mRNA, complete cds, Length = 3705, Score = 5236 bits (2723), Expect = 0.0, Identities = 2745/2751 (99%), Gaps = 4/2751 (0%)

25 >gb|BC038234.1| Homo sapiens, phosphodiesterase 4A, cAMP-specific (phosphodiesterase E2 dunce homolog, Drosophila), clone MGC:46146, IMAGE:5732341, mRNA, complete cds, Length = 2635, Score = 4901 bits (2549), Expect = 0.0, Identities = 2556/2557 (99%), Gaps = 1/2557 (0%)

30 >gb|BC019864.1|BC019864 Homo sapiens, phosphodiesterase 4A, cAMP-specific (phosphodiesterase E2 dunce homolog, Drosophila), clone MGC:30168, IMAGE:4942336, mRNA, complete cds, Length = 2429, Score = 4630 bits (2408), Expect = 0.0, Identities = 2424/2427 (99%), Gaps = 2/2427 (0%)

35 >gb|S75213.1|S75213 type-IVA cyclic AMP specific phosphodiesterase HPDE4A [human, T-cells, mRNA Partial, 2319 nt], Length = 2319, Score = 4197 bits (2183), Expect = 0.0, Identities = 2192/2194 (99%), Gaps = 1/2194 (0%)

40

- 87 -

- >gb|M37744.1|HUMPDEAA Human cyclic AMP (cAMP)-specific phosphodiesterase mRNA, complete cds, Length = 2619, Score = 4142 bits (2154), Expect = 0.0, Identities = 2162/2166 (99%)
- 5 >gb|I22486.1|I22486 Sequence 21 from patent US 5527896, Length = 3131, Score = 3803 bits (1978), Expect = 0.0, Identities = 1987/1989 (99%), Gaps = 1/1989 (0%)
- 10 >gb|L20967.1|HUMPDEC Homo sapiens phosphodiesterase gene, partial cds, Length = 3131, Score = 3803 bits (1978), Expect = 0.0, Identities = 1987/1989 (99%), Gaps = 1/1989 (0%)
- 15 >gb|U97584.1|HSU97584 Homo sapiens cyclic AMP-specific phosphodiesterase HSPDE4A1A (PDE4A), mRNA, complete cds, Length = 1965, Score = 3773 bits (1962), Expect = 0.0, Identities = 1964/1965 (99%)
- 20 >emb|AX348237.1| Sequence 3 from Patent WO0203072, Length = 1949, Score = 3724 bits (1937), Expect = 0.0, Identities = 1941/1943 (99%)
- >emb|AX283141.1|AX283141 Sequence 29 from Patent WO0179526, Length = 2682, Score = 3724 bits (1937), Expect = 0.0, Identities = 1941/1943 (99%)
- 25 >emb|AX283113.1|AX283113 Sequence 1 from Patent WO0179526, Length = 2682, Score = 3724 bits (1937), Expect = 0.0, Identities = 1941/1943 (99%)
- 30 >gb|U18087.1|HSU18087 Human 3',5'-cyclic AMP phosphodiesterase HPDE4A6 mRNA, complete cds, Length = 2091, Score = 3632 bits (1889), Expect = 0.0, Identities = 1893/1895 (99%)
- 35 >gb|AF073745.2|AF073745 Homo sapiens cyclic AMP-specific phosphodiesterase HSPDE4A10 (PDE4A), mRNA, complete cds, Length = 2478, Score = 3609 bits (1877), Expect = 0.0, Identities = 1879/1880 (99%)
- 40 >emb|AX348235.1| Sequence 1 from Patent WO0203072, Length = 2666, Score = 3601 bits (1873), Expect = 0.0, Identities = 1877/1879 (99%)

- 88 -

>emb|AX283143.1|AX283143 Sequence 31 from Patent WO0179526, Length = 3399, Score = 3601 bits (1873), Expect = 0.0, Identities = 1877/1879 (99%)

5 >emb|AX283115.1|AX283115 Sequence 3 from Patent WO0179526, Length = 3399, Score = 3601 bits (1873), Expect = 0.0, Identities = 1877/1879 (99%)

10 >emb|AX283119.1|AX283119 Sequence 7 from Patent WO0179526, Length = 3399, Score = 3590 bits (1867), Expect = 0.0, Identities = 1875/1879 (99%)

15 >gb|AR083686.1|AR083686 Sequence 79 from patent US 5977305, Length = 2041, Score = 3526 bits (1834), Expect = 0.0, Identities = 1836/1837 (99%)

20 >emb|AX283117.1|AX283117 Sequence 5 from Patent WO0179526, Length = 3375, Score = 3257 bits (1694), Expect = 0.0, Identities = 1698/1700 (99%)

>gb|I22481.1|I22481 Sequence 11 from patent US 5527896, Length = 2702, Score = 2661 bits (1384), Expect = 0.0, Identities = 1395/1398 (99%), Gaps = 2/1398 (0%)

25 >gb|AR083659.1|AR083659 Sequence 39 from patent US 5977305, Length = 1230, Score = 2246 bits (1168), Expect = 0.0, Identities = 1206/1218 (99%), Gaps = 8/1218 (0%)

30 >gb|I22496.1|I22496 Sequence 39 from patent US 5527896, Length = 1230, Score = 2246 bits (1168), Expect = 0.0, Identities = 1206/1218 (99%), Gaps = 8/1218 (0%)

35 >gb|AR083693.1|AR083693 Sequence 87 from patent US 5977305, Length = 1481, Score = 2148 bits (1117), Expect = 0.0, Identities = 1119/1120 (99%)

>gb|U18088.1|HSU18088 Human 3',5'-cyclic AMP phosphodiesterase inactive splice variant HSPDE4A7A mRNA, complete cds, Length = 1770, Score = 1586 bits (825), Expect = 0.0, Identities = 829/831 (99%)

Example 2: Expression profiling

Total cellular RNA was isolated from cells by one of two standard methods: 1) 5 guanidine isothiocyanate/Cesium chloride density gradient centrifugation [Kellogg, (1990)]; or with the Tri-Reagent protocol according to the manufacturer's specifications (Molecular Research Center, Inc., Cincinnati, Ohio). Total RNA prepared by the Tri-reagent protocol was treated with DNase I to remove genomic DNA contamination.

10 For relative quantitation of the mRNA distribution of PDE4A, total RNA from each cell or tissue source was first reverse transcribed. 85 µg of total RNA was reverse transcribed using 1 µmole random hexamer primers, 0.5 mM each of dATP, dCTP, dGTP and dTTP (Qiagen, Hilden, Germany), 3000 U RnaseQut (Invitrogen, Groningen, Netherlands) in a final volume of 680 µl. The first strand synthesis 15 buffer and Omniscript reverse transcriptase (2 u/µl) were from (Qiagen, Hilden, Germany). The reaction was incubated at 37°C for 90 minutes and cooled on ice. The volume was adjusted to 6800 µl with water, yielding a final concentration of 12.5 ng/µl of starting RNA.

20 For relative quantitation of the distribution of PDE4A mRNA in cells and tissues the Perkin Elmer ABI Prism RTM. 7700 Sequence Detection system or Biorad iCycler was used according to the manufacturer's specifications and protocols. PCR reactions were set up to quantitate PDE4A and the housekeeping genes HPRT 25 (hypoxanthine phosphoribosyltransferase), GAPDH (glyceraldehyde-3-phosphate dehydrogenase), β-actin, and others. Forward and reverse primers and probes for PDE4A were designed using the Perkin Elmer ABI Primer ExpressTM software and were synthesized by TibMolBiol (Berlin, Germany). The PDE4A forward primer sequence was: Primer1 (SEQ ID NO: 3). The PDE4A reverse primer sequence was 30 Primer2 (SEQ ID NO: 5). Probe1 (SEQ ID NO: 4), labelled with FAM (carboxyfluorescein succinimidyl ester) as the reporter dye and TAMRA

(carboxytetramethylrhodamine) as the quencher, is used as a probe for PDE4A. The following reagents were prepared in a total of 25 μ l : 1x TaqMan buffer A, 5.5 mM MgCl₂, 200 nM of dATP, dCTP, dGTP, and dUTP, 0.025 U/ μ l AmpliTaq GoldTM, 0.01 U/ μ l AmpErase and Probe1 (SEQ ID NO: 4), PDE4A forward and reverse
 5 primers each at 200 nM, 200 nM PDE4A FAM/TAMRA-labelled probe, and 5 μ l of template cDNA. Thermal cycling parameters were 2 min at 50°C, followed by 10 min at 95°C, followed by 40 cycles of melting at 95°C for 15 sec and annealing/extending at 60°C for 1 min.

10 *Calculation of corrected CT values*

The CT (threshold cycle) value is calculated as described in the "Quantitative determination of nucleic acids" section. The CF-value (factor for threshold cycle correction) is calculated as follows :

15

1. PCR reactions were set up to quantitate the housekeeping genes (HKG) for each cDNA sample.

20

2. CT_{HKG}-values (threshold cycle for housekeeping gene) were calculated as described in the "Quantitative determination of nucleic acids" section.

3. CT_{HKG}-mean values (CT mean value of all HKG tested on one cDNAs) of all HKG for each cDNA are calculated (n = number of HKG):

25

$$CT_{HKG-n}\text{-mean value} = (CT_{HKG1}\text{-value} + CT_{HKG2}\text{-value} + \dots + CT_{HKG-n}\text{-value}) / n$$

4. CT_{panel} mean value (CT mean value of all HKG in all tested cDNAs) =

$$(CT_{HKG1}\text{-mean value} + CT_{HKG2}\text{-mean value} + \dots + CT_{HKG-y}\text{-mean value}) / y$$

30

(y = number of cDNAs)

- 91 -

5. CF_{cDNA-n} (correction factor for cDNA n) = $CT_{panel-mean\ value} - CT_{HKG-n}$
mean value
6. CT_{cDNA-n} (CT value of the tested gene for the cDNA n) + CF_{cDNA-n} (correction
5 factor for cDNA n) = $CT_{cor-cDNA-n}$ (corrected CT value for a gene on cDNA n)

Calculation of relative expression

Definition : highest $CT_{cor-cDNA-n} \neq 40$ is defined as $CT_{cor-cDNA} [high]$
10 Relative Expression = $2^{(CT_{cor-cDNA} [high] - CT_{cor-cDNA-n})}$

Tissues

The expression of PDE4A was investigated in the tissues listed in table 1.
15

Expression profile

The results of the the mRNA-quantification (expression profiling) is shown in
Table 1.
20

Table 1: Relative expression of PDE4A in various human tissues

<u>Tissue</u>	<u>Relative Expression</u>
fetal heart	85
25 heart	256
pericardium	367
heart atrium (right)	28
heart atrium (left)	18
heart ventricle (left)	16
30 interventricular septum	33
fetal aorta	1
aorta	4

- 92 -

	aorta sclerotic	8
	artery	10
	coronary artery	10
	coronary artery sclerotic	15
5	vein	6
	coronary artery smooth muscle primary cells	123
	HUVEC cells	89
	skin	40
	adrenal gland	51
10	thyroid	560
	thyroid tumor	261
	pancreas	91
	pancreas liver cirrhosis	422
	esophagus	29
15	esophagus tumor	18
	stomach	124
	stomach tumor	39
	colon	261
	colon tumor	54
20	small intestine	30
	ileum	19
	ileum tumor	4
	ileum chronic inflammation	26
	rectum	174
25	salivary gland	14
	fetal liver	12
	liver	61
	liver liver cirrhosis	320
	liver tumor	11
30	HEP G2 cells	159
	leukocytes (peripheral blood)	131
	Jurkat (T-cells)	23
	bone marrow	27
	erythrocytes	8

- 93 -

	lymphnode	15
	thymus	96
	thrombocytes	8
	bone marrow stromal cells	18
5	bone marrow CD71+ cells	4
	bone marrow CD33+ cells	17
	bone marrow CD34+ cells	4
	bone marrow CD15+ cells	7
	cord blood CD71+ cells	4
10	spleen	143
	spleen liver cirrhosis	765
	skeletal muscle	244
	adipose	172
	fetal brain	434
15	brain	258
	alzheimer brain	588
	cerebellum	26
	cerebellum (right)	54
	cerebellum (left)	64
20	cerebral cortex	123
	alzheimer cerebral cortex	690
	frontal lobe	70
	alzheimer brain frontal lobe	76
	occipital lobe	367
25	parietal lobe	471
	temporal lobe	352
	precentral gyrus	396
	postcentral gyrus	11
	tonsilla cerebelli	62
30	vermis cerebelli	79
	pons	36
	substantia nigra	290
	cerebral meninges	5
	cerebral peduncles	280

- 94 -

	corpus callosum	20
	hippocampus	372
	thalamus	92
	dorsal root ganglia	22
5	spinal cord	191
	neuroblastoma SK-N-MC cells	43
	neuroblastoma SH-SY5Y cells	19
	neuroblastoma IMR32 cells	32
	glial tumor H4 cells	16
10	retina	21
	fetal lung	419
	fetal lung fibroblast IMR-90 cells	13
	lung	352
	lung right upper lobe	23
15	lung right mid lobe	23
	lung right lower lobe	32
	lung tumor	128
	lung COPD	9
	trachea	98
20	cervix	13
	testis	137
	HeLa cells (cervix tumor)	49
	placenta	82
	uterus	68
25	uterus tumor	10
	ovary	23
	ovary tumor	14
	breast	80
	breast tumor	136
30	MDA MB 231 cells (breast tumor)	116
	mammary gland	202
	prostate	47
	prostate BPH	6
	bladder	55

- 95 -

	ureter	60
	penis	8
	corpus cavernosum	10
	fetal kidney	108
5	kidney	147
	kidney tumor	30
	HEK 293 cells	180

Example 3: Antisense Analysis

10

Knowledge of the correct, complete cDNA sequence coding for PDE4A enables its use as a tool for antisense technology in the investigation of gene function. Oligonucleotides, cDNA or genomic fragments comprising the antisense strand of a polynucleotide coding for PDE4A are used either in vitro or in vivo to inhibit translation of the mRNA. Such technology is now well known in the art, and antisense molecules can be designed at various locations along the nucleotide sequences. By treatment of cells or whole test animals with such antisense sequences, the gene of interest is effectively turned off. Frequently, the function of the gene is ascertained by observing behavior at the intracellular, cellular, tissue or organismal level (e.g., lethality, loss of differentiated function, changes in morphology, etc.).

15

20

In addition to using sequences constructed to interrupt transcription of a particular open reading frame, modifications of gene expression is obtained by designing antisense sequences to intron regions, promoter/enhancer elements, or even to trans-acting regulatory genes.

25

Example 4: Expression of PDE4A

30

Expression of PDE4A is accomplished by subcloning the cDNAs into appropriate expression vectors and transfecting the vectors into expression hosts such as, e.g., *E. coli*. In a particular case, the vector is engineered such that it contains a promoter for

- 96 -

β -galactosidase, upstream of the cloning site, followed by sequence containing the amino-terminal Methionine and the subsequent seven residues of β -galactosidase. Immediately following these eight residues is an engineered bacteriophage promoter useful for artificial priming and transcription and for providing a number of unique endonuclease restriction sites for cloning.

Induction of the isolated, transfected bacterial strain with Isopropyl- β -D-thiogalactopyranoside (IPTG) using standard methods produces a fusion protein corresponding to the first seven residues of β -galactosidase, about 15 residues of "linker", and the peptide encoded within the cDNA. Since cDNA clone inserts are generated by an essentially random process, there is probability of 33% that the included cDNA will lie in the correct reading frame for proper translation. If the cDNA is not in the proper reading frame, it is obtained by deletion or insertion of the appropriate number of bases using well known methods including in vitro mutagenesis, digestion with exonuclease III or mung bean nuclease, or the inclusion of an oligonucleotide linker of appropriate length.

The PDE4A cDNA is shuttled into other vectors known to be useful for expression of proteins in specific hosts. Oligonucleotide primers containing cloning sites as well as a segment of DNA (about 25 bases) sufficient to hybridize to stretches at both ends of the target cDNA is synthesized chemically by standard methods. These primers are then used to amplify the desired gene segment by PCR. The resulting gene segment is digested with appropriate restriction enzymes under standard conditions and isolated by gel electrophoresis. Alternately, similar gene segments are produced by digestion of the cDNA with appropriate restriction enzymes. Using appropriate primers, segments of coding sequence from more than one gene are ligated together and cloned in appropriate vectors. It is possible to optimize expression by construction of such chimeric sequences.

Suitable expression hosts for such chimeric molecules include, but are not limited to, mammalian cells such as Chinese Hamster Ovary (CHO) and human 293 cells.,

- 97 -

insect cells such as Sf9 cells, yeast cells such as *Saccharomyces cerevisiae* and bacterial cells such as *E. coli*. For each of these cell systems, a useful expression vector also includes an origin of replication to allow propagation in bacteria, and a selectable marker such as the β -lactamase antibiotic resistance gene to allow plasmid selection in bacteria. In addition, the vector may include a second selectable marker such as the neomycin phosphotransferase gene to allow selection in transfected eukaryotic host cells. Vectors for use in eukaryotic expression hosts require RNA processing elements such as 3' polyadenylation sequences if such are not part of the cDNA of interest.

Additionally, the vector contains promoters or enhancers which increase gene expression. Such promoters are host specific and include MMTV, SV40, and metallothionine promoters for CHO cells; trp, lac, tac and T7 promoters for bacterial hosts; and alpha factor, alcohol oxidase and PGH promoters for yeast. Transcription enhancers, such as the rous sarcoma virus enhancer, are used in mammalian host cells. Once homogeneous cultures of recombinant cells are obtained through standard culture methods, large quantities of recombinantly produced PDE4A are recovered from the conditioned medium and analyzed using chromatographic methods known in the art. For example, PDE4A can be cloned into the expression vector pcDNA3, as exemplified herein. This product can be used to transform, for example, HEK293 or COS by methodology standard in the art. Specifically, for example, using Lipofectamine (Gibco BRL catalog no. 18324-020) mediated gene transfer.

Example 5: Isolation of Recombinant PDE4A

PDE4A is expressed as a chimeric protein with one or more additional polypeptide domains added to facilitate protein purification. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals [Appa Rao, 1997] and the domain utilized in the FLAGS extension/affinity purification system

- 98 -

(Immunex Corp., Seattle, Washington). The inclusion of a cleavable linker sequence such as Factor Xa or enterokinase (Invitrogen, Groningen, The Netherlands) between the purification domain and the PDE4A sequence is useful to facilitate expression of PDE4A.

5

The following example provides a method for purifying PDE4A.

PDE4A is generated using the baculovirus expression system BAC-TO-BAC (GIBCO BRL) based on *Autographa californica* nuclear polyhedrosis virus (AcNPV) infection of *Spodoptera frugiperda* insect cells (Sf9 cells).

10

cDNA encoding PDE is cloned into either the donor plasmid pFASTBAC1 or pFASTBAC-HT which contain a mini-Tn7 transposition element. The recombinant plasmid is transformed into DH10BAC competent cells which contain the parent bacmid bMON14272 (AcNPV infectious DNA) and a helper plasmid. The mini-Tn7 element on the pFASTBAC donor can transpose to the attTn7 attachment site on the bacmid thus introducing the PDE gene into the viral genome. Colonies containing recombinant bacmids are identified by disruption of the *lacZ* gene. The PDE/bacmid construct can then be isolated and infected into insect cells (Sf9 cells) resulting in the production of infectious recombinant baculovirus particles and expression of either unfused recombinant enzyme (pFastbac1) or PDE4A-His fusion protein (pFastbacHT).

15

20

Cells are harvested and extracts prepared 24, 48 and 72 hours after transfection. Expression of PDE4A is confirmed by coomassie staining after sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) and western blotting onto a PVDF membrane of an unstained SDS-PAGE. The PDE-His fusion protein is detected due to the interaction between the Ni-NTA HRP conjugate and the His-tag which is fused to PDE4A.

25

30

Example 6: Production of PDE4A Specific Antibodies

Two approaches are utilized to raise antibodies to PDE4A, and each approach is useful for generating either polyclonal or monoclonal antibodies. In one approach, denatured protein from reverse phase HPLC separation is obtained in quantities up to 75 mg. This denatured protein is used to immunize mice or rabbits using standard protocols; about 100 µg are adequate for immunization of a mouse, while up to 1 mg might be used to immunize a rabbit. For identifying mouse hybridomas, the denatured protein is radioiodinated and used to screen potential murine B-cell hybridomas for those which produce antibody. This procedure requires only small quantities of protein, such that 20 mg is sufficient for labeling and screening of several thousand clones.

In the second approach, the amino acid sequence of an appropriate PDE4A domain, as deduced from translation of the cDNA, is analyzed to determine regions of high antigenicity. Oligopeptides comprising appropriate hydrophilic regions are synthesized and used in suitable immunization protocols to raise antibodies. The optimal amino acid sequences for immunization are usually at the C-terminus, the N-terminus and those intervening, hydrophilic regions of the polypeptide which are likely to be exposed to the external environment when the protein is in its natural conformation.

Typically, selected peptides, about 15 residues in length, are synthesized using an Applied Biosystems Peptide Synthesizer Model 431A using fmoc-chemistry and coupled to keyhole limpet hemocyanin (KLH; Sigma, St. Louis, MO) by reaction with M-maleimidobenzoyl-N-hydroxysuccinimide ester, MBS. If necessary, a cysteine is introduced at the N-terminus of the peptide to permit coupling to KLH. Rabbits are immunized with the peptide-KLH complex in complete Freund's adjuvant. The resulting antisera are tested for antipeptide activity by binding the peptide to plastic, blocking with 1% bovine serum albumin, reacting with antisera,

- 100 -

washing and reacting with labeled (radioactive or fluorescent), affinity purified, specific goat anti-rabbit IgG.

Hybridomas are prepared and screened using standard techniques. Hybridomas of
5 interest are detected by screening with labeled PDE4A to identify those fusions
producing the monoclonal antibody with the desired specificity. In a typical
protocol, wells of plates (FAST; Becton-Dickinson, Palo Alto, CA) are coated during
incubation with affinity purified, specific rabbit anti-mouse (or suitable antispecies
1 g) antibodies at 10 mg/ml. The coated wells are blocked with 1% bovine serum
10 albumin, (BSA), washed and incubated with supernatants from hybridomas. After
washing the wells are incubated with labeled PDE4A at 1 mg/ml. Supernatants with
specific antibodies bind more labeled PDE4A than is detectable in the background.
Then clones producing specific antibodies are expanded and subjected to two cycles
of cloning at limiting dilution. Cloned hybridomas are injected into pristane-treated
15 mice to produce ascites, and monoclonal antibody is purified from mouse ascitic
fluid by affinity chromatography on Protein A. Monoclonal antibodies with affinities
of at least

10^8 M^{-1} , preferably 10^9 to 10^{10} M^{-1} or stronger, are typically made by standard
20 procedures.

Example 7: Diagnostic Test Using PDE4A Specific Antibodies

Particular PDE4A antibodies are useful for investigating signal transduction and the
25 diagnosis of infectious or hereditary conditions which are characterized by
differences in the amount or distribution of PDE4A or downstream products of an
active signaling cascade.

Diagnostic tests for PDE4A include methods utilizing antibody and a label to detect
30 PDE4A in human body fluids, membranes, cells, tissues or extracts of such. The
polypeptides and antibodies of the present invention are used with or without

modification. Frequently, the polypeptides and antibodies are labeled by joining them, either covalently or noncovalently, with a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and have been reported extensively in both the scientific and patent literature.

5 Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent agents, chemiluminescent agents, chromogenic agents, magnetic particles and the like.

A variety of protocols for measuring soluble or membrane-bound PDE4A, using either polyclonal or monoclonal antibodies specific for the protein, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and fluorescent activated cell sorting (FACS). A two-site monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on PDE4A is preferred, but a competitive binding assay may be employed.

15

Example 8: Purification of Native PDE4A Using Specific Antibodies

Native or recombinant PDE4A is purified by immunoaffinity chromatography using antibodies specific for PDE4A. In general, an immunoaffinity column is constructed by covalently coupling the anti-TRH antibody to an activated chromatographic resin.

20

Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate or by purification on immobilized Protein A (Pharmacia LKB Biotechnology, Piscataway N.J.). Likewise, monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated Sepharose (Pharmacia LKB Biotechnology). The antibody is coupled to the resin, the resin is blocked, and the derivative resin is washed according to the manufacturer's instructions.

25
30

Such immunoaffinity columns are utilized in the purification of PDE4A by preparing a fraction from cells containing PDE4A in a soluble form. This preparation is derived by solubilization of whole cells or of a subcellular fraction obtained via differential centrifugation (with or without addition of detergent) or by other methods well known in the art. Alternatively, soluble PDE4A containing a signal sequence is secreted in useful quantity into the medium in which the cells are grown.

A soluble PDE4A-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PDE4A (e.g., high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions that disrupt antibody/protein binding (e.g., a buffer of pH 2-3 or a high concentration of a chaotrope such as urea or thiocyanate ion), and PDE4A is collected.

Example 9: Drug Screening

This invention is particularly useful for screening therapeutic compounds by using PDE4A or fragments thereof in any of a variety of drug screening techniques.

The following example provides a system for drug screening measuring the phosphodiesterase activity.

The phosphodiesterase activity of the crude extracts is measured and confirmed that the PDE cDNA encodes a phosphodiesterase which is able to hydrolyze cAMP or cGMP or both.

The recombinant PDE-His fusion protein can be purified from the crude lysate by metal-affinity chromatography using Ni-NTA agarose. This allows the specific retention of the recombinant material (since this is fused to the His-tag) whilst the endogenous insect proteins are washed off. The recombinant material is then eluted by competition with imidazol.

The phosphodiesterase activity of the recombinant protein is assayed using a commercially available SPA (scintillation proximity assay) kit (Amersham Pharmacia). The PDE enzyme hydrolyzes cyclic nucleotides, e.g. cAMP and cGMP to their linear counterparts. The SPA assay utilizes the tritiated cyclic nucleotides $[^3\text{H}]\text{cAMP}$ or $[^3\text{H}]\text{cGMP}$, and is based upon the selective interaction of the tritiated non cyclic product with the SPA beads whereas the cyclic substrates are not effectively binding. Radiolabelled product bound to the scintillation beads generates light that can be analyzed in a scintillation counter.

Example 10: Rational Drug Design

The goal of rational drug design is to produce structural analogs of biologically active polypeptides of interest or of small molecules with which they interact, agonists, antagonists, or inhibitors. Any of these examples are used to fashion drugs which are more active or stable forms of the polypeptide or which enhance or interfere with the function of a polypeptide in vivo.

In one approach, the three-dimensional structure of a protein of interest, or of a protein-inhibitor complex, is determined by x-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the polypeptide must be ascertained to elucidate the structure and to determine active site(s) of the molecule. Less often, useful information regarding the structure of a polypeptide is gained by modeling based on the structure of homologous proteins. In both cases, relevant structural information is used to design efficient inhibitors. Useful examples of rational drug design include molecules which have improved activity or stability or which act as inhibitors, agonists, or antagonists of native peptides.

It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle,

- 104 -

yields a pharmacore upon which subsequent drug design is based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids is expected to be an analog of the original receptor. The anti-id is then used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides then act as the pharmacore.

By virtue of the present invention, sufficient amount of polypeptide are made available to perform such analytical studies as X-ray crystallography. In addition, knowledge of the PDE4A amino acid sequence provided herein provides guidance to those employing computer modeling techniques in place of or in addition to x-ray crystallography.

Example 11: Identification of Other Members of the Signal Transduction Complex

Labeled PDE4A is useful as a reagent for the purification of molecules with which it interacts. In one embodiment of affinity purification, PDE4A is covalently coupled to a chromatography column. Cell-free extract derived from synovial cells or putative target cells is passed over the column, and molecules with appropriate affinity bind to PDE4A. PDE4A-complex is recovered from the column, and the PDE4A-binding ligand disassociated and subjected to N-terminal protein sequencing. The amino acid sequence information is then used to identify the captured molecule or to design degenerate oligonucleotide probes for cloning the relevant gene from an appropriate cDNA library.

In an alternate method, antibodies are raised against PDE4A, specifically monoclonal antibodies. The monoclonal antibodies are screened to identify those which inhibit the binding of labeled PDE4A. These monoclonal antibodies are then used therapeutically.

Example 12: Use and Administration of Antibodies, Inhibitors, or Antagonists

Antibodies, inhibitors, or antagonists of PDE4A or other treatments and compounds
5 that are limiters of signal transduction (LSTs), provide different effects when
administered therapeutically. LSTs are formulated in a nontoxic, inert, pharma-
ceutically acceptable aqueous carrier medium preferably at a pH of about 5 to 8,
more preferably 6 to 8, although pH may vary according to the characteristics of the
antibody, inhibitor, or antagonist being formulated and the condition to be treated.
10 Characteristics of LSTs include solubility of the molecule, its half-life and
antigenicity/immunogenicity. These and other characteristics aid in defining an
effective carrier. Native human proteins are preferred as LSTs, but organic or
synthetic molecules resulting from drug screens are equally effective in particular
situations.

15 LSTs are delivered by known routes of administration including but not limited to
topical creams and gels; transmucosal spray and aerosol; transdermal patch and
bandage; injectable, intravenous and lavage formulations; and orally administered
liquids and pills particularly formulated to resist stomach acid and enzymes. The
20 particular formulation, exact dosage, and route of administration is determined by the
attending physician and varies according to each specific situation.

Such determinations are made by considering multiple variables such as the
condition to be treated, the LST to be administered, and the pharmacokinetic profile
25 of a particular LST. Additional factors which are taken into account include severity
of the disease state, patient's age, weight, gender and diet, time and frequency of LST
administration, possible combination with other drugs, reaction sensitivities, and
tolerance/response to therapy. Long acting LST formulations might be administered
every 3 to 4 days, every week, or once every two weeks depending on half-life and
30 clearance rate of the particular LST.

- 106 -

Normal dosage amounts vary from 0.1 to 10^5 μ g, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature; see U.S. Pat. Nos. 4,657,760; 5,206,344; or 5,225,212. Those skilled in the art employ different formulations for different LSTs. Administration to cells such as nerve cells necessitates delivery in a manner different from that to other cells such as vascular endothelial cells.

It is contemplated that abnormal signal transduction, trauma, or diseases which trigger PDE4A activity are treatable with LSTs. These conditions or diseases are specifically diagnosed by the tests discussed above, and such testing should be performed in suspected cases of viral, bacterial or fungal infections, allergic responses, mechanical injury associated with trauma, hereditary diseases, lymphoma or carcinoma, or other conditions which activate the genes of lymphoid or neuronal tissues.

Example 13: Production of Non-human Transgenic Animals

Animal model systems which elucidate the physiological and behavioral roles of the PDE4A are produced by creating nonhuman transgenic animals in which the activity of the PDE4A is either increased or decreased, or the amino acid sequence of the expressed PDE4A is altered, by a variety of techniques. Examples of these techniques include, but are not limited to: 1) Insertion of normal or mutant versions of DNA encoding a PDE4A, by microinjection, electroporation, retroviral transfection or other means well known to those skilled in the art, into appropriately fertilized embryos in order to produce a transgenic animal or 2) homologous recombination of mutant or normal, human or animal versions of these genes with the native gene locus in transgenic animals to alter the regulation of expression or the structure of these PDE4A sequences. The technique of homologous recombination is well known in the art. It replaces the native gene with the inserted gene and hence is useful for producing an animal that cannot express native PDE4As but does express, for example, an inserted mutant PDE4A, which has replaced the native PDE4A in the

animal's genome by recombination, resulting in underexpression of the transporter. Microinjection adds genes to the genome, but does not remove them, and the technique is useful for producing an animal which expresses its own and added PDE4A, resulting in overexpression of the PDE4A.

5

One means available for producing a transgenic animal, with a mouse as an example, is as follows: Female mice are mated, and the resulting fertilized eggs are dissected out of their oviducts. The eggs are stored in an appropriate medium such as cesiumchloride M2 medium. DNA or cDNA encoding PDE4A is purified from a vector by methods well known to the one skilled in the art. Inducible promoters may be fused with the coding region of the DNA to provide an experimental means to regulate expression of the transgene. Alternatively or in addition, tissue specific regulatory elements may be fused with the coding region to permit tissue-specific expression of the transgene. The DNA, in an appropriately buffered solution, is put into a microinjection needle (which may be made from capillary tubing using a piper puller) and the egg to be injected is put in a depression slide. The needle is inserted into the pronucleus of the egg, and the DNA solution is injected. The injected egg is then transferred into the oviduct of a pseudopregnant mouse which is a mouse stimulated by the appropriate hormones in order to maintain false pregnancy, where it proceeds to the uterus, implants, and develops to term. As noted above, microinjection is not the only method for inserting DNA into the egg but is used here only for exemplary purposes.

10

15

20

References

25

U.S. 4,522,811

U.S. 5,283,317

U.S. 5,527,896

U.S. 5,565,332

30

U.S. 5,977,305

WO 84/03564

- WO 92/01810
WO 93/03151
WO 94/13804
WO 01/04297
5 WO 01/79526
WO 01/94629
WO 02/03072
WO 02/10436
WO 02/29103
10 Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ;
Nucleic Acids Res 1997 Sep 1; 25(17): 3389-402
Appa Rao et al., 1997, Protein Expr Purif Nov, 11(2): 201-8
Bang, Y.J. et al. (1994)Proc. Natl. Acad. Sci. USA 91:5330-5334
Bardelle, C. et al. (1999) Anal. Biochem. 275: 148-155
15 Barnes, 2000, Chest, 117:10S14S
Beavo, Physiol. Rev . 75:725-748 (1995);
Botstein et al., 1980, Am J Hum Genet. 32: 314-31
Charbonneau, et al., Proc. Natl. Acad. Sci . (USA) 83:9308-9312 (1986)
Colbere-Garapin *et al.*, 1981, *J. Mol. Biol.* 150, 1-14
20 Conti, et al., Physiol. Rev . 75:723-748 (1995)].
Conti et al. (1995) Endocrine Rev. 16:370-389
Engelhard *et al.*, 1994, *Proc. Nat. Acad. Sci.* 91, 3224-3227
Francis, et al., J. Biol. Chem . 269:22477-22480 (1994).
Fujishige, K. et al. (1999) J. Biol. Chem. 274:18438-18445
25 Gergen and Weiss , 1992, Am Rev Respir Dis 146:823-824
Gibson et al., 1996, Genome Research 6: 995-1001
Han, P. et al. (1997) J. Biol. Chem. 272:16152-16157
Haseloff *et al.*, 1988, *Nature* 334, 585-591
Harbinson, et al., Eur. Respir. J . 10:1008-1014 (1997)
30 Heid et al., 1996, Genome Research 6: 986-994

- Hetman JM, Robas N, Baxendale R, Fidock M, Phillips SC, Soderling SH, Beavo JA
Proc. Natl. Acad. Sci. U S A 2000 Nov 7;97(23):12891-5
- Hansen, R.S., and Beavo, J.A., PNAS USA 1982;79: 2788-92
- Holland et al., 1991, PNAS 88: 7276-7280
- 5 Iwabuchi *et al.*, 1993, *Oncogene* 8, 1693-1696
- Jeffreys et al., 1985, *Nature* 316: 76-9
- Johnson *et al.*, 1989, *Endoc. Rev.* 10, 317-331
- Kakkar, R. et al. (1999) *Cell Mol Life Sci* 55:1164-1186
- Kellogg et al., 1990, *Anal. Biochem.* 189:202-208
- 10 Lam , 1997, *Anticancer Drug Res.* 12(3):145-67
- Lamontagne S et al., *Brain Res* 2001 Nov 30;920(1-2):84-96
- Livak et al., 1995 , *PCR Methods and Applications* 357-362
- Livi GP et al., *Mol Cell Biol* 1990 Jun;10(6):2678-86
- Logan, Shenk, 1984, *Proc. Natl. Acad. Sci.* 81, 3655-3659
- 15 Loughney and Ferguson, in *Phosphodiesterase Inhibitors*, Schudt, et al. (Eds.),
Academic Press: New York, N.Y. (1996) pp. 1- 19
- Loughney, K. et al (1999) *Gene* 234:109-117
- Lowy *et al.*, 1980, *Cell* 22, 817-23
- Maddox *et al.*, 1983, *J. Exp. Med.* 158, 1211-1216
- 20 McAllister-Lucas, L.M. (1995) *J. Biol. Chem.* 270:30671-30679
- McConnell *et al.*, 1992, *Science* 257, 1906-1912
- Manganiello, et al., *Arch. Biochem. Acta* 322:1-13 (1995);
- Nicholls *et al.*, 1993, *J. Immunol. Meth.* 165, 81-91
- Perry, M.J. and G.A. Higgs (1998) *Curr. Opin. Chem. Biol.* 2:472-481
- 25 Piatak et al., 1993, *BioTechniques* 14:70-81
- Piatak et al., 1993, *Science* 259:1749-1754
- Porath *et al.*, 1992, *Prot. Exp. Purif.* 3, 263-281
- Roberge *et al.*, 1995, *Science* 269, 202-204
- Sadhu, K. et al. (1999) *J. Histochem Cytochem* 47:895-906
- 30 Scott and Smith (1990) *Science* 249:386-390
- Sjolander, Urbaniczky, 1991, *Anal. Chem.* 63, 2338-2345

- 110 -

- Soderling, S.H. et al. (1999) *Proc. Natl. Acad. Sci. USA* 96:7071-7076
- Sullivan M et al., *Biochem J* 1998 Aug 1;333 (Pt 3):693-703
- Szabo *et al.*, 1995, *Curr. Opin. Struct. Biol.* 5, 699-705
- Terrett, N. et al. (1996) *Bioorg. Med. Chem. Lett.* 6:1819-1824
- 5 Thomas, 1980, *Proc. Nat. Acad. Sci.*, 77:5201-5205
- Turko, I.V. et al. (1998) *Biochemistry* 37:4200-4205
- Uhlmann *et al.*, 1987, *Tetrahedron. Lett.* 215, 3539-3542
- Weber et al., 1990, *Genomics* 7: 524-30
- Wigler *et al.*, 1977, *Cell* 11, 223-32
- 10 Wigler *et al.*, 1980, *Proc. Natl. Acad. Sci.* 77, 3567-70

Claims

1. A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising the steps of
- 5
- i) contacting a test compound with a PDE4A polypeptide,
- 10
- ii) detect binding of said test compound to said PDE4A polypeptide.
2. A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising the steps of
- 15
- i) determining the activity of a PDE4A polypeptide at a certain concentration of a test compound or in the absence of said test compound,
- 20
- ii) determining the activity of said polypeptide at a different concentration of said test compound.
- 25
3. A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising the steps of
- 30

- 112 -

- i) determining the activity of a PDE4A polypeptide at a certain concentration of a test compound,
 - ii) determining the activity of a PDE4A polypeptide at the presence of a compound known to be a regulator of a PDE4A polypeptide.
- 4. The method of any of claims 1 to 3, wherein the step of contacting is in or at the surface of a cell.
- 5. The method of any of claims 1 to 3, wherein the cell is in vitro.
- 6. The method of any of claims 1 to 3, wherein the step of contacting is in a cell-free system.
- 7. The method of any of claims 1 to 3, wherein the polypeptide is coupled to a detectable label.
- 8. The method of any of claims 1 to 3, wherein the compound is coupled to a detectable label.
- 9. The method of any of claims 1 to 3, wherein the test compound displaces a ligand which is first bound to the polypeptide.
- 10. The method of any of claims 1 to 3, wherein the polypeptide is attached to a solid support.
- 11. The method of any of claims 1 to 3, wherein the compound is attached to a solid support.
- 12. A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of disorders of the

peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising the steps of

- 5 i) contacting a test compound with a PDE4A polynucleotide,
- ii) detect binding of said test compound to said PDE4A polynucleotide.

10 13. The method of claim 12 wherein the nucleic acid molecule is RNA.

14. The method of claim 12 wherein the contacting step is in or at the surface of a cell.

15 15. The method of claim 12 wherein the contacting step is in a cell-free system.

16. The method of claim 12 wherein polynucleotide is coupled to a detectable label.

20 17. The method of claim 12 wherein the test compound is coupled to a detectable label.

25 18. A method of diagnosing a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising the steps of

- i) determining the amount of a PDE4A polynucleotide in a sample taken from said mammal,
- 30 ii) determining the amount of PDE4A polynucleotide in healthy and/or diseased mammals.

19. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising a therapeutic agent which binds to a PDE4A polypeptide.
20. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising a therapeutic agent which regulates the activity of a PDE4A polypeptide.
21. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising a therapeutic agent which regulates the activity of a PDE4A polypeptide, wherein said therapeutic agent is
- i) a small molecule,
 - ii) an RNA molecule,
 - iii) an antisense oligonucleotide,
 - iv) a polypeptide,
 - v) an antibody, or
 - vi) a ribozyme.
22. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation,

- 115 -

gastroenterological diseases and endocrinological disease in a mammal comprising a PDE4A polynucleotide.

23. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising a PDE4A polypeptide.
24. Use of regulators of a PDE4A for the preparation of a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal.
25. Method for the preparation of a pharmaceutical composition useful for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal comprising the steps of
- i) identifying a regulator of PDE4A,
 - ii) determining whether said regulator ameliorates the symptoms of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease in a mammal; and
 - iii) combining of said regulator with an acceptable pharmaceutical carrier.

- 116 -

26. Use of a regulator of PDE4A for the regulation of PDE4A activity in a mammal having a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, inflammation, gastroenterological diseases and endocrinological disease.
- 5

- 1/6 -

Fig. 1**SEQ ID NO: 1**

CAGCCTCCTCCTGGGACCCTTGCCCTGCCCCCTCCCATGGGCACGGA
CCCCCACCGCCTCCACCCACTGCCGCGGGGGGGCCCGTTGGGGCCCA
GGGCTGGCGGGCCATGTAACCAGGGCTGCTGCTGGGAGCGCGGAGGGG
AAGGGAGCCCCCAGCCCTGCTGGGCGGGCCAGGCCCTCCGCGGCTC
CCCCTTCCACTACCCACCTGCCCGGCACCCCTCCCCAGTG GTTGTTA
ACCCCGGGACTCCCCAAGCCCAGCCTCTGTGTGCAGCAGCCCCAGGCG
GGCTAAGTCTCCAAGATGCCCTTGGTGGATTTCTTCTGCGAGACCTGC
TCTAAGCCTTGGCTGGTGGGCTGGTGGGACCAGTTCAAAAGGATGTTG
AACCGTGAGCTCACACACCTGTCAGAAATGAGCAGGTCCGGAAACCAG
GTCTCAGAGTACATTTCCACAACATTCCTGGACAAACAGAATGAAGTG
GAGATCCCATCACCCACGATGAAGGAACGAGAAAAACAGCAAGCGCCG
CGACCAAGACCCTCCCAGCCGCCCCCGCCCCCTGTACCACACTTACAG
CCCATGTCCCAAATCACAGGGTTGAAAAAGTTGATGCATAGTAACAGC
CTGAACAACCTCTAACATTCCCCGATTTGGGGTGAAGACCGATCAAGAA
GAGCTCCTGGCCCAAGAACTGGAGAACCTGAACAAGTGGGGCCTGAAC
ATCTTTTTCGTGTCGGATTACGCTGGAGGCCGCTCACTCACCTGCATC
ATGTACATGATATTCCAGGAGCGGGACCTGCTGAAGAAATTCCGCATC
CCTGTGGACACGATGGTGACATACATGCTGACGCTGGAGGATCACTAC
CACGCTGACGTGGCCTACCATAACAGCCTGCACGCAGCTGACGTGCTG
CAGTCCACCCACGTACTGCTGGCCACGCCTGCACTAGATGCAGTGTTT
ACGGACCTGGAGATTCTCGCCGCCCTCTTCGCGGCTGCCATCCACGAT
GTGGATCACCTGGGGTCTCCAACCAGTTCCTCATCAACACCAATTTCG
GAGCTGGCGCTCATGTACAACGATGAGTCGGTGCTCGAGAATCACCA
CTGGCCGTGGGCTTCAAGCTGCTGCAGGAGGACAACTGCGACATCTTC
CAGAACCTCAGCAAGCGCCAGCGGCAGAGCCTACGCAAGATGGTCATC
GACATGGTGCTGGCCACGGACATGTCCAAGCACATGACCCTCCTGGCT
GACCTGAAGACCATGGTGGAGACCAAGAAAGTGACCAGCTCAGGGGTC

- 2/6 -

Fig. 1 (continued)

CTCCTGCTAGATAACTACTCCGACCGCATCCAGGTCCTCCGGAACATG
GTGCACTGTGCCGACCTCAGCAACCCACCAAGCCGCTGGAGCTGTAC
CGCCAGTGGACAGACCGCATCATGGCCGAGTTCTTCCAGCAGGGTGAC
CGAGAGCGCGAGCGTGGCATGGAAATCAGCCCCATGTGTGACAAGCAC
ACTGCCTCCGTGGAGAAGTCTCAGGTGGGTTTTATTGACTACATTGTG
CACCCATTGTGGGAGACCTGGGCGGACCTTGTCCACCCAGATGCCCAG
GAGATCTTGGACACTTTGGAGGACAACCGGGACTGGTACTACAGCGCC
ATCCGGCAGAGCCCATCTCCGCCACCCGAGGAGGAGTCAAGGGGGCCA
GGCCACCCACCCCTGCCTGACAAGTTCCAGTTTGAGCTGACGCTGGAG
GAGGAAGAGGAGGAAGAAATATCAATGGCCCAGATAACCGTGACAGCC
CAAGAGGCATTGACTGCGCAGGGATTGTGAGGAGTCGAGGAAGCTCTG
GATGCAACCATAGCCTGGGAGGCATCCCCGGCCCAGGAGTCGTTGGAA
GTTATGGCACAGGAAGCATCCCTGGAGGCCGAGCTGGAGGCAGTGTAT
TTGACACAGCAGGCACAGTCCACAGGCAGTGACCTGTGGCTCCGGAT
GAGTTCTCGTCCCGGGAGGAATTCGTGGTTGCTGTAAGCCACAGCAGC
CCCTCTGCCCTGGCTCTTCAAAGCCCCCTTCTCCCTGCTTGGAGGACC
CTGTCTGTTTCAGAGCATGCCCCGGGCCTCCCGGGCCTCCCCTCCACG
GCGGCCGAGGTGGAGGCCCAACGAGAGCACCAAGGCTGCCAAGAGGGCT
TGCAGTGCCTGCGCAGGGACATTTGGGGAGGACACATCCGCACTCCCA
GCTCCTGGTGGCGGGGGGTGAGGTGGAGACCCTACCTGATCCCCAGAC
CTCTGTCCCTGTTCCCCCTCCACTCCTCCCCCTCACTCCCCTGCTCCCCC
GACCACCTCCTCCTCTGCCTCAAAGACTCTTGTCCTCTTGTCCCTCCT
GAGAAAAAAGAAAACGAAAAGTGGGGTTTTTTTTCTGTTTTCTTTTTTT
CCCCTTTCCCCCTGCCCCACCCACGGGGCCTTTTTTTGGAGGTGGGG
GCTGGGGAATGAGGGGCTGAGGTCCCGGAAGGATTTTATTTTTTTGAA
TTTTAATTGTAACATTTTTAGAAAAAGAACAAAAAAGAAAAA
GAAAGAAACACAGCAACTGTAGATGCTCCTGTTCCCTGGTTCCCGCTTT

- 3/6 -

Fig. 1 (continued)

CCACTTCCAAATCCCTCCCCTCACCTTCCCCCACTGCCCCCAAGTTC
CAGGCTCAGTCTTCCAGCCGCCTGGGGAGTCTCTACCTGGGCCCCAAGC
AGGTGTGGGGCCTCCTTCTGGGCTTTTCTTCTGAATTTAGAGGATTTTC
TAGAACGTGGTCAGGAATAGCCATTCTAGGCGGGGCTGGGGCCAGGGT
GGGGGGCAGTCACTGTGGGAGGTCCCAGCTCCAGCCCCCCTCTGGTTT
GCTGCCTCCTCTCCCCTCTAAAAAAGTCTTCCGCTTGATTTTGCACAA
TCCCGGCGATACTCCTGGCGATACTGACTAGAAAGTCAGGGAGCTGGG
GGAGCTGTTCACTTTAGGATACGGGGGTGGTATGGAAGGGAGCGTTCA
CACCGCCAGCCTCGGGCCTGGGATTTGAGGAGGGCCCTAGACCTCCTC
CACTCTCCATCCCCTTTCCCTTCCACTTTGGGTTCACCTTTGAATTTTC
TCCGTTTTTTTGGGGCAGTGGCTCTGATCCACTCACCCCCCGCCCCC
GCCCCACTTCTAGCTGCTTCTCCTCTTGTTTCTGCCTTAATAATTCCC
ACGGCCACAGGCAAGGGGGTTCAGTGGCCGCCTGCACCTTGGATGAG
GCAGGGCCAGGCGCCCAGAACCCCCATCCTGGCCGCACCCCCCTTTCC
AGGGTCCTCCGGACCCACCTTCCACACTCTGATCACAGCCCCCTAC
CTTTTGCCCTAGGAGGAAGCAATAATGGTGTATACCCTCATTCTCATT
CCTGGGCAGCCCTTCCCTTCCACCCTGGCACCAAATAATTTCTCCTCC
ATCCGTACCTTGCCTAGCCTCTCCCTCTCCCCCAGCTAGTCCCCTGAG
CAATACGGCAGACAGATGCAAGACCATTTTTCTCCAAGCCATGGGGGA
CTGTTTGGAAGGAAAGCCCCCTCTCTCCCTCCTCCCCTCGCCCTCGGC
CTGGTTCTGCAGCTGGACCGACCTCATTCATCGCCTGCCCCCTACCCA
ATTCTGAGCACACGGTACTGTAGCCCCCAGTTCCTCCCTAGCCTTCCA
TCCCTCTGTCCACCCAGGGGGAGGTAACCCCGCACTCACACTCCCTT
GATGCTGTCTGTACAGGGTTCATATTTTGTAGCGAAAGTCGTTTTTGT
CCCAGCCGGCGATCGGAGTGGGCCTTTTCTTTCTTTTGTTCATTCTT
TACCTTTTTTTCTTTTCTTTCTTTCTTTTTGTACATACTGTAAGGTT
GGTTTGTAATTATTCTACGGAGGCAAAAAGGGAAAATAAAAACCTGC

- 4/6 -

Fig. 1 (continued)

CCTTCCCTGGCTGACCCAGTCGGGAAGGTAGGGAAGGAGGTCTCCCGT
TGGGAGAGTCTCTGTTCCCTGCTGTATTATACAACGTACCATAGTCCT
GGGAAAAGGGTGGACTCACCGCTGTTGTTTTATGGGAAGTCGTGTCAT
CCTAGGGGTTGGGGCTGGGCAGAGCCTGTCCCCTCCCCCTTCTCCAG
GAGCCAGGGGGTGACTGGAGAGACAGACCCACCCCAAGCAGGGCTCC
TCTCCCCAGGGTGAGCACAGGACCTCTGTAAGCTGCTTGTGTATTGTC
CACTTTGACGATCAGTCATTCCGTCCGTTGATCAATAATCCTTCGATC
TTGTCTCCAATTAAACCGAGGCTTTCACCGA

- 5/6 -

Fig. 2**SEQ ID NO: 2**

MEPPTVPSERSLSLSLPGPREGQATLKPPPQHLWRQPRTPIRIQQRGY
SDSAERAERERQPHRPIERADAMDTSDRPGLRTTRMSWPSSFHGTGTG
SGGAGGGSSRRFEAENGPTPSPGRSPLDSQASPLVLHAGAATSQRRE
SFLYRSDSDYDMSPKTMSRNSSVTSEAHAE DLIVTPFAQVLASLRSVR
SNFSLLTNPVPVPSNKRSP LGGPTPVCKATLSEETCQQLARETLEELD W
CLEQLETMQTYRSVSEMASHKFKRMLNRELTHLSEMSRSGNQVSEYIS
TTF LDKQNEVEI PSPTMKEREKQQAPRPRPSQPPPPVPVPHLQPM SQIT
GLKKLMHSNSLNN SNIPRFGVKTDQEELLAQELENLNKWGLNIFCVSD
YAGGRSLTCIMYMI FQERDLLKKFRIPVD TMVTYMLTLEDHYHADVAY
HNSLHAADV LQSTHVLLATPALDAVFTDLEILAALFAAAIHDVDHPGV
SNQFLINTNSELALMYNDESVLENHHLAVGFKLLQEDNCDIFQNLSKR
QRQSLRKMVIDMVLATDMSKHMTLLADLKT MVETKKVTSSGVLLLDNY
SDRIQVLRNMVHCADLSNPTKPLELYRQWTD RIMAEFFQOGDRERERG
MEISPMCDKHTASVEKSQVGFIDYIVHPLWETWADLVHPDAQEILD TL
EDNRDWYYSAIRQSPSPPEEEESRGPGHPPLPDKFQFELTLEEEEEEE
ISMAQIPCTAQEALTEQGLSGVEEALDATIAWEASPAQESLEVMAQEA
SLEAELEAVYLTQQAQSTGSAPVAPDEFSSREEFVVAVSHSSPSALAL
QSPLLPAWRTL SVSEHAPGLPGLPSTAAEVEAQREHQAAKRACSACAG
TFGEDTSALPAPGGGGSGGDPT

Fig. 3**SEQ ID NO: 3**

5' - CAGGAGATCTTGGACACTTTGGA - 3'

- 6/6 -

Fig. 4

SEQ ID NO: 4

5' -GCATCCAGAGCTTCCTCGAC-3'

Fig. 5

SEQ ID NO: 5

5' -AGCCCAAGAGGCATTGACTGCGC-3'

- 1 -

SEQUENCE LISTING

<110> Bayer AG, BHC

<120> Diagnostics and Therapeutics for Diseases Associated with Human Phosphodiesterase 4A (PDE4A)

<130> Le A 36 427

<160> 5

<170> PatentIn version 3.1

<210> 1

<211> 4255

<212> DNA

<213> Homo sapiens

<400> 1

```
cagcctctctc ctgggaccct tgccctgccc ccctcccatg ggcacggacc cccaccgcc      60
tccaccact gccgcggggg ggcccgttgg ggcccagggc tggcgggcca tgtaaccagg      120
gctgctgctg ggagcgcgga ggggaaggga gccccagcc ctgctgggcc ggcccaggcc      180
cctccgcggc tcccccttcc actaccacc tgcccggcac cccctcccca gtggttggtta      240
accccgggac tcccaagcc cagcctctgt gtgcagcagc ccagggcggg ctaagtctcc      300
aagatgccct tgggtggattt cttctgcgag acctgctcta agccttggct ggtgggctgg      360
tgggaccagt tcaaaaggat gttgaaccgt gagctcacac acctgtcaga aatgagcagg      420
tccggaaacc aggtctcaga gtacatttcc acaacattcc tggacaaaca gaatgaagtg      480
gagatcccat ccccacgat gaaggaacga gaaaaacagc aagcgccgcg accaagaccc      540
```

- 2 -

tcccagccgc ccccgccccc tgtaccacac ttacagccca tgtcccaaat cacaggggtg 600

aaaaagttga tgcataagtaa cagcctgaac aactctaaca ttccccgatt tgggggtgaag 660

accgatcaag aagagctcct ggcccaagaa ctggagaacc tgaacaagtg gggcctgaac 720

atcttttgcg tgtcggatta cgctggaggc cgctcactca cctgcatcat gtacatgata 780

ttccaggagc gggacctgct gaagaaattc cgcacccctg tggacacgat ggtgacatac 840

atgctgacgc tggaggatca ctaccacgct gacgtggcct accataacag cctgcacgca 900

gctgacgtgc tgcagtccac ccacgtactg ctggccacgc ctgcactaga tgcagtgttc 960

acggacctgg agattctcgc cgccctcttc ggggtgcca tccacgatgt ggatcacccct 1020

ggggtctcca accagttcct catcaacacc aattcggagc tggcgctcat gtacaacgat 1080

gagtcggtgc tcgagaatca ccacctggcc gtgggcttca agctgctgca ggaggacaac 1140

tgcgacatct tccagaacct cagcaagcgc cagcggcaga gcctacgcaa gatgggtcatc 1200

gacatgggtgc tggccacgga catgtccaag cacatgaccc tcctggctga cctgaagacc 1260

atggtggaga ccaagaaagt gaccagctca ggggtcctcc tgctagataa ctactccgac 1320

cgcacccagg tcctccggaa catggtgcac tgtgccgacc tcagcaaccc caccaagccg 1380

ctggagctgt accgccagtg gacagaccgc atcatggccg agttcttcca gcagggtgac 1440

cgagagcgcg agcgtggcat ggaaatcagc cccatgtgtg acaagcacac tgcctccgtg 1500

gagaagtctc aggtgggttt tattgactac attgtgcacc cattgtggga gacctgggcg 1560

gacctgtgcc acccagatgc ccaggagatc ttggacactt tggaggacaa cggggactgg 1620

tactacagcg ccatccggca gagcccatct ccgccacccg aggaggagtc aagggggcca 1680

ggccaccac ccctgcctga caagttccag tttgagctga cgctggagga ggaagaggag 1740

gaagaaatat caatggccca gataccgtgc acagcccaag aggcattgac tgcgcaggga 1800

ttgtcaggag tcgaggaagc tctggatgca accatagcct gggaggcatc cccggcccag 1860

gagtcgttgg aagttatggc acaggaagca tccttgaggc ccgagctgga ggcagtgtat 1920

- 3 -

ttgacacagc aggcacagtc cacaggcagt gcacctgtgg ctccggatga gttctcgtcc 1980
cgggaggaat tcgtggttgc tgtaagccac agcagcccct ctgccctggc tcttcaaagc 2040
ccccctctcc ctgcttggag gaccctgtct gtttcagagc atgccccggg cctccccggc 2100
ctcccccca cggcgccga ggtggaggcc caacgagagc accaggctgc caagagggt 2160
tgcagtgcct gcgcaggac atttggggag gacacatccg cactcccagc tcctgggtggc 2220
ggggggtcag gtggagacc tacctgatcc ccagacctct gtccctgttc cctccactc 2280
ctccccctac tcccctgtc ccccgaccac ctctctctct gcctcaaaga ctcttgcct 2340
cttgccctc ctgagaaaa agaaaacgaa aagtgggggt tttttctgtt ttctttttt 2400
ccccctccc cctgccccca ccacggggc ctttttttgg aggtgggggc tggggaatga 2460
ggggtgagg tcccggaagg attttatttt tttgaatttt aattgtaaca ttttagaaa 2520
aagaacaaaa aaagaaaaa aaaagaaaga aacacagcaa ctgtagatgc tcctgttcct 2580
ggttcccgt ttcacttcc aaatccctcc cctcaccttc cccactgcc cccaagtcc 2640
caggctcagt cttccagccg cctggggagt ctctacctgg gcccaagcag gtgtggggcc 2700
tccttctggg cttttctct gaatttagag gatttctaga acgtggtcag gaatagccat 2760
tctaggcggg gctggggcca ggtgggggg cagtcactgt gggaggtccc agctccagcc 2820
cccctctggt ttgctgcctc ctctcccctc taaaaagtc ttccgcttga ttttgacaa 2880
tcccgcgat actcctggcg atactgacta gaaagtcagg gagctggggg agctgttcac 2940
tttaggatac gggggtggtg tggaaggag cgttcacacc gccagcctcg ggctgggat 3000
ttgaggagg ccctagacct cctccactct ccacccctt tccctccac tttgggttca 3060
ctttgaattt tctccgtttt ttggggcagt ggctctgac cactacccc cccgcccccc 3120
gccccacttc tagctgcttc tcctcttgtt tctgccttaa taattccac ggccacaggc 3180
aagggggttg cagtggccgc ctgcaccttg gatgaggcag ggccaggcgc ccagaacccc 3240
catcctggcc gcacccccct ttccagggtc ctccggacc cacttccac actctgatca 3300

- 4 -

cagccccctt accttttgcc ctaggaggaa gcaataatgg tgtataccct cattctcatt 3360
cctgggcagc ccttccttcc accctggcac caaaataatt tctcctccat ccgtaccttg 3420
cctagcctct ccctctcccc cagctagtcc cctgagcaat acggcagaca gatgcaagac 3480
cattttttct caagccatgg gggactgttt ggaaggaaag cccctctctt ccctcctccc 3540
ctcgccctcg gcctggttct gcagctggac cgacctcatt catcgctgc cccctacca 3600
attctgagca cacggtactg tagccccag ttctcccta gccttccat cctctgtcca 3660
ccccaggggg aggttaaccc gactcacac tcccttgatg ctgtctgtac agggttcata 3720
ttttgtagcg aaagtcgttt ttgtcccagc cggcgatcgg agtgggcctt ttctttcttt 3780
ttgttcattc ttacctttt ttcttttctt ttctttcttt ttgtacata ctgtaagggt 3840
ggtttgtaaa ttattctacg gaggcaaaaa gggaaaataa aaacttgccc ttccctggct 3900
gaaccagtcg ggaaggtagg gaaggaggtc tcccgttggg agagtctctg ttctgtctgt 3960
attatacaac tgtaccatag tcctgggaaa aggggtggact caccgctgtt gttttatggg 4020
aagtcgtgtc atcctagggg ttggggctgg gcagagcctg tccctcccc ccttctccag 4080
gagccagggg gtgactggag agacagacc accccaagc agggctcctc tcccagggg 4140
gagcacagga cctctgtaag ctgcttgtgt attgtccact ttgacgatca gtcattcggg 4200
ccgttgatca ataaccttc gatcttgtct ccaattaaac cgaggcttcc accga 4255

<210> 2

<211> 886

<212> PRT

<213> Homo sapiens

Ser Pro Leu Gly Gly Pro Thr Pro Val Cys Lys Ala Thr Leu Ser Glu
210 215 220

Glu Thr Cys Gln Gln Leu Ala Arg Glu Thr Leu Glu Glu Leu Asp Trp			
225	230	235	240
Cys Leu Glu Gln Leu Glu Thr Met Gln Thr Tyr Arg Ser Val Ser Glu			
245	250		255
Met Ala Ser His Lys Phe Lys Arg Met Leu Asn Arg Glu Leu Thr His			
260	265		270
Leu Ser Glu Met Ser Arg Ser Gly Asn Gln Val Ser Glu Tyr Ile Ser			
275	280		285
Thr Thr Phe Leu Asp Lys Gln Asn Glu Val Glu Ile Pro Ser Pro Thr			
290	295		300
Met Lys Glu Arg Glu Lys Gln Gln Ala Pro Arg Pro Arg Pro Ser Gln			
305	310	315	320
Pro Pro Pro Pro Pro Val Pro His Leu Gln Pro Met Ser Gln Ile Thr			
325	330		335
Gly Leu Lys Lys Leu Met His Ser Asn Ser Leu Asn Asn Ser Asn Ile			
340	345		350
Pro Arg Phe Gly Val Lys Thr Asp Gln Glu Glu Leu Leu Ala Gln Glu			
355	360		365
Leu Glu Asn Leu Asn Lys Trp Gly Leu Asn Ile Phe Cys Val Ser Asp			
370	375		380
Tyr Ala Gly Gly Arg Ser Leu Thr Cys Ile Met Tyr Met Ile Phe Gln			
385	390	395	400
Glu Arg Asp Leu Leu Lys Lys Phe Arg Ile Pro Val Asp Thr Met Val			
405	410		415
Thr Tyr Met Leu Thr Leu Glu Asp His Tyr His Ala Asp Val Ala Tyr			
420	425		430
His Asn Ser Leu His Ala Ala Asp Val Leu Gln Ser Thr His Val Leu			
435	440		445
Leu Ala Thr Pro Ala Leu Asp Ala Val Phe Thr Asp Leu Glu Ile Leu			
450	455		460

- 7 -

Ala Ala Leu Phe Ala Ala Ala Ile His Asp Val Asp His Pro Gly Val
 465 470 475 480

Ser Asn Gln Phe Leu Ile Asn Thr Asn Ser Glu Leu Ala Leu Met Tyr
 485 490 495

Asn Asp Glu Ser Val Leu Glu Asn His His Leu Ala Val Gly Phe Lys
 500 505 510

Leu Leu Gln Glu Asp Asn Cys Asp Ile Phe Gln Asn Leu Ser Lys Arg
 515 520 525

Gln Arg Gln Ser Leu Arg Lys Met Val Ile Asp Met Val Leu Ala Thr
 530 535 540

Asp Met Ser Lys His Met Thr Leu Leu Ala Asp Leu Lys Thr Met Val
 545 550 555 560

Glu Thr Lys Lys Val Thr Ser Ser Gly Val Leu Leu Leu Asp Asn Tyr
 565 570 575

Ser Asp Arg Ile Gln Val Leu Arg Asn Met Val His Cys Ala Asp Leu
 580 585 590

Ser Asn Pro Thr Lys Pro Leu Glu Leu Tyr Arg Gln Trp Thr Asp Arg
 595 600 605

Ile Met Ala Glu Phe Phe Gln Gln Gly Asp Arg Glu Arg Glu Arg Gly
 610 615 620

Met Glu Ile Ser Pro Met Cys Asp Lys His Thr Ala Ser Val Glu Lys
 625 630 635 640

Ser Gln Val Gly Phe Ile Asp Tyr Ile Val His Pro Leu Trp Glu Thr
 645 650 655

Trp Ala Asp Leu Val His Pro Asp Ala Gln Glu Ile Leu Asp Thr Leu
 660 665 670

Glu Asp Asn Arg Asp Trp Tyr Tyr Ser Ala Ile Arg Gln Ser Pro Ser
 675 680 685

Pro Pro Pro Glu Glu Glu Ser Arg Gly Pro Gly His Pro Pro Leu Pro
 690 695 700

- 8 -

Asp Lys Phe Gln Phe Glu Leu Thr Leu Glu Glu Glu Glu Glu Glu
 705 710 715 720

Ile Ser Met Ala Gln Ile Pro Cys Thr Ala Gln Glu Ala Leu Thr Glu
 725 730 735

Gln Gly Leu Ser Gly Val Glu Glu Ala Leu Asp Ala Thr Ile Ala Trp
 740 745 750

Glu Ala Ser Pro Ala Gln Glu Ser Leu Glu Val Met Ala Gln Glu Ala
 755 760 765

Ser Leu Glu Ala Glu Leu Glu Ala Val Tyr Leu Thr Gln Gln Ala Gln
 770 775 780

Ser Thr Gly Ser Ala Pro Val Ala Pro Asp Glu Phe Ser Ser Arg Glu
 785 790 795 800

Glu Phe Val Val Ala Val Ser His Ser Ser Pro Ser Ala Leu Ala Leu
 805 810 815

Gln Ser Pro Leu Leu Pro Ala Trp Arg Thr Leu Ser Val Ser Glu His
 820 825 830

Ala Pro Gly Leu Pro Gly Leu Pro Ser Thr Ala Ala Glu Val Glu Ala
 835 840 845

Gln Arg Glu His Gln Ala Ala Lys Arg Ala Cys Ser Ala Cys Ala Gly
 850 855 860

Thr Phe Gly Glu Asp Thr Ser Ala Leu Pro Ala Pro Gly Gly Gly Gly
 865 870 875 880

Ser Gly Gly Asp Pro Thr
 885

<210> 3

<211> 23

<212> DNA

<213> Homo sapiens

- 9 -

<400> 3
caggagatct tggacacttt gga 23

<210> 4

<211> 20

<212> DNA

<213> Homo sapiens

<400> 4
gcatccagag cttcctcgac 20

<210> 5

<211> 23

<212> DNA

<213> Homo sapiens

<400> 5
agcccaagag gcattgactg cgc 23