



(86) Date de dépôt PCT/PCT Filing Date: 2016/03/22
 (87) Date publication PCT/PCT Publication Date: 2016/09/29
 (85) Entrée phase nationale/National Entry: 2017/09/27
 (86) N° demande PCT/PCT Application No.: US 2016/023582
 (87) N° publication PCT/PCT Publication No.: 2016/154203
 (30) Priorité/Priority: 2015/03/23 (US62/136,813)

(51) Cl.Int./Int.Cl. *A61K 38/17* (2006.01),
C07J 3/00 (2006.01), *C07J 41/00* (2006.01),
C07J 51/00 (2006.01), *G01N 33/574* (2006.01)
 (71) Demandeur/Applicant:
 EVESTRA, INC., US
 (72) Inventeurs/Inventors:
 NAIR, HAREESH, US;
 SANTHAMMA, BINDU, US;
 NICKISCH, KLAUS, DE
 (74) Agent: GOUDREAU GAGE DUBUC

(54) Titre : NOUVEAUX AGENTS CYTOTOXIQUES QUI CIBLENT PREFERENTIELLEMENT LE FACTEUR D'INHIBITION DE LA LEUCEMIE (LIF) POUR LE TRAITEMENT DE TUMEURS MALIGNES ET COMME NOUVEAUX AGENTS CONTRACEPTIFS
 (54) Title: NOVEL CYTOTOXIC AGENTS THAT PREFERENTIALLY TARGET LEUKEMIA INHIBITORY FACTOR (LIF) FOR THE TREATMENT OF MALIGNANCIES AND AS NEW CONTRACEPTIVE AGENTS




FIG. 4A

(57) **Abrégé/Abstract:**

Described herein are new anti-cancer compounds and methods of using such compounds, acting through a new mechanism of action by simultaneous inhibition of leukemia inhibitory factor (LIF) and MDM2. In one embodiment, a cytotoxic compound has the structure (I) or (II). In an embodiment, a method of treating cancer in a subject comprising administering to a subject a medicament comprising an effective amount of a cytotoxic small molecule compound that inhibits leukemia inhibitory factor or leukemia inhibitory factor receptor.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(43) International Publication Date
29 September 2016 (29.09.2016)

WIPO | PCT

(10) International Publication Number
WO 2016/154203 A1

(51) International Patent Classification:

A61K 38/17 (2006.01) C07J 51/00 (2006.01)
C07J 3/00 (2006.01) G01N 33/574 (2006.01)
C07J 41/00 (2006.01)

(21) International Application Number:

PCT/US2016/023582

(22) International Filing Date:

22 March 2016 (22.03.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/136,813 23 March 2015 (23.03.2015) US

(71) Applicant: **EVESTRA, INC.** [US/US]; 14805 Omicron Drive, San Antonio, Texas 78245 (US).(72) Inventors: **NAIR, Hareesh**; 14805 Omicron Drive, San Antonio, Texas 78245 (US). **SANTHAMMA, Bindu**; 619 Colosseo Way, San Antonio, Texas 78253 (US). **NICK-
ISCH, Klaus**; Zeseher Strasse 14, D-12307 Berlin (DE).(74) Agent: **DELUCA, Mark R.**; Meyertons, Hood, Kivlin, Kowert & Goetzel, P.C., P.O. Box 398, Austin, Texas 78767-0398 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: NOVEL CYTOTOXIC AGENTS THAT PREFERENTIALLY TARGET LEUKEMIA INHIBITORY FACTOR (LIF) FOR THE TREATMENT OF MALIGNANCIES AND AS NEW CONTRACEPTIVE AGENTS




FIG. 4A

(57) Abstract: Described herein are new anti-cancer compounds and methods of using such compounds, acting through a new mechanism of action by simultaneous inhibition of leukemia inhibitory factor (LIF) and MDM2. In one embodiment, a cytotoxic compound has the structure (I) or (II). In an embodiment, a method of treating cancer in a subject comprising administering to a subject a medicament comprising an effective amount of a cytotoxic small molecule compound that inhibits leukemia inhibitory factor or leukemia inhibitory factor receptor.

WO 2016/154203 A1

**TITLE: NOVEL CYTOTOXIC AGENTS THAT PREFERENTIALLY TARGET
LEUKEMIA INHIBITORY FACTOR (LIF) FOR THE TREATMENT OF
MALIGNANCIES AND AS NEW CONTRACEPTIVE AGENTS**

5

BACKGROUND OF THE INVENTION

1. Field of the Invention

[0001] The invention generally relates to new anti-cancer compounds, acting through a new
10 mechanism of action by simultaneous inhibition of leukemia inhibitory factor (LIF) and
MDM2.

2. Description of the Relevant Art

[0002] Cancer is a group of diseases characterized by the uncontrolled growth and spread of
abnormal cells. If the spread is not controlled, it can result in death. Cancer may affect
15 people at all ages, but risk for the more common varieties tends to increase with age. Cancer
is caused by external factors, such as tobacco, infectious organisms, and an unhealthy diet,
and internal factors, such as inherited genetic mutations, hormones, and immune conditions.
These factors may act together or in sequence to cause cancer. Treatments include surgery,
radiation, chemotherapy, hormone therapy, immune therapy, and targeted therapy (drugs that
20 specifically interfere with cancer cell growth). According to American Cancer Society, about
1,685,210 new cancer cases are expected to be diagnosed in 2016 and about 595,690
Americans are expected to die of cancer in 2016, which translates to about 1,630 people per
day (Cancer Facts and Figures, 2016).

[0003] Conventional cancer diagnosis and therapies to date have attempted to selectively
25 detect and eliminate cancer cells that are largely fast-growing. Standard oncology regimens
have often been largely designed to administer the highest dose of irradiation or a
chemotherapeutic agent without undue toxicity, i.e., often referred to as the “maximum
tolerated dose” (MTD) or “no observed adverse effect level” (NOAEL). Chemotherapeutic
strategies often involve administration of a combination of chemotherapeutic agents in order
30 to increase the efficacy of treatment. Despite the availability of a large variety of
chemotherapeutic agents, these therapies have many drawbacks includes bone marrow
depression, immunosuppression, gastrointestinal distress, etc.

[0004] Other novel therapeutic approaches seek to utilize targeted therapies with increased
selectivity and efficacy in preselected patient populations. A recent molecularly targeted

therapy is established by inhibiting the enzyme poly (ADP-ribose) polymerase (PARP) by small molecule inhibitors such as Olaparib on tumors that have a defect in the homologous DNA recombination due to BRCA mutations.

5 [0005] Cancer stem cells have been identified in a large variety of cancer types. Many different cancers including breast, prostate, lung, pancreas etc. have showed the presence of stem cell populations that are resistant to conventional chemotherapies. Therapies that could target cancer stem cells could be of great therapeutic potential in hormone/chemotherapy refractory cancers.

10 [0006] Leukemia inhibitory factor, or LIF, is an interleukin 6 class cytokine that affects cell growth by inhibiting cell differentiation. LIF binds to the specific LIF receptor (LIFR- α) which forms a heterodimer with a specific subunit common to all members of that family of receptors, the GP130 signal transducing subunit. This leads to activation of the JAK/STAT (Janus kinase/signal transducer and activator of transcription) and MAPK (mitogen activated protein kinase) cascades. LIF promotes STAT3 phosphorylation.

15 [0007] LIF promotes tumorigenesis in many solid tumors and mediates pro-invasive activation of stromal fibroblasts in cancer. LIF mediates TGF beta dependent actinomycin contractility, extracellular matrix remodeling leading to cancer cell invasion in fibroblasts. It is established that paracrine molecules such as TGF-beta, growth factors, and proinflammatory molecules (such as the IL-6 family of cytokines that includes LIF) are
20 secreted by cancer cells and promote tumorigenesis. TGF-beta-mediated phosphorylation of Smad3 potentiates transcriptional regulation of many genes that assist in the proliferation of cancer cells. The role of TGF-beta/SMAD and JAK/STAT3 in signaling in tumor cell dependent proinvasive fibroblast activation and expression of alpha-smooth muscle actin (α -SMA) producing carcinoma associated fibroblast (CAF) hallmark is well known.

25 [0008] Leukemia Inhibitory Factor (LIF) is, thus important in sustaining pluripotency and stemcellness and embryogenesis. A critical point during mammalian pregnancy is the implantation of the blastocyst when the embryo attaches to the wall of the uterus. Females lacking a functional LIF gene are fertile, but their blastocysts fail to implant and do not develop. LIF may also be critical to endometrial receptivity in humans, as well as a wide
30 range of other mammals, with reduced LIF expression being linked to several cases of female infertility.

[0009] LIF induces many genes that over express in cancer. One gene LIF induces over expression for is breast cancer antiestrogen resistance protein (p13Cas/BCAR1).

Overexpression of p130Cas/BCAR1 has been detected in human breast cancer, prostate cancer, ovarian cancer, lung cancer, colorectal cancer, hepatocellular carcinoma, glioma, melanoma, anaplastic large cell lymphoma and chronic myelogenous leukemia. The presence of aberrant levels of hyperphosphorylated p130Cas/BCAR1 strongly promotes cell proliferation, migration, invasion, survival, angiogenesis and drug resistance.

[0010] Carcinoma-associated fibroblasts (CAF) are the most abundant population of non-cancer cells found in tumors, and their presence is often associated with poor clinical prognosis. LIF drives cancer cell-dependent pro-invasive extracellular matrix remodeling in carcinoma associated fibroblasts. It has been established that under the influence of bioactive molecules, such as LIF, within the tumor stroma, resident fibroblast are activated and promote tumorigenesis.

[0011] LIF is an important negative regulator of tumor suppressor gene p53. Down regulation of p53 by LIF is mediated by the activation of STAT3, which transcriptionally induces inhibitor of DNA binding 1 (ID1). ID1 upregulates MDM2, a natural negative regulator of p53 and promotes p53 degradation. EC330 was found to indirectly diminish the phosphorylation of SMAD thorough blocking TGF-beta. Overexpression of LIF is associated with poor prognosis and increase incidence of chemoresistance. Targeting LIF and MDM2 to reactivate p53 is a potential therapeutic strategy for chemotherapy as well as in combination with other agents to alleviate chemoresistance.

[0012] There have been few discoveries made in the field of LIF regarding the inhibition of LIF in medicine. Monoclonal antibodies against LIF have been described. For example, U.S. Patent No. 6,156,729 claimed the use of leukemia inhibitory factor (LIF) antagonists to prevent or lessen hypertrophy.

[0013] EP Patent Application No. EP2371860 A1 claimed that LIF specific monoclonal antibody could be useful for the treatment for proliferative diseases such as cancer.

[0014] U.S. Patent No. 9194872 B2 and U.S. Published Patent Application No. 2015/0133376 A1 taught the use of a leukemia inhibitory factor receptor inhibitor for the potentiation of cancer radiotherapy. U.S. Patent No. 9,194,872 also claimed rapamycin and substituted quinoline as cancer therapy sensitizers that modulate LIF.

WO 2016/154203

PCT/US2016/023582

[0015] A receptor protein (DNA encoding fusion receptor) comprising a gp130 polypeptide linked to a single-chain leukemia inhibitory factor receptor (LIF-R) polypeptide is capable of binding both oncostatin M and leukemia inhibitory factor (LIF) has reported in U.S. Patent No. 5,426,048.

5 [0016] A method for treating a mammal experiencing heart failure to prevent or lessen cardiac hypertrophy comprising administering therapeutically effective amount of LIF antagonist (antibody) and an endothelin antagonist to a mammal in need of such treatment was described in U.S. Patent No. 6,156,733, U.S. Patent No. 5,573,762 and U.S. Patent No. 5,837,241.


10 [0017] The use of recombinant LIF from mammalian species to enhance implantation and development of embryos was described in U.S. Patent No. 5,962,321.

[0018] In summary, all prior art among the area of LIF and LIFR targeting agents only included monoclonal antibodies (mAbs) and glycosylated or non-glycosylated antibody fragments. Some of these agents are in clinical trials and none has been approved to use in
15 patients yet. Generally mAbs are expensive to produce and recognize only specific epitope(s) on an antigen. This drawback could lead to miss some variants. Moreover limited clones are available. It is therefore desirable to develop compounds that are small molecule inhibitors of LIF/LIFR and have targeted therapeutic advantage in treating cancers.


20

SUMMARY OF THE INVENTION

[0019] In one embodiment, a cytotoxic compound has the structure (I) or (II):




(I)



(II)

5 where:



X is O, NH, or S;

10

$n = 1-18$; $m = 1-18$;

R² is H, F, Cl, $-\text{C(O)-R}^6$, or $-\text{CH}_2(\text{OH})$;

15

R³ is H, F, Cl, $-\text{C(O)-R}^6$, or $-\text{CH}_2(\text{OR}^6)$;

R⁴ is H, alkyl, $-\text{CH}_2\text{-OH}$, $-\text{CO}_2\text{R}^6$, $-\text{CON(R}^6)_2$;

R⁵ is alkyl, alkenyl, alkylacyl, cycloalkyl, heterocycle, $-\text{CN}$, alkoxy, $-\text{N(R}^6)_2$, $-\text{CON(R}^6)_2$,

20


$-\text{S(O)R}^6$, $-\text{SR}_6$, $-\text{SO}_2\text{R}^6$; or $-(\text{CH}_2)_p\text{-CH}_2\text{-Y}$;

Y is H, OR^6 , SCH_3 , CF_3 , $-\text{N(R}^6)_2$; $p = 1-18$; and

25

R⁶ is H, alkyl, or cycloalkyl.

[0020] In an embodiment, a cytotoxic compound has the structure (III):



5

where:

R^2 is H, F, Cl, -CO-, or -C(OH)-;

10

R^3 is H, F, Cl, -CO-, or -C(OH)-;

R^4 is H, alkyl, -CH₂-OH, -CO₂R⁶, -CON(R⁶)₂;

15


R^5 is alkyl, alkenyl, alkylacyl, cycloalkyl, heterocycle, -CN, alkoxy, -N(R⁶)₂, -CON(R⁶)₂, -S(O)R⁶, -SR₆, -SO₂R⁶; or -(CH₂)_p-CH₂-Y; and

Y is H, OR⁶, SCH₃, CF₃, -N(R⁶)₂; p = 1-18; and

20

R^6 is H, alkyl, or cycloalkyl.

[0021] In an embodiment, a cytotoxic compound has the structure (IV):



(IV)

5

where:


R⁵ is alkyl, alkenyl, alkylacyl, cycloalkyl, heterocycle, -CN, alkoxy, -N(R⁶)₂, -
CON(R⁶)₂,

10 -S(O)R⁶, -SR⁶, or -SO₂R⁶; and

R⁶ is H, alkyl, or cycloalkyl.

[0022] In an embodiment, a cytotoxic compound has the structure (IV):

15



(IV)

where:

20

R⁵ is alkyl, alkenyl, alkylacyl, cycloalkyl, 1,3-imidazolyl, alkoxy, -N(R⁶)₂, -SR⁶, or -
SO₂R⁶; and

R^6 is H, lower alkyl, or cycloalkyl.


[0023] In some embodiments, a cytotoxic compound has the structure (IV), where R^5 is
 5 alkyl, alkenyl, or cycloalkyl. In some embodiments, a cytotoxic compound has the
 structure (IV), where R^5 is 1,3-imidazolyl. In some embodiments, a cytotoxic compound
 has the structure (IV), where R^5 is alkoxy. In some embodiments, a cytotoxic compound
 has the structure (IV), where R^5 is $-N(R^6)_2$. In some embodiments, a cytotoxic compound
 has the structure (IV), where R^5 is $-SR_6$. In some embodiments, a cytotoxic compound has
 10 the structure (IV), where R^5 is $-SO_2R^6$.

[0024] In an embodiment, a method of treating cancer in a subject comprising
 administering to a subject a medicament comprising an effective amount of a cytotoxic
 small molecule compound that inhibits leukemia inhibitory factor or leukemia inhibitory
 15 factor receptor. The cancer may be a cancer that overexpresses leukemia inhibitory factor.
 The cancer may be a cancer that exhibits a desmoplastic stromal response. The cancer
 may be a cancer that exhibits cancer initiating stem cells (CISC) or cancer associated stem
 cells (CASC). The cytotoxic small molecule compound may have the structures as set forth
 above. .

20 [0025] In some embodiments, in addition to inhibiting leukemia inhibitory factor or
 leukemia inhibitory factor receptor, the cytotoxic small molecule compound inhibits
 MDM2 and/or inhibits carcinoma associated fibroblast and/or stabilizes P53 levels.

[0026] In an embodiment, a synthetic intermediate useful for the formation of cytotoxic
 small molecule compounds has the structure (V):

25



where:

R^5 is alkyl, alkenyl, alkylacyl, cycloalkyl, heterocycle, -CN, alkoxy, $-N(R^6)_2$, -
CON(R^6)₂,

5 -S(O) R^6 , -SR₆, -SO₂ R^6 ; or $-(CH_2)_p-CH_2-Y$; and

Y is H, OR⁶, SCH₃, CF₃, $-N(R^6)_2$; p = 1-18; and

R^6 is H, alkyl, or cycloalkyl.

10

[0027] In an embodiment, the synthetic intermediate has the structure (V) where:

R^5 is alkyl, alkenyl, alkylacyl, cycloalkyl, 1,3-imidazolyl, alkoxy, $-N(R^6)_2$, -SR₆, or -
SO₂ R^6 ; and

15 R^6 is H, lower alkyl, or cycloalkyl.

[0028] In an embodiment, a method of treating hypertrophic fibroblasts in a subject
comprises administering to a subject a medicament comprising an effective amount of a
cytotoxic small molecule compound, as described herein, that reduces the amount of
20 hypertrophic fibroblasts in the subject.

[0029] In an embodiment, a cytotoxic small molecule compound, as described herein, as a
biomarker/companion diagnostic (CDx) for selecting a suitable population to treat with LIF
inhibitors by using the cytotoxic compounds to down-regulate phosphorylation of STAT3.

BRIEF DESCRIPTION OF THE DRAWINGS

25 [0030] Advantages of the present invention will become apparent to those skilled in the art
with the benefit of the following detailed description of embodiments and upon reference to
the accompanying drawings in which:

[0031] FIG. 1 depicts *in vitro* tumorigenicity potential of EC332 in T47D breast cancer
cells

30 FIG. 2 shows that EC330 inhibited angiogenesis *in vitro* (tube formation assay);

FIG. 3 shows that alpha-smooth muscle mediated cytoskeletal disruption of fibroblast in human endometrial stromal cells (HESC) cells treated with EC330/332;

FIG. 4A shows the percent of apoptosis induced by treatment with EC330/332;

5 FIG. 4B shows the effect of EC330 on P53 for mutant vs. wild type glioma cells;

FIG. 5 shows that EC330/332 restore P53 levels by inhibiting MDM2 in MCF-7 cells;

FIG. 6A depicts a graph of tumor volume vs. time for the administration of EC330 at 0.5 mg/kg 5 days per week in the MDA-MB-231 (TNBC) Xenograft (*p<0.001);

10 FIG. 6B depicts a graph of tumor volume vs. time for the administration of EC330 at 2.5 mg/kg twice weekly in the MDA-MB-231 (TNBC) Xenograft (*p<0.001);

FIG. 7 depicts a graph of tumor volume vs. time for the administration of EC330 at 5 mg/kg 5 days per week in the IGROV1 (Ovarian) Xenograft (*p<0.001);

15 FIG. 8 depicts the percentage of apoptosis induced by EC330 measured in IGROV1 ovarian cancer xenograft tumors (**p<0.001);

FIG. 9 depicts the percentage of apoptosis induced by EC330 measured in MDA-MB-231 breast cancer xenograft tumors (**p<0.001);

FIG. 10 shows the results of immunohistochemical analysis of various cells treated with EC330 and EC332; and

20 FIG. 11 shows the proposed mechanism of action of EC330/EC332 on cancer cells.

[0032] While the invention may be susceptible to various modifications and alternative forms, specific embodiments thereof are shown by way of example in the drawings and will herein be described in detail. The drawings may not be to scale. It should be understood, however, that the drawings and detailed description thereto are not intended to limit the invention to the particular form disclosed, but to the contrary, the intention is to cover all modifications, equivalents, and alternatives falling within the spirit and scope of the present invention as defined by the appended claims.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0033] It is to be understood the present invention is not limited to particular devices or biological systems, which may, of course, vary. It is also to be understood that the

terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting. As used in this specification and the appended claims, the singular forms “a”, “an”, and “the” include singular and plural referents unless the content clearly dictates otherwise. Thus, for example, reference to “a linker” includes one or more
5 linkers.

[0034] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art.

[0035] Compounds described herein embrace both racemic and optically active compounds. Chemical structures depicted herein that do not designate specific stereochemistry are
10 intended to embrace all possible stereochemistries.

[0036] It will be appreciated by those skilled in the art that compounds having one or more chiral center(s) may exist in and be isolated in optically active and racemic forms. Some compounds may exhibit polymorphism. It is to be understood that the present invention encompasses any racemic, optically-active, polymorphic, or stereoisomeric form, or mixtures
15 thereof, of a compound. As used herein, the term “single stereoisomer” refers to a compound having one or more chiral center that, while it can exist as two or more stereoisomers, is isolated in greater than about 95% excess of one of the possible stereoisomers. As used herein a compound that has one or more chiral centers is considered to be “optically active” when isolated or used as a single stereoisomer.

[0037] The term “alkyl” as used herein generally refers to a radical substituent containing the monovalent group C_nH_{2n} , where n is an integer greater than zero. In some embodiments n is 1 to 12. The term “alkyl” includes a branched or unbranched monovalent hydrocarbon radical. Examples of alkyl radicals include, but are not limited to: methyl, ethyl, propyl, isopropyl, butyl, iso-butyl, sec-butyl, pentyl, 3-pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl,
25 dodecyl. When the alkyl group has from 1-6 carbon atoms, it is referred to as a “lower alkyl.” Suitable lower alkyl radicals include, but are not limited to, methyl, ethyl, n-propyl, i-propyl, 2-propenyl (or allyl), n-butyl, t-butyl, and i-butyl (or 2-methylpropyl).

[0038] The term “cycloalkyl” as used herein generally refers to a radical substituent containing the monovalent group C_nH_{2n-1} , where n is an integer greater than zero and wherein
30 the carbons C_1 and C_n are coupled to each other to form a ring. In some embodiments n is 3 to 8. Examples of cycloalkyl radicals include, but are not limited to: cyclopropyl ($n=3$),

cyclobutyl (n=4), cyclopentyl (n=5), cyclohexyl (n=6), cycloheptyl (n=7), and cyclooctyl (n=8).

[0039] The term “alkoxy” generally refers to an -OR group, where R is a lower alkyl, substituted lower alkyl, aryl, substituted aryl, aralkyl or substituted aralkyl. Suitable alkoxy radicals include, but are not limited to, methoxy, ethoxy, phenoxy, t-butoxy, methoxyethoxy, and methoxymethoxy.

[0040] The term “alkylacyl” denotes groups -C(O)R where R is alkyl as defined herein.

[0041] The term “cycloalkylacyl” denotes groups -C(O)R where R is a cycloalkyl. Examples of cycloalkylacyl compounds include, but are not limited to, cyclopropylacyl-, cyclopentylacyl and cyclohexylacyl.

[0042] The term “heterocycle” as used herein generally refers to a closed-ring structure, in which one or more of the atoms in the ring is an element other than carbon. Heterocycle may include aromatic compounds or non-aromatic compounds. Heterocycles may include rings such as thiophene, pyridine, isoxazole, phthalimide, pyrazole, indole, furan, or benzo-fused analogs of these rings. Examples of heterocycles include tetrahydrofuran, morpholine, piperidine, pyrrolidine, and others. In some embodiments, “heterocycle” is intended to mean a stable 5- to 7-membered monocyclic or bicyclic or 7- to 10-membered bicyclic heterocyclic ring which is either saturated or unsaturated, and which consists of carbon atoms and from 1 to 4 heteroatoms (e.g., N, O, and S) and wherein the nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen may optionally be quaternized, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. In some embodiments, heterocycles may include cyclic rings including boron atoms. The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom that results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. Examples of such heterocycles include, but are not limited to, 1H-indazole, 2-pyrrolidonyl, 2H,6H-1,5,2-dithiazinyl, 2H-pyrrolyl, 3H-indolyl, 4-piperidonyl, 4aH-carbazole, 4H-quinoliziny, 6H-1,2,5-thiadiazinyl, acridinyl, azocinyl, benzofuranyl, benzothiophenyl, carbazole, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, furanyl, furazanyl, imidazolidinyl, imidazoliny, imidazolyl, indolinyl, indoliziny, indolyl, isobenzofuranyl, isochromanyl, isoindolinyl, isoindolyl, isoquinolinyl (benzimidazolyl), isothiazolyl, isoxazolyl, morpholinyl, naphthyridinyl, octahydroisoquinolinyl, oxazolidinyl, oxazolyl, phenanthridinyl,

WO 2016/154203

PCT/US2016/023582

phenanthrolyl, phenarsaziny, phenaziny, phenothiaziny, phenoxathiiny, phenoxaziny, phthalaziny, piperaziny, piperidiny, pteridiny, puriny, pyran, pyraziny, pyrazolidiny, pyrazoliny, pyrazoly, pyridaziny, pyridiny, pyridyl, pyrimidiny, pyrrolidiny, pyrroliny, pyrroly, quinazoliny, quinoliny, quinoxaliny, quinuclidiny, carboliny, tetrahydrofurany, tetrahydroisoquinoliny, tetrahydroquinoliny, tetrazoly, thianthreny, thiazoly, thieny, thiopheny, triaziny, xanthy. Also included are fused ring and spiro compounds containing, for example, the above heterocycles.

[0043] As used herein the terms “alkenyl” and “olefin” generally refer to any structure or moiety having the unsaturation C=C. Examples of alkenyl radicals include, but are not limited to vinyl, 1-propenyl, 2-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-heptenyl, 2-heptenyl, 3-heptenyl, 4-heptenyl, 5-heptenyl, 1-nonenyl, 2-nonenyl, 3-nonenyl, 4-nonenyl, 5-nonenyl, 6-nonenyl, 7-nonenyl, 8-nonenyl, 1-decenyl, 2-decenyl, 3-decenyl, 4-decenyl, 5-decenyl, 6-decenyl, 7-decenyl, 8-decenyl, 9-decenyl; 1-undecenyl, 2-undecenyl, 3-undecenyl, 4-undecenyl, 5-undecenyl, 6-undecenyl, 7-undecenyl, 8-undecenyl, 9-undecenyl, 10-undecenyl, 1-dodecenyl, 2-dodecenyl, 3-dodecenyl, 4-dodecenyl, 5-dodecenyl, 6-dodecenyl, 7-dodecenyl, 8-dodecenyl, 9-dodecenyl, 10-dodecenyl, and 11-dodecenyl.

[0044] The term “pharmaceutically acceptable salts” includes salts prepared from by reacting pharmaceutically acceptable non-toxic bases or acids, including inorganic or organic bases, with inorganic or organic acids. Pharmaceutically acceptable salts may include salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic salts, manganous, potassium, sodium, zinc, etc. Examples include the ammonium, calcium, magnesium, potassium, and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, and basic ion exchange resins, such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-dibenzylethylenediamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethyl-morpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine, etc.



Cytotoxic Agents

WO 2016/154203

PCT/US2016/023582

[0045] Described herein are novel cytotoxic agent, which may be used in the treatment of cancer by acting through a new mechanism of action by simultaneous inhibition of leukemia inhibitory factor (LIF) and MDM2. One significant advantage of the described compounds is that they act directly on the tumor cells/tumor stem cells and on the surrounding stromal fibroblasts (tumors with desmoplastic stroma or hypertrophic cell mass) as well.

[0046] In one embodiment, a cytotoxic agent has the structure (I) or (II):




10

(I)

(II)

where:



15

X is O, NH, or S;

n = 1-18; m = 1-18;

20

R² is H, F, Cl, $-C(O)-R^6$, or $-CH_2(OH)$;

R³ is H, F, Cl, $-C(O)-R^6$, or $-CH_2(OR^6)$;

R⁴ is H, alkyl, $-CH_2-OH$, $-CO_2R^6$, $-CON(R^6)_2$;

25

R⁵ is alkyl, alkenyl, alkylacyl, cycloalkyl, heterocycle, $-CN$, alkoxy, $-N(R^6)_2$, $-CON(R^6)_2$,

WO 2016/154203


PCT/US2016/023582

$-\text{S}(\text{O})\text{R}^6$, $-\text{SR}_6$, $-\text{SO}_2\text{R}^6$; or $-(\text{CH}_2)_p\text{-CH}_2\text{-Y}$;

Y is H, OR^6 , SCH_3 , CF_3 , $-\text{N}(\text{R}^6)_2$; p = 1-18; and

5 R^6 is H, alkyl, or cycloalkyl.

[0047] In one embodiment, a cytotoxic agent has the structure (III):



10

Where:

R^2 is H, F, Cl, $-\text{CO}-$, or $-\text{C}(\text{OH})-$;

15

R^3 is H, F, Cl, $-\text{CO}-$; or $-\text{C}(\text{OH})-$;


R^4 is H, alkyl, $-\text{CH}_2\text{-OH}$, $-\text{CO}_2\text{R}^6$, $-\text{CON}(\text{R}^6)_2$;

20 R^5 is alkyl, alkenyl, alkylacyl, cycloalkyl, heterocycle, $-\text{CN}$, alkoxy, $-\text{N}(\text{R}^6)_2$, $-\text{CON}(\text{R}^6)_2$, $-\text{S}(\text{O})\text{R}^6$, $-\text{SR}_6$, $-\text{SO}_2\text{R}^6$; or $-(\text{CH}_2)_p\text{-CH}_2\text{-Y}$; and

Y is H, OR^6 , SCH_3 , CF_3 , $-\text{N}(\text{R}^6)_2$; p = 1-18; and

25 R^6 is H, alkyl, or cycloalkyl.

[0048] In one embodiment, a cytotoxic agent has the structure (IV):



Where:


5

R⁵ is alkyl, alkenyl, alkylacyl, cycloalkyl, heterocycle, -CN, alkoxy, -N(R⁶)₂, -CON(R⁶)₂, -S(O)R⁶, -SR₆, or -SO₂R⁶; and

R⁶ is H, alkyl, or cycloalkyl.

10

[0049] In a specific embodiment, a cytotoxic agent has the structure **(IV)**:



15

Where:













R⁵ is alkyl, alkenyl, alkylacyl, cycloalkyl, 1,3-imidazolyl, alkoxy, -N(R⁶)₂, -SR₆, or -SO₂R⁶; and

20

R⁶ is H, lower alkyl, or cycloalkyl.

[0050] In a specific embodiment, a cytotoxic agent has the structure (IV), where R^5 is alkyl, alkenyl, or cycloalkyl. In a specific embodiment, a cytotoxic agent has the structure (IV), where R^5 is 1,3-imidazolyl. In a specific embodiment, a cytotoxic agent has the structure (IV), where R^5 is alkoxy. In a specific embodiment, a cytotoxic agent has the structure (IV), where R^5 is $-N(R^6)_2$. In a specific embodiment, a cytotoxic agent has the structure (IV), where R^5 is $-SR_6$. In a specific embodiment, a cytotoxic agent has the structure (IV), where R^5 is $-SO_2R^6$. For each of these embodiments, where appropriate, R^6 is H, lower alkyl, or cycloalkyl.

[0051] Specific examples of cytotoxic agents include:




10

Synthesis of Cytotoxic Agents

[0052] Compounds having general formula (I) may be synthesized as outlined in the following general scheme (Scheme 1).

15




Scheme 1. i) Epoxidation; ii) Grignard reaction; ii) Ac_2O , DMAP, Py; iii) 3-bromo-3,3-difluoro-1-triisopropylsilylpropyne, n-BuLi, THF; iv) 4N, HCl; v) TBAF, THF.

5

[0053] The general synthesis of the cytotoxic compounds described herein is generally accomplished by the use of a triisopropylsilyl (TIPS) protecting group to protect the 17a acetylenic hydrogen. Thus, an important intermediate in the synthesis of compounds having an acetylenic hydrogen is the compound of structure (V):

10



where:

15 R^5 is alkyl, alkenyl, alkylacyl, cycloalkyl, heterocycle, -CN, alkoxy, $-\text{N}(\text{R}^6)_2$, $-\text{CON}(\text{R}^6)_2$, $-\text{S}(\text{O})\text{R}^6$, $-\text{SR}^6$, $-\text{SO}_2\text{R}^6$; or $-(\text{CH}_2)_p-\text{CH}_2-\text{Y}$; and

Y is H, OR^6 , SCH_3 , CF_3 , $-\text{N}(\text{R}^6)_2$; $p = 1-18$; and


20

R^6 is H, alkyl, or cycloalkyl.

[0054] Specific examples of synthetic intermediates include compounds having the structure (V), where: R^5 is alkyl, alkenyl, alkylacyl, cycloalkyl, 1,3-imidazolyl, alkoxy, $-N(R^6)_2$, $-SR_6$, or $-SO_2R^6$; and R^6 is H, lower alkyl, or cycloalkyl.

5 Synthesis of EC 330

[0055] EC330 may be synthesized by following the scheme outlined below (Scheme 2).



Scheme 2

10

Intermediate **2** may be synthesized following the procedure reported in Rao et al. "New 11β -aryl-substituted Steroids Exhibit Both Progestational and Antiprogestational Activity", *Steroids*, 1998, 63, 523, which is incorporated herein by reference. Conjugate Grignard addition of the aryl cuprate reagent, generated by the reaction of 4-bromo-4'-

15


cyclopropylbenzene with magnesium and cuprous chloride, at the 11 position of **2** afforded intermediate **3a**. The cyclopropyl aryl derivative **3a** was subjected to 5-hydroxy elimination using acetic anhydride and pyridine in presence of DMAP to afford **4a**. The 17 addition of 3-lithium 3,3-difluoro-1-triisopropylsilylpropyne, generated by the reaction of 3-bromo-3,3-difluoro-1-triisopropylsilyl propyne with n-butyl lithium at -78°C afforded compound **5a**,

20

which was hydrolyzed by 4N hydrochloric acid to give the key intermediate **6a**. Removal of the triisopropylsilyl (TIPS) group by tetrabutylammonium fluoride (TBAF) afforded the final compound EC330.

25 Synthesis of EC 332

[0056] EC332 may be synthesized by following the scheme outlined below (Scheme 3).




Scheme 3

5 Synthesis of intermediates up to **4b** may be accomplished following the procedure reported in Nickisch et al. "Synthesis and Biological Evaluation of Partially Fluorinated Antiprogestins and Mesoprogestins", *Steroids*, 2013, 78, 255, which is incorporated herein by reference. The 17 addition of lithium 3,3-difluoro-1-triisopropylsilylpropyne on **4b** afforded **5b**, which upon 3-ketal hydrolysis followed by removal of the TIPS group using TBAF provided EC-
10 332.

Synthesis of EC 351

[0057] EC351 may be synthesized by following the scheme outlined below (Scheme 4).



Scheme 4


15

The conjugate Grignard addition of 1,4-diiodobenzene in the presence of isopropyl magnesium chloride and cuprous chloride on compound **2** afforded **3c**. Ullman coupling of

3c with imidazole in presence of cuprous iodide afforded **4c** according to the procedure set forth in Nickisch et al. "Synthesis and Biological Evaluation of 11' Imidazolyl Antiproggestins and Mesoproggestins, Steroids, 2014, 92, 45-55, which is incorporated herein by reference. Subsequent 17 addition of lithium 3,3-difluoro-1-triisopropylpropyne followed by 3-ketal hydrolysis and the TIPS removal by TBAF furnished EC-351.

Synthesis of EC 359

[0058] EC359 may be synthesized by following the scheme outlined below (Scheme 5).



10


15

EC359 synthesis followed the same pattern as the previous schemes, except the synthesis of compound **3k** which was prepared by the Suzuki coupling of **3c** with the corresponding aryl boronic acid in presence of a palladium catalyst.

Synthesis of EC352, EC356, EC357, EC358, EC360, EC362 and EC363

[0059] EC352, EC356, EC357, EC358, EC360, EC362 and EC363 may be synthesized by following the scheme outlined below (Scheme 6).

20




Scheme 6

- 5 Epoxide **2** was subjected to conjugate Grignard addition with the corresponding 4-bromobenzene derivative in presence of magnesium and cuprous chloride to afford compounds **3d-j**. Acetic anhydride and pyridine mediated 5-hydroxy elimination gave compounds **4d-j**. Subsequent 1,7 addition of lithium 3,3-difluoro-1-triisopropylpropyne followed by 3-ketal hydrolysis and TIPS removal by TBAF furnished the respective EC
- 10 compounds as shown in the scheme.

Synthesis of EC361

[0060] EC361 may be synthesized by following the scheme outlined below (Scheme 7).



Scheme 7

Conjugate Grignard reaction of 4-bromo-4'-cyclopropylphenyl sulfide and epoxide **2** in the presence of magnesium and cuprous chloride furnished compound **3I**. Acetic anhydride and pyridine mediated 5-hydroxy elimination gave **4I**, which upon addition of lithium 3,3-difluoro-1-triisopropylpropyne afforded **5I**. The 3-ketal hydrolysis of **5I** by 4N hydrochloric acid followed by TIPS removal by TBAF gave EC361.

10 Cytotoxicity Studies – Use As Anticancer Agents

[0061] Compounds having the above described structures showed potent cytotoxicity in routine screening. These compounds were further tested to confirm the cytotoxicity in various cancer cell lines. The activity was confirmed to be dose dependent in a NCI-60 cell line panel that includes leukemia, non-small cell lung cancer, colon cancer, CNS cancer, melanoma, ovarian cancer, renal cancer, breast and prostate cancer cell lines. The high cytotoxic effects of these compounds were totally unexpected. Further studies showed the compound to reduce the tumor burden in human breast cancer and ovarian cancer xenograft models in mice. The compound exhibited specificity towards artificially induced LIF overexpressing cells over regular cancer cells. One significant advantage of the described compounds is that they act directly on the tumor cells/tumor stem cells and on the surrounding stromal fibroblasts (tumors with desmoplastic stroma or hypertrophic cell mass) as well.

Mechanism of action studies


[0062] Decidualization studies: The above compounds were investigated originally for the treatment of endometriosis. The compounds were tested in a decidualization assay using

human endometrial stromal fibroblast (HESC) cells. In preparation for the possibility of embryonic attachment, the stromal cells in the endometrium undergo extensive waves of proliferation, remodeling, and terminal differentiation that transforms the endometrium into an endocrine gland, called decidua. This transformation is called decidualization and it is
5 under the control by ovarian hormones.

[0063] Apoptosis has been shown to be important for endometrial function. The level of apoptosis increases from the proliferative phase through menstrual cycle and peaks at menses. This experiment will demonstrate that disruption of actin filaments will induce apoptosis in endometrial stromal fibroblast cells. However if the cells are subjected to conditions that
10 induce decidualization, stromal cells will begin to differentiate instead of undergoing apoptosis. The above described compounds showed potent actin cytoskeletal disruption evident from phalloidin staining.

[0064] The above opens up various possibilities to study the mechanism of these compounds. First of all, the most common type of stromal cells are fibroblasts. In preparation for the
15 implanting blastocyst, the endometrium becomes increasingly vascular with prominent increase in the levels of VEGF and PDGF and other cytokines and chemokines as well as alpha-smooth muscle actin is also induced in fibroblasts. Leukemia inhibitory factor (LIF) is a pleotropic cytokine from interleukin-6 (IL-6) family and has been shown to enhance oocyte maturation and preimplantation development. Recently evidence shows that LIF mediates
20 proinvasive activation of tumor associated fibroblasts (CAF) in cancers. Moreover, additional evidence proved that LIF negatively regulates tumor suppressor p53 through STAT3 /ID1/MDM2 axis in colorectal cancers. These recent studies prompt us to investigate whether the compounds above (specific examples of which are referred to herein as EC330/332) possess anticancer activity that mediated through LIF.

25



[0065] EC330 and EC332 exhibited IC₅₀'s in the low nanomolar concentrations in cytotoxicity assays (Table 1). They also exhibited 50% inhibition in the number of colonies formed at 5 nM concentration and a complete inhibition or no colonies were formed at 100nM concentration (FIG.2) in the soft agar colony formation assay.

5 [0066] EC330 and EC332 also showed complete abrogation of tube formation at 1μM concentration at all measured time points in a human umbilical vein endothelial cells (HUVEC) tube formation assay. These results clearly underscore the anti-angiogenic activity of the compounds. Vascular endothelial growth factor (VEGF) is the most prominent among the angiogenic cytokines and is believed to play a central role in the process of
10 neovascularization, both in cancer as well as other inflammatory diseases. A compound that inhibits angiogenesis can be used as monotherapy or in combination with conventional chemotherapy.

[0067] The compounds described herein exhibit indirect diminishment of the phosphorylation of SMAD thorough blocking TGF-beta. p53 levels in tumors were increased
15 with treatment and levels of MDM2 were reduced. The downstream effector of LIF, STAT3 phosphorylation was found reduced in the treated samples when compared to untreated control.

[0068] STAT3 phosphorylation may be utilized as a diagnostic maker/companion diagnostic to identify suitable patient population those could get benefit of LIF inhibition.
20 Overexpression of LIF is associated with poor prognosis and increase incidence of chemoresistance. Targeting LIF and MDM2 to reactivate p53 is a potential therapeutic strategy for chemotherapy as well as in combination with other agents to alleviate chemoresistance. The dual inhibition of LIF and MDM2 would benefit a complete inhibition of tumor cells by inhibiting both tumor epithelium and it surrounding stromal fibroblast or
25 desmoplastic stroma.

[0069] Clinical presentation of a lump in the breast is histologically viewed as a collagenous tumor or desmoplastic response created by myofibroblasts of the tumor stroma. The stroma of the prostate is characteristically muscular and diagnosis of reactive stroma associated with prostate cancer is one of poor prognosis. Recent studies show that targeting the stromal
30 compartment in pancreatic cancer may have antitumor effects and may enhance sensitivity to radiation and chemotherapy. Disease progression in pancreatic cancer is associated with a

robust fibrotic response, or desmoplasia, that promotes tumor progression and inhibits the entrance of therapeutic agents.

[0070] However, when we artificially induced LIF in human breast cancer cells (MCF-7), the compounds showed specificity towards these cells over regular breast cancer cells. The compounds prepared in the EC330 series showed specificity towards LIF in a range of 2 to 20 fold in term of cytotoxicity. We have found that the disclosed compounds showed antagonistic/agonistic property towards PR *in vitro*. Further studies revealed that the compound (EC330) significantly reduced the tumor growth in human triple negative breast cancer (TNBC) and ovarian cancer models.

10 Biological testing

The following assays were performed:

Cytotoxicity assays

[0071] In order to identify the mechanism of action of EC330/332, we checked cytotoxicity of these compounds in various cancer cells lines and derived IC50 values. Briefly, 5×10^3 cells were seeded in 96-well plates and incubated with compounds (0.0001–10 $\mu\text{mol/L}$) or dimethyl sulfoxide (DMSO; 0.02% v/v) for 24, 48, and 72 hours at 37°C and cell viability was measured using a Fluoroscan plate reader. Results of cytotoxicity testing is presented in Table 11.

[0072] EC 330 was also tested using the NCI-60 Screening Methodology. The human tumor cell lines of the cancer screening panel are grown in RPMI 1640 medium containing 5% fetal bovine serum and 2 mM L-glutamine. For a typical screening experiment, cells are inoculated into 96 well microtiter plates in 100 μL at plating densities ranging from 5,000 to 40,000 cells/well depending on the doubling time of individual cell lines. After cell inoculation, the microtiter plates are incubated at 37° C, 5 % CO₂, 95 % air and 100 % relative humidity for 24 h prior to addition of experimental drugs.

[0073] After 24 h, two plates of each cell line are fixed in situ with TCA, to represent a measurement of the cell population for each cell line at the time of drug addition (Tz). EC330 was solubilized in dimethyl sulfoxide at 400-fold the desired final maximum test concentration and stored frozen prior to use. At the time of drug addition, an aliquot of frozen concentrate is thawed and diluted to twice the desired final maximum test

WO 2016/154203

PCT/US2016/023582

concentration with complete medium containing 50 µg/ml gentamicin. Additional four, 10-fold or ½ log serial dilutions are made to provide a total of five drug concentrations plus control. Aliquots of 100 µl of these different drug dilutions are added to the appropriate microtiter wells already containing 100 µl of medium, resulting in the required final drug concentrations.

[0074] Following drug addition, the plates are incubated for an additional 48 h at 37°C, 5 % CO₂, 95 % air, and 100 % relative humidity. For adherent cells, the assay is terminated by the addition of cold TCA. Cells are fixed in situ by the gentle addition of 50 µl of cold 50 % (w/v) TCA (final concentration, 10 % TCA) and incubated for 60 minutes at 4°C. The supernatant is discarded, and the plates are washed five times with tap water and air dried. Sulforhodamine B (SRB) solution (100 µl) at 0.4 % (w/v) in 1 % acetic acid is added to each well, and plates are incubated for 10 minutes at room temperature. After staining, unbound dye is removed by washing five times with 1 % acetic acid and the plates are air dried. Bound stain is subsequently solubilized with 10 mM trizma base, and the absorbance is read on an automated plate reader at a wavelength of 515 nm. For suspension cells, the methodology is the same except that the assay is terminated by fixing settled cells at the bottom of the wells by gently adding 50 µl of 80 % TCA (final concentration, 16 % TCA). Using the seven absorbance measurements [time zero, (Tz), control growth, (C), and test growth in the presence of drug at the five concentration levels (Ti)], the percentage growth is calculated at each of the drug concentrations levels. Percentage growth inhibition is calculated as:

$$[(Ti-Tz)/(C-Tz)] \times 100 \text{ for concentrations for which } Ti \geq Tz$$

$$[(Ti-Tz)/Tz] \times 100 \text{ for concentrations for which } Ti < Tz.$$

[0075] Three dose response parameters are calculated for each experimental agent. Growth inhibition of 50 % (GI₀) is calculated from $[(Ti-Tz)/(C-Tz)] \times 100 = 50$, which is the drug concentration resulting in a 50% reduction in the net protein increase (as measured by SRB staining) in control cells during the drug incubation. The drug concentration resulting in total growth inhibition (TGI) is calculated from $Ti = Tz$. The LC₅₀ (concentration of drug resulting in a 50% reduction in the measured protein at the end of the drug treatment as compared to that at the beginning) indicating a net loss of cells following treatment is calculated from $[(Ti-Tz)/Tz] \times 100 = -50$. Values are calculated for each of these three parameters if the level of activity is reached; however, if the effect is not reached or is

exceeded, the value for that parameter is expressed as greater or less than the maximum or minimum concentration tested. Results of testing for EC330 are presented in Tables 2-10.

Panel/Cell Line	GI50 (log ₁₀ GI50)	TGI (log ₁₀ TGI)	LC50 (log ₁₀ LC50)
Leukemia			
CCRF-CEM	<-8.00	<-8.00	<-8.00
HL-60(TB)	<-8.00	<-8.00	<-8.00
K-562	-5.66	>-4.00	>-4.00
MOLT-4	-5.77	-5.30	>-4.00
RPMI-8226	-6.95	-6.24	>-4.00

TABLE 2

Panel/Cell Line	GI50 (log ₁₀ GI50)	TGI (log ₁₀ TGI)	LC50 (log ₁₀ LC50)
Non-Small Cell Lung Cancer			
A549/ATCC	-5.61	-4.61	>-4.00
EKVX	<-8.00	<-8.00	-7.44
HOP-62	-7.85	-6.79	-5.52
HOP-92	<-8.00	<-8.00	-7.72
NCI-H226	-5.58	-5.04	>-4.00
NCI-H23	<-8.00	<-8.00	>-4.00
NCI-H332M	-5.42	>-4.00	>-4.00
NCI-H460	-5.57	-4.80	>-4.00
NCI-H522	<-8.00	<-8.00	<-8.00

TABLE 3

Panel/Cell Line	GI50 (log ₁₀ GI50)	TGI (log ₁₀ TGI)	LC50 (log ₁₀ LC50)
Colon Cancer			
COLO 205	-6.10	-5.53	-5.05
HCC-2998	-5.53	-4.37	>-4.00
HCT-116	-5.55	-4.55	>-4.00
HCT-15	-5.83	>-4.00	>-4.00
HT29	<-8.00	-6.72	-6.16
KM12	<-8.00	-5.91	-4.35
SW-620	-5.60	-4.79	>-4.00

TABLE 4

Panel/Cell Line	GI50 (log ₁₀ GI50)	TGI (log ₁₀ TGI)	LC50 (log ₁₀ LC50)
CNS Cancer			
SF-268	-6.90	-6.43	-5.81
SF-295	<-8.00	<-8.00	<-8.00
SF-539	<-8.00	-7.77	-7.28
SNB-19	-5.42	>-4.00	>-4.00

U251	-6.82	-5.78	-5.27
------	-------	-------	-------

TABLE 5

Panel/Cell Line	GI50 (log ₁₀ GI50)	TGI (log ₁₀ TGI)	LC50 (log ₁₀ LC50)
Melanoma			
LOX IMVI	<-8.00	<-8.00	<-8.00
MALME-3M	<-8.00	<-8.00	-7.09
M14	-5.56	-4.45	>-4.00
MDA-MB-435	-5.73	-5.03	>-4.00
SK-MEL-2	-5.67	-5.19	-4.52
SK-MEL-28	-5.75	-5.43	-5.11
SK-MEL-5	-5.33	-4.79	-4.36
UACC-257	-5.15	>-4.00	>-4.00
UACC-62	-5.71	-5.27	>-4.00

TABLE 6

Panel/Cell Line	GI50 (log ₁₀ GI50)	TGI (log ₁₀ TGI)	LC50 (log ₁₀ LC50)
Ovarian Cancer			
IGROV1	<-8.00	<-8.00	
OVCAR-3	-7.89	-6.85	-6.03
OVCAR-4	-5.41	>-4.00	>-4.00
OVCAR-5	-5.97	-5.25	>-4.00
OVCAR-8	-5.81	-4.43	>-4.00
NCI/ADR-RES	-7.59	-5.41	>-4.00
SK-OV-3	-6.93	-6.55	-6.18

TABLE 7

Panel/Cell Line	GI50 (log ₁₀ GI50)	TGI (log ₁₀ TGI)	LC50 (log ₁₀ LC50)
Prostate Cancer			
PC-3	-6.29	-5.53	>-4.00
DU-145	-5.72	-4.75	>-4.00

TABLE 8

Panel/Cell Line	GI50 (log ₁₀ GI50)	TGI (log ₁₀ TGI)	LC50 (log ₁₀ LC50)
Renal Cancer			
786-0	<-8.00	<-8.00	<-8.00
A498	-6.58	-6.01	-5.41
ACHN	-5.82	-5.09	>-4.00
CAKI-1	-7.53	-6.41	>-4.00
RXF 393	-5.97	-5.09	>-4.00
SN12C	-5.42	-4.58	>-4.00
TK-10	-5.58	-4.91	>-4.00
UO-31	<-8.00	<-8.00	<-8.00

TABLE 9

Panel/Cell Line	GI50 (log ₁₀ GI50)	TGI (log ₁₀ TGI)	LC50 (log ₁₀ LC50)
Breast Cancer			
MCF7	-5.91	-4.96	>-4.00
MDA-MB-231 ATCC	-6.26	-5.46	>-4.00
HS 578T	-5.64	-4.99	>-4.00
BT-549	-5.68	-5.30	>-4.00
T-47D	<-8.00	-5.00	>-4.00
MDA-MB-468	-6.69	-6.09	-4.30

TABLE 10

Soft agar colony formation assay

[0076] Colonies of cancer cells formed in soft agar in the presence and absence of the testing compounds is a standard assay to interpret *in-vitro* tumorigenic potential in the basal layer. Agar was prepared by mixing 1% DNA grade agar melted and cooled to 40°C with an equal volume of (2×) Dulbecco's Modified Eagle's Medium (DMEM) to obtain 0.5% agar that was dispersed in a 6-well plate and allowed to solidify. A total of 0.6% agar was prepared in RPMI medium and mixed together with T47D cells (0.5 × 10⁶ cells/mL) and immediately plated on the basal layer in the presence or absence of testing compounds. The cultures were incubated at 37°C in a CO₂ incubator for 2 weeks, and colonies were stained with 0.005% crystal violet and observed under a light microscope. FIG. 1 depicts *in vitro* tumorigenicity potential of EC332 in T47D breast cancer cells.

Angiogenesis assay

[0077] Matrigel *in vitro* HUVECs tube formation assay: The tube formation assay was performed using 12-well plate coated with 100 μL Matrigel basement membrane matrix (BD Bioscience) per well and polymerized at 37 °C for 30 min. Human umbilical vascular endothelial cells (HUVECs) suspended in M199 medium containing 2% FBS were plated on the Matrigel at a density of 2 × 10⁵ cells/well. Compounds (0.0001- 10 μM) were then added together with VEGF. After 8 h, the Matrigel-induced morphological changes were photographed and the extent of capillary tube formation was evaluated by measuring the total tube length per field. FIG. 2 shows that EC330 inhibited angiogenesis *in vitro* (tube formation assay).

Decidualization assay

WO 2016/154203

PCT/US2016/023582

[0078] Rationale of this assay is described above in *mechanism of action* studies section. Briefly, human stromal endometrial cells were plated in the presence and absence of progesterone and allow undergoing decidualization. The cells treated with test compounds of different concentrations and stained for actin cytoskeletal morphology using phalloidin staining and monitored under fluorescent microscope. The compounds that inhibited actin cytoskeletal polymerization was detected and compared with cytochalasin D. FIG. 3 shows that alpha-smooth muscle mediated cytoskeletal disruption of fibroblast in human endometrial stromal cells (HESC) cells treated with EC330/332.

Apoptosis assay

[0079] Caspase-3/7 activity in HESE cells was measured using Caspase-Glo assay kit (Promega), as described before. Briefly, cells were homogenized in homogenization buffer (25 mmol/L HEPES, pH 7.5, 5 mmol/L -MgCl₂, and 1 mmol/L EGTA), protease inhibitors, and the homogenate was centrifuged at 13,000 rpm at 4°C for 15 minutes. To 10 µL of the supernatant containing protein was added to an equal volume of the assay reagent and incubated at room temperature for 2 hours. The luminescence was measured using a luminometer. The percent of apoptosis induced by treatment with EC330/332 is shown in FIG. 4A. FIG. 4B shows the effect of EC330 on P53 for mutant vs. wild type glioma cells. FIG. 5 shows that EC330/332 restore P53 levels by inhibiting MDM2 in MCF-7 cells.

20 Immunohistochemistry analysis

[0080] Patient derived tumor (melanoma) was treated with the compounds at 10nM and 1µM for 5 days in RPMI medium and harvested on day 6 using established protocol and immunohistochemistry was performed for different antibodies including p53, LIF, MDM2, pSTAT3, Ki67. Results of immunohistochemical analysis are shown in FIG. 10. The results show that EC332 & EC330 induce apoptosis and restore p53 by inhibiting MDM2 and LIF mediated STAT3 phosphorylation.

Western blotting

[0081] In brief, T47D cells were treated with compounds for 3 days at different concentrations (10, 100nM) and cell lysates were separated by 8% SDS-PAGE and transferred to polyvinylidene difluoride membranes. Membranes were then incubated with primary antibodies including phosphorylated and/or total p53, MDM2, actin, pSTAT3. After

overnight incubation at 4 °C, membranes were incubated with secondary antibodies.

Immunoreactive bands were then visualized by the enhanced chemiluminescence (ECL) detection system (GE healthcare).

Tumor xenograft study

5 [0082] Uniform suspensions of human breast cancer MDA-MB-231 and human ovarian
cancer IGROV1 cells (2×10^6) in 100 μ L (0.02 carboxymethyl cellulose in phosphate
buffered saline) were injected subcutaneously into the right and left flanks of 4- to 5-week-
old female athymic nude mice (Charles River Laboratories). After 10 days, when the tumor
diameter reaches 100 mm^3 , the mice were randomly allocated to 3 groups of each containing
10 6 animals. Group 1 served as the untreated control, groups 2 and 3 received EC330
intraperitoneally at 0.5mg/kg (daily dose for 5 days/week for 4 weeks) and 2.5mg/kg (twice
a week for 4 weeks), respectively. All drug was suspended in PBS followed by sonication
(15 seconds with an intermittent interval of 5 seconds) for 1 minute. Tumors were allowed
to reach palpability before drug intervention. Tumor size was measured every 3 days using
15 digital Vernier Calipers and tumor volume was calculated using the ellipsoid formula $[D \times$
 $(d^2)]/2$, where D is the large diameter of the tumor and d represents the small diameter. On
day 31, the mice were euthanized and tumors were harvested for protein and gene
expression studies. FIG. 6A depicts a graph of tumor volume vs. time for the administration
of EC330 at 0.5 mg/kg 5 days per week in the MDA-MB-231 (TNBC) Xenograft. FIG. 6B
20 depicts a graph of tumor volume vs. time for the administration of EC300 at 2.5 mg/kg twice
weekly in the MDA-MB-231 (TNBC) Xenograft. It is believed that STAT3
phosphorylation is reduced in tumors that underwent treatment with EC330 compared to
control as an indication of LIF downstream targets. FIG. 7 depicts a graph of tumor volume
vs. time for the administration of EC300 at 5 mg/kg 5 days per week in the IGROV1
25 (Ovarian) Xenograft. The compound EC330 showed potent antitumor activity at doses of 5
mg/kg (5 days per week) and post tumor treatment for 4 weeks. STAT3 phosphorylation is
reduced in tumors that underwent treatment compared to control which is an indication of
LIF downstream target. FIG. 8 depicts the percentage of apoptosis induced by EC330
measured in IGR-OV1 ovarian cancer xenograft tumors. FIG. 9 depicts the percentage of
30 apoptosis induced by EC330 measured in MDA-MB-231 breast cancer xenograft tumors.
FIGS. 8 and 9 show that EC330 induced apoptosis dose dependently.

[0083] The proposed mechanism of action of EC330/EC332 on cancer cells is shown in FIG. 11.

[0084] Table 11 depicts short term (72h) cytotoxicity of various cytotoxic compounds described herein in various cancer cell lines.

Compounds	IC50 (nM)	Cell line tested
EC330	35 34 25 20 25 80 100 30 53 5	IGROV-1 (Ovarian cancer cells) MDA-MB-231 (TNBC cells) U87-MU (Glioma cells) U251 (Glioma) A549 (Lung cancer) T47D (ER+ PR+ Breast Cancer) MCF-7 (ER+ Breast Cancer) PC3 (AR- Prostate Cancer) LNCaP (AR+ Prostate) Ishikawa (Endometrial Cancer)
EC332	90 40 75 110 45 200 300 96 10	U87-MU (Glioma cells) U251 (Glioma) A549 (Lung cancer) MDA-MB-231 (TNBC Cells) MCF-7 (ER+ Breast Cancer) T47D (ER+ PR+ Breast Cancer) PC3 (AR- Prostate Cancer) LNCaP (AR+ Prostate) Ishikawa (Endometrial Cancer)
EC358	35 40 25	IGROV-1 MDA-MB-231 (TNBC cells) U87-MU (Glioma cells)
EC359	25 30 10	IGROV-1 MDA-MB-231 (TNBC cells) U87-MU (Glioma cells)
EC360	22 40 73	IGROV-1 MDA-MB-231 (TNBC cells) U87-MU (Glioma cells)
EC361	350 750 230	IGROV-1 MDA-MB-231 (TNBC cells) U87-MU (Glioma cells)
EC351	500	U251 (Glioma)
EC352	40 45 50	U251 (Glioma) A549 (Lung cancer) U87-MU (Glioma)
EC355	>10 (uM) >10 (uM)	U87-MU (Glioma) MDA-MB-231 (TNBC)

WO 2016/154203

PCT/US2016/023582

Compounds	IC50 (nM)	Cell line tested
EC356	40 45 45	A549 (Lung cancer) U251 (Glioma) U87-MU (Glioma cells)
EC362	25 7.5	T47D (Breast cancer) IGROV1 (Ovarian cancer)
EC363	30 30	IGROV-1 (Ovarian cancer cells) U87-MU (Glioma)
EC357	35 30 35	A549 (Lung cancer) U251 (Glioma) U87-MU (Glioma cells)

TABLE 11

LIF Specificity Assay

[0085] LIF specificity of EC330 and related compounds synthesized in this series was conducted by comparing cytotoxicity of LIF-overexpressing MCF-7 (“MCF-7 LIF”) vs. unmodified MCF-7 (“MCF-7”) human breast cancer cells. Table 12 presents the results of this assay. The screening results show that LIF overexpressing cells were more vulnerable to cell death by EC330 and related compounds.


Compound	MCF-7 IC50 (nm)	MCF-7 LIF IC50 (nm)	Fold Change
EC330	250	80	3
EC352	>10000	>10000	0
EC356	500	250	2
EC358	100	50	2
EC360	500	250	2
EC359	240	30	8
EC361	>10000	500	20
EC362	50	25	2

TABLE 12

Embryo implantation is a critical step in the establishment of pregnancy in humans and other higher vertebrates. It has been showed in the literature that uterine LIF is expressed in the luminal epithelium on the day 3 of pregnancy and mediates via STAT3 in mouse and LIF overexpression during implantation in women. Hence inhibitors that block LIF action during early pregnancy would block embryo implantation and terminate the pregnancy. The above compounds, as shown, can act as LIF inhibitors which will terminate pregnancy and, therefore, can be used as a contraceptive.

10 Synthesis – Detailed Procedures

[0086] All the reagents and solvents were analytical grade and used without further purification. Thin-layer chromatography (TLC) analyses were carried out on silica gel GF (Analtech) glass plates (2.5 cm × 10 cm with 250 μM layer and pre-scored) and visualized by UV light (254 nm). Flash column chromatography was performed on 32-64 μM silica gel obtained from EM Science, Gibbstown, New Jersey. Melting points were determined on an Electro thermal MEL-TEMP apparatus and are uncorrected. Nuclear magnetic resonance spectra were recorded on a Bruker ARX (300 MHz) spectrometer as deuteriochloroform (CDCl₃) solutions using tetramethylsilane (TMS) as an internal standard (δ=0) unless noted otherwise. IR spectra were recorded on an Avatar spectrophotometer 370 FT-IR.



[0087] *3,3-Ethylenedioxy-5α-hydroxy-11β-[4'-cyclopropylphenyl]-estr-9-ene-17-one (3a)*

A three neck dried flask was charged with Mg turnings (674 mg, 28.1 mmol), a crystal of I₂ was added and swirled over the Mg and kept for 5 min. 30 mL of anhydrous THF was added followed by 1 mL of 1,2-dibromoethane. The reaction was slightly warmed with a heat gun. When Mg starts reacting, a solution of 4-bromocyclopropylbenzene (5.36 g, 27.2 mmol) in


WO 2016/154203

PCT/US2016/023582

50 mL of THF was added dropwise over a period of 10 min. The reaction was stirred under reflux for 1h. Afterward, the reaction was cooled to room temperature and CuCl (816 mg, 8.16 mmol) was added. The reaction was stirred for 30 min and then a solution of the epoxide **2** (2 g, 9.07 mmol) in THF (30 mL) was added dropwise and stirred for 1h. The TLC showed a more polar product with respect to the epoxide. The reaction was cooled and quenched by the addition of sat. solution of NH₄Cl and extracted with ethyl acetate (3x100mL); the organic layers were washed with water and brine and the solvent was removed under vacuum. The crude was purified by column chromatography using 50% of ethyl acetate in hexane to get 3.6 g of a white foam (89% yield). mp 181-182 °C. UV(nm): 200, 231. R_f: 0.15 (5:5, Hex:EtOAc). FT IR (ATR, cm⁻¹): 3521, 2927, 1735, 1512.

¹H NMR (CDCl₃, 300 MHz) δ 0.48 (s, 3H, H-18), 0.64 (m, 2H, cyclopropyl), 0.91 (m, 2H, cyclopropyl), 3.9 (m, 4H, ketal), 4.28 (d, *J*= 6.3 Hz, H-11), 4.3 (s, 1 H, -OH), 6.9 (d, *J*= 8.1 Hz, 2H, H-Ar), 7.1 (d, *J*= 8.1 Hz, 2H, H-Ar).

¹³C NMR (CDCl₃, 75 MHz) δ 9.10 (cyclopropyl), 9.21 (cyclopropyl), 14.28 (C-18), 14.87 (cyclopropyl), 64.02 (ketal), 64.64 (ketal), 69.98 (C-5), 108.64 (C-3), 125.50 (C-Ar), 126.94 (C-Ar), 133.66 (C-10), 135.02 (C-9), 141.10 (C-Ar), 143.04 (C-Ar), 219.93 (C-17).



[0088] 3,3-Ethylendioxy-11β-[4-p-(cyclopropyl)phenyl]estra-4,9-dien-17-one (4a)


To a solution of compound **3a** (3 g, 6.68 mmol) in pyridine (30 mL), acetic anhydride (3.2 mL, 33.4 mmol) and DMAP (81.6 mg, 0.66 mmol) were added and the mixture was heated at 65 °C for 36 h. TLC showed a less polar product compared to the starting material. Solvents were distilled off under high vacuum. The purification of this compound was done by column chromatography using basic alumina with a mixture of 20% ethyl acetate in hexane to get 2.41 g of a white powder of **4a** (67 % yield), mp 182-186 °C. R_f: 0.66 (5:5, Hex:EtOAc). UV(nm): 200,250. FT IR (ATR, cm⁻¹): 2959, 2921, 1737, 1624.

WO 2016/154203

PCT/US2016/023582

^1H NMR (CDCl_3 , 300 MHz) δ 0.50 (s, 3H, H-18), 0.64 (m, 2H, cyclopropyl), 0.91 (m, 2H, cyclopropyl), 3.9 (m, 4H, ketal), 4.26 (d, $J= 6.3$ Hz, H-11), 5.38 (s, 1 H, H-4), 6.9 (d, $J= 8.1$ Hz, 2H, H-Ar), 7.1 (d, $J= 8.1$ Hz, 2H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 9.13 (cyclopropyl), 9.21 (cyclopropyl), 14.42 (C-18), 14.88 (cyclopropyl), 64.39 (ketal), 64.53 (ketal), 106.13 (C-3), 121.7 (C-4), 125.61 (C-Ar), 126.99 (C-Ar), 130.10 (C-10), 137.63 (C-9), 139.33 (C-5), 141.16 (C-Ar), 141.75 (C-Ar), 219.69 (C-17).



6a

10 [0089] *17 α -[1,1-difluoro-3-[tris(1-methylethyl)silyl]-2-propyn-1-yl]-17 β -hydroxy-11 β -[4-p-(cyclopropyl)phenyl]estra-4,9-dien-3-one. (6a)*


2.3 g of the steroid **4a** (5.3 mmol) and 3.92 g (12.6 mmol) of the 3-bromo-3,3-difluoro-1-triisopropylsilylpropyne were dissolved in THF (230 mL) and cooled to -78 °C. After this, 5 mL (12.6 mmol) of 2.5 M solution of n-BuLi was added dropwise and the reaction was stirred for 1h in which time the TLC showed a less polar product compared to the starting material. The reaction was quenched by adding sat solution of NH_4Cl and extracted with ethyl acetate and the organic layer was washed with water and brine. The solvent was removed under vacuum to afford crude **5a** which was dissolved in 75 mL of methanol and 75 mL of THF. 3.7 mL (15 mmol) of a 4N solution of HCl was added dropwise and stirred for 1 h. The TLC showed completion of the reaction. The reaction was quenched by adding a sat solution of sodium bicarbonate and extracted with ethyl acetate (3x100mL). The combined organic layers were washed with water and brine and the solvent was removed under vacuum. The crude was purified by column chromatography using 30% of ethyl acetate in hexane to get a beige powder of **6a** (quantitative yield). mp $99-101$ °C. R_f : 0.45 (7:3, Hex:EtOAc). UV (nm): 200, 230, 299. FT IR (ATR, cm^{-1}): 3412, 2940, 2873, 1651, 1591.

^1H NMR (CDCl_3 , 300 MHz) δ 0.60 (s, 3H, H-18), 0.64 (m, 2H, cyclopropyl), 0.91 (m, 2H, cyclopropyl), 1.03 (s, 3H, $\text{Si}(\text{CH}_3)_3(\text{CH}_3)_6$), 1.08 (s, 18H, $\text{Si}(\text{CH}_3)_3(\text{CH}_3)_6$), 4.3 (s, H-11), 5.76 (s, 1 H, H-4), 6.9 (d, $J= 8.4$ Hz, 2H, H-Ar), 7.04 (d, $J= 8.4$ Hz, 2H, H-Ar).

WO 2016/154203

PCT/US2016/023582

^{13}C NMR (CDCl_3 , 75 MHz) δ 9.17 (cyclopropyl), 9.26 (cyclopropyl), 10.92 (C-18), 14.51 (cyclopropyl), 18.51 ($\text{Si}(\text{CH}_3)_3(\underline{\text{C}}\text{H}_3)_6$), 24.40 ($\text{Si}(\underline{\text{C}}\text{H})_3(\text{CH}_3)_6$), 86.63 (t, $J=24$ Hz, ($\underline{\text{C}}\text{F}_2\text{CC}$)), 122.90 (C-4), 125.74 (C-Ar), 126.66 (C-Ar), 129.46 (C-10), 141.29 (C-Ar), 141.39 (C-Ar), 145.24 (C-9), 156.50 (C-5), 199.45 (C-3).



EC330

5


[0090] *17 α -(1,1-difluoro-2-propyn-1-yl)-17 β -hydroxy-11 β -[4-p-(cyclopropyl)phenyl]estra-4,9-dien-3-one (EC330)*

To a solution of compound **6a** (2g, 3.3 mmol) in 200 mL of THF was added a 1 M solution of TBAF (6.7 mL) and the mixture was stirred for 30 min. TLC showed a more polar product. The reaction was quenched by adding water and extracted with ethyl acetate (3x100mL). The combined organic layers were washed with water and brine and the solvent was removed under vacuum. The crude was purified by column chromatography using 50% of ethyl acetate in hexane to get 1.08g of a white powder of EC330. (71 % yield), mp 187-188 °C. R_f : 0.26 (7:3, Hex:EtOAc). UV(nm): 200, 230, 303. FT IR (ATR, cm^{-1}): 3284, 3217, 2947, 2124, 1645, 1604.

^1H NMR (CDCl_3 , 300 MHz) δ 0.61 (s, 3H, H-18), 0.65 (m, 2H, cyclopropyl), 0.93 (m, 2H, cyclopropyl), 2.90 (t, $J=5.4$ Hz, 1H, acetylenic hydrogen), 4.3 (d, $J=7.2$ Hz, 1H, H-11), 5.76 (s, 1 H, H-4), 6.9 (d, $J=8.4$ Hz, 2H, H-Ar), 7.06 (d, $J=8.4$ Hz, 2H, H-Ar).


^{13}C NMR (CDCl_3 , 75 MHz) δ 9.17 (cyclopropyl), 9.26 (cyclopropyl), 14.89 (C-18), 16.61 (cyclopropyl), 86.17 (t, $J=24$ Hz, ($\underline{\text{C}}\text{F}_2\text{CC}$)), 123.01 (C-4), 125.76 (C-Ar), 126.67 (C-Ar), 129.56 (C-10), 141.18 (C-Ar), 141.44 (C-Ar), 145.05 (C-9), 156.34 (C-5), 199.46 (C-3).

20



[0091] *17α-[1,1-difluoro-3-[tris(1-methylethyl)silyl]-2-propyn-1-yl]-17β-hydroxy-11β-[4-p-(acetyl)phenyl]estra-4,9-dien-3-one. (6b)*

- 5 Following the procedure described for compound **6a**, 0.77 g of compound **4b** (1.61 mmol) was reacted with 2.5 g of 3-bromo-3,3-difluoro-1-triisopropylsilylpropyne (8.1 mmol) and 3.1 mL of 2.5M solution of n-BuLi to afford crude **5b** which was hydrolysed by 1.4 ml 4N hydrochloric acid to afford 440 mg of **6b** as an amorphous solid in 60% yield. UV(nm): 200, 230, 290 FT IR (ATR, cm^{-1}): 3284, 3217, 2947, 2124, 1732, 1645, 1604.
- 10 ^1H NMR (CDCl_3 , 300 MHz) δ 0.58 (s, 3H, H-18), 1.11 (s, 18H, $\text{Si}(\text{CH}_3)_3$), 2.73 (s, 3H), 4.45 (m, H-11), 5.79 (s, 1 H, H-4), , 7.28 (d, $J= 8.3$ Hz, 2H, H-Ar), 7.88 (d, $J= 8.3$ Hz, 2H, H-Ar).



- 15 [0092] *17α-(1,1-difluoro-2-propyn-1-yl)-17β-hydroxy-11β-[4-p-(acetyl)phenyl]estra-4,9-dien-3-one (EC332)*

To a solution of compound **6b** (750mg, 1.03 mmol) in 50 mL of methanol was cooled to 0°C as 0.7 ml of 4N HCl was added dropwise. The reaction was stirred for an hour during which time TLC showed complete conversion of the starting material to the product. Reaction was

20 quenched by the addition of sat. sodium bicarbonate and extracted with EtOAc (3x100ml). The combined organic layers were washed with water and brine and the solvent was removed under vacuum to get the crude which was dissolved in 50 ml of THF and was treated with a 1M solution of TBAF (1.25 mL) and the mixture was stirred for 30 min. TLC showed a more polar product. The reaction was quenched by adding water and extracted with ethyl acetate


25 (3x100mL). The combined organic layers were washed with water and brine and the solvent

WO 2016/154203

PCT/US2016/023582

was removed under vacuum. The crude was purified by column chromatography using 50% of ethyl acetate in hexane to get 400 mg of EC332 as an off white powder. (83 % yield), UV(nm): 200, 230, 303 FT IR (ATR, cm^{-1}): 3284, 3217, 2947, 2124, 1732, 1645, 1604.

^1H NMR (CDCl_3 , 300 MHz) δ 0.57 (s, 3H, H-18), 2.73 (s, 3H), 2.91 (t, $J= 5.3$ Hz, 1H, acetylenic hydrogen), 4.1 (d, $J= 7.2$ Hz, 1H, H-11), 5.79 (s, 1 H, H-4), 7.28 (d, $J= 8.3$ Hz, 2H, H-Ar), 7.87 (d, $J= 8.3$ Hz, 2H, H-Ar).




[0093] *3,3-Ethylenedioxy-17α-[1,1-difluoro-3-[tris(1-methylethyl)silyl]-2-propyn-1-yl]-5α-hydroxy-11β-[4-p-(1,3-imidazolyl)phenyl]estra-9-ene(5c)*

Following the procedure described for compound **6a**, 1g of compound **4c** (2.1 mmol) was reacted with 3.3 g of 3-bromo-3,3-difluoro-1-triisopropylsilylpropyne (10.5 mmol) and 4.3 mL of 2.5M solution of n-BuLi to afford 300 mg of **5c** as a brown oil, yield: 21%. R_f : 0.61 (7:3, EtAc:Acetone). UV (nm): 200, 244. FT IR (ATR, cm^{-1}): 3517, 2938, 1519, 1456.

^1H NMR (CDCl_3 , 300 MHz) δ 0.56 (s, 3H, H-18), 1.01 (s, 3H, $\text{Si}(\text{CH}_3)_3(\text{CH}_3)_6$), 1.11 (s, 18 H, $\text{Si}(\text{CH}_3)_3(\text{CH}_3)_6$), 3.8 (s, 1H, -OH), 3.9 (m, 4H, ketal), 4.2 (s, 1 H, -OH), 4.3 (d, $J= 6.3$ Hz, H-11), 7.3 (m, 6H, H-Ar), 7.83 (s, 1H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 16.48 (C-18), 18.50 ($\text{Si}(\text{CH}_3)_3(\text{CH}_3)_6$), 23.25 ($\text{Si}(\text{CH}_3)_3(\text{CH}_3)_6$), 64.04 (ketal), 64.66 (ketal), 69.78 (C-5), 86.5 (t, $J= 24$ Hz, (CF_2CC)), 108.51 (C-3), 121.11 (C-Ar), 128.50 (C-Ar), 132.61 (C-10), 134.82 (C-Ar), 135.18 (C-9), 146.71 (C-Ar).




[0094] *3,3-Ethylendioxy-17 α -[1,1-difluoro-3-[tris(1-methylethyl)silyl]-2-propyn-1-yl]-5 α -hydroxy-11 β -[4-*p*-(1,3-imidazolyl)phenyl]estra-2,9-dien-3-one. (6c)*

225 mg of compound **5c** was dissolved in 5 mL of methanol and cooled to -10 °C. A 4N solution of HCl (0.32 mL, 1.3 mmol) was added dropwise. The mixture was stirred at room temperature for 1 hr. The reaction was quenched by the careful addition of saturated sodium bicarbonate solution and extracted with ethyl acetate. Combined organic layers were washed with water, brine and dried over anhydrous sodium sulfate. The crude was purified by column chromatography to get 210 mg of **6c** as a yellow powder (quantitative yield), mp 127-129 °C. R_f : 0.43 (5:5, EtAc:Acetone). UV (nm): 200, 244. FT IR (ATR, cm^{-1}): 3116, 2947, 2852, 1651.

^1H NMR (CDCl_3 , 300 MHz) δ 0.57 (s, 3H, H-18), 1.02 (s, 3H, $\text{Si}(\text{CH}_3)_3(\text{CH}_3)_6$), 1.09 (s, 18H, $\text{Si}(\text{CH}_3)_3(\text{CH}_3)_6$), 4.45 (d, $J=5.7$ Hz, H-11), 5.7 (s, 1H, H-5), 7.27 (m, 6H, H-Ar), 7.82 (s, 1H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 16.63 (C-18), 18.87 (Si(CH)₃(CH₃)₆), 24.46 (Si(CH)₃(CH₃)₆), 60.36 (CF₂CC), 76.3 (CF₂CC), 86.22 (t, $J=24$ Hz, (CF₂CC)), 118.17 (C-17), 121.54 (C-Ar), 123.35 (C-4), 128.30 (C-Ar), 130.04 (C-10), 130.17 (C-Ar), 135.10 (C-Ar), 135.44 (C-Ar), 143.98 (C-9), 144.26 (C-Ar), 155.96 (C-5), 199.05 (C-3).



EC351

20

[0095] *3,3-Ethylendioxy-17 α -(1,1-difluoro-2-propyn-1-yl)-5 α -hydroxy-11 β -[4-*p*-(1,3-imidazolyl)phenyl]estra-2,9-dien-3-one (EC351)*

Following the procedure described for EC330, 180 mg of compound **6c** (0.28 mmol) was treated with 0.56 mL of a 1M solution of TBAF to obtain 70 mg of EC351 as a beige powder. (yield: 51%), mp 175-178 °C. R_f : 0.42 (5:5, EtOAc:Acetone). UV (nm): 200, 242, 302. FT IR (ATR, cm^{-1}): 3230, 2947, 2117, 1651, 1523.


25

WO 2016/154203

PCT/US2016/023582

^1H NMR (CDCl_3 , 300 MHz) δ 0.65 (s, 3H, H-18), 2.89 (t, J = 5.4 Hz, 1H, acetylenic hydrogen), 4.49 (d, J = 5.7 Hz, H-11), 5.8 (s, 1H, H-5), 7.27 (m, 6H, H-Ar), 7.82 (s, 1H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 16.83 (C-18), 59.1 (CF_2CC), 76.16 (CF_2CC), 85.73 (t, J = 24 Hz, (CF_2CC)), 118.1 (C-17), 121.57 (C-Ar), 123.44 (C-4), 128.34 (C-Ar), 130.14 (C-10), 130.28 (C-Ar), 135.14 (C-Ar), 143.9 (C-9), 144.25 (C-Ar), 155.92 (C-5), 199.13 (C-3).

**3k**

[0096] *3,3-Ethylenedioxy-5 α -hidroxy-11 β -[4-p-(2,4,6-trimethylphenyl)phenyl]estra-9-en-17-one (3k)*

Compound **3c** (5.8 g, 10.8 mmol), 2,4,6-trimethylphenylboronic acid (2.6 g, 16.2 mmol), Pd(dppfCl₂) (394 mg, 0.54 mmol) and potassium carbonate (2.2 g, 16.2 mmol) were introduced in a flask fitted with a condenser and the system was connected to nitrogen-vacuum inlet; dioxane (120 mL) and water (12 mL) were added and the flask was evacuated and backfilled with nitrogen 7-10 times. The flask was immersed in a pre-heated oil bath at 100 °C and refluxed overnight. The TLC showed complete conversion of the starting material to the product. The reaction was cooled in an ice bath and water was added, the reaction was extracted with ethyl acetate and the organic layer was washed with water and brine and dried over sodium sulfate. The crude was purified by column chromatography using 50% ethyl acetate in hexane to get 4.61 g of **3k** as a white powder (81% yield), mp 126-128 °C. R_f : 0.37 (5:5, Hex:EtOAc). UV (nm): 204. FT IR (ATR, cm^{-1}): 3520, 2920, 2866, 1732, 1618.


^1H NMR (CDCl_3 , 300 MHz) δ 0.53 (s, 3H, H-18), 1.95 (s, 6H, Ar- CH_3), 2.32 (s, 3H, Ar- CH_3), 3.9 (m, 4H, ketal), 4.39 (s, 1H, H-11), 6.92 (s, 2H, H-Ar), 7.01 (d, J = 8.1 Hz, 2H, H-Ar), 7.26 (d, J = 5.4 Hz, 2H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 13.98 (C-18), 22.12 (Ar- CH_3), 23.36 (Ar- CH_3), 23.39 (Ar- CH_3), 64.07 (ketal), 64.65 (ketal), 70.02 (C-5), 108.68 (C-3), 127.17 (C-Ar), 127.92 (C-Ar),

WO 2016/154203

PCT/US2016/023582

127.96 (C-Ar), 129.15 (C-Ar), 133.65 (C-10), 135.23(C-Ar), 135.99 (C-Ar), 136.42 (C-Ar),
138.32 (C-Ar), 138.74 (C-Ar), 144.49 (C-9), 219.85 (C-17).




5

[0097] *3,3-Ethylenedioxy-11β-[4-p-(2,4,6-trimethylphenyl)phenyl]estra-4,9-dien-17-one*
(4k)

The synthesis of compound **4k** was done following the procedure described for the synthesis of **4a** where 4.96 g of **3k** (9.4 mmol) was treated with 5 mL of acetic anhydride (47 mmol),
10 114.8 mg of DMAP (0.94 mmol) and 50 mL of pyridine, to get 4g of **4k** as a white powder (84% yield), mp 201-203 °C. R_f : 0.57 (5:5, Hex:EtOAc). UV (nm): 200, 249. FT IR (ATR, cm^{-1}): 2920, 2852, 1739, 1631.

1H NMR ($CDCl_3$, 300 MHz) δ 0.54 (s, 3H, H-18), 1.95 (s, 6H, Ar- \underline{CH}_3), 2.32 (s, 3H, Ar- \underline{CH}_3), 3.9 (m, 4H, ketal), 4.42 (d, $J=6.9$ Hz, 1H, H-11), 5.40 (s, 1H, H-4), 6.92 (s, 2H, H-
15 Ar), 7.01 (d, $J=8.1$ Hz, 2H, H-Ar), 7.26 (d, $J=5.4$ Hz, 2H, H-Ar).

^{13}C NMR ($CDCl_3$, 75 MHz) δ 14.17 (C-18), 21.00 (Ar- \underline{CH}_3), 24.32 (Ar- \underline{CH}_3), 25.79 (Ar- \underline{CH}_3), 64.07 (ketal), 64.65 (ketal), 107.42 (C-3), 122.98 (C-4), 126.5 (C-Ar), 126.7 (C-Ar), 128.94 (C-Ar), 129.58 (C-10), 136.82 (C-Ar), 137.15 (C-Ar), 142.85 (C-Ar), 144.95 (C-Ar), 145.0 (C-Ar), 145.13 (C-9), 199.36 (C-17).




20

[0098] *17 α -[1,1-difluoro-3-[tris(1-methylethyl)silyl]-2-propyn-1-yl]-17 β -hydroxy-11 β -[4-p-(2,4,6-trimethylphenyl)phenyl]estra-4,9-dien-3-one (6k)*

This reaction was done following the same procedure described for the synthesis of compound **6a** where 3.9 g of compound **6k** (7.6 mmol) was treated with 9.4 g of 3-bromo-3,3-difluoro-1-triisopropylsilylpropyne (30.4 mmol), and 15.2 mL of a 2M solution of n-BuLi (38 mmol). The crude **5k** obtained was hydrolyzed with 5.7 mL (22.8 mmol) of a 4N solution of HCl to afford 4.1 g of **6k** as a beige powder, yield: 79%, mp 112-115 °C. R_f : 0.77 (5:5, Hex:EtOAc). UV (nm): 200, 231, 312. FT IR (ATR, cm^{-1}): 3385, 2947, 2852, 1658, 1597.

^1H NMR (CDCl_3 , 300 MHz) δ 0.65 (s, 3H, H-18), 1.04 (s, 3H, $\text{Si}(\text{CH}_2)_3(\text{CH}_3)_6$), 1.05 (s, 18H, $\text{Si}(\text{CH}_2)_3(\text{CH}_3)_6$), 1.96 (s, 6H, Ar- CH_3), 2.32 (s, 3H, Ar- CH_3), 4.48 (s, 1H, H-11), 5.78 (s, 1H, H-4), 6.92 (s, 2H, H-Ar), 7.01 (d, $J=8.1$ Hz, 2H, H-Ar), 7.2 (d, $J=7.8$ Hz, 2H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 10.94 (C-18), 18.53 ($\text{Si}(\text{CH}_2)_3(\text{CH}_3)_6$), 20.99 (Ar- CH_3), 21.32 (- CCH_2CH_3), 24.45 ($\text{Si}(\text{CH}_2)_3(\text{CH}_3)_6$), 25.79 (Ar- CH_3), 27.72 (Ar- CH_3), 86.4 (t, $J=23$ Hz, (CF_2CC)), 122.98 (C-4), 126.87 (C-Ar), 128.0 (C-Ar), 129.40 (C-Ar), 129.61 (C-10), 135.98 (C-Ar), 136.5 (C-Ar), 138.48 (C-Ar), 138.63 (C-Ar), 142.63 (C-Ar), 145.21 (C-9), 156.54 (C-5), 199.46 (C-3).



EC359

20

[0099] *17 α -(1,1-difluoro-2-propyn-1-yl)-17 β -hydroxy-11 β -[4-p-(2,4,6-trimethylphenyl)phenyl]estra-4,9-dien-3-one (EC359)*

Following the same procedure reported for **EC330**, 4.1 g of compound **6k** (5.8 mmol) was treated with 8.7 mL of a 1M solution of TBAF to get 1.62 g of **EC359** as a white powder, yield: 52%, mp 255-256 °C. R_f : 0.33 (5:5, Hex:EtOAc). UV (nm): 300. FT IR (ATR, cm^{-1}): 3305, 2947, 2873, 2130, 1638.


25

WO 2016/154203

PCT/US2016/023582

^1H NMR (CDCl_3 , 300 MHz) δ 0.65 (s, 3H, H-18), 1.96 (s, 6H, Ar- CH_3), 2.32 (s, 3H, Ar- CH_3), 2.9 (t, $J=5.4$ Hz, 1H, acetylenic hydrogen), 4.52 (d, $J=6.9$ Hz, 1H, H-11), 5.78 (s, 1H, H-4), 6.93 (s, 2H, H-Ar), 7.05 (d, $J=8.4$ Hz, 2H, H-Ar), 7.23 (d, $J=8.1$ Hz, 2H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 16.40 (C-18), 20.61 (Ar- CH_3), 25.84 (Ar- CH_3), 27.69 (Ar- CH_3), 85.92 (t, $J=23$ Hz, (CF_2CC)), 123.08 (C-4), 126.87 (C-Ar), 128.0 (C-Ar), 129.43 (C-Ar), 129.70 (C-10), 135.99 (C-Ar), 136.54 (C-Ar), 138.53 (C-Ar), 138.61 (C-Ar), 142.51 (C-Ar), 145.04 (C-9), 156.45 (C-5), 199.52 (C-3).




10 [00100] *3,3-Ethylenedioxy-5 α -hidroxy-11 β -[4-p-(dimethylamino)phenyl]estra-9-en-17-one (3d)*

Following the procedure described for compound **3a**, 4g of compound **2** (12.1 mmol) was reacted with 1.75g of Mg (72.6 mmol), 12g of 4-Br-N,N-dimethylanilin (60.4 mmol) and 600 mg of CuCl (6.04 mmol) to afford **3d** as a white powder (5g), yield: 93%, mp 116-118 °C. R_f : 0.28 (5:5, Hex:EtOAc). UV (nm): 201, 259, 302. FT IR (ATR, cm^{-1}): 3507, 2927, 2873, 1749, 1604.

^1H NMR (CDCl_3 , 300 MHz) δ 0.51 (s, 3H, H-18), 2.90 (s, 6H, $-\text{N}(\text{CH}_3)_2$), 3.9 (m, 4H, ketal), 4.2 (d, $J=6.3$ Hz, H-11), 4.3 (s, 1 H, $-\text{OH}$), 6.65 (d, $J=8.7$ Hz, 2H, H-Ar), 7.07 (d, $J=8.4$ Hz, 2H, H-Ar).

20 ^{13}C NMR (CDCl_3 , 75 MHz) δ 14.26 (C-18), 64.02 (ketal), 64.63 (ketal), 70.03 (C-5), 108.73 (C-3), 112.60 (C-Ar), 127.61 (C-Ar), 133.57 (C-10), 134.08 (C-Ar), 134.65 (C-9), 148.44 (C-Ar), 220.17 (C-17).




[00101] **3,3-Ethylendioxy-11 β -[4-*p*-(dimethylamino)phenyl]estra-4,9-dien-17-one (4d)**

The synthesis of compound **4d** was done following the procedure described for compound **4a** where 4.55g of **3d** (10 mmol) was reacted with 4.8 mL of acetic anhydride (50 mmol), 122.1 mg of DMAP (1 mmol) and 30 mL of pyridine, to get 2.76 g of a beige powder of **4d** (64% yield), mp 125-127 °C. R_f : 0.6 (5:5, Hex:EtOAc). UV (nm): 202,255. FT IR (ATR, cm^{-1}): 2906, 1739, 1620.

^1H NMR (CDCl_3 , 300 MHz) δ 0.53 (s, 3H, H-18), 2.93 (s, 6H, $-\text{N}(\text{CH}_3)_2$), 3.9 (m, 4H, ketal), 4.2 (d, $J=6.3$ Hz, 1H, H-11), 6.37 (s, 1H, H-4), 6.64 (d, $J=8.7$ Hz, 2H, H-Ar), 7.07 (d, $J=8.4$ Hz, 2H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 14.16 (C-18), 64.34 (ketal), 64.51 (ketal), 106.20 (C-3), 112.60 (C-Ar), 121.52 (C-4), 127.61 (C-Ar), 129.79 (C-Ar), 132.40 (C-10), 138.15 (C-9), 139.49 (C-5), 148.49 (C-Ar), 219.95 (C-17).



[00102] **17 α -[1,1-difluoro-3-[tris(1-methylethyl)silyl]-2-propyn-1-yl]-17 β -hydroxy-11 β -[4-*p*-(dimethylamino)phenyl]estra-4,9-dien-3-one (6d)**


Following the same procedure described for the synthesis of compound **5a**, 0.5 g of compound **4d** (1.15 mmol) was treated with 1.7g of 3-bromo-3,3-difluoro-1-triisopropylsilylpropyne (5.75 mmol), and 2.3 mL (5.75 mmol) of a 2M solution of n-BuLi. The crude obtained was hydrolysed with 0.3 mL (1.5 mmol) of a 4N solution of HCl to afford **6d** as a brown powder, yield: 63%, mp 113-116 °C. R_f : 0.52 (7:3, Hex:EtOAc). FT IR (ATR, cm^{-1}): 3399, 2947, 2866, 1658, 1608.

^1H NMR (CDCl_3 , 300 MHz) δ 0.64 (s, 3H, H-18), 1.03 (s, 3H, $\text{Si}(\text{CH}_3)_3(\text{CH}_3)_6$), 1.08 (s, 18H, $\text{Si}(\text{CH}_3)_3(\text{CH}_3)_6$), 2.91 (s, 6H, $-\text{N}(\text{CH}_3)_2$), 4.34 (s, 1H, H-11), 5.75 (s, 1H, H-4), 6.66 (d, $J=9$ Hz, 2H, H-Ar), 7.01 (d, $J=8.7$ Hz, 2H, H-Ar).

WO 2016/154203

PCT/US2016/023582

^{13}C NMR (CDCl_3 , 75 MHz) δ 16.38 (C-18), 18.51 ($\text{Si}(\text{CH}_3)_3(\underline{\text{C}}\text{H}_3)_6$), 24.40 ($\text{Si}(\underline{\text{C}}\text{H})_3(\text{CH}_3)_6$), 86.30 (t, $J=21$ Hz, ($\underline{\text{C}}\text{F}_2\text{CC}$)), 102.20 (C-17), 112.76 (C-Ar), 122.75 (C-4), 127.43 (C-Ar), 129.22 (C-Ar), 131.89 (C-10), 145.84 (C-9), 148.55 (C-Ar), 156.63 (C-5), 199.54 (C-3).



EC352


5

[00103] *17 α -(1,1-difluoro-2-propyn-1-yl)-17 β -hydroxy-11 β -[4-*p*-(dimethylamino)phenyl]estra-4,9-dien-3-one (EC352)*

The reaction was done following the same procedure described for EC330 where 80mg of compound **5d** (0.128 mmol) was treated with 0.15 mL of a 1M solution of TBAF to get 60 mg of EC352 as a yellow powder (yield 99%), mp 138-140 °C. R_f : 0.15 (7:3, Hex:EtOAc). UV (nm): 203, 259, 301. FT IR (ATR, cm^{-1}): 3999, 3284, 2130, 1645, 1604.

^1H NMR (CDCl_3 , 300 MHz) δ 0.64 (s, 3H, H-18), 2.89 (t, $J=5.4$ Hz, 1H, acetylenic hydrogen), 2.91 (s, 6H, $-\text{N}(\text{CH}_3)_2$), 4.37 (d, $J=6.3$ Hz, 1H, H-11), 5.29 (s, 1H, H-4), 5.75 (s, 1H, H-4), 6.67 (d, $J=8.7$ Hz, 2H, H-Ar), 7.02 (d, $J=8.7$ Hz, 2H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 16.53 (C-18), 85.86 (t, $J=24$ Hz, ($\underline{\text{C}}\text{F}_2\text{CC}$)), 112.78 (C-Ar), 122.81 (C-4), 127.37 (C-Ar), 129.29 (C-Ar), 131.86 (C-10), 145.78 (C-9), 148.55 (C-Ar), 156.64 (C-5), 199.63 (C-3).



3e

20 [00104] *3,3-Ethylenedioxy-5 α -hidroxy-11 β -[4-*p*-(methoxy)phenyl]estra-9-en-17-one (3e)*

This compound was synthesized following the procedure described for compound **3a**, where 3g of compound **2** (9 mmol) was reacted with 1.3g of Mg (54 mmol), 8.4 g of 4-Br-anisole (45 mmol) and 445 mg of CuCl (4.5 mmol) to afford **3e** as a white powder (3.81g), yield:


WO 2016/154203

PCT/US2016/023582

97%, mp 104-108 °C. R_f: 0.22 (5:5, Hex:EtOAc). UV (nm): 200,230. FT IR (ATR, cm⁻¹): 3507, 2927, 2879, 1739, 1611.

¹H NMR (CDCl₃, 300 MHz) δ 0.49 (s, 3H, H-18), 3.77 (s, 3H, -OCH₃), 3.9 (m, 4H, ketal), 4.28 (d, *J*= 6.9 Hz, H-11), 4.37 (s, 1 H, -OH), 6.8 (d, *J*= 9 Hz, 2H, H-Ar), 7.1 (d, *J*= 8.7 Hz, 2H, H-Ar).

¹³C NMR (CDCl₃, 75 MHz) δ 14.37 (C-18), 64.0 (ketal), 64.6 (ketal), 70.1 (C-5), 112.3 (C-3), 114.01 (C-Ar), 130.02 (C-Ar), 133.5 (C-10), 135.76 (C-Ar), 134.6 (C-9), 157.71 (C-Ar), 218.99 (C-17).



10


[00105] ***3,3-Ethylenedioxy-11β-[4-p-(methoxy)phenyl]estra-4,9-dien-17-one (4e)***

The synthesis of compound **4e** was done following the procedure described for compound **4a** where 3.7 g of **3e** (8.4 mmol) was heated at 70°C with 4.8 mL of acetic anhydride (45.5 mmol), 111.2 mg of DMAP (0.91 mmol) and 40 mL of pyridine, to get 1.6 g of **4e** as a white powder (45% yield), mp 107-110 °C. R_f: 0.63 (5:5, Hex:EtOAc). UV (nm): 200, 250. FT IR (ATR, cm⁻¹): 2940, 2866, 1732, 1604.

¹H NMR (CDCl₃, 300 MHz) δ 0.51 (s, 3H, H-18), 3.76 (s, 3H, -OCH₃), 3.9 (m, 4H, ketal), 4.12 (d, *J*= 6.9 Hz, H-11), 5.38 (s, 1H, H-4), 6.77 (d, *J*= 8.7 Hz, 2H, H-Ar), 7.13 (d, *J*= 8.7 Hz, 2H, H-Ar).

¹³C NMR (CDCl₃, 75 MHz) δ 14.37 (C-18), 55.17 (-OCH₃), 64.0 (ketal), 64.6 (ketal), 114.02 (C-3), 123.34 (C-4), 130.02 (C-10), 135.76 (C-Ar), 144.99 (C-9), 156.01 (C-Ar), 157.71 (C-Ar), 218.96 (C-17).

20




[00106] *17α-[1,1-difluoro-3-[tris(1-methylethyl)silyl]-2-propyn-1-yl]-17β-hydroxy-11β-[4-p-(methoxy)phenyl]estra-4,9-dien-3-one (6e)*

5 Following the same procedure described for **5a**, 1.4 g of compound **4e** (3.3 mmol) was reacted with 4.1g of 3-bromo-3,3-difluoro-1-triisopropylsilylpropyne (13.2 mmol), and 6.6 mL of a 2M solution of n-BuLi (16.5 mmol). The crude obtained was hydrolysed with 2.6 mL (10.6 mmol) of a 4N solution of HCl to get 1.25g of **6e** as a brown powder, yield: 98%, mp 99-102 °C. R_f : 0.18 (9:1, Hex:EtOAc). UV (nm): 229, 300. FT IR (ATR, cm^{-1}): 3412, 2940, 2866, 1665, 1618.

^1H NMR (CDCl_3 , 300 MHz) δ 0.60 (s, 3H, H-18), 1.03 (s, 3H, $\text{Si}(\text{CH}_3)_3$), 1.06 (s, 18H, $\text{Si}(\text{CH}_3)_3$), 3.77 (s, 3H, $-\text{OCH}_3$), 4.36 (d, $J=6.9$ Hz, H-11), 5.75 (s, 1H, H-4), 6.80 (d, $J=8.7$ Hz, 2H, H-Ar), 7.06 (d, $J=8.7$ Hz, 2H, H-Ar).

15 ^{13}C NMR (CDCl_3 , 75 MHz) δ 10.91 (C-18), 18.51 ($\text{Si}(\text{CH}_3)_3$), 24.41 ($\text{Si}(\text{CH}_3)_3$), 55.1 ($-\text{OCH}_3$), 86.32 (t, $J=21$ Hz, (CF_2CC)), 113.89 (C-Ar), 122.93 (C-4), 127.72 (C-Ar), 129.48 (C-10), 136.30 (C-Ar), 145.31 (C-9), 156.47 (C-5), 157.53 (C-Ar), 199.42 (C-3).



20 **[00107]** *17α-(1,1-difluoro-2-propyn-1-yl)-17β-hydroxy-11β-[4-p-(methoxy)phenyl]estra-4,9-dien-3-one (EC356)*


The reaction was done following the same procedure reported for **EC330** where 1.2 g of compound **5e** (1.97 mmol) was treated with 3 mL of a 1M solution of TBAF to get 530 mg of **EC356** as a beige powder, yield: 60%, mp 173-174 °C. R_f : 0.33 (5:5, Hex:EtOAc). UV (nm): 228, 303. FT IR (ATR, cm^{-1}): 3325, 3271, 2940, 2130, 1638, 1597.

WO 2016/154203

PCT/US2016/023582

^1H NMR (CDCl_3 , 300 MHz) δ 0.61 (s, 3H, H-18), 2.89 (t, $J= 5.4$ Hz, 1H, acetylenic hydrogen), 3.77 (s, 3H, $-\text{OCH}_3$), 4.3 (d, $J= 7.2$ Hz, 1H, H-11), 5.76 (s, 1 H, H-4), 6.8 (d, $J= 8.4$ Hz, 2H, H-Ar), 7.06 (d, $J= 8.4$ Hz, 2H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 16.55 (C-18), 55.1 ($-\text{OCH}_3$), 85.85 (t, $J= 24$ Hz, (CF_2CC)),
 5 113.89 (C-Ar), 123.04 (C-4), 127.74 (C-Ar), 129.57 (C-10), 136.47 (C-Ar), 145.12 (C-9),
 156.35 (C-5), 157.56 (C-Ar), 199.44 (C-3).




[00108] *3,3-Ethylendioxy-5α-hidroxy-11β-[4-p-(t-butyl)phenyl]estra-9-en-17-one*
 10 (3f)

1-tert-butyl-4-iodobenzene (3g, 11.53 mmol) was dissolved in 25 mL of THF and cooled to -
 10 °C as a 2M solution of isopropyl magnesium chloride (7.9 mL, 15.72 mmol) was added
 dropwise over a period of 2-3 min. The resulting solution was stirred for 30 min. Solid CuCl
 (466 mg, 4.71 mmol) was added and stirred for another 30 min at 0 °C. A solution of the
 15 epoxide **2** (2.5g, 7.56 mmol) in 20 mL of THF was added dropwise and the flask was taken
 out of the cooling bath after 15 min and allowed to stir for 1 hr at room temperature. The
 TLC showed complete conversion of starting material to product. The reaction was
 quenched by adding sat NH_4Cl , extracted with ethyl acetate and washed with water and brine.
 The organic layer was dried over sodium sulfate and evaporated under vacuum to get the
 20 crude. The crude was purified by column chromatography using 60% ethyl acetate in hexane
 to get 2.91g of **3f** as a white powder (83% yield), mp 94-96°C. UV (nm): 200,230. R_f : 0.4
 (5:5, Hex:EtOAc). FT IR (ATR, cm^{-1}): 3514, 2947, 2873, 1736.

^1H NMR (CDCl_3 , 300 MHz) δ 0.46 (s, 3H, H-18), 1.27 (s, 9H, $-\text{C}(\text{CH}_3)_3$), 3.9 (m, 4H, ketal),
 4.27 (d, $J= 6.9$ Hz, H-11), 7.13 (d, $J= 8.4$ Hz, 2H, H-Ar), 7.22 (d, $J= 8.7$ Hz, 2H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 14.4 (C-18), 63.69 (ketal), 64.0 (ketal), 108.73 (C-3), 123.25
 25 (C-Ar), 126.43 (C-Ar), 140.47 (C-10), 145.15 (C-Ar), 148.88 (C-Ar), 219.03 (C-17).




4f

[00109] *3,3-Ethylendioxy-11β-[4-p-(t-butyl)phenyl]estra-4,9-dien-17-one (4f)*

The synthesis of compound **4f** was accomplished following the procedure described **4a** where
 5 2.8 g of **3f** (6 mmol) was heated at 70 °C with 3.2 mL of acetic anhydride (30 mmol), 73.3
 mg of DMAP (0.6 mmol) and 30 mL of pyridine, to get 1.7 g of **4f** as a white powder (65%
 yield), mp 85-88 °C. R_f : 0.72 (5:5, Hex:EtOAc). UV (nm): 200, 250. FT IR (ATR, cm^{-1}):
 2947, 2873, 1732, 1631.

^1H NMR (CDCl_3 , 300 MHz) δ 0.52 (s, 3H, H-18), 1.28 (s, 9H, $-\text{C}(\text{CH}_3)_3$), 3.9 (m, 4H, ketal),
 10 4.3 (d, $J=6.9$ Hz, H-11), 5.78 (s, 1H, H-4), 7.1 (d, $J=8.4$ Hz, 2H, H-Ar), 7.28 (d, $J=8.7$ Hz,
 2H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 14.42 (C-18), 31.29 ($-\text{C}(\underline{\text{C}}\text{H}_3)_3$), 34.26 ($-\underline{\text{C}}(\text{CH}_3)_3$), 64.1 (ketal),
 64.16 (ketal), 107.95 (C-3), 122.49 (C-4), 123.26 (C-Ar), 124.67 (C-Ar), 125.49 (C-Ar),
 126.42 (C-Ar), 130.81 (C-10), 140.47 (C-5), 141.94 (C-9), 145.09 (C-Ar), 148.86 (C-Ar),
 15 218.93 (C-17).



6f

[00110] *17α-[1,1-difluoro-3-[tris(1-methylethyl)silyl]-2-propyn-1-yl]-17β-hydroxy-11β-[4-p-(t-butyl)phenyl]estra-4,9-dien-3-one (6f)*

20 This reaction was done following the same procedure described for **6a** where 1.6 g of
 compound **4f** (3.5 mmol) was reacted with 4.3 g of 3-bromo-3,3-difluoro-1-
 triisopropylsilylpropyne (14 mmol), and 7 mL of a 2M solution of n-BuLi (17.5 mmol). The
 crude obtained was hydrolysed with 2.6 mL (10.5 mmol) of a 4N solution of HCl to get 1.87g


WO 2016/154203

PCT/US2016/023582

of **6f** as a brown powder, yield: 85%, mp 89-91 °C. UV (nm): 222, 296. R_f : 0.72 (5:5, Hex:EtOAc). FT IR (ATR, cm^{-1}): 3392, 2940, 2859, 1651.

^1H NMR (CDCl_3 , 300 MHz) δ 0.59 (s, 3H, H-18), 1.08 (s, 3H, $\text{Si}(\text{CH}_3)_3$), 1.1 (s, 18H, $\text{Si}(\text{CH}_3)_3$), 1.28 (s, 9H, $-\text{C}(\text{CH}_3)_3$), 4.38 (d, $J=6.9$ Hz, H-11), 5.76 (s, 1H, H-4), 7.08 (d, $J=8.4$ Hz, 2H, H-Ar), 7.26 (d, $J=8.7$ Hz, 2H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 16.47 (C-18), 18.51 ($\text{Si}(\text{CH}_3)_3$), 24.39 ($\text{Si}(\text{CH}_3)_3$), 31.17 ($-\text{C}(\text{CH}_3)_3$), 34.25 ($-\text{C}(\text{CH}_3)_3$), 86.34 (t, $J=23$ Hz, (CF_2CC)), 122.86 (C-4), 125.36 (C-Ar), 126.35 (C-Ar), 129.42 (C-10), 140.98 (C-Ar), 145.39 (C-9), 148.56 (C-Ar), 156.55 (C-5), 199.48 (C-3).



EC357

10


[00111] *17 α -(1,1-difluoro-2-propyn-1-yl)-17 β -hydroxy-11 β -[4-*p*-(*t*-butyl)phenyl]estra-4,9-dien-3-one (EC357)*

Following the procedure reported for EC330, 1.8 g of compound **6f** (2.8 mmol) was treated with 4.5 mL of a 1M solution of TBAF to get 420 mg of **EC357** as a beige powder (yield: 32%), mp 127-129 °C. R_f : 0.43 (5:5, Hex:EtOAc). UV (nm): 222, 301. FT IR (ATR, cm^{-1}): 3406, 3271, 2947, 2866, 2130, 1651, 1597.

^1H NMR (CDCl_3 , 300 MHz) δ 0.60 (s, 3H, H-18), 1.28 (s, 9H, $-\text{C}(\text{CH}_3)_3$), 2.90 (t, $J=5.1$ Hz, 1H, acetylenic hydrogen), 4.42 (d, $J=6.9$ Hz, H-11), 5.76 (s, 1H, H-4), 7.09 (d, $J=8.1$ Hz, 2H, H-Ar), 7.26 (d, $J=7.2$ Hz, 2H, H-Ar).

20

^{13}C NMR (CDCl_3 , 75 MHz) δ 16.63 (C-18), 31.33 ($-\text{C}(\text{CH}_3)_3$), 34.26 ($-\text{C}(\text{CH}_3)_3$), 85.86 (t, $J=25$ Hz, (CF_2CC)), 122.97 (C-4), 125.38 (C-Ar), 126.38 (C-Ar), 129.52 (C-10), 140.93 (C-Ar), 145.21 (C-9), 148.62 (C-Ar), 156.46 (C-5), 199.53 (C-3).




3g

[00112] ***3,3-Ethylendioxy-5 α -hidroxy-11 β -[4-*p*-(1-methylethenyl)phenyl]estra-9-en-17-one (3g)***

Compound **3g** was synthesized following the same procedure described for compound **3a**,
 5 using 3g of compound **2** (9 mmol), 875 g of Mg (36.4 mmol), 6 g of 4-bromoisopropenylbenzene (30.4 mmol) dissolved in 20 mL of THF and 300 mg of CuCl (3 mmol). The product was white powder (3.3g), yield: 83%, mp 98-101 °C. R_f : 0.31 (5:5, Hex:EtOAc). UV (nm): 201, 254. FT IR (ATR, cm^{-1}): 3500, 2933, 2866, 1739, 1624.

^1H NMR (CDCl_3 , 300 MHz) δ 0.49 (s, 3H, H-18), 2.12 (s, 3H, $-\text{CCH}_2\text{CH}_3$), 3.9 (m, 4H, ketal), 4.3 (d, $J=7.2$ Hz, H-11), 4.37 (s, 1 H, $-\text{OH}$), 5.04 (s, 1H, $-\text{CCH}_2\text{CH}_3$), 5.37 (s, 1H, $-\text{CCH}_2\text{CH}_3$), 7.19 (d, $J=8.4$ Hz, 2H, H-Ar), 7.38 (d, $J=6.6$ Hz, 2H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 14.30 (C-18), 64.02 (ketal), 64.65 (ketal), 69.98 (C-5), 108.59 (C-3), 111.74 ($-\text{CCH}_2\text{CH}_3$), 125.32 (C-Ar), 126.91 (C-Ar), 133.43 (C-10), 135.24 (C-Ar), 138.15 (C-9), 142.52 ($-\text{CCH}_2\text{CH}_3$), 145.45 (C-Ar), 219.86 (C-17).



15

4g


[00113] ***3,3-Ethylendioxy-11 β -[4-*p*-(1-methylethenyl)phenyl]estra-4,9-dien-17-one (4g)***

The synthesis of compound **4g** was accomplished following the procedure described **4a**
 20 where 3.2 g of **4f** (7.1 mmol) was heated at 70 °C with 3.78 mL of acetic anhydride (36 mmol), 87 mg of DMAP (0.7 mmol) and 35 mL of pyridine, to get 2.4 g of **4g** as a white powder (80% yield), mp 89-91 °C. R_f : 0.57 (5:5, Hex:EtOAc). UV (nm): 203, 251. FT IR (ATR, cm^{-1}): 2927, 2879, 1739, 1631.

^1H NMR (CDCl_3 , 300 MHz) δ 0.51 (s, 3H, H-18), 2.12 (s, 3H, $-\text{CCH}_2\text{CH}_3$), 3.9 (m, 4H, ketal), 4.3 (d, $J=6.9$ Hz, H-11), 5.04 (s, 1H, $-\text{CCH}_2\text{CH}_3$), 5.36 (s, 1H, $-\text{CCH}_2\text{CH}_3$), 5.39 (s, 1H, H-4), 7.19 (d, $J=8.4$ Hz, 2H, H-Ar), 7.38 (d, $J=8.1$ Hz, 2H, H-Ar).

25

¹³C NMR (CDCl₃, 75 MHz) δ 14.44 (C-18), 21.91 (-CCH₂CH₃), 64.40 (ketal), 64.55 (ketal), 106.09 (C-3), 111.76 (-CCH₂CH₃), 121.92 (C-4), 125.42 (C-Ar), 125.47 (C-Ar), 126.97 (C-Ar), 130.28 (C-10), 137.36 (C-Ar), 138.27 (C-5), 139.25 (C-9), 142.62 (-CCH₂CH₃), 144.15 (C-Ar), 219.65 (C-17).



5


6h

[00114] *17α-[1,1-difluoro-3-[tris(1-methylethyl)silyl]-2-propyn-1-yl]-17β-hydroxy-11β-[4-p-(1-methylethenyl)phenyl]estra-4,9-dien-3-one (6h)*

This reaction was done following the same procedure described for the synthesis of compound **6a** using 2.4 g of compound **4g** (5.6 mmol), 6.9 g of 3-bromo-3,3-difluoro-1-triisopropylsilylpropyne (22.4 mmol), and 13.3 mL of a 2M solution of n-BuLi (28 mmol). The crude obtained was hydrolyzed with 1.8 mL (7.2 mmol) of a 4N solution of HCl to get 450 mg of **6e** as a brown powder, yield: 20%, mp 88-91 °C. R_f: 0.77 (5:5, Hex:EtOAc). UV (nm): 202, 252, 296. FT IR (ATR, cm⁻¹): 3392, 2940, 2852, 1665, 1597.

¹H NMR (CDCl₃, 300 MHz) δ 0.61 (s, 3H, H-18), 1.03 (s, 3H, Si(CH₃)₃(CH₃)₆), 1.06 (s, 18H, Si(CH₃)₃(CH₃)₆), 2.13 (s, 3H, -CCH₂CH₃), 4.41 (s, 1H, H-11), 5.06 (s, 1H, -CCH₂CH₃), 5.37 (d, J= 10.2 Hz, 1H, -CCH₂CH₃), 5.77 (s, 1H, H-4), 7.13 (d, J= 8.1 Hz, 2H, H-Ar), 7.40 (d, J= 8.4 Hz, 2H, H-Ar).

¹³C NMR (CDCl₃, 75 MHz) δ 10.92 (C-18), 18.51 (Si(CH₃)₃(CH₃)₆), 21.35 (-CCH₂CH₃), 24.40 (Si(CH₃)₃(CH₃)₆), 86.30 (t, J= 23.2 Hz, (CF₂CC)), 111.96 (-CCH₂CH₃), 123.02 (C-4), 125.59 (C-Ar), 126.65 (C-Ar), 129.62 (C-10), 138.48 (C-9), 142.51 (-CCH₂CH₃), 143.64 (C-Ar), 144.89 (C-Ar), 156.37 (C-5), 199.37 (C-3).




[00115] *17 α -(1,1-difluoro-2-propyn-1-yl)-17 β -hydroxy-11 β -[4-*p*-(1-methylethenyl)phenyl]estra-4,9-dien-3-one (EC358)*

The reaction was done following the same procedure described for **EC330** where 450 mg of compound **6e** (0.72 mmol) was treated with 2.18 mL of a 1M solution of TBAF to get 190 mg of **EC358** as a white powder, yield: 55%, mp 113-116 °C. R_f : 0.64 (5:5, Hex:EtOAc). UV (nm): 204, 252, 300. FT IR (ATR, cm^{-1}): 3379, 3291, 2954, 2866, 2130, 1651, 1597.

^1H NMR (CDCl_3 , 300 MHz) δ 0.62 (s, 3H, H-18), 2.9 (t, J = 5.4 Hz, 1H, acetylenic hydrogen), 4.41 (d, J = 7.2 Hz, 1H, H-11), 5.06 (s, 1H, $-\text{CCH}_2\text{CH}_3$), 5.37 (s, 1H, $-\text{CCH}_2\text{CH}_3$), 5.77 (s, 1H, H-4), 7.13 (d, J = 8.1 Hz, 2H, H-Ar), 7.40 (d, J = 8.4 Hz, 2H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 16.65 (C-18), 21.65 ($-\text{CCH}_2\text{CH}_3$), 85.83 (t, J = 24 Hz, (CF_2CC)), 111.97 ($-\text{CCH}_2\text{CH}_3$), 123.09 (C-4), 125.6 (C-Ar), 126.65 (C-Ar), 129.7 (C-10), 138.45 (C-9), 142.49 ($-\text{CCH}_2\text{CH}_3$), 143.51 (C-Ar), 144.75 (C-Ar), 156.3 (C-5), 199.41 (C-3).




3h

15 [00116] *3,3-Ethylenedioxy-5 α -hydroxy-11 β -[4-*p*-(cyclohexyl)phenyl]estra-9-en-17-one (3h)*

Compound **3h** was synthesized following the procedure described for compound **3a**, where 3g of compound **2** (9 mmol) was reacted with 875 g of Mg (36.4 mmol), 7 g of 4-bromocyclohexylbenzene (29.2 mmol) and 261 mg of CuCl (2.6 mmol) to afford **3h** as a white powder (3.46g), yield: 80%, mp 112-113 °C. R_f : 0.27 (5:5, Hex:EtOAc). UV (nm): 220. FT IR (ATR, cm^{-1}): 3507, 2920, 2859, 1732, 1442.

^1H NMR (CDCl_3 , 300 MHz) δ 0.47 (s, 3H, H-18), 3.9 (m, 4H, ketal), 4.3 (d, J = 6.9 Hz, H-11), 4.37 (s, 1 H, $-\text{OH}$), 7.08 (d, J = 8.4 Hz, 2H, H-Ar), 7.13 (d, J = 8.1 Hz, 2H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 14.25 (C-18), 64.03 (ketal), 64.65 (ketal), 70.02 (C-5), 108.69 (C-3), 126.64 (C-Ar), 126.85 (C-Ar), 133.80 (C-10), 134.95 (C-Ar), 143.1 (C-9), 145.29 (C-Ar), 219.97 (C-17).

**4h**


5

[00117] **3,3-Ethylenedioxy-11 β -[4-p-(cyclohexyl)phenyl]estra-4,9-dien-17-one (4h)**

The synthesis of compound **4h** was done following the procedure described for the synthesis of compound **4a** where 3.4 g of **3h** (6.9 mmol) was treated with 3.6 mL of acetic anhydride (34.6 mmol), 81.8 mg of DMAP (0.69 mmol) and 40 mL of pyridine, to get 2.5 g of **4h** as a white powder (78% yield), mp 110-112 °C. R_f : 0.66 (5:5, Hex:EtOAc). UV (nm): 200, 220, 300. FT IR (ATR, cm^{-1}): 2913, 2846, 1732, 1503.

^1H NMR (CDCl_3 , 300 MHz) δ 0.49 (s, 3H, H-18), 3.9 (m, 4H, ketal), 4.3 (d, $J= 6.9$ Hz, H-11), 5.38 (s, 1H, H-4), 7.07 (d, $J= 8.4$ Hz, 2H, H-Ar), 7.13 (d, $J= 8.4$ Hz, 2H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 14.39 (C-18), 64.40 (ketal), 64.54 (ketal), 106.18 (C-3), 121.65 (C-4), 126.75 (C-Ar), 126.89 (C-Ar), 130.03 (C-10), 137.81 (C-Ar), 139.38 (C-5), 139.42 (C-9), 141.82 (C-Ar), 145.32 (C-Ar), 219.65 (C-17).

**6h**

[00118] **17 α -[1,1-difluoro-3-[tris(1-methylethyl)silyl]-2-propyn-1-yl]-17 β -hydroxy-11 β -[4-p-(cyclohexyl)phenyl]estra-4,9-dien-3-one (6h)**

This reaction was done following the same procedure described for the synthesis of compound **6a** where 2.4 g of compound **4h** (4.89 mmol) was reacted with 6.0 g of 3-bromo-

WO 2016/154203


PCT/US2016/023582

3,3-difluoro-1-triisopropylsilylpropyne (19.5 mmol), and 9.78 mL of a 2M solution of n-BuLi (24.45 mmol). The crude product **5h** obtained was hydrolyzed with 3.6 mL (14.67 mmol) of a 4N solution of HCl to get 2.4 g of **6h** as a beige powder, yield: 75%, mp 99-102 °C. R_f : 0.64 (7:3, Hex:EtOAc). FT IR (ATR, cm^{-1}): 3406, 2920, 2859, 2184, 1651, 1597.

5 ^1H NMR (CDCl_3 , 300 MHz) δ 0.60 (s, 3H, H-18), 1.03 (s, 3H, $\text{Si}(\text{CH}_3)_3$), 1.1 (s, 18H, $\text{Si}(\text{CH}_3)_3$), 4.40 (d, $J=4.5$ Hz, 1H, H-11), 5.77 (s, 1H, H-4), 7.07 (d, $J=8.7$ Hz, 2H, H-Ar), 7.11 (d, $J=8.4$ Hz, 2H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 10.82 (C-18), 18.35 ($\text{Si}(\text{CH}_3)_3$), 24.40 ($\text{Si}(\text{CH}_3)_3$), 86.34 (t, $J=23.2$ Hz, (CF_2CC)), 122.88 (C-4), 126.58 (C-Ar), 126.88 (C-Ar), 129.43 (C-10),

10 141.43 (C-9), 145.38 (C-Ar), 145.51 (C-Ar), 156.55 (C-5), 199.48 (C-3).




EC360

[00119] *17 α -(1,1-difluoro-2-propyn-1-yl)-17 β -hydroxy-11 β -[4-p-(cyclohexyl)phenyl]estra-4,9-dien-3-one (EC360)*

15 The reaction was done following the same procedure as described for **EC330** where 2.3g of compound **6h** (3.5 mmol) was treated with 5.22 mL of a 1M solution of TBAF to get 1.27 g of **EC360** as a white powder, yield: 72%, mp 133-135 °C. R_f : 0.77 (5:5, Hex:EtOAc). UV (nm): 200, 223, 301. FT IR (ATR, cm^{-1}): 3406, 3284, 2927, 2839, 2124, 1645, 1591.

20 ^1H NMR (CDCl_3 , 300 MHz) δ 0.61 (s, 3H, H-18), 2.92 (t, $J=5.4$ Hz, 1H, acetylenic hydrogen), 4.42 (d, $J=6.6$ Hz, 1H, H-11), 5.77 (s, 1H, H-4), 7.09 (s, 4H, H-Ar).

^{13}C NMR (CDCl_3 , 75 MHz) δ 16.61 (C-18), 85.87 (t, $J=24.9$ Hz, (CF_2CC)), 122.97 (C-4), 126.60 (C-Ar), 126.90 (C-Ar), 129.53 (C-10), 141.36 (C-9), 145.24 (C-Ar), 145.56 (C-Ar), 156.49 (C-5), 199.55 (C-3).




[00120] *3,3-Ethylendioxy-5α-hidroxy-11β-[4-p-(n-heptyl)phenyl]estra-9-en-17-one (3i)*

5 Compound **3i** was synthesized following the procedure described for compound **3a**, where 4g of compound **2** (12.10 mmol) was reacted with 1.76 g of Mg (72.6 mmol), 15.5 g of 4-bromo 4'-n-heptylbenzene (60.5 mmol) and 598 mg of CuCl (6.05 mmol) to afford **3i** as a white amorphous solid (4.73 g), yield: 77%

¹H NMR (CDCl₃, 300 MHz) δ 0.48 (s, 3H, H-18), 0.85 (t, *J*= 6 Hz, 3H, -CH₃), 1.27 (m, 10H, -CH₂), 2.54 (t, *J*= 9 Hz, 3H, -CH₃), 3.98 (m, 4H, ketal), 4.29 (d, *J*= 6 Hz, H-11), 4.3 (s, 1 H, -OH), 7.05 (d, *J*= 9 Hz, 2H, H-Ar), 7.11 (d, *J*= 9 Hz, 2H, H-Ar).

10




[00121] *3,3-Ethylendioxy-11β-[4-p-(n-heptyl)phenyl]estra-9-en-17-one (4i)*

15 The synthesis of compound **4i** was done following the procedure described for the synthesis of compound **4a** where 4.5 g of **3i** (8.89 mmol) was treated with 4.2 mL of acetic anhydride (44.4 mmol), 108.6 mg of DMAP (0.89 mmol) and 40 mL of pyridine, to get 1.76 g of **4i** as an amorphous solid (41% yield)


¹H NMR (CDCl₃, 300 MHz) δ 0.5 (s, 3H, H-18), 0.87 (t, *J*= 6 Hz, 3H, -CH₃), 1.27 (m, 10H, -CH₂), 3.95 (m, 4H, ketal), 4.3 (d, *J*= 6 Hz, H-11), 5.39 (s, 1 H, H-4), 7.06 (d, *J*= 9 Hz, 2H, H-Ar), 7.11 (d, *J*= 9 Hz, 2H, H-Ar).

20



[00122] *17α-[1,1-difluoro-3-[tris(1-methylethyl)silyl]-2-propyn-1-yl]-17β-hydroxy-11β-[4-p-(n-heptyl)phenyl]estra-4,9-dien-3-one (6i)*


- 5 This reaction was done following the same procedure described for the synthesis of compound **6a** where 0.61 g of compound **4i** (4.89 mmol) was reacted with 1.6 g of 3-bromo-3,3-difluoro-1-triisopropylsilylpropyne (4.9 mmol), and 2.6 mL of a 2M solution of n-BuLi (4.9 mmol). The crude product **5i** obtained was hydrolyzed with 1.2 mL (4.8 mmol) of a 4N solution of HCl to get 0.52 g of **6i** as an amorphous solid yield: 64%
- 10 ¹H NMR (CDCl₃, 300 MHz) δ 0.6 (s, 3H, H-18), 0.87 (m, 3H, -CH₃), 1.1(s, 18H, Si(CH)₃(CH₃)₆), 1.28 (m, 10H, -CH₂), 4.39 (m, 1H, H-11), 5.77(s, 1 H, H-4), 7.06 (d, *J*= 9 Hz, 2H, H-Ar), 7.11 (d, *J*= 9 Hz, 2H, H-Ar).



- 15 [00123] *17α-(1,1-difluoro-2-propyn-1-yl)-17β-hydroxy-11β-[4-p-(n-heptyl)phenyl]estra-4,9-dien-3-one (EC362)*

The reaction was done following the same procedure as described for **EC330** where 513 mg of compound **6i** (0.75 mmol) was treated with 1.52 mL of a 1M solution of TBAF to get 300 mg g of **EC362** as a white powder, yield: 76%.

- 20 ¹H NMR (CDCl₃, 300 MHz) δ 0.61 (s, 3H, H-18), 0.87 (m, 3H, -CH₃), 1.27 (m, 10H, -CH₂), 2.90 (t, *J*= 6 Hz, 1H), 4.39 (d, *J*= 6 Hz, 1H, H-11), 5.77(s, 1 H, H-4), 7.07 (m, 4H, H-Ar).
- ¹³C NMR (CDCl₃, 75 MHz) δ 14.05, 16.50, 22.56, 24.25, 25.72, 27.60, 29.07, 29.11, 31.01, 31.28, 31.72, 38.58, 39.15, 40.26, 47.64, 51.14, 85.74 (t, *J*=25Hz), 115.98 (t, *J*=241 Hz), 122.85, 126.56, 128.46, 129.42, 140.28, 141.25, 145.33, 156.60, 199.59




[00124] *3,3-Ethylendioxy-5α-hidroxy-11β-[4-p-(methylthio)phenyl]estra-4,9-dien-17-one (3j)*

5 Following the procedure described for compound **3a**, 3g of compound **2** (9.07 mmol) was reacted with 772 mg of Mg (31.77 mmol), 6.5g of 4-bromo-thioanisole (31.7 mmol) and 454 mg of CuCl (4.5 mmol) to afford **3j** as an amorphous solid (2.83g), yield: 69%.

¹H NMR (CDCl₃, 300 MHz) δ 0.51 (s, 3H, H-18), 2.64 (s, 3H, -SCH₃), 3.97 (m, 4H, ketal), 4.28 (d, *J*= 7 Hz, H-11), 4.38 (s, 1 H, -OH), 7.14 (m, 4H, H-Ar).


10



[00125] *3,3-Ethylendioxy-5α-hidroxy-11β-[4-p-(methylthio)phenyl]estra-4,9-dien-17-one (4j)*

15 The synthesis of compound **4j** was done following the procedure described for compound **4a** where 2.83 g of **3j** (6.22 mmol) was heated at 70°C with 2.93 mL of acetic anhydride (31.11 mmol), 380 mg of DMAP (3.11 mmol) and 30 mL of pyridine, to get 2.01 g of **4j** as an off white amorphous solid. (75% yield)


20 ¹H NMR (CDCl₃, 300 MHz) δ 0.52 (s, 3H, H-18), 2.64 (s, 3H, -SCH₃), 3.97 (m, 4H, ketal), 4.28 (d, *J*= 7 Hz, H-11), 5.4 (s, 1H, H-4), 7.14 (m, 4H, H-Ar).

**6j**

[00126] *17α-[1,1-difluoro-3-[tris(1-methylethyl)silyl]-2-propyn-1-yl]-17β-hydroxy-11β-[4-p-(methylthio)phenyl]estra-4,9-dien-3-one (6j)*

5 Following the same procedure described for **6a**, 2g of compound **4j** (4.58 mmol) was reacted with 3.56g of 3-bromo-3,3-difluoro-1-triisopropylsilylpropyne (11.4 mmol), and 5.8 mL of a 2M solution of n-BuLi (11.9 mmol). The crude **5j** obtained was hydrolyzed with 4.6 mL (18.3 mmol) of a 4N solution of HCl to get 1.68g of **6j** as an amorphous solid, yield: 58%.


¹H NMR (CDCl₃, 300 MHz) δ 0.61 (s, 3H, H-18), 1.03 (m, 3H, Si(CH₃)₃(CH₃)₆), 1.11 (s, 18H, Si(CH₃)₃(CH₃)₆), 2.46 (s, 3H, -SCH₃), 4.37 (m, 1H, H-11), 5.77 (s, 1H, H-4), 7.08 (d, *J*= 9 Hz, 2H, H-Ar), 7.16 (d, *J*= 9 Hz, 2H, H-Ar).

**EC363**

[00127] *17α-(1,1-difluoro-2-propyn-1-yl)-17β-hydroxy-11β-[4-p-(methylthio)phenyl]estra-4,9-dien-3-one (EC363)*

The reaction was done following the same procedure reported for **EC330** where 1.68 g of compound **6j** (2.62 mmol) was treated with 5.2 mL of a 1M solution of TBAF to get 530 mg of **EC356** as a beige powder, yield: 60%

¹H NMR (CDCl₃, 300 MHz) δ 0.61 (s, 3H, H-18), 2.45 (s, 3H, -SCH₃), 2.91 (t, *J*= 5.4 Hz, 1H, acetylenic hydrogen), 4.39 (d, *J*= 7.2 Hz, 1H, H-11), 5.77 (s, 1H, H-4), 7.11 (d, *J*= 9 Hz, 2H, H-Ar), 7.16 (d, *J*= 9 Hz, 2H, H-Ar).



[00128] *3,3-Ethylendioxy-5α-hidroxy-11β-[4-p-(cyclopropylthio)phenyl]estra-9-en-17-one (31)*


5 A three neck dried flask was charged with Mg turnings (350 mg, 14.52 mmol), a crystal of I₂ was added and swirled over the Mg and kept for 5 min. 5 mL of anhydrous THF was added followed by 0.5 mL of 1,2-dibromoethane. The reaction was slightly warmed with a heat gun. When Mg starts reacting, a solution of the 4-bromophenyl-cyclopropyl sulfide (2.5 g, 10.9 mmol) in 10 mL of THF was added drop by drop, the I₂ color gets discharged during the

10 addition, after the addition was over, the reaction was stirred for 15 min and then refluxed at 60°C (oil bath temperature) for 30 min. Afterward, the reaction was cooled to room temperature and CuCl (107.8 mg, 1.1 mmol) was added. The reaction was stirred for 6 hrs and then a solution of the epoxide **2** (1.2 g, 3.63 mmol) in THF (15 mL) was added dropwise and stirred for 1h. The TLC showed a more polar product. The reaction was cooled and

15 quenched by the addition of sat solution of NH₄Cl and extracted with ethyl acetate, the organic layer was washed with water and brine, the solvent was removed under vacuum. The crude was purified by column chromatography using 40% of ethyl acetate in hexane to get 0.85 g of white foam (44% yield), mp 93-95°C. UV (nm): 200, 257. R_f: 0.35 (5:5, Hex:EtOAc). FT IR (ATR, cm⁻¹): 3514, 2927, 2873, 1739, 1489. ¹H NMR (CDCl₃, 300 MHz) δ 0.50 (s, 3H, H-18), 0.67 (m, 2H, cyclopropyl), 3.9 (m, 4H, ketal), 4.29 (d, *J*= 6.9 Hz, 1H, H-11), 4.37 (s, 1H, -OH), 7.15 (d, *J*= 8.4 Hz, 2H, H-Ar), 7.26 (d, *J*= 10.2 Hz, 2H, H-Ar).


20 ¹³C NMR (CDCl₃, 75 MHz) δ 8.45 (cyclopropyl), 12.04 (cyclopropyl), 14.32 (C-18), 64.05 (ketal), 64.67 (ketal), 69.97 (C-5), 108.59 (C-3), 126.47 (C-Ar), 127.53 (C-Ar), 133.37 (C-10), 135.32 (C-Ar), 135.58 (C-9), 143.05 (C-Ar), 219.82 (C-17).

25



[00129] *3,3-Ethylenedioxy-11 β -[4-p-(cyclopropylthio)phenyl]estra-4,9-dien-17-one(4I)*

- 5 The synthesis of compound **4I** was done following the procedure described for the synthesis of compound **4a** where 1.7 g of **3I** (3.4 mmol) was reacted with 1.8 mL of acetic anhydride (17 mmol), 41 mg of DMAP (0.34 mmol) and 20 mL of pyridine, to get 1.22 g of **4I** as a white powder (78% yield), mp 88-90 °C. R_f: 0.55 (5:5, Hex:EtOAc). UV (nm): 200, 256. FT IR (ATR, cm⁻¹): 2933, 2873, 1739, 1638, 1489.
- 10 ¹H NMR (CDCl₃, 300 MHz) δ 0.013 (s, 3H, H-18), 0.53 (m, 2H, cyclopropyl), 0.68 (m, 2H, cyclopropyl), 3.9 (m, 4H, ketal), 4.3 (d, *J*= 6.9 Hz, H-11), 5.41 (s, 1H, H-4), 7.16 (d, *J*= 8.4 Hz, 2H, H-Ar), 7.27 (d, *J*= 9.3 Hz, 2H, H-Ar).
- ¹³C NMR (CDCl₃, 75 MHz) δ 8.45 (cyclopropyl), 12.07 (cyclopropyl), 14.46 (C-18), 64.42 (ketal), 64.57 (ketal), 106.09 (C-3), 121.98 (C-4), 126.18 (C-Ar), 126.65 (C-Ar), 127.59 (C-Ar), 130.34 (C-10), 135.61 (C-Ar), 137.30 (C-5), 139.23 (C-9), 141.77 (C-Ar), 219.6 (C-17).
- 15



[00130] *17 α -[1,1-difluoro-3-[tris(1-methylethyl)silyl]-2-propyn-1-yl]-17 β -hydroxy-11 β -[4-p-(cyclopropyl sulfonyl)phenyl]estra-4,9-dien-3-one (6I)*

- 20 This reaction was done following the same procedure described for the synthesis of compound **6a** where 1.2 g of compound **4I** (2.6 mmol) was reacted with 3.2 g of 3-bromo-3,3-difluoro-1-triisopropylsilylpropyne (10.4 mmol), and 6.5 mL of a 2M solution of n-BuLi (13 mmol). The crude was used for the next step oxidation using Oxone. The crude was


WO 2016/154203

PCT/US2016/023582

dissolved in a mixture of 50 mL of THF and 50 mL of methanol. A solution of 6.4g of Oxone (20.8 mmol) in 30 mL of water was slowly added dropwise at 0 °C and was stirred for 3.5 hrs at 0 °C. The TLC showed a more polar product. The reaction was quenched by adding water and extracted with ethyl acetate, the organic layer was washed with water and brine, the solvent was removed under vacuum. The crude was purified by column chromatography using 50% ethyl acetate in hexane to get 700 mg of a beige product (45% yield), mp 123-125°C. R_f: 0.5 (7:3, Hex:EtOAc). UV (nm): 200, 230, 297. FT IR (ATR, cm⁻¹): 3466, 2933, 2879, 1726, 1631, 1483.

¹H NMR (CDCl₃, 300 MHz) δ 0.56 (s, 3H, H-18), 1.02 (s, 3H, Si(CH₃)₃(CH₃)₆), 1.1 (s, 18H, Si(CH₃)₃(CH₃)₆), 4.40 (d, *J*= 5.4 Hz, 1H, H-11), 5.80 (s, 1H, H-4), 7.38 (d, *J*= 8.4 Hz, 2H, H-Ar), 7.8 (d, *J*= 8.4 Hz, 2H, H-Ar).

¹³C NMR (CDCl₃, 75 MHz) δ 5.96 (cyclopropyl), 10.91 (C-18), 18.51 (Si(CH₃)₃(CH₃)₆), 24.40 (Si(CH₃)₃(CH₃)₆), 86.18 (t, *J*= 24.3 Hz, (CF₂CC)), 123.57 (C-4), 127.73 (C-Ar), 127.93 (C-Ar), 130.36 (C-10), 138.31 (C-9), 143.21 (C-Ar), 151.0 (C-Ar), 155.76 (C-5), 198.93 (C-3).



EC361

[00131] *17α-(1,1-difluoro-2-propyn-1-yl)-17β-hydroxy-11β-[4-p-(cyclopropylsulfonyl)phenyl]estra-4,9-dien-3-one (EC361)*

Following the same procedure reported for EC330, 700 mg of compound **61** (1.03 mmol) was treated with 1.53 mL of a 1M solution of TBAF to get 320 mg of EC361 as a white powder, yield: 59%, mp 145-148 °C. R_f: 0.23 (5:5, Hex:EtOAc). UV (nm): 200, 203, 301. FT IR (ATR, cm⁻¹): 3453, 3244, 2933, 2873, 2124, 1732, 1645.

¹H NMR (CDCl₃, 300 MHz) δ 0.54 (s, 3H, H-18), 2.92 (t, *J*= 5.4 Hz, 1H, acetylenic hydrogen), 4.51 (d, *J*= 6 Hz, 1H, H-11), 5.80 (s, 1H, H-4), 7.40 (d, *J*= 8.4 Hz, 2H, H-Ar), 7.82 (d, *J*= 8.7 Hz, 2H, H-Ar).



¹³C NMR (CDCl₃, 75 MHz) δ 5.99 (cyclopropyl), 14.16 (C-18), 85.71 (t, *J*= 24.9 Hz, (CF₂CC)), 123.60 (C-4), 127.78 (C-Ar), 127.94 (C-Ar), 130.41 (C-10), 138.27 (C-9), 143.18 (C-Ar), 150.93 (C-Ar), 155.82 (C-5), 199.08 (C-3).

In this patent, certain U.S. patents, U.S. patent applications, and other materials (e.g., articles) have been incorporated by reference. The text of such U.S. patents, U.S. patent applications, and other materials is, however, only incorporated by reference to the extent that no conflict exists between such text and the other statements and drawings set forth
5 herein. In the event of such conflict, then any such conflicting text in such incorporated by reference U.S. patents, U.S. patent applications, and other materials is specifically not incorporated by reference in this patent.

Further modifications and alternative embodiments of various aspects of the invention
10 will be apparent to those skilled in the art in view of this description. Accordingly, this description is to be construed as illustrative only and is for the purpose of teaching those skilled in the art the general manner of carrying out the invention. It is to be understood that the forms of the invention shown and described herein are to be taken as examples of
15 embodiments. Elements and materials may be substituted for those illustrated and described herein, parts and processes may be reversed, and certain features of the invention may be utilized independently, all as would be apparent to one skilled in the art after having the benefit of this description of the invention. Changes may be made in the elements described herein without departing from the spirit and scope of the invention as described in the following claims.

20

1. A cytotoxic compound having the structure (I) or (II):




(I)

(II)

where:

10



X is O, NH, or S;

15

n = 1-18; m = 1-18;

R² is H, F, Cl, $-C(O)-R^6$, or $-CH_2(OH)$;

R³ is H, F, Cl, $-C(O)-R^6$, or $-CH_2(OR^6)$;

20

R⁴ is H, alkyl, $-CH_2-OH$, $-CO_2R^6$, $-CON(R^6)_2$;

R⁵ is alkyl, alkenyl, alkylacyl, cycloalkyl, heterocycle, $-CN$, alkoxy, $-N(R^6)_2$, $-CON(R^6)_2$,

25


$-S(O)R^6$, $-SR^6$, $-SO_2R^6$; or $-(CH_2)_p-CH_2-Y$;

Y is H, OR^6 , SCH_3 , CF_3 , $-N(R^6)_2$; p = 1-18; and

R^6 is H, alkyl, or cycloalkyl.

2. The cytotoxic compound of claim 1, wherein the cytotoxic compound has the structure

5 (III):



10 where:

R^2 is H, F, Cl, -CO-, or -C(OH)-;

R^3 is H, F, Cl, -CO-, or -C(OH)-;

15

R^4 is H, alkyl, -CH₂-OH, -CO₂R⁶, -CON(R⁶)₂;

R^5 is alkyl, alkenyl, alkylacyl, cycloalkyl, heterocycle, -CN, alkoxy, -N(R⁶)₂, -CON(R⁶)₂,

20


-S(O)R⁶, -SR⁶, -SO₂R⁶; or -(CH₂)_p-CH₂-Y; and

Y is H, OR⁶, SCH₃, CF₃, -N(R⁶)₂; p = 1-18; and

R^6 is H, alkyl, or cycloalkyl.

25

3. The cytotoxic compound of claim 1, wherein the cytotoxic compound has the structure (IV):




where:

10 R^5 is alkyl, alkenyl, alkylacyl, cycloalkyl, heterocycle, -CN, alkoxy, $-N(R^6)_2$, -
 $CON(R^6)_2$,
 $-S(O)R^6$, $-SR^6$, or $-SO_2R^6$; and

R^6 is H, alkyl, or cycloalkyl.

15 4. The cytotoxic compound of claim 1, wherein the cytotoxic compound has the structure (IV):



20

where:

WO 2016/154203

PCT/US2016/023582

R^5 is alkyl, alkenyl, alkylacyl, cycloalkyl, 1,3-imidazolyl, alkoxy, $-N(R^6)_2$, $-SR_6$, or $-SO_2R^6$; and

R^6 is H, lower alkyl, or cycloalkyl.

5

5. The cytotoxic compound of claim 4, wherein R^5 is alkyl, alkenyl, or cycloalkyl.

6. The cytotoxic compound of claim 4, wherein R^5 is 1,3-imidazolyl.

10 7. The cytotoxic compound of claim 4, wherein R^5 is alkoxy.

8. The cytotoxic compound of claim 4, wherein R^5 is $-N(R^6)_2$.

9. The cytotoxic compound of claim 4, wherein R^5 is $-SR_6$.

15

10. The cytotoxic compound of claim 4, wherein R^5 is $-SO_2R^6$.

11. A method of treating cancer in a subject comprising administering to a subject a medicament comprising an effective amount of a cytotoxic small molecule compound that
20 inhibits leukemia inhibitory factor or leukemia inhibitory factor receptor.

12. The method of claim 11, wherein the cancer is a cancer that overexpresses leukemia inhibitory factor.

25 13. The method of claim 11, wherein the cytotoxic small molecule compound inhibits MDM2.

14. The method of claim 11, wherein the cytotoxic small molecule compound inhibits carcinoma associated fibroblast.

30


15. The method of claim 11, wherein the cytotoxic small molecule compound stabilize P53 levels.

16. The method of claim 11, wherein the cancer exhibits a desmoplastic stromal response.

17. The method of claim 11, wherein the cytotoxic small molecule compound has the structure as set forth in any one of claims 1-10.

5 18. The method of claim 11, wherein the cancer exhibits cancer initiating stem cells (CISC) or cancer associated stem cells (CASC).

19. A compound having the structure (V):



where:

15 R^5 is alkyl, alkenyl, alkylacyl, cycloalkyl, heterocycle, -CN, alkoxy, -N(R^6)₂, -CON(R^6)₂, -S(O) R^6 , -SR₆, -SO₂ R^6 ; or -(CH₂)_p-CH₂-Y; and

Y is H, OR⁶, SCH₃, CF₃, -N(R^6)₂; p = 1-18; and

20 R^6 is H, alkyl, or cycloalkyl.

20. The compound of claim 19, where:

where:

25 R^5 is alkyl, alkenyl, alkylacyl, cycloalkyl, 1,3-imidazolyl, alkoxy, -N(R^6)₂, -SR₆, or -SO₂ R^6 ; and

R^6 is H, lower alkyl, or cycloalkyl.

WO 2016/154203**PCT/US2016/023582**

21. A method of treating hypertrophic fibroblasts in a subject comprising administering to a subject a medicament comprising an effective amount of a cytotoxic small molecule compound that reduces the amount of hypertrophic fibroblasts in the subject.
- 5 22. Use of a cytotoxic small molecule compound as described in any one of claims 1 to 10 as a biomarker/companion diagnostic (CDx) for selecting a suitable population to treat with LIF inhibitors by using the cytotoxic compounds to down-regulate phosphorylation of STAT3.




FIG. 1

2 / 10




FIG. 2

3 / 10




FIG. 3




FIG. 4A





FIG. 4B

5 / 10




FIG. 5

6 / 10




FIG. 6A




FIG. 6B

7 / 10




FIG. 7




FIG. 8

8 / 10




FIG. 9




FIG. 10




FIG.11




FIG. 4A