



(43) International Publication Date
8 November 2012 (08.11.2012)

(10) International Publication Number
WO 2012/151324 A1

- (51) International Patent Classification:
C12N 15/11 (2006.01) C07H 21/04 (2006.01)
- (21) International Application Number:
PCT/US2012/036189
- (22) International Filing Date:
2 May 2012 (02.05.2012)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
61/481,654 2 May 2011 (02.05.2011) US
61/587,074 16 January 2012 (16.01.2012) US
- (71) Applicant (for all designated States except US): **ISIS PHARMACEUTICALS, INC.** [US/US]; 2855 Gazelle Court, Carlsbad, CA 92010 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **HASTINGS, Michelle, L.** [US/US]; 3333 Green Bay Road, North Chicago, IL 60064 (US). **RIGO, Frank** [ES/US]; 2855 Gazelle Court, Carlsbad, CA 92010 (US). **BENNETT, C., Frank** [US/US]; 2855 Gazelle Court, Carlsbad, CA 92010 (US).
- (74) Agents: **GINGRICH, Brenden** et al.; Knobbe Martens Olson & Bear LLP, 2040 Main Street, 14th Floor, Irvine, CA 92614 (US).

- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: ANTISENSE COMPOUNDS TARGETING GENES ASSOCIATED WITH USHER SYNDROME

(57) Abstract: The present invention provides compounds comprising oligonucleotides complementary to an Usher transcript. Certain such compounds are useful for hybridizing to an Usher transcript, including but not limited, to an Usher transcript in a cell. In certain embodiments, such hybridization results in modulation of splicing of the Usher transcript. In certain such embodiments, the Usher transcript includes a mutation that results in cryptic splicing and hybridization of the oligonucleotide results in a decrease in the amount of cryptic splicing. In certain embodiments, such compounds are used to treat Usher Syndrome.



WO 2012/151324 A1

**ANTISENSE COMPOUNDS TARGETING GENES
ASSOCIATED WITH USHER SYNDROME**

SEQUENCE LISTING

5 The present application is being filed along with a Sequence Listing in electronic format. The Sequence Listing is provided as a file entitled CORE0095WOSEQ.txt, created May 2, 2012, which is 72 Kb in size. The information in the electronic format of the sequence listing is incorporated herein by reference in its entirety.

BACKGROUND OF THE INVENTION

10 Usher Syndrome is a leading genetic cause of combined blindness and deafness. In certain instances, Usher Syndrome results from a mutation in an Usher gene. See, e.g., Millan JM et al., An Update on the Genetics of Usher Syndrome, Journal of Ophthalmology, V 2001, Article ID 417217 (2001).

 Antisense compounds have been used to modulate target nucleic acids. Antisense compounds comprising a variety of chemical modifications and motifs have been reported. In certain instances, such
15 compounds are useful as research tools, diagnostic reagents, and as therapeutic agents. In certain instances antisense compounds have been shown to modulate protein expression by binding to a target messenger RNA (mRNA) encoding the protein. In certain instances, such binding of an antisense compound to its target mRNA results in cleavage of the mRNA. Antisense compounds that modulate processing of a pre-mRNA have also been reported. Such antisense compounds alter splicing, interfere with polyadenylation or prevent
20 formation of the 5'-cap of a pre-mRNA.

 Certain antisense compounds have been described previously. See for example United States Patent No. 7,399,845 and published International Patent Application No. WO 2008/049085, which are hereby incorporated by reference herein in their entirety.

25 **SUMMARY OF THE INVENTION**

 In certain embodiments, the present invention provides compounds comprising oligonucleotides. In certain embodiments, such oligonucleotides are complementary to an Usher transcript. In certain such embodiments, the oligonucleotide is complementary to a target region of the Usher transcript comprising exon 2, intron 2, exon 3, and intron 3. In certain embodiments, the Usher transcript comprises a mutation that
30 results in a cryptic splice site. In certain embodiments, oligonucleotides inhibit cryptic splicing. In certain such embodiments, normal splicing is increased. In certain embodiments, truncated, but non-cryptic splicing is increased.

 The present disclosure provides the following non-limiting numbered embodiments:

Embodiment 1: A compound comprising a modified oligonucleotide consisting of 8 to 30 linked nucleosides and having a nucleobase sequence comprising a complementary region comprising at least 8 contiguous nucleobases complementary to a target region of equal length of an Usher transcript.

- 5 Embodiment 2: The compound of embodiment 1, wherein the complementary region of the modified oligonucleotide is 100% complementary to the target region.

Embodiment 3: The compound of embodiment 1 or 2, the complementary region of the modified oligonucleotide comprises at least 10 contiguous nucleobases.

- 10 Embodiment 4: The compound of embodiment 1 or 2, the complementary region of the modified oligonucleotide comprises at least 15 contiguous nucleobases.

- 15 Embodiment 5: The compound of embodiment 1 or 2, the complementary region of the modified oligonucleotide comprises at least 20 contiguous nucleobases.

- Embodiment 6: The compound of any of embodiments 1-5, wherein the nucleobase sequence of the oligonucleotide is at least 80% complementary to an equal-length region of the Usher transcript, as measured over the entire length of the oligonucleotide.

- 20 Embodiment 7: The compound of any of embodiments 1-5, wherein the nucleobase sequence of the oligonucleotide is at least 90% complementary to an equal-length region of the Usher transcript, as measured over the entire length of the oligonucleotide.

- 25 Embodiment 8: The compound of any of embodiments 1-5, wherein the nucleobase sequence of the oligonucleotide is 100% complementary to an equal-length region of the Usher transcript, as measured over the entire length of the oligonucleotide.

- 30 Embodiment 9: The compound of any of embodiments 1-8, wherein the target region is within exon 3 of the Usher transcript.

Embodiment 10: The compound of any of embodiments 1-9, wherein the target region is within nucleobase 138475 and nucleobase 138660 of SEQ ID NO.: 1.

Embodiment 11: The compound of any of embodiments 1-9, wherein the target region is within nucleobase 138480 and nucleobase 138620 of SEQ ID NO.: 1.

Embodiment 12: The compound of any of embodiments 1-9, wherein the target region is within nucleobase 138550 and nucleobase 138620 of SEQ ID NO.: 1.

Embodiment 13: The compound of any of embodiments 1-9, wherein the target region is within nucleobase 138577 and nucleobase 138600 of SEQ ID NO.: 1.

Embodiment 14: The compound of any of embodiments 1-13, wherein the antisense oligonucleotide comprises SEQ ID NO: 30.

Embodiment 15: The compound of any of embodiments 1-13, wherein the antisense oligonucleotide comprises SEQ ID NO: 29.

Embodiment 16: The compound of any of embodiments 1-13, wherein the antisense oligonucleotide comprises SEQ ID NO: 27.

Embodiment 17: The compound of any of embodiments 1-13, wherein the antisense oligonucleotide comprises SEQ ID NO: 50.

Embodiment 18: The compound of any of embodiments 1-13, wherein the antisense oligonucleotide comprises SEQ ID NO: 56.

Embodiment 19: The compound of any of embodiments 1-18, wherein the modified oligonucleotide comprises at least one modified nucleoside.

Embodiment 20: The compound of embodiment 19, wherein at least one modified nucleoside comprises a modified sugar moiety.

Embodiment 21: The compound of embodiment 20, wherein at least one modified sugar moiety is a 2'-substituted sugar moiety.

Embodiment 22: The compound of embodiment 21, wherein the 2'-substituted of at least one 2'-substituted sugar moiety is selected from among: 2'-OMe, 2'-F, and 2'-MOE.

Embodiment 23: The compound of embodiment 22, wherein the 2'-substituent of at least one 2'-substituted sugar moiety is a 2'-MOE.

5 Embodiment 24: The compound of any of embodiments 20-23, wherein at least one modified sugar moiety is a bicyclic sugar moiety.

Embodiment 25: The compound of embodiment 24, wherein at least one bicyclic sugar moiety is LNA or cEt.

10

Embodiment 26: The compound of any of embodiments 20-25, wherein at least one sugar moiety is a sugar surrogate.

15

Embodiment 27: The compound of embodiment 26, wherein at least one sugar surrogate is a morpholino.

Embodiment 28: The compound of embodiment 26, wherein at least one sugar surrogate is a modified morpholino.

20

Embodiment 29: The compound of any of embodiment 1-28, wherein the modified oligonucleotide comprises at least 5 modified nucleosides, each independently comprising a modified sugar moiety.

Embodiment 30: The compound of embodiment 29, wherein the modified oligonucleotide comprises at least 10 modified nucleosides, each independently comprising a modified sugar moiety.

25

Embodiment 31: The compound of embodiment 29, wherein the modified oligonucleotide comprises at least 15 modified nucleosides, each independently comprising a modified sugar moiety.

30

Embodiment 32: The compound of embodiment 29, wherein each nucleoside of the modified oligonucleotide is a modified nucleoside, each independently comprising a modified sugar moiety

Embodiment 33: The compound of any of embodiments 1-32, wherein the modified oligonucleotide comprises at least two modified nucleosides comprising modified sugar moieties that are the same as one another.

35

Embodiment 34: The compound of any of embodiments 1-33, wherein the modified oligonucleotide comprises at least two modified nucleosides comprising modified sugar moieties that are different from one another.

5 Embodiment 35: The compound of any of embodiments 1-34, wherein the modified oligonucleotide comprises a modified region of at least 5 contiguous modified nucleosides.

Embodiment 36: The compound of embodiment 35, wherein the modified oligonucleotide comprises a modified region of at least 10 contiguous modified nucleosides.

10

Embodiment 37: The compound of embodiment 35, wherein the modified oligonucleotide comprises a modified region of at least 15 contiguous modified nucleosides.

15

Embodiment 38: The compound of embodiment 35, wherein the modified oligonucleotide comprises a modified region of at least 20 contiguous modified nucleosides.

Embodiment 39: The compound of any of embodiments 34-38, wherein each modified nucleoside of the modified region has a modified sugar moiety independently selected from among: 2'-F, 2'-OMe, 2'-MOE, cEt, LNA, morpholino, and modified morpholino.

20

Embodiment 40: The compound of any of embodiments 35-39, wherein the modified nucleosides of the modified region each comprise the same modification as one another.

25

Embodiment 41: The compound of embodiment 40, wherein the modified nucleosides of the modified region each comprise the same 2'-substituted sugar moiety.

Embodiment 42: The compound of embodiment 40, wherein the 2'-substituted sugar moiety of the modified nucleosides of the region of modified nucleosides is selected from 2'-F, 2'-OMe, and 2'-MOE.

30

Embodiment 43: The compound of embodiment 41, wherein the 2'-substituted sugar moiety of the modified nucleosides of the region of modified nucleosides is 2'-MOE.

Embodiment 44: The compound of embodiment 40, wherein the modified nucleosides of the region of modified nucleosides each comprise the same bicyclic sugar moiety.

35

Embodiment 45: The compound of embodiment 44, wherein the bicyclic sugar moiety of the modified nucleosides of the region of modified nucleosides is selected from LNA and cEt.

Embodiment 46: The compound of embodiment 40, wherein the modified nucleosides of the region of modified nucleosides each comprises a sugar surrogate.

Embodiment 47: The compound of embodiment 46, wherein the sugar surrogate of the modified nucleosides of the region of modified nucleosides is a morpholino.

Embodiment 48: The compound of embodiment 46, wherein the sugar surrogate of the modified nucleosides of the region of modified nucleosides is a modified morpholino.

Embodiment 49: The compound of any of embodiments 1-48, wherein the modified nucleotide comprises no more than 4 contiguous naturally occurring nucleosides.

Embodiment 50: The compound of any of embodiments 1-48, wherein each nucleoside of the modified oligonucleotide is a modified nucleoside.

Embodiment 51: The compound of embodiment 50 wherein each modified nucleoside comprises a modified sugar moiety.

Embodiment 52: The compound of embodiment 51, wherein the modified nucleosides of the modified oligonucleotide comprise the same modification as one another.

Embodiment 53: The compound of embodiment 52, wherein the modified nucleosides of the modified oligonucleotide each comprise the same 2'-substituted sugar moiety.

Embodiment 54: The compound of embodiment 53, wherein the 2'-substituted sugar moiety of the modified oligonucleotide is selected from 2'-F, 2'-OMe, and 2'-MOE.

Embodiment 55: The compound of embodiment 54, wherein the 2'-substituted sugar moiety of the modified oligonucleotide is 2'-MOE.

Embodiment 56: The compound of embodiment 52, wherein the modified nucleosides of the modified oligonucleotide each comprise the same bicyclic sugar moiety.

Embodiment 57: The compound of embodiment 56, wherein the bicyclic sugar moiety of the modified oligonucleotide is selected from LNA and cEt.

5 Embodiment 58: The compound of embodiment 52, wherein the modified nucleosides of the modified oligonucleotide each comprises a sugar surrogate.

Embodiment 59: The compound of embodiment 58, wherein the sugar surrogate of the modified oligonucleotide is a morpholino.

10

Embodiment 60: The compound of embodiment 58, wherein the sugar surrogate of the modified oligonucleotide is a modified morpholino.

15

Embodiment 61: The compound of any of embodiments 1-60, wherein the modified oligonucleotide comprises at least one modified internucleoside linkage.

Embodiment 62: The compound of embodiment 61, wherein each internucleoside linkage is a modified internucleoside linkage.

20

Embodiment 63: The compound of embodiment 61 or 62, comprising at least one phosphorothioate internucleoside linkage.

25

Embodiment 64: The compound of embodiment 62, wherein each internucleoside linkage is a modified internucleoside linkage and wherein each internucleoside linkage comprises the same modification.

Embodiment 65: The compound of embodiment 64, wherein each internucleoside linkage is a phosphorothioate internucleoside linkage.

30

Embodiment 66: The compound of any of embodiments 1-65 comprising at least one conjugate.

Embodiment 67: The compound of any of embodiments 1-66 consisting of the modified oligonucleotide.

- Embodiment 68: The compound of any of embodiments 1-67, wherein the compound modulates splicing of the Usher transcript.
- Embodiment 69: The compound of any of embodiments 1-68, having a nucleobase sequence comprising any of the sequences as set forth in SEQ ID NOs. 2-57.
- Embodiment 70: The compound of any of embodiments 1-68, having a nucleobase sequence comprising any of the sequences as set forth in SEQ ID NOs. 2-31 or 49-57.
- Embodiment 71: The compound of any of embodiments 1-68, having a nucleobase sequence comprising any of the sequences as set forth in SEQ ID NOs. 2-31.
- Embodiment 72: The compound of any of embodiments 1-68, having a nucleobase sequence comprising any of the sequences as set forth in SEQ ID NOs. 49-57.
- Embodiment 73: A method of modulating splicing an Usher transcript in a cell comprising contacting the cell with a compound according to any of embodiments 1-72.
- Embodiment 74: The method of embodiment 73, wherein the cell is in vitro.
- Embodiment 75: The method of embodiment 73, wherein the cell is in an animal.
- Embodiment 76: A method of modulating the expression of harmonin in a cell, comprising contacting the cell with a compound according to any of embodiments 1-72.
- Embodiment 77: The method of embodiment 76, wherein harmonin expression is increased.
- Embodiment 78: The method of embodiment 76, wherein the cell is in vitro.
- Embodiment 79: The method of embodiment 76, wherein the cell is in an animal.
- Embodiment 80: A method of increasing the total number of correctly shaped stereocilia bundles in the ear, comprising contacting a cell with a compound according to any of embodiments 1-72.
- Embodiment 81: The method of embodiment 80, wherein the cell is in an animal.

Embodiment 82: A pharmaceutical composition comprising a compound according to any of embodiments 1-72 and a pharmaceutically acceptable carrier or diluent.

5 Embodiment 83: The pharmaceutical composition of embodiment 82, wherein the pharmaceutically acceptable carrier or diluent is sterile saline.

Embodiment 84: A method comprising administering the pharmaceutical composition of embodiments 82 or 83 to an animal.

10

Embodiment 85: The method of embodiment 84, wherein the administration is by injection.

Embodiment 86: The method of embodiment 84 or 85, wherein the administration is systemic.

15 Embodiment 87: The method of embodiment 84 or 85 wherein the administration is local.

Embodiment 88: The method of embodiment 87, wherein the administration is to the eye of the animal.

20 Embodiment 89: The method of embodiment 87, wherein the administration is to the ear of an animal.

Embodiment 90: The method of any of embodiments 84-89, wherein the animal has one or more symptom of Usher Syndrome.

25 Embodiment 91: The method of embodiment 90, wherein the administration results in amelioration of at least one symptom of Usher Syndrome.

Embodiment 92: The method of embodiment 91, wherein the symptom is deafness.

30 Embodiment 93: The method of embodiment 91, wherein the symptom is blindness.

Embodiment 94: The method of any of embodiments 84-93, wherein the animal is a mouse.

Embodiment 95: The method of any of embodiments 84-93, wherein the animal is a human.

35

Embodiment 96: The method of any of embodiments 84-93, wherein the first administration occurs within 1-15 days after birth.

Embodiment 97: The method of any of embodiments 84-93, wherein the first administration occurs within 1-10 days after birth.

Embodiment 98: The method of any of embodiments 84-93, wherein the first administration occurs within 1-5 days after birth.

Embodiment 99: The method of any of embodiments 84-93, wherein the first administration occurs within 3-5 days after birth.

Embodiment 100: The method of any of embodiments 84-93, wherein the first administration is in utero.

Embodiment 101: Use of the compound of any of embodiments 1 to 72 or the composition of embodiments 82-83 for the preparation of a medicament for use in the treatment of Usher syndrome.

Embodiment 102: Use of the compound of any of embodiments 1 to 72 or the composition of embodiments 82-83 for the preparation of a medicament for use in the amelioration of one or more symptoms of Usher syndrome.

In certain embodiments, including, but not limited to any of the above numbered embodiments, the Usher transcript is in a human having Usher Syndrome. In certain such embodiments, the Usher gene of the human comprises a 216 mutation. In certain such embodiments, the Usher gene of the human comprises a 216AA mutation.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1a shows a schematic of Ush1c splicing and protein products encoded by expressed mRNAs. A portion of the Ush1c gene is shown. Exons are represented as boxes and lines are introns. Diagonal lines indicate splicing pathways. The location of the 216A mutation and the cryptic splice site are shown. Hatched lines indicate the cryptic splicing pathway activated by the 216A mutation. The three PDZ1 domains and the coiled-coil domain (CC1) of the harmonin protein are shown. Exon skipping removes the first 12 amino acids of the first PDZ domain. The truncated protein terminates within the PDZ1 domain.

Figure 1b shows a model of the minigene expression system (top) discussed in Example 1 and location of ASO targets on Ush1c exon 3 (bottom) discussed in Example 2. Figure 2(B) shows results of RT-

PCR analysis of splicing in the transfected HeLa cells, discussed in Example 2.

Figure 1c shows a schematic of how each of Isis No. 527133, 527134, 535401, and 535407 target the cryptic splice site of the Ush1c gene having the 216A mutation.

Figure 1d shows RT-PCR analysis of RNA isolated from HeLa cells was transfected with either a WT or mutant (Ush1c.216a) minigene and treated with saline (-) or increasing amounts of Ush1c-specific 2'MOE (final concentration 5, 10, 20, 40, 80 mM) discussed in Example 4. Cryptic splicing inhibition is quantitated in the histogram as the % cryptic splicing $[\text{cryptic}/(\text{cryptic} + \text{correct} + \text{skipped}) \times 100]$.

Figure 1e shows RT-PCR analysis of the dose response of Isis No. 527133, 527134, 535401, and 535407 after administration to Ush1c.2166AA mice.

Figure 1f shows RT-PCR analysis of mice kidney after the administration of Isis No. 527134 at different dosages.

Figure 1g shows Western-Blot analysis of lysates from Ush1c.216AA mice administered Isis No. 527134 at different dosages.

Figure 2a shows the results of Ush1c.2166AA mice treated with a mismatched control, Ush1c.2166AA mice treated with Isis No. 527134, and heterozygous mice in an open-field pathway trace.

Figure 2b shows the number of rotations of Ush1c.2166AA mice treated with a mismatched control, Ush1c.2166AA mice treated with Isis No. 527134, and heterozygous mice in an open-field pathway trace.

Figure 2c shows the number of rotations of 6 month old Ush1c.2166AA mice treated with a mismatched control, Ush1c.2166AA mice treated with Isis No. 527134, and heterozygous mice in an open-field pathway trace.

Figures 3a-d show auditory-evoked brainstem response (ABR) analysis at 8 kHz, 16 kHz, and 32 kHz and BBN, for Ush1c.2166AA mice treated with a mismatched control, Ush1c.2166AA mice treated with Isis No. 527134, wild-typ mice, and heterozygous mice.

Figure 4a shows RT-PCR analysis of Ush1c.2166AA mice treated with a mismatched control, Ush1c.2166AA mice treated with Isis No. 527134, heterozygous mice treated with Isis No. 527134, and wild-type mice treated with treated with Isis No. 527134.

Figure 4b shows a Western-Blot of harmonin abundance in Ush1c.2166AA mice treated with a mismatched control, Ush1c.2166AA mice treated with Isis No. 527134, and heterozygous mice treated with Isis No. 527134.

Figures 4c and 4d show immunohistochemistry analysis of whole cochleae in Ush1c.2166AA mice treated with a mismatched control, Ush1c.2166AA mice treated with Isis No. 527134, heterozygous mice treated with a mismatched control, and wild-type mice treated with treated with a mismatched control.

Figure 4e illustrates the stereocilia and a Scanning Electron Microscopy (SEM) image of the stereocilia.

Figure 4f illustrates SEM images of the stereocilia of Ush1c.2166AA mice treated with a mismatched

control, Ush1c.2166AA mice treated with Isis No. 527134, and heterozygous mice treated with Isis No. 527134.

Figure 5a shows a schematic of how Isis No. 527134 targets the cryptic splice site of the Ush1c gene having the 216A mutation.

5 Figure 5b shows RT-PCR analysis of liver cells treated with different concentrations of Isis No. 527134.

Figure 5c shows Western-Blot analysis of harmonin expression in liver cells treated with different concentrations of Isis No. 527134.

10 Figure 6 shows audiograms of broad-band noise and pure-tone stimuli at 8 kHz, 16 kHz, 32 kHz at one month of age for Ush1c.216AA mutant mice treated with Isis No. 527134, Ush1c.216AA mutant mice treated with a mismatched control, and heterozygous mice.

Figure 7 shows RT-PCR analysis of RNA isolated from cultured lymphoblasts derived from an Ush1c patient and treated with Isis No. 527134.

15 Figure 8 shows audiograms of broad-band noise and pure-tone stimuli at 8 kHz, 16 kHz, 32 kHz for Ush1c.216AA mutant mice treated with Isis No. 527134, Ush1c.216AA mutant mice treated with a mismatched control, and heterozygous mice.

Figure 8 shows audiograms of broad-band noise and pure-tone stimuli at 8 kHz, 16 kHz, 32 kHz at two months of age for Ush1c.216AA mutant mice treated with Isis No. 527134, Ush1c.216AA mutant mice treated with a mismatched control, and heterozygous mice.

20 Figure 9 shows audiograms of broad-band noise and pure-tone stimuli at 8 kHz, 16 kHz, 32 kHz at three months of age for Ush1c.216AA mutant mice treated with Isis No. 527134, Ush1c.216AA mutant mice treated with a mismatched control, and heterozygous mice.

25 Figure 10a shows audiograms of broad-band noise and pure-tone stimuli at 8 kHz, 16 kHz, 32 kHz at six months of age for Ush1c.216AA mutant mice treated with Isis No. 527134, Ush1c.216AA mutant mice treated with a mismatched control, and heterozygous mice.

Figure 10b shows the quantification of the audiograms of broad-band noise and pure-tone stimuli at 8 kHz, 16 kHz, 32 kHz at six months of age for Ush1c.216AA mutant mice treated with Isis No. 527134, Ush1c.216AA mutant mice treated with a mismatched control, and heterozygous mice.

30 Figure 10c shows the RT-PCR quantification of correct splicing at P12, P30, P120, and P180 in Ush1c.216AA mutant mice treated with Isis No. 527134.

Figure 11a shows SEM images of the apex tip, the middle apex, and the middle turn region of the organ of Corti of Ush1c.216AA mutant mice treated with Isis No. 527134, Ush1c.216AA mutant mice treated with a mismatched control, and heterozygous mice.

35 Figure 11b shows the quantitation of SEM images of the apex tip, the middle apex, and the middle turn region of the organ of Corti of Ush1c.216AA mutant mice treated with Isis No. 527134, Ush1c.216AA

mutant mice treated with a mismatched control, and heterozygous mice.

Figure 12a shows RT-PCR analysis of RNA isolated from the retina of Ush1c.216AA mice treated with Isis No. 527134.

Figure 12b shows Western Blot analysis of harmonin protein isolated from the retina of Ush1c.216AA mice treated with Isis No. 527134, Ush1c.216AA mutant mice treated with a mismatched control, and heterozygous mice.

Figure 12c shows RT-PCR analysis of RNA isolated from the retina of Ush1c.216AA mice treated with Isis No. 527134 at P12, P30, and P120.

Figure 12d shows quantification of the RT-PCR analysis of RNA isolated from the retina of Ush1c.216AA mice treated with Isis No. 527134 at P12, P30, and P120.

DETAILED DESCRIPTION OF THE INVENTION

Unless specific definitions are provided, the nomenclature used in connection with, and the procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques may be used for chemical synthesis, and chemical analysis. Certain such techniques and procedures may be found for example in "Carbohydrate Modifications in Antisense Research" Edited by Sangvi and Cook, American Chemical Society, Washington D.C., 1994; "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, Pa., 21st edition, 2005; and "Antisense Drug Technology, Principles, Strategies, and Applications" Edited by Stanley T. Crooke, CRC Press, Boca Raton, Florida; and Sambrook et al., "Molecular Cloning, A laboratory Manual," 2nd Edition, Cold Spring Harbor Laboratory Press, 1989, which are hereby incorporated by reference for any purpose. Where permitted, all patents, applications, published applications and other publications and other data referred to throughout in the disclosure are incorporated by reference herein in their entirety.

Unless otherwise indicated, the following terms have the following meanings:

As used herein, "nucleoside" means a compound comprising a nucleobase moiety and a sugar moiety. Nucleosides include, but are not limited to, naturally occurring nucleosides (as found in DNA and RNA) and modified nucleosides. Nucleosides may be linked to a phosphate moiety.

As used herein, "chemical modification" means a chemical difference in a compound when compared to a naturally occurring counterpart. In reference to an oligonucleotide, chemical modification does not include differences only in nucleobase sequence. Chemical modifications of oligonucleotides include nucleoside modifications (including sugar moiety modifications and nucleobase modifications) and internucleoside linkage modifications.

As used herein, "furanosyl" means a structure comprising a 5-membered ring comprising four carbon atoms and one oxygen atom.

As used herein, "naturally occurring sugar moiety" means a ribofuranosyl as found in naturally occurring RNA or a deoxyribofuranosyl as found in naturally occurring DNA.

As used herein, "sugar moiety" means a naturally occurring sugar moiety or a modified sugar moiety of a nucleoside.

5 As used herein, "modified sugar moiety" means a substituted sugar moiety, a bicyclic or tricyclic sugar moiety, or a sugar surrogate.

As used herein, "substituted sugar moiety" means a furanosyl comprising at least one substituent group that differs from that of a naturally occurring sugar moiety. Substituted sugar moieties include, but are not limited to furanosyls comprising substituents at the 2'-position, the 3'-position, the 5'-position and/or the
10 4'-position.

As used herein, "2'-substituted sugar moiety" means a furanosyl comprising a substituent at the 2'-position other than H or OH. Unless otherwise indicated, a 2'-substituted sugar moiety is not a bicyclic sugar moiety (i.e., the 2'-substituent of a 2'-substituted sugar moiety does not form a bridge to another atom of the furanosyl ring).

15 As used herein, "MOE" means $-OCH_2CH_2OCH_3$.

As used herein, "bicyclic sugar moiety" means a modified sugar moiety comprising a 4 to 7 membered ring (including but not limited to a furanosyl) comprising a bridge connecting two atoms of the 4 to 7 membered ring to form a second ring, resulting in a bicyclic structure. In certain embodiments, the 4 to 7 membered ring is a sugar ring. In certain embodiments the 4 to 7 membered ring is a furanosyl. In certain
20 such embodiments, the bridge connects the 2'-carbon and the 4'-carbon of the furanosyl.

As used herein the term "sugar surrogate" means a structure that does not comprise a furanosyl and that is capable of replacing the naturally occurring sugar moiety of a nucleoside, such that the resulting nucleoside is capable of (1) incorporation into an oligonucleotide and (2) hybridization to a complementary nucleoside. Such structures include rings comprising a different number of atoms than furanosyl (e.g., 4, 6,
25 or 7-membered rings); replacement of the oxygen of a furanosyl with a non-oxygen atom (e.g., carbon, sulfur, or nitrogen); or both a change in the number of atoms and a replacement of the oxygen. Such structures may also comprise substitutions corresponding to those described for substituted sugar moieties (e.g., 6-membered carbocyclic bicyclic sugar surrogates optionally comprising additional substituents). Sugar surrogates also include more complex sugar replacements (e.g., the non-ring systems of peptide nucleic acid). Sugar
30 surrogates include without limitation morpholino, modified morpholinos, cyclohexenyls and cyclohexitols.

As used herein, "nucleotide" means a nucleoside further comprising a phosphate linking group. As used herein, "linked nucleosides" may or may not be linked by phosphate linkages and thus includes, but is not limited to "linked nucleotides." As used herein, "linked nucleosides" are nucleosides that are connected in a continuous sequence (i.e. no additional nucleosides are present between those that are linked).

35 As used herein, "nucleobase" means a group of atoms that can be linked to a sugar moiety to create a

nucleoside that is capable of incorporation into an oligonucleotide, and wherein the group of atoms is capable of bonding with a complementary naturally occurring nucleobase of another oligonucleotide or nucleic acid. Nucleobases may be naturally occurring or may be modified.

As used herein, "heterocyclic base" or "heterocyclic nucleobase" means a nucleobase comprising a heterocyclic structure.

As used herein the terms, "unmodified nucleobase" or "naturally occurring nucleobase" means the naturally occurring heterocyclic nucleobases of RNA or DNA: the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) (including 5-methyl C), and uracil (U).

As used herein, "modified nucleobase" means any nucleobase that is not a naturally occurring nucleobase.

As used herein, "modified nucleoside" means a nucleoside comprising at least one chemical modification compared to naturally occurring RNA or DNA nucleosides. Modified nucleosides comprise a modified sugar moiety and/or a modified nucleobase.

As used herein, "bicyclic nucleoside" or "BNA" means a nucleoside comprising a bicyclic sugar moiety.

As used herein, "constrained ethyl nucleoside" or "cEt" means a nucleoside comprising a bicyclic sugar moiety comprising a 4'-CH(CH₃)-O-2' bridge.

As used herein, "locked nucleic acid nucleoside" or "LNA" means a nucleoside comprising a bicyclic sugar moiety comprising a 4'-CH₂-O-2' bridge.

As used herein, "2'-substituted nucleoside" means a nucleoside comprising a substituent at the 2'-position other than H or OH. Unless otherwise indicated, a 2'-substituted nucleoside is not a bicyclic nucleoside.

As used herein, "2'-deoxynucleoside" means a nucleoside comprising 2'-H furanosyl sugar moiety, as found in naturally occurring deoxyribonucleosides (DNA). In certain embodiments, a 2'-deoxynucleoside may comprise a modified nucleobase or may comprise an RNA nucleobase (e.g., uracil).

As used herein, "oligonucleotide" means a compound comprising a plurality of linked nucleosides. In certain embodiments, an oligonucleotide comprises one or more unmodified ribonucleosides (RNA) and/or unmodified deoxyribonucleosides (DNA) and/or one or more modified nucleosides.

As used herein "oligonucleoside" means an oligonucleotide in which none of the internucleoside linkages contains a phosphorus atom. As used herein, oligonucleotides include oligonucleosides.

As used herein, "modified oligonucleotide" means an oligonucleotide comprising at least one modified nucleoside and/or at least one modified internucleoside linkage.

As used herein "internucleoside linkage" means a covalent linkage between adjacent nucleosides in an oligonucleotide.

As used herein "naturally occurring internucleoside linkage" means a 3' to 5' phosphodiester linkage.

As used herein, "modified internucleoside linkage" means any internucleoside linkage other than a naturally occurring internucleoside linkage.

As used herein, "oligomeric compound" means a polymeric structure comprising two or more sub-structures. In certain embodiments, an oligomeric compound comprises an oligonucleotide. In certain
5 embodiments, an oligomeric compound comprises one or more conjugate groups and/or terminal groups. In certain embodiments, an oligomeric compound consists of an oligonucleotide.

As used herein, "terminal group" means one or more atom attached to either, or both, the 3' end or the 5' end of an oligonucleotide. In certain embodiments a terminal group is a conjugate group. In certain
10 embodiments, a terminal group comprises one or more terminal group nucleosides.

As used herein, "conjugate" means an atom or group of atoms bound to an oligonucleotide or oligomeric compound. In general, conjugate groups modify one or more properties of the compound to which they are attached, including, but not limited to pharmacodynamic, pharmacokinetic, binding, absorption, cellular distribution, cellular uptake, charge and/or clearance properties.

As used herein, "conjugate linking group" means any atom or group of atoms used to attach a
15 conjugate to an oligonucleotide or oligomeric compound.

As used herein, "antisense compound" means a compound comprising or consisting of an oligonucleotide at least a portion of which is complementary to a target nucleic acid to which it is capable of hybridizing, resulting in at least one antisense activity.

As used herein, "antisense activity" means any detectable and/or measurable change attributable to
20 the hybridization of an antisense compound to its target nucleic acid.

As used herein, "detecting" or "measuring" means that a test or assay for detecting or measuring is performed. Such detection and/or measuring may result in a value of zero. Thus, if a test for detection or measuring results in a finding of no activity (activity of zero), the step of detecting or measuring the activity has nevertheless been performed.

As used herein, "detectable and/or measureable activity" means a statistically significant activity that
25 is not zero.

As used herein, "essentially unchanged" means little or no change in a particular parameter, particularly relative to another parameter which changes much more. In certain embodiments, a parameter is essentially unchanged when it changes less than 5%. In certain embodiments, a parameter is essentially
30 unchanged if it changes less than two-fold while another parameter changes at least ten-fold. For example, in certain embodiments, an antisense activity is a change in the amount of a target nucleic acid. In certain such embodiments, the amount of a non-target nucleic acid is essentially unchanged if it changes much less than the target nucleic acid does, but the change need not be zero.

As used herein, "expression" means the process by which a gene ultimately results in a protein.
35 Expression includes, but is not limited to, transcription, post-transcriptional modification (e.g., splicing,

polyadenylation, addition of 5'-cap), and translation.

As used herein, "target nucleic acid" means a nucleic acid molecule to which an antisense compound hybridizes.

As used herein, "mRNA" means an RNA molecule that encodes a protein.

5 As used herein, "pre-mRNA" means an RNA transcript that has not been fully processed into mRNA. Pre-mRNA includes one or more intron.

As used herein, "transcript" means an RNA molecule transcribed from DNA. Transcripts include, but are not limited to mRNA, pre-mRNA, and partially processed RNA.

As used herein, "Usher Transcript" means a transcript transcribed from an Usher Gene.

10 As used herein, "Usher gene" means a gene as described in Verpy, E, Leibovici, M, Zwaenepoel, I, Liu, XZ, Gal, A, Salem, N, Mansour, A, Blanchard, S, Kobayashi, I, Keats, BJ, Slim, R, Petit, C. 2000., *A defect in harmonin, a PDZ domain-containing protein expressed in the inner ear sensory hair cells, underlies Usher syndrome type 1C*. Nat. Genet. 26:51-55 and having a sequence Accession Number ENSG00000006611, provided herein as SEQ ID NO. 1, or a variant thereof. In certain embodiments, an
15 Usher gene is at least 90% identical to Accession Number ENSG00000006611, set forth as SEQ ID NO 1. In certain embodiments, an Usher gene is at least 95% identical to Accession Number ENSG00000006611, set forth as SEQ ID NO 1. In certain embodiments, an Usher gene is 100% identical to Accession Number ENSG00000006611, set forth as SEQ ID NO 1.

As used herein, "Ush1c" means an Usher transcript transcribed from the Usher gene.

20 As used herein, "harmonin" means Ush1c or a protein encoded therefrom.

As used herein, "targeting" or "targeted to" means the association of an antisense compound to a particular target nucleic acid molecule or a particular region of a target nucleic acid molecule. An antisense compound targets a target nucleic acid if it is sufficiently complementary to the target nucleic acid to allow hybridization under physiological conditions.

25 As used herein, "nucleobase complementarity" or "complementarity" when in reference to nucleobases means a nucleobase that is capable of base pairing with another nucleobase. For example, in DNA, adenine (A) is complementary to thymine (T). For example, in RNA, adenine (A) is complementary to uracil (U). In certain embodiments, complementary nucleobase means a nucleobase of an antisense compound that is capable of base pairing with a nucleobase of its target nucleic acid. For example, if a
30 nucleobase at a certain position of an antisense compound is capable of hydrogen bonding with a nucleobase at a certain position of a target nucleic acid, then the position of hydrogen bonding between the oligonucleotide and the target nucleic acid is considered to be complementary at that nucleobase pair. Nucleobases comprising certain modifications may maintain the ability to pair with a counterpart nucleobase and thus, are still capable of nucleobase complementarity.

35 As used herein, "non-complementary" in reference to nucleobases means a pair of nucleobases that

do not form hydrogen bonds with one another.

As used herein, "complementary" in reference to oligomeric compounds (e.g., linked nucleosides, oligonucleotides, or nucleic acids) means the capacity of such oligomeric compounds or regions thereof to hybridize to another oligomeric compound or region thereof through nucleobase complementarity under stringent conditions. Complementary oligomeric compounds need not have nucleobase complementarity at each nucleoside. Rather, some mismatches are tolerated. In certain embodiments, complementary oligomeric compounds or regions are complementary at 70% of the nucleobases (70% complementary). In certain embodiments, complementary oligomeric compounds or regions are 80% complementary. In certain embodiments, complementary oligomeric compounds or regions are 90% complementary. In certain
10 embodiments, complementary oligomeric compounds or regions are 95% complementary. In certain embodiments, complementary oligomeric compounds or regions are 100% complementary.

As used herein, "hybridization" means the pairing of complementary oligomeric compounds (e.g., an antisense compound and its target nucleic acid). While not limited to a particular mechanism, the most common mechanism of pairing involves hydrogen bonding, which may be Watson-Crick, Hoogsteen or
15 reversed Hoogsteen hydrogen bonding, between complementary nucleobases.

As used herein, "specifically hybridizes" means the ability of an oligomeric compound to hybridize to one nucleic acid site with greater affinity than it hybridizes to another nucleic acid site. In certain embodiments, an antisense oligonucleotide specifically hybridizes to more than one target site.

As used herein, "percent complementarity" means the percentage of nucleobases of an oligomeric
20 compound that are complementary to an equal-length portion of a target nucleic acid. Percent complementarity is calculated by dividing the number of nucleobases of the oligomeric compound that are complementary to nucleobases at corresponding positions in the target nucleic acid by the total length of the oligomeric compound.

As used herein, "percent identity" means the number of nucleobases in a first nucleic acid that are the
25 same type (independent of chemical modification) as nucleobases at corresponding positions in a second nucleic acid, divided by the total number of nucleobases in the first nucleic acid.

As used herein, "modulation" means a change of amount or quality of a molecule, function, or activity when compared to the amount or quality of a molecule, function, or activity prior to modulation. For example, modulation includes the change, either an increase (stimulation or induction) or a decrease
30 (inhibition or reduction) in gene expression. As a further example, modulation of expression can include a change in splice site selection of pre-mRNA processing, resulting in a change in the absolute or relative amount of a particular splice-variant compared to the amount in the absence of modulation.

As used herein, "motif" means a pattern of chemical modifications in an oligomeric compound or a region thereof. Motifs may be defined by modifications at certain nucleosides and/or at certain linking
35 groups of an oligomeric compound.

As used herein, “nucleoside motif” means a pattern of nucleoside modifications in an oligomeric compound or a region thereof. The linkages of such an oligomeric compound may be modified or unmodified. Unless otherwise indicated, motifs herein describing only nucleosides are intended to be nucleoside motifs. Thus, in such instances, the linkages are not limited.

5 As used herein, “sugar motif” means a pattern of sugar modifications in an oligomeric compound or a region thereof.

As used herein, “linkage motif” means a pattern of linkage modifications in an oligomeric compound or region thereof. The nucleosides of such an oligomeric compound may be modified or unmodified. Unless otherwise indicated, motifs herein describing only linkages are intended to be linkage motifs. Thus, in such
10 instances, the nucleosides are not limited.

As used herein, “nucleobase modification motif” means a pattern of modifications to nucleobases along an oligonucleotide. Unless otherwise indicated, a nucleobase modification motif is independent of the nucleobase sequence.

As used herein, “sequence motif” means a pattern of nucleobases arranged along an oligonucleotide
15 or portion thereof. Unless otherwise indicated, a sequence motif is independent of chemical modifications and thus may have any combination of chemical modifications, including no chemical modifications.

As used herein, “type of modification” in reference to a nucleoside or a nucleoside of a “type” means the chemical modification of a nucleoside and includes modified and unmodified nucleosides. Accordingly, unless otherwise indicated, a “nucleoside having a modification of a first type” may be an unmodified
20 nucleoside.

As used herein, “differently modified” mean chemical modifications or chemical substituents that are different from one another, including absence of modifications. Thus, for example, a MOE nucleoside and an unmodified DNA nucleoside are “differently modified,” even though the DNA nucleoside is unmodified. Likewise, DNA and RNA are “differently modified,” even though both are naturally-occurring unmodified
25 nucleosides. Nucleosides that are the same but for comprising different nucleobases are not differently modified. For example, a nucleoside comprising a 2'-OMe modified sugar and an unmodified adenine nucleobase and a nucleoside comprising a 2'-OMe modified sugar and an unmodified thymine nucleobase are not differently modified.

As used herein, “the same type of modifications” refers to modifications that are the same as one
30 another, including absence of modifications. Thus, for example, two unmodified DNA nucleoside have “the same type of modification,” even though the DNA nucleoside is unmodified. Such nucleosides having the same type modification may comprise different nucleobases.

As used herein, “pharmaceutically acceptable carrier or diluent” means any substance suitable for use in administering to an animal. In certain embodiments, a pharmaceutically acceptable carrier or diluent is
35 sterile saline. In certain embodiments, such sterile saline is pharmaceutical grade saline.

As used herein, "substituent" and "substituent group," means an atom or group that replaces the atom or group of a named parent compound. For example a substituent of a modified nucleoside is any atom or group that differs from the atom or group found in a naturally occurring nucleoside (e.g., a modified 2'-substituent is any atom or group at the 2'-position of a nucleoside other than H or OH). Substituent groups can be protected or unprotected. In certain embodiments, compounds of the present invention have substituents at one or at more than one position of the parent compound. Substituents may also be further substituted with other substituent groups and may be attached directly or via a linking group such as an alkyl or hydrocarbyl group to a parent compound.

Likewise, as used herein, "substituent" in reference to a chemical functional group means an atom or group of atoms differs from the atom or a group of atoms normally present in the named functional group. In certain embodiments, a substituent replaces a hydrogen atom of the functional group (c.g., in certain embodiments, the substituent of a substituted methyl group is an atom or group other than hydrogen which replaces one of the hydrogen atoms of an unsubstituted methyl group). Unless otherwise indicated, groups amenable for use as substituents include without limitation, halogen, hydroxyl, alkyl, alkenyl, alkynyl, acyl ($-C(O)R_{aa}$), carboxyl ($-C(O)O-R_{aa}$), aliphatic groups, alicyclic groups, alkoxy, substituted oxy ($-O-R_{aa}$), aryl, aralkyl, heterocyclic radical, heteroaryl, heteroarylalkyl, amino ($-N(R_{bb})(R_{cc})$), imino ($=NR_{bb}$), amido ($-C(O)N(R_{bb})(R_{cc})$ or $-N(R_{bb})C(O)R_{aa}$), azido ($-N_3$), nitro ($-NO_2$), cyano ($-CN$), carbamido ($-OC(O)N(R_{bb})(R_{cc})$ or $-N(R_{bb})C(O)OR_{aa}$), ureido ($-N(R_{bb})C(O)N(R_{bb})(R_{cc})$), thioureido ($-N(R_{bb})C(S)N(R_{bb})(R_{cc})$), guanidiny ($-N(R_{bb})C(=NR_{bb})N(R_{bb})(R_{cc})$), amidiny ($-C(=NR_{bb})N(R_{bb})(R_{cc})$ or $-N(R_{bb})C(=NR_{bb})(R_{aa})$), thiol ($-SR_{bb}$), sulfinyl ($-S(O)R_{bb}$), sulfonyl ($-S(O)_2R_{bb}$) and sulfonamidyl ($-S(O)_2N(R_{bb})(R_{cc})$ or $-N(R_{bb})S(O)_2R_{bb}$). Wherein each R_{aa} , R_{bb} and R_{cc} is, independently, H, an optionally linked chemical functional group or a further substituent group with a preferred list including without limitation, alkyl, alkenyl, alkynyl, aliphatic, alkoxy, acyl, aryl, aralkyl, heteroaryl, alicyclic, heterocyclic and heteroarylalkyl. Selected substituents within the compounds described herein are present to a recursive degree.

As used herein, "alkyl," as used herein, means a saturated straight or branched hydrocarbon radical containing up to twenty four carbon atoms. Examples of alkyl groups include without limitation, methyl, ethyl, propyl, butyl, isopropyl, n-hexyl, octyl, decyl, dodecyl and the like. Alkyl groups typically include from 1 to about 24 carbon atoms, more typically from 1 to about 12 carbon atoms (C_1 - C_{12} alkyl) with from 1 to about 6 carbon atoms being more preferred.

As used herein, "alkenyl," means a straight or branched hydrocarbon chain radical containing up to twenty four carbon atoms and having at least one carbon-carbon double bond. Examples of alkenyl groups include without limitation, ethenyl, propenyl, butenyl, 1-methyl-2-buten-1-yl, dienes such as 1,3-butadiene and the like. Alkenyl groups typically include from 2 to about 24 carbon atoms, more typically from 2 to about 12 carbon atoms with from 2 to about 6 carbon atoms being more preferred. Alkenyl groups as used herein may optionally include one or more further substituent groups.

As used herein, "alkynyl," means a straight or branched hydrocarbon radical containing up to twenty four carbon atoms and having at least one carbon-carbon triple bond. Examples of alkynyl groups include, without limitation, ethynyl, 1-propynyl, 1-butyne, and the like. Alkynyl groups typically include from 2 to about 24 carbon atoms, more typically from 2 to about 12 carbon atoms with from 2 to about 6 carbon atoms being more preferred. Alkynyl groups as used herein may optionally include one or more further substituent groups.

As used herein, "acyl," means a radical formed by removal of a hydroxyl group from an organic acid and has the general Formula -C(O)-X where X is typically aliphatic, alicyclic or aromatic. Examples include aliphatic carbonyls, aromatic carbonyls, aliphatic sulfonyls, aromatic sulfonyls, aliphatic sulfonyls, aromatic phosphates, aliphatic phosphates and the like. Acyl groups as used herein may optionally include further substituent groups.

As used herein, "alicyclic" means a cyclic ring system wherein the ring is aliphatic. The ring system can comprise one or more rings wherein at least one ring is aliphatic. Preferred alicyclics include rings having from about 5 to about 9 carbon atoms in the ring. Alicyclic as used herein may optionally include further substituent groups.

As used herein, "aliphatic" means a straight or branched hydrocarbon radical containing up to twenty four carbon atoms wherein the saturation between any two carbon atoms is a single, double or triple bond. An aliphatic group preferably contains from 1 to about 24 carbon atoms, more typically from 1 to about 12 carbon atoms with from 1 to about 6 carbon atoms being more preferred. The straight or branched chain of an aliphatic group may be interrupted with one or more heteroatoms that include nitrogen, oxygen, sulfur and phosphorus. Such aliphatic groups interrupted by heteroatoms include without limitation, polyalkoxys, such as polyalkylene glycols, polyamines, and polyimines. Aliphatic groups as used herein may optionally include further substituent groups.

As used herein, "alkoxy" means a radical formed between an alkyl group and an oxygen atom wherein the oxygen atom is used to attach the alkoxy group to a parent molecule. Examples of alkoxy groups include without limitation, methoxy, ethoxy, propoxy, isopropoxy, *n*-butoxy, sec-butoxy, *tert*-butoxy, *n*-pentoxy, neopentoxy, *n*-hexoxy and the like. Alkoxy groups as used herein may optionally include further substituent groups.

As used herein, "aminoalkyl" means an amino substituted C₁-C₁₂ alkyl radical. The alkyl portion of the radical forms a covalent bond with a parent molecule. The amino group can be located at any position and the aminoalkyl group can be substituted with a further substituent group at the alkyl and/or amino portions.

As used herein, "aralkyl" and "arylalkyl" mean an aromatic group that is covalently linked to a C₁-C₁₂ alkyl radical. The alkyl radical portion of the resulting aralkyl (or arylalkyl) group forms a covalent bond with a parent molecule. Examples include without limitation, benzyl, phenethyl and the like. Aralkyl groups

as used herein may optionally include further substituent groups attached to the alkyl, the aryl or both groups that form the radical group.

As used herein, "aryl" and "aromatic" mean a mono- or polycyclic carbocyclic ring system radicals having one or more aromatic rings. Examples of aryl groups include without limitation, phenyl, naphthyl, tetrahydronaphthyl, indanyl, idenyl and the like. Preferred aryl ring systems have from about 5 to about 20 carbon atoms in one or more rings. Aryl groups as used herein may optionally include further substituent groups.

As used herein, "halo" and "halogen," mean an atom selected from fluorine, chlorine, bromine and iodine.

As used herein, "heteroaryl," and "heteroaromatic," mean a radical comprising a mono- or polycyclic aromatic ring, ring system or fused ring system wherein at least one of the rings is aromatic and includes one or more heteroatoms. Heteroaryl is also meant to include fused ring systems including systems where one or more of the fused rings contain no heteroatoms. Heteroaryl groups typically include one ring atom selected from sulfur, nitrogen or oxygen. Examples of heteroaryl groups include without limitation, pyridinyl, pyrazinyl, pyrimidinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, thiadiazolyl, oxadiazolyl, thiophenyl, furanyl, quinolinyl, isoquinolinyl, benzimidazolyl, benzooxazolyl, quinoxalinyl and the like. Heteroaryl radicals can be attached to a parent molecule directly or through a linking moiety such as an aliphatic group or hetero atom. Heteroaryl groups as used herein may optionally include further substituent groups.

Oligomeric Compounds

In certain embodiments, the present invention provides oligomeric compounds. In certain embodiments, such oligomeric compounds comprise oligonucleotides optionally comprising one or more conjugate and/or terminal groups. In certain embodiments, an oligomeric compound consists of an oligonucleotide. In certain embodiments, oligonucleotides comprise one or more chemical modifications. Such chemical modifications include modifications one or more nucleoside (including modifications to the sugar moiety and/or the nucleobase) and/or modifications to one or more internucleoside linkage.

Certain Sugar Moieties

In certain embodiments, oligomeric compounds of the invention comprise one or more modified nucleosides comprising a modified sugar moiety. Such oligomeric compounds comprising one or more sugar-modified nucleosides may have desirable properties, such as enhanced nuclease stability or increased binding affinity with a target nucleic acid relative to oligomeric compounds comprising only nucleosides comprising naturally occurring sugar moieties. In certain embodiments, modified sugar moieties are substituted sugar moieties. In certain embodiments, modified sugar moieties are bicyclic or tricyclic sugar moieties. In certain

embodiments, modified sugar moieties are sugar surrogates. Such sugar surrogates may comprise one or more substitutions corresponding to those of substituted sugar moieties.

In certain embodiments, modified sugar moieties are substituted sugar moieties comprising one or more substituent, including but not limited to substituents at the 2' and/or 5' positions. Examples of sugar substituents suitable for the 2'-position, include, but are not limited to: 2'-F, 2'-OCH₃ ("OMe" or "O-methyl"), and 2'-O(CH₂)₂OCH₃ ("MOE"). In certain embodiments, sugar substituents at the 2' position is selected from allyl, amino, azido, thio, O-allyl, O-C₁-C₁₀ alkyl, O-C₁-C₁₀ substituted alkyl; O-C₁-C₁₀ alkoxy; O-C₁-C₁₀ substituted alkoxy, OCF₃, O(CH₂)₂SCH₃, O(CH₂)₂-O-N(R_m)(R_n), and O-CH₂-C(=O)-N(R_m)(R_n), where each R_m and R_n is, independently, H or substituted or unsubstituted C₁-C₁₀ alkyl. Examples of sugar substituents at the 5'-position, include, but are not limited to: 5'-methyl (R or S); 5'-vinyl, and 5'-methoxy. In certain embodiments, substituted sugars comprise more than one non-bridging sugar substituent, for example, 2'-F-5'-methyl sugar moieties (*see, e.g.*, PCT International Application WO 2008/101157, for additional 5', 2'-bis substituted sugar moieties and nucleosides).

Nucleosides comprising 2'-substituted sugar moieties are referred to as 2'-substituted nucleosides. In certain embodiments, a 2'-substituted nucleoside comprises a 2'-substituent group selected from halo, allyl, amino, azido, O-C₁-C₁₀ alkoxy; O-C₁-C₁₀ substituted alkoxy, SH, CN, OCN, CF₃, OCF₃, O-alkyl, S-alkyl, N(R_m)-alkyl; O-alkenyl, S-alkenyl, or N(R_m)-alkenyl; O-alkynyl, S-alkynyl, N(R_m)-alkynyl; O-alkylenyl-O-alkyl, alkynyl, alkaryl, aralkyl, O-alkaryl, O-aralkyl, O(CH₂)₂SCH₃, O-(CH₂)₂-O-N(R_m)(R_n) or O-CH₂-C(=O)-N(R_m)(R_n), where each R_m and R_n is, independently, H, an amino protecting group or substituted or unsubstituted C₁-C₁₀ alkyl. These 2'-substituent groups can be further substituted with one or more substituent groups independently selected from hydroxyl, amino, alkoxy, carboxy, benzyl, phenyl, nitro (NO₂), thiol, thioalkoxy (S-alkyl), halogen, alkyl, aryl, alkenyl and alkynyl.

In certain embodiments, a 2'-substituted nucleoside comprises a 2'-substituent group selected from F, NH₂, N₃, OCF₃, O-CH₃, O(CH₂)₃NH₂, CH₂-CH=CH₂, O-CH₂-CH=CH₂, OCH₂CH₂OCH₃, O(CH₂)₂SCH₃, O-(CH₂)₂-O-N(R_m)(R_n), O(CH₂)₂O(CH₂)₂N(CH₃)₂, and N-substituted acetamide (O-CH₂-C(=O)-N(R_m)(R_n)) where each R_m and R_n is, independently, H, an amino protecting group or substituted or unsubstituted C₁-C₁₀ alkyl.

In certain embodiments, a 2'-substituted nucleoside comprises a sugar moiety comprising a 2'-substituent group selected from F, OCF₃, O-CH₃, OCH₂CH₂OCH₃, O(CH₂)₂SCH₃, O-(CH₂)₂-O-N(CH₃)₂, -O(CH₂)₂O(CH₂)₂N(CH₃)₂, and O-CH₂-C(=O)-N(H)CH₃.

In certain embodiments, a 2'-substituted nucleoside comprises a sugar moiety comprising a 2'-substituent group selected from F, O-CH₃, and OCH₂CH₂OCH₃.

Certain modified sugar moieties comprise a bridging sugar substituent that forms a second ring resulting in a bicyclic sugar moiety. In certain such embodiments, the bicyclic sugar moiety comprises a bridge between the 4' and the 2' furanose ring atoms. Examples of such 4' to 2' sugar substituents, include,

but are not limited to: $-[C(R_a)(R_b)]_n-$, $-[C(R_a)(R_b)]_n-O-$, $-C(R_aR_b)-N(R)-O-$ or $-C(R_aR_b)-O-N(R)-$; 4'-CH₂-2', 4'-(CH₂)₂-2', 4'-(CH₂)₃-2', 4'-(CH₂)-O-2' (LNA); 4'-(CH₂)-S-2'; 4'-(CH₂)₂-O-2' (ENA); 4'-CH(CH₃)-O-2' (cEt) and 4'-CH(CH₂OCH₃)-O-2', and analogs thereof (*see, e.g.*, U.S. Patent 7,399,845, issued on July 15, 2008); 4'-C(CH₃)(CH₃)-O-2' and analogs thereof, (*see, e.g.*, WO2009/006478, published January 8, 2009); 4'-CH₂-N(OCH₃)-2' and analogs thereof (*see, e.g.*, WO2008/150729, published December 11, 2008); 4'-CH₂-O-N(CH₃)-2' (*see, e.g.*, US2004/0171570, published September 2, 2004); 4'-CH₂-O-N(R)-2', and 4'-CH₂-N(R)-O-2', wherein each R is, independently, H, a protecting group, or C₁-C₁₂ alkyl; 4'-CH₂-N(R)-O-2', wherein R is H, C₁-C₁₂ alkyl, or a protecting group (*see*, U.S. Patent 7,427,672, issued on September 23, 2008); 4'-CH₂-C(H)(CH₃)-2' (*see, e.g.*, Chattopadhyaya, *et al.*, *J. Org. Chem.*, 2009, 74, 118-134); and 4'-CH₂-C(=CH₂)-2' and analogs thereof (*see*, published PCT International Application WO 2008/154401, published on December 8, 2008).

In certain embodiments, such 4' to 2' bridges independently comprise from 1 to 4 linked groups independently selected from $-[C(R_a)(R_b)]_n-$, $-C(R_a)=C(R_b)-$, $-C(R_a)=N-$, $-C(=NR_a)-$, $-C(=O)-$, $-C(=S)-$, $-O-$, $-Si(R_a)_2-$, $-S(=O)_x-$, and $-N(R_a)-$;

wherein:


x is 0, 1, or 2;

n is 1, 2, 3, or 4;

each R_a and R_b is, independently, H, a protecting group, hydroxyl, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₅-C₂₀ aryl, substituted C₅-C₂₀ aryl, heterocycle radical, substituted heterocycle radical, heteroaryl, substituted heteroaryl, C₅-C₇ alicyclic radical, substituted C₅-C₇ alicyclic radical, halogen, OJ₁, NJ₁J₂, SJ₁, N₃, COOJ₁, acyl (C(=O)-H), substituted acyl, CN, sulfonyl (S(=O)₂-J₁), or sulfoxyl (S(=O)-J₁); and

each J₁ and J₂ is, independently, H, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₅-C₂₀ aryl, substituted C₅-C₂₀ aryl, acyl (C(=O)-H), substituted acyl, a heterocycle radical, a substituted heterocycle radical, C₁-C₁₂ aminoalkyl, substituted C₁-C₁₂ aminoalkyl, or a protecting group.

Nucleosides comprising bicyclic sugar moieties are referred to as bicyclic nucleosides or BNAs. Bicyclic nucleosides include, but are not limited to, (A) α-L-Methyleneoxy (4'-CH₂-O-2') BNA, (B) β-D-Methyleneoxy (4'-CH₂-O-2') BNA (also referred to as locked nucleic acid or LNA), (C) Ethyleneoxy (4'-(CH₂)₂-O-2') BNA, (D) Aminooxy (4'-CH₂-O-N(R)-2') BNA, (E) Oxyamino (4'-CH₂-N(R)-O-2') BNA, (F) Methyl(methyleneoxy) (4'-CH(CH₃)-O-2') BNA (also referred to as constrained ethyl or cEt), (G) methylene-thio (4'-CH₂-S-2') BNA, (H) methylene-amino (4'-CH₂-N(R)-2') BNA, (I) methyl carbocyclic (4'-CH₂-CH(CH₃)-2') BNA, and (J) propylene carbocyclic (4'-(CH₂)₃-2') BNA as depicted below.



5 wherein Bx is a nucleobase moiety and R is, independently, H, a protecting group, or C₁-C₁₂ alkyl.

Additional bicyclic sugar moieties are known in the art, for example: Singh et al., *Chem. Commun.*, 1998, 4, 455-456; Koshkin et al., *Tetrahedron*, 1998, 54, 3607-3630; Wahlestedt et al., *Proc. Natl. Acad. Sci. U. S. A.*, 2000, 97, 5633-5638; Kumar et al., *Bioorg. Med. Chem. Lett.*, 1998, 8, 2219-2222; Singh et al., *J. Org. Chem.*, 1998, 63, 10035-10039; Srivastava et al., *J. Am. Chem. Soc.*, 129(26) 8362-8379 (Jul. 4, 2007);
 10 Elayadi et al., *Curr. Opinion Invens. Drugs*, 2001, 2, 558-561; Braasch et al., *Chem. Biol.*, 2001, 8, 1-7; Orum et al., *Curr. Opinion Mol. Ther.*, 2001, 3, 239-243; U.S. Patent Nos. 7,053,207, 6,268,490, 6,770,748, 6,794,499, 7,034,133, 6,525,191, 6,670,461, and 7,399,845; WO 2004/106356, WO 1994/14226, WO 2005/021570, and WO 2007/134181; U.S. Patent Publication Nos. US2004/0171570, US2007/0287831, and US2008/0039618; U.S. Patent Serial Nos. 12/129,154, 60/989,574, 61/026,995, 61/026,998, 61/056,564,
 15 61/086,231, 61/097,787, and 61/099,844; and PCT International Applications Nos. PCT/US2008/064591, PCT/US2008/066154, and PCT/US2008/068922.


In certain embodiments, bicyclic sugar moieties and nucleosides incorporating such bicyclic sugar moieties are further defined by isomeric configuration. For example, a nucleoside comprising a 4'-2' methylene-oxy bridge, may be in the α -L configuration or in the β -D configuration. Previously, α -L-

methyleneoxy (4'-CH₂-O-2') bicyclic nucleosides have been incorporated into antisense oligonucleotides that showed antisense activity (Frieden *et al.*, *Nucleic Acids Research*, 2003, 21, 6365-6372).

In certain embodiments, substituted sugar moieties comprise one or more non-bridging sugar substituent and one or more bridging sugar substituent (e.g., 5'-substituted and 4'-2' bridged sugars). (*see*,
 5 PCT International Application WO 2007/134181, published on 11/22/07, wherein LNA is substituted with, for example, a 5'-methyl or a 5'-vinyl group).

In certain embodiments, modified sugar moieties are sugar surrogates. In certain such embodiments, the oxygen atom of the naturally occurring sugar is substituted, e.g., with a sulfur, carbon or nitrogen atom. In certain such embodiments, such modified sugar moiety also comprises bridging and/or non-bridging
 10 substituents as described above. For example, certain sugar surrogates comprise a 4'-sulfur atom and a substitution at the 2'-position (*see*, e.g., published U.S. Patent Application US2005/0130923, published on June 16, 2005) and/or the 5' position. By way of additional example, carbocyclic bicyclic nucleosides having a 4'-2' bridge have been described (*see, e.g.*, Freier *et al.*, *Nucleic Acids Research*, 1997, 25(22), 4429-4443 and Albaek *et al.*, *J. Org. Chem.*, 2006, 71, 7731-7740).

In certain embodiments, sugar surrogates comprise rings having other than 5-atoms. For example, in certain embodiments, a sugar surrogate comprises a six-membered tetrahydropyran. Such tetrahydropyrans may be further modified or substituted. Nucleosides comprising such modified tetrahydropyrans include, but are not limited to, hexitol nucleic acid (HNA), anitol nucleic acid (ANA), manitol nucleic acid (MNA) (*see*
 15 Leumann, *CJ. Bioorg. & Med. Chem.* (2002) 10:841-854), fluoro HNA (F-HNA), and those compounds having Formula VII:



VII

wherein independently for each of said at least one tetrahydropyran nucleoside analog of Formula VII:

Bx is a nucleobase moiety;

T₃ and T₄ are each, independently, an internucleoside linking group linking the tetrahydropyran nucleoside analog to the antisense compound or one of T₃ and T₄ is an internucleoside linking group linking the tetrahydropyran nucleoside analog to the antisense compound and the other of T₃ and T₄ is H, a hydroxyl
 25 protecting group, a linked conjugate group, or a 5' or 3'-terminal group;


q₁, q₂, q₃, q₄, q₅, q₆ and q₇ are each, independently, H, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C₂-C₆ alkenyl, substituted C₂-C₆ alkenyl, C₂-C₆ alkynyl, or substituted C₂-C₆ alkynyl; and

each of R_1 and R_2 is independently selected from among: hydrogen, halogen, substituted or unsubstituted alkoxy, NJ_1J_2 , SJ_1 , N_3 , $OC(=X)J_1$, $OC(=X)NJ_1J_2$, $NJ_3C(=X)NJ_1J_2$, and CN , wherein X is O , S or NJ_1 , and each J_1 , J_2 , and J_3 is, independently, H or C_1 - C_6 alkyl.

In certain embodiments, the modified THP nucleosides of Formula VII are provided wherein q_1 , q_2 , q_3 , q_4 , q_5 , q_6 and q_7 are each H . In certain embodiments, at least one of q_1 , q_2 , q_3 , q_4 , q_5 , q_6 and q_7 is other than H . In certain embodiments, at least one of q_1 , q_2 , q_3 , q_4 , q_5 , q_6 and q_7 is methyl. In certain embodiments, THP nucleosides of Formula VII are provided wherein one of R_1 and R_2 is F . In certain embodiments, R_1 is fluoro and R_2 is H , R_1 is methoxy and R_2 is H , and R_1 is methoxyethoxy and R_2 is H .

Many other bicyclic and tricyclic sugar and sugar surrogate ring systems are known in the art that can be used to modify nucleosides (*see, e.g.*, review article: Leumann, J. C, *Bioorganic & Medicinal Chemistry*, 2002, 10, 841-854).

In certain embodiments, sugar surrogates comprise rings having more than 5 atoms and more than one heteroatom. For example nucleosides comprising morpholino sugar moieties and their use in oligomeric compounds has been reported (*see for example*: Braasch et al., *Biochemistry*, 2002, 41, 4503-4510; and U.S. Patents 5,698,685; 5,166,315; 5,185,444; and 5,034,506). As used here, the term "morpholino" means a sugar surrogate having the following structure:



In certain embodiments, morpholinos may be modified, for example by adding or altering various substituent groups from the above morpholino structure. Such sugar surrogates are referred to herein as "modified morpholinos."

Combinations of modifications are also provided without limitation, such as 2'-F-5'-methyl substituted nucleosides (*see* PCT International Application WO 2008/101157 Published on 8/21/08 for other disclosed 5', 2'-bis substituted nucleosides) and replacement of the ribosyl ring oxygen atom with S and further substitution at the 2'-position (*see* published U.S. Patent Application US2005-0130923, published on June 16, 2005) or alternatively 5'-substitution of a bicyclic nucleic acid (*see* PCT International Application WO 2007/134181, published on 11/22/07 wherein a 4'- CH_2 -O-2' bicyclic nucleoside is further substituted at the 5' position with a 5'-methyl or a 5'-vinyl group). The synthesis and preparation of carbocyclic bicyclic nucleosides along with their oligomerization and biochemical studies have also been described (*see, e.g.*, Srivastava et al., *J. Am. Chem. Soc.* 2007, 129(26), 8362-8379).

Certain Nucleobases

In certain embodiments, nucleosides of the present invention comprise one or more unmodified nucleobases. In certain embodiments, nucleosides of the present invention comprise one or more modified nucleobases.

In certain embodiments, modified nucleobases are selected from: universal bases, hydrophobic bases, promiscuous bases, size-expanded bases, and fluorinated bases as defined herein. 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil; 5-propynylcytosine; 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl ($-C\equiv C-CH_3$) uracil and cytosine and other alkynyl derivatives of pyrimidine bases, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 2-F-adenine, 2-amino-adenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine, 3-deazaguanine and 3-deazaadenine, universal bases, hydrophobic bases, promiscuous bases, size-expanded bases, and fluorinated bases as defined herein. Further modified nucleobases include tricyclic pyrimidines such as phenoxazine cytidine([5,4-b][1,4]benzoxazin-2(3H)-one), phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one), G-clamps such as a substituted phenoxazine cytidine (e.g. 9-(2-aminoethoxy)-H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), carbazole cytidine (2H-pyrimido[4,5-b]indol-2-one), pyridoindole cytidine (H-pyrido[3',2':4,5]pyrrolo[2,3-d]pyrimidin-2-one). Modified nucleobases may also include those in which the purine or pyrimidine base is replaced with other heterocycles, for example 7-deaza-adenine, 7-deazaguanosine, 2-aminopyridine and 2-pyridone. Further nucleobases include those disclosed in United States Patent No. 3,687,808, those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, Kroschwitz, J.I., Ed., John Wiley & Sons, 1990, 858-859; those disclosed by Englisch *et al.*, *Angewandte Chemie*, International Edition, 1991, 30, 613; and those disclosed by Sanghvi, Y.S., Chapter 15, *Antisense Research and Applications*, Crooke, S.T. and Lebleu, B., Eds., CRC Press, 1993, 273-288.

Representative United States patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include without limitation, U.S. 3,687,808; 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121; 5,596,091; 5,614,617; 5,645,985; 5,681,941; 5,750,692; 5,763,588; 5,830,653 and 6,005,096, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

Certain Internucleoside Linkages

In certain embodiments, the present invention provides oligomeric compounds comprising linked nucleosides. In such embodiments, nucleosides may be linked together using any internucleoside linkage. The two main classes of internucleoside linking groups are defined by the presence or absence of a phosphorus atom. Representative phosphorus containing internucleoside linkages include, but are not limited to, phosphodiester (P=O), phosphotriesters, methylphosphonates, phosphoramidate, and phosphorothioates (P=S). Representative non-phosphorus containing internucleoside linking groups include, but are not limited to, methylenemethylimino (-CH₂-N(CH₃)-O-CH₂-), thiodiester (-O-C(O)-S-), thionocarbamate (-O-C(O)(NH)-S-); siloxane (-O-Si(H)₂-O-); and N,N'-dimethylhydrazine (-CH₂-N(CH₃)-N(CH₃)-). Modified linkages, compared to natural phosphodiester linkages, can be used to alter, typically increase, nuclease resistance of the oligomeric compound. In certain embodiments, internucleoside linkages having a chiral atom can be prepared as a racemic mixture, or as separate enantiomers. Representative chiral linkages include, but are not limited to, alkylphosphonates and phosphorothioates. Methods of preparation of phosphorous-containing and non-phosphorous-containing internucleoside linkages are well known to those skilled in the art.

The oligonucleotides described herein contain one or more asymmetric centers and thus give rise to enantiomers, diastereomers, and other stereoisomeric configurations that may be defined, in terms of absolute stereochemistry, as (R) or (S), α or β such as for sugar anomers, or as (D) or (L) such as for amino acids etc. Included in the antisense compounds provided herein are all such possible isomers, as well as their racemic and optically pure forms.

Neutral internucleoside linkages include without limitation, phosphotriesters, methylphosphonates, MMI (3'-CH₂-N(CH₃)-O-5'), amide-3 (3'-CH₂-C(=O)-N(H)-5'), amide-4 (3'-CH₂-N(H)-C(=O)-5'), formacetal (3'-O-CH₂-O-5'), and thioformacetal (3'-S-CH₂-O-5'). Further neutral internucleoside linkages include nonionic linkages comprising siloxane (dialkylsiloxane), carboxylate ester, carboxamide, sulfide, sulfonate ester and amides (See for example: *Carbohydrate Modifications in Antisense Research*; Y.S. Sanghvi and P.D. Cook, Eds., ACS Symposium Series 580; Chapters 3 and 4, 40-65). Further neutral internucleoside linkages include nonionic linkages comprising mixed N, O, S and CH₂ component parts.

Certain Motifs

In certain embodiments, the present invention provides oligomeric compounds comprising oligonucleotides. In certain embodiments, such oligonucleotides comprise one or more chemical modification. In certain embodiments, chemically modified oligonucleotides comprise one or more modified nucleosides. In certain embodiments, chemically modified oligonucleotides comprise one or more modified nucleosides comprising modified sugars. In certain embodiments, chemically modified oligonucleotides comprise one or more modified nucleosides comprising one or more modified nucleobases. In certain

embodiments, chemically modified oligonucleotides comprise one or more modified internucleoside linkages. In certain embodiments, the chemical modifications (sugar modifications, nucleobase modifications, and/or linkage modifications) define a pattern or motif. In certain embodiments, the patterns of chemical modifications of sugar moieties, internucleoside linkages, and nucleobases are each independent of one another. Thus, an oligonucleotide may be described by its sugar modification motif, internucleoside linkage motif and/or nucleobase modification motif (as used herein, nucleobase modification motif describes the chemical modifications to the nucleobases independent of the sequence of nucleobases).

Certain sugar motifs

In certain embodiments, oligonucleotides comprise one or more type of modified sugar moieties and/or naturally occurring sugar moieties arranged along an oligonucleotide or region thereof in a defined pattern or sugar modification motif. Such motifs may include any of the sugar modifications discussed herein and/or other known sugar modifications.

In certain embodiments, the oligonucleotides comprise or consist of a region having a gapmer sugar modification motif, which comprises two external regions or “wings” and an internal region or “gap.” The three regions of a gapmer motif (the 5'-wing, the gap, and the 3'-wing) form a contiguous sequence of nucleosides wherein at least some of the sugar moieties of the nucleosides of each of the wings differ from at least some of the sugar moieties of the nucleosides of the gap. Specifically, at least the sugar moieties of the nucleosides of each wing that are closest to the gap (the 3'-most nucleoside of the 5'-wing and the 5'-most nucleoside of the 3'-wing) differ from the sugar moiety of the neighboring gap nucleosides, thus defining the boundary between the wings and the gap. In certain embodiments, the sugar moieties within the gap are the same as one another. In certain embodiments, the gap includes one or more nucleoside having a sugar moiety that differs from the sugar moiety of one or more other nucleosides of the gap. In certain embodiments, the sugar modification motifs of the two wings are the same as one another (symmetric gapmer). In certain embodiments, the sugar modification motifs of the 5'-wing differs from the sugar modification motif of the 3'-wing (asymmetric gapmer). In certain embodiments, oligonucleotides comprise 2'-MOE modified nucleosides in the wings and 2'-F modified nucleosides in the gap.

In certain embodiments, oligonucleotides are fully modified. In certain such embodiments, oligonucleotides are uniformly modified. In certain embodiments, oligonucleotides are uniform 2'-MOE. In certain embodiments, oligonucleotides are uniform 2'-F. In certain embodiments, oligonucleotides are uniform morpholino. In certain embodiments, oligonucleotides are uniform BNA. In certain embodiments, oligonucleotides are uniform LNA. In certain embodiments, oligonucleotides are uniform cEt.

In certain embodiments, oligonucleotides comprise a uniformly modified region and additional nucleosides that are unmodified or differently modified. In certain embodiments, the uniformly modified region is at least 5, 10, 15, or 20 nucleosides in length. In certain embodiments, the uniform region is a 2'-

MOE region. In certain embodiments, the uniform region is a 2'-F region. In certain embodiments, the uniform region is a morpholino region. In certain embodiments, the uniform region is a BNA region. In certain embodiments, the uniform region is a LNA region. In certain embodiments, the uniform region is a cEt region.

In certain embodiments, the oligonucleotide does not comprise more than 4 contiguous unmodified 2'-deoxynucleosides. In certain circumstances, antisense oligonucleotides comprising more than 4 contiguous 2'-deoxynucleosides activate RNase H, resulting in cleavage of the target RNA. In certain embodiments, such cleavage is avoided by not having more than 4 contiguous 2'-deoxynucleosides, for example, where alteration of splicing and not cleavage of a target RNA is desired.

Certain Internucleoside Linkage Motifs

In certain embodiments, oligonucleotides comprise modified internucleoside linkages arranged along the oligonucleotide or region thereof in a defined pattern or modified internucleoside linkage motif. In certain embodiments, internucleoside linkages are arranged in a gapped motif, as described above for sugar modification motif. In such embodiments, the internucleoside linkages in each of two wing regions are different from the internucleoside linkages in the gap region. In certain embodiments the internucleoside linkages in the wings are phosphodiester and the internucleoside linkages in the gap are phosphorothioate. The sugar modification motif is independently selected, so such oligonucleotides having a gapped internucleoside linkage motif may or may not have a gapped sugar modification motif and if it does have a gapped sugar motif, the wing and gap lengths may or may not be the same.

In certain embodiments, oligonucleotides comprise a region having an alternating internucleoside linkage motif. In certain embodiments, oligonucleotides of the present invention comprise a region of uniformly modified internucleoside linkages. In certain such embodiments, the oligonucleotide comprises a region that is uniformly linked by phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide is uniformly linked by phosphorothioate. In certain embodiments, each internucleoside linkage of the oligonucleotide is selected from phosphodiester and phosphorothioate. In certain embodiments, each internucleoside linkage of the oligonucleotide is selected from phosphodiester and phosphorothioate and at least one internucleoside linkage is phosphorothioate.

In certain embodiments, the oligonucleotide comprises at least 6 phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least 8 phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least 10 phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least one block of at least 6 consecutive phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least one block of at least 8 consecutive phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least one block of at least 10 consecutive phosphorothioate internucleoside

linkages. In certain embodiments, the oligonucleotide comprises at least one block of at least one 12 consecutive phosphorothioate internucleoside linkages. In certain such embodiments, at least one such block is located at the 3' end of the oligonucleotide. In certain such embodiments, at least one such block is located within 3 nucleosides of the 3' end of the oligonucleotide.

5

Certain Nucleobase Modification Motifs

In certain embodiments, oligonucleotides comprise chemical modifications to nucleobases arranged along the oligonucleotide or region thereof in a defined pattern or nucleobases modification motif. In certain such embodiments, nucleobase modifications are arranged in a gapped motif. In certain embodiments, nucleobase modifications are arranged in an alternating motif. In certain embodiments, each nucleobase is modified. In certain embodiments, none of the nucleobases is chemically modified.

In certain embodiments, oligonucleotides comprise a block of modified nucleobases. In certain such embodiments, the block is at the 3'-end of the oligonucleotide. In certain embodiments the block is within 3 nucleotides of the 3'-end of the oligonucleotide. In certain such embodiments, the block is at the 5'-end of the oligonucleotide. In certain embodiments the block is within 3 nucleotides of the 5'-end of the oligonucleotide.

In certain embodiments, nucleobase modifications are a function of the natural base at a particular position of an oligonucleotide. For example, in certain embodiments each purine or each pyrimidine in an oligonucleotide is modified. In certain embodiments, each adenine is modified. In certain embodiments, each guanine is modified. In certain embodiments, each thymine is modified. In certain embodiments, each cytosine is modified. In certain embodiments, each uracil is modified.

In certain embodiments, some, all, or none of the cytosine moieties in an oligonucleotide are 5-methyl cytosine moieties. Herein, 5-methyl cytosine is not a "modified nucleobase." Accordingly, unless otherwise indicated, unmodified nucleobases include both cytosine residues having a 5-methyl and those lacking a 5 methyl. In certain embodiments, the methylation state of all or some cytosine nucleobases is specified.

Certain Overall Lengths

In certain embodiments, the present invention provides oligomeric compounds including oligonucleotides of any of a variety of ranges of lengths. In certain embodiments, the invention provides oligomeric compounds or oligonucleotides consisting of X to Y linked nucleosides, where X represents the fewest number of nucleosides in the range and Y represents the largest number of nucleosides in the range. In certain such embodiments, X and Y are each independently selected from 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, and 50; provided that $X \leq Y$. For example, in certain embodiments, the invention provides

oligomeric compounds which comprise oligonucleotides consisting of 8 to 9, 8 to 10, 8 to 11, 8 to 12, 8 to 13, 8 to 14, 8 to 15, 8 to 16, 8 to 17, 8 to 18, 8 to 19, 8 to 20, 8 to 21, 8 to 22, 8 to 23, 8 to 24, 8 to 25, 8 to 26, 8 to 27, 8 to 28, 8 to 29, 8 to 30, 9 to 10, 9 to 11, 9 to 12, 9 to 13, 9 to 14, 9 to 15, 9 to 16, 9 to 17, 9 to 18, 9 to 19, 9 to 20, 9 to 21, 9 to 22, 9 to 23, 9 to 24, 9 to 25, 9 to 26, 9 to 27, 9 to 28, 9 to 29, 9 to 30, 10 to 11, 10 to 12, 10 to 13, 10 to 14, 10 to 15, 10 to 16, 10 to 17, 10 to 18, 10 to 19, 10 to 20, 10 to 21, 10 to 22, 10 to 23, 10 to 24, 10 to 25, 10 to 26, 10 to 27, 10 to 28, 10 to 29, 10 to 30, 11 to 12, 11 to 13, 11 to 14, 11 to 15, 11 to 16, 11 to 17, 11 to 18, 11 to 19, 11 to 20, 11 to 21, 11 to 22, 11 to 23, 11 to 24, 11 to 25, 11 to 26, 11 to 27, 11 to 28, 11 to 29, 11 to 30, 12 to 13, 12 to 14, 12 to 15, 12 to 16, 12 to 17, 12 to 18, 12 to 19, 12 to 20, 12 to 21, 12 to 22, 12 to 23, 12 to 24, 12 to 25, 12 to 26, 12 to 27, 12 to 28, 12 to 29, 12 to 30, 13 to 14, 13 to 15, 13 to 16, 13 to 17, 13 to 18, 13 to 19, 13 to 20, 13 to 21, 13 to 22, 13 to 23, 13 to 24, 13 to 25, 13 to 26, 13 to 27, 13 to 28, 13 to 29, 13 to 30, 14 to 15, 14 to 16, 14 to 17, 14 to 18, 14 to 19, 14 to 20, 14 to 21, 14 to 22, 14 to 23, 14 to 24, 14 to 25, 14 to 26, 14 to 27, 14 to 28, 14 to 29, 14 to 30, 15 to 16, 15 to 17, 15 to 18, 15 to 19, 15 to 20, 15 to 21, 15 to 22, 15 to 23, 15 to 24, 15 to 25, 15 to 26, 15 to 27, 15 to 28, 15 to 29, 15 to 30, 16 to 17, 16 to 18, 16 to 19, 16 to 20, 16 to 21, 16 to 22, 16 to 23, 16 to 24, 16 to 25, 16 to 26, 16 to 27, 16 to 28, 16 to 29, 16 to 30, 17 to 18, 17 to 19, 17 to 20, 17 to 21, 17 to 22, 17 to 23, 17 to 24, 17 to 25, 17 to 26, 17 to 27, 17 to 28, 17 to 29, 17 to 30, 18 to 19, 18 to 20, 18 to 21, 18 to 22, 18 to 23, 18 to 24, 18 to 25, 18 to 26, 18 to 27, 18 to 28, 18 to 29, 18 to 30, 19 to 20, 19 to 21, 19 to 22, 19 to 23, 19 to 24, 19 to 25, 19 to 26, 19 to 27, 19 to 28, 19 to 29, 19 to 30, 20 to 21, 20 to 22, 20 to 23, 20 to 24, 20 to 25, 20 to 26, 20 to 27, 20 to 28, 20 to 29, 20 to 30, 21 to 22, 21 to 23, 21 to 24, 21 to 25, 21 to 26, 21 to 27, 21 to 28, 21 to 29, 21 to 30, 22 to 23, 22 to 24, 22 to 25, 22 to 26, 22 to 27, 22 to 28, 22 to 29, 22 to 30, 23 to 24, 23 to 25, 23 to 26, 23 to 27, 23 to 28, 23 to 29, 23 to 30, 24 to 25, 24 to 26, 24 to 27, 24 to 28, 24 to 29, 24 to 30, 25 to 26, 25 to 27, 25 to 28, 25 to 29, 25 to 30, 26 to 27, 26 to 28, 26 to 29, 26 to 30, 27 to 28, 27 to 29, 27 to 30, 28 to 29, 28 to 30, or 29 to 30 linked nucleosides. In embodiments where the number of nucleosides of an oligomeric compound or oligonucleotide is limited, whether to a range or to a specific number, the oligomeric compound or oligonucleotide may, nonetheless further comprise additional other substituents. For example, an oligonucleotide comprising 8-30 nucleosides excludes oligonucleotides having 31 nucleosides, but, unless otherwise indicated, such an oligonucleotide may further comprise, for example one or more conjugates, terminal groups, or other substituents. In certain embodiments, a gapmer oligonucleotide has any of the above lengths.

One of skill in the art will appreciate that certain lengths may not be possible for certain motifs. For example: a gapmer having a 5'-wing region consisting of four nucleotides, a gap consisting of at least six nucleotides, and a 3'-wing region consisting of three nucleotides cannot have an overall length less than 13 nucleotides. Thus, one would understand that the lower length limit is 13 and that the limit of 10 in "10-20" has no effect in that embodiment.

Further, where an oligonucleotide is described by an overall length range and by regions having specified lengths, and where the sum of specified lengths of the regions is less than the upper limit of the overall length range, the oligonucleotide may have additional nucleosides, beyond those of the specified regions, provided that the total number of nucleosides does not exceed the upper limit of the overall length range. For example, an oligonucleotide consisting of 20-25 linked nucleosides comprising a 5'-wing consisting of 5 linked nucleosides; a 3'-wing consisting of 5 linked nucleosides and a central gap consisting of 10 linked nucleosides ($5+5+10=20$) may have up to 5 nucleosides that are not part of the 5'-wing, the 3'-wing, or the gap (before reaching the overall length limitation of 25). Such additional nucleosides may be 5' of the 5'-wing and/or 3' of the 3' wing.

Certain Oligonucleotides

In certain embodiments, oligonucleotides of the present invention are characterized by their sugar motif, internucleoside linkage motif, nucleobase modification motif and overall length. In certain embodiments, such parameters are each independent of one another. Thus, each internucleoside linkage of an oligonucleotide having a gapmer sugar motif may be modified or unmodified and may or may not follow the gapmer modification pattern of the sugar modifications. Thus, the internucleoside linkages within the wing regions of a sugar-gapmer may be the same or different from one another and may be the same or different from the internucleoside linkages of the gap region. Likewise, such sugar-gapmer oligonucleotides may comprise one or more modified nucleobase independent of the gapmer pattern of the sugar modifications. Herein if a description of an oligonucleotide or oligomeric compound is silent with respect to one or more parameter, such parameter is not limited. Thus, an oligomeric compound described only as having a gapmer sugar motif without further description may have any length, internucleoside linkage motif, and nucleobase modification motif. Unless otherwise indicated, all chemical modifications are independent of nucleobase sequence.

Certain Conjugate Groups

In certain embodiments, oligomeric compounds are modified by attachment of one or more conjugate groups. In general, conjugate groups modify one or more properties of the attached oligomeric compound including but not limited to pharmacodynamics, pharmacokinetics, stability, binding, absorption, cellular distribution, cellular uptake, charge and clearance. Conjugate groups are routinely used in the chemical arts and are linked directly or via an optional conjugate linking moiety or conjugate linking group to a parent compound such as an oligomeric compound, such as an oligonucleotide. Conjugate groups includes without limitation, intercalators, reporter molecules, polyamines, polyamides, polyethylene glycols, thioethers, polyethers, cholesterol, thiocholesterol, cholic acid moieties, folate, lipids, phospholipids, biotin, phenazine, phenanthridine, anthraquinone, adamantane, acridine, fluoresceins, rhodamines, coumarins and

dyes. Certain conjugate groups have been described previously, for example: cholesterol moiety (Letsinger et al., Proc. Natl. Acad. Sci. USA, 1989, 86, 6553-6556), cholic acid (Manoharan et al., Bioorg. Med. Chem. Lett., 1994, 4, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., Ann. N.Y. Acad. Sci., 1992, 660, 306-309; Manoharan et al., Bioorg. Med. Chem. Lett., 1993, 3, 2765-2770), a thiocholesterol
5 (Oberhauser et al., Nucl. Acids Res., 1992, 20, 533-538), an aliphatic chain, e.g., do-decan-diol or undecyl residues (Saison-Behmoaras et al., EMBO J., 1991, 10, 1111-1118; Kabanov et al., FEBS Lett., 1990, 259, 327-330; Svinarchuk et al., Biochimie, 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethyl-ammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., Tetrahedron Lett., 1995, 36, 3651-3654; Shea et al., Nucl. Acids Res., 1990, 18, 3777-3783), a polyamine or a polyethylene
10 glycol chain (Manoharan et al., Nucleosides & Nucleotides, 1995, 14, 969-973), or adamantane acetic acid (Manoharan et al., Tetrahedron Lett., 1995, 36, 3651-3654), a palmityl moiety (Mishra et al., Biochim. Biophys. Acta, 1995, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., J. Pharmacol. Exp. Ther., 1996, 277, 923-937).

In certain embodiments, a conjugate group comprises an active drug substance, for example,
15 aspirin, warfarin, phenylbutazone, ibuprofen, suprofen, fen-bufen, ketoprofen, (*S*)-(+)-pranoprofen, carprofen, dansylsarcosine, 2,3,5-triiodobenzoic acid, flufenamic acid, folinic acid, a benzothiadiazide, chlorothiazide, a diazepine, indo-methicin, a barbiturate, a cephalosporin, a sulfa drug, an antidiabetic, an antibacterial or an antibiotic.

In certain embodiments, conjugate groups are directly attached to oligonucleotides in oligomeric
20 compounds. In certain embodiments, conjugate groups are attached to oligonucleotides by a conjugate linking group. In certain such embodiments, conjugate linking groups, including, but not limited to, bifunctional linking moieties such as those known in the art are amenable to the compounds provided herein. Conjugate linking groups are useful for attachment of conjugate groups, such as chemical stabilizing groups, functional groups, reporter groups and other groups to selective sites in a parent compound such as for
25 example an oligomeric compound. In general a bifunctional linking moiety comprises a hydrocarbyl moiety having two functional groups. One of the functional groups is selected to bind to a parent molecule or compound of interest and the other is selected to bind essentially any selected group such as chemical functional group or a conjugate group. In some embodiments, the conjugate linker comprises a chain structure or an oligomer of repeating units such as ethylene glycol or amino acid units. Examples of
30 functional groups that are routinely used in a bifunctional linking moiety include, but are not limited to, electrophiles for reacting with nucleophilic groups and nucleophiles for reacting with electrophilic groups. In some embodiments, bifunctional linking moieties include amino, hydroxyl, carboxylic acid, thiol, unsaturations (e.g., double or triple bonds), and the like.

Some nonlimiting examples of conjugate linking moieties include pyrrolidine, 8-amino-3,6-
35 dioxaoctanoic acid (ADO), succinimidyl 4-(*N*-maleimidomethyl) cyclohexane-1-carboxylate (SMCC) and 6-

aminohexanoic acid (AHEX or AHA). Other linking groups include, but are not limited to, substituted C₁-C₁₀ alkyl, substituted or unsubstituted C₂-C₁₀ alkenyl or substituted or unsubstituted C₂-C₁₀ alkynyl, wherein a nonlimiting list of preferred substituent groups includes hydroxyl, amino, alkoxy, carboxy, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl and alkynyl.

5 Conjugate groups may be attached to either or both ends of an oligonucleotide (terminal conjugate groups) and/or at any internal position.

In certain embodiments, conjugate groups are at the 3'-end of an oligonucleotide of an oligomeric compound. In certain embodiments, conjugate groups are near the 3'-end. In certain embodiments, conjugates are attached at the 3' end of an oligomeric compound, but before one or more terminal group
10 nucleosides. In certain embodiments, conjugate groups are placed within a terminal group. In certain embodiments, the present invention provides oligomeric compounds. In certain embodiments, oligomeric compounds comprise an oligonucleotide. In certain embodiments, an oligomeric compound comprises an oligonucleotide and one or more conjugate and/or terminal groups. Such conjugate and/or terminal groups may be added to oligonucleotides having any of the chemical motifs discussed above. Thus,
15 for example, an oligomeric compound comprising an oligonucleotide having region of alternating nucleosides may comprise a terminal group.

Antisense Compounds

In certain embodiments, oligomeric compounds of the present invention are antisense compounds.
20 Such antisense compounds are capable of hybridizing to a target nucleic acid, resulting in at least one antisense activity. In certain embodiments, antisense compounds specifically hybridize to one or more target nucleic acid. In certain embodiments, a specifically hybridizing antisense compound has a nucleobase sequence comprising a region having sufficient complementarity to a target nucleic acid to allow hybridization and result in antisense activity and insufficient complementarity to any non-target so as to avoid
25 non-specific hybridization to any non-target nucleic acid sequences under conditions in which specific hybridization is desired (e.g., under physiological conditions for *in vivo* or therapeutic uses, and under conditions in which assays are performed in the case of *in vitro* assays).

In certain embodiments, the present invention provides antisense compounds comprising oligonucleotides that are fully complementary to the target nucleic acid over the entire length of the
30 oligonucleotide. In certain embodiments, oligonucleotides are 99% complementary to the target nucleic acid. In certain embodiments, oligonucleotides are 95% complementary to the target nucleic acid. In certain embodiments, such oligonucleotides are 90% complementary to the target nucleic acid.

In certain embodiments, such oligonucleotides are 85% complementary to the target nucleic acid. In certain embodiments, such oligonucleotides are 80% complementary to the target nucleic acid. In certain
35 embodiments, an antisense compound comprises a region that is fully complementary to a target nucleic acid

and is at least 80% complementary to the target nucleic acid over the entire length of the oligonucleotide. In certain such embodiments, the region of full complementarity is from 6 to 14 nucleobases in length.

Certain Target Nucleic Acids and Mechanisms

In certain embodiments, antisense compounds comprise or consist of an oligonucleotide comprising a region that is complementary to a target nucleic acid. In certain embodiments, the target nucleic acid is an endogenous RNA molecule. In certain embodiments, the target nucleic acid is a pre-mRNA. In certain embodiments, the target nucleic acid is an Usher transcript. In certain embodiments, the target RNA is an Usher pre-mRNA.

In certain embodiments, an antisense compound is complementary to a region of an Usher pre-mRNA. In certain embodiments, an antisense compound is complementary within a region of an Usher pre-mRNA comprising exon 2, intron 2, exon 3, and intron 3. In certain embodiments, an antisense compound is complementary to a region of an Usher pre-mRNA consisting of exon 2, intron 2, exon 3, and intron 3. In certain embodiments, an antisense compound is complementary to a region of an Usher pre-mRNA consisting of intron 2, exon 3, and intron 3. In certain embodiments, an antisense compound is complementary to a region of an Usher pre-mRNA consisting of exon 3 and intron 3. In certain embodiments, an antisense compound is complementary to a region of an Usher pre-mRNA consisting of exon 3.

In certain embodiments, an antisense compound comprises a modified oligonucleotide consisting of 8 to 30 linked nucleosides and having a nucleobase sequence comprising a complementary region comprising at least 8 contiguous nucleobases complementary to a target region of equal length of an Usher transcript. In certain embodiments, the target region is within nucleobase 138475 and nucleobase 138660 of SEQ ID NO.: 1. In certain embodiments, the target region is within nucleobase 138480 and nucleobase 138500 of SEQ ID NO.: 1. In certain embodiments, the target region is within nucleobase 138515 and nucleobase 138535 of SEQ ID NO.: 1. In certain embodiments, the target region is within nucleobase 138550 and nucleobase 138575 of SEQ ID NO.: 1. In certain embodiments, the target region is within nucleobase 138577 and nucleobase 138597 of SEQ ID NO.: 1.

In certain embodiments, an antisense oligonucleotide modulates splicing of a pre-mRNA. In certain embodiments, an antisense oligonucleotide modulates splicing an Usher pre-mRNA. In certain such embodiments, the Usher pre-mRNA is transcribed from a mutant variant of Usher. In certain embodiments, the mutant variant comprises a cryptic splice site. In certain embodiments, an antisense oligonucleotide reduces cryptic splicing of an Usher pre-mRNA. In certain embodiments, an antisense oligonucleotide increases that amount of normally spliced Usher mRNA. In certain embodiments, an antisense oligonucleotide increases that amount of exon 3 skipped Usher mRNA.

In certain embodiments, an antisense oligonucleotide alters the amount of harmonin mRNA. In certain embodiments, an antisense oligonucleotide alters the amount of harmonin RNA. In certain

embodiments, an antisense oligonucleotide alters the amount of harmonin protein. In certain embodiments, an antisense oligonucleotide alters the amount of harmonin mRNA in the ear. In certain embodiments, an antisense oligonucleotide alters the amount of harmonin RNA in the ear. In certain embodiments, an antisense oligonucleotide alters the amount of harmonin protein in the ear. In certain embodiments, an antisense oligonucleotide alters the amount of harmonin mRNA in the cochlea. In certain embodiments, an antisense oligonucleotide alters the amount of harmonin RNA in the cochlea. In certain embodiments, an antisense oligonucleotide alters the amount of harmonin protein in the cochlea. In certain embodiments, an antisense oligonucleotide alters the amount of harmonin mRNA in the retina. In certain embodiments, an antisense oligonucleotide alters the amount of harmonin RNA in the retina. In certain embodiments, an antisense oligonucleotide alters the amount of harmonin protein in the retina.

In certain embodiments, an antisense oligonucleotide increases the amount of harmonin mRNA. In certain embodiments, an antisense oligonucleotide increases the amount of harmonin RNA. In certain embodiments, an antisense oligonucleotide increases the amount of harmonin protein. In certain embodiments, an antisense oligonucleotide increases the amount of harmonin mRNA in the ear. In certain embodiments, an antisense oligonucleotide increases the amount of harmonin RNA in the ear. In certain embodiments, an antisense oligonucleotide increases the amount of harmonin protein in the ear. In certain embodiments, an antisense oligonucleotide increases the amount of harmonin mRNA in the cochlea. In certain embodiments, an antisense oligonucleotide increases the amount of harmonin RNA in the cochlea. In certain embodiments, an antisense oligonucleotide increases the amount of harmonin protein in the cochlea. In certain embodiments, an antisense oligonucleotide increases the amount of harmonin mRNA in the retina. In certain embodiments, an antisense oligonucleotide increases the amount of harmonin RNA in the retina. In certain embodiments, an antisense oligonucleotide increases the amount of harmonin protein in the retina.

In certain embodiments, an antisense oligonucleotide increases the amount of correctly shaped stereocilia in the ear of an animal with Usher syndrome.

Certain Pharmaceutical Compositions

In certain embodiments, the present invention provides pharmaceutical compositions comprising one or more antisense compound. In certain embodiments, such pharmaceutical composition comprises a suitable pharmaceutically acceptable diluent or carrier. In certain embodiments, a pharmaceutical composition comprises a sterile saline solution and one or more antisense compound. In certain embodiments, such pharmaceutical composition consists of a sterile saline solution and one or more antisense compound. In certain embodiments, the sterile saline is pharmaceutical grade saline. In certain embodiments, a pharmaceutical composition comprises one or more antisense compound and sterile water. In certain embodiments, a pharmaceutical composition consists of one or more antisense compound and sterile water. In certain embodiments, the sterile saline is pharmaceutical grade water. In certain embodiments, a

pharmaceutical composition comprises one or more antisense compound and phosphate-buffered saline (PBS). In certain embodiments, a pharmaceutical composition consists of one or more antisense compound and sterile phosphate-buffered saline (PBS). In certain embodiments, the sterile saline is pharmaceutical grade PBS.

5 In certain embodiments, antisense compounds may be admixed with pharmaceutically acceptable active and/or inert substances for the preparation of pharmaceutical compositions or formulations. Compositions and methods for the formulation of pharmaceutical compositions depend on a number of criteria, including, but not limited to, route of administration, extent of disease, or dose to be administered.

10 Pharmaceutical compositions comprising antisense compounds encompass any pharmaceutically acceptable salts, esters, or salts of such esters. In certain embodiments, pharmaceutical compositions comprising antisense compounds comprise one or more oligonucleotide which, upon administration to an animal, including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to pharmaceutically acceptable salts of antisense compounds, prodrugs, pharmaceutically acceptable salts of such prodrugs, and other
15 bioequivalents. Suitable pharmaceutically acceptable salts include, but are not limited to, sodium and potassium salts.

A prodrug can include the incorporation of additional nucleosides at one or both ends of an oligomeric compound which are cleaved by endogenous nucleases within the body, to form the active antisense oligomeric compound.

20 Lipid moieties have been used in nucleic acid therapies in a variety of methods. In certain such methods, the nucleic acid is introduced into preformed liposomes or lipoplexes made of mixtures of cationic lipids and neutral lipids. In certain methods, DNA complexes with mono- or poly-cationic lipids are formed without the presence of a neutral lipid. In certain embodiments, a lipid moiety is selected to increase distribution of a pharmaceutical agent to a particular cell or tissue. In certain embodiments, a lipid moiety is
25 selected to increase distribution of a pharmaceutical agent to fat tissue. In certain embodiments, a lipid moiety is selected to increase distribution of a pharmaceutical agent to muscle tissue.

In certain embodiments, pharmaceutical compositions provided herein comprise one or more modified oligonucleotides and one or more excipients. In certain such embodiments, excipients are selected from water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylase, magnesium stearate, talc,
30 silicic acid, viscous paraffin, hydroxymethylcellulose and polyvinylpyrrolidone.

In certain embodiments, a pharmaceutical composition provided herein comprises a delivery system. Examples of delivery systems include, but are not limited to, liposomes and emulsions. Certain delivery systems are useful for preparing certain pharmaceutical compositions including those comprising hydrophobic compounds. In certain embodiments, certain organic solvents such as dimethylsulfoxide are
35 used.

In certain embodiments, a pharmaceutical composition provided herein comprises one or more tissue-specific delivery molecules designed to deliver the one or more pharmaceutical agents of the present invention to specific tissues or cell types. For example, in certain embodiments, pharmaceutical compositions include liposomes coated with a tissue-specific antibody.

5 In certain embodiments, a pharmaceutical composition provided herein comprises a co-solvent system. Certain of such co-solvent systems comprise, for example, benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer, and an aqueous phase. In certain embodiments, such co-solvent systems are used for hydrophobic compounds. A non-limiting example of such a co-solvent system is the VPD co-solvent system, which is a solution of absolute ethanol comprising 3% w/v benzyl alcohol, 8% w/v of the nonpolar
10 surfactant Polysorbate 80™ and 65% w/v polyethylene glycol 300. The proportions of such co-solvent systems may be varied considerably without significantly altering their solubility and toxicity characteristics. Furthermore, the identity of co-solvent components may be varied: for example, other surfactants may be used instead of Polysorbate 80™; the fraction size of polyethylene glycol may be varied; other biocompatible polymers may replace polyethylene glycol, e.g., polyvinyl pyrrolidone; and other sugars or polysaccharides
15 may substitute for dextrose.

In certain embodiments, a pharmaceutical composition provided herein is prepared for oral administration. In certain embodiments, pharmaceutical compositions are prepared for buccal administration.

In certain embodiments, a pharmaceutical composition is prepared for administration by injection (e.g., intravenous, subcutaneous, intramuscular, etc.). In certain of such embodiments, a pharmaceutical
20 composition comprises a carrier and is formulated in aqueous solution, such as water or physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. In certain embodiments, other ingredients are included (e.g., ingredients that aid in solubility or serve as preservatives). In certain embodiments, injectable suspensions are prepared using appropriate liquid carriers, suspending agents and the like. Certain pharmaceutical compositions for injection are presented in unit dosage form, e.g.,
25 in ampoules or in multi-dose containers. Certain pharmaceutical compositions for injection are suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Certain solvents suitable for use in pharmaceutical compositions for injection include, but are not limited to, lipophilic solvents and fatty oils, such as sesame oil, synthetic fatty acid esters, such as ethyl oleate or triglycerides, and liposomes. Aqueous injection suspensions may contain
30 substances that increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, such suspensions may also contain suitable stabilizers or agents that increase the solubility of the pharmaceutical agents to allow for the preparation of highly concentrated solutions.

In certain embodiments, a pharmaceutical composition is prepared for transmucosal administration. In certain of such embodiments penetrants appropriate to the barrier to be permeated are used in the
35 formulation. Such penetrants are generally known in the art.

In certain embodiments, a pharmaceutical composition provided herein comprises an oligonucleotide in a therapeutically effective amount. In certain embodiments, the therapeutically effective amount is sufficient to prevent, alleviate or ameliorate symptoms of a disease or to prolong the survival of the subject being treated. Determination of a therapeutically effective amount is well within the capability of those skilled in the art.

In certain embodiments, one or more modified oligonucleotide provided herein is formulated as a prodrug. In certain embodiments, upon *in vivo* administration, a prodrug is chemically converted to the biologically, pharmaceutically or therapeutically more active form of an oligonucleotide. In certain embodiments, prodrugs are useful because they are easier to administer than the corresponding active form. For example, in certain instances, a prodrug may be more bioavailable (e.g., through oral administration) than is the corresponding active form. In certain instances, a prodrug may have improved solubility compared to the corresponding active form. In certain embodiments, prodrugs are less water soluble than the corresponding active form. In certain instances, such prodrugs possess superior transmittal across cell membranes, where water solubility is detrimental to mobility. In certain embodiments, a prodrug is an ester. In certain such embodiments, the ester is metabolically hydrolyzed to carboxylic acid upon administration. In certain instances the carboxylic acid containing compound is the corresponding active form. In certain embodiments, a prodrug comprises a short peptide (polyaminoacid) bound to an acid group. In certain of such embodiments, the peptide is cleaved upon administration to form the corresponding active form.

In certain embodiments, the present invention provides compositions and methods for reducing the amount or activity of a target nucleic acid in a cell. In certain embodiments, the cell is in an animal. In certain embodiments, the animal is a mammal. In certain embodiments, the animal is a rodent. In certain embodiments, the animal is a primate. In certain embodiments, the animal is a non-human primate. In certain embodiments, the animal is a human.

In certain embodiments, the present invention provides methods of administering a pharmaceutical composition comprising an oligomeric compound of the present invention to an animal. Suitable administration routes include, but are not limited to, oral, rectal, transmucosal, intestinal, enteral, topical, suppository, through inhalation, intrathecal, intracerebroventricular, intraperitoneal, intranasal, intraocular, intratumoral, and parenteral (e.g., intravenous, intramuscular, intramedullary, and subcutaneous). In certain embodiments, pharmaceutical intrathecal are administered to achieve local rather than systemic exposures. For example, pharmaceutical compositions may be injected directly in the area of desired effect (e.g., into the eyes, ears).

In certain embodiments, a pharmaceutical composition is administered to an animal having at least one symptom associated with Usher Syndrome. In certain embodiments, such administration results in amelioration of at least one symptom. In certain embodiments, administration of a pharmaceutical composition to an animal results in a decrease of cryptic spliced Usher mRNA in a cell of the animal. In

certain embodiments, such administration results in an increase in normally spliced Usher mRNA and/or an increase in exon 3 skipped mRNA. In certain embodiments, such administration results in a decrease in cryptic Usher protein and an increase in normal Usher protein and/or truncated Usher protein lacking exon 3 amino acids. In certain embodiments, an Usher protein lacking exon 3 amino acids is preferred over cryptic Usher protein. In certain embodiments, administration of a pharmaceutical composition results in amelioration of auditory and/or visual defects. In certain embodiments, such amelioration is the reduction in severity of such defects. In certain embodiments, amelioration is the delayed onset of such defects. In certain embodiments, amelioration is the slowed progression of such defects. In certain embodiments, amelioration is the prevention of such defects. In certain embodiments, amelioration is the slowed progression of such defects. In certain embodiments, amelioration is the reversal of such defects.

In certain embodiments, one tests an animal for defects in the Usher gene. In certain embodiments, one identifies an animal having one or more splicing defects in the Usher gene. In certain embodiments, a pharmaceutical composition is administered to an animal identified as having a defect in the Usher gene. In certain embodiments, the animal is tested following administration.

In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is administered while the subject is in utero or very young. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is administered while the subject is in utero or very young. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is administered to the subject in utero. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is administered to the subject from 0-15 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is administered to the subject from 0-12 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is administered to the subject from 0-10 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is administered to the subject from 0-9 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is administered to the subject from 0-8 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is administered to the subject from 0-7 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is administered to the subject from 0-6 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is administered to the subject from 0-5 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is administered to the subject from 0-4 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is administered to the subject from 0-3 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is administered to the subject from 0-2 days after birth. In certain embodiments, a

pharmaceutical composition comprising an antisense oligonucleotide is administered to the subject from 0-1 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is administered to the subject from 2-5 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is administered to the subject from 3-5 days after birth.

In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is first administered while the subject is in utero or very young. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is first administered while the subject is in utero or very young. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is first administered to the subject in utero. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is first administered to the subject from 0-15 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is first administered to the subject from 0-12 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is first administered to the subject from 0-10 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is first administered to the subject from 0-9 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is first administered to the subject from 0-8 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is first administered to the subject from 0-7 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is first administered to the subject from 0-6 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is first administered to the subject from 0-5 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is first administered to the subject from 0-4 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is first administered to the subject from 0-3 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is first administered to the subject from 0-2 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is first administered to the subject from 0-1 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is first administered to the subject from 2-5 days after birth. In certain embodiments, a pharmaceutical composition comprising an antisense oligonucleotide is first administered to the subject from 3-5 days after birth.

Nonlimiting disclosure and incorporation by reference

While certain compounds, compositions and methods described herein have been described with specificity in accordance with certain embodiments, the following examples serve only to illustrate the

compounds described herein and are not intended to limit the same. Each of the references, GenBank accession numbers, and the like recited in the present application is incorporated herein by reference in its entirety.

Although the sequence listing accompanying this filing identifies each sequence as either “RNA” or “DNA” as required, in reality, those sequences may be modified with any combination of chemical modifications. One of skill in the art will readily appreciate that such designation as “RNA” or “DNA” to describe modified oligonucleotides is, in certain instances, arbitrary. For example, an oligonucleotide comprising a nucleoside comprising a 2'-OH sugar moiety and a thymine base could be described as a DNA having a modified sugar (2'-OH for the natural 2'-H of DNA) or as an RNA having a modified base (thymine (methylated uracil) for natural uracil of RNA).

Accordingly, nucleic acid sequences provided herein, including, but not limited to those in the sequence listing, are intended to encompass nucleic acids containing any combination of natural or modified RNA and/or DNA, including, but not limited to such nucleic acids having modified nucleobases. By way of further example and without limitation, an oligomeric compound having the nucleobase sequence “ATCGATCG” encompasses any oligomeric compounds having such nucleobase sequence, whether modified or unmodified, including, but not limited to, such compounds comprising RNA bases, such as those having sequence “AUCGAUCG” and those having some DNA bases and some RNA bases such as “AUCGATCG” and oligomeric compounds having other modified or naturally occurring bases, such as “AT^{me}CGAUCG,” wherein ^{me}C indicates a cytosine base comprising a methyl group at the 5-position.

Examples

The following examples illustrate certain embodiments of the present invention and are not limiting. Moreover, where specific embodiments are provided, the inventors have contemplated generic application of those specific embodiments. For example, disclosure of an oligonucleotide having a particular motif provides reasonable support for additional oligonucleotides having the same or similar motif. And, for example, where a particular high-affinity modification appears at a particular position, other high-affinity modifications at the same position are considered suitable, unless otherwise indicated.

Example 1: Ush1c.216a minigene

A plasmid comprising an Usher 1C mini-gene having a 216A mutation (Ush1c.216a) was prepared using standard molecular biology techniques. The Ush1c.216a plasmid comprised exons 2, 3, and 4, and introns 2 and 3. The mini-gene was under control of the CMV promoter. A schematic of the Ush1c.216a plasmid appears in Figure 1a.

Example 2: Antisense Modulation of Usher Transcript

Antisense oligonucleotides complementary to different regions the Usher transcript were synthesized and tested for their ability to modulate splicing of the Usher mini-gene. Position on the Usher transcript is represented in Figure 1b. Each nucleoside of the oligonucleotides was a 2'-MOE modified nucleoside (i.e., uniform 2'-MOE oligonucleotides). All internucleoside linkages were phosphorothioate linkages (i.e., uniform PS). All of the nucleobases were unmodified and cytosine bases were 5-meC.

To test the ability of the antisense oligonucleotides to modulate splicing, HeLa cells were co-transfected with the Ush1c.216a plasmid from Example 1 and an antisense oligonucleotide (or no antisense oligonucleotide in the case of the untreated control). RNA was collected from the cells and RT-PCR was used to identify Usher transcripts. The results are summarized in Table 1 below and in Figure 1b. Start site is the position relative to SEQ ID NO 1.

Table 1

Modulation of splicing of USH1C pre-mRNA levels by modified oligonucleotides

ISIS NO	5'-Start Site	Sequence	Region	% cryptic splicing	SEQ ID NO
N/A	N/A	N/A	Untreated Control	100	N/A
527106	138475	ACGGCCACGTCCATGGTC	USH1C exon 3	13.39	2
527107	138480	CGAGCACGGCCACGTCCA	USH1C exon 3	6.29	3
527108	138485	TCCCACGAGCACGGCCAC	USH1C exon 3	9.93	4
527109	138490	AGGTCTCCCACGAGCACG	USH1C exon 3	32.49	5
527110	138495	GCTTCAGGTCTCCCACGA	USH1C exon 3	34.79	6
527111	138500	GACCAGCTTCAGGTCTCC	USH1C exon 3	64.21	7
527112	138505	TTGATGACCAGCTTCAGG	USH1C exon 3	23.89	8
527113	138510	G TTCATTGATGACCAGCT	USH1C exon 3	34.68	9
527114	138515	GCTGGGTTTCATTGATGAC	USH1C exon 3	41.71	10
527115	138520	AGACGGCTGGGTTTCATTG	USH1C exon 3	12.15	11
527116	138525	GAGGCAGACGGCTGGGTT	USH1C exon 3	36.97	12
527117	138530	AAACAGAGGCAGACGGCT	USH1C exon 3	26.32	13

527118	138535	GCATCAAACAGAGGCAGA	USH1C exon 3	22.23	14
527119	138540	GAATGGCATCAAACAGAG	USH1C exon 3	29.63	15
527120	138545	CGGCCGAATGGCATCAAA	USH1C exon 3	63.65	16
527121	138550	ATCAGCGGCCGAATGGCA	USH1C exon 3	15.79	17
527122	138555	GTGGGATCAGCGGCCGAA	USH1C exon 3	57.54	18
527123	138560	CTTCAGTGGGATCAGCGG	USH1C exon 3	5.27	19
527124	138563	TGCTTCAGTGGGATCAGC	USH1C exon 3	3.61	20
527125	138566	TGGTGCTTCAGTGGGATC	USH1C exon 3	9.68	21
527126	138569	ACCTGGTGCTTCAGTGGG	USH1C exon 3	21.75	22
527127	138569	ATATTCTACCTGGTGCTTCAGTGGG	USH1C exon 3 (G to A mt)	16.77	23
527128	138571	CTACCTGGTGCTTCAGTG	USH1C exon 3 (G to A mt)	22.39	24
527129	138573	TTCTACCTGGTGCTTCAG	USH1C exon 3 (G to A mt)	24.45	25
527130	138576	ATATTCTACCTGGTGCTT	USH1C exon 3 (G to A mt)	14.89	26
527131	138577	AGCTGATCATATTCTACCTGGTGCT	USH1C exon 3 (G to A mt)	2.35	27
527132	138579	ATCATATTCTACCTGGTG	USH1C exon 3 (G to A mt)	12.13	28
527133	138581	TGATCATATTCTACCTGG	USH1C exon 3 (G to A mt)	2.85	29
527134	138584	AGCTGATCATATTCTACC	USH1C exon 3 (G to A mt)	2.70	30
527135	138586	TCAGCTGATCATATTCTA	USH1C exon 3 (G to A mt)	19.98	31

527136	138589	GGGTCAGCTGATCATATT	USH1C exon 3	98.82	32
527137	138591	GGGGGTCAGCTGATCATA	USH1C exon 3	99.28	33
527138	138593	CGCCGGGGGGTCAGCTGA	USH1C exon 3	99.60	34
527139	138598	TGGAGCGCCGGGGGGTCA	USH1C exon 3	90.93	35
527140	138603	GCACCTGGAGCGCCGGGG	USH1C exon 3/intron3	97.59	36
527141	138608	CCTCTGCACCTGGAGCGC	USH1C exon 3/intron3	99.81	37
527142	138613	GGCTTCCTCTGCACCTGG	USH1C exon 3/intron3	99.54	38
527143	138618	CTGGTGGCTTCCTCTGCA	USH1C intron 3	97.64	39
527144	138623	CCAGCCTGGTGGCTTCCT	USH1C intron 3	96.34	40
527145	138628	TGCCTCCAGCCTGGTGGC	USH1C intron 3	94.86	41
527146	138633	CCCCCTGCCTCCAGCCTG	USH1C intron 3	96.78	42
527147	138638	CTCCACCCCCTGCCTCCA	USH1C intron 3	98.2	43
527148	138643	GATCTCTCCACCCCCTGC	USH1C intron 3	97.94	44
527149	138648	AGGGTGATCTCTCCACCC	USH1C intron 3	97.82	45
527150	138653	CGCCCAGGGTGATCTCTC	USH1C intron 3	94.03	46
527151	138658	TGCCCCGCCAGGGTGAT	USH1C intron 3	97.74	47
527152	138663	AGCACTGCCCCGCCAGG	USH1C intron 3	97.83	48

As is evident from the table above, oligonucleotides having a 5'-start site between positions 138475 and 138587 demonstrated modulation of splicing of the Usher transcript. Thus, in certain embodiments active oligonucleotides have a 5'-start site between 138475 and 138587. In certain embodiments active oligonucleotides are complementary to a target sequence from 138475 to 138603 (the 3'-end of ISIS 527135). In certain embodiments active oligonucleotides are complementary to a target sequence from

138475 to 138607 (the 3'-end of a 20-mer having the same 5'-start site as ISIS 527135). In certain embodiments active oligonucleotides are complementary to a target sequence from 138475 to 138617 (the 3'-end of a 30-mer having the same 5'-start site as ISIS 527135).

In certain embodiments, active oligonucleotides have a 5'-start site between 138475 and 138495. In certain embodiments, active oligonucleotides have a 5'-start site between 138475 and 138495. In certain embodiments, active oligonucleotides have a 5'-start site between 138505 and 138540. In certain embodiments, active oligonucleotides have a 5'-start site between 138550 and 138586. In certain embodiments, active oligonucleotides have a 5'-start site between 138560 and 138586. In certain embodiments, active oligonucleotides have a 5'-start site between 138577 and 138584.

In certain embodiments active oligonucleotides are complementary to a target sequence from 138475 to 138415. In certain embodiments active oligonucleotides are complementary to a target sequence from 138475 to 138505. In certain embodiments active oligonucleotides are complementary to a target sequence from 138505 to 138515. In certain embodiments active oligonucleotides are complementary to a target sequence from 138505 to 138560. In certain embodiments active oligonucleotides are complementary to a target sequence from 138550 to 138606. In certain embodiments active oligonucleotides are complementary to a target sequence from 138560 to 138606. In certain embodiments active oligonucleotides are complementary to a target sequence from 138577 to 138604.

Example 3: Antisense Modulation of Usher Transcript

Additional antisense oligonucleotides complementary to Usher were synthesized and tested in HeLa cells transfected with the Ush1c.216a plasmid for their ability to modulate splicing as in Example 2. The antisense oligonucleotides were uniform 2'-MOE/uniform PS and all nucleobases were unmodified (all cytosines 5-mC) as in Example 2. Results are summarized in Table 2, below.

Table 2

Modulation of splicing of USH1C pre-mRNA levels by modified oligonucleotides

ISIS NO	Start Site	Sequence	Target	% cryptic splicing	SEQ ID NO
535400	138579	ATATTCTACCTGGTG	USH1C exon 3 (G to A mt)	53.03	49
535401	138580	CATATTCTACCTGGT	USH1C exon 3 (G to A mt)	61.03	50
535402	138581	TCATATTCTACCTGG	USH1C exon 3 (G to A mt)	66.12	51
535403	138582	ATCATATTCTACCTG	USH1C exon 3 (G to A mt)	41.61	52

535404	138583	GATCATATTCTACCT	USH1C exon 3 (G to A mt)	22.64	53
535405	138584	TGATCATATTCTACC	USH1C exon 3 (G to A mt)	27.35	54
535406	138585	CTGATCATATTCTAC	USH1C exon 3 (G to A mt)	20.08	55
535407	138586	GCTGATCATATTCTA	USH1C exon 3 (G to A mt)	16.79	56
535408	138587	AGCTGATCATATTCT	USH1C exon 3 (G to A mt)	72.49	57
535409	138588	ATCATATTCTAC	USH1C exon 3 (G to A mt)	98.38	58

As is evident from the table above, oligonucleotides having a 5'-start site between positions 138579 and 138587 demonstrated modulation of splicing of the Usher transcript. Thus, in certain embodiments active oligonucleotides have a 5'-start site between 138579 and 138587. In certain embodiments active oligonucleotides are complementary to a target sequence from 138579 to 138601 (the 3'-end of ISIS 535408). In certain embodiments active oligonucleotides are complementary to a target sequence from 138579 to 138607 (the 3'-end of a 20-mer having the same 5'-start site as ISIS 535408). In certain embodiments active oligonucleotides are complementary to a target sequence from 138475 to 138617 (the 3'-end of a 30-mer having the same 5'-start site as ISIS 535408).

In certain embodiments, active oligonucleotides have a 5'-start site between 138582 and 138586. In certain embodiments, active oligonucleotides have a 5'-start site between 138583 and 138586.

In certain embodiments active oligonucleotides are complementary to a target sequence from 138582 to 138606. In certain embodiments active oligonucleotides are complementary to a target sequence from 138583 to 138606.

Example 4: Antisense Modulation of Usher Transcript (Dose-Response)

Four of the antisense oligonucleotides above were separately tested at varying doses (0 (control), 5nM, 10nM, 20nM, 40nM, and 80nM) as described above in Examples 2 and 3. RNA was collected and analyzed by RT-PCR, as above. Results are summarized in Figure 1d. Each antisense oligonucleotide reduced the amount of cryptic spliced transcript and increased the amount of correctly spliced or exon 3-skipped transcript in a dose-dependent manner.

Example 5: In vitro dose response

Ush1c.216AA knock-in mouse kidney cell lines were cultured and were treated with increasing concentrations of Isis No. 527134. RNA was isolated and then analyzed by radiolabeled RT-PCR.

Additionally, Western-blot analysis of harmonin protein in lysates from cells treated with increasing

concentrations of Isis No. 527134 were measured. Results are summarized in Figures 5a-c. This example shows that antisense compounds blocked cryptic splicing and promoted correct splicing of the endogenous Ush1c.216A gene transcript when transfected into cultured cells derived from a Ush1c.216AA mouse kidney. This example also shows that antisense compounds increased harmonin protein expression when transfected into cultured cells derived from an Ush1c.216AA mouse kidney (see Figure 5c).

Example 6: In vitro modulation of the Usher Transcript in Ush1c.216AA patient-derived lymphoblasts

Cultured Ush1c.216AA patient-derived lymphoblasts were treated for 48 hours with 3 μ M or 6 μ M of Isis No. 527134 using Lipofectamine™ 2000. Primers specific to human Ush1c exon 3 and exon 4 were used to analyze the percent of correct splicing by RT-PCR. The results of the RT-PCR are shown in Table 3 below and in Figure 7. This example demonstrates that antisense compounds can correct splicing in patient-derived Ush1c cells.

Table 3

Correction of splicing in patient-derived lymphoblasts

Isis No.	Dose (μ M)	% Correct Splicing
527134	3	26.4
527134	6	45.8

Example 7: In vivo modulation of the Usher Transcript

Mice having the 216A mutation to the Ush1c gene have been described. Such mice have congenital hearing loss and retinal degeneration. Four of the above described antisense oligonucleotides (527133, 527134, 535401, and 535407) were administered to such mice to test their ability to modulate splicing in vivo.

Doses of 50 mg/kg were administered by intraperitoneal injection twice each week for two weeks. Two days after the final injection, the mice were euthanized and RNA was isolated from various tissues. RNA was analyzed by radiolabeled RT-PCR. Splicing modulation was detected in the kidney and in the eye of treated mice. Results of correct kidney splicing are summarized in Figure 1e.

Example 8: Tissue specific in vivo modulation of the Usher Transcript

Mice having the 216A mutation to the Ush1c gene have been described. Such mice have congenital hearing loss and retinal degeneration. Ush1c.216AA mice were injected with Isis No. 527134 or a mismatched control oligonucleotide. Heterozygous Ush1c.216GA or wild-type Ush1c.216GG mice were also injected with Isis No. 527134.

RT-PCR analysis of RNA isolated from the retina of the mice was obtained at 32 days post birth (Figure 12a). An increase in correct splicing was observed in the retina samples from Ush1c.216AA mice injected with Isis No. 527134. Western Blot analysis was also performed on protein isolated from the retina of the mice (Figure 12b). Harmonin, encoded by the targeted Ush1c gene transcript was detected using an Ush1c specific antibody. B-actin was used as a loading control. Results of the Western Blot show that Ush1c.216AA mice injected with Isis No. 527134 had more harmonin protein in the retina than Ush1c.216AA mice injected with a mismatched control (Figure 12b).

Additionally, Ush1c.216AA mice were injected with Isis No. 527134 or a mismatched control oligonucleotide at 5 days after birth. Samples of RNA isolated from the retina of the Ush1c.216AA mice injected with Isis No. 527134 were collected from mice at ages 12, 30, and 120 days. Samples of RNA isolated from the retina of the Ush1c.216AA mice injected with the mismatched control were collected from mice at 30 days of age. RT-PCR analysis was then performed on the samples. The results are shown in Figure 12c-d and in Table 4. This example demonstrates that antisense oligonucleotides correct splicing in the retina.

Table 4

Quantitation of correct Ush1c splicing in the retina

Age	Dose	% Correct	STDEV
30	Control	0	0.006934
12	527134	1.9	1.734585
30	527134	0.9	1.287369
120	527134	0.2	0.098783

Example 9: In vivo modulation of the Usher Transcript—Dose response

Isis No. 527134 was administered via intraperitoneal injection in doses of 0 mg/kg, 50 mg/kg, 100 mg/kg, and 200 mg/kg twice weekly for two weeks to Ush1c.216AA mice. 24 hours after the final intraperitoneal injection, total RNA samples were prepared from the kidney and analyzed using radio labeled RT-PCR. The RT-PCR analysis showed that doses of Isis No. 527134 increased the amount of correct splicing. Results from the RT-PCR analysis are shown in Table 3 below and in Figure 1f.

Additionally, harmonin protein levels were measured. Results from the Western-Blot analysis are shown in Figure 1g. Western-Blot analysis of lysates from each of the Ush1c.216AA mice demonstrated that doses of Isis No. 527134 increased harmonin protein expression. This example demonstrates that Isis No. 527134 increases correct splicing of the Usher transcript in a dose-dependent manner.

Table 5
RT-PCR Dose Response Study

Isis No./ SEQ ID	Mouse	Dose (mg/kg)	% Correct	Average % Correct
527134/30	1	0	0.4	0.5
527134/30	2	0	0.6	
527134/30	3	0	0.6	
527134/30	4	50	14.5	13
527134/30	5	50	11.3	
527134/30	6	50	13.2	
527134/30	7	100	16.3	18.5
527134/30	8	100	16.3	
527134/30	9	100	23	
527134/30	10	200	27.4	20.2
527134/30	11	200	4.6	
527134/30	12	200	28.6	

Example 10. Restoration of Vestibular Function in Ush1c.216AA Mice

5 a. Open Field Pathway Trace

Neonatal Ush1c.216AA mice aged at 3, 5, 10, or 16 postnatal days (P3, P5, P10, or P16 respectively) were given a single 300 mg/kg intraperitoneal injection of either a mismatched 2'-MOE control antisense compound, or Isis No. 527134. Each mouse was then placed in an open field chamber and its behavior analyzed using ANY-maze behavioral tracking software (Stoelting Co, Wood Dale, IL). Untreated mice, or mice treated with a mismatched 2'-MOE compound were observed to display general hyperactivity and circular behavior when observed in an open-field pathway trace, which is characteristic of a severe vestibular effect. In contrast, the behavior and activity of mice treated with Isis No. 527134 was similar to heterozygote (216GA), or wild type (216GG) mice, with no circling, head-tossing, or hyperactivity observed in an open-field pathway trace. The results of the open-field pathway trace are shown in Figure 2a. Additionally, there was no discernible behavioral difference between mice treated at P3, P5, or P10, whereas P16 treated mice exhibited circling behavior similar to untreated mice or mice treated with the a mismatched 2'-MOE compound. Mice treated at P5 with Isis No. 527134 were not observed to exhibit hyperactivity or circling behavior at 8 months of age. This example illustrates that antisense compounds can effectively cure vestibular function associated with Usher when delivered neonatally.

20 b. Open Field Trace-Quantitation of Rotations

The number of rotations in 120 seconds was measured for each of the mice in the open-field pathway trace (see Example 10a). As illustrated in Figure 2b and in Table 6 below, untreated mice, or mice treated with a mismatch antisense oligonucleotide rotated many times in a 120 second time period. Similarly, mice treated with Isis No. 527134 at 16 days postnatal were also observed to rotate many times within a 120 second time period. Mice treated at P3, P5, or P10 with Isis No. 527134 rotated far fewer times within a 120s period, and the number of rotations observed for mice treated at P3, P5, or P10 with Isis No. 527134 was consistent with the number of rotations observed for heterozygote (216GA) or wild type (216GG) mice. This example illustrates that antisense compounds can effectively cure vestibular function associated with Usher when delivered neonatally.

Table 6

Table-rotations per 120s—Open-field pathway trace

Treatment Day	Untreated	P3, P5, P16	P3	P5	P10	P16	P5	Untreated	Untreated
Compound	None	control	527134	527134	527134	527134	527134	Untreated	Untreated
Genotype	AA	AA	AA	AA	AA	AA	AG	AG	GG
Average Rotations per 120s	34.2	37.0	7.6	9.4	8.1	28.9	8.5	9.1	7.2

c. Open Field Trace-Quantitation of Rotations 6-months post treatment

Untreated heterozygote (216GA) mice, or Ush1c.216AA mice treated with Isis No. 527134 or a mismatched 2'-MOE compound were observed at 180 days. Ush1c.216AA mice treated with a mismatched control displayed general hyperactivity and circular behavior when observed in an open-field pathway trace, which is characteristic of a severe vestibular effect. In contrast, the behavior and activity of mice treated with Isis No. 527134 was similar to heterozygote (216GA) mice, with no circling, head-tossing, or hyperactivity observed in an open-field pathway trace. The results of the open-field pathway trace of mice at 180 days of age are given in Table 7 below and in Figure 2c. This example demonstrates that antisense compounds can improve vestibular function and that improvements in vestibular function are maintained.

Table 7

Table-rotations per 120s—Open-field pathway trace

Compound	Genotype	Number of Mice	Average Rotations per 120s	STDEV
----------	----------	----------------	----------------------------	-------

None	216GA	13	1	0.7
527134	Ush1c.216AA	4	2.5	1.29
Control	Ush1c.216AA	3	11.333	2.1

d. Vestibular Phenotyping in Young and Aged Ush1c.216AA Mice Treated With Isis No. 527134

Three additional independent behavioral tests were used to quantify vestibular function in neonatal Ush1c.216AA mice or heterozygote (216GA) mice given a single 300 mg/kg intraperitoneal injection at 5 days after birth of either a mismatched 2'-MOE control antisense compound or Isis No. 527134. Quantitation of heterozygote (216GA) mouse behavior and Ush1c.216AA mouse behavior was performed on mice that were 2-3 months and 6-9 months old using a swim test, a trunk-curl test, and contact righting test.

Swim scores are based on a scoring system of 0-3. A score of 0 indicates that the animal swims, a score of 1 indicates that irregular swimming was observed, a score of 2 indicates immobile floating, and a score of 3 indicates underwater tumbling and an inability to swim. The results of the swim test are given in Table 8 below. As illustrated in Table 8, Ush1c.216AA mice that received a dose of Isis No. 527134 and heterozygote (216GA) mice that received a dose of the mismatched control were observed to swim. Ush1c.216AA mice that received a dose of a mismatched control were observed tumbling underwater and were unable to swim.

The curl test indicates the number of animals in the group that curl towards their trunk when held by their tail and presented with a landing surface. The results of the curl test are given in Table 8. As illustrated in Table 8, Ush1c.216AA mice that received a dose of Isis No. 527134 and heterozygote (216GA) mice that received a dose of the mismatched control did not curl when held by their tail and presented with a landing surface, indicating a lack of vestibular dysfunction. Ush1c.216AA mice that received a dose of a mismatched control were observed curl when held by their tail and presented with a landing surface, indicating vestibular dysfunction.

Righting indicates the time an animal spent to right itself in an enclosed area when inverted. The results of the righting test are given in Table 8. As illustrated in Table 8, Ush1c.216AA mice that received a dose of Isis No. 527134 and heterozygote (216GA) mice that received a dose of the mismatched control were able to right themselves rapidly. Ush1c.216AA mice that received a dose of a mismatched control were observed to take a much longer amount of time to right themselves.

These examples show that antisense oligonucleotides can effectively curb the vestibular dysfunction associated with Usher, and that the lack of vestibular dysfunction is maintained.

Table 8

Vestibular Phenotyping in Young and Aged Ush1c.216AA Mice

Genotype	Treatment	Age (months)	N	Test		
				Swim Test	Curl (%)	Righting (secs)
Ush1c.216AA	Control	2-3	9	3	100	37.3 ± 6.9
Ush1c.216AA	Control	6-9	6	3	100	18.3 ± 10.2
Ush1c.216AA	527134	2-3	9	0	0	1 ± 0.2
Ush1c.216AA	527134	6-9	6	0	0	1 ± 0.3
Ush1c.216GA	Control	2-3	6	0	0	12 ± 4.7
Ush1c.216GA	Control	6-9	9	0	0	1 ± 0.3

Example 11: Correction of deafness

5 Neonatal Ush1c.216AA mice were treated with either Isis No. 527134 or with a mismatched control oligonucleotide. Responses to high amplitude sound were then measured. Mice treated with a mismatched control oligonucleotide exhibited neither an initial startle response, defined as an ear-twitch and rapid head and body movement, nor a subsequent freezing response response after the acoustic stimulus.

10 To quantitatively assess hearing function, auditory-evoked brainstem response (ABR) analysis was performed. ABR thresholds to broad-band noise (BBN) and pure-tone stimuli at 8 kHz, 16 kHz, and 32kHz were compared in 1-month old Ush1c.216AA mutant mice treated with Isis No. 527134 and those of age matched control mice. Two types of control mice were used: (i) treated and untreated wild type (wt, 216GG) and heterozygote (het, 216GA) mice; and (ii) Ush1c.216AA mutants treated with a mismatched control antisense oligonucleotide. Wild type (wt, 216GG) and heterozygote (het, 216GA) mice, whether untreated
15 or treated with antisense oligonucleotides, had the expected thresholds of mice with normal hearing, as seen in Figures 3a – 3d, Figure 6, and Table 9. Like untreated Ush1c.216AA mutants, Ush1c.216AA mutants treated with the mismatched antisense oligonucleotide had an abnormal response or no response to broad-band noise or pure tones at 90dB sound pressure level (see Figures 3a – 3d). In contrast, Ush1c.216AA mutant mice treated between P3 and P5 with a single 300 mg/kg dose of Isis No. 527134 had normal
20 audiograms with the expected 4-5 peaks and near normal thresholds to broad-band noise and pure-tone stimuli at 8 kHz and 16 kHz when compared to wild type (wt, 216GG) and heterozygote (het, 216GA) mice at 1 month, 2 months, and 3 months. These results are illustrated in Figures 3a-d, Figures 6, 8-9 and in Table

9. As illustrated in Figure 3b, Figure 6, and in Table 9, the treatment effect at one month on thresholds at 32kHz for Ush1c.216AA mutant mice treated between P3 and P5 with a single 300 mg/kg dose of Isis No. 527134 was lower than Ush1c.216AA mutants treated with a mismatched control. Additionally, the treatment effect on thresholds at 32kHz for Ush1c.216AA mutant mice treated between P3 and P5 with a single 300 mg/kg dose of Isis No. 527134 were higher when compared to wild type (wt, 216GG) and heterozygote (het, 216GA) mice at 32kHz.

At 6 months of age, Ush1c.216AA mutant mice treated with Isis No. 527134 at P5 still exhibited thresholds that were lower than untreated mutants, but higher than the wild-type and heterozygous control mice. These results are shown in Figure 10a and 10b and in Table 9. Quantitation of RT-PCR experiments measuring the amount of correctly spliced Ush1c transcripts showed that amount of correctly spliced Ush1c RNA at P30 was similar to that at P120 and P180 (Figure 10c). This indicates that the effect of antisense oligonucleotides on splicing is stable and correlates with the observed ABR results. This example illustrates that the mice injected with a single antisense oligonucleotide-treatment early in life can hear at 1, 2, 3 and 6 months of age, indicating a long-term therapeutic correction of deafness.

Additionally, Ush1c.216AA mutant mice treated with a single 300 mg/kg dose of Isis No. 527134 at P10 had more variable responses with higher thresholds to broad-band noise and pure-tone stimuli than those treated between P3 and P5. Ush1c.216AA mutant mice treated with a single 300 mg/kg dose of Isis No. 527134 at P10 also had lower thresholds to broad-band noise and pure-tone stimuli than untreated mutants or mutants treated with a mismatched control antisense oligonucleotide. The results of mice treated at P10 are shown in Figures 3a-b, Figure 6, and Table 9.

This example illustrates that treatment of Ush1c.216AA mice with antisense oligonucleotides rescued hearing over a broad range of frequencies.

Table 9

Auditory threshold in Ush1c.216AA mice treated with Isis No. 527134 and a mismatched control

Month	Compound/SEQ ID		Frequency			
			8 kHz	16 kHz	32 kHz	BBN
1	AG or GG	Threshold (average)	36	35	35	35
	527134/30 (P5)		36	35	35	35
	527134/30 (P10)		72	73	93	78
	Control		81	86	90	89
2	AG or GG		42	40	50	39
	527134/30 (P5)		47	52	90	49

	Control		90	90	90	90
3	AG or GG		34	36	48	38
	527134/30 (P5)		48	60	90	48
	Control		90	90	90	90
6	AG or GG		38	34	37	42
	527134/30 (P5)		63	77	90	69
	Control		90	90	90	90

Example 12: Restoration of splicing, protein expression, and hair cells in mice

a. Cochlea

Cochleae from mice injected at P5 with 300 mg/kg Isis No. 527134 or a mismatched control were harvested at P30 and subjected to RT-PCR and western blot analyses. As shown in Table 10, correct exon 3 splicing was observed in the Ush1c.216AA mice treated with Isis No. 527134. Correct exon 3 splicing was not seen in Ush1c.216AA mice treated with a mismatched control (Fig. 4a). Likewise, harmonin protein abundance was higher in cochleae isolated from Ush1c.216AA mice treated with Isis No. 527134 compared to Ush1c.216AA mice treated with a mismatched control. Additionally, harmonin protein abundance in the Ush1c.216AA mice treated with Isis No. 527134 was similar to harmonin protein levels in samples from control 216GA mice (Fig. 4b). A low level of harmonin protein was detected in Ush1c.216AA mice that were treated with the mismatched control.

Table 10
Cochlea RNA

Genotype	Treatment	Mouse	% correct splicing	Average % of correct splicing
AA	control	361	0.2	0.4
		366	0.5	
		367	0.5	
		368	0.8	
	untreated	261	0.1	
		257	0.1	
	527134/30	350	51.7	5.6
		353	14.7	
		357	NA	
		233	6.4	
		232	0.8	

		231	0.4	
AG	527134/30	351	47.1	50.8
		352	52.4	
		354	72.2	
		356	59.9	
		359	58.0	
		360	62.4	
		264	30.2	
		263	28.2	
		248	46.4	
		358	75.3	
	control	362	43.3	41.1
		363	11.8	
		369	34.4	
		370	74.9	
GG		262	99.9	

Example 13: Immunohistochemistry analysis of whole cochleae

5 Immunohistochemistry analysis of whole cochleae showed rescue of hair cell morphology with antisense oligonucleotide treatment, consistent with the ABR data in Example 11. Hair cells labeled with anti-parvalbumin antibodies (green) from Ush1c.216AA mice treated with Isis No. 527134 have normal morphology and organized inner and outer rows at the apex-middle turn region similar to control cochleae of 216GA heterozygous animals (Fig. 4c). At P30, when hearing development is complete, harmonin (red) can be seen mostly at the apex of the hair cells (Fig. 4c). Using a pan-specific antibody, harmonin isoforms were detected in mice given the mismatched control to recognize the truncated protein. Hair cells of the mid-basal region of the cochlea of Ush1c.216AA mice treated with Isis No. 527134 were disorganized relative to the heterozygote 216GA mice but exhibited less evidence of degeneration compared to Ush1c.216AA mutants treated with a mismatched control antisense oligonucleotide, where hair cell loss and degeneration was pronounced. These results are shown in Figure 4d.

Example 14: Stereocilia morphology in Ush1c.216AA and 216GA heterozygous mice

Scanning electron microscopy (SEM) was used to analyze the inner ear morphology of P18 mutant Ush1c.216AA mice treated with Isis No. 527134, heterozygous 216GA mice treated with a mismatched

control, and mutant Ush1c.216AA mice treated with a mismatched control. SEM analysis of the organ of Corti from the mutant Ush1c.216AA mice treated with Isis No. 527134 showed some correction of the mutant abnormal hair cell bundles with fewer splayed, disorganized and disconnected groups of stereocilia (Fig. 4f). To quantify this effect, stereocilia bundles that resembled a characteristic “U” or “W”-shape were counted from blinded images of at least 100 hair cells from each experimental group taken from the apex tip, the middle apex, and the middle turn region of the organ of Corti (Fig. 4e, 11a). Images from at least 3 animals in each group were evaluated and the percent of total cells that resembled a “U” or “W”-shape were calculated. The mutant Ush1c.216AA mice treated with a mismatched control had significantly less “U”-shaped bundles in the apex tip and middle apex region compared to mutant mice treated with Isis No. 527134 (Fig. 11a and Fig. 11b). Table 11 below shows the results of the SEM quantitation.

In the middle turn region, the rescue of stereocilia morphology in the Ush1c.216AA mice treated with Isis No. 527134 was not significantly different from Ush1c.216AA mice treated with a mismatched control (Fig. 11a and Fig. 11b). Heterozygous 216GA mice had significantly more normal shaped bundles than 216AA mice treated with Isis No. 527134 (Fig. 11a and Fig. 11b). This pattern of stereocilia rescue in the apical to basal direction is consistent with the more robust rescue of hearing at the frequencies detected by hair cells at the apex (8 kHz). This example demonstrates that rescue of hearing in mice can be achieved with partial rescue of stereocilia bundle structure and organization.

Table 11

Stereocilia morphology in Ush1c.216AA and 216GA heterozygous mice

Treatment	Animal	Average Amount of “U” or “W” Stereocilia Bundles		
		Apex	Mid-Apex	Middle-Turn
Isis No. 527134	AA	30	61	70
Control	AA	9	32	63
Control	GA	100	100	100

Claims:

1. A compound comprising a modified oligonucleotide consisting of 8 to 30 linked nucleosides and having a nucleobase sequence comprising a complementary region comprising at least 8 contiguous nucleobases complementary to a target region of equal length of an Usher transcript.
2. The compound of claim 1, wherein the complementary region of the modified oligonucleotide is 100% complementary to the target region.
3. The compound of claim 1 or 2, the complementary region of the modified oligonucleotide comprises at least 10 contiguous nucleobases.
4. The compound of claim 1 or 2, the complementary region of the modified oligonucleotide comprises at least 15 contiguous nucleobases.
5. The compound of claim 1 or 2, the complementary region of the modified oligonucleotide comprises at least 20 contiguous nucleobases.
6. The compound of any of claims 1-5, wherein the nucleobase sequence of the oligonucleotide is at least 80% complementary to an equal-length region of the Usher transcript, as measured over the entire length of the oligonucleotide.
7. The compound of any of claims 1-5, wherein the nucleobase sequence of the oligonucleotide is at least 90% complementary to an equal-length region of the Usher transcript, as measured over the entire length of the oligonucleotide.
8. The compound of any of claims 1-5, wherein the nucleobase sequence of the oligonucleotide is 100% complementary to an equal-length region of the Usher transcript, as measured over the entire length of the oligonucleotide.
9. The compound of any of claims 1-8, wherein the target region is within exon 3 of the Usher transcript.

10. The compound of any of claims 1-9, wherein the target region is within nucleobase 138475 and nucleobase 138660 of SEQ ID NO.: 1.
11. The compound of any of claims 1-9, wherein the target region is within nucleobase 138480 and nucleobase 138620 of SEQ ID NO.: 1.
12. The compound of any of claims 1-9, wherein the target region is within nucleobase 138550 and nucleobase 138620 of SEQ ID NO.: 1.
13. The compound of any of claims 1-9, wherein the target region is within nucleobase 138577 and nucleobase 138600 of SEQ ID NO.: 1.
14. The compound of any of claims 1-13, wherein the antisense oligonucleotide comprises SEQ ID NO: 30.
15. The compound of any of claims 1-13, wherein the antisense oligonucleotide comprises SEQ ID NO: 29.
16. The compound of any of claims 1-13, wherein the antisense oligonucleotide comprises SEQ ID NO: 27.
17. The compound of any of claims 1-13, wherein the antisense oligonucleotide comprises SEQ ID NO: 50.
18. The compound of any of claims 1-13, wherein the antisense oligonucleotide comprises SEQ ID NO: 56.
19. The compound of any of claims 1-18, wherein the modified oligonucleotide comprises at least one modified nucleoside.
20. The compound of claim 19, wherein at least one modified nucleoside comprises a modified sugar moiety.

21. The compound of claim 20, wherein at least one modified sugar moiety is a 2'-substituted sugar moiety.
22. The compound of claim 21, wherein the 2'-substituent of at least one 2'-substituted sugar moiety is selected from among: 2'-OMe, 2'-F, and 2'-MOE.
23. The compound of claim 22, wherein the 2'-substituent of at least one 2'-substituted sugar moiety is a 2'-MOE.
24. The compound of any of claims 20-23, wherein at least one modified sugar moiety is a bicyclic sugar moiety.
25. The compound of claim 24, wherein at least one bicyclic sugar moiety is LNA or cEt.
26. The compound of any of claims 20-25, wherein at least one sugar moiety is a sugar surrogate.
27. The compound of claim 26, wherein at least one sugar surrogate is a morpholino.
28. The compound of claim 26, wherein at least one sugar surrogate is a modified morpholino.
29. The compound of any of claim 1-28, wherein the modified oligonucleotide comprises at least 5 modified nucleosides, each independently comprising a modified sugar moiety.
30. The compound of claim 29, wherein the modified oligonucleotide comprises at least 10 modified nucleosides, each independently comprising a modified sugar moiety.
31. The compound of claim 29, wherein the modified oligonucleotide comprises at least 15 modified nucleosides, each independently comprising a modified sugar moiety.
32. The compound of claim 29, wherein each nucleoside of the modified oligonucleotide is a modified nucleoside, each independently comprising a modified sugar moiety
33. The compound of any of claims 1-32, wherein the modified oligonucleotide comprises at least two modified nucleosides comprising modified sugar moieties that are the same as one another.

34. The compound of any of claims 1-33, wherein the modified oligonucleotide comprises at least two modified nucleosides comprising modified sugar moieties that are different from one another.
35. The compound of any of claims 1-34, wherein the modified oligonucleotide comprises a modified region of at least 5 contiguous modified nucleosides.
36. The compound of claim 35, wherein the modified oligonucleotide comprises a modified region of at least 10 contiguous modified nucleosides.
37. The compound of claim 35, wherein the modified oligonucleotide comprises a modified region of at least 15 contiguous modified nucleosides.
38. The compound of claim 35, wherein the modified oligonucleotide comprises a modified region of at least 20 contiguous modified nucleosides.
39. The compound of any of claims 34-38, wherein each modified nucleoside of the modified region has a modified sugar moiety independently selected from among: 2'-F, 2'-OMe, 2'-MOE, cEt, LNA, morpholino, and modified morpholino.
40. The compound of any of claims 35-39, wherein the modified nucleosides of the modified region each comprise the same modification as one another.
41. The compound of claim 40, wherein the modified nucleosides of the modified region each comprise the same 2'-substituted sugar moiety.
42. The compound of claim 40, wherein the 2'-substituted sugar moiety of the modified nucleosides of the region of modified nucleosides is selected from 2'-F, 2'-OMe, and 2'-MOE.
43. The compound of claim 41, wherein the 2'-substituted sugar moiety of the modified nucleosides of the region of modified nucleosides is 2'-MOE.
44. The compound of claim 40, wherein the modified nucleosides of the region of modified nucleosides each comprise the same bicyclic sugar moiety.

45. The compound of claim 44, wherein the bicyclic sugar moiety of the modified nucleosides of the region of modified nucleosides is selected from LNA and cEt.
46. The compound of claim 40, wherein the modified nucleosides of the region of modified nucleosides each comprises a sugar surrogate.
47. The compound of claim 46, wherein the sugar surrogate of the modified nucleosides of the region of modified nucleosides is a morpholino.
48. The compound of claim 46, wherein the sugar surrogate of the modified nucleosides of the region of modified nucleosides is a modified morpholino.
49. The compound of any of claims 1-48, wherein the modified nucleotide comprises no more than 4 contiguous naturally occurring nucleosides.
50. The compound of any of claims 1-48, wherein each nucleoside of the modified oligonucleotide is a modified nucleoside.
51. The compound of claim 50 wherein each modified nucleoside comprises a modified sugar moiety.
52. The compound of claim 51, wherein the modified nucleosides of the modified oligonucleotide comprise the same modification as one another.
53. The compound of claim 52, wherein the modified nucleosides of the modified oligonucleotide each comprise the same 2'-substituted sugar moiety.
54. The compound of claim 53, wherein the 2'-substituted sugar moiety of the modified oligonucleotide is selected from 2'-F, 2'-OMe, and 2'-MOE.
55. The compound of claim 54, wherein the 2'-substituted sugar moiety of the modified oligonucleotide is 2'-MOE.

56. The compound of claim 52, wherein the modified nucleosides of the modified oligonucleotide each comprise the same bicyclic sugar moiety.
57. The compound of claim 56, wherein the bicyclic sugar moiety of the modified oligonucleotide is selected from LNA and cEt.
58. The compound of claim 52, wherein the modified nucleosides of the modified oligonucleotide each comprises a sugar surrogate.
59. The compound of claim 58, wherein the sugar surrogate of the modified oligonucleotide is a morpholino.
60. The compound of claim 58, wherein the sugar surrogate of the modified oligonucleotide is a modified morpholino.
61. The compound of any of claims 1-60, wherein the modified oligonucleotide comprises at least one modified internucleoside linkage.
62. The compound of claim 61, wherein each internucleoside linkage is a modified internucleoside linkage.
63. The compound of claim 61 or 62, comprising at least one phosphorothioate internucleoside linkage.
64. The compound of claim 62, wherein each internucleoside linkage is a modified internucleoside linkage and wherein each internucleoside linkage comprises the same modification.
65. The compound of claim 64, wherein each internucleoside linkage is a phosphorothioate internucleoside linkage.
66. The compound of any of claims 1-65 comprising at least one conjugate.
67. The compound of any of claims 1-66 consisting of the modified oligonucleotide.

68. The compound of any of claims 1-67, wherein the compound modulates splicing of the Usher transcript.
69. The compound of any of claims 1-68, having a nucleobase sequence comprising any of the sequences as set forth in SEQ ID NOs. 2-57.
70. The compound of any of claims 1-68, having a nucleobase sequence comprising any of the sequences as set forth in SEQ ID NOs. 2-31 or 49-57.
71. The compound of any of claims 1-68, having a nucleobase sequence comprising any of the sequences as set forth in SEQ ID NOs. 2-31.
72. The compound of any of claims 1-68, having a nucleobase sequence comprising any of the sequences as set forth in SEQ ID NOs. 49-57.
73. A method of modulating splicing an Usher transcript in a cell comprising contacting the cell with a compound according to any of claims 1-72.
74. The method of claim 73, wherein the cell is in vitro.
75. The method of claim 73, wherein the cell is in an animal.
76. A method of modulating the expression of harmonin in a cell, comprising contacting the cell with a compound according to any of claims 1-72.
77. The method of claim 76, wherein harmonin expression is increased.
78. The method of claim 76, wherein the cell is in vitro.
79. The method of claim 76, wherein the cell is in an animal.
80. A method of increasing the total number of correctly shaped stereocilia bundles in the ear, comprising contacting a cell with a compound according to any of claims 1-72.

81. The method of claim 80, wherein the cell is in an animal.
82. A pharmaceutical composition comprising a compound according to any of claims 1-72 and a pharmaceutically acceptable carrier or diluent.
83. The pharmaceutical composition of claim 82, wherein the pharmaceutically acceptable carrier or diluent is sterile saline.
84. A method comprising administering the pharmaceutical composition of claims 82 or 83 to an animal.
85. The method of claim 84, wherein the administration is by injection.
86. The method of claim 84 or 85, wherein the administration is systemic.
87. The method of claim 84 or 85 wherein the administration is local.
88. The method of claim 87, wherein the administration is to the eye of the animal.
89. The method of claim 87, wherein the administration is to the ear of an animal.
90. The method of any of claims 84-89, wherein the animal has one or more symptom of Usher Syndrome.
91. The method of claim 90, wherein the administration results in amelioration of at least one symptom of Usher Syndrome.
92. The method of claim 91, wherein the symptom is deafness.
93. The method of claim 91, wherein the symptom is blindness.
94. The method of any of claims 84-93, wherein the animal is a mouse.
95. The method of any of claims 84-93, wherein the animal is a human.

96. The method of any of claims 84-93, wherein the first administration occurs within 1-15 days after birth.
97. The method of any of claims 84-93, wherein the first administration occurs within 1-10 days after birth.
98. The method of any of claims 84-93, wherein the first administration occurs within 1-5 days after birth.
99. The method of any of claims 84-93, wherein the first administration occurs within 3-5 days after birth.
100. The method of any of claims 84-93, wherein the first administration is in utero.
101. Use of the compound of any of claims 1 to 72 or the composition of claims 82-83 for the preparation of a medicament for use in the treatment of Usher syndrome.
102. Use of the compound of any of claims 1 to 72 or the composition of claims 82-83 for the preparation of a medicament for use in the amelioration of one or more symptoms of Usher syndrome.




Figure 1a




Figure 1b

ASO - 535407	ATCTTATACTAGTCG	
ASO - 535401	TGGTCCCATCTTATAC	
ASO - 527134	CCATCTTATACTAGTCGA	
ASO - 527133	GGTCCATCTTATACTAGT	
USH1C	CCCACTGAAGCACCAAGGTAGAAATATGATCAGCTGACCCCCCGCGCTCCAGgtgcag	
		cryptic 5' splice site 216A
		authentic 5' splice site

Figure 1c




Figure 1d



Figure 1e




Figure 1f




Figure 1g




Figure 2a




Figure 2b




Figure 2c




Figure 3a




Figure 3 b-d




Figure 4a




Figure 4b




Figure 4c




Figure 4d




Figure 4e




Figure 4f




Figure 5a




Figure 5b




Figure 5c




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10a




Figure 10b






Figure 10c

**Figure 11a****Figure 11b**




Figure 12 a-b




Figure 12 c-d

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 12/36189

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - C12N 15/11; C07H 21/04 (2012.01)

USPC - 514/44A; 536/24.5

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

USPC- 514/44A; 536/24.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
USPC- 514/44R, 536/23.1Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
PubWEST(PGPB,USPT,USOC,EPAB,JPAB); Google Patents; Google Scholar: USH1C, exon 3, USH1C intron 3, 216GA, USH1G, alternative splicing, USH2A, USH1E, usher transcript, usher syndrome, usher's syndrome, silencing, antisense, siNA, short interfering nucleic acid, siRNA, dsRNA, shRNA, expression, oligonucleotide, modified

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Jodelka, et al. Antisense oligonucleotide correction of splicing in a mouse model of Usher syndrome. 2010 Rustbelt RNA Meeting. RRM [Retrieved from the Internet 01 July 2012: < http://www.rustbeltma.org/2010/talks.php?all >]; Abstract	1-5
X	Gibbs, et al. Function of MYO7A in the Human RPE and the Validity of Shaker 1 Mice as a Model for Usher syndrome 1B. Investigative Ophthalmology & Visual Science February 2010, 51(2):1130-1135	1-5
A	US 2009/0088721 A1 (de Bizemont, et al.) 02 April 2009 (02.04.2009) para [0011], [0043], [0047], [0048], [0081], [0118], [0119], TABLE 1	1-5
A	US 2010/0203076 A1 (Fotin-Mleczeck, et al.) 12 August 2010 (12.08.2010)	1-5
A	US 2010/0190841 A1 (Farrar, et al.) 29 July 2010 (29.07.2010) Abstract, para [0068], Table 5	1-5
A	US 2009/0163435 A1 (Bader, et al.) 25 June 2009 (25.06.2009) Abstract, para [0059], TABLE 3, NM_173477 Usher syndrome 1G protein USH2A; para [0180], [0203], [0219]-[0228]	1-5
A	US 2011/0021605 A1 (Schulte, et al.) 27 January 2011 (27.01.2011) Abstract, claims 78, 79	1-5
A	Lopez, et al. Mutational frequencies in usherin(USH2A gene) in 26 Colombian individuals with Usher syndrome type II. Biomedica. March 2011, 31(1):82-90	1-5
A	Verpy, et al. A defect in harmonin, a PDZ domain-containing protein expressed in the inner ear sensory hair cells, underlies Usher syndrome type 1C. Nat Genet. 2000, 26(1):51-55	1

☒ Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

25 June 2012 (25.06.2012)

Date of mailing of the international search report

05 OCT 2012

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 12/36189

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	McCaffrey, et al. Blocking Cryptic Splicing In Usher Syndrome Using Antisense Oligonucleotides. Abstract. 03 May 2011. [Retrieved from the Internet 26 June 2012: < http://www.abstractsonline.com/plan/ViewAbstract.aspx?mID=2684&sKey=9456c86d-1d58-492c-ab4f-bc1dfbebebee&cKey=06470cb6-03a8-4dc6-ade5-579e12fd0d93&mKey=%7B6F224A2D-AF6A-4533-8BBB-6A8D7B26EDB3%7D >]	1-5
X,P	Vache, et al. Usher syndrome type 2 caused by activation of an USH2A pseudoexon: implications for diagnosis and therapy. Hum Mutat. Epub 16 November 2011, 33(1):104-108	1-5
A	Lentz, et al. Ush1c216A knock-in mouse survives Katrina. Mutat Res. 2007, 616(1-2):139-44	1-5
A	US 2005/0009771 A1 (Levanon, et al.) 13 January 2005 (13.01.2005) Abstract, para [0769], Table 10, USH1C Usher Syndrome Type 1	1-5
A	Goldmann, et al. Beneficial Read-Through of a USH1C Nonsense Mutation by Designed Aminoglycoside NB30 in the Retina. Invest Ophthalmol Vis Sci. 2010, 51(12):6671-80	1-5
A	Ayuso, et al. Retinitis pigmentosa and allied conditions today: a paradigm of translational research. Genome Med. 2010, 2(5):34	1-5

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 12/36189

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☒ Claims Nos.: 6-102
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.