




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : <b>C07K 14/00</b></p>	<p><b>A2</b></p>	<p>(11) International Publication Number: <b>WO 99/09054</b></p> <p>(43) International Publication Date: 25 February 1999 (25.02.99)</p>
<p>(21) International Application Number: PCT/BE98/00124</p> <p>(22) International Filing Date: 20 August 1998 (20.08.98)</p> <p>(30) Priority Data: 9700692 20 August 1997 (20.08.97) BE</p> <p>(71) Applicants (for all designated States except US): UNIVERSITE CATHOLIQUE DE LOUVAIN [BE/BE]; Halles Universitaires, Place de l'Université 1, B-1348 Louvain-la-Neuve (BE). UNIVERSITE DE MONS-HAINAUT [BE/BE]; Place du Parc 20, B-7000 Mons (BE).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): KNOOPS, Bernard [BE/BE]; Rue Chapelle Notre-Dane 3/1, B-1341 Ceroux-Mousty (BE). HERMANS, Cedric [BE/BE]; Avenue des Glycines 42, B-1030 Brussels (BE). BERNARD, Alfred [BE/BE]; Avenue de la Chapelle 6, B-1200 Brussels (BE). WATTIEZ, Ruddy [BE/BE]; Chemin du Sauvelon 17, B-7022 Hyon (BE). FALMAGNE, Paul [BE/BE]; Rue du Point du Jour 8, B-7022 Mesvin (BE).</p> <p>(74) Agents: VAN MALDEREN, Eric et al.; Office Van Malderen, Place Reine Fabiola 6/1, B-1083 Brussels (BE).</p>		<p>(81) Designated States: CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p><b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i></p>

(54) Title: PEROXISOME-ASSOCIATED POLYPEPTIDE, NUCLEOTIDE SEQUENCE ENCODING SAID POLYPEPTIDE AND THEIR USES IN THE DIAGNOSIS AND/OR THE TREATMENT OF LUNG INJURIES AND DISEASES, AND OF OXIDATIVE STRESS-RELATED DISORDERS



## (57) Abstract

The present invention is related to an isolated and purified polypeptide which amino acid sequence presents more than 70 % with the sequence SEQ ID NO 1. The present invention is also related to the nucleotide sequence encoding said amino acid sequence, the inhibitor directed against said sequences and their use in the diagnosis, treatment and/or prevention of lung injuries or diseases and oxidative stress-related disorders.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

<b>AL</b>	Albania	<b>ES</b>	Spain	<b>LS</b>	Lesotho	<b>SI</b>	Slovenia
<b>AM</b>	Armenia	<b>FI</b>	Finland	<b>LT</b>	Lithuania	<b>SK</b>	Slovakia
<b>AT</b>	Austria	<b>FR</b>	France	<b>LU</b>	Luxembourg	<b>SN</b>	Senegal
<b>AU</b>	Australia	<b>GA</b>	Gabon	<b>LV</b>	Latvia	<b>SZ</b>	Swaziland
<b>AZ</b>	Azerbaijan	<b>GB</b>	United Kingdom	<b>MC</b>	Monaco	<b>TD</b>	Chad
<b>BA</b>	Bosnia and Herzegovina	<b>GE</b>	Georgia	<b>MD</b>	Republic of Moldova	<b>TG</b>	Togo
<b>BB</b>	Barbados	<b>GH</b>	Ghana	<b>MG</b>	Madagascar	<b>TJ</b>	Tajikistan
<b>BE</b>	Belgium	<b>GN</b>	Guinea	<b>MK</b>	The former Yugoslav Republic of Macedonia	<b>TM</b>	Turkmenistan
<b>BF</b>	Burkina Faso	<b>GR</b>	Greece			<b>TR</b>	Turkey
<b>BG</b>	Bulgaria	<b>HU</b>	Hungary	<b>ML</b>	Mali	<b>TT</b>	Trinidad and Tobago
<b>BJ</b>	Benin	<b>IE</b>	Ireland	<b>MN</b>	Mongolia	<b>UA</b>	Ukraine
<b>BR</b>	Brazil	<b>IL</b>	Israel	<b>MR</b>	Mauritania	<b>UG</b>	Uganda
<b>BY</b>	Belarus	<b>IS</b>	Iceland	<b>MW</b>	Malawi	<b>US</b>	United States of America
<b>CA</b>	Canada	<b>IT</b>	Italy	<b>MX</b>	Mexico	<b>UZ</b>	Uzbekistan
<b>CF</b>	Central African Republic	<b>JP</b>	Japan	<b>NE</b>	Niger	<b>VN</b>	Viet Nam
<b>CG</b>	Congo	<b>KE</b>	Kenya	<b>NL</b>	Netherlands	<b>YU</b>	Yugoslavia
<b>CH</b>	Switzerland	<b>KG</b>	Kyrgyzstan	<b>NO</b>	Norway	<b>ZW</b>	Zimbabwe
<b>CI</b>	Côte d'Ivoire	<b>KP</b>	Democratic People's Republic of Korea	<b>NZ</b>	New Zealand		
<b>CM</b>	Cameroon		Republic of Korea	<b>PL</b>	Poland		
<b>CN</b>	China	<b>KR</b>	Republic of Korea	<b>PT</b>	Portugal		
<b>CU</b>	Cuba	<b>KZ</b>	Kazakstan	<b>RO</b>	Romania		
<b>CZ</b>	Czech Republic	<b>LC</b>	Saint Lucia	<b>RU</b>	Russian Federation		
<b>DE</b>	Germany	<b>LI</b>	Liechtenstein	<b>SD</b>	Sudan		
<b>DK</b>	Denmark	<b>LK</b>	Sri Lanka	<b>SE</b>	Sweden		
<b>EE</b>	Estonia	<b>LR</b>	Liberia	<b>SG</b>	Singapore		

5

PEROXISOME-ASSOCIATED POLYPEPTIDE, NUCLEOTIDE SEQUENCE  
10 ENCODING SAID POLYPEPTIDE AND THEIR USES IN THE DIAGNOSIS  
AND/OR THE TREATMENT OF LUNG INJURIES AND DISEASES, AND OF  
OXIDATIVE STRESS-RELATED DISORDERS

Field of the invention

15           The present invention is related to a new  
peroxisome-associated polypeptide, the nucleotide sequence  
encoding said polypeptide and portions thereof as well as  
their uses in the diagnosis of several diseases, especially  
the diagnosis and/or the treatment of lung injuries and  
20 diseases, and of oxidative stress-related disorders.

Background of the invention

          The peroxisomes are organelles nearly  
ubiquitous in eukaryotic cells. They contain enzymes  
25 essential for various catabolic and anabolic pathways. Some  
of these enzymes are expressed constitutively while others  
can be induced under appropriate conditions. Peroxisomes  
carry out a variety of essential reactions such as  
peroxisomal oxidation and respiration, fatty acid beta-  
30 oxidation, cholesterol and dolichol metabolism, ether-  
phospholipid synthesis, and glyoxylate and pipercolic acid  
metabolism.

The peroxisomal respiratory pathway is based upon the formation of hydrogen peroxide by a collection of oxidases and the decomposition of the H<sub>2</sub>O<sub>2</sub> by catalase. These reactions are responsible for 20% of oxygen consumption in liver, and several oxidases have been identified in peroxisomes. Ethanol elimination via catalase in peroxisomes may be significant in addition to the oxidation via cytosolic alcohol dehydrogenase.

The peroxisomal beta-oxidation system catalyses the beta-oxidative chain shortening of a specific set of compounds which can not be handled by mitochondria : very long chain fatty acids, di- and trihydroxycholestanic acids, pristanic acid, long chain dicarboxylic acids, several prostaglandins, several leukotrienes, 12- and 15-hydroxyeicosatetraenoic acid, and several mono- and polyunsaturated fatty acids, which are of direct diagnostic relevance for some peroxisomal disorders.

Peroxisomes play also a major role in the synthesis of cholesterol and other isoprenoids. Fibroblasts from patients affected by disorders of peroxisome biogenesis show low capacity to synthesise cholesterol.

Two enzyme activities responsible for introduction of the characteristic ether linkage in ether-linked phospholipids (dihydroacetonephosphate acyltransferase (DHAPAT) and alkyl-dihydroxyacetonephosphate synthase (alkyl-DHAP synthase)) are localised in peroxisomes. These enzymes are not yet cloned. As demonstrated by the identification of patients with deficiency of either DHAPAT or alkyl-DHAP synthase with severe clinical abnormalities, ether-phospholipids are of major importance in humans.

Peroxisomes are able to detoxify glyoxylate via alanine/glyoxylate aminotransferase. The deficiency of this cloned enzyme causes hyperoxaluria type I.

L-pipecolate is a minor metabolite of L-lysine and is  
5 catabolised by the L-pipecolate oxidase localised in peroxisomes. The enzyme is deficient in cerebro-hepato-renal (Zellweger) syndrome.

In human, the importance of peroxisomes was emphasised by a number of inherited diseases involving  
10 either a defect in the biogenesis of peroxisomes or a deficiency of one (or more) peroxisomal enzymes. So far, 12 different peroxisomal disorders have been described and most of them are lethal.

A wide variety of chemicals have been showed  
15 to produce peroxisome proliferation and induction of peroxisomal and microsomal fatty acids-oxidising enzymes activities in rats and mice. Several peroxisomes proliferators have been shown to increase the incidence of liver tumours in these species. Proposed mechanisms of  
20 liver tumour formation by peroxisomes proliferators include induction of sustained oxidative stress.

Therefore, newly identified molecules associated with peroxisomes could be used for the development of diagnostic tools and possibly for the  
25 improvement of several therapeutical applications of various diseases associated with peroxisomal disorders. In addition, it is useful to identify the molecules present in specific organs like the lung and which may be used as specific markers of inflammatory diseases as well as lung  
30 injuries or diseases.

Summary of the invention

The Inventors have isolated and purified a new sequence of a low molecular weight human broncho-alveolar polypeptide. Said mammal, preferably human, protein or polypeptide (hereafter identified as B18hum  
5 protein) has been sequenced and its corresponding genomic DNA (SEQ ID NO 8) and cDNA (SEQ ID NO 1) have been identified. Similarly, the corresponding nucleotide and amino acid sequence from a rat (SEQ ID NO 3 and 4) and from  
10 a mouse (SEQ ID NO 5 and 6) have been obtained.

Said sequences present several homologies with other peroxisomal proteins of yeast and comprise a carboxy-terminal tripeptide SQL which is necessary for the specific targeting and translocation of several proteins  
15 into the peroxisome.

Therefore, the present invention is related to a new isolated and purified polypeptide sequence having a amino acid sequence which presents more than 70% homology, advantageously more than 85% homology, more  
20 preferably more than 95% homology, with the amino acid sequence SEQ ID NO 2., Said amino acid sequence is advantageously obtained from a mammal, preferably from a rat, a mouse or a human.

The present invention is also related to the  
25 isolated and purified polypeptide sequence corresponding to the amino acid sequence SEQ ID NO 2 or a portion thereof, preferably an immunoreactive portion (putative immunogenic domain or T or B cell epitopes).

Said portions are advantageously comprised  
30 between :

- Glutamic acid position 13 - Glutamic acid position 27
- Alanine position 26 - Leucine position 36

- Alanine position 42 - Glutamic acid position 57
- Glutamic acid position 57 - Valine position 69
- Valine position 80 - Leucine position 97
- Arginine position 95 - Leucine position 112
- 5 - Serine position 118 - Serine position 129
- Valine position 137 - Threonine position 150

Preferably, said portion has more than 10, 20, 30, 50 or 70 amino acids. Specific portions of the amino acid sequence SEQ ID NO 2 are also portions of more than 70 amino acids which present at least 80% of the proteinic activity (see example 5) of the complete SEQ ID NO 2 sequence. Therefore, the amino acid sequence according to the invention can be partially deleted while maintaining its activity, preferably its anti-oxidative activity, which will be described hereafter.

According to the invention, the amino acid sequence SEQ ID NO 2 presents a pI of 7.16 and a molecular weight of 17047 Dalton as hereafter defined by bidimensional electrophoresis.

The present invention is also related to the nucleotide sequence encoding the amino acid sequence according to the invention and its regulatory sequences upstream said coding sequence. A nucleotide sequence encoding the polypeptide according to the invention is a genomic DNA (see SEQ ID NO 10), a cDNA (see SEQ ID NO 1) or a mRNA, possibly comprising said upstream regulatory sequence. Advantageously, said nucleotide sequence presents more than 70%, advantageously more than 85%, more preferably more than 95% homology with SEQ ID NO 1 or its complementary strand.

According to a preferred embodiment of the present invention, said nucleotide sequence corresponds to the nucleotide sequence SEQ ID NO 1, its complementary strand or a portion thereof.

5 "A portion of the nucleotide sequence SEQ ID NO 1" means any nucleotide sequence of more than 15 base pairs (such as a primer, a probe or an antisense nucleotide sequence) which allow the specific identification, reconstitution or blocking of the complete nucleotide  
10 sequence SEQ ID NO 1 (including its regulatory sequences upstream the coding sequence).

Said portions allow the specific identification, reconstitution or blocking by specific hybridisation with the nucleotidic sequence SEQ ID NO 1,  
15 preferably under standard stringent conditions, with sequences like probes or primers possibly labelled with a compound (radioactive compound, enzyme, fluorescent marker, etc.), and can be used in a specific diagnostic or dosage method like probe hybridisation (see Sambrook et al., §§  
20 9.47-9.51 in *Molecular Cloning : A Laboratory Manual*, Cold Spring Harbor, Laboratory Press, Cold Spring Harbor, New York (1989)), genetic amplification (like PCR (US patent 4,683,195), LCR (Wu et al., *Genomics* 4, pp. 560-569), CPR (US patent 5,011,769)).

25 Exemplary stringent hybridisation conditions are as follows : hybridisation at 42 °C in 50% formamide 5x SSC, 20 mM sodium phosphate, pH 6.8 washing in 0.2x SSC at 55 °C. It is understood by those skilled in the art that variation of these conditions occur based on the length and  
30 GC nucleotide content of the sequence to be hybridised. Formulas standard in the art are appropriated for

determining exact hybridisation conditions (see Sambrook et al.

Preferred examples of said nucleotide portions are as follows :

5	<u>Sequence</u>	<u>Position</u>
5'	-gccatcccagcagtgagggtgtttg-3'	(SEQ ID NO 11) 217-241
5'	-ttgaacagctctgccaggttcacc-3'	(SEQ ID NO 12) 261-284
5'	-tggaggtgtttgaaggggagccag-3'	(SEQ ID NO 13) 230-253
5'	-caggttcaccttgttccctggctc-3'	(SEQ ID NO 14) 247-270
10	5'-gggtatgggactagctggcg-3'	(SEQ ID NO 15) 33-52
5'	-ctggccaacattccaattgcag-3'	(SEQ ID NO 16) 747-768

and the sequences of respectively 601 (SEQ ID NO 8), 604 (SEQ ID NO 9) and 469 (SEQ ID NO 7) base pairs corresponding to specific mRNA alternative splicing of the  
 15 B18 human nucleotide sequence as described in Figure 4 (the known genomic sequence incorporating several introns and exons is represented in the sequence SEQ ID NO 10).

Said sequences may be used for a genetic amplification or a probe hybridisation as above-described.

20 The present invention is also related to a vector comprising the necessary elements for the injection, transfection or transduction of cells and having incorporated one or more of the nucleotide sequences according to the invention. The vector according to the  
 25 invention is selected from the group consisting of viruses, plasmids, phagemides, cationic vesicles, liposomes or a mixture thereof. Said vector may comprise also one or more adjacent regulatory sequences (such as promoter(s), secretion and termination signal sequence(s)),  
 30 advantageously operably linked to the nucleotide sequence according to the invention.

The present invention is also related to the cell transformed by said vector and expressing the polypeptide according to the invention.

The nucleotide sequence according to the invention can be also introduced in said cell by the formation of CaPO<sub>4</sub>-nucleic acid precipitate, DEAE-dextran-nucleic acid complex or by electroporation.

Another aspect of the present invention is related to an inhibitor of the polypeptide according to the invention or the nucleotide sequence according to the invention (including the upstream sequences like promoter-operator regulatory sequence which may be inhibited by a cis- and/or transactivating repressor). Said inhibitor is advantageously an antibody or a fragment of said antibody such as an hypervariable portion of said antibody directed against the amino acid or nucleotide sequence of the polypeptide according to the invention. Other examples of inhibitors according to the invention are antisense nucleotide sequences which allow the blocking of the expression of the nucleotide sequence according to the invention.

Another aspect of the present invention is related to a diagnostic device (such as a diagnostic kit or a chromatographic column) comprising an element selected from the group consisting of the amino acid sequence of said polypeptide, its nucleotide sequence, and/or the inhibitor according to the invention or a fragment thereof as above-described. Said diagnostic device may comprise also necessary reactants and media for the diagnostic and/or dosage of the nucleotide and/or amino acid sequence of the polypeptide according to the invention, which are based upon the method selected from the group consisting of

in situ hybridisation, hybridisation by labelled antibodies, especially RIA (Radio Immuno Assay) or ELISA (Enzymes Linked Immuno-Sorbent Assay) technologies, detection upon filter, upon solid support, in solution, in sandwich, upon gel, dot blot hybridisation, Northern blot hybridisation, Southern blot hybridisation, isotopic or non-isotopic labelling (by immunofluorescence or biotinilised probes), genetic amplification, (especially by PCR or LCR), double immunodiffusion technique, counter-electrophoresis technique, haemagglutination or a mixture thereof.

Another aspect of the present invention concerns a diagnosis method wherein a biological sample from the patient, such as cephalo-rachidian fluid, serum, blood, plasma, urine, broncho-alveolar lavage, stomach lavage, etc., is isolated from the patient, and is put in contact with the diagnostic device according to the invention for the diagnosis or the monitoring of an injury or a disease, preferably a lung injury or an oxidative stress-related disorder, affected by the presence of pro-oxidant agent or oxidative stress such as specific cardiovascular diseases like arteriosclerosis, neurodegenerative disorders (Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis), apoptosis, inflammatory reactions, allergic reactions such as asthma, hay fever and eczema, high bone mass syndrome, osteopetrosis, osteoporosis-pseudoglioma syndrome, and Bardet-Biedl syndrome 1. Said diagnosis and monitoring upon one or more biological samples obtained from several tissues from the patient can be advantageously obtained by one or more of the methods above-described, which could be adapted

according to the specific biological sample by the person skilled in the art.

Therefore, the product according to the invention could be used as a marker for the above-  
5 identified injuries, diseases or disorders in a broad spectrum of tissues as shown in the enclosed Figure 1.

A further aspect of the present invention is related to a pharmaceutical composition comprising a pharmaceutically acceptable carrier and an element selected  
10 from the group consisting of the nucleotide sequence, the amino acid sequence of the polypeptide according to the invention, the inhibitor directed against said sequences and/or one or more portions thereof.

A last aspect of the present invention is  
15 related to the use of the pharmaceutical composition according to the invention for the manufacture of a medicament in the treatment and/or the prevention of lung injuries and/or diseases or of oxidative stress-related disorders.

The present invention is also related to a  
20 prevention and/or treatment method of a patient, especially a human patient, preferably affected by lung injuries and/or diseases or by oxidative stress-related disorders, wherein a sufficient amount of the pharmaceutical  
25 composition according to the invention is administered to said patient in order to treat, avoid and/or reduce the symptoms of said injuries and/or diseases.

Other injuries and/or diseases which can be prevented and/or treated are injuries and/or diseases  
30 affected by the presence of pro-oxidant agents or oxidative stress, such as specific cardio-vascular diseases like arteriosclerosis, neurodegenerative disorders such as

Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, apoptosis and inflammatory reactions and some allergic reactions such as asthma, hay fever and eczema, high bone mass syndrome, osteopetrosis, 5 osteoporosis-pseudoglioma syndrome, and Bardet-Biedl syndrome 1.

The pharmaceutically acceptable carrier according to the invention is any compatible non-toxic substance suitable for administering the composition 10 according to the invention to a human patient. Pharmaceutically acceptable carriers according to the invention suitable for oral administration are the ones well known by the person skilled in the art, such as tablets, coated or non-coated pills, capsules, spray-gas, 15 patches, gels, solutions or syrups. Pharmaceutically acceptable carriers vary according to the mode of administration (intravenous, intramuscular, subcutaneous, parenteral, etc.), and may comprise also adjuvants well known by the person skilled in the art to increase, reduce 20 and/or regulate humoral, local and/or cellular response of the immune system.

The pharmaceutical composition according to the invention may be prepared by the methods, generally applied by the person skilled in the art in the preparation 25 of various pharmaceutical compositions, wherein the percentage of the active compound/pharmaceutically acceptable carrier can vary within very large ranges, only limited by the tolerance of the patient to said pharmaceutical composition, and wherein the limits are 30 particularly determined by the frequency of administration and the possible side-effects of the active compounds or its pharmaceutically acceptable carrier.

Another aspect of the invention is related to the use of the diagnostic device according to the invention for performing upon the patient or upon a biological fluid obtained from the patient, a diagnosis, a dosage and/or a  
5 monitoring of the above-mentioned injuries or diseases or oxidative stress-related disorders affecting the patient.

A further aspect of the present invention is related to a cell or a non-human animal, preferably a mammal such as a mouse or a rat, transformed by the vector  
10 according to the invention and overexpressing the polypeptide according to the invention, or a non-human animal, preferably a mammal such as a mouse or a rat, genetically modified by a partial or total deletion of its genomic sequence encoding the polypeptide according to the  
15 invention (knock-out non-human mammal) and obtained by methods well known by the person skilled in the art such as the one described by Kahn et al. (*Cell*, Vol. 92, pp. 593-596 (March 1998)).

Other examples of genetically modified non-  
20 human animals according to the invention may be a transgenic non-human animal comprising an inhibitor according to the invention, preferably an antisense nucleic acid sequence complementary to the nucleotide sequence according to the invention so placed as to be transcribed  
25 into antisense mRNA which is complementary to the nucleotide sequence according to the invention and which hybridises to said nucleotide sequence, thereby reducing or blocking its translation.

Further aspects of the present invention will  
30 be described in the enclosed non-limiting examples in reference to the following Figures.

**Brief description of the drawings**

- Figure 1 represents a dot blot analysis of mRNA encoding the polypeptide according to the invention in various types of human tissues.
- 5 Figure 2 represents a Northern blot analysis of mRNA encoding the polypeptide according to the invention in a rat lung after administration of lipopolysaccharides (LPS) inducing an inflammatory reaction of the lung.
- 10 Figure 3 represents a Northern blot analysis of mRNA encoding the polypeptide according to the invention in a rat lung after intraperitoneal injection of pneumotoxicants.
- Figure 4 is a schematic representation of the human genomic sequence, the complete cDNA sequence and the corresponding amino acid sequence.
- 15 Figure 5 represents respectively the alignment of the sequences of the human B18 polypeptide according to the invention with the corresponding rat and mouse sequences.
- 20

**Example 1 : Homology between the B18 polypeptide according to the invention with other known nucleotide or amino acid sequences**

- 25 The BLAST 2.0 software (gapped BLAST at the NCBI Internet site) was used for searching for homologies between human B18 (162 amino acids) and known polypeptides in databases (GenBank, SwissProt). Said search did not give perfect alignment with known peptides from different
- 30 species (Table 1). Homologies of the human B18 cDNA (805 nucleotides) with GenBank, EMBL, DDBJ and PDB deposited

nucleotide sequences (Table 2) and GenBank Expression Sequence TAGS (ESTs) were noted.

**Table 1 :** Homologies of the B18 proteins (162 amino acid) with other proteins

Name	NCBI ID	Identity (%) Homology (%)
Membrane protein (synechocystis sp.)	1652859	57/129 (44%) 81/129 (62%)
Peroxisomal-like protein (Aspergillus fumigatus)	2769700	56/176 (31%) 90/176 (50%)
Haein HI0572 hypothetical protein (Haemophilus influenzae)	1723174	53/146 (36%) 80/146 (54%)
PMP20 (Schizosaccharomyces pombe)	AJ002536	54/161 (33%) 85/161 (52%)
Peroxisomal membrane protein A (PMP 20) (Candida boidinii)	130360	59/170 (34%) 89/170 (51%)
Peroxisomal membrane protein B (PMP 20) (Candida boidinii)	130361	58/170 (34%) 88/170 (51%)
Putative peroxisomal protein PMP from yeast (Saccharomyces cerevisiae)	1709682	41/138 (29%) 72/138 (51%)
Alkylhydroperoxide reductase C22 protein (Escherichia coli)	P26427	36/126 (28%) 58/126 (45%)

**Table 2**

Name	Access NO	Identity
Human mRNA down-regulated in cells infected by adenovirus 5	U82616	259/263 (98%)
Human mRNA down-regulated in cells infected by adenovirus 5	U82615	300/321 (93%)

In the Table 2, an identity of 98% has been obtained with the alignment of 259 nucleotides of CDNA B18, which comprises in its totality 805 nucleotides, with 263 nucleotides of U82616 CDNA. A similar identity has been  
5 obtained with the U82615 sequence.

The sequence SEQ ID NO 1 comprising 805 nucleotides presents a homology with several EST sequences obtained from a human and from a mouse, having the following references :

10 **Human :**

AA130751, N42215, W38597, N91311, N68467, AA187737,  
N68916, W00593, R88950, AA181884, H20154, H66666

**Mouse :**

AA220019, AA123351, AA087129, AA255021, AA249897, W71344

15

**Example 2 : Tissue detection**

A human RNA master Blot (Clontech) containing 100-500 ng of poly-A + human RNA in each dot (normalised to the mRNA expression levels of eight different housekeeping  
20 genes) was hybridised with a 554 bp-long B18 probe labelled with  $^{32}\text{P}$ , and quantified, using Phosphorimaging Technology. As shown in Figure 1, B18 mRNA is present in all tissues examined but predominantly in trachea, lung, kidney, thyroid gland, stomach, colon, heart and some regions of  
25 the brain. Highest expression has been noted in the thyroid tissue. This presence is probably correlated with the possible antioxidant activity of the B18 polypeptide according to the invention.

30 **Example 3 : Inflammatory reaction**

Figure 2 represents a Northern blot analysis of rat lung mRNA after 6, 48 and 72 hours after

lipopolysaccharides (LPS) instillation inducing an inflammatory reaction in the lung.

A Northern blot containing 15  $\mu$ g of total RNA in each lane was hybridised with a 225 bp-long rat B18 probe, stripped and reprobated with a 572 bp-long rat  $\beta$ -actin probe, both labelled with  $^{32}$ P. Northern blot was quantified using Phosphorimaging Technology and the B18 mRNA data were normalised to  $\beta$ -actin mRNA level.

10 **Example 4 : Pneumotoxic reaction**

Figure 3 represents a Northern blot analysis of rat lung mRNA after intraperitoneal injection of pneumotoxicants (4-ipomeanol, 1-(3-furyl)-4-hydroxypentanone (IPO), methylcyclopentadienyl manganese tricarbonyl (MMT) or alpha naphthylthiourea (ANTU)). These agents are known to induce in the lung acute lesions of Clara (IPO) and alveolar cells (MMT) as well as increasing the permeability of the alveolar/blood barrier (ANTU). A Northern blot containing 15  $\mu$ g of total RNA in each lane was hybridised with a 225 bp-long rat B18 probe, stripped and reprobated with a 572 bp-long  $\beta$ -actin probe both labelled with  $^{32}$ P. The Northern blot was quantified using Phosphorimaging Technology and rat B18 mRNA data were normalised to  $\beta$ -actin mRNA level.

25

**Example 5 : Proteinic activity of the B18 polypeptide**

An amino analysis of the complete human B18 amino acid sequence shows that said polypeptide presents specific portions showing an homology with other anti-oxidant enzymes (starting from a Leucine at position 36 until a Cysteine at position 47) and an other portion

having an important homology with beta chains of ATP synthase (starting from a Glutamic acid at position 13 until a Glycine in position 38).

Furthermore, the B18 amino acid sequence according to the invention shows an important homology with an *Aspergillus fumigatus* allergen (34% identity and 60% homology by using clustal V sequence alignment), especially upon the portion of said B18 polypeptide having possible antioxidant properties. Therefore, it is possible that a peroxisomal protein (possibly homologous to B18 protein) is able to induce and to bind IgE from patients sensitised to *Aspergillus fumigatus* peroxisomal proteins after an induction of the patient immune system with *Aspergillus fumigatus* allergen. This mechanism can be compared to a reaction obtained with the manganese superoxide dismutase (MnSOD) wherein the human MnSOD is able to bind to IgE from patients sensitised to *Aspergillus fumigatus* MnSOD.

Furthermore, the Inventors have identified a portion of the B18 human polypeptide which presents an homology with a Cyclophilin-binding domain of *Candida boidinii* PMP20 (receptor, of the immuno-suppressant drug cyclosporine A). Said possible Cyclophilin-binding domain is starting from the Threonine in position 150 until the Leucine in position 161.

25

**Example 6 : B18 human gene and mRNA alternative splicing**

As represented in the enclosed Figure 4, the Inventors have identified upon the genomic DNA (SEQ ID NO 10) 5 exons and 5 introns. By RT-PCR (using primers 5'-gggtatgggactagctggcg-3' and 5'-ctggccaacattccaattgcag-3') and according to the genomic sequence, 4 different cDNAs corresponding to the transcription of the said genomic DNA

30

have been identified in human lung and in human brain. A first cDNA of 736 bp corresponds to the cDNA encoding the complete amino acid sequence of the B18 protein according to the invention. However, 3 other cDNAs of 601, 604 and  
5 469 bp were also identified, and comprise specific splicings of one or more exons.

Therefore, another aspect of the present invention is related to said specific portions of the complete genomic or CDNA nucleotide sequence according to  
10 the invention or to specific portions of the complete amino acid sequence of the B18 protein according to the invention, which could be used also as specific markers of the B18 activity, preferably the anti-oxidative activity.

15 **Example 7 : Knock-out mouse**

Exons of a mouse genomic sequence encoding the B18 polypeptide according to the invention have been deleted by homologous recombination. Said homologous recombination has been obtained with a genetic sequence  
20 comprising a neomycin resistant gene. The targeting vector with said gene and a ,thymidine kinase (in order to eliminate non-homologous recombinants with ganciclovir) has been prepared. Said recombination was used for the deletion of one or more exons of the B18 polypeptide. After  
25 electroporation of ES cells with the targeting vector, positive clones having incorporated homologous recombination were identified by Southern blot with labelled probes. Aggregation of said positive clones with a morula from a Swiss pseudo-pregnant mouse produces several  
30 chimeric mice which survive after birth. Several homozygote mice are obtained by cross-breeding and are used as a model for the above-mentioned diseases.

Similar experiments may be done with another mammal whose B18 sequence is known (the B18 sequence of a mouse and a rat and their alignment with the human sequence is shown in the enclosed Figure 5).

5

**Example 8 : Chromosome localisation of human B18 gene**

Radiation hybrid clones (GeneBridge 4 Radiation Hybrid Panel, Research Genetics) were used for performing chromosome localisation by PCR with two  
10 different pairs of primers (5'-caggttcaccttggtccctggctc-3' (SEQ ID NO 14), 5'-atgttatgcaaccctttgcgacac-3' (SEQ ID NO 17) and 5'-gtgtttgaaggggagccagggaac-3' (SEQ ID NO 18), 5'-agagacagggtttcaccatcttgg-3' (SEQ ID NO 19)).

The Inventors have located B18 genomic  
15 sequence on human chromosome 11q13. B18 gene has been located 7.15-6.1 cR from marker D11S913 between markers D11S1963 and D11S4407 (Genome Database internet site).

Unknown genes linked to different disorders have been localised in the same region of chromosome 11.  
20 Therefore, B18 gene is possibly associated with these disorders:

- atopy (atopic hypersensitivity: asthma, hay fever and eczema; MIM n°147050 at OMIM of NCBI internet site),
- high bone mass syndrome (MIM n°601884),
- 25 - osteopetrosis (MIM n°259700),
- osteoporosis-pseudoglioma syndrome (MIM n°259770) and
- Bardet-Biedl syndrome 1 (MIM n°209901).

CLAIMS

1. Amino acid sequence having more than 70% homology with the sequence SEQ ID NO 2.
2. Amino acid sequence according to claim 1,  
5 having more than 85% homology with the sequence SEQ ID NO 2.
3. Amino acid sequence according to claim 1 or 2, having more than 95% homology with the sequence SEQ ID NO 2.
- 10 4. Amino acid sequence according to any one of the preceding claims, corresponding to SEQ ID NO 2 or an immunoreactive portion thereof.
5. Nucleotide sequence encoding the amino acid sequence according to any one of the preceding claims  
15 and presenting more than 70% homology with SEQ ID NO 1 or its complementary strand.
6. Nucleotide sequence according to claim 5, having more than 85% homology with the sequence SEQ ID NO 1 or its complementary strand.
- 20 7. Nucleotide sequence according to claim 5 more than 95% homology with the sequence SEQ ID NO 1 or its complementary strand.
8. Nucleotide sequence according to any one of the claims 5 to 7, corresponding to the sequence SEQ ID  
25 NO 1, its complementary strand or a portion thereof specific for SEQ ID NO 1 and comprising more than 15 base pairs.
9. Vector comprising the nucleotide sequence according to any one of the claims 5 to 8.
- 30 10. Inhibitor directed against the amino acid or nucleotide sequence according to any one of the claims 1 to 8.

11. Inhibitor according to claim 10, being an antibody, preferably a monoclonal antibody, or a portion of said antibody.

12. Diagnostic device comprising an element  
5 selected from the group consisting of the amino acid sequence according to any one of the claims 1 to 4, the nucleotide sequence according to any one of the claims 5 to 8, the inhibitor according to claim 10 or 11, their portions or a mixture thereof.

10 13. Method for the in vitro detection of lung injuries and diseases or oxidative stress-related diseases and disorders, especially inflammatory diseases, comprising the steps of :

- 15 - isolating a sample from a body fluid of a patient, preferably a human patient,
- possibly inhibiting the contaminants present in said sample,
- put in contact said sample with an element selected from the group consisting of the amino acid sequence  
20 according to any one of the claims 1 to 4, the nucleotide sequence according to any one of the claims 5 to 8, the inhibitor according to claim 10 or 11, their portions or a mixture thereof, and
- detecting a reaction of a molecule present in said  
25 sample with said element.


14. Pharmaceutical composition comprising a pharmaceutically acceptable carrier and an element selected from the group consisting of the amino acid sequence according to any one of the claims 1 to 4, the nucleotide  
30 sequence according to any one of the claims 5 to 8, the

inhibitor according to claim 10 or 11, their portions or a mixture thereof.

15 15. Use of the pharmaceutical composition according to claim 14 for the manufacture of a medicament  
5 for the prevention and/or the treatment of lung injuries or diseases, and of oxidative stress-related diseases or disorders, such as specific cardio-vascular diseases like arteriosclerosis, neurodegenerative disorders such as  
10 Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, apoptosis and inflammatory reactions, allergic reactions such as asthma, hay fever and eczema, high bone mass syndrome, osteopetrosis, osteoporosis-pseudoglioma syndrome, and Bardet-Biedl syndrome 1.

15 16. Cell transformed by the vector according to claim 9 or comprising a partial or total deletion of its nucleotide sequence according to any one of the claims 5 to  
8.

20 17. Non-human animal, preferably a mammal, transformed by the vector according to claim 9 or comprising a partial or total deletion of its nucleotide sequence according to any, one of the claims 5 to 8.






FIG.2






FIG. 3

Gene (chromosome 11q12-13)



mRNAs



Proteins




FIG. 4





CLUSTAL V alignment of human and mouse cDNA sequences (Identity: 552/675, 81.8%):

FIG.5c

```

B18hum      GCCAGGAGGCGGAGTGGAAAGTGGCCGTGGGGCGGGTATGGGACTAGCTGG
B18mouse    -----

B18hum      CGTGTGCGCCCTGAGACGCTCAGCGGGCTATATACTCGTCGGTGGGGCCG
B18mouse    -----TGCTCCGTG-----CATCGACGTGCTTG
                    **** * * * * * * * * * *

B18hum      GCGGTCAGTCTGCGGCAGCGGCAGCAAGACGGTGCAGTGAAGGAGAGTGG
B18mouse    GCAGGCAG-----AGCAGGCCGG---AAAGAAGCAGGTTGG
                    ** * ***          **** * *** * **** * * **

B18hum      GCGTCTGGCGGGGTCCGCAGTTTCAGCAGAGCCGCTGCAGCCATGGCCCC
B18mouse    GAGTGTGGCGGAGCCCCGAGCTTCAGCAGCTCCGCGGTGACCATGGCCCC
                    * ** * * * * * * * * * * * * * * * * * * * * * * * * * *

B18hum      AATCAAGGTGGGAGATGCCATCCCAGCAGTGGAGGTGTTTGAAGGGGAGC
B18mouse    GATCAAGGTGGGAGATGCCATTCCTCAGTGGAGGTATTTGAAGGGGAAC
                    * * * * * * * * * * * * * * * * * * * * * * *

B18hum      CAGGGAACAAGGTGAACCTGGCAGAGCTGTTCAAGGGCAAGAAGGGTGTG
B18mouse    CGGGAAGAAGGTGAACCTGGCAGAGCTGTTCAAGGGCAAGAAAGGTGTT
                    * ** * * * * * * * * * * * * * * * * * * * * * * * * * *

B18hum      CTGTTTGGAGTTCCTGGGGCCTTCACCCCTGGATGTTCCAAGACACACCT
B18mouse    TTGTTTGGAGTCCCTGGGGCATTTACACCTGGCTGTTCTAAGACCCACCT
                    * * * * * * * * * * * * * * * * * * * * * * * * * * * *

B18hum      GCCAGGGTTTGTGGAGCAGGCTGAGGCTCTGAAGGCCAAGGGAGTCCAGG
B18mouse    GCCTGGGTTTGTGGAGCAAGCTGGAGCTCTGAAGGCTAAGGGAGCGCAGG
                    *** * * * * * * * * * * * * * * * * * * * * * * * * * *

B18hum      TGGTGGCCTGTCTGAGTGTTAATGATGCCTTTGTGACTGGCGAGTGGGGC
B18mouse    TGGTGGCCTGTCTGAGCGTTAATGACGTCTTTGTGATTGAAGAGTGGGGT
                    * * * * * * * * * * * * * * * * * * * * * * * * * * * *

B18hum      CGAGCCCACAAGGCGGAAGGCAAGGTTCCGGCTCCTGGCTGATCCCCTGG
B18mouse    CGAGCCCACCAGGCAGAAGGCAAGGTTCCGGCTCCTGGCTGACCCCCTGG
                    * * * * * * * * * * * * * * * * * * * * * * * * * * * *

B18hum      GGCCTTTGGGAAGGAGACAGACTTATTACTAGATGATTGCTGGTGTCCA
B18mouse    AGCCTTTGGGAAGGCGACAGACTTATTATTGGATGATTCTTTGGTGTCTC
                    * * * * * * * * * * * * * * * * * * * * * * * * * * * *

B18hum      TCTTTGGGAATCGACGTCTCAAGAGGTTCTCCATGGTGGTACAGGATGGC
B18mouse    TCTTTGGGAATCGTCCGGCTGAAAAGGTTCTCCATGGTGGTACAGGATGGC
                    * * * * * * * * * * * * * * * * * * * * * * * * * * * *

B18hum      ATAGTGAAGGCCCTGAATGTGGAACCAGATGGCACAGGCCTCACCTGCAG
B18mouse    ATAGTGAAGGCACTGAACGTGGAGCCAGATGGCACAGGCCTCACCTGCAG
                    * * * * * * * * * * * * * * * * * * * * * * * * * * * *

B18hum      CCTGGCACCCAATATCATCTCACAGCTCTGAGGCCCTGGGCCAGATTACT
B18mouse    CCTGGCCCCCAACATCCTCTCCCACTCTGAGGCCCTGG-CCAGATG---
                    * * * * * * * * * * * * * * * * * * * * * * * * * * * *

B18hum      TCCTCCACCCCTCCCTATCTCACCTGCCAGCCCTGTGCTGGGGCCCTGC
B18mouse    TCCTCTGACTCTCCC-ATCTCTCCACCCGGCTCT-----AGGCC----
                    * * * * * * * * * * * * * * * * * * * * * * * * * * * *

B18hum      AATTGGAATGTTGGCCAGATTTCTGCAATAAACACTTGTGGTTTGGCGAA
B18mouse    ----AAAAGGCTCGGTACCTCCTTACTGGGAGC-CACGT-----
                    * * * * * * * * * * * * * * * * * * * * * * * * * * * *

```

1  
SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

(A) NAME: UNIVERSITE CATHOLIQUE DE LOUVAIN  
Halles Universitaires

(B) STREET: Place de l' Universite, 1

(C) CITY: LOUVAIN-LA-NEUVE

(E) COUNTRY: BELGIUM

(F) POSTAL CODE (ZIP): B-1348

(A) NAME: UNIVERSITE DE MONS-HAINAUT

(B) STREET: Place du Parc 20

(C) CITY: MONS

(E) COUNTRY: BELGIUM

(F) POSTAL CODE (ZIP): B-7000

(ii) TITLE OF INVENTION: PEROXISOME-ASSOCIATED PEPTIDE, NUCLEOTIDE  
SEQUENCE ENCODING SAID PEPTIDE AND THEIR USES IN THE  
DIAGNOSTIC AND/OR THE TREATMENT OF LUNG INJURIES AND  
DISEASES, AND OF OXIDATIVE STRESS-RELATED DISORDERS

(iii) NUMBER OF SEQUENCES: 19

## (iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 805 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

## (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION:193..681

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

GCCAGGAGGC GGAGTGAAG TGGCCGTGGG GCGGGTATGG GACTAGCTGG CGTGTGCGCC 60

CTGAGACGCT CAGCGGGCTA TATACTCGTC GGTGGGGCCG GCGGTCAGTC TGCGGCAGCG 120





4

```

AGTGCCGCGG TGA CTATGGC CCCGATCAAG GTGGGAGACA CCATTCCCTC AGTGGAGGTA      180
TTTGRAGGGG AACCTGGAAA GAAGGTGAAC TTGGCAGAGC TGTTC AAGGA CAAGAAAGGT      240
GTTTTGTTTG GAGTCCCTGG GGCATTTACA CCTGGCTGTT CCAAGACCCA TCTGCCTGGG      300
TTTGTGGAGC AAGCCGGAGC TCYGAAGGCC AAGGGAGCAC AAGTGGTGGC CTGTCTGAGT      360
GTTAATGATG YCTTCGTGAC TGCAGAGTGG GGTCGAGCCC ACCAGGCAGA AGGCAAGGTT      420
CAGCTCCTGG CTGACCCAC TGGAGCTTTT GGAAAGGAGA CAGATTTACT ACTAGATGAT      480
TCTTTGGTGT CTCTCTTTGG GAATCGTCGG CTA AAAAGGT TCTCCATGGT GATAGACAAG      540
GGCGTAGTAA AGGCACTGAA CGTGGAGCCG GATGGCACAG GCCTCACCTG CAGCCTGGCC      600
CCCAACATCC TCTCACA ACT CTGAGGCCCT GACCAGAATG TCCTCTGACT CTCCATCTC      660
CTCCACCCAG CTCTGGGCCA AAGGCCCAGT ACCTCCTTAC CTGAGGGCCA CTGGAATGGA      720
ACCTTGACAA TATTTCTGCA ATAAACAGTT TAATTTGTGA AAAAAAAAAA AAAAAAAAAA      780
    
```

(2) INFORMATION FOR SEQ ID NO: 4:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 162 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Rattus Rattus
- (ix) FEATURE:
  - (A) NAME/KEY: Modified-site
  - (B) LOCATION:17
  - (D) OTHER INFORMATION:/product= "Glu or Gly"
- (ix) FEATURE:
  - (A) NAME/KEY: Modified-site
  - (B) LOCATION:63
  - (D) OTHER INFORMATION:/product= "Leu or Pro"
- (ix) FEATURE:
  - (A) NAME/KEY: Modified-site
  - (B) LOCATION:79
  - (D) OTHER INFORMATION:/product= "Ala or Val"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

```

Met Ala Pro Ile Lys Val Gly Asp Thr Ile Pro Ser Val Glu Val Phe
1           5           10           15
    
```

5

Xaa Gly Glu Pro Gly Lys Lys Val Asn Leu Ala Glu Leu Phe Lys Asp  
 20 25 30

Lys Lys Gly Val Leu Phe Gly Val Pro Gly Ala Phe Thr Pro Gly Cys  
 35 40 45

Ser Lys Thr His Leu Pro Gly Phe Val Glu Gln Ala Gly Ala Xaa Lys  
 50 55 60

Ala Lys Gly Ala Gln Val Val Ala Cys Leu Ser Val Asn Asp Xaa Phe  
 65 70 75 80

Val Thr Ala Glu Trp Gly Arg Ala His Gln Ala Glu Gly Lys Val Gln  
 85 90 95

Leu Leu Ala Asp Pro Thr Gly Ala Phe Gly Lys Glu Thr Asp Leu Leu  
 100 105 110

Leu Asp Asp Ser Leu Val Ser Leu Phe Gly Asn Arg Arg Leu Lys Arg  
 115 120 125

Phe Ser Met Val Ile Asp Lys Gly Val Val Lys Ala Leu Asn Val Glu  
 130 135 140

Pro Asp Gly Thr Gly Leu Thr Cys Ser Leu Ala Pro Asn Ile Leu Ser  
 145 150 155 160

Gln Leu

(2) INFORMATION FOR SEQ ID NO: 5:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 675 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Mouse
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION:99..588

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

TGCTCCGTGC ATCGACGTGC TTGGCAGGCA GAGCAGGCCG GAAAGAAGCA GGTGGGAGT 60

GTGGCGGAGC CCGCAGCTTC AGCAGCTCCG CGGTGACCAT GGCCCCGATC AAGGTGGGAG 120

ATGCCATTCC CTCAGTGGAG GTATTTGAAG GGGAAACCGGG AAAGAAGGTG AACTTGGCAG 180

AGCTGTTCAA GGGCAAGAAA GGTGTTTTGT TTGGAGTCCC TGGGGCATT ACACCTGGCT 240

GTTCTAAGAC CCACCTGCCT GGGTTTGTGG AGCAAGCTGG AGCTCTGAAG GCTAAGGGAG 300  
 CGCAGGTGGT GGCCTGTCTG AGCGTTAATG ACGTCTTTGT GATTGAAGAG TGGGGTCGAG 360  
 CCCACCAGGC AGAAGGCAAG GTTCGGCTCC TGGCTGACCC CACTGGAGCC TTTGGGAAGG 420  
 CGACAGACTT ATTATTGGAT GATTCTTTGG TGTCTCTCTT TGGGAATCGT CGGCTGAAAA 480  
 GGTCTCCAT GGTGATAGAC AACGGCATAG TGAAGGCACT GAACGTGGAG CCAGATGGCA 540  
 CAGGCCTCAC CTGCAGCCTG GCCCCAACA TCCTCTCCCA ACTCTGAGGC CCTGGCCAGA 600  
 TGTCTCTGA CTCTCCCATC TCTCCACCC GGCTCTAGGC CAAAAGGCTC GGTACCTCCT 660  
 TACTGGGAGC CACGT 675

(2) INFORMATION FOR SEQ ID NO: 6:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 162 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
 

- (A) ORGANISM: Mouse

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Met	Ala	Pro	Ile	Lys	Val	Gly	Asp	Ala	Ile	Pro	Ser	Val	Glu	Val	Phe
1				5					10					15	
Glu	Gly	Glu	Pro	Gly	Lys	Lys	Val	Asn	Leu	Ala	Glu	Leu	Phe	Lys	Gly
			20					25					30		
Lys	Lys	Gly	Val	Leu	Phe	Gly	Val	Pro	Gly	Ala	Phe	Thr	Pro	Gly	Cys
		35					40					45			
Ser	Lys	Thr	His	Leu	Pro	Gly	Phe	Val	Glu	Gln	Ala	Gly	Ala	Leu	Lys
	50					55					60				
Ala	Lys	Gly	Ala	Gln	Val	Val	Ala	Cys	Leu	Ser	Val	Asn	Asp	Val	Phe
65				70						75				80	
Val	Ile	Glu	Glu	Trp	Gly	Arg	Ala	His	Gln	Ala	Glu	Gly	Lys	Val	Arg
				85					90					95	
Leu	Leu	Ala	Asp	Pro	Thr	Gly	Ala	Phe	Gly	Lys	Ala	Thr	Asp	Leu	Leu
			100					105					110		
Leu	Asp	Asp	Ser	Leu	Val	Ser	Leu	Phe	Gly	Asn	Arg	Arg	Leu	Lys	Arg
		115					120					125			

Phe Ser Met Val Ile Asp Asn Gly Ile Val Lys Ala Leu Asn Val Glu  
 130 135 140

Pro Asp Gly Thr Gly Leu Thr Cys Ser Leu Ala Pro Asn Ile Leu Ser  
 145 150 155 160

Gln Leu

## (2) INFORMATION FOR SEQ ID NO: 7:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 469 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:  
 (A) ORGANISM: Homo sapiens

(ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION:161..382

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

GGGTATGGGA CTAGCTGGCG TGTGCGCCCT GAGACGCTCA GCGGGCTATA TACTCGTCGG	60
TGGGGCCGGC GGTCACTCTG CGGCAGCGGC AGCAAGACGG TGCAGTGAAG GAGAGTGGGC	120
GTCTGGCGGG GTCGCAGTT TCAGCAGAGC CGCTGCAGCC ATGGCCCCAA TCAAGTTTCG	180
GCTCCTGGCT GATCCCACTG GGGCCTTTGG GAAGGAGACA GACTTATTAC TAGATGATTC	240
GCTGGTGTCC ATCTTTGGGA ATCGACGTCT CAAGAGGTTT TCCATGGTGG TACAGGATGG	300
CATAGTGAAG GCCCTGAATG TGGAACCAGA TGGCACAGGC CTCACCTGCA GCCTGGCACC	360
CAATATCATC TCACAGCTCT GAGGCCCTGG GCCAGATTAC TTCCTCCACC CCTCCCTATC	420
TCACCTGCCC AGCCGTGTGC TGGGGCCCTG CAATTGGAAT GTTGGCCAG	469

## (2) INFORMATION FOR SEQ ID NO: 8:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 601 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION:161..514

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

```
GGGTATGGGA CTAGCTGGCG TGTGCGCCCT GAGACGCTCA GCGGGCTATA TACTCGTCGG      60
TGGGGCCGGC GGTCAGTCTG CGGCAGCGGC AGCAAGACGG TGCAGTGAAG GAGAGTGGGC      120
GTCTGGCGGG GTCCGCAGTT TCAGCAGAGC CGCTGCAGCC ATGGCCCCAA TCAAGACACA      180
CCTGCCAGGG TTTGTGGAGC AGGCTGAGGC TCTGAAGGCC AAGGGAGTCC AGGTGGTGGC      240
CTGTCTGAGT GTTAATGATG CCTTTGTGAC TGGCGAGTGG GGCCGAGCCC ACAAGGCGGA      300
AGGCAAGGTT CGGCTCCTGG CTGATCCCAC TGGGGCCTTT GGAAGGAGA CAGACTTATT      360
ACTAGATGAT TCGCTGGTGT CCATCTTTGG GAATCGACGT CTCAGAGGT TCTCCATGGT      420
GGTACAGGAT GGCATAGTGA AGGCCCTGAA TGTGGAACCA GATGGCACAG GCCTCACCTG      480
CAGCCTGGCA CCAATATCA TCTCACAGCT CTGAGGCCCT GGGCCAGATT ACTTCCTCCA      540
CCCCTCCCTA TCTCACCTGC CCAGCCCTGT GCTGGGGCCC TGCAATTGGA ATGTTGGCCA      600
G                                                                                   601
```

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 604 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION:161..517

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

```
GGGTATGGGA CTAGCTGGCG TGTGCGCCCT GAGACGCTCA GCGGGCTATA TACTCGTCGG      60
```

9

TGGGGCCGGC GGTCACTCTG CGGCAGCGGC AGCAAGACGG TGCAGTGAAG GAGAGTGGGC	120
GTCTGGCGGG GTCCGCAGTT TCAGCAGAGC CGCTGCAGCC ATGGCCCCAA TCAAGGTGGG	180
AGATGCCATC CCAGCAGTGG AGGTGTTTGA AGGGGAGCCA GGGAACAAGG TGAACCTGGC	240
AGAGCTGTTC AAGGGCAAGA AGGGTGTGCT GTTTGGAGTT CCTGGGGCCT TCACCCCTGG	300
ATGTTCCAAG GTTCGGCTCC TGGCTGATCC CACTGGGGCC TTTGGGAAGG AGACAGACTT	360
ATTACTAGAT GATTCGCTGG TGTCCATCTT TGGGAATCGA CGTCTCAAGA GGTCTCCAT	420
GGTGGTACAG GATGGCATAG TGAAGGCCCT GAATGTGGAA CCAGATGGCA CAGGCCTCAC	480
CTGCAGCCTG GCACCCAATA TCATCTCACA GCTCTGAGGC CCTGGGCCAG ATTACTTCTT	540
CCACCCCTCC CTATCTCACC TGCCCAGCCC TGTGCTGGGG CCCTGCAATT GGAATGTTGG	600
CCAG	604

## (2) INFORMATION FOR SEQ ID NO: 10:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 2710 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo sapiens
- (ix) FEATURE:
  - (A) NAME/KEY: exon
  - (B) LOCATION:2516..2710
- (ix) FEATURE:
  - (A) NAME/KEY: exon
  - (B) LOCATION:2074..2135
- (ix) FEATURE:
  - (A) NAME/KEY: exon
  - (B) LOCATION:1932..1970
- (ix) FEATURE:
  - (A) NAME/KEY: exon
  - (B) LOCATION:1728..1859
- (ix) FEATURE:
  - (A) NAME/KEY: exon
  - (B) LOCATION:802..936
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

TCTGTCCCTT	AGCGCCCCCG	CGGGGGCTTA	CCCCATCCCA	CTCCATGACC	TCCCCTCCCC	60
CCATGGCGAA	TTCCCACCTT	TCTGTCTTTC	ACTCACTTCC	TGGAACCGTC	CCCAGGGCCT	120
TGGACCTTCC	CCCTTCTCCT	CCCAAACCTT	GTGAGACCCC	ATTCCCTTTC	TACTTCATCC	180
TGCTCTCAAC	TTTTGGGCTC	CTCAGAGGCC	CTCACCCCTG	ACTCTCTCTC	CCTACCCACT	240
CTGGTCCCAT	GAAGCCCTCA	AGTACTCTGG	GGATGGATCC	TTCCCCCTTC	AAAAGATTCC	300
TTCTTTTGT	CTACACCTCC	TGGGTGTAGG	GGCCTGGACA	CCCTCCCCCA	ACGTTCCACC	360
TGCCGCTGCC	CTTCCTCTTC	CTCCTCCTGA	GGGTGGGACC	CTCAGACCTG	GCCAAGATCC	420
TCTCCCTCCA	TGTTGTGAGG	GACTCCTCCT	CACCCCCAAA	TACAGCCCTC	TAGCCCCTGT	480
CCATTTTATT	CCACTCCTTT	CCTGTAACCT	AGACAGCATG	TTATGCAACC	CTTTGCGACA	540
CATGGGGAAA	CCTTCCCTCC	CTTCCTCTGT	TGTCACCAAT	GGCCCTTAA	GAGGAGCAGG	600
GCCACCTTGA	AACTTGGAGG	ATATGGGGTA	ACCCAGTGGG	AGCGGGCAGG	GAGGGCCCTT	660
GGAAACTGAC	AGGGCTGGAG	TATCCTGCTG	GGTTTCAGCC	CCGGTTCCTG	CAGGCACAGC	720
TGCCAGGCTC	TCTGTTACAC	TTCCTGCCTC	TGGTTTGCCC	CGGCTCCCTC	ACCCCCCTTA	780
CCCTGGAGTC	CTTCCTTCTA	GGTGGGAGAT	GCCATCCCAG	CAGTGGAGGT	GTTTGAAGGG	840
GAGCCAGGGA	ACAAGGTGAA	CCTGGCAGAG	CTGTTCAAGG	GCAAGAAGGG	TGTGCTGTTT	900
GGAGTTCCTG	GGCCCTTAC	CCCTGGATGT	TCCAAGGTGA	GGCCCTTCCC	CTTCTGAAGA	960
TCAGGACCTG	GGGATCTTTT	GTGTTGCTCT	TAAGTCTCC	ACATAGTCCT	GATAGGACTC	1020
CTAAAAAGCA	TTTCAGTGCC	ATCACAAAAC	AAGTAGAGCT	GGGTAGAGCT	GGGCGCGGTG	1080
GCTCACGCCT	GTAATCCCAG	CACTTTGGGA	GGCCAAGGCG	GGTGGATCAC	GAGGTCAGGA	1140
GTCCAAAACC	AGCCTGGCCA	AGATGGTGAA	ACCCTGTCTC	TACTAAAAAT	GCAAAAAAAT	1200
CAGCCGGATA	TGGTGGCGGG	CGCCTGTAAT	CCCAGGTATT	GGGGAGGCTG	AGGCAGAGAA	1260
TTGCTTGAAC	CCAGGAGGCG	TAGGTTGCAG	TGAGTGGAGA	TCGTGCCTCT	GCAGTCCAGC	1320
CTGGGTGAAA	GAGCGAGACT	CCGTCTCAA	ATGAAAAAAA	AAAAAGAAAA	CAAGTAGAGA	1380
CTGCAAAAAG	GGAACAGTAC	CGGGAATGTT	GGAGAAAAAC	ATACTACAAT	TAAATCCAAC	1440
ACCCCTGTTG	GTCCTGCTAA	ATGACAGGCA	CTGTGGAAGG	TGCTTGGGAC	TCAGATAAAT	1500
AAGACAAAGA	TCTGCCCATG	GAAAGTTCAC	GTCTGGACCA	TAAGGCATTA	GGTTTCATTC	1560
TGAGCTTCCCT	AGTGGCCAAG	GCAAAAAGGA	AATAGAATGG	TTAGACAGC	TCTCATTGTC	1620
TGATCAAAGG	TGTTGAGGCA	GAGCACTGAG	GAGGGCCTGG	AGATAAAGGG	TGGGCTGGGG	1680
GTCAGATGCA	GTTATCCCTT	TGCCGACCCT	TTGTTCCCTT	TCCTCAGACA	CACCTGCCAG	1740
GGTTTGTGGA	GCAGGCTGAG	GCTCTGAAGG	CCAAGGGAGT	CCAGGTGGTG	GCCTGTCTGA	1800
GTGTTAATGA	TGCCTTTGTG	ACTGGCGAGT	GGGGCCGAGC	CCACAAGGCG	GAAGGCAAGG	1860

TGAGGTGAGG GGCCTGCAGG GAGTCAGGAC CAGGTAGGAT ATTCTTCTTG TGACCTCTAC	1920
TTTCTCTGCA GGTTCGGCTC CTGGCTGATC CCACTGGGGC CTTTGGGAAG GTGAGTGTTT	1980
CCCTGACCGC CACAGGGACA TGGCGGTGCG GGGAGCAGTG GGGGCCCTTG GCCTCTTCAA	2040
GGATTTCTGA CACTTTTCTC TGTCTCTTCT TAGGAGACAG ACTTATTACT AGATGATTCTG	2100
CTGGTGTCCA TCTTTGGGAA TCGACGTCTC AAGAGGTAAG AGTGGAGAGT CCTCTGTGGA	2160
GAAAGTCCTC TGTGGGAGAG AGTCCTCTGT GGGAGAGAGT CCTCTGTGGA GAGGGTCCTC	2220
TGTGGGAAGA GTCGTCTGTG GGGGAGATGT GTGGGAGAGA GTCCTGTGTG GGGAGAGTCT	2280
TCTGTAGGGG AGAGTCCTCT GGGGAGAGAG TCCTGTGTGG GGGAGAGTCC TCTGTGGGGA	2340
GAGTCCTCTG TGTGGAGAGA GTCCTGTGTG GTGGTGAGTC CTCTGTGGGG GAGAGTCCTC	2400
TGTGGGGGGA GTCCTCTCTG GAGTTCTCTT GGGCCCCTGG CTGTTCACTG CCTGTCTCCA	2460
TGCCCAGCCT CCAAGCCCAG GCTGATGCAG CTGGCTGGGC CCCTCTTTCC GGCAGGTTCT	2520
CCATGGTGGT ACAGGATGGC ATAGTGAAGG CCCTGAATGT GGAACCAGAT GGCACAGGCC	2580
TCACCTGCAG CCTGGCACCC AATATCATCT CACAGCTCTG AGGCCCTGGG CCAGATTACT	2640
TCCTCCACCC CTCCCTATCT CACCTGCCCA GCCCTGTGCT GGGGCCCTGC AATTGGAATG	2700
TTGGCCAGAT	2710

(2) INFORMATION FOR SEQ ID NO: 11:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 25 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

GCCATCCCAG CAGTGGAGGT GTTTG 25

(2) INFORMATION FOR SEQ ID NO: 12:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 24 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

## 12

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

TTGAACAGCT CTGCCAGGTT CACC

24

(2) INFORMATION FOR SEQ ID NO: 13:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 24 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

TGGAGGTGTT TGAAGGGGAG CCAG

24

(2) INFORMATION FOR SEQ ID NO: 14:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 24 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

CAGGTTACC TTGTTCCCTG GCTC

24

(2) INFORMATION FOR SEQ ID NO: 15:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

GGGTATGGGA CTAGCTGGCG

20

(2) INFORMATION FOR SEQ ID NO: 16:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 22 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

CTGGCCAACA TTCCAATTGC AG

22

(2) INFORMATION FOR SEQ ID NO: 17:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

ATGTTATGCA ACCCTTTGCG ACAC

24

(2) INFORMATION FOR SEQ ID NO: 18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

GTGTTTGAAG GGGAGCCAGG GAAC

24

(2) INFORMATION FOR SEQ ID NO: 19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

AGAGACAGGG TTCACCATC TTGG

24