



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61B 5/055		A1	(11) International Publication Number: WO 00/25671 (43) International Publication Date: 11 May 2000 (11.05.00)
(21) International Application Number: PCT/US99/24993 (22) International Filing Date: 26 October 1999 (26.10.99)			(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(30) Priority Data: 09/184,835 2 November 1998 (02.11.98) US			
(71) Applicant: EAGLE VISION PHARMACEUTICAL CORPORATION [US/US]; North Commercial Unit, 531 Pottstown Pike, Chester Springs, PA 19425 (US).			
(72) Inventors: SEOANE, Peter, R.; 602 Indigo Lane, Downingtown, PA 19335 (US). HARNISH, Phillip, P.; 175 Krauser Road, Downingtown, PA 19335 (US). VESSEY, Adele, R.; Conestoga Road, Elverson, PA 19520 (US).			Published <i>With international search report.</i>
(74) Agent: HANSEN, Philip, E.; Heslin & Rothenberg, P.C., 5 Columbia Circle, Albany, NY 12203 (US).			
(54) Title: MANGANESE COMPOSITIONS AND METHODS FOR MRI			
(57) Abstract			
An MRI contrast medium of improved safety and efficacy is disclosed. The composition includes a source of calcium ions and a source of manganese ions in a ratio of from 2:1 to 40:1 in a vehicle suitable for parenteral administration. A method of enhancing an MRI signal in a mammalian tissue with the foregoing composition is also provided.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

MANGANESE COMPOSITIONS AND METHODS FOR MRI

Field of the Invention

The invention relates to compositions and methods for enhancing magnetic resonance images of tissues, systems and organs.

Background of the Invention

5 Magnetic resonance imaging enables noninvasive visualization of tissues and organs of the human body. Contrast in the images generated may be enhanced through the use of an agent that alters the relaxation of water in the tissues of interest relative to bulk water. Species with unpaired electrons, such as the paramagnetic transition and lanthanide metal ions, may be used for this purpose.

10 Manganese chloride was investigated as a contrast agent by Lauterbur and by Wolf in animal models. Both investigators demonstrated significant image enhancement of the liver and other organs (but not blood) through the use of manganese chloride, but determined that the potential clinical utility of the agent was limited by acute cardiac toxicity. Development of contrast agents based on other paramagnetic metal ions is similarly constrained by toxicity and solubility. For instance, gadolinium chloride, acetate and sulfate demonstrate poor tolerability, including symptoms of heavy metal poisoning and accumulation of gadolinium in the liver, spleen and bone.

15

Chelates of paramagnetic transition and lanthanide metals have been used with some success in diagnostic imaging to overcome both the toxicity and solubility problems. (See US patent 4,647,447.) Application of this technology has enabled the development of several gadolinium based MR contrast agents including Gd-DTPA (MagnevistTM, Schering), Gd-DTPA-BMA (OmniscanTM, Nycomed Amersham PL), Gd-HP-DO3A (ProhanceTM, Bracco Diagnostics) and Gd-DOTA (DotaremTM, Guerbet), as well as the manganese-based contrast agent MnDPDP (TeslascanTM, Nycomed Amersham PL). There are, however, drawbacks associated with the use of chelation to solve problems of toxicity. Metals ions are not

irreversibly held in chelate complexes, but are subject to equilibrium between bound and free states. *In vivo*, this equilibrium is further complicated by equilibria between the chelator and endogenous metal ions as well as between the paramagnetic metal ion and endogenous ligands. Greis, in US Patent 5,098,692 and Bosworth, in US 5 patent 5,078,986 disclose the use of excess chelator to minimize the amount of free metal ion in chelate-based diagnostic compositions. Nonetheless, the potential for the dissociation of metal ion from the complex remains. The free chelator, either excess or released, may introduce additional toxicity on its own or through chelation of those endogenous metal ions that are required as cofactors for essential 10 enzymes or for other biological functions. Unfortunately, chelates also demonstrate reduced solution relaxivity relative to the free metal ions. The interaction of paramagnetic metal ions with water molecules, which shortens proton relaxation time relative to bulk water and gives rise to signal enhancement, is obtunded by chelation, since the same sites of metal-water interaction are used to form the non- 15 covalent associations between metal and chelate. Thus chelation provides safety at the price of reduced imaging efficacy (compared to the free metal ion). In practice this loss of efficacy may be as high as 60 -80%.

Manganese chelate image enhancement agents are known: e.g. MnDPDP, MnDTPA, MnEDTA and derivatives, Mn porphyrins such as MnTPPS₄, and fatty 20 acyl DTPA derivatives. These manganese chelates are not known to bind to endogenous macromolecules, as is the case for manganese ion. As a consequence, the enhanced efficacy seen for Mn ion following macromolecular association is seen for Mn chelates only as a function of the rate at which and the extent to which the manganese ion dissociates from the complex. This results in the need for increased 25 dosing of Mn chelates relative to free Mn ion. The dose must be additionally increased to make up for losses due to renal excretion of the chelate during the time course of the diagnostic examination. In a variation on chelation, Quay (European patent application 308983) has described the use of manganese amino acid coordination complex solutions. This application also discusses the addition of

calcium ions to the manganese amino acid solutions at levels up to 0.75 mole equivalents relative to manganese.

About ten years ago, Schaefer et al investigated a mixture of Mn⁺⁺ and Ca⁺⁺ salts in the form of manganese gluconate and calcium gluconate in a one to one mole ratio, administered intravenously to dogs, for cardiac perfusion imaging. Although the agent discriminated normally perfused from ischemic tissue, Schaefer et al also noted acute cardiotoxicity similar to that seen with manganese chloride alone. The authors suggested that a possible way around the observed adverse cardiac effects might be to employ a chelate rather than a salt of manganese. No further studies employing manganese gluconate and calcium gluconate or other salts or complexes providing Mn⁺⁺ and Ca⁺⁺ in a higher ratio than one- to-one have subsequently appeared..

US patents 5,525,326 and 5,716,598 describe oral manganese formulations for imaging of the gastrointestinal tract and for liver imaging; the latter takes advantage of the fact that the blood supply from the GI tract passes through the liver, which removes the manganese from the blood stream prior to return of the blood to the heart. Additional oral agents have been investigated, including manganese polymers, manganese impregnated molecular sieves, manganese clays and foodstuffs with high manganese content, such as blueberry juice. In general, cardiovascular safety is achieved for these agents at the expense of limiting the diagnostic utility of the agents to MR examination of the GI tract and in some cases, the liver. The administration of manganese in nanoparticulate form has been described in US patent 5,401,492. Other particulate approaches include sequestration of Mn compounds in liposomes and metal clusters, such as manganese oxalate and manganese hydroxyapatite. Particulate agents are useful for a limited number of diagnostic applications, namely, the gastrointestinal tract and organs, such as the liver and the spleen, that are involved in the uptake and sequestration of blood borne particles.

Thus it would be useful to have an agent for diagnostic imaging of tissues, systems and organs, particularly in humans, that would increase the contrast in an MR image without giving rise to problems of toxicity. It would also be useful to have an agent that could be employed for a wide variety of tissues, systems and organs that are physiologically remote from the gastrointestinal tract.

Summary of the Invention

This need is satisfied, the limitations of the prior art overcome, and other benefits realized in accordance with the principles of the present invention, which in one aspect relates to a diagnostic composition comprising a source of a 5 diagnostically effective quantity of Mn⁺⁺ ion, a source of Ca⁺⁺ ion and a pharmaceutically acceptable carrier for parenteral administration. The Ca⁺⁺ ion is present in a molar ratio of from 2:1 to 40:1 with respect to the Mn⁺⁺ ion. Preferred sources of Mn⁺⁺ are manganese salts, such as manganese acetate, chloride, gluconate, gluceptate, lactate and sulfate or mixtures thereof. Manganese gluconate 10 or manganese gluceptate are most preferred sources of Mn⁺⁺. Preferred sources of Ca⁺⁺ are calcium salts, such as calcium acetate, chloride, gluconate, gluceptate, and lactate or mixtures thereof. Calcium gluconate and calcium gluceptate are most 15 preferred sources of Ca⁺⁺. The molar ratio of calcium to manganese is preferably from 4:1 to 20:1 and most preferably from 8:1 to 10:1. One embodiment of the 20 composition aspect of the invention is a unit dosage form comprising a salt of manganese containing from 5 mg to 200 mg of manganese, a salt of calcium containing from 20 mg to 3g of calcium, and a vehicle suitable for parenteral injection.

In another aspect, the invention relates to a method for enhancing a 25 magnetic resonance image of a mammalian tissue, organ or system. The method comprises administering to a mammal a diagnostically effective amount of a source of Mn⁺⁺ ion together with from 2 to 200 molar equivalents of a source of Ca⁺⁺

ions. The preferred sources of Mn⁺⁺ and Ca⁺⁺ are as before. The sources of manganese and calcium may be administered intravenously at 1 μ mol of Mn⁺⁺ per kg body weight to 100 μ mol of Mn⁺⁺ per kg body weight and 2 μ mol of Ca⁺⁺ per kg body weight to 1400 μ mol of Ca⁺⁺ per kg body weight. The method is
5 applicable to the visualization of liver, kidney, pancreas, adrenal glands, heart, brain, salivary glands, gastrointestinal mucosa, uterus, tumors, the biliary system and the circulatory system. The source of Mn⁺⁺ and the source of Ca⁺⁺ may be administered simultaneously or separately, with Ca administration preceding Mn administration by up to 30 minutes. The method allows one to assess the metabolic
10 activity of a tissue, distinguishing hyperactive or hypoactive states from normal. The method also allows one to assess tissue vascularity, blood flow and tissue perfusion.

Brief Description of the Drawings

Numerous advantages and features of the present invention will become
15 readily apparent from the following detailed description of preferred embodiments, the appended claims and the accompanying drawings wherein:

Figure 1 is a graph of normalized systolic blood pressure in which the change from baseline (in percent) is plotted as a function of time (in minutes);

20 Figure 2 is also a graph of normalized systolic blood pressure in which the change from baseline (in percent) is plotted as a function of time (in minutes); and

25 Figure 3 is a graph of the ratio of MRI signal-to-noise as a function of time (in minutes) following administration of a composition of the invention.

Detailed Description of the Invention

Although preferred embodiments of the invention are described below, it should be understood that the present disclosure is to be considered as an exemplification of the principles of this invention and is not intended to limit the invention to the embodiments illustrated.

We have found that, for the case of manganese salts, formulation and/or administration with sources of calcium (II) results in a contrast agent with significantly improved safety, as long as the ratio of calcium to manganese is at least 2:1. Although in principle one could use any ratio higher than 2:1, in practice, 10 however, the ratio of Ca ion to Mn ion employed for any given dose of Mn ion will be limited by Ca toxicity. For example, assuming LD50 values following rapid IV administration of $138\mu\text{mol/kg}$ and $1819\mu\text{mol/kg}$ for Mn ion and Ca ion respectively, for low doses of Mn, one may contemplate the use of quite large molar excesses of Ca ion before toxicity is seen. Thus at a Mn ion dose of $1\mu\text{mol/kg}$, a Ca ion to Mn 15 ion ratio of as high 1800:1 could be considered without exceeding the Ca ion LD50; more conservatively, a ratio of 180:1 could be contemplated by approaching $1/10^{\text{th}}$ of the Ca ion LD50. If, however, the Mn ion dose is increased to $10\mu\text{mol/kg}$, the ratio of Ca to Mn will not exceed 18:1 if the conservative approach of remaining below $1/10^{\text{th}}$ of the Ca ion LD50 is followed. In practice, it is not necessary to 20 approach maximal amounts of added Ca to achieve a significant safety benefit. The particular ratio of Ca ion to Mn ion depends on the indication, the dose required to achieve efficacy and the desired safety margin (therapeutic index) for the Mn/Ca composition. As a consequence of the invention, several advantages are realized. For example, improved utilization of the paramagnetic properties of manganese is 25 achieved without the reduction in relaxivity or loss of agent through renal excretion associated with chelation. Manganese chelates demonstrate different pharmacokinetics and pharmacodynamics for the manganese metal and the chelator components, suggesting dechelation. In such cases, one may see toxicity mediated

through both the metal and the chelator. In contrast, in the present invention, levels of Mn ion that would otherwise not be considered to be physiologically tolerable may be safely reached.

The compositions and methods of the invention are of utility in imaging a variety of metabolically active organs, in particular the liver, kidney, pancreas, heart, and adrenal glands. Manganese contrast enhancement is also useful for imaging of the gastrointestinal mucosa, uterus, salivary glands, brain, the biliary tree, and tumors. Although applicants do not intend that the invention be restricted to a particular theory, it appears from the results that a macromolecular association between Mn ion and one or more native body protein(s) / component(s) is responsible for the superior relaxivity noted. It is known that Mn ion binds to human serum albumin, alpha.2-macroglobulin, transferrin, and other blood proteins with a concomitant increase in Mn relaxivity due to macromolecular association. This relaxivity increase, coupled with the increased concentration of free manganese in the blood available for macromolecular association through the use of the compositions of this invention, provides an increase in signal intensity sufficient to allow even vascular imaging at biologically tolerable doses of Mn++. In fact, on an equimolar basis, the Mn++/Ca++ compositions of the invention are approximately 20-fold more effective than the current gold standard gadolinium chelates in enhancing MR signal intensity in the blood, without suffering from contrast agent extravasation and rapid signal to noise degradation typical of extracellular fluid agents.

The compositions of the present invention may be used in place of manganese chelates to detect and assess myocardial ischemia and reperfusion using the technique described for the chelates in WO 99/01162. Following administration, Mn is taken up by mitochondria, and Mn uptake correlates with dose across the clinically relevant range of dosing. Manganese uptake also correlates with blood flow to and metabolic activity of the relevant tissues. As a

result, tissue uptake of Mn allows for the use of the compositions of the invention as viability markers (distinguish viable from non-viable tissue). The vascular enhancement due to Mn binding to plasma proteins also aids in defining blood flow and tissue perfusion (mL of blood flow per gram of tissue). Taken together the 5 vascular and tissue phases of MR imaging with the Mn/Ca compositions of the invention allow for evaluation of the status of a variety tissues, organs and organ systems.

Of the many alternative diagnostic uses for these compositions, evaluation of myocardial ischemia is particularly noteworthy. The compositions of the invention 10 provide diagnostic utility for: (1) characterization of viable vs. non-viable tissue including the myocardium and tumors; (2) sizing of infarcts; (3) characterization of ischemic tissue or tissue at risk including the myocardium (reversible vs. irreversible damage); (4) assessment of tissue perfusion; and (5) characterization of vascular lesions proximal to areas of ischemia/infarct. With this knowledge, one may 15 establish the need for and planning of interventional revascularization procedures (PTCA, stenting, by-pass); one may predict the likelihood of success of interventional procedures (salvage of at risk tissue); one may evaluate and follow up treated patients (including for reperfusion); and one may assess the effect of therapy, including new drug therapy.

20 The compositions of the invention are also useful for the assessment of organ function before and after organ transplant, for example of heart, kidney and liver; for characterization of tumor (vascularity, metabolic activity, patterns of enhancement); for the assessment of liver disease and liver abnormalities, such as tumors (particularly to distinguish hepatocellular vs. non-hepatocellular tumor and 25 benign lesions), cirrhosis and hemangioma; and for characterizing tissue vascularity (e.g., angiogenesis, tumor) and vascular lesions. One may thereby establish the need for therapy (including but not limited to drug, genetic, surgical, revascularization) and evaluate the response to therapy on ischemic tissue, vascular

lesions or tumor (including but not limited to surgery, revascularization, transplantation, drug therapy, genetic therapy, new drug evaluation, radiation therapy, chemotherapy, tissue ablation therapy).

An improved therapeutic ratio is demonstrated through an increase of the

5 LD₅₀ for Mn ion from 138 μ mol/kg to 220 μ mol/kg on administering a 10:1 Ca⁺⁺/Mn⁺⁺ composition. It appears that, although unnecessary, toxicity modifiers could be used with the compositions of the invention. Thus one could contemplate, for example, liposomal sequestration, etc, where there is the potential for equilibration between bound and free metal ion *in vivo*. In addition, various rates of

10 administration could provide different advantages. For example, by effectively reducing the instantaneous Mn concentration, slower administration of the agents described in this invention may further enhance the therapeutic (diagnostic) index of the agents or allow for the safe administration of larger doses. Thus the

15 compositions of the invention may be administered intravenously as a bolus or as an infusion over a period of time. Commonly, though not necessarily, the infusion will be over a period of 1 minute to 30 minutes. Larger doses improve the imaging of organs, such as the heart, that take up manganese less efficiently than does liver.

Similarly, since Mn is effectively and efficiently cleared from the blood, primarily by the liver, one may slow the rate of administration to increase the duration of signal

20 intensity enhancement of blood without increasing total dose.

In a preferred embodiment, Mn gluconate /Ca gluconate (1:8), is parenterally administered over periods ranging from 10 seconds to 20 minutes. Dosing is related to target organ of interest and may range from 1 μ mol/kg body weight to 100 μ mol/kg body weight of a source of Mn⁺⁺ ion together with

25 from 2 μ mol/kg body weight to 1400 μ mol/kg body weight of a source of Ca⁺⁺ ions. Preferably the source of manganese is administered at 2 μ mol/kg body weight to 30 μ mol/kg body weight and the source of calcium is administered at 4 μ mol/kg body weight to 400 μ mol/kg body weight. Most preferably the source of

manganese is administered at 3 $\mu\text{mol}/\text{kg}$ body weight to 15 $\mu\text{mol}/\text{kg}$ body weight and the source of calcium is administered at 6 $\mu\text{mol}/\text{kg}$ body weight to 200 $\mu\text{mol}/\text{kg}$ body weight. MRI is performed, according to methods well known to those skilled in the art, from during or immediately post dosing to 24 hours post dosing (vascular indications excepted). The rate of administration may be varied to further improve the cardiovascular tolerability of the contrast agent without an adverse effect on image quality, to increase the duration of the vascular phase of the agent, or to increase the dose without reducing the therapeutic index of the agent in order to enable imaging of target organs that accumulate manganese less efficiently than does liver.

Advantages of the invention include: (1) Less of the administered dose is excreted renally (essentially none vs. 15 or so percent for MnDPDP); this effectively reduces the dose of Mn required for optimal image enhancement; (2) The solution relaxivity of manganese salts is maintained; (3) The risk of toxicity associated with the presence of chelator is avoided. For instance, loss of bioactive metals such as plasma Zn through chelation and renal excretion is avoided; (4) Rapid tissue enhancement is achieved relative to Mn chelates, allowing for earlier imaging; (5) Additional indications, for example imaging of tumors, as well as vascular and cardiac imaging are possible; and (6) The cost of the composition is lower than most chelate compositions and most compositions that employ rarer elements, such as gadolinium.

The invention relates to a diagnostic composition comprising a source of a diagnostically effective quantity of Mn⁺⁺ ion, a source of Ca⁺⁺ ion and a pharmaceutically acceptable carrier for parenteral administration. The term "diagnostically effective" refers to an amount of Mn ion sufficient to increase the signal-to-noise ratio for MRI of the tissue in question. With present day instrumentation an increase of at least 5% is required to be diagnostically effective. The term "source of Mn⁺⁺ or Ca⁺⁺ ion" means any chemical species that can

furnish a measurable concentration of Mn⁺⁺ or Ca⁺⁺ ion in normal saline or blood. Thus, when the source is a simple salt, soluble at the concentration employed, such as manganese chloride, manganese gluconate, or calcium gluconate, the molar concentration of the salt will be the molar concentration of the Mn⁺⁺ or Ca⁺⁺ ion.

5 When the source is a salt or complex that is not fully dissociated at the concentration employed, the molar concentration of Mn⁺⁺ or Ca⁺⁺ ions will be less than the molar concentration of the salt or complex, but its effective molarity can be readily calculated from the solubility product of the components by methods well known in the art, and the effective molarity of the metal ion can be experimentally

10 determined with an ion-specific electrode, as well as similar methods well known in the art. It is to be noted that the concentrations of interest in the invention are those of the ions, not the concentrations of the source species, although the two are the same for simple, soluble salts. A soluble salt, for the purpose of the present invention, refers to a salt which is essentially fully dissociated at the concentration

15 being used.

Antioxidants, such as ascorbate, and stabilizers, such as calcium saccharate and borate complexes may be added to the compositions of the invention, as may other substances known in the pharmaceutical art to be useful in parenteral formulations. Formulations for parenteral administration include aqueous and non-
20 aqueous sterile injection solutions which may contain anti-oxidants, buffers, pH modifiers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient. Formulations for parenteral administration also include aqueous and non-aqueous sterile suspensions, which may include suspending agents and thickening agents. The formulations may be presented in
25 unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of a sterile liquid carrier, for example saline, water-for-injection (WFI) or the like, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders and granules according to methods well known in

the pharmaceutical art. Sterile solutions of a unit dose for injection are preferred. A preferred composition for one embodiment of the invention is a sterile solution of 25 mM manganese salt and 200 mM calcium salt in WFI adjusted to pH 6.0 to 8.2 with sodium hydroxide and/or hydrochloric acid. About 4% to 8%, preferably 6%, 5 by weight of the calcium is provided by calcium saccharate and the remainder by calcium gluconate or gluceptate.

The source of manganese is preferably administered intravenously at 1 $\mu\text{mol}/\text{kg}$ body weight to 100 $\mu\text{mol}/\text{kg}$ body weight, more preferably at 2 $\mu\text{mol}/\text{kg}$ body weight to 30 $\mu\text{mol}/\text{kg}$ body weight. In many preferred embodiments, the 10 source of manganese is administered intravenously at 3 $\mu\text{mol}/\text{kg}$ body weight to 15 $\mu\text{mol}/\text{kg}$ body weight. Similarly, the source of calcium is preferably administered intravenously at 2 $\mu\text{mol}/\text{kg}$ body weight to 1400 $\mu\text{mol}/\text{kg}$ body weight, more preferably at 4 $\mu\text{mol}/\text{kg}$ body weight to 400 $\mu\text{mol}/\text{kg}$ body weight. In many preferred embodiments, the source of calcium is administered intravenously at 6 15 $\mu\text{mol}/\text{kg}$ body weight to 200 $\mu\text{mol}/\text{kg}$ body weight.

The terms "tissue", "organ" and "system" are used in their normal sense. Thus, "tissue" refers to such biological materials as gastrointestinal mucosa, tumor and the like. "Organ" refers to such biological organs as the liver, the kidney, the pancreas, the adrenal glands, the heart, the brain, salivary glands, and the uterus. 20 "System" refers to such biological systems as the biliary system, the gastrointestinal system and the cardiovascular or circulatory system.

In accordance with the method of the invention, the source of Mn⁺⁺ and the source of Ca⁺⁺ may be administered separately or in a single composition. As will be evident to the person of skill, the administration of a single parenteral dosage 25 form will usually be simplest, but the clinician may employ any means known in the art to achieve the desired ratio of Mn⁺⁺ to Ca⁺⁺ in the individual being treated and to achieve the desired level of Mn⁺⁺ in the tissue, organ or system of interest. As

will be seen below, administration of the two ions within 30 minutes of each other will generally be successful, but it is necessary to administer the calcium first, when sequential administration is employed.

Example 1: Preparation of Manganese gluconate / Calcium gluconate 1:X

5 A 27.9mM Manganese gluconate stock solution was prepared as described below. The Mn gluconate used contained 5.5% water by assay. Ca gluconate and water for injection were added to the stock to prepare solutions of Mn gluconate and Ca gluconate of fixed molar ratio, as noted below. Without the addition of solubility enhancers, the concentration of the prepared solutions was limited to
10 approximately 3% by weight Ca gluconate. The 27.9 mM Mn stock solution was prepared by adding 6.56g of Mn gluconate to 500mL of water for injection. Because the stock solution was 27.9mM, rather than 30.0 mM, the ratios shown below in the examples, tables and figures are not exactly 1:1 etc, but they fall within 10% of the nominal ratio. The conclusions to be drawn are not substantively
15 affected by this discrepancy, and the absolute values may be recalculated by the reader if need be.

Mn/Ca 1:1 - To 50mL of stock solution add 0.646g of Ca gluconate.

Mn/Ca 1:2 - To 50mL of stock solution add 1.292 g of Ca gluconate

Mn/Ca 1:4 - To 50mL of stock solution add 50mL of water for injection and
20 2.582g of Ca gluconate.

Mn/Ca 1:6 - To 16.6 mL of stock solution add 33.3mL of water for injection and
1.29g of Ca gluconate.

Mn/Ca 1:8 - To 12.5 mL of stock solution add 38.5mL of water for injection and
1.29g of Ca gluconate.

25 Alternatively, the desired Mn/Ca compositions can be made by adding Mn gluconate to commercially available Calcium Gluconate for Injection, 10% (232mM) which contains calcium saccharate as a solubility enhancing agent. For

example: Mn/Ca 1:8 [29 mM in Mn, 232 mM in Ca(II)] Ten milliliters of Calcium Gluconate Injection, 10%, is added to 136 mg of Mn gluconate.

Example 2 Calcium Gluconate Dose Range in Rabbits

Contrast media of the invention, with molar Ca to Mn ratios of 1: 1 to 8: 1, 5 prepared according to the description in Example 1, were studied for their effects on systolic blood pressure following intravenous administration in New Zealand White Rabbits. The Mn⁺⁺ dose for all compositions was held constant at about 30 μ mol/kg, a dose previously shown to induce a minimum 20% drop in systolic blood pressure. Changes in blood pressure were monitored via an in-dwelling 10 femoral artery catheter. Blood pressure baselines were re-established between doses. Table 1 summarizes the findings in a study in which the Ca/Mn ratio was doubled with each successive dose. Table 2 summarizes findings in a second experiment focused on the more effective range of ratios studied in the exploratory experiment. The data presented in Tables 1 and 2 together demonstrate that the 15 combination of Ca and Mn in 2:1 or greater ratio exhibits weaker cardiovascular side effects than intravenously administered Mn. The effect is dose related, as measured by both peak effect and area under the curve, which reflect the extent and duration of the Mn response. Of particular note in these data is the result for the 1: 1 Mn/Ca composition, which was substantially similar to that of Mn alone. This 20 result is consistent with the findings of Schaefer et al, and stands in surprising contrast to the results obtained at higher Ca / Mn ratios.

Table 1

Normalized Systolic Blood Pressure 0-5 min Post Administration				
	Peak % Change from Baseline	% Difference Between Mn and Mn plus Ca	Area under Curve	% Difference Between Mn and Mn plus Ca
Mn (30 μ mol/kg)*	-27%	NA	-0.96	NA
1X Ca added	-25%	7%	-1.04	-8%
2X Ca added	-17%	37%	-0.70	29%
4X Ca added	-11%	59%	-0.33	66%
8X Ca added	-7%	74%	-0.24	75%

* Mean Normalized Data (n=4)

Table 2

Normalized Systolic Blood Pressure 0-15 Minutes Post Administration				
	Peak % Change from Baseline	% Difference Between Mn and Mn plus Ca	Area under Curve	% Difference Between Mn and Mn plus Ca
Mn (30 μ mol/kg)	-53%	NA	-6.45	NA
4X Ca added	-34%	36%	-4.07	37%
6X Ca added	-32%	40%	-2.86	56%
8X Ca added	-23%	57%	-1.22	81%

Figure 1 illustrates the change in normalized systolic blood pressure over time for Mn and for several Mn /Ca compositions of different molar ratios, all administered intravenously in the rabbit at a constant Mn++ ion dose of about 30 μ mol/kg. The 4:1 mixture of calcium gluconate and manganese gluconate depressed systolic blood pressure only to about half the extent seen with pure manganese gluconate; the 8:1 mixture produced very little and very short lived

depression of systolic blood pressure; 6:1 was better than 4:1 and much better than pure manganese gluconate, but not as good as 8:1 in this test.

Example 3 Rabbit Confirmatory Study

Table 3 shows data collected in a naive rabbit for Mn dosed at about 5 $30\mu\text{mol/kg}$ and Ca at $240\mu\text{mol/kg}$, individually and as a mixture, following the same protocol used in Example 2.

Table 3

Blood Pressure Effects for Mn/Ca 1:8 Compared to Mn and Ca Alone				
Normalized Systolic Blood Pressure			Normalized Mean Blood Pressure	
	Peak % Change from Baseline	Area under Curve (% Difference Between Mn and Mn plus Ca)	Peak % Change from Baseline	Area Under Curve (% Difference Between Mn and Mn plus Ca)
10	Mn ($30\mu\text{mol/kg}$)	-15%	-0.90	-18% -1.26
	Mn / Ca (1:8)	-15%	-0.135 (85%)	-12% -0.57 (55%)
	Ca ($240\mu\text{mol/kg}$)	+7%	0.43	-9% -0.10

The data demonstrate that the cardiovascular effects of the Mn/Ca combination are not predictable from the effects of either agent alone. For example, using area 15 under the curve (AUC) as a measure of both extent and duration of change in systolic blood pressure, one would predict an AUC of -0.47 (by adding 0.43 to -0.90). for the 1:8 Mn/Ca composition; in fact a significantly smaller AUC of -0.135 is seen experimentally. This is clearly visible through examination of Figure 2, in which the changes in normalized systolic blood pressure for a 1:8 mole ratio mixture 20 of Mn gluconate and Ca gluconate, a representative embodiment of this invention, are compared to the effects noted following intravenous administration of Mn or Ca

alone. As in the AUC comparison, the results for the Mn/Ca composition differ from those predictable via simple addition of the individual Mn and Ca curves.

Example 4 Mn/Ca compositions in the rat

A 1: 8 mole ratio Mn / Ca contrast agent of this invention, prepared 5 according to the description in Example 1, was studied for effects on systolic and mean blood pressure relative to a Mn gluconate control in Wistar rats. Agents were administered intravenously at a Mn dose of 100 μ mol /kg. Changes in blood pressure were monitored via an in-dwelling femoral artery catheter. The results, shown in Table 4, confirm that in the rat, as well as the rabbit, the combination of 10 Mn and Ca produces less depression of hemodynamic parameters than does Mn alone, thus permitting the safe administration of a diagnostically effective dose.

Table 4

1:8 Mn/Ca, Effect Measured by Peak Change in Pressure and Time-matched AUC's			
	Mn @ 100 μ mol/kg	Mn @ 100 μ mol/kg Ca @ 800 μ mol/kg	% Difference Between Mn and Mn plus Ca
Systolic Pressure, Peak	-36%	-18%	50%
Mean Pressure, Peak	-33%	-23%	30%
Systolic Pressure, AUC	-0.37	-0.15	59%
Mean Pressure, AUC	-0.40	-0.24	40%

Example 5 Toxicity as measured by Median Lethal Dose

Acute intravenous toxicity testing in the Swiss Webster mouse was 20 conducted on Mn gluconate, Ca gluconate and a mixture of Mn gluconate and Ca gluconate, 1: 10 on a molar basis. All agents were administered via tail vein injection over 30 seconds. For each agent The Median Lethal Dose was established via the Up and Down Method of Bruce (Fundamental and Applied Toxicology 5, 151-157 (1985)). Although the animals were observed for a period of 24 hrs,

manganese-related deaths, when they occurred, were noted within minutes of injection, presumably as the result of cardiovascular collapse. The data are shown in Table 5:

Table 5

Median Lethal Dose Studies in the Mouse		
	MLD	Protective Effect**
Mn	138umol/kg Mn	0
Mn / Ca (1:10)	220umol/kg Mn	60% Based on Mn
Ca	1819umol/kg Ca	NA

**[(MLD / Mn or Ca MLD) - 1] X 100%

10 Example 6 Efficacy Studies, Rabbit Liver Imaging

Pre and post-contrast axial T1 -weighted spin-echo and spoiled gradient echo images of the liver were obtained in two rabbits. The post contrast images were obtained up to 30 minutes after injection of Mn / Ca (1:8) (10 μ mol/kg Mn). Images were obtained on a GE Signa Imager operating at 1.5T. Signal intensity and signal-to- noise ratios were determined via region of interest (ROI) analysis. Good enhancement of the liver was seen in all cases, as shown in Table 6 below. In addition, the data presented in Figure 3, collected via ROI analysis of the liver as seen in the vascular images obtained in Example 7, show that Mn / Ca (1:8) provides fairly rapid enhancement of the liver, with measured signal to noise greater than 50% of the maximal SNR by the end of the one minute injection of contrast.

Table 6
Contrast Enhanced Liver Imaging, Mn / Ca (1:8)

Mn /Ca (1:8): Liver Imaging				
	Liver Signal Intensity		Liver Signal to Noise	
	Pre	Post	Pre	Post
Rabbit 1: T1 Spin Echo	404	635	9.4	12.0
Spoiled Gradient Echo	688	1012	13.1	16.0
Rabbit 2: T1 Spin Echo	544	769	10.1	13.9
Spoiled Gradient Echo	705	1170	6.9	15.5

Example 7 Efficacy Studies, Rabbit Vascular Imaging:

The blood pharmacokinetics of Mn / Ca (1: 8) were examined by MRI in the 10 rabbit. The contrast agent was administered intravenously in the ear vein over 1 minute at a dose of 10 μ mol/kg Mn. Blood samples (3mL each in heparinized tubes) were taken from the ear artery prior to and immediately, 2, 10, 20, 30 and 60 minutes post administration of contrast. The tubes were imaged in cross-section via T1 -weighted spin echo imaging (TR=300, TE=15,4NEX, FOV=8) on a GE Signa 15 1.5T Imager. The signal intensity data, shown in Table 7, demonstrate pharmacokinetics consistent with those found for Mn++, when measured with a 56 Mn tracer. (Borg, D.C.; Cotzias, G.C. J Clin Invest. 37:1269)

Table 7
Vascular Image Enhancement with Mn / Ca (1:8)

Mn / Ca 1:8 Evaluation of Vascular Enhancement over Time		
	Blood Signal Intensity	Percent Enhancement of the Blood
Pre-Contrast	363	0%
Immediately Post Contrast	733	102%
2 Min Post Contrast	594	64%
10 Min Post Contrast	464	28%
25 20 Min Post Contrast	419	15%

Example 8 Ex-vivo imaging - Saline vs. Whole Blood

The T1 shortening seen with Mn/Ca (1:8) in saline was compared to that seen in blood. Tubes of saline and blood (3mL each) were spiked with 100%, 50% and 10% of the intravascular concentration of Mn / Ca (1:8) that would be expected if it were administered as a bolus at a dose of 10 μ mol/kg Mn. For comparison, Gd DTPA was spiked in 3mL of saline and blood at 100% of the expected intravascular concentration following bolus dosing at 100 μ mol/kg, a dose typically used in the clinic for vascular imaging. The tubes were imaged in cross section via T1 - weighted spin echo imaging (TR=300, TE=15,4NEX, FOV=8) on a GE Signa 1.5T Imager. The results are summarized in Table 8.

Table 8

MRI Signal Intensity in Saline and Whole Rabbit Blood				
Spike with Mn/Ca (1:8) or Gd DTPA				
	Signal Intensity in Saline	Signal Intensity in Blood	Percent Enhancement in Blood	Blood / Saline Signal Intensity
Mn, 0%	NA	363	0%	NA
Mn, 10%	262	768	112%	2.93
Mn, 50%	563	1658	357%	2.94
Mn, 100%	685	1817	401%	2.65
Gd, 100%	2162	1449	299%	0.67

It is interesting to note the significant increase in signal intensity seen for Mn / Ca (1:8) in the blood relative to saline. For instance, the signal intensity seen for Mn 10% in blood is essentially equivalent to that seen for Mn 100% in saline. This indicates interaction with blood components. This increase in efficacy is not seen with Gd DTPA, which demonstrates reduced signal enhancement in blood relative to saline. The signal intensity seen for Mn / Ca (1:8) at 50% of the expected blood concentration was greater than that seen for Gd DTPA at 100% of its expected

blood concentration, despite the fact that Gd DTPA was administered at 20 times the Mn/Ca dose on molar basis. This corresponds to at least a 40-fold efficacy advantage over Gd DTPA in blood. An even greater molar efficacy advantage is expected over Mn administered as a chelate, since chelated Mn has a lower relaxivity than does chelate Gd. For example MnDPDP, which does not bind to plasma proteins, has a solution R1 of $1.90 \text{ s}^{-1}\text{mM}^{-1}$ at 40°C and 20MHz. GdDTPA has an R1 of $3.84 \text{ s}^{-1}\text{mM}^{-1}$ under similar conditions.

Claims

1. A diagnostic composition comprising a source of a diagnostically effective quantity of Mn⁺⁺ ion, a source of Ca⁺⁺ ion and a pharmaceutically acceptable carrier for parenteral administration, wherein said Ca⁺⁺ ion is present in a molar ratio of from 2:1 to 40:1 with respect to said Mn⁺⁺ ion.
2. A diagnostic composition according to claim 1 wherein said source of Mn⁺⁺ is a manganese salt chosen from manganese acetate, chloride, gluconate, gluceptate, lactate and sulfate or a mixture thereof.
3. A diagnostic composition according to claim 2 wherein said source of Mn⁺⁺ is manganese gluconate or manganese gluceptate.
4. A diagnostic composition according to claim 1 wherein said source of Ca⁺⁺ is a calcium salt chosen from calcium acetate, chloride, gluconate, gluceptate, and lactate or a mixture thereof.
5. A diagnostic composition according to claim 4 wherein said source of Ca⁺⁺ is calcium gluconate or calcium gluceptate.
6. A diagnostic composition according to claim 1 wherein said molar ratio of calcium to manganese is from 4:1 to 20:1.
7. A diagnostic composition according to claim 6 wherein said molar ratio of calcium to manganese is from 8:1 to 10:1.
8. A diagnostic composition according to claim 1 additionally comprising an antioxidant.

9. A diagnostic composition according to claim 8 wherein said antioxidant is ascorbate.
10. A diagnostic composition according to claim 1 additionally comprising a stabilizer.
11. A diagnostic composition according to claim 10 wherein said stabilizer is calcium saccharate or a borate complex.
12. A unit dosage form comprising a salt of manganese containing from 5 mg to 200 mg of manganese, a salt of calcium containing from 20 mg to 3g of calcium, and a vehicle suitable for parenteral injection, wherein said calcium ion is present in the molar ratio from 2:1 to 40:1 with respect to said manganese ion.
13. A unit dosage form according to claim 12 comprising from 60 mg to 1.5 g of manganese gluconate or manganese gluceptate and from 250 mg to 32g of calcium gluconate or gluceptate.
14. A unit dosage form according to claim 13 in a water-for-injection vehicle adjusted to pH 6.0 to 8.2, additionally comprising calcium saccharate in an amount such that about 4% to about 8% of the calcium present in the dosage form is in the form of calcium saccharate.
15. A method for enhancing a magnetic resonance image of a mammalian tissue, organ or system comprising administering to a mammal a diagnostically effective amount of a source of Mn⁺⁺ ion together with from 2 to 200 molar equivalents of a source of Ca⁺⁺ ions, wherein said calcium ion is present in the molar ratio from 2:1 to 40:1 with respect to said manganese ion.


16. A method according to claim 15 wherein said source of Mn⁺⁺ is a manganese salt chosen from manganese acetate, chloride, gluconate, gluceptate, lactate and sulfate or a mixture thereof and said source of Ca⁺⁺ is a calcium salt chosen from calcium acetate, chloride, gluconate, gluceptate, and lactate or a mixture thereof.
17. A method according to claim 16 wherein said source of Mn⁺⁺ is manganese gluconate or manganese gluceptate and said source of Ca⁺⁺ is calcium gluconate or calcium gluceptate.
18. A method according to claim 15 wherein the molar ratio of calcium to manganese is from 4:1 to 20:1.
19. A method according to claim 15 wherein the molar ratio of calcium to manganese is from 8:1 to 10:1.
20. A method for enhancing a magnetic resonance image of a mammalian tissue, organ or system comprising administering to a mammal from 1 μ mol/kg body weight to 100 μ mol/kg body weight of a source of Mn⁺⁺ ion together with from 2 μ mol/kg body weight to 1400 μ mol/kg body weight of a source of Ca⁺⁺ ions, wherein said calcium ion is present in the molar ratio from 2:1 to 40:1 with respect to said manganese ion.
21. A method according to claim 20 wherein said source of manganese and said source of calcium are administered intravenously.
22. A method according to claim 21 wherein said source of manganese is administered at 2 μ mol/kg body weight to 30 μ mol/kg body weight and said source of calcium is administered at 4 μ mol/kg body weight to 400 μ mol/kg body weight.


23. A method according to claim 22 wherein said source of manganese is administered at 3 $\mu\text{mol}/\text{kg}$ body weight to 15 $\mu\text{mol}/\text{kg}$ body weight and said source of calcium is administered at 6 $\mu\text{mol}/\text{kg}$ body weight to 200 $\mu\text{mol}/\text{kg}$ body weight.
24. A method according to claim 20 wherein the tissue, organ or system is chosen from liver, kidney, pancreas, adrenal glands, heart, brain, salivary glands, gastrointestinal mucosa, uterus, the biliary system and tumor.
25. A method according to claim 20 wherein the tissue, organ or system is the circulatory system.
26. A method according to claim 20 wherein the source of Mn^{++} is administered separately within 30 minutes following the administration of the source of Ca^{++} .
27. A method according to claim 20 wherein the source of Mn^{++} and the source of Ca^{++} are administered simultaneously.
28. A method for assessing the metabolic activity of a tissue comprising administering to a mammal a diagnostically effective amount of a source of Mn^{++} ion together with from 2 to 200 molar equivalents of a source of Ca^{++} ions.
29. A method according to claim 28 wherein said tissue is chosen from cardiac, renal, hepatic, pancreatic and adrenal tissue.
30. A method according to claim 28 wherein said tissue is a tumor.
31. A method for assessing tissue vascularity and tissue perfusion comprising administering to a mammal a diagnostically effective amount of a source of Mn^{++} ion together with from 2 to 200 molar equivalents of a source of Ca^{++} ions.

32. A method according to claim 31 wherein said tissue is chosen from cardiac, renal, hepatic, pancreatic and adrenal tissue.

33. A method according to claim 31 wherein said tissue is a tumor.

32. A method for assessing the relative blood flow in the vasculature comprising administering to a mammal a diagnostically effective amount of a source of Mn⁺⁺ ion together with from 2 to 200 molar equivalents of a source of Ca⁺⁺ ions.








fig. 3

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/24993

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) :A61B 5/055

US CL :424/9.36

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/9.36; 436/173; 600/420

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS

search terms: manganese, calcium, magnetic resonance imaging

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,534,240 A (HASEGAWA et al) 09 July 1996, table 1.	1-34
A	EP 0 308 983 A2 (SALUTAR, INC.) 29 March 1989, Table F	1-34
A	SCHAEFER, S. et al. In Vivo Nuclear Magnetic Resonance Imaging of Myocardial Perfusion Using the Paramagnetic Contrast Agent Manganese Gluconate. J. Am. Coll. Cardiol. August 1989, Vol 14. No. 2. pages 472-480, especially page 473.	1-34

<input type="checkbox"/>	Further documents are listed in the continuation of Box C.	<input type="checkbox"/>	See patent family annex.
A	Special categories of cited documents:	*T*	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A	document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E	earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*a*	document member of the same patent family
O	document referring to an oral disclosure, use, exhibition or other means		
P	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

24 JANUARY 2000

Date of mailing of the international search report

07 FEB 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

GARY E. HOLLINDEN

Telephone No. (703) 308-1235

