

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
3 July 2008 (03.07.2008)

PCT

(10) International Publication Number  
**WO 2008/079600 A1**

- (51) **International Patent Classification:**  
C07D 213/53 (2006.01) C07D 401/14 (2006.01)
- (21) **International Application Number:**  
PCT/US2007/086319
- (22) **International Filing Date:**  
4 December 2007 (04.12.2007)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**  
60/870,851 20 December 2006 (20.12.2006) US
- (71) **Applicant (for all designated States except US):** **ELI LILLY AND COMPANY** [US/US]; Lilly Corporate Center, Indianapolis, Indiana 46285 (US).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** **KOBIERSKI, Michael, Edward** [US/US]; 3969 Shadow Hill Court, Greenwood, Indiana 46142 (US). **KOPACH, Michael, E.** [US/US]; 843 Breaside Lane, Greenwood, Indiana 46143 (US). **CHEN, Pingyun** [US/US]; 105 Westchester Place, Chapel Hill, North Carolina 27514 (US).
- (74) **Agents:** **DESAI, Manisha** et al; ELI LILLY AND COMPANY, Patent Division, P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US).


- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW
- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declarations under Rule 4.17:**

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(U))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(Hi))

[Continued on next page]

(54) **Title:** NOVEL INTERMEDIATE AND PROCESS USEFUL IN THE PREPARATION OF {2-[1-(3,5-BIS-TRIFLUOROMETHYL-BENZYL)-5-PYRIDIN-4-YL-IH-[1,2,3]TRIAZOL-4-YL]-PYRIDIN-3-YL}-(2-CHLOROPHENYL)-METHANONE



(I)

(57) **Abstract:** The present invention relates to novel compounds, (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone benzoate, and (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone toluate, which are useful intermediates for the preparation of the compound of Formula I: (I) The present invention further relates to novel processes for preparing a crystalline compound, which is {2-[1-(3,5-bis-trifluoromethyl-benzyl)-5-pyridin-4-yl-IH-[1,2,3]triazol-4-yl]-pyridin-3-yl}-(2-chlorophenyl)-methanone, Form IV

WO 2008/079600 A1



**Published:**

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*


-1-

**NOVEL INTERMEDIATE AND PROCESS USEFUL IN THE PREPARATION OF {2-[1-(3,5-BIS-TRIFLUOROMETHYL-BENZYL)-5-PYRIDIN-4-YL-1H-[1,2,3]TRIAZOL-4-YL]-PYRIDIN-3-YL}-(2-CHLOROPHENYL)-METHANONE**

5 The present invention relates to novel intermediates and a novel process for the preparation of antagonists of the NK1 subtype of tachykinin receptor. Specifically, the present invention provides novel intermediates and a novel process for the preparation of {2-[1-(3,5-bis-trifluoromethyl-benzyl)-5-pyridin-4-yl-1H-[1,2,3]triazol-4-yl]-pyridin-3-yl}-(2-chlorophenyl)-methanone, Form IV.

10 **BACKGROUND OF THE INVENTION**

The compound {2-[1-(3,5-bis-trifluoromethyl-benzyl)-5-pyridin-4-yl-1H-[1,2,3]triazol-4-yl]-pyridin-3-yl}-(2-chlorophenyl)-methanone, depicted below as the compound of Formula I, was first described in PCT published application WO2003/091226.



15

(I)

Because the compound of Formula I is an antagonist of the NK-I subtype of tachykinin receptor, it is useful for the treatment of disorders associated with an excess of tachykinins. Such disorders include depression, including major depressive disorder; anxiety, including generalized anxiety disorder, panic disorder, obsessive compulsive disorder, and social phobia or social anxiety disorder; schizophrenia and other psychotic disorders, including bipolar disorder; neurodegenerative disorders such as dementia, including senile dementia of the Alzheimer's type or Alzheimer's disease; disorders of bladder function such as bladder detrusor hyper-reflexia and incontinence, including urge incontinence; emesis, including chemotherapy-induced nausea and acute or delayed emesis; pain or nociception; disorders associated with blood pressure, such as hypertension; disorders of blood flow caused by vasodilation and vasospastic diseases, such as angina, migraine, and Reynaud's disease; hot flushes; acute and chronic obstructive airway diseases such as adult respiratory distress syndrome, bronchopneumonia, bronchospasm, chronic bronchitis, drivercough, and asthma; inflammatory diseases such as inflammatory bowel disease; gastrointestinal disorders or diseases associated with the neuronal control of viscera such as ulcerative colitis, Crohn's disease, functional dyspepsia, and irritable bowel syndrome (including constipation-predominant, diarrhea-

20

25

-?-

predominant, and mixed irritable bowel syndrome); and cutaneous diseases such as contact dermatitis, atopic dermatitis, urticaria, and other eczematoid dermatitis.

In PCT published application, WO2005/0425 15, novel crystalline forms of the compound of Formula I, identified as Form IV and Form V, are identified. Also described in WO2005/0425 15 is a process for preparation of the compound of Formula I, comprising reacting (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone or a phosphate salt thereof with 1-azidomethyl-3,5-bistrifluoromethylbenzene in the presence of a suitable base and a solvent. Use of this procedure results in several shortcomings for synthesis on a commercial scale. For example, use of the solvent DMSO, with (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone phosphate, requires a complex work-up that has a propensity to emulsify. This process also requires extraction with  $\text{CH}_2\text{Cl}_2$ , the use of which is discouraged due to its potential as an occupational carcinogen, as well as the use of  $\text{MgSCl}_4$  and acid-washed carbon, which can generate large volumes of waste on a commercial scale. Conducting the reaction with (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone in isopropyl alcohol, as also described in WO2005/042515, is also undesirable due to the need to incorporate a free base step. Furthermore, variable levels of residual 1-azidomethyl-3,5-bistrifluoromethylbenzene, a known mutagen, are obtained from use of the procedures described in WO2005/0425 15.

An improved process for preparing the compound of Formula I would control the level of 1-azidomethyl-3,5-bistrifluoromethylbenzene impurity, and improve the yield. We have discovered that use of the novel salt, (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone benzoate, as well as use of tert-butanol as the reaction solvent, improves reaction times and final yield, and decreases impurities in the final product. In addition, a novel process for the preparation of (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone benzoate, in which a pre-formed enolate of 4-acetyl pyridine is added to (2-phenylsulfonyl-pyridine-3-yl)-(2-chlorophenyl)methanone, results in an overall improved yield and improved purity, and is useful on a commercial scale.

25

#### SUMMARY OF THE INVENTION

The present invention relates to novel compounds, (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone benzoate, and (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone toluate, which are useful intermediates for the preparation of the compound of Formula I.

The present invention further relates to a process for preparing a crystalline compound, which is {2-[1-(3,5-bis-trifluoromethyl-benzyl)-5-pyridin-4-yl-1H-1,2,3-triazol-4-yl]-pyridin-3-yl}-(2-chlorophenyl)-methanone, Form IV, comprising: crystallizing the product from a solvent mixture of isopropyl acetate and heptanes.

The present invention further relates to a process for preparing a compound of Formula I comprising reacting a salt of (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone with 1-azidomethyl-3,5-bistrifluoromethylbenzene in the presence of tert-butanol.

35

**DETAILED DESCRIPTION OF THE INVENTION**

The terms and abbreviations used in the preparations and examples have their normal meanings unless otherwise designated. Other abbreviations include the following: "h" refers to hour or hours; "HPLC" refers to high performance liquid chromatography; "HRMS" refers to high resolution mass spectrometry; "*i*-PrOAc" refers to isopropyl acetate; "KOtBu" refers to potassium tert-butoxide; "min" refers to minute or minutes; "MS" refers to mass spectrometry; "NMR" refers to nuclear magnetic resonance spectroscopy; "ppm" refers to parts per million; "RT" refers to room temperature; "TLC" refers to thin layer chromatography; "DMF" refers to N,N-dimethylformamide; "DMSO" refers to dimethylsulfoxide; "MeOH" refers to methanol; "THF" refers to tetrahydrofuran.


As used herein, the term "heptanes" refers to a solution of monovalent, saturated aliphatic chains of 7 carbon atoms. The solution may contain straight chains (n-heptane), or a combination of straight and branched heptanes.

One of ordinary skill in the art will recognize that an alternate name for the compound of Formula I is: Methanone, [2-[1-[[3,5-bis(trifluoromethyl)phenyl]methyl]-5-(4-pyridinyl)-1H-1,2,3-triazol-4-yl]-3-pyridinyl](2-chlorophenyl)-.

The following examples further illustrate the improved process for preparing the compound, {2-[1-(3,5-bistrifluoromethylbenzyl)-5-pyridin-4-yl-1H-[1,2,3]triazol-4-yl]-pyridin-3-yl}-(2-chlorophenyl)-methanone, Form IV.

**EXAMPLES**Example 1

{2-[1-(3,5-bistrifluoromethylbenzyl)-5-pyridin-4-yl-1H-[1,2,3]triazol-4-yl]-pyridin-3-yl}-(2-chlorophenyl)-methanone (Form IV)



Suspend (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl] methanone benzoate (204.7 g; 1.04 equiv; 445 mmoles) in t-butanol (614 mL) and treat the slurry with potassium carbonate (124.2 g; 898.6 mmoles). Heat to 70°C with mechanical stirring for 1 hour. Add 1-azidomethyl-3,5-

-4-

bistrifluoromethylbenzene (115.6 g; 1.00 equiv; 429.4 mmol) in a single portion, then heat the mixture to reflux. A circulating bath is used to maintain a condenser temperature of 30°C. After 18 hours at reflux, HPLC reveals that the reaction is complete (<2% 1-azidomethyl-3,5-bistrifluoromethylbenzene remaining). The mixture is cooled to 70°C, isopropanol (818 mL) is added, then the mixture is stirred at 70°C for 1 hour. The mixture is filtered, and the waste filter cake is rinsed with isopropanol (409 mL). The combined filtrate and washes are transferred to a reactor, and the mechanically stirred contents are heated to 70°C. To the dark purple solution, water (1.84 L) is added slowly over 35 minutes. The solution is cooled to 60°C, then stirred for 1 hour, during which time a thin precipitate forms. The mixture is slowly cooled to RT, then the solid is filtered, washed with 1:1 isopropanol/water (614 mL), subsequently washed with isopropanol (410 mL), then dried *in vacuo* at 45°C to produce 200.3 g of crude {2-[1-(3,5-bistrifluoromethylbenzyl)-5-pyridin-4-yl-1H-[1,2,3]triazol-4-yl]-pyridin-3-yl}-(2-chlorophenyl)-methanone as a white solid. Crude {2-[1-(3,5-bistrifluoromethylbenzyl)-5-pyridin-4-yl-1H-[1,2,3]triazol-4-yl]-pyridin-3-yl}-(2-chlorophenyl)-methanone (200.3 g) and isopropyl acetate (600 mL) are charged to a 5L 3-neck jacketed flask, then the contents heated to 75°C. After dissolution is achieved, the vessel contents are cooled to 55°C, then the solution polished through a 5 micron filter, and the filter rinsed with a volume of isopropyl acetate (200 mL). After the polish filtration operation is complete, the filtrates are combined, and the vessel contents are adjusted to 50°C. After stirring for at least 15 minutes at 50°C, 0.21 grams of {2-[1-(3,5-bistrifluoromethylbenzyl)-5-pyridin-4-yl-1H-[1,2,3]triazol-4-yl]-pyridin-3-yl}-(2-chlorophenyl)-methanone Form IV seed (d90 = 40 microns) is added, and the mixture stirred at 50°C for at least 2 h. Heptanes (1.90 L) are then added over at least 2 h. After the heptanes addition is completed, the slurry is stirred for an hour at 50°C, cooled to 23°C at a rate less than 2°C per hour, then aged at 23°C for an hour prior to isolation. The mixture is then filtered in portions through the bottom outlet valve in the reactor into a 600 mL filter. The resulting wetcake is washed portionwise with a solution containing heptanes (420 mL) and isopropyl acetate (180 mL), which is passed directly through the 5L crystallization vessel. The wetcake is blown dry for 5 minutes with nitrogen, then transferred to a 500 mL plastic bottle. The product is dried at 50°C for 4 h. to produce 190.3g of pure {2-[1-(3,5-bistrifluoromethylbenzyl)-5-pyridin-4-yl-1H-[1,2,3]triazol-4-yl]-pyridin-3-yl}-(2-chlorophenyl)-methanone, Form IV in 75.0% yield with 100% purity, as determined by HPLC analysis. Particle size is reduced via pin or jet mill. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): 5.46 (s, 2H); 7.19 (m, 5H); 7.36 (dd, 1H, *J* = 4.9, 7.8); 7.45 (s, 2H); 7.59 (m, 1H); 7.83 (s, 1H); 7.93 (dd, 1H, *J* = 1.5, 7.8); 8.56 (dd, 1H, *J* = 1.5, 4.9); 8.70 (d, 2H, *J* = 5.9).

#### Preparation 1-A

(2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone benzoate

Charge powdered KOfBu (221.1 g, 1.93 moles, 1.40 eq.) to Reactor A, then charge DMSO (2 L) at 25°C over 10 min. The KOfBu/DMSO solution is stirred for 30 min at 23°C, then a solution of 4-acetyl pyridine (92 mL, 2.07 moles, 1.50 eq) in DMSO (250 mL) is prepared in reactor B. The contents of reactor

-5-


B are added to Reactor A over 10 minutes, then the Reactor A enolate solution is stirred at 23°C for 1h. In a separate 12-L flask (Reactor C), solid LiOH (84.26 g, 3.45 moles, 2.0 eq) is poured into a mixture of (2-phenylsulfonyl-pyridin-3-yl)-(2-chlorophenyl)methanone (500.0 g, 1.34 moles, 1.0 eq) and DMSO (2L), with stirring, at 23°C. The enolate solution in reactor A is then added to Reactor C over a period of at least 5 15 minutes, and the red suspension warmed to 40°C. The reaction is stirred for 3h, after which time HPLC analysis reveals less than 2% (2-phenylsulfonyl-pyridin-3-yl)-(2-chlorophenyl)methanone. Toluene (2.5 L) is charged, and the reactor temperature cooled to 30°C. The mixture is quenched by addition of glacial acetic acid (316 mL, 5.52 moles, 4.0 eq), followed by 10 % NaCl (2.5 L). The biphasic mixture is transferred to a 22-L bottom-outlet Morton flask, and the aqueous layer is removed. The aqueous layer is then extracted with toluene (750 mL). The combined organic layers are washed with 10 % NaCl (750 mL), then concentrated to 4 volumes and transferred to a 12-L Morton flask and rinsed with isopropyl acetate (4 vol, 2 L). The opaque amber solution is warmed to 75 degrees to 75°C over 40 min. Benzoic acid (171 .Ig, 1.34 moles, 1.0 eq) is dissolved in hot isopropyl acetate (1.5 L), and charged to the crude free base solution over at least 30 min. The crude solution containing benzoate salt is stirred for 0.5 h at 75 °C then cooled to 15 23 °C. When solids are first observed, the cooling is stopped and the mixture is aged for an hour at the temperature at which crystals are first observed. Alternatively, if seed crystal is available, the mixture may be seeded with (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone benzoate (2.25g) at 75°C, followed by stirring for 0.5 h at 75°C, then cooling to 23°C over at least 1.5 h. The mixture is then cooled to <5 °C, then filtered through paper on a 24cm single-plate filter. The filtercake is then 20 rinsed with cold *i*-PrOAc (750 mL) to produce granular crystals of bright orange-red color. The wet solid is dried at 55°C to produce 527.3 g (83% yield) with 99.9% purity. (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone benzoate. Anal. Calcd. for C<sub>26</sub>H<sub>19</sub>N<sub>2</sub>ClO<sub>4</sub>: C, 68.05; H, 4.17; N, 7.13. Found: C, 67.89; H, 4.15; N 6.05. HRMS: calcd for C<sub>19</sub>H<sub>13</sub>ClN<sub>2</sub>O<sub>2</sub>, 336.0666; found 336.0673.

The synthesis of (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone 25 benzoate proceeds optimally when the potassium enolate of 4-acetyl pyridine is pre-formed using KOfBu in DMSO. Pre-formation of the enolate allows the SNAR (nucleophilic aromatic substitution) reaction to be performed between room temperature and 40°C, which minimizes the amount of degradation. Under these conditions, the SNAR is highly regioselective, resulting in a ratio of approximately 95:5 preferential C - acylation. In all cases, less polar solvents such as THF or toluene, or co-solvents of these solvents mixed 30 with DMSO, results in a substantial increase of acylation at the oxygen in the SNAR, and leads to a lower yield of product. This is a substantial improvement over the procedures described in WO2005/0425 15 for synthesis of the free base or the phosphate salt, in which the SNAR is performed at 60-70 °C, resulting in a substantial increase in chemical impurity. Using the conditions described in WO2005/042515, when scaled to 2kg, results in maximum yields of 55%, with sub-optimal potency. In comparison, the improved 35 conditions described herein can be run reproducibly from 0.4 to 2kg scale to give yields of 77-83%, with >99% purity. In addition, the reaction can be held overnight at 40°C with minimal degradation, whereas holding the reaction for 1 h past completion at 60-70°C results in substantial aromatized impurity. The

-6-

reaction may also be performed using sodium tert-amylate as the base, in combination with an aprotic solvent, such as DMSO or DMF.

The title compound exists as a mixture of tautomers and geometric isomers. It is understood that each of these forms is encompassed within the scope of the invention.



#### Preparation 1-B

(2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone toluate

The procedure described in Preparation 1-A is followed, with the following exception. Solid toluic acid (1.0 eq) is added to the crude free base solution at 55°C, then the solution cooled to 45 °C. The solution is stirred for one hour at 45 °C, then slowly cooled to 23 °C. When solids are first observed, the cooling is stopped and the mixture is aged for an hour at the temperature at which crystals are first observed. Alternatively, if seed crystal is available, the mixture may be seeded, aged for 3 h at 45°C, then cooled to 0°C over 4 h. The isolation slurry is filtered, and the wetcake washed with MeOH (3 volumes). The wetcake is dried at 50°C to provide 14.0 g (76.4%) of (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone toluate as a light red powder.

10

15

As with the benzoate salt, the toluate salt can also exist as a mixture of tautomers and geometric isomers, each of which is encompassed within the scope of the invention. (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone toluate. <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>) δ 194.5, 167.8, 167.4, 155.5, 150.7 (2C), 147.4, 144.0, 143.4, 142.7, 138.6, 133.0, 130.8, 130.7, 130.5, 129.8(2C), 129.5(2C), 128.5, 128.0, 127.9, 119.9 (2C), 118.6, 92.6, 21.5.

20

#### Preparation 1-C

(2-phenylsulfonyl-pyridin-3-yl)-(2-chlorophenyl)methanone

A solution of 1.3 eq of diisopropylamine (based on 2-benzenesulfonyl pyridine) in 5 volumes of THF in a mechanically stirred 3-necked flask is cooled to -70 to -75 °C. To this solution is added 1.05 eq of w-butyllithium (1.6M in hexanes) at such a rate as to maintain the temperature below -60°C. The light yellow solution is stirred at -60 to -70 °C for 30 minutes. Once the temperature has cooled back down to -60 to -65°C, 1.0 eq of 2-benzene-sulfonyl pyridine, as a solution in 3 volumes of THF, is added at the fastest rate that will maintain the reaction temperature under -60°C. A yellow suspension forms during the addition that becomes yellow-orange upon longer stirring. This mixture is stirred for 3 hours at -60 to -75°C, and then 1.06 eq of 2-chlorobenzaldehyde, as a solution in 1 volume of THF, is added dropwise at a

25

30

-7-

sufficient rate to keep the temperature under  $-55^{\circ}\text{C}$ . The suspension gradually turns orange-red, thins out, and then becomes a clear red solution. The reaction mixture is allowed to stir at  $-60$  to  $-70^{\circ}\text{C}$  for 1 hour, 3N aqueous HCl (7 volumes) is added over 20-30 minutes, and the temperature is allowed to exotherm to  $0$ - $10^{\circ}\text{C}$ . The color largely disappears, leaving a biphasic yellow solution. The solution is

5 warmed to at least  $10^{\circ}\text{C}$ , the layers are separated, and the aqueous layer is back-extracted with 10 volumes of ethyl acetate. The combined organic layers are washed with 10 volumes of saturated sodium bicarbonate solution and concentrated to about 2 volumes. Ethyl acetate (10 volumes) is added, and the solution is once again concentrated to 2 volumes. The thick solution is allowed to stand overnight and is taken to the next step with no purification of the crude alcohol intermediate. The crude alcohol

10 intermediate is transferred to a 3-necked flask with enough ethyl acetate to make the total solution about 10 volumes. The yellow solution is treated with 3.2 volumes of 10% aqueous (w/w) potassium bromide, followed by 0.07 eq of 2,2,6,6-Tetramethylpiperidine-N-oxide (TEMPO). The orange mixture is cooled to  $0$ - $5^{\circ}\text{C}$  and treated with a solution of 1.25 eq of sodium bicarbonate in 12% w/w sodium hypochlorite (9 volumes) and 5 volumes of water over 30-60 minutes while allowing the temperature to exotherm to a

15 maximum of  $20^{\circ}\text{C}$ . The mixture turns dark brown during the addition, but becomes yellow, and a thick precipitate forms. The biphasic light yellow mixture is allowed to stir at ambient temperature for 1-3 hours, at which time the reaction is generally completed. The biphasic mixture is cooled to  $0$ - $5^{\circ}\text{C}$  and stirred for 3 hours at that temperature. The solid is filtered off, washed with 4 volumes of cold ethyl acetate, followed by 4 volumes of water, and dried *in vacuo* at  $45^{\circ}\text{C}$  to constant weight. Typical yield is 80-83% with a

20 purity of greater than 98%.  $^1\text{H NMR}$  (600 MHz,  $\text{CDCl}_3$ -*d*)  $\delta$  ppm 7.38 (td,  $J=7.52$ , 1.28 Hz, 1 H) 7.47 (dd,  $J=7.80$ , 1.30 Hz, 1 H) 7.51 (td,  $J=7.79$ , 1.60 Hz, 1 H) 7.51 (t,  $J=7.89$  Hz, 2 H) 7.50 - 7.54 (m,  $J=7.75$ , 4.63 Hz, 1 H) 7.60 (t,  $J=7.43$  Hz, 1 H) 7.73 (dd,  $J=7.75$ , 1.60 Hz, 1 H) 7.81 (dd,  $J=7.79$ , 1.56 Hz, 1 H) 8.00 (dd,  $J=8.44$ , 1.10 Hz, 2 H) 8.76 (dd,  $J=4.63$ , 1.61 Hz, 1 H).

#### Preparation 1-D

25 1-azidomethyl-3,5-bis(trifluoromethyl)benzene

Sodium azide (74.3 g, 1.14 mol) is suspended in water (125 mL), then DMSO (625 mL) is added. After stirring for 30 minutes, a solution consisting of 3,5-Bis(trifluoromethyl)benzyl chloride (255.3 g, 0.97 moles) and DMSO (500 mL) is added over 30 minutes. (The 3,5-Bis(trifluoromethyl)benzyl chloride is heated to  $35^{\circ}\text{C}$  to liquefy prior to dispensing (MP =  $30$ - $32^{\circ}\text{C}$ )). The benzyl chloride feed vessel is rinsed

30 with DMSO (50 mL) into the sodium azide solution, the mixture is heated to  $40^{\circ}\text{C}$ , and then maintained for an hour at  $40^{\circ}\text{C}$ , then cooled to  $23^{\circ}\text{C}$ .

In Process Analysis: A drop of the reaction mixture is dissolved in  $d_6$ -DMSO and the relative intensities of the methylene signals are integrated (NMR verified as a 0.35% limit test for 3,5-Bis(trifluoromethyl)benzyl Chloride).

35 Work-up: After the mixture reaches  $23^{\circ}\text{C}$ , it is diluted with heptanes (1500 mL), then water (1000 mL) is added, and the mixture exotherms to  $35^{\circ}\text{C}$  against a jacket setpoint of  $23^{\circ}\text{C}$ .

-8-

The aqueous layer is removed (-2200 mL), then the organic layer (approximately 1700 mL) is washed with water (2 X 750 mL). The combined aqueous layers (-3700 mL) are analyzed and discarded.

The solvent is then partially removed via vacuum distillation with a jacket set point of 85°C, pot temperature of 60-65°C and distillate head temperature of 50-55°C to produce 485g (94.5% yield) of  
5 51Wt% solution title compound as a clear liquid. Heptanes can be either further removed by vacuum distillation or wiped film evaporation technology. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): 4.58 (s, 2H); 7.81 (s, 2H); 7.90 (s, 1H).

#### Preparation 1-E

##### 2-benzene-sulfonyl pyridine

10 Charge 2-chloropyridine (75 mL, 790 mmol), thiophenol (90 mL, 852 mmol), and DMF (450 mL) to a 2L flask. Add K<sub>2</sub>CO<sub>3</sub> (134.6 g, 962 mmol), then heat to 100°C and stir for 18 hours. Filter the mixture, then rinse the waste cake with DMF (195 mL). The combined crude sulfide solution and rinses are transferred to a 5-L flask, and the waste filtercake is discarded. Glacial acetic acid (57 mL, 995 mmol) is added to the filtrate, then the solution is heated to 40°C, and 13 wt % NaOCl solution (850 mL, 1.7 mol) is  
15 added over 2 hours. After the reaction is complete, water (150 mL) is added, then the pH of the mixture adjusted to 9 with 20 % (w/v) NaOH solution (250 mL). The resulting slurry is cooled to <5 °C, stirred for 1.5 h, then filtered, and the cake washed with water (3 x 200 mL). The product wetcake is dried in a 55°C vacuum oven to provide 2-benzene-sulfonyl pyridine (149 g, 676 mmol) in 86 % yield: <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 8.66 (d, *J* = 5.5 Hz, 1H), 8.19 (d, *J* = 1.1 Hz, 1H), 8.05 (m, 2H), 7.92 (ddd, *J* = 9.3, 7.7, 1.6  
20 Hz, 1H), 7.60 (m, 1H), 7.54 (m, 2H), 7.44 (m, 1H); IR (KBr) 788, 984, 1124, 1166, 1306, 1424, 1446, 1575, 3085 cm<sup>-1</sup>; MS (TOF) *m/z* 220.0439 (220.0427 calcd for C<sub>11</sub>H<sub>10</sub>NO<sub>2</sub>S, MH); Anal. calcd for C<sub>11</sub>H<sub>9</sub>NO<sub>2</sub>S: C, 60.26; H, 4.14; N, 6.39; S, 14.62. Found: C, 60.40; H, 4.02; N, 6.40; S, 14.76.

As noted above, use of the improved process of the present invention results in an improved habit  
25 of the crystalline Form IV compound of Formula I. The improved habit reduces surface area of the crystal, improves the filtration, and washing, and improves the efficiency of azide mutagen rejection. These improvements are described in greater detail below.

In patent application WO2005/0425 15, the polish filtration is carried out in 7 volumes (L/kg) of isopropanol near its boiling point (65-83 °C), a process that is difficult and hazardous to execute in  
30 commercial manufacturing because of the high risk of crystallization on the filter and/or vessel transfer lines due to supersaturation. In the preferred crystallization solvent, isopropyl acetate, the polish filtration is conducted in four volumes of isopropyl acetate at temperatures from 45 to 55 °C. This temperature range is 35 to 45 °C lower than the boiling point of isopropyl acetate, which provides a key safety advantage. Under these conditions, {2-[1-(3,5-bistrifluoromethylbenzyl)-5-pyridin-4-yl-1H-[1,2,3]triazol-4-yl]-pyridin-  
35 3-yl}-(2-chlorophenyl)-methanone will remain in solution for days, and there is no danger of crystallization and clogging of filter and transfer lines. Controlled crystallization is conducted by seeding with Form IV

crystals at 40-45°C, followed by a ripening period of at least 2 hrs, then addition of 12-15 volumes of an anti-solvent, preferably heptanes, which maximizes process yields.

Larger API crystals with the preferred habit (e.g., prisms and rods, as opposed to needles) provide favorable processing characteristics, including fast filtration, efficient washing and good powder flow properties. Data from four sample filtrations, conducted in the FLT-37 Hastelloy® filter-drier on a 2-3 kg scale, are provided below. The data show an average increase in filtration rate of three- to four-fold (Flux data) for the preferred isopropyl acetate/heptanes system relative to the prior isopropyl alcohol solvent system.

#### 10 Isopropyl acetate/heptanes filtration

Lot Number	A	B	C	D
Flux (L/m <sup>2</sup> ·h.)	8,858	11,106	12,930	10,807
Filtration Time (min)	11	7	8	9
Wetcake thickness (mm)	130	97	125	100

#### Isopropyl alcohol filtration

Lot Number	A	B	C	D
Flux (L/m <sup>2</sup> ·h.)	3,424	2,133	2,712	2,002
Filtration Time (min)	24	43	31	32
Wetcake thickness (mm)	145	140	140	138

A unimodal particle size distribution is optimally achieved with seeding where seed particle size consists of a  $d_{90} < 20$  microns, seed load = 0.1wt%, seed temperature = 50°C, seed age time = 2 h, and heptanes feed rate = 2 hr. In another preferred embodiment, the seed particle size consists of a  $d_{90} = 40$  microns, seeding temperature = 55°C, seed load = 0.1-2%.

In the preferred embodiment, the final product is milled to a target particle size for optimal use in the drug product. The preferred methods of milling include, but are not limited to, pin mill, turbo rotor, jet mill and slurry mill. Jet mill technology produces final product with  $d_{90}$  of approximately 10 microns.

The preferred crystallization system using isopropyl acetate/heptanes is found to be optimal to remove methanone positional isomer impurities and azide. Spiking studies have revealed that these impurities are completely purged from the final product under the preferred processing conditions described herein.

The free base of (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone does not crystallize directly from the crude reaction mixture, and conversion to a salt form for purification is essential. The benzoate and toluate salts of (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone, described herein, are readily prepared from the crude free base, as described above. Both the benzoate and toluate salts have an improved morphology relative to the phosphate salt. Both the benzoate and toluate salts exist in a large plated morphology as compared to the needle-like morphology of the phosphate salt. This difference in morphology results in a substantially improved isolation rate for the benzoate and toluate salts. The improved morphology translates to a higher bulk density of isolated solid

-10-

(0.4 -0.5 g/L) for the benzoate or toluate salts, as compared to the phosphate salt (0.1 -0.2 g/L). This results in increased drier capacity. In addition, both the benzoate and toluate salts are free flowing solids, whereas the phosphate salt appears to contain considerable static. Generally, the benzoate salt is the most preferred embodiment

**WHAT IS CLAIMED IS:**

1. A process for preparing a compound that is {2-[1-(3,5-bis(trifluoromethyl)benzyl)-5-pyridin-4-yl-1H-[1,2,3]triazol-4-yl]-pyridin-3-yl}-(2-chlorophenyl)-methanone, Form IV, comprising:  
5 crystallizing the product from a solvent mixture of isopropyl acetate and heptanes.
2. The process of **Claim 1** further comprising reacting a salt of (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone with 1-azidomethyl-3,5-bis(trifluoromethyl)benzene in  
10 the presence of tert-butanol.
3. The process of **Claim 2** wherein the salt of (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone is a benzoate salt or a toluate salt.
4. The process of **Claim 3** wherein the salt of (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone is a benzoate salt.  
15
5. A compound selected from the group consisting of (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone benzoate and (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone toluate.  
20
6. The compound of **Claim 5** which is (2-chlorophenyl)-[2-(2-hydroxy-2-pyridin-4-yl-vinyl)pyridin-3-yl]methanone benzoate.

**INTERNATIONAL SEARCH REPORT**

International application No  
PCT/US2007/086319

**A. CLASSIFICATION OF SUBJECT MATTER**  
INV. C07D213/53 C07D401/14

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
**C07D**

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal , WPI Data, BEILSTEIN Data, CHEM ABS Data

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
<b>X</b>	WO 2005/042515 A (LILLY CO ELI [US]; BORGHESE ALFIO [BE]; COFFEY DAVID SCOTT [US]; FOOTM) 12 May 2005 (2005-05-12) cited in the application claim 1; examples 1-3 -----	<b>1-6</b>

**D** Further documents are listed in the continuation of Box C.

See patent family annex.

\* Special categories of cited documents :

- 'A' document defining the general state of the art which is not considered to be of particular relevance
- 'E' earlier document but published on or after the international filing date
- 'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- 'O' document referring to an oral disclosure, use, exhibition or other means
- 'P' document published prior to the international filing date but later than the priority date claimed

- 'T' later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- 'X' document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- 'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- '&' document member of the same patent family

Date of the actual completion of the international search

**25 April 2008**

Date of mailing of the international search report

08/05/2008

Name and mailing address of the ISA/  
 European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

**Seelmann, Ingo**

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2007/086319

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2005042515 A	12-05-2005	AU 2004285855 A1	12-05-2005
		BR PI0415010 A	07-11-2006
		CA 2542140 A1	12-05-2005
		CN 1863791 A	15-11-2006
		EP 1675846 A1	05-07-2006
		JP 2007509143 T	12-04-2007
		KR 20060061388 A	07-06-2006

---