



(51) International Patent Classification:

C07D 213/34 (2006.01) *C07D 407/12* (2006.01)
C07D 213/62 (2006.01) *C07D 417/12* (2006.01)
C07D 233/64 (2006.01) *A61K 31/343* (2006.01)
C07D 307/18 (2006.01) *A61P 31/00* (2006.01)
C07D 307/82 (2006.01) *C07C 311/00* (2006.01)
C07D 333/34 (2006.01)

(21) International Application Number:

PCT/US2011/064233

(22) International Filing Date:

9 December 2011 (09.12.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/423,940 16 December 2010 (16.12.2010) US

(71) Applicant (*for all designated States except US*): **ALLERGAN, INC.** [US/US]; 2525 Dupont Drive, Irvine, California 92886 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **YUAN, Haiqing** [US/US]; 8 San Garin, Irvine, California 92606 (US). **BEARD, Richard L.** [US/US]; 2341 Azure Avenue, Newport Beach, California 92660 (US). **LIU, Xiaoxia** [US/US]; 22212 Newbridge Drive, Lake Forest, California 92630 (US). **DONELLO, John E.** [US/US]; 34041 Pequito Drive, Dana Point, California 92629 (US). **VISWANATH, Veena** [US/US]; 3 Pollena, Irvine, California 92602 (US). **GARST, Michael E.** [US/US]; 2627 Raqueta Drive, Newport Beach, California 92660 (US).

(74) Agents: **ENE, Doina** et al.; Allergan, Inc., 2525 Dupont Drive, Irvine, California 92612 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

- *with international search report (Art. 21(3))*

(54) Title: SULFUR DERIVATIVES AS CHEMOKINE RECEPTOR MODULATORS

(57) Abstract: The present invention relates to novel sulfur derivatives, processes for preparing them, pharmaceutical compositions containing them and their use as pharmaceuticals as modulators of chemokine receptors.



**By inventors: Haiqing Yuan, Richard L. Beard, Xiaoxia Liu, John E. Donello,
Veena Viswanath, and Michael E. Garst**

This application claims the benefit of U.S. Provisional Application Serial No. 61/423,940, filed December 16, 2011, the disclosure of which is hereby incorporated in its entirety herein by reference.

The present invention relates to novel sulfur derivatives, processes for preparing them, pharmaceutical compositions containing them and their use as pharmaceuticals as modulators of chemokine receptors. The invention relates specifically to the use of these compounds and their pharmaceutical compositions to treat disorders associated with chemokine receptor (CCR) modulation.

Chemokines are a group of 7- to 14-kd peptides that play an important role in orchestrating leukocyte recruitment and migration during inflammation, and therefore represent an important target for anti-inflammatory therapies (Wells et al., 2006). They act by binding to seven-transmembrane, G protein-coupled receptors, the chemokine receptors. The chemokine system is complex, with about ~50 chemokines and 20 chemokine receptors identified in humans, often acting with redundancy, making selection of specific antagonists difficult (Gerard and Rollins, 2001). Genetic knockout strategies have confirmed the importance of chemokines as regulators of immune function, but the deletion of specific chemokines has led to only specific and relatively mild defects in the inflammatory response further emphasizing the complex redundancy of the system. Selectivity is crucial for use of chemokine receptor antagonists in systemic diseases where a single chemokine-receptor system is implicated such as atherosclerosis where the macrophage/monocyte system is the major player in order to allow a subtle and specific control over immune function (Weisberg et al., 2006; Feraia and Diaz Gonzalez et al., 2006).

Many ocular conditions are characterized by inappropriate migration and infiltration of cells such as leukocytes and endothelial cells into the eye with deleterious effects to ocular structures (Wallace et al., 2004). Chemokines have been identified in such diseases and misregulation of the chemokine system is apparent in

5 corneal graft rejection, diabetic retinopathy, age-related macular degeneration (ARMD), chronic inflammatory diseases such as uveitis, dry eye etc. Mice lacking CCR2 or MCP-1 develop features of ARMD with age, including drusen deposits, choroidal neovascularization and photoreceptor atrophy indicating a crucial role for this chemokine and its receptor signaling (Amabati et al., 2003). Thus CCR2 receptor-

10 specific inhibitor might have potential therapeutic benefit in ocular diseases like ARMD. In contrast, various human and animal studies have identified several chemokines in different forms of uveitis, produced both by resident and infiltrating cells, that strongly suggests a prominent role for these molecules in its pathogenesis. Studies in rat and mice models of uveitis have demonstrated up-regulation of

15 monocyte chemoattractant protein-1 (MCP-1), macrophage inflammatory protein-1 (MIP-1), RANTES, stromal derived factor-1 (SDF-1) which are powerful chemoattractants for monocytes and T-cells (Fang et al., 2004; Keino et al., 2003). Similar findings have been reported in peripheral blood mononuclear cells in patients with acute anterior uveitis (AAU), the most common form of human uveitis (Klitgaard

20 et al., 2004). MCP-1 knockout mice and CCR5 knockout mice show reduced endotoxin-induced uveitis, which is the animal model for AAU (Takeuchi et al., 2005; Tuallion et al., 2002). It has also been demonstrated that blocking the chemokine system upstream with the use of NF- κ B blockers significantly attenuates experimental AAU in rats (Yang et al., 2005). Blockage of NF- κ B results in transcriptional inhibition

25 of multiple chemokines. Given the complexity of pathogenesis in uveitis it is unlikely that a selective inhibition of a chemokine receptor in monotherapy will offer therapeutic benefit. A similar role of multiple chemokines have been shown to be correlated with clinical stage of disease in diabetic retinopathy and dry eye (Meleth et al., 2005; Yamagami et al., 2005). In these ocular diseases the use of broad spectrum

30 chemokine receptor inhibitor which inhibits the function of a wide range of chemokines may be beneficial.

The first broad spectrum chemokine inhibitor (BSCI) to be reported was termed Peptide 3, which was derived from the sequence of human chemokine MCP-1 and

was shown to block the migration of monocytes in response to MCP-1, MIP-1, RANTES and SDF-1 (Reckless and Grainger. 1999). A cyclic retro inverse analogue of Peptide 3, constructed of D-amino acids in the reverse sequence, called NR58-3.14.3 was observed to be a more potent chemokine inhibitor (Beech et al., 2001).

5 NR58-3.14.3 has been used to test for anti-inflammatory activities in animal models of atherosclerosis, lung inflammation, irritable bowel syndrome etc (Beech et al., 2001; Grainger and Reckless. 2003; Tokuyama et al., 2005). However there are several disadvantages to using these BSCI as a long-term therapeutic strategy. The known BSCIs which are peptides which have relatively low potency, poor pharmacokinetics, 10 and are unstable in vivo. In addition, systemic use of broad spectrum chemokine receptor inhibitors could potentially lead to deleterious side effects due to their systemic anti-inflammatory activity. However in ocular diseases, a local or topical application would prevent the broad spectrum inhibitor to be taken up systemically. Identification of a small molecule inhibitor of several chemokine receptors could be 15 very useful for treatment of inflammatory ocular diseases. Given the evidence for the role of multiple chemokines in several ocular diseases and these results, we propose that the use of small and large molecule broad spectrum chemokine receptor inhibitors will have utility in the local treatment of ocular inflammatory diseases including, but not limited to, uveitis, dry eye, diabetic retinopathy, allergic eye disease and proliferative 20 retinopathies. Manipulation of multiple chemokines therefore represents a novel therapeutic approach in treating ocular diseases.

WO2008008374 discloses CCR2 inhibitors and methods of use thereof.

WO03/099773 discloses CCR9 inhibitors and methods of use thereof.

US7622583 discloses heteroaryl sulfonamides as antagonists of the CCR2 25 receptor.

US 2008/0293720 discloses pyridinyl sulfonamide modulators of chemokine receptors.

US7393873 discloses arylsulfonamide derivatives.


SUMMARY OF THE INVENTION

30 We have now discovered a group of novel sulphur derivatives which are potent and selective chemokine receptor modulators. As such, the compounds described herein are useful in treating a wide variety of disorders associated with modulation of

chemokine receptors. The term “modulator” as used herein, includes but is not limited to: receptor agonist, antagonist, inverse agonist, inverse antagonist, partial agonist, partial antagonist.

This invention describes compounds of Formula I, which have chemokine
 5 receptor biological activity. The compounds in accordance with the present invention are thus of use in medicine, for example in the treatment of humans with diseases and conditions that are alleviated by CCR modulation.

In one aspect, the invention provides a compound having **Formula I** or a
 pharmaceutically acceptable salt thereof or stereoisomeric forms thereof, or the
 10 individual geometrical isomers, enantiomers, diastereoisomers, tautomers, zwitterions and pharmaceutically acceptable salts thereof:



Formula I

15 wherein:

R¹ is H

R² is substituted or unsubstituted C₁₋₆ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl or is substituted or unsubstituted C₃₋₈ cycloalkenyl;

R⁵ is -S-, -S(O)-, or -S(O)₂-;

20 R⁶ is substituted or unsubstituted heterocycle, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted C₃₋₈ cycloalkenyl or is substituted or unsubstituted C₆₋₁₀ aryl;

R¹⁷ is H, substituted or unsubstituted C₁₋₆ alkyl, halogen, substituted or unsubstituted -OC₁₋₆ alkyl, CN, C(O)R¹⁹, NR²⁰R²¹ or hydroxyl;

25 R¹⁸ is H, substituted or unsubstituted C₁₋₆ alkyl, halogen, substituted or unsubstituted -OC₁₋₆ alkyl, CN, C(O)R²², NR²³R²⁴ or hydroxyl;

R^7 is H, halogen, CN, substituted or unsubstituted $-OC_{1-6}$ alkyl, substituted or unsubstituted C_{1-6} alkyl or is substituted or unsubstituted C_{3-8} cycloalkyl;

R^8 is H, substituted or unsubstituted C_{1-6} alkyl, halogen, substituted or unsubstituted $-OC_{1-6}$ alkyl, or hydroxyl;

5 R^{19} is H, OH or substituted or unsubstituted C_{1-6} alkyl

R^{20} is H or substituted or unsubstituted C_{1-6} alkyl;

R^{21} is H or substituted or unsubstituted C_{1-6} alkyl;

R^{22} is H, OH or substituted or unsubstituted C_{1-6} alkyl

R^{23} is H or substituted or unsubstituted C_{1-6} alkyl;

10 R^{24} is H or substituted or unsubstituted C_{1-6} alkyl; and including compounds:

N-{2-[(3-aminobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide;

N-{2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;

N-{2-[(3-aminobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;

N-{2-[(3-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide; and


15 N-{2-[(3-aminobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide; and


with the provisos that

a). R^7 , R^8 , R^{17} and R^{18} cannot all be H in same time;

b). when R^8 is substituted or unsubstituted C_{1-6} alkyl, the alkyl group does not contain a carbonyl group directly attached to the phenyl ring;

20 c). the compound is not of the following structures:





In another aspect, the invention provides a compound having Formula I

wherein

R¹ is H;

R² is substituted or unsubstituted C₁₋₆ alkyl, substituted or unsubstituted C₃₋₈

cycloalkyl or substituted or unsubstituted C₃₋₈ cycloalkenyl;

R⁵ is S;

R^6 is substituted or unsubstituted heterocycle, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted C_{3-8} cycloalkenyl or substituted or unsubstituted C_{6-10} aryl;

R^{17} is H, substituted or unsubstituted C_{1-6} alkyl, halogen, substituted or unsubstituted -
5 OC_{1-6} alkyl, CN, $C(O)R^{19}$, $NR^{20}R^{21}$ or hydroxyl;

R^{18} is H, C_{1-6} alkyl, halogen, substituted or unsubstituted $-OC_{1-6}$ alkyl, CN, $C(O)R^{22}$, $NR^{23}R^{24}$ or hydroxyl;

R^7 is H, halogen, CN, substituted or unsubstituted $-OC_{1-6}$ alkyl, substituted or unsubstituted C_{1-6} alkyl or substituted or unsubstituted C_{3-8} cycloalkyl;

10 R^8 is H, substituted or unsubstituted C_{1-6} alkyl, halogen, substituted or unsubstituted $-OC_{1-6}$ alkyl, or hydroxyl;

R^{19} is H, OH or substituted or unsubstituted C_{1-6} alkyl;

R^{20} is H or substituted or unsubstituted C_{1-6} alkyl;

R^{21} is H or substituted or unsubstituted C_{1-6} alkyl;

15 R^{22} is H, OH or substituted or unsubstituted C_{1-6} alkyl;

R^{23} is H or substituted or unsubstituted C_{1-6} alkyl; and


R^{24} is H or substituted or unsubstituted C_{1-6} alkyl; and


with the provisos that

a). R^7 , R^8 , R^{17} and R^{18} cannot all be H in same time;

20 b). when R^8 is substituted or unsubstituted C_{1-6} alkyl, the alkyl group does not contain a carbonyl group directly attached to the phenyl ring;

c). the compound is not of the following structures:





- 5 In another aspect, the invention provides a compound having Formula I wherein:
- R^1 is H ;
- R^2 is substituted or unsubstituted C_{1-6} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl or substituted or unsubstituted C_{3-8} cycloalkenyl;
- 10 R^5 is -S(O)- ;
- R^6 is substituted or unsubstituted heterocycle, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted C_{3-8} cycloalkenyl or substituted or unsubstituted C_{6-10} aryl;
- R^{17} is H, substituted or unsubstituted C_{1-6} alkyl, halogen, substituted or unsubstituted -
- 15 OC_{1-6} alkyl, CN, $C(O)R^{19}$, $NR^{20}R^{21}$ or hydroxyl;

R^{18} is H, substituted or unsubstituted C_{1-6} alkyl, halogen, substituted or unsubstituted -
 OC_{1-6} alkyl, CN, $C(O)R^{22}$, $NR^{23}R^{24}$ or hydroxyl;

R^7 is H, halogen, CN, substituted or unsubstituted - OC_{1-6} alkyl, substituted or
 unsubstituted C_{1-6} alkyl or substituted or unsubstituted C_{3-8} cycloalkyl;

5 R^8 is H, substituted or unsubstituted C_{1-6} alkyl, halogen, substituted or unsubstituted -
 OC_{1-6} alkyl, or hydroxyl;

R^{19} is H, OH or substituted or unsubstituted C_{1-6} alkyl;

R^{20} is H or substituted or unsubstituted C_{1-6} alkyl;

R^{21} is H or substituted or unsubstituted C_{1-6} alkyl;

10 R^{22} is H, OH or substituted or unsubstituted C_{1-6} alkyl;

R^{23} is H or substituted or unsubstituted C_{1-6} alkyl; and


R^{24} is H or substituted or unsubstituted C_{1-6} alkyl; and

with the provisos that

a). R^7 , R^8 , R^{17} and R^{18} cannot all be H in same time;

15 b). when R^8 is substituted or unsubstituted C_{1-6} alkyl, the alkyl group does not
 contain a carbonyl group directly attached to the phenyl ring;

c). the compound is not of the following structure:



20 In another aspect, the invention provides a compound having Formula I
 wherein:

R^1 is H ;

R^2 is substituted or unsubstituted C_{1-6} alkyl, substituted or unsubstituted C_{3-8}
 cycloalkyl or substituted or unsubstituted C_{3-8} cycloalkenyl;

25 R^5 is $-S(O)_2-$;

R^6 is heterocycle, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or
 unsubstituted C_{3-8} cycloalkenyl or substituted or unsubstituted C_{6-10} aryl;

R^{17} is H, substituted or unsubstituted C_{1-6} alkyl, halogen, - OC_{1-6} alkyl, CN, $C(O)R^{19}$,
 $NR^{20}R^{21}$ or hydroxyl;

R^{18} is H, substituted or unsubstituted C_{1-6} alkyl, halogen, substituted or unsubstituted $-OC_{1-6}$ alkyl, CN, $C(O)R^{22}$, $NR^{23}R^{24}$ or hydroxyl;

R^7 is H, halogen, CN, substituted or unsubstituted $-OC_{1-6}$ alkyl, substituted or unsubstituted C_{1-6} alkyl or substituted or unsubstituted C_{3-8} cycloalkyl;

5 R^8 is H, substituted or unsubstituted C_{1-6} alkyl, halogen, $-OC_{1-6}$ alkyl, or hydroxyl;

R^{19} is H, OH or substituted or unsubstituted C_{1-6} alkyl;

R^{20} is H or substituted or unsubstituted C_{1-6} alkyl;

R^{21} is H or substituted or unsubstituted C_{1-6} alkyl;

R^{22} is H, OH or substituted or unsubstituted C_{1-6} alkyl;

10 R^{23} is H or substituted or unsubstituted C_{1-6} alkyl; and


R^{24} is H or substituted or unsubstituted C_{1-6} alkyl; and

with the provisos that

a). R^7 , R^8 , R^{17} and R^{18} cannot all be H.in same time;

b). when R^8 is substituted or unsubstituted C_{1-6} alkyl, the alkyl group does not
15 contain a carbonyl group directly attached to the phenyl ring;

c). the compound is not of the following structures:



In another aspect, the invention provides a compound having Formula I

20 wherein:

R^1 is H;

R^2 is substituted or unsubstituted C_{1-6} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl or substituted or unsubstituted C_{3-8} cycloalkenyl;

R^5 is $-S-$, $-S(O)-$, or $-S(O)_2-$;

25 R^6 is substituted or unsubstituted heterocycle or substituted or unsubstituted C_{6-10} aryl;

R^{17} is H, substituted or unsubstituted C_{1-6} alkyl or halogen;


R^{18} is H, substituted or unsubstituted C_{1-6} alkyl or halogen;

R^7 is halogen, CN, $-OC_{1-6}$ alkyl, substituted or unsubstituted C_{1-6} alkyl or substituted or unsubstituted C_{3-8} cycloalkyl;

30 R^8 is H, substituted or unsubstituted C_{1-6} alkyl or halogen; and

with the provisos that:

- b). when R^8 is substituted or unsubstituted C_{1-6} alkyl, the alkyl group does not contain a carbonyl group directly attached to the phenyl ring;
- c). the compound is not of the following structures:




In another aspect, the invention provides a compound having Formula I

wherein:

- 10 R^1 is H
- R^2 is substituted or unsubstituted C_{1-6} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl or substituted or unsubstituted C_{3-8} cycloalkenyl;
- R^5 is -S-;
- R^6 is substituted or unsubstituted heterocycle or substituted or unsubstituted C_{6-10} aryl;
- 15 R^{17} is H, substituted or unsubstituted C_{1-6} alkyl or halogen;
- R^{18} is H, substituted or unsubstituted C_{1-6} alkyl or halogen;
- R^7 is halogen, CN, substituted or unsubstituted $-OC_{1-6}$ alkyl, substituted or unsubstituted C_{1-6} alkyl or substituted or unsubstituted C_{3-8} cycloalkyl;
- R^8 is H, substituted or unsubstituted C_{1-6} alkyl or halogen.; and
- 20 with the provisos that

- b). when R^8 is substituted or unsubstituted C_{1-6} alkyl, the alkyl group does not contain a carbonyl group directly attached to the phenyl ring;
- c). the compound is not of the following structures:



In another aspect, the invention provides a compound having Formula I

wherein:

R^1 is H;

- 10 R^2 is substituted or unsubstituted C_{1-6} alkyl, substituted or unsubstituted C_{3-8} cycloalkyl or substituted or unsubstituted C_{3-8} cycloalkenyl;

R^5 is -S(O)-;

R^6 is substituted or unsubstituted heterocycle or substituted or unsubstituted C_{6-10} aryl;

R^{17} is H, substituted or unsubstituted C_{1-6} alkyl or halogen;

- 15 R^{18} is H, substituted or unsubstituted C_{1-6} alkyl or halogen;

R^7 is halogen, CN, -OC $_{1-6}$ alkyl, substituted or unsubstituted C_{1-6} alkyl or substituted or unsubstituted C_{3-8} cycloalkyl;

R^8 is H, substituted or unsubstituted C_{1-6} alkyl or halogen; and

with the proviso that

- 20 b). when R^8 is substituted or unsubstituted C_{1-6} alkyl, the alkyl group does not contain a carbonyl group directly attached to the phenyl ring.

In another aspect, the invention provides a compound having Formula I

wherein:

R^1 is H;

R^2 is substituted or unsubstituted C_{1-6} alkyl, substituted or unsubstituted C_{3-8}

5 cycloalkyl or substituted or unsubstituted C_{3-8} cycloalkenyl;

R^5 is $-S(O)_2-$;

R^6 is substituted or unsubstituted heterocycle or substituted or unsubstituted C_{6-10} aryl;

R^{17} is H, substituted or unsubstituted C_{1-6} alkyl or halogen;

R^{18} is H, substituted or unsubstituted C_{1-6} alkyl or halogen;

10 R^7 is halogen, CN, substituted or unsubstituted $-OC_{1-6}$ alkyl, substituted or unsubstituted C_{1-6} alkyl or substituted or unsubstituted C_{3-8} cycloalkyl;

R^8 is H, substituted or unsubstituted C_{1-6} alkyl or halogen; and with the proviso that

b). when R^8 is substituted or unsubstituted C_{1-6} alkyl, the alkyl group does not contain a carbonyl group directly attached to the phenyl ring.

15 In another aspect, the invention provides a compound having Formula I

wherein:

R^1 is H ;

R^2 is substituted or unsubstituted C_{1-6} alkyl;

R^5 is $-S-$, $-S(O)-$, or $-S(O)_2-$;

20 R^6 is substituted or unsubstituted heterocycle or substituted or unsubstituted phenyl;

R^{17} is H;


R^{18} is H;


R^7 is halogen, CN, substituted or unsubstituted $-OC_{1-6}$ alkyl;

R^8 is H; and

25 with the proviso that

c). the compound is not of the following structures:





In another aspect, the invention provides a compound having Formula I

wherein:

5 R^1 is H;

R^2 is C_{1-6} alkyl;

R^5 is -S;

R^6 is substituted or unsubstituted heterocycle or substituted or unsubstituted phenyl;

R^{17} is H;


10 R^{18} is H;


R^7 is halogen, CN or substituted or unsubstituted $-OC_{1-6}$ alkyl;

R^8 is H; and

with the proviso that

c). the compound is not of the following structures:





In another aspect, the invention provides a compound having Formula I

wherein:

- 5 R^1 is H;
 R^2 is substituted or unsubstituted C_{1-6} alkyl;
 R^5 is $-S(O)$;
 R^6 is substituted or unsubstituted heterocycle or substituted or unsubstituted phenyl;
 R^{17} is H;
 10 R^{18} is H;
 R^7 is halogen, CN or $-OC_{1-6}$ alkyl;
 R^8 is H.

In another aspect, the invention provides a compound having Formula I
 wherein:

- 15 R^1 is H;
 R^2 is substituted or unsubstituted C_{1-6} alkyl;
 R^5 is $-S(O)_2$;
 R^6 is substituted or unsubstituted heterocycle or substituted or unsubstituted phenyl;
 R^{17} is H;
 20 R^{18} is H;
 R^7 is halogen, CN or $-OC_{1-6}$ alkyl;
 R^8 is H.

In another aspect, the invention provides a compound having Formula I wherein:

R¹ is H;

R² is methyl, isopropyl, 2-hydroxyethyl, methylpropionate, 2-methylpyridine,
 5 ethylacetate, N,N-dimethylpropanamide, N-isopropylpropanamide, propamide,
 hydroxycyclopentyl, ethyl, N,N-dimethylacetamide, N-methylacetamide, 2-aminoethyl,
 H-imidazol-2-ylmethyl, 1H-imidazol-4-ylmethyl, 2-[(aminocarbonyl)amino]ethyl, tert-
 butyl pyridin-2-yl-carbamate, 6-aminopyridin-2-yl, 2-oxo-1,3-oxazolidin-5-yl-methyl,
 2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl, 3-aminopropyl, N,N-
 10 dimethylbutanamide, 1H-pyrazol-3-ylmethyl, methyl-1,3-thiazol-2-yl-acetamide, tert-
 butyl-1,3-thiazol-2-yl-carbamate, 2-amino-1,3-thiazol-4-yl)methyl, 3-methylpyridine, 3-
 nitrobenzyl, 3-methoxybenzyl, 5-nitro-1H-pyrazol-3-yl-methyl, 5-amino-1H-pyrazol-3-
 yl-methyl, 1-propyl-1H-imidazol-4-yl)methyl, tert-butyl -1-oxidopyridin-2-yl-carbamate,
 3-hydroxybenzyl, 5-amino-4H-1,2,4-triazol-3-yl)methyl, 2-pyridin-2-ylethyl, 2-(1H-
 15 pyrazol-4-yl)ethyl, 2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl, (2-fluoropyridin-3-yl)methyl,
 trifluoromethyl, benzyl, 3-nitrobenzyl, 4-nitrobenzyl, 2-aminobenzyl, 3-aminobenzyl or
 pyrimidin-2-ylmethyl;

R⁵ is -S(O)₂-, -S- or -S(O)-;

R⁶ is 4-chloro-3-trifluoromethyl-phenyl, 3,4-dichlorophenyl, 3-methoxyphenyl, 4-
 20 methyl-3-nitrophenyl, 4-chlorophenyl, 4-chloro-3-methylphenyl, 4-nitro-3-
 trifluoromethylphenyl, 2,4-difluorophenyl, 4-chloro-2-fluorophenyl, 3-chloro-2-
 fluorophenyl, 4-isopropylphenyl, 4-bromophenyl, 4-iodophenyl, 3-chlorophenyl , 2-
 benzofuran, 5-methyl-2-furan, 2-furan or phenyl ;

R⁷ is chlorine, cyano, methoxy or fluorine;

25 R¹⁷ is H;

R¹⁸ is H; and

R⁸ is H.

The term "alkyl", as used herein, refers to saturated, monovalent or divalent hydrocarbon moieties having linear or branched moieties or combinations thereof and
 30 containing 1 to 6 carbon atoms. One or more methylene (-CH₂-) groups, of the alkyl
 can be replaced by oxygen, sulfur, carbonyl, sulfoxide, nitrogen, sulfonyl, or by a
 divalent C₃₋₆ cycloalkyl. Hydrogen atoms on alkyl groups can be substituted by
 groups including, but not limited to: halogen, -OH, C₃₋₈ cycloalkyl, non-aromatic
 heterocycles, aromatic heterocycles, C₆₋₁₀ aryl, -O(C₁₋₆ alkyl), amine groups, amino

groups, NO₂, amide groups, sulfonamide groups, ester groups, aldehyde groups, carboxylic acids, ketone groups.

The term "cycloalkyl", as used herein, refers to a monovalent or divalent group of 3 to 8 carbon atoms, derived from a saturated cyclic hydrocarbon. Cycloalkyl groups can be monocyclic or polycyclic. Cycloalkyl can be substituted by groups including, but not limited to: halogen, -OH, C₃₋₈ cycloalkyl, non-aromatic heterocycles, aromatic heterocycles, C₆₋₁₀ aryl, -O(C₁₋₆ alkyl), amine groups, amino groups, NO₂, amide groups, carboxylic acids, sulfonamide groups, ester groups, aldehyde groups, ketone groups.

The term "cycloalkenyl", as used herein, refers to a monovalent or divalent group of 3 to 8 carbon atoms, derived from a saturated cycloalkyl having one or more double bonds. Cycloalkenyl groups can be monocyclic or polycyclic. Cycloalkenyl groups can be substituted by groups including, but not limited to: halogen, -OH, C₃₋₈ cycloalkyl, non-aromatic heterocycles, aromatic heterocycles, C₆₋₁₀ aryl, -O(C₁₋₆ alkyl), amine groups, amino groups, NO₂, amide groups, sulfonamide groups, carboxylic acids, ester groups, aldehyde groups, ketone groups.

The term "halogen", as used herein, refers to an atom of chlorine, bromine, fluorine, iodine.

The term "alkenyl", as used herein, refers to a monovalent or divalent hydrocarbon radical having 2 to 6 carbon atoms, derived from a saturated alkyl, having at least one double bond. C₂₋₆ alkenyl can be in the E or Z configuration. Alkenyl groups can be substituted by C₁₋₃ alkyl.

The term "alkynyl", as used herein, refers to a monovalent or divalent hydrocarbon radical having 2 to 6 carbon atoms, derived from a saturated alkyl, having at least one triple bond.

The term "heterocycle" as used herein, refers to a 3 to 10 membered ring, which can be aromatic or non-aromatic, saturated or non-saturated, containing at least one heteroatom selected from O or N or S or combinations of at least two thereof, interrupting the carbocyclic ring structure. The heterocyclic ring can be interrupted by a C=O; the S heteroatom can be oxidized. Heterocycles can be monocyclic or polycyclic. Heterocyclic ring moieties can be substituted groups including, but not limited to: halogens, -OH, C₃₋₈ cycloalkyl, non-aromatic heterocycles, aromatic heterocycles, -OC₁₋₆ alkyl, -NH₂, -NO₂, amides, ethers, esters, ketones, carboxylic acids, aldehydes, sulfonamides groups.

Preferred substituted heterocycle groups are, but not limited to: pyridine, furan, azetidine, thiazol, thiophene, oxazol, pyrazol, isoxazole, 2-oxoindoline, 2-oxo-2,3-dihydro-1,3-benzoxazole, 2-oxo-2H-chromene, imidazole[2,1-b]thiazole, 1-H-pyrazole, indole, imidazole, quinoline, 2-thiophene, 2-benzofuran, 5-methyl-2-furan, 5-oxazolidine-2-one, pyrimidine-2,4(1*H*,3*H*)-dione, pyrimidine.

The term "aryl" as used herein, refers to an organic moiety derived from an aromatic hydrocarbon consisting of a ring containing 6 to 10 carbon atoms by removal of one hydrogen. Aryl can be substituted by groups including, but not limited to: halogens, -OH, C₃₋₈ cycloalkyl, non-aromatic heterocycles, aromatic heterocycles, -OC₁₋₆ alkyl, -NH₂, -NO₂, amides, ethers, esters, carboxylic acids, aldehydes, ketones, sulfonamides groups. Aryl can be monocyclic or bicyclic. Preferred substituted phenyl groups are, but not limited to: 4-chloro-3-trifluoromethyl-phenyl, 3,4-dichlorophenyl, 3-methoxyphenyl, 4-methyl-3-nitrophenyl, 4-chlorophenyl, 4-chloro-3-methylphenyl, nitro-3-trifluoromethylphenyl, 2,4-difluorophenyl, 4-chloro-2-fluorophenyl, 3-chloro-2-fluorophenyl, 4-isopropylphenyl, 4-bromophenyl, 4-iodophenyl, 3-chlorophenyl.

The term "amine" as used herein, represents a group of formula "-NR^xR^y", wherein R^x and R^y can be the same or independently H, alkyl, aryl, cycloalkyl, cycloalkenyl, heterocycle as defined above.

The term "amide" as used herein, represents a group of formula "-C(O)NR^xR^y", or "-C(O)N(R^x)(R^y)" or "NR^xC(O)R^y" wherein R^x and R^y can be the same or independently H, alkyl, aryl, cycloalkyl, cycloalkenyl, heterocycle as defined above.

The term "sulfonamide" as used herein, represents a group of formula "-S(O)₂NR^xR^y" or "NR^xR^yS(O)₂" or "-NR^xS(O)₂R^y" wherein R^x and R^y can be the same or independently H, alkyl, aryl, cycloalkyl, cycloalkenyl, heterocycle as defined above.

The term "ester" as used herein, represents a group of formula "-C(O)O(R^x)", wherein R^x is alkyl, aryl, cycloalkyl, cycloalkenyl, heterocycle as defined above.

The term "aldehyde" as used herein, represents a group of formula "-C(O)H".

The term "ketone" as used herein, represents a group of formula "-C(O)R^x", wherein R^x is C₁₋₆ alkyl.

The term "hydroxyl" as used herein, represents a group of formula "-OH".

The term "amino" as used herein, represents a group of formula "-NH₂".

The term "carbonyl" as used herein, represents a group of formula "-C(O)".

The term "carboxyl" as used herein, represents a group of formula "-C(O)O-".

The term "sulfonyl" as used herein, represents a group of formula "-SO₂".

The term "sulfate" as used herein, represents a group of formula " $-\text{O}-\text{S}(\text{O})_2-\text{O}-$ ".

The term "carboxylic acid" as used herein, represents a group of formula " $-\text{C}(\text{O})\text{OH}$ ".

The term "sulfoxide" as used herein, represents a group of formula " $-\text{S}=\text{O}$ ".

5 The term "phosphonic acid" as used herein, represents a group of formula " $-\text{P}(\text{O})(\text{OH})_2$ ".

The term "phosphoric acid" as used herein, represents a group of formula " $-(\text{O})\text{P}(\text{O})(\text{OH})_2$ ".

10 The term "sulphonic acid" as used herein, represents a group of formula " $-\text{S}(\text{O})_2\text{OH}$ ".

The formula "H", as used herein, represents a hydrogen atom.

The formula "O", as used herein, represents an oxygen atom.

The formula "N", as used herein, represents a nitrogen atom.

The formula "S", as used herein, represents a sulfur atom.

15 Some compounds of the invention are:

4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide;

4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide;

4-chloro-N-[5-chloro-2-(methylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide;

4-chloro-N-[5-chloro-2-(isopropylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide;

20 4-chloro-N-[5-chloro-2-[(2-hydroxyethyl)thio]phenyl]-3-(trifluoromethyl)benzenesulfonamide;

4-chloro-N-[5-chloro-2-(isopropylsulfinyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide;

25 4-chloro-N-[5-chloro-2-[(2-hydroxyethyl)sulfinyl]phenyl]-3-(trifluoromethyl)benzenesulfonamide;

4-chloro-N-[5-chloro-2-[(2-hydroxyethyl)sulfonyl]phenyl]-3-(trifluoromethyl)benzenesulfonamide;

methyl 3-[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl]propanoate;

30 4-chloro-N-[5-chloro-2-[(pyridin-2-ylmethyl)thio]phenyl]-3-(trifluoromethyl)benzenesulfonamide;

4-chloro-N-[5-chloro-2-[(pyridin-3-ylmethyl)thio]phenyl]-3-(trifluoromethyl)benzenesulfonamide;

- 4-chloro-N-{5-chloro-2-[(pyridin-2-ylmethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-{5-chloro-2-[(pyridin-3-ylmethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
- 5 ethyl {[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}acetate;
3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio}-N-isopropylpropanamide;
3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio}-N,N-
- 10 dimethylpropanamide;
3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}-N-isopropylpropanamide;
3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}-N,N-
- 15 dimethylpropanamide;
3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}propanamide;
N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide;
3,4-dichloro-N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide;
N-[5-chloro-2-(methylthio)phenyl]thiophene-2-sulfonamide;
- 20 N-[5-chloro-2-(methylsulfinyl)phenyl]thiophene-2-sulfonamide;
4-chloro-N-{5-chloro-2-[(3-hydroxycyclopentyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-{5-chloro-2-[(3-hydroxycyclopentyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
- 25 4-chloro-N-[5-chloro-2-(ethylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-[5-chloro-2-(ethylsulfinyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-[5-chloro-2-(ethylsulfonyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
N-[5-chloro-2-(methylthio)phenyl]-1-benzofuran-2-sulfonamide;
N-[5-chloro-2-(methylsulfinyl)phenyl]-1-benzofuran-2-sulfonamide;
- 30 N-[5-chloro-2-(methylsulfonyl)phenyl]-1-benzofuran-2-sulfonamide;
N-[5-chloro-2-(methylthio)phenyl]-4-methyl-3-nitrobenzenesulfonamide;
4-chloro-N-[5-chloro-2-(methylthio)phenyl]benzenesulfonamide;
4-chloro-N-[5-chloro-2-(methylthio)phenyl]-3-methylbenzenesulfonamide;
N-[5-chloro-2-(methylsulfinyl)phenyl]-4-methyl-3-nitrobenzenesulfonamide;

- N-[5-chloro-2-(methylsulfonyl)phenyl]-4-methyl-3-nitrobenzenesulfonamide;
4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide;
4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]benzenesulfonamide;
4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-3-methylbenzenesulfonamide;
5 4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-3-methylbenzenesulfonamide;
2-[[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino]phenyl]thio}-N,N-
dimethylacetamide;
2-[[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino]phenyl]thio}-N-
methylacetamide;
10 2-[[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino]phenyl]sulfinyl}-N,N-
dimethylacetamide;
2-[[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino]phenyl]sulfonyl}-
N,N-dimethylacetamide;
2-[[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino]phenyl]sulfinyl}-N-
15 methylacetamide;
2-[[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino]phenyl]sulfonyl}-N-
methylacetamide;
N-{2-[(2-aminoethyl)thio]-5-chlorophenyl}-4-chloro-3-
(trifluoromethyl)benzenesulfonamide;
20 N-[5-chloro-2-(methylthio)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide;
N-[5-chloro-2-(methylsulfonyl)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide;
N-[5-chloro-2-(methylthio)phenyl]-2,4-difluorobenzenesulfonamide;
N-[5-chloro-2-(methylsulfonyl)phenyl]-2,4-difluorobenzenesulfonamide;
N-[5-chloro-2-(methylsulfinyl)phenyl]-2,4-difluorobenzenesulfonamide;
25 N-[5-chloro-2-(methylsulfinyl)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide;
N-[5-chloro-2-(methylthio)phenyl]-5-methylfuran-2-sulfonamide;
N-[5-chloro-2-(methylsulfinyl)phenyl]-5-methylfuran-2-sulfonamide;
N-[5-chloro-2-(methylsulfonyl)phenyl]-5-methylfuran-2-sulfonamide;
N-[5-chloro-2-(methylthio)phenyl]furan-2-sulfonamide;
30 N-[5-chloro-2-(methylsulfinyl)phenyl]furan-2-sulfonamide;
N-[5-chloro-2-(methylsulfonyl)phenyl]furan-2-sulfonamide;
N-{2-[(2-aminoethyl)sulfonyl]-5-chlorophenyl}-4-chloro-3-
(trifluoromethyl)benzenesulfonamide;
4-chloro-N-[5-chloro-2-(methylthio)phenyl]-2-fluorobenzenesulfonamide;

- 3-chloro-N-[5-chloro-2-(methylthio)phenyl]-2-fluorobenzenesulfonamide;
3-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-2-fluorobenzenesulfonamide;
3-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-2-fluorobenzenesulfonamide;
4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-2-fluorobenzenesulfonamide;
5 4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-2-fluorobenzenesulfonamide;
N-{2-[(2-aminoethyl)sulfinyl]-5-chlorophenyl}-4-chloro-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
10 4-chloro-N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
15 4-chloro-N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
20 N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
25 N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
3-({2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}thio)-N,N-dimethylpropanamide;
3-({2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfinyl)-N,N-dimethylpropanamide;
30 3-({2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfonyl)-N,N-dimethylpropanamide;
N-[2-({2-[(aminocarbonyl)amino]ethyl}thio)-5-chlorophenyl]-1-benzofuran-2-sulfonamide;

- N-[2-({2-[(aminocarbonyl)amino]ethyl}sulfinyl)-5-chlorophenyl]-1-benzofuran-2-sulfonamide;
- N-[2-({2-[(aminocarbonyl)amino]ethyl}sulfonyl)-5-chlorophenyl]-1-benzofuran-2-sulfonamide;
- 5 tert-butyl {6-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfinyl)methyl]pyridin-2-yl}carbamate;
- N-(2-[(6-aminopyridin-2-yl)methyl]thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- N-(2-[(6-aminopyridin-2-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- 10 N-(2-[(6-aminopyridin-2-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- N-(2-[(6-amino-1-oxidopyridin-2-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-{[3-(dimethylamino)propyl]thio}phenyl)-1-benzofuran-2-sulfonamide;
- 15 N-(5-chloro-2-{[3-(dimethylamino)propyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-{[3-(dimethylamino)propyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-{[3-(dimethylnitro)propyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;
- N-(2-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfonyl}ethyl)acetamide;
- 20 N-(2-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}ethyl)acetamide;
- N-(5-chloro-2-{[(2-oxo-1,3-oxazolidin-5-yl)methyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-{[(2-oxo-1,3-oxazolidin-5-yl)methyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;
- 25 N-(5-chloro-2-{[(2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl]thio}phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-{[(2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;
- 30 N-(5-chloro-2-{[(2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminopropyl)thio]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminopropyl)sulfinyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminopropyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;

- 4-({2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfinyl)-N,N-dimethylbutanamide;
- 4-({2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfonyl)-N,N-dimethylbutanamide;
- 5 5-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-1-benzofuran-2-sulfonamide;
- 5-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(1H-pyrazol-3-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(1H-pyrazol-3-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(1H-pyrazol-3-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
- 10 N-(5-chloro-2-[(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)methyl]sulfinyl)phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)methyl]sulfonyl)phenyl)-1-benzofuran-2-sulfonamide;
- N-{4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl]-1,3-thiazol-2-yl}acetamide;
- 15 N-{4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfinyl)methyl]-1,3-thiazol-2-yl}acetamide;
- N-{4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]-1,3-thiazol-2-yl}acetamide;
- 20 N-[5-chloro-2-(methylsulfinyl)phenyl]-4-isopropylbenzenesulfonamide;
- 4-bromo-N-[5-chloro-2-(methylthio)phenyl]benzenesulfonamide;
- N-[5-chloro-2-(methylthio)phenyl]-4-iodobenzenesulfonamide;
- N-[5-chloro-2-(methylsulfonyl)phenyl]-4-isopropylbenzenesulfonamide;
- 4-bromo-N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide;
- 25 N-[5-chloro-2-(methylsulfinyl)phenyl]-4-iodobenzenesulfonamide;
- 4-bromo-N-[5-chloro-2-(methylsulfonyl)phenyl]benzenesulfonamide;
- N-[5-chloro-2-(methylsulfonyl)phenyl]-4-iodobenzenesulfonamide;
- tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl]-1,3-thiazol-2-yl}carbamate;
- 30 N-(2-[(2-amino-1,3-thiazol-4-yl)methyl]thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfinyl)methyl]-1,3-thiazol-2-yl}carbamate;

N-(2-[[2-amino-1,3-thiazol-4-yl)methyl]sulfinyl]-5-chlorophenyl)-1-benzofuran-2-sulfonamide;

tert-butyl {4-[[2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl]sulfonyl)methyl]-1,3-thiazol-2-yl}carbamate;

5 N-(2-[[2-amino-1,3-thiazol-4-yl)methyl]sulfonyl]-5-chlorophenyl)-1-benzofuran-2-sulfonamide;

N-{5-chloro-2-[(pyridin-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;

N-{5-chloro-2-[(pyridin-3-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;

N-{5-chloro-2-[(pyridin-3-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;

10 N-(5-chloro-2-[(1-oxidopyridin-3-yl)methyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;

N-{5-chloro-2-[(pyridin-3-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;

N-(5-chloro-2-[(1-oxidopyridin-3-yl)methyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;

15 N-{5-chloro-2-[(3-nitrobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide;

N-{5-chloro-2-[(3-methoxybenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide;

N-(5-chloro-2-[(5-nitro-1H-pyrazol-3-yl)methyl]thio}phenyl)-1-benzofuran-2-sulfonamide;

20 N-(5-chloro-2-[(5-nitro-1H-pyrazol-3-yl)methyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;

N-(2-[(5-amino-1H-pyrazol-3-yl)methyl]thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;

N-(2-[(5-amino-1H-pyrazol-3-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;

25 N-(2-[(5-amino-1H-pyrazol-3-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;

N-(5-chloro-2-[(1-propyl-1H-imidazol-4-yl)methyl]thio}phenyl)-1-benzofuran-2-sulfonamide;

30 N-(5-chloro-2-[(1-propyl-1H-imidazol-4-yl)methyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;

N-(5-chloro-2-[(1-propyl-1H-imidazol-4-yl)methyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;

N-{5-chloro-2-[(pyridin-2-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;

N-{5-chloro-2-[(pyridin-2-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;

- N-(5-chloro-2-[(1-oxidopyridin-2-yl)methyl]sulfonyl)phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[(1-oxidopyridin-2-yl)methyl]sulfinyl)phenyl)-1-benzofuran-2-sulfonamide;
- 5 tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl]pyridin-2-yl}carbamate;
- N-(2-[(2-aminopyridin-4-yl)methyl]thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]pyridin-2-yl}carbamate;
- 10 tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]-1-oxidopyridin-2-yl}carbamate;
- N-(2-[(2-aminopyridin-4-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- N-(2-[(2-aminopyridin-4-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- 15 N-(2-[(2-amino-1-oxidopyridin-4-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-1-benzofuran-2-sulfonamide
- N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- 20 N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(3-hydroxybenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(3-hydroxybenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(3-hydroxybenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
- 3-[4-chloro-2-[(4-chloro-3-(trifluoromethyl)phenyl)sulfonyl]amino]phenyl)sulfonyl]-
- 25 N,N-dimethylpropanamide;
- 3-[(4-chloro-2-[(2,4-difluorophenyl)sulfonyl]amino)phenyl]thio]-N,N-dimethylpropanamide;
- N-(2-[(5-amino-4H-1,2,4-triazol-3-yl)methyl]thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- 30 N-(2-[(5-amino-4H-1,2,4-triazol-3-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- 3-[(4-chloro-2-[(2,4-difluorophenyl)sulfonyl]amino)phenyl)sulfinyl]-N,N-dimethylpropanamide;

- 3-[(4-chloro-2-[(2,4-difluorophenyl)sulfonyl]amino}phenyl)sulfonyl]-N,N-dimethylpropanamide;
- 3-[(4-chloro-2-[(4-chloro-2-fluorophenyl)sulfonyl]amino}phenyl)thio]-N,N-dimethylpropanamide;
- 5 3-[(4-chloro-2-[(4-chloro-2-fluorophenyl)sulfonyl]amino}phenyl)sulfinyl]-N,N-dimethylpropanamide;
- 3-[(4-chloro-2-[(4-chloro-2-fluorophenyl)sulfonyl]amino}phenyl)sulfonyl]-N,N-dimethylpropanamide;
- N-{5-chloro-2-[(2-pyridin-2-ylethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
- 10 N-(5-chloro-2-{[2-(1H-pyrazol-4-yl)ethyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(2-pyridin-2-ylethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(2-pyridin-2-ylethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-{[2-(1H-pyrazol-4-yl)ethyl]thio}phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-{[2-(1H-pyrazol-4-yl)ethyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;
- 15 N-(5-chloro-2-{[2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl]thio}phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-{[2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-{[2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl]sulfonyl}phenyl)-1-benzofuran-2-
- 20 sulfonamide;
- N-(5-chloro-2-{[(2-fluoropyridin-3-yl)methyl]thio}phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-{[(2-fluoropyridin-3-yl)methyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-{[(2-fluoropyridin-3-yl)methyl]sulfonyl}phenyl)-1-benzofuran-2-
- 25 sulfonamide;
- N-{5-chloro-2-[(trifluoromethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(trifluoromethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(trifluoromethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-[2-(benzylthio)-5-chlorophenyl]-1-benzofuran-2-sulfonamide;
- 30 N-[2-(benzylsulfinyl)-5-chlorophenyl]-1-benzofuran-2-sulfonamide;
- N-[2-(benzylsulfonyl)-5-chlorophenyl]-1-benzofuran-2-sulfonamide;
- N-[2-[(3-aminobenzyl)thio]-5-fluorophenyl]-1-benzofuran-2-sulfonamide;
- N-{5-fluoro-2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-[2-[(3-aminobenzyl)sulfinyl]-5-fluorophenyl]-1-benzofuran-2-sulfonamide;

- N-{5-fluoro-2-[(3-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
N-{2-[(3-aminobenzyl)sulfonyl]-5-fluorophenyl}-1-benzofuran-2-sulfonamide;
N-{2-[(3-aminobenzyl)thio]-5-methoxyphenyl}-1-benzofuran-2-sulfonamide;
N-{2-[(3-aminobenzyl)sulfonyl]-5-methoxyphenyl}-1-benzofuran-2-sulfonamide;
5 N-{2-[(3-aminobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-{2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
N-{2-[(3-aminobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-methoxy-2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
N-{2-[(3-aminobenzyl)sulfinyl]-5-methoxyphenyl}-1-benzofuran-2-sulfonamide;
10 N-{2-[(3-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
N-{2-[(3-aminobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(4-nitrobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-{2-[(2-aminobenzyl)thio]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(4-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
15 N-{2-[(4-aminobenzyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(2-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
N-{2-[(2-aminobenzyl)sulfinyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
N-{2-[(2-aminobenzyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-2,4-difluorobenzenesulfonamide;
20 N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-4-chloro-2-fluorobenzenesulfonamide;
N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-2,4-difluorobenzenesulfonamide;
N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-2,4-difluorobenzenesulfonamide;
N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-4-chloro-2-fluorobenzenesulfonamide;
N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-4-chloro-2-fluorobenzenesulfonamide;
25 N-{5-chloro-2-[(pyrimidin-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(pyrimidin-2-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(pyrimidin-2-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}benzenesulfonamide;
N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-4-chlorobenzenesulfonamide;
30 N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-3-chlorobenzenesulfonamide;
N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}benzenesulfonamide;
N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}benzenesulfonamide;
N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-4-chlorobenzenesulfonamide;
N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-4-chlorobenzenesulfonamide;

- N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-3-chlorobenzenesulfonamide;
N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-3-chlorobenzenesulfonamide;
N-[2-(benzylthio)-5-cyanophenyl]-1-benzofuran-2-sulfonamide;
N-[2-(benzylsulfinyl)-5-cyanophenyl]-1-benzofuran-2-sulfonamide;
5 N-[2-(benzylsulfonyl)-5-cyanophenyl]-1-benzofuran-2-sulfonamide.

Some compounds of Formula I and some of their intermediates have at least one stereogenic center in their structure. This stereogenic center may be present in an R or S configuration, said R and S notation is used in correspondence with the rules described in Pure Appl. Chem. (1976), 45, 11-13.

- 10 The term "pharmaceutically acceptable salts" refers to salts or complexes that retain the desired biological activity of the above identified compounds and exhibit minimal or no undesired toxicological effects. The "pharmaceutically acceptable salts" according to the invention include therapeutically active, non-toxic base or acid salt forms, which the compounds of Formula I are able to form.

- 15 The acid addition salt form of a compound of Formula I that occurs in its free form as a base can be obtained by treating the free base with an appropriate acid such as an inorganic acid, for example, a hydrohalic acid, such as hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; or an organic acid such as for example, acetic, hydroxyacetic, propanoic, lactic, pyruvic, malonic,
20 fumaric acid, maleic acid, oxalic acid, tartaric acid, succinic acid, malic acid, ascorbic acid, benzoic acid, tannic acid, pantoic acid, citric, methylsulfonic, ethanesulfonic, benzenesulfonic, formic and the like (Handbook of Pharmaceutical Salts, P.Heinrich Stahl& Camille G. Wermuth (Eds), Verlag Helvetica Chemica Acta- Zürich, 2002, 329-345).

- 25 The base addition salt form of a compound of Formula I that occurs in its acid form can be obtained by treating the acid with an appropriate base such as an inorganic base, for example, sodium hydroxide, magnesium hydroxide, potassium hydroxide, Calcium hydroxide, ammonia and the like; or an organic base such as for example, L-Arginine, ethanolamine, betaine, benzathine, morpholine and the like.
30 (Handbook of Pharmaceutical Salts, P.Heinrich Stahl& Camille G. Wermuth (Eds), Verlag Helvetica Chemica Acta- Zürich, 2002, 329-345).

Compounds of Formula I and their salts can be in the form of a solvate, which is included within the scope of the present invention. Such solvates include for example hydrates, alcoholates and the like.

With respect to the present invention reference to a compound or compounds,
5 is intended to encompass that compound in each of its possible isomeric forms and mixtures thereof unless the particular isomeric form is referred to specifically.

Compounds according to the present invention may exist in different polymorphic forms. Although not explicitly indicated in the above formula, such forms are intended to be included within the scope of the present invention.

10 The compounds of the invention are indicated for use in treating or preventing conditions in which there is likely to be a component involving the chemokine receptors.

In another embodiment, there are provided pharmaceutical compositions including at least one compound of the invention in a pharmaceutically acceptable
15 carrier.

In a further embodiment of the invention, there are provided methods for treating disorders associated with modulation of chemokine receptors. Such methods can be performed, for example, by administering to a subject in need thereof a pharmaceutical composition containing a therapeutically effective amount of at least
20 one compound of the invention.

These compounds are useful for the treatment of mammals, including humans, with a range of conditions and diseases that are alleviated by CCR modulation. Therapeutic utilities of CCR modulators are skin inflammatory diseases and conditions, including, but are not limited to: rosacea (dilation of the blood vessels just
25 under the skin), sunburn, chronic sun damage, discrete erythemas, psoriasis, atopic dermatitis, menopause-associated hot flashes, hot flashes resulting from orchiectomy atopic dermatitis, photoaging, seborrheic dermatitis, acne, allergic dermatitis, irritant dermatitis, telangiectasia (dilations of previously existing small blood vessels) of the face, rhinophyma (hypertrophy of the nose with follicular dilation), red
30 bulbous nose, acne-like skin eruptions (may ooze or crust), burning or stinging

sensation of the face, irritated and bloodshot and watery eyes, cutaneous hyperactivity with dilation of blood vessels of the skin, Lyell's syndrome, Stevens-Johnson syndrome, erythema multiforme minor, erythema multiforme major and other inflammatory skin diseases, actinic keratoses, arsenic keratoses, inflammatory and
5 non-inflammatory acne, ichthyoses and other keratinization and hyperproliferative disorders of the skin, eczema, wound healing.

Therapeutic utilities of CCR modulators are ocular inflammatory diseases including, but not limited to, uveitis, retinal degenerative conditions, angiogenesis, dry eye, Keratitis, allergic eye disease and conditions affecting the posterior part of the
10 eye, such as maculopathies and retinal degeneration including non-exudative age related macular degeneration, exudative age related macular degeneration, choroidal neovascularization, diabetic retinopathy, acute macular neuroretinopathy, central serous chorioretinopathy, cystoid macular edema, and diabetic macular edema; uveitis, retinitis, and choroiditis such as acute multifocal placoid pigment
15 epitheliopathy, Behcet's disease, birdshot retinochoroidopathy, infectious (syphilis, lyme, tuberculosis, toxoplasmosis), intermediate uveitis (pars planitis), multifocal choroiditis, multiple evanescent white dot syndrome (mewds), ocular sarcoidosis, posterior scleritis, serpiginous choroiditis, subretinal fibrosis and uveitis syndrome, Vogt-Koyanagi-and Harada syndrome; vasuclar diseases/ exudative diseases such as
20 retinal arterial occlusive disease, central retinal vein occlusion, disseminated intravascular coagulopathy, branch retinal vein occlusion, hypertensive fundus changes, ocular ischemic syndrome, retinal arterial microaneurysms, Coat's disease, parafoveal telangiectasis, hemi-retinal vein occlusion, papillophlebitis, central retinal artery occlusion, branch retinal artery occlusion, carotid artery disease (CAD), frosted
25 branch angiitis, sickle cell retinopathy and other hemoglobinopathies, angioid streaks, familial exudative vitreoretinopathy, and Eales disease; traumatic/ surgical conditions such as sympathetic ophthalmia, uveitic retinal disease, retinal detachment, trauma, conditions caused by laser, conditions caused by photodynamic therapy, photocoagulation, hypoperfusion during surgery, radiation retinopathy, and bone
30 marrow transplant retinopathy; proliferative disorders such as proliferative vitreal retinopathy and epiretinal membranes, and proliferative diabetic retinopathy; infectious disorders such as ocular histoplasmosis, ocular toxocariasis, presumed ocular histoplasmosis syndrome (POHS), endophthalmitis, toxoplasmosis, retinal diseases

associated with HIV infection, choroidal disease associate with HIV infection, uveitic disease associate with HIV infection, viral retinitis, acute retinal necrosis, progressive outer retinal necrosis, fungal retinal diseases, ocular syphilis, ocular tuberculosis, diffuse unilateral subacute neuroretinitis, and myiasis; genetic disorders such as
5 retinitis pigmentosa, systemic disorders with accosiated retinal dystrophies, congenital stationary night blindness, cone dystrophies, Stargardt's disease and fundus flavimaculatus, Best's disease, pattern dystrophy of the retinal pigmented epithelium, X-linked retinoschisis, Sorsby's fundus dystrophy, benign concentric maculopathy, Bietti's crystalline dystrophy, and pseudoxanthoma elasticum; retinal tears/ holes such
10 as retinal detachment, macular hole, and giant retinal tear; tumors such as retinal disease associated with tumors, congenital hypertrophy of the retinal pigmented epithelium, posterior uveal melanoma, choroidal hemangioma, choroidal osteoma, choroidal metastasis, combined hamartoma of the retina and retinal pigmented epithelium, retinoblastoma, vasoproliferative tumors of the ocular fundus, retinal
15 astrocytoma, and intraocular lymphoid tumors; and miscellaneous other diseases affecting the posterior part of the eye such as punctate inner choroidopathy, acute posterior multifocal placoid pigment epitheliopathy, myopic retinal degeneration, and acute retinal pigement epitheliitis.

In still another embodiment of the invention, there are provided methods for
20 treating disorders associated with modulation of chemokine receptors. Such methods can be performed, for example, by administering to a subject in need thereof a therapeutically effective amount of at least one compound of the invention, or any combination thereof, or pharmaceutically acceptable salts, hydrates, solvates, crystal forms and individual isomers, enantiomers, and diastereomers thereof.

25 The present invention concerns the use of a compound of Formula I or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of ocular inflammatory diseases including, but not limited to, uveitis, dry eye, Keratitis, allergic eye disease and conditions affecting the posterior part of the eye, such as maculopathies and retinal degeneration including non-exudative age related
30 macular degeneration, exudative age related macular degeneration, choroidal neovascularization, diabetic retinopathy, acute macular neuroretinopathy, central serous chorioretinopathy, cystoid macular edema, and diabetic macular edema;

uveitis, retinitis, and choroiditis such as acute multifocal placoid pigment
 epitheliopathy, Behcet's disease, birdshot retinochoroidopathy, infectious (syphilis,
 lyme, tuberculosis, toxoplasmosis), intermediate uveitis (pars planitis), multifocal
 choroiditis, multiple evanescent white dot syndrome (mewds), ocular sarcoidosis,
 5 posterior scleritis, serpiginous choroiditis, subretinal fibrosis and uveitis syndrome,
 Vogt-Koyanagi-and Harada syndrome; vasuclar diseases/ exudative diseases such as
 retinal arterial occlusive disease, central retinal vein occlusion, disseminated
 intravascular coagulopathy, branch retinal vein occlusion, hypertensive fundus
 changes, ocular ischemic syndrome, retinal arterial microaneurysms, Coat's disease,
 10 parafoveal telangiectasis, hemi-retinal vein occlusion, papillophlebitis, central retinal
 artery occlusion, branch retinal artery occlusion, carotid artery disease (CAD), frosted
 branch angiitis, sickle cell retinopathy and other hemoglobinopathies, angioid streaks,
 familial exudative vitreoretinopathy, and Eales disease; traumatic/ surgical conditions
 such as sympathetic ophthalmia, uveitic retinal disease, retinal detachment, trauma,
 15 conditions caused by laser, conditions caused by photodynamic therapy,
 photocoagulation, hypoperfusion during surgery, radiation retinopathy, and bone
 marrow transplant retinopathy; proliferative disorders such as proliferative vitreal
 retinopathy and epiretinal membranes, and proliferative diabetic retinopathy; infectious
 disorders such as ocular histoplasmosis, ocular toxocariasis, presumed ocular
 20 histoplasmosis syndrome (POHS), endophthalmitis, toxoplasmosis, retinal diseases
 associated with HIV infection, choroidal disease associate with HIV infection, uveitic
 disease associate with HIV infection, viral retinitis, acute retinal necrosis, progressive
 outer retinal necrosis, fungal retinal diseases, ocular syphilis, ocular tuberculosis,
 diffuse unilateral subacute neuroretinitis, and myiasis; genetic disorders such as
 25 retinitis pigmentosa, systemic disorders with accosiated retinal dystrophies, congenital
 stationary night blindness, cone dystrophies, Stargardt's disease and fundus
 flavimaculatus, Best's disease, pattern dystrophy of the retinal pigmented epithelium,
 X-linked retinoschisis, Sorsby's fundus dystrophy, benign concentric maculopathy,
 Bietti's crystalline dystrophy, and pseudoxanthoma elasticum; retinal tears/ holes such
 30 as retinal detachment, macular hole, and giant retinal tear; tumors such as retinal
 disease associated with tumors, congenital hypertrophy of the retinal pigmented
 epithelium, posterior uveal melanoma, choroidal hemangioma, choroidal osteoma,
 choroidal metastasis, combined hamartoma of the retina and retinal pigmented
 epithelium, retinoblastoma, vasoproliferative tumors of the ocular fundus, retinal

astrocytoma, and intraocular lymphoid tumors; and miscellaneous other diseases affecting the posterior part of the eye such as punctate inner choroidopathy, acute posterior multifocal placoid pigment epitheliopathy, myopic retinal degeneration, and acute retinal pigment epitheliitis.

5 The actual amount of the compound to be administered in any given case will be determined by a physician taking into account the relevant circumstances, such as the severity of the condition, the age and weight of the patient, the patient's general physical condition, the cause of the condition, and the route of administration.

10 The patient will be administered the compound orally in any acceptable form, such as a tablet, liquid, capsule, powder and the like, or other routes may be desirable or necessary, particularly if the patient suffers from nausea. Such other routes may include, without exception, transdermal, parenteral, subcutaneous, intranasal, via an implant stent, intrathecal, intravitreal, topical to the eye, back to the eye, intramuscular, intravenous, and intrarectal modes of delivery. Additionally, the
15 formulations may be designed to delay release of the active compound over a given period of time, or to carefully control the amount of drug released at a given time during the course of therapy.

20 In another embodiment of the invention, there are provided pharmaceutical compositions including at least one compound of the invention in a pharmaceutically acceptable carrier thereof. The phrase "pharmaceutically acceptable" means the carrier, diluent or excipient must be compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

25 Pharmaceutical compositions of the present invention can be used in the form of a solid, a solution, an emulsion, a dispersion, a patch, a micelle, a liposome, and the like, wherein the resulting composition contains one or more compounds of the present invention, as an active ingredient, in admixture with an organic or inorganic carrier or excipient suitable for enteral or parenteral applications. Invention
30 compounds may be combined, for example, with the usual non-toxic, pharmaceutically acceptable carriers for tablets, pellets, capsules, suppositories, solutions, emulsions, suspensions, and any other form suitable for use. The carriers which can be used include glucose, lactose, gum acacia, gelatin, mannitol, starch paste, magnesium

trisilicate, talc, corn starch, keratin, colloidal silica, potato starch, urea, medium chain length triglycerides, dextrans, and other carriers suitable for use in manufacturing preparations, in solid, semisolid, or liquid form. In addition auxiliary, stabilizing, thickening and coloring agents and perfumes may be used. Invention compounds are
5 included in the pharmaceutical composition in an amount sufficient to produce the desired effect upon the process or disease condition.

Pharmaceutical compositions containing invention compounds may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or
10 syrups or elixirs. Compositions intended for oral use may be prepared according to any method known in the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of a sweetening agent such as sucrose, lactose, or saccharin, flavoring agents such as peppermint, oil of wintergreen or cherry, coloring agents and
15 preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets containing invention compounds in admixture with non-toxic pharmaceutically acceptable excipients may also be manufactured by known methods. The excipients used may be, for example, (1) inert diluents such as calcium carbonate, lactose, calcium phosphate or sodium phosphate; (2) granulating and
20 disintegrating agents such as corn starch, potato starch or alginic acid; (3) binding agents such as gum tragacanth, corn starch, gelatin or acacia, and (4) lubricating agents such as magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period.
25 For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

In some cases, formulations for oral use may be in the form of hard gelatin capsules wherein the invention compounds are mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin. They may also be in the
30 form of soft gelatin capsules wherein the invention compounds are mixed with water or an oil medium, for example, peanut oil, liquid paraffin or olive oil.

The pharmaceutical compositions may be in the form of a sterile injectable suspension. This suspension may be formulated according to known methods using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides, fatty acids (including oleic acid), naturally occurring vegetable oils like sesame oil, coconut oil, peanut oil, cottonseed oil, etc., or synthetic fatty vehicles like ethyl oleate or the like. Buffers, preservatives, antioxidants, and the like can be incorporated as required.

Invention compounds and their pharmaceutically-acceptable salts may be administered through different routes, including but not limited to topical eye drops, direct injection, application at the back of the eye or formulations that may further enhance the long duration of actions such as a slow releasing pellet, suspension, gel, or sustained delivery devices such as any suitable drug delivery system (DDS) known in the art. While topical administration is preferred, this compound may also be used in an intraocular implant as described in U.S. U.S. Patent 7,931,909.

Invention compounds may also be administered in the form of suppositories for rectal administration of the drug. These compositions may be prepared by mixing the invention compounds with a suitable non-irritating excipient, such as cocoa butter, synthetic glyceride esters of polyethylene glycols, which are solid at ordinary temperatures, but liquefy and/or dissolve in the rectal cavity to release the drug.

Since individual subjects may present a wide variation in severity of symptoms and each drug has its unique therapeutic characteristics, the precise mode of administration and dosage employed for each subject is left to the discretion of the practitioner.

The compounds and pharmaceutical compositions described herein are useful as medicaments in mammals, including humans, for treatment of diseases and/or alleviations of conditions which are responsive to treatment by agonists or functional antagonists of chemokine receptors. Thus, in further embodiments of the invention,

there are provided methods for treating a disorder associated with modulation of chemokine receptors. Such methods can be performed, for example, by administering to a subject in need thereof a pharmaceutical composition containing a therapeutically effective amount of at least one invention compound. As used herein, the term

5 "therapeutically effective amount" means the amount of the pharmaceutical composition that will elicit the biological or medical response of a subject in need thereof that is being sought by the researcher, veterinarian, medical doctor or other clinician. In some embodiments, the subject in need thereof is a mammal. In some embodiments, the mammal is human.

10 The present invention concerns also processes for preparing the compounds of Formula I. The compounds of formula I according to the invention can be prepared analogously to conventional methods as understood by the person skilled in the art of synthetic organic chemistry. The synthetic schemes set forth below, illustrate how compounds according to the invention can be made. Those skilled in the art will be
15 able to routinely modify and/or adapt Scheme 1 to synthesize any compounds of the invention covered by Formula I.

The described sulfur derivatives were prepared by general routes as shown in **Scheme 1**. In one route, an appropriately substituted 2-amino-benzenethiol such as **Intermediate A** can react with an electrophile such as a halide, tosylate, mesylate,
20 enone, 2-enoate etc. in the presence of an acid or base to prepare sulfide **Intermediate B**. Alternatively, sulfide **Intermediate B** can be prepared by treatment of **Intermediate A** with an alcohol under Mitsunobu conditions. Reaction of **Intermediate B** with a sulfonyl chloride provides the sulfonamide of **Formula I** wherein R^5 is S. Further, upon treatment with an oxidant such as meta-chloroperoxybenzoic acid, the
25 compound of Formula I wherein R^5 is S affords the compound wherein R^5 is S(O) or R^5 is S(O)₂.

In another route, di-sulfide **Intermediate E** type can be obtained through oxidation of an **Intermediate A** type. Reaction of **Intermediate E** with a, sulfonyl chloride provides sulfonamide **Intermediate F**. In situ or stepwise reduction of
30 **Intermediate F** using polymer bound triphenylphosphine or sodium borohydride, respectively, followed by reaction of the resulting benzenethiol with an electrophile affords the sulfonamide of **Formula I** wherein R^5 is S.

Scheme 1

DETAILED DESCRIPTION OF THE INVENTION

It is to be understood that both the foregoing general description and the
5 following detailed description are exemplary and explanatory only and are not
restrictive of the invention claimed. As used herein, the use of the singular includes
the plural unless specifically stated otherwise.

It will be readily apparent to those skilled in the art that some of the compounds
of the invention may contain one or more asymmetric centers, such that the
10 compounds may exist in enantiomeric as well as in diastereomeric forms. Unless it is
specifically noted otherwise, the scope of the present invention includes all
enantiomers, diastereomers and racemic mixtures. Some of the compounds of the
invention may form salts with pharmaceutically acceptable acids or bases, and such
pharmaceutically acceptable salts of the compounds described herein are also within
15 the scope of the invention.

The present invention includes all pharmaceutically acceptable isotopically
enriched compounds. Any compound of the invention may contain one or more
isotopic atoms enriched or different than the natural ratio such as deuterium ^2H (or D)
in place of protium ^1H (or H) or use of ^{13}C enriched material in place of ^{12}C and the
20 like. Similar substitutions can be employed for N, O and S. The use of isotopes may
assist in analytical as well as therapeutic aspects of the invention. For example, use
of deuterium may increase the in vivo half-life by altering the metabolism (rate) of the
compounds of the invention. These compounds can be prepared in accord with the
preparations described by use of isotopically enriched reagents.

25 As will be evident to those skilled in the art, individual isomeric forms can be
obtained by separation of mixtures thereof in conventional manner. For example, in
the case of diastereoisomeric isomers, chromatographic separation may be employed.

Compound names were generated with ACD version 8 and some
intermediates' and reagents' names used in the examples were generated with
30 software such as Chem Bio Draw Ultra version 12.0 or Auto Nom 2000 from MDL ISIS

Draw 2.5 SP1.. In general, characterization of the compounds is performed according to the following methods:

NMR spectra are recorded on *Varian* 600 or *Varian* 300, in the indicated solvent at ambient temperature; chemical shifts in [ppm], coupling constants in [Hz].

5 All the reagents, solvents, catalysts for which the synthesis is not described are purchased from chemical vendors such as Sigma Aldrich, Fluka, Bio-Blocks, Combi-blocks, TCI, VWR, Lancaster, Oakwood, Trans World Chemical, Alfa, Fisher, Maybridge, Frontier, Matrix, Ukrorgsynth, Toronto, Ryan Scientific, SiliCycle, Anaspec, Syn Chem, Chem-Impex, MIC-scientific, Ltd; however some known intermediates,
10 were prepared according to published procedures. Solvents were purchased from commercial sources in appropriate quality and used as received. Air and/or moisture-sensitive reactions were run under an Ar- or N₂- atmosphere.

Usually the compounds of the invention were purified by chromatography: CombiFlash Companion and RediSep Rf silica gel 60 (0.04-0.063 mm); Preparative
15 thin layer chromatography (PTLC): *Analtech* (silica gel 60 F₂₅₄, 500 or 1000 μm).

The following abbreviations are used in the examples:

	NH ₃	ammonia
	CH ₃ CN	acetonitrile
	CH ₂ Cl ₂	dichloromethane
20	DMF	<i>N,N</i> -dimethylformamide
	NaOH	sodium hydroxide
	MeOH	methanol
	CD ₃ OD	deuterated methanol
	HCl	hydrochloric acid
25	Na ₂ SO ₄	sodium sulfate
	HBTU	2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate)
	DIPEA	<i>N,N</i> -Diisopropylethylamine
	CuI	copper iodide
30	Cs ₂ CO ₃	cesium carbonate
	DMEDA	<i>N,N'</i> -dimethylethylenediamine

	MgSO ₄	magnesium sulfate
	EtOAc	ethyl acetate
	CDCl ₃	deuterated chloroform
	DMSO-d ₆	deuterated dimethyl sulfoxide
5	TFA	trifluoroacetic acid
	THF	tetrahydrofuran
	K ₂ CO ₃	potassium carbonate
	mCPBA	meta-Chloroperoxybenzoic acid
	NaBH ₄	sodium borohydride
10	CaCl ₂	calcium chloride


The following examples are for illustrative purposes only and are not intended, nor should they be construed as limiting the invention in any manner. Those skilled in the art will appreciate that variations and modifications of the following examples can be made without exceeding the spirit or scope of the invention.

15 Specific Examples

GENERAL PROCEDURE A


Intermediate 1

5-chloro-2-[(1H-imidazol-4-ylmethyl)thio]aniline



- 20 A mixture of 2-amino-4-chlorobenzenethiol (CAS 1004-00-8) (1.1 g, 7.1 mmol), 4-(chloromethyl)-1H-imidazole hydrochloride (721 mg, 4.71 mmol) and K₂CO₃ (3.2 g, 23.6 mmol) in DMF (10 ml) was stirred at room temperature over night. The reaction mixture was poured into water (50 ml) and extracted with ethyl acetate (2x50 ml). The organic layer was washed with brine, dried over Na₂SO₄, concentrated in *vacuo*. The
- 25 crude product was purified by column chromatography on silica gel (0→100% ethyl acetate in hexane) to give **Intermediate 1** as a solid (1.0 g, 60%).

¹H NMR (300 MHz, CD₃OD) δ 7.58 (s, 1H), 7.06 (d, *J* = 8.20 Hz, 1H), 6.73 (d, *J* = 2.34 Hz, 1H), 6.65 (s, 1H), 6.48 (dd, *J* = 2.34, 8.20 Hz, 1H), 3.86 (s, 2H).


GENERAL PROCEDURE B**Compound 1****N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide**

5 A mixture of Intermediate 1 (300 mg, 1.26 mmol) and 1-benzofuran-2-sulfonyl chloride (273 mg, 1.26 mmol) in pyridine (4 ml) was heated at 100 °C overnight. Pyridine was removed under reduced pressure and the residue was purified by column chromatography on silica gel (10% MeOH in CH₂Cl₂) to give **Compound 1** (157 mg, 30%).

10 ¹H NMR (600 MHz, acetone-d₆) δ 7.94 (s, 1H), 7.73 (d, *J* = 7.92 Hz, 1H), 7.64 (d, *J* = 2.35 Hz, 1H), 7.56 (d, *J* = 8.51 Hz, 1H), 7.52 (d, *J* = 8.22 Hz, 1H), 7.44 - 7.49 (m, 1H), 7.42 (s, 1H), 7.28 - 7.37 (m, 1H), 7.11 (dd, *J* = 2.35, 8.22 Hz, 1H), 6.97 (s, 1H), 3.91 (s, 2H).

Compound 2

15 **N-[5-chloro-2-(methylthio)phenyl]-1-benzofuran-2-sulfonamide**




Following **General Procedure B**, the title compound (243 mg, 60%) was prepared from 5-chloro-2-(methylthio)aniline (CAS 16423-54-4) (200 mg, 1.15 mmol) and 1-benzofuran-2-sulfonyl chloride (249 mg, 1.152 mmol).

20 ¹H NMR (600 MHz, acetone-d₆) δ 7.75 (d, *J* = 7.92 Hz, 1H), 7.60 (d, *J* = 8.51 Hz, 1H), 7.48 - 7.54 (m, 2H), 7.47 (d, *J* = 2.35 Hz, 1H), 7.29 - 7.38 (m, 2H), 7.22 (dd, *J* = 2.35, 8.51 Hz, 1H), 2.22 (s, 3H).

GENERAL PROCEDURE C**Compound 3**

25 **N-[5-chloro-2-(methylsulfinyl)phenyl]-1-benzofuran-2-sulfonamide**



To a solution of **Compound 2** (194 mg, 0.554 mmol) in CH₂Cl₂ (6 ml) at 0 °C was added *m*CPBA (111 mg, ~0.554 mmol). After it was stirred for 30 min at 0 °C, the mixture was separated into two portions. One portion (2 ml) was concentrated *in vacuo* and purified by column chromatography on silica gel (0→100% ethyl acetate in hexane followed by 0→10% MeOH in CH₂Cl₂) to give the title compound as a solid (35 mg, 52%).

Alternatively, the title compound can be prepared by treating **Compound 2** with 1 equivalent of NaIO₄ in MeOH/CH₃CN and H₂O at 0 °C to room temperature.


In another alternative procedure, the title compound can be prepared by treating **Compound 2** with 1-1.4 equivalent of Oxone[®] in MeOH/CH₃CN and H₂O at room temperature.

¹H NMR (600 MHz, CDCl₃) δ 7.74 (d, *J* = 1.76 Hz, 1H), 7.71 (d, *J* = 8.22 Hz, 1H), 7.52 - 7.59 (m, 2H), 7.49 (td, *J* = 1.17, 7.78 Hz, 1H), 7.32 - 7.39 (m, 1H), 7.25 (d, *J* = 8.22 Hz, 1H), 7.15 (dd, *J* = 1.91, 8.36 Hz, 1H), 2.88 (s, 3H).

GENERAL PROCEDURE D

Compound 4

N-[5-chloro-2-(methylsulfonyl)phenyl]-1-benzofuran-2-sulfonamide



To the other portion (4 ml) of solution from **Compound 3** was added *m*CPBA (111 mg, 0.554 mmol) and the reaction was stirred at room temperature for 1 h. The mixture was concentrated *in vacuo* and the residue was purified by column chromatography on silica gel (0→100% ethyl acetate in hexane) to give the title compound as a solid (75 mg, 53%).


Alternatively, the title compound can be prepared by treating **Compound 2** with 2-3 equivalents of *m*CPBA at room temperature.

In another alternative procedure, the title compound can be prepared by treating **Compound 2** with 3 equivalents of Oxone[®] in MeOH/CH₃CN and H₂O at room temperature.

¹H NMR (600 MHz, CDCl₃) δ 9.43 (s, 1H), 7.88 (d, *J* = 1.76 Hz, 1H), 7.81 (d, *J* = 8.51 Hz, 1H), 7.72 (d, *J* = 7.92 Hz, 1H), 7.61 (s, 1H), 7.47 - 7.57 (m, 2H), 7.34 - 7.40 (m, 1H), 7.26 (s, 1H), 7.23 (dd, *J* = 2.05, 8.51 Hz, 1H), 3.05 (s, 3H).

Intermediate 2

5-chloro-2-(methylsulfinyl)aniline




Following **General Procedure C**, the title compound (914 mg, 84%) was prepared from 5-chloro-2-(methylthio)aniline (1 g, 5.758 mmol).

¹H NMR (300 MHz, CD₃OD) δ 7.32 (d, *J* = 8.20 Hz, 1H), 6.81 (d, *J* = 2.05 Hz, 1H), 6.72 (dd, *J* = 1.76, 8.20 Hz, 1H), 2.86 (s, 3H).

Compound 5

3,4-dichloro-N-[5-chloro-2-(methylthio)phenyl]benzenesulfonamide




Following **General Procedure B**, the title compound (52 mg, 21%) was prepared from 5-chloro-2-(methylsulfinyl)aniline (**Intermediate 2**) (120 mg, 0.635 mmol) and 3,4-dichlorobenzene-1-sulfonyl chloride (156 mg, 0.635 mmol).

¹H NMR (300 MHz, CD₃OD) δ 7.87 (d, *J* = 1.76 Hz, 1H), 7.55 - 7.70 (m, 2H), 7.38 (s, 1H), 7.17 - 7.27 (m, 2H), 2.22 (s, 3H).

Compound 6

3,4-dichloro-N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide




Following **General Procedure C**, the title compound (28 mg, 52%) was prepared from 3,4-dichloro-N-[5-chloro-2-(methylthio)phenyl]benzenesulfonamide (52 mg, 0.136 mmol).

- 5 ^1H NMR (300 MHz, acetone- d_6) δ 10.70 (br. s., 1H), 8.06 (s, 1H), 7.86 (s, 2H), 7.44 - 7.59 (m, 2H), 7.34 (dd, J = 1.90, 8.35 Hz, 1H), 2.84 (s, 3H).

Compound 7

4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-3-methylbenzenesulfonamide




- 10 Following **General Procedure B and C**, the title compound (90 mg) was prepared from 5-chloro-2-(methylthio)aniline and 4-chloro-3-methylbenzene-1-sulfonyl chloride.

- ^1H NMR (600 MHz, CDCl_3) δ 10.62 (s, 1H), 7.82 (d, J = 2.05 Hz, 1H), 7.71 (dd, J = 1.91, 8.36 Hz, 1H), 7.63 (d, J = 2.05 Hz, 1H), 7.48 (d, J = 8.51 Hz, 1H), 7.02 - 7.15 (m, 2H), 2.81 (s, 3H), 2.43 (s, 3H).
- 15

Compound 8

N-[5-chloro-2-(methylsulfinyl)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide




- 20 Following **General Procedure B and C**, the title compound (43 mg) was prepared from 5-chloro-2-(methylthio)aniline and 4-nitro-3-(trifluoromethyl)benzene-1-sulfonyl chloride.

¹H NMR (600 MHz, acetone-d₆) δ 8.25 - 8.35 (m, 2H), 8.11 (d, *J* = 8.22 Hz, 1H), 7.39 - 7.48 (m, 2H), 6.75 (dd, *J* = 2.05, 8.22 Hz, 1H), 2.70 (s, 3H).

Compound 9

4-chloro-N-[5-chloro-2-(methylthio)phenyl]-2-fluorobenzenesulfonamide



5


Following **General Procedure B**, the title compound (380 mg, 66%) was prepared from 5-chloro-2-(methylthio)aniline (273 mg, 1.57 mmol) and 4-chloro-2-fluorobenzene-1-sulfonyl chloride (360 mg, 1.57 mmol).

¹H NMR (300 MHz, acetone-d₆) δ 8.77 (br. s., 1H), 7.84 (t, *J* = 8.06 Hz, 1H), 7.53 (dd, *J* = 1.90, 9.82 Hz, 1H), 7.35 - 7.48 (m, 3H), 7.25 (dd, *J* = 2.20, 8.35 Hz, 1H), 2.34 (s, 3H).

10

Compound 10

4-chloro-N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide




15

Following **General Procedure B and D**, the title compound (95 mg, 36%) was prepared from 5-chloro-2-[(1H-imidazol-4-ylmethyl)thio]aniline (301 mg, 1.26 mmol) and 4-chloro-3-(trifluoromethyl)benzene-1-sulfonyl chloride (351 mg, 1.26 mmol).


¹H NMR (600 MHz, acetone-d₆) δ 8.27 (s, 1H), 8.15 (d, *J* = 8.22 Hz, 1H), 7.81 (br. s., 1H), 7.73 (d, *J* = 8.51 Hz, 1H), 7.68 (d, *J* = 8.51 Hz, 1H), 7.30 (s, 1H), 6.91 (dd, *J* = 1.47, 8.51 Hz, 1H), 4.56 (br. s., 2H).

20

Intermediate 3**tert-butyl [4-(chloromethyl)-1,3-thiazol-2-yl]carbamate**

A solution of 4-(chloromethyl)-1,3-thiazol-2-amine hydrochloride (530 mg, 2.86 mmol),
 5 di-tert-butyl dicarbonate (750 mg, 3.44 mmol), triethylamine (0.6 ml, 4.30 mmol) and
 DMAP (cat. amount) in THF (10 ml) was stirred at room temperature overnight. The
 mixture was diluted with ethyl acetate, washed with water and brine, dried over
 Na₂SO₄, and concentrated in *vacuo*. The residue was purified by column
 chromatography (30% ethyl acetate in hexane) to give the title compound as a solid
 10 (391 mg, 45%).


¹H NMR (600 MHz, CD₃OD) δ 7.02 (s, 1H), 4.56 (s, 2H), 1.54 (s, 9H).

Intermediate 4**tert-butyl (4-[(2-amino-4-chlorophenyl)thio]methyl)-1,3-thiazol-2-yl)carbamate**

15 Following **General Procedure A**, the title compound (588 mg, 67%) was
 prepared from 2-amino-4-chlorobenzenethiol (376 mg, 2.36 mmol), tert-butyl [4-
 (chloromethyl)-1,3-thiazol-2-yl]carbamate (391 mg, 1.57 mmol) and K₂CO₃ (1.08 g,
 3.45 mmol) in DMF (10 ml).

¹H NMR (300 MHz, acetone-d₆) δ 10.18 (br. s., 1H), 7.17 (d, *J* = 8.21 Hz, 1H), 6.79 (d,
 20 *J* = 2.34 Hz, 1H), 6.62 (s, 1H), 6.51 (dd, *J* = 2.05, 8.20 Hz, 1H), 5.30 (br. s., 1H), 3.88
 (s, 2H), 1.53 (s, 9H).

Compound 11**tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio]methyl}-1,3-thiazol-2-yl}carbamate**




Following **General Procedure B**, the title compound (380 mg, 54%) was prepared from tert-butyl (4-[[2-amino-4-chlorophenyl]thio]methyl)-1,3-thiazol-2-yl)carbamate (640 mg, 1.72 mmol) and 1-benzofuran-2-sulfonyl chloride (372 mg, 1.72 mmol) in pyridine (5 ml).

¹H NMR (300 MHz, acetone-d₆) δ 11.20 (br. s., 1H), 9.82 (br. s., 1H), 7.78 (d, *J* = 7.91 Hz, 2H), 7.67 (d, *J* = 2.34 Hz, 1H), 7.62 (s, 1H), 7.45 - 7.60 (m, 3H), 7.31 - 7.43 (m, 1H), 7.07 - 7.22 (m, 1H), 6.77 (s, 1H), 3.93 (s, 2H), 1.55 (s, 9H).

GENERAL PROCEDURE E

Compound 12

N-(2-[[2-amino-1,3-thiazol-4-yl]methyl]thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide




A solution of tert-butyl {4-[[2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl]thio]methyl]-1,3-thiazol-2-yl}carbamate **Compound 11** (62 mg, 0.11 mmol), TFA (0.2 ml) in CH₂Cl₂ (1 ml) was stirred overnight. The solvent was removed in *vacuo* and the crude product was purified by column chromatography on silica gel (50% ethyl acetate in hexane) to afford the title compound (45 mg, 88%).

¹H NMR (600 MHz, acetone-d₆) δ 8.78 (br. s., 1H), 7.77 (d, *J* = 7.92 Hz, 0H), 7.55 - 7.66 (m, 3H), 7.45 - 7.55 (m, 2H), 7.29 - 7.41 (m, 1H), 7.17 (dd, *J* = 2.20, 8.36 Hz, 1H), 6.33 (s, 1H), 3.90 (s, 2H).

Compound 13

4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide



Following **General Procedure B and C**, the title compound was prepared from 5-chloro-2-(methylthio)aniline and 4-chloro-3-(trifluoromethyl)benzene-1-sulfonyl chloride.


- 5 ¹H NMR (300 MHz, CD₃OD) δ 8.07 (s, 1H), 7.96 (d, *J* = 8.50 Hz, 1H), 7.82 (d, *J* = 8.21 Hz, 1H), 7.76 (d, *J* = 8.50 Hz, 1H), 7.41 (d, *J* = 7.91 Hz, 1H), 7.07 (s, 1H), 2.83 (s, 3H).

GENERAL PROCEDURE F

Intermediate 5

N,N'-[dithiobis(5-chloro-2,1-phenylene)]bis[4-chloro-3-(trifluoromethyl)benzenesulfonamide]

10



To a solution of 2,2'-dithiobis(5-chloroaniline) (CAS 29124-55-8)(1.59 g, 5.0 mmol) in pyridine (20 ml) was added 4-chloro-3-(trifluoromethyl)benzene-1-sulfonyl chloride (2.76 g, 10.0 mmol) and the reaction was stirred at room temperature for 16 h. Additional 4-chloro-3-(trifluoromethyl)benzene-1-sulfonyl chloride (2.76 g, 10.0 mmol) and pyridine (20 ml) was added, and the reaction was stirred for 20 h. The mixture was concentrated in vacuo and H₂O was added. A gum-like semi-solid was formed. After decanting H₂O, the semi-solid was rinsed with H₂O (×2), taken in EtOAc, extracted with 1M HCl (×2), washed with H₂O, brine, dried over Na₂SO₄, and concentrated in vacuo to give a brown thick syrup (6.5 g). To a solution of the crude syrup in methanol (100 ml) was added 4M NaOH (12 ml), and the mixture was heated at 100 °C for 15 min, cooled to room temperature, quenched slowly with 1M HCl (~ 50 ml) with stirring and cooling to pH 4-5. The volume of the resulting suspension was

15


20

reduced in vacuo, followed by extraction with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄, and concentrated in vacuo. The crude solid containing sulfonic acid was pulverized in saturated NaHCO₃, and filtered, rinsed with minimal amount of diethyl ether to yield the title compound as an off-white solid (3.11 g).

- 5 ¹H NMR (600 MHz, CD₃OD) δ 8.08 (d, *J* = 2.05 Hz, 2H), 7.93 (dd, *J* = 2.20, 8.36 Hz, 2H), 7.81 (d, *J* = 8.51 Hz, 2H), 7.22 (d, *J* = 2.05 Hz, 2H), 7.18 - 7.20 (m, 2H), 7.10 (d, *J* = 8.51 Hz, 2H).

Compound 14

4-chloro-N-[5-chloro-2-(ethylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide



10

To a solution of N,N'-[dithiobis(5-chloro-2,1-phenylene)]bis[4-chloro-3-(trifluoromethyl)benzenesulfonamide] **Intermediate 5** (0.32 g, 0.40 mmol) in CH₂Cl₂ (10 ml) was added saturated aqueous NaHCO₃ (1.0 ml), polymer-bound triphenylphosphine (~3 mmol/g triphenylphosphine loading, 0.27 g, 0.80 mmol), and ethyl iodide (64 μl, 0.80 mmol). The reaction was stirred at room temperature for 2 h and was diluted with EtOAc, filtered, and washed with brine, dried over Na₂SO₄, and concentrated. The residue was purified by column chromatography on silica gel (0→25% ethyl acetate in hexane) to yield the title compound as a yellow syrup (204 mg, 59%).


15

- 20 ¹H NMR (600 MHz, CDCl₃) δ 8.12 (d, *J* = 2.35 Hz, 1H), 7.88 (dd, *J* = 2.35, 8.22 Hz, 1H), 7.82 (br. s., 1H), 7.64 (d, *J* = 2.35 Hz, 1H), 7.59 (d, *J* = 8.51 Hz, 1H), 7.32 (d, *J* = 8.22 Hz, 1H), 7.06 (dd, *J* = 2.20, 8.36 Hz, 1H), 2.60 (q, *J* = 7.34 Hz, 2H), 1.10 (t, *J* = 7.34 Hz, 3H).

Compound 15

4-chloro-N-[5-chloro-2-(ethylsulfinyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide

25




Following **General Procedure C**, the title compound was prepared from 4-chloro-N-[5-chloro-2-(ethylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide.

¹H NMR (600 MHz, CD₃OD) δ 8.15 (d, *J* = 2.35 Hz, 1H), 8.00 (dd, *J* = 2.20, 8.36 Hz, 1H), 7.72 (d, *J* = 8.51 Hz, 1H), 7.44 (d, *J* = 8.51 Hz, 1H), 7.27 (d, *J* = 2.05 Hz, 1H), 7.01 (dd, *J* = 2.05, 8.51 Hz, 1H), 3.10 - 3.18 (m, 1H), 2.89 - 2.97 (m, 1H), 1.11 (t, *J* = 7.34 Hz, 3H).

Intermediate 6


ethyl 6-[(tert-butoxycarbonyl)amino]pyridine-2-carboxylate



A mixture of ethyl 6-aminopyridine-2-carboxylate (1.57 g, 9.43 mmol), DMAP (1.12 g, 9.18 mmol), and di-tert-butyl dicarbonate (2.46 g, 11.3 mmol) in THF (45 ml) was heated at 60 °C for 16 h. The solvent was removed and the residue was purified by chromatography on silica gel (10→15% ethyl acetate in hexane) to yield the title compound as a white solid (2.50 g, 100%).

Intermediate 7

tert-butyl [6-(chloromethyl)pyridin-2-yl]carbamate




To a solution of ethyl 6-[(tert-butoxycarbonyl)amino]pyridine-2-carboxylate (456 mg, 1.71 mmol) in anhydrous ethanol (20 ml) was added pulverized CaCl₂ (395 mg, 3.42 mmol). The suspension was stirred and cooled to 0 °C and NaBH₄ (325 mg, 8.55 mmol) was added slowly. The reaction was stirred at 0 °C for 2 h, quenched with water, and extracted with CHCl₃ (×3). The combined organic layer was dried over Na₂SO₄ and concentrated. The residue was purified by column chromatography on silica gel (30→50% ethyl acetate in hexane) to yield a colorless syrup (308 mg) as a

mixture containing tert-butyl [6-(hydroxymethyl)pyridin-2-yl]carbamate as the major component (~4:1 ratio). To a solution of the above mixture in CH₂Cl₂ (10 ml) was added pyridine (144 μ l, 1.79 mmol) and SOCl₂ (120 μ l, 1.65 mmol). The reaction was stirred at room temperature for 4 h, quenched with water and saturated Na₂CO₃, and
 5 extracted with CHCl₃. The combined organic layer was dried over Na₂SO₄ and concentrated. The residue was purified by column chromatography on silica gel (10% ethyl acetate in hexane) to yield the title compound as a colorless syrup (208 mg, 50% over 2 steps).

Intermediate 8

10 **N,N'-[dithiobis(5-chloro-2,1-phenylene)]bis(1-benzofuran-2-sulfonamide)**




Following **General Procedure F**, the title compound was prepared from 2,2'-dithiobis(5-chloroaniline) and 1-benzofuran-2-sulfonyl chloride.

¹H NMR (600 MHz, CD₃OD) δ 7.72 (d, *J* = 7.92 Hz, 2H), 7.55 (dd, *J* = 0.59, 8.51 Hz, 2H), 7.48 (s, 2H), 7.32 - 7.37 (m, 4H), 7.22 (d, *J* = 2.05 Hz, 2H), 6.99 - 7.02 (m, 2H),
 15 6.96 - 6.99 (m, 2H).

GENERAL PROCEDURE G

Compound 16

20 **tert-Tutyl {6-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl]pyridin-2-yl}carbamate**




To a solution of N,N'-[dithiobis(5-chloro-2,1-phenylene)]bis(1-benzofuran-2-sulfonamide) (271 mg, 0.40 mmol) in CH₂Cl₂ (4 ml) and dioxane (4 ml) was added

saturated aqueous NaHCO_3 (4 ml), polymer-bound triphenylphosphine (~3 mmol/g triphenylphosphine loading, 0.40 g, 1.20 mmol), tert-butyl [6-(chloromethyl)pyridin-2-yl]carbamate (195 mg, 0.80 mmol), and tetrabutylammonium iodide (30 mg, 0.08 mmol). The reaction was stirred at room temperature for 4 h and was diluted with EtOAc, filtered, and washed with brine, dried over Na_2SO_4 , and concentrated. The residue was purified by column chromatography on silica gel (10→15% ethyl acetate in hexane) to yield the title compound as a yellow solid (260 mg, 59%).

^1H NMR (600 MHz, CD_3OD) δ 7.74 (d, J = 8.22 Hz, 1H), 7.70 (d, J = 7.63 Hz, 1H), 7.50 - 7.57 (m, 2H), 7.44 - 7.50 (m, 2H), 7.41 (s, 1H), 7.31 - 7.36 (m, 2H), 7.09 (d, J = 8.22 Hz, 1H), 6.65 (d, J = 7.34 Hz, 1H), 3.88 (s, 2H), 1.54 (s, 9H).

Compound 17

N-(2-[[6-(aminopyridin-2-yl)methyl]thio]-5-chlorophenyl)-1-benzofuran-2-sulfonamide




Following **General Procedure E**, the title compound (33 mg, 78%) was prepared from tert-butyl {6-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl]thio)methyl]pyridin-2-yl}carbamate.

^1H NMR (600 MHz, CD_3OD) δ 7.63 (d, J = 7.92 Hz, 1H), 7.58 (d, J = 2.35 Hz, 1H), 7.40 - 7.44 (m, 1H), 7.35 - 7.40 (m, 2H), 7.31 - 7.35 (m, 2H), 7.26 (ddd, J = 0.88, 7.04, 7.92 Hz, 1H), 6.95 (dd, J = 2.35, 8.22 Hz, 1H), 6.55 (d, J = 8.51 Hz, 1H), 6.31 (d, J = 7.04 Hz, 1H), 3.78 (s, 2H).

Compound 18

tert-butyl {6-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfinyl)methyl]pyridin-2-yl}carbamate




Following **General Procedure C**, the title compound (98 mg, 91%) was prepared from tert-butyl {6-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl]thio)methyl]pyridin-2-yl}carbamate.

¹H NMR (600 MHz, CD₃OD) δ 7.67 (d, *J* = 8.22 Hz, 1H), 7.64 (d, *J* = 7.04 Hz, 1H), 7.47 (d, *J* = 2.05 Hz, 1H), 7.42 - 7.46 (m, 1H), 7.36 - 7.39 (m, 1H), 7.29 - 7.33 (m, 1H), 7.22 - 7.26 (m, 2H), 7.09 (d, *J* = 8.22 Hz, 1H), 6.77 (dd, *J* = 1.91, 8.36 Hz, 1H), 6.68 (d, *J* = 7.34 Hz, 1H), 4.59 (d, *J* = 12.62 Hz, 1H), 4.16 (d, *J* = 12.62 Hz, 1H), 1.52 (s, 9H).

Compound 19

10 **N-(2-[(6-aminopyridin-2-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide**




Following **General Procedure E**, the title compound (71 mg, 91%) was prepared from tert-butyl {6-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfinyl)methyl]pyridin-2-yl}carbamate.

15 ¹H NMR (300 MHz, CD₃OD) δ 7.65 (d, *J* = 7.91 Hz, 1H), 7.52 - 7.62 (m, 1H), 7.50 (d, *J* = 1.47 Hz, 1H), 7.39 - 7.46 (m, 1H), 7.29 - 7.38 (m, 2H), 7.20 - 7.29 (m, 2H), 6.89 (dd, *J* = 1.47, 8.50 Hz, 1H), 6.79 (d, *J* = 8.79 Hz, 1H), 6.44 (d, *J* = 7.03 Hz, 1H), 4.60 (d, *J* = 13.48 Hz, 1H), 4.25 (d, *J* = 13.19 Hz, 1H).

Compound 20

20 **tert-butyl {6-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]pyridin-2-yl}carbamate**




Following **General Procedure D**, the title compound (60 mg, 56%) was prepared from tert-butyl {6-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl]thio)methyl]pyridin-2-yl}carbamate.

¹H NMR (300 MHz, CD₃OD) δ 7.79 (d, *J* = 1.76 Hz, 1H), 7.65 (d, *J* = 7.62 Hz, 1H), 7.55 (d, *J* = 8.50 Hz, 1H), 7.44 (s, 1H), 7.18 - 7.40 (m, 5H), 6.72 (dd, *J* = 1.76, 8.50 Hz, 1H), 6.60 (d, *J* = 7.33 Hz, 1H), 4.97 (s, 2H), 1.51 (s, 9H).

Compound 21

N-(2-[(6-aminopyridin-2-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide




Following **General Procedure E**, the title compound (47 mg, 94%) was prepared from tert-butyl {6-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]pyridin-2-yl}carbamate.

¹H NMR (300 MHz, CD₃OD) δ 7.73 (d, *J* = 2.05 Hz, 1H), 7.66 (d, *J* = 7.62 Hz, 1H), 7.52 (d, *J* = 8.79 Hz, 1H), 7.20 - 7.44 (m, 4H), 7.14 (t, *J* = 7.77 Hz, 1H), 6.73 (dd, *J* = 1.76, 8.50 Hz, 1H), 6.48 (d, *J* = 8.20 Hz, 1H), 6.28 (d, *J* = 7.33 Hz, 1H), 4.91 (s, 2H).

Compound 22

tert-butyl 3-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl]-1H-pyrazole-1-carboxylate




Following **General Procedure G**, the title compound (206 mg, 36%) was prepared from N,N'-[dithiobis(5-chloro-2,1-phenylene)]bis(1-benzofuran-2-sulfonamide) and tert-butyl 3-(bromomethyl)-1H-pyrazole-1-carboxylate.

¹H NMR (600 MHz, CDCl₃) δ 8.30 (br. s., 1H), 7.93 (d, *J* = 2.64 Hz, 1H), 7.62 - 7.67 (m, 2H), 7.47 - 7.52 (m, 1H), 7.42 - 7.45 (m, 1H), 7.41 (s, 1H), 7.34 (d, *J* = 8.22 Hz,

1H), 7.28 - 7.32 (m, 1H), 7.00 (dd, $J = 2.20, 8.36$ Hz, 1H), 6.03 (d, $J = 2.64$ Hz, 1H), 3.88 (s, 2H), 1.64 (s, 9H).

Compound 23

N-{5-chloro-2-[(1H-pyrazol-3-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide



5

Following **General Procedure E**, the title compound (15 mg, 66%) was prepared from tert-butyl 3-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl]thio)methyl]-1H-pyrazole-1-carboxylate.


^1H NMR (600 MHz, CD_3OD) δ 7.70 (d, $J = 7.92$ Hz, 1H), 7.51 - 7.55 (m, 1H), 7.45 - 7.50 (m, 2H), 7.37 - 7.44 (m, 2H), 7.34 (t, $J = 7.19$ Hz, 1H), 7.25 - 7.31 (m, 1H), 7.11 (dd, $J = 1.76, 8.22$ Hz, 1H), 5.88 (br. s., 1H), 3.89 (s, 2H).

10

GENERAL PROCEDURE H

Intermediate 9

3-[(2-amino-4-chlorophenyl)thio]-N,N-dimethylpropanamide



15

A mixture of 2-amino-4-chlorobenzenethiol (327 mg, 2.05 mmol), N,N-dimethylacrylamide (203 mg, 2.05 mmol) and HOAc (0.5 ml) in CH_2Cl_2 (5 ml) was stirred at room temperature for 3 days. The reaction was quenched with NaHCO_3 (aq.) and then extracted with CH_2Cl_2 (2×10 ml). The organic layer was washed with water and brine and concentrated *in vacuo*. The crude product was purified by chromatography on silica gel (50% ethyl acetate in hexane) to yield the title compound (284 mg, 54%).


20

^1H NMR (300 MHz, CD_3OD) δ 7.26 (d, $J = 8.21$ Hz, 1H), 6.76 (d, $J = 2.05$ Hz, 1H), 6.56 (dd, $J = 2.34, 8.20$ Hz, 1H), 2.85 - 3.02 (m, 8H), 2.57 (t, $J = 7.03$ Hz, 2H).

25

Compound 24

3-[[4-Chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl]-N,N-dimethylpropanamide



5

Following **General Procedure B and C**, the title compound (280 mg) was prepared from 3-[(2-amino-4-chlorophenyl)thio]-N,N-dimethylpropanamide (284 mg, 1.11 mmol).


¹H NMR (600 MHz, CD₃OD) δ 8.16 (d, *J* = 1.76 Hz, 1H), 8.00 (dd, *J* = 2.05, 8.22 Hz, 1H), 7.75 (d, *J* = 8.51 Hz, 1H), 7.57 (d, *J* = 8.51 Hz, 1H), 7.32 (s, 1H), 7.21 (d, *J* = 7.92 Hz, 1H), 3.26 (br. s., 1H), 3.18 (br. s., 1H), 3.02 (s, 3H), 2.95 (s, 3H), 2.83 - 2.91 (m, 1H), 2.62 - 2.76 (m, 1H).

10

Compound 25

tert-Butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl]sulfinyl)methyl]-1,3-thiazol-2-yl}carbamate

15



Following **General Procedure C**, tert-Butyl (4-(((2-(benzofuran-2-sulfonamido)-4-chlorophenyl)thio)methyl)thiazol-2-yl)carbamate (138mg, 0.25 mmol) was oxidized to the title compound (97 mg, 68%).


¹H NMR (600 MHz, acetone-d₆) δ 10.43 (br. s., 1H), 7.75 - 7.84 (m, 2H), 7.66 (d, *J* = 2.05 Hz, 1H), 7.62 (dd, *J* = 0.88, 8.51 Hz, 1H), 7.50 (ddd, *J* = 1.32, 7.26, 8.44 Hz, 1H), 7.32 - 7.41 (m, 1H), 7.11 - 7.24 (m, 2H), 6.74 (s, 1H), 4.47 (d, *J* = 12.91 Hz, 1H), 4.37 (d, *J* = 12.91 Hz, 1H), 1.50 (s, 9H).

20

Compound 26

N-(2-[(2-amino-1,3-thiazol-4-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide

25




Following **General Procedure E**, tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]-1,3-thiazol-2-yl}carbamate (87 mg, 0.153 mmol) was de-Boc to give the title compound (58 mg, 82%).

- 5 ¹H NMR (600 MHz, acetone-d₆) δ 10.43 (br. s., 1H), 7.75 - 7.84 (m, 2H), 7.66 (d, J = 2.05 Hz, 1H), 7.62 (dd, J = 0.88, 8.51 Hz, 1H), 7.50 (ddd, J = 1.32, 7.26, 8.44 Hz, 1H), 7.32 - 7.41 (m, 1H), 7.11 - 7.24 (m, 2H), 6.74 (s, 1H), 4.47 (d, J = 12.91 Hz, 1H), 4.37 (d, J = 12.91 Hz, 1H), 1.50 (s, 9H).

Compound 27

- 10 **tert-Butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]-1,3-thiazol-2-yl}carbamate**




Following **General Procedure D**, the title compound (112 mg, 90%) was prepared from tert-Butyl (4-(((2-(benzofuran-2-sulfonamido)-4-chlorophenyl)thio)methyl)thiazol-2-yl)carbamate (117mg, 0.212 mmol).

- 15 ¹H NMR (600 MHz, acetone-d₆) δ 7.80 (d, J = 7.92 Hz, 2H), 7.78 (d, J = 1.76 Hz, 1H), 7.67 (d, J = 8.51 Hz, 1H), 7.59 (d, J = 7.92 Hz, 1H), 7.50 (t, J = 7.63 Hz, 1H), 7.37 (t, J = 7.48 Hz, 1H), 6.99 (s, 1H), 4.70 (s, 2H), 1.53 (s, 9H)
- 20 ¹H NMR (600 MHz, acetone-d₆) δ 7.80 (d, J = 7.92 Hz, 2H), 7.67 (d, J = 8.51 Hz, 1H), 7.60 (br. s., 1H), 7.50 (t, J = 7.63 Hz, 1H), 7.33 - 7.41 (m, 1H), 6.99 (br. s., 1H), 4.70 (br. s., 2H), 1.53 (s, 9H).

Compound 28

N-(2-[(2-amino-1,3-thiazol-4-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide




Following **General Procedure E**, the title compound (54 mg, 86%) was prepared from tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfinyl)methyl]-1,3-thiazol-2-yl}carbamate (77 mg, 0.153 mmol).

- 5 ¹H NMR (600 MHz, acetone-d₆) δ 7.74 - 7.85 (m, 2H), 7.68 (d, J = 2.35 Hz, 1H), 7.64 (dd, J = 0.88, 8.51 Hz, 1H), 7.53 (ddd, J = 1.32, 7.19, 8.51 Hz, 1H), 7.39 (ddd, J = 0.88, 7.26, 8.00 Hz, 1H), 7.16 - 7.24 (m, 2H), 6.74 (s, 1H), 4.32 - 4.46 (m, 2H), 1.47 - 1.55 (m, 9H).

Intermediate 10

10 5-chloro-2-(((5-nitro-1H-pyrazol-3-yl)methyl)thio)aniline




- (5-Nitro-1H-pyrazol-3-yl)methanol (524 mg, 3.66 mmol) was first treated with SOCl₂ (3 ml) in CH₂Cl₂ (5 ml) at 35 °C for 2 hrs. The solvent was removed to get a crude 3-(chloromethyl)-5-nitro-1H-pyrazole. Then following **General Procedure A**, the title compound (515 mg, 49%) was prepared from 2-amino-4-chlorobenzenethiol (161 mg, 1.01 mmol), crude 3-(chloromethyl)-5-nitro-1H-pyrazole, K₂CO₃ (468 mg, 3.39 mmol) in DMF (3 ml).

¹H NMR (600 MHz, CD₃OD) δ 7.05 (d, J = 8.22 Hz, 1H), 6.77 (d, J = 2.35 Hz, 1H), 6.60 (s, 1H), 6.49 (dd, J = 2.05, 8.22 Hz, 1H), 3.95 (s, 2H).

20

Compound 29

N-(5-chloro-2-(((5-nitro-1H-pyrazol-3-yl)methyl)thio)phenyl)-1-benzofuran-2-sulfonamide




Following **General Procedure B**, the title compound (580 mg, 71%) was prepared from 5-chloro-2-(((5-nitro-1H-pyrazol-3-yl)methyl)thio)aniline (503 mg, 1.77 mmol) and benzofuran-2-sulfonyl chloride (384 mg, 1.77 mmol) in pyridine (5 ml).

- 5 ¹H NMR (600 MHz, CD₃OD) δ 7.70 - 7.73 (m, 1H), 7.53 - 7.56 (m, 1H), 7.49 (d, J = 1.47 Hz, 1H), 7.47 - 7.49 (m, 1H), 7.47 (d, J = 1.17 Hz, 1H), 7.45 (d, J = 2.35 Hz, 1H), 7.44 (d, J = 0.88 Hz, 1H), 7.35 (ddd, J = 0.88, 7.26, 8.00 Hz, 1H), 7.15 (dd, J = 2.35, 8.22 Hz, 1H), 6.42 (s, 1H), 3.97 (s, 2H).

Compound 30

- 10 **N-(5-chloro-2-(((5-nitro-1H-pyrazol-3-yl)methyl)sulfinyl)phenyl)-1-benzofuran-2-sulfonamide**




Following **General Procedure C**, the title compound (155 mg, 61%) was prepared from N-(5-chloro-2-(((5-nitro-1H-pyrazol-3-yl)methyl)thio)phenyl)-1-benzofuran-2-

- 15 sulfonamide (245 mg, 0.528 mmol).
¹H NMR (600 MHz, CD₃OD) δ 7.67 (dt, J = 0.77, 7.56 Hz, 1H), 7.42 - 7.45 (m, 2H), 7.35 (td, J = 1.32, 7.85 Hz, 1H), 7.24 - 7.29 (m, 2H), 7.09 (d, J = 8.22 Hz, 1H), 6.85 (d, J = 8.22 Hz, 1H), 6.33 (s, 1H), 4.61 (d, J = 14.09 Hz, 1H), 4.48 (d, J = 14.09 Hz, 1H).

GENERAL PROCEDURE I

- 20 **Compound 31**

N-(2-(((5-amino-1H-pyrazol-3-yl)methyl)thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide




N-(5-chloro-2-((5-nitro-1H-pyrazol-3-yl)methyl)thio)phenyl)-1-benzofuran-2-sulfonamide (77 mg, 0.166 mmol) was reduced to the title compound under H₂ balloon in the presence of Pd/C (10%wt, 18 mg) in MeOH (3 ml).

- 5 ¹H NMR (600 MHz, CD₃OD) δ 7.70 (d, J = 7.92 Hz, 1H), 7.52 - 7.56 (m, 2H), 7.48 (ddd, J = 1.17, 7.12, 8.44 Hz, 1H), 7.44 (s, 1H), 7.40 (s, 1H), 7.34 (t, J = 7.48 Hz, 1H), 7.28 (d, J = 8.51 Hz, 1H), 7.11 (dd, J = 2.20, 8.36 Hz, 1H), 3.74 (s, 2H).

Compound 32

N-(2-((5-amino-1H-pyrazol-3-yl)methyl)sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide

10




Following the **General Procedure I**, the title compound (39 mg, 91%) was reduced from N-(5-chloro-2-((5-nitro-1H-pyrazol-3-yl)methyl)sulfinyl)phenyl)-1-benzofuran-2-sulfonamide (46 mg, 0.096 mmol).

- 15 ¹H NMR (600 MHz, CD₃OD) δ 7.91 - 7.99 (m, 1H), 7.67 - 7.72 (m, 1H), 7.56 - 7.62 (m, 1H), 7.50 - 7.55 (m, 1H), 7.41 - 7.48 (m, 2H), 7.29 - 7.40 (m, 2H), 7.13 (dd, J = 1.91, 8.36 Hz, 1H), 4.55 (d, J = 13.79 Hz, 1H), 4.05 (d, J = 13.79 Hz, 1H)

Compound 33

N-(2-((5-amino-1H-pyrazol-3-yl)methyl)sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide

20




Following **General Procedure D** and **H**, the title compound (27 mg) was prepared from N-(5-chloro-2-[[[(5-nitro-1H-pyrazol-3-yl)methyl]thio}phenyl)-1-benzofuran-2-sulfonamide.

- 5 ¹H NMR (600 MHz, CD₃OD) δ 7.69 - 7.74 (m, 2H), 7.62 - 7.58 (m, 3H), 7.51 (dd, J = 0.59, 8.51 Hz, 1H), 7.44 (ddd, J = 1.17, 7.12, 8.44 Hz, 1H), 7.26 - 7.35 (m, 1H), 7.09 (dd, J = 1.91, 8.66 Hz, 1H), 4.59 (s, 2H).

Intermediate 11

1-propyl-1H-imidazole-4-carbaldehyde



10

To a suspension of NaH (95%, 231 mg, 9.18 mmol) in THF (20 ml) was added 4-imidazole carboxaldehyde (588 mg, 6.12 mmol) under ice cooling. The mixture was refluxed for 2 hours under nitrogen atmosphere. The mixture was cooled to room temperature and 1-iodopropane (5 ml) was added to the mixture and then refluxed for another 2 hours. The mixture was cooled down to rt and quenched with water. The solution was extracted with EtOAc (2 x 50 ml), washed with brine, dried over Na₂SO₄, concentrated in *vacuo*. The crude product was purified by column chromatography on silica gel (0-10% MeOH in CH₂Cl₂) to give the title compound (487 mg, 58%).


15

¹H NMR (600 MHz, CD₃OD) δ 9.73 (s, 1H), 7.97 (s, 1H), 7.84 (s, 1H), 4.07 (t, J = 7.04 Hz, 2H), 1.73 - 1.92 (m, 2H), 0.93 (t, J = 7.34 Hz, 3H).

20

Intermediate 12

(1-propyl-1H-imidazol-4-yl)methanol




To a solution of LAH (2.0 M in THF, 1.9 ml, 3.882 mmol) in THF (10 ml) was added dropwise a solution of 1-propyl-1H-imidazole-4-carbaldehyde (487 mg, 3.529 mmol) in THF (4 ml) at 0 °C. After stirred at 0 °C for 5 min, water (0.2 ml), 15% NaOH (0.2 ml) was added dropwise to the reaction mixture under 0 °C. The mixture was further stirred at room temperature for 2 hours. MgSO₄ was added to the mixture and the solid was filtered away and the filtrate was concentrated in *vacuo* to get a yellow oil and used without further purification.

¹H NMR (600 MHz, CD₃Cl₃) δ 7.42 (s, 1H), 6.86 (s, 1H), 4.59 (s, 2H), 3.87 (t, *J* = 7.04 Hz, 2H), 1.73 - 1.82 (m, 2H), 0.93 (t, *J* = 7.34 Hz, 3H).

Intermediate 13

5-chloro-2-(((1-propyl-1H-imidazol-4-yl)methyl)thio)aniline




(1-propyl-1H-imidazol-4-yl)methanol (535 mg, 3.11 mmol) was first treated with SOCl₂ (3 ml) in CH₂Cl₂ (5 ml) at 35 °C for 2 hrs. The solvent was removed to get a crude 4-(chloromethyl)-1-propyl-1H-imidazole. Then following **General Procedure A**, the title compound (587 mg, 67%) was prepared from 2-amino-4-chlorobenzenethiol (744 mg, 4.66 mmol), 4-(chloromethyl)-1-propyl-1H-imidazole, K₂CO₃ (2.1g, 15.54 mmol) in DMF (10 ml).

¹H NMR (600 MHz, CDCl₃) δ 7.38 (s, 1H), 7.17 (d, *J* = 8.22 Hz, 1H), 6.67 (d, *J* = 2.05 Hz, 1H), 6.57 (dd, *J* = 2.20, 8.07 Hz, 1H), 6.51 (s, 1H), 3.85 (s, 2H), 3.79 (t, *J* = 7.04 Hz, 2H), 1.67 - 1.76 (m, 2H), 0.88 (t, *J* = 7.34 Hz, 3H).

Compound 34

N-(5-chloro-2-(((1-propyl-1H-imidazol-4-yl)methyl)thio)phenyl)-1-benzofuran-2-sulfonamide




Following **General Procedure B**, the title compound (463 mg, 48%) was prepared from 5-chloro-2-(((1-propyl-1H-imidazol-4-yl)methyl)thio)aniline (587 mg, 2.082 mmol) and benzofuran-2-sulfonyl chloride (451 mg, 2.082 mmol) in pyridine (10 ml).

5 ¹H NMR (600 MHz, CD₃OD) δ 7.71 (dt, J = 1.03, 7.92 Hz, 1H), 7.57 (d, J = 1.17 Hz, 1H), 7.52 - 7.55 (m, 1H), 7.44 - 7.51 (m, 2H), 7.42 (d, J = 0.88 Hz, 1H), 7.34 (td, J = 1.03, 7.56 Hz, 1H), 7.27 (d, J = 8.22 Hz, 1H), 7.07 (dd, J = 2.35, 8.51 Hz, 1H), 6.44 - 6.51 (m, 1H), 3.75 - 3.81 (m, 4H), 1.63 (dquin, J = 7.19, 7.34 Hz, 2H), 0.74 - 0.82 (m, 3H).

10

Compound 35

N-(5-chloro-2-(((1-propyl-1H-imidazol-4-yl)methyl)sulfinyl)phenyl)-1-benzofuran-2-sulfonamide




Following **General Procedure C**, the title compound (148 mg, 75%) was prepared from N-(5-chloro-2-(((1-propyl-1H-imidazol-4-yl)methyl)thio)phenyl)-1-benzofuran-2-sulfonamide.

15

¹H NMR (600 MHz, CD₃OD) δ 8.46 (s, 1H), 7.70 (dd, J = 1.47, 8.22 Hz, 1H), 7.49 (dd, J = 0.88, 8.51 Hz, 1H), 7.38 - 7.43 (m, 2H), 7.35 (d, J = 0.88 Hz, 1H), 7.30 (ddd, J = 1.03, 7.19, 7.92 Hz, 1H), 7.12 (d, J = 8.51 Hz, 1H), 6.91 (dd, J = 1.91, 8.36 Hz, 1H),
20 6.80 (s, 1H), 4.57 (d, J = 14.38 Hz, 1H), 4.43 (d, J = 14.09 Hz, 1H), 3.84 - 3.99 (m, 2H), 1.57 - 1.71 (m, 2H), 0.81 (t, J = 7.34 Hz, 3H).

Compound 36

N-(5-chloro-2-(((1-propyl-1H-imidazol-4-yl)methyl)sulfonyl)phenyl)-1-benzofuran-2-sulfonamide




25

Following **General Procedure D**, the title compound (115 mg, 60%) was prepared from N-(5-chloro-2-[[[(1-propyl-1H-imidazol-4-yl)methyl]thio]phenyl]-1-benzofuran-2-sulfonamide.

¹H NMR (300 MHz, CD₃OD) δ 7.82 (s, 1H), 7.71 (dd, J = 1.32, 7.77 Hz, 1H), 7.67 (d, J = 2.05 Hz, 1H), 7.55 (d, J = 8.50 Hz, 1H), 7.44 - 7.50 (m, 2H), 7.41 (td, J = 1.32, 7.69 Hz, 1H), 7.25 - 7.35 (m, 1H), 6.85 (dd, J = 2.05, 8.79 Hz, 1H), 6.76 (s, 1H), 4.57 (s, 2H), 3.74 (t, J = 7.03 Hz, 2H), 1.54 (quint, J = 6.89, 7.07 Hz, 2H), 0.69 (t, J = 7.47 Hz, 3H).

Intermediate 14

10 **tert-butyl (4-(hydroxymethyl)pyridin-2-yl)carbamate**




A solution of (2-aminopyridin-4-yl)methanol (547 mg, 4.408 mmol), di-tert-butyl dicarbonate (1.25 g, 5.730 mmol), in *t*-BuOH (20 ml) was stirred at room temperature overnight. The solvent was removed and added ethyl acetate, filtered away the solid and the filtrate was concentrated in *vacuo*. The residue was purified by column chromatography (10% MeOH in CH₂Cl₂) to give the title compound as a white solid (703 mg, 71%).

¹H NMR (600 MHz, CDCl₃) δ 8.18 (d, J = 5.28 Hz, 1H), 8.00 (s, 1H), 6.92 - 7.10 (m, 1H), 4.75 (s, 2H), 1.32 - 1.66 (m, 9H).

20 **Intermediate 15**

tert-butyl (4-(((2-amino-4-chlorophenyl)thio)methyl)pyridin-2-yl)carbamate




tert-butyl (4-(hydroxymethyl)pyridin-2-yl)carbamate (600 mg, 2.679 mmol) was first treated with SOCl₂ (3 ml) in CH₂Cl₂ (5 ml) at rt for 2 hrs. The solvent was removed to get a crude tert-butyl (4-(chloromethyl)pyridin-2-yl)carbamate. Then following **General Procedure A**, the title compound (667 mg, 68%) was prepared from 2-amino-4-chlorobenzenethiol (641 mg, 4.019 mmol), K₂CO₃ (1.8g, 13.39 mmol) in DMF (20 ml).

¹H NMR (600 MHz, CDCl₃) δ 9.83 (br. s., 1H), 8.20 (d, *J* = 4.70 Hz, 1H), 7.85 (s, 1H), 6.92 - 7.18 (m, 1H), 6.60 - 6.75 (m, 2H), 6.55 (dd, *J* = 2.05, 8.22 Hz, 1H), 4.42 (br. s., 2H), 3.78 (s, 2H), 1.54 (s, 9H).

Compound 37

5 **tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl]thio)methyl}pyridin-2-yl}carbamate**



Following **General Procedure B**, the title compound (555 mg, 56%) was prepared from tert-butyl (4-(((2-amino-4-chlorophenyl)thio)methyl)pyridin-2-


10 yl)carbamate (667 mg, 1.827 mmol) and benzofuran-2-sulfonyl chloride (396 mg, 1.827 mmol) in pyridine (10 ml).

¹H NMR (600 MHz, CD₃OD) δ 7.99 (d, *J* = 4.99 Hz, 1H), 7.64 - 7.73 (m, 2H), 7.59 (d, *J* = 1.17 Hz, 1H), 7.50 - 7.55 (m, 1H), 7.42 - 7.50 (m, 2H), 7.30 - 7.39 (m, 1H), 7.15 (dd, *J* = 1.03, 8.36 Hz, 1H), 6.96 - 7.05 (m, 1H), 6.56 (d, *J* = 5.28 Hz, 1H), 3.77 (s, 2H),

15 1.48 - 1.61 (m, 9H).

Compound 38

N-(2-[(2-aminopyridin-4-yl)methyl]thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide




20 Following **General Procedure E**, the title compound (76 mg, 100%) was prepared from tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl]thio)methyl}pyridin-2-yl}carbamate (96 mg, 0.176 mmol).

¹H NMR (600 MHz, CD₃OD) δ 7.72 - 7.76 (m, 2H), 7.65 (d, *J* = 6.46 Hz, 2H), 7.58 (dd, *J* = 0.73, 8.36 Hz, 1H), 7.46 - 7.53 (m, 2H), 7.42 (d, *J* = 2.05 Hz, 1H), 7.37 (td, *J* = 1.03, 7.56 Hz, 1H), 7.31 (d, *J* = 8.51 Hz, 1H), 7.16 (dd, *J* = 2.35, 8.51 Hz, 1H), 6.64 (dd, *J* = 1.61, 6.60 Hz, 1H), 6.50 (s, 1H), 3.90 (s, 2H).

Compound 39

tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]pyridin-2-yl}carbamate




Following **General Procedure D**, the title compound (49 mg) was prepared from tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl]pyridin-2-yl}carbamate.

¹H NMR (600 MHz, CD₃OD) δ 7.93 (d, J = 6.75 Hz, 1H), 7.78 (s, 1H), 7.73 (d, J = 2.05 Hz, 1H), 7.63 (d, J = 7.63 Hz, 1H), 7.41 (d, J = 8.80 Hz, 1H), 7.33 (s, 1H), 7.29 (d, J = 3.81 Hz, 1H), 7.23 (dt, J = 3.96, 7.92 Hz, 2H), 6.85 (dd, J = 2.20, 6.60 Hz, 1H), 6.68 (d, J = 7.92 Hz, 1H), 4.56 (br. s., 2H), 1.47 (s, 9H).

Compound 40

tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]-1-oxidopyridin-2-yl}carbamate




Following **General Procedure D**, the title compound (82 mg) was prepared from tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl]pyridin-2-yl}carbamate.

¹H NMR (600 MHz, CD₃OD) δ 8.22 (d, J = 6.46 Hz, 1H), 7.96 (s, 1H), 7.76 (d, J = 16.73 Hz, 1H), 7.60 - 7.68 (m, 1H), 7.43 (d, J = 8.51 Hz, 1H), 7.37 (s, 1H), 7.31 (br. s., 1H), 7.21 - 7.27 (m, 1H), 7.09 (d, J = 6.46 Hz, 1H), 6.85 (d, J = 6.75 Hz, 1H), 6.72 (d, J = 7.63 Hz, 1H), 4.60 (br. s., 2H), 1.55 (s, 9H).

Compound 41

N-(2-[(2-aminopyridin-4-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide




Following **General Procedure E**, the title compound (32 mg, 100%) was prepared from tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]pyridin-2-yl}carbamate.

- 5 ¹H NMR (600 MHz, acetone-d₆) δ 7.74 - 7.85 (m, 3H), 7.61 - 7.70 (m, 2H), 7.52 - 7.58 (m, 1H), 7.45 (td, J = 1.17, 7.92 Hz, 1H), 7.28 - 7.38 (m, 1H), 7.09 (dd, J = 2.05, 8.51 Hz, 1H), 6.88 (s, 1H), 6.57 - 6.66 (m, 1H), 4.89 (s, 2H).

Compound 42

N-(2-[(2-amino-1-oxidopyridin-4-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide

10




Following **General Procedure E**, the title compound (43 mg, 67%) was prepared from tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]-1-oxidopyridin-2-yl}carbamate (77 mg, 0.130 mmol).

- 15 ¹H NMR (600 MHz, acetone-d₆) δ 8.07 (d, J = 4.70 Hz, 3H), 7.76 - 7.81 (m, 2H), 7.74 (s, 1H), 7.64 (d, J = 8.51 Hz, 1H), 7.58 (d, J = 8.51 Hz, 1H), 7.46 - 7.49 (m, 1H), 7.32 - 7.37 (m, 1H), 7.15 (dd, J = 1.32, 8.36 Hz, 1H), 6.94 (br. s., 1H), 6.52 (d, J = 4.99 Hz, 1H), 4.76 (s, 2H).

Compound 43

N-(2-[(2-aminopyridin-4-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide

20




Following **General Procedure C** and **General Procedure E**, the title compound (103 mg) was prepared from tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl]pyridin-2-yl}carbamate.

¹H NMR (600 MHz, acetone-d₆) δ 8.29 (br. s., 1H), 7.74 (dd, J = 7.04, 18.19 Hz, 2H),
5 7.40 - 7.61 (m, 4H), 7.25 - 7.37 (m, 2H), 7.05 (d, J = 7.04 Hz, 1H), 6.84 (s, 1H), 6.49
(d, J = 6.16 Hz, 1H), 4.50 (d, J = 12.62 Hz, 2H), 4.40 (d, J = 12.62 Hz, 1H).

Intermediate 16

3-((2-amino-4-chlorophenyl)thio)-N,N-dimethylpropanamide




10 The solution of 2-amino-4-chlorobenzenethiol (824 mg, 5.126 mmol), N,N-dimethylacrylamide (512 mg, 5.162 mmol), and HOAc (1 ml) in CH₂Cl₂ (10 ml) was stirred at room temperature for 4 days. The solvent was removed and the residue was loaded on silica column and purified to yield a white solid (1.08 g, 83%).

¹H NMR (600 MHz, CD₃OD) δ 7.26 (d, J = 8.22 Hz, 1H), 6.76 (d, J = 2.05 Hz, 1H),
15 6.56 (dd, J = 2.05, 8.22 Hz, 1H), 2.93 - 2.97 (m, 5H), 2.90 (s, 3H), 2.58 (t, J = 7.04 Hz, 2H).

Compound 44

3-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfonyl]-N,N-dimethylpropanamide




Following **General Procedure B** and **D**, the title compound (73mg) was prepared from 3-((2-amino-4-chlorophenyl)thio)-N,N-dimethylpropanamide (104 mg, 0.405 mmol) and 4-chloro-3-(trifluoromethyl)benzene-1-sulfonyl chloride (113 mg, 0.405 mmol).

¹H NMR (600 MHz, CD₃OD) δ 8.31 (d, J = 1.47 Hz, 1H), 8.12 (dd, J = 1.76, 8.51 Hz, 1H), 7.70 (dd, J = 8.51, 18.49 Hz, 2H), 7.58 (s, 1H), 6.90 (s., 1H), 3.73 (br. s., 2H), 3.01 (s, 3H), 2.95 (s, 3H), 2.78 (t, J = 6.75 Hz, 2H).

Compound 45

5 **3-[(4-chloro-2-[(2,4-difluorophenyl)sulfonyl]amino}phenyl)thio]-N,N-dimethylpropanamide**




Following **General Procedure B**, the title compound (364 mg, 72%) was prepared from 3-((2-amino-4-chlorophenyl)thio)-N,N-dimethylpropanamide (300 mg, 1.167 mmol) and 2,4-difluorobenzene-1-sulfonyl chloride (248 mg, 1.167 mmol) in pyridine (5 ml).

¹H NMR (600 MHz, CD₃OD) δ 7.86 (td, J = 6.02, 8.44 Hz, 1H), 7.38 - 7.45 (m, 2H), 7.13 - 7.21 (m, 2H), 7.09 (tdd, J = 0.88, 2.49, 8.44 Hz, 1H), 2.95 - 3.01 (m, 5H), 2.93 (s, 3H), 2.56 (t, J = 7.04 Hz, 2H).

Compound 46

15 **3-[(4-chloro-2-[(2,4-difluorophenyl)sulfonyl]amino}phenyl)sulfinyl]-N,N-dimethylpropanamide**




Following **General Procedure C**, the title compound (130 mg, 88%) was prepared from 3-[(4-chloro-2-[(2,4-difluorophenyl)sulfonyl]amino}phenyl)thio]-N,N-dimethylpropanamide.

¹H NMR (600 MHz, CD₃OD) δ 7.90 (td, J = 5.87, 8.51 Hz, 1H), 7.69 (d, J = 8.51 Hz, 1H), 7.43 (dd, J = 2.05, 8.51 Hz, 1H), 7.31 (d, J = 2.05 Hz, 1H), 7.20 - 7.27 (m, 1H),

7.11 - 7.17 (m, 1H), 3.24 - 3.29 (m, 1H), 3.12 - 3.20 (m, 1H), 3.05(s, 3H), 2.93 (s, 3H), 2.85 - 2.92 (m, J = 4.40 Hz, 1H), 2.72 - 2.80 (m, J = 6.46, 6.46, 17.02 Hz, 1H).

Compound 47

3-[(4-chloro-2-[(2,4-difluorophenyl)sulfonyl]amino}phenyl)sulfonyl]-N,N-dimethylpropanamide




Following **General Procedure D**, the title compound (70 mg, 71%) was prepared from 3-[(4-chloro-2-[(2,4-difluorophenyl)sulfonyl]amino}phenyl)thio]-N,N-dimethylpropanamide.

¹H NMR (600 MHz, CD₃OD) δ 8.01 - 8.12 (m, 1H), 7.76 (d, J = 8.51 Hz, 2H), 7.49 - 7.60 (m, 2H), 6.95 - 7.24 (m, 3H), 3.72 (br. s., 2H), 3.01 (s, 3H), 2.90 (s, 3H), 2.78 (t, J = 7.04 Hz, 2H).

Compound 48

3-[(4-chloro-2-[(4-chloro-2-fluorophenyl)sulfonyl]amino}phenyl)thio]-N,N-dimethylpropanamide




Following **General Procedure B**, the title compound (476 mg, 75%) was prepared from 3-((2-amino-4-chlorophenyl)thio)-N,N-dimethylpropanamide (364 mg, 1.42 mmol) and 4-chloro-2-fluorobenzene-1-sulfonyl chloride (325 mg, 1.42 mmol) in pyridine (5 ml).

¹H NMR (600 MHz, acetone-d₆) δ 9.57 (br. s., 1H), 7.86 (t, J = 8.07 Hz, 1H), 7.50 - 7.56 (m, 2H), 7.45 (dd, J = 1.91, 9.83 Hz, 1H), 7.40 (dd, J = 1.91, 8.36 Hz, 1H), 7.17 (dd, J = 2.35, 8.22 Hz, 1H), 2.94 - 3.01 (m, 5H), 2.91 (s, 3H), 2.54 (t, J = 6.46 Hz, 3H).

Compound 49

3-[(4-chloro-2-[(4-chloro-2-fluorophenyl)sulfonyl]amino}phenyl)sulfinyl]-N,N-dimethylpropanamide




5 Following **General Procedure C**, the title compound (90 mg, 55%) was prepared from 3-[(4-chloro-2-[(4-chloro-2-fluorophenyl)sulfonyl]amino}phenyl)thio]-N,N-dimethylpropanamide.

1H NMR (600 MHz, CD₃OD) δ 7.78 - 7.90 (m, 1H), 7.48 (d, J = 8.51 Hz, 1H), 7.27 - 7.34 (m, 2H), 7.26 (d, J = 2.05 Hz, 1H), 7.03 (d, J = 7.92 Hz, 1H), 3.43 (ddd, J = 6.31, 9.02, 13.43 Hz, 1H), 3.16 (ddd, J = 5.58, 9.10, 13.50 Hz, 1H), 2.99 (s, 3H), 2.92 (s, 3H), 2.79 - 2.88 (m, J = 6.46, 9.17, 16.07 Hz, 1H), 2.52 - 2.64 (m, 1H).

Compound 50

3-[(4-chloro-2-[(4-chloro-2-fluorophenyl)sulfonyl]amino}phenyl)sulfonyl]-N,N-dimethylpropanamide




15 Following **General Procedure D**, the title compound (81 mg, 48%) was prepared from 3-[(4-chloro-2-[(4-chloro-2-fluorophenyl)sulfonyl]amino}phenyl)thio]-N,N-dimethylpropanamide.

1H NMR (600 MHz, CD₃OD) δ 7.89 - 7.98 (m, 1H), 7.69 (d, J = 8.51 Hz, 1H), 7.46 (d, J = 2.05 Hz, 1H), 7.20 - 7.30 (m, 2H), 6.76 (d, J = 7.63 Hz, 1H), 3.86 (t, J = 7.19 Hz, 2H), 2.97 (s, 3H), 2.93 (s, 3H), 2.70 (t, J = 7.34 Hz, 2H).

Intermediate 17

tert-butyl (5-(hydroxymethyl)-4H-1,2,4-triazol-3-yl)carbamate



(5-amino-4H-1,2,4-triazol-3-yl)methanol (467 mg, 4.093 mmol), di-tert-butyl dicarbonate (1.16 g, 5.32 mmol), in *t*-BuOH (20 ml) was stirred at room temperature overnight. The solvent was removed and added ethyl acetate, filtered away the solid
 5 and the filtrate was concentrated in *vacuo*. The residue was purified by column chromatography (10% MeOH in CH₂Cl₂) to give the title compound as a yellow oil (190 mg, 22%).


¹H NMR (300 MHz, CD₃OD) δ 4.44 (s, 2H), 1.63 (s, 9H).

GENERAL PROCEDURE J

10

Intermediate 18

tert-butyl (5-(((2-amino-4-chlorophenyl)thio)methyl)-4H-1,2,4-triazol-3-yl)carbamate




A solution of tert-butyl (5-(hydroxymethyl)-4H-1,2,4-triazol-3-yl)carbamate (190
 15 mg, 0.888 mmol), 2-amino-4-chlorobenzenethiol (213 mg, 1.332 mmol), PPh₃ (466 mg, 1.776 mmol) and di-tert-butylazodicarboxylate (409 mg, 1.776 mmol) in CH₂Cl₂ (10 ml) was stirred at room temperature for 2 days. The solvent was then removed and the crude residues was loaded on silica gel column to get the title compound.

¹H NMR (600 MHz, CD₃OD) δ 7.18 (s, 1H), 6.73 (d, *J* = 2.35 Hz, 1H), 6.49 (dd, *J* =
 20 2.35, 8.22 Hz, 2H), 3.68 (s, 2H), 1.60 (s, 9H).

Intermediate 19

tert-butyl (5-(((2-(benzofuran-2-sulfonamido)-4-chlorophenyl)thio)methyl)-4H-1,2,4-triazol-3-yl)carbamate




Following **General Procedure B**, the title compound was prepared from tert-butyl (5-(((2-amino-4-chlorophenyl)thio)methyl)-4H-1,2,4-triazol-3-yl)carbamate (330 mg, 0.925 mmol) and benzofuran-2-sulfonyl chloride (200 mg, 0.925 mmol) in pyridine (2 ml).

¹H NMR (600 MHz, CD₃OD) δ 7.69 (dd, *J* = 0.88, 7.92 Hz, 1H), 7.45 - 7.57 (m, 4H), 7.40 (s, 1H), 7.29 - 7.35 (m, 1H), 7.16 (dd, *J* = 2.35, 8.22 Hz, 1H), 3.70 (s, 2H), 1.61 (s, 9H).

Compound 51

N-(2-(((5-amino-4H-1,2,4-triazol-3-yl)methyl)thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide




Following **General Procedure E**, the title compound (13 mg) was prepared from tert-butyl (5-(((2-(benzofuran-2-sulfonamido)-4-chlorophenyl)thio)methyl)-4H-1,2,4-triazol-3-yl)carbamate.

¹H NMR (600 MHz, CD₃OD) δ 7.71 (d, *J* = 7.92 Hz, 1H), 7.55 (d, *J* = 8.51 Hz, 1H), 7.39 - 7.51 (m, 5H), 7.31 - 7.37 (m, 1H), 7.16 (d, *J* = 7.04 Hz, 1H), 3.82 (s, 2H).

Compound 52

N-(2-(((5-amino-4H-1,2,4-triazol-3-yl)methyl)sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide




Following **General Procedure C** and **E**, the title compound (15 mg) was prepared from tert-butyl (5-(((2-(benzofuran-2-sulfonamido)-4-chlorophenyl)thio)methyl)-4H-1,2,4-triazol-3-yl)carbamate.

- 5 ¹H NMR (600 MHz, CD₃OD) δ 7.75 (dd, J = 1.17, 7.92 Hz, 1H), 7.57 - 7.64 (m, 2H), 7.51 - 7.55 (m, 1H), 7.50 (d, J = 0.88 Hz, 1H), 7.44 (dd, J = 1.91, 8.36 Hz, 1H), 7.35 - 7.41 (m, 1H), 7.20 (d, J = 2.05 Hz, 1H), 4.45 (d, J = 14.09 Hz, 1H), 4.24 (d, J = 14.09 Hz, 1H).

Intermediate 20

10

5-chloro-2-((2-(pyridin-2-yl)ethyl)thio)aniline




- 2-(Pyridin-2-yl)ethanol (1 g, 8.12 mmol) was first treated with SOCl₂ (3 ml) in CH₂Cl₂ (10 ml) at rt for 3 hrs. The solvent was removed to get a crude 2-(2-chloroethyl)pyridine. Then following **General Procedure A**, the title compound (1.69 g, 79%) was prepared from 2-amino-4-chlorobenzenethiol (1.6 g, 9.74 mmol), K₂CO₃ (3.36 g, 24.36 mmol) in DMF (20 ml).

- 15 ¹H NMR (600 MHz, CD₃OD) δ 8.39 - 8.42 (m, 1H), 7.73 (td, J = 1.76, 7.63 Hz, 1H), 7.28 (d, J = 7.63 Hz, 1H), 7.23 (d, J = 8.22 Hz, 2H), 6.75 (d, J = 2.35 Hz, 1H), 6.55 (dd, J = 2.35, 8.22 Hz, 1H), 3.06 - 3.10 (m, 2H), 2.96 - 3.01 (m, 2H).

20

Compound 53

N-{5-chloro-2-[(2-pyridin-2-ylethyl)thio]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure B**, the title compound was prepared from 5-chloro-2-((2-(pyridin-2-yl)ethyl)thio)aniline (439 mg, 1.663 mmol) and benzofuran-2-sulfonyl chloride (2 x 359 mg, 2x 1.663 mmol) in pyridine (5 ml).

¹H NMR (600 MHz, CD₃OD) δ 8.54 - 8.62 (m, 1H), 7.85 (s, 1H), 7.71 - 7.78 (m, 1H), 7.67 (dt, J = 1.03, 7.92 Hz, 1H), 7.56 (d, J = 2.35 Hz, 1H), 7.39 - 7.48 (m, 2H), 7.36 (s, 1H), 7.27 - 7.32 (m, 2H), 7.15 - 7.22 (m, 2H), 2.98 (t, J = 7.04 Hz, 2H), 2.74 (t, J = 6.90 Hz, 2H).

Compound 54

N-{5-chloro-2-[(2-pyridin-2-ylethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure C**, the title compound (45 mg, 79%) was prepared from N-{5-chloro-2-[(2-pyridin-2-ylethyl)thio]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, CD₃OD) δ 8.52 - 8.60 (m, 1H), 7.81 (td, J = 1.76, 7.63 Hz, 1H), 7.64 - 7.71 (m, 2H), 7.26 - 7.47 (m, 8H), 3.40 - 3.50 (m, 1H), 3.29 - 3.37 (m, 1H), 3.16 - 3.26 (m, 0H), 2.83 - 2.95 (m, J = 7.48, 7.48, 14.97 Hz, 1H).

Compound 55

N-{5-chloro-2-[(2-pyridin-2-ylethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure D**, the title compound (20 mg) was prepared from N-{5-chloro-2-[(2-pyridin-2-ylethyl)thio]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, CD₃OD) δ 8.30 (ddd, J = 1.47, 1.61, 4.55 Hz, 1H), 7.71 - 7.76 (m, 2H), 7.61 - 7.68 (m, 2H), 7.52 (d, J = 0.88 Hz, 1H), 7.31 - 7.36 (m, 1H), 7.24 - 7.31 (m,

2H), 7.18 - 7.23 (m, 2H), 7.05 (dd, $J = 2.05, 8.51$ Hz, 1H), 3.84 - 3.89 (m, 2H), 3.00 - 3.14 (m, 2H).

Intermediate 21

2-((2-(1H-pyrazol-4-yl)ethyl)thio)-5-chloroaniline




5

Following the **General Procedure J**, the title intermediate was prepared from 2-(1H-pyrazol-4-yl)ethanol (472 mg, 4.218 mmol), 2-amino-4-chlorobenzenethiol (1.01 g, 6.327 mmol), PPh₃ (2.21 g, 8.436 mmol) and di-tert-butylazodicarboxylate (1.9 g, 8.436 mmol) in CH₂Cl₂ (20 ml).

10 ¹H NMR (600 MHz, CD₃OD) δ 7.98 (d, $J = 0.88$ Hz, 1H), 7.64 (s, 1H), 7.23 (d, $J = 8.22$ Hz, 1H), 6.76 (d, $J = 2.35$ Hz, 1H), 6.55 (dd, $J = 2.05, 8.22$ Hz, 1H), 2.89 - 3.00 (m, 2H), 2.65 - 2.77 (m, 2H).

Compound 56

N-(5-chloro-2-[[2-(1H-pyrazol-4-yl)ethyl]thio]phenyl)-1-benzofuran-2-sulfonamide




15

Following **General Procedure B**, the title compound (123 mg) was prepared from 2-((2-(1H-pyrazol-4-yl)ethyl)thio)-5-chloroaniline (157 mg, 0.618 mmol) and benzofuran-2-sulfonyl chloride (133 mg, 0.618 mmol) in pyridine (3 ml).

20 ¹H NMR (600 MHz, CD₃OD) δ 7.70 (d, $J = 7.92$ Hz, 1H), 7.43 - 7.53 (m, 3H), 7.39 (s, 1H), 7.28 - 7.36 (m, 4H), 7.17 (dd, $J = 1.76, 8.51$ Hz, 1H), 2.82 (t, $J = 7.63$ Hz, 2H), 2.45 (t, $J = 7.63$ Hz, 2H).

Compound 57

N-(5-chloro-2-[[2-(1H-pyrazol-4-yl)ethyl]sulfinyl]phenyl)-1-benzofuran-2-sulfonamide




Following **General Procedure C**, the title compound (52mg, 40%) was prepared from N-{5-chloro-2-[(2-pyridin-2-ylethyl)thio]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, CD₃OD) δ 7.75 - 7.81 (m, 1H), 7.72 (td, J = 1.76, 7.63 Hz, 1H),
 5 7.67 (d, J = 0.88 Hz, 1H), 7.48 - 7.59 (m, 3H), 7.40 - 7.45 (m, 2H), 7.19 - 7.34 (m, 3H),
 7.17 (d, J = 2.35 Hz, 1H), 3.19 - 3.26 (m, 3H), 2.87 (t, J = 7.63 Hz, 2H).

Compound 58

N-(5-chloro-2-[[2-(1H-pyrazol-4-yl)ethyl]sulfonyl]phenyl)-1-benzofuran-2-sulfonamide




10

Following **General Procedure D**, the title compound (75 mg, 73%) was prepared from N-{5-chloro-2-[(2-pyridin-2-ylethyl)thio]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, CD₃OD) δ 7.67 - 7.74 (m, 2H), 7.62 (d, J = 7.63 Hz, 1H), 7.39 (s,
 15 2H), 7.31 (t, J = 7.19 Hz, 2H), 7.22 - 7.26 (m, 1H), 7.17 (d, J = 8.22 Hz, 1H), 6.80 -
 6.90 (m, 1H), 3.89 (br. s., 2H), 2.71 - 2.81 (m, 2H).

Intermediate 22

5-Chloro-2-[2-(3,5-dimethyl-1H-pyrazol-4-yl)-ethylsulfanyl]-phenylamine



20

Following **General Procedure A**, the title compound (517 mg, 74%) was prepared from 2-Amino-4-chloro-benzenethiol (590 mg, 3.693 mmol) and 4-(2-bromo-


ethyl)-3,5-dimethyl-1H-pyrazole (500mg, 2.462 mmol), K₂CO₃(1.7 g, 12.31 mmol) in DMF (20 ml).

¹H NMR (600 MHz, CD₃OD) δ 7.24 (d, *J* = 8.22 Hz, 1H), 6.77 (d, *J* = 2.05 Hz, 1H), 6.56 (dd, *J* = 2.35, 8.22 Hz, 1H), 2.75 - 2.80 (m, 2H), 2.53 - 2.61 (m, 2H), 2.08 (s, 6H).

5

Compound 59

N-(5-chloro-2-[[2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl]thio]phenyl)-1-benzofuran-2-sulfonamide



Following **General Procedure B**, the title compound (235 mg) was prepared


10 from 2-((2-(1H-pyrazol-4-yl)ethyl)thio)-5-chloroaniline (157 mg, 0.618 mmol) and benzofuran-2-sulfonyl chloride (133 mg, 0.618 mmol) in pyridine (3 ml).

¹H NMR (600 MHz, CD₃OD) δ 7.68 (dd, *J* = 0.88, 7.92 Hz, 1H), 7.52 (d, *J* = 2.35 Hz, 1H), 7.39 - 7.47 (m, 3H), 7.27 - 7.35 (m, 2H), 7.18 (dd, *J* = 2.35, 8.51 Hz, 1H), 2.65 - 2.72 (m, *J* = 7.63 Hz, 2H), 2.28 - 2.35 (m, 2H), 1.97 (s, 6H).

15

Intermediate 23

5-chloro-2-(((2-fluoropyridin-3-yl)methyl)thio)aniline




(2-Fluoropyridin-3-yl)methanol (508 mg, 3.998 mmol) was first treated with SOCl₂ (1.5 ml) in CH₂Cl₂ (5 ml) at rt for 3 hrs. The solvent was removed to get a crude 3-

20 (chloromethyl)-2-fluoropyridine. Then following **General Procedure A**, the title compound (910 mg, 85%) was prepared from 2-amino-4-chlorobenzenethiol (957 mg, 5.995 mmol), K₂CO₃ (2.7 g, 19.98 mmol) in DMF (20 ml).


¹H NMR (600 MHz, CD₃OD) δ 8.01 (dt, *J* = 0.88, 4.99 Hz, 1H), 7.42 (ddd, *J* = 1.91, 7.48, 9.68 Hz, 1H), 7.10 (ddd, *J* = 1.76, 5.14, 7.19 Hz, 1H), 6.91 (d, *J* = 8.22 Hz, 1H), 6.73 (d, *J* = 2.05 Hz, 1H), 6.42 (dd, *J* = 2.35, 8.22 Hz, 1H), 3.90 (s, 2H).

25

Compound 60**N-(5-chloro-2-[[2-(2-fluoropyridin-3-yl)methyl]thio]phenyl)-1-benzofuran-2-sulfonamide**


5 Following **General Procedure B**, the title compound (235 mg) was prepared from 5-chloro-2-(((2-fluoropyridin-3-yl)methyl)thio)aniline (522 mg, 1.948 mmol) and benzofuran-2-sulfonyl chloride (421 mg, 1.948 mmol) in pyridine (5 ml).

¹H NMR (600 MHz, acetone-d₆) δ 9.00 (br. s., 1H), 8.05 (dt, J = 1.47, 4.70 Hz, 1H), 7.79 (d, J = 6.75 Hz, 1H), 7.55 - 7.63 (m, 3H), 7.43 - 7.54 (m, 2H), 7.33 - 7.40 (m, 1H),
 10 7.26 (d, J = 8.22 Hz, 1H), 7.07 - 7.17 (m, 2H), 4.01 (s, 2H).

Compound 61**N-(5-chloro-2-[[2-(2-fluoropyridin-3-yl)methyl]sulfinyl]phenyl)-1-benzofuran-2-sulfonamide**


15 Following **General Procedure C**, the title compound (100mg, 59%) was prepared from N-(5-chloro-2-[[2-(2-fluoropyridin-3-yl)methyl]thio]phenyl)-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, CD₃OD) δ 7.99 (dd, J = 1.91, 4.84 Hz, 1H), 7.65 (d, J = 7.34 Hz, 1H), 7.49 (d, J = 2.05 Hz, 1H), 7.38 (d, J = 8.51 Hz, 1H), 7.28 - 7.33 (m, 1H), 7.19 -
 20 7.26 (m, 3H), 7.00 (ddd, J = 1.47, 5.21, 7.12 Hz, 1H), 6.81 (d, J = 8.22 Hz, 1H), 6.70 (dd, J = 2.05, 8.22 Hz, 1H), 4.43 - 4.59 (m, 2H).

Compound 62**N-(5-chloro-2-[[2-(2-fluoropyridin-3-yl)methyl]sulfonyl]phenyl)-1-benzofuran-2-sulfonamide**


Following **General Procedure D**, the title compound (176mg, 67%) was prepared from N-(5-chloro-2-[[2-(2-fluoropyridin-3-yl)methyl]thio}phenyl)-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, acetone-d₆) δ 8.04 (d, J = 4.70 Hz, 1H), 7.91 (d, J = 2.05 Hz, 1H), 7.70 (d, J = 7.63 Hz, 1H), 7.18 - 7.44 (m, 6H), 6.90 (t, J = 5.43 Hz, 1H), 6.67 (d, J = 7.63 Hz, 1H), 4.96 (s, 2H).

Intermediate 24**2-(benzylthio)-5-chloroaniline**

Following **General Procedure A**, the title compound (819 mg, 100%) was prepared from 2-amino-4-chlorobenzenethiol (700 mg, 4.39 mmol), (bromomethyl)benzene (560 mg, 2.92 mmol), K₂CO₃ (2.0 g, 14.62 mmol) in DMF (20 ml).

¹H NMR (600 MHz, CD₃OD) δ 7.15 - 7.24 (m, 3H), 7.07 - 7.13 (m, 2H), 6.97 (d, J = 8.22 Hz, 1H), 6.73 (d, J = 2.05 Hz, 1H), 6.43 (dd, J = 2.05, 8.22 Hz, 1H), 3.86 (s, 2H).


Compound 63**N-[2-(benzylthio)-5-chlorophenyl]-1-benzofuran-2-sulfonamide**

Following **General Procedure B**, the title compound (235 mg) was prepared from 2-(benzylthio)-5-chloroaniline(509 mg, 2.052 mmol) and benzofuran-2-sulfonyl chloride (443 mg, 2.052 mmol) in pyridine (5 ml).

¹H NMR (600 MHz, CD₃OD) δ 7.73 (d, J = 7.92 Hz, 3H), 7.45 - 7.56 (m, 3H), 7.42 (s, 1H), 7.32 - 7.38 (m, 1H), 7.10 - 7.18 (m, 4H), 7.04 - 7.08 (m, 1H), 6.85 - 6.94 (m, 2H), 3.81 (s, 2H).

Compound 64


N-[2-(benzylsulfinyl)-5-chlorophenyl]-1-benzofuran-2-sulfonamide




Following **General Procedure C**, the title compound (139mg, 84%) was prepared from N-[2-(benzylthio)-5-chlorophenyl]-1-benzofuran-2-sulfonamide. ¹H NMR (600 MHz, CD₃OD) δ 7.65 (d, J = 7.92 Hz, 1H), 7.49 (d, J = 2.05 Hz, 1H), 7.37 (dd, J = 0.73, 8.36 Hz, 1H), 7.28 - 7.33 (m, 1H), 7.16 - 7.27 (m, 3H), 7.08 - 7.14 (m, 2H), 7.01 (d, J = 6.75 Hz, 2H), 6.97 (d, J = 8.51 Hz, 1H), 6.73 (dd, J = 1.91, 8.36 Hz, 1H), 4.50 (d, J = 12.91 Hz, 1H), 4.14 (d, J = 12.91 Hz, 1H).

Compound 65

N-[2-(benzylsulfonyl)-5-chlorophenyl]-1-benzofuran-2-sulfonamide



Following **General Procedure D**, the title compound (212mg, 95%) was prepared from N-[2-(benzylthio)-5-chlorophenyl]-1-benzofuran-2-sulfonamide. ¹H NMR (600 MHz, CD₃OD) δ 7.63 - 7.73 (m, 2H), 7.40 (dd, J = 0.73, 8.36 Hz, 1H), 7.30 - 7.37 (m, 3H), 7.23 - 7.29 (m, 1H), 7.08 - 7.16 (m, 3H), 6.98 - 7.05 (m, 2H), 6.60 (dd, J = 2.05, 8.51 Hz, 1H), 4.99 (s, 2H).


GENERAL PROCEDURE K**Intermediate 25****2-amino-4-fluorobenzenethiol**

5

A solution of 5-fluoro-2-methylbenzo[d]thiazole (1.2 g, 7.18 mmol) in ethylene glycol and NaOH (5N, 2 ml) was degassed under N₂ for 10 min, then refluxed at 129 °C for 3 hours. The solution was cooled to 0 °C and then acidified to pH 3 ~ 4 using c. HCl. The mixture was extracted with EtOAc (2 x 50 ml). The organic layer was washed with brine and dried with Na₂SO₄ and concentrated in vacuo. The crude residue (1.01 g, 98%) was used directly in the next step without purification.

10

¹H NMR (600 MHz, CD₃OD) δ 7.24 (dd, *J* = 6.16, 8.51 Hz, 1H), 6.97 (dd, *J* = 6.46, 8.51 Hz, 1H), 6.20 (td, *J* = 2.79, 8.58 Hz, 1H).


Intermediate 26**5-fluoro-2-((3-nitrobenzyl)thio)aniline**

15

Following **General Procedure A**, the title compound (460 mg, 97%) was prepared from 2-amino-4-fluorobenzenethiol (562 g, 3.93 mmol), 1-(bromomethyl)-3-nitrobenzene (566 mg, 2.62 mmol), K₂CO₃ (1.8 g, 13.10 mmol) in DMF (10 ml).

20

¹H NMR (600 MHz, CD₃OD) δ 8.04 (dt, *J* = 2.09, 7.26 Hz, 1H), 7.89 (s, 1H), 7.34 - 7.54 (m, 2H), 6.93 (dd, *J* = 6.46, 8.51 Hz, 1H), 6.35 - 6.53 (m, 1H), 6.15 (td, *J* = 2.64, 8.51 Hz, 1H), 3.95 (s, 2H).

Compound 66**N-(5-fluoro-2-((3-nitrobenzyl)thio)phenyl)benzofuran-2-sulfonamide**

25


Following **General Procedure B**, the title compound (474 mg, 63%) was prepared from 2-((3-aminobenzyl)thio)-5-fluoroaniline (460 mg, 1.655 mmol) and benzofuran-2-sulfonyl chloride (357 mg, 1.655 mmol) in pyridine (5 ml).

¹H NMR (600 MHz, CD₃OD) δ 7.98 (ddd, J = 0.88, 2.35, 8.22 Hz, 1H), 7.68 - 7.79 (m, 2H), 7.49 - 7.54 (m, 2H), 7.42 - 7.48 (m, 1H), 7.25 - 7.39 (m, 4H), 7.16 (dd, J = 6.16, 8.80 Hz, 1H), 6.78 (td, J = 2.93, 8.36 Hz, 1H), 3.91 (s, 2H).

GENERAL PROCEDURE L

Compound 67

N-{2-[(3-aminobenzyl)thio]-5-fluorophenyl}-1-benzofuran-2-sulfonamide




N-(5-fluoro-2-((3-nitrobenzyl)thio)phenyl)benzofuran-2-sulfonamide (91 mg, 0.199 mmol) was dissolved in MeOH (2 ml). Zn (322 mg, 4.967 mmol) and NH₄Cl (1 ml) was added to the solution. After the mixture was stirred for 30 min at room temperature, the solid was filtered and the filtrate was concentrated in vacuo and then the crude residue was purified by column chromatography (0 ~ 30 % EtOAc in hexane) to afford the title product (50 mg, 59%).

¹H NMR (600 MHz, CD₃OD) δ 7.71 (d, J = 7.92 Hz, 1H), 7.50 - 7.54 (m, 1H), 7.48 (d, J = 0.88 Hz, 1H), 7.46 (ddd, J = 1.32, 7.26, 8.44 Hz, 1H), 7.28 - 7.38 (m, 2H), 7.19 (dd, J = 6.16, 8.80 Hz, 1H), 6.83 - 6.89 (m, 1H), 6.77 (td, J = 2.79, 8.44 Hz, 1H), 6.46 - 6.54 (m, 1H), 6.36 (t, J = 1.76 Hz, 1H), 6.22 (d, J = 7.63 Hz, 1H), 3.63 (s, 2H).

Compound 68

N-{2-[(3-aminobenzyl)sulfinyl]-5-fluorophenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure L**, the title compound (130mg, 84%) was prepared from N-{5-fluoro-2-[(3-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, CD₃OD) δ 7.75 (dd, J = 0.88, 7.92 Hz, 1H), 7.53 - 7.60 (m, 2H), 7.48 (ddd, J = 1.32, 7.26, 8.44 Hz, 1H), 7.33 - 7.38 (m, 1H), 7.28 (dd, J = 6.16, 8.80 Hz, 1H), 7.14 (dd, J = 2.64, 10.27 Hz, 1H), 6.88 - 7.02 (m, 2H), 6.71 (dt, J = 1.06, 8.14 Hz, 1H), 6.57 (s, 1H), 6.41 (d, J = 7.63 Hz, 1H), 4.24 (d, J = 12.62 Hz, 1H), 4.06 (d, J = 12.91 Hz, 1H).

Compound 69

10 **N-{5-fluoro-2-[(3-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide**




Following **General Procedure D**, the title compound (136mg, 74%) was prepared from N-(5-fluoro-2-((3-nitrobenzyl)thio)phenyl)benzofuran-2-sulfonamide.

¹H NMR (600 MHz, CD₃OD) δ 8.10 (ddd, J = 1.17, 2.13, 8.14 Hz, 1H), 7.72 - 7.81 (m, 3H), 7.65 (dd, J = 6.02, 8.95 Hz, 1H), 7.55 (d, J = 8.51 Hz, 1H), 7.45 - 7.51 (m, 3H), 7.41 - 7.45 (m, 1H), 7.33 - 7.39 (m, 1H), 6.80 - 7.09 (m, 1H), 4.70 (s, 2H).

Compound 70

N-{2-[(3-aminobenzyl)sulfonyl]-5-fluorophenyl}-1-benzofuran-2-sulfonamide




20 Following **General Procedure L**, the title compound (91mg, 81%) was prepared from N-{5-fluoro-2-[(3-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, acetone-d₆) δ 7.87 (br. s., 1H), 7.84 (d, J = 7.92 Hz, 2H), 7.57 - 7.65 (m, 2H), 7.50 - 7.56 (m, 2H), 7.39 (t, J = 7.48 Hz, 1H), 6.94 - 7.02 (m, 1H), 6.83 (t, J = 7.78 Hz, 1H), 6.53 (d, J = 8.80 Hz, 1H), 6.38 (s, 1H), 6.13 (d, J = 7.34 Hz, 1H), 4.38 (s, 2H).

5

Intermediate 27

5-methoxy-2-((3-nitrobenzyl)thio)aniline




Following **General Procedure K** and **A**, the title compound (474 mg) was prepared from 5-methoxy-2-methylbenzo[d]thiazole.

10 ¹H NMR (600 MHz, CD₃OD) δ 8.03 (dt, J = 2.05, 7.34 Hz, 1H), 7.84 (d, J = 1.76 Hz, 1H), 7.34 - 7.48 (m, 2H), 6.83 (d, J = 8.51 Hz, 1H), 6.32 (s, 1H), 6.06 (dd, J = 2.79, 8.36 Hz, 1H), 3.90 (s, 2H), 3.69 (s, 3H).

Compound 71

N-{2-[(3-aminobenzyl)thio]-5-methoxyphenyl}-1-benzofuran-2-sulfonamide




15

Following **General Procedure L**, the title compound (82mg, 70%) was prepared from N-(5-methoxy-2-((3-nitrobenzyl)thio)phenyl)benzofuran-2-sulfonamide.

20 ¹H NMR (600 MHz, CD₃OD) δ 7.69 (d, J = 7.92 Hz, 1H), 7.50 (dd, J = 0.88, 8.51 Hz, 1H), 7.40 - 7.47 (m, 2H), 7.31 (ddd, J = 1.03, 7.19, 7.92 Hz, 1H), 7.07 - 7.13 (m, 2H), 6.84 - 6.90 (m, 1H), 6.57 (dd, J = 2.79, 8.66 Hz, 1H), 6.50 - 6.54 (m, 1H), 6.37 (t, J = 1.91 Hz, 1H), 6.25 (d, J = 7.63 Hz, 1H), 3.72 (s, 3H), 3.57 (s, 2H).

Compound 72

N-{5-methoxy-2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure C**, the title compound (95mg, 46%) was prepared from N-(5-methoxy-2-((3-nitrobenzyl)thio)phenyl)benzofuran-2-sulfonamide.

¹H NMR (600 MHz, acetone-d₆) δ 10.74 (br. s., 1H), 8.16 (ddd, J = 0.88, 2.35, 8.22 Hz, 1H), 7.78 - 7.86 (m, 2H), 7.75 (s, 1H), 7.61 (dd, J = 0.88, 8.51 Hz, 1H), 7.56 (t, J = 7.92 Hz, 1H), 7.44 - 7.52 (m, 2H), 7.35 - 7.40 (m, 1H), 7.09 (d, J = 2.35 Hz, 1H), 7.06 (d, J = 8.80 Hz, 1H), 6.69 (dd, J = 2.49, 8.66 Hz, 1H), 4.45 - 4.53 (m, 2H), 3.77 (s, 3H).

Compound 73

N-{2-[(3-aminobenzyl)sulfinyl]-5-methoxyphenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure L**, the title compound (55mg, 65%) was prepared from N-{5-methoxy-2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, acetone-d₆) δ 7.75 - 7.85 (m, 2H), 7.59 (dd, J = 0.59, 8.51 Hz, 1H), 7.44 - 7.53 (m, 1H), 7.32 - 7.40 (m, 1H), 7.25 (s, 1H), 6.93 (d, J = 8.51 Hz, 1H), 6.85 - 6.90 (m, 1H), 6.61 (dd, J = 2.49, 8.66 Hz, 1H), 6.53 - 6.58 (m, 1H), 6.40 (t, J = 1.76 Hz, 1H), 6.21 (d, J = 7.92 Hz, 1H), 4.25 (d, J = 12.32 Hz, 1H), 4.08 (d, J = 11.74 Hz, 2H), 3.78 (s, 3H).

Compound 74


N-(5-methoxy-2-((3-nitrobenzyl)sulfonyl)phenyl)benzofuran-2-sulfonamide



Following **General Procedure D**, the title compound (165mg, 83%) was prepared from N-(5-methoxy-2-((3-nitrobenzyl)thio)phenyl)benzofuran-2-sulfonamide. ¹H NMR (600 MHz, CDCl₃) δ 9.28 (s, 1H), 8.14 - 8.19 (m, 1H), 7.70 (d, *J* = 7.63 Hz, 1H), 7.64 (s, 1H), 7.58 (s, 1H), 7.41 - 7.50 (m, 3H), 7.37 (d, *J* = 9.10 Hz, 1H), 7.33 (td, *J* = 1.03, 7.41 Hz, 1H), 7.29 (d, *J* = 2.35 Hz, 1H), 7.26 (s, 1H), 6.57 (dd, *J* = 2.35, 8.80 Hz, 1H), 4.38 (s, 2H), 3.86 (s, 3H).

Compound 75

N-{2-[(3-aminobenzyl)sulfonyl]-5-methoxyphenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure L**, the title compound (67mg, 45%) was prepared from N-(5-methoxy-2-((3-nitrobenzyl)sulfonyl)phenyl)benzofuran-2-sulfonamide.

¹H NMR (600 MHz, CD₃OD) δ 7.70 (d, *J* = 7.63 Hz, 1H), 7.42 - 7.50 (m, 2H), 7.35 - 7.41 (m, 1H), 7.33 (d, *J* = 8.80 Hz, 1H), 7.27 - 7.31 (m, 1H), 7.22 (d, *J* = 2.35 Hz, 1H), 6.76 (t, *J* = 7.78 Hz, 1H), 6.53 (dd, *J* = 1.32, 8.07 Hz, 1H), 6.45 (s, 1H), 6.39 (d, *J* = 8.51 Hz, 1H), 6.27 (d, *J* = 7.63 Hz, 1H), 4.61 (br. s., 2H), 3.76 (s, 3H).

Compound 76

Benzofuran-2-sulfonic acid [2-(3-nitro-benzylsulfanyl)-phenyl]-amide




Following **General Procedure B**, the title compound (520 mg, 41%) was prepared from 2-((3-nitrobenzyl)thio)aniline (741 mg, 2.85 mmol), benzofuran-2-sulfonyl chloride (616 mg, 2.85 mmol) in pyridine (10 ml).

- 5 ¹H NMR (600 MHz, CD₃OD) δ 7.97 - 8.04 (m, 1H), 7.79 - 7.89 (m, 1H), 7.70 (d, *J* = 0.88 Hz, 1H), 7.50 (d, *J* = 0.88 Hz, 1H), 7.29 - 7.47 (m, 6H), 7.15 - 7.25 (m, 2H), 7.02 - 7.11 (m, 1H), 3.99 (s, 2H).

Compound 77

N-{2-[(3-aminobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide




10

Following **General Procedure L**, the title compound (67mg, 45%) was prepared from benzofuran-2-sulfonic acid [2-(3-nitro-benzylsulfanyl)-phenyl]-amide.

- ¹H NMR (600 MHz, CD₃OD) δ 7.66 (d, *J* = 7.92 Hz, 1H), 7.47 - 7.52 (m, 1H), 7.38 - 7.46 (m, 2H), 7.26 - 7.35 (m, 2H), 7.12 - 7.22 (m, 2H), 7.04 (td, *J* = 1.47, 7.63 Hz, 1H),
 15 6.88 (t, *J* = 7.78 Hz, 1H), 6.54 (dd, *J* = 1.47, 7.92 Hz, 1H), 6.43 (t, *J* = 1.76 Hz, 1H), 6.28 (d, *J* = 7.63 Hz, 1H), 3.69 (s, 2H).

Compound 78


N-{2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide



Following **General Procedure C**, the title compound (172mg, 73%) was prepared from benzofuran-2-sulfonic acid [2-(3-nitro-benzylsulfanyl)-phenyl]-amide. ¹H NMR (600 MHz, acetone-d₆) δ 10.45 (br. s., 1H), 8.15 (d, J = 8.22 Hz, 1H), 7.85 (s, 1H), 7.80 (d, J = 7.92 Hz, 1H), 7.66 (d, J = 0.88 Hz, 1H), 7.59 - 7.63 (m, 1H), 7.42 - 7.57 (m, 5H), 7.34 - 7.40 (m, 1H), 7.13 - 7.24 (m, 2H), 4.54 - 4.60 (m, 1H), 4.45 - 4.51 (m, 1H).

Compound 79


N-{2-[(3-aminobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide



Following **General Procedure L**, the title compound (122 mg, 90%) was prepared from N-{2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide. ¹H NMR (600 MHz, CD₃OD) δ 7.73 (d, J = 7.63 Hz, 1H), 7.59 (d, J = 8.22 Hz, 1H), 7.43 - 7.53 (m, 3H), 7.29 - 7.42 (m, 3H), 7.20 (d, J = 7.63 Hz, 1H), 6.97 (t, J = 7.63 Hz, 1H), 6.66 (d, J = 7.34 Hz, 1H), 6.57 (br. s., 1H), 6.43 (d, J = 7.34 Hz, 1H), 4.27 (d, J = 12.91 Hz, 1H), 3.99 (d, J = 12.91 Hz, 1H).

Compound 80

N-{2-[(3-aminobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure L**, the title compound (75 mg, 82%) was prepared from N-{2-[(3-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide. ¹H NMR (600 MHz, acetone-d₆) δ 7.75 - 7.85 (m, 3H), 7.62 - 7.70 (m, 1H), 7.59 (d, J = 8.51 Hz, 1H), 7.56 (dd, J = 1.76, 7.92 Hz, 1H), 7.49 (td, J = 1.17, 7.92 Hz, 1H), 7.36

(t, $J = 7.48$ Hz, 1H), 7.16 - 7.22 (m, 1H), 6.83 (t, $J = 7.78$ Hz, 1H), 6.57 (dd, $J = 1.47$, 7.92 Hz, 1H), 6.43 (t, $J = 1.91$ Hz, 1H), 6.15 (d, $J = 7.34$ Hz, 1H), 4.37 (s, 2H).

Intermediate 28

5-chloro-2-((4-nitrobenzyl)thio)aniline




5

Following **General Procedure A**, the title compound (624 mg, 92%) was prepared from 2-amino-4-chlorobenzenethiol (555 mg, 3.10 mmol), 1-(bromomethyl)-4-nitrobenzene (490 mg, 2.268 mmol) and K_2CO_3 (1.3 g, 9.50 mmol) in DMF (20 ml). ^1H NMR (600 MHz, CD_3OD) δ 8.07 (d, $J = 8.80$ Hz, 2H), 7.30 (d, $J = 8.80$ Hz, 2H), 6.93 (d, $J = 8.22$ Hz, 1H), 6.73 (d, $J = 2.35$ Hz, 1H), 6.43 (dd, $J = 2.35$, 8.22 Hz, 1H), 3.97 (s, 2H).

10

Compound 81

N-{5-chloro-2-[(4-nitrobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide




15

Following **General Procedure B**, the title compound (475 mg, 72%) was prepared from 5-chloro-2-((4-nitrobenzyl)thio)aniline (411 mg, 1.40 mmol) and benzofuran-2-sulfonyl chloride (303 mg, 1.40 mmol) in pyridine (5 ml). ^1H NMR (600 MHz, CD_3OD) δ 7.94 - 8.00 (m, 2H), 7.72 (d, $J = 7.92$ Hz, 1H), 7.50 - 7.55 (m, 1H), 7.45 - 7.49 (m, 2H), 7.43 (s, 1H), 7.31 - 7.38 (m, 1H), 7.11 - 7.19 (m, 3H), 7.07 (dd, $J = 2.35$, 8.51 Hz, 1H), 3.96 (s, 2H).

20

Compound 82


N-{5-chloro-2-[(4-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide



Following **General Procedure D**, the title compound (120mg, 85%) was prepared from N-{5-chloro-2-[(4-nitrobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide. ¹H NMR (600 MHz, acetone-d₆) δ 9.59 (br. s., 1H), 8.01 (d, J = 7.34 Hz, 2H), 7.86 (s, 1H), 7.76 (d, J = 7.92 Hz, 1H), 7.72 (s, 1H), 7.52 (d, J = 8.51 Hz, 1H), 7.39 - 7.49 (m, 2H), 7.25 - 7.36 (m, 3H), 7.00 (s, 1H), 4.96 (s, 2H).

Compound 83

N-{2-[(4-aminobenzyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure L**, the title compound (68 mg, 81%) was prepared from N-{5-chloro-2-[(4-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, acetone-d₆) δ 7.75 - 7.88 (m, 3H), 7.61 (d, J = 8.51 Hz, 1H), 7.44 - 7.54 (m, 2H), 7.34 - 7.41 (m, 1H), 7.22 (dd, J = 1.76, 8.51 Hz, 1H), 6.69 (d, J = 8.51 Hz, 2H), 6.49 (d, J = 8.51 Hz, 2H), 4.38 (s, 2H).

Intermediate 29


5-chloro-2-((2-nitrobenzyl)thio)aniline



Following **General Procedure A**, the title compound (548 mg, 95%) was prepared from 2-amino-4-chlorobenzenethiol (441 mg, 2.76 mmol), 1-(bromomethyl)-2-nitrobenzene (398 mg, 1.84 mmol) and K₂CO₃ (1.27 g, 9.21 mmol) in DMF (20 ml).
1H NMR (600 MHz, CD₃OD) δ 7.89 - 7.98 (m, 1H), 7.37 - 7.46 (m, 2H), 6.99 - 7.04 (m, 1H), 6.81 (d, J = 8.22 Hz, 1H), 6.71 (d, J = 2.05 Hz, 1H), 6.37 (dd, J = 2.05, 8.22 Hz, 1H), 4.22 (s, 2H).

Compound 84


N-(5-chloro-2-((2-nitrobenzyl)thio)phenyl)benzofuran-2-sulfonamide



Following **General Procedure B**, the title compound (485 mg, 69%) was prepared from 5-chloro-2-((2-nitrobenzyl)thio)aniline (435 mg, 1.48 mmol) and benzofuran-2-sulfonyl chloride (320 mg, 1.48 mmol) in pyridine (5 ml).
1H NMR (600 MHz, CDCl₃) δ 8.03 (dd, J = 1.47, 7.92 Hz, 1H), 7.65 - 7.71 (m, 2H), 7.48 - 7.53 (m, 2H), 7.43 - 7.48 (m, 1H), 7.31 - 7.42 (m, 3H), 7.03 (d, J = 8.22 Hz, 1H), 6.90 (dd, J = 2.05, 8.22 Hz, 1H), 6.76 (dd, J = 1.32, 7.48 Hz, 1H), 4.16 (s, 2H).

Compound 85

N-{2-[(2-aminobenzyl)thio]-5-chlorophenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure L**, the title compound (44 mg, 40%) was prepared from N-(5-chloro-2-((2-nitrobenzyl)thio)phenyl)benzofuran-2-sulfonamide.
1H NMR (600 MHz, acetone-d₆) δ 7.77 (d, J = 7.92 Hz, 1H), 7.59 (dd, J = 0.88, 8.51 Hz, 1H), 7.47 - 7.55 (m, 3H), 7.31 - 7.41 (m, 2H), 7.10 (dd, J = 2.35, 8.51 Hz, 1H),

6.86 - 6.95 (m, 1H), 6.72 (dd, J = 0.88, 7.92 Hz, 1H), 6.57 (dd, J = 1.32, 7.48 Hz, 1H), 6.38 (td, J = 1.03, 7.41 Hz, 1H), 3.97 (s, 2H).

Compound 86

N-{5-chloro-2-[(2-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide




5

Following **General Procedure C**, the title compound (160mg, 79%) was prepared from N-(5-chloro-2-((2-nitrobenzyl)thio)phenyl)benzofuran-2-sulfonamide. ¹H NMR (600 MHz, acetone-d₆) δ 7.92 (d, J = 8.22 Hz, 1H), 7.71 (dd, J = 0.73, 7.78 Hz, 1H), 7.59 (d, J = 2.05 Hz, 1H), 7.47 (d, J = 8.51 Hz, 1H), 7.39 - 7.44 (m, 1H), 7.33 - 7.38 (m, 2H), 7.19 - 7.30 (m, 2H), 6.71 - 6.80 (m, 2H), 6.67 (d, J = 8.22 Hz, 1H), 5.13 (d, J = 12.91 Hz, 2H), 4.73 (d, J = 12.62 Hz, 2H).

10

Compound 87

N-{2-[(2-aminobenzyl)sulfinyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide




15


Following **General Procedure L**, the title compound (67 mg, 50%) was prepared from N-{5-chloro-2-[(2-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, CD₃OD) δ 7.72 (d, J = 7.92 Hz, 1H), 7.56 (d, J = 7.63 Hz, 1H), 7.44 - 7.52 (m, 2H), 7.41 (d, J = 8.51 Hz, 2H), 7.31 - 7.37 (m, 1H), 7.28 (d, J = 1.47 Hz, 1H), 7.21 (dd, J = 1.76, 8.22 Hz, 1H), 7.06 - 7.13 (m, 1H), 6.88 (d, J = 7.92 Hz, 1H), 6.75 (d, J = 7.04 Hz, 1H), 6.60 - 6.67 (m, 1H), 4.50 (d, J = 13.50 Hz, 2H), 4.23 (d, J = 12.91 Hz, 2H).


20

Compound 88**N-{2-[(2-aminobenzyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide**

Following **General Procedure D** and **K**, the title compound (52 mg) was prepared from N-(5-chloro-2-((2-nitrobenzyl)thio)phenyl)benzofuran-2-sulfonamide. ¹H NMR (300 MHz, CD₃OD) δ 7.69 - 7.78 (m, 2H), 7.56 - 7.65 (m, 2H), 7.50 - 7.55 (m, 1H), 7.45 (td, J = 1.17, 7.77 Hz, 1H), 7.28 - 7.38 (m, 1H), 6.99 - 7.15 (m, 2H), 6.88 (d, J = 7.03 Hz, 1H), 6.67 - 6.75 (m, 1H), 6.58 - 6.66 (m, 1H), 4.68 (s, 2H).

Intermediate 30**5-chloro-2-((pyrimidin-2-ylmethyl)thio)aniline**

Following **General Procedure A**, the title compound (284 mg, 39%) was prepared from 2-amino-4-chlorobenzenethiol (529 mg, 3.33 mmol), 2-(chloromethyl)pyrimidine (366 mg, 2.22 mmol) and K₂CO₃ (1.53 g, 11.09 mmol) in DMF (10 ml). ¹H NMR (600 MHz, CD₃OD) δ 8.65 (d, J = 4.99 Hz, 2H), 7.31 (t, J = 4.99 Hz, 1H), 7.03 (d, J = 8.22 Hz, 1H), 6.70 (s, 1H), 6.43 (dd, J = 2.05, 8.22 Hz, 1H), 4.07 (s, 2H).


Compound 89**N-{5-chloro-2-[(pyrimidin-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide**

Following **General Procedure B**, the title compound (243 mg, 51%) was prepared from 5-chloro-2-((pyrimidin-2-ylmethyl)thio)aniline (280 mg, 1.11 mmol) and benzofuran-2-sulfonyl chloride (240 mg, 1.11 mmol) in pyridine (3 ml).

¹H NMR (600 MHz, acetone-d₆) δ 11.69 (br. s., 1H), 8.87 (d, J = 4.99 Hz, 2H), 7.74 (dd, J = 0.88, 7.92 Hz, 1H), 7.66 (s, 1H), 7.62 (d, J = 8.22 Hz, 1H), 7.59 (d, J = 0.59 Hz, 1H), 7.45 - 7.52 (m, 3H), 7.33 (ddd, J = 1.76, 6.38, 8.00 Hz, 1H), 7.14 (dd, J = 2.20, 8.36 Hz, 1H), 4.24 (s, 2H).

Compound 90

N-{5-chloro-2-[(pyrimidin-2-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure C**, the title compound (82mg) was prepared from N-{5-chloro-2-[(pyrimidin-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, CD₃OD) δ 8.68 (dd, J = 1.47, 4.99 Hz, 2H), 7.73 (d, J = 7.92 Hz, 1H), 7.53 - 7.60 (m, 2H), 7.45 - 7.52 (m, 2H), 7.30 - 7.41 (m, 4H), 4.63 (d, J = 13.50 Hz, 1H), 4.46 (d, J = 13.50 Hz, 1H).

Compound 91

N-{5-chloro-2-[(pyrimidin-2-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide



Following **General Procedure D**, the title compound (60mg) was prepared from N-{5-chloro-2-[(pyrimidin-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide.


¹H NMR (600 MHz, DMSO-d₆) δ 8.68 (dd, *J* = 1.47, 4.99 Hz, 2H), 7.76 (s, 1H), 7.68 (br. s., 1H), 7.61 (d, *J* = 8.22 Hz, 1H), 7.54 - 7.58 (m, 2H), 7.46 (t, *J* = 7.34 Hz, 1H), 7.41 (t, *J* = 4.84 Hz, 1H), 7.30 - 7.37 (m, 1H), 7.19 (d, *J* = 7.92 Hz, 1H), 5.11 - 5.19 (m, 2H).

5

GENERAL PROCEDURE M

Intermediate 31

4-mercapto-3-nitrobenzonitrile




To a solution of 4-chloro-3-nitrobenzonitrile (530 mg, 2.88 mmol) in dioxane (5 ml) / water (1 ml) was added Na₂S·9H₂O 9692 mg, 2.88 mmol). After it was stirred at room temperature for 2 hours, the reaction was quenched with HCl (1N). The mixture was extracted with EtOAc (2 x 50 ml). The organic layer was washed with water, brine, dried over Na₂SO₄ and concentrated in *vacuo*. The title compound was purified by silica gel column chromatography (0 ~ 30% EtOAc in hexane).

¹H NMR (600 MHz, acetone-d₆) δ 8.67 (d, *J* = 1.76 Hz, 1H), 8.01 (s, 1H), 7.93 (dd, *J* = 1.76, 8.51 Hz, 1H), 5.30 (s, 1H).

15

Intermediate 32

4-(benzylthio)-3-nitrobenzonitrile




Following **General Procedure A**, the title compound (580 mg, 82%) was prepared from 4-mercapto-3-nitrobenzonitrile (473 mg, 2.63 mmol), (bromomethyl)benzene (449 mg, 2.63 mmol), K₂CO₃ (1.81 g, 13.14 mmol) in DMF (10 ml).


¹H NMR (600 MHz, DMSO-d₆) δ 8.68 (d, *J* = 1.76 Hz, 1H), 8.10 (dd, *J* = 1.47, 8.51 Hz, 1H), 7.88 (d, *J* = 8.51 Hz, 1H), 7.43 (d, *J* = 7.34 Hz, 2H), 7.34 (t, *J* = 7.63 Hz, 2H), 7.25 - 7.30 (m, 1H), 4.44 (s, 2H).

20

25

Intermediate 33**3-amino-4-(benzylthio)benzonitrile**


Following **General Procedure L**, the title compound (372 mg, 81%) was
5 prepared from 4-(benzylthio)-3-nitrobenzonitrile (518 mg, 1.92 mmol).
1H NMR (600 MHz, CD₃OD) δ 7.11 - 7.25 (m, 6H), 6.96 (d, J = 1.76 Hz, 1H), 6.71 -
6.78 (m, 1H), 4.00 (s, 2H).

Compound 92**N-[2-(benzylthio)-5-cyanophenyl]-1-benzofuran-2-sulfonamide**

10

Following **General Procedure B**, the title compound (419 mg, 65%) was prepared
from 3-amino-4-(benzylthio)benzonitrile (370 mg, 1.54 mmol) and benzofuran-2-
sulfonyl chloride (333 mg, 1.54 mmol) in pyridine (3 ml).

1H NMR (600 MHz, CD₃OD) δ 7.73 (d, J = 8.51 Hz, 1H), 7.65 (d, J = 1.17 Hz, 1H),
15 7.50 - 7.56 (m, 2H), 7.48 (dd, J = 1.47, 8.22 Hz, 1H), 7.39 (ddd, J = 2.05, 6.16, 7.92
Hz, 1H), 7.31 - 7.36 (m, 2H), 7.09 - 7.20 (m, 3H), 6.92 (d, J = 7.04 Hz, 2H), 3.97 (s,
2H).

Compound 93**N-[2-(benzylsulfinyl)-5-cyanophenyl]-1-benzofuran-2-sulfonamide**


20

Following **General Procedure C**, the title compound (109 mg) was prepared from N-[2-(benzylthio)-5-cyanophenyl]-1-benzofuran-2-sulfonamide.

1H NMR (600 MHz, CD₃OD) δ 7.74 (d, J = 7.63 Hz, 1H), 7.66 (s, 1H), 7.53 (d, J = 8.51 Hz, 1H), 7.50 (s, 1H), 7.45 (t, J = 7.48 Hz, 1H), 7.30 - 7.39 (m, 2H), 7.28 (d, J = 7.92 Hz, 1H), 7.20 - 7.25 (m, 1H), 7.16 (t, J = 7.48 Hz, 2H), 6.96 (d, J = 7.34 Hz, 2H), 4.47 (d, J = 13.21 Hz, 1H), 4.21 (d, J = 12.91 Hz, 1H).

Compound 94

N-[2-(benzylsulfonyl)-5-cyanophenyl]-1-benzofuran-2-sulfonamide




10 Following **General Procedure D**, the title compound (230 mg) was prepared from N-[2-(benzylthio)-5-cyanophenyl]-1-benzofuran-2-sulfonamide.

1H NMR (600 MHz, CD₃OD) δ 7.97 (d, J = 1.47 Hz, 1H), 7.68 (d, J = 7.63 Hz, 1H), 7.52 (d, J = 8.22 Hz, 1H), 7.43 (s, 1H), 7.32 - 7.38 (m, 2H), 7.24 - 7.30 (m, 1H), 7.09 - 7.18 (m, 3H), 6.98 - 7.06 (m, 2H), 6.87 (dd, J = 1.47, 8.22 Hz, 1H), 5.05 (s, 2H).

15

Compound 95

4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide




20 Following **General Procedure B and D**, the title compound was prepared from 5-chloro-2-(methylthio)aniline and 4-chloro-3-(trifluoromethyl)benzene-1-sulfonyl chloride.

1H NMR (300 MHz, CD₃OD) δ 8.26 (s, 1H), 8.11 (d, J = 8.50 Hz, 1H), 7.74 - 7.87 (m, 2H), 7.58 (d, J = 1.47 Hz, 1H), 7.16 (d, J = 8.50 Hz, 1H), 3.19 (s, 3H).

Compound 96

4-chloro-N-[5-chloro-2-(methylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide




5 Following **General Procedure B**, the title compound was prepared from 5-chloro-2-(methylthio)aniline and 4-chloro-3-(trifluoromethyl)benzene-1-sulfonyl chloride.

1H NMR (300 MHz, CDCl₃) δ 8.13 (d, J = 1.47 Hz, 1H), 7.88 (dd, J = 1.90, 8.35 Hz, 1H), 7.60 (dd, J = 2.64, 12.60 Hz, 2H), 7.28 (d, J = 8.50 Hz, 1H), 7.10 (dd, J = 2.05, 8.50 Hz, 1H), 2.21 (s, 3H).

Compound 97

4-chloro-N-[5-chloro-2-(isopropylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide




15 Following **General Procedure A and B**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 2-iodo-propane, and 4-chloro-3-(trifluoromethyl)benzene-1-sulfonyl chloride.

1H NMR (300 MHz, CD₃OD) δ 8.13 (d, J = 1.47 Hz, 1H), 7.92 (dd, J = 2.05, 8.50 Hz, 1H), 7.76 (d, J = 8.50 Hz, 1H), 7.53 (d, J = 2.05 Hz, 1H), 7.33 (d, J = 8.50 Hz, 1H), 7.15 - 7.23 (m, 1H), 3.03 - 3.21 (m, 1H), 1.06 (d, J = 6.45 Hz, 6H).

Compound 98

4-chloro-N-{5-chloro-2-[(2-hydroxyethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide




Following **General Procedure A and B**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 2-bromo-ethanol, and 4-chloro-3-(trifluoromethyl)benzene-1-sulfonyl chloride.

- 5 ¹H NMR (300 MHz, CD₃OD) δ 8.09 (d, J = 1.47 Hz, 1H), 7.92 (dd, J = 1.90, 8.35 Hz, 1H), 7.76 (d, J = 8.50 Hz, 1H), 7.51 (d, J = 2.34 Hz, 1H), 7.41 (d, J = 8.50 Hz, 1H), 7.20 (dd, J = 2.20, 8.35 Hz, 1H), 3.49 (t, J = 6.15 Hz, 2H), 2.80 (t, J = 6.15 Hz, 2H).

Compound 99

4-chloro-N-[5-chloro-2-(isopropylsulfinyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide

10




Following **General Procedure C**, the title compound (55 mg) was prepared from 4-chloro-N-[5-chloro-2-(isopropylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide.

15

Compound 100

4-chloro-N-[5-chloro-2-[(2-hydroxyethyl)sulfinyl]phenyl]-3-(trifluoromethyl)benzenesulfonamide



Following **General Procedure C**, the title compound (109 mg) was prepared from 4-chloro-N-[5-chloro-2-[(2-hydroxyethyl)thio]phenyl]-3-(trifluoromethyl)benzenesulfonamide.


20

¹H NMR (300 MHz, CD₃OD) δ 8.09 (d, J = 1.17 Hz, 1H), 7.96 - 8.03 (m, 1H), 7.84 (d, J = 8.20 Hz, 1H), 7.71 (d, J = 8.50 Hz, 1H), 7.44 (dd, J = 1.76, 8.50 Hz, 1H), 7.21 (d, J = 1.76 Hz, 1H), 3.98 (td, J = 4.10, 8.06 Hz, 1H), 3.77 - 3.88 (m, 1H), 3.19 (ddd, J = 5.27, 8.42, 13.26 Hz, 1H), 2.95 - 3.06 (m, 1H).

5

Compound 101

4-chloro-N-{5-chloro-2-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide



Following **General Procedure D**, the title compound was prepared from 4-


10 chloro-N-{5-chloro-2-[(2-hydroxyethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide.

¹H NMR (600 MHz, CD₃OD) δ 8.27 (d, J = 1.47 Hz, 1H), 8.11 (dd, J = 2.05, 8.51 Hz, 1H), 7.74 (dd, J = 8.51, 14.97 Hz, 2H), 7.54 - 7.59 (m, 1H), 6.96 - 7.08 (m, 1H), 3.82 (t, J = 6.02 Hz, 2H), 3.57 - 3.68 (m, 2H).

15

Compound 102


methyl 3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}propanoate



Following **General Procedure H, B, and C**, the title compound was prepared

20 from 2-amino-4-chlorobenzenethiol.


¹H NMR (300 MHz, CD₃OD) δ 8.09 (s, 1H), 7.99 (dd, J = 1.76, 8.50 Hz, 1H), 7.85 (d, J = 8.20 Hz, 1H), 7.71 (d, J = 8.50 Hz, 1H), 7.47 (dd, J = 1.61, 8.35 Hz, 1H), 7.11 (d, J = 1.76 Hz, 1H), 3.63 (s, 3H), 3.24 - 3.39 (m, 1H), 3.08 - 3.22 (m, 1H), 2.54 - 2.82 (m, 2H).

Compound 103**4-chloro-N-{5-chloro-2-[(pyridin-2-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide**

5 Following **General Procedure A and B**, the title compound was prepared from 2-amino-4-chlorobenzenethiol.

¹H NMR (600 MHz, CD₃OD) δ 8.63 (d, J = 3.81 Hz, 1H), 8.03 (d, J = 2.35 Hz, 1H), 7.87 (dd, J = 2.20, 8.36 Hz, 1H), 7.75 (td, J = 1.76, 7.63 Hz, 1H), 7.70 (d, J = 8.51 Hz, 1H), 7.54 (d, J = 2.35 Hz, 1H), 7.37 (d, J = 8.51 Hz, 1H), 7.32 - 7.35 (m, 1H), 7.21 (d, J = 7.63 Hz, 1H), 7.13 (dd, J = 2.35, 8.22 Hz, 1H), 4.00 (s, 2H).

10

Compound 104**4-chloro-N-{5-chloro-2-[(pyridin-3-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide**


15 Following **General Procedure A and B**, the title compound was prepared from 2-amino-4-chlorobenzenethiol.

¹H NMR (600 MHz, CD₃OD) δ 8.38 (d, J = 3.82 Hz, 1H), 8.25 (br. s., 1H), 8.11 (d, J = 2.35 Hz, 1H), 7.93 (dd, J = 2.05, 8.51 Hz, 1H), 7.78 (d, J = 8.51 Hz, 1H), 7.60 (dt, J = 1.91, 7.92 Hz, 1H), 7.39 (d, J = 2.35 Hz, 1H), 7.33 (dd, J = 4.84, 7.78 Hz, 1H), 7.20 (d, J = 8.51 Hz, 1H), 7.14 (s, J = 2.35, 2.35, 8.51, 8.51 Hz, 1H), 3.95 (s, 2H).

20

Compound 105

4-chloro-N-{5-chloro-2-[(pyridin-2-ylmethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide




5 Following **General Procedure C**, the title compound was prepared from 4-chloro-N-{5-chloro-2-[(pyridin-2-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide.

¹H NMR (600 MHz, DMSO-d₆) δ 8.44 (d, J = 4.40 Hz, 1H), 8.03 (d, J = 2.05 Hz, 1H), 7.98 (dd, J = 2.05, 8.22 Hz, 1H), 7.88 (d, J = 8.22 Hz, 1H), 7.74 (t, 1H), 7.25 - 7.33 (m, 2H), 7.20 (d, J = 7.63 Hz, 1H), 7.14 (br. s., 1H), 7.03 (d, J = 1.47 Hz, 1H), 4.49 (d, J = 12.62 Hz, 1H), 4.06 (d, J = 12.91 Hz, 1H).

Compound 106


4-chloro-N-{5-chloro-2-[(pyridin-3-ylmethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide



15 Following **General Procedure C**, the title compound was prepared from 4-chloro-N-{5-chloro-2-[(pyridin-3-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide.

¹H NMR (600 MHz, CD₃OD) δ 8.35 (dd, J = 1.61, 4.84 Hz, 1H), 8.24 (d, J = 2.35 Hz, 1H), 8.07 (d, J = 2.05 Hz, 1H), 8.03 (d, J = 1.76 Hz, 1H), 7.71 (d, J = 8.51 Hz, 1H), 7.38 (s, 1H), 7.27 (d, J = 2.05 Hz, 1H), 7.23 - 7.26 (m, 1H), 6.75 (d, J = 8.22 Hz, 1H), 6.65 (dd, J = 1.91, 8.36 Hz, 1H), 4.45 (d, J = 13.50 Hz, 1H), 4.37 (d, J = 13.21 Hz, 1H).

GENERAL PROCEDURE N


Intermediate 34**4-Chloro-N-(5-chloro-2-mercapto-phenyl)-3-trifluoromethyl-benzenesulfonamide**

To a solution of N,N'-[dithiobis(5-chloro-2,1-phenylene)]bis[4-chloro-3-(trifluoromethyl)benzenesulfonamide] **Intermediate 5** (106 mg, 0.13 mmol) in THF (3 ml) at 0 °C was added NaBH₄ (20 mg, 0.53 mmol), the mixture was stirred at room temperature for 1 h, diluted with H₂O, acidified carefully with 6 M HCl, extracted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄, and concentrated *in vacuo* to yield 109 mg off-white solid as the title compound.

¹H NMR (600 MHz, METHANOL-d₄) δ 8.05 (s, 1H), 7.91 (d, *J* = 8.51 Hz, 1H), 7.78 (d, *J* = 8.51 Hz, 1H), 7.30 (dd, *J* = 1.47, 8.22 Hz, 1H), 7.23 - 7.25 (m, 1H), 7.14 (dd, *J* = 2.20, 8.36 Hz, 1H).

Compound 107

ethyl {[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}acetate




Following **General Procedure A and C**, the title compound was prepared from 4-Chloro-N-(5-chloro-2-mercapto-phenyl)-3-trifluoromethyl-benzenesulfonamide

Intermediate 34 and ethyl 2-bromoacetate.

¹H NMR (300 MHz, CD₃OD) δ 8.06 (s, 1H), 7.96 (d, *J* = 8.50 Hz, 1H), 7.84 (dd, *J* = 8.50, 10.84 Hz, 2H), 7.52 (dd, *J* = 1.61, 8.64 Hz, 1H), 7.06 (d, *J* = 1.47 Hz, 1H), 4.17 (q, *J* = 7.23 Hz, 2H), 4.06 (d, *J* = 14.65 Hz, 1H), 3.88 (d, *J* = 14.36 Hz, 1H), 1.22 (t, *J* = 7.18 Hz, 3H).

GENERAL PROCEDURE O

Intermediate 35**4-Chloro-N-[5-chloro-2-(3-oxo-cyclopentyl)sulfanyl]-phenyl]-3-trifluoromethyl-benzenesulfonamide**

5

To a solution of N,N'-[dithiobis(5-chloro-2,1-phenylene)]bis[4-chloro-3-(trifluoromethyl)benzenesulfonamide] **Intermediate 5** (105 mg, 0.13 mmol) in CH₂Cl₂ (2 ml), MeOH (0.5 ml), and H₂O (0.25 ml) was added polymer-bound triphenylphosphine (~3 mmol/g triphenylphosphine loading, 87 mg, 0.26 mmol), cyclopent-2-enone (33 μ l, 0.39 mmol), and PTSA (catalytic amount). The reaction was stirred at room temperature for 4 h and was directly purified by column chromatography on silica gel (0→100% ethyl acetate in hexane) to yield the title compound as an off-white solid (133 mg, 100%).


10

15

¹H NMR (600 MHz, CHLOROFORM-d) δ 8.16 (d, J = 2.35 Hz, 1H), 7.93 (dd, J = 2.35, 8.51 Hz, 1H), 7.89 (s, 1H), 7.62 - 7.65 (m, 2H), 7.35 (d, J = 8.22 Hz, 1H), 7.07 (dd, J = 2.20, 8.36 Hz, 1H), 3.46 (quin, J = 6.53 Hz, 1H), 2.44 - 2.49 (m, 1H), 2.38 - 2.44 (m, 1H), 2.21 (s, 2H), 2.05 - 2.11 (m, 1H), 1.84 - 1.90 (m, 1H).

Compound 108**4-chloro-N-{5-chloro-2-[(3-hydroxycyclopentyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide**

20



Following **General Procedure D**, followed by treatment of the crude product with NaBH₄ in MeOH, the title compound was prepared from 4-Chloro-N-[5-chloro-2-

(3-oxo-cyclopentylsulfanyl)-phenyl]-3-trifluoromethyl-benzenesulfonamide


(Intermediate 35).

1H NMR (600 MHz, CD₃OD) δ 8.13 (d, J = 2.05 Hz, 1H), 7.99 (dd, J = 2.20, 8.36 Hz, 1H), 7.73 (d, J = 8.51 Hz, 1H), 7.46 (d, J = 8.51 Hz, 1H), 7.34 (d, J = 2.05 Hz, 1H),
 5 7.06 (dd, J = 2.05, 8.51 Hz, 1H), 4.17 - 4.22 (m, 1H), 3.70 - 3.77 (m, 1H), 2.19 - 2.26 (m, 1H), 1.90 - 2.00 (m, 1H), 1.78 - 1.86 (m, 1H), 1.62 - 1.76 (m, 2H), 1.29 - 1.38 (m, 1H).

Compound 109

4-chloro-N-[5-chloro-2-(ethylsulfonyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide

10




Following **General Procedure D**, the title compound was prepared from 4-chloro-N-[5-chloro-2-(ethylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide.

1H NMR (600 MHz, CD₃OD) δ 8.30 (d, J = 1.76 Hz, 1H), 8.14 (dd, J = 2.20, 8.36 Hz, 1H), 7.72 - 7.80 (m, 2H), 7.64 (d, J = 2.05 Hz, 1H), 7.10 (br. s., 1H), 3.30 - 3.38 (m, 2H), 1.09 (t, J = 7.34 Hz, 3H)

15

Compound 110

2-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio]-N,N-dimethylacetamide




20

Following **General Procedure G**, the title compound was prepared from N,N'-[dithiobis(5-chloro-2,1-phenylene)]bis[4-chloro-3-(trifluoromethyl)benzenesulfonamide] **Intermediate 5** and 2-chloro-N,N-dimethyl-acetamide.

¹H NMR (600 MHz, CD₃OD) δ 8.09 (d, J = 2.35 Hz, 1H), 7.94 (dd, J = 2.05, 8.51 Hz, 1H), 7.76 (d, J = 8.22 Hz, 1H), 7.47 - 7.50 (m, 2H), 7.18 (dd, J = 2.35, 8.51 Hz, 1H), 3.68 (s, 2H), 2.97 (s, 3H), 2.93 (s, 3H).

Compound 111

2-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio]-N-methylacetamide




Following **General Procedure G**, the title compound was prepared from N,N'-[dithiobis(5-chloro-2,1-phenylene)]bis[4-chloro-3-(trifluoromethyl)benzenesulfonamide]

Intermediate 5 and 2-chloro-N-methyl-acetamide.

¹H NMR (600 MHz, CD₃OD) δ 8.09 (d, J = 2.35 Hz, 1H), 7.93 (dd, J = 2.35, 8.51 Hz, 1H), 7.76 (d, J = 8.51 Hz, 1H), 7.48 (d, J = 2.35 Hz, 1H), 7.44 (d, J = 8.22 Hz, 1H), 7.20 (dd, J = 2.35, 8.51 Hz, 1H), 3.40 (s, 2H), 2.69 (s, 3H).

Compound 112

2-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl]-N,N-dimethylacetamide



Following **General Procedure C**, the title compound was prepared from 2-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio]-N,N-


dimethylacetamide.

¹H NMR (600 MHz, CD₃OD) δ 8.16 (d, J = 2.35 Hz, 1H), 8.01 (dd, J = 2.05, 8.22 Hz, 1H), 7.68 (d, J = 8.51 Hz, 1H), 7.45 (d, J = 8.51 Hz, 1H), 7.28 (d, J = 1.76 Hz, 1H),

6.90 (dd, $J = 2.05, 8.22$ Hz, 1H), 4.65 - 4.71 (m, 1H), 3.51 - 3.57 (m, 1H), 3.11 (s, 3H), 2.98 (s, 3H).

Compound 113

2-[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino)phenyl]sulfonyl]-N,N-dimethylacetamide




Following **General Procedure D**, the title compound was prepared from 2-[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino)phenyl]thio]-N,N-dimethylacetamide.

^1H NMR (600 MHz, CD_3OD) δ 8.29 (d, $J = 2.05$ Hz, 1H), 8.09 - 8.14 (m, 1H), 7.72 (d, $J = 8.51$ Hz, 1H), 7.68 (d, $J = 8.51$ Hz, 1H), 7.48 (d, $J = 1.76$ Hz, 1H), 6.81 (d, $J = 8.22$ Hz, 1H), 3.16 (s, 3H), 3.00 (s, 3H).

Compound 114

2-[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino)phenyl]sulfonyl]-N-methylacetamide




Following **General Procedure C**, the title compound was prepared from 2-[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino)phenyl]thio]-N-methylacetamide.

^1H NMR (600 MHz, CD_3OD) δ 8.15 (d, $J = 2.05$ Hz, 1H), 8.01 (dd, $J = 2.20, 8.36$ Hz, 1H), 7.72 (d, $J = 8.51$ Hz, 1H), 7.51 (d, $J = 8.22$ Hz, 1H), 7.22 (d, $J = 2.05$ Hz, 1H), 7.02 (dd, $J = 1.91, 8.36$ Hz, 1H), 4.21 (d, $J = 13.50$ Hz, 1H), 3.49 (d, $J = 13.21$ Hz, 1H), 2.76 (s, 3H)

Compound 115

2-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfonyl]-N-methylacetamide




Following **General Procedure D**, the title compound was prepared from 2-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio]-N-methylacetamide.

^1H NMR (600 MHz, CD_3OD) δ 8.29 (d, J = 2.05 Hz, 1H), 8.09 - 8.13 (m, 1H), 7.69 (d, J = 8.51 Hz, 1H), 7.66 (d, J = 8.22 Hz, 1H), 7.44 (d, J = 2.05 Hz, 1H), 6.75 (dd, J = 1.91, 8.66 Hz, 1H), 2.77 (s, 3H).

Compound 116

N-{2-[(2-aminoethyl)thio]-5-chlorophenyl}-4-chloro-3-(trifluoromethyl)benzenesulfonamide




Following **General Procedure G**, the title compound was prepared from N,N'-[dithiobis(5-chloro-2,1-phenylene)]bis[4-chloro-3-(trifluoromethyl)benzenesulfonamide] **Intermediate 5** and 2-bromo-ethylamine hydrobromide.

^1H NMR (600 MHz, CD_3OD) δ 8.18 (d, J = 2.05 Hz, 1H), 7.99 (dd, J = 2.05, 8.22 Hz, 1H), 7.65 (d, J = 8.51 Hz, 1H), 7.38 (d, J = 8.22 Hz, 1H), 7.30 (d, J = 2.35 Hz, 1H), 6.72 (dd, J = 2.35, 8.22 Hz, 1H), 2.96 - 3.05 (m, 4H).

Compound 117

N-{2-[(2-aminoethyl)sulfonyl]-5-chlorophenyl}-4-chloro-3-(trifluoromethyl)benzenesulfonamide




Following **General Procedure D**, followed by treatment of the crude product with excess zinc dust in MeOH, aqueous NH₄Cl and HOAc at room temperature for 2 h, the title compound was prepared from N-{2-[(2-aminoethyl)thio]-5-chlorophenyl}-4-chloro-3-(trifluoromethyl)benzenesulfonamide.

¹H NMR (600 MHz, CD₃OD) δ 8.27 (d, *J* = 2.05 Hz, 1H), 8.10 (dd, *J* = 1.91, 8.36 Hz, 1H), 7.74 (d, *J* = 8.51 Hz, 1H), 7.67 (d, *J* = 8.51 Hz, 1H), 7.42 (d, *J* = 1.76 Hz, 1H), 6.80 (dd, *J* = 2.05, 8.51 Hz, 1H), 3.88 - 3.92 (m, 2H), 3.45 - 3.51 (m, 2H).

Compound 118

N-{2-[(2-aminoethyl)sulfinyl]-5-chlorophenyl}-4-chloro-3-(trifluoromethyl)benzenesulfonamide




Following **General Procedure C**, the title compound was prepared from N-{2-[(2-aminoethyl)thio]-5-chlorophenyl}-4-chloro-3-(trifluoromethyl)benzenesulfonamide.

¹H NMR (300 MHz, CD₃OD) δ 8.15 (d, *J* = 1.76 Hz, 1H), 8.00 (dd, *J* = 1.90, 8.35 Hz, 1H), 7.69 (d, *J* = 8.21 Hz, 1H), 7.46 (d, *J* = 8.50 Hz, 1H), 7.25 (d, *J* = 2.05 Hz, 1H), 6.94 (dd, *J* = 1.90, 8.35 Hz, 1H), 3.53 - 3.67 (m, 1H), 3.36 - 3.50 (m, 1H), 3.17 - 3.31 (m, 2H).

Compound 119

3-({2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}thio)-N,N-dimethylpropanamide




Following **General Procedure O**, the title compound was prepared from N,N'-[dithiobis(5-chloro-2,1-phenylene)]bis(1-benzofuran-2-sulfonamide) **Intermediate 8** and N,N-dimethyl-acrylamide.

¹H NMR (600 MHz, CD₃OD) δ 7.69 (d, *J* = 7.92 Hz, 1H), 7.53 - 7.56 (m, 1H), 7.45 - 7.49 (m, 2H), 7.39 (s, 1H), 7.37 (d, *J* = 8.51 Hz, 1H), 7.31 - 7.35 (m, 1H), 7.17 - 7.20 (m, 1H), 2.90 (t, *J* = 7.19 Hz, 2H), 2.87 (s, 3H), 2.85 (s, 3H), 2.33 (t, *J* = 7.19 Hz, 2H).

Compound 120

3-({2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfinyl)-N,N-dimethylpropanamide




Following **General Procedure C**, the title compound was prepared from 3-({2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}thio)-N,N-dimethylpropanamide.

¹H NMR (600 MHz, CD₃OD) δ 7.72 (d, *J* = 7.34 Hz, 1H), 7.64 (d, *J* = 8.22 Hz, 1H), 7.58 (d, *J* = 8.51 Hz, 1H), 7.46 - 7.51 (m, 1H), 7.46 (s, 1H), 7.39 (d, *J* = 1.76 Hz, 1H), 7.32 - 7.37 (m, 2H), 3.27 - 3.34 (m, 1H), 3.16 - 3.24 (m, 1H), 2.96 (s, 3H), 2.90 (s, 3H), 2.77 - 2.86 (m, 1H), 2.59 - 2.66 (m, 1H).

Compound 121

3-({2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfonyl)-N,N-dimethylpropanamide




Following **General Procedure D**, the title compound was prepared from 3-({2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}thio)-N,N-dimethylpropanamide.

¹H NMR (300 MHz, CD₃OD) δ 7.59 - 7.75 (m, 3H), 7.40 - 7.49 (m, 1H), 7.30 - 7.41 (m, 1H), 7.20 - 7.30 (m, 2H), 6.79 (dd, *J* = 1.76, 8.50 Hz, 1H), 3.94 (t, *J* = 7.47 Hz, 2H), 2.87 (s, 3H), 2.81 (s, 3H), 2.59 (t, *J* = 7.47 Hz, 2H).

Compound 122

N-(2-[[[6-amino-1-oxidopyridin-2-yl)methyl]sulfonyl]-5-chlorophenyl)-1-benzofuran-2-sulfonamide




Following **General Procedure D and E**, the title compound was prepared from tert-butyl {6-[[[2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl]thio)methyl]pyridin-2-yl}carbamate.

¹H NMR (300 MHz, CD₃OD) δ 7.77 (d, *J* = 1.76 Hz, 1H), 7.59 - 7.71 (m, 2H), 7.29 - 7.44 (m, 3H), 7.20 - 7.29 (m, 1H), 7.10 (t, *J* = 7.91 Hz, 1H), 6.77 - 6.88 (m, 2H), 6.57 (d, *J* = 7.33 Hz, 1H), 5.41 (br. s., 2H), 4.57 (br. s., 1H).

GENERAL PROCEDURE P

Compound 123

N-(2-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfonyl}ethyl)acetamide




To a solution of N-{2-[(2-aminoethyl)sulfonyl]-5-chlorophenyl}-4-chloro-3-(trifluoromethyl)benzenesulfonamide (23 mg, 0.048 mmol) in CH₂Cl₂ (2 ml) was added Et₃N (14 μl, 0.10 mmol), acetic anhydride (5 μl, 0.053 mmol), and catalytic amount of DMAP. The reaction was stirred at room temperature for 1 h and was concentrated. The crude product was purified by column chromatography on silica gel (0→10% MeOH in ethyl acetate), followed by PTLC (10% MeOH in ethyl acetate) to give the title compound.

¹H NMR (600 MHz, CD₃OD) δ 8.29 (d, *J* = 2.05 Hz, 1H), 8.13 (dd, *J* = 2.05, 8.51 Hz, 1H), 7.65 - 7.75 (m, 2H), 7.51 (d, *J* = 2.05 Hz, 1H), 6.82 (dd, *J* = 2.05, 8.51 Hz, 1H), 3.75 (t, *J* = 6.60 Hz, 2H), 3.51 (t, *J* = 6.60 Hz, 2H), 1.86 (s, 3H).

Compound 124

5

N-(2-([4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino)phenyl]sulfinyl)ethyl)acetamide



Following **General Procedure P and C**, the title compound was prepared from N-{2-[(2-aminoethyl)thio]-5-chlorophenyl}-4-chloro-3-(trifluoromethyl)benzenesulfonamide.


10

¹H NMR (600 MHz, CD₃OD) δ 8.12 (s, 1H), 8.01 (dd, *J* = 1.91, 8.36 Hz, 1H), 7.77 (d, *J* = 8.51 Hz, 1H), 7.56 (d, *J* = 8.22 Hz, 1H), 7.17 (s, 1H), 7.13 (d, *J* = 7.63 Hz, 1H), 3.60 - 3.71 (m, 1H), 3.44 - 3.54 (m, 1H), 3.33 - 3.42 (m, 1H), 3.04 - 3.15 (m, 1H), 1.98 (s, 3H).

15

Intermediate 36

4-Chloro-N,N-dimethyl-butylamide




To a solution of 4-chloro-butyl chloride (1.12 ml, 10.0 mmol) and dimethylamine hydrochloride (4.1 g, 50.0 mmol) in THF at 5 °C was added 2M NaOH (30 ml, 60.0 mmol) dropwise over 30 minutes while maintaining reaction temperature between 5-10 °C. The reaction was stirred for additional 1.5 h and was concentrated, extracted with EtOAc (×2). The combined organic layer was washed with 1M HCl (×2), brine, dried over Na₂SO₄, and concentrated *in vacuo* to yield the title compound as colorless oil (1.5 g, ~100%). The crude was used without further purification.

25

Compound 125

4-({2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfinyl)-N,N-dimethylbutanamide




5 Following **General Procedure G and C**, the title compound was prepared from N,N'-[dithiobis(5-chloro-2,1-phenylene)]bis(1-benzofuran-2-sulfonamide) **Intermediate 8** and 4-chloro-N,N-dimethyl-butylamide.

¹H NMR (300 MHz, CD₃OD) δ 7.68 (d, *J* = 7.62 Hz, 1H), 7.53 (d, *J* = 2.05 Hz, 1H), 7.35 - 7.49 (m, 3H), 7.32 (d, *J* = 0.88 Hz, 1H), 7.25 - 7.32 (m, 1H), 6.95 (dd, *J* = 2.05, 10 8.20 Hz, 1H), 3.33 - 3.49 (m, 1H), 2.98 (s, 3H), 2.91 - 3.08 (m, 1H), 2.90 (s, 3H), 2.38 - 2.57 (m, 2H), 1.92 - 2.09 (m, 2H).

Compound 126

4-({2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfonyl)-N,N-dimethylbutanamide




15

Following **General Procedure G and D**, the title compound was prepared from N,N'-[dithiobis(5-chloro-2,1-phenylene)]bis(1-benzofuran-2-sulfonamide) **Intermediate 8** and 4-chloro-N,N-dimethyl-butylamide.

¹H NMR (600 MHz, CD₃OD) δ 7.76 (s, 1H), 7.74 (d, *J* = 8.51 Hz, 1H), 7.70 (d, *J* = 7.92 20 Hz, 1H), 7.38 - 7.48 (m, 3H), 7.31 (t, *J* = 7.34 Hz, 1H), 6.86 - 7.05 (m, 1H), 3.46 - 3.69 (m, 2H), 2.97 (s, 3H), 2.91 (s, 3H), 2.43 - 2.59 (m, 2H), 1.93 - 2.09 (m, 2H).

Compound 127

5-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-1-benzofuran-2-sulfonamide




Following **General Procedure B and C**, the title compound was prepared from 5-chloro-2-(methylthio)aniline and 5-chloro-benzofuran-2-sulfonyl chloride (CAS #: 128852-02-8) (Bioorganic & Medicinal Chemistry 13 (2005) 3927–3954).

- 5 ^1H NMR (300 MHz, CDCl_3) δ 11.00 (br. s., 1H), 7.80 (d, J = 1.47 Hz, 1H), 7.68 (d, J = 1.76 Hz, 1H), 7.40 - 7.54 (m, 3H), 7.05 - 7.16 (m, 2H), 2.91 (s, 3H).

Compound 128

5-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-1-benzofuran-2-sulfonamide




- 10 Following **General Procedure B and D**, the title compound was prepared from 5-chloro-2-(methylthio)aniline and 5-chloro-benzofuran-2-sulfonyl chloride (CAS #: 128852-02-8) (Bioorganic & Medicinal Chemistry 13 (2005) 3927–3954).

^1H NMR (600 MHz, CD_3OD) δ 7.74 (s, 1H), 7.74 (d, J = 5.58 Hz, 1H), 7.64 (d, J = 1.76 Hz, 1H), 7.43 (d, J = 8.80 Hz, 1H), 7.30 - 7.35 (m, 2H), 6.88 (dd, J = 2.05, 8.51 Hz,

- 15 1H), 3.33 (s, 3H).

Compound 129


N-{5-chloro-2-[(1H-pyrazol-3-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide



- 20 Following **General Procedure C and E**, the title compound was prepared from tert-butyl 3-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl]-1H-pyrazole-1-carboxylate (**Compound 22**).


^1H NMR (600 MHz, CD_3OD) δ 7.67 (d, J = 7.63 Hz, 1H), 7.42 - 7.47 (m, 2H), 7.41 (d, J = 2.05 Hz, 1H), 7.37 (td, J = 1.47, 7.78 Hz, 1H), 7.32 (d, J = 0.88 Hz, 1H), 7.28 (td, J = 0.88, 7.48 Hz, 1H), 7.22 (d, J = 8.51 Hz, 1H), 6.96 (dd, J = 1.76, 8.51 Hz, 1H), 5.92 (s, 1H), 4.56 (d, J = 13.50 Hz, 1H), 4.28 (d, J = 13.50 Hz, 1H).

5

Compound 130**N-[5-chloro-2-(methylsulfinyl)phenyl]-4-isopropylbenzenesulfonamide**

Following **General Procedure B and C**, the title compound was prepared from 5-chloro-2-(methylthio)aniline and 4-isopropyl-benzenesulfonyl chloride.

10 ^1H NMR (600 MHz, CHLOROFORM-d) δ 10.42 (br. s., 1H), 7.80 - 7.86 (m, 2H), 7.68 (s, 1H), 7.35 (s, 2H), 7.05 - 7.08 (m, 1H), 7.01 - 7.03 (m, 1H), 2.94 (spt, J = 6.90 Hz, 1H), 2.69 (s, 3H), 1.22 (d, J = 7.04 Hz, 6H).


Compound 131**4-bromo-N-[5-chloro-2-(methylthio)phenyl]benzenesulfonamide**

15

Following **General Procedure B**, the title compound was prepared from 5-chloro-2-(methylthio)aniline and 4-bromo-benzenesulfonyl chloride.

^1H NMR (600 MHz, CHLOROFORM-d) δ 7.65 - 7.68 (m, 2H), 7.61 - 7.63 (m, 2H), 7.58 - 7.60 (m, 2H), 7.29 (d, J = 8.22 Hz, 1H), 7.04 (dd, J = 2.20, 8.36 Hz, 1H), 2.17 (s, 3H).

20


Compound 132**N-[5-chloro-2-(methylthio)phenyl]-4-iodobenzenesulfonamide**

Following **General Procedure B**, the title compound was prepared from 5-chloro-2-(methylthio)aniline and 4-iodo-benzenesulfonyl chloride.

¹H NMR (600 MHz, CHLOROFORM-d) δ 7.80 (d, *J* = 8.80 Hz, 2H), 7.62 - 7.64 (m, 1H), 7.61 (d, *J* = 2.05 Hz, 1H), 7.49 - 7.53 (m, 2H), 7.30 (d, *J* = 8.22 Hz, 1H), 7.04 (dd, *J* = 2.20, 8.36 Hz, 1H), 2.17 (s, 3H).

Compound 133

4-bromo-N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide




Following **General Procedure C**, the title compound was prepared from 4-bromo-N-[5-chloro-2-(methylthio)phenyl]benzenesulfonamide.

¹H NMR (600 MHz, CHLOROFORM-d) δ 10.63 - 10.77 (m, 1H), 7.79 - 7.82 (m, 2H), 7.64 - 7.68 (m, 3H), 7.05 (dd, *J* = 1.17, 2.35 Hz, 2H), 2.79 (s, 3H).

Compound 134

N-[5-chloro-2-(methylsulfinyl)phenyl]-4-iodobenzenesulfonamide




Following **General Procedure C**, the title compound was prepared from N-[5-chloro-2-(methylthio)phenyl]-4-iodobenzenesulfonamide.

¹H NMR (600 MHz, CHLOROFORM-d) δ 10.69 (br. s., 1H), 7.83 - 7.91 (m, 2H), 7.61 - 7.69 (m, 3H), 7.01 - 7.08 (m, 2H), 2.79 (s, 3H).

Compound 135

4-bromo-N-[5-chloro-2-(methylsulfonyl)phenyl]benzenesulfonamide




Following **General Procedure D**, the title compound was prepared from 4-bromo-N-[5-chloro-2-(methylthio)phenyl]benzenesulfonamide.

¹H NMR (600 MHz, CHLOROFORM-d) δ 9.20 - 9.27 (m, 1H), 7.80 (dd, *J* = 1.32, 8.36 Hz, 2H), 7.76 (d, *J* = 8.51 Hz, 1H), 7.68 (dd, *J* = 1.47, 8.80 Hz, 2H), 7.25 (d, *J* = 1.47 Hz, 1H), 7.18 - 7.21 (m, 1H), 2.93 (d, *J* = 1.47 Hz, 3H).

Compound 136

N-[5-chloro-2-(methylsulfonyl)phenyl]-4-iodobenzenesulfonamide




10 Following **General Procedure D**, the title compound was prepared from N-[5-chloro-2-(methylthio)phenyl]-4-iodobenzenesulfonamide.

¹H NMR (600 MHz, CHLOROFORM-d) δ 9.17 - 9.25 (m, 1H), 7.90 (d, *J* = 8.51 Hz, 2H), 7.76 (d, *J* = 8.51 Hz, 1H), 7.67 (d, *J* = 1.76 Hz, 1H), 7.64 (d, *J* = 8.51 Hz, 2H), 7.19 (dd, *J* = 1.76, 8.51 Hz, 1H), 2.93 (s, 3H).

15

Compound 137

N-{5-chloro-2-[(pyridin-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure A and B**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 2-bromomethyl-pyridine hydrobromide, and benzofuran-2-sulfonyl chloride.

20

¹H NMR (600 MHz, CHLOROFORM-d) δ 12.32 (br. s, 1H), 8.87 (dt, *J* = 0.77, 4.92 Hz, 1H), 7.76 (d, *J* = 2.05 Hz, 1H), 7.58 - 7.65 (m, 2H), 7.44 (dd, *J* = 0.88, 8.22 Hz, 1H), 7.34 - 7.42 (m, 3H), 7.25 - 7.30 (m, 2H), 7.06 (d, *J* = 7.63 Hz, 1H), 6.96 - 7.01 (m, 1H), 3.98 (s, 2H).

5


Compound 138**N-{5-chloro-2-[(pyridin-3-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide**

Following **General Procedure A and B**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 3-bromomethyl-pyridine hydrobromide, and benzofuran-2-sulfonyl chloride.

10

¹H NMR (600 MHz, CHLOROFORM-d) δ 8.42 (dd, *J* = 1.61, 4.84 Hz, 1H), 8.12 (d, *J* = 1.76 Hz, 2H), 7.66 - 7.70 (m, 2H), 7.50 (d, *J* = 1.17 Hz, 1H), 7.46 - 7.49 (m, 1H), 7.42 - 7.46 (m, 1H), 7.33 (ddd, *J* = 1.17, 6.97, 8.00 Hz, 1H), 7.26 - 7.29 (m, 1H), 7.13 (ddd, *J* = 0.88, 4.70, 7.92 Hz, 1H), 7.00 (d, *J* = 8.51 Hz, 1H), 6.89 (dd, *J* = 2.20, 8.36 Hz, 1H), 3.75 (s, 2H).

15

Compound 139**N-{5-chloro-2-[(pyridin-3-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide**

Following **General Procedure C**, the title compound was prepared from N-{5-chloro-2-[(pyridin-3-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide.


20

¹H NMR (600 MHz, METHANOL-d₄) δ 8.28 (d, *J* = 4.11 Hz, 1H), 7.86 (br. s., 1H), 7.68 - 7.71 (m, *J* = 0.73, 1.32 Hz, 1H), 7.54 (d, *J* = 1.76 Hz, 1H), 7.36 - 7.41 (m, 2H), 7.31 - 7.35 (m, 1H), 7.25 - 7.29 (m, *J* = 1.03, 7.78 Hz, 1H), 7.24 (d, *J* = 7.92 Hz, 1H), 7.12 (dd, *J* = 4.99, 7.63 Hz, 1H), 6.73 (d, *J* = 8.51 Hz, 1H), 6.68 (dd, *J* = 2.05, 8.51 Hz, 1H), 4.55 (s, 2H).

25

Compound 140

N-(5-chloro-2-[[1-(1-oxido-2,3-dihydro-1H-benzofuran-3-yl)methyl]sulfinyl]phenyl)-1-benzofuran-2-sulfonamide




5 Following **General Procedure C**, the title compound was prepared from N-{5-chloro-2-[(pyridin-3-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, METHANOL-d₄) δ 8.10 - 8.14 (m, 1H), 8.01 (t, *J* = 1.47 Hz, 1H), 7.66 - 7.68 (m, 1H), 7.42 - 7.49 (m, 2H), 7.34 (ddd, *J* = 1.17, 7.12, 8.44 Hz, 1H), 7.23 - 7.29 (m, 2H), 7.14 (dd, *J* = 6.60, 7.78 Hz, 1H), 6.87 (d, *J* = 7.92 Hz, 1H), 6.83 (d, *J* = 8.22 Hz, 1H), 6.71 (dd, *J* = 2.05, 8.22 Hz, 1H), 4.63 (d, *J* = 13.21 Hz, 1H), 4.37 (d, *J* = 13.21 Hz, 1H).

Compound 141

N-{5-chloro-2-[(pyridin-3-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide




15 Following **General Procedure D**, the title compound was prepared from N-{5-chloro-2-[(pyridin-3-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, METHANOL-d₄) δ 8.27 (d, *J* = 4.11 Hz, 1H), 8.04 (br. s., 1H), 7.78 (t, *J* = 1.76 Hz, 1H), 7.65 (d, *J* = 7.92 Hz, 1H), 7.46 (d, *J* = 0.88 Hz, 1H), 7.31 - 7.37 (m, 2H), 7.27 - 7.31 (m, 1H), 7.21 - 7.26 (m, 2H), 7.06 (dd, *J* = 5.14, 7.48 Hz, 1H), 6.67 (dt, *J* = 1.76, 8.51 Hz, 1H), 4.98 (s, 2H).

Compound 142

N-(5-chloro-2-[[1-(1-oxido-2,3-dihydro-1H-benzofuran-3-yl)methyl]sulfonyl]phenyl)-1-benzofuran-2-sulfonamide




Following **General Procedure D**, the title compound was prepared from N-{5-chloro-2-[(pyridin-3-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, DMSO-d₆) δ 8.01 - 8.08 (m, 2H), 7.68 (d, *J* = 7.92 Hz, 1H), 7.53 (d, *J* = 8.22 Hz, 1H), 7.46 - 7.48 (m, 1H), 7.31 - 7.37 (m, 2H), 7.23 - 7.27 (m, 2H), 7.11 - 7.15 (m, 1H), 6.98 (d, *J* = 7.92 Hz, 1H), 6.58 - 6.61 (m, 1H), 5.09 (s, 2H).

Compound 143

N-{5-chloro-2-[(3-nitrobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure G**, the title compound was prepared from N,N'-[dithiobis(5-chloro-2,1-phenylene)]bis(1-benzofuran-2-sulfonamide) **Intermediate 8** and 1-bromomethyl-3-nitro-benzene.

¹H NMR (600 MHz, CHLOROFORM-d) δ 8.08 (s, 1H), 8.02 (ddd, *J* = 1.17, 2.35, 8.22 Hz, 1H), 7.77 (t, *J* = 2.05 Hz, 1H), 7.65 - 7.69 (m, 2H), 7.50 (d, *J* = 0.88 Hz, 1H), 7.45 - 7.48 (m, 1H), 7.41 - 7.45 (m, 1H), 7.30 - 7.36 (m, 2H), 7.21 (dq, *J* = 0.88, 7.63 Hz, 1H), 7.03 (d, *J* = 8.22 Hz, 1H), 6.89 (dd, *J* = 2.05, 8.22 Hz, 1H), 3.84 (s, 2H).

Compound 144

N-{5-chloro-2-[(3-methoxybenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide




20

Following **General Procedure G**, the title compound was prepared from N,N'-[dithiobis(5-chloro-2,1-phenylene)]bis(1-benzofuran-2-sulfonamide) **Intermediate 8** and 1-bromomethyl-3-methoxy-benzene.

¹H NMR (600 MHz, CHLOROFORM-d) δ 8.11 (s, 1H), 7.64 - 7.68 (m, 2H), 7.40 - 7.50 (m, 3H), 7.31 (ddd, *J* = 0.88, 7.04, 7.92 Hz, 1H), 7.07 - 7.13 (m, 2H), 6.90 (dd, *J* = 2.05, 8.22 Hz, 1H), 6.73 (ddd, *J* = 1.17, 2.49, 8.36 Hz, 1H), 6.53 - 6.56 (m, 1H), 6.48 (t, *J* = 2.35 Hz, 1H), 3.74 (s, 2H), 3.66 (s, 3H).

Compound 145

N-{5-chloro-2-[(pyridin-2-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure C**, the title compound was prepared from N-{5-chloro-2-[(pyridin-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide .

¹H NMR (600 MHz, METHANOL-d₄) δ 8.46 (d, *J* = 4.70 Hz, 1H), 7.64 - 7.73 (m, 2H), 7.48 (d, *J* = 8.51 Hz, 1H), 7.38 - 7.45 (m, 3H), 7.25 - 7.34 (m, 3H), 7.07 - 7.17 (m, 2H), 4.58 (d, *J* = 12.91 Hz, 1H), 4.35 (d, *J* = 12.91 Hz, 1H).

Compound 146

N-{5-chloro-2-[(pyridin-2-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure D**, the title compound was prepared from N-{5-chloro-2-[(pyridin-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide .

¹H NMR (600 MHz, METHANOL-d₄) δ 8.42 (dt, *J* = 0.84, 4.77 Hz, 1H), 7.75 (d, *J* = 2.05 Hz, 1H), 7.67 (d, *J* = 7.63 Hz, 1H), 7.41 - 7.49 (m, 3H), 7.31 - 7.39 (m, 2H), 7.22 - 7.29 (m, 2H), 7.07 (d, *J* = 7.63 Hz, 1H), 6.76 (dd, *J* = 1.91, 8.66 Hz, 1H), 5.07 (s, 2H).

Compound 147


N-(5-chloro-2-[[1-(1-oxido-2-pyridin-2-yl)methyl]sulfonyl]phenyl)-1-benzofuran-2-sulfonamide



Following **General Procedure D**, the title compound was prepared from N-{5-chloro-2-[(pyridin-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide
¹H NMR (600 MHz, METHANOL-*d*₄) δ 8.35 (d, *J* = 6.46 Hz, 1H), 7.74 (d, *J* = 2.05 Hz, 1H), 7.66 (d, *J* = 7.92 Hz, 1H), 7.60 (d, *J* = 8.51 Hz, 1H), 7.38 (s, 2H), 7.34 (d, *J* = 10.56 Hz, 3H), 7.25 (s, 2H), 6.77 (dd, *J* = 1.32, 8.36 Hz, 1H), 5.54 (br. s, 2H).

Compound 148


N-(5-chloro-2-[[1-(1-oxido-2-pyridin-2-yl)methyl]sulfinyl]phenyl)-1-benzofuran-2-sulfonamide



Following **General Procedure D**, the title compound was prepared from N-{5-chloro-2-[(pyridin-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide
¹H NMR (600 MHz, METHANOL-*d*₄) δ 8.31 (d, *J* = 6.16 Hz, 1H), 7.66 (d, *J* = 7.92 Hz, 1H), 7.46 - 7.50 (m, 1H), 7.38 - 7.42 (m, 1H), 7.30 - 7.38 (m, 2H), 7.28 (s, 3H), 7.19 (d, *J* = 8.51 Hz, 1H), 7.05 (d, *J* = 7.63 Hz, 1H), 6.82 (d, *J* = 8.22 Hz, 1H), 5.14 (d, *J* = 12.62 Hz, 1H), 4.68 (d, *J* = 12.32 Hz, 1H).

Compound 149

N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure L**, the title compound was prepared from N-{5-chloro-2-[(3-nitrobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, CHLOROFORM-d) δ 7.64 - 7.68 (m, 2H), 7.47 - 7.51 (m, 1H), 7.40 - 7.46 (m, 2H), 7.29 - 7.34 (m, 1H), 7.14 (d, *J* = 8.22 Hz, 1H), 6.97 (t, *J* = 7.78 Hz, 1H),
5 6.92 (dd, *J* = 2.35, 8.22 Hz, 1H), 6.48 - 6.52 (m, 1H), 6.35 (dd, *J* = 0.59, 7.63 Hz, 1H),
6.28 (t, *J* = 1.91 Hz, 1H), 3.67 (s, 2H).

Compound 150

N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide




10 Following **General Procedure C and K**, the title compound was prepared from N-{5-chloro-2-[(3-nitrobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, METHANOL-d₄) δ 7.69 (d, *J* = 7.92 Hz, 1H), 7.48 (d, *J* = 8.51 Hz, 1H), 7.37 - 7.43 (m, 3H), 7.30 (td, *J* = 0.88, 7.48 Hz, 1H), 7.20 (d, *J* = 8.51 Hz, 1H), 7.01 (dd, *J* = 2.05, 8.51 Hz, 1H), 6.95 (t, *J* = 7.78 Hz, 1H), 6.70 - 6.74 (m, 1H), 6.64 (t, *J* = 1.76 Hz, 1H), 6.45 (d, *J* = 7.63 Hz, 1H), 4.40 (d, *J* = 12.62 Hz, 1H), 4.00 (d, *J* =
15 12.62 Hz, 1H).


Compound 151

N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide




20 Following **General Procedure D and K**, the title compound was prepared from N-{5-chloro-2-[(3-nitrobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, METHANOL-d₄) δ 7.80 (d, *J* = 2.05 Hz, 1H), 7.69 (d, *J* = 7.63 Hz, 1H), 7.52 (d, *J* = 0.59 Hz, 1H), 7.40 (d, *J* = 8.51 Hz, 1H), 7.35 (td, *J* = 1.32, 7.70 Hz, 1H), 7.31 (d, *J* = 8.51 Hz, 1H), 7.26 - 7.30 (m, *J* = 7.92 Hz, 1H), 6.78 (dd, *J* = 2.05, 8.51 Hz, 1H), 6.65 (t, *J* = 7.78 Hz, 1H), 6.49 (dd, *J* = 1.91, 7.78 Hz, 1H), 6.33 (t, *J* = 1.76 Hz, 1H), 6.05 (d, *J* = 7.34 Hz, 1H), 4.66 (s, 2H).

Compound 152**N-{5-chloro-2-[(3-hydroxybenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide**


To a solution of N-{5-chloro-2-[(3-methoxybenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide (315 mg, 0.69 mmol) in CH₂Cl₂ (5 ml) at room temperature was added BBr₃ (1M solution in CH₂Cl₂, 2.1 ml, 2.1 mmol) and the reaction was stirred for 2 h, diluted with EtOAc, washed with brine, dried over Na₂SO₄, concentrated in *vacuo*. The crude product was purified by column chromatography on silica gel (25% ethyl acetate in hexane) to give the title compound (196 mg, 64%).

¹H NMR (600 MHz, CHLOROFORM-d) δ 8.10 (br. s., 1H), 7.63 - 7.69 (m, 2H), 7.41 - 7.51 (m, 3H), 7.32 (td, *J* = 0.88, 7.48 Hz, 1H), 7.12 (d, *J* = 8.22 Hz, 1H), 7.05 (t, *J* = 7.92 Hz, 1H), 6.91 (dd, *J* = 2.35, 8.22 Hz, 1H), 6.66 (ddd, *J* = 0.73, 2.57, 8.14 Hz, 1H), 6.51 (d, *J* = 7.92 Hz, 1H), 6.45 - 6.48 (m, 1H), 5.07 (br. s., 1H), 3.71 (s, 2H).

Compound 153**N-{5-chloro-2-[(3-hydroxybenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide**


Following **General Procedure C**, the title compound was prepared from N-{5-chloro-2-[(3-hydroxybenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide

¹H NMR (600 MHz, CHLOROFORM-d) δ 10.27 - 11.14 (m, 1H), 7.73 (d, *J* = 2.05 Hz, 1H), 7.67 (d, *J* = 7.92 Hz, 1H), 7.53 (s, 1H), 7.43 - 7.47 (m, 1H), 7.38 - 7.42 (m, 1H), 7.28 - 7.33 (m, 1H), 7.04 (t, *J* = 7.92 Hz, 1H), 6.92 (dd, *J* = 1.47, 8.22 Hz, 1H), 6.79 (d, *J* = 8.22 Hz, 1H), 6.74 (dd, *J* = 2.05, 8.22 Hz, 1H), 6.59 (s, 1H), 6.40 (d, *J* = 7.34 Hz, 1H), 4.33 (d, *J* = 12.32 Hz, 1H), 4.16 (d, *J* = 12.62 Hz, 1H).


Compound 154**N-{5-chloro-2-[(3-hydroxybenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide**

Following **General Procedure D**, the title compound was prepared from N-{5-chloro-2-[(3-hydroxybenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide

¹H NMR (600 MHz, CHLOROFORM-d) δ 7.76 (br. s., 1H), 7.65 - 7.72 (m, 1H), 7.56 (br. s., 1H), 7.44 (br. s., 3H), 7.33 (s, 1H), 7.02 (br. s., 2H), 6.77 (br. s., 1H), 6.58 (br. s., 1H), 6.39 - 6.47 (m, 1H), 4.28 (br. s., 2H).

Intermediate 37**benzofuran-2-sulfonic acid (5-chloro-2-mercapto-phenyl)-amide**

Following **General Procedure N**, the title compound was prepared from N,N'-[dithiobis(5-chloro-2,1-phenylene)]bis(1-benzofuran-2-sulfonamide) **Intermediate 8** and was used without further purification.


Compound 155**N-{5-chloro-2-[(trifluoromethyl)thio]phenyl}-1-benzofuran-2-sulfonamide**

To a solution of benzofuran-2-sulfonic acid (5-chloro-2-mercapto-phenyl)-amide (332 mg, 0.98 mmol) in CH₂Cl₂ (10 ml) at -78 °C was added 3,3-dimethyl-1-(trifluoromethyl)-1,2-benziodoxole (CAS #: 887144-97-0, 354 mg, 1.07 mmol) and the reaction was stirred for 1 h, and was concentrated *in vacuo*. The crude product was purified by column chromatography on silica gel (CH₂Cl₂) to give the title compound (375 mg, 94%).

¹H NMR (600 MHz, CHLOROFORM-d) δ 7.88 (s, 1H), 7.82 (d, *J* = 2.05 Hz, 1H), 7.67 (dd, *J* = 0.59, 7.92 Hz, 1H), 7.49 - 7.54 (m, 3H), 7.44 - 7.49 (m, 1H), 7.33 (td, *J* = 1.03, 7.56 Hz, 1H), 7.11 (dd, *J* = 2.20, 8.36 Hz, 1H).

Compound 156

N-{5-chloro-2-[(trifluoromethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure D** using 10 equivalents of mCPBA in refluxing CH₂Cl₂, the title compound was prepared from N-{5-chloro-2-[(trifluoromethyl)thio]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, acetone-d₆) δ 7.90 (d, *J* = 2.05 Hz, 1H), 7.78 (d, *J* = 8.80 Hz, 1H), 7.71 (d, *J* = 7.92 Hz, 1H), 7.51 (d, *J* = 8.22 Hz, 1H), 7.36 - 7.43 (m, 2H), 7.29 (t, *J* = 7.48 Hz, 1H), 6.92 (d, *J* = 6.46 Hz, 1H).

Compound 157

N-{5-chloro-2-[(trifluoromethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure D** using 10 equivalents of mCPBA in refluxing CH₂Cl₂, the title compound was prepared from N-{5-chloro-2-[(trifluoromethyl)thio]phenyl}-1-benzofuran-2-sulfonamide.

¹H NMR (600 MHz, acetone-d₆) δ 10.00 (br. s., 1H), 7.89 (d, *J* = 8.22 Hz, 1H), 7.78 - 7.81 (m, 1H), 7.63 - 7.66 (m, 1H), 7.50 - 7.59 (m, 4H), 7.39 (ddd, *J* = 0.88, 7.26, 8.00 Hz, 1H).

Compound 158

N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-2,4-difluorobenzenesulfonamide




Following **General Procedure A, B, and K**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 1-bromomethyl-3-nitro-benzene, and 2,4-difluorobenzenesulfonyl chloride.

- 5 ^1H NMR (600 MHz, CHLOROFORM- d) δ 7.92 (td, J = 6.16, 8.51 Hz, 1H), 7.47 (d, J = 2.35 Hz, 1H), 7.19 (d, J = 8.22 Hz, 1H), 7.01 - 7.06 (m, 1H), 6.95 - 7.01 (m, 1H), 6.88 - 6.94 (m, 2H), 6.57 (dd, J = 1.47, 8.80 Hz, 1H), 6.40 - 6.45 (m, 2H), 3.72 (s, 2H).

Compound 159

N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-2,4-difluorobenzenesulfonamide




10

Following **General Procedure A, B, D, and K**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 1-bromomethyl-3-nitro-benzene, and 2,4-difluorobenzenesulfonyl chloride.

- 15 ^1H NMR (600 MHz, METHANOL- d_4) δ 8.05 (td, J = 6.16, 8.51 Hz, 1H), 7.48 - 7.54 (m, 2H), 7.14 - 7.25 (m, 2H), 7.09 (dd, J = 2.05, 8.51 Hz, 1H), 6.96 (t, J = 7.78 Hz, 1H), 6.66 - 6.71 (m, 1H), 6.54 (t, J = 1.91 Hz, 1H), 6.34 (d, J = 7.63 Hz, 1H), 4.44 (s, 2H).

Compound 160

N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-4-chloro-2-fluorobenzenesulfonamide




20

Following **General Procedure A, B, D, and K**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 1-bromomethyl-3-nitro-benzene, and 4-chloro-2-fluorobenzenesulfonyl chloride.

¹H NMR (600 MHz, METHANOL-d₄) δ 7.97 (t, *J* = 8.51 Hz, 1H), 7.54 (d, *J* = 8.51 Hz, 1H), 7.43 - 7.52 (m, 3H), 7.13 - 7.17 (m, 1H), 6.98 (t, *J* = 7.78 Hz, 1H), 6.67 - 6.72 (m, 1H), 6.53 (s, 1H), 6.33 (d, *J* = 7.63 Hz, 1H), 4.42 (s, 2H).

Compound 161

N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-4-chloro-2-fluorobenzenesulfonamide




Following **General Procedure A, B, C, and K**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 1-bromomethyl-3-nitro-benzene, and 4-chloro-2-fluorobenzenesulfonyl chloride.

¹H NMR (600 MHz, METHANOL-d₄) δ 7.85 (t, *J* = 8.07 Hz, 1H), 7.49 (dd, *J* = 1.76, 9.98 Hz, 1H), 7.40 (dd, *J* = 1.91, 8.36 Hz, 1H), 7.33 (d, *J* = 8.22 Hz, 1H), 7.23 (dd, *J* = 1.91, 8.36 Hz, 1H), 7.16 (d, *J* = 2.05 Hz, 1H), 7.04 (t, *J* = 7.78 Hz, 1H), 6.72 - 6.75 (m, 1H), 6.63 (t, *J* = 1.91 Hz, 1H), 6.50 (d, *J* = 7.34 Hz, 1H), 4.30 (d, *J* = 12.91 Hz, 1H), 4.04 (d, *J* = 12.62 Hz, 1H).

Compound 162


N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}benzenesulfonamide



Following **General Procedure A, B, and K**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 1-bromomethyl-3-nitro-benzene, and benzenesulfonyl chloride.

¹H NMR (600 MHz, METHANOL-d₄) δ 7.75 - 7.79 (m, 2H), 7.57 - 7.62 (m, 1H), 7.48 - 7.53 (m, 2H), 7.44 (d, *J* = 2.05 Hz, 1H), 7.14 (d, *J* = 8.22 Hz, 1H), 6.99 (dd, *J* = 2.35, 8.22 Hz, 1H), 6.95 (t, *J* = 7.78 Hz, 1H), 6.56 - 6.60 (m, 1H), 6.45 (t, *J* = 1.91 Hz, 1H), 6.35 (d, *J* = 7.34 Hz, 1H), 3.58 (s, 2H).

5


Compound 163**N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-4-chlorobenzenesulfonamide**

Following **General Procedure A, B, and K**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 1-bromomethyl-3-nitro-benzene, and 4-chloro-benzenesulfonyl chloride.

10

¹H NMR (600 MHz, CHLOROFORM-d) δ 7.69 - 7.72 (m, 2H), 7.54 (d, *J* = 2.35 Hz, 1H), 7.40 - 7.44 (m, 2H), 7.17 (d, *J* = 8.22 Hz, 1H), 7.03 (t, *J* = 7.78 Hz, 1H), 6.95 (dd, *J* = 2.20, 8.36 Hz, 1H), 6.58 (dd, *J* = 1.76, 7.92 Hz, 1H), 6.34 - 6.39 (m, 2H), 3.61 (s, 2H).

15


Compound 164**N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-3-chlorobenzenesulfonamide**

Following **General Procedure A, B, and K**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 1-bromomethyl-3-nitro-benzene, and 3-chloro-benzenesulfonyl chloride.

20


¹H NMR (600 MHz, CHLOROFORM-d) δ 7.80 (t, *J* = 1.91 Hz, 1H), 7.63 - 7.66 (m, 1H), 7.56 (d, *J* = 2.35 Hz, 1H), 7.52 (ddd, *J* = 0.88, 1.98, 8.00 Hz, 1H), 7.39 (t, *J* = 7.92 Hz, 1H), 7.16 (d, *J* = 8.51 Hz, 1H), 7.02 (t, *J* = 7.78 Hz, 1H), 6.96 (dd, *J* = 2.05, 8.22 Hz, 1H), 6.55 - 6.59 (m, 1H), 6.35 (d, *J* = 7.63 Hz, 1H), 6.32 (t, *J* = 1.91 Hz, 1H), 3.56 (s, 2H).

25

Compound 165**N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}benzenesulfonamide**

Following **General Procedure A, B, D, and K**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 1-bromomethyl-3-nitro-benzene, and benzenesulfonyl chloride.


¹H NMR (600 MHz, METHANOL-d₄) δ 7.90 - 7.93 (m, 2H), 7.63 - 7.68 (m, 1H), 7.56 - 7.62 (m, 3H), 7.42 (d, *J* = 8.51 Hz, 1H), 7.11 (dd, *J* = 2.05, 8.51 Hz, 1H), 6.93 (t, *J* = 7.78 Hz, 1H), 6.66 (ddd, *J* = 0.73, 2.27, 8.14 Hz, 1H), 6.39 (t, *J* = 1.91 Hz, 1H), 6.16 (d, *J* = 7.34 Hz, 1H), 4.18 (s, 2H).

Compound 166**N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}benzenesulfonamide**

Following **General Procedure A, B, C, and K**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 1-bromomethyl-3-nitro-benzene, and benzenesulfonyl chloride.

¹H NMR (600 MHz, METHANOL-d₄) δ 7.82 - 7.86 (m, 2H), 7.64 - 7.69 (m, 1H), 7.56 - 7.62 (m, 2H), 7.27 - 7.31 (m, 1H), 7.22 - 7.27 (m, 1H), 7.08 (d, *J* = 2.05 Hz, 1H), 6.99 (t, *J* = 7.78 Hz, 1H), 6.67 (ddd, *J* = 0.88, 2.35, 8.22 Hz, 1H), 6.50 (t, *J* = 1.91 Hz, 1H), 6.36 (d, *J* = 7.63 Hz, 1H), 4.12 (d, *J* = 12.62 Hz, 1H), 3.95 (d, *J* = 12.91 Hz, 1H).

Compound 167**N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-4-chlorobenzenesulfonamide**




Following **General Procedure A, B, D, and K**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 1-bromomethyl-3-nitro-benzene, and 4-chloro-benzenesulfonyl chloride.

- 5 ^1H NMR (600 MHz, METHANOL- d_4) δ 7.85 - 7.88 (m, 2H), 7.58 - 7.62 (m, 2H), 7.55 (d, J = 2.05 Hz, 1H), 7.49 (d, J = 8.51 Hz, 1H), 7.15 (dd, J = 1.91, 8.66 Hz, 1H), 6.95 (t, J = 7.78 Hz, 1H), 6.68 (ddd, J = 0.88, 2.20, 8.07 Hz, 1H), 6.45 (t, J = 1.91 Hz, 1H), 6.21 (d, J = 7.63 Hz, 1H), 4.28 (s, 2H).

Compound 168

- 10 **N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-4-chlorobenzenesulfonamide**




Following **General Procedure A, B, C, and K**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 1-bromomethyl-3-nitro-benzene, and 4-chloro-benzenesulfonyl chloride.

- 15 ^1H NMR (600 MHz, METHANOL- d_4) δ 7.79 (d, J = 8.51 Hz, 2H), 7.59 (d, J = 8.80 Hz, 2H), 7.32 (d, J = 8.51 Hz, 1H), 7.24 - 7.29 (m, 1H), 7.06 (d, J = 1.76 Hz, 1H), 6.99 (t, J = 7.78 Hz, 1H), 6.65 - 6.69 (m, 1H), 6.52 (s, 1H), 6.38 (d, J = 7.34 Hz, 1H), 4.16 (d, J = 12.91 Hz, 1H), 3.97 (d, J = 12.91 Hz, 1H).

Compound 169

- 20 **N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-3-chlorobenzenesulfonamide**




Following **General Procedure A, B, D, and K**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 1-bromomethyl-3-nitro-benzene, and 3-chloro-benzenesulfonyl chloride.

¹H NMR (600 MHz, METHANOL-d₄) δ 8.00 (t, *J* = 1.91 Hz, 1H), 7.89 - 7.93 (m, 1H), 7.42 - 7.51 (m, 3H), 7.38 (d, *J* = 8.51 Hz, 1H), 6.89 (t, *J* = 7.78 Hz, 1H), 6.66 (dd, *J* = 1.76, 8.51 Hz, 1H), 6.59 (d, *J* = 8.22 Hz, 1H), 6.55 (s, 1H), 6.37 (d, *J* = 7.63 Hz, 1H), 4.69 (s, 2H).

Compound 170

N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-3-chlorobenzenesulfonamide




Following **General Procedure A, B, C, and K**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 1-bromomethyl-3-nitro-benzene, and 3-chlorobenzenesulfonyl chloride.

¹H NMR (600 MHz, METHANOL-d₄) δ 7.85 (s, 1H), 7.77 (d, *J* = 7.63 Hz, 1H), 7.51 - 7.55 (m, 1H), 7.47 (t, *J* = 7.92 Hz, 1H), 7.22 (d, *J* = 8.22 Hz, 1H), 7.19 (d, *J* = 1.76 Hz, 1H), 6.93 - 7.00 (m, 2H), 6.65 (dd, *J* = 2.20, 8.07 Hz, 1H), 6.58 (s, 1H), 6.43 (d, *J* = 7.63 Hz, 1H), 4.35 (d, *J* = 12.91 Hz, 1H), 3.82 (d, *J* = 12.91 Hz, 1H).

Compound 171

3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio}-N-isopropylpropanamide




Following **General Procedure H, B**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, N-isopropyl-acrylamide, and 4-chloro-3-trifluoromethyl-benzenesulfonyl chloride.

^1H NMR (300 MHz, CD_3OD) δ 8.19 (d, J = 1.47 Hz, 1H), 7.98 (d, J = 8.20 Hz, 1H), 7.60 (d, J = 8.20 Hz, 1H), 7.23 (d, J = 2.05 Hz, 1H), 7.06 (d, J = 8.20 Hz, 1H), 6.75 (dd, J = 2.20, 8.35 Hz, 1H), 3.85 - 4.03 (m, 1H), 3.00 (t, J = 7.33 Hz, 2H), 2.39 (t, J = 7.47 Hz, 2H), 1.12 (d, 6H).

5

Compound 172

3-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio]-N,N-dimethylpropanamide



Following **General Procedure H, B**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, N,N-dimethyl-acrylamide, and 4-chloro-3-trifluoromethyl-benzenesulfonyl chloride.


10

^1H NMR (300 MHz, CD_3OD) δ 8.11 (s, 1H), 7.90 (d, J = 7.91 Hz, 1H), 7.71 - 7.79 (m, 1H), 7.49 (d, J = 2.34 Hz, 1H), 7.39 (d, J = 8.50 Hz, 1H), 7.21 (none, J = 2.34, 8.50 Hz, 1H), 2.96 (d, J = 7.62 Hz, 6H), 2.90 (t, J = 7.03 Hz, 2H), 2.51 (t, J = 7.03 Hz, 2H).

15

Compound 173

3-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl]-N-isopropylpropanamide



isopropylpropanamide

Following **General Procedure C**, the title compound was prepared from 3-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio]-N,N-dimethylpropanamide

20


^1H NMR (300 MHz, CD_3OD) δ 8.17 (d, J = 1.76 Hz, 1H), 8.01 (d, J = 8.20 Hz, 1H), 7.67 (d, J = 8.50 Hz, 1H), 7.39 (d, J = 8.20 Hz, 1H), 7.24 (d, J = 1.76 Hz, 1H), 6.88

(dd, $J = 1.90, 8.35$ Hz, 1H), 3.91 (quin, $J = 6.59$ Hz, 1H), 3.37 - 3.52 (m, 1H), 3.09 - 3.25 (m, 1H), 2.58 (ddd, $J = 5.71, 9.89, 15.16$ Hz, 1H), 2.26 (ddd, $J = 5.86, 9.74, 15.16$ Hz, 1H), 1.10 (dd, 4H).

Compound 174

5

3-[[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino]phenyl]sulfinyl]propanamide



Following **General Procedure H, B**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, acrylamide, and 4-chloro-3-trifluoromethyl-benzenesulfonyl chloride.


10

^1H NMR (600 MHz, CD_3OD) δ 8.16 (d, $J = 1.76$ Hz, 1H), 8.00 (dd, $J = 2.05, 8.22$ Hz, 1H), 7.75 (d, $J = 8.51$ Hz, 1H), 7.57 (d, $J = 8.51$ Hz, 1H), 7.32 (s, 1H), 7.21 (d, $J = 7.92$ Hz, 1H), 3.26 (br. s., 1H), 3.18 (br. s., 1H), 2.99 - 3.07 (m, 3H), 2.95 (s, 3H), 2.88 (dt, $J = 7.37, 16.95$ Hz, 1H), 2.62 - 2.77 (m, 1H).

15

Compound 175

N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide




Following **General Procedure B, C**, the title compound was prepared from 5-chloro-2-(methylsulfinyl)aniline and benzenesulfonyl chloride.

20

^1H NMR (300 MHz, acetone- d_6) δ 10.64 (br. s., 1H), 7.93 (d, $J = 7.03$ Hz, 2H), 7.59 - 7.79 (m, 3H), 7.55 (d, $J = 1.76$ Hz, 1H), 7.45 (d, $J = 8.50$ Hz, 1H), 7.26 (dd, $J = 1.90, 8.35$ Hz, 1H), 2.74 (s, 3H).

Compound 176

N-[5-chloro-2-(methylthio)phenyl]thiophene-2-sulfonamide




Following **General Procedure B**, the title compound was prepared from 5-chloro-2-(methylsulfinyl)aniline and thiophene-2-sulfonyl chloride.

¹H NMR (300 MHz, CD₃OD) δ 7.75 (dd, *J* = 1.17, 4.98 Hz, 1H), 7.50 (dd, *J* = 1.32, 3.66 Hz, 1H), 7.40 (d, *J* = 2.05 Hz, 1H), 7.26 - 7.34 (m, 1H), 7.15 - 7.24 (m, 1H), 7.08 (dd, *J* = 3.81, 4.98 Hz, 1H), 2.23 (s, 3H).

Compound 177

N-[5-chloro-2-(methylsulfinyl)phenyl]thiophene-2-sulfonamide




Following **General Procedure C**, the title compound was prepared from N-[5-chloro-2-(methylthio)phenyl]thiophene-2-sulfonamide

¹H NMR (600 MHz, CD₃OD) δ 7.81 (d, *J* = 4.40 Hz, 1H), 7.75 (d, *J* = 8.51 Hz, 1H), 7.51 - 7.57 (m, 1H), 7.43 (d, *J* = 7.92 Hz, 1H), 7.15 (dd, *J* = 4.84, 8.66 Hz, 2H), 2.79 (s, 3H).

Compound 178

N-{2-[(3-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure D**, the title compound (136mg, 82%) was prepared from benzofuran-2-sulfonic acid [2-(3-nitro-benzylsulfonyl)-phenyl]-amide.

¹H NMR (600 MHz, acetone-d₆) δ 9.57 (br. s., 1H), 8.17 (dt, *J* = 2.53, 6.38 Hz, 1H), 7.93 (d, *J* = 0.88 Hz, 1H), 7.74 - 7.83 (m, 3H), 7.69 (td, *J* = 1.47, 7.92 Hz, 1H), 7.63

(dd, J = 1.47, 7.92 Hz, 1H), 7.52 - 7.59 (m, 3H), 7.47 (ddd, J = 1.17, 7.41, 8.44 Hz, 1H), 7.30 - 7.38 (m, 1H), 7.23 (t, J = 7.63 Hz, 1H), 4.85 (s, 2H).

Compound 179

N-(5-chloro-2-[[2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl]sulfinyl]phenyl)-1-benzofuran-2-sulfonamide




Following **General Procedure C**, the title compound (79mg, 73%) was prepared from N-(5-chloro-2-[[2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl]thio]phenyl)-1-benzofuran-2-sulfonamide.

1H NMR (600 MHz, CD₃OD) δ 7.61 (d, J = 7.92 Hz, 1H), 7.46 (d, J = 2.05 Hz, 1H), 7.44 (d, J = 8.22 Hz, 1H), 7.28 - 7.32 (m, 1H), 7.21 - 7.26 (m, 1H), 7.19 (dd, J = 0.73, 8.36 Hz, 1H), 7.15 (d, J = 0.88 Hz, 1H), 6.92 (dd, J = 1.91, 8.36 Hz, 1H), 3.52 - 3.62 (m, 1H), 2.80 - 2.91 (m, 2H), 2.56 - 2.65 (m, 1H), 2.11 (s, 6H).

Compound 180

4-chloro-N-{5-chloro-2-[(3-hydroxycyclopentyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide




Following **General Procedure D**, followed by treatment of the crude product with NaBH₄ in MeOH, the title compound was prepared from 4-Chloro-N-[5-chloro-2-(3-oxo-cyclopentylsulfonyl)-phenyl]-3-trifluoromethyl-benzenesulfonamide (**Intermediate 35**).

^1H NMR (600 MHz, CD_3OD) δ 8.34 (s, 1H), 8.18 (dd, J = 1.91, 8.36 Hz, 1H), 7.70 - 7.79 (m, 2H), 7.65 (s, 1H), 7.00 (br. s., 1H), 4.12 - 4.18 (m, 2H), 1.98 - 2.10 (m, 2H), 1.72 - 1.84 (m, 2H), 1.63 - 1.71 (m, 2H).

Compound 181

5 N-[5-chloro-2-(methylthio)phenyl]-4-methyl-3-nitrobenzenesulfonamide




Following **General Procedure B**, the title compound (197 mg, 61%) was prepared from 5-Chloro-2-methylsulfanyl-phenylamine (150 mg, 0.87 mmol) and 4-Methyl-3-nitro-benzenesulfonyl chloride (235 mg, 0.87 mmol).

10 ^1H NMR (600 MHz, CD_3OD) δ 8.29 (d, J = 1.76 Hz, 1H), 7.85 (dd, J = 1.91, 8.07 Hz, 1H), 7.57 (d, J = 8.22 Hz, 1H), 7.39 (d, J = 2.05 Hz, 1H), 7.17 - 7.25 (m, 2H), 2.60 (s, 3H), 2.20 (s, 3H).

Compound 182

chloro-2-(methylsulfinyl)phenyl]-4-methyl-3-nitrobenzenesulfonamide



15


Following **General Procedure C**, the title compound was prepared from N-[5-chloro-2-(methylthio)phenyl]-4-methyl-3-nitrobenzenesulfonamide

^1H NMR (600 MHz, CDCl_3) δ 10.88 (br. s., 1H), 8.53 (d, J = 2.05 Hz, 1H), 8.05 (dd, J = 1.91, 8.07 Hz, 1H), 7.64 (d, J = 1.76 Hz, 1H), 7.54 (d, J = 8.22 Hz, 1H), 7.04 - 7.16 (m, 2H), 2.87 (s, 3H), 2.67 (s, 3H).

20

Compound 183

N-[5-chloro-2-(methylsulfonyl)phenyl]-4-methyl-3-nitrobenzenesulfonamide




Following **General Procedure D**, the title compound was prepared from N-[5-chloro-2-(methylthio)phenyl]-4-methyl-3-nitrobenzenesulfonamide

¹H NMR (600 MHz, CDCl₃) δ 9.36 (br. s., 1H), 8.52 (d, *J* = 2.05 Hz, 1H), 8.05 (dd, *J* = 2.05, 8.22 Hz, 1H), 7.79 (d, *J* = 8.51 Hz, 1H), 7.67 (d, *J* = 2.05 Hz, 1H), 7.58 (d, *J* = 7.92 Hz, 1H), 7.23 (dd, *J* = 1.76, 8.51 Hz, 1H), 3.04 (s, 3H), 2.68 (s, 3H).

Compound 184

4-chloro-N-[5-chloro-2-(methylthio)phenyl]benzenesulfonamide




Following **General Procedure B**, the title compound (150 mg, 68%) was prepared from 5-Chloro-2-methylsulfanyl-phenylamine (110 mg, 0.64 mmol) and 4-Chloro-benzenesulfonyl chloride (134mg, 0.64 mmol).

¹H NMR (600 MHz, CD₃OD) δ 7.72 (d, *J* = 9.10 Hz, 2H), 7.51 (d, *J* = 8.80 Hz, 2H), 7.37 (d, *J* = 2.05 Hz, 1H), 7.25 (d, *J* = 8.51 Hz, 1H), 7.19 (dd, *J* = 2.05, 8.22 Hz, 1H), 2.20 (s, 3H).

Compound 185

4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide




Following **General Procedure C**, the title compound was prepared from 4-chloro-N-[5-chloro-2-(methylthio)phenyl]benzenesulfonamide

^1H NMR (600 MHz, CDCl_3) δ 10.67 (br. s., 1H), 7.89 (d, J = 8.80 Hz, 2H), 7.66 (d, J = 1.76 Hz, 1H), 7.50 (d, J = 8.80 Hz, 2H), 6.91 - 7.16 (m, 2H), 2.80 (none, 3H).

Compound 186

4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]benzenesulfonamide



5


Following **General Procedure D**, the title compound was prepared from 4-chloro-N-[5-chloro-2-(methylthio)phenyl]benzenesulfonamide

^1H NMR (600 MHz, CDCl_3) δ 7.89 (d, J = 8.80 Hz, 2H), 7.78 (d, J = 8.51 Hz, 1H), 7.68 (d, J = 1.76 Hz, 1H), 7.53 (d, J = 8.80 Hz, 2H), 7.20 (dd, J = 2.05, 8.51 Hz, 1H), 2.96 (s, 3H).

10

Compound 187

4-chloro-N-[5-chloro-2-(methylthio)phenyl]-3-methylbenzenesulfonamide



Following **General Procedure B**, the title compound (205 mg, 59%) was prepared from 5-chloro-2-methylsulfanyl-phenylamine (165 mg, 0.95 mmol) and 4-chloro-3-methyl-benzenesulfonyl chloride (215mg, 0.95 mmol).


15

^1H NMR (600 MHz, CDCl_3) δ 7.71 (d, 1H), 7.65 (s, 1H), 7.62 (d, J = 2.05 Hz, 1H), 7.57 (dd, J = 2.05, 8.22 Hz, 1H), 7.41 (d, J = 8.22 Hz, 1H), 7.32 (d, J = 8.51 Hz, 1H), 7.04 (dd, J = 2.20, 8.36 Hz, 1H), 2.39 (s, 3H), 2.19 (s, 3H).

20

Compound 188

4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-3-methylbenzenesulfonamide




Following **General Procedure C**, the title compound was prepared from 4-chloro-N-[5-chloro-2-(methylthio)phenyl]-3-methylbenzenesulfonamide

¹H NMR (600 MHz, CDCl₃) δ 10.62 (s, 1H), 7.82 (d, *J* = 2.05 Hz, 1H), 7.71 (dd, *J* = 1.91, 8.36 Hz, 1H), 7.63 (d, *J* = 2.05 Hz, 1H), 7.48 (d, *J* = 8.51 Hz, 1H), 7.02 - 7.15 (m, 2H), 2.81 (s, 3H), 2.43 (s, 3H).

Compound 189

4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-3-methylbenzenesulfonamide




Following **General Procedure D**, the title compound was prepared from 4-chloro-N-[5-chloro-2-(methylthio)phenyl]-3-methylbenzenesulfonamide

¹H NMR (600 MHz, CDCl₃) δ 9.24 (br. s., 1H), 7.83 (d, *J* = 2.05 Hz, 1H), 7.78 (d, *J* = 8.51 Hz, 1H), 7.71 (dd, *J* = 2.05, 8.51 Hz, 1H), 7.67 (d, *J* = 1.76 Hz, 1H), 7.50 (d, *J* = 8.22 Hz, 1H), 7.19 (dd, *J* = 1.76, 8.51 Hz, 1H), 2.97 (s, 3H), 2.44 (s, 3H).

Compound 190

N-[5-chloro-2-(methylthio)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide




Following **General Procedure B**, the title compound (176 mg, 43%) was prepared from 5-chloro-2-methylsulfanyl-phenylamine (167 mg, 0.95 mmol) and 4-nitro-3-trifluoromethyl-benzenesulfonyl chloride (278mg, 0.95 mmol).

¹H NMR (300 MHz, CD₃OD) δ 8.04 - 8.22 (m, 3H), 7.42 (d, *J* = 2.05 Hz, 1H), 7.16 - 7.32 (m, 2H), 2.20 (s, 3H).

Compound 191

N-[5-chloro-2-(methylsulfonyl)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide




5 Following **General Procedure D**, the title compound was prepared from N-[5-chloro-2-(methylthio)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide ^1H NMR (600 MHz, acetone- d_6) δ 9.64 (br. s., 1H), 8.49 - 8.65 (m, 2H), 8.33 (d, J = 8.51 Hz, 1H), 7.89 (d, J = 8.51 Hz, 1H), 7.70 (d, J = 1.76 Hz, 1H), 7.39 (d, J = 8.22 Hz, 1H), 3.24 (s, 3H).

10

Compound 192


N-[5-chloro-2-(methylsulfinyl)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide



15 Following **General Procedure D**, the title compound was prepared from N-[5-chloro-2-(methylthio)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide ^1H NMR (600 MHz, acetone- d_6) δ 8.25 - 8.35 (m, 2H), 8.11 (d, J = 8.22 Hz, 1H), 7.39 - 7.48 (m, 2H), 6.75 (dd, J = 2.05, 8.22 Hz, 1H), 2.70 (s, 3H).

Compound 193

N-[5-chloro-2-(methylthio)phenyl]-2,4-difluorobenzenesulfonamide




20

Following **General Procedure B**, the title compound (358 mg, 94%) was prepared from 5-chloro-2-methylsulfonyl-phenylamine (189 mg, 1.09 mmol) and 2,4-difluoro-benzenesulfonyl chloride (231 mg, 1.09 mmol).

¹H NMR (300 MHz, CD₃OD) δ 7.82 (td, *J* = 6.15, 8.50 Hz, 1H), 7.34 (d, *J* = 2.05 Hz, 1H), 7.16 - 7.30 (m, 3H), 7.00 - 7.16 (m, 1H), 2.27 (s, 3H).

Compound 194

N-[5-chloro-2-(methylsulfonyl)phenyl]-2,4-difluorobenzenesulfonamide




Following **General Procedure D**, the title compound was prepared from N-[5-chloro-2-(methylthio)phenyl]-2,4-difluorobenzenesulfonamide

¹H NMR (300 MHz, acetone-d₆) δ 9.81 (br. s., 1H), 8.15 - 8.32 (m, 1H), 7.91 (d, *J* = 8.50 Hz, 1H), 7.63 (d, *J* = 1.76 Hz, 1H), 7.24 - 7.46 (m, 3H), 3.27 (s, 3H).

Compound 195

N-[5-chloro-2-(methylsulfinyl)phenyl]-2,4-difluorobenzenesulfonamide




Following **General Procedure D**, the title compound was prepared from N-[5-chloro-2-(methylthio)phenyl]-2,4-difluorobenzenesulfonamide

¹H NMR (600 MHz, acetone-d₆) δ 10.97 (br. s., 1H), 8.10 (td, *J* = 6.16, 8.51 Hz, 1H), 7.51 (d, *J* = 8.22 Hz, 1H), 7.46 (d, *J* = 2.05 Hz, 1H), 7.30 - 7.38 (m, 1H), 7.24 - 7.31 (m, 2H), 2.90 (s, 3H).

Compound 196

N-[5-chloro-2-(methylthio)phenyl]-5-methylfuran-2-sulfonamide




Following **General Procedure B**, the title compound (212 mg, 48%) was prepared from 5-chloro-2-methylsulfanyl-phenylamine (189 mg, 1.09 mmol) and 5-methyl-furan-2-sulfonyl chloride (250 mg, 1.38 mmol).

- 5 ^1H NMR (300 MHz, acetone- d_6) δ 8.53 (br. s., 1H), 7.37 - 7.49 (m, 2H), 7.25 (dd, J = 2.34, 8.50 Hz, 1H), 6.96 (d, J = 3.22 Hz, 1H), 6.24 (d, J = 2.64 Hz, 1H), 2.38 (s, 3H), 2.34 (s, 3H).

Compound 197

N-[5-chloro-2-(methylsulfinyl)phenyl]-5-methylfuran-2-sulfonamide




10

Following **General Procedure C**, the title compound was prepared from N-[5-chloro-2-(methylthio)phenyl]-5-methylfuran-2-sulfonamide

- 15 ^1H NMR (600 MHz, acetone- d_6) δ 10.74 (br. s., 1H), 7.55 (d, J = 2.05 Hz, 1H), 7.51 (d, J = 8.22 Hz, 1H), 7.30 (dd, J = 1.91, 8.36 Hz, 1H), 7.18 (d, J = 3.23 Hz, 1H), 6.24 - 6.35 (m, 1H), 2.90 (s, 3H), 2.34 (s, 3H).


Compound 198

N-[5-chloro-2-(methylsulfonyl)phenyl]-5-methylfuran-2-sulfonamide




- 20 Following **General Procedure D**, the title compound was prepared from N-[5-chloro-2-(methylthio)phenyl]-5-methylfuran-2-sulfonamide

^1H NMR (600 MHz, acetone- d_6) δ 9.54 (br. s., 1H), 7.90 (d, J = 8.51 Hz, 1H), 7.73 (d, J = 2.05 Hz, 1H), 7.39 (dd, J = 1.76, 8.51 Hz, 1H), 7.31 (d, J = 3.52 Hz, 1H), 6.32 (d, J = 3.23 Hz, 1H), 3.24 (s, 3H), 2.33 (s, 3H).

Compound 199**N-[5-chloro-2-(methylthio)phenyl]furan-2-sulfonamide**


Following **General Procedure B**, the title compound (215 mg, 47%) was prepared from 5-chloro-2-methylsulfanyl-phenylamine (260 mg, 1.50 mmol) and furan-2-sulfonyl chloride (250mg, 1.50 mmol).

¹H NMR (300 MHz, acetone-d₆) δ 8.67 (br. s., 1H), 7.84 (s, 1H), 7.32 - 7.46 (m, 2H), 7.26 (dd, *J* = 2.20, 8.35 Hz, 1H), 7.07 (d, *J* = 3.52 Hz, 1H), 6.63 (dd, *J* = 1.76, 3.52 Hz, 1H), 2.36 (s, 3H).

Compound 200**N-[5-chloro-2-(methylsulfinyl)phenyl]furan-2-sulfonamide**

Following **General Procedure C**, the title compound was prepared from N-[5-chloro-2-(methylthio)phenyl]furan-2-sulfonamide

¹H NMR (600 MHz, acetone-d₆) δ 10.85 (br. s., 1H), 7.89 (s, 1H), 7.49 - 7.59 (m, 2H), 7.28 - 7.36 (m, 2H), 6.68 (dd, *J* = 1.76, 3.52 Hz, 1H), 2.88 (s, 3H).

Compound 201**N-[5-chloro-2-(methylsulfonyl)phenyl]furan-2-sulfonamide**


Following **General Procedure D**, the title compound was prepared from N-[5-chloro-2-(methylthio)phenyl]furan-2-sulfonamide

^1H NMR (600 MHz, acetone- d_6) δ 9.62 (br. s., 1H), 7.90 (d, J = 8.51 Hz, 2H), 7.74 (d, J = 2.05 Hz, 1H), 7.27 - 7.48 (m, 2H), 6.70 (dd, J = 1.76, 3.52 Hz, 1H), 3.22 (s, 3H).

5

Compound 202

4-chloro-N-[5-chloro-2-(methylthio)phenyl]-2-fluorobenzenesulfonamide



Following **General Procedure B**, the title compound (380 mg, 66%) was prepared from 5-chloro-2-methylsulfanyl-phenylamine (273 mg, 1.57 mmol) and 4-chloro-2-fluoro-benzenesulfonyl chloride (360mg, 1.57 mmol).


10

^1H NMR (300 MHz, acetone- d_6) δ 8.77 (br. s., 1H), 7.84 (t, J = 8.06 Hz, 1H), 7.53 (dd, J = 1.90, 9.82 Hz, 1H), 7.35 - 7.48 (m, 3H), 7.25 (dd, J = 2.20, 8.35 Hz, 1H), 2.34 (s, 3H).

Compound 203

15

4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-2-fluorobenzenesulfonamide




Following **General Procedure C**, the title compound was prepared from 4-chloro-N-[5-chloro-2-(methylthio)phenyl]-2-fluorobenzenesulfonamide

^1H NMR (300 MHz, acetone- d_6) δ 10.95 (br. s., 1H), 8.03 (t, J = 8.20 Hz, 1H), 7.39 - 7.61 (m, 4H), 7.28 (dd, J = 1.90, 8.35 Hz, 1H), 2.93 (s, 3H).

20

Compound 204

4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-2-fluorobenzenesulfonamide




Following **General Procedure D**, the title compound was prepared from 4-chloro-N-[5-chloro-2-(methylthio)phenyl]-2-fluorobenzenesulfonamide

¹H NMR (300 MHz, acetone-d₆) δ 9.84 (br. s., 1H), 8.12 (t, *J* = 8.20 Hz, 1H), 7.90 (d, *J* = 8.50 Hz, 1H), 7.46 - 7.69 (m, 3H), 7.33 (dd, *J* = 1.47, 8.50 Hz, 1H), 3.27 (s, 3H).

Compound 205

3-chloro-N-[5-chloro-2-(methylthio)phenyl]-2-fluorobenzenesulfonamide




Following **General Procedure B**, the title compound (386 mg, 62%) was prepared from 5-chloro-2-methylsulfanyl-phenylamine (297 mg, 1.71 mmol) and 3-chloro-2-fluoro-benzenesulfonyl chloride (392mg, 1.71 mmol).

¹H NMR (300 MHz, acetone-d₆) δ 8.87 (br. s., 1H), 7.70 - 7.93 (m, 2H), 7.32 - 7.47 (m, 3H), 7.20 - 7.31 (m, 1H), 2.32 (s, 3H).


Compound 206

3-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-2-fluorobenzenesulfonamide




Following **General Procedure D**, the title compound was prepared from 3-chloro-N-[5-chloro-2-(methylthio)phenyl]-2-fluorobenzenesulfonamide

¹H NMR (300 MHz, acetone-d₆) δ 7.89 - 8.04 (m, 1H), 7.69 (d, *J* = 8.50 Hz, 1H), 7.55 - 7.65 (m, 2H), 7.23 - 7.37 (m, 1H), 6.72 (dd, *J* = 1.90, 8.64 Hz, 1H), 3.22 (s, 3H).


Compound 207**3-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-2-fluorobenzenesulfonamide**

Following **General Procedure C**, the title compound was prepared from 3-chloro-N-[5-chloro-2-(methylthio)phenyl]-2-fluorobenzenesulfonamide
¹H NMR (300 MHz, acetone-d₆) δ 7.94 - 8.04 (m, 1H), 7.83 - 7.93 (m, 1H), 7.42 - 7.59 (m, 3H), 7.29 (dd, *J* = 1.90, 8.35 Hz, 1H), 2.93 (s, 3H).

Compound 208**4-chloro-N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide**

Following **General Procedure A, B**, the title compound was prepared from 2-Amino-4-chloro-benzenethiol, 2-chloromethyl-1H-imidazole and 4-chloro-3-trifluoromethyl-benzenesulfonyl chloride.

¹H NMR (300 MHz, CD₃OD) δ 8.06 (d, *J* = 1.76 Hz, 1H), 7.89 (dd, *J* = 2.05, 8.50 Hz, 1H), 7.63 - 7.78 (m, 2H), 7.45 (d, *J* = 2.05 Hz, 1H), 7.30 (d, *J* = 8.50 Hz, 1H), 7.14 (dd, *J* = 2.34, 8.50 Hz, 1H), 6.73 (s, 1H), 3.83 (s, 2H).


Compound 209**4-chloro-N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide**

Following **General Procedure C**, the title compound was prepared from 4-chloro-N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide.

¹H NMR (600 MHz, CD₃OD) δ 8.18 (d, 1H), 8.03 (dd, *J* = 1.91, 8.36 Hz, 1H), 7.72 (d, *J* = 8.22 Hz, 1H), 7.41 (s, 2H), 7.20 (d, *J* = 1.76 Hz, 1H), 7.06 (d, *J* = 8.51 Hz, 1H), 6.84 (dd, *J* = 1.91, 8.36 Hz, 1H), 4.76 (d, *J* = 8.22 Hz, 2H).

Compound 210


4-chloro-N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide



¹H NMR (600 MHz, acetone-d₆) δ 8.33 (d, *J* = 1.76 Hz, 1H), 8.18 (dd, *J* = 1.76, 8.22 Hz, 1H), 7.70 (d, *J* = 8.22 Hz, 1H), 7.66 (d, *J* = 8.51 Hz, 1H), 7.58 (d, *J* = 2.05 Hz, 1H), 7.10 (s, 2H), 6.72 (dd, *J* = 1.91, 8.36 Hz, 1H), 4.86 (s, 2H).

Compound 211

4-chloro-N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide




Following **General Procedure A, B**, the title compound was prepared from 2-Amino-4-chloro-benzenethiol, 4-chloromethyl-1H-imidazole hydrogen chloride and 4-chloro-3-trifluoromethyl-benzenesulfonyl chloride.

¹H NMR (600 MHz, acetone-d₆) δ 8.07 (d, *J* = 2.05 Hz, 1H), 7.89 - 7.97 (m, 2H), 7.77 (d, *J* = 8.51 Hz, 1H), 7.61 (d, *J* = 2.35 Hz, 1H), 7.56 (d, *J* = 8.22 Hz, 1H), 7.16 (dd, *J* = 2.35, 8.22 Hz, 1H), 7.02 (s, 1H), 3.87 (s, 2H).

Compound 212

4-chloro-N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide




Following **General Procedure C**, the title compound was prepared from 4-chloro-N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide.

¹H NMR (600 MHz, acetone-d₆) δ 8.26 (br. s., 1H), 8.17 (s, 1H), 8.08 (dd, J = 1.76, 8.51 Hz, 1H), 7.82 (d, J = 8.51 Hz, 1H), 7.47 (br. s., 1H), 7.38 (d, J = 8.51 Hz, 1H), 7.30 (s, 1H), 7.09 (d, J = 7.92 Hz, 1H), 4.48 (dd, J = 2.35, 13.79 Hz, 1H), 4.20 (d, J = 14.09 Hz, 1H).

Compound 213

N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure A, B**, the title compound was prepared from 2-Amino-4-chloro-benzenethiol, 2-chloromethyl-1H-imidazole and 1-benzofuran-2-sulfonyl chloride.

¹H NMR (600 MHz, acetone-d₆) δ 7.74 (d, J = 7.92 Hz, 1H), 7.64 (d, J = 2.35 Hz, 1H), 7.61 (d, J = 8.51 Hz, 1H), 7.53 (d, J = 8.51 Hz, 1H), 7.46 (td, J = 1.17, 7.78 Hz, 1H), 7.41 (s, 1H), 7.33 (t, J = 7.48 Hz, 1H), 7.17 (s, 2H), 7.09 (dd, J = 2.35, 8.22 Hz, 1H), 4.10 (s, 2H).

Compound 214

N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure C**, the title compound was prepared from N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide

¹H NMR (300 MHz, DMSO-d₆) δ 7.68 (d, J = 7.62 Hz, 1H), 7.51 - 7.61 (m, 3H), 7.41 (d, J = 2.05 Hz, 1H), 7.32 - 7.40 (m, 1H), 7.21 - 7.32 (m, 1H), 7.17 (s, 1H), 6.92 (d, J = 8.50 Hz, 1H), 6.74 (dd, J = 1.90, 8.06 Hz, 1H), 4.89 (d, J = 14.07 Hz, 1H), 4.56 (d, J = 13.77 Hz, 1H).

Compound 215

N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure D**, the title compound was prepared from N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide

¹H NMR (300 MHz, acetone-d₆) δ 7.69 - 7.79 (m, 3H), 7.66 (s, 2H), 7.46 - 7.54 (m, 1H), 7.41 (s, 2H), 7.24 - 7.34 (m, 1H), 6.88 (dd, J = 1.90, 8.64 Hz, 1H), 5.62 (s, 2H).

Compound 216

N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure C**, the title compound was prepared from N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide .

¹H NMR (300 MHz, acetone-d₆) δ 8.34 (s, 1H), 7.75 (d, J = 7.91 Hz, 1H), 7.60 (d, J = 2.05 Hz, 1H), 7.47 - 7.56 (m, 2H), 7.43 (t, J = 7.33 Hz, 1H), 7.24 - 7.38 (m, 3H), 7.03 (d, J = 10.55 Hz, 1H), 4.58 (d, J = 14.36 Hz, 1H), 4.24 (d, J = 13.77 Hz, 1H)

Compound 217

5 **N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide**




Following **General Procedure D**, the title compound was prepared from N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide

10 ¹H NMR (300 MHz, acetone-d₆) δ 8.20 (br. s., 1H), 7.66 - 7.81 (m, 3H), 7.35 - 7.53 (m, 4H), 7.23 - 7.35 (m, 1H), 6.95 (dd, J = 2.05, 8.50 Hz, 1H), 4.87 (s, 2H).

Compound 218

N-[2-({2-[(aminocarbonyl)amino]ethyl}thio)-5-chlorophenyl]-1-benzofuran-2-sulfonamide




15

Following **General Procedure A, B**, the title compound was prepared from 2-Amino-4-chloro-benzenethiol, (2-chloro-ethyl)-urea and 1-benzofuran-2-sulfonyl chloride.

1H NMR (300 MHz, acetone-d₆) δ 7.77 (d, J = 7.91 Hz, 1H), 7.45 - 7.66 (m, 5H), 7.31
20 - 7.42 (m, 1H), 7.22 (dd, J = 2.05, 8.50 Hz, 1H), 6.02 (br. s., 1H), 5.40 (br. s., 2H), 3.14 (q, J = 6.15 Hz, 2H), 2.84 (t, J = 6.45 Hz, 2H).

Compound 219

N-[2-({2-[(aminocarbonyl)amino]ethyl}sulfinyl)-5-chlorophenyl]-1-benzofuran-2-sulfonamide




Following **General Procedure C**, the title compound was prepared from N-[2-
 5 ({2-[(aminocarbonyl)amino]ethyl}thio)-5-chlorophenyl]-1-benzofuran-2-sulfonamide .
¹H NMR (600 MHz, CD₃OD) δ 7.68 (d, J = 7.92 Hz, 1H), 7.54 (dd, J = 5.28, 8.22 Hz,
 2H), 7.42 (t, J = 7.92 Hz, 1H), 7.37 (s, 1H), 7.27 - 7.34 (m, 2H), 7.10 (d, J = 6.75 Hz,
 1H), 3.58 - 3.67 (m, 1H), 3.44 - 3.53 (m, 1H), 3.36 - 3.43 (m, 1H), 2.98 - 3.09 (m, 1H).

Compound 220

10

N-[2-({2-[(aminocarbonyl)amino]ethyl}sulfonyl)-5-chlorophenyl]-1-benzofuran-2-sulfonamide




Following **General Procedure D**, the title compound was prepared from N-[2-
 15 ({2-[(aminocarbonyl)amino]ethyl}thio)-5-chlorophenyl]-1-benzofuran-2-sulfonamide .
¹H NMR (600 MHz, CD₃OD) δ 7.85 (d, J = 8.51 Hz, 1H), 7.80 (d, J = 2.05 Hz, 1H),
 7.77 (d, J = 7.92 Hz, 1H), 7.72 (s, 1H), 7.61 (d, J = 8.51 Hz, 1H), 7.52 (t, J = 7.34 Hz,
 1H), 7.31 - 7.40 (m, 2H), 3.36 - 3.47 (m, 4H).

Compound 221

20

N-(5-chloro-2-{[3-(dimethylamino)propyl]thio}phenyl)-1-benzofuran-2-sulfonamide




Following **General Procedure A, B**, the title compound was prepared from 2-Amino-4-chloro-benzenethiol, (3-chloro-propyl)-dimethyl-amine hydrochloride and 1-benzofuran-2-sulfonyl chloride.

- 5 ^1H NMR (600 MHz, acetone- d_6) δ 7.62 - 7.72 (m, 2H), 7.49 (d, J = 8.22 Hz, 1H), 7.31 - 7.38 (m, 2H), 7.25 (t, J = 7.48 Hz, 1H), 7.18 (s, 1H), 6.69 (dd, J = 2.05, 8.22 Hz, 1H), 3.44 (t, J = 5.58 Hz, 2H), 3.30 (s, 3H), 3.12 (s, 3H), 2.97 (t, J = 6.31 Hz, 2H), 1.96 - 2.10 (m, 2H)

Compound 222

- 10 **N-(5-chloro-2-[[3-(dimethylamino)propyl]sulfonyl]phenyl)-1-benzofuran-2-sulfonamide**




Following **General Procedure C**, the title compound was prepared from N-(5-chloro-2-[[3-(dimethylamino)propyl]thio]phenyl)-1-benzofuran-2-sulfonamide .

- 15 ^1H NMR (600 MHz, acetone- d_6) δ 7.67 (d, J = 7.92 Hz, 1H), 7.60 (d, J = 1.76 Hz, 1H), 7.50 (d, J = 8.22 Hz, 1H), 7.41 (d, J = 8.22 Hz, 1H), 7.33 (t, J = 7.92 Hz, 1H), 7.20 - 7.28 (m, 2H), 6.83 (dd, J = 1.76, 8.22 Hz, 1H), 3.69 (ddd, J = 4.99, 8.58, 13.43 Hz, 1H), 3.61 (ddd, J = 4.11, 10.71, 14.23 Hz, 1H), 3.48 (ddd, J = 5.58, 5.72, 13.35 Hz, 1H), 3.07 - 3.21 (m, 7H), 2.30 - 2.42 (m, 1H), 2.16 - 2.30 (m, 1H).

Compound 223

- 20 **N-(5-chloro-2-[[3-(dimethylamino)propyl]sulfonyl]phenyl)-1-benzofuran-2-sulfonamide**




Following **General Procedure D**, the title compound was prepared from N-(5-chloro-2-[[3-(dimethylamino)propyl]thio]phenyl)-1-benzofuran-2-sulfonamide .

¹H NMR (600 MHz, acetone-d₆) δ 7.63 (d, J = 7.63 Hz, 1H), 7.53 (d, J = 1.76 Hz, 1H),
 5 7.48 (d, J = 8.22 Hz, 1H), 7.43 (d, J = 8.22 Hz, 1H), 7.30 (t, J = 7.34 Hz, 1H), 7.21 (t, J = 7.21 Hz, 1H), 7.17 (s, 1H), 6.74 (dd, J = 2.05, 8.22 Hz, 1H), 3.60 – 3.69 (m, 2H),
 3.36 - 3.48 (m, 1H), 3.20 - 3.33 (m, 6H), 2.90 - 3.01 (br. s., 1H), 2.29 - 2.42 (m, J = 6.75 Hz, 1H), 2.08 - 2.21 (m, 1H).

Compound 224

10 **N-(5-chloro-2-[[3-(dimethylnitroroyl)propyl]sulfonyl]phenyl)-1-benzofuran-2-sulfonamide**




Following **General Procedure D**, the title compound was prepared from N-(5-chloro-2-[[3-(dimethylamino)propyl]thio]phenyl)-1-benzofuran-2-sulfonamide .

15 ¹H NMR (600 MHz, acetone-d₆) d 7.79 (s, 1H), 7.70 (d, J = 8.51 Hz, 1H), 7.64 (d, J = 7.63 Hz, 1H), 7.52 (d, J = 8.51 Hz, 1H), 7.29 - 7.38 (m, 2H), 7.19 - 7.28 (m, 1H), 6.74 (d, J = 8.51 Hz, 1H), 3.74 – 3.84 (m, 2H), 3.60 – 3.65 (m., 2H), 3.26 (s, 6H), 2.27 – 2.39(m, 2H).

Compound 225

20 **N-(5-chloro-2-[[[(2-oxo-1,3-oxazolidin-5-yl)methyl]sulfinyl]phenyl]-1-benzofuran-2-sulfonamide**




Following **General Procedure A, B, C**, the title compound was prepared from 2-Amino-4-chloro-benzenethiol, 5-chloromethyl-oxazolidin-2-one and 1-benzofuran-2-sulfonyl chloride.

- 5 ¹H NMR (600 MHz, acetone-d₆) δ 7.62 (dd, J = 7.92, 11.15 Hz, 1H), 7.44 - 7.54 (m, 2H), 7.26 - 7.44 (m, 3H), 7.20 (ddd, J = 7.48, 7.63, 11.30 Hz, 1H), 6.87 - 7.05 (m, 1H), 6.51 (br. s., 1H), 4.69 - 5.03 (m, 1H), 3.60 (q, J = 8.31 Hz, 1H), 3.45 - 3.55 (m, 1H), 3.23 - 3.45 (m, 1H), 3.06 - 3.17 (m, 1H).

Compound 226

10

N-(5-chloro-2-[(2-oxo-1,3-oxazolidin-5-yl)methyl]sulfonyl)phenyl)-1-benzofuran-2-sulfonamide




Following **General Procedure D**, the title compound was prepared from N-(5-chloro-2-[(2-oxo-1,3-oxazolidin-5-yl)methyl]thio)phenyl)benzofuran-2-sulfonamide.

- 15 ¹H NMR (600 MHz, acetone-d₆) δ 7.81 (d, J = 1.76 Hz, 1H), 7.68 (d, J = 8.22 Hz, 2H), 7.54 (d, J = 8.22 Hz, 1H), 7.29 - 7.40 (m, 2H), 7.22 - 7.29 (m, 1H), 6.79 (br. s., 1H), 6.47 (br. s., 1H), 4.94 (dt, J = 6.93, 14.01 Hz, 1H), 4.17 (dd, J = 5.58, 14.09 Hz, 1H), 3.96 (dd, J = 6.90, 14.23 Hz, 1H), 3.65 (t, J = 8.66 Hz, 1H), 3.32 - 3.47 (m, 1H).

20

Compound 227

N-(5-chloro-2-[(2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl]thio)phenyl)-1-benzofuran-2-sulfonamide




Following **General Procedures A and B** the title compound was prepared from 2-amino-4-chloro-benzenethiol, 6-chloromethyl-1H-pyrimidine-2,4-dione and 1-benzofuran-2-sulfonyl chloride.

- 5 ^1H NMR (600 MHz, CD_3OD) δ 7.72 (d, J = 7.92 Hz, 1H), 7.57 (d, J = 8.22 Hz, 1H), 7.45 - 7.51 (m, 1H), 7.44 (d, J = 2.05 Hz, 1H), 7.42 (s, 1H), 7.27 - 7.37 (m, 2H), 7.09 (br. s., 1H), 5.05 (s, 1H), 3.64 (s, 2H).

Compound 228

N-(5-chloro-2-[(2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl]sulfinyl)phenyl)-1-benzofuran-2-sulfonamide

10




Following **General Procedure C**, the title compound was prepared from N-(5-chloro-2-[(2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl]thio)phenyl)-1-benzofuran-2-sulfonamide.

- 15 ^1H NMR (600 MHz, CD_3OD) δ 7.68 (d, J = 7.92 Hz, 1H), 7.52 (d, J = 8.51 Hz, 1H), 7.49 (d, J = 8.51 Hz, 1H), 7.41 (t, J = 7.92 Hz, 1H), 7.32 - 7.37 (m, 2H), 7.27 - 7.32 (m, 1H), 7.09 (d, J = 8.22 Hz, 1H), 5.32 (s, 1H), 4.40 (d, J = 13.21 Hz, 1H), 3.89 (d, J = 12.91 Hz, 1H).

Compound 229

N-(5-chloro-2-[(2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl]sulfonyl)phenyl)-1-benzofuran-2-sulfonamide

20




Following **General Procedure D**, the title compound was prepared from N-(5-chloro-2-[(2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl]thio)phenyl)-1-benzofuran-2-sulfonamide.

- 5 ^1H NMR (600 MHz, acetone- d_6) δ 10.36 (br. s., 1H), 10.15 (br. s., 1H), 7.78 (br. s., 1H), 7.70 (d, J = 8.51 Hz, 1H), 7.65 (d, J = 7.92 Hz, 1H), 7.42 - 7.51 (m, 2H), 7.35 (t, J = 7.63 Hz, 1H), 7.26 (t, J = 7.48 Hz, 1H), 6.87 (d, J = 6.75 Hz, 1H), 5.48 (br. s., 1H), 4.76 (br. s., 2H).

Compound 230

- 10 **N-{2-[(3-aminopropyl)thio]-5-chlorophenyl}-1-benzofuran-2-sulfonamide**




Following **General Procedure F, G**, the title compound was prepared from 2,2'-dithiobis(5-chloroaniline), 1-benzofuran-2-sulfonyl chloride and 3-bromo-propylamine hydrochloride.

- 15 ^1H NMR (600 MHz, DMSO- d_6) δ 8.03 (br. s., 2H), 7.69 (d, J = 7.63 Hz, 1H), 7.56 (d, J = 8.22 Hz, 1H), 7.37 (t, J = 7.78 Hz, 1H), 7.23 - 7.32 (m, 2H), 7.05 - 7.20 (m, 2H), 6.60 (br. s., 1H), 3.03 (br. s., 2H), 2.91 (t, J = 6.60 Hz, 2H), 1.74 - 1.92 (m, 2H).

Intermediate 38

(3-Bromo-propyl)-carbamic acid tert-butyl ester



20

A mixture of 3-bromo-propylamine hydrochloride (523 mg, 2.39 mmol), di-tert-butyl dicarbonate (573 mg, 2.63 mmol), NaOH (1N, 1.5 ml) in MeOH was stirred at rt overnight. The mixture was added water and extracted with EtOAc. The organic layer


was washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by column chromatography in silica gel.

¹H NMR (600 MHz, CDCl₃) δ 4.65 (br. s., 1H), 3.38 - 3.56 (m, 2H), 3.28 (d, *J* = 5.28 Hz, 2H), 1.95 - 2.14 (m, 2H), 1.45 (br. s., 9H).

5

Compound 231

N-{2-[(3-aminopropyl)sulfinyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure F, G, C, E** the title compound was prepared from 2,2'-dithiobis(5-chloroaniline), 1-benzofuran-2-sulfonyl chloride and (3-bromo-
10 propyl)-carbamic acid tert-butyl ester.

¹H NMR (600 MHz, acetone-d₆) δ 7.78 (d, *J* = 7.92 Hz, 1H), 7.63 (d, *J* = 8.51 Hz, 1H), 7.54 - 7.59 (m, 2H), 7.46 - 7.52 (m, 2H), 7.32 - 7.40 (m, 1H), 7.21 (d, *J* = 7.92 Hz, 1H), 4.03 - 4.14 (m, 1H), 3.88 - 4.00 (m, 1H), 3.35 (ddd, *J* = 6.97, 7.19, 13.72 Hz, 1H), 3.10 (ddd, *J* = 6.46, 6.60, 13.35 Hz, 1H), 2.16 - 2.36 (m, 2H).

15

Compound 232

N-{2-[(3-aminopropyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure F, G, D, E** the title compound was prepared from 2,2'-dithiobis(5-chloroaniline), 1-benzofuran-2-sulfonyl chloride and (3-bromo-
20 propyl)-carbamic acid tert-butyl ester.

¹H NMR (600 MHz, CD₃OD) δ 7.88 (d, *J* = 8.51 Hz, 1H), 7.79 (d, *J* = 7.92 Hz, 1H), 7.72 (d, *J* = 1.76 Hz, 1H), 7.68 (s, 1H), 7.61 (d, *J* = 8.51 Hz, 1H), 7.53 (t, *J* = 7.78 Hz, 1H), 7.40 (t, *J* = 7.63 Hz, 1H), 7.34 (d, *J* = 8.51 Hz, 1H), 3.47 (t, *J* = 7.04 Hz, 2H), 3.02 (t, *J* = 7.63 Hz, 2H), 2.06 (quin, 2H).

Compound 233

N-(5-chloro-2-[(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)methyl]sulfinyl)phenyl)-1-benzofuran-2-sulfonamide




5 Following **General Procedure A, B, C**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 5-(chloromethyl)uracil and 1-benzofuran-2-sulfonyl chloride.

¹H NMR (600 MHz, DMSO-d₆) δ 10.99 (s, 1H), 10.75 (br. s., 1H), 7.61 (d, J = 7.63 Hz, 1H), 7.51 (d, J = 8.22 Hz, 1H), 7.32 (t, J = 7.48 Hz, 1H), 7.16 - 7.27 (m, 3H), 7.01 - 7.10 (m, 1H), 6.86 (br. s., 1H), 6.64 - 6.79 (m, 1H), 4.02 (d, J = 13.21 Hz, 1H), 3.85 (d, J = 13.21 Hz, 1H).

Compound 234

N-(5-chloro-2-[(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)methyl]sulfonyl)phenyl)-1-benzofuran-2-sulfonamide




15 Following **General Procedure A, B, D**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 5-(chloromethyl)uracil and 1-benzofuran-2-sulfonyl chloride.

¹H NMR (600 MHz, CD₃OD) δ 7.69 (d, J = 2.05 Hz, 1H), 7.63 - 7.67 (m, 2H), 7.44 (t, J = 8.36 Hz, 1H), 7.32 - 7.39 (m, 2H), 7.27 (t, J = 7.04 Hz, 1H), 7.18 (s, 1H), 6.82 (d, J = 7.92 Hz, 1H), 4.70 (s, 2H).

Compound 235

N-{4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl]-1,3-thiazol-2-yl}acetamide




Following **General Procedure A, B**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, N-(4-Chloromethyl-thiazol-2-yl)-acetamide and 1-benzofuran-2-sulfonyl chloride.

- 5 ^1H NMR (600 MHz, CD_3OD) δ 7.70 (d, J = 7.92 Hz, 1H), 7.49 - 7.54 (m, 2H), 7.45 - 7.50 (m, 1H), 7.42 (s, 1H), 7.29 - 7.36 (m, 2H), 7.11 (dd, J = 2.20, 8.36 Hz, 1H), 6.45 (s, 1H), 3.84 (s, 2H), 2.21 (s, 3H).

Compound 236

N-{4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfinyl]methyl}-1,3-
thiazol-2-yl}acetamide

10




Following **General Procedure C**, the title compound was prepared from N-{4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl]-1,3-thiazol-2-yl}acetamide .

- 15 ^1H NMR (600 MHz, acetone- d_6) δ 10.93 (br. s., 1H), 7.83 (d, J = 7.92 Hz, 1H), 7.80 (s, 1H), 7.62 - 7.67 (m, 2H), 7.49 - 7.58 (m, 1H), 7.39 (t, J = 7.48 Hz, 1H), 7.08 - 7.22 (m, 2H), 6.74 (s, 1H), 4.29 - 4.46 (m, 2H), 2.23 (s, 3H).

Compound 237

N-{4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl]methyl}-
1,3-thiazol-2-yl}acetamide

20




Following **General Procedure D**, the title compound was prepared from N-4-[[{2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}thio)methyl]-1,3-thiazol-2-yl}acetamide.

- 5 ¹H NMR (600 MHz, acetone-d₆) δ 10.90 (br. s., 1H), 9.65 (br. s., 1H), 7.81 (d, J = 7.34 Hz, 2H), 7.75 (d, J = 2.05 Hz, 1H), 7.65 (d, J = 8.51 Hz, 1H), 7.60 (s., 1H), 7.51 (s., 1H), 7.38 (t, J = 7.19 Hz, 1H), 7.27 (s., 1H), 6.95 (s, 1H), 4.67 (br. s., 2H), 2.21 (s, 3H).

Compound 238

- 10 **N-(5-chloro-2-[[2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl]sulfonyl]phenyl)-1-benzofuran-2-sulfonamide**




Following **General Procedure D**, the title compound (53mg, 38%) was prepared from N-(5-chloro-2-[[2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl]thio}phenyl)-1-benzofuran-2-sulfonamide.

- 15 ¹H NMR (600 MHz, CD₃OD) δ 7.79 (d, J = 1.76 Hz, 1H), 7.76 (d, J = 8.51 Hz, 1H), 7.67 (d, J = 7.92 Hz, 1H), 7.57 (d, J = 0.59 Hz, 1H), 7.37 (ddd, J = 1.17, 7.19, 8.36 Hz, 1H), 7.26 - 7.31 (m, 1H), 7.20 - 7.24 (m, 1H), 7.15 (dd, J = 1.76, 8.51 Hz, 1H), 3.40 - 3.50 (m, 2H), 2.60 - 2.72 (m, 2H), 2.01 (s, 6H).

Compound 239


N-(5-fluoro-2-[(3-nitrobenzyl)sulfinyl]phenyl)-1-benzofuran-2-sulfonamide



Following **General Procedure C**, the title compound (174mg, 79%) was prepared from N-(5-fluoro-2-((3-nitrobenzyl)thio)phenyl)benzofuran-2-sulfonamide. ¹H NMR (600 MHz, CD₃OD) δ 7.98 (d, J = 9.10 Hz, 1H), 7.67 (d, J = 7.63 Hz, 1H), 7.49 (br. s., 1H), 7.40 (s, 1H), 7.35 (d, J = 8.51 Hz, 1H), 7.26 - 7.32 (m, 2H), 7.21 - 7.26 (m, 2H), 7.12 (d, J = 7.63 Hz, 1H), 6.74 - 6.80 (m, 1H), 6.40 (t, J = 7.34 Hz, 1H), 4.52 - 4.64 (m, 2H).

Compound 240

N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-2,4-difluorobenzenesulfonamide




Following **General Procedure A, B, C, and K**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 1-bromomethyl-3-nitro-benzene, and 2,4-difluoro-benzenesulfonyl chloride.

¹H NMR (600 MHz, METHANOL-d₄) δ 7.95 (d, J = 6.16 Hz, 1H), 7.19 - 7.30 (m, 3H), 7.11 - 7.19 (m, 2H), 7.00 (t, J = 7.78 Hz, 1H), 6.72 (dd, J = 1.47, 7.92 Hz, 1H), 6.61 (s, 1H), 6.44 (d, J = 7.04 Hz, 1H), 4.34 (d, J = 12.91 Hz, 1H), 4.07 (d, J = 12.90 Hz, 1H).

Compound 241

N-[5-chloro-2-(methylsulfonyl)phenyl]-4-isopropylbenzenesulfonamide




Following **General Procedure B and D**, the title compound was prepared from 5-chloro-2-(methylthio)aniline and 4-isopropyl-benzenesulfonyl chloride.

¹H NMR (600 MHz, CHLOROFORM-d) δ 9.13 (s, 1H), 7.80 - 7.85 (m, 2H), 7.71 - 7.76 (m, 2H), 7.34 - 7.39 (m, 2H), 7.16 (dd, *J* = 2.05, 8.51 Hz, 1H), 2.95 (spt, *J* = 6.90 Hz, 1H), 2.81 (s, 3H), 1.22 (d, *J* = 7.04 Hz, 6H).

Compound 242

N-(5-methoxy-2-((3-nitrobenzyl)thio)phenyl)benzofuran-2-sulfonamide




Following **General Procedure B**, the title compound (406 mg, 56%) was prepared from 5-methoxy-2-((3-nitrobenzyl)thio)aniline (450 mg, 1.552 mmol) and benzofuran-2-sulfonyl chloride (335 mg, 1.552 mmol) in pyridine (5 ml).

¹H NMR (600 MHz, CD₃OD) δ 8.53 (dd, *J* = 1.76, 5.87 Hz, 1H), 7.96 (ddd, *J* = 1.03, 2.27, 8.14 Hz, 1H), 7.70 (d, *J* = 7.92 Hz, 1H), 7.63 (t, *J* = 1.91 Hz, 1H), 7.51 (s, 1H), 7.42 (dd, *J* = 1.17, 8.22 Hz, 1H), 7.24 - 7.37 (m, 3H), 7.11 (s, 1H), 7.00 (s, 1H), 6.55 (dd, *J* = 2.79, 8.66 Hz, 1H), 3.83 (s, 2H), 3.71 (s, 3H).

Compound 243

N-{5-chloro-2-[(1H-pyrazol-3-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide




Following **General Procedure D and E**, the title compound was prepared from tert-butyl 3-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl]-1H-pyrazole-1-carboxylate (**Compound 22**).

^1H NMR (600 MHz, CD_3OD) δ 7.71 (d, J = 1.76 Hz, 1H), 7.66 (d, J = 7.92 Hz, 1H), 7.51 (d, J = 8.51 Hz, 1H), 7.31 - 7.43 (m, 4H), 7.24 - 7.29 (m, 1H), 6.79 (d, J = 7.04 Hz, 1H), 5.82 (br. s., 1H), 4.99 (br. s., 2H).

Compound 244

N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-4-chloro-2-fluorobenzenesulfonamide



Following **General Procedure A, B, and K**, the title compound was prepared from 2-amino-4-chloro-benzenethiol, 1-bromomethyl-3-nitro-benzene, and 4-chloro-2-fluoro-benzenesulfonyl chloride.

^1H NMR (600 MHz, CHLOROFORM-d) δ 7.84 (t, J = 8.22 Hz, 1H), 7.47 (d, J = 2.05 Hz, 1H), 7.25 (dt, J = 1.03, 8.51 Hz, 1H), 7.17 - 7.22 (m, 2H), 7.04 (t, J = 7.63 Hz, 1H), 6.93 (dd, J = 2.35, 8.22 Hz, 1H), 6.58 (dt, J = 1.17, 7.92 Hz, 1H), 6.40 - 6.45 (m, 2H), 3.72 (s, 2H).

Biological Data

HEK-Gqi5 cells stably expressing CCR2 were cultured in (DMEM high glucose, 10% FBS, 1% PSA, 400 $\mu\text{g/ml}$ geneticin and 50 $\mu\text{g/ml}$ hygromycin. Appropriate positive control chemokines (MCP-1, MIP1A or RANTES) was used as the positive control agonist for screening compound-induced calcium activity assayed on the FLIPR^{Tetra}. The drug plates were prepared in 384-well microplates using the EP3 and the MultiPROBE robotic liquid handling systems. Compounds were synthesized and tested for CCR2 activity.

Table 1 shows activity at CCR2 receptor (IC_{50} nM)

Table 1

Compound Name	CCR2 IC_{50} (nM)	CCR2 % ANTAGONISM
4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide	14	78
4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide	64	90

4-chloro-N-[5-chloro-2-(methylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide	134	82
4-chloro-N-[5-chloro-2-(isopropylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide	379	72
4-chloro-N-{5-chloro-2-[(2-hydroxyethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide	153	88
4-chloro-N-{5-chloro-2-[(2-hydroxyethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide	146	101
4-chloro-N-{5-chloro-2-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide	95	96
methyl 3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}propionate	108	82
4-chloro-N-{5-chloro-2-[(pyridin-2-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide	199	79
4-chloro-N-{5-chloro-2-[(pyridin-3-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide	94	85
4-chloro-N-{5-chloro-2-[(pyridin-2-ylmethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide	92	83
4-chloro-N-{5-chloro-2-[(pyridin-3-ylmethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide	112	97
ethyl {[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}acetate	141	79
3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio}-N-isopropylpropanamide	1033	59
3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio}-N,N-dimethylpropanamide	506	80
3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}-N-isopropylpropanamide	1788	93
3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}-N,N-dimethylpropanamide	96	89
3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}propanamide	797	99
N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide	NA	48
3,4-dichloro-N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide	44	87
N-[5-chloro-2-(methylthio)phenyl]thiophene-2-sulfonamide	NA	0
N-[5-chloro-2-(methylsulfinyl)phenyl]thiophene-2-sulfonamide	1113	53
4-chloro-N-{5-chloro-2-[(3-nitrophenyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide	509	85
4-chloro-N-{5-chloro-2-[(3-	107	101

hydroxycyclopentyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide		
4-chloro-N-{5-chloro-2-[(3-hydroxycyclopentyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide	76	97
4-chloro-N-[5-chloro-2-(ethylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide	193	81
4-chloro-N-[5-chloro-2-(ethylsulfinyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide	50	78
4-chloro-N-[5-chloro-2-(ethylsulfonyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide	88	90
N-[5-chloro-2-(methylthio)phenyl]-1-benzofuran-2-sulfonamide	26	99
N-[5-chloro-2-(methylsulfinyl)phenyl]-1-benzofuran-2-sulfonamide	11	98
N-[5-chloro-2-(methylsulfonyl)phenyl]-1-benzofuran-2-sulfonamide	42	93
N-[5-chloro-2-(methylthio)phenyl]-4-methyl-3-nitrobenzenesulfonamide	680	92
4-chloro-N-[5-chloro-2-(methylthio)phenyl]benzenesulfonamide	406	94
4-chloro-N-[5-chloro-2-(methylthio)phenyl]-3-methylbenzenesulfonamide	453	93
N-[5-chloro-2-(methylsulfinyl)phenyl]-4-methyl-3-nitrobenzenesulfonamide	181	101
N-[5-chloro-2-(methylsulfonyl)phenyl]-4-methyl-3-nitrobenzenesulfonamide	207	89
4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide	62	91
4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]benzenesulfonamide	155	98
4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-3-methylbenzenesulfonamide	44	88
2-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio}-N,N-dimethylacetamide	118	92
2-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}-N,N-dimethylacetamide	268	100
2-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfonyl}-N,N-dimethylacetamide	462	98
2-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}-N-methylacetamide	329	91
2-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfonyl}-N-methylacetamide	469	96
N-{2-[(2-aminoethyl)thio]-5-chlorophenyl}-4-chloro-3-	1067	97

(trifluoromethyl)benzenesulfonamide		
N-[5-chloro-2-(methylthio)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide	491	90
N-[5-chloro-2-(methylsulfonyl)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide	213	97
N-[5-chloro-2-(methylthio)phenyl]-2,4-difluorobenzenesulfonamide	NA	87
N-[5-chloro-2-(methylsulfonyl)phenyl]-2,4-difluorobenzenesulfonamide	354	91
N-[5-chloro-2-(methylsulfinyl)phenyl]-2,4-difluorobenzenesulfonamide	236	98
N-[5-chloro-2-(methylsulfinyl)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide	54	97
N-[5-chloro-2-(methylthio)phenyl]-5-methylfuran-2-sulfonamide	70	77
N-[5-chloro-2-(methylsulfinyl)phenyl]-5-methylfuran-2-sulfonamide	194	76
N-[5-chloro-2-(methylsulfonyl)phenyl]-5-methylfuran-2-sulfonamide	990	90
N-[5-chloro-2-(methylthio)phenyl]furan-2-sulfonamide	940	93
N-[5-chloro-2-(methylsulfinyl)phenyl]furan-2-sulfonamide	2318	96
N-[5-chloro-2-(methylsulfonyl)phenyl]furan-2-sulfonamide	345	99
4-chloro-N-[5-chloro-2-(methylthio)phenyl]-2-fluorobenzenesulfonamide	25	95
3-chloro-N-[5-chloro-2-(methylthio)phenyl]-2-fluorobenzenesulfonamide	1924	94
3-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-2-fluorobenzenesulfonamide	116	101
3-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-2-fluorobenzenesulfonamide	61	101
4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-2-fluorobenzenesulfonamide	171	98
4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-2-fluorobenzenesulfonamide	85	103
N-{2-[(2-aminoethyl)sulfinyl]-5-chlorophenyl}-4-chloro-3-(trifluoromethyl)benzenesulfonamide	3531	84
4-chloro-N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide	190	84
4-chloro-N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide	163	109
4-chloro-N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide	249	113
4-chloro-N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide	68	83

4-chloro-N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide	158	111
4-chloro-N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide	51	108
N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide	60	93
N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide	1898	108
N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide	1350	109
N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide	10	100
N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide	296	108
N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide	81	106
3-{2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}thio)-N,N-dimethylpropanamide	146	100
3-{2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfinyl)-N,N-dimethylpropanamide	160	104
3-{2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfonyl)-N,N-dimethylpropanamide	76	99
N-[2-{2-[(aminocarbonyl)amino]ethyl}thio]-5-chlorophenyl]-1-benzofuran-2-sulfonamide	80	103
N-[2-{2-[(aminocarbonyl)amino]ethyl}sulfinyl]-5-chlorophenyl]-1-benzofuran-2-sulfonamide	1891	64
N-[2-{2-[(aminocarbonyl)amino]ethyl}sulfonyl]-5-chlorophenyl]-1-benzofuran-2-sulfonamide	1995	68
tert-butyl {6-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfinyl)methyl}pyridin-2-yl}carbamate	527	82
N-(2-[(6-aminopyridin-2-yl)methyl]thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide	32	94
N-(2-[(6-aminopyridin-2-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide	28	99
N-(2-[(6-aminopyridin-2-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide	15	102
N-(2-[(6-amino-1-oxidopyridin-2-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide	264	95
N-(5-chloro-2-{[3-(dimethylamino)propyl]thio}phenyl)-1-benzofuran-2-sulfonamide	2043	106
N-(5-chloro-2-{[3-(dimethylamino)propyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide	NA	81
N-(5-chloro-2-{[3-(dimethylamino)propyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide	NA	96
N-(5-chloro-2-{[3-(dimethylnitro)propyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide	NA	83

N-(2-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfonyl]ethyl)acetamide	5190	98
N-(2-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl]ethyl)acetamide	NA	97
N-(5-chloro-2-{{(2-oxo-1,3-oxazolidin-5-yl)methyl}sulfinyl}phenyl)-1-benzofuran-2-sulfonamide	2128	79
N-(5-chloro-2-{{(2-oxo-1,3-oxazolidin-5-yl)methyl}sulfonyl}phenyl)-1-benzofuran-2-sulfonamide	2561	98
N-(5-chloro-2-{{(2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl}thio}phenyl)-1-benzofuran-2-sulfonamide	2368	98
N-(5-chloro-2-{{(2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl}sulfinyl}phenyl)-1-benzofuran-2-sulfonamide	NA	62
N-(5-chloro-2-{{(2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl}sulfonyl}phenyl)-1-benzofuran-2-sulfonamide	NA	70
N-{2-[(3-aminopropyl)thio]-5-chlorophenyl}-1-benzofuran-2-sulfonamide	NA	102
N-{2-[(3-aminopropyl)sulfinyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide	NA	63
N-{2-[(3-aminopropyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide	NA	82
4-{2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfinyl-N,N-dimethylbutanamide	NA	54
4-{2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfonyl-N,N-dimethylbutanamide	1003	98
5-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-1-benzofuran-2-sulfonamide	655	92
5-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-1-benzofuran-2-sulfonamide	2047	74
N-{5-chloro-2-[(1H-pyrazol-3-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide	16	99
N-{5-chloro-2-[(1H-pyrazol-3-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide	634	99
N-{5-chloro-2-[(1H-pyrazol-3-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide	367	102
N-(5-chloro-2-{{(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)methyl}sulfinyl}phenyl)-1-benzofuran-2-sulfonamide	3077	21
N-(5-chloro-2-{{(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)methyl}sulfonyl}phenyl)-1-benzofuran-2-sulfonamide	NA	0
N-{4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio]methyl}-1,3-thiazol-2-yl}acetamide	848	102
N-{4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfinyl]methyl}-1,3-thiazol-2-yl}acetamide	2685	79
N-{4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl]methyl}-1,3-thiazol-2-yl}acetamide	NA	87
N-[5-chloro-2-(methylsulfinyl)phenyl]-4-isopropylbenzenesulfonamide	934	68

4-bromo-N-[5-chloro-2-(methylthio)phenyl]benzenesulfonamide	1329	76
N-[5-chloro-2-(methylthio)phenyl]-4-iodobenzenesulfonamide	2032	71
N-[5-chloro-2-(methylsulfonyl)phenyl]-4-isopropylbenzenesulfonamide	NA	0
4-bromo-N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide	116	94
N-[5-chloro-2-(methylsulfinyl)phenyl]-4-iodobenzenesulfonamide	258	95
4-bromo-N-[5-chloro-2-(methylsulfonyl)phenyl]benzenesulfonamide	403	84
N-[5-chloro-2-(methylsulfonyl)phenyl]-4-iodobenzenesulfonamide	1068	92
tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio]methyl}-1,3-thiazol-2-yl}carbamate	2425	28
N-(2-[(2-amino-1,3-thiazol-4-yl)methyl]thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide	42	102
tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfinyl]methyl}-1,3-thiazol-2-yl}carbamate	2402	101
N-(2-[(2-amino-1,3-thiazol-4-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide	262	100
tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl]methyl}-1,3-thiazol-2-yl}carbamate	NA	72
N-(2-[(2-amino-1,3-thiazol-4-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide	83	102
N-{5-chloro-2-[(pyridin-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide	35	91
N-{5-chloro-2-[(pyridin-3-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide	22	97
N-{5-chloro-2-[(pyridin-3-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide	76	100
N-(5-chloro-2-[(1-oxidopyridin-3-yl)methyl]sulfinyl)phenyl)-1-benzofuran-2-sulfonamide	646	101
N-{5-chloro-2-[(pyridin-3-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide	32	93
N-(5-chloro-2-[(1-oxidopyridin-3-yl)methyl]sulfonyl)phenyl)-1-benzofuran-2-sulfonamide	1753	81
N-{5-chloro-2-[(3-nitrobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide	NA	63
N-{5-chloro-2-[(3-methoxybenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide	651	85
N-(5-chloro-2-[(5-nitro-1H-pyrazol-3-yl)methyl]thio)phenyl)-1-benzofuran-2-sulfonamide	NA	91
N-(5-chloro-2-[(5-nitro-1H-pyrazol-3-yl)methyl]sulfinyl)phenyl)-1-benzofuran-2-sulfonamide	752	94
N-(2-[(5-amino-1H-pyrazol-3-yl)methyl]thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide	25	105

N-(2-[(5-amino-1H-pyrazol-3-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide	1381	99
N-(2-[(5-amino-1H-pyrazol-3-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide	1867	98
N-(5-chloro-2-[(1-propyl-1H-imidazol-4-yl)methyl]thio}phenyl)-1-benzofuran-2-sulfonamide	1283	79
N-(5-chloro-2-[(1-propyl-1H-imidazol-4-yl)methyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide	806	93
N-(5-chloro-2-[(1-propyl-1H-imidazol-4-yl)methyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide	804	98
N-{5-chloro-2-[(pyridin-2-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide	29	100
N-{5-chloro-2-[(pyridin-2-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide	29	100
N-(5-chloro-2-[(1-oxidopyridin-2-yl)methyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide	663	102
N-(5-chloro-2-[(1-oxidopyridin-2-yl)methyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide	846	101
tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl}pyridin-2-yl}carbamate	1999	40
N-(2-[(2-aminopyridin-4-yl)methyl]thio}-5-chlorophenyl)-1-benzofuran-2-sulfonamide	80	100
tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl}pyridin-2-yl}carbamate	2260	76
tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]-1-oxidopyridin-2-yl}carbamate	2872	29
N-(2-[(2-aminopyridin-4-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide	865	101
N-(2-[(2-aminopyridin-4-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide	364	101
N-(2-[(2-amino-1-oxidopyridin-4-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide	9959	54
N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-1-benzofuran-2-sulfonamide	42	92
N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide	39	96
N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide	23	90
N-{5-chloro-2-[(3-hydroxybenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide	77	98
N-{5-chloro-2-[(3-hydroxybenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide	112	103
N-{5-chloro-2-[(3-hydroxybenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide	56	96
3-{[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino]phenyl}sulfonyl}-N,N-dimethylpropanamide	217	93


3-[(4-chloro-2-[(2,4-difluorophenyl)sulfonyl]amino}phenyl)thio]-N,N-dimethylpropanamide	2154	43
N-(2-[(5-amino-4H-1,2,4-triazol-3-yl)methyl]thio}-5-chlorophenyl)-1-benzofuran-2-sulfonamide	32	103
N-(2-[(5-amino-4H-1,2,4-triazol-3-yl)methyl]sulfinyl}-5-chlorophenyl)-1-benzofuran-2-sulfonamide	NA	48
3-[(4-chloro-2-[(2,4-difluorophenyl)sulfonyl]amino}phenyl)sulfinyl]-N,N-dimethylpropanamide	1477	93
3-[(4-chloro-2-[(2,4-difluorophenyl)sulfonyl]amino}phenyl)sulfonyl]-N,N-dimethylpropanamide	1577	89
3-[(4-chloro-2-[(4-chloro-2-fluorophenyl)sulfonyl]amino}phenyl)thio]-N,N-dimethylpropanamide	771	78
3-[(4-chloro-2-[(4-chloro-2-fluorophenyl)sulfonyl]amino}phenyl)sulfinyl]-N,N-dimethylpropanamide	221	99
3-[(4-chloro-2-[(4-chloro-2-fluorophenyl)sulfonyl]amino}phenyl)sulfonyl]-N,N-dimethylpropanamide	211	97
N-{5-chloro-2-[(2-pyridin-2-ylethyl)thio]phenyl}-1-benzofuran-2-sulfonamide	713	84
N-(5-chloro-2-[[2-(1H-pyrazol-4-yl)ethyl]sulfinyl]phenyl)-1-benzofuran-2-sulfonamide	1061	67
N-{5-chloro-2-[(2-pyridin-2-ylethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide	153	86
N-{5-chloro-2-[(2-pyridin-2-ylethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide	197	93
N-(5-chloro-2-[[2-(1H-pyrazol-4-yl)ethyl]thio]phenyl)-1-benzofuran-2-sulfonamide	412	91
N-(5-chloro-2-[[2-(1H-pyrazol-4-yl)ethyl]sulfonyl]phenyl)-1-benzofuran-2-sulfonamide	570	93
N-(5-chloro-2-[[2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl]thio]phenyl)-1-benzofuran-2-sulfonamide	279	89
N-(5-chloro-2-[[2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl]sulfinyl]phenyl)-1-benzofuran-2-sulfonamide	3047	83
N-(5-chloro-2-[[2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl]sulfonyl]phenyl)-1-benzofuran-2-sulfonamide	1088	96
N-(5-chloro-2-[(2-fluoropyridin-3-yl)methyl]thio}phenyl)-1-benzofuran-2-sulfonamide	94	76
N-(5-chloro-2-[(2-fluoropyridin-3-yl)methyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide	79	95
N-(5-chloro-2-[(2-fluoropyridin-3-yl)methyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide	79	88
N-{5-chloro-2-[(trifluoromethyl)thio]phenyl}-1-benzofuran-2-sulfonamide	333	92
N-{5-chloro-2-[(trifluoromethyl)sulfonyl]phenyl}-1-	73	91

benzofuran-2-sulfonamide		
N-{5-chloro-2-[(trifluoromethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide	19	89
N-[2-(benzylthio)-5-chlorophenyl]-1-benzofuran-2-sulfonamide	129	90
N-[2-(benzylsulfinyl)-5-chlorophenyl]-1-benzofuran-2-sulfonamide	27	85
N-[2-(benzylsulfonyl)-5-chlorophenyl]-1-benzofuran-2-sulfonamide	34	90
N-{2-[(3-aminobenzyl)thio]-5-fluorophenyl}-1-benzofuran-2-sulfonamide	55	97
N-{5-fluoro-2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide	314	96
N-{2-[(3-aminobenzyl)sulfinyl]-5-fluorophenyl}-1-benzofuran-2-sulfonamide	38	91
N-{5-fluoro-2-[(3-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide	693	81
N-{2-[(3-aminobenzyl)sulfonyl]-5-fluorophenyl}-1-benzofuran-2-sulfonamide	22	90
N-{2-[(3-aminobenzyl)thio]-5-methoxyphenyl}-1-benzofuran-2-sulfonamide	1397	100
N-{2-[(3-aminobenzyl)sulfonyl]-5-methoxyphenyl}-1-benzofuran-2-sulfonamide	176	107
N-{2-[(3-aminobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide	676	105
N-{2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide	653	97
N-{2-[(3-aminobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide	58	110
N-{5-methoxy-2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide	845	84
N-{2-[(3-aminobenzyl)sulfinyl]-5-methoxyphenyl}-1-benzofuran-2-sulfonamide	85	100
N-{2-[(3-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide	2131	83
N-{2-[(3-aminobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide	41	94
N-{5-chloro-2-[(4-nitrobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide	147	82
N-{2-[(2-aminobenzyl)thio]-5-chlorophenyl}-1-benzofuran-2-sulfonamide	100	91
N-{5-chloro-2-[(4-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide	2159	20
N-{2-[(4-aminobenzyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide	90	104
N-{5-chloro-2-[(2-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide	75	95
N-{2-[(2-aminobenzyl)sulfinyl]-5-chlorophenyl}-1-	115	93

benzofuran-2-sulfonamide		
N-{2-[(2-aminobenzyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide	129	98
N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-2,4-difluorobenzenesulfonamide	3244	64
N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-4-chloro-2-fluorobenzenesulfonamide	2824	88
N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-2,4-difluorobenzenesulfonamide	718	89
N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-2,4-difluorobenzenesulfonamide	443	99
N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-4-chloro-2-fluorobenzenesulfonamide	248	96
N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-4-chloro-2-fluorobenzenesulfonamide	113	95
N-{5-chloro-2-[(pyrimidin-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide	549	100
N-{5-chloro-2-[(pyrimidin-2-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide	506	104
N-{5-chloro-2-[(pyrimidin-2-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide	151	104
N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}benzenesulfonamide	3102	40
N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-4-chlorobenzenesulfonamide	3894	95
N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-3-chlorobenzenesulfonamide	1458	95
N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}benzenesulfonamide	1239	52
N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}benzenesulfonamide	2564	62
N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-4-chlorobenzenesulfonamide	421	99
N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-4-chlorobenzenesulfonamide	159	99
N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-3-chlorobenzenesulfonamide	246	91
N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-3-chlorobenzenesulfonamide	159	93
N-[2-(benzylthio)-5-cyanophenyl]-1-benzofuran-2-sulfonamide	3837	78
N-[2-(benzylsulfinyl)-5-cyanophenyl]-1-benzofuran-2-sulfonamide	50	105
N-[2-(benzylsulfonyl)-5-cyanophenyl]-1-benzofuran-2-sulfonamide	56	95

What is claimed is:

1. A compound having Formula I, its enantiomers, diastereoisomers, hydrates, solvates, crystal forms and individual isomers, tautomers or a pharmaceutically acceptable salt thereof:



Formula I

wherein:

R¹ is H;

R² is substituted or unsubstituted C₁₋₆ alkyl, substituted or unsubstituted C₃₋₈

10 cycloalkyl or is substituted or unsubstituted C₃₋₈ cycloalkenyl;

R⁵ is -S-, -S(O)-, or -S(O)₂-;

R⁶ is substituted or unsubstituted heterocycle, substituted or unsubstituted C₃₋₈ cycloalkyl, substituted or unsubstituted C₃₋₈ cycloalkenyl or is substituted or unsubstituted C₆₋₁₀ aryl;

15 R¹⁷ is H, substituted or unsubstituted C₁₋₆ alkyl, halogen, substituted or unsubstituted -OC₁₋₆ alkyl, CN, C(O)R¹⁹, NR²⁰R²¹ or hydroxyl;

R¹⁸ is H, substituted or unsubstituted C₁₋₆ alkyl, halogen, substituted or unsubstituted -OC₁₋₆ alkyl, CN, C(O)R²², NR²³R²⁴ or hydroxyl;

20 R⁷ is H, halogen, CN, substituted or unsubstituted -OC₁₋₆ alkyl, substituted or unsubstituted C₁₋₆ alkyl or is substituted or unsubstituted C₃₋₈ cycloalkyl;

R⁸ is H, substituted or unsubstituted C₁₋₆ alkyl, halogen, substituted or unsubstituted -OC₁₋₆ alkyl, or hydroxyl;

R¹⁹ is H, OH or substituted or unsubstituted C₁₋₆ alkyl;

R²⁰ is H or substituted or unsubstituted C₁₋₆ alkyl;

25 R²¹ is H or substituted or unsubstituted C₁₋₆ alkyl;

R²² is H, OH or substituted or unsubstituted C₁₋₆ alkyl;

R²³ is H or substituted or unsubstituted C₁₋₆ alkyl;

R^{24} is H or substituted or unsubstituted C_{1-6} alkyl; and including compounds:

N-{2-[(3-aminobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide;

N-{2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;

N-{2-[(3-aminobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;

5 N-{2-[(3-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide; and


N-{2-[(3-aminobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide; and


with the provisos that

a). R^7 , R^8 , R^{17} and R^{18} cannot all be H in same time;

b). when R^8 is substituted or unsubstituted C_{1-6} alkyl, the alkyl group does not
10 contain a carbonyl group directly attached to the phenyl ring;

c). the compound is not of the following structures:





5 2. A compound according to claim 1, wherein:

R^5 is S.

3. A compound according to claim 1, wherein:

R^5 is -S(O)- .

10

4. A compound according to claim 1, wherein:

R^5 is -S(O)₂- .

5. A compound according to claim 1, wherein:

15 R^1 is H;

R^2 is substituted or unsubstituted C₁₋₆ alkyl, substituted or unsubstituted C₃₋₈ cycloalkyl or substituted or unsubstituted C₃₋₈ cycloalkenyl;

R^5 is -S-, -S(O)-, or -S(O)₂-;

R^6 is substituted or unsubstituted heterocycle or is substituted or unsubstituted phenyl;

20 R^{17} is H, substituted or unsubstituted C₁₋₆ alkyl or halogen;

R¹⁸ is H, substituted or unsubstituted C₁₋₆ alkyl or halogen;

R⁷ is halogen, CN, -OC₁₋₆ alkyl, substituted or unsubstituted C₁₋₆ alkyl or substituted or unsubstituted C₃₋₈ cycloalkyl; and

R⁸ is H, substituted or unsubstituted C₁₋₆ alkyl or halogen.

5

6. A compound according to claim 5, wherein:

R⁵ is -S-.

7. A compound according to claim 5, wherein:

10 R⁵ is -S(O)-.

8. A compound according to claim 5, wherein:

R⁵ is -S(O)₂-.

15 9. A compound according to claim 5, wherein:

R¹ is H;

R² is substituted or unsubstituted C₁₋₆ alkyl;

R⁵ is -S-, -S(O)-, or -S(O)₂-;

R⁶ is substituted or unsubstituted heterocycle or substituted or unsubstituted phenyl;

20 R¹⁷ is H;

R¹⁸ is H;

R⁷ is halogen, CN or -OC₁₋₆ alkyl; and

R⁸ is H.

25 10. A compound according to claim 9, wherein:

R⁵ is -S-.

11. A compound according to claim 9, wherein:

R⁵ is -S(O).

30

12. A compound according to claim 9, wherein:

R⁵ is -S(O)₂-.

13. A compound according to claim 9, wherein:

R¹ is H;

R² is methyl, isopropyl, 2-hydroxyethyl, methylpropionate, 2-methylpyridine, ethylacetate, N,N-dimethylpropanamide, N-isopropylpropanamide, propamide, -hydroxycyclopentyl, ethyl, N,N-dimethylacetamide, N-methylacetamide, 2-aminoethyl, H-imidazol-2-ylmethyl, 1H-imidazol-4-ylmethyl, 2-[(aminocarbonyl)amino]ethyl, tert-butyl pyridin-2-yl-carbamate, 6-aminopyridin-2-yl, 2-oxo-1,3-oxazolidin-5-yl-methyl, 2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl, 3-aminopropyl, N,N-dimethylbutanamide, 1H-pyrazol-3-ylmethyl, methyl-1,3-thiazol-2-yl-acetamide, tert-butyl-1,3-thiazol-2-yl-carbamate, 2-amino-1,3-thiazol-4-yl)methyl, 3-methylpyridine, 3-nitrobenzyl, 3-methoxybenzyl, 5-nitro-1H-pyrazol-3-yl-methyl, 5-amino-1H-pyrazol-3-yl-methyl, 1-propyl-1H-imidazol-4-yl)methyl, tert-butyl -1-oxidopyridin-2-yl-carbamate, 3-hydroxybenzyl, 5-amino-4H-1,2,4-triazol-3-yl)methyl, 2-pyridin-2-ylethyl, 2-(1H-pyrazol-4-yl)ethyl, 2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl, (2-fluoropyridin-3-yl)methyl, trifluoromethyl, benzyl, 3-nitrobenzyl, 4-nitrobenzyl, 2-aminobenzyl, 3-aminobenzyl or pyrimidin-2-ylmethyl;

R⁵ is -S(O)₂-, S or -S(O)-;

R⁶ is 4-chloro-3-trifluoromethyl-phenyl, 3,4-dichlorophenyl, 3-methoxyphenyl, 4-methyl-3-nitrophenyl, 4-chlorophenyl, 4-chloro-3-methylphenyl, nitro-3-trifluoromethylphenyl, 2,4-difluorophenyl, 4-chloro-2-fluorophenyl, 3-chloro-2-fluorophenyl, 4-isopropylphenyl, 4-bromophenyl, 4-iodophenyl, 3-chlorophenyl, 2-benzofuran, 5-methyl-2-furan, 2-furan or phenyl ;

R⁷ is chlorine, cyano, methoxy or fluorine;

R¹⁷ is H;

R¹⁸ is H; and

R⁸ is H.

14. A compound according to claim 1, selected from:

4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
 4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
 4-chloro-N-[5-chloro-2-(methylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
 4-chloro-N-[5-chloro-2-(isopropylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
 4-chloro-N-[5-chloro-2-[(2-hydroxyethyl)thio]phenyl]-3-(trifluoromethyl)benzenesulfonamide;

- 4-chloro-N-[5-chloro-2-(isopropylsulfinyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
- 4-chloro-N-[5-chloro-2-[(2-hydroxyethyl)sulfinyl]phenyl]-3-(trifluoromethyl)benzenesulfonamide;
- 5 4-chloro-N-[5-chloro-2-[(2-hydroxyethyl)sulfonyl]phenyl]-3-(trifluoromethyl)benzenesulfonamide;
- methyl 3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}propanoate;
- 4-chloro-N-[5-chloro-2-[(pyridin-2-ylmethyl)thio]phenyl]-3-
- 10 (trifluoromethyl)benzenesulfonamide;
- 4-chloro-N-[5-chloro-2-[(pyridin-3-ylmethyl)thio]phenyl]-3-(trifluoromethyl)benzenesulfonamide;
- 4-chloro-N-[5-chloro-2-[(pyridin-2-ylmethyl)sulfinyl]phenyl]-3-(trifluoromethyl)benzenesulfonamide;
- 15 4-chloro-N-[5-chloro-2-[(pyridin-3-ylmethyl)sulfinyl]phenyl]-3-(trifluoromethyl)benzenesulfonamide;
- ethyl {[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}acetate;
- 3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio}-N-
- 20 isopropylpropanamide;
- 3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio}-N,N-dimethylpropanamide;
- 3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}-N-isopropylpropanamide;
- 25 3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}-N,N-dimethylpropanamide;
- 3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}propanamide;
- N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide;
- 30 3,4-dichloro-N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide;
- N-[5-chloro-2-(methylthio)phenyl]thiophene-2-sulfonamide;
- N-[5-chloro-2-(methylsulfinyl)phenyl]thiophene-2-sulfonamide;
- 4-chloro-N-[5-chloro-2-[(3-hydroxycyclopentyl)sulfinyl]phenyl]-3-(trifluoromethyl)benzenesulfonamide;

- 4-chloro-N-{5-chloro-2-[(3-hydroxycyclopentyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
- 4-chloro-N-[5-chloro-2-(ethylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
- 4-chloro-N-[5-chloro-2-(ethylsulfinyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
- 5 4-chloro-N-[5-chloro-2-(ethylsulfonyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
- N-[5-chloro-2-(methylthio)phenyl]-1-benzofuran-2-sulfonamide;
- N-[5-chloro-2-(methylsulfinyl)phenyl]-1-benzofuran-2-sulfonamide;
- N-[5-chloro-2-(methylsulfonyl)phenyl]-1-benzofuran-2-sulfonamide;
- N-[5-chloro-2-(methylthio)phenyl]-4-methyl-3-nitrobenzenesulfonamide;
- 10 4-chloro-N-[5-chloro-2-(methylthio)phenyl]benzenesulfonamide;
- 4-chloro-N-[5-chloro-2-(methylthio)phenyl]-3-methylbenzenesulfonamide;
- N-[5-chloro-2-(methylsulfinyl)phenyl]-4-methyl-3-nitrobenzenesulfonamide;
- N-[5-chloro-2-(methylsulfonyl)phenyl]-4-methyl-3-nitrobenzenesulfonamide;
- 4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide;
- 15 4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]benzenesulfonamide;
- 4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-3-methylbenzenesulfonamide;
- 4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-3-methylbenzenesulfonamide;
- 2-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio]-N,N-dimethylacetamide;
- 20 2-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio]-N-methylacetamide;
- 2-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl]-N,N-dimethylacetamide;
- 2-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfonyl]-N,N-dimethylacetamide;
- 25 N,N-dimethylacetamide;
- 2-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl]-N-methylacetamide;
- 2-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfonyl]-N-methylacetamide;
- 30 N-{2-[(2-aminoethyl)thio]-5-chlorophenyl}-4-chloro-3-(trifluoromethyl)benzenesulfonamide;
- N-[5-chloro-2-(methylthio)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide;
- N-[5-chloro-2-(methylsulfonyl)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide;
- N-[5-chloro-2-(methylthio)phenyl]-2,4-difluorobenzenesulfonamide;

- N-[5-chloro-2-(methylsulfonyl)phenyl]-2,4-difluorobenzenesulfonamide;
N-[5-chloro-2-(methylsulfinyl)phenyl]-2,4-difluorobenzenesulfonamide;
N-[5-chloro-2-(methylsulfinyl)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide;
N-[5-chloro-2-(methylthio)phenyl]-5-methylfuran-2-sulfonamide;
5 N-[5-chloro-2-(methylsulfinyl)phenyl]-5-methylfuran-2-sulfonamide;
N-[5-chloro-2-(methylsulfonyl)phenyl]-5-methylfuran-2-sulfonamide;
N-[5-chloro-2-(methylthio)phenyl]furan-2-sulfonamide;
N-[5-chloro-2-(methylsulfinyl)phenyl]furan-2-sulfonamide;
N-[5-chloro-2-(methylsulfonyl)phenyl]furan-2-sulfonamide;
10 N-{2-[(2-aminoethyl)sulfonyl]-5-chlorophenyl}-4-chloro-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-[5-chloro-2-(methylthio)phenyl]-2-fluorobenzenesulfonamide;
3-chloro-N-[5-chloro-2-(methylthio)phenyl]-2-fluorobenzenesulfonamide;
3-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-2-fluorobenzenesulfonamide;
15 3-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-2-fluorobenzenesulfonamide;
4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-2-fluorobenzenesulfonamide;
4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-2-fluorobenzenesulfonamide;
N-{2-[(2-aminoethyl)sulfinyl]-5-chlorophenyl}-4-chloro-3-(trifluoromethyl)benzenesulfonamide;
20 4-chloro-N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
25 (trifluoromethyl)benzenesulfonamide;
4-chloro-N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
30 4-chloro-N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;

- N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
 N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
 N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
 3-{2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}thio)-N,N-
 5 dimethylpropanamide;
 3-{2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfinyl)-N,N-
 dimethylpropanamide;
 3-{2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfonyl)-N,N-
 dimethylpropanamide;
 10 N-[2-({2-[(aminocarbonyl)amino]ethyl}thio)-5-chlorophenyl]-1-benzofuran-2-
 sulfonamide;
 N-[2-({2-[(aminocarbonyl)amino]ethyl}sulfinyl)-5-chlorophenyl]-1-benzofuran-2-
 sulfonamide;
 N-[2-({2-[(aminocarbonyl)amino]ethyl}sulfonyl)-5-chlorophenyl]-1-benzofuran-2-
 15 sulfonamide;
 tert-butyl {6-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-
 chlorophenyl}sulfinyl)methyl]pyridin-2-yl}carbamate;
 N-(2-[(6-aminopyridin-2-yl)methyl]thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
 N-(2-[(6-aminopyridin-2-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-
 20 sulfonamide;
 N-(2-[(6-aminopyridin-2-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-
 sulfonamide;
 N-(2-[(6-amino-1-oxidopyridin-2-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-
 sulfonamide;
 25 N-(5-chloro-2-{[3-(dimethylamino)propyl]thio}phenyl)-1-benzofuran-2-sulfonamide;
 N-(5-chloro-2-{[3-(dimethylamino)propyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;
 N-(5-chloro-2-{[3-(dimethylamino)propyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;
 N-(5-chloro-2-{[3-(dimethylnitro)propyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;
 N-(2-[4-chloro-2-([4-chloro-3-
 30 (trifluoromethyl)phenyl]sulfonyl)amino]phenyl)sulfonyl)ethyl)acetamide;
 N-(2-[4-chloro-2-([4-chloro-3-
 (trifluoromethyl)phenyl]sulfonyl)amino]phenyl)sulfinyl)ethyl)acetamide;
 N-(5-chloro-2-[(2-oxo-1,3-oxazolidin-5-yl)methyl]sulfinyl)phenyl)-1-benzofuran-2-
 sulfonamide;

- N-(5-chloro-2-[(2-oxo-1,3-oxazolidin-5-yl)methyl]sulfonyl)phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[(2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl]thio)phenyl)-1-benzofuran-2-sulfonamide;
- 5 N-(5-chloro-2-[(2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl]sulfinyl)phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[(2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl]sulfonyl)phenyl)-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminopropyl)thio]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- 10 N-{2-[(3-aminopropyl)sulfinyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminopropyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- 4-{2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfinyl)-N,N-dimethylbutanamide;
- 4-{2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfonyl)-N,N-
- 15 dimethylbutanamide;
- 5-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-1-benzofuran-2-sulfonamide;
- 5-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(1H-pyrazol-3-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(1H-pyrazol-3-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- 20 N-{5-chloro-2-[(1H-pyrazol-3-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)methyl]sulfinyl)phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)methyl]sulfonyl)phenyl)-1-benzofuran-2-sulfonamide;
- 25 N-{4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl]-1,3-thiazol-2-yl}acetamide;
- N-{4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfinyl)methyl]-1,3-thiazol-2-yl}acetamide;
- N-{4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]-1,3-
- 30 thiazol-2-yl}acetamide;
- N-[5-chloro-2-(methylsulfinyl)phenyl]-4-isopropylbenzenesulfonamide;
- 4-bromo-N-[5-chloro-2-(methylthio)phenyl]benzenesulfonamide;
- N-[5-chloro-2-(methylthio)phenyl]-4-iodobenzenesulfonamide;
- N-[5-chloro-2-(methylsulfonyl)phenyl]-4-isopropylbenzenesulfonamide;

- 4-bromo-N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide;
N-[5-chloro-2-(methylsulfinyl)phenyl]-4-iodobenzenesulfonamide;
4-bromo-N-[5-chloro-2-(methylsulfonyl)phenyl]benzenesulfonamide;
N-[5-chloro-2-(methylsulfonyl)phenyl]-4-iodobenzenesulfonamide;
- 5 tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl]-1,3-thiazol-2-yl}carbamate;
N-(2-[(2-amino-1,3-thiazol-4-yl)methyl]thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- 10 tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfinyl)methyl]-1,3-thiazol-2-yl}carbamate;
N-(2-[(2-amino-1,3-thiazol-4-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]-1,3-thiazol-2-yl}carbamate;
- 15 N-(2-[(2-amino-1,3-thiazol-4-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(pyridin-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(pyridin-3-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(pyridin-3-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- 20 N-(5-chloro-2-[(1-oxidopyridin-3-yl)methyl]sulfinyl)phenyl)-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(pyridin-3-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
N-(5-chloro-2-[(1-oxidopyridin-3-yl)methyl]sulfonyl)phenyl)-1-benzofuran-2-sulfonamide;
- 25 N-{5-chloro-2-[(3-nitrobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(3-methoxybenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-(5-chloro-2-[(5-nitro-1H-pyrazol-3-yl)methyl]thio)phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[(5-nitro-1H-pyrazol-3-yl)methyl]sulfinyl)phenyl)-1-benzofuran-2-sulfonamide;
- 30 N-(2-[(5-amino-1H-pyrazol-3-yl)methyl]thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- N-(2-[(5-amino-1H-pyrazol-3-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;

- N-(2-[[5-amino-1H-pyrazol-3-yl)methyl]sulfonyl]-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[[1-propyl-1H-imidazol-4-yl)methyl]thio}phenyl)-1-benzofuran-2-sulfonamide;
- 5 N-(5-chloro-2-[[1-propyl-1H-imidazol-4-yl)methyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[[1-propyl-1H-imidazol-4-yl)methyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(pyridin-2-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- 10 N-{5-chloro-2-[(pyridin-2-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[[1-oxidopyridin-2-yl)methyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[[1-oxidopyridin-2-yl)methyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;
- 15 tert-butyl {4-[[2-[(1-benzofuran-2-yl)sulfonyl]amino]-4-chlorophenyl]thio)methyl}pyridin-2-yl}carbamate;
- N-(2-[[2-aminopyridin-4-yl)methyl]thio]-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- tert-butyl {4-[[2-[(1-benzofuran-2-yl)sulfonyl]amino]-4-chlorophenyl]sulfonyl)methyl}pyridin-2-yl}carbamate;
- 20 tert-butyl {4-[[2-[(1-benzofuran-2-yl)sulfonyl]amino]-4-chlorophenyl]sulfonyl)methyl}-1-oxidopyridin-2-yl}carbamate;
- N-(2-[[2-aminopyridin-4-yl)methyl]sulfinyl]-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- N-(2-[[2-aminopyridin-4-yl)methyl]sulfonyl]-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- 25 sulfonamide;
- N-(2-[[2-amino-1-oxidopyridin-4-yl)methyl]sulfonyl]-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-1-benzofuran-2-sulfonamide
- N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- 30 N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(3-hydroxybenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(3-hydroxybenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(3-hydroxybenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;

- 3-[[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfonyl]-N,N-dimethylpropanamide;
- 3-[(4-chloro-2-{{(2,4-difluorophenyl)sulfonyl}amino}phenyl)thio]-N,N-dimethylpropanamide;
- 5 N-(2-{{[(5-amino-4H-1,2,4-triazol-3-yl)methyl]thio}-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- N-(2-{{[(5-amino-4H-1,2,4-triazol-3-yl)methyl]sulfinyl}-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- 3-[(4-chloro-2-{{(2,4-difluorophenyl)sulfonyl}amino}phenyl)sulfinyl]-N,N-dimethylpropanamide;
- 10 3-[(4-chloro-2-{{(2,4-difluorophenyl)sulfonyl}amino}phenyl)sulfonyl]-N,N-dimethylpropanamide;
- 3-[(4-chloro-2-{{(4-chloro-2-fluorophenyl)sulfonyl}amino}phenyl)thio]-N,N-dimethylpropanamide;
- 15 3-[(4-chloro-2-{{(4-chloro-2-fluorophenyl)sulfonyl}amino}phenyl)sulfinyl]-N,N-dimethylpropanamide;
- 3-[(4-chloro-2-{{(4-chloro-2-fluorophenyl)sulfonyl}amino}phenyl)sulfonyl]-N,N-dimethylpropanamide;
- N-{5-chloro-2-[(2-pyridin-2-ylethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
- 20 N-(5-chloro-2-{{[2-(1H-pyrazol-4-yl)ethyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(2-pyridin-2-ylethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(2-pyridin-2-ylethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-{{[2-(1H-pyrazol-4-yl)ethyl]thio}phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-{{[2-(1H-pyrazol-4-yl)ethyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;
- 25 N-(5-chloro-2-{{[2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl]thio}phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-{{[2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-{{[2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;
- 30 N-(5-chloro-2-{{[(2-fluoropyridin-3-yl)methyl]thio}phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-{{[(2-fluoropyridin-3-yl)methyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;

- N-(5-chloro-2-[(2-fluoropyridin-3-yl)methyl]sulfonyl]phenyl)-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(trifluoromethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(trifluoromethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
- 5 N-{5-chloro-2-[(trifluoromethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-[2-(benzylthio)-5-chlorophenyl]-1-benzofuran-2-sulfonamide;
- N-[2-(benzylsulfinyl)-5-chlorophenyl]-1-benzofuran-2-sulfonamide;
- N-[2-(benzylsulfonyl)-5-chlorophenyl]-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminobenzyl)thio]-5-fluorophenyl}-1-benzofuran-2-sulfonamide;
- 10 N-{5-fluoro-2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminobenzyl)sulfinyl]-5-fluorophenyl}-1-benzofuran-2-sulfonamide;
- N-{5-fluoro-2-[(3-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminobenzyl)sulfonyl]-5-fluorophenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminobenzyl)thio]-5-methoxyphenyl}-1-benzofuran-2-sulfonamide;
- 15 N-{2-[(3-aminobenzyl)sulfonyl]-5-methoxyphenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-methoxy-2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- 20 N-{2-[(3-aminobenzyl)sulfinyl]-5-methoxyphenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(3-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(4-nitrobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(2-aminobenzyl)thio]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- 25 N-{5-chloro-2-[(4-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(4-aminobenzyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(2-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(2-aminobenzyl)sulfinyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(2-aminobenzyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- 30 N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-2,4-difluorobenzenesulfonamide;
- N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-4-chloro-2-fluorobenzenesulfonamide;
- N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-2,4-difluorobenzenesulfonamide;
- N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-2,4-difluorobenzenesulfonamide;
- N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-4-chloro-2-fluorobenzenesulfonamide;

- N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-4-chloro-2-fluorobenzenesulfonamide;
 N-{5-chloro-2-[(pyrimidin-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
 N-{5-chloro-2-[(pyrimidin-2-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
 N-{5-chloro-2-[(pyrimidin-2-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
 5 N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}benzenesulfonamide;
 N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-4-chlorobenzenesulfonamide;
 N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-3-chlorobenzenesulfonamide;
 N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}benzenesulfonamide;
 N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}benzenesulfonamide;
 10 N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-4-chlorobenzenesulfonamide;
 N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-4-chlorobenzenesulfonamide;
 N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-3-chlorobenzenesulfonamide;
 N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-3-chlorobenzenesulfonamide;
 N-[2-(benzylthio)-5-cyanophenyl]-1-benzofuran-2-sulfonamide;
 15 N-[2-(benzylsulfinyl)-5-cyanophenyl]-1-benzofuran-2-sulfonamide; and
 N-[2-(benzylsulfonyl)-5-cyanophenyl]-1-benzofuran-2-sulfonamide.

15. A pharmaceutical composition comprising as active ingredient a therapeutically effective amount of a compound according to claim 1 and a pharmaceutically acceptable adjuvant, diluents or carrier.

16. A pharmaceutical composition according to claim 15 wherein the compound is selected from:
- 4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
 25 4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
 4-chloro-N-[5-chloro-2-(methylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
 4-chloro-N-[5-chloro-2-(isopropylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
 4-chloro-N-[5-chloro-2-[(2-hydroxyethyl)thio]phenyl]-3-(trifluoromethyl)benzenesulfonamide;
 30 4-chloro-N-[5-chloro-2-(isopropylsulfinyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
 4-chloro-N-[5-chloro-2-[(2-hydroxyethyl)sulfinyl]phenyl]-3-(trifluoromethyl)benzenesulfonamide;

- 4-chloro-N-{5-chloro-2-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
methyl 3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}propanoate;
- 5 4-chloro-N-{5-chloro-2-[(pyridin-2-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-{5-chloro-2-[(pyridin-3-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-{5-chloro-2-[(pyridin-2-ylmethyl)sulfinyl]phenyl}-3-
- 10 (trifluoromethyl)benzenesulfonamide;
4-chloro-N-{5-chloro-2-[(pyridin-3-ylmethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
ethyl {[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}acetate;
- 15 3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio}-N-isopropylpropanamide;
3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]thio}-N,N-dimethylpropanamide;
3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}-N-
- 20 isopropylpropanamide;
3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}-N,N-dimethylpropanamide;
3-{[4-chloro-2-({[4-chloro-3-(trifluoromethyl)phenyl]sulfonyl}amino)phenyl]sulfinyl}propanamide;
- 25 N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide;
3,4-dichloro-N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide;
N-[5-chloro-2-(methylthio)phenyl]thiophene-2-sulfonamide;
N-[5-chloro-2-(methylsulfinyl)phenyl]thiophene-2-sulfonamide;
4-chloro-N-{5-chloro-2-[(3-hydroxycyclopentyl)sulfinyl]phenyl}-3-
- 30 (trifluoromethyl)benzenesulfonamide;
4-chloro-N-{5-chloro-2-[(3-hydroxycyclopentyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-[5-chloro-2-(ethylthio)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-[5-chloro-2-(ethylsulfinyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide;

- 4-chloro-N-[5-chloro-2-(ethylsulfonyl)phenyl]-3-(trifluoromethyl)benzenesulfonamide;
 N-[5-chloro-2-(methylthio)phenyl]-1-benzofuran-2-sulfonamide;
 N-[5-chloro-2-(methylsulfinyl)phenyl]-1-benzofuran-2-sulfonamide;
 N-[5-chloro-2-(methylsulfonyl)phenyl]-1-benzofuran-2-sulfonamide;
 5 N-[5-chloro-2-(methylthio)phenyl]-4-methyl-3-nitrobenzenesulfonamide;
 4-chloro-N-[5-chloro-2-(methylthio)phenyl]benzenesulfonamide;
 4-chloro-N-[5-chloro-2-(methylthio)phenyl]-3-methylbenzenesulfonamide;
 N-[5-chloro-2-(methylsulfinyl)phenyl]-4-methyl-3-nitrobenzenesulfonamide;
 N-[5-chloro-2-(methylsulfonyl)phenyl]-4-methyl-3-nitrobenzenesulfonamide;
 10 4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide;
 4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]benzenesulfonamide;
 4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-3-methylbenzenesulfonamide;
 4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-3-methylbenzenesulfonamide;
 2-[[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino]phenyl]thio}-N,N-
 15 dimethylacetamide;
 2-[[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino]phenyl]thio}-N-
 methylacetamide;
 2-[[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino]phenyl]sulfinyl}-N,N-
 dimethylacetamide;
 20 2-[[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino]phenyl]sulfonyl}-
 N,N-dimethylacetamide;
 2-[[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino]phenyl]sulfinyl}-N-
 methylacetamide;
 2-[[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino]phenyl]sulfonyl}-N-
 25 methylacetamide;
 N-{2-[(2-aminoethyl)thio]-5-chlorophenyl}-4-chloro-3-
 (trifluoromethyl)benzenesulfonamide;
 N-[5-chloro-2-(methylthio)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide;
 N-[5-chloro-2-(methylsulfonyl)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide;
 30 N-[5-chloro-2-(methylthio)phenyl]-2,4-difluorobenzenesulfonamide;
 N-[5-chloro-2-(methylsulfonyl)phenyl]-2,4-difluorobenzenesulfonamide;
 N-[5-chloro-2-(methylsulfinyl)phenyl]-2,4-difluorobenzenesulfonamide;
 N-[5-chloro-2-(methylsulfinyl)phenyl]-3-nitro-4-(trifluoromethyl)benzenesulfonamide;
 N-[5-chloro-2-(methylthio)phenyl]-5-methylfuran-2-sulfonamide;

- N-[5-chloro-2-(methylsulfinyl)phenyl]-5-methylfuran-2-sulfonamide;
N-[5-chloro-2-(methylsulfonyl)phenyl]-5-methylfuran-2-sulfonamide;
N-[5-chloro-2-(methylthio)phenyl]furan-2-sulfonamide;
N-[5-chloro-2-(methylsulfinyl)phenyl]furan-2-sulfonamide;
5 N-[5-chloro-2-(methylsulfonyl)phenyl]furan-2-sulfonamide;
N-{2-[(2-aminoethyl)sulfonyl]-5-chlorophenyl}-4-chloro-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-[5-chloro-2-(methylthio)phenyl]-2-fluorobenzenesulfonamide;
3-chloro-N-[5-chloro-2-(methylthio)phenyl]-2-fluorobenzenesulfonamide;
10 3-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-2-fluorobenzenesulfonamide;
3-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-2-fluorobenzenesulfonamide;
4-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-2-fluorobenzenesulfonamide;
4-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-2-fluorobenzenesulfonamide;
N-{2-[(2-aminoethyl)sulfinyl]-5-chlorophenyl}-4-chloro-3-(trifluoromethyl)benzenesulfonamide;
15 4-chloro-N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
20 4-chloro-N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)thio]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
4-chloro-N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfinyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
25 4-chloro-N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfonyl]phenyl}-3-(trifluoromethyl)benzenesulfonamide;
N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
30 N-{5-chloro-2-[(1H-imidazol-2-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(1H-imidazol-4-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;

- 3-({2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}thio)-N,N-dimethylpropanamide;
- 3-({2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfinyl)-N,N-dimethylpropanamide;
- 5 3-({2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfonyl)-N,N-dimethylpropanamide;
- N-[2-({2-[(aminocarbonyl)amino]ethyl}thio)-5-chlorophenyl]-1-benzofuran-2-sulfonamide;
- N-[2-({2-[(aminocarbonyl)amino]ethyl}sulfinyl)-5-chlorophenyl]-1-benzofuran-2-
- 10 sulfonamide;
- N-[2-({2-[(aminocarbonyl)amino]ethyl}sulfonyl)-5-chlorophenyl]-1-benzofuran-2-sulfonamide;
- tert-butyl {6-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfinyl)methyl]pyridin-2-yl}carbamate;
- 15 N-(2-[(6-aminopyridin-2-yl)methyl]thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- N-(2-[(6-aminopyridin-2-yl)methyl}sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- N-(2-[(6-aminopyridin-2-yl)methyl}sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- 20 N-(2-[(6-amino-1-oxidopyridin-2-yl)methyl}sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[(3-(dimethylamino)propyl]thio)phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[(3-(dimethylamino)propyl}sulfinyl)phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[(3-(dimethylamino)propyl}sulfonyl)phenyl)-1-benzofuran-2-sulfonamide;
- 25 N-(5-chloro-2-[(3-(dimethylnitro)propyl}sulfonyl)phenyl)-1-benzofuran-2-sulfonamide;
- N-(2-[[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino]phenyl]sulfonyl)ethyl)acetamide;
- N-(2-[[4-chloro-2-([4-chloro-3-(trifluoromethyl)phenyl]sulfonyl)amino]phenyl]sulfinyl)ethyl)acetamide;
- 30 N-(5-chloro-2-[(2-oxo-1,3-oxazolidin-5-yl)methyl}sulfinyl)phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[(2-oxo-1,3-oxazolidin-5-yl)methyl}sulfonyl)phenyl)-1-benzofuran-2-sulfonamide;

- N-(5-chloro-2-[(2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl]thio}phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[(2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;
- 5 N-(5-chloro-2-[(2,6-dioxo-1,2,3,6-tetrahydropyrimidin-4-yl)methyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminopropyl)thio]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminopropyl)sulfinyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminopropyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- 10 4-{2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfinyl)-N,N-dimethylbutanamide;
- 4-{2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl}sulfonyl)-N,N-dimethylbutanamide;
- 5-chloro-N-[5-chloro-2-(methylsulfinyl)phenyl]-1-benzofuran-2-sulfonamide;
- 15 5-chloro-N-[5-chloro-2-(methylsulfonyl)phenyl]-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(1H-pyrazol-3-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(1H-pyrazol-3-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(1H-pyrazol-3-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)methyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;
- 20 N-(5-chloro-2-[(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)methyl]sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;
- N-{4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl]-1,3-thiazol-2-yl}acetamide;
- 25 N-{4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfinyl)methyl]-1,3-thiazol-2-yl}acetamide;
- N-{4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]-1,3-thiazol-2-yl}acetamide;
- N-[5-chloro-2-(methylsulfinyl)phenyl]-4-isopropylbenzenesulfonamide;
- 30 4-bromo-N-[5-chloro-2-(methylthio)phenyl]benzenesulfonamide;
- N-[5-chloro-2-(methylthio)phenyl]-4-iodobenzenesulfonamide;
- N-[5-chloro-2-(methylsulfonyl)phenyl]-4-isopropylbenzenesulfonamide;
- 4-bromo-N-[5-chloro-2-(methylsulfinyl)phenyl]benzenesulfonamide;
- N-[5-chloro-2-(methylsulfinyl)phenyl]-4-iodobenzenesulfonamide;


- 4-bromo-N-[5-chloro-2-(methylsulfonyl)phenyl]benzenesulfonamide;
N-[5-chloro-2-(methylsulfonyl)phenyl]-4-iodobenzenesulfonamide;
tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl]-1,3-thiazol-2-yl}carbamate;
- 5 N-(2-[(2-amino-1,3-thiazol-4-yl)methyl]thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfinyl)methyl]-1,3-thiazol-2-yl}carbamate;
N-(2-[(2-amino-1,3-thiazol-4-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-
- 10 sulfonamide;
tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]-1,3-thiazol-2-yl}carbamate;
N-(2-[(2-amino-1,3-thiazol-4-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-
- 15 sulfonamide;
N-{5-chloro-2-[(pyridin-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(pyridin-3-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(pyridin-3-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
N-(5-chloro-2-[(1-oxidopyridin-3-yl)methyl]sulfinyl)phenyl)-1-benzofuran-2-
- 20 sulfonamide;
N-{5-chloro-2-[(pyridin-3-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
N-(5-chloro-2-[(1-oxidopyridin-3-yl)methyl]sulfonyl)phenyl)-1-benzofuran-2-
- 25 sulfonamide;
N-{5-chloro-2-[(3-nitrobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(3-methoxybenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
- 30 N-(5-chloro-2-[(5-nitro-1H-pyrazol-3-yl)methyl]thio)phenyl)-1-benzofuran-2-sulfonamide;
N-(5-chloro-2-[(5-nitro-1H-pyrazol-3-yl)methyl]sulfinyl)phenyl)-1-benzofuran-2-sulfonamide;
N-(2-[(5-amino-1H-pyrazol-3-yl)methyl]thio)-5-chlorophenyl)-1-benzofuran-2-
- 30 sulfonamide;
N-(2-[(5-amino-1H-pyrazol-3-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
N-(2-[(5-amino-1H-pyrazol-3-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;

- N-(5-chloro-2-[(1-propyl-1H-imidazol-4-yl)methyl]thio)phenyl)-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[(1-propyl-1H-imidazol-4-yl)methyl]sulfinyl)phenyl)-1-benzofuran-2-sulfonamide;
- 5 N-(5-chloro-2-[(1-propyl-1H-imidazol-4-yl)methyl]sulfonyl)phenyl)-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(pyridin-2-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(pyridin-2-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-(5-chloro-2-[(1-oxidopyridin-2-yl)methyl]sulfonyl)phenyl)-1-benzofuran-2-
- 10 sulfonamide;
- N-(5-chloro-2-[(1-oxidopyridin-2-yl)methyl]sulfinyl)phenyl)-1-benzofuran-2-sulfonamide;
- tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)thio)methyl]pyridin-2-yl}carbamate;
- 15 N-(2-[(2-aminopyridin-4-yl)methyl]thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]pyridin-2-yl}carbamate;
- tert-butyl {4-[(2-[(1-benzofuran-2-ylsulfonyl)amino]-4-chlorophenyl)sulfonyl)methyl]-1-oxidopyridin-2-yl}carbamate;
- 20 N-(2-[(2-aminopyridin-4-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- N-(2-[(2-aminopyridin-4-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
- N-(2-[(2-amino-1-oxidopyridin-4-yl)methyl]sulfonyl)-5-chlorophenyl)-1-benzofuran-2-
- 25 sulfonamide;
- N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-1-benzofuran-2-sulfonamide
- N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(3-hydroxybenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
- 30 N-{5-chloro-2-[(3-hydroxybenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
- N-{5-chloro-2-[(3-hydroxybenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
- 3-[4-chloro-2-({4-chloro-3-(trifluoromethyl)phenyl}sulfonyl)amino]phenyl)sulfonyl)-N,N-dimethylpropanamide;

- 3-[(4-chloro-2-[(2,4-difluorophenyl)sulfonyl]amino}phenyl)thio]-N,N-dimethylpropanamide;
N-(2-[(5-amino-4H-1,2,4-triazol-3-yl)methyl]thio)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
5 N-(2-[(5-amino-4H-1,2,4-triazol-3-yl)methyl]sulfinyl)-5-chlorophenyl)-1-benzofuran-2-sulfonamide;
3-[(4-chloro-2-[(2,4-difluorophenyl)sulfonyl]amino}phenyl)sulfinyl]-N,N-dimethylpropanamide;
3-[(4-chloro-2-[(2,4-difluorophenyl)sulfonyl]amino}phenyl)sulfonyl]-N,N-dimethylpropanamide;
10 3-[(4-chloro-2-[(4-chloro-2-fluorophenyl)sulfonyl]amino}phenyl)thio]-N,N-dimethylpropanamide;
3-[(4-chloro-2-[(4-chloro-2-fluorophenyl)sulfonyl]amino}phenyl)sulfinyl]-N,N-dimethylpropanamide;
15 3-[(4-chloro-2-[(4-chloro-2-fluorophenyl)sulfonyl]amino}phenyl)sulfonyl]-N,N-dimethylpropanamide;
N-{5-chloro-2-[(2-pyridin-2-ylethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-(5-chloro-2-{2-(1H-pyrazol-4-yl)ethyl}sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(2-pyridin-2-ylethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
20 N-{5-chloro-2-[(2-pyridin-2-ylethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
N-(5-chloro-2-{2-(1H-pyrazol-4-yl)ethyl}thio}phenyl)-1-benzofuran-2-sulfonamide;
N-(5-chloro-2-{2-(1H-pyrazol-4-yl)ethyl}sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;
N-(5-chloro-2-{2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl}thio}phenyl)-1-benzofuran-2-sulfonamide;
25 N-(5-chloro-2-{2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl}sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;
N-(5-chloro-2-{2-(3,5-dimethyl-1H-pyrazol-4-yl)ethyl}sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;
N-(5-chloro-2-{[(2-fluoropyridin-3-yl)methyl]thio}phenyl)-1-benzofuran-2-sulfonamide;
30 N-(5-chloro-2-{[(2-fluoropyridin-3-yl)methyl]sulfinyl}phenyl)-1-benzofuran-2-sulfonamide;
N-(5-chloro-2-{[(2-fluoropyridin-3-yl)methyl}sulfonyl}phenyl)-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(trifluoromethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;

- N-{5-chloro-2-[(trifluoromethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(trifluoromethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
N-[2-(benzylthio)-5-chlorophenyl]-1-benzofuran-2-sulfonamide;
N-[2-(benzylsulfinyl)-5-chlorophenyl]-1-benzofuran-2-sulfonamide;
5 N-[2-(benzylsulfonyl)-5-chlorophenyl]-1-benzofuran-2-sulfonamide;
N-[2-[(3-aminobenzyl)thio]-5-fluorophenyl]-1-benzofuran-2-sulfonamide;
N-{5-fluoro-2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
N-[2-[(3-aminobenzyl)sulfinyl]-5-fluorophenyl]-1-benzofuran-2-sulfonamide;
N-{5-fluoro-2-[(3-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
10 N-[2-[(3-aminobenzyl)sulfonyl]-5-fluorophenyl]-1-benzofuran-2-sulfonamide;
N-[2-[(3-aminobenzyl)thio]-5-methoxyphenyl]-1-benzofuran-2-sulfonamide;
N-[2-[(3-aminobenzyl)sulfonyl]-5-methoxyphenyl]-1-benzofuran-2-sulfonamide;
N-[2-[(3-aminobenzyl)thio]phenyl]-1-benzofuran-2-sulfonamide;
N-[2-[(3-nitrobenzyl)sulfinyl]phenyl]-1-benzofuran-2-sulfonamide;
15 N-[2-[(3-aminobenzyl)sulfinyl]phenyl]-1-benzofuran-2-sulfonamide;
N-{5-methoxy-2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
N-[2-[(3-aminobenzyl)sulfinyl]-5-methoxyphenyl]-1-benzofuran-2-sulfonamide;
N-[2-[(3-nitrobenzyl)sulfonyl]phenyl]-1-benzofuran-2-sulfonamide;
N-[2-[(3-aminobenzyl)sulfonyl]phenyl]-1-benzofuran-2-sulfonamide;
20 N-{5-chloro-2-[(4-nitrobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-[2-[(2-aminobenzyl)thio]-5-chlorophenyl]-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(4-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
N-[2-[(4-aminobenzyl)sulfonyl]-5-chlorophenyl]-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(2-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;
25 N-[2-[(2-aminobenzyl)sulfinyl]-5-chlorophenyl]-1-benzofuran-2-sulfonamide;
N-[2-[(2-aminobenzyl)sulfonyl]-5-chlorophenyl]-1-benzofuran-2-sulfonamide;
N-[2-[(3-aminobenzyl)thio]-5-chlorophenyl]-2,4-difluorobenzenesulfonamide;
N-[2-[(3-aminobenzyl)thio]-5-chlorophenyl]-4-chloro-2-fluorobenzenesulfonamide;
N-[2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl]-2,4-difluorobenzenesulfonamide;
30 N-[2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl]-2,4-difluorobenzenesulfonamide;
N-[2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl]-4-chloro-2-fluorobenzenesulfonamide;
N-[2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl]-4-chloro-2-fluorobenzenesulfonamide;
N-{5-chloro-2-[(pyrimidin-2-ylmethyl)thio]phenyl}-1-benzofuran-2-sulfonamide;
N-{5-chloro-2-[(pyrimidin-2-ylmethyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;

- N-{5-chloro-2-[(pyrimidin-2-ylmethyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide;
 N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}benzenesulfonamide;
 N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-4-chlorobenzenesulfonamide;
 N-{2-[(3-aminobenzyl)thio]-5-chlorophenyl}-3-chlorobenzenesulfonamide;
 5 N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}benzenesulfonamide;
 N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}benzenesulfonamide;
 N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-4-chlorobenzenesulfonamide;
 N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-4-chlorobenzenesulfonamide;
 N-{2-[(3-aminobenzyl)sulfonyl]-5-chlorophenyl}-3-chlorobenzenesulfonamide;
 10 N-{2-[(3-aminobenzyl)sulfinyl]-5-chlorophenyl}-3-chlorobenzenesulfonamide;
 N-[2-(benzylthio)-5-cyanophenyl]-1-benzofuran-2-sulfonamide;
 N-[2-(benzylsulfinyl)-5-cyanophenyl]-1-benzofuran-2-sulfonamide; and
 N-[2-(benzylsulfonyl)-5-cyanophenyl]-1-benzofuran-2-sulfonamide.
- 15 17. A method of treating a disorder associated with chemokine receptor modulation, which comprises administering to a mammal in need thereof, a pharmaceutical composition comprising a therapeutically effective amount of at least one compound of Formula I



Formula I

wherein:

R^1 is H;

R^2 is substituted or unsubstituted C_{1-6} alkyl, substituted or unsubstituted C_{3-8}

25 cycloalkyl or is substituted or unsubstituted C_{3-8} cycloalkenyl;

R^5 is $-\text{S}-$, $-\text{S}(\text{O})-$, or $-\text{S}(\text{O})_2-$;

R^6 is substituted or unsubstituted heterocycle, substituted or unsubstituted C_{3-8} cycloalkyl, substituted or unsubstituted C_{3-8} cycloalkenyl or is substituted or unsubstituted C_{6-10} aryl;

R^{17} is H, substituted or unsubstituted C_{1-6} alkyl, halogen, substituted or unsubstituted -
 5 OC_{1-6} alkyl, CN, $C(O)R^{19}$, $NR^{20}R^{21}$ or hydroxyl;

R^{18} is H, substituted or unsubstituted C_{1-6} alkyl, halogen, substituted or unsubstituted -
 OC_{1-6} alkyl, CN, $C(O)R^{22}$, $NR^{23}R^{24}$ or hydroxyl;

R^7 is H, halogen, CN, substituted or unsubstituted $-OC_{1-6}$ alkyl, substituted or unsubstituted C_{1-6} alkyl or is substituted or unsubstituted C_{3-8} cycloalkyl;

10 R^8 is H, substituted or unsubstituted C_{1-6} alkyl, halogen, substituted or unsubstituted -
 OC_{1-6} alkyl, or hydroxyl;

R^{19} is H, OH or substituted or unsubstituted C_{1-6} alkyl;

R^{20} is H or substituted or unsubstituted C_{1-6} alkyl;

R^{21} is H or substituted or unsubstituted C_{1-6} alkyl;

15 R^{22} is H, OH or substituted or unsubstituted C_{1-6} alkyl;

R^{23} is H or substituted or unsubstituted C_{1-6} alkyl;

R^{24} is H or substituted or unsubstituted C_{1-6} alkyl; and including compounds:

N-{2-[(3-aminobenzyl)thio]phenyl}-1-benzofuran-2-sulfonamide;

N-{2-[(3-nitrobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;

20 N-{2-[(3-aminobenzyl)sulfinyl]phenyl}-1-benzofuran-2-sulfonamide;

N-{2-[(3-nitrobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide; and





N-{2-[(3-aminobenzyl)sulfonyl]phenyl}-1-benzofuran-2-sulfonamide; and


with the provisos that

a). R^7 , R^8 , R^{17} and R^{18} cannot all be H in same time;

25 b). when R^8 is substituted or unsubstituted C_{1-6} alkyl, the alkyl group does not
 contain a carbonyl group directly attached to the phenyl ring;

c). the compound is not of the following structures:





18. A method of claim 17, wherein the pharmaceutical composition is administered
5 to the mammal to treat ocular inflammatory diseases or skin inflammatory diseases.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2011/064233

A. CLASSIFICATION OF SUBJECT MATTER INV. C07D213/34 C07D213/62 C07D233/64 C07D307/18 C07D307/82 C07D333/34 C07D407/12 C07D417/12 A61K31/343 A61P31/00 C07C311/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) C07D C07C A61K A61P Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, CHEM ABS Data, WPI Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 947 500 A1 (ONO PHARMACEUTICAL CO [JP]) 6 October 1999 (1999-10-06) example 21(5)	1,2,15
X	----- MASAKI HIDEKAZU ET AL: "Structure-activity relationship of benzo(b)thiophene-2-sulfonamide derivatives as novel human chymase inhibitors", BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, PERGAMON, ELSEVIER SCIENCE, GB, vol. 13, no. 22, 17 November 2003 (2003-11-17), pages 4085-4088, XP002523990, ISSN: 0960-894X, DOI: 10.1016/J.BMCL.2003.08.040 table 2; compounds 21,22 ----- -/--	1,4
☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search 7 February 2012		Date of mailing of the international search report 17/02/2012
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016		Authorized officer Usuelli, Ambrogio

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2011/064233

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE 196 29 144 A1 (BAYER AG [DE]) 22 January 1998 (1998-01-22) examples 1-3 -----	1-3
X	US 5 163 995 A (VAN HEERTUM JOHN C [US] ET AL) 17 November 1992 (1992-11-17) table 1; compound 6 -----	1
X	EP 1 325 920 A1 (TOA EIYO LTD [JP]) 9 July 2003 (2003-07-09) example 13 -----	1
X	EP 0 142 152 A2 (DOW CHEMICAL CO [US]) 22 May 1985 (1985-05-22) compounds 69,70 -----	1
X	EP 0 244 098 A2 (SCHERING AGROCHEMICALS LTD [GB]) 4 November 1987 (1987-11-04) compounds 34,40,44,45,53,78,79 -----	1
X	EP 0 246 749 A2 (SCHERING AGROCHEMICALS LTD [GB]) 25 November 1987 (1987-11-25) compounds A48,A59 -----	1
X	US 5 177 206 A (JOHNSON TIMOTHY C [US] ET AL) 5 January 1993 (1993-01-05) table 1 -----	1
X	EP 0 244 097 A2 (SCHERING AGROCHEMICALS LTD [GB]) 4 November 1987 (1987-11-04) compound D39 -----	1
X	DATABASE CAPLUS [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; 1998, XP002668923, retrieved from STN accession no. 1999:104638 Database accession no. 130:248286 Compound with RN 98967-44-3 & GAODENG XUEXIAO HUAXUE XUEBAO, vol. 19, no. 12, 1998, - 1998, pages 1950-1953, -----	1
Y	US 2007/037794 A1 (UNGASHE SOLOMON [US] ET AL) 15 February 2007 (2007-02-15) paragraph [0074] - paragraph [0097] -----	1-18
Y	WO 2007/067875 A2 (GLAXO GROUP LIMITED [GB]; GOODMAN KRISTA B [US]; SEHON CLARK A [US]; CL) 14 June 2007 (2007-06-14) page 1, line 25 - page 3, line 5 -----	1-18

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2011/064233

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0947500	A1	06-10-1999	AT 499339 T 15-03-2011
		AU 733493 B2 17-05-2001	
		AU 5411598 A 15-07-1998	
		CA 2274954 A1 25-06-1998	
		CN 1247529 A 15-03-2000	
		EP 0947500 A1 06-10-1999	
		JP 3426252 B2 14-07-2003	
		KR 20000057576 A 25-09-2000	
		NO 992935 A 16-08-1999	
		TW 523506 B 11-03-2003	
		US 6448290 B1 10-09-2002	
		WO 9827053 A1 25-06-1998	
DE 19629144	A1	22-01-1998	AU 733250 B2 10-05-2001
		AU 3620297 A 10-02-1998	
		BR 9710385 A 17-08-1999	
		CA 2261136 A1 29-01-1998	
		CN 1225638 A 11-08-1999	
		DE 19629144 A1 22-01-1998	
		EP 0915878 A2 19-05-1999	
		JP 2000515518 A 21-11-2000	
		US 6297195 B1 02-10-2001	
		WO 9803508 A2 29-01-1998	
US 5163995	A	17-11-1992	NONE
EP 1325920	A1	09-07-2003	AT 478857 T 15-09-2010
		AU 8805301 A 26-03-2002	
		CA 2422807 A1 18-03-2003	
		CN 1458928 A 26-11-2003	
		EP 1325920 A1 09-07-2003	
		JP 3847711 B2 22-11-2006	
		WO 0222595 A1 21-03-2002	
EP 0142152	A2	22-05-1985	AR 244230 A1 29-10-1993
		AU 583799 B2 11-05-1989	
		AU 613665 B2 08-08-1991	
		AU 2290088 A 05-01-1989	
		AU 3533084 A 23-05-1985	
		BR 8405797 A 17-09-1985	
		CA 1231708 A1 19-01-1988	
		DE 3484294 D1 25-04-1991	
		DE 3486287 D1 07-04-1994	
		DE 3486287 T2 01-06-1994	
		DE 19775042 I2 03-05-2001	
		DK 541384 A 15-05-1985	
		EP 0142152 A2 22-05-1985	
		EP 0330137 A1 30-08-1989	
		GB 2149792 A 19-06-1985	
		GB 2196627 A 05-05-1988	
		GB 2196628 A 05-05-1988	
		GR 80917 A1 13-03-1985	
		IL 73486 A 13-05-1993	
		JP 1904055 C 08-02-1995	
		JP 6035459 B 11-05-1994	
		JP 60116684 A 24-06-1985	
		NZ 210192 A 30-06-1988	
		ZA 8408844 A 30-07-1986	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2011/064233

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0244098	A2	04-11-1987	AU 574016 B2 23-06-1988
			AU 7221087 A 10-12-1987
			BR 8702111 A 09-02-1988
			CN 87103150 A 23-03-1988
			DD 260433 A5 28-09-1988
			DK 218387 A 31-10-1987
			EP 0244098 A2 04-11-1987
			FI 871877 A 31-10-1987
			JP 62258385 A 10-11-1987
			PH 23354 A 14-07-1989
			PT 84778 A 01-05-1987
			SU 1678212 A3 15-09-1991
			TR 23214 A 21-06-1989
			US 4795483 A 03-01-1989
			ZA 8703073 A 24-02-1988
EP 0246749	A2	25-11-1987	AU 583728 B2 04-05-1989
			AU 7300087 A 26-11-1987
			BR 8702527 A 23-02-1988
			DK 245087 A 18-11-1987
			EP 0246749 A2 25-11-1987
			FI 872124 A 18-11-1987
			HU 204974 B 30-03-1992
			PT 84872 A 01-06-1987
			TR 24082 A 04-03-1991
			US 4889553 A 26-12-1989
US 5177206	A	05-01-1993	AT 169626 T 15-08-1998
			AU 649358 B2 19-05-1994
			BR 9203919 A 27-04-1993
			CA 2080097 A1 09-04-1993
			CN 1071429 A 28-04-1993
			DE 69226590 D1 17-09-1998
			DE 69226590 T2 24-12-1998
			DK 0537611 T3 10-05-1999
			EP 0537611 A1 21-04-1993
			ES 2118775 T3 01-10-1998
			FI 924522 A 09-04-1993
			HU 214216 B 28-01-1998
			IL 103374 A 27-11-1995
			JP 3342057 B2 05-11-2002
			JP 5239060 A 17-09-1993
			RU 2061692 C1 10-06-1996
			US 5177206 A 05-01-1993
EP 0244097	A2	04-11-1987	AU 573129 B2 26-05-1988
			AU 7220887 A 10-12-1987
			BR 8702112 A 09-02-1988
			CN 87103904 A 10-02-1988
			DD 260212 A5 21-09-1988
			DK 218287 A 31-10-1987
			EP 0244097 A2 04-11-1987
			FI 871878 A 31-10-1987
			IL 82356 A 18-07-1991
			JP 62263180 A 16-11-1987
			PH 23571 A 11-09-1989
			PT 84779 A 01-05-1987

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2011/064233

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
		SU 1528322 A3	07-12-1989	
		ZA 8703072 A	24-02-1988	

US 2007037794	A1	15-02-2007	US 2007037794 A1	15-02-2007
			US 2010056509 A1	04-03-2010
			WO 2008008431 A2	17-01-2008

WO 2007067875	A2	14-06-2007	EP 1962847 A2	03-09-2008
			JP 2009518444 A	07-05-2009
			US 2008293720 A1	27-11-2008
			WO 2007067875 A2	14-06-2007

15 If available, please enter the **Chinese abstract** text in the space below.

本發明涉及新的硫衍生物、它們的製備方法、包含它們的藥物組合物以及它們用作趨化因子受體調節劑的藥物的用途。