



(86) Date de dépôt PCT/PCT Filing Date: 2006/04/30
(87) Date publication PCT/PCT Publication Date: 2006/11/09
(45) Date de délivrance/Issue Date: 2011/03/15
(85) Entrée phase nationale/National Entry: 2007/11/05
(86) N° demande PCT/PCT Application No.: CN 2006/000895
(87) N° publication PCT/PCT Publication No.: 2006/116950
(30) Priorité/Priority: 2005/05/04 (US11/121,721)

(51) Cl.Int./Int.Cl. *A61K 36/8984* (2006.01),
A61K 39/00 (2006.01), *A61K 39/39* (2006.01),
A61P 27/02 (2006.01), *A61P 37/02* (2006.01)
(72) Inventeur/Inventor:
WU, RONG-TSUN, CN
(73) Propriétaire/Owner:
NATIONAL YANG-MING UNIVERSITY, TW
(74) Agent: MARKS & CLERK

(54) Titre : POLYSACCHARIDE EXTRAIT DE DENBROBII HERBA ET SON UTILISATION DANS LE TRAITEMENT DE
L'UVEITE AUTO-IMMUNE
(54) Title: POLYSACCHARIDE EXTRACT OF DENDROBII HERBA AND USE THEREOF FOR TREATING
AUTOIMMUNE UVEITIS

(57) **Abrégé/Abstract:**

A composition contains a polysaccharide from Dendrobium. It is used for treating autoimmune diseases and mucosal diseases by inducing oral tolerance and accelerating innate immunity.

Abstract

A composition contains a polysaccharide from Dendrobium. It is used for treating autoimmune diseases and mucosal diseases by inducing oral tolerance and accelerating innate immunity.

DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME __1__ DE __2__

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**

THIS IS VOLUME __1__ OF __2__

NOTE: For additional volumes please contact the Canadian Patent Office.

POLYSACCHARIDE EXTRACT OF DENDROBII HERBA AND USE THEREOF
FOR TREATING AUTOIMMUNE UVEITIS

FIELD OF THE INVENTION

[0001] This invention relates to a composition and a method for treating an autoimmune disease and mucosal disorder, and more particularly to a composition including the polysaccharides derived from Dendrobium and a method including the administration of the polysaccharides derived from Dendrobium for treating the autoimmune disease and mucosal disorder via oral-tolerance induction and innate immunity promotion.

BACKGROUND OF THE INVENTION

[0002] A basic property of the immune system is the immunologic tolerance that provides for self/non-self discrimination, so that the immune system can protect the host from external pathogens without reacting against itself. When the immune system reacts against itself, autoimmune disease results. (Annu. Rev. Med. 48:341-351, 1997)

[0003] Oral tolerance has been proven to be of therapeutic benefit in autoimmune diseases, including uveitis, collagen-induced arthritis, adjuvant arthritis, systemic

lupus erythematosus, multiple sclerosis, thyroiditis, myasthenia gravis, inflammatory bowel disease and type 1 diabetes. (Annals of the New York Academy of Sciences 778(1): 217-227, 1996)

[0004] An orally administered antigen encounters the gut-associated lymphoid tissue (GALT), which is a well-developed immune network. The GALT consists of villi, lamina propria, intraepithelial lymphocytes and Peyer's patch, wherein Peyer's patch consists of lymphoid nodules interspersed among the villi. The epithelium that lines on the gut contains M cells for transporting antigens and microorganisms. Antigens and pathogens in the gut can only penetrate the barrier through M cells, specialized to deliver antigen directly to underlying immune cells. Then, the intraepithelial lymphocytes recognizing the antigens translocate to Peyer's patches to trigger immune responses. In Peyer's patches, B cells and macrophages present the antigens to lymphocytes, and then the antigen-specific lymphocytes circulate to the lamina propria in the mucosal system (such as mucosal surface of respiratory tract) of the body. (Science 277:949-952, 1997; Science 277:910-911, 1997)

[0005] The intestinal immune system includes the innate immunity and the adaptive immunity for protecting against diseases mediated via the oral administration. The tissue portion infected by the pathogens secretes chemokine for inducing the non-specific innate immune responses of macrophages, monocytes, granulocytes and natural killer cells. The macrophages and granulocytes can directly take the pathogens by phagocytosis. The natural killer cells are the immune cells in the body firstly responding to the viral infections or cells infected by other pathogens. The natural killer cells not only attack the target cells, but also secrete cytokines for regulating immune responses such as IFN- γ , TGF- β , TNF- α , IL-5 and IL-10. The macrophages and other antigen presenting cells, such as dendritic cells, present antigens to T cells via MHC class II molecules and co-stimulators on the cell surface, so as to further activate immune responses, i.e. cell-mediated immunity and humoral immunity respectively performed by T cells and B cells. The mucosal immunity is mainly regulated by the cytokines secreted from CD4⁺ T cells. IL-4, IL-5, IL-6 and IL-10 secreted from type 2 helper T cells involve in stimulating B cells for antibody production. TGF- β secreted from type 3 helper T cells induces B cells to produce IgA. (Trends Immunol. 22:244-247, 2004; J. Clin. Invest. 106:935-937, 2000)

[0006] It is to be noted that the mucosal immune responses are not only induced in the gut, but also in the mucosal system of the whole body. Once there are antigen-specific lymphocytes activated in the inductive site-Peyer's patch in intestine, the lymphocytes travel through lymph drain and blood to effector site include rectum, genital tract, lung and so on. Then the antibodies were produced and detected at these effector sites. This homing phenomenon is so called common mucosal immune system. Due to the common mucosal immune response, the antibodies induced by the oral administration are distributed in the whole body. If there is the antigen mediated via the oral administration, the systematic tolerance or IgA, i.e. the oral tolerance is induced. The oral tolerance includes both the systematic unresponsiveness and the mucosal responsiveness, wherein the former prevents the whole body from autoimmune diseases, and the later induces active suppression to specific antigen. Therefore, the mucosal immune system tends to inhibit the immune responses of the inflammations, and further to enhance the presentation of IgA in the non-inflammatory local mucosa. (Immunol. Today 18:335-343, 1997)

[0007] The surface of the gastrointestinal tract is lined by a simple columnar epithelium to form a barrier against the excessive absorption of bacteria, food antigens and large molecules, and moreover the transportation of small molecules is controlled by the tight junction formed in the barrier. The crypts embedded in the connective tissue include stem cells for regenerating the intestinal epithelial cells. In addition to the intestinal epithelia cells, the intestinal stem cells also can differentiate into Goblet cells and enteroendocrine cells for secreting mucin, and into Paneth cells for secreting antimicrobial peptides. In addition to the Paneth cells staying in the stem cells region, the differentiated cells migrate to the top of the villi. (J. Clin. Invest. 105:1493-1499, 2000; Science 294:2115-2116, 2001)

[0008] It is a reasonable suggestion that the intestinal epithelia cells play a critical role for influencing immune responses to various antigens in the intestinal lumen due to the special location of the epithelial cells. The intestinal epithelial cells can express antigen-presenting molecules as antigen-presenting cells (APCs) so as to regulate T-cell responses in the intestinal mucosa. (Immunol. Today 21:123-128, 2000) It is investigated that the antigen-presenting molecules expressed by the intestinal epithelial cells include MHC class I, MHC class II and CDId.

(Gastroenterology 124:1420-1431, 2003) The MHC class I and MHC class II are

expressed on the basolateral membrane of epithelial cells, wherein MHC class I is responsible for coupling with CD8+ cells, and MHC class II is responsible for coupling with CD4+ cells. CD1d activates NK T cells via glycolipid. These evidences show that the intestinal epithelial cells play a critical role in the mucosal immune system.

[0009] In addition to being the physical barrier, the intestinal epithelial cells secrete antimicrobial peptides against microbes in the gastrointestinal tract and provide signals to other cells. (Immunol. Today 21:123-128, 2000; J. Clin. Invest. 95:55-65, 1995) It is found that there are specific antimicrobial peptides, defensins. Defensins are 3-5 kD proteins, which include .alpha.-defensin and .beta.-defensin family having 8 peptides. These defensins have organ-specific expression patterns in the epithelial cells of oral mucosa, lung and gastrointestinal tract. (Eur. J. Gastroenterol. Hepatol. 13:771-776, 2001; Nat. Rev. Immunol. 3:710-720, 2003)

When the gastrointestinal tract is infected, the intestinal epithelial cells express defensins, the defensins function as chemokines to induce NK cells and dendritic cells to the infected areas, and to perform the so-called innate immunity. In addition to the induction of innate immunity, defensins induce the dendritic cells to express co-stimulator (B7.2) through the toll-like receptor 4, so as to induce the

proliferation of T cells. Therefore, defensins involve the link between the innate immunity and the adaptive immunity.

[0010] Other antimicrobial proteins, angiogenins, were considered to involve the angiogenesis of cancer cells; however, in 2003, Hooper disclosed that angiogenins are produced by the Paneth cells in the gastrointestinal tract under normal physiological conditions, and stored in the cellular granules. The angiogenins are secreted into the gastrointestinal tract in response to the lipopolysaccharide stimulation. Comparing the Germ free mice and the mice colonized with intestinal bacteria, it is found that angiogenins are significantly increased due to the presence of the intestinal bacteria. Like defensins, the angiogenins are bactericidal and modulated by local environmental conditions encountered at infected sites. (Nat. Immunol. 4:269-273, 2003)

[0011] In addition to secreting antimicrobial peptides against the pathogen infection, the intestinal epithelial cells produce many signals for regulating the immune responses in the gastrointestinal tract. In order to protect the host by destroying the invading microbes, the intestinal epithelial cells should immediately respond to the invading microbes via the defense mechanism. The defense

mechanism is known as the innate immunity system, which uses germline-encoded pattern-recognition receptors for the recognition of the macromolecules of the microbial pathogens, such as lipopolysaccharide present in the cell wall of Gram-negative bacteria. In recent years, how the pattern-recognition receptors activate cells has been disclosed. In adult *Drosophila*, it is shown that Toll induces antifungal and antibacterial peptides upon infection. Medzhitov and Janeway et al. disclose that in mammals, the activation of Toll-like receptor (TLR) results in induction of cytokines and costimulatory molecules required for the activation of the adaptive immune response. (Cell 91:295-298, 1997) The intracellular signaling pathways activated by TLRs share much in common with IL-1R signaling due to their conserved TIR (Toll/IL-1R homology) domains. When the endotoxin is recognized by TLRs, the nuclear translocation of NF- κ B translocates is triggered to regulate the transcriptions of other genes involved in immune responses. Subsequently, the proinflammatory cytokine genes are activated, antimicrobial peptides are secreted, and the chemokines, such as IL-8, MIP and MCP-1, are secreted. The macrophages and NK cells are induced by these chemokines so as to translocate to the infected site and to destroy the infected cells. (Cell 91:295-298, 1997; Curr. Opin. Immunol 14:103-110, 2002) Hence, the toll-like receptors not only induce the innate

immunity system, but also indirectly induce the adaptive immunity system due to the secretion of proinflammatory cytokines triggered by NF-kB activation.

[0012] In human cells, there are at least 10 different TLRs, wherein the TLR-2, TLR-4 and TLR-9 are more characterized members of the TLR family. TLRs have been shown to mediate the recognition of many types of pathogens, including bacteria and viruses. TLR-4 is the receptor for Gram-negative bacterial LPS, TLR-2 is the receptor for Gram-positive bacterial peptidoglycan and lipopeptide, and TLR-9 is the receptor for the unmethylated and phosphorylated cytosine-guanine oligonucleotide, CpG. (Curr. Opin. Immunol. 14:103-110, 2002) TLR3 can recognize double strand RNA virus, (Nature 433:887-892, 2005) TLR7 can recognize single strand RNA virus. (Proc Natl Acad Sci USA. 101(15):5598-5603, 2004) When TLR signal pathway was activated, the innate immunity or antiviral response can be induced for host protection.

[0013] In normal intestinal epithelial cells, TLR-3 and TLR-5 are constitutively expressed, while TLR2 and TLR4 are barely detectable. (Infection and Immunity 68:7010-7017, 2000) IL-8 is produced by the epithelial cells in response to the stimulation of the bacterial DNA. (FASEB J 17:1319-1321, 2003) In intestinal

epithelial cells, the bacterial CpG oligonucleotides are the ligands for the TLR-9.

Furthermore, the intestinal epithelial cells do not induce the inflammation in response to all foreign substances, i.e. the intestinal epithelial cells tolerate commensal microflora, while the intestinal epithelial cells provide danger signals to APCs under potential pathogenic conditions or autoimmune diseases, so as to induce the inflammation formed by T cells mediated by APCs. (Cell 118:229-241, 2004)

[0014] Under normal steady-state conditions, commensal bacteria recognizing by TLRs plays a crucial role in protection against gut injury. (Cell 118: 229-241, 2004)

But in human inflammatory bowel disease (IBD), a Th1-mediated pathological effect is thought to be due to aberrant mucosal immune response to the microflora.

(Gastroenterol. Clin. North Am. 31: 41-62, 2002) These findings reveal that TLRs control mucosal homeostasis between host-commensal.

[0015] It is well-known that the oral administration of Chinese herbal extract regulates the development of the immunity. It is very possible the Chinese herbal extract regulates the development of the immunity via the intestinal mucosal immunity. Dendrobium species is considered to be the most precious Chinese

herb. A Dendrobium species belongs to an orchid family, and its stem is the mainly medicinal part. It tastes a little sweet and brackish. Some Chinese medical codices disclose that the Dendrobium species is the curative for some illnesses such as mucosal disorders, stomach disorders and ophthalmic disorders. According to our previous research experience, it appears that the Dendrobii Herba is the most curative medicinal species.

[0016] The present invention provides a method for preparing the polysaccharide derived from Dendrobium and further provides the pharmaceutical use of the polysaccharide derived from Dendrobium.

SUMMARY OF THE INVENTION

[0017] It is an aspect of the present invention to provide a method for preparing a polysaccharide from a plant. The method includes steps of a) extracting the plant with a first alcohol to obtain a first extract, b) extracting the first extract with a solvent to obtain a second extract, and c) precipitating the second extract with a second alcohol to obtain the polysaccharide.

[0018] Preferably, the plant belongs to Genus Dendrobium.

[0019] Preferable, the first alcohol is a methanol.

[0020] Preferably the solvent is water.

[0021] Preferably, the second alcohol is an ethanol.

[0022] In accordance with the present invention, after extracting the first extract with the solvent in the step b), the method further includes steps of centrifuging and filtering so as to obtain the second extract.

[0023] It is another aspect of the present invention to provide a composition for treating an autoimmune disease and mucosal disorder. The composition includes a polysaccharide prepared from a plant, wherein the plant belongs to Genus Dendrobium, and an antigen associated with an induction of the autoimmune disease.

[0024] Preferably, the polysaccharide is prepared by the method provided in the present invention.

[0025] In accordance with the present invention, the autoimmune disease can be uveitis.

[0026] It is another aspect of the present invention to provide a method for treating an autoimmune disease and mucosal disorder in a mammal. The method includes a step of administering an antigen and a polysaccharide to the mammal, wherein the antigen is associated with an induction of the autoimmune disease, and the polysaccharide is prepared from a plant belonging to Genus Dendrobium.

[0027] In accordance with present invention, the administration of the polysaccharide can be performed in a manner of an oral administration.

[0028] In accordance with the present invention, the autoimmune disease can be uveitis.

[0029] It is another aspect of the present invention to provide a method for treating an autoimmune disease and mucosal disorder in a mammal. The method includes a step of administering a polysaccharide to the mammal, wherein the

polysaccharide is prepared from a plant belonging to Genus Dendrobium.

[0030] In accordance with the present invention, the administration of the polysaccharide is performed in a manner of an oral administration.

[0031] In accordance with the present invention, the autoimmune disease can be uveitis.

[0034] The above aspects and advantages of the present invention will become more readily apparent to those ordinarily skilled in the art after reviewing the following detailed description and accompanying drawings, in which:

BRIEF DESCRIPTION OF THE DRAWINGS

[0035] FIG. 1 is the PCR analysis showing mRNA expression of cytokines and antimicrobial peptides in the IEC-6 cells treated with the DeCaPS;

[0036] FIG. 2 is the PCR analysis showing the mRNA expressions of toll-like receptors in the IEC-6 cells treated with the DeCaPS;

[0037] FIG. 3 is a chart showing the effects on the mitogenic responses of ConA-stimulated splenocytes from the C3H mice orally treated with the DeCaPS;

[0038] FIG. 4 is a chart showing the effects on the mitogenic responses of GM-CSF co-stimulated bone marrow cells from the C3H mice orally treated with the DeCaPS;

[0039] FIG. 5 is the PCR analysis showing the cytokine-specific mRNA in Peyer's patch and splenocytes obtained from the C3H mice orally treated with the DeCaPS;

[0040] FIG. 6 is a chart showing the titers of the ovalbumin specific IgA in the intestinal lavage solution from the C57BL/6j mice orally immunized by the ovalbumin with or without administration of DeCaPS;

[0041] FIG. 7 is a chart showing the titers of the ovalbumin specific IgM in serum from the C57BL/6j mice orally immunized by the ovalbumin with or without administration of DeCaPS;

[0042] FIG. 8 is a chart showing the titers of the ovalbumin specific IgG in serum from the C57BL/6j mice orally immunized by the ovalbumin with or without administration of DeCaPS;

[0043] FIG. 9 is the PCR analysis showing the cytokine expressions of lymphocytes in Peyer's patch from the C57BL/6j mice;

[0044] FIG. 10 is a chart showing the DTH responses of the C57BL/6j mice;

[0045] FIG. 11A is a chart showing the electro-retinograms to a range of flash intensities for the mice of the normal group;

[0046] FIG. 11B is a diagram showing the histopathology of eyes of the mice in the normal group;

[0047] FIG. 12A is a chart showing the electro-retinograms to a range of flash intensities for the mice of the control group;

[0048] FIG. 12B is a diagram showing the histopathology of eyes of the mice in the control group;

[0049] FIG. 13A is a chart showing the electro-retinograms to a range of flash intensities for the mice of the DC-10 group;

[0050] FIG. 13B is a diagram showing the histopathology of eyes of the mice in the DC-10 group;

[0051] FIG. 14A is a chart showing the electro-retinograms to a range of flash intensities for the mice of the DC-40 group;

[0052] FIG. 14B is a diagram showing the histopathology of eyes of the mice in the DC-40 group;

[0053] FIG. 15A is a chart showing the electro-retinograms to a range of flash intensities for the mice of the DC-160 group;

[0054] FIG. 15B is a diagram showing the histopathology of eyes of the mice in the DC-160 group;

[0055] FIG. 16A is a chart showing the amplitudes of the a-waves in the FIGS. 11A, 12A, 13A, 14A and 15A; and

[0056] FIG. 16B is a chart showing the amplitudes of the b-waves in the FIGS. 11A, 12A, 13A, 14A and 15A.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

[0057] The invention is described more specifically with reference to the following embodiments. It is to be noted that the following descriptions of preferred embodiments of this invention are presented herein for the purpose of illustration and description only; it is not intended to be exhaustive or to be limited to the precise form disclosed.

[0058] The present provides a method for preparing a polysaccharide from a plant, including steps of a) extracting the plant with a first alcohol to obtain a first extract, b) extracting the first extract with a solvent to obtain a second extract, and c)

precipitating the second extract with a second alcohol to obtain the polysaccharide.

EXAMPLE I

Preparation of Polysaccharides from Dendrobium

[0059] 10 kg of fresh Dendrobii Herba was dried to form 2.92 kg of the dry material, and then the dry material was immersed and extracted with 73.3 L of methanol to form 136.79 gram of crude extract and residual. The crude extract was treated with a de-methanol process, and then immersed in water overnight. The reaction solution was extracted with warm water at the temperature ranged from 55.degree. C. to 60.degree. C. for 30 minutes, and the centrifuged by the centrifuge (ER-RC13 C-124, HITACH) at 5,000 rpm and 10.degree. C. After the centrifugation, the supernatant was filtered through 6 .mu.m filter paper. The filtrate was precipitated with ethanol, so that 20.684 g of polysaccharides (hereafter named DeCaPS) were obtained.

EXAMPLE II**Immune Responses of IEC-6 Cells Treated by the DeCaPS****[0060] a. Culture of IEC-6 Cells**

[0061] IEC cells, which were originated from the normal rat small intestine were cultured in DMEM medium containing 5% fetal bovine serum, 4.5 g/L glucose, 5 .mu.g/ml bovine insulin and 2 mM L-glutamine at the incubator with 5% CO.sub.2 at 37.degree. C.

[0062] b. Treating the IEC-6 Cells with the DeCaPS

[0063] When the IEC-6 cells were confluent in the 6-well plate, the IEC cells were treated with the DeCaPS at the concentrations of 1 .mu.g/ml, 10 .mu.g/ml and 100 .mu.g/ml, respectively, for 6 hours.

[0064] c. The Isolation of Total RNA from the IEC Cells Treated with the DeCaPS

[0065] After the treatment with the DeCaPS, the IEC cells were harvested and suspended in the 1 ml Ultraspec TM RNA isolation Kit (Biotex laboratories Inc.(USA)), and the total RNA was obtained by following the standard protocol of the kit. The obtained total RNA is quantitatively determined.

[0066] d. Reverse-Transcriptions and PCR Analysis of Cytokines and Toll-Like Receptors

[0067] The reaction having a reaction volume of 26.5 .mu.l and including 0.1 g of oligo-dT, 5 .mu.g of the obtained total RNA and DEPC-treated sterilized water was performed at 70.degree. C. for 10 minutes. Then, the reaction was added with 4 .mu.l of 10 mM dNTP, 0.5 .mu.l of rRNasin, 1 .mu.l of AMV (Avian Myeloblastosis virus) reverse transcriptase (10 unit) and 8 .mu.l of 5.times.RT buffer so as to have the total volume of 40 .mu.l, and to be incubated at 42.degree. C. for 60 minutes, then at 90.degree. C. for 5 minutes, and thereby the cDNA products were obtained by the reverse transcription reaction. Subsequently, 0.5 mu of 10 mM dNTP, 0.5 .mu.l of 10.times. Prozyme buffer, the primers (0.8 .mu.l of 5 .mu.M sense DNA and 0.8 .mu.l of 5 .mu.M antisense DNA) and the sterilized water were added to 2.5 .mu.l of the cDNA products to form a total volume of

25 .mu.l, and to be incubated in the DNA thermal cycler (Perkin-Elmer-Cetus), and thereby the PCR was performed for 35 cycles. The primer sequences including sense and antisense primers [SEQ ID NOS: 1-32] for the target genes and the corresponding annealing temperatures are listed in the following table I.

TABLE I

SEQ ID NO.	Target gene (PCR product size)	Annealing temperature (.degree. C.)
1, 2	β -actin (510 bps)	57
3, 4	IL-1 β 563 bps	61
5, 6	IL-2 (502 bps)	61
7, 8	IL-4 (399 bps)	61
9, 10	IL-6 (638 bps)	61
11, 12	IL-10 (455 bps)	61
13, 14	TNF- α (308 bps)	57
15, 16	IFN- γ (460 bps)	61
17, 18	TGF- β (525 bps)	61
19, 20	Rat α -defensin (900 bps)	59
21, 22	Rat angiogenin (900 bps)	59 (hot start)
23, 24	TLR2 (495 bps)	55
25, 26	TLR4 (508 bps)	57
27, 28	TLR5 (737 bps)	59
29, 30	TLR7 (729 bps)	59
31, 32	TLR9 (725bps)	60

[0068] Please refer to FIG. 1, which shows the mRNA expressions of cytokines and antimicrobial peptides in the IEC-6 cells treated with the DeCaPS. After being treated with 1 .mu.g/ml, 10 .mu.g/ml and 100 .mu.g/ml of the DeCaPS for 6 hours, the mRNA expressions of TNF-.alpha. [SEQ ID NO: 33] in the IEC-6 cells were suppressed about 29%, 20% and 30%; however, the mRNA expressions of IL-10 [SEQ. ID NO: 34] in the IEC-cells were enhanced up to 9.89-fold, 6.55-fold and 1.77-fold, respectively. It is shown that the intestinal epithelial cells can be induced by the DeCaPS to provide tolerance signals to other cells in the immune system, so as to trigger the TH2/3 pathways. In addition, after being treated with 1 .mu.g/ml, 10 .mu.g/ml and 100 .mu.g/ml of the DeCaPS for 6 hours, the mRNA expressions of a-defensin [SEQ ID NO: 35] were suppressed about 42%, 33% and 51%; however, the MRNA expressions of angiogenin [SEQ ID NO: 36] were enhanced up to 1.96-fold, 1.42-fold and 1.70-fold, respectively.

[0069] Please refer to FIG. 2, which shows the mRNA expressions of toll-like receptors in the IEC-6 cells treated with the DeCaPS. In the IEC-6 cells, toll-like receptors, TLR2 [SEQ ID NO: 37], TNR4 [SEQ ID NO: 38], TLR7 [SEQ ID NO: 39] and TLR9 [SEQ ID NO: 40], were expressed without the DeCaPS treatment. After

being treated with 1 .mu.g/ml, 10 .mu.g/ml and 100 .mu.g/ml of the DeCaPS for 6 hours, the mRNA expressions of TLR4 [SEQ ID NO: 38] in the IEC-6 cells were suppressed about 6%, 16% and 13%, the mRNA expressions of TLR5 [SEQ ID NO: 41] in the IEC-6 cells were suppressed about 13%, 37% and 35%, and the mRNA expressions of TLR7 [SEQ ID NO: 39] in the IEC-6 cells were suppressed about 11%, 7% and 48%, respectively; however, the mRNA expressions of TLR9 [SEQ ID NO: 40] were enhanced up to 2.21-fold, 1.09-fold and 3.56-fold, respectively. It is shown that the polysaccharides derived from Dendrobium not only suppress the expression of TNF-.alpha. [SEQ ID NO: 33], but also suppress the expressions of TLR4 [SEQ ID NO: 38], TLR5 [SEQ ID NO: 41], TLR7 [SEQ ID NO: 39] and TLR9 [SEQ ID NO: 40].

EXAMPLE III

Effects on Immune System of Animal Model by Oral Administration of DeCaPS

[0070] The C3H mice (13 weeks old) were respectively fed with the DeCaPS, prepared from the Example I, via the drinking water at the dosages of 10 mg/kg/day, 50 mg/kg/day and 250 mg/kg/day for 5 days, and then were sacrificed to obtain the spleens and the bone marrows thereof.

[0071] a. MTT Assays for Concanavalin A-Stimulated Splenocytes from the C3H Mice

[0072] The splenocytes (4×10^5 cells/well) obtained from the C3H mice treated with the DeCaPS were cultured in RPMI-1640 medium containing 10% fetal bovine serum and 1 and 5 $\mu\text{g/ml}$ concanavalin A (ConA) for 72 hours, and then were treated with 1 mg/ml MTT for 3 hours. Then, the cell cultures were respectively added with the lysis buffer containing 50% DMF and 20% SDS, and the reactions were performed for 16 hours. The absorptions at 570 nm for the reactions were analyzed to denote the growth index as shown in FIG. 3.

[0073] Please refer to FIG. 3, which shows the effects on the mitogenic responses of ConA-stimulated splenocytes from the C3H mice orally treated with the DeCaPS. After the C3H mice were orally treated with the DeCaPS at the dosages of 10 mg/kg/day, 50 mg/kg/day and 250 mg/kg/day for 5 days, the responses to the 1 $\mu\text{g/ml}$ ConA stimulation for the splenocytes were suppressed about 28%, 49% and 30%.

[0074] b. MTT Assays for GM-CSF-Stimulated Bone Marrow Cells Obtained from the C3H Mice

[0075] The bone marrow cells were obtained from the legs of the C3H mice treated with the DeCaPS. The bone marrow cells were cultured in α -MEM medium containing 2% FCS and 4 ng/ml GM-CSF for 72 hours, and the positive control group of the bone marrow cells was cultured in α -MEM medium containing 2% FCS and 20 ng/ml GM-CSF for 72 hours. Then, the medium was replaced by RPMI-1640 containing 1mg/ml MTT and 2% FCS for treating the bone marrow cells for 3 hours, and the cells were further treated with MTT lysis buffer (20% SDS and 50% DMF at pH4.5) and incubated overnight. The absorptions at 570 nm for the reactions were analyzed to denote the growth index as shown in FIG. 4.

[0076] Please refer to FIG. 4, which shows the effects on the mitogenic responses of GM-CSF-stimulated bone marrow cells obtained from the C3H mice orally treated with the DeCaPS. After the C3H mice were respectively and orally treated with the DeCaPS at the dosages of 10 mg/kg/day, 50 mg/kg/day and 250

mg/kg/day for 5 days, the mitogenic responses to the GM-CSF stimulation for the bone marrow cells were enhanced up to 1.71-fold, 1.62-fold and 1.76-fold.

[0077] c. mRNA Expressions of Cytokines in Peyer's Patch and Splenocytes

Obtained from the C3H Mice

[0078] The Peyer's patch and splenocytes were obtained from the C3H mice treated with the DeCaPS. The PCR analysis of the cytokine-specific mRNA in Peyer's patch and the splenocytes was performed by the method described in Example II, and the PCR results were shown in FIG. 5.

[0079] Please refer to FIG. 5, which shows the PCR results of cytokine-specific mRNA in Peyer's patch and splenocytes obtained from the C3H mice orally treated with the DeCaPS, wherein the lanes 1-4 denote the C3H mice orally treated with the DeCaPS at the dosages of 0 mg/kg/day, 10 mg/kg/day, 50 mg/kg/day and 250 mg/kg/day, respectively.

[0080] After the C3H mice were orally treated with the DeCaPS for 5 days, the mRNA expressions of IL-4 [SEQ ID NO: 42], IL-6 [SEQ ID NO: 43], IL-1.beta.

[SEQ ID NO: 44], IFN- γ . [SEQ ID NO: 45] and TGF- β . [SEQ ID NO: 46] in Peyer's patch were increased as shown in FIG. 5. After the C3H mice were orally treated with the DeCaPS at the dosages of 10 mg/kg/day and 50 mg/kg/day for 5 days, the mRNA expressions of IL-4 [SEQ ID NO: 42] were enhanced up to 2.64-fold and 2.46-fold, respectively. After the C3H mice were orally treated with the DeCaPS at the dosages of 10 mg/kg/day, 50 mg/kg/day and 250 mg/kg/day for 5 days, the mRNA expressions of IL-6 [SEQ ID NO: 43] were respectively enhanced up to 2.61-fold, 3.99-fold and 5.35-fold, the mRNA expressions of IL- β . [SEQ ID NO: 44] were respectively enhanced up to 1.83-fold, 2.00-fold and 1.11-fold, the mRNA expressions of IFN- γ . [SEQ ID NO: 45] were respectively enhanced up to 3.47-fold, 5.47-fold and 5.57-fold, and the mRNA expressions of TGF- β . [SEQ ID NO: 46] were respectively enhanced up to 2.38-fold, 2.54-fold and 1.89-fold. Since the Peyer's patch is the main site for the determination of immune responses to the antigens, the Th2 pathway is triggered in response to the increment of IL-4 [SEQ ID NO: 42] and IL-6 [SEQ ID NO: 43], T helper cells are activated by the increments of IL- β . [SEQ ID NO: 44] to secrete cytokines, such as TGF- β . [SEQ ID NO: 46], and the IgA class switched B cells are induced by TGF- β . [SEQ ID NO: 46] to secrete IgA so as to suppress certain immune responses.

[0081] After the C3H mice were orally treated with the DeCaPS for 5 days, the mRNA expressions of IL-1. β . [SEQ ID NO: 44], IL-4 [SEQ ID NO: 42] and TGF- β . [SEQ ID NO: 46] in the splenocytes were suppressed as shown in FIG.

5. After the C3H mice were orally treated with the DeCaPS at the dosages of 10 mg/kg/day, 50 mg/kg/day and 250 mg/kg/day for 5 days, the mRNA expressions of IL-1. β . [SEQ ID NO: 44] were suppressed about 32%, 43% and 79%, and the mRNA expressions of IL-4 [SEQ ID NO: 42] were suppressed about 7%, 67% and 14%. After the C3H mice were orally treated with the DeCaPS at the dosages of 10 mg/kg/day, 50 mg/kg/day and 250 mg/kg/day for 5 days, the mRNA expressions of TGF- β . [SEQ ID NO: 46] were suppressed about 27% and 44%, and the mRNA expressions of IFN- γ . [SEQ ID NO: 45] were suppressed about 33% and 97%, respectively. However, after the C3H mice were orally treated with the DeCaPS at the dosages of 10 mg/kg/day, 50 mg/kg/day and 250 mg/kg/day for 5 days, the mRNA expressions of IL-6 [SEQ ID NO: 43] were enhanced up to 1.84-fold, 1.4-fold and 1.23-fold. It is to be noted that there are not significant cellular responses to the DeCaPS treatment, and this phenomenon is consistent with the responses to the ConA stimulations.

EXAMPLE IV

Enhancement of Oral Tolerance in Animal Model by DeCaPS

[0082] Since it is well-known that animals can be immunized by ovalbumin (OVA) to establish the animal model for the autoimmune disease (The Journal of Pharmacology and Experimental Therapeutics 288:849-857, 1999), the present invention establishes the mice with the OVA-induced autoimmune disease, and the enhancement of Oral tolerance in the mice with the OVA-induced autoimmune disease by DeCaPS is studied as follows. The C57/BL6j mice (65 weeks old) were orally treated with 0.5 mg/ml ovalbumin (OVA) in the drinking water at the day 4 and day 5, and the DeCaPS was administrated to the C57/BL6j mice as the adjuvant at the dosages of 10 mg/kg/day, 40 mg/kg/day and 160 mg/kg/day at the days 3-7. 50 .mu.g of OVA adjuvanted with CFA was administrated to the mice as the antigen via i.p. injection at the day 8, IgG and IgM in response to OVA in the blood collection were detected at the day 22, 50 .mu.g of OVA was administrated to the mice as the antigen via i.p. injection at the day 24, the antibodies in response to OVA were detected at the day 30, the mice were sacrificed to obtain the intestinal lavage solution of the intestinal mucosa and the lung mucosa at day

32, and then IgA in response to OVA was detected. The detailed experimental procedures are illustrated in the following table II.

TABLE II

Group of Mice	Mice Number	Blood Collected	Oral OVA	Oral DeCaPS	OVA + CFA injection	OVA injection
Normal	5	Day 0, 22,30	-	-	Day 8	Day 24
Control	6	Day 0, 22, 30	Day 4-5	-	Day 8	Day 24
Positive Control	6	Day 0, 22, 30	Day 1-5	-	Day 8	Day 24
DC-1 (10 mg/kg/day)	5	Day 0, 22, 30	Day 4-5	Day 3-7	Day 8	Day 24
DC-2 (40 mg/kg/day)	5	Day 0, 22, 30	Day 4-5	Day 3-7	Day 8	Day 24
DC-3 (160 mg/kg/day)	5	Day 0, 22, 30	Day 4-5	Day 3-7	Day 8	Day 24

[0083] The intestinal lavage solution of the intestinal mucosa was obtained from the mice at the day 32, and the titer of the ovalbumin-specific IgA in the intestinal lavage solution was determined as shown in FIG. 6. Referring to FIG. 6, the IgA

secreted from the intestinal mucosa was enhanced up to 1.9-fold ($p<0.01$) in the C57/BL6j mice treated by the OVA with DeCaPS (40 mg/kg/day), and the IgA secreted from the intestinal mucosa was enhanced up to 4.35-fold ($p<0.001$) in the C57/BL6j mice treated by the OVA with DeCaPS (160 mg/kg/day).

[0084] The IgM antibodies in the blood collection at the day 22 and the day 30 were determined as shown in FIG. 7. With comparison to the IgM antibody level expressed in the control group at the day 22, the IgM antibody level expressed in the normal group was suppressed about 49% ($p<0.05$), the IgM antibody level expressed in the positive control group was suppressed about 87% ($p<0.05$), the IgM antibody level expressed in the DC-1 group was suppressed about 46% ($p<0.02$), the IgM antibody level expressed in the DC-2 group was suppressed about 88% ($p<0.001$), and the IgM antibody level expressed in the DC-3 group was suppressed about 86% ($p<0.001$). Furthermore, with comparison to the IgM antibody level expressed in the control group at the day 30, the IgM antibody level expressed in the normal group was suppressed about 47% ($p<0.01$), the IgM antibody level expressed in the positive control group was suppressed about 72% ($p<0.001$), the IgM antibody level expressed in the DC-1 group was suppressed about 38% ($p<0.01$), the IgM antibody level expressed in the DC-2 group was

suppressed about 58% ($p<0.01$), and the IgM antibody level expressed in the DC-3 group was suppressed about 71% ($p<0.001$).

[0085] The IgG antibodies in the blood collection at the day 22 and the day 30 were determined as shown in FIG. 8. With comparison to the IgG antibody level expressed in the control group at the day 22, the IgG antibody level expressed in the positive control group was suppressed about 94% ($p<0.01$), the IgG antibody level expressed in the DC-1 group was suppressed about 67% ($p<0.01$), the IgG antibody level expressed in the DC-2 group was suppressed about 76% ($p<0.01$), and the IgG antibody level expressed in the DC-3 group was suppressed about 88% ($p<0.001$). Furthermore, with comparison to the IgG antibody level expressed in the control group at the day 30, the IgG antibody level expressed in the positive control group was suppressed about 75% ($p<0.001$), the IgG antibody level expressed in the DC-2 group was suppressed about 62% ($p<0.001$), and the IgG antibody level expressed in the DC-3 group was suppressed about 63% ($p<0.001$).

[0086] In addition, Peyer's patch was obtained from the mice at the day 32, and the RNA of the lymphocytes in Peyer's patch was extracted and analyzed for the

cytokines. The cytokine expressions of lymphocytes in Peyer's patch were determined as shown in FIG. 9, wherein the lane A denotes the normal group, lane B denotes the control group, lane C denotes the positive control group, lane D denotes the DC-1 group, lane E denotes the DC-2 group, and lane F denotes the DC-3 group. The PCR amplification was programmed for denature at 94.degree. C. for 45 seconds, the annealing was performed at 61.degree. C. for 45 seconds and the extension was performed at 72.degree. C. for 1 minute. There were 37 cycles in the PCR program. The PCR products were separated by electrophoresis in 2% agarose gel, and the gel was stained with ethidium bromide and visualized under UV light.

[0087] It is to be noted that with comparison to the control group, the TGF- β . [SEQ ID NO: 46] level expressed in the DC-3 group was suppressed about 30%, the IL-4 [SEQ ID NO: 42] level expressed in the DC-3 group was enhanced up to 2.2-fold, and the IL-10 [SEQ ID NO: 34] level expressed in the DC-3 group was enhanced up to 1.3-fold. With comparison to the control group, the IL-2 [SEQ ID NO: 47] level expressed in the DC-2 group was enhanced up to 1.4-fold, and the IL-4 [SEQ ID NO: 42] level expressed in the DC-2 group was enhanced up to 1.6-fold. Moreover, with comparison to the control group, the IL-4 [SEQ ID NO: 42]

level expressed in the DC-1 group was enhanced up to 1.4-fold, and the IL-6 [SEQ ID NO: 43] level expressed in the DC-1 group was enhanced up to 1.3-fold.

[0088] The mRNA expressions of IL-4 [SEQ ID NO: 42] and IL-6 [SEQ ID NO: 43] were enhanced in the lymphocytes in Peyer's patch. It is to be emphasized that since Payer's patch is the site for the determination of immune responses to the antigens, when the antigen-primed T or B cells back to intestine through homing mechanism, the Th2 pathway is activated to produce antibodies in response to the increments of IL-4 [SEQ ID NO: 42] and IL-6 [SEQ ID NO: 43], the IgG and IgM expressions in the serum are suppressed so as to suppress the allergenic responses, and the oral tolerance is achieved.

[0089] In view of aforesaid descriptions, the mitogenic response of T cells in the spleen is suppressed by the polysaccharides of the Dendrobium; however, the granulocytes and macrophages in bone marrow are activated by the polysaccharides of the Dendrobium. In other words, T cells activated in autoimmune diseases can be suppressed by the oral administration of the polysaccharides of the Dendrobium; however, the granulocyte lineage cells are

activated by the oral administration of the polysaccharides of the Dendrobium for preventing from pathogen invasions by innate immunity promotion.

[0090] With regard to the intestinal immunity, the mRNA expression of IL-4 [SEQ ID NO: 42], IL-6 [SEQ ID NO: 43], IL-1 β . [SEQ ID NO: 44], IFN- γ . [SEQ ID NO: 45] and TGF- β . [SEQ ID NO: 46] are enhanced to trigger the Th2/Th3 pathways in the intestine, so as to form the oral tolerance. According to the expressions of TNF- α . [SEQ ID NO: 33], TLR2 [SEQ ID NO: 37], TLR4 [SEQ ID NO: 38], TLR5 [SEQ ID NO: 41] and TLR7 [SEQ ID NO: 39] in the IEC-6 cells treated with the polysaccharides of Dendrobium, it is shown that the signals mediated by the toll-like receptors can be regulated to be immuno-supressed or anti-inflammatory response by the polysaccharides of Dendrobium, so as to enhance the oral tolerance-associated signals, i.e. the Th2/3 pathways and maintain mucosal homeostasis during pathogenic status such as autoimmune disease and mucosal disorder. Interestingly, the expression of angiogenin, which has the antimicrobial activities, can be induced by the administration of the polysaccharides of Dendrobium as described in Example II. Accordingly, the oral administration of the polysaccharides of Dendrobium not only enhances the oral tolerance, but also triggers the innate immunity mechanisms for preventing from

pathogen invasions. According to these findings, polysaccharides of *Dendrobium* may have beneficial use of autoimmune disease and mucosal disorder such as inflammatory bowel disease (IBD) for releasing prolonged inflammation and further injury.

[0091] It has been disclosed that the angiogenins have the activities against not only enteric microbes such as *Enterococcus faecalis* but also microbes that cause systematic infections in humans such as *Candida albicans* and *Streptococcus pneumoniae*. (Nat. Immunol. 4:269-273, 2003) In the clinical studies, leucorrhea in women is a polymicrobial, superficial vaginal infection. The infection frequently caused by bacteria such as *G. vaginalis*, *Candida* spp, *C. albicans*, *T. vaginalis*, *Streptococcus* group D, *Streptococcus* β hemolytic, *E. coli*, and *Klebsiella* spp etc. (Salud publica Mex vol.45 suppl. 5, pS694-S697, 2003). It is obvious that the leucorrhea in women caused by microbes such as *Candida albicans* could be relieved by the activities of angiogenins, which are induced by the oral administration of the polysaccharides of *Dendrobium*.

EXAMPLE IV

Treatment of Autoimmune Disease by DeCaPS

[0092] Experimental autoimmune uveitis is a T-cell-mediated autoimmune disease that serves as a model for several ocular autoimmune diseases. It is identified that the experimental autoimmune uveitis is induced by the immunization with interphotoreceptor retinoid binding protein (IRBP), a 140-kD glycolipoprotein. It is also identified that the autoimmune uveitis can be induced by the IRBP peptide 1-20 (SEQ ID NO. 48: GPTHLFQPSLVLDMAKVLLD), the amino acids 1-20 of IRBP. In addition, the IRBP peptide 1-20 is conserved in mice and human. (Investigative Ophthalmology & Visual Science 41(1):127-131, 2000) Hence, the IRBP peptide was administrated to the C57BL/6j mice to establish the autoimmune disease model in the present invention.

[0093] a. Synthesis of IRBP Peptide

[0094] The IRBP peptide (SEQ ID NO. 48: GPTHLFQPSLVLDMAKVLLD) was synthesized on the peptide synthesizer (PS3) using Fmoc Chemistry, purified by the Agilent HPLC and identified by the Bruker esquire 2000 MS.

[0095] b. Administrations of IRBP and DeCaPS

[0096] The specific-pathogen-free (SPF) mice, C57BL/6j mice, were randomly grouped into the normal group, the control group, the DC-10 group, the DC-40 group and the DC-160 group, wherein there were 6 mice in each group. The human IRBP peptide [SEQ ID NO: 48] (150 .mu.g/mouse) and CFA (1:1, vol/vol) in 0.2 ml emulsion was administrated to each SPF mouse in the groups, except the normal group, via i.p. injection. The mice in the DC-10 group were fed with the DeCaPS at the dosage of 10 mg/kg/day for 28 days, the mice in the DC-40 group were fed with the DeCaPS at the dosage of 40 mg/kg/day for 28 days, and the mice in the DC-160 group were fed with the DeCaPS at the dosage of 160 mg/kg/day via the drinking water for 28 days. The retinal functions of the mice were determined by the electro-retinogram (ERG) analysis, and the mice were sacrificed at the day 28 to obtain the eyes thereof for the bioassays and histopathology determination. The experimental procedures are illustrated in the following table III.

TABLE III Treatment of autoimmune disease with/without DeCaPS in animal model

Group	Amount of Mice	ERG determination At day 0	Immunized with IRBP At day 1	DeCaPS (days 1-28) mg/kg/day	DTH assay (days 21 and 23)	ERG & histopathology At day 28
Normal	6	+	—	—	+	+
Control	6	+	+	—	+	+
DC-10	6	+	+	10	+	+
DC-40	6	+	+	40	+	+
DC-160	6	+	+	160	+	+

[0097] c. Delayed-Type Hypersensitivity (DTH) Test

[0098] On the day 21 after immunization of IRBP peptide, the mice were injected subcutaneously with 20 .mu.g of the IRBP peptide emulsion in IFA (20 .mu.l), into

the left footpad. The right footpad was injected with IFA. After 48 hours, the thicknesses of the footpads were measured with a caliper.

[0099] Please refer to FIG. 10, which shows the DTH responses of the mice. The DTH responses were shown by the footpad increments in water volume relative to the footpad increment of the mice injected with the phosphate buffer saline (PBS). It is to be noted that the footpad increment of the mice co-treated with the DeCaPS (160 mg/kg/day) is significantly decreased.

[0100] d. Electro-Retinogram (ERG) Analysis

[0101] The mice were dark-adapted for 2 hours, and then anesthetized with sodium pentobarbital. Then, the corneas of the mice were anesthetized, and the pupils were dilated. The retinas were stimulated with a flash of light, and the responses of the retina to the flash of light were recorded as the electro-retinograms. The electro-retinograms show the action of the photoreceptors and the functions of the proximal retina such as bioplar and Muller cells. The electro-retinograms are used for reflecting the state of the entire retina. In the electro-retinogram, there is typically a negative-going a-wave, followed by a

positive-going b-wave. The leading edge of the a-wave provides a direct measure of the activities of the cone and rod cells in the photoreceptor layer, while the b-wave provides a reflection of the action of bipolar cells in the inner nuclear layer.

[0102] For the two mice randomly selected from the normal group at the day 28, the results of the electro-retinogram analysis were respectively shown in the upper part and the lower part of FIG. 11A, wherein the flash onset was at time 100 millisecond and the flash duration was approximately 600 milliseconds. For the two mice randomly selected from the normal group at the day 28, the histopathology of the eyes were respectively shown in the upper part and the lower part of FIG. 11B (400.times. magnification), wherein the photoreceptor layer was denoted as "P", the outer nuclear layer was denoted as "ONL", and the inner nuclear layer was denoted as "INL".

[0103] For the two mice randomly selected from the control group at the day 28, the results of the electro-retinogram analysis were respectively shown in the upper part and the lower part of FIG. 12A, wherein the flash onset was at time 100 millisecond and the flash duration was approximately 600 milliseconds. For the two mice randomly selected from the control group at the day 28, the

histopathology of the eyes were respectively shown in the upper part and the lower part of FIG. 12B (400.times. magnification). It was shown that due to the immunization of the IRBP peptide, the photoreceptor layer (P), the outer nuclear layer (ONL) and the inner nuclear layer (INL) of the mice in the control group were severely damaged.

[0104] For the two mice randomly selected from the DC-10 group at the day 28, the results of the electro-retinogram analysis were respectively shown in the upper part and the lower part of FIG. 13A, wherein the flash onset was at time 100 millisecond and the flash duration was approximately 600 milliseconds. For the two mice randomly selected from the DC-10 group at the day 28, the histopathology of the eyes were respectively shown in the upper part and the lower part of FIG. 13B (400.times. magnification). It was shown that after the immunization of the IRBP peptide and the oral administration of the DeCaPS (10 mg/kg/day) for 28 days, the photoreceptor layer (P), the outer nuclear layer (ONL) and the inner nuclear layer (INL) of the mice in the DC-10 group were severely damaged.

[0105] For the two mice randomly selected from the DC-40 group at the day 28, the results of the electro-retinogram analysis were respectively shown in the upper part and the lower part of FIG. 14A, wherein the flash onset was at time 100 millisecond and the flash duration was approximately 600 milliseconds. For the two mice randomly selected from the DC-40 group at the day 28, the histopathology of the eyes were respectively shown in the upper part and the lower part of FIG. 14B (400.times. magnification). It was shown that after the immunization of the IRBP peptide and the oral administration of the DeCaPS (40 mg/kg/day) for 28 days, the photoreceptor layer (P), the outer nuclear layer (ONL) and the inner nuclear layer (INL) of the mice in the DC-40 group were slightly damaged.

[0106] For the two mice randomly selected from the DC-160 group at the day 28, the results of the electro-retinogram analysis were respectively shown in the upper part and the lower part of FIG. 15A, wherein the flash onset was at time 100 millisecond and the flash duration was approximately 600 milliseconds. For the two mice randomly selected from the DC-160 group at the day 28, the histopathology of the eyes were respectively shown in the upper part and the lower part of FIG. 15B (400.times. magnification). It was shown that after the

immunization of the IRBP peptide and the oral administration of the DeCaPS (160 mg/kg/day) for 28 days, the photoreceptor layer (P), the outer nuclear layer (ONL) and the inner nuclear layer (INL) of the mice in the DC-160 group were not damaged.

[0107] Accordingly, after the immunization with the IRBP peptide, it was observed that there were more cell deaths in the inner nuclear layer, but the dead cell did not sink on the outer nuclear layer. It was also observed that the cells in the inner nuclear layer were damaged by the IRBP peptide treatment, but the cell peeling off from the inner nuclear layer was significantly prevented by the co-treatment with the DeCaPS at the dosage of 160 mg/kg/day.

[0108] In addition, the a-waves respectively shown in FIGS. 11A, 12A, 13A, 14A and 15A were compared in FIG. 16A ($p < 0.02$ for the statistics), and the b-waves respectively shown in FIGS. 11A, 12A, 13A, 14A and 15A were compared in FIG. 16B ($p < 0.02$ for the statistics). It is to be emphasized that the a-wave of the mice in the control group immunized with IRBP peptide were much lower than that of the mice in the normal group; however, with comparison to the control group, the cell peeling off from the inner nuclear layer of the mice co-treated with the

DeCaPS (40 mg/kg/day) is significantly prevented. Furthermore, the amplitudes of the a-wave and the b-wave of the mice co-treated with the DeCaPS (160 mg/kg/day) are nearly equal to those of the mice of the normal group. It is to be further emphasized that the damages from the immunization of IRBP peptide can be rescued by the co-treatment with the polysaccharides (160/mg/kg/day) prepared from the Dendrobium. In other words, the oral administration of the polysaccharides prepared from the Dnedrobium can prevent the retinal from the inflammation induced by the IRBP.

[0109] According to the foregoing experiments, the present invention provides a composition for treating an autoimmune disease and mucosal disorder, wherein the composition includes polysaccharides prepared from Dendrobium and an antigen associated with an induction of the autoimmune disease due to that the polysaccharides prepared from Dendrobium can enhance the oral tolerance induction and innate immunity promotion. Furthermore, the present invention provide a composition just including the polysaccharides prepared from Dendrobium for treating an autoimmune disease and mucosal disorder due to that the polysaccharides prepared from Dendrobium can induce the oral tolerance and promote innate immunity alone.

[0110] While the invention has been described in terms of what is presently considered to be the most practical and preferred embodiments, it is to be understood that the invention needs not be limited to the disclosed embodiment. On the contrary, it is intended to cover various modifications and similar arrangements included within the spirit and scope of the appended claims which are to be accorded with the broadest interpretation so as to encompass all such modifications and similar structures.

DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME __1__ DE __2__

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**

THIS IS VOLUME __1__ OF __2__

NOTE: For additional volumes please contact the Canadian Patent Office.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. Use of polysaccharides extracted from *Dendrobii Herba* in the manufacture of a medicament for treating autoimmune uveitis in a mammal, wherein the polysaccharides are prepared by a process comprising the steps of:
 - (a) extracting *Dendrobii Herba* with methanol to obtain a first extract;
 - (b) extracting the first extract with water to obtain a second extract; and
 - (c) precipitating the second extract with ethanol to obtain the polysaccharides.
2. The use of claim 1, wherein the medicament is formulated for oral administration.
3. A pharmaceutical composition for treating autoimmune uveitis in a mammal, comprising polysaccharides extracted from *Dendrobii Herba* in admixture with a pharmaceutically acceptable carrier or diluent, wherein the polysaccharides are prepared by a process comprising the steps of:
 - (a) extracting *Dendrobii Herba* with methanol to obtain a first extract;
 - (b) extracting the first extract with water to obtain a second extract; and
 - (c) precipitating the second extract with ethanol to obtain the polysaccharides.
4. The pharmaceutical composition of claim 3, which is formulated for oral administration.
5. A commercial package comprising polysaccharides extracted from *Dendrobii Herba*, together with instructions for use to treat autoimmune uveitis in a mammal, wherein the polysaccharides are prepared by a process comprising the steps of:
 - (a) extracting *Dendrobii Herba* with methanol to obtain a first extract;
 - (b) extracting the first extract with water to obtain a second extract; and
 - (c) precipitating the second extract with ethanol to obtain the polysaccharides.

6. The commercial package of claim 5, wherein the polysaccharides are formulated for oral administration.

Application number / numéro de demande: CN06/00895

Figures: 3, 4, 6, 11b, 12b, 13b, 14b, 15b

Pages: _____

Unscannable item(s)

received with this application

To inquire if you can order a copy of the unscannable items, please visit the
CIPO WebSite at [HTTP://CIPO.GC.CA](http://CIPO.GC.CA)

Item(s) ne pouvant être balayés

Documents reçus avec cette demande ne pouvant être balayés.

Pour vous renseigner si vous pouvez commander une copie des items ne
pouvant être balayés, veuillez visiter le site web de l'OPIC au [HTTP://CIPO.GC.CA](http://CIPO.GC.CA)




Fig. 1

BEST AVAILABLE COPY

DeCaPS	0	1	10	100 (μ g/ml)
TLR2	1	0.94	1.03	0.94
TLR4	1	0.94	0.84	0.69
TLR5	1	0.87	0.63	0.65
TLR7	1	0.89	0.93	0.52
TLR9	1	2.21	1.09	3.56
β -actin	1	0.87	0.94	0.91

Fig. 2




Fig. 5




Fig. 7




Fig. 8

	A	B.	C.	D.	E.	F.
IL-2						
	2.1	1.0	0.5	1.2	1.4	1.1
IL-4						
	2.0	1.0	2.3	1.4	1.6	2.2
IL-6						
	1.5	1.0	1.9	1.3	1.1	1.2
IL-10						
	1.2	1.0	1.5	1.2	1.2	1.3
TGF- β						
	1.0	1.0	1.5	1.0	0.9	0.7
β -actin						

Fig. 9




Fig. 10




Fig. 11A

13 of 22





Fig. 12A





Fig. 13A





Fig. 14A






Fig. 15A

a wave**Fig. 16A**




Fig. 16A

b wave




Fig. 16B