



(86) Date de dépôt PCT/PCT Filing Date: 2014/05/27  
(87) Date publication PCT/PCT Publication Date: 2014/12/04  
(45) Date de délivrance/Issue Date: 2021/09/07  
(85) Entrée phase nationale/National Entry: 2015/11/19  
(86) N° demande PCT/PCT Application No.: IL 2014/050477  
(87) N° publication PCT/PCT Publication No.: 2014/191995  
(30) Priorité/Priority: 2013/05/27 (US61/827,685)

(51) Cl.Int./Int.Cl. A61K 8/66 (2006.01),  
A61K 8/22 (2006.01), A61K 8/60 (2006.01),  
A61Q 19/02 (2006.01), A61Q 5/08 (2006.01)


(72) Inventeurs/Inventors:  
BELINKY, PAULA, IL;  
KARMON, YORAM, IL;  
KRINFELD, BELLA, IL;  
LASSER, HAIM, IL

(73) Propriétaire/Owner:  
RAKUTO BIO TECHNOLOGIES LTD., IL

(74) Agent: INTEGRAL IP

(54) Titre : COMPOSITIONS COSMETIQUES CONTENANT UN SYSTEME ENZYMATIQUE

(54) Title: ENZYMATIC SYSTEM-CONTAINING COSMETIC COMPOSITIONS



(57) Abrégé/Abstract:

Cosmetic or pharmaceutical compositions are disclosed herein for application to skin, hair and/or a nail of a subject, as well as methods utilizing same for administering hydrogen peroxide to skin, hair and/or a nail, and kits for applying same. The compositions comprise a hydrogen peroxide-producing enzyme, and a substrate of the hydrogen peroxide-producing enzyme. Further disclosed herein are cosmetic compositions and kits for lightening skin and/or hair, which further comprise a lignin peroxidase, as well as cosmetic methods utilizing same.

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



## (10) International Publication Number

WO 2014/191995 A3

(43) International Publication Date  
4 December 2014 (04.12.2014)

(51) International Patent Classification:  
*A61Q 3/00* (2006.01)      *A61K 8/66* (2006.01)  
*A61Q 5/08* (2006.01)      *A61Q 19/02* (2006.01)

(21) International Application Number:  
PCT/IL2014/050477

(22) International Filing Date:  
27 May 2014 (27.05.2014)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
61/827,685      27 May 2013 (27.05.2013)      US

(71) Applicant: RAKUTO BIO TECHNOLOGIES LTD.  
[IL/IL]; Kochav Yokneam, P.O. Box 219, 20692 Yokneam Ilit (IL).

(72) Inventors: BELINKY, Paula; 25 HaNarkis Street, 10292 Metulla (IL). KARMON, Yoram; 6 Martin Buber Street, 49407 Petah Tikva (IL). COHEN, Shaul; Kibbutz Ashdot Yaakov Meuhad, 1515000 Doar-Na Emek HaYarden (IL). KRINFELD, Bella; P.O. Box 2117, 1156915 Kiryat-Shmona (IL). LASER, Haim; 12 Moran Street, 44288 Kfar Saba (IL).

(74) Agents: EHRLICH, Gal et al.; G.E. Ehrlich (1995) Ltd., 11 Menachem Begin Road, 5268104 Ramat Gan (IL).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

## Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

[Continued on next page]

(54) Title: ENZYMIC SYSTEM-CONTAINING COSMETIC COMPOSITIONS




FIG. 10A

WO 2014/191995 A3

(57) Abstract: Cosmetic or pharmaceutical compositions are disclosed herein for application to skin, hair and/or a nail of a subject, as well as methods utilizing same for administering hydrogen peroxide to skin, hair and/or a nail, and kits for applying same. The compositions comprise a hydrogen peroxide-producing enzyme, and a substrate of the hydrogen peroxide-producing enzyme. Further disclosed herein are cosmetic compositions and kits for lightening skin and/or hair, which further comprise a lignin peroxidase, as well as cosmetic methods utilizing same.

**WO 2014/191995 A3** 

---

— *with sequence listing part of description (Rule 5.2(a))*

**(88) Date of publication of the international search report:**  
19 March 2015

## ENZYMATIC SYSTEM-CONTAINING COSMETIC COMPOSITIONS

### FIELD AND BACKGROUND OF THE INVENTION

The present invention, in some embodiments ther eof, relates to novel cosmetic compositions, kits and methods and, more particularly, but not exclusively, to cosmetic compositions, kits and methods, which utilize an enzymatic system for producing hydrogen peroxide *in situ*, and to uses thereof in, for example, lightening skin and/or hair.

The color of human skin is governed by the quantity, quality, and distribution of melanin, a pigment which is also present in plants and microorganisms.

The synthesis of melanin initiates from the precursor L-tyrosine, which is transformed into a second precursor dopaquinone via the action of tyrosinase. Synthesis of melanin takes place in melanosomes, which are present in melanocyte cells present in the epidermal basal layer; synthesis of melanin in these cells is induced by ultraviolet (UV) light. Following synthesis, melanin migrates to epidermal cells and is dispersed therein, where melanin is decolored following dermal metabolism and then scaled off in the form of dirt at the time of skin renewal. Melanin has a clinical importance since it protects the skin from adverse effects caused by UV light. However, high levels of melanin can result in unwanted skin darkening, while the heterogeneous distribution thereof can lead to chloasma and freckling which can be aesthetically displeasing.

Skin lightening products have become increasingly popular in the past few years. The main purpose of skin lightening products is to lighten or whiten the skin or to treat pigmentation disorders such as chloasma, freckles, pregnancy marks and age spots. Several types of skin lightening products are presently available.

Products based on the degeneration and death of pigment cells typically include harsh chemicals, such as hydroquinone, 4-isopropylcatechol, and hydroquinone monobenzyl ether, that promote skin whitening, lightening or fade out skin pigmentation. Other agents which are commonly-available for skin-whitening include kojic acid and derivatives thereof, which inhibits melanin production, and a variety of extracts such as licorice (*Glycyrrhiza glabra*) which include glabridin that decreases oxidization activity of the melanin creating cells in the skin in similar way as Kojic acid does.

Other lightening products are based on the inhibition of tyrosinase, the enzyme that transforms the precursor L-tyrosine into a second precursor dopaquinone. This group of products includes arbutin, a glucose hydroquinone compound, which is capable of inhibiting tyrosinase by chelating copper ions thereby suppressing the 5 tautomerization from dopachrome to 5,6-dihydroxyindole (DHI) and 5,6-dihydroxyindole-2-carboxylic acid (DHICA). Melanostat is another lightening product that acts through tyrosinase. Melanostat is a synthetic peptide that functions in deactivating melanogenesis in melanocytes.

Several antioxidant compounds that can inhibit the production of melanin are 10 also utilized in lightening products. Since the synthesis of melanin involves an oxidation reaction, blocking the oxidation at various points from tyrosine/DOPA to melanin ultimately inhibits the synthesis of melanin. For example, L-ascorbic acid (vitamin C) acts as a reducing agent on melanin intermediates and blocks oxidative reactions; other antioxidants utilized by lightening products include bioflavonoids, 15 which are typically extracted from mulberry or licorice.

However, the currently available skin whitening products are typically inefficient and may be harmful to the skin since a continuous external application of these products can lead to permanent leucoderma and side effects such as dyschromatosis and rash.

Consumers frequently use cosmetic products to care for their skin. Rough and/or 20 broken skin and hyperpigmentations (such as age spots, freckles, and brown patches associated with sunlight exposure, skin aging or environmental damage to the human skin) are areas consumers typically seek to treat. Skin whitening is of particular interest in certain Asian populations.

U.S. Patent Application Publication No. 20050013784 discloses a treatment 25 mask comprising a water insoluble substrate and a liquid composition comprising a skin tone changing agent and a water-soluble thickening agent.

Treatment masks, which include hydrogen peroxide, are currently available from Reviva Labs (e.g., Oxygen Mask, Green Papaya & Hydrogen Peroxide Oxygen Mask, REV-11305).

30 International Patent Application Publication WO 2004/052275 discloses methods of producing lignin peroxidase, and methods and cosmetic compositions suitable for

skin and hair lightening as well as kits and an article-of-manufacturing including active ingredients for skin and hair lightening.

International Patent Application Publication WO 2012/153336 discloses kits and methods for lightening a skin of a subject, using a mask comprising an oxidizing activator such as hydrogen peroxide, and a cosmetic composition comprising a lignin peroxidase; or a mask comprising a lignin peroxidase and a cosmetic composition comprising the oxidizing activator.

U.S. Patent No. 8,426,158 describes methods for increasing hydrolysis of a cellulosic material by hydrolyzing the cellulosic material with an enzyme composition in the presence of a polypeptide having peroxidase activity, the peroxidase activity being for reducing the inhibition of hydrolysis by peroxides generated during fermentation.

Hydrogen peroxide is formed in saliva by oxidation of glucose by glucose oxidase. The hydrogen peroxide serves as a substrate for lactoperoxidase in the saliva, which forms bactericidal oxidized intermediates such as hypothiocyanate, hypobromite and hypoiodite from thiocyanate, bromide and iodide, respectively.

International Patent Application Publication WO 88/002600 describes a bactericidal composition comprising a hydrogen peroxide-forming enzyme such as glucose oxidase, a peroxidase such as lactoperoxidase, a thiocyanate and lysozyme. Hydrogen peroxide generated during use is intended, in combination with the peroxidase, to convert the thiocyanate into hypothiocyanate, thereby attenuating bacteria such that they are lysed by the lysozyme.

Midda & Cooksey [*J Clin Periodontol* 1986, 13:950-956] describe mouthrinses and dentrifices which contain amyloglucosidase and glucose oxidase for producing hydrogen peroxide from dietary fermentable carbohydrates, which in turn converts thiocyanate to hypothiocyanate in the presence of salivary lactoperoxidase, thereby inhibiting bacteria.

Enzycal™ toothpaste (Curaprox) contains lactoperoxidase, glucose oxidase and amyloglucosidase, for strengthening the antibacterial effect of saliva [*Br Dent J* 2008, 205:681].

Biotene™ toothpaste and mouthwash (GlaxoSmithKline) contain glucose oxidase, lactoferrin, lactoperoxidase, lysozyme.

U.S. Patent No. 5,607,681 describes anti-microbial compositions containing iodide, thiocyanate, glucose oxidase, D-glucose and optionally also a peroxidase such as lactoperoxidase. The compositions are for use as preservatives or as active agents in oral hygiene, deodorant and anti-dandruff products. The compositions may be provided 5 as non-reacting dry powders and non-aqueous solutions which may be diluted in order to obtain anti-microbial activity.

U.S. Patent No. 6,214,339 describes treatment of otitis externa in dogs and cats by administering to the outer ear of a non-aqueous composition containing an oxidizable substrate such as glucose and an oxidoreductase enzyme such as glucose oxidase, for 10 producing hydrogen peroxide upon encountering the environment of the outer ear. The composition further contains an iodide salt and a peroxidase such as lactoperoxidase for producing hypioiodite from the hydrogen peroxide.

U.S. Patent No. 7,833,290 describes a method of dyeing keratinous fibers by treatment with a compound such as hematoxylin, hematein, brazilin and brazilein; a 15 metal salt; a bicarbonate; and hydrogen peroxide or a system which generates hydrogen peroxide, such as urea peroxide, metal peroxides, perborates, percarborates and oxidases in the presence of a suitable substrate.

Additional background art includes Hugoson et al. [*Odontol Rev* 1974, 25:69-80]; U.S. Patent No. 7,981,166; and U.S. Patent Application Publication Nos. 20 20040161435 and 20090130040.

### SUMMARY OF THE INVENTION

The present inventors have now devised and successfully practiced a methodology which circumvents the need to include hydrogen peroxide in cosmetic 25 compositions and kits, and in lignin peroxidase-containing compositions and kits in particular.

According to an aspect of some embodiments of the invention, there is provided a cosmetic composition for lightening a skin and/or hair of a subject, the composition comprising a lignin peroxidase, a hydrogen peroxide-producing enzyme, and a substrate 30 of the hydrogen peroxide-producing enzyme.

According to an aspect of some embodiments of the invention, there is provided a kit for lightening a skin and/or hair of a subject, the kit comprising a hydrogen

peroxide-producing enzyme, a substrate of the hydrogen peroxide-producing enzyme and a lignin peroxidase, wherein at least one of the hydrogen peroxide-producing enzyme, the substrate of the hydrogen peroxide-producing enzyme and the lignin peroxidase forms a part of a composition which is packaged individually within the kit.

5 According to an aspect of some embodiments of the invention, there is provided a cosmetic method of lightening a skin and/or hair of a subject, the method comprising contacting a skin and/or hair region of the subject with a composition comprising a lignin peroxidase, a hydrogen peroxide-producing enzyme and a substrate of the hydrogen peroxide-producing enzyme, thereby lightening the skin and/or hair of the 10 subject.

According to an aspect of some embodiments of the invention, there is provided a cosmetic or pharmaceutical composition for application to skin, hair and/or a nail of a subject, the composition comprising a hydrogen peroxide-producing enzyme, and a substrate of the hydrogen peroxide-producing enzyme.

15 According to an aspect of some embodiments of the invention, there is provided a kit for applying a cosmetic or pharmaceutical composition to skin, hair and/or a nail of a subject, the cosmetic or pharmaceutical composition comprising hydrogen peroxide, the kit comprising a first composition comprising a hydrogen peroxide-producing enzyme, and a second composition comprising a substrate of the hydrogen peroxide-producing enzyme. 20

According to an aspect of some embodiments of the invention, there is provided a method of administering hydrogen peroxide to skin, hair and/or a nail of a subject, the method comprising contacting a skin, hair and/or a nail region of the subject with a composition comprising a hydrogen peroxide-producing enzyme and a substrate of the 25 hydrogen peroxide-producing enzyme, thereby generating hydrogen peroxide on the skin, hair and/or nail.

According to some of any of the embodiments of the invention, and any combination thereof, a concentration of lignin peroxidase in the composition is in a range of from 1.25 to 125 units/gram.

30 According to some of any of the embodiments of the invention, and any combination thereof, a concentration of the hydrogen peroxide-producing enzyme in the composition is in a range of from 0.125 to 1250 units/gram.

According to some of any of the embodiments of the invention, and any combination thereof, a concentration of the substrate of the hydrogen peroxide-producing enzyme in the composition is in a range of from 0.0175  $\mu$ mole/gram to 175  $\mu$ mole/gram.

5 According to some of any of the embodiments of the invention, and any combination thereof, the composition further comprises an oxidizing mediator.

According to some of any of the embodiments of the invention, and any combination thereof, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene.

10 According to some of any of the embodiments of the invention, and any combination thereof, a concentration of the oxidizing mediator in the composition is in a range of from 0.03  $\mu$ mole/gram to 300  $\mu$ mole/gram.

According to some of any of the embodiments of the invention, and any combination thereof, the hydrogen peroxide-producing enzyme is selected from the 15 group consisting of a glucose oxidase, a hexose oxidase, a cholesterol oxidase, an aryl alcohol oxidase, an L-gulonolactone oxidase, a galactose oxidase, a pyranose-2-oxidase, a pyridoxine-4-oxidase, an alcohol oxidase, a 2-hydroxy-acid oxidase, a choline oxidase, a long-chain-alcohol oxidase, a glycerol-3-phosphate oxidase, a D-arabinono-1,4-lactone oxidase, a vanillyl alcohol oxidase, an alditol oxidase, a prosolanapyrone II oxidase, a 20 paromamine 6'-oxidase, a glyoxal oxidase, a veratryl alcohol oxidase, an alcohol dehydrogenase, a cellobiose dehydrogenase, an aldehyde oxidase, an oxalate oxidase, an aryl aldehyde oxidase, a dihydroorotate oxidase, a pyrroloquinoline-quinone synthase, an L-amino acid oxidase, an L-glutamate oxidase, a polyamine oxidase, an NAD(P)H oxidase, a urate oxidase, a hydroxylamine oxidase, a thiol oxidase, a glutathione 25 oxidase, a cytochrome c oxidase, a xanthine oxidase and a superoxide dismutase.

According to some of any of the embodiments of the invention, and any combination thereof, the hydrogen peroxide-producing enzyme comprises glucose oxidase and the substrate comprises D-glucose.

According to some of any of the embodiments of the invention, and any 30 combination thereof, the lignin peroxidase is isoenzyme H1 or a modified form of isoenzyme H2.

According to some of any of the embodiments of the invention, and any combination thereof, the lignin peroxidase is a lignin peroxidase of a white rot fungus.

According to some of any of the embodiments of the invention, and any combination thereof, the lignin peroxidase is an extract of a *Phanerochaete chrysosporium* fungus.

According to some of any of the embodiments of the invention, and any combination thereof, the lignin peroxidase is recombinantly expressed.

According to some of any of the embodiments of the invention, and any combination thereof, the lignin peroxidase comprises a polypeptide sequence set forth in  
10 SEQ ID NO:1.

According to some of any of the embodiments of the invention, and any combination thereof, the cosmetic composition is in a form selected from the group consisting of a mask, a serum, a lotion, a cream, a shampoo, a gel and a toner.

According to some of any of the embodiments of the invention, and any combination thereof, the composition further comprises a cosmetically acceptable carrier  
15 and/or pharmaceutically acceptable carrier.

According to some of any of the embodiments of the invention, and any combination thereof, the composition further comprises a cosmetically acceptable carrier.

20 According to some of any of the embodiments of the invention, and any combination thereof, the composition is packaged in an air-tight container.

According to some of any of the embodiments of the invention, and any combination thereof, the air-tight container is configured for dispensing the composition without entry of air into the container.

25 According to some of any of the embodiments of the invention, and any combination thereof, the hydrogen peroxide-producing enzyme forms a part of a first composition of the kit and the substrate forms a part of a second composition of the kit, the first and second compositions being packaged individually within the kit.

According to some of any of the embodiments of the invention, and any combination thereof, the first composition further comprises lignin peroxidase.

30 According to some of any of the embodiments of the invention, and any combination thereof, the second composition further comprises lignin peroxidase.

According to some of any of the embodiments of the invention, and any combination thereof, the lignin peroxidase forms a part of a third composition within the kit, the third composition being packaged individually within the kit.

According to some of any of the embodiments of the invention, and any combination thereof, the hydrogen peroxide-producing enzyme and the substrate of the hydrogen peroxide-producing enzyme are packaged together in an air-tight package within the kit.

According to some of any of the embodiments of the invention, and any combination thereof, the air-tight package is a container configured for dispensing the composition without entry of air into the container.

According to some of any of the embodiments of the invention, and any combination thereof, a concentration of lignin peroxidase in a composition comprising the lignin peroxidase is in a range of from 2.5 to 250 units/gram.

According to some of any of the embodiments of the invention, and any combination thereof, the kit comprises an oxidizing mediator, the hydrogen peroxide-producing enzyme, the substrate of said hydrogen peroxide-producing enzyme and the lignin peroxidase.

According to some of any of the embodiments of the invention, and any combination thereof, the oxidizing mediator forms a part of a composition that comprises the lignin peroxidase.

According to some of any of the embodiments of the invention, and any combination thereof, a concentration of the oxidizing mediator in the composition is in a range of from 0.06  $\mu$  mole/gram to 600  $\mu$  mole/gram.

According to some of any of the embodiments of the invention, and any combination thereof, a concentration of the hydrogen peroxide-producing enzyme in a composition comprising the enzyme is in a range of from 0.25 to 2500 units/gram.

According to some of any of the embodiments of the invention, and any combination thereof, a concentration of the substrate in a composition comprising the substrate is in a range of from 0.035  $\mu$  mole/gram to 350  $\mu$  mole/gram.

According to some of any of the embodiments of the invention, and any combination thereof, the kit comprises:

a first composition comprising lignin peroxidase at a concentration in a range of from 2.5 to 250 units/gram, and glucose oxidase at a concentration in a range of from 2.5 to 250 units/gram; and

5 a second composition comprising D-glucose at a concentration in a range of from 0.35  $\mu$ mole/gram to 35  $\mu$ mole/gram.

According to some of any of the embodiments of the invention, and any combination thereof, the composition which is packaged individually within the kit is in a form selected from the group consisting of a mask, a serum, a lotion, a cream, a shampoo, a gel and a toner.

10 According to some of any of the embodiments of the invention, and any combination thereof, the composition which is packaged individually within the kit further comprises a cosmetically acceptable carrier.

15 According to some of any of the embodiments of the invention, and any combination thereof, the composition is formed by a lignin peroxidase, a hydrogen peroxide-producing enzyme and a substrate of the hydrogen peroxide-producing enzyme as packaged within a kit described hereinabove.

According to some of any of the embodiments of the invention, and any combination thereof, the contacting is effected at least five times.

20 According to some of any of the embodiments of the invention, and any combination thereof, the contacting is effected once or twice per day.

25 According to some of any of the embodiments of the invention, and any combination thereof, the composition is formed by contacting the skin and/or hair region with the hydrogen peroxide-producing enzyme and/or the substrate of the hydrogen peroxide-producing enzyme subsequent to contacting the skin and/or hair region with the lignin peroxidase.

According to some of any of the embodiments of the invention, and any combination thereof, the skin comprises facial skin.

According to some of any of the embodiments of the invention, and any combination thereof, the skin comprises a skin region exposed to the sun.

30 According to some of any of the embodiments of the invention, and any combination thereof, lightening the skin comprises lightening whole skin complexion.

10

According to some of any of the embodiments of the invention, and any combination thereof, the skin comprises an uneven skin tone, dark spot(s), freckle(s), melasma, hyperpigmentation, skin discoloration, age spot(s), acne mark(s) and/or a scar.

According to some of any of the embodiments of the invention, and any combination thereof, the composition further comprises at least one agent which exhibits a cosmetic or pharmaceutical activity in a presence of hydrogen peroxide.

According to some of any of the embodiments of the invention, and any combination thereof, the at least one agent is selected from the group consisting of a peroxidase and an oxidative dye precursor.

According to some of any of the embodiments of the invention, and any combination thereof, the composition is selected from the group consisting of a bleaching composition, a hair-dye composition, an anti-microbial composition and an odor-removal composition.

According to some of any of the embodiments of the invention, and any combination thereof, the kit further comprises at least one agent which exhibits a cosmetic or pharmaceutical activity in a presence of hydrogen peroxide.

According to some of any of the embodiments of the invention, and any combination thereof, the first composition comprises glucose oxidase at a concentration in a range of from 2.5 to 250 units/gram, and the second composition comprises D-glucose at a concentration in a range of from 0.35  $\mu$ mole/gram to 35  $\mu$ mole/gram.

According to some of any of the embodiments of the invention, and any combination thereof, the first composition and second composition are each independently in a form selected from the group consisting of a mask, a serum, a lotion, a cream, a shampoo, a gel and a toner.

According to some of any of the embodiments of the invention, and any combination thereof, the first composition and second composition further comprise a cosmetically acceptable carrier and/or pharmaceutically acceptable carrier.

According to some of any of the embodiments of the invention, and any combination thereof, the composition is formed by a hydrogen peroxide-producing enzyme and a substrate of the hydrogen peroxide-producing enzyme as packaged within a kit described hereinabove.

According to some of any of the embodiments of the invention, and any combination thereof, the composition of a method as described hereinabove is any of the compositions described herein, including any one of the embodiments thereof, in any combination.

Unless otherwise defined, all technical and/or scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of embodiments of the invention, exemplary methods and/or materials are described below. In case of conflict, the patent specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and are not intended to be necessarily limiting.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Some embodiments of the invention are herein described, by way of example only, with reference to the accompanying drawings. With specific reference now to the drawings in detail, it is stressed that the particulars shown are by way of example and for purposes of illustrative discussion of embodiments of the invention. In this regard, the description taken with the drawings makes apparent to those skilled in the art how embodiments of the invention may be practiced.

In the drawings:

FIG. 1 presents images showing an exemplary enzyme cream comprising 25 units/gram lignin peroxidase (LIP) isoenzyme H1 fraction, 25 units/gram glucose oxidase (GOX) and 6 mmol/kg veratryl alcohol (VA) with 140  $\mu$ g/ml melanin (LIP/VA/GOX) and an exemplary activator cream comprising 3.5 mmol/kg glucose (GLU), a control enzyme cream comprising 25 units/gram LIP isoenzyme H1 fraction and 6 mmol/kg VA with 140  $\mu$ g/ml melanin (LIP/VA) and control activator cream comprising 3.53 mmol/kg  $H_2O_2$ , as well as a mixture of the exemplary enzyme and activator cream (LIP/VA/GOX/GLU) and a mixture of the control enzyme and activator cream (LIP/VA/ $H_2O_2$ );

FIGs. 2A and 2B present bar graphs showing values for a melanin index (FIG. 2A) and erythema index (FIG. 2B) in various facial regions of a subject on days 0, 7 and 21 of a treatment comprising administration twice per day of LIP isoenzyme H1,

glucose oxidase and glucose, according to an exemplary embodiment of the invention (\* p<0.05 vs. day 0);

FIGs. 3A and 3B present bar graphs showing values for a melanin index (FIG. 3A) and erythema index (FIG. 3B) in various facial regions of a subject on days 0 and 5 21 of a treatment comprising administration twice per day of LIP isoenzyme H1, glucose oxidase and glucose, according to an exemplary embodiment of the invention (\* p<0.05 vs. day 0);

FIGs. 4A and 4B present bar graphs showing values for a melanin index (FIG. 4A) and erythema index (FIG. 4B) in various facial regions of a subject on days 0, 7 10 and 21 of a treatment comprising administration twice per day of LIP isoenzyme H1, glucose oxidase and glucose, according to an exemplary embodiment of the invention (\* p<0.05 vs. day 0);

FIGs. 5A and 5B present bar graphs showing values for a melanin index (FIG. 5A) and erythema index (FIG. 5B) in various facial regions of a subject on days 0, 7 15 and 21 of a treatment comprising administration twice per day of LIP isoenzyme H1, glucose oxidase and glucose, according to an exemplary embodiment of the invention (\* p<0.05 vs. day 0, # p<0.05 vs. day 7);

FIGs. 6A and 6B present bar graphs showing values for a melanin index (FIG. 6A) and erythema index (FIG. 6B) in various facial regions of a subject on days 0, 7 20 and 21 of a treatment comprising administration twice per day of LIP isoenzyme H1, glucose oxidase and glucose, according to an exemplary embodiment of the invention (\* p<0.05 vs. day 0, # p<0.05 vs. day 7);

FIGs. 7A and 7B present bar graphs showing values for a melanin index (FIG. 7A) and erythema index (FIG. 7B) in various facial regions of a subject on days 0, 7 25 and 21 of a treatment comprising administration twice per day of LIP isoenzyme H1, glucose oxidase and glucose, according to an exemplary embodiment of the invention (\* p<0.05 vs. day 0, # p<0.05 vs. day 7);

FIGs. 8A and 8B present bar graphs showing values for a melanin index (FIG. 8A) and erythema index (FIG. 8B) in various facial regions of a subject on days 0 and 7 30 of a treatment comprising administration twice per day of LIP isoenzyme H1, glucose oxidase and glucose, according to an exemplary embodiment of the invention (\* p<0.05 vs. day 0);

FIGs. 9A and 9B present bar graphs showing values for a melanin index (FIG. 9A) and erythema index (FIG. 9B) in various facial regions of a subject on days 0, 7 and 21 of a treatment comprising administration twice per day of LIP isoenzyme H1, glucose oxidase and glucose, according to an exemplary embodiment of the invention  
 5 (\* p<0.05 vs. day 0, # p<0.05 vs. day 7; increase for erythema index in cheeks on day 21 due to irritation unrelated to treatment);

FIGs. 10A and 10B present graphs showing the melanin index of pigmented (FIG. 10A) and non-pigmented (FIG. 10B) skin after 0, 4, 6 and 8 weeks of a treatment comprising administration twice per day of LIP isoenzyme H1, glucose oxidase and glucose, according to an exemplary embodiment of the invention (\*\* p<0.01 (p=0.001 in FIG. 10A and p=0.007 in FIG. 10B), \*\*\* p<0.001);  
 10

FIG. 11 presents a visual analogue scale showing uniformity of pigmentation of skin, as judged by a dermatologist, after 0, 4, 6 and 8 weeks of a treatment comprising administration twice per day of LIP isoenzyme H1, glucose oxidase and glucose, according to an exemplary embodiment of the invention (0 = uniform, even, 10 = uneven, blotchy, mottled, \* p<0.05);  
 15

FIG. 12 presents a visual analogue scale showing clarity of skin, as judged by a dermatologist, after 0, 4, 6 and 8 weeks of a treatment comprising administration twice per day of LIP isoenzyme H1, glucose oxidase and glucose, according to an exemplary embodiment of the invention (0 = clear, even, 10 = dull, opaque, \* p<0.05, \*\* p<0.01);  
 20

FIG. 13 presents a visual analogue scale showing lightening/brightening of skin, as judged by a dermatologist, after 0, 4, 6 and 8 weeks of a treatment comprising administration twice per day of LIP isoenzyme H1, glucose oxidase and glucose, according to an exemplary embodiment of the invention (0 = light/bright appearance, 10 = dull, matte, \*\* p<0.01, \*\*\* p<0.001);  
 25

FIG. 14 presents a visual analogue scale showing skin tone, as judged by a dermatologist, after 0, 4, 6 and 8 weeks of a treatment comprising administration twice per day of LIP isoenzyme H1, glucose oxidase and glucose, according to an exemplary embodiment of the invention (0 = even, healthy skin color, 10 = uneven, discolored appearance, \* p<0.05, \*\*\* p<0.001);  
 30

FIG. 15 presents a visual analogue scale showing radiance of skin, as judged by a dermatologist, after 0, 4, 6 and 8 weeks of a treatment comprising administration

twice per day of LIP isoenzyme H1, glucose oxidase and glucose, according to an exemplary embodiment of the invention (0 = radiant, luminous appearance, 10 = dull, matte, sallow appearance, \*\*\* p<0.001);

5 FIG. 16 presents a visual analogue scale showing overall appearance of skin, as judged by a dermatologist, after 0, 4, 6 and 8 weeks of a treatment comprising administration twice per day of LIP isoenzyme H1, glucose oxidase and glucose, according to an exemplary embodiment of the invention (0 = healthy looking, 10 = unhealthy looking, \*\*\* p<0.001); and

10 FIG. 17 presents photographs of subjects before (baseline) and after 8 weeks of a treatment comprising administration twice per day of LIP isoenzyme H1, glucose oxidase and glucose, according to an exemplary embodiment of the invention.

#### DESCRIPTION OF SPECIFIC EMBODIMENTS OF THE INVENTION

15 The present invention, in some embodiments thereof, relates to novel cosmetic compositions, kits and methods and, more particularly, but not exclusively, to hydrogen peroxide-producing cosmetic compositions, kits and methods, and to uses thereof in, for example, lightening skin and/or hair.

20 Before explaining at least one embodiment of the invention in detail, it is to be understood that the invention is not necessarily limited in its application to the details set forth in the following description or exemplified by the Examples. The invention is capable of other embodiments or of being practiced or carried out in various ways.

25 Methods for lightening a skin of a subject, using hydrogen peroxide and a lignin peroxidase, have been previously disclosed (see, for example, International Patent Application Publication WO 2012/153336, discussed hereinabove). Such methods require utilization of compositions containing hydrogen peroxide.

Many other cosmetic compositions and methods include or utilize hydrogen peroxide.

However, utilizing hydrogen peroxide in cosmetic applications has recently been recognized as undesired.

30 In a search for methodologies in which the need to utilize compositions containing hydrogen peroxide is circumvented, the present inventors have envisioned and successfully devised and practiced a novel methodology which utilizes an enzymatic

system for generating hydrogen peroxide upon application. More specifically, the present inventors have utilized a hydrogen peroxide-producing enzyme and a substrate thereof, for generating hydrogen peroxide *in situ*, immediately prior to application of a composition or system containing same and/or during application of a composition or system containing same. The enzymatic system is readily controllable by keeping the enzyme and substrate separate, until hydrogen peroxide generation is desired and/or by keeping the enzyme and the substrate inactive towards one another, by storage in the absence of oxygen. The herein disclosed methodology is advantageous by being capable of generating the hydrogen peroxide gradually *in situ*, where it is consumed, such that a concentration of hydrogen peroxide remains low and stable over time.

While reducing the present invention to practice, the inventors have demonstrated the efficacy of an enzymatic system for generating hydrogen peroxide for activating lignin peroxidase in order to lighten skin.

Thus, as shown in the Examples section which follows, the present inventors have designed a cosmetic composition which comprises lignin peroxidase, and an activator composition which comprises glucose, which when applied together (e.g., on a skin region of a subject) results in a significant whitening effect as a result of reduction in melanin content (Figures 1-9B), due to the interaction of lignin peroxidase with hydrogen peroxide generated by the glucose oxidase and glucose.

These results demonstrate the efficacy of using an enzyme/substrate system for generating hydrogen peroxide *in situ* (i.e., in or on a region of the body where it is intended to react (e.g., with melanin in a peroxidase-catalyzed reaction).

According to an aspect of some embodiments of the invention, there is provided a cosmetic or pharmaceutical composition for topical application (e.g., for application to skin, hair and/or nail of a subject). According to some embodiments of the present invention, the composition comprises a hydrogen peroxide-producing enzymatic system, namely, a hydrogen peroxide-producing enzyme and a substrate of the hydrogen peroxide-producing enzyme.

As used herein throughout, the term “cosmetic” with regard to any of the compositions, kits and methods (and any other aspect) disclosed herein, refers to compositions, kits and method which enhance the appearance or odor of the human body.

In some embodiments, the term “cosmetic” follows the definition regulatory authorities worldwide. For example, in some embodiments, the term “cosmetic” follows the definition of the U.S Food and Drug Administration (FDA), as being “intended to be applied to the human body for cleansing, beautifying, promoting 5 attractiveness, or altering the appearance without affecting the body's structure or functions”.

***The hydrogen peroxide-producing enzymatic system:***

As used herein throughout, the phrase “hydrogen peroxide-producing enzyme” refers to an enzyme which catalyzes a reaction in which hydrogen peroxide is a product.

As used herein throughout, the “substrate” of a hydrogen peroxide-producing enzyme refers to any compound other than O<sub>2</sub> or water which serves as a substrate for the abovementioned reaction catalyzed by the hydrogen peroxide-producing enzyme in which hydrogen peroxide is a product. The “substrate” may be a single compound or a plurality of compounds. The plurality of compounds may interact with the enzyme as 10 part of different reactions, or the enzyme catalyzes a single reaction which utilizes a plurality of compounds as substrates.

In some embodiments, the enzyme catalyzes a reaction in which the substrate is oxidized and O<sub>2</sub> is reduced to hydrogen peroxide.

In some embodiments, the enzyme catalyzes a reaction in which the substrate is 20 reduced and water is oxidized to hydrogen peroxide.

Examples of hydrogen peroxide-producing enzymes which may be used in embodiments of the invention include, without limitation:

an EC 1.1.3 oxidoreductase such as, for example, a glucose oxidase (EC 1.1.3.4), a hexose oxidase (EC 1.1.3.5), a cholesterol oxidase (EC 1.1.3.6), an aryl alcohol 25 oxidase (EC 1.1.3.7), an L-gulonolactone oxidase (EC 1.1.3.8), a galactose oxidase (EC 1.1.3.9), a pyranose-2-oxidase (EC 1.1.3.10), a pyridoxine-4-oxidase (EC 1.1.3.12), an alcohol oxidase (EC 1.1.3.13), a 2-hydroxy-acid oxidase (EC 1.1.3.15), a choline oxidase (EC 1.1.3.17), a long-chain-alcohol oxidase (EC 1.1.3.20), a glycerol-3-phosphate oxidase (EC 1.1.3.21), a D-arabinono-1,4-lactone oxidase (EC 1.1.3.37), a 30 vanillyl alcohol oxidase (EC 1.1.3.38), an alditol oxidase (EC 1.1.3.41), a prosolanapyrone II oxidase (EC 1.1.3.42), a paromamine 6'-oxidase (EC 1.1.3.43), a glyoxal oxidase and a veratryl alcohol oxidase;

- an alcohol dehydrogenase (EC 1.1.99.8);
- a cellobiose dehydrogenase (EC 1.1.99.18);
- an EC 1.2.3 oxidoreductase such as, for example, an aldehyde oxidase (EC 1.2.3.1), an oxalate oxidase (EC 1.2.3.4), and an aryl aldehyde oxidase (EC 1.2.3.9);
- 5 an EC 1.3.3 oxidoreductase such as, for example, a dihydroorotate oxidase (EC 1.3.3.1), and a pyrroloquinoline-quinone synthase (EC 1.3.3.11);
- an EC 1.4.3 oxidoreductase such as, for example, an L-amino acid oxidase (EC 1.4.3.2), and an L-glutamate oxidase (1.4.3.11);
- 10 an EC 1.5.3 oxidoreductase such as, for example, a polyamine oxidase (1.5.3.11);
- an EC 1.6.3 oxidoreductase such as, for example, an NAD(P)H oxidase (1.6.3.1);
- an EC 1.7.3 oxidoreductase such as, for example, a urate oxidase (EC 1.7.3.3), and a hydroxylamine oxidase (EC 1.7.3.4);
- 15 an EC 1.8.3 oxidoreductase such as, for example, a thiol oxidase (EC 1.8.3.2), and a glutathione oxidase (EC 1.8.3.3);
- an EC 1.9.3 oxidoreductase, such as, for example, a cytochrome c oxidase (EC 1.9.3.1);
- an EC 1.17.3 oxidoreductase such as, for example, a xanthine oxidase (EC 1.17.3.2); and
- 20 a superoxide dismutase (EC 1.15.1.1).

The EC nomenclature herein is according to the International Union of Biochemistry and Molecular Biology.

In some embodiments, the hydrogen peroxide-producing enzyme is one or more of an EC 1.1.3 oxidoreductase, an EC 1.2.3 oxidoreductase, an EC 1.3.3 oxidoreductase, 25 an EC 1.4.3 oxidoreductase, an EC 1.5.3 oxidoreductase, an EC 1.6.3 oxidoreductase, an EC 1.7.3 oxidoreductase, an EC 1.8.3 oxidoreductase, an EC 1.9.3 oxidoreductase and an EC 1.17.3 oxidoreductase (e.g., such as an oxidoreductase described herein).

In some embodiments a superoxide dismutase is included for producing hydrogen peroxide from superoxide, in combination with a source of superoxide. In 30 some embodiments, the source of superoxide is an enzyme which produces superoxide, for example, an NAD(P)H oxidase and/or a xanthine oxidase.

In some embodiments, the hydrogen peroxide-producing enzyme produces superoxide. As superoxide normally reacts to a considerable degree to form hydrogen peroxide (e.g., by dismutation), production of superoxide is considered herein to encompass production of hydrogen peroxide.

5 Suitable substrates of the hydrogen peroxide-producing enzyme will be apparent to the skilled person.

In exemplary embodiments, the enzyme is a glucose oxidase (EC 1.1.3.4) and the substrate comprises D-glucose.

10 In some embodiments, the enzyme is a hexose oxidase (EC 1.1.3.5) and the substrate comprises D-glucose, D-galactose, D-mannose, maltose, lactose and/or cellobiose.

In some embodiments, the enzyme is a cholesterol oxidase (EC 1.1.3.6) and the substrate comprises cholesterol.

15 In some embodiments, the enzyme is an aryl alcohol oxidase (EC 1.1.3.7) and the substrate comprises an aromatic primary alcohol.

In some embodiments, the enzyme is an L-gulonolactone oxidase (EC 1.1.3.8) and the substrate comprises L-gulono-1,4-lactone.

20 In some embodiments, the enzyme is a galactose oxidase (EC 1.1.3.9) and the substrate comprises a primary alcohol (e.g., methanol, ethanol, n-propanol, n-butanol, isobutanol), a monosaccharide, an oligosaccharide and/or a polysaccharide.

25 In some embodiments, the enzyme is a pyranose-2-oxidase (EC 1.1.3.10) and the substrate comprises an aldopyranose (e.g., D-glucose, D-xylose, D-galactose), L-sorbose and/or 1,5-anhydroglucitol, and/or disaccharides thereof. It is to be appreciated that pyranose-2-oxidase is a natural source for hydrogen peroxide for activating lignin cellulose.

In some embodiments, the enzyme is a pyridoxine-4-oxidase (EC 1.1.3.12) and the substrate comprises pyridoxine.

30 In some embodiments, the enzyme is an alcohol oxidase (EC 1.1.3.13) and the substrate comprises a primary alcohol (e.g., methanol, ethanol, n-propanol, n-butanol, isobutanol).

In some embodiments, the enzyme is a 2-hydroxy-acid oxidase (EC 1.1.3.15) and the substrate comprises an (S)-2-hydroxy acid (e.g., glycolate, 2-hydroxyglutarate) and/or an (S)-2-amino acid.

5 In some embodiments, the enzyme is a choline oxidase (EC 1.1.3.17) and the substrate comprises choline and/or glycine betaine aldehyde.

In some embodiments, the enzyme is a long-chain-alcohol oxidase (EC 1.1.3.20) and the substrate is a long-chain alcohol (e.g., a fatty alcohol).

In some embodiments, the enzyme is a glycerol-3-phosphate oxidase (EC 1.1.3.21) and the substrate comprises glycerol-3-phosphate.

10 In some embodiments, the enzyme is a D-arabinono-1,4-lactone oxidase (EC 1.1.3.37) and the substrate comprises L-gulono-1,4-lactone, D-arabinono-1,4-lactone, and/or L-galactono-1,4-lactone.

15 In some embodiments, the enzyme is a vanillyl alcohol oxidase (EC 1.1.3.38) and the substrate comprises a 4-allylphenol, a 4-hydroxybenzyl alcohol (e.g., vanillyl alcohol), a 4-hydroxybenzamine and/or a 4-(methoxymethyl)phenol.

In some embodiments, the enzyme is an alditol oxidase (EC 1.1.3.41) and the substrate comprises an alditol (e.g., xylitol, D-sorbitol).

In some embodiments, the enzyme is a prosolanapyrone II oxidase (EC 1.1.3.42) and the substrate comprises prosolanapyrone II.

20 In some embodiments, the enzyme is a paromamine 6'-oxidase (EC 1.1.3.43) and the substrate comprises paromamine and/or 6'''-deamino-6'''hydroxyneomycin C.

In some embodiments, the enzyme is a veratryl alcohol oxidase and the substrate is a primary (4-methoxy)-aromatic alcohol (e.g., veratryl alcohol).

25 In some embodiments, the enzyme is a glyoxal oxidase and the substrate is glyoxal, an alkyl glyoxal (e.g., methyl glyoxal) and/or cellobiose.

In some embodiments, the enzyme is an alcohol dehydrogenase (EC 1.1.99.8) and the substrate comprises a primary alcohol (e.g., methanol, ethanol, n-propanol, n-butanol, isobutanol).

30 In some embodiments, the enzyme is a cellobiose dehydrogenase (EC 1.1.99.18) and the substrate comprises cellobiose, a cello-oligosaccharide, lactose, and/or D-glucosyl-1,4- $\beta$ -D-mannose.

In some embodiments, the enzyme is an aldehyde oxidase (EC 1.2.3.1) and the substrate comprises an aldehyde.

In some embodiments, the enzyme is an oxalate oxidase (EC 1.2.3.4) and the substrate comprises oxalate.

5 In some embodiments, the enzyme is an aryl aldehyde oxidase (EC 1.2.3.9) and the substrate comprises an aromatic aldehyde (e.g., benzaldehyde, vanillin).

In some embodiments, the enzyme is a dihydroorotate oxidase (EC 1.3.3.1) and the substrate comprises dihydroorotate.

10 In some embodiments, the enzyme is a pyrroloquinoline-quinone synthase (EC 1.3.3.11) and the substrate comprises 6-(2-amino-2-carboxyethyl)-7,8-dioxo-1,2,3,4,7,8-hexahydroquinoline-2,4-dicarboxylate.

In some embodiments, the enzyme is an L-amino acid oxidase (EC 1.4.3.2) and the substrate comprises an L-amino acid.

15 In some embodiments, the enzyme is an L-glutamate oxidase (1.4.3.11) and the substrate comprises L-glutamate.

In some embodiments, the enzyme is a polyamine oxidase (1.5.3.11) and the substrate comprises N<sub>1</sub>-acetylspermine.

In some embodiments, the enzyme is an NAD(P)H oxidase (1.6.3.1) and the substrate comprises NADH and/or NADPH.

20 In some embodiments, the enzyme is a urate oxidase (EC 1.7.3.3) and the substrate comprises urate.

In some embodiments, the enzyme is a hydroxylamine oxidase (EC 1.7.3.4) and the substrate comprises a hydroxylamine.

25 In some embodiments, the enzyme is a thiol oxidase (EC 1.8.3.2) and the substrate comprises a thiol group.

In some embodiments, the enzyme is a glutathione oxidase (EC 1.8.3.3) and the substrate comprises glutathione and/or cysteine.

In some embodiments, the enzyme is a cytochrome c oxidase (EC 1.9.3.1) and the substrate comprises ferrocytocrome c.

30 In some embodiments, the enzyme is a xanthine oxidase (EC 1.17.3.2) and the substrate comprises xanthine and/or hypoxanthine.

The hydrogen peroxide-producing enzyme described herein (e.g., glucose oxidase) may be obtained, for example, from a commercial supplier, by extraction from a natural source, and/or by recombinant expression. General procedures for expressing a protein are described in detail hereinafter, and may be used for expressing a hydrogen peroxide-producing enzyme described herein.

***The hydrogen peroxide-producing compositions:***

A hydrogen peroxide-producing composition as described herein produces hydrogen peroxide whenever the hydrogen peroxide-producing enzyme and its substrate contact one another in the presence of oxygen (or air) and/or water.

It is to be understood that a hydrogen peroxide-producing composition according to any one of the embodiments relating to such compositions (e.g., as described in this section) may utilize a hydrogen peroxide-producing system according to any one of the embodiments described herein relating to such systems.

In some embodiments of any of the embodiments described herein, any of the composition described herein is prepared *in situ*, that is, on the site of application (e.g., skin, hair and/or nail) of the subject, by contacting different components of the composition with each other. In some embodiments, hydrogen peroxide generation begins only upon contacting the hydrogen peroxide-producing enzyme with the substrate thereof *in situ*.

In some embodiments of any of the embodiments described herein, the composition is prepared on the site of application (e.g., skin, hair and/or nail) of the subject, by contacting the application site with a part of the composition that comprises the enzyme and thereafter contacting the application site with a part of the composition that comprises the enzyme's substrate, or *vice versa*, contacting the application site with a part of the composition that comprises the enzyme's substrate and thereafter contacting the application site with a part of the composition that comprises the enzyme.

In some embodiments of any of the embodiments described herein, the composition is prepared shortly before being contacted with the application site (e.g., skin, hair and/or nail) of the subject, by contacting a part of the composition that comprises the enzyme's substrate with a part of the composition that comprises the enzyme's substrate, and thereafter contacting the resulting composition with the application site (e.g., skin, hair and/or nail) of a subject.

The composition may be prepared from components packaged in a kit, as described in further detail hereinunder.

In some embodiments of any of the embodiments described herein, the composition is packaged in a container, and can be used well after packaging.

5 In some of any of these embodiments, a composition comprises the hydrogen peroxide-producing enzyme in contact with its substrate, but the enzyme in the composition is inhibited from producing hydrogen peroxide due to a lack of oxygen in the container.

10 In some embodiments of any of the embodiments described herein, the container is air-tight, in order to limit the amount of oxygen which can be used by the enzyme to produce hydrogen peroxide.

15 Herein, the term “air-tight” refers to a packaging and/or container sealed in such a manner such that there are no openings which allow gases such as oxygen to enter or escape, and wherein packaging and/or container is formed of a material which is substantially impermeable to such gases.

In some embodiments of any of the embodiments described herein, air is evacuated from the packaging and/or container before sealing, for example, by subjecting the packaging and/or container to a reduced pressure and/or by purging the packaging and/or container with a gas which does not contain oxygen (e.g., nitrogen).

20 In some embodiments of any of the embodiments described herein, the air-tight container is configured for dispensing the composition (e.g., dispensing more than once) without entry of air into the container, so as to avoid production of hydrogen peroxide in the container after dispensing some of the composition.

25 In some embodiments of any of the embodiments described herein, the container has a volume which decreases upon dispensing, such that the container has little or no room for air to enter, even when the volume of the composition in the container decreases upon dispensing. Examples of such containers include, for example, a flexible tube (e.g., similar to a toothpaste tube) which dispenses a composition when subjected to pressure, and a container comprising a piston which moves when dispensing a composition.

30 In some embodiments of any of the embodiments described herein, an opening of the container (e.g., a nozzle) comprises a valve configured so as to be open only when a

composition is exiting the container through the opening, at which time the composition being dispensed obstructs air from entering through the opening.

In some embodiments of any of the embodiments described herein, a composition comprises the hydrogen peroxide-producing enzyme in contact with its substrate, but the enzyme in the composition is inhibited from producing hydrogen peroxide due to a lack of water or humidity in the container. Means for packaging in non-humid environment are well known in the art.

In embodiments where the composition is packaged in a container, the container can be identified for use in any cosmetic application that requires hydrogen peroxide, as described in further detail hereinafter.

The concentrations of the hydrogen peroxide-producing enzyme and the substrate thereof may be selected so as to result in a suitable rate of hydrogen peroxide production. The skilled person can determine a suitable rate of hydrogen peroxide based on a desired concentration (e.g., steady state concentration) of hydrogen peroxide (which will depend on an intended use of the composition) and on a rate at which the hydrogen peroxide is consumed, which will depend, for example, on a concentration and reactivity of compounds which react with hydrogen peroxide (e.g., agents such as described herein).

In some embodiments of any of the embodiments described in the context of a hydrogen peroxide-producing composition, a concentration of the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) in the composition is in a range of from 0.125 to 1250 units/gram. In some embodiments, the concentration is in a range of from 1.25 to 125 units/gram. In some embodiments, the concentration is in a range of from 4 to 40 units/gram. In some embodiments, the concentration is in a range of from 6 to 20 units/gram. In some embodiments, the concentration is about 12.5 units/gram.

In some embodiments of any of the embodiments described in the context of a hydrogen peroxide-producing composition, a concentration of the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) in the composition is in a range of from 1.25 to 1250 units/gram. In some embodiments, the concentration is in a range of from 4 to 1250 units/gram. In some embodiments, the concentration is in a range of from 12.5 to 1250 units/gram.

In some embodiments of any of the embodiments described in the context of a hydrogen peroxide-producing composition, a concentration of the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) in the composition is in a range of from 0.125 to 125 units/gram. In some embodiments, the concentration is in a range of from 0.125 to 40 units/gram. In some embodiments, the concentration is in a range of from 0.125 to 12.5 units/gram.

As used herein and in the art, a “unit” of an enzyme refers to an amount of the enzyme which catalyzes a conversion of 1 micromole of a substrate (e.g., a substrate as described herein) per minute, at a temperature of 25 °C and at substrate concentrations which yield a maximal conversion rate.

In some embodiments of any of the embodiments described in the context of a hydrogen peroxide-producing composition, a concentration of the substrate of the hydrogen peroxide-producing enzyme (e.g., glucose) in the composition is in a range of from 0.0175 μmole/gram to 175 μmole/gram. In some embodiment, the concentration is in a range of from 0.175 to 17.5 μmole/gram. In some embodiment, the concentration is in a range of from 0.5 to 6 μmole/gram. In some embodiment, the concentration is in a range of from 1 to 3.5 μmole/gram. In some embodiment, the concentration is about 1.75 μmole/gram.

In some embodiments of any of the embodiments described in the context of a hydrogen peroxide-producing composition, a concentration of the substrate of the hydrogen peroxide-producing enzyme (e.g., glucose) in the composition is in a range of from 0.175 μmole/gram to 175 μmole/gram. In some embodiment, the concentration is in a range of from 0.5 to 175 μmole/gram. In some embodiment, the concentration is in a range of from 1.75 to 175 μmole/gram.

In some embodiments of any of the embodiments described in the context of a hydrogen peroxide-producing composition, a concentration of the substrate of the hydrogen peroxide-producing enzyme (e.g., glucose) in the composition is in a range of from 0.0175 μmole/gram to 17.5 μmole/gram. In some embodiment, the concentration is in a range of from 0.0175 to 6 μmole/gram. In some embodiment, the concentration is in a range of from 0.0175 to 1.75 μmole/gram.

In some embodiments of any of the embodiments described in the context of a hydrogen peroxide-producing composition, a concentration of the hydrogen peroxide-

producing enzyme (e.g., glucose oxidase) in the composition is in a range of from 0.125 to 1250 units/gram, and a concentration of the substrate of the hydrogen peroxide-producing enzyme (e.g., glucose) in the composition is in a range of from 0.0175  $\mu$ mole/gram to 175  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 1.25 to 125 units/gram. In some embodiments, the enzyme concentration is in a range of from 4 to 40 units/gram. In some embodiments, the enzyme concentration is in a range of from 6 to 20 units/gram. In some embodiments, the enzyme concentration is about 12.5 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is glucose.

10 In some embodiments of any of the embodiments described in the context of a hydrogen peroxide-producing composition, a concentration of the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) in the composition is in a range of from 0.125 to 1250 units/gram, and a concentration of the substrate of the hydrogen peroxide-producing enzyme (e.g., glucose) in the composition is in a range of from 0.175  $\mu$ mole/gram to 175  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 1.25 to 125 units/gram. In some embodiments, the enzyme concentration is in a range of from 4 to 40 units/gram. In some embodiments, the enzyme concentration is in a range of from 6 to 20 units/gram. In some embodiments, the enzyme concentration is about 12.5 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is glucose.

15 In some embodiments of any of the embodiments described in the context of a hydrogen peroxide-producing composition, a concentration of the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) in the composition is in a range of from 0.125 to 1250 units/gram, and a concentration of the substrate of the hydrogen peroxide-producing enzyme (e.g., glucose) in the composition is in a range of from 0.0175  $\mu$ mole/gram to 17.5  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 1.25 to 125 units/gram. In some embodiments, the enzyme concentration is in a range of from 4 to 40 units/gram. In some embodiments, the enzyme concentration is in a range of from 6 to 20 units/gram. In some embodiments, the enzyme concentration is about 12.5 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is glucose.

In some embodiments of any of the embodiments described in the context of a hydrogen peroxide-producing composition, a concentration of the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) in the composition is in a range of from 0.125 to 1250 units/gram, and a concentration of the substrate of the hydrogen peroxide-producing enzyme (e.g., glucose) in the composition is in a range of from 0.175 to 17.5  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 1.25 to 125 units/gram. In some embodiments, the enzyme concentration is in a range of from 4 to 40 units/gram. In some embodiments, the enzyme concentration is in a range of from 6 to 20 units/gram. In some embodiments, the enzyme concentration is about 10 12.5 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is glucose.

In some embodiments of any of the embodiments described in the context of a hydrogen peroxide-producing composition, a concentration of the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) in the composition is in a range of from 0.125 to 1250 units/gram, and a concentration of the substrate of the hydrogen peroxide-producing enzyme (e.g., glucose) in the composition is in a range of from 0.5 to 175  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 1.25 to 125 units/gram. In some embodiments, the enzyme concentration is in a range of from 4 to 40 units/gram. In some embodiments, the enzyme concentration is in a range of from 6 to 20 units/gram. In some embodiments, the enzyme concentration is about 15 12.5 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is glucose.

In some embodiments of any of the embodiments described in the context of a hydrogen peroxide-producing composition, a concentration of the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) in the composition is in a range of from 0.125 to 1250 units/gram, and a concentration of the substrate of the hydrogen peroxide-producing enzyme (e.g., glucose) in the composition is in a range of from 0.0175 to 6  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 1.25 to 125 units/gram. In some embodiments, the enzyme concentration is in a range of from 4 to 40 units/gram. In some embodiments, the enzyme concentration is in a range of from 6 to 20 units/gram. In some embodiments, the enzyme concentration is about 20 25 30 30 units/gram.

12.5 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is glucose.

In some embodiments of any of the embodiments described in the context of a hydrogen peroxide-producing composition, a concentration of the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) in the composition is in a range of from 0.125 to 1250 units/gram, and a concentration of the substrate of the hydrogen peroxide-producing enzyme (e.g., glucose) in the composition is in a range of from 0.5 to 6  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 1.25 to 125 units/gram. In some embodiments, the enzyme concentration is in a range of from 4 to 40 units/gram. In some embodiments, the enzyme concentration is in a range of from 6 to 20 units/gram. In some embodiments, the enzyme concentration is about 12.5 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is glucose.

In some embodiments of any of the embodiments described in the context of a hydrogen peroxide-producing composition, a concentration of the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) in the composition is in a range of from 0.125 to 1250 units/gram, and a concentration of the substrate of the hydrogen peroxide-producing enzyme (e.g., glucose) in the composition is in a range of from 1.75 to 175  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 1.25 to 125 units/gram. In some embodiments, the enzyme concentration is in a range of from 4 to 40 units/gram. In some embodiments, the enzyme concentration is in a range of from 6 to 20 units/gram. In some embodiments, the enzyme concentration is about 12.5 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is glucose.

In some embodiments of any of the embodiments described in the context of a hydrogen peroxide-producing composition, a concentration of the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) in the composition is in a range of from 0.125 to 1250 units/gram, and a concentration of the substrate of the hydrogen peroxide-producing enzyme (e.g., glucose) in the composition is in a range of from 0.0175 to 1.75  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 1.25 to 125 units/gram. In some embodiments, the enzyme concentration is in a range of from 4 to 40 units/gram. In some embodiments, the enzyme concentration is in a range

of from 6 to 20 units/gram. In some embodiments, the enzyme concentration is about 12.5 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is glucose.

In some embodiments of any of the embodiments described in the context of a 5 hydrogen peroxide-producing composition, a concentration of the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) in the composition is in a range of from 0.125 to 1250 units/gram, and a concentration of the substrate of the hydrogen peroxide-producing enzyme (e.g., glucose) in the composition is in a range of from 1 to 3.5  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 10 1.25 to 125 units/gram. In some embodiments, the enzyme concentration is in a range of from 4 to 40 units/gram. In some embodiments, the enzyme concentration is in a range of from 6 to 20 units/gram. In some embodiments, the enzyme concentration is about 12.5 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is glucose.

15 It is to be noted that for any one of the embodiments pertaining to a composition as described herein, any one of the hydrogen peroxide producing enzymes, including any of the embodiments described herein with respect thereto, in any combination, can be utilized (unless other indicated).

In some embodiments of any of the embodiments and any of the aspects 20 described herein, the composition is for obtaining hydrogen peroxide *per se* as an active agent. The hydrogen peroxide may be intended, for example, for lightening a topical bodily site of a subject (e.g., a region of the skin, hair and/or nail of a subject) by direct reaction (e.g., bleaching) with the topical bodily site, for exhibiting an anti-microbial activity, or for exhibiting an odor removal activity (e.g., removal of skunk odor).

25 In some embodiments of any of the embodiments and any of the aspects described herein, the composition further comprises at least one agent which exhibits a cosmetic or pharmaceutical activity in a presence of hydrogen peroxide. In some embodiment, the composition is for obtaining the activity of such an agent, by generating hydrogen peroxide in a presence of the agent.

30 Herein, “an agent that exhibits a cosmetic or pharmaceutical activity in the presence of hydrogen peroxide” encompasses agents that are activated in the presence of hydrogen peroxide. By “activated” it is meant that the agent is chemically modified

(e.g., oxidized) in the presence of hydrogen peroxide and its oxidized form exhibits a desired cosmetic or pharmaceutical activity; or the agent is an enzyme which requires hydrogen peroxide for exhibiting its activity (e.g., requires hydrogen peroxide as a substrate or as a co-activator).

5 Non-limiting examples of agents which exhibit a cosmetic or pharmaceutical activity in a presence of hydrogen peroxide include, for example, peroxidases and oxidative dye precursors.

10 Oxidative dye precursors are compounds which react to form a colorant (e.g., a permanent dye) upon oxidation, for example, for use in coloring hair. Hydrogen peroxide is commonly used for oxidizing such dye precursors, and many suitable oxidative dye precursors and their precise uses will be known to the skilled person. In some embodiments of any of the embodiments described herein, the composition comprises at least one oxidative dye precursor for coloring hair.

15 Peroxidases are enzymes which catalyze a formation of a reactive compound, such as a reactive oxygen species (ROS), in a presence of hydrogen peroxide. A peroxidase thereby enhances a reactivity of the hydrogen peroxide, by converting it to a more reactive compound. Such a composition may be useful, for example, for lightening a topical bodily site by reaction of the reactive compound with the bodily site, for exhibiting an anti-microbial activity, or for exhibiting an enhanced odor removal 20 activity (e.g., removal of skunk odor).

25 In some embodiments of any of the embodiments described herein, a peroxidase oxidizes thiocyanate to hypothiocyanate and/or at least one halide (e.g., chloride, bromide and/or iodide) to a hypohalite (e.g., hypochlorite, hypobromite and/or hypoiodite). Examples of such peroxidases include, without limitation, myeloperoxidase and eosinophil peroxidase (which can each oxidize, for example, thiocyanate, chloride, bromide and iodide), lactoperoxidase (which can oxidize, for example, thiocyanate, bromide and iodide), ovoperoxidase and vanadium bromoperoxidase (which can each oxidize, for example, bromide and iodide), *Murex* bromoperoxidase (which can oxidize, for example, bromide), horseradish peroxidase 30 (which can oxidize, for example, thiocyanate and iodide) and thyroid peroxidase (which can oxidize, for example, iodide).

In some embodiments of any of the embodiments described herein, the composition further comprises a substrate (e.g., thiocyanate, chloride, bromide, iodide, and/or an oxidizing mediator as described herein) which is oxidized in the presence of hydrogen peroxide and the peroxidase.

5       Herein, the phrase “oxidizing mediator” refers to a molecule which facilitates a redox reaction catalyzed by the peroxidase, for example, by increasing the oxidative potential and/or the stability of a peroxidase, and/or by acting as a co-factor for the peroxidase, and/or by acting as a substrate of the peroxidase which forms a reactive intermediate product (e.g., a free radical). Oxidizing mediators include compounds  
10      (e.g., aromatic compounds) capable of stabilizing electrons (e.g., by delocalization) transferred during a reaction catalyzed by the peroxidase.

In some embodiments of any of the embodiments described herein, the peroxidase oxidizes a substrate (e.g., thiocyanate, chloride, bromide, iodide) present in or on the topical bodily site in the presence of hydrogen peroxide.

15       Other cosmetic or pharmaceutical activities of peroxidases may also be utilized in the context of any of the embodiments of this aspect of the invention.

In some embodiments of any of the embodiments described herein, the peroxidase is capable, in a presence of hydrogen peroxide, of degrading melanin in a subject (e.g., in skin and/or hair). In some embodiments, a presence of an oxidizing mediator is necessary for the peroxidase to degrade melanin in the presence of hydrogen  
20      peroxide. Degradation of melanin is useful for lightening skin and/or hair. Lignin peroxidase is an exemplary peroxidase capable of degrading melanin.

Without being bound by any particular theory, it is believed that some peroxidases (e.g., lignin peroxidase) degrade melanin by oxidizing an oxidizing mediator (which serves as a substrate) so as to form a free radical, and the free radical then reacts so as to degrade melanin.

In some embodiments of any of the embodiments and any of the aspects described herein, the composition is a hair-dye composition. Herein, the phrase “hair-dye composition” refers to any composition used in the coloring (e.g., permanent  
30      dyeing) of hair, including compositions which comprise a hair dye and/or precursor thereof and compositions which do not comprise a hair dye or precursor thereof, but

which are intended to be contacted with a hair dye and/or precursor thereof (e.g., on the hair).

In some embodiments of any of the embodiments and any of the aspects described herein, the composition is an odor-removal composition, for example, for removal of skunk odor (e.g., from skin, hair and/or nail). In some embodiments, the odor-removal composition further comprises a detergent (e.g., sodium lauryl sulfate) and/or an alkaline agent (e.g., sodium bicarbonate).

In some embodiment of any of the embodiments and any of the aspects described herein, the composition is an anti-microbial composition. Herein, the phrase “anti-microbial composition” refers to any composition for killing or otherwise inhibiting microscopic animals, plants, fungi, bacteria, archaeabacteria and/or viruses.

In some embodiments of any of the embodiments and any of the aspects described herein, the composition is a bleaching composition. Herein, the phrase “bleaching composition” refers to any composition for removing at least a portion of a color, and includes, for example, compositions for lightening skin and/or portions of skin, hair bleaching compositions, and compositions for removing stains from nails.

As exemplified herein, lignin peroxidase is capable of degrading melanin, thereby lightening skin and/or hair.

Lignin peroxidase is an exemplary agent which exhibits a cosmetic activity in a presence of hydrogen peroxide.

In some embodiments, lignin peroxidase is utilized in combination with any one of the embodiments described herein for a composition and/or hydrogen peroxide producing enzyme, and any combination thereof.

In some embodiments of any of the embodiments described herein, the skin to be lightened comprises facial skin (e.g., forehead and/or cheeks).

In some embodiments of any of the embodiments described herein, the skin to be lightened comprises a skin region exposed to the sun (e.g., a tanned region).

In some embodiments of any of the embodiments described herein, lightening skin comprises lightening a whole skin complexion.

In some embodiments of any of the embodiments described herein, the skin to be lightened comprises a relatively dark portion of skin, for example, skin comprising an

uneven tone, one or more dark spots, one or more freckles, melasma, hyperpigmentation of skin, skin discoloration, one or more age spots, one or more acne marks and/or a scar.

As described herein, a desired concentration of hydrogen peroxide may be determined by selecting suitable concentrations of hydrogen peroxide-producing enzyme and the substrate thereof and compounds which react with hydrogen peroxide (if any described herein).

In some embodiments of any of the embodiments described herein, a composition for obtaining hydrogen peroxide *per se* (e.g., a bleaching composition) has concentrations of ingredients selected so as to result in a relatively high hydrogen peroxide concentration, for example, from 0.5 to 12 weight percents. In some embodiments, the hydrogen peroxide concentration is no more than 6 weight percents. In some embodiments, the hydrogen peroxide concentration is no more than 3 weight percents. A relatively high hydrogen peroxide concentration may be obtained, for example, by minimizing a presence of compounds which react with hydrogen peroxide and/or by selecting higher concentration of the component of the enzymatic system, as described herein.

In some embodiments of any of the embodiments described herein, a composition comprising an agent exhibiting an activity in a presence of hydrogen peroxide (e.g., as described herein) has concentrations of ingredients selected so as to result in a hydrogen peroxide concentration suitable for obtaining the activity of the agent. In some embodiments, the hydrogen peroxide concentration is relatively low, for example, no more than 0.5 weight percents. In some embodiments, the hydrogen peroxide concentration is no more than 0.1 weight percents. In some embodiments, the hydrogen peroxide concentration is no more than 0.02 weight percents. It is to be appreciated that the agent may help to maintaining a relatively low hydrogen peroxide concentration by reacting with the hydrogen peroxide. The agent may continuously react with the generated hydrogen peroxide such that a low concentration of hydrogen peroxide is continuously maintained.

Low concentration of hydrogen peroxide may be especially advantageous, for example, in a composition comprises an agent (e.g., a protein such as a peroxidase) which may be sensitive to (e.g., adversely affected by) high peroxide concentrations and/or which may be highly efficient at reacting with hydrogen peroxide. Such agents

which may be adversely affected by high hydrogen peroxide concentrations include, for example, certain peroxidases (e.g., lignin peroxidase), which are known for, for example, undergoing “self-suicidality” in the presence of high concentrations of hydrogen peroxide.

5

***Kits for obtaining the hydrogen peroxide-producing composition:***

It is to be understood that a hydrogen peroxide-producing composition according to any one of the embodiments relating to such compositions may be obtained using a kit according to any one of the embodiments described herein relating to such kits (e.g., as described in this section). The skilled person will be capable of selecting a suitable kit 10 as described herein so as to obtain any desired composition described herein, based on the guidance provided herein.

For any one of the embodiments describing a kit, any one of the hydrogen peroxide-producing enzymes and/or compositions, including any one of the respective embodiments thereof, and any combination thereof, can be utilized.

15

In some embodiments of any of the embodiments described herein, any of the compositions as described herein can be considered as a two (or more)-part composition, in which one part of the composition comprises a hydrogen peroxide-producing enzyme as described herein and one part of the composition comprises a substrate of the enzyme.

20

In some embodiments of any of the embodiments described herein, the two-part composition is packaged in a kit and is identified for use in any cosmetic or pharmaceutical application that utilizes hydrogen peroxide, either *per se*, or in combination with another agent, as described herein.

25

According to an aspect of some embodiments of the invention, there is provided a kit for forming and/or applying (topically) to a topical bodily site of a subject a cosmetic or pharmaceutical composition, as described herein, wherein the cosmetic or pharmaceutical composition comprises hydrogen peroxide (e.g., in accordance with a method described herein).

30

In some embodiments of any of the embodiments described herein, the composition is packaged in the kit such that all of its components are packaged together in a container, under conditions that inhibit hydrogen peroxide formation, as described herein. In some of these embodiments, once the composition is dispensed from the

container and applied on the bodily site, its exposure to air and/or humidity promotes a reaction between the hydrogen peroxide-producing enzyme and its substrate and hydrogen peroxide is produced.

In some embodiments of any of the embodiments described herein, a part of the 5 composition which comprises a hydrogen peroxide-producing enzyme and a part of the composition which comprise the substrate are packaged individually within the kit, such that the composition is formed once these two parts of the composition are mixed.

By "packaged individually" it is meant that the there is no direct contact between the parts of the composition (or the first and second compositions as described herein) 10 when in the kit.

For example, each part of the composition (e.g., a first composition and a second composition, as described herein) is individually packaged within a sealed container in the kit. The container can be made of e.g., plastic, wood, nylon, glass, fabric, metal, leather and/or metal foil.

15 One or more of the containers may include means for dispensing its content, either directly to the topical bodily site or to a receptacle for mixing with other parts of the composition which are packaged individually within the kit. Alternatively, the kit further comprises means for dispensing the contents of one or more of the containers, and/or means for mixing the contents of the container and means for dispensing the 20 obtained mixture on the topical bodily site.

In some embodiments of any of the embodiments described herein, the kit 25 comprises a first composition comprising a hydrogen peroxide-producing enzyme (e.g., as described herein) and a second composition comprising a substrate of the hydrogen peroxide-producing enzyme (e.g., as described herein). In some embodiments, the first and second compositions of the kit are packaged individually in the kit. The first composition and second composition of the kit may be combined so as to prepare the cosmetic or pharmaceutical composition (e.g., a composition described herein) in which the enzyme contacts the substrate, thereby producing hydrogen peroxide in the composition.

30 In some embodiments of any of the embodiments described herein, the kit further comprises at least one agent which exhibits a cosmetic or pharmaceutical activity in a presence of hydrogen peroxide (e.g., as described herein). The agent(s)

may be included within the first composition, the second composition and/or in a third composition within the kit, which is individually packaged in the kit. The first composition and second composition and optional third composition of the kit may be combined so as to prepare the cosmetic or pharmaceutical composition comprising the 5 agent(s) (e.g., as described herein).

Combining the first and second compositions of the kit can be effected immediately before application to the topical bodily site (e.g., between 1 minute and 10 minutes before application), or during application to the topical bodily site.

Minimizing a time interval between combination of the compositions may be 10 advantageous in avoiding a situation in which hydrogen peroxide is being produced in the presence of an agent, such as a peroxidase or an oxidative dye precursor, which reacts with the hydrogen peroxide (e.g., as described herein) when the composition is not in contact with the topical body site, as such a situation could result in non-productive consumption (e.g., a waste) of the hydrogen peroxide.

15 Combining the first and second compositions of the kit during application can be effected, for example, by co-administering the first and second compositions of the kit to the topical bodily site.

By “co-administering” it is meant that the first and second compositions are 20 topically administered to the bodily site at the same time (concomitantly) or, that the first and second compositions of the kit are administered sequentially, in any order, whereby the time interval between administration of the first composition and administration of the second composition may range from 0.1 minute to 5 minute. In some embodiments, the time interval is no more than 2 minutes.

If a third composition is included in the kit, it can be combined with the first and 25 second composition before application to the bodily site, or be co-administered to the bodily site with the first and second compositions.

The third composition can be co-administered concomitantly with the first and/or second composition, before administration of the first and/or second composition, or thereafter, with the time interval between administrations is as 30 described hereinabove.

In some embodiments of any of the embodiments described herein, the kit further comprises written instructions for how to use a composition(s) therein, for example, an

amount of a composition to be applied, a sequence in which different compositions are to be applied, a timing and/or frequency of applications, and/or a duration of treatment (e.g., as described herein).

5 In some embodiments of any of the embodiments described herein, the kit further comprises a wash composition, such as a facial wash, for cleaning a topical body site prior to contact with a composition described herein.

According to exemplary embodiments, the facial wash comprises ingredients as presented in Table 1:

10

**Table 1: Facial wash according to exemplary embodiments**

Ingredient	Weight percentage range
Water	25 – 50 %
Propanediol	10 – 25 %
Glycerin	10 – 25 %
Stearic acid	10 – 25 %
Myristic acid	5 – 10 %
Potassium hydroxide	5 – 10 %
Lauric acid	1 – 5 %
Decyl glucoside	1 – 5 %
Glycol distearate	1 – 5 %
Glyceryl stearate	1 – 5 %
PEG-100 stearate	1 – 5 %
Synthetic beeswax	1 – 5 %
Potassium cocoyl glycinate	0 - 0.1 %
Butylated hydroxytoluene (BHT)	0 - 0.1 %
Lignin peroxidase	0 - 0.1 %

In some embodiments of any of the embodiments described herein, the kit includes appropriate instructions for use and labels indicating approval by a regulatory agency for the indicated use, as described herein.

***Cosmetic applications of a hydrogen peroxide-producing composition:***

As exemplified herein, by providing a hydrogen peroxide-producing enzyme in contact with its substrate, the compositions described herein are effective at administering hydrogen peroxide to a surface.

5 According to an aspect of embodiments of the invention, there is provided a method of topically administering hydrogen peroxide to a topical bodily site of a subject (e.g., to skin, hair and/or nail of a subject). The method, according to some of these embodiments, is effected by contacting a topical bodily site, or a region thereof, of the subject with a composition (e.g., a composition described herein) comprising a hydrogen 10 peroxide-producing enzyme and a substrate of the hydrogen peroxide-producing enzyme. In some embodiments, the composition is formed by a hydrogen peroxide-producing enzyme and a substrate thereof as packaged within a kit as described herein.

15 It is to be understood that a method of topically administering hydrogen peroxide according to any one of the embodiments relating to such a method (e.g., as described in this section) may utilize a hydrogen peroxide-producing system and/or a kit according to any one of the embodiments described herein relating to such systems and/or kits.

20 As used herein the terms “topical” and “topically” refer to a topical body site, as defined herein, including, for example, to administration of a composition by application onto a topical body site, and to compositions for administration in such a manner.

25 Herein, the phrase “topical body site” or “topical bodily site” refers to a skin surface (including any particular region of the skin surface), and/or hair (including all hair and/or any particular region of hair) from any part of the body, and/or a nail surface (including all fingernails and/or toenails, and/or any individual nail and/or a region thereof). The skin may be of any part of a body, for example, facial skin, neck skin, arm skin, hand skin, torso skin and/or leg skin. The hair may be, for example, head hair, facial hair and/or pubic hair. This phrase encompasses also a certain region or part of a skin surface (e.g., a part of a facial skin, a part of an arm or leg skin), or of a hair (e.g., a part of facial, head or pubic hair), as described in further detail elsewhere herein.

30 In some embodiments of any of the embodiments described herein, the skin of a topical body site does not include mucous membranes such as those of the mouth, lips, eyelids, ears, genital area and anus.

In some embodiments of any of the embodiments described herein, the composition is formed by contacting the topical bodily site or a region thereof with the hydrogen peroxide-producing enzyme and/or the substrate thereof subsequent to contacting the topical bodily site or a region thereof with the agent which exhibits a cosmetic or pharmaceutical activity in a presence of hydrogen peroxide. In such a sequence, hydrogen peroxide will only be generated in a presence of the agent which exhibits a cosmetic or pharmaceutical activity in a presence of hydrogen peroxide, which may allow for a more efficient utilization of the hydrogen peroxide.

In some embodiments of any of the embodiments described herein, the composition is formed by contacting the topical body site with the hydrogen peroxide-producing enzyme before contacting the topical body site with the substrate.

In some embodiments of any of the embodiments described herein, the composition is formed by contacting the topical body site with the hydrogen peroxide-producing enzyme after contacting the topical body site with the substrate.

In some embodiments of any of the embodiments described herein, the composition is formed by contacting the topical body site with the hydrogen peroxide-producing enzyme concomitantly with contacting the topical body site with the substrate.

The agent which exhibits a cosmetic or pharmaceutical activity in a presence of hydrogen peroxide may be contacted with the topical body site (or region thereof) before, after or concomitantly with contacting the topical region with any of the other components, for example, by co-administering (as defined herein) the agent with the hydrogen peroxide-producing enzyme and/or the substrate thereof.

In some embodiments of any of the embodiments described herein, the method is for coloring hair, and the method further comprising contacting the hair with at least one oxidative dye precursor for coloring hair, as described herein.

In some embodiments of any of the embodiments described herein, the method further comprises contacting the topical body site with a peroxidase (e.g., as described herein). Such a method, may be useful, for example, for lightening a topical bodily site, for exhibiting an anti-microbial activity, or for exhibiting an enhanced odor removal activity (e.g., as described herein).

In some embodiments of any of the embodiments described herein, the method further comprises contacting the topical body site with a substrate and/or oxidizing mediator of the peroxidase (e.g., as described herein).

5 In some embodiments of any of the embodiments described herein, the method is for removal of skunk odor (e.g., from a topical body site). In some embodiments, the method further comprises contacting the topical body site with a detergent (e.g., sodium lauryl sulfate) and/or an alkaline agent (e.g., sodium bicarbonate).

10 In some embodiments of any of the embodiments described herein, the method is for obtaining an anti-microbial activity, for example, by forming an anti-microbial composition (e.g., as described herein).

15 In some embodiments of any of the embodiments described herein, the method is for bleaching a topical body site or a region thereof, for example, by forming a bleaching composition (e.g., as described herein). The method may comprise, for example, lightening skin and/or portions of skin, bleaching hair, and/or removing stains from nails.

In some embodiments of any of the embodiments described herein, the method is for lightening facial skin (e.g., forehead and/or cheeks).

In some embodiments of any of the embodiments described herein, the method is for lightening a skin region exposed to the sun (e.g., a tanned region).

20 In some embodiments of any of the embodiments described herein, the method is for lightening a whole skin complexion.

25 In some embodiments of any of the embodiments described herein, the method is for lightening a relatively dark portion of skin, for example, skin comprising an uneven tone, one or more dark spots, one or more freckles, melasma, hyperpigmentation of skin, skin discoloration, one or more age spots, one or more acne marks and/or a scar.

In some embodiments of any of the embodiments described herein, the method is effected repeatedly, optionally even indefinitely, in order to respond to repeated color formation, such as newly formed melanin in skin, new growth of dark hair, chronic stain formation in nails, and so forth.

30 In some embodiments of any of the embodiments described herein, the method is effected by contacting the topical region with the composition at least 5 times. In some embodiments, the method is effected by contacting the topical region with the

composition at least 10 times. In some embodiments, the method is effected by contacting the topical region with the composition at least 20 times. In some embodiments, the method is effected by contacting the topical region with the composition at least 50 times. In some embodiments, the method is effected by 5 contacting the topical region with the composition at least 100 times. In some embodiments, the method is effected by contacting the topical region with the composition at least 200 times.

In some embodiments of any of the embodiments described herein, the method is effected by contacting the topical region once or twice per day. In exemplary 10 embodiments, contacting is effected twice per day.

In some embodiments of any of the embodiments described herein, the method is effected for at least one week. In some embodiments, the method is effected for at least two weeks. In some embodiments, the method is effected for at least three weeks. In some embodiments, the method is effected for at least four weeks. In some 15 embodiments, the method is effected for at least six weeks. In some embodiments, the method is effected for at least eight weeks. In some embodiments, the method is effected for at least three months. In some embodiments, the method is effected for at least four months. In some embodiments, the method is effected for at least six months. In some embodiments, the method is effected for at least one year.

20 The dosing and course of treatment will depend on the type of treatment and the severity of an affliction being treated, with course of treatment lasting from one day to several weeks or longer, or until cure is effected or diminution of the condition being treated is achieved.

The amount of a composition to be administered will, of course, be dependent 25 on the subject being treated, the degree of a condition being treated, the manner of administration, the judgment of a prescribing physician or a cosmetician or the consumer, the desired effect of the composition, etc.

In some embodiments of any of the embodiments described herein, the topical body site is cleaned (e.g., with a wash, as described herein) prior to contact with a 30 composition described herein.

In embodiments where the method is, for example, for bleaching or coloring hair, the method can be effected once, for achieving a desired effect and can be repeated once the desired effect is reduced or diminished.

5 In embodiments where the method is, for example, for treating an infection, the treatment is effected once, until the infection is eliminated, and the number of administrations and treatment duration is effected until cure is effected.

***Hydrogen peroxide-producing compositions, methods and kits for lightening skin and/or hair:***

10 An exemplary embodiment of the compositions, methods and kits described herein is a composition for lightening a topical body site such as skin and/or hair in which a hydrogen peroxide-producing composition as described herein is used in combination with lignin peroxidase.

15 It is to be understood that a composition, method and/or kit according to any one of the embodiments relating to such compositions, methods and/or kits may be utilized for lightening kin and/or hair, as described herein (e.g., in this section).

20 In some embodiments of any of the embodiments described herein, there is provided a cosmetic composition for lightening a topical body site such as skin and/or hair of a subject (e.g., as described herein), the composition comprising a lignin peroxidase, a hydrogen peroxide-producing enzyme (e.g., as described herein) and a substrate of the hydrogen peroxide-producing enzyme (e.g., as described herein).

25 According to another aspect of embodiments described herein, there is provided a cosmetic method of lightening a topical body site such as skin and/or hair of a subject (e.g., as described herein), the method comprising contacting a topical region of a subject with a composition comprising a lignin peroxidase, a hydrogen peroxide-producing enzyme and a substrate of the hydrogen peroxide-producing enzyme (e.g., a lignin peroxidase-containing composition described herein). In some embodiments, the composition is formed by a lignin peroxidase, a hydrogen peroxide-producing enzyme and a substrate thereof as packaged within a kit described herein.

30 In some embodiments of any of the embodiments described herein, the composition is formed by contacting the topical region with the hydrogen peroxide-producing enzyme and/or the substrate thereof subsequent to contacting the topical region with the lignin peroxidase. In such a sequence, hydrogen peroxide will only be

generated in a presence of the lignin peroxidase, which may allow for a more efficient utilization of the hydrogen peroxide.

Compositions comprising lignin peroxidase, as described herein, can be provided at higher concentrations and be prescribed by a physician as a pharmaceutical or a 5 cosmetic composition to treat skin pigmentation disorders such as melasma, chloasma, ochronosis, and lentigo, and/or to lighten whole skin complexion.

In some embodiments of any of the embodiments described herein, the method is effected by contacting the topical region with the composition at least 5 times. In some 10 embodiments, the method is effected by contacting the topical region with the composition at least 10 times. In some embodiments, the method is effected by contacting the topical region with the composition at least 20 times. In some embodiments, the method is effected by contacting the topical region with the composition at least 50 times. In some embodiments, the method is effected by contacting the topical region with the composition at least 100 times. In some 15 embodiments, the method is effected by contacting the topical region with the composition at least 100 times.

In some embodiments of any of the embodiments described herein, the method is effected by contacting the topical region once or twice per day. In exemplary 20 embodiments, contacting is effected twice per day.

In some embodiments of any of the embodiments described herein, the method is effected for at least one week. In some embodiments, the method is effected for at least 25 two weeks. In some embodiments, the method is effected for at least three weeks. In some embodiments, the method is effected for at least four weeks. In some embodiments, the method is effected for at least six weeks. In some embodiments, the method is effected for at least eight weeks. In some embodiments, the method is effected for at least three months. In some embodiments, the method is effected for at least four months. In some embodiments, the method is effected for at least six months. In some 30 embodiments, the method is effected for at least one year.

The dosing and course of treatment will depend on the degree of color (e.g., 35 pigmentation) in the region to be treated, for example, a degree of skin pigmentation in a skin pigmentation disorder (e.g., chloasma, melasma, ochronosis and lentigo), the tone of skin color, skin complexion and/or the responsiveness of the skin, with course of

treatment lasting from several days to several weeks or longer, or until cure is effected or diminution of the skin disorder is achieved.

The amount of a composition to be administered will, of course, be dependent on the subject being treated, the degree of a condition being treated, the manner of 5 administration, the judgment of a prescribing physician or a cosmetician or the consumer, etc.

***Lignin peroxidase:***

It is to be understood that lignin peroxidase according to any one of the 10 embodiments described herein relating to lignin peroxidase (e.g., as described in this section) may be used in combination with a hydrogen peroxide-producing enzymatic system according to any one of the embodiments described herein relating to such a system.

As used herein the phrase "lignin peroxidase" refers to an enzyme which plays a major role in lignin degradation. Lignin peroxidase is able to catalyze the oxidation of 15 substrates with high redox potential. This unique ability is consistent with a heme active site of low electron density, which is indicated by high redox potential [Cai and Tien, *J Biotechnol* 1993, 30: 79-90].

Examples of lignin peroxidases are classified as EC 1.11.1.14.

Manganese peroxidase (EC 1.11.1.13) is involved in degradation of lignin in 20 Basidiomycetes (e.g., *Phanerochaete chrysosporium*), and is also defined herein as a lignin degrading peroxidase.

Lignin peroxidase is produced in various organisms and the coding sequences of the lignin peroxidase enzymes are available from GenBank via the Hypertext Transfer Protocol ://World Wide Web (dot) ncbi (dot) nlm (dot) nih (dot) gov/. For example, 25 lignin peroxidase from *Phanerochaete chrysosporium* (SEQ ID NO:1); *Mycobacterium tuberculosis* H37Rv [GenBank Accession No. NP\_216416.1 (SEQ ID NO:2 (polypeptide) and SEQ ID NO:3 (polynucleotide)]; *Mycobacterium bovis* AF2122/97 [GenBank Accession No. NP\_855586.1 (SEQ ID NO:4 (polypeptide) and SEQ ID NO:5 (polynucleotide)]; *Mycobacterium bovis* BCG str. Pasteur 1173P2 [GenBank 30 Accession No. YP\_978029.1 (SEQ ID NO:6 (polypeptide) and SEQ ID NO:7 (polynucleotide)]; *Mycobacterium bovis* BCG str. Tokyo 172 [GenBank Accession No. YP\_002644977.1 (SEQ ID NO:8 (polypeptide) and SEQ ID NO:9 (polynucleotide)];

*Mycobacterium tuberculosis* H37Ra [GenBank Accession No. YP\_001283231.1 (SEQ ID NO:10 (polypeptide) and SEQ ID NO:11 (polynucleotide)]; *Mycobacterium ulcerans* Agy99 [GenBank Accession No. YP\_908214.1 (SEQ ID NO:12 (polypeptide) and SEQ ID NO:13 (polynucleotide)]; *Mycobacterium marinum* M [GenBank Accession No. YP\_001848608.1 (SEQ ID NO:14 (polypeptide) and SEQ ID NO:15 (polynucleotide)]; *Mycobacterium tuberculosis* KZN 1435 [GenBank Accession No. YP\_003032055.1 (SEQ ID NO:16 (polypeptide) and SEQ ID NO:17 (polynucleotide)]; *Mycobacterium tuberculosis* F11 [GenBank Accession No. YP\_001287866.1 (SEQ ID NO:18 (polypeptide) and SEQ ID NO:19 (polynucleotide)]; *P. chrysosporium* ligninase (CKG4) [GenBank Accession No. M18743.1 (SEQ ID NO:20 (polypeptide) and SEQ ID NO:21 (polynucleotide)]; *Phanerochaete chrysosporium* (anamorph: *Sporotrichum pruiniosum*) [GenBank Accession No. M80213.1 (SEQ ID NO:22 (polypeptide) and SEQ ID NO:23 (polynucleotide)]; *Phanerochaete sordida* ylpB [GenBank Accession No. AB455007.1 (SEQ ID NO:24 (polypeptide) and SEQ ID NO:25 (polynucleotide)]; *Phanerochaete chrysosporium* (anamorph: *Sporotrichum pruiniosum*) [GenBank Accession No. M77508.1 (SEQ ID NO:26 (polypeptide) and SEQ ID NO:27 (polynucleotide)].

In some embodiments of any of the embodiments described herein, the lignin peroxidase is a lignin peroxidase of a white rot fungus, for example, a lignin peroxidase extracted from a white rot fungus or a recombinant protein derived from a white rot fungus lignin peroxidase.

Herein and in the art, the phrase “white rot fungus” refers to fungi capable of breaking down lignin in wood. As such a process often leaves the light-colored cellulose of the wood, such fungi typically cause the attacked wood to assume a lighter color.

Examples of white rot fungi producing a suitable lignin peroxidase include, without limitation, *Phanerochaete* (e.g., *Phanerochaete chrysosporium*, *Phanerochaete sordida*), *Trametes* (e.g., *Trametes versicolor*) and *Ganoderma*.

According to some embodiments of any of the embodiments of the invention, the lignin peroxidase enzyme used according to some embodiments of the invention is the lignin peroxidase H1 isoform, which exhibits melanin oxidation activities both *in vitro* and *in vivo* [WO 2004/052275].

45

In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in the composition is in a range of from 1.25 to 125 units/gram. In some embodiments, the concentration is in a range of from 4 to 40 units/gram. In some embodiments, the concentration is in a range of from 6 to 20 units/gram. In some embodiments, the concentration is about 12.5 units/gram.

In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in the composition is in a range of from 4 to 125 units/gram. In some embodiments, the concentration is in a range of from 6 to 125 units/gram. In some embodiments, the concentration is in a range of from 12.5 to 125 units/gram.

In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in the composition is in a range of from 1.25 to 40 units/gram. In some embodiments, the concentration is in a range of from 1.25 to 20 units/gram. In some embodiments, the concentration is in a range of from 1.25 to 12.5 units/gram.

In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in the composition is in a range of from 1.25 to 125 units/gram, and a concentration of the hydrogen peroxide-producing enzyme in the composition is in a range of from 0.125 to 1250 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 1.25 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 4 to 125 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 4 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 6 to 20 units/gram. In some embodiments, the lignin peroxidase concentration is about 12.5 units/gram. In some embodiments, the hydrogen peroxide-producing enzyme is glucose oxidase.

In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in the composition is in a range of from 1.25 to 125 units/gram, and a concentration of the hydrogen peroxide-producing enzyme in the composition is in a range of from 1.25 to 1250 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 1.25 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 4 to 125

units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 4 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 6 to 20 units/gram. In some embodiments, the lignin peroxidase concentration is about 12.5 units/gram. In some embodiments, the hydrogen peroxide-producing enzyme is glucose oxidase.

In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in the composition is in a range of from 1.25 to 125 units/gram, and a concentration of the hydrogen peroxide-producing enzyme in the composition is in a range of from 0.125 to 125 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 1.25 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 4 to 125 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 4 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 6 to 20 units/gram. In some embodiments, the lignin peroxidase concentration is about 12.5 units/gram. In some embodiments, the hydrogen peroxide-producing enzyme is glucose oxidase.

In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in the composition is in a range of from 1.25 to 125 units/gram, and a concentration of the hydrogen peroxide-producing enzyme in the composition is in a range of from 1.25 to 125 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 1.25 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 4 to 125 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 4 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 6 to 20 units/gram. In some embodiments, the lignin peroxidase concentration is about 12.5 units/gram. In some embodiments, the hydrogen peroxide-producing enzyme is glucose oxidase.

In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in the composition is in a range of from 1.25 to 125 units/gram, and a concentration of the hydrogen peroxide-producing enzyme in the composition is in a range of from 4 to 1250 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 1.25 to 40 units/gram. In some

embodiments, the lignin peroxidase concentration is in a range of from 4 to 125 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 4 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 6 to 20 units/gram. In some embodiments, the lignin peroxidase 5 concentration is about 12.5 units/gram. In some embodiments, the hydrogen peroxide-producing enzyme is glucose oxidase.

In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in the composition is in a range of from 1.25 to 125 units/gram, and a concentration of the hydrogen peroxide-producing enzyme in the 10 composition is in a range of from 0.125 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 1.25 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 4 to 125 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 4 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is 15 in a range of from 6 to 20 units/gram. In some embodiments, the lignin peroxidase concentration is about 12.5 units/gram. In some embodiments, the hydrogen peroxide-producing enzyme is glucose oxidase.

In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in the composition is in a range of from 1.25 to 125 units/gram, and a concentration of the hydrogen peroxide-producing enzyme in the 20 composition is in a range of from 4 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 1.25 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 4 to 125 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 4 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is 25 in a range of from 6 to 20 units/gram. In some embodiments, the lignin peroxidase concentration is about 12.5 units/gram. In some embodiments, the hydrogen peroxide-producing enzyme is glucose oxidase.

According to some embodiments of any of the embodiments described herein, 30 lignin peroxidase is used in combination with an oxidizing mediator. In some embodiments of any of the embodiments described herein, a composition comprising lignin peroxidase further comprises an oxidizing mediator (as defined herein).

Oxidizing mediators for lignin peroxidase include aromatic molecules, such as methoxylated aromatic compounds.

Examples of suitable oxidizing mediators include, without limitation, phenolic compounds such as veratryl alcohol (3,4-dimethoxybenzyl alcohol), veratrole (1,2-dimethoxybenzene) and 1,4-dimethoxybenzene. Veratryl alcohol is an exemplary oxidizing mediator.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition is in a range of from 0.03  $\mu$ mole/gram to 300  $\mu$ mole/gram. In some embodiments, the concentration is in a range of from 0.3 to 30  $\mu$ mole/gram. In some embodiments, the concentration is in a range of from 1 to 10  $\mu$ mole/gram. In some embodiments, the concentration is in a range of from 2 to 5  $\mu$ mole/gram. In some embodiments, the concentration is about 3  $\mu$ mole/gram.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator is in a range of from 0.3  $\mu$ mole/gram to 300  $\mu$ mole/gram. In some embodiments, the concentration is in a range of from 1 to 300  $\mu$ mole/gram. In some embodiments, the concentration is in a range of from 3 to 300  $\mu$ mole/gram.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator is in a range of from 0.03  $\mu$ mole/gram to 30  $\mu$ mole/gram. In some embodiments, the concentration is in a range of from 0.03 to 10  $\mu$ mole/gram. In some embodiments, the concentration is in a range of from 0.03 to 3  $\mu$ mole/gram.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition is in a range of from 0.03  $\mu$ mole/gram to 300  $\mu$ mole/gram, and a concentration of lignin peroxidase in the composition is in a range of from 1.25 to 125 units/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.03 to 30  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 300  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 30  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 1 to 10  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 5  $\mu$ mole/gram. In some

embodiments, the oxidizing mediator concentration is about 3  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing mediator is veratryl alcohol.

5 In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition is in a range of from 0.03  $\mu$ mole/gram to 300  $\mu$ mole/gram, and a concentration of lignin peroxidase in the composition is in a range of from 4 to 125 units/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.03 to 30  $\mu$ mole/gram. In some 10 embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 300  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 30  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 1 to 10  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 5  $\mu$ mole/gram. In some 15 embodiments, the oxidizing mediator concentration is about 3  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing mediator is veratryl alcohol.

20 In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition is in a range of from 0.03  $\mu$ mole/gram to 300  $\mu$ mole/gram, and a concentration of lignin peroxidase in the composition is in a range of from 1.25 to 40 units/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.03 to 30  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 300  $\mu$ mole/gram. In some 25 embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 30  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 1 to 10  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 5  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is about 3  $\mu$ mole/gram. In some 30 embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing mediator is veratryl alcohol.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition is in a range of from 0.03  $\mu$ mole/gram to 300  $\mu$ mole/gram, and a concentration of lignin peroxidase in the composition is in a range of from 4 to 40 units/gram. In some embodiments, the 5 oxidizing mediator concentration is in a range of from 0.03 to 30  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 300  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 30  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 1 to 10  $\mu$ mole/gram. In some embodiments, the 10 oxidizing mediator concentration is in a range of from 2 to 5  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is about 3  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing 15 mediator is veratryl alcohol.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition is in a range of from 0.03  $\mu$ mole/gram to 300  $\mu$ mole/gram, and a concentration of lignin peroxidase in the composition is in a range of from 6 to 125 units/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.03 to 30  $\mu$ mole/gram. In some 20 embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 300  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 30  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 1 to 10  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 5  $\mu$ mole/gram. In some 25 embodiments, the oxidizing mediator concentration is about 3  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing mediator is veratryl alcohol.

In some embodiments of any of the embodiments described herein, a 30 concentration of the oxidizing mediator in the composition is in a range of from 0.03  $\mu$ mole/gram to 300  $\mu$ mole/gram, and a concentration of lignin peroxidase in the composition is in a range of from 1.25 to 20 units/gram. In some embodiments, the

oxidizing mediator concentration is in a range of from 0.03 to 30  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 300  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 30  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 1 to 10  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 5  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is about 3  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing mediator is veratryl alcohol.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition is in a range of from 0.03  $\mu$ mole/gram to 300  $\mu$ mole/gram, and a concentration of lignin peroxidase in the composition is in a range of from 6 to 20 units/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.03 to 30  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 300  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 30  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 1 to 10  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 5  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is about 3  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing mediator is veratryl alcohol.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition is in a range of from 0.03  $\mu$ mole/gram to 300  $\mu$ mole/gram, and a concentration of lignin peroxidase in the composition is in a range of from 12.5 to 125 units/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.03 to 30  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 300  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 30  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 1 to 10  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 5  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is about 3  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing mediator is veratryl alcohol.

concentration is in a range of from 1 to 10  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 5  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is about 3  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl 5 alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing mediator is veratryl alcohol.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition is in a range of from 0.03  $\mu$ mole/gram to 300  $\mu$ mole/gram, and a concentration of lignin peroxidase in the 10 composition is in a range of from 1.25 to 12.5 units/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.03 to 30  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 300  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.3 to 30  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 1 to 10  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 5  $\mu$ mole/gram. In some 15 embodiments, the oxidizing mediator concentration is about 3  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing mediator is veratryl 20 alcohol.

Several approaches can be used to produce the lignin peroxidase enzyme utilized according to some embodiments of the present invention.

For example, lignin peroxidase isoenzyme H1 can be prepared from the fungus *Phanerochaete chrysosporium*. High levels of enzymatic activity of lignin peroxidase 25 can be produced from the above fungus when grown in a stirred tank reactor (STR) fermentor while being immobilized on polyurethane foam or in suspension [Dosoretz et al., *Appl Environ Microbiol* 1993, 59: 1919-26].

According to some embodiments of the invention the fermentor is connected to a cooling system to maintain a culturing temperature of 37 °C and is stirred at speed of 30 50-300 rpm (rounds per minute), more preferably, 100-200 rpm, most preferably at 160 rpm. In order to increase the yield of lignin peroxidase activity the fermentor is aerated at an aeration rate of 0.1-1 liter of air per liter of culture medium per minutes.

According to presently preferred configurations the fermentor is aerated at an aeration rate of 0.2 liter of air per liter of culture medium per minute.

The *Phanerochaete chrysosporium* is cultured under culturing conditions devoid of manganese ions and containing glycerol as a source of carbon. According to some 5 embodiments of the invention, the glycerol is provided at a concentration range of 3-20 grams per liter. According to some embodiments of the invention the glycerol is provided at a concentration of 6 grams per liter.

During the purification process of lignin peroxidase isoenzyme H1 from the above fungus the enzymatic activity of the purified protein is been further tested by a 10 change in absorbance at 310 nm that occurs due to the oxidation of veratryl alcohol to veratryl aldehyde.

As lignin peroxidase isoenzyme H1 can result from a post-translational dephosphorylation of isoenzyme H2 [Kuan and Tien, *J Biol Chem* 1989, 264:20350-20355], the lignin peroxidase used according to some embodiments of the invention can 15 be prepared by dephosphorylating the lignin peroxidase isoenzyme H2.

According to some embodiments of any of the embodiments of the invention, the lignin peroxidase can be recombinantly expressed using a nucleic acid construct designed to express the coding sequence of lignin peroxidase in a host cell. Non-limiting examples of such nucleic acid sequences are provided in SEQ ID NOs: 3, 5, 7, 20 9, 11, 13, 15, 17, 19, 21, 23, 25 and 27.

General procedures for expressing a protein such as lignin peroxidase are described in detail herein.

According to some embodiments of any of the embodiments of the invention, the lignin peroxidase is expressed in bacterial cells modified to express the lignin peroxidase as disclosed in US Patent No. 5,200,338. For example, a bacterial cell, such as, *E. coli*, can be transformed with an expression vector including the lignin peroxidase coding sequence positioned under the regulatory control of a strong constitutive promoter (e.g., SP6) (e.g., coding for a polypeptide having SEQ ID NO: 1). Following expression, the bacterial cells can be lysed and the lignin peroxidase can be collected 25 30 using chromatographic techniques (see, Billman-Jacobe [*Curr Opin Biotechnol* 1996, 7: 500-4]; Harris and Emtage [*Microbiol. Sci.* 1986, 3: 28-31], for further details).

The lignin peroxidase according to some embodiments of any of the embodiments of the invention can be extracted from mammalian cell lines such as HeLa cells. In this case the lignin peroxidase coding sequence is positioned under a strong mammalian promoter (e.g., CMV) in a suitable expression vector (e.g., pCDNA3.1, 5 Invitrogen Life Technologies, Frederick, MD, USA). Following transfection of HeLa cells with the expression vector, the lignin peroxidase expression product can be extracted from the cells or medium (e.g., by modifying the lignin peroxidase sequence to include a secretion signal) by conventional purification and chromatography techniques (see Cunha and Aires-Barros [*Mol. Biotechnol.* 2002, 20: 29-40] for further 10 details).

***Protein expression:***

A nucleic acid construct (also referred to herein as an "expression vector") used in some embodiments of the invention (e.g., for expressing a lignin peroxidase or hydrogen peroxide-producing enzyme described herein) includes additional sequences 15 which render this vector suitable for replication and integration in prokaryotes, eukaryotes, or preferably both (e.g., shuttle vectors). In addition, typical cloning vectors may also contain a transcription and translation initiation sequence, transcription and translation terminator and a polyadenylation signal. By way of example, such constructs will typically include a 5' LTR, a tRNA binding site, a 20 packaging signal, an origin of second-strand DNA synthesis, and a 3' LTR or a portion thereof.

The nucleic acid construct of some embodiments of the invention typically includes a signal sequence for secretion of the peptide from a host cell in which it is placed. Preferably the signal sequence for this purpose is a mammalian signal sequence 25 or the signal sequence of the polypeptide variants of some embodiments of the invention.

Eukaryotic promoters typically contain two types of recognition sequences, the TATA box and upstream promoter elements. The TATA box, located 25-30 base pairs upstream of the transcription initiation site, is thought to be involved in directing RNA 30 polymerase to begin RNA synthesis. The other upstream promoter elements determine the rate at which transcription is initiated.

Enhancer elements can stimulate transcription up to 1,000 fold from linked homologous or heterologous promoters. Enhancers are active when placed downstream or upstream from the transcription initiation site. Many enhancer elements derived from viruses have a broad host range and are active in a variety of tissues. For example, 5 the SV40 early gene enhancer is suitable for many cell types. Other enhancer/promoter combinations that are suitable for some embodiments of the invention include those derived from polyoma virus, human or murine cytomegalovirus (CMV), the long term repeat from various retroviruses such as murine leukemia virus, murine or Rous sarcoma virus and HIV. See, Enhancers and Eukaryotic Expression, Cold Spring 10 Harbor Press, Cold Spring Harbor, N.Y. 1983

In the construction of the expression vector, the promoter is preferably positioned approximately the same distance from the heterologous transcription start site as it is from the transcription start site in its natural setting. As is known in the art, 15 however, some variation in this distance can be accommodated without loss of promoter function.

Polyadenylation sequences can also be added to the expression vector in order to increase the efficiency of mRNA translation (e.g., lignin peroxidase mRNA translation). Two distinct sequence elements are required for accurate and efficient polyadenylation: 20 GU or U rich sequences located downstream from the polyadenylation site and a highly conserved sequence of six nucleotides, AAUAAA, located 11-30 nucleotides upstream. Termination and polyadenylation signals that are suitable for some embodiments of the invention include those derived from SV40.

In addition to the elements already described, the expression vector of some 25 embodiments of the invention may typically contain other specialized elements intended to increase the level of expression of cloned nucleic acids or to facilitate the identification of cells that carry the recombinant DNA. For example, a number of animal viruses contain DNA sequences that promote the extra chromosomal replication of the viral genome in permissive cell types. Plasmids bearing these viral replicons are 30 replicated episomally as long as the appropriate factors are provided by genes either carried on the plasmid or with the genome of the host cell.

The vector may or may not include a eukaryotic replicon. If a eukaryotic replicon is present, then the vector is amplifiable in eukaryotic cells using the appropriate selectable marker. If the vector does not comprise a eukaryotic replicon, no episomal amplification is possible. Instead, the recombinant DNA integrates into the 5 genome of the engineered cell, where the promoter directs expression of the desired nucleic acid.

The expression vector of some embodiments of the invention can further include additional polynucleotide sequences that allow, for example, the translation of several proteins from a single mRNA such as an internal ribosome entry site (IRES) and 10 sequences for genomic integration of the promoter-chimeric polypeptide.

As mentioned hereinabove, a variety of prokaryotic or eukaryotic cells can be used as host-expression systems to express the polypeptides of some embodiments of the invention. These include, but are not limited to, microorganisms, such as bacteria transformed with a recombinant bacteriophage DNA, plasmid DNA or cosmid DNA 15 expression vector containing the coding sequence; yeast transformed with recombinant yeast expression vectors containing the coding sequence; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors, such as Ti plasmid, containing the coding sequence. Mammalian expression 20 systems can also be used to express the polypeptides of some embodiments of the invention.

Examples for mammalian expression vectors include, but are not limited to, pcDNA3, pcDNA3.1(+/-), pGL3, pZeoSV2(+/-), pSecTag2, pDisplay, pEF/myc/cyto, pCMV/myc/cyto, pCR3.1, pSinRep5, DH26S, DHBB, pNMT1, pNMT41, pNMT81, 25 which are available from Invitrogen, pCI which is available from Promega, pMbac, pPbac, pBK-RSV and pBK-CMV which are available from Strategene, pTRES which is available from Clontech, and their derivatives.

Expression vectors containing regulatory elements from eukaryotic viruses such as retroviruses can be also used. SV40 vectors include pSVT7 and pMT2. Vectors 30 derived from bovine papilloma virus include pBV-1MTHA, and vectors derived from Epstein Bar virus include pHEBO, and p2O5. Other exemplary vectors include pMSG, pAV009/A<sup>+</sup>, pMTO10/A<sup>+</sup>, pMAMneo-5, baculovirus pDSVE, and any other vector

allowing expression of proteins under the direction of the SV-40 early promoter, SV-40 later promoter, metallothionein promoter, murine mammary tumor virus promoter, Rous sarcoma virus promoter, polyhedrin promoter, or other promoters shown effective for expression in eukaryotic cells.

5 Various methods can be used to introduce the expression vector of some embodiments of the invention into stem cells. Such methods are generally described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Springs Harbor Laboratory, New York (1989, 1992), in Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley and Sons, Baltimore, Md. (1989), Chang et al., *Somatic Gene Therapy*, CRC Press, Ann Arbor, Mich. (1995), Vega et al., *Gene Targeting*, CRC Press, Ann Arbor Mich. (1995), *Vectors: A Survey of Molecular Cloning Vectors and Their Uses*, Butterworths, Boston Mass. (1988) and Gilboa et al. [Biotechniques 4 (6): 10 504-512, 1986] and include, for example, stable or transient transfection, electroporation and infection with recombinant viral vectors. In addition, see U.S. Pat. 15 Nos. 5,464,764 and 5,487,992 for positive-negative selection methods.

Introduction of nucleic acids by viral infection offers several advantages over other methods such as electroporation, since higher transfection efficiency can be obtained due to the infectious nature of viruses.

Other than containing the necessary elements for the transcription and 20 translation of the inserted coding sequence, the expression construct of some embodiments of the invention can also include sequences engineered to enhance stability, production, purification, yield or toxicity of the expressed peptide. For example, the expression of a fusion protein or a cleavable fusion protein comprising the lignin peroxidase H1 or H2 protein of some embodiments of the invention and a 25 heterologous protein can be engineered. Such a fusion protein can be designed so that the fusion protein can be readily isolated by affinity chromatography; e.g., by immobilization on a column specific for the heterologous protein. Where a cleavage site is engineered between the lignin peroxidase H1 or H2 protein and the heterologous protein, the lignin peroxidase protein can be released from the chromatographic column 30 by treatment with an appropriate enzyme or agent that disrupts the cleavage site [e.g., see Booth et al. (1988) *Immunol. Lett.* 19:65-70; and Gardella et al., (1990) *J. Biol. Chem.* 265:15854-15859].

As mentioned hereinabove, a variety of prokaryotic or eukaryotic cells can be used as host-expression systems to express the polypeptides of some embodiments of the invention. These include, but are not limited to, microorganisms, such as bacteria transformed with a recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vector containing the coding sequence; yeast transformed with recombinant yeast expression vectors containing the coding sequence; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors, such as Ti plasmid, containing the coding sequence. Mammalian expression systems can also be used to express the polypeptides of some embodiments of the invention.

Examples of bacterial constructs include the pET series of *E. coli* expression vectors [Studier et al. (1990) *Methods in Enzymol.* 185:60-89].

In yeast, a number of vectors containing constitutive or inducible promoters can be used, as disclosed in U.S. Pat. Application No: 5,932,447. Alternatively, vectors can be used which promote integration of foreign DNA sequences into the yeast chromosome.

In cases where plant expression vectors are used, the expression of the coding sequence can be driven by a number of promoters. For example, viral promoters such as the 35S RNA and 19S RNA promoters of CaMV [Brisson et al. (1984) *Nature* 310:511-514], or the coat protein promoter to TMV [Takamatsu et al. (1987) *EMBO J.* 6:307-311] can be used. Alternatively, plant promoters such as the small subunit of RUBISCO [Coruzzi et al. (1984) *EMBO J.* 3:1671-1680 and Brogli et al., (1984) *Science* 224:838-843] or heat shock promoters, e.g., soybean hsp17.5-E or hsp17.3-B [Gurley et al. (1986) *Mol. Cell. Biol.* 6:559-565] can be used. These constructs can be introduced into plant cells using Ti plasmid, Ri plasmid, plant viral vectors, direct DNA transformation, microinjection, electroporation and other techniques well known to the skilled artisan. See, for example, Weissbach & Weissbach, 1988, *Methods for Plant Molecular Biology*, Academic Press, NY, Section VIII, pp 421-463.

Other expression systems such as insects and mammalian host cell systems which are well known in the art and are further described hereinbelow can also be used by some embodiments of the invention.

Recovery of the recombinant polypeptide is effected following an appropriate time in culture. The phrase "recovering the recombinant polypeptide" refers to collecting the whole fermentation medium containing the polypeptide and need not imply additional steps of separation or purification. Notwithstanding the above, 5 polypeptides of some embodiments of the invention can be purified using a variety of standard protein purification techniques, such as, but not limited to, affinity chromatography, ion exchange chromatography, filtration, electrophoresis, hydrophobic interaction chromatography, gel filtration chromatography, reverse phase chromatography, concanavalin A chromatography, chromatofocusing and differential 10 solubilization.

*A lignin peroxidase-containing kit:*

According to another aspect of some embodiments of the invention, there is provided a kit for lightening a topical body site such as skin and/or hair of a subject (e.g., in accordance with a method described herein). The kit comprises a lignin 15 peroxidase (e.g., as described herein), a hydrogen peroxide-producing enzyme (e.g., as described herein) and a substrate of the hydrogen peroxide-producing enzyme (e.g., as described herein). At least one of the lignin peroxidase, hydrogen peroxide-producing enzyme and substrate thereof forms a part of a composition which is packaged individually within the kit.

20 It is to be understood that a lignin peroxidase-containing kit described herein (e.g., in this section) may contain a lignin peroxidase according to any one of the embodiments described herein relating to lignin peroxidase, and may contain a hydrogen peroxide-producing system (e.g., enzyme and/or substrate) according to any one of the embodiments described herein relating to such a system and/or to a kit containing such a 25 system.

In some embodiments of any of the embodiments described herein, the hydrogen peroxide-producing enzyme forms a part of a first composition and the substrate forms a part of a second composition, wherein the first composition and second composition are packaged individually within the kit. That is, the enzyme and substrate thereof are 30 in separate compositions within the kit, and there is no contact between the enzyme and substrate while they are in the kit.

In some embodiments of any of the embodiments described herein, the second composition further comprises the lignin peroxidase (e.g., the lignin peroxidase is packaged within the same composition as the substrate of the hydrogen peroxide-producing enzyme). In such embodiments, the first composition is also referred to as an 5 “activator”, as it results in activation of lignin peroxidase when contacted with the lignin peroxidase-containing second composition.

In exemplary embodiments, the first composition further comprises the lignin peroxidase (e.g., the lignin peroxidase is packaged within the same composition as the hydrogen peroxide-producing enzyme). In such embodiments, the first composition is 10 also referred to as the “enzyme” formulation (e.g., “enzyme cream”), as it comprises the two active enzymes. The second composition is also referred to as an “activator”, as it results in activation of lignin peroxidase when contacted with the lignin peroxidase-containing first composition.

In some embodiments of any of the embodiments described herein, the lignin peroxidase forms a part of a third composition within the kit, wherein the third 15 composition is packaged individually within the kit.

In any of the embodiments described herein, each of the first composition described herein, the second composition described herein, and any other part of the hydrogen peroxide-producing composition described herein may be referred to herein 20 also as a formulation, which is packaged within the kit, whereby mixing these formulations results in formation of any of the hydrogen peroxide-producing composition, as described herein.

The first composition and second composition and optional third composition of the kit may be combined so as to prepare a cosmetic composition comprising a lignin peroxidase, a hydrogen peroxide-producing enzyme and a substrate thereof (e.g., as 25 described herein).

In some embodiments of any of the embodiments described herein, the hydrogen peroxide-producing enzyme and the substrate thereof are packaged together (e.g., as part of a single composition) in an air-tight package within the kit (e.g., an air-tight 30 container described herein). The lignin peroxidase may be packaged together with the hydrogen peroxide-producing enzyme and substrate thereof or packaged separately.

In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 2.5 to 250 units/gram. In some embodiments, the concentration is in a range of from 8 to 80 units/gram. In some embodiments, the 5 concentration is in a range of from 12 to 40 units/gram. In some embodiments, the concentration is about 25 units/gram.

In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 8 to 250 units/gram. In some embodiments, the 10 concentration is in a range of from 12 to 250 units/gram. In some embodiments, the concentration is in a range of from 25 to 250 units/gram.

In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 2.5 to 80 units/gram. In some embodiments, the 15 concentration is in a range of from 2.5 to 40 units/gram. In some embodiments, the concentration is in a range of from 2.5 to 25 units/gram.

According to some embodiments of any of the embodiments described herein, the kit further comprises an oxidizing mediator (e.g., as described herein).

In some embodiments of any of the embodiments described herein, the oxidizing 20 mediator is part of a composition in the kit which comprises the lignin peroxidase. In some embodiments, the lignin peroxidase and oxidizing mediator are both part of the first composition. In some embodiments, the lignin peroxidase and oxidizing mediator are both part of the second composition. In some embodiments, the lignin peroxidase and oxidizing mediator are both part of the third composition.

In some embodiments of any of the embodiments described herein, the oxidizing 25 mediator is packaged separately from the composition in the kit which comprises lignin peroxidase. In some embodiments, the lignin peroxidase is a part of the first composition or third composition described herein, and the oxidizing mediator is a part of the second composition. In some embodiments, the lignin peroxidase a part of the 30 second composition or third composition described herein and the oxidizing mediator is a part of the first composition.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition in the kit which comprises the oxidizing mediator is in a range of from 0.06  $\mu$ mole/gram to 600  $\mu$ mole/gram. In some embodiments, the concentration is in a range of from 0.6 to 60  $\mu$ mole/gram. In 5 some embodiments, the concentration is in a range of from 2 to 20  $\mu$ mole/gram. In some embodiments, the concentration is in a range of from 4 to 10  $\mu$ mole/gram. In some embodiments, the concentration is about 6  $\mu$ mole/gram.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition in the kit which comprises 10 the oxidizing mediator is in a range of from 0.6  $\mu$ mole/gram to 600  $\mu$ mole/gram. In some embodiments, the concentration is in a range of from 2 to 600  $\mu$ mole/gram. In some embodiments, the concentration is in a range of from 6 to 600  $\mu$ mole/gram.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition in the kit which comprises 15 the oxidizing mediator is in a range of from 0.06  $\mu$ mole/gram to 60  $\mu$ mole/gram. In some embodiments, the concentration is in a range of from 0.06 to 60  $\mu$ mole/gram. In some embodiments, the concentration is in a range of from 0.06 to 6  $\mu$ mole/gram.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition in the kit which comprises 20 the oxidizing mediator is in a range of from 0.06  $\mu$ mole/gram to 600  $\mu$ mole/gram, and a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 2.5 to 250 units/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.06 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 600  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 20  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 4 to 10  $\mu$ mole/gram. In some 25 embodiments, the oxidizing mediator concentration is about 6  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing 30

mediator is veratryl alcohol. In some embodiments, the oxidizing mediator and the lignin peroxidase are comprised by the same composition.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition in the kit which comprises the oxidizing mediator is in a range of from 0.06  $\mu$ mole/gram to 600  $\mu$ mole/gram, and a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 8 to 250 units/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.06 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 600  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 20  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 4 to 10  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is about 6  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing mediator is veratryl alcohol. In some embodiments, the oxidizing mediator and the lignin peroxidase are comprised by the same composition.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition in the kit which comprises the oxidizing mediator is in a range of from 0.06  $\mu$ mole/gram to 600  $\mu$ mole/gram, and a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 2.5 to 80 units/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.06 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 600  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 20  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 4 to 10  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is about 6  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing

mediator is veratryl alcohol. In some embodiments, the oxidizing mediator and the lignin peroxidase are comprised by the same composition.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition in the kit which comprises the oxidizing mediator is in a range of from 0.06  $\mu$ mole/gram to 600  $\mu$ mole/gram, and a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 8 to 80 units/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.06 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 600  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 20  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 4 to 10  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is about 6  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing mediator is veratryl alcohol. In some embodiments, the oxidizing mediator and the lignin peroxidase are comprised by the same composition.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition in the kit which comprises the oxidizing mediator is in a range of from 0.06  $\mu$ mole/gram to 600  $\mu$ mole/gram, and a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 12 to 250 units/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.06 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 600  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 20  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 4 to 10  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is about 6  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing

mediator is veratryl alcohol. In some embodiments, the oxidizing mediator and the lignin peroxidase are comprised by the same composition.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition in the kit which comprises the oxidizing mediator is in a range of from 0.06  $\mu$ mole/gram to 600  $\mu$ mole/gram, and a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 2.5 to 40 units/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.06 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 600  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 20  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 4 to 10  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is about 6  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing mediator is veratryl alcohol. In some embodiments, the oxidizing mediator and the lignin peroxidase are comprised by the same composition.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition in the kit which comprises the oxidizing mediator is in a range of from 0.06  $\mu$ mole/gram to 600  $\mu$ mole/gram, and a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 12 to 40 units/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.06 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 600  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 20  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 4 to 10  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is about 6  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing

mediator is veratryl alcohol. In some embodiments, the oxidizing mediator and the lignin peroxidase are comprised by the same composition.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition in the kit which comprises the oxidizing mediator is in a range of from 0.06  $\mu$ mole/gram to 600  $\mu$ mole/gram, and a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 25 to 250 units/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.06 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 600  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 20  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 4 to 10  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is about 6  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing mediator is veratryl alcohol. In some embodiments, the oxidizing mediator and the lignin peroxidase are comprised by the same composition.

In some embodiments of any of the embodiments described herein, a concentration of the oxidizing mediator in the composition in the kit which comprises the oxidizing mediator is in a range of from 0.06  $\mu$ mole/gram to 600  $\mu$ mole/gram, and a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 2.5 to 25 units/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.06 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 600  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 0.6 to 60  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 2 to 20  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is in a range of from 4 to 10  $\mu$ mole/gram. In some embodiments, the oxidizing mediator concentration is about 6  $\mu$ mole/gram. In some embodiments, the oxidizing mediator is selected from the group consisting of veratryl alcohol, veratrole and 1,4-dimethoxybenzene. In some embodiments, the oxidizing

mediator is veratryl alcohol. In some embodiments, the oxidizing mediator and the lignin peroxidase are comprised by the same composition.

In some embodiments of any of the embodiments described herein, a concentration of the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) in a composition in any of the kits described herein (e.g., a first composition described herein) is in a range of from 0.25 to 2500 units/gram. In some embodiments, the concentration is in a range of from 2.5 to 250 units/gram. In some embodiments, the concentration is in a range of from 8 to 80 units/gram. In some embodiments, the concentration is in a range of from 12 to 40 units/gram. In some embodiments, the concentration is about 25 units/gram.

In some embodiments of any of the embodiments described herein, a concentration of the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) in a composition in the kit (e.g., a first composition described herein) is in a range of from 2.5 to 2500 units/gram. In some embodiments, the concentration is in a range of from 8 to 2500 units/gram. In some embodiments, the concentration is in a range of from 25 to 2500 units/gram.

In some embodiments of any of the embodiments described herein, a concentration of the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) in a composition in the kit (e.g., a first composition described herein) is in a range of from 0.25 to 250 units/gram. In some embodiments, the concentration is in a range of from 0.25 to 80 units/gram. In some embodiments, the concentration is in a range of from 0.25 to 25 units/gram.

In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 2.5 to 250 units/gram, and a concentration of the hydrogen peroxide-producing enzyme in a composition in the kit (e.g., a first composition described herein) is in a range of from 0.25 to 2500 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 2.5 to 80 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 8 to 250 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 8 to 80 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 12 to 40 units/gram. In some embodiments, the

lignin peroxidase concentration is about 25 units/gram. In some embodiments, the hydrogen peroxide-producing enzyme and the lignin peroxidase are comprised by the same composition. In some embodiments, the hydrogen peroxide-producing enzyme is glucose oxidase.

5 In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 2.5 to 250 units/gram, and a concentration of the hydrogen peroxide-producing enzyme in a composition in the kit (e.g., a first composition described herein) is in a range of from 2.5 to 2500 units/gram. In some 10 embodiments, the lignin peroxidase concentration is in a range of from 2.5 to 80 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 8 to 250 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 8 to 80 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 12 to 40 units/gram. In some 15 embodiments, the lignin peroxidase concentration is about 25 units/gram. In some embodiments, the hydrogen peroxide-producing enzyme and the lignin peroxidase are comprised by the same composition. In some embodiments, the hydrogen peroxide-producing enzyme is glucose oxidase.

20 In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 2.5 to 250 units/gram, and a concentration of the hydrogen peroxide-producing enzyme in a composition in the kit (e.g., a first composition described herein) is in a range of from 0.25 to 250 units/gram. In some 25 embodiments, the lignin peroxidase concentration is in a range of from 2.5 to 80 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 8 to 250 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 8 to 80 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 12 to 40 units/gram. In some 30 embodiments, the lignin peroxidase concentration is about 25 units/gram. In some embodiments, the hydrogen peroxide-producing enzyme and the lignin peroxidase are comprised by the same composition. In some embodiments, the hydrogen peroxide-producing enzyme is glucose oxidase.

In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 2.5 to 250 units/gram, and a concentration of the hydrogen peroxide-producing enzyme in a composition in the kit (e.g., a first 5 composition described herein) is in a range of from 2.5 to 250 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 2.5 to 80 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 8 to 250 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 8 to 80 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 12 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is about 25 units/gram. In some embodiments, the hydrogen peroxide-producing enzyme and the lignin peroxidase are comprised by the same composition. In some embodiments, the hydrogen peroxide-producing enzyme is glucose oxidase.

15 In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 2.5 to 250 units/gram, and a concentration of the hydrogen peroxide-producing enzyme in a composition in the kit (e.g., a first composition described herein) is in a range of from 8 to 2500 units/gram. In some 20 embodiments, the lignin peroxidase concentration is in a range of from 2.5 to 80 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 8 to 250 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 8 to 80 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 12 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is about 25 units/gram. In some embodiments, the hydrogen peroxide-producing enzyme and the lignin peroxidase are comprised by the same composition. In some embodiments, the hydrogen peroxide-producing enzyme is glucose oxidase.

30 In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 2.5 to 250 units/gram, and a concentration of the hydrogen peroxide-producing enzyme in a composition in the kit (e.g., a first

composition described herein) is in a range of from 0.25 to 80 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 2.5 to 80 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 8 to 250 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 8 to 80 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 12 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is about 25 units/gram. In some embodiments, the hydrogen peroxide-producing enzyme and the lignin peroxidase are comprised by the same composition. In some embodiments, the hydrogen peroxide-producing enzyme is glucose oxidase.

In some embodiments of any of the embodiments described herein, a concentration of lignin peroxidase in a composition in the kit which comprises the lignin peroxidase is in a range of from 2.5 to 250 units/gram, and a concentration of the hydrogen peroxide-producing enzyme in a composition in the kit (e.g., a first composition described herein) is in a range of from 8 to 80 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 2.5 to 80 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 8 to 250 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 8 to 80 units/gram. In some embodiments, the lignin peroxidase concentration is in a range of from 12 to 40 units/gram. In some embodiments, the lignin peroxidase concentration is about 25 units/gram. In some embodiments, the hydrogen peroxide-producing enzyme and the lignin peroxidase are comprised by the same composition. In some embodiments, the hydrogen peroxide-producing enzyme is glucose oxidase.

In some embodiments of any of the embodiments described herein, a concentration of the substrate of the hydrogen peroxide-producing enzyme (e.g., glucose) in a composition in any of the kits described herein (e.g., a second composition described herein) is in a range of from 0.035  $\mu$ mole/gram to 350  $\mu$ mole/gram. In some embodiment, the concentration is in a range of from 0.35 to 35  $\mu$ mole/gram. In some embodiment, the concentration is in a range of from 1 to 12  $\mu$ mole/gram. In some embodiment, the concentration is in a range of from 2 to 7  $\mu$ mole/gram. In some embodiment, the concentration is about 3.5  $\mu$ mole/gram.

In some embodiments of any of the embodiments described herein, a concentration of the substrate of the hydrogen peroxide-producing enzyme (e.g., glucose) in a composition in the kit (e.g., a second composition described herein) is in a range of from 0.35  $\mu$ mole/gram to 350  $\mu$ mole/gram. In some embodiment, the 5 concentration is in a range of from 1 to 350  $\mu$ mole/gram. In some embodiment, the concentration is in a range of from 3.5 to 350  $\mu$ mole/gram.

In some embodiments of any of the embodiments described herein, a concentration of the substrate of the hydrogen peroxide-producing enzyme (e.g., glucose) in a composition in the kit (e.g., a second composition described herein) is in a 10 range of from 0.035  $\mu$ mole/gram to 35  $\mu$ mole/gram. In some embodiment, the concentration is in a range of from 0.035 to 12  $\mu$ mole/gram. In some embodiment, the concentration is in a range of from 0.035 to 3.5  $\mu$ mole/gram.

In some embodiments of any of the embodiments described herein, a concentration of the hydrogen peroxide-producing enzyme in a composition in any of 15 the kits described herein (e.g., a first composition described herein) is in a range of from 0.25 to 2500 units/gram, and a concentration of the substrate of the hydrogen peroxide-producing enzyme in a composition in the kit (e.g., a second composition described herein) is in a range of from 0.035  $\mu$ mole/gram to 350  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 0.25 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 2.5 to 2500 units/gram. In some embodiments, the enzyme concentration is in a range of from 2.5 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 8 to 80 units/gram. In some embodiments, the enzyme concentration is in a range of from 12 to 40 units/gram. In some embodiments, the enzyme concentration is about 25 20 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is glucose. In some embodiments, the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) and the substrate (e.g., glucose) are comprised by separate compositions in the kit.

In some embodiments of any of the embodiments described herein, a 30 concentration of the hydrogen peroxide-producing enzyme in a composition in any of the kits described herein (e.g., a first composition described herein) is in a range of from 0.25 to 2500 units/gram, and a concentration of the substrate of the hydrogen peroxide-

producing enzyme in a composition in the kit (e.g., a second composition described herein) is in a range of from 0.35  $\mu$ mole/gram to 350  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 0.25 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 2.5 to 2500 units/gram. In some embodiments, the enzyme concentration is in a range of from 2.5 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 8 to 80 units/gram. In some embodiments, the enzyme concentration is in a range of from 12 to 40 units/gram. In some embodiments, the enzyme concentration is about 25 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is glucose. In some embodiments, the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) and the substrate (e.g., glucose) are comprised by separate compositions in the kit.

In some embodiments of any of the embodiments described herein, a concentration of the hydrogen peroxide-producing enzyme in a composition in any of the kits described herein (e.g., a first composition described herein) is in a range of from 0.25 to 2500 units/gram, and a concentration of the substrate of the hydrogen peroxide-producing enzyme in a composition in the kit (e.g., a second composition described herein) is in a range of from 0.035  $\mu$ mole/gram to 35  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 0.25 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 2.5 to 2500 units/gram. In some embodiments, the enzyme concentration is in a range of from 2.5 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 8 to 80 units/gram. In some embodiments, the enzyme concentration is in a range of from 12 to 40 units/gram. In some embodiments, the enzyme concentration is about 25 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is glucose. In some embodiments, the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) and the substrate (e.g., glucose) are comprised by separate compositions in the kit.

In some embodiments of any of the embodiments described herein, a concentration of the hydrogen peroxide-producing enzyme in a composition in any of the kits described herein (e.g., a first composition described herein) is in a range of from 0.25 to 2500 units/gram, and a concentration of the substrate of the hydrogen peroxide-

producing enzyme in a composition in the kit (e.g., a second composition described herein) is in a range of from 0.35  $\mu$ mole/gram to 35  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 0.25 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 2.5 to 2500 units/gram. In 5 some embodiments, the enzyme concentration is in a range of from 2.5 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 8 to 80 units/gram. In some embodiments, the enzyme concentration is in a range of from 12 to 40 units/gram. In some embodiments, the enzyme concentration is about 25 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is 10 glucose. In some embodiments, the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) and the substrate (e.g., glucose) are comprised by separate compositions in the kit.

In some embodiments of any of the embodiments described herein, a concentration of the hydrogen peroxide-producing enzyme in a composition in any of 15 the kits described herein (e.g., a first composition described herein) is in a range of from 0.25 to 2500 units/gram, and a concentration of the substrate of the hydrogen peroxide-producing enzyme in a composition in the kit (e.g., a second composition described herein) is in a range of from 1  $\mu$ mole/gram to 350  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 0.25 to 250 units/gram. In some 20 embodiments, the enzyme concentration is in a range of from 2.5 to 2500 units/gram. In some embodiments, the enzyme concentration is in a range of from 2.5 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 8 to 80 units/gram. In some embodiments, the enzyme concentration is in a range of from 12 to 40 units/gram. In some embodiments, the enzyme concentration is about 25 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is 25 glucose. In some embodiments, the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) and the substrate (e.g., glucose) are comprised by separate compositions in the kit.

In some embodiments of any of the embodiments described herein, a 30 concentration of the hydrogen peroxide-producing enzyme in a composition in any of the kits described herein (e.g., a first composition described herein) is in a range of from 0.25 to 2500 units/gram, and a concentration of the substrate of the hydrogen peroxide-

producing enzyme in a composition in the kit (e.g., a second composition described herein) is in a range of from 0.035  $\mu$ mole/gram to 12  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 0.25 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 2.5 to 2500 units/gram. In some embodiments, the enzyme concentration is in a range of from 2.5 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 8 to 80 units/gram. In some embodiments, the enzyme concentration is in a range of from 12 to 40 units/gram. In some embodiments, the enzyme concentration is about 25 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is glucose. In some embodiments, the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) and the substrate (e.g., glucose) are comprised by separate compositions in the kit.

In some embodiments of any of the embodiments described herein, a concentration of the hydrogen peroxide-producing enzyme in a composition in any of the kits described herein (e.g., a first composition described herein) is in a range of from 0.25 to 2500 units/gram, and a concentration of the substrate of the hydrogen peroxide-producing enzyme in a composition in the kit (e.g., a second composition described herein) is in a range of from 1  $\mu$ mole/gram to 12  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 0.25 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 2.5 to 2500 units/gram. In some embodiments, the enzyme concentration is in a range of from 2.5 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 8 to 80 units/gram. In some embodiments, the enzyme concentration is in a range of from 12 to 40 units/gram. In some embodiments, the enzyme concentration is about 25 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is glucose. In some embodiments, the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) and the substrate (e.g., glucose) are comprised by separate compositions in the kit.

In some embodiments of any of the embodiments described herein, a concentration of the hydrogen peroxide-producing enzyme in a composition in any of the kits described herein (e.g., a first composition described herein) is in a range of from 0.25 to 2500 units/gram, and a concentration of the substrate of the hydrogen peroxide-

producing enzyme in a composition in the kit (e.g., a second composition described herein) is in a range of from 3.5  $\mu$ mole/gram to 350  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 0.25 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 2.5 to 2500 units/gram. In 5 some embodiments, the enzyme concentration is in a range of from 2.5 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 8 to 80 units/gram. In some embodiments, the enzyme concentration is in a range of from 12 to 40 units/gram. In some embodiments, the enzyme concentration is about 25 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is 10 glucose. In some embodiments, the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) and the substrate (e.g., glucose) are comprised by separate compositions in the kit.

In some embodiments of any of the embodiments described herein, a concentration of the hydrogen peroxide-producing enzyme in a composition in any of 15 the kits described herein (e.g., a first composition described herein) is in a range of from 0.25 to 2500 units/gram, and a concentration of the substrate of the hydrogen peroxide-producing enzyme in a composition in the kit (e.g., a second composition described herein) is in a range of from 0.035  $\mu$ mole/gram to 3.5  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 0.25 to 250 units/gram. In some 20 embodiments, the enzyme concentration is in a range of from 2.5 to 2500 units/gram. In some embodiments, the enzyme concentration is in a range of from 2.5 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 8 to 80 units/gram. In some embodiments, the enzyme concentration is in a range of from 12 to 40 units/gram. In some embodiments, the enzyme concentration is about 25 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is 25 glucose. In some embodiments, the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) and the substrate (e.g., glucose) are comprised by separate compositions in the kit.

In some embodiments of any of the embodiments described herein, a 30 concentration of the hydrogen peroxide-producing enzyme in a composition in any of the kits described herein (e.g., a first composition described herein) is in a range of from 0.25 to 2500 units/gram, and a concentration of the substrate of the hydrogen peroxide-

producing enzyme in a composition in the kit (e.g., a second composition described herein) is in a range of from 2  $\mu$ mole/gram to 7  $\mu$ mole/gram. In some embodiments, the enzyme concentration is in a range of from 0.25 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 2.5 to 2500 units/gram. In 5 some embodiments, the enzyme concentration is in a range of from 2.5 to 250 units/gram. In some embodiments, the enzyme concentration is in a range of from 8 to 80 units/gram. In some embodiments, the enzyme concentration is in a range of from 12 to 40 units/gram. In some embodiments, the enzyme concentration is about 25 units/gram. In some embodiments, the enzyme is glucose oxidase and the substrate is 10 glucose. In some embodiments, the hydrogen peroxide-producing enzyme (e.g., glucose oxidase) and the substrate (e.g., glucose) are comprised by separate compositions in the kit.

In some embodiments of any of the embodiments described herein, the first composition of the kit comprises glucose oxidase at a concentration in a range of from 15 2.5 to 250 units/gram (e.g., a glucose oxidase concentration described herein), and the second composition of the kit comprises D-glucose at a concentration in a range of from 0.35  $\mu$ mole/gram to 35  $\mu$ mole/gram (e.g., a glucose concentration described herein).

In some embodiments of any of the embodiments described herein, the first composition of the kit comprises glucose oxidase at a concentration in a range of from 20 2.5 to 250 units/gram (e.g., a glucose oxidase concentration described herein), and lignin peroxidase at a concentration of 2.5 to 250 units/gram (e.g., a lignin peroxidase concentration described herein); and the second composition of the kit comprises D-glucose at a concentration in a range of from 0.35  $\mu$ mole/gram to 35  $\mu$ mole/gram (e.g., a glucose concentration described herein).

25 The kits described herein may include additional compositions (e.g., cosmetic compositions), besides the compositions comprising one or more active ingredients as described herein (e.g., an agent which exhibits an activity in a presence of hydrogen peroxide, a lignin peroxidase, a hydrogen peroxide-producing enzyme, a substrate of hydrogen peroxide-producing enzyme and/or an oxidizing mediator).

30 In some embodiments of any of the embodiments described herein, the kit further comprises a wash composition (e.g., as described herein), such as a facial wash, for cleaning a topical body site prior to contact with a composition described herein.

In some embodiments of any of the embodiments described herein, the kit includes appropriate instructions for use and labels indicating approval by a regulatory agency for the indicated use, as described herein.

***Formulation of compositions:***

5 According to some embodiments of any of the embodiments and any of the aspects of embodiments of the invention described herein, the active ingredient(s) described herein (e.g., an agent which exhibits an activity in a presence of hydrogen peroxide, a lignin peroxidase, a hydrogen peroxide-producing enzyme, a substrate of hydrogen peroxide-producing enzyme and/or an oxidizing mediator) may be used in a  
10 composition comprising the active ingredient(s) *per se* or in combination with additional chemical components such as physiologically suitable carriers and excipients. Compositions include a cosmetic or pharmaceutical composition described herein (which exhibit a cosmetic or pharmaceutical activity as described herein) and compositions within a kit (e.g., a first composition, a second composition, and/or a third  
15 composition, as described herein) which may be used to prepare a cosmetic or pharmaceutical composition described herein (although they may be inactive by themselves).

The purpose of a cosmetic or pharmaceutical composition comprising carriers and excipients is to facilitate administration of the active ingredient to an organism. In  
20 embodiments wherein an activity exhibited by the composition is cosmetic, as defined herein, the composition is also referred to herein as a “cosmetic composition”. In embodiments wherein an activity exhibited by the composition is pharmaceutical, the composition is also referred to herein as a “pharmaceutical composition”.

Hereinthroughout, the term “pharmaceutical” means that the composition or kit  
25 are for exhibiting a therapeutic effect when applied on a subject’s bodily site as described herein.

Hereinafter, the phrase “suitable carrier” refers to a carrier or a diluent that does not cause significant irritation to an organism and does not abrogate the biological activity and properties of the active ingredient(s). In embodiments wherein an activity  
30 exhibited by the active ingredient(s) is cosmetic, such a carrier is also referred to herein as a “cosmetically or cosmeceutically acceptable carrier”. In embodiments wherein an

activity exhibited by the active ingredient(s) is pharmaceutical, such a carrier is also referred to herein as a “pharmaceutically acceptable carrier”.

Herein the term “excipient” refers to an inert substance added to a composition to further facilitate administration of an active ingredient(s). Examples, without limitation, of excipients include calcium carbonate, calcium phosphate, various sugars and types of starch, cellulose derivatives, gelatin, vegetable oils and polyethylene glycols.

Compositions of the present invention may be manufactured by processes well known in the art, e.g., by means of conventional mixing, dissolving, granulating, 10 dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

Compositions for use in accordance with some embodiments of the invention thus may be formulated in conventional manner using one or more suitable carriers comprising excipients and auxiliaries, which facilitate processing of the active 15 ingredient(s) into preparations. Proper formulation is dependent upon the administration approach chosen.

Alternatively, the active ingredient(s) may be in a powder form for constitution with a suitable vehicle, e.g., sterile, pyrogen-free water based solution, before use.

Compositions according to some embodiments of any of the embodiments and 20 any of the aspects described herein may be formulated in any of a variety of forms utilized by the cosmetic industry (e.g., for topical use), such as, for example, a mask, a serum, a lotion, a cream, a shampoo, a gel, and/or a toner. As will be apparent to the skilled person, suitability of a formulations will depend on the type of surface to be contacted with the active ingredient(s) – thus, for example, a shampoo formulation is 25 suitable for contacting hair, a facial mask is suitable for contacting facial skin, and so forth.

Methods for preparing compositions having properties such as described herein are well known to those skilled in the art, and are described in detail in Remington's Pharmaceutical Sciences, 1990 (supra); and Pharmaceutical Dosage Forms and Drug 30 Delivery Systems, 6th ed., Williams & Wilkins (1995).

According to some embodiments of any of the embodiments of the invention, the cosmetic composition is formulated viscous enough to remain on the treated area, does not readily evaporate, and/or is not easily removed by rinsing with water.

As used herein the term “serum” refers to an aqueous solution which is 5 formulated for topical application.

According to some embodiments of any of the embodiments of the invention, the serum is a thickened aqueous cosmetic solution. According to some embodiments of the invention, the serum is a clear to slight hazy aqueous solution.

According to some embodiments of any of the embodiments of the invention, 10 the serum is formulated to remain on a topical body site and does not need to be removed by water, soap and/or cleansers.

According to some embodiments of any of the embodiments of the invention, upon topical application of the serum (e.g., to skin), the serum is absorbed rapidly, leaving a cosmetically elegant silky smooth feel.

15 According to some embodiments of any of the embodiments of the invention, the serum is incorporated within a mask (a cosmetic mask).

In some embodiments of any of the embodiments described herein, the serum comprises ingredients as presented in Table 2:

20

**Table 2: Serum composition according to some embodiments**

Ingredient	Weight percentage range
Water	75 – 100 %
Active ingredient(s)	As described herein
Glycerin	5 – 10 %
Polyacrylate-13	0.1 – 1 %
Dimethicone	0.1 – 1 %
Dicaprylyl carbonate	0.1 – 1 %
Cetyl alcohol	0.1 – 1 %
Stearyl alcohol	0.1 – 1 %
Phenoxyethanol	0.1 – 1 %
Portulaca oleracea	0 – 1 %
Polyisobutene	0.1 – 1 %

Ingredient	Weight percentage range
Butylene glycol	0 – 1 %
Pullulan	0 – 0.1 %
Polysorbate 20	0 – 0.1 %
PEG-4 Laurate	0 – 0.1 %
PEG-4 Dilaurate	0 – 0.1 %
PEG-4	0 – 0.1 %
Iodopropynyl butylcarbamate	0 – 0.1 %

In some embodiments of any of the embodiments described herein, a composition as described herein (e.g., serum, lotion, cream) comprises a 5 dermatologically acceptable emollient. Such compositions preferably contain from about 2% to about 50% of the emollient. As used herein, “emollient” refers to a material useful for the prevention or relief of dryness, as well as for the protection of skin. A wide variety of suitable emollients are known and may be used herein. See, e.g., Sagarin, Cosmetics, Science and Technology, 2nd Edition, Vol. 1, pp. 3243 (1972), 10 which contains numerous examples of materials suitable as an emollient. A preferred emollient is glycerin. Glycerin is preferably used in an amount of from or about 0.001 to or about 20 %, more preferably from or about 0.01 to or about 10 %, most preferably from or about 0.1 to or about 5 %, e.g., 3 %.

15 Lotions and creams according to the present invention generally comprise a solution carrier system and one or more emollients. Lotions typically comprise from about 1 % to about 20 %, preferably from about 5 % to about 10 % of emollient; from about 50 % to about 90 %, preferably from about 60 % to about 80 % water; and a pharmaceutically effective amount of an agent described herein. A cream typically 20 comprises from about 5 % to about 50 %, preferably from about 10 % to about 20 % of emollient; from about 45 % to about 85 %, preferably from about 50 % to about 75 % water; and an effective amount of active ingredient(s) described herein.

In some embodiments of any of the embodiments described herein, the cream or lotion comprises ingredients as presented in Table 3:

**Table 3: Cream or lotion composition according to some embodiments**

Ingredient	Weight percentage range
Water	50 – 75 %
Active ingredient(s)	As described herein
Glycerin	5 – 10 %
<i>Butyrospermum Parkii</i> butter	0 – 5 %
Dimethicone	1 – 5 %
Dicaprylyl carbonate	1 – 5 %
Trehalose	0 – 5 %
Caprylic capric triglyceride	0 – 1 %
Cetyl alcohol	0.1 – 1 %
Jojoba ester	0 – 1 %
Stearyl alcohol	0.1 – 1 %
Polyacrylate-13	0.1 – 1 %
Cetyl palmitate	0 – 1 %
Myristyl myristate	0 – 1 %
Phenoxyethanol	0.1 – 1 %
Polyisobutene	0.1 – 1 %
Pullulan	0 – 1 %
Polysorbate 20	0 – 0.1 %
PEG- 4 laurate	0 – 0.1 %

Ingredient	Weight percentage range
PEG-4 dilaurate	0 – 0.1 %
PEG-4	0 – 0.1 %
Iodopropynyl butylcarbamate	0 – 0.1 %

**Table 3 (Cont.)**

In exemplary embodiments, the cream or lotion comprises ingredients as presented in Table 4:

5

**Table 4: Cream or lotion composition according to exemplary embodiments**

Ingredient	Weight percentage range
Water	75 – 100 %
Active ingredient(s)	As described herein
Glycerin	5 – 10 %
Propylene glycol	1 – 5 %
Dicaprylyl carbonate	0.1 – 1 %
Dimethicone	0.1 – 1 %
Polyacrylate-13	0.1 – 1 %
Cetyl alcohol	0.1 – 1 %
Paraffinum liquidum	0.1 – 1 %
Stearyl alcohol	0.1 – 1 %
Phenoxyethanol	0.1 – 1 %

Ingredient	Weight percentage range
Polyisobutene	0.1 – 1 %
Polysorbate 20	0 – 0.1 %
PEG- 4 laurate	0 – 0.1 %
PEG-4 dilaurate	0 – 0.1 %
PEG-4	0 – 0.1 %
Iodopropynyl butylcarbamate	0 – 0.1 %
Chlorphenesin	0 – 1 %
<i>Caesalpinia spinosa</i> gum	0 – 1 %
Citric acid	0 – 0.1 %

**Table 4 (Cont.)**

As used herein the term “mask” refers to a cosmetic mask which comprises the active ingredient(s) as described herein and which is topically applied in order to 5 maintain contact of the active ingredient(s) with a topical region, for example, in order to lighten the topical region.

Various types of cosmetic masks are known in the cosmetic industry. These include, but are not limited to a mask having a solid support, a rinse-off mask, a peeled-off mask, or a mask which hardens or polymerizes after a set period of time.

10 As used herein the phrase “rinse-off mask” refers to a cosmetic composition formulated in order to be topically applied in a relatively thick layer and left for a certain period of time, generally from a few minutes to several tens of minutes, and in order to be, at the end of this period of time, rinsed off with water or also simply wiped off.

15 Suitable masks and techniques for preparing masks with an active ingredient(s) such as described herein are described in International Patent Application Publication WO 2012/153336.

In some embodiments of any of the embodiments described herein, the form of the composition is determined, at least in part, by selecting a suitable carrier.

In some embodiments of any of the embodiments described herein, the carrier comprises an emulsion. Such carriers include, without limitation, oil-in-water, water-in-oil, water-in-oil-in-water, and oil-in-water-in-silicone emulsions, a cream, an ointment, an aqueous solution, a lotion or an aerosol. As will be understood by the skilled artisan, a given component will distribute primarily into either the water or oil/silicone phase, depending on the water solubility/dispersibility of the component in the composition.

Emulsions according to some embodiments of any of the embodiments of the invention generally contain an effective amount of active ingredient(s) disclosed herein and oil. The active ingredient(s) (e.g., an agent which exhibits an activity in a presence of hydrogen peroxide, a lignin peroxidase, a hydrogen peroxide-producing enzyme, a substrate of hydrogen peroxide-producing enzyme and/or an oxidizing mediator) and oils may be derived from animals, plants, or petroleum and may be natural or synthetic (i.e., man-made). Preferred emulsions also contain a humectant, such as glycerin. Emulsions will preferably further contain from about 1 % to about 10 %, more preferably from about 2 % to about 5 %, of an emulsifier, based on the weight of the carrier. Emulsifiers may be nonionic, anionic or cationic. Suitable emulsifiers are described in, for example, U.S. Pat. No. 3,755,560, issued to Dickert, et al. Aug. 28, 1973; U.S. Pat. No. 4,421,769, issued to Dixon, et al., Dec. 20, 1983; and McCutcheon's Detergents and Emulsifiers, North American Edition, pages 317-324 (1986).

The emulsion may also contain an anti-foaming agent to minimize foaming upon application to the keratinous tissue. Anti-foaming agents include high molecular weight silicones and other materials well known in the art for such use.

Suitable emulsions may have a wide range of viscosities, depending on the desired product form. Exemplary low viscosity emulsions, which are preferred, have a viscosity of about 50 centistokes or less, more preferably about 10 centistokes or less, most preferably about 5 centistokes or less. The emulsion may also contain an anti-foaming agent to minimize foaming upon topical application. Anti-foaming agents include high molecular weight silicones and other materials well known in the art for such use.

In some embodiments of any of the embodiments described herein, the emulsion is an oil-in-water emulsion, having a continuous aqueous phase and a hydrophobic, water-insoluble phase (“oil phase”) dispersed therein. Examples of suitable carriers comprising oil-in-water emulsions are described in U.S. Pat. Nos. 5,073,371 and 5,073,372.

In some embodiments of any of the embodiments described herein, the oil-in-water emulsion comprises a structuring agent to assist in the formation of a liquid crystalline gel network structure. Without being limited by theory, it is believed that the structuring agent assists in providing rheological characteristics to the composition which contribute to the stability of the composition. The structuring agent may also function as an emulsifier or surfactant. Preferred compositions of this invention comprise from about 0.5 % to about 20 %, more preferably from about 1 % to about 10 %, most preferably from about 1 % to about 5 %, by weight of the composition, of a structuring agent. The preferred structuring agents of the present invention are selected from the group consisting of stearic acid, palmitic acid, stearyl alcohol, cetyl alcohol, behenyl alcohol, the polyethylene glycol ether of stearyl alcohol having an average of about 1 to about 21 ethylene oxide units, the polyethylene glycol ether of cetyl alcohol having an average of about 1 to about 5 ethylene oxide units, and mixtures thereof.

A wide variety of anionic surfactants are also useful herein, as described, for example, in U.S. Pat. No. 3,929,678. In addition, amphoteric and zwitterionic surfactants are also useful herein.

In some embodiments of any of the embodiments described herein, the oil-in-water emulsion comprises from about 25 % to about 98 %, preferably from about 65 % to about 95 %, more preferably from about 70 % to about 90 % water by weight of the topical carrier.

In some embodiments of any of the embodiments described herein, the emulsion is a water-in-silicone emulsion. Water-in-silicone emulsions contain a continuous silicone phase and a dispersed aqueous phase. Preferred water-in-silicone emulsions of the present invention comprise from about 1 % to about 60 %, preferably from about 5% to about 40%, more preferably from about 10 % to about 20 %, by weight of a continuous silicone phase. The continuous silicone phase exists as an external phase

that contains or surrounds the discontinuous aqueous phase described hereinafter.

The continuous silicone phase may contain a polyorganosiloxane oil.

The continuous silicone phase may contain one or more non-silicone oils. Concentrations of non-silicone oils in the continuous silicone phase are preferably 5 minimized or avoided altogether so as to further enhance oxidative stability of the active ingredient(s) in the compositions. Suitable non-silicone oils have a melting point of about 25 °C or less under about one atmosphere of pressure. Examples of non-silicone oils suitable for use in the continuous silicone phase are those well known in the chemical arts in topical personal care products in the form of water-in-oil emulsions, 10 e.g., mineral oil, vegetable oils, synthetic oils, semisynthetic oils, etc.

Useful topical compositions according to some embodiments of any of the embodiments of the present invention comprise from about 30 % to about 90 %, more preferably from about 50 % to about 85 %, and most preferably from about 70 % to about 80 % of a dispersed aqueous phase. The term “dispersed phase” is well-known to 15 one skilled in the art it implies that the phase exists as small particles or droplets that are suspended in and surrounded by a continuous phase. The dispersed phase is also known as the internal or discontinuous phase. The dispersed aqueous phase is a dispersion of small aqueous particles or droplets suspended in and surrounded by the continuous oil and/or silicone phase (e.g., as described herein). The aqueous phase can be water, or a 20 combination of water and one or more water soluble or dispersible ingredients. Non-limiting examples of such optional ingredients include thickeners, acids, bases, salts, chelants, gums, water-soluble or dispersible alcohols and polyols, buffers, preservatives, sunscreening agents, colorings, and the like.

The topical compositions of some embodiments of the invention typically 25 comprise from about 25 % to about 90 %, preferably from about 40 % to about 80 %, more preferably from about 60 % to about 80 %, water in the dispersed aqueous phase by weight of the composition.

The emulsions of some embodiments of the invention comprise an emulsifier. In a preferred embodiment, the composition contains from about 0.1 % to about 10 % 30 emulsifier, more preferably from about 0.5 % to about 7.5 %, most preferably from about 1 % to about 5 %, emulsifier by weight of the composition. The emulsifier helps disperse and suspend the aqueous phase within the oil and/or silicone phase. Known or

conventional emulsifying agents can be used in the composition, provided that the selected emulsifying agent is chemically and physically compatible with essential components of the composition, and provides the desired dispersion characteristics. Suitable emulsifiers include emulsifiers known by those skilled in the art for use in 5 topical personal care products.

Suitable emulsifiers include any of a wide variety of known cationic, anionic, zwitterionic, and amphoteric surfactants. See, McCutcheon's. Detergents and Emulsifiers, North American Edition (1986), published by Allured Publishing Corporation; U.S. Pat. No. 5,011,681 to Ciotti et al., issued Apr. 30, 1991; U.S. Pat. No. 10 4,421,769 to Dixon et al. issued to Dec. 20, 1983; and U.S. Pat. No. 3,755,560. The exact surfactant chosen depends upon the pH of the composition and the other components present. Preferred are cationic surfactants, especially dialkyl quaternary ammonium compounds, examples of which are described in U.S. Pat. No. 5,151,209 to McCall et al. issued to Sep. 29, 1992; U.S. Pat. No. 5,151,210 to Steuri et al., issued to 15 Sep. 29, 1992; U.S. Pat. Nos. 5,120,532; U.S. Pat. No. 4,387,090; U.S. Pat. No. 3,155,591; U.S. Pat. No. 3,929,678; U.S. Pat. No. 3,959,461; McCutcheon's, Detergents & Emulsifiers (North American edition 1979) M.C. Publishing Co.; and Schwartz, et al., Surface Active Agents, Their chemistry and Technology, New York: Interscience Publishers, 1949.

20 Alternatively, other useful cationic emulsifiers include amino-amides. Nonlimiting examples of these cationic emulsifiers include stearamidopropyl PG-dimonium chloride phosphate, behenamidopropyl PG dimonium chloride, stearamidopropyl ethyldimonium ethosulfate, stearamidopropyl dimethyl (myristyl acetate) ammonium chloride, stearamidopropyl dimethyl cetearyl ammonium tosylate, 25 stearamidopropyl dirnethyl ammonium chloride, stearamidopropyl dimethyl ammonium lactate, and mixtures thereof.

Determination of an effective amount of active ingredient(s) is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

30 For any preparation used in the methods of the invention, the therapeutically effective amount or dose can be estimated initially from *in vitro* assays. In addition, a dose can be formulated in *ex vivo* systems (e.g., *ex vivo* skin, hair and/or nail) or in

animal models to achieve a desired concentration or titer. Such information can be used to more accurately determine useful doses in humans.

In order to enhance the percutaneous absorption of the active ingredients (e.g., lignin peroxidase, and/or oxidizing mediator), one or more of a number of agents can be added to the composition including, but not limited to, dimethylsulfoxide, dimethylacetamide, dimethylformamide, surfactants, azone, alcohol, acetone, propylene glycol, polyethylene glycol, and butylene glycol.

The compositions described herein may also include additional components which are added, for example, in order to enrich the compositions with fragrance and nutrition factors (e.g., skin or hair nutrition factors).

Such components are selected suitable for topical use on a human without inducing toxicity, incompatibility, instability, allergic response, and the like within the scope of sound medical judgment. In addition, such optional components are useful provided that they do not unacceptably alter the benefits of the active compounds of the invention.

The CTFA Cosmetic Ingredient Handbook, Second Edition (1992) describes a wide variety of non-limiting cosmetic ingredients commonly used in the skin care industry, which are suitable for use in the compositions of the present invention. Examples of these ingredient classes include: abrasives, absorbents, aesthetic components such as fragrances, pigments, colorings/colorants, essential oils, skin sensates, astringents, etc. (e.g., clove oil, menthol, camphor, eucalyptus oil, eugenol, menthol lactate, witch hazel distillate), anti-acne agents, anti-caking agents, antifoaming agents, antimicrobial agents (e.g., iodopropyl butylcarbamate), antioxidants, binders, biological additives, buffering agents, bulking agents, chelating agents, chemical additives, colorants, cosmetic astringents, cosmetic biocides, denaturants, drug astringents, external analgesics, film formers or materials, e.g., polymers, for aiding the film-forming properties and substantivity of the composition (e.g., copolymer of eicosene and vinyl pyrrolidone), opacifying agents, pH adjusters, propellants, reducing agents, sequestrants, skin-conditioning agents (e.g., humectants, including miscellaneous and occlusive), skin soothing and/or healing agents (e.g., panthenol and derivatives (e.g., ethyl panthenol), aloe vera, pantothenic acid and its derivatives,

allantoin, bisabolol, and dipotassium glycyffhizinate), skin treating agents, thickeners, and vitamins and derivatives thereof.

As compositions according to some embodiments described herein are for utilization *in vivo*, the composition is preferably of high purity and substantially free of 5 potentially harmful contaminants, e.g., at least National Food (NF) grade, generally at least analytical grade, and preferably at least pharmaceutical grade. To the extent that a given compound must be synthesized prior to use, such synthesis or subsequent purification shall preferably result in a product that is substantially free of any potentially contaminating toxic agents that may have been used during the synthesis or 10 purification procedures.

As used herein the term "about" refers to  $\pm 10\%$ .

The terms "comprises", "comprising", "includes", "including", "having" and their conjugates mean "including but not limited to".

The term "consisting of" means "including and limited to".

15 The term "consisting essentially of" means that the composition, method or structure may include additional ingredients, steps and/or parts, but only if the additional ingredients, steps and/or parts do not materially alter the basic and novel characteristics of the claimed composition, method or structure.

As used herein, the singular form "a", "an" and "the" include plural references 20 unless the context clearly dictates otherwise. For example, the term "a compound" or "at least one compound" may include a plurality of compounds, including mixtures thereof.

Throughout this application, various embodiments of this invention may be 25 presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically disclosed all the possible subranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed subranges such as 30 from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 3, 4, 5, and 6. This applies regardless of the breadth of the range.

Whenever a numerical range is indicated herein, it is meant to include any cited numeral (fractional or integral) within the indicated range. The phrases "ranging/ranges between" a first indicate number and a second indicate number and "ranging/ranges from" a first indicate number "to" a second indicate number are used herein 5 interchangeably and are meant to include the first and second indicated numbers and all the fractional and integral numerals therebetween.

As used herein the term "method" refers to manners, means, techniques and procedures for accomplishing a given task including, but not limited to, those manners, means, techniques and procedures either known to, or readily developed from known 10 manners, means, techniques and procedures by practitioners of the chemical, pharmacological, biological, biochemical and medical arts.

As used herein, the term "treating" includes abrogating, substantially inhibiting, slowing or reversing the progression of a condition, substantially ameliorating clinical or aesthetical symptoms of a condition or substantially preventing the appearance of 15 clinical or aesthetical symptoms of a condition.

It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, may also be provided in combination in a single embodiment. Conversely, various features of the invention, which are, for brevity, described in the context of a single embodiment, may also be provided 20 separately or in any suitable subcombination or as suitable in any other described embodiment of the invention. Certain features described in the context of various embodiments are not to be considered essential features of those embodiments, unless the embodiment is inoperative without those elements.

Various embodiments and aspects of the present invention as delineated 25 hereinabove and as claimed in the claims section below find experimental support in the following examples.

## EXAMPLES

Reference is now made to the following examples, which together with the above 30 descriptions illustrate some embodiments of the invention in a non-limiting fashion.

Generally, the nomenclature used herein and the laboratory procedures utilized in the present invention include molecular, biochemical, microbiological and

recombinant DNA techniques. Such techniques are thoroughly explained in the literature. See, for example, "Molecular Cloning: A laboratory Manual" Sambrook et al., (1989); "Current Protocols in Molecular Biology" Volumes I-III Ausubel, R. M., ed. (1994); Ausubel et al., "Current Protocols in Molecular Biology", John Wiley and Sons, 5 Baltimore, Maryland (1989); Perbal, "A Practical Guide to Molecular Cloning", John Wiley & Sons, New York (1988); Watson et al., "Recombinant DNA", Scientific American Books, New York; Birren et al. (eds) "Genome Analysis: A Laboratory Manual Series", Vols. 1-4, Cold Spring Harbor Laboratory Press, New York (1998); methodologies as set forth in U.S. Pat. Nos. 4,666,828; 4,683,202; 4,801,531; 5,192,659 10 and 5,272,057; "Cell Biology: A Laboratory Handbook", Volumes I-III Cellis, J. E., ed. (1994); "Current Protocols in Immunology" Volumes I-III Coligan J. E., ed. (1994); Stites et al. (eds), "Basic and Clinical Immunology" (8th Edition), Appleton & Lange, Norwalk, CT (1994); Mishell and Shiigi (eds), "Selected Methods in Cellular 15 Immunology", W. H. Freeman and Co., New York (1980); available immunoassays are extensively described in the patent and scientific literature, see, for example, U.S. Pat. Nos. 3,791,932; 3,839,153; 3,850,752; 3,850,578; 3,853,987; 3,867,517; 3,879,262; 3,901,654; 3,935,074; 3,984,533; 3,996,345; 4,034,074; 4,098,876; 4,879,219; 20 5,011,771 and 5,281,521; "Oligonucleotide Synthesis" Gait, M. J., ed. (1984); "Nucleic Acid Hybridization" Hames, B. D., and Higgins S. J., eds. (1985); "Transcription and Translation" Hames, B. D., and Higgins S. J., Eds. (1984); "Animal Cell Culture" Freshney, R. I., ed. (1986); "Immobilized Cells and Enzymes" IRL Press, (1986); "A Practical Guide to Molecular Cloning" Perbal, B., (1984) and "Methods in Enzymology" Vol. 1-317, Academic Press; "PCR Protocols: A Guide To Methods And Applications", Academic Press, San Diego, CA (1990); Marshak et al., "Strategies for 25 Protein Purification and Characterization - A Laboratory Course Manual" CSHL Press (1996) . Other general references are provided throughout this document. The procedures therein are believed to be well known in the art and are provided for the convenience of the reader.

**MATERIALS AND METHODS****Materials:**

Glucose oxidase (stabilizer-free) was obtained from Sigma-Aldrich;

Lignin peroxidase isoenzyme H1 (from *P. chrysosporium*) was prepared as described in International Patent Application Publication WO2004/052275, with the following minor modifications: a spore solution with  $7.5 \times 10^4$  spores/ml was directly spiked to the growth medium; the growth medium further comprised yeast extract and nutrient broth; and aeration of the fermentor was 4 liters/minute with agitation of 150 rotations per minute (lignin peroxidases from of other white rot fungi are also contemplated for use in embodiments of the invention);

Veratryl alcohol was obtained from Fluka.

**Cream formulation:**

Exemplary formulations were prepared in the form of two complementary creams: an enzyme cream comprising 25 units/gram LIP isoenzyme H1 fraction, 25 units/gram GOX and 0.10 weight percent veratryl alcohol (6 mmol/kg of cream), and the complementary activator cream comprised glucose (3.5 mmol/kg of cream).

For comparison, a complementary enzyme cream and activator cream were formulated for activation of LIP by non-enzymatically produced  $\text{H}_2\text{O}_2$ . This enzyme cream comprised the LIP isoenzyme H1 fraction (25 units/gram) and veratryl alcohol (6 mmol/kg of cream) without GOX, and the complementary activator cream comprised  $\text{H}_2\text{O}_2$  (3.53 mmol/kg of cream).

Other ingredients (e.g., the carrier) in the creams were as described in Table 5 below.

**Table 5: Non-active ingredients of exemplary creams**

Ingredient	Weight percentage	Trade names and other details
Glycerin	8.0 %	Superol® V
Propylene glycol	2.0 %	
Dimethicone	1.0 %	xiameter® pmx-200
Dicaprylyl carbonate	1.0 %	Cetiol® CC

Paraffinum liquidum	0.50 %	Draekol® 19
Cetyl alcohol	0.50 %	Lanette® C16 98-100MY
Stearyl alcohol	0.50 %	Lanette® C18 98-100MY
Polyisobutene	0.200 – 0.213 %	0.80 - 0.85 % Sepiplus™ 400 (25 % polyisobutene, 5 %
Polysorbate 20	0.040 - 0.043 %	polysorbate 20 and 70 % polyacrylate 13)
Polyacrylate 13	0.560 - 0.595 %	
Phenoxyethanol	0.50 %	
Chlorphenesin	0.30 %	Elestab® CPN
<i>Caesalpinia spinosa</i> gum	0.30 %	Solagum™ Tara
Citric acid	0.011- 0.040 %	Added as aqueous solution
Deionized water	To 100 %	

***Visual analogue scale (VAS):***

A visual analogue scale (VAS) is a method that is used to measure a characteristic or attitude that is believed to range across a continuum of values which 5 cannot easily be directly measured. The VAS was presented as a horizontal line, anchored by word descriptors at each end, as described below. The ends of the scale were labeled by the values 0 (for most desirable outcomes) and 10 (for least desirable outcome). Statistical significance was determined by calculating p values according to repeated measures ANOVA. A VAS was used to evaluate each of the following 10 variables:

Uniformity of pigmentation (anchored by the descriptors "uniform, even" and ""uneven, blotchy, mottled");

Clarity (anchored by the descriptors "clear, even" and "dull, opaque");

Lightening/Brightening (anchored by the descriptors "light/bright appearance" 15 and "dull, matte");

Skin tone (anchored by the descriptors "even, healthy skin color" and "uneven, discolored appearance");

Radiance (anchored by the descriptors "radiant, luminous appearance" and "dull, matte, sallow appearance"); and

5 Overall appearance (anchored by the descriptors "healthy looking" and "unhealthy looking").

***Digital photography:***

10 Digital photography of the faces of treated subjects was performed using a VISIA®-CR facial imaging system, placed in an environment with stable and consistent ambient lighting. The system remained in the same location/orientation for different visits by each subject.

***EXAMPLE 1***

***Oxidation of melanin by lignin peroxidase, glucose oxidase and glucose in cream formulation***

15 In order to evaluate the ability of lignin peroxidase (LIP) to lighten skin upon activation by an enzymatic source of hydrogen peroxide, an enzyme cream comprising LIP, veratryl alcohol and GOX (glucose oxidase) and an activator cream comprising glucose were prepared as described hereinabove, and tested on solubilized melanin. The results were compared with those obtained with an enzyme cream comprising LIP 20 and veratryl alcohol and an activator cream comprising hydrogen peroxide, prepared as described hereinabove.

0.3 gram of each enzyme cream was first mixed with melanin (140 µg/ml), resulting in a grayish black color. 0.3 gram of the complementary activator cream was then added.

25 As shown in Figure 1, the color of each enzyme cream formulation changed from grayish black to beige immediately after addition of the activator cream.

These results indicate that LIP is efficiently activated by the H<sub>2</sub>O<sub>2</sub> produced enzymatically by the GOX-glucose system, to a comparable degree as is obtained with the conventional non-enzymatic H<sub>2</sub>O<sub>2</sub> activation system.

**EXAMPLE 2*****Skin lightening by lignin peroxidase, glucose oxidase and glucose in cream formulation upon treatment for 21 days***

The exemplary enzyme and activator creams described herein were tested for  
5 their ability to lighten skin *in vivo*.

Lignin peroxidase, glucose oxidase and glucose were administered to 10 female subjects (aged 23-65), of which 8 completed the study. Compositions were applied to facial skin according to the following regimen:

**Morning**

10 Step 1 - ELURE<sup>TM</sup> facial wash  
Step 2 – cream comprising lignin peroxidase and glucose oxidase (as described in Materials and Methods section)  
Step 3 – activator cream comprising glucose (as described in Materials and Methods section)

15 Step 4 - SPF 30 sunscreen (for reducing formation of new melanin)

**Evening**

Step 1 - facial wash  
Step 2 – cream comprising lignin peroxidase and glucose oxidase  
Step 3 – activator cream comprising glucose  
20 After the cream comprising lignin peroxidase and glucose oxidase was applied and absorbed by the skin, the activator cream was applied.

Skin color was measured using a Mexameter<sup>®</sup> skin colorimeter, which was used to evaluate melanin and erythema (hemoglobin) contributions to the skin color. The skin color was measured in various regions (forehead, right cheek, left cheek, and spots of interest). In each region skin color was measured at different points and the results were averaged. Skin color was assessed on day 0 (i.e., prior to treatment) and on days 7 and 25 21 of the treatment.

As shown in Figures 2A-9B, the treatment resulted in at least some reduction in melanin content, in all 8 subjects tested, and the reductions in melanin content were 30 usually treatment duration-dependent (Figures 2A, 6A, 7A, 9A).

As further shown therein, the treatment did not have any consistent effect on erythema levels.

As shown in Figures 3A, 5A and 6A, the treatment resulted in reduction in melanin content of most spots on the skin.

In one subject (Figure 9B), an increase of erythema of the cheeks on day 21 was observed, but judged to be caused by irritation unrelated to the treatment (irritation by toner used to remove smudged eye shadow).

These results indicate that treatment with lignin peroxidase, glucose oxidase and glucose is effective at lightening skin by decreasing melanin content, without causing significant irritation of the skin.

10

### **EXAMPLE 3**

#### ***Skin lightening by lignin peroxidase, glucose oxidase and glucose in cream formulation upon treatment for 8 weeks***

The formulation and daily regimen described in Example 2 were tested on 28 female volunteer subjects (aged 33-50) with apparent but mild, diffuse hyperpigmentation including melasma, for a period of 8 weeks. Subjects fulfilling any of the following criteria were excluded: sensitive, hypersensitive, or with atopic skin; history of allergy to cosmetic products or skin cleansers; history of a disease/condition or concurrent illness that could interfere with the outcome of the study; pregnant, planning to be pregnant, or breast feeding; skin problems in the area for testing, including skin diseases, moles, acne, red spots, telangiectasis, etc; dermatologist treatments, or procedures with potential interference within the last 3 months; currently involved in, or participated in any clinical skin care study within the last 3 months; judged by a physician to be unsuitable for enrollment in this clinical trial.

The effect of the formulation was assessed by evaluating skin pigmentation at 0, 4, 6 and 8 weeks from the beginning of treatment, using the following techniques:

- 1) instrumental measurement using a Mexameter® skin colorimeter;
- 2) visual analogue scale (VAS) assessment by a dermatologist;
- 3) digital photography with VISIA®-CR facial imaging system; and
- 4) self-assessment by subjects.

As shown in Figures 10A and 10B, the formulation progressively lowered the melanin index in pigmented skin (Figure 10A) and non-pigmented skin (Figure 10B)

throughout the 8 weeks of treatment (as determined using a Mexameter® skin colorimeter).

Furthermore, after 4 weeks, the melanin index in pigmented skin was lowered in 22 of the 28 subjects (78.6 %), and the melanin index in non-pigmented skin was lowered in 20 of the 28 subjects (71.4 %). After 6 and 8 weeks, the melanin index in each of pigmented and non-pigmented skin was lowered in 27 of the 28 subjects (96.4 %).

A visual analogue scale (VAS) was used, as described in the Materials and Methods section, to evaluate uniformity of pigmentation, clarity, lightness/brightness, tone, radiance and overall appearance of treated skin, at various time points over the course of the treatment.

As shown in Figures 11-16, as well as in Tables 6 and 7, the formulation progressively enhanced the uniformity of pigmentation (Figure 11), clarity (Figure 12), lightness/brightness (Figure 13), tone (Figure 14), radiance (Figure 15) and overall appearance (Figure 16) of treated skin, over the course of 8 weeks of treatment.

The average degree of improvements at each time point shown in Figures 11-16 are presented in Table 6:

**Table 6: Average improvement and statistical significance of skin appearance relative to beginning of treatment (n=28)**

	Improvement (%) (p value)		
	after 4 weeks	after 6 weeks	after 8 weeks
Uniformity of pigmentation	1.26 % (0.223)	2.96 % (0.106)	5.47 % (0.046)
Clarity	4.00 % (0.004)	4.65 % (0.033)	8.59 % (0.002)
Lightening/Brightening	9.28 % (0.000)	8.09 % (0.001)	11.52 % (0.000)
Tone	8.03 % (0.000)	6.13 % (0.015)	12.53 %

			(0.000)
Radiance	11.64 % (0.000)	12.15 % (0.000)	18.38 % (0.000)
Overall Appearance	14.25 % (0.000)	13.21 % (0.000)	21.57 % (0.000)

Figure 17 shows the changes in facial complexion in individual subjects treated for 8 weeks with the formulation.

The percentages of the 28 subjects which showed significant improvement in clarity, lightening/brightening, skin tone, radiance and/or overall appearance of whole face complexion after 4, 6 and 8 weeks of treatment are presented in Table 7:

**Table 7: Percentage of subjects showing improvement of skin appearance relative to beginning of treatment (n=28)**

	after 4 weeks	after 6 weeks	after 8 weeks
Uniformity of pigmentation	46.4 % (13/28)	60.7 % (17/28)	67.9 % (19/28)
Clarity	67.9 % (19/28)	64.3 % (18/28)	67.9 % (19/28)
Lightening/Brightening	89.3 % (25/28)	78.6 % (22/28)	75.0 % (21/28)
Tone	78.6 % (22/28)	67.9 % (19/28)	82.1 % (23/28)
Radiance	92.9 % (26/28)	92.9 % (26/28)	96.4 % (27/28)
Overall Appearance	92.9 % (26/28)	82.1 % (23/28)	89.3 % (25/28)

As shown in Table 8, satisfaction of the subjects with the formulation progressively increased over the course of 8 weeks of treatment (as determined according to self-assessment by questionnaire).

**Table 8: Responses of subjects to question "Overall, are you satisfied with the product you used?"**

5

Response	0 weeks		4 weeks		6 weeks		8 weeks	
	n	%	n	%	n	%	n	%
Strongly disagree (1)	0	0.0	0	0.0	0	0.0	0	0.0
Disagree (2)	0	0.0	1	3.6	0	0.0	1	3.6
Fair (3)	27	96.4	9	32.1	8	28.6	6	21.4
Agree (4)	1	3.6	18	64.3	16	57.1	18	64.3
Strongly agree (5)	0	0.0	0	0.0	4	14.3	3	10.7

Furthermore, as shown in Table 9, satisfaction of the subjects with the formulation was expressed over the course of 8 weeks of treatment with respect to every topic asked about in the questionnaire.

**Table 9: Responses of subjects to specific questionnaire items (n=28)**

Questionnaire Item	Response	0 weeks		4 weeks		6 weeks		8 weeks	
		n	%	n	%	n	%	n	%
<i>Overall, have you felt improvement effect of skin brightness after application?</i>	Strongly disagree (1)	0	0.0	0	0.0	0	0.0	0	0.0
	Disagree (2)	0	0.0	0	0.0	0	0.0	1	3.6
	Fair (3)	26	92.9	11	39.3	4	14.3	3	10.7
	Agree (4)	2	7.1	15	53.6	23	82.1	22	78.6
	Strongly agree (5)	0	0.0	2	7.1	1	3.6	2	7.1
<i>Does the skin color on the pigmented area lighten?</i>	Strongly disagree (1)	0	0.0	0	0.0	0	0.0	0	0.0
	Disagree (2)	0	0.0	0	0.0	0	0.0	2	7.1
	Fair (3)	27	96.4	17	60.7	11	39.3	8	28.6
	Agree (4)	1	3.6	10	35.7	16	57.1	17	60.7
	Strongly agree (5)	0	0.0	1	3.6	1	3.6	1	3.6
<i>Do you feel the change of pigmented area?</i>	Strongly disagree (1)	0	0.0	0	0.0	0	0.0	0	0.0
	Disagree (2)	1	3.6	0	0.0	0	0.0	1	3.6
	Fair (3)	27	96.4	19	67.9	15	53.6	15	53.6
	Agree (4)	0	0.0	9	32.1	12	42.9	10	35.7
	Strongly agree (5)	0	0.0	0	0.0	1	3.6	2	7.1
<i>Does your skin feel smoother after application?</i>	Strongly disagree (1)	0	0.0	0	0.0	0	0.0	0	0.0
	Disagree (2)	0	0.0	0	0.0	0	0.0	1	3.6
	Fair (3)	23	82.1	10	35.7	8	28.6	6	21.4
	Agree (4)	5	17.9	18	64.3	18	64.3	19	67.9
	Strongly agree (5)	0	0.0	0	0.0	2	7.1	2	7.1
<i>Is this product</i>	Strongly disagree (1)	0	0.0	0	0.0	0	0.0	0	0.0

<i>more effective than the products you previously used?</i>	Disagree (2)	0	0.0	0	0.0	0	0.0	1	3.6
	Fair (3)	27	96.4	11	39.3	9	32.1	9	32.1
	Agree (4)	0	0.0	17	60.7	17	60.7	16	57.1
	Strongly agree (5)	1	3.6	0	0.0	2	7.1	2	7.1
<i>Would you recommend this product to others?</i>	Strongly disagree (1)	0	0.0	1	3.6	0	0.0	0	0.0
	Disagree (2)	0	0.0	1	3.6	0	0.0	1	3.6
	Fair (3)	27	96.4	9	32.1	10	35.7	8	28.6
	Agree (4)	0	0.0	16	57.1	16	57.1	14	50.0
	Strongly agree (5)	1	3.6	1	3.6	2	7.1	5	17.9
<i>Would you purchase this product if it is released?</i>	Strongly disagree (1)	0	0.0	2	7.1	0	0.0	0	0.0
	Disagree (2)	0	0.0	0	0.0	0	0.0	1	3.6
	Fair (3)	25	89.3	8	28.6	8	28.6	7	25.0
	Agree (4)	3	10.7	17	60.7	17	60.7	16	57.1
	Strongly agree (5)	0	0.0	1	3.6	3	10.7	4	14.3

As further determined by self-assessment, none of the 28 subjects reported any discomfort or allergic reaction while using the formulation.

These results confirm that treatment with lignin peroxidase, glucose oxidase and 5 glucose is effective at lightening skin by decreasing melanin content, without causing significant irritation of the skin, and further indicate that the complexion of the treated skin continues to improve during the course of treatment.

Although the invention has been described in conjunction with specific 10 embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such alternatives, modifications and variations that fall within the spirit and broad scope of the appended claims.

In addition, citation or identification of any reference in this application shall not be construed as an admission that such reference is available as prior art to the present invention. To the extent that section headings are used, they should not be construed as necessarily limiting.

WHAT IS CLAIMED IS:

1. A cosmetic composition for lightening a skin and/or hair of a subject, the composition comprising a lignin peroxidase, a hydrogen peroxide-producing enzyme, and a substrate of said hydrogen peroxide-producing enzyme, wherein said hydrogen peroxide-producing enzyme comprises glucose oxidase and said substrate comprises D-glucose, and wherein a concentration of said glucose oxidase is in a range of from 1.25 to 125 units/gram and a concentration of said D-glucose is in a range of from 0.175  $\mu$ mole/gram to 17.5  $\mu$ mole/gram.
2. The composition of claim 1, wherein a concentration of said lignin peroxidase is in a range of from 1.25 units/gram to 125 units/gram.
3. The composition of claim 1, wherein a concentration of said lignin peroxidase is in a range of from 1.25 units/gram to 12.5 units/gram.
4. The composition of any one of claims 1 to 3, wherein a concentration of said hydrogen peroxide-producing enzyme is in a range of from 4 to 40 units/gram.
5. The composition of any one of claims 1 to 4, wherein a concentration of said substrate of said hydrogen peroxide-producing enzyme is in a range of from 0.5  $\mu$ mole/gram to 6  $\mu$ mole/gram.
6. The composition of any one of claims 1 to 5, further comprising an oxidizing mediator.
7. The composition of claim 6, wherein said oxidizing mediator is veratryl alcohol.
8. The composition of claim 6, wherein said oxidizing mediator is veratrole.

9. The composition of claim 6, wherein said oxidizing mediator is 1,4-dimethoxybenzene.

10. The composition of any one of claims 6 to 9, wherein a concentration of said oxidizing mediator is in a range of from 0.03  $\mu$ mole/gram to 300  $\mu$ mole/gram.

11. The composition of any one of claims 6 to 9, wherein a concentration of said oxidizing mediator is in a range of from 0.3  $\mu$ mole/gram to 30  $\mu$ mole/gram.

12. The composition of any one of claims 1 to 11, wherein said lignin peroxidase is isoenzyme H1.

13. The composition of any one of claims 1 to 11, wherein said lignin peroxidase is a modified form of isoenzyme H2.

14. The composition of claim 13, wherein said lignin peroxidase is a dephosphorylated form of isoenzyme H2.

15. The composition of any one of claims 1 to 14, wherein said lignin peroxidase is a lignin peroxidase of a white rot fungus.

16. The composition of any one of claims 1 to 14, wherein said lignin peroxidase is an extract of a *Phanerochaete chrysosporium* fungus.

17. The composition of any one of claims 1 to 16, wherein said cosmetic composition is in a form of a mask.

18. The composition of any one of claims 1 to 16, wherein said cosmetic composition is in a form of a serum.

19. The composition of any one of claims 1 to 16, wherein said cosmetic composition is in a form of a lotion.

20. The composition of any one of claims 1 to 16, wherein said cosmetic composition is in a form of a cream.

21. The composition of any one of claims 1 to 16, wherein said cosmetic composition is in a form of a shampoo.

22. The composition of any one of claims 1 to 16, wherein said cosmetic composition is in a form of a gel.

23. The composition of any one of claims 1 to 16, wherein said cosmetic composition is in a form of a toner.

24. The composition of any one of claims 1 to 23, further comprising a cosmetically acceptable carrier.

25. The composition of claim 24, wherein said carrier comprises an emulsion.

26. The composition of any one of claims 1 to 25, being packaged in an air-tight container.

27. The composition of any one of claims 1 to 26, being for lightening a skin of a subject.

28. A kit for lightening a skin and/or hair of a subject, the kit comprising a hydrogen peroxide-producing enzyme, a substrate of said hydrogen peroxide-producing enzyme and a lignin peroxidase, wherein said hydrogen peroxide-producing enzyme comprises glucose oxidase and said substrate comprises D-glucose, wherein said hydrogen peroxide-producing enzyme forms a part of a first composition and said substrate forms a part of a second composition, said first and said second compositions being packaged individually within the kit, and wherein a concentration of said glucose oxidase in said first composition is in a range of from 2.5 to 250 units/gram and a concentration of said D-

glucose in said second composition is in a range of from 0.35  $\mu$ mole/gram to 35  $\mu$ mole/gram.

29. The kit of claim 28, wherein said first composition further comprises said lignin peroxidase.

30. The kit of claim 28, wherein said second composition further comprises said lignin peroxidase.

31. The kit of claim 28, wherein said lignin peroxidase forms a part of a third composition within the kit, said third composition being packaged individually within the kit.

32. The kit of any one of claims 28 to 31, wherein a concentration of said lignin peroxidase in a composition comprising said lignin peroxidase is in a range of from 2.5 to 250 units/gram.

33. The kit of claim 32, wherein a concentration of said lignin peroxidase is in a range of from 2.5 units/gram to 25 units/gram.

34. The kit of any one of claims 28 to 33, comprising an oxidizing mediator, said hydrogen peroxide-producing enzyme, said substrate of said hydrogen peroxide-producing enzyme and said lignin peroxidase.

35. The kit of claim 34, wherein said oxidizing mediator forms a part of a composition that comprises said lignin peroxidase.

36. The kit of claim 34 or claim 35, wherein said oxidizing mediator is veratryl alcohol.

37. The kit of claim 34 or claim 35, wherein said oxidizing mediator is veratrole.

38. The kit of claim 34 or claim 35, wherein said oxidizing mediator is 1,4-dimethoxybenzene.

39. The kit of any one of claims 34 to 38, wherein a concentration of said oxidizing mediator in a composition comprising said oxidizing mediator is in a range of from 0.06  $\mu$ mole/gram to 600  $\mu$ mole/gram.

40. The kit of any one of claims 34 to 38, wherein a concentration of said oxidizing mediator in a composition comprising said oxidizing mediator is in a range of from 0.6  $\mu$ mole/gram to 60  $\mu$ mole/gram.

41. The kit of any one of claims 28 to 40, wherein a concentration of said hydrogen peroxide-producing enzyme in said first composition is in a range of from 8 to 80 units/gram.

42. The kit of any one of claims 28 to 41, wherein a concentration of said substrate in said second composition is in a range of from 1  $\mu$ mole/gram to 12  $\mu$ mole/gram.

43. The kit of any one of claims 28 to 42, wherein said lignin peroxidase is isoenzyme H1.

44. The kit of any one of claims 28 to 42, wherein said lignin peroxidase is a modified form of isoenzyme H2.

45. The kit of claim 44, wherein said lignin peroxidase is a dephosphorylated form of isoenzyme H2.

46. The kit of any one of claims 28 to 45, wherein said lignin peroxidase is a lignin peroxidase of a white rot fungus.

47. The kit of any one of claims 28 to 45, wherein said lignin peroxidase is an extract of a *Phanerochaete chrysosporium* fungus.

48. The kit of any one of claims 28 to 47, wherein said composition which is packaged individually within the kit is in a form of a mask.

49. The kit of any one of claims 28 to 47, wherein said composition which is packaged individually within the kit is in a form of a serum.

50. The kit of any one of claims 28 to 47, wherein said composition which is packaged individually within the kit is in a form of a lotion.

51. The kit of any one of claims 28 to 47, wherein said composition which is packaged individually within the kit is in a form of a cream.

52. The kit of any one of claims 28 to 47, wherein said composition which is packaged individually within the kit is in a form of a shampoo.

53. The kit of any one of claims 28 to 47, wherein said composition which is packaged individually within the kit is in a form of a gel.

54. The kit of any one of claims 28 to 47, wherein said composition which is packaged individually within the kit is in a form of a toner.

55. The kit of any one of claims 28 to 54, wherein said composition which is packaged individually within the kit further comprises a cosmetically acceptable carrier.

56. The kit of claim 55, wherein said carrier comprises an emulsion.

57. The kit of any one of claims 1 to 56, being for lightening a skin of a subject.

58. A cosmetic method of lightening a skin and/or hair of a subject, the method comprising contacting a skin and/or hair region of the subject with the composition of any one of claims 1 to 25, thereby lightening the skin and/or hair of the subject.

59. The method of claim 58, wherein said composition is formed by a lignin peroxidase, a hydrogen peroxide-producing enzyme and a substrate of said hydrogen peroxide-producing enzyme as packaged within the kit of any one of claims 28 to 56.

60. The method of any one of claims 58 to 59, wherein said contacting is effected at least five times.

61. The method of any one of claims 58 to 60, wherein said contacting is effected once or twice per day.

62. The method of any one of claims 58 to 61, being for lightening a skin of a subject.

63. The composition of any one of claims 1 to 27, the kit of any one of claims 28 to 57 or the method of any one of claims 58 to 62, wherein said lightening said skin comprises lightening whole skin complexion.

64. The composition of any one of claims 1 to 27, the kit of any one of claims 28 to 57 or the method of any one of claims 58 to 62, wherein said skin comprises an uneven skin tone.

65. The composition of any one of claims 1 to 27, the kit of any one of claims 28 to 57 or the method of any one of claims 58 to 62, wherein said skin comprises a dark spot.

66. The composition of any one of claims 1 to 27, the kit of any one of claims 28 to 57 or the method of any one of claims 58 to 62, wherein said skin comprises a freckle.

67. The composition of any one of claims 1 to 27, the kit of any one of claims 28 to 57 or the method of any one of claims 58 to 62, wherein said skin comprises melasma.

68. The composition of any one of claims 1 to 27, the kit of any one of claims 28 to 57 or the method of any one of claims 58 to 62, wherein said skin comprises hyperpigmentation.

69. The composition of any one of claims 1 to 27, the kit of any one of claims 28 to 57 or the method of any one of claims 58 to 62, wherein said skin comprises skin discoloration.

70. The composition of any one of claims 1 to 27, the kit of any one of claims 28 to 57 or the method of any one of claims 58 to 62, wherein said skin comprises an age spot.

71. The composition of any one of claims 1 to 27, the kit of any one of claims 28 to 57 or the method of any one of claims 58 to 62, wherein said skin comprises an acne mark.

72. The composition of any one of claims 1 to 27, the kit of any one of claims 28 to 57 or the method of any one of claims 58 to 62, wherein said skin comprises a scar.




FIG. 1




FIG. 2A




FIG. 2B




FIG. 3A




FIG. 3B




FIG. 4A




FIG. 4B




FIG. 5A




FIG. 5B




FIG. 6A




FIG. 6B




FIG. 7A




FIG. 7B




FIG. 8A




FIG. 8B




FIG. 9A




FIG. 9B




FIG. 10A




FIG. 10B




FIG. 11.




FIG. 12




FIG. 13








FIG. 17

