



(22) **Date de dépôt/Filing Date:** 2003/08/08
(41) **Mise à la disp. pub./Open to Public Insp.:** 2004/04/29
(45) **Date de délivrance/Issue Date:** 2017/04/11
(62) **Demande originale/Original Application:** 2 495 082
(30) **Priorité/Priority:** 2002/08/09 (US60/402,228)

(51) **Cl.Int./Int.Cl.** **A23K 50/30** (2016.01),
A23K 10/30 (2016.01), **A23K 20/189** (2016.01),
A61K 38/48 (2006.01)
(72) **Inventeur/Inventor:**
SHIH, JASON C.H., US
(73) **Propriétaire/Owner:**
NORTH CAROLINA STATE UNIVERSITY, US
(74) **Agent:** AIRD & MCBURNEY LP

(54) **Titre :** PROCÉDES ET COMPOSITIONS POUR AMÉLIORER LA CROISSANCE D'UNE VOLAILLE D'ABATTAGE
(54) **Title:** METHODS AND COMPOSITIONS FOR IMPROVING GROWTH OF MEAT-TYPE POULTRY

(57) **Abrégé/Abstract:**

The present invention provides methods of improving growth performance, improving the efficiency of feed utilization, increasing feed digestibility, and decreasing mortality of immature and developing animals receiving animal feed. Methods of producing a crude keratinase enzyme extract and animal feed supplements for achieving the same are also provided.



Abstract

The present invention provides methods of improving growth performance, improving the efficiency of feed utilization, increasing feed digestibility, and decreasing mortality of immature and developing animals receiving animal feed. Methods of producing a crude keratinase enzyme extract and animal feed supplements for achieving the same are also provided.

METHODS AND COMPOSITIONS FOR IMPROVING GROWTH OF MEAT-TYPE POULTRY

Cross Reference to Related Applications

5 This application claims the benefit of U.S. Publication No. 2004/0166196 filed
August 9, 2002.

Statement of Federal Support

10 Research directed to this invention is supported in part by US Department of
Agriculture Small Business Innovation Research Grant No. 2002-33610-11850. The
Government has certain rights in this invention.

Field of the Invention

15 The present invention relates to methods of improving growth performance of
immature and developing animals receiving animal feed and animal feed supplements
for achieving the same.

Background of the Invention

20 Broiler chick starter diets contain a considerable amount of crude protein. Most
of the crude protein is obtained from traditional feed ingredients such as soybean meal.
Approximately 90% of the crude protein present in soybean meal (48% crude protein
content) is highly digestible for poultry (National Research Council (1994). Nutrient
requirements of poultry. 9th revised Ed. National Academy Press, Washington, DC).
Although the traditional corn-soybean meal broiler starter diets are considered highly
25 digestible, they often contain a variety of complex proteins that are not easily digested
by a young chick due to the lack of necessary innate enzymes at early stages of life
(Uni, et al. (1999) *Poultry Sci.* 78: 215-222). Inclusion of proteases in broiler diets has
been suggested, but much of the early work with protease addition to cereal grain-based
diets did not result in any improvements in bird performance (Jensen, et al. (1957)
30 *Poultry Sci.* 36: 919-921).

More recently, enzyme supplementation of poultry diets with enzyme mixtures, including proteases and amylases, has produced some improvements in growth performance (Greenwood, et al. (2002) *Poultry Sci.* 81(Suppl. 1): 25; Burrows, et al. (2002) *Poultry Sci.* 81(Suppl. 1): 29; Short, et al. (2002) *Poultry Sci.* 81(Suppl. 1): 136).
 5 Supplementing a corn-soybean broiler starter diet with an enzyme preparation containing a mixture of xylanase, protease, and amylase resulted in improvements in body weight at 14 and 42 days of age with no significant effects on feed conversion ratio (Greenwood, et al. (2002) *supra*). Upon supplementing corn-soy based duck diets with the same enzyme mixture, the enzyme supplementation resulted in improvements in body weight
 10 gain and feed conversion ratio (Burrows, et al. (2002) *supra*).

Poultry feed further contains some complex antinutritional and/or indigestible compounds. Some of these compounds, such as the nonstarch polysaccharides, absorb water into a viscous mass within the chyme from which nutrients are not readily absorbed (Odetallah, 2000; Odetallah, et al. (2002) *supra*). As chyme
 15 viscosity increases, the rate of diffusion of digestive enzymes and nutrients decreases, thus impeding nutrient absorption by the enterocyte. Fat micelle formation and absorption also decrease as chyme viscosity increases, thus impairing absorption of many of the fat-soluble compounds, including fat-soluble vitamins, pigments, and lipids (Ferket and Veldkamp (1999) *In: Proceedings of the 1998 World Poultry
 20 Science Association*, pgs 43-52). Therefore, viscosity reduction achieved by endolytic enzyme activity may play a role in the improvement seen in young chicks fed high-viscosity cereals, and the relative effectiveness of various enzymes appears to be related to their viscosity-reducing capability (Rotter, et al. (1990) *J. Sci. Food Agric.* 50:19-27).

25 PWD-1 keratinase is an enzyme that was originally purified from the growth medium of *Bacillus licheniformis* PWD-1 (Williams, et al. (1990) *Appl. Environ. Microbiol.* 56:1509-1515; Lin, et al. (1992) *Appl. Environ. Microbiol.* 58:3271-3275). PWD-1 keratinase hydrolyzes a broad range of protein substrates including casein, collagen, elastin and keratin (Shih (2001) *In: Proceedings International Conference of
 30 Agricultural Science and Technology*, Beijing, China, pgs 244-247). PWD-1 keratinase has been used to produce hydrolyzed feather meal by incubating commercial feather meal with cell-free keratinase overnight (Carter (1998) *Bacterial Keratinase: Assay development and nutritional application*. Ph.D. Thesis, North Carolina State University,

Raleigh, NC). *See also* U.S. Patent Nos. 4,908,220; 5,186,961; and 5,063,161 to Shih et al.

In spite of the foregoing, there remains a need for additional methods of enhancing growth performance of broiler chicks and animal feed supplements that
5 accomplish the same.

Summary of the Invention

The present invention provides methods and compositions that enhance growth performance of immature and developing animals receiving animal feed.

10 An aspect of the invention relates to a method of growing meat-type poultry comprising feeding meat-type poultry a corn-soybean meal feed as a poultry diet wherein the feed further comprises keratinase in an amount effective to enhance the weight gain of meat-type poultry.

Another aspect of the invention relates to a method of growing meat-type
15 poultry comprising feeding meat-type poultry a corn-soybean meal feed as a starter diet wherein the feed further comprises keratinase in an amount effective to enhance the weight gain of the meat-type poultry.

A further aspect of the invention relates to a method of improving the efficiency of feed utilization of an animal feed in meat-type poultry comprising
20 feeding meat-type poultry a corn-soybean meal feed as a poultry diet wherein the feed further comprises keratinase in an amount effective to improve the efficiency of feed utilization of an animal feed in meat-type poultry.

An additional aspect of the present invention relates to a method of increasing the digestibility of an animal feed in meat-type poultry comprising feeding meat-type
25 poultry a corn-soybean meal feed as a poultry diet wherein the feed further comprises keratinase an amount effective to increase the digestibility of an animal feed in meat-type poultry.

Another aspect of the invention relates to a method of reducing mortality in meat-type poultry comprising feeding meat-type poultry a corn-soybean meal feed as
30 a starter diet wherein the feed further comprises keratinase in an amount effective to reduce mortality of meat-type poultry.

A further aspect of the invention relates to an animal feed consisting essentially of keratinase, protein, and carbohydrate.

An additional aspect of the invention relates to a method of producing crude keratinase enzyme extract.

The present invention further relates to improving the nutritional status of a hatchling, and thereby increasing disease resistance and survivability of the immature
5 bird to achieve a greater level of growth performance meat-type poultry.

In accordance with an aspect, there is provided a method of feeding a pig, comprising:

feeding said pig an animal feed comprising a corn-soybean meal feed as a pig diet, wherein said pig diet is essentially free of keratin, and

10 said animal feed further comprises *Bacillus licheniformis* PWD-1 keratinase in an amount effective to enhance the weight gain of said pig.

In accordance with another aspect, there is provided a method of supplementing an animal feed for a pig, the method consisting essentially of:

feeding said pig a corn-soybean meal feed,

15 said animal feed comprising *Bacillus licheniformis* PWD-1 keratinase in an amount effective to enhance the weight gain of said pig, wherein said animal feed is essentially free of keratin.

In accordance with a further aspect, there is provided a method of improving the efficiency of feed utilization of an animal feed in a pig, comprising:

20 feeding said pig a corn-soybean meal feed as a pig diet, wherein said pig diet is essentially free of keratin, and

said animal feed further comprises *Bacillus licheniformis* PWD-1 keratinase in an amount effective to improve the efficiency of feed utilization of an animal feed in said pig.

25 In accordance with another aspect, there is provided a method of increasing the digestibility of an animal feed in a pig, comprising:

feeding said pig a corn-soybean meal feed as a pig diet, wherein said pig diet is essentially free of keratin, and

30 said animal feed further comprises *Bacillus licheniformis* PWD-1 keratinase an amount effective to increase the digestibility of an animal feed in said pig.

In accordance with a further aspect, there is provided a method of reducing mortality in a pig, comprising:

feeding said pig an animal feed comprising corn-soybean meal feed as a pig diet, wherein said pig diet is essentially free of keratin, and

said animal feed further comprises *Bacillus licheniformis* PWD-1 keratinase in an amount effective to reduce mortality of said pig.

5 In accordance with another aspect, there is provided a pig feed consisting essentially of soybean meal, corn, and keratinase, wherein said pig feed is essentially free of keratin.

In accordance with another aspect, there is provided an animal feed consisting essentially of:

- 10 (a) a *Bacillus licheniformis* PWD-1 keratinase;
 (b) one or more protein sources; and
 (c) one or more carbohydrate sources,
 wherein the animal feed is essentially free of keratin.

In accordance with another aspect, there is provided an animal feed consisting
 15 essentially of:

- (a) a *Bacillus licheniformis* PWD-1 keratinase;
 (b) one or more protein sources; and
 (c) one or more carbohydrate sources; and
 (d) one or more other substances selected from the group consisting of one or
 20 more vitamins, one or more minerals, one or more fats, and one or more antibiotics,
 wherein the animal feed is essentially free of keratin.

In accordance with another aspect, there is provided a poultry feed comprising *Bacillus licheniformis* PWD-1 keratinase, wherein the poultry feed:

- (a) is essentially free of keratin, and
 25 (b) comprises less than about 23% crude protein.

In accordance with another aspect, there is provided a poultry feed comprising *Bacillus licheniformis* PWD-1 keratinase, wherein the poultry feed:

- (a) is essentially free of keratin, and
 (b) comprises less than about 3201 kcal/kg of metabolizable energy.

30 In accordance with another aspect, there is provided a method for improving the digestibility of an animal feed essentially free of keratin, the method comprising contacting the feed with *Bacillus licheniformis* PWD-1 keratinase.

In accordance with another aspect, there is provided a method for improving the feed conversion ratio of an animal feed essentially free of keratin, the method comprising contacting the feed with *Bacillus licheniformis* PWD-1 keratinase.

In accordance with another aspect, there is provided a method for reducing the anti-nutritional content of an animal feed essentially free of keratin, the method comprising contacting the feed with *Bacillus licheniformis* PWD-1 keratinase.

In accordance with another aspect, there is provided a method for increasing the metabolizable energy of an animal feed essentially free of keratin, the method comprising contacting the feed with *Bacillus licheniformis* PWD-1 keratinase.

In accordance with another aspect, there is provided a method for improving the nutritional value of an animal feed essentially free of keratin, the method comprising contacting the feed with *Bacillus licheniformis* PWD-1 keratinase.

Detailed Description of the Invention

The foregoing and other aspects of the present invention will now be described in more detail with respect to other embodiments described herein. It should be appreciated that the invention can be embodied in different forms and should not be construed as limited to the embodiments set forth herein. Rather, these embodiments are provided so that this disclosure will be thorough and complete, and will fully convey the scope of the invention to those skilled in the art.

The terminology used in the description of the invention herein is for the purpose of describing particular embodiments only and is not intended to be limiting of the invention. As used in the description of the invention and the appended claims, the singular forms "a", "an" and "the" are intended to include the plural forms as well, unless the context clearly indicates otherwise.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs.

As used herein, the term "meat-type poultry" refers to any avian species that is produced or used for meat consumption as understood by one skilled in the art. Examples of such avian species include, but are not limited to, chickens, turkeys, ducks, geese, quail, pheasant, ratites, and the like.

As used herein, the term "immature bird" refers to a member of the avian species that lacks complete growth, differentiation, or development. Such members

can have the potential capacity to attain a definite mature form or state. An immature bird can be from about 1 to about 50 days old, preferably about 1 to about 21 days old, and more preferably about 1 to about 5 days old, or can have a body weight comparable to birds within these ranges.

As used herein, the term “developing bird” refers to a member of the avian species that is older or weighs more than an immature bird.

As used herein, the term “mature bird” refers to a member of the avian species that is older or weighs more than a developing bird.

5 As used herein, the term “broiler chick” refers to any immature chicken produced or eventually used for meat consumption.

As used herein, the term “poultry diet” refers to a diet that can be administered to a member of the avian species to promote and maintain growth of the bird. A poultry diet can contain sources of protein, vitamins, minerals, energy such as fat,
10 carbohydrates, and additional protein, antibiotics, and other substances or compounds known to be included in animal feeds, in particular, poultry feeds. Poultry diet is inclusive of, but not limited to, a starter diet, a grower-type diet, and a finisher-type diet. A “starter diet” refers to a diet that can be administered to an animal starting from birth or hatch until a desired age and/or weight is obtained. A “grower-type
15 diet” refers to a diet that can be administered to an animal upon completion of the starter growth phase. A “finisher-type diet” refers to a diet that can be administered to an animal during the period of development through the time of slaughter.

As used herein, the terms “growth” or “growth performance” refer to increases in either, or both, weight and size (e.g., height, width, diameter, circumference, etc.)
20 over that which would otherwise occur without implementation of the methods and/or administration of the compositions of the present invention. Growth can refer to an increase in the mass (e.g., weight or size) of the entire animal or of a particular tissue (e.g., muscle tissue in general or a specific muscle). Alternatively, growth can indicate a relative increase in the mass of one tissue in relation to another, in
25 particular, an increase in muscle tissue relative to other tissues (e.g., adipose tissue). Growth further relates to nutritional status and disease resistance wherein improvement of nutritional status and/or increase in disease resistance is also indicative of improved growth performance.

In view of the foregoing, embodiments according to the present invention
30 relate to methods of growing meat-type poultry, comprising feeding meat-type poultry an animal feed poultry diet wherein the feed further comprises keratinase and is added to the poultry diet in an amount effective to enhance the weight gain of the meat-type poultry. The poultry diet can be an animal feed which includes sources of protein, for example, soybean meal, fish meal, blood meal, poultry by-product (ground poultry

offal), meat meal, wheat-meal, rapeseed, canola and combinations of the same. The animal feed further includes carbohydrates, for example, corn, oats, barley, sorghum, or combinations of the same that can be ground into a meal for use in the animal feed. Additionally, the animal feed can include vitamins, minerals, fat, antibiotics, and
5 other substances or compounds as necessary or desired. Non-limiting examples of animal feed poultry diets include cereal-based feeds including cereals such as barley, corn, soya, wheat, triticale, and rye. Corn-soybean, wheat-soybean, and wheat-corn-soybean, sorghum-soybean, and corn-sorghum-soybean represent other non-limiting examples of suitable animal feeds according to the present invention. When the
10 poultry diet is a corn-soybean meal feed, the corn-soybean meal feed comprises from about 60 to about 70% corn by weight and from about 20 to about 30% soybean by weight.

The poultry diet can further be categorized as a starter diet, a grower-type diet, or a finisher-type diet. The precise composition and physical characteristics of the
15 animal feed, and thus the poultry diet, will depend upon the species for which the feed is intended, the age and/or weight of the animal, and the duration of feeding, and can be readily determined by those skilled in the art.

According to embodiments of the present invention, the methods of growing meat-type poultry do not require concurrently providing a specific keratin-containing
20 substrate along with the keratinase. For example, in embodiments of the present invention, the keratinase can directly supplement a poultry diet as a feed additive in contrast to producing a hydrolyzed feather meal as described in Carter, 1998. Thus, the animal feed can be essentially free of keratin (*e.g.*, not more than 1 or 2% by weight keratin.)

25 A keratinase suitable for practicing the present invention is obtained from *Bacillus licheniformis* strain PWD-1, which is described in U.S. Pat. Nos. 4,908,220 and 4,959,311. This bacterium was deposited with the American Type Culture Collection (ATCC) in Rockville, MD, USA in accordance with the Budapest Treaty on Mar. 23, 1988, and assigned ATCC Accession No. 53757. Other keratinases that
30 can be used to practice the present invention are available from a variety of bacterial sources, such as *Streptomyces fradiae*. See generally U.S. Pat. No. 2,988,487 to Nickerson; See also Goktan, D., "Decomposition Rates of Keratinous Material Used by Certain Microorganisms," (Abstract No. 207369b), Microbial

Biochem. 101, 333 (1984); Daniels, G., "The Digestion of Human Hair Keratin by *Microsporum Canis*," J. Gen. Microbiol. 8, 289 (1953); Koh, W. et al., "Keratinolytic Enzymes from *Aspergillus flavus* and *Aspergillus niger*," Bacillus. Aust. J. Biol. Sci. 274 (1959); Molyneaux, G. S., "The Digestion of Wool by a Keratinolytic Bacillus," Aust. J. Biol. Sci. 274 (1959); Noval, J. and Nickerson, W., "Decomposition of Native Keratin by *Streptomyces Fradiae*," J. Bacteriol. 77, 251 (1959); Kapica, L. and Blank, F., "Growth of *Candida Parapsilosis* with Keratin as Sole Source of Nitrogen," Dermatologica 117, 433 (1958); Kapica, L. and Blank, F., "Growth of *Albicans* on Keratin as Sole Source of Nitrogen," Dermatologica 115, 81 (1957).

10 Keratinase for practicing the present invention can be obtained by growing a host cell which contains nucleic acid sequences encoding a keratinase, under conditions which permit expression of the encoded keratinase, filtering the medium to remove the cells and collecting and concentrating the remaining supernatant by ultrafiltration to obtain the keratinase. Beneficiary co-factor(s) can also be obtained.

15 While strains of *B. licheniformis* are exemplified herein, it is contemplated that other eukaryotic and prokaryotic microbes containing nucleic acid sequences encoding a keratinase may also be useful in producing an animal feed supplement of the present invention. Eukaryotic and prokaryotic microbes containing nucleic acid sequences encoding a keratinase may include those which naturally produce the enzyme as well as strains genetically modified to express keratinase. In general, recombinant production of a protein may require incorporation of nucleic acid sequences encoding said protein into a recombinant expression vector in a form suitable for expression of the protein in a host cell. A suitable form for expression provides that the recombinant expression vector includes one or more regulatory sequences operatively-linked to the nucleic acids encoding the a keratinase protein in a manner which allows for transcription of the nucleic acids into mRNA and translation of the mRNA into the protein. Regulatory sequences may include promoters, enhancers and other expression control elements (e.g., polyadenylation signals). Such regulatory sequences are known to those skilled in the art and are described in Goeddel D.D., ed., Gene Expression Technology, Academic Press, San Diego, CA (1991). It should be understood that the design of the expression vector may depend on such factors as the choice of the host cell to be transfected and/or the level of expression required. Nucleic acid sequences or expression vectors harboring nucleic acid sequences encoding a keratinase protein may be introduced into a host

cell, which may be of eukaryotic or prokaryotic origin, by standard techniques for transforming cells. Suitable methods for transforming host cells may be found in Sambrook, *et al.* (Molecular Cloning: A Laboratory Manual, 3rd Edition, Cold Spring Harbor Laboratory Press (2000)) and other laboratory manuals. The number of host
 5 cells transformed with a nucleic acid sequence encoding a keratinase protein will depend, at least in part, upon the type of recombinant expression vector used and the type of transformation technique used. Nucleic acids may be introduced into a host cell transiently, or more typically, for long-term expression of a keratinase protein the nucleic acid sequence is stably integrated into the genome of the host cell or remains
 10 as a stable episome in the host cell. Once produced, a keratinase protein may be recovered from culture medium as a secreted polypeptide, although it also may be recovered from host cell lysates when directly expressed without a secretory signal.

Eukaryotic microbes such as yeast cultures may be transformed with vectors carrying nucleic acid sequences encoding a keratinase. See, *e.g.*, U.S. Patent No.
 15 4,745,057. *Saccharomyces cerevisiae* is the most commonly used among lower eukaryotic host microorganisms, although a number of other strains are commonly available. Yeast vectors may contain an origin of replication from the 2 micron yeast plasmid or an autonomously replicating sequence (ARS), a promoter, DNA encoding a keratinase such as that provided in U.S. Patent No. 5,712,147, sequences for
 20 polyadenylation, and transcription termination, and a selection gene. An exemplary plasmid is YRp7, (Stinchcomb, *et al.* (1979) *Nature* 282:39; Kingsman, *et al.* (1979) *Gene* 7:141; Tschemper, *et al.* (1980) *Gene* 10:157). Suitable promoting sequences in yeast vectors include the promoters for metallothionein, 3-phosphoglycerate kinase (Hitzeman, *et al.* (1980) *J. Biol. Chem.* 255:2073) or other glycolytic enzymes (Hess,
 25 *et al.* (1968) *J. Adv. Enzyme Reg.* 7:149; Holland, *et al.* (1978) *Biochemistry* 17:4900). Suitable vectors and promoters for use in yeast expression are further described in EPO Publication. No. 73,657. Further, fungal strains such as of *Trichoderma* (*e.g.*, *T. longibrachiatum*, *T. reesei* or *T. viride*) are particularly useful in expressing secreted enzymes.

30 Prokaryote host cells which may be used to produce a keratinase include gram negative or gram-positive organisms, for example *Escherichia coli* (*E. coli*) or *Bacilli*. Exemplary host cells are *E. coli* W3110 (ATCC 27,325), *E. coli* B, *E. coli* X1776 (ATCC 31,537), *E. coli* 294 (ATCC 31,446). A broad variety of suitable prokaryotic and microbial vectors are available. *E. coli* is typically transformed using pBR322.

Promoters most commonly used in recombinant microbial expression vectors include the beta-lactamase (penicillinase) and lactose promoter systems (Chang, et al. (1978) *Nature* 275:615; Goeddel, et al. (1979) *Nature* 281:544), a tryptophan (trp) promoter system (Goeddel, et al. (1980) *Nucleic Acids Res.* 8:4057; EPO Publication No. 36,776) and the tac promoter (De Boer, et al. (1983) *Proc. Natl. Acad. Sci. USA* 80:21). The promoter and Shine-Delgarno sequence (for prokaryotic host expression) are operable linked to the DNA encoding the keratinase, *i.e.*, they are positioned so as to promote transcription of keratinase messenger RNA from the DNA. A species of *Bacillus* is preferably used in the production of a keratinase. Recombinant expression vectors for *Bacillus* are well known to those of skill in the art. *Bacillus* strains can be *B. alkalophilus*, *B. amyloliquefaciens*, *B. brevis*, *B. circulans*, *B. coagulans*, *B. firmus*, *B. lautus*, *B. lentus*, *B. licheniformis*, *B. megaterium*, *B. pumilus*, *B. stearothermophilus*, *B. subtilis*, and *B. thuringiensis*. In a preferred embodiment, strains of *B. licheniformis* are utilized. In some embodiments, *B. licheniformis* strains T399D or PWD-1 are utilized.

As provided herein, a keratinase enzyme may be produced by culturing a host cell as described above under conditions that permit expression of the encoded keratinase, and collecting the expressed keratinase. The host cell may be cultured under conditions in which the cell grows, and then cultured under conditions which cause the expression of the encoded keratinase, or the cells may be caused to grow and express the encoded keratinase at the same time. Such conditions are well known to one of skill in the art and may vary with the host cell and the amount of enzyme expression level desired.

In some embodiments, the medium used to cultivate the transformed host cells may be any medium suitable for keratinase production. The keratinase is recovered from the medium by conventional techniques including separation of the cells from the medium by centrifugation, or filtration, and concentration of the proteins in the supernatant or filtrate by ultrafiltration or evaporation followed by drying via lyophilization or spray-drying.

Alternatively, the culture supernatant may be spray-dried or lyophilized after separation without being concentrated.

The keratinase should be present in an amount at least sufficient to achieve the intended effect, but the upper limit to the amount of keratinase can be determined based upon achieving the intended effect. In some embodiments, the animal feed

comprises from about 0.01% to about 20% *Bacillus licheniformis* PWD-1 keratinase by weight. Additionally, keratinases used in practicing the present invention can be in crude form or in pure form. Keratinases in crude form can be prepared, for example, by separating bacterial cells which produce the keratinase from their liquid growth media, the liquid growth media comprising crude keratinase. Alternatively, the cells can be lysed (chemically or physically) in a liquid growth media to produce a crude, cell free extract. Other means of preparing such an extract will be apparent to persons skilled in the art. The crude keratinase can be included in the feed in any form compatible therewith, such as in an aqueous form or in lyophilized form. In some embodiments, the crude keratinase is in the lyophilized form.

Pure (or substantially pure) keratinases can be obtained by separating the crude keratinase described above into its individual constituents, in accordance with known techniques. *See generally* W. Jakoby, Ed., *Enzyme purification and Related Techniques, Methods in Enzymology*, vol. 22 (1971) and vol. 104, pt. C (1984), Academic Press, NY. Numerous suitable separation procedures, such as column chromatography, are known to persons skilled in the art. The individual constituent proteins can be screened for their ability to degrade keratinaceous material, and that constituent which best degrades keratinaceous material comprises the keratinase. Like the crude keratinase, the pure keratinase can be employed in any suitable form, including aqueous form and lyophilized form.

Embodiments of the present invention further relate to methods of improving the efficiency of feed utilization of an animal feed in meat-type poultry comprising feeding meat-type poultry an animal feed poultry diet wherein the feed further comprises keratinase in an amount effective to improve the efficiency of feed utilization of an animal feed provided to meat-type poultry. The animal feed can include the animal feeds as described above and, in particular embodiments can be corn-soybean meal. The keratinase can include keratinases as described above including, but not limited to, *Bacillus licheniformis* PWD-1 keratinase. As described above, the keratinase can be a crude extract or pure form enzyme.

Improving the efficiency of feed utilization refers to a reduction in the Feed Conversion Ratio (FCR) as compared with that which would otherwise occur without implementation of the methods and/or administration of the compositions of the present invention. The FCR is the ratio of the amount of feed consumed relative to the weight gain of an animal. In one embodiment of the present invention, the

improved efficiency of feed utilization can occur by increasing gastrointestinal nutrient absorption without a concomitant increase in intestinal energy expenditure. In another embodiment of the present invention, the improved efficiency of feed utilization can occur by increasing the digestibility of the animal feed. In another
5 embodiment of the present invention, the improved efficiency of feed utilization can occur by decreasing the viscosity of the animal feed. In particular embodiments, the present invention relates to methods of increasing the digestibility of an animal feed in a meat-type poultry comprising feeding meat-type poultry an animal feed poultry diet wherein the feed further comprises *Bacillus licheniformis* PWD-1 keratinase in an
10 amount effective to increase the digestibility of an animal feed in meat-type poultry. The animal feed can include the animal feeds as described above and, in particular embodiments, can be corn-soybean meal. The keratinase can include keratinases as described above including, but not limited to, *Bacillus licheniformis* PWD-1 keratinase. As described above, the keratinase can be a crude extract or pure form
15 enzyme. Increasing the digestibility of an animal feed refers to increasing the availability of nutrients absorbed from the animal's gut without a concurrent increase in feed intake or nutrient ingestion. In some embodiments of the present invention, the viscosity of materials present in the animal's gut or digesta viscosity is reduced. In other embodiments, the entrapment of nutrients rendering them nutritionally
20 unavailable to the animal is reduced.

In other embodiments, the present invention relates to methods of reducing mortality in meat-type poultry comprising feeding meat-type poultry an animal feed poultry diet wherein the feed further comprises a keratinase in an amount effective to reduce the mortality of meat-type poultry, for example immature birds, and more
25 specifically, broiler chicks. The animal feed can include the animal feeds as described above and, in particular embodiments, can be corn-soybean meal. The keratinase can include keratinases as described above including, but not limited to, *Bacillus licheniformis* PWD-1 keratinase. As described above, the keratinase can be a crude extract or pure form enzyme. Reducing mortality refers to increasing the
30 survivability or decreasing the death rate in animals after birth or hatch as compared with that which would otherwise occur in the absence of implementation of the methods and/or administration of the compositions of the present invention. Mortality can be from any cause, in particular, stress, stunting, "starveouts", and disease. In some embodiments, the present invention reduces mortality in immature birds. In

other embodiments, the birds are about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 31, 32, 33, 34, or 35 days old, preferably about 1 to about 21 days old, and more preferably about 1 to about 5 days old.

5 In some embodiments, the present invention relates to an animal feed comprising protein, carbohydrate, and keratinase as the major components. The keratinase is a major component supplementing the animal feed. The animal feed can include the animal feeds as described above and, in particular embodiments, can be corn-soybean meal. The keratinase can include keratinases as described above
10 including, but not limited to, *Bacillus licheniformis* PWD-1 keratinase. As described above, the keratinase can be a crude extract or pure form enzyme.

The animal feed supplement provided by the present invention can be mixed directly with the animal feed, such as one comprising barley, to prepare the final feed. Alternatively, the animal feed supplement can be mixed with one or more other
15 animal feed supplements such as a vitamin animal feed supplement, a mineral animal feed supplement and an amino acid animal feed supplement. The resulting animal feed supplement including several different types of components may then be mixed in an appropriate amount with the animal feed.

The animal feed of the present invention comprises keratinase in an amount at
20 least sufficient to achieve the intended effect, wherein the upper limit to the amount of keratinase can be determined based upon achieving the intended effect. Intended effects include, but are not limited to, enhancing animal growth performance, such as weight gain, improving the efficiency of feed utilization, increasing feed digestibility, and decreasing mortality. The animal feed supplement added to the animal feed can
25 comprise up to 100% keratinase by weight. The animal feed comprising the supplement comprises from about 5% to about 25% keratinase by weight. In some embodiments, the keratinase is *Bacillus licheniformis* PWD-1 keratinase.

Any animal is a suitable subject for the present invention, including cows, sheep, pigs, cats, dogs, ferrets, and avians, however, the present invention is
30 preferably employed with monogastric animals. Suitable subjects can be of any age range including neonatal animals, developing animals, and mature animals. In some embodiments, the suitable subject can be an avian, preferably a chicken, and more preferably a broiler chick. In other embodiments the suitable subject can be a chicken. In still other embodiments, the suitable subject can be an immature,

developing, or mature bird. In other embodiments, the suitable subject can be a chicken that is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, or 65 days old, or within any range of these numbers. Thus, the present invention provides a variety of different feeds, including pet feed, poultry feed, and pig feed.

The animal feed supplement of the present invention can also enable a conventional animal feed to be modified by reducing its energy, and/or protein, and/or amino acid content while simultaneously maintaining the same nutritional levels of energy, protein, and amino acids available to the animal. Consequently, the amounts of costly energy and protein supplements typically included in an animal feed can be reduced as compared to conventional feeds.

The following Examples are provided to illustrate the present invention, and should not be construed as limiting thereof.

15

EXAMPLE 1

Production of Keratinase From Recombinant *B. licheniformis* PWD-1 Strain

A fermentation scale-up strategy was designed for the production of keratinase, using the wild-type *B. licheniformis* strain PWD-1.

Flask Culture in LB Medium. Flask culture was carried out in Luria-Bertani (LB) medium that was prepared according to the manufacturer's specification, containing: 1.0 L of distilled water, 15 g BACTO[®] agar, 10 g NaCl, 10 g BACTO[®] tryptone, and 5.0 g yeast extract. *B. licheniformis* strain PWD-1 was streaked from a glycerol stock onto an LB plate and grown at 50°C for 8-12 hours. A single colony of *B. licheniformis* strain PWD-1 was then transferred from the LB plate into a flask that contained 500 ml LB medium, and grown at 50°C for 6 hours.

Seed Cultures. Seed cultures for *B. licheniformis* strain PWD-1 were conducted in a medium containing: 0.7 g/L KH₂PO₄, 1.4 g/L K₂HPO₄, 0.1 g/L MgSO₄•7H₂O, 10 g/L defatted soy flour, and 0.1 g/L antifoaming agent. The initial pH of the seed culture was adjusted to 7.0 by adding 1M HCl or NaOH.

The 500 ml flask culture was transferred into a first stage seed fermentor of about 10 L to 20 L that contained the seed culture medium, and was grown therein at

50°C for 8-12 hours to reach 2.5% to 5% inoculum size. The first stage seed culture was then transferred to a second stage seed fermentor of 100 L, 250 L or 800 L, and was grown therein at 50°C for 8 hours and then shifted to 37°C

For the seed culture, the cell density reached at least 3×10^8 CFU/mL at about 8 or 10 hours of the culture process.

Production Media. The production culture medium used for *B. licheniformis* strain PWD-1 contained 0.7 g/L KH_2PO_4 , 1.4 g/L K_2HPO_4 , 0.1 g/L $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 10 g/L defatted soy flour and 0.1 g/L antifoaming agent. The initial pH of the production culture was adjusted to 7.0 by adding 1M HCl or NaOH.

The second stage seed culture was transferred to a production fermentor that contained the production culture medium for final stage culturing. The final stage culture was carried out at 50°C for 8 hours, reaching a total culturing time of about 24 to 30 hours before harvesting.

During the above culturing steps, the initial pH of the culture medium was adjusted to 7.0, but no pH control during the culture process was provided. The optimal level of dissolved oxygen was about 20% for *B. licheniformis* strain PWD-1. The inoculum size was about 2.5 to 5%, and the inoculum age was about 8-12 hours.

For the production culture, the peak cell density reached 1.2×10^9 CFU/mL at about 20 or 24 hours of the culture process. The peak enzyme activity, as measured by azocasein assay, reached 35-40 A_{450} per mL at about 24 to 30 hours of the culture process. The pH value of the production culture medium changed from 7.0 to 8.3, but the enzyme activity and productivity stayed at high levels, which indicated that no pH control was necessary.

Recovery and Downstream Processing. The enzyme activity in the production culture was checked before harvesting. The culture supernatant was separated from the cell mass via centrifuge, and then concentrated via ultrafiltration or evaporation. The concentrated liquid enzyme was then spray-dried.

Alternatively, the culture supernatant was directly spray-dried after separation from the cell mass, without being concentrated.

Enzyme Yield and Enzyme Activity. For 100 L production culture, the enzyme activity measured by azocasein assay before harvesting was 3,000 to 3,500 U/mL, and the cell number was 1.3×10^9 CFU/mL. The total dry weight of the 100 L production culture was 9.12 g/L, including 2.15 g/L insoluble dry weight and 6.88 g/L soluble dry weight.

The crude enzyme yield was about 1.75-2.0 g/L. The crude enzyme was prepared by concentration of the fermentation supernatant via Pellicon filtration with a 5 kDa molecular weight cut-off, and then freeze dried. The enzyme activity of the crude dry enzyme was about 1,000,000 to about 1,400,000 U/g, as measured by the azocasein assay. The total protein content of the crude dry enzyme was about 30-36%, of which approximately 14-20% consisted of pure keratinase.

EXAMPLE 2

Production of Keratinase From Recombinant *B. licheniformis* T399D Strain

10 A fermentation scale-up strategy was designed for the production of keratinase, using a recombinant *Bacillus licheniformis* T399D strain (hereinafter the “*Bacillus licheniformis* strain T1”).

Flask Culture in LB Medium. Flask culture was carried out in LB medium that was prepared according to the manufacturer’s specification, containing: 1.0 L of distilled water, 15 g BACTO® agar, 10 g NaCl, 10 g BACTO® tryptone, and 5.0 g yeast extract. *B. licheniformis* strain T1 was streaked from a glycerol stock onto LB plates and grown at 37°C for 18 hours. A single colony of *B. licheniformis* strain T1 was then transferred from the LB plate into a flask that contained 500 mL LB medium, and grown at 37°C for 6 hours. Cell growth was monitored by measuring the optical density at 660 nm (Beckman DU Series 660 Spectrophotometer, Fullerton, CA). After 6 hours of growth, the OD₆₆₀ measured above 1.0.

Seed Cultures. Seed cultures for *B. licheniformis* strain T1 were grown in a medium containing: 0.7 g/L KH₂PO₄, 1.4 g/L K₂HPO₄, 0.1 g/L MgSO₄•7H₂O, 10 g/L defatted soy flour, and 0.1 g/L antifoaming agent. The initial pH of the seed culture was adjusted to 7.0 by adding 1 M HCl or NaOH.

The 500 mL flask culture was transferred into a first stage seed fermentor of about 10 L to 20 L that contained the seed culture medium, and was grown therein at 37°C for 8 hours to reach 2.5% to 5% inoculum size. The first stage seed culture was then transferred to a second stage seed fermentor of 100 L, 250 L or 800 L, and was grown therein at 37°C for 8 hours.

Production Media. The production culture medium used for *B. licheniformis* strain T1 contained 0.7 g/L KH₂PO₄, 1.4 g/L K₂HPO₄, 0.1 g/L MgSO₄•7H₂O, 13 g/L defatted soy flour, 40 g/L starch, 13 g/L feather meal, and 0.1 g/L antifoaming agent.

The initial pH of the production culture was adjusted to 7.0 by adding 1 M HCl or NaOH.

The second stage seed culture was transferred to a production fermentor that contained the production culture medium for final stage culturing. The final stage culture was carried out at 37°C for 48 hours before harvesting.

During the above culturing steps, the initial pH of the culture medium was adjusted to 7.0, but no pH control was provided. The optimal level of dissolved oxygen was about 30% for *B. licheniformis* strain T1. The inoculum size was about 2.5 to 5%, and the inoculum age was about 12 hours.

Recovery and Downstream Processing. Enzyme activity in the production culture was checked before harvesting. The culture supernatant was separated from the cell mass via centrifugation, and then concentrated via ultrafiltration or evaporation. The concentrated liquid enzyme was then spray-dried.

Alternatively, the culture supernatant was directly spray-dried after separation from the cell mass, without being concentrated.

Enzyme Yield and Enzyme Activity. For 100 L production culture, the enzyme activity measured by azocasein assay before harvesting was 30,000 to 35,000 U/mL, and the cell number was 6×10^9 CFU/mL. The total dry weight of the 100 L production culture was 40 g/L, including 15 g/L insoluble dry weight and 25 g/L soluble dry weight.

The crude enzyme yield from the directly dried culture supernatant was 20 g/L, while the crude enzyme yield from a culture concentrate, as obtained via Pellicon filtration with a 10 kDa molecular weight cut-off, was 16 g/L. The enzyme activity of the crude dry enzyme was greater than 1,000,000 U/g, as measured by the azocasein assay.

EXAMPLE 3

Materials and Methods of Supplementation of Poultry Feed with Keratinase

Birds and Housing. Three experiments were conducted. In each experiment, 192 day-old broiler chicks were weighed and randomly assigned to 24 cage-pens in a completely randomized design to two Alternate Design batteries (Wilvec, Billerica, MA). Birds were weighed, wing-banded and introduced to the experimental treatments at five (experiments one and two) or one (experiment three) day of age.

Each treatment was replicated five times with eight birds per pen except for the control treatment that was replicated four times with eight birds per pen. The birds were housed in a room with controlled temperature, ventilation, and lighting (24 hours/day). During the experimental period the birds were feed *ad libitum* in trough
 5 feeders and water by nipple drinkers.

The PWD-1 Keratinase Enzyme. The enzyme, PWD-1 keratinase was produced with a 150-L fermentor using standard methods (Wang and Shih (1998) *J. Indust. Microb. Biotech.* 22:608-616). Briefly, *Bacillus licheniformis* PWD-1 (Williams, et al. (1990) *supra*) was grown in the fermentor at 50°C for 48 hours. The cell-free
 10 media were concentrated by membrane ultrafiltration and dried by a freeze-dryer. Typically, the yield of the crude enzyme was 2.0 g/L. The crude keratinase had an activity of 300,000 U/g as measured by the hydrolysis of azo-keratin (Lin, et al. (1992) *supra*).

Dietary Treatments. All diets were formulated using least-cost linear
 15 programming software and are presented in **Table 1**.

TABLE 1

Ingredient	Dietary treatment		
	High protein ¹	Control ²	Low Protein
Corn	49.60	59.00	49.00
Soybean meal, 48% CP	41.44	32.00	26.60
Limestone	1.32	1.40	1.32
Dical phosphate	1.75	1.70	1.82
Poultry fat	5.34	5.00	4.20
DL-Methionine	0.15	0.16	0.13
Salt	0.40	0.50	0.42
Choline chloride	0.10	0.10	0.08
Minerals ³ (TM-90)	0.12	0.12	0.10
Vitamins ⁴ (NCSU-90)	0.07	0.07	0.06
Selenium premix ⁵	0.07	0.07	0.06
Sodium Bicarbonate	0.10	0.10	0.08
Starch, corn	0	0	16.60
Total (kg)	100.46	100.22	100.47

	Analysis ⁶	% NRC	Analysis	% NRC	Analysis	% NRC
Crude protein, %	25.00	108.70	20.23	88.00	16.80	72.90
ME, kcal/kg	3,050	95.31	3,201	100.00	3,257	101.80
Met + Cys, %	0.933	103.67	0.85	99.00	0.70	77.90
Lysine, %	1.437	130.60	1.17	106.30	0.97	88.10
Calcium, %	1.00	100.00	1.03	103.30	1.01	100.10
Available Phosphate, %	0.45	100.00	0.45	100.70	0.45	100.70

¹ Provided to birds *ad libitum* for the first 5 days of age in experiment three only.

² Provided to birds *ad libitum* for the first 5 days of age in experiments one and two. In all experiments, birds on the control treatment continued to receive the same diet after the first 5 days of age while other birds were subjected to corresponding treatments.

³ The mineral premix was obtained from Eastern Minerals, Inc., Henderson, NC and provided the following (per kg of diet): 120 mg Zn from ZnSO₄; 120 mg Mn from MnSO₄; 80 mg Fe from FeSO₄C5H2O; 10 mg Cu from CuSO₄; 2.5 mg I from CaIO₄; and 1 mg Co from CoSO₄.

⁴ The vitamin premix was obtained from Roche, Nutley, NJ and provided the following (per kg of diet): 13,200 IU vitamin A; 4,000 ICU vitamin D; 66 IU vitamin E; 39.6 Fg vitamin B₁₂; 13.2 mg riboflavin; 110 mg niacin; 22 mg d-pantothenate; 0.4 mg vitamin K; 2.2 mg folic acid; 4.0 mg thiamin; 7.9 mg pyridoxine; 0.253 mg biotin; 100 mg ethoxyquin.

⁵ The selenium premix provided 0.2 mg Se/kg diet as Na₂SeO₃.

⁶ Calculated analysis.

All feed was fed in mash form throughout the experiments. Birds received a basal diet at one day of age and were subsequently switched to the corresponding experimental diets at five days of age. In experiment one, the basal diet fed for the first five days was about 93% of the National Research Council's (NRC) recommendations for crude protein ((1994) *supra*) but provided 100% for essential amino acids, energy, and calcium and phosphorus contents. In experiments two and three, the basal diet fed for the first 5 days was about 95% of the NRC recommendations ((1994) *supra*) for energy and 100% of the calcium and phosphorus but provided 105% crude protein content. Subsequently, a pen of broiler chicks was subjected to one of five dietary treatments through to the end of each experiment (21 days in experiments one and three, 26 days in experiment two). The five dietary treatments in experiment one and two were: 1) unsupplemented control diet (C, 21.39% crude protein); 2) low protein diet (LP, 18% crude protein); 3) low protein diet supplemented with 0.05% (wt/wt) enzyme preparation (LP+0.05E); 4) low protein diet supplemented with 0.10% (wt/wt) enzyme preparation (LP+0.10E); and

5) low protein diet supplemented with 0.15% (wt/wt) enzyme preparation (LP+0.15E). The control diet was the same basal diet fed to the birds for the first five days of age in experiment one. Birds on treatment one continued to receive the same diet after five days of age, while the rest of the treatments were switched to the experimental diets at five days of age. The dietary treatments in experiment three were: 1) unsupplemented control (C, 21.39% crude protein); 2) control diet supplemented with 0.10% (wt/wt) enzyme preparation (C+0.10E); 3) low protein diet (LP, 18% crude protein); 4) low protein diet supplemented with 0.10% (wt/wt) enzyme preparation (LP+0.10E); and 5) same as treatment two but fed to birds starting at one day of age rather than five days of age.

The enzyme dosage was dissolved in 0.10 *N* sodium carbonate solution in a ratio of 1 gram enzyme/10 ml solution prior to feed application. Thereafter, the enzyme solution was sprayed on top of the feed using a spray bottle in a ratio of 10 ml enzyme preparation/kg diet and mixed using a small bowl mixer (The Hobart Manufacturing Company, Troy, OH).

Viscosity of Digestae. At the end of each experiment, all of the birds but two per pen were euthanized using CO₂ gas. The two remaining birds per pen were kept for the next day. Early the next day, feeders were removed from the pens to produce a 2-hour period with no access to feed. After the 2 hours, birds were once again allowed access to the feed *ad libitum*. An hour later, the birds were removed, two at a time, and were subsequently euthanized using CO₂ gas. Necropsy was carried out immediately after euthanization, and the jejunal contents were emptied into 1-ml Eppendorf tubes. Two samples were obtained per bird. The tubes were immediately centrifuged at 12,000 x *g* for 5 minutes and immediately placed in ice until viscosity was measured using a commercial type viscometer (Brookfield Digital Viscometer, Model DV-II Version 2.0, Brookfield Engineering Laboratories, Inc., Stoughton, MA). Viscosity reading was conducted under conditions that avoided any bacterial growth in the solution.

Data Analysis. Body weights and feed consumption were recorded at 5-day intervals starting day one of age and through the end of each experiment. Feed conversion ratio (feed-to-weight gain), corrected for mortality and culls, was calculated. Mortality was recorded daily. Body weight, feed consumption, feed conversion ratio, and viscosity readings of each experiment were analyzed separately using one-way analysis of the general linear model procedures of SAS software (SAS

Institute (1996) SAS/STAT User's Guide: Statistics. Release 6.11. SAS Institute, Inc., Cary, NC). The percentage data were subjected to ANOVA after arcsine square root percentage transformation. The means were separated using least significant difference. Statements of significance were based on $P \leq 0.05$.

5

EXAMPLE 4

Supplementation of Poultry Feed with Keratinase: Experiment 1

Final body weight, cumulative feed consumption, and feed conversion ratios are presented in Table 2.

10

TABLE 2

Treatment	Body Weight (g)	Feed Consumption (g)	Feed Conversion ratio
Control (C)	709 ± 16	901 ^a ± 26	1.56 ± 0.05
Low protein (LP)	668 ± 14	835 ^{ab} ± 23	1.57 ± 0.04
LP + 0.05 E	691 ± 14	860 ^{ab} ± 23	1.56 ± 0.04
LP + 0.10 E	700 ± 14	847 ^{ab} ± 23	1.51 ± 0.04
LP + 0.15 E	677 ± 14	826 ^b ± 23	1.54 ± 0.04

^{a,b} Means within a column with differing superscripts differ significantly ($P < 0.05$) according to the least squares means function of SAS software (SAS Institute (1996) *supra*).

15 ¹ Values represent means of four to five pens of eight broiler chicks per pen. Values represent means ± standard error of the mean.

² E = enzyme.

The enzyme treatments generally improved body weight with a probability value of $P > 0.05$. The low protein + 0.10% enzyme treatment had higher body weight than the low protein treatment (700 vs. 668 grams for low protein + 0.10% enzyme vs. low protein, respectively, $P = 0.08$). The low protein + 0.10% enzyme treatment had the highest body weight among the enzyme treatments and was not different from the control treatment (700 vs. 709 grams for the low protein + 0.10% enzyme vs. control, respectively).

There were no significant differences among the treatments in feed consumption except between the low protein + 0.15% enzyme and the control treatment (826 vs. 901 grams for the low protein + 0.15% enzyme vs. control,

respectively, $P < 0.05$). There were no significant differences in feed consumption among the enzyme treatment groups.

There was only one dead bird throughout the whole experiment. The weight of this dead bird and the culls were included in the calculation of feed conversion ratio, which is presented in Table 2. Dietary enzyme supplementation had marginal effects on feed conversion ratio. On a cumulative basis, the low protein + 0.10% enzyme treatment had the lowest feed conversion ratio (1.51 vs. 1.57 for the low protein + 0.10% enzyme vs. low protein, respectively, $P > 0.05$).

The results of the first experiment revealed a trend in the response for the enzyme treatment. To further analyze the positive effect of the enzyme for a longer time, a second experiment was conducted growing the birds 5 more days (to 26 days of age).

EXAMPLE 5

Supplementation of Poultry Feed with Keratinase: Experiment 2

This experiment was a repeat of experiment 1 except that birds were grown 5 more days (to 26 days of age). Final body weight, cumulative feed consumption, and feed conversion ratio are presented in Table 3.

TABLE 3

Treatment	Body Weight (g)	Feed Consumption (g)	Feed Conversion ratio
Control (C)	1089 ^a ± 15	1717 ^c ± 16	1.83 ^a ± 0.04
Low protein (LP)	964 ^c ± 13	1734 ^b ± 14	2.14 ^c ± 0.04
LP + 0.05 E	1019 ^b ± 13	1796 ^a ± 14	2.08 ^{bc} ± 0.04
LP + 0.10 E	1025 ^b ± 13	1764 ^{ab} ± 14	2.02 ^b ± 0.04
LP + 0.15 E	1032 ^b ± 13	1794 ^a ± 14	2.04 ^{bc} ± 0.04

^{a,b,c} Means within a column with differing superscripts differ significantly ($P < 0.05$) according to the least squares means function of SAS software (SAS Institute (1996) *supra*).

¹ Values represent means of four to five pens of eight broiler chicks per pen. Values represent means ± standard error of the mean.

² E = enzyme.

There was an improvement in body weight ($P < 0.05$) upon supplementing the low protein diet with all three levels of the enzyme at 26 days of age. The low protein + 0.10% enzyme treatment and the low protein + 0.15% enzyme treatment gave the highest body weight improvement (1,032 and 1,025 vs. 964 grams for the low protein

+ 0.15% enzyme and low protein + 0.10 vs. low protein, respectively, $P < 0.05$). However, all the enzyme treatments had lower ($P < 0.05$) body weight than the control treatment (1,032, 1,025, and 1,016 vs. 1,089 grams for the low protein + 0.15% enzyme, low protein + 0.10% enzyme and the low protein + 0.05% enzyme vs. control, respectively).

All birds receiving the low protein diet consumed more feed than the control group ($P < 0.05$). Enzyme treatment groups also consumed more feed than the low protein diet ($P < 0.05$; Table 2). There were no significant differences in feed consumption among the enzyme treatments. Enzyme supplementation at 0.05 and 0.15% levels resulted in numerically better feed conversion ratios than the low protein treatment, while the low protein + 0.10% enzyme treatment showed significantly ($P < 0.05$) a better feed conversion ratio than the low protein treatment (2.02 vs. 2.14 for the low protein + 0.10% enzyme vs. low protein, respectively). In this experiment, supplementing the low protein diet with enzyme did not improve the performance of the chicks to a level equivalent to that of the control diet. However, supplementing the low protein diet with the 0.10% enzyme (wt/wt) level did ($P < 0.05$) improve the performance of chicks over that of the low protein diet.

The birds in both experiments 1 and 2 were provided the control diet for the first 5 days of age prior to being subjected to the treatment diets. Although the control diet provided adequate energy, calcium and phosphorus, and essential amino acids, it provided only 93% of the NRC ((1994) *supra*) crude protein recommendation, which made it marginally adequate and sensitive to protease supplementation.

25

EXAMPLE 6

Supplementation of Poultry Feed with Keratinase: Experiment 3

In experiment 3, birds were provided a high protein prestarter diet providing 105% of the NRC ((1994) *supra*) recommendation for crude protein and slightly higher than requirements for all other nutrients except for energy (95% of NRC recommendations; Table 1). Experiment 3 was conducted to determine whether or not the enzyme would continue to exert its effect even after the chicks had received adequate nutrient requirements.

In this experiment, only one level of the enzyme was used (0.10% wt/wt). However, two new treatments were introduced to test the enzyme's ability to exert an

effect upon supplementation to marginally adequate broiler starter diets. The two new treatments consisted of supplementing the same control diet (21.39% crude protein) used in experiments 1 and 2 with 0.10% enzyme (wt/wt) and introducing the treated feed to the chicks either at five days (treatment 2) or one day (treatment 5) of age.

- 5 This provided information whether enzyme supplementation at one day of age would have any further improvements in performance.

Final body weight, cumulative feed consumption, and feed conversion ratio are presented in Table 4.

TABLE 4

Treatment	Body Weight (g)	Feed Consumption (g)	Feed Conversion ratio
Control (C)	695 ^b ± 14	974 ^b ± 18	1.49 ^{ab} ± 0.03
C + 0.10 E	767 ^a ± 13	1046 ^a ± 16	1.45 ^a ± 0.03
Low protein (LP)	651 ^c ± 13	1043 ^a ± 16	1.71 ^c ± 0.03
LP + 0.10% E	679 ^{bc} ± 13	978 ^b ± 16	1.53 ^b ± 0.03
C + 0.10% E ³	764 ^a ± 13	1022 ^{ab} ± 16	1.42 ^a ± 0.03

- 10 ^{a,b,c} Means within a column with differing superscripts differ significantly ($P < 0.05$) according to the least squares means function of SAS software (SAS Institute (1996) *supra*).

¹ Values represent means of four to five pens of eight broiler chicks per pen. Values represent means ± standard error of the mean.

- 15 ² E = enzyme.

³ The enzyme was added at one day of age in this treatment. All others were added at five days of age.

- Although the data in Table 2, Table 3, and Table 4 show only the final body weight numbers of the birds in the different experiments, the birds were weighed every 5 days in each of the experiments. Looking at the 5-day interval numbers for this experiment, it was clear that supplementing the low protein diet with the enzyme in experiment 3 showed similar effect on body weight gain to those of experiments 1 and 2. Supplementing the low protein diet with the enzyme increased the 21-day body weight of the birds, but the effect could not be detected at $P < 0.05$ (679 vs. 651 grams for the low protein + 0.10% enzyme vs. low protein, respectively, $P > 0.05$). However, supplementing the control diet with the enzyme (control + 0.10% enzyme preparation, treatments 2 and 5) showed higher body weight than the control treatment (767 and 764 vs. 695 grams for treatments 2 and 5 vs. control, respectively, $P < 0.05$). Unexpectedly, supplementation of the control diet with the enzyme showed

significantly higher improvements in body weight than when the low protein diet was supplemented with the enzyme whether the enzyme was supplemented at one or five days of age (Table 4). This may have been due to the higher protein and/or amino acids content of the control diet vs. the low protein diet. Keratinase is a broad-spectrum protease enzyme that attacks proteins of different sources and breaks them down into smaller polypeptide components. These polypeptides become easier to degrade by the digestive enzymes in the lumen of the intestines. Higher crude protein and/or amino acids content of the diet (in this case the control diet) means higher substrate content for the enzyme to work on releasing more protein components and making it more available to the young chick which, in turn, will be reflected in higher body weight gain.

EXAMPLE 7

Supplementation of Poultry Feed with Keratinase: Digestae Viscosity

The viscosity readings (mPas) of the jejunal contents of 22-day-old (experiments 1 and 2) and 27-day-old (experiment 3) broiler chicks from all three experiments are presented in Table 5.

TABLE 5

Treatment	Experiment 1	Experiment 2	Experiment 3
Control (C)	3.65 ^b ± 0.42	2.31 ^b ± 0.15	2.55 ^{ab} ± 0.17
C + 0.10 E	----	----	2.18 ^b ± 0.15
C + 0.10 E ³	----	----	1.99 ^b ± 0.15
Low protein (LP)	3.59 ^b ± 0.38	2.36 ^{ab} ± 0.14	2.97 ^a ± 0.15
LP + 0.05 E	2.98 ^{ab} ± 0.38	2.78 ^a ± 0.14	----
LP + 0.10 E	2.88 ^{ab} ± 0.38	2.21 ^{bc} ± 0.14	2.20 ^b ± 0.15
LP + 0.15 E	2.27 ^a ± 0.38	1.98 ^c ± 0.14	----

^{a,b,c} Means within a column with differing superscripts differ significantly ($P < 0.05$) according to the least squares means function of SAS software (SAS Institute (1996) *supra*).

¹ Values represent means of four to five pens (16-20 numbers). Values represent means ± standard error of the mean.

² E = enzyme.

³ The enzyme was added at one day of age in this treatment. All others were added at five days of age.

Supplementing both the low protein and the control diets in all the experiments with keratinase reduced the viscosity of jejunal contents. The reduction was directly proportional to the level of enzyme supplementation. Supplementing the low protein diet with 0.15% enzyme (low protein + 0.15% enzyme) reduced the viscosity of the jejunal contents in experiments 1 and 2 (2.27 and 1.98 mPas vs. 3.59 and 2.36 mPas for low protein + 0.15% enzyme vs. low protein in experiments 1 and 2, respectively, $P < 0.05$). The low protein + 0.15% enzyme treatment also had a lower jejunal viscosity when compared to the control treatment (2.27 and 1.98 mPas vs. 3.65 and 2.31 mPas for the low protein + 0.15% enzyme vs. control in experiments 1 and 2, respectively, $P < 0.05$).

When supplementing the control diet with keratinase at 5 days of age, jejunal viscosity was also reduced (2.18 mPas vs. 2.55 mPas for the control + 0.10% enzyme [experiment 3, treatment 2] vs. control, respectively, $P > 0.05$). However, the reduction was significant only when the diet supplemented with the enzyme starting at 1 day of age (1.99 mPas vs. 2.55 mPas for the control + 0.10% enzyme [experiment 3, treatment 5] vs. control, respectively, $P < 0.05$).

EXAMPLE 8

Feeding Trials Using Varying Enzymatic Activity

A dry crude enzyme extract produced according to methods described herein is composed primarily of keratinase enzyme, but may also contain other types of compounds, including other enzymes, carbohydrates, non-enzymatic peptides, nucleotide fragments, etc., which have a molecular weight of greater than 5 kDa and are therefore retained upon ultrafiltration.

Additional experiments were conducted to study the correlation between the growth performance of broiler chicks and the keratinase activity of the crude enzyme extracts added to their diets.

Crude enzyme extracts obtained from fermentation production of *B. licheniformis* strains P1, P2, and T399 were used in these studies and the results are provided in Table 1.

TABLE 1

Trial #	Crude Enzyme Extract Type	Enzyme Activity	Inclusion Rate (%)	Growth Performance
---------	---------------------------	-----------------	--------------------	--------------------

		(Unit/g)		
MT 301	P1 strain (in 0.5% soy flour) ¹	1,000,000	0.05	No effect
			0.10	Improvement (0.05<P<0.1)
			0.15	Improvement (0.05<P<0.1)
MT 401	P1 strain (in 0.5% soy flour) ²	1,000,000	0.05	Significant improvement in low protein
			0.10	Significant improvement in low protein
			0.15	Significant improvement in low protein
MT 501¹	P1 strain (in 0.5% soy flour) ¹	1,000,000	0.10	Significant improvement in low protein and control
MT 701	P1 strain (in 0.5% soy flour) ¹	300,000	0.25	Negative effect
	P2 strain (in 0.5% soy flour) ¹	450,000	0.17	Slight improvement
MT 801	P1 strain (inhibited by PMSF) ³	300,000	0.25	No effect
	P1 strain (inactivated by heat) ⁴	300,000	0.25	No effect
MT 901	P2 strain (in 1% soy flour) ¹	450,000	0.10	Significant improvement
	P2 strain (in 2%, 1% soy flour) ¹	800,000	0.10	No effect
	T399 strain (in 2%, 1% soy flour) ¹	<5000	0.10	No effect
KE 202	P2 strain (in 1% soy flour) ¹	600,000	0.10	Improvement in 23% diets
	P2 strain (in 2% soy flour) ¹	600,000	0.10	Improvement in 21 & 23% diets
KE 302	P1 strain ⁵	150,000	0.10	Significant improvement in all
Pilot Test	P1 strain ¹	300,000	0.10	Improvement in 21% diet
	P1 strain ¹	300,000	0.25	Improvement in 21% diet

¹ Enzyme supplement fed from 6 to 21 days of age.

² Enzyme supplement fed from 6 to 27 days of age.

³ >90% enzymatic activity is inhibited.

⁴ >98% enzymatic activity is inhibited.

⁵ Enzyme supplement fed from 1 to 21 days of age.

The foregoing examples are illustrative of the present invention, and are not to be construed as limiting thereof. The invention is described by the following claims, with equivalents of the claims to be included therein.

WHAT IS CLAIMED IS:

1. A method of feeding a pig, comprising:
feeding said pig an animal feed comprising a corn-soybean meal feed as a pig diet, wherein said pig diet is essentially free of keratin, and
said animal feed further comprises *Bacillus licheniformis* PWD-1 keratinase in an amount effective to enhance the weight gain of said pig.
2. The method of claim 1, wherein the pig is a neonatal pig.
3. The method of claim 1, wherein the pig is a developing pig.
4. The method of claim 1, wherein the pig is a mature pig.
5. A method of supplementing an animal feed for a pig, the method consisting essentially of:
feeding said pig a corn-soybean meal feed,
said animal feed comprising *Bacillus licheniformis* PWD-1 keratinase in an amount effective to enhance the weight gain of said pig, wherein said animal feed is essentially free of keratin.
6. The method of claim 5, wherein the pig is a neonatal pig.
7. The method of claim 5, wherein the pig is a developing pig.
8. The method of claim 5, wherein the pig is a mature pig.
9. A method of improving the efficiency of feed utilization of an animal feed in a pig, comprising:
feeding said pig a corn-soybean meal feed as a pig diet, wherein said pig diet is essentially free of keratin, and

said animal feed further comprises *Bacillus licheniformis* PWD-1 keratinase in an amount effective to improve the efficiency of feed utilization of an animal feed in said pig.

10. A method of increasing the digestibility of an animal feed in a pig, comprising:

feeding said pig a corn-soybean meal feed as a pig diet, wherein said pig diet is essentially free of keratin, and

said animal feed further comprises *Bacillus licheniformis* PWD-1 keratinase in an amount effective to increase the digestibility of an animal feed in said pig.

11. A method of reducing mortality in a pig, comprising:

feeding said pig an animal feed comprising corn-soybean meal feed as a pig diet, wherein said pig diet is essentially free of keratin, and

said animal feed further comprises *Bacillus licheniformis* PWD-1 keratinase in an amount effective to reduce mortality of said pig.

12. A pig feed consisting essentially of soybean meal, corn, and keratinase, wherein said pig feed is essentially free of keratin.

13. The pig feed of claim 12, wherein the keratinase is a crude extract or pure enzyme.

14. The pig feed of claim 12, wherein the keratinase is *Bacillus licheniformis* PWD-1 keratinase.

15. Use of pig feed of claim 12, wherein the pig feed is added to a diet of a neonatal pig.

16. Use of pig feed of claim 12, wherein the pig feed is added to a diet of a developing pig.

17. Use of pig feed of claim 12, wherein the pig feed is added to a diet of a mature pig.

18. A poultry feed comprising *Bacillus licheniformis* PWD-1 keratinase, wherein the poultry feed:

- (a) is essentially free of keratin, and
- (b) comprises less than about 23% crude protein.

19. The poultry feed of claim 18, wherein the poultry feed:

- (a) is essentially free of keratin, and
- (b) comprises between 20.23% and about 23% crude protein.

20. The poultry feed of claim 18, wherein the poultry feed:

- (a) is essentially free of keratin, and
- (b) comprises between about 16.80% and 20.23% crude protein.

21. A poultry feed comprising *Bacillus licheniformis* PWD-1 keratinase, wherein the poultry feed:

- (a) is essentially free of keratin, and
- (b) comprises less than about 3201 kcal/kg of metabolizable energy.

22. The poultry feed of claim 21, wherein the poultry feed:

- (a) is essentially free of keratin, and
- (b) comprises between about 3050 kcal/kg to about 3201 kcal/kg of metabolizable energy.

23. A method for improving the digestibility of an animal feed essentially free of keratin, the method comprising contacting the feed with *Bacillus licheniformis* PWD-1 keratinase.

24. A method for improving the feed conversion ratio of an animal feed essentially free of keratin, the method comprising contacting the feed with *Bacillus licheniformis* PWD-1 keratinase.

25. A method for reducing the anti-nutritional content of an animal feed essentially free of keratin, the method comprising contacting the feed with *Bacillus licheniformis* PWD-1 keratinase.

26. A method of increasing the metabolizable energy of an animal feed essentially free of keratin, the method comprising contacting the feed with *Bacillus licheniformis* PWD-1 keratinase.

27. A method of improving the nutritional value of an animal feed essentially free of keratin, the method comprising contacting the feed with *Bacillus licheniformis* PWD-1 keratinase.

28. The animal feed of claim 12 or 18, wherein the animal feed is added to a starter diet, wherein said starter diet has not more than 18% crude protein.

29. The animal feed of claim 12 or 18, wherein the animal feed is added to a starter diet, wherein said starter diet is essentially free of keratin.

30. The animal feed of claim 12 or 18, wherein the animal feed is a grower diet, wherein said grower diet has not more than 18% crude protein.

31. The animal feed of claim 12 or 18, wherein the animal feed is a grower diet, wherein said grower diet is essentially free of keratin.

32. The animal feed of claim 12 or 18, wherein the animal feed is a finisher diet, wherein said finisher diet has not more than 18% crude protein.

33. The animal feed of claim 12 or 18, wherein the animal feed is a finisher diet, wherein said finisher diet is essentially free of keratin.