

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 May 2006 (11.05.2006)

PCT

(10) International Publication Number
WO 2006/049618 A1

(51) International Patent Classification⁷:
1/21, 1/20, C12P 13/00, 07/40

C12N 9/78,

(74) Agent: FLOYD, Linda, Axamethy; E. I. du Pont de Nemours and Company, Legal Patent Records Center, 4417 Lancaster Pike, Wilmington, Delaware 19805 (US).

(21) International Application Number:

PCT/US2004/036352

(22) International Filing Date:

1 November 2004 (01.11.2004)

(25) Filing Language:

English

(26) Publication Language:

English

(71) Applicant (for all designated States except US): E. I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, Delaware 19898 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): PAYNE, Mark, S. [US/US]; 4617 Old Linden Hill Road, Wilmington, Delaware 19808 (US). DICOSIMO, Robert [US/US]; 1607 Masters Way, Chadds Ford, Pennsylvania 19317 (US). GAVAGAN, John, E. [US/US]; 2300 Dorval Road, Wilmington, Delaware 19808 (US). FALLON, Robert, D. [US/US]; 104 Black Oak Drive, Elkton, Maryland 21921 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

[Continued on next page]

(54) Title: NITRILE HYDRATASE AND AMIDASE FROM COMAMONAS TESTOTERONI 5-MGAM-4D



(57) Abstract: The invention relates to the isolation, sequencing, and recombinant expression of genes encoding either a nitrile hydratase (NHase) or amidase (Am) from *Comamonas testosteroni* 5-MGAM-4D, where the NHase is useful for catalyzing the hydration of nitriles to the corresponding amides, and the amidase is useful for hydrolysis of amides to the corresponding carboxylic acids. Also provided are transformed host cells containing polynucleotides for expressing the nitrile hydratase or amidase enzymes from *Comamonas testosteroni* 5-MGAM-4D.



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

TITLE

Nitrile hydratase and Amidase from *Comamonas Testosteroni* 5-MGAM-4D.

5

This application claims the benefit of U.S. Application No. 10/431,966 filed May 8, 2003.

FIELD OF THE INVENTION


10 The present invention relates to the field of molecular biology and methods for the isolation and expression of foreign genes in recombinant microorganisms. More specifically, the invention relates to the isolation, sequencing, and recombinant expression of nucleic acid fragments (genes) encoding either a nitrile hydratase (NHase) or amidase (Am) from
15 *Comamonas testosteroni* 5-MGAM-4D, where the NHase is useful for catalyzing the hydration of nitriles to the corresponding amides, and the amidase is likewise useful for hydrolysis of amides to the corresponding carboxylic acids.

20

BACKGROUND OF THE INVENTION

Nitrile hydratases catalyze the addition of one molecule of water to the nitrile, resulting in the formation of the corresponding amide according to Reaction 1:

Reaction 1



25 Similarly, methods for producing carboxylic acids are known and use microorganisms that produce an enzyme that possesses amidase (Am) activity. In general, amidases convert the amide product of Reaction 1 to the corresponding carboxylic acid plus ammonia according to Reaction 2:

Reaction 2



30 A wide variety of bacterial genera are known to possess a diverse spectrum of nitrile hydratase and amidase activities, including *Rhodococcus*, *Pseudomonas*, *Alcaligenes*, *Arthrobacter*, *Bacillus*, *Bacteridium*, *Brevibacterium*, *Corynebacterium*, and *Micrococcus*

(Martinkova and Kren, *Biocatalysis and Biotransformation*, 20:73-93 (2002); Cowan *et al.*, *Extremophiles*, 2:207-216 (1998)). For example, nitrile hydratase enzymes have been isolated from *Pseudomonas chlororaphis* B23 (Nishiyama *et al.*, *J. Bacteriol.*, 173:2465-2472 (1991)),
5 *Rhodococcus rhodochrous* J1 (Kobayashi *et al.*, *Biochem. Biophys. Acta*, 1129:23-33 (1991)), *Brevibacterium* sp. 312 (Mayaux *et al.*, *J. Bacteriol.*, 172:6764-6773 (1990)), *Rhodococcus* sp. N-774 (Ikehata *et al.*, *Eur. J. Biochem.*, 181:563-570 (1989)), and *Pseudomonas putida* 5B NRRL-18668 (Payne *et al.*, *Biochemistry*, 36:5447-5454 (1997)).

10 Wild-type microorganisms known to possess nitrile hydratase activity have been used to convert nitriles to the corresponding amides. Nagasawa *et al.* (*Appl. Microbiol. Biotechnol.*, 40:189-195 (1993)) have compared three microbial nitrile hydratase catalysts which have been used for commercial production of acrylamide from acrylonitrile; the nitrile
15 hydratase activities of *Brevibacterium* R312 and *Pseudomonas chlororaphis* B23 were not stable above 10 °C, compared to the nitrile hydratase activity of *Rhodococcus rhodochrous* J1. Cowan *et al.* (*supra*) reported that many mesophilic nitrile hydratases are remarkably unstable, having very short enzyme activity half-lives in the growth temperature
20 range of 20-35 °C. In addition to temperature instability, microbial catalysts containing a nitrile hydratase can be susceptible to inactivation by high concentrations of certain substrates such as acrylonitrile. In commercial use, the concentration of acrylonitrile was maintained at 1.5-2 wt % when using *Brevibacterium* R312 and *P. chlororaphis* B23 catalysts,
25 while a concentration of up to 7 wt % was used with *R. rhodochrous* J1 (Nagasawa *et al.*, *supra*). Similarly, Padmakumar and Oriel (*Appl. Biochem. Biotechnol.*, 77-79:671-679 (1999)) reported that *Bacillus* sp. BR449 expresses a thermostable nitrile hydratase, but when used for hydration of acrylonitrile to acrylamide, inactivation of the enzyme
30 occurred at concentration of acrylonitrile of only 2 wt %, making this catalyst unsuitable for commercial applications. Webster *et al.* (*Biotechnology Letters*, 23:95-101 (2001)) compare two *Rhodococcus* isolates as catalysts for ammonium acrylate production (one with only a nitrilase activity, and one with only a combination of nitrile hydratase and
35 amidase activities), and concluded that the catalyst having a combination of nitrile hydratase and amidase activities was less preferred due to (a) difficulty in inducing the two enzymes in the required ratio, (b) the susceptibility of the two enzymes (nitrile hydratase and amidase) to

deactivation by acrylonitrile, and (c) inhibition of the two enzymes by the respective products.

The hydration of aromatic and heteroaromatic nitriles to the corresponding amides has been reported using the nitrile hydratase activity of *Rhodococcus rhodochrous* AJ270 (A. Meth-Cohn and M. Wang, *J. Chem Soc., Perkin Trans. 1*, (8):1099-1104 (1997)), where significant subsequent conversion of the amide to the corresponding acid by amidase was also observed. The nitrile hydratase activity of *Rhodococcus rhodochrous* J1 was used to convert a variety of aromatic and heteroaromatic nitriles to the corresponding amides with 100 % molar conversion (J. Mauger *et al.*, *Tetrahedron*, 45:1347-1354 (1989); J. Mauger *et al.*, *J. Biotechnol.*, 8:87-96 (1988)); an inhibitory affect of certain nitriles on the nitrile hydratase was overcome by maintaining a low concentration of the nitrile over the course of the reaction.

US 20040142447 describes the use of several *Rhodococcus* strains for the conversion of 3-cyanopyridine to nicotinamide, where the *Rhodococcus* strains were relatively stable and had a relatively low Km value for 3-cyanopyridine when compared to previously-reported microbial cell catalysts.

In addition to the use of wild-type organisms, recombinant organisms containing heterologous genes for the expression of nitrile hydratase are also known for the conversion of nitriles. For example, Cerebelaud *et al.* (WO 9504828) teach the isolation and expression in *E. coli* of nitrile hydratase genes isolated from *C. testosteroni*. The transformed hosts effectively convert nitriles to amides, including substrates which consist of one nitrile and one carboxylate group. Endo *et al.* disclose the production of an *E. coli* transformant which expresses the nitrile hydratase of *Rhodococcus* N-771 (US 6,316,242 B1). Similarly, Beppu *et al.*, (EP 5024576) disclose plasmids carrying both nitrile hydratase and amidase genes from *Rhodococcus* capable of transforming *E. coli* where the transformed host is then able to use isobutyronitrile and isobutyramide as enzymatic substrates. A stereoselective nitrile hydratase from *Pseudomonas putida* 5B has been overproduced in *E. coli* (Wu *et al.*, *Appl. Microbiol. Biotechnol.*, 48:704-708 (1997); US 5,811,286).

Genes encoding enzymes having amidase activity have also been cloned, sequenced, and expressed in recombinant organisms. For example, Azza *et al.*, (*FEMS Microbiol. Lett.*, 122:129 (1994)) disclose the

cloning and over-expression in *E. coli* of an amidase gene from *Brevibacterium* sp. R312 under the control of the native promoter. Similarly, Kobayashi *et al.*, (*Eur. J. Biochem.*, 217:327 (1993)) teach the cloning of both a nitrile hydratase and amidase gene from *R. rhodococcus* J1 and their co-expression in *E. coli*. Wu *et al.* (*DNA Cell Biol.*, 17:915-920 (1998); US 6,251,650) report the cloning and overexpressing of a gene for amidase from *Pseudomonas putida* 5B in *E. coli*.

Applicants have previously isolated *Comamonas testosteroni* 5-MGAM-4D (ATCC 55744; US 5,858,736 and US 5,922,589).

10 *Comamonas testosteroni* 5-MGAM-4D has been shown to contain thermally-stable, regiospecific nitrile hydratase (EC 4.2.1.84) and amidase (EC 3.5.1.4) activities useful in the conversion of a variety of nitriles to their corresponding amides and carboxylic acids. Methods illustrating the utility of the *Comamonas testosteroni* 5-MGAM-4D nitrile hydratase and
15 amidase activities have been described previously by the Applicants. These uses include regio-selective preparation of lactams from aliphatic α,ω -dinitriles (US 5,858,736), bioconversion of 3-hydroxy nitriles to 3-hydroxyacids (US 2002/0039770 A1), and bioconversion of methacrylonitrile and acrylonitrile to their corresponding carboxylic acids
20 (US SN10/067,652), hereby incorporated by reference in their entirety. However, the isolation and recombinant expression of the nucleic acid fragments encoding the nitrile hydratase and amidase from *Comamonas testosteroni* 5-MGAM-4D has been elusive.

The problem to be solved is to provide the genes and encoding for
25 the thermally-stable, regio-selective nitrile hydratase and amidase enzymes from *Comamonas testosteroni* 5-MGAM-4D and to provide transformants expressing these catalysts.

Additionally, the development of industrial processes which employ microbial catalysts having nitrile hydratase/amidase activities to efficiently
30 manufacture amides or carboxylic acids has proved difficult. Many methods using enzyme catalysts to prepare these products from the corresponding nitriles do not produce and accumulate the product at a sufficiently high concentration to meet commercial needs, or are subject to enzyme inactivation (requiring a low concentration of nitrile over the
35 course of the reaction) or product inhibition during the course of the reaction.

The additional problem to be solved continues to be the lack of facile microbial catalysts to convert nitriles to the corresponding amides or

acids in a process characterized by high yield, high concentration, and high selectivity, and with the added advantages of low temperature and energy requirements and low waste production when compared to known chemical methods of nitrile hydrolysis. *Comamonas testosteroni* 5-MGAM-4D expresses a thermally-stable, regio-selective nitrile hydratase as well as a thermally-stable amidase. An enzyme catalyst having only the nitrile hydratase activity of *Comamonas testosteroni* 5-MGAM-4D would be highly useful in applications where only the amide product from nitrile hydration is desired.

SUMMARY OF THE INVENTION

The Applicants have isolated and sequenced the genes necessary to express thermally-stable, regio-selective nitrile hydratase and amidase from *Comamonas testosteroni* 5-MGAM-4D. The corresponding amino acid sequences for each enzyme are also disclosed. The invention also encompasses 1) an isolated polynucleotide encoding a polypeptide having at least 98 % identity to a polypeptide alpha-subunit of the nitrile hydratase enzyme from *Comamonas testosteroni* 5-MGAM-4D as represented by SEQ ID No:4; 2) an isolated polynucleotide encoding a polypeptide having at least 95 % identity to a polypeptide beta-subunit of the nitrile hydratase enzyme from *Comamonas testosteroni* 5-MGAM-4D as represented by SEQ ID No:6; and 3) an isolated polynucleotide encoding a polypeptide having amidase activity and having at least 95 % identity to the polypeptide from *Comamonas testosteroni* 5-MGAM-4D as represented by SEQ ID No:17.

The invention further provides a region of the *Comamonas testosteroni* 5-MGAM-4D genome encompassed within a 0.9 kb fragment (SEQ ID NO:9) which encodes a polypeptide (designated "PK7"; and represented by SEQ ID NO:14) that is necessary for optimum activity of the nitrile hydratase enzyme.

Transformants are provided that express either the nitrile hydratase or amidase enzymes separately or that co-express both enzymes. Also provided are methods to produce the nitrile hydratase and amidase catalysts in a recombinant host. The present invention further provides recombinant hosts, transformed with the polynucleotides encoding the amidase and/or the nitrile hydratase in combination with the PK7 accessory protein.

A particular embodiment of the invention is *Escherichia coli* transformed with the nucleic acid sequence represented by SEQ ID NO:11.

The Applicants also provide methods for converting a variety of aliphatic nitriles, aromatic nitriles, heterocyclic aromatic nitriles, unsaturated nitriles, aliphatic dinitriles, and 2-, 3-, or 4-hydroxynitriles to the corresponding amides using a transformed host cell expressing the nitrile hydratase from *Comamonas testosteroni* 5-MGAM-4D. *Comamonas testosteroni* 5-MGAM-4D expresses a thermally-stable, regioselective nitrile hydratase as well as a thermally-stable amidase. The present application describes the preparation of microbial transformants that have only the nitrile hydratase activity of *Comamonas testosteroni* 5-MGAM-4D for use in applications where the production of only the amide product from nitrile hydration would be desirable.

15

BRIEF DESCRIPTION OF THE DRAWINGS, SEQUENCE DESCRIPTIONS, AND BIOLOGICAL DEPOSITS

The invention can be more fully understood from the Figure, the Sequence Listing, the Biological Deposits, and the detailed description that together form this application.

Figure 1 shows the nucleic acid fragments inserted in several plasmids created for recombinant expression of genes cloned from *Comamonas testosteroni* 5-MGAM-4D (ATCC 55744).

The following sequences comply with 37 C.F.R. 1.821-1.825 ("Requirements for Patent Applications Containing Nucleotide Sequences and/or Amino Acid Sequence Disclosures - the Sequence Rules") and are consistent with World Intellectual Property Organization (WIPO) Standard ST.25 (1998) and the sequence listing requirements of the EPO and PCT (Rules 5.2 and 49.5(a-bis), and Section 208 and Annex C of the Administrative Instructions). The symbols and format used for nucleotide and amino acid sequence data comply with the rules set forth in 37 C.F.R. §1.822.

SEQ ID NO:1 is the nucleic acid sequence encoding the α -subunit of a nitrile hydratase from *Pseudomonas putida* 5B (NRRL-18668) used to probe genomic DNA fragments from *Comamonas testosteroni* 5-MGAM-4D (ATCC 55744).

SEQ ID NO:2 is the nucleic acid sequence encoding the β -subunit of a nitrile hydratase from *Pseudomonas putida* 5B (NRRL-18668) used to

probe genomic DNA fragments from *Comamonas testosteroni* 5-MGAM-4D (ATCC 55744).

SEQ ID NO:3 is the nucleic acid sequence encoding the α -subunit of a nitrile hydratase from *Comamonas testosteroni* 5-MGAM-4D.

5 SEQ ID NO:4 is the deduced amino acid sequence for the α -subunit of a nitrile hydratase from *Comamonas testosteroni* 5-MGAM-4D.

SEQ ID NO:5 is the nucleic acid sequence encoding the β -subunit of a nitrile hydratase from *Comamonas testosteroni* 5-MGAM-4D.

10 SEQ ID NO:6 is the deduced amino acid sequence for the β -subunit of a nitrile hydratase from *Comamonas testosteroni* 5-MGAM-4D.

SEQ ID NO:7 is the nucleic acid sequence encoding the α - and β -subunits of the nitrile hydratase from *Comamonas testosteroni* 5-MGAM-4D used in the creation of pSW131.

15 SEQ ID NO:8 is the first of two primers ("Primer 1") useful for amplifying a nucleic acid fragment (SEQ ID NO:7) for creation of plasmid pSW131 and for amplifying a nucleic acid fragment (SEQ ID NO:11) for creation of pSW137.

20 SEQ ID NO:9 is the second of two primers ("Primer 2") useful for amplifying a nucleic acid fragment (SEQ ID NO:7) for creation of plasmid pSW131.

SEQ ID NO:10 is the nucleic acid sequence of a 0.9 kb nucleic acid fragment from *Comamonas testosteroni* 5-MGAM-4D containing a small open reading frame (ORF) which encodes an accessory protein (denoted as "P7K") useful in the expression of active nitrile hydratase.

25 SEQ ID NO:11 is the nucleic acid sequence encoding the α - and β -subunits of the nitrile hydratase plus 0.9 kb of downstream DNA (SEQ ID NO. 10) encoding the accessory protein P7K from *Comamonas testosteroni* 5-MGAM-4D used in the creation of pSW132 and pSW137.

30 SEQ ID NO:12 is the second of two primers ("Primer 3") useful for amplifying a nucleic acid fragment (SEQ ID NO:11) for creation of pSW137 and for amplifying a nucleic acid fragment (SEQ ID NO:23) for creation of pSW136.

35 SEQ ID NO:13 is the nucleic acid sequence encoding the accessory protein P7K, and found within the 0.9 kb downstream DNA sequence (SEQ ID NO:10) from *Comamonas testosteroni* 5-MGAM-4D.

SEQ ID NO:14 is the deduced amino acid sequence for the accessory protein P7K useful in the recombinant expression of *Comamonas testosteroni* 5-MGAM-4D nitrile hydratase.

SEQ ID NO:15 is the nucleic acid sequence of a nucleic acid fragment comprising the first 0.6 kb of the pKP57 insert useful as a probe to identify an amidase from *Comamonas testosteroni* 5-MGAM-4D.

5 SEQ ID NO:16 is the nucleic acid sequence encoding an amidase from *Comamonas testosteroni* 5-MGAM-4D.

SEQ ID NO:17 is the deduced amino acid sequence of an amidase from *Comamonas testosteroni* 5-MGAM-4D.

10 SEQ ID NO:18 is the nucleic acid sequence of 7.4 kb nucleic acid fragment comprising the complete coding sequences for an amidase and a nitrile hydratase and the P7K accessory protein from *Comamonas testosteroni* 5-MGAM-4D.

SEQ ID NO:19 is the first of two primers ("Primer 4") useful for amplifying a nucleic acid fragment encoding an amidase from *Comamonas testosteroni* 5-MGAM-4D for creation of plasmid pKP60.

15 SEQ ID NO:20 is the second of two primers ("Primer 5") useful for amplifying a nucleic acid fragment encoding an amidase from *Comamonas testosteroni* 5-MGAM-4D for creation of plasmid pKP60.

20 SEQ ID NO:21 is the first of two primers ("Primer 6") useful for amplifying a nucleic acid fragment encoding an amidase from *Comamonas testosteroni* 5-MGAM-4D for creation of plasmid pSW133 and for amplifying a nucleic acid fragment (SEQ ID NO:23) for creation of pSW136.

25 SEQ ID NO:22 is the second of two primers ("Primer 7") useful for amplifying a nucleic acid fragment encoding an amidase from *Comamonas testosteroni* 5-MGAM-4D for creation of plasmid pSW133.

SEQ ID NO:23 is the nucleic acid fragment encoding an amidase, a nitrile hydratase (α - and β -subunits), and the accessory protein P7K from *Comamonas testosteroni* 5-MGAM-4D used in the creation of plasmid pSW136.

30 Applicants have made the following biological deposits under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure:

Depositor Identification Reference	Int'l. Depository Designation	Date of Deposit
<i>Comamonas testosteroni</i> 5-MGAM-4D	ATCC 55744	8 March 1996
<i>Pseudomonas putida</i> 5B	NRRL 18668	6 July 1990
<i>Escherichia coli</i> SW132	ATCC PTA-5073	21 March 2003
<i>Escherichia coli</i> SW137	ATCC PTA-5074	21 March 2003

As used herein, "ATCC" refers to the American Type Culture Collection International Depository Authority located at ATCC, 10801 University Blvd., Manassas, VA 20110-2209, USA. The "International Depository Designation" is the accession number to the culture on deposit with ATCC.

As used herein, "NRRL" refers to the Northern Regional Research Laboratory, Agricultural Research Service Culture Collection International Depository Authority located at 11815 N. University Street, Peoria, IL 61604 U.S.A. The "NRRL No." is the accession number to cultures on deposit at the NRRL.

The listed deposits will be maintained in the indicated international depository for at least thirty (30) years and will be made available to the public upon the grant of a patent disclosing it. The availability of a deposit does not constitute a license to practice the subject invention in derogation of patent rights granted by government action.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides isolated polynucleotides and the nucleic acid sequences that encode three polypeptides (α - and β -subunits of a nitrile hydratase and an amidase) from *Comamonas testosteroni* 5-MGAM-4D (ATCC 55744) that act as catalysts. When coexpressed, the α - and β -subunits of the nitrile hydratase selectively hydrate nitriles into the corresponding amides, and the amidase hydrolyzes amides into the corresponding carboxylic acids. The invention also provides transformed microbial host cells expressing the polypeptides. The invention further provides a method for producing the polypeptide catalysts using the transformed microbes and a method for using the catalysts for converting nitriles to the corresponding amides and/or carboxylic acids, or for converting amides to the corresponding carboxylic acids.

Definitions:

In this disclosure, a number of terms and abbreviations are used. The following definitions apply unless specifically stated otherwise.

The terms "catalyst", "enzyme catalyst" or "microbial cell catalyst" refer to polypeptides (or proteins) having a nitrile hydratase activity, an amidase activity, or having a combination of nitrile hydratase and amidase

activities. The catalyst may be in the form of an intact microbial cell, permeabilized microbial cell(s), one or more cell components of a microbial cell extract, partially purified enzyme(s), or purified enzyme(s).

As used herein, the term "comprising" means the presence of the stated features, integers, steps, or components as referred to in the claims, and it does not preclude the presence or addition of one or more other features, integers, steps, components, or groups thereof.

The term "thermally-stable" characterizes an enzyme that retains activity despite exposure to a given temperature.

The terms "*Comamonas testosteroni*" and "*C. testosteroni*" are used interchangeably and refer to *Comamonas testosteroni* 5-MGAM-4D (ATCC 55744).

The terms "*Psuedomonas putida* 5B" and "*P. putida* 5B" are used interchangeably and refer to *Psuedomonas putida* NRRL-18668.

The terms "*Escherichia coli* SW132" and "*E. coli* SW132" are used interchangeably and refer to an *E. coli* strain transformed with plasmid pSW132 and having ATCC accession number PTA-5073.

The term "pSW132" refers to a plasmid containing a DNA fragment encoding the *C. testosteroni* 5-MGAM-4D nitrile hydratase α - and β -subunits plus 0.9 kb of downstream DNA under the control of the T7 promoter. The 0.9 kb of downstream DNA encodes an accessory protein ("P7K") useful in recombinant expression of nitrile hydratase. *E. coli* strain SW132 harbors plasmid pSW132 and has ATCC accession number PTA-5073.

The terms "*Escherichia coli* SW137" and "*E. coli* SW137" are used interchangeably and refer to an *E. coli* strain transformed with plasmid pSW137 and having ATCC accession number PTA-5074.

The term "pSW137" refers to a plasmid containing a DNA fragment encoding the *C. testosteroni* 5-MGAM-4D nitrile hydratase α - and β -subunits plus 0.9 kb of downstream DNA under the control of the *trc* promoter. The 0.9 kb of downstream DNA encodes an accessory protein ("P7K") useful in recombinant expression of nitrile hydratase. *E. coli* strain SW137 harbors plasmid pSW137 and has ATCC accession number PTA-5074.

"Open reading frame" is abbreviated ORF.

"Polymerase chain reaction" is abbreviated PCR.

As used herein, an "isolated nucleic acid fragment" or "isolated polynucleotide" is a polymer of RNA or DNA that is single- or double-stranded, optionally containing synthetic, non-natural, or altered nucleotide bases. An isolated nucleic acid fragment in the form of a polymer of DNA may be comprised of one or more segments of cDNA, genomic DNA, or synthetic DNA.

The term "accessory nucleic acid" refers to the 0.9 kb sequence (SEQ ID NO:10), located downstream of the nitrile hydratase α (alpha) and β (beta) subunit genes, which contains an open reading frame (SEQ ID NO:13) encoding a polypeptide (SEQ ID NO:14) useful in increased recombinant expression of the nitrile hydratase.

The term "complementary" is used to describe the relationship between nucleotide bases that are capable to hybridizing to one another. For example, with respect to DNA, adenosine is complementary to thymine and cytosine is complementary to guanine. Accordingly, the instant invention also includes isolated nucleic acid fragments that are complementary to the complete sequences as reported in the accompanying Sequence Listing.

The term "percent identity", as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" and "similarity" can be readily calculated by known methods, including but not limited to those described in: Computational Molecular Biology (Lesk, A. M., ed.) Oxford University Press, NY (1988); Biocomputing: Informatics and Genome Projects (Smith, D. W., ed.) Academic Press, NY (1993); Computer Analysis of Sequence Data, Part I (Griffin, A. M., and Griffin, H. G., eds.) Humana Press, NJ (1994); Sequence Analysis in Molecular Biology (von Heinje, G., ed.) Academic Press (1987); and Sequence Analysis Primer (Gribskov, M. and Devereux,

J., eds.) Stockton Press, NY (1991). Preferred methods to determine identity are designed to give the best match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Sequence alignments and percent identity
5 calculations may be performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences may be performed using the CLUSTAL method of alignment (Higgins and Sharp *CABIOS*. 5:151-153 (1989)) with the default parameters (GAP PENALTY=10, GAP LENGTH
10 PENALTY=10). Default parameters typically used for pairwise alignments using the CLUSTAL method are KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5.

“Codon degeneracy” refers to the nature in the genetic code permitting variation of the nucleotide sequence without effecting the amino
15 acid sequence of an encoded polypeptide. Accordingly, the instant invention relates to any nucleic acid fragment that encodes all or a substantial portion of the amino acid sequence encoding the instant microbial polypeptides as set forth in SEQ ID NOs:4, 6, 14, and 17. The skilled artisan is well aware of the “codon-bias” exhibited by a specific host
20 cell in usage of nucleotide codons to specify a given amino acid. Therefore, when synthesizing a gene for improved expression in a host cell, it is desirable to design the gene such that its frequency of codon usage approaches the frequency of preferred codon usage of the host cell.

25 “Synthetic genes” can be assembled from oligonucleotide building blocks that are chemically synthesized using procedures known to those skilled in the art. These building blocks are ligated and annealed to form gene segments which are then enzymatically assembled to construct the entire gene. “Chemically synthesized”, as related to a sequence of DNA,
30 means that the component nucleotides were assembled *in vitro*. Manual chemical synthesis of DNA may be accomplished using well-established procedures, or automated chemical synthesis can be performed using one of a number of commercially available machines. Accordingly, the genes can be tailored for optimal gene expression based on optimization of

nucleotide sequence to reflect the codon bias of the host cell. The skilled artisan appreciates the likelihood of successful gene expression if codon usage is biased towards those codons favored by the host. Determination of preferred codons can be based on a survey of genes derived from the host cell where sequence information is available.

"Gene" refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. "Native gene" refers to a gene as found in nature with its own regulatory sequences. "Chimeric gene" refers to any gene that is not a native gene, comprising regulatory and coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature. "Endogenous gene" refers to a native gene in its natural location in the genome of an organism. A "foreign" gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A "transgene" is a gene that has been introduced into the genome by a transformation procedure.

"Coding sequence" refers to a DNA sequence that codes for a specific amino acid sequence. "Suitable regulatory sequences" refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, polyadenylation recognition sequences, RNA processing site, effector binding site and stem-loop structure.

"Promoter" refers to a DNA sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. Promoters may be

derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic DNA segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental or physiological conditions. Promoters which cause a gene to be expressed in most cell types at most times are commonly referred to as "constitutive promoters". It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, DNA fragments of different lengths may have identical promoter activity.

The term "operably linked" refers to the association of nucleic acid sequences on a single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in sense or antisense orientation.

The term "expression", as used herein, refers to the transcription and stable accumulation of sense (mRNA) or antisense RNA derived from the nucleic acid fragment of the invention. Expression may also refer to translation of mRNA into a polypeptide.

"Transformation" refers to the transfer of a nucleic acid fragment into the genome of a host organism, resulting in genetically stable inheritance. In the present invention, the host cell's genome includes chromosomal and extrachromosomal (e.g., plasmid) genes. Host organisms containing the transformed nucleic acid fragments are referred to as "transgenic" or "recombinant" or "transformed" organisms.

The term "carbon substrate" refers to a carbon source capable of being metabolized by host organisms of the present invention, and particularly refers to carbon sources selected from, but not limited to, the group consisting of aliphatic carboxylic acids or dicarboxylic acids, monosaccharides, oligosaccharides, polysaccharides, and one-carbon substrates or mixtures thereof.

Nitriles particularly pertinent to the invention are nitriles of




- 5 wherein N is Nitrogen, C is Carbon, and R is selected from the group consisting of: a) C_1-C_9 alkyl that is linear, branched or cyclic, optionally substituted with a hydroxyl group, an amino group, a carboxylic acid group, a carboxamide group, a halogen atom, or an oxo group; b) C_1-C_9 alkenyl, linear, branched or cyclic, optionally substituted with a hydroxyl group, an amino group, a carboxylic acid group, a carboxamide group, a halogen atom, or an oxo group; and c) C_6-C_9 aryl, optionally substituted with a hydroxyl group, an amino group, a carboxylic acid group, a carboxamide group, or a halogen atom. Particularly useful nitriles in the invention include, but are not limited to, acrylonitrile, methacrylonitrile, 3-hydroxypropionitrile, 3-hydroxybutyronitrile, 3-hydroxyvaleronitrile, butyronitrile, adiponitrile, benzonitrile, and glycolonitrile.


Additional nitriles particularly pertinent to the invention are nitriles of Formula 3:




wherein N is Nitrogen, C is Carbon, and R^2 is selected from the group consisting of the general formulae 4, 5, 6, 7, and 8:




Formula 4




Formula 5



Formula 6



Formula 7



Formula 8

wherein Formulae 4, 5, and 6 X is N, in Formula 7 X is NH, O, or S, and in Formula 8 X is NH and Y is N, O, or S, and where R³ and R⁴ are,
 5 independently, selected from the group consisting of: a) a hydrogen atom, b) a halogen atom, c) a C₁-C₉ alkyl group that is linear, branched, or cyclic and optionally substituted with a hydroxyl group, an amino group, a carboxylic acid group, a carboxamide group, a halogen atom, or an oxo group, d) a C₁-C₉ alkenyl group that is linear, branched, or cyclic and
 10 optionally substituted with a hydroxyl group, an amino group, a carboxylic acid group, a carboxamide group, a halogen atom, or an oxo group, and e) a C₆-C₉ aryl, optionally substituted with a hydroxyl group, an amino group, a carboxylic acid group, a carboxamide group, or a halogen atom. Halogen atoms may be independently F, Cl, Br, or I. Particularly useful
 15 heterocyclic nitriles in the invention include, but are not limited to, 3-cyanopyridine, 4-cyanopyridine, pyrazinecarbonitrile, 2-furancarbonitrile, 2-thiophenecarbonitrile, and 4-thiazolecarbonitrile.

The terms "plasmid", "vector" and "cassette" refer to an extra chromosomal element often carrying genes which are not part of the
 20 central metabolism of the cell, and usually in the form of circular double-stranded DNA fragments. Such elements may be autonomously replicating sequences, genome integrating sequences, phage or nucleotide sequences, linear or circular, of a single- or double-stranded DNA or RNA, derived from any source, in which a number of nucleotide
 25 sequences have been joined or recombined into a unique construction which is capable of introducing a promoter fragment and DNA sequence for a selected gene product along with appropriate 3' untranslated sequence into a cell. "Transformation cassette" refers to a specific vector containing a foreign gene and having elements in addition to the foreign
 30 gene that facilitate transformation of a particular host cell. "Expression cassette" refers to a specific vector containing a foreign gene and having

elements in addition to the foreign gene that allow for enhanced expression of that gene in a foreign host.

The term "altered biological activity" will refer to an activity, associated with a polypeptide (protein) encoded by a microbial nucleotide sequence which can be measured by an assay method, where that activity is either greater than or less than the activity associated with the native microbial sequence. "Enhanced biological activity" refers to an altered activity that is greater than that associated with the native sequence. "Diminished biological activity" is an altered activity that is less than that associated with the native sequence.

The terms "suitable aqueous reaction mixture" or "suitable reaction mixture" refer to the materials and water in which the nitrile and/or amide substrate and enzyme catalyst come into contact. Components of suitable aqueous reaction mixtures are referred to herein and those skilled in the art appreciate the range of component variations suitable for this process.

The term "sequence analysis software" refers to any computer algorithm or software program that is useful for the analysis of nucleotide or amino acid sequences. "Sequence analysis software" may be commercially available or independently developed. Typical sequence analysis software will include but is not limited to the GCG suite of programs (Wisconsin Package Version 9.0, Genetics Computer Group (GCG), Madison, WI), BLASTP, BLASTN, BLASTX (Altschul *et al.*, *J. Mol. Biol.* 215:403-410 (1990), and DNASTAR (DNASTAR, Inc. 1228 S. Park St. Madison, WI 53715 USA), and the FASTA program incorporating the Smith-Waterman algorithm (W. R. Pearson, *Comput. Methods Genome Res.*, [Proc. Int. Symp.] (1994), Meeting Date 1992, 111-20. Editor(s): Suhai, Sándor. Publisher: Plenum, New York, NY). The term "MEME" refers to a software program used to identify conserved diagnostic motifs based on a hidden Markov model (Timothy L. Bailey and Charles Elkan, Fitting a mixture model by expectation maximization to discover motifs in biopolymers, Proceedings of the Second International Conference on Intelligent Systems for Molecular Biology, pp. 28-36, AAAI Press, Menlo Park, CA (1994)). "MAST" (Timothy L. Bailey and Michael Gribskov,

"Combining evidence using p-values: application to sequence homology searches" *Bioinformatics*, Vol. 14, pp. 48-54 (1998)) is a program that takes the output from the MEME program and searches the identified motifs against the protein databases such as EMBL and SwissProt.

- 5 Within the context of this application it will be understood that where sequence analysis software is used for analysis, that the results of the analysis will be based on the "default values" of the program referenced, unless otherwise specified. As used herein "default values" will mean any set of values or parameters which originally load with the software when
10 first initialized.

- Standard recombinant DNA and molecular cloning techniques used here are well known in the art and are described by Sambrook, J., Fritsch, E. F. and Maniatis, T., Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
15 (1989) (hereinafter "Maniatis"); and by Silhavy, T. J., Bennan, M. L. and Enquist, L. W., Experiments with Gene Fusions, Cold Spring Harbor Laboratory Cold Press Spring Harbor, NY (1984) (hereinafter "Silhavy"); and by Ausubel, F. M. *et al.*, Current Protocols in Molecular Biology, published by Greene Publishing Assoc. and Wiley-Interscience (1987)
20 (hereinafter "Ausubel").

Sequence Identification

- Comparison of the present amino acid sequences for the nitrile hydratase α and β -subunits and the amidase from *Comamonas testosteroni* 5-MGAM-4D to public databases revealed that the most
25 similar-known sequences were all from *Pseudomonas putida* 5B (NRRL-18668; see: US 5,811,286, and Wu *et al.*, *supra*). The nitrile hydratase α -subunit was about 97.1 % identical to the corresponding nitrile hydratase α -subunit from *P. putida* 5B (Table 1). The nitrile hydratase β -subunit was about 82.0 % identical to the corresponding nitrile hydratase β -subunit
30 from *P. putida* 5B. Lastly, the amidase was about 92.3 % identical to the corresponding amidase from *P. putida* 5B.

Table 1
Sequence Analysis Results

Gene Name	Similarity Identified	% Identity ^a	% Similarity ^b	E-value ^c	Citation
Amidase	Gi 6225048 sp O69768 AMID_PSEPU	92.3	92.5	0	Wu <i>et al.</i> , <i>DNA Cell Biol.</i> 17 (10), 915-920 (1998)
nitrile hydratase α -subunit	Gi 2499193 sp P97051 NHAA_PSEPU	97.1	97.1	10e-108	Payne <i>et al.</i> , <i>Biochemistry</i> 36 (18), 5447-5454 (1997)
nitrile hydratase β -subunit	Gi 2499195 sp P97052 NHAB_PSEPU	82.0	82.5	6e-92	Payne <i>et al.</i> , <i>Biochemistry</i> 36 (18), 5447-5454 (1997)

- ^a Identity is defined as percentage of amino acids that are identical between the two proteins.
- ^b Similarity is defined as percentage of amino acids that are identical or conserved between the two proteins.
- ^c Expect value. The Expect value estimates the statistical significance of the match, specifying the number of matches, with a given score, that are expected in a search of a database of this size absolutely by chance.

Despite the sequence similarities between the nitrile hydratase and amidase enzymes of *Comamonas testosteroni* 5-MGAM-4D to those of *Pseudomonas putida* 5B (NRRL-18668), the accompanying Examples demonstrate that both the thermal stability, and stability under reaction conditions, of the *Comamonas testosteroni* 5-MGAM-4D nitrile hydratase enzyme (expressed in *E. coli* transformant SW132) are both different and markedly superior to the *Pseudomonas putida* 5B nitrile hydratase (expressed in *E. coli* transformant SW30, Wu *et al.*, *supra*).

5 Identification of homologs

The instant sequences may be employed as hybridization reagents for the identification of homologs. The basic components of a nucleic acid hybridization test include a probe, a sample suspected of containing the gene or gene fragment of interest, and a specific hybridization method.

- 10 Probes of the present invention are typically single stranded nucleic acid sequences which are complementary to the nucleic acid sequences to be detected. Probes are "hybridizable" to the nucleic acid sequence to be

detected. The probe length can vary from 5 bases to tens of thousands of bases, and will depend upon the specific test to be done. Typically a probe length of about 15 bases to about 30 bases is suitable. Only part of the probe molecule need be complementary to the nucleic acid sequence to be detected. In addition, the complementarity between the probe and the target sequence need not be perfect. Hybridization does occur between imperfectly complementary molecules with the result that a certain fraction of the bases in the hybridized region are not paired with the proper complementary base.

Hybridization methods are well defined. Typically the probe and sample must be mixed under conditions which will permit nucleic acid hybridization. This involves contacting the probe and sample in the presence of an inorganic or organic salt under the proper concentration and temperature conditions. The probe and sample nucleic acids must be in contact for a long enough time that any possible hybridization between the probe and sample nucleic acid may occur. The concentration of probe or target in the mixture will determine the time necessary for hybridization to occur. The higher the probe or target concentration the shorter the hybridization incubation time needed. Optionally a chaotropic agent may be added. The chaotropic agent stabilizes nucleic acids by inhibiting nuclease activity. Furthermore, the chaotropic agent allows sensitive and stringent hybridization of short oligonucleotide probes at room temperature (Van Ness and Chen, *Nucl. Acids Res.*, 19:5143-5151, (1991)). Suitable chaotropic agents include guanidinium chloride, guanidinium thiocyanate, sodium thiocyanate, lithium tetrachloroacetate, sodium perchlorate, rubidium tetrachloroacetate, potassium iodide, and cesium trifluoroacetate, among others. Typically, the chaotropic agent will be present at a final concentration of about 3M. If desired, one can add formamide to the hybridization mixture, typically 30-50 % (v/v).

Various hybridization solutions can be employed. Typically, these comprise from about 20 to 60 % volume, preferably 30 %, of a polar organic solvent. A common hybridization solution employs about 30-50 % v/v formamide, about 0.15 to 1M sodium chloride, about 0.05 to 0.1M buffers, such as sodium citrate, Tris-HCl, PIPES or HEPES (pH range

about 6-9), about 0.05 to 0.2 % detergent, such as sodium dodecylsulfate, or between 0.5-20 mM EDTA, FICOLL (about 300-500 kilodaltons), polyvinylpyrrolidone (about 250-500 kdal), and serum albumin. Also included in the typical hybridization solution will be unlabeled carrier
5 nucleic acids from about 0.1 to 5 mg/mL, fragmented nucleic DNA, e.g., calf thymus or salmon sperm DNA, or yeast RNA, and optionally from about 0.5 to 2 % wt./vol. glycine. Other additives may also be included, such as volume exclusion agents which include a variety of polar water-soluble or swellable agents, such as polyethylene glycol, anionic polymers
10 such as polyacrylate or polymethylacrylate, and anionic saccharidic polymers, such as dextran sulfate.

Nucleic acid hybridization is adaptable to a variety of assay formats. One of the most suitable is the sandwich assay format. The sandwich assay is particularly adaptable to hybridization under non-
15 denaturing conditions. A primary component of a sandwich-type assay is a solid support. The solid support has adsorbed to it or covalently coupled to it immobilized nucleic acid probe that is unlabeled and complementary to one portion of the sequence.

Microbial Recombinant Expression

20 The genes and gene products of the instant sequences may be produced in heterologous host cells, particularly in the cells of microbial hosts. Expression in recombinant microbial hosts may be useful for the expression of various pathway intermediates; for the modulation of pathways already existing in the host, or for the synthesis of new products
25 heretofore not possible using the host.

Preferred heterologous host cells for expression of the instant genes and nucleic acid fragments are microbial hosts that can be found broadly within the fungal or bacterial families and which grow over a wide range of temperature, pH values, and solvent tolerances. For example, it
30 is contemplated that any of bacteria, yeast, and filamentous fungi will be suitable hosts for expression of the present nucleic acid fragments. Because of transcription, translation and the protein biosynthetic apparatus is the same irrespective of the cellular feedstock, functional genes are expressed irrespective of carbon feedstock used to generate

cellular biomass. Large-scale microbial growth and functional gene expression may utilize a wide range of simple or complex carbohydrates, organic acids and alcohols, saturated hydrocarbons such as methane or carbon dioxide in the case of photosynthetic or chemoautotrophic hosts.

5 However, the functional genes may be regulated, repressed or depressed by specific growth conditions, which may include the form and amount of nitrogen, phosphorous, sulfur, oxygen, carbon or any trace micronutrient including small inorganic ions. In addition, the regulation of functional genes may be achieved by the presence or absence of specific regulatory
10 molecules that are added to the culture and are not typically considered nutrient or energy sources.

Examples of host strains include but are not limited to bacterial, fungal, or yeast species such as *Aspergillus*, *Trichoderma*, *Saccharomyces*, *Pichia*, *Candida*, *Hansenula*, *Salmonella*, *Bacillus*,
15 *Acinetobacter*, *Zymomonas*, *Agrobacterium*, *Erythrobacter*, *Chlorobium*, *Chromatium*, *Flavobacterium*, *Cytophaga*, *Rhodobacter*, *Rhodococcus*, *Streptomyces*, *Brevibacterium*, *Corynebacteria*, *Mycobacterium*, *Deinococcus*, *Escherichia*, *Erwinia*, *Pantoea*, *Pseudomonas*, *Sphingomonas*, *Methylomonas*, *Methylobacter*, *Methylococcus*,
20 *Methylosinus*, *Methylomicrobium*, *Methylocystis*, *Methylobacterium*, *Alcaligenes*, *Synechocystis*, *Synechococcus*, *Anabaena*, *Thiobacillus*, *Methanobacterium*, *Klebsiella*, *Myxococcus*, and *Staphylococcus*. In another embodiment, suitable host strains are selected from the group consisting of *Aspergillus*, *Saccharomyces*, *Pichia*, *Candida*, *Hansuela*,
25 *Bacillus*, *Rhodococcus*, *Streptomyces*, *Brevibacterium*, *Corynebacteria*, *Escherichia*, *Pseudomonas*, *Methylomonas*, *Synechocystis*, and *Klebsiella*. In a further embodiment, suitable host strains are selected from the group consisting of *Bacillus*, *Rhodococcus*, *Escherichia*, *Pseudomonas*, *Klebsiella*, and *Methylomonas*.

30 Microbial expression systems and expression vectors containing regulatory sequences that direct high level expression of foreign proteins are well known to those skilled in the art. Any of these could be used to construct chimeric genes for expression of the present nitrile hydratase, amidase, and PK7. These chimeric genes could then be introduced into

appropriate microorganisms via transformation to provide high-level expression of the enzymes

Accordingly it is expected, for example, that introduction of chimeric genes encoding the instant bacterial enzyme under the control of the appropriate promoter, will demonstrate increased nitrile to amide and/or carboxylic acid conversion. It is contemplated that it will be useful to express the instant genes both in natural host cells as well as in a heterologous host. Introduction of the present genes into native hosts will result in altered levels of existing nitrile hydratase and amidase activity. Additionally, the instant genes may also be introduced into non-native host bacteria where an existing nitrile-amide-carboxylic acid pathway may be manipulated.

Vectors or cassettes useful for the transformation of suitable host cells are well known in the art. Typically the vector or cassette contains sequences directing transcription and translation of the relevant gene, a selectable marker, and sequences allowing autonomous replication or chromosomal integration. Suitable vectors comprise a region 5' of the gene which harbors transcriptional initiation controls and a region 3' of the DNA fragment which controls transcriptional termination. It is most preferred when both control regions are derived from genes homologous to the transformed host cell, although it is to be understood that such control regions need not be derived from the genes native to the specific species chosen as a production host.

Initiation control regions or promoters, which are useful to drive expression of the instant ORF in the desired host cell are numerous and familiar to those skilled in the art. Virtually any promoter capable of driving these genes is suitable for the present invention including but not limited to *CYC1*, *HIS3*, *GAL1*, *GAL10*, *ADH1*, *PGK*, *PHO5*, *GAPDH*, *ADC1*, *TRP1*, *URA3*, *LEU2*, *ENO*, *TPI* (useful for expression in *Saccharomyces*); *AOX1* (useful for expression in *Pichia*); and *lac*, *ara*, *tet*, *trp*, *IP_L*, *IP_R*, *T7*, *tac*, and *trc* (useful for expression in *Escherichia coli*) as well as the *amy*, *apr*, *npr* promoters and various phage promoters useful for expression in *Bacillus*. Additionally, the deoxy-xylulose phosphate synthase or methanol dehydrogenase operon promoter (Springer *et al.*,

FEMS Microbiol Lett 160:119-124 (1998)), the promoter for polyhydroxyalkanoic acid synthesis (Foellner *et al.*, *Appl. Microbiol. Biotechnol.* 40:284-291 (1993)), promoters identified from native plasmids in methylotrophs (EP 296484), promoters identified from methanotrophs (PCT/US03/33698), and promoters associated with antibiotic resistance [e.g., kanamycin (Springer *et al.*, *supra*; Ueda *et al.*, *Appl. Environ. Microbiol.* 57:924-926 (1991)) or tetracycline (US 4,824,786)] are suitable for expression of the present coding sequences, especially in C1 metabolizers.

10 Termination control regions may also be derived from various genes native to the preferred hosts. Optionally, a termination site may be unnecessary; however, it is most preferred if included.

Methods of manipulating genetic pathways are common and well known in the art. Selected genes in a particular pathway may be up-regulated or down-regulated by variety of methods. Additionally, 15 competing pathways may be eliminated or sublimated by gene disruption and similar techniques.

Once a key genetic pathway has been identified and sequenced, specific genes may be up-regulated to increase the output of the pathway. 20 For example, additional copies of the targeted genes may be introduced into the host cell on multicopy plasmids such as pBR322. Alternatively the target genes may be modified so as to be under the control of non-native promoters. Where it is desired that a pathway operate at a particular point in a cell cycle or during a fermentation run, regulated or inducible 25 promoters may be used to replace the native promoter of the target gene. Similarly, in some cases the native or endogenous promoter may be modified to increase gene expression. For example, endogenous promoters can be altered *in vivo* by mutation, deletion, and/or substitution (see, Kmiec, US 5,565,350; Zarling *et al.*, PCT/US93/03868).

30 Alternatively, it may be necessary to reduce or eliminate the expression of certain genes in a pathway or in competing pathways that may serve as competing sinks for energy or carbon. Methods of down-regulating genes for this purpose have been explored. Where sequence of the gene to be disrupted is known, one of the most effective methods of

gene down regulation is targeted gene disruption where foreign DNA is inserted into a structural gene so as to disrupt transcription. This can be effected by the creation of genetic cassettes comprising the DNA to be inserted (often a genetic marker) flanked by sequence having a high
5 degree of homology to a portion of the gene to be disrupted. Introduction of the cassette into the host cell results in insertion of the foreign DNA into the structural gene via the native DNA replication mechanisms of the cell (Hamilton *et al.*, *J. Bacteriol.* 171:4617-4622 (1989); Balbas *et al.*, *Gene* 136:211-213 (1993); Gueldener *et al.*, *Nucleic Acids Res.* 24:2519-2524
10 (1996); and Smith *et al.*, *Methods Mol. Cell. Biol.* 5:270-277(1996)).

Antisense technology is another method of down regulating genes where the sequence of the target gene is known. To accomplish this, a nucleic acid segment from the desired gene is cloned and operably linked to a promoter such that the anti-sense strand of RNA will be transcribed.
15 This construct is then introduced into the host cell and the antisense strand of RNA is produced. Antisense RNA inhibits gene expression by preventing the accumulation of mRNA which encodes the protein of interest. The person skilled in the art will know that special considerations are associated with the use of antisense technologies in order to reduce
20 expression of particular genes. For example, the proper level of expression of antisense genes may require the use of different chimeric genes utilizing different regulatory elements known to the skilled artisan.

Although targeted gene disruption and antisense technology offer effective means of down regulating genes where the sequence is known,
25 other less specific methodologies have been developed that are not sequence based. For example, cells may be exposed to a UV radiation and then screened for the desired phenotype. Mutagenesis with chemical agents is also effective for generating mutants and commonly used substances include chemicals that affect non-replicating DNA such as
30 HNO₂ and NH₂OH, as well as agents that affect replicating DNA such as acridine dyes, notable for causing frameshift mutations. Specific methods for creating mutants using radiation or chemical agents are well documented in the art. (See for example, Thomas D. Brock in *Biotechnology: A Textbook of Industrial Microbiology*, Second Edition

(1989) Sinauer Associates, Inc., Sunderland, MA. (hereinafter "Brock"), or Deshpande, Mukund V., *Appl. Biochem. Biotechnol.*, 36:227 (1992) (hereinafter "Deshpande")).

Another non-specific method of gene disruption is the use of transposable elements or transposons. Transposons are genetic elements that insert randomly in DNA but can be later retrieved on the basis of sequence to determine where the insertion has occurred. Both *in vivo* and *in vitro* transposition methods are known. Both methods involve the use of a transposable element in combination with a transposase enzyme. When the transposable element or transposon, is contacted with a nucleic acid fragment in the presence of the transposase, the transposable element will randomly insert into the nucleic acid fragment. The technique is useful for random mutagenesis and for gene isolation, since the disrupted gene may be identified on the basis of the sequence of the transposable element. Kits for *in vitro* transposition are commercially available (see for example The Primer Island Transposition Kit, available from Perkin Elmer Applied Biosystems, Branchburg, NJ, based upon the yeast Ty1 element; The Genome Priming System, available from New England Biolabs, Beverly, MA; based upon the bacterial transposon Tn7; and the EZ::TN Transposon Insertion Systems, available from Epicentre Technologies, Madison, WI, based upon the Tn5 bacterial transposable element).

Industrial Production of Biocatalyst

Commercial production of biocatalyst for preparing amides using transformants harboring the nitrile hydratase catalyst disclosed herein (encoded by genes for the α - and β -subunits, and optionally, accessory protein P7K) and for preparing carboxylic acids from amides using the amidase catalyst disclosed herein may be conducted using a variety of culture methodologies. Large-scale production of a specific gene product may be produced by both batch and continuous culture methodologies.

A classical batch culturing method is a closed system where the composition of the media is set at the beginning of the culture and not subject to artificial alterations during the culturing process. Thus, at the beginning of the culturing process the media is inoculated with the desired organism or organisms and growth or metabolic activity is permitted to

occur adding nothing to the system. Typically, however, a "batch" culture is batch with respect to the addition of carbon source and attempts are often made at controlling factors such as pH and oxygen concentration. In batch systems the metabolite and biomass compositions of the system change constantly up to the time the culture is terminated. Within batch cultures cells moderate through a static lag phase to a high growth log phase and finally to a stationary phase where growth rate is diminished or halted. If untreated, cells in the stationary phase will eventually die. Cells in log phase are often responsible for the bulk of production of end product or intermediate in some systems. Stationary or post-exponential phase production can be obtained in other systems.

A variation on the standard batch system is the Fed-Batch system. Fed-Batch culture processes are also suitable in the present invention and comprise a typical batch system with the exception that the substrate is added in increments as the culture progresses. Fed-Batch systems are useful when catabolite repression is apt to inhibit the metabolism of the cells and where it is desirable to have limited amounts of substrate in the media. Measurement of the actual substrate concentration in Fed-Batch systems is difficult and is therefore estimated on the basis of the changes of measurable factors such as pH, dissolved oxygen and the partial pressure of waste gases such as CO₂. Batch and Fed-Batch culturing methods are common and well known in the art and examples may be found in Brock (*supra*) and Deshpande (*supra*).

Commercial production of biocatalysts may also be accomplished with a continuous culture. Continuous cultures are an open system where a defined culture media is added continuously to a bioreactor and an equal amount of conditioned media is removed simultaneously for processing. Continuous cultures generally maintain the cells at a constant high-liquid-phase density where cells are primarily in log phase growth.

Continuous or semi-continuous culture allows for the modulation of one factor or any number of factors that affect cell growth or end product concentration. For example, one method will maintain a limiting nutrient such as the carbon source or nitrogen level at a fixed rate and allow all other parameters to moderate. In other systems a number of factors

affecting growth can be altered continuously while the cell concentration, measured by media turbidity, is kept constant. Continuous systems strive to maintain steady state growth conditions and thus the cell loss due to media being drawn off must be balanced against the cell growth rate in the culture. Methods of modulating nutrients and growth factors for continuous culture processes as well as techniques for maximizing the rate of product formation are well known in the art of industrial microbiology and a variety of methods are detailed by Brock, *supra*.

Fermentation media in the present invention must contain suitable carbon substrates. Suitable substrates may include but are not limited to aliphatic carboxylic acid and dicarboxylic acids (such as lactic acid or succinic acid), glycerol, monosaccharides (such as glucose and fructose), disaccharides (such as lactose or sucrose), oligosaccharides (such as soluble starch), polysaccharides (such as starch or cellulose or mixtures thereof), and unpurified mixtures from renewable feedstocks (such as cheese whey permeate, cornsteep liquor, sugar beet molasses, and barley malt). Additionally the carbon substrate may also be one-carbon substrates such as carbon dioxide, methane, or methanol for which metabolic conversion into key biochemical intermediates has been demonstrated. In addition to one and two carbon substrates, methylotrophic organisms are also known to utilize a number of other carbon-containing compounds such as methylamine, glucosamine, and a variety of amino acids for metabolic activity. For example, methylotrophic yeast are known to utilize the carbon from methylamine to form trehalose or glycerol (Bellion *et al.*, *Microb. Growth C1 Compd.*, [Int. Symp.], 7th (1993), 415-32. Editor(s): Murrell, J. Collin; Kelly, Don P. Publisher: Intercept, Andover, UK). Similarly, various species of *Candida* will metabolize alanine or oleic acid (Sulter *et al.*, *Arch. Microbiol.* 153:485-489 (1990)). Therefore, it is contemplated that the source of carbon utilized in the present invention may encompass a wide variety of carbon-containing substrates and will only be limited by the microorganism employed.

Biocatalytic Conversion of Nitriles to Amides or Carboxylic Acids

An aqueous reaction mixture containing the aliphatic, aromatic, or heterocyclic aromatic nitrile is prepared by mixing the nitrile with an aqueous suspension of the appropriate enzyme catalyst. Intact microbial cells can be used as catalyst without any pretreatment, such as permeabilization or heating. Alternatively, the cells can be immobilized in a polymer matrix (e.g., alginate, carrageenan, polyvinyl alcohol, or polyacrylamide gel (PAG)) or on a soluble or insoluble support (e.g., celite, silica) to facilitate recovery and reuse of the catalyst. Methods to immobilize cells in a polymer matrix or on a soluble or insoluble support have been widely reported and are well known to those skilled in the art. The enzyme can also be isolated from the microbial cells and used directly as catalyst, or the enzyme can be immobilized in a polymer matrix or on a soluble or insoluble support. These methods have also been widely reported and are well known to those skilled in the art (Methods in Biotechnology, Vol. 1: Immobilization of Enzymes and Cells; Gordon F. Bickerstaff, Editor; Humana Press, Totowa, NJ, USA; 1997).

Intact microbial cells, either immobilized or unimmobilized, containing genes that encode a polypeptide having nitrile hydratase activity or amidase activity, or containing genes that encode a combination of polypeptides separately having nitrile hydratase and amidase activities, can be used as catalyst without any pretreatment, such as permeabilization, freeze thawing or heating. Alternatively, the microbial cells may be permeabilized by methods familiar to those skilled in the art (e.g., treatment with toluene, detergents, or freeze-thawing) to improve the rate of diffusion of materials into and out of the cells. Methods for permeabilization of microbial cells are well-known to those skilled in the art (Felix, H., *Anal. Biochem.*, 120:211-234 (1982)).

Some of the aliphatic or aromatic nitriles used as starting material in the present invention are only moderately water soluble. Their solubility also depends on the temperature of the solution and the salt concentration in the aqueous phase; the optional inclusion of a buffer, or the production of the ammonium salt of a carboxylic acid by hydrolysis of the corresponding amide are two possible sources of salt in a reaction

mixture. In the present case, producing a hydrated or hydrolyzed reaction product at a concentration greater than the solubility limit of the starting aliphatic or aromatic nitrile is accomplished using a reaction mixture that is initially composed of two phases: an aqueous phase (containing the
5 enzyme catalyst and dissolved aliphatic or aromatic nitrile) and an organic phase (the undissolved aliphatic or aromatic nitrile, optionally dissolved in an organic solvent not miscible with the aqueous phase). As the reaction progresses, the aliphatic or aromatic nitrile dissolves into the aqueous phase, eventually yielding a product mixture which may be a single phase, depending on the solubility of the products in water, and on the presence
10 or absence of an optional organic solvent not miscible with water.

The aqueous phase of a two-phase reaction mixture can contain, at a minimum, only as much water as is sufficient to result in a) complete conversion of the aliphatic or aromatic nitrile to the corresponding amide
15 or carboxylic acid (dependent on whether only active nitrile hydratase or a combination of active nitrile hydratase and amidase enzyme are present), and b) maintenance of the hydrolytic activity of the enzyme catalyst. The reaction may also be run by adding the aliphatic or aromatic nitrile to the reaction mixture at a rate approximately equal to the enzymatic hydration
20 or hydrolysis reaction rate, thereby maintaining a single-phase aqueous reaction mixture, thereby avoiding the potential problem of substrate inhibition of the enzyme at high starting material concentrations.

The final concentration of aliphatic or aromatic amide or carboxylic acid in solution in the product mixture at complete conversion of the
25 corresponding aliphatic or aromatic nitrile may range from 0.001 M to the solubility limit of the aliphatic or aromatic nitrile in the product mixture. Product may precipitate from the reaction mixture during the course of the reaction, allowing for the production of amide or carboxylic acid in excess of the solubility of said product in the reaction mixture. Typically, the
30 concentration of the aliphatic or aromatic amide or carboxylic acid product in solution in the product mixture ranges from 0.001 M to 7.0 M. The aliphatic or aromatic amide or carboxylic acid may also be isolated from the product mixture (after removal of the catalyst) by optionally adjusting the pH of the reaction mixture to between 2.0 and 2.5 with concentrated

HCl when the product of the reaction is a carboxylic acid, saturating the resulting solution with sodium chloride, and extracting the aliphatic or aromatic amide or carboxylic acid with a suitable organic solvent, such as ethyl acetate, ethyl ether, methyl isobutyl ketone or dichloromethane. The organic extracts are then combined, stirred with a suitable drying agent (e.g., magnesium sulfate), filtered, and the solvent removed (e.g., by rotary evaporation) to produce the desired product in high yield and in high purity (typically 98-99 % pure). If desired, the product can be further purified by recrystallization or distillation.

10 The concentration of enzyme catalyst in the reaction mixture depends on the specific catalytic activity of the enzyme catalyst and is chosen to obtain the desired rate of reaction. The wet cell weight of the microbial cells used as catalyst in hydrolysis reactions typically ranges from 0.001 grams to 0.300 grams of wet cells per mL of total reaction
15 volume, preferably from 0.002 grams to 0.050 grams of wet cells per mL; the cells may be optionally immobilized as described above. The specific activity of the microbial cells (IU/gram dry cell weight) is determined by measuring the rate of conversion of a 0.10-0.50 M solution of a nitrile substrate to the desired amide or carboxylic acid product at 25 °C, using a
20 known weight of microbial cell catalyst. An IU of enzyme activity is defined as the amount of enzyme activity required to convert one micromole of substrate to product per minute.

 The temperature of the hydrolysis reaction is chosen to optimize both the reaction rate and the stability of enzyme catalyst. The
25 temperature of the reaction may range from just above the freezing point of the reaction mixture (ca. 0 °C) to 65 °C, with a preferred range of reaction temperature of from 5 °C to 45 °C. An enzyme catalyst solution or suspension may be prepared by suspending the unimmobilized or immobilized cells in distilled water, or in an aqueous reaction mixture of a
30 buffer that will maintain the initial pH of the reaction between 5.0 and 10.0, preferably between 6.0 and 8.0, or by suspending the immobilized enzyme catalyst in a similar mixture, or by preparing a solution of a cell extract, partially purified or purified enzyme(s), or a soluble form of the immobilized enzymes in a similar mixture. After the nitrile is added and a s

the reaction proceeds, the pH of the reaction mixture may change due to the formation of an ammonium salt of the carboxylic acid from the corresponding nitrile functionality of the aliphatic or aromatic nitrile (when using a combination of nitrile hydratase and amidase enzymes). The
5 reaction can be run to completely convert the nitrile with no pH control, or a suitable acid or base can be added over the course of the reaction to maintain the desired pH.

EXAMPLES

The present invention is further defined in the following Examples
10 that indicate preferred embodiments of the invention. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various uses and conditions.

15 In the following Examples, the percent recovery of nitrile, and the percent yields of the corresponding amide and carboxylic acid products were based on the initial concentration of nitrile present in the reaction mixture, and were determined by HPLC. Analyses of 3-hydroxyvaleronitrile, adiponitrile, butyronitrile, benzonitrile, and
20 methacrylonitrile were performed by HPLC using a refractive index detector in combination with a Supelco LC-18-DB column (15 cm x 4.6 mm diameter) with precolumn at 25 °C and 10 mM acetic acid, 10 mM sodium acetate in 7.5 % methanol in water as eluent at 1.5 mL/min. Analyses for glycolonitrile, acrylonitrile, 3-HPN, 3-HBN, and their
25 corresponding reaction products were performed by HPLC using a Bio-Rad HPX-87H organic acid analysis column (30 cm x 7.8 mm dia.) with precolumn at 50 °C and 0.010 N H₂SO₄ as eluent at 1 mL/min. Analyses for 3-cyanopyridine, pyrazinecarbonitrile, 2-furancarbonitrile, 2-thiophenecarbonitrile, 4-thiazolecarbonitrile and their corresponding
30 reaction products, were performed by HPLC using a UV detector at 254 nm in combination with a 10-cm x 4-mm ID, 5 µm C8 Discovery column (Supelco) with precolumn, 1.0 mL/min of 5% CH₃CN/95% 10mM NaOAc,

10mM AcOH as solvent, and N,N-dimethylbenzamide as external standard.

GENERAL METHODS

5 Standard recombinant DNA and molecular cloning techniques used in the Examples are well known in the art and are described by Maniatis (*supra*) and Ausubel (*supra*).

Materials and methods suitable for the maintenance and growth of bacterial cultures are well known in the art. Techniques suitable for use in
10 the following examples may be found as set out in Manual of Methods for General Bacteriology (Phillip Gerhardt, R. G. E. Murray, Ralph N. Costilow, Eugene W. Nester, Willis A. Wood, Noel R. Krieg and G. Briggs Phillips, eds., American Society for Microbiology, Washington, DC (1994)) or in Brock, *supra*.

15 The following abbreviations in the specification correspond to units of measure, techniques, properties, or compounds as follows: "sec" means second(s), "min" means minute(s), "h" means hour(s), "d" means day(s), "mL" means milliliters, "L" means liters, "mM" means millimolar, "M" means molar, "mmol" means millimole(s), "rpm" means revolutions per
20 minute, "slpm" means standard liters per minute, "psig" means pounds per square inch, and "wt" means weight. "HPLC" means high performance liquid chromatography, "ca" means approximately, "O.D." means optical density at the designated wavelength, "dcw" means dry cell weight, and "IU" means International Units.

25

EXAMPLE 1

Identification of a genomic DNA fragment encoding *C. testosteroni* 5-MGAM-4D nitrile hydratase

Comamonas testosteroni 5-MGAM-4D (ATCC 55744) was grown in LB media at 37 °C with shaking. Genomic DNA was prepared using a
30 Puregene DNA Isolation Kit according to the manufacturer (Gentra Systems, Minneapolis, MN). A Southern analysis (Southern *et al.*, *J. Mol. Biol.*, 98:503 (1975)) was performed on *EcoRI* restricted genomic DNA using *Psuedomonas putida* NRRL-18668 genes (SEQ ID NOs: 1 and 2) encoding nitrile hydratase alpha and beta subunits (US 5,811,286) as

probes. Probe labeling, hybridization, and detection were done using ECL random prime labeling and detection systems version II according to the manufacturer (Amersham International, Buckinghamshire, UK). The alpha (SEQ ID NO:1) and beta (SEQ ID NO:2) probes each showed positive
5 hybridization to the same 5.7 kb *EcoRI* DNA fragment.

EXAMPLE 2

Cloning of a genomic DNA fragment encoding *C. testosteroni* 5-MGAM-4D nitrile hydratase

10 Genomic DNA from *C. testosteroni* 5-MGAM-4D was prepared (Example 1), restricted with *EcoRI*, and subjected to standard agarose gel electrophoresis. DNA fragments in the size range of approximately 5-7 kb were isolated and ligated to *EcoRI* restricted pUC19 (New England Biolabs, Beverly, MA). This plasmid library was plated and screened with
15 the *P. putida* NRRL-18668 nitrile hydratase α -subunit gene probe (SEQ ID NO:1). Probe labeling, hybridization and detection were done using ECL random prime labeling and detection systems version II according to the manufacturer (Amersham International). A positively hybridizing colony was isolated and determined to contain an insert of 5.7
20 kb (pKP57).

EXAMPLE 3

Determination of the nucleotide sequence of the genes encoding *C. testosteroni* 5-MGAM-4D nitrile hydratase

25 The nucleotide sequence of the pKP57 (EXAMPLE 2) insert was determined using an ABI 377-XL DNA sequencer and BigDye Terminator Cycle Sequencing chemistry. A BlastN analysis (Altschul *et al.*, *Nucleic Acids Res.*, 25:3389-3402 (1997)) of the obtained sequence to the GenBank® database confirmed the presence of complete genes encoding
30 nitrile hydratase α and β -subunits and a partial gene encoding amidase. Nucleotide sequences of the pKP57 insert encoding nitrile hydratase α - and β -subunits are given in SEQ ID NO:3 and SEQ ID NO:5, respectively.

Deduced amino acid sequences of the pKP57 insert for the α - and β -subunits are given in SEQ ID NO:4 and SEQ ID NO:6, respectively.

BlastP analysis was conducted using the deduced amino acid sequence for the nitrile hydratase α - and β -subunits. Results are shown in Table 1. The closest match to the instant amino acid sequence of the nitrile hydratase α -subunit was the nitrile hydratase α -subunit from *Pseudomonas putida* NRRL-18668 (97.1 % identity, 97.1 % similarity, E value= $10e^{-108}$). The closest match to the instant amino acid sequence of the nitrile hydratase β -subunit was the nitrile hydratase β -subunit from *Pseudomonas putida* NRRL-18668 (82.0 % identity, 82.5 % similarity, E value= $6e^{-92}$).

EXAMPLE 4

Production of *C. testosteroni* 5-MGAM-4D nitrile hydratase in *E. coli*

A DNA fragment encoding the nitrile hydratase alpha and beta subunits from *C. testosteroni* 5-MGAM-4D (SEQ ID NO:7) was obtained from pKP57 by standard PCR via a GeneAmp Kit according to the manufacturer (Roche, Branchburg, NJ) using Primer 1 (SEQ ID NO:8) and Primer 2 (SEQ ID NO:9) and subcloned into pGEM-T (Promega, Madison, WI) under control of the T7 promoter to generate pSW131. *E. coli* BL21(DE3) (Novagen, Madison, WI) was transformed with pSW131 using standard procedures. Growth and induction of *E. coli* BL21(DE3) harboring pSW131 was carried out essentially as recommended by Novagen. Modifications included the addition of cobalt chloride and sodium citrate at the time of induction to a final concentration of 0.01 mg/ml and 0.1 mg/ml respectively. Induction was carried out at 30 C for 16 hrs. Production of alpha (23 kDa) and beta (24 kDa) proteins was confirmed by standard SDS-PAGE analysis.

EXAMPLE 5

Conversion of 3-hydroxyvaleronitrile (3-HVN) by transformed *E. coli*

Growth and induction of *E. coli* BL21(DE3) cells harboring pSW131 was carried out as described in EXAMPLE 4. Cells were then harvested by centrifugation, washed twice in buffer (0.1 M potassium phosphate pH

7.0) and suspended at 100 mg wet cells/ml in buffer. The nitrilase activity assay mix included cells (50 mg/mL), 3-hydroxyvaleronitrile (0.3 M) and buffer (0.1 M potassium phosphate, pH 7.0) stirred at ambient temperature. HPLC analysis demonstrated 17 % conversion of 3-HVN to the corresponding amide (3-hydroxyvaleramide) in 15 min.

EXAMPLE 6

Production of high-level *C. testosteroni* 5-MGAM-4D nitrile hydratase activity in *E. coli* requires downstream sequence

10 A plasmid (pSW132; ATCC PTA-5073), containing a DNA fragment encoding the *C. testosteroni* 5-MGAM-4D nitrile hydratase alpha and beta subunits plus 0.9 kb of downstream DNA encoding the accessory protein P7K (SEQ ID NO:11) under control of the T7 promoter was constructed by replacing the smaller *Bam*HI/*Pst*I fragment in pSW131 with the
15 corresponding *Bam*HI/*Pst*I fragment from pKP57. *E. coli* BL21(DE3) (Novagen) was transformed with pSW132 using standard procedures. Growth and induction of *E. coli* BL21(DE3) harboring pSW132 ("*E. coli* strain SW132") was carried out as described in EXAMPLE 4, and production of alpha and beta proteins was confirmed by standard SDS-
20 PAGE analysis.

A DNA fragment (SEQ ID NO:11) encoding the *C. testosteroni* 5-MGAM-4D nitrile hydratase alpha and beta subunits plus 0.9 kb of downstream DNA encoding the accessory protein P7K was also obtained from pKP57 by standard PCR via a GeneAmp Kit according to the
25 manufacturer (Roche) using Primer 1 (SEQ ID NO:8) and Primer 3 (SEQ ID NO:12) and subcloned into pTrcHis2-TOPO under control of the trc promoter to generate pSW137 in *E. coli* TOP10 according to the manufacturer (Invitrogen). Growth and induction of *E. coli* TOP10 harboring pSW137 was carried out as described in EXAMPLE 4, and
30 production of alpha and beta proteins was confirmed by standard SDS-PAGE analysis. Production of the alpha and beta proteins in *E. coli* BL21(DE3) harboring pSW132 and in *E. coli* TOP10 harboring pSW137 was qualitatively indistinguishable to that obtained from *E. coli* BL21(DE3) harboring pSW131 (EXAMPLE 4).

Growth and induction of *E. coli* BL21(DE3) harboring pSW132 was carried out as described in EXAMPLE 4. Cells were then harvested by centrifugation, washed twice in buffer (0.1 M potassium phosphate pH 7.0) and suspended at 100 mg wet cells/ml in buffer. The nitrile hydratase activity assay mix included cells (2 mg/mL), 3-hydroxyvaleronitrile (0.3 M) and buffer (0.1 M potassium phosphate, pH 7.0) stirred at ambient temperature. HPLC analysis demonstrated 100 % conversion of 3-hydroxyvaleronitrile to 3-hydroxyvaleramide in 5 min. Similarly, nitrile hydratase activity assay of *E. coli* TOP10 harboring pSW137 demonstrated 100 % conversion of 3-hydroxyvaleronitrile to 3-hydroxyvaleramide in 5 min. Comparing these results to those obtained from pSW131 (Example 5) demonstrated the importance of the DNA downstream of the nitrile hydratase beta gene, in obtaining maximal nitrile hydratase activity (Table 2). The downstream nucleotide sequence (SEQ ID NO:10) contains a small open reading frame, the sequence of which is given in SEQ ID NO:13. The deduced amino acid sequence (called P7K) is given in SEQ ID NO:14.

Table 2. Comparison of Expression Vectors pSW131 and pSW132

Expression Vector	Expressed Genetic Components from <i>C. testosteroni</i> 5-MGAM-4D	% Conversion of 0.35 M 3-HVN to 3-HVAm at Room Temperature
PSW131	Nitrile hydratase α - and β -subunits	17 % in 15 min
PSW132	Nitrile hydratase α - and β -subunits plus accessory protein "P7K"	100 % in 5 min

EXAMPLE 7

Determination of the nucleotide sequence of the gene encoding *C. testosteroni* 5-MGAM-4D amidase

Genomic DNA from *C. testosteroni* 5-MGAM-4D was prepared (EXAMPLE 1), restricted with *Pst*I, and subjected to Southern analysis using a standard PCR product comprising the first 0.6 kb of the pKP57 (EXAMPLE 2) insert as a probe (SEQ ID NO:15). Probe labeling, hybridization and detection were done using ECL random prime labeling

and detection systems version II according to the manufacturer (Amersham International). This probe gave hybridized to a 2.4 kb *Pst*I fragment. Genomic DNA digested with *Pst*I was subjected to standard agarose gel electrophoresis. DNA fragments in the size range of approximately 2-4 kb were isolated and ligated into *Pst*I restricted pUC19. This plasmid library was plated and screened with the same 0.6 kb probe (SEQ ID NO:15). Probe labeling, hybridization and detection were done using ECL random prime labeling and detection systems version II according to the manufacturer (Amersham International). A positively hybridizing colony was isolated and determined to contain an insert of 2.4 kb (pKP59).

Nucleotide sequencing confirmed that the insert is a DNA fragment that overlaps the *Eco*RI DNA fragment previously cloned (pKP57). Thus, by combining the nucleotide sequences from pKP57 and pKP59, the complete nucleotide sequence for the amidase gene was determined (SEQ ID NO:16). The deduced amidase amino acid sequence is given in SEQ ID NO:17. The nucleotide sequence of a 7.4 kb DNA fragment from *C. testosteroni* 5-MGAM-4D comprising complete coding sequences for amidase and nitrile hydratase is given in SEQ ID NO:18.

BlastP analysis was conducted using the deduced amidase amino acid sequence (SEQ ID NO:17; Table 1). The closest publicly known match was to that of the amidase from *Pseudomonas putida* NRRL-18668 (92.3 % identity, 92.5 % similarity, E value= 0).

EXAMPLE 8

Production of *C. testosteroni* 5-MGAM-4D amidase in *E. coli*

A DNA fragment encoding the amidase from *C. testosteroni* 5-MGAM-4D was obtained from genomic DNA by standard PCR via a GeneAmp Kit according to the manufacturer (Roche), using Primer 4 (SEQ ID NO:19) and Primer 5 (SEQ ID NO:20) and subcloned into pGEM-T (Promega) under control of the T7 promoter to generate pKP60. *E. coli* BL21(DE3) (Novagen) was transformed with pKP60 using standard procedures. Growth and induction of *E. coli* BL21(DE3) harboring pKP60

was carried out according to Novagen, and production of amidase protein was confirmed by standard SDS-PAGE analysis

A DNA fragment encoding the amidase from *C. testosteroni* 5-MGAM-4D was also obtained by standard PCR using Primer 6 (SEQ ID NO:21) and Primer 7 (SEQ ID NO:22) and subcloned into pTrcHis2 TOPO under control of the trc promoter to generate pSW133 in *E. coli* TOP10 according to the manufacturer (Invitrogen). Growth and induction of *E. coli* TOP10 harboring pSW133 was carried out according to Invitrogen, and production of amidase protein was confirmed by standard SDS-PAGE analysis.

EXAMPLE 9

Co-production of *C. testosteroni* 5-MGAM-4D nitrile hydratase and amidase in *E. coli*

A DNA fragment encoding the amidase, NHase alpha and beta, and accessory protein P7K from *C. testosteroni* 5-MGAM-4D (SEQ ID NO:23) was obtained from genomic DNA by standard PCR via a GeneAmp Kit according to the manufacturer (Roche), using Primer 6 (SEQ ID NO:21) and Primer 3 (SEQ ID NO:12), and subcloned into pTrcHis2-TOPO (Invitrogen) under control of the Trc promoter to generate pSW136.

EXAMPLE 10

Fermentation of *Escherichia coli* SW132 Cells

The production of nitrile hydratase in a 14 L Braun Biostat C fermentor (B. Braun Biotech International GmbH, Melsungen, Germany) was made in mineral medium with glucose, ammonia, and yeast extract. *E. coli* strain SW132 (*E. Coli* BL21(DE3) harboring plasmid pSW132 as described in Example 6) was grown in a seed culture for 10 h prior to inoculation of the fermentor. IPTG (1 mM) was added to the fermentor at 30-35 OD_{λ=550} and cells were harvested 5 h after IPTG addition.

Fermentation protocol: vessel medium was prepared in an initial batch of 7.5 L containing 32 g KH₂PO₄, 8.0 g MgSO₄·7H₂O, 8.0 g (NH₄)₂SO₄, 50 g yeast extract, and 10 mL Mazu DF204 antifoam (BASF Corporation,

Mount Olive, NJ). Following sterilization, 369 g glucose solution (60 % w/w), 160 mL trace element solution (Table 3), and 100 mg/L ampicillin were added. NH_4OH (40 % w/v) and 20 % w/v H_2SO_4 were used for pH control. The set points for agitation, aeration, pH, pressure, dissolved oxygen concentration (DO), and temperature are described in Table 4 below. The dissolved oxygen concentration was controlled at 25 % of air saturation with the agitation to rise first with increase oxygen demand and the aeration to follow. The 500 mL seed culture was grown in a 2 L flask at 36 °C, 300 rpm for 10 h to an $\text{OD}_{\lambda=550}$ of >2.0. In the fermentor at culture densities of 20-30 OD additional AMP was added to 100 mg/L. IPTG was added to 1 mM at culture densities of 30-35 OD. Glucose feed was started at <5 g/L and the scheduled rates are described in Table 5. Glucose feed rate was reduced if glucose accumulated above 2 g/L. Five hours after IPTG addition the cells were chilled to 5-10 °C and harvested by centrifugation; 490 g (wet cells) was harvested. The kinetics of growth and nitrile hydratase production are presented in Table 6.

Table 3 Trace elements solution:

Chemical	Concentration g/L
Citric acid	10.0
$\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$	1.50
$\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$	5.00
$\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$	0.39
$\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$	0.38
$\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$	0.20
$\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$	0.30

Table 4
Fermentation Run Conditions

	Initial Set-Point	Minimum	Maximum
Stirrer (rpm)	400	400	850
Airflow (slpm)	2	2	16
pH	6.8	6.8	6.8
Pressure (psig)	0.5	0.5	0.5
DO	25 %	25 %	25 %
Temp. C	36	36	36

5

Table 5
Glucose feed protocol

Time, h	Rate (g/min)
0-2	0.39
2-8	0.78
8- End	0.60

10

Table 6
The kinetics of growth and nitrile hydratase production

Time, h	OD _{λ=550}	Glucose (g/L)	Nitrile Hydratase Production (U/g dry cell wt.)
2.9	5.8	26.2	
4.9	15.7	20.2	
6.3	28	8.5	9157
8.1	44	6	10220
11.2	57.6	0.04	10888

EXAMPLE 11

15

Hydration of Nitriles to Corresponding Amides by Unimmobilized *E. coli* SW132 Cells

20

25

To a 20-mL reaction vessel equipped with magnetic stirring was added 0.04, 0.4, 2.0, or 5.0 mmol of acrylonitrile, methacrylonitrile, 3-hydroxypropionitrile, 3-hydroxybutyronitrile, 3-hydroxyvaleronitrile, butyronitrile, adiponitrile, benzonitrile, or glycolonitrile and distilled, deionized water was added to adjust the final volume of the mixture to 3.0 mL. To the reaction vessel was next added 1.0 mL of an aqueous suspension of 0.44 – 8.8 mg dry cell weight (dcw)/mL of *E. coli* SW132 cells (prepared as described in Example 10) in 0.10 M potassium phosphate buffer (pH 7.0, except for glycolonitrile, which was run at pH 6.0), and the mixture was stirred at 25 °C. Samples (0.100 mL) of the reaction mixture were mixed with 0.400 mL of water, and then 0.200 mL of the diluted sample was either (a) mixed with 0.200 mL of 0.200 M sodium butyrate (acrylonitrile and 3-hydroxyvaleronitrile HPLC standard), 0.200 M

N-ethylacetamide (methacrylonitrile HPLC standard), 0.200 M isobutyric acid (butyronitrile and 3-hydroxypropionitrile HPLC standard), or 0.200 M malonic acid (3-hydroxybutyronitrile HPLC standard) in water, or (b) measured against a calibration curve for product at 100 % nitrile conversion (5-cyanovaleramide, adipamide, benzamide, glycolamide). The resulting mixture was centrifuged, and the supernatant analyzed by HPLC.

All reactions produced only the amide as the hydration product at 100 % conversion of nitrile, with no hydrolysis of the nitrile to the corresponding carboxylic acid.

Table 7
Hydration of Nitriles to Corresp. Amides by *E. coli* SW132 Cells

Nitrile	conc. (M)	mg dcw/mL	time (h)	amide	Amide yield (%)
Acrylonitrile	0.50	2.2	3	acrylamide	100
Acrylonitrile	1.25	8.8	5	acrylamide	98
Methacrylonitrile	0.50	2.2	20	methacrylamide	100
3-hydroxypropionitrile	0.50	2.2	3	3-hydroxypropionamide	100
3-hydroxybutyronitrile	0.50	4.4	4	3-hydroxybutyramide	99
3-hydroxyvaleronitrile	0.51	2.2	1	3-hydroxyvaleramide	99
Butyronitrile	0.50	2.2	16	butyramide	100
Adiponitrile	0.50	0.44	1.5	5-cyanovaleramide	93
				adipamide	8
Adiponitrile	0.50	4.4	1	adipamide	100
Benzonitrile	0.010	2.2	2	benzamide	99
Glycolonitrile	0.10	2.2	2	glycolamide	63

15

EXAMPLE 12

Hydration of 3-Hydroxyvaleronitrile using a Partially-Purified Protein

Extract of *E. coli* SW132 Cells

E. coli SW132 cells (0.4874 g) were suspended in 2.0 mL of cold breaking buffer consisting of 1 mM DTT and 0.1 mM PMSF in 0.1 M

20

potassium phosphate buffer (pH 7). The suspension (200 mg wet cell weight/mL) was loaded in a French Pressure Mini Cell, and the cells were ruptured at 16000 - 17000 psi. Cell debris was removed from the resulting mixture by centrifugation at 38000 RCF for 15 min. Approximately 1.68
5 mL of extract supernatant was recovered, having a nitrile hydratase activity equivalent to a 199 mg wet cell weight/mL cell suspension.

To a 20-mL reaction vessel equipped with magnetic stirring was added 0.4 mL of the *E. coli* SW132 extract supernatant, 0.6 mL of deionized water, and 3 mL of 0.667 M 3-hydroxyvaleronitrile in water. The
10 mixture was stirred at 25 °C. Samples (0.100 mL) of the reaction mixture were mixed with 0.100 mL of water, and then 0.200 mL of the diluted sample was mixed with 0.200 mL of 0.200 M sodium butyrate (external standard) and analyzed by HPLC. After 120 min, the yield of 3-hydroxyvaleramide was 99 % at 100 % conversion of 3-
15 hydroxyvaleronitrile.

EXAMPLE 13

Comparison of Thermal Stability of Nitrile Hydratase from *Comamonas testosteroni* 5-MGAM-4D and *Pseudomonas putida* 5B Cells

A 44 mg dry cell weight/mL suspension of either *E. coli* SW30 wet
20 cells (having active nitrile hydratase from *Pseudomonas putida* 5B) or *E. coli* SW132 wet cells (having active nitrile hydratase from *Comamonas testosteroni* 5-MGAM-4D) in 0.50 M phosphate buffer was heated to 50 °C in a water bath. At predetermined times, aliquots of the 50 °C cell suspensions were rapidly cooled to 25 °C in a water bath, and these
25 suspensions were assayed for remaining nitrile hydratase activity by adding 1.0 mL aliquots of the heated/cooled cell suspensions with stirring to 3.0 mL of 0.667 M 3-hydroxyvaleronitrile in water at 25 °C. Samples (0.100 mL) of the reaction mixture were withdrawn at predetermined times and mixed with 0.100 mL of water, then 0.200 mL of the diluted sample
30 was mixed with 0.200 mL of 0.200 M sodium butyrate (external standard), centrifuged, and the supernatant and analyzed by HPLC. The rate of hydration of 3-hydroxyvaleronitrile in each reaction was determined, and remaining nitrile hydratase specific activity of the cells calculated. The nitrile hydratase specific activity and % of enzyme recovered for *E. coli*

SW30 and *E. coli* SW132, respectively, as a function of time at 50 °C is listed in Table 8, below.

Table 8
Comparison of *E. coli* SW30 and *E. coli* SW132 Nitrile Hydratase
Thermostability.

Time At 50 °C (min)	SW30 Nitrile Hydratase (IU/g dcw)	Nitrile Hydratase Recovery (%)	SW132 Nitrile Hydratase (IU/g dcw)	Nitrile Hydratase Recovery (%)
0	103	100	7752	100
30	1.6	1.6	6940	90
60	0	0	6437	83

EXAMPLE 14

10 Comparison of *Comamonas testosteroni* 5-MGAM-4D and
Pseudomonas putida 5B Nitrile Hydratase for Hydration of Acrylonitrile to
Acrylamide

To a 20-mL reaction vessel (equipped with magnetic stirring) was added 0.270 g (5.1 mmol) of acrylonitrile and a suspension of either 107.6 mg dry cell weight *E. coli* SW30 wet cells (expressing the active nitrile hydratase from *Pseudomonas putida* 5B) or 4.41 mg dry cell weight *E. coli* SW132 wet cells (expressing the active nitrile hydratase from *Comamonas testosteroni* 5-MGAM-4D) in a total volume of 9.664 mL of 0.10 M potassium phosphate buffer (pH 7.0); the amount of dry cell weight present in each reaction was chosen to provide ca. equivalent nitrile hydratase activities (IU/mL) in the reaction mixtures. The final concentration of acrylonitrile was 0.51 M. The mixture was stirred at 25 °C. Samples (0.200 mL) of the reaction mixture were mixed with 0.200 mL of 0.200 M sodium butyrate (HPLC standard) in water, and 0.020 mL of 6 N acetic acid. The resulting sample was centrifuged, and the supernatant analyzed by HPLC. The reaction time, % acrylonitrile conversion, and % yield of acrylamide are listed in Table 9. For reactions using *E. coli* SW30 as biocatalyst, a loss of nitrile hydratase activity was observed over the course of the reaction, and incomplete conversion of nitrile was obtained at extended reaction times.

Table 9
Comparison of *E. coli* SW30 and *E. coli* SW132 in Acrylonitrile Hydration Reactions

<i>E. coli</i> Construct	mg dcw/mL	time (h)	acrylonitrile conv. (%)	acrylamide yield (%)
SW30	10.8	1.5	42	38
SW30	10.8	20	80	82
SW30	10.8	45	96	97
SW132	0.44	1.5	100	98

EXAMPLE 15

Comparison of *Comamonas testosteroni* 5-MGAM-4D and *Pseudomonas putida* 5B Nitrile Hydratase for Hydration of 3-Hydroxyvaleronitrile to 3-Hydroxyvaleramide

To a 20-mL reaction vessel (equipped with magnetic stirring) was added 3.0 mL of a solution of 0.204 g (2.0 mmol) of 3-hydroxyvaleronitrile in distilled, deionized water and a 1.0 mL suspension of 44 mg dry cell weight of either *E. coli* SW30 wet cells (expressing the active nitrile hydratase from *Pseudomonas putida* 5B) or *E. coli* SW132 wet cells (expressing the active nitrile hydratase from *Comamonas testosteroni* 5-MGAM-4D) in 0.10 M potassium phosphate buffer (pH 7.0). The final concentration of 3-hydroxyvaleronitrile was 0.50 M. The mixture was stirred at 25 °C. Samples (0.100 mL) of the reaction mixture were mixed with 0.100 mL of water, 0.200 mL of 0.200 M sodium butyrate (HPLC standard) in water, and 0.020 mL of 6 N HCl. The resulting mixture was centrifuged, and the supernatant analyzed by HPLC. The reaction time, % 3-hydroxyvaleronitrile conversion, and % yield of 3-hydroxyvaleramide are listed in Table 10. For reactions using SW30 as biocatalyst, a loss of nitrile hydratase activity was observed over the course of the reaction, and incomplete conversion of nitrile was obtained at extended reaction times.

Table 10
Comparison of *E. coli* SW30 and *E. coli* SW132 in 3-Hydroxyvaleronitrile Reactions

<i>E. coli</i> Construct	mg dcw/mL	Time (h)	3-hydroxyvaleronitrile conv. (%)	3-hydroxyvaleramide yield (%)
SW30	11	1	12	12
<i>E. coli</i> Construct	mg dcw/mL	Time (h)	3-hydroxyvaleronitrile conv. (%)	3-hydroxyvaleramide yield (%)
SW30	11	7	42	40
SW30	11	24	61	61
SW132	11	0.25	100	100

EXAMPLE 16

Hydration of 3-Hydroxyvaleronitrile to 3-Hydroxyvaleramide by *E. coli* SW137

To a 4-mL reaction vessel (equipped with magnetic stirring) was added 0.75 mL of an aqueous solution containing 0.404 M 3-hydroxyvaleronitrile and 0.25 mL of a suspension of 18 mg dry cell weight *E. coli* SW137 wet cells (prepared as described in Example 6) in 0.10 M potassium phosphate buffer (pH 7.0). The *E. coli* SW137 cells express the polypeptide having nitrile hydratase activity from *Comamonas testosteroni* 5-MGAM-4D. The final concentration of 3-hydroxyvaleronitrile was 0.303 M. The mixture was stirred at 25 °C. Samples (0.100 mL) of the reaction mixture were mixed with 0.100 mL of water, 0.200 mL of 0.200 M sodium butyrate (HPLC standard) in water, and 0.020 mL of 6 N HCl. The resulting mixture was centrifuged, and the supernatant analyzed by HPLC. After 10 min, the conversion of 3-hydroxyvaleronitrile was 100 %, and the yield of 3-hydroxyvaleramide was 100 %.

EXAMPLE 17

Immobilization of *Escherichia coli* SW132 Cells in Calcium Cross-linked Alginate

Into a 250-mL media bottle (equipped with magnetic stir bar and containing 59.7 g of distilled, deionized water at 50 °C) was slowly added 3.30 g of FMC BioPolymer Protanal® LF 10/60 alginate with rapid stirring. The mixture was heated to 75-80 °C with rapid stirring until the alginate was completely dissolved, and the resulting solution cooled to 25 °C in a water bath. To the alginate suspension was added 40.8 g of *Escherichia*

coli SW132 wet cell paste (22 % dry cell weight) and 16.2 mL of distilled water with stirring. The cell/alginate mixture was added dropwise by syringe to 640 mL of 0.20 M calcium acetate buffer (pH 7.0) at 25 °C with stirring. After stirring for 2 h, the buffer was decanted from the resulting beads (82 g), which were resuspended in 200 mL of 0.20 M calcium acetate buffer (pH 7.0) at 25 °C. With stirring, 4.10 g of 25 wt % glutaraldehyde (GA) in water was added and the beads mixed for 1.0 h at 25 °C. To the suspension was then added 16.4 g of 12.5 wt % polyethylenimine (PEI) (BASF Lupasol® PR971L, average molecular weight ca. 750,000) in water, and the beads mixed for an additional 1 h at 25 °C. The GA/PEI-crosslinked beads were then washed twice with 250 mL of 0.05 M calcium acetate buffer (pH 7.0) at 25 °C, and stored in this same buffer at 5 °C.

EXAMPLE 18

Hydration of Nitriles (0.50 M to 3.0 M) to Corresponding Amides by Alginate-Immobilized *Escherichia coli* SW132 Cells in Consecutive Batch Reactions with Biocatalyst Recycle

Into a 50-mL jacketed reaction vessel (equipped with an overhead stirrer (temperature-controlled at 25 °C or 35 °C) with a recirculating temperature bath) was placed 4.0 g of GA/PEI-crosslinked *Escherichia coli* SW132 cell/alginate beads prepared as described in Example 17. To the reaction vessel was added 0.2 mL of 0.20 M calcium acetate buffer (pH 7.0, 2.0 mM final calcium ion concentration in reaction mixture), 10, 20, 40, or 60 mmol of acrylonitrile, methacrylonitrile, or 3-hydroxyvaleronitrile, and the final volume of the reaction mixture adjusted to 20 mL by the addition of distilled, deionized water. The mixture was stirred at 25 °C or 35 °C. Samples (0.100 mL) of the reaction mixture were mixed with 0.400 mL of water, and then 0.200 mL of the diluted sample was mixed with 0.200 mL of 0.200 M sodium butyrate (acrylonitrile and 3-hydroxyvaleronitrile HPLC external standard) or 0.200 M N-ethylacetamide (methacrylonitrile HPLC external standard) in water. The resulting mixture was centrifuged, and the supernatant analyzed by HPLC.

At the completion of the reaction (100 % conversion of nitrile), the product mixture was decanted from the biocatalyst beads, and additional distilled, deionized water, 0.2 mL of 0.20 M calcium acetate buffer (pH 7.0, 2.0 mM final calcium ion concentration in reaction mixture) and 10, 20, 40 or 60 mmol of acrylonitrile, methacrylonitrile, or 3-hydroxyvaleronitrile mixed with the reaction heel (immobilized-cell catalyst and remaining

product mixture from the first reaction) at 25 °C or 35 °C. At the completion of the second reaction, the product mixture was decanted and a third reaction performed as before. The reaction time, product yield for acrylamide, methacrylamide, or 3-hydroxyvaleramide, and the percent recovered biocatalyst activity for each recycle reaction is listed in Table 11 below.

Table 11
Hydration of Nitriles (0.50 M to 3.0 M) to Corresponding Amides by Immobilized *E. coli* SW132 Cells in Consecutive Batch Reactions with Biocatalyst Recycle

Nitrile	Conc. (M)	Temp. (°C)	Rxn #	Time (h)	Amide Yield (%)	Recovered Biocatalyst Activity (%)
Acrylonitrile	0.53	25	1	0.18	100	100
			2	0.18	100	94
			3	0.12	100	112
Acrylonitrile	1.02	25	1	0.25	100	100
			2	0.25	100	118
			3	0.25	100	124
acrylonitrile	2.05	25	1	0.5	100	100
			2	0.5	100	95
			3	0.5	100	100
acrylonitrile	3.04	25	1	0.75	100	100
			2	0.83	100	98
			3	0.5	100	106
acrylonitrile	1.06	35	1	0.15	100	100
			2	0.25	100	107
			3	0.15	100	110
methacrylonitrile	1.00	25	1	1.0	100	100
			2	1.0	100	106
			3	1.5	99	103
3-hydroxyvaleronitrile	1.05	25	1	0.75	100	100
			2	0.75	100	93
			3	0.75	100	84
3-hydroxyvaleronitrile	2.05	25	1	1.5	100	100
			2	2.0	100	102
			3	2.0	100	93

EXAMPLE 19

Immobilization of *E. coli* SW132 Cells in Carrageenan

Into a 250 mL media bottle equipped with magnetic stir bar and containing 54.6 g of water at 50 °C is slowly added 2.88 g of kappa-carrageenan (FMC RG300) with rapid stirring. The mixture is heated to 75-80 °C with rapid stirring until the carrageenan is completely dissolved, and the resulting solution cooled to 55-56 °C (ca. 52 °C gelling temperature) in a thermostated water bath. A suspension of 18.6 g of *E. coli* SW132 wet cell paste (22.0 % dry cell wt) in 19.7 g of 0.35 M sodium phosphate buffer (pH 7.3) is heated to 50 °C for 15 min, then added to the carrageenan solution at 55 - 56 °C with stirring. The cell/carrageenan mixture is immediately added slowly to 383 mL of soybean oil at 50 °C with stirring using an overhead stirrer. After cell/carrageenan droplets of the desired size are produced in the oil by controlling the stirring rate, the temperature of the oil is reduced to 40-42 °C to gel the droplets, and the oil decanted from the resulting beads. The beads are washed with 150 mL of 0.1 M potassium bicarbonate buffer (pH 7.0), then suspended in 182 mL of this same buffer, and 1.9 g of 25 wt % glutaraldehyde in water is added and the beads mixed for 1 h at 25 °C. To the mixture is then added 7.6 g of 12.5 wt % polyethylenimine (BASF Lupasol PR971L) average Mw ca. 750,000) in water, and the beads mixed for 1 h at 25 °C. The beads are then washed twice with 0.30 M ammonium bicarbonate (pH 7.0), and stored in this same at 5 °C.

EXAMPLE 20

Hydration of Acrylonitrile by Carrageenan-Immobilized *E. coli* SW132 Cells

Into a 50-mL jacketed reaction vessel equipped with an overhead stirrer (temperature-controlled at 35 °C with a recirculating temperature bath) is placed 4.0 g of GA/PEI-crosslinked *E. coli* SW132 cell/carrageenan beads prepared as described in Example 19. To the reaction vessel is then added 1.06 g of acrylonitrile (1.0 M final concentration), the final volume of the reaction mixture adjusted to 20 mL by the addition of distilled, deionized water, and the mixture stirred at 35 °C. Samples (0.100 mL) of the reaction mixture are mixed with

0.400 mL of water, and then 0.200 mL of the diluted sample is mixed with 0.200 mL of 0.200 M sodium butyrate (HPLC external standard) in water. The resulting mixture is centrifuged, and the supernatant analyzed by HPLC. At complete conversion of acrylonitrile, there is a quantitative yield
5 of acrylamide.

EXAMPLE 21

Hydration of Acrylonitrile Using Immobilized Nitrile Hydratase from Partially-Purified Protein Extract of *E. coli* SW132 Cells

Into a 25-mL Erlenmeyer flask is weighed 1.0 g of oxirane acrylic
10 beads (Sigma). To the flask is then added ca. 7.5 mL of a solution containing potassium phosphate buffer (50 mM, pH 8.0), and the oxirane acrylic beads suspended in the buffer by briefly mixing the contents of the flask. After cessation of mixing, the beads settle to the bottom of the flask, and the fine particles which float to the top of the mixture are
15 removed by pipette, along with as much of the supernatant which can be removed without disturbing the settled beads. This washing procedure is repeated a second time. To the flask is then added 1.0 mL of the *E. coli* SW132 cell extract supernatant described in Example 12 and the final volume of the mixture adjusted to 10 mL with additional potassium
20 phosphate buffer. The resulting mixture is mixed on a rotary platform shaker for 16 h at 25 °C. The mixture is then transferred to a chromatography column equipped with a fritted bed support, and the immobilized nitrile hydratase washed three times with 10 mL of potassium phosphate buffer and stored at 5 °C in this same buffer.

25 Into a 50-mL jacketed reaction vessel equipped with an overhead stirrer (temperature-controlled at 35 °C with a recirculating temperature bath) is placed 1.0 g of immobilized *E. coli* SW132 nitrile hydratase prepared as described above. To the reaction vessel is then added 1.06 g of acrylonitrile (1.0 M final concentration), the final volume of the reaction
30 mixture adjusted to 20 mL by the addition of distilled, deionized water, and the mixture stirred at 35 °C. Samples (0.100 mL) of the reaction mixture are mixed with 0.400 mL of water, and then 0.200 mL of the diluted sample is mixed with 0.200 mL of 0.200 M sodium butyrate (HPLC external standard) in water. The resulting mixture is centrifuged, and the

supernatant analyzed by HPLC. At complete conversion of acrylonitrile, there is a quantitative yield of acrylamide.

EXAMPLE 22

Hydration of 3-Cyanopyridine (0.5 M) to Nicotinamide by Unimmobilized

E. coli SW132 Cells

5 To a 15-mL polypropylene centrifuge tube was added 3.73 mL of 50 mM potassium phosphate buffer (pH 7.0), 1.0 mL of a suspension of 22.1 mg dry cell weight *E. coli* SW132 wet cells (prepared as described in Example 10) in 50 mM potassium phosphate buffer (pH 7.0), and 0.2660
10 g of 3-cyanopyridine. The final concentration of 3-cyanopyridine was 0.501 M. The reaction mixture was mixed on a rotating platform at 23 °C. After 15 minutes, 7.50 mL of 95:5 acetonitrile/water containing 0.30 M N,N-dimethylbenzamide (HPLC external standard) was added to the reaction, the resulting mixture centrifuged, and a 0.100 mL of the
15 supernatant mixed with 0.900 mL of acetonitrile and analyzed by HPLC. The conversion of 3-cyanopyridine was 100 %, and the yields of nicotinamide and nicotinic acid were 100 % and 0%, respectively.

EXAMPLE 23

Hydration of 3-Cyanopyridine (1.0 M) to Nicotinamide by Unimmobilized

E. coli SW132 Cells

20 To a 15-mL polypropylene centrifuge tube was added 3.47 mL of 50 mM potassium phosphate buffer (pH 7.0), 1.0 mL of a suspension of 55.2 mg dry cell weight *E. coli* SW132 wet cells (prepared as described in Example 10) in 50 mM potassium phosphate buffer (pH 7.0), and 0.5339
25 g of 3-cyanopyridine. The final concentration of 3-cyanopyridine was 1.00 M. The reaction mixture was mixed on a rotating platform at 23 °C. After 30 min, 7.50 mL of 95:5 acetonitrile/water containing 0.30 M N,N-dimethylbenzamide (HPLC external standard) was added to the reaction, the resulting mixture centrifuged, and a 0.100 mL of the supernatant
30 mixed with 0.900 mL of acetonitrile and analyzed by HPLC. The conversion of 3-cyanopyridine was 100 %, and the yields of nicotinamide and nicotinic acid were 99 % and 0%, respectively.

EXAMPLE 24

Hydration of 3-Cyanopyridine using a Partially-Purified Protein Extract of *E. coli* SW132 Cells

E. coli SW132 cells (0.5377 g, (prepared as described in Example 10)) were suspended in 5.377 mL of cold breaking buffer consisting of 1 mM DTT and 0.1 mM PMSF in 0.50 mM potassium phosphate buffer (pH 7.0). The suspension (100 mg wet cell weight/mL) was loaded in a French Pressure Mini Cell, and the cells were ruptured at 16000 - 17000 psi (approximately 110.3 – 117.2 megapascal (Mpa)). Cell debris was removed from the resulting mixture by centrifugation at 38000 RCF for 15 min. Approximately 4.20 mL of extract supernatant was recovered, having a nitrile hydratase activity equivalent to a 100 mg wet cell weight/mL (22.06 mg dry cell weight/mL) cell suspension.

To a 15-mL polypropylene centrifuge tube was added 3.73 mL of 50 mM potassium phosphate buffer (pH 7.0), 1.0 mL of an *E. coli* SW132 cell extract suspension prepared as described above, and 0.2626 g of 3-cyanopyridine. The final concentration of 3-cyanopyridine was 0.494 M. The reaction mixture was mixed on a rotating platform at 23 °C. After 15 min, 7.50 mL of 95:5 acetonitrile/water containing 0.30 M N,N-dimethylbenzamide (HPLC external standard) was added to the reaction, the resulting mixture centrifuged, and a 0.100 mL of the supernatant mixed with 0.900 mL of acetonitrile and analyzed by HPLC. The conversion of 3-cyanopyridine was 100 %, and the yields of nicotinamide and nicotinic acid were 100 % and 0%, respectively.

EXAMPLE 25

Hydration of Pyrazinecarbonitrile (0.5 M) to Pyrazinamide by Unimmobilized *E. coli* SW132 Cells

To a 15-mL polypropylene centrifuge tube was added 3.73 mL of 50 mM potassium phosphate buffer (pH 7.0), 1.0 mL of a suspension of 22.1 mg dry cell weight *E. coli* SW132 wet cells (prepared as described in Example 10) in 50 mM potassium phosphate buffer (pH 7.0), and 0.2694 g of pyrazinecarbonitrile. The final concentration of pyrazinecarbonitrile was 0.512 M. The reaction mixture was mixed on a rotating platform at 23 °C. After 15 min, 7.50 mL of 95:5 acetonitrile/water containing 0.30 M

N,N-dimethylbenzamide (HPLC external standard) was added to the reaction, the resulting mixture centrifuged, and a 0.100 mL of the supernatant mixed with 0.900 mL of acetonitrile and analyzed by HPLC. The conversion of pyrazinecarbonitrile was 100 %, and the yields of pyrazinamide and pyrazinecarboxylic acid were 100 % and 0%, respectively.

EXAMPLE 26

Hydration of Pyrazinecarbonitrile (1.0 M) to Pyrazinamide by Unimmobilized *E. coli* SW132 Cells

To a 15-mL polypropylene centrifuge tube was added 3.47 mL of 50 mM potassium phosphate buffer (pH 7.0), 1.0 mL of a suspension of 55.2 mg dry cell weight *E. coli* SW132 wet cells (prepared as described in Example 10) in 50 mM potassium phosphate buffer (pH 7.0), and 0.5330 g of pyrazinecarbonitrile. The final concentration of pyrazinecarbonitrile was 1.00 M. The reaction mixture was mixed on a rotating platform at 23 °C. After 30 min, 7.50 mL of 95:5 acetonitrile/water containing 0.30 M N,N-dimethylbenzamide (HPLC external standard) was added to the reaction, the resulting mixture centrifuged, and a 0.100 mL of the supernatant mixed with 0.900 mL of acetonitrile and analyzed by HPLC. The conversion of pyrazinecarbonitrile was 100 %, and the yields of pyrazinamide and pyrazinecarboxylic acid were 100 % and 0%, respectively.

EXAMPLE 27

Hydration of 2-Furancarbonitrile (0.5 M) to 2-Furancarboxamide by Unimmobilized *E. coli* SW132 Cells

To a 15-mL polypropylene centrifuge tube was added 3.78 mL of 50 mM potassium phosphate buffer (pH 7.0), 1.0 mL of a suspension of 22.1 mg dry cell weight *E. coli* SW132 wet cells (prepared as described in Example 10) in 50 mM potassium phosphate buffer (pH 7.0), and 0.2381 g of 2-furancarbonitrile. The final concentration of 2-furancarbonitrile was 0.506 M. The reaction mixture was mixed on a rotating platform at 23 °C. After 30 min, 7.50 mL of 95:5 acetonitrile/water containing 0.30 M N,N-dimethylbenzamide (HPLC external standard) was added to the reaction, the resulting mixture centrifuged, and a 0.100 mL of the supernatant

mixed with 0.900 mL of acetonitrile and analyzed by HPLC. The conversion of 2-furancarboxitrile was 100 %, and the yields of 2-furancarboxamide and 2-furancarboxylic acid were 99 % and 0%, respectively.

5

EXAMPLE 28

Hydration of 2-Thiophenecarbonitrile (0.3 M) to 2-Thiophenecarboxamide by Unimmobilized *E. coli* SW132 Cells

To a 15-mL polypropylene centrifuge tube was added 3.86 mL of 50 mM potassium phosphate buffer (pH 7.0), 1.0 mL of a suspension of 22.1 mg dry cell weight *E. coli* SW132 wet cells (prepared as described in Example 10) in 50 mM potassium phosphate buffer (pH 7.0), and 0.1691 g of 2-thiophenecarbonitrile. The final concentration of 2-thiophenecarbonitrile was 0.307 M. The reaction mixture was mixed on a rotating platform at 27 °C. After 30 min, 7.50 mL of 95:5 acetonitrile/water containing 0.30 M N,N-dimethylbenzamide (HPLC external standard) was added to the reaction, the resulting mixture centrifuged, and a 0.100 mL of the supernatant mixed with 0.900 mL of acetonitrile and analyzed by HPLC. The conversion of 2-thiophenecarbonitrile was 99.5 %, and the yields of 2-thiophenecarboxamide and 2-thiophenecarboxylic acid were 98 % and 0%, respectively.

20

EXAMPLE 29

Hydration of 4-Thiazolecarbonitrile (0.5 M) to 4-Thiazolecarboxamide by Unimmobilized *E. coli* SW132 Cells

To a 15-mL polypropylene centrifuge tube is added 3.70 mL of 50 mM potassium phosphate buffer (pH 7.0), 1.0 mL of a suspension of 22.1 mg dry cell weight *E. coli* SW132 wet cells (prepared as described in Example 10) in 50 mM potassium phosphate buffer (pH 7.0), and 0.2754 g of 4-thiazolecarbonitrile. The final concentration of 4-thiazolecarbonitrile is 0.500 M. The reaction mixture is mixed on a rotating platform at 25 °C. After 30 min, 7.50 mL of 95:5 acetonitrile/water containing 0.30 M N,N-dimethylbenzamide (HPLC external standard) is added to the reaction, the resulting mixture centrifuged, and a 0.100 mL of the supernatant mixed with 0.900 mL of acetonitrile and analyzed by HPLC. The conversion of 4-

25

30

thiazolecarbonitrile is 100 %, and the yields of 4-thiazolecarboxamide and 4-thiazolecarboxylic acid are 100 % and 0%, respectively.

EXAMPLE 29

Hydration of 3-Cyanopyridine (1.0 M) to Nicotinamide by

Alginate-Immobilized *Escherichia coli* SW132 Cells in Consecutive Batch

Reactions with Biocatalyst Recycle at 25 °C

Into a 50-mL jacketed reaction vessel (equipped with an overhead stirrer (temperature-controlled at 25 °C with a recirculating temperature bath) was placed 1.0 g of GA/PEI-crosslinked *Escherichia coli* SW132 cell/alginate beads prepared as described in Example 17. To the reaction vessel was added 0.2 mL of 0.20 M calcium acetate buffer (pH 7.0, 2.0 mM final calcium ion concentration in reaction mixture), 16.67 mL of distilled, deionized water, and 2.124 g of 3-cyanopyridine. The final concentration of 3-cyanopyridine was 1.00 M in 20 mL of reaction mixture. Samples (0.100 mL) of the reaction mixture were mixed with 0.400 mL of 95:5 acetonitrile/water containing 0.30 M N,N-dimethylbenzamide (HPLC external standard), and a 0.100 mL of the resulting solution mixed with 0.900 mL of acetonitrile and analyzed by HPLC. After 24 h, the conversion of 3-cyanopyridine was 100 %, and the yields of nicotinamide and nicotinic acid were 100 % and 0%, respectively. The initial reaction rate for production of nicotinamide, measured during the first 30 min of reaction, was 7.65 mM/minute.

At the completion of the reaction (100 % conversion of nitrile), the product mixture was decanted from the biocatalyst beads, and 0.2 mL of 0.20 M calcium acetate buffer (pH 7.0, 2.0 mM final calcium ion concentration in reaction mixture), 16.56 mL of distilled, deionized water, and 2.127 g of 3-cyanopyridine was added to the catalyst beads in the reaction vessel. The final concentration of 3-cyanopyridine was 1.06 M in 20 mL of reaction mixture. Samples (0.100 mL) of the reaction mixture were mixed with 0.400 mL of 95:5 acetonitrile/water containing 0.30 M N,N-dimethylbenzamide (HPLC external standard), and a 0.100 mL of the resulting solution mixed with 0.900 mL of acetonitrile and analyzed by HPLC. After 23 h, the conversion of 3-cyanopyridine was 100 %, and the yields of nicotinamide and nicotinic acid were 100 % and 0%, respectively.

The initial reaction rate for production of nicotinamide, measured during the first 30 min of reaction, was 4.60 mM/minute.

EXAMPLE 30

Hydration of 3-Cyanopyridine (1.0 M) to Nicotinamide by 5 Alginate-Immobilized *Escherichia coli* SW132 Cells in Consecutive Batch Reactions with Biocatalyst Recycle at 10 °C

Into a 50-mL jacketed reaction vessel (equipped with an overhead stirrer (temperature-controlled at 10 °C with a recirculating temperature bath) was placed 4.0 g of GA/PEI-crosslinked *Escherichia coli* SW132
10 cell/alginate beads prepared as described in Example 17. To the reaction vessel was added 0.2 mL of 0.20 M calcium acetate buffer (pH 7.0, 2.0 mM final calcium ion concentration in reaction mixture), 13.63 mL of distilled, deionized water, and 2.127 g of 3-cyanopyridine. The final concentration of 3-cyanopyridine was 1.00 M in 20 mL of reaction mixture.
15 Samples (0.100 mL) of the reaction mixture were mixed with 0.400 mL of 95:5 acetonitrile/water containing 0.30 M N,N-dimethylbenzamide (HPLC external standard), and a 0.100 mL of the resulting solution mixed with 0.900 mL of acetonitrile and analyzed by HPLC. After 1 h, the conversion of 3-cyanopyridine was 100 %, and the yields of nicotinamide and nicotinic
20 acid were 100 % and 0%, respectively. The initial reaction rate for production of nicotinamide, measured during the first 30 min of reaction, was 23.4 mM/minute.

At the completion of the reaction (100 % conversion of nitrile), the product mixture was decanted from the biocatalyst beads, and 0.2 mL of
25 0.20 M calcium acetate buffer (pH 7.0, 2.0 mM final calcium ion concentration in reaction mixture), 13.14 mL of distilled, deionized water, and 2.125 g of 3-cyanopyridine was added to the catalyst beads in the jacketed reaction vessel at 10 °C. The final concentration of 3-cyanopyridine was 1.00 M in 20 mL of reaction mixture. Samples
30 (0.100 mL) of the reaction mixture were mixed with 0.400 mL of 95:5 acetonitrile/water containing 0.30 M N,N-dimethylbenzamide (HPLC external standard), and a 0.100 mL of the resulting solution mixed with 0.900 mL of acetonitrile and analyzed by HPLC. After 1 h, the conversion of 3-cyanopyridine was 94 %, and the yields of nicotinamide and nicotinic

acid were 94 % and 0%, respectively. The initial reaction rate for production of nicotinamide, measured during the first 30 min of reaction, was 18.5 mM/minute.

EXAMPLE 31

5 Hydration of 3-Cyanopyridine (3.0 M) to Nicotinamide by
 Alginate-Immobilized *Escherichia coli* SW132 Cells in Consecutive Batch
 Reactions with Biocatalyst Recycle at 25 °C

 Into a 50-mL jacketed reaction vessel (equipped with an overhead
stirrer (temperature-controlled at 25 °C with a recirculating temperature
10 bath) was placed 2.0 g of GA/PEI-crosslinked *Escherichia coli* SW132
cell/alginate beads prepared as described in Example 17. To the reaction
vessel was added 0.2 mL of 0.20 M calcium acetate buffer (pH 7.0,
2.0 mM final calcium ion concentration in reaction mixture), 11.42 mL of
distilled, deionized water, and 6.3648 g of 3-cyanopyridine. The final
15 concentration of 3-cyanopyridine was 3.00 M in 20 mL of reaction mixture.
Samples (0.100 mL) of the reaction mixture were mixed with 1.400 mL of
95:5 acetonitrile/water containing 0.30 M N,N-dimethylbenzamide (HPLC
external standard), and a 0.100 mL of the resulting solution mixed with
0.900 mL of acetonitrile and analyzed by HPLC. After 48 h, the conversion
20 of 3-cyanopyridine was 100 %, and the yields of nicotinamide and nicotinic
acid were 100 % and 0%, respectively.

CLAIMS

WHAT IS CLAIMED IS:

1. An isolated polynucleotide encoding a polypeptide comprising the alpha-subunit of a nitrile hydratase enzyme, said polypeptide having the amino acid sequence as represented in SEQ ID NO:4.
2. An isolated polynucleotide encoding a polypeptide having at least 98 % identity to a polypeptide alpha-subunit of the nitrile hydratase enzyme from *Comamonas testosteroni* 5-MGAM-4D as represented by SEQ ID NO:4.
3. An isolated polynucleotide encoding a polypeptide comprising the alpha-subunit of a nitrile hydratase enzyme, said isolated polynucleotide having the nucleic acid sequence represented in SEQ ID NO:3.
4. A polypeptide having the amino acid sequence represented in SEQ ID NO:4.
5. An isolated polynucleotide encoding a polypeptide comprising the beta-subunit of a nitrile hydratase enzyme, said polypeptide having the amino acid sequence as represented in SEQ ID NO:6.
6. An isolated polynucleotide encoding a polypeptide comprising the beta-subunit of a nitrile hydratase enzyme, said isolated polynucleotide having the nucleic acid sequence represented in SEQ ID NO:5.
7. An isolated polynucleotide encoding a polypeptide having at least 95 % identity to a polypeptide as represented by SEQ ID NO:6, wherein the polypeptide is the beta-subunit of the nitrile hydratase enzyme from *Comamonas testosteroni* 5-MGAM-4D.
8. A polypeptide having the amino acid sequence represented in SEQ ID NO:6.
9. An isolated polynucleotide encoding the alpha- and beta-subunits of a nitrile hydratase enzyme, said isolated polynucleotide having the nucleic acid sequence as represented in SEQ ID NO:7.
10. An isolated polynucleotide encoding the alpha- and beta-subunits of a nitrile hydratase enzyme and an accessory protein, said

isolated polynucleotide having the nucleic acid sequence as represented in SEQ ID NO:11.

11. An isolated polynucleotide encoding a polypeptide comprising an amidase, said polypeptide having the amino acid sequence as represented in SEQ ID NO:17.

12. An isolated polynucleotide encoding a polypeptide having amidase activity and having at least 95 % identity to the polypeptide from *Comamonas testosteroni* 5-MGAM-4D represented by SEQ ID NO:17.

13. An isolated polynucleotide encoding a polypeptide comprising an amidase enzyme, said isolated polynucleotide having the nucleic acid sequence represented in SEQ ID NO:16.

14. A polypeptide having the amino acid sequence represented in SEQ ID NO:17.

15. An isolated polynucleotide encoding a polypeptide comprising an accessory protein, said polypeptide having the amino acid sequence as represented in SEQ ID NO:14.

16. An isolated polynucleotide encoding a polypeptide comprising an accessory protein, said isolated polynucleotide having the nucleic acid sequence represented in SEQ ID NO:13.

17. A polypeptide having the amino acid sequence represented in SEQ ID NO:14.

18. An isolated polynucleotide encoding the alpha- and beta-subunits of a nitrile hydratase enzyme, an accessory protein, and an amidase, said alpha- and beta-subunits of a nitrile hydratase enzyme, the accessory protein, and the amidase encoded by the nucleic acid sequence as represented in SEQ ID NO:23.

19. An expression vector comprising any one of the nucleic acid sequences of Claims 1-3, 5-7, 9-13, 15-16, or 18.

20. The expression vector as contained in *E. coli* SW132 designated ATCC PTA-5073.

21. The expression vector as contained in *E. coli* SW137 designated ATCC PTA-5074

22. A transformed microbial host cell comprising the expression vector of Claim 19.

23. A transformed microbial host cell comprising the expression vector of Claims 20 or 21.

24. The transformed microbial host cell of Claim 23 wherein the microbial host cell is a bacterium, yeast, or filamentous fungi.

25. The transformed microbial host cell of Claim 24 wherein the microbial host cell is

- a) a bacterium selected from the group consisting of the genera *Escherichia*, *Pseudomonas*, *Rhodococcus*, *Acinetobacter*, *Bacillus*, *Methylobacter*, and *Streptomyces*;
- b) a yeast selected from the group consisting of the genera *Pichia*, *Hansenula*, and *Saccharomyces*; or
- c) a filamentous fungi selected from the group consisting of the genera *Aspergillus*, *Neurospora*, and *Penicillium*.

26. The transformed microbial host cell of Claim 25 wherein the microbial host cell is *Escherichia coli*.

27. A purified transformed microbial host cell selected from the group consisting of *Escherichia coli* SW132 designated ATCC PTA-5073 and a purified microbial host cell *Escherichia coli* SW137 designated ATCC PTA-5074.

28. A method for producing polypeptides comprising

- a) culturing, under suitable conditions, a transformed microbial host cell containing the expression vector of Claim 19; and
- b) recovering the polypeptides produced in step a).

29. A method for converting a substrate containing one or more nitrile functional groups to an amide, the method comprising:

- a) contacting, under suitable reaction conditions, a transformed microbial host cell expressing a nitrile hydratase polypeptide encoded by the nucleic acid sequence of SEQ ID NO:11 with a substrate containing one or more nitrile functional groups; and
- b) recovering the amide produced in step (a).

30. The method of Claim 29 wherein the substrate comprises at least one nitrile functional group and is a nitrile of

Formula 1 $R-C\equiv N$ or Formula 2 $N\equiv C-R-C\equiv N$

wherein N is Nitrogen, C is Carbon, and R is:

a) C_1-C_9 alkyl, linear, branched, or cyclic, optionally substituted with a hydroxyl group, an amino group, a carboxylic acid group, a carboxamide group, a halogen atom, or an oxo group;

b) C_1-C_9 alkenyl, linear, branched, or cyclic, optionally substituted with a hydroxyl group, an amino group, a carboxylic acid group, a carboxamide group, a halogen atom, or an oxo group; or

c) C_1-C_9 aryl, optionally substituted with a hydroxyl group, an amino group, a carboxylic acid group, a carboxamide group, or a halogen atom.

31. The method of Claim 30 wherein the nitrile is 2-hydroxynitrile, 3-hydroxynitrile, or 4-hydroxynitrile.

32. The method of Claim 30 wherein the R of Formula 2 is C_1-C_4 alkyl, linear, or branched.

33. The method of Claim 32 wherein the nitrile is selected from the group consisting of malononitrile, adiponitrile, glutaronitrile, and 2-methylglutaronitrile.

34. The method of Claim 30 wherein the R of Formula 1 is C_1-C_4 alkenyl, linear, or branched.

35. The method of Claim 34 wherein the nitrile is acrylonitrile or methacrylonitrile.

36. A method for hydrating methacrylonitrile to methacrylamide comprising

a) contacting methacrylonitrile, under suitable reaction conditions, with a catalyst having nitrile hydratase activity from *Comamonas testosteroni* 5-MGAM-4D.

37. The method of Claim 36, further comprising

b) recovering the methacrylamide produced in step a).

38. A method of hydrating acrylonitrile to acrylamide comprising

a) contacting acrylonitrile, under suitable reaction conditions, with a catalyst having nitrile hydratase activity from *Comamonas testosteroni* 5-MGAM-4D.

39. The method of Claim 38, further comprising

b) recovering the acrylamide formed in step a).

40. The method of Claims 29, 36, or 38, wherein the catalyst is in the form of whole cells, permeabilized microbial cells, one or more components of a microbial cell extract, partially purified enzyme, or purified enzyme.


41. The method of Claim 40, wherein the catalyst is immobilized on or in a soluble or insoluble support.

42. The method of Claim 41, wherein the catalyst is immobilized in alginate or carrageenan.


43. The method of Claim 29 wherein the substrate comprises at least one nitrile functional group and is a nitrile of:

Formula 3 $R^2-C\equiv N$


wherein N is Nitrogen, C is Carbon, and R^2 is selected from the group consisting of the general formulae 4, 5, 6, 7, and 8:




Formula 4




Formula 5



Formula 6



Formula 7



Formula 8

wherein in Formulae 4, 5, and 6 X is N, wherein in Formula 7 X is NH, O or S, and wherein in Formula 8 X is NH and Y is N, O, or S, and where R^3 and R^4 are independently selected from the group comprised of a) a hydrogen atom, b) a halogen atom, c) a C_1 - C_9 alkyl group that is linear, branched or cyclic and optionally substituted with a

hydroxyl group, an amino group, a carboxylic acid group, a carboxamide group, a halogen atom, or an oxo group, d) a C₁-C₉ alkenyl group that is linear, branched, or cyclic and optionally substituted with a hydroxyl group, an amino group, a carboxylic acid group, a carboxamide group, a halogen atom, or an oxo group, and e) a C₆-C₉ aryl group, optionally substituted with a hydroxyl group, an amino group, a carboxylic acid group, a carboxamide group, or a halogen atom.

44. The method of Claim 43 wherein the nitrile is 3-cyanopyridine, 4-cyanopyridine, pyrazinecarbonitrile, 2-furancarbonitrile, 2-thiophenecarbonitrile, or 4-thiazolecarbonitrile.

45. The method of Claim 43 wherein R³ and R⁴ are independently of a hydrogen atom.

46. The method of Claim 43 wherein R³ and R⁴ are independently of a halogen atom.

47. The method of Claim 43 wherein R³ and R⁴ are independently of a C₁-C₉ alkyl group that is linear, branched, or cyclic and optionally substituted with a hydroxyl group, an amino group, a carboxylic acid group, a carboxamide group, a halogen atom, or an oxo group.

48. The method of Claim 43 wherein R³ and R⁴ are independently of a C₁-C₉ alkenyl group that is linear, branched, or cyclic and optionally substituted with a hydroxyl group, an amino group, a carboxylic acid group, a carboxamide group, a halogen atom, or an oxo group.

49. The method of Claim 43 wherein R³ and R⁴ are independently of a C₆-C₉ aryl group, optionally substituted with a hydroxyl group, an amino group, a carboxylic acid group, a carboxamide group, or a halogen atom.

50. A method for hydrating 3-cyanopyridine to nicotinamide comprising

- a) contacting 3-cyanopyridine, under suitable reaction conditions, with a catalyst having nitrile hydratase activity from *Comamonas testosteroni* 5-MGAM-4D.

51. The method of Claim 50, further comprising

- b) recovering the nicotinamide produced in step a).

52. A method of hydrating pyrazinecarbonitrile to pyrazinecarboxamide comprising

- a) contacting pyrazinecarbonitrile, under suitable reaction conditions, with a catalyst having nitrile hydratase activity from *Comamonas testosteroni* 5-MGAM-4D.


53. The method of Claim 52, further comprising

- b) recovering the pyrazinecarboxamide formed in step a).

54. The method of Claims 43, 50, or 52, wherein the catalyst is in the form of whole cells, permeabilized microbial cells, one or more components of a microbial cell extract, partially purified enzyme, or purified enzyme.

55. The method of Claims 43, 50, or 52 wherein the catalyst is immobilized on or in a soluble or insoluble support.

56. The method of Claims 43, 50, or 52 wherein the catalyst is immobilized in alginate or carrageenan.



SEQUENCE LISTING

<110> E.I. duPont de Nemours and Company, Inc .

<120> Nucleic Acid Fragments Encoding Nitrile Hydratase and Amidase Enzymes from Comamonas Testosteroni 5-MGAM-4D and Recombinant Organisms Expressing Those Enzymes Useful for the Production of Amides and Acids

<130> CL-2200 US PCT1

<150> US 10/431966

<151> 2003-05-08

<160> 23

<170> PatentIn version 3.3

<210> 1

<211> 633

<212> DNA

<213> Pseudomonas putida 5B

<400> 1

```

atgggggaat cacacacgca tgaccacat cagcaggggt acCaggcacc gcccgaagac      60
atcgcgctgc ggggtcaaggc cttggagtct ctgctgatcg agaaaggtct tgtcgaccca      120
gcggccatgg acttggtcgt ccaaactat gaacacaagg taGggccccc aaacggcgcc      180
aaagtcgtgg ccaaggcctg ggtggaccct gcctacaagg ccCgtctgct ggcagacgca      240
actgcggaac ttgccgagct gggcttctcc ggggtacagg gcGaggacat ggtcattctg      300
gaaaacaccc ccgccgtcca caacgtcttc gtttgacact tgTgctcttg ctacccatgg      360
ccgacgctgg gcttgcccc tgccctgtac aaggccgccg ccTaccgggc ccgcatggtg      420
agcgacccgc gtgggggttct cgcgaggttc ggcctggtga tcCccgcca caaggaaatc      480
cgcgctctggg acaccacggc cgaattgcgc tacatggtgc tgCcggaacg gcccggaact      540
gaagcctaca gcgaagaaca actggccgaa ctcgttaccc gcGattcgat gatcggcacc      600
ggcctgcca cccaaccac cccatctcat taa                                     633

```

<210> 2

<211> 654

<212> DNA

<213> Pseudomonas putida 5B

<400> 2

```

atgaatggca ttcacgatac tggcggagca catggttatg ggCcggttta cagagaaccg      60
aacgaacccg tctttcgcta cgactgggaa aaaacgggtca tgTccctgct cccggccctg      120
ctcgccaacg cgaacttcaa cctcgatgaa tttcggcatt cgatcgagcg aatgggcccg      180
gccactatc tggaggggaa ctactacgaa cactggcttc atGtctttga gaacctgctg      240
gtcgagaagg gtgtgtcac ggccacggaa gtcgcgaccg gcAaggctgc gtctggcaag      300
acggcgacgc gcgtgtgac gccggccatc gtggacgact cgTcagcacc ggggcttctg      360
cgcccgggag gaggggttct ttttttctc gtgggggaca agGttcgcgt cctcaacaag      420

```

aacccggtgg gccatacccg catgccgcgc tacacgcggg caaagtgggg acagtgggtca 480
 tcgaccatgg tgtgtttcgt gacgccggac accgcggcac acggaaaggg cgagcagccc 540
 cagcacgttt acaccgtgag ttccacgtcg gtcgaactgt gggggcaaga cgcttcctcg 600
 ccgaaggaca cgattcgcgt cgacttgtgg gatgactacc tggagccagc gtga 654

<210> 3
 <211> 633
 <212> DNA
 <213> Comamonas testosteroni 5-MGAM-4D

<220>
 <221> CDS
 <222> (1)..(633)

<400> 3
 atg ggg caa tca cac acg cat gac cac cat cac gac ggg tac cag gca 48
 Met Gly Gln Ser His Thr His Asp His His His Asp Gly Tyr Gln Ala
 1 5 10 15
 ccg ccc gaa gac atc gcg ctg cgg gtc aag gcc ttg gag tct ctg ctg 96
 Pro Pro Glu Asp Ile Ala Leu Arg Val Lys Ala Leu Glu Ser Leu Leu
 20 25 30
 atc gag aaa ggt ctt gtc gac cca gcg gcc atg gac ttg gtc gtc caa 144
 Ile Glu Lys Gly Leu Val Asp Pro Ala Ala Met Asp Leu Val Val Gln
 35 40 45
 acg tat gaa cac aag gta ggc ccc cga aac ggc gcc aaa gtc gtg gcc 192
 Thr Tyr Glu His Lys Val Gly Pro Arg Asn Gly Ala Lys Val Val Ala
 50 55 60
 aag gcc tgg gtg gac cct gcc tac aag gcc cgt ctg ctg gca gac ggc 240
 Lys Ala Trp Val Asp Pro Ala Tyr Lys Ala Arg Leu Leu Ala Asp Gly
 65 70 75 80
 act gcc ggc att gcc gag ctg ggc ttc tcc ggg gta cag ggc gag gac 288
 Thr Ala Gly Ile Ala Glu Leu Gly Phe Ser Gly Val Gln Gly Glu Asp
 85 90 95
 atg gtc att ctg gaa aac acc ccc gcc gtc cac aac gtc gtc gtt tgc 336
 Met Val Ile Leu Glu Asn Thr Pro Ala Val His Asn Val Val Val Cys
 100 105 110
 acc ttg tgc tct tgc tac cca tgg ccg acg ctg ggc ttg ccc cct gcc 384
 Thr Leu Cys Ser Cys Tyr Pro Trp Pro Thr Leu Gly Leu Pro Pro Ala
 115 120 125
 tgg tac aag gcc ccg ccc tac cgg tcc cgc atg gtg agc gac ccg cgt 432
 Trp Tyr Lys Ala Pro Pro Tyr Arg Ser Arg Met Val Ser Asp Pro Arg
 130 135 140
 ggg gtt ctc gcg gag ttc ggc ctg gtg atc ccc gcg aag gaa atc cgc 480
 Gly Val Leu Ala Glu Phe Gly Leu Val Ile Pro Ala Lys Glu Ile Arg
 145 150 155 160
 gtc tgg gac acc acg gcc gaa ttg cgc tac atg gtg ctg ccg gaa cgg 528
 Val Trp Asp Thr Thr Ala Glu Leu Arg Tyr Met Val Leu Pro Glu Arg
 165 170 175
 ccc gcg gga act gaa gcc tac agc gaa gaa caa ctg gcc gaa ctc gtt 576
 Pro Ala Gly Thr Glu Ala Tyr Ser Glu Glu Gln Leu Ala Glu Leu Val
 180 185 190

acc cgc gat tcg atg atc ggc acc ggc ctg ccc atc caa ccc acc cca 624
 Thr Arg Asp Ser Met Ile Gly Thr Gly Leu Pro Ile Gln Pro Thr Pro
 195 200 205

tct cat taa 633
 Ser His
 210

<210> 4
 <211> 210
 <212> PRT
 <213> Comamonas testosteroni 5-MGAM-4D

<400> 4

Met Gly Gln Ser His Thr His Asp His His His Asp Gly Tyr Gln Ala
 1 5 10 15

Pro Pro Glu Asp Ile Ala Leu Arg Val Lys Ala Leu Glu Ser Leu Leu
 20 25 30

Ile Glu Lys Gly Leu Val Asp Pro Ala Ala Met Asp Leu Val Val Gln
 35 40 45

Thr Tyr Glu His Lys Val Gly Pro Arg Asn Gly Ala Lys Val Val Ala
 50 55 60

Lys Ala Trp Val Asp Pro Ala Tyr Lys Ala Arg Leu Leu Ala Asp Gly
 65 70 75 80

Thr Ala Gly Ile Ala Glu Leu Gly Phe Ser Gly Val Gln Gly Glu Asp
 85 90 95

Met Val Ile Leu Glu Asn Thr Pro Ala Val His Asn Val Val Val Cys
 100 105 110

Thr Leu Cys Ser Cys Tyr Pro Trp Pro Thr Leu Gly Leu Pro Pro Ala
 115 120 125

Trp Tyr Lys Ala Pro Pro Tyr Arg Ser Arg Met Val Ser Asp Pro Arg
 130 135 140

Gly Val Leu Ala Glu Phe Gly Leu Val Ile Pro Ala Lys Glu Ile Arg
 145 150 155 160

Val Trp Asp Thr Thr Ala Glu Leu Arg Tyr Met Val Leu Pro Glu Arg
 165 170 175

Pro Ala Gly Thr Glu Ala Tyr Ser Glu Glu Gln Leu Ala Glu Leu Val
 180 185 190

Thr Arg Asp Ser Met Ile Gly Thr Gly Leu Pro Ile Gln Pro Thr Pro
 195 200 205

Ser His
210

<210> 5
<211> 657
<212> DNA
<213> Comamonas testosteroni 5-MGAM-4D

<220>
<221> CDS
<222> (1)..(657)

<400> 5
atg aat ggc att cac gat act ggg gga gca cat ggt tat ggg ccg gtt 48
Met Asn Gly Ile His Asp Thr Gly Gly Ala His Gly Tyr Gly Pro Val
1 5 10 15

tac aga gaa ccg aac gaa ccc gtc ttt cgc tac gac tgg gaa aaa acg 96
Tyr Arg Glu Pro Asn Glu Pro Val Phe Arg Tyr Asp Trp Glu Lys Thr
20 25 30

gtc atg tcc ctg ttc ccg gcg ctg ttc gcc aac ggc aac ttc aac ctg 144
Val Met Ser Leu Phe Pro Ala Leu Phe Ala Asn Gly Asn Phe Asn Leu
35 40 45

gat gag ttt cga cac ggc atc gag cgc atg aac ccc atc gac tac ctg 192
Asp Glu Phe Arg His Gly Ile Glu Arg Met Asn Pro Ile Asp Tyr Leu
50 55 60

aag gga acc tac tac gaa cac tgg atc cat tcc atc gaa acc ttg ctg 240
Lys Gly Thr Tyr Tyr Glu His Trp Ile His Ser Ile Glu Thr Leu Leu
65 70 75 80

gtc gaa aag ggt gtg ctc acg gca acg gaa ctc gcg acc ggc aag gca 288
Val Glu Lys Gly Val Leu Thr Ala Thr Glu Leu Ala Thr Gly Lys Ala
85 90 95

tct ggc aag aca gcg aca ccg gtg ctg acg ccg gcc atc gtg gac gga 336
Ser Gly Lys Thr Ala Thr Pro Val Leu Thr Pro Ala Ile Val Asp Gly
100 105 110

ctg ctc agc acc ggg gct tct gcc gcc cgg gag gag ggt gcg cgg gcg 384
Leu Leu Ser Thr Gly Ala Ser Ala Ala Arg Glu Glu Gly Ala Arg Ala
115 120 125

cgg ttc gct gtg ggg gac aag gtt cgc gtc ctc aac aag aac ccg gtg 432
Arg Phe Ala Val Gly Asp Lys Val Arg Val Leu Asn Lys Asn Pro Val
130 135 140

ggc cat acc cgc atg ccg cgc tac acg cgg ggc aaa gtg ggg aca gtg 480
Gly His Thr Arg Met Pro Arg Tyr Thr Arg Gly Lys Val Gly Thr Val
145 150 155 160

gtc atc gac cat ggt gtg ttc gtg acg ccg gac acc gcg gca cac gga 528
Val Ile Asp His Gly Val Phe Val Thr Pro Asp Thr Ala Ala His Gly
165 170 175

aag ggc gag cac ccc cag cac gtt tac acc gtg agt ttc acg tcg gtc 576
Lys Gly Glu His Pro Gln His Val Tyr Thr Val Ser Phe Thr Ser Val
180 185 190

gaa ctg tgg ggg caa gac gcc tcc tcg ccg aag gac acg att cgc gtc 624
Glu Leu Trp Gly Gln Asp Ala Ser Ser Pro Lys Asp Thr Ile Arg Val
195 200 205

gac ttg tgg gat gac tac ctg gag cca gcg tga 657

Asp Leu Trp Asp Asp Tyr Leu Glu Pro Ala
210 215

<210> 6

<211> 218

<212> PRT

<213> Comamonas testosteroni 5-MGAM-4D

<400> 6

Met Asn Gly Ile His Asp Thr Gly Gly Ala His Gly Tyr Gly Pro Val
1 5 10 15

Tyr Arg Glu Pro Asn Glu Pro Val Phe Arg Tyr Asp Trp Glu Lys Thr
20 25 30

Val Met Ser Leu Phe Pro Ala Leu Phe Ala Asn Gly Asn Phe Asn Leu
35 40 45

Asp Glu Phe Arg His Gly Ile Glu Arg Met Asn Pro Ile Asp Tyr Leu
50 55 60

Lys Gly Thr Tyr Tyr Glu His Trp Ile His Ser Ile Glu Thr Leu Leu
65 70 75 80

Val Glu Lys Gly Val Leu Thr Ala Thr Glu Leu Ala Thr Gly Lys Ala
85 90 95

Ser Gly Lys Thr Ala Thr Pro Val Leu Thr Pro Ala Ile Val Asp Gly
100 105 110

Leu Leu Ser Thr Gly Ala Ser Ala Ala Arg Glu Glu Gly Ala Arg Ala
115 120 125

Arg Phe Ala Val Gly Asp Lys Val Arg Val Leu Asn Lys Asn Pro Val
130 135 140

Gly His Thr Arg Met Pro Arg Tyr Thr Arg Gly Lys Val Gly Thr Val
145 150 155 160

Val Ile Asp His Gly Val Phe Val Thr Pro Asp Thr Ala Ala His Gly
165 170 175

Lys Gly Glu His Pro Gln His Val Tyr Thr Val Ser Phe Thr Ser Val
180 185 190

Glu Leu Trp Gly Gln Asp Ala Ser Ser Pro Lys Asp Thr Ile Arg Val
195 200 205

Asp Leu Trp Asp Asp Tyr Leu Glu Pro Ala
210 215

<210> 7

<211> 1386

<212> DNA

<213> Comamonas testosteroni 5-MGAM-4D

<400> 7

tcgatgccga gttgaagtcg ctgtaccct tttttcaacc acaccaggag aaccgcacca	60
tggggcaatc acacacgcat gaccaccatc acgacgggta ccaggcaccg cccgaagaca	120
tcgcgctgcg ggtcaaggcc ttggagtctc tgctgatcga gaaagggtctt gtcgaccag	180
cggccatgga cttggtcgctc caaacgtatg aacacaaggt agggccccga aacggcgcca	240
aagtcgtggc caaggcctgg gtggaccctg cctacaaggc ccgtctgctg gcagacggca	300
ctgccggcat tgccgagctg ggcttctccg gggtagagg cgaggacatg gtcattctgg	360
aaaacacccc cgccgtccac aacgtcgtcg tttgcacctt gtgctcttgc taccatggc	420
cgacgtggg cttgccccct gcctggtaga agggcccgcc ctaccgggtcc cgcattgtga	480
gcgaccgcg tggggttctc gcggagtctg gcctgggtat ccccggaag gaaatccgcg	540
tctgggacac cacggccgaa ttgcgtaca tgggtgctgcc ggaacggccc gcgggaactg	600
aagcctacag cgaagaacaa ctggccgaac tcgttaccg cgattcgatg atcggcaccg	660
gcctgcccac ccaaccacc ccatctcatt aaggagttcg tcatgaatgg cattcacgat	720
actgggggag cacatggtta tgggccggtt tacagagaac cgaacgaacc cgtctttcgc	780
tacgactggg aaaaaacggt catgtccctg ttcccggcgc tggtcgccaa cggcaacttc	840
aacctcgatg agtttcgaca cggcatcgag cgcattgaacc ccatcgacta cctgaaggga	900
acctactacg aacctggat ccattccatc gaaaccttgc tggtcgaaaa ggggtgtgctc	960
acggcaacgg aactcgcgac cggcaaggca tctggcaaga cagcgacacc ggtgctgacg	1020
ccggccatcg tggacggact gctcagcacc ggggcttctg ccgcccggga ggaggggtcg	1080
cgggcgcggt tcgctgtggg ggacaagggt cgcgctctca acaagaaccc ggtgggcat	1140
acccgcatgc cgcgtacac gcggggcaaa gtggggacag tggcatcga ccatggtgtg	1200
ttcgtgacgc cggacaccgc ggcacacgga aagggcgagc acccccagca cgtttacacc	1260
gtgagtttca cgtcggtcga actgtggggg caagacgcct cctcgccgaa ggacacgatt	1320
cgcgtcgact tgtgggatga ctacctggag ccagcgtgat catgaaagac gaacggtttc	1380
cattgc	1386

<210> 8

<211> 23

<212> DNA

<213> artificial sequence

<220>

<223> Primer 1

<400> 8

tcgatgccga gttgaagtcg ctg	23
---------------------------	----

<210> 9

<211> 23

<212> DNA
 <213> artificial sequence

<220>
 <223> Primer 2

<400> 9
 gcaatggaaa ccgttcgtct ttc 23

<210> 10
 <211> 923
 <212> DNA
 <213> Comamonas testosteroni 5-MGAM-4D

<400> 10
 tcatgaaaga cgaacggttt ccattgccag agggttcgct gaaggacctc gatggccctg 60
 tgtttgacga gccttggcag tcccaggcgt ttgccttggg ggtcagcatg cacaaggccg 120
 gtctctttca gtggaaagac tggggccgaga ctttcaccgc cgaaatcgac gcttccccgg 180
 ctctgcccgg cgaaagcgct aacgacacct actaccggca atgggtgtcg gcgctggaaa 240
 agttggtggc gtcgctgggg cttgtgacgg gtggagacgt caactcgcg gcacaggagt 300
 ggaaacaggc ccacctcaac accccacatg ggcacccgat cctgctggcc catgcgcttt 360
 gcccgccagc gatcgacccc aagcacaagc acgagccaaa acgctcaccg atcaaggctc 420
 ttgccgcaat ggcttgagat ccactgtcct gtttccttac cttgaatgga gtaaaccatg 480
 agttcatttt ctaccactgc cgtccccgcc gccagcgcc tgagcgccac gcgcagctta 540
 ctgttgcaac tgtccgctgg cgccgcgctg ggcttggtcg tcctgtacgg cgtggctttt 600
 gccgaaagcc cgcttgcgca caacgccg cgacgatgttc gccacgtgac ggtcaagcct 660
 tgtcactaac tgtggcccac ccccgccg cgctgcgcga cccctccag ggggcaaacc 720
 gagtggcccc gcggagccgg ttccaccg cgctccggtg ggtgcaactg tggcgcttgcg 780
 ggtcctgttt cggttgatg gtttcttgac ggagccacac cgtcgctcat ctgtatatgg 840
 agaggggtcat gatttttcga cgcttgatct gggcggcgct ggccgtggcc ttgctggtgg 900
 gcagtttgca gtcgggtctg cag 923

<210> 11
 <211> 2223
 <212> DNA
 <213> Comamonas testosteroni 5-MGAM-4D

<400> 11
 atggggcaat cacacacgca tgaccaccat cagcaggggt accaggcacc gcccgaagac 60
 atcgcgctgc ggtcaaggc cttggagtct ctgctgatcg agaaaggctt tgtcgaccca 120
 gcggccatgg acttggtcgt ccaaacgtat gaacacaagg taggcccccg aaacggcgcc 180
 aaagtcgtgg ccaaggcctg ggtggaccct gcctacaagg cccgtctgct ggcagacggc 240
 actgccggca ttgccgagct gggcttctcc ggggtacagg gcgaggacat ggtcattctg 300
 gaaaacaccc ccgccgtcca caacgtcgtc gtttgacact tgtgctcttg ctacccatgg 360
 ccgacgctgg gcttgcccc tgcttggtac aaggccccgc cctaccggtc ccgcatgggtg 420

```

agcgacccgc gtgggggttct cgcgagggttc ggcctggtga tccccgcgaa ggaaatccgc      480
gtctgggaca ccacggccga attgcgctac atggtgctgc cggaacggcc cgcggggaact      540
gaagcctaca gcgaagaaca actggccgaa ctcgttaccc gcgattcgat gatcggcacc      600
ggcctgcccc tccaaccac cccatctcat taaggagttc gtcatgaatg gcattcacga      660
tactggggga gcacatggtt atggggccggt ttacagagaa ccgaacgaac ccgtctttcg      720
ctacgactgg gaaaaaacgg tcatgtccct gtccccggcg ctgttcgcca acggcaactt      780
caacctcgat gagtttcgac acggcatcga gcgcatgaac cccatcgact acctgaaggg      840
aacctactac gaactcgga tccattccat cgaaaccttg ctggtcgaaa aggggtgtgct      900
cacggcaacg gaactcgga ccggcaaggc atctggcaag acagcgacac cggtgctgac      960
gccggccatc gtggacggac tgctcagcac cggggcttct gccgcccggg aggaggggtgc     1020
gcgggcgcgg ttcgctgtgg gggacaaggt tcgcgtcctc aacaagaacc cggtgggcca     1080
taccgcgatg ccgcgctaca cgcggggcaa agtggggaca gtggtcatcg accatggtgt     1140
gttcgtgacg ccggacaccg cggcacacgg aaagggcgag cacccccagc acgtttacac     1200
cgtgagtttc acgtcggtcg aactgtgggg gcaagacgcc tcctcgccga aggacacgat     1260
tcgcgtcgac ttgtgggatg actacctgga gccagcgtga tcatgaaaga cgaacggttt     1320
ccattgccag agggttcgct gaaggacctc gatggccctg tgtttgacga gccttggcag     1380
tcccaggcgt ttgccttggg ggtcagcatg cacaaggccg gtctctttca gtggaaagac     1440
tggggccgaga ctttcaccgc cgaaatcgac gttccccgg ctctgcccgg cgaaagcgtc     1500
aacgacacct actaccggca atgggtgtcg gcgctggaaa agttggtggc gtcgctgggg     1560
cttgtgacgg gtggagacgt caactcgcg gcacaggagt ggaaacaggc ccacctcaac     1620
acccacatg ggcacccgat cctgctggcc catgcgcttt gcccgccagc gatcgacccc     1680
aagcacaagc acgagccaaa acgctcaccg atcaaggctg ttgccgcaat ggcttgagat     1740
ccactgtcct gtttccctac cttgaatgga gtaaaccatg agttcatttt ctaccactgc     1800
cgtccccgcc gccagcgcc tgagcgccac gcgcagctta ctgttgcaac tgtccgctgg     1860
cgccgcgctg ggcctggctg tcctgtacgg cgtggctttt gccgaaagcc cgcttgcgca     1920
caacgccgcg cacgatgttc gccacgtgac ggtcaagcct tgtcactaac tgtggcccac     1980
ccccgccgcg ctgcgcgga cccctccag ggggcaaacc gagtggcccg gcggagccgg     2040
ttccaccgcg ttcccggtgg ggtgcaactgc tggcgttgcg ggtcttgttt cggctggatg     2100
gtttcttgac ggagccacac cgtcgctcat ctgtatatgg agaggggtcat gatttttcga     2160
cgcttgatct gggcggcgct ggccgtggcc ttgctgggtg gcagtttgca gtcgggtctg     2220
cag                                                                    2223

```

<210> 12
 <211> 30
 <212> DNA
 <213> artificial sequence

<220>
 <223> Primer 3

<400> 12
 ctgaagcttc aaggtaggga aacaggacag 30

<210> 13
 <211> 216
 <212> DNA
 <213> Comamonas testosteroni 5-MGAM-4D

<220>
 <221> CDS
 <222> (1)..(216)

<400> 13
 atg gcc ctg tgt ttg acg agc ctt ggc agt ccc agg cgt ttg cct tgg 48
 Met Ala Leu Cys Leu Thr Ser Leu Gly Ser Pro Arg Arg Leu Pro Trp
 1 5 10 15
 tgg tca gca tgc aca agg ccg gtc tct ttc agt gga aag act ggg ccg 96
 Trp Ser Ala Cys Thr Arg Pro Val Ser Phe Ser Gly Lys Thr Gly Pro
 20 25 30
 aga cct tca ccg ccg aaa tcg acg ctt ccc cgg ctc tgc ccg gcg aaa 144
 Arg Pro Ser Pro Pro Lys Ser Thr Leu Pro Arg Leu Cys Pro Ala Lys
 35 40 45
 gcg tca acg aca cct act acc ggc aat ggg tgt cgg cgc tgg aaa agt 192
 Ala Ser Thr Thr Pro Thr Thr Gly Asn Gly Cys Arg Arg Trp Lys Ser
 50 55 60
 tgg tgg cgt cgc tgg ggc ttg tga 216
 Trp Trp Arg Arg Trp Gly Leu
 65 70

<210> 14
 <211> 71
 <212> PRT
 <213> Comamonas testosteroni 5-MGAM-4D

<400> 14
 Met Ala Leu Cys Leu Thr Ser Leu Gly Ser Pro Arg Arg Leu Pro Trp
 1 5 10 15
 Trp Ser Ala Cys Thr Arg Pro Val Ser Phe Ser Gly Lys Thr Gly Pro
 20 25 30
 Arg Pro Ser Pro Pro Lys Ser Thr Leu Pro Arg Leu Cys Pro Ala Lys
 35 40 45
 Ala Ser Thr Thr Pro Thr Thr Gly Asn Gly Cys Arg Arg Trp Lys Ser
 50 55 60
 Trp Trp Arg Arg Trp Gly Leu
 65 70

<210> 15

<211> 669
 <212> DNA
 <213> Comamonas testosteroni 5-MGAM-4D

<400> 15
 gaattcgcct atggcgatcat cactccgaag tcgcgcaacc cctgggaccc gggagaaca 60
 ccgggtggct ccagcggcgg ctcggcggcc acggtcgcag cctgcggcgt ctacttggcg 120
 accggcaccg acaccggtgg atccgttcgc atcccttcgt cgatgtgcaa caccgtaggc 180
 ctgaagccaa cctacgggcg cgtgagccgt gccgggtgtga gttcactttc ctggagcctg 240
 gaccatccag gcccgatcac gcgcaccgtg gaagacacgg cgctcagcct tcagggtgatg 300
 gctggcttcg atccagccga ccgcgggtcg ttggatgagc cggtgcccag ctatgccgaa 360
 gggctcggcc aaggcgtgaa aggcctgcgc gtgggcgtgc cgaagaacta cttcttcgac 420
 cgcgtaggacc cggaagttga aagtgcggtt cgtgccgcca tcgatcaact gaaagagctg 480
 ggcgccgaac tgggtggaagt cgaagtgcgc atggccgagc agatcatccc ggtggagttc 540
 gggatcgtgc taccgaagc cagcgcctac caccgcacga tgctgcgcga gtcacccgag 600
 ctctacaccg ccgatgtccg catactgctg gaactcggaa atctagtcac cgccaccgac 660
 tacctgcag 669

<210> 16
 <211> 1407
 <212> DNA
 <213> Comamonas testosteroni 5-MGAM-4D

<220>
 <221> CDS
 <222> (1)..(1407)

<400> 16
 atg agt tcg cta acc cgc ctc acc ctc gcg caa gtt gcg cag aaa ctt 48
 Met Ser Ser Leu Thr Arg Leu Thr Leu Ala Gln Val Ala Gln Lys Leu
 1 5 10 15
 aag gca cgg gaa gtc tcc gcc gtt gaa gtt ctg gac gcc tgt ctg acg 96
 Lys Ala Arg Glu Val Ser Ala Val Glu Val Leu Asp Ala Cys Leu Thr
 20 25 30
 cag gtg cgc tcc acc gaa aaa cag atc agt gcg tac gtg tgc gtg ctg 144
 Gln Val Arg Ser Thr Glu Lys Gln Ile Ser Ala Tyr Val Cys Val Leu
 35 40 45
 gag gat cag gcc cgt gca gca gcc cag caa gct gac gcc gac atc agc 192
 Glu Asp Gln Ala Arg Ala Ala Ala Gln Gln Ala Asp Ala Asp Ile Ser
 50 55 60
 gcc ggg cgc tgg aaa ggc ccg ctg cat ggc gtg cct gta gcg gtc aag 240
 Ala Gly Arg Trp Lys Gly Pro Leu His Gly Val Pro Val Ala Val Lys
 65 70 75 80
 gac tta tac gac atc gct ggc gta ccc acc acg gca tcg tcg cgc cag 288
 Asp Leu Tyr Asp Ile Ala Gly Val Pro Thr Thr Ala Ser Ser Arg Gln
 85 90 95
 cgc acg aat tgg acg ccg cag caa gac tgc gcc gta gtc cgg cgc ttg 336
 Arg Thr Asn Trp Thr Pro Gln Gln Asp Cys Ala Val Val Arg Arg Leu
 100 105 110

aaa Lys	gac Asp	gca Ala 115	ggt Gly	gcc Ala	gtt Val	atc Ile	ctt Leu 120	ggc Gly	aag Lys	acc Thr	cat His	acg Thr 125	cac His	gaa Glu	ttc Phe	384
gcc Ala	tat Tyr 130	ggc Gly	gtc Val	atc Ile	act Thr	ccg Pro 135	aag Lys	tcg Ser	cgc Arg	aac Asn	ccc Pro 140	tgg Trp	gac Asp	ccg Pro	gga Gly	432
aga Arg 145	aca Thr	ccg Pro	ggt Gly	ggc Gly	tcc Ser 150	agc Ser	ggc Gly	ggc Gly	tcg Ser	ggc Ala 155	gcc Ala	acg Thr	gtc Val	gca Ala	gcc Ala 160	480
tgc Cys	ggc Gly	gtc Val	tac Tyr	ttg Leu 165	gcg Ala	acc Thr	ggc Gly	acc Thr	gac Asp 170	acc Thr	ggg Gly	gga Gly	tcc Ser	gtt Val 175	cgc Arg	528
atc Ile	cct Pro	tcg Ser	tcg Ser 180	atg Met	tgc Cys	aac Asn	acc Thr	gta Val 185	ggc Gly	ctg Leu	aag Lys	cca Pro	acc Thr 190	tac Tyr	ggg Gly	576
cgc Arg	gtg Val	agc Ser 195	cgt Arg	gcc Ala	ggt Gly	gtg Val	agt Ser 200	tca Ser	ctt Leu	tcc Ser	tgg Trp	agc Ser 205	ctg Leu	gac Asp	cat His	624
cca Pro	ggc Gly 210	ccg Pro	atc Ile	acg Thr	cgc Arg	acc Thr 215	gtg Val	gaa Glu	gac Asp	acg Thr	ggc Ala 220	ctc Leu	agc Ser	ctt Leu	cag Gln	672
gtg Val 225	atg Met	gct Ala	ggc Gly	ttc Phe	gat Asp 230	cca Pro	gcc Ala	gac Asp	cgc Arg	ggc Gly 235	tcg Ser	ttg Leu	gat Asp	gag Glu	ccg Pro 240	720
gtg Val	ccc Pro	agc Ser	tat Tyr	gcc Ala 245	gaa Glu	ggg Gly	ctc Leu	ggc Gly	caa Gln 250	ggc Gly	gtg Val	aaa Lys	ggc Gly 255	ctg Leu	cgc Arg	768
gtg Val	ggc Gly	gtg Val	ccg Pro 260	aag Lys	aac Asn	tac Tyr	ttc Phe	ttc Phe 265	gac Asp	cgc Arg	gtg Val	gac Asp	ccg Pro 270	gaa Glu	gtt Val	816
gaa Glu	agt Ser	gcg Ala 275	gtt Val	cgt Arg	gcc Ala	gcc Ala	atc Ile 280	gat Asp	caa Gln	ctg Leu	aaa Lys	gag Glu 285	ctg Leu	ggc Gly	gcc Ala	864
gaa Glu 290	ctg Leu	gtg Val	gaa Glu	gtc Val	gaa Glu	gtg Val 295	ccc Pro	atg Met	gcc Ala	gag Glu	cag Gln 300	atc Ile	atc Ile	ccg Pro	gtg Val	912
gag Glu 305	ttc Phe	ggg Gly	atc Ile	gtg Val	cta Leu 310	ccc Pro	gaa Glu	gcc Ala	agc Ser	gcc Ala 315	tac Tyr	cac His	cgc Arg	acg Thr	atg Met 320	960
ctg Leu	cgc Arg	gag Glu	tca Ser	ccc Pro 325	gag Glu	ctc Leu	tac Tyr	acc Thr	gcc Ala 330	gat Asp	gtc Val	cgc Arg	ata Ile	ctg Leu 335	ctg Leu	1008
gaa Glu	ctc Leu	gga Gly	aat Asn 340	cta Leu	gtc Val	acc Thr	gcc Ala	acc Thr 345	gac Asp	tac Tyr	ctg Leu	cag Gln	gcg Ala 350	cag Gln	cgc Arg	1056
gtc Val	cgt Arg	acg Thr 355	ctg Leu	atg Met	cag Gln	cgc Arg	gcg Ala 360	gtg Val	gcc Ala	gag Glu	atg Met	ttc Phe 365	cag Gln	cgc Arg	atc Ile	1104
gat Asp 370	gtg Val	ctg Leu	atc Ile	gca Ala	ccc Pro	aca Thr 375	ctg Leu	ccc Pro	atc Ile	ccg Pro	gct Ala 380	gct Ala	cgc Arg	agc Ser	ggg Gly	1152

gag gag gtc cac aca tgg ccg gac ggc acg gta gag gcg ttg ttc atg 1200
 Glu Glu Val His Thr Trp Pro Asp Gly Thr Val Glu Ala Leu Phe Met
 385 390 395 400
 gcc tat acg cgc ttc acc tcg ttc ggc aac gtg aca gga tta ccc acg 1248
 Ala Tyr Thr Arg Phe Thr Ser Phe Gly Asn Val Thr Gly Leu Pro Thr
 405 410 415
 ctg aac ctg ccc tgt ggt ttc tcc aag gat ggg ttg ccg atc ggc atg 1296
 Leu Asn Leu Pro Cys Gly Phe Ser Lys Asp Gly Leu Pro Ile Gly Met
 420 425 430
 cag atc acc ggc cgg ccg ctg gac gag aag acc ctg ctg cgt gct ggg 1344
 Gln Ile Thr Gly Arg Pro Leu Asp Glu Lys Thr Leu Leu Arg Ala Gly
 435 440 445
 ctg gcc tac gag aaa gcc acg acc tgg cac cag cgt cat ccg gaa ctg 1392
 Leu Ala Tyr Glu Lys Ala Thr Thr Trp His Gln Arg His Pro Glu Leu
 450 455 460
 atc gga gcg ggc tga 1407
 Ile Gly Ala Gly
 465

<210> 17
 <211> 468
 <212> PRT
 <213> Comamonas testosteroni 5-MGAM-4D

<400> 17

Met Ser Ser Leu Thr Arg Leu Thr Leu Ala Gln Val Ala Gln Lys Leu
 1 5 10 15
 Lys Ala Arg Glu Val Ser Ala Val Glu Val Leu Asp Ala Cys Leu Thr
 20 25 30
 Gln Val Arg Ser Thr Glu Lys Gln Ile Ser Ala Tyr Val Cys Val Leu
 35 40 45
 Glu Asp Gln Ala Arg Ala Ala Ala Gln Gln Ala Asp Ala Asp Ile Ser
 50 55 60
 Ala Gly Arg Trp Lys Gly Pro Leu His Gly Val Pro Val Ala Val Lys
 65 70 75 80
 Asp Leu Tyr Asp Ile Ala Gly Val Pro Thr Thr Ala Ser Ser Arg Gln
 85 90 95
 Arg Thr Asn Trp Thr Pro Gln Gln Asp Cys Ala Val Val Arg Arg Leu
 100 105 110
 Lys Asp Ala Gly Ala Val Ile Leu Gly Lys Thr His Thr His Glu Phe
 115 120 125
 Ala Tyr Gly Val Ile Thr Pro Lys Ser Arg Asn Pro Trp Asp Pro Gly
 130 135 140

Arg Thr Pro Gly Gly Ser Ser Gly Gly Ser Ala Ala Thr Val Ala Ala
 145 150 155 160
 Cys Gly Val Tyr Leu Ala Thr Gly Thr Asp Thr Gly Gly Ser Val Arg
 165 170 175
 Ile Pro Ser Ser Met Cys Asn Thr Val Gly Leu Lys Pro Thr Tyr Gly
 180 185 190
 Arg Val Ser Arg Ala Gly Val Ser Ser Leu Ser Trp Ser Leu Asp His
 195 200 205
 Pro Gly Pro Ile Thr Arg Thr Val Glu Asp Thr Ala Leu Ser Leu Gln
 210 215 220
 Val Met Ala Gly Phe Asp Pro Ala Asp Arg Gly Ser Leu Asp Glu Pro
 225 230 235 240
 Val Pro Ser Tyr Ala Glu Gly Leu Gly Gln Gly Val Lys Gly Leu Arg
 245 250 255
 Val Gly Val Pro Lys Asn Tyr Phe Phe Asp Arg Val Asp Pro Glu Val
 260 265 270
 Glu Ser Ala Val Arg Ala Ala Ile Asp Gln Leu Lys Glu Leu Gly Ala
 275 280 285
 Glu Leu Val Glu Val Glu Val Pro Met Ala Glu Gln Ile Ile Pro Val
 290 295 300
 Glu Phe Gly Ile Val Leu Pro Glu Ala Ser Ala Tyr His Arg Thr Met
 305 310 315 320
 Leu Arg Glu Ser Pro Glu Leu Tyr Thr Ala Asp Val Arg Ile Leu Leu
 325 330 335
 Glu Leu Gly Asn Leu Val Thr Ala Thr Asp Tyr Leu Gln Ala Gln Arg
 340 345 350
 Val Arg Thr Leu Met Gln Arg Ala Val Ala Glu Met Phe Gln Arg Ile
 355 360 365
 Asp Val Leu Ile Ala Pro Thr Leu Pro Ile Pro Ala Ala Arg Ser Gly
 370 375 380
 Glu Glu Val His Thr Trp Pro Asp Gly Thr Val Glu Ala Leu Phe Met
 385 390 395 400
 Ala Tyr Thr Arg Phe Thr Ser Phe Gly Asn Val Thr Gly Leu Pro Thr
 405 410 415

Leu Asn Leu Pro Cys Gly Phe Ser Lys Asp Gly Leu Pro Ile Gly Met
 420 425 430

Gln Ile Thr Gly Arg Pro Leu Asp Glu Lys Thr Leu Leu Arg Ala Gly
 435 440 445

Leu Ala Tyr Glu Lys Ala Thr Thr Trp His Gln Arg His Pro Glu Leu
 450 455 460

Ile Gly Ala Gly
 465

<210> 18

<211> 7415

<212> DNA

<213> Comamonas testosteroni 5-MGAM-4D

<400> 18

ctgcagagtg catggcatcg agctggcggg cgctccagct gctgggggttg cgccgcatgc	60
cccacagcag ctgcctgagc gtcttgcggc cgccagcgcc cagcgcggtt cgacccgcct	120
gcgcgtcggt ggccatctcc gcgcggcgca cctgggtccat cgcttcgatg gccatcgaga	180
cgacgtgaaa gcggtcgtag ctgatctgcy cctcgggcag cgccagcgcc acgccccttg	240
cgtaggctgc gctcatgtcc atgcacacgt gccgcacctg ggcgggatcg ccgccatggg	300
ccttcagatc ctcggcgaac tccaccaccg tgcggtgctc gcgcccctcg gtggcgaaca	360
gcagccgctt gcgatccagg tcgtgcacga cggatgatga gtgctgcccg cgccgcaggc	420
tggctctgct gatgccacc gtgcgcacgc cggcgaagtc ttccagcgca cgcgctgcy	480
cgacgtagaa ctcgatgcyg cgccacagcc gcttgctcct gcagcgcagc aactcggcgg	540
cctggcgcac cggcaggtcc aggcacaagg tcagcgccag cgcttcgaac gccgcggtga	600
agcccagacc cggacgcgcc cagggcacgg ccacctgcgt ggtcttgccg caggcaccgc	660
aggccacgcy cggcacgctc cagtgcagcc aggcctcgaa ctggaagaag tccagggtgc	720
gccaggatcg gcgcagccgg tcgtgcaccg gttgcgtggc cgaccgcat gccgggcagg	780
cgagcctgct ggtgtggcag ccgatctcga agtcgatacg ccgcttgcc gtgtcgagcc	840
tgacgtcatc gacgaccac ggcggtgca agcccagcgc gctggtgaac agagcttcta	900
cggcaatgcc catggctcat cctcctgatc aaacacgcct cgtggacatg tgcacaggcc	960
ggccagcggc ctgcacacat gcccaccggg ctcgacgacc aggagtcatt gtgactgttg	1020
ggttcacac gaaatgacga aggacccta gtgtgcgggg ttatcgtagc ggtgagaacc	1080
cgacacgatg gcaagggtttt gaccaacatg ctaaaaggcc cattcaaacc aactacccc	1140
gagccagcca tgccacgatc cctactcacc gacgctgacg tgcgttcggt cagcgattcc	1200
gtcgccctcg gcctgccgtc cgaacgcgat gcgtcccttg ccgaggcggt caacctgat	1260
gtgctgccga ccctccagca actcgatgcy gtcaataccg gtgaaatcca gccggcacct	1320
gccttcgacc cgcgctggaa ggagggtgcag tcatgagttc gctaaccgcg ctcaccctcg	1380

cgcaagttgc gcagaaactt aaggcacggg aagtctccgc cgttgaagtt ctggacgcct 1440
 gtctgacgca ggtgcgctcc accgaaaaac agatcagtgc gtacgtgtgc gtgctggagg 1500
 atcaggccccg tgcagcagcc cagcaagctg acgccgacat cagcgccggg cgctggaaaag 1560
 gcccgcgtgca tggcgtgcct gtagcgggtca aggacttata cgacatcgct ggcgtaccca 1620
 ccacggcatc gtcgcgccag cgcacgaatt ggacgccgca gcaagactgc gccgtagtcc 1680
 ggcgcttgaa agacgcaggt gccgttatcc ttggcaagac ccatacgac gaattcgct 1740
 atggcgatcat cactccgaag tcgcgcaacc cctgggaccc gggaagaaca ccgggtggct 1800
 ccagcggcgg ctcggcgcc acggtcgcag cctgcggcgt ctacttggcg accggcaccg 1860
 acaccgggtg atccgttcgc atcccttcgt cgatgtgcaa caccgtaggc ctgaagccaa 1920
 cctacgggcg cgtgagccgt gccggtgtga gttcactttc ctggagcctg gaccatccag 1980
 gcccgatcac gcgcaccgtg gaagacacgg cgctcagcct tcagggtgatg gctggcttcg 2040
 atccagccga ccgcggctcg ttggatgagc cgggtgccag ctatgccgaa gggctcggcc 2100
 aaggcgtgaa aggcctgcgc gtgggcgtgc cgaagaacta cttcttcgac cgcgtggacc 2160
 cggaagtga aagtgcggtt cgtgccgcga tcgatcaact gaaagagctg ggcgccgaac 2220
 tggtggaagt cgaagtgcc atggccgagc agatcatccc ggtggagtcc gggatcgtgc 2280
 taccgaagc cagcgcctac caccgcacga tgctgcgcga gtcaccgag ctctacaccg 2340
 ccgatgtccg catactgctg gaactcggaa atctagtcac cgccaccgac tacctgcagg 2400
 cgcagcgcgt ccgtacgtg atgcagcgcg cgggtggccga gatgttccag cgcacgtatg 2460
 tgctgatcgc acccactg cccatcccgg ctgctcgcag cggggaggag gtccacacat 2520
 ggccggacgg cacggtagag gcgttgttca tggcctatac gcgcttcacc tcgttcggca 2580
 acgtgacagg attacccacg ctgaacctgc cctgtggttt ctccaaggat gggttgccga 2640
 tcggcatgca gatcaccggc cggccgtcgg acgagaagac cctgctgcgt gctgggctgg 2700
 cctacgagaa agccacgacc tggcaccagc gtcacccgga actgatcga gcgggctgag 2760
 ctgcaagaaa gcaggcgcgg cggcgcaaga cccattgtgc taccgctgcg tctgcctgaa 2820
 gtggccgaga ccaagcagaa gaaggagggtg acatagcctc ctacaatctg ccgtcttctc 2880
 ggagccctct ggatgctcga agttcttctgc atgggggttc gccgggagcg caatctgcaa 2940
 ggtggcattg gccttcagtg tcgatgcaga gttgaagtgc ctgtaccct ttttcaacc 3000
 acaccaggag aaccgcacca tggggcaatc acacacgcat gaccaccatc acgacgggta 3060
 ccaggcaccg cccgaagaca tcgcgctgcg ggtcaaggcc ttggagtctc tgctgatcga 3120
 gaaaggtctt gtcgacccag cggccatgga cttggtcgtc caaacgtatg aacacaaggt 3180
 aggccccga aacggcgcca aagtcgtggc caaggcctgg gtggaccctg cctacaaggc 3240
 ccgtctgctg gcagacggca ctgccggcat tgccgagctg ggcttctccg gggtagagg 3300
 cgaggacatg gtcattctgg aaaacacccc cgccgtccac aacgtcgtcg tttgcacctt 3360
 gtgctcttgc taccatggc cgacgctggg cttgccccct gcctggtaca aggccccgcc 3420

ctaccggtcc cgcattggtga gcgacccgcg tggggttctc gcggagttcg gcctggtgat 3480
 ccccgcaag gaaatccgcg tctgggaacac cacggccgaa ttgcgtaca tgggtctgcc 3540
 ggaacggccc gcgggaactg aagcctaacag cgaagaacaa ctggccgaac tcgttaccgc 3600
 cgattcgatg atcggcaccg gcctgcccat ccaaccacc ccatctcatt aaggagttcg 3660
 tcatgaatgg cattcacgat actgggggag cacatggtta tgggccggtt tacagagaac 3720
 cgaacgaacc cgtctttcgc tacgactggg aaaaaacggt catgtccctg tttccggcgc 3780
 tgttcgccaa cggcaacttc aacctcgatg agtttcgaca cggcatcgag cgcattgaacc 3840
 ccatcgacta cctgaagga acctactacg aacctggat ccattccatc gaaaccttgc 3900
 tgggtcgaaa ggggtgtgctc acggcaacgg aactcgcgac cggcaaggca tctggcaaga 3960
 cagcgacacc ggtgctgacg ccggccatcg tggacggact gctcagcacc ggggcttctg 4020
 ccgcccggga ggagggtgcg cgggcgcggt tcgctgtggg ggacaagggt cgcgctctca 4080
 acaagaaccc ggtgggcat acccgcatgc cgcgctacac gcggggcaaa gtggggacag 4140
 tgggtcatcga ccatggtgtg ttcgtgacgc cggacaccgc ggcacacgga aaggcgagc 4200
 acccccagca cgtttacacc gtgagtttca cgtcggtcga actgtggggg caagacgcct 4260
 cctcgccgaa ggacacgatt cgcgtcgact tgtgggatga ctacctggag ccagcgtgat 4320
 catgaaagac gaacggtttc cattgccaga gggttcgtg aaggacctcg atggccctgt 4380
 gtttgacgag ccttggcagt cccaggcggt tgccttggtg gtcagcatgc acaaggccgg 4440
 tctctttcag tggaaagact gggccgagac cttaccgcc gaaatcgacg cttccccggc 4500
 tctgcccggc gaaagcgtca acgacaacta ctaccggcaa tgggtgtcgg cgctggaaaa 4560
 gttggtggcg tcgctggggc ttgtgacggg tggagacgtc aactcgcgcg cacaggagtg 4620
 gaaacaggcc cacctcaaca cccacatgg gcacccgatc ctgctggccc atgcgctttg 4680
 cccgccagcg atcgaccca agcacaagca cgagccaaaa cgctcaccga tcaaggctgt 4740
 tgccgcaatg gcttgagatc cactgtcctg tttccctacc ttgaatggag taaacctga 4800
 gttcattttc taccactgcc gtccccgccg cccagcgctt gagcgccacg cgcagcttac 4860
 tgttgcaact gtccgctggc gccgcgctgg gcctggctgt cctgtacggc gtggcttttg 4920
 ccgaaagccc gcttgccgac aacgccgcgc acgatgttcg ccacgtgacg gtcaagcctt 4980
 gtcactaact gtggccacc cccgccgcgc tgcgcgcgac cccctccagg gggcaaaccg 5040
 agtggccccg cgagaccggt tccaccgcgt tcccgggtgg gtgcactgct ggcgttgcgg 5100
 gtcttggttc ggctggatgg tttcttgacg gagccacacc gtcgctcatc tgtatatgga 5160
 gagggcatg atttttcgac gcttgatctg ggcggcgctg gccgtggcct tgctgggtgg 5220
 cagtttgag tcgggtctgc agcagttgca gaccgtgccc atcatcctgg cggccgaggt 5280
 gtttgagggc cagaagggtg ccgcgcccga gccggtggca acaccggccg gtgcggctgc 5340
 gcacgtccat gcggacggtg cgacaacga ccatggcgac gccgccgaag cctgggcgcc 5400
 cgccgacggg gtggagcgcc acttctggac ctgggtggcc aatgtgctgc acgcgttcag 5460

catggcgctg	ctggtgctgg	ccgtgatggc	ggtgtgccag	tggcgcgga	gcgccttgcg	5520
cgctgtgcct	ttggccgcct	gggtggccgc	cgccggttgg	ctgagctttc	acttctggcc	5580
ttcgctgggc	ctgcacgccg	agatccccgg	catggatgcc	gcccggctgg	gttcacgcca	5640
gggctggtgg	gtgctcgcg	cgggcggtgc	ggcgctggcc	tgtgcgtcgg	tggcgctgat	5700
gcgcagcccc	ctgcgctggg	ccgctgcaat	ggcttgtctg	gcgctgccct	tcgtggtggg	5760
cgcgccgcac	atcgtggccg	acccgctggc	cggtttcacg	ggtgaggctc	aggcggcgct	5820
gcgtgaactg	ggccgccagt	tcactctggg	caccacctgg	ttgtcgctgt	cgttctgggt	5880
gagtatgggc	gtggtggccg	gtctggcctt	ccagcgctgg	ctgagccccg	ccgtggcggc	5940
gctgctgcag	cgacaggaca	ccgcccgggc	cccagccatg	gagacgccgc	gataagcgac	6000
gtcaacgcaa	gggtcgtgcg	gggtgctgat	ctgaaccgcc	tttttttttg	aggaggctct	6060
tttccttgag	aggatagagc	catgagcaag	aagtcgaacc	agttttcacc	cgaagtgcgc	6120
gagcgtgctg	ttcgcattgt	gctggagcac	cgaggcgagt	acccatctct	gtgggccgcc	6180
atcgattcga	tcgcgcccaa	gatcggttgt	gtgccgcaga	ccctgcatac	ctgggtcaag	6240
cgtgtcgagg	tcgacagcgg	cgtgcgtgag	ggcgtcagca	cttccgagct	gcagcgtctg	6300
aaggagctcg	aacgcgagaa	caaggagctg	cgcaaggcca	acgagatctt	gaagctggcc	6360
agcgcgtttt	tcgcccaggc	ggagctcgac	cgccgtctga	agccctgaag	ggcttcatcg	6420
accagtatcg	acaggcctac	ggggctgagt	cgatctgcaa	ggtgttgca	gtcgccccgt	6480
caggctattg	gcgccacgcc	gctcaacaac	gcaaccccc	actgcgctgt	ccgtgcgctc	6540
aacgtgacga	caccctggtg	gcgcacattg	aacgcgtctg	gcaggccaac	atgcgggtct	6600
atggcgccga	caaggctctg	aagcaaatga	accgcgaggg	cattgtgggtg	gcgcgctgca	6660
cggtcgagcg	gctgatgcgg	cgcttgcgct	tgcagggcgt	gcgccgtggc	aagggtggtg	6720
gtaccaccat	cagcgtatgg	cgggcgccgt	gcccgtgga	ccgggtcaac	cggtgtgtca	6780
gagcagaccg	gcccaccag	ctctgggtct	cggacttcac	ctacgtgtcg	acctggcagg	6840
gctggctgta	cgtggccttc	gtgatcgatg	tgtacgccc	gcgcatcgtg	ggctggcggg	6900
taagcagttc	gatgcgcacg	gacttcgtgc	tcgatgccct	ggaacaggcg	ctgtatgcgc	6960
gccagcctga	gcgcgatggc	agcctggttt	gccactccga	caggggggtc	caatacgtca	7020
gcatccgcta	caccgagcga	ctggcccagg	ccggtatcga	gccctcggtg	ggcagcaaag	7080
gtgacagcta	cgacaacgcg	ctggccgaga	cgatcaacgg	gctgtacaag	gccgaattga	7140
tccatcggcg	agccccatgg	aggaccaagg	agtcggtgga	gctggctacc	ctcccatggg	7200
tgtcatgggt	caaccaccac	cggctgctcg	aaccatcgg	ctacatccc	ccggccgagg	7260
ctgaggcaaa	ctactaccgg	caactcgcca	atcaggcctc	catggtgggtg	gcctgactta	7320
aaccaaccgg	cctccacgat	tccggggcg	gttcaagagc	ctcctatttc	ctgtccaata	7380
ggagccccta	tgagcagtca	acgttaccgc	aattc			7415

<210> 19
 <211> 31
 <212> DNA
 <213> artificial sequence

<220>
 <223> Primer 4

<400> 19
 ttatctagac ccgcgctgga aggaggtgca g 31

<210> 20
 <211> 29
 <212> DNA
 <213> artificial sequence

<220>
 <223> Primer 5

<400> 20
 cagaagcttt cttgcagctc agcccgcctc 29

<210> 21
 <211> 33
 <212> DNA
 <213> artificial sequence

<220>
 <223> Primer 6

<400> 21
 tatgaattca tgagttcgct aaccgcctc acc 33

<210> 22
 <211> 40
 <212> DNA
 <213> artificial sequence

<220>
 <223> Primer 7

<400> 22
 tagcggccgc ttagcccgct ccgatcagtt ccgatgacg 40

<210> 23
 <211> 3449
 <212> DNA
 <213> Comamonas testosteroni 5-MGAM-4D

<400> 23
 tatgaattca tgagttcgct aaccgcctc accctcgcgc aagttgcgca gaaacttaag 60
 gcacgggaag tctccgccgt tgaagttctg gacgcctgtc tgacgcaggt gcgctccacc 120
 gaaaaacaga tcagtgcgta cgtgtgcgtg ctggaggatc aggcccgctc agcagccag 180
 caagctgacg ccgacatcag cgccggggcgc tggaaaggcc cgctgcatgg cgtgccctgta 240
 gcggtcaagg acttatacga catcgctggc gtaccaccca cggcatcgtc gcgccagcgc 300
 acgaattgga cgccgcagca agactgcgcc gtagtcgggc gcttgaaaga cgcaggctgcc 360
 gttatccttg gcaagacca tacgcacgaa ttcgcctatg gcgtcatcac tccgaagctcg 420

cgcaaccctt	gggacccggg	aagaacaccg	ggtggctcca	gcggcggctc	ggcggccacg	480
gtcgcagcct	gcggcgctta	cttggcgacc	ggcaccgaca	ccggtggatc	cgttcgcata	540
ccttcgtcga	tgtgcaacac	cgtaggcctg	aagccaacct	acgggcgcgt	gagccgtgcc	600
ggtgtgagtt	cacttttctg	gagcctggac	catccaggcc	cgatcacgcg	caccgtggaa	660
gacacggcgc	tcagccttca	ggtgatggct	ggcttcgata	cagccgaccg	cggctcgttg	720
gatgagccgg	tgcccagcta	tgccgaaggg	ctcggccaag	gcgtgaaagg	cctgcgcgtg	780
ggcgtgccga	agaactactt	cttcgaccgc	gtggacccgg	aagttgaaag	tgcggttcgt	840
gccgccatcg	atcaactgaa	agagctgggc	gccgaactgg	tggaagtcga	agtgcccatg	900
gccgagcaga	tcatcccggg	ggagttcggg	atcgtgctac	ccgaagccag	cgcctaccac	960
cgcacgatgc	tgcgcgagtc	acccgagctc	tacaccgccg	atgtccgcat	actgctggaa	1020
ctcggaaatc	tagtcaccgc	caccgactac	ctcgaggcgc	agcgcgtccg	tacgtctgatg	1080
cagcgcgcgg	tggccgagat	gttccagcgc	atcgatgtgc	tgatcgcacc	cacactgccc	1140
atcccggctg	ctcgcagcgg	ggaggaggtc	cacacatggc	cggacggcac	ggtagaggcg	1200
ttgttcatgg	cctatacgcg	cttcacctcg	ttcggcaacg	tgacaggatt	acccacgctg	1260
aacctgccct	gtggtttctc	caaggatggg	ttgccgatcg	gcatgcagat	caccggcccg	1320
ccgctggacg	agaagaccct	gctgcgtgct	gggctggcct	acgagaaagc	cacgacctgg	1380
caccagcgtc	atccggaact	gatcggagcg	ggctgagctg	caagaaagca	ggcgcggcgg	1440
cgcacgaccc	attgtgctac	cgctgcgtct	gcctgaagtg	gccgagacca	agcagaagaa	1500
ggagggtgaca	tagcctccta	caatctgccg	tcttctcgga	gccctctgga	tgctcgaagt	1560
tctttgcatg	gggttgcgcc	gggagcgcaa	tctgcaaggc	ggcattggcc	ttcagtgtcg	1620
atgccgagtt	gaagtcgctg	tacccctttt	ttcaaccaca	ccaggagaac	cgcaccatgg	1680
ggcaatcaca	cacgcatgac	caccatcacg	acgggtacca	ggcaccgcc	gaagacatcg	1740
cgctgcgggt	caaggccttg	gagtctctgc	tgatcgagaa	aggtcttgtc	gacccagcgg	1800
ccatggactt	ggtcgtccaa	acgtatgaac	acaaggtagg	ccccgaaa	ggcgccaaag	1860
tcgtggccaa	ggcctgggtg	gaccctgcct	acaaggcccg	tctgctggca	gacggcactg	1920
ccggcattgc	cgagctgggc	ttctccgggg	tacaggcgga	ggacatggtc	attctggaaa	1980
acacccccgc	cgccacaac	gtcgtcgttt	gcaccttggt	ctcttgcta	ccatggccga	2040
cgctgggctt	gccccctgcc	tggtacaagg	ccccgcccta	ccggtcccg	atgggtgagcg	2100
acccgcgtgg	ggttctcgcg	gagttcggcc	tggtgatccc	cgcgaaggaa	atccgcgtct	2160
gggacaccac	ggccgaattg	cgctacatgg	tgctgccgga	acggcccgcg	ggaactgaag	2220
cctacagcga	agaacaactg	gccgaactcg	ttaccgcgga	ttcgatgat	ggcaccggcc	2280
tgcccatcca	accacccca	tctcattaag	gagttcgtca	tgaatggcat	tcacgatact	2340
gggggagcac	atggttatgg	gccggtttac	agagaaccga	acgaaccctg	ctttcgctac	2400
gactgggaaa	aaacgggtcat	gtccctgttc	ccggcgctgt	tcgccaacgg	caacttcaac	2460

ctcgatgagt ttcgacacgg catcgagcgc atgaacccca tcgactacct gaaggggaacc	2520
tactacgaac actggatcca ttccatcgaa accttgctgg tcgaaaaggg tgtgctcacg	2580
gcaacggaac tcgcgaccgg caaggcatct ggcaagacag cgacaccggt gctgacgccg	2640
gccatcgtgg acggactgct cagcaccggg gcttctgccg cccgggagga ggggtgcgcg	2700
gcgcggttcg ctgtggggga caaggttcgc gtcctcaaca agaaccggt gggccatacc	2760
cgcgacccgc gctacacgcg gggcaaagtg gggacagtgg tcatcgacca tgggtgtgttc	2820
gtgacgccgg acaccgcggc acacggaaag ggcgagcacc cccagcacgt ttacaccgtg	2880
agtttcacgt cggtcgaact gtgggggcaa gacgcctcct cgccgaagga cacgattcgc	2940
gtcgacttgt gggatgacta cctggagcca gcgtgatcat gaaagacgaa cggtttccat	3000
tgccagaggg ttcgctgaag gacctcgatg gccctgtgtt tgacgagcct tggcagtcct	3060
aggcgtttgc cttggtgggtc agcatgcaca aggccggtct ctttcagtgg aaagactggg	3120
ccgagacctt caccgccgaa atcgacgctt ccccggtctt gcccggcgaa agcgtcaacg	3180
acacctacta ccggcaatgg gtgtcggcgc tggaaaagtt ggtggcgtcg ctggggcttg	3240
tgacgggttg agacgtcaac tcgcgcgcac aggagtggaa acaggccac ctcaacaccc	3300
cacatgggca cccgatcctg ctggcccatg cgctttgcc gccagcgatc gaccccaagc	3360
acaagcacga gccaaaacgc tcaccgatca aggtcgttgc cgcaatggct tgagatccac	3420
tgtcctgttt ccctaccttg aagcttcag	3449

INTERNATIONAL SEARCH REPORT

International application No. 7

PCT/US04/36352²

A. CLASSIFICATION OF SUBJECT MATTER IPC(7) : C12N 9/78, 1/21, 1/20, C12P 13/00, 7/40 US CL : 435/227, 252.3, 128, 136, 252.1 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/227, 252.3, 128, 136, 252.1 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 6,670, 158 B2 (Dicosimo R. et al.) 30 Dec. 2003 all and WO 95/04828 (Cerbelaud et. al) 16, Feb. 1995	1-14, 19-56
☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.		
* Special categories of cited documents:		
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family	
Date of the actual completion of the international search 12 June 2005 (12.06.2005)		Date of mailing of the international search report 12 JUL 2005
Name and mailing address of the ISA/US Mail Stop PCT, Attn: ISA/US Commissioner for Patents P.O. Box 1450 Alexandria, Virginia 22313-1450 Facsimile No. (703) 305-3230		Authorized officer Achutamurthy ponnathapura Telephone No. 571-272-1600

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US04/3633b

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:
Please See Continuation Sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US04/36352

Continuation of Item 4 of the first sheet:

The title is too long.

Nitrile hydratase and Amidase from Comamonas Testosteroni 5-MGAM-4D.

BOX III. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING

The inventions listed as Groups I-VI do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

Group I claims 1-3, 5-7, 9 and 18-27 are drawn to DNA, vectors, host cells and expression method for the alpha- and beta subunits of nitrile hydratase from Comamonase testosterone 5-MGAM-4D.

Group II, claim(s) 4, 8, drawn to alpha- and beta subunits of nitrile hydratase from Comamonase testosterone 5-MGAM-4D.

Group III, claim(s) 10 in part, 15, 16, 17 are drawn to DNA encoding the accessory gene and the accessory protein.

Group IV claims 11, 12, 13 and 19-28 are drawn to DNA for the the amidase gene.

Group V claim 14 is drawn to the amidase protein.

Group VI claims 29-35, (40-42 in part), 43-49, (54-56 in part) are drawn to a method of converting a substrate containing one or more functional groups to an amide using a transformed microbial cell expressing a nitrile hydratase polypeptide from Comamonase testosterone 5-MGAM-4D..

Group VII claims 36-38, (40-42 in part), 50-53, (54-56 in part) are drawn to a method of converting a substrate containing one or more functional groups using a catalyst having nitrile hydratase activity from Comamonase testosterone 5-MGAM-4D.

The special technical feature linking Groups I-VI is that it teaches a two step bio-catalytic method whereby in a first step one or more nitrile functional groups are converted to the corresponding amides and in a second step the amides are in turn converted to the corresponding carboxylic acid. The DNA of group I from Comamonase testosterone 5-MGAM-4D encodes the Nitrile hydratase of group II and that the nitrile hydratase catalyses the conversion of a substrate containing one or more nitrile functional groups to an amide. The amidase gene of group IV from Comamonase testosterone 5-MGAM-4D encodes for the amidase protein of group V that converts the amide product(s) from the previous to the corresponding carboxylic acid(s) in the method of Group VI or VII. It is recognized that the accessory protein found in Comamonase testosterone 5-MGAM-4D is part of the nitrile hydratase reaction. However US patent 6670,158 B2 teaches the bio-catalytic conversion of compounds with one or more nitrile functional groups to the corresponding carboxylic acid(s) by Comamonase testosterone 5-MGAM-4D. It teaches a method of catalysis having nitrile hydratase and amidase activities of Comamonase testosterone 5-MGAM-4D in the form of intact microbial cells (Group VI), permeabilized microbial cells, one or more cell components of microbial cell extract (Group VII), and a partial purified enzyme(s) (Group VII), , or purified enzyme(s). In addition it teaches that the catalysis does not require monitoring the nitrile concentration over the course of the reaction to maintain a low concentration of nitrile, and does not require running the reaction at a low temperature in order to maintain the stability of the nitrile hydratase activity. It also teaches that the enzyme catalysts can be immobilized

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US04/36352

in a polymer matrix such as alginate, carrageenan, polyvinylalcohol or polyacrylamid gelor on a soluble or insoluble support to facilitate recovery and reuse of the catalyst. As such the only shared technical feature of groups I-VII is not special as defined by PCT rule 13.2 as it does not define a contribution over the ar