



(51) International Patent Classification:

C07D 471/04 (2006.01) **A61P 19/02** (2006.01)
A61K 31/4462 (2006.01)

(21) International Application Number:

PCT/EP2011/056600

(22) International Filing Date:

27 April 2011 (27.04.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

1007203.1 29 April 2010 (29.04.2010) GB

(71) **Applicant** (for all designated States except US): **GLAXO GROUP LIMITED** [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford Middlesex UB6 0NN (GB).

(72) **Inventors; and**

(75) **Inventors/Applicants** (for US only): **ATKINSON, Francis Louis** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **BARKER, Michael David** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **DOUAULT, Clement** [FR/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **GARTON, Neil Stuart** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **LIDDLE, John** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **PATEL, Vipulkumar Kantibhai** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **PRESTON, Alexander George**

Steven [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB). **WILSON, David Matthew** [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage Hertfordshire SG1 2NY (GB).

(74) **Agents**: **PRICE, Susanna Clare Hopley** et al.; GlaxoSmithKline, Global Patents (CN925.1), 980 Great West Road, Brentford Middlesex TW8 9GS (GB).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

[Continued on next page]

(54) **Title**: 7-(1H-PYRAZOL-4-YL)-1,6-NAPHTHYRIDINE COMPOUNDS AS SYK INHIBITORS



(57) **Abstract**: A compound of formula (I) or a salt thereof; which is an inhibitor of spleen tyrosine kinase (Syk) and therefore potentially of use in treating diseases resulting from inappropriate activation of mast and/or basophil cells, macrophages, and B-cells and related inflammatory responses and tissue damage, for instance inflammatory disease and/or allergic disorders, and in cancer therapy, specifically heme malignancies, chronic spontaneous urticaria and autoimmune conditions.



-
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*
- *of inventorship (Rule 4.17(iv))*
- Published:**
- *with international search report (Art. 21(3))*

7-(1H-PYRAZOL-4-YL)-1,6-NAPHTHYRIDINE COMPOUNDS AS SYK INHIBITORS

5 The present invention relates to novel chemical compounds which have activity against the spleen tyrosine kinase (Syk), processes for their preparation, pharmaceutically acceptable formulations containing them and their use in therapy.

10 Syk is a non-receptor tyrosine kinase that is involved in coupling activated immunoreceptors to signal downstream events that mediate diverse cellular responses, including proliferation, differentiation, and phagocytosis. Syk is widely expressed in hematopoietic cells. Syk inhibitors have potential anti-inflammatory and immunomodulating activities. They inhibit Syk-mediated IgG Fc epsilon and gamma receptor and BCR receptor signaling, resulting in inhibition of the activation of mast cells, macrophages, and B-cells and related inflammatory responses and tissue damage. Accordingly, Syk inhibitors have attracted interest in a number of therapeutic areas, including the treatment of rheumatoid arthritis, B-cell lymphoma and asthma/rhinitis.

20 Rheumatoid arthritis (RA) is an auto-immune disease affecting approximately 1% of the population. It is characterised by inflammation of articular joints leading to debilitating destruction of bone and cartilage. Recent clinical studies with rituximab, which causes a reversible B cell depletion, (J.C.W. Edwards *et al.* New Eng. J. Med., 2004, 350, 25: 2572-2581), have shown that targeting B cell function is an appropriate therapeutic strategy in auto-immune diseases such as RA. Clinical benefit correlates with a reduction in auto-reactive antibodies (or rheumatoid factor) and these studies suggest that B cell function and indeed auto-antibody production are central to the ongoing pathology in the disease.

30 Studies using cells from mice deficient in the Syk have demonstrated a non-redundant role of this kinase in B cell function. The deficiency in Syk is characterised by a block in B cell development (M. Turner *et al.* Nature, 1995, 378: 298-302 and Cheng *et al.* Nature, 1995, 378: 303-306). These studies, along with studies on mature B cells deficient in Syk (Kurosaki *et al.* Immunol. Rev. 2000, 176:19-29), demonstrate that Syk is required for the differentiation and activation of B cells. Hence, inhibition of Syk in RA patients is likely to block B cell function and hence to reduce rheumatoid factor production. In addition to the role of Syk in B cell function, of relevance to the treatment of RA, is the requirement for Syk activity in Fc receptor (FcR) signalling. FcR activation by immune complexes in RA has been suggested to contribute to the release of multiple pro-inflammatory mediators.

40 The contribution of Syk dependent processes to the pathology of RA has been reviewed by Wong *et al.* (Expert Opinion Investigational Drugs, 2004, 13 (7): 743-762).

The results of a 12 week clinical trial for the Syk inhibitor R788 (fostamatinib disodium, Rigel) have been published: Treatment of rheumatoid arthritis with a syk kinase inhibitor: A twelve-week, randomized, placebo-controlled trial, *Arthritis & Rheumatism*, 58(11), 2008: 3309-3318.

5

Syk inhibitors may also be useful in cancer therapy, specifically heme malignancies, particularly Non-Hodgkin's Lymphomas including follicular (FL), mantle cell, Burkitt and diffuse large B cell (DLBCL) lymphomas.

10 Studies have shown that Syk is dysregulated by overexpression and/or constitutively activation in a variety of primary B-lymphoma tumors and also in B-lymphoma cell lines. Syk, through the PI3K/AKT pathway, the PLD pathway and AKT independent signalling, activates mTOR (mammalian target of rapamycin) which in turn increases B-cell survival and proliferation. Inhibition of Syk, *in vitro*, results in decreased mTOR
15 activation and a reduction of clonicity in FL cells. Inhibition of Syk kinase with curcumin in a murine model of B lymphoma (BKS-2) gave a significant reduction of tumour burden as measured by the total splenocyte number (Leseux L. *et al.* *Blood* 15 Dec 2006, 108(13): 4156-4162 and Gururajan M. *et al.* *Journal of Immunology*, 2007, 178: 111-121).

20

Results of a Phase 2 clinical trial of R788 (fostamatinib disodium) in patients with relapsed or refractory B-Cell non-Hodgkin's lymphoma (NHL) show that the compound is well-tolerated by these patients, as well as a therapeutic benefit in patients suffering from diffuse large B-Cell lymphoma (DLBCL) and chronic
25 lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL). Despite the fact that the patients enrolled in this trial had advanced disease and had failed treatment with marketed therapies, a significant number of them were particularly responsive to Syk inhibition with R788 (www.Rigel.com; Friedberg J. W. *et al.* *Blood*, 1 April 2010; 115(13): 2578-85).

30

Syk inhibitors may also be useful in the treatment of asthma and rhinitis as they are important in transducing the downstream cellular signals associated with cross-linking FcεR1 and/or FcγR1 receptors, and they are positioned early in the signalling cascade. In mast cells, for example, the early sequence of FcεR1 signalling following
35 allergen cross-linking of receptor-IgE complexes involves first Lyn (a Src family tyrosine kinase) and then Syk.

Allergic rhinitis and asthma are diseases associated with hypersensitivity reactions and inflammatory events involving a multitude of cell types including mast cells,
40 eosinophils, T cells and dendritic cells. Following exposure to allergen, high affinity immunoglobulin receptors for IgE (FcεRI) and IgG (FcγRI) become cross-linked and activate downstream processes in mast cells and other cell types leading to the release of pro-inflammatory mediators and airway spasmogens. In the mast cell, for example, IgE receptor cross-linking by allergen leads to release of mediators

including histamine from pre-formed granules, as well as the synthesis and release of newly synthesised lipid mediators including prostaglandins and leukotrienes.

5 The Syk inhibitor R112 (Rigel), dosed intranasally in a phase I/II study for the treatment of allergic rhinitis, was shown to give a statistically significant decrease in PGD₂, a key immune mediator that is highly correlated with improvements in allergic rhinorrhea, as well as being safe across a range of indicators, thus providing the first evidence for the clinical safety and efficacy of a topical Syk inhibitor (see Meltzer E. O. *et al.* "An intranasal Syk kinase inhibitor (R112) improves the symptoms of
10 seasonal allergic rhinitis in a park environment"; Journal of Allergy and Clinical Immunology, 2005, 115(4): 791-796). In a further phase II clinical trial, for allergic rhinitis, R112 was however shown as having a lack of efficacy versus placebo (www.clinicaltrials.gov Identifier NCT0015089).


15 Acute and chronic urticaria are common skin diseases thought to affect around 25% of the total population within the USA (see "New concepts in chronic urticaria"; Current Opinions in Immunology, 2008, 20: 709-716). Although urticaria can be triggered by allergic reactions many cases have an unclear etiology. Chronic spontaneous urticaria is defined as when wide spread wheals are present for greater
20 than 6 weeks. There are many pathological similarities in chronic urticaria patients, in terms of extent of wheals in the skin, with allergen-induced mast and basophil cell degranulation reactions via IgE activation. Around 40% of chronic spontaneous urticaria patients contain serum IgG auto-antibodies targeting IgE or the IgE receptor (Fc Epsilon Receptor) and these are thought to drive the histamine and other
25 mediator release via mast and basophil degranulation. Syk inhibitors would inhibit the signalling response post IgE mediated Fc Epsilon activation and inhibit the mediator release known to be involved in chronic pruritis in multiple diseases.

WO03/057695A1 (Boehringer Ingelheim Pharmaceuticals, Inc.) describes novel 1,6-naphthyridines that have Syk inhibitory activity. These are further described in
30 "Discovery and SAR of Novel [1,6] Naphthyridines as Potent Inhibitors of Spleen Tyrosine Kinase (SYK)" (Bioorganic & Medicinal Chemistry Letters 13 (2003): 1415 – 1418). This has been followed with two more recent patent applications, WO2010/015518A2 and WO2010/015520A1 (Boehringer Ingelheim International
35 GmbH), describing 4-dimethylamino-phenyl-substituted naphthyridines and substituted naphthyridines, respectively.

WO04/035604A2 (Millennium Pharmaceuticals, Inc.) discloses the structural co-ordinates of the human Syk protein.

40 There remains however the need to identify further compounds which are inhibitors of spleen tyrosine kinase (Syk).

Thus, in a first aspect invention, the present invention provides a compound of formula (I):



5

wherein:

X is O or NH;

R¹ is C₂₋₄ alkyl, C₃₋₇cycloalkyl, hydroxyC₂₋₄ alkyl, C₁₋₂ alkoxyC₁₋₄alkyl, trifluoromethylC₁₋₂ alkyl or benzyl;

10 or a salt thereof (hereinafter "compounds of the invention").

In one aspect, X is NH.

In one embodiment R¹ is ethyl, *iso*-propyl, *t*-butyl, cyclopentyl, methoxymethyl, methoxyethyl, hydroxyethyl, 2,2-hydroxymethylpropyl, 2,2,2-trifluoroethyl or benzyl. In another embodiment, R¹ is C₂₋₄alkyl, C₃₋₇cycloalkyl or C₁₋₂alkoxyC₁₋₄alkyl. In another embodiment, R¹ is ethyl, *t*-butyl, -CH₂OCH₃, -CH₂C(CH₃)₂OH, -CH₂CH₂OH, -CH₂CH₂OCH₃, -CH(CH₃)₂, -CH₂CF₃, benzyl or cyclopentyl. In another embodiment, R¹ is ethyl, *t*-butyl, or methoxymethyl. In another embodiment, R¹ is ethyl or *t*-butyl. In a further embodiment, R¹ is *t*-butyl.


It is to be understood that the present invention covers all combinations of substituent groups described hereinabove.

25 It will be appreciated that the compounds of formula (I) comprise a chiral centre at the 3-carbon of the piperidiny ring. It has been found that those having the *S* absolute stereochemistry are generally preferred, on account of better drug development properties, including better bioavailability (as measured in predictive animal pharmacokinetic studies) and lesser hERG activity.

30

Accordingly, in one embodiment, the present invention provides a compound of formula (I) having the *S* absolute stereochemistry. In another embodiment, the present invention provides a compound of formula (I) having the *S* absolute stereochemistry, that is a compound of formula (II):

35



wherein:

X and R¹ are as hereinbefore defined;

5 or a salt thereof.

Compounds of the invention include the compounds of Examples 1 to 11 and salts thereof.

10 In one embodiment, the present invention provides a compound formula (I) selected from the group consisting of:

7-(1-ethyl-1*H*-pyrazol-4-yl)-*N*-{[(3*S*)-3-fluoro-3-piperidiny]methyl}-1,6-naphthyridin-5-amine;

15 7-[1-(1,1-dimethylethyl)-1*H*-pyrazol-4-yl]-*N*-{[(3*S*)-3-fluoro-3-piperidiny]methyl}-1,6-naphthyridin-5-amine;

N-{[(3*S*)-3-fluoro-3-piperidiny]methyl}-7-{1-[(methyloxy)methyl]-1*H*-pyrazol-4-yl}-1,6-naphthyridin-5-amine;

7-(1-ethyl-1*H*-pyrazol-4-yl)-5-({[(3*S*)-3-fluoro-3-piperidiny]methyl}oxy)-1,6-naphthyridine;

20 1-{4-[5-({[(3*S*)-3-fluoro-3-piperidiny]methyl}amino)-1,6-naphthyridin-7-yl]-1*H*-pyrazol-1-yl}-2-methyl-2-propanol;

2-{4-[5-({[(3*S*)-3-fluoro-3-piperidiny]methyl}amino)-1,6-naphthyridin-7-yl]-1*H*-pyrazol-1-yl}ethanol;

25 *N*-{[(3*S*)-3-fluoro-3-piperidiny]methyl}-7-{1-[2-(methyloxy)ethyl]-1*H*-pyrazol-4-yl}-1,6-naphthyridin-5-amine;

7-(1-cyclopentyl-1*H*-pyrazol-4-yl)-*N*-[(3-fluoro-3-piperidiny)methyl]-1,6-naphthyridin-5-amine;

N-[(3-fluoro-3-piperidiny)methyl]-7-[1-(2,2,2-trifluoroethyl)-1*H*-pyrazol-4-yl]-1,6-naphthyridin-5-amine;

30 *N*-{[(3*S*)-3-fluoro-3-piperidiny]methyl}-7-[1-(2,2,2-trifluoroethyl)-1*H*-pyrazol-4-yl]-1,6-naphthyridin-5-amine;

N-{[(3*R*)-3-fluoro-3-piperidiny]methyl}-7-[1-(2,2,2-trifluoroethyl)-1*H*-pyrazol-4-yl]-1,6-naphthyridin-5-amine;

35 *N*-[(3-fluoro-3-piperidiny)methyl]-7-[1-(phenylmethyl)-1*H*-pyrazol-4-yl]-1,6-naphthyridin-5-amine;

N-{[3-fluoro-3-piperidinyl]methyl}-7-[1-(1-methylethyl)-1*H*-pyrazol-4-yl]-1,6-naphthyridin-5-amine;

N-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-7-[1-(1-methylethyl)-1*H*-pyrazol-4-yl]-1,6-naphthyridin-5-amine; and

- 5 *N*-{[(3*R*)-3-fluoro-3-piperidinyl]methyl}-7-[1-(1-methylethyl)-1*H*-pyrazol-4-yl]-1,6-naphthyridin-5-amine; and
salts thereof.

In another embodiment, the present invention provides a compound formula (I) selected from the group consisting of:

7-(1-ethyl-1*H*-pyrazol-4-yl)-*N*-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine;

7-[1-(1,1-dimethylethyl)-1*H*-pyrazol-4-yl]-*N*-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine;

- 15 *N*-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-7-{1-[(methyloxy)methyl]-1*H*-pyrazol-4-yl}-1,6-naphthyridin-5-amine;

7-(1-ethyl-1*H*-pyrazol-4-yl)-5-({[(3*S*)-3-fluoro-3-piperidinyl]methyl}oxy)-1,6-naphthyridine;

- 1-{4-[5-({[(3*S*)-3-fluoro-3-piperidinyl]methyl}amino)-1,6-naphthyridin-7-yl]-1*H*-pyrazol-20 1-yl}-2-methyl-2-propanol;

2-{4-[5-({[(3*S*)-3-fluoro-3-piperidinyl]methyl}amino)-1,6-naphthyridin-7-yl]-1*H*-pyrazol-1-yl}ethanol;

N-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-7-{1-[2-(methyloxy)ethyl]-1*H*-pyrazol-4-yl}-1,6-naphthyridin-5-amine;

- 25 7-(1-cyclopentyl-1*H*-pyrazol-4-yl)-*N*-[(3-fluoro-3-piperidinyl)methyl]-1,6-naphthyridin-5-amine;

N-[(3-fluoro-3-piperidinyl)methyl]-7-[1-(2,2,2-trifluoroethyl)-1*H*-pyrazol-4-yl]-1,6-naphthyridin-5-amine; and

- 30 *N*-[(3-fluoro-3-piperidinyl)methyl]-7-[1-(phenylmethyl)-1*H*-pyrazol-4-yl]-1,6-naphthyridin-5-amine; and
salts thereof.


In another embodiment, the present invention provides a compound formula (I) selected from the group consisting of:

- 35 7-(1-ethyl-1*H*-pyrazol-4-yl)-*N*-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine; and

7-[1-(1,1-dimethylethyl)-1*H*-pyrazol-4-yl]-*N*-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine; and
salts thereof.


40

In another embodiment, the present invention provides a compound formula (I) which is 7-[1-(1,1-dimethylethyl)-1*H*-pyrazol-4-yl]-*N*-{[3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine



or a salt thereof.

- In a further embodiment, the present invention provides a compound formula (I)
 5 which is 7-[1-(1,1-dimethylethyl)-1H-pyrazol-4-yl]-N-[[[(3S)-3-fluoro-3-piperidyl]methyl]-1,6-naphthyridin-5-amine



or a salt thereof.

- 10 Included within the scope of the "compounds of the invention" are all solvates (including hydrates), complexes, polymorphs, prodrugs, radiolabelled derivatives, stereoisomers and optical isomers of the compounds of formula (I) and salts thereof.

- It is to be understood that the references herein to compounds of formula (I) and
 15 salts thereof covers the compounds of formula (I) as free bases, or as salts thereof, for example as pharmaceutically acceptable salts thereof. Thus, in one embodiment, the invention is directed to compounds of formula (I) as the free base. In another embodiment, the invention is directed to compounds of formula (I) and salts thereof. In a further embodiment, the invention is directed to compounds of formula (I) and
 20 pharmaceutically acceptable salts thereof.

- Compounds of the present invention are useful as inhibitors of Syk. Compounds of the present invention exhibit low activity in the hERG binding assay, a key measure of potential cardiac toxicity. Compounds of the present invention are also negative in
 25 the Bacterial Reverse Mutation Test, an assay designed to detect a wide range of chemical substances that can produce genetic damage that leads to gene mutations (base pair substitutions and frameshift mutations).

- Compounds of the present invention are thus potentially of use in treating some
 30 cancer therapies, in particular heme malignancies, as well as inflammatory conditions

which involve B cells and/or activated macrophages, and also diseases resulting from inappropriate mast cell activation, for instance allergic and inflammatory diseases.

5 When used herein, the term "alkyl" includes all saturated straight chain and branched isomers. For example, C₁₋₄alkyl means a straight or branched chain alkyl group containing at least 1, and at most 4, carbon atoms. Representative examples thereof include, but are not limited to, methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl and *t*-butyl.

10

When used herein, the term "alkoxy" includes all saturated straight chain and branched isomers. For example, C₁₋₂alkoxy means a straight chain alkoxy group containing at least 1, and at most 2, carbon atoms. Representative examples of "alkoxy" as used herein include, but are not limited to, methoxy and ethoxy.

15

When used herein the term "cycloalkyl" includes, unless otherwise defined, carbocyclic rings having from three to seven ring carbon atoms. Representative examples thereof include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. Preferably the cycloalkyl ring comprises five or six ring carbon atoms.

20

When used herein, the term "pharmaceutically acceptable" refers to those compounds, materials, compositions, and dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, or other problem or complication, commensurate with a reasonable benefit/risk ratio. The skilled artisan will appreciate that pharmaceutically acceptable salts of the compound of the present invention may be prepared.

25

30 When used herein, the term "pharmaceutically acceptable salts" refers to salts that retain the desired biological activity of the subject compound and exhibit minimal undesired toxicological effects. These pharmaceutically acceptable salts may be prepared *in situ* during the final isolation and purification of the compound, or by separately reacting the purified compound in its free acid or free base form with a suitable base or acid, respectively. Indeed, in certain embodiments of the invention, pharmaceutically acceptable salts may be preferred over the respective free base or free acid because such salts impart greater stability or solubility to the molecule thereby facilitating formulation into a dosage form.

35

40 The compounds of formula (I) are basic and accordingly generally capable of forming pharmaceutically acceptable acid addition salts by treatment with a suitable acid. Suitable acids include pharmaceutically acceptable inorganic acids and pharmaceutically acceptable organic acids. Representative pharmaceutically

acceptable acid addition salts include hydrochloride, hydrobromide, nitrate, methylnitrate, sulfate, bisulfate, sulfamate, phosphate, acetate, hydroxyacetate, phenylacetate, propionate, butyrate, isobutyrate, valerate, maleate, hydroxymaleate, acrylate, fumarate, malate, tartrate, citrate, salicylate, *p*-aminosalicylate, glycollate, lactate, heptanoate, phthalate, oxalate, succinate, benzoate, *o*-acetoxybenzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, mandelate, tannate, formate, stearate, ascorbate, palmitate, oleate, pyruvate, pantoate, malonate, laurate, glutarate, glutamate, estolate, methanesulfonate (mesylate), ethanesulfonate (esylate), 2-hydroxyethanesulfonate, benzenesulfonate (besylate), *p*-aminobenzenesulfonate, *p*-toluenesulfonate (tosylate), and naphthalene-2-sulfonate. In one embodiment, the present invention provides a pharmaceutically acceptable salt of a compound of formula (I) which is the hydrochloride salt.

The compounds of formula (I) contain a chiral centre and, therefore, exist as individual enantiomers, or as mixtures thereof. Where the stereochemistry of the chiral centre is not specified the structure is intended to encompass each enantiomer and all mixtures thereof. Thus, the compounds of formula (I) may be used as racemic mixtures including racemic mixtures and racemates, enantiomerically enriched mixtures, or as enantiomerically pure individual stereoisomers. The present invention includes all such mixtures as well as pure individual enantiomers. Generally it is preferred to use a compound of formula (I) in the form of a purified single enantiomer, for instance a compound of formula (II).

The individual enantiomers of a compound of formula (I) may be resolved by methods known to those skilled in the art. For example, such resolution may be carried out (1) by formation of diastereoisomeric salts, complexes or other derivatives; (2) by selective reaction with a stereoisomer-specific reagent, for example by enzymatic oxidation or reduction; or (3) by gas-liquid or liquid chromatography in a chiral environment, for example, on a chiral support such as silica with a bound chiral ligand or in the presence of a chiral solvent. The skilled person will appreciate that where the desired stereoisomer is converted into another chemical entity by one of the separation procedures described above, a further step is required to liberate the desired form. Alternatively, specific enantiomers may be synthesized by asymmetric synthesis using optically active reagents, substrates, catalysts or solvents, or by converting one enantiomer to the other by asymmetric transformation.


A compound of the present invention may exist in solid or liquid form. In the solid state, the compound of the present invention may exist in crystalline or non-crystalline (amorphous) form, or as a mixture thereof. For a compound of the present invention that is in crystalline form, the skilled artisan will appreciate that pharmaceutically acceptable solvates may be formed wherein solvent molecules are

incorporated into the crystalline lattice during crystallization. Solvates may involve non-aqueous solvents such as, but not limited to, ethanol, isopropanol, n-butanol, i-butanol, acetone, tetrahydrofuran, dioxane, DMSO, acetic acid, ethanolamine, and ethyl acetate, or they may involve water as the solvent that is incorporated into the crystalline lattice. Solvates wherein water is the solvent incorporated into the crystalline lattice are typically referred to as "hydrates". Hydrates include stoichiometric hydrates as well as compositions containing variable amounts of water. The invention includes all such solvates. Further, the term solvate encompasses solvates of both the free base compound as well as any salt thereof.

The skilled artisan will further appreciate that a compound of the present invention that exists in crystalline form, including the various solvates thereof, may exhibit polymorphism (i.e. the capacity to occur in different crystalline structures). These different crystalline forms are typically known as "polymorphs." The invention includes all such polymorphs. Polymorphs have the same chemical composition but differ in packing, geometrical arrangement, and other descriptive properties of the crystalline solid state. Polymorphs, therefore, may have different physical properties such as shape, density, hardness, deformability, stability, and dissolution properties. Polymorphs typically exhibit different melting points, IR spectra, and X-ray powder diffraction patterns, which may be used for identification. The skilled artisan will appreciate that different polymorphs may be produced, for example, by changing or adjusting the reaction conditions or reagents, used in making the compound. For example, changes in temperature, pressure, or solvent may result in polymorphs. In addition, one polymorph may spontaneously convert to another polymorph under certain conditions.

A compound of formula (I) may be prepared by the general synthetic schemes described hereinafter.

Scheme 1: Synthesis of 1,1-dimethylethyl (3*R*)-3-(aminomethyl)-3-fluoro-1-piperidinecarboxylate



a) 2,6-lutidine, ethyl 2-oxo-3-piperidinecarboxylate, [(S)-(-)-2,2'-bisphosphino)-1,1'-binaphthyl]palladium (II) dihydrate ditriflate, N-fluorobenzenesulfonimide, EtOH;

b) BH_3 -THF;


c) BOC_2O , Et_3N , DCM;

d) Tf_2O , Et_3N , DCM;

e) NaN_3 , DMF;

f) Pd/C, EtOH.


Scheme 2: Synthesis of 1-(1,1-dimethylethyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole



a) conc. HCl, EtOH;


- b) NBS, DCM;
c) Bis(pinacolato)diboron, KOAc, Pd₂(dba)₃, 2-(dicyclohexylphosphino)-2',4',6'-triisopropylbiphenyl, 1,4-dioxane.

5 **Scheme 3:** Synthesis of 1-[(methoxy)methyl]-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazole



- 10 a) Iodomethyl methyl ether, K₂CO₃, acetonitrile.

Scheme 4 (X=NH)




Where R = Et, *t*-Bu, CH₂OMe

- 5 a) 2,6-dichloro-4-pyridinamine, 2,2-dimethylpropanoyl chloride, Et₃N, DCM
 b) *n*BuLi, (2*E*)-3-(dimethylamino)-2-propenal, THF;
 c) 1,1-dimethylethyl (3*R*)-3-(aminomethyl)-3-fluoro-1-piperidinecarboxylate (see
 scheme 1), DIPEA, NMP;
 10 d) Pyrazole boronic ester, Cs₂CO₃, Pd(PPh₃)₄, 1,4-dioxane, H₂O; or pyrazole
 boronic ester, KOH, PEPPSI cat., DME, EtOH, H₂O;
 e) TFA, DCM.


Scheme 5 (X=O): Synthesis of 7-(1-ethyl-1*H*-pyrazol-4-yl)-5-({[(3*S*)-3-fluoro-3-piperidiny]methyl}oxy)-1,6-naphthyridine

15




- a) 1,1-dimethylethyl (3*S*)-3-fluoro-3-(hydroxymethyl)-1-piperidinecarboxylate (see scheme 1), NaH, DMF;
 5 b) 1-ethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazole, Cs₂CO₃, Pd(PPh₃)₄, 1,4-dioxane, H₂O;
 c) TFA, DCM.

Scheme 6: Synthesis of 7-[1-(1,1-dimethylethyl)-1*H*-pyrazol-4-yl]-N-[(3*S*)-3-fluoro-3-piperidinyl]methyl]-1,6-naphthyridin-5-amine



- a) (+)-Menthol, DMAP, toluene, reflux;
 b) [(S)BINAP]Pd(OTf)₂; (PhSO₂)₂NF; 2,6-lutidine; EtOH;
 c) (i) BH₃.DMS; THF; reflux;
 (ii) Boc₂O;
 d) Tf₂O, pyridine;
 e) NaN₃, DMF;
 f) Pt/C, hydrogen, THF, NH₃(aq);
 g) Compound of formula (XIX), DIPEA, NMP;
 h) Compound of formula (XIV), Pd(di-t-bpf)Cl₂, NaHCO₃, dioxane (aq);
 i) HCl/dioxane, toluene;
 j) (i) NaOH(aq), EtOAc;
 (ii) n-BuOAc, TBME.

Scheme 7: Synthesis of 5,7-dichloro-1,6-naphthyridine




a) $\text{Cu}(\text{OAc})_2$, *tert*-BuOK, IPA;

b) TEA, THF, ClCO_2Et ;

c) $\text{NH}_3(\text{aq})$, $\text{HCl}(\text{aq})$;

d) POCl_3 , Me_4NCl , reflux.

Scheme 8: Synthesis of 1-(1,1-dimethylethyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole



a) EtOH, HCl, reflux;


b) NBS, DCM;

c) Reflux;


d) BuLi, THF

Compounds of formulae (IV), (X), (XI), (XV), (XVII), (XXVII), (XXVIII), (XXXV) and (XXXVI) are commercially available from, for example, Sigma-Aldrich UK.

Thus, in one embodiment, the present invention provides a process for preparing a compound of formula (I) which process comprises reacting a 1,6-naphthyridine compound of formula (III):



wherein P is a protecting group and X is as hereinbefore defined;
with a pyrazole boronic ester/acid compound of formula (XXVI):



wherein R^3 and R^4 which may be the same or different, are each hydrogen, C_{1-6} alkyl or R^3 and R^4 may be joined to form a C_{1-3} alkylene group optionally substituted by up to four methyl groups, for instance $-C(Me)_2C(Me)_2-$; and R^1 is as hereinbefore defined;

- 5 in the presence of a catalyst, under conditions typically used for a boronic ester/acid coupling; and thereafter, removing the protecting group P.

10 Examples of protecting groups and the means for their removal can be found in T. W. Greene 'Protective Groups in Organic Synthesis' (J. Wiley and Sons, 1991). Suitable amine protecting groups include, but are not restricted to, sulphonyl (such as tosyl), acyl (such as benzyloxycarbonyl or t-butoxycarbonyl) and arylalkyl (such as benzyl), which may be removed by hydrolysis or hydrogenolysis as appropriate. Other suitable amine protecting groups include trifluoroacetyl ($-C(O)CF_3$), which may be removed by base catalysed hydrolysis, or a solid phase resin bound benzyl group, 15 such as a Merrifield resin bound 2,6-dimethoxybenzyl group (Ellman linker) which may be removed by acid catalysed hydrolysis (using, for example, trifluoroacetic acid).

20 In one embodiment of the present invention the protecting group (P) is selected from *tert*-butoxycarbonyl "BOC" and 9-fluorenylmethyloxycarbonyl "Fmoc".

It will be noted that, for synthetic efficiency, a precursor which is substantially enantiomerically pure is obtained as early as possible in the overall synthesis, in this instance, in the first stage of Scheme 1. Thus, enantioselective fluorination of the 25 piperidine nucleus at the 3-position, using an optically active catalyst ([(*S*)-(-)-2,2'-bisphosphino)-1,1'-binaphthyl]palladium (II) dihydrate ditriflate), leads to an enantiomerically enriched fluoro-product. This is then further resolved to give substantially enantiomerically pure product using, for instance, preparative chiral HPLC.

30 A compound of formula (I) may be prepared by the general synthetic schemes described hereinabove, using the racemic version of the catalyst ([(-)-2,2'-bisphosphino)-1,1'-binaphthyl]palladium (II) dihydrate ditriflate), in the first stage of Scheme 1, so there is no enantiomeric enrichment, and racemic intermediates are then used throughout. A compound of formula (II) may then be prepared from the 35 corresponding, racemic compound of formula (I), or an amine protected precursor, by methods known in the art for chiral resolution, in particular (preparative) chiral HPLC. Alternatively, resolution may be carried out at an earlier stage, if so desired, for instance, in Scheme 4, after stage c), after coupling of racemic amine to the 1,6-naphthyridine 40 core.

Compounds of the present invention are useful as inhibitors of Syk and thus potentially of use in treating some cancer therapies, in particular heme malignancies,

as well as inflammatory conditions which involve B cells, and also diseases resulting from inappropriate mast and basophil cell activation, for instance allergic and inflammatory diseases such as cutaneous mast cell mediated diseases including acute and chronic urticaria, mastocytosis, atopic dermatitis and autoimmune diseases such as cutaneous lupus and autoimmune bullous conditions including pemphigus and pemphigoid.

They may also be of use in treatment of acute conditions involving the mast cell such as post operative ileus.

Thus, in one embodiment, the present invention provides a compound of formula (I), or a pharmaceutically acceptable salt thereof, for use in therapy.

In one embodiment, the present invention provides a compound of formula (I) or a pharmaceutically acceptable salt thereof, for use in inhibiting spleen tyrosine kinase (Syk).

In another embodiment, the present invention provides a method of inhibiting spleen tyrosine kinase (Syk), which comprises administering to a subject in need thereof a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

Syk inhibitors may be useful in cancer therapy, specifically heme malignancies, particularly Non-Hodgkin's Lymphomas including follicular (FL), mantle cell, small lymphocytic lymphoma/chronic lymphocytic lymphoma (SLL/CLL), Burkitt and diffuse large B cell (DLBCL) lymphomas.

Accordingly, in one embodiment, the present invention provides a compound of formula (I) or a pharmaceutically acceptable salt thereof, for use in the treatment of cancer, for example heme malignancies, particularly Non-Hodgkin's lymphomas including follicular (FL), mantle cell, small lymphocytic lymphoma/chronic lymphocytic lymphoma (SLL/CLL), Burkitt and diffuse large B cell (DLBCL) lymphomas.

In another embodiment, the present invention provides a method of treating cancer, specifically heme malignancies, particularly Non-Hodgkin's Lymphomas including follicular (FL), mantle cell, small lymphocytic lymphoma/chronic lymphocytic lymphoma (SLL/CLL), Burkitt and diffuse large B cell (DLBCL) lymphomas, which method comprises administering to a patient in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof.

In a further embodiment, the present invention provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of cancer, for example heme malignancies, particularly Non-Hodgkin's lymphomas including follicular (FL), mantle cell, small lymphocytic

lymphoma/chronic lymphocytic lymphoma (SLL/CLL), Burkitt and diffuse large B cell (DLBCL) lymphomas.

5 Compounds of the present invention may also be used in cancer chemotherapy in combination with other classes of cancer chemotherapy agents which are known in the art. Representative classes of agents for use in such combinations for Non-Hodgkin's Lymphomas include rituximab, BEXXAR (tositumomab and Iodine I 131 tositumomab), pixantrone and chemotherapy. Combination of compounds of the present invention may also be used in combination with the CHOP drug regime
10 (cyclophosphamide, adriamycin, vincristine, prednisone) or CHOP plus rituximab (CHOP+R).

Compounds of the present invention are potentially of use in treating autoimmune conditions which involve B cells and/or macrophage activation, for instance Systemic
15 Lupus Erythematosus (SLE), Sjorgens Syndrome, Wegners granulomatosis and other vasculitides, bullous pemphigoid and pemphigus, idiopathic thrombocytopenic purpura (ITP), giant cell arteriosis, glomerulonephritis, chronic transplant rejection, and rheumatoid arthritis.

20 In one embodiment, the present invention provides a compound of formula (I) or a pharmaceutically acceptable salt thereof, for use in the treatment of an autoimmune condition, for example systemic lupus erythematosus (SLE), discoid (cutaneous) lupus, Sjorgens syndrome, Wegners granulomatosis and other vasculitides, bullous pemphigoid and pemphigus, idiopathic thrombocytopenic purpura (ITP), giant cell
25 arteriosis, glomerulonephritis, chronic transplant rejection, and rheumatoid arthritis. In another embodiment, the present invention provides a compound of formula (I) or a pharmaceutically acceptable salt thereof for use in the treatment of rheumatoid arthritis. In a further embodiment, the present invention provides a compound which is 7-[1-(1,1-dimethylethyl)-1*H*-pyrazol-4-yl]-*N*-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine or a pharmaceutically acceptable salt thereof for use in the
30 treatment of rheumatoid arthritis.

In one embodiment, the present invention provides a method of treating an autoimmune condition, for example systemic lupus erythematosus (SLE), discoid
35 (cutaneous) lupus, Sjorgens syndrome, Wegners granulomatosis and other vasculitides, bullous pemphigoid and pemphigus, idiopathic thrombocytopenic purpura (ITP), giant cell arteriosis, glomerulonephritis, chronic transplant rejection and rheumatoid arthritis, which method comprises administering to a patient in need thereof a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof. In another embodiment, the present invention provides a method of treating rheumatoid arthritis, which method comprises administering to a patient in need thereof a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof. In a further embodiment, the present invention provides a method of treating rheumatoid arthritis,
40

which method comprises administering to a patient in need thereof a therapeutically effective amount of a compound which is 7-[1-(1,1-dimethylethyl)-1*H*-pyrazol-4-yl]-*N*-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine or a pharmaceutically acceptable salt thereof.

5

In one embodiment, the present invention provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of an autoimmune condition, for example systemic lupus erythematosus (SLE), discoid (cutaneous) lupus, Sjorgens syndrome, Wegners granulomatosis and other vasculitides, bullous pemphigoid and pemphigus, idiopathic thrombocytopenic purpura (ITP), giant cell arteriosis, glomerulonephritis, chronic transplant rejection and rheumatoid arthritis. In another embodiment, the present invention provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of rheumatoid arthritis. In a further embodiment, the present invention provides the use of a compound which is 7-[1-(1,1-dimethylethyl)-1*H*-pyrazol-4-yl]-*N*-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of rheumatoid arthritis.

20

Compounds of the present invention are potentially of use in treating chronic idiopathic urticaria with and without auto-antibody status (now known as chronic spontaneous urticaria).

25

In one embodiment, the present invention provides a compound of formula (I) or a pharmaceutically acceptable salt thereof for use in the treatment of chronic spontaneous urticaria. In another embodiment, the present invention provides a compound which is 7-[1-(1,1-dimethylethyl)-1*H*-pyrazol-4-yl]-*N*-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine or a pharmaceutically acceptable salt thereof for use in the treatment chronic spontaneous urticaria.

30

In one embodiment, the present invention provides a method of treating chronic spontaneous urticaria, which method comprises administering to a patient in need thereof a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof. In another embodiment, the present invention provides a method of treating chronic spontaneous urticaria, which method comprises administering to a patient in need thereof a therapeutically effective amount of a compound of which is 7-[1-(1,1-dimethylethyl)-1*H*-pyrazol-4-yl]-*N*-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine or a pharmaceutically acceptable salt thereof.

40

In one embodiment, the present invention provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of chronic spontaneous urticaria. In another embodiment, the

present invention provides the use of a compound which is 7-[1-(1,1-dimethylethyl)-1*H*-pyrazol-4-yl]-*N*-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of chronic spontaneous urticaria.

5

In one embodiment, the present invention provides a compound of formula (I) or a pharmaceutically acceptable salt thereof, for use in the treatment of an inflammatory disease which involves B cells.

10 In one embodiment, the present invention provides a method of treating an inflammatory disease which involves B cells which method comprises administering to a patient in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof.

15 In one embodiment, the present invention provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of an inflammatory disease which involves B cells.

20 Compounds of the present invention are potentially of use in treating diseases resulting from inappropriate mast cell activation, for instance allergic and inflammatory diseases.

25 In one embodiment, the present invention provides a compound of formula (I) or a pharmaceutically acceptable salt thereof, for use in the treatment of a disease associated with inappropriate mast and/or basophil cell activation, including those diseases with skin manifestations.

30 In one embodiment, the present invention provides a method of treating a disease associated with inappropriate mast and/or basophil cell activation which comprises administering to a patient in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In another embodiment, the present invention provides a method of treating a disease associated with inappropriate mast and/or basophil cell activation including those diseases with skin manifestations, which comprises administering to a patient in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof.

40 In one embodiment, the present invention provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of a disease associated with inappropriate mast and/or basophil cell activation.

In one embodiment, the present invention provides a compound of formula (I) or a pharmaceutically acceptable salt thereof, for use in the treatment of an inflammatory

disease and/or allergic disorder for example, chronic obstructive pulmonary disease (COPD), adult respiratory distress syndrome (ARDS), asthma, severe asthma, ulcerative colitis, Crohn's disease, bronchitis, conjunctivitis, psoriasis, scleroderma, dermatitis, allergy, rhinitis, cutaneous lupus, autoimmune bullous conditions including pemphigus and pemphigoid, mastocytosis and anaphylaxis.

In one embodiment, the present invention provides a method of treating an inflammatory disease and/or allergic disorder for example, chronic obstructive pulmonary disease (COPD), adult respiratory distress syndrome (ARDS), asthma, severe asthma, ulcerative colitis, Crohn's disease, bronchitis, conjunctivitis, psoriasis, scleroderma, dermatitis, allergy, rhinitis, cutaneous lupus, autoimmune bullous conditions including pemphigus and pemphigoid, mastocytosis and anaphylaxis, which method comprises administering to a patient in need thereof a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

In one embodiment, the present invention provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of an inflammatory disease and/or allergic disorder for example, chronic obstructive pulmonary disease (COPD), adult respiratory distress syndrome (ARDS), asthma, severe asthma, ulcerative colitis, Crohn's disease, bronchitis, conjunctivitis, psoriasis, scleroderma, dermatitis, allergy, rhinitis, cutaneous lupus, autoimmune bullous conditions including pemphigus and pemphigoid, mastocytosis and anaphylaxis.

In one embodiment, the present invention provides a compound of formula (I) or a pharmaceutically acceptable salt thereof, for use in the treatment of an inflammatory disease which involves inappropriate mast cell activation.

In one embodiment, the present invention provides a method of treating an inflammatory disease which involves inappropriate mast cell activation which method comprises administering to a patient in need thereof an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof. In another embodiment, the present invention provides a method of treating an inflammatory disease which involves inappropriate mast cell activation which method comprises administering to a patient in need thereof an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

In one embodiment, the present invention provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of an inflammatory disease which involves inappropriate mast cell activation.

In one embodiment, the present invention provides a compound of formula (I) or a pharmaceutically acceptable salt thereof, for use in the treatment of an allergic disorder which involves inappropriate mast cell activation.

- 5 In one embodiment, the present invention provides a method of treating an allergic disorder which involves inappropriate mast cell activation which method comprises administering to a patient in need thereof an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof. In another embodiment, the present invention provides a method of treating an allergic disorder which involves
10 inappropriate mast cell activation which method comprises administering to a patient in need thereof an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

- 15 In one embodiment, the present invention provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of an allergic disorder which involves inappropriate mast cell activation.

- 20 Diseases and pathological conditions thought to be mediated by Syk include inflammatory and allergic disorders involving mast cell activation, such as chronic obstructive pulmonary disease (COPD), adult respiratory distress syndrome (ARDS), severe asthma, ulcerative colitis, Crohn's disease, bronchitis, conjunctivitis, psoriasis, scleroderma, chronic and acute urticaria, including chronic spontaneous urticaria and contact and physical urticarias, dermatitis, allergy, rhinitis, mastocytosis and
25 anaphylaxis.

- Compounds of the present invention may also be used in combination with other classes of therapeutic agents, for example selected from anti-inflammatory agents, anticholinergic agents (particularly an $M_1/M_2/M_3$ receptor antagonist), β_2 -
30 adrenoreceptor agonists, antiinfective agents such as antibiotics or antivirals, or antihistamines.

- In another embodiment, compounds of the present invention may also be used in combination with other classes of therapeutic agents which are known in the art for
35 treating autoimmune diseases, for instance disease modifying anti-rheumatic drugs including cyclosporine, methotrexate, sulphasalazine, prednisone, leflunomide, and chloroquine/hydrochloroquine and also biopharmaceutical agents such as humanised monoclonal antibodies (mabs), for example including anti-TNF alpha blockers such as remicade, enbrel and humira and B cell depleting therapies such as rituximab and
40 ofatumumab, and anti-Blys mabs such as belilumab.

The invention thus provides, a combination comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof together with one or more other therapeutically active agents, for example selected from an anti-inflammatory agent

such as a corticosteroid or an NSAID, an anticholinergic agent, a β_2 -adrenoreceptor agonist, an antiinfective agent such as an antibiotic or an antiviral, an antihistamine, a disease modifying anti-rheumatic drug, and a biopharmaceutical agent such as humanised monoclonal antibodies (mabs), B cell depleting therapies and anti-Blys mabs. One embodiment of the invention encompasses combinations comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof together with a β_2 -adrenoreceptor agonist, and/or an anticholinergic, and/or a PDE-4 inhibitor, and/or an antihistamine, and/or a disease modifying anti-rheumatic drug, and/or a biopharmaceutical agent.

One embodiment of the invention encompasses combinations comprising one or two other therapeutic agents.

It will be clear to a person skilled in the art that, where appropriate, the other therapeutic ingredient(s) may be used in the form of salts, for example as alkali metal or amine salts or as acid addition salts, or prodrugs, or as esters, for example lower alkyl esters, or as solvates, for example hydrates to optimise the activity and/or stability and/or physical characteristics, such as solubility, of the therapeutic ingredient. It will be clear also that, where appropriate, the therapeutic ingredients may be used in optically pure form.

Examples of β_2 -adrenoreceptor agonists include salmeterol (which may be a racemate or a single enantiomer such as the *R*-enantiomer), salbutamol (which may be a racemate or a single enantiomer such as the *R*-enantiomer), formoterol (which may be a racemate or a single stereoisomer such as the *R,R*-stereoisomer), salmefamol, fenoterol, carmoterol, etanterol, naminterol, clenbuterol, pirbuterol, flerbuterol, reproterol, bambuterol, indacaterol, terbutaline and salts thereof, for example the xinafoate (1-hydroxy-2-naphthalenecarboxylate) salt of salmeterol, the sulphate salt or free base of salbutamol or the fumarate salt of formoterol. In one embodiment the β_2 -adrenoreceptor agonists are long-acting β_2 -adrenoreceptor agonists, for example, compounds which provide effective bronchodilation for about 12 hours or longer.

Other β_2 -adrenoreceptor agonists include those described in WO02/066422, WO02/070490, WO02/076933, WO03/024439, WO03/072539, WO03/091204, WO04/016578, WO04/022547, WO04/037807, WO04/037773, WO04/037768, WO04/039762, WO04/039766, WO01/42193 and WO03/042160.

Examples of β_2 -adrenoreceptor agonists include:

3-(4-{[6-({(2*R*)-2-hydroxy-2-[4-hydroxy-3-(hydroxymethyl)phenyl]ethyl}amino)hexyl]oxy}butyl)benzenesulfonamide;
3-(3-{[7-({(2*R*)-2-hydroxy-2-[4-hydroxy-3-hydroxymethyl]phenyl]ethyl}-amino)heptyl]oxy}propyl)benzenesulfonamide;

- 4-[(1*R*)-2-[(6-{2-[(2, 6-dichlorobenzyl) oxy] ethoxy} hexyl) amino]-1-hydroxyethyl]-2-(hydroxymethyl) phenol;
 4-[(1*R*)-2-[(6-{4-[3-(cyclopentylsulfonyl)phenyl]butoxy}hexyl)amino]-1-hydroxyethyl]-2-(hydroxymethyl)phenol;
 5 N-[2-hydroxyl-5-[(1*R*)-1-hydroxy-2-[[2-4-[(2*R*)-2-hydroxy-2-phenylethyl]amino]phenyl]ethyl]amino]ethyl]phenyl]formamide;
 N-2{2-[4-(3-phenyl-4-methoxyphenyl)aminophenyl]ethyl}-2-hydroxy-2-(8-hydroxy-2(1*H*)-quinolinon-5-yl)ethylamine; and
 5-[(*R*)-2-(2-{4-[4-(2-amino-2-methyl-propoxy)-phenylamino]-phenyl}-ethylamino)-1-hydroxy-ethyl]-8-hydroxy-1*H*-quinolin-2-one.

The β_2 -adrenoreceptor agonist may be in the form of a salt formed with a pharmaceutically acceptable acid selected from sulphuric, hydrochloric, fumaric, hydroxynaphthoic (for example 1- or 3-hydroxy-2-naphthoic), cinnamic, substituted
 15 cinnamic, triphenylacetic, sulphamic, sulphanilic, naphthaleneacrylic, benzoic, 4-methoxybenzoic, 2- or 4-hydroxybenzoic, 4-chlorobenzoic and 4-phenylbenzoic acid.

Examples of corticosteroids may include those described in WO02/088167,
 20 WO02/100879, WO02/12265, WO02/12266, WO05/005451, WO05/005452, WO06/072599 and WO06/072600.

Anti-inflammatory corticosteroids are well known in the art. Representative examples include fluticasone propionate (e.g. see US patent 4,335,121), fluticasone furoate
 25 (e.g. see US patent 7,101,866), beclomethasone 17-propionate ester, beclomethasone 17,21-dipropionate ester, dexamethasone or an ester thereof, mometasone or an ester thereof (e.g. mometasone furoate), ciclesonide, budesonide, flunisolide, methyl prednisolone, prednisolone, dexamethasone and 6 α ,9 α -difluoro-11 β -hydroxy-16 α -methyl-3-oxo-17 α -(2,2,3,3-
 30 tetramethycyclopropylcarbonyl)oxy-androsta-1,4-diene-17 β -carbothioic acid S-cyanomethyl ester. Further examples of anti-inflammatory corticosteroids are described in WO02/088167, WO02/100879, WO02/12265, WO02/12266, WO05/005451, WO05/005452, WO06/072599 and WO06/072600.

35 Non-steroidal compounds having glucocorticoid agonism that may possess selectivity for transrepression over transactivation and that may be useful in combination therapy include those covered in the following published patent applications and patents: WO03/082827, WO98/54159, WO04/005229, WO04/009017, WO04/018429, WO03/104195, WO03/082787, WO03/082280, WO03/059899,
 40 WO03/101932, WO02/02565, WO01/16128, WO00/66590, WO03/086294, WO04/026248, WO03/061651, WO03/08277, WO06/000401, WO06/000398, WO06/015870, WO06/108699, WO07/000334 and WO07/054294.

Examples of anti-inflammatory agents include non-steroidal anti-inflammatory drugs (NSAID's).

Examples of NSAID's include sodium cromoglycate, nedocromil sodium, phosphodiesterase (PDE) inhibitors (for example, theophylline, PDE4 inhibitors or mixed PDE3/PDE4 inhibitors), leukotriene antagonists, inhibitors of leukotriene synthesis (for example montelukast), iNOS inhibitors, tryptase and elastase inhibitors, beta-2 integrin antagonists and adenosine receptor agonists or antagonists (e.g. adenosine 2a agonists), cytokine antagonists (for example chemokine antagonists, such as a CCR3 antagonist) or inhibitors of cytokine synthesis, or 5-lipoxygenase inhibitors. An iNOS (inducible nitric oxide synthase inhibitor) is preferably for oral administration. Examples of iNOS inhibitors include those disclosed in WO93/13055, WO98/30537, WO02/50021, WO95/34534 and WO 99/62875. Examples of CCR3 inhibitors include those disclosed in WO02/26722.

Examples of PDE4 inhibitors include *cis*-4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexan-1-carboxylic acid, 2-carbomethoxy-4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-one and *cis*-[4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-ol]. Also, *cis*-4-cyano-4-[3-(cyclopentyloxy)-4-methoxyphenyl]cyclohexane-1-carboxylic acid (also known as cilomilast) and its salts, esters, pro-drugs or physical forms (e.g. see U.S. patent 5,552,438).

Other compounds include AWD-12-281 from Elbion (Hofgen, N. *et al.* 15th EFMC Int Symp Med Chem (Sept 6-10, Edinburgh) 1998, Abst P.98; CAS reference No. 247584020-9); a 9-benzyladenine derivative nominated NCS-613 (INSERM); D-4418 from Chiroscience and Schering-Plough; a benzodiazepine PDE4 inhibitor identified as CI-1018 (PD-168787) and attributed to Pfizer; a benzodioxole derivative disclosed by Kyowa Hakko in WO99/16766; K-34 from Kyowa Hakko; V-11294A from Napp (Landells, L.J. *et al.* Eur Resp J [Annu Cong Eur Resp Soc (Sept 19-23, Geneva) 1998] 1998, 12 (Suppl. 28): Abst P2393); roflumilast (CAS reference No 162401-32-3) and a pthalazinone (e.g. see WO99/47505) from Byk-Gulden; Pumafentrine, (-)-p-[(4aR*,10bS*)-9-ethoxy-1,2,3,4,4a,10b-hexahydro-8-methoxy-2-methylbenzo[c][1,6]naphthyridin-6-yl]-N,N-diisopropylbenzamide which is a mixed PDE3/PDE4 inhibitor which has been prepared and published on by Byk-Gulden, now Altana; arofylline under development by Almirall-Prodesfarma; VM554/UM565 from Vernalis; or T-440 (Tanabe Seiyaku; Fuji, K. *et al.* J Pharmacol Exp Ther, 1998, 284(1): 162), and T2585.

Further compounds are disclosed in the published international patent application WO04/024728 (Glaxo Group Ltd), WO04/056823 (Glaxo Group Ltd) and WO04/103998 (Glaxo Group Ltd).

Examples of anticholinergic agents are those compounds that act as antagonists at the muscarinic receptors, in particular those compounds which are antagonists of the M₁ or M₃ receptors, dual antagonists of the M₁/M₃ or M₂/M₃ receptors or pan-antagonists of the M₁/M₂/M₃ receptors. Exemplary compounds for administration via inhalation include ipratropium (for example, as the bromide, CAS 22254-24-6, sold under the name Atrovent), oxitropium (for example, as the bromide, CAS 30286-75-0) and tiotropium (for example, as the bromide, CAS 136310-93-5, sold under the name Spiriva). Also of interest are revatropate (for example, as the hydrobromide, CAS 262586-79-8) and LAS-34273 which is disclosed in WO01/04118. Exemplary compounds for oral administration include pirenzepine (CAS 28797-61-7), darifenacin (CAS 133099-04-4, or CAS 133099-07-7 for the hydrobromide sold under the name Enablex), oxybutynin (CAS 5633-20-5, sold under the name Ditropan), terodiline (CAS 15793-40-5), tolterodine (CAS 124937-51-5, or CAS 124937-52-6 for the tartrate, sold under the name Detrol), otilonium (for example, as the bromide, CAS 26095-59-0, sold under the name Spasmomen), trospium chloride (CAS 10405-02-4) and solifenacin (CAS 242478-37-1, or CAS 242478-38-2 for the succinate also known as YM-905 and sold under the name Vesicare).

Other anticholinergic agents include compounds which are disclosed in US patent application 60/487981 including, for example:

(3-*endo*)-3-(2,2-di-2-thienylethenyl)-8,8-dimethyl-8-azoniabicyclo[3.2.1]octane bromide;
 (3-*endo*)-3-(2,2-diphenylethenyl)-8,8-dimethyl-8-azoniabicyclo[3.2.1]octane bromide;
 (3-*endo*)-3-(2,2-diphenylethenyl)-8,8-dimethyl-8-azoniabicyclo[3.2.1]octane 4-methylbenzenesulfonate;
 (3-*endo*)-8,8-dimethyl-3-[2-phenyl-2-(2-thienyl)ethenyl]-8-azoniabicyclo[3.2.1]octane bromide; and
 (3-*endo*)-8,8-dimethyl-3-[2-phenyl-2-(2-pyridinyl)ethenyl]-8-azoniabicyclo[3.2.1]octane bromide.

Further anticholinergic agents include compounds which are disclosed in US patent application 60/511009 including, for example:

(*endo*)-3-(2-methoxy-2,2-di-thiophen-2-yl-ethyl)-8,8-dimethyl-8-azoniabicyclo[3.2.1]octane iodide;
 3-((*endo*)-8-methyl-8-aza-bicyclo[3.2.1]oct-3-yl)-2,2-diphenyl-propionitrile;
 (*endo*)-8-methyl-3-(2,2,2-triphenyl-ethyl)-8-aza-bicyclo[3.2.1]octane;
 3-((*endo*)-8-methyl-8-aza-bicyclo[3.2.1]oct-3-yl)-2,2-diphenyl-propionamide;
 3-((*endo*)-8-methyl-8-aza-bicyclo[3.2.1]oct-3-yl)-2,2-diphenyl-propionic acid;
 (*endo*)-3-(2-cyano-2,2-diphenyl-ethyl)-8,8-dimethyl-8-azonia-bicyclo[3.2.1]octane iodide;
 (*endo*)-3-(2-cyano-2,2-diphenyl-ethyl)-8,8-dimethyl-8-azonia-bicyclo[3.2.1]octane bromide;
 3-((*endo*)-8-methyl-8-aza-bicyclo[3.2.1]oct-3-yl)-2,2-diphenyl-propan-1-ol;
 N-benzyl-3-((*endo*)-8-methyl-8-aza-bicyclo[3.2.1]oct-3-yl)-2,2-diphenyl-propionamide;

(*endo*)-3-(2-carbamoyl-2,2-diphenyl-ethyl)-8,8-dimethyl-8-azonia-bicyclo[3.2.1]octane iodide;

1-benzyl-3-[3-((*endo*)-8-methyl-8-aza-bicyclo[3.2.1]oct-3-yl)-2,2-diphenyl-propyl]-urea;

5 1-ethyl-3-[3-((*endo*)-8-methyl-8-aza-bicyclo[3.2.1]oct-3-yl)-2,2-diphenyl-propyl]-urea;

N-[3-((*endo*)-8-methyl-8-aza-bicyclo[3.2.1]oct-3-yl)-2,2-diphenyl-propyl]-acetamide;

N-[3-((*endo*)-8-methyl-8-aza-bicyclo[3.2.1]oct-3-yl)-2,2-diphenyl-propyl]-benzamide;

3-((*endo*)-8-methyl-8-aza-bicyclo[3.2.1]oct-3-yl)-2,2-di-thiophen-2-yl-propionitrile;

(*endo*)-3-(2-cyano-2,2-di-thiophen-2-yl-ethyl)-8,8-dimethyl-8-azonia-

10 bicyclo[3.2.1]octane iodide;

N-[3-((*endo*)-8-methyl-8-aza-bicyclo[3.2.1]oct-3-yl)-2,2-diphenyl-propyl]-

benzenesulfonamide;

[3-((*endo*)-8-methyl-8-aza-bicyclo[3.2.1]oct-3-yl)-2,2-diphenyl-propyl]-urea;

N-[3-((*endo*)-8-methyl-8-aza-bicyclo[3.2.1]oct-3-yl)-2,2-diphenyl-propyl]-

15 methanesulfonamide; and

(*endo*)-3-{2,2-diphenyl-3-[(1-phenyl-methanoyl)-amino]-propyl}-8,8-dimethyl-8-

azonia-bicyclo[3.2.1]octane bromide.

Further compounds include:

20 (*endo*)-3-(2-methoxy-2,2-di-thiophen-2-yl-ethyl)-8,8-dimethyl-8-azonia-

bicyclo[3.2.1]octane iodide;

(*endo*)-3-(2-cyano-2,2-diphenyl-ethyl)-8,8-dimethyl-8-azonia-bicyclo[3.2.1]octane

iodide;

(*endo*)-3-(2-cyano-2,2-diphenyl-ethyl)-8,8-dimethyl-8-azonia-bicyclo[3.2.1]octane

25 bromide;

(*endo*)-3-(2-carbamoyl-2,2-diphenyl-ethyl)-8,8-dimethyl-8-azonia-bicyclo[3.2.1]octane

iodide;

(*endo*)-3-(2-cyano-2,2-di-thiophen-2-yl-ethyl)-8,8-dimethyl-8-azonia-

bicyclo[3.2.1]octane iodide; and

30 (*endo*)-3-{2,2-diphenyl-3-[(1-phenyl-methanoyl)-amino]-propyl}-8,8-dimethyl-8-

azonia-bicyclo[3.2.1]octane bromide.

In one embodiment the invention provides a combination comprising a compound of

formula (I), or a pharmaceutically acceptable salt thereof, together with an H1

35 antagonist. Examples of H1 antagonists include, without limitation, methapyrilene,

desloratadine, amexanox, astemizole, azatadine, azelastine, acrivastine,

brompheniramine, cetirizine, levocetirizine, efletirizine, chlorpheniramine, clemastine,

cyclizine, carebastine, cyproheptadine, carbinoxamine, descarboethoxyloratadine,

doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine,

40 hydroxyzine, ketotifen, loratadine, levocabastine, mizolastine, mequitazine,

mianserin, noberastine, meclizine, norastemizole, olopatadine, picumast, pyrilamine,

promethazine, terfenadine, tripelennamine, temelastine, trimeprazine and triprolidine,

particularly cetirizine, levocetirizine, efletirizine and fexofenadine. In a further

embodiment the invention provides a combination comprising a compound of formula

(I), or a pharmaceutically acceptable salt thereof, together with an H3 antagonist (and/or inverse agonist). Examples of H3 antagonists include, for example, those compounds disclosed in WO2004/035556 and in WO2006/045416. Other histamine receptor antagonists which may be used in combination with the compounds of formula (I), or a pharmaceutically acceptable salt thereof, include antagonists (and/or inverse agonists) of the H4 receptor, for example, the compounds disclosed in Jablonowski *et al.*, *J. Med. Chem.* 46:3957-3960 (2003).

In one embodiment there is provided, a combination comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof together with a corticosteroid. In another embodiment there is provided, a combination comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof together with an NSAID. In another embodiment there is provided, a combination comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof together with an anticholinergic. In another embodiment there is provided, a combination comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof together with a β_2 -adrenoreceptor agonist. In another embodiment there is provided, a combination comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof together with an antiinfective. In another embodiment there is provided, a combination comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof together with an antihistamine. In another embodiment there is provided, a combination comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof together with a disease modifying anti-rheumatic drug. In a further embodiment there is provided, a combination comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof together with a biopharmaceutical agent.

A compound of the present invention will normally, but not necessarily, be formulated into pharmaceutical compositions prior to administration to a patient. Accordingly, in one embodiment the invention is directed to pharmaceutical compositions comprising a compound of formula (I), or a pharmaceutically acceptable salt thereof, and one or more pharmaceutically acceptable excipients. In another embodiment the invention is directed to a pharmaceutical composition comprising 7-[1-(1,1-dimethylethyl)-1H-pyrazol-4-yl]-N-[[[(3S)-3-fluoro-3-piperidiny]methyl]-1,6-naphthyridin-5-amine or a pharmaceutically acceptable salt thereof and one or more pharmaceutically acceptable excipients.

The pharmaceutical compositions of the invention may be prepared and packaged in bulk form wherein a safe and effective amount of a compound of the invention can be extracted and then given to the patient, such as with powders or syrups. Alternatively, the pharmaceutical compositions of the invention may be prepared and packaged in unit dosage form wherein each physically discrete unit contains a safe and effective amount of a compound of the invention. The pharmaceutical compositions of the invention may also be prepared and packaged in a sub-unit

dosage form wherein two or more sub-unit dosage forms provide the unit dosage form. When prepared in unit dosage form, the pharmaceutical compositions of the invention typically contain from about 0.1 to 99.9 wt.%, of a compound of the invention, depending on the nature of the formulation.

5

In addition, the pharmaceutical compositions of the invention may optionally further comprise one or more additional pharmaceutically active compounds.

As used herein, "pharmaceutically acceptable excipient" means a pharmaceutically acceptable material, composition or vehicle involved in giving form or consistency to the pharmaceutical composition. Each excipient must be compatible with the other ingredients of the pharmaceutical composition when commingled, such that interactions which would substantially reduce the efficacy of the compound of the invention when administered to a patient and would result in pharmaceutically unacceptable compositions are avoided. In addition, each excipient must of course be of sufficiently high purity to render it pharmaceutically acceptable.

Compositions of the present invention comprising a compound of formula (I), or a pharmaceutically acceptable salt thereof and the pharmaceutically acceptable excipient or excipients will typically be provided as a dosage form adapted for administration to the patient by the desired route of administration. For example, dosage forms include those adapted for: (1) oral administration, such as tablets, capsules, caplets, pills, troches, powders, syrups, elixers, suspensions, solutions, emulsions, sachets, and cachets; (2) topical dermal administration, such as creams, ointments, lotions, solutions, pastes, sprays, foams, and gels, (3) inhalation, such as aerosols and solutions; (4) intranasal administration, such as solutions or sprays; and (5) parenteral administration, such as sterile solutions, suspensions, and powders for reconstitution and.

It will be appreciated that dosage forms adapted for oral administration are commonly used for treating autoimmune disease including rheumatoid arthritis and systemic lupus erythematosus; cancer including heme malignancies; and chronic spontaneous urticaria. Dosage forms adapted for topical administration to the skin are commonly used for treating atopic dermatitis, psoriasis and chronic and acute urticaria conditions, and autoimmune bullous conditions including pemphigus and pemphigoid. Dosage forms adapted for inhalation or oral administration are commonly used for treating COPD and asthma; whilst dosage forms adapted for intranasal administration are commonly used for treating allergic rhinitis.

Suitable pharmaceutically acceptable excipients will vary depending upon the particular dosage form chosen. In addition, suitable pharmaceutically acceptable excipients may be chosen for a particular function that they may serve in the composition. For example, certain pharmaceutically acceptable excipients may be chosen for their ability to facilitate the production of uniform dosage forms. Certain

pharmaceutically acceptable excipients may be chosen for their ability to facilitate the production of stable dosage forms. Certain pharmaceutically acceptable excipients may be chosen for their ability to facilitate the carrying or transporting the compound of the present invention once administered to the patient from one organ, or portion
5 of the body, to another organ, or portion of the body. Certain pharmaceutically acceptable excipients may be chosen for their ability to enhance patient compliance.

Suitable pharmaceutically acceptable excipients include the following types of excipients: diluents, fillers, binders, disintegrants, lubricants, glidants, granulating
10 agents, coating agents, wetting agents, solvents, co-solvents, suspending agents, emulsifiers, sweeteners, flavouring agents, flavour masking agents, colouring agents, anticaking agents, humectants, chelating agents, plasticizers, viscosity increasing agents, antioxidants, preservatives, stabilizers, surfactants and buffering agents. The skilled artisan will appreciate that certain pharmaceutically acceptable excipients
15 may serve more than one function and may serve alternative functions depending on how much of the excipient is present in the formulation and what other ingredients are present in the formulation.

Skilled artisans possess the knowledge and skill in the art to enable them to select
20 suitable pharmaceutically acceptable excipients in appropriate amounts for use in the invention. In addition, there are a number of resources that are available to the skilled artisan which describe pharmaceutically acceptable excipients and may be useful in selecting suitable pharmaceutically acceptable excipients. Examples include Remington's Pharmaceutical Sciences (Mack Publishing Company), Remington: The
25 Science and Practice of Pharmacy, (Lippincott Williams & Wilkins), The Handbook of Pharmaceutical Additives (Gower Publishing Limited), and The Handbook of Pharmaceutical Excipients (the American Pharmaceutical Association and the Pharmaceutical Press).

30 The pharmaceutical compositions of the invention are prepared using techniques and methods known to those skilled in the art. Some of the methods commonly used in the art are described in Remington's Pharmaceutical Sciences (Mack Publishing Company).

35 Oral solid dosage forms such as tablets will typically comprise one or more pharmaceutically acceptable excipients, which may for example help impart satisfactory processing and compression characteristics, or provide additional desirable physical characteristics to the tablet. Such pharmaceutically acceptable excipients may be selected from diluents, binders, glidants, lubricants, disintegrants,
40 colourants, flavourants, sweetening agents, polymers, waxes or other solubility-modulating materials.

Dosage forms for topical administration to the skin may, for example, be in the form of ointments, creams, lotions, eye ointments, eye drops, ear drops, impregnated

dressings, and aerosols, and may contain appropriate conventional additives, including, for example, preservatives, solvents to assist drug penetration, and emollients in ointments and creams. Such topical formulations may also contain compatible conventional carriers, for example cream or ointment bases, and ethanol or oleyl alcohol for lotions. Such carriers may constitute from about 1% to about 98% by weight of the formulation; more usually they will constitute up to about 80% by weight of the formulation.

Dosage forms for parenteral administration will generally comprise fluids, particularly intravenous fluids, i.e., sterile solutions of simple chemicals such as sugars, amino acids or electrolytes, which can be easily carried by the circulatory system and assimilated. Such fluids are typically prepared with water for injection USP. Fluids used commonly for intravenous (IV) use are disclosed in Remington: The Science and Practice of Pharmacy, (Lippincott Williams & Wilkins). The pH of such IV fluids may vary, and will typically be from 3.5 to 8, as known in the art.

Dosage forms for nasal or inhaled administration may conveniently be formulated as aerosols, solutions, drops, gels or dry powders.

Dosage forms for topical administration to the nasal cavity (nasal administration) include pressurised aerosol formulations and aqueous formulations administered to the nose by pressurised pump. Formulations which are non-pressurised and adapted for nasal administration are of particular interest. Suitable formulations contain water as the diluent or carrier for this purpose. Aqueous formulations for administration to the nose may be provided with conventional excipients such as buffering agents, tonicity modifying agents and the like. Aqueous formulations may also be administered to the nose by nebulisation.

Dosage forms for nasal administration are provided in a metered dose device. The dosage form may be provided as a fluid formulation for delivery from a fluid dispenser having a dispensing nozzle or dispensing orifice through which a metered dose of the fluid formulation is dispensed upon the application of a user-applied force to a pump mechanism of the fluid dispenser. Such fluid dispensers are generally provided with a reservoir of multiple metered doses of the fluid formulation, the doses being dispensable upon sequential pump actuations. The dispensing nozzle or orifice may be configured for insertion into the nostrils of the user for spray dispensing of the fluid formulation into the nasal cavity. In one embodiment, the fluid dispenser is of the general type described and illustrated in WO2005/044354A1. The dispenser has a housing which houses a fluid discharge device having a compression pump mounted on a container for containing a fluid formulation. The housing has at least one finger-operable side lever which is movable inwardly with respect to the housing to cam the container upwardly in the housing to cause the pump to compress and pump a metered dose of the formulation out of a pump stem through a nasal nozzle of the

housing. A particularly preferred fluid dispenser is of the general type illustrated in Figures 30-40 of WO2005/044354A1.

Aerosol compositions, e.g. for inhaled administration, can comprise a solution or fine suspension of the active substance in a pharmaceutically acceptable aqueous or non-aqueous solvent. Aerosol formulations can be presented in single or multidose quantities in sterile form in a sealed container, which can take the form of a cartridge or refill for use with an atomising device or inhaler. Alternatively the sealed container may be a unitary dispensing device such as a single dose nasal inhaler or an aerosol dispenser fitted with a metering valve (metered dose inhaler) which is intended for disposal once the contents of the container have been exhausted.

Where the dosage form comprises an aerosol dispenser, it preferably contains a suitable propellant under pressure such as compressed air, carbon dioxide or an organic propellant such as a hydrofluorocarbon (HFC). Suitable HFC propellants include 1,1,1,2,3,3,3-heptafluoropropane and 1,1,1,2-tetrafluoroethane. The aerosol dosage forms can also take the form of a pump-atomiser. The pressurised aerosol may contain a solution or a suspension of the active compound. This may require the incorporation of additional excipients e.g. co-solvents and/or surfactants to improve the dispersion characteristics and homogeneity of suspension formulations. Solution formulations may also require the addition of co-solvents such as ethanol. Other excipient modifiers may also be incorporated to improve, for example, the stability and/or taste and/or fine particle mass characteristics (amount and/or profile) of the formulation.

For pharmaceutical compositions suitable and/or adapted for inhaled administration, it is preferred that the pharmaceutical composition is a dry powder inhalable composition. Such a composition can comprise a powder base such as lactose, glucose, trehalose, mannitol or starch, the compound of formula (I) or (II) or a pharmaceutically acceptable salt thereof (preferably in particle-size-reduced form, e.g. in micronised form), and optionally a performance modifier such as L-leucine or another amino acid, cellobiose octaacetate and/or metals salts of stearic acid such as magnesium or calcium stearate. Preferably, the dry powder inhalable composition comprises a dry powder blend of lactose and the compound of formula (I) or a pharmaceutically acceptable salt thereof. The lactose is preferably lactose hydrate e.g. lactose monohydrate and/or is preferably inhalation-grade and/or fine-grade lactose. Preferably, the particle size of the lactose is defined by 90% or more (by weight or by volume) of the lactose particles being less than 1000 microns (micrometres) (e.g. 10-1000 microns e.g. 30-1000 microns) in diameter, and/or 50% or more of the lactose particles being less than 500 microns (e.g. 10-500 microns) in diameter. More preferably, the particle size of the lactose is defined by 90% or more of the lactose particles being less than 300 microns (e.g. 10-300 microns e.g. 50-300 microns) in diameter, and/or 50% or more of the lactose particles being less than 100 microns in diameter. Optionally, the particle size of the lactose is defined by 90% or

more of the lactose particles being less than 100-200 microns in diameter, and/or 50% or more of the lactose particles being less than 40-70 microns in diameter. Most importantly, it is preferable that about 3 to about 30% (e.g. about 10%) (by weight or by volume) of the particles are less than 50 microns or less than 20 microns in diameter. For example, without limitation, a suitable inhalation-grade lactose is E9334 lactose (10% fines) (Borculo Domo Ingredients, Hanzeplein 25, 8017 JD Zwolle, Netherlands).

Optionally, in particular for dry powder inhalable compositions, a pharmaceutical composition for inhaled administration can be incorporated into a plurality of sealed dose containers (e.g. containing the dry powder composition) mounted longitudinally in a strip or ribbon inside a suitable inhalation device. The container is rupturable or peel-openable on demand and the dose of e.g. the dry powder composition can be administered by inhalation via the device such as the DISKUS[®] device, marketed by GlaxoSmithKline. The DISKUS[®] inhalation device is for example described in GB 2242134 A, and in such a device at least one container for the pharmaceutical composition in powder form (the container or containers preferably being a plurality of sealed dose containers mounted longitudinally in a strip or ribbon) is defined between two members peelably secured to one another; the device comprises: a means of defining an opening station for the said container or containers; a means for peeling the members apart at the opening station to open the container; and an outlet, communicating with the opened container, through which a user can inhale the pharmaceutical composition in powder form from the opened container.

A composition of the present invention, for intranasal administration, may also be adapted for dosing by insufflation, as a dry powder formulation.

For dosage forms for inhaled administration, where the compound or a pharmaceutically acceptable salt of formula (I) is present as a dry powder or in suspension, then it is preferred that it is in a particle-size-reduced form. Preferably the size-reduced form is obtained or obtainable by micronisation. The preferable particle size of the size-reduced (e.g. micronised) compound or pharmaceutically acceptable salt is defined by a D50 value of about 0.5 to about 10 microns (for example as measured using laser diffraction).

It will be appreciated that when the compounds of the present invention are administered in combination with other therapeutic agents normally administered by the inhaled, intravenous, oral, topical or intranasal route, that the resultant pharmaceutical composition may be administered by the same routes.

The compounds of the present invention may conveniently be administered in amounts of, for example, 1µg to 2g. The precise dose will of course depend on the age and condition of the patient and the particular route of administration chosen.

Biological test methods

Compounds of the invention may be tested for *in vitro* activity in accordance with the following assays:

5

1. Basic SYK enzyme activity

3 μ l of SYK lysate diluted 16-fold in assay buffer (20mM TRIS pH 7.4, 0.01% BSA, 0.1% Pluronic F-68) was added to wells containing 0.1 μ l of various concentrations of compound or DMSO vehicle (1.7% final) in a Greiner low volume 384 well black plate. Following 15min pre-incubation at room temperature, the reaction was initiated by the addition of 3 μ l of substrate reagent containing Y7 Sox peptide, (Invitrogen Cat. # KNZ3071, 5 μ M final), ATP (35 μ M final) and MgCl₂ (10mM final) in assay buffer. The reaction was incubated at room temperature before measuring fluorescence intensity (λ_{ex} 360/ λ_{em} 485) on an Envision plate reader (Perkin Elmer Life Sciences, Waltham, MA, USA) at 15min and 55min post-substrate addition.

The compounds of Examples 1, 2, 3, 4, 5, 7, 8, 9A, 9B, 10, 11A and 11B (as the free base) were tested essentially as described above and were each found to have an average IC₅₀ value in this assay of <1 μ M. The hydrochloric acid salts of compounds of Examples 1, 2, 3, 4 and 6 were tested essentially as described above and were each found to have an average IC₅₀ value in this assay of <1 μ M.

The compounds of Examples 1, 2, 3, 4, 5, 7, 8, 9A, 9B, 10, 11A and 11B (as the free base) were tested essentially as described above and were each found to have an average pIC₅₀ value in this assay of ≥ 7.0 . For example, Examples 1, 2, 3 and 4 (as the free base) were found to have average values of 7.6 (n = 7), 7.7 (n = 10), 7.2 (n = 11) and 8.1 (n = 11) respectively. The hydrochloric acid salts of compounds of Examples 1, 2, 3, 4 and 6 were tested essentially as described above and were each found to have an average pIC₅₀ value in this assay of ≥ 6.5 .

Those skilled in the art will recognize that *in vitro* binding assays and cell-based assays for functional activity are subject to variability. Accordingly, it is to be understood that the values for the IC₅₀s and pIC₅₀s recited above are exemplary only.

Preparation of SYK lysate

i. Preparation of Ramos cell lysates

40

Ramos B Cells (human B cells of Burkitt's lymphoma, clone 296.4C10, ATCC) were cultured in suspension in growth medium (RPMI-1640, Sigma; supplemented with 2mM L-glutamine, Gibco; 10mM Hepes, Sigma; 1mM sodium pyruvate, Sigma; 10% v/v heat-inactivated FCS, Gibco). Cells were grown in Corning Cellstacks (6360 cm²)

in 1 litre volume and viability and cell density were monitored daily. Cells were maintained at $<1.5 \times 10^6/\text{ml}$ and $>92\%$ viability.

Large scale production runs were generated from Large Scale Intermediate Aliquots (LSIA's) of frozen Ramos cells as this was found to give greater reproducibility than production from a continuously growing culture of Ramos cells.

The large scale production run cells were generated in four steps:

1. Thaw LSIA into 1 x Cellstack;
2. Expand culture into 4 x Cellstack;
3. Expand from 4 to 12 x Cellstacks;
4. Harvest all 12 Cellstacks.

Cellstacks were harvested in 2L centrifuge bottles using a Sorvall Mistral centrifuge, 2000rpm, 10 minutes, 4°C . ($2\text{L} \times 2 \times 10^6 \text{ cells/ml} = 4 \times 10^9 \text{ cells total}$)

(Notes for cell scale-up: If the cell density exceeded $1.8 \times 10^6/\text{ml}$ or viability dropped below 90% the Syk prep obtained post-stimulation was likely to be of lower activity.)

Also, repeated passage of the Ramos cells seemed to have a detrimental effect on Syk activity when cell growth is done at scale (this did not seem to be the case in small scale cultures) – it is recommended always to use LSIA's and modular scale-up for large scale preps.

ii. Stimulation of Ramos Cells with anti-IgM Ab to produce Syk & Preparation of lysates

Cells were stimulated at $20 \times 10^6 \text{ cells/ml}$ using $15 \mu\text{g/ml}$ (final concentration) anti-IgM antibody. Following harvest (as described above), a total of 4×10^9 cells were resuspended in 180ml pre-warmed (37°C) DPBS in a Corning 500ml centrifuge bottle. 20ml anti-IgM antibody at $150 \mu\text{g/ml}$ were added to each 500ml centrifuge bottle (working stock made up in DPBS pre-warmed to 37°C). Cells were incubated for exactly 5min at 37°C following the addition of anti IgM antibody. Following 5min stimulation, 300ml ice-cold DPBS were added to each bottle to stop the stimulation (temperature drops to $\sim 12^\circ\text{C}$) then cells were centrifuged at 2000rpm (Sorvall Legend RT+ centrifuge - pre-chilled to 4°C). Cells were washed by resuspension in ice-cold DPBS and centrifugation as above. The cell pellet was then lysed in ice-cold lysis buffer containing 1% triton-x-100 at a ratio of $150 \mu\text{l}/1 \times 10^7 \text{ cells}$ (i.e. 48ml lysis buffer). Following the addition of lysis buffer, the cells were pipetted up & down & kept on ice for 15min. The clarified lysate was then obtained by centrifugation (Sorvall Evolution RC (SLA-1500 rotor, $\sim 20,000g$ ($\sim 14,500\text{rpm}$), 45min, 4°C).

Lysate was aliquoted, snap-frozen on dry-ice & stored at -80°C prior to assay.

Materials

Ramos Cells: Human B cells of Burkitts lymphoma, clone 296.4C10 (ATCC).

Growth Media: 500ml RPMI, 10% heat inactivated FCS, 2mM L-Glutamine, 2mM HEPES, 1mM sodium pyruvate.

RPMI: Sigma R0883, stores CT5652

Foetal Calf Serum: Gibco 10099-141, stores CT2509

5 L-Glutamine: 200mM, Gibco 25030, stores CT3005

HEPES: 1M, Sigma H0887, stores CT5637

Sodium Pyruvate: 100mM, Sigma S8636, stores CT7741

Anti-IgM Ab: Goat anti-human IgM ((Fab')₂ fragments) in PBS. Invitrogen, custom-made preparation (azide free and low endotoxin levels). Catalogue no. NON0687,

10 Lot 1411913. 2.74mg/ml.

DPBS: Dulbeccos phosphate buffered saline, Sigma D8537

Lysis Buffer: 50mM TRIS pH7.5 + 150mM NaCl + 1% Triton-X-100 + 2mM EGTA + 1:100 dilution inhibitor cocktails (phosphatase inhibitor cocktail set II, calbiochem cat no. 524625 & protease inhibitor cocktail set V, calbiochem cat no. 539137)

15 Triton-X-100: Roche 10 789 704 001 (GI 198233X, SC/159824). Made up as a 20% stock in water.

EGTA: Sigma E4378. Added solid directly to buffer.

2. B cell activity assays

20

2.1. Ramos pErk assay

Principle of the assay

Ramos B cells (human B cells of Burkitt's Lymphoma) are stimulated using anti-IgM.

25 This results in the recruitment of Syk to the B cell receptor. The subsequent autophosphorylation of Syk leads to initiation of a signalling cascade resulting in B cell activation via the Erk MAP Kinase pathway. As a result Erk is phosphorylated and following cell lysis is detected by an immune capture assay.

30 Stimulation of Ramos cells with anti-IgM

Cells were plated at a density of 2.5×10^5 /well in a volume of 25µl assay medium (RPMI containing 10% heat inactivated foetal calf serum, 1% L-glutamine) in 96 v-well polypropylene plates. 25µl appropriately diluted compound solution was added and the plate incubated for 30min at 37°C with 5% CO₂. Cells were stimulated with
35 5µl Fab'₂ fragments of goat anti-human IgM (5µg/ml final) for 7min at 37°C. Cells are lysed by the addition of 55µL 2x RIPA lysis buffer for 2h at 4°C. Lysate may be frozen at this point at -80°C

pErk MSD assay

40 50µl cell lysate was transferred to a 96 well MSD plate coated with anti-pErk1/2 (Thr/Tyr: 202/204; 185/187) capture antibody and incubated for 16 hours at 4°C or 3 hours at room temperature. The plate was washed and an anti-pErk detection antibody added (25µl/well) for 1 hour at room temperature. This was removed, 150µL

MSD read buffer added and the resultant electrochemiluminescence signal measured.

Compound Preparation

5 Compound was prepared as a 10mM stock in DMSO and a dilution series prepared in DMSO using 9 successive 5-fold dilutions. This dilution series was diluted a further 1:100 with assay medium to give the final concentration range to be tested of 5×10^{-5} to 2.56×10^{-11} M. Compound dilutions were prepared using the Biomek 2000 and Biomek Nx automated robotic pipetting systems.

10 The compounds of Examples 1, 2, 3 and 4 (as the hydrochloride salts) were tested essentially as described above and were each found to have an average IC_{50} value in this assay of $<1 \mu M$.

The compounds of Examples 1, 2, 3 and 4 (as the hydrochloride salts) were tested essentially as described above and were each found to have an average pIC_{50} value
15 in this assay of ≥ 6.5 . For example, Examples 1, 2, 3 and 4 (as the hydrochloride salts) were found to have average values of 7.4 ($n = 2$), 7.2 ($n = 4$), 7.0 ($n = 2$) and 7.5 ($n = 4$) respectively.

Those skilled in the art will recognize that *in vitro* binding assays and cell-based
20 assays for functional activity are subject to variability. Accordingly, it is to be understood that the values for the IC_{50} s and pIC_{50} s recited above are exemplary only.

2.2 CD69 whole blood assay

Principle of the assay

Whole blood B cells are stimulated *ex-vivo* using anti-IgM. This results in the recruitment of Syk to the B cell receptor. The subsequent autophosphorylation of Syk leads to initiation of a signalling cascade resulting in B cell activation as indicated by expression of the activation marker CD69 on the cell surface. CD20/CD69+ve whole
30 blood B cells are detected by flow cytometry.

Stimulation of whole blood B cells with anti-IgM

100 μ l heparinised human blood was added to a 5ml polypropylene tube containing 1 μ l appropriately diluted compound solution and incubated for 25min at 37°C with 5%
35 CO₂. B cells were stimulated with 10 μ l Fab'₂ fragments of goat anti-human IgM (43 μ g/ml final concentration) for a further 3.5h under the conditions previously described. The red blood cells were lysed and all other cells fixed by the addition of 2ml Lyse/Fix buffer for 10min at room temperature.

CD69 assay

40 The cells were stained using a cocktail of mouse anti-human CD20 FITC and mouse anti-human CD69 PE conjugated antibodies. CD20/CD69+ve B cells present in the sample were detected by flow cytometry.

Compound Preparation

Compound was prepared as a 10mM stock in DMSO, further diluted down in DMSO to 1mM and a dilution series prepared in DMSO using 7 successive 3-fold dilutions to give the final concentration range to be tested of 1×10^{-5} to 4.57×10^{-9} M. Compound dilutions were prepared using the Biomek 2000 automated robotic pipetting system.

The compounds of Examples 1, 2, 3, 4, 7, 8, 9A, 9B, 10, 11A and 11B (as the free base) were tested essentially as described above and were each found to have an average IC_{50} value in this assay of $<1 \mu M$. The hydrochloric acid salts of compounds of Examples 1, 2, 3, 4 and 6 were tested essentially as described above and were each found to have an average IC_{50} value in this assay of $<1 \mu M$.

The compounds of Examples 1, 2, 3, 4, 5, 7, 8, 9A, 9B, 10, 11A and 11B (as the free base) were tested essentially as described above and were each found to have an average pIC_{50} value in this assay of ≥ 6.0 . For example, Examples 1, 2, 3 and 4 (as the free base) were found to have average values of 6.5 ($n = 2$), 6.8 ($n = 2$), 6.6 ($n = 2$) and 6.8 ($n = 2$) respectively. The hydrochloric acid salts of compounds of Examples 1, 2, 3, 4 and 6 were tested essentially as described above and were each found to have an average pIC_{50} value in this assay of ≥ 6.0 .

Those skilled in the art will recognize that *in vitro* binding assays and cell-based assays for functional activity are subject to variability. Accordingly, it is to be understood that the values for the IC_{50} s and pIC_{50} s recited above are exemplary only.

3. hERG activity - Cy3B Dofetilide fluoro-ligand binding assay for hERG

Compound potencies were determined by a fluoro-ligand^s (Cy3b-Dofetilide) fluorescence polarisation assay.

hERG-expressing CHO-K1 membranes* (60 $\mu g/ml$) were incubated with 1.0nM fluoro-ligand^s, in assay buffer (25mM HEPES, 1.2mM $MgCl_2$, 100mM KCl and 0.1% pluronic, pH adjusted to 7.4 using 5M KOH). The final potassium concentration in the assay was 100mM. After 70min mixing at room temperature, in the dark, 10 μl was dispensed into each well of a black LV Greiner 384-well plate containing 0.1 μl of test compound in DMSO. The plates were left to equilibrate for 2h before reading on an Acquest™ / Analyst™ imager. pIC_{50} data were generated using from an 11-point inhibition curve (top assay concentration of 50 μM and a 1:3 step-dilution), a six parameter curve-fit being applied using ABase and XC50 to analyse data and generate curve fits.

The compounds of Examples 1, 2, 3, 4, 5, 7, 8, 9A, 9B, 10, 11A and 11B (as the free base) were tested essentially as described above and were each found to have an average IC_{50} value in this assay of greater than 10 μM . The hydrochloric acid salts of

compounds of Examples 1, 2, 3, 4 and 6 were tested essentially as described above and were each found to have an average IC₅₀ value in this assay of greater than 20µM.

- 5 The compounds of Examples 1, 2, 3, 4, 5, 7, 8, 9A, 9B, 10, 11A and 11B were tested essentially as described above and were each found to have an average pIC₅₀ value in this assay of less than 5.0. The hydrochloric acid salts of compounds of Examples 1, 2, 3, 4 and 6 were tested essentially as described above and were each found to have an average pIC₅₀ value in this assay of less than 5.0.

10

Those skilled in the art will recognize that *in vitro* binding assays and cell-based assays for functional activity are subject to variability. Accordingly, it is to be understood that the values for the IC₅₀s and pIC₅₀s recited above are exemplary only.

15 * CHO-K1 membranes

- Chinese Hamster Ovary (CHO) cells stably expressing the human hERG receptor were grown to 80% confluency before being harvested by trypsinisation and subsequent centrifugation at 500g for 10min. Cell pellets were frozen at -80C before membrane production. The frozen pellet was thawed on ice, re-suspended and homogenised in 10 volumes of membrane buffer (50mM HEPES, pH 7.4, 1mM EDTA, 1mM PMSF, 2x10⁻⁶M Pepstatin A). The membrane suspension was centrifuged for 20min at 500g, the pellet discarded and the supernatant spun again at 48,000g for 30min. Following the second centrifugation the remaining pellet containing the membrane fraction was re-suspended in an appropriate volume (4ml for each ml of frozen cell pellet) and assayed for protein concentration.

25

[§]Fluoro-ligand (octahydrobenzo[2",3"]indolizino[8",7":5',6']pyrano[3',2':3,4]pyrido[1,2-a]indol-5-ium-2-sulfonate (International Application no. PCT/EP2010/050228, publication no. WO2010/097248A1 (Glaxo Group Ltd), TFA salt described in J.M.C. 2007, 50(13), 2931-2941).

30

- N-[4-({2-[(6-aminohexyl)(2-{4-[(methylsulfonyl)amino]phenyl}ethyl)amino]ethyl}oxy)phenyl]methanesulfonamide (1.508mg) as a solution in acetonitrile (100µl) was added to solid Cy3B-ONSu (14-{2-[(2,5-dioxo-1-pyrrolidinyl)oxy]-2-oxoethyl}-16,16,18,18-tetramethyl-6,7,7a,8a,9,10,16,18-octahydrobenzo[2",3"]indolizino [8",7":5',6']pyrano[3',2':3,4]pyrido[1,2-a]indol-5-ium-2-sulfonate (1.7 mg, WO9931181) in a silanised 4ml vial. A second portion of acetonitrile (100µl) was added followed by Hunig's base (0.9µl). Two portions (2 x 50µl) of dimethylformamide were added and the reaction mixture was concentrated under reduced pressure. The residue was re-dissolved in dimethylformamide (200µl). Hunig's base (0.9µl) was added and the mixture vortex mixed for 22h. The reaction mixture was evaporated to dryness, re-dissolved in acetonitrile/water/acetic acid (5/4/1, ~500µl), filtered and applied to a semi-preparative Spherisorb ODS2 HPLC column which was eluted with the following gradient (flow rate = 5ml/min, AU 5.0, 214nm, AU 2, 256nm, A=

40

0.1%TFA/water, B= 90% acetonitrile/10% water/0.1% TFA): t=0min: B=5%; t=10min: B=5%; t = 30min: B=25%; t=90min: B=55%; t=105min: B=100%; t=120min: B = 100%. The major component eluted between 46% and 48%B and collected in one fraction which was evaporated to dryness and the purple solid transferred to a vial using methanol as solvent. The methanol was removed under reduced pressure and the purple solid triturated with dry ether. The solid was dried overnight at 1mbar in a drying pistol to give the title compound (1.2mg).

N-[4-({2-[(6-aminohexyl)(2-{4-[(methylsulfonyl)amino]phenyl}ethyl)amino]ethyl}oxy)phenyl)methanesulfonamide

Crude N-[4-({2-[[6-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)hexyl](2-{4-[(methylsulfonyl)amino]phenyl}ethyl)amino]ethyl}oxy)phenyl)methanesulfonamide (142mg) was dissolved in methylamine (33% in ethanol, 10ml, 0.216) and left at 22°C for 48h. Excess reagent was evaporated under reduced pressure and the oily residue azeotroped with two further portions of ethanol. The crude product was dissolved in acetonitrile/water/acetic acid (5/4/, <2ml), half applied to a Phenomenex Jupiter C18 HPLC column and eluted using the following gradient (flow rate = 10ml/min, AU 20.0, 214nm, AU 10, 256nm, A= 0.1%TFA/water, B= 90% acetonitrile/10% water/0.1% TFA): t=0min: B=5%; t=10min: B=5%; t=100min: B=35%; t=115min: B=100%; t=130min: B = 100%. Fractions containing mainly the slower eluting component (>90%) were pooled and evaporated to give the title compound (14.9mg). The remaining crude was applied to the C18 column but with a modified gradient: t=0min: B=5%; t=10min: B=5%; t=15min: B=10%; t=95min: B=30%; t=110min: B=100%; t=125min: B = 100%. Fractions containing mainly the desired product were combined and evaporated as before to yield the title compound (21.3mg ~ 80% purity). The material was used without further purification.

N-[4-({2-[[6-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)hexyl](2-{4-[(methylsulfonyl)amino]phenyl}ethyl)amino]ethyl}oxy)phenyl)methanesulfonamide

2-[6-([2-(4-aminophenyl)ethyl]{2-[(4-aminophenyl)oxy]ethyl}amino)hexyl]-1H-isoindole-1,3(2H)-dione (108.3mg) was dissolved in DCM (5 ml) and cooled to 0-4°C in an ice-bath. Hunig's Base (0.227 ml) was added followed by the dropwise addition of mesylchloride (0.051ml). The reaction was maintained at 0-4°C for 0.5h and then allowed to warm slowly to room temperature. After 3h the reaction mixture was evaporated to dryness and used crude in next step.

2-[6-([2-(4-aminophenyl)ethyl]{2-[(4-aminophenyl)oxy]ethyl}amino)hexyl]-1H-isoindole-1,3(2H)-dione

2-[6-([2-(4-nitrophenyl)ethyl]{2-[(4-nitrophenyl)oxy]ethyl}amino)hexyl]-1H-isoindole-1,3(2H)-dione (0.35g) was dissolved in a mixture of ethanol (40ml), water (5ml) and acetic acid (5ml) and the resulting solution degassed under reduced pressure. 10%

Palladium on carbon (56% paste, 0.27g) was added and the resulting mixture stirred vigorously under a hydrogen atmosphere (atmospheric pressure) for 12h. The reaction mixture was filtered through Celite™ and washed with ethanol. The filtrate and washings were evaporated to dryness to give the title compound (0.313g) which was used without further purification.

2-[6-([2-(4-nitrophenyl)ethyl]{2-[(4-nitrophenyl)oxy]ethyl}amino)hexyl]-1H-isoindole-1,3(2H)-dione

[2-(4-Nitrophenyl)ethyl]{2-[(4-nitrophenyl)oxy]ethyl}amine (253mg) and 2-(6-bromohexyl)-1H-isoindole-1,3(2H)-dione (1186 mg) were dissolved in DMF (4ml) and basified by the addition of DIPEA (0.665ml). The reaction was stirred for 120h. The reaction mixture was evaporated to dryness and the residue dissolved in DCM, the solution was absorbed onto a pad of silica and purified on a silica cartridge (12g) eluting with the following gradient: (A = DCM, B= methanol) t=0min: B=10%; t=7.5min: B=0%; t=22.5min: B=5%. The desired product eluted at ~15%B (isocratically) and evaporation of the solution to dryness gave the title compound (0.364g).

[2-(4-nitrophenyl)ethyl]{2-[(4-nitrophenyl)oxy]ethyl}amine

[2-(4-nitrophenyl)ethyl]amine (498.9mg) and 111-[(2-bromoethyl)oxy]-4-nitrobenzene 2-bromoethyl 4-nitrophenyl ether (513 mg) were dissolved in DMF (5ml) at 22°C and DIPEA (0.872ml) added. The reaction mixture was left for 60h at 22°C, evaporated to dryness and the residue dissolved in DCM. The compound was absorbed onto silica and purified on a silica cartridge (12g) in two batches eluting with a methanol / DCM gradient (0-15%). Fractions containing pure product were pooled and the solvent removed under reduced pressure. The resulting title compound was isolated as a deep yellow oil which partially solidified under high vacuum (253mg).

4. Bacterial Mutation Screening Assay (Ames Test)

The bacterial mutation assay (Ames Test) is a short-term reverse mutation assay specifically designed to detect a wide range of chemical substances that can produce genetic damage that leads to gene mutations (base pair substitutions and frameshift mutations).

The test employs several histidine dependent *Salmonella typhimurium* strains and several tryptophan dependent strains of *Escherichia coli* each carrying different mutations in various genes in the histidine or tryptophan operon respectively. These mutations act as hot spots for mutagens that cause DNA damage via different mechanisms.

Briefly, bacteria are cultured for 10h prior to the test. Top agar is supplemented with trace amounts of histidine, biotin, and tryptophan and aliquoted, before adding the required test article, vehicle or positive control followed by the appropriate bacterial suspension, and either an exogenous mammalian oxidative metabolism system (S9-mix) or buffer solution. The final mixture is poured over minimal agar plates which are then inverted and incubated before scoring for revertant colonies.

When the tester strains are grown on a minimal media agar plate containing a trace of histidine or tryptophan, only those bacteria that revert to histidine or tryptophan independence are able to form colonies. The number of spontaneously induced revertant colonies per plate is relatively constant. However, when a mutagen is added to the plate, the number of revertant colonies per plate is increased, usually in a dose-related manner.

Bacterial Strains routinely used are *Salmonella typhimurium* strains TA98, TA100 TA1535 and TA1537 and *Escherichia coli* WP2 *uvrA* (pKM101).

Assays are performed in the presence and absence of an exogenous mammalian oxidative metabolising system (S9-mix), to mimic mammalian metabolism.

The highest concentration tested is one that allows maximum exposure up to 5000 µg/plate for freely soluble compounds, or the limit of solubility or toxicity (whichever is the lower). If compound solubility is a limiting factor, the maximum concentration chosen will be the lowest concentration at which compound precipitate is observed by eye on treatment plates at the end of the incubation period.

Appropriate vehicle and positive controls are included in all tests.

A single mutation assay (using the plate incorporation methodology) is conducted in the presence and absence of S9-mix (consisting of 4 replicate plates for vehicle controls and 2 replicate plates for the positive controls and test article).

Plates are usually scored electronically for bacterial colony formation following inspection of the plates for signs of toxicity (i.e. reduced growth (diminution) of background lawn, the presence of pin-dot/pseudorevertant colonies and/or a reduction in colony numbers). Plates will also be evaluated (by eye) for compound precipitation. Where compound precipitation occurs, scoring of bacterial colony formation for each strain will stop at the lowest treatment concentration at which compound precipitate is observed on the test plates at the end of the incubation period.

The number of colonies on each plate is recorded and mean values calculated for each concentration of test article used. Mean values are also be expressed as a ratio

of the mean concurrent vehicle control value (i.e. fold increase). In addition, any observed precipitate or toxicity are documented.

5 If the data for any treatment level shows a response ≥ 2 times the concurrent vehicle control value (TA98, TA100 TA102 WP2 uvrA (pKM101) and WP2 (pKM101)), or ≥ 3 times the concurrent vehicle control value (TA1535 and TA1537), in conjunction with a dose related response, the result is considered positive.

10 Where the data for any strain shows a dose related response, approaching the 2 or 3 fold threshold as detailed above, but does not exceed it, the result is considered equivocal and further testing may be required for clarification

15 The test methodology is based on established procedures for bacterial mutagenicity testing [Maron, 1983; Green, 1984; Ames, 1975; Garner, 1972; Green, 1976; Ames, 1973] and is in accordance with the general principles of the following regulatory guidelines.

20 The compounds of Examples 1, 2, 3 and 4 (as the free base) were negative in this assay.

COM (2000) Guidance on a Strategy for Testing of Chemicals for Mutagenicity, Committee on Mutagenicity of Chemicals in Food, Consumer Products and the Environment, December 2000.

25 ICH-S2A (1996) "Specific Aspects of Regulatory Genotoxicity Tests for Pharmaceuticals", in: Federal Register, April 1996 (61 FR 18199).

30 ICH-S2B (1997) "A Standard Battery for Genotoxicity Testing of Pharmaceuticals", in: Federal Register, November 1997 (62 FR 62472).

Gatehouse, D.G., Wilcox, P., Forster, R., Rowland, I.R. and Callander, R.D. (1990) Bacterial Mutation Assays. In: D.J. Kirkland (Ed.), Basic Mutagenicity Test: UKEMS Recommended Procedures. Cambridge University Press, Cambridge, UK, pp. 13 61.

35 OECD (1997) "Bacterial Reverse Mutation Test", in: OECD Guideline for the Testing of Chemicals, Test Guideline 471.

Intermediates and Examples

General

- 5 All temperatures are in °C.

Bis(pinacolato)diboron refers to 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi-1,3,2-dioxaborolane

BH₃.DMS refers to borane dimethylsulphide complex

- 10 BH₃-THF refers to borane tetrahydrofuran complex

BINAP refers to 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl

[(S)BINAP]Pd(OTf)₂ refers to (S)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl bis(acetonitrile)palladium (II) triflate

BOC refers to *tert*-butoxycarbonyl

- 15 BOC₂O refers to Di-*tert*-butyl dicarbonate

BuOH refers to butanol

tert-BuOK refers to potassium *tert*-butoxide

n-BuOAc refers to n-butyl acetate

ClCO₂Et refers to ethyl chloroformate

- 20 Cs₂CO₃ refers to caesium carbonate

Cu(OAc)₂ refers to copper acetate

CV refers to column volume

DCM refers to dichloromethane

DIPEA refers to *N, N*-diisopropylethylamine

- 25 DMAP refers to 4-dimethylaminopyridine

DME refers to dimethoxyethane

DMSO refers to dimethylsulfoxide.

DMF refers to *N,N*-dimethylformamide

Et₃N refers to triethylamine

- 30 Et₂O or Ether refers to diethyl ether

EtOAc refers to ethyl acetate

EtOH refers to ethanol

Fmoc refers to 9-fluorenylmethyloxycarbonyl

h refers to hours

- 35 HCl refers to hydrochloric acid

H₂O refers to water

HPLC refers to high performance liquid chromatography

IPA refers to *iso*-propanol

K₂CO₃ refers to potassium carbonate

- 40 KOAc refers to potassium acetate

KOH refers to potassium hydroxide

LCMS refers to liquid chromatography- mass spectroscopy

MDAP refers to mass directed autoprep

Me₄NCl refers to tetramethylammonium chloride

- MeOD refers to deuterated methanol
 min refers to minutes
 nBuLi refers to N-butyllithium
 NBS refers to N-bromosuccinimide
 5 NaHCO₃ refers to sodium bicarbonate
 Na₂SO₄ refers to sodium sulfate
 NaN₃ refers to sodium azide
 NMP refers to *N*-methylpyrrolidone
 (PhSO₂)₂NF or NFSI refers to N-fluorobenzenesulfonimide
 10 Pd/C refers to palladium on carbon
 PdCl₂.dppf refers to [1, 1'-bis(diphenylphosphino)ferrocene] dichloropalladium
 Pd₂(dba)₃ refers to tris(dibenzylideneacetone)dipalladium (0)
 Pd(PPh₃)₄ refers to tetrakis (triphenylphosphine) palladium (0)
 Pd(di-*t*-bpf)Cl₂ refers to 1, 1'-bis(di-*tert*-butylphosphino)ferrocene palladium dichloride
 15 PEPPSI refers to Pyridine-Enhanced Precatalyst Preparation Stabilization and Initiation
 POCl₃ refers to phosphorus oxychloride
 Pt/C refers to platinum on carbon
 r.t. refers to room temperature
 20 Rt refers to retention time
 TBME refers to *tert*-butyl methyl ether
 TEA refers to triethylamine
 TFA refers to trifluoroacetic acid
 Tf₂O refers to triflic anhydride
 25 THF refers to tetrahydrofuran

¹H NMR spectra were recorded using a Bruker DPX 400MHz, referenced to tetramethylsilane.

- 30 LC/MS (Method A) was conducted on an Acquity UPLC BEH C18 column (50mm x 2.1mm i.d. 1.7µm packing diameter) at 40 degrees centigrade, eluting with 10 mM ammonium bicarbonate in water adjusted to pH 10 with ammonia solution (Solvent A) and acetonitrile (Solvent B) using the following elution gradient 0-1.5min 1 – 97% B, 1.5-1.9min 97% B, 1.9 – 2.0min 100% B at a flow rate of 1ml/min. The UV detection
 35 was a summed signal from wavelength of 210nm to 350nm. The mass spectra were recorded on a Waters ZQ Mass Spectrometer using Alternate-scan Positive and Negative Electrospray. Ionisation data was rounded to the nearest integer.

- 40 LC/MS (Method B) was conducted on an Acquity UPLC BEH C18 column (50mm x 2.1mm i.d. 1.7µm packing diameter) at 40 degrees centigrade, eluting with 0.1% v/v solution of formic Acid in water (Solvent A) and 0.1% v/v solution of formic acid in acetonitrile (Solvent B) using the following elution gradient 0-1.5min 3 – 100% B, 1.5-1.9min 100% B, 1.9 – 2.0min 3% B at a flow rate of 1ml/min. The UV detection was a summed signal from wavelength of 210nm to 350nm. The mass spectra were

recorded on a Waters ZQ Mass Spectrometer using Alternate-scan Positive and Negative Electrospray. Ionisation data was rounded to the nearest integer.

- LC/MS (Method C) was conducted on an Acquity UPLC BEH C18 column (50mm x 2.1mm i.d. 1.7µm packing diameter) at 40 degrees centigrade, eluting with 0.1% v/v solution of trifluoroacetic acid in water (Solvent A) and 0.1% v/v solution of trifluoroacetic acid in acetonitrile (Solvent B) using the following elution gradient 0-1.5min 3 – 100% B, 1.5-1.9min 100% B, 1.9 – 2.0min 3% B at a flow rate of 1ml/min. The UV detection was a summed signal from wavelength of 210nm to 350nm. The mass spectra were recorded on a Waters ZQ Mass Spectrometer using Alternate-scan Positive and Negative Electrospray. Ionisation data was rounded to the nearest integer.

- MDAP (Method A). The HPLC analysis was conducted on an XBridge C18 column (100mm x 30mm i.d. 5µm packing diameter) at ambient temperature, eluting with 10mM ammonium bicarbonate in water adjusted to pH 10 with ammonia solution (Solvent A) and acetonitrile (Solvent B) using the following elution gradient:

Time (min)	Flow Rate (ml/min)	% A	% B
0	40	85	15
1	40	85	15
10	40	45	55
11	40	1	99
15	40	1	99

- The UV detection was an averaged signal from wavelength of 210nm to 350nm. The mass spectra were recorded on a Waters ZQ Mass Spectrometer using Alternate-scan Positive and Negative Electrospray. Ionisation data was rounded to the nearest integer.
- MDAP (Method B). The HPLC analysis was conducted on an XBridge C18 column (100mm x 30mm i.d. 5µm packing diameter) at ambient temperature, eluting with 10mM ammonium bicarbonate in water adjusted to pH 10 with ammonia solution (Solvent A) and acetonitrile (Solvent B) using the following elution gradient:

Time (min)	Flow Rate (ml/min)	% A	% B
0	40	85	15
1	40	85	15
20	40	45	55
21	40	1	99
25	40	1	99

The UV detection was an averaged signal from wavelength of 210nm to 350nm. The mass spectra were recorded on a Waters ZQ Mass Spectrometer using Alternate-

scan Positive and Negative Electrospray. Ionisation data was rounded to the nearest integer.

- 5 MDAP (Method C). The HPLC analysis was conducted on a Sunfire C18 column (150mm x 30mm i.d. 5µm packing diameter) at ambient temperature, eluting with 0.1% v/v solution of trifluoroacetic acid in water (Solvent A) and 0.1% v/v solution of trifluoroacetic acid in acetonitrile (Solvent B) using the following elution gradient:

Time (min)	Flow Rate (ml/min)	% A	% B
0	40	100	0
3	40	100	0
3.5	30	100	0
24.5	30	70	30
25	30	1	99
32	30	1	99

- 10 The UV detection was an averaged signal from wavelength of 210nm to 350nm. The mass spectra were recorded on a Waters ZQ Mass Spectrometer using Alternate-scan Positive and Negative Electrospray. Ionisation data was rounded to the nearest integer.

- 15 MDAP (Method D). The HPLC analysis was conducted on a Sunfire C18 column (150mm x 30mm i.d. 5µm packing diameter) at ambient temperature, eluting with 10mM ammonium bicarbonate in water adjusted to pH 10 with ammonia solution (Solvent A) and acetonitrile (Solvent B) using the following elution gradient:

Time (min)	Flow Rate (ml/min)	% A	% B
0	40	100	0
3	40	100	0
3.5	30	100	0
24.5	30	70	30
25	30	1	99
32	30	1	99

- 20 The UV detection was an averaged signal from wavelength of 210nm to 350nm. The mass spectra were recorded on a Waters ZQ Mass Spectrometer using Alternate-scan Positive and Negative Electrospray. Ionisation data was rounded to the nearest integer.

- 25 Silica chromatography techniques include either automated (Flashmaster, Biotage SP4) techniques or manual chromatography on pre-packed cartridges (SPE) or manually-packed flash columns.

When the name of a commercial supplier is given after the name of a compound or a reagent, for instance "compound X (Aldrich)" or "compound X / Aldrich", this means that compound X is obtainable from a commercial supplier, such as the commercial supplier named.


5

Similarly, when a literature or a patent reference is given after the name of a compound, for instance compound Y (EP 0 123 456), this means that the preparation of the compound is described in the named reference.

10 The names of the above mentioned Examples have been obtained using the compound naming programme "ACD Name Pro 6.02".

Intermediate 1: Ethyl (3S)-3-fluoro-2-oxo-3-piperidinecarboxylate

15



20

2,6-Lutidine (31.7g, 296mmol) (Aldrich) was added drop wise over 30min to a suspension of ethyl 2-oxo-3-piperidinecarboxylate (101.2g, 591mmol) (Aldrich), [(S)-(-)-2,2'-bisphosphino)-1,1'-binaphthyl]palladium (II) dihydrate ditriflate (3.14g, 2.96mmol) (Sodeoka, M et al. *Synlett* **1997**, 463-466; Fujii, A et al. *J. Am. Chem. Soc.* **1999**, 121, 5450-5458) (Aldrich) and *N*-fluorobenzenesulfonimide (242.0g, 768mmol) (Aldrich) in ethanol (500ml) at 0°C in an ice bath. The temperature was maintained at approximately 10°C during addition and then allowed to warm to room temperature overnight as ice melted. Presence of solid around flask (3L) neck suggests possible exotherm may have occurred overnight. LCMS showed presence of product. The reaction was filtered and the solid was washed with ethanol, then DCM (200ml). NMR confirmed no product in solid. The liquors were evaporated and re-dissolved in DCM (3500ml). The organics were washed with saturated ammonium chloride solution (300ml) and the aqueous was re-extracted with DCM (2x200ml). The combined organics were evaporated and re-dissolved in DCM (300ml), filtered through celite and washed with DCM (200ml). The organic solution was left to stand overnight (sealed so no evaporation) – a fine precipitate appeared. The mixture was filtered through celite again and washed with DCM.

35

The combined organic layers were loaded onto a 1500g silica column and purified on the companion XL eluting with 0-100% ethyl acetate in cyclohexane gradient. Appropriate fractions were identified by LCMS, combined and the solvent was removed to give the title compound as a yellow solid, which was dried under high vacuum for 1h (92.2g).

40

LCMS (Method B): R_t = 0.52min, MH^+ 190

Chiral analytical HPLC (25cm Chiralpak IA, col.no.IA00CE-MC024, 15%EtOH/C7, 1ml/min, wavelength 215nm, RT) showed enrichment of the fast eluter - 44% ee.

- 5 The compound was combined with three other batches of varying enantiomeric enrichment and 215g of material was purified further using preparative HPLC to improve the enantiomeric excess of the fast eluter to >99%.


- 10 Samples were prepared in batches according to the following method: ethyl-3-fluoro-2-oxo-3-piperidinecarboxylate (25g) was dissolved in ethanol (450ml) with gentle heating and sonication. The solution was then filtered through a Pall Acrodisc 37mm syringe filter with 1µm glass fibre membrane to remove fines and undissolved material. The filtered solution was adjusted to a total volume of 500ml with ethanol to give a solution with nominal concentration of 50mg/mL.

The preparative HPLC details were as follows:

Column	Chiralpak AD, 330× 50 mm, 20 µm	
Mobile Phase	A: Heptane	B: Ethanol
Gradient Profile	15% B Isocratic	
Run Time	20 min	
Flow Rate	473 mL/min	
Column Temperature	20 °C	
Wavelength	220 nm	


15

Intermediate 2: 1,1-Dimethylethyl (3S)-3-fluoro-3-(hydroxymethyl)-1-piperidinecarboxylate




- 20 Ethyl (3S)-3-fluoro-2-oxo-3-piperidinecarboxylate (50g, 264mmol) was dissolved in THF (100ml) and borane-THF complex (793ml, 793mmol, 1M solution) was added drop wise. The mixture was heated at reflux for 24h, cooled to room temperature and the borane quenched by addition of methanol (150ml). 2M HCl (200ml) was added and the mixture heated to reflux for 20min, then cooled and evaporated *in vacuo*.
- 25 The residue was suspended in DCM (500ml) and triethylamine (111ml, 793mmol) was added, followed by BOC anhydride (73.6ml, 317mmol). The mixture was stirred for 3h, then washed with water (100ml) and 0.5M HCl (100ml), dried and evaporated to give 1,1-dimethylethyl (3S)-3-fluoro-3-(hydroxymethyl)-1-piperidinecarboxylate as pale yellow crystalline solid (52.85g).
- 30 LCMS (Method B): Rt = 0.80min, MH⁺ 234

**Intermediate 3: 1,1-dimethylethyl (3S)-3-fluoro-3-
 ([(trifluoromethyl)sulfonyl]oxy)methyl)-1-piperidinecarboxylate**




- 5 Triflic anhydride (24.06ml, 142mmol) was added to a solution of 1,1-dimethylethyl (3S)-3-fluoro-3-(hydroxymethyl)-1-piperidinecarboxylate (30.2g, 129mmol) and triethylamine (23.46ml, 168mmol) in DCM (100ml) at -10°C over 20min. The mixture was stirred for 2h, allowing to warm to 0°C, then washed with water and brine, dried and evaporated to give 1,1-dimethylethyl (3S)-3-fluoro-3-
 10 ([(trifluoromethyl)sulfonyl]oxy)methyl)-1-piperidinecarboxylate as a dark brown oil (50.2g).
 LCMS (Method B): Rt = 1.23min, MH⁺ 366


**Intermediate 4: 1,1-dimethylethyl (3S)-3-(azidomethyl)-3-fluoro-1-
 15 piperidinecarboxylate**



- Sodium azide (9.79g, 151mmol) was added to a solution of 1,1-dimethylethyl (3S)-3-fluoro-3-([(trifluoromethyl)sulfonyl]oxy)methyl)-1-piperidinecarboxylate (50g,
 20 137mmol) in DMF (200ml) and the mixture was heated to 80°C for 1h. A sample was taken and quenched with water, extracted with ether and the ether layer evaporated *in vacuo*. The residue was analysed by NMR showing complete consumption of starting material.
 The mixture was cooled, diluted with water (1L) and extracted with EtOAc (2 x
 25 300ml). The solvent was washed with water (2 x 300ml), dried and evaporated to give 1,1-dimethylethyl (3S)-3-(azidomethyl)-3-fluoro-1-piperidinecarboxylate as an amber oil (36.7g).
 LCMS (Method B): Rt = 1.12min, MH⁺ 259


Intermediate 5: 1,1-Dimethylethyl (3R)-3-(aminomethyl)-3-fluoro-1-piperidinecarboxylate

- 5 1,1-Dimethylethyl (3S)-3-(azidomethyl)-3-fluoro-1-piperidinecarboxylate (36g, 139mmol) was dissolved in ethanol (500ml) and added under nitrogen to Pd/C (2.6g, 1.222mmol). The mixture was hydrogenated at atmospheric pressure overnight. The suspension was filtered and the filtrate evaporated *in vacuo* to give 1,1-dimethylethyl (3R)-3-(aminomethyl)-3-fluoro-1-piperidinecarboxylate as a pale yellow oil (32.7g).
- 10 ¹H NMR (CDCl₃) 3.75-3.52ppm (2H, 2xm, 2xCH), 3.30ppm (1H, dd, CH), 3.20ppm (1H, m, CH), 2.90-2.73ppm (2H, m, CH₂), 1.96-1.72ppm (2H, 2xm, CH₂), 1.70-1.58ppm (1H, m, CH), 1.57-1.43ppm (10H, m+s, CH + 3xCH₃), 1.32ppm (2H, br.s, NH₂).

Intermediate 6: N-(2,6-Dichloro-4-pyridinyl)-2,2-dimethylpropanamide


- 20 2,6-Dichloro-4-pyridinamine (10g, 61.3mmol) (Aldrich) and triethylamine (10.69ml, 77mmol) were combined in dichloromethane (DCM) (75ml) and cooled in an ice-bath. 2,2-dimethylpropanoyl chloride (8.30ml, 67.5mmol) (Aldrich) in DCM (15ml) was added drop wise and the mixture stirred whilst allowing to warm up overnight producing a clear dark orange solution. LCMS showed good conversion to product. The solution was washed with water and sat. NaHCO₃ (100ml each), dried with
- 25 Na₂SO₄, filtered and concentrated to yield a dark orange solid. The crude product was taken up in the minimum of DCM and applied to a 320g Companion XL silica column and eluted with 0% EtOAc in cyclohexane for 2CVs then 0-15% EtOAc in cyclohexane over 12CVs then held at 15% for 2CV. Appropriate fractions were combined and evaporated to give the title compound as a yellow solid (11.59g).
- 30 LCMS (Method B): Rt = 1.13min, MH⁺ 247/249

Intermediate 7: 5,7-Dichloro-1,6-naphthyridine



- 5 *N*-(2,6-Dichloro-4-pyridinyl)-2,2-dimethylpropanamide (1g, 4.05mmol) was taken up
 in THF (10ml) under nitrogen and cooled to $<-70^{\circ}\text{C}$. *n*-Butyl lithium (4.05ml,
 10.12mmol, 2.5M solution in hexanes) was added over 30min keeping the
 temperature below -60°C and then stirred at below -70°C for 1h. (2*E*)-3-
 (dimethylamino)-2-propenal (0.607ml, 6.07mmol) in THF (2ml) was added over
 30min keeping the temperature below -60°C . The reaction was stirred at below -70°C
 10 for 20min and then allowed to warm to room temperature. LCMS showed that no
 starting material remained so the reaction was quenched with 5M HCl (5ml) and
 refluxed overnight. LCMS showed good conversion to product so the reaction was
 cooled to room temperature. The reaction mixture was basified with solid K_2CO_3 and
 extracted with EtOAc (4x25ml). The combined organics were dried with Na_2SO_4 ,
 15 filtered and concentrated to yield a brown solid. The crude product was applied to a
 samplet and columned using a 40+M eluting with 12% diethyl ether in cyclohexane
 for 2CVs, then with 12%-63% diethyl ether in cyclohexane over 10CVs then held at
 63% for 5CVs. Appropriate fractions were combined and evaporated to give the title
 compound as a yellow solid (0.42g).
 20 LCMS (Method B): $\text{Rt} = 0.89\text{min}$, $\text{MH}^+ 199/201$

Intermediate 8: 1,1-Dimethylethyl (3*R*)-3-[(7-chloro-1,6-naphthyridin-5-yl)amino]methyl}-3-fluoro-1-piperidinecarboxylate




25

- To 5,7-dichloro-1,6-naphthyridine (5.01g, 25.2mmol) and 1,1-dimethylethyl (3*R*)-3-
 (aminomethyl)-3-fluoro-1-piperidinecarboxylate (5.32g, 22.90mmol) in NMP (20ml)
 was added DIPEA (8.00ml, 45.8mmol) and the mixture was stirred at 100°C for 72h
 30 under nitrogen. The reaction was cooled and partitioned between ethyl acetate and
 water (200ml each). The aqueous was washed with ethyl acetate. The combined
 organics were washed with water and the solvent was removed. The residue was
 dissolved in DCM and loaded onto a 100g silica SNAP column and purified on the
 SP4 eluting with 0-50% ethyl acetate in cyclohexane over 17CVs. Appropriate

fractions were combined and the solvent was removed to give the title compound as a pale orange solid which was dried under high vacuum for 2h (7.61g).


LCMS (Method B): R_t = 1.12min, MH^+ 395/397

5 **Intermediate 9: 1-(1,1-Dimethylethyl)-1H-pyrazole**




- 10 A mixture of (1,1-dimethylethyl)hydrazine hydrochloride (1.02g, 8.19mmol) (Aldrich), 1,1,3,3-tetrakis(methyloxy)propane (1.344g, 8.19mmol) (Aldrich) and hydrochloric acid (0.672ml, 8.19mmol) in ethanol (10ml) was heated under reflux. A solution formed. After 16h the reaction had gone to completion and so was concentrated *in vacuo* to yield the crude product. This was dissolved in methanol and passed through an aminopropyl cartridge (20g), eluting with methanol. The combined filtrates were concentrated *in vacuo* to yield the title compound as a cream gum (210mg)
- 15 LCMS (Method B): R_t = 0.74min, MH^+ = 125.6

Intermediate 10: 4-Bromo-1-(1,1-dimethylethyl)-1H-pyrazole



- 20 To an ice cooled solution of 1-(1,1-dimethylethyl)-1H-pyrazole (6.1g, 49.1mmol) in dichloromethane (DCM) (200ml) was added N-bromosuccinimide (8.74g, 49.1mmol). This was warmed to ambient temperature and stirred for 18h. There was no starting material present and so the reaction mixture was quenched with aqueous sodium thiosulfate. The organics were washed with water (2 x 500ml) and passed through a
- 25 hydrophobic frit. The filtrate was concentrated *in vacuo* to yield an orange-brown oil which turned purple on standing (7.402g).
- LCMS (Method B): R_t = 1.03 min, MH^+ = 203, 205


30 **Intermediate 11: 1-(1,1-Dimethylethyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole**



A mixture of 4-bromo-1-(1,1-dimethylethyl)-1H-pyrazole (2.4g, 11.82mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi-1,3,2-dioxaborolane (3.00g, 11.82mmol), potassium acetate (2.90g, 29.5mmol), tris(dibenzylideneacetone)dipalladium (0) (0.108g, 0.118mmol), 2-(dicyclohexylphosphino)-2',4',6'-triisopropylbiphenyl (0.225g, 0.473mmol) in 1,4-dioxane (30ml) was degassed with nitrogen. The reaction mixture was spilt into 2 microwave vials and heated at 110°C in a microwave for 1.5h. There was no starting material remaining so the reactions were filtered through a bond elut reservoir and the residue was washed with ethyl acetate. The filtrate was concentrated *in vacuo* to yield the crude product. This was dissolved in DCM and purified through silica (50g) eluting with 0-50% ethyl acetate in DCM gradient. Appropriate fractions were combined and concentrated *in vacuo* to yield the title compound as a beige solid (1.68g).

LCMS (Method B): Rt = 1.08min, MH⁺ = 250.8

Intermediate 12: 1-[(Methyloxy)methyl]-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole




4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (400mg, 2.061mmol) (Aldrich) was dissolved in acetonitrile (25ml) and stirred for 5min at 35°C. Iodomethyl methyl ether (0.873ml, 10.31mmol) and potassium carbonate (1424mg, 10.31mmol) were added and the mixture was allowed to stir at 35°C for 3h.

The LCMS showed incomplete reaction. The mixture was left to stir at 35°C for one additional hour. The LCMS showed no progression so 2eq (349μl) of the alkylating agent was added and the mixture was allowed to stir at 35°C for 30min. Ammonium chloride was then added. The mixture was partitioned between ethyl acetate and water. The aqueous layer was re-extracted with ethyl acetate and the combined organic phases were washed with water, dried using a hydrophobic frit and concentrated to an oil. The oil was purified on 50g silica using an SP4 and eluted

with a 0-100% ethyl acetate/cyclohexane gradient. The product was found in the waste, which was concentrated to give the title compound as an oil (260mg).

LCMS (Method B): $R_t = 0.84\text{min}$, $MH^+ = 238.85$

5 **Intermediate 13: 1,1-Dimethylethyl (3R)-3-([7-(1-ethyl-1H-pyrazol-4-yl)-1,6-naphthyridin-5-yl]amino)methyl]-3-fluoro-1-piperidinecarboxylate**



10 To a 2-5mL microwave vial under nitrogen was added caesium carbonate (990mg, 3.04mmol) and 1-ethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (292mg, 1.317mmol) (Boron Molecular). 1,1-Dimethylethyl (3R)-3-[[7-chloro-1,6-naphthyridin-5-yl]amino]methyl]-3-fluoro-1-piperidinecarboxylate (400mg, 1.013mmol) was dissolved in 1,4-dioxane (4ml) and water (0.8ml) and added in one aliquot. Nitrogen was bubbled through the resultant suspension for ~2min.

15 Tetrakis(triphenylphosphine)palladium (0) (117mg, 0.101mmol) was then added in one portion and nitrogen bubbled through the yellow suspension for a further ~1min. The microwave vial was sealed and heated at 150°C in a microwave reactor for 1h. LCMS showed the reaction had gone to completion. The reaction was partitioned between water (20ml) and ethyl acetate (20ml). The aqueous layer was further


20 extracted with ethyl acetate (2 x 20ml) and the combined organics washed with brine (10ml). The organics were dried (Na_2SO_4) and concentrated *in vacuo*.

The reaction was repeated as above using the same amounts and reaction conditions. The two crude batches were combined and purified using a 100g Biotage silica column and eluted with a 0-100% ethyl acetate/cyclohexane gradient.

25 Appropriate fractions were combined and concentrated *in vacuo* to give the title compound as a colourless oil (811mg)

LCMS (Method B): $R_t = 0.88\text{min}$, $MH^+ = 455$


30 **Intermediate 14: 1,1-Dimethylethyl (3R)-3-([7-[1-(1,1-dimethylethyl)-1H-pyrazol-4-yl]-1,6-naphthyridin-5-yl]amino)methyl]-3-fluoro-1-piperidinecarboxylate**



To a solution of 1,1-dimethylethyl (3R)-3-[[[(7-chloro-1,6-naphthyridin-5-yl)amino]methyl]-3-fluoro-1-piperidinecarboxylate (1.300g, 3.29mmol) in 1,4-dioxane (10ml) was added 1-(1,1-dimethylethyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (0.906g, 3.62mmol), caesium carbonate (2.145g, 6.58mmol), tetrakis(triphenylphosphine)palladium (0) (0.114g, 0.099mmol) and water (2ml). The reaction mixture was heated in the microwave at 130°C for 2h. LCMS shows main peak of product with no starting material. The reaction mixture was filtered through 10g celite and partitioned between ethyl acetate and water (100ml X 2). The organic layer was washed with brine (100ml), evaporated the solvent and dissolved in minimum amount of DCM. This was loaded on to a 100g silica column and purified on SP4 eluting with 20-90% ethyl acetate in cyclohexane over 22CVs. Appropriate fractions were collected and the solvent was evaporated. The product was dried on high vacuum overnight to give the title compound as a yellow crystalline solid (1.586g).

LCMS (Method B): Rt = 1.00min, MH⁺ 483

Intermediate 15: 1,1-Dimethylethyl (3R)-3-fluoro-3-[[[(7-{1-[(methoxy)methyl]-1H-pyrazol-4-yl}-1,6-naphthyridin-5-yl)amino]methyl]-1-piperidinecarboxylate




To 1,1-dimethylethyl (3R)-3-[[[(7-chloro-1,6-naphthyridin-5-yl)amino]methyl]-3-fluoro-1-piperidinecarboxylate (650mg, 1.646mmol) in DME (5ml), water (2.5ml), ethanol (5.00ml) was added 1-[(methoxy)methyl]-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (1.03g 4.33mmol), potassium hydroxide (3.95ml, 3.95 mmol, 1M aqueous solution) and PEPPSI (112mg, 0.165mmol). The reaction was refluxed at 130°C under nitrogen for 4 nights. LCMS showed main peak as product. The reaction mixture was filtered through celite and the solvent removed. The residue was

dissolved in DCM and loaded onto a 25g silica column and purified on the SP4 eluting with a 50-100% ethyl acetate in cyclohexane gradient. Appropriate fractions were combined and the solvent removed to give a yellow oil which was dried under high vacuum overnight to give the title compound as a yellow solid/film (813mg).

5 LCMS (Method B): Rt = 0.86min, MH⁺ 471

Intermediate 16: 1,1-Dimethylethyl (3S)-3-[(7-chloro-1,6-naphthyridin-5-yl)oxy]methyl}-3-fluoro-1-piperidinecarboxylate



10

To an ice cooled solution of 1,1-dimethylethyl (3S)-3-fluoro-3-(hydroxymethyl)-1-piperidinecarboxylate (1.406g, 6.03mmol) in DMF (20ml) was added sodium hydride (0.313g, 7.84mmol) (60% dispersion in mineral oil). This was stirred for 15min before adding 5,7-dichloro-1,6-naphthyridine (1.2g, 6.03mmol). This was warmed to room temp and stirred for 4h. The reaction had gone to completion and so was cautiously quenched with aqueous ammonium chloride before partitioning between aqueous ammonium chloride and ethyl acetate. The layers were separated and the aqueous was re-extracted with ethyl acetate. The combined organics were washed with brine and concentrated *in vacuo* to give the crude product. This was dissolved in DCM and purified through silica (50g), eluting with a gradient of 0-50% ethyl acetate in DCM. Appropriate fractions were concentrated *in vacuo* to yield the title compound as a cream foamy gum (1.91g).


15

20

LCMS: Rt = 1.18min, MH⁺ = 395.85

25

Intermediate 17: 1,1-Dimethylethyl (3S)-3-([7-(1-ethyl-1H-pyrazol-4-yl)-1,6-naphthyridin-5-yl]oxy)methyl}-3-fluoro-1-piperidinecarboxylate




30

A mixture of 1,1-dimethylethyl (3S)-3-[(7-chloro-1,6-naphthyridin-5-yl)oxy]methyl}-3-fluoro-1-piperidinecarboxylate (1.1g, 2.78mmol), 1-ethyl-4-(4,4,5,5-tetramethyl-1,3,2-

dioxaborolan-2-yl)-1H-pyrazole (0.679g, 3.06mmol) and caesium carbonate (2.263g, 6.95mmol) in 1,4-dioxane (16ml) and water (3ml) was degassed with nitrogen before adding tetrakis(triphenylphosphine)palladium (0) (0.096g, 0.083mmol). This was heated under reflux for 18h. The reaction was partitioned between ethyl acetate and aqueous ammonium chloride. The aqueous was re-extracted with ethyl acetate and the combined organics were washed with brine, passed through a hydrophobic frit and concentrated *in vacuo* to give crude product. This was dissolved in DCM and purified through silica (20g) eluting with a gradient of ethyl acetate in DCM. The product eluted in neat ethyl acetate. Appropriate fractions were combined and concentrated *in vacuo* to yield the title compound as a pinkish cream foam (1.10g). LCMS: Rt = 1.21min, MH⁺ = 456.09

Example 1. 7-(1-Ethyl-1H-pyrazol-4-yl)-N-[(3S)-3-fluoro-3-piperidinyl]methyl]-1,6-naphthyridin-5-amine



To a solution of 1,1-dimethylethyl (3R)-3-({[7-(1-ethyl-1H-pyrazol-4-yl)-1,6-naphthyridin-5-yl]amino}methyl)-3-fluoro-1-piperidinecarboxylate (811mg, 1.784mmol) in DCM (6.1ml) was added trifluoroacetic acid (3.16ml, 41.0mmol) and this was stirred at ambient temperature for 2h. After this time the reaction had gone to completion and so was concentrated *in vacuo* to yield the crude product. This was dissolved in methanol and loaded onto an SCX cartridge (70g) and eluted with methanol (3 column volumes) and 2M ammonia in methanol.

The filtrate from the ammonia fractions was concentrated *in vacuo* to give a yellow oil (653mg). The free base (553mg) was purified by High pH preparative MDAP (Method A). The appropriate vials were combined and concentrated *in vacuo* to afford the title compound as a yellow solid (265mg).

¹H NMR (d₆-DMSO) 8.65ppm (1H, dd, CH), 8.70ppm (1H, br.d, CH), 8.33ppm (1H, s, CH), 8.05ppm (1H, s, CH), 7.77ppm (1H, br.t, NH), 7.37ppm (1H, dd, CH), 7.25ppm (1H, s, CH), 4.19ppm (2H, q, CH₂), 3.95ppm (2H, m, CH₂), 2.85-2.50ppm (4H, 3xm, 2xCH₂), 1.90-1.54ppm (4H, 2xm, 2xCH₂), 1.42ppm (3H, t, CH₃).

HPLC (column: Luna 3u C18(2) , #409663-12, 8-minute run time at 1ml/min, eluent: 0-95% gradient of 0.05% TFA/Acetonitrile in 0.05% TFA/Water, wavelength 220nm): Rt = 2.49min


Optical rotation= -25.9 (c. 1.1 in methanol, T=23.6C)

Preparation of the HCl salt:

100mg of the free base was dissolved in DCM (1ml) and to this was added HCl (1.0M in Et₂O) (0.357mL, 0.357mmol). An orange solid immediately precipitated. The residual solvents were blown off and the resultant orange solid dried *in vacuo* to afford the title compound as the HCl salt (120mg), as an orange solid.

5 LCMS (Method B): Rt = 0.45min, MH⁺ 355

Example 2. 7-[1-(1,1-Dimethylethyl)-1H-pyrazol-4-yl]-N-[(3S)-3-fluoro-3-piperidinyl]methyl]-1,6-naphthyridin-5-amine



- 10 1,1-Dimethylethyl (3R)-3-[(7-[1-(1,1-dimethylethyl)-1H-pyrazol-4-yl]-1,6-naphthyridin-5-yl]amino)methyl]-3-fluoro-1-piperidinecarboxylate (1.586g, 3.29mmol) was dissolved in DCM (5ml) and trifluoroacetic acid (5.06ml, 65.7mmol) was added. The reaction was stirred for 20min at 20°C. LCMS showed main peak of product. The reaction mixture was loaded onto a 20g SCX cartridge and washed with methanol.
- 15 The compound was eluted with 2M methanolic ammonia. The solvent from the ammonia fractions was evaporated and was kept under high vacuum for 2h to give a yellow crystalline solid (1.171g). The free base (1.12g) was purified by High pH preparative MDAP (Method B). The appropriate fractions were combined, concentrated *in vacuo* and kept on high vacuum overnight to give the title compound as a yellow crystalline solid (688.1mg).
- 20 LCMS (Method B): Rt = 0.52min, MH⁺ 383
- ¹H NMR (MeOD) 8.80ppm (1H, dd, CH), 8.58ppm (1H, dd, CH), 8.32ppm (1H, s, CH), 8.08ppm (1H, s, CH), 7.39ppm (1H, dd, CH), 7.25ppm (1H, s, CH), 4.24ppm (1H, dd, CH), 3.82ppm (1H, dd, CH), 2.95-2.63ppm (4H, 2xm, 2xCH₂), 2.00-1.69ppm (4H, 3xm, 2xCH₂), 1.65ppm (9H, s, 3xCH₃).
- 25

Preparation of the HCl salt:

- 50mg of the free base was dissolved in dichloromethane (DCM) (1ml) and to this was added HCl (1.0M in Et₂O) (0.13ml). The solvents were blown down overnight and the resulting product was dried *in vacuo* for 1h to give the title compound as the HCl salt (56.6mg), as an orange solid.
- 30 LCMS (Method B): Rt = 0.53min, MH⁺ 383

Example 2. 7-[1-(1,1-Dimethylethyl)-1H-pyrazol-4-yl]-N-[(3S)-3-fluoro-3-piperidinyl]methyl]-1,6-naphthyridin-5-amine – Alternative Preparation

35


To a solution of 1,1-dimethylethyl (3*R*)-3-[(7-[1-(1,1-dimethylethyl)-1*H*-pyrazol-4-yl]-1,6-naphthyridin-5-yl)amino]methyl-3-fluoro-1-piperidinecarboxylate (2.24g, 4.64mmol) in DCM (14ml) was added TFA (8.22ml, 107mmol) and this was stirred at ambient temperature for 2h. After this time the reaction had gone to completion and so was concentrated *in vacuo* to yield the crude product. This was dissolved in methanol and loaded onto an SCX cartridge (50g). It was eluted with methanol (3CVs) and product eluted as the free base with 2M ammonia in methanol. The filtrate from the ammonia fractions was concentrated *in vacuo* to yield a yellow foam (1.55g).

A portion of this (305mg) was further purified by high pH MDAP to afford the title compound (265mg)

LCMS (Method C): $R_t = 0.56\text{min}$, MH^+ 383

Optical rotation = -34.2 (c. 1.01 in methanol, $T=23.6^\circ\text{C}$).

Example 3. N-[(3*S*)-3-Fluoro-3-piperidinyl]methyl-7-{1-[(methyloxy)methyl]-1*H*-pyrazol-4-yl}-1,6-naphthyridin-5-amine



To 1,1-dimethylethyl (3*R*)-3-fluoro-3-[(7-{1-[(methyloxy)methyl]-1*H*-pyrazol-4-yl}-1,6-naphthyridin-5-yl)amino]methyl-1-piperidinecarboxylate (813mg, 1.728mmol) in DCM (10ml) was added trifluoroacetic acid (2ml, 26.0mmol) and stirred at room temperature for 2h. LCMS showed product as the main peak. The solvent was removed and dissolved in methanol and loaded onto a pre-equilibrated 50g SCX cartridge. Washed with methanol and eluted with 2M methanolic ammonia. The solvent from the ammonia fractions was removed and the residue dried under high vacuum overnight. The residue was purified on the large scale high pH MDAP (Method D). Appropriate fractions were combined and the solvent removed. The resulting product was dried under high vacuum for 1h to give the title compound as a yellow solid (349mg).

LCMS (Method B): $R_t = 0.43\text{min}$, MH^+ 371


^1H NMR (MeOD) 8.84ppm (1H, br.d, CH), 8.60ppm (1H, br.d, CH), 8.44ppm (1H, s, CH), 8.18ppm, (1H, s, CH), 7.44ppm (1H, dd, CH), 7.31ppm (1H, s, CH), 5.49ppm (2H, s, CH_2), 4.25-3.87ppm (2H, 2xdd, CH_2), 3.38ppm (3H, s, CH_3), 3.21-2.75ppm (4H, 3xm, 2x CH_2), 2.10-1.75ppm (4H, 2xm, 2x CH_2).

Optical rotation = -27.7 (c. 0.74 in methanol, $T=23.6^\circ\text{C}$)

Preparation of the HCl salt:

47mg of the free base was dissolved in dichloromethane (DCM) (1ml) and a small amount of methanol and HCl (1.0M in Et₂O) (0.127mL, 1eq) was added. The solvent was removed and the residue was dried under high vacuum overnight to give the title compound as the HCl salt (48.5mg), as a pale orange/yellow solid
LCMS (Method B): Rt = 0.43min, MH⁺ 371

Example 4. 7-(1-Ethyl-1H-pyrazol-4-yl)-5-({[(3S)-3-fluoro-3-piperidinyl]methyl}oxy)-1,6-naphthyridine



To a solution of 1,1-dimethylethyl (3S)-3-({[7-(1-ethyl-1H-pyrazol-4-yl)-1,6-naphthyridin-5-yl]oxy}methyl)-3-fluoro-1-piperidinecarboxylate (1.1g, 2.415mmol) in DCM (3ml) was added trifluoroacetic acid (3ml, 38.9mmol) and this was stirred at room temp for 3h. The reaction had gone to completion and so was concentrated *in vacuo* to yield the crude product. The TFA salt was dissolved in methanol and loaded onto an SCX cartridge (20g). The impurities were eluted with methanol, and the product was eluted with 2M ammonia in methanol. The filtrate was concentrated *in vacuo* to yield a cream gummy solid (620mg). This was purified by large-scale MDAP (Method C). The appropriate vials were combined and concentrated *in vacuo* to give the product as the TFA salt. To form the free base, this was dissolved in methanol and purified through an SCX cartridge (20g) eluting with methanol. The product was eluted with 2M ammonia in methanol. The filtrate was concentrated *in vacuo* to yield the title compound as a clear glass (500mg).

LCMS (Method C): Rt = 0.51min, MH⁺ = 355.9

¹H NMR (MeOD) 8.92ppm (1H, dd, CH), 8.57ppm (1H, br.d, CH), 8.30ppm (1H, s, CH), 8.10ppm (1H, s, CH), 7.58ppm (1H, s, CH), 7.51ppm (1H, dd, CH), 4.68ppm (2H, m, CH₂), 4.26ppm (2H, q, CH₂), 3.30-2.60ppm (4H, 3xm, 2xCH₂), 2.20-1.62ppm (4H, 3xm, 2xCH₂), 1.52ppm (3H, t, CH₃).


Optical rotation= -7.6 (c. 0.79 in methanol, T=23.6°C)

Preparation of the HCl salt:



115mg of the free base was dissolved in DCM (2ml) and to this was added 1M HCl in diethyl ether (0.32ml). This was blown down under nitrogen and dried *in vacuo* to yield the title compound as the HCl salt (74mg), as a cream solid.
LCMS (Method C): Rt = 0.51 min, MH⁺ = 356

Examples 5 to 11

The following further examples were similarly prepared:



Name	R-	Alkylating agent ^s to form boronic ester	Stereo-chemistry	LCMS MH ⁺	LCMS Rt (min)
Ex 5: 1-{4-[5-({[(3S)-3-fluoro-3-piperidinyl]methyl}amino)-1,6-naphthyridin-7-yl]-1H-pyrazol-1-yl}-2-methyl-2-propanol	-CH ₂ C(Me) ₂ OH		homochiral	399	0.42 (Method B)
Ex 6: 2-{4-[5-({[(3S)-3-fluoro-3-piperidinyl]methyl}amino)-1,6-naphthyridin-7-yl]-1H-pyrazol-1-yl}ethanol	-CH ₂ CH ₂ OH		homochiral	371	0.38 (Method B)
Ex 7: N-({[(3S)-3-fluoro-3-piperidinyl]methyl}-7-{1-[2-(methyloxy)ethyl]-1H-pyrazol-4-yl}-1,6-naphthyridin-5-amine	-CH ₂ CH ₂ OMe		homochiral	385	0.43 (Method B)
Ex 8: 7-(1-cyclopentyl-1H-pyrazol-4-yl)-N-({[(3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine	-cyclopentyl		racemic	395	0.59 (Method C)
Ex 9 Enantiomer A: N-({[(3-fluoro-3-piperidinyl]methyl}-7-[1-(2,2,2-trifluoroethyl)-1H-pyrazol-4-yl]-1,6-naphthyridin-5-amine	-CH ₂ CF ₃		homochiral	409	0.50 (Method B)
Ex 9 Enantiomer B: N-({[(3-fluoro-3-piperidinyl]methyl}-7-[1-(2,2,2-trifluoroethyl)-1H-pyrazol-4-yl]-1,6-naphthyridin-5-amine	-CH ₂ CF ₃		homochiral	409	0.51 (Method B)
Ex 10: N-({[(3-fluoro-3-piperidinyl]methyl}-7-[1-(phenylmethyl)-1H-pyrazol-4-	-benzyl	Not synthesised – boronic ester commercially	racemic	417	0.6 (Method C)


yl]-1,6-naphthyridin-5-amine		available (Aldrich)			
Ex 11 Enantiomer A: <i>N</i> -{[3-fluoro-3-piperidinyl)methyl]-7-[1-(1-methylethyl)-1 <i>H</i> -pyrazol-4-yl]-1,6-naphthyridin-5-amine	CH(Me) ₂		homochiral	369	0.46 (Method B)
Ex 11 Enantiomer B: <i>N</i> -{[3-fluoro-3-piperidinyl)methyl]-7-[1-(1-methylethyl)-1 <i>H</i> -pyrazol-4-yl]-1,6-naphthyridin-5-amine	CH(Me) ₂		homochiral	369	0.47 (Method B)

* Example 11 was prepared using an Fmoc protected piperidine, rather than the BOC protected piperidine and separation of enantiomers at Fmoc protected formula (I).

- 5 § - Alkylating agents are commercially available from, for example, Sigma-Aldrich UK (Aldrich).

Process Description for Scheme 6

- 10 **Stage a): Preparation of (1*S*,2*R*,5*S*)-5-methyl-2-(1-methylethyl)cyclohexyl 2-oxo-3-piperidinecarboxylate**



- 15 4-Dimethylaminopyridine (0.35wt), (+)-menthol (0.95wt) and ethyl 2-oxo-3-piperidinecarboxylate (1wt) are dissolved/suspended in toluene (10vol). The mixture is heated to reflux and stirred for seven days with periodic distillation of solvent and replenishment with fresh toluene. The reaction mixture is cooled to 25°C and washed with two portions of 5%w/w aqueous HCl (4 vol) and then water (4 vol). The organic phase is concentrated under reduced pressure to give the title compound as a colourless oil.


20

Preparation of [(*S*)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl]bis(acetonitrile)palladium(II) ditriflate, catalyst for stage b)

- 25 Palladium acetate (1wt) is dissolved in acetonitrile (4vol) and stirred at 20±3°C for 1h. The reaction mixture is cooled to 0±3°C and trifluoromethanesulfonic acid (0.75vol, 1.9eq) is added slowly over 30min. After stirring for 15min at 0±3°C the reaction mixture is heated to 20±3°C and stirred for a further 30min. (*S*)-BINAP (2.77wt, 1eq) is added followed by a wash of acetonitrile (0.6vol) and the mixture is stirred at 20±3°C for 2.5h. Diisopropyl ether (10vol) is added slowly to give a slurry which is aged for 30min. The solid is filtered off under suction and washed with


acetonitrile:diisopropyl ether (1:3, 8vol) and diisopropyl ether (8vol). The product is then dried *in vacuo* at 20±3°C to constant probe temperature.

Stage b): Preparation of (1S,2R,5S)-5-methyl-2-(1-methylethyl)cyclohexyl (3S)-3-fluoro-2-oxo-3-piperidinecarboxylate



(1S,2R,5S)-5-Methyl-2-(1-methylethyl)cyclohexyl 2-oxo-3-piperidinecarboxylate (1wt) is dissolved in ethanol (1.5vol) and N-fluorobenzenesulfonimide (NFSI) (1.07wt) added, followed by [(S)-BINAP]Pd(CH₃CN)₂(OTf)₂ (0.038wt) at 20°C. Residual NFSI and catalyst were washed in with ethanol (3vol). The jacket is cooled to 0°C and 2,6-lutidine (0.21vol) charged slowly, maintaining the temperature below 20°C. The jacket temperature is then raised to 20°C. The reaction mixture is stirred for 24h, cooled to 0°C and stirred for an additional 2h. The solids are isolated by filtration, washing with ethanol (3vol) and dried under vacuum at 40°C. The dried crude solid is charged to a reactor, and dichloromethane (5vol) added. The dichloromethane phase is washed sequentially with 2M aqueous NaOH (3vol), water (3vol), 5%w/w aqueous HCl (3vol) and water (3vol). The dichloromethane extract is transferred to a rotary evaporator and solvent swapped with n-heptane under reduced pressure to give a mobile slurry and a final volume of ca 11vol. The product is isolated by filtration, washed with heptane (3vol) and dried under vacuum at 40°C.


Stage c): Preparation of 1,1-dimethylethyl (3S)-3-fluoro-3-(hydroxymethyl)-1-piperidinecarboxylate



(1S,2R,5S)-5-Methyl-2-(1-methylethyl)cyclohexyl (3S)-3-fluoro-2-oxo-3-piperidinecarboxylate (1eq, 1wt) is suspended in anhydrous THF (6vol) and treated with borane dimethyl sulfide complex (6eq, 1.9vol). The resulting solution is stirred at 63-66°C for 40-46h. The reaction mixture is cooled to 0±5°C and slowly added to cold (0±5°C) methanol (3vol) over 30-90min, maintaining the internal temperature below 15°C. Vigorous gas evolution and foaming is observed. 2M Hydrochloric acid (4vol) is added over 10-15min, maintaining the internal temperature below 15°C. The mixture is stirred at reflux (60-65°C) for 30-90min then cooled to 20-25°C. Toluene (4vol) is added, the mixture stirred for 15min then filtered. The filtrate is separated and the lower acidic aqueous phase run off. The filter cake is washed with 2M hydrochloric acid (2vol) and the wash used for a second extraction of the toluene phase. The lower acidic aqueous phase is run off, combined with the first acidic aqueous and cooled to 5±5°C. The solution is treated with sodium hydroxide (5eq, 0.668wt) and stirred at 10-20°C until the solid dissolved. A solution of di-tert-butyl-


dicarbonate (1.1eq, 0.802wt) in dichloromethane (2vol) is added and rinsed in with more dichloromethane (2vol). The two-phase mixture is stirred vigorously for 15h, filtered and the filter cake washed with dichloromethane (2vol). The lower organic phase is run off. The filter cake is washed with dichloromethane (2vol) and this wash
 5 used to extract the aqueous phase. The dichloromethane extracts are washed with water (4vol) and concentrated *in vacuo*. The residual oil is dried on the Buchi at 40°C. Heptane (4vol) is added and the mixture concentrated *in vacuo*. Towards the end of the evaporation the mixture is seeded with the title compound (0.001wt), the resultant slurry concentrated to a small volume and diluted with heptane (4vol). The
 10 slurry is rotated on Buchi at 40°C for 20-36min, cooled and rotated at 20±5°C for 1h. The solid is collected under suction, washed with heptane (2 x 1vol) and dried *in vacuo* at 35±5°C.

**Stage d): Preparation of 1,1-dimethylethyl (3S)-3-fluoro-3-
 15 {[(trifluoromethyl)sulfonyl]oxy}methyl)-1-piperidinecarboxylate**



1,1-Dimethylethyl (3S)-3-fluoro-3-(hydroxymethyl)-1-piperidinecarboxylate (1wt) in pyridine (3vol) is dissolved at 20°C under a nitrogen atmosphere. The resulting
 20 solution is cooled to -10°C, then triflic anhydride (0.80vol) is added at such a rate as to maintain the internal temperature below 5°C. The reaction mixture is stirred at -10°C for 4h, then monitored for completion by TLC. The batch is warmed to 0°C, isopropyl acetate (8vol) is added, followed by a solution of aqueous citric acid (5.8wt citric acid in 5.8vol of water). The mixture is stirred at 20°C for 15min, then the layers
 25 are separated. The organic layer is washed sequentially at 20°C with another solution of aqueous citric acid (5.8wt citric acid in 5.8vol of water), followed by aqueous sodium bicarbonate solution (0.35wt NaHCO₃ in water 4.65vol) and water (5vol). The organic phase is evaporated to dryness on a rotary evaporator to give the title compound as an orange oil.


**Stage e): Preparation of 1,1-dimethylethyl (3S)-3-(azidomethyl)-3-fluoro-1-
 30 piperidinecarboxylate**



1,1-Dimethylethyl (3S)-3-fluoro-3-[(trifluoromethyl)sulfonyl]oxy-1-piperidinecarboxylate (1 wt) is dissolved in *N,N*-dimethylformamide (3vol) at 20°C and sodium azide (1.1eq, 0.20wt) is added. The resulting suspension is stirred at 20-
 35 30°C for 6h. TBME (5vol) and water (9vol) are added. The mixture is stirred for 5min and the layers are separated. The aqueous phase is extracted with TBME (5vol) and the TBME phases are combined. The combined organics are washed with water (2 x


5vol), filtered and concentrated under reduced pressure to give the title compound as an orange oil.

Stage f): Preparation of 1,1-dimethylethyl (3*R*)-3-(aminomethyl)-3-fluoro-1-piperidinecarboxylate




The hydrogenation vessel is purged with nitrogen. 1,1-Dimethylethyl (3*S*)-3-(azidomethyl)-3-fluoro-1-piperidinecarboxylate (1wt) is dissolved in THF (11vol) and charged to the vessel, rinsing in with THF (2vol). Concentrated aqueous ammonia solution (~33% wt solution, 2vol) is added, followed by 10% platinum on carbon (Johnson Matthey type 128; ~50% wet, 8.3%wt). The vessel is evacuated and purged with nitrogen 3 times. The reaction vessel is purged with hydrogen, stirring is started and hydrogenation carried out at atmospheric pressure maintaining an internal temperature of 20°C for 38.5h (reaction complete by LCMS). The vessel is evacuated and purged with nitrogen three times, and the contents passed through a CUNO Zetacarbon filter (R55SP). The vessel is washed with THF (3vol) and this is passed through the previously used filter. The combined organics are concentrated to dryness on the rotary evaporator.

Stage g): Preparation of 1,1-dimethylethyl (3*R*)-3-[(7-chloro-1,6-naphthyridin-5-yl)amino]methyl}-3-fluoro-1-piperidinecarboxylate



A mixture of 5,7-dichloro-1,6-naphthyridine (1eq, 1wt), 1,1-dimethylethyl (3*R*)-3-(aminomethyl)-3-fluoro-1-piperidinecarboxylate (1.1eq, 1.28wt), and diisopropylethylamine (2eq, 1.75vol) in NMP (6vol) is stirred at 110±5°C for 17h. The solution is cooled to 70-75°C and water (6vol) added over 47min at 65-75°C. The mixture is aged at 65-75°C for 50min, then cooled to 45°C to give a slurry. The mixture is slowly cooled to 20-25°C then aged for 2.25h. The solid is collected under suction. The filter cake is washed with 1:2v/v NMP/water (2vol) then water (2 x 4vol) and dried *in vacuo* at 40±5°C. As the solid still contains significant NMP (16.2%w/w) it is reslurried in 1:1v/v NMP/water (7.3vol). The slurry is heated to 65°C and stirred at 65-66°C for 1h. The slurry is cooled to 20-25°C and aged for 3 days. The solid is collected under suction. The filter cake is washed with 1:2v/v NMP/water (2vol) then water (2 x 4vol) and dried *in vacuo* at 40±5°C.


Stages h) & i): Preparation of 7-[1-(1,1-dimethylethyl)-1H-pyrazol-4-yl]-N-{[(3S)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine dihydrochloride (Hydrochloride salt of Example 2)



- 5 1,1-Dimethylethyl (3R)-3-[[7-chloro-1,6-naphthyridin-5-yl]amino]methyl}-3-fluoro-1-piperidinecarboxylate (1wt), 1-(1,1-dimethylethyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (0.76wt), sodium bicarbonate (0.64wt) and 1,1'-bis(di-*tert*-butylphosphino)ferrocene palladium dichloride (0.00825wt) are dissolved/suspended in 1,4-dioxane (8vol) and water (2vol). The mixture is heated to reflux and stirred for 3h. The mixture is cooled to 20±3°C to form a suspension. The solid is filtered off under vacuum. The filtrate is concentrated to 2vol via vacuum distillation. Toluene (10vol) is added. The solution is washed with water (5vol) then stirred with Silicycle Si-Thiol derivatised silica gel (1wt) at 60-65°C for 2h. The mixture is filtered at 60-65°C and the cake washed with toluene (2vol). The organic filtrate is then heated to 60-63°C and 4M hydrogen chloride in 1,4-dioxane (2.9vol) added over 30min. The mixture is stirred at 60-63°C for 5h. The resulting slurry is cooled to 20±3°C and aged for 15.5h. The solid is filtered off under suction and washed with toluene (2 x 2vol). The product is dried *in vacuo* at 40±3°C to constant probe temperature.

20

Stage j): Preparation of 7-[1-(1,1-dimethylethyl)-1H-pyrazol-4-yl]-N-{[(3S)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine (Example 2)



- 25 7-[1-(1,1-Dimethylethyl)-1H-pyrazol-4-yl]-N-{[(3S)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine dihydrochloride (1wt) is dissolved/suspended in water (10vol) and washed with ethyl acetate (2 x 5vol). The acidic aqueous phase is basified with 32% NaOH (0.6vol) to pH 10 and extracted with ethyl acetate (2 x 5vol). The extracts are washed with 5%w/v sodium chloride in water (5vol). The combined extracts are concentrated on the Buchi to a solid. The residue is suspended in n-butyl acetate (2.8vol) and heated to 73-78°C to complete dissolution. The solution is line filtered through a 5µM domnick hunter filter, with a 0.2vol line wash. The solution is cooled to 45-50°C and seeded with the title compound (0.001wt). The mixture is cooled to 40-

30


45°C and aged for 50min. The slurry is diluted with TBME (3vol) over 30min at 40-45°C, and aged at 40-45°C for 1h. The slurry is cooled to 20-23°C and aged for 16h. The solid is filtered, washed with 1:1v/v n-butyl acetate/TBME (1vol) followed by TBME (2 x 2vol) and dried *in vacuo* at 40±5°C to constant probe temperature.

5

Process Description for Scheme 7

Stage a): Preparation of 2-(2-ethoxy-2-oxoethyl)nicotinic acid and 2-(2-isopropoxy-2-oxoethyl)nicotinic acid

10



ratio ~2:1

The reactor is charged with potassium *tert*-butoxide (3eq) followed by iso-propanol (9vol) under nitrogen atmosphere at 30±5°C. The mixture is stirred to dissolve. After cooling to ~30°C, ethyl acetoacetate (2eq) is slowly added under nitrogen atmosphere at <40°C. After stirring for 1-2h copper acetate (0.1eq) is charged at 35±5°C, followed by 2-chloronicotinic acid (1eq). This is followed by a rinse of isopropyl alcohol (1vol). The reaction mixture is heated to 78±3 °C under nitrogen atmosphere for at least 4h then sampled. On receipt of a successful sample result the reaction is cooled to 30±5°C and quenched with water, followed by dilute HCl to adjust the pH to 6-7.

15

20


The reaction mixture is concentrated by vacuum distillation at <50 °C to 4-5 volumes. 10% NaCl solution is added followed by ethyl acetate. Further dilute HCl is added to adjust the pH to 2.5-3.0. The biphasic mixture is then filtered through a celite bed and the cake washed with ethyl acetate (twice). The ethyl acetate and aqueous layers are separated then the aqueous layer back extracted with ethyl acetate (twice). The ethyl acetate phase is washed with water and 5% aq. NaCl solution. The organic phase is distilled to 1-2 volumes under vacuum at <50°C. Toluene is added and the mixture distilled again down to 1-2 volumes. This toluene addition/distillation step is repeated. The resulting toluene suspension is cooled to 30±5°C and treated with hexane (10vol). The suspension is stirred for 1-2h at 30±5°C then filtered. The solid is washed with hexane (2vol) then offloaded and dried at 52.5±2.5 °C under vacuum (NLT 650 mm of Hg) for 10h.

25

30

Stages b) & c): Preparation of 1,6-naphthyridin-5,7(6H,8H)-dione


35



Tetrahydrofuran (9vol) is charged to the reactor followed by the starting material ester (1eq) and a rinse of tetrahydrofuran (1vol). The mixture is cooled to $-2.5 \pm 7.5^{\circ}\text{C}$ under nitrogen atmosphere then treated with triethylamine (1.35eq) and later ethyl chloroformate (1.25eq). The reaction is stirred for 1h at $-2.5 \pm 7.5^{\circ}\text{C}$ then sampled.

- 5 On receipt of a successful result the reaction is carefully treated with aq. ammonia, stirred, then water is added. Dilute HCl is added at $<5^{\circ}\text{C}$ to adjust the pH to 6.5-7.5. The mixture is distilled under vacuum at $<40^{\circ}\text{C}$ to 12-14 volumes and cooled to $7.5 \pm 2.5^{\circ}\text{C}$. The suspension is stirred $\sim 1\text{h}$ at $7.5 \pm 2.5^{\circ}\text{C}$ then filtered. The solid is washed with water (1vol) then offloaded and dried under vacuum (NLT 650 mm of Hg) at $52.5 \pm 2.5^{\circ}\text{C}$ for 12.0h.
- 10

Stages d) & e): Preparation of 5,7-Dichloro-1,6-naphthyridine




- 15 Phosphorus oxychloride (3vol) is charged to the reactor at $30 \pm 5^{\circ}\text{C}$ followed by 1,6-naphthyridin-5,7(6H,8H)-dione (1eq), tetramethylammonium chloride (1.05eq) and a rinse of phosphorus oxychloride (0.5vol). The mixture is heated to $103.5 \pm 3.5^{\circ}\text{C}$ and stirred for a minimum of 36h. The reaction is then cooled to $60 \pm 5^{\circ}\text{C}$ and sampled. On receipt of a successful sample, toluene (4vol) is added and the mixture
- 20 concentrated to 2-3 volumes by distillation under vacuum (NLT 650 mm of Hg) and temperature $<60^{\circ}\text{C}$. Additional toluene (2vol) is added and again the mixture distilled under vacuum to 2-3 vols. This toluene addition/distillation cycle is repeated once more then the mixture is cooled to $47.5 \pm 2.5^{\circ}\text{C}$ and treated with tetrahydrofuran (8vol) and ethyl acetate (8vol). The mixture is cooled to $30 \pm 5^{\circ}\text{C}$ and poured into 15% aq.
- 25 sodium carbonate (15vol) which has been pre-cooled to $5.0 \pm 5.0^{\circ}\text{C}$. Additional tetrahydrofuran (3vol) is added along with ethyl acetate (3vol) in order to rinse out the initial reaction vessel. Whilst still at $5 \pm 5^{\circ}\text{C}$ the mixture is slowly treated with further 15% aqueous sodium carbonate solution (5vol) and stirred for 15-30min. The temperature is increased to $22.5 \pm 2.5^{\circ}\text{C}$ and the pH adjusted to 7-8 by addition of
- 30 either 15% aqueous sodium carbonate solution or dilute HCl. Ethyl acetate (5vol) is added followed by celite. The mixture is stirred then filtered. The celite bed is washed with ethyl acetate (2x2vol). The filtrate is distilled at $<55^{\circ}\text{C}$ under vacuum (NLT 650 mm of Hg) down to 30 ± 5 volumes. Water (2vol) is added and the distilled again down to ~ 30 vols. This water addition/distillation cycle is repeated until the tetrahydrofuran,
- 35 toluene and ethyl acetate content are each NMT 3.0% by GC (% w/w). At that point, water (10vol) is added and the mixture heated to $42.5 \pm 2.5^{\circ}\text{C}$ and stirred for 30-60min. The product is isolated by filtration at $42.5 \pm 2.5^{\circ}\text{C}$ and washed with water (2x3vol). The cake is sucked dry then offloaded and charged to a separate reactor. Water (20vol) is added and the suspension heated to $42.5 \pm 2.5^{\circ}\text{C}$ and stirred for 30-
- 40 60min $42.5 \pm 2.5^{\circ}\text{C}$. The product is filtered and washed with water (2x3vol), sucked

dry then transferred to a vacuum oven and further dried at $52.5 \pm 2.5^\circ\text{C}$ under vacuum (NLT 650 mm of Hg) for $8 \pm 2\text{h}$.

Process Description for Scheme 8


5

Stage a): Preparation of 1-(1,1-Dimethylethyl)-1H-pyrazole (Intermediate 9)




To a mixture of 1,1,3,3-tetramethoxypropane (3.82kg, 23.27mol) and tert-butylhydrazine hydrochloride (2.9kg, 23.27mol) in ethanol (24.54kg), conc HCl (4.72kg, 46.55mol) was added, keeping the temperature below 50°C . The reaction mixture was then rapidly heated to reflux. After ca. 2h the reaction was sampled and analysed by NMR. Pass criteria was $<3.0\%$ starting material remaining. On receipt of a pass result the solution is cooled, diluted with water (8.29kg) and evaporated *in vacuo* ($T < 50^\circ\text{C}$, $p < -0.08\text{MPa}$) until approximately all of the original ethanol was removed. The solution was basified with 10M NaOH(aq), extracted with EtOAc (11.11kgx2) and the organic phase washed with saturated ammonium chloride solution (4.3ml/g x 2) and brine (4.3ml/g), then evaporated to give the title compound (2.08kg, 72%yield) as a brown liquid (GC purity 99.70%a/a).

20 Stage b): Preparation of 4-Bromo-1-(1,1-dimethylethyl)-1H-pyrazole (Intermediate 10)




To an ice cooled solution (0°C to 10°C) of 1-(tert-butyl)pyrazole (1.75kg, 14.09mol) in dichloromethane (12.9kg) was added NBS (2.63kg, 14.79mol) portionwise. The solution was stirred at 0°C until the content of 1-(tert-butyl)pyrazole $<30\%$ (GC), then warmed to RT and stirred until the sample taken shows $<1.0\%$ by GC. On receipt of a pass result, 10% sodium bisulfite aqueous was added to the reaction mixture until KI-starch did not turn to blue. The organic phase was then washed with 5% NaCl solution and brine in sequence, then evaporated to give the title compound as a brown liquid (2.67kg, 93.4%yield; GC purity 99.1%a/a).

Stage c): Preparation of iso-Propyl pinacol borate



To a 20L four-neck bottle tri-isopropyl borate (261.0g, 1.388mol) and pinacol (142.5g, 1.207mol) were added and heated to ~90°C for 12-16h. The criteria for a reaction completion is pinacol <4.0% by GC. After completion, the reaction was converted to distillation, and the product fractions (boiling at 174-178°C) collected. Thus the title product was obtained as a colourless oil in a yield range of 80-90%th across the six 20L batches which were operated. GC records purity of 87-96%, but ¹H NMR shows the product to be very pure. [discrepancy due to product instability to GC conditions.]


Stage d): Preparation of 1-(1,1-Dimethylethyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (Intermediate 11)



To a 20L four-neck bottle, 4-bromo-(tert-butyl)pyrazole (1.15kg, 5.66mol) and THF (9.2L) were added, then the mixture was cooled to between -78°C and -85°C and treated with nBuLi (6.23mol) dropwise at that temperature. After addition the solution was stirred at the same temperature for 1h and isopropyl pinacol borate (1.47kg, 7.92mol) was added dropwise. The reaction was complete after stirring for ~3h (starting material <1.0%a/a by GC) then water (2.3L) was added to quench the reaction; and the pH adjusted to 8-9 by addition of 3.45kg 1M HCl. The aqueous phase was extracted with TBME (3.45L x 2), and the combined organic phase washed with 5% NaCl (3.45L x 2) and water (3.45L) in sequence. The organic phase was evaporated to give the crude product. After subsequent recrystallization from heptanes the pure product was obtained as a white solid (GC purity 99.7%a/a) in a 63.5%th overall yield. (Two 20L reactions were run, then combined in the heptane recrystallization).

Claims

1. A compound of formula (I):



wherein:

X is O or NH;

R¹ is C₂₋₄ alkyl, C₃₋₇cycloalkyl, hydroxyC₂₋₄ alkyl, C₁₋₂ alkoxyC₁₋₄ alkyl, trifluoromethylC₁₋₂ alkyl or benzyl; or

a salt thereof.

2. A compound or a salt thereof according to claim 1 in which X is NH.

3. A compound or a salt thereof according to claim 1 or 2 in which R¹ is C₂₋₄ alkyl, C₃₋₇cycloalkyl or C₁₋₂alkoxyC₁₋₄ alkyl.

4. A compound or a salt thereof according to claim 1 in which R¹ is ethyl, *t*-butyl, -CH₂OCH₃, -CH₂C(CH₃)₂OH, -CH₂CH₂OH, -CH₂CH₂OCH₃, -CH(CH₃)₂, -CH₂CF₃, benzyl or cyclopentyl.

5. A compound or a salt thereof according to claim 4 in which R¹ is *t*-butyl.

6. A compound or a salt thereof according to one of claims 1 to 5 having the *S* absolute stereochemistry.

7. A compound or a salt thereof selected from the group consisting of:

7-(1-ethyl-1*H*-pyrazol-4-yl)-*N*-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine;

7-[1-(1,1-dimethylethyl)-1*H*-pyrazol-4-yl]-*N*-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine;

N-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-7-[1-[(methyloxy)methyl]-1*H*-pyrazol-4-yl]-1,6-naphthyridin-5-amine;

7-(1-ethyl-1*H*-pyrazol-4-yl)-5-({[(3*S*)-3-fluoro-3-piperidinyl]methyl}oxy)-1,6-naphthyridine;

1-{4-[5-({[(3*S*)-3-fluoro-3-piperidinyl]methyl}amino)-1,6-naphthyridin-7-yl]-1*H*-pyrazol-1-yl}-2-methyl-2-propanol;

2-{4-[5-({[(3S)-3-fluoro-3-piperidinyl]methyl}amino)-1,6-naphthyridin-7-yl]-1H-pyrazol-1-yl}ethanol;

N-{[(3S)-3-fluoro-3-piperidinyl]methyl}-7-[1-[2-(methyloxy)ethyl]-1H-pyrazol-4-yl]-1,6-naphthyridin-5-amine;

5 7-(1-cyclopentyl-1H-pyrazol-4-yl)-*N*-[(3-fluoro-3-piperidinyl)methyl]-1,6-naphthyridin-5-amine;

N-[(3-fluoro-3-piperidinyl)methyl]-7-[1-(2,2,2-trifluoroethyl)-1H-pyrazol-4-yl]-1,6-naphthyridin-5-amine;

10 *N*-{[(3S)-3-fluoro-3-piperidinyl]methyl}-7-[1-(2,2,2-trifluoroethyl)-1H-pyrazol-4-yl]-1,6-naphthyridin-5-amine;

N-{[(3R)-3-fluoro-3-piperidinyl]methyl}-7-[1-(2,2,2-trifluoroethyl)-1H-pyrazol-4-yl]-1,6-naphthyridin-5-amine;

N-[(3-fluoro-3-piperidinyl)methyl]-7-[1-(phenylmethyl)-1H-pyrazol-4-yl]-1,6-naphthyridin-5-amine;

15 *N*-[(3-fluoro-3-piperidinyl)methyl]-7-[1-(1-methylethyl)-1H-pyrazol-4-yl]-1,6-naphthyridin-5-amine;

N-{[(3S)-3-fluoro-3-piperidinyl]methyl}-7-[1-(1-methylethyl)-1H-pyrazol-4-yl]-1,6-naphthyridin-5-amine; and


20 *N*-{[(3R)-3-fluoro-3-piperidinyl]methyl}-7-[1-(1-methylethyl)-1H-pyrazol-4-yl]-1,6-naphthyridin-5-amine; and
salts thereof.

8. A compound or a salt thereof selected from the group consisting of:

25 7-(1-ethyl-1H-pyrazol-4-yl)-*N*-{[(3S)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine; and


7-[1-(1,1-dimethylethyl)-1H-pyrazol-4-yl]-*N*-{[(3S)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine; and
salts thereof.

30 9. A compound which is 7-[1-(1,1-dimethylethyl)-1H-pyrazol-4-yl]-*N*-{[(3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine



or a salt thereof.

35 10. A compound which is 7-[1-(1,1-dimethylethyl)-1H-pyrazol-4-yl]-*N*-{[(3S)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine



or a salt thereof.

11. A compound or a salt thereof according to any one of claims 1 to 10,
5 wherein the salt is a pharmaceutically acceptable salt.
12. A compound or a salt thereof according to claim 11, wherein the
pharmaceutically acceptable salt is the hydrochloride salt.
- 10 13. A pharmaceutical composition comprising a compound or a
pharmaceutically acceptable salt thereof, as defined in claim 11 or claim 12, and one
or more pharmaceutically acceptable excipients.
14. A pharmaceutical composition comprising 7-[1-(1,1-dimethylethyl)-1H-
15 pyrazol-4-yl]-N-[(3S)-3-fluoro-3-piperidiny]methyl]-1,6-naphthyridin-5-amine or a
pharmaceutically acceptable salt thereof and one or more pharmaceutically
acceptable excipients.
15. A compound or a pharmaceutically acceptable salt thereof, as defined in
20 claim 11 or claim 12, for use in therapy.
16. A compound or a pharmaceutically acceptable salt thereof, as defined in
claim 11 or claim 12, for use in inhibiting spleen tyrosine kinase (Syk).
- 25 17. A compound or a pharmaceutically acceptable salt thereof as defined in
claim 11 or claim 12, for use in the treatment of an autoimmune condition.
18. A compound or a pharmaceutically acceptable salt thereof for use according
to claim 17, wherein the autoimmune condition is selected from systemic lupus
30 erythematosus (SLE), discoid (cutaneous) lupus, Sjorgens syndrome, Wegners
granulomatosis and other vasculitides, bullous pemphigoid and pemphigus,
idiopathic thrombocytopenic purpura (ITP), giant cell arteriosis, glomerulonephritis,
chronic transplant rejection and rheumatoid arthritis.
- 35 19. A compound or a pharmaceutically acceptable salt thereof for use
according to claim 18, wherein the autoimmune condition is rheumatoid arthritis.

20. A compound which is 7-[1-(1,1-dimethylethyl)-1*H*-pyrazol-4-yl]-*N*-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine or a pharmaceutically acceptable salt thereof, for use in the treatment of rheumatoid arthritis.
- 5 21. A compound or a pharmaceutically acceptable salt thereof as defined in claim 11 or claim 12, for use in the treatment of chronic spontaneous urticaria.
22. A compound which is 7-[1-(1,1-dimethylethyl)-1*H*-pyrazol-4-yl]-*N*-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine or a pharmaceutically
10 acceptable salt thereof, for use in the treatment of chronic spontaneous urticaria.
23. A compound or a pharmaceutically acceptable salt thereof as defined in claim 11 or claim 12, for use in the treatment of cancer.
- 15 24. A compound or a pharmaceutically acceptable salt thereof for use according to claim 23, wherein the cancer is selected from heme malignancies, particularly Non-Hodgkin's Lymphomas including follicular (FL), mantle cell, small lymphocytic lymphoma/chronic lymphocytic lymphoma (SLL/CLL), Burkitt and diffuse large B cell (DLBCL) lymphomas.
- 20 25. A compound or a pharmaceutically acceptable salt thereof as defined in claim 11 or claim 12, for use in the treatment of a disease associated with inappropriate mast and/or basophil cell activation.
- 25 26. A compound or a pharmaceutically acceptable salt thereof as defined in claim 11 or claim 12, for use in the treatment of an inflammatory disease and/or allergic disorder.
27. A compound or a pharmaceutically acceptable salt thereof for use according
30 to claim 26, wherein the inflammatory disease and/or allergic disorder is selected from chronic obstructive pulmonary disease (COPD), adult respiratory distress syndrome (ARDS), asthma, severe asthma, ulcerative colitis, Crohn's disease, bronchitis, conjunctivitis, psoriasis, scleroderma, dermatitis, allergy, rhinitis, cutaneous lupus, autoimmune bullous conditions including pemphigus and
35 pemphigoid, mastocytosis and anaphylaxis.
28. The use of a compound or a pharmaceutically acceptable salt thereof as defined in claim 11 or claim 12, for the manufacture of a medicament for the treatment of an autoimmune condition.
- 40 29. The use of a compound or a pharmaceutically acceptable salt thereof as defined in claim 11 or claim 12, for the manufacture of a medicament for the treatment of cancer.

30. The use of a compound or a pharmaceutically acceptable salt thereof as defined in claim 11 or claim 12, for the manufacture of a medicament for the treatment of an inflammatory disease and/or allergic disorder.

5 31. A method of inhibiting spleen tyrosine kinase (syk), which comprises administering to a subject in need thereof a therapeutically effective amount of compound or a pharmaceutically acceptable salt thereof, as defined in claim 11 or claim 12.

10 32. A method of treating an autoimmune condition, which comprises administering to a subject in need thereof a therapeutically effective amount of compound or a pharmaceutically acceptable salt thereof, as defined in claim 11 or claim 12.

15 33. A method of treating an autoimmune condition according to claim 32, wherein the autoimmune condition is selected from systemic lupus erythematosus (SLE), discoid (cutaneous) lupus, Sjorgens syndrome, Wegners granulomatosis and other vasculitides, bullous pemphigoid and pemphigus, idiopathic thrombocytopenic purpura (ITP), giant cell arteriosis, glomerulonephritis, chronic transplant rejection
20 and rheumatoid arthritis.

34. A method of treatment according to claim 33, wherein the autoimmune condition is rheumatoid arthritis.

25 35. A method of treating rheumatoid arthritis, which comprises administering to a subject in need thereof a therapeutically effective amount of compound which is 7-[1-(1,1-dimethylethyl)-1*H*-pyrazol-4-yl]-*N*-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine or a pharmaceutically acceptable salt thereof.

30 36. A method of treating chronic spontaneous urticaria, which comprises administering to a subject in need thereof a therapeutically effective amount of compound or a pharmaceutically acceptable salt thereof, as defined in claim 11 or claim 12.

35 37. A method of treating chronic spontaneous urticaria, which comprises administering to a subject in need thereof a therapeutically effective amount of compound which is 7-[1-(1,1-dimethylethyl)-1*H*-pyrazol-4-yl]-*N*-{[(3*S*)-3-fluoro-3-piperidinyl]methyl}-1,6-naphthyridin-5-amine or a pharmaceutically acceptable salt thereof.

40 38. A method of treating cancer which comprises administering to a patient in need thereof a therapeutically effective amount of compound or a pharmaceutically acceptable salt thereof, as defined in claim 11 or claim 12.

39. A method of treating cancer according to claim 38, wherein the cancer is selected from heme malignancies, particularly Non-Hodgkin's Lymphomas including follicular (FL), mantle cell, small lymphocytic lymphoma/chronic lymphocytic lymphoma (SLL/CLL), Burkitt and diffuse large B cell (DLBCL) lymphomas.

5

40. A method of treating a disease associated with inappropriate mast and/or basophil cell activation, which comprises administering to a subject in need thereof a therapeutically effective amount of compound or a pharmaceutically acceptable salt thereof, as defined in claim 11 or claim 12.

10

41. A method of treating an inflammatory disease and/or allergic disorder, which comprises administering to a subject in need thereof a therapeutically effective amount of a compound or a pharmaceutically acceptable salt thereof, as defined in claim 11 or claim 12.

15

42. A method of treating an inflammatory disease and/or allergic disorder according to claim 41, wherein the inflammatory and/or allergic disorder is selected from chronic obstructive pulmonary disease (COPD), adult respiratory distress syndrome (ARDS), asthma, severe asthma, ulcerative colitis, Crohn's disease, bronchitis, conjunctivitis, psoriasis, scleroderma, dermatitis, allergy, rhinitis, cutaneous lupus, autoimmune bullous conditions including pemphigus and pemphigoid, mastocytosis and anaphylaxis.

20

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2011/056600

A. CLASSIFICATION OF SUBJECT MATTER INV. C07D471/04 A61K31/4462 A61P19/02 ADD.		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) C07D A61K A61P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, BEILSTEIN Data, BIOSIS, CHEM ABS Data, WPI Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2010/015520 A1 (BOEHRINGER INGELHEIM INT [DE]; BOUYSSOU THIERRY [DE]; DAHMANN GEORG [D] 11 February 2010 (2010-02-11) cited in the application claims; compound 70 -----	1-42
A	WO 2010/015518 A2 (BOEHRINGER INGELHEIM INT [DE]; FIEGEN DENNIS [DE]; HANDSCHUH SANDRA [D] 11 February 2010 (2010-02-11) cited in the application claims; compounds 14, 15 -----	1-42
☐ Further documents are listed in the continuation of Box C. ☒ See patent family annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search	Date of mailing of the international search report	
4 August 2011	16/08/2011	
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Stroeter, Thomas	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2011/056600

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2010015520 A1	11-02-2010	AR 072913 A1	29-09-2010
		CA 2732087 A1	11-02-2010
		EP 2324022 A1	25-05-2011

WO 2010015518 A2	11-02-2010	CA 2731926 A1	11-02-2010
		EP 2313406 A2	27-04-2011
