



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2020/04/24
(87) Date publication PCT/PCT Publication Date: 2020/10/29
(85) Entrée phase nationale/National Entry: 2021/10/20
(86) N° demande PCT/PCT Application No.: US 2020/029946
(87) N° publication PCT/PCT Publication No.: 2020/219972
(30) Priorité/Priority: 2019/04/25 (US62/838,770)

(51) Cl.Int./Int.Cl. *C12N 9/10* (2006.01),
C12N 15/63 (2006.01), *C12P 21/02* (2006.01)
(71) Demandeur/Applicant:
IMPOSSIBLE FOODS INC., US
(72) Inventeurs/Inventors:
ROY CHAUDHURI, BISWAJOY, US;
SHANKAR, SMITA, US
(74) Agent: SMART & BIGGAR LLP

(54) Titre : COMPOSITIONS ET PROCEDES POUR LA PRODUCTION DE PROTEINES CONTENANT DE L'HEME
(54) Title: STRAINS AND METHODS FOR PRODUCTION OF HEME-CONTAINING PROTEINS

(57) Abrégé/Abstract:

This document relates to materials and methods for the production of protein. In one aspect, this document provides a cell including a first exogenous nucleic acid construct comprising a nucleotide sequence encoding an aminolevulinate synthase (ALAS) protein operably linked to a first promoter element, wherein the ALAS includes a mutation in a first heme responsive motif (HRM) and a mutation in a second HRM, and a second exogenous nucleic acid construct including a nucleotide sequence encoding a heme-binding protein, wherein the second exogenous nucleic acid construct including a nucleotide sequence encoding the heme-binding protein is operably linked to the first promoter element or is operably linked to a second promoter element.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

29 October 2020 (29.10.2020)



(10) International Publication Number

WO 2020/219972 A1

(51) International Patent Classification:

C12N 9/10 (2006.01)

C12P 21/02 (2006.01)

C12N 15/63 (2006.01)

Published:

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))

(21) International Application Number:

PCT/US2020/029946

(22) International Filing Date:

24 April 2020 (24.04.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/838,770

25 April 2019 (25.04.2019)

US

(71) Applicant: **IMPOSSIBLE FOODS INC.** [US/US]; 400
Saginaw Drive, Redwood City, California 94063 (US).

(72) Inventors; and

(71) Applicants: **ROY-CHAUDHURI, Biswajoy** [US/US];
2831 Flores Street, San Mateo, California 94403 (US).
SHANKAR, Smita [IN/US]; 120 S. El Camino Real, Apt
115, Millbrae, California 94030 (US).

(74) Agent: **MAIZE, Kimberly M.** et al.; P.O. Box 1022, Min-
neapolis, Minnesota 55440-1022 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,
KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

(54) Title: STRAINS AND METHODS FOR PRODUCTION OF HEME-CONTAINING PROTEINS

(57) Abstract: This document relates to materials and methods for the production of protein. In one aspect, this document provides a cell including a first exogenous nucleic acid construct comprising a nucleotide sequence encoding an aminolevulinate synthase (ALAS) protein operably linked to a first promoter element, wherein the ALAS includes a mutation in a first heme responsive motif (HRM) and a mutation in a second HRM, and a second exogenous nucleic acid construct including a nucleotide sequence encoding a heme-binding protein, wherein the second exogenous nucleic acid construct including a nucleotide sequence encoding the heme-binding protein is operably linked to the first promoter element or is operably linked to a second promoter element.



WO 2020/219972 A1

STRAINS AND METHODS FOR PRODUCTION OF HEME-CONTAINING PROTEINS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. Provisional Application Serial No. 62/838,770, filed on April 25, 2019, which is incorporated by reference herein in its entirety.

5

DESCRIPTION OF THE TEXT FILE SUBMITTED ELECTRONICALLY

The contents of the text file submitted electronically herewith are incorporated herein by reference in their entirety: A computer readable format copy of the Sequence Listing filename: 38767-0158WO1.txt, date recorded, April 24, 2020, file size $\approx$ 69 kilobytes.

10

TECHNICAL FIELD

This disclosure generally relates to DNA constructs and methods of using such DNA constructs to genetically engineer cells (e.g., yeast cells (e.g., methylotrophic yeast cells)).

BACKGROUND

15

Cells such as *Pichia pastoris* are commonly used for expression of recombinant proteins. Constructs that can be used to efficiently express one or more proteins in a cell (e.g., fungal cell, such as an *Aspergillus* cell, a *Trichoderma* cell, or a yeast cell (e.g., a methylotrophic yeast cell)) are provided herein.

20

SUMMARY

25

This document is based, at least in part, on the identification of mutations in aminolevulinic acid synthase (ALAS) that can confer increased expression of heme-binding proteins. The mutated ALAS proteins described herein can be used for the efficient expression of heme-binding proteins in *Pichia*, for example.

In one aspect, provided herein is a methylotrophic yeast cell including a first exogenous nucleic acid construct comprising a nucleotide sequence encoding an aminolevulinate synthase (ALAS) protein operably linked to a first promoter element, wherein the ALAS includes a first mutation in a first heme responsive motif (HRM) and a
 5 second exogenous nucleic acid construct comprising a nucleotide sequence encoding a heme-binding protein, wherein the second exogenous nucleic acid construct comprising a nucleotide sequence encoding the heme-binding protein is operably linked to the first promoter element or is operably linked to a second promoter element.

Implementations can have one or more of the following features. The methylotrophic
 10 yeast cell can be a *Pichia* cell, a *Candida* cell, a *Hansenula* cell, or a *Torulopsis* cell. The methylotrophic yeast cell can be a *Pichia methanolica* cell, a *Pichia pastoris* cell, a *Candida boidinii* cell, or a *Hansenula polymorpha* cell. The methylotrophic yeast cell can be a *Pichia pastoris* cell. The first mutation can be a cysteine to serine mutation. The first mutation can be a cysteine to alanine mutation. The ALAS can include a second mutation in a second
 15 HRM. The second mutation can be a cysteine to serine mutation. The second mutation can be a cysteine to alanine mutation. The first exogenous nucleic acid construct can include a nucleic acid sequence having at least 90% sequence identity to the nucleic acid sequence in SEQ ID NO: 28. The first exogenous nucleic acid construct can include a nucleic acid sequence having at least 95% sequence identity to the nucleic acid sequence in SEQ ID NO:
 20 28. The ALAS protein can include an amino acid sequence having at least 90% sequence identity to the amino acid sequence in SEQ ID NO: 29. The ALAS protein can include an amino acid sequence having at least 95% sequence identity to the amino acid sequence in SEQ ID NO: 29. The heme-binding protein can be selected from the group consisting of a globin, a cytochrome, a cytochrome c oxidase, a ligninase, a catalase, and a peroxidase. The
 25 heme-binding protein can be selected from the group consisting of an androglobin, a chlorocruorin, a cytoglobin, an erythrocruorin, a flavohemoglobin, a globin E, a globin X, a globin Y, a hemoglobin, a histoglobin, a leghemoglobin, a myoglobin, a neuroglobin, a non-symbiotic hemoglobin, a protoglobin, and a truncated hemoglobin. The heme-binding protein can be a non-symbiotic hemoglobin. The heme-binding protein can be a leghemoglobin. The
 30 heme-binding protein can include an amino acid sequence having at least 90% sequence identity to an amino acid sequence in any one of SEQ ID NOs: 1-27. The methylotrophic yeast cell can further include a third nucleic acid construct comprising a nucleotide sequence

encoding a transcription factor, wherein the third nucleic acid construct is operably linked to the first promoter element, the second promoter element, or a third promoter element. The first promoter element can include a recognition sequence for the transcription factor. The second exogenous nucleic acid construct can be operably linked to a second promoter
5 element, and wherein the second promoter element includes a recognition sequence for the transcription factor. The third nucleic acid construct can be operably linked to the third promoter element, and wherein the third promoter element includes a recognition sequence for the transcription factor. The methylotrophic yeast cell can further include a fourth nucleic acid construct comprising a nucleotide sequence encoding a protein involved in heme
10 biosynthesis, wherein the fourth nucleic acid construct is operably linked to the first promoter element, the second promoter element, the third promoter element, or a fourth promoter element. The protein involved in heme biosynthesis can be selected from the group consisting of ALA dehydratase, porphobilinogen deaminase, UPG III synthase, UPG III decarboxylase, CPG oxidase, PPG oxidase, and ferrochelatase. The first exogenous nucleic acid construct
15 can be a heterologous nucleic acid construct. The second exogenous nucleic acid construct can be a heterologous nucleic acid construct. The heme-binding protein can be an exogenous heme-binding protein. The heme-binding protein can be a heterologous heme-binding protein.

In another aspect, provided herein is a method of producing a heme-binding protein in
20 a methylotrophic yeast cell including expressing a first exogenous nucleic acid construct comprising a nucleotide sequence encoding an aminolevulinate synthase (ALAS) protein operably linked to a first promoter element, wherein the ALAS includes a first mutation in a first heme responsive motif (HRM) and expressing a second exogenous nucleic acid construct comprising a nucleotide sequence encoding a heme-binding protein,
25 wherein the second exogenous nucleic acid construct comprising a nucleotide sequence encoding the heme-binding protein is operably linked to the first promoter element or is operably linked to a second promoter element.

Implementations can have one or more of the following features. The ALAS can include a second mutation in a second HRM. The method can produce the heme-binding
30 protein in a titer that is at least 5% greater than a corresponding method lacking the first exogenous nucleic acid construct. The method can produce the heme-binding protein in a titer that is at least 10% greater than a corresponding method lacking the first exogenous nucleic

acid construct. The method can produce the heme-binding protein in a titer that is at least 15% greater than a corresponding method lacking the first exogenous nucleic acid construct. The method can produce the heme-binding protein in a titer that is at least 20% greater than a corresponding method lacking the first exogenous nucleic acid construct. The method can produce the heme-binding protein in a titer that is at least 5% greater than a corresponding method lacking the first mutation. The method can produce the heme-binding protein in a titer that is at least 10% greater than a corresponding method lacking the first mutation. The method can produce the heme-binding protein in a titer that is at least 15% greater than a corresponding method lacking the first mutation. The method can produce the heme-binding protein in a titer that is at least 20% greater than a corresponding method lacking the first mutation. The method can produce the heme-binding protein in a titer that is at least 5% greater than a corresponding method lacking the first mutation and the second mutation. The method can produce the heme-binding protein in a titer that is at least 10% greater than a corresponding method lacking the first mutation and the second mutation. The method can produce the heme-binding protein in a titer that is at least 15% greater than a corresponding method lacking the first mutation and the second mutation. The method can produce the heme-binding protein in a titer that is at least 20% greater than a corresponding method lacking the first mutation and the second mutation. The method can be carried out in the absence of added methanol. The methylotrophic yeast cell can be a *Pichia* cell, a *Candida* cell, a *Hansenula* cell, or a *Torulopsis* cell. The methylotrophic yeast cell can be a *Pichia methanolica* cell, a *Pichia pastoris* cell, a *Candida boidinii* cell, or a *Hansenula polymorpha* cell. The methylotrophic yeast cell can be a *Pichia pastoris* cell. The first mutation can be a cysteine to serine mutation. The first mutation can be a cysteine to alanine mutation. The second mutation can be a cysteine to serine mutation. The second mutation can be a cysteine to alanine mutation. The first exogenous nucleic acid construct can include a nucleic acid sequence having at least 90% sequence identity to the nucleic acid sequence in SEQ ID NO: 28. The first exogenous nucleic acid construct can include a nucleic acid sequence having at least 95% sequence identity to the nucleic acid sequence in SEQ ID NO: 28. The ALAS protein can include an amino acid sequence having at least 90% sequence identity to the amino acid sequence in SEQ ID NO: 29. The ALAS protein can include an amino acid sequence having at least 95% sequence identity to the amino acid sequence in SEQ ID NO: 29. The heme-binding protein can be selected from the group consisting of a globin, a

cytochrome, a cytochrome c oxidase, a ligninase, a catalase, and a peroxidase. The heme-binding protein can be selected from the group consisting of an androglobin, a chlorocruorin, a cytoglobin, an erythrocruorin, a flavohemoglobin, a globin E, a globin X, a globin Y, a hemoglobin, a histoglobin, a leghemoglobin, a myoglobin, a neuroglobin, a non-symbiotic hemoglobin, a protoglobin, and a truncated hemoglobin. The heme-binding protein can be a non-symbiotic hemoglobin. The heme-binding protein can be a leghemoglobin. The heterologous heme-binding protein can include an amino acid sequence having at least 90% sequence identity to an amino acid sequence in any one of SEQ ID NOs: 1-27. The method can further include expressing a third nucleic acid construct comprising a nucleotide sequence encoding a transcription factor, wherein the third nucleic acid construct is operably linked to the first promoter element, the second promoter element, or a third promoter element. The first promoter element can include a recognition sequence for the transcription factor. The second exogenous nucleic acid construct can be operably linked to a second promoter element, and wherein the second promoter element includes a recognition sequence for the transcription factor. The third nucleic acid construct can be operably linked to the third promoter element, and wherein the third promoter element includes a recognition sequence for the transcription factor. The method can further include expressing a fourth nucleic acid construct comprising a nucleotide sequence encoding a protein involved in heme biosynthesis, wherein the fourth nucleic acid construct is operably linked to the first promoter element, the second promoter element, the third promoter element, or a fourth promoter element. The protein involved in heme biosynthesis can be selected from the group consisting of ALA dehydratase, porphobilinogen deaminase, UPG III synthase, UPG III decarboxylase, CPG oxidase, PPG oxidase, and ferrochelatase. The first exogenous nucleic acid construct can be a heterologous nucleic acid construct. The second exogenous nucleic acid construct can be a heterologous nucleic acid construct. The heme-binding protein can be an exogenous heme-binding protein. The heme-binding protein can be a heterologous heme-binding protein.

In another aspect, provided herein is a *Pichia pastoris* cell including a first exogenous nucleic acid construct comprising a nucleotide sequence encoding an aminolevulinate synthase (ALAS) protein operably linked to a first promoter element, wherein the ALAS includes a first mutation in a first heme responsive motif (HRM) and a second mutation in a second HRM and a second exogenous nucleic acid construct comprising a nucleotide

sequence encoding leghemoglobin, wherein the second exogenous nucleic acid construct comprising a nucleotide sequence encoding leghemoglobin is operably linked to the first promoter element, or the second exogenous nucleic acid construct comprising a nucleotide sequence encoding leghemoglobin is operably linked to a second promoter element.

5 Implementations can include one or more of the following features. The first mutation can be a cysteine to serine mutation. The second mutation can be a cysteine to serine mutation.

In another aspect, provided herein is a method of producing leghemoglobin, the method including expressing a first exogenous nucleic acid construct comprising a nucleotide
10 sequence encoding an aminolevulinate synthase (ALAS) protein operably linked to a first promoter element, wherein the ALAS includes a first mutation in a first heme responsive motif (HRM) and a second mutation in a second HRM and expressing a second exogenous nucleic acid construct comprising a nucleotide sequence encoding leghemoglobin, wherein
15 the second exogenous nucleic acid construct comprising a nucleotide sequence encoding leghemoglobin is operably linked to the first promoter element, or the second exogenous nucleic acid construct comprising a nucleotide sequence encoding leghemoglobin is operably linked to a second promoter element.

Implementations can have one or more of the following features. The method can produce the leghemoglobin in a titer that is at least 5% greater than a corresponding method
20 lacking the first exogenous nucleic acid construct. The method can produce the leghemoglobin in a titer that is at least 10% greater than a corresponding method lacking the first exogenous nucleic acid construct. The method can produce the leghemoglobin in a titer that is at least 15% greater than a corresponding method lacking the first exogenous nucleic acid construct. The method can produce the leghemoglobin in a titer that is at least 20%
25 greater than a corresponding method lacking the first exogenous nucleic acid construct. The method can produce the leghemoglobin in a titer that is at least 5% greater than a corresponding method lacking the first mutation and the second mutation. The method can produce the leghemoglobin in a titer that is at least 10% greater than a corresponding method lacking the first mutation and the second mutation. The method can produce the
30 leghemoglobin in a titer that is at least 15% greater than a corresponding method lacking the first mutation and the second mutation. The method can produce the leghemoglobin in a titer

that is at least 20% greater than a corresponding method lacking the first mutation and the second mutation. The method can be carried out in the absence of added methanol.

In another aspect, provided herein is a *Pichia pastoris* cell including a first exogenous nucleic acid construct comprising a nucleotide sequence encoding a protein with at least 90% sequence identity to SEQ ID NO: 29, wherein the nucleic acid encodes a serine residue in a position corresponding to position 12 of SEQ ID NO: 29, and wherein the nucleic acid encodes a serine residue in a position corresponding to position 39 of SEQ ID NO: 29, operably linked to a first promoter element.

Implementations can include one or more of the following features. The *Pichia pastoris* cell can further include a second exogenous nucleic acid construct comprising a nucleotide sequence encoding a protein with at least 90% sequence identity to any of SEQ ID NOs: 1-27, wherein the second exogenous nucleic acid construct is operably linked to the first promoter element, or the second exogenous nucleic acid construct is operably linked to a second promoter element. The *Pichia pastoris* cell can further include a second exogenous nucleic acid construct comprising a nucleotide sequence encoding a protein with at least 90% sequence identity to SEQ ID NO: 4, wherein the second exogenous nucleic acid construct is operably linked to the first promoter element, or the second exogenous nucleic acid construct is operably linked to a second promoter element.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. Although methods and materials similar or equivalent to those described herein can be used to practice the invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

The details of one or more embodiments of the disclosure are set forth in the accompanying drawings and the description below. Other features, objects, and advantages of the disclosure will be apparent from the description and drawings, and from the claims. The word “comprising” in the claims may be replaced by “consisting essentially of” or with “consisting of,” according to standard practice in patent law.

DESCRIPTION OF THE DRAWINGS

Figure 1 provides a schematic depicting the steps involved in the heme biosynthesis pathway.

Figure 2 provides a structure of Heme b.

Figure 3 provides a structure of hemin.

5 Figure 4 provides the sequences of exemplary heme-binding proteins (SEQ ID NOs: 1-27).

Figure 5 provides an exemplary construct diagram of a mutated ALAS.

Figure 6 provides the nucleic acid sequence (SEQ ID NO: 28) and amino acid sequence (SEQ ID NO: 29) of wild-type ALAS.

10 Figure 7 provides the nucleic acid sequence (SEQ ID NO: 30) and amino acid sequence (SEQ ID NO: 31) of an exemplary mutant ALAS.

Figure 8 provides the sequences of SEQ ID NOs: 32-66.

DETAILED DESCRIPTION

15 This document is related to materials and methods for protein production. In particular, this document is related to materials and method for the production of heme and heme-binding proteins.

Methylotrophic yeast, such as *Pichia pastoris*, are commonly used to produce recombinant proteins. *Pichia* strains are typically able to grow using methanol as the sole carbon source. It will be understood that *Pichia pastoris* has been reclassified as
20 *Komagataella* species, such as *Komagataella phaffii*, *Komagataella pastoris*, or *Komagataella pseudopastoris*, though the term '*Pichia pastoris*' is still in use and may refer to any appropriate *Komagataella* species. Commonly, laboratory strains of *P. pastoris* are *Komagataella phaffii*.

Proteins that bind heme include cytochromes, catalase, myoglobin and hemoglobin
25 among others. Aminolevulinic acid synthase (ALAS) is an enzyme (EC 2.3.1.37) that catalyzes the first step in heme biosynthesis (See Figure 1 for a diagram of a Heme b biosynthesis pathway), catalyzing the transformation of glycine and succinyl-CoA to aminolevulinic acid. ALAS is translocated to the mitochondria for this step of heme biosynthesis. Aminolevulinic acid is transformed by other enzymes (e.g., ALA dehydratase
30 (ALAD), porphobilinogen deaminase (PBGD), uroporphyrinogen III synthase (UPG3S), uroporphyrinogen III decarboxylase (UPG3D), coprotoporphyrinogen oxidase (COPROX),

protoporphyrinogen IX oxidase (PROTOX), and/or ferrochelatase (FC)) to Heme b (Figure 2). In some cases, the action of ALAS can be rate-limiting for the heme biosynthesis pathway. Other hemes (e.g., Heme o, Heme a, Heme c) can be made through the action of enzymes on Heme b. Hemin (Figure 3) is a complex of protoporphyrin IX (e.g., the protoporphyrin of Heme b) with a ferric (+3 oxidation state) iron and a chloride ligand.

As used herein, “intermediate of the heme biosynthesis pathway” refers to one or more of: delta aminolevulinic acid (d-ALA), porphobilinogen, hydroxymethylbilane, uroporphyrinogen III, coproporphyrinogen III, protoporphyrinogen IX, or protoporphyrin IX. In some embodiments, an intermediate of the heme biosynthesis pathway can be selected from the group consisting of delta aminolevulinic acid (d-ALA), porphobilinogen, hydroxymethylbilane, uroporphyrinogen III, coproporphyrinogen III, protoporphyrinogen IX, and protoporphyrin IX. In some embodiments, an intermediate of the heme biosynthesis pathway can be selected from the group consisting of porphobilinogen, hydroxymethylbilane, uroporphyrinogen III, coproporphyrinogen III, protoporphyrinogen IX, or protoporphyrin IX.

Translocation of some ALAS proteins to the mitochondria can be affected by one or more heme regulatory motifs (HRMs; sometimes also called heme responsive motifs) contained within the ALAS protein. Many HRMs include a C-P motif, and the cysteine is often an axial ligand for heme. In some embodiments, a HRM has a sequence of R/L/N/A/C/S/H/I/G/Q-C-P-L/V/I/F/C-L/M/P/V (SEQ ID NO: 32) (e.g., a HRM can have a sequence of A-C-P-F-V (SEQ ID NO: 33), H-C-P-V-V (SEQ ID NO: 34), I-C-P-F-M (SEQ ID NO: 35), or G-C-P-V-V (SEQ ID NO: 36); see, e.g., Figure 8). Some organisms have 3 HRMs in their ALAS sequence, while other organisms (e.g., a methylotrophic yeast such as *Pichia pastoris*) have two. In general, a HRM can be called HRM1, HRM2, HRM3, and so forth, depending on where the HRM occurs in the protein sequence, e.g., the first-occurring HRM in the protein sequence (read as N-terminus to C-terminus) would be called HRM1. Mutation of the cysteine (e.g., to serine or alanine) in the HRMs can increase the translocation of ALAS to the mitochondria (see, e.g., González-Domínguez, et al., Yeast. 2001 Jan 15;18(1):41-8. (PubMed ID (PMID) 11124700); Munakata et al., J Biochem. 2004 Aug;136(2):233-8. (PMID 15496594); Dailey et al, Biochem J. 2005 Mar 1;386(Pt 2):381-6. (PMID 15482256)), each of which is herein incorporated by reference in its entirety).

Without being bound by any particular theory, it is believed that heme binding to the HRMs of wild type ALAS can inhibit the translocation of ALAS to the mitochondria in a negative

feedback manner. Heme is also believed to be involved in the degradation of some ALAS proteins (see, e.g., Kubota, et al., J Biol Chem. 2016 Sep 23;291(39):20516-29. doi: 10.1074/jbc.M116.719161. Epub 2016 Aug 5. (PMID 27496948), incorporated by reference herein in its entirety).

5 It will be appreciated that a “first HRM” can be an HRM in any part of a protein sequence; a “first HRM” can be, but is not necessarily, the first-occurring HRM (HRM1) in a protein sequence. In some embodiments, a first HRM is HRM1, and a second HRM is HRM2. In some embodiments, a first HRM is HRM1, and a second HRM is HRM3. In some
10 embodiments, a first HRM is HRM2, and a second HRM is HRM1. In some embodiments, a first HRM is HRM2, and a second HRM is HRM3. In some embodiments, a first HRM is HRM3, and a second HRM is HRM1. In some embodiments, a first HRM is HRM3, and a second HRM is HRM2.

In some embodiments, a mutation in a nucleic acid can be an insertion, a deletion, or a substitution. In some embodiments, a mutation in a nucleic acid can be a substitution (e.g., a
15 guanosine to cytosine mutation). In some embodiments, a substitution in a coding sequence (e.g., encoding a protein) can be a silent mutation (e.g., the same amino acid is encoded). In some embodiments, a substitution in a coding sequence can be a nonsynonymous mutation (e.g., a missense mutation or a nonsense mutation). In some embodiments, a substitution in a coding sequence can be a missense mutation (e.g., a different amino acid is encoded). In
20 some embodiments, a substitution in a coding sequence can be nonsense mutation (e.g., a premature stop codon is encoded). In some embodiments, a mutation in a nucleic acid can be a deletion. It will be understood that mutations can be used to alter an endogenous nucleic acid, using, for example, CRISPR, TALEN, and/or Zinc-finger nucleases.

In some embodiments, a mutation in a protein sequence can be an insertion, a
25 deletion, or a substitution. It will be understood that a mutation in a nucleic acid that encodes a protein can cause a mutation in a protein sequence. In some embodiments, a mutation in a protein sequence is a substitution (e.g., a cysteine to serine mutation, a cysteine to alanine mutation, a cysteine to valine mutation, a cysteine to leucine mutation, a cysteine to
30 isoleucine mutation, a cysteine to glycine mutation, a cysteine to phenylalanine mutation, a cysteine to threonine mutation, a cysteine to methionine mutation, a cysteine to tryptophan mutation, a cysteine to tyrosine mutation, a cysteine to asparagine mutation, a cysteine to glutamine mutation, a cysteine to proline mutation, a cysteine to arginine mutation, a cysteine

to histidine mutation, a cysteine to lysine mutation, a cysteine to aspartic acid mutation, or a cysteine to glutamic acid mutation). In some embodiments, a mutation in a protein sequence can be a deletion.

In some embodiments, an ALAS protein as described herein can include a mutation in at least one HRM; for example, a first HRM, a second HRM, a third HRM, a fourth HRM, a fifth HRM, and so forth.

In some embodiments, an ALAS protein as described herein can include a mutation in at least one HRM (e.g., a first HRM). In some embodiments, an ALAS protein as described herein can include a mutation in a single HRM (e.g., a first HRM or a second HRM). In some
10 embodiments, an ALAS protein as described herein can include a mutation in a cysteine of at least one HRM (e.g., a first HRM). In some embodiments, an ALAS protein as described herein can include a mutation in a cysteine in a single HRM (e.g., a first HRM or a second HRM).

In some embodiments, an ALAS protein as described herein can include a mutation in at least two HRMs (e.g., a first HRM and a second HRM). In some embodiments, an ALAS
15 protein as described herein can include a mutation in each of two HRMs (e.g., a first HRM and a second HRM). In some embodiments, the mutation in the first HRM and the mutation in the second HRM can be the same (e.g., both cysteine to serine or alanine mutations). In some embodiments, the mutation in the first HRM and the mutation in the second HRM can
20 be different (e.g., one is a cysteine to serine mutation and one is a cysteine to alanine mutation). In some embodiments, an ALAS protein as described herein can include a mutation in a cysteine of at least two HRMs (e.g., a first HRM and a second HRM). In some embodiments, an ALAS protein as described herein can include a mutation in a cysteine in each of two HRMs (e.g., a first HRM and a second HRM). In some embodiments, the
25 mutation in the first HRM and the mutation in the second HRM can be the same (e.g., both cysteine to serine or alanine mutations). In some embodiments, the mutation in the first HRM and the mutation in the second HRM can be different (e.g., one is a cysteine to serine mutation and one is a cysteine to alanine mutation).

In some embodiments, an ALAS protein as described herein can include a mutation in at least three HRMs (e.g., a first HRM, a second HRM, and a third HRM). In some
30 embodiments, an ALAS protein as described herein can include a mutation in each of three HRMs (e.g., a first HRM, a second HRM, and a third HRM). In some embodiments, an

ALAS protein as described herein can include a mutation in a cysteine at least three HRMs (e.g., a first HRM, a second HRM, and a third HRM). In some embodiments, an ALAS protein as described herein can include a mutation in a cysteine in each of three HRMs (e.g., a first HRM, a second HRM, and a third HRM). In some embodiments, the mutation in each of the first HRM, the second HRM, and the third HRM can be the same (e.g., both cysteine to serine or alanine mutations). In some embodiments, the mutation in one or two of the three HRMs can be different.

In some embodiments, a substitution (a mutation to a different amino acid) can be a substitution for an amino acid (e.g., a cysteine) in a HRM. In some embodiments, a mutation in a first HRM can be a substitution for an amino acid (e.g., a cysteine) in the first HRM. In some embodiments, a mutation in a second HRM can be a substitution for an amino acid (e.g., a cysteine) in the second HRM. In some embodiments, a mutation in a third HRM can be a substitution for an amino acid (e.g., a cysteine) in the third HRM. A substitution can be any appropriate substitution. In some embodiments, a different amino acid can be selected from group consisting of arginine, histidine, lysine, serine, threonine, asparagine, glutamine, selenocysteine, glycine, proline, alanine, isoleucine, leucine, methionine, phenylalanine, tryptophan, tyrosine, and valine. In some embodiments, a different amino acid can be selected from group consisting of arginine, histidine, lysine, serine, threonine, asparagine, glutamine, selenocysteine, glycine, proline, alanine, isoleucine, leucine, methionine, aspartic acid, glutamic acid, phenylalanine, tryptophan, tyrosine, and valine. In some embodiments, a different amino acid is selected from a nonpolar aliphatic amino acid (e.g., glycine, proline, alanine, isoleucine, leucine, methionine, or valine), an aromatic amino acid (e.g., phenylalanine, tryptophan, or tyrosine), a polar uncharged amino acid (e.g., serine, threonine, asparagine, or glutamine), or a positively charged amino acid (arginine, histidine, or lysine). In some embodiments, the different amino acid is serine. In some embodiments, the different amino acid is alanine. In some embodiments, the different amino acid is phenylalanine. In some embodiments, the different amino acid is aspartic acid. In some embodiments, the different amino acid is histidine.

Surprisingly, when an ALAS protein with a cysteine to serine mutation in each of its two HRMs is co-expressed with an exogenous heme-binding protein, the titer of the exogenous heme-binding protein can increase significantly.

Generally, a “titer” is the measurement of the amount of a substance in solution. As used herein, the “titer” of a heme-binding protein refers to the overall amount of the polypeptide, whether or not it is bound to heme, unless otherwise specified. The titer of a protein can be measured by suitable method, such as high-performance liquid chromatography (HPLC or UPLC), liquid chromatography-mass spectrometry (LC-MS), an enzyme-linked immunosorbent assay (ELISA), enzyme activity measurements, iron measurement techniques such as atomic absorption spectroscopy, LC-MS, or ultraviolet and/or visible light spectroscopy.

Mutations in ALAS, such as those described herein, can be used to increase heme production. In some embodiments, the titer of a heme-binding protein can be increased. In some embodiments, the titer of a leghemoglobin (LegH) can be increased. The materials and methods of this disclosure can be useful to increase the production of heme.

The materials and methods described herein can have applications in many industries. For example, heme proteins can be used in food. As another example, heme proteins (e.g., hemoglobins, cytochrome P450s) can be produced for research purposes (e.g., to study drug metabolism). Heme proteins can also be used in industry (e.g., catalases, laccases, and/or peroxidases for uses such as washing detergents, pulp bleaching, lignin degradation). Another potential application for heme proteins is in biocatalysis (e.g. cytochrome P450s, lipoxygenases, and/or laccases can be used for the production of pharmaceutical and commodity chemicals). Some heme proteins can be used as therapeutic agents (e.g., nitric oxide synthases or as part of a blood replacement). In some cases, heme proteins can be used for electronic purposes (e.g. for the production of renewable electricity). As another example, heme therapy can be a treatment option for patients with acute porphyrias, a group of eight genetic diseases that result from inability to produce heme.

Nucleic acid constructs (sometimes also called nucleic acid molecules) are provided herein that allow for genetically engineering a cell (e.g., fungal cell, such as an *Aspergillus* cell, a *Trichoderma* cell, or a yeast cell (e.g., a methylotrophic yeast cell)) to produce a mutant ALAS. In addition, nucleic acid constructs are provided herein that allow for genetically engineering a cell (e.g., fungal cell, such as an *Aspergillus* cell, a *Trichoderma* cell, or a yeast cell (e.g., a methylotrophic yeast cell)) to increase the expression of a heme-binding protein. In some embodiments, nucleic acid constructs are provided herein that allow for genetically engineering a cell (e.g., fungal cell, such as an *Aspergillus* cell, a *Trichoderma*

cell, or a yeast cell (e.g., a methylotrophic yeast cell)) to increase the expression of a heme-binding protein from an inducible promoter in the absence of the inducing molecule.

Nucleic acid constructs are provided herein that allow for genetically engineering a cell. A cell can be any appropriate cell. For example, a cell can be a bacterial cell (e.g., an *E. coli* cell, a *B. subtilis* cell, or a *Lactococcus lactis* cell), a fungal cell, an algal cell, a plant cell, an insect cell, or a mammalian cell. In some embodiments, a cell can be a fungal cell. In some embodiments, a cell can be a filamentous fungus cell. In some embodiments, a cell can be an *Aspergillus* or *Trichoderma* cell. In some embodiments, a cell can be a yeast cell. Non-limiting examples of yeast cells include *Pichia* (e.g., *Pichia methanolica*, *Pichia pastoris*), *Candida* (e.g., *Candida boidinii*) cells, *Hansenula* (e.g., *Hansenula polymorpha*) cells, *Torulopsis* cells, and *Sacharomyces* (e.g., *Sacharomyces cerevisiae*) cells. In some embodiments, a cell can be a methylotrophic yeast cell. Non-limiting examples of methylotrophic yeast cells include *Pichia* cells, *Candida* cells, *Hansenula* cells, and *Torulopsis* cells. In some embodiments, a cell can be a *Pichia* cell or a *Sacharomyces* cell. In some embodiments, a cell is a eukaryotic cell. In some embodiments, a cell is a fungal cell. In some embodiments, a cell is a plant cell. In some embodiments, a cell is an algal cell. In some embodiments, a cell is a yeast cell. In some embodiments, a cell is a *Saccharomyces cerevisiae* cell. In some embodiments, a cell is a methylotrophic yeast cell. While the methods are exemplified herein using a *Pichia* species (e.g., *P. pastoris*), other cells can be used, such as other species of the *Pichia* genus or species from any of the *Candida*, *Hansenula*, *Pichia*, and *Torulopsis* genera. Non-limiting examples of species of methylotrophic yeast include *Pichia methanolica*, *Pichia pastoris*, *Candida boidinii*, and *Hansenula polymorpha*.

Accordingly, in one aspect, this document provides materials and methods for expressing protein. In some embodiments, this document provides a cell (e.g., fungal cell, such as an *Aspergillus* cell, a *Trichoderma* cell, or a yeast cell (e.g., a methylotrophic yeast cell)) including a first nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein. In some embodiments, the first nucleic acid encodes an ALAS protein operably linked to a first promoter element. In some embodiments, an ALAS protein includes at least 1 HRM (e.g., 1, 2, 3, or more HRMs). In some embodiments, an ALAS protein includes at least 2 HRMs (e.g., 2, 3 or more HRMs). In some embodiments, an ALAS protein includes at least 3 HRMs (e.g., 3 or more HRMs). In some embodiments, the ALAS protein

includes a mutation in a first HRM (sometimes also called “a first mutation in a first HRM”). In some embodiments, the ALAS protein includes a mutation in a second HRM (sometimes also called “a second mutation in a second HRM”). In some embodiments, the ALAS protein includes a mutation in a first HRM and a mutation in a second HRM. In some embodiments, the first HRM is HRM1. In some embodiments, the second HRM is HRM2. In some
5 embodiments, the ALAS protein includes a mutation in a third HRM (sometimes also called “a third mutation in a third HRM”). In some embodiments, the ALAS protein includes a mutation in a first HRM, a mutation in a second HRM, and a mutation in a third HRM.

10 In some embodiments, the mutation in the first HRM is a substitution. In some embodiments, the mutation in the first HRM is a cysteine substitution. In some embodiments, the mutation in the first HRM is a cysteine to serine mutation. In some embodiments, the mutation in the first HRM is a cysteine to alanine mutation. In some embodiments, the mutation in the first HRM is a cysteine to phenylalanine mutation. In some embodiments, the mutation in the first HRM is a cysteine to aspartic acid mutation. In some embodiments, the
15 mutation in the first HRM is a cysteine to histidine mutation.

20 In some embodiments, the mutation in the second HRM is a substitution. In some embodiments, the mutation in the second HRM is a cysteine substitution. In some embodiments, the mutation in the second HRM is a cysteine to serine mutation. In some embodiments, the mutation in the second HRM is a cysteine to alanine mutation. In some
25 embodiments, the mutation in the second HRM is a cysteine to phenylalanine mutation. In some embodiments, the mutation in the second HRM is a cysteine to aspartic acid mutation. In some embodiments, the mutation in the second HRM is a cysteine to histidine mutation.

30 In some embodiments, the mutation in the third HRM is a substitution. In some embodiments, the mutation in the third HRM is a cysteine substitution. In some embodiments, the mutation in the third HRM is a cysteine to serine mutation. In some
35 embodiments, the mutation in the third HRM is a cysteine to alanine mutation. In some embodiments, the mutation in the third HRM is a cysteine to phenylalanine mutation. In some embodiments, the mutation in the third HRM is a cysteine to aspartic acid mutation. In some embodiments, the mutation in the third HRM is a cysteine to histidine mutation.

40 In some embodiments, the mutation in the first HRM corresponds to a mutation in residue 12 in SEQ ID NO: 29. In some embodiments, the mutation in the second HRM corresponds to a mutation in residue 39 in SEQ ID NO: 29. In some embodiments, the

mutation in the first HRM is a cysteine to serine mutation corresponding to a cysteine to serine mutation in residue 12 in SEQ ID NO: 29. In some embodiments, the mutation in the first HRM is a cysteine to alanine mutation corresponding to a cysteine to alanine mutation in residue 12 in SEQ ID NO: 29. In some embodiments, the mutation in the first HRM is a cysteine to phenylalanine mutation corresponding to a cysteine to phenylalanine mutation in residue 12 in SEQ ID NO: 29. In some embodiments, the mutation in the first HRM is a cysteine to histidine mutation corresponding to a cysteine to histidine mutation in residue 12 in SEQ ID NO: 29. In some embodiments, the mutation in the second HRM is a cysteine to serine mutation corresponding to a cysteine to serine mutation in residue 39 in SEQ ID NO: 29. In some embodiments, the mutation in the second HRM is a cysteine to alanine mutation corresponding to a cysteine to alanine mutation in residue 39 in SEQ ID NO: 29. In some embodiments, the mutation in the second HRM is a cysteine to phenylalanine mutation corresponding to a cysteine to phenylalanine mutation in residue 39 in SEQ ID NO: 29. In some embodiments, the mutation in the second HRM is a cysteine to histidine mutation corresponding to a cysteine to histidine mutation in residue 39 in SEQ ID NO: 29.

In some embodiments, the first nucleic acid includes a mutation corresponding to a guanosine to cytosine mutation at nucleotide position 35 in SEQ ID NO: 28. In some embodiments, the first nucleic acid includes a mutation corresponding to a thymine to guanosine mutation at nucleotide position 34 in SEQ ID NO: 28 and a guanosine to cytosine mutation at nucleotide position 35 in SEQ ID NO: 28. In some embodiments, the first nucleic acid includes a mutation corresponding to a guanosine to cytosine mutation at nucleotide position 116 in SEQ ID NO: 28. In some embodiments, the first nucleic acid includes a mutation corresponding to a thymine to guanosine mutation in nucleotide position 115 in SEQ ID NO: 28 and a guanosine to cytosine mutation at nucleotide position 116 in SEQ ID NO: 28. It will be appreciated that cysteine to serine mutations can be accomplished by mutations at nucleotides other than those specifically disclosed herein. It will be appreciated that cysteine to alanine mutations can be accomplished by mutations at nucleotides other than those specifically disclosed herein. In some embodiments, the first nucleic acid construct includes SEQ ID NO: 30. In some embodiments, the ALAS protein has the sequence of SEQ ID NO: 31. (See, e.g., Figure 7).

As used herein, “operably linked” means that a promoter or other expression element(s) are positioned relative to a nucleic acid coding sequence in such a way as to direct or regulate expression of the coding sequence (e.g., in-frame).

A “corresponding” amino acid position (or substitution) in a protein sequence
5 different from a reference protein sequence (e.g., in the ALAS protein sequence of a different organism compared to a reference ALAS protein sequence, such as SEQ ID NO: 29) can be identified by performing a sequence alignment between the protein sequences of interest. It will be understood that in some cases, a gap exists in a protein alignment. Similarly, a
“corresponding” nucleic acid position (or substitution) in a nucleic acid sequence different
10 from a reference nucleic acid sequence (e.g., in the ALAS nucleic acid sequence of a different organism compared to a reference ALAS nucleic acid sequence, such as SEQ ID NO: 28) can be identified by performing a sequence alignment between the nucleic acid sequences of interest. It will be understood that in some cases, a gap exists in a nucleic acid alignment. As used herein, a nucleotide or amino acid position “relative to” a reference
15 sequence can be the corresponding nucleotide or amino acid position in a reference sequence.

Nucleic acid molecules used in the methods described herein are typically DNA, but RNA molecules can be used under the appropriate circumstances. As used herein, “exogenous” refers to any nucleic acid sequence that is introduced into a cell from, for example, the same or a different organism or a nucleic acid generated synthetically (e.g., a
20 codon-optimized nucleic acid sequence). For example, an exogenous nucleic acid can be a nucleic acid from one microorganism (e.g., one genus or species of methylotrophic yeast) that is introduced into a different genus or species of methylotrophic yeast; however, an exogenous nucleic acid also can be a nucleic acid from a methylotrophic yeast that is introduced recombinantly into a methylotrophic yeast as an additional copy despite the
25 presence of a corresponding native nucleic acid sequence, or a nucleic acid from a methylotrophic yeast that is introduced recombinantly into a methylotrophic yeast containing one or more mutations, insertions, or deletions compared to the sequence native to the methylotrophic yeast. For example, *P. pastoris* contains an endogenous nucleic acid encoding an ALAS; an additional copy of the *P. pastoris* ALAS nucleic acid (e.g., introduced
30 recombinantly into *P. pastoris*) is considered to be exogenous. Similarly, an “exogenous” protein is a protein encoded by an exogenous nucleic acid.

In some instances, an exogenous nucleic acid can be a heterologous nucleic acid. As used herein, a “heterologous” nucleic acid refers to any nucleic acid sequence that is not native to an organism (e.g., a heterologous nucleic acid can be a nucleic acid from one microorganism (e.g., one genus or species of methylotrophic yeast, whether or not it has been
 5 codon-optimized) that is introduced into a different genus or species of methylotrophic yeast)). Similarly, a “heterologous” protein is a protein encoded by a heterologous nucleic acid.

A nucleic acid molecule is considered to be exogenous to a host organism when any portion thereof (e.g., a promoter sequence or a sequence of an encoded protein) is exogenous
 10 to the host organism. A nucleic acid molecule is considered to be heterologous to a host organism when any portion thereof (e.g., a promoter sequence or a sequence of an encoded protein) is heterologous to the host organism.

In some embodiments, this document provides a cell (e.g., fungal cell, such as an *Aspergillus* cell, a *Trichoderma* cell, or a yeast cell (e.g., a methylotrophic yeast cell))
 15 comprising a first nucleic acid construct as disclosed herein and a second nucleic acid construct comprising a nucleotide sequence encoding a heme-binding protein. In some embodiments, the second nucleic acid encodes a heme-binding protein operably linked to the first promoter element. In some embodiments, the second nucleic acid encodes a heme-binding protein operably linked to a second promoter element. In some embodiments, a
 20 heme-binding protein can be an exogenous heme-binding protein. In some embodiments, a heme-binding protein can be a heterologous heme-binding protein. In some embodiments, a heme-binding protein can be selected from the group consisting of a globin (PF00042 in the Pfam database), a cytochrome (e.g., a cytochrome P450, a cytochrome a, a cytochrome b, a cytochrome c), a cytochrome c oxidase, a ligninase, a catalase, and a peroxidase. In some
 25 embodiments, a globin can be selected from the group consisting of an androglobin, a chlorocruorin, a cytoglobin, an erythrocruorin, a flavohemoglobin, a globin E, a globin X, a globin Y, a hemoglobin (e.g., a beta hemoglobin, an alpha hemoglobin), a histoglobin, a leghemoglobin, a myoglobin, a neuroglobin, a non-symbiotic hemoglobin, a protoglobin, and a truncated hemoglobin (e.g., a HbN, a HbO, a Glb3, a cyanoglobin). In some embodiments,
 30 the heme-binding protein can be a non-symbiotic hemoglobin. In some embodiments, the heme-binding protein can be a leghemoglobin. In some embodiments, the heme-binding protein can be soybean leghemoglobin (LegH). A reference amino acid sequence for LegH is

provided in Figure 4 as SEQ ID NO: 4. LegH is a protein that binds to heme, which results in a characteristic absorption at 415 nm and a distinct red color. The LegH protein (also known as LGB2) is naturally found in root nodules of soybean (see, for example, UniprotKB Accession No. P02236). See, also, WO 2014/110539 and WO 2014/110532, each of which

5 is herein incorporated by reference in its entirety. In some embodiments, a heme-binding protein can have an amino acid sequence that is at least 70% (e.g., at least 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%) identical to the amino acid sequence set forth in any of SEQ ID NOs: 1-27 (Figure 4). In some embodiments, a heme-binding protein can have an amino acid sequence set forth in any of SEQ ID NOs: 1-27. In

10 some embodiments, a heme-binding protein uses heme as a cofactor. As used herein, a “cofactor” is a molecule or ion that is directly involved in enzyme catalysis. In some embodiments, a heme-binding protein may or may not be an enzyme. In some embodiments, the heme-binding protein may or may not be part of the heme biosynthesis pathway. In some embodiments, the heme-binding protein may or may not be an ALAS. In some embodiments,

15 the heme-binding protein may or may not be a ferrochelatase. In some embodiments, the heme-binding protein may or may not be a coprotoporphyrinogen oxidase. In some embodiments, the heme binding protein may or may not be ALAD, PBGD, UPG3S, UPG3D, COPROX, PROTOX, or FC. In some embodiments, a heme-binding protein has an amino acid sequence that is at least 70% (e.g., at least 75%, 80%, 85%, 90%, or 95%) identical to an

20 amino acid sequence in any of SEQ ID NOs: 1-27. In some embodiments, a heme-binding protein is a bacteria-derived heme-binding protein, a yeast-derived heme-binding protein, an algae-derived heme-binding protein, a fungus-derived heme-binding protein, or a plant-derived heme-binding protein.

As used herein, a “bacteria-derived protein”, “yeast-derived protein”, “algae-derived protein”, “fungus-derived protein”, or “plant-derived protein” refers to the immediate source of the protein, and can mean any protein that is produced in a bacterium, a yeast, an algae, a fungus, or a plant, respectively, independently of whether the protein is natively expressed in the bacterium, yeast, algae, fungus, or plant, respectively.

Provided herein are methods of producing an ALAS protein. Also provided herein are

30 methods of producing an ALAS protein using any of the cells (e.g., fungal cells, such as *Aspergillus* cells, *Trichoderma* cells, or yeast cells (e.g., methylotrophic yeast cells)) described herein. Also provided herein are methods of producing an ALAS protein using any

of the nucleic acid constructs described herein. In some embodiments, provided herein are methods comprising expressing a first nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM. In some embodiments, the titer of an ALAS protein can be increased by at least 5% (e.g., at least 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 120%, 140%, 150%, 160%, 180%, 200%, 220%, 240%, 250%, or more) compared to a corresponding method lacking a nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM. In some embodiments, the titer of an ALAS protein can be increased by at least 5% (e.g., at least 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 120%, 140%, 150%, 160%, 180%, 200%, 220%, 240%, 250%, or more) compared to a corresponding method expressing a nucleic acid encoding an ALAS protein that does not comprise a mutation in a first HRM. In some embodiments, provided herein are methods comprising expressing a first nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM and a mutation in a second HRM. In some embodiments, the titer of an ALAS protein can be increased by at least 5% (e.g., at least 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 120%, 140%, 150%, 160%, 180%, 200%, 220%, 240%, 250%, or more) compared to a corresponding method lacking a nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM and a mutation in a second HRM. In some embodiments, the titer of an ALAS protein can be increased by at least 5% (e.g., at least 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 120%, 140%, 150%, 160%, 180%, 200%, 220%, 240%, 250%, or more) compared to a corresponding method expressing a nucleic acid encoding an ALAS protein that does not comprise a mutation in a first HRM and a mutation in a second HRM.

Provided herein are methods of producing a tetrapyrrole or a derivative thereof. Also provided herein are methods of producing a tetrapyrrole or a derivative thereof using any of the cells (e.g., fungal cells, such as *Aspergillus* cells, *Trichoderma* cells, or yeast cells (e.g., methylotrophic yeast cells)) described herein. Also provided herein are methods of producing a tetrapyrrole or a derivative thereof using any of the nucleic acid constructs described herein. In some embodiments, provided herein are methods of producing a tetrapyrrole or a derivative thereof comprising expressing a first nucleic acid construct comprising a

nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM. In some embodiments, the titer of a tetrapyrrole or a derivative thereof can be increased by at least 5% (e.g., at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 120%, 140%, 150%, 160%, 180%, 200%, 220%, 240%, 250%, 280%, 300%, 320%, 340%, 350%, or more) compared to a corresponding method lacking a nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM. In some embodiments, the titer of a tetrapyrrole or a derivative thereof can be increased by at least 5% (e.g., at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 120%, 140%, 150%, 160%, 180%, 200%, 220%, 240%, 250%, 280%, 300%, 320%, 340%, 350%, or more) compared to a corresponding method expressing a nucleic acid encoding an ALAS protein that does not comprise a mutation in a first HRM. In some embodiments, provided herein are methods of producing a tetrapyrrole or a derivative thereof comprising expressing a first nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM and a second mutation in a second HRM. In some embodiments, the titer of a tetrapyrrole or a derivative thereof can be increased by at least 5% (e.g., at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 120%, 140%, 150%, 160%, 180%, 200%, 220%, 240%, 250%, 280%, 300%, 320%, 340%, 350%, or more) compared to a corresponding method lacking a nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM and a mutation in a second HRM. In some embodiments, the titer of a tetrapyrrole or a derivative thereof can be increased by at least 5% (e.g., at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 120%, 140%, 150%, 160%, 180%, 200%, 220%, 240%, 250%, 280%, 300%, 320%, 340%, 350%, or more) compared to a corresponding method expressing a nucleic acid encoding an ALAS protein that does not comprise a mutation in a first HRM and a mutation in a second HRM. In some embodiments, a tetrapyrrole or a derivative thereof is a chlorin (e.g., a chlorophyll) or a derivative thereof. In some embodiments, a tetrapyrrole or a derivative thereof is a precorrin, a co-precorrin, a corrin (e.g., vitamin B12), or a derivative thereof. In some embodiments, a tetrapyrrole or a derivative thereof is a porphyrin or a derivative thereof. In some embodiments, a tetrapyrrole or a derivative thereof is a heme or a derivative thereof. In some embodiments, a tetrapyrrole or a derivative thereof is heme B. In some embodiments, a tetrapyrrole or a derivative thereof is a metabolic product of heme (e.g., bilirubin or a derivative thereof).

Provided herein are methods of producing an intermediate of the heme biosynthesis pathway, a heme (e.g., heme B, heme o, heme a, heme c), a corrin (e.g., Vitamin B12), a chlorophyll, or a derivative thereof. Also provided herein are methods of producing an intermediate of the heme biosynthesis pathway, a heme, a corrin, a chlorophyll, or a derivative thereof using any of the cells (e.g., fungal cells, such as *Aspergillus* cells, *Trichoderma* cells, or yeast cells (e.g., methylotrophic yeast cells)) described herein. Also provided herein are methods of producing an intermediate of the heme biosynthesis pathway, a heme, a corrin, a chlorophyll, or a derivative thereof using any of the nucleic acid constructs described herein. In some embodiments, provided herein are methods of producing an intermediate of the heme biosynthesis pathway, a heme, a corrin, a chlorophyll, or a derivative thereof comprising expressing a first nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM. In some embodiments, the titer of an intermediate of the heme biosynthesis pathway, a heme, a corrin, a chlorophyll, or a derivative thereof can be increased by at least 5% (e.g., at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 120%, 140%, 150%, 160%, 180%, 200%, 220%, 240%, 250%, 280%, 300%, 320%, 340%, 350%, or more) compared to a corresponding method lacking a nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM. In some embodiments, the titer of an intermediate of the heme biosynthesis pathway, a heme, a corrin, a chlorophyll, or a derivative thereof can be increased by at least 5% (e.g., at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 120%, 140%, 150%, 160%, 180%, 200%, 220%, 240%, 250%, 280%, 300%, 320%, 340%, 350%, or more) compared to a corresponding method expressing a nucleic acid encoding an ALAS protein that does not comprise a mutation in a first HRM. In some embodiments, provided herein are methods of producing an intermediate of the heme biosynthesis pathway, a heme, a corrin, a chlorophyll, or a derivative thereof comprising expressing a first nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM and a mutation in a second HRM. In some embodiments, the titer of an intermediate of the heme biosynthesis pathway, a heme, a corrin, a chlorophyll, or a derivative thereof can be increased by at least 5% (e.g., at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 120%, 140%, 150%, 160%, 180%, 200%, 220%, 240%, 250%, 280%, 300%, 320%, 340%, 350%, or more) compared to a corresponding method lacking a nucleic acid construct comprising a nucleotide sequence

encoding an ALAS protein comprising a mutation in a first HRM and a mutation in a second HRM. In some embodiments, the titer of an intermediate of the heme biosynthesis pathway, a heme, a corrin, a chlorophyll, or a derivative thereof can be increased by at least 5% (e.g., at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 120%, 140%, 150%, 160%,
 5 180%, 200%, 220%, 240%, 250%, 280%, 300%, 320%, 340%, 350%, or more) compared to a corresponding method expressing a nucleic acid encoding an ALAS protein that does not comprise a mutation in a first HRM and a mutation in a second HRM. In some embodiments, an intermediate of the heme biosynthesis pathway, a heme, or a derivative thereof is a heme. In some embodiments, an intermediate of the heme biosynthesis pathway, a heme, or a
 10 derivative thereof is heme B.

Provided herein are methods of producing a heme (e.g., heme B, heme o, heme a, heme c) or a derivative thereof. Also provided herein are methods of producing a heme or derivative thereof using any of the cells (e.g., fungal cells, such as *Aspergillus* cells, *Trichoderma* cells, or yeast cells (e.g., methylotrophic yeast cells)) described herein. Also
 15 provided herein are methods of producing a heme or derivative thereof using any of the nucleic acid constructs described herein. In some embodiments, provided herein are methods of producing a heme or derivative thereof comprising expressing a first nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM. In some embodiments, the titer of a heme or derivative thereof can be increased by at
 20 least 5% (e.g., at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 120%, 140%, 150%, 160%, 180%, 200%, 220%, 240%, 250%, 280%, 300%, 320%, 340%, 350%, or more) compared to a corresponding method lacking a nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM. In some embodiments, the titer of a heme or derivative thereof can be increased by at least 5%
 25 (e.g., at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 120%, 140%, 150%, 160%, 180%, 200%, 220%, 240%, 250%, 280%, 300%, 320%, 340%, 350%, or more) compared to a corresponding method expressing a nucleic acid encoding an ALAS protein that does not comprise a mutation in a first HRM. In some embodiments, provided herein are methods of producing a heme or derivative thereof comprising expressing a first nucleic acid
 30 construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM and a mutation in a second HRM. In some embodiments, the titer of a heme or derivative thereof can be increased by at least 5% (e.g., at least 10%, 20%, 30%,

40%, 50%, 60%, 70%, 80%, 90%, 100%, 120%, 140%, 150%, 160%, 180%, 200%, 220%, 240%, 250%, 280%, 300%, 320%, 340%, 350%, or more) compared to a corresponding method lacking a nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM and a mutation in a second HRM. In some
5 embodiments, the titer of a heme or derivative thereof can be increased by at least 5% (e.g., at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 120%, 140%, 150%, 160%, 180%, 200%, 220%, 240%, 250%, 280%, 300%, 320%, 340%, 350%, or more) compared to a corresponding method expressing a nucleic acid encoding an ALAS protein that does not comprise a mutation in a first HRM and a mutation in a second HRM. In some embodiments,
10 a heme or a derivative thereof is heme B.

Provided herein are methods of producing a heme-binding protein. Also provided herein are methods of producing a heme-binding protein using any of the cells (e.g., fungal cells, such as *Aspergillus* cells, *Trichoderma* cells, or yeast cells (e.g., methylotrophic yeast cells)) described herein. Also provided herein are methods of producing a heme-binding
15 protein using any of the nucleic acid constructs described herein. In some embodiments, provided herein are methods comprising expressing a first nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM and expressing a second nucleic acid construct comprising a nucleotide sequence encoding a heme-binding protein. In some embodiments, provided herein are methods comprising
20 expressing a first nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM and expressing a second nucleic acid construct comprising a nucleotide sequence encoding a heme-binding protein. In some embodiments, provided herein are methods comprising expressing a first nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM and a
25 mutation in a second HRM and expressing a second nucleic acid construct comprising a nucleotide sequence encoding a heme-binding protein. In some embodiments, provided herein are methods comprising expressing a first nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM and a mutation in a second HRM and expressing a second nucleic acid construct comprising a
30 nucleotide sequence encoding a heme-binding protein. In some embodiments of any of the methods described herein, the methods allow for an increase in the titer of a heme-binding protein. In some embodiments, the titer of a heme-binding protein can be increased by at least

5% (e.g., at least 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 35%, 40%, 45%, 50%, or more) compared to a corresponding method lacking a nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM. In some embodiments, the

5 titer of a heme-binding protein can be increased by at least 5% (e.g., at least 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 35%, 40%, 45%, 50%, or more) compared to a corresponding method expressing a nucleic acid encoding an ALAS protein that does not comprise a mutation in a first HRM.

In some embodiments, the titer of a heme-binding protein can be increased by at least 5%

10 (e.g., at least 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 35%, 40%, 45%, 50%, or more) compared to a corresponding method lacking a nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM and a mutation in a second HRM. In some embodiments, the titer of a heme-binding protein can be increased by at least

15 5% (e.g., at least 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 35%, 40%, 45%, 50%, or more) compared to a corresponding method expressing a nucleic acid encoding an ALAS protein that does not comprise a mutation in a first HRM and a mutation in a second HRM.

Provided herein are methods of producing a tetrapyrrole-binding protein. Also

20 provided herein are methods of producing a tetrapyrrole-binding protein using any of the cells (e.g., fungal cells, such as *Aspergillus* cells, *Trichoderma* cells, or yeast cells (e.g., methylotrophic yeast cells)) described herein. Also provided herein are methods of producing a tetrapyrrole-binding protein using any of the nucleic acid constructs described herein. In some embodiments, provided herein are methods comprising expressing a first

25 nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM and expressing a second nucleic acid construct comprising a nucleotide sequence encoding a tetrapyrrole-binding protein. In some embodiments, provided herein are methods comprising expressing a first nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a

30 mutation in a first HRM and expressing a second nucleic acid construct comprising a nucleotide sequence encoding a tetrapyrrole-binding protein. In some embodiments, provided herein are methods comprising expressing a first nucleic acid construct comprising a

nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM and a mutation in a second HRM and expressing a second nucleic acid construct comprising a nucleotide sequence encoding a tetrapyrrole-binding protein. In some embodiments, provided herein are methods comprising expressing a first nucleic acid construct comprising a

5 nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM and a mutation in a second HRM and expressing a second nucleic acid construct comprising a nucleotide sequence encoding a tetrapyrrole-binding protein. In some embodiments of any of the methods described herein, the methods allow for an increase in the titer of a tetrapyrrole-binding protein. In some embodiments, the titer of a tetrapyrrole-binding protein can be

10 increased by at least 5% (e.g., at least 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 35%, 40%, 45%, 50%, or more) compared to a corresponding method lacking a nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM. In some embodiments, the titer of a tetrapyrrole-binding protein can be increased by at least 5%

15 (e.g., at least 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 35%, 40%, 45%, 50%, or more) compared to a corresponding method expressing a nucleic acid encoding an ALAS protein that does not comprise a mutation in a first HRM. In some embodiments, the titer of a tetrapyrrole-binding protein can be increased by at least 5% (e.g., at least 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%,

20 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 35%, 40%, 45%, 50%, or more) compared to a corresponding method lacking a nucleic acid construct comprising a nucleotide sequence encoding an ALAS protein comprising a mutation in a first HRM and a mutation in a second HRM. In some embodiments, the titer of a tetrapyrrole-binding protein can be increased by at least 5% (e.g., at least 6%, 7%, 8%, 9%, 10%, 11%,

25 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 35%, 40%, 45%, 50%, or more) compared to a corresponding method expressing a nucleic acid encoding an ALAS protein that does not comprise a mutation in a first HRM and a mutation in a second HRM. In some embodiments, a tetrapyrrole is a chlorin (e.g., a chlorophyll) or a derivative thereof. In some embodiments, a tetrapyrrole is a precorrin, a co-precorrin, a corrin

30 (e.g., vitamin B12), or a derivative thereof. In some embodiments, a tetrapyrrole is a porphyrin or a derivative thereof. In some embodiments, a tetrapyrrole is a heme or a

derivative thereof. In some embodiments, a tetrapyrrole is heme B. In some embodiments, a tetrapyrrole is a metabolic product of heme (e.g., bilirubin or a derivative thereof).

As used herein, a “corresponding method” is a method that is essentially identical to a reference method in all ways except for the identified difference. For example, a
5 corresponding method expressing a nucleic acid encoding an ALAS protein that does not comprise a mutation in a first HRM and a mutation in a second HRM would be the same in all aspects (e.g., genetic makeup of cell, temperature and time of culture, and so forth), except that the corresponding method would express a nucleic acid encoding an ALAS protein that does not comprise a mutation in a first HRM and a mutation in a second HRM.

10 Genetically engineering a cell (e.g., fungal cell, such as an *Aspergillus* cell, a *Trichoderma* cell, or a yeast cell (e.g., a methylotrophic yeast cell)) typically includes introducing a recombinant nucleic acid molecule (also called a nucleic acid construct) into the cell. As described herein, a recombinant nucleic acid molecule typically includes an exogenous nucleic acid that encodes a protein (e.g., a protein involved in heme biosynthesis,
15 a heme-binding protein, or a transcription factor) operably linked to at least one promoter element (e.g., an inducible or constitutive promoter element). In some embodiments, a recombinant nucleic acid molecule can include a linear sequence of two or more protein-coding sequences operably linked to the same or separate promoter elements (e.g., a first promoter operably linked to a first nucleic acid encoding a first protein and a second
20 promoter operably linked to a second nucleic acid encoding a second protein, or a promoter operably linked to a first nucleic acid encoding a first protein and a second nucleic acid encoding a second protein). In some cases, a recombinant nucleic acid molecule including at least one promoter operably linked to a nucleotide sequence encoding a protein can be called a cassette.

25 A recombinant nucleic acid can include expression elements. Expression elements include nucleic acid sequences that direct and regulate expression of nucleic acid coding sequences. One example of an expression element is a promoter sequence. Expression elements also can include introns, enhancer sequences, response elements, or inducible elements that modulate expression of a nucleic acid. Expression elements can be of bacterial,
30 yeast, insect, mammalian, or viral origin, and vectors can contain a combination of elements from different origins.

Nucleic acids can be detected using any number of amplification techniques (see, e.g., PCR Primer: A Laboratory Manual, 1995, Dieffenbach & Dveksler, Eds., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; and U.S. Patent Nos. 4,683,195; 4,683,202; 4,800,159; and 4,965,188) with an appropriate pair of oligonucleotides (e.g., primers). A number of modifications to the original PCR method have been developed and can be used to detect selected nucleic acids.

Methanol utilization is typically initiated by the conversion of methanol to formaldehyde by the action of alcohol oxidase. *Pichia pastoris* contains two genes for alcohol oxidases, AOX1 and AOX2. Strains with reduced alcohol oxidase activity (“methanol utilization slow” or MutS strains) can produce more of a recombinant protein expressed from the AOX1 promoter than strains that do not have reduced alcohol oxidase activity. Strains mutated in both AOX genes and completely lacking alcohol oxidase activity cannot metabolize methanol, but can still be induced for expression from the AOX1 promoter by methanol. These strains retain the ability to use other carbon sources for growth, but still express heterologous proteins from the AOX1 promoter upon the addition of methanol. Because these strains do not metabolize methanol (“methanol utilization minus” or Mut- strains), much less methanol is required for induction of protein expression, and strains carrying these mutations avoid issues related to methanol feeding in large-scale fermentations. See, for example, Chiruvolu et al., 1997, *Enzyme Microb. Technol.*, 21:277-83.

Suitable transcription factors, and nucleic acids encoding transcription factors (e.g., exogenous nucleic acids encoding transcription factors), include, for example, Mxr1 from a *P. pastoris*. A representative *K. pastoris* Mxr1 nucleic acid sequence can be found, for example, in GenBank Accession No. DQ395124, while a representative *K. pastoris* Mxr1 protein sequence can be found, for example, in GenBank Accession No. ABD57365. In some embodiments, the transcription factor is a Mit1 sequence from *K. phaffii* (see, for example, UniParc Accession No. UPI0001A4D18B). Suitable transcription factors also can be found in *Hansenula polymorpha* (e.g., the Adr1 sequence; see, for example, GenBank Accession No. AEOI02000005, bases 858873 to 862352, for the nucleic acid sequence and GenBank Accession No. ESX01253 for the amino acid sequence) and *Candida boidinii* (e.g., the Trm1 sequence; see, for example, GenBank Accession No. AB365355 for the nucleic acid sequence and GenBank Accession No. BAF99700 for the amino acid sequence; and Trm2 sequence;

see, for example, GenBank Accession No. AB548760 for the nucleic acid sequence and GenBank Accession No. BAJ07608 for the amino acid sequence).

Transcription factors such as Mxr1 may be normally expressed at low levels. In some embodiments, it is desirable to place an exogenous nucleic acid (e.g., a transcription factor) under control of a promoter that is inducible.

Methanol-regulated transcription factors in *Pichia* can bind to an AOX1 promoter and act cooperatively with Mxr1 to activate transcription from an AOX1 promoter. In some embodiments, two methanol-regulated transcription factors (e.g., Mxr1 and Mit1) can be operably linked to a methanol inducible promoter element.

There are a number of inducible promoters that can be used when genetically engineering cells (e.g., fungal cells such as *Aspergillus*, *Trichoderma* cells, or yeast cells (e.g., methylotrophic yeast cells)). Suitable methanol inducible promoters include pAOX1, (e.g., a pAOX1 such as the promoter for *K. pastoris* AOX1 (see, for example, the promoter for GenBank Accession No. U96967.1) or a promoter described in U.S. Provisional Patent Application No. 62/835,338, filed April 17, 2019, incorporated herein by reference in its entirety), as well as other methanol-inducible promoters, or promoter elements therefrom. These include, without limitation, a pAOX2 promoter (e.g., from *K. phaffii* or *K. pastoris* (see, for example, GenBank Accession No. X79871.1)), an alcohol oxidase (AOD1) promoter from, e.g., *Candida boidinii* (see, for example, GenBank Accession No. E06147.1), the alcohol oxidase (MOX) promoter from *Hansenula polymorpha* (see, for example, GenBank Accession No. AJ313360.1), the MOD1 or MOD2 promoter from *Pichia methanolica* (see, for example, Raymond et al., 1998, Yeast, 14:11-23; and Nakagawa et al., 1999, Yeast, 15:1223-30), the DHAS promoter from *P. pastoris* (see, for example, the promoter for GenBank Accession No. FJ752551) or a promoter element therefrom, the formaldehyde dehydrogenase (FLD1) promoter from *K. pastoris* (see, for example, GenBank Accession No. KJ755994.1), or the PEX8 promoter from *P. pastoris* (see, for example, Kranthi et al., 2010, Yeast, 27:705-11). Typically, these promoters can be induced by methanol. Suitable constitutive promoters and constitutive promoter elements include, without limitation, the *P. pastoris* promoter (or a portion thereof) from the transcriptional elongation factor EF-1 α gene (TEF1), which is strongly transcribed in a constitutive manner. Other suitable constitutive promoters (or promoter elements therefrom) also can be used, including, without limitation, the glyceraldehyde-3-phosphate dehydrogenase (GAPDH) promoter from *K. pastoris* (see,

for example, the promoter for GenBank Accession No. U62648.1), the promoter from the potential glycosyl phosphatidyl inositol (GPI)-anchored protein, GCW14p (PAS_chr1-4_0586) from *K. phaffii* (see, for example, the promoter for GenBank Accession No. XM_002490678), and the promoter from the 3-phosphoglycerate kinase gene (PGK1) from *K. pastoris* (see, for example, GenBank Accession No. AY288296). It will be appreciated that a choice of promoter may be influenced by the expression system. For example, for expression in *K. phaffii*, a *K. phaffii* promoter might be chosen, while for expression in *C. boidinii*, a *C. boidinii* promoter might be chosen. However, in some cases, a promoter from one organism (e.g., *K. phaffii*) may be appropriate to be used in another organism (e.g., *C. boidinii* or *K. pastoris*). Further, it is noted that a combination of inducible (e.g., methanol-inducible) and constitutive promoters (or promoter elements therefrom) can be combined to further increase the expression of any of the nucleic acids operably linked thereto.

Any of the encoded proteins as described herein can be operably linked to an inducible promoter element (e.g., a methanol-inducible promoter element) or a constitutive promoter element. Inducible promoters and elements therefrom are discussed above. Constitutive promoters and constitutive promoter elements are known in the art. For example, a commonly used constitutive promoter from *P. pastoris* is the promoter, or a portion thereof, from the transcriptional elongation factor EF-1 α gene (TEF1), which is strongly transcribed in a constitutive manner. Other constitutive promoters, or promoter elements therefrom, however, can be used, including, without limitation, the glyceraldehyde-3-phosphate dehydrogenase (GAPDH) promoter from *K. pastoris* (see, for example, the promoter for GenBank Accession No. U62648.1), the promoter from the potential glycosyl phosphatidyl inositol (GPI)-anchored protein, GCW14p (PAS_chr1-4_0586), from *K. phaffii* (see, for example, the promoter for GenBank Accession No. XM_002490678), or the promoter from the 3-phosphoglycerate kinase gene (PGK1) from *K. pastoris* (see, for example, the promoter for GenBank Accession No. AY288296).

In some embodiments, any of the cells (e.g., fungal cell, such as an *Aspergillus* cell, a *Trichoderma* cell, or a yeast cell (e.g., a methylotrophic yeast cell)) herein can further include a third nucleic acid construct comprising a nucleotide sequence encoding a third protein operably linked to the first promoter element, the second promoter element, or a third promoter element. In some embodiments, any of the cells (e.g., fungal cell, such as an *Aspergillus* cell, a *Trichoderma* cell, or a yeast cell (e.g., a methylotrophic yeast cell)) herein

can further include a fourth nucleic acid construct comprising a nucleotide sequence encoding a fourth protein operably linked to the first promoter element, the second promoter element, the third promoter element, or a fourth promoter element. In some embodiments, the third protein can be a transcription factor. In some embodiments, the fourth protein can be a transcription factor. In some embodiments, any of the promoter elements herein (e.g., a first promoter element, a second promoter element, a third promoter element, or a fourth promoter element) can contain one or more recognition sequences for a transcription factor. Therefore, in some embodiments, a feedback loop may be constructed such that the transcription factor drives the expression of additional copies of the transcription factor, as well as the expression of one or more of an ALAS and a heme-binding protein. In some embodiments, a transcription factor can be Mxr1 (see, e.g., U.S. Patent No. 9,938,327, which is incorporated by reference in its entirety). In some embodiments, a third protein can be a protein involved in heme biosynthesis. In some embodiments, a fourth protein can be a protein involved in heme biosynthesis. In some embodiments, a protein involved in heme biosynthesis can be selected from the group consisting of δ -aminolevulinic acid dehydratase (ALAD), porphobilinogen deaminase (PBGD), uroporphyrinogen III synthase (UPG3S), uroporphyrinogen III decarboxylase (UPG3D), coprotoporphyrinogen oxidase (COPROX), protoporphyrinogen IX oxidase (PROTOX), and/or ferrochelatase (FC). In some embodiments, a protein involved in heme biosynthesis can be selected from the group consisting of δ -aminolevulinic acid dehydratase (ALAD), porphobilinogen deaminase (PBGD), uroporphyrinogen III synthase (UPG3S), uroporphyrinogen III decarboxylase (UPG3D), coprotoporphyrinogen oxidase (COPROX), and/or protoporphyrinogen IX oxidase (PROTOX).

Previous studies in *Saccharomyces cerevisiae* identified ALA dehydratase and porphobilinogen deaminase as rate limiting enzymes in heme biosynthesis (see, for example, Hoffman et al., 2003, Biochem. Biophys. Res. Commun., 310(4):1247-53). However, heterologous expression of individual heme enzymes in *P. pastoris* from the glyceraldehyde-3-phosphate dehydrogenase (GAP) promoter failed to overcome limitations associated with the expression of a recombinant protein containing a heme (see Krainer et al., 2015, Microb. Cell Fact., 13;14:4). Expression of a recombinant heme containing protein in *P. pastoris* can be achieved by co-expressing one or more enzymes of the heme biosynthetic pathway from methanol-inducible promoters, although it would be appreciated that one or more of the genes

involved in the heme biosynthetic pathway could be expressed from one or more constitutive promoters (see, e.g., U.S. Patent No. 9,938,327, which is incorporated by reference in its entirety).

In addition, it is noted that a first nucleic acid encoding a first protein (e.g., an ALAS protein) operably linked to a promoter element as described herein can be physically separate from a second nucleic acid encoding a second protein (e.g., a heme-binding protein) operably linked to a promoter element (that is, the first and second nucleic acids can be completely separate molecules). Alternatively, a first nucleic acid encoding a first protein operably linked to a promoter element and a second nucleic acid encoding a second protein operably linked to a promoter element can be included in the same nucleic acid construct. In some embodiments, a first nucleic acid encoding a first protein operably linked to a promoter element can be contiguous with a second nucleic acid encoding a second protein operably linked to a promoter element. It would be appreciated by a skilled artisan that, if the second nucleic acid molecule encoding a second protein is contiguous with the first nucleic acid encoding a protein of interest, a single promoter, or promoter element therefrom, can be used to drive transcription of both or all of the genes (e.g., the nucleic acid encoding the first protein as well as the second protein).

Methods of introducing nucleic acids into cells (e.g., fungal cells, such as *Aspergillus* cells, *Trichoderma* cells, or yeast cells (e.g., methylotrophic yeast cells)) are known in the art, and include, without limitation, transduction, electroporation, biolistic particle delivery, and chemical transformation.

In addition, methods of culturing cells (e.g., fungal cells, such as *Aspergillus* cells, *Trichoderma* cells, or yeast cells (e.g., methylotrophic yeast cells)) are known in the art. See, for example, Pichia Protocols, *Methods In Molecular Biology*, 389, Cregg, Ed., 2007, 2nd Ed., Humana Press, Inc. Under some circumstances, it may be desirable to introduce or add methanol to the culture media, although, as demonstrated herein, methanol is not required to obtain efficient expression at high levels of one or more proteins of interest. Under some circumstances (e.g., when one or more nucleic acids encoding enzyme(s) involved in heme biosynthesis are expressed), it may be desirable to supplement the culture media with iron or a pharmaceutically or metabolically acceptable (or GRAS) salt thereof.

The recombinant nucleic acid molecules described herein can be stably integrated into the genome of the cell (e.g., the fungal cell, such as *Aspergillus* cell, *Trichoderma* cell, or

yeast cell (e.g., the methylotrophic yeast cell)), or can be extrachromosomally expressed from a replication-competent plasmid. Methods of achieving both are known and used in the art.

The methods provided herein also can include purifying the expressed protein. As used herein, an “enriched” protein is a protein that accounts for at least 5% (e.g., at least 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, or more) by dry weight, of the mass of the production cell (e.g., fungal cell, such as *Aspergillus* cell, *Trichoderma* cell, or yeast cell (e.g., methylotrophic yeast cell)), or at least 10% (e.g., at least 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 90%, 95%, or 99%) by dry weight, the mass of the production cell lysate (e.g., excluding cell wall or membrane material). As used herein, a “purified” protein is a protein that has been separated or purified from cellular components that naturally accompany it. Typically, the protein is considered “purified” when it is at least 60% (e.g., at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99%) by dry weight, free from the proteins and naturally occurring molecules with which it is naturally associated.

As used herein, nucleic acids can include DNA and RNA, and includes nucleic acids that contain one or more nucleotide analogs or backbone modifications. A nucleic acid can be single stranded or double stranded, which usually depends upon its intended use. Also provided are nucleic acids and polypeptides that differ from a given sequence. Nucleic acids and polypeptides can have at least 50% sequence identity (e.g., at least 55%, 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity) to a given nucleic acid or polypeptide sequence.

In calculating percent sequence identity, two sequences are aligned and the number of identical matches of nucleotides or amino acid residues between the two sequences is determined. The number of identical matches is divided by the length of the aligned region (i.e., the number of aligned nucleotides or amino acid residues) and multiplied by 100 to arrive at a percent sequence identity value. It will be appreciated that the length of the aligned region can be a portion of one or both sequences up to the full-length size of the shortest sequence. It also will be appreciated that a single sequence can align with more than one other sequence and hence, can have different percent sequence identity values over each aligned region.

The alignment of two or more sequences to determine percent sequence identity can be performed using the computer program ClustalW and default parameters, which allows alignments of nucleic acid or polypeptide sequences to be carried out across their entire length (global alignment). Chenna et al., 2003, *Nucleic Acids Res.*, 31(13):3497-500.

5 ClustalW calculates the best match between a query and one or more subject sequences, and aligns them so that identities, similarities and differences can be determined. Gaps of one or more residues can be inserted into a query sequence, a subject sequence, or both, to maximize sequence alignments. For fast pairwise alignment of nucleic acid sequences, the default parameters can be used (i.e., word size: 2; window size: 4; scoring method: percentage;
10 number of top diagonals: 4; and gap penalty: 5); for an alignment of multiple nucleic acid sequences, the following parameters can be used: gap opening penalty: 10.0; gap extension penalty: 5.0; and weight transitions: yes. For fast pairwise alignment of polypeptide sequences, the following parameters can be used: word size: 1; window size: 5; scoring method: percentage; number of top diagonals: 5; and gap penalty: 3. For multiple alignment
15 of polypeptide sequences, the following parameters can be used: weight matrix: blosum; gap opening penalty: 10.0; gap extension penalty: 0.05; hydrophilic gaps: on; hydrophilic residues: Gly, Pro, Ser, Asn, Asp, Gln, Glu, Arg, and Lys; and residue-specific gap penalties: on. ClustalW can be run, for example, at the Baylor College of Medicine Search Launcher website or at the European Bioinformatics Institute website on the World Wide Web.

20 Changes can be introduced into a nucleic acid molecule, thereby leading to changes in the amino acid sequence of the encoded polypeptide. For example, changes can be introduced into nucleic acid coding sequences using mutagenesis (e.g., site-directed mutagenesis, PCR-mediated mutagenesis, transposon mutagenesis, chemical mutagenesis, UV mutagenesis or radiation induced mutagenesis) or by chemically synthesizing a nucleic
25 acid molecule having such changes. Such nucleic acid changes can lead to conservative and/or non-conservative amino acid substitutions at one or more amino acid residues. A “conservative amino acid substitution” is one in which one amino acid residue is replaced with a different amino acid residue having a similar side chain (see, for example, Dayhoff et al., 1978, *Atlas of Protein Sequence and Structure*, 5(Suppl. 3):345-352, which provides
30 frequency tables for amino acid substitutions), and a non-conservative substitution is one in which an amino acid residue is replaced with an amino acid residue that does not have a similar side chain. Nucleic acid and/or polypeptide sequences may be modified as described

herein to improve one or more properties such as, without limitation, increased expression (e.g., transcription and/or translation), tighter regulation, deregulation, loss of catabolite repression, modified specificity, secretion, thermostability, solvent stability, oxidative stability, protease resistance, catalytic activity, and/or color.

5 As used herein, an “isolated” nucleic acid molecule is a nucleic acid molecule that is free of sequences that naturally flank one or both ends of the nucleic acid in the genome of the organism from which the isolated nucleic acid molecule is derived (e.g., a cDNA or genomic DNA fragment produced by PCR or restriction endonuclease digestion). Such an isolated nucleic acid molecule is generally introduced into a vector (e.g., a cloning vector, or
10 an expression vector) for convenience of manipulation or to generate a fusion nucleic acid molecule, discussed in more detail below. In addition, an isolated nucleic acid molecule can include an engineered nucleic acid molecule such as a recombinant or a synthetic nucleic acid molecule.

Vectors as described herein can be introduced into a host cell (e.g., fungal cell, such
15 as an *Aspergillus* cell, a *Trichoderma* cell, or a yeast cell (e.g., a methylotrophic yeast cell)). As used herein, “host cell” refers to the particular cell into which the nucleic acid is introduced and also includes the progeny of such a cell that carry the vector. A host cell can be any prokaryotic or eukaryotic cell. For example, nucleic acids can be expressed in bacterial cells such as *E. coli*, or in insect cells, yeast or mammalian cells (such as Chinese
20 hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art. Many methods for introducing nucleic acids into host cells, both *in vivo* and *in vitro*, are well known to those skilled in the art and include, without limitation, electroporation, calcium phosphate precipitation, polyethylene glycol (PEG) transformation, heat shock, lipofection, microinjection, and viral-mediated nucleic acid transfer.

25 Nucleic acids can be isolated using techniques routine in the art. For example, nucleic acids can be isolated using any method including, without limitation, recombinant nucleic acid technology, and/or the polymerase chain reaction (PCR). General PCR techniques are described, for example in PCR Primer: A Laboratory Manual, Dieffenbach & Dveksler, Eds., Cold Spring Harbor Laboratory Press, 1995. Recombinant nucleic acid techniques include,
30 for example, restriction enzyme digestion and ligation, which can be used to isolate a nucleic acid. Isolated nucleic acids also can be chemically synthesized, either as a single nucleic acid molecule or as a series of oligonucleotides.

Polypeptides can be purified from natural sources (e.g., a biological sample) by known methods such as DEAE ion exchange, gel filtration, and hydroxyapatite chromatography. A polypeptide also can be purified, for example, by expressing a nucleic acid in an expression vector. In addition, a purified polypeptide can be obtained by chemical synthesis. The extent of purity of a polypeptide can be measured using any appropriate method, e.g., column chromatography, polyacrylamide gel electrophoresis, or HPLC analysis.

A construct or vector containing a nucleic acid (e.g., a nucleic acid that encodes a polypeptide) also is provided. Constructs or vectors, including expression constructs or vectors, are commercially available or can be produced by recombinant DNA techniques routine in the art. A construct or vector containing a nucleic acid can have expression elements operably linked to such a nucleic acid, and further can include sequences such as those encoding a selectable marker (e.g., an antibiotic resistance gene). A construct or vector containing a nucleic acid can encode a chimeric or fusion polypeptide (i.e., a polypeptide operatively linked to a heterologous polypeptide, which can be at either the N-terminus or C-terminus of the polypeptide). Representative heterologous polypeptides include those that can be used in purification of the encoded polypeptide (e.g., 6xHis tag, glutathione S-transferase (GST)).

Nucleic acids also can be detected using hybridization. Hybridization between nucleic acids is discussed in detail in Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; Sections 7.37-7.57, 9.47-9.57, 11.7-11.8, and 11.45-11.57). Sambrook et al. discloses suitable Southern blot conditions for oligonucleotide probes less than about 100 nucleotides (Sections 11.45-11.46). The T_m between a sequence that is less than 100 nucleotides in length and a second sequence can be calculated using the formula provided in Section 11.46. Sambrook et al. additionally discloses Southern blot conditions for oligonucleotide probes greater than about 100 nucleotides (see Sections 9.47-9.54). The T_m between a sequence greater than 100 nucleotides in length and a second sequence can be calculated using the formula provided in Sections 9.50-9.51 of Sambrook et al.

The conditions under which membranes containing nucleic acids are prehybridized and hybridized, as well as the conditions under which membranes containing nucleic acids are washed to remove excess and non-specifically bound probe, can play a significant role in the stringency of the hybridization. Such hybridizations and washes can be performed, where

appropriate, under moderate or high stringency conditions. For example, washing conditions can be made more stringent by decreasing the salt concentration in the wash solutions and/or by increasing the temperature at which the washes are performed. Simply by way of example, high stringency conditions typically include a wash of the membranes in 0.2X SSC at 65°C.

In addition, interpreting the amount of hybridization can be affected, for example, by the specific activity of the labeled oligonucleotide probe, by the number of probe-binding sites on the template nucleic acid to which the probe has hybridized, and by the amount of exposure of an autoradiograph or other detection medium. It will be readily appreciated by those of ordinary skill in the art that although any number of hybridization and washing conditions can be used to examine hybridization of a probe nucleic acid molecule to immobilized target nucleic acids, it is more important to examine hybridization of a probe to target nucleic acids under identical hybridization, washing, and exposure conditions. Preferably, the target nucleic acids are on the same membrane.

A nucleic acid molecule is deemed to hybridize to a nucleic acid but not to another nucleic acid if hybridization to a nucleic acid is at least 5-fold (e.g., at least 6-fold, 7-fold, 8-fold, 9-fold, 10-fold, 20-fold, 50-fold, or 100-fold) greater than hybridization to another nucleic acid. The amount of hybridization can be quantitated directly on a membrane or from an autoradiograph using, for example, a PhosphorImager or a Densitometer (Molecular Dynamics, Sunnyvale, CA).

Polypeptides can be detected using antibodies. Techniques for detecting polypeptides using antibodies include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence. An antibody can be polyclonal or monoclonal. An antibody having specific binding affinity for a polypeptide can be generated using methods well known in the art. The antibody can be attached to a solid support such as a microtiter plate using methods known in the art. In the presence of a polypeptide, an antibody-polypeptide complex is formed.

Detection (e.g., of an amplification product, a hybridization complex, or a polypeptide) is usually accomplished using detectable labels. The term "label" is intended to encompass the use of direct labels as well as indirect labels. Detectable labels include enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials.

Methods are described herein that can be used to generate a strain that lacks sequences for selection (i.e., that lacks a selectable marker). These methods include using a circular plasmid DNA vector and a linear DNA sequence; the circular plasmid DNA vector contains a selection marker and an origin of DNA replication (also known as an autonomously replicating sequence (ARS)), and the linear DNA sequence contains sequences for integration into the genome (e.g., the *Pichia* genome) by homologous recombination. The linear DNA molecule additionally can include nucleic acid sequences encoding one or more proteins of interest such as, without limitation, an ALAS, a heme-binding protein, or a third protein (e.g., a transcription factor or a protein involved in heme biosynthesis).

Cells (e.g., *Pichia* cells) can be transformed with both DNA molecules and the transformants selected by the presence of the selectable marker on the circular plasmid. Transformants then can be screened for integration of the linear DNA molecule into the genome using, for example, PCR. Once transformants with the correct integration of the marker-free linear DNA molecule are identified, the cells can be grown in the absence of selection for the circular plasmid. Because the marker-bearing plasmid is not stably maintained in the absence of selection, the plasmid is lost, often very quickly, after selection is relaxed. The resulting strain carries the integrated linear DNA in the absence of heterologous sequences for selection. Therefore, this approach can be used to construct strains (e.g., *Pichia* strains) that lack a selectable marker (e.g., a heterologous selection marker) with little to no impact on recombinant protein yield.

In accordance with the present disclosure, there may be employed conventional molecular biology, microbiology, biochemical, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. The materials and methods of the disclosure will be further described in the following examples, which do not limit the scope of the methods and compositions of matter described in the claims.

The materials and methods of the disclosure will be further described in the following examples, which do not limit the scope the claims.

Exemplary Embodiments

Embodiment 1 is a cell comprising:

a first exogenous nucleic acid construct comprising a nucleotide sequence encoding an aminolevulinate synthase (ALAS) protein operably linked to a first promoter element, wherein the ALAS comprises at least a first heme responsive motif (HRM), and wherein the ALAS comprises a mutation in the first HRM; and

5 a second exogenous nucleic acid construct comprising a nucleotide sequence encoding a heme-binding protein,

wherein the second exogenous nucleic acid construct comprising a nucleotide sequence encoding the heme-binding protein is operably linked to the first promoter element or is operably linked to a second promoter element.

10 Embodiment 2 is the cell of embodiment 1, wherein the cell is a fungal cell.

Embodiment 3 is the cell of embodiment 2, wherein the cell is an *Aspergillus* or *Trichoderma* cell.

Embodiment 4 is the cell of any one of embodiments 1-3, wherein the cell is a yeast cell.

15 Embodiment 5 is the cell of embodiment 4, wherein the yeast cell is a methylotrophic yeast cell.

Embodiment 6 is the cell of embodiment 5, wherein the methylotrophic yeast cell is a *Pichia* cell, a *Candida* cell, a *Hansenula* cell, or a *Torulopsis* cell.

20 Embodiment 7 is the cell of any one of embodiments 5-6, wherein the methylotrophic yeast cell is a *Pichia methanolica* cell, a *Pichia pastoris* cell, a *Candida boidinii* cell, or a *Hansenula polymorpha* cell.

Embodiment 8 is the cell of any one of embodiments 5-7, wherein the methylotrophic yeast cell is a *Pichia pastoris* cell.

25 Embodiment 9 is the cell of any one of embodiments 1-8, wherein the mutation in the first HRM is a mutation from a cysteine to a different amino acid.

Embodiment 10 is the cell of any one of embodiments 1-9, wherein the ALAS protein comprises a second HRM, and wherein the ALAS protein comprises a mutation in the second HRM.

30 Embodiment 11 is the cell of embodiment 10, wherein the mutation in the second HRM is a mutation from a cysteine to a different amino acid.

Embodiment 12 is the cell of embodiment 10 or embodiment 11, wherein the different amino acid is the same for the mutation in the first HRM and the mutation in the second HRM.

Embodiment 13 is the cell of embodiment 10 or embodiment 11, wherein the different amino acid is not the same for the mutation in the first HRM and the mutation in the second HRM.

Embodiment 14 is the cell of any one of embodiments 10-13, wherein the ALAS protein comprises a third HRM, and wherein the ALAS protein comprises a mutation in the third HRM.

Embodiment 15 is the cell of embodiment 14, wherein the mutation in the third HRM is a mutation from a cysteine to a different amino acid.

Embodiment 16 is the cell of embodiment 15, wherein the different amino acid is the same for the mutation in the first HRM, the mutation in the second HRM, and the mutation in the third HRM.

Embodiment 17 is the cell of any one of embodiments 9-16, wherein the different amino acid is selected from the group consisting of arginine, histidine, lysine, serine, threonine, asparagine, glutamine, selenocysteine, glycine, proline, alanine, isoleucine, leucine, methionine, glutamic acid, aspartic acid, phenylalanine, tryptophan, tyrosine, and valine.

Embodiment 18 is the cell of any one of embodiments 9-16, wherein the different amino acid is selected from a nonpolar aliphatic amino acid, an aromatic amino acid, a polar uncharged amino acid, or a positively charged amino acid.

Embodiment 19 is the cell of embodiment 18, wherein the nonpolar aliphatic amino acid is selected from the group consisting of glycine, proline, alanine, isoleucine, leucine, methionine, and valine.

Embodiment 20 is the cell of embodiment 18, wherein the aromatic amino acid is selected from the group consisting of phenylalanine, tryptophan, and tyrosine.

Embodiment 21 is the cell of embodiment 18, wherein the polar uncharged amino acid is selected from the group consisting of a polar uncharged amino acid serine, threonine, asparagine, or glutamine.

Embodiment 22 is the cell of embodiment 18, wherein the positively charged amino acid is selected from the group consisting of arginine, histidine, and lysine.

Embodiment 23 is the cell of any one of embodiments 9-22, wherein the different amino acid is serine.

Embodiment 24 is the cell of any one of embodiments 9-22, wherein the different amino acid is alanine.

5 Embodiment 25 is the cell of any one of embodiments 9-22, wherein the different amino acid is phenylalanine.

Embodiment 26 is the cell of any one of embodiments 9-22, wherein the different amino acid is histidine.

10 Embodiment 27 is the cell of any one of embodiments 1-26, wherein the first HRM is HRM1.

Embodiment 28 is the cell of any one of embodiments 10-27, wherein the second HRM is HRM2.

Embodiment 29 is the cell of any one of embodiments 1-26, wherein the first HRM is HRM2.

15 Embodiment 30 is the cell of any one of embodiments 10-27, wherein the second HRM is HRM1.

Embodiment 31 is the cell of any one of embodiments 1-30, wherein the first exogenous nucleic acid construct comprises a nucleic acid sequence having at least 90% sequence identity to the nucleic acid sequence in SEQ ID NO: 28.

20 Embodiment 32 is the cell of any one of embodiments 1-30, wherein the first exogenous nucleic acid construct comprises a nucleic acid sequence having at least 95% sequence identity to the nucleic acid sequence in SEQ ID NO: 28.

Embodiment 33 is the cell of any one of embodiments 1-30, wherein the first exogenous nucleic acid construct comprises the nucleic acid sequence in SEQ ID NO: 30.

25 Embodiment 34 is the cell of any one of embodiments 1-32, wherein the ALAS protein comprises an amino acid sequence having at least 90% sequence identity to the amino acid sequence in SEQ ID NO: 29.

30 Embodiment 35 is the cell of any one of embodiments 1-32, wherein the ALAS protein comprises an amino acid sequence having at least 95% sequence identity to the amino acid sequence in SEQ ID NO: 29.

Embodiment 36 is the cell of any one of embodiments 1-35, wherein the ALAS protein comprises the amino acid sequence in SEQ ID NO: 31.

Embodiment 37 is the cell of any one of embodiments 1-36, wherein the heme-binding protein is selected from the group consisting of a globin, a cytochrome, a cytochrome c oxidase, a ligninase, a catalase, and a peroxidase.

Embodiment 38 is the cell of any one of embodiments 1-36, wherein the heme-binding protein is selected from the group consisting of an androglobin, a chlorocruorin, a
5 cytoglobin, an erythrocrucorin, a flavohemoglobin, a globin E, a globin X, a globin Y, a hemoglobin, a histoglobin, a leghemoglobin, a myoglobin, a neuroglobin, a non-symbiotic hemoglobin, a protoglobin, and a truncated hemoglobin.

Embodiment 39 is the cell of any one of embodiments 1-36, wherein the heme-binding protein is a non-symbiotic hemoglobin.
10

Embodiment 40 is the cell of any one of embodiments 1-36, wherein the heme-binding protein is a leghemoglobin.

Embodiment 41 is the cell of any one of embodiments 1-40, wherein the heme-binding protein comprises an amino acid sequence having at least 90% sequence identity to
15 an amino acid sequence in any one of SEQ ID NOs: 1-27.

Embodiment 42 is the cell of any one of embodiments 1-41, further comprising a third nucleic acid construct comprising a nucleotide sequence encoding a transcription factor, wherein the third nucleic acid construct is operably linked to the first promoter element, the second promoter element, or a third promoter element.

Embodiment 43 is the cell of embodiment 42, wherein the first promoter element
20 comprises a recognition sequence for the transcription factor.

Embodiment 44 is the cell of any one of embodiments 42 or 43, wherein the second exogenous nucleic acid construct is operably linked to a second promoter element, and wherein the second promoter element comprises a recognition sequence for the transcription
25 factor.

Embodiment 45 is the cell any one of embodiments 42 or 43, wherein the third nucleic acid construct is operably linked to the third promoter element, and wherein the third promoter element comprises a recognition sequence for the transcription factor.

Embodiment 46 is the cell of any one of embodiments 1-45, further comprising a
30 fourth nucleic acid construct comprising a nucleotide sequence encoding a protein involved in heme biosynthesis, wherein the fourth nucleic acid construct is operably linked to the first

promoter element, the second promoter element, the third promoter element, or a fourth promoter element.

Embodiment 47 is the cell of embodiment 46, wherein the protein involved in heme biosynthesis is selected from the group consisting of ALA dehydratase, porphobilinogen
5 deaminase, UPG III synthase, UPG III decarboxylase, CPG oxidase, PPG oxidase, and ferrochelatase.

Embodiment 48 is the cell of any one of embodiments 1-47, wherein the first exogenous nucleic acid construct is a heterologous nucleic acid construct.

Embodiment 49 is the cell of any one of embodiments 1-48, wherein the second
10 exogenous nucleic acid construct is a heterologous nucleic acid construct.

Embodiment 50 is the cell of any one of embodiments 1-47, wherein the heme-binding protein is an exogenous heme-binding protein.

Embodiment 51 is the cell of any one of embodiments 1-47 or 50, wherein the heme-binding protein is a heterologous heme-binding protein.

Embodiment 52 is a method of producing a heme-binding protein in a cell
15 comprising:

expressing a first exogenous nucleic acid construct comprising a nucleotide sequence encoding an aminolevulinate synthase (ALAS) protein operably linked to a first promoter element, wherein the ALAS comprises at least a first heme responsive motif
20 (HRM), and wherein the ALAS comprises a mutation in the first HRM; and

expressing a second exogenous nucleic acid construct comprising a nucleotide sequence encoding a heme-binding protein,

wherein the second exogenous nucleic acid construct comprising a nucleotide sequence encoding the heme-binding protein is operably linked to the first promoter element
25 or is operably linked to a second promoter element.

Embodiment 53 is the method of 52, wherein the method produces the heme-binding protein in a titer that is at least 5% greater than a corresponding method lacking the first exogenous nucleic acid construct.

Embodiment 54 is the method of embodiment 52 or embodiment 53, wherein the
30 method produces the heme-binding protein in a titer that is at least 10% greater than a corresponding method lacking the first exogenous nucleic acid construct.

Embodiment 55 is the method of embodiment 52 or embodiment 53, wherein the method produces the heme-binding protein in a titer that is at least 15% greater than a corresponding method lacking the first exogenous nucleic acid construct.

5 Embodiment 56 is the method of embodiment 52 or embodiment 53, wherein the method produces the heme-binding protein in a titer that is at least 20% greater than a corresponding method lacking the first exogenous nucleic acid construct.

Embodiment 57 is the method of embodiment 52, wherein the method produces the heme-binding protein in a titer that is at least 5% greater than a corresponding method lacking the mutation in the first HRM.

10 Embodiment 58 is the method of embodiment 52, wherein the method produces the heme-binding protein in a titer that is at least 10% greater than a corresponding method lacking the mutation in the first HRM.

Embodiment 59 is the method of embodiment 52, wherein the method produces the heme-binding protein in a titer that is at least 15% greater than a corresponding method lacking the mutation in the first HRM.

15 Embodiment 60 is the method of embodiment 52, wherein the method produces the heme-binding protein in a titer that is at least 20% greater than a corresponding method lacking the mutation in the first HRM.

Embodiment 61 is the method of any one of embodiments 52-60, wherein the method is carried out in the absence of added methanol.

Embodiment 62 is the method of any one of embodiments 52-61, wherein the cell is a fungal cell.

Embodiment 63 is the method of embodiment 62, wherein the cell is an *Aspergillus* or *Trichoderma* cell

25 Embodiment 64 is the method of any one of embodiments 62-63, wherein the cell is a yeast cell.

Embodiment 65 is the method of embodiment 64, wherein the yeast cell is a methylotrophic yeast cell.

30 Embodiment 66 is the method of embodiment 65, wherein the methylotrophic yeast cell is a *Pichia* cell, a *Candida* cell, a *Hansenula* cell, or a *Torulopsis* cell.

Embodiment 67 is the method of any one of embodiments 65-66, wherein the methylotrophic yeast cell is a *Pichia methanolica* cell, a *Pichia pastoris* cell, a *Candida boidinii* cell, or a *Hansenula polymorpha* cell.

Embodiment 68 is the method of any one of embodiments 51-53, wherein the
5 methylotrophic yeast cell is a *Pichia pastoris* cell.

Embodiment 69 is the method of any one of embodiments 52-68, wherein the mutation in the first HRM is a mutation from a cysteine to a different amino acid.

Embodiment 70 is the method of any one of embodiments 52-69, wherein the ALAS protein comprises a second HRM, and wherein the ALAS protein comprises a mutation in the
10 second HRM.

Embodiment 71 is the method of embodiment 70, wherein the method produces the heme-binding protein in a titer that is at least 5% greater than a corresponding method lacking the mutations in the first HRM and second HRM.

Embodiment 72 is the method of embodiment 70, wherein the method produces the
15 heme-binding protein in a titer that is at least 10% greater than a corresponding method lacking the mutations in the first HRM and second HRM.

Embodiment 73 is the method of embodiment 70, wherein the method produces the heme-binding protein in a titer that is at least 15% greater than a corresponding method lacking the mutations in the first HRM and second HRM.

Embodiment 74 is the method of embodiment 70, wherein the method produces the
20 heme-binding protein in a titer that is at least 20% greater than a corresponding method lacking the mutations in the first HRM and second HRM.

Embodiment 75 is the method of any one of embodiments 70-74, wherein the mutation in the second HRM is a mutation from a cysteine to a different amino acid.

Embodiment 76 is the method of any one of embodiments 70-75, wherein the
25 different amino acid is the same for the mutation in the first HRM and the mutation in the second HRM.

Embodiment 77 is the method of any one of embodiments 70-75, wherein the different amino acid is not the same for the mutation in the first HRM and the mutation in the
30 second HRM.

Embodiment 78 is the cell of any one of embodiments 70-77, wherein the ALAS protein comprises a third HRM, and wherein the ALAS protein comprises a mutation in the third HRM.

Embodiment 79 is the cell of embodiment 78, wherein the mutation in the third HRM
5 is a mutation from a cysteine to a different amino acid.

Embodiment 80 is the cell of embodiment 79, wherein the different amino acid is the same for the mutation in the first HRM, the mutation in the second HRM, and the mutation in the third HRM.

Embodiment 81 is the method any one of embodiments 69-80, wherein the different
10 amino acid is selected from the group consisting of arginine, histidine, lysine, serine, threonine, asparagine, glutamine, selenocysteine, glycine, proline, alanine, isoleucine, leucine, methionine, aspartic acid, glutamic acid, phenylalanine, tryptophan, tyrosine, and valine.

Embodiment 82 is the method of any one of embodiments 69-80, wherein the
15 different amino acid is selected from a nonpolar aliphatic amino acid, an aromatic amino acid, a polar uncharged amino acid, or a positively charged amino acid.

Embodiment 83 is the method of embodiment 82, wherein the nonpolar aliphatic amino acid is selected from the group consisting of glycine, proline, alanine, isoleucine, leucine, methionine, and valine.

Embodiment 84 is the method of embodiment 82, wherein the aromatic amino acid is
20 selected from the group consisting of phenylalanine, tryptophan, and tyrosine.

Embodiment 85 is the cell of embodiment 82, wherein the polar uncharged amino acid is selected from the group consisting of a polar uncharged amino acid serine, threonine, asparagine, or glutamine.

Embodiment 86 is the method of embodiment 82, wherein the positively charged
25 amino acid is selected from the group consisting of arginine, histidine, and lysine.

Embodiment 87 is the method of any one of embodiments 69-86, wherein the different amino acid is serine.

Embodiment 88 is the method of any one of embodiments 69-86, wherein the
30 different amino acid is alanine.

Embodiment 89 is the method of any one of embodiments 69-86, wherein the different amino acid is phenylalanine.

Embodiment 90 is the method of any one of embodiments 69-86, wherein the different amino acid is histidine.

Embodiment 91 is the cell of any one of embodiments 52-90, wherein the first HRM is HRM1.

5 Embodiment 92 is the cell of any one of embodiments 70-91, wherein the second HRM is HRM2.

Embodiment 93 is the cell of any one of embodiments 52-90, wherein the first HRM is HRM2.

10 Embodiment 94 is the cell of any one of embodiments 70-91, wherein the second HRM is HRM1.

Embodiment 95 is the method of any one of embodiments 52-94, wherein the first exogenous nucleic acid construct comprises a nucleic acid sequence having at least 90% sequence identity to the nucleic acid sequence in SEQ ID NO: 28.

15 Embodiment 96 is the method of any one of embodiments 52-95, wherein the first exogenous nucleic acid construct comprises a nucleic acid sequence having at least 95% sequence identity to the nucleic acid sequence in SEQ ID NO: 28.

Embodiment 97 is the method of any one of embodiments 52-96, wherein the first exogenous nucleic acid construct comprises the nucleic acid sequence in SEQ ID NO: 30.

20 Embodiment 98 is the method of any one of embodiments 52-97, wherein the ALAS protein comprises an amino acid sequence having at least 90% sequence identity to the amino acid sequence in SEQ ID NO: 29.

Embodiment 99 is the method of any one of embodiments 52-98, wherein the ALAS protein comprises an amino acid sequence having at least 95% sequence identity to the amino acid sequence in SEQ ID NO: 29.

25 Embodiment 100 is the method of any one of embodiments 52-99, wherein the ALAS protein comprises the amino acid sequence in SEQ ID NO: 31.

Embodiment 101 is the method of any one of embodiments 52-100, wherein the heme-binding protein is selected from the group consisting of a globin, a cytochrome, a cytochrome c oxidase, a ligninase, a catalase, and a peroxidase.

30 Embodiment 102 is the method of any one of embodiments 52-100, wherein the heme-binding protein is selected from the group consisting of an androglobin, a chlorocruorin, a cytoglobin, an erythrocrucorin, a flavohemoglobin, a globin E, a globin X, a

globin Y, a hemoglobin, a histoglobin, a leghemoglobin, a myoglobin, a neuroglobin, a non-symbiotic hemoglobin, a protoglobin, and a truncated hemoglobin.

Embodiment 103 is the method of any one of embodiments 52-100, wherein the heme-binding protein is a non-symbiotic hemoglobin.

5 Embodiment 104 is the method of any one of embodiments 52-100, wherein the heme-binding protein is a leghemoglobin.

Embodiment 105 is the method of any one of embodiments 52-100, wherein the heterologous heme-binding protein comprises an amino acid sequence having at least 90% sequence identity to an amino acid sequence in any one of SEQ ID NOs: 1-27.

10 Embodiment 106 is the method of any one of embodiments 52-105, further comprising expressing a third nucleic acid construct comprising a nucleotide sequence encoding a transcription factor, wherein the third nucleic acid construct is operably linked to the first promoter element, the second promoter element, or a third promoter element.

Embodiment 107 is the method of embodiment 106, wherein the first promoter
15 element comprises a recognition sequence for the transcription factor.

Embodiment 108 is the method of embodiment 107, wherein the second exogenous nucleic acid construct is operably linked to a second promoter element, and wherein the second promoter element comprises a recognition sequence for the transcription factor.

Embodiment 109 is the method of any one of embodiments 107-108, wherein the
20 third nucleic acid construct is operably linked to the third promoter element, and wherein the third promoter element comprises a recognition sequence for the transcription factor.

Embodiment 110 is the method of any one of embodiments 52-109, further comprising expressing a fourth nucleic acid construct comprising a nucleotide sequence encoding a protein involved in heme biosynthesis, wherein the fourth nucleic acid construct is
25 operably linked to the first promoter element, the second promoter element, the third promoter element, or a fourth promoter element.

Embodiment 111 is the method of embodiment 110, wherein the protein involved in heme biosynthesis is selected from the group consisting of ALA dehydratase, porphobilinogen deaminase, UPG III synthase, UPG III decarboxylase, CPG oxidase, PPG
30 oxidase, and ferrochelatase.

Embodiment 112 is the method of any one of embodiments 52-111, wherein the first exogenous nucleic acid construct is a heterologous nucleic acid construct.

Embodiment 113 is the method of any one of embodiments 52-112, wherein the second exogenous nucleic acid construct is a heterologous nucleic acid construct.

Embodiment 114 is the method of any one of embodiments 52-113, wherein the heme-binding protein is an exogenous heme-binding protein.

5 Embodiment 115 is the method of any one of embodiments 52-111 or 114, wherein the heme-binding protein is a heterologous heme-binding protein.

EXAMPLES

10

Example 1

Polymerase Chain Reaction

15

Genes of interest were amplified from genomic DNA or plasmid DNA templates using Phusion High-fidelity PCR master mix (New England Biolabs, Cat #M0531), 0.6 μ M each of forward and reverse primers and 10-50 ng of template DNA. The reaction conditions were as follows:

1 cycle	Initial Denaturation	98°C	1min
25 cycles	Denaturation	98°C	10 sec
	Annealing	60°C	20 sec
	Extension	72°C	30 sec per kb
1 cycle	Final Extension	72°C	5min
	Hold	4°C	Forever

20

Pichia pastoris ALAS gene (KEGG identifier PAS_chr2-1_0716, which can be accessed at the Kyoto Encyclopedia of Genes and Genomes (KEGG) website) was amplified using primers (ATGGAGTTTGTGCGCCCGTCAG; SEQ ID NO: 65) and (CTACAATCTGACTCCTGATGAGGTTTC; SEQ ID NO: 66) from genomic DNA. In the examples, wtALAS denotes this native sequence of ALAS (SEQ ID NO: 28).

25

Example 2

Cloning of ALAS and mutagenesis

PCR product was purified using NucleoSpin Gel and PCR Clean-Up (Takara Bio, Cat #740609) and cloned in pCR-BluntII-TOPO vector using Zero Blunt TOPO PCR Cloning Kit (Thermo Fisher Scientific, Cat #K280020) following the manufacturer's recommendation.

Site directed mutagenesis was carried out on the resulting vector (40 ng of purified plasmid) using two sets of primers (set1:

CGTCAGTCCATGAATGCCTCTCCCTTTGTCAGGTCAACTTC; SEQ ID NO: 37 and

GAAGTTGACCTGACAAAGGGAGAGGCATTCATGGACTGACG; SEQ ID NO: 38,

set2: GCTGCTACTGCTAGTCATTCTCCCGTGGTTGGCCCTG; SEQ ID NO: 39 and

CAGGGCCAACCAACGGGAGAATGACTAGCAGTAGCAGC; SEQ ID NO: 40) and

QuikChange II XL Site-Directed Mutagenesis Kit (Agilent, Cat #200521) to create mutALAS carrying mutations of cysteine residues at positions 12 and 39 to serine residues (SEQ ID NO: 30). In the Examples, mutALAS specifically denotes these two mutations unless specified otherwise. The reaction conditions were as follows:

1 cycle	Initial Denaturation	95°C	1 min
18 cycles	Denaturation	95°C	50 sec
	Annealing	60°C	50 sec
	Extension	68°C	11 min
1 cycle	Final Extension	68°C	7 min
	Hold	4°C	Forever

Single cysteine to serine mutation, C12S or C39S was performed by site directed mutagenesis on vector carrying wtALAS using primer set1 (SEQ ID NO: 37, SEQ ID NO: 38) and set2 (SEQ ID NO: 39, SEQ ID NO: 40) respectively.

Example 3

Construction of ALAS (wild-type or mutated, C12S, C39S) gene integration cassette

Integration cassette of wtALAS or mutALAS was designed such that gene integration occurred under the methanol-inducible alcohol oxidase 1 (AOX1) promoter element from *Pichia pastoris* and before the translation stop signal immediately followed by the transcription terminator sequence from the *Pichia pastoris* FDH1 gene. The linear constructs

contained the 3' half of the promoter element, followed by ALAS gene (wild-type or mutant), followed by the FDH1 transcription terminator. This was immediately followed by the selection cassette containing the pTEF promoter element from *Ashbya gossypii*, the acetamidase gene (*amdS*) from *Aspergillus nidulans* and the TEF terminator from *Ashbya gossypii*. Finally, the construct contained the 5' half of the promoter element (See, e.g., Figure 5).

Overlapping PCR was used to create the linear constructs [3'pAOX1-ALAS (wt/mut)-FDH1tt-pTEF-*amdS*-TEFtt-5'pAOX1). Primers Used to Amplify the Linear Constructs:

Overlapping PCR	Primer sequences	SEQ ID NO:
PCR1	AAACGCTGTCTTGGAACCTAATATGAC	41
	GACGGGCGACAACTCCATCGTTTCGAATAATTAGTTG	42
PCR2	CAACTAATTATTCGAAACGATGGAGTTTGTCGCCCCGTCAG	43
	AATTAAATACATTTCAACTACAATCTGACTCCTGATGAGGTTTCG	44
PCR3	CCTCATCAGGAGTCAGATTGTAGTTGAAATGTATTTAATTTG	45
	AAACTGTCAGTTTTGGGCCATTTG	46

Individual PCR reaction conditions were performed as indicated above. PCR products were purified using NucleoSpin Gel and PCR Clean-Up (Takara Bio, Cat #740609). The final overlapping PCR product was performed using 0.5 U Platinum Pfx DNA Polymerase (Thermo Fisher Scientific, Cat#11708039), 1x amplification buffer, 0.3 mM dNTPs and 1 mM MgCl₂ by mixing the three purified PCR amplicons in an equimolar ratio in a 2-part PCR reaction. The PCR conditions were:

Part 1:

1 cycle	Initial Denaturation	94°C	2min
15 cycles	Denaturation	94°C	15 sec
	Annealing	58°C	30 sec
	Extension	68°C	5 min
	Hold	4°C	

Following part 1, the reaction mixes were spiked with primers AAACGCTGTCTTGGAACCTAATATGAC (SEQ ID NO: 41) and AAACGCTGTCTTGGAACCTAATATGAC (SEQ ID NO: 41) and AAACGCTGTCTTGGAACCTAATATGAC (SEQ ID NO: 41) and AAACGCTGTCTTGGAACCTAATATGAC (SEQ ID NO: 41) (0.3 uM final concentration) and 1.5U Pfx and continued to part 2.

Part 2:

1 cycle	Initial Denaturation	94°C	2min
20 cycles	Denaturation	94°C	15 sec
	Annealing	58°C	30 sec
	Extension	68°C	5 min
	Hold	4°C	Forever

Sequences of exemplary mutant nucleic acids (e.g., SEQ ID NO: 30) and proteins (e.g., SEQ ID NO: 31) are provided in Figure 7.

Expression plasmids of wtALAS and mutALAS were constructed in an autonomous replicating vector (panARS) under a modified pAOX1 followed by the transcription terminator sequence from the *Pichia pastoris* FDH1 gene. The vector conferred resistance to G418 (Geneticin). Cloning of inserts GFP, wtALAS and mutALAS in the vector was performed using Gibson Assembly Master Mix (New England Biolabs, catalog # E2611L) following manufacturer's recommendation.

Example 4

Preparation of *P. pastoris* transformation-competent cells

Selected strains of *P. pastoris* (*K. phaffii*) were grown to mid-exponential growth phase (~2 OD) in 25 ml YPD medium. Cells were collected by centrifugation at 930×g for 15 minutes. The cell pellet was resuspended in 2 ml of a solution of 80% YPD and 200 mM HEPES, pH 6.8. 75 µl of 1 M DTT was added. The resuspended cell pellet was mixed at 100 rpm at 30°C for 25 minutes. A 40 ml volume of ice cold, sterile water was added to the suspension, and the cells were collected by centrifugation at 1125×g for 15 minutes and placed on ice. The cell pellet was resuspended in 40 ml ice cold water and collected as before for two additional wash steps. The cell pellet was then resuspended in 20 ml of ice cold 1 M sorbitol and collected by centrifugation as before. The final cell pellet was suspended in 0.3 ml ice cold, sterile 1 M sorbitol, aliquoted and frozen at -80° C.

Example 5

Transformation into *P. pastoris*

For genome integration, 100-300 ng of linearized DNA was transformed into 30 µl of electrocompetent *P. pastoris* cells using a 1 mm gap GenePulser cuvette (Bio-Rad, Cat# 1652083) with a GenePulser (Bio-Rad) set at 1.15 kV. 1 ml of YPD/1M sorbitol (1:1 vol/vol) was added immediately to the cells. The cells were allowed to recover for 3 h at 30°C with shaking at 100 rpm. 100 µl of the recovery mixture was plated on yeast carbon base plates containing 5mM acetamide (Teknova, Cat#Y5216). Plates were incubated at 30° C for 48 hours. Individual clones were streaked onto yeast carbon base plates containing acetamide to obtain single colonies and the isolated colonies were used to do colony PCR or gDNA prep to confirm gene integration into the chromosome and sequence the integration construct.

For plasmid DNA transformation, the same steps were followed as for genome integration except that after electroporation and recovery of cells, 50-100 µl of the recovery mixture was plated on YPD plates containing 300 ug/ml G418 (Geneticin).

Example 6

Construction of Strains St2, St3, St5 and St6

A high yielding parent strain (St1) had pre-existing recombinant ALAS under the methanol-inducible strong promoter pAOX1 (alcohol oxidase 1) in addition to other pAOX1-driven heme enzymes, the carbon-responsive transcription factor MxR1 and multiple copies of LegH. A low yielding parent LegH strain (St4) lacked recombinant MxR1, ALAS and other heme enzymes except aminolevulinate dehydratase (ALAD).

Competent St1 cells (Table 1; a high LegH titer strain) were transformed with each of the linear cassettes (for wtALAS and mutALAS) and transformants containing the amdS selection cassette were selected based on their ability to grow on agar plates containing acetamide as the sole nitrogen source. The resulting strains (St2 and St3 with integrated cassette for wtALAS and mutALAS respectively, Table 1) were purified, isolated and the presence of pAOX1 driven wtALAS or mutALAS was verified by colony PCR and sequencing. Similarly, competent St4 cells (a low LegH titer strain) were transformed with linear cassettes for wtALAS and mutALAS to obtain St5 and St6 respectively.

Table 1. Strains with LegH integrated in genome

Parent	1 extra copy of recombinant wtALAS added	1 extra copy of recombinant mutALAS added
High LegH titer strain (Strain St1)	Strain St2	Strain St3
Low LegH titer strain (Strain St4)	Strain St5	Strain St6

Example 7***ALAS* gene copy number analysis**

ALAS gene copy number in different strains was measured by probe-based qPCR. Briefly, genomic DNA (20ng) was amplified with 1x PrimeTime gene expression mastermix (Integrated DNA Technologies, Cat# 1055770) and PrimeTime qPCR Probe Assays in a real-time qPCR CFX96 machine (Bio-Rad). qPCR normalization was performed with respect to actin. A double delta Ct analysis method was followed to calculate the relative copy number of gene of interest between strains. Primer and probe sequences were:

Gene	Primer sequence	SEQ ID. NO.	Probe sequence	SEQ ID. NO.
ALAS	GCTCTCCAACAGCAGA GATAC	47	6-FAM/AAGCCCAAA/Zen/CC TCCGACATTGCTA/3IABkF Q	51
	GTCCATACGGATCGGA GAAAC	48		
Actin	AGCAACATCCCTGATT CCG	49	HEX/TCGCCGTAA/Zen/GTT CTTGTT TAGACGTTC/3IABkFQ	52
	ATGCGTACCTTCAATCC TGG	50		

The engineered strains (Table 1) contained identical number of extra ALAS gene copies, either one copy of wtALAS or mutALAS, as measured by qPCR of ALAS normalized to actin levels. Hence, ALAS gene dosage was ruled out for any difference in the phenotype of the resulting strains (e.g., St2 vs St3 and St5 vs St6).

Example 8

PCR detection of heme enzyme, MxR1, LegH and Mb genes

Strain characterization was done by PCR for pAOX1-driven recombinant heme biogenesis pathway enzyme, MxR1 and LegH genes. Forward primer sequence for PCR was TAGCGCAGTCTCTCTATCGCTTC (SEQ ID NO: 53) specific to pAOX1. Reverse primer was specific to each gene of interest as shown below.

Gene	Reverse Primer	Amplicon size	SEQ ID NO:
5-aminolevulinate synthase	CACTGGGTTGTGCACATTGG	1995 bp	54
Delta-aminolevulinate dehydratase	ACAATATTCTTCTCTGCCGC	1268 bp	55
Phorphobilinogen deaminase	TTGATCTCGTCAAGAATGCG	1358 bp	56
Uroporphyrinogen III synthase	TAGGTGCCACAACCTTTTGTTTC	1102 bp	57
Uroporphyrinogen decarboxylase	GATCCAATGCGATGACATTCTTGT	1430 bp	58
Coproporphyrinogen III oxidase	ACCTGCAATAACTCCTCTTCTCTG	1301 bp	59
Protoporphyrinogen oxidase	CCACTGAGGGTAGCCGAATC	2027 bp	60
Ferrochelatase	GGGCTCTGAAAACTCTTTTGG	1486 bp	61
MxR1	GCATGTCTCAATAACAGATCTCGACGG	629 bp	62
LegH	AAGCCTCTTGTTTTTCTGTAAATGCAC	382 bp	63
Bovine Mb	TGATGGCGTCCGAGATGAACTC	688 bp	64

Results of these reactions are shown below.

Recombinant Gene (under pAOX1)	St1	St2	St3	St4	St5	St6	St7
5-aminolevulinate synthase	x	x	x	-	x	x	x
Delta-aminolevulinate dehydratase	x	x	x	x	x	x	x
Phorphobilinogen deaminase	x	x	x	-	-	-	x
Uroporphyrinogen III synthase	x	x	x	-	-	-	x
Uroporphyrinogen decarboxylase	x	x	x	-	-	-	x
Coproporphyrinogen III oxidase	x	x	x	-	-	-	x
Protoporphyrinogen oxidase	x	x	x	-	-	-	x
Ferrochelatase	x	x	x	-	-	-	x
MxR1	x	x	x	-	-	-	x

LegH	X	X	X	X	X	X	-
Mb (Bovine Myoglobin)	-	-	-	-	-	-	X

Example 9

5

Shake flask cultivation of strains St4, St5 and St6

The strains were inoculated into growth media (1% yeast extract, 2% peptone, supplemented with 1% glycerol) overnight at 30°C with shaking at 200 rpm. The next day the overnight cultures were diluted to an OD₆₀₀ of 0.5-0.7 with YP media supplemented with 1% methanol and 1% dextrose. The cultures were grown for 48 hours and harvested by centrifugation at ~4000g for 15 mins at 4°C.

The low LegH titer strain, St4, didn't have a pre-existing copy of recombinant ALAS, unlike St1. Integrating a copy of mutALAS (St6) led to a ~30% improvement in LegH titer compared to St5 (wtALAS). The titer calculation is based on LegH content as measured by a liquid chromatography method as described in Example 13.

The relative LegH titer is shown in Table 2.

Table 2

	St5	St6
Relative LegH titer	1.00	1.32

20

Example 10

2L cultivation of strains St1, St2 and St3

Strains St1, St2 and St3 were grown in 2L fermentation tanks in media containing dextrose as the principal carbon source at 30°C. No methanol was used. In the background of a high LegH titer strain (St1) that contained pre-existing recombinant ALAS, St3 overexpressing mutALAS improved LegH titer by >30% over the parent St1. When compared to wtALAS (in St2), mutALAS (in St3) resulted in a 20% improvement in LegH

titer. The titer calculation is based on LegH content as measured by a liquid chromatography method as described in Example 13.

The relative LegH titer is shown in Table 3.

Table 3

	St1	St2	St3
Relative LegH titer	1.00	1.10	1.37

Hence, mutating both the ALAS HRMs and overexpressing the mutated ALAS (mutALAS) in Pichia strains with different LegH titer led to further improvement in LegH titer. Furthermore, this suggested that mutALAS improved LegH titer independent of the strain genetic makeup (MxR1 and heme enzymes other than ALAS), and presence of methanol.

Example 11

mutALAS improved heme levels

A quantitative assay for total heme based on reversed-phase high-performance liquid chromatography indicated that strains with mutALAS accumulated more heme levels than wtALAS containing strains. Shown in Table 4 is heme quantification in strain St3 (mutALAS) vs strain St2 (wtALAS). Additionally, mutALAS increased heme loading in multiple strains.

Table 4

	St2	St3
Relative heme titer	1.00	2.7

Example 12

ALAS protein levels with mutALAS

It is generally believed that heme regulates ALAS levels in a feedback fashion to regulate its own levels. At the protein level, the ALAS level was higher 3-fold in St3 (mutALAS) compared to St2 (wtALAS) (Table 5) as quantitated by shotgun mass spectrometry, when the strains were grown in 2L fermenter tanks with dextrose.

Table 5

	St2	St3

Relative ALAS titer	1.00	3.3
---------------------	------	-----

Example 13

Quantification of Leghemoglobin

Cell broth samples were pelleted down (at 4000 x g, 4°C, 30 min) and decanted. The pellet samples were then diluted four times with lysis buffer (150 mM NaCl, 50mM Potassium Phosphate, pH 7.4). 300 uL of each resuspension was dispensed into a 96 well deep plate with 120 uL of beads (Zirconium/silica beads (0.5 mm)) per well for cell lysis. The lysis was done with a mini bead beater for 3 minutes, then the plate was cooled down on ice for 5 minutes, and followed with another 2 minutes of bead beating. The plate was then spun down (at 4000 x g, 4°C, 30 min). The supernatant was filtered through a 0.2 um filter plate (at 4000 x g, 4°C, 60 min).

The filtered lysate was loaded onto a UHPLC with a size-exclusion column (Acquity BEH SEC column, 200 Å, 1.7 um, 4.6 x 150 mm). Method parameters: 1) Mobile phase: 5mM NaCl, 50mM Potassium Phosphate, (pH 7.4); 2) Flow rate: 0.3 mL/min; 3) Injection volume: 10 uL; 4) Run time: 15 min; 5) Sample tray temperature: 4C. A calibration curve was built with a purified LegH standard using absorbance at 280 nm and 415 nm. The quantification was done using peak area with valley-to-valley peak integration method. The absorbance at 280 nm is proportional to the amount of the polypeptide present and the absorbance at 415 nm is proportional to the amount of heme present. Where a peak is seen at the same elution time at both wavelengths, a heme containing protein is detected.

Example 14

mutALAS improved levels of bovine myoglobin

In addition to LegH, the production of bovine myoglobin was evaluated. Strain, St7, was generated by integrating myoglobin cDNA of *Bos taurus* (NM_173881.2) in a strain containing recombinant copies of heme enzymes and MxR1 (St7 characterization is described in Example 8) integrated under pAOX1.

Three expression plasmids for expression of GFP (control), wtALAS and mutALAS were constructed as described in Example 3. Three strains, St8, St9 and St10 were generated using a method as described in Example 5 by transforming these three plasmids

overexpressing GFP, wtALAS and mutALAS respectively in strain S7, as shown in Table 6. Episomal expression of wtALAS or mutALAS was obtained by growing transformants using a method described in Example 9, except that the growth media was supplemented with 300 ug/ml G418.

5

Table 6

	Plasmid expressing	Host strain	Host strain expressing	Resulting Strain
1	GFP	St7	Mb	St8
2	wtALAS			St9
3	mutALAS (C12S, C39S)			St10

The cells were pelleted, and shotgun proteomics by LC-MS was performed to quantitate Mb. When normalized to the myoglobin level in Strain St8 (GFP), the average myoglobin levels in St9 and St10 were as shown in Table 7.

10

Table 7

Strains	Normalized Myoglobin levels
St8 (GFP)	1
St9 (wt ALAS)	1.60
St10 (mutALAS)	2.58

15

Example 15**Mutations to other amino acids**

In addition to cysteine to serine mutations, additional mutations were evaluated. Expression plasmids of mutALAS variants were constructed by gene synthesis and cloning in an autonomous replicating vector (panARS) under a modified pAOX1 followed by the FDH1 transcription terminator sequence. Furthermore, these plasmids were transformed in strain St1, using a method as described in Example 5 and the resulting transformants produced

20

LegH when cultured, using a method described in Example 9, except that the growth media was supplemented with 300 ug/ml G418. The titer of LegH produced by these strains was normalized to the titer of LegH produced by wtALAS, as determined by the method described in Example 13. The results are shown in Table 8.

5

Table 8

Variant	Normalized LegH titer fold
C12, C39 (wtALAS)	1
C12S, C39S (mutALAS)	1.14
C12A, C39A	1.22
C12D, C39D	0.91
C12F, C39F	1.17
C12H, C39H	1.12

Example 16**Single cysteine to serine mutants in *Pichia pastoris***

10

In addition to evaluating the double mutant (C12S, C39S) of ALAS in *P. pastoris*, single cysteine to serine mutants were also evaluated. Single mutations were created in wtALAS by site directed mutagenesis as described in Example 2. Expression plasmids for expression of wtALAS, mutALAS and single ALAS mutants (C12S and C39S) were constructed as described in Example 3 and were transformed in strain St1, using a method as described in Example 5. The resulting transformants produced LegH when cultured, using a method described in Example 9, except that the growth media was supplemented with 300 ug/ml G418. The titer of LegH produced by these strains was normalized to the titer of LegH

15

produced by wtALAS, as determined by the method described in Example 13. The results are shown in Table 9.

Table 9

Variant	Normalized LegH titer fold
C12, C39 (wtALAS)	1
C12S, C39S	1.27
C12S	1.03
C39S	1.01

5

OTHER EMBODIMENTS

10 It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

WHAT IS CLAIMED IS:

1. A cell comprising:
 - a first exogenous nucleic acid construct comprising a nucleotide sequence encoding an aminolevulinate synthase (ALAS) protein operably linked to a first promoter element, wherein the ALAS comprises at least a first heme responsive motif (HRM), and wherein the ALAS comprises a mutation in the first HRM; and
 - a second exogenous nucleic acid construct comprising a nucleotide sequence encoding a heme-binding protein,
 - wherein the second exogenous nucleic acid construct comprising a nucleotide sequence encoding the heme-binding protein is operably linked to the first promoter element or is operably linked to a second promoter element.
2. The cell of claim 1, wherein the cell is a fungal cell, or is a yeast cell.
3. The cell of claim 2, wherein yeast cell is a *Pichia pastoris* cell.
4. The cell of any one of claims 1-3, wherein the mutation in the first HRM is a mutation from a cysteine to a different amino acid.
5. The cell of any one of claims 1-4, wherein the ALAS protein comprises a second HRM, and wherein the ALAS protein comprises a mutation in the second HRM.
6. The cell of claim 5, wherein the mutation in the second HRM is a mutation from a cysteine to a different amino acid.
7. The cell of claim 5, wherein the different amino acid is the same for the mutation in the first HRM and the mutation in the second HRM.
8. The cell of claim 5, wherein the different amino acid is not the same for the mutation in the first HRM and the mutation in the second HRM.

9. The cell of any one of claims 4-8, wherein the different amino acid is selected from the group consisting of arginine, histidine, lysine, serine, threonine, asparagine, glutamine, selenocysteine, glycine, proline, alanine, isoleucine, leucine, methionine, glutamic acid, aspartic acid, phenylalanine, tryptophan, tyrosine, and valine.
- 5
10. The cell of any one of claims 4-8, wherein the different amino acid is selected from a nonpolar aliphatic amino acid, an aromatic amino acid, a polar uncharged amino acid, or a positively charged amino acid.
- 10
11. The cell of any one of claims 4-9, wherein the different amino acid is serine.
12. The cell of any one of claims 4-9, wherein the different amino acid is alanine.
13. The cell of any one of claims 4-9, wherein the different amino acid is
- 15 phenylalanine.
14. The cell of any one of claims 4-9, wherein the different amino acid is histidine.
- 20
15. The cell of any one of claims 1-14, wherein the first exogenous nucleic acid construct comprises a nucleic acid sequence having at least 90% sequence identity to the nucleic acid sequence in SEQ ID NO: 28.
16. The cell of any one of claims 1-15, wherein the ALAS protein comprises an
- 25 amino acid sequence having at least 90% sequence identity to the amino acid sequence in SEQ ID NO: 29.
17. The cell of any one of claims 1-16, wherein the heme-binding protein is selected from the group consisting of a globin, a cytochrome, a cytochrome c oxidase,
- 30 a ligninase, a catalase, and a peroxidase.

18. The cell of any one of claims 1-16, wherein the heme-binding protein is a leghemoglobin.

19. The cell of any one of claims 1-16, wherein the heme-binding protein
5 comprises an amino acid sequence having at least 90% sequence identity to an amino acid sequence in any one of SEQ ID NOs: 1-27.

20. A method of producing a heme-binding protein in a cell comprising:
expressing a first exogenous nucleic acid construct comprising a nucleotide
10 sequence encoding an aminolevulinate synthase (ALAS) protein operably linked to a first promoter element, wherein the ALAS comprises at least a first heme responsive motif (HRM), and wherein the ALAS comprises a mutation in the first HRM; and
expressing a second exogenous nucleic acid construct comprising a nucleotide sequence encoding a heme-binding protein,
15 wherein the second exogenous nucleic acid construct comprising a nucleotide sequence encoding the heme-binding protein is operably linked to the first promoter element or is operably linked to a second promoter element.

21. The method of claim 20, wherein the method produces the heme-binding
20 protein in a titer that is at least 5% greater than a corresponding method lacking the first exogenous nucleic acid construct.

22. The method of claim 20 or claim 21, wherein the method produces the heme-binding protein in a titer that is at least 10% greater than a corresponding method
25 lacking the first exogenous nucleic acid construct.

23. The method of claim 20, wherein the method produces the heme-binding protein in a titer that is at least 5% greater than a corresponding method lacking the mutation in the first HRM.

30

24. The method of claim 20, wherein the method produces the heme-binding protein in a titer that is at least 10% greater than a corresponding method lacking the mutation in the first HRM.

5 25. The method of any one of claims 20-24, wherein the method is carried out in the absence of added methanol.

26. The method of any one of claims 20-25, wherein the mutation in the first HRM is a mutation from a cysteine to a different amino acid.

10

27. The method of any one of claims 20-26, wherein the ALAS protein comprises a second HRM, and wherein the ALAS protein comprises a mutation in the second HRM.

15 28. The method of claim 26, wherein the method produces the heme-binding protein in a titer that is at least 5% greater than a corresponding method lacking the mutations in the first HRM and second HRM.

20 29. The method of claim 26, wherein the method produces the heme-binding protein in a titer that is at least 10% greater than a corresponding method lacking the mutations in the first HRM and second HRM.

30. The method of any one of claims 27-29, wherein the mutation in the second HRM is a mutation from a cysteine to a different amino acid.

25

31. The method of claim 30, wherein the different amino acid is the same for the mutation in the first HRM and the mutation in the second HRM.

30 32. The method of claim 30, wherein the different amino acid is not the same for the mutation in the first HRM and the mutation in the second HRM.

33. The method any one of claims 26-32, wherein the different amino acid is selected from the group consisting of arginine, histidine, lysine, serine, threonine, asparagine, glutamine, selenocysteine, glycine, proline, alanine, isoleucine, leucine, methionine, aspartic acid, glutamic acid, phenylalanine, tryptophan, tyrosine, and
5 valine.
34. The method of any one of claims 26-33, wherein the different amino acid is serine.
- 10 35. The method of any one of claims 26-33, wherein the different amino acid is alanine.
36. The method of any one of claims 26-33, wherein the different amino acid is phenylalanine.
15
37. The method of any one of claims 26-33, wherein the different amino acid is histidine.
38. The method of any one of claims 20-37, wherein the first exogenous nucleic
20 acid construct comprises a nucleic acid sequence having at least 90% sequence identity to the nucleic acid sequence in SEQ ID NO: 28.
39. The method of any one of claims 20-38, wherein the ALAS protein comprises an amino acid sequence having at least 90% sequence identity to the amino acid
25 sequence in SEQ ID NO: 29.
40. The method of any one of claims 20-39, wherein the heme-binding protein is selected from the group consisting of a globin, a cytochrome, a cytochrome c oxidase, a ligninase, a catalase, and a peroxidase.
30
41. The method of any one of claims 20-39, wherein the heme-binding protein is a leghemoglobin.

42. The method of any one of claims 20-39, wherein the heterologous heme-binding protein comprises an amino acid sequence having at least 90% sequence identity to an amino acid sequence in any one of SEQ ID NOs: 1-27.

5




Figure 1




Figure 2




Figure 3

Figure 4

<110> Impossible Foods Inc.

<210> 1

<211> 161

<212> PRT

<213> *Vigna radiata*

<400> 1

Met	Thr	Thr	Thr	Leu	Glu	Arg	Gly	Phe	Thr	Glu	Glu	Gln	Glu	Ala	Leu	
1				5					10					15		
Val	Val	Lys	Ser	Trp	Asn	Val	Met	Lys	Lys	Asn	Ser	Gly	Glu	Leu	Gly	
			20					25					30			
Leu	Lys	Phe	Phe	Leu	Lys	Ile	Phe	Glu	Ile	Ala	Pro	Ser	Ala	Gln	Lys	
		35					40					45				
Leu	Phe	Ser	Phe	Leu	Arg	Asp	Ser	Thr	Val	Pro	Leu	Glu	Gln	Asn	Pro	
	50					55				60						
Lys	Leu	Lys	Pro	His	Ala	Val	Ser	Val	Phe	Val	Met	Thr	Cys	Asp	Ser	
65					70					75					80	
Ala	Val	Gln	Leu	Arg	Lys	Ala	Gly	Lys	Val	Thr	Val	Arg	Glu	Ser	Asn	
				85					90					95		
Leu	Lys	Lys	Leu	Gly	Ala	Thr	His	Phe	Arg	Thr	Gly	Val	Ala	Asn	Glu	
			100					105					110			
His	Phe	Glu	Val	Thr	Lys	Phe	Ala	Leu	Leu	Glu	Thr	Ile	Lys	Glu	Ala	
		115					120					125				
Val	Pro	Glu	Met	Trp	Ser	Pro	Ala	Met	Lys	Asn	Ala	Trp	Gly	Glu	Ala	
	130					135					140					
Tyr	Asp	Gln	Leu	Val	Asp	Ala	Ile	Lys	Tyr	Glu	Met	Lys	Pro	Pro	Ser	
145					150					155					160	
Ser																

<210> 2

<211> 133

<212> PRT

<213> *Methylobacterium radiatum*

<400> 2

Met	Ile	Asp	Gln	Lys	Glu	Lys	Glu	Leu	Ile	Lys	Glu	Ser	Trp	Lys	Arg	
1				5					10					15		
Ile	Glu	Pro	Asn	Lys	Asn	Glu	Ile	Gly	Leu	Leu	Phe	Tyr	Ala	Asn	Leu	
			20					25					30			
Phe	Lys	Glu	Glu	Pro	Thr	Val	Ser	Val	Leu	Phe	Gln	Asn	Pro	Ile	Ser	
		35					40					45				
Ser	Gln	Ser	Arg	Lys	Leu	Met	Gln	Val	Leu	Gly	Ile	Leu	Val	Gln	Gly	

50						55					60					
Ile	Asp	Asn	Leu	Glu	Gly	Leu	Ile	Pro	Thr	Leu	Gln	Asp	Leu	Gly	Arg	
65					70					75					80	
Arg	His	Lys	Gln	Tyr	Gly	Val	Val	Asp	Ser	His	Tyr	Pro	Leu	Val	Gly	
			85					90					95			
Asp	Cys	Leu	Leu	Lys	Ser	Ile	Gln	Glu	Tyr	Leu	Gly	Gln	Gly	Phe	Thr	
		100						105				110				
Glu	Glu	Ala	Lys	Ala	Ala	Trp	Thr	Lys	Val	Tyr	Gly	Ile	Ala	Ala	Gln	
		115					120					125				
Val	Met	Thr	Ala	Glu												
130																

<210> 3

<211> 139

<212> PRT

<213> Aquifex aeolicus

<400> 3

Met	Leu	Ser	Glu	Glu	Thr	Ile	Arg	Val	Ile	Lys	Ser	Thr	Val	Pro	Leu	
1				5					10					15		
Leu	Lys	Glu	His	Gly	Thr	Glu	Ile	Thr	Ala	Arg	Met	Tyr	Glu	Leu	Leu	
			20					25					30			
Phe	Ser	Lys	Tyr	Pro	Lys	Thr	Lys	Glu	Leu	Phe	Ala	Gly	Ala	Ser	Glu	
		35					40					45				
Glu	Gln	Pro	Lys	Lys	Leu	Ala	Asn	Ala	Ile	Ile	Ala	Tyr	Ala	Thr	Tyr	
50						55					60					
Ile	Asp	Arg	Leu	Glu	Glu	Leu	Asp	Asn	Ala	Ile	Ser	Thr	Ile	Ala	Arg	
65				70						75					80	
Ser	His	Val	Arg	Arg	Asn	Val	Lys	Pro	Glu	His	Tyr	Pro	Leu	Val	Lys	
			85					90					95			
Glu	Cys	Leu	Leu	Gln	Ala	Ile	Glu	Glu	Val	Leu	Asn	Pro	Gly	Glu	Glu	
		100						105					110			
Val	Leu	Lys	Ala	Trp	Glu	Glu	Ala	Tyr	Asp	Phe	Leu	Ala	Lys	Thr	Leu	
		115					120					125				
Ile	Thr	Leu	Glu	Lys	Lys	Leu	Tyr	Ser	Gln	Pro						
130						135										

<210> 4

<211> 145

<212> PRT

<213> Glycine max

<400> 4

Met	Gly	Ala	Phe	Thr	Glu	Lys	Gln	Glu	Ala	Leu	Val	Ser	Ser	Ser	Phe	
1				5					10					15		
Glu	Ala	Phe	Lys	Ala	Asn	Ile	Pro	Gln	Tyr	Ser	Val	Val	Phe	Tyr	Thr	
			20					25					30			

```

Ser Ile Leu Glu Lys Ala Pro Ala Ala Lys Asp Leu Phe Ser Phe Leu
      35              40              45
Ser Asn Gly Val Asp Pro Ser Asn Pro Lys Leu Thr Gly His Ala Glu
      50              55              60
Lys Leu Phe Gly Leu Val Arg Asp Ser Ala Gly Gln Leu Lys Ala Asn
      65              70              75              80
Gly Thr Val Val Ala Asp Ala Ala Leu Gly Ser Ile His Ala Gln Lys
      85              90              95
Ala Ile Thr Asp Pro Gln Phe Val Val Val Lys Glu Ala Leu Leu Lys
      100             105             110
Thr Ile Lys Glu Ala Val Gly Asp Lys Trp Ser Asp Glu Leu Ser Ser
      115             120             125
Ala Trp Glu Val Ala Tyr Asp Glu Leu Ala Ala Ala Ile Lys Lys Ala
      130             135             140
Phe
145

```

```

<210> 5
<211> 162
<212> PRT
<213> Hordeum vulgare

```

```

<400> 5
Met Ser Ala Ala Glu Gly Ala Val Val Phe Ser Glu Glu Lys Glu Ala
  1              5              10              15
Leu Val Leu Lys Ser Trp Ala Ile Met Lys Lys Asp Ser Ala Asn Leu
      20              25              30
Gly Leu Arg Phe Phe Leu Lys Ile Phe Glu Ile Ala Pro Ser Ala Arg
      35              40              45
Gln Met Phe Pro Phe Leu Arg Asp Ser Asp Val Pro Leu Glu Thr Asn
      50              55              60
Pro Lys Leu Lys Thr His Ala Val Ser Val Phe Val Met Thr Cys Glu
      65              70              75              80
Ala Ala Ala Gln Leu Arg Lys Ala Gly Lys Ile Thr Val Arg Glu Thr
      85              90              95
Thr Leu Lys Arg Leu Gly Gly Thr His Leu Lys Tyr Gly Val Ala Asp
      100             105             110
Gly His Phe Glu Val Thr Arg Phe Ala Leu Leu Glu Thr Ile Lys Glu
      115             120             125
Ala Leu Pro Ala Asp Met Trp Gly Pro Glu Met Arg Asn Ala Trp Gly
      130             135             140
Glu Ala Tyr Asp Gln Leu Val Ala Ala Ile Lys Gln Glu Met Lys Pro
      145             150             155             160
Ala Glu

```

```

<210> 6

```

Met 1	Asp	Gly	Ala	Val 5	Arg	Leu	Asp	Trp	Thr 10	Gly	Leu	Asp	Leu	Thr 15	Gly
His	Glu	Ile	His 20	Asp	Gly	Val	Pro	Ile 25	Ala	Ser	Arg	Val	Gln 30	Val	Met
Val	Ser	Phe 35	Pro	Leu	Phe	Lys	Asp 40	Gln	His	Ile	Ile	Met 45	Ser	Ser	Lys
Glu 50	Ser	Pro	Ser	Arg	Lys	Ser 55	Ser	Thr	Ile	Gly	Gln 60	Ser	Thr	Arg	Asn
Gly 65	Ser	Cys	Gln	Ala	Asp 70	Thr	Gln	Lys	Gly	Gln 75	Leu	Pro	Pro	Val	Gly 80
Glu	Lys	Pro	Lys	Pro 85	Val	Lys	Glu	Asn 90	Pro	Met	Lys	Lys	Leu	Lys 95	Glu
Met	Ser	Gln	Arg 100	Pro	Leu	Pro	Thr	Gln 105	His	Gly	Asp	Gly	Thr 110	Tyr	Pro
Thr	Glu	Lys 115	Lys	Leu	Thr	Gly	Ile 120	Gly	Glu	Asp	Leu	Lys 125	His	Ile	Arg
Gly 130	Tyr	Asp	Val	Lys	Thr	Leu 135	Leu	Ala	Met	Val	Lys 140	Ser	Lys	Leu	Lys
Gly 145	Glu	Lys	Leu	Lys 150	Asp	Asp	Lys	Thr	Met	Leu 155	Met	Glu	Arg	Val	Met 160
Gln	Leu	Val	Ala 165	Arg	Leu	Pro	Thr	Glu 170	Ser	Lys	Lys	Arg	Ala 175	Glu	Leu
Thr	Asp	Ser 180	Leu	Ile	Asn	Glu	Leu 185	Trp	Glu	Ser	Leu	Asp 190	His	Pro	Pro
Leu	Asn 195	Tyr	Leu	Gly	Pro	Glu	His 200	Ser	Tyr	Arg	Thr	Pro 205	Asp	Gly	Ser
Tyr 210	Asn	His	Pro	Phe	Asn 215	Pro	Gln	Leu	Gly	Ala	Ala 220	Gly	Ser	Arg	Tyr
Ala 225	Arg	Ser	Val	Ile	Pro 230	Thr	Val	Thr	Pro	Pro	Gly 235	Ala	Leu	Pro	Asp 240
Pro	Gly	Leu	Ile 245	Phe	Asp	Ser	Ile	Met	Gly 250	Arg	Thr	Pro	Asn 255	Ser	Tyr
Arg	Lys	His 260	Pro	Asn	Asn	Val	Ser 265	Ser	Ile	Leu	Trp	Tyr 270	Trp	Ala	Thr
Ile	Ile 275	Ile	His	Asp	Ile	Phe	Trp 280	Thr	Asp	Pro	Arg 285	Asp	Ile	Asn	Thr
Asn 290	Lys	Ser	Ser	Ser	Tyr	Leu 295	Asp	Leu	Ala	Pro	Leu 300	Tyr	Gly	Asn	Ser
Gln 305	Glu	Met	Gln	Asp 310	Ser	Ile	Arg	Thr	Phe	Lys 315	Asp	Gly	Arg	Met	Lys 320
Pro	Asp	Cys	Tyr 325	Ala	Asp	Lys	Arg	Leu	Ala 330	Gly	Met	Pro	Pro	Gly 335	Val
Ser	Val	Leu	Leu 340	Ile	Met	Phe	Asn 345	Arg	Phe	His	Asn 350	His	Val	Ala	Glu

Asn	Leu	Ala	Leu	Ile	Asn	Glu	Gly	Gly	Arg	Phe	Asn	Lys	Pro	Ser	Asp
		355					360					365			
Leu	Leu	Glu	Gly	Glu	Ala	Arg	Glu	Ala	Ala	Trp	Lys	Lys	Tyr	Asp	Asn
		370					375					380			
Asp	Leu	Phe	Gln	Val	Ala	Arg	Leu	Val	Thr	Ser	Gly	Leu	Tyr	Ile	Asn
385					390					395					400
Ile	Thr	Leu	Val	Asp	Tyr	Val	Arg	Asn	Ile	Val	Asn	Leu	Asn	Arg	Val
				405					410						415
Asp	Thr	Thr	Trp	Thr	Leu	Asp	Pro	Arg	Gln	Asp	Ala	Gly	Ala	His	Val
			420						425					430	
Gly	Thr	Ala	Asp	Gly	Ala	Glu	Arg	Gly	Thr	Gly	Asn	Ala	Val	Ser	Ala
		435						440				445			
Glu	Phe	Asn	Leu	Cys	Tyr	Arg	Trp	His	Ser	Cys	Ile	Ser	Glu	Lys	Asp
		450					455					460			
Ser	Lys	Phe	Val	Glu	Ala	Gln	Phe	Gln	Asn	Ile	Phe	Gly	Lys	Pro	Ala
465					470					475					480
Ser	Glu	Val	Arg	Pro	Asp	Glu	Met	Trp	Lys	Gly	Phe	Ala	Lys	Met	Glu
				485					490					495	
Gln	Asn	Thr	Pro	Ala	Asp	Pro	Gly	Gln	Arg	Thr	Phe	Gly	Gly	Phe	Lys
			500					505					510		
Arg	Gly	Pro	Asp	Gly	Lys	Phe	Asp	Asp	Asp	Asp	Leu	Val	Arg	Cys	Ile
		515					520					525			
Ser	Glu	Ala	Val	Glu	Asp	Val	Ala	Gly	Ala	Phe	Gly	Ala	Arg	Asn	Val
	530					535					540				
Pro	Gln	Ala	Met	Lys	Val	Val	Glu	Thr	Met	Gly	Ile	Ile	Gln	Gly	Arg
545					550					555					560
Lys	Trp	Asn	Val	Ala	Gly	Leu	Asn	Glu	Phe	Arg	Lys	His	Phe	His	Leu
				565					570					575	
Lys	Pro	Tyr	Ser	Thr	Phe	Glu	Asp	Ile	Asn	Ser	Asp	Pro	Gly	Val	Ala
			580					585					590		
Glu	Ala	Leu	Arg	Arg	Leu	Tyr	Asp	His	Pro	Asp	Asn	Val	Glu	Leu	Tyr
		595					600					605			
Pro	Gly	Leu	Val	Ala	Glu	Glu	Asp	Lys	Gln	Pro	Met	Val	Pro	Gly	Val
	610					615					620				
Gly	Ile	Ala	Pro	Thr	Tyr	Thr	Ile	Ser	Arg	Val	Val	Leu	Ser	Asp	Ala
625					630					635					640
Val	Cys	Leu	Val	Arg	Gly	Asp	Arg	Phe	Tyr	Thr	Thr	Asp	Phe	Thr	Pro
				645					650					655	
Arg	Asn	Leu	Thr	Asn	Trp	Gly	Tyr	Lys	Glu	Val	Asp	Tyr	Asp	Leu	Ser
			660					665						670	
Val	Asn	His	Gly	Cys	Val	Phe	Tyr	Lys	Leu	Phe	Ile	Arg	Ala	Phe	Pro
		675					680					685			
Asn	His	Phe	Lys	Gln	Asn	Ser	Val	Tyr	Ala	His	Tyr	Pro	Met	Val	Val
	690					695					700				
Pro	Ser	Glu	Asn	Lys	Arg	Ile	Leu	Glu	Ala	Leu	Gly	Arg	Ala	Asp	Leu
705					710					715					720
Phe	Asp	Phe	Glu	Ala	Pro	Lys	Tyr	Ile	Pro	Pro	Arg	Val	Asn	Ile	Thr
				725					730					735	
Ser	Tyr	Gly	Gly	Ala	Glu	Tyr	Ile	Leu	Glu	Thr	Gln	Glu	Lys	Tyr	Lys

Figure 4 (Continued)

			740					745					750				
Val	Thr	Trp	His	Glu	Gly	Leu	Gly	Phe	Leu	Met	Gly	Glu	Gly	Gly	Leu		
		755					760					765					
Lys	Phe	Met	Leu	Ser	Gly	Asp	Asp	Pro	Leu	His	Ala	Gln	Gln	Arg	Lys		
	770					775					780						
Cys	Met	Ala	Ala	Gln	Leu	Tyr	Lys	Asp	Gly	Trp	Thr	Glu	Ala	Val	Lys		
785					790					795					800		
Ala	Phe	Tyr	Ala	Gly	Met	Met	Glu	Glu	Leu	Leu	Val	Ser	Lys	Ser	Tyr		
				805					810					815			
Phe	Leu	Gly	Asn	Asn	Lys	His	Arg	His	Val	Asp	Ile	Ile	Arg	Asp	Val		
			820					825					830				
Gly	Asn	Met	Val	His	Val	His	Phe	Ala	Ser	Gln	Val	Phe	Gly	Leu	Pro		
	835						840					845					
Leu	Lys	Thr	Ala	Lys	Asn	Pro	Thr	Gly	Val	Phe	Thr	Glu	Gln	Glu	Met		
	850					855					860						
Tyr	Gly	Ile	Leu	Ala	Ala	Ile	Phe	Thr	Thr	Ile	Phe	Phe	Asp	Leu	Asp		
865					870					875					880		
Pro	Ser	Lys	Ser	Phe	Pro	Leu	Arg	Thr	Lys	Thr	Arg	Glu	Val	Cys	Gln		
				885					890					895			
Lys	Leu	Ala	Lys	Leu	Val	Glu	Ala	Asn	Val	Lys	Leu	Ile	Asn	Lys	Ile		
			900					905					910				
Pro	Trp	Ser	Arg	Gly	Met	Phe	Val	Gly	Lys	Pro	Ala	Lys	Asp	Glu	Pro		
	915						920					925					
Leu	Ser	Ile	Tyr	Gly	Lys	Thr	Met	Ile	Lys	Gly	Leu	Lys	Ala	His	Gly		
	930					935					940						
Leu	Ser	Asp	Tyr	Asp	Ile	Ala	Trp	Ser	His	Val	Val	Pro	Thr	Ser	Gly		
945					950					955					960		
Ala	Met	Val	Pro	Asn	Gln	Ala	Gln	Val	Phe	Ala	Gln	Ala	Val	Asp	Tyr		
				965					970					975			
Tyr	Leu	Ser	Pro	Ala	Gly	Met	His	Tyr	Ile	Pro	Glu	Ile	His	Met	Val		
			980					985					990				
Ala	Leu	Gln	Pro	Ser	Thr	Pro	Glu	Thr	Asp	Ala	Leu	Leu	Leu	Gly	Tyr		
	995						1000					1005					
Ala	Met	Glu	Gly	Ile	Arg	Leu	Ala	Gly	Thr	Phe	Gly	Ser	Tyr	Arg	Glu		
	1010					1015					1020						
Ala	Ala	Val	Asp	Asp	Val	Val	Lys	Glu	Asp	Asn	Gly	Arg	Gln	Val	Pro		
1025					1030					1035					1040		
Val	Lys	Ala	Gly	Asp	Arg	Val	Phe	Val	Ser	Phe	Val	Asp	Ala	Ala	Arg		
				1045					1050					1055			
Asp	Pro	Lys	His	Phe	Pro	Asp	Pro	Glu	Val	Val	Asn	Pro	Arg	Arg	Pro		
			1060					1065					1070				
Ala	Lys	Lys	Tyr	Ile	His	Tyr	Gly	Val	Gly	Pro	His	Ala	Cys	Leu	Gly		
	1075						1080					1085					
Arg	Asp	Ala	Ser	Gln	Ile	Ala	Ile	Thr	Glu	Met	Phe	Arg	Cys	Leu	Phe		
	1090					1095					1100						
Arg	Arg	Arg	Asn	Val	Arg	Arg	Val	Pro	Gly	Pro	Gln	Gly	Glu	Leu	Lys		
1105				1110					1115					1120			
Lys	Val	Pro	Arg	Pro	Gly	Gly	Phe	Tyr	Val	Tyr	Met	Arg	Glu	Asp	Trp		
				1125					1130					1135			

Figure 4 (Continued)

Gly Gly Leu Phe Pro Phe Pro Val Thr Met Arg Val Met Trp Asp Asp
 1140 1145 1150
 Glu

<210> 7
 <211> 530
 <212> PRT
 <213> *Fusarium oxysporum*

<400> 7
 Met Lys Gly Ser Ala Thr Leu Ala Phe Ala Leu Val Gln Phe Ser Ala
 1 5 10 15
 Ala Ser Gln Leu Val Trp Pro Ser Lys Trp Asp Glu Val Glu Asp Leu
 20 25 30
 Leu Tyr Met Gln Gly Gly Phe Asn Lys Arg Gly Phe Ala Asp Ala Leu
 35 40 45
 Arg Thr Cys Glu Phe Gly Ser Asn Val Pro Gly Thr Gln Asn Thr Ala
 50 55 60
 Glu Trp Leu Arg Thr Ala Phe His Asp Ala Ile Thr His Asp Ala Lys
 65 70 75 80
 Ala Gly Thr Gly Gly Leu Asp Ala Ser Ile Tyr Trp Glu Ser Ser Arg
 85 90 95
 Pro Glu Asn Pro Gly Lys Ala Phe Asn Asn Thr Phe Gly Phe Phe Ser
 100 105 110
 Gly Phe His Asn Pro Arg Ala Thr Ala Ser Asp Leu Thr Ala Leu Gly
 115 120 125
 Thr Val Leu Ala Val Gly Ala Cys Asn Gly Pro Arg Ile Pro Phe Arg
 130 135 140
 Ala Gly Arg Ile Asp Ala Tyr Lys Ala Gly Pro Ala Gly Val Pro Glu
 145 150 155 160
 Pro Ser Thr Asn Leu Lys Asp Thr Phe Ala Ala Phe Thr Lys Ala Gly
 165 170 175
 Phe Thr Lys Glu Glu Met Thr Ala Met Val Ala Cys Gly His Ala Ile
 180 185 190
 Gly Gly Val His Ser Val Asp Phe Pro Glu Ile Val Gly Ile Lys Ala
 195 200 205
 Asp Pro Asn Asn Asp Thr Asn Val Pro Phe Gln Lys Asp Val Ser Ser
 210 215 220
 Phe His Asn Gly Ile Val Thr Glu Tyr Leu Ala Gly Thr Ser Lys Asn
 225 230 235 240
 Pro Leu Val Ala Ser Lys Asn Ala Thr Phe His Ser Asp Lys Arg Ile
 245 250 255
 Phe Asp Asn Asp Lys Ala Thr Met Lys Lys Leu Ser Thr Lys Ala Gly
 260 265 270
 Phe Asn Ser Met Cys Ala Asp Ile Leu Thr Arg Met Ile Asp Thr Val
 275 280 285
 Pro Lys Ser Val Gln Leu Thr Pro Val Leu Glu Ala Tyr Asp Val Arg

290	295	300
Pro Tyr Ile Thr Glu Leu Ser Leu Asn Asn Lys Asn Lys Ile His Phe		
305	310	315
Thr Gly Ser Val Arg Val Arg Ile Thr Asn Asn Ile Arg Asp Asn Asn		
	325	330
Asp Leu Ala Ile Asn Leu Ile Tyr Val Gly Arg Asp Gly Lys Lys Val		
	340	345
Thr Val Pro Thr Gln Gln Val Thr Phe Gln Gly Gly Thr Ser Phe Gly		
	355	360
Ala Gly Glu Val Phe Ala Asn Phe Glu Phe Asp Thr Thr Met Asp Ala		
	370	375
Lys Asn Gly Ile Thr Lys Phe Phe Ile Gln Glu Val Lys Pro Ser Thr		
385	390	395
Lys Ala Thr Val Thr His Asp Asn Gln Lys Thr Gly Gly Tyr Lys Val		
	405	410
Asp Asp Thr Val Leu Tyr Gln Leu Gln Gln Ser Cys Ala Val Leu Glu		
	420	425
Lys Leu Pro Asn Ala Pro Leu Val Val Thr Ala Met Val Arg Asp Ala		
	435	440
Arg Ala Lys Asp Ala Leu Thr Leu Arg Val Ala His Lys Lys Pro Val		
	450	455
Lys Gly Ser Ile Val Pro Arg Phe Gln Thr Ala Ile Thr Asn Phe Lys		
465	470	475
Ala Thr Gly Lys Lys Ser Ser Gly Tyr Thr Gly Phe Gln Ala Lys Thr		
	485	490
Met Phe Glu Glu Gln Ser Thr Tyr Phe Asp Ile Val Leu Gly Gly Ser		
	500	505
Pro Ala Ser Gly Val Gln Phe Leu Thr Ser Gln Ala Met Pro Ser Gln		
	515	520
Cys Ser		525
530		

<210> 8

<211> 358

<212> PRT

<213> Fusarium graminearum

<400> 8

Met Ala Ser Ala Thr Arg Gln Phe Ala Arg Ala Ala Thr Arg Ala Thr		
1	5	10
Arg Asn Gly Phe Ala Ile Ala Pro Arg Gln Val Ile Arg Gln Gln Gly		
	20	25
Arg Arg Tyr Tyr Ser Ser Glu Pro Ala Gln Lys Ser Ser Ser Ala Trp		
	35	40
Ile Trp Leu Thr Gly Ala Ala Val Ala Gly Gly Ala Gly Tyr Tyr Phe		
	50	55
Tyr Gly Asn Ser Ala Ser Ser Ala Thr Ala Lys Val Phe Asn Pro Ser		
65	70	75
		80

Lys Glu Asp Tyr Gln Lys Val Tyr Asn Glu Ile Ala Ala Arg Leu Glu
 85 90 95
 Glu Lys Asp Asp Tyr Asp Asp Gly Ser Tyr Gly Pro Val Leu Val Arg
 100 105 110
 Leu Ala Trp His Ala Ser Gly Thr Tyr Asp Lys Glu Thr Gly Thr Gly
 115 120 125
 Gly Ser Asn Gly Ala Thr Met Arg Phe Ala Pro Glu Ser Asp His Gly
 130 135 140
 Ala Asn Ala Gly Leu Ala Ala Ala Arg Asp Phe Leu Gln Pro Val Lys
 145 150 155 160
 Glu Lys Phe Pro Trp Ile Thr Tyr Ser Asp Leu Trp Ile Leu Ala Gly
 165 170 175
 Val Cys Ala Ile Gln Glu Met Leu Gly Pro Ala Ile Pro Tyr Arg Pro
 180 185 190
 Gly Arg Ser Asp Arg Asp Val Ser Gly Cys Thr Pro Asp Gly Arg Leu
 195 200 205
 Pro Asp Ala Ser Lys Arg Gln Asp His Leu Arg Gly Ile Phe Gly Arg
 210 215 220
 Met Gly Phe Asn Asp Gln Glu Ile Val Ala Leu Ser Gly Ala His Ala
 225 230 235 240
 Leu Gly Arg Cys His Thr Asp Arg Ser Gly Tyr Ser Gly Pro Trp Thr
 245 250 255
 Phe Ser Pro Thr Val Leu Thr Asn Asp Tyr Phe Arg Leu Leu Val Glu
 260 265 270
 Glu Lys Trp Gln Trp Lys Lys Trp Asn Gly Pro Ala Gln Tyr Glu Asp
 275 280 285
 Lys Ser Thr Lys Ser Leu Met Met Leu Pro Ser Asp Ile Ala Leu Ile
 290 295 300
 Glu Asp Lys Lys Phe Lys Pro Trp Val Glu Lys Tyr Ala Lys Asp Asn
 305 310 315 320
 Asp Ala Phe Phe Lys Asp Phe Ser Asn Val Val Leu Arg Leu Phe Glu
 325 330 335
 Leu Gly Val Pro Phe Ala Gln Gly Thr Glu Asn Gln Arg Trp Thr Phe
 340 345 350
 Lys Pro Thr His Gln Glu
 355

<210> 9

<211> 122

<212> PRT

<213> Chlamydomonas eugametos

<400> 9

Met Ser Leu Phe Ala Lys Leu Gly Gly Arg Glu Ala Val Glu Ala Ala
 1 5 10 15
 Val Asp Lys Phe Tyr Asn Lys Ile Val Ala Asp Pro Thr Val Ser Thr
 20 25 30
 Tyr Phe Ser Asn Thr Asp Met Lys Val Gln Arg Ser Lys Gln Phe Ala

		35					40				45						
Phe	Leu	Ala	Tyr	Ala	Leu	Gly	Gly	Ala	Ser	Glu	Trp	Lys	Gly	Lys	Asp		
	50					55					60						
Met	Arg	Thr	Ala	His	Lys	Asp	Leu	Val	Pro	His	Leu	Ser	Asp	Val	His		
65					70					75					80		
Phe	Gln	Ala	Val	Ala	Arg	His	Leu	Ser	Asp	Thr	Leu	Thr	Glu	Leu	Gly		
				85					90					95			
Val	Pro	Pro	Glu	Asp	Ile	Thr	Asp	Ala	Met	Ala	Val	Val	Ala	Ser	Thr		
			100					105					110				
Arg	Thr	Glu	Val	Leu	Asn	Met	Pro	Gln	Gln								
		115					120										

<210> 10

<211> 121

<212> PRT

<213> Tetrahymena pyriformis

<400> 10

Met	Asn	Lys	Pro	Gln	Thr	Ile	Tyr	Glu	Lys	Leu	Gly	Gly	Glu	Asn	Ala		
1				5				10					15				
Met	Lys	Ala	Ala	Val	Pro	Leu	Phe	Tyr	Lys	Lys	Val	Leu	Ala	Asp	Glu		
		20						25					30				
Arg	Val	Lys	His	Phe	Phe	Lys	Asn	Thr	Asp	Met	Asp	His	Gln	Thr	Lys		
		35					40					45					
Gln	Gln	Thr	Asp	Phe	Leu	Thr	Met	Leu	Leu	Gly	Gly	Pro	Asn	His	Tyr		
	50					55					60						
Lys	Gly	Lys	Asn	Met	Thr	Glu	Ala	His	Lys	Gly	Met	Asn	Leu	Gln	Asn		
65				70					75					80			
Leu	His	Phe	Asp	Ala	Ile	Ile	Glu	Asn	Leu	Ala	Ala	Thr	Leu	Lys	Glu		
			85					90						95			
Leu	Gly	Val	Thr	Asp	Ala	Val	Ile	Asn	Glu	Ala	Ala	Lys	Val	Ile	Glu		
			100				105						110				
His	Thr	Arg	Lys	Asp	Met	Leu	Gly	Lys									
		115					120										

<210> 11

<211> 117

<212> PRT

<213> Paramecium caudatum

<400> 11

Met	Ser	Leu	Phe	Glu	Gln	Leu	Gly	Gly	Gln	Ala	Ala	Val	Gln	Ala	Val		
1				5				10					15				
Thr	Ala	Gln	Phe	Tyr	Ala	Asn	Ile	Gln	Ala	Asp	Ala	Thr	Val	Ala	Thr		
		20						25					30				
Phe	Phe	Asn	Gly	Ile	Asp	Met	Pro	Asn	Gln	Thr	Asn	Lys	Thr	Ala	Ala		
		35					40					45					

```

Phe Leu Cys Ala Ala Leu Gly Gly Pro Asn Ala Trp Thr Gly Arg Asn
  50                      55                      60
Leu Lys Glu Val His Ala Asn Met Gly Val Ser Asn Ala Gln Phe Thr
65                      70                      75                      80
Thr Val Ile Gly His Leu Arg Ser Ala Leu Thr Gly Ala Gly Val Ala
                      85                      90                      95
Ala Ala Leu Val Glu Gln Thr Val Ala Val Ala Glu Thr Val Arg Gly
                      100                      105                      110
Asp Val Val Thr Val
                      115

```

```

<210> 12
<211> 147
<212> PRT
<213> Aspergillus niger

```

```

<400> 12
Met Pro Leu Thr Pro Glu Gln Ile Lys Ile Ile Lys Ala Thr Val Pro
  1                      5                      10                      15
Val Leu Gln Glu Tyr Gly Thr Lys Ile Thr Thr Ala Phe Tyr Met Asn
                      20                      25                      30
Met Ser Thr Val His Pro Glu Leu Asn Ala Val Phe Asn Thr Ala Asn
                      35                      40                      45
Gln Val Lys Gly His Gln Ala Arg Ala Leu Ala Gly Ala Leu Phe Ala
                      50                      55                      60
Tyr Ala Ser His Ile Asp Asp Leu Gly Ala Leu Gly Pro Ala Val Glu
65                      70                      75                      80
Leu Ile Cys Asn Lys His Ala Ser Leu Tyr Ile Gln Ala Asp Glu Tyr
                      85                      90                      95
Lys Ile Val Gly Lys Tyr Leu Leu Glu Ala Met Lys Glu Val Leu Gly
                      100                      105                      110
Asp Ala Cys Thr Asp Asp Ile Leu Asp Ala Trp Gly Ala Ala Tyr Trp
                      115                      120                      125
Ala Leu Ala Asp Ile Met Ile Asn Arg Glu Ala Ala Leu Tyr Lys Gln
                      130                      135                      140
Ser Gln Gly
145

```

```

<210> 13
<211> 165
<212> PRT
<213> Zea mays

```

```

<400> 13
Met Ala Leu Ala Glu Ala Asp Asp Gly Ala Val Val Phe Gly Glu Glu
  1                      5                      10                      15
Gln Glu Ala Leu Val Leu Lys Ser Trp Ala Val Met Lys Lys Asp Ala

```

			20					25					30				
Ala	Asn	Leu	Gly	Leu	Arg	Phe	Phe	Leu	Lys	Val	Phe	Glu	Ile	Ala	Pro		
		35					40					45					
Ser	Ala	Glu	Gln	Met	Phe	Ser	Phe	Leu	Arg	Asp	Ser	Asp	Val	Pro	Leu		
	50					55					60						
Glu	Lys	Asn	Pro	Lys	Leu	Lys	Thr	His	Ala	Met	Ser	Val	Phe	Val	Met		
65					70					75					80		
Thr	Cys	Glu	Ala	Ala	Ala	Gln	Leu	Arg	Lys	Ala	Gly	Lys	Val	Thr	Val		
				85					90					95			
Arg	Glu	Thr	Thr	Leu	Lys	Arg	Leu	Gly	Ala	Thr	His	Leu	Arg	Tyr	Gly		
			100					105					110				
Val	Ala	Asp	Gly	His	Phe	Glu	Val	Thr	Gly	Phe	Ala	Leu	Leu	Glu	Thr		
		115						120				125					
Ile	Lys	Glu	Ala	Leu	Pro	Ala	Asp	Met	Trp	Ser	Leu	Glu	Met	Lys	Lys		
	130					135					140						
Ala	Trp	Ala	Glu	Ala	Tyr	Ser	Gln	Leu	Val	Ala	Ala	Ile	Lys	Arg	Glu		
145					150					155					160		
Met	Lys	Pro	Asp	Ala													
				165													

<210> 14

<211> 169

<212> PRT

<213> *Oryza sativa* subsp. *japonica*

<400> 14

Met	Ala	Leu	Val	Glu	Gly	Asn	Asn	Gly	Val	Ser	Gly	Gly	Ala	Val	Ser		
1				5					10					15			
Phe	Ser	Glu	Glu	Gln	Glu	Ala	Leu	Val	Leu	Lys	Ser	Trp	Ala	Ile	Met		
		20						25					30				
Lys	Lys	Asp	Ser	Ala	Asn	Ile	Gly	Leu	Arg	Phe	Phe	Leu	Lys	Ile	Phe		
		35					40					45					
Glu	Val	Ala	Pro	Ser	Ala	Ser	Gln	Met	Phe	Ser	Phe	Leu	Arg	Asn	Ser		
	50					55					60						
Asp	Val	Pro	Leu	Glu	Lys	Asn	Pro	Lys	Leu	Lys	Thr	His	Ala	Met	Ser		
65					70					75					80		
Val	Phe	Val	Met	Thr	Cys	Glu	Ala	Ala	Ala	Gln	Leu	Arg	Lys	Ala	Gly		
				85					90					95			
Lys	Val	Thr	Val	Arg	Asp	Thr	Thr	Leu	Lys	Arg	Leu	Gly	Ala	Thr	His		
			100					105					110				
Phe	Lys	Tyr	Gly	Val	Gly	Asp	Ala	His	Phe	Glu	Val	Thr	Arg	Phe	Ala		
		115					120					125					
Leu	Leu	Glu	Thr	Ile	Lys	Glu	Ala	Val	Pro	Val	Asp	Met	Trp	Ser	Pro		
	130					135					140						
Ala	Met	Lys	Ser	Ala	Trp	Ser	Glu	Ala	Tyr	Asn	Gln	Leu	Val	Ala	Ala		
145					150					155					160		
Ile	Lys	Gln	Glu	Met	Lys	Pro	Ala	Glu									
				165													

<210> 15
 <211> 160
 <212> PRT
 <213> *Arabidopsis thaliana*

<400> 15
 Met Glu Ser Glu Gly Lys Ile Val Phe Thr Glu Glu Gln Glu Ala Leu
 1 5 10 15
 Val Val Lys Ser Trp Ser Val Met Lys Lys Asn Ser Ala Glu Leu Gly
 20 25 30
 Leu Lys Leu Phe Ile Lys Ile Phe Glu Ile Ala Pro Thr Thr Lys Lys
 35 40 45
 Met Phe Ser Phe Leu Arg Asp Ser Pro Ile Pro Ala Glu Gln Asn Pro
 50 55 60
 Lys Leu Lys Pro His Ala Met Ser Val Phe Val Met Cys Cys Glu Ser
 65 70 75 80
 Ala Val Gln Leu Arg Lys Thr Gly Lys Val Thr Val Arg Glu Thr Thr
 85 90 95
 Leu Lys Arg Leu Gly Ala Ser His Ser Lys Tyr Gly Val Val Asp Glu
 100 105 110
 His Phe Glu Val Ala Lys Tyr Ala Leu Leu Glu Thr Ile Lys Glu Ala
 115 120 125
 Val Pro Glu Met Trp Ser Pro Glu Met Lys Val Ala Trp Gly Gln Ala
 130 135 140
 Tyr Asp His Leu Val Ala Ala Ile Lys Ala Glu Met Asn Leu Ser Asn
 145 150 155 160

<210> 16
 <211> 147
 <212> PRT
 <213> *Pisum sativum*

<400> 16
 Met Gly Phe Thr Asp Lys Gln Glu Ala Leu Val Asn Ser Ser Trp Glu
 1 5 10 15
 Ser Phe Lys Gln Asn Leu Ser Gly Asn Ser Ile Leu Phe Tyr Thr Ile
 20 25 30
 Ile Leu Glu Lys Ala Pro Ala Ala Lys Gly Leu Phe Ser Phe Leu Lys
 35 40 45
 Asp Thr Ala Gly Val Glu Asp Ser Pro Lys Leu Gln Ala His Ala Glu
 50 55 60
 Gln Val Phe Gly Leu Val Arg Asp Ser Ala Ala Gln Leu Arg Thr Lys
 65 70 75 80
 Gly Glu Val Val Leu Gly Asn Ala Thr Leu Gly Ala Ile His Val Gln
 85 90 95
 Arg Gly Val Thr Asp Pro His Phe Val Val Val Lys Glu Ala Leu Leu

			100					105					110				
Gln	Thr	Ile	Lys	Lys	Ala	Ser	Gly	Asn	Asn	Trp	Ser	Glu	Glu	Leu	Asn		
			115					120					125				
Thr	Ala	Trp	Glu	Val	Ala	Tyr	Asp	Gly	Leu	Ala	Thr	Ala	Ile	Lys	Lys		
			130				135					140					
Ala	Met	Thr															
145																	

<210> 17
 <211> 145
 <212> PRT
 <213> Vigna unguiculata

<400> 17

Met	Val	Ala	Phe	Ser	Asp	Lys	Gln	Glu	Ala	Leu	Val	Asn	Gly	Ala	Tyr		
1				5					10					15			
Glu	Ala	Phe	Lys	Ala	Asn	Ile	Pro	Lys	Tyr	Ser	Val	Val	Phe	Tyr	Thr		
			20					25					30				
Thr	Ile	Leu	Glu	Lys	Ala	Pro	Ala	Ala	Lys	Asn	Leu	Phe	Ser	Phe	Leu		
			35				40					45					
Ala	Asn	Gly	Val	Asp	Ala	Thr	Asn	Pro	Lys	Leu	Thr	Gly	His	Ala	Glu		
			50				55					60					
Lys	Leu	Phe	Gly	Leu	Val	Arg	Asp	Ser	Ala	Ala	Gln	Leu	Arg	Ala	Ser		
65						70				75					80		
Gly	Gly	Val	Val	Ala	Asp	Ala	Ala	Leu	Gly	Ala	Val	His	Ser	Gln	Lys		
				85					90					95			
Ala	Val	Asn	Asp	Ala	Gln	Phe	Val	Val	Lys	Glu	Ala	Leu	Val	Lys			
			100					105				110					
Thr	Leu	Lys	Glu	Ala	Val	Gly	Asp	Lys	Trp	Ser	Asp	Glu	Leu	Gly	Thr		
			115				120					125					
Ala	Val	Glu	Leu	Ala	Tyr	Asp	Glu	Leu	Ala	Ala	Ala	Ile	Lys	Lys	Ala		
			130				135					140					
Tyr																	
145																	

<210> 18
 <211> 154
 <212> PRT
 <213> Bos taurus

<400> 18

Met	Gly	Leu	Ser	Asp	Gly	Glu	Trp	Gln	Leu	Val	Leu	Asn	Ala	Trp	Gly		
1				5					10					15			
Lys	Val	Glu	Ala	Asp	Val	Ala	Gly	His	Gly	Gln	Glu	Val	Leu	Ile	Arg		
			20					25					30				
Leu	Phe	Thr	Gly	His	Pro	Glu	Thr	Leu	Glu	Lys	Phe	Asp	Lys	Phe	Lys		
			35				40					45					

```

His Leu Lys Thr Glu Ala Glu Met Lys Ala Ser Glu Asp Leu Lys Lys
 50                      55                      60
His Gly Asn Thr Val Leu Thr Ala Leu Gly Gly Ile Leu Lys Lys Lys
65                      70                      75                      80
Gly His His Glu Ala Glu Val Lys His Leu Ala Glu Ser His Ala Asn
                      85                      90                      95
Lys His Lys Ile Pro Val Lys Tyr Leu Glu Phe Ile Ser Asp Ala Ile
                      100                     105                     110
Ile His Val Leu His Ala Lys His Pro Ser Asp Phe Gly Ala Asp Ala
                      115                     120                     125
Gln Ala Ala Met Ser Lys Ala Leu Glu Leu Phe Arg Asn Asp Met Ala
                      130                     135                     140
Ala Gln Tyr Lys Val Leu Gly Phe His Gly
145                      150

```

```

<210> 19
<211> 154
<212> PRT
<213> Sus scrofa

```

```

<400> 19
Met Gly Leu Ser Asp Gly Glu Trp Gln Leu Val Leu Asn Val Trp Gly
 1                      5                      10                      15
Lys Val Glu Ala Asp Val Ala Gly His Gly Gln Glu Val Leu Ile Arg
                      20                      25                      30
Leu Phe Lys Gly His Pro Glu Thr Leu Glu Lys Phe Asp Lys Phe Lys
                      35                      40                      45
His Leu Lys Ser Glu Asp Glu Met Lys Ala Ser Glu Asp Leu Lys Lys
50                      55                      60
His Gly Asn Thr Val Leu Thr Ala Leu Gly Gly Ile Leu Lys Lys Lys
65                      70                      75                      80
Gly His His Glu Ala Glu Leu Thr Pro Leu Ala Gln Ser His Ala Thr
                      85                      90                      95
Lys His Lys Ile Pro Val Lys Tyr Leu Glu Phe Ile Ser Glu Ala Ile
                      100                     105                     110
Ile Gln Val Leu Gln Ser Lys His Pro Gly Asp Phe Gly Ala Asp Ala
                      115                     120                     125
Gln Gly Ala Met Ser Lys Ala Leu Glu Leu Phe Arg Asn Asp Met Ala
                      130                     135                     140
Ala Lys Tyr Lys Glu Leu Gly Phe Gln Gly
145                      150

```

```

<210> 20
<211> 154
<212> PRT
<213> Equus caballus

```

<400> 20

```

Met Gly Leu Ser Asp Gly Glu Trp Gln Gln Val Leu Asn Val Trp Gly
 1          5          10          15
Lys Val Glu Ala Asp Ile Ala Gly His Gly Gln Glu Val Leu Ile Arg
          20          25          30
Leu Phe Thr Gly His Pro Glu Thr Leu Glu Lys Phe Asp Lys Phe Lys
          35          40          45
His Leu Lys Thr Glu Ala Glu Met Lys Ala Ser Glu Asp Leu Lys Lys
 50          55          60
His Gly Thr Val Val Leu Thr Ala Leu Gly Gly Ile Leu Lys Lys Lys
65          70          75          80
Gly His His Glu Ala Glu Leu Lys Pro Leu Ala Gln Ser His Ala Thr
          85          90          95
Lys His Lys Ile Pro Ile Lys Tyr Leu Glu Phe Ile Ser Asp Ala Ile
          100          105          110
Ile His Val Leu His Ser Lys His Pro Gly Asp Phe Gly Ala Asp Ala
          115          120          125
Gln Gly Ala Met Thr Lys Ala Leu Glu Leu Phe Arg Asn Asp Ile Ala
          130          135          140
Ala Lys Tyr Lys Glu Leu Gly Phe Gln Gly
145          150

```

<210> 21

<211> 152

<212> PRT

<213> *Nicotiana benthamiana*

<400> 21

```

Met Ser Ser Phe Thr Glu Glu Gln Glu Ala Leu Val Val Lys Ser Trp
 1          5          10          15
Asp Ser Met Lys Lys Asn Ala Gly Glu Trp Gly Leu Lys Leu Phe Leu
          20          25          30
Lys Ile Phe Glu Ile Ala Pro Ser Ala Lys Lys Leu Phe Ser Phe Leu
          35          40          45
Lys Asp Ser Asn Val Pro Leu Glu Gln Asn Ala Lys Leu Lys Pro His
 50          55          60
Ser Lys Ser Val Phe Val Met Thr Cys Glu Ala Ala Val Gln Leu Arg
65          70          75          80
Lys Ala Gly Lys Val Val Val Arg Asp Ser Thr Leu Lys Lys Leu Gly
          85          90          95
Ala Thr His Phe Lys Tyr Gly Val Ala Asp Glu His Phe Glu Val Thr
          100          105          110
Lys Phe Ala Leu Leu Glu Thr Ile Lys Glu Ala Val Pro Glu Met Trp
          115          120          125
Ser Val Asp Met Lys Asn Ala Trp Gly Glu Ala Phe Asp Gln Leu Val
          130          135          140
Asn Ala Ile Lys Thr Glu Met Lys
145          150

```

<210> 22
 <211> 132
 <212> PRT
 <213> *Bacillus subtilis*

<400> 22
 Met Gly Gln Ser Phe Asn Ala Pro Tyr Glu Ala Ile Gly Glu Glu Leu
 1 5 10 15
 Leu Ser Gln Leu Val Asp Thr Phe Tyr Glu Arg Val Ala Ser His Pro
 20 25 30
 Leu Leu Lys Pro Ile Phe Pro Ser Asp Leu Thr Glu Thr Ala Arg Lys
 35 40 45
 Gln Lys Gln Phe Leu Thr Gln Tyr Leu Gly Gly Pro Pro Leu Tyr Thr
 50 55 60
 Glu Glu His Gly His Pro Met Leu Arg Ala Arg His Leu Pro Phe Pro
 65 70 75 80
 Ile Thr Asn Glu Arg Ala Asp Ala Trp Leu Ser Cys Met Lys Asp Ala
 85 90 95
 Met Asp His Val Gly Leu Glu Gly Glu Ile Arg Glu Phe Leu Phe Gly
 100 105 110
 Arg Leu Glu Leu Thr Ala Arg His Met Val Asn Gln Thr Glu Ala Glu
 115 120 125
 Asp Arg Ser Ser
 130

<210> 23
 <211> 131
 <212> PRT
 <213> *Corynebacterium glutamicum*

<400> 23
 Met Thr Thr Ser Glu Asn Phe Tyr Asp Ser Val Gly Gly Glu Glu Thr
 1 5 10 15
 Phe Ser Leu Ile Val His Arg Phe Tyr Glu Gln Val Pro Asn Asp Asp
 20 25 30
 Ile Leu Gly Pro Met Tyr Pro Pro Asp Asp Phe Glu Gly Ala Glu Gln
 35 40 45
 Arg Leu Lys Met Phe Leu Ser Gln Tyr Trp Gly Gly Pro Lys Asp Tyr
 50 55 60
 Gln Glu Gln Arg Gly His Pro Arg Leu Arg Met Arg His Val Asn Tyr
 65 70 75 80
 Pro Ile Gly Val Thr Ala Ala Glu Arg Trp Leu Gln Leu Met Ser Asn
 85 90 95
 Ala Leu Asp Gly Val Asp Leu Thr Ala Glu Gln Arg Glu Ala Ile Trp
 100 105 110
 Glu His Met Val Arg Ala Ala Asp Met Leu Ile Asn Ser Asn Pro Asp

115 120 125
Pro His Ala
130

<210> 24
<211> 124
<212> PRT
<213> Synechocystis sp.

<400> 24
Met Ser Thr Leu Tyr Glu Lys Leu Gly Gly Thr Thr Ala Val Asp Leu
1 5 10 15
Ala Val Asp Lys Phe Tyr Glu Arg Val Leu Gln Asp Asp Arg Ile Lys
20 25 30
His Phe Phe Ala Asp Val Asp Met Ala Lys Gln Arg Ala His Gln Lys
35 40 45
Ala Phe Leu Thr Tyr Ala Phe Gly Gly Thr Asp Lys Tyr Asp Gly Arg
50 55 60
Tyr Met Arg Glu Ala His Lys Glu Leu Val Glu Asn His Gly Leu Asn
65 70 75 80
Gly Glu His Phe Asp Ala Val Ala Glu Asp Leu Leu Ala Thr Leu Lys
85 90 95
Glu Met Gly Val Pro Glu Asp Leu Ile Ala Glu Val Ala Ala Val Ala
100 105 110
Gly Ala Pro Ala His Lys Arg Asp Val Leu Asn Gln
115 120

<210> 25
<211> 183
<212> PRT
<213> Synechococcus sp.

<400> 25
Met Asp Val Ala Leu Leu Glu Lys Ser Phe Glu Gln Ile Ser Pro Arg
1 5 10 15
Ala Ile Glu Phe Ser Ala Ser Phe Tyr Gln Asn Leu Phe His His His
20 25 30
Pro Glu Leu Lys Pro Leu Phe Ala Glu Thr Ser Gln Thr Ile Gln Glu
35 40 45
Lys Lys Leu Ile Phe Ser Leu Ala Ala Ile Ile Glu Asn Leu Arg Asn
50 55 60
Pro Asp Ile Leu Gln Pro Ala Leu Lys Ser Leu Gly Ala Arg His Ala
65 70 75 80
Glu Val Gly Thr Ile Lys Ser His Tyr Pro Leu Val Gly Gln Ala Leu
85 90 95
Ile Glu Thr Phe Ala Glu Tyr Leu Ala Ala Asp Trp Thr Glu Gln Leu
100 105 110

Ala Thr Ala Trp Val Glu Ala Tyr Asp Val Ile Ala Ser Thr Met Ile
 115 120 125
 Glu Gly Ala Asp Asn Pro Ala Ala Tyr Leu Glu Pro Glu Leu Thr Phe
 130 135 140
 Tyr Glu Trp Leu Asp Leu Tyr Gly Glu Glu Ser Pro Lys Val Arg Asn
 145 150 155 160
 Ala Ile Ala Thr Leu Thr His Phe His Tyr Gly Glu Asp Pro Gln Asp
 165 170 175
 Val Gln Arg Asp Ser Arg Gly
 180

<210> 26

<211> 118

<212> PRT

<213> Nostoc commune

<400> 26

Met Ser Thr Leu Tyr Asp Asn Ile Gly Gly Gln Pro Ala Ile Glu Gln
 1 5 10 15
 Val Val Asp Glu Leu His Lys Arg Ile Ala Thr Asp Ser Leu Leu Ala
 20 25 30
 Pro Val Phe Ala Gly Thr Asp Met Val Lys Gln Arg Asn His Leu Val
 35 40 45
 Ala Phe Leu Ala Gln Ile Phe Glu Gly Pro Lys Gln Tyr Gly Gly Arg
 50 55 60
 Pro Met Asp Lys Thr His Ala Gly Leu Asn Leu Gln Gln Pro His Phe
 65 70 75 80
 Asp Ala Ile Ala Lys His Leu Gly Glu Arg Met Ala Val Arg Gly Val
 85 90 95
 Ser Ala Glu Asn Thr Lys Ala Ala Leu Asp Arg Val Thr Asn Met Lys
 100 105 110
 Gly Ala Ile Leu Asn Lys
 115

<210> 27

<211> 136

<212> PRT

<213> Bacillus megaterium

<400> 27

Met Arg Glu Lys Ile His Ser Pro Tyr Glu Leu Leu Gly Gly Glu His
 1 5 10 15
 Thr Ile Ser Lys Leu Val Asp Ala Phe Tyr Thr Arg Val Gly Gln His
 20 25 30
 Pro Glu Leu Ala Pro Ile Phe Pro Asp Asn Leu Thr Glu Thr Ala Arg
 35 40 45
 Lys Gln Lys Gln Phe Leu Thr Gln Tyr Leu Gly Gly Pro Ser Leu Tyr

50						55					60					
Thr	Glu	Glu	His	Gly	His	Pro	Met	Leu	Arg	Ala	Arg	His	Leu	Pro	Phe	
65					70					75					80	
Glu	Ile	Thr	Pro	Ser	Arg	Ala	Lys	Ala	Trp	Leu	Thr	Cys	Met	His	Glu	
				85					90					95		
Ala	Met	Asp	Glu	Ile	Asn	Leu	Glu	Gly	Pro	Glu	Arg	Asp	Glu	Leu	Tyr	
			100					105					110			
His	Arg	Leu	Ile	Leu	Thr	Ala	Gln	His	Met	Ile	Asn	Ser	Pro	Glu	Gln	
		115					120					125				
Thr	Asp	Glu	Lys	Gly	Phe	Ser	His									
130						135										




Figure 5

Figure 6

SEQ ID NO: 28

```

atggagtttg tggcccgta gtcacgaat gctgtccct ttgtcaggtc aactttctacc      60
caccatttga agaagttggc agcaaacagt tctctagctg ctactgctag tcattgtccc      120
gtggttggcc ctgctctcca acagcagaga tactactctc aaccttccaa gccagcccaa      180
gcccaaacct cggacattgc tactgggato aagaaggatg tttctccgat cgtatggac      240
tctaataaaa ccgcttttga ttacaatgga atgtatgagt ctgactcttg gaataaacgt      300
aaagataact cgtatcgtta tttcaataac atcaaccgtc tagccaagga gtttcccaag      360
gcacatcgcc agaccgaaga tgacaagggt accgtctggt gctctaacga ctacttagga      420
atgggtaggc atcctgagat tatcaaaacc atgaaggcta ccatggacaa gtacggttcc      480
ggagcaggag gaactaggaa cattgcaggc cataaccacg ccgctatcaa tttgaaagc      540
gagttggctt gcttgaacaa gaaggaagcg gctctggtgt tttcatcatg tttcatagct      600
aacgatgcaa tcctctcgtt gttgggacaa aaaatcaaaa atttggtcac tttctctgac      660
cagtcgaatc atgcttccat gatattgggt gtgcgtaact ccaaagcgaa gaagcacatc      720
ttcaagcaca acaatttgaa ggatctggag tcgcagttag ctacgtaccc caagtcgact      780
cctaaactga tggccttcga gtcagtttac tctatgtgtg gatctgtggc tcccattgag      840
aagatttgcg atttggctaa aaggtacggc gccctcacct tcttgatga agttcatgct      900
gttggaatgt atggctctca tggacagggt gtagctgagc atttggactt tgatctgcat      960
ttacagcttg gaatcgccag tctagcgtg gtggacaaac gcaccatatt ggatcgtgac      1020
gacatgatta ctggtacttg cggaaagtca tttggtactg ttggaggtta cgttgcgtgt      1080
agtgcacacc taattgattg gtttaagatc tatgcgccag gtttcatttt cactaccaca      1140
cttcctctct ctatcatggc tggtagcggc acttctgttc gtattgttag ggccgacatt      1200
gaggcccgta tcaagcaaca gcttaatact cgctacgtca aagactcatt tgaaaacctt      1260
ggtattccag tcattccaaa cccaagtcac attgttctctg ttctagttgg aaatgctgca      1320
gatgccaaaga aggcattcga tatgttaatg aacaaacacc gtatttatgt tcaagctatt      1380
aactacccta ctgtgctgtg cgggtgaagaa cgactaagga ttactcctac tccaggtcat      1440
ggaaaggaga tttgtgacca gctgacgagc gctgtcgagc atgtttttac tgagcttaat      1500
ttaccaagaa tcaacaaatg gcagtcacca ggtggtcatt gcggtgttgg tgatgctaat      1560
tacgtaccag aaccgaatct gtggactcag gaccagctca gcttgacaaa ccaagacttg      1620
cactccaatg tgcacaaccc agtgattgag cagatcgaaa cctcatcagg agtcagattg      1680
tag                                                                 1683

```

SEQ ID NO: 29

```

      10      20      30      40      50
MEFVARQSMN ACPFVRSTST HHLKKLAANS SLAATASHCP VVGPAQQQR
      60      70      80      90     100
YYSQPSKPAQ AQTSDIATGI KKDVSPIRMD SNETAFDYNG MYESDLANKR
     110     120     130     140     150
KDNSYRYFNN INRLAKEFPK AHRQTEDDKV TVWCSNDYLG MGRHPEIIKT
     160     170     180     190     200
MKATMDKYGS GAGGTRNIAG HNHAALNLES ELACLNKKEA ALVFSSCFIA
     210     220     230     240     250
NDAIISLLGQ KIKNLVIFSD QSNHASMILG VRNSKAKKHI FKHNNLKDLE
     260     270     280     290     300
SQLAQYPKST PKLIAFESVY SMCGSVAPIE KICDLAKRYG ALTFLDEVHA
     310     320     330     340     350
VGMYGPHGQG VAEHLDFDLH LQSGIASPSV VDKRTILDRV DMITGTGCKS
     360     370     380     390     400
FGTVGGYVAG SANLIDWLRS YAPGFIFTTT LPPAIMAGTA TSVRIVRADI
     410     420     430     440     450

```

EARIKQQLNT	RYVKDSFENL	GIPVIPNPSH	IVPVLVGNA	DAKKASDMLM
460	470	480	490	500
NKHRIYVQAI	NYPTVPVGEE	RLRITPTPGH	GKEICDQLIS	AVDDVFTELN
510	520	530	540	550
LPRINKWQSQ	GGHCGVGDAN	YVPEPNLWTQ	DQLSLTNQDL	HSNVHNPVIE
560				
QIETSSGVRL				

Figure 7

Sequence of an exemplary mutant ALAS gene and protein

Gene sequence (SEQ ID NO: 30):

ATGGAGTTTGTGCGCCCGTCAGTCCATGAATGCCTCTCCCTTTGTCAGGTCAACTTCTACCCACC
ATTTGAAGAAGTTGGCAGCAAACAGTTCTCTAGCTGCTACTGCTAGTCATTCTCCCGTGGTTGG
CCCTGCTCTCCAACAGCAGAGATACTACTCTCAACCTTCCAAGCCAGCCCAAGCCCAAACCTCC
GACATTGCTACTGGGATCAAGAAGGATGTTTCTCCGATCCGTATGGACTCTAATGAAACCGCCT
TTGATTACAATGGAATGTATGAGTCTGATCTTGCGAATAAACGTAAAGATAACTCGTATCGTTA
TTTCAATAACATCAACCGTCTAGCCAAGGAGTTTCCCAAGGCACATCGCCAGACCGAAGATGAC
AAGGTGACCGTCTGGTGTCTAACGACTACTTAGGAATGGGTAGGCATCCTGAGATTATCAAAA
CCATGAAGGCTACCATGGACAAGTACGGTTCGGGAGCAGGAGGAAGTAACTAGGAACATTGCAGGTCA
TAACCACGCCGCTATCAATTTGGAAAGCGAGTTGGCTTGCTTGAACAAGAAGGAAGCGGCTCTG
GTGTTTTTCATCATGTTTCATAGCTAACGATGCAATCATCTCGTTGTTGGGACAAAAAATCAAAA
ATTTGGTCATTTTCTCTGACCAGTCGAATCATGCTTCCATGATATTGGGTGTGCGTAACTCCAA
AGCGAAGAAGCACATCTTCAAGCACAACAATTTGAAGGATCTGGAGTCGCAGTTAGCTCAGTAC
CCCAAGTCGACTCCTAAACTGATCGCCTTCGAGTCAGTTTACTCTATGTGTGGATCTGTGGCTC
CCATTGAGAAGATTTGCGATTTGGCTAAAAGGTACGGTGCCCTCACCTTCTTGGATGAAGTTCA
TGCTGTTGGAATGTATGGTCCTCATGGACAGGGTGTAGCTGAGCATTGGAAGTTTGTATCTGCAT
TTACAGTCTGGAATCGCCAGTCCTAGCGTGGTGGACAAACGCACCATATTGGATCGTGTGCGACA
TGATTACTGGTACTTGCGGAAAGTCATTTGGTACTGTTGGAGGTTACGTTGCTGGTAGTGCCAA
CCTAATTGATTGGTTAAGATCCTATGCGCCAGGTTTCATTTTCACTACCACACTTCCTCCTGCT
ATCATGGCTGGTACAGCCACTTCTGTTTCGTATTGTTAGGGCCGACATTGAGGCCCGTATCAAGC
AACAGCTTAATACTCGCTACGTCAAAGACTCATTTGAAAACCTTGGTATTCCAGTCATTCCAAA
CCCAAGTCACATTGTTTCTGTTCTAGTTGGAAATGCTGCAGATGCCAAGAAGGCATCCGATATG
TTAATGAACAAACACCGTATTTATGTTCAAGCTATTAATACTACCCTACTGTGCCTGTCGGTGAAG
AACGACTAAGGATTACTCCTACTCCAGGTCATGGAAAGGAGATTTGTGACCAGCTGATCAGCGC
TGTCGACGATGTTTTTACTGAGCTTAATTTACCAAGAATCAACAAATGGCAGTCCCAAGGTGGT
CATTGCGGTGTTGGTGATGCTAATTACGTACCAGAACCCAATCTGTGGACTCAGGACCAGCTCA
GCTTGACAAACCAAGACTTGCACTCCAATGTGCACAACCCAGTGATTGAGCAGATCGAAACCTC
ATCAGGAGTCAGATTGTAG

Mutated codons (TGT in wild-type to TCT in mutant) are indicated in bold and double underlined.

Protein sequence (SEQ ID NO: 31):

MEFVARQSMNA**S**PFVVRSTSTHHLKKLAANSSLAATASH**S**PVVGPALQQQRYYSQPSKPAQAQTS
 DIATGIKKDVSPIRMDSNETAFDYNGMYESDLANKRKDNSYRYFNNINRLAKEFPKAHRQTEDD
 KVTVWCSDNYLGMGRHPEIIKTMKATMDKYGSGAGGTRNIAGHNHAAINLESELACLNKKEAAL
 VFSSCFIANDAIISLLGQKIKNLVIFSDQSNHASMILGVRNSKAKKHIFKHNNLKDLESQLAQY
 PKSTPKLIAFESVYSMCGSVAPIEKICDLAKRYGALTFLDEVHAVGMYGPHGQGVAEHLDFDLH
 LQSGIASPSVVDKRTILDRVDMITGTCGKSFGTVGGYVAGSANLIDWLRSYAPGFIFTTTLPPA
 IMAGTATSVRIVRADIEARIKQQLNTRYVKDSFENLGIPVIPNPSHIVPVLVGNAADAKKASDM
 LMNKHRIYVQAINYPTVPVGEERLRITPTPGHGKEICDQLISAVDDVFTELNLPRINKWQSQGG
 HCGVGDANYVPEPNLWTQDQLSLTNQDLHSNVHNPVIEQIETSSGVRL

Length: 560 aa

Mol wt: 61.5 kDa

Two cysteine residues present in wtALAS were mutated to serine (bold and double underlined):
 C12S and C39S.

Figure 8

SEQ ID NO: 32

R/L/N/A/C/S/H/I/G/Q-C-P-L/V/I/F/C-L/M/P/V

SEQ ID NO: 33

A-C-P-F-V

SEQ ID NO: 34

H-C-P-V-V

SEQ ID NO: 35

I-C-P-F-M

SEQ ID NO: 36

G-C-P-V-V

SEQ ID NO: 37

CGTCAGTCCATGAATGCCTCTCCCTTTGTCAGGTCAACTTC

SEQ ID NO: 38

GAAGTTGACCTGACAAAGGGAGAGGCATTCATGGACTGACG

SEQ ID NO: 39

GCTGCTACTGCTAGTCATTCTCCCGTGGTTGGCCCTG

SEQ ID NO: 40

CAGGGCCAACCACGGGAGAATGACTAGCAGTAGCAGC

SEQ ID NO: 41

AAACGCTGTCTTGGAACCTAATATGAC

SEQ ID NO: 42

GACGGGCGACAACTCCATCGTTTCGAATAATTAGTTG

SEQ ID NO: 43

CAACTAATTATTCGAAACGATGGAGTTTGTCGCCCCGTCAG

SEQ ID NO: 44

AATTAAATACATTTCAACTACAATCTGACTCCTGATGAGGTTTCG

SEQ ID NO: 45

CCTCATCAGGAGTCAGATTGTAGTTGAAATGTATTTAATTG

SEQ ID NO: 46

AAACTGTCAGTTTTGGGCCATTG

SEQ ID NO: 47

GCTCTCCAACAGCAGAGATAC

SEQ ID NO: 48

GTCCATACGGATCGGAGAAAC

SEQ ID NO: 49

AGCAACATCCCTGATTCCG

SEQ ID NO: 50

ATGCGTACCTTCAATCCTGG

SEQ ID NO: 51

6-FAM/AAGCCCAAA/Zen/CCTCCGA CATTGCTA/3IABkFQ

SEQ ID NO: 52

HEX/TCGCCGTAA/Zen/GTTCTTGGTT TAGACGTTC/3IABkFQ

SEQ ID NO: 53

TAGCGCAGTCTCTCTATCGCTTC

SEQ ID NO: 54

CACTGGGTTGTGCACATTGG

SEQ ID NO: 55

ACAATATTCTTCTCTGCCGC

SEQ ID NO: 56

TTGATCTCGTCAAGAATGCG

SEQ ID NO: 57

TAGGTGCCACAACCTTTTGGTTTC

SEQ ID NO: 58

GATCCAATGCGATGACATTCTTGT

SEQ ID NO: 59

ACCTGCAATAACTCCTCTTCTCTG

SEQ ID NO: 60

CCACTGAGGGTAGCCGAATC

SEQ ID NO: 61

GGGCTCTGAAAACTCTTTTGG

SEQ ID NO: 62
GCATGTCTCAATAACAGATCTCGACGG

SEQ ID NO: 63
AAGCCTCTTGTTTTTCTGTAAATGCAC

SEQ ID NO: 64
TGATGGCGTCCGAGATGAACTC

SEQ ID NO: 65
ATGGAGTTTGTCGCCCCGTCAG

SEQ ID NO: 66
CTACAATCTGACTCCTGATGAGGTTTC