



(86) Date de dépôt PCT/PCT Filing Date: 1999/05/24
(87) Date publication PCT/PCT Publication Date: 1999/12/02
(45) Date de délivrance/Issue Date: 2011/07/26
(85) Entrée phase nationale/National Entry: 2000/11/24
(86) N° demande PCT/PCT Application No.: JP 1999/002713
(87) N° publication PCT/PCT Publication No.: 1999/061032
(30) Priorité/Priority: 1998/05/25 (JP10/142763)

(51) Cl.Int./Int.Cl. *A61K 48/00* (2006.01),
A61K 31/70 (2006.01), *A61K 31/7088* (2006.01),
A61K 47/16 (2006.01), *A61K 9/10* (2006.01),
A61K 9/19 (2006.01), *C07H 21/00* (2006.01),
C07H 21/02 (2006.01)

(72) Inventeurs/Inventors:
SUGIHARA, KATSUHIRO, JP;
SEKI, JUNZO, JP;
HIRABAYASHI, KAZUKO, JP

(73) Propriétaire/Owner:
NIPPON SHINYAKU CO., LTD., JP

(74) Agent: SIM & MCBURNEY

(54) Titre : PROCEDE DE PRODUCTION D'UNE PREPARATION COMPOSITE CONTENANT DE L'ACIDE NUCLEIQUE
(54) Title: PROCESS FOR PRODUCING COMPOSITE PREPARATION CONTAINING NUCLEIC ACID

(57) **Abrégé/Abstract:**

A process for producing a homogeneous nucleic-acid-containing composite preparation of good quality which is capable of sterilization by filtration and is characterized by containing no coarse particles having a size of 7 µm or larger, which are regarded as unsafe for the human body. The process for producing the preparation, containing a composite of a cationic carrier with a nucleic acid polymer, is characterized in that two single-stranded nucleic acid polymers capable of at least partly forming a double-stranded state are separately added, each in the single-stranded state, to either a cationic carrier or a material from which the cationic carrier is to be formed, and all these ingredients are subjected to a dispersing treatment.



PCT

特許協力条約に基づいて公開された国際出願



(51) 国際特許分類6 A61K 31/70, 47/16, 47/24, 9/10 // C07H 21/02	A1	(11) 国際公開番号 WO99/61032 (43) 国際公開日 1999年12月2日(02.12.99)
(21) 国際出願番号 PCT/JP99/02713 (22) 国際出願日 1999年5月24日(24.05.99) (30) 優先権データ 特願平10/142763 1998年5月25日(25.05.98) JP (71) 出願人 (米国を除くすべての指定国について) 日本新薬株式会社(NIPPON SHINYAKU CO., LTD.)(JP/JP) 〒601-8550 京都府京都市南区吉祥院西ノ庄門口町14番地 Kyoto, (JP) (72) 発明者 ; および (75) 発明者 / 出願人 (米国についてのみ) 杉原勝広(SUGIHARA, Katsuhiko)(JP/JP) 〒617-0844 京都府長岡京市調子一丁目11-4 Kyoto, (JP) 関 純造(SEKI, Junzo)(JP/JP) 〒567-0032 大阪府茨木市西駅前町13番11号 Osaka, (JP) 平林加壽子(HIRABAYASHI, Kazuko)(JP/JP) 〒616-8105 京都府京都市右京区太秦森ヶ前町11-1 グリーンパレス蚕の社307 Kyoto, (JP)		(81) 指定国 CA, CN, JP, KR, RU, US, 欧州特許 (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), ユーラシア特許 (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM) 添付公開書類 国際調査報告書
(54) Title: PROCESS FOR PRODUCING COMPOSITE PREPARATION CONTAINING NUCLEIC ACID (54) 発明の名称 核酸含有複合体製剤の製造方法 (57) Abstract A process for producing a homogeneous nucleic-acid-containing composite preparation of good quality which is capable of sterilization by filtration and is characterized by containing no coarse particles having a size of 7 μm or larger, which are regarded as unsafe for the human body. The process for producing the preparation, containing a composite of a cationic carrier with a nucleic acid polymer, is characterized in that two single-stranded nucleic acid polymers capable of at least partly forming a double-stranded state are separately added, each in the single-stranded state, to either a cationic carrier or a material from which the cationic carrier is to be formed, and all these ingredients are subjected to a dispersing treatment.		



DESCRIPTION

PROCESS FOR PRODUCING COMPOSITE PREPARATION CONTAINING
NUCLEIC ACID

Technical Field

The present invention relates to complex preparations which contain a complex of cationic carriers and nucleic acid polymers. The complex is sometimes called lipoplex.

As used herein, the term "cationic carrier" refers to drug carriers having positive charges in water and effective for transferring drugs, especially transferring anionic drugs into cells. Cationic carriers have been recently studied extensively as drug carriers for transfer of genes and RNA such as poly I:C into cells.

"Nucleic acid polymers" refer to naturally occurring or synthesized or semi-synthesized polynucleotides (DNA, RNA), and naturally occurring or synthesized or semi-synthesized oligonucleotides.

Background Art

The nucleic-acid-containing complex preparations which contain a complex of a cationic carrier and a polyanionic double strand nucleic acid polymer having a double helical structure

can be produced by only mixing the cationic carrier and the double strand nucleic acid polymer.

However, in the nucleic-acid-containing complex preparations produced by this method, their particles are generally coarse with diameters of some micrometers to some hundred micrometers, and are heterogeneous. The nucleic-acid-containing complex preparations with these coarse and heterogeneous particles have disadvantages wherein homogeneous data are not sufficiently obtained in studies on intracellular transfers and signal expressions of nucleic acid polymers. The most critical problems concerning particle coarseness are that sterilization is difficult on an industrial scale and that embolizations and the like may occur in injection needles or capillaries in the case of intravenous administrations despite pharmaceuticals being prepared on assumption that they are safe for administration to humans. These problems are difficult to be solved not only by methods in which the complex preparations are produced only by mixing as described above but also in production methods by applying a dispersion process using appropriate emulsifying dispersion devices.

Aggregation of the particles is also a problem resulting from freeze-drying being attempted for the stabilization of

the complex preparations.

Nucleic acid polymers, in the complex preparations are desirably at high concentrations in order to decrease the dosages and reduce the burden to patients and medical workers at the time of administration as well as in order to achieve productive efficacy of the complex preparations. However, by conventional processes when the total amount of nucleic acid polymers is 0.1mg/mL or more, especially 0.5mg/mL or more, remarkable aggregation occurs under the production process and huge precipitates or suspended solids, easily confirmed with the naked eye, are yielded. These are incapable of being dispersed sufficiently by any dispersion process.

Conventionally, double strand RNA having double helical structures have been commonly employed as genes and RNA such as poly I:C from the view of their physiological features and stability for various nucleases. For example, it has been known that sufficient pharmacological efficacy is not obtained by the separate administration of poly I and poly C instead of poly I:C which has physiological activities such as a strong induction potency of interferon and immunopotentiating actions (Archives of Virology , 51, 199-215 (1976)). Thus, double strands having double helical structures have been believed to be essential for genes and RNA such as poly I:C.

For nucleic-acid-containing complex preparations which contain a complex of a cationic carrier and nucleic acid polymers, necessity for double helical structures has not been discussed at all, and double strand DNA and double strand RNA having the double helical structure have been commonly employed in production procedures of the complex preparations.

The present applicants have applied for the patent for the nucleic-acid-containing complex preparations as nuclease activating preparations in cancer cells since nucleic-acid-containing complex preparations consist of cationic carriers and double strand RNA such as poly I:C activate nucleases in cancer cells which are effective for the treatment of cancers, and have already applied for the patent for the nucleic-acid-containing complex preparations as therapeutic agents for hepatitis since they induce effective amounts of interferons specifically for the liver and spleen for a long time (PCT/JP98/04695, PCT/JP99/01438).

Summary of the Invention

According to one aspect of the present invention there is provided a production method for homogenous nuclear acid containing complex preparations with good quality characterized in that the preparations are capable of being

done with so-called sterilizing filtration and do not contain coarse particles of 7 μm or more which are considered to have a safety problem for human body.

The present inventors were the first to discover that the above problems can be solved with one effort without having an effect on their pharmacological activities by producing using nucleic acid polymers with a single strand which has been separated from a double helical structure or initially formed without a double strand structure, without using double strand DNA or double strand RNA usually having double helical structures, in a production process of nucleic-acid-containing complex preparations which contain a cationic carrier and nucleic acid polymers, and have thus completed the present invention.

Therefore, the present invention can include a production method of the nucleic-acid-containing complex preparations characterized in that two single strand nucleic acid polymers which can at least partly form double strands are separately added, in a single strand form, to a cationic carrier or source materials before forming a cationic carrier, and the two single strand nucleic acid polymers are processed to be dispersed in the production processes of the said nucleic-acid-containing complex preparation which contains a cationic carrier and

nucleic acid polymers (referred to as "nucleic-acid-containing complex preparations" hereinafter).

In accordance with a further aspect of the present invention there is provided a method of producing a nucleic-acid-containing complex preparation, comprising the following steps:

- 1) separately mixing two single-strand nucleic acid polymers, capable of forming a double strand, with a cationic carrier or with source materials for the cationic carrier before the cationic carrier is formed, and

- 2) performing an emulsifying dispersion process on the mixture.

In accordance with an aspect of the present invention there is provided a method of producing a nucleic-acid-containing complex preparation, comprising the following steps:


- 1) separately mixing two single-strand nucleic acid polymers, that form a double strand, with a cationic carrier or with source materials for the cationic carrier before the cationic carrier is formed thus forming a mixture, and

- 2) performing an emulsifying dispersion process on the mixture to provide said preparation.

Detailed Description of the Invention

The present invention will be described in detail below.

"Cationic carriers" applicable to the present invention can include drug carriers disclosed in PCT W094/19314 such as 2-o-(2-diethylaminoethyl) carbamoyl-1,3-o-dioleoyl glycerol (referred to as "compound A" hereinafter) represented as the following structural formula [1] and drug carriers formed by phospholipids as essential component and drug carriers such as polylysine, in addition to commercially available lipofectin (brand name), lipofectamine (brand name), lipofectace (brand name), and DMRIE-C (brand name).



"Two single strand nucleic acid polymers" applicable to the present invention can include, but are not especially limited to them if they are two single strand nucleic acid polymers which can at least partly form double strands, for example, two single strand DNA and RNA which construct natural genes or artificially modified genes (e.g. plasmid), and two single strand RNA such as poly I and poly C,; poly I and poly C₁₂U, poly I with partially chemical medication (e.g. poly

(7-deazainosinic acid)) and poly C, poly I and poly C with partially chemical modification (e.g. poly(bromocytidylic acid), poly(thiocytidylic acid)). The present invention can be applied to two single strand RNA such as poly I and poly C which construct poly I:C with physiological activities such as a strong induction potency of interferons. Herein used "poly I", "poly C", "poly A", "poly U", and "poly C₁₂U" mean polyinosinic acid, polycytidylic acid, polyadenylic acid, polyuridylic acid, and copolymer of cytidylic acid and uridylic acid where one uridylic acid is substituted for about every 12 cytidylic acids, respectively.

"Can at least partly form double strands" refers to those that complementary bases at a degree where they can form double strands in physiological conditions exist in two single strand nucleic acid polymers, and the degree varies depending on kinds of the two single strand nucleic acid polymers and the length of each nucleic acid polymer, and a concrete generality is that the numbers of complementary bases are 20 or more.

The numbers of bases which each single strand nucleic acid polymer has are, but are not especially limited thereto, suitably 10,000 or less, and preferably 2,000 or less. The numbers of the bases can be appropriately selected depending on the natures which each nucleic acid polymer has. Besides,

the two single strand nucleic acid polymers are not necessarily composed of the same number of bases. Each nucleic acid polymer usually exists with a definite distribution composed of a various numbers of bases, but each number of bases means the number of bases for a distributed maximum and herein refers to an "average base number".

Further, for example, the average base number of poly I and poly C in the present invention can be determined based on a balance between efficacy and safety. Specifically, it is suitable in the range of 30 to 3,000 bases, preferably in the range of 60 to 2,000 bases, and more preferably in the range of 100 to 500 bases.

Phospholipids in drug carriers (cationic carriers) formed by the above compound A and phospholipids as essential components are not limited if they are pharmaceutically acceptable. For example, they include phosphatidyl choline, phosphatidyl ethanolamine, phosphatidyl inositol, phosphatidyl serine, sphingomyelin, and lecithin. Also hydrogenated phospholipids can be included. Preferred phospholipids can include egg yolk phosphatidyl choline, egg yolk lecithin, soy lecithin, and egg yolk phosphatide. Two or more kinds of phospholipids can be used. Yet, among these cationic carriers, phosphatidyl choline or lecithin is

superior than phosphatidyl ethanolamine generally used in cationic carriers.

The component ratio of phospholipids to the compound A is varied depending on kinds of the phospholipids and kinds of the applied two single strand nucleic acid polymers, however, the phospholipids are appropriately comprised in the range of 0.1 to 10 weight parts, preferably in the range of 0.5 to 5 weight parts, and more preferably in the range of 1 to 2 weight parts per 1 weight parts of the compound A. This ratio is similar when the phospholipid is replaced by lecithin.

The component ratio of nucleic acid polymers to a cationic carrier varies depending on the kind of source of the cationic carrier and kind of nucleic acid polymer used, however, the total amount of nucleic acid polymers is appropriately in the range of 0.05 to 10 weight parts, preferably in the range of 0.1 to 4 weight parts, and more preferably in the range of 0.5 to 2 weight parts per 10 weight parts of the cationic carrier. Similarly, when the complex is formed by poly I and poly C with the cationic carrier made of the compound A and phospholipids as essential components, provided that the two single strand nucleic acid polymers are poly I and poly C, the total amount of poly I and poly C is appropriately in the range of 0.05 to 10 weight parts, preferably in the range of 0.1 to 4 weight

parts, and more preferably in the range of 0.5 to 2 weight parts per 10 weight parts of the cationic carrier.

The nucleic-acid-containing complex preparations of the present invention (referred to as "present invention preparations" hereinafter) can be produced, for example, by a dispersion process by conventional methods using appropriate emulsifying dispersion machines, after adding the two single strand nucleic acid polymers sequentially or simultaneously to an aqueous solution in which a commercially available cationic carrier is dispersed, or to an aqueous solution in which a cationic carrier produced by a dispersion process with a conventional method of an aqueous solution processed using an appropriate emulsifying dispersion machine in which source materials before making a cationic carrier are dispersed, or to an aqueous solution in which source materials before forming a cationic carrier are dispersed. The present invention preparations can be also produced by processes wherein the two single strand nucleic acid polymers are added to a solid cationic carrier or its source materials and water is added and then the mixture is dispersed by an appropriate emulsifying dispersion machine. The sequences to be added, volumes to be added, concentrations to be added, and concentrations of cationic carriers and their source materials in solutions are

optionally selected and are not especially limited in the present invention.

More specifically, when used the cationic carrier made of the compound A and phospholipids as essential components, the present invention preparation can be produced by the emulsifying dispersion process of the mixture formed by gradually dripping the aqueous solutions of poly I and poly C separately into the aqueous solution in which the cationic carrier is dispersed. Also the present invention preparation can be produced by the process wherein the compound A, phospholipid, poly I and poly C are weighted and taken into a beaker, and are roughly dispersed with a homogenizer after the addition of water, and then dispersed with a pressurized emulsifying dispersion machine.

Two single strand nucleic acid polymers obtained by separating a double strand nucleic acid polymer by conventional manipulations may be used. Specifically, the manipulations can include non-enzymatic treatments such as heating at 60°C or more, or enzymatic treatments.

The above-described commercially available cationic carriers can be used without processing or with appropriate processing.

The above aqueous solutions can include injectable water,

injectable distilled water, and electrolyte solutions such as saline, and glucose solution.

The above emulsifying dispersion machines can include, for example, a homomixer, homogenizer, ultrasonic dispersing machine, ultrasonic homogenizer, high-pressurized emulsifying dispersion machine, Microfluidizer (brand name), Nanomizer (brand name), Ultimizer (brand name), DeBEE2000 (brand name), and Manton-Gaulin type high-pressurized homogenizer, however those appropriately employed for medical uses are sufficient. Processing conditions and time periods, and processing temperatures are selected appropriately.

The present invention preparations may contain pharmaceutically acceptable additives, for example, emulsifying dispersion auxiliary, stabilizer, isotonic agent, lyoprotectant, and pH adjuster at appropriate amounts. Specifically, those can include fatty acids having from 6 to 22 of carbon atoms (e.g. caprylic acid, capric acid, lauric acid, myristic acid, palmitic acid, stearic acid, oleic acid, linoleic acid, arachidonic acid, and docosahexaenoic acid), pharmaceutically acceptable salts thereof (e.g. sodium salts, potassium salts, and calcium salts), emulsifying dispersion auxiliaries such as albumin and dextran, stabilizers such as cholesterol and phosphatidic acid, isotonic agents such as

sodium chloride, glucose, maltose, lactose and sucrose, lyoprotectants, and pH adjusters such as hydrochloric acid, sulfuric acid, phosphoric acid, acetic acid, sodium hydroxide, potassium hydroxide, and triethanolamine.

The above-mentioned pharmaceutically acceptable optional additives may be added in appropriate processes before or after dispersion.

After dispersion, the preparations can be optionally filtrated through a sterilizing filtration membrane of 0.2 μm , and then packed into ampoules and vials. Particle diameters of almost all of the present invention preparations are 200 nm or less. Therefore, approximately 100% of the present invention preparations can pass through the 0.2 μm sterilizing filtration membrane.

By the above-mentioned production methods according to the present invention, the nucleic-acid-containing complex preparations which contain homogenous and fine complex particles can be obtained, and further, the nucleic-acid-containing complex preparations which contain nucleic acid polymers in the solution at 0.1 mg/mL or more can be obtained.

Therefore, the nucleic-acid-containing complex preparations obtained by the above-mentioned production methods and the nucleic-acid-containing complex preparations

obtained by the above production methods and comprising nucleic acid polymers in concentrations in the solution in a range of 0.1 to 10 mg/mL, in a range of 0.5 to 10 mg/mL, in a range of 1 to 10 mg/mL or in a range of 2 to 10 mg/mL can also be included in the present invention. Yet, the present invention does not exclude nucleic acid polymers with concentrations in solutions at 10 mg/mL or more.

Additionally, if the present invention preparations, produced by dispersion as described above, are freeze-dried, these can be freeze-dried preparations of the present invention. Therefore, the freeze-dried preparations can be also included in one of the present invention preparations. Freeze-drying can be carried out by conventional methods.

The freeze-dried preparations of the present invention are, for example, freeze-dried by preliminarily freezing in conditions at about -40 to -20°C for 2 hours after dispensing into vials , followed by primary drying from about 0 to 10°C under reduced pressure, and then secondary drying from about 15 to 25°C under reduced pressure. And generally, the inside of the vials are filled with nitrogen gas, and the vials are plugged to obtain the freeze-dried preparations of the present invention. When applied freeze-drying, the use of lyoprotectants which form freeze-dried cakes is preferred.

Especially, saccharides are suitable and disaccharides, in particular, maltose is the most preferable among others.

The freeze-dried preparations of the invention can be generally reconstituted by the addition of an optional appropriate solution (reconstituting solutions) and used. These reconstituting solutions can include injectable water, glucose solution, electrolyte solutions such as saline, and the other infusion solutions. Fluid volumes of these reconstituting solutions vary depending on use and are not especially limited, but 0.5 to 2 folds of the fluid volume before drying or 500 mL or less are appropriate.

The present invention preparations can be provided in the forms of liquid formulations such as injectable and drip drugs or in the form of freeze-dried preparations.

The present invention preparations can be administered to animals including humans by various routes of administration such as intravenously, intra-arterially, subcutaneously, intra-muscularly, by inhalation, nasal, ophthalmically, orally, and rectally. The dosing forms and dosages can be appropriately selected according to desire. Besides, the preparations can be also used as various reagents and medicines for the cultured cells of animals, plants, mycetes, and bacteria.

Best Mode for Carrying Out the Invention

Hereinafter, the present invention will be described in more detail by representing the examples, comparative examples and study examples.

The single strand RNA of poly I and poly C each with an average base number of approximately 200 were used as sources in Example 1 to 5. Besides, filtrated sterilization could be carried out successfully without clogging the filters during filtered sterilization with a 0.2 μ m membrane filter after the dispersion process in each Example, and the yields of filtrates were in the range of 98 to 102% for all nucleic acid polymers, indicating potential sterilization at approximately 100%.

Example 1

Forty grams of maltose dissolved in 100 mL of injectable water was added to 2 g of the compound A and 2 g of purified egg yolk lecithin, mixed by agitation, and dispersed for 5 minutes by means of a homogenizer to obtain a crude dispersion solution of a cationic carrier. The crude dispersion solution was further dispersed for an hour using a small emulsifying dispersion machine for experiments, and compensated with injectable water to up to 250 mL to obtain the dispersion

solution of the cationic carrier. Seventy five mL of aqueous solution containing 250 mg of poly I was added to 250mL of this dispersion solution with agitation, then 75 mL of aqueous solution containing 250 mg of poly C was added with agitation, and the mixture was further dispersed for an hour using a small emulsifying dispersion machine for experiments, followed by sterilization by filtration with a 0.2 μm membrane filter to obtain the present invention preparation. The average particle diameter of complex particles in this present invention preparation was 138 nm by measurement using a particle diameter measuring apparatus (DLS-700, manufactured by Otsuka Electronics Inc. hereinafter, the same is applied) employing a dynamic light scattering method. Besides, no particle with a diameter of 1 μm or more was included.

Thereafter, 1 mL of this present invention preparation was dispensed into each vial and processed into a freeze-dried preparation according to conventional methods. The obtained freeze-dried preparation was reconstituted by the addition of 0.9 mL of injectable water. The average particle diameter of complex particles in the reconstituted present invention preparation was 140 nm by measurement using the particle diameter measuring apparatus (DLS-700, manufactured by Otsuka Electronics Inc. hereinafter, the same is applied) employing

a dynamic light scattering method. Besides, no particle with a diameter of 1 μm or more was included.

Example 2

Four kg of sucrose dissolved in 10 L of injectable water was added to 50 g of the compound A and 30 g of egg yolk phosphatide, and the mixture was dispersed for 10 minutes by means of a Manton-Gaulin high-pressured homogenizer followed by compensating with injectable water to up to 25 L to obtain a dispersion solution of the cationic carrier. Six L of aqueous solution containing 50 g of poly C was added to 20 L of this dispersion solution with agitation, and then 6 L of aqueous solution containing 50 g of poly I was added with agitation. The pH of this dispersion solution was adjusted to 5.5 using hydrochloric acid, and the dispersion solution was further dispersed for 30 minutes by means of a Manton-Gaulin high-pressured homogenizer followed by sterilization by filtration with a 0.2 μm membrane filter to obtain the present invention preparation. The average particle diameter of complex particles in the present invention preparation was measured to be 150 nm. Besides, no particle with a diameter of 1 μm or more was included.

Thereafter, 20 mL of this present invention preparation

was dispensed into each vial and processed into a freeze-dried preparation according to conventional methods. The obtained freeze-dried preparation was reconstituted by the addition of a commercially available 5% glucose infusion solution (500 mL). The average particle diameter of complex particles in the reconstituted present invention preparation was measured to be 151 nm by means of the particle diameter measuring apparatus employing a dynamic light scattering method. Besides, no particle with a diameter of 1 μm or more was included.

Example 3

Two grams of the compound A, 2g of soy lecithin, 25 mg of poly I and 25 mg of poly C were taken into a beaker, 20 g of glucose dissolved in 100 mL of injectable water was added to the beaker, then the contents in the beaker were mixed by agitation, and dispersed for 5 minutes by means of a homogenizer. The crude dispersion solution was dispersed for an hour using a small high-pressurized emulsifying dispersion machine (800 kg/cm^2) for experiments, and compensated with injectable water to up to 400 mL followed by sterilization by filtration with a 0.2 μm membrane filter to obtain the present invention preparation. The average particle diameter of complex particles in this present invention preparation was 121 nm by

measurement using a particle diameter measuring apparatus employing a dynamic light scattering method. Besides, no particle with a diameter of 1 μm or more was included.

Example 4

Forty grams of maltose dissolved in 100 mL of injectable water was added to 1.2 g of the compound A and 2.0 g of purified egg yolk lecithin, mixed by agitation, and dispersed for 30 minutes by means of a small high-pressurized emulsifying dispersion machine for experiments, and then compensated with injectable water to up to 250 mL to obtain a dispersion solution of the cationic carrier. Seventy five ml of aqueous solution containing 100 mg of poly I and 75 ml of aqueous solution containing 100 mg of poly C were simultaneously gradually dripped into 250 mL of this dispersion with agitation, and further dispersed for 2 hours using a small pressurized emulsifying dispersion machine for experiments (1,100 kg/cm²) followed by sterilization by filtration with a 0.2 μm membrane filter to obtain the present invention preparation. The average particle diameter of complex particles in this present invention preparation was 124 nm by measurement using a particle diameter measuring apparatus employing a dynamic light scattering method. Besides, no particle with a diameter

of 1 μm or more was included.

Beside, when the distribution of particle diameter sizes of this present invention preparation was measured by a particle diameter measuring apparatus (LA-910, Horiba Ltd., hereafter the same is applied) employing a laser diffractive scattering method, the results as shown in Fig.1 was obtained. According to this result, a peak of the distribution of particle diameter sizes was 139 nm, and no coarse particle was detected.

Example 5

Forty grams of maltose dissolved in 100 mL of injectable water was added to 4.8 g of the compound A and 8.0 g of purified egg yolk lecithin, mixed by agitation, dispersed for 30 minutes by means of a small high-pressurized emulsifying dispersion machine for experiments, and then compensated with injectable water to up to 250 mL to obtain a dispersion solution of the cationic carrier. Seventy five ml of aqueous solution containing 400 mg of poly I and 75 ml of aqueous solution containing 400 mg of poly C were simultaneously gradually dripped into 250 mL of this dispersion solution with agitation, and further dispersed for 2 hours using a small high-pressurized emulsifying dispersion machine for experiments (1,100 kg/cm²) followed by sterilization by filtration with

a 0.2 μm membrane filter to obtain the present invention preparation. The average particle diameter of complex particles in this present invention preparation was 138 nm by measurement using a particle diameter measuring apparatus employing a dynamic light scattering method. Besides, no particle with a diameter of 1 μm or more was included.

Example 6

One mL of aqueous solution containing 100 μg of commercially available DNA plasmid vector (pMC1neo) was heated in a water bath at 70°C while its agitation for 3 hours. Two mL of dispersion solution containing 2 mg of commercially available lipofectin (brand name) similarly heated at 70°C was added to this with agitation, and the mixture was dispersed for 10 minutes at 70°C using a probe type ultrasonic dispersion machine followed by sterilization by filtration with a 0.2 μm membrane filter to obtain the present invention preparation. The average particle diameter of complex particles in this present invention preparation was 145 nm by measurement using a particle diameter measuring apparatus employing a dynamic light scattering method. Besides, no particle with a diameter of 1 μm or more was included.

Example 7

The single strand RNA of poly I and poly C both of which average base numbers are about 1,500 were used as the sources.

Forty grams of maltose dissolved in 100 mL of injectable water was added to 1.2 g of the compound A and 2.0 g of purified egg yolk lecithin, mixed by agitation, dispersed for 30 minutes by means of a small high-pressurized emulsifying dispersion machine for experiments, and then compensated with injectable water to up to 250 mL to obtain a dispersion solution of the cationic carrier. Seventy five ml of aqueous solution containing 100 mg of poly I and 75 ml of aqueous solution containing 100 mg of poly C were simultaneously dripped into 250 mL of this dispersion solution by agitation, and further dispersed for 2 hours using a small high-pressurized emulsifying dispersion machine for experiments (1,100 kg/cm²) followed by sterilization by filtration with a 0.2 μ m membrane filter to obtain the present invention preparation. The average particle diameter of complex particles in this present invention preparation was 134 nm by measurement using a particle diameter measuring apparatus employing a dynamic light scattering method. Besides, no particle with a diameter of 1 μ m or more was included.

Example 8

The present invention preparation with the average particle diameter of 130 nm was obtained by the similar processes as those of Example 7 with a dispersion pressure at 800 kg/cm² using 200 mg of poly I with the average base number of about 350 and 200 mg of poly C with the average base number of about 350.

Example 9

The present invention preparation with the average particle diameter of 150 nm was obtained by the similar processes as those of Example 7 with a dispersion pressure at 800 kg/cm² using 200 mg of poly I with the average base number of about 1,450 and 200 mg of poly C with the average base number of about 1,450.

Example 10

The present invention preparation with the average particle diameter of 135 nm was obtained by the similar processes as those of Example 7 with a dispersion pressure at 800 kg/cm² using 400 mg of poly I with the average base number of about 80 and 400 mg of poly C with the average base number of about 80.

Comparative example 1 (production by a conventional method corresponding to Example 4)

Forty grams of maltose dissolved in 100 mL of injectable water was added to 1.2 g of the compound A and 2.0 g of purified egg yolk lecithin, mixed by agitation, and dispersed for 30 minutes by means of a small high-pressurized emulsifying dispersion machine for experiments, and then compensated with injectable water to up to 250 mL to obtain a dispersion solution of the cationic carrier. 150 mL of aqueous solution containing 200 mg of double strand poly I:C having approximately 200 base pairs was gradually dripped into 250 mL of this dispersion solution by agitation, and the mixture was further dispersed for 2 hours using a small high-pressurized emulsifying dispersion machine for experiments (1,100 kg/cm²) to obtain a comparative preparation. The average particle diameter of complex particles in this comparative preparation was 182 nm by measurement using a particle diameter measuring apparatus employing a dynamic light scattering method.

Beside, when a distribution of the particle diameter sizes of this comparative preparation was measured by means of a particle diameter measuring apparatus employing a laser diffractive scattering method in the same way as that in Example

4, the results as shown in Fig.2 were obtained. According to this result, a peak in the distribution of particle diameter sizes was 243 nm, however 20% were detected as coarse particles with 3-20 μm in diameters having a peak of 8000 nm in the distribution, indicating a bimodal distribution of the particles.

Additionally, when an attempt was made to filter this comparative preparation through a 0.2 μm membrane filter, only 50 mL of the preparation passed through the filter resulting in clogging of the filter and the sterilization by filtration was failed.

Comparative example 2 (production by a conventional method corresponding to Example 5)

Forty grams of maltose dissolved in 100 mL of injectable water was added to 4.8 g of the compound A and 8.0 g of purified egg yolk lecithin, mixed by agitation, dispersed for 30 minutes by means of a small high-pressurized emulsifying dispersion machine for experiments, and then compensated with injectable water to up to 250 mL to obtain a dispersion solution of the cationic carrier. One hundred and fifty mL of aqueous solution containing 800 mg of double strand poly I:C having approximately 200 base pairs was gradually dripped into 250

mL of this dispersion solution by agitation, and further dispersed for 2 hours using a small high-pressurized emulsifying dispersion machine for experiments (1,100 kg/cm²) to collect a comparative preparation. This comparative preparation was a white precipitable suspension solution, and precipitated with aggregation within 5 minutes after collection. It was similar to a suspension solution of sake lees. The particle diameters could not be measured because their sizes were above the measuring ranges of a particle diameter measuring apparatus employing a dynamic light scattering method. Their filtrating sterilization through a 0.2 μ m membrane filter was impossible.

Test example 1

Biological activity of the present invention preparation obtained from Example 4 and that of the comparative preparation from Comparative example 1 were evaluated by their suppressive effects on proliferation of uterine cervix cancer cells (Hela S3).

In the experiment, HelaS3 cells were seeded in 96-well plates at concentration of 10^4 cells/well, confirmed to be adhered to the wells after culture for 5 hours or more followed by the adding of each preparation to the culture, then culturing

was continued, and the number of viable cells was counted by MTT method 3 days after the addition of the preparations. Inhibition ratios were obtained from the following formula, and the values of IC_{50} were calculated. The result is shown in Table 1.

$$IC_{50} = [1 - (\text{value of absorbance in the preparation treated-cell group} / \text{value of absorbance in the saline treated-cell group})] \times 100 \%$$

Table 1

	IC_{50} value (ng/mL)
Present invention preparation of Example 4	8.6 ± 2.5
Comparative preparation of Comparative example 1	8.7 ± 1.2

The IC_{50} value was represented as the concentration of total nucleic acid polymers by combining poly I and poly C.

As shown in Table 1, there was no difference in the biological activity between the present invention preparation according to Example 4 and that of the comparative preparation according to the Comparative example 1.

Effects of the Invention

The present invention has, for example, the following effects.

(1) Homogenous nucleic-acid-containing complex preparations of a high quality without coarse complex particles can be produced.

(2) Homogenous nucleic-acid-containing complex preparations of a high quality which do not substantially comprise coarse complex particles can be produced. This effect is more remarkable at higher concentrations of nucleic acid polymers.

(3) When freeze-dried preparations of nucleic-acid-containing complex preparations produced according to the present invention are reconstituted, nucleic-acid-containing complex preparations equivalent to those before freeze-drying can be reconstructed.

(4) Nucleic-acid-containing complex preparations which can pass through a 0.2 μm sterilizing filter with approximately 100% efficacy can be provided.

Brief Description of the Drawings

Fig. 1 shows the distribution of the particle diameters of complex particles in the present invention preparation

according to Example 4. The horizontal axis, left vertical axis and right vertical axis denote particle diameters (μm), frequencies (%) and integrated frequencies (%), respectively.

Fig. 2 shows the distribution of the particle diameters of complex particles in the comparative preparation according to Comparative example 1. The horizontal axis, left vertical axis and right vertical axis denote particle diameters (μm), frequencies (%) and integrated frequencies (%), respectively.

What is claimed is:

1. A method of producing a nucleic-acid-containing complex preparation, comprising the following steps:

1) separately mixing two single-strand nucleic acid polymers, that form a double strand, with a cationic carrier or with source materials for the cationic carrier before the cationic carrier is formed thus forming a mixture, and

2) performing an emulsifying dispersion process on the mixture to provide said preparation.

2. The method of producing the nucleic-acid-containing complex preparation according to Claim 1, wherein the average base numbers of the two single-strand nucleic acid polymers are in the range of 100 to 2,000 respectively.

3. The method of producing the nucleic-acid-containing complex preparation according to Claims 1 or 2, wherein the two single-strand nucleic acid polymers are poly I and poly C.

4. The method of producing the nucleic-acid-containing complex preparation according to any one of Claims 1 to 3, wherein the cationic carrier includes 2-o-(2-diethylaminoethyl)carbamoyl-1, 3-o-dioleoyl glycerol and a phospholipid.

5. The method of producing the nucleic-acid-containing complex preparation according to claim 4 wherein the phospholipid is lecithin.

1/1




FIGURE 1




FIGURE 2