




(86) Date de dépôt PCT/PCT Filing Date: 2015/07/22	(51) Cl.Int./Int.Cl. C07K 14/22(2006.01), A61P 31/04 (2006.01), A61P 37/04 (2006.01)
(87) Date publication PCT/PCT Publication Date: 2016/01/28	
(45) Date de délivrance/Issue Date: 2023/10/31	
(85) Entrée phase nationale/National Entry: 2017/01/19	
(86) N° demande PCT/PCT Application No.: US 2015/041616	
(87) N° publication PCT/PCT Publication No.: 2016/014719	
(30) Priorité/Priority: 2014/07/23 (US62/028,123)	
	(72) Inventeur/Inventor: BEERNINK, PETER T., US
	(73) Propriétaire/Owner: CHILDREN'S HOSPITAL & RESEARCH CENTER AT OAKLAND, US
	(74) Agent: SMART & BIGGAR LP

(54) Titre : VARIANTS DE PROTEINES DE LIAISON AU FACTEUR H ET PROCEDES D'UTILISATION DE CEUX-CI
(54) Title: FACTOR H BINDING PROTEIN VARIANTS AND METHODS OF USE THEREOF



(57) Abrégé/Abstract:

Variant factor H binding proteins that can elicit antibodies that are bactericidal for at least one strain of *Neisseria meningitidis*, compositions comprising such proteins, and methods of use of such proteins, are provided.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(10) International Publication Number

WO 2016/014719 A1

(43) International Publication Date
28 January 2016 (28.01.2016)

(51) International Patent Classification:
C07K 14/22 (2006.01) *A61P 37/04* (2006.01)
A61P 31/04 (2006.01) *A61K 39/095* (2006.01)

(21) International Application Number:
PCT/US2015/041616

(22) International Filing Date:
22 July 2015 (22.07.2015)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
62/028,123 23 July 2014 (23.07.2014) US

(71) Applicant: CHILDREN'S HOSPITAL & RESEARCH CENTER OAKLAND [US/US]; 747 52nd Street, Oakland, California 94609 (US).

(72) Inventor: BEERNINK, Peter T.; 3158 Lippizaner Lane, Walnut Creek, California 94598 (US).

(74) Agent: CHANDRA, Shweta; 1900 University Avenue, Suite 200, East Palo Alto, California 94303 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: FACTOR H BINDING PROTEIN VARIANTS AND METHODS OF USE THEREOF

FIG. 1

(57) Abstract: Variant factor H binding proteins that can elicit antibodies that are bactericidal for at least one strain of *Neisseria meningitidis*, compositions comprising such proteins, and methods of use of such proteins, are provided.



WO 2016/014719 A1



Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*
- *with sequence listing part of description (Rule 5.2(a))*

FACTOR H BINDING PROTEIN VARIANTS AND METHODS OF USE THEREOF**CROSS-REFERENCE**

[0001] This application claims the benefit of U.S. Provisional Patent Application No. 62/028,123, filed July 23, 2014.

INTRODUCTION

[0002] *Neisseria meningitidis* is a Gram-negative bacterium that colonizes the human upper respiratory tract and is responsible for worldwide sporadic and cyclical epidemic outbreaks of, most notably, meningitis and sepsis. The attack and morbidity rates are highest in children under 2 years of age. Like other Gram-negative bacteria, *Neisseria meningitidis* typically possess a cytoplasmic membrane, a peptidoglycan layer, an outer membrane, which together with the capsular polysaccharide constitute the bacterial wall, and pili, which project into the outside environment. Encapsulated strains of *Neisseria meningitidis* are a major cause of bacterial meningitis and septicemia in children and young adults. The prevalence and economic importance of invasive *Neisseria meningitidis* infections have driven the search for effective vaccines that can confer immunity across different strains, and particularly across genetically diverse serogroup B strains with different serotypes or serosubtypes.

[0003] Factor H Binding Protein (fHbp, also referred to in the art as lipoprotein 2086 (Fletcher et al (2004) *Infect Immun* 72:2088-2100), Genome-derived Neisserial antigen (GNA) 1870 (Massignani et al. (2003) *J Exp Med* 197:789-99) or “741”) is an *N. meningitidis* protein that is expressed in the bacterium as a surface-exposed lipoprotein. An important function of fHbp is to bind human complement factor H (fH), which down-regulates complement activation. Binding of fH to the bacterial surface is an important mechanism by which the pathogen survives in non-immune human serum or blood and evades innate host defenses. Recently, genetic variation in the human factor H gene cluster was found to affect susceptibility to developing meningococcal disease (Davila S et al. (2010) *Nat Genetics* doi:10.1038/ng.640). Binding of fH to fHbp is specific for human fH, and several non-human primates and could partially explain why *Neisseria meningitidis* is strictly a human pathogen. fHbp occurs in many natural sequence variants that are designated by identification (ID) numbers as assigned in the fHbp database on the internet at pubmlst(dot)org/neisseria/fHbp.

[0004] There remains a need for an fHbp polypeptide that can elicit effective bactericidal antibody responses.

SUMMARY

[0005] Variant factor H binding proteins that can elicit antibodies that are bactericidal for at least one strain of *Neisseria meningitidis*, compositions comprising such proteins, and methods of use of such proteins, are provided.

FEATURES

[0006] The present disclosure provides variants of factor H binding protein (fHbp) ID 1. The present disclosure provides a variant of fHbp wherein the variant comprises an amino acid substitution selected from at least one of: a) an amino acid substitution of the glutamine at amino acid 38 (Q38); b) an amino acid substitution of the glutamic acid at amino acid 92 (E92); c) a substitution of glycine for arginine at amino acid 130 (R130G); d) an amino acid substitution of the serine at amino acid 223 (S223); and e) a substitution of histidine for leucine at amino acid 248 (H248L), wherein the amino acid substitutions are relative to fHbp ID 1 (SEQ ID NO:1), wherein the variant comprises an amino acid sequence having at least 80% amino acid sequence identity to SEQ ID NO:1, wherein the variant fHbp binds human factor H (fH) with an affinity that is 50% or less of the affinity of fHbp ID 1 for human fH, and wherein the variant induces a bactericidal antibody response to at least one strain of *Neisseria meningitidis* in a mammalian host. In some cases, the amino acid substitution at Q38 is Q38R, Q38K, Q38H, Q38F, Q38Y, or Q38W. In some cases, the amino acid substitution at E92 is E92K, E92R, E92H, E92F, E92Y, or E92W. In some cases, the amino acid substitution at S223 is S223R, S223K, S223H, S223F, S223Y, or S223W. In some cases, the variant fHbp may further include a R41S or a R41A substitution relative to fHbp ID 1. For example the variant fHbp may include a R41S or a R41A substitution and a substitution at S223, e.g., R41S/S223R, relative to fHbp ID 1. In other cases, the variant fHbp may further include a R41S or a R41A substitution and a H248L substitution relative to fHbp ID 1. In certain cases, the variant fHbp may include two, three, or more of the substitutions disclosed herein. In a specific example, the variant fHbp may include the following substitutions: S223R and H248L relative to fHbp ID 1. In some cases, the variant fHbp binds human fH with an affinity that is 25% or less of the affinity of the fHbp ID 1 for human fH. In some cases, the variant fHbp binds human fH with an affinity that is 10% or less of the affinity of the fHbp ID 1 for human fH. In some cases, the variant fHbp binds human fH with an affinity that is 5% or less of the affinity of the fHbp ID 1 for human fH.

[0007] The present disclosure provides variants of fHbp ID 22. The present disclosure provides a variant of fHbp, wherein the variant comprises at least one amino acid substitution selected from: a) a substitution of isoleucine for asparagine at amino acid 115 (N115I); b) a substitution of glycine for aspartic acid at amino acid 121 (D121G); c) a substitution of threonine for serine at amino acid 128 (S128T); d) an amino acid substitution of the valine at position 131 (V131); e) an amino acid substitution of the lysine at position 219 (K219); f) an amino acid substitution of the glycine at position 220 (G220), wherein the amino acid substitutions are relative to fHbp ID 22 (SEQ ID NO:2), wherein the variant comprises an amino acid sequence having greater than 85% amino acid sequence identity to SEQ ID NO:2, wherein the variant fHbp binds human factor H (fH) with an affinity that is 50% or less of the affinity of fHbp ID 22 for human fH, and wherein the variant induces a bactericidal antibody response in a mammalian host. In some cases, the variant fHbp binds human fH with an affinity that is 25% or less of the affinity of the fHbp ID 22 for human fH. In some cases, the variant fHbp binds human fH with an affinity that is 10% or less of the affinity of the fHbp ID 22 for human fH. In some cases, the variant fHbp binds human fH with an affinity that is 5% or less of the affinity of the fHbp ID 22 for human fH. In some cases, the amino acid substitution at V131 is V131D, V131E, V131K, V131R, V131H, V131F, V131Y, or V131W. In some cases, the amino acid substitution at K219 is K219N, K219Q, K219D, K219E, K219F, K219Y, or K219W. In some cases, the amino acid substitution at G220 is G220S, G220N, G220Q, G220D, G220E, G220K, G220R, G220H, G220F, G220Y, or G220W.

[0008] In some cases, the variant fHbp includes a double mutation that increases thermal stability of the variant fHbp compared to thermal stability of wild type (WT) fHbp, e.g., WT fHbp ID 22. In some cases, the variant fHbp may include the substitutions L130R and G133D relative to fHbp ID 22 (SEQ ID NO:2), wherein the variant fHbp comprises an amino acid sequence having greater than 85% amino acid sequence identity to SEQ ID NO:2, wherein the variant fHbp binds human factor H (fH) with an affinity that is 50% or less of the affinity of fHbp ID 22 for human fH, wherein the variant induces a bactericidal antibody response in a mammalian host, and wherein the variant has a higher thermal stability compared to thermal stability of fHbp ID 22. In some cases, the variant fHbp may include a combination of substitutions, such as, L130R, G133D, and at least one amino acid substitution selected from: a) N115I; b) D121G; c) S128T; d) V131; e) K219 (e.g., K219N); and f) G220 (e.g., G220S), wherein the amino acid substitutions are relative to fHbp ID 22 (SEQ ID NO:2), wherein the variant fHbp comprises an amino acid sequence having greater than 85% amino acid sequence identity to SEQ ID NO:2, wherein the variant fHbp binds human factor H (fH) with an affinity

that is 50% or less of the affinity of fHbp ID 22 for human fH, and wherein the variant induces a bactericidal antibody response in a mammalian host. The thermal stability of the variant fHbp may be higher than a WT fHbp (e.g., fHbp ID 22) by at least 5°C, 10°C, 15°C, 20°C, or more, e.g., higher by 5°C-30°C, 5°C-25°C, 5°C-20°C, 10°C-20°C, or 15°C-20°C. As used herein, “thermal stability” refers to stability of a protein when exposed to higher temperature; a thermal stability variant protein maintains its conformation at a higher temperature than a wild type protein. For example, the variant fHbp, that include the double mutation that increases thermal stability compared to thermal stability of wild type (WT) fHbp, e.g., WT fHbp ID 22, may unfold at a higher temperature compared to WT fHbp. In certain cases, the N-terminal domain of the variant fHbp may unfold at a higher temperature than the N-terminal domain of the WT fHbp (e.g., fHbp ID 22).

[0009] Also disclosed herein are fHbp variants that include the mutations that enhance thermal stability as compared to a WT fHbp and further include additional mutations known to reduce binding of fH, such as, those disclosed in US2011/0256180. In certain embodiments, a variant of factor H binding protein (fHbp) is disclosed, wherein the variant comprises amino acid substitutions L130R and G133D and at least one of the substitutions: R80A, D211A, E218A, E248A, G236I, T221A, and H223A relative to fHbp ID 22 (SEQ ID NO:2), wherein the variant comprises an amino acid sequence having greater than 85% amino acid sequence identity to SEQ ID NO:2, wherein the variant fHbp binds human factor H (fH) with an affinity that is 50% or less of the affinity of fHbp ID 22 for human fH, and wherein the variant induces a bactericidal antibody response in a mammalian host.

[0010] The present disclosure provides variants of fHbp ID 55. The present disclosure provides a variant of fHbp, wherein the variant comprises at least one amino acid substitution selected from the group consisting of: a) an amino acid substitution of the glutamic acid at position 92 (E92); b) an amino acid substitution of the serine at position 223 (S223); and c) an amino acid substitution of the histidine at position 248 (H248), wherein the amino acid substitutions are relative to fHbp ID 55 (SEQ ID NO:3), wherein the variant comprises an amino acid sequence having at least 90% amino acid sequence identity to SEQ ID NO:3, wherein the variant fHbp binds human factor H (fH) with an affinity that is less than 50% of the affinity of fHbp ID 55 for human fH, and wherein the variant induces a bactericidal antibody response in a mammalian host. In some cases, the variant fHbp binds human fH with an affinity that is 25% or less of the affinity of the fHbp ID 55 for human fH. In some cases, the variant fHbp binds human fH with an affinity that is 10% or less of the affinity of the fHbp ID 55 for human fH. In some cases, the variant fHbp binds human fH with an affinity that is 5% or less of the affinity of the fHbp ID 55 for human fH. In some cases, the amino acid substitution at E92 is E92K, E92R, E92H, E92F,

E92Y, or E92W. In some cases, the amino acid substitution at S223 is S223R, S223K, S223H, S223F, S223Y, or S223W. In some cases, the amino acid substitution at H248 is H248L, H248I, H248V, H248D, H248E, H248F, H248Y, or H248W.

[0011] The present disclosure provides immunogenic compositions comprising a variant fHbp of the present disclosure. The present disclosure provides an immunogenic composition comprising: a) the variant fHbp according to any one of paragraphs 0006-0010 above; and b) a pharmaceutically acceptable excipient. In some cases, the fHbp variant is in a vesicle preparation prepared from a *Neisseria meningitidis* strain. In some cases, the pharmaceutically acceptable excipient comprises an adjuvant; e.g., where the adjuvant is aluminum phosphate or aluminum hydroxide. In some cases, the pharmaceutical composition further includes Neisserial surface protein A.

[0012] The present disclosure provides a nucleic acid encoding a variant fHbp according to any one of paragraphs 0006-0010 above. The present disclosure provides a recombinant expression vector comprising a nucleic acid encoding a variant fHbp according to any one of paragraphs 0006-0010 above. The present disclosure provides an *in vitro* host cell comprising a nucleic acid encoding a variant fHbp according to any one of paragraphs 0006-0010 above. The present disclosure provides an *in vitro* host cell comprising a recombinant expression vector comprising a nucleic acid encoding a variant fHbp according to any one of paragraphs 0006-0010 above.

[0013] The present disclosure provides a method of eliciting an antibody response in a mammal, the method comprising administering to a mammal an immunogenic composition of paragraph 0011, above. In some cases, the mammal is a human. In some cases, the antibody response is a bactericidal antibody response to one or more strains of *N. meningitidis*.

[0013A] Various embodiments of the claimed invention relate to a variant factor H binding protein (fHbp), wherein the variant fHbp comprises the amino acid leucine (L) or isoleucine (I) at position 248, wherein the amino acid position is relative to the numbering of SEQ ID NO:1 (fHbp ID1), wherein the variant fHbp comprises an amino acid sequence having at least 85% amino acid sequence identity to SEQ ID NO:1, wherein the variant fHbp binds human factor H (fH) with an affinity that is 50% or less of the affinity of fHbp ID 1 for human fH, and wherein the variant induces a

bactericidal antibody response to at least one strain of *Neisseria meningitidis* in a mammalian host.

[0013B] Various embodiments of the claimed invention also relate to a variant factor H binding protein (fHbp), wherein the variant fHbp comprises the amino acid substitution H248L, wherein the amino acid substitutions are relative to fHbp ID 55 (SEQ ID NO:3), wherein numbering of H248 is based on the amino acid sequence of SEQ ID NO:1, wherein the variant comprises an amino acid sequence having at least 90% amino acid sequence identity to SEQ ID NO:3, wherein the variant fHbp binds human factor H (fH) with an affinity that is less than 50% of the affinity of fHbp ID 55 for human fH, and wherein the variant induces a bactericidal antibody response in a mammalian host.

BRIEF DESCRIPTION OF THE DRAWINGS

[0014] **FIG. 1** depicts purified recombinant fHbp ID 1 mutants stained with Coomassie blue on a polyacrylamide gel. Lane 1, Kaleidoscope molecular weight marker (Bio-Rad Laboratories); 2, fHbp ID 1 wild-type; 3, Q38R; 4, E92K; 5, R130G; 6, S223R; 7, H248L.

[0015] **FIGS. 2A and 2B** depict binding of fHbp ID 1 mutants to human fH, measured by ELISA. The mean and range for replicate measurements are shown.

[0016] **FIGS. 3A-3E** depict binding of fHbp ID 1 mutants to human fH, measured by surface plasmon resonance. For reference, the same data for the ID 1 wild-type (WT) protein are shown in each of FIGS. 3A-3E.

[0017] **FIG. 4A-4E** depict binding of murine anti-fHbp monoclonal antibodies (mAb) to fHbp ID 1 mutant proteins, measured by ELISA. The mean and range for duplicate measurements are shown.

[0018] **FIGS. 5A and 5B** depict bactericidal activity of serum from mice immunized with fHbp ID 1 mutants. Each symbol represents the titer of an individual mouse, and the horizontal bars represent the geometric mean titers. **FIG. 5A** depicts bactericidal activity of serum from wild-type mice immunized with fHbp ID 1 mutants. **FIG. 5B** depicts bactericidal activity of serum from human fH transgenic mice immunized with fHbp ID 1 mutants.

[0019] **FIGS. 6A and 6B** depict characterization of fHbp ID 1 single or double mutants. **FIG. 6A** depicts binding of human fH to fHbp ID 1 double mutants. **FIG. 6B** depicts binding of murine anti-fHbp monoclonal antibody (mAb) JAR 4 to mutants.

[0020] **FIG. 7** depicts bactericidal activity of serum from wild-type mice immunized with fHbp ID 1 single or double mutants. Each symbol represents the titer of an individual mouse, and the horizontal bars represent the geometric mean titers.

[0021] **FIGS. 8A and 8B** depict characterization of fHbp ID 55 mutants. **FIG. 8A** shows binding of human fH to immobilized fHbp ID 55 mutants, measured by ELISA and **FIG. 8B** shows binding of murine anti-fHbp monoclonal antibody (mAb) JAR 41 to fHbp mutant ID 55 proteins, measured by ELISA. The mean and range for two to four replicates are shown.

[0022] **FIG. 9** depicts bactericidal activity of serum from wild-type mice immunized with fHbp ID 55 mutants. Each symbol represents the titer of an individual mouse, and the horizontal bars represent the geometric mean titers.

[0023] **FIGS. 10A and 10B** depict bactericidal activity of serum from mice immunized with fHbp ID 55. **FIG. 10A** depicts the bactericidal activity of serum from human fH transgenic mice immunized with the licensed Trumenba vaccine or an investigational fHbp ID 55 mutant S223R. **FIG. 10B** depicts the relationship between serum human fH concentrations in individual transgenic mice and the serum bactericidal antibody titers (circular symbols). For comparison, the titers of wild-type (WT) mice are shown (squares).

[0024] **FIGS. 11A-11D** depict characterization of fHbp ID 22 mutants. **FIGS. 11A-11C** depict binding of fHbp ID 22 mutant to human fH, measured by ELISA. The mean and range of two to four replicates are shown. **FIG. 11D** depicts binding of murine anti-fHbp monoclonal antibody (mAb) JAR 4 to fHbp mutant ID 22 proteins, measured by ELISA. fHbp ID 22 wild-type (WT) and D211A mutant are shown as controls. The mean and range of duplicate measurements are shown.

[0025] FIGS. 12A and 12B show bactericidal activity of serum from mice immunized with fHbp ID 22 mutants. Each symbol represents the titer of an individual mouse, and the horizontal bars represent the geometric mean titers. FIG. 12A and FIG. 12B show bactericidal activity of serum from wild-type mice in two experiments to test different fHbp ID 22 mutants.

[0026] FIG. 13 depicts bactericidal activity of serum from human fH transgenic mice immunized with fHbp ID 22 mutants.

[0027] FIG. 14 depicts thermal unfolding of fHbp ID 22 wild-type (WT) and L130R/G133D double mutant measured by differential scanning microcalorimetry.

[0028] FIG. 15 depicts characterization of fHbp ID 22 triple mutants. FIG. 15A depicts binding of human fH to fHbp ID 22 triple mutants. DM refers to L130R/G133D double mutant. FIG. 15B depicts binding of murine anti-fHbp monoclonal antibody (mAb) JAR 4 to fHbp ID 22 triple mutants.

[0029] FIG. 16 depicts bactericidal activity of serum from human fH transgenic mice immunized with fHbp ID 22 triple mutants. DM refers to L130R/G133D double mutant.

[0030] FIG. 17 provides a table of exemplary fHbp mutants with decreased binding of human fH.

[0031] FIG. 18 provides the amino acid sequence of wild-type Human factor H.

[0032] FIG. 19 provides amino acid sequences of fHbp ID 1, ID 22, and ID 55 from *N. meningitidis* strains.

[0033] FIGS. 20-24 provide amino acid sequences of fHbp ID 1 variants.

[0034] FIGS. 25-30 provide amino acid sequences of fHbp ID 22 variants.

[0035] FIGS. 31-33 provide amino acid sequences of fHbp ID 55 variants.

[0036] FIGS. 34-36 provide amino acid sequences of fHbp ID 1 double mutant variants.

[0037] FIGS. 37-39 provide amino acid sequences of fHbp ID 22 double and triple mutant variants.

[0038] FIG. 40 provides an amino acid sequence of NspA.

DEFINITIONS

[0039] “Factor H Binding Protein” (fHbp), which is also known in the literature as GNA1870, GNA 1870, ORF2086, LP2086 (lipoprotein 2086), and “741” refers to a class of *N. meningitidis* polypeptides. It is found in nature as a lipoprotein on the surface of the bacterium *N. meningitidis*. fHbps have been sub-divided into three fHbp variant groups (referred to as variant group 1 (v.1), variant group 2 (v.2), and variant group 3 (v.3) in some reports (Massignani et al. (2003) *J Exp Med* 197:789-99) and sub-family A and B in other reports (see, e.g., Fletcher et al.

(2004) *Infect Immun* 72:2088-2100)) based on amino acid sequence variability and immunologic cross-reactivity (Massignani et al. (2003) *J Exp Med* 197:789-99). fHbp also can be classified into one of the six most common fHbp modular groups, designated Modular Group I to Modular Group VI, as shown in Figure 2 of Vu et al. (2012) *Sci. Reports* 2:341. Each unique fHbp found in *N. meningitidis* is also assigned an fHbp peptide ID according to the pubmlst.org/neisseria/fHbp/ website. Because the length of variant 2 (v.2) fHbp protein (from strain 8047, fHbp ID 77) and variant 3 (v.3) fHbp (from strain M1239, fHbp ID 28) differ by -1 and +7 amino acid residues, respectively, from that from strain MC58 (fHbp ID 1), the numbering used herein to refer to residues for v.2 and v.3 fHbp proteins differs from numbering based on the actual amino acid sequences of these proteins. Thus, for example, reference to a leucine residue (L) at position 166 of the v.2 or v.3 fHbp sequence refers to the residue at position 165 of the v.2 protein and at position 173 in the v.3 protein. Unless noted otherwise, the numbering of the amino acid substitutions present in the fHbp variants is with reference to the numbering of the amino acid residues in fHbp ID 1.

[0040] Human factor H (“human fH”) as used herein, refers to a protein comprising an amino acid sequence as shown in Figure 18 (SEQ ID NO:4), and naturally-occurring human allelic variants thereof.

[0041] “Derived from” in the context of an amino acid sequence or polynucleotide sequence (e.g., an amino acid sequence “derived from” fHbp ID 1) is meant to indicate that the polypeptide or nucleic acid has a sequence that is based on that of a reference polypeptide or nucleic acid (e.g., a naturally occurring fHbp protein or encoding nucleic acid), and is not meant to be limiting as to the source or method in which the protein or nucleic acid is made. Non-limiting examples of reference polypeptides and reference polynucleotides from which an amino acid sequence or polynucleotide sequence may be “derived from” include a naturally-occurring fHbp, fHbp ID 1, and a non-naturally-occurring fHbp. “Derived from” in the context of bacterial strains is meant to indicate that a strain was obtained through passage *in vivo*, or in *in vitro* culture, of a parental strain and/or is a recombinant cell obtained by modification of a parental strain.

[0042] “Conservative amino acid substitution” refers to a substitution of one amino acid residue for another sharing chemical and physical properties of the amino acid side chain (e.g., charge, size, hydrophobicity/hydrophilicity). “Conservative substitutions” are intended to include substitution within the following groups of amino acid residues: gly, ala; val, ile, leu; asp, glu; asn, gln; ser, thr; lys, arg; and phe, tyr. Guidance for such substitutions can be drawn from alignments of amino acid sequences of polypeptides presenting the epitope of interest.

[0043] The term “protective immunity” means that a vaccine or immunization schedule that is administered to a mammal induces an immune response that prevents, retards the development of, or reduces the severity of a disease that is caused by *Neisseria meningitidis*, or diminishes or altogether eliminates the symptoms of the disease. Protective immunity can be accompanied by production of bactericidal antibodies. It should be noted that production of bactericidal antibodies against *Neisseria meningitidis* is accepted in the field as predictive of a vaccine's protective effect in humans. (Goldschneider et al. (1969) *J. Exp. Med.* 129:1307; Borrow et al. (2001) *Infect Immun.* 69:1568).

[0044] The phrase “a disease caused by a strain of *Neisseria meningitidis*” encompasses any clinical symptom or combination of clinical symptoms that are present in an infection of a human with a *Neisseria meningitidis*. These symptoms include but are not limited to: colonization of the upper respiratory tract (e.g. mucosa of the nasopharynx and tonsils) by a pathogenic strain of *Neisseria meningitidis*, penetration of the bacteria into the mucosa and the submucosal vascular bed, septicemia, septic shock, inflammation, hemorrhagic skin lesions, activation of fibrinolysis and of blood coagulation, organ dysfunction such as kidney, lung, and cardiac failure, adrenal hemorrhaging and muscular infarction, capillary leakage, edema, peripheral limb ischemia, respiratory distress syndrome, pericarditis and meningitis.

[0045] The phrase “specifically binds to an antibody” or “specifically immunoreactive with”, in the context of an antigen (e.g., a polypeptide antigen) refers to a binding reaction that is based on and/or is probative of the presence of the antigen in a sample which may also include a heterogeneous population of other molecules. Thus, under designated conditions, the specified antibody or antibodies bind(s) to a particular antigen or antigens in a sample and do not bind in a significant amount to other molecules present in the sample. “Specifically binds to an antibody” or “specifically immunoreactive with” in the context of an epitope of an antigen (e.g., an epitope of a polypeptide) refers to a binding reaction which is based on and/or is probative of the presence of the epitope in an antigen (e.g., polypeptide) which may also include a heterogeneous population of other epitopes, as well as a heterogeneous population of antigens. Thus, under designated conditions, the specified antibody or antibodies bind(s) to a particular epitope of an antigen and do not bind in a significant amount to other epitopes present in the antigen and/or in the sample.

[0046] The phrase “in a sufficient amount to elicit an immune response” means that there is a detectable difference between an immune response indicator measured before and after administration of a particular antigen preparation. Immune response indicators include but are not limited to: antibody titer or specificity, as detected by an assay such as enzyme-linked immunosorbent assay (ELISA), bactericidal assay, flow cytometry, immunoprecipitation,

Ouchterlony immunodiffusion; binding detection assays of, for example, spot, Western blot or antigen arrays; cytotoxicity assays, etc.

[0047] A “surface antigen” is an antigen that is present in a surface structure of *Neisseria meningitidis* (e.g. the outer membrane, capsule, pili, etc.).

[0048] “Isolated” refers to an entity of interest that is in an environment different from that in which the compound may naturally occur. “Isolated” is meant to include compounds that are within samples that are substantially enriched for the compound of interest and/or in which the compound of interest is partially or substantially purified. In some cases, an isolated component (e.g., a polypeptide, such as an fHbp variant of the present disclosure; a nucleic acid of the present disclosure; a recombinant vector of the present disclosure) is purified, e.g., the isolated component is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, or greater than 99%, pure.

[0049] “Enriched” means that a sample is non-naturally manipulated (e.g., by an experimentalist or a clinician) so that a compound of interest is present in a greater concentration (e.g., at least a three-fold greater, at least 4-fold greater, at least 8-fold greater, at least 64-fold greater, or more) than the concentration of the compound in the starting sample, such as a biological sample (e.g., a sample in which the compound naturally occurs or in which it is present after administration), or in which the compound was made (e.g., as in a bacterial polypeptide, antibody, nucleic acid, and the like).

[0050] Before the present invention is further described, it is to be understood that this invention is not limited to particular embodiments described, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

[0051] Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limit of that range and any other stated or intervening value in that stated range, is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges, and are also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either or both of those included limits are also included in the invention.

[0052] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the present invention, the preferred methods and materials are now described.

[0053] It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a factor H binding protein" includes a plurality of such factor H binding proteins and reference to "the immunogenic composition" includes reference to one or more immunogenic compositions and equivalents thereof known to those skilled in the art, and so forth. It is further noted that the claims may be drafted to exclude any optional element. As such, this statement is intended to serve as antecedent basis for use of such exclusive terminology as "solely," "only" and the like in connection with the recitation of claim elements, or use of a "negative" limitation.

[0054] It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, may also be provided in combination in a single embodiment. Conversely, various features of the invention, which are, for brevity, described in the context of a single embodiment, may also be provided separately or in any suitable sub-combination. All combinations of the embodiments pertaining to the invention are specifically embraced by the present invention and are disclosed herein just as if each and every combination was individually and explicitly disclosed. In addition, all sub-combinations of the various embodiments and elements thereof are also specifically embraced by the present invention and are disclosed herein just as if each and every such sub-combination was individually and explicitly disclosed herein.

[0055] The publications discussed herein are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention. Further, the dates of publication provided may be different from the actual publication dates which may need to be independently confirmed.

DETAILED DESCRIPTION

[0056] The present disclosure provides variant factor H binding proteins (fHbp) that can elicit antibodies that are bactericidal for at least one strain of *Neisseria meningitidis*. The present disclosure provides compositions, including immunogenic compositions, comprising a variant fHbp of the present disclosure. The present disclosure provides methods of use of variant fHbp of the present disclosure, or a composition comprising a variant fHbp of the present disclosure.

VARIANT FHBP

[0057] The present disclosure provides variant fHbp that differ in amino acid sequence from a wild-type *N. meningitidis* fHbp by from 1 to 10 amino acids (e.g., by from 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acids) from 10 amino acids to 15 amino acids, from 15 amino acids to 20 amino acids, from 20 amino acids to 30 amino acids, from 30 amino acids to 40 amino acids, or from 40 amino acids to 50 amino acids, such that the variant fHbp exhibits reduced affinity to human factor H (fH), compared to a reference fHbp, and where the variant fHbp elicits a bactericidal immune response to one or more *N. meningitidis* strains when administered to a mammalian host. In some cases, the variant fHbp differs in amino acid sequence from a reference wild-type *N. meningitidis* fHbp by no more than from 1 to 10 acid substitutions. In some cases, the variant fHbp differs in amino acid sequence from a reference wild-type *N. meningitidis* fHbp by only one amino acid substitution.

[0058] In some cases, variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to a reference fHbp sequence; where the variant fHbp comprises one or more amino acid substitutions relative to the reference fHbp sequence such that the variant fHbp exhibits an affinity for human fH that is 85% or less of the binding affinity of the reference fHbp for human fH, e.g., the variant fHbp exhibits an affinity for human fH that is from about 85% to about 75%, from about 75% to about 65%, from about 65% to about 55%, from about 55% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%, of the binding affinity of the affinity of the reference fHbp for human fH; and the variant fHbp induces a bactericidal immune response to at least one strain of *N. meningitidis* when administered to a mammalian host (e.g., a human; or a non-human animal model).

[0059] A variant fHbp of the present disclosure maintains substantially the same conformation of a reference (e.g., wild-type) fHbp that binds human fH when the reference fHbp is in its native conformation. Whether a variant fHbp of the present disclosure maintains substantially the same

conformation of a reference (e.g., wild-type) fHbp that binds human fH can be determined using antibodies that bind wild-type fHbp when the wild-type fHbp is in its native conformation. Such antibodies include, e.g., JAR 41; JAR 4; and JAR 31. See, e.g., Vu et al. (2012) *Sci. Reports* 2:341. A hybridoma producing JAR 4 monoclonal antibody has the American Type Culture Collection (ATCC) number PTA-8943; see also USPN 8,470,340. For example, in some cases, a variant fHbp of the present disclosure retains binding to JAR 4; e.g., a variant fHbp of the present disclosure retains at least 75%, at least 80%, at least 85%, at least 90%, or at least 95%, binding to JAR 4 of a reference fHbp (e.g., fHbp ID 1, fHbp ID 22, or fHbp ID 55) in its native conformation.

Variants of fHbp ID 1

[0060] A “reference fHbp” from which a variant fHbp of the present disclosure is derived is in some cases fHbp ID 1. The amino acid sequence of fHbp ID 1 is set out below.

[0061] fHbp ID 1:

CSSGGGGVAADIGAGLADALTAPLDHKDKGLQSLTLDQSVRKNEKLKLAQGAEKTY
GNGDSLNTGKLKNDKVSRFDFIRQIEVDGQLITLESGEFQVYKQSHSALTAFQTEQIQDS
EHSGKMKVAKRQFRIGDIAGEHTSFDKLPEGGRATYRGTAFGSDDAGGKLTYTIDFAAKQ
GNGKIEHLKSPELNVDLAAADIKPDGKRHAVISGSVLYNQAEKGSYSLGIFGGKAQEVA
GSAEVKTVNGIRHIGLAAKQ (SEQ ID NO:1).

[0062] In some cases, a variant fHbp of the present disclosure is a variant group 1 fHbp. In some cases, a variant fHbp of the present disclosure is a variant group 1 fHbp, and is a modular group I fHbp. In some cases, variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:1; where the variant fHbp comprises one or more amino acid substitutions relative to fHbp ID 1 such that the variant fHbp exhibits an affinity for human fH that is 85% or less of the binding affinity of fHbp ID 1 for human fH, e.g., the variant fHbp exhibits an affinity for human fH that is from about 85% to about 75%, from about 75% to about 65%, from about 65% to about 55%, from about 55% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%, of the binding affinity of the affinity of fHbp ID 1 for human fH; and the variant fHbp induces a bactericidal immune response to at least one strain of *N. meningitidis* when administered to a mammalian host (e.g., a human; or a non-human animal model).

[0063] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least

99%, amino acid sequence identity to SEQ ID NO:1, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 1 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises an amino acid substitution selected from at least one of: a) an amino acid substitution of the glutamine at amino acid 38 (Q38); b) an amino acid substitution of the glutamic acid at amino acid 92 (E92); c) a substitution of glycine for arginine at amino acid 130 (R130G); d) an amino acid substitution of the serine at amino acid 223 (S223); and e) a substitution of histidine for leucine at amino acid 248 (H248L), based on the numbering of fHbp ID 1.

[0064] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:1, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 1 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises an amino acid substitution of the glutamine at amino acid 38 (Q38). In some cases, the variant fHbp comprises a Q38R substitution. Other amino acids with positively charged or aromatic side chains, such as lysine, histidine, phenylalanine, tyrosine or tryptophan, also may be substituted at this position. Thus, in some cases, the variant fHbp comprises a Q38K substitution, a Q38H substitution, a Q38F substitution, a Q38Y substitution, or a Q38W substitution. As one example, a variant fHbp of the present disclosure can comprise the amino acid sequence depicted in Figure 20 and set forth in SEQ ID NO:5.

[0065] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:1, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 1 for human fH,

where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises an amino acid substitution of the glutamic acid at amino acid 92 (E92). In some cases the fHbp variant comprises an E92K substitution. Other amino acids with positively charged or aromatic side chains, such as arginine, histidine, phenylalanine, tyrosine or tryptophan, also may be substituted at this position. Thus, for example, in some cases the fHbp variant comprises an E92R substitution, an E92H substitution, an E92F substitution, an E92Y substitution, or an E92W substitution. As one example, a variant fHbp of the present disclosure can comprise the amino acid sequence depicted in Figure 21 and set forth in SEQ ID NO:6.

[0066] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:1, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 1 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises a substitution of glycine for arginine at amino acid 130 (R130G). For example, a variant fHbp of the present disclosure can comprise the amino acid sequence depicted in Figure 22 and set forth in SEQ ID NO:7. Other amino acids with negatively charged or aromatic side chains, such as aspartate, glutamate, phenylalanine, tyrosine, or tryptophan, may also be substituted at R130. Thus, for example, in some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:1, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 1 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises an R130D substitution, an R130E substitution, an R130F substitution, an R130Y substitution, or an R130W substitution.

[0067] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:1, where the variant fHbp binds human fH

with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 1 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises an amino acid substitution of the serine at amino acid 223 (S223). In some cases, the fHbp variant comprises an S223R substitution. Other amino acids with positively charged or aromatic side chains, such as lysine, histidine, phenylalanine, tyrosine or tryptophan, also may be substituted at this position. Thus, for example, in some cases, the fHbp variant comprises an S223K substitution, an S223H substitution, an S223F substitution, an S223Y substitution, or an S223W substitution. As one example, a variant fHbp of the present disclosure can comprise the amino acid sequence depicted in Figure 23 and set forth in SEQ ID NO:8.

[0068] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:1, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 1 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises a substitution of histidine for leucine at amino acid 248 (H248L). For example, a variant fHbp of the present disclosure can comprise the amino acid sequence depicted in Figure 24 and set forth in SEQ ID NO:9. Other amino acids with non-polar, negatively charged or aromatic side chains, such as isoleucine, valine, aspartate, glutamate, phenylalanine, tyrosine or tryptophan, also may be substituted at H248. Thus, in some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:1, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 1 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises an H248I substitution,

an H248V substitution, an H248D substitution, an H248E substitution, an H248F substitution, an H248Y substitution, or an H248W substitution.

Combinations of amino acid substitutions

[0069] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:1, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 1 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises an amino acid substitution selected from two or more of: a) an amino acid substitution of the glutamine at amino acid 38 (Q38); b) an amino acid substitution of the glutamic acid at amino acid 92 (E92); c) a substitution of glycine for arginine at amino acid 130 (R130G); d) an amino acid substitution of the serine at amino acid 223 (S223); and e) a substitution of histidine for leucine at amino acid 248 (H248L), based on the numbering of fHbp ID 1.

[0070] Combinations of substitutions may be included wherein the two substitutions are in different structural domains, and each independently decreases binding of fH to fHbp (e.g., one substitution in the N-terminal domain, in combination with an amino acid substitution in the C-terminal domain. In some cases, a variant fHbp of the present disclosure comprises a first amino acid substitution within the N-terminal domain; and a second amino acid substitution within the C-terminal domain. In some cases, a variant fHbp of the present disclosure comprises a first amino acid substitution within the N-terminal domain; and a second amino acid substitution within the N-terminal domain. In some cases, a variant fHbp of the present disclosure comprises a first amino acid substitution within the C-terminal domain; and a second amino acid substitution within the C-terminal domain.

[0071] For example, in some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:1, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 1 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N.*

meningitidis in a mammalian host, and where the variant fHbp comprises: a) an amino acid substitution of the glutamine at amino acid 38 (Q38) and b) an amino acid substitution of the glutamic acid at amino acid 92 (E92); or where the variant fHbp comprises: a) an amino acid substitution of the glutamine at amino acid 38 (Q38) and c) a substitution of glycine for arginine at amino acid 130 (R130G); or where the variant fHbp comprises a) an amino acid substitution of the glutamine at amino acid 38 (Q38) and d) an amino acid substitution of the serine at amino acid 223 (S223); or where the variant fHbp comprises a) an amino acid substitution of the glutamine at amino acid 38 (Q38) and c) a substitution of histidine for leucine at amino acid 248 (H248L), based on the numbering of fHbp ID 1.

[0072] As additional non-limiting examples, in some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:1, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 1 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises: b) an amino acid substitution of the glutamic acid at amino acid 92 (E92) and c) a substitution of glycine for arginine at amino acid 130 (R130G); or where the variant fHbp comprises b) an amino acid substitution of the glutamic acid at amino acid 92 (E92) and d) an amino acid substitution of the serine at amino acid 223 (S223); or where the variant fHbp comprises b) an amino acid substitution of the glutamic acid at amino acid 92 (E92) and e) a substitution of histidine for leucine at amino acid 248 (H248L); or where the variant fHbp comprises c) a substitution of glycine for arginine at amino acid 130 (R130G) and d) an amino acid substitution of the serine at amino acid 223 (S223); or where the variant fHbp comprises c) a substitution of glycine for arginine at amino acid 130 (R130G) and e) a substitution of histidine for leucine at amino acid 248 (H248L); or where the variant fHbp comprises d) an amino acid substitution of the serine at amino acid 223 (S223) and e) a substitution of histidine for leucine at amino acid 248 (H248L), based on the numbering of fHbp ID 1.

[0073] As additional non-limiting examples, in some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:1, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about

15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 1 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises: i) a Q38R substitution; and ii) an R130G substitution, based on the numbering of fHbp ID 1.

[0074] Also provided herein are variant fHbp proteins that include one or more substitutions relative to amino acid sequence of fHbp ID 1 as set forth above and further include the substitution R41S. Exemplary variant fHbp include a R41S substitution and a substitution at S223, e.g., R41S/S223R relative to fHbp ID 1 or a R41S substitution and a H248L substitution relative to fHbp ID 1. In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:1, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises two or more of the following amino acid substitutions: a) a substitution of serine for arginine at amino acid 41 (R41S); b) a substitution of arginine for serine at amino acid 223 (S223R); c) a substitution of leucine for histidine at amino acid 248 (H248L), based on the numbering of fHbp ID 1.

[0075] Also disclosed herein are variant fHbp proteins that include one or more substitutions relative to amino acid sequence of fHbp ID 1 as set forth above and further include the substitutions disclosed in US2011/0256180.

Variants of fHbp ID 22

[0076] A “reference fHbp” from which a variant fHbp of the present disclosure is derived is in some cases fHbp ID 22. The amino acid sequence of fHbp ID 22 is set out below.

[0077] fHbp ID 22:

CSSGGGGVAADIGAGLADALTAPLDHKDKSLQLSLTLDQSVRKNEKLKLAQGAEKTYG
NGDSLNTGKLNDKVSRFDFIRQIEVDGQLITLESGEFQIYKQDHSAVVALQIEKINNPDK
IDSLINQRSFLVSGLGGEHTAFNQLPSGKAELYHGKAFSSDDPNGLHYSIDFTKKQGYG

RIEHLKTPEQNVELASAEKADEKSHAVILGDTRYGEEKGTYHLALFGDRAQEIAAGSA
TVKIREKVHEIGIAGKQ (SEQ ID NO:2).

[0078] In some cases, a variant fHbp of the present disclosure is a variant group 2 fHbp. In some cases, a variant fHbp of the present disclosure is a variant group 2 fHbp, and is a modular group III fHbp. In some cases, variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:2; where the variant fHbp comprises one or more amino acid substitutions relative to fHbp ID 22 such that the variant fHbp exhibits an affinity for human fH that is 85% or less of the binding affinity of fHbp ID 22 for human fH, e.g., the variant fHbp exhibits an affinity for human fH that is from about 85% to about 75%, from about 75% to about 65%, from about 65% to about 55%, from about 55% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%, of the binding affinity of the affinity of fHbp ID 22 for human fH; and the variant fHbp induces a bactericidal immune response to at least one strain of *N. meningitidis* when administered to a mammalian host (e.g., a human; or a non-human animal model).

[0079] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:2, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises an amino acid substitution selected from at least one of: a) a substitution of isoleucine for asparagine at amino acid 115 (N115I); b) a substitution of glycine for aspartic acid at amino acid 121 (D121G); c) a substitution of threonine for serine at amino acid 128 (S128T); d) an amino acid substitution of the valine at position 131 (V131); e) an amino acid substitution of the lysine at position 219 (K219); f) an amino acid substitution of the glycine at position 220 (G220), relative to the amino acid sequence of fHbp ID 22. As noted herein, the numbering of the amino acid residue is based on that of fHbp ID 1.

[0080] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least

99%, amino acid sequence identity to SEQ ID NO:2, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises a substitution of isoleucine for asparagine at amino acid 115 (N115I). For example, a variant fHbp of the present disclosure can comprise the amino acid sequence depicted in Figure 25 and set forth in SEQ ID NO:10. Other amino acids with non-polar, positively charged or aromatic side chains, such as valine, leucine, lysine, arginine, histidine, phenylalanine, tyrosine or tryptophan, also may be substituted at N115. Thus, for example, in some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:2, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises an N115V substitution, an N115L substitution, an N115K substitution, an N115R substitution, an N115H substitution, an N115F substitution, an N115Y substitution, or an N115W substitution relative to amino acid sequence of fHbp ID 22.

[0081] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:2, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises a substitution of glycine for aspartic acid at amino acid 121 (D121G). For example, a variant fHbp of the present disclosure can comprise the amino acid sequence depicted in Figure 26 and set forth in SEQ ID NO:11. Other amino acids with non-polar, positively charged or aromatic side chains, such as

leucine, isoleucine, valine, lysine, arginine, histidine, phenylalanine, tyrosine or tryptophan, also may be substituted at D121. Thus, for example, in some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:2, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises a D121L substitution, a D121I substitution, a D121V substitution, a D121K substitution, a D121R substitution, a D121H substitution, a D121F substitution, a D121Y substitution, or a D121W substitution relative to amino acid sequence of fHbp ID 22.

[0082] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:2, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises a substitution of threonine for serine at amino acid 128 (S128T). For example, a variant fHbp of the present disclosure can comprise the amino acid sequence depicted in Figure 27 and set forth in SEQ ID NO:12. Other amino acids with polar, charged or aromatic side chains, such as methionine, asparagine, glutamine, aspartate, glutamate, lysine, arginine, histidine, phenylalanine, tyrosine or tryptophan, also may be substituted at S128. Thus, for example, in some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:2, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant

fHbp comprises an S128M substitution, an S128N substitution, an S128D substitution, an S128E substitution, an S128K substitution, an S128R substitution, an S128H substitution, an S128F substitution, an S128Y substitution, or an S128W substitution relative to amino acid sequence of fHbp ID 22.

[0083] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:2, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises an amino acid substitution of the valine at position 131 (V131). In some cases, the fHbp variant comprises a V131D substitution. Other amino acids with charged or aromatic side chains, such as glutamate, lysine, arginine, histidine, phenylalanine, tyrosine or tryptophan, also may be substituted at this position. Thus, for example, in some cases, the fHbp variant comprises a V131E substitution, a V131K substitution, a V131R substitution, a V131H substitution, a V131F substitution, a V131Y substitution, or a V131W substitution. As one example, a variant fHbp of the present disclosure can comprise the amino acid sequence depicted in Figure 28 and set forth in SEQ ID NO:13 relative to amino acid sequence of fHbp ID 22.

[0084] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:2, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises an amino acid substitution of the lysine at position 219 (K219). In some cases, the fHbp variant comprises a K219N substitution. Other amino acids with polar, negatively charged or aromatic side chains, such as glutamine, aspartate, glutamate, phenylalanine, tyrosine or tryptophan, also may be substituted at this position. Thus, for example, in some cases, the fHbp variant comprises a K219Q substitution, a K219D substitution, a K219E substitution, a K219F substitution, a K219Y

substitution, or a K219W substitution. As one example, a variant fHbp of the present disclosure can comprise the amino acid sequence depicted in Figure 29 and set forth in SEQ ID NO:14.

[0085] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:2, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises an amino acid substitution of the glycine at position 220 (G220). In some cases, the fHbp variant comprises a G220S substitution. Other amino acids with polar, charged or aromatic side chains, such as asparagine, glutamine, aspartate, glutamate, lysine, arginine, histidine, phenylalanine, tyrosine or tryptophan, also may be substituted at this position. Thus, for example, in some cases, the fHbp variant comprises a G220N substitution, a G220Q substitution, a G220D substitution, a G220E substitution, a G220K substitution, a G220R substitution, a G220H substitution, a G220F substitution, a G220Y substitution, or a G220W substitution. For example, a variant fHbp of the present disclosure can comprise the amino acid sequence depicted in Figure 30 and set forth in SEQ ID NO:15.

Combinations of amino acid substitutions

[0086] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:2, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises an amino acid substitution selected from two or more of: a) a substitution of isoleucine for asparagine at amino acid 115 (N115I); b) a substitution of glycine for aspartic acid at amino acid 121 (D121G); c) a substitution of threonine for serine at amino acid 128 (S128T); d) an amino acid substitution of the valine at position 131 (V131); e) an amino acid substitution of the lysine at position 219 (K219); f) an amino acid substitution of the glycine at position 220 (G220), relative to amino

acid sequence of fHbp ID 22. As noted herein, the numbering of the residues is based on the numbering of the amino acids in fHbp ID 1.

[0087] Combinations of substitutions may be included wherein the two substitutions are in different structural domains, and each independently decreases binding of fH to fHbp (e.g., one substitution in the N-terminal domain, in combination with an amino acid substitution in the C-terminal domain. In some cases, a variant fHbp of the present disclosure comprises a first amino acid substitution within the N-terminal domain; and a second amino acid substitution within the C-terminal domain. In some cases, a variant fHbp of the present disclosure comprises a first amino acid substitution within the N-terminal domain; and a second amino acid substitution within the N-terminal domain. In some cases, a variant fHbp of the present disclosure comprises a first amino acid substitution within the C-terminal domain; and a second amino acid substitution within the C-terminal domain.

[0088] For example, in some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:2, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises: a) a substitution of isoleucine for asparagine at amino acid 115 (N115I) and b) a substitution of glycine for aspartic acid at amino acid 121 (D121G); or where the variant fHbp comprises: a) a substitution of isoleucine for asparagine at amino acid 115 (N115I) and c) a substitution of threonine for serine at amino acid 128 (S128T); or where the variant fHbp comprises: a) a substitution of isoleucine for asparagine at amino acid 115 (N115I) and d) an amino acid substitution of the valine at position 131 (V131); or where the variant fHbp comprises: a) a substitution of isoleucine for asparagine at amino acid 115 (N115I) and e) an amino acid substitution of the lysine at position 219 (K219); or where the variant fHbp comprises: a) a substitution of isoleucine for asparagine at amino acid 115 (N115I) and f) an amino acid substitution of the glycine at position 220 (G220), relative to amino acid sequence of fHbp ID 22, the numbering of the substituted residue based on the numbering of amino acid sequence of fHbp ID 1.

[0089] As additional non-limiting examples, in some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:2, where

the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises: b) a substitution of glycine for aspartic acid at amino acid 121 (D121G) and c) a substitution of threonine for serine at amino acid 128 (S128T); or where the variant fHbp comprises: b) a substitution of glycine for aspartic acid at amino acid 121 (D121G) and d) an amino acid substitution of the valine at position 131 (V131); or where the variant fHbp comprises: b) a substitution of glycine for aspartic acid at amino acid 121 (D121G) and e) an amino acid substitution of the lysine at position 219 (K219); or where the variant fHbp comprises: b) a substitution of glycine for aspartic acid at amino acid 121 (D121G) and f) an amino acid substitution of the glycine at position 220 (G220), relative to amino acid sequence of fHbp ID 22. The numbering of the substituted residue(s) is based on the numbering of amino acid sequence of fHbp ID 1.

[0090] As additional non-limiting examples, in some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:2, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises: c) a substitution of threonine for serine at amino acid 128 (S128T) and d) an amino acid substitution of the valine at position 131 (V131); or where the variant fHbp comprises: c) a substitution of threonine for serine at amino acid 128 (S128T) and e) an amino acid substitution of the lysine at position 219 (K219); or where the variant fHbp comprises: c) a substitution of threonine for serine at amino acid 128 (S128T) and f) an amino acid substitution of the glycine at position 220 (G220); or where the variant fHbp comprises: d) an amino acid substitution of the valine at position 131 (V131) and e) an amino acid substitution of the lysine at position 219 (K219); or where the variant fHbp comprises: d) an amino acid substitution of the valine at position 131 (V131) and f) an amino acid substitution of the glycine at position 220 (G220); or where the variant fHbp comprises: e) an amino acid substitution of the lysine at position 219

(K219) and f) an amino acid substitution of the glycine at position 220 (G220),), relative to amino acid sequence of fHbp ID 22. The numbering of the substituted residue(s) is based on the numbering of amino acid sequence of fHbp ID 1.

[0091] Combinations of substitutions may be included wherein the two substitutions are in different structural domains, and each independently decreases binding of fH to fHbp (e.g. one from the N-terminal domain (e.g., N115I, D121G, S128T or V131D) in combination with one from the C-terminal domain (e.g., D211A, K219N, G220S).

[0092] For example, in some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:2, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises: i) an N115I substitution; and ii) a D211A substitution.

[0093] As another example, in some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:2, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises: i) an N115I substitution; and ii) a K219N substitution.

[0094] As another example, in some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:2, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 22 for human fH, where the variant induces a bactericidal antibody response to at least one strain

of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises: i) an N115I substitution; and ii) a G220S substitution.

[0095] Also disclosed herein are variant fHbp polypeptides that have increased thermal stability compared to wild type fHbp ID22. In some cases, the variant fHbp may include the substitutions L130R and G133D relative to fHbp ID 22 (SEQ ID NO:2), wherein the variant fHbp comprises an amino acid sequence having greater than 85% amino acid sequence identity to SEQ ID NO:2, wherein the variant fHbp binds human factor H (fH) with an affinity that is 50% or less of the affinity of fHbp ID 22 for human fH, wherein the variant induces a bactericidal antibody response in a mammalian host, and the variant has a higher thermal stability than WT fHbp ID 22. The thermal stability of the variant fHbp may be higher than a WT fHbp (e.g., fHbp ID 22) by at least 5°C, 10°C, 15°C, 20°C, or more, e.g., higher by 5°C-30°C, 5°C-25°C, 5°C-20°C, 10°C-20°C, or 15°C-20°C. As used herein, “thermal stability” refers to stability of a protein when exposed to higher temperature; a thermal stability variant protein maintains its conformation at a higher temperature than a wild type protein. For example, the variant fHbp, that include the double mutation that increases thermal stability compared to thermal stability of wild type (WT) fHbp, e.g., WT fHbp ID 22, may unfold at a higher temperature compared to WT fHbp. In certain cases, the N-terminal domain of the variant fHbp may unfold at a higher temperature than the N-terminal domain of the WT fHbp (e.g., fHbp ID 22).

[0096] In certain embodiments, a variant of factor H binding protein (fHbp) is disclosed, wherein the variant comprises amino acid substitutions L130R and G133D and at least one of the substitutions: R80A, N115I, D121G, S128T, V131, D211A, E218A, K219 (e.g., K219N), G220 (e.g., G220S), E248A, G236I, T221A, and H223A relative to fHbp ID 22 (SEQ ID NO:2), wherein the variant comprises an amino acid sequence having greater than 85% amino acid sequence identity to SEQ ID NO:2, wherein the variant fHbp binds human factor H (fH) with an affinity that is 50% or less of the affinity of fHbp ID 22 for human fH, and wherein the variant induces a bactericidal antibody response in a mammalian host.

[0097] In some cases, the variant fHbp may include a combination of substitutions, such as, L130R, G133D, and at least one amino acid substitution selected from: a) N115I; b) D121G; c) S128T; d) V131D; e) K219 (e.g., K219N); and f) G220 (e.g., G220S), wherein the amino acid substitutions are relative to fHbp ID 22 (SEQ ID NO:2), wherein the variant fHbp comprises an amino acid sequence having greater than 85% amino acid sequence identity to SEQ ID NO:2, wherein the variant fHbp binds human factor H (fH) with an affinity that is 50% or less of the affinity of fHbp ID 22 for human fH, and wherein the variant induces a bactericidal antibody response in a mammalian host.

[0098] In certain embodiments, a variant of factor H binding protein (fHbp) is disclosed, wherein the variant comprises amino acid substitutions L130R and G133D and at least one of the substitutions: R80A, D211A, E218A, E248A, G236I, T221A, and H223A relative to fHbp ID 22 (SEQ ID NO:2), wherein the variant comprises an amino acid sequence having greater than 85% amino acid sequence identity to SEQ ID NO:2, wherein the variant fHbp binds human factor H (fH) with an affinity that is 50% or less of the affinity of fHbp ID 22 for human fH, and wherein the variant induces a bactericidal antibody response in a mammalian host.

[0099] Exemplary variant fHbp include a polypeptide having an amino acid sequence having greater than 85% amino acid sequence identity (e.g., at least 90% identity, at least 95%, at least 96% identity, at least 97% identity, at least 98% identity, at least 99% identity to SEQ ID NO:2, and including the following substitutions relative to the amino acid sequence of SEQ ID NO:2: L130R and G133D; L130R, G133D, and K219N; or L130R, G133D, and G220S.

[00100] Also disclosed herein are variant fHbp proteins that include one or more substitutions relative to amino acid sequence of fHbp ID 22 as set forth above and further include the substitutions disclosed in US2011/0256180.

Variants of fHbp ID 55

[00101] A “reference fHbp” from which a variant fHbp of the present disclosure is derived is in some cases fHbp ID 55. The amino acid sequence of fHbp ID 55 is set out below.

[00102] fHbp ID 55:

CSSGGGGSGGGGVADIGTGLADALTAPLDHKDKGLKSLTLEDSISQNGTLTSAQGAE
KTYGNGDSLNTGKLKNDKVSRFDFIRQIEVDGQLITLESGEFQVYKQSHSALTALQTEQE
QDPEHSEKMOVAKRRFRIGDIAGEHTSFDKLPKDVMATYRGTAFGSDDAGGKLTYTIDFA
AKQGHGKIEHLKSPELNVDLAVAYIKPDEKHHAVISGSVLYNQDEKGSYSLGIFGEKAQ
EVAGSAEVETANGIHHIGLAAKQ (SEQ ID NO:3).

[00103] In some cases, a variant fHbp of the present disclosure is a variant group 1 fHbp. In some cases, a variant fHbp of the present disclosure is a variant group 1 fHbp, and is a modular group IV fHbp.

[00104] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:3; where the variant fHbp comprises one or more amino acid substitutions relative to fHbp ID 55 such that the variant fHbp exhibits an affinity for human fH that is 85% or less of the binding affinity of fHbp ID 55 for human fH,

e.g., the variant fHbp exhibits an affinity for human fH that is from about 85% to about 75%, from about 75% to about 65%, from about 65% to about 55%, from about 55% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%, of the binding affinity of the affinity of fHbp ID 55 for human fH; and the variant fHbp induces a bactericidal immune response to at least one strain of *N. meningitidis* when administered to a mammalian host (e.g., a human; or a non-human animal model).

[00105] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:3, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 55 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises an amino acid substitution selected from at least one of: a) an amino acid substitution of the glutamic acid at position 92 (E92); b) an amino acid substitution of the serine at position 223 (S223); and c) an amino acid substitution of the histidine at position 248 (H248), relative to the amino acid sequence of fHbp ID 55, where the numbering of the amino acid residues is based on the numbering of fHbp ID 1.

[00106] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:3, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 55 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises an amino acid substitution of the glutamic acid at position 92 (E92). In some cases, the fHbp variant comprises an E92K substitution. Other amino acids with positively charged or aromatic side chains, such as arginine, histidine, phenylalanine, tyrosine or tryptophan, also may be substituted at this position. Thus, for example, in some cases, the fHbp variant comprises an E92R substitution, an

E92H substitution, an E92F substitution, an E92Y substitution, or an E92W substitution. As one example, a variant fHbp of the present disclosure can comprise the amino acid sequence depicted in Figure 31 and set forth in SEQ ID NO:16.

[00107] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:3, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 55 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises an amino acid substitution of the serine at position 223 (S223). In some cases, the fHbp variant comprises an S223R substitution. Other amino acids with positively charged or aromatic side chains, such as lysine, histidine, phenylalanine, tyrosine or tryptophan, also may be substituted at this position. Thus, for example, in some cases, the fHbp variant comprises an S223K substitution, an S223H substitution, an S223F substitution, an S223Y substitution, or an S223W substitution. As one example, a variant fHbp of the present disclosure can comprise the amino acid sequence depicted in Figure 32 and set forth in SEQ ID NO:17.

[00108] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:3, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 55 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises an amino acid substitution of the histidine at position 248 (H248). In some cases, the fHbp variant comprises an H248L substitution. Other amino acids with non-polar, negatively charged or aromatic side chains, such as isoleucine, valine, aspartate, glutamate, phenylalanine, tyrosine or tryptophan, also may be substituted at this position. Thus, for example, in some cases, the fHbp variant comprises an H248I substitution, an H248V substitution, an H248D substitution, an H248E substitution, an H248F substitution, an H248Y substitution, or an H248W substitution. As one

example, a variant fHbp of the present disclosure can comprise the amino acid sequence depicted in Figure 33 and set forth in SEQ ID NO:18.

Combinations of amino acid substitutions

[00109] In some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:3, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 55 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N. meningitidis* in a mammalian host, and where the variant fHbp comprises two or more amino acid substitutions selected from the group consisting of: a) an amino acid substitution of the glutamic acid at position 92 (E92); b) an amino acid substitution of the serine at position 223 (S223); and c) an amino acid substitution of the histidine at position 248 (H248), relative to fHbp ID 55, where the numbering of the residue is based on the numbering of amino acids in the sequence for fHbp ID 1.

[00110] Combinations of substitutions may be included wherein the two substitutions are in different structural domains, and each independently decreases binding of fH to fHbp (e.g., one substitution in the N-terminal domain, in combination with an amino acid substitution in the C-terminal domain. In some cases, a variant fHbp of the present disclosure comprises a first amino acid substitution within the N-terminal domain; and a second amino acid substitution within the C-terminal domain. In some cases, a variant fHbp of the present disclosure comprises a first amino acid substitution within the N-terminal domain; and a second amino acid substitution within the N-terminal domain. In some cases, a variant fHbp of the present disclosure comprises a first amino acid substitution within the C-terminal domain; and a second amino acid substitution within the C-terminal domain.

[00111] For example, in some cases, a variant fHbp of the present disclosure comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to SEQ ID NO:3, where the variant fHbp binds human fH with an affinity that 50% or less (e.g., from about 50% to about 45%, from about 45% to about 35%, from about 35% to about 25%, from about 25% to about 15%, from about 15% to about 10%, from about 10% to about 5%, from about 5% to about 2%, from about 2% to about 1%, or from about 1% to about 0.1%, or less than 0.1%) of the affinity of fHbp ID 55 for human fH, where the variant induces a bactericidal antibody response to at least one strain of *N.*

meningitidis in a mammalian host, and where the variant fHbp comprises: a) an amino acid substitution of the glutamic acid at position 92 (E92) and b) an amino acid substitution of the serine at position 223 (S223); or where the variant fHbp comprises: a) an amino acid substitution of the glutamic acid at position 92 (E92) and c) an amino acid substitution of the histidine at position 248 (H248); or where the variant fHbp comprises: b) an amino acid substitution of the serine at position 223 (S223) and c) an amino acid substitution of the histidine at position 248 (H248); or where the variant fHbp comprises: a) an amino acid substitution of the glutamic acid at position 92 (E92) and b) an amino acid substitution of the serine at position 223 (S223) and c) an amino acid substitution of the histidine at position 248 (H248), relative to fHbp ID 55, where the numbering of the residue is based on the numbering of amino acids in the sequence for fHbp ID 1.

[00112] Also disclosed herein are variant fHbp proteins that include one or more substitutions relative to amino acid sequence of fHbp ID 55 as set forth above and further include the substitutions disclosed in US2011/0256180.

Fusion polypeptides

[00113] A variant fHbp of the present disclosure can be a fusion polypeptide, e.g., a polypeptide comprising a variant fHbp as described above, and a heterologous polypeptide (e.g., a fusion partner). The fusion partner can be at the N-terminus of the variant fHbp, at the C-terminus of the variant fHbp, or at an internal site within the fHbp.

[00114] Suitable fusion partners include peptides and polypeptides that confer enhanced stability *in vivo* (e.g., enhanced serum half-life); provide ease of purification, e.g., (His)_n, e.g., 6His, and the like; provide for secretion of the fusion protein from a cell; provide an epitope tag, e.g., GST, hemagglutinin (HA; e.g., YPYDVPDYA; SEQ ID NO:26), FLAG (e.g., DYKDDDDK; SEQ ID NO:27), c-myc (e.g., EQKLISEEDL; SEQ ID NO:28), and the like; provide a detectable signal, e.g., an enzyme that generates a detectable product (e.g., β -galactosidase, luciferase), or a protein that is itself detectable, e.g., a green fluorescent protein, a yellow fluorescent protein, etc.; provides for multimerization, e.g., a multimerization domain such as an Fc portion of an immunoglobulin; and the like.

METHODS OF PRODUCTION

[00115] The fHbps of the present disclosure can be produced by any suitable method, including recombinant and non-recombinant methods (e.g., chemical synthesis). Where the subject fHbp is produced using recombinant techniques, the methods can involve any suitable construct and any suitable host cell, which can be a prokaryotic or eukaryotic cell, usually a bacterial or yeast host

cell, more usually a bacterial cell. Methods for introduction of genetic material into host cells include, for example, transformation, electroporation, conjugation, calcium phosphate methods and the like. The method for transfer can be selected so as to provide for stable expression of the introduced fHbp-encoding nucleic acid. The fHbp-encoding nucleic acid can be provided as an inheritable episomal element (e.g., plasmid) or can be genomically integrated.

[00116] The present disclosure provides nucleic acids (including isolated nucleic acids) that comprise a nucleotide sequence encoding a fHbp variant of the present disclosure. In some embodiments, the nucleotide sequence encoding the fHbp variant is operably linked to a transcriptional control element, e.g., a promoter. The promoter is in some cases constitutive. The promoter is in some cases inducible. In some cases, the promoter is suitable for use (e.g., active in) a prokaryotic host cell. In some cases, the promoter is suitable for use (e.g., active in) a eukaryotic host cell.

[00117] In some instances, a nucleic acid comprising a nucleotide sequence encoding a fHbp variant of the present disclosure is present in an expression vector. The present disclosure provides a recombinant expression vector (e.g., an isolated recombinant expression vector) that comprises a nucleotide sequence encoding a fHbp variant of the present disclosure. In some embodiments, the nucleotide sequence encoding the fHbp variant is operably linked to a transcriptional control element, e.g., a promoter. The promoter is in some cases constitutive. The promoter is in some cases inducible. In some cases, the promoter is suitable for use (e.g., active in) a prokaryotic host cell. In some cases, the promoter is suitable for use (e.g., active in) a eukaryotic host cell.

[00118] Suitable vectors for transferring fHbp-encoding nucleic acid can vary in composition. Integrative vectors can be conditionally replicative or suicide plasmids, bacteriophages, and the like. The constructs can include various elements, including for example, promoters, selectable genetic markers (e.g., genes conferring resistance to antibiotics (for instance kanamycin, erythromycin, chloramphenicol, or gentamycin)), origin of replication (to promote replication in a host cell, e.g., a bacterial host cell), and the like. The choice of vector will depend upon a variety of factors such as the type of cell in which propagation is desired and the purpose of propagation. Certain vectors are useful for amplifying and making large amounts of the desired DNA sequence. Other vectors are suitable for expression in cells in culture. Still other vectors are suitable for transfer and expression in cells in a whole animal. The choice of appropriate vector is well within the skill of the art. Many such vectors are available commercially.

[00119] In one example, the vector is an expression vector based on episomal plasmids containing selectable drug resistance markers and elements that provide for autonomous

replication in different host cells (e.g., in both *E. coli* and *N. meningitidis*). One example of such a “shuttle vector” is the plasmid pFP10 (Pagotto et al. (2000) Gene 244:13-19).

[00120] Constructs (recombinant vectors) can be prepared by, for example, inserting a polynucleotide of interest into a construct backbone, typically by means of DNA ligase attachment to a cleaved restriction enzyme site in the vector. Alternatively, the desired nucleotide sequence can be inserted by homologous recombination or site-specific recombination. Typically homologous recombination is accomplished by attaching regions of homology to the vector on the flanks of the desired nucleotide sequence, while site-specific recombination can be accomplished through use of sequences that facilitate site-specific recombination (e.g., cre-lox, att sites, etc.). Nucleic acid containing such sequences can be added by, for example, ligation of oligonucleotides, or by polymerase chain reaction using primers comprising both the region of homology and a portion of the desired nucleotide sequence.

[00121] Vectors can provide for extrachromosomal maintenance in a host cell or can provide for integration into the host cell genome. Vectors are amply described in numerous publications well known to those in the art, including, e.g., Short Protocols in Molecular Biology, (1999) F. Ausubel, et al., eds., Wiley & Sons. Vectors may provide for expression of the nucleic acids encoding the subject fHbp, may provide for propagating the subject nucleic acids, or both.

[00122] Examples of vectors that may be used include but are not limited to those derived from recombinant bacteriophage DNA, plasmid DNA or cosmid DNA. For example, plasmid vectors such as pBR322, pUC 19/18, pUC 118, 119 and the M13 mp series of vectors may be used. pET21 is also an expression vector that may be used. Bacteriophage vectors may include λgt10, λgt11, λgt18-23, λZAP/R and the EMBL series of bacteriophage vectors. Further vectors that may be utilized include, but are not limited to, pJB8, pCV 103, pCV 107, pCV 108, pTM, pMCS, pNNL, pHSG274, COS202, COS203, pWE15, pWE16 and the charomid 9 series of vectors.

[00123] For expression of a subject fHbp, an expression cassette may be employed. Thus, the present disclosure provides a recombinant expression vector comprising a subject nucleic acid. The expression vector provides transcriptional and translational regulatory sequences, and may provide for inducible or constitutive expression, where the coding region is operably linked under the transcriptional control of the transcriptional initiation region, and a transcriptional and translational termination region. These control regions may be native to an fHbp from which the subject fHbp is derived, or may be derived from exogenous sources. In general, the transcriptional and translational regulatory sequences may include, but are not limited to, promoter sequences, ribosomal binding sites, transcriptional start and stop sequences,

translational start and stop sequences, and enhancer or activator sequences. Promoters can be either constitutive or inducible, and can be a strong constitutive promoter (e.g., T7, and the like).

[00124] Expression vectors generally have convenient restriction sites located near the promoter sequence to provide for the insertion of nucleic acid sequences encoding proteins of interest. A selectable marker operative in the expression host may be present to facilitate selection of cells containing the vector. In addition, the expression construct may include additional elements. For example, the expression vector may have one or two replication systems, thus allowing it to be maintained in organisms, for example in mammalian or insect cells for expression and in a prokaryotic host for cloning and amplification. In addition the expression construct may contain a selectable marker gene to allow the selection of transformed host cells. Selection genes are well known in the art and will vary with the host cell used.

[00125] It should be noted that fHbps of the present disclosure may comprise additional elements, such as a detectable label, e.g., a radioactive label, a fluorescent label, a biotin label, an immunologically detectable label (e.g., a hemagglutinin tag, a poly-Histidine tag) and the like. Additional elements of fHbp can be provided to facilitate isolation (e.g., biotin tag, immunologically detectable tag) through various methods (e.g., affinity capture, etc.). The subject fHbp can optionally be immobilized on a support through covalent or non-covalent attachment.

[00126] Isolation and purification of fHbp can be accomplished according to methods known in the art. For example, fHbp can be isolated from a lysate of cells genetically modified to express a fHbp, or from a synthetic reaction mix, by immunoaffinity purification, which generally involves contacting the sample with an anti-fHbp antibody (e.g., an anti-fHbp monoclonal antibody (mAb), such as a JAR 4 MAb or other appropriate JAR MAb known in the art), washing to remove non-specifically bound material, and eluting specifically bound fHbp. Isolated fHbp can be further purified by dialysis and other methods normally employed in protein purification methods. In one example, the fHbp can be isolated using metal chelate chromatography methods.

Host cells

[00127] Any of a number of suitable host cells can be used in the production of fHbp. In general, the fHbp described herein may be expressed in prokaryotes or eukaryotes, e.g., bacteria such as *Escherichia coli* or *Neisseria* (e.g., *N. meningitidis*) in accordance with conventional techniques. Thus, the present disclosure further provides a genetically modified *in vitro* host cell, which contains a nucleic acid encoding a subject fHbp. Host cells for production (including large scale production) of a subject fHbp can be selected from any of a variety of available host cells. Examples of host cells for expression include those of a prokaryotic or eukaryotic unicellular organism, such as bacteria (e.g., *Escherichia coli* strains), yeast (e.g., *Saccharomyces cerevisiae*,

Pichia spp., and the like), and may include host cells originally derived from a higher organism such as insects, vertebrates, e.g., mammals. Suitable mammalian cell lines include, but are not limited to, HeLa cells (e.g., American Type Culture Collection (ATCC) No. CCL-2), CHO cells (e.g., ATCC Nos. CRL9618, CCL61, CRL9096), 293 cells (e.g., ATCC No. CRL-1573), Vero cells, NIH 3T3 cells (e.g., ATCC No. CRL-1658), Huh-7 cells, BHK cells (e.g., ATCC No. CCL10), PC12 cells (ATCC No. CRL1721), COS cells, COS-7 cells (ATCC No. CRL1651), RAT1 cells, mouse L cells (ATCC No. CCL1.3), human embryonic kidney (HEK) cells (ATCC No. CRL1573), HLHepG2 cells, and the like.). In some cases, bacterial host cells and yeast host cells are of particular interest for subject fHbp production.

[00128] Subject fHbps can be prepared in substantially pure or substantially isolated form (i.e., substantially free from other Neisserial or host cell polypeptides) or substantially isolated form. The subject fHbp can be present in a composition that is enriched for the polypeptide relative to other components that may be present (e.g., other polypeptides or other host cell components). Purified subject fHbp can be provided such that the polypeptide is present in a composition that is substantially free of other expressed polypeptides, e.g., less than 90%, usually less than 60% and more usually less than 50% of the composition is made up of other expressed polypeptides.

Host cells for vesicle production

[00129] Where a subject fHbp is to be provided in a membrane vesicle (as discussed in more detail below), a Neisserial host cell is genetically modified to express a subject fHbp. Any of a variety of *Neisseria* spp. strains can be modified to produce a subject fHbp, and, optionally, which produce or can be modified to produce other antigens of interest, such as PorA, can be used in the methods disclosed herein.

[00130] Methods and vectors to provide for genetic modification of Neisserial strains and expression of a desired polypeptide are known in the art. Examples of vectors and methods can be found in WO 02/09746 and O'Dwyer et al. (2004) *Infect Immun* 72:6511-80. Strong promoters, particularly constitutive strong promoters are of particular interest. Examples of promoters include the promoters of porA, porB, lbpB, lbpB, p110, hpuAB, lgtF, opa, p110, lst, hpuAB, and rmp.

[00131] Pathogenic *Neisseria* spp. or strains derived from pathogenic *Neisseria* spp., particularly strains pathogenic for humans or derived from strains pathogenic or commensal for humans, are of particular interest for use in membrane vesicle production. Examples of *Neisseria* spp. include *N. meningitidis*, *N. flavescens*, *N. gonorrhoeae*, *N. lactamica*, *N. polysaccharea*, *N. cinerea*, *N. mucosa*, *N. subflava*, *N. sicca*, *N. elongata*, and the like.

[00132] *N. meningitidis* strains are of particular interest for genetic modification to express the subject fHbps and for use in vesicle production. The strain used for vesicle production can be selected according to a number of different characteristics that may be desired. For example, the strain may be selected according to: a desired PorA type (a “serosubtype”), capsular group, serotype, and the like; decreased capsular polysaccharide production; and the like. For example, the production strain can produce any desired PorA polypeptide, and may express one or more PorA polypeptides (either naturally or due to genetic engineering). Examples of strains include those that produce a PorA polypeptide which confers a serosubtype of P1.7,16; P1.19,15; P1.7,1; P1.5,2; P1.22a,14; P1.14 ; P1.5,10; P1.7,4; P1.12,13; as well as variants of such PorA polypeptides which may or may not retain reactivity with conventional serologic reagents used in serosubtyping. Also of interest are PorA polypeptides characterized according to PorA variable region (VR) typing (see, e.g., Russell et al. (2004) *Emerging Infect Dis* 10:674-678; Sacchi CT et al. (1998) *Clin Diagn Lab Immunol* 5:845-55; Sacchi et al (2000) *J. Infect Dis* 182:1169-1176). A substantial number of distinct VR types have been identified, which can be classified into VR1 and VR2 family “prototypes”. A web-accessible database describing this nomenclature and its relationship to previous typing schemes is found at neisseria.org/nm/typing/pora. Alignments of certain PorA VR1 and VR2 types are provided in Russell et al. (2004) *Emerging Infect Dis* 10:674-678.

[00133] Alternatively or in addition, the production strain can be a capsule deficient strain. Capsule deficient strains can provide vesicle-based vaccines that provide for a reduced risk of eliciting a significant autoantibody response in a subject to whom the vaccine is administered (e.g., due to production of antibodies that cross-react with sialic acid on host cell surfaces). “Capsule deficient” or “deficient in capsular polysaccharide” as used herein refers to a level of capsular polysaccharide on the bacterial surface that is lower than that of a naturally-occurring strain or, where the strain is genetically modified, is lower than that of a parental strain from which the capsule deficient strain is derived. A capsule deficient strain includes strains that are decreased in surface capsular polysaccharide production by at least 10%, 20%, 25%, 30%, 40%, 50%, 60%, 75%, 80%, 85%, 90% or more, and includes strains in which capsular polysaccharide is not detectable on the bacterial surface (e.g., by whole cell enzyme-linked immunosorbent assay (ELISA) using an anti-capsular polysaccharide antibody).

[00134] Capsule deficient strains include those that are capsule deficient due to a naturally-occurring or recombinantly-generated genetic modification. Naturally-occurring capsule deficient strains (see, e.g., Dolan-Livengood et al. (2003) *J. Infect. Dis.* 187:1616-28), as well as methods of identifying and/or generating capsule-deficient strains (see, e.g., Fisseha et al. (2005)

Infect. Immun. 73:4070-4080; Stephens et al. (1991) Infect Immun 59:4097-102; Frosch et al. (1990) Mol Microbiol.4:1215-1218) are known in the art.

[00135] Modification of a Neisserial host cell to provide for decreased production of capsular polysaccharide may include modification of one or more genes involved in capsule synthesis, where the modification provides for, for example, decreased levels of capsular polysaccharide relative to a parent cell prior to modification. Such genetic modifications can include changes in nucleotide and/or amino acid sequences in one or more capsule biosynthesis genes rendering the strain capsule deficient (e.g., due to one or more insertions, deletions, substitutions, and the like in one or more capsule biosynthesis genes). Capsule deficient strains can lack or be non-functional for one or more capsule genes.

[00136] Of particular interest are strains that are deficient in sialic acid biosynthesis. Such strains can provide for production of vesicles that have reduced risk of eliciting anti-sialic acid antibodies that cross-react with human sialic acid antigens, and can further provide for improved manufacturing safety. Strains having a defect in sialic acid biosynthesis (due to either a naturally occurring modification or an engineered modification) can be defective in any of a number of different genes in the sialic acid biosynthetic pathway. Of particular interest are strains that are defective in a gene product encoded by the N-acetylglucosamine-6-phosphate 2-epimerase gene (known as *synX* AAF40537.1 or *siaA* AAA20475), with strains having this gene inactivated being of especial interest. For example, in one embodiment, a capsule deficient strain is generated by disrupting production of a functional *synX* gene product (see, e.g., Swartley et al. (1994) J Bacteriol. 176:1530-4).

[00137] Capsule-deficient strains can also be generated from naturally-occurring strains using non-recombinant techniques, e.g., by use of bactericidal anti-capsular antibodies to select for strains with reduced levels of capsular polysaccharide.

[00138] Where the disclosure involves use of two or more strains (e.g., to produce antigenic compositions containing a subject fHbp-presenting vesicles from different strains), the strains can be selected so as to differ in one or more strain characteristics, e.g., to provide for vesicles that differ in the subject fHbp used, PorA, and the like.

Preparation of Vesicles

[00139] The antigenic compositions contemplated by the present disclosure generally include vesicles prepared from Neisserial cells that express a subject fHbp. As referred to herein “vesicles” is meant to encompass outer membrane vesicles as well as microvesicles (which are also referred to as blebs).

[00140] The antigenic composition can contain outer membrane vesicles (OMV) prepared from the outer membrane of a cultured strain of *Neisseria meningitidis* spp. genetically modified to express a subject fHbp. OMVs may be obtained from *Neisseria meningitidis* grown in broth or solid medium culture, preferably by separating the bacterial cells from the culture medium (e.g. by filtration or by a low-speed centrifugation that pellets the cells, or the like), lysing the cells (e.g. by addition of detergent, osmotic shock, sonication, cavitation, homogenization, or the like) and separating an outer membrane fraction from cytoplasmic molecules (e.g. by filtration; or by differential precipitation or aggregation of outer membranes and/or outer membrane vesicles, or by affinity separation methods using ligands that specifically recognize outer membrane molecules; or by a high-speed centrifugation that pellets outer membranes and/or outer membrane vesicles, or the like); outer membrane fractions may be used to produce OMVs.

[00141] The antigenic composition can contain microvesicles (MV) (or “blebs”) containing subject fHbps, where the MV or blebs are released during culture of a *Neisseria meningitidis* strain genetically modified to express a subject fHbp. For example, MVs may be obtained by culturing a strain of *Neisseria meningitidis* in broth culture medium, separating whole cells from the broth culture medium (e.g. by filtration, or by a low-speed centrifugation that pellets only the cells and not the smaller blebs, or the like), and then collecting the MVs that are present in the cell-free culture medium (e.g. by filtration, differential precipitation or aggregation of MVs, or by a high-speed centrifugation that pellets the blebs, or the like). Strains for use in production of MVs can generally be selected on the basis of the amount of blebs produced in culture (e.g., bacteria can be cultured in a reasonable number to provide for production of blebs suitable for isolation and administration in the methods described herein). An exemplary strain that produces high levels of blebs is described in PCT Publication No. WO 01/34642. In addition to bleb production, strains for use in MV production may also be selected on the basis of NspA production, where strains that produce higher levels of NspA may be of particular interest (for examples of *N. meningitidis* strains having different NspA production levels, see, e.g., Moe et al. (1999) *Infect. Immun.* 67: 5664). Other strains of interest for use in production of blebs include strains having an inactivated GNA33 gene, which encodes a lipoprotein required for cell separation, membrane architecture and virulence (see, e.g., Adu-Bobie et al. (2004) *Infect. Immun.* 72:1914-1919).

[00142] The antigenic compositions of the present disclosure can contain vesicles from one strain, or from 2, 3, 4, 5 or more strains, which strains may be homologous or heterologous, usually heterologous, to one another. For example, the strains may be homologous or heterologous with respect to PorA and/or the fHbp from which the subject fHbp is derived. The vesicles can be prepared from strains that express more than one subject fHbp (e.g., 1, 2, 3, or

more subject fHbp) which may be composed of fHbp amino acid sequences from different variants (v.1, v.2, or v.3) or subvariants (e.g., a subvariant of v.1, v.2, or v.3).

[00143] The antigenic compositions can comprise a mixture of OMVs and MVs presenting the same or different subject fHbps, where the subject fHbps may optionally present epitopes from different combinations of fHbp variants and/or subvariants and where the OMVs and/or MVs may be from the same or different strains. Vesicles from different strains can be administered as a mixture, or can be administered serially.

[00144] Where desired (e.g., where the strains used to produce vesicles are associated with endotoxin or particular high levels of endotoxin), the vesicles are optionally treated to reduce endotoxin, e.g., to reduce toxicity following administration. Although less desirable as discussed below, reduction of endotoxin can be accomplished by extraction with a suitable detergent (for example, BRIJ-96, sodium deoxycholate, sodium lauroylsarcosinate, Empigen BB, Triton X-100TM, non-ionic detergent TWEENTM 20 (sorbitan monolaurate polyoxyethylene), non-ionic detergent TWEENTM 80, at a concentration of 0.1-10%, e.g., 0.5-2%, and sodium dodecyl sulfate (SDS)). Where detergent extraction is used, it is preferable to use a detergent other than deoxycholate.

[00145] The vesicles of the antigenic compositions can be prepared without detergent, e.g., without use of deoxycholate. Although detergent treatment is useful to remove endotoxin activity, it may deplete the native fHbp lipoprotein and/or subject fHbp (including lipidated fHbp) by extraction during vesicle production. Thus it may be particularly desirable to decrease endotoxin activity using technology that does not require a detergent. In one approach, strains that are relatively low producers of endotoxin (lipopolysaccharide, LPS) are used so as to avoid the need to remove endotoxin from the final preparation prior to use in humans. For example, the vesicles can be prepared from *Neisseria* mutants in which lipooligosaccharide or other antigens that may be undesirable in a vaccine (e.g. Rmp) is reduced or eliminated.

[00146] Vesicles can be prepared from *N. meningitidis* strains that contain genetic modifications that result in decreased or no detectable toxic activity of lipid A. For example, such strain can be genetically modified in lipid A biosynthesis (Steeghs et al. (1999) Infect Immun 67:4988-93; van der Ley et al. (2001) Infect Immun 69:5981-90; Steeghs et al. (2004) J Endotoxin Res 10:113-9; Fissha et al. (2005) Infect Immun 73:4070). The immunogenic compositions may be detoxified by modification of LPS, such as downregulation and/or inactivation of the enzymes encoded by lpxL1 or lpxL2, respectively. Production of a penta-acylated lipid A made in lpxL1 mutants indicates that the enzyme encoded by lpxL1 adds the C12 to the N-linked 3-OH C14 at the 2' position of GlcN II. The major lipid A species found in lpxL2 mutants is tetra-acylated, indicating the enzyme encoded by lpxL2 adds the other C12, i.e., to the N-linked 3-OH C14 at

the 2 position of GlcN I. Mutations resulting in a decreased (or no) expression of these genes (or decreased or no activity of the products of these genes) result in altered toxic activity of lipid A (van der Ley et al. (2001) *Infect Immun* 69:5981-90). Tetra-acylated (lpxL2 mutant) and penta acylated (lpxL1 mutant) lipid A are less toxic than the wild-type lipid A. Mutations in the lipid A 4'-kinase encoding gene (lpxK) also decrease the toxic activity of lipid A. Of particular interest for use in production of vesicles (e.g., MV or OMV) are *N. meningitidis* strains genetically modified so as to provide for decreased or no detectable functional LpxL1-encoded protein, e.g., where the Neisseria bacterium (e.g., *N. meningitidis* strain) is genetically modified to provide for decreased or no activity of a gene product of the lpxL1 gene. For example, the Neisseria bacterium can be genetically modified to have an lpxL1 gene knockout, e.g., where the lpxL1 gene is disrupted. See, e.g., US Patent Publication No. 2009/0035328. The Neisseria bacterium can be genetically modified to provide for decreased or no activity of a gene product of the lpxL2 gene. The Neisseria bacterium can be genetically modified to provide for decreased or no activity of a gene product of the lpxL1 gene and the lpxL2 gene. Such vesicles provide for reduced toxicity as compared to *N. meningitidis* strains that are wild-type for LPS production, while retaining immunogenicity of subject fHbp.

[00147] LPS toxic activity can also be altered by introducing mutations in genes/loci involved in polymyxin B resistance (such resistance has been correlated with addition of aminoarabinose on the 4' phosphate of lipid A). These genes/loci could be pmrE that encodes a UDP-glucose dehydrogenase, or a region of antimicrobial peptide-resistance genes common to many enterobacteriaceae which could be involved in aminoarabinose synthesis and transfer. The gene pmrF that is present in this region encodes a dolichol-phosphate manosyl transferase (Gunn J. S., Kheng, B. L., Krueger J., Kim K., Guo L., Hackett M., Miller S. I. 1998. *Mol. Microbiol.* 27: 1171-1182).

[00148] Mutations in the PhoP-PhoQ regulatory system, which is a phospho-relay two component regulatory system (e.g., PhoP constitutive phenotype, PhoPc), or low Mg⁺⁺ environmental or culture conditions (that activate the PhoP-PhoQ regulatory system) lead to the addition of aminoarabinose on the 4'-phosphate and 2-hydroxymyristate replacing myristate (hydroxylation of myristate). This modified lipid A displays reduced ability to stimulate E-selectin expression by human endothelial cells and TNF secretion from human monocytes.

[00149] Polymyxin B resistant strains are also suitable for use, as such strains have been shown to have reduced LPS toxicity (see, e.g., van der Ley et al. (1994) In: *Proceedings of the ninth international pathogenic Neisseria conference*. The Guildhall, Winchester, England). Alternatively, synthetic peptides that mimic the binding activity of polymyxin B may be added

to the antigenic compositions to reduce LPS toxic activity (see, e.g., Rustici et al. (1993) *Science* 259:361-365; Porro et al. (1998) *Prog Clin Biol Res.* 397:315-25).

[00150] Endotoxin can also be reduced through selection of culture conditions. For example, culturing the strain in a growth medium containing 0.1 mg-100 mg of aminoarabinose per liter medium provides for reduced lipid toxicity (see, e.g., WO 02/097646).

COMPOSITIONS AND FORMULATIONS

[00151] “Compositions”, “antigen composition”, “antigenic composition” or “immunogenic composition” is used herein as a matter of convenience to refer generically to compositions comprising a subject fHbp as disclosed herein, which subject fHbp may be optionally conjugated to further enhance immunogenicity. Compositions useful for eliciting antibodies, e.g., antibodies against *Neisseria meningitidis*, e.g., bactericidal antibodies to *Neisseria meningitidis*, in a human are specifically contemplated by the present disclosure. Antigenic compositions can contain 1, 2, or more different subject fHbps. Where there is more than one type of fHbp, each subject fHbps may present epitopes from different combinations of fHbp variants and/or subvariants.

[00152] Antigenic compositions contain an immunologically effective amount of a subject fHbp, and may further include other compatible components, as needed. Compositions of the present disclosure can contain fHbp that are low fH binders. The composition contain one or more fHbp, in which at least one fHbp is a low fH binder. Where there is more than one fHbp in a composition, each fHbp may be different (e.g. in amino acid sequences and/or conjugation).

[00153] In some cases, an antigenic composition of the present disclosure comprises only one fHbp variant of the present disclosure. In some cases, an antigenic composition of the present disclosure comprises two or more different fHbp variants of the present disclosure. As non-limiting examples, in some cases, an antigenic composition of the present disclosure comprises:

[00154] 1) a first variant of fHbp ID 1, where the first fHbp ID 1 variant comprises an amino acid substitution at Q38 (e.g., Q38R); and a second variant of fHbp ID 1, where the second fHbp ID 1 variant comprises an amino acid substitution at E92 (e.g., E92K);

[00155] 2) a first variant of fHbp ID 1, where the first fHbp ID 1 variant comprises an amino acid substitution at Q38 (e.g., Q38R); and a second variant of fHbp ID 1, where the second fHbp ID 1 variant comprises an amino acid substitution at R130 (e.g., R130G);

[00156] 3) a first variant of fHbp ID 1, where the first fHbp ID 1 variant comprises an amino acid substitution at Q38 (e.g., Q38R); and a second variant of fHbp ID 1, where the second fHbp ID 1 variant comprises an amino acid substitution at S223 (e.g., S223R);

[00157] 4) a first variant of fHbp ID 1, where the first fHbp ID 1 variant comprises an amino acid substitution at Q38 (e.g., Q38R); and a second variant of fHbp ID 1, where the second fHbp ID 1 variant comprises an amino acid substitution at H248 (e.g., H248L);

[00158] 5) a variant of fHbp ID 22, where the fHbp ID 22 variant comprises an amino acid substitution at N115 (e.g., N115I); and a variant of fHbp ID 1, where the fHbp ID 1 variant comprises an amino acid substitution at Q38 (e.g., Q38R);

[00159] 6) a variant of fHbp ID 22, where the fHbp ID 22 variant comprises an amino acid substitution at D121 (e.g., D121G); and a variant of fHbp ID 1, where the fHbp ID 1 variant comprises an amino acid substitution at E92 (e.g., E92K);

[00160] 7) a variant of fHbp ID 22, where the fHbp ID 22 variant comprises an amino acid substitution at S128 (e.g., S128T); and a variant of fHbp ID 1, where the fHbp ID 1 variant comprises an amino acid substitution at H248 (e.g., H248L);

[00161] 8) a variant of fHbp ID 22, where the fHbp ID 22 variant comprises an amino acid substitution at V131 (e.g., V131D); and a variant of fHbp ID 1, where the fHbp ID 1 variant comprises an amino acid substitution at Q38 (e.g., Q38R);

[00162] 9) a variant of fHbp ID 22, where the fHbp ID 22 variant comprises an amino acid substitution at K219 (e.g., K219N); and a variant of fHbp ID 1, where the fHbp ID 1 variant comprises an amino acid substitution at Q38 (e.g., Q38R);

[00163] 10) a variant of fHbp ID 22, where the fHbp ID 22 variant comprises an amino acid substitution at G220 (e.g., G220S); and a variant of fHbp ID 1, where the fHbp ID 1 variant comprises an amino acid substitution at Q38 (e.g., Q38R);

[00164] 11) a variant of fHbp ID 22, where the fHbp ID 22 variant comprises an amino acid substitution at N115 (e.g., N115I); and a variant of fHbp ID 55, where the fHbp ID 55 variant comprises an amino acid substitution at E92 (e.g., E92K);

[00165] 12) a variant of fHbp ID 22, where the fHbp ID 22 variant comprises an amino acid substitution at D121 (e.g., D121G); and a variant of fHbp ID 55, where the fHbp ID 55 variant comprises an amino acid substitution at S223 (e.g., S223R);

[00166] 13) a variant of fHbp ID 22, where the fHbp ID 22 variant comprises an amino acid substitution at S128 (e.g., S128T); and a variant of fHbp ID 55, where the fHbp ID 55 variant comprises an amino acid substitution at H248 (e.g., H248L);

[00167] 14) a variant of fHbp ID 22, where the fHbp ID 22 variant comprises an amino acid substitution at V131 (e.g., V131D); and a variant of fHbp ID 55, where the fHbp ID 55 variant comprises an amino acid substitution at E92 (e.g., E92K);

[00168] 15) a variant of fHbp ID 22, where the fHbp ID 22 variant comprises an amino acid substitution at K219 (e.g., K219N); and a variant of fHbp ID 55, where the fHbp ID 55 variant comprises an amino acid substitution at E92 (e.g., E92K);

[00169] 16) a variant of fHbp ID 22, where the fHbp ID 22 variant comprises an amino acid substitution at G220 (e.g., G220S); and a variant of fHbp ID 55, where the fHbp ID 55 variant comprises an amino acid substitution at E92 (e.g., E92K);

[00170] 17) a first variant of fHbp ID 1, where the first fHbp ID 1 variant comprises an amino acid substitution at E92 (e.g., E92K); and a second variant of fHbp ID 1, where the second fHbp ID 1 variant comprises an amino acid substitution at H248 (e.g., H248L);

[00171] 18) a first variant of fHbp ID 1, where the first fHbp ID 1 variant comprises an amino acid substitution at E92 (e.g., E92K); and a second variant of fHbp ID 1, where the second fHbp ID 1 variant comprises an amino acid substitution at S223 (e.g., S223R);

[00172] 19) a first variant of fHbp ID 22, where the first fHbp ID 22 variant comprises an amino acid substitution at N115 (e.g., N115I); and a second variant of fHbp ID 22, where the second fHbp ID 22 variant comprises an amino acid substitution at D211 (e.g., D211A);

[00173] 20) a first variant of fHbp ID 22, where the first fHbp ID 22 variant comprises an amino acid substitution at N115 (e.g., N115I); and a second variant of fHbp ID 22, where the second fHbp ID 22 variant comprises an amino acid substitution at K219 (e.g., K219N);

[00174] 21) a first variant of fHbp ID 22, where the first fHbp ID 22 variant comprises an amino acid substitution at N115 (e.g., N115I); and a second variant of fHbp ID 22, where the second fHbp ID 22 variant comprises an amino acid substitution at G220 (e.g., G220S);

[00175] 22) a first variant of fHbp ID 22, where the first fHbp ID 22 variant comprises an amino acid substitution at D121 (e.g., D121G); and a second variant of fHbp ID 22, where the second fHbp ID 22 variant comprises an amino acid substitution at G220 (e.g., G220S);

[00176] 23) a first variant of fHbp ID 22, where the first fHbp ID 22 variant comprises an amino acid substitution at S128 (e.g., S128T); and a second variant of fHbp ID 22, where the second fHbp ID 22 variant comprises an amino acid substitution at G220 (e.g., G220S);

[00177] 24) a first variant of fHbp ID 22, where the first fHbp ID 22 variant comprises an amino acid substitution at V131 (e.g., V131D); and a second variant of fHbp ID 22, where the second fHbp ID 22 variant comprises an amino acid substitution at G220 (e.g., G220S);

[00178] 25) a first variant of fHbp ID 55, where the first fHbp ID 55 variant comprises an amino acid substitution at E92 (e.g., E92K); and a second variant of fHbp ID 55, where the second fHbp ID 55 variant comprises an amino acid substitution at S223 (e.g., S223R);

[00179] 26) a first variant of fHbp ID 55, where the first fHbp ID 55 variant comprises an amino acid substitution at E92 (e.g., E92K); and a second variant of fHbp ID 55, where the second fHbp ID 55 variant comprises an amino acid substitution at H248 (e.g., H248L);

[00180] 27) a variant of fHbp ID22 comprising amino acid substitutions: L130R and G133D and a variant of fHbp ID 1, where the fHbp ID 1 variant comprises an amino acid substitution at S223 (e.g., S223R);

[00181] 28) a variant of fHbp ID22 comprising amino acid substitutions: L130R and G133D and a variant of fHbp ID 1, where the fHbp ID 1 variant comprises an amino acid substitution at H248 (e.g., H248L);

[00182] 29) a variant of fHbp ID22 comprising amino acid substitutions: L130R, G133D, and K219N and a variant of fHbp ID 1, where the fHbp ID 1 variant comprises an amino acid substitution at S223 (e.g., S223R) or H248 (e.g., H248L); or

[00183] 30) a variant of fHbp ID22 comprising amino acid substitutions: L130R, G133D, and G220S and a variant of fHbp ID 1, where the fHbp ID 1 variant comprises an amino acid substitution at S223 (e.g., S223R) or H248 (e.g., H248L).

[00184] Immunogenic compositions contemplated by the present disclosure include, but are not limited to, compositions comprising:

[00185] 1) at least one variant fHbp of the present disclosure; and

[00186] 2) NspA;

[00187] where the fHbp and/or NspA can be provided as recombinant proteins and/or in a vesicle-based composition (e.g., OMV or MV). It should be noted that where the composition includes both NspA and a fHbp, the bactericidal activity of antibodies elicited by administration of the composition can result from cooperation of antibodies to one or both antigens. Examples of immunogenic compositions provided by the present disclosure include:

[00188] a) an immunogenic composition that comprises a fHbp variant as described above (e.g., where the variant fHbp elicits a bactericidal antibody response to at least one *Neisseria meningitidis* strain);

[00189] b) an immunogenic composition that comprises a fHbp variant as described above (e.g., where the variant fHbp elicits a bactericidal antibody response to at least one *Neisseria meningitidis* strain); and a recombinant NspA protein;

[00190] c) an immunogenic composition that comprises a native OMV obtained from a genetically modified *Neisseria* host cell that is genetically modified with a nucleic acid encoding a variant fHbp of the present disclosure, such that the encoded variant fHbp is produced by the genetically modified host cell, where the OMV comprises the encoded variant fHbp; and

[00191] d) an immunogenic composition that comprises a native OMV obtained from a genetically modified *Neisseria* host cell that is genetically modified with a nucleic acid encoding a variant fHbp of the present disclosure, such that the encoded non-naturally occurring fHbp is produced by the genetically modified host cell, where the OMV comprises the encoded variant fHbp; and where the *Neisseria* host cell also produces higher levels of NspA, such that the OMV also comprises NspA. For example, the *Neisseria* host cell can be one that is genetically modified for increased expression of NspA.

[00192] By “immunologically effective amount” is meant that the administration of that amount to an individual, either in a single dose, as part of a series of the same or different antigenic compositions, is effective to elicit an antibody response effective for treatment or prevention of a symptom of, or disease caused by, for example, infection by *Neisseria*, particularly *N. meningitidis*, more particularly Group B *N. meningitidis*. This amount varies depending upon the health and physical condition of the individual to be treated, age, the capacity of the individual's immune system to produce antibodies, the degree of protection desired, the formulation of the vaccine, the treating clinician's assessment of the medical situation, and other relevant factors. It is expected that the amount will fall in a relatively broad range that can be determined through routine trials.

[00193] Amino acid sequences of NspA polypeptides are known in the art. See, e.g., WO 96/29412; and Martin et al. (1997) *J. Exp. Med.* 185:1173; GenBank Accession No. U52066; and GenBank Accession No. AAD53286. An “NspA polypeptide” can comprise an amino acid sequence having at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or 100%, amino acid sequence identity to a contiguous stretch of from about 75 amino acids to about 100 amino acids, from about 100 amino acids to about 150 amino acids or from about 150 amino acids to about 174 amino acids, of the amino acid sequence depicted in Figure 40 and set forth in SEQ ID NO:25. An “NspA polypeptide” can comprise an amino acid sequence having at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or 100%, amino acid sequence identity to a contiguous stretch of from about 75 amino acids to about 100 amino acids, or from about 100 amino acids to about 155 amino acids, of amino acids 20 to 174 of the amino acid sequence depicted in Figure 40 and set forth in SEQ ID NO:25. In some cases, the NspA polypeptide lacks a signal sequence; in other cases (e.g., for expression in a host cell), the NspA polypeptide includes a signal sequence.

[00194] Dosage regimen may be a single dose schedule or a multiple dose schedule (e.g., including booster doses) with a unit dosage form of the antigenic composition administered at different times. The term “unit dosage form,” as used herein, refers to physically discrete units

suitable as unitary dosages for human and animal subjects, each unit containing a predetermined quantity of the antigenic compositions of the present disclosure in an amount sufficient to produce the desired effect, which compositions are provided in association with a pharmaceutically acceptable excipient (e.g., pharmaceutically acceptable diluent, carrier or vehicle). The antigenic composition may be administered in conjunction with other immunoregulatory agents.

[00195] Antigenic compositions can be provided in a pharmaceutically acceptable excipient, which can be a solution such as a sterile aqueous solution, often a saline solution, or they can be provided in powder form. Such excipients can be substantially inert, if desired.

[00196] In some embodiments, a subject immunogenic composition comprises a subject fHbp present in a vesicle. In some embodiments, a subject immunogenic composition comprises a subject fHbp present in an MV. In some embodiments, a subject immunogenic composition comprises a subject fHbp present in an OMV. In some embodiments, a subject immunogenic composition comprises a mixture of MV and OMV comprising a subject fHbp. Vesicles, such as MV and OMV, are described above.

[00197] The antigenic compositions can further contain an adjuvant. Examples of known suitable adjuvants that can be used in humans include, but are not necessarily limited to, an aluminum adjuvant (e.g., aluminum phosphate, or aluminum hydroxide), MF59 (4.3% w/v squalene, 0.5% w/v Tween 80TM, 0.5% w/v Span 85), a CpG-containing nucleic acid (where the cytosine is unmethylated), QS21, MPL, 3DMPL, extracts from Aquilla, ISCOMS, LT/CT mutants, poly(D,L-lactide-co-glycolide) (PLG) microparticles, Quil A, interleukins, and the like. For experimental animals, one can use Freund's adjuvant (incomplete Freund's adjuvant; complete Freund's adjuvant), N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-nor-muramyl-L-alanyl-D-isoglutamine (CGP 11637, referred to as nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dip- almitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (CGP 19835A, referred to as MTP-PE), and RIBI, which contains three components extracted from bacteria, monophosphoryl lipid A, trehalose dimycolate and cell wall skeleton (MPL+TDM+CWS) in a 2% squalene/Tween 80 emulsion. The effectiveness of an adjuvant may be determined by measuring the amount of antibodies directed against the immunogenic antigen or antigenic epitope thereof.

[00198] Further exemplary adjuvants to enhance effectiveness of the composition include, but are not limited to: (1) oil-in-water emulsion formulations (with or without other specific immunostimulating agents such as muramyl peptides (see below) or bacterial cell wall components), such as for example (a) MF59 (WO 90/14837; Chapter 10 in Vaccine design: the subunit and adjuvant approach, eds. Powell & Newman, Plenum Press 1995), containing 5%

Squalene, 0.5% Tween 80, and 0.5% Span 85 (optionally containing MTP-PE) formulated into submicron particles using a microfluidizer, (b) SAF, containing 10% Squalane, 0.4% Tween 80, 5% pluronic-blocked polymer L121, and thr-MDP either microfluidized into a submicron emulsion or vortexed to generate a larger particle size emulsion, and (c) RIBI adjuvant system (RAS), (Ribi Immunochem, Hamilton, Mont.) containing 2% Squalene, 0.2% Tween 80, and one or more bacterial cell wall components such as monophosphorylipid A (MPL), trehalose dimycolate (TDM), and cell wall skeleton (CWS), e.g., MPL+CWS (Detox TM); (2) saponin adjuvants, such as QS21 or Stimulon™ (Cambridge Bioscience, Worcester, Mass.) may be used or particles generated therefrom such as ISCOMs (immunostimulating complexes), which ISCOMS may be devoid of additional detergent e.g. WO 00/07621; (3) Complete Freund's Adjuvant (CFA) or Incomplete Freund's Adjuvant (IFA); (4) cytokines, such as interleukins (e.g. IL-1, IL-2, IL-4, IL-5, IL-6, IL-7, IL-12 (WO99/44636), etc.), interferons (e.g. gamma interferon), macrophage colony stimulating factor (M-CSF), tumor necrosis factor (TNF), etc.; (5) monophosphoryl lipid A (MPL) or 3-O-deacylated MPL (3dMPL) e.g. GB-2220221, EP-A-0689454, optionally in the substantial absence of alum when used with pneumococcal saccharides e.g. WO 00/56358; (6) combinations of 3dMPL with, for example, QS21 and/or oil-in-water emulsions e.g. EP-A-0835318, EP-A-0735898, EP-A-0761231; (7) oligonucleotides comprising CpG motifs (see, e.g., WO 98/52581), e.g., an oligonucleotide containing at least one CG dinucleotide, where the cytosine is unmethylated; (8) a polyoxyethylene ether or a polyoxyethylene ester (see, e.g. WO 99/52549); (9) a polyoxyethylene sorbitan ester surfactant in combination with an octoxynol (WO 01/21207) or a polyoxyethylene alkyl ether or ester surfactant in combination with at least one additional non-ionic surfactant such as an octoxynol (WO 01/21152); (10) a saponin and an immunostimulatory oligonucleotide (e.g. a CpG oligonucleotide) (WO 00/62800); (11) an immunostimulant and a particle of metal salt e.g. WO 00/23105; (12) a saponin and an oil-in-water emulsion e.g. WO 99/11241; (13) a saponin (e.g. QS21)+3dMPL+IM2 (optionally+a sterol) e.g. WO 98/57659; (14) other substances that act as immunostimulating agents to enhance the efficacy of the composition. Muramyl peptides include N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-25 acetyl-normuramyl-L-alanyl-D-isoglutamine (nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutarninyl-L-alanine-2-(1'-2'-dipalmitoyl-- sn-glycero-3-hydroxyphosphoryloxy)-ethylamine MTP-PE), etc. Adjuvants suitable for administration to a human are of particular interest. In some cases, the adjuvant is an aluminum salt adjuvant (e.g., aluminum phosphate or aluminum hydroxide).

[00199] The antigen compositions may contain other components, such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, magnesium, carbonate, and the like. The compositions may contain pharmaceutically

acceptable auxiliary substances as required to approximate physiological conditions such as pH adjusting and buffering agents, toxicity adjusting agents and the like, for example, sodium acetate, sodium chloride, potassium chloride, calcium chloride, sodium lactate and the like.

[00200] The concentration of the subject fHbp in a formulation can vary widely (e.g., from less than about 0.1%, e.g., at or at least about 2% to as much as 20% to 50% or more by weight) and will usually be selected primarily based on fluid volumes, viscosities, and patient-based factors in accordance with the particular mode of administration selected and the patient's needs.

[00201] The fHbp-containing formulations can be provided in the form of a solution, suspension, tablet, pill, capsule, powder, gel, cream, lotion, ointment, aerosol or the like. It is recognized that oral administration can require protection of the compositions from digestion. This is typically accomplished either by association of the composition with an agent that renders it resistant to acidic and enzymatic hydrolysis or by packaging the composition in an appropriately resistant carrier. Means of protecting from digestion are well known in the art.

[00202] The fHbp-containing formulations can also be provided so as to enhance serum half-life of fHbp following administration. For example, where isolated fHbps are formulated for injection, the fHbp may be provided in a liposome formulation, prepared as a colloid, or other conventional techniques for extending serum half-life. A variety of methods are available for preparing liposomes, as described in, e.g., Szoka et al., Ann. Rev. Biophys. Bioeng. 9:467 (1980), U.S. Pat. Nos. 4,235,871, 4,501,728 and 4,837,028. The preparations may also be provided in controlled release or slow-release forms.

METHODS OF INDUCING AN IMMUNE RESPONSE

[00203] The present disclosure provides a method of inducing an immune response to at least one Neisserial strain in a mammalian host. The methods generally involve administering to an individual in need thereof an effective amount of a subject immunogenic composition.

[00204] The fHbp-containing antigenic compositions are generally administered to a human subject that is at risk of acquiring a Neisserial disease so as to prevent or at least partially arrest the development of disease and its complications. An amount adequate to accomplish this is defined as a "therapeutically effective dose." Amounts effective for therapeutic use will depend on, e.g., the antigenic composition, the manner of administration, the weight and general state of health of the patient, and the judgment of the prescribing physician. Single or multiple doses of the antigenic compositions may be administered depending on the dosage and frequency required and tolerated by the patient, and route of administration.

[00205] The fHbp-containing antigenic compositions are generally administered in an amount effective to elicit an immune response, particularly a humoral immune response, e.g., a bactericidal antibody response, in the host. As noted above, amounts for immunization will vary, and can generally range from about 1 μ g to 100 μ g per 70 kg patient, usually 5 μ g to 50 μ g/70 kg. Substantially higher dosages (e.g. 10 mg to 100 mg or more) may be suitable in oral, nasal, or topical administration routes. The initial administration can be followed by booster immunization of the same or different fHbp-containing antigenic composition. Vaccination in some cases involves at least one booster, and in some cases two boosters.

[00206] In general immunization can be accomplished by administration by any suitable route, including administration of the composition orally, nasally, nasopharyngeally, parenterally, enterically, gastrically, topically, transdermally, subcutaneously, intramuscularly, in tablet, solid, powdered, liquid, aerosol form, locally or systemically, with or without added excipients. Actual methods for preparing parenterally administrable compositions will be known or apparent to those skilled in the art and are described in more detail in such publications as Remington's *Pharmaceutical Science*, 15th ed., Mack Publishing Company, Easton, Pa. (1980).

[00207] An anti-fHbp immune response can be assessed by known methods (e.g. by obtaining serum from the individual before and after the initial immunization, and demonstrating a change in the individual's immune status, for example an immunoprecipitation assay, an ELISA, or a bactericidal assay, a Western blot assay, or flow cytometric assay, or the like).

[00208] Whether a variant fHbp of the present disclosure elicits a bactericidal response to one or more strains of *N. meningitidis* in a mammalian host can be determined using any well-known assay. For example, a human fH transgenic mouse can be used, where the mouse expresses human fH (e.g., human fH is present in serum of the mouse at a concentration of about 100 μ g/ml or greater than 100 μ g/ml). A variant fHbp of the present disclosure is administered to the human fH transgenic mouse. After a period of time, serum from the mouse is tested for bactericidal activity against one or more strains of *N. meningitidis*. Suitable controls include, e.g., fHbp ID 1. An example of a suitable assay is described in Vu et al. (2012) *Sci. Reports* 2:341.

[00209] The antigenic compositions can be administered to a mammalian host (e.g., a human subject) that is immunologically naive with respect to *Neisseria meningitidis*. In a particular embodiment, the subject is a human child about five years or younger, and preferably about two years old or younger, and the antigenic compositions are administered at any one or more of the following times: two weeks, one month, 2, 3, 4, 5, 6, 7, 8, 9, 10, or 11 months, or one year or 15, 18, or 21 months after birth, or at 2, 3, 4, or 5 years of age.

[00210] It may be generally desirable to initiate immunization prior to the first sign of disease symptoms, or at the first sign of possible or actual exposure to infection or disease (e.g., due to exposure or infection by *Neisseria*).

EXAMPLES

[00211] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the present invention, and are not intended to limit the scope of what the inventors regard as their invention nor are they intended to represent that the experiments below are all or the only experiments performed. Efforts have been made to ensure accuracy with respect to numbers used (e.g. amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is weight average molecular weight, temperature is in degrees Celsius, and pressure is at or near atmospheric. Standard abbreviations may be used, e.g., bp, base pair(s); kb, kilobase(s); pl, picoliter(s); s or sec, second(s); min, minute(s); h or hr, hour(s); aa, amino acid(s); kb, kilobase(s); bp, base pair(s); nt, nucleotide(s); i.m., intramuscular(ly); i.p., intraperitoneal(ly); s.c., subcutaneous(ly); and the like.

Example 1: Identification and characterization of fHbp ID 1 mutants

Materials and methods

Library screening

[00212] A random mutant fHbp library was generated by error-prone polymerase chain reaction (PCR), which was followed by cloning of the PCR products into a pET28 expression plasmid that included a signal sequence to allow surface display on *E. coli*. Fluorescence-activated cell sorting was used to isolate mutant clones with low binding of human fH and high binding of a control anti-fHbp monoclonal antibody, which ensured sufficient expression and proper folding of the fHbp mutants. The collected cells were plated on agar plates (LB agar containing 50 µg/ml kanamycin sulfate), which were incubated overnight at 37 °C. Single *E. coli* colonies were used as templates for PCR amplification and the DNA amplicons were purified (PCR Purification Kit; Qiagen™) and subjected to DNA sequencing of the fHbp gene using primers that annealed to the T7 promotor and T7 terminator. The approach of screening a random mutant fHbp library has the potential to identify: 1) positions that affect fH binding that are not predictable from the crystal structure alone; and 2) substitutions other than alanine that affect binding of fH in cases where an alanine substitution would not result in a sufficient decrease in fH binding.

Selection of mutants for further study

[00213] The positions of amino acid substitutions that were identified from the FACS experiment were examined in the crystal structure of fHbp in a complex with a fragment of human fH. Mutants that were in proximity (<5 Å) to the fH binding interface were chosen for site-specific mutagenesis. Recapitulation of the library mutants by site-specific mutagenesis was necessary to create soluble, recombinant fHbps for further characterization. This approach also removed non-desired secondary mutations, which were present in many of the sorted clones and which were distant from the fH binding site. The site-specific mutants were constructed with the Phusion Site-Directed Mutagenesis Kit (Thermo Scientific, Inc.).

Expression and purification of soluble, mutant fHbps

[00214] Soluble, recombinant fHbps were expressed in *E. coli* and lysates were prepared as previously described. fHbps were purified by nickel-affinity chromatography using HiTrap™ Chelating HP columns (5 ml; GE Life Sciences, Inc.) and an Akta Purifier chromatography system (GE Life Sciences). Buffers for binding and elution using an imidazole gradient were prepared according to the column manufacturer's protocols. Fractions containing purified fHbp were combined, dialyzed against PBS containing 3% sucrose and stored at -80 °C prior to use.

[00215] For mouse immunogenicity studies, a second purification step was performed using ion exchange chromatography with HiTrap™ SP HP columns (5 ml; GE Life Sciences). The binding and elution buffers were 25 mM MES, pH 5.5, containing 150 mM and 750 mM NaCl, respectively. Bound fHbp was eluted from the SP column with a linear gradient formed by the binding and elution buffers. Fractions containing purified fHbp were combined, dialyzed against PBS containing 3% sucrose and stored at -80 °C prior to use.

Purification of human Factor H (fH)

[00216] Human fH was purified on a fHbp affinity column. The column was prepared by coupling 5 mg of fHbp ID 1 to an NHS-activated HP column (5 ml; GE Life Sciences) using the manufacturer's protocols. Human serum from a healthy individual donor was diluted 1:1 in phosphate-buffered saline (PBS). The serum was applied to the column and the column was washed with 10 volumes of PBS (i.e. 50 ml). The bound fH was eluted with 5 column volumes of 0.1 M glycine-HCl, pH 2.7. The elution fractions were collected in tubes containing 50 µl of 1 M TRIS-HCl, pH 9.0. Fractions containing fH were identified by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) (4-12% NuPAGE; Invitrogen). Electrophoresis was performed at 200 V for 45 min using 1x MES Running Buffer (Invitrogen). The proteins were visualized by staining with Coomassie G-250 (SimplyBlue SafeStain; Invitrogen). Fractions containing fH were pooled and dialyzed against PBS and aliquots of fH were stored at -30 °C prior to use.

Characterization of fHbp mutants

[00217] SDS-PAGE. The size and purity of the purified fHbp mutant proteins was assessed by SDS-PAGE using 4-12% polyacrylamide gradient gels (NuPAGE; Invitrogen, Inc.). Two μ g of each protein was loaded on the gel. SDS-PAGE was performed as described above for fH.

[00218] Binding of fH to fHbp by enzyme-linked immunosorbent assay (ELISA). The wells of a 96-well microtiter plate (Immulon 2HB; Thermo Scientific) were coated with 2 μ g/ml of purified, recombinant wild-type fHbp (positive control) or mutant fHbp (experimental). Non-specific binding to the wells was blocked with PBS containing 1% BSA (Lifeblood Medical, Inc.) or 5% non-fat dry milk (Carnation; Nestle, Inc.). Five-fold serial dilutions of purified human fH ranging from 25 to 0.0016 μ g/ml in Dilution Buffer (PBS containing 0.1% Tween-20, 0.01% sodium azide and 1% BSA), were added to the wells and the plate was incubated at room temperature for 2 h. After washing three times with PBS containing 0.1% Tween-20 (Sigma) and 0.01% sodium azide (Sigma), bound fH was detected with sheep anti-human fH (1:7,000; Abcam, Inc.) in Dilution Buffer. The plate was incubated at room temperature for 1 h. After washing the wells again, bound primary antibody was detected with donkey anti-sheep IgG conjugated to alkaline phosphatase (1:5,000; Sigma-Aldrich, Inc.) in Dilution Buffer. The plate was incubated at room temperature for 1 h and the wells were washed again. The ELISA was developed with phosphatase substrate (1 mg/ml para-nitrophenyl phosphate; Sigma) in Substrate Buffer (50 mM sodium carbonate, 1 mM MgCl₂, pH 9.8). After incubation at room temperature for 30 min, the absorbance at 405 nm was measured in a UV-VIS plate reader (Spectromax 190; Molecular Devices, Inc.).

[00219] Binding of anti-fHbp monoclonal antibodies to fHbp by ELISA. The wells of a microtiter plate were coated with fHbp, blocked and washed as described above for the fH ELISA. Five-fold serial dilutions of murine anti-fHbp monoclonal antibodies (mAbs) from 25 to 0.0016 μ g/ml in Dilution Buffer were added and the plate was incubated at room temperature for 1 h. After washing the wells, primary antibody was detected with goat anti-mouse IgG conjugated to alkaline phosphatase (1:5,000; Sigma-Aldrich). The ELISA was developed and read as described above.

[00220] Binding of fH to fHbp by surface plasmon resonance (SPR). SPR experiments were performed on a Biacore X100 Plus instrument (GE Life Sciences). Three thousand response units of purified human fH were coupled to a CM5 chip (GE Life Sciences) using the Amine Coupling Kit (GE Life Sciences). fH was immobilized in flow cell 2 and a blank immobilization (no fH) was performed in flow cell 1 as a reference. Three startup cycles consisting of HEPES-buffered saline containing 3 mM EDTA and 0.05% Surfactant P-20 (GE Life Sciences) and regeneration with 100 mM Glycine, 3 M NaCl, pH 2.0, were performed to condition the chip

surface. Dilutions of purified, recombinant fHbp ranging from 100 to 1 nM (wild-type) or 316 to 3.16 nM were injected for 150 seconds. Dissociation was monitored for 300 sec. and the data were analyzed with Biacore X100 Evaluation software.

[00221] Mouse immunogenicity. Groups of wild-type CD-1 mice (N=14 to 21) were immunized with fHbp vaccines adsorbed with aluminum hydroxide. Each dose of vaccine contained 10 µg of fHbp and 600 µg of Alhydrogel (Brenntag Biosector) in 10 mM Histidine, 150 mM NaCl, pH 6.5. Two doses were given three-weeks apart and blood was collected by cardiac puncture three weeks after the second dose. Blood was processed to obtain serum, which was kept at -80 °C for long-term storage (>2 weeks) or 4 °C for short-term storage (< 2 weeks).

[00222] Human fH transgenic BALB/c mice first were screened to identify animals with serum human fH concentrations >240 µg/ml using a fHbp ELISA and a standard curve of purified human fH. The ELISA was performed using purified fHbp ID 1 immobilized on the plate, and the primary and secondary antibodies to detect fH were the same as described above (see “Binding of fH to fHbp by ELISA”).

[00223] Groups of transgenic mice (N=11 to 21) were immunized with fHbp vaccines adsorbed with aluminum hydroxide (same amount of antigen and adjuvant as for wild-type CD-1 mice described above). Three doses were administered at three-week intervals and blood was collected three weeks after the third dose of vaccine. Serum was processed and stored as described above.

[00224] Serum bactericidal antibody (SBA) responses. Human complement-mediated SBA responses were measured against meningococcal strains with an identical or closely matched fHbp sequence compared with the respective vaccine antigen. The bacteria were grown in regular Frantz medium (Frasch et al. “Outer membrane protein vesicle vaccines for meningococcal disease.” In *Methods in Molecular Medicine*, v. 66. *Meningococcal Vaccines: Methods and Protocols*. Edited by Pollard, A.J. and Maiden, M.C. Humana Press Inc. Totowa, NJ) containing 4 mM lactate and 0.02 mM CMP-NANA to mid-exponential phase (OD620 nm =0.6). The bacteria were diluted 1:25,000 in Dulbecco’s PBS containing 1% BSA (Equitech Bio.). Human complement was from a donor with no intrinsic bactericidal antibodies and was depleted of IgG antibodies using a HiTrap Protein G column (5 ml; GE Life Sciences). Each reaction contained 25% human complement, ~400 cfu of bacteria and dilutions of test antisera or control antibodies. The SBA titer was calculated as the serum dilution that resulted in a 50% decrease in cfu relative to negative control wells after 60 min incubation at 37 °C. Protein purification: Recombinant fHbps were expressed in *E. coli* with a C-terminal hexa-histidine tag and purified by metal chelate chromatography (HiTrap Chelating HP; GE Life Sciences) followed by ion exchange chromatography (HiTrap SP; GE Life Sciences). The proteins (2 µg

each) were separated on a 4-12% NuPAGE gel (Invitrogen) using MES running buffer (Invitrogen), and visualized with Coomassie blue staining (Simply Blue Safe Stain; Invitrogen).

Results

[00225] A random mutant library-based approach was developed to identify fHbp mutants with decreased binding of human fH. This approach was able to identify mutations leading to decreased binding of fH that might not be predictable based on structural information alone, and was able to generate multiple amino acid substitutions at any given position. This approach contrasts with the common approach of substitution of alanine at selected positions, which sometimes results in small decreases in binding of fH.

[00226] Using the random mutant library approach, we identified five promising new fHbp ID 1 mutants. The purified, recombinant mutant fHbp ID 1 antigens Q38R, E92K, R130G, S223R and H248L are shown in Figure. 1.

[00227] **Figure 1.** Purity of fHbp ID 1 mutants. Recombinant fHbps were expressed in *E. coli* with a C-terminal hexa-histidine tag and purified by metal chelate chromatography (HiTrap Chelating HP; GE Life Sciences) followed by ion exchange chromatography (HiTrap SP; GE Life Sciences). The proteins (2 µg each) were separated on a 4-12% NuPAGE gel (Invitrogen) using MES running buffer (Invitrogen), and visualized with Coomassie blue staining (Simply Blue Safe Stain; Invitrogen). Lane 1, Kaleidoscope molecular weight marker (Bio-Rad Laboratories); 2, fHbp ID 1 wild-type; 3, Q38R; 4, E92K; 5, R130G; 6, S223R; 7, H248L.

[00228] These mutants exhibit decreased binding of fH ranging from ~10- fold (R130G) to ~20-fold (Q38R) to ~100-fold (E92K, S223R and H248L) (Figure. 2).

[00229] **Figures 2A and 2B.** fH binding of fHbp ID 1 mutants by ELISA. The wells of a microtiter plate were coated with purified recombinant ID 1 wild-type (WT) or one of six different mutant proteins. Different concentrations of purified human fH were added to the wells. Bound fH was detected with sheep anti-human fH (Abcam) and donkey anti-sheep IgG conjugated to alkaline phosphatase (Sigma). A, Positive control fHbp ID 1 wild-type (WT) protein with high binding of human fH. Negative control fHbp ID 1 R41S mutant with low binding of fH. B, New fHbp ID 1 mutants with decreased binding of fH. The R130G mutant showed moderate binding of fH, Q38R showed low binding and E92K, S223R and H248L showed significantly lower binding than that of R41S. The mean and standard deviation for replicate measurements are shown.

[00230] A similar pattern of decreased binding of fH to the mutant proteins was obtained from surface plasmon experiments, in which the R130G and Q38R mutants show some binding, whereas the other three mutants show no detectable binding (Figures 3A and 3B).

[00231] **Figures 3A-3E.** fH binding of fHbp ID 1 mutants by surface plasmon resonance. 3000 response units of purified human fH were coupled to a CM5 chip (GE Life Sciences) and 316 nM of purified, recombinant fHbp was injected for 150 seconds. For reference, the same data for the ID 1 wild-type (WT) protein are shown in each panel. The same pattern of binding was observed as in the ELISA (Figure 2, above); moderate binding to fH for R130G mutant, low binding for Q38R and very low binding for E92K, S223R and H248L. All experiments employed HBS-EP running buffer and a Biacore X100 Plus surface plasmon resonance instrument. Data were analyzed with Biacore X100 Evaluation software.

[00232] All five of the mutant fHbp ID 1 proteins retained conformational epitopes recognized by anti-fHbp monoclonal antibodies. Concentration-dependent binding of five anti-fHbp monoclonal antibodies to wild-type or mutant fHbps is shown in Figure 4.

[00233] **Figures 4A-4E.** Binding of murine anti-fHbp monoclonal antibodies to fHbp mutant proteins as measured by ELISA. Similar concentration-dependent binding of anti-fHbp monoclonal antibodies indicated that the wild-type and mutant fHbps were present in similar amounts in the wells of the microtiter plate and that the mutant fHbps retained conformational epitopes recognized by five distinct monoclonal antibodies. The secondary antibody was goat anti-mouse IgG conjugated to alkaline phosphatase (Sigma). The mean and standard deviation for duplicate measurements are shown.

[00234] The mutant proteins also retain thermal stability similar to the wild-type fHbp ID 1, except the E92K mutant, which has somewhat decreased stability. Finally, the mutants elicited similar bactericidal antibody responses in wild-type CD-1 mice when tested against serogroup B strain H44/76 (Figures 5A-5B).

[00235] **Figures 5A and 5B.** Bactericidal antibody responses to fHbp ID 1 mutants in mice. Figure 5A, Groups of 12 to 14 wild-type mice were immunized intraperitoneally with two doses of purified recombinant fHbp (10 µg per dose) given at three-week intervals. Serum was obtained three weeks after the second dose. Serum bactericidal activity was measured using IgG-depleted human serum as the complement source and serogroup B strain H44/76 as the test strain. H44/76 expresses fHbp ID 1, which matches the control fHbp ID 1 WT vaccine. Each symbol represents the titer of an individual mouse, and the horizontal bars represent the geometric mean titers. The differences between the WT group and each of the mutant groups were not statistically significantly different ($p>0.4$ by t-test). Figure 5B, Groups of 14 to 15 human fH transgenic mice were immunized intraperitoneally with three doses of purified recombinant fHbp (10 µg per dose) given at three-week intervals. Serum was obtained three weeks after the third dose. Serum bactericidal activity was measured as described above for wild-type mice.

[00236] **Figures 6A and 6B.** Binding of human fH to fHbp ID 1 single and double mutants by ELISA. The experiments were performed as described above for Figures 2A-2B. **Figure 7.** Bactericidal antibody responses to fHbp ID 1 single and double mutants in mice. Groups of 20 wild-type mice were immunized and the serum bactericidal antibody responses were determined as described above for Figure 5A.

Example 2: Characterization of fHbp ID 55 mutants

Materials and methods

[00237] The experiments were performed as described in Example 1.

Results

[00238] Three promising mutants that were identified in fHbp ID 1 also were constructed in fHbp ID 55; these included E92K, S223R and H248L. All three of the fHbp ID 55 mutants had significantly decreased binding of fH (Figure 8A). The mutants had preserved conformational integrity as judged by binding of murine anti-fHbp monoclonal antibody JAR 41 (Figure 8B).

[00239] **Figures 8A and 8B.** fH binding of fHbp ID 55 mutants. A. Binding of fH to immobilized fHbp ID 55 mutants by ELISA. The experiment was performed as described in the legend to Figure 2. The mean and range for two to four replicates are shown. B. Concentration dependent binding of anti-fHbp monoclonal antibody (mAb) JAR 41 indicated that the recombinant fHbps were present in similar amounts in the wells of the microtiter plate and had preserved conformation in the region of the epitope recognized by JAR 41 (N-terminal domain; Vu et al. (2012) *Sci. Reports, supra*). The secondary antibody was goat anti-mouse IgG conjugated to alkaline phosphatase (Sigma).

[00240] **Figure 9.** Bactericidal antibody responses to fHbp ID 55 mutants in wild-type mice. Groups of 12 mice were immunized and the serum bactericidal antibody responses were determined as described above for Figure 5A. Bactericidal activity was measured against a mutant of strain H44/76 that expresses fHbp ID 55.

[00241] **Figure 10A,** Bactericidal antibody responses of human fH transgenic mice to fHbp ID 55 S223R mutant. Groups of 11 to 12 transgenic mice were immunized with three doses of purified recombinant fHbp (12 µg per dose) or one-tenth of a human dose of the licensed Trumenba (Pfizer) vaccine containing a total of 12 µg fHbp. The transgenic mice were immunized and the serum bactericidal antibody responses were determined as described above for Figure 5B. **Figure 10B,** Bactericidal antibody responses to Trumenba in wild-type and human fH transgenic mice in relation to serum human fH concentrations. Serum human fH

concentrations were measured by ELISA as described previously (Beernink et al. (2011) Journal of Immunology 186(6):3606-14).

Example 3: Identification and characterization of fHbp ID 22 mutants

Materials and methods

[00242] The experiments were performed as described in Example 1.

Results

[00243] An independent search for random fHbp mutants with decreased binding of fH was performed using fHbp ID 22, which is in variant group 2 (sub-family A). This screen resulted in six promising new mutants (Figure 17). fH binding to the ID 22 wild-type and previously described D211A mutant by ELISA is shown in Figure 11A. fH binding to the six new mutant ID 22 proteins is shown in Figure 11B.

[00244] **Figures 11A-11D.** fH binding of fHbp ID 22 library mutants. Figure 11A, Positive control fHbp ID 22 wild-type (WT) protein with high binding of human fH. Negative control fHbp ID 22 D211A mutant with low binding of fH. Figure 11B, New fHbp ID 22 mutants with decreased binding of fH. All of the mutants showed low binding and V131D showed very low binding similar to D211A. The experiment was performed as described in the legend to Figure 2. The mean and range of two to four replicates are shown. Figure 11C, fH binding of a subset of fHbp ID 22 mutants at fH concentrations up to 100 µg/ml. Figure 11D, Binding of anti-fHbp monoclonal antibody JAR 4 to a subset of the mutants (same symbols used as in Figure 11C). New mutant K219N retains JAR 4 binding, whereas G220S mutant has decreased binding of JAR 4. All of the mutants had normal binding of another anti-fHbp monoclonal antibody JAR 31 (data not shown). The mean and standard deviation of duplicate measurements are shown.

[00245] Bactericidal antibody responses of wild-type CD-1 mice to the new mutants V131D and K219N, along with the control wild-type ID 22 protein and the previously characterized mutant D211A are shown in Figure 12.

[00246] **Figures 12A-12B.** Bactericidal antibody responses to fHbp ID 22 library mutants in wild-type mice. Library mutants with low binding of human fH were selected for immunization. Groups of 10 to 21 mice were immunized with two doses of purified recombinant fHbp (10 µg per dose) given at three-week intervals. Serum was obtained three weeks after the second dose. Serum bactericidal activity was measured using IgG depleted human serum as the complement source and serogroup B strain CH597 as the test strain. This strain expresses fHbp ID 23, which closely matches the control fHbp ID 22 WT vaccine. Figure 12A, Experiment testing new mutants V131D and K219N. The D211A mutant was used as a control mutant fHbp vaccine that did not decrease immunogenicity. Each symbol represents the titer of an individual mouse, and

the horizontal bars represent the geometric mean titers. Figure 12B, Second experiment testing new mutants D121G, S128T, F129S, and G220S. No significant loss of immunogenicity was observed for the new mutant fHbps, except for a modest loss for V131D.

[00247] **Figure 13.** Bactericidal antibody responses to fHbp ID 22 library mutant K219N in human fH transgenic mice. The control mutant D211A gave higher responses and the K219N mutant gave responses similar to the fHbp ID 22 wild-type (WT) antigen.

[00248] **Figure 14.** Thermal stability of fHbp ID 22 measured by differential scanning microcalorimetry. The fHbp ID 22 wild-type (WT, solid line) undergoes unfolding transitions at 38 (N-terminal) and 81 °C (C-terminal domain). An fHbp ID 22 L130R/G133D double mutant exhibits 19 °C higher thermal stability for the N-terminal domain compared with that of the ID 22 WT.

[00249] **Figures 15A-15B.** fHbp ID 22 triple mutants combining stabilizing substitutions L130R and G133D (double mutant, DM) with library derived mutants to decrease fH binding. A, fH binding to fHbp ID 22 triple mutants. B, Control murine anti-fHbp monoclonal antibody (mAb) JAR 4 binding (same symbols as in panel A. All of the stabilized mutants bind JAR 4 better than the fHbp ID 22 WT.

[00250] **Figure 16.** Bactericidal antibody responses to fHbp ID 22 triple mutants in human fH transgenic mice. The two mutants tested combine stability double mutant (DM) with K219N and G220S, respectively. The triple mutants elicited eight- and 18-fold higher responses than the control ID 22 WT antigen.

[00251] A summary of exemplary fHbp ID 1 and ID 22 mutants described above is presented in the table in Figure. 17.

[00252] While the present invention has been described with reference to the specific embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many modifications may be made to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective, spirit and scope of the present invention. All such modifications are intended to be within the scope of the claims appended hereto.

CLAIMS

What is claimed is:

1. A variant factor H binding protein (fHbp), wherein the variant fHbp comprises the amino acid leucine (L) or isoleucine (I) at position 248, wherein the amino acid position is relative to the numbering of SEQ ID NO:1 (fHbp ID1), wherein the variant fHbp comprises an amino acid sequence having at least 85% amino acid sequence identity to SEQ ID NO:1, wherein the variant fHbp binds human factor H (fH) with an affinity that is 50% or less of the affinity of fHbp ID 1 for human fH, and wherein the variant induces a bactericidal antibody response to at least one strain of *Neisseria meningitidis* in a mammalian host.
2. The variant fHbp of claim 1, wherein the variant fHbp binds human fH with an affinity that is 10% or less of the affinity of fHbp ID 1 for human fH.
3. The variant fHbp of claim 1, wherein the variant fHbp binds human fH with an affinity that is 5% or less of the affinity of fHbp ID 1 for human fH.
4. The variant fHbp of any one of claims 1 to 3, further comprising an amino acid substitution selected from the group consisting of R41S and R41A.
5. The variant fHbp of any one of claims 1 to 4, further comprising the amino acid substitution S223R, S223K, S223H, S223F, S223Y, or S223W.
6. The variant fHbp of any one of claims 1 to 5, further comprising the amino acid substitution G130R.
7. The variant fHbp of any one of claims 1 to 6, wherein the amino acid sequence has at least 90% amino acid sequence identity to SEQ ID NO:1.

8. The variant fHbp of any one of claims 1 to 6, wherein the amino acid sequence has at least 95% amino acid sequence identity to SEQ ID NO:1.

9. A variant factor H binding protein (fHbp), wherein the variant fHbp comprises the amino acid substitution H248L, wherein the amino acid substitutions are relative to fHbp ID 55 (SEQ ID NO:3), wherein numbering of H248 is based on the amino acid sequence of SEQ ID NO:1, wherein the variant comprises an amino acid sequence having at least 90% amino acid sequence identity to SEQ ID NO:3, wherein the variant fHbp binds human factor H (fH) with an affinity that is less than 50% of the affinity of fHbp ID 55 for human fH, and wherein the variant induces a bactericidal antibody response in a mammalian host.

10. The variant fHbp of claim 9, wherein the variant fHbp binds human fH with an affinity that is 25% or less of the affinity of fHbp ID 55 for human fH.

11. The variant fHbp of claim 9, wherein the variant fHbp binds human fH with an affinity that is 10% or less of the affinity of fHbp ID 55 for human fH.

12. The variant fHbp of claim 9, wherein the variant fHbp binds human fH with an affinity that is 5% or less of the affinity of fHbp ID 55 for human fH.

13. The variant fHbp of claim 9, further comprising the amino acid substitution S223R, S223K, S223H, S223F, S223Y, or S223W.

14. The variant fHbp of any one of claims 9 to 13, wherein the amino acid sequence has at least 95% amino acid sequence identity to SEQ ID NO:3.

15. An immunogenic composition comprising:

- a) the variant fHbp according to any one of claims 1 to 14; and
- b) a pharmaceutically acceptable excipient.

16. The immunogenic composition of claim 15, wherein the variant fHbp is in a vesicle preparation prepared from a *Neisseria meningitidis* strain.

17. The immunogenic composition of claim 15 or claim 16, wherein said pharmaceutically acceptable excipient comprises an adjuvant.

18. The immunogenic composition of any one of claims 15 to 17, further comprising Neisserial surface protein A.

19. The immunogenic composition of claim 17, wherein the adjuvant is aluminum phosphate or aluminum hydroxide.

20. A nucleic acid encoding the variant fHbp according to any one of claims 1 to 14.

21. A recombinant expression vector comprising the nucleic acid of claim 20.

22. An *in vitro* host cell comprising the nucleic acid of claim 20, or the recombinant expression vector of claim 21.

23. Use of the immunogenic composition of any one of claims 15 to 19 for eliciting an antibody response to *Neisseria meningitidis* in a mammal.


24. Use of the immunogenic composition of any one of claims 15 to 19 in the preparation of a vaccine for eliciting an antibody response to *Neisseria meningitidis* in a mammal.

25. The use of claim 23 or 24, wherein the mammal is a human.


26. The use of any one of claims 23 to 25, wherein the antibody response is a bactericidal antibody response to one or more strains of *N. meningitidis*.

27. Use of the variant fHbp according to any one of claims 1 to 14 in the preparation of a vaccine for eliciting an antibody response to *Neisseria meningitidis* in a mammal.
28. Use of the variant fHbp according to any one of claims 1 to 14 for eliciting an antibody response to *Neisseria meningitidis* in a mammal.
29. The use of claim 27 or 28, wherein the mammal is a human.
30. The use of any one of claims 27 to 29, wherein the antibody response is a bactericidal antibody response to one or more strains of *N. meningitidis*.
31. The variant fHbp according to any one of claims 1 to 14 for use in eliciting an antibody response to *Neisseria meningitidis* in a mammal.
32. The variant fHbp according to claim 31, wherein the mammal is a human.
33. The variant fHbp according to claim 31 or 32, wherein the antibody response is a bactericidal antibody response to one or more strains of *N. meningitidis*.
34. The immunogenic composition according to any one of claims 15 to 19 for use in eliciting an antibody response to *Neisseria meningitidis* in a mammal.
35. The immunogenic composition according to claim 34, wherein the mammal is a human.
36. The immunogenic composition according to claim 34 or 35, wherein the antibody response is a bactericidal antibody response to one or more strains of *N. meningitidis*.


FIG. 1




FIGS. 2A-2B




FIGS. 3A-3E



FIGS. 4A-4E



FIGS. 5A-5B



FIGS. 6A-6B






FIG. 7




FIGS. 8A-8B




9/24

FIG. 9




10/24

FIGS. 10A-10B



11/24

FIGS. 11A-11D



FIGS. 12A-12B




FIG. 13






FIG. 14




FIGS. 15A-15B



16/24

FIG. 16



17/24

FIG. 17

Residue in ID 1	Position ^a	Residue in library mutant	Residue conserved across variant groups	Alternative residue(s) in other variant groups	Hydrogen bond with FH	Position reported as contact residue ^b	Fold-change for Ala mutant ^{c,d}
Q	38	R	No	D	Yes	Yes ^e	n/d
E	92	K	No	A	No	No	n/d
R	138	G	No	L	Yes	Yes ^c	80.8
S	223	R	No	H	No	No	1.2
H	248	I	No	E	No	No	9.6
Residue in ID 22	Position ^a	Residue in library mutant	Residue conserved across variant groups	Alternative residue(s) in other variant groups ^e	Hydrogen bond with FH	Position reported as contact residue ^b	Fold-change for Ala mutant ^{c,d}
N	115	I	No	Q	Yes	Yes	n/d
D	121	G	No	G	No	No	n/d
S	128	T	No	Q, R	Yes	Yes	>100
V	131	D	No	I	No	No	11.5
K	218	N	Yes	n/a	No	No	n/d
G	223	S	Yes	n/a	No	No	n/d

^a Position is according to mature FHbp 1D1 sequence, with the laipidated cysteine as position 1.^b Position reported as contact residue by Schneider et al. (Nature 2009;458:890-3)^c Position tested as alanine mutant for FH binding by Johnson et al. (PLOS Pathog 2012;8:e1002981)^d n/d, not done^e n/a, not applicable

Fig. 18

Human factor H

MRLIAKIICLMLWAICVAEDCNELEPPRRNTEILTGSWSDQTYPEGTQAIYKCRPGYRSLG 60
 NVIMVCRKGEWVALNPLRKCQKRPGHGPDTFGTFTLTGGNVFEYGVKAVYTCNEGQL 120
 LGEINYRECDTDGWTNDIPICEVVKCLPVTAENGKIVSSAMEPDREYHFGQAVRFVCNS 180
 GYKIEGDEEMHCSDDGFWSEKPKCWEISCKSPDVINGSPIQSQKIIYKENERFQYKCNMG 240
 YEYSERGDAVCTESGWRPLPSCEEKSCDNPYIPNGDYSPLRIKHRTGDEITYQCRNGFYP 300
 ATRGNTAKCTSTGWIAPRCTLKPCDYPDIIKGGLYHENMRRPYFPVAVGKYYSYCDEH 360
 FETPSGSYWDHIHCTQDGWSPAVERKCYFPYLENGYNQNYGRKFVQGKSIDVACHPGY 420
 ALPKAQTTVTCMENGWSPTPRCIRVKTCSSIDIENGFISESQYTYALKEKAKYQCKLG 480
 YVTADGETSGSITCGKDGWSAQPTCIKSCDIPVFMNARTKNDFTWFKLNDTLDYECHDGY 540
 ESNTGTTGSIVCGYNGWSDLPLICYERECELPKIDVHLVDRKKDQYKVGEVLKFSCPG 600
 FTIVGPNSVQCYHFGLSPDLPICKEQVQSCGPPPELLNGNVKEKTEEYGHSEVVEYYCN 660
 PRFLMKGPNNKIQCVDGEWTTLPCVIVEESTCGDIPELEHGWQAQLSSPPYYGDSVEFNCS 720
 ESFTMIGHRSITCIHGVWTQLPQCVAILDKLKKCKSSNLIILEEHLKNKEFDHNSNIRYR 780
 CRGKEGWIHTVCINGRWDPEVNCSMAQIQLCPPPQIPNSHNMTTLNYRDGEKVS VLCQ 840
 ENYLIQECEEITCKDGRWQSIPLCVEKIPCSQPPQIEHTINSSRSSQESYAHGTL SYT 900
 CEGGFRISEENETTCYMGKSSPPQCEGLPCKSPPEISHGVVAHMSDSYQYGEETVYKCF 960
 EGFGIDGPAIAKCLGEKWSHPPSCIKDCLSLPSFENAIPMGEKKDVYKAGEQVTTYCAT 1020
 YYKMDGASNVTICINSRWTGRPTCRDTSCVNPTVQNAYIVSRQMSKYPGERVYQCRSP 1080
 YEMFGDEEVMLNGNWTEPPQCKDSTGKCGPPPQIDNGDITSFPLSVYAPASSVEYQCQN 1140
 LYQLEGNKRTICRNGQWSEPPKCLHPCVIISREIMENYNIALRWTAKQKLYSRTGESVEFV 1200
 CKRGYRLSSRSHTLRTTCWDGKLEYPTCAKR 1231 (SEQ ID NO:4)

Fig. 19

	1	10	20	30	40	50	
ID_1	CS-----	SGGGGVAADIGAGLADALTAPLDHKDKGLQSLTLD		SVRKNEKLKLAAQGAEK			55
ID_22	CS-----	SGGGGVAADIGAGLADALTAPLDHKDKSLQSLTLD	QSVRKNEKLKLAAQGAEK				55
ID_55	CSSGGGGSGGGGV	TADIGTGLADALTAPLDHKDKGLKSLTLEDSISQNGTL	LSAQGAEK				60
	60	70	80	90	100	110	
ID_1	TYGNGDSLNTGKLKNDKVSRFDFIRQIEVDGQLITL		SGEFQVYKQSHSALTAFQTEQIQ				115
ID_22	TYGNGDSLNTGKLKNDKVSRFDFIRQIEVDGQLITL	ESGEFQIYKQDHSAVVALQIEKI					115
ID_55	TYGNGDSLNTGKLKNDKVSRFDFIRQIEVDGQLITL		SGEFQVYKQSHSALTALQTEQEQ				120
	120	130	140	150	160	170	
ID_1	DSEHSGKMKVAKRQF	IGDIAGEHTSFDKLPEGGRATYRGTA	FGSDDAGGKLTYTIDFAAK				175
ID_22	NPDKI	SLINQRFL	SGLGGEHTAFNQLPSG-KAEYHGKA	FSSDDPNGLRHYSIDFTKK			174
ID_55	DPEHSEKMKVAKRRFRIGDIAGEHTSFDKLPKDV	MA	TYRGTA	FGSDDAGGKLTYTIDFAAK			180
	180	190	200	210	220	230	
ID_1	QGNKGKIEHLKSP	ELNV	DAADIKPDGKRH	AVISGSVLYNQAEKGSY	LGIFGGKAQEVA		235
ID_22	QGYGRIEHLKTPE	QNVELASAE	ELKADEKSHAVILGDTRYGEE		TYHLALFGDRAQEIA		234
ID_55	QGHGKIEHLKSP	ELNV	DAVAYIKPDEKH	AVISGSVLYNQDEKGSY	LGIFGEKAQEVA		240
	240	250					
ID_1	GSAEVKTVNGIR	IGLAAKQ	255	(SEQ ID NO:1)			
ID_22	GSATVKIREKVHEIGIA	IGAKQ	254	(SEQ ID NO:2)			
ID_55	GSAEVETANGI	IGLAAKQ	260	(SEQ ID NO:3)			

Fig. 20

fHbp ID1 Q38R variant

CSSGGGGVAADIGAGLADALTAPLDHKDKGLQLSLTLDRSVRKNEKLKLAQGA
 EKTYGNGDSLNTGKLNDKVSRFDFIRQIEVDGQLITLESGEFQVYKQSHSALTA
 FQTEQIQDSEHSGKMVAKRQFRIGDIAGEHTSFDKLPEGGRATYRGTAFGSDDA
 GGKLTYTIDFAAKQGNGKIEHLKSPLENVDLAAADIKPDGKRHAVISGSVLYNQ
 AEKGSYSLGIFGGKAQEVAGSAEVKTVNGIRHIGLAAKQ (SEQ ID NO:5)

Fig. 21

fHbp ID1 E92K variant

CSSGGGGVAADIGAGLADALTAPLDHKDKGLQLSLTLDQSVRKNEKLKLAQGA
 EKTYGNGDSLNTGKLNDKVSRFDFIRQIEVDGQLITLEKSGEFQVYKQSHSALTA
 FQTEQIQDSEHSGKMVAKRQFRIGDIAGEHTSFDKLPEGGRATYRGTAFGSDDA
 GGKLTYTIDFAAKQGNGKIEHLKSPLENVDLAAADIKPDGKRHAVISGSVLYNQ
 AEKGSYSLGIFGGKAQEVAGSAEVKTVNGIRHIGLAAKQ (SEQ ID NO:6)

Fig. 22

fHbp ID1 R130G variant

CSSGGGGVAADIGAGLADALTAPLDHKDKGLQLSLTLDQSVRKNEKLKLAQGA
 EKTYGNGDSLNTGKLNDKVSRFDFIRQIEVDGQLITLESGEFQVYKQSHSALTA
 FQTEQIQDSEHSGKMVAKRQGIGDIAGEHTSFDKLPEGGRATYRGTAFGSDDA
 GGKLTYTIDFAAKQGNGKIEHLKSPLENVDLAAADIKPDGKRHAVISGSVLYNQ
 AEKGSYSLGIFGGKAQEVAGSAEVKTVNGIRHIGLAAKQ (SEQ ID NO:7)

Fig. 23

fHbp ID1 S223R variant

CSSGGGGVAADIGAGLADALTAPLDHKDKGLQLSLTLDQSVRKNEKLKLAQGA
 EKTYGNGDSLNTGKLNDKVSRFDFIRQIEVDGQLITLESGEFQVYKQSHSALTA
 FQTEQIQDSEHSGKMVAKRQFRIGDIAGEHTSFDKLPEGGRATYRGTAFGSDDA
 GGKLTYTIDFAAKQGNGKIEHLKSPLENVDLAAADIKPDGKRHAVISGSVLYNQ
 AEKGSYRLGIFGGKAQEVAGSAEVKTVNGIRHIGLAAKQ (SEQ ID NO:8)

Fig. 24

fHbp ID1 H248L variant

CSSGGGGVAADIGAGLADALTAPLDHKDKGLQLSLTLDQSVRKNEKLKLAQGA
 EKTYGNGDSLNTGKLNDKVSRFDFIRQIEVDGQLITLESGEFQVYKQSHSALTA
 FQTEQIQDSEHSGKMVAKRQFRIGDIAGEHTSFDKLPEGGRATYRGTAFGSDDA
 GGKLTYTIDFAAKQGNGKIEHLKSPLENVDLAAADIKPDGKRHAVISGSVLYNQ
 AEKGSYSLGIFGGKAQEVAGSAEVKTVNGIRLIGLAAKQ (SEQ ID NO:9)

Fig. 25

fHbp ID22 N115I variant

CSSGGGGVAADIGAGLADALTAPLDHKDKSLQSLTLDQSVRKNEKLKLAQQGA
 EKTYGNGDSLNTGKLNDKVSRFDFIRQIEVDGQLITLESGEFQIYKQDHSAVVA
 LQIEKIJNPDKIDS LINQRSLVSGLGGEHTAFNQLPSGKAELYHGKAFSSDDPNGR
 LHYSIDFTKKQGYGRIEHLKTPEQNVELASAEKADEKSHAVILGDTRYGEEKG
 TYHLALFGDRAQEIAGSATVKIREKVHEIGIAGKQ (SEQ ID NO:10)

Fig. 26

fHbp ID22 D121G variant

CSSGGGGVAADIGAGLADALTAPLDHKDKSLQSLTLDQSVRKNEKLKLAQQGA
 EKTYGNGDSLNTGKLNDKVSRFDFIRQIEVDGQLITLESGEFQIYKQDHSAVVA
 LQIEKINNPDKIGSLINQRSLVSGLGGEHTAFNQLPSGKAELYHGKAFSSDDPNGR
 LHYSIDFTKKQGYGRIEHLKTPEQNVELASAEKADEKSHAVILGDTRYGEEKG
 TYHLALFGDRAQEIAGSATVKIREKVHEIGIAGKQ (SEQ ID NO:11)

Fig. 27

fHbp ID22 S128T variant

CSSGGGGVAADIGAGLADALTAPLDHKDKSLQSLTLDQSVRKNEKLKLAQQGA
 EKTYGNGDSLNTGKLNDKVSRFDFIRQIEVDGQLITLESGEFQIYKQDHSAVVA
 LQIEKINNPDKIDS LINQRFLVSGLGGEHTAFNQLPSGKAELYHGKAFSSDDPNGR
 LHYSIDFTKKQGYGRIEHLKTPEQNVELASAEKADEKSHAVILGDTRYGEEKG
 TYHLALFGDRAQEIAGSATVKIREKVHEIGIAGKQ (SEQ ID NO:12)

Fig. 28

fHbp ID22 V131D variant

CSSGGGGVAADIGAGLADALTAPLDHKDKSLQSLTLDQSVRKNEKLKLAQQGA
 EKTYGNGDSLNTGKLNDKVSRFDFIRQIEVDGQLITLESGEFQIYKQDHSAVVA
 LQIEKINNPDKIDS LINQRSLDSGLGGEHTAFNQLPSGKAELYHGKAFSSDDPNGR
 LHYSIDFTKKQGYGRIEHLKTPEQNVELASAEKADEKSHAVILGDTRYGEEKG
 TYHLALFGDRAQEIAGSATVKIREKVHEIGIAGKQ (SEQ ID NO:13)

Fig. 29

fHbp ID22 K219N variant

CSSGGGGVAADIGAGLADALTAPLDHKDKSLQSLTLDQSVRKNEKLKLAQQGA
 EKTYGNGDSLNTGKLNDKVSRFDFIRQIEVDGQLITLESGEFQIYKQDHSAVVA
 LQIEKINNPDKIDS LINQRSLVSGLGGEHTAFNQLPSGKAELYHGKAFSSDDPNGR
 LHYSIDFTKKQGYGRIEHLKTPEQNVELASAEKADEKSHAVILGDTRYGEENG
 TYHLALFGDRAQEIAGSATVKIREKVHEIGIAGKQ (SEQ ID NO:14)

Fig. 30

fHbp ID22 G220S variant

CSSGGGGVAADIGAGLADALTAPLDHKDKGLQLSLTLDQSVRKNEKLKLAAGA
 EKTYGNGDSLNTGKLKNDKVSRFDFIRQIEVDGQLITLESGEFQIYKQDHSAVVA
 LQIEKINNPDKIDS LINQRSLVSGLGGEHTAFNQLPSGKAEYHGKAFSSDDPGR
 LHYSIDFTKKQGYGRIEHLKTPEQNVELASAEKADEKSHAVILGTRYGEEKS
 TYHLALFGDRAQEAGSATVKIREKVHEIGIAGKQ (SEQ ID NO:15)

Fig. 31

fHbp ID55 E92K variant

CSSGGGSGGGVTADIGTGLADALTAPLDHKDKGLQLSLTLEDISQNGTLTLSA
 QGAEKTYGNGDSLNTGKLKNDKVSRFDFIRQIEVDGQLITLEKSGEFQVYKQSHS
 ALTALQTEQECDPEHSEKMVAKRRFRIGDIAGEHTSFDKLPKDVMATYRGTAFG
 SDDAGGKLTYTIDFAAKQGHGKIEHLKSPELNVDLAVAYIKPDEKHHAVISGSVL
 YNQDEKGSYSLGIFGEKAQEVASEVETANGIHHIGLAAKQ (SEQ ID NO:16)

Fig. 32

fHbp ID55 S223R variant

CSSGGGSGGGVTADIGTGLADALTAPLDHKDKGLQLSLTLEDISQNGTLTLSA
 QGAEKTYGNGDSLNTGKLKNDKVSRFDFIRQIEVDGQLITLESGEFQVYKQSHSA
 LTALQTEQECDPEHSEKMVAKRRFRIGDIAGEHTSFDKLPKDVMATYRGTAFGS
 DDAGGKLTYTIDFAAKQGHGKIEHLKSPELNVDLAVAYIKPDEKHHAVISGSVL
 YNQDEKGSYRLGIFGEKAQEVASEVETANGIHHIGLAAKQ (SEQ ID NO:17)

Fig. 33

fHbp ID55 H248L variant

CSSGGGSGGGVTADIGTGLADALTAPLDHKDKGLQLSLTLEDISQNGTLTLSA
 QGAEKTYGNGDSLNTGKLKNDKVSRFDFIRQIEVDGQLITLESGEFQVYKQSHSA
 LTALQTEQECDPEHSEKMVAKRRFRIGDIAGEHTSFDKLPKDVMATYRGTAFGS
 DDAGGKLTYTIDFAAKQGHGKIEHLKSPELNVDLAVAYIKPDEKHHAVISGSVL
 YNQDEKGSYSLGIFGEKAQEVASEVETANGIHLIGLAAKQ (SEQ ID NO:18)

Fig. 34

fHbp ID1 R41S/S223R variant

CSSGGGGVAADIGAGLADALTAPLDHKDKGLQLSLTLDQSVSKNEKLKLAAGA
 EKTYGNGDSLNTGKLKNDKVSRFDFIRQIEVDGQLITLESGEFQVYKQSHSALTA
 FQTEQIQDSEHSGKMVAKRQFRIGDIAGEHTSFDKLPEGGRATYRGTAFGSDDA
 GGKLTYTIDFAAKQGNGKIEHLKSPELNVDLAAADIKPDGKRHAVISGSVLYNQ
 AEKGSYRLGIFGGKAQEVASEVKTVNGIRHIGLAAKQ (SEQ ID NO:19)

Fig. 35

fHbp ID1 R41S/H248L variant

CSSGGGGVAADIGAGLADALTAPLDHKDKGLQLSLTLDQSVSKNEKLKLAQQGA
 EKTYGNGDSLNTGKLNDKVSRFDFIRQIEVDGQLITLESGEFQVYKQSHSALTA
 FQTEQIQDSEHSGKMVAKRQFRIGDIAGEHTSFDKLPEGGRATYRGTAFGSDDA
 GGKLTYTIDFAAKQGNGKIEHLKSPELNVDLAAADIKPDGKRHAVISGSVLYNQ
 AEKGSYSLGIFGGKAQEVAGSAEVKTVNGIRLIGLAAKQ (SEQ ID NO:20)

Fig. 36

fHbp ID 1 S223R/H248L variant

CSSGGGGVAADIGAGLADALTAPLDHKDKGLQLSLTLDQSVRKNEKLKLAQQGA
 EKTYGNGDSLNTGKLNDKVSRFDFIRQIEVDGQLITLESGEFQVYKQSHSALTA
 FQTEQIQDSEHSGKMVAKRQFRIGDIAGEHTSFDKLPEGGRATYRGTAFGSDDA
 GGKLTYTIDFAAKQGNGKIEHLKSPELNVDLAAADIKPDGKRHAVISGSVLYNQ
 AEKGSYRLGIFGGKAQEVAGSAEVKTVNGIRLIGLAAKQ (SEQ ID NO:21)

Fig. 37

fHbp ID22 L130R/G133D (stability double mutant “DM”) variant

CSSGGGGVAADIGAGLADALTAPLDHKDKSLQLSLTLDQSVRKNEKLKLAQQGA
 EKTYGNGDSLNTGKLNDKVSRFDFIRQIEVDGQLITLESGEFQIYKQDHSAVVA
 LQIEKINNPKIDSLNQRSFRVSDLGGEHTAFNQLPSGKAEYHGKAFSSDDPNGR
 LHYSIDFTKKQGYGRIEHLKTPEQNVELASAELKADEKSHAVILGDTRYGEEKG
 TYHLALFGDRAQEIAGSATVKIREKVHEIGIAGKQ (SEQ ID NO:22)

Fig. 38

fHbp ID22 L130R/G133D/K219N (DM-K219N) variant

CSSGGGGVAADIGAGLADALTAPLDHKDKSLQLSLTLDQSVRKNEKLKLAQQGA
 EKTYGNGDSLNTGKLNDKVSRFDFIRQIEVDGQLITLESGEFQIYKQDHSAVVA
 LQIEKINNPKIDSLNQRSFRVSDLGGEHTAFNQLPSGKAEYHGKAFSSDDPNGR
 LHYSIDFTKKQGYGRIEHLKTPEQNVELASAELKADEKSHAVILGDTRYGEENG
 TYHLALFGDRAQEIAGSATVKIREKVHEIGIAGKQ (SEQ ID NO:23)

Fig. 39

fHbp ID22 L130R/G133D/G220S (DM-G220S) variant

CSSGGGGVAADIGAGLADALTAPLDHKDKSLQLSLTLDQSVRKNEKLKLAQQGA
 EKTYGNGDSLNTGKLNDKVSRFDFIRQIEVDGQLITLESGEFQIYKQDHSAVVA
 LQIEKINNPKIDSLNQRSFRVSDLGGEHTAFNQLPSGKAEYHGKAFSSDDPNGR
 LHYSIDFTKKQGYGRIEHLKTPEQNVELASAELKADEKSHAVILGDTRYGEEKS
 TYHLALFGDRAQEIAGSATVKIREKVHEIGIAGKQ (SEQ ID NO:24)

Fig. 40**NspA**

GenBank Accession No. AAD53286

1 mkkalatlia lalpaaalae gasgfyvqad aahakasssl gsakgfspri sagyrindlr
61 favdytrykn ykapstdfkl ysigasaiyd fdtqspvkpy lgarlsinra svdlggdsf
121 sqtsiglgvl tgvtsyavtpn vdldagyrn yigkvntvkn vrsgelsvgv rvkf
(SEQ ID NO:25)

