



(86) Date de dépôt PCT/PCT Filing Date: 2000/12/08

(87) Date publication PCT/PCT Publication Date: 2001/06/28

(45) Date de délivrance/Issue Date: 2011/04/19

(85) Entrée phase nationale/National Entry: 2002/05/30

(86) N° demande PCT/PCT Application No.: EP 2000/012532

(87) N° publication PCT/PCT Publication No.: 2001/046357

(30) Priorité/Priority: 1999/12/22 (EP99310428.0)

(51) Cl.Int./Int.Cl. *C07K 19/00* (2006.01),
C07K 16/00 (2006.01), *C07K 16/12* (2006.01),
C07K 16/26 (2006.01), *C07K 16/44* (2006.01),
C11D 3/38 (2006.01), *C11D 3/386* (2006.01),
C12N 9/24 (2006.01), *C12N 9/42* (2006.01),
D06M 16/00 (2006.01)

(72) Inventeurs/Inventors:
DAVIS, PAUL JAMES, GB;
PARRY, NEIL JAMES, GB

(73) Propriétaire/Owner:
UNILEVER PLC, GB

(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : COMPOSITIONS DETERGENTES COMPRENANT DES AGENTS AMELIORANTS

(54) Title: DETERGENT COMPOSITIONS COMPRISING BENEFIT AGENTS

(57) **Abrégé/Abstract:**

There is provided a fusion protein comprising a cellulose binding domain and a domain having a high binding affinity for another ligand. This high affinity binding domain is preferably directed at a Benefit Agent. The latter is preferably selected from the group consisting of a fabric softening agents, fragrances, perfumes, polymeric lubricants, photoprotective agents, latexes, resins, dye fixative agents, encapsulated materials, antioxidants, insecticides, soil repelling agents or a soil release agents. The fusion protein is advantageously used in a detergent composition to deliver the Benefit Agent to a fabric.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 June 2001 (28.06.2001)

PCT

(10) International Publication Number
WO 01/46357 A2

- (51) International Patent Classification⁷: **C11D** Research Colworth, Colworth House, Sharnbrook, Bedford, Bedfordshire MK44 1LQ (GB).
- (21) International Application Number: PCT/EP00/12532
- (22) International Filing Date: 8 December 2000 (08.12.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
99310428.0 22 December 1999 (22.12.1999) EP
- (71) Applicant (for all designated States except AG, AU, BB, BZ, CA, CY, GB, GD, GH, GM, IE, IL, IN, KE, LK, LS, MN, MW, MZ, NZ, SD, SG, SL, SZ, TT, TZ, UG, ZA, ZW): **UNILEVER N.V.** [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL).
- (71) Applicant (for AG, AU, BB, BZ, CA, CY, GB, GD, GH, GM, IE, IL, KE, LK, LS, MN, MW, MZ, NZ, SD, SG, SL, SZ, TT, TZ, UG, ZA, ZW only): **UNILEVER PLC** [GB/GB]; Unilever House, Blackfriars, London, Greater London EC4P 4BQ (GB).
- (71) Applicant (for IN only): **HINDUSTAN LEVER LTD** [IN/IN]; Hindustan Lever House, 165-166 Backbay Reclamation, 400 020 Mumbai (IN).
- (72) Inventors: **DAVIS, Paul, James**; Unilever Research Colworth, Colworth House, Sharnbrook, Bedford, Bedfordshire MK44 1LQ (GB). **PARRY, Neil, James**; Unilever
- (74) Agent: **KAN, Jacob, Hendrik**; Unilever Research Vlaardingen, Patent Department, Olivier van Noortlaan 120, NL-3133 AT Vlaardingen (NL).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— Without international search report and to be republished upon receipt of that report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DETERGENT COMPOSITIONS COMPRISING BENEFIT AGENTS

(57) Abstract: There is provided a fusion protein comprising a cellulose binding domain and a domain having a high binding affinity for another ligand. This high affinity binding domain is preferably directed at a Benefit Agent. The latter is preferably selected from the group consisting of a fabric softening agents, fragrances, perfumes, polymeric lubricants, photoprotective agents, latexes, resins, dye fixative agents, encapsulated materials, antioxidants, insecticides, soil repelling agents or a soil release agents. The fusion protein is advantageously used in a detergent composition to deliver the Benefit Agent to a fabric.

WO 01/46357 A2

DETERGENT COMPOSITIONS COMPRISING BENEFIT AGENTSTECHNICAL FIELD

The present invention generally relates to detergent
5 compositions comprising benefit agents. More in particular,
it relates to detergent compositions comprising fusion
proteins which are useful for delivering a benefit agent to
a fabric.

10 BACKGROUND AND PRIOR ART

WO-A-98/00500 (Unilever) discloses detergent compositions
wherein a Benefit Agent is delivered onto fabric by means of
peptide or protein Deposition Aid having a high affinity for
fabric. The Benefit Agent can be a fabric softening agent, a
15 perfume, a polymeric lubricant, a photoprotective agent, a
latex, a resin, a dye fixative agent, an encapsulated
material, an antioxidant, an insecticide, a soil repelling
agent or a soil release agent. The Benefit Agent is attached
or adsorbed to a peptide or protein Deposition Aid having a
20 high affinity for fabric. Preferably, the deposition aid is
the cellulose binding domain of a cellulase enzyme. The
compositions are said to effectively deposit the Benefit
Agent onto the fabric during the wash cycle.

25 The detergent compositions disclosed in WO-A-98/00500 have a
number of drawbacks. The Benefit Agent has to be
attached/adsorbed to the Deposition Aid. If it is to be
attached, it is preferably done by means of a linking agent.
However, not all benefit agents can be derivatised.
30 Furthermore, linking the Benefit Agent to the Deposition Aid
may negatively affect the properties of the Benefit Agent.
For example, perfumes cannot be covalently bound to larger
molecules without negatively affecting their perfume
properties.

It is therefore an object of the present invention to provide an alternative or improved detergent composition which is capable of delivering a benefit agent to a fabric during a washing or rinsing process.

5 Surprisingly, we have now found that these and other objects of the invention may be achieved by the detergent compositions of the present invention, which are characterised in that they comprise a fusion protein
10 comprising a cellulose binding domain and a domain having a high binding affinity for another ligand. This fusion molecule is bi-functional in its binding ability, whereby the cellulose binding domain region binds to cellulosic based materials and the second domain binds to another
15 ligand.

According to DE-A-196 21 224 (Henkel), the transfer of textile dyes from one garment to another during a washing or rinsing process may be inhibited by adding antibodies
20 against the textile dye to the wash or rinse liquid.

WO-A-98/07820 (P&G) discloses rinse treatment compositions containing antibodies directed at cellulase and standard softener actives (such as DEQA).

25

DEFINITION OF THE INVENTION

According to a first aspect of the invention there is provided a detergent composition comprising one or more surfactants and a fusion protein comprising a cellulose binding domain and a domain
30 having a high binding affinity for another ligand, wherein the domain having a high binding affinity is an antibody or antibody fragment which is directed at a Benefit Agent selected from the group consisting of fabric softening agents, fragrances, perfumes, polymeric lubricants, photoprotective agents, latexes,
35 resins, dye fixative agents, encapsulated materials, antioxidants, insecticides, soil repelling agents or soil release agents.

According to a second aspect there is provided a process for delivering a benefit agent to a fabric by treating said fabric with a composition comprising a fusion protein as defined above and a benefit agent selected from the group consisting of softening agents, finishing agents/ protective agents, fragrances and bleaching agents.

DETAILED DESCRIPTION OF THE INVENTION

1.1 The cellulose binding domain

In its first aspect, the invention relates to a fusion protein comprising a cellulose binding domain (CBD) and a domain having a high binding affinity for another ligand. The cellulose binding domain is a part of most cellulase enzymes and can be obtained therefrom. Cellulose binding domains are also obtainable from xylanase and other hemicellulase degrading enzymes. Preferably, the cellulose binding domain is obtainable from a fungal enzyme origin such as Humicola, Trichoderma, Thermomonospora, Phanerochaete, Aspergillus or from a bacterial enzyme origin such as Bacillus, Clostridium, Streptomyces, Cellulomonas and Pseudomonas. Especially preferred is the cellulose binding domain which is obtainable from Trichoderma reesei.


In the fusion protein according to the invention, the cellulose binding domain is fused to a second domain having a high binding affinity for another ligand. Preferably, the cellulose binding domain is connected to the domain having a high binding affinity for another ligand by means of a linker consisting of about 0-20, preferably about 2-15, more preferably of 2-5 amino acid residues.

The second domain having a high binding affinity for another ligand may, for instance, be an antibody or an antibody fragment. Especially preferred are heavy chain antibodies such as found in Camelidae.

The fusion protein according to the invention may comprise more than two recognition domains. It is for example possible to produce a CBD fusion protein with more than one antibody domain, in which the antibody domains may bind to the same or bind to different antigens. Conversely, it is also possible in the CBD antibody fusion format to produce a molecule with one antibody domain with more than one CBD, whereby the CBD's incorporated may be identical sequences or from more than one source, or modified varieties thereof.

10

Generally speaking, the degree of binding of a molecule A to another molecule B can be generally expressed by the chemical equilibrium constant K_d resulting from the following reaction:



15 The chemical equilibrium constant K_d is then given by:

$$K_d = \frac{[A] \times [B]}{[A \equiv B]}$$

Whether the binding of a molecule to the fabric / ligand is specific or not can be judged from the difference between the binding (K_d value) of the molecule to one type of fabric / ligand, versus the binding to another type of fabric / ligand material. For applications in laundry, said ligand material will form part of or be associated with a benefit agent. In this aspect of the invention, the CBD region of the fusion protein binds to the fabric and the high affinity domain region binds to the benefit agent. Alternatively, this approach can be reversed whereby the fabric material, or a ligand bound to the fabric is targeted by the high affinity domain of the fusion protein and the cellulose binding domain region binds to a cellulosic based or cellulosic containing benefit agent. However, it will usually be more convenient to measure K_d values and differences in K_d values on other materials such as a

30

5

polystyrene microtitre plate or a specialised surface in an analytical biosensor. The difference between the two binding constants should be minimally 10, preferably more than 100, and more preferably, more than 1000. Typically, the reagent should bind to the ligand / fabric, with a K_d lower than 10^{-4} M, preferably lower than 10^{-6} M and could be 10^{-10} M or even less. Higher binding affinities (K_d of less than 10^{-5} M) and/or a larger difference between the one type of ligand / fabric and another type of ligand / fabric (or background) binding would increase the deposition of the benefit agent. Also, the weight efficiency of the reagent in the total rinse composition would be increased and smaller amounts of the reagent would be required.

Several classes of reagent or molecules can be envisaged which deliver the capability of specific binding to fabrics / ligands, to which one would like to deliver the benefit agent. In the following we will give a number of examples of such molecules having such capabilities, without pretending to be exhaustive.

1.2.1. Antibodies.

Antibodies are specific binding proteins. Their function in nature is to protect against disease by recognising (and binding) foreign bodies, such as viruses or Bacteria, but not self-cells. Furthermore, methods are well-known in the art to generate antibodies that are specific for almost any protein, organic molecule, or cell surface, that is likely to be encountered. This binding specificity has been exploited in the Biotechnology industry, principally for medical diagnostics. For example, many home-based pregnancy test kits comprise an antibody that specifically binds to the pregnancy marker hormone, human chorionic gonadotropin (hCG), but not to other hormones present in urine.

More recently, the use of antibodies in laundry products has been described (Henkel, Procter and Gamble, Unilever). In particular, Unilever has described the use of stain-specific antibodies to target bleaching enzymes exclusively to stains
5 but not to dyes - thus achieving efficient stain removal without damaging surrounding fabric.

Antibodies are well known examples of molecules which are capable of binding specifically to compounds against which
10 they were raised. Antibodies can be derived from several sources. From mice, monoclonal antibodies can be obtained which possess very high binding affinities. From such antibodies, Fab, Fv or scFv fragments, can be prepared which have retained their binding properties. Such antibodies or
15 fragments can be produced through recombinant DNA technology by microbial fermentation. Well known production hosts for antibodies and their fragments are yeast, moulds or bacteria.

20 A class of antibodies of particular interest is formed by the Heavy Chain antibodies as found in Camelidae, like the camel or the llama. The binding domains of these antibodies consist of a single polypeptide fragment, namely the variable region of the heavy chain polypeptide (HC-V). In
25 contrast, in the classic antibodies (murine, human, etc.), the binding domain consists of two polypeptide chains (the variable regions of the heavy chain (V_h) and the light chain (V_l)). Procedures to obtain heavy chain immunoglobulins from Camelidae, or (functionalized) fragments thereof, have been
30 described in WO-A-94/04678 (Casterman and Hamers) and WO-A-94/25591 (Unilever and Free University of Brussels).

Alternatively, binding domains can be obtained from the V_h fragments of classical antibodies by a procedure termed
35 "camelization". Hereby the classical V_h fragment is transformed, by substitution of a number of amino acids,

- into a HC-V-like fragment, whereby its binding properties are retained. This procedure has been described by Riechmann et al. in a number of publications (J. Mol. Biol. (1996) **259**, 957-969; Protein. Eng. (1996) **9**, 531-537, Bio/Technology (1995) **13**, 475-479). Also HC-V fragments can be produced through recombinant DNA technology in a number of microbial hosts (bacterial, yeast, mould), as described in WO-A-94/29457 (Unilever).
- 10 Methods for producing fusion proteins that comprise an enzyme and an antibody or that comprise an enzyme and an antibody fragment are already known in the art. One approach is described by Neuberger and Rabbits (EP-A-194 276). A method for producing a fusion protein comprising an enzyme and an antibody fragment that was derived from an antibody originating in *Camelidae* is described in WO-A-94/25591. A method for producing bispecific antibody fragments is described by Holliger et al. (1993) PNAS **90**, 6444-6448.
- 20 A particularly attractive feature of antibody binding behaviour is their reported ability to bind to a "family" of structurally related molecules. For example, in Gani et al. (J. Steroid Biochem. Molec. Biol. **48**, 277-282) an antibody is described that was raised against progesterone but also binds to the structurally-related steroids, pregnanediolone, pregnanolone and 6-hydroxy-progesterone. Therefore, using the same approach, antibodies could be isolated that bind to a whole "family" of stain chromophores (such as the polyphenols, porphyrins, or carotenoids as described below).
- 30 A broad action antibody such as this could be used to treat several different stains when coupled to a bleaching enzyme.

1.2.2. Peptides.

- Peptides usually have lower binding affinities to the substances of interest than antibodies. Nevertheless, the binding properties of carefully selected or designed

peptides can be sufficient to deliver the desired selectivity in an oxidation process. A peptide which is capable of binding selectively to a fabric / ligand to which one would like to deliver a benefit agent, can for instance
5 be obtained from a protein which is known to bind to that specific fabric / ligand. An example of such a peptide would be a binding region extracted from an antibody raised against that fabric / ligand. A suitable peptide could be analogous to the active center of a protein analogous to a
10 non-catalytic binding domain of a protein, e.g. a receptor.

Alternatively, peptides that bind to such substance can be obtained by the use of peptide combinatorial libraries. Such a library may contain up to 10^{10} peptides, from which the
15 peptide with the desired binding properties can be isolated. (R.A. Houghten, Trends in Genetics, Vol 9, no 6, 235-239). Several embodiments have been described for this procedure (J. Scott et al., Science (1990) **249**, 386-390; Fodor et al., Science (1991) **251**, 767-773; K. Lam et al., Nature (1991)
20 **354**, 82-84; R.A. Houghten et al., Nature (1991) **354**, 84-86).

Suitable peptides can be produced by organic synthesis, using for example the Merrifield procedure (Merrifield (1963) J.Am.Chem.Soc. **85**, 2149-2154). Alternatively, the
25 peptides can be produced by recombinant DNA technology in microbial hosts (yeast, moulds, bacteria) (K.N. Faber et al. (1996) Appl. Microbiol. Biotechnol. **45**, 72-79).

1.2.3. Peptidomimics.

30 In order to improve the stability and/or binding properties of a peptide, the molecule can be modified by the incorporation of non-natural amino acids and/or non-natural chemical linkages between the amino acids. Such molecules are called peptidomimics (H.U. Saragovi et al. (1991)
35 Bio/Technology **10**, 773-778; S. Chen et al. (1992)

Proc.Natl.Acad. Sci. USA **89**, 5872-5876). The production of such compounds is restricted to chemical synthesis.

1.2.4. Other organic molecules.

5 It can be readily envisaged that other molecular structures, which need not be related to proteins, peptides or derivatives thereof, can be found which bind selectively to fabrics / ligands to which one would like to deliver a benefit agent. For example, certain polymeric RNA molecules
10 which have been shown to bind small synthetic dye molecules (A. Ellington et al. (1990) Nature **346**, 818-822). Such binding compounds can be obtained by the combinatorial approach, as described for peptides (L.B. McGown et al. (1995), Analytical Chemistry, 663A-668A).

15

This approach can also be applied for purely organic compounds which are not polymeric. Combinatorial procedures for synthesis and selection for the desired binding properties have been described for such compounds (Weber et
20 al. (1995) Angew.Chem.Int.Ed.Engl. **34**, 2280-2282; G. Lowe (1995), Chemical Society Reviews **24**, 309-317; L.A. Thompson et al. (1996) Chem. Rev. **96**, 550-600). Once suitable binding compounds have been identified, they can be produced on a larger scale by means of organic synthesis.

25

A further embodiment would be for the domain with a high binding affinity to be a bispecific reagent. Such a reagent could fulfil the requirement of accumulating the benefit agent on the fabric either by supplying said reagent
30 together with the benefit agent as a pre-formed non-covalent complex or by supplying the two separately and allowing them to self-assemble either in the wash liquor or on the fabric.

In the detergent composition according to the invention, the
35 fusion protein can be used to deposit a Benefit Agent onto the fabric by means of the cellulose binding domain.

Alternatively, the domain having a high binding affinity for the fabric / ligand can be used to target the benefit agent.

In a further embodiment, the detergent compositions of the invention comprise micro-particles sensitised with CBD-antibody fusions, and configured such that the micro-particles are loaded with the benefit agent and the antibody domain of the fusion protein has a high affinity or specificity for a substance (or "marker molecule") typically found on some regions of fabrics but not on others. Examples of such marker molecules include bleach-damaged dyes and microbes known to be associated with malodour. The antibody domain targets the benefit agent to its intended site of action and binds it there. For example, Microbe-specific antibody based fusions may target fragrance-containing particles to the regions of malodour. Thus, a more efficient use of expensive ingredients is achieved. Alternatively, antibody based fusions specific for bleach-damaged dyes can target dyed particles to faded regions, thus replenishing the colour lost in the main wash cycle.

The CBD antibody fusion protein binds to the fabric via the CBD region, thereby allowing the antibody domain to bind to corresponding antigens / ligands that comprise or form part of the benefit agent. The fusion protein can be dosed in conjunction with the benefit agent or can be added to the fabric prior to the said fabric coming into contact with the benefit agent.

A further aspect of the invention is a process for delivering a benefit agent to a fabric, by treating said fabric with a composition comprising the fusion protein and a benefit agent. The benefit agent can be selected from softening agents, finishing agents/ protective agents, fragrances (perfumes), bleaching agents.

Examples of softening agents are clays, cationic surfactants or silicon compounds. Examples of finishing agents/ protective agents are polymeric lubricants, soil repelling agents, soil release agents, photo-protective agents
5 (sunscreens), anti-static agents, dye-fixing agents, anti-bacterial agents and anti-fungal agents. The fragrances or perfumes may be encapsulated, e.g. in latex microcapsules or gelatine based coacervates.

10 Suitable examples of bleaches are photobleaches. Examples of photobleaches are given in EP-A-379 312 (British Petroleum), which discloses a water-insoluble photobleach derived from anionically substituted porphine, and in EP-A-035 470 (Ciba Geigy), which discloses a textile treatment composition
15 comprising a photobleaching component.

Another advantage of the present invention is that it is possible to target some benefit molecules to particular regions of fabric. For example, dyes can be targeted to
20 colour-bleached regions to replenish dye lost in the main wash or fragrance can be targeted to regions where it is most needed, in particular to those regions where microbes associated with malodour are present, such as the "underarm" regions.

25

1.3 The fabrics

For laundry detergent applications, several classes of natural or man-made fabrics can be envisaged, in particular cotton. In the embodiment of the invention whereby the
30 antibody region of the fusion protein targets the fabric, such macromolecular compounds have the advantage that they can have a more immunogenic nature, i.e. that it is easier to raise antibodies against them. Furthermore, they are more accessible at the surface of the fabric than for instance
35 coloured substances in stains, which generally have a low molecular weight.

An important embodiment of the invention is to use binding domains (as described above) that bind to several different types of fabrics. This would have the advantage of enabling
5 a Benefit Agent to be deposited to several different types of fabric using the CBD-antibody fusion molecule.

2. The Detergent Composition.

The fusion proteins of the invention can be used in a
10 detergent composition which is specifically suited for the purpose, and this constitutes a second aspect of the invention. When formulating a detergent composition, it is important to ensure that the other ingredients of the product are compatible with the activity of the fusion
15 protein. WO-A-98/07820 (P&G) discloses *inter alia* rinse treatment compositions containing antibodies directed at cellulase and standard softener actives such as DEQA. The detergent product according to the present invention preferably contains no softener or low levels of softener
20 active (e.g. HEQ).

To that extent, the detergent composition comprises one or more benefit agents and optionally other conventional detergent ingredients. The invention in its second aspect
25 provides a detergent composition which comprises from 0.1 - 50 % by weight, based on the total composition, of one or more surfactants. This surfactant system may in turn comprise 0 - 95 % by weight of one or more anionic surfactants and 5 - 100 % by weight of one or more nonionic
30 surfactants. The surfactant system may additionally contain amphoteric or zwitterionic detergent compounds, but this is not normally desired owing to their relatively high cost. It may be advantageous to also include cationic surfactants into the composition. Examples of suitable cationic
35 surfactants are given in WO-A-97/03160 and WO-A-98/17767 (Procter&Gamble).

In general, the nonionic and anionic surfactants of the surfactant system may be chosen from the surfactants described "Surface Active Agents" Vol. 1, by Schwartz & Perry, Interscience 1949, Vol. 2 by Schwartz, Perry & Berch, 5 Interscience 1958, in the current edition of "McCutcheon's Emulsifiers and Detergents" published by Manufacturing Confectioners Company or in "Tenside-Taschenbuch", H. Stache, 2nd Edn., Carl Hauser Verlag, 1981.

10 Suitable nonionic detergent compounds which may be used include, in particular, the reaction products of compounds having a hydrophobic group and a reactive hydrogen atom, for example, aliphatic alcohols, acids, amides or alkyl phenols with alkylene oxides, especially ethylene oxide either alone 15 or with propylene oxide. Specific nonionic detergent compounds are C₆-C₂₂ alkyl phenol-ethylene oxide condensates, generally 5 to 25 EO, i.e. 5 to 25 units of ethylene oxide per molecule, and the condensation products of aliphatic C₈-C₁₈ primary or secondary linear or branched alcohols with 20 ethylene oxide, generally 5 to 40 EO.

Suitable anionic detergent compounds which may be used are usually water-soluble alkali metal salts of organic sulphates and sulphonates having alkyl radicals containing 25 from about 8 to about 22 carbon atoms, the term alkyl being used to include the alkyl portion of higher acyl radicals. Examples of suitable synthetic anionic detergent compounds are sodium and potassium alkyl sulphates, especially those obtained by sulphating higher C₈-C₁₈ alcohols, produced for 30 example from tallow or coconut oil, sodium and potassium alkyl C₉-C₂₀ benzene sulphonates, particularly sodium linear secondary alkyl C₁₀-C₁₅ benzene sulphonates; and sodium alkyl glyceryl ether sulphates, especially those ethers of the higher alcohols derived from tallow or coconut oil and 35 synthetic alcohols derived from petroleum. The preferred anionic detergent compounds are sodium C₁₁-C₁₅ alkyl benzene

14

5 sulphonates and sodium C₁₂-C₁₈ alkyl sulphates. Also applicable are surfactants such as those described in EP-A-328 177 (Unilever), which show resistance to salting-out, the alkyl polyglycoside surfactants described in EP-A-070 074, and alkyl monoglycosides.

Preferred surfactant systems are mixtures of anionic with nonionic detergent active materials, in particular the groups and examples of anionic and nonionic surfactants pointed out in EP-A-346 995 (Unilever). Especially preferred is surfactant system which is a mixture of an alkali metal salt of a C₁₆-C₁₈ primary alcohol sulphate together with a C₁₂-C₁₅ primary alcohol 3-7 EO ethoxylate.

15 The nonionic detergent is preferably present in amounts greater than 10%, e.g. 25-90% by weight of the surfactant system. Anionic surfactants can be present for example in amounts in the range from about 5% to about 40% by weight of the surfactant system.

20

The detergent composition may take any suitable physical form, such as a powder, a tablet, an aqueous or non aqueous liquid, a paste or a gel. The fusion protein according to the invention will generally be used as a dilution in water of about 0.05 to 2%.

25

The detergent composition in accordance with the invention comprising the fusion protein can have any suitable form, i.e. the form of a granular composition, a liquid or a slurry of the enzyme, or with carrier material (e.g. as in EP-A-258 068 and the Savinase (TM) and Lipolase (TM) products of Novo Nordisk). A good way of adding the fusion protein to a liquid detergent product is in the form of a slurry containing from 0.005 to 50 % by weight of the complex in an ethoxylated alcohol nonionic surfactant.

30

35

The detergent compositions of the invention comprise about 0.001 to 50 mg, preferably from 0.01 to 10 mg of fusion protein per liter of the rinse liquor in use. A concentrated detergent composition before use will comprise about 1 to 5 1000 mg/l, preferably from 10 mg to 100 mg per liter of the detergent product.

The invention will now be further illustrated in the following, non-limiting Examples, with reference to the 10 following drawing figures:

Figure 1 shows the sequence of the expressed fusion protein (SEQ ID NOS: 1&2) of Example 4;

15 Figure 2 demonstrates the enhanced binding of perfumed latex in the presence of the fusion protein referred to in Example 5; and,

20 Figures 3 and 4 demonstrate the enhanced level of perfume deposition achieved using the novel cross linking fusion protein approach referred to in Example 6.

25

30

35

15a

Example 1CBD-anti *Streptococcus sanguis* fusion (CBD-4715)

The ability of the CBD-fusion to capture bacteria onto a cellulosic matrix was investigated. The following protocol describes the method used to crosslink bacteria to cellulose via a CBD-antibody crosslinker, whereby the CBD region bound to the cellulose and the antibody domain binds to the bacteria: 1% phosphoric acid swollen cellulose was used to which 0-500µl 0.1 mg/ml CBD-4715 was added and volume made to 1ml with PBS pH 7. The samples were incubated for 30 minutes on a end over end shaker at 25°C. After each incubation step the samples were washed. To each tube 100µl 2.56×10^8 *Streptococcus sanguis* cells were added. Resulting samples were mixed for 10 minutes and then the samples were centrifuged and washed in PBS. An aliquot of each solution was removed and 10µl 0.1mg/ml carboxyfluorescein diacetate was added. This is a fluorescent marker for viable bacterial cells. Captured bacteria were visualised by green fluorescence (Excitation 488nm, Emission 525nm). In the presence of the CBD fusion, bacteria were specifically captured onto the cellulosic material. This was in stark contrast to samples in the absence of the crosslinker, whereby bacteria were not associated with the cellulosic.

25

30

Example 2**Latex capture on cellulose fibres (CBD-anti HCG fusion)**

The ability of the CBD-anti HCG fusion to capture latex particles onto cotton yarns was tested: cotton yarns were washed thoroughly in 0.5ml 0.1M sodium carbonate pH 9.0. To these 0-12µl 0.4 mg/ml CBD-anti HCG fusion was added and incubated at 25°C for 10 minutes. The supernatants were removed and the yarns washed in 0.1M sodium carbonate pH 9.0. Yarns were then resuspended in 0.1M sodium carbonate pH 9.0 containing 0.25% Tween™. 0-15µl duke green-GCG latex was added and samples were then incubated at 25°C for 20 minutes. Yarns were then washed with PBS-Tween and the yarns removed and dried. Latex remaining on the yarns was identified by fluorescence.

15

Single cotton yarns were used as the target surface. The incorporation of Tween was essential to remove the non-specific entrapment of latex on cotton. There was a good correlation between CBD fusion dose and degree of particle capture. This correlation is described in the following table:

		µl CBD Fusion			
		0	4µ	8µl	12µl
25	0µl latex	-	-	-	-
	5µl latex	-	+	+	+++
	10µl latex	-	++	++	+++
	15µl latex	-	+++	++++	+++++

Scale:

30 - = No capture of latex onto cotton
 ++ = Capture of latex onto cotton clearly visible
 +++++ = Significant capture of latex onto cotton

Example 3

Capture of coacervate particles containing a model benefit agent onto fabric via a CBD-antibody fusion protein crosslinker.

5 The next step was to investigate the ability to capture coacervate microcapsules onto cotton fabric using a CBD-fusion. The benefit agent in the capsule to be delivered was sunflower oil / β -carotene sample as a model. Coacervate particles allow the incorporation of many hydrophobic and
 10 insoluble molecules. There was clear indication of enhanced deposition of particles to fabric, and the qualitative data produced suggested a dose response with the concentration of fusion protein. The untreated fabric did show entrapment of particles but there was a considerable increase in the
 15 delivery and deposition of capsules in the presence of the fusion protein. This correlation is shown in the following table:

	mg CBD Fusion	0	0.05	0.125
20	Capture	+	++++	+++++

Scale:

+ = Capture of coacervates onto cotton visible

+++++ = Significant capture of coacervates onto cotton.

25

Example 4Production of the fusion protein

To demonstrate the feasibility of the concept we have
 30 combined an existing llama antibody gene sequence (anti-Reactive Red 6) with a CBD sequence to produce a single fusion protein. In this manner, we have put Reactive Red 6 dye onto perfume particles and show the deposition using the fusion proteins. It will be clear that we can generate any
 35 molecule of choice for the antibody region e.g. an anti-gelatine coacervate specificity.

The antibody gene which confers binding to the dye Reactive Red 6 was combined with a CBD gene (taken from *Tricoderma reesei*) with a small peptide linker in between the two sequences. The combination of the two genes was performed via convention molecular biology techniques. To demonstrate that the gene was successfully produced, the sequence of the expressed fusion protein is shown in Figure 1 (SEQ ID NOS: 1&2).

10 The vector containing the selected anti-Reactive Red 6 Vhh-CBD-gene was then used to transfer cells for a larger scale production of the fusion protein. From a 5 litre fermentation broth 60 mg/l fusion protein was isolated.

15 1.1 Perfumed latex particles.

Positively charged latex containing perfume fragrance was derivatised with 0.1% Reactive Red 6. Binding to the dye was effective by incubation in 0.1 M sodium borohydrate buffer / 0.05 M NaCl for 2 hours. Resulting latex particles were spun and washed to remove any free dye.

1.2 Coacervate particles

Coacervate particles were constructed via the simple coacervation method: 3 gm gum arabic dissolved in 100ml water and maintained at 40°C. To this solution 0.5ml perfume was added and an emulsion was formed using a polytron. This solution was mixed with 100ml gelatine (also at 40°C). The pH of the resulting solution was reduced to pH 4 using 1M HCl. Sample was left to stir at room temperature and finally chilled on ice for 2 hours. The resulting particles were then derivatised with Reactive Red 6 as described for the latex particles (section 1.1).

Example 5

Binding perfumed latex particles to cotton.

Assay: 100µl 0.1% Reactive Red 6 coated perfume latex was added to a reaction vial in a total volume of 5ml. This
5 contained 50mM HEPES buffer and 200µl PBST. The solution also contained either no CBD fusion or 200µg CBD-anti Reactive Red 6 protein.

Solutions were mixed for 30 minutes at room temperature.
10 Cotton strands (2 x 3cm - prewashed in buffer) were then added to the vials and incubated for a further 30 minutes at room temperature. The cotton strands were then removed and placed into glass vials containing 25 ml water and were rinsed. The resulting strands were then placed into glass
15 vials containing 1ml ethyl acetate and capped. These vials were sonicated and then analysed by GC-MS following specific perfume components. Figure 2 demonstrates the enhanced binding of perfumed latex in the presence of the fusion protein.

20

Example 6

Binding perfume containing coacervate particles to cotton

Assay: 50µl 0.375% coacervate particles were added to a
25 reaction volume containing 2ml PBST either in the presence or absence of 100µg fusion protein. The samples were mixed for 30 minutes prior to the addition of 2cm cotton yarns. Samples were incubated for a further 30 minutes at room temperature and were rinsed at the end of of this period in
30 25 ml PBST. The cotton strands were then added to glass vials containing 1ml ethyl acetate and were incubated at 50°C for 30 minutes and sonicated. Perfume components of the encapsulated perfume were then analysed by GC-MS. Figures 3 and 4 below clearly demonstrate the enhanced level of
35 perfume deposition using this novel cross linking fusion protein approach.

20

SEQUENCE LISTING

<110> Unilever PLC

<120> Detergent Compositions Comprising Benefit Agents

<130> 748-2961

<140> CA 2,392,158

<141> 2001-05-30

<150> EP99310428

<151> 1999-12-22

<160> 2

<170> PatentIn. version 3.1

<210> 1

<211> 558

<212> DNA

<213> Trichoderma reesei

<220>

<221> CDS

<222> (3)..(551)

<400> 1

tc	gag	aaa	aga	cag	gtg	cag	ctg	cag	gag	tca	ggg	gga	gga	ttg	gtg	47
	Glu	Lys	Arg	Gln	Val	Gln	Leu	Gln	Glu	Ser	Gly	Gly	Gly	Leu	Val	
1					5					10					15	

cag	gct	ggg	ggc	tct	ctg	aga	ctc	tcc	tgt	gca	gcc	tcc	gga	cgc	gcc	95
Gln	Ala	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Arg	Ala	
				20					25					30		

acc	agt	ggt	cat	ggt	cac	tac	ggt	atg	ggc	tgg	ttc	cgc	cag	gtt	cca	143
Thr	Ser	Gly	His	Gly	His	Tyr	Gly	Met	Gly	Trp	Phe	Arg	Gln	Val	Pro	
			35					40					45			

ggg	aag	gag	cgt	gag	ttt	gtc	gca	gct	att	agg	tgg	agt	ggc	aaa	gag	191
Gly	Lys	Glu	Arg	Glu	Phe	Val	Ala	Ala	Ile	Arg	Trp	Ser	Gly	Lys	Glu	
		50					55					60				

aca	tgg	tat	aaa	gac	tcc	gtg	aag	ggc	cga	ttc	acc	atc	tcc	aga	gat	239
Thr	Trp	Tyr	Lys	Asp	Ser	Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	
	65					70				75						

aac	gcc	aag	act	acg	gtt	tac	ctg	caa	atg	aac	agc	ctg	aaa	cct	gaa	287
Asn	Ala	Lys	Thr	Thr	Val	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Lys	Pro	Glu	
80					85				90					95		

gat	acg	gcc	gtt	tat	tat	tgt	gcc	gct	cga	cgg	gtc	cgc	gtg	gat	gat	335
Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Ala	Arg	Pro	Val	Arg	Val	Asp	Asp	
				100					105					110		

att	tcc	ctg	ccg	gtt	ggg	ttt	gac	tac	tgg	ggc	cag	ggg	acc	cag	gtc	383
Ile	Ser	Leu	Pro	Val	Gly	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Gln	Val	
			115				120						125			

21

acc gtc tcc tca gga tct cat cac cat cac cat cac gga tcc acc tcc 431
 Thr Val Ser Ser Gly Ser His His His His His His Gly Ser Thr Ser
 130 135 140

att gaa ggt cgt acc cag tct cac tac ggt cag tgt ggt ggt att ggt 479
 Ile Glu Gly Arg Thr Gln Ser His Tyr Gly Gln Cys Gly Gly Ile Gly
 145 150 155

tac tcc ggt cca acc gtc tgt gcc tct ggt acc acc tgt cag gtt ctg 527
 Tyr Ser Gly Pro Thr Val Cys Ala Ser Gly Thr Thr Cys Gln Val Leu
 160 165 170 175

aac cct tac tac tcc cag tgt ctg taataag 558
 Asn Pro Tyr Tyr Ser Gln Cys Leu
 180

<210> 2
 <211> 183
 <212> PRT
 <213> Trichoderma reesei

<400> 2
 Glu Lys Arg Gln Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln
 1 5 10 15
 Ala Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Arg Ala Thr
 20 25 30
 Ser Gly His Gly His Tyr Gly Met Gly Trp Phe Arg Gln Val Pro Gly
 35 40 45
 Lys Glu Arg Glu Phe Val Ala Ala Ile Arg Trp Ser Gly Lys Glu Thr
 50 55 60
 Trp Tyr Lys Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn
 65 70 75 80
 Ala Lys Thr Thr Val Tyr Leu Gln Met Asn Ser Leu Lys Pro Glu Asp
 85 90 95
 Thr Ala Val Tyr Tyr Cys Ala Ala Arg Pro Val Arg Val Asp Asp Ile
 100 105 110
 Ser Leu Pro Val Gly Phe Asp Tyr Trp Gly Gln Gly Thr Gln Val Thr
 115 120 125
 Val Ser Ser Gly Ser His His His His His His Gly Ser Thr Ser Ile
 130 135 140
 Glu Gly Arg Thr Gln Ser His Tyr Gly Gln Cys Gly Gly Ile Gly Tyr
 145 150 155 160
 Ser Gly Pro Thr Val Cys Ala Ser Gly Thr Thr Cys Gln Val Leu Asn
 165 170 175
 Pro Tyr Tyr Ser Gln Cys Leu
 180

Claims:

1. Detergent composition comprising one or more surfactants and a fusion protein comprising a cellulose binding domain and a domain having a high binding affinity for another ligand, wherein the domain having a high binding affinity is an antibody or antibody fragment which is directed at a Benefit Agent selected from the group consisting of fabric softening agents, fragrances, perfumes, polymeric lubricants, photoprotective agents, latexes, resins, dye fixative agents, encapsulated materials, antioxidants, insecticides, soil repelling agents and soil release agents.
2. Detergent composition according to claim 1, wherein the cellulose binding domain is obtained from a fungal enzyme origin or from a bacterial enzyme origin.
3. Detergent composition according to claim 2, wherein the origin of the fungal enzyme is selected from Humicola, Trichoderma, Thermomonospora, Phanerochaete, and Aspergillus and wherein the origin of the bacterial enzyme is selected from Bacillus, Clostridium, Streptomyces, Cellulomonas and Pseudomonas.
4. Detergent composition according to claim 2 or 3, wherein the cellulose binding domain is obtained from Trichoderma reesei.
5. Detergent composition according to any one of claims 1 to 4, wherein the domain having a high binding affinity is a Heavy Chain antibody which is found in Camelidae.
6. Detergent composition according to any one of claims 1 to 5, wherein the cellulose binding domain is connected to the domain having a high binding affinity for another ligand by means of a linker consisting of 2-15 amino acids.


7. Detergent composition according to claim 6 wherein the linker consists of 2 to 5 amino acids.

8. Detergent composition according to any one of claims 1 to 7, wherein the domain having a high binding affinity is directed at micro-particles which are loaded with the benefit agent.

9. Detergent composition according to any one of claims 1 to 8, wherein the domain having a high binding affinity is a multi-specific antibody or antibody fragment or an analogous structure, whereby at least one specificity is directed to the fabric and the others are directed to one or more benefit agents.

10. Process for delivering a benefit agent to a fabric by treating said fabric with a composition comprising a fusion protein according to any one of claims 1 to 9 and a benefit agent selected from the group consisting of softening agents, finishing agents/ protective agents, fragrances and bleaching agents


Fig.1.



2/3

Fig.2.

GC-MS On Cotton Yarns Washed With Perfumed Latex



3/3

Fig.3.




Fig.4.

