



US006150134A

# United States Patent [19]

[11] **Patent Number:** **6,150,134**

**Maertens et al.**

[45] **Date of Patent:** **Nov. 21, 2000**

[54] **PURIFIED HEPATITIS C VIRUS ENVELOPE PROTEINS FOR DIAGNOSTIC AND THERAPEUTIC USE**

PCT/IT92/  
00081 1/1992 WIPO .  
PCT/US92/  
07189 1/1992 WIPO .  
PCT/US93/  
00907 1/1993 WIPO .

[75] Inventors: **Geert Maertens**, Bruges; **Fons Bosman**, Opwijk; **Guy De Martynoff**, Waterloo; **Marie-Ange Buysel**, Merelbeke, all of Belgium

### OTHER PUBLICATIONS

[73] Assignee: **Innogenetics, N.V.**, Ghent, Belgium

Ralston et al., "Characterization of Hepatitis C Virus . . .," *J. Virol* 67: 6753-6761 (1993).

[21] Appl. No.: **08/612,973**

Nishihara et al., "Secretion and Purification of Hepatitis C . . .," *Gene* 129: 207-214 (1993).

[22] PCT Filed: **Jul. 31, 1995**

Choo et al., "Vaccination of Chimpanzees Against . . . Hepatitis C Virus," *Proc Natl Acad Sci* 91:1294-1298 (1994).

[86] PCT No.: **PCT/EP95/03031**

Lanford et al., "Analysis of Hepatitis C Virus . . .," *Virology* 197: 225-235 (1993).

§ 371 Date: **Mar. 11, 1996**

§ 102(e) Date: **Mar. 11, 1996**

[87] PCT Pub. No.: **WO96/04385**

*Primary Examiner*—Donna C. Wortman  
*Assistant Examiner*—Mary K Zeman  
*Attorney, Agent, or Firm*—Nixon & Vanderhye P.C.

PCT Pub. Date: **Feb. 15, 1996**

### [30] Foreign Application Priority Data

### [57] ABSTRACT

Jul. 29, 1994 [EP] European Pat. Off. . . . . 94870132

The present invention relates to a method for purifying recombinant HCV single or specific oligomeric envelope proteins selected from the group consisting of E1 and/or E2 and/or E1/E2, characterized in that upon lysing the transformed host cells to isolate the recombination expressed protein a disulphide bond cleavage or reduction step is carried out with a disulphide bond cleavage agent. The present invention also relates to a composition isolated by such a method. The present invention also relates to the diagnostic and therapeutic application of these compositions. Furthermore, the invention relates to the use of HCV E1 protein and peptides for prognosing and monitoring the clinical effectiveness and/or clinical outcome of HCV treatment.

[51] **Int. Cl.<sup>7</sup>** . . . . . **C12N 15/09**; A61K 39/29

[52] **U.S. Cl.** . . . . . **435/69.3**; 435/69.1; 435/235.1; 435/803; 530/350; 424/228.1

[58] **Field of Search** . . . . . 435/69.3, 69.1, 435/235.1, 803; 530/350; 424/228.1

### [56] References Cited

#### U.S. PATENT DOCUMENTS

5,514,539 5/1996 Bukh et al. . . . . 435/5  
5,610,009 3/1997 Watanabe et al. . . . . 435/5

#### FOREIGN PATENT DOCUMENTS

PCT/US91/  
08272 1/1991 WIPO .

**26 Claims, 58 Drawing Sheets**




Fig. 1




Fig. 2




Fig. 3




Fig. 4

Anti-E1 levels in NON-responders to IFN treatment




Fig. 5

Anti-E1 levels in RESPONDERS to IFN treatment




Fig. 6

Anti-E1 levels in patients with COMPLETE response to IFN

SERIES 2





Fig. 7



Anti-E1 levels in INCOMPLETE responders to IFN treatment


SERIES 2



months after start of treatment

Fig. 8

Anti-E2 levels in NON-RESPONDERS to IFN treatment



weeks after start of treatment

Fig. 9

Anti-E2 levels in RESPONDERS to IFN treatment





Fig.10

Anti-E2 levels in INCOMPLETE responders to IFN treatment

SERIES 2




months after start of treatment

Fig.11

Anti-E2 levels in COMPLETE responders to IFN treatment

SERIES 2



months after start of treatment

Fig.12

# Human anti-E1 reactivity competed with peptides




Fig.13

Competition of reactivity of anti-E1 Mabs with peptides




Fig.14

Anti-E1 (epitope 1) levels in NON-RESPONDERS to IFN treatment




Fig.15



Anti-E1 (epitope 1) levels in RESPONDERS to IFN treatment




Fig.16

Anti-E1 (epitope 2) levels in NON-RESPONDERS to IFN treatment




Fig.17

Anti-E1 (epitope 2) levels in RESPONDERS to IFN treatment




Fig.18

# Competition of reactivity of anti-E2 Mabs with peptides




Fig.19

# Human anti-E2 reactivity competed with peptides




Fig. 20

## Fig. 21A

5' GGCATGCAAGCTTAATTAATT3' (SEQ ID NO 1)  
3'ACGTCCGTACGTTTCGAATTAATTAATCGA5' (SEQ ID NO 94)

5'CCGGGGAGGCCTGCACGTGATCGAGGGCAGACACCATCACCACCATCACTAATAGT  
TAATTAAGTCA 3' (SEQ ID NO 2)  
3'CCTCCGGACGTGCACTAGCTCCCGTCTGTGGTAGTGGTGGTAGTGATTATCAATTAATTG  
5' (SEQ ID NO 95)

SEQ ID NO 3 (HCCI9A)

ATGCCCGGTTGCTCTTTCTCTATCTTCCTCTTGGCTTTACTGTCCTGTCTGACCATTCCA  
GCTTCCGCTTATGAGGTGCGCAACGTGTCCGGGATGTACCATGTCACGAACGACTGCT  
CCAAGTCAAGCATTGTGTATGAGGCAGCGGACATGATCATGCACACCCCCGGGTGCGT  
GCCCTGCGTTTCGGGAGAACAACCTCTCCCGCTGCTGGGTAGCGCTCACCCCCACGCTC  
GCAGCTAGGAACGCCAGCGTCCCCACCACGACAATACGACGCCACGTCGATTTGCTCG  
TTGGGGCGGCTGCTCTCTGTTCCGCTATGTACGTGGGGGATCTCTGCGGATCTGTCTTC  
CTCGTCTCCAGCTGTTACCATCTCGCCTCGCCGGCATGAGACGGTGCAGGACTGCA  
ATTGCTCAATCTATCCCGGCCACATAACAGGTCACCGTATGGCTTGGGATATGATGAT  
GAACTGGTGCCTACAACGGCCCTGGTGGTATCGCAGCTGCTCCGGATCCCACAAGCT  
GTCGTGGACATGGTGGCGGGGGCCATTGGGGAGTCTGGCGGGCCTCGCTACTATT  
CCATGGTGGGGAAGTGGGCTAAGGTTTTGATTGTGATGCTACTCTTTGCTCTCTAATAG

SEQ ID NO 5 (HCCI10A)

ATGTTGGGTAAGGTCATCGATACCCTTACATGCGGCTTCGCCGACCTCGTGGGGTACA  
TTCCGCTCGTCGGCGCCCCCTAGGGGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCG  
GGTCTGGAGGACGGCGTGAAGTATGCAACAGGGAATTTGCCCGGTTGCTCTTTCTCT  
ATCTTCCTCTTGGCTTTGCTGTCTGTCTGACCGTTCAGCTTCCGCTTATGAAGTGC  
CAACGTGTCCGGGATGTACCATGTCACGAACGACTGCTCCAAGTCAAGCATTGTGTAT  
GAGGCAGCGGACATGATCATGCACACCCCCGGGTGCGTGCCCTGCGTTTCGGGAGAAC  
AACTCTTCCCGCTGCTGGGTAGCGCTCACCCCCACGCTCGCAGCTAGGAACGCCAGCG  
TCCCCACCACGACAATACGACGCCACGTCGATTTGCTCGTTGGGGCGGCTGCTTTCTG

## Fig. 21B

TTCCGCTATGTACGTGGGGGACCTCTGCGGATCTGTCTTCCTCGTCTCCCAGCTGTTCA  
CCATCTCGCCTCGCCGGCATGAGACGGTGCAGGACTGCAATTGCTCAATCTATCCCGG  
CCACATAACGGGTCACCGTATGGCTTGGGATATGATGATGAACTGGTCGCCTACAACG  
GCCCTGGTGGTATCGCAGCTGCTCCGGATCCCACAAGCTGTCTGGACATGGTGGCGG  
GGGCCATTGGGGAGTCTGGCGGGTCTCGCCTACTATTCCATGGTGGGGAACTGGGC  
TAAGGTTTTGATTGTGATGCTACTCTTTGCTCCCTAATAG

SEQ ID NO 7 (HCCI11A)

ATGTTGGGTAAGGTCATCGATACCCTTACGTGCGGCTTCGCCGACCTCATGGGGTACA  
TTCCGCTCGTCGGCGCCCCCTAGGGGGTGTGCCAGAGCCCTGGCGCATGGCGTCCG  
GGTTCTGGAAGACGGCGTGAACATGCAACAGGGAATTTGCCTGGTTGCTCTTTCTCTA  
TCTTCCTCTTGGCTTTACTGTCCTGTCTGACCATTCCAGCTTCCGCTTATGAGGTGCGC  
AACGTGTCCGGGATGTACCATGTCACGAACGACTGCTCCAACCAAGCATTGTGTATG  
AGGCAGCGGACATGATCATGCACACCCCCGGGTGCGTGCCCTGCGTTCGGGAGAACA  
ACTCTTCCCGCTGCTGGGTAGCGCTCACCCCCACGCTCGCAGCTAGGAACGCCAGCGT  
CCCCACTACGACAATACGACGCCACGTCGATTTGCTCGTTGGGGCGGCTGCTTTCTGTT  
CCGCTATGTACGTGGGGGATCTCTGCGGATCTGTCTTCCTCGTCTCCCAGCTGTTACC  
ATCTCGCCTCGCCGGCATGAGACGGTGCAGGACTGCAATTGCTCAATCTATCCCGGCC  
ACATAACAGGTCACCGTATGGCTTGGGATATGATGATGAACTGGTAATAG

SEQ ID NO 9 (HCCI12A)

ATGCCCGTTGCTCTTTCTCTATCTTCCTCTTGGCCCTGCTGTCCTGTCTGACCATACCA  
GCTTCCGCTTATGAAGTGCGCAACGTGTCCGGGGTGTACCATGTCACGAACGACTGCT  
CCAACCAAGCATAGTGTATGAGGCAGCGGACATGATCATGCACACCCCCGGGTGCGT  
GCCCTGCGTTCGGGAGGGCAACTCCTCCCGTTGCTGGGTGGCGCTCACTCCCACGCTC  
GCGGCCAGGAACGCCAGCGTCCCCACAACGACAATACGACGCCACGTCGATTTGCTC  
GTTGGGGCTGCTGCTTTCTGTTCCGCTATGTACGTGGGGGATCTCTGCGGATCTGTTTT  
CCTTGTTCACGCTGTTACCTTCTCACCTCGCCGGCATCAAACAGTACAGGACTGCA  
ACTGCTCAATCTATCCCGGCCATGTATCAGGTCACCGCATGGCTTGGGATATGATGAT  
GAACTGGTCCTAATAG

SEQ ID NO 11 (HCCI13A)

ATGTCCGGTTGCTCTTTCTCTATCTTCCTCTTGGCCCTGCTGTCCTGTCTGACCATACCA  
GCTTCCGCTTATGAAGTGCGCAACGTGTCCGGGGTGTACCATGTCACGAACGACTGCT  
CCAACCAAGCATAGTGTATGAGGCAGCGGACATGATCATGCACACCCCCGGGTGCGT

## Fig. 21C

GCCCTGCGTTTCGGGAGGGCAACTCCTCCCGTTGCTGGGTGGCGCTCACTCCCACGCTC  
GCGGCCAGGAACGCCAGCGTCCCCACAACGACAATACGACGCCACGTCGATTTGCTC  
GTTGGGGCTGCTGCTTTTCTGTTCCGCTATGTACGTGGGGGATCTCTGCGGATCTGTTTT  
CCTTGTTTTCCAGCTGTTACCTTCTCACCTCGCCGGCATCAAACAGTACAGGACTGCA  
ACTGCTCAATCTATCCCGGCCATGTATCAGGTCACCGCATGGCTTGGGATATGATGAT  
GAACTGGTAATAG

SEQ ID NO 13 (HCCI17A)

ATGCTGGGTAAGGCCATCGATACCCTTACGTGCGGCTTCGCCGACCTCGTGGGGTACA  
TTCCGCTCGTCGGCGCCCCCTAGGGGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCG  
GGTTCTGGAAGACGGCGTGAACATGCAACAGGGAATTTGCCTGGTTGCTCTTTCTCTA  
TCTTCTCTTGGCTTTACTGTCCTGTCTAACCATTCCAGCTTCCGCTTACGAGGTGCGC  
AACGTGTCCGGGATGTACCATGTACGAACGACTGCTCCAACCTCAAGCATTGTGTATG  
AGGCAGCGGACATGATCATGCACACCCCCGGGTGCGTGCCCTGCGTTCCGGGAGAACA  
ACTCTTCCCCTGCTGGGTAGCGCTCACCCCCACGCTCGCGGCTAGGAACGCCAGCAT  
CCCCACTACAACAATACGACGCCACGTGATTTGCTCGTTGGGGCGGCTGCTTTCTGTT  
CCGCTATGTACGTGGGGGATCTCTGCGGATCTGTCTTCTCGTCTCCCAGCTGTTACCC  
ATCTCGCCTCGCCGGCATGAGACGGTGCAGGACTGCAATTGCTCAATCTATCCCGGCC  
ACATAACGGGTCACCGTATGGCTTGGGATATGATGATGAACTGGTACTAATAG

SEQ ID NO 15 (HCPPr51)

ATGCCCGGTTGCTCTTTCTCTATCTT

SEQ ID NO 16 (HCPPr52)

ATGTTGGGTAAGGTCATCGATACCCT

SEQ ID NO 17 (HCPPr53)

CTATTAGGACCAGTTCATCATCATATCCCA

SEQ ID NO 18 (HCPPr54)

CTATTACCAGTTCATCATCATATCCCA

SEQ ID NO 19 (HCPPr107)

ATACGACGCCACGTGCGATTCCCAGCTGTTACCATC



## Fig. 21D

SEQ ID NO 20 (HCP108)

GATGGTGAACAGCTGGAATCGACGTGGCGTCGTAT

SEQ ID NO 21 (HCCI37)

ATGTTGGGTAAGGTCATCGATACCCTTACATGCGGCTTCGCCGACCTCGTGGGGTACA  
TTCCGCTCGTCGGCGCCCCCTAGGGGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCG  
GGTTCTGGAGGACGGCGTGAACATGCAACAGGGAATTTGCCCGGTTGCTCTTTCTCT  
ATCTTCCTCTTGGCTTTGCTGTCTGTCTGACCGTTCCAGCTTCCGCTTATGAAGTGCG  
CAACGTGTCCGGGATGTACCATGTCACGAACGACTGCTCCAACCTCAAGCATTGTGTAT  
GAGGCAGCGGACATGATCATGCACACCCCCGGGTGCGTGCCCTGCGTTCGGGAGAAC  
AACTCTTCCCGCTGCTGGGTAGCGCTCACCCCCACGCTCGCAGCTAGGAACGCCAGCG  
TCCCCACCACGACAATACGACGCCACGTTCGATTCCCAGCTGTTCAACCATCTCGCCTCG  
CCGGCATGAGACGGTGCAGGACTGCAATTGCTCAATCTATCCCGGCCACATAACGGGT  
CACCGTATGGCTTGGGATATGATGATGAACTGGTGCCTACAACGGCCCTGGTGGTAT  
CGCAGCTGCTCCGGATCCACAAGCTGTCTGTGACATGGTGGCGGGGGCCATTGGGG  
AGTCCTGGCGGGTCTCGCCTACTATTCCATGGTGGGGAACCTGGGCTAAGGTTTTGATTG  
TGATGCTACTCTTTGCTCCCTAATAG

SEQ ID NO 23 (HCCI38)

ATGTTGGGTAAGGTCATCGATACCCTTACATGCGGCTTCGCCGACCTCGTGGGGTACA  
TTCCGCTCGTCGGCGCCCCCTAGGGGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCG  
GGTTCTGGAGGACGGCGTGAACATGCAACAGGGAATTTGCCCGGTTGCTCTTTCTCT  
ATCTTCCTCTTGGCTTTGCTGTCTGTCTGACCGTTCCAGCTTCCGCTTATGAAGTGCG  
CAACGTGTCCGGGATGTACCATGTCACGAACGACTGCTCCAACCTCAAGCATTGTGTAT  
GAGGCAGCGGACATGATCATGCACACCCCCGGGTGCGTGCCCTGCGTTCGGGAGAAC  
AACTCTTCCCGCTGCTGGGTAGCGCTCACCCCCACGCTCGCAGCTAGGAACGCCAGCG  
TCCCCACCACGACAATACGACGCCACGTTCGATTCCCAGCTGTTCAACCATCTCGCCTCG  
CCGGCATGAGACGGTGCAGGACTGCAATTGCTCAATCTATCCCGGCCACATAACGGGT  
CACCGTATGGCTTGGGATATGATGATGAACTGGTAA  
TAG

SEQ ID NO 25 (HCCI39)

ATGTTGGGTAAGGTCATCGATACCCTTACATGCGGCTTCGCCGACCTCGTGGGGTACA  
TTCCGCTCGTCGGCGCCCCCTAGGGGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCG  
GGTTCTGGAGGACGGCGTGAACATGCAACAGGGAATTTGCCCGGTTGCTCTTTCTCT

## Fig. 21E

ATCTTCCTCTTGGCTTTGCTGTCCTGTCTGACCGTTCCAGCTTCCGCTTATGAAGTGCG  
CAACGTGTCCGGGATGTACCATGTCACGAACGACTGCTCCAACCTCAAGCATTGTGTAT  
GAGGCAGCGGACATGATCATGCACACCCCCGGGTGCGTGCCCTGCGTTCGGGAGAAC  
AACTCTTCCCGCTGCTGGGTAGCGCTCACCCCCACGCTCGCAGCTAGGAACGCCAGCG  
TCCCCACCACGACAATACGACGCCACGTGCGATTCCCAGCTGTTACCATCTCGCCTCG  
CCGGCATGAGACGGTGCAGGACTGCAATTGCTCAATCTATCCCGGCCACATAACGGGT  
CACCGTATGGCTTGGGATATGATGATGAACTGGTCGCCTACAACGGCCCTGGTGGTAT  
CGCAGCTGCTCCGGATCCTCTAATAG

SEQ ID NO 27 (HCCI40)

ATGTTGGGTAAGGTCATCGATACCCTTACATGCGGCTTCGCCGACCTCGTGGGGTACA  
TTCCGCTCGTCGGCGCCCCCTAGGGGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCG  
GTTTCTGGAGGACGGCGTGAACCTATGCAACAGGGAATTTGCCCGTTGCTCTTTCTCT  
ATCTTCCTCTTGGCTTTGCTGTCCTGTCTGACCGTTCCAGCTTCCGCTTATGAAGTGCG  
CAACGTGTCCGGGATGTACCATGTCACGAACGACTGCTCCAACCTCAAGCATTGTGTAT  
GAGGCAGCGGACATGATCATGCACACCCCCGGGTGCGTGCCCTGCGTTCGGGAGAAC  
AACTCTTCCCGCTGCTGGGTAGCGCTCACCCCCACGCTCGCAGCTAGGAACGCCAGCG  
TCCCCACCACGACAATACGACGCCACGTGCGATTCCCAGCTGTTACCATCTCGCCTCG  
CCGGCATGAGACGGTGCAGGACTGCAATTGCTCAATCTATCCCGGCCACATAACGGGT  
CACCGTATGGCTTGGGATATGATGATGAACTGGTCGCCTACAACGGCCCTGGTGGTAT  
CGCAGCTGCTCCGGATCGTGATCGAGGGCAGACACCATCACACCATCACTAATAG

SEQ ID NO 29 (HCCI62)

ATGGGTAAGGTCATCGATACCCTTACGTGCGGATTCGCCGATCTCATGGGGTACATCC  
CGCTCGTCGGCGCTCCCGTAGGAGGCGTCGCAAGAGCCCTTGCGCATGGCGTGAGGGC  
CCTTGAAGACGGGATAAATTTGCAACAGGGAATTTGCCCGTTGCTCCTTTTCTATTT  
TCCTTCTCGCTCTGTTCTCTTGCTTAATTCATCCAGCAGCTAGTCTAGAGTGCGGGAAT  
ACGTCTGGCCTCTATGTCCTTACCAACGACTGTTCCAATAGCAGTATTGTGTACGAGGC  
CGATGACGTTATTCTGCACACACCCGGCTGCATACCTTGTGTCCAGGACGGCAATACA  
TCCACGTGCTGGACCCAGTGACACCTACAGTGGCAGTCAAGTACGTCCGAGCAACCA  
CCGCTTCGATACGCAGTCATGTGGACCTATTAGTGGGCGCGGCCACGATGTGCTCTGC  
GCTCTACGTGGGTGACATGTGTGGGGCTGTCTTCCCTCGTGGGACAAGCCTTACGTTCA  
GACCTCGTCGCCATCAAACGGTCCAGACCTGTAAGTCTCGCTGTACCCAGGCCATCT  
TTCAGGACATCGAATGGCTTGGGATATGATGATGAACTGGTAATAG

## Fig. 21F

SEQ ID NO 31 (HCCI63)

ATGGGTAAGGTCATCGATACCCTAACGTGCGGATTCGCCGATCTCATGGGGTATATCC  
CGCTCGTAGGCGGCCCCATTGGGGGCGTCGCAAGGGCTCTCGCACACGGTGTGAGGGT  
CCTTGAGGACGGGGTAAACTATGCAACAGGGGAATTTACCCGGTTGCTCTTTCTCTATCT  
TTATTCTTGCTCTTCTCTCGTGTCTGACCGTTCGGCCTCTGCAGTTCCTACCGAAATG  
CCTCTGGGATTTATCATGTTACCAATGATTGCCCAAACCTTTCCATAGTCTATGAGGCCA  
GATAACCTGATCCTACACGCACCTGGTTGCGTGCCTTGTGTGCATGACAGGTAATGTGA  
GTAGATGCTGGGTCCAAATTACCCCTACACTGTCAGCCCCGAGCCTCGGAGCAGTCAC  
GGCTCCTCTTCGGAGAGCCGTTGACTACCTAGCGGGAGGGGCTGCCCTCTGCTCCGCG  
TTATACGTAGGAGACGCGTGTGGGGCACTATTCTTGGTAGGCCAAATGTTACCTATA  
GGCCTCGCCAGCACGCTACGGTGCAGAACTGCAACTGTTCCATTTACAGTGGCCATGT  
TACCGGCCACCGGATGGCATGGGATATGATGATGAACTGGTAATAG

SEQ ID NO 33 (HCP109)

TGGGATATGATGATGAACTGGTC

SEQ ID NO 34 (HCP172)

CTATTATGGTGGTAAKGCCARCARGAGCAGGAG

SEQ ID NO 35 (HCCL22A)

TGGGATATGATGATGAACTGGTCGCCTACAACGGCCCTGGTGGTATCGCAGCTGCTCC  
GGATCCCACAAGCTGTGCGTGGACATGGTGGCGGGGGCCATTGGGGAGTCCTGGCGG  
GCCTCGCCTACTATTCCATGGTGGGGAACCTGGGCTAAGGTTTTGGTTGTGATGCTACTC  
TTTGCCGGCGTCGACGGGCATACCCGCGTGTGAGGAGGGGCAGCAGCCTCCGATACCA  
GGGGCCTTGTGTCCCTCTTTAGCCCCGGGTGCGCTCAGAAAATCCAGCTCGTAAACAC  
CAACGGCAGTTGGCACATCAACAGGACTGCCCTGAACTGCAACGACTCCCTCCAAAC  
AGGGTTCTTTGCCGCACTATTCTACAAACACAAATTCAACTCGTCTGGATGCCAGAG  
CGCTTGGCCAGCTGTGCTCCATCGACAAGTTCGCTCAGGGGTGGGGTCCCCTCACTT  
ACACTGAGCCTAACAGCTCGGACCAGAGGCCCTACTGCTGGCACTACGCGCCTCGACC  
GTGTGGTATTGTACCCGCGTCTCAGGTGTGCGGTCCAGTGTATTGCTTACCCCCGAGCC  
CTGTTGTGGTGGGGACGACCGATCGGTTTGGTGTCCCCACGTATAACTGGGGGGCGAA  
CGACTCGGATGTGCTGATTCTCAACAACACGCGGCCGCGGAGGCAACTGGTTCGGC  
TGTACATGGATGAATGGCACTGGGTTACCAAGACGTGTGGGGGGCCCCCGTGCAACA  
TCGGGGGGGGCCGGCAACAACACCTTGACCTGCCCACTGACTGTTTTCGGAAGCACCC  
CGAGGCCACCTACGCCAGATGCGGTTCTGGGCCCTGGCTGACACCTAGGTGTATGGTT

## Fig. 21G

CATTACCCATATAGGCTCTGGCACTACCCCTGCACTGTCAACTTCACCATCTTCAAGGT  
TAGGATGTACGTGGGGGGCGTGGAGCACAGGTTTGAAGCCGCATGCAATTGGACTCG  
AGGAGAGCGTTGTGACTTGGAGGACAGGGATAGATCAGAGCTTAGCCCGCTGCTGCTG  
TCTACAACAGAGTGGCAGATACTGCCCTGTTCTTACCACCCTGCCGGCCCTATCCA  
CCGGCCTGATCCACCTCCATCAGAACATCGTGGACGTGCAATACCTGTACGGTGTAGG  
GTCCGGCGGTTGTCTCCCTTGTTCATCAAATGGGAGTATGTCTGTTGCTCTTCTTCTCT  
GGCAGACGCGCGCATCTGCGCCTGCTTATGGATGATGCTGCTGATAGCTCAAGCTGAG  
GCCGCCTTAGAGAACCTGGTGGTCCCAATGCGGCGGCCGTGGCCGGGGCGCATGGC  
ACTCTTTCTTCTTGTGTTCTTCTGTGCTGCCTGGTACATCAAGGGCAGGCTGGTCCC  
TGGTGCGGCATACGCCTTCTATGGCGTGTGGCCGCTGCTCCTGCTTCTGCTGGCCTTAC  
CACCACGAGCTTATGCCTAGTAA

SEQ ID NO 37 (HCCI41)

GATCCCAACAAGCTGTCGTGGACATGGTGGCGGGGGCCATTGGGGAGTCTGGCGGG  
CCTCGCCTACTATTCCATGGTGGGGAAGTGGGCTAAGGTTTTGGTTGTGATGCTACTCT  
TTGCCGGCGTCGACGGGCATACCCGCGTGTCAAGAGGGGCAGCAGCCTCCGATACCA  
GGGGCCTTGTGTCCCTCTTTAGCCCCGGGTCGGCTCAGAAAATCCAGCTCGTAAACAC  
CAACGGCAGTTGGCACATCAACAGGACTGCCCTGAACTGCAACGACTCCCTCCAAAC  
AGGGTTCTTTGCCGCACTATTCTACAAACACAAATTCAACTCGTCTGGATGCCAGAG  
CGCTTGGCCAGCTGTCGCTCCATCGACAAGTTCGCTCAGGGGTGGGGTCCCCTCACTT  
ACACTGAGCCTAACAGCTCGGACCAGAGGCCCTACTGCTGGCACTACGCGCCTCGACC  
GTGTGGTATTGTACCCGCGTCTCAGGTGTGCGGTCCAGTGTATTGCTTACCCCCGAGCC  
CTGTTGTGGTGGGGACGACCGATCGGTTTGGTGTCCCCACGTATAACTGGGGGGCGAA  
CGACTCGGATGTGCTGATTCTCAACAACACGCGGCCGCCGCGAGGCAACTGGTTCGGC  
TGACATGGATGAATGGCACTGGGTTACCAAGACGTGTGGGGGGCCCCCGTGAACA  
TCGGGGGGGGCCGGCAACAACACCTTGACCTGCCCCACTGACTGTTTTCGGAAGCACCC  
CGAGGCCACCTACGCCAGATGCGGTTCTGGGCCCTGGCTGACACCTAGGTGTATGGTT  
CATTACCCATATAGGCTCTGGCACTACCCCTGCACTGTCAACTTCACCATCTTCAAGGT  
TAGGATGTACGTGGGGGGCGTGGAGCACAGGTTTGAAGCCGCATGCAATTGGACTCG  
AGGAGAGCGTTGTGACTTGGAGGACAGGGATAGATCAGAGCTTAGCCCGCTGCTGCTG  
TCTACAACAGAGTGGCAGAGTGGCAGAGCTTAATTAATTAG

SEQ ID NO 39 (HCCI42)

GATCCCAACAAGCTGTCGTGGACATGGTGGCGGGGGCCATTGGGGAGTCTGGCGGG  
CCTCGCCTACTATTCCATGGTGGGGAAGTGGGCTAAGGTTTTGGTTGTGATGCTACTCT

## Fig. 21H

TTGCCGGCGTCGACGGGCATACCCGCGTGTTCAGGAGGGGCAGCAGCCTCCGATACCA  
GGGGCCTTGTGTCCCTCTTTAGCCCCGGGTTCGGCTCAGAAAATCCAGCTCGTAAACAC  
CAACGGCAGTTGGCACATCAACAGGACTGCCCTGAACTGCAACGACTCCCTCCAAAC  
AGGGTTCTTTGCCGCACTATTCTACAAACACAAATTCAACTCGTCTGGATGCCAGAG  
CGTTGGCCAGCTGTTCGCTCCATCGACAAGTTCGCTCAGGGGTGGGGTCCCCTCACTT  
AACTGAGCCTAACAGCTCGGACCAGAGGCCCTACTGCTGGCACTACGCGCCTCGACC  
GTGTGGTATTGTACCCGCGTCTCAGGTGTGCGGTCCAGTGTATTGCTTACCCCGAGCC  
CTGTTGTGGTGGGGACGACCGATCGGTTTTGGTGTCCCCACGTATAACTGGGGGGCGAA  
CGACTCGGATGTGCTGATTCTCAACAACACGCGGCCGCCGCGAGGCAACTGGTTCGGC  
TGTACATGGATGAATGGCACTGGGTTACCAAGACGTGTGGGGGGCCCCCGTGCAACA  
TCGGGGGGGCCGCAACAACACCTTGACCTGCCCCACTGACTGTTTTCGGAAGCACCC  
CGAGGCCACCTACGCCAGATGCGGTTCTGGGCCCTGGCTGACACCTAGGTGTATGGTT  
CATTACCCATATAGGCTCTGGCACTACCCCTGCACTGTCAACTTCACCATCTTCAAGGT  
TAGGATGTACGTGGGGGGCGTGGAGCACAGGTTTGAAGCCGCATGCAATTGGACTCG  
AGGAGAGCGTTGTGACTTGGAGGACAGGGATAGATCAGAGCTTAGCCCGCTGCTGCTG  
TCTACAACAGGTGATCGAGGGCAGACACCATCACCACCATCACTAATAG

SEQ ID NO 41 (HCCI43)

ATGGTGGGGAAGTGGGCTAAGTTTTGGTGTGATGCTACTCTTTGCCGGCGTCGACG  
GGCATAACCCGCGTGTTCAGGAGGGGCAGCAGCCTCCGATACCAGGGGCCTTGTGTCCCT  
CTTTAGCCCCGGGTTCGGCTCAGAAAATCCAGCTCGTAAACACCAACGGCAGTTGGCAC  
ATCAACAGGACTGCCCTGAACTGCAACGACTCCCTCCAAACAGGGTTCTTTGCCGCAC  
TATTCTACAAACACAAATTCAACTCGTCTGGATGCCAGAGCGCTTGGCCAGCTGTTCG  
CTCCATCGACAAGTTCGCTCAGGGGTGGGGTCCCCTCACTTAACTGAGCCTAACAGC  
TCGGACCAGAGGCCCTACTGCTGGCACTACGCGCCTCGACCGTGTGGTATTGTACCCG  
CGTCTCAGGTGTGCGGTCCAGTGTATTGCTTACCCCGAGCCCTGTTGTGGTGGGGAC  
GACCGATCGGTTTGGTGTCCCCACGTATAACTGGGGGGCGAACGACTCGGATGTGCTG  
ATTCTCAACAACACGCGGCCGCCGCGAGGCAACTGGTTCGGCTGTACATGGATGAATG  
GCACTGGGTTACCAAGACGTGTGGGGGGCCCCCGTGCAACATCGGGGGGGCCGGCA  
ACAACACCTTGACCTGCCCCACTGACTGTTTTCGGAAGCACCCCGAGGCCACCTACGC  
CAGATGCGGTTCTGGGCCCTGGCTGACACCTAGGTGTATGGTTCACTTACCCATATAGG  
CTCTGGCACTACCCCTGCACTGTCAACTTCACCATCTTCAAGGTTAGGATGTACGTGGG  
GGGCGTGGAGCACAGGTTTGAAGCCGCATGCAATTGGACTCGAGGAGAGCGTTGTGA  
CTTGGAGGACAGGGATAGATCAGAGCTTAGCCCGCTGCTGCTGTCTACAACAGAGTGG  
CAGAGCTTAATTAATTAG

## Fig. 21I

SEQ ID NO 43 (HCCI44)

ATGGTGGGGAAC TGGGCTAAGGTTTTGGTTGTGATGCTACTCTTTGCCGGCGTTCGACG  
GGCATACCCGCGTGT CAGGAGGGGCAGCAGCCTCCGATACCAGGGGCCTTGTGTCCCT  
CTTTAGCCCCGGGTCGGCTCAGAAAATCCAGCTCGTAAACACCAACGGCAGTTGGCAC  
ATCAACAGGACTGCCCTGAACTGCAACGACTCCCTCCAAACAGGGTTCTTTGCCGCAC  
TATTCTACAAACACAAATTCAACTCGTCTGGATGCCAGAGCGCTTGGCCAGCTGTCTG  
CTCCATCGACAAGTTCGCTCAGGGGTGGGGTCCCCTCACTTACACTGAGCCTAACAGC  
TCGGACCAGAGGCCCTACTGCTGGCACTACGCGCCTCGACCGTGTGGTATTGTACCCG  
CGTCTCAGGTGTGCGGTCCAGTGTATTGCTTACCCCGAGCCCTGTTGTGGTGGGGAC  
GACCGATCGGTTTGGTGTCCCCACGTATAACTGGGGGGCGAACGACTCGGATGTGCTG  
ATTCTCAACAACACGCGGCCCGCCGAGGCAACTGGTTCGGCTGTACATGGATGAATG  
GCACTGGGTTACCAAGACGTGTGGGGGCCCCCGTGCAACATCGGGGGGGCCGGCA  
ACAACACCTTGACCTGCCCCACTGACTGTTTTCGGAAGCACCCCGAGGCCACCTACGC  
CAGATGCGGTTCTGGGCCCTGGCTGACACCTAGGTGTATGGTTCATTACCCATATAGG  
CTCTGGCACTACCCCTGCACTGTCAACTTCACCATCTTCAAGGTTAGGATGTACGTGGG  
GGGCGTGGAGCACAGGTTTCAAGCCGCATGCAATTGGACTCGAGGAGAGCGTTGTGA  
CTTGGAGGACAGGGATAGATCAGAGCTTAGCCCGCTGCTGCTGTCTACAACAGGTGAT  
CGAGGGCAGACACCATCACCACCATCACTAATAG

SEQ ID NO 45 (HCCL64)

ATGGTGGCGGGGGCCCATTTGGGGAGTCCTGGCGGGCCTCGCCTACTATTCCATGGTGG  
GGAAC TGGGCTAAGGTTTTGGTTGTGATGCTACTCTTTGCCGGCGTTCGACGGGCATAC  
CCGCGTGT CAGGAGGGGCAGCAGCCTCCGATACCAGGGGCCTTGTGTCCCTCTTTAGC  
CCCGGGTCGGCTCAGAAAATCCAGCTCGTAAACACCAACGGCAGTTGGCACATCAAC  
AGGACTGCCCTGAACTGCAACGACTCCCTCCAAACAGGGTTCTTTGCCGCACTATTCT  
ACAAACACAAATTCAACTCGTCTGGATGCCAGAGCGCTTGGCCAGCTGTCTGCTCCAT  
CGACAAGTTCGCTCAGGGGTGGGGTCCCCTCACTTACACTGAGCCTAACAGCTCGGAC  
CAGAGGCCCTACTGCTGGCACTACGCGCCTCGACCGTGTGGTATTGTACCCGCGTCTC  
AGGTGTGCGGTCCAGTGTATTGCTTACCCCGAGCCCTGTTGTGGTGGGGACGACCGA  
TCGGTTTGGTGTCCCCACGTATAACTGGGGGGCGAACGACTCGGATGTGCTGATTCTC  
AACAACACGCGGCCCGCCGCGAGGCAACTGGTTCGGCTGTACATGGATGAATGGCACT  
GGTTTACCAAGACGTGTGGGGGCCCCCGTGCAACATCGGGGGGGCCGGCAACAAC  
ACCTTGACCTGCCCCACTGACTGTTTTCGGAAGCACCCCGAGGCCACCTACGCCAGAT  
GCGGTTCTGGGCCCTGGCTGACACCTAGGTGTATGGTTCATTACCCATATAGGCTCTGG  
CACTACCCCTGCACTGTCAACTTCACCATCTTCAAGGTTAGGATGTACGTGGGGGGCG

## Fig. 21J

TGGAGCACAGGTTCGAAGCCGCATGCAATTGGACTCGAGGAGAGCGTTGTGACTTGGA  
GGACAGGGATAGATCAGAGCTTAGCCCGCTGCTGCTGTCTACAACAGAGTGGCAGATA  
CTGCCCTGTTCTTACCACCCTGCCGGCCCTATCCACCGGCCTGATCCACCTCCATCA  
GAACATCGTGGACGTGCAATACCTGTACGGTGTAGGGTCGGCGGTTGTCTCCCTTGTC  
ATCAAATGGGAGTATGTCCTGTTGCTCTTCTTCTCCTGGCAGACGCGCGCATCTGCGC  
CTGCTTATGGATGATGCTGCTGATAGCTCAAGCTGAGGCCGCCTTAGAGAACCTGGTG  
GTCCTCAATGCGGCGGCCGTGGCCGGGGCGCATGGCACTCTTTCCTTCTTGTGTTCTT  
CTGTGCTGCCTGGTACATCAAGGGCAGGCTGGTCCCTGGTGCGGCATACGCCTTCTAT  
GGCGTGTGGCCGCTGCTCCTGCTTCTGCTGGCCTTACCACCACGAGCTTATGCCTAGTAA

SEQ ID NO 47 (HCCI65)

AATTTGGGTAAGGTCATCGATACCCTTACATGCGGCTTCGCCGACCTCGTGGGGTACA  
TTCCGCTCGTCGGCGCCCCCTAGGGGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCG  
GGTTCTGGAGGACGGCGTGAACATGCAACAGGGAATTTGCCCGGTTGCTCTTCTCT  
ATCTTCTCTTGGCTTTGCTGTCTGCTGACCGTTCCAGCTTCCGCTTATGAAGTGCG  
CAACGTGTCCGGGATGTACCATGTACGAACGACTGCTCCAACTCAAGCATTGTGTAT  
GAGGCAGCGGACATGATCATGCACACCCCCGGGTGCGTGCCCTGCGTTCGGGAGAAC  
AACTCTTCCCGCTGCTGGGTAGCGCTCACCCCCACGCTCGCAGCTAGGAACGCCAGCG  
TCCCCACCACGACAATACGACGCCACGTGATTTGCTCGTTGGGGCGGCTGCTTTCTG  
TTCCGCTATGTACGTGGGGGACCTCTGCGGATCTGTCTTCTCGTCTCCCAGCTGTTCA  
CCATCTCGCCTCGCCGGCATGAGACGGTGCAGGACTGCAATTGCTCAATCTATCCCGG  
CCACATAACGGGTACCGTATGGCTTGGGATATGATGATGAACTGGTCGCCTACAACG  
GCCCTGGTGGTATCGCAGCTGCTCCGGATCCCACAAGCTGTCGTGGACATGGTGGCGG  
GGGCCATTGGGGAGTCTGGCGGGCCTCGCCTACTATTCCATGGTGGGGAACTGGGC  
TAAGGTTTTGGTTGTGATGCTACTCTTTGCCGGCGTCGACGGGCATACCCGCGTGTGAG  
GAGGGGCAGCAGCCTCCGATACCAGGGGCCTTGTGTCCCTCTTTAGCCCCGGGTGCGC  
TCAGAAAATCCAGCTCGTAAACACCAACGGCAGTTGGCACATCAACAGGACTGCCCT  
GAACTGCAACGACTCCCTCAAACAGGGTTCTTTGCCGCACTATTCTACAAACACAAA  
TTCAACTCGTCTGGATGCCAGAGCGCTTGGCCAGCTGTGCTCCATCGACAAGTTCG  
CTCAGGGGTGGGGTCCCCTCACTTACACTGAGCCTAACAGCTCGGACCAGAGGCCCTA  
CTGCTGGCACTACGCGCCTCGACCGTGTGGTATTGTACCCGCGTCTCAGGTGTGCGGT  
CCAGTGTATTGCTTACCCCCGAGCCCTGTTGTGGTGGGGACGACCGATCGGTTTGGTGT  
CCCCACGTATAACTGGGGGGCGAACGACTCGGATGTGCTGATTCTCAACAACACGCGG  
CCGCCGCGAGGGCAACTGGTTCCGGCTGTACATGGATGAATGGCACTGGGTTACCAAGA  
CGTGTGGGGGCCCCCCGTGCAACATCGGGGGGGCCGGCAACAACACCTTGACCTGCC

## Fig. 21K

CCACTGACTGTTTTCGGAAGCACCCCGAGGCCACCTACGCCAGATGCGGTTCTGGGCC  
CTGGCTGACACCTAGGTGTATGGTTCATTACCCATATAGGCTCTGGCACTACCCCTGCA  
CTGTCAACTTCACCATCTTCAAGGTTAGGATGTACGTGGGGGGCGTGGAGCACAGGTT  
CGAAGCCGCATGCAATTGGACTCGAGGAGAGCGTTGTGACTTGGAGGACAGGGATAG  
ATCAGAGCTTAGCCCCGTGCTGCTGTCTACAACAGAGTGGCAGATACTGCCCTGTTCC  
TTCACCACCCTGCCGGCCCTATCCACCGGCCTGATCCACCTCCATCAGAACATCGTGG  
ACGTGCAATACCTGTACGGTGTAGGGTCGGCGGTTGTCTCCCTTGTCATCAAATGGGA  
GTATGTCCTGTTGCTCTTCTTCTCCTGGCAGACGCGCGCATCTGCGCCTGCTTATGGA  
TGATGCTGCTGATAGCTCAAGCTGAGGCCGCCTTAGAGAACCTGGTGGTCTCAATGC  
GGCGGCCGTGGCCGGGGCGCATGGCACTTTTCTTCTTGTGTTCTTCTGTGCTGCCT  
GGTACATCAAGGGCAGGCTGGTCCCTGGTGGCAGATACGCCTTCTATGGCGTGTGGCC  
GCTGCTCCTGCTTCTGCTGGCCTTACCACCACGAGCTTATGCCTAGTAAGCTT

SEQ ID NO 49 (HCCI66)

ATGAGCACGAATCCTAAACCTCAAAGAAAAACCAAACGTAACACCAACCGCCGCCCA  
CAGGACGTCAAGTTCCCGGGCGGTGGTCAGATCGTTGGTGGAGTTTACCTGTTGCCGC  
GCAGGGGCCCCAGGTTGGGTGTGCGCGCGACTAGGAAGACTTCCGAGCGGTGCGAAC  
CTCGTGGGAGGCGACAACCTATCCCCAAGGCTCGCCGACCCGAGGGTAGGGCCTGGG  
CTCAGCCCGGGTACCCTTGGCCCCTCTATGGCAATGAGGGCATGGGGTGGGCAGGATG  
GCTCCTGTACCCCCGCGGCTCTCGGCCTAGTTGGGGCCCTACAGACCCCCGGCGTAGG  
TCGCGTAATTTGGGTAAGGTCATCGATACCCTTACATGCGGCTTCGCCGACCTCGTGG  
GGTACATTCCGCTCGTGGCGCCCCCTAGGGGGCGCTGCCAGGGCCCTGGCGCATGG  
CGTCCGGGTTCTGGAGGACGGCGTGAACATGCAACAGGGAATTTGCCCGGTTGCTCT  
TTCTCTATCTTCTTCTTGGCTTTGCTGTCCTGTCTGACCGTTCCAGCTTCCGCTTATGAA  
GTGCGCAACGTGTCCGGGATGTACCATGTCACGAACGACTGCTCCAACCAAGCATTG  
TGTATGAGGCAGCGGACATGATCATGCACACCCCCGGGTGCGTGCCCTGCGTTCGGGA  
GAACAACCTTCCCGCTGCTGGGTAGCGCTCACCCCCACGCTCGCAGCTAGGAACGCC  
AGCGTCCCCACCACGACAATACGACGCCACGTCGATTTGCTCGTTGGGGCGGCTGCTT  
TCTGTTCCGCTATGTACGTGGGGGACCTCTGCGGATCTGTCTTCTCGTCTCCAGCTG  
TTCACCATCTCGCCTCGCCGGCATGAGACGGTGCAGGACTGCAATTGCTCAATCTATC  
CCGGCCACATAACGGGTCACCGTATGGCTTGGGATATGATGATGAACTGGTCGCCTAC  
AACGGCCCTGGTGGTATCGCAGCTGCTCCGATCCCACAAGCTGTGCTGGACATGGTG  
GCGGGGGCCATTGGGGAGTCTGGCGGGCCTCGCCTACTATTCCATGGTGGGGAAC  
GGGCTAAGGTTTTGGTTGTGATGCTACTCTTTGCCGGCGTGCACGGGCATACCCGCGT  
GTCAGGAGGGGCAGCAGCCTCCGATACCAGGGCCTTGTGTCCCTCTTACCCCGGG




## Fig. 21L

TCGGCTCAGAAAATCCAGCTCGTAAACACCAACGGCAGTTGGCACATCAACAGGACT  
GCCCTGAACTGCAACGACTCCCTCCAAACAGGGTTCTTTGCCGCACTATTCTACAAAC  
ACAAATTCAACTCGTCTGGATGCCAGAGCGCTTGGCCAGCTGTCGCTCCATCGACAA  
GTTCGCTCAGGGGTGGGGTCCCCTCACTTACACTGAGCCTAACAGCTCGGACCAGAGG  
CCCTACTGCTGGCACTACGCGCCTCGACCGTGTGGTATTGTACCCGCGTCTCAGGTGT  
GCGGTCCAGTGTATTGCTTCACCCCGAGCCCTGTTGTGGTGGGGACGACCGATCGGTT  
TGGTGTCCCCACGTATAACTGGGGGGCGAACGACTCGGATGTGCTGATTCTCAACAAC  
ACGCGGCCGCGCGAGGCAACTGGTTCGGCTGTACATGGATGAATGGCACTGGGTTCA  
CCAAGACGTGTGGGGGCCCCCGTGCAACATCGGGGGGGCCGGCAACAACACCTTGA  
CCTGCCCCACTGACTGTTTTCGGAAGCACCCCGAGGCCACCTACGCCAGATGCGGTTT  
TGGGCCCTGGCTGACACCTAGGTGTATGGTTCATTACCCATATAGGCTCTGGCACTAC  
CCCTGCACTGTCAACTTCACCATCTTCAAGGTTAGGATGTACGTGGGGGGCGTGGAGC  
ACAGGTTTGAAGCCGCATGCAATTGGACTCGAGGAGAGCGTTGTGACTTGGAGGACA  
GGGATAGATCAGAGCTTAGCCCGCTGCTGCTGTCTACAACAGAGTGGCAGATACTGCC  
CTGTTCCCTTACCACCCTGCCGGCCCTATCCACCGGCCTGATCCACCTCCATCAGAAC  
ATCGTGGACGTGCAATACCTGTACGGTGTAGGGTCGGCGGTTGTCTCCCTTGTCAATCA  
AATGGGAGTATGTCCTGTTGCTCTTCCCTTCTCCTGGCAGACGCGCGCATCTGCGCCTGC  
TTATGGATGATGCTGCTGATAGCTCAAGCTGAGGCCGCCTTAGAGAACCTGGTGGTCC  
TCAATGCGGCGGCCGTGGCCGGGGCGCATGGCACTCTTTCCTTCTTGTGTTCTTCTGT  
GCTGCCTGGTACATCAAGGGCAGGCTGGTCCCTGGTGCGGCATACGCCTTCTATGGCG  
TGTGGCCGCTGCTCCTGCTTCTGCTGGCCTTACCACCACGAGCTTATGCCTAGTAA

Fig. 22

OD measured at 450 nm  
construct

Fraction	volume	dilution	39 Type 1b	40 Type 1b	62 Type 3a	63 Type 5a
START	23 ml	1/20	2.517	1.954	1.426	1.142
FLOW THROUGH	23 ml	1/20	0.087	0.085	0.176	0.120
1	0.4 ml	1/200	0.102	0.051	0.048	0.050
2			0.396	0.550	0.090	0.067
3			2.627	2.603	2.481	2.372
4			3	2.967	3	2.694
5			3	2.810	2.640	2.154
6			2.694	2.499	1.359	1.561
7			2.408	2.481	0.347	1.390
8			2.176	1.970	1.624	0.865
9			1.461	1.422	0.887	0.604
10			1.286	0.926	0.543	0.519
11			0.981	0.781	0.294	0.294
12			0.812	0.650	0.249	0.199
13			0.373	0.432	0.239	0.209
14			0.653	0.371	0.145	0.184
15			0.441	0.348	0.151	0.151
16			0.321	0.374	0.098	0.106
17			0.525	0.186	0.099	0.108
18			0.351	0.171	0.083	0.090
19			0.192	0.164	0.084	0.087



FRACIES  
Fig. 23

Fig. 24

Fraction	volume	dilution	OD measured at 450 nm construct			
			39 Type 1b	40 Type 1b	62 Type 3a	63 Type 5a
20	250 $\mu$ l	1/200	0.072	0.130	0.096	0.051
21			0.109	0.293	0.084	0.052
22			0.279	0.249	0.172	0.052
23			0.093	0.151	0.297	0.054
24			0.080	0.266	0.438	0.056
25			0.251	0.100	0.457	0.048
26			3	1.649	0.722	0.066
27			3	3	2.528	0.889
28			3	3	3	2.345
29			3	3	2.849	2.580
30			2.227	1.921	1.424	1.333
31			0.263	0.415	0.356	0.162
32			0.071	0.172	0.154	0.064
33			0.103	0.054	0.096	0.057
34			0.045	0.045	0.044	0.051
35			0.043	0.047	0.045	0.046
36			0.045	0.045	0.049	0.040
37			0.045	0.047	0.046	0.048
38			0.046	0.048	0.047	0.057
39			0.045	0.048	0.050	0.057
40			0.046	0.049	0.048	0.049




Fig. 25




Fig. 26




Fig. 27




Fig.28




Fig.29

Lane 1: Crude Lysate  
Lane 2: Flow through Lentil Chromatography  
Lane 3: Wash with EMPiGEN Lentil Chromatography  
Lane 4: Eluate Lentil Chromatography  
Lane 5: Flow through during concentration lentil eluate  
Lane 6: Pool of EI after Size Exclusion Chromatography






Fig. 30



NON - REDUCED **Fig. 31A**  
E2 + CONTAMINANTS (AGGREGATES)



**REDUCED**



**Fig. 31B**






Fig. 32

### SILVER STAIN OF PURIFIED E2



1. 30 mM IMIDAZOLE WASH Ni-IMAC
2. 0.5 ug E2

Fig. 33



No.	Ret. (ml)	Peak start (ml)	Peak end (ml)	Dur (ml)	Area (ml*mAU)	Height (mAU)
1	-0.45	-0.46	-0.43	0.04	0.0976	4.579
2	1.55	0.75	3.26	2.51	796.4167	889.377
3	3.27	3.26	3.31	0.05	0.0067	0.224
4	3.33	3.32	3.33	0.02	0.0002	0.018

Total number of detected peaks = 4  
 Total Area above baseline = 0.796522 ml\*AU  
 Total area in evaluated peaks = 0.796521 ml\*AU  
 Ratio peak area / total area = 0.999999  
 Total peak duration = 2.613583 ml

Fig. 34




Fig. 35A-1




Fig. 35A-2




Fig. 35A-3




Fig. 35A-4




Fig. 35A-5




Fig. 35A-6




Fig. 35A-7




Fig. 35A-8




Fig. 35B-1




Fig. 35B-2




Fig. 35B-3




Fig. 35B-4




Fig. 35B-5




Fig. 35B-6




Fig. 35B-7




Fig. 35B-8



Fig. 36A

E1 Ab






Fig. 36B


E2 Ab




**Fig. 37B**  
Long Term Responders




**Fig. 37D**  
Type 3a



**Fig. 37A**  
Non Responders




**Fig. 37C**  
Type 1b




# Fig. 38

Relative Map Positions of  
anti-E2 monoclonal antibodies



**PARTIAL DEGLYCOSYLATION  
OF HCV E1 ENVELOPE PROTEIN**

Fig.39



**PARTIAL TREATMENT OF HCV E2\E2s ENVELOPE PROTEINS  
BY PNGase F**




Fig. 40

Fig. 41 *In Vitro* Mutagenesis of HCV E1 glycoprotein






Fig. 42A *In Vitro* Mutagenesis of HCV E1 glycoprotein




1. First step of PCR amplification (Gly-# and Ovr-# primers)



2. Overlap extension and nested PCR

Fig. 42B

a. Overlap extension



b. Nested PCR amplification (GPT-2 and TKr-2 primers)







Fig. 43 *In Vitro* Mutagenesis of HCV E1 glycoprotein






Fig. 44A




Fig. 44B




Fig. 45




Fig. 46

**PURIFIED HEPATITIS C VIRUS ENVELOPE  
PROTEINS FOR DIAGNOSTIC AND  
THERAPEUTIC USE**

**FIELD OF THE INVENTION**

The present invention relates to the general fields of recombinant protein expression, purification of recombinant proteins, synthetic peptides, diagnosis of HCV infection, prophylactic treatment against HCV infection and to the prognosis/monitoring of the clinical efficiency of treatment of an individual with chronic hepatitis, or the prognosis/monitoring of natural disease.

More particularly, the present invention relates to purification methods for hepatitis C virus envelope proteins, the use in diagnosis, prophylaxis or therapy of HCV envelope proteins purified according to the methods described in the present invention, the use of single or specific oligomeric E1 and/or E2 and/or E1/E2 envelope proteins in assays for monitoring disease, and/or diagnosis of disease, and/or treatment of disease. The invention also relates to epitopes of the E1 and/or E2 envelope proteins and monoclonal antibodies thereto, as well their use in diagnosis, prophylaxis or treatment.

**BACKGROUND OF THE INVENTION**

The E2 protein purified from cell lysates according to the methods described in the present invention reacts with approximately 95% of patient sera. This reactivity is similar to the reactivity obtained with E2 secreted from CHO cells (Spaete et al., 1992). However, the intracellularly expressed form of E2 may more closely resemble the native viral envelope protein because it contains high mannose carbohydrate motifs, whereas the E2 protein secreted from CHO cells is further modified with galactose and sialic acid sugar moieties. When the aminoterminal half of E2 is expressed in the baculovirus system, only about 13 to 21% of sera from several patient groups can be detected (Inoue et al., 1992). After expression of E2 from *E. coli*, the reactivity of HCV sera was even lower and ranged from 14 (Yokosuka et al., 1992) to 17% (Mita et al., 1992).

About 75% of HCV sera (and 95% of chronic patients) are anti-E1 positive using the purified, vaccinia-expressed recombinant E1 protein of the present invention, in sharp contrast with the results of Kohara et al. (1992) and Hsu et al. (1993). Kohara et al. used a vaccinia-virus expressed E1 protein and detected anti-E1 antibodies in 7 to 23% of patients, while Hsu et al. only detected 14/50 (28%) sera using baculovirus-expressed E1.

These results show that not only a good expression system but also a good purification protocol are required to reach a high reactivity of the envelope proteins with human patient sera. This can be obtained using the proper expression system and/or purification protocols of the present invention which guarantee the conservation of the natural folding of the protein and the purification protocols of the present invention which guarantee the elimination of contaminating proteins and which preserve the conformation, and thus the reactivity of the HCV envelope proteins. The amounts of purified HCV envelope protein needed for diagnostic screening assays are in the range of grams per year. For vaccine purposes, even higher amounts of envelope protein would be needed. Therefore, the vaccinia virus system may be used for selecting the best expression constructs and for limited upscaling, and large-scale expression and purification of single or specific oligomeric envelope proteins containing high-mannose carbohydrates may be achieved

when expressed from several yeast strains. In the case of hepatitis B for example, manufacturing of HBsAg from mammalian cells was much more costly compared with yeast-derived hepatitis B vaccines.

**Aims of the Invention**

It is an aim of the present invention to provide a new purification method for recombinantly expressed E1 and/or E2 and/or E1/E2 proteins such that said recombinant proteins are directly usable for diagnostic and vaccine purposes as single or specific oligomeric recombinant proteins free from contaminants instead of aggregates.

It is another aim of the present invention to provide compositions comprising purified (single or specific oligomeric) recombinant E1 and/or E2 and/or E1/E2 glycoproteins comprising conformational epitopes from the E1 and/or E2 domains of HCV.

It is yet another aim of the present invention to provide novel recombinant vector constructs for recombinantly expressing E1 and/or E2 and/or E1/E2 proteins, as well as host cells transformed with said vector constructs.

It is also an aim of the present invention to provide a method for producing and purifying recombinant HCV E1 and/or E2 and/or E1/E2 proteins.

It is also an aim of the present invention to provide diagnostic and immunogenic uses of the recombinant HCV E1 and/or E2 and/or E1/E2 proteins of the present invention, as well as to provide kits for diagnostic use, vaccines or therapeutics comprising any of the recombinant HCV E1 and/or E2 and/or E1/E2 proteins of the present invention.

It is further an aim of the present invention to provide for a new use of E1, E2, and/or E1/E2 proteins, or suitable parts thereof, for monitoring/prognosing the response to treatment of patients (e.g. with interferon) suffering from HCV infection.

It is also an aim of the present invention to provide for the use of the recombinant E1, E2, and/or E1/E2 proteins of the present invention in HCV screening and confirmatory antibody tests.

It is also an aim of the present invention to provide E1 and/or E2 peptides which can be used for diagnosis of HCV infection and for raising antibodies. Such peptides may also be used to isolate human monoclonal antibodies.

It is also an aim of the present invention to provide monoclonal antibodies, more particularly human monoclonal antibodies or mouse monoclonal antibodies which are humanized, which react specifically with E1 and/or E2 epitopes, either comprised in peptides or conformational epitopes comprised in recombinant proteins.

It is also an aim of the present invention to provide possible uses of anti-E1 or anti-E2 monoclonal antibodies for HCV antigen detection or for therapy of chronic HCV infection.

It is also an aim of the present invention to provide kits for monitoring/prognosing the response to treatment (e.g. with interferon) of patients suffering from HCV infection or monitoring/prognosing the outcome of the disease.

All the aims of the present invention are considered to have been met by the embodiments as set out below.

**Definitions**

The following definitions serve to illustrate the different terms and expressions used in the present invention.

The term 'hepatitis C virus single envelope protein' refers to a polypeptide or an analogue thereof (e.g. mimotopes) comprising an amino acid sequence (and/or amino acid

analogues) defining at least one HCV epitope of either the E1 or the E2 region. These single envelope proteins in the broad sense of the word may be both monomeric or homo-oligomeric forms of recombinantly expressed envelope proteins. Typically, the sequences defining the epitope correspond to the amino acid sequence of either the E1 or the E2 region of HCV (either identically or via substitution of analogues of the native amino acid residue that do not destroy the epitope). In general, the epitope-defining sequence will be 3 or more amino acids in length, more typically, 5 or more amino acids in length, more typically 8 or more amino acids in length, and even more typically 10 or more amino acids in length. With respect to conformational epitopes, the length of the epitope-defining sequence can be subject to wide variations, since it is believed that these epitopes are formed by the three-dimensional shape of the antigen (e.g. folding). Thus, the amino acids defining the epitope can be relatively few in number, but widely dispersed along the length of the molecule being brought into the correct epitope conformation via folding. The portions of the antigen between the residues defining the epitope may not be critical to the conformational structure of the epitope. For example, deletion or substitution of these intervening sequences may not affect the conformational epitope provided sequences critical to epitope conformation are maintained (e.g. cysteines involved in disulfide bonding, glycosylation sites, etc.). A conformational epitope may also be formed by 2 or more essential regions of subunits of a homooligomer or heterooligomer.

The HCV antigens of the present invention comprise conformational epitopes from the E1 and/or E2 (envelope) domains of HCV. The E1 domain, which is believed to correspond to the viral envelope protein, is currently estimated to span amino acids 192–383 of the HCV polyprotein (Hijikata et al., 1991). Upon expression in a mammalian system (glycosylated), it is believed to have an approximate molecular weight of 35 kDa as determined via SDS-PAGE. The E2 protein, previously called NS1, is believed to span amino acids 384–809 or 384–746 (Grakoui et al., 1993) of the HCV polyprotein and to also be an envelope protein. Upon expression in a vaccinia system (glycosylated), it is believed to have an apparent gel molecular weight of about 72 kDa. It is understood that these protein endpoints are approximations (e.g. the carboxy terminal end of E2 could lie somewhere in the 730–820 amino acid region, e.g. ending at amino acid 730, 735, 740, 742, 744, 745, preferably 746, 747, 748, 750, 760, 770, 780, 790, 800, 809, 810, 820). The E2 protein may also be expressed together with the E1, P7 (aa 747–809), NS2 (aa 810–1026), NS4A (aa 1658–1711) or NS4B (aa 1712–1972). Expression together with these other HCV proteins may be important for obtaining the correct protein folding.

It is also understood that the isolates used in the examples section of the present invention were not intended to limit the scope of the invention and that any HCV isolate from type 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or any other new genotype of HCV is a suitable source of E1 and/or E2 sequence for the practice of the present invention.

The E1 and E2 antigens used in the present invention may be full-length viral proteins, substantially full-length versions thereof, or functional fragments thereof (e.g. fragments which are not missing sequence essential to the formation or retention of an epitope). Furthermore, the HCV antigens of the present invention can also include other sequences that do not block or prevent the formation of the conformational epitope of interest. The presence or absence of a conformational epitope can be readily determined

though screening the antigen of interest with an antibody (polyclonal serum or monoclonal to the conformational epitope) and comparing its reactivity to that of a denatured version of the antigen which retains only linear epitopes (if any). In such screening using polygonal antibodies, it may be advantageous to adsorb the polygonal serum first with the denatured antigen and see if it retains antibodies to the antigen of interest.

The HCV antigens of the present invention can be made by any recombinant method that provides the epitope of interest. For example, recombinant intracellular expression in mammalian or insect cells is a preferred method to provide glycosylated E1 and/or E2 antigens in 'native' conformation as is the case for the natural HCV antigens. Yeast cells and mutant yeast strains (e.g. mnn 9 mutant (Kniskern et al., 1994) or glycosylation mutants derived by means of vanadate resistance selection (Ballou et al., 1991)) may be ideally suited for production of secreted high-mannose-type sugars; whereas proteins secreted from mammalian cells may contain modifications including galactose or sialic acids which may be undesirable for certain diagnostic or vaccine applications. However, it may also be possible and sufficient for certain applications, as it is known for proteins, to express the antigen in other recombinant hosts (such as *E. coli*) and renature the protein after recovery.

The term 'fusion polypeptide' intends a polypeptide in which the HCV antigen(s) are part of a single continuous chain of amino acids, which chain does not occur in nature. The HCV antigens may be connected directly to each other by peptide bonds or be separated by intervening amino acid sequences. The fusion polypeptides may also contain amino acid sequences exogenous to HCV.

The term 'solid phase' intends a solid body to which the individual HCV antigens or the fusion polypeptide comprised of HCV antigens are bound covalently or by noncovalent means such as hydrophobic adsorption.

The term 'biological sample' intends a fluid or tissue of a mammalian individual (e.g. an anthropoid, a human) that commonly contains antibodies produced by the individual, more particularly antibodies against HCV. The fluid or tissue may also contain HCV antigen. Such components are known in the art and include, without limitation, blood, plasma, serum, urine, spinal fluid, lymph fluid, secretions of the respiratory, intestinal or genitourinary tracts, tears, saliva, milk, white blood cells and myelomas. Body components include biological liquids. The term 'biological liquid' refers to a fluid obtained from an organism. Some biological fluids are used as a source of other products, such as clotting factors (e.g. Factor VIII;C), serum albumin, growth hormone and the like. In such cases, it is important that the source of biological fluid be free of contamination by virus such as HCV.

The term 'immunologically reactive' means that the antigen in question will react specifically with anti-HCV antibodies present in a body component from an HCV infected individual.

The term 'immune complex' intends the combination formed when an antibody binds to an epitope on an antigen.

'E1' as used herein refers to a protein or polypeptide expressed within the first 400 amino acids of an HCV polyprotein, sometimes referred to as the E, ENV or S protein. In its natural form it is a 35 kDa glycoprotein which is found in strong association with membranes. In most natural HCV strains, the E1 protein is encoded in the viral polyprotein following the C (core) protein. The E1 protein extends from approximately amino acid (aa) 192 to about aa 383 of the full-length polyprotein.

The term 'E1' as used herein also includes analogs and truncated forms that are immunologically cross-reactive with natural E1, and includes E1 proteins of genotypes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or any other newly identified HCV type or subtype.

'E2' as used herein refers to a protein or polypeptide expressed within the first 900 amino acids of an HCV polyprotein, sometimes referred to as the NS1 protein. In its natural form it is a 72 kDa glycoprotein that is found in strong association with membranes. In most natural HCV strains, the E2 protein is encoded in the viral polyprotein following the E1 protein. The E2 protein extends from approximately amino acid position 384 to amino acid position 746, another form of E2 extends to amino acid position 809. The term 'E2' as used herein also includes analogs and truncated forms that are immunologically cross-reactive with natural E2. For example, insertions of multiple codons between codon 383 and 384, as well as deletions of amino acids 384-387 have been reported by Kato et al. (1992).

'E1/E2' as used herein refers to an oligomeric form of envelope proteins containing at least one E1 component and at least one E2 component.

The term 'specific oligomeric' E1 and/or E2 and/or E1/E2 envelope proteins refers to all possible oligomeric forms of recombinantly expressed E1 and/or E2 envelope proteins which are not aggregates. E1 and/or E2 specific oligomeric envelope proteins are also referred to as homo-oligomeric E1 or E2 envelope proteins (see below).

The term 'single or specific oligomeric' E1 and/or E2 and/or E1 E2 envelope proteins refers to single monomeric E1 or E2 proteins (single in the strict sense of the word) as well as specific oligomeric E1 and/or E2 and/or E1/E2 recombinantly expressed proteins. These single or specific oligomeric envelope proteins according to the present invention can be further defined by the following formula (E1), (E2), wherein x can be a number between 0 and 100, and y can be a number between 0 and 100, provided that x and y are not both 0. With x=1 and y=0 said envelope proteins include monomeric E1.

The term 'homo-oligomer' as used herein refers to a complex of E1 and/or E2 containing more than one E1 or E2 monomer, e.g. E1/E1 dimers, E1/E1/E1 trimers or E1/E1/E1/E1 tetramers and E2/E2 dimers, E2/E2/E2 trimers or E2/E2/E2/E2 tetramers, E1 pentamers and hexamers, E2 pentamers and hexamers or any higher-order homo-oligomers of E1 or E2 are all 'homo-oligomers' within the scope of this definition. The oligomers may contain one, two, or several different monomers of E1 or E2 obtained from different types or subtypes of hepatitis C virus including for example those described in an international application published under WO 94/25601 and European application No. 94870166.9 both by the present applicants. Such mixed oligomers are still homo-oligomers within the scope of this invention, and may allow more universal diagnosis, prophylaxis or treatment of HCV.

The term 'purified' as applied to proteins herein refers to a composition wherein the desired protein comprises at least 35% of the total protein component in the composition. The desired protein preferably comprises at least 40%, more preferably at least about 50%, more preferably at least about 60%, still more preferably at least about 70%, even more preferably at least about 80%, even more preferably at least about 90%, and most preferably at least about 95% of the total protein component. The composition may contain other compounds such as carbohydrates, salts, lipids, solvents, and the like, without affecting the determination of the percent-

age purity as used herein. An 'isolated' HCV protein intends an HCV protein composition that is at least 35% pure.

The term 'essentially purified proteins' refers to proteins purified such that they can be used for in vitro diagnostic methods and as a therapeutic compound. These proteins are substantially free from cellular proteins, vector-derived proteins or other HCV viral components. Usually these proteins are purified to homogeneity (at least 80% pure, preferably, 90%, more preferably 95%, more preferably 97%, more preferably 98%, more preferably 99%, even more preferably 99.5%, and most preferably the contaminating proteins should be undetectable by conventional methods like SDS-PAGE and silver staining.

The term 'recombinantly expressed' used within the context of the present invention refers to the fact that the proteins of the present invention are produced by recombinant expression methods be it in prokaryotes, or lower or higher eukaryotes as discussed in detail below.

The term 'lower eukaryote' refers to host cells such as yeast, fungi and the like. Lower eukaryotes are generally (but not necessarily) unicellular. Preferred lower eukaryotes are yeasts, particularly species within *Saccharomyces*, *Schizosaccharomyces*, *Kluveromyces*, *Pichia* (e.g. *Pichia pastoris*), *Hansenula* (e.g. *Hansenula polymorpha*, *Yarrowia*, *Schwaniomyces*, *Schizosaccharomyces*, *Zygosaccharomyces* and the like. *Saccharomyces cerevisiae*, *S. carlsbergensis* and *K. lactis* are the most commonly used yeast hosts, and are convenient fungal hosts.

The term 'prokaryotes' refers to hosts such as *E.coli*, *Lactobacillus*, *Lactococcus*, *Salmonella*, *Streptococcus*, *Bacillus subtilis* or *Streptomyces*. Also these hosts are contemplated within the present invention.

The term 'higher eukaryote' refers to host cells derived from higher animals, such as mammals, reptiles, insects, and the like. Presently preferred higher eukaryote host cells are derived from Chinese hamster (e.g. CHO), monkey (e.g. COS and Vero cells), baby hamster kidney (BHK), pig kidney (PK15), rabbit kidney 13 cells (RK13), the human osteosarcoma cell line 143 B, the human cell line HeLa and human hepatoma cell lines like Hep G2, and insect cell lines (e.g. *Spodoptera frugiperda*). The host cells may be provided in suspension or flask cultures, tissue cultures, organ cultures and the like. Alternatively the host cells may also be transgenic animals.

The term 'polypeptide' refers to a polymer of amino acids and does not refer to a specific length of the product; thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide. This term also does not refer to or exclude post-expression modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like. Included within the definition are, for example, polypeptides containing one or more analogues of an amino acid (including, for example, unnatural amino acids, PNA, etc.), polypeptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring.

The term 'recombinant polynucleotide or nucleic acid' intends a polynucleotide or nucleic acid of genomic, cDNA, semisynthetic, or synthetic origin which, by virtue of its origin or manipulation : (1) is not associated with all or a portion of a polynucleotide with which it is associated in nature, (2) is linked to a polynucleotide other than that to which it is linked in nature, or (3) does not occur in nature.

The term 'recombinant host cells', 'host cells', 'cells', 'cell lines', 'cell cultures', and other such terms denoting microorganisms or higher eukaryotic cell lines cultured as

unicellular entities refer to cells which can be or have been, used as recipients for a recombinant vector or other transfer polynucleotide, and include the progeny of the original cell which has been transfected. It is understood that the progeny of a single parental cell may not necessarily be completely identical in morphology or in genomic or total DNA complement as the original parent, due to natural, accidental, or deliberate mutation.

The term 'replicon' is any genetic element, e.g., a plasmid, a chromosome, a virus, a cosmid, etc., that behaves as an autonomous unit of polynucleotide replication within a cell; i.e., capable of replication under its own control.

The term 'vector' is a replicon further comprising sequences providing replication and/or expression of a desired open reading frame.

The term 'control sequence' refers to polynucleotide sequences which are necessary to effect the expression of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences generally include promoter, ribosomal binding site, and terminators; in eukaryotes, generally, such control sequences include promoters, terminators and, in some instances, enhancers. The term 'control sequences' is intended to include, at a minimum, all components whose presence is necessary for expression, and may also include additional components whose presence is advantageous, for example, leader sequences which govern secretion.

The term 'promoter' is a nucleotide sequence which is comprised of consensus sequences which allow the binding of RNA polymerase to the DNA template in a manner such that mRNA production initiates at the normal transcription initiation site for the adjacent structural gene.

The expression 'operably linked' refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. A control sequence 'operably linked' to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

An 'open reading frame' (ORF) is a region of a polynucleotide sequence which encodes a polypeptide and does not contain stop codons; this region may represent a portion of a coding sequence or a total coding sequence.

A 'coding sequence' is a polynucleotide sequence which is transcribed into mRNA and/or translated into a polypeptide when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a translation start codon at the 5'-terminus and a translation stop codon at the 3'-terminus. A coding sequence can include but is not limited to mRNA, DNA (including cDNA), and recombinant polynucleotide sequences.

As used herein, 'epitope' or 'antigenic determinant' means an amino acid sequence that is immunoreactive. Generally an epitope consists of at least 3 to 4 amino acids, and more usually, consists of at least 5 or 6 amino acids, sometimes the epitope consists of about 7 to 8, or even about 10 amino acids. As used herein, an epitope of a designated polypeptide denotes epitopes with the same amino acid sequence as the epitope in the designated polypeptide, and immunologic equivalents thereof. Such equivalents also include strain, subtype (=genotype), or type(group)-specific variants, e.g. of the currently known sequences or strains belonging to genotypes 1a, 1b, 1c, 1d, 1e, 1f, 2a, 2b, 2c, 2d, 2e, 2f, 2g, 2h, 2i, 3a, 3b, 3c, 3d, 3e, 3f, 3g, 4a, 4b, 4c, 4d,

4e, 4f, 4g, 4h, 4i, 4j, 4k, 4l, 5a, 5b, 6a, 6b, 6c, 7a, 7b, 7c, 8a, 8b, 9a, 9b, 10a, or any other newly defined HCV (sub)type. It is to be understood that the amino acids constituting the epitope need not be part of a linear sequence, but may be interspersed by any number of amino acids, thus forming a conformational epitope.

The term 'immunogenic' refers to the ability of a substance to cause a humoral and/or cellular response, whether alone or when linked to a carrier, in the presence or absence of an adjuvant. 'Neutralization' refers to an immune response that blocks the infectivity, either partially or fully, of an infectious agent. A 'vaccine' is an immunogenic composition capable of eliciting protection against HCV, whether partial or complete. A vaccine may also be useful for treatment of an individual, in which case it is called a therapeutic vaccine.

The term 'therapeutic' refers to a composition capable of treating HCV infection.

The term 'effective amount' refers to an amount of epitope-bearing polypeptide sufficient to induce an immunogenic response in the individual to which it is administered, or to otherwise detectably immunoreact in its intended system (e.g., immunoassay). Preferably, the effective amount is sufficient to effect treatment, as defined above. The exact amount necessary will vary according to the application. For vaccine applications or for the generation of polyclonal antiserum/antibodies, for example, the effective amount may vary depending on the species, age, and general condition of the individual, the severity of the condition being treated, the particular polypeptide selected and its mode of administration, etc. It is also believed that effective amounts will be found within a relatively large, non-critical range. An appropriate effective amount can be readily determined using only routine experimentation. Preferred ranges of E1 and/or E2 and/or E1/E2 single or specific oligomeric envelope proteins for prophylaxis of HCV disease are 0.01 to 100  $\mu\text{g}/\text{dose}$ , preferably 0.1 to 50  $\mu\text{g}/\text{dose}$ . Several doses may be needed per individual in order to achieve a sufficient immune response and subsequent protection against HCV disease.

#### DETAILED DESCRIPTION OF THE INVENTION

More particularly, the present invention contemplates a method for isolating or purifying recombinant HCV single or specific oligomeric envelope protein selected from the group consisting of E1 and/or E2 and/or E1/E2, characterized in that upon lysing the transformed host cells to isolate the recombinantly expressed protein a disulphide bond cleavage or reduction step is carried out with a disulphide bond cleaving agent.

The essence of these 'single or specific oligomeric' envelope proteins of The invention is that they are free from contaminating proteins and that they are not disulphide bond linked with contaminants.

The proteins according to the present invention are recombinantly expressed in lower or higher eukaryotic cells or in prokaryotes. The recombinant proteins of the present invention are preferably glycosylated and may contain high-mannose-type, hybrid, or complex glycosylations. Preferentially said proteins are expressed from mammalian cell lines as discussed in detail in the Examples section, or in yeast such as in mutant yeast strains also as detailed in the Examples section.

The proteins according to the present invention may be secreted or expressed within components of the cell, such as

the ER or the Golgi Apparatus. Preferably, however, the proteins of the present invention bear high-mannose-type glycosylations and are retained in the ER or Golgi Apparatus of mammalian cells or are retained in or secreted from yeast cells, preferably secreted from yeast mutant strains such as the *mnn9* mutant (Kniskern et al., 1994), or from mutants that have been selected by means of vanadate resistance (Ballou et al., 1991).

Upon expression of HCV envelope proteins, the present inventors could show that some of the free thiol groups of cysteines not involved in intra- or inter-molecular disulphide bridges, react with cysteines of host or expression-system-derived (e.g. vaccinia) proteins or of other HCV envelope proteins (single or oligomeric), and form aspecific intermolecular bridges. This results in the formation of 'aggregates' of HCV envelope proteins together with contaminating proteins. It was also shown in WO 92/08734 that 'aggregates' were obtained after purification, but it was not described which protein interactions were involved. In patent application WO 92/08734, recombinant E1/E2 protein expressed with the vaccinia virus system were partially purified as aggregates and only found to be 70% pure, rendering the purified aggregates not useful for diagnostic, prophylactic or therapeutic purposes.

Therefore, a major aim of the present invention resides in the separation of single or specific-oligomeric HCV envelope proteins from contaminating proteins, and to use the purified proteins (>95% pure) for diagnostic, prophylactic and therapeutic purposes. To those purposes, the present inventors have been able to provide evidence that aggregated protein complexes ('aggregates') are formed on the basis of disulphide bridges and non-covalent protein-protein interactions. The present invention thus provides a means for selectively cleaving the disulphide bonds under specific conditions and for separating the cleaved proteins from contaminating proteins which greatly interfere with diagnostic, prophylactic and therapeutic applications. The free thiol groups may be blocked (reversibly or irreversibly) in order to prevent the reformation of disulphide bridges, or may be left to oxidize and oligomerize with other envelope proteins (see definition homo-oligomer). It is to be understood that such protein oligomers are essentially different from the 'aggregates' described in WO 92/08734 and WO 94/01778, since the level of contaminating proteins is undetectable.

Said disulphide bond cleavage may also be achieved by:

(1) performic acid oxidation by means of cysteic acid in which case the cysteine residues are modified into cysteic acid (Moore et al., 1963).

(2) Sulfitolysis ( $R-S-S-R \rightarrow 2R-SO_3^-$ ) for example by means of sulphite ( $SO_3^{2-}$ ) together with a proper oxidant such as  $Cu^{2+}$  in which case the cysteine is modified into S-sulpho-cysteine (Bailey and Cole, 1959).

(3) Reduction by means of mercaptans, such as dithiothreitol (DDT),  $\beta$ -mercapto-ethanol, cysteine, glutathione Red, E-mercapto-ethylamine, or thioglycolic acid, of which DTT and  $\beta$ -mercapto-ethanol are commonly used (Cleland, 1964), is the preferred method of this invention because the method can be performed in a water environment and because the cysteine remains unmodified.

(4) Reduction by means of a phosphine (e.g.  $Bu_2P$ ) (Ruegg and Rudinger, 1977).

All these compounds are thus to be regarded as agents or means for cleaving disulphide bonds according to the present invention.

Said disulphide bond cleavage (or reducing) step of the present invention is preferably a partial disulphide bond

cleavage (reducing) step (carried out under partial cleavage or reducing conditions).

A preferred disulphide bond cleavage or reducing agent according to the present invention is dithiothreitol (DTT). Partial reduction is obtained by using a low concentration of said reducing agent, i.e. for DTT for example in the concentration range of about 0.1 to about 50 mM, preferably about 0.1 to about 20 mM, preferably about 0.5 to about 10 mM, preferably more than 1 mM, more than 2 mM or more than 5 mM, more preferably about 1.5 mM, about 2.0 mM, about 2.5 mM, about 5 mM or about 7.5 mM.


Said disulphide bond cleavage step may also be carried out in the presence of a suitable detergent (as an example of a means for cleaving disulphide bonds or in combination with a cleaving agent) able to dissociate the expressed proteins, such as DECYLPEG (Decyl polyethylene glycol (Kwant, the Netherlands), EMPIGEN-BB (N-Dodecyl-N,N-dimethylglycine (Allbright & Wilson, UK)), NP-40 (nonidet P40 (Pasture, Belgium)), sodium cholate, TRITON X-100 (polyethylene glycol-p-isooctylphenyl ether; octylphenoxy-polyethoxyethanol (Merck, Belgium)).

Said reduction or cleavage step (preferably a partial reduction or cleavage step) is carried out preferably in the presence of (with) a detergent. A preferred detergent according to the present invention is Empigen-BB. The amount of detergent used is preferably in the range of 1 to 10%, preferably more than 3%, more preferably about 3.5% of a detergent such as Empigen-BB.

A particularly preferred method for obtaining disulphide bond cleavage employs a combination of a classical disulphide bond cleavage agent as detailed above and a detergent (also as detailed above). As contemplated in the Examples section, the particular combination of a low concentration of DTT (1.5 to 7.5 mM) and about 3.5% of Empigen-BB is proven to be a particularly preferred combination of reducing agent and detergent for the purification of recombinantly expressed E1 and E2 proteins. Upon gel filtration chromatography, said partial reduction is shown to result in the production of possibly dimeric E1 protein and separation of this E1 protein from contaminating proteins that cause false reactivity upon use in immunoassays.

It is, however, to be understood that also any other combination of any reducing agent known in the art with any detergent or other means known in the art to make the cysteines better accessible is also within the scope of the present invention, insofar as said combination reaches the same goal of disulphide bridge cleavage as the preferred combination exemplified in the present invention.

Apart from reducing the disulphide bonds, a disulphide bond cleaving means according to the present invention may also include any disulphide bridge exchanging agents (competitive agent being either organic or proteinaceous, see for instance Creighton, 1988) known in the art which allows the following type of reaction to occur:



R1, R2: compounds of protein aggregates

R3 SH: competitive agent (organic, proteinaceous)

The term 'disulphide bridge exchanging agent' is to be interpreted as including disulphide bond reforming as well as disulphide bond blocking agents.

The present invention also relates to methods for purifying or isolating HCV single or specific oligomeric envelope proteins as set out above further including the use of any SH group blocking or binding reagent known in the art such as chosen from the following list;



Glutathion

5,5'-dithiobis-(2-nitrobenzoic acid) or bis-(3-carboxy-4-nitrophenyl)-disulphide (DTNB or Ellman's reagent) (Elmann, 1959)

N-ethylmaleimide (NEM; Benesch et al., 1956)

N-(4-dimethylamino-3,5-dinitrophenyl) maleimide or Tuppy's maleimide which provides a color to the protein

P-chloromercuribenzoate (Grassetti et al., 1969)

4-vinylpyridine (Friedman and Krull, 1969) can be liberated after reaction by acid hydrolysis

acrylonitrile, can be liberated after reaction by acid hydrolysis (Weil and Seibles, 1961)

NEM-biotin (e.g. obtained from Sigma B1267)

2,2'-dithiopyridine (Grassetti and Murray, 1967)

4,4'-dithiopyridine (Grassetti and Murray, 1967)

6,6'-dithiodinicotinic acid (DTDNA; Brown and Cunningham, 1970)

2,2'-dithiobis-(5'-nitropyridine) (DTNP; U.S. Pat. No. 3,597,160) or other dithiobis (heterocyclic derivative) compounds (Grassetti and Murray, 1969)

A survey of the publications cited shows that often different reagents for sulphhydryl groups will react with varying numbers of thiol groups of the same protein or enzyme molecule. One may conclude that this variation in reactivity of the thiol groups is due to the steric environment of these groups, such as the shape of the molecule and the surrounding groups of atoms and their charges, as well as to the size, shape and charge of the reagent molecule or ion. Frequently the presence of adequate concentrations of denaturants such as sodium dodecylsulfate, urea or guanidine hydrochloride will cause sufficient unfolding of the protein molecule to permit equal access to all of the reagents for thiol groups. By varying the concentration of denaturant, the degree of unfolding can be controlled and in this way thiol groups with different degrees of reactivity may be revealed. Although up to date most of the work reported has been done with p-chloromercuribenzoate, N-ethylmaleimide and DTNB, it is likely that the other more recently developed reagents may prove equally useful. Because of their varying structures, it seems likely, in fact, that they may respond differently to changes in the steric environment of the thiol groups.

Alternatively, conditions such as low pH (preferably lower than pH 6) for preventing free SH groups from oxidizing and thus preventing the formation of large intermolecular aggregates upon recombinant expression and purification of E1 and E2 (envelope) proteins are also within the scope of the present invention.

A preferred SH group blocking reagent according to the present invention is N-ethylmaleimide (NEM). Said SH group blocking reagent may be administered during lysis of the recombinant host cells and after the above-mentioned partial reduction process or after any other process for cleaving disulphide bridges. Said SH group blocking reagent may also be modified with any group capable of providing a detectable label and/or any group aiding in the immobilization of said recombinant protein to a solid substrate, e.g. biotinylated NEM.

Methods for cleaving cysteine bridges and blocking free cysteines have also been described in Darbre (1987), Means and Feeney (1971), and by Wong (1993).

A method to purify single or specific oligomeric recombinant E1 and/or E2 and/or E1/E2 proteins according to the present invention as defined above is further characterized as comprising the following steps:

lysing recombinant E1 and/or E2 and/or E1/E2 expressing host cells, preferably in the presence of an SH group blocking agent, such as N-ethylmaleimide (NEM), and possibly a suitable detergent, preferably Empigen-BB, recovering said HCV envelope protein by affinity purification for instance by means lectin-chromatography, such as lentil-lectin chromatography, or immunoaffinity chromatography using anti-E1 and/or anti-E2 specific monoclonal antibodies, followed by,

reduction or cleavage of disulphide bonds with a disulphide bond cleaving agent, such as DTT, preferably also in the presence of an SH group blocking agent, such as NEM or Biotin-NEM, and,

recovering the reduced HCV E1 and/or E2 and/or E1/E2 envelope proteins for instance by gel filtration (size exclusion chromatography or molecular sieving) and possibly also by an additional Ni<sup>2+</sup>-IMAC chromatography and desalting step.

It is to be understood that the above-mentioned recovery steps may also be carried out using any other suitable technique known by the person skilled in the art.

Preferred lectin-chromatography systems include *Galanthus nivalis* agglutinin (GNA)-chromatography, or *Lens culinaris* agglutinin (LCA) (lentil) lectin chromatography as illustrated in the Examples section. Other useful lectins include those recognizing high-mannose type sugars, such as *Narcissus pseudonarcissus* agglutinin (NPA), *Pisum sativum* agglutinin (PSA), or *Allium ursinum* agglutinin (AUA).

Preferably said method is usable to purify single or specific oligomeric HCV envelope protein produced intracellularly as detailed above.

For secreted E1 or E2 or E1/E2 oligomers, lectins binding complex sugars such as *Ricinus communis* agglutinin I (RCA I), are preferred lectins.

The present invention: more particularly contemplates essentially purified recombinant HCV single or specific oligomeric envelope proteins, selected from the group consisting of E1 and/or E2 and/or E1/E2, characterized as being isolated or purified by a method as defined above.

The present invention more particularly relates to the purification or isolation of recombinant envelope proteins which are expressed from recombinant mammalian cells such as vaccinia.

The present invention also relates to the purification or isolation of recombinant envelope proteins which are expressed from recombinant yeast cells.

The present invention equally relates to the purification or isolation of recombinant envelope proteins which are expressed from recombinant bacterial (prokaryotic) cells.

The present invention also contemplates a recombinant vector comprising a vector sequence, an appropriate prokaryotic, eukaryotic or viral or synthetic promoter sequence followed by a nucleotide sequence allowing the expression of the single or specific oligomeric E1 and/or E2 and/or E1/E2 of the invention.

Particularly, the present invention contemplates a recombinant vector comprising a vector sequence, an appropriate prokaryotic, eukaryotic or viral or synthetic promoter sequence followed by a nucleotide sequence allowing the expression of the single E1 or E1 of the invention.

Particularly, the present invention contemplates a recombinant vector comprising a vector sequence, an appropriate prokaryotic, eukaryotic or viral or synthetic promoter sequence followed by a nucleotide sequence allowing the expression of the single E1 or E2 of the invention.

The segment of the HCV cDNA encoding the desired E1 and/or E2 sequence inserted into the vector sequence may be

attached to a signal sequence. Said signal sequence may be that from a non-HCV source, e.g. the IgG or tissue plasminogen activator (tpa) leader sequence for expression in mammalian cells, or the  $\alpha$ -mating factor sequence for expression into yeast cells, but particularly preferred constructs according to the present invention contain signal sequences appearing in the HCV genome before the respective start points of the E1 and E2 proteins. The segment of the HCV cDNA encoding the desired E1 and/or E2 sequence inserted into the vector may also include deletions e.g. of the hydrophobic domain(s) as illustrated in the examples section, or of the E2 hypervariable region 1.

More particularly, the recombinant vectors according to the present invention encompass a nucleic acid having an HCV cDNA segment encoding the polyprotein starting in the region between amino acid positions 1 and 192 and ending in the region between positions 250 and 400 of the HCV polyprotein, more preferably ending in the region between positions 250 and 341, even more preferably ending in the region between positions 290 and 341 for expression of the HCV single E1 protein. Most preferably, the present recombinant vector encompasses a recombinant nucleic acid having a HCV cDNA segment encoding part of the HCV polyprotein starting in the region between positions 117 and 192, and ending at any position in the region between positions 263 and 326, for expression of HCV single E1 protein. Also within the scope of the present invention are forms that have the first hydrophobic domain deleted (positions 264 to 293 plus or minus 8 amino acids), or forms to which a 5'-terminal ATG codon and a 3'-terminal stop codon has been added, or forms which have a factor Xa cleavage site and/or 3 to 10, preferably 6 Histidine codons have been added.

More particularly, the recombinant vectors according to the present invention encompass a nucleic acid having an HCV cDNA segment encoding the polyprotein starting in the region between amino acid positions 290 and 406 and ending in the region between positions 600 and 820 of the HCV polyprotein, more preferably starting in the region between positions 322 and 406, even more preferably starting in the region between positions 347 and 406, even still more preferably starting in the region between positions 364 and 406 for expression of the HCV single E2 protein. Most preferably, the present recombinant vector encompasses a recombinant nucleic acid having a HCV cDNA segment encoding the polyprotein starting in the region between positions 290 and 406, and ending at any position of positions 623, 650, 661, 673, 710, 715, 720, 746 or 809, for expression of HCV single E2 protein. Also within the scope of the present invention are forms to which a 5'-terminal ATG codon and a 3'-terminal stop codon has been added, or forms which have a factor Xa cleavage site and/or 3 to 10, preferably 6 Histidine codons have been added.

A variety of vectors may be used to obtain recombinant expression of HCV single or specific oligomeric envelope proteins of the present invention. Lower eukaryotes such as yeasts and glycosylation mutant strains are typically transformed with plasmids, or are transformed with a recombinant virus. The vectors may replicate within the host independently, or may integrate into the host cell genome.

Higher eukaryotes may be transformed with vectors, or may be infected with a recombinant virus, for example a recombinant vaccinia virus. Techniques and vectors for the insertion of foreign DNA into vaccinia virus are well known in the art, and utilize, for example homologous recombination. A wide variety of viral promoter sequences, possibly terminator sequences and poly(A)-addition sequences, pos-

sibly enhancer sequences and possibly amplification sequences, all required for the mammalian expression, are available in the art. Vaccinia is particularly preferred since vaccinia halts the expression of host cell proteins. Vaccinia is also very much preferred since it allows the expression of E1 and E2 proteins of HCV in cells or individuals which are immunized with the live recombinant vaccinia virus. For vaccination of humans the avipox and Ankara Modified Virus (AMV) are particularly useful vectors.

Also known are insect expression transfer vectors derived from baculovirus *Autographa californica* nuclear polyhedrosis virus (AcNPV), which is a helper-independent viral expression vector. Expression vectors derived from this system usually use the strong viral polyhedrin gene promoter to drive the expression of heterologous genes. Different vectors as well as methods for the introduction of heterologous DNA into the desired site of baculovirus are available to the man skilled in the art for baculovirus expression. Also different signals for posttranslational modification recognized by insect cells are known in the art.

Also included within the scope of the present invention is a method for producing purified recombinant single or specific oligomeric HCV E1 or E2 or E1/E2 proteins, wherein the cysteine residues involved in aggregates formation are replaced at the level of the nucleic acid sequence by other residues such that aggregate formation is prevented. The recombinant proteins expressed by recombinant vectors carrying such a mutated E1 and/or E2 protein encoding nucleic acid are also within the scope of the present invention.

The present invention also relates to recombinant E1 and/or E2 and/or E1/E2 proteins characterized in that at least one of their glycosylation sites has been removed and are consequently termed glycosylation mutants. As explained in the Examples section, different glycosylation mutants may be desired to diagnose (screening, confirmation, prognosis, etc.) and prevent HCV disease according to the patient in question. An E2 protein glycosylation mutant lacking the GLY4 has for instance been found to improve the reactivity of certain sera in diagnosis. These glycosylation mutants are preferably purified according to the method disclosed in the present invention. Also contemplated within the present invention are recombinant vectors carrying the nucleic acid insert encoding such a E1 and/or E2 and/or E1/E2 glycosylation mutant as well as host cells transformed with such a recombinant vector.

The present invention also relates to recombinant vectors including a polynucleotide which also forms part of the present invention. The present invention relates more particularly to the recombinant nucleic acids as represented in SEQ ID NO 3, 5, 7, 9, 11, 13, 21, 23, 25, 27, 29, 31, 35, 37, 39, 41, 43, 45, 47 and 49, or parts thereof.

The present invention also contemplates host cells transformed with a recombinant vector as defined above, wherein said vector comprises a nucleotide sequence encoding HCV E1 and/or E2 and/or E1/E2 protein as defined above in addition to a regulatory sequence operably linked to said HCV E1 and/or E2 and/or E1/E2 sequence and capable of regulating the expression of said HCV E1 and/or E2 and/or E1/E2 protein.

Eukaryotic hosts include lower and higher eukaryotic hosts as described in the definitions section. Lower eukaryotic hosts include yeast cells well known in the art. Higher eukaryotic hosts mainly include mammalian cell lines known in the art and include many immortalized cell lines available from the ATCC, including HeLa cells, Chinese hamster ovary (CHO) cells, Baby hamster kidney (BHK) cells, PK15, RK13 and a number of other cell lines.

The present invention relates particularly to a recombinant E1 and/or E2 and/or E1/E2 protein expressed by a host cell as defined above containing a recombinant vector as defined above. These recombinant proteins are particularly purified according to the method of the present invention.

A preferred method for isolating or purifying HCV envelope proteins as defined above is further characterized as comprising at least the following steps:

growing a host cell as defined above transformed with a recombinant vector according to the present invention or with a known recombinant vector expressing E1 and/or E2 and/or E1/E2 HCV envelope proteins in a suitable culture medium,

causing expression of said vector sequence as defined above under suitable conditions, and,

lysing said transformed host cells, preferably in the presence of a SH group blocking agent, such as N-ethylmaleimide (NEM), and possibly a suitable detergent, preferably Empigen-BB,

recovering said HCV envelope protein by affinity purification such as by means of lectin-chromatography or immunoaffinity chromatography using anti-E1 and/or anti-E2 specific monoclonal antibodies, with said lectin being preferably lentil-lectin or GNA, followed by,

incubation of the eluate of the previous step with a disulphide bond cleavage means, such as DTT, preferably followed by incubation with an SH group blocking agent, such as NEM or Biotin-NEM, and,

isolating the HCV single or specific oligomeric E1 and/or E2 and/or E1/E2 proteins such as by means of gel-filtration and possibly also by a subsequent Ni<sup>2+</sup>-IMAC chromatography followed by a desalting step.

As a result of the above-mentioned process, E1 and/or E2 and/or E1/E2 proteins may be produced in a form which elute differently from the large aggregates containing vector-derived components and/or cell components in the void volume of the gel-filtration column or the IMAC column as illustrated in the Examples section. The disulphide bridge cleavage step advantageously also eliminates the false reactivity due to the presence of host and/or expression-system-derived proteins. The presence of NEM and a suitable detergent during lysis of the cells may already partly or even completely prevent the aggregation between the HCV envelope proteins and contaminants.

Ni<sup>2+</sup>-IMAC chromatography followed by a desalting step is preferably used for constructs bearing a (His)<sub>6</sub> as described by Janknecht et al., 1991, and Hochuli et al., 1988.

The present invention also relates to a method for producing monoclonal antibodies in small animals such as mice or rats, as well as a method for screening and isolating human B-cells that recognize anti-HCV antibodies, using the HCV single or specific oligomeric envelope proteins of the present invention.

The present invention further relates to a composition comprising at least one of the following E1 peptides as listed in Table 3:

E1-31 (SEQ ID NO 56) spanning amino acids 181 to 200 of the Core/E1 V1 region,

E1-33 (SEQ ID NO 57) spanning amino acids 193 to 212 of the E1 region,

E1-35 (SEQ ID NO 58) spanning amino acids 205 to 224 of the E1 V2 region (epitope B),

E1-35A (SEQ ID NO 59) spanning amino acids 208 to 227 of the E1 V2 region (epitope B),

1bE1 (SEQ ID NO 53) spanning amino acids 192 to 228 of E1 regions (V1, C1, and V2 regions (containing epitope B)),

E1-51 (SEQ ID NO 66) spanning amino acids 301 to 320 of the E1 region, E1-53 (SEQ ID NO 67) spanning amino acids 313 to 332 of the E1 C4 region (epitope A), E1-55 (SEQ ID NO 68) spanning amino acids 325 to 344 of the E1 region.

The present invention also relates to a composition comprising at least one of the following E2 peptides as listed in Table 3:

Env 67 or E2-67 (SEQ ID NO 72) spanning amino acid positions 397 to 416 of the E2 region (epitope A, recognized by monoclonal antibody 2F10H10, see FIG. 19),

Env 69 or E2-69 (SEQ ID NO 73) spanning amino acid positions 409 to 428 of the E2 region (epitope A),

Env 23 or E2-23 (SEQ ID NO 86) spanning positions 583 to 602 of the E2 region (epitope E),

Env 25 or E2-25 (SEQ ID NO 87) spanning positions 595 to 614 of the E2 region (epitope E),

Env 27 or E2-27 (SEQ ID NO 88) spanning positions 607 to 626 of the E2 region (epitope E),

Env 17B or E2-17B (SEQ ID NO 83) spanning positions 547 to 566 of the E2 region (epitope D),

Env 13B or E2-13B (SEQ ID NO 82) spanning positions 523 to 542 of the E2 region (epitope C; recognized by monoclonal antibody 16A6E7, see FIG. 19).

The present invention also relates to a composition comprising at least one of the following E2 conformational epitopes:

epitope F recognized by monoclonal antibodies 15C8C1, 12D11F1 and 8G10D1H9,

epitope G recognized by monoclonal antibody 9G3E6, epitope H (or C) recognized by monoclonal antibody 10D3C4 and 4H6B2, or, epitope I recognized by monoclonal antibody 17F2C2.

The present invention also relates to an E1 or E2 specific antibody raised upon immunization with a peptide or protein composition, with said antibody being specifically reactive with any of the polypeptides or peptides as defined above, and with said antibody being preferably a monoclonal antibody.

The present invention also relates to an E1 or E2 specific antibody screened from a variable chain library in plasmids or phages or from a population of human B-cells by means of a process known in the art, with said antibody being reactive with any of the polypeptides or peptides as defined above, and with said antibody being preferably a monoclonal antibody.

The E1 or E2 specific monoclonal antibodies of the invention can be produced by any hybridoma liable to be formed according to classical methods from splenic cells of an animal, particularly from a mouse or rat, immunized against the HCV polypeptides or peptides according to the invention, as defined above on the one hand, and of cells of a myeloma cell line on the other hand, and to be selected by the ability of the hybridoma to produce the monoclonal antibodies recognizing the polypeptides which has been initially used for the immunization of the animals.

The antibodies involved in the invention can be labelled by an appropriate label of the enzymatic, fluorescent, or radioactive type.

The monoclonal antibodies according to this preferred embodiment of the invention may be humanized versions of mouse monoclonal antibodies made by means of recombinant DNA technology, departing from parts of mouse and/or human genomic DNA sequences coding for H and L chains from cDNA or genomic clones coding for H and L chains.

Alternatively the monoclonal antibodies according to this preferred embodiment of the invention may be human monoclonal antibodies. These antibodies according to the present embodiment of the invention can also be derived from human peripheral blood lymphocytes of patients infected with HCV, or vaccinated against HCV. Such human monoclonal antibodies are prepared, for instance, by means of human peripheral blood lymphocytes (PBL) repopulation of severe combined immune deficiency (SCID) mice (for recent review, see Duchosal et al., 1992).

The invention also relates to the use of the proteins or peptides of the invention, for the selection of recombinant antibodies by the process of repertoire cloning (Persson et al., 1991).

Antibodies directed to peptides or single or specific oligomeric envelope proteins derived from a certain genotype may be used as a medicament, more particularly for incorporation into an immunoassay for the detection of HCV genotypes (for detecting the presence of HCV E1 or E2 antigen), for prognosing/monitoring of HCV disease, or as therapeutic agents.

Alternatively, the present invention also relates to the use of any of the above-specified E1 or E2 specific monoclonal antibodies for the preparation of an immunoassay kit for detecting the presence of E1 or E2 antigen in a biological sample, for the preparation of a kit for prognosing/monitoring of HCV disease or for the preparation of a HCV medicament.

The present invention also relates to the a method for in vitro diagnosis or detection of HCV antigen present in a biological sample, comprising at least the following steps:

- (i) contacting said biological sample with any of the E1 and/or E2 specific monoclonal antibodies as defined above, preferably in an immobilized form under appropriate conditions which allow the formation of an immune complex,
- (ii) removing unbound components,
- (iii) incubating the immune complexes formed with heterologous antibodies, which specifically bind to the antibodies present in the sample to be analyzed, with said heterologous antibodies having conjugated to a detectable label under appropriate conditions,
- (iv) detecting the presence of said immune complexes visually or mechanically (e.g. by means of densitometry, fluorimetry, colorimetry).

The present invention also relates to a kit for in vitro diagnosis of HCV antigen present in a biological sample, comprising:

- at least one monoclonal antibody as defined above, with said antibody being preferentially immobilized on a solid substrate,
- a buffer or components necessary for producing the buffer enabling binding reaction between these antibodies and the HCV antigens present in the biological sample,
- a means for detecting the immune complexes formed in the preceding binding reaction,
- possibly also including an automated scanning and interpretation device for inferring the HCV antigens present in the sample from the observed binding pattern.

The present invention also relates to a composition comprising E1 and/or E2 and/or E1, E2 recombinant HCV proteins purified according to the method of the present invention or a composition comprising at least one peptides as specified above for use as a medicament.

The present invention more particularly relates to a composition comprising at least one of the above-specified

envelope peptides or a recombinant envelope protein composition as defined above, for use as a vaccine for immunizing a mammal, preferably humans, against HCV, comprising administering a sufficient amount of the composition possibly accompanied by pharmaceutically acceptable adjuvant(s), to produce an immune response.

More particularly, the present invention relates to the use of any of the compositions as described here above for the preparation of a vaccine as described above.

Also, the present invention relates to a vaccine composition for immunizing a mammal, preferably humans, against HCV, comprising HCV single or specific oligomeric proteins or peptides derived from the E1 and/or the E2 region as described above.

Immunogenic compositions can be prepared according to methods known in the art. The present compositions comprise an immunogenic amount of a recombinant E1 and/or E2 and/or E1/E2 single or specific oligomeric proteins as defined above or E1 or E2 peptides as defined above, usually combined with a pharmaceutically acceptable carrier, preferably further comprising an adjuvant.

The single or specific oligomeric envelope proteins of the present invention, either E1 and/or E2 and/or E1/E2, are expected to provide a particularly useful vaccine antigen, since the formation of antibodies to either E1 or E2 may be more desirable than to the other envelope protein, and since the E2 protein is cross-reactive between HCV types and the E1 protein is type-specific. Cocktails including type 1 E2 protein and E1 proteins derived from several genotypes may be particularly advantageous. Cocktails containing a molar excess of E1 versus E2 or E2 versus E1 may also be particularly useful. Immunogenic compositions may be administered to animals to induce production of antibodies, either to provide a source of antibodies or to induce protective immunity in the animal.

Pharmaceutically acceptable carriers include any carrier that does not itself induce the production of antibodies harmful to the individual receiving the composition. Suitable carriers are typically large, slowly metabolized macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers; and inactive virus particles. Such carriers are well known to those of ordinary skill in the art.

Preferred adjuvants to enhance effectiveness of the composition include, but are not limited to: aluminum hydroxide (alum), N-acetyl-muramyl-L-threonyl-D-isoglutamine (th-MDP) as found in U.S. Pat. No. No. 4,606,918, N-acetyl-normuramyl-L-alanyl-D-isoglutamine (nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (MTP-PE) and RIBI, which contains three components extracted from bacteria, monophosphoryl lipid A, trehalose dimycolate, and cell wall skeleton (MPL+TDM+CWS) in a 2% squalene/Tween 80 emulsion. Any of the 3 components MPL, TDM or CWS may also be used alone or combined 2 by 2. Additionally, adjuvants such as Stimulon (Cambridge Bioscience, Worcester, Mass.) or SAF-1 (Syntex) may be used. Further, Complete Freund's Adjuvant (CFA) and Incomplete Freund's Adjuvant (IFA) may be used for non-human applications and research purposes.

The immunogenic compositions typically will contain pharmaceutically acceptable vehicles, such as water, saline, glycerol, ethanol, etc. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, preservatives, and the like, may be included in such vehicles.

Typically, the immunogenic compositions are prepared as injectables, either as liquid solutions or suspensions; solid

forms suitable for solution in, or suspension in, liquid vehicles prior to injection may also be prepared. The preparation also may be emulsified or encapsulated in liposomes for enhanced adjuvant effect. The E1 and E2 proteins may also be incorporated into Immune Stimulating Complexes together with saponins, for example Quil A (ISCOMS).

Immunogenic compositions used as vaccines comprise a 'sufficient amount' or 'an immunologically effective amount' of the envelope proteins of the present invention, as well as any other of the above mentioned components, as needed. 'Immunologically effective amount', means that the administration of that amount to an individual, either in a single dose or as part of a series, is effective for treatment, as defined above. This amount varies depending upon the health and physical condition of the individual to be treated, the taxonomic group of individual to be treated (e.g. non-human primate, primate, etc.), the capacity of the individual's immune system to synthesize antibodies, the degree of protection desired, the formulation of the vaccine, the treating doctor's assessment of the medical situation, the strain of infecting HCV, and other relevant factors. It is expected that the amount will fall in a relatively broad range that can be determined through routine trials. Usually, the amount will vary from 0.01 to 1000  $\mu\text{g}/\text{dose}$ , more particularly from 0.1 to 100  $\mu\text{g}/\text{dose}$ .

The single or specific oligomeric envelope proteins may also serve as vaccine carriers to present homologous (e.g. T cell epitopes or B cell epitopes from the core, NS2, NS3, NS4 or NS5 regions) or heterologous (non-HCV) haptens, in the same manner as Hepatitis B surface antigen (see European Patent Application 174,444). In this use, envelope proteins provide an immunogenic carrier capable of stimulating an immune response to haptens or antigens conjugated to the aggregate. The antigen may be conjugated either by conventional chemical methods, or may be cloned into the gene encoding E1 and/or E2 at a location corresponding to a hydrophilic region of the protein. Such hydrophilic regions include the V1 region (encompassing amino acid positions 191 to 202), the V2 region (encompassing amino acid positions 213 to 223), the V3 region (encompassing amino acid positions 230 to 242), the V4 region (encompassing amino acid positions 230 to 242), the V5 region (encompassing amino acid positions 294 to 303) and the V6 region (encompassing amino acid positions 329 to 336). Another useful location for insertion of haptens is the hydrophobic region (encompassing approximately amino acid positions 264 to 293). It is shown in the present invention that this region can be deleted without affecting the reactivity of the deleted E1 protein with antisera. Therefore, haptens may be inserted at the site of the deletion.

The immunogenic compositions are conventionally administered parenterally, typically by injection, for example, subcutaneously or intramuscularly. Additional formulations suitable for other methods of administration include oral formulations and suppositories. Dosage treatment may be a single dose schedule or a multiple dose schedule. The vaccine may be administered in conjunction with other immunoregulatory agents.

The present invention also relates to a composition comprising peptides or polypeptides as described above, for in vitro detection of HCV antibodies present in a biological sample.

The present invention also relates to the use of a composition as described above for the preparation of an immunoassay kit for detecting HCV antibodies present in a biological sample.

The present invention also relates to a method for in vitro diagnosis of HCV antibodies present in a biological sample, comprising at least the following steps

- (i) contacting said biological sample with a composition comprising any of the envelope peptide or proteins as defined above, preferably in an immobilized form under appropriate conditions which allow the formation of an immune complex, wherein said peptide or protein can be a biotinylated peptide or protein which is covalently bound to a solid substrate by means of streptavidin or avidin complexes,
- (ii) removing unbound components,
- (iii) incubating the immune complexes formed with heterologous antibodies, with said heterologous antibodies having conjugated to a detectable label under appropriate conditions,
- (iv) detecting the presence of said immune complexes visually or mechanically (e.g. by means of densitometry, fluorimetry, colorimetry).

Alternatively, the present invention also relates to competition immunoassay formats in which recombinantly produced purified single or specific oligomeric protein E1 and/or E2 and/or E1/E2 proteins as disclosed above are used in combination with E1 and/or E2 peptides in order to compete for HCV antibodies present in a biological sample.

The present invention also relates to a kit for determining the presence of HCV antibodies, in a biological sample, comprising:

- at least one peptide or protein composition as defined above, possibly in combination with other polypeptides or peptides from HCV or other types of HCV, with said peptides or proteins being preferentially immobilized on a solid substrate, more preferably on different microwells of the same ELISA plate, and even more preferentially on one and the same membrane strip,
- a buffer or components necessary for producing the buffer enabling binding reaction between these polypeptides or peptides and the antibodies against HCV present in the biological sample,
- means for detecting the immune complexes formed in the preceding binding reaction,
- possibly also including an automated scanning and interpretation device for inferring the HCV genotypes present in the sample from the observed binding pattern.

The immunoassay methods according to the present invention utilize single or specific oligomeric antigens from the E1 and/or E2 domains that maintain linear (in case of peptides) and conformational epitopes (single or specific oligomeric proteins) recognized by antibodies in the sera from individuals infected with HCV. It is within the scope of the invention to use for instance single or specific oligomeric antigens, dimeric antigens, as well as combinations of single or specific oligomeric antigens. The HCV E1 and E2 antigens of the present invention may be employed in virtually any assay format that employs a known antigen to detect antibodies. Of course, a format that denatures the HCV conformational epitope should be avoided or adapted. A common feature of all of these assays is that the antigen is contacted with the body component suspected of containing HCV antibodies under conditions that permit the antigen to bind to any such antibody present in the component. Such conditions will typically be physiologic temperature, pH and ionic strength using an excess of antigen. The incubation of the antigen with the specimen is followed by detection of immune complexes comprised of the antigen.

Design of the immunoassays is subject to a great deal of variation, and many formats are known in the art. Protocols may, for example, use solid supports, or immunoprecipita-

tion. Most assays involve the use of labeled antibody or polypeptide; the labels may be, for example, enzymatic, fluorescent, chemiluminescent, radioactive, or dye molecules. Assays which amplify the signals from the immune complex are also known; examples of which are assays which utilize biotin and avidin or streptavidin, and enzyme-labeled and mediated immunoassays, such as ELISA assays.

The immunoassay may be, without limitation, in a heterogeneous or in a homogeneous format, and of a standard or competitive type. In a heterogeneous format, the polypeptide is typically bound to a solid matrix or support to facilitate separation of the sample from the polypeptide after incubation. Examples of solid supports that can be used are nitrocellulose (e.g., in membrane or microtiter well form), polyvinyl chloride (e.g., in sheets or microtiter wells), polystyrene latex (e.g., in beads or microtiter plates, polyvinylidene fluoride (known as Immunolon™), diazotized paper, nylon membranes, activated beads, and Protein A beads. For example, Dynatech Immunolon™1 or Immunolon™ 2 microtiter plates or 0.25 inch polystyrene beads (Precision Plastic Ball) can be used in the heterogeneous format. The solid support containing the antigenic polypeptides is typically washed after separating it from the test sample, and prior to detection of bound antibodies. Both standard and competitive formats are known in the art.

In a homogeneous format, the test sample is incubated with the combination of antigens in solution. For example, it may be under conditions that will precipitate any antigen-antibody complexes which are formed. Both standard and competitive formats for these assays are known in the art.

In a standard format, the amount of HCV antibodies in the antibody-antigen complexes is directly monitored. This may be accomplished by determining whether labeled anti-xenogeneic (e.g. anti-human) antibodies which recognize an epitope on anti-HCV antibodies will bind due to complex formation. In a competitive format, the amount of HCV antibodies in the sample is deduced by monitoring the competitive effect on the binding of a known amount of labeled antibody (or other competing ligand) in the complex.

Complexes formed comprising anti-HCV antibody (or in the case of competitive assays, the amount of competing antibody) are detected by any of a number of known techniques, depending on the format. For example, unlabeled HCV antibodies in the complex may be detected using a conjugate of anti-xenogeneic Ig complexed with a label (e.g. an enzyme label).

In an immunoprecipitation or agglutination assay format the reaction between the HCV antigens and the antibody forms a network that precipitates from the solution or suspension and forms a visible layer or film of precipitate. If no anti-HCV antibody is present in the test specimen, no visible precipitate is formed.

There currently exist three specific types of particle agglutination (PA) assays. These assays are used for the detection of antibodies to various antigens when coated to a support. One type of this assay is the hemagglutination assay using red blood cells (RBCs) that are sensitized by passively adsorbing antigen (or antibody) to the RBC. The addition of specific antigen antibodies present in the body component, if any, causes the RBCs coated with the purified antigen to agglutinate.

To eliminate potential non-specific reactions in the hemagglutination assay, two artificial carriers may be used instead of RBC in the PA. The most common of these are latex particles. However, gelatin particles may also be used. The assays utilizing either of these carriers are based on passive agglutination of the particles coated with purified antigens.

The HCV single or specific oligomeric E1 and/or E2 and/or E1e/E2 antigens of the present invention comprised of conformational epitopes will typically be packaged in the form of a kit for use in these immunoassays. The kit will normally contain in separate containers the native HCV antigen, control antibody formulations (positive and/or negative), labeled antibody when the assay format requires the same and signal generating reagents (e.g. enzyme substrate) if the label does not generate a signal directly. The native HCV antigen may be already bound to a solid matrix or separate with reagents for binding it to the matrix. Instructions (e.g. written, tape, CD-ROM, etc.) for carrying out the assay usually will be included in the kit.

Immunoassays that utilize the native HCV antigen are useful in screening blood for the preparation of a supply from which potentially infective HCV is lacking. The method for the preparation of the blood supply comprises the following steps. Reacting a body component, preferably blood or a blood component, from the individual donating blood with HCV E1 and/or E2 proteins of the present invention to allow an immunological reaction between HCV antibodies, if any, and the HCV antigen. Detecting whether anti-HCV antibody—HCV antigen complexes are formed as a result of the reacting. Blood contributed to the blood supply is from donors that do not exhibit antibodies to the native HCV antigens, E1 or E2.

In cases of a positive reactivity to the HCV antigen, it is preferable to repeat the immunoassay to lessen the possibility of false positives. For example, in the large scale screening of blood for the production of blood products (e.g. blood transfusion, plasma, Factor VIII, immunoglobulin, etc.) 'screening' tests are typically formatted to increase sensitivity (to insure no contaminated blood passes) at the expense of specificity; i.e. the false-positive rate is increased. Thus, it is typical to only defer for further testing those donors who are 'repeatedly reactive'; i.e. positive in two or more runs of the immunoassay on the donated sample. However, for confirmation of HCV-positivity, the 'confirmation' tests are typically formatted to increase specificity (to insure that no false-positive samples are confirmed) at the expense of sensitivity. Therefore the purification method described in the present invention for E1 and E2 will be very advantageous for including single or specific oligomeric envelope proteins into HCV diagnostic assays.

The solid phase selected can include polymeric or glass beads, nitrocellulose, microparticles, microwells of a reaction tray, test tubes and magnetic beads. The signal generating compound can include an enzyme, a luminescent compound, a chromogen, a radioactive element and a chemiluminescent compound. Examples of enzymes include alkaline phosphatase, horseradish peroxidase and beta-galactosidase. Examples of enhancer compounds include biotin, anti-biotin and avidin. Examples of enhancer compounds binding members include biotin, anti-biotin and avidin. In order to block the effects of rheumatoid factor-like substances, the test sample is subjected to conditions sufficient to block the effect of rheumatoid factor-like substances. These conditions comprise contacting the test sample with a quantity of anti-human IgG to form a mixture, and incubating the mixture for a time and under conditions sufficient to form a reaction mixture product substantially free of rheumatoid factor-like substance.

The present invention further contemplates the use of E1 proteins, or parts thereof, more particularly HCV single or specific oligomeric E1 proteins as defined above, for in vitro monitoring HCV disease or prognosing the response to treatment (for instance with Interferon) of patients suffering from HCV infection comprising:

incubating a biological sample from a patient with hepatitis C infection with an E1 protein or a suitable part thereof under conditions allowing the formation of an immunological complex,

removing unbound components,

calculating the anti-E1 titers present in said sample (for example at the start of and/or during the course of (interferon) therapy),

monitoring the natural course of HCV disease, or prognosing the response to treatment of said patient on the basis of the amount anti-E1 titers found in said sample at the start of treatment and/or during the course of treatment.

Patients who show a decrease of 2, 3, 4, 5, 7, 10, 15, or preferably more than 20 times of the initial anti-E1 titers could be concluded to be long-term, sustained responders to HCV therapy, more particularly to interferon therapy. It is illustrated in the Examples section, that an anti-E1 assay may be very useful for prognosing long-term response to IFN treatment, or to treatment of Hepatitis C virus disease in general.

More particularly the following E1 peptides as listed in Table 3 were found to be useful for in vitro monitoring HCV disease or prognosing the response to interferon treatment of patients suffering from HCV infection:

E1-31 (SEQ ID NO 56) spanning amino acids 181 to 200 of the Core/E1 V1 region,

E1-33 (SEQ ID NO 57) spanning amino acids 193 to 212 of the E1 region,

E1-35 (SEQ ID NO 58) spanning amino acids 205 to 224 of the E1 V2 region (epitope B),

E1-35A (SEQ ID NO 59) spanning amino acids 208 to 227 of the E1 V2 region (epitope B),

1bE1 (SEQ ID NO 53) spanning amino acids 192 to 228 of E1 regions (V1, C1, and V2 regions (containing epitope B)),

E1-51 (SEQ ID NO 66) spanning amino acids 301 to 320 of the E1 region,

E1-53 (SEQ ID NO 67) spanning amino acids 313 to 332 of the E1 C4 region (epitope A),

E1-55 (SEQ ID NO 68) spanning amino acids 325 to 344 of the E1 region.

It is to be understood that smaller fragments of the above-mentioned peptides also fall within the scope of the present invention. Said smaller fragments can be easily prepared by chemical synthesis and can be tested for their ability to be used in an assay as detailed above and in the Examples section.

The present invention also relates to a kit for monitoring HCV disease or prognosing the response to treatment (for instance to interferon) of patients suffering from HCV infection comprising:

at least one E1 protein or E1 peptide, more particularly an E1 protein or E1 peptide as defined above,

a buffer or components necessary for producing the buffer enabling the binding reaction between these proteins or peptides and the anti-E1 antibodies present in a biological sample,

means for detecting the immune complexes formed in the preceding binding reaction,

possibly also an automated scanning and interpretation device for inferring a decrease of anti-E1 titers during the progression of treatment.

It is to be understood that also E2 protein and peptides according to the present invention can be used to a certain

degree to monitor/prognose HCV treatment as indicated above for the E1 proteins or peptides because also the anti-E2 levels decrease in comparison to antibodies to the other HCV antigens. It is to be understood, however, that it might be possible to determine certain epitopes in the E2 region which would also be suited for use in a test for monitoring/prognosing HCV disease.

The present invention also relates to a serotyping assay for detecting one or more serological types of HCV present in a biological sample, more particularly for detecting antibodies of the different types of HCV to be detected combined in one assay format, comprising at least the following steps:

(i) contacting the biological sample to be analyzed for the presence of HCV antibodies of one or more serological types, with at least one of the E1 and/or E2 and/or E1/E2 protein compositions or at least one of the E1 or E2 peptide compositions as defined above, preferentially in an immobilized form under appropriate conditions which allow the formation of an immune complex,

(ii) removing unbound components,

(iii) incubating the immune complexes formed with heterologous antibodies, with said heterologous antibodies being conjugated to a detectable label under appropriate conditions,

(iv) detecting the presence of said immune complexes visually or mechanically (e.g. by means of densitometry, fluorimetry, calorimetry) and inferring the presence of one or more HCV serological types present from the observed binding pattern.

It is to be understood that the compositions of proteins or peptides used in this method are recombinantly expressed type-specific envelope proteins or type-specific peptides.

The present invention further relates to a kit for serotyping one or more serological types of HCV present in a biological sample, more particularly for detecting the antibodies to these serological types of HCV comprising:

at least one E1 and/or E2 and/or E1/E2 protein or E1 or E2 peptide, as defined above,

a buffer or components necessary for producing the buffer enabling the binding reaction between these proteins or peptides and the anti-E1 antibodies present in a biological sample,

means for detecting the immune complexes formed in the preceding binding reaction,

possibly also an automated scanning and interpretation device for detecting the presence of one or more serological types present from the observed binding pattern.

The present invention also relates to the use of a peptide or protein composition as defined above, for immobilization on a solid substrate and incorporation into a reversed phase hybridization assay, preferably for immobilization as parallel lines onto a solid support such as a membrane strip, for determining the presence or the genotype of HCV according to a method as defined above. Combination with other type-specific antigens from other HCV polyprotein regions also lies within the scope of the present invention.

#### FIGURE AND TABLE LEGENDS

FIG. 1: Restriction map of plasmid pgpt ATA 18

FIG. 2: Restriction map of plasmid pgs ATA 18

FIG. 3: Restriction map of plasmid pMS 66

FIG. 4: Restriction map of plasmid pv HCV-11A

FIG. 5: Anti-E1 levels in non-responders to IFN treatment

FIG. 6: Anti-E1 levels in responders to IFN treatment

FIG. 7: Anti-E1 levels in patients with complete response to IFN treatment

FIG. 8: Anti-E1 levels in incomplete responders to IFN treatment

FIG. 9: Anti-E2 levels in non-responders to IFN treatment

FIG. 10: Anti-E2 levels in responders to IFN treatment

FIG. 11: Anti-E2 levels in incomplete responders to IFN treatment

FIG. 12: Anti-E2 levels in complete responders to IFN treatment

FIG. 13: Human anti-E1 reactivity competed with peptides

FIG. 14: Competition of reactivity of anti-E1 monoclonal antibodies with peptides

FIG. 15: Anti-E1 (epitope 1) levels in non-responders to IFN treatment

FIG. 16: Anti-E1 (epitope 1) levels in responders to IFN treatment

FIG. 17: Anti-E1 (epitope 2) levels in non-responders to IFN treatment

FIG. 18: Anti-E1 (epitope 2) levels in responders to IFN treatment

FIG. 19: Competition of reactivity of anti-E2 monoclonal antibodies with peptides

FIG. 20: Human anti-E2 reactivity competed with peptides

FIG. 21: FIGS. 21A–L provide nucleic acid sequences of the present invention. The nucleic acid sequences encoding an E1 or E2 protein according to the present invention may be translated (SEQ ID NO 3 to 13, 21–31, 35 and 41–49 are translated in a reading frame starting from residue number 1, SEQ ID NO 37–39 are translated in a reading frame starting from residue number 2), into the amino acid sequences of the respective E1 or E2 proteins as shown in the sequence listing.

FIG. 22: ELISA results obtained from lentil lectin chromatography eluate fractions of 4 different E1 purifications of cell lysates infected with vvHCV39 (type 1b), vvHCV40 (type 1b), vvHCV62 (type 3a), and vvHCV63 (type 5a).

FIG. 23: Elution profiles obtained from the lentil lectin chromatography of the 4 different E1 constructs on the basis of the values as shown in FIG. 22.

FIG. 24: ELISA results obtained from fractions obtained after gel filtration chromatography of 4 different E1 purifications of cell lysates infected with vvHCV39 (type 1b), vvHCV40 (type 1b), vvHCV62 (type 3a), and vvHCV63 (type 5a).

FIG. 25: Profiles obtained from purifications of E1 proteins of type 1b (1), type 3a (2), and type 5a (3) (from RK13 cells infected with vvHCV39, vvHCV62, and vvHCV63, respectively; purified on lentil lectin and reduced as in example 5.2–5.3) and a standard (4). The peaks indicated with '1', '2', and '3', represent pure E1 protein peaks (see FIG. 24, E1 reactivity mainly in fractions 26 to 30).

FIG. 26: Silver staining of an SDS-PAGE as described in example 4 of a raw lysate of E1 vvHCV40 (type 1b) (lane 1), pool 1 of the gel filtration of vvHCV40 representing fractions 10 to 17 as shown in FIG. 25 (lane 2), pool 2 of the gel filtration of vvHCV40 representing fractions 18 to 25 as shown in FIG. 25 (lane 3), and E1 pool (fractions 26 to 30) (lane 4).

FIG. 27: Streptavidine-alkaline phosphatase blot of the fractions of the gel filtration of E1 constructs 39 (type 1b) and 62 (type 3a). The proteins were labelled with NEM-biotin. Lane 1: start gel filtration construct 39, lane 2: fraction 26 construct 39, lane 3: fraction 27 construct 39, lane 4: fraction 28 construct 39, lane 5: fraction 29 construct 39, lane 6: fraction 30 construct 39, lane 7 fraction 31 construct 39, lane 8: molecular weight marker, lane 9: start gel filtration construct 62, lane 10: fraction 26 construct 62, lane 11: fraction 27 construct 62, lane 12: fraction 28 construct 62, lane 13: fraction 29 construct 62, lane 14: fraction 30 construct 62, lane 15: fraction 31 construct 62.

FIG. 28: Silver staining of an SDS-PAGE gel of the gel filtration fractions of vvHCV-39 (E1s, type 1b) and vvHCV-62 (E1s, type 3a) run under identical conditions as FIG. 26. Lane 1: start gel filtration construct 39, lane 2: fraction 26 construct 39, lane 3: fraction 27 construct 39, lane 4: fraction 28 construct 39, lane 5: fraction 29 construct 39, lane 6: fraction 30 construct 39, lane 7 fraction 31 construct 39, lane 8: molecular weight marker, lane 9: start gel filtration construct 62, lane 10: fraction 26 construct 62, lane 11: fraction 27 construct 62, lane 12: fraction 28 construct 62, lane 13: fraction 29 construct 62, lane 14: fraction 30 construct 62, lane 15: fraction 31 construct 62.

FIG. 29: Western Blot analysis with anti-E1 mouse monoclonal antibody 5E1A10 giving a complete overview of the purification procedure. Lane 1: crude lysate, Lane 2: flow through of lentil chromatography, Lane 3: wash with Empigen BB after lentil chromatography, Lane 4: Eluate of lentil chromatography, Lane 5: Flow through during concentration of the lentil eluate, Lane 6: Pool of E1 after Size Exclusion Chromatography (gel filtration).

FIG. 30: OD<sub>280</sub> profile (continuous line) of the lentil lectin chromatography of E2 protein from RK13 cells infected with vvHCV44. The dotted line represents the E2 reactivity as detected by ELISA (as in example 6).

FIG. 31A: OD<sub>280</sub> profile (continuous line) of the lentil-lectin gel filtration chromatography E2 protein pool from RK13 cells infected with vvHCV44 in which the E2 pool is applied immediately on the gel filtration column (non-reduced conditions). The dotted line represents the E2 reactivity as detected by ELISA (as in example 6).

FIG. 31B: OD<sub>280</sub> profile (continuous line) of the lentil-lectin gel filtration chromatography E2 protein pool from RK13 cells infected with vvHCV44 in which the E2 pool was reduced and blocked according to Example 5.3 (reduced conditions). The dotted line represents the E2 reactivity as detected by ELISA (as in example 6).

FIG. 32: Ni<sup>2+</sup>-IMAC chromatography and ELISA reactivity of the E2 protein as expressed from vvHCV44 after gel filtration under reducing conditions as shown in FIG. 31B.

FIG. 33: Silver staining of an SDS-PAGE of 0.5 μg of purified E2 protein recovered by a 200 mM imidazole elution step (lane 2) and a 30 mM imidazole wash (lane 1) of the Ni<sup>2+</sup>-IMAC chromatography as shown in FIG. 32.

FIG. 34: OD profiles of a desalting step of the purified E2 protein recovered by 200 mM imidazole as shown in FIG. 33, intended to remove imidazole.

FIGS. 35A–I to 35A–8: Antibody levels to the different HCV antigens (Core 1, Core 2, E2HCVR, NS3) for NR and LTR followed during treatment and over a period of 6 to 12 months after treatment determined by means of the LIAscan method. The average values are indicated by the curves with the open squares.



FIGS. 35B-1 to 35B-8: Antibody levels to the different HCV antigens (NS4, NS5, E1 and E2) for NR and LTR followed during treatment and over a period of 6 to 12 months after treatment determined by means of the LIAScan method. The average values are indicated by the curve with the open squares.

FIGS. 36A and 36B: Average E1 antibody (E1Ab) and E2 antibody (E2Ab) levels in the LTR and NR groups.

FIGS. 37A-D: Averages E1 antibody (E1Ab) levels for non-responders (NR) and long term responders (LTR) for type 1b and type 3a.

FIG. 38: Relative map positions of the anti-E2 monoclonal antibodies.

FIG. 39: Partial deglycosylation of HCV E1 envelope protein. The lysate of vvHCV10A-infected RK13 cells were incubated with different concentrations of glycosidases according to the manufacturer's instructions. Right panel: Glycopeptidase F (PNGase F). Left panel: Endoglycosidase H (Endo H).

FIG. 40: Partial deglycosylation of HCV E2 envelope proteins. The lysate of vvHCV64-infected (E2) and vvHCV41-infected (E2s) RK13 cells were incubated with different concentrations of Glycopeptidase F (PNGase F) according to the manufacturer's instructions.

FIG. 41: In vitro mutagenesis of HCV E1 glycoproteins. Map of the mutated sequences and the creation of new restriction sites.

FIG. 42A: In vitro mutagenesis of HCV E1 glycoprotein (part 1). First step of PCR amplification.

FIG. 42B: In vitro mutagenesis of HCV E1 glycoprotein (part 2). Overlap extension and nested PCR.

FIG. 43: In vitro mutagenesis of HCV E1 glycoproteins. Map of the PCR mutated fragments (GLY-# and OVR-#) synthesized during the first step of amplification.

FIG. 44A: Analysis of E1 glycoprotein mutants by Western blot expressed in HeLa (left) and RK13 (right) cells. Lane 1: wild type VV (vaccinia virus), Lane 2: original E1 protein (vvHCV-10A), Lane 3: E1 mutant Gly-1 (vvHCV-81), Lane 4: E1 mutant Gly-2 (vvHCV-82), Lane 5: E1 mutant Gly-3 (vvHCV-83), Lane 6: E1 mutant Gly-4 (vvHCV-84), Lane 7: E1 mutant Gly-5 (vvHCV-85), Lane 8: E1 mutant Gly-6 (vvHCV-86).

FIG. 44B: Analysis of E1 glycosylation mutant vaccinia viruses by PCR amplification/restriction. Lane 1: E1 (vvHCV-10A), BspE I, Lane 2: E1.GLY-1 (vvHCV-81), BspE I, Lane 4: E1 (vvHCV-10A), Sac I, Lane 5: E1.GLY-2 (vvHCV-82), Sac I, Lane 7: E1 (vvHCV-10A), Sac I, Lane 8: E1 .GLY-3 (vvHCV-83), Sac I, Lane 10: E1 (vvHCV-10A), Stu I, Lane 11: E1.GLY-4 (vvHCV-84), Stu I, Lane 13: E1 (vvHCV-10A), Sma I, Lane 14: E1.GLY-5 (vvHCV-85), Sma I, Lane 16: E1 (vvHCV-10A), Stu I, Lane 17: E1.GLY-6 (vvHCV-86), Stu I, Lane 3-6-9-12-15: Low Molecular Weight Marker, pBluescript SK+, Msp I.

FIG. 45: SDS polyacrylamide gel electrophoresis of recombinant E2 expressed in *S. cerevisiae*. Inoculates were grown in leucine selective medium for 72 hrs. and diluted 1/15 in complete medium. After 10 days of culture at 28° C., medium samples were taken. The equivalent of 200 µl of culture supernatant concentrated by speedvac was loaded on the gel. Two independent transformants were analysed.

FIG. 46: SDS polyacrylamide gel electrophoresis of recombinant E2 expressed in a glycosylation deficient *S. cerevisiae* mutant. Inoculae were grown in leucine selective medium for 72 hrs. and diluted 1/15 in complete medium. After 10 days of culture at 28° C., medium samples were taken. The equivalent of 350 µl of culture supernatant, concentrated by ion exchange chromatography, was loaded on the gel.

Table 1 : Features of the respective clones and primers used for amplification for constructing the different forms of the E1 protein as depicted in Example 1.

Table 2 : Summary of Anti-E1 tests

Table 3 : Synthetic peptides for competition studies

Table 4: Changes of envelope antibody levels over time.

Table 5: Difference between LTR and NR

Table 6: Competition experiments between murine E2 monoclonal antibodies

Table 7: Primers for construction of E1 glycosylation mutants

Table 8: Analysis of E1 glycosylation mutants by ELISA

## EXAMPLE 1

### Cloning and Expression of the Hepatitis C Virus E1 Protein

#### 1. Construction of Vaccinia Virus Recombination vectors

The pgptATA18 vaccinia recombination plasmid is a modified version of pATA18 (Stunnenberg et al, 1988) with an additional insertion containing the *E. coli* xanthine guanine phosphoribosyl transferase gene under the control of the vaccinia virus 13 intermediate promoter (FIG. 1). The plasmid pgsATA18 was constructed by inserting an oligonucleotide linker with SEQ ID NO 1/94, containing stop codons in the three reading frames, into the Pst I and HindIII-cut pATA18 vector. This created an extra Pac I restriction site (FIG. 2). The original HindIII site was not restored.

Oligonucleotide linker with SEQ ID NO 1/94:  
 5' G GCATGC AAGCTT AATTAATT 3'  
 3' ACGTC CGTACG TTCGAA TTAATTAA TCGA 5'  
 PstI SphI HindIII Pac I (HindIII)

In order to facilitate rapid and efficient purification by means of Ni<sup>2+</sup>-chelation of engineered histidine stretches fused to the recombinant proteins, the vaccinia recombination vector pMS66 was designed to express secreted proteins with an additional carboxy-terminal histidine tag. An oligonucleotide linker with SEQ ID NO 2/95, containing unique sites for 3 restriction enzymes generating blunt ends (Sma I, Stu I and PmlI/BbrPI) was synthesized in such a way that the carboxy-terminal end of any cDNA could be inserted in frame with a sequence encoding the protease factor Xa cleavage site followed by a nucleotide sequence encoding 6 histidines and 2 stop codons (a new Pac I restriction site was also created downstream the 3' end). This oligonucleotide

with SEQ ID NO 2/95 was introduced between the Xma I and Pst I sites of pgpATA18 (FIG. 3).

Oligonucleotide linker with SEQ ID NO 2/95:

Oligonucleotide linker with SEQ ID NO 2/95:

'5' CCGGG GAGGCGTGACACGTGATCGAGGGCAGACACCATCACCACCATCACAATAGTTAATTAA CTGCA3

3' C CTCCGGACGTGCACTAGCTCCCGTCTGTGGTAGTGGTGGTAGTATTATCAATTAATT G  
XmaI PstI

## EXAMPLE 2

### Construction of HCV Recombinant Plasmids

#### 2.1. Constructs Encoding Different Forms of the E1 Protein

Polymerase Chain Reaction (PCR) products were derived from the serum samples by RNA preparation and subsequent reverse-transcription and PCR as described previously (Stuyver et al., 1993b). Table 1 shows the features of the respective clones and the primers used for amplification. The PCR fragments were cloned into the Sma I-cut pSP72 (Promega) plasmids. The following clones were selected for insertion into vaccinia recombination vectors: HCC19A (SEQ ID NO 3), HCC11 OA (SEQ ID NO 5), HCC111A (SEQ ID NO 7), HCC112A (SEQ ID NO 9), HCC113A (SEQ ID NO 11), and HCC117A (SEQ ID NO 13) as depicted in FIG. 21. cDNA fragments containing the E1-coding regions were cleaved by EcoRI and HindIII restriction from the respective pSP72 plasmids and inserted into the EcoRI/HindIII-cut pgpATA18 vaccinia recombination vector (described in example 1), downstream of the 11K vaccinia virus late promoter. The respective plasmids were designated pvHCV-9A, pvHCV-10A, pvHCV-11A, pvHCV-12A, pvHCV-13A and pvHCV-17A, of which pvHCV-11A is shown in FIG. 4.

#### 2.2. Hydrophobic Region E1 Deletion Mutants

Clone HCC137, containing a deletion of codons Asp264 to Val287 (nucleotides 790 to 861, region encoding hydrophobic domain 1) was generated as follows: 2 PCR fragments were generated from clone HCC110A with primer sets HCPPr52 (SEQ ID NO 16)/HCPPr107 (SEQ ID NO 19) and HCPPr108 (SEQ ID NO 20)/HCPPr54 (SEQ ID NO 18). These primers are shown in FIG. 21. The two PCR fragments were purified from agarose gel after electrophoresis and 1 ng of each fragment was used together as template for PCR by means of primers HCPPr52 (SEQ ID NO 16) and HCPPr54 (SEQ ID NO 18). The resulting fragment was cloned into the Sma I-cut pSP72 vector and clones containing the deletion were readily identified because of the deletion of 24 codons (72 base pairs). Plasmid pSP72HCC137 containing clone HCC137 (SEQ ID 15) was selected. A recombinant vaccinia plasmid containing the full-length E1 cDNA lacking hydrophobic domain I was constructed by inserting the HCV sequence surrounding the deletion (fragment cleaved by Xma I and BamH I from the vector pSP72-HCC137) into the Xma I-BamHI sites of the vaccinia plasmid pvHCV-10A. The resulting plasmid was named pvHCV-37. After confirmatory sequencing, the amino-terminal region containing the internal deletion was isolated from this vector pvHCV-37 (cleavage by EcoR I and BstE II) and reinserted into the Eco RI and Bst EII-cut pvHCV-11A plasmid. This construct was expected to express an E1 protein with both hydrophobic domains deleted and was named pvHCV-38. The E1-coding region of clone HCC138 is represented by SEQ ID NO 23.

As the hydrophilic region at the E1 carboxyterminus (theoretically extending to around amino acids 337-340)

was not completely included in construct pvHCV-38, a larger E1 region lacking hydrophobic domain I was isolated from the pvHCV-37 plasmid by EcoRI/BamHI cleavage and

cloned into an EcoRII/BamHI-cut pgsATA-18 vector. The resulting plasmid was named pvHCV-39 and contained clone HCC139 (SEQ ID NO 25). The same fragment was cleaved from the pvHCV-37 vector by BamHI (of which the sticky ends were filled with Klenow DNA Polymerase I (Boehringer)) and subsequently by EcoRI (5' cohesive end). This sequence was inserted into the EcoRI and Bbr PI-cut vector pMS-66. This resulted in clone HCC140 (SEQ ID NO 27) in plasmid pvHCV-40, containing a 6 histidine tail at its carboxy-terminal end.

#### 2.3. E1 of Other Genotypes

Clone HCC162 (SEQ ID NO 29) was derived from a type 3a-infected patient with chronic hepatitis C (serum BR36, clone BR36-9-13, SEQ ID NO 19 in WO 94/25601, and see also Stuyver et al. 1993a) and HCC163 (SEQ ID NO 31) was derived from a type 5a-infected child with post-transfusion hepatitis (serum BE95, clone PC-4-1, SEQ ID NO 45 in WO 94/25601).

#### 2.4. E2 Constructs

The HCV E2 PCR fragment 22 was obtained from serum BE11 (genotype 1b) by means of primers HCPPr109 (SEQ ID NO 33) and HCPPr72 (SEQ ID NO 34) using techniques of RNA preparation, reverse-transcription and PCR, as described in Stuyver et al., 1993b, and the fragment was cloned into the Sma I-cut pSP72 vector. Clone HCC122A (SEQ ID NO 35) was cut with NcoI/AlwNI or by BamHI/AlwNI and the sticky ends of the fragments were blunted (NcoI and BamHI sites with Klenow DNA Polymerase I (Boehringer), and AlwNI with T4 DNA polymerase (Boehringer)). The BamHI/AlwNI cDNA fragment was then inserted into the vaccinia pgsATA-18 vector that had been linearized by EcoRI and Hind III cleavage and of which the cohesive ends had been filled with Klenow DNA Polymerase (Boehringer). The resulting plasmid was named pvHCV-41 and encoded the E2 region from amino acids Met347 to Gln673, including 37 amino acids (from Met347 to Gly383) of the E1 protein that can serve as signal sequence. The same HCV cDNA was inserted into the EcoR I and Bbr PI-cut vector pMS66, that had subsequently been blunt ended with Klenow DNA Polymerase. The resulting plasmid was named pvHCV-42 and also encoded amino acids 347 to 683. The NcoI/AlwNI fragment was inserted in a similar way into the same sites of pgsATA-18 (pvHCV-43) or pMS-66 vaccinia vectors (pvHCV-44). pvHCV-43 and pvHCV-44 encoded amino acids 364 to 673 of the HCV polyprotein, of which amino acids 364 to 383 were derived from the natural carboxyterminal region of the E1 protein encoding the signal sequence for E2, and amino acids 384 to 673 of the mature E2 protein.

#### 2.5. Generation of Recombinant HCV-Vaccinia Viruses

Rabbit kidney RK13 cells (ATCC CCL 37), human osteosarcoma 143B thymidine kinase deficient (TK-) (ATCC CRL 8303), HeLa (ATCC CCL 2), and Hep G2 (ATCC HB 8065) cell lines were obtained from the American Type Culture Collection (ATCC, Rockville, Md., USA).

## 31

The cells were grown in Dulbecco's modified Eagle medium (DMEM) supplemented with 10% foetal calf serum, and with Earle's salts (EMEM) for RK13 and 143 B (TK-), and with glucose (4 g/l) for Hep G2. The vaccinia virus WR strain (Western Reserve, ATTC VRI 19) was routinely propagated in either 143B or RK13 cells, as described previously (Panicali & Paoletti, 1982; Piccini et al., 1987; Mackett et al., 1982, 1984, and 1986). A confluent monolayer of 143B cells was infected with wild type vaccinia virus at a multiplicity of infection (m.o.i.) of 0.1 (=0.1 plaque forming unit (PFU) per cell). Two hours later, the vaccinia recombination plasmid was transfected into the infected cells in the form of a calcium phosphate coprecipitate containing 500 ng of the plasmid DNA to allow homologous recombination (Graham & van der Eb, 1973; Mackett et al., 1985). Recombinant viruses expressing the *Escherichia coli* xanthine-guanine phosphoribosyl transferase (gpt) protein were selected on rabbit kidney RK13 cells incubated in selection medium (EMEM containing 25 µg/ml mycophenolic acid (MPA), 250 µg/ml xanthine, and 15 µg/ml hypoxanthine; Falkner and Moss, 1988; Janknecht et al., 1991). Single recombinant viruses were purified on fresh monolayers of RK13 cells under a 0.9% agarose overlay in selection medium. Thymidine kinase deficient (TK-) recombinant viruses were selected and then plaque purified on fresh monolayers of human 143B cells (TK-) in the presence of 25 µg/ml 5-bromo-2'-deoxyuridine. Stocks of purified recombinant HCV-vaccinia viruses were prepared by infecting either human 143 B or rabbit RK13 cells at an m.o.i. of 0.05 (Mackett et al, 1988). The insertion of the HCV cDNA fragment in the recombinant vaccinia viruses was confirmed on an aliquot (50 µl) of the cell lysate after the MPA selection by means of PCR with the primers used to clone the respective HCV fragments (see Table 1). The recombinant vaccinia-HCV viruses were named according to the vaccinia recombination plasmid number, e.g. the recombinant vaccinia virus vvHCV-10A was derived from recombining the wild type WR strain with the pvHCV-10A plasmid.

## EXAMPLE 3

## Infection of Cells with Recombinant Vaccinia Viruses

A confluent monolayer of RK13 cells was infected at a m.o.i. of 3 with the recombinant HCV-vaccinia viruses as described in example 2. For infection, the cell monolayer was washed twice with phosphate-buffered saline pH 7.4 (PBS) and the recombinant vaccinia virus stock was diluted in MEM medium. Two hundred µl of the virus solution was added per 10<sup>5</sup> cells such that the m.o.i. was 3, and incubated for 45 min at 24° C. The virus solution was aspirated and 2 ml of complete growth medium (see example 2) was added per 10<sup>5</sup> cells. The cells were incubated for 24 hr at 37° C. during which expression of the HCV proteins took place.

## EXAMPLE 4

## Analysis of Recombinant Proteins by Means of Western Blotting

The infected cells were washed two times with PBS, directly lysed with lysis buffer (50 mM Tris.HCl pH 7.5, 150 mM NaCl, 1% Triton X-100, 5 mM MgCl<sub>2</sub>, 1 µg/ml aprotinin (Sigma, Bornem, Belgium)) or detached from the flasks by incubation in 50 mM Tris-HCl pH 7.5/10 mM EDTA/150 mM NaCl for 5 min, and collected by centrifugation (5 min at 1000 g). The cell pellet was then resuspended in 200

## 32

µl lysis buffer (50 mM Tris.HCl pH 8.0, 2 mM EDTA, 150 mM NaCl, 5 mM MgCl<sub>2</sub>, aprotinin, 1% Triton X-100) per 10<sup>5</sup> cells. The cell lysates were cleared for 5 min at 14,000 rpm in an Eppendorf centrifuge to remove the insoluble debris. Proteins of 20 µl lysate were separated by means of sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE). The proteins were then electro-transferred from the gel to a nitrocellulose sheet (Amersham) using a Hoefer HSI transfer unit cooled to 4° C. for 2 hr at 100 V constant voltage, in transfer buffer (25 mM Tris.HCl pH 8.0, 192 mM glycine, 20% (v/v) methanol). Nitrocellulose filters were blocked with Blotto (5% (w/v) fat-free instant milk powder in PBS; Johnson et al., 1981) and incubated with primary antibodies diluted in Blotto/0.1% Tween 20. Usually, a human negative control serum or serum of a patient infected with HCV were 200 times diluted and preincubated for 1 hour at room temperature with 200 times diluted wild type vaccinia virus-infested cell lysate in order to decrease the non-specific binding. After washing with Blotto/0.1% Tween 20, the nitrocellulose filters were incubated with alkaline phosphatase substrate solution diluted in Blotto/0.1% Tween 20. After washing with 0.1%, Tween 20 in PBS, the filters were incubated with alkaline phosphatase substrate solution (100 mM Tris.HCl pH 9.5, 100 mM NaCl, 5 mM MgCl<sub>2</sub>, 0.38 µg/ml nitroblue tetrazolium, 0.165 µg/ml 5-bromo-4-chloro-3-indolylphosphate). All steps, except the electrotransfer, were performed at room temperature.

## EXAMPLE 5

## Purification of Recombinant E1 or E2 Protein

## 5.1. Lysis

Infected RK13 cells (carrying E1 or E2 constructs) were washed 2 times with phosphate-buffered saline (PBS) and detached from the culture recipients by incubation in PBS containing 10 mM EDTA. The detached cells were washed twice with PBS and 1 ml of lysis buffer (50 mM Tris.HCl pH 7.5, 150 mM NaCl, 1% Triton X-100, 5 mM MgCl<sub>2</sub>, 1 µg/ml aprotinin (Sigma, Bornem, Belgium) containing 2 mM biotinylated N-ethylmaleimide (biotin-NEM) (Sigma) was added per 10<sup>5</sup> cells at 4° C. This lysate was homogenized with a type B douncer and left at room temperature for 0.5 hours. Another 5 volumes of lysis buffer containing 10 mM N-ethylmaleimide (NEM, Aldrich, Bornem, Belgium) was added to the primary lysate and the mixture was left at room temperature for 15 min. Insoluble cell debris was cleared from the solution by centrifugation in a Beckman JA-14 rotor at 14,000 rpm (30100 g at  $r_{max}$ ) for 1 hour at 4° C.

## 5.2. Lectin Chromatography

The cleared cell lysate was loaded at a rate of 1 ml/min on a 0.8 by 10 cm Lentil-lectin Sepharose 4B column (Pharmacia) that had been equilibrated with 5 column volumes of lysis buffer at a rate of 1 ml/min. The lentil-lectin column was washed with 5 to 10 column volumes of buffer 1 (0.1M potassium phosphate pH 7.3, 500 mM KCl, 5% glycerol, 1 mM 6-NH<sub>2</sub>-hexanoic acid, 1 mM MgCl<sub>2</sub>, and 1% DecylPEG (KWANT, Bedum, The Netherlands). In some experiments, the column was subsequently washed with 10 column volumes of buffer 1 containing 0.5% Empigen-BB (Calbiochem, San Diego, Calif., USA) instead of 1% DecylPEG. The bound material was eluted by applying elution buffer (10 mM potassium phosphate pH 7.3, 5% glycerol, 1 mM hexanoic acid, 1 mM MgCl<sub>2</sub>, 0.5% Empigen-BB, and 0.5 M α-methyl-mannopyranoside). The eluted material was fractionated and fractions were screened for the presence of E1 or E2 protein by means of ELISA as

described in example 6. FIG. 22 shows ELISA results obtained from lentil lectin eluate fractions of 4 different E1 purifications of cell lysates infected with vvHCV39 (type 1b), vvHCV40 (type 1b), vvHCV62 (type 3a), and vvHCV63 (type 5a). FIG. 23 shows the profiles obtained from the values shown in FIG. 22. These results show that the lectin affinity column can be employed for envelope proteins of the different types of HCV.

### 5.3. Concentration and Partial Reduction

The E1- or E2-positive fractions were pooled and concentrated on a Centricon 30 kDa (Amicon) by centrifugation for 3 hours at 5,000 rpm in a Beckman JA-20 rotor at 4° C. In some experiments the E1- or E2-positive fractions were pooled and concentrated by nitrogen evaporation. An equivalent of 3.108 cells was concentrated to approximately 200  $\mu$ l. For partial reduction, 30% Empigen-BB (Calbiochem, San Diego, Calif., USA) was added to this 200  $\mu$ l to a final concentration of 3.5%, and 1 M DTT in H<sub>2</sub>O was subsequently added to a final concentration of 1.5 to 7.5 mM and incubated for 30 min at 37° C. NEM (1M in dimethylsulphoxide) was subsequently added to a final concentration of 50 mM and left to react for another 30 min at 37° C. to block the free sulphhydryl groups.

### 5.4. Gel Filtration Chromatography

A Superdex-200 HR 10/20 column (Pharmacia) was equilibrated with 3 column volumes PBS13% Empigen-BB. The reduced mixture was infected in a 500  $\mu$ l sample loop of the Smart System (Pharmacia) and PBS/3% Empigen-BB buffer was added for gelfiltration. Fractions of 250  $\mu$ l were collected from V<sub>C</sub> to V<sub>1</sub>. The fractions were screened for the presence of E1 or E2 protein as described in example 6.

FIG. 24 shows ELISA results obtained from fractions obtained after gelfiltration chromatography of 4 different E1 purifications of cell lysates infected with vvHCV39 (type 1b), vvHCV40 (type 1b), vvHCV62 (type 3a), and vvHCV63 (type 5a). FIG. 25 shows the profiles obtained from purifications of E1 proteins of types 1b, 3a, and 5a (from RK13 cells infected with vvHCV39, vvHCV62, and vvHCV63, respectively; purified on lentil lectin and reduced as in the previous examples). The peaks indicated with '1', '2', and '3', represent pure E1 protein peaks (E1 reactivity mainly in fractions 26 to 30). These peaks show very similar molecular weights of approximately 70 kDa, corresponding to dimeric E1 protein. Other peaks in the three profiles represent vaccinia virus and/or cellular proteins which could be separated from E1 only because of the reduction step as outlined in example 5.3. and because of the subsequent gelfiltration step in the presence of the proper detergent. As shown in FIG. 26 pool 1 (representing fractions 10 to 17) and pool 2 (representing fractions 18 to 25) contain contaminating proteins not present in the E1 pool (fractions 26 to 30). The E1 peak fractions were ran on SDS/PAGE and blotted as described in example 4. Proteins labelled with NEM-biotin were detected by streptavidin-alkaline phosphatase as shown in FIG. 27. It can be readily observed that, amongst others, the 29 kDa and 45kDa contaminating proteins present before the gelfiltration chromatography (lane 1) are only present at very low levels in the fractions 26 to 30. The band at approximately 65 kDa represents the E1 dimeric form that could not be entirely disrupted into the monomeric E1 form. Similar results were obtained for the type 3a E1 protein (lanes 10 to 15), which shows a faster mobility on SDS/PAGE because of the presence of only 5 carbohydrates instead of 6. FIG. 28 shows a silver stain of an SDS/PAGE gel run in identical conditions as in FIG. 26. A complete overview of the purification procedure is given in FIG. 29.

The presence of purified E1 protein was further confirmed by means of western blotting as described in example 4. The dimeric E1 protein appeared to be non-aggregated and free of contaminants. The subtype 1b E1 protein purified from vvHCV40-infected cells according to the above scheme was aminoterminally sequenced on an 477 Perkins-Elmer sequencer and appeared to contain a tyrosine as first residue. This confirmed that the E1 protein had been cleaved by the signal peptidase at the correct position (between A191 and Y192) from its signal sequence. This confirms the finding of Hijikata et al. (1991) that the aminotermis of the mature E1 protein starts at amino acid position 192.

### 5.5. Purification of the E2 Protein

The E2 protein (amino acids 384 to 673) was purified from RK13 cells infected with vvHCV44 as indicated in Examples 5.1 to 5.4. FIG. 30 shows the OD<sub>280</sub> profile (continuous line) of the lentil lectin chromatography. The dotted line represents the E2 reactivity as detected by ELISA (see example 6). FIG. 31 shows the same profiles obtained from gelfiltration chromatography of the lentil-lectin E2 pool (see FIG. 30), part of which was reduced and blocked according to the methods as set out in example 5.3., and part of which was immediately applied to the column. Both parts of the E2 pool were run on separate gelfiltration columns. It could be demonstrated that E2 forms covalently-linked aggregates with contaminating proteins if no reduction has been performed. After reduction and blocking, the majority of contaminating proteins segregated into the V<sub>0</sub> fraction. Other contaminating proteins copurified with the E2 protein, were not covalently linked to the E2 protein any more because these contaminants could be removed in a subsequent step. FIG. 32 shows an additional Ni<sup>2+</sup>-IMAC purification step carried out for the E2 protein purification. This affinity purification step employs the 6 histidine residues added to the E2 protein as expressed from vvHCV44. Contaminating proteins either run through the column or can be removed by a 30 mM; imidazole wash. FIG. 33 shows a silver-stained SDS/PAGE of 0.5  $\mu$ g of purified E2 protein and a 30 mM imidazole wash. The pure E2 protein could be easily recovered by a 200 mM imidazole elution step. FIG. 34 shows an additional desalting step intended to remove imidazole and to be able to switch to the desired buffer, e.g. PBS, carbonate buffer, saline.

Starting from about 50,000 cm<sup>2</sup> of RK13 cells infected with vvHCV11 A (or vvHCV40) for the production of E1 or vvHCV41, vvHCV42, vvHCV43, or vvHCV44 for production of E2 protein, the procedures described in examples 5.1 to 5.5 allow the purification of approximately 1.3 mg of E1 protein and 0.6 mg of E2 protein.

It should also be remarked that secreted E2 protein (constituting approximately 30–40%, 60–70% being in the intracellular form) is characterized by aggregate formation (contrary to expectations). The same problem is thus posed to purify secreted E2. The secreted E2 can be purified as disclosed above.

## EXAMPLE 6

### ELISA for the Detection of Anti-E1 or Anti-E2 Antibodies or for the Detection of E1 or E2 Proteins

Maxisorb microwell plates (Nunc, Roskilde, Denmark) were coated with 1 volume (e.g. 50  $\mu$ l or 100  $\mu$ l or 200  $\mu$ l) per well of a 5  $\mu$ g/ml solution of Streptavidin (Boehringer Mannheim) in PBS for 16 hours at 4° C. or for 1 hour at 37° C. Alternatively, the wells were coated with 1 volume of 5

$\mu\text{g/ml}$  of *Galanthus nivalis* agglutinin (GNA) in 50 mM sodium carbonate buffer pH 9.6 for 16 hours at 4° C. or for 1 hour at 37° C. In the case of coating with GNA, the plates were washed 2 times with 400  $\mu\text{l}$  of Washing Solution of the innotest HCV Ab III kit (Innogenetics, Zwijndrecht, Belgium). Unbound coating surfaces were blocked with 1.5 to 2 volumes of blocking solution (0.1% casein and 0.1%  $\text{NaN}_3$  in PBS) for 1 hour at 37° C. or for 16 hours at 4° C. Blocking solution was aspirated. Purified E1 or E2 was diluted to 100–1000 ng/ml (concentration measured at  $A=280$  nm) or column fractions to be screened for E1 or E2 (see example 5), or E1 or E2 in non-purified cell lysates (example 5.1.) were diluted 20 times in blocking solution, and 1 volume of the E1 or E2 solution was added to each well and incubated for 1 hour at 37° C. on the Streptavidin- or GNA-coated plates. The microwells were washed 3 times with 1 volume of Washing Solution of the Innotest HCV Ab III kit (Innogenetics, Zwijndrecht, Belgium). Serum samples were diluted 20 times or monoclonal anti-E1 or anti-E2 antibodies were diluted to a concentration of 20 ng/ml in Sample Diluent of the Innotest HCV Ab III kit and 1 volume of the solution was left to react with the E1 or E2 protein for 1 hour at 37° C. The microwells were washed 5 times with 400  $\mu\text{l}$  of Washing Solution of the Innotest HCV Ab III kit (Innogenetics, Zwijndrecht, Belgium). The bound antibodies were detected by incubating each well for 1 hour at 37° C. with a goat anti-human or anti-mouse IgG, peroxidase-conjugated secondary antibody (DAKO, Glostrup, Denmark) diluted  $\frac{1}{80,000}$  in 1 volume of Conjugate Diluent of the Innotest HCV Ab III kit (Innogenetics, Zwijndrecht, Belgium), and color development was obtained by addition of substrate of the Innotest HCV Ab III kit (Innogenetics, Zwijndrecht, Belgium) diluted 100 times in 1 volume of Substrate Solution of the Innotest HCV Ab III kit (Innogenetics, Zwijndrecht, Belgium) for 30 min at 24° C. after washing of the plates 3 times with 400  $\mu\text{l}$  of Washing Solution of the Innotest HCV Ab III kit (Innogenetics, Zwijndrecht, Belgium).

#### EXAMPLE 7

##### Follow up of Patient Groups with Different Clinical Profiles

###### 7.1. Monitoring of Anti-E1 and Anti-E2 Antibodies

The current hepatitis C virus (HCV) diagnostic assays have been developed for screening and confirmation of the presence of HCV antibodies. Such assays do not seem to provide information useful for monitoring of treatment or for prognosis of the outcome of disease. However, as is the case for hepatitis B, detection and quantification of anti-envelope antibodies may prove more useful in a clinical setting. To investigate the possibility of the use of anti-E1 antibody titer and anti-E2 antibody titer as prognostic markers for outcome of hepatitis C disease, a series of IFN- $\alpha$  treated patients with long-term sustained response (defined as patients with normal transaminase levels and negative HCV-RNA test (PCR in the 5' non-coding region) in the blood for a period of at least 1 year after treatment) was compared with patients showing no response or showing biochemical response with relapse at the end of treatment.

A group of 8 IFN- $\alpha$  treated patients with long-term sustained response (LTR, follow up 1 to 3.5 years, 3 type 3a and 5 type 1b) was compared with 9 patients showing non-complete responses to treatment (NR, follow up 1 to 4 years, 6 type 1b and 3 type 3a). Type 1b (vvHCV-39, see example 2.5.) and 3a E1 (vvHCV-62, see example 2.5.) proteins were expressed by the vaccinia virus system (see

examples 3 and 4) and purified to homogeneity (example 5). The samples derived from patients infected with a type 1b hepatitis C virus were tested for reactivity with purified type 1b E1 protein, while samples of a type 3a infection were tested for reactivity of anti-type 3a E1 antibodies in an ELISA as described in example 6. The genotypes of hepatitis C viruses infecting the different patients were determined by means of the Inno-LiPA genotyping assay (innogenetics, Zwijndrecht, Belgium). FIG. 5 shows the anti-E1 signal-to-noise ratios of these patients followed during the course of interferon treatment and during the follow-up period after treatment. LTR cases consistently showed rapidly declining anti-E1 levels (with complete negativation in 3 cases), while anti-E1 levels of NR cases remained approximately constant. Some of the obtained anti-E1 data are shown in Table 2 as average S/N ratios  $\pm$ SD (mean anti-E1 titer). The anti-E1 titer could be deduced from the signal to noise ratio as show in FIGS. 5, 6, 7, and 8.

Already at the end of treatment, marked differences could be observed between the 2 groups. Anti-E1 antibody titers had decreased 6.9 times in LTR but only 1.5 times in NR. At the end of follow up, the anti-E1 titers had declined by a factor of 22.5 in the patients with sustained response and even slightly increased in NR. Therefore, based on these data, decrease of anti-E1 antibody levels during monitoring of IFN- $\alpha$  therapy correlates with long-term, sustained response to treatment. The anti-E1 assay may be very useful for prognosis of long-term response to IFN treatment, or to treatment of the hepatitis C disease in general.

This finding was not expected. On the contrary, the inventors had expected the anti-E1 antibody levels to increase during the course of IFN treatment in patients with long term response. As is the case for hepatitis B, the virus is cleared as a consequence of the seroconversion for anti-HBsAg antibodies. Also in many other virus infections, the virus is eliminated when anti-envelope antibodies are raised. However, in the experiments of the present invention, anti-E1 antibodies clearly decreased in patients with a long-term response to treatment, while the antibody-level remained approximately at the same level in non-responding patients. Although the outcome of these experiments was not expected, this non-obvious finding may be very important and useful for clinical diagnosis of HCV infections. As shown in FIGS. 9, 10, 11, and 12, anti-E2 levels behaved very differently in the same patients studied and no obvious decline in titers was observed as for anti-E1 antibodies. FIG. 35 gives a complete overview of the pilot study.

As can be deduced from Table 2, the anti-E1 titers were on average at least 2 times higher at the start of treatment in long term responders compared with incomplete responders to treatment. Therefore, measuring the titer of anti-E1 antibodies at the start of treatment, or monitoring the patient during the course of infection and measuring the anti-E1 titer, may become a useful marker for clinical diagnosis of hepatitis C. Furthermore, the use of more defined regions of the E1 or E2 proteins may become desirable, as shown in example 7.3.

###### 7.2. Analysis of E1 and E2 Antibodies in a Larger Patient Cohort

The pilot study lead the inventors to conclude that, in case infection was completely cleared, antibodies to the HCV envelope proteins changed more rapidly than antibodies to the more conventionally studied HCV antigens, with E1 antibodies changing most vigorously. We therefore included more type 1b and 3a-infected LTR and further supplemented the cohort with a matched series of NR, such that both

groups included 14 patients each. Some partial responders (PR) and responders with relapse (RR) were also analyzed.

FIG. 36 depicts average E1 antibody (E1Ab) and E2 antibody (E2Ab) levels in the LTR and NR groups and Tables 4 and 5 show the statistical analyses. In this larger cohort, higher E1 antibody levels before IFN- $\alpha$  therapy were associated with LTR ( $P < 0.03$ ). Since much higher E1 antibody levels were observed in type 3a-infected patients compared with type 1b-infected patients (FIG. 37), the genotype was taken into account (Table 4). Within the type 1b-infected group, LTR also had higher E1 antibody levels than NR at the initiation of treatment [ $P < 0.05$ ]; the limited number of type 3a-infected NR did not allow statistical analysis.

Of antibody levels monitored in LTR during the 1.5-year follow up period, only E1 antibodies cleared rapidly compared with levels measured at initiation of treatment [ $P = 0.0058$ , end of therapy;  $P = 0.0047$  and  $P = 0.0051$  at 6 and 12 months after therapy, respectively]. This clearance remained significant within type 1- or type 3-infected LTR (average  $P$  values  $< 0.05$ ). These data confirmed the initial finding that E1Ab levels decrease rapidly in the early phase of resolution. This feature seems to be independent of viral genotype. In NR, PR, or RR, no changes in any of the antibodies measured were observed throughout the follow up period. In patients who responded favourably to treatment with normalization of ALT levels and HCV-RNA negative during treatment, there was a marked difference between sustained responders (LTR) and responders with a relapse (RR). In contrast to LTR, RR did not show any decreasing E1 antibody levels, indicating the presence of occult HCV infection that could neither be demonstrated by PCR or other classical techniques for detection of HCV-RNA, nor by raised ALT levels. The minute quantities of viral RNA, still present in the RR group during treatment, seemed to be capable of anti-E1 B cell stimulation. Anti-E1 monitoring may therefore not only be able to discriminate LTR from NR, but also from RR.

### 7.3. Monitoring of Antibodies of Defined Regions of the E1 Protein

Although the molecular biological approach of identifying HCV antigens resulted in unprecedented breakthrough in the development of viral diagnostics, the method of immune screening of  $\lambda$ gt11 libraries predominantly yielded linear epitopes dispersed throughout the core and non-structural regions, and analysis of the envelope regions had to await cloning and expression of the E1/E2 region in mammalian cells. This approach sharply contrasts with many other viral infections of which epitopes to the envelope regions had already been mapped long before the deciphering of the genomic structure. Such epitopes and corresponding antibodies often had neutralizing activity useful for vaccine development and/or allowed the development of diagnostic assays with clinical or prognostic significance (e.g. antibodies to hepatitis B surface antigen). As no HCV vaccines or tests allowing clinical diagnosis and prognosis of hepatitis C disease are available today, the characterization of viral envelope regions exposed to immune surveillance may significantly contribute to new directions in HCV diagnosis and prophylaxis.

Several 20-mer peptides (Table 3) that overlapped each other by 8 amino acids, were synthesized according to a previously described method (EP-A-0 489 968) based on the HC-J1 sequence (Okamoto et al., 1990). None of these, except peptide env35 (also referred to as E1-35), was able to detect antibodies in sera of approximately 200 HCV cases.

Only 2 sera reacted slightly with the env35 peptide. However, by means of the anti-E1 ELISA as described in example 6, it was possible to discover additional epitopes as follows: The anti-E1 ELISA as described in example 6 was modified by mixing 50  $\mu$ g/ml of E1 peptide with the  $\frac{1}{20}$  diluted human serum in sample diluent. FIG. 13 shows the results of reactivity of human sera to the recombinant E1 (expressed from vvHCV-40) protein, in the presence of single or of a mixture of E1 peptides. While only 2% of the sera could be detected by means of E1 peptides coated on strips in a Line Immunoassay format, over half of the sera contained anti-E1 antibodies which could be competed by means of the same peptides, when tested on the recombinant E1 protein. Some of the murine monoclonal antibodies obtained from Balb/C mice after injection with purified E1 protein were subsequently competed for reactivity to E1 with the single peptides (FIG. 14). Clearly, the region of env53 contained the predominant epitope, as the addition of env53 could substantially compete reactivity of several sera with E1, and antibodies to the env31 region were also detected. This finding was surprising, since the env53 and env31 peptides had not shown any reactivity when coated directly to the solid phase.

Therefore peptides were synthesized using technology described by applicant previously (in WO 93/18054). The following peptides were synthesized:

peptide env35A-biotin NH<sub>2</sub>-SNSSEAADMIMHTPGCV-GKbiotin (SEQ ID NO 51) spanning amino acids 208 to 227 of the HCV polyprotein in the E1 region

peptide biotin-env53 ('epitope A') biotin-GG-ITGHRMAWDMMMNWSPTTAL-COOH (SEQ ID NO 52) spanning amino acids to 313 of 332 of the HCV polyprotein in the E1 region

peptide 1bE1 ('epitope B') H<sub>2</sub>N- YEVNRNVS GIYHVTNDCSNSSIVYEAAD-MIMHTPGCGK -biotin (SEQ ID NO 53)

spanning amino acids 192 to 228 of the HCV polyprotein in the E1 region and compared with the reactivities of peptides E1a-BB (biotin-GG-TPTVATRDKLTPATQLRRHIDL, SEQ ID NO 54) and E1b-BB (biotin-GG-TPTLAARDASVPTTTIRRHVDLL, SEQ ID NO 55) which are derived from the same region of sequences of genotype 1a and 1b respectively and which have been described at the IXth international virology meeting in Glasgow, 1993 ('epitope C').

Reactivity of a panel of HCV sera was tested on epitopes A, B and C and epitope B was also compared with env35A (of 47 HCV-positive sera, 8 were positive on epitope B and none reacted with env35A). Reactivity towards epitopes A, B, and C was tested directly to the biotinylated peptides (50  $\mu$ g/ml) bound to streptavidin-coated plates as described in example 6. Clearly, epitopes A and B were most reactive while epitopes C and env35A-biotin were much less reactive. The same series of patients that had been monitored for their reactivity towards the complete E1 protein (example 7.1.) was tested for reactivity towards epitopes A, B, and C. Little reactivity was seen to epitope C, while as shown in FIGS. 15, 16, 17, and 18, epitopes A and B reacted with the majority of sera. However, antibodies to the most reactive epitope (epitope A) did not seem to predict remission of disease, while the anti-1bE1 antibodies (epitope B) were present almost exclusively in long term responders at the start of IFN treatment. Therefore, anti-1bE1 (epitope B) antibodies and anti-env53 (epitope A) antibodies could be shown to be useful markers for prognosis of hepatitis C disease. The env53 epitope may be advantageously used for the detection of cross-reactive antibodies (antibodies that cross-react between major

genotypes) and antibodies to the env53 region may be very useful for universal E1 antigen detection in serum or liver tissue. Monoclonal antibodies that recognized the env53 region were reacted with a random epitope library. In 4 clones that reacted upon immunoscreening with the monoclonal antibody 5E1A10, the sequence —GWD— was present. Because of its analogy with the universal HCV sequence present in all HCV variants in the env53 region, the sequence AWD is thought to contain the essential sequence of the env53 cross-reactive murine epitope. The env31 clearly also contains a variable region which may contain an epitope in the amino terminal sequence —YQVRNSTGL— (SEQ ID NO 93) and may be useful for diagnosis. Env31 or E1-31 as shown in Table 3, is a part of the peptide 1bE1. Peptides E1-33 and E1-51 also reacted to some extent with the murine antibodies, and peptide E1-55 (containing the variable region 6 (V6); spanning amino acid positions 329–336) also reacted with some of the patient sera.

Anti-E2 antibodies clearly followed a different pattern than the anti-E1 antibodies, especially in patients with a long-term response to treatment. Therefore, it is clear that the decrease in anti-envelope antibodies could not be measured as efficiently with an assay employing a recombinant E1/E2 protein as with a single anti-E1 or anti-E2 protein. The anti-E2 response would clearly blur the anti-E1 response in an assay measuring both kinds of antibodies at the same time. Therefore, the ability to test anti-envelope antibodies to the single E1 and E2 proteins, was shown to be useful.

#### 7.4. Mapping of Anti-E2 Antibodies

Of the 24 anti-E2 Mabs only three could be competed for reactivity to recombinant E2 by peptides, two of which reacted with the HVRI region (peptides E2-67 and E2-69, designated as epitope A) and one which recognized an epitope competed by peptide E2-13B (epitope C). The majority of murine antibodies recognized conformational anti-E2 epitopes (FIG. 19). A human response to HVRI (epitope A), and to a lesser extent HVRII (epitope B) and a third linear epitope region (competed by peptides E2-23, E2-25 or E2-27, designated epitope E1 and a fourth linear epitope region (competed by peptide E2-17B, epitope D) could also frequently be observed, but the majority of sera reacted with conformational epitopes (FIG. 20). These conformational epitopes could be grouped according to their relative positions as follows: the IgG antibodies in the supernatant of hybridomas 15C8C 1, 12D11F1, 9G3E6, 8G10D1H9, 10D3C4, 4H6B2, 17F2C2, 5H6A7, 1 5B7A2 recognizing conformational epitopes were purified by means of protein A affinity chromatography and 1 mg/ml of the resulting IgG's were biotinylated in borate buffer in the presence of biotin. Biotinylated antibodies were separated from free biotin by means of gelfiltration chromatography. Pooled biotinylated antibody fractions were diluted 100 to 10,000 times. E2 protein bound to the solid phase was detected by the biotinylated IgG in the presence of 100 times the amount of non-biotinylated competing antibody and subsequently detected by alkaline phosphatase labeled streptavidin.

Percentages of competition are given in Table 6. Based on these results, 4 conformational anti-E2 epitope regions (epitopes F, G, H and I) could be delineated (FIG. 38). Alternatively, these Mabs may recognize mutant linear

epitopes not represented by the peptides used in this study. Mabs 4H682 and 10D3C4 competed reactivity of 16A6E7, but unlike 16A6E7, they did not recognize peptide E2-13B. These Mabs may recognize variants of the same linear epitope (epitope C) or recognize a conformational epitope which is sterically hindered or changes conformation after binding of 16A6E7 to the E2-13B region (epitope H).

### EXAMPLE 8

#### E1 Glycosylation Mutants

##### 8.1. Introduction

The E1 protein encoded by vvHCV10A, and the E2 protein encoded by vvHCV41 to 44 expressed from mammalian cells contain 6 and 11 carbohydrate moieties, respectively. This could be shown by incubating the lysate of vvHCV1 OA-infected or vvHCV44-infected RK13 cells with decreasing concentrations of glycosidases (PNGase F or Endoglycosidase H, (Boehringer Mannheim Biochemica) according to the manufacturer's instructions), such that the proteins in the lysate (including E1) are partially deglycosylated (FIG. 39 and 40, respectively).

Mutants devoid of some of their glycosylation sites could allow the selection of envelope proteins with improved immunological reactivity. For HIV for example, gp120 proteins lacking certain selected sugar-addition motifs, have been found to be particularly useful for diagnostic or vaccine purpose. The addition of a new oligosaccharide side chain in the hemagglutinin protein of an escape mutant of the A/Hong Kong/3/68 (H3N2) influenza virus prevents reactivity with a neutralizing monoclonal antibody (Skehel et al., 1984). When novel glycosylation sites were introduced into the influenza hemagglutinin protein by site-specific mutagenesis, dramatic antigenic changes were observed, suggesting that the carbohydrates serve as a modulator of antigenicity (Gallagher et al., 1988). In another analysis, the 8 carbohydrate-addition motifs of the surface protein gp70 of the Friend Murine Leukemia Virus were deleted. Although seven of the mutations did not affect virus infectivity, mutation of the fourth glycosylation signal with respect to the amino terminus resulted in a non-infectious phenotype (Kayman et al., 1991). Furthermore, it is known in the art that addition of N-linked carbohydrate chains is important for stabilization of folding intermediates and thus for efficient folding, prevention of misfolding and degradation in the endoplasmic reticulum, oligomerization, biological activity, and transport of glycoproteins (see reviews by Rose et al., 1988; Doms et al., 1993; Helenius, 1994).

After alignment of the different envelope protein sequences of HCV genotypes, it may be inferred that not all 6 glycosylation sites on the HCV subtype 1b E1 protein are required for proper folding and reactivity, since some are absent in certain (sub)types. The fourth carbohydrate motif (on Asn251), present in types 1b, 6a, 7, 8, and 9, is absent in all other types known today. This sugar-addition motif may be mutated to yield a type 1b E1 protein with improved reactivity. Also the type 2b sequences show an extra glycosylation site in the V5 region (on Asn299). The isolate S83, belonging to genotype 2c, even lacks the first carbohydrate motif in the V1 region (on Asn), while it is present on all other isolates (Stuyver et al., 1994). However, even among

the completely conserved sugar-addition motifs, the presence of the carbohydrate may not be required for folding, but may have a role in evasion of immune surveillance. Therefore, identification of the carbohydrate addition motifs which are not required for proper folding (and reactivity) is not obvious, and each mutant has to be analyzed and tested for reactivity. Mutagenesis of a glycosylation motif (NXS or NXT sequences) can be achieved by either mutating the codons for N, S, or T, in such a way that these codons encode amino acids different from N in the case of N, and/or amino acids different from S or T in the case of S and in the case of T. Alternatively, the X position may be mutated into P since it is known that NPS or NPT are not frequently modified with carbohydrates. After establishing which carbohydrate-addition motifs are required for folding and/or reactivity and which are not, combinations of such mutations may be made.

### 8.2. Mutagenesis of the E1 Protein

All mutations were performed on the E1 sequence of clone HCC110A (SEQ ID NO. 5). The first round of PCR was performed using sense primer 'GPT' (see Table 7) targeting the GPT sequence located upstream of the vaccinia 11K late promoter, and an antisense primer (designated GLY#, with # representing the number of the glycosylation site, see FIG. 41) containing the desired base change to obtain the mutagenesis. The six GLY# primers (each specific for a given glycosylation site) were designed such that:

Modification of the codon encoding for the N-glycosylated Asn (AAC or AAT) to a Gin codon (CAA or CAG). Glutamine was chosen because it is very similar to asparagine (both amino acids are neutral and contain non-polar residues, glutamine has a longer side chain (one more —CH<sub>2</sub>— group).

The introduction of silent mutations in one or several of the codons downstream of the glycosylation site, in order to create a new unique or rare (e.g. a second SmaI site for E1 Gly5) restriction enzyme site. Without modifying the amino acid sequence, this mutation will provide a way to distinguish the mutated sequences from the original E1 sequence (pvHCV-10A) or from each other (FIG. 41). This additional restriction site may also be useful for the construction of new hybrid (double, triple, etc.) glycosylation mutants.

18 nucleotides extend 5' of the first mismatched nucleotide and 12 to 16 nucleotides extend to the 3' end. Table 7 depicts the sequences of the six GLY# primers overlapping the sequence of N-linked glycosylation sites.

For site-directed mutagenesis, the 'mispriming' or 'overlap extension' (Horton, 1993) was used. The concept is illustrated in FIGS. 42 and 43. First, two separate fragments were amplified from the target gene for each mutated site. The PCR product obtained from the 5' end (product GLY#) was amplified with the 5' sense GPT primer (see Table 7) and with the respective 3' antisense GLY# primers. The second fragment (product OVR#) was amplified with the 3' antisense TK<sub>R</sub> primer and the respective 5' sense primers (OVR# primers, see Table 7, FIG. 43).

The OVR# primers target part of the GLY# primer sequence. Therefore, the two groups of PCR products share an overlap region of identical sequence. When these intermediate products are mixed (GLY-1 with OVR-1, GLY-2

with OVR-2, etc.), melted at high temperature, and reannealed, the top sense strand of product GLY# can anneal to the antisense strand of product DVR# (and vice versa) in such a way that the two strands act as primers for one another (see FIG. 42.B.). Extension of the annealed overlap by Taq polymerase during two PCR cycles created the full-length mutant molecule E1 Gly#, which carries the mutation destroying the glycosylation site number #. Sufficient quantities of the E1 GLY# products for cloning were generated in a third PCR by means of a common set of two internal nested primers. These two new primers are respectively overlapping the 3' end of the vaccinia 11K promoter (sense GPT-2 primer) and the 5' end of the vaccinia thymidine kinase locus (antisense TK<sub>R</sub>-2 primer, see Table 7). All PCR conditions were performed as described in Stuyver et al. (1993).

Each of these PCR products was cloned by EcoRI/BamHI cleavage into the EcoRI/BamHI-cut vaccinia vector containing the original E1 sequence (pvHCV-10A).

The selected clones were analyzed for length of insert by EcoRII/BamH I cleavage and for the presence of each new restriction site. The sequences overlapping the mutated sites were confirmed by double-stranded sequencing.

### 8.3. Analysis of E1 Glycosylation Mutants

Starting from the 6 plasmids containing the mutant E1 sequences as described in example 8.2, recombinant vaccinia viruses were generated by recombination with wt vaccinia virus as described in example 2.5. Briefly, 175 cm<sup>2</sup>-flasks of subconfluent RK13 cells were infected with the 6 recombinant vaccinia viruses carrying the mutant E1 sequences, as well as with the vvHCV-10A (carrying the non-mutated E1 sequence) and wt vaccinia viruses. Cells were lysed after 24 hours of infection and analyzed on western blot as described in example 4 (see FIG. 44A). All mutants showed a faster mobility (corresponding to a smaller molecular weight of approximately 2 to 3 kDa) on SDS-PAGE than the original E1 protein; confirming that one carbohydrate moiety was not added. Recombinant viruses were also analyzed by PCR and restriction enzyme analysis to confirm the identity of the different mutants. FIG. 44B shows that all mutants (as shown in FIG. 41) contained the expected additional restriction sites. Another part of the cell lysate was used to test the reactivity of the different mutant by ELISA. The lysates were diluted 20 times and added to micro well plates coated with the lectin GNA as described in example 6. Captured (mutant) E1 glycoproteins were left to react with 20-times diluted sera of 24 HCV-infected patients as described in example 6. Signal to noise (S/N) values (OD of GLY#10D of wt) for the six mutants and E1 are shown in Table 8. The table also shows the ratios between S/N values of GLY# and E1 proteins. It should be understood that the approach to use cell lysates of the different mutants for comparison of reactivity with patient sera may result in observations that are the consequence of different expression levels rather than reactivity levels. Such difficulties can be overcome by purification of the different mutants as described in example 5, and by testing identical quantities of all the different E1 proteins. However, the results shown in table 5 already indicate that removal of the 1st (GLY1), 3rd (GLY3), and 6th (GLY6) glycosylation motifs reduces reactivity of some sera, while removal of the 2nd and 5th site



does not. Removal of GLY4 seems to improve the reactivity of certain sera. These data indicate that different patients react differently to the glycosylation mutants of the present invention. Thus, such mutant E1 proteins may be useful for the diagnosis (screening, confirmation, prognosis, etc.) and prevention of HCV disease.

#### EXAMPLE 9

##### Expression of HCV E2 Protein in Glycosylation-Deficient Yeasts

The E2 sequence corresponding to clone HCCL41 was provided with the  $\alpha$ -mating factor pre/pro signal sequence, inserted in a yeast expression vector and *S. cerevisiae* cells transformed with this construct secreted E2 protein into the growth medium. It was observed that most glycosylation sites were modified with high-mannose type glycosylations upon expression of such a construct in *S. cerevisiae* strains (FIG. 45). This resulted in a too high level of heterogeneity and in shielding of reactivity, which is not desirable for either vaccine or diagnostic purposes. To overcome this problem, *S. cerevisiae* emutants with modified glycosylation pathways were generated by means of selection of vanadate-resistant clones. Such clones were analyzed for modified glycosylation pathways by analysis of the molecular weight and heterogeneity of the glycoprotein invertase. This allowed us to identify different glycosylation deficient *S. cerevisiae* mutants. The E2 protein was subsequently expressed in some of the selected mutants and left to react with a monoclonal antibody as described in example 7, on western blot as described in example 4 (FIG. 46).

#### EXAMPLE 10

##### General Utility

The present results show that not only a good expression system but also a good purification protocol are required to

reach a high reactivity of the HCV envelope proteins with human patient sera. This can be obtained using the proper HCV envelope protein expression system and/or purification protocols of the present invention which guarantee the conservation of the natural folding of the protein and the purification protocols of the present invention which guarantee the elimination of contaminating proteins and which preserve the conformation, and thus the reactivity of the HCV envelope proteins. The amounts of purified HCV envelope protein needed for diagnostic screening assays are in the range of grams per year. For vaccine purposes, even higher amounts of envelope protein would be needed. Therefore, the vaccinia virus system may be used for selecting the best expression constructs and for limited upscaling, and large-scale expression and purification of single or specific oligomeric envelope proteins containing high-mannose carbohydrates may be achieved when expressed from several yeast strains. In the case of hepatitis B for example, manufacturing of HBsAg from mammalian cells was much more costly compared with yeast-derived hepatitis B vaccines.

The purification method disclosed in the present invention may also be used for 'viral envelope proteins' in general. Examples are those derived from Flaviviruses, the newly discovered GB-A, GB-B and GB-C Hepatitis viruses, Pestiviruses (such as Bovine viral Diarrhoea Virus (BVDV), Hog Cholera Virus (HCV), Border Disease Virus (BDV)), but also less related viruses such as Hepatitis B Virus (mainly for the purification of HBsAg).

The envelope protein purification method of the present invention may be used for intra- as well as extracellularly expressed proteins in lower or higher eukaryotic cells or in prokaryotes as set out in the detailed description section.

TABLE 1

Recombinant vaccinia plasmids and viruses				
Plasmid name	Name	cDNA subclone construction	Length (nt/aa)	Vector used for insertion
pvHCV-13A	E1s	EcoR I - Hind III	472/157	pgptATA-18
pvHCV-12A	E1s	EcoR I - Hind III	472/158	pgptATA-18
pvHCV-9A	E1	EcoR I - Hind III	631/211	pgptATA-18
pvHCV-11A	E1s	EcoR I - Hind III	625/207	pgptATA-18
pvHCV-17A	E1s	EcoR I - Hind III	625/208	pgptATA-18
pvHCV-10A	E1	EcoR I - Hind III	783/262	pgptATA-18
pvHCV-18A	COREs	Acc I (KI) - EcoR I (KI)	403/130	pgptATA-18
pvHCV-34	CORE	Acc I (KI) - Fsp I	595/197	pgptATA-18
pvHCV-33	CORE-E1	Acc I (KI)	1150/380	pgptATA-18
pvHCV-35	CORE-E1b.his	EcoR I - BamH I (KI)	1032/352	pMS-66
pvHCV-36	CORE-E1n.his	EcoR I - Nco I (KI)	1106/376	pMS-66
pvHCV-37	E1A	Xma I - BamH I	711/239	pvHCV-10A
pvHCV-38	E1As	EcoR I - BstE II	553/183	pvHCV-11A
pvHCV-39	E1Ab	EcoR I - BamH I	960/313	pgsATA-18
pvHCV-40	E1Ab.his	EcoR I - BamH I (KI)	960/323	pMS-66
pvHCV-41	E2bs	BamH I (KI)-AlwN I (T4)	1005/331	pgsATA-18
pvHCV-42	E2bs.his	BamH I (KI)-AlwN I (T4)	1005/341	pMS-66
pvHCV-43	E2ns	Nco I (KI) - AlwN I (T4)	932/314	pgsATA-18
pvHCV-44	E2ns.his	Nco I (KI) - AlwN I (T4)	932/321	pMS-66
pvHCV-62	E1s (type 3a)	EcoR I - Hind III	625/207	pgsATA-18
pvHCV-63	E1s (type 5)	EcoR I - Hind III	625/207	pgsATA-18
pvHCV-64	E2	BamH I - Hind III	1410/463	pgsATA-18

TABLE 1-continued

Recombinant vaccinia plasmids and viruses				
Plasmid name	Name	cDNA subclone construction	Length (nt/aa)	Vector used for insertion
pvHCV-65	E1-E2	BamH I - Hind III	2072/691	pvHCV-10A
pvHCV-66	CORE-E1-E2	BamH I - Hind III	2427/809	pvHCV-33
pvHCV-81	E1*-GLY 1	EcoRI - BamH I	783/262	pvHCV-10A
pvHCV-82	E1*-GLY 2	EcoRI - BamH I	783/262	pvHCV-10A
pvHCV-83	E1*-GLY 3	EcoRI - BamH I	783/262	pvHCV-10A
pvHCV-84	E1*-GLY 4	EcoRI - BamH I	783/262	pvHCV-10A
pvHCV-85	E1*-GLY 5	EcoRI - BamH I	783/262	pvHCV-10A
pvHCV-86	E1*-GLY 6	EcoRI - BamH I	783/262	pvHCV-10A

nt: nucleotide

aa: aminoacid

KI: Klenow DNA Pol filling

T4:T4 DNA Pol filling

Position: aminoacid position in the HCV polyprotein sequence

TABLE 2

Summary of anti-E1 tests			
S/N $\pm$ SD (mean anti-E1 titer)			
	Start of treatment	End of treatment	Follow-up
LTR	6.94 $\pm$ 2.29 (1:3946)	4.48 $\pm$ 2.69 (1:568)	2.99 $\pm$ 2.69 (1:175)
NR	5.77 $\pm$ 3.77 (1:1607)	5.29 $\pm$ 3.99 (1:1060)	6.08 $\pm$ 3.73 (1:1978)

LTR: Long-term, sustained response for more than 1 year

NR: No response, response with relapse, or partial response

TABLE 3

Synthetic peptides for competition studies				
PROTEIN	PEPTIDE	AMINO ACID SEQUENCE	POSITION	SEQ ID
E1	E1-31	LLSCLTVPASAYQVRNSTGL	181-200	56
	E1-33	QVRNSTGLYHVTNDPCNSSI	193-212	57
	E1-35	NDCPNSIVYEAHDAILHTP	205-224	58
	E1-35A	SNSIVYEAADMIMHTPGCV	208-227	59
	E1-37	HDAILHTPGCVPCVREGNVS	217-236	60
	E1-39	CVREGNVSRCWVAMTPTVAT	229-248	61
	E1-41	AMTPTVATRDGKLPATQLRR	241-260	62
	E1-43	LPATQLRRHIDLVGVSATLC	253-272	63
	E1-45	LVGSATLCSALYVGDLCGSV	265-284	64
	E1-49	QLFTFSRRHWTQGCNCSI	289-308	65
	E1-51	TQGCNCSIYPGHITGHRMAW	301-320	66
	E1-53	ITGHRMAWDMMNWSPPTAAL	313-332	67
	E1-55	NWSEPTAALVMAQLLRIPQAI	325-344	68
	E1-57	LLRJPQAILDMIAGAHWGL	337-356	69
	E1-59	AGAHWGLAGIAYFSMVGNM	349-368	70

TABLE 3-continued

Synthetic peptides for competition studies				
PROTEIN	PEPTIDE	AMINO ACID SEQUENCE	POSITION	SEQ ID
E2	E1-63	VVLLLFAGVDAETIVSGGQA	373-392	71
	E2-67	SGLVSLFTPGAKQNJQLINT	397-416	72
	E2-69	QNIQLINTNGSWHINSTALN	409-428	73
	E2- $\$$ 3B	LNCNESLNTGWWLAGLIYQHK	427-446	74
	E2- $\$$ 1B	AGLIYQHKFNSSGCPERLAS	439-458	75
	E2-1B	GCPERLASCRPLTDFDQGWG	451-470	76
	E2-3B	TDFDQGWGPISYANGSGPDQ	463-482	77
	E2-5B	ANGSGPDQRPCYWHYPPKPC	475-494	78
	E2-7B	WHYPPKPCGIVPAKSVCGPV	487-506	79
	E2-9B	AKSVCGPVYCFTPSPVVVGT	499-518	80
	E2-11B	PSPVVVGTTRDSGAPTYSWG	511-530	81
	E2-13B	GAPTYSWGENTDVFLNNT	523-542	82
	E2-17B	GNWFCTWMNSTGFTKVCGA	547-566	83
	E2-19B	GFTKVCGAPPVCIGGAGNNT	559-578	84
	E2-21	IGGAGNNTLHCPTDCFRKHP	571-590	85
	E2-23	TDCFRKHPDATYSRCGSGPW	583-602	86
	E2-25	SRCSGSGPWITPRCLVDYPYR	595-614	87
E2-27	CLVDYPYRLWHYPCNTYTI	607-626	88	
E2-29	PCTINYTIFKIRMYVGGVEH	619-638	89	
E2-31	MYVGGVEHRLEAACNWTPE	631-650	90	
E2-33	ACNWTPEGERCDLEDRDRSEL	643-662	91	
E2-35	EDRDRSELSPLLTTTQWQV	655-674	92	

TABLE 4

Change of Envelope Antibody levels over time (complete study, 28 patients)								
Wilcoxon Signed Rank test (P values)	E1Ab NR All	E1Ab NR type 1b	E1Ab NR type 3a	E1Ab LTR II	E1Ab LTR type 1b	E1Ab LTR type 3a	E2Ab NR All	E1Ab LTR All
End of therapy*	0.1167	0.2604	0.285	0.0058**	0.043**	0.0499**	0.0186**	0.0640
6 months follow up*	0.86	0.7213	0.5930	0.0047**	0.043**	0.063	0.04326	0.0464**
12 months follow up*	0.7989	0.3105	1	0.0051**	0.0679	0.0277**	0.0869	0.0058**

\*Data were compared with values obtained at initiation of therapy

\*\*P values &lt; 0.05

TABLE 5

Difference between LTR and NR (complete study)					
	E1Ab	E1Ab	E1Ab	E1Ab	E2Ab
Mann-Whitney	S/N	titers	S/N	S/N	S/N
U test (P values)	All	all	type 1b	type 3a	All
Initiation of therapy	0.0257*		0.05*	0.68	0.1078
End of therapy	0.1742				0.1295
6 months follow up,	1		0.6099	0.425	0.3081
12 months follow up	0.67		0.23	0.4386	0.6629

\*P values &lt; 0.05

TABLE 6

Competition experiments between murine E2 monoclonal antibodies										
Decrease (%) of anti-E2 reactivity of biotinylated anti-E2 mabs										
	17H10F4D10	2F10H10	16A6E7	10D3C4	4H6B2	17C2F2	9G3E6	12D11F1	15C8C1	8G10D1H9
<u>competitor</u>										
17H10F4D10		62	10	ND	11	ND	5	6	30	ND
2F10H10	90	—	1	ND	30	ND	0	4	12	ND
16A6E7	ND	ND	—	ND	ND	ND	ND	ND	ND	ND
10D3C4	11	50	92	—	94	26	28	43	53	30
4H6B2	ND	ND	82	ND	—	ND	ND	ND	ND	ND
17C2F2	2	ND	75	ND	56	—	11	10	0	0
9G3E6	ND	ND	68	ND	11	ND	—	60	76	ND
12D11F1	ND	ND	26	ND	13	ND	ND	—	88	ND
15C8C1	ND	ND	18	ND	10	ND	ND	—	—	ND
8G10D1H9	2	2	11	ND	15	ND	67	0.82	81	—
<u>competitor controls</u>										
15B7A2	0	0	9	15	10	9	0	0	0	5
5H6A7	0	2	0	12	8	0	0	4	0	0
23C12H9	ND	ND	2	12	ND	4	ND	ND	ND	2

ND, not done

TABLE 7

Primers			
SEQ ID NO. 96	GPT	5'	<u>-GTTTAAACCACTGCATGATG-3'</u>
SEQ ID NO. 97	TK <sub>R</sub>	5'	<u>-GTCCCATCGAGTGGCGCTAC-3'</u>
SEQ ID NO. 98	GLY1	5'	<u>-CGTGACATGGTACATTCGGACACTTGGCGCACTTCATAAGCGGA-3'</u>
SEQ ID NO. 99	GLY2	5'	<u>-TGCCCTACACAAATGGAGCTCTGGGACGAGTCGTTCTGTGAC-3'</u>
SEQ ID NO. 100	GLY3	5'	<u>-TACCCAGCAGCGGGAGCTCTGTGTGCTCCCGAACGCAGGGCAC-3'</u>
SEQ ID NO. 101	GLY4	5'	<u>-TGTCGTGGTGGGACGGAGGCTGCCTAGCTGCGAGCGTGGG-3'</u>
SEQ ID NO. 102	GLY5	5'	<u>-CGTTATGTGGCCCGGGTAGATTGAGCACTGGCAGTCTGCACCCGTCTC-3'</u>
SEQ ID NO. 103	GLY6	5'	<u>-CAGGGCCGTTGTAGGCCCTCCACTGCATCATATCCCAAGC-3'</u>
SEQ ID NO. 104	OVR1	5'	<u>-CCGGAATGTACCATGTCACGAACGAC-3'</u>
SEQ ID NO. 105	OVR2	5'	<u>-GCTCCATTGTGTATGAGGACGGC-3'</u>
SEQ ID NO. 106	OVR3	5'	<u>-GAGCTCCCGCTGCTGGGTAGCGC-3'</u>
SEQ ID NO. 107	OVR4	5'	<u>-CCTCCGTCCCAACACGACAATACG-3'</u>
SEQ ID NO. 108	OVR5	5'	<u>-CTACCCGGCCACATAACGGGTACCG-3'</u>
SEQ ID NO. 109	OVR6	5'	<u>-GGAGGCCCTACAACGGCCCTGGTGG-3'</u>
SEQ ID NO. 110	GPT-2	5'	<u>-TTCTATCGATTAATAGAAATTC-3'</u>
SEQ ID NO. 111	TK <sub>R</sub> -2	5'	<u>-GCCATACGCTCACAGCCGATCCC-3'</u>

nucleotides underlined represent additional restriction site  
nucleotides in bold represent mutations with respect to the original HCC110A sequence

TABLE 8

Analysis of EI glycosylation mutants by ELISA

	SERUM												Sum	Average S/N	S/N
	1	2	3	4	5	6	7	8	9	10	11	12			
SN GLY 1	1.802462	2.120971	1.403871	1.205597	2.120191	2.866913	1.950345	1.866183	1.730193	2.468162	1.220654	1.629403			
SN GLY 2	2.400795	1.76818	2.325495	2.639308	2.459019	5.043993	2.146302	1.595477	1.688973	2.482212	1.467582	2.070524			
SN GLY 3	1.642718	1.71547	2.261646	2.354748	1.591818	4.833742	1.96692	1.482099	1.602222	1.464216	1.721164	1.464216			
SN GLY 4	2.578154	3.824038	3.874605	1.499387	3.15	4.71302	4.198751	3.959542	3.710507	5.170841	4.250784	3.955153			
SN GLY 5	2.482051	1.793761	2.409344	2.627358	1.715311	4.964765	2.13912	1.576336	1.708937	3.021807	1.562092	2.07278			
SN GLY 6	2.031487	1.495737	2.131613	2.527925	2.494833	4.784027	2.02069	1.496489	1.704976	2.677757	1.529608	1.744221			
SN EI	2.828205	2.227036	2.512792	2.790881	3.131579	4.869128	2.287753	1.954198	1.805556	2.616822	1.55719	2.593886			
													Sum	Average S/N	S/N
SN GLY 1	5.685561	3.233684	3.763498	1.985105	2.317721	6.675179	1.93476	2.47171	4.378633	1.188748	2.158889	1.786992	59.88534	2.495223	
SN GLY 2	7.556682	2.567613	3.621928	3.055649	2.93792	7.65433	2.127712	2.921288	4.680101	1.158781	1.661914	1.632785	69.65243	2.982185	
SN GLY 3	7.930538	2.763055	3.016099	2.945628	2.515305	5.775357	1.980185	2.557384	4.268633	0.97767	1.336775	1.20376	62.89872	2.587447	
SN GLY 4	8.176816	6.561122	5.707668	5.684498	5.604813	6.4125	3.813321	3.002535	4.293038	2.393011	3.68213	2.481585	102.6978	4.279076	
SN GLY 5	8.883408	2.940334	3.125561	3.338912	2.642224	5.424107	3.126761	4.64557	1.153656	1.817901	1.638211	1.638211	69.26511	2.886846	
SN GLY 6	8.005561	2.499952	2.621704	2.572385	2.363301	5.194107	1.586716	2.665433	2.781063	1.280743	1.475862	1.716423	61.32181	2.555075	
SN EI	8.825112	3.183771	3.067265	3.280335	2.980354	7.191964	2.771218	3.678068	5.35443	1.167286	2.083333	1.78252	76.54068	3.189195	
													Sum	Average S/N	S/N
GLY 1/E1	0.675316	0.952374	0.55869	0.431977	0.677036	0.588794	0.852516	0.954961	0.958261	0.94319	0.783882	0.628171			
GLY 2/E1	0.848876	0.793961	0.925463	0.94569	0.785233	1.035913	0.93817	0.816436	0.935431	0.94856	0.942455	8.798232			
GLY 3/E1	0.580834	0.770296	0.900053	0.84373	0.508312	0.992733	0.859761	0.758418	0.887385	0.837488	0.940294	8.663547			
GLY 4/E1	0.911587	1.717097	1.541952	0.537245	1.005882	0.967939	1.8355317	2.026172	2.05505	1.976	2.72978	1.524798			
GLY 5/E1	0.877607	0.805447	0.958831	0.941488	0.547746	1.019642	0.935031	0.806641	0.946488	1.154762	1.883148	8.799182			
GLY 6/E1	0.718296	0.671626	0.848305	0.90578	0.796669	0.982522	0.883264	0.765781	0.944294	1.023286	0.982288	8.672435			
													Sum	Average S/N	S/N
GLY 1/E1	0.644248	1.815652	1.226988	1.605153	0.777666	0.928144	0.698162	0.672013	0.817759	1.018386	1.830267	0.957628	19.36524	0.806885	
GLY 2/E1	8.85627	6.806469	1.188833	0.931505	8.984377	1.064289	8.76779	0.794245	0.874061	0.98586	8.797719	8.915998	21.67384	0.903077	
GLY 3/E1	0.898633	0.867856	8.983319	0.897966	0.843962	0.883029	0.714554	0.695306	0.797215	0.837558	0.641652	0.675314	19.19921	0.799967	
GLY 4/E1	0.92654	2.060802	1.860833	1.732902	1.880587	0.89162	8.8016335	8.816335	8.801773	2.058864	1.767422	1.392178	36.38592	1.51608	
GLY 5/E1	1.006606	0.923538	1.019006	1.817857	0.890574	8.75419	0.881491	0.850109	8.867612	8.988323	0.872593	0.919042	21.78679	8.907783	
GLY 6/E1	0.907134	0.785217	8.854737	0.784184	0.79296	8.72221	0.543702	8.724683	0.519395	1.097197	0.70803	0.962919	19.59691	0.816538	

- Bailey, J. and Cole, R. (1959) *J. Biol. Chem.* 234, 1733-1739.
- Bailou, L., Hitzeman, R., Lewis, M. & Ballou, C. (1991) *PNAS* 88, 3209-3212.
- Benesch, R., Benesch, R. E. Gutcho, M. & Lanter, L. (1956) *Science* 123, 981.
- Cavins, J. & Friedman, (1970) *Anal. Biochem.* 35, 489.
- Cleland, W. (1964) *Biochemistry* 3, 480.
- Creighton., E. (1988) *BioEssays* 8, 57.
- Darbre, A., John Wiley & Sons Ltd. (1987) *Practical Protein Chemistry—A Handbook*.
- Darbre, A., John Wiley & Sons Ltd. (1987) *Practical Proteinchemistry* p. 69-79.
- Doms et al., (1993), *Virology* 193, 545-562.
- Ellman, G. (1959) *Arch. Biochem. Biophys.* 82, 70.
- Falkner, F. & Moss, B. (1988) *J. Virol.* 62, 1849-1854.
- Friedman, M., & Krull, (1969) *Biochem. Biophys. Res. Commun.* 37, 630.
- Gallagher J. (1988) *J. Cell Biol.* 107, 2059-2073.
- Glazer, A., Delange, R., Sigman, D. (1975) North Holland publishing company, Elsevier, Biomedical, Part : Modification of protein (p. 116).
- Graham, F. & van der Eb, A. (1973) *Virology* 52, 456-467.
- Grakoui et al. (1993) *Journal of Virology* 67:1385-1395.
- Grassetti, D. & Murray, J. (1969) *Analyt. Chim. Acta.* 46, 139.
- Grassetti, D. & Murray, J. (1967) *Arch. Biochem Biophys.* 119, 41.
- Helenius, *Mol. Biol. Cell* (1994), 5: 253-265.
- Hijikata, M., Kato, N., Ootusyama, Y., Nakagawa, M. & Shimotohno, K. (1991) *Proc. Natl. Acad. Sci. U.S.A.* 88(13):5547-51.
- Hochuli, E., Bannwarth, W., Dëbeli, H., Gentz, R., Stüber, D. (1988) *Biochemistry* 88, 8976.
- Hsu, H., Donets, M., Greenberg, H. & Feinstone, S. (1993) *Hepatology* 17:763-771.
- Inoue, Y., Suzuki, R., Matsuura, Y., Harada, S., Chiba, J., Watanabe, Y., Saito, I. & Miyamura, T. (1992) *J. Gen. Birol.* 73:2541-2154.
- Jankecht, R., de Martynoff, G. et al., (1991) *Proc. Natl. Acad. Sci.* 88, 8972-8976.
- Kayman (1991) *J. Virology* 65, 5323-5332.
- Kato, N., Oostuyama, Y., Tanaka, T., Nakagawa, M., Muraio, K., Ohkoshi, S., Hijikata, M., Shimitohno, K. (1992) *Virus Res.* 22:107-123.
- Kriskern, P., Hagopian, A., Burke, P., Schultz, L., Montgomery, D., Hurmi, W., Yu Ip. C., Schulman, C., Maigetter, R., Wampler, D., Kubek, C., Sitrin, R., West, D., Ellis, R., Miller, W. (1994) *Vaccine* 12:1021-1025.
- Kohara, M., Tsukiyama-Kohara, K., Maki, N., Asano, K., Yoshizawa, K., Miki, K., Tanaka, S., Hattori, N., Matsuura, Y., Saito, I., Miyamura, T. & Nomoto, A. (1992) *J. Gen. Virol.* 73:2313-2318.

- Mackett, M., Smith, G. & Moss, B. (1985) In: *DNA cloning: a practical approach* (Ed. Glover, D.) IRL Press, Oxford.
- Mackett, M., & Smith, G. (1986) *J. Gen. Virol.* 67, 2067-2082.
- 5 Mackett, M., Smith, C & Moss, B. (1984) *J. Virol.* 49, 857-864.
- Mackett, M., Smith, G. & Moss, B. (1984) *Proc. Natl. Acad. Sci. USA* 79, 7415-7419.
- Means, G. (1971) Holden Day, Inc.
- Means, G. & Feeney, R. (1971) Holden Day p.105 & p. 217.
- 10 Mita, E., Hayashi, N., Ueda, K., Kasahara, A., Fusamoto, H., Takamizawa, A., Matsubara, K., Okayama, H. & Kamada T. (1992) *Biochem. Biophys. Res. Comm.* 183:925-930.
- Moore, S. (1963) *J. Biol. Chem.* 238, 235-237.
- Okamoto, H., Okada, S., Sugiyama, Y., Yotsumoto, S., Tanaka, T., Yoshizawa, H., Tsuda, F., Miyakawa, Y. & Mayumi, M. (1990) *Jpn. J. Exp. Med.* 60:167-177.
- 15 Panicali & Paoletti (1982) *Proc. Natl. Acad. Sci. USA* 79, 4927-4931.
- Piccini, A., Perkus, M. & Paoletti, E. (1987) *Meth. Enzymol.* 153, 545-563.
- 20 Rose (1988) *Annu. Rev. Cell Biol.* 1988, 4: 257-288;
- Ruegg, V. and Rudinger, J. (1977) *Methods Enzymol.* 47, 111-116.
- Shan, S. & Wong (1993) *CRC-press* p. 30-33.
- 25 Spaete, R., Alexander, D., Rugroden, M., Choo, Q., Berger, K., Crawford, K., Kuo, C., Leng, S., Lee, C., Ralston, R., et al. (1992) *Virology* 188(2):819-30.
- Skehel, J., (1984) *Proc. Natl. Acad. Sci. USA* 81, 1179-1783.
- 30 Stunnenberg, H., Lange, H., Philipson, L., Miltenburg, R. & van der Vliet, R. (1988) *Nucl. Acids Res.* 16, 2431-2444.
- Stuyver, L., Van Arnhem, W., Wyseur, A., DeLeys, R. & Maertens, G. (1993a) *Biochem. Biophys. Res. Commun.* 192, 635-641.
- 35 Stuyver, L., Rossau, R., Wyseur, A., Duhamel, M., Vanderborgh, B., Van Heuverswyn, H., & Maertens, G. (1993b) *J. Gen. Virol.* 74, 1093-1102.
- Stuyver, L., Van Arnhem, W., Wyseur, A., Hernandez, F., Delaporte, E., Maertens, G. (1994), *Proc. Natl. Acad. Sci. USA* 91:10134-10138.
- 40 Weil, L. & Seibler, S. (1 961) *Arch. Biochem. Biophys.* 95, 470.
- Yokosuka, O., Ito, Y., Imazeki, F., Ohto, M. & Omata, M. (1992) *Biochem. Biophys. Res. Commun.* 189:565-571.
- 45 Miller P, Yano J, Yano E, Carroll C, Jayaram K, Ts'o P (1979) *Biochemistry* 18:51 34-43.
- Nielsen P, Egholm M, Berg R, Buchardt O (1991) *Science* 254:1497-500.
- Nielsen P, Egholm M, Berg R, Buchardt O (1993) *Nucleic-Acids-Res.* 21:197-200.
- 50 Asseline U, Delarue M, Lancelot G, Toulme F, Thuong N (1984) *Proc. Natl. Acad. Sci. USA* 81:3297-301.
- Matsukura M, Shinozuka K, Zon G, Mitsuya H, Reitz M, Cohen J, Broder S (1987) *Proc. Natl. Acad. Sci. USA* 84:7706-10.

## SEQUENCE LISTING

(1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 111

(2) INFORMATION FOR SEQ ID NO: 1:

-continued

---

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 21 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

GGCATGCAAG CTTAATTAAT T 21

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 68 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

CCGGGGAGGC CTGCACGTGA TCGAGGGCAG ACACCATCAC CACCATCACT AATAGTTAAT 60  
 TAACTGCA 68

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 642 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 1..639

(ix) FEATURE:  
 (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 1..636

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

ATG CCC GGT TGC TCT TTC TCT ATC TTC CTC TTG GCT TTA CTG TCC TGT 48  
 Met Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu Leu Ser Cys  
 1 5 10 15

CTG ACC ATT CCA GCT TCC GCT TAT GAG GTG CGC AAC GTG TCC GGG ATG 96  
 Leu Thr Ile Pro Ala Ser Ala Tyr Glu Val Arg Asn Val Ser Gly Met  
 20 25 30

TAC CAT GTC ACG AAC GAC TGC TCC AAC TCA AGC ATT GTG TAT GAG GCA 144  
 Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val Tyr Glu Ala  
 35 40 45

GCG GAC ATG ATC ATG CAC ACC CCC GGG TGC GTG CCC TGC GTT CGG GAG 192  
 Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys Val Arg Glu  
 50 55 60

-continued

AAC AAC TCT TCC CGC TGC TGG GTA GCG CTC ACC CCC ACG CTC GCA GCT Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr Leu Ala Ala 65 70 75 80	240
AGG AAC GCC AGC GTC CCC ACC ACG ACA ATA CGA CGC CAC GTC GAT TTG Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His Val Asp Leu 85 90 95	288
CTC GTT GGG GCG GCT GCT CTC TGT TCC GCT ATG TAC GTG GGG GAT CTC Leu Val Gly Ala Ala Ala Leu Cys Ser Ala Met Tyr Val Gly Asp Leu 100 105 110	336
TGC GGA TCT GTC TTC CTC GTC TCC CAG CTG TTC ACC ATC TCG CCT CGC Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Ile Ser Pro Arg 115 120 125	384
CGG CAT GAG ACG GTG CAG GAC TGC AAT TGC TCA ATC TAT CCC GGC CAC Arg His Glu Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His 130 135 140	432
ATA ACA GGT CAC CGT ATG GCT TGG GAT ATG ATG ATG AAC TGG TCG CCT Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp Ser Pro 145 150 155 160	480
ACA ACG GCC CTG GTG GTA TCG CAG CTG CTC CGG ATC CCA CAA GCT GTC Thr Thr Ala Leu Val Val Ser Gln Leu Leu Arg Ile Pro Gln Ala Val 165 170 175	528
GTG GAC ATG GTG GCG GGG GCC CAT TGG GGA GTC CTG GCG GGC CTC GCC Val Asp Met Val Ala Gly Ala His Trp Gly Val Leu Ala Gly Leu Ala 180 185 190	576
TAC TAT TCC ATG GTG GGG AAC TGG GCT AAG GTT TTG ATT GTG ATG CTA Tyr Tyr Ser Met Val Gly Asn Trp Ala Lys Val Leu Ile Val Met Leu 195 200 205	624
CTC TTT GCT CTC TAATAG Leu Phe Ala Leu 210	642

## (2) INFORMATION FOR SEQ ID NO: 4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 212 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Met Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu Leu Ser Cys 1 5 10 15
Leu Thr Ile Pro Ala Ser Ala Tyr Glu Val Arg Asn Val Ser Gly Met 20 25 30
Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val Tyr Glu Ala 35 40 45
Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys Val Arg Glu 50 55 60
Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr Leu Ala Ala 65 70 75 80
Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His Val Asp Leu 85 90 95
Leu Val Gly Ala Ala Ala Leu Cys Ser Ala Met Tyr Val Gly Asp Leu 100 105 110
Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Ile Ser Pro Arg 115 120 125
Arg His Glu Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His 130 135 140

-continued

Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp Ser Pro  
 145 150 155 160  
 Thr Thr Ala Leu Val Val Ser Gln Leu Leu Arg Ile Pro Gln Ala Val  
 165 170 175  
 Val Asp Met Val Ala Gly Ala His Trp Gly Val Leu Ala Gly Leu Ala  
 180 185 190  
 Tyr Tyr Ser Met Val Gly Asn Trp Ala Lys Val Leu Ile Val Met Leu  
 195 200 205  
 Leu Phe Ala Leu  
 210

## (2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 795 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 1..792

(ix) FEATURE:  
 (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 1..789

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

ATG TTG GGT AAG GTC ATC GAT ACC CTT ACA TGC GGC TTC GCC GAC CTC	48
Met Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu	
1 5 10 15	
GTG GGG TAC ATT CCG CTC GTC GGC GCC CCC CTA GGG GGC GCT GCC AGG	96
Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg	
20 25 30	
GCC CTG GCG CAT GGC GTC CGG GTT CTG GAG GAC GGC GTG AAC TAT GCA	144
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala	
35 40 45	
ACA GGG AAT TTG CCC GGT TGC TCT TTC TCT ATC TTC CTC TTG GCT TTG	192
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu	
50 55 60	
CTG TCC TGT CTG ACC GTT CCA GCT TCC GCT TAT GAA GTG CGC AAC GTG	240
Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Glu Val Arg Asn Val	
65 70 75 80	
TCC GGG ATG TAC CAT GTC ACG AAC GAC TGC TCC AAC TCA AGC ATT GTG	288
Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val	
85 90 95	
TAT GAG GCA GCG GAC ATG ATC ATG CAC ACC CCC GGG TGC GTG CCC TGC	336
Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys	
100 105 110	
GTT CGG GAG AAC AAC TCT TCC CGC TGC TGG GTA GCG CTC ACC CCC ACG	384
Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr	
115 120 125	
CTC GCA GCT AGG AAC GCC AGC GTC CCC ACC ACG ACA ATA CGA CGC CAC	432
Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His	
130 135 140	



-continued

GTC GAT TTG CTC GTT GGG GCG GCT GCT TTC TGT TCC GCT ATG TAC GTG	480
Val Asp Leu Leu Val Gly Ala Ala Ala Phe Cys Ser Ala Met Tyr Val	
145 150 155 160	
GGG GAC CTC TGC GGA TCT GTC TTC CTC GTC TCC CAG CTG TTC ACC ATC	528
Gly Asp Leu Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Ile	
165 170 175	
TCG CCT CGC CGG CAT GAG ACG GTG CAG GAC TGC AAT TGC TCA ATC TAT	576
Ser Pro Arg Arg His Glu Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr	
180 185 190	
CCC GGC CAC ATA ACG GGT CAC CGT ATG GCT TGG GAT ATG ATG ATG AAC	624
Pro Gly His Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn	
195 200 205	
TGG TCG CCT ACA ACG GCC CTG GTG GTA TCG CAG CTG CTC CGG ATC CCA	672
Trp Ser Pro Thr Thr Ala Leu Val Val Ser Gln Leu Leu Arg Ile Pro	
210 215 220	
CAA GCT GTC GTG GAC ATG GTG GCG GGG GCC CAT TGG GGA GTC CTG GCG	720
Gln Ala Val Val Asp Met Val Ala Gly Ala His Trp Gly Val Leu Ala	
225 230 235 240	
GGT CTC GCC TAC TAT TCC ATG GTG GGG AAC TGG GCT AAG GTT TTG ATT	768
Gly Leu Ala Tyr Tyr Ser Met Val Gly Asn Trp Ala Lys Val Leu Ile	
245 250 255	
GTG ATG CTA CTC TTT GCT CCC TAATAG	795
Val Met Leu Leu Phe Ala Pro	
260	

## (2) INFORMATION FOR SEQ ID NO: 6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 263 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Met Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu	
1 5 10 15	
Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg	
20 25 30	
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala	
35 40 45	
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu	
50 55 60	
Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Glu Val Arg Asn Val	
65 70 75 80	
Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val	
85 90 95	
Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys	
100 105 110	
Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr	
115 120 125	
Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His	
130 135 140	
Val Asp Leu Leu Val Gly Ala Ala Ala Phe Cys Ser Ala Met Tyr Val	
145 150 155 160	
Gly Asp Leu Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Ile	
165 170 175	

-continued

Ser Pro Arg Arg His Glu Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr  
 180 185 190

Pro Gly His Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn  
 195 200 205

Trp Ser Pro Thr Thr Ala Leu Val Val Ser Gln Leu Leu Arg Ile Pro  
 210 215 220

Gln Ala Val Val Asp Met Val Ala Gly Ala His Trp Gly Val Leu Ala  
 225 230 235 240

Gly Leu Ala Tyr Tyr Ser Met Val Gly Asn Trp Ala Lys Val Leu Ile  
 245 250 255

Val Met Leu Leu Phe Ala Pro  
 260

## (2) INFORMATION FOR SEQ ID NO: 7:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 633 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

- (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 1..630

- (ix) FEATURE:  
 (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 1..627

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

ATG TTG GGT AAG GTC ATC GAT ACC CTT ACG TGC GGC TTC GCC GAC CTC	48
Met Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu	
1 5 10 15	
ATG GGG TAC ATT CCG CTC GTC GGC GCC CCC CTA GGG GGT GCT GCC AGA	96
Met Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg	
20 25 30	
GCC CTG GCG CAT GGC GTC CGG GTT CTG GAA GAC GGC GTG AAC TAT GCA	144
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala	
35 40 45	
ACA GGG AAT TTG CCT GGT TGC TCT TTC TCT ATC TTC CTC TTG GCT TTA	192
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu	
50 55 60	
CTG TCC TGT CTG ACC ATT CCA GCT TCC GCT TAT GAG GTG CGC AAC GTG	240
Leu Ser Cys Leu Thr Ile Pro Ala Ser Ala Tyr Glu Val Arg Asn Val	
65 70 75 80	
TCC GGG ATG TAC CAT GTC ACG AAC GAC TGC TCC AAC TCA AGC ATT GTG	288
Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val	
85 90 95	
TAT GAG GCA GCG GAC ATG ATC ATG CAC ACC CCC GGG TGC GTG CCC TGC	336
Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys	
100 105 110	
GTT CGG GAG AAC AAC TCT TCC CGC TGC TGG GTA GCG CTC ACC CCC ACG	384
Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr	
115 120 125	
CTC GCA GCT AGG AAC GCC AGC GTC CCC ACT ACG ACA ATA CGA CGC CAC	432
Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His	
130 135 140	

-continued

---

GTC GAT TTG CTC GTT GGG GCG GCT GCT TTC TGT TCC GCT ATG TAC GTG 480  
 Val Asp Leu Leu Val Gly Ala Ala Ala Phe Cys Ser Ala Met Tyr Val  
 145 150 155 160

GGG GAT CTC TGC GGA TCT GTC TTC CTC GTC TCC CAG CTG TTC ACC ATC 528  
 Gly Asp Leu Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Ile  
 165 170 175

TCG CCT CGC CGG CAT GAG ACG GTG CAG GAC TGC AAT TGC TCA ATC TAT 576  
 Ser Pro Arg Arg His Glu Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr  
 180 185 190

CCC GGC CAC ATA ACA GGT CAC CGT ATG GCT TGG GAT ATG ATG ATG AAC 624  
 Pro Gly His Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn  
 195 200 205

TGG TAATAG 633  
 Trp  
 210

## (2) INFORMATION FOR SEQ ID NO: 8:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 209 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Met Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu  
 1 5 10 15

Met Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg  
 20 25 30

Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala  
 35 40 45

Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu  
 50 55 60

Leu Ser Cys Leu Thr Ile Pro Ala Ser Ala Tyr Glu Val Arg Asn Val  
 65 70 75 80

Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val  
 85 90 95

Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys  
 100 105 110

Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr  
 115 120 125

Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His  
 130 135 140

Val Asp Leu Leu Val Gly Ala Ala Ala Phe Cys Ser Ala Met Tyr Val  
 145 150 155 160

Gly Asp Leu Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Ile  
 165 170 175

Ser Pro Arg Arg His Glu Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr  
 180 185 190

Pro Gly His Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn  
 195 200 205

Trp

## (2) INFORMATION FOR SEQ ID NO: 9:

## (i) SEQUENCE CHARACTERISTICS:

-continued

(A) LENGTH: 483 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: CDS  
 (B) LOCATION: 1..480

(ix) FEATURE:

(A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 1..477

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

ATG CCC GGT TGC TCT TTC TCT ATC TTC CTC TTG GCC CTG CTG TCC TGT	48
Met Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu Leu Ser Cys	
1 5 10 15	
CTG ACC ATA CCA GCT TCC GCT TAT GAA GTG CGC AAC GTG TCC GGG GTG	96
Leu Thr Ile Pro Ala Ser Ala Tyr Glu Val Arg Asn Val Ser Gly Val	
20 25 30	
TAC CAT GTC ACG AAC GAC TGC TCC AAC TCA AGC ATA GTG TAT GAG GCA	144
Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val Tyr Glu Ala	
35 40 45	
GCG GAC ATG ATC ATG CAC ACC CCC GGG TGC GTG CCC TGC GTT CGG GAG	192
Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys Val Arg Glu	
50 55 60	
GGC AAC TCC TCC CGT TGC TGG GTG GCG CTC ACT CCC ACG CTC GCG GCC	240
Gly Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr Leu Ala Ala	
65 70 75 80	
AGG AAC GCC AGC GTC CCC ACA ACG ACA ATA CGA CGC CAC GTC GAT TTG	288
Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His Val Asp Leu	
85 90 95	
CTC GTT GGG GCT GCT GCT TTC TGT TCC GCT ATG TAC GTG GGG GAT CTC	336
Leu Val Gly Ala Ala Ala Phe Cys Ser Ala Met Tyr Val Gly Asp Leu	
100 105 110	
TGC GGA TCT GTT TTC CTT GTT TCC CAG CTG TTC ACC TTC TCA CCT CGC	384
Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Phe Ser Pro Arg	
115 120 125	
CGG CAT CAA ACA GTA CAG GAC TGC AAC TGC TCA ATC TAT CCC GGC CAT	432
Arg His Gln Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His	
130 135 140	
GTA TCA GGT CAC CGC ATG GCT TGG GAT ATG ATG ATG AAC TGG TCC TAATAG	483
Val Ser Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp Ser	
145 150 155 160	

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 159 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Met Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu Leu Ser Cys	
1 5 10 15	
Leu Thr Ile Pro Ala Ser Ala Tyr Glu Val Arg Asn Val Ser Gly Val	
20 25 30	



-continued

---

TGC GGA TCT GTT TTC CTT GTT TCC CAG CTG TTC ACC TTC TCA CCT CGC	384
Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Phe Ser Pro Arg	
115 120 125	
CGG CAT CAA ACA GTA CAG GAC TGC AAC TGC TCA ATC TAT CCC GGC CAT	432
Arg His Gln Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His	
130 135 140	
GTA TCA GGT CAC CGC ATG GCT TGG GAT ATG ATG AAC TGG TAATAG	480
Val Ser Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp	
145 150 155	

## (2) INFORMATION FOR SEQ ID NO: 12:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 158 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

Met Ser Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu Leu Ser Cys	
1 5 10 15	
Leu Thr Ile Pro Ala Ser Ala Tyr Glu Val Arg Asn Val Ser Gly Val	
20 25 30	
Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val Tyr Glu Ala	
35 40 45	
Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys Val Arg Glu	
50 55 60	
Gly Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr Leu Ala Ala	
65 70 75 80	
Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His Val Asp Leu	
85 90 95	
Leu Val Gly Ala Ala Ala Phe Cys Ser Ala Met Tyr Val Gly Asp Leu	
100 105 110	
Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Phe Ser Pro Arg	
115 120 125	
Arg His Gln Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His	
130 135 140	
Val Ser Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp	
145 150 155	

## (2) INFORMATION FOR SEQ ID NO: 13:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 636 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (iii) HYPOTHETICAL: NO

## (iii) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..633

## (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide
- (B) LOCATION: 1..630

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

-continued

ATG CTG GGT AAG GCC ATC GAT ACC CTT ACG TGC GGC TTC GCC GAC CTC	48
Met Leu Gly Lys Ala Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu	
1 5 10 15	
GTG GGG TAC ATT CCG CTC GTC GGC GCC CCC CTA GGG GGC GCT GCC AGG	96
Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg	
20 25 30	
GCC CTG GCG CAT GGC GTC CGG GTT CTG GAA GAC GGC GTG AAC TAT GCA	144
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala	
35 40 45	
ACA GGG AAT TTG CCT GGT TGC TCT TTC TCT ATC TTC CTC TTG GCT TTA	192
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu	
50 55 60	
CTG TCC TGT CTA ACC ATT CCA GCT TCC GCT TAC GAG GTG CGC AAC GTG	240
Leu Ser Cys Leu Thr Ile Pro Ala Ser Ala Tyr Glu Val Arg Asn Val	
65 70 75 80	
TCC GGG ATG TAC CAT GTC ACG AAC GAC TGC TCC AAC TCA AGC ATT GTG	288
Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val	
85 90 95	
TAT GAG GCA GCG GAC ATG ATC ATG CAC ACC CCC GGG TGC GTG CCC TGC	336
Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys	
100 105 110	
GTT CGG GAG AAC AAC TCT TCC CGC TGC TGG GTA GCG CTC ACC CCC ACG	384
Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr	
115 120 125	
CTC GCG GCT AGG AAC GCC AGC ATC CCC ACT ACA ACA ATA CGA CGC CAC	432
Leu Ala Ala Arg Asn Ala Ser Ile Pro Thr Thr Thr Ile Arg Arg His	
130 135 140	
GTC GAT TTG CTC GTT GGG GCG GCT GCT TTC TGT TCC GCT ATG TAC GTG	480
Val Asp Leu Leu Val Gly Ala Ala Ala Phe Cys Ser Ala Met Tyr Val	
145 150 155 160	
GGG GAT CTC TGC GGA TCT GTC TTC CTC GTC TCC CAG CTG TTC ACC ATC	528
Gly Asp Leu Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Ile	
165 170 175	
TCG CCT CGC CGG CAT GAG ACG GTG CAG GAC TGC AAT TGC TCA ATC TAT	576
Ser Pro Arg Arg His Glu Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr	
180 185 190	
CCC GGC CAC ATA ACG GGT CAC CGT ATG GCT TGG GAT ATG ATG ATG AAC	624
Pro Gly His Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn	
195 200 205	
TGG TAC TAATAG	636
Trp Tyr	
210	

## (2) INFORMATION FOR SEQ ID NO: 14:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 210 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

Met Leu Gly Lys Ala Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu	
1 5 10 15	
Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg	
20 25 30	
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala	
35 40 45	
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu	

-continued

50	55	60
Leu Ser Cys Leu Thr Ile	Pro Ala Ser Ala Tyr Glu Val Arg Asn Val	
65	70	75 80
Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val		
	85	90 95
Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys		
	100	105 110
Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr		
	115	120 125
Leu Ala Ala Arg Asn Ala Ser Ile Pro Thr Thr Thr Ile Arg Arg His		
	130	135 140
Val Asp Leu Leu Val Gly Ala Ala Ala Phe Cys Ser Ala Met Tyr Val		
	145	150 155 160
Gly Asp Leu Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Ile		
	165	170 175
Ser Pro Arg Arg His Glu Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr		
	180	185 190
Pro Gly His Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn		
	195	200 205
Trp Tyr		
	210	

## (2) INFORMATION FOR SEQ ID NO: 15:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 26 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

ATGCCCGGTT GCTCTTTCTC TATCTT

26

## (2) INFORMATION FOR SEQ ID NO: 16:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 26 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

ATGTTGGGTA AGGCATCGA TACCCT

26

## (2) INFORMATION FOR SEQ ID NO: 17:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 30 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear



-continued

---

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

CTATTAGGAC CAGTTCATCA TCATATCCCA 30

(2) INFORMATION FOR SEQ ID NO: 18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

CTATTACCAG TTCATCATCA TATCCCA 27

(2) INFORMATION FOR SEQ ID NO: 19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

ATACGACGCC ACGTCGATTC CCAGCTGTTC ACCATC 36

(2) INFORMATION FOR SEQ ID NO: 20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

GATGGTGAAC AGCTGGGAAT CGACGTGGCG TCGTAT 36

(2) INFORMATION FOR SEQ ID NO: 21:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 723 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

-continued

(iii) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: CDS  
(B) LOCATION: 1..720

(ix) FEATURE:

(A) NAME/KEY: mat\_peptide  
(B) LOCATION: 1..717

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

ATG	TTG	GGT	AAG	GTC	ATC	GAT	ACC	CTT	ACA	TGC	GGC	TTC	GCC	GAC	CTC	48
Met	Leu	Gly	Lys	Val	Ile	Asp	Thr	Leu	Thr	Cys	Gly	Phe	Ala	Asp	Leu	
1				5					10					15		
GTG	GGG	TAC	ATT	CCG	CTC	GTC	GGC	GCC	CCC	CTA	GGG	GGC	GCT	GCC	AGG	96
Val	Gly	Tyr	Ile	Pro	Leu	Val	Gly	Ala	Pro	Leu	Gly	Gly	Ala	Ala	Arg	
			20					25					30			
GCC	CTG	GCG	CAT	GGC	GTC	CGG	GTT	CTG	GAG	GAC	GGC	GTG	AAC	TAT	GCA	144
Ala	Leu	Ala	His	Gly	Val	Arg	Val	Leu	Glu	Asp	Gly	Val	Asn	Tyr	Ala	
			35				40					45				
ACA	GGG	AAT	TTG	CCC	GGT	TGC	TCT	TTC	TCT	ATC	TTC	CTC	TTG	GCT	TTG	192
Thr	Gly	Asn	Leu	Pro	Gly	Cys	Ser	Phe	Ser	Ile	Phe	Leu	Leu	Ala	Leu	
	50					55					60					
CTG	TCC	TGT	CTG	ACC	GTT	CCA	GCT	TCC	GCT	TAT	GAA	GTG	CGC	AAC	GTG	240
Leu	Ser	Cys	Leu	Thr	Val	Pro	Ala	Ser	Ala	Tyr	Glu	Val	Arg	Asn	Val	
	65				70					75					80	
TCC	GGG	ATG	TAC	CAT	GTC	ACG	AAC	GAC	TGC	TCC	AAC	TCA	AGC	ATT	GTG	288
Ser	Gly	Met	Tyr	His	Val	Thr	Asn	Asp	Cys	Ser	Asn	Ser	Ser	Ile	Val	
				85					90					95		
TAT	GAG	GCA	GCG	GAC	ATG	ATC	ATG	CAC	ACC	CCC	GGG	TGC	GTG	CCC	TGC	336
Tyr	Glu	Ala	Ala	Asp	Met	Ile	Met	His	Thr	Pro	Gly	Cys	Val	Pro	Cys	
			100					105					110			
GTT	CGG	GAG	AAC	AAC	TCT	TCC	CGC	TGC	TGG	GTA	GCG	CTC	ACC	CCC	ACG	384
Val	Arg	Glu	Asn	Asn	Ser	Ser	Arg	Cys	Trp	Val	Ala	Leu	Thr	Pro	Thr	
	115						120					125				
CTC	GCA	GCT	AGG	AAC	GCC	AGC	GTC	CCC	ACC	ACG	ACA	ATA	CGA	CGC	CAC	432
Leu	Ala	Ala	Arg	Asn	Ala	Ser	Val	Pro	Thr	Thr	Thr	Ile	Arg	Arg	His	
	130					135						140				
GTC	GAT	TCC	CAG	CTG	TTC	ACC	ATC	TCG	CCT	CGC	CGG	CAT	GAG	ACG	GTG	480
Val	Asp	Ser	Gln	Leu	Phe	Thr	Ile	Ser	Pro	Arg	Arg	His	Glu	Thr	Val	
	145				150					155					160	
CAG	GAC	TGC	AAT	TGC	TCA	ATC	TAT	CCC	GGC	CAC	ATA	ACG	GGT	CAC	CGT	528
Gln	Asp	Cys	Asn	Cys	Ser	Ile	Tyr	Pro	Gly	His	Ile	Thr	Gly	His	Arg	
			165						170					175		
ATG	GCT	TGG	GAT	ATG	ATG	ATG	AAC	TGG	TCG	CCT	ACA	ACG	GCC	CTG	GTG	576
Met	Ala	Trp	Asp	Met	Met	Met	Asn	Trp	Ser	Pro	Thr	Thr	Ala	Leu	Val	
			180					185						190		
GTA	TCG	CAG	CTG	CTC	CGG	ATC	CCA	CAA	GCT	GTC	GTG	GAC	ATG	GTG	GCG	624
Val	Ser	Gln	Leu	Leu	Arg	Ile	Pro	Gln	Ala	Val	Val	Asp	Met	Val	Ala	
	195						200					205				
GGG	GCC	CAT	TGG	GGA	GTC	CTG	GCG	GGT	CTC	GCC	TAC	TAT	TCC	ATG	GTG	672
Gly	Ala	His	Trp	Gly	Val	Leu	Ala	Gly	Leu	Ala	Tyr	Tyr	Ser	Met	Val	
	210					215					220					
GGG	AAC	TGG	GCT	AAG	GTT	TTG	ATT	GTG	ATG	CTA	CTC	TTT	GCT	CCC	TAATAG	723
Gly	Asn	Trp	Ala	Lys	Val	Leu	Ile	Val	Met	Leu	Leu	Phe	Ala	Pro		
	225				230					235					240	

(2) INFORMATION FOR SEQ ID NO: 22:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 239 amino acids  
(B) TYPE: amino acid

-continued

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

Met Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu  
 1 5 10 15  
 Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg  
 20 25 30  
 Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala  
 35 40 45  
 Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu  
 50 55 60  
 Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Glu Val Arg Asn Val  
 65 70 75 80  
 Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val  
 85 90 95  
 Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys  
 100 105 110  
 Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr  
 115 120 125  
 Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His  
 130 135 140  
 Val Asp Ser Gln Leu Phe Thr Ile Ser Pro Arg Arg His Glu Thr Val  
 145 150 155 160  
 Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His Ile Thr Gly His Arg  
 165 170 175  
 Met Ala Trp Asp Met Met Met Asn Trp Ser Pro Thr Thr Ala Leu Val  
 180 185 190  
 Val Ser Gln Leu Leu Arg Ile Pro Gln Ala Val Val Asp Met Val Ala  
 195 200 205  
 Gly Ala His Trp Gly Val Leu Ala Gly Leu Ala Tyr Tyr Ser Met Val  
 210 215 220  
 Gly Asn Trp Ala Lys Val Leu Ile Val Met Leu Leu Phe Ala Pro  
 225 230 235

(2) INFORMATION FOR SEQ ID NO: 23:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 561 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 1..558

(ix) FEATURE:  
 (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 1..555

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

ATG TTG GGT AAG GTC ATC GAT ACC CTT ACA TGC GGC TTC GCC GAC CTC  
 Met Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu  
 1 5 10 15

48



-continued

115		120		125											
Leu	Ala	Ala	Arg	Asn	Ala	Ser	Val	Pro	Thr	Thr	Thr	Ile	Arg	Arg	His
130							135					140			
Val	Asp	Ser	Gln	Leu	Phe	Thr	Ile	Ser	Pro	Arg	Arg	His	Glu	Thr	Val
145					150					155					160
Gln	Asp	Cys	Asn	Cys	Ser	Ile	Tyr	Pro	Gly	His	Ile	Thr	Gly	His	Arg
			165						170					175	
Met	Ala	Trp	Asp	Met	Met	Met	Asn	Trp							
			180					185							

## (2) INFORMATION FOR SEQ ID NO: 25:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 606 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

(A) NAME/KEY: CDS  
 (B) LOCATION: 1..603

## (ix) FEATURE:

(A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 1..600

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

ATG	TTG	GGT	AAG	GTC	ATC	GAT	ACC	CTT	ACA	TGC	GGC	TTC	GCC	GAC	CTC	48
Met	Leu	Gly	Lys	Val	Ile	Asp	Thr	Leu	Thr	Cys	Gly	Phe	Ala	Asp	Leu	
1				5					10					15		
GTG	GGG	TAC	ATT	CCG	CTC	GTC	GGC	GCC	CCC	CTA	GGG	GGC	GCT	GCC	AGG	96
Val	Gly	Tyr	Ile	Pro	Leu	Val	Gly	Ala	Pro	Leu	Gly	Gly	Ala	Ala	Arg	
			20					25					30			
GCC	CTG	GCG	CAT	GGC	GTC	CGG	GTT	CTG	GAG	GAC	GGC	GTG	AAC	TAT	GCA	144
Ala	Leu	Ala	His	Gly	Val	Arg	Val	Leu	Glu	Asp	Gly	Val	Asn	Tyr	Ala	
			35				40						45			
ACA	GGG	AAT	TTG	CCC	GGT	TGC	TCT	TTC	TCT	ATC	TTC	CTC	TTG	GCT	TTG	192
Thr	Gly	Asn	Leu	Pro	Gly	Cys	Ser	Phe	Ser	Ile	Phe	Leu	Leu	Ala	Leu	
		50				55					60					
CTG	TCC	TGT	CTG	ACC	GTT	CCA	GCT	TCC	GCT	TAT	GAA	GTG	CGC	AAC	GTG	240
Leu	Ser	Cys	Leu	Thr	Val	Pro	Ala	Ser	Ala	Tyr	Glu	Val	Arg	Asn	Val	
		65			70					75					80	
TCC	GGG	ATG	TAC	CAT	GTC	ACG	AAC	GAC	TGC	TCC	AAC	TCA	AGC	ATT	GTG	288
Ser	Gly	Met	Tyr	His	Val	Thr	Asn	Asp	Cys	Ser	Asn	Ser	Ser	Ile	Val	
				85					90					95		
TAT	GAG	GCA	GCG	GAC	ATG	ATC	ATG	CAC	ACC	CCC	GGG	TGC	GTG	CCC	TGC	336
Tyr	Glu	Ala	Ala	Asp	Met	Ile	Met	His	Thr	Pro	Gly	Cys	Val	Pro	Cys	
			100					105					110			
GTT	CGG	GAG	AAC	AAC	TCT	TCC	CGC	TGC	TGG	GTA	GCG	CTC	ACC	CCC	ACG	384
Val	Arg	Glu	Asn	Asn	Ser	Ser	Arg	Cys	Trp	Val	Ala	Leu	Thr	Pro	Thr	
		115					120					125				
CTC	GCA	GCT	AGG	AAC	GCC	AGC	GTC	CCC	ACC	ACG	ACA	ATA	CGA	CGC	CAC	432
Leu	Ala	Ala	Arg	Asn	Ala	Ser	Val	Pro	Thr	Thr	Thr	Ile	Arg	Arg	His	
					130										140	
GTC	GAT	TCC	CAG	CTG	TTC	ACC	ATC	TCG	CCT	CGC	CGG	CAT	GAG	ACG	GTG	480
Val	Asp	Ser	Gln	Leu	Phe	Thr	Ile	Ser	Pro	Arg	Arg	His	Glu	Thr	Val	
145					150					155					160	

-continued

---

CAG GAC TGC AAT TGC TCA ATC TAT CCC GGC CAC ATA ACG GGT CAC CGT	528
Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His Ile Thr Gly His Arg	
165 170 175	
ATG GCT TGG GAT ATG ATG ATG AAC TGG TCG CCT ACA ACG GCC CTG GTG	576
Met Ala Trp Asp Met Met Met Asn Trp Ser Pro Thr Thr Ala Leu Val	
180 185 190	
GTA TCG CAG CTG CTC CGG ATC CTC TAATAG	606
Val Ser Gln Leu Leu Arg Ile Leu	
195 200	

## (2) INFORMATION FOR SEQ ID NO: 26:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 200 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

Met	Leu	Gly	Lys	Val	Ile	Asp	Thr	Leu	Thr	Cys	Gly	Phe	Ala	Asp	Leu
1				5					10					15	
Val	Gly	Tyr	Ile	Pro	Leu	Val	Gly	Ala	Pro	Leu	Gly	Gly	Ala	Ala	Arg
			20					25					30		
Ala	Leu	Ala	His	Gly	Val	Arg	Val	Leu	Glu	Asp	Gly	Val	Asn	Tyr	Ala
		35				40						45			
Thr	Gly	Asn	Leu	Pro	Gly	Cys	Ser	Phe	Ser	Ile	Phe	Leu	Leu	Ala	Leu
		50				55					60				
Leu	Ser	Cys	Leu	Thr	Val	Pro	Ala	Ser	Ala	Tyr	Glu	Val	Arg	Asn	Val
65					70					75				80	
Ser	Gly	Met	Tyr	His	Val	Thr	Asn	Asp	Cys	Ser	Asn	Ser	Ser	Ile	Val
				85					90					95	
Tyr	Glu	Ala	Ala	Asp	Met	Ile	Met	His	Thr	Pro	Gly	Cys	Val	Pro	Cys
		100						105					110		
Val	Arg	Glu	Asn	Asn	Ser	Ser	Arg	Cys	Trp	Val	Ala	Leu	Thr	Pro	Thr
		115					120					125			
Leu	Ala	Ala	Arg	Asn	Ala	Ser	Val	Pro	Thr	Thr	Thr	Ile	Arg	Arg	His
	130					135						140			
Val	Asp	Ser	Gln	Leu	Phe	Thr	Ile	Ser	Pro	Arg	Arg	His	Glu	Thr	Val
145					150					155				160	
Gln	Asp	Cys	Asn	Cys	Ser	Ile	Tyr	Pro	Gly	His	Ile	Thr	Gly	His	Arg
			165						170					175	
Met	Ala	Trp	Asp	Met	Met	Met	Asn	Trp	Ser	Pro	Thr	Thr	Ala	Leu	Val
			180					185					190		
Val	Ser	Gln	Leu	Leu	Arg	Ile	Leu								
		195					200								

## (2) INFORMATION FOR SEQ ID NO: 27:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 636 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (iii) HYPOTHETICAL: NO

## (iii) ANTI-SENSE: NO

## (ix) FEATURE:

-continued

(A) NAME/KEY: CDS  
(B) LOCATION: 1..633

## (ix) FEATURE:

(A) NAME/KEY: mat\_peptide  
(B) LOCATION: 1..630

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

```

ATG TTG GGT AAG GTC ATC GAT ACC CTT ACA TGC GGC TTC GCC GAC CTC      48
Met Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu
  1                               5                               10          15

GTG GGG TAC ATT CCG CTC GTC GGC GCC CCC CTA GGG GGC GCT GCC AGG      96
Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg
                20                               25          30

GCC CTG GCG CAT GGC GTC CGG GTT CTG GAG GAC GGC GTG AAC TAT GCA     144
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala
                35                               40          45

ACA GGG AAT TTG CCC GGT TGC TCT TTC TCT ATC TTC CTC TTG GCT TTG     192
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu
  50                               55                               60

CTG TCC TGT CTG ACC GTT CCA GCT TCC GCT TAT GAA GTG CGC AAC GTG     240
Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Glu Val Arg Asn Val
  65                               70                               75          80

TCC GGG ATG TAC CAT GTC ACG AAC GAC TGC TCC AAC TCA AGC ATT GTG     288
Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val
                85                               90          95

TAT GAG GCA GCG GAC ATG ATC ATG CAC ACC CCC GGG TGC GTG CCC TGC     336
Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys
                100                              105          110

GTT CGG GAG AAC AAC TCT TCC CGC TGC TGG GTA GCG CTC ACC CCC ACG     384
Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr
                115                              120          125

CTC GCA GCT AGG AAC GCC AGC GTC CCC ACC ACG ACA ATA CGA CGC CAC     432
Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His
                130                              135          140

GTC GAT TCC CAG CTG TTC ACC ATC TCG CCT CGC CGG CAT GAG ACG GTG     480
Val Asp Ser Gln Leu Phe Thr Ile Ser Pro Arg Arg His Glu Thr Val
                145                              150          155          160

CAG GAC TGC AAT TGC TCA ATC TAT CCC GGC CAC ATA ACG GGT CAC CGT     528
Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His Ile Thr Gly His Arg
                165                              170          175

ATG GCT TGG GAT ATG ATG ATG AAC TGG TCG CCT ACA ACG GCC CTG GTG     576
Met Ala Trp Asp Met Met Met Asn Trp Ser Pro Thr Thr Ala Leu Val
                180                              185          190

GTA TCG CAG CTG CTC CGG ATC GTG ATC GAG GGC AGA CAC CAT CAC CAC     624
Val Ser Gln Leu Leu Arg Ile Val Ile Glu Gly Arg His His His His
                195                              200          205

CAT CAC TAATAG                                                         636
His His
  210

```

## (2) INFORMATION FOR SEQ ID NO: 28:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 210 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

```

Met Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu
  1                               5                               10          15

```

-continued

Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg  
 20 25 30  
 Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala  
 35 40 45  
 Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu  
 50 55 60  
 Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Glu Val Arg Asn Val  
 65 70 75 80  
 Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val  
 85 90 95  
 Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys  
 100 105 110  
 Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr  
 115 120 125  
 Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His  
 130 135 140  
 Val Asp Ser Gln Leu Phe Thr Ile Ser Pro Arg Arg His Glu Thr Val  
 145 150 155 160  
 Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His Ile Thr Gly His Arg  
 165 170 175  
 Met Ala Trp Asp Met Met Met Asn Trp Ser Pro Thr Thr Ala Leu Val  
 180 185 190  
 Val Ser Gln Leu Leu Arg Ile Val Ile Glu Gly Arg His His His His  
 195 200 205  
 His His  
 210

## (2) INFORMATION FOR SEQ ID NO: 29:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 630 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS  
 (B) LOCATION: 1..627

## (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 1..624

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

ATG GGT AAG GTC ATC GAT ACC CTT ACG TGC GGA TTC GCC GAT CTC ATG 48  
 Met Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu Met  
 1 5 10 15  
 GGG TAC ATC CCG CTC GTC GGC GCT CCC GTA GGA GGC GTC GCA AGA GCC 96  
 Gly Tyr Ile Pro Leu Val Gly Ala Pro Val Gly Gly Val Ala Arg Ala  
 20 25 30  
 CTT GCG CAT GGC GTG AGG GCC CTT GAA GAC GGG ATA AAT TTC GCA ACA 144  
 Leu Ala His Gly Val Arg Ala Leu Glu Asp Gly Ile Asn Phe Ala Thr  
 35 40 45  
 GGG AAT TTG CCC GGT TGC TCC TTT TCT ATT TTC CTT CTC GCT CTG TTC 192  
 Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu Phe





-continued

Asp	Leu	Leu	Val	Gly	Ala	Ala	Thr	Met	Cys	Ser	Ala	Leu	Tyr	Val	Gly	
145					150					155					160	
Asp	Met	Cys	Gly	Ala	Val	Phe	Leu	Val	Gly	Gln	Ala	Phe	Thr	Phe	Arg	
				165					170					175		
Pro	Arg	Arg	His	Gln	Thr	Val	Gln	Thr	Cys	Asn	Cys	Ser	Leu	Tyr	Pro	
			180					185					190			
Gly	His	Leu	Ser	Gly	His	Arg	Met	Ala	Trp	Asp	Met	Met	Met	Asn	Trp	
	195						200					205				

## (2) INFORMATION FOR SEQ ID NO: 31:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 630 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 1..627

(ix) FEATURE:  
 (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 1..624

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

ATG	GGT	AAG	GTC	ATC	GAT	ACC	CTA	ACG	TGC	GGA	TTC	GCC	GAT	CTC	ATG	48
Met	Gly	Lys	Val	Ile	Asp	Thr	Leu	Thr	Cys	Gly	Phe	Ala	Asp	Leu	Met	
1				5					10					15		
GGG	TAT	ATC	CCG	CTC	GTA	GGC	GGC	CCC	ATT	GGG	GGC	GTC	GCA	AGG	GCT	96
Gly	Tyr	Ile	Pro	Leu	Val	Gly	Gly	Pro	Ile	Gly	Gly	Val	Ala	Arg	Ala	
			20					25					30			
CTC	GCA	CAC	GGT	GTG	AGG	GTC	CTT	GAG	GAC	GGG	GTA	AAC	TAT	GCA	ACA	144
Leu	Ala	His	Gly	Val	Arg	Val	Leu	Glu	Asp	Gly	Val	Asn	Tyr	Ala	Thr	
			35				40						45			
GGG	AAT	TTA	CCC	GGT	TGC	TCT	TTC	TCT	ATC	TTT	ATT	CTT	GCT	CTT	CTC	192
Gly	Asn	Leu	Pro	Gly	Cys	Ser	Phe	Ser	Ile	Phe	Ile	Leu	Ala	Leu	Leu	
	50					55					60					
TCG	TGT	CTG	ACC	GTT	CCG	GCC	TCT	GCA	GTT	CCC	TAC	CGA	AAT	GCC	TCT	240
Ser	Cys	Leu	Thr	Val	Pro	Ala	Ser	Ala	Val	Pro	Tyr	Arg	Asn	Ala	Ser	
	65				70				75					80		
GGG	ATT	TAT	CAT	GTT	ACC	AAT	GAT	TGC	CCA	AAC	TCT	TCC	ATA	GTC	TAT	288
Gly	Ile	Tyr	His	Val	Thr	Asn	Asp	Cys	Pro	Asn	Ser	Ser	Ile	Val	Tyr	
				85				90						95		
GAG	GCA	GAT	AAC	CTG	ATC	CTA	CAC	GCA	CCT	GGT	TGC	GTG	CCT	TGT	GTC	336
Glu	Ala	Asp	Asn	Leu	Ile	Leu	His	Ala	Pro	Gly	Cys	Val	Pro	Cys	Val	
			100					105						110		
ATG	ACA	GGT	AAT	GTG	AGT	AGA	TGC	TGG	GTC	CAA	ATT	ACC	CCT	ACA	CTG	384
Met	Thr	Gly	Asn	Val	Ser	Arg	Cys	Trp	Val	Gln	Ile	Thr	Pro	Thr	Leu	
		115					120						125			
TCA	GCC	CCG	AGC	CTC	GGA	GCA	GTC	ACG	GCT	CCT	CTT	CGG	AGA	GCC	GTT	432
Ser	Ala	Pro	Ser	Leu	Gly	Ala	Val	Thr	Ala	Pro	Leu	Arg	Arg	Ala	Val	
		130					135					140				
GAC	TAC	CTA	GCG	GGA	GGG	GCT	GCC	CTC	TGC	TCC	GCG	TTA	TAC	GTA	GGA	480
Asp	Tyr	Leu	Ala	Gly	Gly	Ala	Ala	Leu	Cys	Ser	Ala	Leu	Tyr	Val	Gly	
	145				150					155					160	
GAC	GCG	TGT	GGG	GCA	CTA	TTC	TTG	GTA	GGC	CAA	ATG	TTC	ACC	TAT	AGG	528
Asp	Ala	Cys	Gly	Ala	Leu	Phe	Leu	Val	Gly	Gln	Met	Phe	Thr	Tyr	Arg	

-continued

	165	170	175	
CCT CGC CAG CAC GCT ACG GTG CAG AAC TGC AAC TGT TCC ATT TAC AGT				576
Pro Arg Gln His Ala Thr Val Gln Asn Cys Asn Cys Ser Ile Tyr Ser	180	185	190	
GGC CAT GTT ACC GGC CAC CGG ATG GCA TGG GAT ATG ATG AAC TGG				624
Gly His Val Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp	195	200	205	
TAATAG				630

## (2) INFORMATION FOR SEQ ID NO: 32:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 208 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

Met Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu Met  
 1 5 10 15

Gly Tyr Ile Pro Leu Val Gly Gly Pro Ile Gly Gly Val Ala Arg Ala  
 20 25 30

Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala Thr  
 35 40 45

Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Ile Leu Ala Leu Leu  
 50 55 60

Ser Cys Leu Thr Val Pro Ala Ser Ala Val Pro Tyr Arg Asn Ala Ser  
 65 70 75 80

Gly Ile Tyr His Val Thr Asn Asp Cys Pro Asn Ser Ser Ile Val Tyr  
 85 90 95

Glu Ala Asp Asn Leu Ile Leu His Ala Pro Gly Cys Val Pro Cys Val  
 100 105 110

Met Thr Gly Asn Val Ser Arg Cys Trp Val Gln Ile Thr Pro Thr Leu  
 115 120 125

Ser Ala Pro Ser Leu Gly Ala Val Thr Ala Pro Leu Arg Arg Ala Val  
 130 135 140

Asp Tyr Leu Ala Gly Gly Ala Ala Leu Cys Ser Ala Leu Tyr Val Gly  
 145 150 155 160

Asp Ala Cys Gly Ala Leu Phe Leu Val Gly Gln Met Phe Thr Tyr Arg  
 165 170 175

Pro Arg Gln His Ala Thr Val Gln Asn Cys Asn Cys Ser Ile Tyr Ser  
 180 185 190

Gly His Val Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp  
 195 200 205

## (2) INFORMATION FOR SEQ ID NO: 33:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (iii) HYPOTHETICAL: NO

## (iii) ANTI-SENSE: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:

-continued

---

TGGGATATGA TGATGAACTG GTC 23

(2) INFORMATION FOR SEQ ID NO: 34:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 30 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 34:

CTATTATGGT GGTAAGCCAC AGAGCAGGAG 30

(2) INFORMATION FOR SEQ ID NO: 35:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1476 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 1..1473

(ix) FEATURE:  
 (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 1..1470

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 35:

TGG GAT ATG ATG ATG AAC TGG TCG CCT ACA ACG GCC CTG GTG GTA TCG 48  
 Trp Asp Met Met Met Asn Trp Ser Pro Thr Thr Ala Leu Val Val Ser  
 1 5 10 15

CAG CTG CTC CGG ATC CCA CAA GCT GTC GTG GAC ATG GTG GCG GGG GCC 96  
 Gln Leu Leu Arg Ile Pro Gln Ala Val Val Asp Met Val Ala Gly Ala  
 20 25 30

CAT TGG GGA GTC CTG GCG GGC CTC GCC TAC TAT TCC ATG GTG GGG AAC 144  
 His Trp Gly Val Leu Ala Gly Leu Ala Tyr Tyr Ser Met Val Gly Asn  
 35 40 45

TGG GCT AAG GTT TTG GTT GTG ATG CTA CTC TTT GCC GGC GTC GAC GGG 192  
 Trp Ala Lys Val Leu Val Val Met Leu Leu Phe Ala Gly Val Asp Gly  
 50 55 60

CAT ACC CGC GTG TCA GGA GGG GCA GCA GCC TCC GAT ACC AGG GGC CTT 240  
 His Thr Arg Val Ser Gly Gly Ala Ala Ala Ser Asp Thr Arg Gly Leu  
 65 70 75 80

GTG TCC CTC TTT AGC CCC GGG TCG GCT CAG AAA ATC CAG CTC GTA AAC 288  
 Val Ser Leu Phe Ser Pro Gly Ser Ala Gln Lys Ile Gln Leu Val Asn  
 85 90 95

ACC AAC GGC AGT TGG CAC ATC AAC AGG ACT GCC CTG AAC TGC AAC GAC 336  
 Thr Asn Gly Ser Trp His Ile Asn Arg Thr Ala Leu Asn Cys Asn Asp  
 100 105 110

TCC CTC CAA ACA GGG TTC TTT GCC GCA CTA TTC TAC AAA CAC AAA TTC 384  
 Ser Leu Gln Thr Gly Phe Phe Ala Ala Leu Phe Tyr Lys His Lys Phe  
 115 120 125

-continued

AAC Asn 130	TCG Ser 130	TCT Ser 130	GGA Gly 130	TGC Cys 130	CCA Pro 130	GAG Glu 135	CGC Arg 135	TTG Leu 135	GCC Ala 140	AGC Ser 140	TGT Cys 140	CGC Arg 140	TCC Ser 140	ATC Ile 140	GAC Asp 140	432
AAG Lys 145	TTC Phe 145	GCT Ala 145	CAG Gln 145	GGG Gly 150	TGG Trp 150	GGT Gly 150	CCC Pro 150	CTC Leu 150	ACT Thr 155	TAC Tyr 155	ACT Thr 155	GAG Glu 155	CCT Pro 155	AAC Asn 160	AGC Ser 160	480
TCG Ser 165	GAC Asp 165	CAG Gln 165	AGG Arg 165	CCC Pro 165	TAC Tyr 165	TGC Cys 165	TGG Trp 170	CAC His 170	TAC Tyr 170	GCG Ala 170	CCT Pro 170	CGA Arg 175	CCG Pro 175	TGT Cys 175	GGT Gly 175	528
ATT Ile 180	GTA Val 180	CCC Pro 180	GCG Ala 180	TCT Ser 180	CAG Gln 185	GTG Val 185	TGC Cys 185	GGT Gly 185	CCA Pro 185	GTG Val 190	TAT Tyr 190	TGC Cys 190	TTC Phe 190	ACC Thr 190	CCG Pro 190	576
AGC Ser 195	CCT Pro 195	GTT Val 195	GTG Val 195	GTG Val 200	GGG Gly 200	ACG Thr 200	ACC Thr 200	GAT Asp 200	CGG Arg 205	TTT Phe 205	GGT Gly 205	GTC Val 205	CCC Pro 205	ACG Thr 205	TAT Tyr 205	624
AAC Asn 210	TGG Trp 210	GGG Gly 210	GCG Ala 210	AAC Asn 215	GAC Asp 215	TCG Ser 215	GAT Asp 215	GTG Val 220	CTG Leu 220	ATT Ile 220	CTC Leu 220	AAC Asn 220	AAC Asn 220	ACG Thr 220	CGG Arg 220	672
CCG Pro 225	CCG Pro 225	CGA Arg 230	GGC Gly 230	AAC Asn 230	TGG Trp 230	TTC Phe 230	GGC Gly 235	TGT Cys 235	ACA Thr 235	TGG Trp 235	ATG Met 235	AAT Asn 235	GGC Gly 240	ACT Thr 240	GGG Gly 240	720
TTC Phe 245	ACC Thr 245	AAG Lys 245	ACG Thr 245	TGT Cys 245	GGG Gly 245	GGC Gly 245	CCC Pro 250	CCG Pro 250	TGC Cys 250	AAC Asn 250	ATC Ile 250	GGG Gly 255	GGG Gly 255	GCC Ala 255	GGC Gly 255	768
AAC Asn 260	AAC Asn 260	ACC Thr 260	TTG Leu 260	ACC Thr 260	TGC Cys 265	CCC Pro 265	ACT Thr 265	GAC Asp 265	TGT Cys 265	TTT Phe 270	CGG Arg 270	AAG Lys 270	CAC His 270	CCC Pro 270	GAG Glu 270	816
GCC Ala 275	ACC Thr 275	TAC Tyr 275	GCC Ala 275	AGA Arg 275	TGC Cys 280	GGT Gly 280	TCT Ser 280	GGG Gly 280	CCC Pro 285	TGG Trp 285	CTG Leu 285	ACA Thr 285	CCT Pro 285	AGG Arg 285	TGT Cys 285	864
ATG Met 290	GTT Val 290	CAT His 290	TAC Tyr 290	CCA Pro 295	TAT Tyr 295	AGG Arg 295	CTC Leu 295	TGG Trp 295	CAC His 300	TAC Tyr 300	CCC Pro 300	TGC Cys 300	ACT Thr 300	GTC Val 300	AAC Asn 300	912
TTC Phe 305	ACC Thr 305	ATC Ile 305	TTC Phe 310	AAG Lys 310	GTT Val 310	AGG Arg 310	ATG Met 315	TAC Tyr 315	GTG Val 315	GGG Gly 315	GGC Gly 315	GTG Val 320	GAG Glu 320	CAC His 320	AGG Arg 320	960
TTC Phe 325	GAA Glu 325	GCC Ala 325	GCA Ala 325	TGC Cys 325	AAT Asn 330	TGG Trp 330	ACT Thr 330	CGA Arg 330	GGA Glu 330	GAG Glu 330	CGT Arg 330	TGT Cys 335	GAC Asp 335	TTG Leu 335	GAG Glu 335	1008
GAC Asp 340	AGG Arg 340	GAT Asp 340	AGA Arg 340	TCA Ser 340	GAG Glu 345	CTT Leu 345	AGC Ser 345	CCG Pro 345	CTG Leu 345	CTG Leu 345	CTG Leu 345	TCT Ser 350	ACA Thr 350	ACA Thr 350	GAG Glu 350	1056
TGG Trp 355	CAG Gln 355	ATA Ile 355	CTG Leu 355	CCC Pro 360	TGT Cys 360	TCC Ser 360	TTC Phe 360	ACC Thr 365	ACC Thr 365	CTG Leu 365	CCG Pro 365	GCC Ala 365	CTA Leu 365	TCC Ser 365	ACC Thr 365	1104
GGC Gly 370	CTG Leu 370	ATC Ile 370	CAC His 370	CTC Leu 375	CAT His 375	CAG Gln 375	AAC Asn 375	ATC Ile 380	GTG Val 380	GAC Asp 380	GTG Val 380	CAA Gln 380	TAC Tyr 380	CTG Leu 380	TAC Tyr 380	1152
GGT Gly 385	GTA Val 385	GGG Gly 385	TCG Ser 385	GCG Ala 390	GTT Val 390	GTC Val 390	TCC Ser 390	CTT Leu 395	GTC Val 395	ATC Ile 395	AAA Lys 395	TGG Trp 395	GAG Glu 395	TAT Tyr 395	GTC Val 395	1200
CTG Leu 405	TTG Leu 405	CTC Leu 405	TTC Phe 405	CTT Leu 405	CTC Leu 410	CTG Leu 410	GCA Ala 410	GAC Asp 410	GCG Ala 410	CGC Arg 415	ATC Ile 415	TGC Cys 415	GCC Ala 415	TGC Cys 415	TTA Leu 415	1248
TGG Trp 420	ATG Met 420	ATG Met 420	CTG Leu 420	CTG Leu 425	ATA Ile 425	GCT Ala 425	CAA Gln 425	GCT Ala 425	GAG Glu 425	GCC Ala 430	GCC Ala 430	TTA Leu 430	GAG Glu 430	AAC Asn 430	CTG Leu 430	1296
GTG Val 435	GTC Val 435	CTC Leu 435	AAT Asn 435	GCG Ala 440	GCG Ala 440	GCC Ala 440	GTG Val 440	GCC Ala 440	GGG Gly 445	GCG Ala 445	CAT His 445	GGC Gly 445	ACT Thr 445	CTT Leu 445	TCC Ser 445	1344

-continued

---

TTC CTT GTG TTC TTC TGT GCT GCC TGG TAC ATC AAG GGC AGG CTG GTC 1392  
Phe Leu Val Phe Phe Cys Ala Ala Trp Tyr Ile Lys Gly Arg Leu Val  
450 455 460

CCT GGT GCG GCA TAC GCC TTC TAT GGC GTG TGG CCG CTG CTC CTG CTT 1440  
Pro Gly Ala Ala Tyr Ala Phe Tyr Gly Val Trp Pro Leu Leu Leu Leu  
465 470 475 480

CTG CTG GCC TTA CCA CCA CGA GCT TAT GCC TAGTAA 1476  
Leu Leu Ala Leu Pro Pro Arg Ala Tyr Ala  
485 490

## (2) INFORMATION FOR SEQ ID NO: 36:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 490 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 36:

Trp Asp Met Met Met Asn Trp Ser Pro Thr Thr Ala Leu Val Val Ser  
1 5 10 15

Gln Leu Leu Arg Ile Pro Gln Ala Val Val Asp Met Val Ala Gly Ala  
20 25 30

His Trp Gly Val Leu Ala Gly Leu Ala Tyr Tyr Ser Met Val Gly Asn  
35 40 45

Trp Ala Lys Val Leu Val Val Met Leu Leu Phe Ala Gly Val Asp Gly  
50 55 60

His Thr Arg Val Ser Gly Gly Ala Ala Ala Ser Asp Thr Arg Gly Leu  
65 70 75 80

Val Ser Leu Phe Ser Pro Gly Ser Ala Gln Lys Ile Gln Leu Val Asn  
85 90 95

Thr Asn Gly Ser Trp His Ile Asn Arg Thr Ala Leu Asn Cys Asn Asp  
100 105 110

Ser Leu Gln Thr Gly Phe Phe Ala Ala Leu Phe Tyr Lys His Lys Phe  
115 120 125

Asn Ser Ser Gly Cys Pro Glu Arg Leu Ala Ser Cys Arg Ser Ile Asp  
130 135 140

Lys Phe Ala Gln Gly Trp Gly Pro Leu Thr Tyr Thr Glu Pro Asn Ser  
145 150 155 160

Ser Asp Gln Arg Pro Tyr Cys Trp His Tyr Ala Pro Arg Pro Cys Gly  
165 170 175

Ile Val Pro Ala Ser Gln Val Cys Gly Pro Val Tyr Cys Phe Thr Pro  
180 185 190

Ser Pro Val Val Val Gly Thr Thr Asp Arg Phe Gly Val Pro Thr Tyr  
195 200 205

Asn Trp Gly Ala Asn Asp Ser Asp Val Leu Ile Leu Asn Asn Thr Arg  
210 215 220

Pro Pro Arg Gly Asn Trp Phe Gly Cys Thr Trp Met Asn Gly Thr Gly  
225 230 235 240

Phe Thr Lys Thr Cys Gly Gly Pro Pro Cys Asn Ile Gly Gly Ala Gly  
245 250 255

Asn Asn Thr Leu Thr Cys Pro Thr Asp Cys Phe Arg Lys His Pro Glu  
260 265 270

Ala Thr Tyr Ala Arg Cys Gly Ser Gly Pro Trp Leu Thr Pro Arg Cys  
275 280 285

-continued

Met	Val	His	Tyr	Pro	Tyr	Arg	Leu	Trp	His	Tyr	Pro	Cys	Thr	Val	Asn
	290					295					300				
Phe	Thr	Ile	Phe	Lys	Val	Arg	Met	Tyr	Val	Gly	Gly	Val	Glu	His	Arg
	305				310					315					320
Phe	Glu	Ala	Ala	Cys	Asn	Trp	Thr	Arg	Gly	Glu	Arg	Cys	Asp	Leu	Glu
				325					330					335	
Asp	Arg	Asp	Arg	Ser	Glu	Leu	Ser	Pro	Leu	Leu	Leu	Ser	Thr	Thr	Glu
			340					345					350		
Trp	Gln	Ile	Leu	Pro	Cys	Ser	Phe	Thr	Thr	Leu	Pro	Ala	Leu	Ser	Thr
		355					360					365			
Gly	Leu	Ile	His	Leu	His	Gln	Asn	Ile	Val	Asp	Val	Gln	Tyr	Leu	Tyr
	370					375					380				
Gly	Val	Gly	Ser	Ala	Val	Val	Ser	Leu	Val	Ile	Lys	Trp	Glu	Tyr	Val
	385				390					395					400
Leu	Leu	Leu	Phe	Leu	Leu	Leu	Ala	Asp	Ala	Arg	Ile	Cys	Ala	Cys	Leu
			405						410					415	
Trp	Met	Met	Leu	Leu	Ile	Ala	Gln	Ala	Glu	Ala	Ala	Leu	Glu	Asn	Leu
			420					425					430		
Val	Val	Leu	Asn	Ala	Ala	Ala	Val	Ala	Gly	Ala	His	Gly	Thr	Leu	Ser
		435					440					445			
Phe	Leu	Val	Phe	Phe	Cys	Ala	Ala	Trp	Tyr	Ile	Lys	Gly	Arg	Leu	Val
	450					455					460				
Pro	Gly	Ala	Ala	Tyr	Ala	Phe	Tyr	Gly	Val	Trp	Pro	Leu	Leu	Leu	Leu
	465				470					475					480
Leu	Leu	Ala	Leu	Pro	Pro	Arg	Ala	Tyr	Ala						
			485					490							

## (2) INFORMATION FOR SEQ ID NO: 37:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1021 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

- (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 2..1018

- (ix) FEATURE:  
 (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 2..1015

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 37:

G	ATC	CCA	CAA	GCT	GTC	GTG	GAC	ATG	GTG	GCG	GGG	GCC	CAT	TGG	GGA	46
Ile	Pro	Gln	Ala	Val	Val	Asp	Met	Val	Ala	Gly	Ala	His	Trp	Gly		
	1			5					10					15		
GTC	CTG	GCG	GGC	CTC	GCC	TAC	TAT	TCC	ATG	GTG	GGG	AAC	TGG	GCT	AAG	94
Val	Leu	Ala	Gly	Leu	Ala	Tyr	Tyr	Ser	Met	Val	Gly	Asn	Trp	Ala	Lys	
			20					25						30		
GTT	TTG	GTT	GTG	ATG	CTA	CTC	TTT	GCC	GGC	GTC	GAC	GGG	CAT	ACC	CGC	142
Val	Leu	Val	Val	Met	Leu	Leu	Phe	Ala	Gly	Val	Asp	Gly	His	Thr	Arg	
			35					40					45			
GTG	TCA	GGA	GGG	GCA	GCA	GCC	TCC	GAT	ACC	AGG	GGC	CTT	GTG	TCC	CTC	190
Val	Ser	Gly	Gly	Ala	Ala	Ala	Ser	Asp	Thr	Arg	Gly	Leu	Val	Ser	Leu	

-continued

	50		55		60											
TTT	AGC	CCC	GGG	TCG	GCT	CAG	AAA	ATC	CAG	CTC	GTA	AAC	ACC	AAC	GGC	238
Phe	Ser	Pro	Gly	Ser	Ala	Gln	Lys	Ile	Gln	Leu	Val	Asn	Thr	Asn	Gly	
	65					70					75					
AGT	TGG	CAC	ATC	AAC	AGG	ACT	GCC	CTG	AAC	TGC	AAC	GAC	TCC	CTC	CAA	286
Ser	Trp	His	Ile	Asn	Arg	Thr	Ala	Leu	Asn	Cys	Asn	Asp	Ser	Leu	Gln	
	80				85					90					95	
ACA	GGG	TTC	TTT	GCC	GCA	CTA	TTC	TAC	AAA	CAC	AAA	TTC	AAC	TCG	TCT	334
Thr	Gly	Phe	Phe	Ala	Ala	Leu	Phe	Tyr	Lys	His	Lys	Phe	Asn	Ser	Ser	
				100					105					110		
GGA	TGC	CCA	GAG	CGC	TTG	GCC	AGC	TGT	CGC	TCC	ATC	GAC	AAG	TTC	GCT	382
Gly	Cys	Pro	Glu	Arg	Leu	Ala	Ser	Cys	Arg	Ser	Ile	Asp	Lys	Phe	Ala	
			115					120					125			
CAG	GGG	TGG	GGT	CCC	CTC	ACT	TAC	ACT	GAG	CCT	AAC	AGC	TCG	GAC	CAG	430
Gln	Gly	Trp	Gly	Pro	Leu	Thr	Tyr	Thr	Glu	Pro	Asn	Ser	Ser	Asp	Gln	
	130						135							140		
AGG	CCC	TAC	TGC	TGG	CAC	TAC	GCG	CCT	CGA	CCG	TGT	GGT	ATT	GTA	CCC	478
Arg	Pro	Tyr	Cys	Trp	His	Tyr	Ala	Pro	Arg	Pro	Cys	Gly	Ile	Val	Pro	
	145					150						155				
GCG	TCT	CAG	GTG	TGC	GGT	CCA	GTG	TAT	TGC	TTC	ACC	CCG	AGC	CCT	GTT	526
Ala	Ser	Gln	Val	Cys	Gly	Pro	Val	Tyr	Cys	Phe	Thr	Pro	Ser	Pro	Val	
	160				165					170					175	
GTG	GTG	GGG	ACG	ACC	GAT	CGG	TTT	GGT	GTC	CCC	ACG	TAT	AAC	TGG	GGG	574
Val	Val	Gly	Thr	Thr	Asp	Arg	Phe	Gly	Val	Pro	Thr	Tyr	Asn	Trp	Gly	
					180					185					190	
GCG	AAC	GAC	TCG	GAT	GTG	CTG	ATT	CTC	AAC	AAC	ACG	CGG	CCG	CCG	CGA	622
Ala	Asn	Asp	Ser	Asp	Val	Leu	Ile	Leu	Asn	Asn	Thr	Arg	Pro	Pro	Arg	
			195					200					205			
GGC	AAC	TGG	TTC	GGC	TGT	ACA	TGG	ATG	AAT	GGC	ACT	GGG	TTC	ACC	AAG	670
Gly	Asn	Trp	Phe	Gly	Cys	Thr	Trp	Met	Asn	Gly	Thr	Gly	Phe	Thr	Lys	
	210					215						220				
ACG	TGT	GGG	GGC	CCC	CCG	TGC	AAC	ATC	GGG	GGG	GCC	GGC	AAC	AAC	ACC	718
Thr	Cys	Gly	Gly	Pro	Pro	Cys	Asn	Ile	Gly	Gly	Ala	Gly	Asn	Asn	Thr	
	225					230					235					
TTG	ACC	TGC	CCC	ACT	GAC	TGT	TTT	CGG	AAG	CAC	CCC	GAG	GCC	ACC	TAC	766
Leu	Thr	Cys	Pro	Thr	Asp	Cys	Phe	Arg	Lys	His	Pro	Glu	Ala	Thr	Tyr	
	240				245					250					255	
GCC	AGA	TGC	GGT	TCT	GGG	CCC	TGG	CTG	ACA	CCT	AGG	TGT	ATG	GTT	CAT	814
Ala	Arg	Cys	Gly	Ser	Gly	Pro	Trp	Leu	Thr	Pro	Arg	Cys	Met	Val	His	
				260					265					270		
TAC	CCA	TAT	AGG	CTC	TGG	CAC	TAC	CCC	TGC	ACT	GTC	AAC	TTC	ACC	ATC	862
Tyr	Pro	Tyr	Arg	Leu	Trp	His	Tyr		Cys	Thr	Val	Asn	Phe	Thr	Ile	
			275					280					285			
TTC	AAG	GTT	AGG	ATG	TAC	GTG	GGG	GGC	GTG	GAG	CAC	AGG	TTC	GAA	GCC	910
Phe	Lys	Val	Arg	Met	Tyr	Val	Gly	Gly	Val	Glu	His	Arg	Phe	Glu	Ala	
	290					295						300				
GCA	TGC	AAT	TGG	ACT	CGA	GGA	GAG	CGT	TGT	GAC	TTG	GAG	GAC	AGG	GAT	958
Ala	Cys	Asn	Trp	Thr	Arg	Gly	Glu	Arg	Cys	Asp	Leu	Glu	Asp	Arg	Asp	
	305					310					315					
AGA	TCA	GAG	CTT	AGC	CCG	CTG	CTG	CTG	TCT	ACA	ACA	GAG	TGG	CAG	AGT	1006
Arg	Ser	Glu	Leu	Ser	Pro	Leu	Leu	Leu	Ser	Thr	Thr	Glu	Trp	Gln	Ser	
	320				325					330					335	
GGC	AGA	GCT	TAATTA													1021
Gly	Arg	Ala														

(2) INFORMATION FOR SEQ ID NO: 38:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 338 amino acids

(B) TYPE: amino acid



-continued

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 38:

Ile Pro Gln Ala Val Val Asp Met Val Ala Gly Ala His Trp Gly Val  
 1 5 10 15  
 Leu Ala Gly Leu Ala Tyr Tyr Ser Met Val Gly Asn Trp Ala Lys Val  
 20 25 30  
 Leu Val Val Met Leu Leu Phe Ala Gly Val Asp Gly His Thr Arg Val  
 35 40 45  
 Ser Gly Gly Ala Ala Ala Ser Asp Thr Arg Gly Leu Val Ser Leu Phe  
 50 55 60  
 Ser Pro Gly Ser Ala Gln Lys Ile Gln Leu Val Asn Thr Asn Gly Ser  
 65 70 75 80  
 Trp His Ile Asn Arg Thr Ala Leu Asn Cys Asn Asp Ser Leu Gln Thr  
 85 90 95  
 Gly Phe Phe Ala Ala Leu Phe Tyr Lys His Lys Phe Asn Ser Ser Gly  
 100 105 110  
 Cys Pro Glu Arg Leu Ala Ser Cys Arg Ser Ile Asp Lys Phe Ala Gln  
 115 120 125  
 Gly Trp Gly Pro Leu Thr Tyr Thr Glu Pro Asn Ser Ser Asp Gln Arg  
 130 135 140  
 Pro Tyr Cys Trp His Tyr Ala Pro Arg Pro Cys Gly Ile Val Pro Ala  
 145 150 155 160  
 Ser Gln Val Cys Gly Pro Val Tyr Cys Phe Thr Pro Ser Pro Val Val  
 165 170 175  
 Val Gly Thr Thr Asp Arg Phe Gly Val Pro Thr Tyr Asn Trp Gly Ala  
 180 185 190  
 Asn Asp Ser Asp Val Leu Ile Leu Asn Asn Thr Arg Pro Pro Arg Gly  
 195 200 205  
 Asn Trp Phe Gly Cys Thr Trp Met Asn Gly Thr Gly Phe Thr Lys Thr  
 210 215 220  
 Cys Gly Gly Pro Pro Cys Asn Ile Gly Gly Ala Gly Asn Asn Thr Leu  
 225 230 235 240  
 Thr Cys Pro Thr Asp Cys Phe Arg Lys His Pro Glu Ala Thr Tyr Ala  
 245 250 255  
 Arg Cys Gly Ser Gly Pro Trp Leu Thr Pro Arg Cys Met Val His Tyr  
 260 265 270  
 Pro Tyr Arg Leu Trp His Tyr Pro Cys Thr Val Asn Phe Thr Ile Phe  
 275 280 285  
 Lys Val Arg Met Tyr Val Gly Gly Val Glu His Arg Phe Glu Ala Ala  
 290 295 300  
 Cys Asn Trp Thr Arg Gly Glu Arg Cys Asp Leu Glu Asp Arg Asp Arg  
 305 310 315 320  
 Ser Glu Leu Ser Pro Leu Leu Leu Ser Thr Thr Glu Trp Gln Ser Gly  
 325 330 335  
 Arg Ala

(2) INFORMATION FOR SEQ ID NO: 39:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1034 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

-continued

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 2..1032

(ix) FEATURE:  
 (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 2..1029

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 39:

```

G ATC CCA CAA GCT GTC GTG GAC ATG GTG GCG GGG GCC CAT TGG GGA      46
  Ile Pro Gln Ala Val Val Asp Met Val Ala Gly Ala His Trp Gly
    1             5             10             15

GTC CTG GCG GGC CTC GCC TAC TAT TCC ATG GTG GGG AAC TGG GCT AAG      94
  Val Leu Ala Gly Leu Ala Tyr Tyr Ser Met Val Gly Asn Trp Ala Lys
    20             25             30

GTT TTG GTT GTG ATG CTA CTC TTT GCC GGC GTC GAC GGG CAT ACC CGC     142
  Val Leu Val Val Met Leu Leu Phe Ala Gly Val Asp Gly His Thr Arg
    35             40             45

GTG TCA GGA GGG GCA GCA GCC TCC GAT ACC AGG GGC CTT GTG TCC CTC     190
  Val Ser Gly Gly Ala Ala Ala Ser Asp Thr Arg Gly Leu Val Ser Leu
    50             55             60

TTT AGC CCC GGG TCG GCT CAG AAA ATC CAG CTC GTA AAC ACC AAC GGC     238
  Phe Ser Pro Gly Ser Ala Gln Lys Ile Gln Leu Val Asn Thr Asn Gly
    65             70             75

AGT TGG CAC ATC AAC AGG ACT GCC CTG AAC TGC AAC GAC TCC CTC CAA     286
  Ser Trp His Ile Asn Arg Thr Ala Leu Asn Cys Asn Asp Ser Leu Gln
    80             85             90             95

ACA GGG TTC TTT GCC GCA CTA TTC TAC AAA CAC AAA TTC AAC TCG TCT     334
  Thr Gly Phe Phe Ala Ala Leu Phe Tyr Lys His Lys Phe Asn Ser Ser
    100            105            110

GGA TGC CCA GAG CGC TTG GCC AGC TGT CGC TCC ATC GAC AAG TTC GCT     382
  Gly Cys Pro Glu Arg Leu Ala Ser Cys Arg Ser Ile Asp Lys Phe Ala
    115            120            125

CAG GGG TGG GGT CCC CTC ACT TAC ACT GAG CCT AAC AGC TCG GAC CAG     430
  Gln Gly Trp Gly Pro Leu Thr Tyr Thr Glu Pro Asn Ser Ser Asp Gln
    130            135            140

AGG CCC TAC TGC TGG CAC TAC GCG CCT CGA CCG TGT GGT ATT GTA CCC     478
  Arg Pro Tyr Cys Trp His Tyr Ala Pro Arg Pro Cys Gly Ile Val Pro
    145            150            155

GCG TCT CAG GTG TGC GGT CCA GTG TAT TGC TTC ACC CCG AGC CCT GTT     526
  Ala Ser Gln Val Cys Gly Pro Val Tyr Cys Phe Thr Pro Ser Pro Val
    160            165            170            175

GTG GTG GGG ACG ACC GAT CGG TTT GGT GTC CCC ACG TAT AAC TGG GGG     574
  Val Val Gly Thr Thr Asp Arg Phe Gly Val Pro Thr Tyr Asn Trp Gly
    180            185            190

GCG AAC GAC TCG GAT GTG CTG ATT CTC AAC AAC ACG CGG CCG CCG CGA     622
  Ala Asn Asp Ser Asp Val Leu Ile Leu Asn Asn Thr Arg Pro Pro Arg
    195            200            205

GGC AAC TGG TTC GGC TGT ACA TGG ATG AAT GGC ACT GGG TTC ACC AAG     670
  Gly Asn Trp Phe Gly Cys Thr Trp Met Asn Gly Thr Gly Phe Thr Lys
    210            215            220

ACG TGT GGG GGC CCC CCG TGC AAC ATC GGG GGG GCC GGC AAC AAC ACC     718
  Thr Cys Gly Gly Pro Pro Cys Asn Ile Gly Gly Ala Gly Asn Asn Thr
    225            230            235

TTG ACC TGC CCC ACT GAC TGT TTT CGG AAG CAC CCC GAG GCC ACC TAC     766
  Leu Thr Cys Pro Thr Asp Cys Phe Arg Lys His Pro Glu Ala Thr Tyr

```

-continued

240	245	250	255	
GCC AGA TGC GGT TCT	GGG CCC TGG CTG	ACA CCT AGG TGT	ATG GTT CAT	814
Ala Arg Cys Gly Ser	Gly Pro Trp Leu Thr	Pro Arg Cys Met	Val His	
	260	265	270	
TAC CCA TAT AGG CTC	TGG CAC TAC CCC	TGC ACT GTC AAC	TTC ACC ATC	862
Tyr Pro Tyr Arg Leu	Trp His Tyr Pro	Cys Thr Val Asn	Phe Thr Ile	
	275	280	285	
TTC AAG GTT AGG ATG	TAC GTG GGG GGC	GTG GAG CAC AGG	TTC GAA GCC	910
Phe Lys Val Arg Met	Tyr Val Gly Gly	Val Glu His Arg	Phe Glu Ala	
	290	295	300	
GCA TGC AAT TGG ACT	CGA GGA GAG CGT	TGT GAC TTG GAG	GAC AGG GAT	958
Ala Cys Asn Trp Thr	Arg Gly Glu Arg	Cys Asp Leu Glu	Asp Arg Asp	
	305	310	315	
AGA TCA GAG CTT AGC	CCG CTG CTG CTG	TCT ACA ACA GGT	GAT CGA GGG	1006
Arg Ser Glu Leu Ser	Pro Leu Leu Leu	Ser Thr Thr Gly	Asp Arg Gly	
	320	325	330	335
CAG ACA CCA TCA CCA	CCA TCA CTA AT AG			1034
Gln Thr Pro Ser Pro	Pro Ser Leu			
	340			

## (2) INFORMATION FOR SEQ ID NO: 40:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 343 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 40:

Ile	Pro	Gln	Ala	Val	Val	Asp	Met	Val	Ala	Gly	Ala	His	Trp	Gly	Val
1				5					10					15	
Leu	Ala	Gly	Leu	Ala	Tyr	Tyr	Ser	Met	Val	Gly	Asn	Trp	Ala	Lys	Val
			20					25					30		
Leu	Val	Val	Met	Leu	Leu	Phe	Ala	Gly	Val	Asp	Gly	His	Thr	Arg	Val
		35				40						45			
Ser	Gly	Gly	Ala	Ala	Ala	Ser	Asp	Thr	Arg	Gly	Leu	Val	Ser	Leu	Phe
	50					55					60				
Ser	Pro	Gly	Ser	Ala	Gln	Lys	Ile	Gln	Leu	Val	Asn	Thr	Asn	Gly	Ser
	65				70				75					80	
Trp	His	Ile	Asn	Arg	Thr	Ala	Leu	Asn	Cys	Asn	Asp	Ser	Leu	Gln	Thr
			85					90					95		
Gly	Phe	Phe	Ala	Ala	Leu	Phe	Tyr	Lys	His	Lys	Phe	Asn	Ser	Ser	Gly
			100				105					110			
Cys	Pro	Glu	Arg	Leu	Ala	Ser	Cys	Arg	Ser	Ile	Asp	Lys	Phe	Ala	Gln
	115					120					125				
Gly	Trp	Gly	Pro	Leu	Thr	Tyr	Thr	Glu	Pro	Asn	Ser	Ser	Asp	Gln	Arg
	130				135					140					
Pro	Tyr	Cys	Trp	His	Tyr	Ala	Pro	Arg	Pro	Cys	Gly	Ile	Val	Pro	Ala
	145				150				155					160	
Ser	Gln	Val	Cys	Gly	Pro	Val	Tyr	Cys	Phe	Thr	Pro	Ser	Pro	Val	Val
			165				170						175		
Val	Gly	Thr	Thr	Asp	Arg	Phe	Gly	Val	Pro	Thr	Tyr	Asn	Trp	Gly	Ala
			180				185						190		
Asn	Asp	Ser	Asp	Val	Leu	Ile	Leu	Asn	Asn	Thr	Arg	Pro	Pro	Arg	Gly
	195				200							205			
Asn	Trp	Phe	Gly	Cys	Thr	Trp	Met	Asn	Gly	Thr	Gly	Phe	Thr	Lys	Thr
	210				215						220				

-continued

Cys Gly Gly Pro Pro Cys Asn Ile Gly Gly Ala Gly Asn Asn Thr Leu  
 225 230 235 240  
 Thr Cys Pro Thr Asp Cys Phe Arg Lys His Pro Glu Ala Thr Tyr Ala  
 245 250 255  
 Arg Cys Gly Ser Gly Pro Trp Leu Thr Pro Arg Cys Met Val His Tyr  
 260 265 270  
 Pro Tyr Arg Leu Trp His Tyr Pro Cys Thr Val Asn Phe Thr Ile Phe  
 275 280 285  
 Lys Val Arg Met Tyr Val Gly Gly Val Glu His Arg Phe Glu Ala Ala  
 290 295 300  
 Cys Asn Trp Thr Arg Gly Glu Arg Cys Asp Leu Glu Asp Arg Asp Arg  
 305 310 315 320  
 Ser Glu Leu Ser Pro Leu Leu Leu Ser Thr Thr Gly Asp Arg Gly Gln  
 325 330 335  
 Thr Pro Ser Pro Pro Ser Leu  
 340

## (2) INFORMATION FOR SEQ ID NO: 41:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 945 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (iii) HYPOTHETICAL: NO

## (iii) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..942

## (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide
- (B) LOCATION: 1..939

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 41:

ATG GTG GGG AAC TGG GCT AAG GTT TTG GTT GTG ATG CTA CTC TTT GCC	48
Met Val Gly Asn Trp Ala Lys Val Leu Val Val Met Leu Leu Phe Ala	
1 5 10 15	
GGC GTC GAC GGG CAT ACC CGC GTG TCA GGA GGG GCA GCA GCC TCC GAT	96
Gly Val Asp Gly His Thr Arg Val Ser Gly Gly Ala Ala Ala Ser Asp	
20 25 30	
ACC AGG GGC CTT GTG TCC CTC TTT AGC CCC GGG TCG GCT CAG AAA ATC	144
Thr Arg Gly Leu Val Ser Leu Phe Ser Pro Gly Ser Ala Gln Lys Ile	
35 40 45	
CAG CTC GTA AAC ACC AAC GGC AGT TGG CAC ATC AAC AGG ACT GCC CTG	192
Gln Leu Val Asn Thr Asn Gly Ser Trp His Ile Asn Arg Thr Ala Leu	
50 55 60	
AAC TGC AAC GAC TCC CTC CAA ACA GGG TTC TTT GCC GCA CTA TTC TAC	240
Asn Cys Asn Asp Ser Leu Gln Thr Gly Phe Phe Ala Ala Leu Phe Tyr	
65 70 75 80	
AAA CAC AAA TTC AAC TCG TCT GGA TGC CCA GAG CGC TTG GCC AGC TGT	288
Lys His Lys Phe Asn Ser Ser Gly Cys Pro Glu Arg Leu Ala Ser Cys	
85 90 95	
CGC TCC ATC GAC AAG TTC GCT CAG GGG TGG GGT CCC CTC ACT TAC ACT	336
Arg Ser Ile Asp Lys Phe Ala Gln Gly Trp Gly Pro Leu Thr Tyr Thr	
100 105 110	
GAG CCT AAC AGC TCG GAC CAG AGG CCC TAC TGC TGG CAC TAC GCG CCT	384

-continued

Glu	Pro	Asn	Ser	Ser	Asp	Gln	Arg	Pro	Tyr	Cys	Trp	His	Tyr	Ala	Pro		
		115					120					125					
CGA	CCG	TGT	GGT	ATT	GTA	CCC	GCG	TCT	CAG	GTG	TGC	GGT	CCA	GTG	TAT		432
Arg	Pro	Cys	Gly	Ile	Val	Pro	Ala	Ser	Gln	Val	Cys	Gly	Pro	Val	Tyr		
		130				135					140						
TGC	TTC	ACC	CCG	AGC	CCT	GTT	GTG	GTG	GGG	ACG	ACC	GAT	CGG	TTT	GGT		480
Cys	Phe	Thr	Pro	Ser	Pro	Val	Val	Val	Gly	Thr	Thr	Asp	Arg	Phe	Gly		
		145			150				155						160		
GTC	CCC	ACG	TAT	AAC	TGG	GGG	GCG	AAC	GAC	TCG	GAT	GTG	CTG	ATT	CTC		528
Val	Pro	Thr	Tyr	Asn	Trp	Gly	Ala	Asn	Asp	Ser	Asp	Val	Leu	Ile	Leu		
				165					170					175			
AAC	AAC	ACG	CGG	CCG	CCG	CGA	GGC	AAC	TGG	TTC	GGC	TGT	ACA	TGG	ATG		576
Asn	Asn	Thr	Arg	Pro	Pro	Arg	Gly	Asn	Trp	Phe	Gly	Cys	Thr	Trp	Met		
			180					185						190			
AAT	GGC	ACT	GGG	TTC	ACC	AAG	ACG	TGT	GGG	GGC	CCC	CCG	TGC	AAC	ATC		624
Asn	Gly	Thr	Gly	Phe	Thr	Lys	Thr	Cys	Gly	Gly	Pro	Pro	Cys	Asn	Ile		
		195				200						205					
GGG	GGG	GCC	GGC	AAC	AAC	ACC	TTG	ACC	TGC	CCC	ACT	GAC	TGT	TTT	CGG		672
Gly	Gly	Ala	Gly	Asn	Asn	Thr	Leu	Thr	Cys	Pro	Thr	Asp	Cys	Phe	Arg		
		210				215					220						
AAG	CAC	CCC	GAG	GCC	ACC	TAC	GCC	AGA	TGC	GGT	TCT	GGG	CCC	TGG	CTG		720
Lys	His	Pro	Glu	Ala	Thr	Tyr	Ala	Arg	Cys	Gly	Ser	Gly	Pro	Trp	Leu		
		225			230				235						240		
ACA	CCT	AGG	TGT	ATG	GTT	CAT	TAC	CCA	TAT	AGG	CTC	TGG	CAC	TAC	CCC		768
Thr	Pro	Arg	Cys	Met	Val	His	Tyr	Pro	Tyr	Arg	Leu	Trp	His	Tyr	Pro		
				245					250					255			
TGC	ACT	GTC	AAC	TTC	ACC	ATC	TTC	AAG	GTT	AGG	ATG	TAC	GTG	GGG	GGC		816
Cys	Thr	Val	Asn	Phe	Thr	Ile	Phe	Lys	Val	Arg	Met	Tyr	Val	Gly	Gly		
			260					265					270				
GTG	GAG	CAC	AGG	TTC	GAA	GCC	GCA	TGC	AAT	TGG	ACT	CGA	GGA	GAG	CGT		864
Val	Glu	His	Arg	Phe	Glu	Ala	Ala	Cys	Asn	Trp	Thr	Arg	Gly	Glu	Arg		
		275				280						285					
TGT	GAC	TTG	GAG	GAC	AGG	GAT	AGA	TCA	GAG	CTT	AGC	CCG	CTG	CTG	CTG		912
Cys	Asp	Leu	Glu	Asp	Arg	Asp	Arg	Ser	Glu	Leu	Ser	Pro	Leu	Leu	Leu		
		290				295					300						
TCT	ACA	ACA	GAG	TGG	CAG	AGC	TTA	ATT	AAT	TAG							945
Ser	Thr	Thr	Glu	Trp	Gln	Ser	Leu	Ile	Asn								
				305			310										

## (2) INFORMATION FOR SEQ ID NO: 42:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 314 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 42:

Met	Val	Gly	Asn	Trp	Ala	Lys	Val	Leu	Val	Val	Met	Leu	Leu	Phe	Ala		
1				5					10					15			
Gly	Val	Asp	Gly	His	Thr	Arg	Val	Ser	Gly	Gly	Ala	Ala	Ala	Ser	Asp		
			20					25						30			
Thr	Arg	Gly	Leu	Val	Ser	Leu	Phe	Ser	Pro	Gly	Ser	Ala	Gln	Lys	Ile		
		35					40						45				
Gln	Leu	Val	Asn	Thr	Asn	Gly	Ser	Trp	His	Ile	Asn	Arg	Thr	Ala	Leu		
		50				55						60					
Asn	Cys	Asn	Asp	Ser	Leu	Gln	Thr	Gly	Phe	Phe	Ala	Ala	Leu	Phe	Tyr		
		65			70					75					80		
Lys	His	Lys	Phe	Asn	Ser	Ser	Gly	Cys	Pro	Glu	Arg	Leu	Ala	Ser	Cys		

-continued

		85			90			95							
Arg	Ser	Ile	Asp	Lys	Phe	Ala	Gln	Gly	Trp	Gly	Pro	Leu	Thr	Tyr	Thr
			100					105					110		
Glu	Pro	Asn	Ser	Ser	Asp	Gln	Arg	Pro	Tyr	Cys	Trp	His	Tyr	Ala	Pro
		115					120					125			
Arg	Pro	Cys	Gly	Ile	Val	Pro	Ala	Ser	Gln	Val	Cys	Gly	Pro	Val	Tyr
		130					135					140			
Cys	Phe	Thr	Pro	Ser	Pro	Val	Val	Val	Gly	Thr	Thr	Asp	Arg	Phe	Gly
145					150					155					160
Val	Pro	Thr	Tyr	Asn	Trp	Gly	Ala	Asn	Asp	Ser	Asp	Val	Leu	Ile	Leu
				165					170					175	
Asn	Asn	Thr	Arg	Pro	Pro	Arg	Gly	Asn	Trp	Phe	Gly	Cys	Thr	Trp	Met
			180					185					190		
Asn	Gly	Thr	Gly	Phe	Thr	Lys	Thr	Cys	Gly	Gly	Pro	Pro	Cys	Asn	Ile
		195					200					205			
Gly	Gly	Ala	Gly	Asn	Asn	Thr	Leu	Thr	Cys	Pro	Thr	Asp	Cys	Phe	Arg
		210					215					220			
Lys	His	Pro	Glu	Ala	Thr	Tyr	Ala	Arg	Cys	Gly	Ser	Gly	Pro	Trp	Leu
225					230					235					240
Thr	Pro	Arg	Cys	Met	Val	His	Tyr	Pro	Tyr	Arg	Leu	Trp	His	Tyr	Pro
				245					250					255	
Cys	Thr	Val	Asn	Phe	Thr	Ile	Phe	Lys	Val	Arg	Met	Tyr	Val	Gly	Gly
			260					265					270		
Val	Glu	His	Arg	Phe	Glu	Ala	Ala	Cys	Asn	Trp	Thr	Arg	Gly	Glu	Arg
		275					280					285			
Cys	Asp	Leu	Glu	Asp	Arg	Asp	Arg	Ser	Glu	Leu	Ser	Pro	Leu	Leu	Leu
	290					295					300				
Ser	Thr	Thr	Glu	Trp	Gln	Ser	Leu	Ile	Asn						
305					310										

## (2) INFORMATION FOR SEQ ID NO: 43:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 961 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS  
 (B) LOCATION: 1..958

## (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 1..955

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 43:

ATG	GTG	GGG	AAC	TGG	GCT	AAG	GTT	TTG	GTT	GTG	ATG	CTA	CTC	TTT	GCC	48
Met	Val	Gly	Asn	Trp	Ala	Lys	Val	Leu	Val	Val	Met	Leu	Leu	Phe	Ala	
1				5				10						15		
GGC	GTC	GAC	GGG	CAT	ACC	CGC	GTG	TCA	GGA	GGG	GCA	GCA	GCC	TCC	GAT	96
Gly	Val	Asp	Gly	His	Thr	Arg	Val	Ser	Gly	Gly	Ala	Ala	Ala	Ser	Asp	
			20					25					30			
ACC	AGG	GGC	CTT	GTG	TCC	CTC	TTT	AGC	CCC	GGG	TCG	GCT	CAG	AAA	ATC	144
Thr	Arg	Gly	Leu	Val	Ser	Leu	Phe	Ser	Pro	Gly	Ser	Ala	Gln	Lys	Ile	

-continued

35			40			45										
CAG	CTC	GTA	AAC	ACC	AAC	GGC	AGT	TGG	CAC	ATC	AAC	AGG	ACT	GCC	CTG	192
Gln	Leu	Val	Asn	Thr	Asn	Gly	Ser	Trp	His	Ile	Asn	Arg	Thr	Ala	Leu	
	50					55					60					
AAC	TGC	AAC	GAC	TCC	CTC	CAA	ACA	GGG	TTC	TTT	GCC	GCA	CTA	TTC	TAC	240
Asn	Cys	Asn	Asp	Ser	Leu	Gln	Thr	Gly	Phe	Phe	Ala	Ala	Leu	Phe	Tyr	
	65				70					75					80	
AAA	CAC	AAA	TTC	AAC	TCG	TCT	GGA	TGC	CCA	GAG	CGC	TTG	GCC	AGC	TGT	288
Lys	His	Lys	Phe	Asn	Ser	Ser	Gly	Cys	Pro	Glu	Arg	Leu	Ala	Ser	Cys	
				85					90					95		
CGC	TCC	ATC	GAC	AAG	TTC	GCT	CAG	GGG	TGG	GGT	CCC	CTC	ACT	TAC	ACT	336
Arg	Ser	Ile	Asp	Lys	Phe	Ala	Gln	Gly	Trp	Gly	Pro	Leu	Thr	Tyr	Thr	
		100						105						110		
GAG	CCT	AAC	AGC	TCG	GAC	CAG	AGG	CCC	TAC	TGC	TGG	CAC	TAC	GCG	CCT	384
Glu	Pro	Asn	Ser	Ser	Asp	Gln	Arg	Pro	Tyr	Cys	Trp	His	Tyr	Ala	Pro	
		115					120						125			
CGA	CCG	TGT	GGT	ATT	GTA	CCC	GCG	TCT	CAG	GTG	TGC	GGT	CCA	GTG	TAT	432
Arg	Pro	Cys	Gly	Ile	Val	Pro	Ala	Ser	Gln	Val	Cys	Gly	Pro	Val	Tyr	
	130					135					140					
TGC	TTC	ACC	CCG	AGC	CCT	GTT	GTG	GTG	GGG	ACG	ACC	GAT	CGG	TTT	GGT	480
Cys	Phe	Thr	Pro	Ser	Pro	Val	Val	Val	Gly	Thr	Thr	Asp	Arg	Phe	Gly	
	145				150					155					160	
GTC	CCC	ACG	TAT	AAC	TGG	GGG	GCG	AAC	GAC	TCG	GAT	GTG	CTG	ATT	CTC	528
Val	Pro	Thr	Tyr	Asn	Trp	Gly	Ala	Asn	Asp	Ser	Asp	Val	Leu	Ile	Leu	
				165					170					175		
AAC	AAC	ACG	CGG	CCG	CCG	CGA	GGC	AAC	TGG	TTC	GGC	TGT	ACA	TGG	ATG	576
Asn	Asn	Thr	Arg	Pro	Pro	Arg	Gly	Asn	Trp	Phe	Gly	Cys	Thr	Trp	Met	
			180					185					190			
AAT	GGC	ACT	GGG	TTC	ACC	AAG	ACG	TGT	GGG	GGC	CCC	CCG	TGC	AAC	ATC	624
Asn	Gly	Thr	Gly	Phe	Thr	Lys	Thr	Cys	Gly	Gly	Pro	Pro	Cys	Asn	Ile	
		195				200						205				
GGG	GGG	GCC	GGC	AAC	AAC	ACC	TTG	ACC	TGC	CCC	ACT	GAC	TGT	TTT	CGG	672
Gly	Gly	Ala	Gly	Asn	Asn	Thr	Leu	Thr	Cys	Pro	Thr	Asp	Cys	Phe	Arg	
	210					215						220				
AAG	CAC	CCC	GAG	GCC	ACC	TAC	GCC	AGA	TGC	GGT	TCT	GGG	CCC	TGG	CTG	720
Lys	His	Pro	Glu	Ala	Thr	Tyr	Ala	Arg	Cys	Gly	Ser	Gly	Pro	Trp	Leu	
	225				230				235					240		
ACA	CCT	AGG	TGT	ATG	GTT	CAT	TAC	CCA	TAT	AGG	CTC	TGG	CAC	TAC	CCC	768
Thr	Pro	Arg	Cys	Met	Val	His	Tyr	Pro	Tyr	Arg	Leu	Trp	His	Tyr	Pro	
				245				250						255		
TGC	ACT	GTC	AAC	TTC	ACC	ATC	TTC	AAG	GTT	AGG	ATG	TAC	GTG	GGG	GGC	816
Cys	Thr	Val	Asn	Phe	Thr	Ile	Phe	Lys	Val	Arg	Met	Tyr	Val	Gly	Gly	
			260					265					270			
GTG	GAG	CAC	AGG	TTC	GAA	GCC	GCA	TGC	AAT	TGG	ACT	CGA	GGA	GAG	CGT	864
Val	Glu	His	Arg	Phe	Glu	Ala	Ala	Cys	Asn	Trp	Thr	Arg	Gly	Glu	Arg	
		275				280						285				
TGT	GAC	TTG	GAG	GAC	AGG	GAT	AGA	TCA	GAG	CTT	AGC	CCG	CTG	CTG	CTG	912
Cys	Asp	Leu	Glu	Asp	Arg	Asp	Arg	Ser	Glu	Leu	Ser	Pro	Leu	Leu	Leu	
	290					295						300				
TCT	ACA	ACA	GGT	GAT	CGA	GGG	CAG	ACA	CCA	TCA	CCA	CCA	TCA	CTA	A	958
Ser	Thr	Thr	Gly	Asp	Arg	Gly	Gln	Thr	Pro	Ser	Pro	Pro	Ser	Leu		
	305				310					315						
TAG																961

(2) INFORMATION FOR SEQ ID NO: 44:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 319 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

-continued

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 44:

```

Met Val Gly Asn Trp Ala Lys Val Leu Val Val Met Leu Leu Phe Ala
 1          5          10          15
Gly Val Asp Gly His Thr Arg Val Ser Gly Gly Ala Ala Ala Ser Asp
          20          25          30
Thr Arg Gly Leu Val Ser Leu Phe Ser Pro Gly Ser Ala Gln Lys Ile
          35          40          45
Gln Leu Val Asn Thr Asn Gly Ser Trp His Ile Asn Arg Thr Ala Leu
          50          55          60
Asn Cys Asn Asp Ser Leu Gln Thr Gly Phe Phe Ala Ala Leu Phe Tyr
          65          70          75          80
Lys His Lys Phe Asn Ser Ser Gly Cys Pro Glu Arg Leu Ala Ser Cys
          85          90          95
Arg Ser Ile Asp Lys Phe Ala Gln Gly Trp Gly Pro Leu Thr Tyr Thr
          100          105          110
Glu Pro Asn Ser Ser Asp Gln Arg Pro Tyr Cys Trp His Tyr Ala Pro
          115          120          125
Arg Pro Cys Gly Ile Val Pro Ala Ser Gln Val Cys Gly Pro Val Tyr
          130          135          140
Cys Phe Thr Pro Ser Pro Val Val Val Gly Thr Thr Asp Arg Phe Gly
          145          150          155          160
Val Pro Thr Tyr Asn Trp Gly Ala Asn Asp Ser Asp Val Leu Ile Leu
          165          170          175
Asn Asn Thr Arg Pro Pro Arg Gly Asn Trp Phe Gly Cys Thr Trp Met
          180          185          190
Asn Gly Thr Gly Phe Thr Lys Thr Cys Gly Gly Pro Pro Cys Asn Ile
          195          200          205
Gly Gly Ala Gly Asn Asn Thr Leu Thr Cys Pro Thr Asp Cys Phe Arg
          210          215          220
Lys His Pro Glu Ala Thr Tyr Ala Arg Cys Gly Ser Gly Pro Trp Leu
          225          230          235          240
Thr Pro Arg Cys Met Val His Tyr Pro Tyr Arg Leu Trp His Tyr Pro
          245          250          255
Cys Thr Val Asn Phe Thr Ile Phe Lys Val Arg Met Tyr Val Gly Gly
          260          265          270
Val Glu His Arg Phe Glu Ala Ala Cys Asn Trp Thr Arg Gly Glu Arg
          275          280          285
Cys Asp Leu Glu Asp Arg Asp Arg Ser Glu Leu Ser Pro Leu Leu Leu
          290          295          300
Ser Thr Thr Gly Asp Arg Gly Gln Thr Pro Ser Pro Pro Ser Leu
          305          310          315

```

(2) INFORMATION FOR SEQ ID NO: 45:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1395 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO



-continued

## (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..1392

## (ix) FEATURE:

(A) NAME/KEY: mat\_peptide

(B) LOCATION: 1..1389

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 45:

ATG	GTG	GCG	GGG	GCC	CAT	TGG	GGA	GTC	CTG	GCG	GGC	CTC	GCC	TAC	TAT	48
Met	Val	Ala	Gly	Ala	His	Trp	Gly	Val	Leu	Ala	Gly	Leu	Ala	Tyr	Tyr	
1				5					10					15		
TCC	ATG	GTG	GGG	AAC	TGG	GCT	AAG	GTT	TTG	GTT	GTG	ATG	CTA	CTC	TTT	96
Ser	Met	Val	Gly	Asn	Trp	Ala	Lys	Val	Leu	Val	Val	Met	Leu	Leu	Phe	
			20					25					30			
GCC	GGC	GTC	GAC	GGG	CAT	ACC	CGC	GTG	TCA	GGA	GGG	GCA	GCA	GCC	TCC	144
Ala	Gly	Val	Asp	Gly	His	Thr	Arg	Val	Ser	Gly	Gly	Ala	Ala	Ala	Ser	
		35					40					45				
GAT	ACC	AGG	GGC	CTT	GTG	TCC	CTC	TTT	AGC	CCC	GGG	TCG	GCT	CAG	AAA	192
Asp	Thr	Arg	Gly	Leu	Val	Ser	Leu	Phe	Ser	Pro	Gly	Ser	Ala	Gln	Lys	
	50					55					60					
ATC	CAG	CTC	GTA	AAC	ACC	AAC	GGC	AGT	TGG	CAC	ATC	AAC	AGG	ACT	GCC	240
Ile	Gln	Leu	Val	Asn	Thr	Asn	Gly	Ser	Trp	His	Ile	Asn	Arg	Thr	Ala	
65					70					75					80	
CTG	AAC	TGC	AAC	GAC	TCC	CTC	CAA	ACA	GGG	TTC	TTT	GCC	GCA	CTA	TTC	288
Leu	Asn	Cys	Asn	Asp	Ser	Leu	Gln	Thr	Gly	Phe	Phe	Ala	Ala	Leu	Phe	
				85					90					95		
TAC	AAA	CAC	AAA	TTC	AAC	TCG	TCT	GGA	TGC	CCA	GAG	CGC	TTG	GCC	AGC	336
Tyr	Lys	His	Lys	Phe	Asn	Ser	Ser	Gly	Cys	Pro	Glu	Arg	Leu	Ala	Ser	
			100					105					110			
TGT	CGC	TCC	ATC	GAC	AAG	TTC	GCT	CAG	GGG	TGG	GGT	CCC	CTC	ACT	TAC	384
Cys	Arg	Ser	Ile	Asp	Lys	Phe	Ala	Gln	Gly	Trp	Gly	Pro	Leu	Thr	Tyr	
		115					120					125				
ACT	GAG	CCT	AAC	AGC	TCG	GAC	CAG	AGG	CCC	TAC	TGC	TGG	CAC	TAC	GCG	432
Thr	Glu	Pro	Asn	Ser	Ser	Asp	Gln	Arg	Pro	Tyr	Cys	Trp	His	Tyr	Ala	
	130					135						140				
CCT	CGA	CCG	TGT	GGT	ATT	GTA	CCC	GCG	TCT	CAG	GTG	TGC	GGT	CCA	GTG	480
Pro	Arg	Pro	Cys	Gly	Ile	Val	Pro	Ala	Ser	Gln	Val	Cys	Gly	Pro	Val	
145					150					155					160	
TAT	TGC	TTC	ACC	CCG	AGC	CCT	GTT	GTG	GTG	GGG	ACG	ACC	GAT	CGG	TTT	528
Tyr	Cys	Phe	Thr	Pro	Ser	Pro	Val	Val	Gly	Thr	Thr	Asp	Arg	Phe		
				165					170					175		
GGT	GTC	CCC	ACG	TAT	AAC	TGG	GGG	GCG	AAC	GAC	TCG	GAT	GTG	CTG	ATT	576
Gly	Val	Pro	Thr	Tyr	Asn	Trp	Gly	Ala	Asn	Asp	Ser	Asp	Val	Leu	Ile	
			180					185					190			
CTC	AAC	AAC	ACG	CGG	CCG	CCG	CGA	GGC	AAC	TGG	TTC	GGC	TGT	ACA	TGG	624
Leu	Asn	Asn	Thr	Arg	Pro	Pro	Arg	Gly	Asn	Trp	Phe	Gly	Cys	Thr	Trp	
			195				200						205			
ATG	AAT	GGC	ACT	GGG	TTC	ACC	AAG	ACG	TGT	GGG	GGC	CCC	CCG	TGC	AAC	672
Met	Asn	Gly	Thr	Gly	Phe	Thr	Lys	Thr	Cys	Gly	Gly	Pro	Pro	Cys	Asn	
	210						215					220				
ATC	GGG	GGG	GCC	GGC	AAC	AAC	ACC	TTG	ACC	TGC	CCC	ACT	GAC	TGT	TTT	720
Ile	Gly	Gly	Ala	Gly	Asn	Asn	Thr	Leu	Thr	Cys	Pro	Thr	Asp	Cys	Phe	
225					230					235					240	
CGG	AAG	CAC	CCC	GAG	GCC	ACC	TAC	GCC	AGA	TGC	GGT	TCT	GGG	CCC	TGG	768
Arg	Lys	His	Pro	Glu	Ala	Thr	Tyr	Ala	Arg	Cys	Gly	Ser	Gly	Pro	Trp	
				245					250					255		
CTG	ACA	CCT	AGG	TGT	ATG	GTT	CAT	TAC	CCA	TAT	AGG	CTC	TGG	CAC	TAC	816
Leu	Thr	Pro	Arg	Cys	Met	Val	His	Tyr	Pro	Tyr	Arg	Leu	Trp	His	Tyr	
			260					265						270		

-continued

CCC Pro	TGC Cys	ACT Thr	GTC Val	AAC Asn	TTC Phe	ACC Thr	ATC Ile	TTC Phe	AAG Lys	GTT Val	AGG Arg	ATG Met	TAC Tyr	GTG Val	GGG Gly	864
	275						280					285				
GGC Gly	GTG Val	GAG Glu	CAC His	AGG Arg	TTC Phe	GAA Glu	GCC Ala	GCA Ala	TGC Cys	AAT Asn	TGG Trp	ACT Thr	CGA Arg	GGA Gly	GAG Glu	912
	290					295					300					
CGT Arg	TGT Cys	GAC Asp	TTG Leu	GAG Glu	GAC Asp	AGG Arg	GAT Asp	AGA Arg	TCA Ser	GAG Glu	CTT Leu	AGC Ser	CCG Pro	CTG Leu	CTG Leu	960
	305				310					315				320		
CTG Leu	TCT Ser	ACA Thr	ACA Thr	GAG Glu	TGG Trp	CAG Gln	ATA Ile	CTG Leu	CCC Pro	TGT Cys	TCC Ser	TTC Phe	ACC Thr	ACC Thr	CTG Leu	1008
				325					330					335		
CCG Pro	GCC Ala	CTA Leu	TCC Ser	ACC Thr	GGC Gly	CTG Leu	ATC Ile	CAC His	CTC Leu	CAT His	CAG Gln	AAC Asn	ATC Ile	GTG Val	GAC Asp	1056
			340					345					350			
GTG Val	CAA Gln	TAC Tyr	CTG Leu	TAC Tyr	GGT Gly	GTA Val	GGG Gly	TCG Ser	GCG Ala	GTT Val	GTC Val	TCC Ser	CTT Leu	GTC Val	ATC Ile	1104
		355				360						365				
AAA Lys	TGG Trp	GAG Glu	TAT Tyr	GTC Val	CTG Leu	TTG Leu	CTC Leu	TTC Phe	CTT Leu	CTC Leu	CTG Leu	GCA Ala	GAC Asp	GCG Ala	CGC Arg	1152
	370					375					380					
ATC Ile	TGC Cys	GCC Ala	TGC Cys	TTA Leu	TGG Trp	ATG Met	ATG Met	CTG Leu	CTG Leu	ATA Ile	GCT Ala	CAA Gln	GCT Ala	GAG Glu	GCC Ala	1200
	385				390					395				400		
GCC Ala	TTA Leu	GAG Glu	AAC Asn	CTG Leu	GTG Val	GTC Val	CTC Leu	AAT Asn	GCG Ala	GCG Ala	GCC Ala	GTG Val	GCC Ala	GGG Gly	GCG Ala	1248
			405					410						415		
CAT His	GGC Gly	ACT Thr	CTT Ser	TCC Ser	TTC Phe	CTT Leu	GTG Val	TTC Phe	TTC Phe	TGT Cys	GCT Ala	GCC Ala	TGG Trp	TAC Tyr	ATC Ile	1296
			420					425					430			
AAG Lys	GGC Gly	AGG Arg	CTG Leu	GTC Val	CCT Pro	GGT Gly	GCG Ala	GCA Ala	TAC Tyr	GCC Ala	TTC Phe	TAT Tyr	GGC Gly	GTG Val	TGG Trp	1344
		435				440						445				
CCG Pro	CTG Leu	CTC Leu	CTG Leu	CTT Leu	CTG Leu	CTG Leu	GCC Ala	TTA Leu	CCA Pro	CCA Pro	CGA Arg	GCT Ala	TAT Tyr	GCC Ala	TAGTAA	1395
	450					455					460					

(2) INFORMATION FOR SEQ ID NO: 46:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 463 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 46:

Met	Val	Ala	Gly	Ala	His	Trp	Gly	Val	Leu	Ala	Gly	Leu	Ala	Tyr	Tyr	
1				5					10					15		
Ser	Met	Val	Gly	Asn	Trp	Ala	Lys	Val	Leu	Val	Val	Met	Leu	Leu	Phe	
			20					25					30			
Ala	Gly	Val	Asp	Gly	His	Thr	Arg	Val	Ser	Gly	Gly	Ala	Ala	Ala	Ser	
		35					40						45			
Asp	Thr	Arg	Gly	Leu	Val	Ser	Leu	Phe	Ser	Pro	Gly	Ser	Ala	Gln	Lys	
	50					55					60					
Ile	Gln	Leu	Val	Asn	Thr	Asn	Gly	Ser	Trp	His	Ile	Asn	Arg	Thr	Ala	
65				70						75				80		
Leu	Asn	Cys	Asn	Asp	Ser	Leu	Gln	Thr	Gly	Phe	Phe	Ala	Ala	Leu	Phe	
			85						90					95		
Tyr	Lys	His	Lys	Phe	Asn	Ser	Ser	Gly	Cys	Pro	Glu	Arg	Leu	Ala	Ser	

-continued

100					105					110					
Cys	Arg	Ser	Ile	Asp	Lys	Phe	Ala	Gln	Gly	Trp	Gly	Pro	Leu	Thr	Tyr
		115					120					125			
Thr	Glu	Pro	Asn	Ser	Ser	Asp	Gln	Arg	Pro	Tyr	Cys	Trp	His	Tyr	Ala
	130					135					140				
Pro	Arg	Pro	Cys	Gly	Ile	Val	Pro	Ala	Ser	Gln	Val	Cys	Gly	Pro	Val
145					150					155					160
Tyr	Cys	Phe	Thr	Pro	Ser	Pro	Val	Val	Val	Gly	Thr	Thr	Asp	Arg	Phe
				165						170				175	
Gly	Val	Pro	Thr	Tyr	Asn	Trp	Gly	Ala	Asn	Asp	Ser	Asp	Val	Leu	Ile
			180					185					190		
Leu	Asn	Asn	Thr	Arg	Pro	Pro	Arg	Gly	Asn	Trp	Phe	Gly	Cys	Thr	Trp
	195						200					205			
Met	Asn	Gly	Thr	Gly	Phe	Thr	Lys	Thr	Cys	Gly	Gly	Pro	Pro	Cys	Asn
	210						215				220				
Ile	Gly	Gly	Ala	Gly	Asn	Asn	Thr	Leu	Thr	Cys	Pro	Thr	Asp	Cys	Phe
225					230					235					240
Arg	Lys	His	Pro	Glu	Ala	Thr	Tyr	Ala	Arg	Cys	Gly	Ser	Gly	Pro	Trp
				245					250					255	
Leu	Thr	Pro	Arg	Cys	Met	Val	His	Tyr	Pro	Tyr	Arg	Leu	Trp	His	Tyr
			260						265				270		
Pro	Cys	Thr	Val	Asn	Phe	Thr	Ile	Phe	Lys	Val	Arg	Met	Tyr	Val	Gly
		275					280					285			
Gly	Val	Glu	His	Arg	Phe	Glu	Ala	Ala	Cys	Asn	Trp	Thr	Arg	Gly	Glu
	290					295					300				
Arg	Cys	Asp	Leu	Glu	Asp	Arg	Asp	Arg	Ser	Glu	Leu	Ser	Pro	Leu	Leu
305					310					315					320
Leu	Ser	Thr	Thr	Glu	Trp	Gln	Ile	Leu	Pro	Cys	Ser	Phe	Thr	Thr	Leu
				325					330					335	
Pro	Ala	Leu	Ser	Thr	Gly	Leu	Ile	His	Leu	His	Gln	Asn	Ile	Val	Asp
		340						345					350		
Val	Gln	Tyr	Leu	Tyr	Gly	Val	Gly	Ser	Ala	Val	Val	Ser	Leu	Val	Ile
		355					360					365			
Lys	Trp	Glu	Tyr	Val	Leu	Leu	Leu	Phe	Leu	Leu	Leu	Ala	Asp	Ala	Arg
	370					375						380			
Ile	Cys	Ala	Cys	Leu	Trp	Met	Met	Leu	Leu	Ile	Ala	Gln	Ala	Glu	Ala
385					390					395					400
Ala	Leu	Glu	Asn	Leu	Val	Val	Leu	Asn	Ala	Ala	Ala	Val	Ala	Gly	Ala
			405					410						415	
His	Gly	Thr	Leu	Ser	Phe	Leu	Val	Phe	Phe	Cys	Ala	Ala	Trp	Tyr	Ile
			420					425					430		
Lys	Gly	Arg	Leu	Val	Pro	Gly	Ala	Ala	Tyr	Ala	Phe	Tyr	Gly	Val	Trp
	435						440					445			
Pro	Leu	Leu	Leu	Leu	Leu	Leu	Ala	Leu	Pro	Pro	Arg	Ala	Tyr	Ala	
	450					455					460				

## (2) INFORMATION FOR SEQ ID NO: 47:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 2082 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

-continued

---

```

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:
      (A) NAME/KEY: CDS
      (B) LOCATION: 1..2079

(ix) FEATURE:
      (A) NAME/KEY: mat_peptide
      (B) LOCATION: 1..2076

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 47:

AAT TTG GGT AAG GTC ATC GAT ACC CTT ACA TGC GGC TTC GCC GAC CTC      48
Asn Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu
  1             5             10             15

GTG GGG TAC ATT CCG CTC GTC GGC GCC CCC CTA GGG GGC GCT GCC AGG      96
Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg
  20             25             30

GCC CTG GCG CAT GGC GTC CGG GTT CTG GAG GAC GGC GTG AAC TAT GCA      144
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala
  35             40             45

ACA GGG AAT TTG CCC GGT TGC TCT TTC TCT ATC TTC CTC TTG GCT TTG      192
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu
  50             55             60

CTG TCC TGT CTG ACC GTT CCA GCT TCC GCT TAT GAA GTG CGC AAC GTG      240
Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Glu Val Arg Asn Val
  65             70             75             80

TCC GGG ATG TAC CAT GTC ACG AAC GAC TGC TCC AAC TCA AGC ATT GTG      288
Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val
  85             90             95

TAT GAG GCA GCG GAC ATG ATC ATG CAC ACC CCC GGG TGC GTG CCC TGC      336
Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys
  100            105            110

GTT CGG GAG AAC AAC TCT TCC CGC TGC TGG GTA GCG CTC ACC CCC ACG      384
Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr
  115            120            125

CTC GCA GCT AGG AAC GCC AGC GTC CCC ACC ACG ACA ATA CGA CGC CAC      432
Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His
  130            135            140

GTC GAT TTG CTC GTT GGG GCG GCT GCT TTC TGT TCC GCT ATG TAC GTG      480
Val Asp Leu Leu Val Gly Ala Ala Ala Phe Cys Ser Ala Met Tyr Val
  145            150            155            160

GGG GAC CTC TGC GGA TCT GTC TTC CTC GTC TCC CAG CTG TTC ACC ATC      528
Gly Asp Leu Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Ile
  165            170            175

TCG CCT CGC CGG CAT GAG ACG GTG CAG GAC TGC AAT TGC TCA ATC TAT      576
Ser Pro Arg Arg His Glu Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr
  180            185            190

CCC GGC CAC ATA ACG GGT CAC CGT ATG GCT TGG GAT ATG ATG ATG AAC      624
Pro Gly His Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn
  195            200            205

TGG TCG CCT ACA ACG GCC CTG GTG GTA TCG CAG CTG CTC CGG ATC CCA      672
Trp Ser Pro Thr Thr Ala Leu Val Val Ser Gln Leu Leu Arg Ile Pro
  210            215            220

CAA GCT GTC GTG GAC ATG GTG GCG GGG GCC CAT TGG GGA GTC CTG GCG      720
Gln Ala Val Val Asp Met Val Ala Gly Ala His Trp Gly Val Leu Ala
  225            230            235            240

GGC CTC GCC TAC TAT TCC ATG GTG GGG AAC TGG GCT AAG GTT TTG GTT      768
Gly Leu Ala Tyr Tyr Ser Met Val Gly Asn Trp Ala Lys Val Leu Val
  245            250            255

GTG ATG CTA CTC TTT GCC GGC GTC GAC GGG CAT ACC CGC GTG TCA GGA      816

```



-continued

CAG AAC ATC GTG GAC GTG CAA TAC CTG TAC GGT GTA GGG TCG GCG GTT	1776
Gln Asn Ile Val Asp Val Gln Tyr Leu Tyr Gly Val Gly Ser Ala Val	
580 585 590	
GTC TCC CTT GTC ATC AAA TGG GAG TAT GTC CTG TTG CTC TTC CTT CTC	1824
Val Ser Leu Val Ile Lys Trp Glu Tyr Val Leu Leu Leu Phe Leu Leu	
595 600 605	
CTG GCA GAC GCG CGC ATC TGC GCC TGC TTA TGG ATG ATG CTG CTG ATA	1872
Leu Ala Asp Ala Arg Ile Cys Ala Cys Leu Trp Met Met Leu Leu Ile	
610 615 620	
GCT CAA GCT GAG GCC GCC TTA GAG AAC CTG GTG GTC CTC AAT GCG GCG	1920
Ala Gln Ala Glu Ala Ala Leu Glu Asn Leu Val Val Leu Asn Ala Ala	
625 630 635 640	
GCC GTG GCC GGG GCG CAT GGC ACT CTT TCC TTC CTT GTG TTC TTC TGT	1968
Ala Val Ala Gly Ala His Gly Thr Leu Ser Phe Leu Val Phe Phe Cys	
645 650 655	
GCT GCC TGG TAC ATC AAG GGC AGG CTG GTC CCT GGT GCG GCA TAC GCC	2016
Ala Ala Trp Tyr Ile Lys Gly Arg Leu Val Pro Gly Ala Ala Tyr Ala	
660 665 670	
TTC TAT GGC GTG TGG CCG CTG CTC CTG CTT CTG CTG GCC TTA CCA CCA	2064
Phe Tyr Gly Val Trp Pro Leu Leu Leu Leu Ala Leu Pro Pro	
675 680 685	
CGA GCT TAT GCC TAGTAA	2082
Arg Ala Tyr Ala	
690	

## (2) INFORMATION FOR SEQ ID NO: 48:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 692 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 48:

Asn Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu	1 5 10 15
Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg	20 25 30
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala	35 40 45
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu	50 55 60
Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Glu Val Arg Asn Val	65 70 75 80
Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val	85 90 95
Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys	100 105 110
Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr	115 120 125
Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His	130 135 140
Val Asp Leu Leu Val Gly Ala Ala Ala Phe Cys Ser Ala Met Tyr Val	145 150 155 160
Gly Asp Leu Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Ile	165 170 175
Ser Pro Arg Arg His Glu Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr	180 185 190

-continued

---

Pro Gly His Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn  
 195 200 205

Trp Ser Pro Thr Thr Ala Leu Val Val Ser Gln Leu Leu Arg Ile Pro  
 210 215 220

Gln Ala Val Val Asp Met Val Ala Gly Ala His Trp Gly Val Leu Ala  
 225 230 235 240

Gly Leu Ala Tyr Tyr Ser Met Val Gly Asn Trp Ala Lys Val Leu Val  
 245 250 255

Val Met Leu Leu Phe Ala Gly Val Asp Gly His Thr Arg Val Ser Gly  
 260 265 270

Gly Ala Ala Ala Ser Asp Thr Arg Gly Leu Val Ser Leu Phe Ser Pro  
 275 280 285

Gly Ser Ala Gln Lys Ile Gln Leu Val Asn Thr Asn Gly Ser Trp His  
 290 295 300

Ile Asn Arg Thr Ala Leu Asn Cys Asn Asp Ser Leu Gln Thr Gly Phe  
 305 310 315 320

Phe Ala Ala Leu Phe Tyr Lys His Lys Phe Asn Ser Ser Gly Cys Pro  
 325 330 335

Glu Arg Leu Ala Ser Cys Arg Ser Ile Asp Lys Phe Ala Gln Gly Trp  
 340 345 350

Gly Pro Leu Thr Tyr Thr Glu Pro Asn Ser Ser Asp Gln Arg Pro Tyr  
 355 360 365

Cys Trp His Tyr Ala Pro Arg Pro Cys Gly Ile Val Pro Ala Ser Gln  
 370 375 380

Val Cys Gly Pro Val Tyr Cys Phe Thr Pro Ser Pro Val Val Val Gly  
 385 390 395 400

Thr Thr Asp Arg Phe Gly Val Pro Thr Tyr Asn Trp Gly Ala Asn Asp  
 405 410 415

Ser Asp Val Leu Ile Leu Asn Asn Thr Arg Pro Pro Arg Gly Asn Trp  
 420 425 430

Phe Gly Cys Thr Trp Met Asn Gly Thr Gly Phe Thr Lys Thr Cys Gly  
 435 440 445

Gly Pro Pro Cys Asn Ile Gly Gly Ala Gly Asn Asn Thr Leu Thr Cys  
 450 455 460

Pro Thr Asp Cys Phe Arg Lys His Pro Glu Ala Thr Tyr Ala Arg Cys  
 465 470 475 480

Gly Ser Gly Pro Trp Leu Thr Pro Arg Cys Met Val His Tyr Pro Tyr  
 485 490 495

Arg Leu Trp His Tyr Pro Cys Thr Val Asn Phe Thr Ile Phe Lys Val  
 500 505 510

Arg Met Tyr Val Gly Gly Val Glu His Arg Phe Glu Ala Ala Cys Asn  
 515 520 525

Trp Thr Arg Gly Glu Arg Cys Asp Leu Glu Asp Arg Asp Arg Ser Glu  
 530 535 540

Leu Ser Pro Leu Leu Leu Ser Thr Thr Glu Trp Gln Ile Leu Pro Cys  
 545 550 555 560

Ser Phe Thr Thr Leu Pro Ala Leu Ser Thr Gly Leu Ile His Leu His  
 565 570 575

Gln Asn Ile Val Asp Val Gln Tyr Leu Tyr Gly Val Gly Ser Ala Val  
 580 585 590

Val Ser Leu Val Ile Lys Trp Glu Tyr Val Leu Leu Leu Phe Leu Leu  
 595 600 605

-continued

Leu Ala Asp Ala Arg Ile Cys Ala Cys Leu Trp Met Met Leu Leu Ile  
 610 615 620

Ala Gln Ala Glu Ala Ala Leu Glu Asn Leu Val Val Leu Asn Ala Ala  
 625 630 635 640

Ala Val Ala Gly Ala His Gly Thr Leu Ser Phe Leu Val Phe Phe Cys  
 645 650 655

Ala Ala Trp Tyr Ile Lys Gly Arg Leu Val Pro Gly Ala Ala Tyr Ala  
 660 665 670

Phe Tyr Gly Val Trp Pro Leu Leu Leu Leu Leu Ala Leu Pro Pro  
 675 680 685

Arg Ala Tyr Ala  
 690

## (2) INFORMATION FOR SEQ ID NO: 49:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 2433 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 1..2430

(ix) FEATURE:  
 (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 1..2427

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 49:

ATG AGC ACG AAT CCT AAA CCT CAA AGA AAA ACC AAA CGT AAC ACC AAC	48
Met Ser Thr Asn Pro Lys Pro Gln Arg Lys Thr Lys Arg Asn Thr Asn	
1 5 10 15	
CGC CGC CCA CAG GAC GTC AAG TTC CCG GGC GGT GGT CAG ATC GTT GGT	96
Arg Arg Pro Gln Asp Val Lys Phe Pro Gly Gly Gly Gln Ile Val Gly	
20 25 30	
GGA GTT TAC CTG TTG CCG CGC AGG GGC CCC AGG TTG GGT GTG CGC GCG	144
Gly Val Tyr Leu Leu Pro Arg Arg Gly Pro Arg Leu Gly Val Arg Ala	
35 40 45	
ACT AGG AAG ACT TCC GAG CGG TCG CAA CCT CGT GGG AGG CGA CAA CCT	192
Thr Arg Lys Thr Ser Glu Arg Ser Gln Pro Arg Gly Arg Arg Gln Pro	
50 55 60	
ATC CCC AAG GCT CGC CGA CCC GAG GGT AGG GCC TGG GCT CAG CCC GGG	240
Ile Pro Lys Ala Arg Arg Pro Glu Gly Arg Ala Trp Ala Gln Pro Gly	
65 70 75 80	
TAC CCT TGG CCC CTC TAT GGC AAT GAG GGC ATG GGG TGG GCA GGA TGG	288
Tyr Pro Trp Pro Leu Tyr Gly Asn Glu Gly Met Gly Trp Ala Gly Trp	
85 90 95	
CTC CTG TCA CCC CGC GGC TCT CGG CCT AGT TGG GGC CCT ACA GAC CCC	336
Leu Leu Ser Pro Arg Gly Ser Arg Pro Ser Trp Gly Pro Thr Asp Pro	
100 105 110	
CGG CGT AGG TCG CGT AAT TTG GGT AAG GTC ATC GAT ACC CTT ACA TGC	384
Arg Arg Arg Ser Arg Asn Leu Gly Lys Val Ile Asp Thr Leu Thr Cys	
115 120 125	
GGC TTC GCC GAC CTC GTG GGG TAC ATT CCG CTC GTC GGC GCC CCC CTA	432
Gly Phe Ala Asp Leu Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu	
130 135 140	



-continued

GGG Gly 145	GCC Ala 150	GCT Ala	AGC Arg	GCC Ala	CTG Leu	GCG Ala	CAT His	GGC Gly	GTC Val 155	CGG Arg	GTT Val	CTG Leu	GAG Glu	GAC Asp 160	480	
GGC Gly 165	GTG Val	AAC Asn	TAT Tyr	GCA Ala 165	ACA Thr	GGG Gly	AAT Asn	TTG Leu	CCC Pro 170	GGT Gly	TGC Cys	TCT Ser	TTC Phe	TCT Ser 175	528	
TTC Phe	CTC Leu	TTG Leu	GCT Ala 180	TTG Leu	CTG Leu	TCC Ser	TGT Cys 185	CTG Leu	ACC Thr	GTT Val	CCA Pro	GCT Ala	TCC Ser 190	GCT Ala	TAT Tyr	576
GAA Glu	GTG Val	CGC Arg	AAC Asn 195	GTG Val	TCC Ser	GGG Gly	ATG Met 200	TAC Tyr	CAT His	GTC Val	ACG Thr	AAC Asn 205	GAC Asp	TGC Cys	TCC Ser	624
AAC Asn 210	TCA Ser	AGC Ser	ATT Ile	GTG Val	TAT Tyr	GAG Glu 215	GCA Ala	GCG Ala	GAC Asp	ATG Met 220	ATC Ile	ATG Met	CAC His	ACC Thr	CCC Pro	672
GGG Gly 225	TGC Cys	GTG Val	CCC Pro	TGC Cys 230	GTT Val	CGG Arg	GAG Glu	AAC Asn	AAC Asn	TCT Ser 235	TCC Ser	CGC Arg	TGC Cys	TGG Trp 240	GTA Val	720
GCG Ala	CTC Leu	ACC Thr	CCC Pro	ACG Thr 245	CTC Leu	GCA Ala	GCT Ala	AGG Arg	AAC Asn 250	GCC Ala	AGC Ser	GTC Val	CCC Pro	ACC Thr 255	ACG Thr	768
ACA Thr	ATA Ile	CGA Arg	CGC Arg	CAC His 260	GTC Val	GAT Asp	TTG Leu	CTC Leu	GTT Val	GGG Gly 265	GCG Ala	GCT Ala	GCT Ala	TTC Ala 270	TGT Phe Cys	816
TCC Ser	GCT Ala	ATG Met	TAC Tyr	GTG Val	GGG Gly	GAC Asp 280	CTC Leu	TGC Cys	GGA Gly	TCT Ser	GTC Val	TTC Phe	CTC Leu	GTC Val	TCC Ser	864
CAG Gln	CTG Leu	TTC Phe	ACC Thr	ATC Ile	TCG Ser	CCT Pro	CGC Arg	CGG Arg	CAT His	GAG Glu	ACG Thr	GTG Val	CAG Gln	GAC Asp	TGC Cys	912
AAT Asn 305	TGC Cys	TCA Ser	ATC Ile	TAT Tyr	CCC Pro 310	GGC Gly	CAC His	ATA Ile	ACG Thr	GGT Gly 315	CAC His	CGT Arg	ATG Met	GCT Ala	TGG Trp 320	960
GAT Asp	ATG Met	ATG Met	ATG Met	AAC Asn 325	TGG Trp	TCG Ser	CCT Pro	ACA Thr	ACG Thr	GCC Ala 330	CTG Leu	GTG Val	GTA Val	TCG Ser 335	CAG Gln	1008
CTG Leu	CTC Leu	CGG Arg	ATC Ile	CCA Pro 340	CAA Gln	GCT Ala	GTC Val	GTG Val	GAC Asp	ATG Met	GTG Val	GCG Ala	GGG Gly 350	GCC Ala	CAT His	1056
TGG Trp	GGA Gly	GTC Val	CTG Leu	GCG Ala	GGC Gly	CTC Leu	GCC Ala 360	TAC Tyr	TAT Tyr	TCC Ser	ATG Met	GTG Val	GGG Gly 365	AAC Asn	TGG Trp	1104
GCT Ala	AAG Lys	GTT Val	TTG Leu	GTT Val	GTG Val	ATG Met	CTA Leu	CTC Leu	TTT Phe	GCC Ala 380	GGC Gly	GTC Val	GAC Asp	GGG Gly	CAT His	1152
ACC Thr 385	CGC Arg	GTG Val	TCA Ser	GGA Gly	GGG Ala 390	GCA Ala	GCA Ala	GCC Ala	TCC Ser	GAT Asp 395	ACC Thr	AGG Arg	GGC Gly	CTT Leu	GTG Val 400	1200
TCC Ser	CTC Leu	TTT Phe	AGC Ser	CCC Pro 405	GGG Gly	TCG Ser	GCT Ala	CAG Gln	AAA Lys 410	ATC Ile	CAG Gln	CTC Leu	GTA Val	AAC Asn 415	ACC Thr	1248
AAC Asn 420	GGC Gly	AGT Ser	TGG Trp	CAC His 420	ATC Ile	AAC Asn	AGG Arg	ACT Thr	GCC Ala 425	CTG Leu	AAC Asn	TGC Cys	AAC Asn 430	GAC Asp	TCC Ser	1296
CTC Leu	CAA Gln	ACA Thr	GGG Gly	TTC Phe	TTT Phe	GCC Ala	GCA Ala	CTA Leu	TTC Phe	TAC Tyr	AAA Lys	CAC His	AAA Lys	TTC Phe	AAC Asn	1344
TCG Ser	TCT Ser	GGA Gly	TGC Cys	CCA Pro	GAG Glu	CGC Arg	TTG Leu	GCC Ala	AGC Ser	TGT Cys	CGC Arg	TCC Ser	ATC Ile	GAC Asp	AAG Lys	1392

-continued

TTC Phe 465	GCT Ala Gln	CAG Gln Gly	GGG Gly Trp	TGG Trp 470	GGT Gly Pro	CCC Pro Leu	CTC Thr Tyr	ACT Thr Tyr	TAC Thr 475	ACT Glu Pro	GAG Pro Arg	CCT Pro Cys	AAC Asn Ser	AGC Ser 480	TCG Ser 480	1440
GAC Asp	CAG Gln	AGG Arg	CCC Pro	TAC Tyr 485	TGC Cys	TGG Trp	CAC His	TAC Tyr	GCG Ala	CCT Pro	CGA Arg	CCG Pro	TGT Cys	GGT Gly	ATT Ile 495	1488
GTA Val	CCC Pro	GCG Ala	TCT Ser	CAG Gln	GTG Val	TGC Cys	GGT Gly	CCA Pro	GTG Val	TAT Tyr	TGC Cys	TTC Phe	ACC Thr	CCG Pro	AGC Ser	1536
CCT Pro	GTT Val	GTG Val	GTG Val	GGG Gly	ACG Thr	ACC Thr	GAT Asp	CGG Arg	TTT Phe	GGT Gly	GTC Val	CCC Pro	ACG Thr	TAT Tyr	AAC Asn	1584
TGG Trp	GGG Gly	GCG Ala	AAC Asn	GAC Asp	TCG Ser	GAT Asp	GTG Val	CTG Leu	ATT Ile	CTC Leu	AAC Asn	AAC Asn	ACG Thr	CGG Arg	CCG Pro	1632
CCG Pro	CGA Arg	GGC Gly	AAC Asn	TGG Trp	TTC Phe	GGC Gly	TGT Cys	ACA Thr	TGG Trp	ATG Met	AAT Asn	GGC Gly	ACT Thr	GGG Gly	TTC Phe	1680
ACC Thr	AAG Lys	ACG Thr	TGT Cys	GGG Gly	CCC Pro	CCG Pro	TGC Cys	AAC Asn	ATC Ile	GGG Gly	GGG Gly	GCC Ala	GGC Gly	AAC Asn	1728	
AAC Asn	ACC Thr	TTG Leu	ACC Thr	TGC Cys	CCC Pro	ACT Thr	GAC Asp	TGT Cys	TTT Phe	CGG Arg	AAG Lys	CAC His	CCC Pro	GAG Glu	GCC Ala	1776
ACC Thr	TAC Tyr	GCC Ala	AGA Arg	TGC Cys	GGT Gly	TCT Ser	GGG Gly	CCC Pro	TGG Trp	CTG Leu	ACA Thr	CCT Pro	AGG Arg	TGT Cys	ATG Met	1824
GTT Val	CAT His	TAC Tyr	CCA Pro	TAT Tyr	AGG Arg	CTC Leu	TGG Trp	CAC His	TAC Tyr	CCC Pro	TGC Cys	ACT Thr	GTC Val	AAC Asn	TTC Phe	1872
ACC Thr	ATC Ile	TTC Phe	AAG Lys	GTT Val	AGG Arg	ATG Met	TAC Tyr	GTG Val	GGG Gly	GGC Gly	GTG Val	GAG Glu	CAC His	AGG Arg	TTC Phe	1920
GAA Glu	GCC Ala	GCA Ala	TGC Cys	AAT Asn	TGG Trp	ACT Thr	CGA Arg	GGA Gly	GAG Glu	CGT Arg	TGT Cys	GAC Asp	TTG Leu	GAG Glu	GAC Asp	1968
AGG Arg	GAT Asp	AGA Arg	TCA Ser	GAG Glu	CTT Leu	AGC Ser	CCG Pro	CTG Leu	CTG Leu	TCT Leu	ACA Thr	ACA Thr	GAG Glu	TGG Trp	2016	
CAG Gln	ATA Ile	CTG Leu	CCC Pro	TGT Cys	TCC Ser	TTC Phe	ACC Thr	ACC Thr	CTG Leu	CCG Pro	GCC Ala	CTA Leu	TCC Ser	ACC Thr	GGC Gly	2064
CTG Leu	ATC Ile	CAC His	CTC Leu	CAT His	CAG Gln	AAC Asn	ATC Ile	GTG Val	GAC Asp	GTG Val	CAA Gln	TAC Tyr	CTG Leu	TAC Tyr	GGT Gly	2112
GTA Val	GGG Gly	TCG Ser	GCG Ala	GTT Val	GTC Val	TCC Ser	CTT Leu	GTC Val	ATC Ile	AAA Lys	TGG Trp	GAG Glu	TAT Tyr	GTC Val	CTG Leu	2160
TTG Leu	CTC Leu	TTC Phe	CTT Leu	CTC Leu	CTG Leu	GCA Ala	GAC Asp	GCG Ala	CGC Arg	ATC Ile	TGC Cys	GCC Ala	TGC Cys	TTA Leu	TGG Trp	2208
ATG Met	ATG Met	CTG Leu	CTG Leu	ATA Ile	GCT Ala	CAA Gln	GCT Ala	GAG Gly	GCC Ala	GCC Ala	TTA Leu	GAG Glu	AAC Asn	CTG Leu	GTG Val	2256
GTC Val	CTC Leu	AAT Asn	GCG Ala	GCG Ala	GCC Ala	GTG Val	GCC Ala	GGG Gly	GCG Ala	CAT His	GGC Gly	ACT Thr	CTT Leu	TCC Ser	TTC Phe	2304
CTT Leu	GTG Val	TTC Phe	TTC Phe	TGT Cys	GCT Ala	GCC Ala	TGG Trp	TAC Tyr	ATC Ile	AAG Lys	GGC Gly	AGG Arg	CTG Leu	GTC Val	CCT Pro	2352

-continued

770	775	780	
GGT GCG GCA TAC GCC TTC TAT GGC GTG TGG CCG CTG CTC CTG CTT CTG			2400
Gly Ala Ala Tyr Ala Phe Tyr Gly Val Trp Pro Leu Leu Leu Leu Leu			
785	790	795	800
CTG GCC TTA CCA CCA CGA GCT TAT GCC TAGTAA			2433
Leu Ala Leu Pro Pro Arg Ala Tyr Ala			
	805	810	

## (2) INFORMATION FOR SEQ ID NO: 50:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 809 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 50:

Met	Ser	Thr	Asn	Pro	Lys	Pro	Gln	Arg	Lys	Thr	Lys	Arg	Asn	Thr	Asn
1				5					10					15	
Arg	Arg	Pro	Gln	Asp	Val	Lys	Phe	Pro	Gly	Gly	Gly	Gln	Ile	Val	Gly
			20					25					30		
Gly	Val	Tyr	Leu	Leu	Pro	Arg	Arg	Gly	Pro	Arg	Leu	Gly	Val	Arg	Ala
		35					40					45			
Thr	Arg	Lys	Thr	Ser	Glu	Arg	Ser	Gln	Pro	Arg	Gly	Arg	Arg	Gln	Pro
	50					55					60				
Ile	Pro	Lys	Ala	Arg	Arg	Pro	Glu	Gly	Arg	Ala	Trp	Ala	Gln	Pro	Gly
	65				70					75					80
Tyr	Pro	Trp	Pro	Leu	Tyr	Gly	Asn	Glu	Gly	Met	Gly	Trp	Ala	Gly	Trp
				85					90					95	
Leu	Leu	Ser	Pro	Arg	Gly	Ser	Arg	Pro	Ser	Trp	Gly	Pro	Thr	Asp	Pro
			100					105					110		
Arg	Arg	Arg	Ser	Arg	Asn	Leu	Gly	Lys	Val	Ile	Asp	Thr	Leu	Thr	Cys
		115					120					125			
Gly	Phe	Ala	Asp	Leu	Val	Gly	Tyr	Ile	Pro	Leu	Val	Gly	Ala	Pro	Leu
	130					135					140				
Gly	Gly	Ala	Ala	Arg	Ala	Leu	Ala	His	Gly	Val	Arg	Val	Leu	Glu	Asp
	145				150					155					160
Gly	Val	Asn	Tyr	Ala	Thr	Gly	Asn	Leu	Pro	Gly	Cys	Ser	Phe	Ser	Ile
				165					170					175	
Phe	Leu	Leu	Ala	Leu	Leu	Ser	Cys	Leu	Thr	Val	Pro	Ala	Ser	Ala	Tyr
			180					185						190	
Glu	Val	Arg	Asn	Val	Ser	Gly	Met	Tyr	His	Val	Thr	Asn	Asp	Cys	Ser
	195						200					205			
Asn	Ser	Ser	Ile	Val	Tyr	Glu	Ala	Ala	Asp	Met	Ile	Met	His	Thr	Pro
	210					215					220				
Gly	Cys	Val	Pro	Cys	Val	Arg	Glu	Asn	Asn	Ser	Ser	Arg	Cys	Trp	Val
	225				230					235					240
Ala	Leu	Thr	Pro	Thr	Leu	Ala	Ala	Arg	Asn	Ala	Ser	Val	Pro	Thr	Thr
				245					250					255	
Thr	Ile	Arg	Arg	His	Val	Asp	Leu	Leu	Val	Gly	Ala	Ala	Ala	Phe	Cys
			260					265						270	
Ser	Ala	Met	Tyr	Val	Gly	Asp	Leu	Cys	Gly	Ser	Val	Phe	Leu	Val	Ser
		275					280					285			
Gln	Leu	Phe	Thr	Ile	Ser	Pro	Arg	Arg	His	Glu	Thr	Val	Gln	Asp	Cys
	290					295					300				

-continued

Asn	Cys	Ser	Ile	Tyr	Pro	Gly	His	Ile	Thr	Gly	His	Arg	Met	Ala	Trp	305	310	315	320
Asp	Met	Met	Met	Asn	Trp	Ser	Pro	Thr	Thr	Ala	Leu	Val	Val	Ser	Gln	325	330	335	
Leu	Leu	Arg	Ile	Pro	Gln	Ala	Val	Val	Asp	Met	Val	Ala	Gly	Ala	His	340	345	350	
Trp	Gly	Val	Leu	Ala	Gly	Leu	Ala	Tyr	Tyr	Ser	Met	Val	Gly	Asn	Trp	355	360	365	
Ala	Lys	Val	Leu	Val	Val	Met	Leu	Leu	Phe	Ala	Gly	Val	Asp	Gly	His	370	375	380	
Thr	Arg	Val	Ser	Gly	Gly	Ala	Ala	Ala	Ser	Asp	Thr	Arg	Gly	Leu	Val	385	390	395	400
Ser	Leu	Phe	Ser	Pro	Gly	Ser	Ala	Gln	Lys	Ile	Gln	Leu	Val	Asn	Thr	405	410	415	
Asn	Gly	Ser	Trp	His	Ile	Asn	Arg	Thr	Ala	Leu	Asn	Cys	Asn	Asp	Ser	420	425	430	
Leu	Gln	Thr	Gly	Phe	Phe	Ala	Ala	Leu	Phe	Tyr	Lys	His	Lys	Phe	Asn	435	440	445	
Ser	Ser	Gly	Cys	Pro	Glu	Arg	Leu	Ala	Ser	Cys	Arg	Ser	Ile	Asp	Lys	450	455	460	
Phe	Ala	Gln	Gly	Trp	Gly	Pro	Leu	Thr	Tyr	Thr	Glu	Pro	Asn	Ser	Ser	465	470	475	480
Asp	Gln	Arg	Pro	Tyr	Cys	Trp	His	Tyr	Ala	Pro	Arg	Pro	Cys	Gly	Ile	485	490	495	
Val	Pro	Ala	Ser	Gln	Val	Cys	Gly	Pro	Val	Tyr	Cys	Phe	Thr	Pro	Ser	500	505	510	
Pro	Val	Val	Val	Gly	Thr	Thr	Asp	Arg	Phe	Gly	Val	Pro	Thr	Tyr	Asn	515	520	525	
Trp	Gly	Ala	Asn	Asp	Ser	Asp	Val	Leu	Ile	Leu	Asn	Asn	Thr	Arg	Pro	530	535	540	
Pro	Arg	Gly	Asn	Trp	Phe	Gly	Cys	Thr	Trp	Met	Asn	Gly	Thr	Gly	Phe	545	550	555	560
Thr	Lys	Thr	Cys	Gly	Gly	Pro	Pro	Cys	Asn	Ile	Gly	Gly	Ala	Gly	Asn	565	570	575	
Asn	Thr	Leu	Thr	Cys	Pro	Thr	Asp	Cys	Phe	Arg	Lys	His	Pro	Glu	Ala	580	585	590	
Thr	Tyr	Ala	Arg	Cys	Gly	Ser	Gly	Pro	Trp	Leu	Thr	Pro	Arg	Cys	Met	595	600	605	
Val	His	Tyr	Pro	Tyr	Arg	Leu	Trp	His	Tyr	Pro	Cys	Thr	Val	Asn	Phe	610	615	620	
Thr	Ile	Phe	Lys	Val	Arg	Met	Tyr	Val	Gly	Gly	Val	Glu	His	Arg	Phe	625	630	635	640
Glu	Ala	Ala	Cys	Asn	Trp	Thr	Arg	Gly	Glu	Arg	Cys	Asp	Leu	Glu	Asp	645	650	655	
Arg	Asp	Arg	Ser	Glu	Leu	Ser	Pro	Leu	Leu	Leu	Ser	Thr	Thr	Glu	Trp	660	665	670	
Gln	Ile	Leu	Pro	Cys	Ser	Phe	Thr	Thr	Leu	Pro	Ala	Leu	Ser	Thr	Gly	675	680	685	
Leu	Ile	His	Leu	His	Gln	Asn	Ile	Val	Asp	Val	Gln	Tyr	Leu	Tyr	Gly	690	695	700	
Val	Gly	Ser	Ala	Val	Val	Ser	Leu	Val	Ile	Lys	Trp	Glu	Tyr	Val	Leu	705	710	715	720
Leu	Leu	Phe	Leu	Leu	Leu	Ala	Asp	Ala	Arg	Ile	Cys	Ala	Cys	Leu	Trp				



-continued

---

Tyr Glu Val Arg Asn Val Ser Gly Ile Tyr His Val Thr Asn Asp Cys  
 1 5 10 15  
 Ser Asn Ser Ser Ile Val Tyr Glu Ala Ala Asp Met Ile Met His Thr  
 20 25 30  
 Pro Gly Cys Gly Lys  
 35

(2) INFORMATION FOR SEQ ID NO: 54:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 25 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:  
 (A) NAME/KEY: Modified-site  
 (B) LOCATION: 1..25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 54:

Gly Gly Thr Pro Thr Val Ala Thr Arg Asp Gly Lys Leu Pro Ala Thr  
 1 5 10 15  
 Gln Leu Arg Arg His Ile Asp Leu Leu  
 20 25

(2) INFORMATION FOR SEQ ID NO: 55:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 25 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:  
 (A) NAME/KEY: Modified-site  
 (B) LOCATION: 1..25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 55:

Gly Gly Thr Pro Thr Leu Ala Ala Arg Asp Ala Ser Val Pro Thr Thr  
 1 5 10 15  
 Thr Ile Arg Arg His Val Asp Leu Leu  
 20 25

(2) INFORMATION FOR SEQ ID NO: 56:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 56:

Leu Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Gln Val Arg Asn  
 1 5 10 15  
 Ser Thr Gly Leu  
 20

(2) INFORMATION FOR SEQ ID NO: 57:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids

-continued

(B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 57:

Gln Val Arg Asn Ser Thr Gly Leu Tyr His Val Thr Asn Asp Cys Pro  
 1                   5                           10                           15

Asn Ser Ser Ile  
           20

(2) INFORMATION FOR SEQ ID NO: 58:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 58:

Asn Asp Cys Pro Asn Ser Ser Ile Val Tyr Glu Ala His Asp Ala Ile  
 1                   5                           10                           15

Leu His Thr Pro  
           20

(2) INFORMATION FOR SEQ ID NO: 59:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 59:

Ser Asn Ser Ser Ile Val Tyr Glu Ala Ala Asp Met Ile Met His Thr  
 1                   5                           10                           15

Pro Gly Cys Val  
           20

(2) INFORMATION FOR SEQ ID NO: 60:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 19 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 60:

His Asp Ala Ile Leu His Thr Pro Gly Val Pro Cys Val Arg Glu Gly  
 1                   5                           10                           15

Asn Val Ser

(2) INFORMATION FOR SEQ ID NO: 61:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

-continued

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 61:

Cys Val Arg Glu Gly Asn Val Ser Arg Cys Trp Val Ala Met Thr Pro  
 1                   5                   10                   15  
 Thr Val Ala Thr  
                   20

(2) INFORMATION FOR SEQ ID NO: 62:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 62:

Ala Met Thr Pro Thr Val Ala Thr Arg Asp Gly Lys Leu Pro Ala Thr  
 1                   5                   10                   15  
 Gln Leu Arg Arg  
                   20

(2) INFORMATION FOR SEQ ID NO: 63:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 63:

Leu Pro Ala Thr Gln Leu Arg Arg His Ile Asp Leu Leu Val Gly Ser  
 1                   5                   10                   15  
 Ala Thr Leu Cys  
                   20

(2) INFORMATION FOR SEQ ID NO: 64:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 64:

Leu Val Gly Ser Ala Thr Leu Cys Ser Ala Leu Tyr Val Gly Asp Leu  
 1                   5                   10                   15  
 Cys Gly Ser Val  
                   20

(2) INFORMATION FOR SEQ ID NO: 65:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 65:



-continued

Gln Leu Phe Thr Phe Ser Pro Arg Arg His Trp Thr Thr Gln Gly Cys  
 1 5 10 15

Asn Cys Ser Ile  
 20

(2) INFORMATION FOR SEQ ID NO: 66:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 66:

Thr Gln Gly Cys Asn Cys Ser Ile Tyr Pro Gly His Ile Thr Gly His  
 1 5 10 15

Arg Met Ala Trp  
 20

(2) INFORMATION FOR SEQ ID NO: 67:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 67:

Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp Ser Pro  
 1 5 10 15

Thr Ala Ala Leu  
 20

(2) INFORMATION FOR SEQ ID NO: 68:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 68:

Asn Trp Ser Pro Thr Ala Ala Leu Val Met Ala Gln Leu Leu Arg Ile  
 1 5 10 15

Pro Gln Ala Ile  
 20

(2) INFORMATION FOR SEQ ID NO: 69:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 69:

Leu Leu Arg Ile Pro Gln Ala Ile Leu Asp Met Ile Ala Gly Ala His  
 1 5 10 15

-continued

Trp Gly Val Leu  
20

## (2) INFORMATION FOR SEQ ID NO: 70:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 70:

Ala Gly Ala His Trp Gly Val Leu Ala Gly Ile Ala Tyr Phe Ser Met  
 1 5 10 15

Val Gly Asn Met  
20

## (2) INFORMATION FOR SEQ ID NO: 71:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 71:

Val Val Leu Leu Leu Phe Ala Gly Val Asp Ala Glu Thr Ile Val Ser  
 1 5 10 15

Gly Gly Gln Ala  
20

## (2) INFORMATION FOR SEQ ID NO: 72:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 72:

Ser Gly Leu Val Ser Leu Phe Thr Pro Gly Ala Lys Gln Asn Ile Gln  
 1 5 10 15

Leu Ile Asn Thr  
20

## (2) INFORMATION FOR SEQ ID NO: 73:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 73:

Gln Asn Ile Gln Leu Ile Asn Thr Asn Gly Gln Trp His Ile Asn Ser  
 1 5 10 15

Thr Ala Leu Asn  
20

-continued

## (2) INFORMATION FOR SEQ ID NO: 74:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 21 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 74:

Leu Asn Cys Asn Glu Ser Leu Asn Thr Gly Trp Trp Leu Ala Gly Leu  
 1                   5                           10                           15

Ile Tyr Gln His Lys  
           20

## (2) INFORMATION FOR SEQ ID NO: 75:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 75:

Ala Gly Leu Ile Tyr Gln His Lys Phe Asn Ser Ser Gly Cys Pro Glu  
 1                   5                           10                           15

Arg Leu Ala Ser  
           20

## (2) INFORMATION FOR SEQ ID NO: 76:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 76:

Gly Cys Pro Glu Arg Leu Ala Ser Cys Arg Pro Leu Thr Asp Phe Asp  
 1                   5                           10                           15

Gln Gly Trp Gly  
           20

## (2) INFORMATION FOR SEQ ID NO: 77:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 77:

Thr Asp Phe Asp Gln Gly Trp Gly Pro Ile Ser Tyr Ala Asn Gly Ser  
 1                   5                           10                           15

Gly Pro Asp Gln  
           20

## (2) INFORMATION FOR SEQ ID NO: 78:

-continued

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 78:

Ala Asn Gly Ser Gly Pro Asp Gln Arg Pro Tyr Cys Trp His Tyr Pro  
 1                   5                   10                   15

Pro Lys Pro Cys  
           20

(2) INFORMATION FOR SEQ ID NO: 79:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 79:

Trp His Tyr Pro Pro Lys Pro Cys Gly Ile Val Pro Ala Lys Ser Val  
 1                   5                   10                   15

Cys Gly Pro Val  
           20

(2) INFORMATION FOR SEQ ID NO: 80:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 80:

Ala Lys Ser Val Cys Gly Pro Val Tyr Cys Phe Thr Pro Ser Pro Val  
 1                   5                   10                   15

Val Val Gly Thr  
           20

(2) INFORMATION FOR SEQ ID NO: 81:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 81:

Pro Ser Pro Val Val Val Gly Thr Thr Asp Arg Ser Gly Ala Pro Thr  
 1                   5                   10                   15

Tyr Ser Trp Gly  
           20

(2) INFORMATION FOR SEQ ID NO: 82:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids

-continued

(B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 82:

Gly Ala Pro Thr Tyr Ser Trp Gly Glu Asn Asp Thr Asp Val Phe Val  
 1                   5                   10                   15

Leu Asn Asn Thr  
           20

(2) INFORMATION FOR SEQ ID NO: 83:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 83:

Gly Asn Trp Phe Gly Cys Thr Trp Met Asn Ser Thr Gly Phe Thr Lys  
 1                   5                   10                   15

Val Cys Gly Ala  
           20

(2) INFORMATION FOR SEQ ID NO: 84:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 84:

Gly Phe Thr Lys Val Cys Gly Ala Pro Pro Val Cys Ile Gly Gly Ala  
 1                   5                   10                   15

Gly Asn Asn Thr  
           20

(2) INFORMATION FOR SEQ ID NO: 85:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 19 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 85:

Ile Gly Gly Ala Gly Asn Asn Thr Leu His Cys Pro Thr Asp Cys Arg  
 1                   5                   10                   15

Lys His Pro

(2) INFORMATION FOR SEQ ID NO: 86:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

-continued

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 86:

Thr Asp Cys Phe Arg Lys His Pro Asp Ala Thr Tyr Ser Arg Cys Gly  
 1                   5                   10                   15  
 Ser Gly Pro Trp  
                   20

(2) INFORMATION FOR SEQ ID NO: 87:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 87:

Ser Arg Cys Gly Ser Gly Pro Trp Ile Thr Pro Arg Cys Leu Val Asp  
 1                   5                   10                   15  
 Tyr Pro Tyr Arg  
                   20

(2) INFORMATION FOR SEQ ID NO: 88:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 88:

Cys Leu Val Asp Tyr Pro Tyr Arg Leu Trp His Tyr Pro Cys Thr Ile  
 1                   5                   10                   15  
 Asn Tyr Thr Ile  
                   20

(2) INFORMATION FOR SEQ ID NO: 89:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 89:

Pro Cys Thr Ile Asn Tyr Thr Ile Phe Lys Ile Arg Met Tyr Val Gly  
 1                   5                   10                   15  
 Gly Val Glu His  
                   20

(2) INFORMATION FOR SEQ ID NO: 90:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 90:

-continued

Met Tyr Val Gly Gly Val Glu His Arg Leu Glu Ala Ala Cys Asn Trp  
 1 5 10 15

Thr Pro Gly Glu  
 20

## (2) INFORMATION FOR SEQ ID NO: 91:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 91:

Ala Cys Asn Trp Thr Pro Gly Glu Arg Cys Asp Leu Glu Asp Arg Asp  
 1 5 10 15

Arg Ser Glu Leu  
 20

## (2) INFORMATION FOR SEQ ID NO: 92:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 92:

Glu Asp Arg Asp Arg Ser Glu Leu Ser Pro Leu Leu Leu Thr Thr Thr  
 1 5 10 15

Gln Trp Gln Val  
 20

## (2) INFORMATION FOR SEQ ID NO: 93:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 9 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 93:

Tyr Gln Val Arg Asn Ser Thr Gly Leu  
 1 5

## (2) INFORMATION FOR SEQ ID NO: 94:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 29 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 94:

ACGTCCGTAC GTTCGAATTA ATTAATCGA

-continued

## (2) INFORMATION FOR SEQ ID NO: 95:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 60 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 95:

CCTCCGGACG TGCAGTAGCT CCCGTCTGTG GTAGTGGTGG TAGTGATTAT CAATTAATTG 60

## (2) INFORMATION FOR SEQ ID NO: 96:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 19 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 96:

GTTTAACCAC TGCATGATG 19

## (2) INFORMATION FOR SEQ ID NO: 97:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 97:

GTCCCATCGA GTGCGGCTAC 20

## (2) INFORMATION FOR SEQ ID NO: 98:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 45 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 98:

CGTGACATGG TACATTCCGG ACACTGGCG CACTTCATAA GCGGA 45

## (2) INFORMATION FOR SEQ ID NO: 99:



-continued

- 
- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 42 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iii) ANTI-SENSE: NO
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 99:  
 TGCCCTCATAC ACAATGGAGC TCTGGGACGA GTCGTTCGTG AC 42
- (2) INFORMATION FOR SEQ ID NO: 100:
- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 42 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iii) ANTI-SENSE: NO
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 100:  
 TACCCAGCAG CGGGAGCTCT GTTGCTCCCG AACGCAGGGC AC 42
- (2) INFORMATION FOR SEQ ID NO: 101:
- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 42 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iii) ANTI-SENSE: NO
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 101:  
 TGTCGTGGTG GGGACGGAGG CCTGCCTAGC TGCAGCGTG GG 42
- (2) INFORMATION FOR SEQ ID NO: 102:
- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 48 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iii) ANTI-SENSE: NO
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 102:  
 CGTTATGTGG CCCGGGTAGA TTGAGCACTG GCAGTCCTGC ACCGTCTC 48
- (2) INFORMATION FOR SEQ ID NO: 103:
- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 42 base pairs

-continued

---

(B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 103:  
 CAGGGCCGTT CTAGGCCTCC ACTGCATCAT CATATCCCAA GC 42

(2) INFORMATION FOR SEQ ID NO: 104:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 26 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 104:  
 CCGGAATGTA CCATGTCACG AACGAC 26

(2) INFORMATION FOR SEQ ID NO: 105:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 24 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 105:  
 GCTCCATTGT GTATGAGGCA GCGG 24

(2) INFORMATION FOR SEQ ID NO: 106:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 23 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 106:  
 GAGCTCCCGC TGCTGGGTAG CGC 23

(2) INFORMATION FOR SEQ ID NO: 107:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 25 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

-continued

---

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 107:

CCTCCGTCCC CACCACGACA ATACG 25

(2) INFORMATION FOR SEQ ID NO: 108:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 108:

CTACCCGGGC CACATAACGG GTCACCG 27

(2) INFORMATION FOR SEQ ID NO: 109:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 109:

GGAGGCCTAC AACGGCCCTG GTGG 24

(2) INFORMATION FOR SEQ ID NO: 110:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 110:

TTCTATCGAT TAAATAGAAT TC 22

(2) INFORMATION FOR SEQ ID NO: 111:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

-continued

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 111:

GCCATACGCT CACAGCCGAT CCC

23

We claim:

1. A recombinant HCV single or specifically oligomerized envelope viral protein selected from the group consisting of E1, E2 or E1/E2 viral proteins obtained from a method comprising the steps of
  - (a) optionally lysing host cells expressing said viral protein;
  - (b) optionally recovering said viral protein;
  - (c) cleaving disulphide bonds of said viral protein with a disulphide bond cleaving agent to form a cleaved protein;
  - (d) preventing disulphide bond reformation of said cleaved viral protein with at least one of an SH group blocking agent and low pH condition; and
  - (e) purifying the cleaved viral protein obtained in step (d) to produce a viral protein which is at least 80% purified.
2. A recombinant protein according to claim 1 wherein said method further comprises desalting said purified viral protein of step (e).
3. A recombinant protein according to any one of claims 1 or 2 wherein step (b) or step (e) further comprises a chromatographic recovery.
4. A recombinant protein according to claim 3 wherein said affinity chromatography comprises lectin-chromatography or immunoaffinity chromatography with at least one of an anti-E1 specific monoclonal antibody or an anti-E2 specific monoclonal antibody.
5. A recombinant protein according to any one of claims 1 or 2 wherein said SH group blocking agent is N-ethylmaleimide.
6. A recombinant protein according to any one of claims 1 or 2, wherein said purified protein is at least 95% pure.
7. A recombinant protein according to any one of claims 1 or 2, wherein said purified protein is at least 90% pure.
8. A recombinant protein according to claim 1 wherein step (b) further comprises an affinity chromatography.
9. A recombinant protein according to claim 1 wherein step (a) further comprises addition of an SH group blocking agent.
10. A recombinant protein according to claim 1 wherein said cleaving comprises partial cleaving conditions including addition of a detergent.
11. A recombinant protein according to claim 10 wherein said detergent comprises -N-Dodecyl-N,N-dimethylglycine.
12. A recombinant protein according to claim 11 wherein said detergent is present at a concentration of 1 to 10%.
13. A recombinant protein according to claim 11 wherein said detergent is present at a concentration of 3.5%.
14. A recombinant protein according to claim 1 wherein said disulphide bond cleaving agent is dithiothreitol.
15. A recombinant protein according to claim 14 wherein said dithiothreitol is present at a concentration of 0.1 to 50 mM.
16. A recombinant protein according to claim 15 wherein said dithiothreitol is present at a concentration of 0.1 to 20 mM.
17. A recombinant protein according to claim 15 wherein said dithiothreitol is present at a concentration of 0.5 to 10 mM.
18. A recombinant protein according to claim 1 wherein said purified protein is at least 97% pure.
19. A recombinant protein according to claim 1 wherein said purified protein is at least 98% pure.
20. A recombinant protein according to claim 1 wherein said purified protein is at least 99% pure.
21. A recombinant HCV single or specifically oligomerized envelope protein selected from the group consisting of E1, E2 or E1/E2 viral proteins, which is at least 80% pure.
22. A recombinant HCV single or specifically oligomerized envelope protein according to claim 21 which is at least 90% pure.
23. A recombinant HCV single or specifically oligomerized envelope protein according to claim 21 which is at least 95% pure.
24. A recombinant HCV single or specifically oligomerized envelope protein according to claim 21, which is at least 97% pure.
25. A recombinant HCV single or specifically oligomerized envelope protein according to claim 21, which is at least 98% pure.
26. A recombinant HCV single or specifically oligomerized envelope protein according to claim 21, which is at least 99% pure.

\* \* \* \* \*