



(86) Date de dépôt PCT/PCT Filing Date: 2005/05/04
(87) Date publication PCT/PCT Publication Date: 2005/11/10
(85) Entrée phase nationale/National Entry: 2006/10/20
(86) N° demande PCT/PCT Application No.: EP 2005/052041
(87) N° publication PCT/PCT Publication No.: 2005/105810
(30) Priorités/Priorities: 2004/05/04 (EP04101913.4);
2004/08/25 (US60/604,275)

(51) Cl.Int./Int.Cl. *C07D 498/04* (2006.01),
A61K 31/538 (2006.01), *C07D 231/00* (2006.01),
C07D 265/00 (2006.01)
(71) Demandeur/Applicant:
TIBOTEC PHARMACEUTICALS LTD., IE
(72) Inventeurs/Inventors:
TAHRI, ABDELLAH, BE;
MICHIELS, LIEVE EMMA JAN, BE;
VAN ACKER, KOENRAAD LODEWIJK AUGUST, BE;
SURLERAUX, DOMINIQUE LOUIS NESTOR GHISLAIN,
BE
(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : DERIVES DE (1,10B-DIHYDRO-2-(AMINOCARBONYL-PHENYL)-5H-PYRAZOLO[1,5- C][1,3]BENZOXAZIN-5-YL)PHENYLMETHANONE UTILISES EN TANT QU'INHIBITEURS DE REPLICATION VIRALE DU VIH
(54) Title: (1,10B-DIHYDRO-2-(AMINOCARBONYL-PHENYL)-5H-PYRAZOLO[1,5- C][1,3]BENZOXAZIN-5-YL)PHENYL
METHANONE DERIVATIVES AS HIV VIRAL REPLICATION INHIBITORS

(57) **Abrégé/Abstract:**

The present invention relates to 5H-pyrazolo[1,5-c] [1,3]benzoxazin-5-yl)phenyl methanone derivatives as inhibitors of the viral replication of the HIV virus, processes for their preparation as well as pharmaceutical compositions, their use as medicines, and diagnostic kits comprising them. The present invention also concerns combinations of the present HIV inhibitors with other anti-retroviral agents. It further relates to their use in assays as reference compounds or as reagents. The compounds of the present invention are useful for preventing or treating infection by HIV and for treating AIDS.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 November 2005 (10.11.2005)

PCT

(10) International Publication Number
WO 2005/105810 A3

(51) International Patent Classification⁷: **C07D 498/04**,
A61K 31/538 // C07D 498/04, 265:00, 231:00

SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN,
YU, ZA, ZM, ZW.

(21) International Application Number:
PCT/EP2005/052041

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(22) International Filing Date: 4 May 2005 (04.05.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
04101913.4 4 May 2004 (04.05.2004) EP
60/604,275 25 August 2004 (25.08.2004) US

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations*
- *of inventorship (Rule 4.17(iv)) for US only*

(71) Applicant (*for all designated States except US*): **TI-BOTEC PHARMACEUTICALS LTD.** [IE/IE]; Little Island, Co Cork (IE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **TAHRI, Abdellah** [BE/BE]; Rue Maurice Albert Raskin 30, B-1070 Anderlecht (BE). **MICHIELS, Lieve, Emma, Jan** [BE/BE]; Schoolstraat 95, B-2400 Mol (BE). **VAN ACKER, Koenraad, Lodewijk, August** [BE/BE]; Uilenberg 30, B-9140 Temse (BE). **SURLERAUX, Dominique, Louis, Nestor, Ghislain** [BE/BE]; rue de la scaillée 13, B-1440 Braine-le-château (BE).

(74) Agent: **DAELEMANS, Frank**; Tibotec-Virco Comm. VA, Generaal De Wittelaan L 11B 3, B-2800 Mechelen (BE).

Published:

- *with international search report*

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM,

(88) Date of publication of the international search report:
2 March 2006

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: (1,10B-DIHYDRO-2-(AMINOCARBONYL-PHENYL)-5H-PYRAZOLO[1,5-C][1,3]BENZOXAZIN-5-YL)PHENYL METHANONE DERIVATIVES AS HIV VIRAL REPLICATION INHIBITORS

(57) Abstract: The present invention relates to 5H-pyrazolo[1,5-c][1,3]benzoxazin-5-yl)phenyl methanone derivatives as inhibitors of the viral replication of the HIV virus, processes for their preparation as well as pharmaceutical compositions, their use as medicines, and diagnostic kits comprising them. The present invention also concerns combinations of the present HIV inhibitors with other anti-retroviral agents. It further relates to their use in assays as reference compounds or as reagents. The compounds of the present invention are useful for preventing or treating infection by HIV and for treating AIDS.



WO 2005/105810 A3


- 1 -

(1,10B-DIHYDRO-2-(AMINOCARBONYL-PHENYL)-5H-PYRAZOLO[1,5-C][1,3]BENZOXAZIN-5-YL)PHENYL METHANONE
DERIVATIVES AS HIV VIRAL REPLICATION INHIBITORS

5 The present invention relates to 5H-pyrazolo[1,5-c][1,3]benzoxazin-5-yl)phenyl methanone derivatives as inhibitors of the viral replication of the HIV virus, processes for their preparation as well as pharmaceutical compositions, their use as medicines, and diagnostic kits comprising them. The present invention also concerns combinations of the present HIV inhibitors with other anti-retroviral agents. It further
 10 relates to their use in assays as reference compounds or as reagents. The compounds of the present invention are useful for preventing or treating infection by HIV and for treating AIDS.

The number of people living with HIV/AIDS totalled in December 2001 about 40
 15 million of which more than 37 million adults and about 2.7 million children under 15 years old. The people newly infected with HIV in 2001 alone rose to 5 million whereas there were in 2001 3 million AIDS deaths. Current chemotherapy for these people infected with HIV/AIDS employs the inhibitors of the viral fusion as well as reverse transcriptase (RT) and protease enzymes. In view of the emergence of HIV strains
 20 resistant to the current generation of fusion, RT and protease inhibitors, there exists an increasing need for the development of new antivirals with novel mechanisms of action.

The present invention concerns the compounds having formula (I)



and the N-oxides, stereoisomeric forms and salts thereof,
 wherein

a is zero, 1, 2, 3, 4 or 5;

- 2 -

- R^1 is hydrogen, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{3-12} cycloalkyl, Het, aryl or C_{1-10} alkyl substituted with a substituent selected from the group consisting of Het, aryl, C_{3-12} cycloalkyl, amino and mono-or disubstituted amino wherein the substituents on the amino group are each individually selected from C_{1-10} alkyl, C_{2-10} alkenyl, C_{3-12} cycloalkyl, Het and aryl;
- R^2 is hydrogen, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{3-12} cycloalkyl, Het, aryl or C_{1-10} alkyl substituted with a substituent selected from the group consisting of Het, aryl, C_{3-12} cycloalkyl, amino and mono-or disubstituted amino wherein the substituents on the amino group are each individually selected from C_{1-10} alkyl, C_{2-10} alkenyl, C_{3-12} cycloalkyl, Het and aryl; or
- R^1 and R^2 taken together with the nitrogen atom to which they are attached form a 5 to 12 membered saturated or partially saturated heterocycle which heterocycle may optionally be substituted with C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{3-12} cycloalkyl, C_{1-6} alkyloxycarbonyl, Het, aryl or C_{1-10} alkyl substituted with a substituent selected from the group consisting of Het, aryl, C_{3-12} cycloalkyl, amino and mono-or disubstituted amino wherein the substituents on the amino group are each individually selected from C_{1-10} alkyl, C_{2-10} alkenyl, C_{3-12} cycloalkyl, Het and aryl;
- R^3 is carboxyl, halogen, nitro, C_{1-10} alkyl, C_{3-12} cycloalkyl, polyhalo C_{1-10} alkyl, cyano, amino, mono-or disubstituted amino, aminocarbonyl, mono-or disubstituted aminocarbonyl, C_{1-10} alkyloxy, C_{1-10} alkylthio, C_{1-10} alkylsulfonyl, pyrrolidinyl, piperidinyl, homopiperidinyl, morpholinyl, thiomorpholinyl or piperazinyl optionally substituted with C_{1-10} alkyl, wherein the substituents on any of the amino groups are each individually selected from C_{1-10} alkyl, C_{2-10} alkenyl, C_{3-12} cycloalkyl, Het and aryl;
- aryl is phenyl optionally substituted with one or more substituents selected from the group consisting of C_{1-10} alkyl, polyhalo C_{1-10} alkyl, C_{1-10} alkyloxy, C_{1-10} alkylthio, C_{1-10} alkylsulfonyl, nitro, cyano, halo, C_{3-7} cycloalkyl, C_{1-10} alkylcarbonyl, carboxyl, C_{1-10} alkyloxycarbonyl, amino, mono-or disubstituted amino, aminocarbonyl, mono-or disubstituted aminocarbonyl, wherein the substituents on any of the amino groups are each individually selected from phenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{3-7} cycloalkyl, pyrrolidinyl, piperidinyl, homopiperidinyl or piperazinyl optionally substituted with C_{1-10} alkyl;
- Het is a 5 or 6 membered aromatic, saturated or partially saturated, monocyclic or bicyclic heterocycle with one or more heteroatoms each individually selected from nitrogen, oxygen or sulfur, and which heterocycle may optionally be substituted by one, or where possible, more than one substituent selected from the group consisting of C_{1-10} alkyl, polyhalo C_{1-10} alkyl, C_{1-10} alkyloxy, C_{1-10} alkylthio,

- 3 -

5 C₁₋₁₀alkylsulfonyl, nitro, cyano, halo, C₃₋₇cycloalkyl, C₁₋₁₀alkylcarbonyl, carboxyl, C₁₋₁₀alkyloxycarbonyl, amino, mono-or disubstituted amino, aminocarbonyl, mono-or disubstituted aminocarbonyl, wherein the substituents on any of the amino groups are each individually selected from phenyl, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₃₋₇cycloalkyl, pyrrolidinyl, piperidinyl, homopiperidinyl or piperazinyl optionally substituted with C₁₋₁₀alkyl.

10 As used herein, the term "halo" or "halogen" as a group or part of a group is generic for fluoro, chloro, bromo or iodo. The term "polyhalo" used as a prefix means substituted with one or more halogen atoms. Examples of the use of polyhalo as a prefix include for instance polyhaloC₁₋₁₀alkyl, polyhaloC₁₋₆alkyl, polyhaloC₁₋₄alkyl and the like. Particular interesting polyhaloalkyls are difluoromethyl and trifluoromethyl.

15 The term C₁₋₄alkyl as a group or part of a group means straight and branched chained saturated monovalent hydrocarbon radicals containing from 1 to 4 carbon atoms. Examples of such C₁₋₄alkyl radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl and the like.

20 The term C₁₋₆alkyl as a group or part of a group means straight and branched chained saturated monovalent hydrocarbon radicals containing from 1 to 6 carbon atoms. Examples of such C₁₋₆alkyl radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, 2-methylbutyl, pentyl, iso-amyl, hexyl, 3-methylpentyl and the like.

25 The term C₁₋₁₀alkyl as a group or part of a group means straight and branched chained saturated monovalent hydrocarbon radicals containing from 1 to 10 carbon atoms. Examples of such C₁₋₁₀alkyl radicals include the examples of C₁₋₆alkyl radicals and heptyl, octyl, nonyl, decyl, 3-ethyl-heptyl and the like.

30 C₂₋₆alkenyl as a group or part of a group means straight and branched chained monovalent hydrocarbon radicals having at least one double bond and containing from 2 to 6 carbon atoms. Examples of such C₂₋₆alkenyl radicals include ethenyl, propenyl, 1-butenyl, 2-butenyl, isobutenyl, 2-methyl-1-butenyl, 1-pentenyl, 2-pentenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 3-methyl-2-pentenyl and the like.

35 C₂₋₁₀alkenyl as a group or part of a group means straight and branched chained monovalent hydrocarbon radicals having at least one double bond and containing from 2 to 10 carbon atoms. Examples of such C₂₋₁₀alkenyl radicals include the examples of C₂₋₆alkenyl and 2-heptenyl, 3-heptenyl, 3-octenyl, 4-octenyl, 4-nonenyl, 4-decenyl and

- 4 -

the like.

C₂₋₆alkynyl as a group or part of a group means straight and branched chained monovalent hydrocarbon radicals having at least one triple bond and containing from
 5 2 to 6 carbon atoms. Examples of such C₂₋₆alkynyl radicals include ethynyl, propynyl, 1-butynyl, 2-butynyl, isobutynyl, 2-methyl-1-butynyl, 1-pentynyl, 2-pentynyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 3-methyl-2-pentynyl and the like.

C₂₋₁₀alkynyl as a group or part of a group means straight and branched chained
 10 monovalent hydrocarbon radicals having at least one triple bond and containing from 2 to 10 carbon atoms. Examples of such C₂₋₁₀alkynyl radicals include the examples of C₂₋₆alkynyl and 2-heptynyl, 3-heptynyl, 3-octynyl, 4-octynyl, 4-nonyl, 4-decynyl and the like.

15 The term C₃₋₇cycloalkyl as a group or part of a group means carbocyclic or spiro carbocyclic monovalent hydrocarbon radicals having from 3 to 7 carbon atoms in the backbone of the carbocycle or spirocarbocycle. Examples of such C₃₋₇cycloalkyl radicals include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, spiro[2.4]heptanyl, cycloheptyl and the like.

20 The term C₃₋₁₂cycloalkyl as a group or part of a group means carbocyclic or spiro carbocyclic monovalent hydrocarbon radicals having from 3 to 12 carbon atoms in the backbone of the carbocycle or spirocarbocycle. Examples of such C₃₋₁₂cycloalkyl radicals include the examples of C₃₋₇cycloalkyl and cyclooctyl, cyclononyl,
 25 spiro[4.4]nonyl, spiro[4.5]decyl, spiro[5.5]undecyl, cyclododecyl, spiro[5.6]dodecyl and the like.

Examples of 5 or 6 membered heterocycles as defined by Het include but are not limited to pyridine, pyrimidine, pyridazine, pyrazine, triazine, imidazole, thiazole,
 30 oxazole, oxadiazole, thiadiazole, isothiazole, isoxazole, pyrazole, furane, thiophene, pyrrol, quinoline, isoquinoline, benzoxazole, isobenzoxazole, benzothiazole, isobenzothiazole, benzimidazole, benzotriazole, tetrahydroquinoline, tetrahydro-isoquinoline, piperidine, piperazine, morpholine, thiomorpholine, pyrrolidine, homopiperidine, homopiperazine, tetrahydrofurane and tetrahydrothienyl.

35 As used herein, the term C(=O) is meant to define a carbonyl moiety and the term S(=O)₂ is meant to define a sulfonyl moiety. As used herein, the term hydroxy means -OH, the term nitro means -NO₂, the term cyano means -CN, the term thio means -S, the term oxo means =O.

- 5 -

Whenever the terms "one or more substituents" or "substituted" are used in defining the compounds of formula (I), it is meant to indicate that one or more hydrogens on the atom indicated in the expressions using "one or more substituents" or "substituted" is replaced with a selection from the indicated group, provided that the indicated atom's normal valency is not exceeded, and that the substitution results in a chemically stable compound, i.e. a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into a therapeutic agent.

For therapeutic use, the salts of the compounds of the present invention are those wherein the counter-ion is pharmaceutically or physiologically acceptable. However, salts having a pharmaceutically unacceptable counter-ion may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound of the present invention. All salts, whether pharmaceutically acceptable or not are included within the ambit of the present invention.

The pharmaceutically acceptable or physiologically tolerable acid addition salt forms which the compounds of the present invention are able to form can conveniently be prepared using the appropriate acids, such as, for example, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid, sulfuric, nitric, phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, p-toluenesulfonic, cyclamic, salicylic, p-aminosalicylic, pamoic and the like acids.

Conversely said acid addition salt forms can be converted by treatment with an appropriate base into the free base form.

The compounds of the present invention containing an acidic proton may also be converted into their non-toxic metal or amine addition salt form by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts, quaternary ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. the benzathine, N-methyl, -D-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

Conversely said base addition salt forms can be converted by treatment with an appropriate acid into the free acid form.

The term "salts" also comprises the hydrates and the solvent addition forms that the

- 6 -

compounds of the present invention are able to form. Examples of such forms are e.g. hydrates, alcoholates and the like. The term "salts" also comprises the quaternization of the nitrogen atoms of the present compounds. A basic nitrogen can be quaternized with any agent known to those of ordinary skill in the art including, for instance, lower alkyl halides, dialkyl sulfates, long chain halides and arylalkyl halides.

The N-oxide forms of the present compounds are meant to comprise the compounds wherein one or several nitrogen atoms are oxidized to the so-called N-oxide.

The present compounds may also exist in their tautomeric forms. Such forms, although not explicitly indicated in the above formula, are intended to be included within the scope of the present invention.

Whenever used in the present patent application, the terms "present compounds", "compounds of the present invention", "compounds of formula (I)" or similar terms are meant to comprise the compounds of formula (I), their N-oxides, their stereoisomeric forms, their salt forms or any subgroup thereof.

Interesting compounds are those compounds of formula (I) or any subgroup thereof wherein a is zero, 1 or 2; more in particular wherein a is zero or 1.

Interesting compounds are those compounds of formula (I) wherein R^1 is hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-7} cycloalkyl, Het, aryl or C_{1-6} alkyl substituted with a substituent selected from the group consisting of Het, aryl, C_{3-7} cycloalkyl, amino and mono-or disubstituted amino wherein the substituents on the amino group are each individually selected from C_{1-6} alkyl or C_{3-7} cycloalkyl; in particular wherein R^1 is C_{1-6} alkyl, C_{3-7} cycloalkyl, Het, aryl or C_{1-6} alkyl substituted with a substituent selected from the group consisting of Het, aryl, C_{3-7} cycloalkyl and mono-or disubstituted amino wherein the substituents on the amino group are each individually selected from C_{1-6} alkyl.

Interesting compounds are those compounds of formula (I) or any subgroup thereof wherein R^2 is hydrogen, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{3-12} cycloalkyl, or C_{1-10} alkyl substituted with a substituent selected from the group consisting of Het, aryl, C_{3-7} cycloalkyl, amino and mono-or disubstituted amino wherein the substituents on the amino group are each individually selected from C_{1-6} alkyl; in particular wherein R^2 is hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, or C_{1-6} alkyl substituted with a substituent selected from the group consisting of aryl.

- 7 -

Interesting compounds are those compounds of formula (I) or any subgroup thereof wherein R^1 and R^2 taken together with the nitrogen atom to which they are attached form a 5 to 12 membered saturated or partially saturated heterocycle which heterocycle may optionally be substituted with C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxycarbonyl, Het, aryl or C_{1-6} alkyl substituted with a substituent selected from the group consisting of Het, aryl, C_{3-7} cycloalkyl, amino and mono-or disubstituted amino wherein the substituents on the amino group are each individually selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-7} cycloalkyl, Het and aryl;

in particular wherein R^1 and R^2 taken together with the nitrogen atom to which they are attached form a 5 to 12 membered saturated or partially saturated heterocycle which heterocycle may optionally be substituted with C_{1-6} alkyl, C_{1-6} alkyloxycarbonyl or C_{1-6} alkyl substituted with aryl;

more in particular wherein R^1 and R^2 taken together with the nitrogen atom to which they are attached form a heterocycle selected from the group consisting of piperazinyl, homopiperazinyl, piperidinyl, homopiperidinyl, morpholinyl, thiomorpholinyl, pyrrolidinyl, indolinyl, 1,2,3,4-tetrahydroquinolinyl, 1,2,3,4-tetrahydro-isoquinolinyl, which heterocycle may optionally be substituted with C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxycarbonyl, Het, aryl or C_{1-6} alkyl substituted with a substituent selected from the group consisting of Het, aryl, C_{3-7} cycloalkyl, amino and mono-or disubstituted amino wherein the substituents on the amino group are each individually selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-7} cycloalkyl, Het and aryl;

and even more in particular wherein R^1 and R^2 taken together with the nitrogen atom to which they are attached form a heterocycle selected from the group consisting of piperazinyl, homopiperazinyl, piperidinyl, homopiperidinyl, morpholinyl, thiomorpholinyl, pyrrolidinyl, indolinyl, 1,2,3,4-tetrahydroquinolinyl, 1,2,3,4-tetrahydro-isoquinolinyl, which heterocycle may optionally be substituted with C_{1-6} alkyl, C_{1-6} alkyloxycarbonyl or C_{1-6} alkyl substituted with aryl.

Interesting compounds are those compounds of formula (I) or any subgroup thereof wherein R^3 is carboxyl, halogen, nitro, C_{1-6} alkyl, C_{3-7} cycloalkyl, polyhalo C_{1-4} alkyl, cyano, amino, mono-or disubstituted amino, aminocarbonyl, mono-or disubstituted aminocarbonyl, C_{1-6} alkyloxy, C_{1-6} alkylthio, C_{1-6} alkylsulfonyl, pyrrolidinyl, piperidinyl, homopiperidinyl, morpholinyl, thiomorpholinyl or piperazinyl optionally substituted with C_{1-6} alkyl, wherein the substituents on any of the amino groups are each individually selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{3-7} cycloalkyl, Het and aryl;

in particular wherein R^3 is halogen, nitro, C_{1-6} alkyl, C_{3-7} cycloalkyl, polyhalo C_{1-4} alkyl, cyano, C_{1-6} alkyloxy, C_{1-6} alkylthio, C_{1-6} alkylsulfonyl, piperidinyl, morpholinyl.

- 8 -

- Interesting compounds are those compounds of formula (I) or any subgroup thereof wherein aryl is phenyl optionally substituted with one or more substituents selected from the group consisting of C₁₋₆alkyl, polyhaloC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkylsulfonyl, nitro, cyano, halo, C₃₋₇cycloalkyl, C₁₋₆alkylcarbonyl, carboxyl,
- 5 C₁₋₆alkyloxycarbonyl, amino, mono-or disubstituted amino, aminocarbonyl, mono-or disubstituted aminocarbonyl, wherein the substituents on any of the amino groups are each individually selected from phenyl, C₁₋₆alkyl, C₂₋₆alkenyl, C₃₋₇cycloalkyl, pyrrolidinyl, piperidinyl, homopiperidinyl or piperazinyl optionally substituted with C₁₋₆alkyl;
- 10 in particular wherein aryl is phenyl optionally substituted with one, two or three substituents selected from the group consisting of C₁₋₆alkyl, C₁₋₆alkyloxy, nitro, cyano, halo, amino, mono-or disubstituted amino, wherein the substituents on any of the amino groups are each individually selected from C₁₋₆alkyl.
- 15 Interesting compounds are those compounds of formula (I) or any subgroup thereof wherein Het is a 5 or 6 membered aromatic, saturated or partially saturated, monocyclic or bicyclic heterocycle with one or more heteroatoms each individually selected from nitrogen, oxygen or sulfur, and which heterocycle may optionally be substituted by one, or where possible, more than one substituent selected from the group consisting of
- 20 C₁₋₆alkyl, polyhaloC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkylsulfonyl, nitro, cyano, halo, C₃₋₇cycloalkyl, C₁₋₆alkylcarbonyl, carboxyl, C₁₋₆alkyloxycarbonyl, amino, mono-or disubstituted amino, aminocarbonyl, mono-or disubstituted aminocarbonyl, wherein the substituents on any of the amino groups are each individually selected from phenyl, C₁₋₆alkyl, C₂₋₆alkenyl, C₃₋₇cycloalkyl, pyrrolidinyl, piperidinyl, homopiperidinyl or
- 25 piperazinyl optionally substituted with C₁₋₆alkyl;
- in particular wherein Het is a 5 or 6 membered aromatic, saturated or partially saturated, monocyclic or bicyclic heterocycle with one or more heteroatoms each individually selected from nitrogen, oxygen or sulfur, and which heterocycle may optionally be substituted with C₁₋₆alkyl;
- 30 more in particular wherein Het is a heterocycle selected from the group consisting of pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, pyranyl, pyrrolyl, furanyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, tetrahydrofuranyl, tetrahydrothienyl, imidazolyl, dioxolanyl, pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, homopiperazinyl, dioxanyl, morpholinyl, thiomorpholinyl, which
- 35 heterocycle may optionally be substituted by one, or where possible, more than one substituent selected from the group consisting of C₁₋₆alkyl, polyhaloC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkylsulfonyl, nitro, cyano, halo, C₃₋₇cycloalkyl, C₁₋₆alkylcarbonyl, carboxyl, C₁₋₆alkyloxycarbonyl, amino, mono-or disubstituted

- 9 -

amino, aminocarbonyl, mono-or disubstituted aminocarbonyl, wherein the substituents on any of the amino groups are each individually selected from phenyl, C₁₋₆alkyl, C₂₋₆alkenyl, C₃₋₇cycloalkyl, pyrrolidiny, piperidiny, homopiperidiny or piperaziny optionally substituted with C₁₋₆alkyl;

- 5 even more in particular wherein Het is a heterocycle selected from the group consisting of pyridiny, pyrimidiny, pyraziny, pyridaziny, triaziny, pyran, pyrroly, furany, thieny, oxazol, isoxazol, thiazol, isothiazol, imidazol, tetrahydrofurany, tetrahydrothieny, imidazoliny, dioxolany, pyrrolidiny, piperidiny, homopiperidiny, piperaziny, homopiperaziny, dioxany, morpholin, thiomorpholin, which
10 heterocycle may optionally be substituted by C₁₋₆alkyl.

- Further interesting compounds are those compounds of formula (I) or any subgroup thereof wherein Het is a heterocycle selected from the group consisting of pyridiny, furany, thieny, tetrahydrofurany, pyrrolidiny, piperidiny, which heterocycle may
15 optionally be substituted by one, or where possible, more than one substituent selected from the group consisting of C₁₋₆alkyl, polyhaloC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkylsulfonyl, nitro, cyano, halo, C₃₋₇cycloalkyl, C₁₋₆alkylcarbonyl, carboxyl, C₁₋₆alkyloxycarbonyl, amino, mono-or disubstituted amino, aminocarbonyl, mono-or disubstituted aminocarbonyl, wherein the substituents on any of the amino groups are
20 each individually selected from phenyl, C₁₋₆alkyl, C₂₋₆alkenyl, C₃₋₇cycloalkyl, pyrrolidiny, piperidiny, homopiperidiny or piperaziny optionally substituted with C₁₋₆alkyl;

- in particular wherein Het is a heterocycle selected from the group consisting of pyridiny, furany, thieny, tetrahydrofurany, pyrrolidiny, piperidiny, which
25 heterocycle may optionally be substituted by C₁₋₆alkyl.


An interesting group of compounds are those compounds of formula (I) where one or more of the following restrictions apply :

1. a is zero or 1;
- 30 2. R¹ is C₁₋₁₀alkyl, C₃₋₁₂cycloalkyl, Het, aryl or C₁₋₁₀alkyl substituted with a substituent selected from the group consisting of Het, aryl, C₃₋₁₂cycloalkyl and mono-or disubstituted amino wherein the substituents on the amino group are each individually selected from C₁₋₁₀alkyl;
3. R² is hydrogen, C₁₋₁₀alkyl, C₂₋₁₀alkenyl or C₁₋₁₀alkyl substituted with aryl;
- 35 4. R¹ and R² taken together with the nitrogen atom to which they are attached form a 5 to 12 membered saturated or partially saturated heterocycle which heterocycle may optionally be substituted with C₁₋₁₀alkyl, C₁₋₆alkyloxycarbonyl or C₁₋₁₀alkyl substituted with aryl;

- 10 -

5. R^3 is halogen, nitro, C_{1-10} alkyl, C_{3-12} cycloalkyl, polyhalo C_{1-10} alkyl, cyano, C_{1-10} alkyloxy, C_{1-10} alkylthio, C_{1-10} alkylsulfonyl, piperidinyl, morpholinyl;
6. aryl is phenyl optionally substituted with one or more substituents selected from the group consisting of C_{1-10} alkyloxy, halo, mono-or disubstituted amino, wherein the substituents on any of the amino groups are each individually selected from C_{1-10} alkyl;
7. Het is a 5 or 6 membered aromatic, saturated or partially saturated, monocyclic or bicyclic heterocycle with one or more heteroatoms each individually selected from nitrogen, oxygen or sulfur, and which heterocycle may optionally be substituted by C_{1-10} alkyl.

Particular compounds are those compounds of formula (I) or any subgroup thereof, such as the interesting compounds defined above, wherein the $-C(=O)-NR^1R^2$ moiety is in the para position on the phenyl ring which is linked to the 2-position of the 1,9b-dihydro-5-oxa-3,3a-diaza-cyclopenta[a]naphthalene scaffold (also named 1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazine scaffold) as depicted in the figure below by the compound of formula (Ia)



Particular compounds are those compounds of formula (I) or any subgroup thereof, such as the interesting compounds defined above, wherein a is zero or 1 and R^3 is halogen, nitro, C_{1-10} alkyl, C_{3-12} cycloalkyl, polyhalo C_{1-10} alkyl, cyano, C_{1-10} alkyloxy, C_{1-10} alkylthio, C_{1-10} alkylsulfonyl, piperidinyl, morpholinyl.

Particular compounds are those compounds of formula (I) or (Ia) or any subgroup thereof, such as the interesting compounds defined above, wherein R^1 is C_{1-10} alkyl, C_{3-12} cycloalkyl, Het, aryl or C_{1-10} alkyl substituted with a substituent selected from the group consisting of Het, aryl, C_{3-12} cycloalkyl and mono-or disubstituted amino wherein the substituents on the amino group are each individually selected from C_{1-10} alkyl; R^2 is hydrogen, C_{1-10} alkyl, C_{2-10} alkenyl or C_{1-10} alkyl substituted with aryl; or R^1 and R^2 taken

- 11 -

together with the nitrogen atom to which they are attached form a 5 to 12 membered saturated or partially saturated heterocycle which heterocycle may optionally be substituted with C₁₋₁₀alkyl, C₁₋₆alkyloxycarbonyl or C₁₋₁₀alkyl substituted with aryl.

- 5 Preferred compounds are those compounds of formula (I) or (Ia) or any subgroup thereof, such as the interesting and particular compounds defined above, wherein a is zero or 1;

R¹ is C₁₋₁₀alkyl, C₃₋₁₂cycloalkyl, Het, aryl or C₁₋₁₀alkyl substituted with a substituent selected from the group consisting of Het, aryl, C₃₋₁₂cycloalkyl and mono-or
10 disubstituted amino wherein the substituents on the amino group are each individually selected from C₁₋₁₀alkyl;

R² is hydrogen, C₁₋₁₀alkyl, C₂₋₁₀alkenyl or C₁₋₁₀alkyl substituted with aryl; or

R¹ and R² taken together with the nitrogen atom to which they are attached form a 5 to
15 12 membered saturated or partially saturated heterocycle which heterocycle may optionally be substituted with C₁₋₁₀alkyl, C₁₋₆alkyloxycarbonyl or C₁₋₁₀alkyl substituted with aryl;

R³ is halogen, nitro, C₁₋₁₀alkyl, C₃₋₁₂cycloalkyl, polyhaloC₁₋₁₀alkyl, cyano, C₁₋₁₀alkyloxy, C₁₋₁₀alkylthio, C₁₋₁₀alkylsulfonyl, piperidinyl, morpholinyl.

- 20 Preferred compounds are those compounds of formula (I) or (Ia) or any subgroup thereof, such as the interesting and particular compounds defined above, wherein a is zero or 1;

R¹ is C₁₋₁₀alkyl, C₃₋₁₂cycloalkyl, Het, aryl or C₁₋₁₀alkyl substituted with a substituent selected from the group consisting of Het, aryl, C₃₋₁₂cycloalkyl and mono-or
25 disubstituted amino wherein the substituents on the amino group are each individually selected from C₁₋₁₀alkyl;

R² is hydrogen, C₁₋₁₀alkyl, C₂₋₁₀alkenyl or C₁₋₁₀alkyl substituted with aryl; or

R¹ and R² taken together with the nitrogen atom to which they are attached form a 5 to
30 12 membered saturated or partially saturated heterocycle which heterocycle may optionally be substituted with C₁₋₁₀alkyl, C₁₋₆alkyloxycarbonyl or C₁₋₁₀alkyl substituted with aryl;

R³ is halogen, nitro, C₁₋₁₀alkyl, C₃₋₁₂cycloalkyl, polyhaloC₁₋₁₀alkyl, cyano, C₁₋₁₀alkyloxy, C₁₋₁₀alkylthio, C₁₋₁₀alkylsulfonyl, piperidinyl, morpholinyl;

aryl is phenyl optionally substituted with one or more substituents selected from the
35 group consisting of C₁₋₁₀alkyloxy, halo, mono-or disubstituted amino, wherein the substituents on any of the amino groups are each individually selected from C₁₋₁₀alkyl;

- 12 -

Het is a 5 or 6 membered aromatic, saturated or partially saturated, monocyclic or bicyclic heterocycle with one or more heteroatoms each individually selected from nitrogen, oxygen or sulfur, and which heterocycle may optionally be substituted by C₁₋₁₀alkyl.

5

More preferred compounds are those compounds of formula (I) or (Ia) or any subgroup thereof, such as the interesting and particular compounds defined above, wherein a is zero or 1;

10 R¹ is C₁₋₆alkyl, C₃₋₇cycloalkyl, Het, aryl or C₁₋₆alkyl substituted with a substituent selected from the group consisting of Het, aryl, C₃₋₇cycloalkyl and mono-or disubstituted amino wherein the substituents on the amino group are each individually selected from C₁₋₆alkyl;

R² is hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl, or C₁₋₆alkyl substituted with a substituent selected from the group consisting of aryl or

15 R¹ and R² taken together with the nitrogen atom to which they are attached form a heterocycle selected from the group consisting of piperazinyl, homopiperazinyl, piperidinyl, homopiperidinyl, morpholinyl, thiomorpholinyl, pyrrolidinyl, indolinyl, 1,2,3,4-tetrahydroquinolinyl, 1,2,3,4-tetrahydro-isoquinolinyl, which heterocycle may optionally be substituted with C₁₋₆alkyl, C₁₋₆alkyloxycarbonyl or C₁₋₆alkyl substituted with aryl;

20

R³ is halogen, nitro, C₁₋₆alkyl, C₃₋₇cycloalkyl, polyhaloC₁₋₄alkyl, cyano, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkylsulfonyl, piperidinyl, morpholinyl;

aryl is phenyl optionally substituted with one, two or three substituents selected from the group consisting of C₁₋₆alkyl, C₁₋₆alkyloxy, nitro, cyano, halo, amino, mono-or 25 disubstituted amino, wherein the substituents on any of the amino groups are each individually selected from C₁₋₆alkyl;

25

Het is a heterocycle selected from the group consisting of pyridinyl, furanyl, thienyl, tetrahydrofuranyl, pyrrolidinyl, piperidinyl, which heterocycle may optionally be substituted by C₁₋₆alkyl.

30

Other more preferred compounds are the compounds of formula (I) or (Ia) or any subgroup thereof, such as the interesting, particular and preferred compounds defined above, wherein R¹ is cyclopentyl and R² is hydrogen.

35 Other more preferred compounds are the compounds of formula (I) or any subgroup thereof, such as the interesting, particular and preferred compounds defined above, wherein a is 1 and R³ is halogen, cyano, C₁₋₄alkyl, C₁₋₄alkylthio; morpholinyl, C₁₋₄alkyloxy, nitro, C₁₋₄alkylsulfonyl, trifluoromethyl.

- 13 -

Most preferred compounds are

- 4-[5-(4-fluorobenzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-[(4-methoxyphenyl)methyl]-N-methyl-benzamide;
- 4-[5-(4-cyanobenzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-cyclopentyl-benzamide;
- 4-[5-(4-propylbenzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-cyclopentyl-benzamide;
- 4-[5-(4-methylsulfanyl-benzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-cyclopentyl-benzamide;
- 4-[5-(4-ethyl-benzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-cyclopentyl-benzamide;
- 4-[5-(4-tert-butyl-benzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-cyclopentyl-benzamide;
- 4-[5-(4-nitro-benzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-cyclopentyl-benzamide;
- 4-[5-(4-methanesulfonyl-benzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-cyclopentyl-benzamide;
- 4-[5-(4-trifluoromethyl-benzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-cyclopentyl-benzamide
- or a N-oxide, stereoisomeric form or salt thereof.

Due to the fact that the compounds of formula (I) are inhibitors of the replication of HIV, the compounds of formula (I) are useful in the treatment of warm-blooded animals, in particular humans, infected with HIV. Conditions associated with HIV which may be prevented or treated with the compounds of the present invention include AIDS, AIDS-related complex (ARC), progressive generalized lymphadenopathy (PGL), as well as chronic CNS diseases caused by retroviruses, such as, for example HIV mediated dementia and multiple sclerosis.

The compounds of the present invention may therefore be used as medicines against or in a method of treating above-mentioned conditions. Said use as a medicine or method of treatment comprises the systemic administration of an effective therapeutic amount of a compound of formula (I) to HIV-infected warm-blooded animals, in particular HIV-infected humans. Consequently, the compounds of the present invention can be used in the manufacture of a medicament useful for treating conditions associated with HIV infection.

In one embodiment, the invention relates to the use of the compounds of the present

- 14 -

invention in the manufacture of a medicament for preventing HIV transmission or HIV infection or disease associated with HIV infection of warm-blooded animals, in particular humans, and in particular transmission or infection via sexual intercourse or related intimate contact between partners. Thus, the invention also relates to a method of preventing HIV transmission or HIV infection or a disease associated with HIV infection comprising administering to a warm-blooded animal, in particular a human, an effective preventive amount of a compound of formula (I), and in particular transmission or infection via sexual intercourse or related intimate contact between partners.

10

The term stereochemically isomeric forms of compounds of the present invention, as used hereinbefore, defines all possible compounds made up of the same atoms bonded by the same sequence of bonds but having different three-dimensional structures which are not interchangeable, which the compounds of the present invention may possess.

15

Unless otherwise mentioned or indicated, the chemical designation of a compound encompasses the mixture of all possible stereochemically isomeric forms which said compound may possess. Said mixture may contain all diastereomers and/or enantiomers of the basic molecular structure of said compound. All stereochemically isomeric forms of the compounds of the present invention both in pure form or in admixture with each other are intended to be embraced within the scope of the present invention.

20

Pure stereoisomeric forms of the compounds and intermediates as mentioned herein are defined as isomers substantially free of other enantiomeric or diastereomeric forms of the same basic molecular structure of said compounds or intermediates. In particular, the term 'stereoisomerically pure' concerns compounds or intermediates having a stereoisomeric excess of at least 80% (i. e. minimum 80% of one isomer and maximum 20% of the other possible isomers) up to a stereoisomeric excess of 100% (i.e. 100% of one isomer and none of the other), more in particular, compounds or intermediates having a stereoisomeric excess of 90% up to 100%, even more in particular having a stereoisomeric excess of 94% up to 100% and most in particular having a stereoisomeric excess of 97% up to 100%. The terms 'enantiomerically pure' and 'diastereomerically pure' should be understood in a similar way, but then having regard to the enantiomeric excess and the diastereomeric excess respectively, of the mixture in question.

30

35

Pure stereoisomeric forms of the compounds and intermediates of this invention may be obtained by the application of art-known procedures. For instance, enantiomers may

- 15 -

- be separated from each other by the selective crystallization of their diastereomeric salts with optically active acids. Alternatively, enantiomers may be separated by chromatographic techniques using chiral stationary phases. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically. Preferably, if a specific stereoisomer is desired, said compound will be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.
- 10 The diastereomeric racemates of the compounds of the present invention can be obtained separately by conventional methods. Appropriate physical separation methods which may advantageously be employed are, for example, selective crystallization and chromatography, e.g. column chromatography.
- 15 The compounds may contain one or more asymmetric centers and thus may exist as different stereoisomeric forms. The absolute configuration of each asymmetric center that may be present in the compounds may be indicated by the stereochemical descriptors R and S, this R and S notation corresponding to the rules described in Pure Appl. Chem. 1976, 45, 11-30.
- 20 The present invention is also intended to include all isotopes of atoms occurring on the present compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen include tritium and deuterium. Isotopes of carbon include C-13 and C-14.
- 25 The favourable properties of the present compounds with respect to their ability to inhibit HIV viral replication can be demonstrated using an anti-viral replication assay that directly measures the ongoing replication of a wild-type HIV virus in MT4 cells via the specific interaction of HIV-tat with LTR sequences coupled to GFP (MT4-LTR-EGFP cells). It has also been found, using the above anti-viral replication assay, that
- 30 the present compounds inhibit the replication a panel of reverse transcriptase resistant viruses, protease resistant viruses or combined reverse transcriptase and protease resistant viruses (so-called multi-drug resistant viruses).
- 35 With respect to the mechanism of action by which the present compounds inhibit viral replication, it has been demonstrated the present compounds inhibit the early stage of the viral replication process in a time of addition experiment. Time of addition experiments provide an indication at what time point a test compound inhibits the viral replication in a cellular environment. In particular, test compounds of the present

- 16 -

invention were added at different time intervals to HIV-1 infected MT4 cells expressing HIV-1-LTR-luciferase reporter gene (MT4-LTR-Luc) or MT4 cells expressing HIV-1-LTR-EGFP reporter gene (MT4-LTR-EGFP). The first time point for addition of a test compound was 30 minutes post virus synchronisation.

5


Furthermore, it has been found that the present compounds have little or no activity in an entry reporter assay (ERA) that measures inhibition of cell-cell fusion between cell line persistently expressing HIV (effector cell line) and a cell line expressing CD4 and CXCR4 (target cell line) equipped with LTR-EGFP using FACS read-out.

10

A toxicity assay wherein a reduced expression of the GFP reporter protein (MT4-CMV-EGFP cells) serves as a marker for cellular toxicity of a test compound can be used to measure the toxicity of the present compounds.


15 In general, the compounds of the present invention can be obtained by the synthetic sequence depicted in Scheme 1.

Scheme 1



- 17 -


In scheme 1, the intermediate of formula 1.2 can be prepared via the coupling of an amine of formula $\text{NH}(\text{R}^1)(\text{R}^2)$ and a benzoic acid of formula 1.1. This coupling reaction may be performed in an appropriate reaction-inert solvent such as acetonitrile, dichloromethane, tetrahydrofuran or any other solvent that solubilizes the reagents, in the presence of a coupling agent such as EDCI (ethyl dimethylaminopropyl carbodiimide) and HOBt (hydroxy benzotriazol) or a functional equivalent thereof. The intermediates of formula 1.2 can then be condensed with a salicylaldehyde of formula 1.3 in the presence of suitable base such as mineral base like potassium hydroxide or sodium hydroxide in a suitable solvent system such as for instance a mixture of an alcohol and water, e.g. a mixture of ethanol and water. Alternatively, an organic base such as pyrrolidine in a suitable solvent such as tetrahydrofuran or dichloromethane can be used. The variable "PG" in intermediate 1.3 is meant to be a protecting group for the hydroxygroup such as for instance methoxymethyl, tert-butoxymethyl, tetrahydropyranyl or methoxyethoxymethyl (MEM). Better yields were observed when using the protected salicylaldehydes of formula 1.3. A particularly suitable protecting group is for instance MEM. Scheme 2 depicts the reaction procedure to prepare an intermediate of formula 1.3 wherein the protecting group is MEM. The resulting intermediate of the above condensation reaction is a chalcone of formula 1.4 (also described in J. Med. Chem, 2000, 43, 4868-4876). The protected chalcone of formula 1.4 can be deprotected in acidic media for instance by using hydrochloric acid in an appropriate solvent such as for example tetrahydrofuran, dichloromethane or an alcohol, thus yielding an intermediate of formula 1.5. Intermediate 1.5 can then be further reacted with hydrazine in a suitable water-miscible solvent such as for instance dioxane, toluene or an alcohol like ethanol, to yield the dihydropyrazol of formula 1.6 (J. Ind. Chem. Soc., 1989, 66, 893-896). The compound of formula (I) can then be prepared via the coupling of the dihydropyrazol of formula 1.6 with an appropriate glyoxal of formula 1.7 in a suitable solvent such as toluene or dioxane, and in the presence of a catalytic amount of an acid such as para-toluenesulfonic acid.

Scheme 2

The preparation of intermediate of formula 1.3 with the MEM as protecting group can be obtained by reacting salicylaldehyde 2.1, which is commercially available, with chloromethoxymethoxy ethane in the presence of a base such as sodium hydride or

- 18 -


potassium hydride, and in the presence of suitable solvent like *N,N*-dimethylformamide or tetrahydrofuran.

Scheme 3

- 5 The preparation of a glyoxal of formula 1.7 can be carried out starting from an intermediate of formula 3.1 in a manner or analogous to a manner described in J. Het. Chem. 1987, 24, 441-451. More in particular, the acetyl of formula 3.1 can be oxidized to a glyoxal of formula 1.7 in a solvent such as dioxane using selenium oxide as oxidating agent.

10

Compounds of formula (I) can be also obtained by the synthetic sequence depicted in Scheme 4.

Scheme 4

- 19 -

According to scheme 4, an intermediate of formula 4.1 can be condensed with a salicylaldehyde of formula 4.2 in the presence of a suitable base such as a mineral base like potassium hydroxide or sodium hydroxide and in a suitable solvent system such as for instance an alcohol or a mixture of an alcohol and water, e.g. a mixture of ethanol and water. This condensation yields a chalcone of formula 4.3. This chalcone of formula 4.3 can then be reacted with hydrazine in a suitable water-miscible solvent such as for instance dioxane, toluene or an alcohol like ethanol, to yield a dihydropyrazol of formula 4.4. Intermediates of formula 4.5 can be prepared via the coupling of the dihydropyrazol of formula 4.4 and an appropriate glyoxal of 1.7 in a suitable solvent such as toluene, *N,N*-dimethylformamide or dioxane, and in the presence of a catalytic amount of an acid such as para-toluenesulfonic acid. Subsequently, the benzoic acid of formula 4.5 can be coupled to an amine of formula $\text{NH}(\text{R}^1)(\text{R}^2)$ in an appropriate reaction-inert solvent such as acetonitrile, dichloromethane, tetrahydrofuran or any other solvent that solubilizes the reagents, and in the presence of a coupling agent such as for example EDCI and HOBt or a functional equivalent thereof, to yield a compound of formula (I).

In preparations presented above, the reaction products may be isolated from the reaction medium and, if necessary, further purified according to methodologies generally known in the art such as, for example, extraction, crystallization, distillation, trituration and chromatography.

The compounds of formula (I) as prepared in the hereinabove described processes may be synthesized as a mixture of stereoisomeric forms, in particular in the form of racemic mixtures of enantiomers which can be separated from one another following art-known resolution procedures. The racemic compounds of formula (I) may be converted into the corresponding diastereomeric salt forms by reaction with a suitable chiral acid. Said diastereomeric salt forms are subsequently separated, for example, by selective or fractional crystallization and the enantiomers are liberated therefrom by alkali. An alternative manner of separating the enantiomeric forms of the compounds of formula (I) involves liquid chromatography using a chiral stationary phase. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically. Preferably if a specific stereoisomer is desired, said compound will be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.

- 20 -

The compounds of the present invention can thus be used in animals, preferably in mammals, and in particular in humans as pharmaceuticals per se, in mixtures with one another or in the form of pharmaceutical preparations.

5 Furthermore, the present invention relates to pharmaceutical preparations which as active constituents contain an effective dose of at least one of the compounds of formula (I) in addition to customary pharmaceutically innocuous excipients and auxiliaries. The pharmaceutical preparations normally contain 0.1 to 90% by weight of the compound. The pharmaceutical preparations can be prepared in a manner known
10 per se to one of skill in the art. For this purpose, at least one of a compound of the present invention, together with one or more solid or liquid pharmaceutical excipients and/or auxiliaries and, if desired, in combination with other pharmaceutical active compounds, are brought into a suitable administration form or dosage form which can then be used as a pharmaceutical in human medicine or veterinary medicine.

15 Pharmaceuticals which contain a compound according to the invention can be administered orally, parenterally, e.g., intravenously, rectally, by inhalation, or topically, the preferred administration being dependent on the individual case, e.g., the particular course of the disorder to be treated. Oral administration is preferred.

20 The person skilled in the art is familiar on the basis of his expert knowledge with the auxiliaries which are suitable for the desired pharmaceutical formulation. Beside solvents, gel-forming agents, suppository bases, tablet auxiliaries and other active compound carriers, antioxidants, dispersants, emulsifiers, antifoams, flavor corrigents, preservatives, solubilizers, agents for achieving a depot effect, buffer substances or
25 colorants are also useful.

The compounds of the present invention may also find use in inhibiting ex vivo samples containing HIV or expected to be exposed to HIV. Hence, the present
30 compounds may be used to inhibit HIV present in a body fluid sample which contains or is suspected to contain or be exposed to HIV.

Also, the combination of an antiretroviral compound and a compound of the present invention can be used as a medicine. Thus, the present invention also relates to a
35 product containing (a) a compound of the present invention, and (b) another antiretroviral compound, as a combined preparation for simultaneous, separate or sequential use in treatment of retroviral infections. Thus, to combat or treat HIV infections, or the infection and disease associated with HIV infections, such as Acquired Immunodeficiency Syndrome (AIDS) or AIDS Related Complex (ARC), the

compounds of this invention may be co-administered in combination with for instance, binding inhibitors, such as, for example, dextran sulfate, suramine, polyanions, soluble CD4, gp120 binder such as BMS378806, anti CD4 Ab compounds such as PRO-542 or TNX-355; fusion inhibitors, such as, for example, T20, T1249, co-receptor binding inhibitors, such as, for example, AK-602, SCH-C, SCH-D, AMD 3100 (Bicyclams), AMD-070, TAK 779, TAK 220, UK-427-857, PRO-140; RT inhibitors, such as, for example, foscarnet and prodrugs; nucleoside RTIs, such as, for example, AZT, 3TC, DDC, DDI, D4T, Abacavir, FTC, Emtricitabine, DAPD, dOTC; nucleotide RTIs, such as, for example, PMEA, PMPA, tenofovir; NNRTIs, such as, for example, nevirapine, delavirdine, efavirenz, zalcitabine, zidovudine, zalcitabine, zalcitabine, R278474, MKC-442, UC 781, Capravirine, DPC 961, DPC963, DPC082, DPC083, calanolide A, SJ-3366, TSAO, 4"-deaminated TSAO; RNase H inhibitors, such as, for example, SP1093V, PD126338; TAT inhibitors, such as, for example, RO-5-3335, K12, K37; integrase inhibitors, such as, for example, L 708906, L 731988; protease inhibitors, such as, for example, darunavir, amprenavir, ritonavir, nelfinavir, saquinavir, indinavir, lopinavir, lasinavir, atazanavir, BMS 186316, DPC 681, DPC 684, tipranavir, AG1776, DMP 450, L 756425, PD178390, PNU 140135; glycosylation inhibitors, such as, for example, castanospermine, deoxynojirimycin.

20 Of particular interest are products comprising (i) a compound of formula (I), and (ii) darunavir, and (iii) ritanovir, as a combined preparation for simultaneous, separate or sequential use in treatment of retroviral infections.

Of particular interest are products comprising (i) a compound of formula (I), and (ii)
25 etravirine, as a combined preparation for simultaneous, separate or sequential use in
treatment of retroviral infections.

Of particular interest are products comprising (i) a compound of formula (I), and (ii) dapivirine, as a combined preparation for simultaneous, separate or sequential use in
30 treatment of retroviral infections.

Of particular interest are products comprising (i) a compound of formula (I), and (ii) R278474, as a combined preparation for simultaneous, separate or sequential use in treatment of retroviral infections.

The compounds of the present invention may also be administered in combination with immunomodulators (e.g., bropirimine, anti-human alpha interferon antibody, IL-2, methionine enkephalin, interferon alpha, and naltrexone) or with antibiotics (e.g.,

- 22 -

pentamidine isothiorate) to ameliorate, combat, or eliminate HIV infection and its symptoms.

For an oral administration form, compounds of the present invention are mixed with
5 suitable additives, such as excipients, stabilizers or inert diluents, and brought by means
of the customary methods into the suitable administration forms, such as tablets, coated
tablets, hard capsules, aqueous, alcoholic, or oily solutions. Examples of suitable inert
carriers are gum arabic, magnesia, magnesium carbonate, potassium phosphate, lactose,
glucose, or starch, in particular, corn starch. In this case the preparation can be carried
10 out both as dry and as moist granules. Suitable oily excipients or solvents are vegetable
or animal oils, such as sunflower oil or cod liver oil. Suitable solvents for aqueous or
alcoholic solutions are water, ethanol, sugar solutions, or mixtures thereof.
Polyethylene glycols and polypropylene glycols are also useful as further auxiliaries for
other administration forms.

15 For subcutaneous or intravenous administration, the active compounds, if desired with
the substances customary therefor such as solubilizers, emulsifiers or further
auxiliaries, are brought into solution, suspension, or emulsion. The compounds can
also be lyophilized and the lyophilizates obtained used, for example, for the production
20 of injection or infusion preparations. Suitable solvents are, for example, water,
physiological saline solution or alcohols, e.g. ethanol, propanol, glycerol, in addition
also sugar solutions such as glucose or mannitol solutions, or alternatively mixtures of
the various solvents mentioned.

25 Suitable pharmaceutical formulations for administration in the form of aerosols or
sprays are, for example, solutions, suspensions or emulsions of the compounds or their
physiologically tolerable salts in a pharmaceutically acceptable solvent, such as ethanol
or water, or a mixture of such solvents. If required, the formulation can also
30 additionally contain other pharmaceutical auxiliaries such as surfactants, emulsifiers
and stabilizers as well as a propellant. Such a preparation customarily contains the
active compound in a concentration from approximately 0.1 to 50%, in particular from
approximately 0.3 to 3% by weight.

35 In order to enhance the solubility and/or the stability of the compounds in
pharmaceutical compositions, it can be advantageous to employ α -, β - or γ -cyclo-
dextrins or their derivatives. Also co-solvents such as alcohols may improve the
solubility and/or the stability of the compounds in pharmaceutical compositions. In the

- 23 -

preparation of aqueous compositions, addition salts of the subject compounds are obviously more suitable due to their increased water solubility.

Appropriate cyclodextrins are α -, β - or γ -cyclodextrins (CDs) or ethers and mixed
5 ethers thereof wherein one or more of the hydroxy groups of the anhydroglucose units of the cyclodextrin are substituted with alkyl, particularly methyl, ethyl or isopropyl, e.g. randomly methylated β -CD; hydroxyalkyl, particularly hydroxyethyl, hydroxypropyl or hydroxybutyl; carboxyalkyl, particularly carboxymethyl or carboxyethyl; alkylcarbonyl, particularly acetyl; alkyloxycarbonylalkyl or carboxyalkyloxyalkyl,
10 particularly carboxymethoxypropyl or carboxyethoxypropyl; alkylcarbonyloxyalkyl, particularly 2-acetyloxypropyl. Especially noteworthy as complexants and/or solubilizers are β -CD, randomly methylated β -CD, 2,6-dimethyl- β -CD, 2-hydroxyethyl- β -CD, 2-hydroxyethyl- γ -CD, 2-hydroxypropyl- γ -CD and (2-carboxymethoxy)-propyl- β -CD, and in particular 2-hydroxypropyl- β -CD (2-HP- β -CD).

15 The term mixed ether denotes cyclodextrin derivatives wherein at least two cyclodextrin hydroxy groups are etherified with different groups such as, for example, hydroxy-propyl and hydroxyethyl.

20 An interesting way of formulating the present compounds in combination with a cyclodextrin or a derivative thereof has been described in EP-A-721,331. Although the formulations described therein are with antifungal active ingredients, they are equally interesting for formulating the compounds of the present invention. The formulations described therein are particularly suitable for oral administration and comprise an
25 antifungal as active ingredient, a sufficient amount of a cyclodextrin or a derivative thereof as a solubilizer, an aqueous acidic medium as bulk liquid carrier and an alcoholic co-solvent that greatly simplifies the preparation of the composition. Said formulations may also be rendered more palatable by adding pharmaceutically acceptable sweeteners and/or flavors.

30 Other convenient ways to enhance the solubility of the compounds of the present invention in pharmaceutical compositions are described in WO-94/05263, PCT application No. PCT/EP98/01773, EP-A-499299 and WO 97/44014, all incorporated herein by reference.

35 More in particular, the present compounds may be formulated in a pharmaceutical composition comprising a therapeutically effective amount of particles consisting of a solid dispersion comprising (a) a compound of the present invention, and (b) one or more pharmaceutically acceptable water-soluble polymers.

- 24 -

The term "a solid dispersion" defines a system in a solid state (as opposed to a liquid or gaseous state) comprising at least two components, wherein one component is dispersed more or less evenly throughout the other component or components. When
5 said dispersion of the components is such that the system is chemically and physically uniform or homogenous throughout or consists of one phase as defined in thermodynamics, such a solid dispersion is referred to as "a solid solution". Solid solutions are preferred physical systems because the components therein are usually readily bioavailable to the organisms to which they are administered.

10

The term "a solid dispersion" also comprises dispersions which are less homogenous throughout than solid solutions. Such dispersions are not chemically and physically uniform throughout or comprise more than one phase.

15 The water-soluble polymer in the particles is conveniently a polymer that has an apparent viscosity of 1 to 100 mPa.s when dissolved in a 2 % aqueous solution at 20°C solution.

Preferred water-soluble polymers are hydroxypropyl methylcelluloses or HPMC.
20 HPMC having a methoxy degree of substitution from about 0.8 to about 2.5 and a hydroxypropyl molar substitution from about 0.05 to about 3.0 are generally water soluble. Methoxy degree of substitution refers to the average number of methyl ether groups present per anhydroglucose unit of the cellulose molecule. Hydroxy-propyl molar substitution refers to the average number of moles of propylene oxide which
25 have reacted with each anhydroglucose unit of the cellulose molecule.

The particles as defined hereinabove can be prepared by first preparing a solid dispersion of the components, and then optionally grinding or milling that dispersion. Various techniques exist for preparing solid dispersions including melt-extrusion,
30 spray-drying and solution-evaporation.

It may further be convenient to formulate the present compounds in the form of nanoparticles which have a surface modifier adsorbed on the surface thereof in an amount sufficient to maintain an effective average particle size of less than 1000 nm.
35 Useful surface modifiers are believed to include those which physically adhere to the surface of the antiretroviral agent but do not chemically bond to the antiretroviral agent.

Suitable surface modifiers can preferably be selected from known organic and inorganic pharmaceutical excipients. Such excipients include various polymers, low

- 25 -

molecular weight oligomers, natural products and surfactants. Preferred surface modifiers include nonionic and anionic surfactants.

Yet another interesting way of formulating the present compounds involves a pharmaceutical composition whereby the compounds of this invention are incorporated in hydrophilic polymers and applying this mixture as a coat film over many small beads, thus yielding a composition with good bioavailability which can conveniently be manufactured and which is suitable for preparing pharmaceutical dosage forms for oral administration.

10

Said beads comprise (a) a central, rounded or spherical core, (b) a coating film of a hydrophilic polymer and an antiretroviral agent and (c) a seal-coating polymer layer.

15

Materials suitable for use as cores in the beads are manifold, provided that said materials are pharmaceutically acceptable and have appropriate dimensions and firmness. Examples of such materials are polymers, inorganic substances, organic substances, and saccharides and derivatives thereof.

20

Another aspect of the present invention concerns a kit or container comprising a compound of the present invention, in an amount effective for use as a standard or reagent in a test or assay for determining the ability of a potential pharmaceutical to inhibit HIV entry, HIV growth, or both. This aspect of the invention may find its use in pharmaceutical research programs.

25

The dose of the present compounds or of the physiologically tolerable salt(s) thereof to be administered depends on the individual case and, as customary, is to be adapted to the conditions of the individual case for an optimum effect. Thus it depends, of course, on the frequency of administration and on the potency and duration of action of the compounds employed in each case for therapy or prophylaxis, but also on the nature and severity of the infection and symptoms, and on the sex, age, weight and individual responsiveness of the human or animal to be treated and on whether the therapy is acute or prophylactic. Customarily, the daily dose of a compound of the present invention, in the case of administration to a patient approximately 75kg in weight ranges between 1mg and 5g, preferably between 10mg and 2 g, more preferably between 20 mg and 1g.

30

The dose can be administered in the form of an individual dose, or divided into several, e.g. two, three, four or even more, individual doses.

Experimental Part

Example 1 : synthesis of 4-[5-(4-cyano-benzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-cyclopentyl-benzamide according to Scheme 1 (compound 44)

Synthesis of 4-Acetyl-N-cyclopentyl-benzamide 1.2.a

- 5 A mixture of 4-acetyl benzoic acid **1.1.a** (10 mmol), cyclopentyl amine (11 mmol), EDCI (11 mmol) and HOBt (2 mmol) in CH₂Cl₂ (100 ml) was stirred at room temperature for 20 hours. Water was added to the resulting reaction mixture and the organic layer was separated. The water layer was extracted three times with dichloromethane. The combined organic layers were dried over MgSO₄, filtered and
10 the solvent was evaporated till dryness. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/methanol : 95/5). The pure fractions were collected and the solvent was evaporated yielding 90% of **1.2.a**.

Synthesis of 2-(2-methoxy-ethoxymethoxy)-benzaldehyde 1.3.a

- Intermediate **2.1** (20 mmol) was dissolved in *N,N*-dimethylformamide (150 ml). The
15 resulting mixture was cooled to 0°C. Sodium hydride (30 mmol) was added and the mixture was stirred at room temperature for 1 hour and chloromethoxymethoxy ethane (20 mol) was added. The mixture was stirred at room temperature for 16 hours. Water (200 ml) was added and the mixture was extracted with dichloromethane. The organic layer was separated, dried (over MgSO₄), filtered and the solvent was evaporated till
20 dryness. The residue was filtered on silica gel using dichloromethane as eluent. The solvent was removed to yield 95% of compound **1.3.a**.

Synthesis of N-cyclopentyl-4-{3-[2-(2-methoxy-ethoxymethoxy)-phenyl]-acryloyl}-benzamide 1.4.a

- Intermediate **1.2.a** (5 mmol) was dissolved in ethanol (100 ml) and then intermediate
25 **1.3.a** (5 mmol) was added. The mixture was stirred at room temperature and a solution of potassium hydroxide (10 mmol) in water (50 ml) was added dropwise. The mixture was warmed to 80°C for 12 hours. After cooling to room temperature a solution of ammonium chloride was added (50 ml) and ethanol was removed and replaced by ethyl acetate. The mixture was extracted with ethyl acetate and dried with magnesium
30 sulphate. Removal of the solvent and purification on silica gel using CH₂Cl₂ and methanol as eluent yielded product **1.4.a** (70%) as an oil.

Synthesis of N-Cyclopentyl-4-[3-(2-hydroxy-phenyl)-acryloyl]-benzamide 1.5.a

- Intermediate **1.4.a** (5 mmol) was dissolved in tetrahydrofuran (100 ml) and a solution
35 of hydrochloric acid (2 molar) was added dropwise. The mixture was stirred at room temperature overnight. Water was added (100 ml) and the mixture was extracted with ethyl acetate. The organic layer was washed with brine, dried on magnesium sulphate

- 27 -

and the solvent was evaporated till dryness yielding intermediate **1.5.a** (80 %) as a powder.

Synthesis of N-Cyclopentyl-4-[5-(2-hydroxy-phenyl)-4,5-dihydro-1H-pyrazol-3-yl]-benzamide **1.6.a**

- 5 Intermediate **1.5.a** (4 mmol) was dissolved in ethanol (100 ml). Hydrazine (4.4 mmol) was added dropwise to the reaction mixture. The mixture was then stirred at 70 °C for 4 hours. After cooling to room temperature the mixture was filtered, yielding intermediate **1.6.a** as white powder (90%).

Synthesis of 4-(2-oxo-acetyl)-benzonitrile **1.7.a**

- 10 To a mixture of dioxane (75 ml) and water (7.5 ml), selenium oxide (100 mmol) was added. The mixture was stirred at 50°C until the selenium oxide was dissolved. Then para-cyano-acetophenone (100 mmol) was added. The mixture was stirred at 90°C for 12 hours. The warm solution was filtered and the solvent was removed. The residue obtained was dissolved in dichloromethane and water was added. The organic layer
15 was separated, dried on magnesium sulphate and the solvent was removed, yielding intermediate **1.7.a** as a solid (85%).

Synthesis of compound 44

- Intermediate **1.6.a** (4 mmol) was dissolved in toluene (50 ml), then intermediate **1.7.a** (4.4 mmol) and para-toluene sulfonic acid (5%) were added. The reaction mixture was
20 refluxed for 8 hours. After cooling to room temperature, water was added and the mixture was extracted with ethyl acetate. The organic layer was washed with brine, dried on magnesium sulphate, filtered and the solvent was evaporated till dryness. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/methanol : 95/5). The product fractions were collected and the solvent was evaporated,
25 yielding 85% of compound 44.

Example 2 : synthesis of 4-[5-(4-cyano-benzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-cyclopentyl-benzamide according to Scheme 4 (compound 44)

Synthesis of 4-[3-(2-Hydroxy-phenyl)-acryloyl]-benzoic acid **4.3.a**

- 30 **Method A** 4-acetyl benzoic acid **4.1.a** (10 mmol) was dissolved in ethanol (100 ml) and then salicylaldehyde **4.2** (10 mmol) was added. The mixture was stirred at room temperature and a solution of potassium hydroxide (50 mmol) in water (30 ml) was added dropwise. The mixture was warmed to 80°C for 12 hours. After cooling to room temperature water (100 ml) was added and ethanol was removed and replaced
35 by ethyl acetate. The mixture was acidified until pH 4 by adding aqueous hydrochloric acid. The reaction mixture was then extracted with ethyl acetate and

- 28 -

the organic layer was dried with magnesium sulphate. The solvent was evaporated and the residue was purified by column chromatography on silica gel (eluent : CH_2Cl_2 /methanol : 80/20), yielding intermediate **4.3.a** as a white powder (20%).


Method B 4-acetyl benzoic acid **4.1.a** (10 mmol) was dissolved in tetrahydrofuran (100 ml) and then salicylaldehyde **4.2** (10 mmol) was added. The mixture was stirred at room temperature and pyrrolidine (20 mmol) was added drop wise. The mixture was warmed to 80°C for 12 hours. After cooling to room temperature, water (100 ml) and ethyl acetate (100 ml) were added. The mixture was extracted with ethyl acetate and dried with magnesium sulphate. The solvent was evaporated and the residue was purified by column chromatography on silica gel (eluent : CH_2Cl_2 /MeOH : 80/20), yielding intermediate **4.3.a** as a white powder (35%).

Synthesis of 4-[5-(2-Hydroxy-phenyl)-4,5-dihydro-1H-pyrazol-3-yl]-benzoic acid **4.4.a**
Intermediate **4.3.a** (33 mmol) was dissolved in ethanol (150 ml) and hydrazine (134 mmol) was added dropwise. The mixture was stirred at 70 °C for 4 hours. After cooling to room temperature the mixture was filtered, yielding intermediate **4.4.a** as white powder (90%).

Synthesis of 4-[5-(4-cyano-benzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-benzoic acid **4.5.a**
Intermediate **4.4.a** (5 mmol) was dissolved in toluene (50 ml), intermediate **1.7.a** (6 mmol) and para-toluene sulfonic acid (5%) were then added. The mixture was refluxed for 8 hours. After cooling to room temperature, water was added and the mixture was extracted with ethyl acetate. The organic layer was dried on magnesium sulphate, filtered and the solvent was removed. The residue was purified on silica gel using CH_2Cl_2 and methanol as eluent, yielding intermediate **4.5.a** as a powder (81%).


















Synthesis of compound 44
The carboxylic acid **4.5.a** (2.4 mmol) was dissolved in tetrahydrofuran (20 ml). EDCI (2.64 mmol) and HOBt (10%) were then added. The mixture was stirred 10 minutes at room temperature and cyclopentyl amine (2.64 mmol) was added. The mixture was then stirred at room temperature overnight. Water was added and the mixture was extracted with ethyl acetate, dried with magnesium sulphate and the solvent was removed. The crude residue was purified by column chromatography on silica gel using dichloromethane and methanol as eluent, yielding compound 44 as a powder (82%).

The compounds listed in the following tables were prepared analogous to the procedures described in examples 1 or 2.

















Table 1

Compound	L	R ³
1		
2		
3		
4		
5		
6		
7		












- 30 -

Compound	L	R ³
8		
9		
10		
11		
12		
13		
14		
15		
16		
















- 31 -

Compound	L	R ³
17		
18		
19		
20		
21		
22		
23		
24		
25		

- 32 -

Compound	L	R ³
26		
27		
28		
29		
30		
31		
32		
33		

- 33 -

Compound	L	R ³
34		
35		
36		
37		
38		
39		
40		
41		

- 34 -

Compound	L	R ³
42		
43		
44		
45		
46		
47		
48		
49		
50		
51		

- 35 -















Compound	L	R ³
52		
53		
54		
55		
56		
57		

Table 2

Compound	Q
58	
59	

- 36 -

Example 3 : Virological properties of the compounds of the present invention

The compounds were tested in a cellular assay using the MT4-LTR-EGFP cells for anti-viral activity. The assay demonstrated that these compounds exhibit potent anti-HIV activity against a wild type laboratory HIV strain (HIV-1 strain LAI) and a multi-
 5 drug resistant HIV-1 virus (HIV-1-MDR). The cellular assay was performed according to the following procedure.

HIV- or mock-infected MT4-LTR-EGFP cells were incubated for three days in the presence of various concentrations of the inhibitor. Upon infection, the GFP reporter is
 10 activated by the viral tat protein. At the end of the incubation period, the GFP signal was measured. In the virus control samples (in the absence of any inhibitor) the maximal fluorescent signal was obtained. The inhibitory activity of the compound was monitored on the virus-infected cells and was expressed as EC₅₀ and EC₉₀. These values represent the amount of the compound required to protect 50% and 90%,
 15 respectively, of the cells from virus infection. The toxicity of the compound was measured on the mock-infected cells and was expressed as CC₅₀, which represents the concentration of compound required to inhibit the growth of the cells by 50%. The selectivity index (SI) (ratio CC₅₀/EC₅₀) is an indication of the selectivity of the anti-HIV activity of the inhibitor.

20

Table 3

Compounds having a pEC₅₀ value (defined as the negative logarithm of the result expressed as EC₅₀) higher than 6 belong to the activity class A. Compounds having a
 25 pEC₅₀ value of higher than or equal to 5 but lower than or equal to 6 belong to the activity class B. Compounds having a pEC₅₀ value lower than 5 belong to the activity class C. For some of the compounds of the present invention, more than one test run was performed. In such a case, the average EC₅₀ value was used to determine the activity class. An empty cell in table 3 means that the data is not available.

Compound Number	Activity class for HIV-1-LAI test	Activity class for HIV-1-MDR test
1	B	C
2	C	C
3	C	C
4	B	
5	A	C
6	A	C
7	B	C

- 37 -

Compound Number	Activity class for HIV-1-LAI test	Activity class for HIV-1-MDR test
8		C
9	A	C
10	C	C
11	C	C
12	B	C
13	B	C
14	C	C
15	C	C
16	C	C
17	A	C
18	A	
19	B	C
20	B	B
21	C	C
22	A	C
23	C	C
24	B	B
25	B	B
26	B	B
27	A	A
28	A	A
29	B	A
30	C	C
31	B	B
32	C	B
33	A	A
34	B	C
35	A	C
36	A	
37	B	C
38	A	B
39	A	B
40	B	C
41	B	C
42	B	C

- 38 -

Compound Number	Activity class for HIV-1-LAI test	Activity class for HIV-1-MDR test
43	B	C
44	A	A
45	A	A
46	A	A
47	A	A
48	B	B
49	C	
50	A	A
51		A
52	A	A
53	A	A
54	B	B
55	A	A
58	A	A

Example 4 : Time of Addition experiment

- MT4-LTR-EGFP or MT4-LTR-Luc cells were infected at a high multiplicity of infection (MOI) by centrifugation for 10 minutes at 1200 g. Unbound virus was removed by two washing steps at 4°C in order to synchronize the infection. From 30 minutes post infection on, the test compound was added to parallel cultures in microtiter plates at different times. In case of MT4-LTR-EGFP cells, the cultures were scored microscopically for fluorescence 24 hours after infection and supernatant was collected. HIV replication in the supernatant samples was quantified by measuring the concentration of the p24 viral antigen using a commercial kit, according to the manufacturer protocol (NEN). Because of the high MOI used in this type of experiments, concentrations of inhibitors were at least 100 fold higher than their EC₅₀ value in the cellular antiviral assay. In case of MT4-LTR-Luc cells, HIV inhibitors were added at EC90 concentration at different time intervals to HIV-1 infected MT4 cells expressing HIV-1-LTR-luciferase reporter gene. HIV-1 infection was revealed by luciferase detection. 24 Hours post incubation, the luciferase substrate was added and the chemiluminescence signal was measured. At the time points where compounds showed activity, the luciferase signal was reduced.
- The compounds of the present invention that were tested inhibited viral replication earlier than three hours after synchronisation.

*Example 5 : pharmaceutical compositions*Capsules

Active ingredient, *in casu* a compound of formula (I), is dissolved in organic solvent such as ethanol, methanol or methylene chloride, preferably, a mixture of ethanol and methylene chloride. Polymers such as polyvinylpyrrolidone copolymer with vinyl acetate (PVP-VA) or hydroxypropylmethylcellulose (HPMC), typically 5 mPa.s, are dissolved in organic solvents such as ethanol, methanol methylene chloride. Suitably the polymer is dissolved in ethanol. The polymer and compound solutions are mixed and subsequently spray dried. The ratio of compound/polymer was selected from 1/1 to 1/6. Intermediate ranges are 1/1.5 and 1/3. A suitable ratio is 1/6. The spray-dried powder, a solid dispersion, is subsequently filled in capsules for administration. The drug load in one capsule ranges between 50 and 100 mg depending on the capsule size used.

15 Film-coated Tablets*Preparation of Tablet Core*


A mixture of 100 g of active ingredient, *in casu* a compound of formula (I), 570 g lactose and 200 g starch is mixed well and thereafter humidified with a solution of 5 g sodium dodecyl sulfate and 10 g polyvinylpyrrolidone in about 200 ml of water. The wet powder mixture is sieved, dried and sieved again. Then there was added 100 g microcrystalline cellulose and 15 g hydrogenated vegetable oil. The whole is mixed well and compressed into tablets, giving 10.000 tablets, each comprising 10 mg of the active ingredient.

Coating

25 To a solution of 10 g methylcellulose in 75 ml of denaturated ethanol there is added a solution of 5 g of ethylcellulose in 150 ml of dichloromethane. Then there are added 75 ml of dichloromethane and 2.5 ml 1,2,3-propanetriol. 10 g of polyethylene glycol is molten and dissolved in 75 ml of dichloromethane. The latter solution is added to the former and then there are added 2.5 g of magnesium octadecanoate, 5 g of polyvinylpyrrolidone and 30 ml of concentrated color suspension and the whole is homogenated. The tablet cores are coated with the thus obtained mixture in a coating apparatus.

Claims

1. A compound having the formula



a N-oxide, stereoisomeric form or salt thereof,

5 wherein

a is zero, 1, 2, 3, 4 or 5;

10 R¹ is hydrogen, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, C₃₋₁₂cycloalkyl, Het, aryl or C₁₋₁₀alkyl substituted with a substituent selected from the group consisting of Het, aryl, C₃₋₁₂cycloalkyl, amino and mono-or disubstituted amino wherein the substituents on the amino group are each individually selected from C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₃₋₁₂cycloalkyl, Het and aryl;

15 R² is hydrogen, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, C₃₋₁₂cycloalkyl, Het, aryl or C₁₋₁₀alkyl substituted with a substituent selected from the group consisting of Het, aryl, C₃₋₁₂cycloalkyl, amino and mono-or disubstituted amino wherein the substituents on the amino group are each individually selected from C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₃₋₁₂cycloalkyl, Het and aryl; or

20 R¹ and R² taken together with the nitrogen atom to which they are attached form a 5 to 12 membered saturated or partially saturated heterocycle which heterocycle may optionally be substituted with C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, C₃₋₁₂cycloalkyl, C₁₋₆alkyloxycarbonyl, Het, aryl or C₁₋₁₀alkyl substituted with a substituent selected from the group consisting of Het, aryl, C₃₋₁₂cycloalkyl, amino and mono-or disubstituted amino wherein the substituents on the amino group are each individually selected from C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₃₋₁₂cycloalkyl, Het and aryl;

25 R³ is carboxyl, halogen, nitro, C₁₋₁₀alkyl, C₃₋₁₂cycloalkyl, polyhaloC₁₋₁₀alkyl, cyano, amino, mono-or disubstituted amino, aminocarbonyl, mono-or disubstituted aminocarbonyl, C₁₋₁₀alkyloxy, C₁₋₁₀alkylthio, C₁₋₁₀alkylsulfonyl, pyrrolidinyl, piperidinyl, homopiperidinyl, morpholinyl, thiomorpholinyl or piperazinyl optionally substituted with C₁₋₁₀alkyl, wherein the substituents on any of the

- 41 -

amino groups are each individually selected from C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₃₋₁₂cycloalkyl, Het and aryl;

aryl is phenyl optionally substituted with one or more substituents selected from the group consisting of C₁₋₁₀alkyl, polyhaloC₁₋₁₀alkyl, C₁₋₁₀alkyloxy, C₁₋₁₀alkylthio, C₁₋₁₀alkylsulfonyl, nitro, cyano, halo, C₃₋₇cycloalkyl, C₁₋₁₀alkylcarbonyl, carboxyl, C₁₋₁₀alkyloxycarbonyl, amino, mono-or disubstituted amino, aminocarbonyl, mono-or disubstituted aminocarbonyl, wherein the substituents on any of the amino groups are each individually selected from phenyl, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₃₋₇cycloalkyl, pyrrolidinyl, piperidinyl, homopiperidinyl or piperazinyl optionally substituted with C₁₋₁₀alkyl;

Het is a 5 or 6 membered aromatic, saturated or partially saturated, monocyclic or bicyclic heterocycle with one or more heteroatoms each individually selected from nitrogen, oxygen or sulfur, and which heterocycle may optionally be substituted by one, or where possible, more than one substituent selected from the group consisting of C₁₋₁₀alkyl, polyhaloC₁₋₁₀alkyl, C₁₋₁₀alkyloxy, C₁₋₁₀alkylthio, C₁₋₁₀alkylsulfonyl, nitro, cyano, halo, C₃₋₇cycloalkyl, C₁₋₁₀alkylcarbonyl, carboxyl, C₁₋₁₀alkyloxycarbonyl, amino, mono-or disubstituted amino, aminocarbonyl, mono-or disubstituted aminocarbonyl, wherein the substituents on any of the amino groups are each individually selected from phenyl, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₃₋₇cycloalkyl, pyrrolidinyl, piperidinyl, homopiperidinyl or piperazinyl optionally substituted with C₁₋₁₀alkyl.


2. A compound as claimed in claim 1 wherein a is zero or 1.

3. A compound as claimed in claim 1 or 2 wherein R¹ is C₁₋₁₀alkyl, C₃₋₁₂cycloalkyl, Het, aryl or C₁₋₁₀alkyl substituted with a substituent selected from the group consisting of Het, aryl, C₃₋₁₂cycloalkyl and mono-or disubstituted amino wherein the substituents on the amino group are each individually selected from C₁₋₁₀alkyl; R² is hydrogen, C₁₋₁₀alkyl, C₂₋₁₀alkenyl or C₁₋₁₀alkyl substituted with aryl; or R¹ and R² taken together with the nitrogen atom to which they are attached form a 5 to 12 membered saturated or partially saturated heterocycle which heterocycle may optionally be substituted with C₁₋₁₀alkyl, C₁₋₆alkyloxycarbonyl or C₁₋₁₀alkyl substituted with aryl.

4. A compound as claimed in any one of claims 1 to 3 wherein R³ is halogen, nitro, C₁₋₁₀alkyl, C₃₋₁₂cycloalkyl, polyhaloC₁₋₁₀alkyl, cyano, C₁₋₁₀alkyloxy, C₁₋₁₀alkylthio, C₁₋₁₀alkylsulfonyl, piperidinyl, morpholinyl.

- 42 -

5. A compound as claimed in any one of claims 1 to 4 wherein aryl is phenyl optionally substituted with one or more substituents selected from the group consisting of C₁₋₁₀alkyloxy, halo, mono-or disubstituted amino, wherein the substituents on any of the amino groups are each individually selected from C₁₋₁₀alkyl.
6. A compound as claimed in any one of claims 1 to 5 wherein Het is a 5 or 6 membered aromatic, saturated or partially saturated, monocyclic or bicyclic heterocycle with one or more heteroatoms each individually selected from nitrogen, oxygen or sulfur, and which heterocycle may optionally be substituted by C₁₋₁₀alkyl.
7. A compound as claimed in any one of claims 1 to 6 wherein Het is a heterocycle selected from the group consisting of pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, pyranyl, pyrrolyl, furanyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, tetrahydrofuranyl, tetrahydrothienyl, imidazolinyl, dioxolanyl, pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, homopiperazinyl, dioxanyl, morpholinyl, thiomorpholinyl, which heterocycle may optionally be substituted by one, or where possible, more than one substituent selected from the group consisting of C₁₋₆alkyl, polyhaloC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkylsulfonyl, nitro, cyano, halo, C₃₋₇cycloalkyl, C₁₋₆alkyl-carbonyl, carboxyl, C₁₋₆alkyloxycarbonyl, amino, mono-or disubstituted amino, aminocarbonyl, mono-or disubstituted aminocarbonyl, wherein the substituents on any of the amino groups are each individually selected from phenyl, C₁₋₆alkyl, C₂₋₆alkenyl, C₃₋₇cycloalkyl, pyrrolidinyl, piperidinyl, homopiperidinyl or piperazinyl optionally substituted with C₁₋₆alkyl.
8. A compound as claimed in any one of claims 1 to 7 wherein the compound has the formula (Ia)



- 43 -

9. A compound as claimed in any one of claims 1 to 8 wherein
a is zero or 1;
R¹ is C₁₋₁₀alkyl, C₃₋₁₂cycloalkyl, Het, aryl or C₁₋₁₀alkyl substituted with a
substituent selected from the group consisting of Het, aryl, C₃₋₁₂cycloalkyl and
5 mono-or disubstituted amino wherein the substituents on the amino group are
each individually selected from C₁₋₁₀alkyl;
R² is hydrogen, C₁₋₁₀alkyl, C₂₋₁₀alkenyl or C₁₋₁₀alkyl substituted with aryl; or
R¹ and R² taken together with the nitrogen atom to which they are attached form a
5 to 12 membered saturated or partially saturated heterocycle which heterocycle
10 may optionally be substituted with C₁₋₁₀alkyl, C₁₋₆alkyloxycarbonyl or
C₁₋₁₀alkyl substituted with aryl;
R³ is halogen, nitro, C₁₋₁₀alkyl, C₃₋₁₂cycloalkyl, polyhaloC₁₋₁₀alkyl, cyano,
C₁₋₁₀alkyloxy, C₁₋₁₀alkylthio, C₁₋₁₀alkylsulfonyl, piperidinyl, morpholinyl.
10. A compound as claimed in any one of claims 1 to 9 wherein
15 a is zero or 1;
R¹ is C₁₋₁₀alkyl, C₃₋₁₂cycloalkyl, Het, aryl or C₁₋₁₀alkyl substituted with a
substituent selected from the group consisting of Het, aryl, C₃₋₁₂cycloalkyl and
mono-or disubstituted amino wherein the substituents on the amino group are
each individually selected from C₁₋₁₀alkyl;
20 R² is hydrogen, C₁₋₁₀alkyl, C₂₋₁₀alkenyl or C₁₋₁₀alkyl substituted with aryl; or
R¹ and R² taken together with the nitrogen atom to which they are attached form a 5
to 12 membered saturated or partially saturated heterocycle which heterocycle
may optionally be substituted with C₁₋₁₀alkyl, C₁₋₆alkyloxycarbonyl or
C₁₋₁₀alkyl substituted with aryl;
25 R³ is halogen, nitro, C₁₋₁₀alkyl, C₃₋₁₂cycloalkyl, polyhaloC₁₋₁₀alkyl, cyano,
C₁₋₁₀alkyloxy, C₁₋₁₀alkylthio, C₁₋₁₀alkylsulfonyl, piperidinyl, morpholinyl;
aryl is phenyl optionally substituted with one or more substituents selected from the
group consisting of C₁₋₁₀alkyloxy, halo, mono-or disubstituted amino, wherein
the substituents on any of the amino groups are each individually selected from
30 C₁₋₁₀alkyl;
Het is a 5 or 6 membered aromatic, saturated or partially saturated, monocyclic or
bicyclic heterocycle with one or more heteroatoms each individually selected
from nitrogen, oxygen or sulfur, and which heterocycle may optionally be
substituted by C₁₋₁₀alkyl.
- 35 11. A compound as claimed in any one of claims 1 to 10 wherein
a is zero or 1;

- 44 -

- R^1 is C_{1-6} alkyl, C_{3-7} cycloalkyl, Het, aryl or C_{1-6} alkyl substituted with a substituent selected from the group consisting of Het, aryl, C_{3-7} cycloalkyl and mono-or disubstituted amino wherein the substituents on the amino group are each individually selected from C_{1-6} alkyl;
- 5 R^2 is hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, or C_{1-6} alkyl substituted with a substituent selected from the group consisting of aryl or
- R^1 and R^2 taken together with the nitrogen atom to which they are attached form a heterocycle selected from the group consisting of piperazinyl, homopiperazinyl, piperidinyl, homopiperidinyl, morpholinyl, thiomorpholinyl, pyrrolidinyl,
- 10 indolinyl, 1,2,3,4-tetrahydroquinolinyl, 1,2,3,4-tetrahydro-isoquinolinyl, which heterocycle may optionally be substituted with C_{1-6} alkyl, C_{1-6} alkyloxycarbonyl or C_{1-6} alkyl substituted with aryl;
- R^3 is halogen, nitro, C_{1-6} alkyl, C_{3-7} cycloalkyl, polyhalo C_{1-4} alkyl, cyano, C_{1-6} alkyloxy, C_{1-6} alkylthio, C_{1-6} alkylsulfonyl, piperidinyl, morpholinyl;
- 15 aryl is phenyl optionally substituted with one, two or three substituents selected from the group consisting of C_{1-6} alkyl, C_{1-6} alkyloxy, nitro, cyano, halo, amino, mono-or disubstituted amino, wherein the substituents on any of the amino groups are each individually selected from C_{1-6} alkyl;
- Het is a heterocycle selected from the group consisting of pyridinyl, furanyl,
- 20 thienyl, tetrahydrofuranyl, pyrrolidinyl, piperidinyl, which heterocycle may optionally be substituted by C_{1-6} alkyl.
12. A compound as claimed in any one of claims 1 to 11 wherein the compound has the formula (Ia) and R^1 is cyclopentyl and R^2 is hydrogen.
13. A compound as claimed in any one of claims 1 to 12 wherein a is 1 and R^3 is
- 25 halogen, cyano, C_{1-4} alkyl, C_{1-4} alkylthio; morpholinyl, C_{1-4} alkyloxy, nitro, C_{1-4} alkylsulfonyl, trifluoromethyl.
14. A compound as claimed in claim 1 having the structure
- 4-[5-(4-fluorobenzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-[(4-methoxyphenyl)methyl]-N-methyl-benzamide;
- 30 4-[5-(4-cyanobenzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-cyclopentyl-benzamide;
- 4-[5-(4-propylbenzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-cyclopentyl-benzamide;
- 4-[5-(4-methylsulfanyl-benzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-
- 35 yl]-N-cyclopentyl-benzamide;

- 45 -

- 7 4-[5-(4-ethyl-benzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-cyclopentyl-benzamide;
4-[5-(4-tert-butyl-benzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-cyclopentyl-benzamide;
- 5 4-[5-(4-nitro-benzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-cyclopentyl-benzamide;
4-[5-(4-methanesulfonyl-benzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-cyclopentyl-benzamide;
4-[5-(4-trifluoromethyl-benzoyl)-1,10b-dihydro-5H-pyrazolo[1,5-c][1,3]benzoxazin-2-yl]-N-cyclopentyl-benzamide
10 or a N-oxide, stereoisomeric form or salt thereof.
15. A compound as claimed in any one of claims 1 to 14 for use as a medicine.
16. Use of a compound as claimed in any one of claims 1 to 14 in the manufacture of a medicament useful for treating conditions associated with HIV infection.
- 15 17. Use of a compound as claimed in any one of claims 1 to 14 in the manufacture of a medicament useful for preventing HIV transmission or preventing HIV infection or preventing a disease associated with HIV infection.
18. A method of treatment comprising the systemic administration of an effective therapeutic amount of a compound as claimed in any one of claims 1 to 14 to HIV-
20 infected warm-blooded animals, in particular, HIV-infected humans.
19. A pharmaceutical preparation which as active constituents contain an effective dose of a compound as claimed in any one of claims 1 to 14 in addition to customary pharmaceutically innocuous excipients and auxiliaries.
20. A product containing (a) a compound as claimed in any one of claims 1 to 14, and
25 (b) another antiretroviral compound, as a combined preparation for simultaneous, separate or sequential use in treatment of retroviral infections.