

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 199946784 B2**
(10) Patent No. **764474**

(54) Title
Use of optically pure (-) norcisapride in the treatment of apnea, bulimia, and other disorders

(51)⁶ International Patent Classification(s)
A61K 031/445

(21) Application No: **199946784** (22) Application Date: **1999.06.10**

(87) WIPO No: **WO99/65490**

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/089225	1998.06.15	US
60/122236	1999.03.01	US

(43) Publication Date : **2000.01.05**

(43) Publication Journal Date : **2000.03.16**

(44) Accepted Journal Date : **2003.08.21**

(71) Applicant(s)
Sepracor, Inc.

(72) Inventor(s)
Paul D. Rubin; Timothy J Barberich

(74) Agent/Attorney
PHILLIPS ORMONDE and FITZPATRICK, 367 Collins Street, MELBOURNE VIC 3000

(56) Related Art
WO 96/40133
US 5663173
WO 98/03173

46784/99



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/445		A2	(11) International Publication Number: WO 99/65490
		(43) International Publication Date: 23 December 1999 (23.12.99)	
(21) International Application Number: PCT/US99/13100		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 10 June 1999 (10.06.99)			
(30) Priority Data: 60/089,225 15 June 1998 (15.06.98) US 60/122,236 1 March 1999 (01.03.99) US			
(71) Applicant: SEPRACOR, INC. [US/US]; 111 Locke Drive, Marlborough, MA 01752 (US).			
(72) Inventors: RUBIN, Paul, D.; 37 Greystone Lane, Sudbury, MA 01776 (US). BARBERICH, Timothy, J.; 40 Elm Street, Concord, MA 01742 (US).			
(74) Agents: LAWRENCE, Stanton, T., III et al.; Pennie & Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US).		Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(54) Title: USE OF OPTICALLY PURE (-) NORCISAPRIDE IN THE TREATMENT OF APNEA, BULIMIA, AND OTHER DISORDERS			
(57) Abstract			
<p>Methods for the prevention, treatment, or management of apnea, apnea disorders, bulimia nervosa, irritable bowel syndrome, urinary incontinence, bradycardia, bradyarrhythmia, syncope, other disorders, or symptoms thereof using (-) norcispapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.</p>			

USE OF OPTICALLY PURE (-) NORCISAPRIDE IN THE TREATMENT OF APNEA, BULIMIA, AND OTHER DISORDERS

1. FIELD OF THE INVENTION

5 The invention relates to methods of prevention, treatment, or management, of apnea, apnea disorders, bulimia, other disorders, or symptoms thereof.

2. BACKGROUND OF THE INVENTION

10 Apnea is defined in *Stedman's Medical Dictionary*, 26th Edition, Williams and Wilkins (1995), as the absence of breathing. There are a number of disorders associated with apnea, which are characterized by interrupted breathing in which a person stops breathing long enough to decrease the amount of oxygen and increase the amount of carbon dioxide in the blood and brain. Each type of apnea involves the absence of airflow at the nose or the mouth, typically for at least 10 seconds.

15 Various apnea disorders exist, including: central apnea, which results from medullary depression and inhibits respiratory movement; deglutition apnea, which is the inhibition of breathing during swallowing; obstructive or peripheral apnea, which is either a result of obstruction of air passages or inadequate respiratory muscle activity; sleep apnea, which is central and/or obstructive apnea during sleep; and sleep induced apnea, which
20 results from failure of the respiratory center to stimulate adequate respiration during sleep.

 Obstructive apneas usually occur in obese men and are much less common in women. The obesity, perhaps in combination with aging body tissues and other factors, leads to narrowing of the upper airways. Tobacco smoking, excessive alcohol use, and lung diseases, such as emphysema, increase the risk of developing obstructive apneas.

25 For those suffering from sleep apnea, quitting smoking, avoiding excessive use of alcohol, and losing weight are commonly the first behavioral steps for treating the disorder. In order to inhibit or avoid apnea, heavy snorers and people who often choke in their sleep should not take tranquilizers, sleep aids, or other sedating drugs.

30 Sleep apnea is one of the most common forms of apnea. Rarely, a person who has severe sleep apnea needs a tracheostomy, a surgical procedure that creates a permanent opening into the windpipe through the neck. Sometimes other surgical procedures are performed to widen the upper airway and alleviate the problem. However, such extreme measures are seldom needed and never desired.

Apnea can also be treated by non-invasive means by administering to a patient a therapeutic agent. U.S. Patent No. 5,075,290 discloses the medical treatment of obstructive sleep apnea and associated symptoms, such as snoring, by the administration of the nucleoside uptake blocker, dipyridamole, during sleep. U.S. Patent Nos. 5,502,067 and 5,407,953 disclose a method of treating sleep apnea, hyponea and snoring in a human patient by administering a pilocarpine compound. U.S. Patent No. 5,422,374 discloses a method of treating sleep apnea by the administration of ubidecarenone to a patient. U.S. Patent No. 5,356,934 discloses a method of employing (R)-fluoxetine to treat sleep apnea.

Bulimia nervosa, or bulimia, is a disorder described in the *Diagnostic and Statistical Manual of Mental Disorders*, Fourth Edition, American Psychiatric Association, 1996 ("DSM-IV"), which is characterized in part by recurrent episodes of binge eating during which the patient experiences a loss of control resulting in over eating and self-induced vomiting. The disorder primarily afflicts females of upper and middle socioeconomic status, especially college-age women.

Currently, two approaches for treating bulimia are used: cognitive-behavioral and pharmacological. Traditional pharmacological treatments involved antidepressants. More recent research into the fundamental causes of bulimia, however, has suggested other pharmacological treatments. In particular, some researchers have argued that the pathophysiological characteristics driving the behaviors characteristic of bulimia involve an increase in the basal tone of the vagal nerve, and have suggested that racemic ondansetron may be useful for the treatment of bulimia. Faris, P. L. *et al.*, *Biol. Psychiatry*, 32:462-466 (1992); Dumuis *et al.*, *N.S. Arch. Pharmacol.*, 340:403-410 (1989).

Cisapride, chemically named *cis*-4-amino-5-chloro-N-[1-[3-(4-fluorophenoxy)propyl]-3-methoxy-4-piperidinyl]-2-methoxybenzamide, is a benzamide derivative, the parent compound being metoclopramide. Schapira *et al.*, *Acta Gastroenterolog. Belg.*, LIII:446-457 (1990). Benzamide derivatives have several prominent pharmacological actions due to their effects on neuronal systems modulated by the neurotransmitter serotonin.

Because of their modulation of the serotonin neuronal system in the gastrointestinal tract, many benzamide derivatives are effective antiemetic agents and are commonly used to control vomiting during cancer chemotherapy or radiotherapy. Costall *et al.*, *Neuropharmacology*, 26:1321-1326 (1987). This action is almost certainly the result of an ability to block serotonin at specific sites, particularly type-3 5-hydroxytryptamine

(5-HT₃) receptors. Clarke *et al.*, *Trends in Pharmacological Sciences*, 10:385-386 (1989).

In theory, chemotherapy and radiation therapy can induce nausea and vomiting by damaging enterochromaffin cells in the gastrointestinal tract. As a result, the

neurotransmitter serotonin is released and stimulates both afferent vagal nerve fibers (thus

- 5 initiating the vomiting reflex) and serotonin receptors in the chemoreceptor trigger zone of the area postrema region of the brain. The anatomical site for this action of the benzamide derivatives, and whether such action is central (CNS), peripheral, or a combination thereof, remains unresolved. Barnes *et al.*, *J. Pharm. Pharmacol.*, 40:586-588 (1988).

- Racemic cisapride is used primarily to treat gastroesophageal reflux disease
10 ("GERD"), which is characterized as the backward flow of the stomach contents into the esophagus. Cisapride is commercially available as the racemic mixture of the *cis* diastereomeric enantiomers of cisapride known as PROPULSID®.

- The co-administration of racemic cisapride with other therapeutic agents causes inhibitory problems with the metabolism of cisapride by the liver. For example,
15 ketoconazole has a pronounced effect on cisapride kinetics resulting from the inhibition of the metabolic elimination of cisapride and leads to an 8-fold increase of the steady-state plasma levels. *Physician's Desk Reference*®, 52nd Edition, Medical Economics Co., Inc., 1998. Interaction of racemic cisapride and other therapeutic agents can also potentiate cardiovascular side effects, such as cardiotoxicity. This potentiation occurs when other
20 drugs present in the patient's system interfere with the metabolism of cisapride, thereby building up racemic cisapride in the body.

- These interactions are a significant drawback to the use of racemic cisapride; in particular, because racemic cisapride is often used before, with or immediately after another therapeutic agent. In addition, administration of racemic cisapride to a human has
25 been found to cause adverse effects such as cardiac arrhythmia, including ventricular tachycardia, ventricular fibrillation, Q_T prolongation, and torsades de pointes, central nervous system ("CNS") effects, increased systolic pressure, interactions with other drugs, diarrhea, abdominal cramping and cardiac depression.

- Racemic cisapride is almost completely absorbed after oral administration to
30 humans, but bioavailability of cisapride is only 40-50% due to rapid first-pass metabolism in the liver. Van Peer *et al.*, in *Progress in the Treatment of Gastrointestinal Motility Disorders: The Role of Cisapride*, Proceedings of a Symposium in Frankfurt, Germany, November 1986, *Excerpta Medica*, A.G. Johnson and G Lux Eds., Amsterdam, pp. 23-29

(1988). More than 90% of a dose of racemic cisapride in humans is metabolized mainly by oxidative N-dealkylation of the piperidine nitrogen or by aromatic hydroxylation occurring on either the 4-fluorophenoxy or benzamide rings. Meuldermans *et al.*, *Drug Metab. Dispos.*, 16(3):410-419 (1988); and
5 Meuldermans *et al.*, *Drug Metab. Dispos.*, 16(3):403-409 (1988). Norcisapride, chemically named 4-amino-5-chloro-N-(3-methoxy-4-piperidinyl)-2-methoxybenzamide, is a metabolite of cisapride.

It is desirable to provide safe and effective methods of preventing, treating, or managing apnea, apnea disorders, bulimia nervosa and related
10 disorders, or symptoms thereof, particularly a treatment that allows the patient to undergo other related therapies without adverse effects or drug-drug interactions.

3. SUMMARY OF THE INVENTION

15 The present invention encompasses the use of the optically pure (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer, in preventing, treating, or managing apnea, apnea disorders, bulimia, irritable bowel syndrome, asthma, urinary incontinence, syncope, bradycardia, bradyarrhythmia, or symptoms thereof. It should be
20 understood that the invention encompasses any combination of preventing, treating, or managing each disorder or multiple disorders.

This invention also relates to pharmaceutical compositions adapted for the prevention, treatment, or management of a patient suffering from a vagal nerve mediated disorder or symptoms thereof, which comprises a
25 therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

This invention also relates to pharmaceutical compositions adapted for the prevention, treatment, or management of a patient suffering from apnea, apnea disorders, or symptoms thereof, which comprises a
30 therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

This invention further relates to pharmaceutical compositions adapted for the prevention, treatment, or management of bulimia, irritable bowel syndrome, asthma, urinary incontinence, bradycardia, bradyarrhythmia, syncope, related disorders, and symptoms thereof in a mammal, which comprises a therapeutically effective amount of (-)

5 norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer, said amount being sufficient to alleviate symptoms of said conditions while reducing or avoiding adverse effects associated with administration of racemic cisapride.

The invention also encompasses single unit dosage forms of optically pure (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer, which comprise from about 0.5 mg to about 500 mg of active ingredient in a compressed tablet. This dosage form is particularly suitable for the prevention, treatment, or management of apnea, apnea disorders, bulimia, irritable bowel syndrome, asthma, urinary incontinence, bradycardia, bradyarrhythmia, syncope, related disorders, or symptoms thereof.

4. DETAILED DESCRIPTION OF THE INVENTION

The present invention encompasses the use of optically pure (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer, in preventing, treating, or managing disorders, including, but not limited to, apnea, apnea disorders, bulimia, irritable bowel syndrome, asthma, urinary incontinence, bradycardia, bradyarrhythmia, syncope, and related disorders, or symptoms thereof. Apnea or apnea disorders include, but are not limited to, central apnea, deglutition apnea, obstructive or peripheral apnea, sleep apnea, and sleep induced apnea, or a combination thereof.

The present invention also encompasses the use of optically pure (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer, in preventing, treating, or managing apnea, apnea disorders, bulimia, irritable bowel syndrome, asthma, urinary incontinence, bradycardia, bradyarrhythmia, syncope, and related disorders, or symptoms thereof, preferably while reducing or avoiding adverse effects associated with administration of racemic cisapride.

30 In one embodiment, the present invention relates to a method of preventing, treating, or managing bulimia comprising administering to a patient a therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

In another embodiment, the present invention relates to a method of preventing, treating, or managing apnea or apnea disorders comprising administering to a patient a therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

5 In another embodiment, the present invention relates to a method for preventing, treating, or managing conditions mediated by vagal activity in a patient comprising administering a therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

The present invention also encompasses a method of preventing, treating, or
10 managing irritable bowel syndrome comprising administering to a patient a therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

In another embodiment, the present invention relates to methods of preventing, treating, or managing syncope, and in particular vasovagal syncope and cardiac
15 or carotid sinus syncope, which comprises administering to a patient a therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

The present invention further encompasses methods of preventing, treating, or managing bradycardia or bradyarrhythmia, which comprises administering to a patient a
20 therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

In another embodiment, the present invention relates to a method of preventing, treating, or managing asthma or asthma symptoms, which comprises administering to a patient a therapeutically effective amount of (-) norcisapride, or a
25 pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

The present invention also encompasses a method of preventing, treating, or managing urinary incontinence, which comprises administering to a patient a therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

30 In another embodiment, this invention encompasses single unit dosage forms of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer, which comprise from about 0.5 mg to about 500 mg of active ingredient in a compressed tablet. This dosage form is particularly suitable for the prevention, treatment,

or management of apnea, apnea disorders, bulimia, irritable bowel syndrome, asthma, urinary incontinence, bradycardia, bradyarrhythmia, syncope, related disorders, or symptoms thereof.

The vagus nerve is the largest nerve of the cranial nerves. There are two
5 main branches of the vagus nerve, each of which act to provide both motor and sensory functions. The vagus nerves contain efferent fibers, which carry impulses from the nerve's origin in the medulla oblongata of the brain to a tissue or visceral organ, and afferent fibers, which carry impulses from the organ back to the brain. It is present in a large portion of the body, extending from the brain stem to the organs of the neck, chest, and abdomen. Vagal
10 stimulation occurs in a number of organs, including the heart, lungs, bronchia, trachea, esophagus, stomach, pancreas, small intestine, large intestine, colon, liver, gall bladder, and portions of the urinary tract.

Without being limited by theory, it is believed that symptoms of bulimia, irritable bowel syndrome, urinary incontinence, bradycardia, bradyarrhythmia, asthma, and
15 syncope, particularly vasovagal syncope and cardiac or carotid sinus syncope, are affected by the basal tone of the vagus, or vagal, nerve.

Without being limited by theory, it is further believed that by blocking 5-HT-induced depolarization in the vagus, or vagal, nerve, (-) norcisapride lessens or inhibits symptoms of these disorders. Therefore, in one embodiment, the present invention relates
20 to the use of optically pure (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer, in preventing, treating, or managing bulimia, irritable bowel syndrome, asthma, urinary incontinence, bradycardia, bradyarrhythmia, syncope, and related disorders, or symptoms thereof.

Additionally, the invention includes the use of (-) norcisapride, or a
25 pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer, in combination with other therapeutic agents. Examples of other therapeutic agents include, but are not limited to, fluoxetine or the R or S stereoisomer thereof; descarboethoxyloratidine; ondansetron or the R or S stereoisomer thereof, preferably R ondansetron; ubidecarenone; dipyridamole; pilocarpine or the stereoisomers thereof;
30 primidone or the R or S stereoisomer thereof; orphenadrine citrate; and the like, as well as any active metabolites thereof. Administration of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer, in combination with these other therapeutic agents for the prevention, treatment, or management of apnea, apnea

disorders, bulimia, irritable bowel syndrome, urinary incontinence, bradycardia, bradyarrhythmia, asthma, syncope, or symptoms thereof in the methods of the present invention, may be made either concurrently or sequentially, *i.e.*, (-) norcispapride and at least one other therapeutic agent may be administered as a combination, concurrently but
5 separately, or by sequential administration. The compositions administered in each of these methods may be concurrent, sequential, or in any combination of concurrent or sequential.

The methods and compositions of this invention also include the benefit of reducing or avoiding adverse effects associated with administration of racemic cisapride. The invention also allows the concurrent or sequential use of antidepressant drugs, such as
10 tricyclic antidepressants, fluoxetine or its R or S stereoisomer, Zoloft®, and the like, and other drugs, such as anti-anxiety drugs.

The term "patient" as used herein refers to mammals, particularly humans.

The methods of the present invention for the prevention, treatment, or management of bulimia are particularly useful in adolescents and young adults. In a
15 preferred embodiment, the method of preventing, treating, or managing bulimia is directed to females from the ages of 13 to 25. It should be recognized that the methods of the present invention can be used to prevent, treat, or manage bulimia in males and females, including children and adults, notwithstanding the preferences mentioned above.

The methods of the present invention for the prevention, treatment, or management of apnea or apnea disorders are particularly useful in obese men. In a
20 preferred embodiment, the methods are directed to the prevention, treatment, or management of obstructive apnea in obese men. It should be recognized that the methods can be used to prevent, treat, or manage apnea or apnea disorders in males and females, including children and adults, notwithstanding the preferences mentioned above.

As used herein, the terms "adverse effects" and "adverse side effects"
25 include, but are not limited to, cardiac arrhythmia, cardiac conduction disturbances, appetite stimulation, weight gain, sedation, gastrointestinal distress, headache, dry mouth, constipation, diarrhea, and drug-drug interactions. See, for example, *Physician's Desk Reference*®, 52nd Edition, Medical Economics Co., Inc., pp. 1308-1309, 1998. The term
30 "cardiac arrhythmia" includes, but is not limited to, ventricular tachyarrhythmia, torsades de pointes, Q_T prolongation, and ventricular fibrillation.

The term "racemic" as used herein means a mixture of the (-) and (+) enantiomers of a compound wherein the (-) and (+) enantiomers are present in approximately a 1:1 ratio.

The terms "substantially optically pure," "optically pure," and "optically pure enantiomers," as used herein, mean that the composition contains greater than about 90% of the (-) norcisapride stereoisomer by weight, preferably greater than about 95% of the desired enantiomer by weight, and more preferably greater than about 99% of the desired enantiomer by weight, based upon the total weight of norcisapride. In other words, the term "substantially free" means less than about 10 weight percent, preferably less than about 5 weight percent, and more preferably less than about 1 weight percent of (+) norcisapride is present according to the invention.

The terms "5-hydroxytryptamine receptor antagonist," "serotonin receptor antagonist," and "5-HT₂ receptor antagonist," as used herein, mean a compound capable of binding reversibly to a 5-hydroxytryptamine receptor, whether on the vagal nerve or elsewhere in a mammal.

The phrases "bulimia" and "bulimia nervosa" are used herein consistently with the definition according to DSM-IV.

The terms "apnea" and "apnea disorder," as used herein, include, but are not limited to, a disorder characterized by interrupted breathing, in which a person stops breathing long enough to decrease the amount of oxygen and increase the amount of carbon dioxide in the blood and brain.

The term "asthma," as used herein, is defined as a disorder characterized by increased responsiveness of the trachea and bronchi to various stimuli, which results in symptoms that include, but are not limited to, wheezing, cough, shortness of breath, dyspnea, and the like. Asthma includes, for example, allergic asthma.

The term "syncope," as used herein, is defined as a disorder characterized by loss of consciousness and postural tone caused by diminished cerebral blood flow. Syncope includes, for example, Adams-Stokes syncope, cardiac syncope, carotid sinus syncope, hysterical syncope, laryngeal syncope, local syncope, micturition syncope, orthostatic syncope, postural syncope, swallow syncope, syncope due to seizures, syncope due to pulmonary embolism, syncope of gradual onset, tussive syncope, vasodepressor syncope, or vasovagal syncope.

The phrase "therapeutically effective amount of (-) norcisapride," as used herein, means that amount of substantially optically pure (-) norcisapride, or a pharmaceutically acceptable salt thereof, which, alone or in combination with other drugs, provides a therapeutic benefit in the prevention, treatment, or management, or of apnea, apnea disorders, bulimia, irritable bowel syndrome, urinary incontinence, bradycardia, bradyarrhythmia, asthma, syncope, or one or more symptoms thereof. Different therapeutically effective amounts may be applicable for each disorder, as will be readily known by those of ordinary skill in the art.

Substantially pure (-) norcisapride may be obtained from a racemic mixture of cisapride, the chemical synthesis of which can be performed according to the method described in European Patent Application No. 0,076,530 A2 published April 13, 1983, U.S. Patent Nos. 4,962,115, 5,057,525, 5,137,896, the disclosures of which are each hereby incorporated herein by express reference thereto. *See also*, and Van Daele, *et al.*, *Drug Development Res.*, 8:225-232 (1986) The metabolism of cisapride to norcisapride is described in Meuldermans, W., *et al.*, *Drug Metab. Dispos.*, 16(3):410-419 (1988) and Meuldermans, W., *et al.*, *Drug Metab. Dispos.*, 16(3):403-409 (1988). The preparation of racemic norcisapride is also known to those of ordinary skill in the art, particularly in view of EP 0,076,530 A2 and U.S. Patent No. 5,137,896 to Van Daele, the disclosures of which are hereby incorporated herein by express reference thereto.

Optically pure (-) norcisapride may also be obtained from racemic norcisapride by HPLC separation or resolution of the enantiomers using conventional means, for example, from an optically active resolving acid. The resolution of racemic norcisapride is also known to those of ordinary skill in the art, particularly from Jacques, J., *et al.*, *Enantiomers, Racemates and Resolutions* (Wiley-Interscience, New York, 1981); Wilen, S. H., *et al.*, *Tetrahedron*, 33:2725 (1977); Eliel, E. L. *Stereochemistry of Carbon Compounds* (McGraw-Hill, NY, 1962); Wilen, S. H. *Tables of Resolving Agents and Optical Resolutions*, p. 268 (E.L. Eliel, Ed. Univ. of Notre Dame Press, Notre Dame, IN, 1972).

In addition to separation techniques, such as those described above, (-) norcisapride may be synthesized by stereospecific synthesis using methodology well known to those of ordinary skill in the art. Chiral synthesis can result in products of high enantiomeric purity. However, in some cases, the enantiomeric purity of the product is not sufficiently high. The skilled artisan will appreciate that the separation methods described

above may be used to further enhance the enantiomeric purity of (-) norcisapride obtained by chiral synthesis.

Optically pure (-) norcisapride may also be prepared from the racemic mixture by enzymatic biocatalytic resolution. See, for example, U.S. Patent Nos. 5,057,427 and 5,077,217, the disclosures of which are incorporated herein by express reference thereto.

The magnitude of a prophylactic or therapeutic dose of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer, in the acute or chronic management of diseases and disorders described herein will vary with the severity of the condition to be prevented, treated, or managed and the route of administration. For example, oral, mucosal (including rectal), parenteral (including subcutaneous, intramuscular, bolus injection, and intravenous), sublingual, transdermal, nasal, buccal, and like may be employed. Dosage forms include tablets, troches, lozenges, dispersions, suspensions, suppositories, solutions, capsules, soft elastic gelatin capsules, patches, and the like. The dose, and perhaps the dose frequency, will also vary according to the age, body weight, and response of the individual patient. Suitable dosing regimens can be readily selected by those skilled in the art with due consideration of such factors. In general, the total daily dosage for the conditions described herein, is from about 0.5 mg to about 500 mg of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer. Preferably, a daily dose range is from about 1 mg to about 250 mg and more preferably, a daily dose range is between about 1 mg to about 100 mg. Preferably, the active ingredient is administered in single or divided doses orally from one to four times a day, or by slow intravenous injection. The most preferred route of administration for the present invention is oral. The oral dosage forms may be conveniently presented in unit dosage forms and prepared by any methods well known in the art of pharmacy.

In managing the patient, the therapy may be initiated at a lower dose, *e.g.*, from about 0.5 mg to about 10 mg, and increased up to the recommended daily dose or higher depending on the patient's global response. It is further recommended that children, patients over 65 years, and those with impaired renal or hepatic function, initially receive low doses, and that they be titrated based on individual response(s) and blood level(s). It may be necessary to use dosages outside these ranges in some cases, as will be apparent to those of ordinary skill in the art. Furthermore, it is noted that the clinician or treating

physician will know how and when to interrupt, adjust, or terminate therapy in conjunction with individual patient response.

Any suitable route of administration may be employed for providing the patient with an effective dosage of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer. The most suitable route in any given case will depend on the nature and severity of the condition being prevented, treated, or managed.

The pharmaceutical compositions for use in the present invention comprise optically pure (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer, as the active ingredient, and may also contain a pharmaceutically acceptable carrier, and optionally, other therapeutic ingredients. As used herein, the term "pharmaceutically acceptable salt" refers to a salt prepared from pharmaceutically acceptable non-toxic acids including inorganic acids, organic acids, solvates, hydrates, or clathrates thereof. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, sulfuric, and phosphoric. Appropriate organic acids may be selected, for example, from aliphatic, aromatic, carboxylic and sulfonic classes of organic acids, examples of which are formic, acetic, propionic, succinic, camphorsulfonic, citric, fumaric, gluconic, isethionic, lactic, malic, mucic, tartaric, para-toluenesulfonic, glycolic, glucuronic, maleic, furoic, glutamic, benzoic, anthranilic, salicylic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, pantothenic, benzenesulfonic (besylate), stearic, sulfanilic, alginic, galacturonic, and the like. Particularly preferred acids are hydrobromic, hydrochloric, phosphoric, and sulfuric acids. In a preferred embodiment, (-) norcisapride is administered as the free base or hydrate.

In practical use, (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer, can be combined as the active ingredient in intimate admixture with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms and may comprise a number of components depending on the form of preparation desired for administration. The compositions of the present invention include, but are not limited to, suspensions, solutions and elixirs; aerosols; or carriers, including, but not limited to, starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders, disintegrating agents, and the like. Because of their ease of administration, tablets and capsules are preferred and represent the most advantageous oral dosage unit form, in which

case solid pharmaceutical carriers are employed. If desired, tablets may be coated by standard aqueous or nonaqueous techniques.

Pharmaceutical compositions of the present invention suitable for oral administration may be presented as discrete pharmaceutical unit dosage forms, such as capsules, cachets, soft elastic gelatin capsules, tablets, or aerosols sprays, each containing a predetermined amount of the active ingredient, as a powder or granules, or as a solution or a suspension in an aqueous liquid, a non-aqueous liquid, an oil-in-water emulsion, or a water-in-oil liquid emulsion. Such compositions may be prepared by any of the methods of pharmacy, but all methods include the step of bringing into association the active ingredient with the pharmaceutically acceptable carrier which constitutes one or more necessary ingredients. In general, the compositions are prepared by uniformly and intimately admixing the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product into the desired presentation. Oral solid preparations are preferred over oral liquid preparations. One preferred oral solid preparation is capsules, but the most preferred oral solid preparation is tablets.

For example, a tablet may be prepared by compression or molding, optionally, with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as powder or granules, optionally mixed with a binder, lubricant, inert diluent, granulating agent, surface active agent, dispersing agent, or the like. Molded tablets may be made by molding, in a suitable machine, a mixture of the powdered compound moistened with an inert liquid diluent. Preferably, each tablet contains from about 0.5 mg to about 500 mg of the active ingredient, more preferably from about 1 mg to about 250 mg. Preferably, each cachet or capsule contains from about 0.5 mg to about 500 mg of the active ingredient, more preferably from about 1 mg to about 250 mg. However, the amount of active ingredient found in the composition may vary depending on the amount of active ingredient to be administered to the patient.

Optically pure (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer, may be formulated as a pharmaceutical composition in a soft elastic gelatin capsule unit dosage form by using conventional methods well known in the art, such as in Ebert, *Pharm. Tech.*, 1(5):44-50 (1977). Soft elastic gelatin capsules have a soft, globular gelatin shell somewhat thicker than that of hard gelatin capsules, wherein a gelatin is plasticized by the addition of plasticizing agent, e.g.,

glycerin, sorbitol, or a similar polyol. The hardness of the capsule shell may be changed by varying the type of gelatin used and the amounts of plasticizer and water. The soft gelatin shells may contain a preservative, such as methyl- and propylparabens and sorbic acid, to prevent the growth of fungi. The active ingredient may be dissolved or suspended in a liquid vehicle or carrier, such as vegetable or mineral oils, glycols such as polyethylene glycol and propylene glycol, triglycerides, surfactants such as polysorbates, or a combination thereof.

In addition to the common dosage forms set out above, the compounds of the present invention may also be administered by controlled release means, delivery devices, or both, as are well known to those of ordinary skill in the art, such as those described in U.S. Patent Nos.: 3,845,770; 3,916,899; 3,536,809; 3,598,123; 4,008,719; 5,674,533; 5,059,595; 5,591,767; 5,120,548; 5,073,543; 5,639,476; 5,354,556; and 5,733,566, the disclosures of which are hereby incorporated herein by express reference thereto. These pharmaceutical compositions can be used to provide slow or controlled-release of the active ingredient therein using, for example, hydropropylmethyl cellulose in varying proportions to provide the desired release profile, other polymer matrices, gels, permeable membranes, osmotic systems, multilayer coatings, microparticles, liposomes, microspheres, or the like, or a combination thereof. Suitable controlled-release formulations known to those of ordinary skill in the art, including those described herein, may be readily selected for use with the (-) norcisapride compositions of the invention. Thus, single unit dosage forms suitable for oral administration, such as tablets, capsules, gencaps, caplets, and the like, that are adapted for controlled-release are encompassed by the present invention.

All controlled-release pharmaceutical products have a common goal of improving drug therapy over that achieved by their non-controlled counterparts. Ideally, the use of an optimally designed controlled-release preparation in medical treatment is characterized by a minimum of drug substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled-release formulations may include: 1) extended activity of the drug; 2) reduced dosage frequency; and 3) increased patient compliance.

Most controlled-release formulations are designed to initially release an amount of drug that promptly produces the desired therapeutic effect, and gradual and continual release of other amounts of drug to maintain this level of therapeutic effect over an extended period of time. In order to maintain this constant level of drug in the body, the

drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body.

The controlled-release of the active ingredient may be stimulated by various inducers, for example pH, temperature, enzymes, water, or other physiological conditions or compounds. The term "controlled-release component" in the context of the present invention is defined herein as a compound or compounds, including polymers, polymer matrices, gels, permeable membranes, liposomes, microspheres, or the like, or a combination thereof, that facilitates the controlled-release of the active ingredient (*e.g.*, (-) norcisapride) in the pharmaceutical composition.

Optically pure (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer, may also be formulated for parenteral administration by injection (subcutaneous, bolus injection, intramuscular, or intravenous), and may be dispensed in a unit dosage form, such as a multidose container or an ampule. Compositions of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer, for parenteral administration may be in the form of suspensions, solutions, emulsions, or the like in aqueous or oily vehicles, and in addition to the active ingredient may contain one or more formulary agents, such as dispersing agents, suspending agents, stabilizing agents, preservatives, and the like.

In the case where an intravenous injection or infusion composition is employed, a suitable daily dosage range is, *e.g.*, from about 0.5 mg to about 500 mg total daily dose, preferably from about 1 mg to about 250 mg, more preferably from about 1 mg to about 100 mg.

Another preferred route of administration is transdermal delivery, for example, via an abdominal skin patch.

The invention is further defined by reference to the following examples, describing in detail the preparation of the compound and the compositions of the present invention, as well as their utility. It will be apparent to those skilled in the art that many modifications, both to materials and methods, may be practiced without departing from the purpose and interest of this invention.

5 EXAMPLES

5.1 EXAMPLE 1: Bioavailability

A single dose of test substance or vehicle is administered to male beagle dogs either intravenously as a bolus over one minute using a 23 gauge butterfly needle into the saphenous vein, or as a single dose via oral gavage. 2.0 mL of whole blood is collected from each dog prior to and at intervals of 0.083, 0.25, 0.5, 1, 2, 3, 4, 6, 9, 12, and 24 hours following the intravenous or oral administration of the optical isomers or racemic mixture of cisapride or of norcisapride. The dogs are placed in sling-restraint prior to administration of test substance and are transferred to metabolic cages following collection of the 0.083 hour blood sample. All blood samples are collected from an angiocatheter placed in a cephalic vein on the morning of the experiment.

The blood is drawn into a 3 cc syringe. The first 1.0-2.0 mL of blood is discarded. The next 2.0 mL of whole blood is quickly transferred to a heparinized tube. The heparinized tubes are kept on ice until the blood is added. After adding the blood to the tube, the contents of the tube are mixed and centrifuged to obtain plasma. The plasma is carefully decanted and transferred to a test tube labeled with: the animal number, the dose of test substance administered, the route of administration, the date of administration, and the time of blood collection. The tubes are stored at -20°C until analysis.

Analysis of the concentration of the optical isomers or racemates of norcisapride in each plasma sample is determined using high performance liquid chromatography. For each test substance the plasma concentration with respect to sample time is plotted for both routes of administration. The oral bioavailability of each test substance is determined by comparing the C_{max} and AUC for the oral route of administration versus those for the intravenous route. The $t_{1/2}$ for each test substance by both routes is calculated as an indicator of duration of action.

5.2 EXAMPLE 2: Receptor Activity

5-HT_{1A} Receptor Activity

Receptor selection and amplification technology (R-SAT) is used (Receptor Technologies Inc., Winooski, VT) to determine potential agonist and/or antagonist activity of racemic norcisapride, cisapride, and their enantiomers on cloned human serotonin 5-HT_{1A} receptor subtypes expressed in NIH 3T3 cells, such as in Burstein *et al.*, *J. Biol Chem.*, 270:3141-3146 (1995); and Messier *et al.*, *Pharmacol. Toxicol.*, 76(5):308-311 (1995).

The assay involves co-expression of a marker enzyme, β -galactosidase, with the serotonin receptor of interest. Ligands stimulate proliferation of cells that express the receptor and, therefore, the marker. Ligand-induced effects can be determined by assay of the marker.

5 NIH 3T3 cells are incubated, plated, and then transfected using human 5-HT_{1A} serotonin receptors, pSV- β -galactosidase, and salmon sperm DNA. The medium is changed one day later, and after 2 days, aliquots of the trypsinized cells are placed in wells of a 96 well plate. After five days in culture in the presence of the ligands, the levels of β -galactosidase are measured. The cells are then rinsed and incubated with the substrate, *o*-nitrophenyl β -D-galactopyranoside. After 16 hours, the plates are read at 405 nm on a
10 plate-reader. Each compound is tested for activity in triplicate at seven different concentrations (10, 2.5, 0.625, 0.156, 0.039, 0.0098, and 0.0024 nM).

None of the compounds tested show agonist activity at human 5-HT_{1A} serotonin receptors. Data from antagonist inhibition of the compounds are fit to the
15 equation:

$$\text{Response} = \text{Max Response} + \frac{(\text{Min Response})}{1 + (\text{Ligand Conc}/\text{EC}_{50})}$$

20 IC₅₀ values (concentration required to inhibit 50% of specific binding) are calculated for antagonist activity against a concentration of 2 μ M 5-HT using the non-linear least squares analysis of KalcidaGraph, the results of which are set forth in Tables 1 and 2.

5-HT₂ Receptor Activity

25 Receptor selection and amplification technology (R-SAT) is used (Receptor Technologies Inc., Winooski, VT) to determine potential agonist and/or antagonist activity of racemic norcispripide, cispripide, and their enantiomers on cloned human serotonin 5-HT₂ receptor subtypes expressed in NIH 3T3 cells, such as in Burstein *et al.*, *J. Biol Chem.*, 270:3141-3146 (1995); and Messier *et al.*, *Pharmacol. Toxicol.*, 76(5):308-311 (1995).

30 The assay involves co-expression of a marker enzyme, β -galactosidase, with the serotonin receptor of interest. Ligands stimulate proliferation of cells that express the receptor and, therefore, the marker. Ligand-induced effects can be determined by assay of the marker.

NIH 3T3 cells are incubated, plated, and then transfected using human 5-HT₂ serotonin receptors, pSV-β-galactosidase, and salmon sperm DNA. The medium is changed one day later, and after 2 days, aliquots of the trypsinized cells are placed in wells of a 96 well plate. After five days in culture in the presence of the ligands, the levels of β-galactosidase are measured. The cells are then rinsed and incubated with the substrate, *o*-nitrophenyl β-D-galactopyranoside. After 16 hours, the plates are read at 405 nm on a plate-reader. Each compound is tested for activity in triplicate at seven different concentrations (10, 2.5, 0.625, 0.156, 0.039, 0.0098, and 0.0024 nM).

None of the compounds tested show agonist activity at human 5-HT₂ serotonin receptors. Data from antagonist inhibition of the compounds are fit to the equation:

$$\text{Response} = \text{Max Response} + \frac{(\text{Min Response})}{1 + (\text{Ligand Conc}/\text{EC}_{50})}$$

IC₅₀ values are calculated for antagonist activity against a concentration of 2 μM 5-HT using the non-linear least squares analysis of KaleidaGraph, the results of which are set forth in Tables 1 and 2.

TABLE 1
Calculated IC₅₀ Values (μM) at 5-HT_{1A} and 5-HT₂ Receptors

Compound	5-HT _{1A}	5-HT ₂
(±) Norcisapride	7.48	2.21
(+) Norcisapride	0.0054	0.38
(-) Norcisapride	1.30	----

TABLE 2
Calculated IC₅₀ Values (μM) at 5-HT_{1A} and 5-HT₂ Receptors

Compound	5-HT _{1A}	5-HT ₂
(±) Cisapride	----	0.26
(+) Cisapride	----	0.0050
(-) Cisapride	----	7.08

5.3 EXAMPLE 3: Receptor Binding

5-HT₂ Receptor

Racemic norcisapride, racemic cisapride and their (+)- and (-)- stereoisomers are tested (Cerep, Celle l'Evescault, France) for binding to 5-HT₂ receptor subtypes derived from N1E-115 cells.

Following incubation with the appropriate ligands, the preparations are rapidly filtered under vacuum through GF/B glass fiber filters and washed with ice-cold buffer using a Brandel or Packard cell harvester. Bound radioactivity is determined with a liquid scintillation counter (LS 6000, Beckman) using a liquid scintillation cocktail (Formula 989).

Specific radioligand binding to the receptor is defined as the difference between total binding and nonspecific binding determined in the presence of an excess of unlabeled ligand. Results are expressed as a percent inhibition of specific binding obtained in the presence of the compounds. IC₅₀ are determined using concentrations ranging from 3×10^{-10} M to 10^{-5} M to obtain full competition curves and are calculated by non-linear regression analysis. The results are shown in Tables 3 and 4 below.

5-HT_{1A} Receptor

Racemic norcisapride, racemic cisapride and their (+)- and (-)- stereoisomers are tested (Cerep, Celle l'Evescault, France) for binding to 5-HT_{1A} receptor subtypes derived from guinea-pig striata.

Following incubation with the appropriate ligands, the preparations are rapidly filtered under vacuum through GF/B glass fiber filters and washed with ice-cold buffer using a Brandel or Packard cell harvester. Bound radioactivity is determined with a

liquid scintillation counter (LS 6000, Beckman) using a liquid scintillation cocktail (Formula 989).

Specific radioligand binding to the receptor is defined as the difference between total binding and nonspecific binding determined in the presence of an excess of unlabeled ligand. Results are expressed as a percent inhibition of specific binding obtained in the presence of the compounds. IC_{50} are determined using concentrations ranging from 3×10^{-10} M to 10^{-5} M to obtain full competition curves and are calculated by non-linear regression analysis. The results are shown in Tables 3 and 4 below.

TABLE 3

IC_{50} (nM) Values for Binding to 5-HT₃ and 5-HT₄ Sites

Compound	5-HT ₃	5-HT ₄	5-HT ₃ /5-HT ₄ Ratio
(±) Norcisapride	8.2	686	0.012
(+) Norcisapride	4.5	331	0.014
(-) Norcisapride	30.4	1350	0.023

TABLE 4

IC_{50} (nM) Values for Binding to 5-HT₃ and 5-HT₄ Sites

Compound	5-HT ₃	5-HT ₄	5-HT ₃ /5-HT ₄ Ratio
(±) Cisapride	365	169	2.2
(+) Cisapride	310	340	0.9
(-) Cisapride	2790	199	14.0

Agonist activity at 5-HT₄ receptor sites may also be assessed using an assay based on the ability of active compounds to increase cyclic AMP production in mouse embryo colliculi neurones grown in tissue culture, such as in Dumuis *et al.*, *N. S. Arch. Pharmacol.*, 340:403-410 (1989).

5.4 EXAMPLE 4: Oral FormulationTablets

	Formula	Quantity per Tablet in mg.		
		A	B	C
5	Active Ingredient (-) Norcisapride	5.0	10.0	25.0
	Lactose BP	62.0	57.0	42.0
	Starch BP	20.0	20.0	20.0
	Microcrystalline Cellulose	10.0	10.0	10.0
10	Hydrogenated Vegetable Oil	1.5	1.5	1.5
	Polyvinylpyrrolidinone	1.5	1.5	1.5
	Compression Weight	100.0	100.0	100.0

The active ingredient, (-) norcisapride, is sieved through a suitable sieve and
 15 blended with the lactose until a uniform blend is formed. Suitable volumes of water are
 added and the powders are granulated. After drying, the granules are then screened and
 blended with the remaining excipients. The resulting granules are then compressed into
 tablets of desired shape. Tablets of other strengths may be prepared by altering the ratio of
 active ingredient to the excipient(s) or the compression weight.

20

5.5 EXAMPLE 5: Oral FormulationTablets

	Formula	Quantity per Tablet in mg.		
		A	B	C
25	Active Ingredient (-) Norcisapride	5.0	10.0	25.0
	Lactose BP	48.5	43.5	28.5
	Starch BP	30.0	30.0	30.0
	Pregelatinized Maize Starch BP	15.0	15.0	15.0
30	Magnesium Stearate BP	1.5	1.5	1.5
	Compression Weight	100.0	100.0	100.0

The active ingredient, (-) norcisapride, is sieved through a suitable sieve and blended with lactose, starch, and pregelatinized maize starch until a uniform blend is formed. Suitable volumes of water are added and the powders are granulated. After drying, the granules are then screened and blended with the remaining excipients. The resulting granules are then compressed into tablets of desired shape. Tablets of other strengths may be prepared by altering the ratio of active ingredient to the excipient(s) or the compression weight.

5.6 EXAMPLE 6: Oral Formulation

10 Capsules

Formula	Quantity per Capsule in mg.		
	A	B	C
Active Ingredient (-) Norcisapride	5.0	10.0	25.0
15 Starch 1500	94.0	89.0	74.0
Magnesium Stearate BP	1.0	1.0	1.0
Total Weight	100.0	100.0	100.0

The active ingredient is sieved and blended with the excipients. The mix is filled into size No. 2 hard gelatin capsules using suitable machinery. Other doses may be prepared by altering the fill weight, and if necessary, changing the capsule size to suit.

5.7. EXAMPLE 7

INTRAVENOUS FORMULATION

25

Formula	
Active Ingredient (-) norcisapride	1000 µg/mL
Dilute Hydrochloric Acid BP	to pH 3.5
Sodium Chloride Injection BP	1 mL

30

The active ingredient is dissolved in dilute hydrochloric acid BP to form a solution having a concentration of 1000 µg/mL (-) norcisapride. The solution is then mixed with sodium chloride injection BP prior to use.

5 While the present invention has been described with respect to the particular embodiments, it will be apparent to those skilled in the art that various changes and modifications may be made without departing from the spirit and scope of the invention as defined in the claims. Such modifications are also intended to fall within the scope of the appended claims.

10

 The above discussion of documents, acts, materials, devices, articles and the like is included in this specification solely for the purpose of providing a context for the present invention. It is not suggested or represented that any or all of these matters formed part of the prior art base or were
15 common general knowledge in the field relevant to the present invention as it existed in Australia before the priority date of each claim of this application.

 Throughout the description and claims of the specification the word
20 "comprise" and variations of the word, such as "comprising" and "comprises", is not intended to exclude other additives, components, integers or steps.



THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A method of treating bulimia in a patient which comprises administering to said patient in need of such treatment a therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.
2. A method according to claim 1, wherein the mammal is a human.
3. A method according to either claim 1 or 2, wherein (-) norcisapride is administered orally.
4. A method according to any one of claims 1 to 3, wherein (-) norcisapride is administered as a tablet or a capsule.
5. A method according to any one of claims 1 to 4, wherein the amount administered is from about 0.5 mg to about 500 mg.
6. A method according to claim 5, wherein the amount administered is from about 1 mg to about 250 mg.
7. A method according to any one of claims 1 to 6, wherein (-) norcisapride is administered together with a pharmaceutically acceptable carrier.
8. A method according to any one of claims 1 to 7, wherein said (-) norcisapride is administered from one to four times per day.
9. A method according to either claim 1 or 2, wherein (-) norcisapride is administered parenterally, transdermally, rectally or sublingually.
10. A method of treating disorders mediated by vagal activity in a patient which comprises administering to said patient in need of such treatment a therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

JMN W:\SPECIM\0704-00.doc

11. A method according to claim 10, wherein the mammal is a human.
12. A method according to either claim 10 or 11, wherein (-) norcisapride is
5 administered orally.
13. A method according to any one of claims 10 to 12, wherein (-) norcisapride
is administered as a tablet or a capsule.
- 10 14. A method according to any one of claims 10 to 13, wherein the amount
administered is from about 0.5 mg to about 500 mg.
- 15 15. A method according to claim 14, wherein the amount administered is from
about 1 mg to about 250 mg.
16. A method according to any one of claims 10 to 15, wherein (-) norcisapride
is administered together with a pharmaceutically acceptable carrier.
17. A method according to any one of claims 10 to 16, wherein said (-)
20 norcisapride is administered from one to four times per day.
18. A method according to either claim 10 or 11, wherein (-) norcisapride is
administered parenterally, transdermally, rectally or sublingually.
- 25 19. A method of treating irritable bowel syndrome in a patient which comprises
administering to said patient in need of such treatment a therapeutically
effective amount of (-) norcisapride, or a pharmaceutically acceptable salt
thereof, substantially free of its (+) stereoisomer.
- 30 20. A method according to claim 19, wherein the mammal is a human.
21. A method according to either claim 19 or 20, wherein (-) norcisapride is
administered orally.

22. A method according to any one of claims 19 to 21, wherein (-) norcisapride is administered as a tablet or a capsule.
23. A method according to any one of claims 19 to 22, wherein the amount administered is from about 0.5 mg to about 500 mg.
24. A method according to claim 23, wherein the amount administered is from about 1 mg to about 250 mg.
25. A method according to any one of claims 19 to 24, wherein (-) norcisapride is administered together with a pharmaceutically acceptable carrier.
26. A method according to any one of claims 19 to 25, wherein said (-) norcisapride is administered from one to four times per day.
27. A method according to claim 19, wherein (-) norcisapride is administered parenterally, transdermally, rectally or sublingually.
28. A method of treating bradycardia or bradyarrhythmia in a patient which comprises administering to said patient in need of such treatment a therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.
29. A method according to claim 28, wherein the patient is a human.
30. A method according to either claim 28 or 29, wherein (-) norcisapride is administered orally.
31. A method according to any one of claims 28 to 30, wherein (-) norcisapride is administered as a tablet or a capsule.
32. A method according to any one of claims 28 to 31, wherein the amount administered is from about 0.5 mg to about 500 mg.

33. A method according to claim 32, wherein the amount administered is from about 1 mg to about 250 mg.
34. A method according to any one of claims 28 to 33, wherein (-) norcisapride
5 is administered together with a pharmaceutically acceptable carrier.
35. A method according to any one of claims 28 to 34, wherein said (-) norcisapride is administered from one to four times per day.
- 10 36. A method according to either claim 28 or 29, wherein (-) norcisapride is administered parenterally, transdermally, rectally or sublingually.
37. A method of treating asthma in a patient which comprises administering to said patient in need of such treatment a therapeutically effective amount of (-)
15 norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.
38. A method according to claim 37, wherein the patient is a human.
- 20 39. A method according to either claim 37 or 38, wherein (-) norcisapride is administered orally.
40. A method according to any one of claims 37 to 40, wherein (-) norcisapride is administered as a tablet or a capsule.
- 25 41. A method according to any one of claims 37 to 40, wherein the amount administered is from about 0.5 mg to about 500 mg.
42. A method according to claim 41, wherein the amount administered is from
30 about 1 mg to about 250 mg.
43. A method according to any one of claims 37 to 42, wherein (-) norcisapride is administered together with a pharmaceutically acceptable carrier.

44. A method according to any one of claims 37 to 43, wherein said (-) norcisapride is administered from one to four times per day.

45. A method according to either claim 37 or 38, wherein (-) norcisapride is administered parenterally, transdermally, rectally or sublingually.

46. A method of treating urinary incontinence in a patient which comprises administering to said patient in need of such treatment a therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

47. A method according to claim 46, wherein the patient is a mammal.

48. A method according to either claim 46 or 47, wherein (-) norcisapride is administered orally.

49. A method according to any one of claims 46 to 48, wherein (-) norcisapride is administered as a tablet or a capsule.

50. A method according to any one of claims 46 to 49, wherein the amount administered is from about 0.5 mg to about 500 mg.

51. A method according to claim 50, wherein the amount administered is from about 1 mg to about 250 mg.

52. A method according to any one of claims 46 to 51, wherein (-) norcisapride is administered together with a pharmaceutically acceptable carrier.

53. A method according to any one of claims 46 to 52, wherein said (-) norcisapride is administered from one to four times per day.

54. A method according to either claim 46 or 47, wherein (-) norcisapride is administered parenterally, transdermally, rectally or sublingually.

55. A method of treating apnea or apnea disorders in a patient which comprises administering to said patient in need of such treatment a therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

5

56. A method according to claim 55, wherein the apnea being treated comprises central apnea, deglutition apnea, obstructive apnea, sleep apnea, or sleep induced apnea.

10 57. A method according either claim 55 or 56, wherein the patient is a human.

58. A method according to any one of claims 55 to 57, wherein (-) norcisapride is administered orally.

15 59. A method according to any one of claims 55 to 58, wherein (-) norcisapride is administered as a tablet or a capsule.

60. A method according to any one of claims 55 to 59, wherein the amount administered is from about 0.5 mg to about 500 mg.

20

61. A method according to claim 60, wherein the amount administered is from about 1 mg to about 250 mg.

62. A method according to any one of claims 55 to 61, wherein (-) norcisapride is administered together with a pharmaceutically acceptable carrier.

25

63. A method according to any one of claims 55 to 62, wherein said (-) norcisapride is administered from one to four times per day.

30 64. A method according to any one of claims 55 to 57, wherein (-) norcisapride is administered parenterally, transdermally, rectally or sublingually.

65. A method of preventing or managing bulimia in a patient which comprises administering to said patient in need of such prevention or management a

therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

5 66. A method of preventing or managing disorders mediated by vagal activity in a patient which comprises administering to said patient in need of such prevention or management a therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

10 67. A method of preventing or managing irritable bowel syndrome in a patient which comprises administering to said patient in need of such prevention or management a therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

15 68. A method of preventing or managing bradycardia or bradyarrhythmia in a patient which comprises administering to said patient in need of such prevention or management a therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

20 69. A method of preventing or managing asthma in a patient which comprises administering to said patient in need of such prevention or management a therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

25 70. A method of preventing or managing urinary incontinence in a patient which comprises administering to said patient in need of such prevention or management a therapeutically effective amount of (-) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

71. A method of preventing or managing apnea or apnea disorders in a patient which comprises administering to said patient in need of such prevention or management a therapeutically effective amount of (-) norcisapride, or a
5 pharmaceutically acceptable salt thereof, substantially free of its (+) stereoisomer.

DATED : 13 June, 2001

10

PHILLIPS ORMONDE & FITZPATRICK

Attorneys For:

SEPRACOR, INC.

5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189