



US 20090181399A1

(19) **United States**

(12) **Patent Application Publication**
Negrotto et al.

(10) **Pub. No.: US 2009/0181399 A1**

(43) **Pub. Date: Jul. 16, 2009**

(54) **INSECT RESISTANT COTON PLANTS AND
METHODS OF DETECTING THE SAME**

(22) Filed: **Mar. 12, 2009**

Related U.S. Application Data

(75) Inventors: **David Vincent Negrotto**, Research
Triangle Park, NC (US); **Frank**
Arthur Shotkoski, Ithaca, NY
(US); **Wenjin Yu**, Cary, NC (US)

(62) Division of application No. 10/580,596, filed on May
25, 2006, now Pat. No. 7,521,550, filed as application
No. PCT/EP2004/012662 on Nov. 9, 2004.

(60) Provisional application No. 60/526,112, filed on Dec.
1, 2003.

Correspondence Address:

SYNGENTA BIOTECHNOLOGY, INC.
PATENT DEPARTMENT
3054 CORNWALLIS ROAD, P.O. BOX 12257
RESEARCH TRIANGLE PARK, NC 27709-2257
(US)

Publication Classification

(51) **Int. Cl.**
C12Q 1/68 (2006.01)

(52) **U.S. Cl.** **435/6**

(57) **ABSTRACT**

(73) Assignee: **SYNGENTA PARTICIPATIONS**
AG, Research Triangle Park, NC
(US)

The present application relates to an insect resistant trans-
genic cotton event designated COT202. The application also
relates to polynucleotides which are characteristic of the
COT202 event, plants comprising said polynucleotides, and
methods of detecting the COT202 event.

(21) Appl. No.: **12/402,957**

INSECT RESISTANT COTON PLANTS AND METHODS OF DETECTING THE SAME

[0001] This is a divisional of U.S. patent application Ser. No. 10/580,596 filed May 25, 2006, which is a §371 of PCT/EP2004/012662, filed Nov. 9, 2004, and published Jun. 16, 2005, as WO 05/054479, which claims priority to U.S. Provisional Application No. 60/526,112, filed Dec. 1, 2003, which is hereby incorporated by reference in its entirety.

[0002] The present invention relates to genetic engineering of plants and in particular to an insect resistant transgenic cotton plant. Specifically, the invention relates to a cotton plant designated COT202 which comprises a VIP3A gene. It also relates to methods of detecting material derived from the plant.

[0003] Plant pests are a major factor in the loss of the world's important agricultural crops. About \$8 billion is lost every year in the U.S. due to infestations of plants by non-mammalian pests including insects. In addition to losses in field crops, insect pests are also a burden to vegetable and fruit growers, to producers of ornamental flowers, and to home gardeners.

[0004] Insect pests are mainly controlled by intensive applications of chemical pesticides, which are active through inhibition of insect growth, prevention of insect feeding or reproduction, or cause death. Good control of insect pests can thus be reached, but these chemicals can sometimes also affect other, beneficial insects. Another problem resulting from the wide use of chemical pesticides is the appearance of resistant insect varieties. This has been partially alleviated by various resistance management practices, but there is an increasing need for alternative pest control agents. Biological pest control agents, such as *Bacillus thuringiensis* strains expressing pesticidal toxins like δ -endotoxins, have also been applied to crop plants with satisfactory results, offering an alternative or complement to chemical pesticides. The genes coding for some of these δ -endotoxins have been isolated and their expression in heterologous hosts has been shown to provide another tool for the control of economically important insect pests. In particular, the expression of insecticidal toxins such as *Bacillus thuringiensis* δ -endotoxins in transgenic plants, has provided efficient protection against selected insect pests, and transgenic plants expressing such toxins have been commercialised, allowing farmers to reduce applications of chemical insect control agents.

[0005] Recently, a new family of insecticidal proteins produced by *Bacillus* sp. during the vegetative stages of growth (vegetative insecticidal proteins (VIPs)) has been identified. U.S. Pat. Nos. 5,877,012, 6,107,279, and 6,137,033 describe vip3A toxin genes isolated from *Bacillus* species. The VIP3A toxins possess insecticidal activity against a wide spectrum of lepidopteran insects including but not limited to fall armyworm, *Spodoptera frugiperda*, black cutworm, *Agrotis ipsilon*, sugarcane borer, *Diatraea saccharalis*, and lesser cornstalk borer, *Elasmopalpus lignosellus*, and when expressed in transgenic plants, for example cotton, confer protection on the plant from insect feeding damage.

[0006] The cotton family, genus *Gossypium*, a member of the Malvaceae, consists of 39 species, of which *Gossypium hirsutum* is the most commonly cultivated species. Three other species are also cultivated: *G. arboreum*, *G. barbadense*, and *G. herbaceum*. These cultivated species are grown primarily for the seed hairs that are made into textiles.

Cotton is suitable as a textile fibre because the mature dry hairs twist in such a way that fine strong threads can be spun from them. Other products, such as cottonseed oil, cake, and cotton linters are by-products of fibre production.

[0007] Damage to cotton crops by insect pests throughout the world results in a significant yield loss each year. Effective control of these pests to minimise yield loss is of great economic importance. Examples of insect pests of cotton include Beet armyworm (*Spodoptera exigua*), Boll weevil (*Anthonomus grandis grandis*), Cabbage looper (*Trichoplusia ni*), Clouded plant bug (*Neurocolpus nubilus*), Cotton aphid (*Aphis gossypii*), Cotton bollworm (*Helioverpa zea*), Cutworms (*Feltia subterranea*, *Peridroma saucia*, *Agrotis ipsilon*), European corn borer (*Ostrinia nubilalis*), Fall armyworm (*Spodoptera frugiperda*), Seedling thrips (*Frankliniella* spp.), Soybean looper (*Pseudoplusia includens*), Stink bugs (*Nezara viridula*, *Acrosternum hilare*, *Euschistus servus*), Tarnished plant bug (*Lygus lineolaris*), Tobacco budworm (*Heliothis virescens*) and Whiteflies (*Trialeurodes abutilonea*, *Bemisia tabaci*).

[0008] Transformation and regeneration of cotton plants is now a well-established procedure, typically based on *Agrobacterium tumefaciens* mediated transfer of foreign DNA into cotton plant parts and regeneration of said plant parts in tissue culture into fully fertile, transgenic cotton plants.

[0009] There exists a requirement to generate a cotton plant that is insect resistant so that yield loss through damage to cotton crops by insect pests is reduced. An insect resistant cotton plant could reduce the need to apply chemical pesticides, which may be detrimental to other, beneficial insects and the environment. Further, it is desirable to provide an insect resistant plant that comprises a VIP gene, as an alternative to transgenic plants comprising crystal proteins from *Bacillus thuringiensis*. This may be of use in insect resistance management.

[0010] Therefore, the present invention relates to an insect resistant transgenic cotton event, designated COT202. It also relates to methods of detecting plant material derived therefrom. "COT202 event" in the context of this application refers to the original insecticidal transgenic cotton plant described herein. "Insecticidal" as used herein refers to any inhibitory effect on an insect, including but not limited to reduced feeding, retarded growth, reduced fecundity, paralysis or death. "Fecundity" comprises all aspects related to reproduction such as reproductive ability, reproductive frequency and number of offspring. Also embraced by this invention is any plant material derived from the COT202 event, including seeds.

[0011] The COT202 event exhibits a novel genotype comprising at least one expression cassette. The cassette comprises a suitable promoter for expression in plants operably linked to a gene that encodes a VIP3A insecticidal toxin, useful in controlling a wide spectrum of lepidopteran insect pests, and a suitable polyadenylation signal. Suitable promoters may be isolated from, inter alia, plants. Numerous plant promoters have been isolated and characterised including constitutive, switchable and/or tissue specific promoters. Suitable promoters may be selected from the following, non-limiting group: CaMV35S, FMV35S, Ubiquitin, Act2, NOS, OCS, Cestrum yellow leaf curl virus promoter, Patatin, E9, alcA/alcR switch, GST switch, RMS switch, oleosin, Gelvin, ribulose biphosphate carboxylase-oxygenase small subunit, actin 7, MR7 promoter (maize), Gos 9 (rice), GOS2 promoters, MasOcs (or super promoter), RolD promoter

(*Agrobacterium rhizogenes*), SuperMAS promoter, and Suc2 promoter (*Arabidopsis*). In one embodiment of the present invention, the promoter is the Ubiquitin promoter, UBQ3, from *Arabidopsis thaliana*. Additional elements such as enhancer sequences may also be incorporated into the expression cassette in order to boost levels of gene expression, for example transcriptional or translational enhancers, such as tobacco etch virus (TEV) translation activator, CaMV35S enhancer, and FMV35S enhancer. Alternatively it may be desirable to include a targeting sequence, for example, to direct transportation of the VIP3A toxin to a particular cellular compartment. For example if it is desired to provide the protein outside of the cell then an extracellular targeting sequence may be ligated to the polynucleotide encoding the VIP protein. Other examples of targeting include targeting to a specific intracellular organelle or compartment, for example to the endoplasmic reticulum using a 'KDEL' retention sequence. Numerous polyadenylation signals have been isolated and characterised. Examples of suitable polyadenylation signals functional in plants include that from the nopaline synthase gene (nos) of *Agrobacterium tumefaciens*, from the proteinase inhibitor II gene and from the alpha-tubulin gene (EP-A 652,286). In one embodiment of the present invention, the polyadenylation signal is that from the nos gene of *Agrobacterium tumefaciens*.

[0012] According to the invention, the polynucleotide encoding the VIP3A protein may also be codon-optimised or otherwise altered to enhance for example, transcription once it is incorporated into plant material. Such codon optimisation may also be used to alter the predicted secondary structure of the RNA transcript produced in any transformed cell, or to destroy cryptic RNA instability elements present in the unaltered transcript, thereby increasing the stability and/or availability of the transcript in the transformed cell (Abler and Green (1996) Plant Molecular Biology (32) pp. 63-78).

[0013] In a precursor to the COT202 event, a second cassette is present that comprises a gene which, when expressed, can be used as a selectable marker. Numerous selectable markers have been characterised, including some that confer tolerance to antibiotics and others that confer tolerance to herbicides. Examples of suitable selectable marker genes include those that confer tolerance to hygromycin, kanamycin or gentamycin. Further suitable selectable markers include genes that confer resistance to herbicides such as glyphosate-based herbicides or resistance to toxins such as eutypine. Other forms of selection are also available such as hormone based selection systems such as the Multi Auto Transformation (MAT) system of Hiroyasu Ebinuma et al. (1997) PNAS Vol. 94 pp. 2117-2121; visual selection systems which use the known green fluorescence protein, β glucuronidase; and any other selection system such as mannose isomerase (Positech™), xylose isomerase and 2-deoxyglucose (2-DOG). In one embodiment of the present invention, the selectable marker gene is one that confers tolerance to hygromycin. This second expression cassette is useful for selecting transformants during and following plant transformation. Optionally, it may be segregated away from the COT202 event precursor after transformation to leave the COT202 event itself. The COT202 event per se does not comprise a selectable marker cassette. Further expression cassettes are optionally comprised in the COT202 event. For example these may provide other desirable benefits such as herbicide resistance.

[0014] The expression cassettes may be introduced into the plant on the same or different plasmids. If the expression cassettes are present on the same plasmid and introduced into the plant via an *Agrobacterium*-mediated transformation method, they may be present within the same or different T-DNA regions. In one embodiment of the present invention, two expression cassettes are present on different T-DNA regions within the same plasmid.

[0015] According to the first aspect of the invention, there is provided a polynucleotide comprising at least 17 contiguous nucleotides from the 26-nucleotide sequence of SEQ ID NO: 1. In one embodiment said polynucleotide comprises at least 18 contiguous nucleotides from SEQ ID NO: 1. In a further embodiment said polynucleotide comprises at least 20 contiguous nucleotides from SEQ ID NO: 1. In a still further embodiment said polynucleotide comprises at least 22 contiguous nucleotides from SEQ ID NO: 1. In a further embodiment said polynucleotide comprises at least 23 contiguous nucleotides from SEQ ID NO: 1. In yet a further embodiment said polynucleotide comprises at least 24 contiguous nucleotides from SEQ ID NO: 1. In a further embodiment said polynucleotide comprises at least 25 contiguous nucleotides from SEQ ID NO: 1. In a still further embodiment there is provided a polynucleotide comprising the sequence of SEQ ID NO: 1.

[0016] In a further aspect of the invention, there is provided a polynucleotide comprising at least 17 contiguous nucleotides from the 26-nucleotide sequence of SEQ ID NO: 2. In one embodiment said polynucleotide comprises at least 18 contiguous nucleotides from SEQ ID NO: 2. In a further embodiment said polynucleotide comprises at least 20 contiguous nucleotides from SEQ ID NO: 2. In a still further embodiment said polynucleotide comprises at least 22 contiguous nucleotides from SEQ ID NO: 2. In a further embodiment said polynucleotide comprises at least 23 contiguous nucleotides from SEQ ID NO: 2. In yet a further embodiment said polynucleotide comprises at least 24 contiguous nucleotides from SEQ ID NO: 2. In a further embodiment said polynucleotide comprises at least 25 contiguous nucleotides from SEQ ID NO: 2. In a still further embodiment there is provided a polynucleotide comprising the sequence of SEQ ID NO: 2.

[0017] In a further aspect of the present invention there is provided a polynucleotide as described above further comprising the sequence of SEQ ID NO: 7. In a further aspect of the present invention there is provided a polynucleotide as described above further comprising the sequence of SEQ ID NO: 8.

[0018] In another aspect of the present invention there is provided a plant comprising a polynucleotide which comprises at least 17 contiguous nucleotides of SEQ ID NO: 1 and/or SEQ ID NO: 2. In one embodiment said plant comprises at least 18 contiguous nucleotides of SEQ ID NO: 1 and/or SEQ ID NO: 2. In a further embodiment said plant comprises at least 20 contiguous nucleotides of SEQ ID NO: 1 and/or SEQ ID NO: 2. In a further embodiment said plant comprises at least 22 contiguous nucleotides of SEQ ID NO: 1 and/or SEQ ID NO: 2. In a further embodiment said plant comprises at least 23 contiguous nucleotides of SEQ ID NO: 1 and/or SEQ ID NO: 2. In a still further embodiment said plant comprises at least 24 contiguous nucleotides of SEQ ID NO: 1 and/or SEQ ID NO: 2. In a further embodiment said plant comprises at least 25 contiguous nucleotides of SEQ ID NO: 1 and/or SEQ ID NO: 2. In yet a further embodiment said

plant comprises the sequence of SEQ ID NO: 1 and/or SEQ ID NO: 2. In a further embodiment, said plant additionally comprises the sequence of SEQ ID NO: 7. In a further embodiment still, said plant additionally comprises the sequence of SEQ ID NO: 8. In one embodiment of the present invention, said plant is a cotton plant. In a further embodiment, said plant is an insecticidal cotton plant which is the COT202 event, or a plant derived therefrom.

[0019] The skilled man is familiar with plant transformation methods. In particular, two principal techniques have been characterised across a wide range of plant species: transformation by *Agrobacterium* and transformation by direct DNA transfer.

[0020] *Agrobacterium*-mediated transformation is a commonly used method for transformation of dicotyledonous plants. The foreign DNA to be introduced into the plant is cloned into a binary vector in between left and right border consensus sequences. This is the T-DNA region. The binary vector is transferred into an *Agrobacterium* cell, which is subsequently used to infect plant tissue. The T-DNA region of the vector comprising the foreign DNA is inserted into the plant genome. The marker gene cassette and trait gene cassette may be present on the same T-DNA region, different T-DNA regions in the same vector, or even different T-DNA regions in different vectors. In one embodiment of the present invention, the cassettes are present on different T-DNA regions in the same vector.

[0021] Alternatively, direct DNA transfer can be used to introduce the DNA directly into a plant cell. One suitable method of direct transfer may be bombardment of plant cells with a vector comprising the DNA for insertion using a particle gun (particle-mediated biolistic transformation); another established method, 'whiskers', involves coating the DNA onto silicon carbide fibres onto which cells are impaled. Other methods for transforming plant cells include protoplast transformation (optionally in the presence of polyethylene glycols); sonication of plant tissues, cells or protoplasts in a medium comprising the polynucleotide or vector; micro-insertion of the polynucleotide or vector into plant material (optionally employing the known silicon carbide "whiskers" technique), electroporation and the like.

[0022] Following transformation, transgenic plants must be regenerated from the transformed plant tissue, and progeny possessing the foreign DNA selected using an appropriate marker such as resistance to hygromycin. The skilled man is familiar with the composition of suitable regeneration media. The selectable marker can be segregated away from transgenic events by conventional plant breeding methods, thus resulting in, for example, the COT202 event.

[0023] A plant of the invention, as described herein, has an insecticidal effect on insects from one or more species from the group comprising *Heliothis* sp., *Helicoverpa* sp. and *Spodoptera* sp. which may infest it. "Infest" as used herein refers to attack, colonisation, feeding or damage in any way by one or more insects. Thus, for example, the plant of the present invention will provide a self-defence mechanism against infestation by pest insects such as *Helicoverpa zea* (cotton boll worm). As a result, a reduced number of insecticide sprays are required during the cultivation of said plant compared to a non-transgenic cotton plant of the same variety and yield loss through insect pests is kept at a minimal level.

[0024] The present invention is not limited to the COT202 event itself, but is further extended to include any plant material derived therefrom, including seeds in so far as they con-

tain at least one of the present inventive polynucleotides. The present invention includes, but is not limited to plants that are derived from a breeding cross with the COT202 event or a derivative therefrom by conventional breeding or other methods. The invention also includes plant material derived from the COT202 event that may comprise additional, modified or fewer polynucleotide sequences compared to the COT202 event or exhibit other phenotypic characteristics. For example it may be desirable to transform plant material derived from the COT202 event to generate a new event that possesses an additional trait, such as a second insect resistance gene. This process is known as gene stacking. The second insect resistance gene may encode, for example insecticidal lectins, insecticidal protease inhibitors and insecticidal proteins derived from species of the *Bacillus thuringiensis*, *Xenorhabdus nematophilus*, or *Photorhabdus luminescens*. In one aspect, the second insect resistance gene encodes an insecticidal gene from *Bacillus thuringiensis*. Preferably, the second insect resistance gene encodes a Cry gene from the bacterium *Bacillus thuringiensis*, which Cry gene produces a toxin with a different mode of action or binding site in the insect gut to VIP for the control of different insect species. The Cry gene may, for example, be Cry 1 Ab.

[0025] The present invention further provides plant material derived from the COT202 event which possesses an additional trait such as herbicide resistance, nematode resistance or fungal resistance. In one embodiment, said additional trait is herbicide resistance. The herbicide resistance trait may be provided, for example, by a herbicide degradation enzyme, or a target-site specific resistant enzyme. In a further embodiment, said herbicide resistance trait provides resistance to a herbicide which comprises glyphosate acid or an agriculturally acceptable salt thereof. In a further embodiment still, said herbicide resistance trait is provided by a gene encoding EPSP synthase or a mutant thereof.

[0026] The present invention further provides a method of controlling insects comprising providing the COT202 event or plant material derived from the COT202 event at a locus where said insects feed. The invention yet further provides a method of controlling insects comprising providing the COT202 event or plant material derived from the COT202 event at a locus where said insects feed, and applying other agrochemicals to said plant material such as herbicides, fungicides and other insecticidal compounds including other insecticidal proteins. Examples of possible insecticidal compounds include insecticidal lectins, insecticidal protease inhibitors and insecticidal proteins derived from species of the *Bacillus thuringiensis*, *Xenorhabdus nematophilus*, or *Photorhabdus luminescens*. Examples of possible chemicals include pyrethroids, carbamates, imidacloprid, organochlorines, and macromolecules such as spinosad, abamectin or emamectin.

[0027] According to yet a further aspect of the present invention, there is provided a method of detecting the COT202 event or plant material derived from the COT202 transgenic event comprising obtaining a sample for analysis; extracting DNA from the sample; providing a pair of primers designed to bind to a polynucleotide comprising at least 17 contiguous nucleotides of SEQ ID NO: 1 and/or SEQ ID NO: 2; amplifying the region which lies between the sites at which the primers bind; and detecting the presence of the amplification product. Suitable pairs of primers for use in this method of detection can be designed using parameters well known to those skilled in the art of molecular biology now

that SEQ ID NOs 1 and 2 are made available. For example, one or both primers of the pair may be designed to be vector-specific, trait gene specific, promoter specific, and/or specific to the sequence of the junction between the inserted DNA and the genomic DNA. Preferably one of the primers is designed to be specific to the inserted sequence, and the other primer specific to the genomic DNA upstream or downstream of the insertion site. In one embodiment, the sequence of said primers is depicted as SEQ ID NO: 3 and SEQ ID NO: 4.

[0028] In an embodiment of the present invention, the region amplified by said method (the 'amplicon') is between 100 and 1000 base pairs in length. In a further embodiment the amplicon is between 100 and 400 base pairs in length. In a still further embodiment the amplicon is 181 base pairs in length. In a further embodiment the amplicon is produced using the above method in conjunction with the primers of the sequence of SEQ ID NO: 3 and SEQ ID NO: 4, and is 181 base pairs in length. These primers are specific for the COT202 event.

[0029] Alternative primers which may be used in combination to detect the COT202 event include SEQ ID NOs 13 and 14 which are specific for the COT202 event and produce an 86 bp amplicon, and SEQ ID NOs 5 and 6 which are specific for the VIP gene and produce a 556 bp amplicon.

[0030] There are many amplification methods that may be used in accordance with this aspect of the invention. The underlying principle, a known technique to those skilled in the art, is the polymerase chain reaction (PCR). The amplification product from a PCR reaction may be visualised by staining with ethidium bromide and excitation with UV light, typically after size separation using agarose gel electrophoresis.

[0031] An embodiment of the present invention employs variations of the PCR principle such as TaqMan™. This involves labelling at least one of the primers involved in the amplification process with a fluorescent dye. When unbound, the primer adopts a conformation such that no fluorescence can be detected. However, when the primer is bound to a piece of DNA, the conformation changes and fluorescence can be detected. In this way, the amplification process can be monitored in real-time, the intensity of fluorescence corresponding directly to the level of amplification. Suitable primers for use in TaqMan™ PCR are depicted as SEQ ID NOs 13 to 15. These may be used in conjunction with internal control primers such as those depicted as SEQ ID NOs 10 to 12. TaqMan™ analysis may be useful for example, for detecting the presence of the COT202 event in a background of wild type cotton, or for detecting the adventitious presence of COT202 in other germplasm. Further embodiments of the present invention include, but are not limited to, RACE PCR.

[0032] A further embodiment of the present invention involves the use of multiplex PCR for distinguishing between homozygous COT202 plant material and heterozygous COT202 plant material. This is known to those skilled in the art as zygosity testing, and involves the use of three PCR primers which bind to specific parts of the cotton genome and/or inserted DNA. The presence or absence of each of two amplification products of particular sizes indicates whether the test sample is heterozygous or homozygous for COT202. Suitable primers for use in such a zygosity test are depicted as SEQ ID NOs 16 to 18.

[0033] In another aspect of the invention there is provided a method of detecting plant material derived from the COT202 event comprising obtaining a sample for analysis; providing a

probe designed to bind to the complement of a polynucleotide which comprises at least 17 contiguous nucleotides of SEQ ID NO: 1 and/or SEQ ID NO: 2 when said polynucleotide is single stranded; hybridising said probe with the sample; and detecting whether the probe has hybridised. In one embodiment, said probe comprises the sequence of SEQ ID NO: 1 and/or SEQ ID NO: 2. In an embodiment of the present invention there is provided a method of detecting plant material derived from the COT202 event using a probe comprising SEQ ID NO: 7 or SEQ ID NO: 8. In one embodiment, said probe comprises SEQ ID NO: 7. In a further embodiment, said probe consists of SEQ ID NO: 7. In one embodiment, said probe comprises SEQ ID NO: 8. In a further embodiment, said probe consists of SEQ ID NO: 8. The probe may be, for example, a PCR product or restriction digestion fragment. In a further embodiment, the probe as described herein may be tagged with a fluorescent, radioactive, enzymatic or other suitable label to enable hybridisation to be detected. The skilled man will know how to design suitable probes, now that he has the benefit of the present disclosure.

[0034] In a further embodiment of the present invention, there is provided a method of hybridising a probe to the sample under stringent conditions and detecting whether the probe has hybridised. Stringent hybridisation conditions are well known to the skilled man and comprise, for example: hybridisation at a temperature of about 60° C. in a solution containing 6×SSC, 0.01% SDS and 0.25% skimmed milk powder, followed by rinsing at the same temperature in a solution containing 1×SSC and 0.1% SDS. More stringent hybridisation conditions may comprise: hybridisation at a temperature of about 65° C. in a solution containing 6×SSC, 0.01% SDS and 0.25% skimmed milk powder, followed by rinsing at the same temperature in a solution containing 0.2×SSC and 0.1% SDS.

[0035] Suitable techniques for detecting plant material derived from the COT202 event based on the hybridisation principle include, but are not limited to Southern Blots, Northern Blots and in-situ hybridisation. The skilled man is familiar with techniques such as these. Typically, they involve incubating a probe with a sample, washing to remove unbound probe, and detecting whether the probe has hybridised. Said detection method is dependent on the type of tag attached to the probe—for example, a radioactively labelled probe can be detected by exposure to and development of x-ray film. Alternatively, an enzymatically labelled probe may be detected by conversion of a substrate to effect a colour change.

[0036] In a further aspect of the invention there is provided a method of detecting plant material derived from the COT202 event comprising obtaining a sample for analysis; providing an antibody or binding protein designed to bind to a VIP protein contained within a plant comprising at least 17 contiguous nucleotides from SEQ ID NO: 1 and/or SEQ ID NO: 2; incubating said antibody or binding protein with the sample; and detecting whether the antibody or binding protein has bound. In one embodiment of the present invention said VIP protein comprises the sequence of SEQ ID NO: 9.

[0037] Suitable methods of detecting plant material derived from the COT202 event based on said antibody binding include, but are not limited to Western Blots, Enzyme-Linked ImmunoSorbent Assays (ELISA) and SELDI mass spectrometry. The skilled man is familiar with these immunological techniques. Typical steps include incubating a sample with an antibody that binds to the VIP protein, washing to remove

unbound antibody, and detecting whether the antibody has bound. Many such detection methods are based on enzymatic reactions—for example the antibody may be tagged with an enzyme such as horse radish peroxidase, and on application of a suitable substrate, a colour change detected. Suitable antibodies may be monoclonal or polyclonal.

[0038] In another aspect of the invention there is provided a method of detecting plant material derived from the COT202 event comprising obtaining a sample for analysis; making a protein extract of the sample; providing a test strip designed to detect the presence of a VIP protein present within the sample; incubating the test strip with the sample; and detecting whether VIP protein is present. In one embodiment of the present invention said VIP protein comprises the sequence of SEQ ID NO: 9.

[0039] An alternative antibody-based detection method for COT202 uses of dipsticks or test strips. Typical steps include incubating a test strip with a sample and observing the presence or absence of coloured bands on the test strip. The coloured bands are indicative of the presence of a protein in the sample. Such dipstick or test strip tests are protein specific, and may be used for rapid testing of samples in the field.

[0040] In a further aspect of the present invention there is provided a method of detecting plant material derived from the COT202 event comprising obtaining a sample for analysis; subjecting one or more insects of the species *Spodoptera frugiperda* (susceptible to VIP3A) to the sample; subjecting one or more insects of species *Ostrinia nubilalis* (not susceptible to VIP3A) to the sample as a control; detecting whether the sample has an insecticidal effect on insects from each species; and comparing the results with an authentic COT202 bioassay profile. The results are compared against an authentic COT202 bioassay profile that is produced using insects of the same condition (including insect age and culture conditions) which have been subjected to the same dose and type of COT202 plant material (including plant age, plant variety and tissue type) and where the insecticidal effect is detected the same length of time after subjecting the insects to the COT202 sample. Detection of an insecticidal effect may be, for example, an assessment of insect mortality, or of the growth stage of the insects. *Spodoptera frugiperda* is a positive control for COT202 as it is susceptible to a suitable dose of VIP3A, while *Ostrinia nubilalis* is a negative control for COT202 as it is not susceptible to a suitable dose of VIP3A. Alternative insect species that are either susceptible or not susceptible to VIP3A may be substituted in an assay as described above as appropriate, provided that the results are compared against an authentic profile generated using the same insect species.

[0041] In one embodiment of the invention, the method of detecting plant material derived from the COT202 event includes but is not limited to leaf-feeding bioassays in which a leaf or other suitable plant part from the COT202 event or any plant material derived from the COT202 event, is infested with one or more pest insects. Detection may be through assessment of damage to the leaf or plant part after set time periods, assessment of mortality or another insecticidal effect on the insects. Alternative plant parts which may be used for such bioassays include bolls and squares. Such bioassays may, for example, be carried out in the laboratory, field, or glasshouse, and may be subject to natural or artificial insect infestation.

[0042] In another aspect of the invention, there is provided a kit of parts comprising a means for detecting the presence in a sample of plant material derived from the COT202 event. Preferably, said kit of parts comprises a means for detecting the presence in a sample of a polynucleotide comprising at

least 17 contiguous nucleotides from the sequence of SEQ ID NO: 1 and/or SEQ ID NO: 2, or a protein encoded by a polynucleotide as described above, or a VIP protein. In an embodiment of the present invention, said kit of parts may comprise DNA amplification-detection technology such as PCR or TaqMan™. In a further embodiment of the present invention, said kit of parts may comprise probe hybridisation-detection technology such as Southern Blots, Northern Blots or in-situ Hybridisation. In another embodiment of the present invention, said kit of parts may comprise antibody binding-detection technology such as Western Blots, ELISA's, SELDI mass spectrometry or test strips. In a further embodiment of the present invention, said kit of parts may comprise insect bioassay-detection technology such as leaf feeding bioassays or mortality bioassays. Each of these detection technologies may be used as described above. In a further embodiment of the present invention, said kit of parts may comprise any combination of the afore-mentioned detection technologies. In a still further embodiment, said kit of parts may comprise in the form of instructions one or more of the methods described above.

[0043] According to the present invention, there is provided the use of one or more of the polynucleotides of the invention as described above for detecting the COT202 event. In one embodiment, said polynucleotides may be used in a method for detecting the COT202 event as described above.

EXAMPLES

[0044] The invention will be further apparent from the following non-limiting examples in conjunction with the associated sequence listings as described below:

[0045] SEQ ID NO 1: Polynucleotide sequence which extends across the junction where the 5' end of the COT202 insert is inserted into the cotton genome in event COT202.

[0046] SEQ ID NO 2: Polynucleotide sequence which extends across the junction where the 3' end of the COT202 insert is inserted into the cotton genome in event COT202.

[0047] SEQ ID NOs 3-6: Polynucleotide sequences suitable for use as primers in the detection of the COT202 event.

[0048] SEQ ID NOs 7-8: Polynucleotide sequences suitable for use as probes in the detection of the COT202 event.

[0049] SEQ ID NO 9: Amino acid sequence of the VIP3A toxin protein.

[0050] SEQ ID NOs 10-15: Polynucleotide sequences suitable for use as TaqMan™ primers in the detection of the COT202 event.

[0051] SEQ ID NOs 16-18: Polynucleotide sequences suitable for use as primers in the detection of the COT202 event via zygosity testing.

Example 1

Cloning and Transformation

[0052] 1.1 Vector Cloning

[0053] Standard gene cloning techniques of restriction digestion and ligation of fragments from in-house vectors were used to construct the transformation vector, pNOV103. The vector included a selectable marker cassette comprising a Ubiquitin (UBQ3) promoter, the UBQ3 intron, a gene sequence which encodes a protein conferring resistance to hygromycin, and a nos polyadenylation sequence. The vector also included the expression cassette of the target gene, which cassette comprised a Ubiquitin (UBQ3) promoter, the UBQ3 intron, a sequence encoding the VIP3A gene that had been codon optimised for expression in maize, and a nos polyade-

nylation sequence. The selectable marker cassette and VIP3A containing cassette were cloned between the left and right border sequences within different T-DNA regions of vector pNOV103. The vector also comprised a gene conferring resistance to an antibiotic, kanamycin, for prokaryotic selection.

[0054] The vector was transformed into *Agrobacterium tumefaciens* strain GV3101 using standard *Agrobacterium* transformation techniques, and transformed cells selected through their resistance to kanamycin.

[0055] 1.2 Plant Transformation

[0056] The COT202 event was produced by *Agrobacterium*-mediated transformation of *Gossypium hirsutum* L. cv Coker 312.

[0057] Coker 312 seeds were sown in the glasshouse. Tender petioles were cut from 3 to 5 weeks old plants, and sterilized by immersion in 70% ethanol. The petioles were then immersed in a 5% Clorox+2 ml/L Tween 20 solution for 20 minutes. Petioles were washed 3 times in ddH₂O. The ends of petioles were cut off, and petioles transferred to petiole pre-culture medium (4.3 g/L MS salts, 200×B5 vitamins, 30 g/L glucose, 2.4 g/L phytogel, pH 7.0) and allowed to pre-culture in the light at 30° C. for 3 days.

[0058] 2 ml cultures of *Agrobacterium* containing the pNOV103 construct were grown overnight in appropriate antibiotics and then diluted with liquid MMS1 medium (4.3 g/L MS salts, 200×B5 vitamins, 0.05 mg/L 2,4-D, 0.1 mg/L kinetin, 30 g/L glucose, pH 6.5) to an OD₆₆₀ of between 0.1 and 0.2.

[0059] The ends were cut off the petioles and placed in 3 to 5 ml of bacterial solution in a sterile petri dish. Once in the solution, the petioles were cut lengthwise and then cut into 2 cm sections. After the petiole explants had soaked in bacterial solution for 5 to 10 minutes, they were transferred to co-culture plates, and allowed to co-culture at 24° C. for 48 hours under low light intensity. Co-cultured explants were transferred to MMS1 medium (recipe as for MMS1 liquid medium, additionally with 2.4 g/L phytogel) containing 500 mg/L Cefotaxime and 10 mg/L Hygromycin, and incubated at 30° C. under a light cycle of 16 hours light and 8 hours dark. Explants were transferred to fresh medium after 2 weeks, and every 4 to 6 weeks thereafter until callus was formed.

[0060] Once calli were the size of a garden pea, they were removed from the explants and transferred to fresh MMS1 medium containing 500 mg/L Cefotaxime and 10 mg/L Hygromycin, and maintained in tissue culture by subculturing every 4 weeks as appropriate.

[0061] 1.5 g callus tissue was broken up thoroughly and placed in a 50 ml Erlenmeyer flask containing 10 ml of liquid MMS2 medium (4.3 g/L MS salts, 200×B5 vitamins, 1.9 g/L KNO₃, 30 g/L glucose, pH 6.5). The suspended callus was shaken at 100 rpm in the light at 30° C. until small white slightly round cell clusters were visible. These clusters indicate that the tissue is embryogenic. The suspension culture cells were rinsed 3 times in MMS2 liquid medium, resuspended and plated onto solid MMS2 medium (recipe as per liquid MMS2 medium, additionally with 2.4 g/L phytogel). Once plated, excess liquid MMS2 medium was removed, and the plates incubated at 30° C. in the light. Plates were checked for somatic embryo development each week. Somatic embryos formed within 1 to 2 months. Somatic embryos were transferred to EG (embryoid germination) medium (10×EG stock (consisting of 1×10 L pack of Musashige and Skoog Modified Basal Salt Mixture (Sigma), 19 g KNO₃, 50 ml 200×B5 vitamins, water to 1 L), 1 g/L glutamine, 0.5 g/L asparagine, recipe), and sub-cultured to fresh EG medium every 3 to 4 weeks.

[0062] Once somatic embryos turned green and were larger than 2 cm, they were plated root down in EG medium. At all stages of regeneration, growing plantlets were prevented from reaching the lids or sides of their containers. Germinated embryos with 1 to 2 true leaves were transferred to EG medium in 175 ml Greiners. Strong plantlets with true leaves were transferred to sterile peat plugs expanded with dH₂O in 175 ml Greiners and transferred to a growth cabinet under conditions of 14 hours daylight at 30° C. and 10 hours darkness at 20° C. Thereafter, plantlets were transplanted into pots and grown in the glasshouse.

[0063] 1.3 Identification and Selection of Transgenics

[0064] Putative transgenic plants were screened by PCR for the presence of the VIP3A gene. Positive events were identified and screened using insect bioassays for insecticidal activity against Fall Armyworm (*Spodoptera frugiperda*) (see Example 7). Insecticidal lines were characterized for copy number by TaqMan™ analysis (see Example 2). T1 seed from several events were observed in a field trial for insect resistance and agronomic quality. Two events, COT202 and COT203, were chosen based on having a single copy of the transgene, good protein expression as identified by ELISA (see Example 4), good insecticidal activity against Cotton Boll Worm (*Helicoverpa zea*) and field performance. The hygromycin selectable marker cassette was segregated away using conventional plant breeding to result in the COT202 event and the COT203 event.

[0065] 1.4 Verification of Sequence of COT202

[0066] Genomic DNA was isolated from the COT202 event. This was used in the sequencing of the junctions of the DNA insertion site with the cotton genomic DNA in the COT202 event, using standard DNA sequencing techniques.

Example 2

COT202 Event Specific Detection via TaqMan™

[0067] 2.1 DNA Extraction

[0068] DNA was extracted from leaf tissue using the Wizard™ Magnetic 96 DNA Plant System (Promega, #FF3760), according to the manufacturers instructions, with an additional step at the beginning of the protocol: following grinding of the leaf material, 0.9 ml Cotton Extraction Buffer (0.2M Tris pH 8.0, 50 mM EDTA, 0.25M NaCl, 0.1% v/v 2-mercaptoethanol, 2.5% w/v polyvinyl-pyrrolidone) was added to each well, the plant tissue resuspended and the plate centrifuged at 4,000 rpm (2755 g) for 10 minutes. After aspirating and discarding the supernatant, 300 ul Lysis Buffer A (Promega) was added and the manufacturers protocol was followed from this point. This procedure resulted in approximately 85 ul of purified genomic DNA at a concentration of approximately 10 ng/ul.

[0069] 2.2 TaqMan™ PCR Reactions

[0070] TaqMan™ PCR reactions were setup using a standard reaction mix comprising:

[0071] 5 ul 2× Jumpstart Master Mix for Q-PCR (Sigma, #P2893), supplemented with 15 mM MgCl₂ and 200 nM Strata-ROX

[0072] 0.2 ul 50× FAM primer/probe mix

[0073] 0.2 ul 50× VIC primer/probe mix

[0074] 1.6 ul Water.

[0075] 50× primer/probe mixes comprised 45 ul of each primer at a concentration of 1 mM, 50 ul of the probe at a concentration of 100 uM and 860 ul nuclease free water, and were stored in an amber tube at 4° C. Examples of suitable primer/probe sequence combinations which were used are:

Primer Name	Primer Sequence 5'-3'	SEQ ID
GhCHI2b-F Forward	GGTCCCTGGATACGGTGTCA	SEQ ID NO: 10
GhCHI2b-R Reverse	TTGAGGGTTGGATCCTTTGC	SEQ ID NO: 11
GhCHI2bNEW-VIC Probe	CACCAACATCATCAATGGTGGCATCG (5' label = VIC, 3' label = TAMRA)	SEQ ID NO: 12
COT202-F Forward	GGAATGTGGCGAATGGTGAT	SEQ ID NO: 13
COT202-R Reverse	TGTCGTTTCCCGCCTTCA	SEQ ID NO: 14
COT202-FAM Probe	CAAATGCCCATTTTCATTCACAAAAGC (5' label = FAM, 3' label = TAMRA)	SEQ ID NO: 15

[0076] 7 μ l of master mix was dispensed into each well of a 384-well TaqMan™ assay plate. 3 μ l DNA template was added to the appropriate wells. 3 μ l of copy control dilution series was added to specific wells as a control. The reactions were run in an ABI7900HT (Applied Biosystems) using the following cycling conditions:

Step	Temperature	Time
1	50° C.	2 min
2	95° C.	10 min
3	95° C.	15 sec
4	60° C.	1 min
5	Goto step 3, repeat 40 times	

[0077] Data was analysed using SDS2.0 version A, software (Applied Biosystems).

Example 3

COT202 Detection via Multiplex PCR Zygosity Test

[0078] 3.1 Genomic DNA Extraction

[0079] Genomic DNA from COT202 was extracted as described in Example 2.1.

[0080] 3.2 Multiplex PCR

[0081] PCR primers were designed to bind to cotton genomic DNA sequence upstream of the site at which the COT202 cassette inserted (SEQ ID NO: 16); the cotton genomic DNA sequence downstream of the site at which the COT202 cassette inserted (SEQ ID NO: 17); and the COT202 cassette sequence itself (SEQ ID NO: 18). A 25 μ l PCR reaction was set up for each sample to be tested as follows:

[0082] 1 $\times$ JumpState ReadyMix REDTaq PCR (Sigma P-1107)

[0083] 0.5 μ M primer 1 (SEQ ID NO: 16)

[0084] 0.5 μ M primer 2 (SEQ ID NO: 17)

[0085] 0.5 μ M primer 3 (SEQ ID NO: 18)

[0086] 0.2% BSA

[0087] 20 ng genomic DNA

[0088] ddH₂O to 25 μ l

[0089] The PCR reactions were heated in a thermocycler at 94° C. for 5 minutes, followed by 30 cycles as follows: 94° C.

for 30 seconds, 55° C. for 45 seconds, 72° C. for 1 minute. The reaction was completed by heating at 72° C. for 5 minutes.

[0090] 3.3 Analysis

[0091] PCR reactions were run on an agarose gel, and DNA bands visualised under UV light after staining with ethidium bromide. The presence of 2 bands indicated that the sample was from a COT202 heterozygote plant; 1 band of 181 bp in size indicated that the sample was from a COT202 homozygote plant; and 1 band of approximately 400 bp in size indicated that the sample was from a homozygote wild type cotton plant.

[0092] 3.4 COT202 Detection via Standard PCR

[0093] As an alternative to the multiplex PCR, the COT202 event can be detected in a simple PCR reaction using the primers depicted as SEQ ID NO: 3 and 4, SEQ ID NO: 13 and 14, or SEQ ID NO: 16 and 18. The composition of the PCR reaction mixture is the same as described in example 3.2 above. The PCR reactions are heated in a thermocycler at 94° C. for 5 minutes, followed by 30 cycles as follows: 94° C. for 30 seconds, 55° C. for 30 seconds, 72° C. for 20 seconds. The reaction is completed by heating at 72° C. for 5 minutes. A DNA fragment of 181 bp, 86 bp or 181 bp in size respectively indicates the presence of the COT202 event.

Example 4

COT202 Detection via Southern Blot

[0094] 4.1 DNA Extraction for Use in Southern Blotting

[0095] Approximately 2 to 3 g fresh weight of young leaf tissue was ground in a chilled mortar and pestle to a fine powder and added to 15 ml of ice-cold Nuclei extraction buffer (0.35M glucose, 0.1 M Tris-HCl pH8, 50 mM Na₂EDTA, 2% Polyvinyl-pyrrolidone-10, 0.1% ascorbic acid, 0.2% B-mercaptoethanol) in a labelled tube. The sample was incubated on ice for 15-20 minutes. The tube was mixed gently and centrifuged at 2700 g for 20 minutes at 4° C. The supernatant was discarded and 8 ml of nuclei lysis buffer (0.14M sorbitol, 0.22M Tris-Cl pH8, 0.8M NaCl, 0.22M Na₂EDTA, 0.8% w/v CTAB, 1% Sarkosyl, 1% Polyvinyl-pyrrolidone-10, 0.1% ascorbic acid, 0.2% B-mercaptoethanol, 5 μ g/ml proteinase K) was added. After mixing, the tubes were incubated at 65° C. for 30 minutes. 10 ml chloroform was added, and the tube mixed gently by inversion until an

emulsion formed followed by centrifugation at 4600 rpm for 10 minutes at room temperature.

[0096] The aqueous layer was removed into a new tube containing 10 μ l RNase A (10 mg sigma R4642), and the tube incubated for 30 minutes at 37° C. The chloroform and centrifugation steps were repeated once. The aqueous layer was removed into a new tube containing 10 ml propan-2-ol. After approximately 15 minutes incubation at room temperature, a gelatinous precipitate was observed in the middle of the tube. The tube was mixed gently to precipitate out the DNA. The DNA was spooled out using a sterile loop into a falcon tube containing 70% ethanol. The DNA was air-dried to remove the ethanol and resuspended in 200-400 μ l TE.

[0097] 4.2 Alternative Method for DNA Extraction

[0098] 2-3 young cotton leaves (approximately 1 g fresh weight) are ground to a paste in a mortar and pestle at room temperature, with 2 ml of grinding buffer (100 mM NaOAc pH 4.8, 50 mM EDTA pH8.0, 500 mM NaCl, 2% PVP (10,000 MW), 1.4 % SDS) and a little sand. The ground tissue is transferred to a 15 ml falcon tube, and the remnants in the mortar rinsed with 1 ml of grinding buffer into the tube. The sample is incubated at 65° C. for 15 minutes, shaking occasionally. 4 ml 10M ammonium acetate is added, and the sample mixed well and incubated at 65° C. for 10 minutes to precipitate proteins. The samples are incubated at room temperature at 4600 rpm for 10 minutes. The aqueous phase is transferred to a fresh 15 ml tube.

[0099] 0.6 volumes of cold isopropanol are added and the sample is incubated at room temperature for approximately 30 minutes. After mixing by slowly inverting the tube several times, the DNA is spooled out and dissolved in 500 μ l TE. 10 μ l of 10 mg/ml RNase are added and incubated for 15 minutes at room temperature. Following extraction with 500 μ l of phenol:chloroform:isoamyl alcohol (25:24:1), the sample is mixed gently and centrifuged at 13000 rpm for 5 min.

[0100] The supernatant is transferred to a fresh tube using a fine Pasteur pipette and re-extracted with chloroform:isoamyl alcohol (24:1) as above. The supernatant is transferred to fresh tubes, $\frac{1}{10}$ volume 3M NaOAc (pH4.8) added and mixed, and then one volume cold isopropanol is added. The sample may be incubated at room temperature for up to 30 minutes to precipitate the DNA. The DNA is spooled out and resuspended in 70% ethanol. The DNA is air-dried to remove the ethanol and resuspended in 200 μ l water.

[0101] 4.3 Restriction Enzyme Digests

[0102] The DNA was quantified using a spectrophotometer and running out on a gel. Suitable enzyme digests were prepared using 5 μ g DNA per digest in a total volume of 40 μ l. Digests included HindIII, XmaI, BamHI, NheI, and SacI, both alone and in combination. In particular, a HindIII and XmaI double digest was used to detect the intactness of the VIP3A gene; a NheI digest was used to detect VIP3A locus number. Digests were incubated for 6 hours at the appropriate temperature for each enzyme.

[0103] 4.4 Gel Electrophoresis

[0104] Bromophenol blue loading dye was added to each sample from 4.2 above, and each sample loaded on a 0.8% TBE agarose gel. The gel was run at 50 volts overnight.

[0105] After running, the gel was washed in 0.25M HCl for 10 minutes to depurinate the DNA, incubated in denaturing solution (0.5M NaOH, 1.5M NaCl) with gentle agitation for 30 minutes, rinsed with distilled water and then incubated in neutralising solution (0.5M Tris, 1.5M NaCl) for 30 minutes.

[0106] A Southern Blot was prepared as follows: A glass plate was placed over a tray containing 20 $\times$ SSC and a strip of 3M paper was placed onto the glass plate such that both ends dipped into the 20 $\times$ SSC solution (to act as a wick). A piece of

3M paper the same size as the gel was placed on the wick, and the gel placed on this. Strips of nescofilm were laid around the edges of the gel to form a seal. A Hybond membrane was placed on top of the gel, followed by two further pieces of 3M paper. Throughout the assembly of the blot, care was taken to ensure that no air bubbles were trapped between the membrane, gel and 3M paper. A 5 cm-10 cm stack of absorbent paper towels was placed on top of the 3M paper and held in place with a weight.

[0107] The DNA was allowed to transfer to the Hybond membrane overnight. After transfer the Southern Blot stack was disassembled and the DNA was bound to the membrane via UV cross-linking.

[0108] 4.5 Hybridisation

[0109] A suitable DNA probe was prepared by PCR or restriction digest of binary plasmid. 25 ng probe DNA in 45 μ l TE was boiled for 5 minutes, placed on ice for 5 minutes then transferred to a Rediprime II (Amersham Pharmacia Biotech, #RPN1633) tube. After addition of 5 μ l P32-labelled dCTP to the Rediprime tube, the probe was incubated at 37° C. for 1 hour. The probe was purified by centrifugation through a microspin G-50 column (Amersham Pharmacia Biotech, #27-5330-01) according to the manufacturers instructions to remove unincorporated dNTPs. The activity of the probe was measured roughly by comparing the amount of radioactive component remaining in the column to the amount in the sample tube, with a ratio of at least 50:50 being acceptable. The Hybond membrane was pre-hybridised by wetting with 40 ml pre-warmed Rapid-Hyb buffer (Amersham-Pharmacia), at 65° C. for 30 minutes. The labelled probe was boiled for 5 minutes, and placed on ice for 5 minutes. An appropriate amount of probe (1 million counts per 1 ml pre-hybridisation buffer) was added to the pre-hybridisation buffer and hybridisation occurred at 65° C. overnight. The following day, the hybridisation buffer was discarded, and following a rinse with 50 ml 2 $\times$ SSC/1% SDS solution the membrane washed in 150 ml 2 $\times$ SSC/1% SDS solution at 65° C. for 30-45 minutes. This process was repeated twice with 0.1 $\times$ SSC/1% SDS solution. The membrane was exposed to a phosphor screen or X-ray film to detect where the probe had bound.

Example 5

COT202 Detection via ELISA

[0110] 5.1 Protein Extraction

[0111] Cotton tissue for analysis was harvested and frozen at -70° C. Fresh tissue was ground to a fine powder and weighed into a labelled polypropylene tube. Extraction buffer (100 mM Tris, 100 mM Sodium Borate, 5 mM MgCl, 0.05% Tween 20, 0.2% Sodium Ascorbate, Water, pH 7.8, 1 mM AEBSE, 0.001 mM Leupeptin) was added to the sample in a ratio of 2:1 (volume extraction buffer:sample fresh weight) for fresh tissue or 30:1 (volume extraction buffer:sample dry weight) for lyophilised tissue. The sample was vortexed and homogenised using a Brinkman PT 10/35 Polytron equipped with a PTA 10TS foam-reducing generator, until the mixture became liquefied. Extracts were centrifuged at 10,000 $\times$ g for 15 minutes. The protein extract supernatant was stored at 2-8° C.

[0112] 5.2 ELISA Protocol

[0113] The ELISA procedure used standard techniques as follows. A 96-well plate was soaked in ethanol for 2 hours, and air-dried. The plate was coated with 50 μ l goat anti-VIP3A antibody per well and incubated overnight at 2-8° C. After washing three times with 1 $\times$ ELISA wash solution (100 mM Tris, 0.5% Tween-20, 75 mM NaCl, pH8.5), the plate was dried briefly by tapping upside down on a paper towel.

150 ul blocking solution (10 mM NaPO₄, 140 mM NaCl, 1% BSA, 0.02% Sodium Azide, titrated to pH7.4 with monobasic NaPi and dibasic NaPi) was added to each well followed by incubation at room temperature for 45 minutes. The plate was washed 3 times as described above.

[0114] VIP3A standards and protein extract samples were applied to appropriate wells of the plate in triplicate, 50 ul total volume per well. The plate was incubated at 2-8° C. for 1 hour 30 minutes, followed by room temperature for a further 30 minutes. The plate was washed three times with ELISA wash solution, and then incubated at 35-39° C. for 1 hour with 50 ul rabbit anti-VIP3A antibody per well. The plate was washed three times with ELISA wash solution, and incubated at room temperature for 30 minutes with 50 ul donkey anti-rabbit alkaline phosphatase per well. Following a further three washes with ELISA wash solution, 50 ul phosphatase substrate solution was added per well and the plate incubated for 30 minutes at room temperature. The reaction was stopped by addition of 50 ul 3M NaOH per well. The absorbance of the solution in each well was measured at 405 nm using a Ceres 900C multiwell plate reader and the results analysed using KC3 Curve fitting software (Bio-Tek Instruments Inc.). The concentration of VIP3A in the samples was calculated by reference to the VIP3A protein standards.

Example 6

COT202 Detection Via DipStick

[0115] 6.1 Protein Extraction

[0116] A piece of leaf tissue approximately 2 cm² was placed in a tube containing extraction buffer. A plastic stirrer was used to extract protein from the tissue, by cutting into and mascerating the tissue.

[0117] 6.2 Dipstick Test

[0118] A test strip was placed into the tube and incubated for 5 to 10 minutes for the result to develop. The test strip comprised a first band at which anti-VIP3A antibody was bound, and a second band at which a control antibody was bound. After incubation, a double red line in the result window of the test strip indicated that VIP3A was present. The lower line indicated the presence of VIP3A protein while the upper line was a control indicating that the assay was working correctly.

Example 7

COT202 Detection Via Insect Bioassay

[0119] 7.1 Leaf Biosassays

[0120] Leaf assays were performed on Fall Army Worm (*Spodoptera frugiperda*), Cotton Boll Worm (*Helicoverpa zea*) and Tobacco Budworm (*Heliothis virescens*) as follows: Pads were soaked with 300 ul to 500 ul distilled water and placed into Gelman dishes. Leaf pieces measuring between approximately 0.5 square inches and 0.75 square inches were excised from cotton plants 8 to 12 inches in height, and placed on the pads. Between 8 and 10 insect larvae were placed in each dish and a lid fitted. The dishes were incubated at 28° C. On the third and sixth days after infestation, damage to the leaf in each dish was scored and compared with the control plants.

[0121] 7.2 Boll Bioassays

[0122] Four absorbent pads were saturated with water and placed inside a large plastic cup. Three extra thick glass filters, each soaked with 100 ul distilled water, were placed in a smaller plastic cup, which was then seated inside the larger cup. A 1.25 inch long boll was excised, immersed in 10 mg/ml to 20 mg/ml Nystatin and placed on the filters in the small cup.

50 insect larvae were placed on the square or boll and a lid attached to the larger cup. The squares or bolls were re-infested with 50 more larvae after 7 days.

[0123] The experiment was incubated at room temperature for approximately 3 weeks. The bolls were then cut open to determine damage. Damage to the boll was compared to the control samples.

[0124] 7.3 Lyophilised Leaf Bioassays

[0125] Bioassays using freeze-dried leaf tissue were performed on *Heliothis virescens* as follows:

[0126] Terminal leaves were snap-frozen on dry-ice at time of picking and lyophilised overnight. The freeze dried tissue was ground in a mortar and pestle to a fine powder and resuspended in 0.2% agar solution to make an 8% (0.08 g/ml) suspension of leaf powder. The suspension was overlaid on top of artificial insect diet in 96-well plates and left to dry. A single neonate insect larva was introduced into each well and the plates sealed. The plates were incubated at 28° C. On the sixth day after infestation, larval mortality was scored and compared with control samples.

Example 8

COT202 Field Trials

[0127] 8.1 Field Trial Design

[0128] The efficacy of the COT202 event against *Heliothis virescens* (Tobacco Bud Worm) and *Helicoverpa zea* (Cotton Boll Worm) was tested by conducting field trials at three locations in the US, namely Leland (Miss.), Quitman (Ga.) and Beasley (Tex.). Trials in each location were set up using a randomised complete block design, with four entry plots comprising four rows of 40 feet in length and four repetitions per trial. Seed was planted to obtain a plant stand of approximately 3 plants per foot of row length. Each field trial included non transgenic Coker 312 plants for control purposes, and two other transgenic events designated event A and event B for comparison purposes.

[0129] 8.2 Field Trial Assessment

[0130] An assessment of the natural insect populations was made at each trial location at the first white flower stage, approximately 80 days after planting. Where insect pressure was below the US economic threshold of 10% damage, artificial infestation of Cotton Boll Worm and Tobacco Bud Worm was made. The artificial infestation method was designed to obtain a rate of 10 eggs per foot per insect species. Assessment of damage to cotton squares and bolls was made by visual inspection of 50 fruiting forms per plot at 5-7 days after artificial infestation. When relying on a natural infestation, damage ratings were made when the non transgenic Coker 312 control plants showed fruiting form damage above the economic threshold level of 10% in all control plots.

[0131] 8.3 Field Trial Results

[0132] The results presented below show percentage damage to cotton squares and bolls at each field trial location, for each plant category. The data below represents an average of 200 fruiting forms (squares or bolls) per event per trial.

[0133] Location: Leland, Miss.

	Square Damage	Boll damage
Control	77	No data
COT202 event	6.6	No data
Event A	3.6	No data
Event B	29.6	No data

[0134] Location: Quitman, Ga.

[0135] Location: Beasley, Tex., USA

	Square Damage	Boll damage
Control	80	52
COT202 event	2	0
Event A	1.5	0
Event B	24.5	2

	Square Damage	Boll damage
Control	24.4	13.2
COT202 event	4	2
Event A	2	1.2
Event C	5.2	3.2

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 18

<210> SEQ ID NO 1

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: COT202 nucleotide motif

<400> SEQUENCE: 1

aacaacacaca aaatcttttc accagt 26

<210> SEQ ID NO 2

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: COT202 nucleotide motif

<400> SEQUENCE: 2

ttcccgctt cagatcttct gcaaca 26

<210> SEQ ID NO 3

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: COT202 nucleotide motif

<400> SEQUENCE: 3

ggtgtccatc gggtagtcca taa 23

<210> SEQ ID NO 4

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: COT202 nucleotide motif

<400> SEQUENCE: 4

tctatgttac tagatcgga attg 24

<210> SEQ ID NO 5

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: COT202 nucleotide motif

<400> SEQUENCE: 5

-continued

gatcggggtc aggaaggtct 20

<210> SEQ ID NO 6
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: COT202 nucleotide motif

<400> SEQUENCE: 6

cagcatcatg aacgagcact 20

<210> SEQ ID NO 7
 <211> LENGTH: 290
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: COT202 nucleotide motif

<400> SEQUENCE: 7

gacaaggaca gcttgagcga ggtgatctac ggcgacatgg acaagctgct gtgtccggac 60

cagagcgagc aaatctacta caccaacaac atcgtgttcc cgaacgagta cgtgatcacc 120

aagatcgact tcaccaagaa gatgaagacc ctgcgctacg aggtgaccgc caacttctac 180

gacagcagca ccggcgagat cgacctgaac aagaagaagg tggagagcag cgaggccgag 240

taccgcacc tgagcgcgaa cgacgacggc gtctacatgc cactgggct 290

<210> SEQ ID NO 8
 <211> LENGTH: 4382
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: COT202 nucleotide motif

<400> SEQUENCE: 8

gatttggagc caagtctcat aaacgccatt gtggaagaaa gtcttgagtt ggtggtaatg 60

taacagagta gtaagaacag agaagagaga gagtgtgaga tacatgaatt gtcgggcaac 120

aaaaatcctg aacatcttat tttagcaaa agaaagagtt ccgagtctgt agcagaagag 180

tgaggagaaa tttaagctct tggacttggt aattgttccg cctcttgaat acttcttcaa 240

tctcatata ttcttcttct atgttacctg aaaaccggca tttaatctcg cgggtttatt 300

ccggttcaac attttttttg ttttgagtta ttatctgggc ttaataacgc aggcctgaaa 360

taaatccaag gcccaactgt tttttttttt aagaagttgc tgttaaaaaa aaaaaaagg 420

aattaacaac aacaacaaaa aaagataaag aaaataataa caattacttt aattgtagac 480

taaaaaaaca tagattttat catgaaaaaa agagaaaaga aataaaaaact tggatcaaaa 540

aaaaacatac agatcttcta attattaact tttcttaaaa attaggtcct ttttcccaac 600

aattagggtt agagtttttg aattaaccca aaaagattgt tctaaaaaat actcaaat 660

ggtagataag tttccttatt ttaattagtc aatggtagat actttttttt cttttcttta 720

ttagagtaga ttagaatctt ttatgccaa tattgataaa ttaaatcaag aagataaact 780

atcataatca acatgaattt aaaagaaaaa tctcatatat agtattagta ttctctatat 840

atattatgat tgcttattct taatgggttg ggtaaccaa gacatagtct taatggaaag 900

aatctttttt gaactttttc cttattgatt aaattcttct atagaaaaga aagaaattat 960

-continued

ttgaggaaaa	gtatatataca	aaagaaaaat	agaaaaatgt	cagtgaagca	gatgtaatgg	1020
atgacctaat	ccaaccacca	ccataggatg	tttctacttg	agtcgggtctt	ttaaaaacgc	1080
acgggtggaaa	atatgacacg	tatcatatga	ttccttcctt	tagtttcgtg	ataataatcc	1140
tcaactgata	tcttcctttt	tttgttttgg	ctaaagatat	tttattctca	ttaatagaaa	1200
agacggtttt	gggcttttgg	tttgcgatat	aaagaagacc	ttcgtgtgga	agataataat	1260
tcatcctttc	gtctttttct	gactcttcaa	tctctcccaa	agcctaaagc	gatctctgca	1320
aatctctcgc	gactctctct	ttcaaggat	atcttctgat	tctttttggt	tttgattcgt	1380
atctgatctc	caatttttgt	tatgtggatt	attgaatctt	ttgtataaat	tgcttttgac	1440
aatattgttc	gtttcgtcaa	tccagcttct	aaattttgtc	ctgattacta	agatatcgat	1500
tcgtagtggt	tacatctgtg	taatttcttg	cttgattgtg	aaattaggat	tttcaaggac	1560
gatctattca	atctttgtgt	tttctttgtt	cgattctctc	tgtttttaggt	ttcttatgtt	1620
tagatccgtt	tctcttttgt	gttgttttga	tttctcttac	ggcttttgat	ttggatatatg	1680
ttcgtctgatt	ggtttctact	tggtctattg	ttttatttca	ggcggatcca	ccatgaacaa	1740
gaacaacacc	aagctgagca	cccgcgcct	gccgagcttc	atcgactact	tcaacggcat	1800
ctacggcttc	gccaccggca	tcaaggacat	catgaacatg	atcttcaaga	ccgacaccgg	1860
cggcgacctg	accctggacg	agatcctgaa	gaaccagcag	ctgctgaacg	acatcagcgg	1920
caagctggac	ggcgtgaacg	gcagcctgaa	cgacctgatc	gccagggca	acctgaacac	1980
cgagctgagc	aaggagatcc	ttaagatcgc	caacgagcag	aaccaggtgc	tgaacgacgt	2040
gaacaacaag	ctggacgccca	tcaacaccat	gctgcgcgtg	tacctgccga	agatcaccag	2100
catgctgagc	gacgtgatga	agcagaacta	cgccctgagc	ctgcagatcg	agtacctgag	2160
caagcagctg	caggagatca	gcgacaagct	ggacatcatc	aacgtgaacg	tctgatcaa	2220
cagcaccctg	accgagatca	ccccggccta	ccagcgcata	aagtacgtga	acgagaagtt	2280
cgaagagctg	accttcgccca	ccgagaccag	cagcaagggtg	aagaaggacg	gcagcccggc	2340
cgacatcctg	gacgagctga	ccgagctgac	cgagctggcg	aagagcgtga	ccaagaacga	2400
cgtggacggc	ttcgagtctt	acctgaacac	cttcacgac	gtgatggtgg	gcaacaacct	2460
gttcggccgc	agcgccttga	agaccgccag	cgagctgatc	accaaggaga	acgtgaagac	2520
cagcggcagc	gaggtgggca	acgtgtacaa	cttcctgatc	gtgctgaccg	cctgcaggc	2580
ccaggccttc	ctgaccttga	ccacctgtcg	caagctgctg	ggcctggccg	acatcgacta	2640
caccagcatc	atgaacgagc	acttgaacaa	ggagaaggag	gagttccgcg	tgaacatcct	2700
gccgaccctg	agcaaacctt	tcagcaaccc	gaactacgcc	aaggtgaagg	gcagcgacga	2760
ggacgccaa	atgatcgtgg	aggctaagcc	gggccacgcg	ttgatcggct	tcgagatcag	2820
caacgacagc	atcaccgtgc	tgaagggtga	cgaggccaag	ctgaagcaga	actaccaggt	2880
ggacaaggac	agcttgagcg	agggtgatcta	cggcgacatg	gacaagctgc	tgtgtccgga	2940
ccagagcgag	caaatctact	acaccaacaa	catcgtgttc	ccgaacgagt	acgtgatcac	3000
caagatcgac	ttaccaaga	agatgaagac	cctgcgctac	gaggtgaccg	ccaacttcta	3060
cgacagcagc	accggcgaga	tcgacctgaa	caagaagaag	gtggagagca	gcgaggccga	3120
gtaccgcacc	ctgagcgcga	acgacgacgg	cgtctacatg	ccactgggcg	tgatcagcga	3180
gaccttcctg	accccgatca	acggcttttg	cctgcaggcc	gacgagaaca	gccgcctgat	3240

-continued

```

caccctgacc tgtaagagct acctgcgcga gctgctgcta gccaccgacc tgagcaacaa 3300
ggagaccaag ctgatcgtgc caccgagcgg cttcatcagc aacatcgtgg agaacggcag 3360
catcgaggag gacaacctgg agccgtggaa ggccaacaac aagaacgcct acgtggacca 3420
caccggcggc gtgaacggca ccaaggccct gtacgtgcac aaggacggcg gcatcagcca 3480
gttcatcggc gacaagctga agccaagac cgagtacgtg atccagtaca cctgaagggg 3540
caagccatcg attcacctga aggacgagaa caccggctac atccactacg aggacaccaa 3600
caacaacctg gaggactacc agaccatcaa caagcgcttc accaccggca ccgacctgaa 3660
gggcgtgtac ctgatcctga agagccagaa cggcgacgag gcctggggcg acaacttcat 3720
catcctggag atcagcccga gcgagaagct gctgagcccg gagctgatca acaccaacaa 3780
ctggaccagc accggcagca ccaacatcag cggcaacacc ctgaccctgt accagggcgg 3840
ccgcggcatc ctgaagcaga acctgcagct ggacagcttc agcacctacc gcgtgtactt 3900
cagcgtgagc ggcgacgcca acgtgcgcat ccgcaactcc cgcgaggtgc tgttcgagaa 3960
gaggtagatg agcggcgcca aggacgtgag cgagatgttc accaccaagt tcgagaagga 4020
caacttttac atcgagctga gccagggcaa caacctgtac ggcgggccga tcgtgcactt 4080
ctacgacgtg agcatcaagt aggagctcta gatccccgga atttccccga tcgttcaaac 4140
atttggcaat aaagtttctt aagattgaat cctgttgccg gtcttgcgat gattatcata 4200
taattttctg tgaattacgt taagcatgta ataattaaca tgtaatgcat gacgttat 4260
atgagatggg tttttatgat tagagtcccg caattataca tttaatacgc gatagaaaac 4320
aaaatatagc gcgcaaaacta ggataaatta tcgcgcgcgg tgtcatctat gttactagat 4380
cg 4382

```

```

<210> SEQ ID NO 9
<211> LENGTH: 789
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: VIP3A protein motif

```

```

<400> SEQUENCE: 9

```

```

Met Asn Lys Asn Asn Thr Lys Leu Ser Thr Arg Ala Leu Pro Ser Phe
1             5             10             15

Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile Lys Asp
20            25            30

Ile Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asp Leu Thr Leu
35            40            45

Asp Glu Ile Leu Lys Asn Gln Gln Leu Leu Asn Asp Ile Ser Gly Lys
50            55            60

Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Ile Ala Gln Gly Asn
65            70            75            80

Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn Glu Gln
85            90            95

Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asp Ala Ile Asn Thr
100           105           110

Met Leu Arg Val Tyr Leu Pro Lys Ile Thr Ser Met Leu Ser Asp Val
115           120           125

Met Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu Ser Lys
130           135           140

```

-continued

Gln Leu Gln Glu Ile	Ser Asp Lys Leu Asp	Ile Ile Asn Val Asn Val
145	150	155 160
Leu Ile Asn Ser Thr	Leu Thr Glu Ile Thr	Pro Ala Tyr Gln Arg Ile
165	170	175
Lys Tyr Val Asn Glu	Lys Phe Glu Glu Leu	Thr Phe Ala Thr Glu Thr
180	185	190
Ser Ser Lys Val Lys	Lys Asp Gly Ser Pro	Ala Asp Ile Leu Asp Glu
195	200	205
Leu Thr Glu Leu Thr	Glu Leu Ala Lys Ser	Val Thr Lys Asn Asp Val
210	215	220
Asp Gly Phe Glu Phe	Tyr Leu Asn Thr Phe	His Asp Val Met Val Gly
225	230	235 240
Asn Asn Leu Phe Gly	Arg Ser Ala Leu Lys	Thr Ala Ser Glu Leu Ile
245	250	255
Thr Lys Glu Asn Val	Lys Thr Ser Gly Ser	Glu Val Gly Asn Val Tyr
260	265	270
Asn Phe Leu Ile Val	Leu Thr Ala Leu Gln	Ala Gln Ala Phe Leu Thr
275	280	285
Leu Thr Thr Cys Arg	Lys Leu Leu Gly Leu	Ala Asp Ile Asp Tyr Thr
290	295	300
Ser Ile Met Asn Glu	His Leu Asn Lys Glu	Lys Glu Glu Phe Arg Val
305	310	315 320
Asn Ile Leu Pro Thr	Leu Ser Asn Thr Phe	Ser Asn Pro Asn Tyr Ala
325	330	335
Lys Val Lys Gly Ser	Asp Glu Asp Ala Lys	Met Ile Val Glu Ala Lys
340	345	350
Pro Gly His Ala Leu	Ile Gly Phe Glu Ile	Ser Asn Asp Ser Ile Thr
355	360	365
Val Leu Lys Val Tyr	Glu Ala Lys Leu Lys	Gln Asn Tyr Gln Val Asp
370	375	380
Lys Asp Ser Leu Ser	Glu Val Ile Tyr Gly	Asp Met Asp Lys Leu Leu
385	390	395 400
Cys Pro Asp Gln Ser	Glu Gln Ile Tyr Tyr	Thr Asn Asn Ile Val Phe
405	410	415
Pro Asn Glu Tyr Val	Ile Thr Lys Ile Asp	Phe Thr Lys Lys Met Lys
420	425	430
Thr Leu Arg Tyr Glu	Val Thr Ala Asn Phe	Tyr Asp Ser Ser Thr Gly
435	440	445
Glu Ile Asp Leu Asn	Lys Lys Lys Val Glu	Ser Ser Glu Ala Glu Tyr
450	455	460
Arg Thr Leu Ser Ala	Asn Asp Asp Gly Val	Tyr Met Pro Leu Gly Val
465	470	475 480
Ile Ser Glu Thr Phe	Leu Thr Pro Ile Asn	Gly Phe Gly Leu Gln Ala
485	490	495
Asp Glu Asn Ser Arg	Leu Ile Thr Leu Thr	Cys Lys Ser Tyr Leu Arg
500	505	510
Glu Leu Leu Leu Ala	Thr Asp Leu Ser Asn	Lys Glu Thr Lys Leu Ile
515	520	525
Val Pro Pro Ser Gly	Phe Ile Ser Asn Ile	Val Glu Asn Gly Ser Ile
530	535	540

-continued

Glu	Glu	Asp	Asn	Leu	Glu	Pro	Trp	Lys	Ala	Asn	Asn	Lys	Asn	Ala	Tyr
545					550					555					560
Val	Asp	His	Thr	Gly	Gly	Val	Asn	Gly	Thr	Lys	Ala	Leu	Tyr	Val	His
565					570					575					
Lys	Asp	Gly	Gly	Ile	Ser	Gln	Phe	Ile	Gly	Asp	Lys	Leu	Lys	Pro	Lys
580					585					590					
Thr	Glu	Tyr	Val	Ile	Gln	Tyr	Thr	Val	Lys	Gly	Lys	Pro	Ser	Ile	His
595					600					605					
Leu	Lys	Asp	Glu	Asn	Thr	Gly	Tyr	Ile	His	Tyr	Glu	Asp	Thr	Asn	Asn
610					615					620					
Asn	Leu	Glu	Asp	Tyr	Gln	Thr	Ile	Asn	Lys	Arg	Phe	Thr	Thr	Gly	Thr
625					630					635					640
Asp	Leu	Lys	Gly	Val	Tyr	Leu	Ile	Leu	Lys	Ser	Gln	Asn	Gly	Asp	Glu
645					650					655					
Ala	Trp	Gly	Asp	Asn	Phe	Ile	Ile	Leu	Glu	Ile	Ser	Pro	Ser	Glu	Lys
660					665					670					
Leu	Leu	Ser	Pro	Glu	Leu	Ile	Asn	Thr	Asn	Asn	Trp	Thr	Ser	Thr	Gly
675					680					685					
Ser	Thr	Asn	Ile	Ser	Gly	Asn	Thr	Leu	Thr	Leu	Tyr	Gln	Gly	Gly	Arg
690					695					700					
Gly	Ile	Leu	Lys	Gln	Asn	Leu	Gln	Leu	Asp	Ser	Phe	Ser	Thr	Tyr	Arg
705					710					715					720
Val	Tyr	Phe	Ser	Val	Ser	Gly	Asp	Ala	Asn	Val	Arg	Ile	Arg	Asn	Ser
725					730					735					
Arg	Glu	Val	Leu	Phe	Glu	Lys	Arg	Tyr	Met	Ser	Gly	Ala	Lys	Asp	Val
740					745					750					
Ser	Glu	Met	Phe	Thr	Thr	Lys	Phe	Glu	Lys	Asp	Asn	Phe	Tyr	Ile	Glu
755					760					765					
Leu	Ser	Gln	Gly	Asn	Asn	Leu	Tyr	Gly	Gly	Pro	Ile	Val	His	Phe	Tyr
770					775					780					
Asp	Val	Ser	Ile	Lys											
785															

<210> SEQ ID NO 10
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: COT202 nucleotide motif

<400> SEQUENCE: 10

ggtcacctgga tacggtgtca

20

<210> SEQ ID NO 11
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: COT202 nucleotide motif

<400> SEQUENCE: 11

ttgagggttg gatcctttgc

20

<210> SEQ ID NO 12
 <211> LENGTH: 26
 <212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: COT202 nucleotide motif
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: VIC label at 5' end
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (26)..(26)
<223> OTHER INFORMATION: TAMRA label at 3' end

<400> SEQUENCE: 12

caccaacatc atcaatggtg gcatcg 26

<210> SEQ ID NO 13
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: COT202 nucleotide motif

<400> SEQUENCE: 13

ggaatgtggc gaatggtgat 20

<210> SEQ ID NO 14
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: COT202 nucleotide motif

<400> SEQUENCE: 14

tgtcgtttcc cgcttca 18

<210> SEQ ID NO 15
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: COT202 nucleotide motif
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: FAM label at 5' end
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (29)..(29)
<223> OTHER INFORMATION: TAMRA label at 3' end

<400> SEQUENCE: 15

caaattgccc atttcattca tccaaaagc 29

<210> SEQ ID NO 16
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: COT202 nucleotide motif

<400> SEQUENCE: 16

gggtgccatc gggtagtcca taa 23

<210> SEQ ID NO 17
<211> LENGTH: 24
<212> TYPE: DNA

-continued

```

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: COT202 nucleotide motif

<400> SEQUENCE: 17

```

```

tgagtaggag atgtaagttg gcgc

```

24

```

<210> SEQ ID NO 18
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: COT202 nucleotide motif

<400> SEQUENCE: 18

```

```

tctatgttac tagatcgga attg

```

24

What is claimed is:

1. A method of detecting plant material derived from the COT202 event comprising:

- (a) obtaining a sample for analysis;
- (b) providing DNA from the sample;
- (c) providing a first primer and a second designed to bind to a polynucleotide comprising the sequence of SEQ ID NO: 1 or SEQ ID NO: 2 when said polynucleotide is single stranded;
- (d) amplifying the region which lies between the sites at which the primers bind; and
- (e) detecting the presence of the amplification product;

whereby the presence of the amplification product is indicative that the sample is derived from the COT202 event.

2. The method according to claim 1 wherein the first primer has the sequence of SEQ ID NO: 3 and the second primer has the sequence of SEQ ID NO: 4.

3. A method of detecting plant material derived from the COT202 event comprising:

- (a) obtaining a sample for analysis;
- (b) providing a probe designed to bind to the complement of a polynucleotide comprising the sequence of SEQ ID

NO: 1 or SEQ ID NO: 2 when said polynucleotide is single stranded;

- (c) hybridising said probe with the sample; and
- (d) detecting whether the probe has hybridised;

whereby the hybridisation of the probe is indicative that the sample is derived from the COT202 event.

4. The method according to claim 3 wherein the sequence of the probe is selected from the group comprising SEQ ID NO: 7 and SEQ ID NO: 8.

5. The method according to claim 3 wherein the probe hybridises to the sample under stringent hybridisation conditions.

6. A kit of parts comprising a means for detecting the presence in a sample of plant material derived from event COT202.

7. The kit of parts according to claim 6 comprising a means for detecting the presence in a sample of a polynucleotide comprising the sequence of SEQ ID NO: 1 or SEQ ID NO: 2 or a protein encoded by said polynucleotide.

* * * * *