



(22) Date de dépôt/Filing Date: 1998/10/14

(41) Mise à la disp. pub./Open to Public Insp.: 1999/04/21

(45) Date de délivrance/Issue Date: 2008/04/29

(30) Priorités/Priorities: 1997/10/21 (JP288526/1997);
1997/10/21 (JP288629/1997);
1997/11/14 (JP313077/1997);
1997/11/14 (JP313763/1997);
1997/11/21 (JP320805/1997);
1997/12/11 (JP341154/1997); 1998/01/16 (JP6410/1998);
1998/02/05 (JP24614/1998); 1998/02/05 (JP24613/1998)

(51) Cl.Int./Int.Cl. *A61K 49/00* (2006.01),
A61K 49/18 (2006.01), *A61K 51/12* (2006.01),
G01N 33/58 (2006.01)


(72) Inventeurs/Inventors:
KOHNO, TADASHI, JP;
HOSOI, ISABURO, JP;
OHSHIMA, JUNKO, JP;
ITO, ASUKA, JP;
SHIBATA, KUNIHIKO, JP

(73) Propriétaire/Owner:
TOKYO GAS CO., LTD., JP

(74) Agent: HILL & SCHUMACHER

(54) Titre : AGENT DE DIAGNOSTIC DU DIABETE

(54) Title: DIAGNOSTIC AGENT FOR DIABETES



Increase of ¹³C₂ in Exhalation after
Administration of 1-¹³C-Galactose

1-¹³C-galactose (100 mg/kg) was intravenously administered to normal rats (male SD; 9-week-old; fasting blood sugar level 74.8±5.1 mg/dl; n=4) and diabetic rats (male SD; 9-week-old; casual blood sugar level 423.3±87.5 mg/dl; fasting blood sugar level 94±18.5 mg/dl; n=4) at time 0. Then, degrees of increase of ¹³C levels in exhaled CO₂ (Δ¹³C (‰)) were measured. All the rats were fasted overnight before the experiment.

(57) Abrégé/Abstract:

The present invention relates to a diagnostic agent for diabetes, comprising a compound labelled with ¹³C at least at one specific position selected from the group consisting of the following (a) to (g): (a) galactose, fructose or xylose labelled with ¹³C at least at

(57) Abrégé(suite)/Abstract(continued):

one specific position, or a starch composed of glucose units labelled with ^{13}C at least at one specific position; (b) an amino acid labelled with ^{13}C at least at one specific position; (c) lactic acid or citric acid labelled with ^{13}C at least at one specific position; (d) a fatty acid labelled with ^{13}C at least at one specific position; (e) a glyceride labelled with ^{13}C at least at one specific position; (f) glycerol labelled with ^{13}C at least at one specific position; and (g) aminopyrin of which the dimethylamino group at position 4 is labelled with ^{13}C , phenacetin of which the ethoxy group is labelled with ^{13}C at position 1, or methacetin of which the methoxy group is labelled with ^{13}C . According to the present invention, there is provided a diagnostic agent for diabetes which does not impose a heavy physical burden on a subject, can give accurate test results immediately and can be used safely without side effects. The diagnostic agent of the invention makes it possible to discriminate patients with diabetes from normal subjects even under circumstances where patients are easily missed.

ABSTRACT

The present invention relates to a diagnostic agent for diabetes, comprising a compound labelled with ^{13}C at least at one specific position selected from the group consisting of the following (a) to (g):

- (a) galactose, fructose or xylose labelled with ^{13}C at least at one specific position, or a starch composed of glucose units labelled with ^{13}C at least at one specific position;
 - (b) an amino acid labelled with ^{13}C at least at one specific position;
 - (c) lactic acid or citric acid labelled with ^{13}C at least at one specific position;
 - (d) a fatty acid labelled with ^{13}C at least at one specific position;
 - (e) a glyceride labelled with ^{13}C at least at one specific position;
 - (f) glycerol labelled with ^{13}C at least at one specific position;
- and
- (g) aminopyrin of which the dimethylamino group at position 4 is labelled with ^{13}C , phenacetin of which the ethoxy group is labelled with ^{13}C at position 1, or methacetin of which the methoxy group is labelled with ^{13}C .

According to the present invention, there is provided a diagnostic agent for diabetes which does not impose a heavy physical burden on a subject, can give accurate test results immediately and can be used safely without side effects. The diagnostic agent of the invention makes it possible to discriminate patients with diabetes from normal subjects even under circumstances where patients are easily missed.

DIAGNOSTIC AGENT FOR DIABETES

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to a diagnostic agent for diabetes.

More specifically, the present invention relates to a diagnostic agent for diabetes which comprises a compound labelled with ^{13}C at least at one specific position.

2. Description of the Prior Art

Test methods generally used in the primary screening in diagnosis of diabetes are urine sugar test and fasting blood sugar level test. These tests are simple and high in specificity, but are low in sensitivity and give negative results for patients with light diabetes. Thus, 70% or more patients are missed and these tests are considered inadequate as screening tests for diabetes (Sekikawa et al., Medical Practice 10:63, 1933). On the other hand, the glucose tolerance test used for diagnosis of diabetes causes side effects due to administration of a large amount of glucose, and this test requires restraint of a subject for several hours and repeated collection of blood, imposing a heavy physical burden on the subject. Further, the procedures of this test are troublesome. Therefore, this test is actually impossible to carry out as a screening test for diabetes. Recently, blood HbA1C and fructosamine tests, which reflect the average blood sugar level of a subject for a certain period in the past, have been introduced as screening tests for diabetes in some facilities. Under the existing circumstances, however, even these tests cannot be said to be adequate in

sensitivity and specificity for light diabetes, and there remains the problem of difference in measurement results among facilities.

Blood sugar level, HbA1C and fructosamine tests have been used widely for management of outpatients with diabetes and evaluation of therapeutic effects. However, since blood sugar levels will drop at the time of fasting in the case of light diabetes, they cannot be a criterion for the evaluation. HbA1C and fructosamine tests have, in addition to the above-described problems, the following problem: the results of these tests cannot be known until the next visit to the hospital, so instructions are given to the patient on the basis of the past test results.

Under such circumstances, there is a demand for development of a test method for diagnosis of diabetes, management of patients with diabetes and evaluation of therapeutic effects, which method is effective even for patients with light diabetes, does not impose a heavy burden on subjects, and gives results immediately and accurately.

OBJECTS AND SUMMARY OF THE INVENTION

It is an object of the present invention to provide a diagnostic agent for diabetes which is effective even for patients with light diabetes, does not impose a heavy burden on subjects, and can give accurate test results immediately.

As a result of their eager researches toward the solution of the above problems, the present inventors have found that it is possible to diagnose diabetes accurately by administering to a subject a compound labelled with ^{13}C at least at one specific position and measuring the degree of increase of ^{13}C levels in the exhaled CO_2 . Thus, the present invention has been achieved.

The present invention relates to a diagnostic agent for diabetes, comprising a compound labelled with ^{13}C at least at one specific position selected from the group consisting of the following (a) to (g):

- (a) galactose, fructose or xylose labelled with ^{13}C at least at one specific position, or a starch composed of glucose units labelled with ^{13}C at least at one specific position;
 - (b) an amino acid labelled with ^{13}C at least at one specific position;
 - (c) lactic acid or citric acid labelled with ^{13}C at least at one specific position;
 - (d) a fatty acid labelled with ^{13}C at least at one specific position;
 - (e) a glyceride labelled with ^{13}C at least at one specific position;
 - (f) glycerol labelled with ^{13}C at least at one specific position;
- and
- (g) aminopyrin of which the dimethylamino group at position 4 is labelled with ^{13}C , phenacetin of which the ethoxy group is labelled with ^{13}C at position 1, or methacetin of which the methoxy group is labelled with ^{13}C .

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows a method of sampling an exhalation from a rat.

Fig. 2 shows the increase of $^{13}\text{CO}_2$ in exhalation after administration of 1- ^{13}C -galactose.

Fig. 3 shows the increase of $^{13}\text{CO}_2$ in exhalation after administration of 2- ^{13}C -galactose.

Fig. 4 shows the increase of $^{13}\text{CO}_2$ in exhalation after administration of 1- ^{13}C -fructose.

Fig. 5 shows the increase of $^{13}\text{CO}_2$ in exhalation after administration of 2- ^{13}C -fructose.

Fig. 6 shows the increase of $^{13}\text{CO}_2$ in exhalation after administration of U- ^{13}C -starch.

Fig. 7 shows the increase of $^{13}\text{CO}_2$ in exhalation after administration of 1- ^{13}C -isoleucine.

Fig. 8 shows the increase of $^{13}\text{CO}_2$ in exhalation after administration of 1- ^{13}C -alanine.

Fig. 9 shows the increase of $^{13}\text{CO}_2$ in exhalation after administration of 1- ^{13}C -histidine.

Fig. 10 shows the increase of $^{13}\text{CO}_2$ in exhalation after administration of 1- ^{13}C -valine.

Fig. 11 shows the increase of $^{13}\text{CO}_2$ in exhalation after administration of 1,2- ^{13}C -ornithine hydrochloride.

Fig. 12 shows the increase of $^{13}\text{CO}_2$ in exhalation after administration of 1- ^{13}C -methionine.

Fig. 13 shows the increase of $^{13}\text{CO}_2$ in exhalation after administration of 1- ^{13}C -arginine.

Fig. 14 shows the increase of $^{13}\text{CO}_2$ in exhalation after administration of 1- ^{13}C -tryptophan.

Fig. 15 shows the increase of $^{13}\text{CO}_2$ in exhalation after administration of 1- ^{13}C -serine.

Fig. 16 shows the increase of $^{13}\text{CO}_2$ in exhalation after administration of 1- ^{13}C -tyrosine.

Fig. 17 shows the increase of $^{13}\text{CO}_2$ in exhalation after administration of 1- ^{13}C -glutamine.

Fig. 18 shows the increase of $^{13}\text{CO}_2$ in exhalation after administration of 1- ^{13}C -lysine hydrochloride.

Fig. 19 shows the increase of $^{13}\text{CO}_2$ in exhalation after

administration of 1-¹³C-glutamic acid.

Fig. 20 shows the increase of ¹³CO₂ in exhalation after administration of 1-¹³C-proline.

Fig. 21 shows the increase of ¹³CO₂ in exhalation after administration of 1-¹³C-glycine.

Fig. 22 shows the increase of ¹³CO₂ in exhalation after administration of 1-¹³C-aspartic acid.

Fig. 23 shows the increase of ¹³CO₂ in exhalation after administration of 1,1-¹³C-cystine.

Fig. 24 shows the increase of ¹³CO₂ in exhalation after administration of 3-¹³C-lactic acid.

Fig. 25 shows the increase of ¹³CO₂ in exhalation after administration of 1-¹³C-acetic acid.

Fig. 26 shows the increase of ¹³CO₂ in exhalation after administration of 1-¹³C-oleic acid.

Fig. 27 shows the increase of ¹³CO₂ in exhalation after administration of 1-¹³C-palmitic acid.

Fig. 28 shows the increase of ¹³CO₂ in exhalation after administration of 1-¹³C-octanoic acid.

Fig. 29 shows the increase of ¹³CO₂ in exhalation after administration of 1,1,1-¹³C-trioctanoin.

Fig. 30 shows the increase of ¹³CO₂ in exhalation after administration of 1,1,1-¹³C-triacetin.

Fig. 31 shows the increase of ¹³CO₂ in exhalation after administration of 2-¹³C-glycerol.

Fig. 32 shows the increase of ¹³CO₂ in exhalation after administration of N,N-dimethyl-¹³C-aminopyrin.

Fig. 33 shows the increase of ¹³CO₂ in exhalation after administration of ethoxy-1-¹³C-phenacetin.

DETAILED DESCRIPTION OF THE INVENTION

Hereinbelow, the present invention will be described in detail.

In the diagnostic agent of the present invention, galactose, fructose or xylose labelled with ^{13}C at least at one specific position or a starch composed of glucose units labelled with ^{13}C at least at one specific position may be used. The position of the labelling is not particularly limited.

Alternatively, an aliphatic amino acid, aromatic amino acid, sulfur-containing amino acid, heterocyclic amino acid, acidic amino acid or basic amino acid labelled with ^{13}C at least at one specific position may be used in the diagnostic agent of the present invention.

The position of the labelling is not particularly limited.

Preferable examples of the aliphatic amino acid include, but are not limited to, glycine, alanine, serine, threonine, valine, leucine and isoleucine. Preferable examples of the aromatic amino acid include, but are not limited to, phenylalanine and tyrosine. Preferable examples of the sulfur-containing amino acid include, but are not limited to, cysteine, cystine and methionine. Preferable examples of the heterocyclic amino acid include, but are not limited to, tryptophan, proline and histidine. Preferable examples of the acidic amino acid include, but are not limited to, aspartic acid, asparagine, glutamic acid and glutamine. Preferable examples of the basic amino acid include, but are not limited to, arginine, lysine and ornithine.

Alternatively, lactic acid or citric acid labelled with ^{13}C at least at one specific position may be used in the diagnostic agent of the present invention. The position of the labelling is not

particularly limited.

Alternatively, a fatty acid labelled with ^{13}C at least at one specific position may be used in the diagnostic agent of the present invention. The position of the labelling is not particularly limited.




Preferable examples of the fatty acid include, but are not limited to, acetic acid, octanoic acid, palmitic acid, oleic acid, linolic acid and linolenic acid.

Alternatively, a glyceride labelled with ^{13}C at least at one specific position may be used in the diagnostic agent of the present invention. The position of the labelling is not particularly limited.

Preferable examples of the glyceride include, but are not limited to, trioctanoin, tripalmitin, triolein and triacetin.

Alternatively, glycerol labelled with ^{13}C at least at one specific position may be used in the diagnostic agent of the present invention. The position of the labelling is not particularly limited.

Alternatively, aminopyrin of which the dimethylamino group at position 4 is labelled with ^{13}C (N,N-dimethyl- ^{13}C -aminopyrin), phenacetin of which the ethoxy group is labelled with ^{13}C at position 1 (ethoxy-1- ^{13}C -phenacetin), or methacetin of which the methoxy group is labelled with ^{13}C (methoxy- ^{13}C -methacetin) may be used in the diagnostic agent of the present invention. The structures of N,N-dimethyl- ^{13}C -aminopyrin, ethoxy-1- ^{13}C -phenacetin and methoxy- ^{13}C -methacetin are shown in the following formulas (I) to (III), respectively.



Any of the above-mentioned compounds used in the present invention is contained in foods, or the safety thereof as a medicine has been established. Further, unlike radioisotopes, ^{13}C is a stable isotope, and thus there is no danger of exposure to radiation. Accordingly, the diagnostic agent of the invention has no problem in its safety.

The test using the diagnostic agent of the invention is a breath test in which the agent is administered to a subject once or continuously and then the increase of ^{13}C levels in the exhaled CO_2 is measured. Specifically, ^{13}C levels in the exhaled CO_2 after administration of the agent are measured, followed by evaluation of

diabetes from data on the degree of increase of ^{13}C levels in the exhaled CO_2 ($\Delta^{13}\text{C} (\text{‰})$) at predetermined intervals (e.g., 5 min, 10 min, 15 min) after the administration, total amount of $^{13}\text{CO}_2$ exhalation for a predetermined time after administration of the reagent, or data on the time course (slope at the start, change in the slope, peak time, etc.) of the degree of increase of ^{13}C levels in the exhaled CO_2 ($\Delta^{13}\text{C} (\text{‰})$) for a predetermined time period after the administration. The results of this breath test are useful by themselves, but it is more preferable to use these results in combination with blood sugar levels, HbA1C values and fructosamine values for diagnosis.

^{13}C levels in exhaled CO_2 can be determined using gas chromatography mass spectrometry (GC-MS), infrared spectrophotometry, mass spectrometry, photoelectric acoustic spectrophotometry and NMR (nuclear magnetic resonance).

The diagnostic agent of the invention for diabetes can be formulated into pharmaceutical preparations such as parenteral agents (tablets, capsules, powder, granules, liquid, etc.), injections and the like, depending on the administration route, by using the above-described compound labelled with ^{13}C at least at one specific position alone or mixing it with fillers or carriers. The fillers or carriers may be any of those conventionally used in this field as long as they are pharmaceutically acceptable. The type and composition of such fillers or carriers are altered appropriately according to the route and method of administration. For example, water is used as a liquid carrier. As solid carriers, cellulose derivatives such as hydroxypropyl cellulose and organic acid salts such as magnesium stearate are used. Water, physiological saline and various buffer solutions are generally desirable in the preparation of

injections. Such preparations may be lyophilized for use as oral agents, or the lyophilized preparations may be dissolved in suitable injection solvents e.g. liquids for intravenous administration (such as sterilized water, physiological saline, electrolyte, etc.) just before use.

The content of the labelled compound in the pharmaceutical preparation varies depending to the type of the preparation, and is usually in the range of 1 to 100% by weight, preferably 50 to 100% by weight. In the case of injections, for example, the labelled compound is added usually in an amount of 1 to 40% by weight. In the case of capsules, tablets, granules and powder, the content of the labelled compound is in the range from about 10 to 100% by weight, preferably 50 to 100% by weight, with the remainder being carriers.

The diagnostic agent of the invention for diabetes should be administered at such a dosage that enables the confirmation of an increase of $^{13}\text{CO}_2$ in an exhalation after administration of the diagnostic agent. Depending on the age and weight of the patient and the object of the test, the dosage for each administration ranges from 1 to 1000 mg/kg body weight in the case of an adult.

EFFECT OF THE INVENTION

According to the present invention, there is provided a diagnostic agent for diabetes which does not impose a heavy physical burden on a subject, can give accurate test results immediately and can be used safely without side effects. With the diagnostic agent of the invention, patients with light diabetes who exhibit normal blood sugar levels at the time of fasting can be screened. Thus, the diagnostic agent of the invention makes it possible to discriminate

patients with diabetes from normal subjects even under circumstances where patients are easily missed. Further, the diagnostic agent of the invention is useful for management of outpatient with diabetes and evaluation of therapeutic effects.

PREFERRED EMBODIMENTS OF THE INVENTION

Hereinbelow, the present invention will be described more specifically below with reference to the following Examples. However, the scope of the present invention is not limited to these Examples.

The ^{13}C purity at the position of labelling in each of the compounds used in the present invention is 99% or more. Unless otherwise indicated, all the reagents used were guaranteed reagents.

EXAMPLE 1

Method of Breath Test

(1) Preparation of rats with light diabetes

As test animals, male Sprague-Dawley strain (SD) neonatal rats were purchased from Nippon Charles River K.K. together with their lactating rats. They were bred at 23 ± 2 °C under $55 \pm 10\%$ humidity before use.

Insulin secretion-deficient type diabetes was generated by administering streptozotocin (STZ) to neonatal rats ("Saibokogaku (Cell Engineering)", Extra Issue, medical Experiment Manual Series, Strategy for Study of Diabetes, edited by Susumu Seino and Yoshikazu Oka, published by Shujunsha Co., Japan). STZ was subcutaneously administered to 2-day-old rats at a dose of 90 mg/kg. The STZ had been dissolved in a citrate buffer (pH 4.5) and was administered

within 5 min after the dissolution. Two days after the administration, blood was collected by cardiac puncture, and the casual blood sugar level of each rat was measured with Terumo™ Mediacce (blood sugar measurement set). Those rats showing a casual blood sugar level of 275 mg/dl or above were selected. In these rats, their casual blood sugar levels begin to rise at 5 weeks after the administration STZ. At 7 weeks after the administration, almost all rats show a high casual blood level. However, their fasting blood sugar levels increase only slightly and remain at almost normal levels.

According to the standards stipulated by the Japan Diabetes Society (1982), those individuals with a fasting blood sugar level (in venous whole blood) of 120 mg/kg or more are regarded as having diabetes ("Tonyobyō Kensa Manyuaru (Diabetes Test Manual)", edited by Yukio Shigeta, published by Nanko-Do Co., Japan). Then, those rats with a fasting blood sugar level of less than 120 mg/dl (and with a casual blood sugar level of 250 mg/dl or more) which are not regarded as having diabetes in a fasting blood sugar level test were selected from the above-described rats with insulin secretion-deficient type diabetes and used as a model of light diabetes.

(2) ^{13}C breath test

Breath tests were carried out according to the method as described in (2)-1 or (2)-2 below on diabetic rats prepared in (1) above (8-10-week-old) and normal rats (8-10-week-old).

(2)-1 Intravenous administration

Rats fasted overnight were anesthetized by intraperitoneal administration of Nembutal (50 mg/kg) and fixed on an operation table (Fig. 1). A blood sample was collected from the tail vein, and its sugar level was determined using Terumo Mediacce (blood sugar measurement set). The head was covered with a cap for sucking the

exhalation. A specific amount of the labelled compound was administered from the femoral vein. The exhalation was sucked with a stroke pump (variable stroke pump VS-500; Shibata Scientific Technology) at a rate of 100 ml/min and introduced directly into a flow cell in $^{13}\text{CO}_2$ Analyzer EX-130S (Japan Spectroscopic Co., Ltd.). A Perma Pure Drier (MD-050-12P; Perma Pure Inc.) was located between the cap and the stroke pump to remove moisture in the exhalation (Fig. 1).

The data output from the $^{13}\text{CO}_2$ analyzer were incorporated into a personal computer (Apple Power Macintosh™8500) after AD conversion. Using the data processing software Lab VIEW (National Instruments), data on 10 points at every 100 msec were added up and averaged at intervals of 5 sec and then converted into ^{13}C atom %, $\Delta^{13}\text{C}$ (‰) and CO_2 gas concentration (%), to thereby perform a continuous measurement ^{13}C breath test. The converted data were displayed on the screen in real time and then stored in the hard disk. During the measurement, the rectum temperature in the rat was monitored and maintained at 37 ± 0.5 °C using a body temperature controller for small animals (TR-100; Fine Science Tools Inc.). The CO_2 gas concentration in the sucked exhalation was maintained at $3 \pm 0.5\%$.

$\Delta^{13}\text{C}$ (‰) was calculated from the ^{13}C level in exhaled CO_2 at each time point (^{13}C t min) and the ^{13}C level in standard CO_2 gas (^{13}C std) using the following formula:

$$\Delta^{13}\text{C} (\text{‰}) = [(^{13}\text{C} \text{ t min} - ^{13}\text{C} \text{ 0 min}) / ^{13}\text{C} \text{ std}] \times 1000$$

(2)-2 Oral administration

Rats fasted overnight were fixed individually in a rat holder of a microwave irradiation apparatus without anesthesia. A blood sample

was collected from the tail vein, and its sugar level was determined using Terumo Mediatec (blood sugar measurement set). The exhalation was sucked with a stroke pump (variable stroke pump VS-500; Shibata Scientific Technology) at a rate of 100-300 ml/min and introduced directly into a flow cell in $^{13}\text{CO}_2$ Analyzer EX-130S (Japan Spectroscopic Co., Ltd.). A Perma Pure Drier (MD-050-12P; Perma Pure Inc.) was located between the rat holder and the stroke pump to remove moisture in the exhalation (Fig. 1). When the CO_2 gas concentration was stabilized, the rat was once released from the rat holder, and then a specific amount of the labelled compound was administered into its stomach using a syringe for oral administration.

The data output from the $^{13}\text{CO}_2$ analyzer were incorporated into a personal computer (Apple Power Macintosh™ 8500) after AD conversion. Using the data processing software Lab VIEW (National Instruments), data on 10 points at every 100 msec were added up and averaged at intervals of 5 sec and then converted into ^{13}C atom %, $\Delta^{13}\text{C}$ (‰) and CO_2 gas concentration (%), to thereby perform a continuous measurement ^{13}C breath test. The converted data were displayed on the screen in real time and then stored in the hard disk. The CO_2 gas concentration in the sucked exhalation was maintained at $3 \pm 0.5\%$.

$\Delta^{13}\text{C}$ (‰) was calculated by the formula described above.

[EXAMPLE 2] $1\text{-}^{13}\text{C}$ -Galactose Breath Test

$1\text{-}^{13}\text{C}$ -galactose (purchased from ICON) dissolved in physiological saline was administered to normal rats (9-week-old; fasting blood sugar level 74.8 ± 5.1 mg/dl; $n=4$) and diabetic rats (9-week-old; casual blood sugar level 423.3 ± 87.5 mg/dl; fasting blood sugar level 94 ± 18.5 mg/dl; $n=4$) from the femoral vein at a dose of 100 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$

(‰)) were measured according to the method described in Example 1.

$\Delta^{13}\text{C}$ values (‰) continued increasing up to 30 min after the administration of $1\text{-}^{13}\text{C}$ -galactose in both the normal and the diabetic rats (Fig. 2).

The $\Delta^{13}\text{C}$ value (‰) at 20 min after the administration was $64.90 \pm 5.75\%$ in the diabetic rats, while the corresponding value was $37.31 \pm 3.37\%$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 5 to 15 min after the administration was $35.53 \pm 4.34\%$ /10 min in the diabetic rats, while the corresponding slope was $20.59 \pm 2.26\%$ /10 min in the normal rats. Thus, the slope in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of $1\text{-}^{13}\text{C}$ -galactose or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 3] $2\text{-}^{13}\text{C}$ -Galactose Breath Test

$2\text{-}^{13}\text{C}$ -galactose (purchased from ICON) dissolved in physiological saline was administered to normal rats (9-week-old; fasting blood sugar level 63.8 ± 13.4 mg/dl; $n=4$) and diabetic rats (9-week-old; casual blood sugar level 371.5 ± 63.4 mg/dl; fasting blood sugar level 86.5 ± 22.6 mg/dl; $n=4$) from the femoral vein at a dose of 100 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured according to the method described in Example 1.

$\Delta^{13}\text{C}$ values (‰) continued increasing up to 20 min after the administration of $2\text{-}^{13}\text{C}$ -galactose in both the normal and the diabetic rats (Fig. 3).

The $\Delta^{13}\text{C}$ value (‰) at 20 min after the administration was $29.68 \pm 8.55\%$ in the diabetic rats, while the corresponding value was $15.00 \pm 1.63\%$ in the normal rats. Thus, the value in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of $2\text{-}^{13}\text{C}$ -galactose. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 4] $1\text{-}^{13}\text{C}$ -Fructose Breath Test

$1\text{-}^{13}\text{C}$ -fructose (purchased from ICON) dissolved in physiological saline was administered to normal rats (8-week-old; fasting blood sugar level 75.3 ± 4.1 mg/dl; $n=4$) and diabetic rats (8-week-old; casual blood sugar level 498.5 ± 75.1 mg/dl; fasting blood sugar level 99.3 ± 11.8 mg/dl; $n=4$) from the femoral vein at a dose of 100 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured according to the method described in Example 1.

$\Delta^{13}\text{C}$ values (‰) continued increasing up to 20 min after the administration of $1\text{-}^{13}\text{C}$ -fructose in both the normal and the diabetic rats (Fig. 4).

The $\Delta^{13}\text{C}$ value (‰) at 15 min after the administration was $83.51 \pm 7.30\%$ in the diabetic rats, while the corresponding value was $63.90 \pm 7.42\%$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 2 to 7 min after the administration was $47.87 \pm 8.34\%$ / 5 min in the diabetic rats, while the corresponding slope was $29.07 \pm 4.08\%$ / 5 min in the normal rats. Thus, the slope in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value ($\%$) at a specific time after administration of $1\text{-}^{13}\text{C}$ -fructose or the slope of increase of $\Delta^{13}\text{C}$ values ($\%$) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 5] $2\text{-}^{13}\text{C}$ -Fructose Breath Test

$2\text{-}^{13}\text{C}$ -fructose (purchased from ICON) dissolved in physiological saline was administered to normal rats (8-week-old; fasting blood sugar level 75 ± 3.1 mg/dl; $n=4$) and diabetic rats (8-week-old; casual blood sugar level 533.5 mg/dl; fasting blood sugar level 86 mg/dl; $n=2$) from the femoral vein at a dose of 100 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ ($\%$)) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values ($\%$) continued increasing up to 20 min after the administration of $2\text{-}^{13}\text{C}$ -fructose. In the diabetic rats, $\Delta^{13}\text{C}$ values ($\%$) increased sharply up to about 8 min after the administration, but thereafter increased gradually up to 20 min (Fig. 5).

The $\Delta^{13}\text{C}$ value ($\%$) at 20 min after the administration was 94.68% in the diabetic rats, while the corresponding value was $58.75 \pm 2.73\%$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 2 to 6 min after the administration was $47.66\text{‰}/4$ min in the diabetic rats, while the corresponding slope was $21.4 \pm 1.84\text{‰}/4$ min in the normal rats. Thus, the slope in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of $2\text{-}^{13}\text{C}$ -fructose or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 6] $1\text{-}^{13}\text{C}$ -Xylose Breath Test

$1\text{-}^{13}\text{C}$ -xylose (purchased from ICON) dissolved in distilled water was administered orally to normal rats (9-week-old; fasting blood sugar level 75.5 ± 2.1 mg/dl; $n=4$) and diabetic rats (9-week-old; casual blood sugar level 370.3 ± 39.0 mg/dl; fasting blood sugar level 77 ± 9.1 mg/dl; $n=4$) at a dose of 300 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured according to the method described in Example 1.

The $\Delta^{13}\text{C}$ value (‰) at 30 min after the administration of $1\text{-}^{13}\text{C}$ -xylose was $23.40 \pm 7.60\text{‰}$ in the diabetic rats, while the corresponding value was $6.82 \pm 3.02\text{‰}$ in the normal rats. Thus, the value in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 20 to 30 min after the administration was $12.80 \pm 2.18\text{‰}/10$ min in the diabetic rats, while the corresponding slope was $4.61 \pm 1.40\text{‰}/10$ min in the normal rats. Thus, the slope in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of $1\text{-}^{13}\text{C}$ -xylose or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 7] $\text{U-}^{13}\text{C}$ -Starch Breath Test

$\text{U-}^{13}\text{C}$ -starch (purchased from Chlorella Industry) dissolved in distilled water was administered orally to normal rats (9-week-old; fasting blood sugar level 57.7 ± 4.5 mg/dl; $n=3$) and diabetic rats (9-week-old; casual blood sugar level 398.3 ± 24.2 mg/dl; fasting blood sugar level 76.5 ± 7.9 mg/dl; $n=4$) at a dose of 30 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured according to the method described in Example 1.

$\Delta^{13}\text{C}$ values (‰) continued increasing up to 20 min after the administration of $\text{U-}^{13}\text{C}$ -starch in both the normal and the diabetic rats (Fig. 6).

The $\Delta^{13}\text{C}$ value (‰) at 30 min after the administration was $144.50 \pm 23.27\%$ in the diabetic rats, while the corresponding value was $213.89 \pm 4.66\%$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) lower than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of $\text{U-}^{13}\text{C}$ -starch. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 8] $1\text{-}^{13}\text{C}$ -Isoleucine Breath Test

$1\text{-}^{13}\text{C}$ -isoleucine (purchased from mass Trace) dissolved in

physiological saline was administered to normal rats (9-week-old; fasting blood sugar level 80.8 ± 5.5 mg/dl; $n=4$) and diabetic rats (9-week-old; casual blood sugar level 368.7 ± 55.7 mg/dl; fasting blood sugar level 81.3 ± 6.2 mg/dl; $n=3$) from the femoral vein at a dose of 20 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C} (\text{‰})$) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values (‰) continued increasing up to 20 min after the administration of $1\text{-}^{13}\text{C}$ -isoleucine. In the diabetic rats, $\Delta^{13}\text{C}$ values (‰) increased sharply up to about 9 min after the administration, but thereafter remained at an almost constant level up to 20 min (Fig. 7).

The $\Delta^{13}\text{C}$ value (‰) at 5 min after the administration was $83.92 \pm 5.53\text{‰}$ in the diabetic rats, while the corresponding value was $46.98 \pm 3.24\text{‰}$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.001$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 2 to 5 min after the administration was $56.21 \pm 2.09\text{‰} / 3 \text{ min}$ in the diabetic rats, while the corresponding slope was $33.21 \pm 2.44\text{‰} / 3 \text{ min}$ in the normal rats. Thus, the slope in the diabetic rats was very significantly ($p < 0.0001$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of $1\text{-}^{13}\text{C}$ -isoleucine or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 9] 1-¹³C-Alanine Breath Test

1-¹³C-alanine (purchased from mass Trace) dissolved in physiological saline was administered to normal rats (8-week-old; fasting blood sugar level 79.5 ± 4.5 mg/dl; n=4) and diabetic rats (8-week-old; casual blood sugar level 445 ± 20.9 mg/dl; fasting blood sugar level 100.8 ± 15.0 mg/dl; n=4) from the femoral vein at a dose of 10 mg/kg. Then, degrees of increase of ¹³C levels in exhaled CO₂ ($\Delta^{13}\text{C} (\text{‰})$) were measured according to the method described in Example 1.

In both the normal and the diabetic rats, $\Delta^{13}\text{C}$ values (‰) sharply increased up to 4 min after the administration of 1-¹³C-alanine, but thereafter decreased gradually up to 20 min (Fig. 8).

The $\Delta^{13}\text{C}$ value (‰) at 10 min after the administration was $161.35 \pm 6.67\%$ in the diabetic rats, while the corresponding value was $132.16 \pm 3.53\%$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 5 to 15 min after the administration was $-52.09 \pm 19.38\%$ /10 min in the diabetic rats, while the corresponding slope was $-18.12 \pm 7.28\%$ /10 min in the normal rats. Thus, the slope in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) smaller than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of 1-¹³C-alanine or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 10] 1-¹³C-Histidine Breath Test

1-¹³C-histidine (purchased from ICON) dissolved in physiological saline was administered to normal rats (9-week-old; fasting blood sugar level 73.8 ± 2.3 mg/dl; n=4) and diabetic rats (9-week-old; casual blood sugar level 417.7 ± 36.6 mg/dl; fasting blood sugar level 106.7 ± 5.4 mg/dl; n=3) from the femoral vein at a dose of 30 mg/kg. Then, degrees of increase of ¹³C levels in exhaled CO₂ ($\Delta^{13}\text{C}$ (‰)) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values (‰) continued increasing up to 20 min after the administration of 1-¹³C-histidine. In the diabetic rats, $\Delta^{13}\text{C}$ values (‰) continued increasing up to about 17 min after the administration, but thereafter decreased gradually up to 20 min (Fig. 9).

The $\Delta^{13}\text{C}$ value (‰) at 15 min after the administration was $133.57 \pm 2.60\%$ in the diabetic rats, while the corresponding value was $81.56 \pm 10.84\%$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 5 to 10 min after the administration was $81.60 \pm 2.25\% / 5$ min in the diabetic rats, while the corresponding slope was $35.91 \pm 6.47\% / 5$ min in the normal rats. Thus, the slope in the diabetic rats was very significantly ($p < 0.001$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of 1-¹³C-histidine or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 11] 1-¹³C-Valine Breath Test

1-¹³C-valine (purchased from mass Trace) dissolved in physiological saline was administered to normal rats (9-week-old; fasting blood sugar level 76.5 ± 9.9 mg/dl; n=4) and diabetic rats (9-week-old; casual blood sugar level 430 ± 30.5 mg/dl; fasting blood sugar level 99.8 ± 11.1 mg/dl; n=4) from the femoral vein at a dose of 20 mg/kg. Then, degrees of increase of ¹³C levels in exhaled CO₂ ($\Delta^{13}\text{C} (\text{‰})$) were measured according to the method described in Example 1.

In both the normal and diabetic rats, $\Delta^{13}\text{C}$ values (‰) sharply increased up to about 6 min after the administration of 1-¹³C-valine, and thereafter increased gradually up to 20 min (Fig. 10).

The $\Delta^{13}\text{C}$ value (‰) at 5 min after the administration was $73.54 \pm 7.30\text{‰}$ in the diabetic rats, while the corresponding value was $45.99 \pm 5.62\text{‰}$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 2 to 4 min after the administration was $38.55 \pm 3.15\text{‰} / 2 \text{ min}$ in the diabetic rats, while the corresponding slope was $24.17 \pm 2.32\text{‰} / 2 \text{ min}$ in the normal rats. Thus, the slope in the diabetic rats was very significantly ($p < 0.001$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of 1-¹³C-valine or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 12] 1,2- ^{13}C -Ornithine Hydrochloride Breath Test

1,2- ^{13}C -ornithine hydrochloride (purchased from ICON) dissolved in physiological saline was administered to normal rats (8-week-old; fasting blood sugar level 67 ± 5.8 mg/dl; $n=4$) and diabetic rats (8-week-old; casual blood sugar level 411.8 ± 102.5 mg/dl; fasting blood sugar level 77 ± 5.7 mg/dl; $n=4$) from the femoral vein at a dose of 20 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values (‰) continued increasing up to 20 min after the administration of 1,2- ^{13}C -ornithine hydrochloride. In the diabetic rats, $\Delta^{13}\text{C}$ values (‰) increased up to about 14 min after the administration, and thereafter remained at an almost constant level up to 20 min (Fig. 11).

The $\Delta^{13}\text{C}$ value (‰) at 10 min after the administration was $155.70 \pm 17.99\%$ in the diabetic rats, while the corresponding value was $100.71 \pm 5.97\%$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 2 to 7 min after the administration was $87.79 \pm 12.93\%$ / 5 min in the diabetic rats, while the corresponding slope was $53.56 \pm 4.72\%$ / 5 min in the normal rats. Thus, the slope in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of 1,2- ^{13}C -ornithine hydrochloride or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level

at the time of fasting.

[EXAMPLE 13] 1-¹³C-Methionine Breath Test

1-¹³C-methionine (purchased from ICON) dissolved in physiological saline was administered to normal rats (10-week-old; fasting blood sugar level 73.3 ± 2.8 mg/dl; n=4) and diabetic rats (10-week-old; casual blood sugar level 590.5 ± 9.9 mg/dl; fasting blood sugar level 84 ± 5.8 mg/dl; n=4) from the femoral vein at a dose of 40 mg/kg. Then, degrees of increase of ¹³C levels in exhaled CO₂ ($\Delta^{13}\text{C} (\text{‰})$) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values (‰) continued increasing up to 20 min after the administration of 1-¹³C-methionine. In the diabetic rats, $\Delta^{13}\text{C}$ values (‰) increased up to about 10 min after the administration, but thereafter decreased gradually up to 20 min (Fig. 12).

The $\Delta^{13}\text{C}$ value (‰) at 10 min after the administration was $216.30 \pm 30.02\text{‰}$ in the diabetic rats, while the corresponding value was $120.31 \pm 18.84\text{‰}$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 2 to 7 min after the administration was $161.58 \pm 24.17\text{‰} / 5 \text{ min}$ in the diabetic rats, while the corresponding slope was $88.17 \pm 17.04 \text{‰} / 5 \text{ min}$ in the normal rats. Thus, the slope in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of 1-¹³C-methionine or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light

diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 14] 1-¹³C-Arginine Breath Test

1-¹³C-arginine (purchased from ICON) dissolved in physiological saline was administered to normal rats (8-week-old; fasting blood sugar level 68.8 ± 6.9 mg/dl; n=4) and diabetic rats (8-week-old; casual blood sugar level 419.5 ± 69.3 mg/dl; fasting blood sugar level 83.3 ± 11.8 mg/dl; n=4) from the femoral vein at a dose of 50 mg/kg. Then, degrees of increase of ¹³C levels in exhaled CO₂ ($\Delta^{13}\text{C} (\text{‰})$) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values (‰) continued increasing up to 20 min after the administration of 1-¹³C-arginine. In the diabetic rats, $\Delta^{13}\text{C}$ values (‰) increased up to about 12 min after the administration, but thereafter decreased gradually up to 20 min (Fig. 13).

The $\Delta^{13}\text{C}$ value (‰) at 10 min after the administration was $192.40 \pm 30.23\text{‰}$ in the diabetic rats, while the corresponding value was $111.68 \pm 14.15\text{‰}$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 2 to 7 min after the administration was $146.90 \pm 18.97\text{‰} / 5 \text{ min}$ in the diabetic rats, while the corresponding slope was $75.02 \pm 10.01\text{‰} / 5 \text{ min}$ in the normal rats. Thus, the slope in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of 1-¹³C-arginine or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration.

With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 15] 1-¹³C-Tryptophan Breath Test

1-¹³C-tryptophan (purchased from ICON) dissolved in physiological saline was administered to normal rats (9-week-old; fasting blood sugar level 61 ± 1.6 mg/dl; n=4) and diabetic rats (9-week-old; casual blood sugar level 487 ± 56.5 mg/dl; fasting blood sugar level 94.7 ± 10.4 mg/dl; n=3) from the femoral vein at a dose of 10 mg/kg. Then, degrees of increase of ¹³C levels in exhaled CO₂ ($\Delta^{13}\text{C} (\text{‰})$) were measured according to the method described in Example 1.

In both the normal and the diabetic rats, $\Delta^{13}\text{C}$ values (‰) continued increasing up to 20 min after the administration of 1-¹³C-tryptophan (Fig. 14).

The $\Delta^{13}\text{C}$ value (‰) at 20 min after the administration was $19.42 \pm 4.37\text{‰}$ in the diabetic rats, while the corresponding value was $8.10 \pm 0.77\text{‰}$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 10 to 15 min after the administration was $4.46 \pm 0.49\text{‰} / 5 \text{ min}$ in the diabetic rats, while the corresponding slope was $1.20 \pm 0.93\text{‰} / 5 \text{ min}$ in the normal rats. Thus, the slope in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of 1-¹³C-tryptophan or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of

fasting.

[EXAMPLE 16] 1-¹³C-Serine Breath Test

1-¹³C-serine (purchased from ICON) dissolved in physiological saline was administered to normal rats (10-week-old; fasting blood sugar level 69.8 ± 3.8 mg/dl; n=4) and diabetic rats (10-week-old; casual blood sugar level 569 ± 50.3 mg/dl; fasting blood sugar level 93.3 ± 7.6 mg/dl; n=4) from the femoral vein at a dose of 50 mg/kg. Then, degrees of increase of ¹³C levels in exhaled CO₂ ($\Delta^{13}\text{C} (\text{‰})$) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values (‰) increased up to about 8 min after the administration of 1-¹³C-serine, and thereafter remained at an almost constant level. In the diabetic rats, $\Delta^{13}\text{C}$ values (‰) increased sharply up to about 8 min after the administration, but thereafter decreased gradually up to 20 min (Fig. 15).

The $\Delta^{13}\text{C}$ value (‰) at 10 min after the administration was $228.08 \pm 43.45\%$ in the diabetic rats, while the corresponding value was $127.79 \pm 10.69\%$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 1 to 4 min after the administration was $158.13 \pm 41.24\%$ / 3 min in the diabetic rats, while the corresponding slope was $84.17 \pm 11.38\%$ / 3 min in the normal rats. Thus, the slope in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of 1-¹³C-serine or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light diabetes which

shows a normal blood sugar level at the time of fasting.

[EXAMPLE 17] 1-¹³C-Tyrosine Breath Test

1-¹³C-tyrosine (purchased from mass Trace) suspended in 0.5% aqueous solution of sodium carboxymethylcellulose was administered orally to normal rats (9-week-old; fasting blood sugar level 79.8 ± 3.9 mg/dl; n=4) and diabetic rats (9-week-old; casual blood sugar level 442 ± 26.8 mg/dl; fasting blood sugar level 86.5 ± 7.4 mg/dl; n=4) at a dose of 60 mg/kg. Then, degrees of increase of ¹³C levels in exhaled CO₂ ($\Delta^{13}\text{C} (\text{‰})$) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values (‰) continued increasing up to 30 min after the administration of 1-¹³C-tyrosine. In the diabetic rats, $\Delta^{13}\text{C}$ values (‰) increased up to about 20 min after the administration, but thereafter decreased gradually up to 30 min (Fig. 16).

The $\Delta^{13}\text{C}$ value (‰) at 20 min after the administration was $205.30 \pm 51.43\%$ in the diabetic rats, while the corresponding value was $73.96 \pm 35.10\%$ in the normal rats. Thus, the value in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 5 to 10 min after the administration was $79.62 \pm 19.90\%$ / 5 min in the diabetic rats, while the corresponding slope was $27.52 \pm 17.10\%$ / 5 min in the normal rats. Thus, the slope in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of 1-¹³C-tyrosine or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration.

With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 18] 1-¹³C-Glutamine Breath Test

1-¹³C-glutamine (purchased from mass Trace) dissolved in physiological saline was administered to normal rats (8-week-old; fasting blood sugar level 72.8 ± 4.6 mg/dl; n=4) and diabetic rats (8-week-old; casual blood sugar level 491 mg/dl; fasting blood sugar level 88.5 mg/dl; n=2) from the femoral vein at a dose of 20 mg/kg. Then, degrees of increase of ¹³C levels in exhaled CO₂ ($\Delta^{13}\text{C} (\text{‰})$) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values (‰) increased sharply up to about 3 min after the administration of 1-¹³C-glutamine, and thereafter increased gradually up to 20 min. In the diabetic rats, $\Delta^{13}\text{C}$ values (‰) increased sharply up to about 3 min after the administration, but thereafter decreased gradually up to 20 min (Fig. 17).

The $\Delta^{13}\text{C}$ value (‰) at 5 min after the administration was 168.18‰ in the diabetic rats, while the corresponding value was $107.17 \pm 15.42\%$ in the normal rats. Thus, the value in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 1 to 2 min after the administration was 105.59‰/min in the diabetic rats, while the corresponding slope was $69.88 \pm 11.51\%$ /min in the normal rats. Thus, the slope in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of 1-¹³C-glutamine

or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 19] 1- ^{13}C -Lysine Hydrochloride Breath Test

1- ^{13}C -lysine hydrochloride (purchased from mass Trace) dissolved in physiological saline was administered orally to normal rats (8-week-old; fasting blood sugar level 71 ± 1.6 mg/dl; $n=3$) and diabetic rats (8-week-old; casual blood sugar level 431 ± 71.7 mg/dl; fasting blood sugar level 90.3 ± 4.9 mg/dl; $n=3$) from the femoral vein at a dose of 50 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values (‰) increased up to about 11 min after the administration of 1- ^{13}C -lysine hydrochloride, and thereafter remained at an almost constant level up to 20 min. In the diabetic rats, $\Delta^{13}\text{C}$ values (‰) increased up to about 10 min after the administration, but thereafter decreased gradually up to 20 min (Fig. 18).

The $\Delta^{13}\text{C}$ value (‰) at 8 min after the administration was $181.53 \pm 23.82\text{‰}$ in the diabetic rats, while the corresponding value was $96.95 \pm 27.48\text{‰}$ in the normal rats. Thus, the value in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 15 to 20 min after the administration was $-26.64 \pm 5.36\text{‰} / 5$ min in the diabetic rats, while the corresponding slope was $-4.58 \pm 6.91\text{‰} / 5$ min in the normal rats. Thus, the slope in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) smaller than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of 1- ^{13}C -lysine hydrochloride or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 20] 1- ^{13}C -Glutamic Acid Breath Test

1- ^{13}C -glutamic acid (purchased from mass Trace) dissolved in physiological saline was administered to normal rats (8-week-old; fasting blood sugar level 70.8 ± 8.3 mg/dl; $n=4$) and diabetic rats (8-week-old; casual blood sugar level 487.7 ± 38.4 mg/dl; fasting blood sugar level 108 ± 8.8 mg/dl; $n=3$) from the femoral vein at a dose of 10 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured according to the method described in Example 1.

In both the normal and the diabetic rats, $\Delta^{13}\text{C}$ values (‰) increased sharply up to about 4 min after the administration of 1- ^{13}C -glutamic acid, but thereafter decreased gradually up to 20 min (Fig. 19).

The $\Delta^{13}\text{C}$ value (‰) at 4 min after the administration was $170.62 \pm 12.18\%$ in the diabetic rats, while the corresponding value was $139.95 \pm 11.79\%$ in the normal rats. Thus, the value in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 10 to 20 min after the administration was $-43.29 \pm 3.47\%$ /10 min in the diabetic rats, while the corresponding slope was $-27.97 \pm 5.05\%$ /10 min in the normal rats. Thus, the slope in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD))

smaller than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of $1\text{-}^{13}\text{C}$ -glutamic acid or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 21] $1\text{-}^{13}\text{C}$ -Proline Breath Test

$1\text{-}^{13}\text{C}$ -proline (purchased from mass Trace) dissolved in physiological saline was administered to normal rats (10-week-old; fasting blood sugar level 77.6 ± 7.3 mg/dl; $n=5$) and diabetic rats (10-week-old; casual blood sugar level 430.3 ± 45.2 mg/dl; fasting blood sugar level 97.3 ± 12.5 mg/dl; $n=4$) from the femoral vein at a dose of 20 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured according to the method described in Example 1.

In both the normal and the diabetic rats, $\Delta^{13}\text{C}$ values (‰) increased up to about 9 min after the administration of $1\text{-}^{13}\text{C}$ -proline, but thereafter decreased gradually up to 20 min (Fig. 20).

The $\Delta^{13}\text{C}$ value (‰) at 20 min after the administration was $115.77 \pm 6.61\text{‰}$ in the diabetic rats, while the corresponding value was $92.27 \pm 11.10\text{‰}$ in the normal rats. Thus, the value in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of $1\text{-}^{13}\text{C}$ -proline. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 22] 1-¹³C-Glycine Breath Test

1-¹³C-glycine (purchased from mass Trace) dissolved in physiological saline was administered to normal rats (9-week-old; fasting blood sugar level 75.5 ± 5.8 mg/dl; n=4) and diabetic rats (9-week-old; casual blood sugar level 410 ± 10.7 mg/dl; fasting blood sugar level 90 ± 11.9 mg/dl; n=3) from the femoral vein at a dose of 20 mg/kg. Then, degrees of increase of ¹³C levels in exhaled CO₂ ($\Delta^{13}\text{C} (\text{‰})$) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values (‰) increased sharply up to about 5 min after the administration of 1-¹³C-glycine, remained at an almost constant level up to about 12 min, but thereafter decreased gradually up to 20 min. In the diabetic rats, $\Delta^{13}\text{C}$ values (‰) increased sharply up to about 5 min after the administration, but thereafter decreased gradually up to 20 min (Fig. 21).

The $\Delta^{13}\text{C}$ value (‰) at 7 min after the administration was $156.62 \pm 10.83\text{‰}$ in the diabetic rats, while the corresponding value was $119.06 \pm 12.19\text{‰}$ in the normal rats. Thus, the value in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of 1-¹³C-glycine. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 23] 1-¹³C-Cysteine Breath Test

1-¹³C-cysteine (purchased from ICON) dissolved in physiological saline was administered to normal rats (9-week-old; fasting blood sugar level 74 ± 7.7 mg/dl; n=4) and diabetic rats (9-week-old;

casual blood sugar level 457 ± 4.3 mg/dl; fasting blood sugar level 93.7 ± 16.6 mg/dl; $n=3$) from the femoral vein at a dose of 20 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C} (\text{‰})$) were measured according to the method described in Example 1.

The $\Delta^{13}\text{C}$ value (‰) at 2 min after the administration of 1- ^{13}C -cysteine was $42.87 \pm 2.09\text{‰}$ in the diabetic rats, while the corresponding value was $78.97 \pm 13.98\text{‰}$ in the normal rats. Thus, the value in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) lower than that in the normal rats.

The slope from 1 to 2 min after the administration was $38.50 \pm 2.33\text{‰}/\text{min}$ in the diabetic rats, while the corresponding slope was $65.72 \pm 10.66 \text{‰}/\text{min}$ in the normal rats. Thus, the slope in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) smaller than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of 1- ^{13}C -cysteine or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 24] 1- ^{13}C -Leucine Breath Test

1- ^{13}C -leucine (purchased from mass Trace) dissolved in physiological saline was administered to normal rats (8-week-old; fasting blood sugar level 82.5 ± 4.5 mg/dl; $n=4$) and diabetic rats (8-week-old; casual blood sugar level 399.5 mg/dl; fasting blood sugar level 85.5 mg/dl; $n=2$) from the femoral vein at a dose of 10 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C} (\text{‰})$) were measured according to the method described in Example 1.

The $\Delta^{13}\text{C}$ value (‰) at 5 min after the administration of 1-

^{13}C -leucine was $52.89 \pm 10.83\%$ in the diabetic rats, while the corresponding value was $42.56 \pm 2.43\%$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) lower than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value ($\%$) at a specific time after administration of $1\text{-}^{13}\text{C}$ -leucine. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 25] $1\text{-}^{13}\text{C}$ -Aspartic Acid Breath Test

$1\text{-}^{13}\text{C}$ -aspartic acid (purchased from mass Trace) dissolved in physiological saline was administered to normal rats (8-week-old; fasting blood sugar level 56.3 ± 6.8 mg/dl; $n=4$) and diabetic rats (8-week-old; casual blood sugar level 389 ± 52.1 mg/dl; fasting blood sugar level 80.5 ± 6.1 mg/dl; $n=4$) from the femoral vein at a dose of 10 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ ($\%$)) were measured according to the method described in Example 1.

In both the normal and the diabetic rats, $\Delta^{13}\text{C}$ values ($\%$) increased sharply up to about 2 min after the administration of $1\text{-}^{13}\text{C}$ -aspartic acid, but thereafter decreased gradually up to 20 min (Fig. 22).

The $\Delta^{13}\text{C}$ value ($\%$) at 3 min after the administration was $172.66 \pm 8.65\%$ in the diabetic rats, while the corresponding value was $126.56 \pm 13.45\%$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 3 to 8 min after the administration was $-53.79 \pm 11.42\%$ / 5 min in the diabetic rats, while the corresponding slope was -

$8.13 \pm 10.84\%$ /5 min in the normal rats. Thus, the slope in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) smaller than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value ($\%$) at a specific time after administration of $1\text{-}^{13}\text{C}$ -aspartic acid or the slope of increase of $\Delta^{13}\text{C}$ values ($\%$) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 26] $1,1\text{-}^{13}\text{C}$ -Cystine Breath Test

$1\text{-}^{13}\text{C}$ -cystine (purchased from mass Trace) suspended in 0.5% aqueous solution of sodium carboxymethylcellulose was administered orally to normal rats (9-week-old; fasting blood sugar level 65.5 ± 6.4 mg/dl; $n=4$) and diabetic rats (9-week-old; casual blood sugar level 324.5 ± 7.7 mg/dl; fasting blood sugar level 81.8 ± 11.5 mg/dl; $n=4$) at a dose of 45 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ ($\%$)) were measured according to the method described in Example 1.

In both the normal and the diabetic rats, $\Delta^{13}\text{C}$ values ($\%$) continued increasing up to 30 min after the administration of $1\text{-}^{13}\text{C}$ -cystine (Fig. 23).

The $\Delta^{13}\text{C}$ value ($\%$) at 10 min after the administration was $69.15 \pm 6.44\%$ in the diabetic rats, while the corresponding value was $45.26 \pm 8.68\%$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 5 to 10 min after the administration was $46.82 \pm 2.94\%$ /5 min in the diabetic rats, while the corresponding slope was

$30.86 \pm 10.06\%$ /5 min in the normal rats. Thus, the slope in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value ($\%$) at a specific time after administration of $1\text{-}^{13}\text{C}$ -cystine or the slope of increase of $\Delta^{13}\text{C}$ values ($\%$) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 27] $3\text{-}^{13}\text{C}$ -Lactic Acid Breath Test

$3\text{-}^{13}\text{C}$ -sodium lactate (purchased from mass Trace) dissolved in physiological saline was administered to normal rats (8-week-old; fasting blood sugar level 67.8 ± 6.5 mg/dl; $n=4$) and diabetic rats (8-week-old; casual blood sugar level 412.8 ± 71.7 mg/dl; fasting blood sugar level 88.5 ± 17.2 mg/dl; $n=4$) from the femoral vein at a dose of 50 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ ($\%$)) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values ($\%$) increased sharply up to about 5 min after the administration of $3\text{-}^{13}\text{C}$ -sodium lactate, and thereafter increased gradually up to 20 min. In the diabetic rats, $\Delta^{13}\text{C}$ values ($\%$) increased sharply up to about 7 min after the administration, but thereafter decreased gradually up to 20 min (Fig. 24).

The $\Delta^{13}\text{C}$ value ($\%$) at 10 min after the administration was $202.08 \pm 6.05\%$ in the diabetic rats, while the corresponding value was $130.30 \pm 11.47\%$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.0001$ (ANOVA with Fischer

LSD)) higher than that in the normal rats.

The slope from 2 to 4 min after the administration was $96.83 \pm 17.69\%$ / 2 min in the diabetic rats, while the corresponding slope was $60.69 \pm 6.80\%$ / 2 min in the normal rats. Thus, the slope in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value ($\%$) at a specific time after administration of $3\text{-}^{13}\text{C}$ -sodium lactate or the slope of increase of $\Delta^{13}\text{C}$ values ($\%$) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 28] $1,6\text{-}^{13}\text{C}$ -Citric Acid Breath Test

$1,6\text{-}^{13}\text{C}$ -citric acid (purchased from ICON) dissolved in physiological saline was administered to normal rats (8-week-old; fasting blood sugar level 66.3 ± 9.6 mg/dl; $n=4$) and diabetic rats (8-week-old; casual blood sugar level 376.5 ± 57.3 mg/dl; fasting blood sugar level 76.5 ± 16.6 mg/dl; $n=4$) from the femoral vein at a dose of 5 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ ($\%$)) were measured according to the method described in Example 1.

The $\Delta^{13}\text{C}$ value ($\%$) at 20 min after the administration of $1,6\text{-}^{13}\text{C}$ -citric acid was $54.38 \pm 3.34\%$ in the diabetic rats, while the corresponding value was $67.49 \pm 1.82\%$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) lower than that in the normal rats.

The slope from 10 to 14 min after the administration was $-10.45 \pm 1.30\%$ / 4 min in the diabetic rats, while the corresponding slope was $-5.81 \pm 0.92\%$ / 4 min in the normal rats. Thus, the slope in the

diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) smaller than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of 1,6- ^{13}C -citric acid or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 29] 1- ^{13}C -Acetic Acid Breath Test

1- ^{13}C -acetic acid (purchased from mass Trace) dissolved in physiological saline was administered to normal rats (8-week-old; fasting blood sugar level 73.8 ± 5.1 mg/dl; $n=4$) and diabetic rats (8-week-old; casual blood sugar level 398.3 ± 57.5 mg/dl; fasting blood sugar level 75.0 ± 8.3 mg/dl; $n=3$) from the femoral vein at a dose of 10 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values (‰) increased sharply up to about 5 min after the administration of 1- ^{13}C -acetic acid, and thereafter remained at an almost constant level up to 20 min. In the diabetic rats, $\Delta^{13}\text{C}$ values (‰) increased sharply up to about 4 min after the administration, but thereafter decreased gradually up to 20 min (Fig. 25).

The $\Delta^{13}\text{C}$ value (‰) at 3 min after the administration was $193.09 \pm 25.69 \text{‰}$ in the diabetic rats, while the corresponding value was $117.50 \pm 8.74 \text{‰}$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 10 to 20 min after the administration was $-31.25 \pm 13.00\%$ /10 min in the diabetic rats, while the corresponding slope was $-4.67 \pm 8.49\%$ /10 min in the normal rats. Thus, the slope in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) smaller than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value ($\%$) at a specific time after administration of $1\text{-}^{13}\text{C}$ -acetic acid or the slope of increase of $\Delta^{13}\text{C}$ values ($\%$) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 30] $1\text{-}^{13}\text{C}$ -Oleic Acid Breath Test

$1\text{-}^{13}\text{C}$ -oleic acid (purchased from ICON) emulsified with Tween 20 (0.1%) and physiological saline was administered to normal rats (9-week-old; fasting blood sugar level 74.8 ± 9.8 mg/dl; $n=4$) and diabetic rats (9-week-old; casual blood sugar level 402.5 ± 31.9 mg/dl; fasting blood sugar level 83.5 ± 4.4 mg/dl; $n=4$) from the femoral vein at a dose of 70 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ ($\%$))) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values ($\%$) continued increasing up to 20 min after the administration of $1\text{-}^{13}\text{C}$ -oleic acid. In the diabetic rats, $\Delta^{13}\text{C}$ values ($\%$) increased sharply up to about 8 min after the administration, but thereafter decreased gradually up to 20 min (Fig. 26).

The $\Delta^{13}\text{C}$ value ($\%$) at 8 min after the administration was $132.93 \pm 7.09\%$ in the diabetic rats, while the corresponding value was $86.51 \pm 14.72\%$ in the normal rats. Thus, the value in the diabetic

rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 10 to 20 min after the administration was $-12.79 \pm 3.72\%$ /10 min in the diabetic rats, while the corresponding slope was $13.61 \pm 4.58\%$ /10 min in the normal rats. Thus, the slope in the diabetic rats was significantly ($p < 0.001$ (ANOVA with Fischer LSD)) smaller than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value ($\%$) at a specific time after administration of $1\text{-}^{13}\text{C}$ -oleic acid or the slope of increase of $\Delta^{13}\text{C}$ values ($\%$) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 31] $1\text{-}^{13}\text{C}$ -Palmitic Acid Breath Test

$1\text{-}^{13}\text{C}$ -palmitic acid (purchased from mass Trace) emulsified with Tween 20 (0.2%) and physiological saline was administered to normal rats (9-week-old; fasting blood sugar level 74.8 ± 7.1 mg/dl; $n=4$) and diabetic rats (9-week-old; casual blood sugar level 422.8 ± 42.0 mg/dl; fasting blood sugar level 80.8 ± 5.1 mg/dl; $n=4$) from the femoral vein at a dose of 50 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ ($\%$)) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values ($\%$) continued increasing up to 20 min after the administration of $1\text{-}^{13}\text{C}$ -palmitic acid. In the diabetic rats, $\Delta^{13}\text{C}$ values ($\%$) increased sharply up to about 5 min after the administration, but thereafter decreased gradually up to 20 min (Fig. 27).

The slope from 10 to 20 min after the administration was $-4.40 \pm$

2.21‰/10 min in the diabetic rats, while the corresponding slope was 1.43 ± 1.33 ‰/10 min in the normal rats. Thus, the slope in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) smaller than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration of $1\text{-}^{13}\text{C}$ -palmitic acid. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 32] $1\text{-}^{13}\text{C}$ -Octanoic Acid Breath Test

$1\text{-}^{13}\text{C}$ -octanoic acid (purchased from mass Trace) emulsified with Tween 20 (0.2%) and physiological saline was administered to normal rats (9-week-old; fasting blood sugar level 78.3 ± 4.7 mg/dl; $n=4$) and diabetic rats (9-week-old; casual blood sugar level 440.3 ± 38.1 mg/dl; fasting blood sugar level 89.8 ± 9.4 mg/dl; $n=4$) from the femoral vein at a dose of 30 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values (‰) increased sharply up to about 5 min after the administration of $1\text{-}^{13}\text{C}$ -octanoic acid, and thereafter remained at an almost constant level up to 20 min. In the diabetic rats, $\Delta^{13}\text{C}$ values (‰) increased sharply up to about 5 min after the administration, but thereafter decreased gradually up to 20 min (Fig. 28).

The slope from 10 to 20 min after the administration was -20.55 ± 7.75 ‰/10 min in the diabetic rats, while the corresponding slope was -4.76 ± 4.41 ‰/10 min in the normal rats. Thus, the slope in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) smaller than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration of $1\text{-}^{13}\text{C}$ -octanoic acid. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 33] $1,1,1\text{-}^{13}\text{C}$ -Trioctanoin Breath Test

$1,1,1\text{-}^{13}\text{C}$ -trioctanoin (purchased from mass Trace) dissolved in olive oil was administered orally to normal rats (8-week-old; fasting blood sugar level 75.5 ± 8.4 mg/dl; $n=4$) and diabetic rats (8-week-old; casual blood sugar level 406.8 ± 61.4 mg/dl; fasting blood sugar level 76.8 ± 6.1 mg/dl; $n=4$) at a dose of 50 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured according to the method described in Example 1.

In both the normal and the diabetic rats, $\Delta^{13}\text{C}$ values (‰) continued increasing up to 30 min after the administration of $1,1,1\text{-}^{13}\text{C}$ -trioctanoin (Fig. 29).

The $\Delta^{13}\text{C}$ value (‰) at 20 min after the administration was $150.72 \pm 22.48\%$ in the diabetic rats, while the corresponding value was $110.16 \pm 16.08\%$ in the normal rats. Thus, the value in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of $1,1,1\text{-}^{13}\text{C}$ -trioctanoin. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 34] $1,1,1\text{-}^{13}\text{C}$ -Triacetin Test

$1,1,1\text{-}^{13}\text{C}$ -triacetin emulsified with Tween™ 20 (0.4%) and

physiological saline was administered to normal rats (9-week-old; fasting blood sugar level 67.7 ± 6.8 mg/dl; $n=3$) and diabetic rats (9-week-old; casual blood sugar level 332 ± 44.2 mg/dl; fasting blood sugar level 80.3 ± 15.0 mg/dl; $n=4$) from the femoral vein at a dose of 10 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C} (\text{‰})$) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values (‰) continued increasing up to 20 min after the administration of 1,1,1- ^{13}C -triacetin. In the diabetic rats, $\Delta^{13}\text{C}$ values (‰) increased sharply up to about 8 min after the administration, but thereafter decreased gradually up to 20 min (Fig. 30).

The $\Delta^{13}\text{C}$ value (‰) at 4 min after the administration was $41.97 \pm 2.79\text{‰}$ in the diabetic rats, while the corresponding value was $27.29 \pm 2.31\text{‰}$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 6 to 10 min after the administration was $-6.26 \pm 1.50\text{‰}/4$ min in the diabetic rats, while the corresponding slope was $-0.83 \pm 0.90\text{‰}/4$ min in the normal rats. Thus, the slope in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) smaller than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of 1,1,1- ^{13}C -triacetin or from the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 35] 2-¹³C-Glycerol Breath Test

2-¹³C-glycerol (purchased from CIL) dissolved in physiological saline was administered to normal rats (9-week-old; fasting blood sugar level 70.3 ± 4.1 mg/dl; n=4) and diabetic rats (9-week-old; casual blood sugar level 440 ± 18.0 mg/dl; fasting blood sugar level 83 ± 9.3 mg/dl; n=3) from the femoral vein at a dose of 50 mg/kg. Then, degrees of increase of ¹³C levels in exhaled CO₂ ($\Delta^{13}\text{C} (\text{‰})$) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values (‰) continued increasing gradually up to 20 min after the administration of 2-¹³C-glycerol, and thereafter remained at an almost constant level. In the diabetic rats, $\Delta^{13}\text{C}$ values (‰) increased sharply up to about 8 min after the administration, and thereafter increased gradually up to 20 min (Fig. 31).

The $\Delta^{13}\text{C}$ value (‰) at 20 min after the administration was $112.44 \pm 5.12\%$ in the diabetic rats, while the corresponding value was $54.77 \pm 4.23\%$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.0001$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 2 to 7 min after the administration was $62.72 \pm 3.22\%$ / 5 min in the diabetic rats, while the corresponding slope was $16.99 \pm 1.61\%$ / 5 min in the normal rats. Thus, the slope in the diabetic rats was very significantly ($p < 0.0001$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of 2-¹³C-glycerol or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 36] N,N-Dimethyl- ^{13}C -Aminopyrin Breath Test

N,N-dimethyl- ^{13}C -aminopyrin (purchased from ICON) dissolved in physiological saline was administered to normal rats (8-week-old; fasting blood sugar level 68.8 ± 4.0 mg/dl; $n=4$) and diabetic rats (8-week-old; casual blood sugar level 414.5 ± 29.9 mg/dl; fasting blood sugar level 94.8 ± 10.0 mg/dl; $n=4$) from the femoral vein at a dose of 40 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C} (\text{‰})$) were measured according to the method described in Example 1.

In the normal rats, $\Delta^{13}\text{C}$ values (‰) increased sharply up to about 3 min after the administration of N,N-dimethyl- ^{13}C -aminopyrin, and thereafter increased gradually up to 20 min. In the diabetic rats, $\Delta^{13}\text{C}$ values (‰) increased sharply up to about 5 min after the administration, and thereafter remained at an almost constant level up to 20 min (Fig. 32).

The $\Delta^{13}\text{C}$ value (‰) at 20 min after the administration was $193.62 \pm 22.21\text{‰}$ in the diabetic rats, while the corresponding value was $92.2 \pm 12.64\text{‰}$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.001$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 1 to 2 min after the administration was $95.68 \pm 16.33 \text{ ‰/min}$ in the diabetic rats, while the corresponding slope was $41.16 \pm 8.89\text{‰/min}$ in the normal rats. Thus, the slope in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of N,N-dimethyl- ^{13}C -aminopyrin or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the

administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 37] Ethoxy-1- ^{13}C -Phenacetin Breath Test

Ethoxy-1- ^{13}C -phenacetin (purchased from ICON) suspended in 0.5% aqueous solution of carboxymethylcellulose was administered orally to normal rats (9-week-old; fasting blood sugar level 72 ± 8.9 mg/dl; $n=4$) and diabetic rats (9-week-old; casual blood sugar level 504.7 ± 55.0 mg/dl; fasting blood sugar level 101.7 ± 3.3 mg/dl; $n=3$) at a dose of 90 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C} (\text{‰})$) were measured according to the method described in Example 1.

In both the normal and the diabetic rats, $\Delta^{13}\text{C}$ values (‰) continued increasing up to 30 min after the administration of ethoxy-1- ^{13}C -phenacetin (Fig. 33).

The $\Delta^{13}\text{C}$ value (‰) at 30 min after the administration was $119.85 \pm 13.87\text{‰}$ in the diabetic rats, while the corresponding value was $70.99 \pm 12.34\text{‰}$ in the normal rats. Thus, the value in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) higher than that in the normal rats.

The slope from 10 to 15 min after the administration was $30.88 \pm 5.62\text{‰} / 5 \text{ min}$ in the diabetic rats, while the corresponding slope was $16.3 \pm 4.47\text{‰} / 5 \text{ min}$ in the normal rats. Thus, the slope in the diabetic rats was significantly ($p < 0.05$ (ANOVA with Fischer LSD)) greater than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the $\Delta^{13}\text{C}$ value (‰) at a specific time after administration of ethoxy-1- ^{13}C -phenacetin or the slope of increase of $\Delta^{13}\text{C}$ values (‰) after the

administration. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[EXAMPLE 38] Ethoxy- ^{13}C -Methacetin Breath Test

Ethoxy- ^{13}C -methacetin (purchased from CIL) suspended in 0.5% aqueous solution of carboxymethylcellulose was administered orally to normal rats (8-week-old; fasting blood sugar level 86 ± 2.9 mg/dl; $n=4$) and diabetic rats (8-week-old; casual blood sugar level 418.5 ± 24.9 mg/dl; fasting blood sugar level 90.3 ± 6.6 mg/dl; $n=4$) at a dose of 90 mg/kg. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C} (\text{‰})$) were measured according to the method described in Example 1.

The slope from 12 to 17 min after the administration of ethoxy- ^{13}C -methacetin was $-8.40 \pm 7.67\text{‰} / 5$ min in the diabetic rats, while the corresponding slope was $16.05 \pm 7.71\text{‰} / 5$ min in the normal rats. Thus, the slope in the diabetic rats was very significantly ($p < 0.01$ (ANOVA with Fischer LSD)) smaller than that in the normal rats.

Accordingly, it is possible to diagnose diabetes from the slope of increase of $\Delta^{13}\text{C}$ values (‰) after administration of ethoxy- ^{13}C -methacetin. With this test, it is possible to diagnose even light diabetes which shows a normal blood sugar level at the time of fasting.

[FORMULATION EXAMPLE 1] Injection

10 parts by weight of 1- ^{13}C -galactose was dissolved in 90 parts by weight of physiological saline and sterilized by filtration with a Millipore™ filter. The filtrate was put into a vial and sealed to give an injection.

[FORMULATION EXAMPLE 2] Internal Liquid Agent

10 parts by weight of 1-¹³C-fructose was dissolved in 90 parts by weight of deionized and distilled water (DDW) and sterilized by filtration with a Millipore™ filter. The filtrate was put into a vial and sealed to give an internal liquid agent.

[FORMULATION EXAMPLE 3] Injection

10 parts by weight of 1-¹³C-isoleucine was dissolved in 90 parts by weight of physiological saline and sterilized by filtration with a Millipore™ filter. The filtrate was put into a vial and sealed to give an injection.

[FORMULATION EXAMPLE 4] Internal Liquid Agent

10 parts by weight of 1-¹³C-alanine was dissolved in 90 parts by weight of DDW and sterilized by filtration with a Millipore™ filter. The filtrate was put into a vial and sealed to give an internal liquid agent.

[FORMULATION EXAMPLE 5] Injection

10 parts by weight of 3-¹³C-sodium lactate was dissolved in 90 parts by weight of physiological saline and sterilized by filtration with a Millipore™ filter. The filtrate was put into a vial and sealed to give an injection.

[FORMULATION EXAMPLE 6] Internal Liquid Agent

10 parts by weight of 1,6-¹³C-citric acid was dissolved in 90 parts by weight of DDW and sterilized by filtration with a Millipore™ filter. The filtrate was put into a vial and sealed to give an internal liquid agent.

[FORMULATION EXAMPLE 7] Injection

10 parts by weight of 1-¹³C-oleic acid, 89 parts by weight of physiological saline and 1 part by weight of Polysorbate 80 (all sterilized in advance) were mixed aseptically and emulsified with a ultrasonic homogenizer. The resultant emulsion was put into a vial and sealed to give an injection.

[FORMULATION EXAMPLE 8] Internal Liquid Agent

10 parts by weight of 1-¹³C-acetic acid was dissolved in 90 parts by weight of DDW and sterilized by filtration with a Millipore™ filter. The filtrate was put into a vial and sealed to give an internal liquid agent.

[FORMULATION EXAMPLE 9] Injection

10 parts by weight of 1,1,1-¹³C-trioctanoin, 89 parts by weight of physiological saline and 1 part by weight of Polysorbate 80 (all sterilized in advance) were mixed aseptically and emulsified with a ultrasonic homogenizer. The resultant emulsion was put into a vial and sealed to give an injection.

[FORMULATION EXAMPLE 10] Internal Liquid Agent

10 parts by weight of 1,1,1-¹³C-trioctanoin, 89 parts by weight of DDW and 1 part by weight of Tween™ 80 (all sterilized in advance) were mixed aseptically and emulsified with a ultrasound homogenizer. The resultant emulsion was put into a vial and sealed to give an internal liquid agent.

[FORMULATION EXAMPLE 11] Injection

10 parts by weight of 2-¹³C-glycerol was dissolved in 90 parts by weight of physiological saline and sterilized by filtration with a Millipore™ filter. The filtrate was put into a vial and sealed to give an injection.

[FORMULATION EXAMPLE 12] Internal Liquid Agent

10 parts by weight of 2-¹³C-glycerol was dissolved in 90 parts by weight of DDW and sterilized by filtration with a Millipore™ filter. The filtrate was put into a vial and sealed to give an internal liquid agent.

[FORMULATION EXAMPLE 13] Injection

10 parts by weight of N,N-dimethyl-¹³C-aminopyrin was dissolved in 90 parts by weight of physiological saline and sterilized by filtration with a Millipore™ filter. The filtrate was put into a vial and sealed to give an injection.

[FORMULATION EXAMPLE 14] Internal Liquid Agent

10 parts by weight of N,N-dimethyl-¹³C-aminopyrin was dissolved in 90 parts by weight of DDW and sterilized by filtration with a Millipore™ filter. The filtrate was put into a vial and sealed to give an internal liquid agent.

What is claimed is:

1. A diagnostic agent for diabetes, comprising an amino acid labelled with ^{13}C at least at one specific position.
2. The diagnostic agent according to claim 1, wherein the amino acid is an aliphatic amino acid.
3. The diagnostic agent according to claim 1, wherein the amino acid is an aromatic amino acid.
4. The diagnostic agent according to claim 1, wherein the amino acid is a sulfur-containing amino acid.
5. The diagnostic agent according to claim 1, wherein the amino acid is a heterocyclic amino acid.
6. The diagnostic agent according to claim 1, wherein the amino acid is an acidic amino acid.
7. The diagnostic agent according to claim 1, wherein the amino acid is a basic amino acid.
8. The diagnostic agent according to claim 2, wherein the aliphatic amino acid is glycine, alanine, serine, threonine, valine, leucine or isoleucine.
9. The diagnostic agent according to claim 3, wherein the aromatic amino acid is phenylalanine or tyrosine.
10. The diagnostic agent according to claim 4, wherein the sulfur-containing amino acid is cysteine, cystine or methionine.

11. The diagnostic agent according to claim 5, wherein the heterocyclic amino acid is tryptophan, proline or histidine.
12. The diagnostic agent according to claim 6, wherein the acidic amino acid is aspartic acid, asparagine, glutamic acid or glutamine.
13. The diagnostic agent according to claim 7, wherein the basic amino acid is arginine, lysine or ornithine.
14. A method for diagnosing diabetes, comprising:
administering to a subject an amino acid labelled with ^{13}C at least at one specific position;
measuring ^{13}C levels in the exhaled CO_2 for a specific period of time, and
comparing the degree of increase of ^{13}C levels of the subject to the corresponding degrees of normal subjects.
15. The method according to claim 14, wherein the amino acid is an aliphatic amino acid.
16. The method according to claim 14, wherein the amino acid is an aromatic amino acid.
17. The method according to claim 14, wherein the amino acid is a sulfur-containing amino acid.
18. The method according to claim 14, wherein the amino acid is a heterocyclic amino acid.
19. The method according to claim 14, wherein the amino acid is an acidic amino acid.

20. The method according to claim 14, wherein the amino acid is a basic amino acid.

21. The diagnostic agent according to claim 15, wherein the aliphatic amino acid is glycine, alanine, serine, threonine, valine, leucine or isoleucine.

22. The diagnostic agent according to claim 16, wherein the aromatic amino acid is phenylalanine or tyrosine.


23. The diagnostic agent according to claim 17, wherein the sulfur-containing amino acid is cysteine, cystine or methionine.

24. The diagnostic agent according to claim 18, wherein the heterocyclic amino acid is tryptophan, proline or histidine.

25. The diagnostic agent according to claim 19, wherein the acidic amino acid is aspartic acid, asparagine, glutamic acid or glutamine.

26. The diagnostic agent according to claim 20, wherein the basic amino acid is arginine, lysine or ornithine.


FIG. 1



Breath Sampling System from Rat

A rat anesthetized by intraperitoneal administration of Nembutal is fixed on its back. The head is covered with a cylindrical tube. A test sample is administered from the femoral vein. The exhalation is sucked with a stroke pump and introduced directly into a $^{13}\text{CO}_2$ analyzer. A Perma Pure Drier is located in the passage to remove moisture in the exhalation.


FIG. 2



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Galactose

$1\text{-}^{13}\text{C}$ -galactose (100 mg/kg) was intravenously administered to normal rats (male SD; 9-week-old; fasting blood sugar level 74.8 ± 5.1 mg/dl; $n=4$) and diabetic rats (male SD; 9-week-old; casual blood sugar level 423.3 ± 87.5 mg/dl; fasting blood sugar level 94 ± 18.5 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}(\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 3



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $2\text{-}^{13}\text{C}$ -Galactose

$2\text{-}^{13}\text{C}$ -galactose (100 mg/kg) was intravenously administered to normal rats (male SD; 9-week-old; fasting blood sugar level 63.8 ± 13.4 mg/dl; $n=4$) and diabetic rats (male SD; 9-week-old; casual blood sugar level 371.5 ± 63.4 mg/dl; fasting blood sugar level 86.5 ± 22.6 mg/dl; $n=3$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured. All the rats were fasted overnight before the experiment.


FIG. 4



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Fructose

$1\text{-}^{13}\text{C}$ -fructose (100 mg/kg) was intravenously administered to normal rats (male SD; 8-week-old; fasting blood sugar level 75.3 ± 4.1 mg/dl; $n=4$) and diabetic rats (male SD; 8-week-old; casual blood sugar level 498.5 ± 75.1 mg/dl; fasting blood sugar level 99.3 ± 11.8 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}(\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 5



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $2\text{-}^{13}\text{C}$ -Fructose

$2\text{-}^{13}\text{C}$ -fructose (100 mg/kg) was intravenously administered to normal rats (male SD; 8-week-old; fasting blood sugar level 75 ± 3.1 mg/dl; $n=4$) and diabetic rats (male SD; 8-week-old; casual blood sugar level 533.5 mg/dl; fasting blood sugar level 86 mg/dl; $n=2$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C} (\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 6



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $\text{U-}^{13}\text{C}$ -Starch

$\text{U-}^{13}\text{C}$ -starch (30 mg/kg) was orally administered to normal rats (male SD; 9-week-old; fasting blood sugar level 57.7 ± 4.5 mg/dl; $n=3$) and diabetic rats (male SD; 9-week-old; casual blood sugar level 398.3 ± 24.2 mg/dl; fasting blood sugar level 76.5 ± 7.9 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured. All the rats were fasted overnight before the experiment.


FIG. 7



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Isoleucine

$1\text{-}^{13}\text{C}$ -isoleucine (20 mg/kg) was intravenously administered to normal rats (male SD; 9-week-old; fasting blood sugar level 80.8 ± 5.5 mg/dl; $n=4$) and diabetic rats (male SD; 9-week-old; casual blood sugar level 368.7 ± 55.7 mg/dl; fasting blood sugar level 81.3 ± 6.2 mg/dl; $n=3$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}(\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 8



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Alanine

$1\text{-}^{13}\text{C}$ -alanine (10 mg/kg) was intravenously administered to normal rats (male SD; 8-week-old; fasting blood sugar level 79.5 ± 4.5 mg/dl; $n=4$) and diabetic rats (male SD; 8-week-old; casual blood sugar level 445 ± 20.9 mg/dl; fasting blood sugar level 100.8 ± 15.0 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}(\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 9



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Histidine

$1\text{-}^{13}\text{C}$ -histidine (30 mg/kg) was intravenously administered to normal rats (male SD; 9-week-old; fasting blood sugar level 73.8 ± 2.3 mg/dl; $n=4$) and diabetic rats (male SD; 9-week-old; casual blood sugar level 417.7 ± 36.6 mg/dl; fasting blood sugar level 106.7 ± 5.4 mg/dl; $n=3$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}(\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 10



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Valine

$1\text{-}^{13}\text{C}$ -valine (20 mg/kg) was intravenously administered to normal rats (male SD; 9-week-old; fasting blood sugar level 76.5 ± 9.9 mg/dl; $n=4$) and diabetic rats (male SD; 9-week-old; casual blood sugar level 430 ± 30.5 mg/dl; fasting blood sugar level 99.8 ± 11.1 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured. All the rats were fasted overnight before the experiment.


FIG. 11



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of 1,2- ^{13}C -Ornithine

1,2- ^{13}C -ornithine (20 mg/kg) was intravenously administered to normal rats (male SD; 8-week-old; fasting blood sugar level 67 ± 5.8 mg/dl; $n=4$) and diabetic rats (male SD; 8-week-old; casual blood sugar level 411.8 ± 102.5 mg/dl; fasting blood sugar level 77 ± 5.7 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}(\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 12



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Methionine

$1\text{-}^{13}\text{C}$ -methionine (40 mg/kg) was intravenously administered to normal rats (male SD; 10-week-old; fasting blood sugar level 73.3 ± 2.8 mg/dl; $n=4$) and diabetic rats (male SD; 10-week-old; casual blood sugar level 590.5 ± 9.9 mg/dl; fasting blood sugar level 84 ± 5.8 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured. All the rats were fasted overnight before the experiment.


FIG. 13



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Arginine

$1\text{-}^{13}\text{C}$ -arginine (50 mg/kg) was intravenously administered to normal rats (male SD; 8-week-old; fasting blood sugar level 68.8 ± 6.9 mg/dl; $n=4$) and diabetic rats (male SD; 8-week-old; casual blood sugar level 419.5 ± 69.3 mg/dl; fasting blood sugar level 83.3 ± 11.8 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C} (\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 14



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Tryptophan

$1\text{-}^{13}\text{C}$ -tryptophan (10 mg/kg) was intravenously administered to normal rats (male SD; 9-week-old; fasting blood sugar level 61 ± 1.6 mg/dl; $n=4$) and diabetic rats (male SD; 9-week-old; casual blood sugar level 487 ± 56.5 mg/dl; fasting blood sugar level 94.7 ± 10.4 mg/dl; $n=3$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}(\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 15



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Serine

$1\text{-}^{13}\text{C}$ -serine (50 mg/kg) was intravenously administered to normal rats (male SD; 10-week-old; fasting blood sugar level 69.8 ± 3.8 mg/dl; $n=4$) and diabetic rats (male SD; 10-week-old; casual blood sugar level 569 ± 50.3 mg/dl; fasting blood sugar level 93.3 ± 7.6 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}(\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 16



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Tyrosine

$1\text{-}^{13}\text{C}$ -tyrosine (60 mg/kg) was orally administered to normal rats (male SD; 9-week-old; fasting blood sugar level 79.8 ± 3.9 mg/dl; $n=4$) and diabetic rats (male SD; 9-week-old; casual blood sugar level 442 ± 28.6 mg/dl; fasting blood sugar level 86.5 ± 7.4 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured. All the rats were fasted overnight before the experiment.


FIG. 17



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Glutamine

$1\text{-}^{13}\text{C}$ -glutamine (20 mg/kg) was intravenously administered to normal rats (male SD; 8-week-old; fasting blood sugar level 72.8 ± 4.6 mg/dl; $n=4$) and diabetic rats (male SD; 8-week-old; casual blood sugar level 491 mg/dl; fasting blood sugar level 88.5 mg/dl; $n=2$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured. All the rats were fasted overnight before the experiment.


FIG. 18



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Lysine

$1\text{-}^{13}\text{C}$ -lysine (50 mg/kg) was intravenously administered to normal rats (male SD; 8-week-old; fasting blood sugar level 71 ± 1.6 mg/dl; $n=3$) and diabetic rats (male SD; 8-week-old; casual blood sugar level 431 ± 71.1 mg/dl; fasting blood sugar level 90.3 ± 4.9 mg/dl; $n=3$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}(\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 19



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Glutamic Acid

$1\text{-}^{13}\text{C}$ -glutamic acid (10 mg/kg) was intravenously administered to normal rats (male SD; 8-week-old; fasting blood sugar level 70.8 ± 8.3 mg/dl; $n=4$) and diabetic rats (male SD; 8-week-old; casual blood sugar level 487.7 ± 38.4 mg/dl; fasting blood sugar level 108 ± 8.8 mg/dl; $n=3$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}(\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 20



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Proline

$1\text{-}^{13}\text{C}$ -proline (20 mg/kg) was intravenously administered to normal rats (male SD; 10-week-old; fasting blood sugar level 77.6 ± 7.3 mg/dl; $n=5$) and diabetic rats (male SD; 10-week-old; casual blood sugar level 430.3 ± 45.2 mg/dl; fasting blood sugar level 97.3 ± 12.5 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}(\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 21



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Glycine

$1\text{-}^{13}\text{C}$ -glycine (20 mg/kg) was intravenously administered to normal rats (male SD; 9-week-old; fasting blood sugar level 75.5 ± 5.8 mg/dl; $n=4$) and diabetic rats (male SD; 9-week-old; casual blood sugar level 410 ± 10.7 mg/dl; fasting blood sugar level 90 ± 11.9 mg/dl; $n=3$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}(\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 22



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Aspartic Acid

$1\text{-}^{13}\text{C}$ -aspartic acid (10 mg/kg) was intravenously administered to normal rats (male SD; 8-week-old; fasting blood sugar level 56.3 ± 6.8 mg/dl; $n=4$) and diabetic rats (male SD; 8-week-old; casual blood sugar level 389 ± 52.1 mg/dl; fasting blood sugar level 80.5 ± 6.1 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C} (\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 23



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1,1\text{-}^{13}\text{C}$ -Cystine

$1,1\text{-}^{13}\text{C}$ -cystine (45 mg/kg) was orally administered to normal rats (male SD; 9-week-old; fasting blood sugar level 65.5 ± 6.4 mg/dl; $n=4$) and diabetic rats (male SD; 9-week-old; casual blood sugar level 324.5 ± 7.7 mg/dl; fasting blood sugar level 81.8 ± 11.5 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C} (\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 24



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $3\text{-}^{13}\text{C}$ -Lactic Acid

$3\text{-}^{13}\text{C}$ -sodium lactate (50 mg/kg) was intravenously administered to normal rats (male SD; 8-week-old; fasting blood sugar level 67.8 ± 6.5 mg/dl; $n=4$) and diabetic rats (male SD; 8-week-old; casual blood sugar level 412.8 ± 71.7 mg/dl; fasting blood sugar level 88.5 ± 17.2 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C} (\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 25



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Acetic Acid

$1\text{-}^{13}\text{C}$ -acetic acid (10 mg/kg) was intravenously administered to normal rats (male SD; 8-week-old; fasting blood sugar level 73.8 ± 5.1 mg/dl; $n=4$) and diabetic rats (male SD; 8-week-old; casual blood sugar level 398.3 ± 57.5 mg/dl; fasting blood sugar level 75.0 ± 8.3 mg/dl; $n=3$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C} (\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 26



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Oleic Acid

$1\text{-}^{13}\text{C}$ -oleic acid (70 mg/kg) was intravenously administered to normal rats (male SD; 9-week-old; fasting blood sugar level 74.8 ± 9.8 mg/dl; $n=4$) and diabetic rats (male SD; 9-week-old; casual blood sugar level 402.5 ± 31.9 mg/dl; fasting blood sugar level 83.5 ± 4.4 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured. All the rats were fasted overnight before the experiment.


FIG. 27



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Palmitic Acid

$1\text{-}^{13}\text{C}$ -palmitic acid (50 mg/kg) was intravenously administered to normal rats (male SD; 9-week-old; fasting blood sugar level 74.8 ± 7.1 mg/dl; $n=4$) and diabetic rats (male SD; 9-week-old; casual blood sugar level 422.8 ± 42.0 mg/dl; fasting blood sugar level 80.8 ± 5.1 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}(\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 28



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Octanoic Acid

$1\text{-}^{13}\text{C}$ -octanoic acid (30 mg/kg) was intravenously administered to normal rats (male SD; 9-week-old; fasting blood sugar level 78.3 ± 4.7 mg/dl; $n=4$) and diabetic rats (male SD; 9-week-old; casual blood sugar level 440.3 ± 38.1 mg/dl; fasting blood sugar level 89.8 ± 9.4 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C} (\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 29



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of 1,1,1- ^{13}C -Trioctanoin

1,1,1- ^{13}C -trioctanoin (50 mg/kg) was orally administered to normal rats (male SD; 8-week-old; fasting blood sugar level 75.5 ± 8.4 mg/dl; $n=4$) and diabetic rats (male SD; 8-week-old; casual blood sugar level 406.8 ± 61.4 mg/dl; fasting blood sugar level 76.8 ± 6.1 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C} (\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 30



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1,1,1\text{-}^{13}\text{C}$ -Triacetin

$1,1,1\text{-}^{13}\text{C}$ -triacetin (10 mg/kg) was intravenously administered to normal rats (male SD; 9-week-old; fasting blood sugar level 67.7 ± 6.8 mg/dl; $n=3$) and diabetic rats (male SD; 9-week-old; casual blood sugar level 332 ± 44.2 mg/dl; fasting blood sugar level 80.3 ± 15.0 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured. All the rats were fasted overnight before the experiment.


FIG. 31



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $2\text{-}^{13}\text{C}$ -Glycerol

$2\text{-}^{13}\text{C}$ -glycerol (50 mg/kg) was intravenously administered to normal rats (male SD; 9-week-old; fasting blood sugar level 70.3 ± 4.1 mg/dl; $n=4$) and diabetic rats (male SD; 9-week-old; casual blood sugar level 440 ± 18.0 mg/dl; fasting blood sugar level 83 ± 9.3 mg/dl; $n=3$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}(\text{‰})$) were measured. All the rats were fasted overnight before the experiment.


FIG. 32



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of N,N-Dimethyl- ^{13}C -Aminopyrin


N,N-dimethyl- ^{13}C -aminopyrin (40 mg/kg) was intravenously administered to normal rats (male SD; 8-week-old; fasting blood sugar level 68.8 ± 4.0 mg/dl; $n=4$) and diabetic rats (male SD; 8-week-old; casual blood sugar level 414.5 ± 29.9 mg/dl; fasting blood sugar level 94.8 ± 10.0 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured. All the rats were fasted overnight before the experiment.

FIG. 33



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of Ethoxy-1- ^{13}C -Phenacetin

Ethoxy-1- ^{13}C -phenacetin (90 mg/kg) was orally administered to normal rats (male SD; 9-week-old; fasting blood sugar level 72 ± 8.9 mg/dl; $n=4$) and diabetic rats (male SD; 9-week-old; casual blood sugar level 504.7 ± 55.0 mg/dl; fasting blood sugar level 101.7 ± 3.3 mg/dl; $n=3$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}(\text{‰})$) were measured. All the rats were fasted overnight before the experiment.



Increase of $^{13}\text{CO}_2$ in Exhalation after
Administration of $1\text{-}^{13}\text{C}$ -Galactose

$1\text{-}^{13}\text{C}$ -galactose (100 mg/kg) was intravenously administered to normal rats (male SD; 9-week-old; fasting blood sugar level 74.8 ± 5.1 mg/dl; $n=4$) and diabetic rats (male SD; 9-week-old; casual blood sugar level 423.3 ± 87.5 mg/dl; fasting blood sugar level 94 ± 18.5 mg/dl; $n=4$) at time 0. Then, degrees of increase of ^{13}C levels in exhaled CO_2 ($\Delta^{13}\text{C}$ (‰)) were measured. All the rats were fasted overnight before the experiment.