

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 December 2003 (18.12.2003)

PCT

(10) International Publication Number
WO 03/103680 A1

(51) International Patent Classification⁷: **A61K 31/555**,
31/409, C07D 487/22

(74) Agent: **WILSON, Mary, J.**; Nixon & Vanderhye P.C.,
Suite 800, 1100 North Glebe Road, Arlington, VA 22201-
4714 (US).

(21) International Application Number: PCT/US03/18099

(22) International Filing Date: 9 June 2003 (09.06.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/386,454 7 June 2002 (07.06.2002) US

(71) Applicant (for all designated States except US): **DUKE UNIVERSITY** [US/US]; Office of Science and Technology, P.O. Box 90083, Durham, NC 27708-0083 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BATINIC-HABERLE, Ines** [HR/US]; c/o Duke University, Office of Science and Technology, P.O. Box 90083, Durham, NC 27708-0083 (US). **SPASOJEVIC, Ivan** [HR/US]; c/o Duke University, Office of Science and Technology, P.O. Box 90083, Durham, NC 27708-0083 (US). **FRIDOVICH, Irwin** [US/US]; c/o Duke University, Office of Science and Technology, P.O. Box 90083, Durham, NC 27708-0083 (US).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SUBSTITUTED PORPHYRINS

(57) Abstract: A series of ortho isomers of meso tetrakis N-alkylpyridylporphyrins (alkyl being methyl, ethyl, n-propyl, n-butyl, n-hexyl, and n-octyl) and their Mn(III) complexes were synthesized and characterized by elemental analysis, uv/vis spectroscopy, electrospray ionization mass spectrometry and electrochemistry. An increase in the number of carbon atoms in the alkyl chains from 1 to 8 is accompanied by an increase in: (a) lipophilicity measured by the chromatographic retention factor, R_f ; (b) metal-centered redox potential, $E_{1/2}$ from +220 to +367 mV vs NHE, and (c) proton dissociation constant, pK_{a2} from 10.9 to 13.2. A linear correlation was found between $E_{1/2}$ and R_f of the Mn(III) porphyrins and between the pK_{a2} and R_f of the metal-free compounds. As the porphyrins become increasingly more lipophilic, the decrease in hydration disfavors the separation of charges, while enhancing the electron-withdrawing effect of the positively charged pyridyl nitrogen atoms. Consequently, the $E_{1/2}$ increases linearly with the increase in pK_{a2} , a trend in porphyrin basicity opposite from the one we previously reported for other water-soluble Mn(III) porphyrins. All of these Mn(III) porphyrins are potent catalysts for superoxide dismutation (disproportionation). Despite the favorable increase of $E_{1/2}$ with the increase in chain length, the catalytic rate constant decreases from methyl ($\log k_{cat} = 7.79$) to n-butyl, and then increases such that the n-octyl is as potent an SOD mimic as are the methyl and ethyl compounds. The observed behavior originates from an interplay of hydration and steric effects that modulate electronic effects.



WO 03/103680 A1

This application claims priority from Provisional Application No. 60/386,454, filed June 7, 2002, the content of which is incorporated herein by reference.

Introduction

Low-molecular weight catalytic scavengers of reactive oxygen and nitrogen species, aimed at treating oxidative stress injuries, have been actively sought. Three major groups of manganese complexes have been developed and tested *in vitro* and *in vivo*; Mn porphyrins,¹⁻⁹ Mn cyclic polyamines¹⁰ and Mn salen derivatives.¹¹ Based on a structure-activity relationships that we developed for water-soluble Mn(III) and Fe(III) porphyrins,²⁻⁴ Mn(III) *meso* tetrakis(*N*-methylpyridinium-2-yl)porphyrin (Mn^{III}TM-2-PyP⁵⁺, AEOL-10112) and *meso* tetrakis(*N*-ethylpyridinium-2-yl)porphyrins (Mn^{III}TE-2-PyP⁵⁺, AEOL-10113) were proposed and then shown to be potent catalysts for superoxide dismutation.^{4,12} The alkyl substitutions at the *ortho* positions restrict the rotation of the pyridyl rings with respect to the porphyrin plane. Consequently both compounds exist as mixtures of four atropoisomers, all of which were shown to be equally potent catalysts for O₂⁻ dismutation.¹³ These Mn porphyrins also allow SOD-deficient *Escherichia coli* to grow under aerobic conditions,^{4,12} and offer protection in rodent models of oxidative stress such as stroke,¹⁴ diabetes,¹⁵ sickle cell disease,¹⁶ and cancer/radiation.¹⁷ The high formal +5 charge of these metalloporphyrins could influence their tissue distribution, transport across biological membranes, and binding to other biomolecules and their low lipophilicities may restrict their protective effects. With the aim of modulating metalloporphyrin subcellular distribution, higher *N*-alkylpyridylporphyrin analogues (Scheme I) with increased lipophilicity were synthesized. We anticipate that their comparative kinetic and thermodynamic characterization will deepen our insight into the modes of action of porphyrin-based catalytic antioxidants and the mechanisms of oxidative stress injuries.

BRIEF DESCRIPTION OF THE DRAWINGS**Scheme**

Scheme I. Structures of the most hydrophilic ($\text{Mn}^{\text{III}}\text{TM}-2\text{-PyP}^{5+}$) and the most lipophilic ($\text{Mn}^{\text{III}}\text{TnOct}-2\text{-PyP}^{5+}$) members of the series studied. The $\alpha\beta\alpha\beta$ atropoisomers are shown.

Figures

Figure 1. The lipophilicity, R_f of $\text{H}_2\text{T}(\text{alkyl})-2\text{-PyP}^{4+}$ (A) and $\text{Mn}^{\text{III}}\text{T}(\text{alkyl})-2\text{-PyP}^{5+}$ compounds (B) vs the number of CH_2 groups.

Figure 2. Proton dissociation constants pK_{a2} of the metal-free porphyrins, $\text{H}_2\text{T}(\text{alkyl})-2\text{-PyP}^{4+}$ (A), and the metal-centered redox potentials $E_{1/2}$ for the $\text{Mn}(\text{III})/\text{Mn}(\text{II})$ couple of $\text{Mn}^{\text{III}}\text{T}(\text{alkyl})-2\text{-PyP}^{5+}$ porphyrins (B) as a function of R_f . Inserts: pK_{a2} (Figure 2A) and $E_{1/2}$ (Figure 2B) vs the number of CH_2 groups.

Figure 3. The reactivity of water-soluble $\text{Mn}(\text{III})$ porphyrins (A) (ref 4) and $\text{Mn}^{\text{III}}\text{T}(\text{alkyl})-2\text{-PyP}^{5+}$ porphyrins (B) as catalysts for O_2^- dismutation, expressed in terms of $\log k_{\text{cat}}$ vs $E_{1/2}$.

Figure 4. $E_{1/2}$ for the $\text{Mn}(\text{III})/\text{Mn}(\text{II})$ couple of $\text{Mn}^{\text{III}}\text{T}(\text{alkyl})-2\text{-PyP}^{5+}$ porphyrins vs pK_{a2} of the corresponding metal-free ligands. Insert: $E_{1/2}$ of water-soluble $\text{Mn}(\text{III})$ porphyrins vs pK_{a3} (data from ref 4); $\text{Mn}^{\text{III}}\text{TE}-2\text{-PyP}^{5+}$ (1), $\text{Mn}^{\text{III}}\text{TM}-2\text{-PyP}^{5+}$ (2), $\text{Mn}^{\text{III}}\text{PTnM}-2\text{-PyP}^{4+}$

(3), $\text{Mn}^{\text{III}}\text{TM-4-PyP}^{5+}$ (4), $\text{Mn}^{\text{III}}\text{TM-3-PyP}^{5+}$ (5), $\text{Mn}^{\text{III}}\text{T(2,6-Cl}_2\text{-3-SO}_3\text{-P)P}^{3-}$ (6),
 $\text{Mn}^{\text{III}}\text{T(TFTMA)P}^{5+}$ (7), $\text{Mn}^{\text{III}}\text{T(}\alpha\alpha\alpha\alpha\text{-2-MINP)P}^{5+}$ (8), $\text{Mn}^{\text{III}}\text{T(2,6-Cl}_2\text{-3-SO}_3\text{-P)P}^{3-}$ (9),
 $\text{Mn}^{\text{III}}\text{T(2,4,6-Me}_3\text{-3,5-(SO}_3\text{)}_2\text{-P)P}^{7-}$ (10), $\text{Mn}^{\text{III}}\text{(TMA)P}^{5+}$ (11), MnTSP^{3-} (12), MnTCPP^{3-}
(13), $\text{Mn}^{\text{III}}\text{hematoP}^-$ (14).


Figure S1. Cyclic voltammetry of 0.5 mM $\text{Mn}^{\text{III}}\text{TE-2-PyP}^{5+}$ and $\text{Mn}^{\text{III}}\text{TnHex-2-PyP}^{5+}$ porphyrins in a 0.05 M phosphate buffer (pH 7.8, 0.1 M NaCl) at a scan rate of 0.1 V/s.

Figure S2. Electrospray mass spectrometry of 0.5 mM solutions of $\text{H}_2\text{T(alkyl)2-PyP}^{4+}$ compounds in 1:1 water : acetonitrile at a cone voltage of 20 V.


Figure S3. Electrospray mass spectrometry of 0.5 mM solutions of $\text{Mn}^{\text{III}}\text{T(alkyl)2-PyP}^{5+}$ compounds in 1:1 water : acetonitrile at a cone voltage of 20 V.

Detailed Description of the Invention


The present invention relates to a compound of formula



I




II



III

or



IV,

wherein

each R is, independently, an C1-C12 alkyl (preferably, a C8 to C12 alkyl),

each A is, independently, hydrogen or an electron withdrawing group ,

M is a metal selected from the group consisting of

manganese, iron, copper, cobalt, nickel and zinc, and

Z^- is a counterion. In one embodiment, at least one A is a halogen.

The invention further relates to a method of protecting cells (eg mammalian cells) from oxidant-induced toxicity comprising contacting the cells with a protective amount of a compound as described above. The invention further relates to a method of treating a pathological condition of a patient resulting from oxidant-induced toxicity comprising administering to the patient an effective amount of such a compound. The invention also relates to a method of treating a pathological condition of a patient resulting from degradation of $NO^\cdot$, comprising administering to the patient an effective amount of a compound as described above. Additionally, the invention relates to a method of treating a patient for inflammatory lung disease comprising administering to the patient an effective amount of a compound as described above. The inflammatory lung disease can be a hyper-reactive airway disease. The disease can be asthma.

The entire content of all documents cited herein are incorporated herein by reference. Also incorporated herein by reference is Batinic-Haberle et al, J. Chem. Soc., Dalton Trans. 2002, 2689-2696.

Also incorporated by reference is US Appln. 09/880,125, filed June 14, 2001.

EXAMPLE**Experimental****Materials and Methods**

General. $\text{MnCl}_2 \times 4 \text{H}_2\text{O}$, and Baker-flex silica gel IB TLC plates were purchased from J. T. Baker. *N, N'*-dimethylformamide, ethyl *p*-toluenesulfonate, 2-propanol (99.5+ %), NH_4PF_6 (99.99 %), NaCl, sodium L-ascorbate (99+ %) and tetrabutylammonium chloride were from Aldrich, while xanthine, and ferricytochrome *c* were from Sigma. The *n*-propyl, *n*-butyl, *n*-hexyl and *n*-octyl esters of *p*-toluenesulfonic acid were from TCI America. Methanol (anhydrous, absolute), ethanol (absolute), acetone, ethyl ether (anhydrous), chloroform, EDTA and KNO_3 were from Mallinckrodt and acetonitrile was from Fisher Scientific. Xanthine oxidase was prepared by R. Wiley and was supplied by K. V. Rajagopalan.¹⁸ Catalase was from Boehringer, ultrapure argon from National Welders Supply Co., and tris buffer (ultrapure) was from ICN Biomedicals, Inc.

$\text{H}_2\text{T(alkyl)-2-PyP}^{4+}$. Tetrakis(2-pyridyl)porphyrin, $\text{H}_2\text{T-2-PyP}$ was purchased from Mid-Century Chemicals, Chicago, IL. The increased lipophilicity of the *n*-propyl, *n*-butyl, *n*-hexyl, and *n*-octyl analogues required a slight modification of the synthetic approach used for methyl and ethyl compounds.^{4,12} Typically, 100 mg of $\text{H}_2\text{T-2-PyP}$ was dissolved in 20 mL of DMF at 100 °C, followed by the addition of 4 mL of the corresponding *p*-toluenesulfonate. The course of *N*-alkylation was followed by thin-layer chromatography on silica gel TLC plates using 1:1:8 KNO_3 -saturated $\text{H}_2\text{O} : \text{H}_2\text{O} :$ acetonitrile as a mobile phase. While complete *N*-alkylation is achieved within a few

hours for the methyl analogue, the required time gradually increases and it took three and five days to prepare the n-hexyl and n-octyl compounds, respectively. Upon completion, for the methyl, ethyl and n-propyl compounds, the reaction mixture was poured into a separatory funnel containing 200 mL each of water and chloroform and shaken well. The chloroform layer was discarded and the extraction with CHCl_3 was repeated several times. The n-butyl, n-hexyl and n-octyl analogues are more lipophilic and tended to remain in the chloroform layer. Therefore, increasing amounts of methanol were added to the water/ CHCl_3 mixture in order to force the porphyrin into the aqueous/methanol layer. This layer was filtered and the porphyrin was precipitated as the PF_6^- salt by the addition of a concentrated aqueous solution of NH_4PF_6 . The precipitate was thoroughly washed with 1:1 2-propanol : diethylether in the case of methyl and ethyl compounds and with pure diethylether for the others. The precipitate was then dissolved in acetone, filtered and precipitated as the chloride salt by the addition of tetrabutylammonium chloride dissolved in acetone. The precipitate was washed thoroughly with acetone, and dried *in vacuo* at room temperature. Elemental analysis: $\text{H}_2\text{TnPr-2-PyPCL}_4 \times 12.5 \text{ H}_2\text{O}$ ($\text{C}_{52}\text{H}_{71}\text{N}_8\text{O}_{12.5}\text{Cl}_4$): Found: C, 54.2; H, 6.42; N, 9.91; Cl, 12.04. Calculated: C, 54.20; H, 6.18; N, 9.68; Cl, 12.25. $\text{H}_2\text{TnBut-2-PyPCL}_4 \times 10.5 \text{ H}_2\text{O}$ ($\text{C}_{56}\text{H}_{75}\text{N}_8\text{O}_{10.5}\text{Cl}_4$): Found: C, 57.16; H, 6.94; N, 9.513; Cl, 1.77. Calculated: C, 57.10; H, 6.41; N, 9.51; Cl, 12.03. $\text{H}_2\text{TnHex-2-PyPCL}_4 \times 11 \text{ H}_2\text{O}$ ($\text{C}_{64}\text{H}_{100}\text{N}_8\text{O}_{11}\text{Cl}_4$): Found: C, 59.19; H, 7.31; N, 8.61; Cl, 11.09. Calculated: C, 59.16; H, 7.751; N, 8.60; Cl, 10.91. $\text{H}_2\text{TnOct-2-PyPCL}_4 \times 13.5 \text{ H}_2\text{O}$ ($\text{C}_{64}\text{H}_{121}\text{N}_8\text{O}_{13.5}\text{Cl}_4$): Found: C, 59.37; H, 7.41; N, 7.73. Calculated: C, 59.37; H, 8.37; N, 7.69.

$\text{Mn}^{\text{III}}\text{T(alkyl)-2-PyP}^{5+}$. Metalation of the N-alkylated porphyrins was achieved

as described previously for the methyl and ethyl compounds.^{4,12} Metal incorporation became slower as the alkyl chains lengthened. Under same conditions (20-fold excess metal, 25 °C, pH 12.3) it occurs almost instantaneously for methyl and ethyl, within minutes for n-propyl, in ~30 minutes for n-butyl, in ~1 hour with the n-hexyl, and took several hours at 100 °C for the n-octyl porphyrin. The formation of the Mn(II) porphyrin and its oxidation to Mn(III) were clearly distinguishable steps when the *n*-hexyl and n-octyl analogues were metalated. As was the case with the metal-free ligands, the PF₆⁻ salts of Mn(III) n-propyl, n-butyl, n-hexyl and n-octyl compounds were washed only with diethylether. Elemental analysis: Mn^{III}TnPr-2-PyPCL₅ x 11.5 H₂O

(MnC₅₂H₇₅N₈O_{11.5}Cl₅): Found: C, 50.90; H, 6.07; N, 9.27; Cl, 13.48. Calculated: C, 50.85; H, 6.16; N, 9.12; Cl, 14.43. Mn^{III}TnBut-2-PyPCL₅ x 12.5 H₂O

(MnC₅₆H₈₅N₈O_{12.5}Cl₅): Found: C, 51.58; H, 6.33; N, 9.55; Cl, 15.53. Calculated: C, 51.64; H, 6.58; N, 8.60; Cl, 13.61. Mn^{III}TnHex-2-PyPCL₅ x 10.5 H₂O

(MnC₆₄H₉₇N₈O_{12.5}Cl₅): Found: C, 55.64; H, 7.14; N, 8.23; Cl, 12.60. Calculated: C, 55.76; H, 7.09; N, 8.13; Cl, 12.86. Mn^{III}TnOct-2-PyPCL₅ x 10 H₂O x 2.5 NH₄Cl

(MnC₆₄H₁₂₂N_{10.5}O₁₀Cl_{7.5}): Found: C, 53.56; H, 7.13; N, 9.12; Cl, 16.84. Calculated: C, 53.53; H, 7.60; N, 9.10; Cl, 16.46.

Thin-layer chromatography. All ligands and their Mn(III) complexes were chromatographed on silica gel TLC plates using 1:1:8 KNO₃-saturated H₂O : H₂O : acetonitrile. The atropoisomers could not be separated for the methyl¹⁹ and ethyl analogues,²⁻⁴ they begin to separate for the n-propyl and n-butyl species and were clearly resolved with the n-hexyl and n-octyl compounds.

Uv/vis spectroscopy. The uv/vis spectra were taken on a Shimadzu UV-2501 PC

spectrophotometer at 25 °C. The proton dissociation constants (pK_{a2}), were determined spectrophotometrically at 25 °C, at an ionic strength of 0.1 M (NaOH/NaNO₃), as previously described.⁴

Electrochemistry. Measurements were performed on a CH Instruments Model 600 Voltammetric Analyzer.^{3,4} A three-electrode system in a small volume cell (0.5 mL to 3 mL), with a 3 mm-diameter glassy carbon button working electrode (Bioanalytical Systems), plus the Ag/AgCl reference and Pt auxilliary electrodes was used. Solutions contained 0.05 M phosphate buffer, pH 7.8, 0.1 M NaCl, and 0.5 mM metalloporphyrin. The scan rates were 0.01-0.5 V/s, typically 0.1 V/s. The potentials were standardized against the potassium ferrocyanide/ferricyanide²⁰ and/or against Mn^{III}TE-2-PyP⁵⁺. All voltammograms were reversible.

Electrospray mass spectrometry. ESMS measurements were performed on a Micromass Quattro LC triple quadrupole mass spectrometer equipped with a pneumatically assisted electrostatic ion source operating at atmospheric pressure. Typically, the 0.5 mM 50% aqueous acetonitrile solutions of chloride salts of metal-free porphyrins or their Mn(III) complexes were introduced by loop injection into a stream of 50% aqueous acetonitrile flowing at 8 μ L/min. Mass spectra were acquired in continuum mode, scanning from 100-500 m/z in 5 s, with cone voltages of 20 V and 24 V. The mass scale was calibrated using polyethylene glycol.

Catalysis of O₂⁻ dismutation. We have previously shown that the O₂⁻/cytochrome c reduction assay gives the same catalytic rate constants as does pulse radiolysis for Mn^{III}TE-2-PyP⁵⁺, {Mn^{III}BVDME}₂, {Mn^{III}BV}₂ and MnCl₂.²¹ Therefore the convenient cytochrome c assay was used to characterize the series of Mn(III) N-

alkylpyridylporphyrins. The xanthine/xanthine oxidase reaction was the source of $O_2^{\cdot -}$ and ferricytochrome *c* was used as the indicating scavenger for $O_2^{\cdot -}$.²² The reduction of cytochrome *c* was followed at 550 nm. Assays were conducted at $(25 \pm 1)^\circ\text{C}$, in 0.05 M phosphate buffer, pH 7.8, 0.1 mM EDTA, in the presence and absence of 15 $\mu\text{g/mL}$ of catalase. Rate constants for the reaction of metalloporphyrins with $O_2^{\cdot -}$ were based upon competition with 10 μM cytochrome *c*, $k_{\text{cyt } c} = 2.6 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$ as described elsewhere.²¹ The $O_2^{\cdot -}$ was produced at the rate of 1.2 μM per minute. Any possible interference through inhibition of the xanthine/ xanthine oxidase reaction by the test compounds was examined by following the rate of urate accumulation at 295 nm in the absence of cytochrome *c*. No reoxidation of cytochrome *c* by the metalloporphyrins was observed.

Results

Thin Layer Chromatography. The increase in the length of the alkyl chains is accompanied by an increase in the lipophilicity of the compounds as indicated by the increase in the retention factor R_f (porphyrin path/solvent path) (Table 1, Figure 1). The apparent lag that was observed in the case of shorter chains with Mn(III) complexes (Figure 1B), is presumably due to their higher overall formal charge (+5 for the Mn(III) complexes, +4 for the ligand). As the chains lengthen, their contribution to the overall lipophilicity increases, and eventually the n-octyl porphyrin and its Mn(III) complex are more alike in R_f than are methyl analogues.

Uv/vis spectroscopy. Molar Absorptivities. The porphyrins obeyed the Beer-Lambert law from 10^{-7} M to 10^{-5} M , and the uv/vis data are given in Table 2. As the

length of alkyl chains increased from methyl to n-butyl a red shift of the Soret absorption maxima was generally observed, as well as an increase in the molar absorptivities, and these effects plateau beyond butyl compound. Such trends may be understood in terms of the interplay of porphyrin nucleus distortion (red shifts) and the electron-withdrawing (blue shifts) effect of the *N*-alkylpyridyls groups.^{12,23}

Metalation behavior and proton dissociation constants. The rates of Mn^{2+} incorporation at pH ~12.3 decreased with an increase in chain length. The same was found for the kinetics of Zn^{2+} and Cu^{2+} insertion into these compounds below pH 7, where the kinetics were first order in metal and porphyrin concentration.²⁴ Since the free-base porphyrin H_2P^{4+} reactants were mixtures of the four atropoisomers, each isomer has a similar metalation rate constant. As noted before for both water soluble and insoluble porphyrins, compounds with substituents in the *ortho* positions tend to metalate more slowly than derivatives with the same groups in the *meta* or *para* positions.²⁵⁻³⁴

The proton dissociation constants, K_{a2} and K_{a3} are defined as follows:



The $\text{p}K_{a2}$ values for the *N*-alkylpyridyl series are given in Table 1. As the alkyl chains lengthen the porphyrins become less hydrated and the separation of charges (eq [1]) becomes less favorable, *i.e.* $\text{p}K_{a2}$ increases (Figure 2 insert). Figure 2A shows the linear relationship between $\text{p}K_{a2}$ and R_f .

Equilibrium constants $\text{p}K_{a3}$ for reaction [2] are 1.8 for the *meta* $\text{H}_2\text{TM-3-PyP}^{4+}$,

1.4 for the *para* H₂TM-4-PyP⁴⁺, and -0.9 for *ortho* H₂TM-2-PyP⁴⁺.^{4,25} While the *meta* and *para* *N*-methylpyridylporphyrins are mixtures of protonated H₃P⁵⁺ and H₄P⁶⁺ species in 1.0 M HCl, the *ortho* substituted H₂TM-2-PyP⁴⁺ to H₂TnOct-2-PyP⁴⁺ compounds remain as the unprotonated free base H₂P⁴⁺ in 1.0 M HCl and in 1.0 M HNO₃. With *ortho*, *meta* and *para* *N*-methylpyridylporphyrins the pK_{a2} increases as the pK_{a3} increases.

The half-lives for the acid and anion-catalyzed removal of zinc from Zn *N*-methylated derivatives³⁵ in 1.0 M HNO₃ were 89 s for the *meta*, 165 s for the *para*, and 19 hours for the *ortho* ZnTM-2-PyP⁴⁺. No indication of zinc loss was found within a week for the ZnTnHex-2-PyP⁴⁺ compound.³⁶ Similar behavior is found in 1.0 M HCl, with t_{1/2} ranging from 21 s for the *meta* methyl to 76 hours for ZnTnOct-2-PyP⁴⁺.²⁴ In accord are the observations that when solid MnTnHex-2-PyP⁵⁺ was dissolved in 12 M HCl, the spectra did not change within 3 months, while over 50% of the Mn from Mn^{III}TM-2-PyP⁵⁺ species was lost within a month. In addition to porphyrin ring distortion,²⁹⁻³² the steric hindrance and solvation effects imposed by the progressively longer alkyl chains may also contribute to the differences in metalation/demetalation behavior.

Due to their high metal-centered redox potentials, the Mn(III) *meso* tetrakis *ortho* *N*-alkylpyridylporphyrins *in vivo* will be readily reduced with cell reductants such as ascorbic acid.^{2,3,12} The reduced Mn(II) porphyrins will also be transiently formed in the catalysis of O₂⁻ dismutation. Therefore, we also examined the behavior of the reduced and more biologically relevant Mn^{II}T(alkyl)-2-PyP⁴⁺ compounds. We compared the methyl, n-hexyl and n-octyl derivatives (6 μM) aerobically and anaerobically in the

presence of a 70-fold excess of ascorbic acid (pH 7.8, 0.1 M tris buffer) and in the presence and absence of a 150-fold excess of EDTA. Under anaerobic conditions both Mn(II) porphyrins were stable to Mn loss and porphyrin decomposition inside 24 hours. Aerobically, ~40% of Mn methyl but none of the Mn n-hexyl and n-octyl compounds underwent degradation within 125 min. The absorption spectral changes indicate that the degradation occurred through the Mn porphyrin catalyzed reduction of oxygen by ascorbate resulting in the formation of H₂O₂. The peroxide in turn causes porphyrin destruction. These observations are consistent with previous results which indicate that a more electron rich compound (Mn^{III}TM-2-PyP⁴⁺) reduces O₂ faster than does a more electron deficient species (Mn^{III}TnOct-2-PyP⁴⁺).^{2,3} EDTA did not significantly influence porphyrin degradation or Mn loss.

Electrochemistry. Cyclic voltammetry of the Mn(III) porphyrins shows a reversible voltammogram that we ascribe to the Mn(III)/Mn(II) redox couple. The metal-centered redox potentials, E_{1/2} are in Table 1 and the representative voltammograms of the Mn^{III/II}TE-2-PyP^{5+/4+} and Mn^{III/II}TnHex-2-PyP^{5+/4+} compounds are shown in the Supporting Material, Figure S1. Both lipophilicity (Figure 1B) and E_{1/2} (Figure 2B, insert) increase exponentially with the number of CH₂ groups in the alkyl chains. Consequently, the increase in E_{1/2} is a linear function of R_f (Figure 2B).

Electrospray mass spectrometry. The ESMS proved to be a valuable tool for accessing the properties of the free base porphyrins and their Mn complexes whereby the impact of structure on solvation, ion-pairing, redox properties, protonation/deprotonation, dealkylation, and catalytic properties are clearly depicted.

H₂T(alkyl)-2-PyP⁴⁺. The ESMS of the metal-free porphyrins obtained at the low cone voltage of 20 V showed dominant molecular ions assigned to $H_2P^{4+}/4$ and/or its mono-deprotonated analogue, $H_2P^{4+}-H^+/3$ (Table 3 and Supporting Material, Figures S2A-E). Negligible double deprotonation ($H_2P^{4+}-2H^+/2$) was noted. Only *H₂TM-2-PyP⁴⁺* gave rise to a high-intensity $H_2P^{4+}+H^+/5$ peak.

The ESMS shows a pronounced decrease in solvation by acetonitrile as the alkyl chains lengthen. Compared to the base peak, the relative intensities of the mono-solvated molecular ions range from 40% for methyl, 15% for ethyl, and <10% for the higher analogues. Only with the n-hexyl and n-octyl porphyrins are small peaks (<5%) from ions associated with chloride found.

From methyl to n-butyl, the ratio of the molecular ion to mono-deprotonated ion peaks decreases, consistent with the trend in pK_{a2} . Thus, the base peak for methyl is that of the molecular ion, while the base peak for the n-propyl and n-butyl porphyrins is the mono-deprotonated ion. This pK_{a2} trend is overcome by the higher lipophilicities of the n-hexyl and n-octyl compounds, where roughly equal-intensity molecular ion (100%) and mono-deprotonated ion (98%) peaks are observed. The loss of one alkyl group ($H_2P^{4+}-a^+/3$) was noted for all derivatives (except for the methyl), and either no or negligible loss of a second alkyl group ($H_2P^{4+}-2a^+/2$) was found.

Mn^{III}T(alkyl)-2-PyP⁵⁺. The ESMS of the Mn(III) complexes was done at a lower cone voltage (20 V) than in our previous study (30-58 V).³⁷ Therefore, less fragmentation occurs and more solvent-associated and ion-paired species could be observed (Table 4 and Supporting Material, Figures S3A-E). Solvation and ion pairing are more pronounced when compared with the metal-free ligands. The more lipophilic

Mn(III) porphyrins are more easily desolvated in the electrospray ionization source. In accordance with our previous observations, the ESMS also clearly reflects the redox properties of these compounds.^{37,38} The higher the $E_{1/2}$ the more reduced porphyrins are noted. Species solvated with acetonitrile or associated with chloride were observed with both Mn(III) and Mn(II) compounds. Two chlorides were associated only with Mn(III) porphyrins.

In the ESMS of the *n*-hexyl and *n*-octyl porphyrins we observed strong signals at m/z 337 and 375 that are assigned to compounds doubly reduced either at the metal ($Mn^I P^{3+}/3$) or at both the metal and porphyrin ring ($Mn^{II} P^{3+}/3$). Such doubly reduced manganese porphyrins should have a higher tendency to lose the metal, and indeed peaks for the metal-free species were found for the *n*-hexyl and *n*-octyl derivatives, while only traces of doubly reduced and demetalated species were found for *n*-butyl.

The ESMS behavior of Mn porphyrins changes sharply once the alkyl chains lengthen beyond butyl, as observed with corresponding metal-free analogues. No loss of methyl groups was detected.³⁷ As the chains lengthen up to butyl the loss of an alkyl group from Mn(III) and Mn(II) porphyrins becomes more pronounced and then the tendency decreases with *n*-hexyl and *n*-octyl. The same trend, but of lower intensity was noted for the loss of two alkyl groups. The ratio of mono-chlorinated Mn(III) to mono-chlorinated Mn(II) species decreases from methyl to *n*-butyl and then increases up to *n*-octyl. Thus the base peak of the methyl and ethyl porphyrins relates to $Mn^{III} P^{5+} + Cl^-/4$, while for the *n*-propyl and *n*-butyl derivatives it relates to $Mn^{II} P^{4+} + Cl^-/3$. Yet, with the *n*-hexyl, the $Mn^{III} P^{5+} + Cl^-/4$ and $Mn^{II} P^{4+} + Cl^-/3$ peaks are both of 100% intensity, and the di-chlorinated species ($Mn^{III} P^{5+} + 2Cl^-/3$) is of 86% intensity. With the *n*-octyl analogue, the

mono- and di-chlorinated species give rise to 100% $\text{Mn}^{\text{III}}\text{P}^{5+} + \text{Cl}^-/4$ and 89% $\text{Mn}^{\text{III}}\text{P}^{5+} + 2\text{Cl}^-/3$ peaks, and the third most intense (59%) signal relates to $\text{Mn}^{\text{II}}\text{P}^{4+} + \text{Cl}^-/3$. The lack of significant association of metal-free porphyrins with chloride observed here and elsewhere,³⁷ strongly supports the idea that chloride is bound to the metal. Furthermore, at the same cone voltage, the base peak of *ortho* MnTM-2-PyP^{5+} is the mono-chlorinated species, which was only 35% for *para* isomer. This suggests that the longer the chains, the more defined the cavity, which can hold up to two chloride ions, and the more stable is the Mn(III) state. While a species bearing two chlorides is hardly noted in $\text{Mn}^{\text{III}}\text{TM-2-PyP}^{5+}$, it is the second major peak in the ESMS of $\text{Mn}^{\text{III}}\text{TnOct-2-PyP}^{5+}$.


Catalysis of $\text{O}_2^{\cdot -}$ dismutation. None of the parent metal-free porphyrins exhibit any $\text{O}_2^{\cdot -}$ dismuting activity. All of the manganese compounds are potent catalysts of $\text{O}_2^{\cdot -}$ dismutation with $\log k_{\text{cat}}$ between 7.79 and 7.25. As shown in Table 1, $\log k_{\text{cat}}$ decreases from methyl to n-butyl and then increases, making n-octyl and methyl of comparable antioxidant potency.

Discussion

When designing metalloporphyrin SOD mimics we are aiming at approximating the redox properties of the enzyme active site. Superoxide dismutases catalyse the dismutation (disproportionation) of $\text{O}_2^{\cdot -}$ to H_2O_2 and O_2 at $\sim +300$ V vs NHE (pH 7.0).^{39,40} This potential is roughly midway ($+360$ mV vs NHE) between the potential for the reduction ($+890$ V vs NHE)⁴¹ and the oxidation of $\text{O}_2^{\cdot -}$ (-160 V vs NHE)⁴¹ thus

providing an equal driving force for both half-reactions in the catalytic cycle. The $O_2^{\cdot -}$ dismutation by Cu,Zn-SOD occurs with catalytic rate constant, $k_{cat} = k_{red} = k_{ox} \simeq 2 \times 10^9 M^{-1} s^{-1}$ ($\log k_{cat} \simeq 9.3$).⁴²⁻⁴⁴

We previously demonstrated a structure-activity relationship between $\log k_{cat}$ and the metal-centered $E_{1/2}$ of the Mn(III)/Mn(II) couple for a variety of water-soluble *meso* substituted porphyrins (Figure 3A).²⁻⁴ Electron-withdrawing substituents on the porphyrin ring shift $E_{1/2}$ towards more positive values resulting in higher values for k_{cat} .²⁻⁴ Each 120 mV increase in $E_{1/2}$ gave a 10-fold increase in k_{cat} ,⁴ consistent with the Marcus equation⁴⁵ for outer-sphere electron transfer reactions (Figure 3A). The Marcus equation is valid as long as one of the two steps in the catalytic dismutation cycle is



rate-limiting.

On the basis of such structure-activity relationships, the *ortho* isomers of Mn(III) *meso* tetrakis *N*-methyl- and *N*-ethylpyridylporphyrins were tested and proved to be potent catalysts of $O_2^{\cdot -}$ dismutation. Their $\log k_{cat}$ values are 7.79 and 7.76 and they operate at potentials (+220 and +228 V) similar to the potential of the enzyme itself. These two metalloporphyrins also exhibit protection in *in vivo* models of oxidative stress injuries.¹⁴⁻¹⁷ We have now extended our work to a series of $Mn^{III}T(alkyl)-2-PyP^{5+}$ compounds where alkyl is methyl, ethyl, *n*-propyl, *n*-butyl, *n*-hexyl, and *n*-octyl (Scheme I). The significant differences in lipophilicity along the series (Figure 1A), with retention

of catalytic potency (Table 1), might lead to favorably selective subcellular distributions of these new $\text{Mn}^{\text{III}}\text{T(alkyl)-2-PyP}^{5+}$ compounds and hence broader their utility.

$E_{1/2}$ vs $\text{pK}_{\text{a}2}$. We did not expect a profound change in $E_{1/2}$ along the series based on the fact that the increase in alkyl chain length from methyl to n-hexyl is without effect on the basicity of alkylamines.⁴⁶ However, we found that the metal-centered redox potentials varied from +220 mV for methyl to +367 mV (vs NHE) for the n-octyl compound. Such an increase in $E_{1/2}$ may originate from progressively unshielded positive charges at pyridyl nitrogens which would then exert stronger electron-withdrawing effect on the coordinated Mn as the compounds increase in lipophilicity. This reasoning is supported by the ESMS data (Table 4 and Supporting Material, Figures S3A-E) which show that the susceptibility to desolvation is accompanied by a greater preponderance of reduced Mn(II) porphyrin ions as the alkyl chains of the Mn complexes lengthen. We have previously reported⁴ that mainly electronic effects determine the relation between the pK_{a} of the metal-free porphyrin and the $E_{1/2}$ of the corresponding metal complex such that the decrease in $\text{pK}_{\text{a}3}$ is accompanied by a linear increase in $E_{1/2}$ (Figure 4, insert). However as the compounds become increasingly more lipophilic, the lack of solvation disfavors separation of charges (higher $\text{pK}_{\text{a}2}$ values), while the electron-withdrawing effects of the positively charged pyridyl nitrogens are enhanced. Thus the electronic $\text{pK}_{\text{a}2}$ effects are overcome by solvation/steric effects resulting in an inverted trend, *i. e.* the $E_{1/2}$ now increases in a linear fashion with an increase in $\text{pK}_{\text{a}2}$ (Figure 4).

$\text{Log } k_{\text{cat}}$ vs $E_{1/2}$. Based on a previously established structure-activity relationship for water-soluble Mn(III) porphyrins,⁴ we expected the 147 mV increase in $E_{1/2}$ to be accompanied by a ~12-fold increase in k_{cat} (Figure 3A).⁴ We actually found that k_{cat}

decreased from methyl to n-butyl, and then increased by the same factor of ~3 to n-octyl (Table 1, Figure 3B). One explanation is that the Mn porphyrins are solvated to different extents, as indicated by the ESMS data, and this in turn affects the magnitude of k_{cat} . The trend in k_{cat} may also be influenced by the electrostatic/steric effects originating from the shielding of the single positive charge on the Mn(III) center. Thus the difference in the magnitude of lipophilicity between the metal-free ligands (formally +4) and the Mn(III) complexes (formally +5) becomes less noticeable as the alkyl chains get longer (Table 1). These H_2P^{4+} compounds of formal +4 charge behave in solution kinetically as +1.6 to +1.8 electrolytes.²⁴ From methyl to n-butyl, $\log k_{cat}$ decreases almost linearly (Figure 3B, insert). Due to the exponential increase in $E_{1/2}$ along the series of Mn porphyrins (Figure 2B, insert), the unfavorable electrostatic/steric effects are in part opposed and finally overcome by the progressively more favorable redox potentials that originate from increased desolvation (lipophilicity). Consequently, the very lipophilic n-octyl compound is essentially as potent an SOD mimic as the less lipophilic methyl and ethyl derivatives.

Regan *et al*⁴⁷ were able to uncouple the steric and solvation effects in reactions of chloride ions with methyl- and *tert*-butyl-substituted chloroacetonitrile, and showed that both were of comparable magnitudes. Similarly, the reactivity of *N*-alkylpyridylporphyrins are the result of the interplay of electronic, steric and solvation effects, the latter dominating with the more lipophilic members of the series.

Recent findings indicate that biologically relevant reactions, other than $O_2^{\cdot -}$ dismutation, can occur at the metal center in Mn porphyrins.^{2,3,5,7,8,48-52} The same has been reported for the enzyme active site,^{20,53-57} thus raising the complexity of the free

radical chemistry and biology of the enzymes and their mimics. Reactive oxygen and nitrogen species are involved in direct damage of key biological targets such as nucleic acids, proteins and fatty acids, and there is an increasing amount of evidence that such species are also involved in the modulation of signaling processes.^{14,58,59} Thus, it is important to understand the mechanisms of action of Mn porphyrins and related compounds. Based on the electrostatic, steric, solvation, and lipophilic effects observed in this study, we expect the members of *N*-alkylpyridyl series to differ one from another in *in vivo* models of oxidative stress injuries with respect to their specificity towards reactive oxygen and nitrogen species as well as with regard to their pharmacokinetics. Such work is in progress.

Abbreviations

SOD, superoxide dismutase; AN, acetonitrile; DMF, *N,N'*-dimethylformamide; NHE, normal hydrogen electrode; TLC, thin-layer chromatography; H_2P^{4+} , any *meso* tetrakis *N*-alkylpyridylporphyrin ligand; $Mn^{III/II}P^{4+/5+}$, any Mn(III/II) *meso* tetrakis *N*-alkylpyridylporphyrin; *meso* refers to the substituents at the 5,10,15, and 20 (*meso* carbon) position of the porphyrin core. $Mn^{III}T(alkyl)-2(3,4)-PyP^{5+}$, manganese(III) *meso* tetrakis(*N*-methyl, *N*-ethyl, *N*-n-propyl, *N*-n-butyl, *N*-n-hexyl, *N*-n-octyl)pyridinium-2(3,4)-ylporphyrin; alkyl is M, methyl; E, ethyl; nPr, n-propyl; nBu, n-butyl; nHex, n-hexyl; nOct, n-octyl on the pyridyl ring; 2 is the *ortho*, 3, the *meta* and 4 the *para* isomer; $Mn^{III}TM-2-PyP^{5+}$ is AEOL-10112, and $Mn^{III}TE-2-PyP^{5+}$ is AEOL-10113; $Mn^{III}PTr(M-2-PyP^{4+})$, manganese(III) 5-phenyl-10,15,20-tris(*N*-methylpyridinium-2-yl)porphyrin; $Mn^{III}BM-2-PyP^{3+}$, manganese(III) *meso* bis(2-pyridyl)-bis(*N*-methylpyridinium-2-yl)porphyrin; $Mn^{III}TrM-2-PyP^{4+}$, 5-(2-pyridyl)-10,15,20-tris(*N*-methylpyridinium-2-yl)porphyrin; $Mn^{III}T(TMA)P^{5+}$, manganese(III) *meso* tetrakis(*N,N,N*-trimethylanilinium-4-yl)porphyrin; $Mn^{III}T(TFTMA)P^{5+}$, manganese(III) *meso* tetrakis(2,3,5,6-tetrafluoro-*N,N,N*-trimethylanilinium-4-yl)porphyrin; $Mn^{III}TCPP^{3+}$, manganese *meso* tetrakis(4-carboxylatophenyl)porphyrin; $MnTSPP^{3+}$,

manganese(III) *meso* tetrakis(4-sulfonatophenyl)porphyrin; $\text{Mn}^{\text{III}}\text{T}(2,6\text{-Cl}_2\text{-3-SO}_3\text{-P})\text{P}^{3-}$, manganese(III) *meso* tetrakis(2,6-dichloro-3-sulfonatophenyl)porphyrin; $\text{Mn}^{\text{III}}\text{T}(2,6\text{-F}_2\text{-3-SO}_3\text{-P})\text{P}^{3-}$, manganese(III) *meso* tetrakis(2,6-difluoro-3-sulfonatophenyl)porphyrin; $\text{Mn}^{\text{III}}\text{T}(2,4,6\text{-Me}_3\text{-3,5-(SO}_3)_2\text{-P})\text{P}^{7-}$, manganese(III) 5,10,15,20-tetrakis(2,4,6-trimethyl-3,5-disulfonatophenyl)porphyrin; $\text{Mn}^{\text{III}}\text{hematoP}^-$, manganese(III) hematoporphyrin IX.

References

1. R. F. Pasternack, A. Banth, J. M. Pasternack and C. S. Johnson, *J. Inorg. Biochem.* 1981, **15**, 261 (b) R. F. Pasternack and B. J. Halliwell, *J. Am. Chem. Soc.* 1979, **101**, 1026.
2. I. Batinić-Haberle, *Methods Enzymol.* 2002, **349**, 223.
3. I. Spasojević and I. Batinić-Haberle, *Inorg. Chim. Acta*, 2001, **317**, 230.
4. I. Batinić-Haberle, I. Spasojević, P. Hambright, L. Benov, A. L. Crumbliss and I. Fridovich, *Inorg. Chem.* 1999, **38**, 4011.
5. G. Ferrer-Sueta, I. Batinić-Haberle, I. Spasojević, I. Fridovich and R. Radi, *Chem. Res. Toxicol.* 1999, **12**, 42.
6. R. Kachadourian, I. Batinić-Haberle and I. Fridovich, *Inorg. Chem.* 1999, **38**, 391.
7. J. Lee, J. A. Hunt and J. T. Groves, *J. Am. Chem. Soc.* 1998, **120**, 6053.
8. J. P. Crow, *Arch. Biochem. Biophys.* 1999, **371**, 41.
9. M. Patel and B. J. Day, *Trends Pharmacol.* 1999, **20**, 359.
10. (a) K. Aston, N. Rath, A. Naik, U. Slomczynska, O. F. Schall and D. P. Riley, *Inorg. Chem.* 2001, **40**, 1779. (b) S. Cuzzocrea, E. Mazzon, L. Dugo, A. P. Caputi, K. Aston, D. P. Riley and D. Salvemini, *Br. J. Pharmacol.* 2001, **132**, 19.
11. (a) S. Melov, J. Ravenscroft, S. Malik, M. S. Gill, D. W. Walker, P. E. Clayton, D. C. Wallace, B. Malfroy, S. R. Doctrow and G. J. Lithgow, *Science*, 2000, **289**, 1567. (b) K. Baker, C. Bucay Marcus, K. Huffman, H. Kruk, B. Malfroy and S. R. Doctrow, *J. Pharmacol. Exp. Ther.*, 1998, **284**, 215.

12. I. Batinić-Haberle, L. Benov, I. Spasojević and I. Fridovich, *J. Biol. Chem.*, 1998, **273**, 24521.
13. I. Spasojević, R. Menzeleev, P. S. White and I. Fridovich, *Inorg. Chem.*, 2002, submitted.
14. G. B. Mackensen, M. Patel, H. Sheng, C. C. Calvi, I. Batinić-Haberle, B. J. Day, L. P. Liang, I. Fridovich, J. D. Crapo, R. D. Pearlstein and D. S. Warner, *J. Neurosci.*, 2001, **21**, 4582.
15. J. D. Piganelli, S. C. Flores, C. Cruz, J. Koepp, I. Batinić-Haberle, J. Crapo, B. J. Day, R. Kachadourian, R. Young, B. Bradley and K. Haskins, *Diabetes*, 2002, **51**, 347.
16. M. Aslan, T. M. Ryan, B. Adler, T. M. Townes, D. A. Parks, J. A. Thompson, A. Tousson, M. T. Gladwin, M. M. Tarpey, M., R. P. Patel, I. Batinić-Haberle, C. R. White and B. A. Freeman, *Proc. Natl. Acad. Sci. USA*, 2001, **98**, 15215.
17. (a) I. Batinić-Haberle, I. Spasojević, I. Fridovich, M. S. Anscher and Ž. Vujaskovic, Proc of the 43rd Annual Meeting of American Society for Therapeutics in Radiation Onciology, San Francisco 2001, 235-236. (b) Z. Vujasković, I. Batinić-Haberle, I. Spasojević, T. V. Samulski, M. W. Dewhirst and M. S. Anscher, Annual Meeting of Radiation Research Society, San Juan, Puerto Rico 2001. (c) Ž. Vujasković, I. Batinić-Haberle, I. Spasojević, Irwin Fridovich, M. S. Anscher and M. W. Dewhirst, *Free Rad. Biol. Med.* 2001, S128.
18. W. R. Waud, F. O. Brady, R. D. Wiley and K. V. Rajagopalan, *Arch. Biochem. Biophys.* 1975, **19**, 695.

19. T. Kaufmann, T., B. Shamsai, R. S. Lu, R. Bau and G. M. Miskelly, *Inorg. Chem.* 1995, **34**, 5073.
20. I. M. Kolthof and W. J. Tomsicek, W. J., *J. Phys. Chem.* 1935, **39**, 945.
21. I. Batinić-Haberle, I. Spasojević, R. D. Stevens, P. Hambright, A. N. Thorpe, J. Grodkowski, P. Neta and I. Fridovich, *Inorg. Chem.* 2001, **40**, 726.
22. J. M. McCord and I. Fridovich, *J. Biol. Chem.* 1969, **244**, 6049.
23. I. Batinić-Haberle, S. I. Liochev, I. Spasojević and I. Fridovich, *Arch. Biochem. Biophys.* 1997, **343**, 225.
24. P. Hambright, I. Spasojević, I. Fridovich and I. Batinić-Haberle, in preparation.
25. P. Hambright, *Water-Soluble Metalloporphyrins* in The Porphyrin Handbook, K. M. Kadish, K. M. Smith, R. Guillard, Eds. Academic Press, N. Y. 2000, Chapter 18.
26. T. P. G. Sutter and P. Hambright, *J. Coord. Chem.* 1993, **30**, 317.
27. L. R. Robinson and P. Hambright, *Inorg. Chem.*, 1992, **31**, 652.
28. J. B. Reid and P. Hambright, *Inorg. Chem.* 1977, **16**, 968.
29. M. Inamo, N. Kamiya, Y. Inada, M. Nomura and S. Funahashi, *Inorg. Chem.*, 2001, **40**, 5636.
30. P. B. Chock and P. Hambright, *J. Am. Chem. Soc.* 1974, **96**, 3123.
31. S. Funahashi, Y. Inada and M. Inamo, *Anal. Sci.* 2001, **17**, 917.
32. T. P. G. Sutter, R. Rahimi, P. Hambright, J. Bommer, M. Kumar and P. Neta, *J. Chem. Soc. Faraday Trans.* 1993, **84**, 495.
33. B. Cheng, O. Q. Munro, H. M. Marques and W. R. Scheidt, *J. Am. Chem. Soc.* 1997, **119**, 10732.


34. R. F. Pasternack, N. Sutin and D. H. Turner, *J. Am. Chem. Soc.* 1976, **98**, 1908.
35. P. Hambright, T. Gore and M. Burton, *Inorg. Chem.* 1976, **15**, 2314.
36. J. Davilla, A. Harriman, M.-G. Richoux and L. R. Milgrom, *J. Chem. Soc. Chem. Commun.* 1987, 525.
37. I. Batinić-Haberle, R. D. Stevens and I. Fridovich, *J. Porphyrins Phthalocyanines*, 2000, **4**, 217.
38. R. Kachadourian, N. Srinivasan, C. A. Haney and R. D. Stevens, *J. Porphyrins Phthalocyanines*, 2001, **5**, 507.
39. Vance, C. K. and Miller, A.-F., *Biochemistry*, 2001, **40**, 13079.
40. (a) G. D. Lawrence and D. T. Sawyer, *Biochemistry*, 1979, **18**, 3045. (b) W. C. Jr. Barrette, D. T. Sawyer, J. A. Free and K. Asada, *Biochemistry* 1983, **22**, 624.
41. Wood, P. M., *Biochem. J.*, 1988, **253**, 287.
42. Vance, C. K. and Miller, A. – F., *J. Am. Chem. Soc.*, 1998, **120**, 461.
43. R. M. Ellerby, D. E. Cabelli, J. A. Graden and J. S. Valentine, *J. Am. Chem. Soc.*, 1996, **118**, 6556.
44. D. Klug-Roth, I. Fridovich and J. Rabani, *J. Am. Chem. Soc.*, 1973, **95**, 2786.
45. R. A. Marcus, *Annu. Rev. Phys. Chem.*, 1964, **15**, 155.
46. *CRC Handbook of Chemistry and Physics*, D. R. Lide, Editor-in-Chief, 74th Edition, 1993-1994, CRC Press, Boca Raton.
47. I. Spasojević, I. Batinić-Haberle and I. Fridovich, *Nitric Oxide: Biology and Chemistry* 2000, **4**, 526.
48. (a) C. K. Regan, S. L. Craig and J. I. Brauman, *Science*, 2002, **295**, 2245.
49. R. Shimanovich and J. T. Groves, *Arch. Biochem. Biophys.*, 2001, **387**, 307.

50. N. Jin, J. L. Bourassa, S. C. Tizio and J. T. Groves, *Angew. Chem. Int. Ed.*, 2000, **39**, 3849.
51. H. Zhang, J. Joseph, M. Gurney, D. Becker and B. Kalyanaraman, *J. Biol. Chem.*, 2002, **277**, 1013.
52. N. Motohashi and Y. Saito, *Chem. Pharm. Bull.*, 1995, **43**, 505.
53. C. Quijano, D. Hernandez-Saavedra, L. Castro, J. M. McCord, B. A. Freeman and R. Radi, *J. Biol. Chem.*, 2001, **276**, 11631.
54. (a) S. L. Jewet, A. M. Rocklin, M. Ghanevati, J. M. Abel and J. A. Marach, *Free Rad. Biol. Med.*, 1999, **26**, 905. (b) S. P. A. Goss, R. J. Singh and B. Kalyanaraman, *J. Biol. Chem.*, 1999, **274**, 28233. (c) S. I Liochev and I. Fridovich, *Free Rad. Biol. Med.*, 199, **27**, 1444.
55. (a) A. G. Estevez, J. P. Crow, J. B. Sampson, L. Reither, J. Zhuang, G. J. Richardson, M. M. Tarpey, L. Barbeito and J. S. Beckman, *Science*, 1999, **286**, 2498. (b) S. I. Liochev and I. Fridovich, *J. Biol. Chem.*, 2001, **276**, 35253.
56. S. I. Liochev and I. Fridovich *J. Biol. Chem.*, 2000, **275**, 38482.
57. E. D. Coulter, J. P. Emerson, D. M. Jr., Kurtz and D. E. Cabelli, *J. Am. Chem. Soc.*, 2000, **122**, 11555.
58. B. M. Matata and M. Galinanes, *J. Biol. Chem.*, 2002, **277**, 2330.
59. (a) Ž. Vujašković, I. Batinić-Haberle, M. S. Anscher, Z. N. Rabbani, T. V. Samulski, K. Amin, M. W. Dewhirst and Z. Haroon, Proc. of the 43rd Annual Meeting of American Society for Therapeutics in Radiation Oncology, San Francisco 2001, 88-89. (b) Ž. Vujašković, I. Batinić-Haberle, Z. N. Rabbani, Q.-


F. Feng, S. K. Kang, I. Spasojević, T. V. Samulski, I. Fridovich, M. W. Dewhirst,
M. S. Anscher, *Free Rad. Biol. Med.* In press.

What is claimed is:


1. A compound of formula



I




II



III

or



IV,

wherein

each R is, independently, an alkyl group of greater than 8 carbons,

each A is, independently, hydrogen or an electron withdrawing group,

M is a metal selected from the group consisting of manganese, iron, copper, cobalt, nickel and zinc, and

Z⁻ is a counterion.

2. The compound according to claim 1 wherein at least one A is a halogen.

3. The compound according to claim 1 wherein said compound is of Formula I or II and M is manganese.

4. The compound according to claim 1 wherein said compound is of Formula I or III.

5. The compound according to claim 4 wherein said compound is of Formula I and M is manganese.

6. A method of protecting cells from oxidant-induced toxicity comprising contacting said cells with a protective amount of the compound of claim 1 under conditions such that the protection is effected.

7. The method according to claim 6 wherein said cells are mammalian cells.

8. A method of treating a pathological condition of a patient resulting from oxidant-induced toxicity comprising administering to said patient an effective amount of the compound of claim 1 under conditions such that the treatment is effected.

9. A method of treating a pathological condition of a patient resulting from degradation of NO[•], comprising administering to said patient an effective amount of the compound of claim 1 under conditions such that the treatment is effected.

10. A method of treating a patient for inflammatory lung disease comprising administering to said patient an effective amount of the compound of claim 1 under conditions such that the treatment is effected.

Tables

Table 1. Metal-Centered Redox Potentials $E_{1/2}$, $\log k_{cat}$ for $O_2^{\cdot -}$ Dismutation, and Chromatographic R_f values.

Porphyrin	R_f^a	pK_{a2}^b	$E_{1/2}^c$ mV vs NHE	$\log k_{cat}^d$
Mn ^{III} TM-2-PyP ^{S+}	0.09 (0.13)	10.9	+220	7.79
Mn ^{III} TE-2-PyP ^{S+}	0.13 (0.21)	10.9	+228	7.76
Mn ^{III} TnPr-2-PyP ^{S+}	0.20 (0.31)	11.4	+238	7.38
Mn ^{III} TnBut-2-PyP ^{S+}	0.33 (0.46)	11.7	+254	7.25
Mn ^{III} TnHex-2-PyP ^{S+}	0.57 (0.63)	12.2	+314	7.48
Mn ^{III} TnOct-2-PyP ^{S+}	0.80 (0.86)	13.2	+367	7.71

^a R_f (compound path/solvent path) on silica gel TLC plates in 1:1:8 KNO₃-saturated H₂O:H₂O:acetonitrile. R_f for the metal-free porphyrins are in parentheses. ^b pK_{a2} determined at 25 °C ionic strength 0.10 (NaNO₃/NaOH). ^c $E_{1/2}$ determined in 0.05 M phosphate buffer (pH 7.8, 0.1 M NaCl). ^d k_{cat} determined using the cytochrome c assay, in 0.05 M phosphate buffer, pH 7.8, at (25±1) °C.

Table 2. Molar Absorptivities of Tetrakis (*N*-alkylpyridinium-2-yl)porphyrin chlorides and their Mn(III) Complexes.

Porphyrin	λ_{nm} (log ϵ) ^a
H ₂ TM-2-PyP ⁴⁺	413.2(5.32); 510.4(4.13); 544.4(3.49); 581.4(3.72); 634.6(3.13)
H ₂ TE-2-PyP ⁴⁺	414(5.33); 511(4.20); 545(3.58); 582(3.80); 635(3.38);
H ₂ TnPr-2-PyP ⁴⁺	415(5.38); 511.5(4.24); 545(3.62); 583(3.84); 635(3.37)
H ₂ TnBut-2-PyP ⁴⁺	415(5.37); 511(4.24); 544(3.60); 583(3.84); 636(3.39)
H ₂ TnHex-2-PyP ⁴⁺	415.5(5.34); 510.5(4.24); 543(3.62); 584.5(3.84); 638(3.43)
H ₂ TnOct-2-PyP ⁴⁺	416.5(5.31); 510(4.25); 542(3.59); 585(3.82); 639.5(3.43)
Mn ^{III} TM-2-PyP ⁵⁺	363.5(4.64); 411(4.27); 453.4(5.11); 499(3.66); 556(4.03); 782(3.15)
Mn ^{III} TE-2-PyP ⁵⁺	363.5(4.68); 409(4.32); 454(5.14); 499(3.75); 558(4.08); 782(3.26)
Mn ^{III} TnPr-2-PyP ⁵⁺	363(4.70); 411(4.37); 454(5.21); 498(3.81); 559(4.12); 782(3.35)
Mn ^{III} TnBut-2-PyP ⁵⁺	364(4.70); 410(4.35); 454(5.23); 498(3.83); 559(4.14); 781(3.33)
Mn ^{III} TnHex-2-PyP ⁵⁺	364.5(4.70); 415(4.57); 454.5(5.21); 507(3.85); 560(4.12); 780(3.30)
Mn ^{III} TnOct-2-PyP ⁵⁺	364(4.72); 414(4.44); 454.5(5.24); 500.5(3.84); 559.5(4.14); 781(3.25)

^aThe molar absorptivities were determined in water at room temperature.

Table 3. Electrospray Mass Spectrometry Results for $H_2T(\text{alkyl})-2\text{-PyP}^{4+}$ Compounds. ^a

Species ^b	m/z					
	M	E	nPr	nBu	nHex	nOct
$H_2P^{4+}/4$	169	184	198	212	239	268
$H_2P^{4+}+AN/4$	180	194	208	222	250	
$H_2P^{4+}+2AN/4$	190					
$H_2P^{4+}-H^+/3$	226	245	263	282	319	357
$H_2P^{4+}-H^++AN/3$	240	258	278			
$H_2P^{4+}-H^++H_2O/3$				288		
$H_2P^{4+}-H^++Cl^-/2$					496	
$H_2P^{4+}-a^+/3$		235	249	263	291	319
$H_2P^{4+}-a^+-H^+/2$		352	374	394	436	
$H_2P^{4+}-a^++H_2O/3$			255			
$H_2P^{4+}-2a^+/2$			352	366		
$H_2P^{4+}+H^+/5$	136					
$H_2P^{4+}+H^++AN/5$	143					
$H_2P^{4+}+H^++2AN/5$	152					
$H_2P^{4+}+H^++2Cl^-/3$					343	381
$H_2P^{4+}+2H^++2Cl^-/4$						286
$H_2P^{4+}-2H^+/2$	339	367	395	423	479	



^a~0.5 mM solutions of H_2P^{4+} in 1:1 acetonitrile:water, 20 V cone voltage. ^bAN denotes acetonitrile and a is an alkyl group.

Table 4. Electrospray Mass Spectrometry for Mn^{III}T(alkyl)-2-PyP⁵⁺ Porphyrins.^a

Species ^b	m/z					
	M	E	nPr	nBu	nHex	nOct
Mn ^{III} P ⁵⁺ /5	146	157				
Mn ^{III} P ⁵⁺ +AN/4	155	166	177	188		
Mn ^{III} P ⁵⁺ +2AN/5	163	174	185	196		
Mn ^{III} P ⁵⁺ +3AN/5	171	182	193	205		
Mn ^{III} P ⁵⁺ +4AN/5	179	190		213		
Mn ^{III} P ⁵⁺ +5AN/5	187	198				
Mn ^{III} P ⁵⁺ +6AN/5	195					
Mn ^{III} P ⁵⁺ +H ₂ O/5	150					
Mn ^{III} P ⁵⁺ +Cl ⁻ /4	192	206		234	262	290
Mn ^{III} P ⁵⁺ +2Cl ⁻ /3	267	286	305	323	361	398
Mn ^{III} P ⁵⁺ +Cl ⁻ +AN/4	202	216	230	244	272	
Mn ^{III} P ⁵⁺ -a/4			200			
Mn ^{III} P ⁵⁺ -a+AN/4		200		221	242	
Mn ^{III} P ⁵⁺ -a+Cl ⁻ /3		264	279	293	321	349
Mn ^{III} P ⁵⁺ -2a/3		243	252	262		299
Mn ^{III} P ⁵⁺ -2a+AN/3				275	294	
Mn ^{II} P ⁴⁺ /4	183	197	211			281
Mn ^{II} P ⁴⁺ +AN/4	193	207	221	235	263	
Mn ^{II} P ⁴⁺ +2AN/4	204					
Mn ^{II} P ⁴⁺ +Cl ⁻ /3	255	274	293	312	349	387
Mn ^{II} P ⁴⁺ -a/3		253	266	281	309	337
Mn ^{III} P ⁵⁺ -Mn ³⁺ +H ⁺ /3				281	319	357
M ^{II} P ³⁺ /3 or Mn ^I P ³⁺ /3				294	337	375

^a-0.5 mM solutions of Mn^{III}P⁵⁺ in 1:1 acetonitrile:water, 20 V cone voltage. ^bAN denotes acetonitrile and a is an alkyl group.

1/9

*Scheme I*

2/9

Figure 1A





Figure 1B



3/9

Figure 2A





Figure 2B



4/9

Figure 3A






Figure 3B




5/9

Figure 4




6/9

Figure S1


7/9

Figure S2




8/9

Figure S3



9/9

As the alkyl chains of Mn(III) porphyrins lengthen, the favorable increase in $E_{1/2}$ overcomes the unfavorable steric/electrostatic effects such that the n-octyl is as potent a catalyst of $O_2^{\cdot -}$ dismutation as are the methyl and ethyl compounds.



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/18099

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/555, 31/409; C07D 487/22

US CL : 514/185, 410; 540/145

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/185, 410; 540/145

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	US 6,566,517 B2 (MIURA et al.) 20 May 2003 (20.05.2003), columns 4-5.	1-10
A,P	US 6,548,045 B2 (SAKATA et al) 15 April 2003 (15.04.2002), columns 4-6.	1-10
A,E	US 6,602,998 B2 (KOBUEKE et al.) 05 August 2003 (05.08.2003), columns 3-6.	1-10
A,E	US 6,624,187 B1 (PANDEY et al) 23 September 2003 (23.09.2003), columns 3-4.	1-10

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"B" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

24 September 2003 (24.09.2003)

Date of mailing of the international search report

03 NOV 2003

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US

Commissioner for Patents

P.O. Box 1450

Alexandria, Virginia 22313-1450

Facsimile No. (703)305-3230

Authorized officer

Deborah C Lambkin

Telephone No. 703-308-1235