




(86) Date de dépôt PCT/PCT Filing Date: 2000/10/12
(87) Date publication PCT/PCT Publication Date: 2001/04/19
(45) Date de délivrance/Issue Date: 2014/02/11
(85) Entrée phase nationale/National Entry: 2002/04/11
(86) N° demande PCT/PCT Application No.: IB 2000/001538
(87) N° publication PCT/PCT Publication No.: 2001/027268
(30) Priorités/Priorities: 1999/10/12 (US60/158,774);
2000/02/14 (US09/503,954)

(51) Cl.Int./Int.Cl. *C12N 15/12* (2006.01),
A61K 38/17 (2006.01), *C07K 14/47* (2006.01),
C12N 15/62 (2006.01), *A61K 38/00* (2006.01)
(72) Inventeur/Inventor:
BONNY, CHRISTOPHE, CH
(73) Propriétaire/Owner:
XIGEN INFLAMMATION LTD., CY
(74) Agent: SMART & BIGGAR

(54) Titre : INHIBITEURS PEPTIDIQUES, A PERMEABILITE CELLULAIRE, DU MECANISME DE TRANSDUCTION DU
SIGNAL DE JNK
(54) Title: CELL-PERMEABLE PEPTIDE INHIBITORS OF THE JNK SIGNAL TRANSDUCTION PATHWAY

Effects of TAT-IB Peptides on JNK Mediated Phosphorylation



(57) Abrégé/Abstract:

The invention provides cell-permeable peptides that bind to JNK proteins and inhibit JNK-mediated effects in JNK-expressing cells.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 April 2001 (19.04.2001)


PCT

(10) International Publication Number
WO 01/27268 A3

- (51) International Patent Classification⁷: C12N 15/12, 15/62, C07K 14/47, A61K 38/17
- (71) Applicant (for all designated States except US): UNIVERSITY OF LAUSANNE [CH/CH]; Rue de Bugnon 21, CH-1005 Lausanne (CH).
- (21) International Application Number: PCT/IB00/01538
- (22) International Filing Date: 12 October 2000 (12.10.2000)
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): BONNY, Christophe [CH/CH]; 5, rue du Docteur Yersin, CH-1110 Morges (CH).
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
- | | | |
|------------|-------------------------------|----|
| 60/158,774 | 12 October 1999 (12.10.1999) | US |
| 09/503,954 | 14 February 2000 (14.02.2000) | US |
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:
- | | |
|----------|-------------------------------|
| US | 60/158,774 (CIP) |
| Filed on | 12 October 1999 (12.10.1999) |
| US | 09/503,954 (CIP) |
| Filed on | 14 February 2000 (14.02.2000) |
- (74) Agent: SCHLICH, George, William; Mathys & Squire, 100 Gray's Inn Road, London WC1X 8AL (GB).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

[Continued on next page]

(54) Title: CELL-PERMEABLE PEPTIDE INHIBITORS OF THE JNK SIGNAL TRANSDUCTION PATHWAY



(57) Abstract: The invention provides cell-permeable peptides that bind to JNK proteins and inhibit JNK-mediated effects in JNK-expressing cells.



WO 01/27268 A3

50115-3

CELL-PERMEABLE PEPTIDE INHIBITORS OF THE JNK SIGNAL TRANSDUCTION PATHWAY

FIELD OF THE INVENTION

This invention relates generally to protein kinase inhibitors and more specifically to
5 inhibitors of the protein kinase c-Jun amino terminal kinase.

BACKGROUND OF THE INVENTION

The c-Jun amino terminal kinase (JNK) is a member of the stress-activated group of
mitogen-activated protein (MAP) kinases. These kinases have been implicated in the control of
cell growth and differentiation, and, more generally, in the response of cells to environmental
10 stimuli. The JNK signal transduction pathway is activated in response to environmental stress
and by the engagement of several classes of cell surface receptors. These receptors can include
cytokine receptors, serpentine receptors, and receptor tyrosine kinases. In mammalian cells, JNK
has been implicated in such biological processes as oncogenic transformation and in mediating
adaptive responses to environmental stress. JNK has also been associated with modulating
15 immune responses, including maturation and differentiation of immune cells, as well effecting
programmed cell death in cells identified for destruction by the immune system.

SUMMARY OF THE INVENTION

The present invention is based in part on the discovery of peptides that are effective
20 inhibitors of JNK proteins. The peptides, referred to herein as JNK peptide inhibitors, decrease
the downstream cell-proliferative effects of c-Jun amino terminal kinase (JNK).

Accordingly, the invention includes novel JNK inhibitor peptides, as well as chimeric
peptides which include a JNK peptide inhibitor linked a trafficking peptide that can be used to
direct a peptide on which it is present to a desired cellular location. The trafficking sequence
25 can be used to direct transport of the peptide across the plasma membrane. Alternatively, or in
addition, the trafficking peptide can be used to direct the peptide to desired intracellular location,
such as the nucleus.

53493-1

In one aspect, the invention relates a chimeric peptide, comprising a first domain and a second domain linked by a covalent bond, wherein the first domain comprises a trafficking sequence directing the chimeric peptide across the plasma membrane and/or to a desired cellular location, and the second domain comprises a retro-inverso isomer of a JNK inhibitor sequence, wherein said chimeric peptide is composed of D enantiomeric amino acids in the JNK inhibitor peptide of the second domain, and wherein the JNK inhibitor peptide is less than 280 amino acids in length, wherein the JNK inhibitor sequence comprises the amino acid sequence of SEQ ID NO: 3 or an amino acid sequence having a sequence identity of at least about 80% with the full length sequence of SEQ ID NO: 3.

In another aspect, the invention relates to a composition, comprising the peptide as described herein, and a carrier.

In another aspect, the invention relates to use of the peptide as described herein, in the manufacture of a medicament for treating or preventing a pathophysiology associated with activation of JNK in a subject, wherein the pathophysiology is selected from: (1) a malignancy selected from leukemia, and adenocarcinoma; (2) a non-malignant or immunologically-related cell proliferative disease selected from psoriasis, restenosis, vasculitis, pemphigus vulgaris, Behcet's syndrome, rheumatoid arthritis, acute respiratory distress syndrome (ARDS), ischemic heart disease, post-dialysis syndrome, acquired immunodeficiency syndrome, lipid histiocytosis, septic shock and apoptosis; (3) a pathological state induced by ionizing radiation, free radicals, DNA damaging agents or UV light; and (4) diabetes.

53493-1

In another aspect, the invention relates to use of the peptide as described herein, for treating or preventing a pathophysiology associated with activation of JNK in a subject, wherein the pathophysiology is selected from: (1) a malignancy
5 selected from leukemia, and adenocarcinoma; (2) a non-malignant or immunologically-related cell proliferative disease selected from psoriasis, restenosis, vasculitis, pemphigus vulgaris, Behcet's syndrome, rheumatoid arthritis, acute respiratory distress syndrome (ARDS), ischemic heart disease, post-dialysis
10 syndrome, acquired immunodeficiency syndrome, lipid histiocytosis, septic shock and apoptosis; (3) a pathological state induced by ionizing radiation, free radicals, DNA damaging agents or UV light; and (4) diabetes.

In another aspect, the invention relates to the
15 composition as described herein for use in treating or preventing a pathophysiology associated with activation of JNK in a subject, wherein the pathophysiology is selected from: (1) a malignancy selected from leukemia, and adenocarcinoma; (2) a non-malignant or immunologically-related cell proliferative
20 disease selected from psoriasis, restenosis, vasculitis, pemphigus vulgaris, Behcet's syndrome, rheumatoid arthritis, acute respiratory distress syndrome (ARDS), ischemic heart disease, post-dialysis syndrome, acquired immunodeficiency syndrome, lipid histiocytosis, septic shock and apoptosis; (3)
25 a pathological state induced by ionizing radiation, free radicals, DNA damaging agents or UV light; and (4) diabetes.

The JNK inhibitor peptides can be present as polymers of L-amino acids. Alternatively,

50115-3

the peptides can be present as polymers of D-amino acids.

Also included in the invention are pharmaceutical compositions that include the JNK-binding peptides, as well as antibodies that specifically recognize the JNK-binding peptides.

The invention also includes a method of inhibiting expression of a JNK kinase in a cell.

5 In another aspect, the invention includes a method of treating a pathophysiology associated with activation of JNK in a cell or cells. For example, the target cells may be, *e.g.*, cultured animal cells, human cells or micro-organisms. Delivery can be carried out *in vivo* by administering the chimeric peptide to an individual in whom it is to be used for diagnostic, preventative or therapeutic purposes. The target cells may be *in vivo* cells, *i.e.*, cells composing
10 the organs or tissues of living animals or humans, or microorganisms found in living animals or humans.

Among the advantages provided by the invention is that the JNK inhibitor peptides are small, and can be produced readily in bulk quantities and in high purity. The inhibitor peptides are also resistant to intracellular degradation, and are weakly immunogenic. Accordingly, the
15 peptides are well suited for *in vitro* and *in vivo* applications in which inhibition of JNK-expression is desired.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be
20 used in the practice or testing of the present invention, suitable methods and materials are described below. In the case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

25 Other features and advantages of the invention will be apparent from the following detailed description and claims.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A-C are diagrams showing alignments of conserved JBD domain regions in the

indicated transcription factors.

FIG. 2 is a diagram showing alignments of generic TAT-IB fusion peptides.

FIG. 3 is a histogram depicting inhibition of β -cell death by the minimal 23 amino acid long JBD domain of IB1 compared to the full 280 amino acid JBD domain.

5 FIG. 4 is an illustration demonstrating the effects of TAT, TAT-IB1 and TAT-IB2 peptides on phosphorylation of recombinant JNKs. Panel A shows inhibition of c-Jun, ATF2 and Elk1 phosphorylation by recombinant JNKs *in vitro*. Panel B shows dose response experiments similar to Panel A.

10 FIG. 5 is a histogram depicting L-TAT-IB inhibition of phosphorylation by recombinant JNKs. Panel A shows L-TAT-IB inhibition of c-Jun, ATF2 and Elk1 phosphorylation by recombinant JNKs *in vitro* in the presence of MKK4. Panel B shows similar dose response experiments with MKK7.

FIG. 6 is an illustration demonstrating the inhibition of c-Jun phosphorylation by activated JNKs.

15 FIG. 7 is a histogram depicting short term inhibition of IL-1 β induced pancreatic β -cell death by the L-TAT-IB peptides.

FIG. 8 is a histogram depicting short term inhibition of IL-1 β induced pancreatic β -cell death by the D-TAT-IB peptides.

20 FIG. 9 is a histogram depicting long term inhibition of IL-1 β induced pancreatic β -cell death by L-TAT-IB1 and D-TAT-IB1 peptides.

FIG. 10 is a histogram depicting inhibition of irradiation induced human colon cancer WiDr cell death by L-TAT-IB1 and D-TAT-IB1 peptides.

FIG. 11 is an illustration of the modulation of JNK kinase activity by L-TAT, TAT-IB1 and D-TAT-IB1 peptides.

25 FIG. 12 are graphs depicting the protective effects of the TAT-IB1 peptides in mice. Panel A shows the effect of irradiation on weight. Panel B shows the effect of irradiation on oedemus and erythemus status.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is based in part on the discovery of cell permeable peptides that inhibit the activated c-Jun amino terminal kinase (JNK) signaling pathway. These peptides are referred to herein as JNK inhibitor peptides. Additionally, the discovery provides methods and pharmaceutical compositions for treating pathophysiologies associated with JNK signaling.

JNK inhibitor peptides were identified by inspecting sequence alignments between kJNK Binding Domains in various insulin binding (IB) proteins. The results of this alignment are shown in FIGS. 1A-1C. FIG. 1A depicts the region of highest homology between the JBDs of IB1, IB2, c-Jun and ATF2. Panel B depicts the amino acid sequence alignment of the JBDs of IB1 and IB2. Fully conserved residues are indicated by asterisks, while residues changed to Ala in the GFP-JBD_{23Mut} vector are indicated by open circles. FIG. 1C shows the amino acid sequences of chimeric proteins that include a JNK inhibitor peptide domain and a trafficking domain. In the example shown, the trafficking domain is derived from the human immunodeficiency virus (HIV) TAT polypeptide, and the JNK inhibitor peptide is derived from an IB1 polypeptide. Human, mouse, and rat sequences are identical in Panels B and C.

Sequence comparison between the JNK binding domains of IB1 [SEQ ID NO: 17], IB2 [SEQ ID NO: 18], c-Jun [SEQ ID NO: 19] and ATF2 [SEQ ID NO: 20] revealed a partially conserved 8 amino acid sequence (FIG. 1A). A comparison of the JBDs of IB1 and IB2 further revealed two blocks of seven and three amino acids that are highly conserved between the two sequences. These two blocks are contained within a peptide sequence of 23 amino acids in IB1 [SEQ ID NO: 1] and 21 amino acids in IB2 [SEQ ID NO: 2].

The JNK inhibitor peptides of the invention can be used in any situation in which inhibition of JNK activity is desired. This can include *in vitro* applications, *ex vivo*, and *in vivo* applications. As JNKs and all its isoforms participate in the development and establishment of pathological states or in pathways, the JNK peptides can be used to prevent or inhibit the occurrence of such pathological states. This includes prevention and treatment of diseases and prevention and treatment of conditions secondary to therapeutic actions. For example, the peptides of the present invention can be used to treat or prevent, *e.g.*, diabetes, ionizing radiation, immune responses (including autoimmune diseases), ischemia/reperfusion injuries, heart and cardiovascular hypertrophies, and some cancers (*e.g.*, *Bcr-Abl* transformation).

The peptides can also be used to inhibit expression of genes whose expression increases in the presence of an active JNK polypeptide. These genes and gene products includes, *e.g.*, proinflammatory cytokines. Such cytokines are found in all forms of inflammatory, auto-inflammatory, immune and autoimmune diseases, degenerative diseases, myopathies, cardiomyopathies, and graft rejection.

The JNK inhibitor peptides described herein can also be used to treat or prevent effects associated with cellular shear stress, such as in pathological states induced by arterial hypertension, including cardiac hypertrophy and arteriosclerotic lesions, and at bifurcations of blood vessels, and the like; ionizing radiation, as used in radiotherapy and UV lights; free radicals; DNA damaging agents, including chemotherapeutic drugs; oncogenic transformation; ischemia and reperfusion; hypoxia; and hypo- and hyperthermia.

The polynucleotides provided by the present invention can be used to express recombinant peptides for analysis, characterization or therapeutic use; as markers for tissues in which the corresponding peptides is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in disease states). Other uses for the nucleic acids include, *e.g.*, molecular weight markers in gel electrophoresis-based analysis of nucleic acids.

The JNK inhibitor peptides disclosed herein are presented in Table 1. The table presents the name of the JNK inhibitor peptide, as well as its sequence identifier number, length, and amino acid sequence.

TABLE 1

PEPTIDE NAME	SEQ ID	AA	Sequence
L-IB1	1	23	DTYRPKRPTT LNLFPQVPRS QDT
L-IB2	2	21	EEPHKHRPTT LRLTTLGAQD S
D-IB1	3	23	TDQSRPVQPF LNLTPRKPR YTD
D-IB2	4	21	SDQAGLTTLR LTPRHKHPE E
L-IB (generic)	5	19	XRPTTLXLXX XXXXXQDS/TX
D-IB (generic)	6	19	XS/TDQXXXXXX XLXLTPRX
L-TAT	7	10	GRKKRRQRRR
D-TAT	8	10	RRRQRRKKRG
L-generic-TAT	9	17	XXXXRKKRRQ RRRXXXX
D-generic-TAT	10	17	XXXXRRRQRR KKRXXXX
L-TAT-IB1	11	35	GRKKRRQRRR PPDTYRPKRP TTLNLFPQVP RSQDT
L-TAT-IB2	12	33	GRKKRRQRRR PPEPHKHRP TTLRLTTLGA QDS

WO 01/27268

PCT/IB00/01538

L-TAT-IB (generic)	13	42	XXXXXXXXRKK RRQRRRXXXX XXXXRPTTLX LXXXXXXXXXD S/TX
D-TAT-IB1	14	35	TDQSRPVQPF LNLTPRKPR YDPPRRRQR RKKRG
D-TAT-IB2	15	33	SDQAGLTTLR LTPRHKHPE EPRRRQRK KRG
D-TAT-IB (generic)	16	42	XT/SDQXXXXXX XLXLTPRXX XXXXXRRRQ RRKKRXXXXX XX
IB1-long	17	29	PGTGCGDTYR PKRPTTLNLF PQVPRSQDT
IB2-long	18	27	IPSPSVEEPH KHRPTTLRLT TLGAQDS
c-Jun	19	29	GAYGYSNPKE LKQSMNLNLA DPVGNLKP
ATF2	20	29	TNEDHLAVHK HKHEMTLKFG PARNSVIV

JNK INHIBITOR PEPTIDES

In one aspect, the invention provides a JNK inhibitor peptide. No particular length is implied by the term "peptide." In some embodiments, the JNK-inhibitor peptide is less than 280 amino acids in length, *e.g.*, less than or equal to 150, 100, 75, 50, 35, or 25 amino acids in length. In various embodiment, the JNK-binding inhibitor peptide includes the amino acid sequence of one or more of SEQ ID NOs: 1-6. In one embodiment, the JNK inhibitor peptide peptides bind JNK. In another embodiment the peptide inhibits the activation of at least one JNK activated transcription factor, *e.g.* c-Jun, ATF2 or Elk1.

Examples of JNK inhibitor peptides include a peptide which includes (in whole or in part) the sequence NH_2 -DTYRPKRPTTLNLFQVPRSQDT-COOH [SEQ ID NO:1]. In another embodiment, the peptide includes the sequence NH_2 -EEPHKHRPTTLRLTTLGAQDS-COOH [SEQ ID NO:2]

The JNK inhibitor peptides can be polymers of L-amino acids, D-amino acids, or a combination of both. For example, in various embodiments, the peptides are D retro-inverso peptides. The term "retro-inverso isomer" refers to an isomer of a linear peptide in which the direction of the sequence is reversed and the chirality of each amino acid residue is inverted. *See, e.g.*, Jameson *et al.*, *Nature*, 368, 744-746 (1994); Brady *et al.*, *Nature*, 368, 692-693 (1994). The net result of combining D-enantiomers and reverse synthesis is that the positions of carbonyl and amino groups in each amide bond are exchanged, while the position of the side-chain groups at each alpha carbon is preserved. Unless specifically stated otherwise, it is presumed that any given L-amino acid sequence of the invention may be made into an D retro-inverso peptide by synthesizing a reverse of the sequence for the corresponding native L-amino acid sequence.

In one embodiment, a D retro-inverso peptide has the sequence NH_2 -TDQSRPVQPFLNLTPRKPRYTD-COOH [SEQ ID NO:3]. In another embodiment the D

retro-inverso peptide has the sequence $\text{NH}_2\text{-SDQAGLTTLRLTTPRHKHPEE-COOH}$ [SEQ ID NO: 4]. It has been unexpectedly found that D-retro-inverso TAT-IB peptides have a variety of useful properties. For example, D-TAT and D-TAT-IB peptides enter cells as efficiently as L-TAT and L-TAT-IB peptides, and D-TAT and D-TAT-IB peptides are more stable than the
 5 corresponding L-peptides. Further, while D-TAT-IB1 are $\sim 10\text{-}20$ fold less efficient in inhibiting JNK than L-TAT-IB, they are ~ 50 fold more stable *in vivo*. Finally, as is discussed further below, D-TAT-IB peptides protect interleukin-1 treated and ionizing irradiated cells from apoptosis.

In another embodiment, a JNK inhibitor peptide according to the invention includes the
 10 amino acid sequence $\text{NH}_2\text{-X}_n\text{-RPTTLXLXXXXXXXXXQDS/T -X}_n\text{-COOH}$ [SEQ ID NO: 5, and residues 17-42 of L-TAT-IB, SEQ ID NO: 13, as shown in FIG.2]. As used herein, X_n may be zero residues in length, or may be a contiguous stretch of peptide residues derived from SEQ ID NOS:1-2, preferably a stretch of between 1 and 7 amino acids in length, or may be 10, 20, 30 or more amino acids in length. The single residue represented by S/T may be either Ser or Thr in
 15 the generic sequence. In a further embodiment, the generic IB peptide may be a D retro-inverso peptide having the sequence $\text{NH}_2\text{-X}_n\text{-S/TDQXXXXXXXXLXLTPR-X}_n\text{-COOH}$ [SEQ ID NO: 6, and residues 17-42 of L-TAT-IB, SEQ ID NO:16, as shown in FIG.2].

JNK-inhibitor peptides may be obtained or produced by methods well-known in the art, *e.g.* chemical synthesis, genetic engineering methods as discussed below. For example, a peptide
 20 corresponding to a portion of a JNK inhibitor peptide including a desired region or domain, or that mediates the desired activity *in vitro*, may be synthesized by use of a peptide synthesizer.

A candidate JNK inhibitor peptide may also be analyzed by hydrophilicity analysis (*see, e.g.,* Hopp and Woods, 1981. *Proc Natl Acad Sci USA* 78: 3824-3828) that can be utilized to identify the hydrophobic and hydrophilic regions of the peptides, thus aiding in the design of
 25 substrates for experimental manipulation, such as in binding experiments, antibody synthesis. Secondary structural analysis may also be performed to identify regions of a JNK inhibitor peptide that assume specific structural motifs. *See e.g.,* Chou and Fasman, 1974. *Biochem* 13: 222-223. Manipulation, translation, secondary structure prediction, hydrophilicity and hydrophobicity profiles, open reading frame prediction and plotting, and determination of
 30 sequence homologies can be accomplished using computer software programs available in the

50115-3

art. Other methods of structural analysis including, *e.g.*, X-ray crystallography (*see, e.g.*, Engstrom, 1974. *Biochem Exp Biol* 11: 7-13); mass spectroscopy and gas chromatography (*see, e.g.*, METHODS IN PROTEIN SCIENCE, 1997. J. Wiley and Sons, New York, NY) and computer modeling (*see, e.g.*, Fletterick and Zoller, eds., 1986. Computer Graphics and Molecular Modeling, In: CURRENT COMMUNICATIONS IN MOLECULAR BIOLOGY, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY) may also be employed.

The present invention additionally relates to nucleic acids that encode JNK-binding peptides having L-form amino acids, *e.g.*, those L-peptides indicated in Table 1, as well as the complements of these sequences. Suitable sources of nucleic acids encoding JNK inhibitor peptides include the human IB1 nucleic acid (and the encoded protein sequences) available as GenBank Accession Nos. AF074091 and AAD20443, respectively. Other sources include rat IB1 nucleic acid and protein sequences are shown in GenBank Accession No. AF108959 and AAD22543, respectively. Human IB2 nucleic acid and protein sequences are shown in GenBank Accession No AF218778.

Nucleic acids encoding the JNK inhibitor peptides may be obtained by any method known in the art (*e.g.*, by PCR amplification using synthetic primers hybridizable to the 3'- and 5'-termini of the sequence and/or by cloning from a cDNA or genomic library using an oligonucleotide sequence specific for the given gene sequence).

For recombinant expression of one or more JNK inhibitor peptides, the nucleic acid containing all or a portion of the nucleotide sequence encoding the peptide may be inserted into an appropriate expression vector (*i.e.*, a vector that contains the necessary elements for the transcription and translation of the inserted peptide coding sequence). In some embodiments, the regulatory elements are heterologous (*i.e.*, not the native gene promoter). Alternately, the necessary transcriptional and translational signals may also be supplied by the native promoter for the genes and/or their flanking regions.

A variety of host-vector systems may be utilized to express the peptide coding sequence(s). These include, but are not limited to: (i) mammalian cell systems that are infected with vaccinia virus, adenovirus, and the like; (ii) insect cell systems infected with baculovirus and the like; (iii) yeast containing yeast vectors or (iv) bacteria transformed with bacteriophage,

WO 01/27268

PCT/IB00/01538

DNA, plasmid DNA, or cosmid DNA. Depending upon the host-vector system utilized, any one of a number of suitable transcription and translation elements may be used.

Promoter/enhancer sequences within expression vectors may utilize plant, animal, insect, or fungus regulatory sequences, as provided in the invention. For example, promoter/enhancer elements can be used from yeast and other fungi (*e.g.*, the GAL4 promoter, the alcohol dehydrogenase promoter, the phosphoglycerol kinase promoter, the alkaline phosphatase promoter). Alternatively, or in addition, they may include animal transcriptional control regions, *e.g.*, (i) the insulin gene control region active within pancreatic β -cells (*see, e.g.*, Hanahan, *et al.*, 1985. *Nature* 315: 115-122); (ii) the immunoglobulin gene control region active within lymphoid cells (*see, e.g.*, Grosschedl, *et al.*, 1984. *Cell* 38: 647-658); (iii) the albumin gene control region active within liver (*see, e.g.*, Pinckert, *et al.*, 1987. *Genes and Dev* 1: 268-276; (iv) the myelin basic protein gene control region active within brain oligodendrocyte cells (*see, e.g.*, Readhead, *et al.*, 1987. *Cell* 48: 703-712); and (v) the gonadotropin-releasing hormone gene control region active within the hypothalamus (*see, e.g.*, Mason, *et al.*, 1986. *Science* 234: 1372-1378), and the like.

Expression vectors or their derivatives include, *e.g.* human or animal viruses (*e.g.*, vaccinia virus or adenovirus); insect viruses (*e.g.*, baculovirus); yeast vectors; bacteriophage vectors (*e.g.*, lambda phage); plasmid vectors and cosmid vectors.

A host cell strain may be selected that modulates the expression of inserted sequences of interest, or modifies or processes expressed peptides encoded by the sequences in the specific manner desired. In addition, expression from certain promoters may be enhanced in the presence of certain inducers in a selected host strain; thus facilitating control of the expression of a genetically-engineered peptides. Moreover, different host cells possess characteristic and specific mechanisms for the translational and post-translational processing and modification (*e.g.*, glycosylation, phosphorylation, and the like) of expressed peptides. Appropriate cell lines or host systems may thus be chosen to ensure the desired modification and processing of the foreign peptide is achieved. For example, peptide expression within a bacterial system can be used to produce an unglycosylated core peptide; whereas expression within mammalian cells ensures "native" glycosylation of a heterologous peptide.

Also included in the invention are derivatives, fragments, homologs, analogs and variants

WO 01/27268

PCT/IB00/01538

of JNK inhibitor peptides and nucleic acids encoding these peptides. For nucleic acids, derivatives, fragments, and analogs provided herein are defined as sequences of at least 6 (contiguous) nucleic acids, and which have a length sufficient to allow for specific hybridization. For amino acids, derivatives, fragments, and analogs provided herein are defined as sequences of
5 at least 4 (contiguous) amino acids, a length sufficient to allow for specific recognition of an epitope.

The length of the fragments are less than the length of the corresponding full-length nucleic acid or polypeptide from which the JNK inhibitor peptide, or nucleic acid encoding same, is derived. Derivatives and analogs may be full length or other than full length, if the
10 derivative or analog contains a modified nucleic acid or amino acid. Derivatives or analogs of the JNK inhibitor peptides include, *e.g.*, molecules including regions that are substantially homologous to the peptides, in various embodiments, by at least about 30%, 50%, 70%, 80%, or 95%, 98%, or even 99%, identity over an amino acid sequence of identical size or when compared to an aligned sequence in which the alignment is done by a computer homology
15 program known in the art. For example sequence identity can be measured using sequence analysis software (Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, Wis. 53705), with the default parameters therein.

In the case of polypeptide sequences, which are less than 100% identical to a reference
20 sequence, the non-identical positions are preferably, but not necessarily, conservative substitutions for the reference sequence. Conservative substitutions typically include substitutions within the following groups: glycine and alanine; valine, isoleucine, and leucine; aspartic acid and glutamic acid; asparagine and glutamine; serine and threonine; lysine and arginine; and phenylalanine and tyrosine. Thus, included in the invention are peptides having
25 mutated sequences such that they remain homologous, *e.g.* in sequence, in function, and in antigenic character or other function, with a protein having the corresponding parent sequence. Such mutations can, for example, be mutations involving conservative amino acid changes, *e.g.*, changes between amino acids of broadly similar molecular properties. For example, interchanges within the aliphatic group alanine, valine, leucine and isoleucine can be considered
30 as conservative. Sometimes substitution of glycine for one of these can also be considered conservative. Other conservative interchanges include those within the aliphatic group aspartate

and glutamate; within the amide group asparagine and glutamine; within the hydroxyl group serine and threonine; within the aromatic group phenylalanine, tyrosine and tryptophan; within the basic group lysine, arginine and histidine; and within the sulfur-containing group methionine and cysteine. Sometimes substitution within the group methionine and leucine can also be
5 considered conservative. Preferred conservative substitution groups are aspartate-glutamate; asparagine-glutamine; valine-leucine-isoleucine; alanine-valine; phenylalanine- tyrosine; and lysine-arginine.

Where a particular polypeptide is said to have a specific percent identity to a reference polypeptide of a defined length, the percent identity is relative to the reference peptide. Thus, a
10 peptide that is 50% identical to a reference polypeptide that is 100 amino acids long can be a 50 amino acid polypeptide that is completely identical to a 50 amino acid long portion of the reference polypeptide. It might also be a 100 amino acid long polypeptide, which is 50% identical to the reference polypeptide over its entire length. Of course, other polypeptides will meet the same criteria.

15 The invention also encompasses allelic variants of the disclosed polynucleotides or peptides; that is, naturally-occurring alternative forms of the isolated polynucleotide that also encode peptides that are identical, homologous or related to that encoded by the polynucleotides. Alternatively, non-naturally occurring variants may be produced by mutagenesis techniques or by direct synthesis.

20 Species homologs of the disclosed polynucleotides and peptides are also provided by the present invention. "Variant" refers to a polynucleotide or polypeptide differing from the polynucleotide or polypeptide of the present invention, but retaining essential properties thereof. Generally, variants are overall closely similar, and in many regions, identical to the polynucleotide or polypeptide of the present invention. The variants may contain alterations in
25 the coding regions, non-coding regions, or both.

In some embodiments, altered sequences include insertions such that the overall amino acid sequence is lengthened while the protein retains trafficking properties. Additionally, altered sequences may include random or designed internal deletions that shorten the overall amino acid sequence while the protein retains transport properties.

30 The altered sequences can additionally or alternatively be encoded by polynucleotides

50115-3

that hybridize under stringent conditions with the appropriate strand of the naturally-occurring polynucleotide encoding a polypeptide or peptide from which the JNK inhibitor peptide is derived. The variant peptide can be tested for JNK-binding and modulation of JNK-mediated activity using the herein described assays. 'Stringent conditions' are sequence dependent and will
 5 be different in different circumstances. Generally, stringent conditions can be selected to be about 5°C lower than the thermal melting point (T_M) for the specific sequence at a defined ionic strength and pH. The T_M is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically, stringent conditions will be those in which the salt concentration is at least about 0.02 molar at pH 7 and the
 10 temperature is at least about 60°C. As other factors may affect the stringency of hybridization (including, among others, base composition and size of the complementary strands), the presence of organic solvents and the extent of base mismatching, the combination of parameters is more important than the absolute measure of any one.

High stringency can include, *e.g.*, Step 1: Filters containing DNA are pretreated for 8
 15 hours to overnight at 65°C in buffer composed of 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll*, 0.02% BSA, and 500 µg/ml denatured salmon sperm DNA. Step 2: Filters are hybridized for 48 hours at 65°C in the above prehybridization mixture to which is added 100 mg/ml denatured salmon sperm DNA and 5-20 x 10⁶ cpm of ³²P-labeled probe. Step 3: Filters are washed for 1 hour at 37°C in a solution containing 2X SSC, 0.01%
 20 PVP, 0.01% Ficoll*, and 0.01% BSA. This is followed by a wash in 0.1X SSC at 50°C for 45 minutes. Step 4: Filters are autoradiographed. Other conditions of high stringency that may be used are well known in the art. *See, e.g.*, Ausubel *et al.*, (eds.), 1993, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley and Sons, NY; and Kriegler, 1990, GENE TRANSFER AND EXPRESSION, A LABORATORY MANUAL, Stockton Press, NY.

25 Moderate stringency conditions can include the following: Step 1: Filters containing DNA are pretreated for 6 hours at 55°C in a solution containing 6X SSC, 5X Denhardt's solution, 0.5% SDS and 100 mg/ml denatured salmon sperm DNA. Step 2: Filters are hybridized for 18-20 hours at 55°C in the same solution with 5-20 x 10⁶ cpm ³²P-labeled probe added. Step 3: Filters are washed at 37°C for 1 hour in a solution containing 2X SSC, 0.1% SDS, then washed
 30 twice for 30 minutes at 60°C in a solution containing 1X SSC and 0.1% SDS. Step 4: Filters are blotted dry and exposed for autoradiography. Other conditions of moderate stringency that may

*Trade-mark

WO 01/27268

PCT/IB00/01538

be used are well-known in the art. *See, e.g.,* Ausubel *et al.*, (eds.), 1993, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley and Sons, NY; and Kriegler, 1990, GENE TRANSFER AND EXPRESSION, A LABORATORY MANUAL, Stockton Press, NY.

Low stringency can include: Step 1: Filters containing DNA are pretreated for 6 hours at 40°C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 µg/ml denatured salmon sperm DNA. Step 2: Filters are hybridized for 18-20 hours at 40°C in the same solution with the addition of 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 µg/ml salmon sperm DNA, 10% (wt/vol) dextran sulfate, and 5-20 x 10⁶ cpm ³²P-labeled probe. Step 3: Filters are washed for 1.5 hours at 55°C in a solution containing 2X SSC, 25 mM Tris-HCl (pH 7.4), 5 mM EDTA, and 0.1% SDS. The wash solution is replaced with fresh solution and incubated an additional 1.5 hours at 60°C. Step 4: Filters are blotted dry and exposed for autoradiography. If necessary, filters are washed for a third time at 65-68°C and reexposed to film. Other conditions of low stringency that may be used are well known in the art (*e.g.*, as employed for cross-species hybridizations). *See, e.g.,* Ausubel *et al.*, (eds.), 1993, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley and Sons, NY; and Kriegler, 1990, GENE TRANSFER AND EXPRESSION, A LABORATORY MANUAL, Stockton Press, NY.

Chimeric Peptides Including a JNK Inhibitor Domain and a Trafficking Domain

In another aspect the invention provides a chimeric peptide that includes a first and second domain. The first domain includes a trafficking sequence, while the second domain includes a JNK inhibitor sequence linked by a covalent bond, *e.g.* peptide bond, to the first domain. The first and second domains can occur in any order in the peptide, and the peptide can include one or more of each domain.

A trafficking sequence is any sequence of amino acids that directs a peptide in which it is present to a desired cellular destination. Thus, the trafficking sequence can direct the peptide across the plasma membrane, *e.g.*, from outside the cell, through the plasma membrane, and into the cytoplasm. Alternatively, or in addition, the trafficking sequence can direct the peptide to a desired location within the cell, *e.g.*, the nucleus, the ribosome, the ER, a lysosome, or peroxisome.

50115-3

In some embodiments, the trafficking peptide is derived from a known membrane-translocating sequence. For example, the trafficking peptide may include sequences from the human immunodeficiency virus (HIV)1 TAT protein. This protein is described in, *e.g.*, U.S. Patent Nos. 5,804,604 and 5,674,980. The JNK inhibitor

5 peptide may be linked to some or all of the entire 86 amino acids that make up the TAT protein. For example, a functionally effective fragment or portion of a TAT protein that has fewer than 86 amino acids, which exhibits uptake into cells, and optionally uptake into the cell nucleus, can be used. In one embodiment, the fragment includes a peptide containing TAT residues 48 -57, *e.g.* $\text{NH}_2\text{-GRKKRRQRRR-COOH}$ [SEQ ID NO: 7] or a generic TAT sequence $\text{NH}_2\text{-X}_n\text{-RKKRRQRRR-}$
 10 $\text{X}_n\text{-COOH}$ [SEQ ID NO: 9]. A TAT peptide that includes the region that mediates entry and uptake into cells can be further defined using known techniques. *See, e.g.*, Franked *et al.*, *Proc. Natl. Acad. Sci. USA* 86: 7397-7401 (1989).

The TAT sequence may be linked either to the N-terminal or the C-terminal end of JNK inhibitor sequence. A hinge of two proline residues may be added between the TAT and JNK
 15 inhibitor peptide to create the full fusion peptide. For example, L-amino acid fusion peptides may be the L-TAT-IB1 peptide [SEQ ID NO:11], the L-TAT-IB2 peptide [SEQ ID NO:12], or the generic L-TAT-IB peptide [SEQ ID NO:13]. D retro-inverso fusion peptides may be the D-TAT-IB1 peptide [SEQ ID NO:14], the D-TAT-IB2 peptide [SEQ ID NO:15], or the generic D-TAT-IB peptide [SEQ ID NO:16]. The TAT peptide may be a D retro-inverso peptide having
 20 the sequence $\text{NH}_2\text{-X}_n\text{-RRRQRRKKR-X}_n\text{-COOH}$ [SEQ ID NO:10]. In SEQ ID NOs:5-6, 9-10, 13 and 16, the number of "X" residues is not limited to the one depicted, and may vary as described above.

The trafficking sequence can be a single (*i.e.*, continuous) amino acid sequence present in the TAT sequence. Alternatively it can be two or more amino acid sequences, which are present
 25 in TAT protein, but in the naturally-occurring protein are separated by other amino acid sequences. As used herein, TAT protein includes a naturally-occurring amino acid sequence that is the same as that of naturally-occurring TAT protein, or its functional equivalent protein or functionally equivalent fragments thereof (peptides). Such functional equivalent proteins or functionally equivalent fragments possess uptake activity into the cell and into the cell nucleus
 30 that is substantially similar to that of naturally-occurring TAT protein. TAT protein can be obtained from naturally-occurring sources or can be produced using genetic engineering

techniques or chemical synthesis.

The amino acid sequence of naturally-occurring HIV TAT protein can be modified, for example, by addition, deletion and/or substitution of at least one amino acid present in the naturally-occurring TAT protein, to produce modified TAT protein (also referred to herein as
5 TAT protein). Modified TAT protein or TAT peptide analogs with increased or decreased stability can be produced using known techniques. In some embodiments TAT proteins or peptides include amino acid sequences that are substantially similar, although not identical, to that of naturally-occurring TAT protein or portions thereof. In addition, cholesterol or other lipid derivatives can be added to TAT protein to produce a modified TAT having increased membrane
10 solubility.

Variants of the TAT protein can be designed to modulate intracellular localization of TAT- JNK inhibitor peptide. When added exogenously, such variants are designed such that the ability of TAT to enter cells is retained (*i.e.*, the uptake of the variant TAT protein or peptide into the cell is substantially similar to that of naturally-occurring HIV TAT). For example, alteration
15 of the basic region thought to be important for nuclear localization (*see, e.g.*, Dang and Lee, *J. Biol. Chem.* 264: 18019-18023 (1989); Hauber *et al.*, *J. Virol.* 63: 1181-1187 (1989); Ruben *et al.*, *J. Virol.* 63: 1-8 (1989)) can result in a cytoplasmic location or partially cytoplasmic location of TAT, and therefore, of the JNK inhibitor peptide. Alternatively, a sequence for binding a cytoplasmic or any other component or compartment (*e.g.*, endoplasmic reticulum, mitochondria,
20 Golgi apparatus, lysosomal vesicles,) can be introduced into TAT in order to retain TAT and the JNK inhibitor peptide in the cytoplasm or any other compartment to confer regulation upon uptake of TAT and the JNK inhibitor peptide.

Other sources for the trafficking peptide include, *e.g.*, VP22 (described in, *e.g.*, WO 97/05265; Elliott and O'Hare, *Cell* 88: 223-233 (1997)), or non-viral proteins (Jackson *et al.*,
25 *Proc. Natl. Acad. Sci. USA* 89: 10691-10695 (1992)).

The JNK inhibitor sequence and the trafficking sequence can be linked by chemical coupling in any suitable manner known in the art. Many known chemical cross-linking methods are non-specific, *i.e.*; they do not direct the point of coupling to any particular site on the transport polypeptide or cargo macromolecule. As a result, use of non-specific cross-linking
30 agents may attack functional sites or sterically block active sites, rendering the conjugated

proteins biologically inactive.

One way to increasing coupling specificity is to directly chemical coupling to a functional group found only once or a few times in one or both of the polypeptides to be cross-linked. For example, in many proteins, cysteine, which is the only protein amino acid containing a thiol group, occurs only a few times. Also, for example, if a polypeptide contains no lysine residues, a cross-linking reagent specific for primary amines will be selective for the amino terminus of that polypeptide. Successful utilization of this approach to increase coupling specificity requires that the polypeptide have the suitably rare and reactive residues in areas of the molecule that may be altered without loss of the molecule's biological activity.

Cysteine residues may be replaced when they occur in parts of a polypeptide sequence where their participation in a cross-linking reaction would otherwise likely interfere with biological activity. When a cysteine residue is replaced, it is typically desirable to minimize resulting changes in polypeptide folding. Changes in polypeptide folding are minimized when the replacement is chemically and sterically similar to cysteine. For these reasons, serine is preferred as a replacement for cysteine. As demonstrated in the examples below, a cysteine residue may be introduced into a polypeptide's amino acid sequence for cross-linking purposes. When a cysteine residue is introduced, introduction at or near the amino or carboxy terminus is preferred. Conventional methods are available for such amino acid sequence modifications, whether the polypeptide of interest is produced by chemical synthesis or expression of recombinant DNA.

Coupling of the two constituents can be accomplished via a coupling or conjugating agent. There are several intermolecular cross-linking reagents which can be utilized, See *for example*, Means and Feeney, CHEMICAL MODIFICATION OF PROTEINS, Holden-Day, 1974, pp. 39-43. Among these reagents are, for example, J-succinimidyl 3-(2-pyridyldithio) propionate (SPDP) or N, N'- (1,3-phenylene) bismaleimide (both of which are highly specific for sulfhydryl groups and form irreversible linkages); N, N'-ethylene-bis- (iodoacetamide) or other such reagent having 6 to 11 carbon methylene bridges (which relatively specific for sulfhydryl groups); and 1,5-difluoro-2, 4-dinitrobenzene (which forms irreversible linkages with amino and tyrosine groups). Other cross-linking reagents useful for this purpose include: p,p'-difluoro-m,m'-dinitrodiphenylsulfone (which forms irreversible cross-linkages with amino and phenolic

groups); dimethyl adipimidate (which is specific for amino groups); phenol-1,4-disulfonylchloride (which reacts principally with amino groups); hexamethylenediisocyanate or diisothiocyanate, or azophenyl-p-diisocyanate (which reacts principally with amino groups); glutaraldehyde (which reacts with several different side chains) and disdiazobenzidine (which
5 reacts primarily with tyrosine and histidine).

Cross-linking reagents may be homobifunctional, *i.e.*, having two functional groups that undergo the same reaction. A preferred homobifunctional cross-linking reagent is bismaleimidohexane ("BMH"). BMH contains two maleimide functional groups, which react specifically with sulfhydryl-containing compounds under mild conditions (pH 6.5-7.7). The two
10 maleimide groups are connected by a hydrocarbon chain. Therefore, BMH is useful for irreversible cross-linking of polypeptides that contain cysteine residues.

Cross-linking reagents may also be heterobifunctional. Heterobifunctional cross-linking agents have two different functional groups, for example an amine-reactive group and a thiol-reactive group, that will cross-link two proteins having free amines and thiols, respectively.
15 Examples of heterobifunctional cross-linking agents are succinimidyl 4-(N-maleimidomethyl) cyclohexane-1-carboxylate ("SMCC"), m-maleimidobenzoyl-N-hydroxysuccinimide ester ("MBS"), and succinimide 4-(p-maleimidophenyl) butyrate ("SMPB"), an extended chain analog of MBS. The succinimidyl group of these cross-linkers reacts with a primary amine, and the thiol-reactive maleimide forms a covalent bond with the thiol of a cysteine residue.

20 Cross-linking reagents often have low solubility in water. A hydrophilic moiety, such as a sulfonate group, may be added to the cross-linking reagent to improve its water solubility. Sulfo-MBS and sulfo-SMCC are examples of cross-linking reagents modified for water solubility.

Many cross-linking reagents yield a conjugate that is essentially non-cleavable under
25 cellular conditions. However, some cross-linking reagents contain a covalent bond, such as a disulfide, that is cleavable under cellular conditions. For example, Traut's reagent, dithiobis (succinimidylpropionate) ("DSP"), and N-succinimidyl 3-(2-pyridyldithio) propionate ("SPDP") are well-known cleavable cross-linkers. The use of a cleavable cross-linking reagent permits the cargo moiety to separate from the transport polypeptide after delivery into the target cell. Direct
30 disulfide linkage may also be useful.

Numerous cross-linking reagents, including the ones discussed above, are commercially available. Detailed instructions for their use are readily available from the commercial suppliers. A general reference on protein cross-linking and conjugate preparation is: Wong, CHEMISTRY OF PROTEIN CONJUGATION AND CROSS-LINKING, CRC Press (1991).

5 Chemical cross-linking may include the use of spacer arms. Spacer arms provide intramolecular flexibility or adjust intramolecular distances between conjugated moieties and thereby may help preserve biological activity. A spacer arm may be in the form of a polypeptide moiety that includes spacer amino acids, *e.g.* proline. Alternatively, a spacer arm may be part of the cross-linking reagent, such as in "long-chain SPDP" (Pierce Chem. Co., Rockford, IL., cat.
10 No. 21651 H).

Alternatively, the chimeric peptide can be produced as a fusion peptide that includes the trafficking sequence and the JNK inhibitor sequence which can conveniently be expressed in known suitable host cells. Fusion peptides, as described herein, can be formed and used in ways analogous to or readily adaptable from standard recombinant DNA techniques, as describe
15 above.

PRODUCTION OF ANTIBODIES SPECIFIC FOR JNK INHIBITOR PEPTIDES

JNK inhibitor peptides, including chimeric peptides including the JNK inhibitor peptides (*e.g.*, peptides including the amino acid sequences shown in Table 1), as well peptides, or derivatives, fragments, analogs or homologs thereof, may be utilized as immunogens to generate
20 antibodies that immunospecifically-bind these peptide components. Such antibodies include, *e.g.*, polyclonal, monoclonal, chimeric, single chain, Fab fragments and a Fab expression library. In a specific embodiment, antibodies to human peptides are disclosed. In another specific embodiment, fragments of the JNK inhibitor peptides are used as immunogens for antibody production. Various procedures known within the art may be used for the production of
25 polyclonal or monoclonal antibodies to a JNK inhibitor peptide, or derivative, fragment, analog or homolog thereof.

For the production of polyclonal antibodies, various host animals may be immunized by injection with the native peptide, or a synthetic variant thereof, or a derivative of the foregoing. Various adjuvants may be used to increase the immunological response and include, but are not

limited to, Freund's (complete and incomplete), mineral gels (*e.g.*, aluminum hydroxide), surface active substances (*e.g.*, lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, dinitrophenol, etc.) and human adjuvants such as *Bacille Calmette-Guerin* and *Corynebacterium parvum*.

5 For preparation of monoclonal antibodies directed towards a JNK inhibitor peptide, or derivatives, fragments, analogs or homologs thereof, any technique that provides for the production of antibody molecules by continuous cell line culture may be utilized. Such techniques include, but are not limited to, the hybridoma technique (*see*, Kohler and Milstein, 1975. *Nature* 256: 495-497); the trioma technique; the human B-cell hybridoma technique (*see*,
10 Kozbor, *et al.*, 1983. *Immunol Today* 4: 72) and the EBV hybridoma technique to produce human monoclonal antibodies (*see*, Cole, *et al.*, 1985. In: *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96). Human monoclonal antibodies may be utilized in the practice of the present invention and may be produced by the use of human hybridomas (*see*, Cote, *et al.*, 1983. *Proc Natl Acad Sci USA* 80: 2026-2030) or by transforming human B-cells
15 with Epstein Barr Virus *in vitro* (*see*, Cole, *et al.*, 1985. In: *Monoclonal Antibodies and Cancer Therapy* (Alan R. Liss, Inc., pp. 77-96).

According to the invention, techniques can be adapted for the production of single-chain antibodies specific to a JNK inhibitor peptide (*see, e.g.*, U.S. Patent No. 4,946,778). In addition, methodologies can be adapted for the construction of Fab expression libraries (*see, e.g.*, Huse, *et al.*, 1989. *Science* 246: 1275-1281) to allow rapid and effective identification of monoclonal Fab
20 fragments with the desired specificity for a JNK inhibitor peptide or derivatives, fragments, analogs or homologs thereof. Non-human antibodies can be "humanized" by techniques well known in the art. *See e.g.*, U.S. Patent No. 5,225,539. Antibody fragments that contain the idiotypes to a JNK inhibitor peptide may be produced by techniques known in the art including,
25 *e.g.*, (i) an F(ab')₂ fragment produced by pepsin digestion of an antibody molecule; (ii) an Fab fragment generated by reducing the disulfide bridges of an F(ab')₂ fragment; (iii) an Fab fragment generated by the treatment of the antibody molecule with papain and a reducing agent and (iv) Fv fragments.

In one embodiment, methodologies for the screening of antibodies that possess the
30 desired specificity include, but are not limited to, enzyme-linked immunosorbent assay (ELISA)

and other immunologically-mediated techniques known within the art. In a specific embodiment, selection of antibodies that are specific to a particular domain of a JNK inhibitor peptide is facilitated by generation of hybridomas that bind to the fragment of a JNK inhibitor peptide possessing such a domain. Antibodies that are specific for a domain within a JNK inhibitor peptide, or derivative, fragments, analogs or homologs thereof, are also provided herein.

The anti-JNK inhibitor peptide antibodies may be used in methods known within the art relating to the localization and/or quantitation of a JNK inhibitor peptide (*e.g.*, for use in measuring levels of the peptide within appropriate physiological samples, for use in diagnostic methods, for use in imaging the peptide, and the like). In a given embodiment, antibodies for the JNK inhibitor peptides, or derivatives, fragments, analogs or homologs thereof that contain the antibody derived binding domain, are utilized as pharmacologically active compounds [hereinafter "Therapeutics"].

METHODS OF TREATING OR PREVENTING DISORDERS ASSOCIATED UNDESIRE JNK ACTIVITY

Also included in the invention also are methods of treating cell-proliferative disorders associated with JNK activation in a subject by administering to a subject a biologically-active therapeutic compound (hereinafter "Therapeutic"). The subject can be *e.g.*, any mammal, *e.g.*, a human, a primate, mouse, rat, dog, cat, cow, horse, pig.

The Therapeutics include, *e.g.*: (i) any one or more of the JNK inhibitor peptides, and derivative, fragments, analogs and homologs thereof; (ii) antibodies directed against the JNK inhibitor peptides; (iii) nucleic acids encoding a JNK inhibitor peptide, and derivatives, fragments, analogs and homologs thereof; (iv) antisense nucleic acids to sequences encoding a JNK inhibitor peptide, and (v) modulators (*i.e.*, inhibitors, agonists and antagonists).

The term "therapeutically effective" means that the amount of inhibitor peptide, for example, which is used, is of sufficient quantity to ameliorate the JNK associated disorder. The term "cell-proliferative disorder" denotes malignant as well as non-malignant cell populations that often appear to differ morphologically and functionally from the surrounding tissue. For example, the method may be useful in treating malignancies of the various organ systems, in which activation of JNK has often been demonstrated, *e.g.*, lung, breast, lymphoid,

WO 01/27268

PCT/IB00/01538

gastrointestinal, and genito-urinary tract as well as adenocarcinomas which include malignancies such as most colon cancers, renal-cell carcinoma, prostate cancer, non-small cell carcinoma of the lung, cancer of the small intestine and cancer of the esophagus. Cancers with Bcr-Abl oncogenic transformations that clearly require activation of JNK are also included.

5 The method is also useful in treating non-malignant or immunological-related cell-proliferative diseases such as psoriasis, pemphigus vulgaris, Behcet's syndrome, acute respiratory distress syndrome (ARDS), ischemic heart disease, post-dialysis syndrome, leukemia, rheumatoid arthritis, acquired immune deficiency syndrome, vasculitis, septic shock and other types of acute inflammation, and lipid histiocytosis. Especially preferred are
10 immunopathological disorders. Essentially, any disorder, which is etiologically linked to JNK kinase activity, would be considered susceptible to treatment.

Treatment includes administration of a reagent that modulates JNK kinase activity. The term "modulate" includes the suppression of expression of JNK when it is over-expressed. It also includes suppression of phosphorylation of c-jun, ATF2 or NFAT4, for example, by using a
15 peptide of any one or more of SEQ ID NOs: 1-6 and SEQ ID NOs: 11-16 as a competitive inhibitor of the natural c-jun ATF2 and NFAT4 binding site in a cell. Thus also includes suppression of hetero- and homo-meric complexes of transcription factors made up of c-jun, ATF2, or NFAT4 and their related partners, such as for example the AP-1 complex that is made up of c-jun, AFT2 and c-fos. When a cell proliferative disorder is associated with JNK
20 overexpression, such suppressive JNK inhibitor peptides can be introduced to a cell. In some instances, "modulate" may include the increase of JNK expression, for example by use of an IB peptide-specific antibody that blocks the binding of an IB-peptide to JNK, thus preventing JNK inhibition by the IB-related peptide.

The JNK inhibitor, peptides, fusion peptides and nucleic acids of the invention can be
25 formulated in pharmaceutical compositions. These compositions may comprise, in addition to one of the above substances, a pharmaceutically acceptable excipient, carrier, buffer, stabiliser or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material may depend on the route of administration, *e.g.* oral, intravenous, cutaneous or
30 subcutaneous, nasal, intramuscular, intraperitoneal or patch routes.

Pharmaceutical compositions for oral administration may be in tablet, capsule, powder or liquid form. A tablet may include a solid carrier such as gelatin or an adjuvant. Liquid pharmaceutical compositions generally include a liquid carrier such as water, petroleum, animal or vegetable oils, mineral oil or synthetic oil. Physiological saline solution, dextrose or other
5 saccharide solution or glycols such as ethylene glycol, propylene glycol or polyethylene glycol may be included.

For intravenous, cutaneous or subcutaneous injection, or injection at the site of affliction, the active ingredient will be in the form of a parenterally acceptable aqueous solution which is pyrogen-free and has suitable pH, isotonicity and stability. Those of relevant skill in the art are
10 well able to prepare suitable solutions using, for example, isotonic vehicles such as Sodium Chloride Injection, Ringer's Injection, Lactated Ringer's Injection. Preservatives, stabilisers, buffers, antioxidants and/or other additives may be included, as required.

Whether it is a polypeptide, peptide, or nucleic acid molecule, other pharmaceutically useful compound according to the present invention that is to be given to an individual,
15 administration is preferably in a "prophylactically effective amount" or a "therapeutically effective amount" (as the case may be, although prophylaxis may be considered therapy), this being sufficient to show benefit to the individual. The actual amount administered, and rate and time-course of administration, will depend on the nature and severity of what is being treated. Prescription of treatment, *e.g.* decisions on dosage etc, is within the responsibility of general
20 practitioners and other medical doctors, and typically takes account of the disorder to be treated, the condition of the individual patient, the site of delivery, the method of administration and other factors known to practitioners. Examples of the techniques and protocols mentioned above can be found in REMINGTON'S PHARMACEUTICAL SCIENCES, 16th edition, Osol, A. (ed), 1980.

Alternatively, targeting therapies may be used to deliver the active agent more
25 specifically to certain types of cell, by the use of targeting systems such as antibody or cell specific ligands. Targeting may be desirable for a variety of reasons; for example if the agent is unacceptably toxic, or if it would otherwise require too high a dosage, or if it would not otherwise be able to enter the target cells.

Instead of administering these agents directly, they could be produced in the target cells
30 by expression from an encoding gene introduced into the cells, *e.g.* in a viral vector (a variant of

the VDEPT technique - *see* below). The vector could be targeted to the specific cells to be treated, or it could contain regulatory elements, which are switched on more or less selectively by the target cells.

Alternatively, the agent could be administered in a precursor form, for conversion to the active form by an activating agent produced in, or targeted to, the cells to be treated. This type of approach is sometimes known as ADEPT or VDEPT; the former involving targeting the activating agent to the cells by conjugation to a cell-specific antibody, while the latter involves producing the activating agent, *e.g.* a JNK inhibitor peptide, in a vector by expression from encoding DNA in a viral vector (*see* for example, EP-A-415731 and WO 90/07936).

In a specific embodiment of the present invention, nucleic acids include a sequence that encodes a JNK inhibitor peptide, or functional derivatives thereof, are administered to modulate activated JNK signaling pathways by way of gene therapy. In more specific embodiments, a nucleic acid or nucleic acids encoding a JNK inhibitor peptide, or functional derivatives thereof, are administered by way of gene therapy. Gene therapy refers to therapy that is performed by the administration of a specific nucleic acid to a subject. In this embodiment of the present invention, the nucleic acid produces its encoded peptide(s), which then serve to exert a therapeutic effect by modulating function of the disease or disorder. Any of the methodologies relating to gene therapy available within the art may be used in the practice of the present invention. *See e.g., Goldspiel, et al., 1993. Clin Pharm 12: 488-505.*

In a preferred embodiment, the Therapeutic comprises a nucleic acid that is part of an expression vector expressing any one or more of the IB-related peptides, or fragments, derivatives or analogs thereof, within a suitable host. In a specific embodiment, such a nucleic acid possesses a promoter that is operably-linked to coding region(s) of a JNK inhibitor peptide. The promoter may be inducible or constitutive, and, optionally, tissue-specific. In another specific embodiment, a nucleic acid molecule is used in which coding sequences (and any other desired sequences) are flanked by regions that promote homologous recombination at a desired site within the genome, thus providing for intra-chromosomal expression of nucleic acids. *See e.g., Koller and Smithies, 1989. Proc Natl Acad Sci USA 86: 8932-8935.*

Delivery of the Therapeutic nucleic acid into a patient may be either direct (*i.e.*, the patient is directly exposed to the nucleic acid or nucleic acid-containing vector) or indirect (*i.e.*,

cells are first transformed with the nucleic acid *in vitro*, then transplanted into the patient).

These two approaches are known, respectively, as *in vivo* or *ex vivo* gene therapy. In a specific embodiment of the present invention, a nucleic acid is directly administered *in vivo*, where it is expressed to produce the encoded product. This may be accomplished by any of numerous

5 methods known in the art including, *e.g.*, constructing the nucleic acid as part of an appropriate nucleic acid expression vector and administering the same in a manner such that it becomes intracellular (*e.g.*, by infection using a defective or attenuated retroviral or other viral vector; *see* U.S. Patent No. 4,980,286); directly injecting naked DNA; using microparticle bombardment (*e.g.*, a "Gene Gun[®]"; Biolistic, DuPont); coating the nucleic acids with lipids; using associated
10 cell-surface receptors/transfecting agents; encapsulating in liposomes, microparticles, or microcapsules; administering it in linkage to a peptide that is known to enter the nucleus; or by administering it in linkage to a ligand predisposed to receptor-mediated endocytosis (*see, e.g.*, Wu and Wu, 1987. *J Biol Chem* 262: 4429-4432), which can be used to "target" cell types that specifically express the receptors of interest, etc.

15 An additional approach to gene therapy in the practice of the present invention involves transferring a gene into cells in *in vitro* tissue culture by such methods as electroporation, lipofection, calcium phosphate-mediated transfection, viral infection, or the like. Generally, the method of transfer includes the concomitant transfer of a selectable marker to the cells. The cells are then placed under selection pressure (*e.g.*, antibiotic resistance) so as to facilitate the isolation
20 of those cells that have taken up, and are expressing, the transferred gene. Those cells are then delivered to a patient. In a specific embodiment, prior to the *in vivo* administration of the resulting recombinant cell, the nucleic acid is introduced into a cell by any method known within the art including, *e.g.*, transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences of interest, cell fusion,
25 chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, and similar methodologies that ensure that the necessary developmental and physiological functions of the recipient cells are not disrupted by the transfer. *See e.g.*, Loeffler and Behr, 1993. *Meth Enzymol* 217: 599-618. The chosen technique should provide for the stable transfer of the nucleic acid to the cell, such that the nucleic acid is expressible by the cell. Preferably, the
30 transferred nucleic acid is heritable and expressible by the cell progeny.

In preferred embodiments of the present invention, the resulting recombinant cells may

WO 01/27268

PCT/IB00/01538

be delivered to a patient by various methods known within the art including, *e.g.*, injection of epithelial cells (*e.g.*, subcutaneously), application of recombinant skin cells as a skin graft onto the patient, and intravenous injection of recombinant blood cells (*e.g.*, hematopoietic stem or progenitor cells). The total amount of cells that are envisioned for use depend upon the desired effect, patient state, and the like, and may be determined by one skilled within the art.

Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and may be xenogeneic, heterogeneic, syngeneic, or autogeneic. Cell types include, but are not limited to, differentiated cells such as epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes and blood cells, or various stem or progenitor cells, in particular embryonic heart muscle cells, liver stem cells (International Patent Publication WO 94/08598), neural stem cells (Stemple and Anderson, 1992, *Cell* 71: 973-985), hematopoietic stem or progenitor cells, *e.g.*, as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, and the like. In a preferred embodiment, the cells utilized for gene therapy are autologous to the patient.

IMMUNOASSAYS

The peptides and antibodies of the present invention may be utilized in assays (*e.g.*, immunoassays) to detect, prognose, diagnose, or monitor various conditions, diseases, and disorders characterized by aberrant levels of JNK, or a JNK inhibitor peptide, or monitor the treatment thereof. An "aberrant level" means an increased or decreased level in a sample relative to that present in an analogous sample from an unaffected part of the body, or from a subject not having the disorder. The immunoassay may be performed by a method comprising contacting a sample derived from a patient with an antibody under conditions such that immunospecific-binding may occur, and subsequently detecting or measuring the amount of any immunospecific-binding by the antibody. In a specific embodiment, an antibody specific for a JNK inhibitor peptide may be used to analyze a tissue or serum sample from a patient for the presence of JNK or a JNK inhibitor peptide; wherein an aberrant level of JNK or a JNK inhibitor peptide is indicative of a diseased condition. The immunoassays that may be utilized include, but are not limited to, competitive and non-competitive assay systems using techniques such as Western Blots, radioimmunoassays (RIA), enzyme linked immunosorbent assay (ELISA), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion

50115-3

precipitin reactions, immunodiffusion assays, agglutination assays, fluorescent immunoassays, complement-fixation assays, immunoradiometric assays, and protein-A immunoassays, etc.

KITS

The present invention additionally provides kits for diagnostic or therapeutic use that
5 include one or more containers containing an anti-JNK inhibitor peptide antibody and, optionally, a labeled binding partner to the antibody. The label incorporated into the antibody may include, but is not limited to, a chemiluminescent, enzymatic, fluorescent, colorimetric or radioactive moiety. In another specific embodiment, kits for diagnostic use that are comprised of one or more containers containing modified or unmodified nucleic acids that encode, or
10 alternatively, that are the complement to, a JNK inhibitor peptide and, optionally, a labeled binding partner to the nucleic acids, are also provided. In an alternative specific embodiment, the kit may comprise, in one or more containers, a pair of oligonucleotide primers (*e.g.*, each 6-30 nucleotides in length) that are capable of acting as amplification primers for polymerase chain reaction (PCR; *see, e.g.*, Innis, *et al.*, 1990. PCR PROTOCOLS, Academic Press, Inc., San
15 Diego, CA), ligase chain reaction, cyclic probe reaction, and the like, or other methods known within the art. The kit may, optionally, further comprise a predetermined amount of a purified JNK inhibitor peptide, or nucleic acids thereof, for use as a diagnostic, standard, or control in the assays.

The present invention is not to be limited in scope by the specific embodiments described
20 herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications fall within the scope of the appended claims.

25 SPECIFIC EXAMPLES

Example 1: Identification of JNK Inhibitor Peptides

Amino acid sequences important for efficient interaction with JNK were identified by

sequence alignments between known JBDs. Sequence comparison between the JBDs of IB1 [SEQ ID NO:17], IB2 [SEQ ID NO:18], c-Jun [SEQ ID NO:19] and ATF2 [SEQ ID NO:20] defined a weakly conserved 8 amino acid sequence (FIG. 1A). Since the JBDs of IB1 and IB2 are approximately 100 fold as efficient as c-Jun or ATF2 in binding JNK (Dickens *et al. Science* 277: 693 (1997)), it was reasoned that conserved residues between IB1 and IB2 must be important to confer maximal binding. The comparison between the JBDs of IB1 and IB2 defined two blocks of seven and three amino acids that are highly conserved between the two sequences. These two blocks are contained within a peptide sequence of 23 amino acids in IB1 [SEQ ID NO:1] and 21 amino acid IB2 [SEQ ID NO:2]. These sequences are shown in FIG. 1B, dashes in the IB2 sequence indicate a gap in the sequence in order to align the conserved residues.

Example 2: Preparation of JNK Inhibitor Fusion Proteins

JNK inhibitor fusion proteins were synthesized by covalently linking the C-terminal end of JBD₂₃ or the 21 amino acid sequence derived from the JBD of IB2 (JBD₂₁) to a N-terminal 10 amino acid long carrier peptide derived from the HIV-TAT₄₈₋₅₇ (Vives *et al., J. Biol. Chem.* 272: 16010 (1997)) via a spacer consisting of two proline residues. This spacer was used to allow for maximal flexibility and prevent unwanted secondary structural changes. As shown in FIG. 1C, these preparations were designated L-TAT [SEQ ID NO:7], L-TAT-IB1 [SEQ ID NO:11] and L-TAT-IB2 [SEQ ID NO:12], respectively. All-D retro-inverso peptides TAT-fusion peptides were also synthesized and were designated D-TAT [SEQ ID NO:8] and D-TAT-IB1 [SEQ ID NO:14], respectively. All D and L peptides were produced by classical F-mock synthesis and further analysed by Mass Spectrometry. They were finally purified by HPLC. To determine the effects of the proline spacer, two types of TAT peptide were produced one with and one without two prolines. The addition of the two prolines did not appear to modify the entry or the localization of the TAT peptide inside cells.

Generic peptides showing the conserved amino acid residues are given in FIG. 2. An "X" indicates any amino acid. The number of Xs in a given peptide is not limited to the one depicted, and may vary. See above for a more detailed description of the generic sequences.

Example 3: Inhibition of β Cell Death By JBD₂₃

Effects of the 23 a.a. long JBD sequence of IB1 on JNK biological activities were then studied. The 23 a.a. sequence was linked N-terminal to the Green Fluorescent Protein (GFP-JBD₂₃ construct), and the effect of this construct on pancreatic β -cell apoptosis induced by IL-1 β was evaluated. See FIG. 3. This mode of apoptosis was previously shown to be blocked by transfection with JBD₁₋₂₈₀, whereas specific inhibitors of ERK1/2 or p38 did not protect. See Ammendrup *et al*, *supra*.

Oligonucleotides corresponding to the 23 amino acid sequence (JBD₂₃; FIG 1B) and a sequence mutated at the fully conserved regions (JBD_{23mut}) were synthesized and directionally inserted into the EcoRI and Sall sites of the pEGFP-N1 vector encoding the Green Fluorescent Protein (GFP) (from Clontech). Insulin producing β TC-3 cells were cultured in RPMI 1640 medium supplemented with 10% Fetal Calf Serum, 100 μ g/mL Streptomycin, 100 units/mL Penicillin and 2 mM Glutamine. Insulin producing β TC-3 cells were transfected with the indicated vectors and IL-1 β (10 ng/mL) was added to the cell culture medium. The number of apoptotic cells were counted at 48 hours after the addition of IL-1 β using an inverted fluorescence microscope. Apoptotic cells were discriminated from normal cells by the characteristic "blebbing out" of the cytoplasm were counted after two days.

As indicated in FIG. 3, GFP is Green Fluorescent protein expression vector used as a control; JBD₂₃ is the vector expressing a chimeric GFP linked to the 23 a.a. sequence from the JBD of IB1; JBD_{23Mut} is the same vector as GFP-JBD₂₃, but with a JBD mutated at four conserved residues shown as FIG 1B; and JBD₂₈₀ is the GFP vector linked to the entire JBD (a.a. 1-280). The GFP-JBD₂₃ expressing construct prevented IL-1 β induced pancreatic β -cell apoptosis as efficiently as the entire JBD₁₋₂₈₀ (FIG. 3, JBD₂₃/IL-1 compared to JBD₂₈₀/IL-1). As additional controls, sequences mutated at fully conserved IB1 residues had greatly decreased ability to prevent apoptosis (FIG. 3, JBD_{23Mut}/IL-1).

Example 4: Cellular Import Of TAT-IB1 And TAT-IB2 Peptides

The ability of the L- and D-enantiomeric forms of TAT, TAT-IB1 and TAT-IB2 peptides ("TAT-IB peptides") to enter cells were evaluated.

L-TAT, D-TAT, L-TAT-IB1, L-TAT-IB2 and D-TAT-IB1 peptides [SEQ ID NOs:7, 8, 11, 12 and 14, respectively] were labeled by N-terminal addition of a glycine residue conjugated to fluorescein. Labeled peptides (1 μ M) were added to β TC-3 cell cultures, which were maintained as described in Example 3. At predetermined times, cells were washed with PBS and
5 fixed for five minutes in ice-cold methanol-acetone (1:1) before being examined under a fluorescence microscope. Fluorescein-labeled BSA (1 μ M, 12 moles/mole BSA) was used as a control. Results demonstrated that all the above fluorescein labeled peptides had efficiently and rapidly (less than five minutes) entered cells once added to the culture medium. Conversely, fluorescein labeled bovine serum albumin (1 μ M BSA, 12 moles fluorescein/mole BSA) did not
10 enter the cells.

A time course study indicated that the intensity of the fluorescent signal for the L-enantiomeric peptides decreased by 70% following a 24 hours period. Little to no signal was present at 48 hours. In contrast, D-TAT and D-TAT-IB1 were extremely stable inside the cells. Fluorescent signals from these all-D retro-inverso peptides were still very strong 1 week later,
15 and the signal was only slightly diminish at 2 weeks post treatment.

Example 5: *In Vitro* Inhibition Of c-JUN, ATF2 and Elk1 Phosphorylation

The effects of the peptides on JNKs-mediate phosphorylation of their target transcription factors were investigated *in vitro*. Recombinant and nonactivated JNK1, JNK2 and JNK3 were produced using a TRANSCRIPTION AND TRANSLATION rabbit reticulocyte lysate kit (Promega) and
20 used in solid phase kinase assays with c-Jun, ATF2 and Elk1, either alone or fused to glutathione-S-transferase (GST), as substrates. Dose response studies were performed wherein L-TAT, L-TAT-IB1 or L-TAT-IB2 peptides (0-25 μ M) were mixed with the recombinant JNK1, JNK2, or JNK3 kinases in reaction buffer (20 mM Tris-acetate, 1mM EGTA, 10mM p-nitrophenyl-phosphate (pNPP), 5 mM sodium pyrophosphate, 10 mM p-glycerophosphate, 1mM
25 dithiothreitol) for 20 minutes. The kinase reactions were then initiated by the addition of 10 mM $MgCl_2$ and 5 μ Ci ^{33}P - γ -dATP and 1 μ g of either GST-Jun (a.a. 1-89), GST-AFT2 (a.a. 1-96) or GST-ELK1 (a.a. 307-428). GST-fusion proteins were purchased from Stratagene (La Jolla, CA). Ten μ L of glutathione-agarose beads were also added to the mixture. Reaction products were then separated by SDS-PAGE on a denaturing 10 % polyacrylamide gel. Gels were dried and

50115-3

subsequently exposed to X-ray films (Kodak). Nearly complete inhibition of c-Jun, ATF2 and Elk1 phosphorylation by JNKs was observed at TAT-IB peptide doses as low as 2.5 μ M. However, a marked exception was the absence of TAT-IB inhibition of JNK3 phosphorylation of Elk1. Overall, the TAT-IB1 peptide appeared slightly superior to TAT-IB2 in inhibiting JNK family phosphorylation of their target transcription factors. (See, FIG. 4A).

The ability of D-TAT, D-TAT-IB1 and L-TAT-IB1 peptides (0-250 μ M dosage study) to inhibit GST-Jun (a.a. 1-73) phosphorylation by recombinant JNK1, JNK2, and JNK3 by were analyzed as described above. Overall, D-TAT-IB1 peptide decreased JNK-mediated phosphorylation of c-Jun, but at levels approximately 10-20 fold less efficiently than L-TAT-IB1. (See, FIG. 4B).

Example 6: Inhibition of c-JUN Phosphorylation By Activated JNKs

The effects of the L-TAT, L-TAT-IB1 or L-TAT-IB2 peptides on JNKs activated by stressful stimuli were evaluated using GST-Jun to pull down JNKs from UV-light irradiated HeLa cells or IL-1 β treated β TC cells. β TC cells were cultured as described above. HeLa cells were cultured in DMEM medium supplemented with 10 % Fetal Calf Serum, 100 μ g/mL Streptomycin, 100 units/ml Penicillin and 2 mM Glutamine. One hour prior to being used for cell extract preparation, β TC cells were activated with IL-1 β as described above, whereas HeLa cells were activated by UV-light (20 J/m²). Cell extracts were prepared from control, UV-light irradiated HeLa cells and IL-1 β treated β TC-3 cells by scraping the cell cultures in lysis buffer (20 mM Tris-acetate, 1mM EGTA, 1% Triton^{*} X-100, 10 mM p-nitrophenyl-phosphate, 5 mM sodium pyrophosphate, 10 mM β -glycerophosphate, 1 mM dithiothreitol). Debris was removed by centrifugation for five minutes at 15,000 rpm in an SS-34 Beckman rotor. One-hundred μ g extracts were incubated for one hour at room temperature with one μ g GST-jun (amino acids 1-89) and 10 μ L of glutathione-agarose beads (Sigma). Following four washes with the scraping buffer, the beads were resuspended in the same buffer supplemented with L-TAT, L-TAT-IB1 or L-TAT-IB2 peptides (25 μ M) for 20 minutes. Kinase reactions were then initiated by the addition of 10 mM MgCl₂ and 5 μ Ci ³³P- γ -dATP and incubated for 30 minutes at 30°C. Reaction products were then separated by SDS-PAGE on a denaturing 10 % polyacrylamide gel. Gels were dried and subsequently exposed to X-ray films (Kodak). The TAT-IB peptides efficiently prevented phosphorylation of c-Jun by activated JNKs in these experiments. . (See,

*Trade-mark

FIG. 6).

Example 7: *In vivo* Inhibition of c-JUN Phosphorylation By TAT-IB Peptides

To determine whether the cell-permeable peptides could block JNK signaling *in vivo*, we used a heterologous GAL4 system. HeLa cells, cultured as described above, were co-transfected with the 5xGAL-LUC reporter vector together with the GAL-Jun expression construct (Stratagene) comprising the activation domain of c-Jun (amino acids 1-89) linked to the GAL4 DNA-binding domain. Activation of JNK was achieved by the co-transfection of vectors expressing the directly upstream kinases MKK4 and MKK7 (*See, Whitmarsh et al., Science* 285: 1573 (1999)). Briefly, 3×10^5 cells were transfected with the plasmids in 3.5-cm dishes using DOTAP (Boehringer Mannheim) following instructions from the manufacture. For experiments involving GAL-Jun, 20 ng of the plasmid was transfected with 1 μ g of the reporter plasmid pFR-Luc (Stratagene) and 0.5 μ g of either MKK4 or MKK7 expressing plasmids. Three hours following transfection, cell media were changed and TAT, TAT-IB1, and TAT-IB2 peptides (1 μ M) were added. The luciferase activities were measured 16 hours later using the "Dual Reporter System" from Promega after normalization to protein content. As shown in FIG. 5, addition of both the TAT-IB1 and TAT-IB2 peptides blocked activation of c-Jun following MKK4 and MKK7 mediated activation of JNK. Because HeLa cells express both JNK1 and JNK2 isoforms but not JNK3, we transfected cells with JNK3. Again, the two TAT-IB peptides inhibited JNK2 mediated activation of c-Jun.

Example 8: Inhibition Of IL-1 β Induced Pancreatic β -Cell Death By TAT-IB Peptides

We investigated the effects of the L-TAT-IB peptides on the promotion of pancreatic β -cell apoptosis elicited by IL-1 β . β TC-3 cell cultures were incubated for 30 minutes with 1 μ M of either L-TAT-IB1 or L-TAT-IB2 peptides followed by 10 ng/mL of IL-1 β . A second addition of peptide (1 μ M) was performed 24 hours later. Apoptotic cells were counted after two days of incubation with IL-1 β using Propidium Iodide (red stained cell are dead cells) and Hoechst 33342 (blue stained cell are cells with intact plasma membrane) nuclear staining. As shown in FIG. 5, addition of the TAT-IB peptides inhibited IL-1 β -induced apoptosis of β TC-3 cells cultured in the presence of IL-1 β for two days.

Long term inhibition of IL-1 β induced cells death was examined by treating β TC-3 cells as described above, except that incubation of the cells with the peptides and IL-1 β was sustained for 12 days. Additional peptides (1 μ M) were added each day and additional IL-1 β (10 ng/mL) was added every 2 days. The TAT-IB1 peptide confers strong protection against apoptosis in these conditions. Taken together, these experiments establish that TAT-IB peptides are biologically active molecules able to prevent the effects of JNK signaling on cell fate.

Example 9: Synthesis of an All-D-retro-inverso Peptides

Peptides of the invention may be all-D amino acid peptides synthesized in reverse to prevent natural proteolysis (*i.e.*, all-D-retro-inverso peptides). An all-D retro-inverso peptide of the invention would provide a peptide with functional properties similar to the native peptide, wherein the side groups of the component amino acids would correspond to the native peptide alignment, but would retain a protease resistant backbone.

Retro-inverso peptides of the invention are analogs synthesized using D-amino acids by attaching the amino acids in a peptide chain such that the sequence of amino acids in the retro-inverso peptide analog is exactly opposite of that in the selected peptide which serves as the model. To illustrate, if the naturally occurring TAT protein (formed of L-amino acids) has the sequence GRKKRRQRRR [SEQ ID NO:7], the retro-inverso peptide analog of this peptide (formed of D-amino acids) would have the sequence RRRQRRKKRG [SEQ ID NO:8]. The procedures for synthesizing a chain of D-amino acids to form the retro-inverso peptides are known in the art. *See, e.g.*, Jameson *et al.*, *Nature*, 368, 744-746 (1994); Brady *et al.*, *Nature*, 368, 692-693 (1994)); Guichard *et al.*, *J. Med. Chem.* 39, 2030-2039 (1996). Specifically, the retro- peptides were produced by classical F-mock synthesis and further analysed by Mass Spectrometry. They were finally purified by HPLC.

Since an inherent problem with native peptides is degradation by natural proteases and inherent immunogenicity, the heterobivalent or heteromultivalent compounds of this invention will be prepared to include the "retro-inverso isomer" of the desired peptide. Protecting the peptide from natural proteolysis should therefore increase the effectiveness of the specific heterobivalent or heteromultivalent compound, both by prolonging half-life and decreasing the extent of the immune response aimed at actively destroying the peptides.

Example 10: Long term biological activity of all-D-retro-inverso IB Peptides

Long term biological activity is predicted for the D-TAT-IB retro-inverso containing peptide heteroconjugate when compared to the native L-amino acid analog owing to protection of the D-TAT-IB peptide from degradation by native proteases, as shown in Example 5.

5 Inhibition of IL-1 β induced pancreatic β -cell death by the D-TAT-IB1 peptide was analyzed. As shown in FIG. 10, β TC-3 cells were incubated as described above for 30 minutes with one single addition of the indicated peptides (1 μ M), then IL-1 β (10 ng/ml) was added. Apoptotic cells were then counted after two days of incubation with IL-1 β by use of Propidium Iodide and Hoechst 33342 nuclear staining. A minimum of 1,000 cells were counted for each
10 experiment. Standard Error of the Means (SEM) are indicated, n=5. The D-TAT-IB1 peptide decreased IL-1 induced apoptosis to a similar extent as L-TAT-IB peptides (compare FIG. 5 and FIG. 10).

Long term inhibition of IL-1 β induced cell-death by the D-TAT-IB1 peptide was also analyzed. β TC-3 cells were incubated as above for 30 minutes with one single addition of the
15 indicated peptides (1 μ M), then IL-1 β (10 ng/ml) was added, followed by addition of the cytokine every two days. Apoptotic cells were then counted after 15 days of incubation with IL-1 β by use of Propidium Iodide and Hoechst 33342 nuclear staining. Note that one single addition of the L-TAT-IB1 peptide does not confer long-term protection. A minimum of 1,000 cells were counted for each experiment. Standard Error of the Means (SEM) are indicated, n=5. Results are shown
20 in FIG.9. D-TAT-IB1, but not L-TAT-IB1, was able to confer long term (15 day) protection.

Example 11: Inhibition Of Irradiation Induced Pancreatic β -Cell Death By TAT-IB Peptides

JNK is also activated by ionizing radiation. To determine whether TAT-IB peptides would provide protection against radiation-induced JNK damage, "WiDr" cells were irradiated
25 (30Gy) in presence or absence of D-TAT, L-TAT-IB1 or D-TAT-IB1 peptides (1 μ M added 30 minutes before irradiation), as indicated in FIG. 10. Control cells (CTRL) were not irradiated. Cells were analyzed 48 hours later by mean of PI and Hoechst 33342 staining, as described above. n=3, SEM are indicated. L-TAT-IB1 and D-TAT-IB1 peptides were both able to prevent irradiation induced apoptosis in this human colon cancer cell line.

Example 12: Radioprotection to Ionizing Radiation By TAT-IB Peptides

To determine the radioprotective effects of the TAT-IB peptides, C57 Bl/6 mice (2 to 3 months old) were irradiated with a Phillips RT 250 R-ray at a dose rate of 0.74 Gy/min (17 mA, 0.5 mm Cu filter). Thirty minutes prior to irradiation, the animals were injected i.p. with either
 5 the TAT, L-TAT-IB1 and D-TAT-IB1 peptides (30 μ l of a 1mM solution). Briefly, mice were irradiated as follows: mice were placed in small plastic boxes with the head lying outside the box. The animals were placed on their back under the irradiator, and their neck fixed in a small plastic tunnel to maintain their head in a correct position. The body was protected with lead. Prior to irradiation mice were maintained on standard pellet mouse chow, however post
 10 irradiation mice were fed with a semi-liquid food that was renewed each day.

The reaction of the lip mucosa was then scored by 2 independent observers according to the scoring system developed by Parkins *et al.* (Parkins *et al.*, *Radiotherapy & Oncology*, 1: 165-173, 1983), in which the erythema status as well as the presence of edema, desquamation and exudation was quoted. Additionally, animals were weighed before each recording of their
 15 erythema/edema status.

FIG. 12A: illustrated the weight of the mice following irradiation. Values are reported to the initial weight of the mice that was set to 100. CTRL: control mice injected with 30 μ l of a saline solution. n=2 for each values reported, S.D. are indicated. x values are days

FIG. 12B is illustrative of the erythema/edema scoring following irradiation. The edema
 20 and erythema status of the ventral lip of the same mice as in FIG. 12A was quantified. n=2 for each value reported. x values are days

The results of these experiments indicate that the TAT-IB Peptides can protect against weight loss and erythema/edema associated with ionizing radiation.

Example 13: Suppression of JNK Transcription Factors by L-TAT-IB1 peptides

25 Gel retardation assays were carried out with an AP-1 doubled labeled probe (5'- CGC TTG ATG AGT CAG CCG GAA-3'. HeLa cell nuclear extracts that were treated or not for one hour with 5 ng/ml TNF- α , as indicated. TAT and L-TAT-IB1 peptides were added 30 minutes before TNF- α . Only the part of the gel with the specific AP-1 DNA complex (as demonstrated by competition experiments with non-labeled specific and non-specific competitors) is shown.

WO 01/27268

PCT/IB00/01538

L-TAT -IB1 peptides decrease the formation of the AP-1 DNA binding complex in the presence of TNF- α . (See, FIG 11).

EQUIVALENTS

From the foregoing detailed description of the specific embodiments of the invention, it
5 should be apparent that unique cell-permeable bioactive peptides have been described. Although
particular embodiments have been disclosed herein in detail, this has been done by way of
example for purposes of illustration only, and is not intended to be limiting with respect to the
scope of the appended claims which follow. In particular, it is contemplated by the inventor that
various substitutions, alterations, and modifications may be made to the invention without
10 departing from the spirit and scope of the invention as defined by the claims.

1

SEQUENCE LISTING

<110> Bonny, Christophe

<120> CELL-PERMEABLE PEPTIDE INHIBITORS OF THE JNK SIGNAL
TRANSDUCTION PATHWAY

<130> 20349-501-008

<140> Not Yet Assigned

<141> 2000-10-12

<150> PCT/IB00/01538

<151> 2000-10-12

<150> 09/503,954

<151> 2000-02-14

<150> 60/158,774

<151> 1999-10-12

<160> 20

<170> PatentIn Ver. 2.0

<210> 1

<211> 23

<212> PRT

<213> Artificial Sequence

<220>

<223> chemically synthesized

<400> 1

Asp Thr Tyr Arg Pro Lys Arg Pro Thr Thr Leu Asn Leu Phe Pro Gln
1 5 10 15Val Pro Arg Ser Gln Asp Thr
20

<210> 2

<211> 21

<212> PRT

<213> Artificial Sequence

<220>

<223> chemically synthesized

<400> 2

Glu Glu Pro His Lys His Arg Pro Thr Thr Leu Arg Leu Thr Thr Leu
1 5 10 15Gly Ala Gln Asp Ser
20

2

<210> 3
 <211> 23
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> chemically synthesized

<400> 3

Thr Asp Gln Ser Arg Pro Val Gln Pro Phe Leu Asn Leu Thr Thr Pro
 1 5 10 15

Arg Lys Pro Arg Tyr Thr Asp
 20

<210> 4
 <211> 21
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> chemically synthesized

<400> 4

Ser Asp Gln Ala Gly Leu Thr Thr Leu Arg Leu Thr Thr Pro Arg His
 1 5 10 15

Lys His Pro Glu Glu

20

<210> 5
 <211> 279
 <212> PRT
 <213> Artificial Sequence

<220>
 <221> VARIANT

<222> (1) .. (131)

<223> Each Xaa is any amino acid, or absent

<220>
 <221> VARIANT

<222> (137)

<223> Xaa is any amino acid

<220>
 <221> VARIANT

<222> (139)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (140)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (141)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (142)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (143)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (144)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (145)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (148)

<223> Xaa is S or T

<220>

<221> VARIANT

<222> (149) .. (279)

<223> Each Xaa is any amino acid, or absent

<220>

<223> chemically synthesized

<400> 5

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	1	5	10	15
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	20	25	30	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	35	40	45	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	50	55	60	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	65	70	75	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	80	85	90	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	95	100	105	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	110	115	120	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Arg	Pro	Thr	Thr	125	130	135	
Leu	Xaa	Leu	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Gln	Asp	Xaa	Xaa	Xaa	140	145	150	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	155	160	165	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	170	175	180	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	185	190	195	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	200	205	210	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	215	220	225	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	230	235	240	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	245	250	255	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	260	265	270	

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
275

<210> 6

<211> 279

<212> PRT

<213> Artificial Sequence

<220>

<221> VARIANT

<222> (1) .. (131)

<223> Xaa is any amino acid, or absent

<220>

<221> VARIANT

<222> (132)

<223> Xaa is Ser or Thr

<220>

<221> VARIANT

<222> (135)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (136)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (137)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (138)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (139)

6

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (140)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (141)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (143)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (149) .. (279)

<223> Xaa is any amino acid, or absent

<220>

<223> chemically synthesized

<400> 6

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
1				5					10						15

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				20					25						30

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				35					40						45

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				50					55						60

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				65					70						75

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				80					85						90

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				95					100						105

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				110					115						120

7

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Asp	Gln	Xaa
				125					130					135
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Leu	Xaa	Leu	Thr	Thr	Pro	Arg	Xaa	Xaa
				140					145					150
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				155					160					165
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				170					175					180
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				185					190					195
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				200					205					210
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				215					220					225
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				230					235					240
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				245					250					255
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				260					265					270
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa						
				275										

<210> 7
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> chemically synthesized

<400> 7
 Gly Arg Lys Lys Arg Arg Gln Arg Arg Arg
 1 5 10

<210> 8
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> chemically synthesized

<400> 8
 Arg Arg Arg Gln Arg Arg Lys Lys Arg Gly

1

5

10

8

<210> 9

<211> 279

<212> PRT

<213> Artificial Sequence

<220>

<221> VARIANT

<222> (1) .. (132)

<223> Xaa is any amino acid, or absent

<220>

<221> VARIANT

<222> (133)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (134)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (135)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (145)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (146)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (147)

<223> Xaa is any amino acid

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa

10

230	235	240
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa	Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa	Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
245	250	255
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa	Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa	Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
260	265	270
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa		
275		

<210> 10
 <211> 279

<212> PRT
 <213> Artificial Sequence

<220>
 <221> VARIANT

<222> (1) .. (132)

<223> Xaa is any amino acid, or absent

<220>
 <221> VARIANT

<222> (133)

<223> Xaa is any amino acid

<220>
 <221> VARIANT

<222> (134)

<223> Xaa is any amino acid

<220>
 <221> VARIANT

<222> (135)

<223> Xaa is any amino acid

<220>
 <221> VARIANT

<222> (145)

<223> Xaa is any amino acid

<220>
 <221> VARIANT

<222> (146)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (147)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (148) .. (279)

<223> Xaa is any amino acid, or absent

<220>

<223> chemically synthesized

<400> 10

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
1				5					10					15

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				20					25					30

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				35					40					45

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				50					55					60

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				65					70					75

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				80					85					90

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				95					100					105

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				110					115					120

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				125					130					135

Arg	Arg	Arg	Gln	Arg	Arg	Lys	Lys	Arg	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				140					145					150

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				155					160					165

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				170					175					180

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				185					190					195

12

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 200 205 210
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 215 220 225
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 230 235 240
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 245 250 255
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 260 265 270
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 275

<210> 11
 <211> 35
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> chemically synthesized

<400> 11
 Gly Arg Lys Lys Arg Arg Gln Arg Arg Arg Pro Pro Asp Thr Tyr Arg
 1 5 10 15
 Pro Lys Arg Pro Thr Thr Leu Asn Leu Phe Pro Gln Val Pro Arg Ser
 20 25 30

Gln Asp Thr
 35

<210> 12
 <211> 33
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> chemically synthesized

<400> 12
 Gly Arg Lys Lys Arg Arg Gln Arg Arg Arg Pro Pro Glu Glu Pro His
 1 5 10 15
 Lys His Arg Pro Thr Thr Leu Arg Leu Thr Thr Leu Gly Ala Gln Asp
 20 25 30

Ser

<210> 13
<211> 42
<212> PRT
<213> Artificial Sequence

<220>
<221> VARIANT

<222> (1)
<223> may be any amino acid

<220>
<221> VARIANT

<222> (2)
<223> may be any amino acid

<220>
<221> VARIANT

<222> (3)
<223> may be any amino acid

<220>
<221> VARIANT

<222> (4)
<223> may be any amino acid

<220>
<221> VARIANT

<222> (5)
<223> may be any amino acid

<220>
<221> VARIANT

<222> (6)
<223> may be any amino acid

<220>
<221> VARIANT

<222> (7)
<223> may be any amino acid

<220>
<221> VARIANT

<222> (17)
<223> may be any amino acid

<220>
<221> VARIANT

<222> (18)
<223> may be any amino acid

<220>

<221> VARIANT

<222> (19)

<223> may be any amino acid

<220>

<221> VARIANT

<222> (20)

<223> may be any amino acid

<220>

<221> VARIANT

<222> (21)

<223> may be any amino acid

<220>

<221> VARIANT

<222> (22)

<223> may be any amino acid

<220>

<221> VARIANT

<222> (23)

<223> may be any amino acid

<220>

<221> VARIANT

<222> (24)

<223> may be any amino acid

<220>

<221> VARIANT

<222> (30)

<223> may be any amino acid

<220>

<221> VARIANT

<222> (32)

<223> many any amino acid

<220>

<221> VARIANT

<222> (33)

<223> may be any amino acid

<220>

<221> VARIANT

<222> (34)

<223> may be any amino acid

15

<220>

<221> VARIANT

<222> (35)

<223> may be any amino acid

<220>

<221> VARIANT

<222> (36)

<223> may be any amino acid

<220>

<221> VARIANT

<222> (37)

<223> may be any amino acid

<220>

<221> VARIANT

<222> (38)

<223> may be any amino acid

<220>

<221> VARIANT

<222> (41)

<223> may be S or T

<220>

<221> VARIANT

<222> (42)

<223> may be any amino acid

<220>

<223> chemically synthesized

<400> 13

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Arg	Lys	Lys	Arg	Arg	Gln	Arg	Arg	Arg
1				5					10					15	

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Arg	Pro	Thr	Thr	Leu	Xaa	Leu	Xaa
			20					25					30		

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Gln	Asp	Xaa	Xaa
			35				40		

<210> 14

<211> 35

<212> PRT

<213> Artificial Sequence

<220>

<223> chemically synthesized

<400> 14

Thr Asp Gln Ser Arg Pro Val Gln Pro Phe Leu Asn Leu Thr Thr Pro

1 5 10 15 16
 Arg Lys Pro Arg Tyr Thr Asp Pro Pro Arg Arg Arg Gln Arg Arg Lys
 20 25 30

Lys Arg Gly

35

<210> 15
 <211> 33
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> chemically synthesized

<400> 15
 Ser Asp Gln Ala Gly Leu Thr Thr Leu Arg Leu Thr Thr Pro Arg His
 1 5 10 15

Lys His Pro Glu Glu Pro Pro Arg Arg Arg Gln Arg Arg Lys Lys Arg
 20 25 30

Gly

<210> 16
 <211> 279
 <212> PRT
 <213> Artificial Sequence

<220>
 <221> VARIANT
 <222> (1) .. (120)

<223> Xaa is any amino acid, or absent

<220>
 <221> VARIANT

<222> (121)
 <223> Xaa is Ser or Thr

<220>
 <221> VARIANT

<222> (124)
 <223> Xaa is any amino acid

<220>
 <221> VARIANT

<222> (125)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (126)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (127)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (128)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (129)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (130)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (132)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (138)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (139)

<223> Xaa is any amino acid

<220>
<221> VARIANT

<222> (140)

<223> Xaa is any amino acid

<220>
<221> VARIANT

<222> (141)

<223> Xaa is any amino acid

<220>
<221> VARIANT

<222> (142)

<223> Xaa is any amino acid

<220>
<221> VARIANT

<222> (143)

<223> Xaa is any amino acid

<220>
<221> VARIANT

<222> (144)

<223> Xaa is any amino acid

<220>
<221> VARIANT

<222> (145)

<223> Xaa is any amino acid

<220>
<221> VARIANT

<222> (155)

<223> Xaa is any amino acid

<220>
<221> VARIANT

<222> (156)

<223> Xaa is any amino acid

<220>
<221> VARIANT

<222> (157)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (158)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (159)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (160)

<223> Xaa is any amino acid

<220>

<221> VARIANT

<222> (161) .. (279)

<223> Xaa is any amino acid, or absent

<220>

<223> chemically synthesized

<400> 16

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
1				5					10					15

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				20					25					30

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				35					40					45

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				50					55					60

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				65					70					75

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				80					85					90

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				95					100					105

:

20

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				110						115					120
Xaa	Asp	Gln	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Leu	Xaa	Leu	Thr	Thr	
				125						130					135
Pro	Arg	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Arg	Arg	Arg	Gln	Arg	
				140						145					150
Arg	Lys	Lys	Arg	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				155						160					165
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				170						175					180
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				185						190					195
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				200						205					210
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				215						220					225
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				230						235					240
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				245						250					255
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				260						265					270
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa						
				275											

<210> 17

<211> 29

<212> PRT

<213> Artificial Sequence

<220>

<223> chemically synthesized

<400> 17

Pro	Gly	Thr	Gly	Cys	Gly	Asp	Thr	Tyr	Arg	Pro	Lys	Arg	Pro	Thr	Thr
1				5					10					15	

Leu	Asn	Leu	Phe	Pro	Gln	Val	Pro	Arg	Ser	Gln	Asp	Thr
			20					25				

<210> 18

<211> 27

<212> PRT

<213> Artificial Sequence

<220>

21

<223> chemically synthesized

<400> 18

Ile	Pro	Ser	Pro	Ser	Val	Glu	Glu	Pro	His	Lys	His	Arg	Pro	Thr	Thr
1				5					10					15	

Leu	Arg	Leu	Thr	Thr	Leu	Gly	Ala	Gln	Asp	Ser
			20					25		

<210> 19

<211> 29

<212> PRT

<213> Artificial Sequence

<220>

<223> chemically synthesized

<400> 19

Gly	Ala	Tyr	Gly	Tyr	Ser	Asn	Pro	Lys	Ile	Leu	Lys	Gln	Ser	Met	Thr
1				5					10					15	

Leu	Asn	Leu	Ala	Asp	Pro	Val	Gly	Asn	Leu	Lys	Pro	His
			20					25				

<210> 20

<211> 29

<212> PRT

<213> Artificial Sequence

<220>

<223> chemically synthesized

<400> 20

Thr	Asn	Glu	Asp	His	Leu	Ala	Val	His	Lys	His	Lys	His	Glu	Met	Thr
1				5					10					15	

Leu	Lys	Phe	Gly	Pro	Ala	Arg	Asn	Asp	Ser	Val	Ile	Val
			20					25				

53493-1

CLAIMS:

1. A chimeric peptide, comprising a first domain and a second domain linked by a covalent bond, wherein the first domain comprises a trafficking sequence directing the chimeric peptide across the plasma membrane and/or to a desired cellular location, and the second domain comprises a retro-inverso isomer of a JNK inhibitor sequence, wherein said chimeric peptide is composed of D enantiomeric amino acids in the JNK inhibitor peptide of the second domain, and wherein the JNK inhibitor peptide is less than 280 amino acids in length, wherein the JNK inhibitor sequence comprises the amino acid sequence of SEQ ID NO: 3 or an amino acid sequence having a sequence identity of at least about 80% with the full length sequence of SEQ ID NO: 3.
2. The peptide of claim 1, wherein the JNK inhibitor sequence comprises an amino acid sequence having a sequence identity of at least about 95% with the full length sequence of SEQ ID NO: 3.
3. The peptide of claim 1 or 2, wherein the trafficking sequence comprises the amino acid sequence of SEQ ID NO: 7, 8, 9, or 10.
4. The peptide of claim 1, wherein the trafficking sequence augments cellular uptake of the peptide.
5. The peptide of any one of claims 1 to 4, wherein the trafficking sequence directs nuclear localization of the peptide.
6. The peptide of any one of claims 1 to 5, wherein the trafficking sequence comprises the amino acid sequence of a human immunodeficiency virus TAT polypeptide.
7. The peptide of any one of claims 1 to 6, wherein the JNK inhibitor sequence comprises the amino acid sequence of SEQ ID NO: 3.

53493-1

8. The peptide of claim 7, wherein the JNK inhibitor sequence binds JNK.
9. The peptide of claim 7 or 8, wherein the JNK inhibitor sequence inhibits activation of at least one JNK targeted transcription factor.
10. The peptide of claim 9, wherein the JNK targeted transcription factor is
5 C-Jun, ATF2, or Elk1.
11. The peptide of claim 7, wherein the JNK inhibitor sequence alters JNK induced effects when introduced into a JNK expressing cell.
12. The peptide of claim 11, wherein the JNK induced effect is induced by
10 restenosis, oncogenic transformation, maturation and differentiation of immune cells, proinflammatory cytokines, ionizing radiation as used in radiotherapy, ultraviolet light, free radicals, DNA damaging agents, chemotherapeutic drugs, ischemia, reperfusion, hypoxia, hypothermia, hyperthermia, apoptosis, a response to stressful stimuli, or a combination thereof.
13. The peptide of any one of claims 1 to 6, wherein said peptide comprises
15 the amino acid sequence of SEQ ID NO: 14 or a variant sequence of SEQ ID NO: 14 wherein the variant sequence has at least 80% sequence identity over amino acids 1 to 23 of SEQ ID NO: 14.
14. The peptide of any one of claims 1 to 6, wherein the peptide is less than 50 amino acids in length.
- 20 15. A composition, comprising the peptide of any one of claims 1 to 14, and a carrier.
16. Use of the peptide as defined in any one of claims 1 to 14, in the manufacture of a medicament for treating or preventing a pathophysiology associated with activation of JNK in a subject, wherein the pathophysiology is selected from: (1)
25 a malignancy selected from leukemia, and adenocarcinoma; (2) a non-malignant or immunologically-related cell proliferative disease selected from psoriasis, restenosis,

53493-1

vasculitis, pemphigus vulgaris, Behcet's syndrome, rheumatoid arthritis, acute respiratory distress syndrome (ARDS), ischemic heart disease, post-dialysis syndrome, acquired immunodeficiency syndrome, lipid histiocytosis, septic shock and apoptosis; (3) a pathological state induced by ionizing radiation, free radicals, DNA
 5 damaging agents or UV light; and (4) diabetes.

17. The use of claim 16, wherein the malignancy is a malignancy of the lung, breast, lymphoid, gastrointestinal, or genitourinary tract.

18. The use of claim 16, wherein the adenocarcinoma is colon cancer, renal-cell carcinoma, prostate cancer, non small cell carcinoma of the lung, cancer of
 10 the small intestine, cancer of the esophagus, or cancer with Bcr-Abl oncogenic transformation.

19. The use of any one of claims 16 to 18, wherein the peptide blocks activation of at least one JNK targeted transcription factor.

20. The use of claim 19, wherein the JNK targeted transcription factor is C-
 15 Jun, ATF2, or Elk1.

21. The use of claims 19, wherein the JNK targeted transcription factor is C-Jun.

22. The use of any one of claims 16 to 21, wherein the medicament is adapted for intraperitoneal, nasal, intravenous, oral or patch delivery.

20 23. Use of the peptide as defined in any one of claims 1 to 14, for treating or preventing a pathophysiology associated with activation of JNK in a subject, wherein the pathophysiology is selected from: (1) a malignancy selected from leukemia, and adenocarcinoma; (2) a non-malignant or immunologically-related cell proliferative disease selected from psoriasis, restenosis, vasculitis, pemphigus
 25 vulgaris, Behcet's syndrome, rheumatoid arthritis, acute respiratory distress syndrome (ARDS), ischemic heart disease, post-dialysis syndrome, acquired

53493-1

immunodeficiency syndrome, lipid histiocytosis, septic shock and apoptosis; (3) a pathological state induced by ionizing radiation, free radicals, DNA damaging agents or UV light; and (4) diabetes.

24. The use of claim 23, wherein the malignancy is a malignancy of the
5 lung, breast, lymphoid, gastrointestinal, or genitourinary tract.

25. The use of claim 23, wherein the adenocarcinoma is colon cancer, renal-cell carcinoma, prostate cancer, non small cell carcinoma of the lung, cancer of the small intestine, cancer of the esophagus, or cancer with Bcr-Abl oncogenic transformation.

10 26. The use of any one of claims 23 to 25, wherein the peptide blocks activation of at least one JNK targeted transcription factor.

27. The use of claim 26, wherein the JNK targeted transcription factor is C-Jun, ATF2, or Elk1.

28. The use of claim 26, wherein the JNK targeted transcription factor is
15 C-Jun.

29. The use of any one of claims 23 to 28, for intraperitoneal, nasal, intravenous, oral or patch delivery.

30. The composition of claim 15 for use in treating or preventing a pathophysiology associated with activation of JNK in a subject, wherein the
20 pathophysiology is selected from: (1) a malignancy selected from leukemia, and adenocarcinoma; (2) a non-malignant or immunologically-related cell proliferative disease selected from psoriasis, restenosis, vasculitis, pemphigus vulgaris, Behcet's syndrome, rheumatoid arthritis, acute respiratory distress syndrome (ARDS), ischemic heart disease, post-dialysis syndrome, acquired immunodeficiency
25 syndrome, lipid histiocytosis, septic shock and apoptosis; (3) a pathological state

53493-1

induced by ionizing radiation, free radicals, DNA damaging agents or UV light; and
(4) diabetes.

31. The composition of claim 30, for use in treating or preventing malignancies of the lung, breast, lymphoid, gastrointestinal, or genitourinary tract.

5 32. The composition of claim 30, for use in treating or preventing adenocarcinoma of colon cancer, renal-cell carcinoma, prostate cancer, non small cell carcinoma of the lung, cancer of the small intestine, cancer of the esophagus, or cancer with Bcr-Abl oncogenic transformation.

10 33. The composition of any one of claims 30 to 32, for use in blocking activation of at least one JNK targeted transcription factor.

34. The composition of claim 33, for use in blocking the JNK targeted transcription factor C-Jun, ATF2, or Elk1.

35. The composition of claim 33, for use in blocking the JNK targeted transcription factor C-Jun.


15 36. The composition of any one of claims 30 to 35 for use in intraperitoneal, nasal, intravenous, oral or patch delivery.

FIG. 2 Generic Sequences, Human, Mouse and Rat

L-TAT-IB	:	NH ₂ -	XXXXXXXXXXRRRQRRXXXXXXXXXXRRPTFLXXXXXXXXQDS/TX	-COOH
D-TAT	:	NH ₂ -	XXXXXXXXXXRRRQRRXXXXX	-COOH
D-TAT-IB	:	NH ₂ -	XT/SDQXXXXXXXXXLTLTPRXXXXXXXXXXRRRQRRKKRXXXXXXXXX	-COOH

3/ 9

FIG. 3
Effects of GFP-JBD₂₃ Construct on Pancreatic β -Cell Apoptosis



4/9

FIG. 4a

Effects of TAT-IB Peptides on JNK Mediated Phosphorylation






FIG. 4b

Effects of TAT-IB Peptides on JNK Mediated Phosphorylation



5/9

FIG. 5 *L-TAT -IB Inhibition of Phosphorylation
By Recombinant JNKs*



6/9



FIG. 6*In Vivo Inhibition of c-Jun Phosphorylation by TAT-IB Peptides***FIG. 11***Suppression of JNK Transcription Factor Phosphorylation by L-TAT-IB1 Peptides*

FIG. 8
*The D-TAT-IB1 peptide decreases
IL-1-induced apoptosis in β TC-3 cells*




FIG. 7
*Inhibition of IL-1 β Induced Pancreatic
 β -Cell Death by TAT-IB Peptides*




FIG. 9
The D-TAT-IB1 peptide confers long-term
protection (15 days) against
IL-1-induced apoptosis in β TTC-3 cells






FIG. 10
L-TAT-IB1 and D-TAT-IB1 peptides
prevent IR-induced apoptosis
in a human colon cancer cell line



9/9

FIG. 12 Radioprotection to Ionizing Radiation by TAT-IB1 Peptides



Effects of TAT-IB Peptides on JNK Mediated Phosphorylation

