



(86) Date de dépôt PCT/PCT Filing Date: 2001/04/28

(87) Date publication PCT/PCT Publication Date: 2001/11/08

(85) Entrée phase nationale/National Entry: 2002/10/24

(86) N° demande PCT/PCT Application No.: US 2001/013932

(87) N° publication PCT/PCT Publication No.: 2001/083529

(30) Priorité/Priority: 2000/04/28 (60/200,298) US

(51) Cl.Int.⁷/Int.Cl.⁷ C07K 14/705, A61K 38/00


(71) Demandeur/Applicant:
PLANET BIOTECHNOLOGY, INC., US

(72) Inventeurs/Inventors:
LARRICK, JAMES WILLIAM, US;
WYCOFF, KEITH LYNN, US

(74) Agent: SMART & BIGGAR


(54) Titre : NOUVELLE IMMUNOADHESINE DESTINEE A LA PREVENTION D'INFECTIONS A RHINOVIRUS

(54) Title: IMMUNOADHESIN FOR THE PREVENTION OF RHINOVIRUS INFECTION



(57) Abrégé/Abstract:

The immunoadhesions of the present invention are useful in treating rhinovirus infections. The immunoadhesions contain a chimeric ICAM molecule and may optionally also contain J chain and secretory compounds. The chimeric ICAM molecule is a fusion protein that has a rhinovirus receptor protein linked to an immunoglobulin protein. This invention also includes the greatly increased and improved method of producing immunoadhesions in plants. Each of the components of an immunoadhesin is produced in a plant cell and thereby assembles within the plant cell. This method of producing the immunoadhesions of the present invention results in the efficient and economic production of these molecules. The present invention also contemplates the production of immunoadhesions in a variety of eukaryotic cells including plants and mammalian cells. The immunoadhesions of the present invention are useful as a therapeutic against the common cold in humans which is caused by rhinoviruses.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
8 November 2001 (08.11.2001)

PCT

(10) International Publication Number
WO 01/083529 A3(51) International Patent Classification⁷: **C07K 14/705**,
A61K 38/00(74) Agents: **GUISE, Jeffrey, W.** et al.; Brobeck, Phleger &
Harrison LLP, 12390 El Camino Real, San Diego, CA
92130 (US).

(21) International Application Number: PCT/US01/13932

(22) International Filing Date: 28 April 2001 (28.04.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/200,298 28 April 2000 (28.04.2000) US(71) Applicant (for all designated States except US): **PLANET
BIOTECHNOLOGY, INC.** [US/US]; 25571 Clawiter
Road, Hayward, CA 94545 (US).(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **LARRICK, James,
William** [US/US]; Star Route Box #48, Woodside, CA
94062 (US). **WYCOFF, Keith, Lynn** [US/US]; 2399
Carmel Drive, Palo Alto, CA 94303 (US).

Published:


— with international search report

(88) Date of publication of the international search report:

12 September 2002

[Continued on next page]

(54) Title: IMMUNOADHESIN FOR THE PREVENTION OF RHINOVIRUS INFECTION



(57) **Abstract:** The immunoadhesions of the present invention are useful in treating rhinovirus infections. The immunoadhesions contain a chimeric ICAM molecule and may optionally also contain J chain and secretory compounds. The chimeric ICAM molecule is a fusion protein that has a rhinovirus receptor protein linked to an immunoglobulin protein. This invention also includes the greatly increased and improved method of producing immunoadhesions in plants. Each of the components of an immunoadhesin is produced in a plant cell and thereby assembles within the plant cell. This method of producing the immunoadhesions of the present invention results in the efficient and economic production of these molecules. The present invention also contemplates the production of immunoadhesions in a variety of eukaryotic cells including plants and mammalian cells. The immunoadhesions of the present invention are useful as a therapeutic against the common cold in humans which is caused by rhinoviruses.



WO 01/083529 A3

WO 01/083529 A3



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**NOVEL IMMUNOADHESIN FOR THE
PREVENTION OF RHINOVIRUS INFECTION**

PRIORITY

This application claims benefit under §119(e) of U.S. *Provisional* Patent Application No. 60/200,298, filed April 28, 2000, entitled NOVEL IMMUNOADHESIN FOR THE PREVENTION OF RHINOVIRUS INFECTION, and naming J. W. Larrick and K. L. Wycoff as inventors. This application is incorporated herein by reference in its entirety and for all purposes.

**STATEMENT REGARDING FEDERALLY SPONSORED
RESEARCH OR DEVELOPMENT**

Federal research support was provided in the form of an SBIR Phase I grant (R43 AI43122) and SBIR Phase II grant (2R44AI43122-02).

FIELD OF THE INVENTION

The present invention relates to immunoadhesins, fusions of the human rhinovirus receptor protein and immunoglobulin, and the expression of immunoadhesins in plants. The therapeutic use of immunoadhesins for the prevention and treatment of human rhinovirus infection is also contemplated.

BACKGROUND TO THE INVENTION

The common cold is generally a relatively mild disease, however significant complications resulting from colds, such as otitis media, sinusitis and asthma exacerbations are common. Human rhinoviruses (HRV) cause up to 50% of all adult colds and 25% of colds in children (Bella and Rossmann, *J Struct Biol.* 128:69-74, 1999, and Sperber and Hayden, *Antimicrob Agents Chemother.* 32:409-19, 1988). The cost to society runs into billions of dollar per year. These small, nonenveloped RNA viruses represent a subgroup of

picornavirus (Rueckert, *Virology*, pp. 507-548, eds. Fields, *et al.*, Raven Press, Ltd. New York, 1990) X-ray crystallography of rhinovirus identified a capsid 300 Å in diameter (1 Å = 0.1 nm) with icosahedral symmetry, constructed from sixty copies each of the viral coat proteins VP1, VP2, and VP3 (Rossmann, *Nature* 317:145-153, 1985). A surface depression or "canyon" on HRV was suggested as the receptor binding site (Colonno, *et al.*, *Proc Natl Acad Sci U S A.* 85:5449-5453, 1985; Rossmann, *et al.* *Nature* 317:145-153, 1985). Of the 102 characterized HRV serotypes, 91 (known as the major group) share as their receptor a cell surface glycoprotein known as intercellular adhesion molecule-1 (ICAM-1) (Greve, *et al.*, *Cell* 56:839-847, 1989; Staunton, *et al.*, *Cell* 56:849-853, 1989); the binding site is located within N-terminal domain 1 (Greve, *et al.*, *J Virol.* 65:6015-6023, 1991; Staunton, *et al.*, *Cell* 61:243-254, 1990).

ICAM-1 is a membrane protein with five extracellular domains, a hydrophobic transmembrane domain, and a short cytoplasmic domain. ICAM-1 is expressed on many cells important in immune and inflammatory responses, and is inducible on others (Casasnovas, *et al.*, *Proc Natl Acad Sci U S A.* 95:4134-9, 1998). ICAM-1 functions as a ligand for the leukocyte integrins LFA-1 and Mac-1 (Springer, *Cell.* 76:301-14, 1994; Staunton *et al.*, *Cell* 61:243-254, 1990). On the cell surface, ICAM-1 is primarily a dimer due to association of the transmembrane domains (Miller, *et al.*, *J Exp Med.* 182:1231-41, 1995; Reilly, *et al J Immunol.* 155:529-32, 1995).

Recombinant, soluble forms of ICAM-1 (sICAM-1) consisting of the five extracellular domains were shown to be effective in blocking rhinovirus infection of human cells *in vitro* (Greve, *et al.*, *J Virol.* 65:6015-6023, 1991; Marlin, *et al.*, *Nature.* 344:70-2, 1990). Evaluation of sICAM-1 activity against a spectrum of laboratory strains and field isolates showed that all major strains of HRV are sensitive to sICAM-1. Minor strains, which do not use ICAM as a receptor, were unaffected by sICAM-1 (Crump *et al.*, *Antiviral Chem Chemother.* 4:323-327, 1993; Ohlin, *et al.*, *Antimicrob Agents Chemother.* 38:1413-5, 1994).

The anti-viral activity of soluble ICAM-1 *in vitro* appears to be mediated by more than one mechanism. These mechanisms include competition with cell-surface ICAM-1 for binding sites, interference with virus entry or uncoating, and direct inactivation by premature release of viral RNA and formation of empty capsids (Arruda, *et al.*, *Antimicrob Agents Chemother.* 36:1186-1191, 1992; Greve, *et al.*, *J Virol.* 65:6015-6023, 1991; Marlin, *et al.*, *Nature* 344:70-2, 1990; Martin *et al.*, *J Virol.* 67:3561-8, 1993).

The host range of HRV is restricted to primates. A recent study showed that soluble ICAM-1 was effective in preventing rhinovirus infection in chimpanzees (Huguenel, *et al.*, *Am J Respir Crit Care Med.* 155:1206-10, 1997). Although chimpanzees do not show clinical symptoms, infection was demonstrated by measuring seroconversion and virus shedding. A single dose of 10 mg of soluble ICAM-1 as an intranasal spray was effective at preventing infection by HRV-16 when co-administered with HRV, or when the virus was administered ten minutes later.

A human clinical trial with soluble ICAM-1 showed that it reduced the severity of experimental HRV colds (Turner, *et al.*, *JAMA* 281:1797-804, 1999). In this trial a total of 196 subjects received either soluble ICAM-1 or placebo in various formulations. Some subjects were given soluble ICAM-1 or placebo starting seven hours before inoculation with HRV 39 and others were started twelve hours after virus inoculation. Medications were administered as either an intranasal solution or powder, given in six daily doses for seven days (a total of 4.4 mg per day). In this study, soluble ICAM-1 did not prevent infection, as measured by either virus isolation or seroconversion (infection rate of 92% for placebo-treated vs. 85% of soluble ICAM-1 treated). However, soluble ICAM-1 did have an impact on all measures of illness. The total symptom score was reduced by 45%, the proportion of subjects with clinical colds was reduced 23% and nasal mucus weight was reduced by 56%. There was not a significant difference between the use of powder or solution formulations, or between pre- and post-inoculation groups. Treatment with soluble ICAM-1 did not result in any adverse effects or evidence of absorption through the nasal

mucosa. Also, there was no inhibition of the development of anti-HRV type-specific antibodies.

As discussed, ICAM-1 is dimeric on the cell surface. Martin *et al.*, in *J Virol.* 67:3561-8, (1993) first proposed that multivalent binding to HRV by a multimeric soluble
5 ICAM might result in a higher effective affinity, termed avidity, and thus facilitate uncoating of the virus. They constructed multivalent, ICAM-1/immunoglobulin molecules, postulating that these would be more effective than monovalent soluble ICAM-1 in neutralizing HRV and thus would have increased therapeutic utility. These ICAM-1/immunoglobulin molecules included ICAM-1 amino-terminal domains 1 and 2 fused to the hinge and constant domains of
10 the heavy chains of IgA1 (IC1-2D/IgA), IgM (IC1-2D/IgM) and IgG1(IC1-2D/IgG). In addition, five extracellular domains were fused to IgA1 (IC1-5D/IgA). These ICAM-1/immunoglobulin molecules were compared with soluble forms of ICAM-1 having two (sIC1-2D) and five (sIC1-5D) domains in assays of HRV binding, infectivity and conformation. The ICAM-1/IgA immunoglobulin (IC1-5D/IgA) was 200 times, and the
15 ICAM-1/IgM immunoglobulin (IC1-2D/IgM) and ICAM-1/IgG immunoglobulin molecules (IC1-2D/IgG) were 25 and 10 times, more effective than soluble ICAM-1. These molecules were highly effective in inhibiting rhinovirus binding to cells and disrupting the conformation of the virus capsid. The ICAM-1/IgA immunoglobulin molecules were effective in the nanomolar concentration range. Comparison of IC1-2D/IgA and IC1-2D/IgG showed that the
20 class of Ig constant region used had a large impact on efficacy.

A subsequent study compared the inhibitory activities of soluble ICAM-1 and IC1-5D/IgA against nine major HRV serotypes and a variant of HRV-39 selected for moderate resistance to soluble ICAM-1 (Crump, *et al.*, *Antimicrob Agents Chemother.* 38:1425-7, 1993). IC1-5D/IgA was more potent than monomeric soluble ICAM-1 by 50 to
25 143 times on a weight basis and by 60 to 170 times on a molar basis against the standard serotypes. The HRV-39 variant was 38-fold more resistant to soluble ICAM-1 than the wild-type, and it was only 5-fold more resistant to IC1-5D/IgA. This is consistent with the hypothesis that virus escape from inhibition by multivalent molecules would be expected to

occur at lower frequency than virus escape from inhibition by monomeric soluble receptor (Martin, *et al.*, *J Virol.* 67:3561-8, 1993). An assay designed to measure viral inactivation showed that HRV-39 and HRV-13 were not directly inactivated to a significant extent by soluble ICAM-1 ($<0.5 \log_{10}$ reduction in infectivity). However, incubation with IC1-5D/IgA
5 resulted in a reduction of infectivity of these same viruses by about $1.0 \log_{10}$ (Crump, *et al.*, *Antimicrob Agents Chemother.* 38:1425-7, 1994). Results by Martin *et al.* (*J Virol.* 67:3561-8, 1993) suggest that the greater the valence, the greater the effectiveness of the molecules. Dimeric and decameric forms of IC1-2/IgM were separable by sucrose gradient sedimentation. The decameric form was five times more effective than the dimeric form at
10 blocking binding of HRV to HeLa cells.

The ICAM-1/immunoglobulin molecules that have been described suffer from several drawbacks, including the laborious production techniques and high costs associated with those production methods. In addition, the previously described ICAM-1/immunoglobulin molecules have limited stability as multimers in the harsh environment in which the molecule
15 must inactivate rhinoviruses.

The immunoadhesins of the present invention have significant advantages over what has been described in the art. The immunoadhesins of the present invention that are expressed in plants would be tetrameric, rather than only dimeric. Immunoadhesins having multiple binding sites have a higher effective affinity for the virus, thereby increasing the
20 effectiveness of the immunoadhesin. In addition, the association of secretory component and immunoglobulin J chain with the immunoadhesin of the present invention increases the stability of the immunoadhesin in the mucosal environment (Corthesy, *Biochem Soc Trans.* 25:471-475, 1997). Secretory IgA, which is associated with secretory component, is the antibody isotype normally found in mucosal secretions, including milk and colostrum. Unlike
25 other antibody isotypes, SIgA can pass through the gut with very little proteolytic degradation. It also is very stable in crude plant preparations at room temperature. A function of the secretory component appears to be to protect the antibody from the harsh environment of the mucosa (Paul, *Fundamental Immunology*, Raven Press, NY, Third

Edition, pp. 303-304, 1993). Furthermore, the immunoadhesin of the present invention are significantly less expensive to produce in plants than in animal cell culture, and production in plants would make it safer for human use, since plants are not known to harbor any animal viruses.

5

SUMMARY OF THE INVENTION

The present invention contemplates an immunoadhesin comprising a chimeric ICAM-1 molecule having a rhinovirus receptor protein linked to at least a portion of an immunoglobulin heavy chain, wherein J chain and secretory component are associated with the chimeric ICAM-1 molecule.

10

In preferred embodiments, the immunoadhesin of the present invention is comprised of a rhinovirus receptor protein made of any combination of extracellular domains 1, 2, 3, 4 and 5 of the rhinovirus receptor protein, ICAM-1, linked to an immunoglobulin heavy chain. Also contemplated by the present invention are immunoadhesins of the present invention in which the immunoglobulin is IgA, IgA₁, IgA₂, IgG₁, IgG₂, IgG₃, IgG₄, IgM, IgD, IgE or a chimeric immunoglobulin heavy chain made up of domains or segments from different immunoglobulin isotypes.

15

In other preferred embodiments of the present invention, the immunoadhesin comprises multiple chimeric ICAM-1 molecules associated with J chain and secretory component. The increase in valency results in a higher effective affinity for the rhinovirus, thereby increasing the effectiveness of the immunoadhesin.

20

In a preferred embodiment of the present invention, all proteins used to make the immunoadhesin of the present invention are human proteins. In addition to production in plants or plant cells, the present invention contemplates an immunoadhesin expressed in mammalian cells, hairy root cultures, plant cells in tissue culture, and heterologous cells derived from plants, vertebrates or invertebrates.

25

In preferred embodiments of the present invention, the immunoadhesins are expressed, in plants, including monocotyledonous plants and dicotyledonous plants as a part of the plants genome. Expression in plants, as opposed to expression in cultured cells, allows for a significant reduction in the cost of producing the immunoadhesin.

5 The present invention contemplates an immunoadhesin having plant-specific glycosylation. A gene coding for a polypeptide having within its amino acid sequence, the glycosylation signal asparagine-X-serine/threonine, where X can be any amino acid residue, is glycosylated via oligosaccharides linked to the asparagine residue of the sequence when expressed in a plant cell. See Marshall, *Ann. Rev. Biochem.*, 41:673 (1972) and Marshall,
10 *Biochem. Soc. Symp.*, 40:17 (1974) for a general review of the polypeptide sequences that function as glycosylation signals. These signals are recognized in both mammalian and in plant cells. At the end of their maturation, proteins expressed in plants or plant cells have a different pattern of glycosylation than do proteins expressed in other types of cells, including mammalian cells and insect cells. Detailed studies characterizing plant-specific glycosylation
15 and comparing it with glycosylation in other cell types have been performed, for example, in studies described by Cabanes-Macheteau *et al.*, *Glycobiology* 9(4):365-372 (1999), and Altmann, *Glycoconjugate J.* 14:643-646 (1997). These groups and others have shown that plant-specific glycosylation generates glycans that have xylose linked $\beta(1,2)$ to mannose, but xylose is not linked $\beta(1,2)$ to mannose as a result of glycosylation in mammalian and insect
20 cells. Plant-specific glycosylation results in a fucose linked $\alpha(1,3)$ to the proximal GlcNAc, while glycosylation in mammalian cells results in a fucoselinked $\alpha(1,6)$ to the proximal GlcNAc. Furthermore, plant-specific glycosylation does not result in the addition of a sialic acid to the terminus of the protein glycan, whereas in glycosylation in mammalian cells, sialic acid is added.

25 In other embodiments, the immunoadhesin of the present invention is part of a composition comprising plant material and the immunoadhesin, associated with J chain and secretory component. The plant material present may be plant cell walls, plant organelles, plant cytoplasm, intact plant cells, viable plants, and the like. The particular plant materials

or plant macromolecules that may be present include ribulose biphosphate carboxylase, light harvesting complex, pigments, secondary metabolites or chlorophyll. Compositions of the present invention may have an immunoadhesin concentration of between 0.001% and 99.9% mass excluding water. In other embodiments, the immunoadhesin is present in a concentration of 0.01% to 99% mass excluding water. In other embodiments, the compositions of the present invention have plant material or plant macromolecules present at a concentration of 0.01% to 99% mass excluding water.

The present invention also contemplates methods for the treatment or prevention of human rhinovirus infection in a subject, including reducing the infection by human rhinovirus of host cells susceptible to infection by the virus, or reducing the initiation or spread of the common cold due to human rhinovirus, by a method comprising contacting the virus with an immunoadhesin of the present invention, wherein the immunoadhesin binds to the human rhinovirus and reduces infectivity. The immunoadhesin could mediate infection by competition with cell-surface ICAM-1 for binding sites, interference with virus entry or uncoating, and/or direct inactivation by premature release of viral RNA and formation of empty capsids (Arruda, *et al.*, *Antimicrob. Agents Chemother.* 36:1186-1191, 1992; Greve, *et al.*, *J. Virol.* 65:6015-6023, 1991; Martin, *et al.*, *Nature* 344:70-2, 1990; Martin, *et al.*, *J. Virol.* 67:3561-8, 1993). In another embodiment, human rhinovirus infection in a subject is treated by a method comprising intranasally administering to the subject an effective amount of an immunoadhesin of the present invention, wherein the immunoadhesin reduces human rhinovirus infectivity.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 illustrates pSSPHuA2, vector in which DNAs encoding a chimeric ICAM-1 molecule containing the first five domains of human ICAM-1 and the Fc region of human IgA2m(2) were fused. This vector contains the SuperMas promoter for driving the expression of a signal peptide and the constant regions of the human IgA2m(2) heavy-chain. Sequences encoding ICAM domains 1-5 were amplified, by PCR, to contain convenient restriction sites

(5' SpeI and 3' Spe I) for insertion between the signal peptide and the C α 1 domain. DNA encoding an ER retention signal (RSEKDEL) was appended to the 3' end of the heavy-chain in order to boost the expression level of the construct.

FIG. 2 illustrates a chimeric ICAM molecule. **2A** shows the DNA expression cassette from which the chimeric ICAM-1 molecule was produced. **2B** shows the amino acid sequence, after signal peptide cleavage, of the mature form of the fusion protein. Amino acids introduced by the cloning procedure are underlined and mark the junction between the five extracellular domains of ICAM-1 and the C α 1-C α 3 domains of the IgA2m(2) heavy chain. The bolded N's indicate the fifteen potential glycosylation sites.

FIG. 3 illustrates the expression of the immunoadhesin in independently transformed tobacco calli. **3A** shows immunoblots of non-reducing SDS-polyacrylamide gels on which samples containing different transformed tobacco calli (**C**) and aqueous extracts (**Aq**) were run and probed for the presence of human ICAM. The molecular weight markers are indicated, and the reference standard (**R**) was a mixture (~75 ng each) of human ICAM (~75 kD) and human SigA (>>250 kD). **3B** shows immunoblots of nonreducing SDS-polyacrylamide gels containing various fractions of partially purified immunoadhesin from callus Rhi107-11. The purification fractions analyzed were juice (**J**), G-100 fraction (**G**), sterile filtered G-100 fraction (**SG**), and a mixture of reference standards of human SigA (75 ng) and human ICAM-1 (75ng) (**RS**).

Blots were probed with antibodies against human ICAM (~**ICAM**), human IgA heavy chain (~ α), human secretory component (~**SC**) and human J chain (~**J**). Secondary, enzyme-conjugated antibodies were employed as necessary to label immuno-positive bands with alkaline phosphatase.

FIG. 4 illustrates the results of an enzyme-linked immunosorbent assay (ELISA) showing competition between plant extract and soluble ICAM-1 for binding to an anti-ICAM mAb. For the assay, 96-well plates were coated with 0.25 μ g soluble ICAM-1/ml. The squares represent the increasing concentrations of sICAM and the circles represent the

increasing amounts of callus extract (sterile filtered fraction from G-100) used to compete with the adhered ICAM for a constant amount of a mouse (anti-human ICAM) antibody.

FIG. 5 illustrates the results of an assay showing the ability of an immunoadhesin to inhibit human rhinovirus killing of HeLa cells (cytopathic effect, or CPE, assay). **5A** shows the results of an assay comparing the CPE of human rhinovirus on HeLa cells in the presence of partially purified extracts containing either the immunoadhesin in the ICAM-Fc fusion (IC1-5D/IgA) or containing an antibody against doxorubicin. (The right side-up and upside-down triangles represent two extracts derived from Rhi107-11, containing the immunoadhesin.) **5B** shows the results of an assay comparing the CPE of human rhinovirus on HeLa cells in the presence of soluble human ICAM-1 or an extract from the immunoadhesin in the ICAM-Fc fusion (IC1-5D/IgA). The Inset shows scale expansion in the range of the IC50 for soluble ICAM (1.35 µg/ml) and for IC1-5D/IgA (0.12 µg/ml; 11.3 fold-less).

FIG. 6 shows an evaluation of the production necessities for making 1 gram of finished immunoadhesin. In this diagram, the number of plants needed for 1 g of immunoadhesin, at 20% yield, at expected levels of expression and plant weight is illustrated. At different levels of immunoadhesin expression (mg/kg fresh weight) and overall recovery (set at 20%), the weight of each plant, and so the total number of plants, may be determined for a specified production target (1 g/harvest) within a window (dotted square) of reasonable possibilities. The number of required plants decreases, inversely, with the number of specified growth and re-growth periods. The expected biomass production, a function of time and growth conditions, influences the time to harvest and the time between harvests. These growth periods can be adjusted to the realities of the purification schedule by staggering planting and harvesting dates.

FIG. 7 shows the coding and amino acid sequences of each of the immunoglobulin genes and proteins listed in Table 2.

FIG. 8 shows the sequences of plasmids used to transform plants, as described in Example 2, for use in studies of the expression of immunoadhesins of the present invention.

FIG. 8 A shows the nucleotide and protein sequences for plasmids PSSpICAMHuA2

FIG 8 B shows the nucleotide protein sequences for the bean legumin signal peptide.

5 **FIG 8 C** shows the nucleotide and amino acid sequence of the protein coding region of pSHuJ.

FIG 8 D shows the nucleotide and amino acid sequence of protein coding region of pSHuSC.

FIG 8 E shows the nucleotide sequence of plasmids pBMSP-1.

10 **FIG 8 F** shows the nucleotide sequence of plasmids pBMSP-1spJSC.

FIG. 9 contains nucleotide and protein sequences SEQ ID NO: 1; SEQ ID NO: 2; SEQ ID NO: 3; SEQ ID NO: 4; SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7; SEQ ID NO: 8, for ICAM-1, and human IgA₂ and other nucleotide sequences.

DETAILED DESCRIPTION OF THE INVENTION

15 **A. Definitions.**

As used herein, the following abbreviations and terms include, but are not necessarily limited to, the following definitions.

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of immunology, molecular biology, microbiology, cell biology and
20 recombinant DNA, which are within the skill of the art. See, e.g., Sambrook, *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd edition (1989); *Current Protocols In Molecular Biology* (F.M. Ausubel, *et al.* eds., (1987)); the series *Methods In Enzymology* (Academic Press, Inc.); M.J. MacPherson, *et al.*, eds. *Pcr 2: A Practical Approach* (1995);

Harlow and Lane, eds, *Antibodies: A Laboratory Manual* (1988), and H. Jones, *Methods In Molecular Biology vol. 49, "Plant Gene Transfer And Expression Protocols"* (1995).

Immunoglobulin molecule or Antibody. A polypeptide or multimeric protein containing the immunologically active portions of an immunoglobulin heavy chain and immunoglobulin light chain covalently coupled together and capable of specifically combining with antigen. The immunoglobulins or antibody molecules are a large family of molecules that include several types of molecules such as IgD, IgG, IgA, secretory IgA (SIgA), IgM, and IgE.

Construct or Vector. An artificially assembled DNA segment to be transferred into a target plant tissue or cell. Typically, the construct will include the gene or genes of a particular interest, a marker gene and appropriate control sequences. The term "plasmid" refers to an autonomous, self-replicating extrachromosomal DNA molecule. In a preferred embodiment, the plasmid constructs of the present invention contain sequences coding for heavy and light chains of an antibody. Plasmid constructs containing suitable regulatory elements are also referred to as "expression cassettes." In a preferred embodiment, a plasmid construct can also contain a screening or selectable marker, for example an antibiotic resistance gene.

Selectable marker. A gene that encodes a product that allows the growth of transgenic tissue on a selective medium. Non-limiting examples of selectable markers include genes encoding for antibiotic resistance, e.g., ampicillin, kanamycin, or the like. Other selectable markers will be known to those of skill in the art.

Transgenic plant. Genetically engineered plant or progeny of genetically engineered plants. The transgenic plant usually contains material from at least one unrelated organism, such as a virus, another plant or animal.

Chimeric ICAM-1 molecule: The fusion of any combination of the extracellular domains 1, 2, 3, 4 and 5 of ICAM-1 with at least a part of an immunoglobulin heavy chain

protein, made by linking ICAM-1 sequence upstream of an immunoglobulin heavy chain gene sequence and expressing the encoded protein from the construct.

Chimeric immunoglobulin heavy chain: An immunoglobulin derived heavy chain having at least a portion of its amino acid sequence derived from an immunoglobulin heavy chain of a different isotype or subtype or some other peptide, polypeptide or protein. Typically, a chimeric immunoglobulin heavy chain has its amino acid residue sequence derived from at least two different isotypes or subtypes of immunoglobulin heavy chain.

Dicotyledonous plants (dicots): Flowering plants whose embryos have two seed halves or cotyledons. Examples of dicots are: tobacco; tomato; the legumes including alfalfa; oaks; maples; roses; mints; squashes; daisies; walnuts; cacti; violets and buttercups.

Effective amount: An effective amount of an immunoadhesin of the present invention is sufficient to detectably inhibit rhinovirus infection, cytotoxicity or replication; or to reduce the severity or length of rhinovirus infection.

Human rhinovirus (HRV): A nonenveloped RNA virus representing a subgroup of picornavirus, that is a major cause of the common cold in humans. Rhinoviruses are described in Rhinoviruses, Reoviruses, and Parvoviruses, pp. 1057-1059, *Zinsser Microbiology*, Joklik *et al.*, eds. Appleton and Lange (1992).

Immunoadhesin : A complex containing a chimeric ICAM-1 molecule, and optionally containing secretory component, and J chain.

Immunoglobulin heavy chain: A polypeptide that contains at least a portion of the antigen binding domain of an immunoglobulin and at least a portion of a variable region of an immunoglobulin heavy chain or at least a portion of a constant region of an immunoglobulin heavy chain. Thus, the immunoglobulin derived heavy chain has significant regions of amino acid sequence homology with a member of the immunoglobulin gene superfamily. For example, the heavy chain in an Fab fragment is an immunoglobulin-derived heavy chain.

Immunoglobulin light chain: A polypeptide that contains at least a portion of the antigen binding domain of an immunoglobulin and at least a portion of the variable region or at least a portion of a constant region of an immunoglobulin light chain. Thus, the immunoglobulin-derived light chain has significant regions of amino acid homology with a member of the immunoglobulin gene superfamily.

Immunoglobulin molecule: A protein containing the immunologically-active portions of an immunoglobulin heavy chain and immunoglobulin light chain covalently coupled together and capable of specifically combining with antigen.

ICAM-1: Intercellular adhesion molecule-1. In humans, ICAM-1 functions as the receptor for human rhinovirus.

J chain: A polypeptide that is involved in the polymerization of immunoglobulins and transport of polymerized immunoglobulins through epithelial cells. See, *The Immunoglobulin Helper: The J Chain in Immunoglobulin Genes*, at pg. 345, Academic Press (1989). J chain is found in pentameric IgM and dimeric IgA and typically attached via disulphide bonds. J chain has been studied in both mouse and human.

Monocotyledonous plants (monocots): Flowering plants whose embryos have one cotyledon or seed leaf. Examples of monocots are: lilies; grasses; corn; grains, including oats, wheat and barley; orchids; irises; onions and palms.

Glycosylation: The modification of a protein by oligosaccharides. See, Marshall, *Ann. Rev. Biochem.*, 41:673 (1972) and Marshall, *Biochem. Soc. Symp.*, 40:17 (1974) for a general review of the polypeptide sequences that function as glycosylation signals. These signals are recognized in both mammalian and in plant cells.

Plant-specific glycosylation: The glycosylation pattern found on plant-expressed proteins, which is different from that found in proteins made in mammalian or insect cells. Proteins expressed in plants or plant cells have a different pattern of glycosylation than do proteins expressed in other types of cells, including mammalian cells and insect cells.

Detailed studies characterizing plant-specific glycosylation and comparing it with glycosylation in other cell types have been performed by Cabanes-Macheteau *et al.*, *Glycobiology* 9(4):365-372 (1999), Lerouge *et al.*, *Plant Molecular Biology* 38:31-48 (1998) and Altmann, *Glycoconjugate J.* 14:643-646 (1997). Plant-specific glycosylation generates
5 glycans that have xylose linked $\beta(1,2)$ to mannose. Neither mammalian nor insect glycosylation generate xylose linked $\beta(1,2)$ to mannose. Plants do not have a sialic acid linked to the terminus of the glycan, whereas mammalian cells do. In addition, plant-specific glycosylation results in a fucose linked $\alpha(1,3)$ to the proximal GlcNAc, while glycosylation in mammalian cells results in a fucose linked $\alpha(1,6)$ to the proximal GlcNAc.

10 Secretory component (SC): A component of secretory immunoglobulins that helps to protect the immunoglobulin against inactivating agents thereby increasing the biological effectiveness of secretory immunoglobulin. The secretory component may be from any mammal or rodent including mouse or human.

15 sICAM: A naturally-occurring soluble-truncated form of ICAM-1 lacking both the hydrophobic transmembrane domain and the carboxy-terminal cytoplasmic domain of ICAM.

The articles, patents and patent applications cited in this document are incorporated into this document as if set forth in full.

B. Immunoadhesins Containing Chimeric ICAM Molecules.

The present invention provides novel methods for producing immunoadhesin
20 molecules containing chimeric ICAM molecules. The immunoadhesins of the present invention contain chimeric ICAM-1 molecules made up of a rhinovirus receptor protein linked to a portion of an immunoglobulin heavy chain molecule in association with J chain and secretory component. The chimeric ICAM-1 molecules of the present invention contain two molecules derived from different sources: a rhinovirus receptor protein portion and an
25 immunoglobulin chain portion. The rhinovirus receptor protein of the present invention is derived from the intercellular adhesion molecule 1 (ICAM-1). The nucleotide sequence for

the human rhinovirus receptor ICAM-1 has been determined and characterized by Staunton, *et al.*, *Cell* 52:925-933 (1988); Greve, *et al.* *Cell* 56:839-847 (1989); Greve, *et al.* *J. Virology* 65:6015-6023 (1991); Staunton, *et al.*, *Cell*, 61:243-254 (1990) and described in Sequence ID No. 3 and GenBank accession no. M24283.

5 The ICAM-1 molecule is a membrane protein (SEQ ID NOS: 1 and 2) that has 5 extracellular domains, a hydrophobic transmembrane domain and a short cytoplasmic domain. These features have been described by Casasnovas, *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 95:4134-4139 (1998) and Staunton, *et al.*, *Cell* 52:925-933 (1988). Of particular use in the present invention are the domains of the ICAM-1 molecule that are responsible for the
10 binding of human rhinoviruses which have been localized to the N-terminal domains 1 and 2 (Greve, *et al.*, *J. Virol.*, 65:6015-6023 1991, and Staunton, *et al.*, *Cell*, 61:243-245 1990. The present invention also contemplates rhinovirus receptor protein portions which include any combination of extracellular domains 1, 2, 3, 4, and 5 of the ICAM-1 molecule. In particular preferred embodiments, the rhinovirus receptor protein portion includes domains 1 and 2 of
15 the ICAM-1 molecule and in other preferred embodiments domains 1, 2, 3, 4 and 5 of the ICAM-1 molecule are present.

The boundaries of the 5 extracellular domains are well known in the art and described in Staunton, *et al.*, *Cell* 52:925-933 (1988). The approximated domain boundaries are shown in Table 1 below.

20

Table 1

<u>ICAM-1 Domains</u>	<u>Amino Acids</u>
1	1-88
2	89-105
3	106-284
4	285-385
5	386-453

As used in the present invention, the ICAM-1 domain 1 is from about residue 1 to about residue 88; domain 2 is from about residue 89 to about residue 105; domain 3 is from about residue 106 to about residue 284; domain 4 is from about residue 285 to about 385; and domain 5 is from about residue 386 to 453. One of skill in the art will understand that the exact boundaries of these domains may vary.

The chimeric ICAM-1 molecules of the present invention preferably contain at least a portion of an IgM or IgA heavy chain which allows that immunoglobulin heavy chain to bind to immunoglobulin J chain and thereby binds to the secretory component. It is contemplated that the portion of the chimeric ICAM-1 molecule derived from the immunoglobulin heavy chain of the present invention may be comprised of individual domains selected from the IgA heavy chain or the IgM heavy chain or from some other isotype of heavy chain. It is also contemplated that an immunoglobulin domain derived from an immunoglobulin heavy chain other than IgA or IgM or from an immunoglobulin light chain may be molecularly engineered to bind immunoglobulin J chain and thus may be used to produce immunoglobulins of the present invention.

One skilled in the art will understand that immunoglobulins consist of domains which are approximately 100-110 amino acid residues. These various domains are well known in the art and have known boundaries. The removal of a single domain and its replacement with a domain of another antibody molecule is easily achieved with modern molecular biology.

The domains are globular structures which are stabilized by intrachain disulfide bonds. This confers a discrete shape and makes the domains a self-contained unit that can be replaced or interchanged with other similarly shaped domains. The heavy chain constant region domains of the immunoglobulins confer various properties known as antibody effector functions on a particular molecule containing that domain. Example effector functions include complement fixation, placental transfer, binding to staphylococcal protein, binding to streptococcal protein G, binding to mononuclear cells, neutrophils or mast cells and basophils. The association of particular domains and particular immunoglobulin isotypes with these effector functions is

well known and for example, described in Immunology, Roitt *et al.*, Mosby St. Louis, Mo. (1993 3rd Ed.)

One of skill in the art will be able to identify immunoglobulin heavy chain constant region sequences. For example, a number of immunoglobulin DNA and protein sequences are available through GenBank. Table 2 shows the GenBank Accession numbers of immunoglobulin heavy chain genes and the proteins encoded by the genes. The sequences listed in Table 2 are shown in Fig. 7.

Table 2

GENBANK ACCESSION NO.	HUMAN IMMUNOGLOBULIN SEQUENCE NAME	SEQ ID NO.
J00220	Ig α ₁ Heavy Chain Constant Region Coding Sequence	15
J00220	Ig α ₁ Heavy Chain Constant Region Amino Acid Sequence	16
J00221	IgA ₂ Heavy Chain Constant Region Coding Sequence	17
J00221	IgA ₂ Heavy Chain Constant Region Amino Acid Sequence	18
J00228	Ig γ ₁ Heavy Chain Constant Region Coding Sequence	19
J00228	Ig γ ₁ Heavy Chain Constant Region Amino Acid Sequence	20
J00230 V00554	IgG ₂ Heavy Chain Constant Region Coding Sequence	21
J00230 V00554	IgG ₂ Heavy Chain Constant Region Amino Acid Sequence	22
X03604 M12958	IgG ₃ Heavy Chain Constant Region Coding Sequence	23
X03604 M12958	IgG ₃ Heavy Chain Constant Region Amino Acid Sequence	24
K01316	IgG ₄ Heavy Chain Constant Region Coding Sequence	25
K01316	IgG ₄ Heavy Chain Constant Region Amino Acid Sequence	26
K02876	IgD Heavy Chain Constant Region Coding Sequence	27
K02876	IgD Heavy Chain Constant Region Amino Acid Sequence	28
K02877	IgD Heavy Chain Constant Region Coding Sequence	29
K02877	IgD Heavy Chain Constant Region Amino Acid Sequence	30

GENBANK ACCESSION NO.	HUMAN IMMUNOGLOBULIN SEQUENCE NAME	SEQ ID NO.
K02878	Germline IgD Heavy Chain Coding Sequence	31
K02878	Germline IgD Heavy Chain Amino Acid Sequence	32
K02879	Germline IgD Heavy Chain C- δ -3 Domain Coding Sequence	33
K02879	Germline IgD Heavy Chain C- δ -3 Amino Acid Sequence	34
K01311	Germline IgD Heavy Chain J- δ Region: C- δ CH1 Amino Acid Sequence	35
K02880	Germline IgD Heavy Chain Gene, C-Region, Secreted Terminus Coding Sequence	36
K02880	Germline IgD Heavy Chain Gene, C-Region, Secreted Terminus Amino Acid Sequence	37
K02881	Germline IgD-Heavy Chain Gene, C-Region, First Domain of Membrane Terminus Coding Sequence	38
K02881	Germline IgD-Heavy Chain Gene, C-Region, First Domain of Membrane Terminus Amino Acid Sequence	39
K02882	Germline IgD Heavy Chain Coding Sequence	40
K02882	Germline IgD Heavy Chain Amino Acid Sequence	41
K02875	Germline IgD Heavy Chain Gene, C-Region, C- δ -1 Domain Coding Sequence	42
K02875	Germline IgD Heavy Chain Gene, C-Region, C- δ -1 Domain Amino Acid Sequence	43
L00022 J00227 V00555	IgE Heavy Chain Constant Region Coding Sequence	44
L00022 J00227 V00555	IgE Heavy Chain Constant Region Amino Acid Sequence	45
X17115	IgM Heavy Chain Complete Sequence Coding Sequence	46
X17115	IgM Heavy Chain Complete Sequence mRNA	47

The immunoadhesins of the present invention may, in addition to the chimeric ICAM-1 molecule, contain immunoglobulin light chains, or immunoglobulin J chain bound to the immunoglobulin derived heavy chains. In preferred embodiments, the immunoadhesin of the present invention comprises two or four chimeric ICAM-1 molecules and an immunoglobulin J chain bound to at least one of the chimeric ICAM-1 molecules. The J chain is described and known in the art. See, for example, M. Koshland, *The Immunoglobulin Helper: The J Chain*, in *Immunoglobulin Genes*, Academic Press, London, pg. 345, (1989) and Matsuuchi *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 83:456-460 (1986). The sequence of the immunoglobulin J chain is available on various databases in the United States.

The immunoadhesin of the present invention may have a secretory component associated with the chimeric ICAM-1 molecule. This association may occur by hydrogen bonds, disulfide bonds, covalent bonds, ionic interactions or combinations of these various bonds. Typically, chimeric ICAM-1 molecules are held together by disulfide bonds between the molecules. The interaction of the chimeric ICAM-1 molecules may be non-covalent or disulfide bonding. The present invention contemplates the use of secretory component from a number of different species, including human, rat, rabbit, bovine and the like. The nucleotide sequences for these molecules are well known in the art. For example, U.S. Patent 6,046,037 contains many of the sequences and this patent is incorporated herein by reference.

The immunoadhesins of the present invention containing the secretory component, the chimeric ICAM-1 molecule and J chain are typically bonded together by one of the following: hydrogen bonds, disulfide bonds, covalent bonds, ionic interactions or combinations of these bonds.

The present invention also contemplates immunoadhesins which comprise more than one chimeric ICAM-1 molecule. The immunoadhesin may contain chimeric ICAM-1 molecules that are monomeric units and not disulfide bonded to other chimeric ICAM-1 molecules. In preferred embodiments, the immunoadhesion does contain chimeric ICAM-1

5 molecules that are in association with other chimeric ICAM-1 molecules to form dimers and other multivalent molecules. Typically the chimeric ICAM-1 molecule is present as a dimer because of the association of the immunoglobulin portion of the chimeric molecule. The immunoglobulin portion of the chimeric ICAM-1 molecule allows the association of two
10 chimeric ICAM-1 molecules to form a dimeric molecule having two active binding portions made up of the rhinovirus receptor protein portion. In preferred embodiments, dimerization occurs via the disulfide bonding regions that normally occur between the immunoglobulin domains as part of a naturally-occurring immunoglobulin molecule and the native immunoglobulin protein. One of skill in the art will understand that these disulfide bonds that
15 are normally present in the native immunoglobulin molecule can be modified, moved and removed while still maintaining the ability to form a dimer of the chimeric ICAM-1 molecules.

In other preferred embodiments, the immunoadhesin contains multimeric forms of the chimeric ICAM-1 molecule due to the association of J chain with the immunoglobulin portion
20 of the chimeric ICAM molecule. The association of J chain with the dimer of two chimeric ICAM-1 molecules allows the formation of tetrameric forms of the immunoadhesin. In a preferred embodiment, the immunoglobulin portion of the chimeric ICAM-1 molecule is derived from the IgA molecule, and the addition of J chain allows the formation of a tetrameric complex containing four chimeric ICAM-1 molecules and four binding sites. In
25 other preferred embodiments, the immunoglobulin heavy-chain portion of the chimeric molecule is derived from IgM and multivalent complexes containing ten or twelve molecules may be formed. In other preferred embodiments, in which the chimeric ICAM-1 molecule uses a chimeric immunoglobulin heavy-chain, the chimeric ICAM-1 molecule may form dimers or other higher order multivalent complexes through the domains from either IgA or
30 IgM that are responsible for J chain binding. In other chimeric immunoglobulin molecules the portions of the immunoglobulin responsible for the disulfide bonding between the two immunoglobulin heavy-chains and/or the disulfide bonding between an immunoglobulin light-chain and heavy-chain may be placed in the chimeric immunoglobulin molecule to allow the formation of dimers or other high order multivalent complexes.

The present invention contemplates immunoadhesins containing a chimeric ICAM-1 molecule in which the immunoglobulin domains comprising the heavy chain are derived from different isotypes of either heavy or light chain immunoglobulins. One skilled in the art will understand that using molecular techniques, these domains can be substituted for a similar domain and thus produce an immunoglobulin that is a hybrid between two different immunoglobulin molecules. These chimeric immunoglobulins allow immunoadhesins containing secretory component to be constructed that contain a variety of different and desirable properties that are conferred by different immunoglobulin domains.

The present invention also contemplates chimeric ICAM-1 molecules in which the portion of the chimeric molecule derived from immunoglobulin, heavy or light J chain may contain less than an entire domain derived from a different immunoglobulin molecule. The same molecular techniques may be employed to produce such chimeric ICAM-1 molecules.

In preferred embodiments, the chimeric ICAM-1 molecules of the present invention contain at least the C_{H1} , C_{H2} , C_{H3} , domain of mouse or human IgA₁, IgA₂ or IgM. Other preferred embodiments of the present invention contain immunoglobulin domains that include at least the $C_{\mu 1}$, $C_{\mu 2}$, $C_{\mu 3}$, or $C_{\mu 4}$ domains of IgM.

Preferred chimeric ICAM-1 molecules contain domains from two different isotypes of human immunoglobulin. Preferred chimeric ICAM-1 molecules that include immunoglobulins that contain immunoglobulin domains including at least the C_{H1} , C_{H2} , or C_{H3} of human IgG, IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, IgE, or IgD. Other preferred immunoglobulins for use as part of chimeric ICAM-1 molecules include immunoglobulins that contain domains from at least the C_{H1} , C_{H2} , C_{H3} , or C_{H4} domain of IgM or IgE. The present invention also contemplates chimeric ICAM-1 molecules that contain immunoglobulin domains derived from at least two different isotypes of mammalian immunoglobulins. Generally, any of the mammalian immunoglobulins can be used in the preferred embodiments, such as the following isotypes: any isotype of IgG, any isotype of IgA, IgE, IgD or IgM. The present invention also contemplates chimeric ICAM-1 molecules

derived from a species such as human, mouse or other mammals. In preferred embodiments, the chimeric ICAM-1 molecule contains the portion of IgA or IgM responsible for the association of J chain with the IgA and IgM. Thus, by using a chimeric immunoglobulin in the chimeric ICAM-1 molecule, the J chain may associate with a chimeric immunoglobulin that is predominantly of an isotype that does not bind J chain or secretory component.

The present invention also contemplates chimeric ICAM-1 molecules that contain immunoglobulin domains derived from two different isotypes of rodent or primate immunoglobulin. The isotypes of rodent or primate immunoglobulin are well known in the art. The chimeric ICAM-1 molecules of the present invention may contain immunoglobulin derived heavy chains that include at least one of the following immunoglobulin domains: the C_H1, C_H2, or C_H3 domains of a mouse IgG, IgG1, IgG2a, IgG2b, IgG3, IgA, IgE, or IgD; the C_H1, C_H2, C_H3 or C_H4 domain of mouse IgE or IgM; the C_H1 domain of a human IgG, IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, or IgD; the C_H1, C_H2, C_H3, C_H4 domain of human IgM or IgE; the C_H1, C_H2, or C_H3 domain of an isotype of mammalian IgG, an isotype of IgA, IgE, or IgD; the C_H1, C_H2, C_H3 or C_H4 domain of a mammalian IgE or IgM; the C_H1, C_H2, or C_H3 domain of an isotype of rodent IgG, IgA, IgE, or IgD; the C_H1, C_H2, C_H3 or C_H4 domain of a rodent IgE or IgM; the C_H1, C_H2, or C_H3 domain of an isotype of animal IgG, an isotype of IgA, IgE, or IgD; and the C_H1, C_H2, C_H3, or C_H4 domain of an animal IgE or IgM. The present invention also contemplates the replacement or addition of protein domains derived from molecules that are members of the immunoglobulin superfamily into the chimeric ICAM-1 molecules. The molecules that belong to the immunoglobulin superfamily have amino acid residue sequence and nucleic acid sequence homology to immunoglobulins. The molecules that are part of the immunoglobulin superfamily can be identified by amino acid or nucleic acid sequence homology. See, for example, p. 361 of Immunoglobulin Genes, Academic Press (1989).

In preferred embodiments of the present invention, the immunoadhesin is expressed by methods that generate an immunoadhesin having plant-specific glycosylation. It is well-known in the art that glycosylation is a major modification of proteins in plant cells

(Lerouge *et al.*, *Plant Molecular Biology* 38:31-48, 1998). Glycosylation of proteins also occurs in other cell types, including mammalian and insect cells. The glycosylation process starts in the endoplasmic reticulum by the co-translational transfer of a precursor oligosaccharide to specific residues of the nascent polypeptide chain. Processing of this oligosaccharide into different types of glycans, which differ in the types of residues present and the nature of the linkages between the residues, occurs in the secretory pathway as the glycoprotein moves from the endoplasmic reticulum to its final destination. One of skill in the art will understand that at the end of their maturation, proteins expressed in plants or plant cells have a different pattern of glycosylation than do proteins expressed in other types of cells, including mammalian cells and insect cells. Detailed studies characterizing plant-specific glycosylation and comparing it with glycosylation in other cell types have been performed, for example, in studies described by Cabanes-Macheteau *et al.*, *Glycobiology* 9(4):365-372 (1999), and Altmann, *Glycoconjugate J.* 14:643-646 (1997). These groups and others have shown that plant-specific glycosylation generates glycans that have xylose linked $\beta(1,2)$ to mannose, but xylose is not linked $\beta(1,2)$ to mannose as a result of glycosylation in mammalian and insect cells. Plant-specific glycosylation results in a fucose linked $\alpha(1,3)$ to the proximal GlcNAc, while glycosylation in mammalian cells results in a fucose linked $\alpha(1,6)$ to the proximal GlcNAc. Furthermore, plant-specific glycosylation does not result in the addition of a sialic acid to the terminus of the protein glycan, whereas in glycosylation in mammalian cells, sialic acid is added.

The immunoadhesin of the present invention that is glycosylated in a plant-specific manner can contain a chimeric ICAM-1 molecule that includes any combination of extracellular domains 1, 2, 3, 4, and 5 of the ICAM-1 molecule. FIG. 2B shows the amino acid sequence of the chimeric ICAM-1/IgA2 molecule (SEQ ID NO: 8) of the present invention, that contains all five domains of ICAM-1. The bolded N's represent asparagine residues to which oligosaccharide moieties are linked during glycosylation in plant cells, as well as mammalian and insect cells. One of skill in the art will know that the glycosylation sites are the tripeptide Asn-X-Ser/Thr where X can be any amino acid except proline and

aspartic acid (Kornfeld and Kornfeld, *Annu Rev Biochem* 54:631-664, 1985). It will therefore be known to one of skill in the art that which amino acids of the protein having plant-specific glycosylation would depend on which domains of ICAM-1 are present. Because the sequence and domain boundaries of ICAM-1 are known (see Staunton *et. al.*, *Cell* 52:925-933, 1988), it would be evident to one of skill in the art how to determine the plant-specific glycosylation sites on any potential combination of any of the five ICAM-1 domains.

In other preferred embodiments of the present invention, the immunoadhesin having plant-specific glycosylation and containing a chimeric ICAM-1 molecule having any combination of ICAM-1 extracellular domains 1, 2, 3, 4 and 5 further comprises a J chain and secretory component associated with said chimeric ICAM-1 molecule. As was true with respect to the chimeric ICAM-1 molecule, one of skill in the art will be able to identify the sites for plant-specific glycosylation in the J chain and secretory component sequences.

The present invention contemplates immunoadhesins having plant-specific glycosylation, that contain a chimeric ICAM-1 molecule in which the immunoglobulin heavy chain is selected from the group of IgA (SEQ ID NOS: 15-18), IgA₁ (SEQ ID NOS: 15-16), IgA₂ (SEQ ID NO: 17), IgG₁ (SEQ ID NOS: 19-20), IgG₂ (SEQ ID NOS: 21-22), IgG₃ (SEQ ID NOS: 23-24), IgG₄ (SEQ ID NOS: 25-26), IgM (SEQ ID NOS: 46-47), IgD (SEQ ID NO: 27-32 and 36-41-43), IgE (SEQ ID NOS: 44-45), and a chimeric immunoglobulin heavy chain. One of skill in the art will know that which of these immunoglobulin heavy chain sequences, or which combination of immunoglobulin heavy chain sequences are combined in a chimeric immunoglobulin heavy chain, will have an effect on the number and location of glycosylation sites in the chimeric ICAM-1 molecule of the immunoadhesin. As was true with respect to the chimeric ICAM-1 molecule, one of skill in the art will be able to identify the sites for plant-specific glycosylation in the immunoglobulin heavy chain sequences, including the various chimeric immunoglobulin heavy chain sequences that can be constructed.

Also provided herein are immunoadhesin functional derivatives. By “functional derivative” is meant a “chemical derivative,” “fragment,” or “variant,” of the polypeptide or nucleic acid of the invention which retains at least a portion of the function of the protein, for example reactivity with an antibody specific for the protein, enzymatic activity or binding activity, which permits its utility in accordance with the present invention. It is well known in the art that due to the degeneracy of the genetic code numerous different nucleic acid sequences can code for the same amino acid sequence. It is also well known in the art that conservative changes in amino acid can be made to arrive at a protein or polypeptide that retains the functionality of the original. In both cases, all permutations are intended to be covered by this disclosure.

The derivatives may also be engineered according to routine methods to include an affinity purification tag such that large quantities and/or relatively pure or isolated quantities of immunoadhesin may be produced. Many different versions of tag exist that can be incorporated into one or more components of the immunoadhesin, preferably not destroying the desired binding activity of the immunoadhesin in the absence of tag. Such tags can be engineered as expressible encoded nucleic acid sequence fused with nucleic acid sequences encoding the immunoadhesins of the invention. The tags may further be engineered to be removable, *e.g.*, with a commercially available enzyme.

Further, it is possible to delete codons or to substitute one or more codons with codons other than degenerate codons to produce a structurally modified polypeptide, but one which has substantially the same utility activity as the polypeptide produced by the unmodified nucleic acid molecule. As recognized in the art, the two polypeptides can be functionally equivalent, as are the two nucleic acid molecules that give rise to their production, even though the differences between the nucleic acid molecules are not related to the degeneracy of the genetic code.

Manipulations of this sort, and post-production chemical derivatization may be implemented, *e.g.*, to improve stability, solubility, absorption, biological or therapeutic effect,

and/or biological half-life. Moieties capable of mediating such effects are disclosed, for example, in Remington's Pharmaceutical Sciences, 18th ed., Mack Publishing Co., Easton, PA (1990). A functional derivative intended to be within the scope of the present invention is a "variant" polypeptide which either lacks one or more amino acids or contains additional or substituted amino acids relative to the native polypeptide. The variant may be derived from a naturally occurring complex component by appropriately modifying the protein DNA coding sequence to add, remove, and/or to modify codons for one or more amino acids at one or more sites of the C-terminus, N-terminus, and/or within the native sequence. It is understood that such variants having added, substituted and/or additional amino acids retain one or more characterizing portions of the native protein, as described above.

A functional derivative of a protein with deleted, inserted and/or substituted amino acid residues may be prepared using standard techniques well-known to those of ordinary skill in the art. For example, the modified components of the functional derivatives may be produced using site-directed mutagenesis techniques (as exemplified by Adelman *et. al.*, 1983, DNA 2:183) wherein nucleotides in the DNA coding sequence are modified such that a modified coding sequence is produced, and thereafter expressing this recombinant DNA in a prokaryotic or eukaryotic host cell, using techniques such as those described above. Alternatively, proteins with amino acid deletions, insertions and/or substitutions may be conveniently prepared by direct chemical synthesis, using methods well-known in the art. The functional derivatives of the proteins typically exhibit the same qualitative biological activity as the native proteins.

In addition, the immunoadhesins of the invention may be not just modified ICAM-1/Ig immunoadhesins, but may also embrace other native ICAM family members, isotypes, and/or other homologous amino acid sequences, *e.g.*, human, primate, rodent, canine, feline, bovine, avian, etc. Furthermore, the Ig type used in the immunoadhesins can vary, *e.g.*, may assume a different Ig family member identity, within or without a given species. ICAMs and Igs are diverse and have well-known sequences that one of ordinary skill can exploit to create different immunoadhesins having more or less different utility in a given organism to undergo

treatment. An illustrative, nonexhaustive list of examples of molecules having ICAM-1 homology that can be used to create other immunoadhesins include those in the following table.

Table 3

ACCESSION NO.	ICAM NAME	SPECIES
NP 000192	Intercellular Adhesion Molecule-1 (CD54)	Homo sapiens
AAH03097	Intercellular Adhesion Molecule ICAM-2	Homo sapiens
NP 002153	Intercellular Adhesion Molecule 3 Precursor	Homo sapiens
BAB20325	TCAM-1	Homo sapiens
NP 003250	Intercellular Adhesion Molecule 5 (Telencephalin)	Homo sapiens
NM 007164	Mucosal Vascular Address in Cell Adhesion Molecule (MADCAM1)	Homo sapiens
NM 001078	Vascular Cell Adhesion Molecule 1 (VCAM1)	Homo sapiens
AAA37875	MALA-2	Mus musculus
AAA37876	Intercellular Adhesion Molecule-1 Precursor	Mus musculus
AAG30280	Intracellular Adhesion Molecule 1	Cricetulus griseus
AAB39264	Intercellular Adhesion Molecule-3	Bos taurus
AAF80287	Intercellular Adhesion Molecule-1 Precursor	Sus scrofa
AAA18478	Telecephalin	Oryctolagus cuniculus

ACCESSION NO.	ICAM NAME	SPECIES
NP 032345	Intercellular Adhesion Molecule 5, telencephalin	Mus musculus
BAB41106	Cell adhesion molecule TCAM-1	Mus musculus
NP 067705	Testicular Cell Adhesion Molecule 1	Rattus norvegicus
AAG35584	Nectin-Like Protein 1	Mus musculus
AAC18956	CD22 Protein	Homo sapiens
AAA35415	Intercellular Adhesion Molecule 1	Pan troglodytes
AAA83206	89 kDa Protein	Mus musculus
AAA92551	Intercellular Adhesion Molecule-1	Canis familiaris
AAB06749	Intercellular Adhesion Molecule-1	Bos taurus
AAD13617	Intercellular Adhesion Molecule-1 Precursor	Ovis aries
NP 037099	Intercellular Adhesion Molecule-1	Rattus norvegicus
AAE22202	ICAM-4	Rattus norvegicus
AAA60392	cell surface glycoprotein	Homo sapiens
AAF91086	nephrin	Rattus norvegicus
AAF91087	nephrin	Mus musculus

Likewise, numerous heavy chain constant regions of different Ig molecules, both in humans and other species, are known that can be substituted in for those specific Ig regions of the chimeras described herein.

C. Vectors, Cells and Plants Containing Immuno adhesins

5 The present invention also contemplates expression and cloning vectors, cells and plants containing the immuno adhesins of the present invention. Technology for isolating the genes encoding the various portions of the immuno adhesions are well-known to one of skill in the art and can be applied to insert the various required genes into expression vectors and cloning vectors such as those vectors can be introduced into cells and into transgenic plants.

10 The present invention contemplates a method of assembling an immuno adhesin comprising the steps of: introducing into an organism a DNA segment encoding a chimeric ICAM-1 molecule, immunoglobulin J chain, and introducing into the same organism a DNA encoding a secretory component. The preferred secretory component contains at least a segment of the amino acid residues 1 to residue about 606 of the human polyimmunoglobulin
15 receptor (pIgR) amino acid residue sequence or analogous amino acid residues from other species (Mostov, *Ann Dev. Immu.* 12:63-84 1994).

The present invention contemplates eukaryotic cells, including plant cells, containing immuno adhesins of the present invention. The present invention also contemplates plant cells that contain nucleotide sequences encoding the various components of the immuno adhesin of
20 the present invention. One skilled in the art will understand that the nucleotide sequences that encode the secretory component protection protein and the chimeric ICAM-1 molecule and J chain will typically be operably linked to a promoter and present as part of an expression vector or cassette. Typically, if the eukaryotic cell used is a plant cell than the promoter used will be a promoter that is able to operate in a plant cell. After the chimeric ICAM-1 genes,
25 secretory component genes and J chain genes are isolated, they are typically operatively linked to a transcriptional promoter in an expression vector. The present invention also contemplates expression vectors containing a nucleotide sequence encoding a chimeric

ICAM-1 molecule which has been operatively linked to a regulatory sequence for expression. These expression vectors place the nucleotide sequence to be expressed in a particular cell 3' of a promoter sequence which causes the nucleotide sequence to be transcribed and expressed. The expression vector may also contain various enhancer sequences which improve the efficiency of this transcription. In addition, such sequences as terminators, polyadenylation (poly A) sites and other 3' end processing signals may be included to enhance the amount of nucleotide sequence transcribed within a particular cell.

Expression of the components in the organism of choice can be derived from an independently replicating plasmid, or from a permanent component of the chromosome, or from any piece of DNA which may transiently give rise to transcripts encoding the components. Organisms suitable for transformation can be either prokaryotic or eukaryotic. Introduction of the components of the complex can be by direct DNA transformation, by biolistic delivery into the organism, or mediated by another organism as for example by the action of recombinant *Agrobacterium* on plant cells. Expression of proteins in transgenic organisms usually requires co-introduction of an appropriate promoter element and polyadenylation signal. In one embodiment of the invention, the promoter element potentially results in the constitutive expression of the components in all of the cells of a plant. Constitutive expression occurring in most or all of the cells will ensure that precursors can occupy the same cellular endomembrane system as might be required for assembly to occur.

Expression vectors compatible with the host cells, preferably those compatible with plant cells are used to express the genes of the present invention. Typical expression vectors useful for expression of genes in plants are well known in the art and include vectors derived from the tumor-inducing (Ti) plasmid of *Agrobacterium tumefaciens* described by Rogers *et al.*, *Meth. in Enzymol.*, 153:253-277 (1987). However, several other expression vector systems are known to function in plants. See for example, Verma *et al.*, PCT Publication No. WO87/00551; and Cocking and Davey, *Science*, 236:1259-1262 (1987).

The expression vectors described above contain expression control elements including the promoter. The genes to be expressed are operatively linked to the expression vector to allow the promoter sequence to direct RNA polymerase binding and synthesis of the desired polypeptide coding gene. Useful in expressing the genes are promoters which are inducible, viral, synthetic, constitutive, and regulated. The choice of which expression vector is used and ultimately to which promoter a nucleotide sequence encoding part of the immunoadhesin of the present invention is operatively linked depends directly, as is well known in the art, on the functional properties desired, *e.g.* the location and timing of protein expression, and the host cell to be transformed, these being limitations inherent in the art of constructing recombinant DNA molecules. However, an expression vector useful in practicing the present invention is at least capable of directing the replication, and preferably also the expression of the polypeptide coding gene included in the DNA segment to which it is operatively linked.

In preferred embodiments, the expression vector used to express the genes includes a selection marker that is effective in a plant cell, preferably a drug resistance selection marker.

A preferred drug resistance marker is the gene whose expression results in kanamycin resistance, *i.e.*, the chimeric gene containing the nopaline synthase promoter, Tn5 neomycin phosphotransferase II and nopaline synthase 3' nontranslated region described by Rogers *et al.*, in *Methods For Plant Molecular Biology*, a Weissbach and H. Weissbach, eds., Academic Press Inc., San Diego, Calif. (1988). A useful plant expression vector is commercially available from Pharmacia, Piscataway, N.J. Expression vectors and promoters for expressing foreign proteins in plants have been described in U.S. Pat. Nos. 5,188,642; 5,349,124; 5,352,605, and 5,034,322 which are hereby incorporated by reference.

A variety of methods have been developed to operatively link DNA to vectors via complementary cohesive termini. For instance, complementary homopolymer tracks can be added to the DNA segment to be inserted into the vector DNA. The vector and DNA segment are then joined by hydrogen bonding between the complementary homopolymeric tails to form recombinant DNA molecules. Alternatively, synthetic linkers containing one or more restriction endonuclease sites can be used to join the DNA segment to the expression vector.

The synthetic linkers are attached to blunt-ended DNA segments by incubating the blunt-ended DNA segments with a large excess of synthetic linker molecules in the presence of an enzyme that is able to catalyze the ligation of blunt-ended DNA molecules, such as bacteriophage T4 DNA ligase. Thus, the products of the reaction are DNA segments carrying synthetic linker sequences at their ends. These DNA segments are then cleaved with the appropriate restriction endonuclease and ligated into an expression vector that has been cleaved with an enzyme that produces termini compatible with those of the synthetic linker. Synthetic linkers containing a variety of restriction endonuclease sites are commercially available from a number of sources including New England BioLabs, Beverly, Mass.

The nucleotide sequences encoding the secretory component, J chain, the chimeric ICAM-1 molecules of the present invention are introduced into the same plant cell either directly or by introducing each of the components into a plant cell and regenerating a plant and cross-hybridizing the various components to produce the final plant cell containing all the required components.

Any method may be used to introduce the nucleotide sequences encoding the components of the immunoadhesins of the present invention into a eukaryotic cell. For example, methods for introducing genes into plants include *Agrobacterium*-mediated plant transformation, protoplast transformation, gene transfer into pollen, injection into reproductive organs and injection into immature embryos. Each of these methods has distinct advantages and disadvantages. Thus, one particular method of introducing genes into a particular eukaryotic cell or plant species may not necessarily be the most effective for another eukaryotic cell or plant species.

Agrobacterium tumefaciens-mediated transfer is a widely applicable system for introducing genes into plant cells because the DNA can be introduced into whole plant tissues, bypassing the need for regeneration of an intact plant from a protoplast. The use of *Agrobacterium*-mediated expression vectors to introduce DNA into plant cells is well known in the art. See, for example, the methods described by Fraley *et al.*, *Biotechnology*, 3:629

(1985) and Rogers *et al.*, *Methods in Enzymology*, 153:253-277 (1987). Further, the integration of the Ti-DNA is a relatively precise process resulting in few rearrangements. The region of DNA to be transferred is defined by the border sequences and intervening DNA is usually inserted into the plant genome as described by Spielmann *et al.*, *Mol. Gen. Genet.*, 205:34 (1986) and Jorgensen *et al.*, *Mol. Gen. Genet.*, 207:471 (1987). Modern *Agrobacterium* transformation vectors are capable of replication in *Escherichia coli* as well as *Agrobacterium*, allowing for convenient manipulations as described by Klee *et al.*, in *Plant DNA Infectious Agents*, T. Hohn and J. Schell, eds., *Springer-Verlag*, New York, pp. 179-203 (1985). Further recent technological advances in vectors for *Agrobacterium*-mediated gene transfer have improved the arrangement of genes and restriction sites in the vectors to facilitate construction of vectors capable of expressing various polypeptide coding genes. The vectors described by Rogers *et al.*, *Methods in Enzymology*, 153:253 (1987), have convenient multi-linker regions flanked by a promoter and a polyadenylation site for direct expression of inserted polypeptide coding genes and are suitable for present purposes.

In those plant species where *Agrobacterium*-mediated transformation is efficient, it is the method of choice because of the facile and defined nature of the gene transfer. *Agrobacterium*-mediated transformation of leaf disks and other tissues appears to be limited to plant species that *Agrobacterium tumefaciens* naturally infects. Thus, *Agrobacterium*-mediated transformation is most efficient in dicotyledonous plants.

Few monocots appear to be natural hosts for *Agrobacterium*, although transgenic plants have been produced in asparagus using *Agrobacterium* vectors as described by Bytebier *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 84:5345 (1987). Therefore, commercially important cereal grains such as rice, corn, and wheat must be transformed using alternative methods. Transformation of plant protoplasts can be achieved using methods based on calcium phosphate precipitation, polyethylene glycol treatment, electroporation, and combinations of these treatments. See, for example, Potrykus *et al.*, *Mol. Gen. Genet.*, 199:183 (1985); Lorz *et al.*, *Mol. Gen. Genet.*, 199:178 (1985); Fromm *et al.*, *Nature*, 319:791

(1986); Uchimiya *et al.*, *Mol. Gen. Genet.*, 204:204 (1986); Callis *et al.*, *Genes and Development*, 1:1183 (1987); and Marcotte *et al.*, *Nature*, 335:454 (1988).

Application of these methods to different plant species depends upon the ability to regenerate that particular plant species from protoplasts. Illustrative methods for the regeneration of cereals from protoplasts are described in Fujimura *et al.*, *Plant Tissue Culture Letters*, 2:74 (1985); Toriyama *et al.*, *Theor Appl. Genet.*, 73:16 (1986); Yamada *et al.*, *Plant Cell Rep.*, 4:85 (1986); Abdullah *et al.*, *Biotechnology*, 4:1087 (1986).

To transform plant species that cannot be successfully regenerated from protoplasts, other ways to introduce DNA into intact cells or tissues can be utilized. For example, regeneration of cereals from immature embryos or explants can be effected as described by Vasil, *Biotechnology*, 6:397 (1988). In addition, "particle gun" or high-velocity microprojectile technology can be utilized. Using such technology, DNA is carried through the cell wall and into the cytoplasm on the surface of small (0.525 μm) metal particles that have been accelerated to speeds of one to several hundred meters per second as described in Klein *et al.*, *Nature*, 327:70 (1987); Klein *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 85:8502 (1988); and McCabe *et al.*, *Biotechnology*, 6:923 (1988). The metal particles penetrate through several layers of cells and thus allow the transformation of cells within tissue explants. Metal particles have been used to successfully transform corn cells and to produce fertile, stably transformed tobacco and soybean plants. Transformation of tissue explants eliminates the need for passage through a protoplast stage and thus speeds the production of transgenic plants.

DNA can also be introduced into plants by direct DNA transfer into pollen as described by Zhou *et al.*, *Methods in Enzymology*, 101:433 (1983); D. Hess, *Intern Rev. Cytol.*, 107:367 (1987); Luo *et al.*, *Plant Mol. Biol. Reporter*, 6:165 (1988). Expression of polypeptide coding genes can be obtained by injection of the DNA into reproductive organs of a plant as described by Pena *et al.*, *Nature*, 325:274 (1987). DNA can also be injected directly into the cells of immature embryos and the rehydration of desiccated embryos as

described by Neuhaus *et al.*, *Theor. Appl. Genet.*, 75:30 (1987); and Benbrook *et al.*, in *Proceedings Bio Expo* 1986, Butterworth, Stoneham, Mass., pp. 27-54 (1986).

The regeneration of plants from either single plant protoplasts or various explants is well known in the art. See, for example, *Methods for Plant Molecular Biology*, A. Weissbach and H. Weissbach, eds., Academic Press, Inc., San Diego, Calif. (1988). This regeneration and growth process includes the steps of selection of transformant cells and shoots, rooting the transformant shoots and growth of the plantlets in soil.

The regeneration of plants containing the foreign gene introduced by *Agrobacterium tumefaciens* from leaf explants can be achieved as described by Horsch *et al.*, *Science*, 227:1229-1231 (1985). In this procedure, transformants are grown in the presence of a selection agent and in a medium that induces the regeneration of shoots in the plant species being transformed as described by Fraley *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 80:4803 (1983). This procedure typically produces shoots within two to four weeks and these transformant shoots are then transferred to an appropriate root-inducing medium containing the selective agent and an antibiotic to prevent bacterial growth. Transformant shoots that rooted in the presence of the selective agent to form plantlets are then transplanted to soil to allow the production of roots. These procedures will vary depending upon the particular plant species employed, such variations being well known in the art.

The immunoadhesins of the present invention may be produced in any plant cell including plant cells derived from plants that are dicotyledonous or monocotyledonous, solanaceous, alfalfa, legumes, or tobacco.

Transgenic plants of the present invention can be produced from any sexually crossable plant species that can be transformed using any method known to those skilled in the art. Useful plant species are dicotyledons including tobacco, tomato, the legumes, alfalfa, oaks, and maples; monocotyledons including grasses, corn, grains, oats, wheat, and barley; and lower plants including gymnosperms, conifers, horsetails, club mosses, liverworts, hornworts, mosses, algae, gametophytes, sporophytes or pteridophytes.

The present invention also contemplates expressing the immunoadhesins within eukaryotic cells including mammalian cells. One of skill in the art will understand the various systems available for expression of the immunoadhesin in mammalian cells and can readily modify those system to express the immunoadhesions and chimeric ICAM-1 molecules of the present invention in those cells. In preferred embodiments, the chimeric ICAM-1, J chain and secretory component molecules of the present invention are placed in a vector pCDM8 which has been previously described by Aruffo, *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 84:8573-8577 (1987). The use of the PCDM8 construct is by no means unique and numerous other systems are available that do not utilize the cog cell expression system. For example, the following United States Patents describe useful eukaryotic expression systems that may be used with the chimeric ICAM-1 and other molecules of the immunoadhesin.

D. Compositions Containing Immunoadhesins

The present invention also contemplates compositions containing an immunoadhesin of the present invention together with plant macromolecules or material. Typically these plant macromolecules or plant materials are derived from any plant useful in the present invention. The plant macromolecules are present together with an immunoadhesin of the present invention for example, in a plant cell, in an extract of a plant cell, or in a plant. Typical plant macromolecules associated with the immunoadhesin of the present invention in a composition are ribulose bisphosphate carboxylase, light harvesting complex pigments (LHCP), secondary metabolites or chlorophyll. The compositions of the present invention have plant material or plant macromolecules in a concentration of between 0.01% and 99% mass excluding water. Other compositions include compositions having the immunoadhesins of the present invention present at a concentration of between 1% and 99% mass excluding water. Other compositions include immunoadhesins at a concentration of 50% to 90% mass excluding water.

The compositions of the present invention may contain plant macromolecules at a concentration of between 0.1% and 5% mass excluding water. Typically the mass present in the composition will consist of plant macromolecules and immunoadhesins of the present

invention. When the immunoadhesins of the present invention are present at a higher or lower concentration the concentration of plant macromolecules present in the composition will vary inversely. In other embodiments the composition of plant macromolecules are present in a concentration of between 0.12% and 1% mass excluding water.

5 The present invention contemplates a composition of matter comprising all or part of the following: a chimeric ICAM-1 molecule, a J chain or a secretory component. These components form a complex and are associated as was previously described. Typically, the composition also contains molecules derived from a plant. This composition may also be obtained after an extraction process yielding functional immunoadhesin and plant-derived
10 molecules.

The extraction method comprises the steps of applying a force to a plant containing the complex whereby the apoplastic compartment of the plant is ruptured releasing said complex. The force involves shearing as the primary method of releasing the apoplastic liquid.

15 The whole plant or plant extract contains an admixture of immunoadhesin and various other macromolecules of the plant. Among the macromolecules contained in the admixture is ribulose biphosphate carboxylase (RuBisCo) or fragments of RuBisCo. Another macromolecule is LHCP. Another molecule is chlorophyll.

Other useful methods for preparing compositions containing immunoadhesins having
20 chimeric ICAM-1 molecule include extraction with various solvents and application of vacuum to the plant material. The compositions of the present invention may contain plant macromolecules in a concentration of between about 0.1% and 5% mass excluding water.

The present invention also contemplates therapeutic compositions which may be used in the treatment of a patient or animal. Administration of the therapeutic composition can be
25 before or after extraction from the plant or other transgenic organism. Once extracted the immunoadhesins may also be further purified by conventional techniques such as size

exclusion, ion exchange, or affinity chromatography. Plant molecules may be co-administered with the complex.

The present invention also contemplates that the relative proportion of plant-derived molecules and animal-derived molecules can vary. Quantities of specific plant proteins, such as RuBisCo or chlorophyll may be as little as 0.01% of the mass or as much as 99.9% of the mass of the extract, excluding water.

The present invention also contemplates the direct use of the therapeutic plant extract containing immunoadhesins without any further purification of the specific therapeutic component. Administration may be by topical application, oral ingestion, nasal spray or any other method appropriate for delivering the antibody to the mucosal target pathogen.

E. Pharmaceutical Compositions, Formulations, And Routes Of Administration

The immunoadhesins described herein can be administered to a patient, preferably in the form of a suitable pharmaceutical composition. Such composition may include components in addition to, or in lieu of, those described above. The composition preferably exhibits either or both of a therapeutic and prophylactic property when administered. The preparation of such compositions can be done according to routine methodologies in the art, and may assume any of a variety of forms, e.g., liquid solutions, suspensions or emulsifications, and solid forms suitable for inclusion in a liquid prior to ingestion.

Techniques for the formulation and administration of polypeptides and proteins may be found in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, latest edition. Using these procedures, one of ordinary skill can utilize the immunoadhesins of the invention to achieve success without undue experimentation.

1. Administration Routes

Suitable routes of administration for the invention include, e.g., oral, nasal, inhalation, intraocular, pharyngeal, bronchial, transmucosal, or intestinal administration. Alternatively,

one may administer the compound in a local manner, *e.g.*, via injection or other application of the compound to a preferred site of action.

2. Formulations

The pharmaceutical compositions of the present invention may be manufactured in a manner that is itself known, *e.g.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes. One or more physiologically acceptable carriers comprising excipients and/or other auxiliaries can be used to facilitate processing of the active compounds into pharmaceutical preparations. Proper formulation is dependent upon the particular route of administration chosen.

For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Suitable carriers include excipients such as, *e.g.*, fillers such as sugars, including lactose, sucrose, mannitol, and/or sorbitol; cellulose preparations such as, *e.g.*, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

For administration by inhalation, the compounds for use according to the present invention may be conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of *e.g.* gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use.

In addition, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

Additionally, the compounds may be delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein stabilization may be employed.

The pharmaceutical compositions also may comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

Pharmaceutically compatible salts may be formed with many acids, including but not limited to hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, citric, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In solutions, manipulation of pH is also routinely employed for optimizing desired properties.

3. Determining Effective Dosages and Dosage Regimens

Pharmaceutical compositions suitable for use in the present invention include compositions where the active ingredients are contained in an amount effective to achieve an intended purpose, *e.g.*, a therapeutic and/or prophylactic use. A pharmaceutically effective

amount means an amount of compound effective to prevent, alleviate or ameliorate symptoms of disease or prolong the survival of the subject being treated. Determination of a pharmaceutically effective amount is well within the capability of those skilled in the art, and will typically assume an amount of between about 0.5 $\mu\text{g/kg/day}$ and about 500g/kg/day, with
5 individual dosages typically comprising between about 1 nanogram and several grams of immunoadhesin.

For any compound used in the methods of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. For example, varying dosages can be administered to different animals or cell cultures and compared for effect. In this way, one
10 can identify a desired concentration range, and prepare and administer such amount accordingly. Optimization is routine for one of ordinary skill in the art.

The person of skill, in addition to considering pharmaceutical efficacy, also considers toxicity according to standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD50 (the dose lethal to 50% of the population) and the
15 ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between LD50 and ED50. Compounds which exhibit high therapeutic indices are preferred. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a
20 range of concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See *e.g.*, Fingl *et al.*, 1975, in "*The Pharmacological Basis of Therapeutics*," Ch. 1 p.1).

25 Dosage amount and frequency may be adjusted to provide mucosal levels of immunadhesin sufficient to maintain or provide a pharmaceutical effect, *e.g.*, therapeutic and/or prophylactic. The minimal effective concentration (MEC) will vary for each

immunadhesin and immunoadhesin formulation, but can be estimated from *in vitro* and/or *in vivo* data. Dosages necessary to achieve MEC will depend on individual characteristics and route of administration. However, assays as described herein can be used to determine mucosal concentrations, which can then be further optimized in amount and precise
5 formulation.

Dosage intervals can also be determined using MEC value. Compounds can be administered using a regimen which maintains mucosal levels above the MEC for 10-90% of the time, 30-90% of the time, or, most preferably, 50-90% of the time.

The compositions may, if desired, be presented in a pack or dispenser device which
10 may contain one or more unit dosage forms containing the active ingredient. The pack may for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. The pack or dispenser may also be accompanied with a notice associated with the container in form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceuticals, which
15 notice is reflective of approval by the agency of the form of the immunoadhesin for human or veterinary administration. Such notice, for example, may be the labeling approved by the U.S. Food and Drug Administration for prescription drugs, or the approved product insert. Compositions comprising a compound of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled
20 for treatment of an indicated condition, *e.g.* treatment or prophylaxis of a disease mediated by host organism/patient ICAM molecules.

F. Methods of Treatment and Prevention of ICAM-mediated Afflictions

A patient in need of therapeutic and/or prophylactic immunoadhesin chimeras of the invention, *e.g.*, to counter rhinovirus infection and/or symptoms such as occur with colds, can
25 be administered a pharmaceutically effective amount of desired immunoadhesin, preferably as part of a pharmaceutical composition determined, produced, and administered as described above. These formulations and delivery modalities can vary widely. Described following are

preliminary procedures that can be used to deduce effective amounts and toxicity, and which can then be conveniently used to determine treatment and prophylaxis parameters and regimens, both in humans and other animals. These procedures are illustrative only and are not intended to be limiting of the invention. Further, these procedures are routine for one of
5 ordinary skill in the art.

1. Ability of the Immunoadhesin to Reduce Rhinovirus Infectivity in Humans: Dose Escalation Tolerance Study

Immunoadhesins of the invention may be tested, *e.g.*, using randomized controlled trials to determine the effect of administration, *e.g.*, intranasal, of immunoadhesin on
10 infection. Other administration routes can be used. Various assays exist that can be used to monitor effect, *e.g.*, IL-8 response assays that evaluate illness symptoms, *e.g.*, cold symptoms caused by rhinovirus infection. These studies can evaluate the extent to which an immunoadhesin taken by a patient subjects can prevent or treat rhinovirus infection. For example, healthy or unhealthy subjects can be administered the immunoadhesin and
15 evaluated over a time course, *e.g.*, in tandem with rhinovirus inoculation and/or illness progression. The clinical protocols used may be based on protocols previously used in evaluation of a recombinant soluble ICAM-1 molecule for efficacy against rhinovirus infection, or modifications thereto (Turner, *et. al.*, *JAMA* 281:1797-804, 1999).

Male and female subjects of any species, age, health, or disease state can be evaluated
20 The subjects may exhibit a serum neutralizing antibody titer in advance of study, which titer may fluctuate in response to infection and immunoadhesin administration.

The immunoadhesin of the present invention may be formulated as a buffered saline with varying amounts of immunoadhesin within and administered at various intervals to a patient. Single ascending dose and multiple ascending dose studies can be used to evaluate
25 the safety of the immunoadhesin. In each case, one or more subjects may be evaluated at each dosage level, some receiving the immunoadhesin, and one or more optionally receiving

placebo. In either study, multiple dosage levels may be evaluated. Dosage levels can vary, but are typically in the nanogram to gram range.

Dosages may be administered over seconds, minutes, hours, weeks, and months, and evaluated for toxicity and/or pharmaceutical effect.

5 Safety and toxicity may be assessed, e.g., by visual examination of the nasal mucosa for signs of irritation or inflammation. Blood safety evaluations can also be employed according to routine methods and using sensitive assays such as ELISA to determine various blood components, including circulating immunoadhesin and rhinovirus quantities. Naval lavage testing may similarly be done according to routine methodologies.

10 Routine statistical analyses and calculations may be employed to determine efficacy and toxicity predicted over time courses for single patients and/or for populations of patient-recipients..

15 Challenge studies as well known in the art can be used to demonstrate that treatment protects against clinical colds and/or reduces cold symptoms after viral challenge, and using commercially available starting materials such virus, cells, and animals. *See, e.g., Gwaltney, et. al., Prog. Med. Virol. 39:256-263, 1992.*

The following examples illustrate the disclosed invention. These examples in no way limit the scope of the claimed invention.

EXAMPLES

1. Construction of Immunoadhesin Expression Cassettes

A cassette encoding ICAM-1 extracellular domains D1 through D5 was prepared by PCR cloning. Specifically, a fragment containing all five extracellular Ig-like domains of ICAM-1 was amplified from plasmid pCDIC1-5D/IgA (insert Martin, *et al.* reference) using the following oligonucleotide primers:

5'-TCTGTTCCCAGGAACTAGTTTGGCACAGACATCTGTGTCCCCCTCAAAGTC-3'
(SEQ ID NO: 6)

5'-CATACCGGGGACTAGTCACATTACGGTCACCTCGCGG-3'
(SEQ ID NO: 7)

These two primers were designed to introduce SpeI sites at the 5' and 3' ends of the PCR fragment (underlined nucleotides). PCR was performed with Pfu polymerase (Stratagene) to reduce accumulation of errors. The PCR fragment was cloned into the vector PCRScript (Stratagene), and sequenced before fusing to the human IgA2 cassettes (with and without SEKDEL at the carboxy-terminus).

Constructs for the expression in plants of human J chain and secretory component, as well as a human IgA2 heavy chain, were developed. A heavy chain expression cassette vector was made and called pSSpHuA2 (See FIG. 1). It contains sequence encoding a bean legumin signal peptide (Baumlein *et al.*, *Nucleic Acids Res.* 14 (6), 2707-2720, 1986). The sequence of bean legumin is provided as GenBank Accession No. X03677, and the sequence of the bean legumin signal peptide is SEQ ID NO: 10 (also see Fig. 8) and the IgA2m(2) constant region with SpeI and SacI sites in between, and the SuperMas promoter for driving the expression of a signal peptide and the constant regions of the human IgA2m(2) heavy-chain.

The amplified DNAs encoding the first five domains of human ICAM-1, and the Fc region of human IgA2m(2) were fused in a plant-expression cassette to make a chimeric

ICAM-1 molecule expression construct, shown in FIG. 2A. This was done by cloning the fragment encoding the five extracellular domains of ICAM-1 into vector pSSPHuA2 to generate pSSPICAMHuA2. The convenient restriction sites (5' SpeI and 3' Spe I) allowed the amplified fragment to be inserted between the signal peptide and the C α 1 domain. In the
5 resulting construct, expression of the chimeric ICAM-1 molecule is under the control of the constitutive promoter "superMAS" (Ni *et al.*, 1995) and the nos 3' terminator region.

The resulting chimeric ICAM-1 molecule construct contains no variable region. Upon translation of the mRNA, signal peptide cleavage is predicted to deposit the ICAM-1-heavy chain fusion into the plant cell's endoplasmic reticulum (ER). DNA encoding an ER retention
10 signal (RSEKDEL, SEQ ID NO: 5) was appended to the 3' end of the heavy-chain in order to boost the expression level of the construct. The amino acid sequence SEKDEL (SEQ ID NO: 4) is the consensus signal sequence for retention of proteins in the endoplasmic reticulum in plant cells. This sequence has been shown to enhance accumulation levels of antibodies in plants (Schouten *et al.*, *Plant Molecular Biology* 30:781-793,1996). The amino acid sequence
15 of the chimeric ICAM-1 molecule construct is shown in FIG. 2B. The DNA sequence and translational frame of the construct was verified before it was used for particle bombardment.

It has been shown recently that assembly of J chain with IgA takes place in the Golgi apparatus (Yoo *et al.*, *J. Biol. Chem.* 274:33771-33777, 1999), and so constructions of heavy
chain without SEKDEL have been made as well. The ICAM-1 fragment was cloned into an
20 expression cassette containing the IgA2m(2) constant region without SEKDEL.

2. Expression of Assembled Immunoadhesin in Plants

A. Immunoadhesin Expression Vectors

The plasmid pSSPICAMHuA2 (SEQ ID NO:9 and FIG. 8) is 6313 bp in length. Nucleotides 49-1165 represent the Superpromoter (Ni *et al.*, *Plant Journal* 7:661-676, 1995).
25 Nucleotides 1166-3662 comprise a sequence encoding a human ICAM-1/human IgA2m(2) constant hybrid with linker sequences. A consensus Kozak sequence (Kozak, *Cell* 44(2):283-92, 1986) is included (nt 1186-1192) to enhance translation initiation, as well as the signal

peptide from *V. faba* legumin (nt 1189-1257; Bäumlein *et al.*, *Nucleic Acids Res.* 14(6):2707-2720 (1986). The sequence of the human IgA2m(2) constant region (nt 3663-3633) has been previously published (Chintalacharuvu, *et al.*, *J. Imm.* 152: 5299-5304, 1994). A sequence encoding the endoplasmic reticulum retention signal SEKDEL is appended to the end of the heavy Chain (nt 3634-3654). Nucleotides 3663-3933 derive from the nopaline synthase 3' end (transcription termination and polyadenylation signal; Depicker *et al.*, 1982). The remainder of the plasmid derives from the vector pSP72 (Promega).

The plasmid pSHuJ (SEQ ID NO: 11 and FIG. 8) is 4283 bp in length. Nucleotides 14-1136 represent the Superpromoter (Ni *et al.*, *Plant Journal* 7:661-676, 1995) and nucleotides 1137-1648 are shown in FIG. 8 and comprise a sequence encoding the human J Chain including the native signal peptide (Max and Korsmeyer, *J Imm.* 152:5299-5304, 1985) along with linker sequences. A consensus Kozak sequence (Kozak, *Cell* 44(2):283-92, 1986) is included (nt 1162-1168) to enhance translation initiation. Nucleotides 1649-1902 derive from the nopaline synthase 3' end (transcription termination and polyadenylation signal; Depicker *et al.*, *J Mol Appl Genet* 1(6):561-73, 1982). The remainder of the plasmid derives from the vector pSP72 (Promega).

The plasmid pSHuSC (SEQ ID NO:12 and FIG. 8) is 5650 bp in length. Nucleotides 13-1136 are derived from the Superpromoter (Ni *et al.*, *Plant Journal* 7:661-676, 1995), and nucleotides 1137-2981 comprise a sequence encoding the human Secretory Component including the native signal peptide (Krajci, *et al.*, *Biochem. and Biophys. Res. Comm* 158:783, 1994) along with linker sequences. A consensus Kozak sequence (Kozak, *Cell* 44(2):283-92, 1986) is included (nt 1151-1157) to enhance translation initiation. Nucleotides 2982-3236 derive from the nopaline synthase 3' end, providing a transcription termination and polyadenylation signal, described in Depicker *et al.*, *J Mol Appl Genet* 1(6):561-73 (1982). The remainder of the plasmid derives from the vector pSP72 (Promega).

The plasmid pBMSP-1 (SEQ ID NO:13 and FIG. 8) is derived from pGPTV-KAN. Becker *et al.*, in *Plant Molecular Biology* 20, 1195-1197, (1992), describe new plant binary

vectors with selectable markers located proximal to the left T-DNA border, and the sequences outside of the left and right borders. Nucleotides 18-187 of pBMSP-1 represent the right T-DNA border, and nucleotides 1811-775 represent the superMAS promoter. Nucleotides 2393-2663 represent the NOS promoter (Depicker *et al.*, *J Mol Appl Genet* 1(6):561-73, 1982), nucleotides 2698-3492 encode the NPTII gene (conferring resistance to kanamycin), and nucleotides 3511-3733 are the polyadenylation signal from *A. tumefaciens* gene 7 (Gielen *et al.*, *Embo J* 3:835-46, 1984). Nucleotides 1768-976 encode the NPTII gene, and nucleotides 4317-4464 represent the left T-DNA border.

The plasmid pBMSP-1spJSC (SEQ ID NO:14 and FIG. 8) is a derivative of pBMSP, containing both J and SC under control of superpromoter. In this plasmid, nucleotides 1-149 represent the left T-DNA border. Nucleotides 955-733 are the polyadenylation signal from *A. tumefaciens* gene, nucleotides 1768-976 encode the NPTII gene (conferring resistance to kanamycin), and nucleotides 2073-1803 represent the NOS promoter. Nucleotides 2635-3768 represent the superMAS promoter, nucleotides 3774-5595 encode the Human Secretory component, and nucleotides 5603-5857 represent the NOS polyadenylation signal. Nucleotides 5880-6991 represent the superMAS promoter, nucleotides 7007-7490 encode the Human Joining Chain, and nucleotides 7504-7757 represent the NOS polyadenylation signal. Nucleotides 7886-8057 represent the right T-DNA border.

The plasmid pGPTV-HPT, encoding the enzyme conferring hygromycin resistance, is available commercially from the Max-Planck-Institut für Züchtungsforschung (Germany). It is described by Becker in *Plant Molecular Biology* 20, 1195-1197 (1992).

B. Plant Transformation and Immunoadhesin Expression in Plants

The expression cassettes described above were used to produce the assembled immunoadhesin in plants. Plasmids pSSPICAMHuA2, pSHuJ, pSHuSC and pBMSP-1 were co-bombarded into tobacco leaf tissue (*N. tabacum* cultivar Xanthi) and transformed microcalli were selected on nutrient agar in the presence of kanamycin. Individual microcalli,

indicative of independent transformation events, were dissected from the parent tissue and propagated on nutrient agar with kanamycin.

The callus tissues were screened for transgene expression. Callus #7132 was shown to express a chimeric ICAM-1 immunoadhesin and J chain by immunoblotting and PCR (data not shown). This callus did not possess DNA encoding the SC. The callus grew well in culture and, upon accumulation of sufficient mass, #7132 was bombarded again, this time with two of the plasmids described above, PBMS-1 SpJSC, containing expression cassettes for both the J chain and SC and pGPTV-HPT, containing an expression cassette for the hpt I gene which confers hygromycin resistance. After a period of selection and growth on nutrient agar, several independent transformants were identified, by immunoblotting, that expressed the chimeric ICAM-1 molecule, the J chain and SC in several states of assembly.

FIG. 3 illustrates the expression of the chimeric ICAM-1 molecule in independently transformed tobacco calli. FIG. 3A shows immunoblots of non-reducing SDS-polyacrylamide gels on which samples containing different transformed tobacco calli (C) and aqueous extracts (Aq) were run and probed for the presence of human ICAM. The solubility of the immunoadhesin assured us that extraction could be easily performed, and the similarity of signals leads us to believe in the reproducibility of expression. FIG. 3B shows immunoblots of nonreducing SDS-polyacrylamide gels containing various fractions of partially purified immunoadhesin from callus Rhi107-11. The blots were probed with antibodies against human ICAM (~ICAM), human IgA heavy chain (~ α), human secretory component (~SC) and human J chain (~J). Secondary, enzyme-conjugated antibodies were employed as necessary to label immuno-positive bands with alkaline phosphatase. The specificity of immuno-blotting was further verified by a failure to detect immuno-positive bands in extracts of non-expressing calli (not shown). The expected MW for a dimerized chimeric ICAM-1 molecule, without glycosylation, is 173,318; this form is likely present in the band migrating just below the 250kD marker since it is immuno-positive for ICAM-1 and heavy-chain. This band is also immuno-positive for SC (total expected MW of ~248 kD) but not for J chain which is somewhat unexpected given the canonical pathway for SIgA

assembly, which involves 2 cell types (in mammalian) and requires the presence of J chain prior to assembly of SC. A tetrameric immunoadhesin, containing a single molecule of J chain and a single molecule of SC, has an expected MW of ~440 kD, prior to glycosylation. Several species with molecular weights well in excess of 200 kD, immuno-positive with all
5 four probes, are readily apparent.

Bombardment with DNA-coated microprojectiles is used to produce stable transformants in both plants and animals (reviewed by Sanford *et al.*, *Meth. Enz.* 217:483-509,1993). Particle-mediated transformation with the vectors encoding the immunoadhesin of the present invention was performed using the PDS-1000/He particle acceleration device,
10 manufactured by Bio-Rad. The PDS-1000/He particle acceleration device system uses Helium pressure to accelerate DNA-coated microparticles toward target cells. The physical nature of the technique makes it extremely versatile and easy to use. We have successfully transformed tobacco with all four components of a secretory IgA simultaneously.

The basic biolistic procedure was performed as follows: A stock suspension of
15 microprojectiles was prepared by mixing 60 mg of particles in 1 ml of absolute ethanol. This suspension was vortexed and 25-50 μ l was removed and added to a sterile microcentrifuge tube. After microcentrifuging for 30 seconds the ethanol was removed and the pellet resuspended in 1 ml sterile water and centrifuged for 5 minutes. The water was then removed and the pellet resuspended in 25-50 μ l of DNA solution containing a mixture of plasmid
20 DNAs, usually, but not always in equimolar amounts. The amount of plasmid added varied between 0.5 ng and 1 μ g per preparation. The following were added sequentially: 220 μ l of sterile water, 250 μ l of 2.5M CaCl_2 , and 50 μ l of 0.1M spermidine. This mixture was vortexed for at least 10 min and then centrifuged for 5 min. The supernatant was removed and the DNA/microprojectile precipitated in 600 μ l of absolute ethanol, mixed and
25 centrifuged 1min. The ethanol was removed and the pellet resuspended in 36 μ l of ethanol. Ten μ l of the suspension was applied as evenly as possible onto the center of a macrocarrier sheet made of Kapton (DuPont) and the ethanol was evaporated. The macrocarrier sheet and a rupture disk were placed in the unit. A petri dish containing surface-sterilized tobacco

leaves was placed below the stopping screen. The chamber was evacuated to 28-29mm Hg and the target was bombarded once. The protocol has been optimized for tobacco, but is optimized for other plants as well by varying parameters such as He pressure, quantity of coated particles, distance between the macrocarrier and the stopping screen and flying
5 distance from the stopping screen to the tissue.

Expression cassettes for chimeric ICAM-1 molecules were also assembled in binary vectors for use in *Agrobacterium*-mediated transformation. An *Agrobacterium* binary vector designed for expression of both human J chain and human secretory component, as well as kanamycin resistance, was introduced into *A. tumefaciens* strain LBA4404. The chimeric
10 ICAM/IgA molecule in another binary vector was also used to transform LBA4404. Overnight cultures of both strains were used for simultaneous "co-cultivation" with leaf pieces of tobacco, according to a standard protocol (Horsch *et al.*, *Science* 227:1229-1231, 1985).

A standard protocol for regeneration of both bombarded and *Agrobacterium*-
15 transformed tobacco leaf disks was used (Horsch *et al.*, *Science* 227:1229-1231, 1985). Because transformed plants, regenerated from bombarded tissue, frequently undergo gene-silencing upon maturation, transgenic tobacco plants were prepared via *Agrobacterium*-mediated transformation, which gives a higher yield of expressing, mature plants.

3. Purification of Assembled Immunoadhesin

20 The immunoadhesin expressed according to Examples 3 was purified. Calli were grown in large amounts to facilitate the development of extraction procedures. A partial purification schedule provided a stable concentrate, available in a variety of buffer conditions, for investigation of subsequent chromatographic techniques for the further purification of the immunoadhesin (See FIG. 3). Calli were extracted in a juicer, which crushes tissue between
25 two stainless-steel gears, while bathed in a buffer containing sodium citrate (0.6 M, pH 7.4) and urea (final concentration of 2 M). The juice (~1 ml/g fresh weight) was precipitated, after coarse filtration through cheesecloth, with 0.67 volumes of saturated ammonium sulfate. A

green pellet was collected after centrifugation and thoroughly extracted, in a small volume of 50 mM sodium citrate (pH 6.6), with a Dounce homogenizer. After additional centrifugation, a clear brown supernatant was collected and partially purified, during buffer exchange in a de-salting mode, by passage through a Sephadex G-100 column. The desalting/buffer exchange step has allowed preparation of a partially purified concentrate (~0.2 ml/ g fresh weight callus) in a desirable buffer; the G-100 column was eluted with 0.25 X phosphate buffered saline. This eluate appeared to be stable for at least 10 days at 2-8°C.

4. The Immunoadhesin Inhibits Human Rhinovirus Infectivity

The infectivity of cells by human rhinovirus was demonstrated to be inhibited by the immunoadhesin prepared according to Example 3. Callus extract prepared according to Example 3 successfully competed for binding of an anti-ICAM monoclonal antibody to soluble ICAM-1. FIG. 4 shows the data from an enzyme-linked immunosorbent assay (ELISA). For the assay, 96-well plates were coated with 0.25 µg soluble ICAM-1/ml. The squares represent the increasing concentrations of sICAM and the circles represent the increasing amounts of callus extract (sterile filtered fraction from G-100) used to compete with the adhered ICAM for a constant amount of a mouse (anti-human ICAM) antibody. After washing the wells, adherent mouse antibody was detected with an anti-mouse antibody conjugated to horseradish peroxidase. Adherent enzyme activity was measured at 490 nm, with ortho-phenylene diamine as a substrate. The data (squares, sICAM; circles, Extract) are well described by sigmoids of the form $OD_{490} = y = y_0 + a/[1 + e^{-\{(x-x_0)/b\}}]$, where $a = y_{\max} - y_0$, $y_0 = y_{\min}$, b = the slope of the rapidly changing portion of the curve and x_0 = the value of x at the 50% response level. Relative to soluble ICAM-1, the immunoadhesin extract tested here contains the equivalent of ~250 µg ICAM/ml; this is an overestimate due to expected avidity effects of the dimeric and tetrameric assemblies of the ICAM-1-heavy-chain fusions. Thus, this ELISA demonstrated that the immunoadhesin competes with soluble ICAM-1 for binding to an anti-ICAM mAb.

The competitive ELISA allows for quantitative assessment of the recovery of activity by comparing the normalized amounts of various fractions required to give a 50% response.

Upon purification, the titer of an immunoadhesin preparation may be expressed as a reciprocal dilution, or the number of milliliters to which a milligram of immunoadhesin must be diluted in order to give a 50 % response. This ELISA will facilitate the development of a purification process for the immunoadhesin.

5 A cytopathic effect assay (CPE) demonstrated the specific ability of the partially purified immunoadhesin to inhibit the infectivity of human cells by human rhinovirus (FIG. 5). Rhinovirus serotype HRV-39 was pre-incubated with human ICAM-1, an ICAM/IgA fusion (gift of Dr. Tim Springer), or with extracts from calli either expressing our ICAM-1/SlgA immunoadhesin or another, different, antibody before plating each of the
10 mixtures with HeLa S3 cells at 33°C. After 3 days, viable cells were fixed and stained with a methanolic solution of Crystal Violet; the optical density at 570 nm provides a proportional measure of cell viability.

Two extracts derived from Rhi107-11, containing the immunoadhesin, clearly inhibited the virus' ability to infect and kill HeLa S3 cells (FIG. 5A, right side-up and
15 upside-down triangles). Because the extracts were only partially purified, we also assayed a similarly prepared extract that contained a human IgA2m(2) directed against Doxorubicin, a chemotherapeutic agent. That extract, containing a similar immunoglobulin with an unrelated binding specificity, was unable to inhibit the infectivity of the rhinovirus and demonstrates that expression of the ICAM-1-heavy-chain fusion confers specificity to the inhibition. The
20 CPE assay demonstrated, as expected, the differential ability of soluble ICAM-1 and an (IC1-5/IgA; Martin, *et al.*, 1993) to inhibit viral infectivity (FIG. 5B). The insert in Figure 5B is the scale expansion in the range of the IC50 for soluble ICAM-1 (1.35 µg/ml) and for the ICI-5/IgA (0.12 µg/ml; 11.3 fold less).

25 5. Production and Purification of Immunoadhesins for Clinical and Toxicological Studies

Production of sufficient immunoadhesin for the proposed clinical and toxicological needs is performed by making transgenic tobacco plants. The transgenic plants which express

the immunoadhesin (without an ER retention signal) are generated by *Agrobacterium*-mediated transformation. The absence of an ER retention signal is anticipated to enhance assembly since the nascent SIgA is processed through the entire Golgi apparatus, including, in particular, the trans-Golgi, where SC is covalently linked to dIgA as suggested by pulse-chase experiments (Chintalacharuvu & Morrison, *Immunotechnology* 4:165-174, 1999). Because *Agrobacterium*-mediated transformation is much more likely to generate plants with consistent levels of transgene expression, it is likely that progeny of these plants will be used for the production of clinical grade immunoadhesin.

In order to maximize expression levels, and create a true-breeding line, it is desirable to create homozygous plants. The highest producing plants (generation T₀) can self-fertilize in the greenhouse before seed is collected. One quarter of the T₁ plants are expected to be homozygous. These are grown in the greenhouse and seed samples from several plants are separately germinated on medium containing kanamycin. All the progeny (T₂) from homozygous positive plants are expected to be green. Some of the progeny of heterozygous plants are expected to be white or yellowish. Homozygosity is confirmed by back-crossing to wild-type and immunoblotting extracts of the progeny.

Harvesting and processing may be continuously meshed during a production campaign, especially since multiple harvests may be obtained from a single planting, i.e. plants cut to soil level for one harvest are regrown for subsequent harvests. In developing a sense of scale for the production of immunoadhesin it is necessary to decide on the required amount of finished immunoadhesin, account for expression levels (mg immunoadhesin present/ kg fresh weight tobacco), know the growth rate of the plants and the expected weight of the average plant, and the overall yield of the purification schedule (set at 20%). Setting the overall need at 3 g of finished immunoadhesin requires preparing for 4 harvests, each with an expected yield of 1 g of finished immunoadhesin.

Given these targets and parameters, the necessary number of plants and hence the space requirements for plant growth is determined. **FIG. 6** shows an evaluation of the

production necessities for making 1 gram of finished Immunoadhesin. In this diagram, the number of plants needed for 1 g of immunoadhesin, at 20% yield, at expected levels of expression and plant weight is illustrated. At different levels of immunoadhesin expression (mg/kg fresh weight) and overall recovery (set at 20%), the weight of each plant, and so the total number of plants, may be determined for a specified production target (1 g/harvest) within a window (dotted square) of reasonable possibilities. The number of required plants decreases, inversely, with the number of specified growth and re-growth periods. The expected biomass production, a function of time and growth conditions, influences the time to harvest and the time between harvests. These growth periods can be adjusted to the realities of the purification schedule by staggering planting and harvesting dates. From our experience, production requires x number of plants. For example, 1 g of finished immunoadhesin from plants with a reasonable expression level, of 100 mg of immunoadhesin/kg fresh weight, require 250 plants when harvested at a weight of 200 g/plant (~80 days post germination). At this scale, these plants require about 10 m² of growing space and are harvested twice over 150 days.

Processing 50+ kg of biomass at a time requires several moderately large-scale operations which all have counter-parts in the food-processing industry. These include bulk materials handling, size reduction, juicing and filtration. A Vincent Press and a Durco filtration system are used to efficiently process these quantities. The juicing step employs a proven and simple buffer of sodium citrate and urea. These components buffer the extract, help prevent the oxidation of phenolics and their association with proteins (Gegenheimer, *Methods in Enzymology* 182:174-193, 1990; Loomis, *Methods in Enzymology*, 31:528-544, 1974; Van Sumere, *et al.*, *The Chemistry and Biochemistry of Plant Proteins*, 1975.) and ensure the solubility of the immunoadhesin during a subsequent acid precipitation.

Filtration of acid-insoluble lipid and protein (~90% of the total) is followed by tangential flow ultrafiltration to concentrate the immunoadhesin and to remove small proteins, especially phenolics. Diafiltration enhances the removal of small molecules and exchanges the buffer in preparation for short-term storage and subsequent chromatography. Either

SP-Sepharose (binding at pH 5.0 or below) or Q-Sepharose (binding at pH 5.5 or above) are among the ion-exchanges that can be used for filtering immunoadhesin. They are readily available, scalable, robust and have high capacities. In particular, they are available for expanded-bed formats, which reduce the stringency of prior filtration steps. Cation-exchange chromatography, which can be more selective than anion-exchange chromatography, is used first. The immunoadhesin is purified from the several species of protein potentially present, to the point where at least 95% of the protein is in the form of ICAM-1/IgA, ICAM-1/dIgA or ICAM-1/SIgA, as the presence of di- and tetra-valent ICAM-1 domains are critical for potent anti-viral activity. Purified immunoadhesin is then tested for acceptable levels of endotoxin, alkaloids such as nicotine and for bio-burden. In addition, potency levels (defined by ELISA and CPE assays), protein concentration, pH and appearance are monitored. Subsequently, the stability of the clinical lots of immunoadhesin is determined, to ensure that patients receive fully potent immunoadhesin. Even partially purified extracts have been found to be stable for 10 days when refrigerated. The titer and potency of clinically formulated immunoadhesin (in phosphate-buffered saline), when stored at -20°C , $2-8^{\circ}\text{C}$, and at 37°C , over a period of 3 to 6 months, is also tested.

6. The Immunoadhesins Have Plant-Specific Glycosylation

The immunoadhesins produced are analyzed to determine the pattern of glycosylation present. Cabanes-Macheteau *et al.*, *Glycobiology* 9(4):365-372 (1999), demonstrated the presence of several glycosyl moieties, typical of plants, on a plant-expressed antibody construct. Their methods are used to demonstrate that the immunoadhesins produced according to Example 1, 2 and 3 have a plant-specific glycosylation pattern. We anticipate that this diversity will also be a source of variability for immunoadhesin. Since crude extracts have been shown to have anti-viral activity *in vitro* (data not shown), glycosylation, as such, does not appear to affect potency. N-linked glycosylation (FIG. 2 shows that there are fifteen potential sites on the chimeric ICAM-1 molecule alone) probably contributes to the diversity of bands seen in immuno-blot. Immunoadhesin preparations are digested with N-Glycosidase A, before blotting, showing that the difference in banding patterns collapse

into fewer, discrete bands. In this way, glycoforms are initially characterized with reducing and non-reducing polyacrylamide gels. In addition, digested and mock-digested fractions are tested in the CPE assay and competition ELISA, demonstrating the effect of N-linked glycosylation on potency and titer *in vitro*.

7. The Immunoadhesin Inactivates Human Rhinovirus

The immunoadhesin prepared according to Examples 1, 2 and 3 is assayed for its ability and to inactivate HRV by binding to the virus, blocking virus entry, and inducing the formation of empty virus capsids. To measure binding of the immunoadhesin to HRV, the immunoadhesin is incubated with [³H]leucine-labeled HRV-39 for 30 min and then added to HeLa cells for 1 hr. After washing, cells and bound virus are detached with Triton X-100 and [³H] measured in a scintillation counter.

Inactivation of HRV-39 by incubation with the immunoadhesin is compared with HRV inactivation by sICAM-1. HRV-39 is not directly inactivated to a significant extent (<0.5 log₁₀ reduction in infectivity) by incubation with monomeric sICAM-1, while incubation with IC1-5D/IgA reduced infectivity approximately 1.0 log₁₀ (Arruda, *et al.*, *Antimicrob. Agents Chemother.* 36:1186-1191, 1992; Crump, *et al.*, *Antimicrob. Agents Chemother.* 38:1425-7, 1994). In order to test the ability of the immunoadhesin to inactivate HRV-39, 10⁶ 50% tissue culture infective doses (TCID₅₀) of HRV-39 are incubated in medium containing a concentration of sICAM-1 or immunoadhesin equal to ten times the IC₅₀ of each molecule for that virus, or in plain medium, for 1 hr at 33°C on a rocker platform. Each virus-immunoadhesin or virus-medium mixture are then diluted serially in ten-fold dilutions, and the titer determined on HeLa cells in 96-well plates.

The effect of the immunoadhesin on HRV attachment to host cells is tested by inoculating HeLa cells with HRV-39 at a MOI of 0.3 in the presence or absence of the immunoadhesin. Absorbance proceeds for one hour at 4°C, the cells are washed, and media is replaced plus or minus the immunoadhesin. Cells are incubated for ten hours at 33°C (to

allow one round of replication), and virus are harvested by freeze/thawing the cells. The virus is titered on HeLa cells.

ICAM-IgA (IC1-5D/IgA) is more efficient than sICAM-1 at inducing conformational changes in HRV, leading to the formation of empty, non-infectious viral particles (Martin, *et al. J. Virol.* 67:3561-8, 1993). To examine the ability of the immunoadhesin produced according to Examples 1, 2 and 3 to induce conformational changes in HRV, causing release of viral RNA, purified immunoadhesin is incubated with [³H]leucine-labeled HRV-39 for 30 min and then the virus is overlaid onto a 5 to 30% sucrose gradient. Following centrifugation for 90 min at 40,000 rpm, fractions are collected, [³H] measured, and fractions assessed for infectivity. (Intact HRV sediments at 149S on a sucrose gradient while empty capsids lacking RNA sediments at 75S (Martin, *et al. J. Virol.* 67:3561-8, 1993)). Due to its increased valence, we expect the ICAM/SlgA immunoadhesin is more efficient at inducing empty non-infectious particles than ICAM-IgA.

The inhibitory effect of purified immunoadhesin on a panel of both major and minor (that do not use ICAM-1 as a receptor) HRV serotypes will be examined using the CPE assay. The ability of ICAM-1 to inhibit HRV infection varies among viral isolates. It has been shown (Crump, *et al., Antimicrob. Agents Chemother.* 38:1425-7, 1994) that the EC₅₀ for sICAM-1 varies from 0.6 µg/ml to >32 µg/ml when tested on a panel of HRV major receptor serotypes assay using HeLa cells. Our panel includes nine major serotypes (HRV-3, -13, -14, -16, -23, -39, -68, -73, and -80) and the minor receptor serotype HRV-1A.

8. Clinical Studies Demonstrating the Ability of the Immunoadhesin to Reduce Infectivity in Humans: Dose Escalation Tolerance Study

The immunoadhesin of the present invention is tested in two randomized controlled trials to determine the effect of intranasal administration of the immunoadhesin on infection, IL-8 response, and illness in experimental rhinovirus colds. These two studies evaluate the immunoadhesin taken by subjects before or after rhinovirus inoculation. The clinical protocols used here are based on protocols previously used by in evaluation of a recombinant

soluble ICAM-1 molecule for efficacy against rhinovirus infection (Turner, *et al.*, *JAMA* 281:1797-804, 1999).

A. Subjects.

Subjects are recruited from university communities at the University of Virginia,
5 Charlottesville. Subjects are required to be in good health, non-smokers, and between the
ages of 18 and 60 years. Subjects are excluded if they have a history of allergic disease or
nonallergic rhinitis, abnormal nasal anatomy or mucosa, or a respiratory tract infection in the
previous 2 weeks. Pregnant or lactating women or women not taking medically approved
10 birth control are also excluded. In the experimental virus challenge study (Phase I/II, see
below), subjects are required to be susceptible to the study virus as evidenced by a serum
neutralizing antibody titer of 1:4 or less to the virus, determined within 90 days of the start of
the trial.

B. Study Medication.

The immunoadhesin of the present invention is formulated as a phosphate-buffered
15 saline (PBS) spray solution containing 2.6 mg/ml. The placebo consists of PBS and is
identical in appearance to the active preparation. The solutions are administered using a
medication bottle equipped with a metered nasal spray pump. The pump delivers 70 μ l of
solution containing 183 μ g of the immunoadhesin with each spray. The medication is
administered as two sprays per nostril, six times daily (at 3-hour intervals) for a total daily
20 dose of 4.4 mg. This is the same dose, in mg protein/day, as was used for soluble ICAM-1 in
the tremacamra study infection (Turner, *et al.*, *JAMA* 281:1797-804, 1999). A mole of the
immunoadhesin has about twice the mass as a mole of sICAM-1. However, given the
differences in *in vitro* activity between sICAM-1 and ICAM/IgA fusions, the immunoadhesin
is many fold more effective on a molar basis than sICAM-1. Thus, this amount is a
25 conservative calculation of what is necessary. This amount is used, except in the event that
the dose escalation study reveals problems at this dose.

C. Study Design

Single ascending dose and multiple ascending dose studies are used to evaluate the safety of the immunoadhesin. In each case, three subjects are evaluated at each dosage level, two receiving the immunoadhesin and one receiving placebo. In the single ascending dose study, four dosage levels are evaluated. The lowest individual dose is half the anticipated dose to be used in the challenge study, and the highest individual dose is twice the anticipated challenge study dose. The dosage levels are as follows: one spray in each nostril (366 µg total), two sprays in each nostril (732 µg total), three sprays in each nostril (1098 µg total), four sprays in each nostril (1464 µg total).

The same dosage levels are used in the multiple ascending dose study. Subjects receive doses every three hours (six times per day) for five days. In both studies subjects are evaluated at each dosage level, staggering the start of each subsequent level until it is clear that there is no acute toxicity at the previous level. All subjects return for a single dose 21 days after the first dose, and then for a follow-up at six weeks (for determination of serum antibody against the immunoadhesin).

A separate group of twelve subjects is given one dose of two sprays in each nostril (732 µg total), and nasal lavage is done at 1, 2, 4, 8 and 16 hours (two subjects at each time point). Washings are assayed at Panorama Research by ELISA for the immunoadhesin in order to calculate its *in vivo* half-life. The total amount of the immunoadhesin to be used in the dose escalation and half-life determination studies (on a total of 28 subjects) will be approximately 270 mg.

D. Safety Evaluations.

In addition to routine adverse event recording, the safety of the immunoadhesin is assessed in three ways. First, prior to the first dose and after the last dose the investigators perform a visual examination of the nasal mucosa, in particular looking for signs of irritation or inflammation. Any visible changes are noted. Second, standard blood safety evaluations are done on samples collected prior to treatment and after the last dose on study days 1, 4, and

8 (and 21 in the multiple ascending dose study). Third, serum samples are saved, frozen, and used to determine if the immunoadhesin is able to pass through the nasal mucosa into the blood. This is accomplished in two ways. First, the presence the immunoadhesin in serum samples is measured by ELISA. In this assay, anti-human IgA antibodies adsorbed to microtiter plates capture any the immunoadhesin in the serum, which are detected by an anti-ICAM antibody. The sensitivity of the assay is determined using normal human serum samples spiked with known concentrations of the immunoadhesin. Alternatively, anti-ICAM antibodies can be adsorbed to plates to capture the immunoadhesin in the serum, that would be detected by anti-IgA. Second, the presence of an immune response to the immunoadhesin is assayed with an ELISA method that uses the immunoadhesin adsorbed to microtiter plates. Any anti-immunoadhesin antibodies in the serum bind, and are detected with anti-human IgG or anti-human IgM. Pre-treatment and post-treatment serum samples are compared, and any change in titer is considered evidence of uptake of the immunoadhesin. If there is any positive evidence of anti-immunoadhesin antibodies, additional assays will be done to distinguish between anti-ICAM-1 and anti-IgA activity.

Patients are screened for the development of an allergic reaction to the immunoadhesin. (In previous studies, there were no episodes of adverse reactions with soluble ICAM applied topically in the nose or plantibodies applied topically in the oral cavity.) Individuals exhibiting symptoms of nasal allergy are tested for anti-immunoadhesin-specific IgE antibodies in nasal lavage fluids using a sensitive two-step ELISA (R & D Systems).

E. Statistical Analysis.

The sample size for these studies is based on previous studies using the rhinovirus challenge model. The sample size planned for the protection studies should be adequate to detect a reduction in the incidence of clinical colds from 75% in the placebo groups to 25% in the active treatment groups at 1-sided levels of $\alpha = .05$ and $1-\beta = .80$. In addition, the sample size should be adequate to detect a change in the total symptom score of 5 units assuming an SD of 5.8 units.

9. **Clinical Studies Demonstrating the Ability of the Immunoadhesin to Reduce Infectivity in Humans: Challenge Studies**

Challenge studies are used to demonstrate that treatment with the immunoadhesin of the present invention protect against clinical colds or reduce cold symptoms after viral
5 challenge.

A. **Challenge Virus.**

The challenge virus used for this study is rhinovirus 39 (HRV-39). Rhinovirus type 39 is a major group of rhinovirus that requires ICAM-1 for attachment to cells. The challenge virus pool is safety-tested according to consensus guidelines (Gwaltney, *et al.*,
10 *Prog. Med. Virol.* 39:256-263, 1992). All subjects are inoculated with approximately 200 median tissue culture infective dose (TCID₅₀). The virus are administered as drops in two inocula of 250 µl per nostril given approximately 15 minutes apart while the subjects are supine.

TABLE 1

				Pre-inoculation study timetable					
				Day					
	0	1	2	3	4	5	6	7 - 14	21
Medications		6 doses	6 doses	6 doses	6 doses	6 doses			
Inoculation		hour 4							
Symptom scores		m/e	m/e	m/e	m/e	m/e	m/e	e	
Nasal lavage		m	m	m	m	m	m		
Serum sample	X								X
				Post-inoculation study timetable					
				Day					
	0	1	2	3	4	5	6	7 - 14	21
Medications		6 doses	6 doses	6 doses	6 doses	6 doses			
Inoculation	hour 0								
Symptom scores		m/e	m/e	m/e	m/e	m/e	m/e	e	
Nasal lavage		m	m	m	m	m	m		
Serum sample	X								X
Note: In both studies on days 1-5, doses are given at hours 0, 3, 6, 9, 12, and 15									
m = morning									
e = evening									

B. Study Design.

5 Two randomized rhinovirus challenge studies are performed (see Table 1). The same formulation of the immunoadhesin of the present invention is evaluated in pre-inoculation and post-inoculation studies. In both studies, medication is administered as six doses each day for five days. Subjects are randomly assigned to receive either the immunoadhesin or matching placebo at the time of enrollment into each study. The study is blinded and all clinical trial personnel, subjects, and employees of Panorama Research remain blinded until all data are collected.

In the pre-inoculation study, medications are started four hours (two doses) prior to viral challenge. The virus challenge is administered one hour after the second dose of the

immunoadhesin (or placebo) and the four remaining doses of study medication for the first day are given as scheduled. In this study eighteen subjects receive the active treatment and eighteen subjects receive placebo.

5 In the post-inoculation study, medications begin 24 hours after virus challenge. This timepoint was chosen because it has been used in other studies of protection from virus challenge, and because cold symptoms are clearly present (Harris & Gwaltney, *Clin. Infect. Dis.* 23:1287-90, 1996). Virus challenge in this study is administered in the morning of study day 0 approximately 24 hours prior to the first dose of study medication on the morning of study day 1. In this study, 36 subjects receive the active treatment and 18 subjects receive
10 placebo.

Subjects are isolated in individual hotel rooms from study day 0 (the day of virus challenge) to study day 6. On each of these days a symptom score and a nasal lavage for virus isolation are done in the morning prior to the first dose of medication and a second symptom score is done each evening. On study day 6, subjects are released from isolation but
15 continue to record symptom scores each evening through day 14. The subjects return to the study site on study day 21, when a final serum sample for detection of anti-immunoadhesin antibodies will be collected. The total amount of immunoadhesin to be used in the two virus challenge studies (on a total of 54 subjects) is approximately 1200 mg.

C. Viral Isolation.

20 Virus shedding is detected by virus isolation in cell culture. Nasal wash specimens are collected by instillation of 5 ml of 0.9% saline into each nostril. This wash is then expelled into a plastic cup and kept chilled for one to two hours until it is processed for viral cultures. Immunoadhesin is removed from the specimens by treatment with anti-ICAM-1 antibody adsorbed to an agarose support (Affi-Gel 10, Bio-Rad Laboratories, Hercules, CA). A portion
25 of each processed specimen is stored at -80°C , and another portion is inoculated into two tubes of HeLa-1 cells, a HeLa cell line enriched for the production of ICAM-1 Arruda, *et al.*, *J. Clin. Microb.* 34:1277-1279, 1996). Rhinovirus are identified by the development of

typical cytopathic effect. Subjects with a positive viral culture on any of the postchallenge study days are considered infected. Viral titers in the specimens stored at -80°C are determined by culturing serial ten-fold dilutions in microtiter plates of HeLa-1 cells.

Antibody to the challenge virus are detected by serum neutralizing titers done using standard methods (Gwaltney, *et al*, *Diagnostic Procedures for Viral Rickettsial and Chlamydial Infections*, p. 579-614, American Public Health Association). Serum specimens for antibody testing are collected during screening, immediately prior to virus challenge (acute), and again 21 days later (convalescent). Subjects with at least a four-fold rise in antibody titer to the challenge virus when the convalescent serum sample is compared with the acute serum sample are considered infected.

D. Evaluation of Illness Severity

Illness severity is assessed as previously described (Turner, *et al.*, *JAMA* 281:1797-804, 1999). Symptom scores are recorded prior to virus challenge (baseline) and twice each day at approximately twelve-hour intervals for the next 6 days. On study days 7 through 14 each subject records his/her symptom score once per day in the evening. At each evaluation, subjects are asked to judge the maximum severity of the following eight symptoms in the interval since the last symptom evaluation: sneezing, rhinorrhea, nasal obstruction, sore throat, cough, headache, malaise, and chilliness. Each symptom is assigned a severity score of 0 to 3 corresponding to a report of symptom severity of absent, mild, moderate, or severe. If symptoms are present at baseline, the baseline symptom score will be subtracted from the reported symptom score. The higher of the two daily evaluations are taken as the daily symptom score for each symptom. The daily symptom scores for the eight individual symptoms are summed to yield the total daily symptom score. The total daily symptom scores for the first 5 days after virus challenge (study days 1-5) are summed and on the evening of study day 5, all subjects are asked, "Do you feel you have had a cold?" Subjects who had a total symptom score of at least 6 and either at least three days of

rhinorrhea or the subjective impression that they had a cold are defined as having a clinical cold.

The weight of expelled nasal secretions is determined on days 1-7 by providing all subjects with packets of preweighed nasal tissues. After the tissues are used they are stored in
5 an airtight plastic bag. Each morning the used tissues, together with any unused tissues from the original packet, are collected and weighed.

E. IL-8 Assay.

Recent studies have suggested that the host inflammatory response, particularly interleukin 8 (IL-8), may play a role in the pathogenesis of common cold symptoms due to
10 rhinovirus infection. Concentrations of IL-8 in nasal lavage are determined with a commercially available ELISA (R&D Systems, Minneapolis, Minn) as previously described (Turner, *et al.*, *JAMA* 281:1797-804, 1999).

F. Safety Evaluations.

The same evaluations are done in the challenge study as in the dose escalation study
15 described in Example 8.

G. Statistical Analysis.

Statistical analysis is performed similarly as to that described for the dose escalation study described in Example 8.

Claims

1. An immunoadhesin comprising a chimeric ICAM-1 molecule, said chimeric ICAM-1 molecule comprising:

a rhinovirus receptor protein linked to at least a portion of an immunoglobulin heavy chain;
5 and

J chain and secretory component associated with said chimeric ICAM-1 molecule.
2. The immunoadhesin of claim 1 wherein said rhinovirus receptor protein is comprised of any combination of extracellular domains 1, 2, 3, 4 and 5 of ICAM-1.
3. The immunoadhesin of claim 1 wherein said immunoglobulin is selected from the group of
10 IgA, IgA₁, IgA₂, IgM, and chimeric immunoglobulin heavy chains.
4. The immunoadhesin of claim 1 further comprising at least one additional chimeric ICAM-1 molecule.
5. The immunoadhesin of claim 1 wherein said rhinovirus receptor protein is comprised of any combination of extracellular domains 1, 2, 3, 4 and 5 of ICAM-1; and said
15 immunoglobulin heavy chain comprises at least a portion of an IgA₂ heavy chain.
6. The immunoadhesin of claim 1 expressed in transgenic plants.
7. The immunoadhesin of claim 1 expressed in monocotyledonous plants.
8. The immunoadhesin of claim 1 expressed in dicotyledonous plants.
9. The immunoadhesin of claim 1 wherein all proteins are human.
- 20 10. The immunoadhesin of claim 1 expressed in heterologous cells derived from plants, vertebrates, or invertebrates.

11. The immunoadhesin of claim 1 expressed in mammalian cells.
12. The immunoadhesin of claim 1 expressed in hairy root cultures.
13. The immunoadhesin of claim 1 expressed in plant cells in tissue culture.
14. An immunoadhesin comprising a chimeric ICAM-1 molecule, said chimeric ICAM-1
5 molecule comprising: a rhinovirus receptor protein linked to at least a portion of an immunoglobulin heavy chain, wherein immunoadhesin has plant-specific glycosylation.
15. The immunoadhesin of claim 14 wherein said immunoadhesin further comprises a J chain and secretory component associated with said chimeric ICAM-1 molecule.
16. The immunoadhesin of claim 14 wherein said rhinovirus receptor protein is comprised of
10 any combination of extracellular domains 1, 2, 3, 4 and 5 of ICAM-1.
17. The immunoadhesin of claim 14 wherein said immunoglobulin heavy chain is selected from the group of IgA, IgA₁, IgA₂, IgG₁, IgG₂, IgG₃, IgG₄, IgM, IgD, IgE, and a chimeric
immunoglobulin heavy chain.
18. The immunoadhesin of claim 14 further comprising at least one additional chimeric
15 ICAM-1 molecule.
19. The immunoadhesin of claim 14 wherein said rhinovirus receptor protein is comprised of any combination of extracellular domains 1, 2, 3, 4 and 5 of ICAM-1; and said immunoglobulin heavy chain comprises at least a portion of an IgA₂ heavy chain.
20. The immunoadhesin of claim 14 wherein all proteins are human.
- 20 21. The immunoadhesin of claim 14 expressed in heterologous cells derived from plants.
22. The immunoadhesin of claim 14 expressed in hairy root cultures.
23. The immunoadhesin of claim 14 expressed in plant cells in tissue culture.

24. The immunoadhesin of claim 14 expressed in plants.
25. The immunoadhesin of claim 14 expressed in monocotyledonous plants.
26. The immunoadhesin of claim 14 expressed in dicotyledonous plants.
27. A composition comprising an immunoadhesin and plant material, wherein
- 5 said immunoadhesin comprises a chimeric ICAM-1 molecule, said chimeric ICAM-1 molecule comprising a rhinovirus receptor protein linked to at least a portion of an immunoglobulin heavy chain.
28. The composition of claim 27 further comprising a J chain and secretory component associated with said chimeric ICAM-1 molecule.
- 10 29. A composition of claim 27 wherein said rhinovirus receptor protein is comprised of any combination of extracellular domains 1, 2, 3, 4 and 5 of ICAM-1; and said immunoadhesin has plant-specific glycosylation.
30. A composition of claim 27 wherein said immunoglobulin is selected from the group of IgA, IgA₁, IgA₂, IgG₁, IgG₂, IgG₃, IgG₄, IgM, IgD, IgE, and a chimeric immunoglobulin
- 15 heavy chain.
31. A composition of claim 27 further comprising at least one additional chimeric ICAM-1 molecule.
32. A composition of claim 27 wherein said rhinovirus receptor protein is comprised of any combination of extracellular domains 1, 2, 3, 4 and 5 of ICAM-1; and
- 20 said immunoglobulin heavy chain is an IgA₂ heavy chain
33. A method for reducing the infection by human rhinovirus of host cells susceptible to infection by human rhinovirus, said method comprising:

contacting the virus with an immunoadhesin of claim 1, 14, or 27, and wherein
said immunoadhesin binds to human rhinovirus and reduces infectivity thereof.

34. A method for reducing the initiation or spread of the common cold due to human
rhinovirus, said method comprising:

5 contacting the virus with an immunoadhesin of claim 1, 14 or 27, and wherein
said immunoadhesin binds to human rhinovirus and reduces infectivity thereof.

35. A method for the treatment or prevention of human rhinovirus infection in a human
subject, said method comprising:

administering to said subject an effective amount of an immunoadhesin of claim 1, 14 or 27,
10 and wherein said immunoadhesin reduces human rhinovirus infectivity thereof.

36. A method for the treatment or prevention of human rhinovirus infection in a subject, said
method comprising:

intranasally administering to said subject an effective amount of an immunoadhesin of claim
1, 14 or 27, and wherein said immunoadhesin reduces human rhinovirus infectivity thereof.

15 37. A method for the treatment or prevention of human rhinovirus infection in a subject, said
method comprising: administering through the oral cavity to said subject an effective amount
of an immunoadhesin of claim 1, 14 or 27, and wherein said immunoadhesin reduces human
rhinovirus infectivity thereof.

38. A pharmaceutical composition comprising an immunoadhesin of claims 1, 14, or 27 in a
20 pharmaceutically acceptable buffer.

39. An expression vector comprising a gene encoding a chimeric ICAM-1 molecule
operatively linked to a plant promoter, said chimeric ICAM-1 molecule comprising a
rhinovirus receptor protein linked to at least a portion of an immunoglobulin heavy chain.

FIGURE 1





Figure 2


A.


	Spe I		Spe I	
SuperMas Promoter	Signal Peptide	ICAM-1 Extracellular Domains	Cα1-Cα3 of Human IgA2m(2)	NOS 3' Terminator

B.

QTSVSPSKVILPRGGSVLVTCSTSCDQPKLLGIETPLPKKELLLPGNNRKVYELSNVQEDSQPMC
 YSNCPDGQSTAKTFLTVYWTPERVELAPLPSWQPVGKNLTLRCQVEGGAPRANLTVVLLRGEKELKREPA
 VGEPAEVTTTVLVRRDHHGANFSCRTELDLRPQGLELFENTSAPYQLQTFVLPATPPQLVSPRVLEVDTQ
 GTVVCSLDGLFPVSEAQVHLALGDQRLNPTVTYGNDSFSAKASVSVTAEDEGTQRLTCAVILGNQSQETL
 QTVTIYSFPAPNVILTKEVSEGTEVTVKCEAHPRAKVTLNGVPAQPLGPRAQLLLKATPEDNGRSFSCS
 ATLEVAGQLIHKNQTRRLRVLYGPRLDERDCPGNWTWPENSQQTPMCQAWGNPLPELKCLKDGTFFPLPIG
 ESVTVTRDLEGTYLCRARSTQGEVTREVTVNVTSGSSASPTSPKVFPLSLDSTPQDGNVVVACLVOGFFP
 QEPLSVTWSESGQNVITARNFPPSQDASGDLYTTSSQLTLPATQCPDGKSVTCHVKHYTNSSQDVTVPCR
 VPPPPCCHPRLSLHRPALEDLLLGSEANLTCTLTGLRDASGATFTWTPSSGKSAVQGPPERDLCCGCYSVS
 RVLPGCAQPWNHGETFTCTAAHPELKTPLTANITKSGNTFRPEVHLLPPPSEELALNELVTLTCLARGFS
 PKDVLVRWLQGSQELPREKYLTWASRQEPSQGTTTYAVTSILRVAAEDWKKGETFSCMVGHEALPLAFTQ
 KTIDRLAGKPTHINVSVMMAEADGTCYRSEKDEL

FIGURE 3

FIGURE 4

**FIGURE 5**




FIGURE 6

FIGURE 7 A

I. HUMAN IG ALPHA-1 CHAIN C REGION - HOMO SAPIENS (HUMAN).

AMINO ACID SEQUENCE

>sp|P01876|ALC1_HUMAN IG ALPHA-1 CHAIN C REGION - Homo sapiens (Human).

```

      10      20      30      40      50      60
      |      |      |      |      |      |
ASPTSPKVFP LSLCSTQPDG NVVIACLVQG FEPQEPLSVT WSESGQGVTA RNFPPSQDAS

      70      80      90     100     110     120
      |      |      |      |      |      |
GDLYTTSSQL TLPATQCLAG KSVTCHVKHY TNPSQDVTVP CPVPSTPPTP SPSTPPTPSP

     130     140     150     160     170     180
      |      |      |      |      |      |
SCCHPRLSLH RPALEDLLLG SEANLTCTLT GLRDASGVTF TWTPSSGKSA VQGPPERDLC

     190     200     210     220     230     240
      |      |      |      |      |      |
GCYSVSSVLP GCAEPWNHGK TFTCTAAYPE SKTPLTATLS KSGNTERPEV HLLPPPSEEL

     250     260     270     280     290     300
      |      |      |      |      |      |
ALNELVTLTC LARGFSPKDV LVRWLQGSQE LPREKYLTWA SRQEPSQGT TFAVTSILRV

     310     320     330     340     350
      |      |      |      |      |
AAEDWKKGDT FSCMVGHEAL PLAFTQKTID RLAGKPTHVN VSVVMAEVDG TCY

```

CODING SEQUENCE

```

catccccgac cagccccaa gtcctccgc tgagcctctg cagcaccag ccagatggga      g      -1
acgtggatcat cgcctgcctg gtccagggt tcttcccca ggagccactc agtgtgacct      60
ggagcgaaag cggacagggc gtgaccgcca gaaacttccc acccagccag gatgcctccg      120
gggacctgta caccacgagc agccagctga ccctgccggc cacacagtgc ctagccggca      180
agtcctgtgac atgccacgtg aagcactaca cgaatcccag ccaggatgtg actgtgccct      240
gcccagttcc ctcaactcca cctaccccat ctccctcaac tccacctacc ccatctccct      300
catgctgcca ccccgaactg tcaactgcac gaccggccct cgaggacctg ctcttaggtt      360
cagaagcgaa cctcacgtgc acaactgacc gcctgagaga tgccctcagg gtcaccttca      420
cctggacgcc ctcaagtggg aagagcgctg ttcaaggacc acctgagcgt gacctctgtg      480
gctgctacag cgtgtccagt gtccctgccg gctgtgccga gccatggaac catgggaaga      540
ccttcacttg cactgctgcc taccctgagt ccaagacccc gctaaccgcc accctctcaa      600
aatccggaaa cacattccgg cccgagggtc acctgctgcc gccgccgtcg gaggagctgg      660
ccctgaacga gctggtgacg ctgacgtgcc tggcacgcgg cttcagcccc aaggacgtgc      720
tggttcgctg gctgcagggg tcacaggagc tgccccgcga gaagtacctg acttgggcat      780
cccggcagga gccagccag ggcaccacca ccttcgctgt gaccagcata ctgogcgtgg      840
cagccgagga ctggaagaag ggggacacct tctcctgcat ggtggggcac gaggccctgc      900
cgctggcctt cacacagaag accatcgacc gcttggcggg taaaccacc catgtcaatg      960
tgtctgttgt catggcggag gtggacggca cctgctactg a      1020
                                         1061

```

GenBank

J00220
LOCUS HUMIGCC8 2533 bp DNA PRI 02-DEC-1998
DEFINITION Homo sapiens immunoglobulin alpha-1 heavy chain constant region
(IGHA1) gene, partial cds.
ACCESSION J00220
VERSION J00220.1 GI:184743
KEYWORDS .
SOURCE human.

FIGURE 7 B

ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.

REFERENCE 1 (bases 1 to 2533)

AUTHORS Takahashi,N., Ueda,S., Obata,M., Nikaido,T., Nakai,S. and Honjo,T.

TITLE Structure of human immunoglobulin gamma genes: Implications for
 evolution of a gene family

JOURNAL Cell 29, 671-679 (1982)

MEDLINE 83001943

COMMENT This sequence is part of a multigene region containing the
 immunoglobulin heavy chain gamma-3, gamma-1, pseudo-epsilon, and
 alpha-1 genes.

FEATURES

	Location/Qualifiers
source	1..2533 /organism="Homo sapiens" /db_xref="taxon:9606" /chromosome="14" /map="14q32.33" /clone="cosmid Ig13; Ig-gamma3-122" /tissue_type="placenta; liver" /germline
intron	<1..141 /note="alpha-1 intron J-C"
exon	142..447 /gene="IGHA1"
gene	<142..>1638 /gene="IGHA1"
CDS	join(<142..447,662..1021,1244..1638) /gene="IGHA1" /codon_start=3 /product="immunoglobulin alpha-1 heavy chain constant region" /protein_id="AAC82528.1" /db_xref="GI:184749" /translation="SPTSPKVFPLSLCSTQPDGNVVIACLVQGFFPQEPLSVTWSESG QGV TARNFPPSQDASGDLYTTSSQLTLPATQCLAGKSVTCHVKHYTNPSQDVTVP PCPV PSTPPTPSPSTPPTPSPSCCHPRLSLHRPALEDLLLGSEANLTCTLTGLRDASGVTFT WTPSSGKSAVQGPPELDLCGCYSVSSVLPGCAEPWNHGKTFTCTAAYPESKTPLTATL SKSGNTFRPEVHLLPPPSEELALNELVTLTCLARGFSPKDVLRWLQGSQELPREKYL TWASRQEPSQGTTFFAVTSILRVAEDWKKGDTFSCMVGHEALPLAFTQKTIDRLAGK PTHVNVSVVMAEVDGTCY"
intron	448..661 /gene="IGHA1" /note="A"
exon	662..1021 /gene="IGHA1"
intron	1022..1243 /gene="IGHA1" /note="B"
exon	1244..>1638 /gene="IGHA1"

BASE COUNT 490 a 866 c 753 g 424 t

ORIGIN

```

1 ggtccaactg caggcctgtg gtgcaggagc tgtgtgacca tggggctgtc accaggcctc
61 tctgtgctgg gttcctccag tatagaggag aggcagtata gaggagaggg ccgcgtcctc
121 acagtgcatt ctgtgttcca gcatccccga ccagcccca ggtcttcccg ctgagcctct
181 gcagcaccca gccagatggg aacgtggtca tcgcctgcct ggtccagggc ttcttcccc
241 aggagccact cagtgtgacc tggagcgaaa gcggacaggg cgtgaccgcc agaaacttcc
301 caccagcca ggatgcctcc ggggacctgt acaccacgag cagccagctg accctgccgg
361 ccacacagtg cctagccggc aagtcctgta catgccacgt gaagcactac acgaatcca
421 gccaggatgt gactgtgccc tgcccaggtc agagggcagg ctggggagtg gggcggggcc
481 acccgtcgt gccctgacac tgcgcctgca cccgtgttcc ccacaggag cgcctcttc
541 actcacacca gaggggagcc cggggcagac cccaggaggt ggtggtggac aggccaggag

```

FIGURE 7 C

```

601 gggcgaggcg ggggcatggg gaagtatgtg ctgaccagct caggccatct ctccactcca
661 gtccctcaa ctccacctac cccatctccc tcaactccac ctaccccatc tccctcatgc
721 tgccaccccc gactgtcact gcaccgaccg gccctcgagg acctgctctt aggttcagaa
781 gcgaacctca cgtgcacact gaccggcctg agagatgcct cagggtgtcac ctccacctgg
841 acgcccctcaa gtgggaagag cgctgttcaa ggaccacctg agcgtgacct ctgtggctgc
901 tacagcgtgt ccagtgtcct gccgggctgt gccgagccat ggaacctatg gaagaccttc
961 acttgactg ctgcctaccc cgagtccaag accccgctaa cggccaccct ctcaaatcc
1021 ggtgggtcca gacctgctc ggggcctgc tcagtgtctt ggtttgcaa gcatattcct
1081 ggctgcctc ctccctccca atcctgggt ccagtgtctc tgccaagtac acagggaaac
1141 tgaggcaggc tgaggggcca ggacacagcc cggggtgcc accagagcag aggggctctc
1201 tcatccctg cccagccccc tgacctggct ctctaccctc caggaaacac attccggccc
1261 gaggtccacc tgctgccgc gccgtcggag gagctggccc tgaacgagct ggtgacgctg
1321 acgtgcctgg cacgcggctt cagccccaag gacgtgctgg ttgctggct gcaggggtca
1381 caggagctgc cccgcgagaa gtacctgact tgggcatccc ggcaggagcc cagccagggc
1441 accaccact tcgctgtgac cagcatactg cgcgtggcag ccgaggactg gaagaagggg
1501 gacaccttct cctgcatggt gggccacgag gccctgccgc tggccttcac acagaagacc
1561 atcgaccgct tggcgggtaa accaccccat gtcaatgtgt ctggtgtcat ggcggaggtg
1621 gacggcacct gctactgagc cgcccgctg tccccacccc tgaataaact ccatgctccc
1681 ccaagcagcc ccacgcttcc atccggcgcc tgtctgtcca tcctcagggg ctccagcactt
1741 gggaaagggc cagggcatgg acaggggaaga atacccctg ccctgagcct cggggggccc
1801 ctggcacccc catgagactt tccacctgg tgtgagtgtg agttgtgagt gtgagagtgt
1861 gtggtgcagg aggcctcgct ggtgtgagat cttaggctctg ccaaggcagg cacagcccag
1921 gatgggttct gagagacgca catgccccgg acagttctga gtgagcagtg gcatggccgt
1981 ttgtccctga gagagccgcc tctggctgta gctgggaggg aataggagg gtaaaaggag
2041 caggctagcc aagaaaggcg caggtagtgg caggagcggc gaggagtgta ggggctggac
2101 tccagggccc cactgggagg acaagctcca ggaggcccc accacctag tgggtgggccc
2161 tcaggacgtc ccactgacgc atgcaggaag gggcacctcc ccttaaccac actgctctgt
2221 acggggcacg tgggcacagg tgcacactca cactcacata tatgcctgag ccctgcagga
2281 gcggaacgtt cacagcccag acccagttcc agaaaagcca ggggagtccc ctccaagcc
2341 cccaagctca gcctgctccc ctaggccct ctggcttccc tgtgtttcca ctgtgcacag
2401 atcaggcacc aactccacag acccctccca ggcagccct gctccctgcc tggccaagtc
2461 tccatccct tcctaagccc aactaggacc caaagcatag acaggagggg gccacgtggg
2521 gtggcatcag aag

```

II. HUMAN IG ALPHA-2 CHAIN C REGION - HOMO SAPIENS (HUMAN).**AMINO ACID SEQUENCE**

>sp|P01877|ALC2_HUMAN IG ALPHA-2 CHAIN C REGION - Homo sapiens (Human).

```

      10      20      30      40      50      60
      |      |      |      |      |      |
ASPTSPKVFP LSLDSTPQDG NVVVACL VQG FFPQEPLSVT WSESGQNVTA RNFPPSQDAS
      70      80      90     100     110     120
      |      |      |      |      |      |
GDLYTTSSQL TLPATQCPDG KSVTCHVKHY TNPSQDVTVP CPVPPPPPC HPRLSLHRPA
     130     140     150     160     170     180
      |      |      |      |      |      |
LEDLLLGSEA NLTCTLTGLR DASGATFTWT PSSGKSAVQG PPERDLGCGY SVSSVLPGCA
     190     200     210     220     230     240
      |      |      |      |      |      |
QPWNHGETFT CTAAPHELKT PLTANITKSG NTFRPEVHLL PPPSEELALN ELVTLTCLAR
     250     260     270     280     290     300
      |      |      |      |      |      |
GFSPKDVLR WLQGSQELPR EKYL TWASRQ EPSQGTTF VTSILRVAE DWKKGDTFSC

```

FIGURE 7 D

310 320 330 340
 | | | |
 MVGHEALPLA FTQKTIDRLA GKPTHVNVSV VMAEVDGTCY

CODING SEQUENCE

catccccgac	cagccccaag	gtcttcccg	tgagcctcga	cagcaccccc	caagatggga	g	-1
acgtggtcgt	cgcctgcctg	gtccagggct	tcttccccc	ggagccactc	agtgtgacct		60
ggagcgaaag	cggacagAAC	gtgaccgcca	gaaacttccc	acctagccag	gatgcctccg		120
gggacctgta	caccacgagc	agccagctga	ccctgcccgc	cacacagtgc	ccagacggca		180
agtccgtgac	atgccacgtg	aagcactaca	cgaatcccag	ccaggatgtg	actgtgccct		240
gccacgttcc	cccacctccc	ccatgctgcc	acccccgact	gtcgtgtcac	cgaccggccc		300
tccaggacct	gctcttaggt	tcagaagcga	acctcactgt	cacactgacc	ggcctgagag		360
atgcctctgg	tgccaccttc	acctggacgc	cctcaagtgg	gaagagcgct	gttcaaggac		420
cacctgagcg	tgacctctgt	ggctgctaca	gcgtgtccag	tgtcctgcct	ggctgtgccc		480
agccatggaa	ccatggggag	accttcacct	gcactgctgc	ccaccccgag	ttgaagacct		540
cactaaccgc	caacatcaca	aaatccggaa	acacattccg	gcccgaggtc	cacctgctgc		600
cgccgcgcgc	ggaggagctg	gccctgaacg	agctggtgac	gctgacgtgc	ctggcacgtg		660
gcttcagccc	caaggatgtg	ctgggttcgct	ggctgcaggg	gtcacaggag	ctgccccgcg		720
agaagtacct	gacttgggca	tcccggcagg	agcccagcca	gggcaccacc	accttcgctg		780
tgaccagcat	actgcgcgtg	gcagccgagg	actggaagaa	gggggacacc	ttctcctgca		840
tggtgggcca	cgaggccctg	ccgctggcct	tcacacagaa	gaccatcgac	cgcttggcgg		900
gtaaaccac	ccatgtcaat	gtgtctgttg	tcatggcggg	ggtggacggc	acctgctact		960
ga							1020
							1022

GenBank

J00221 Human Ig germline
 LOCUS HUMIGCD7 2516 bp DNA PRI 11-APR-2001
 DEFINITION Human Ig germline H-chain G-E-A region B: alpha-2 A2m(1) allele
 constant region, 3' end.
 ACCESSION J00221
 VERSION J00221.1 GI:184756
 KEYWORDS
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.
 REFERENCE 1 (bases 1 to 2516)
 AUTHORS Ellison, J. and Hood, L.
 TITLE Linkage and sequence homology of two human immunoglobulin gamma
 heavy chain constant region genes
 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 79 (6), 1984-1988 (1982)
 MEDLINE 82197621
 PUBMED 6804948
 REFERENCE 2 (bases 737 to 1016)
 AUTHORS Flanagan, J.G. and Rabbitts, T.H.
 TITLE Arrangement of human immunoglobulin heavy chain constant region
 genes implies evolutionary duplication of a segment containing
 gamma, epsilon and alpha genes
 JOURNAL Nature 300 (5894), 709-713 (1982)
 MEDLINE 83088998
 PUBMED 6817141
 REFERENCE 3 (bases 49 to 229; 425 to 514)
 AUTHORS Hisajima, H., Nishida, Y., Nakai, S., Takahashi, N., Ueda, S. and
 Honjo, T.
 TITLE Structure of the human immunoglobulin C epsilon 2 gene, a truncated
 pseudogene: implications for its evolutionary origin
 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 80 (10), 2995-2999 (1983)
 MEDLINE 83221529
 PUBMED 6407005
 REFERENCE 4 (bases 1 to 2516)
 AUTHORS Flanagan, J.G., Lefranc, M.P. and Rabbitts, T.H.

FIGURE 7

E

TITLE Mechanisms of divergence and convergence of the human immunoglobulin alpha 1 and alpha 2 constant region gene sequences

JOURNAL Cell 36 (3), 681-688 (1984)

MEDLINE 84130179

PUBMED 6421489

COMMENT [3] also reports the complete alpha-1 gene and part of the A2m(2) alpha-2 allele (bases 737-2516; see Sites table). Comparison of the three sequences suggests that the A2m(1) alpha-2 allele might be a hybrid of the alpha-1 gene and A2m(2) alpha-2 allele. The hinge region in the alpha genes occurs at beginning of the CH2 domain. The alpha-1 hinge region is 13 amino acids longer than that in alpha-2. Both hinge regions consist of approximate tandem repeats of a 15 bp sequence. The first repeat occurs 5' to the mRNA splice site and is non-coding. The authors [3] suggest that this repetitive structure provides a possible mechanism for the large number of variations observed in hinge regions. There is a coupled 30 bp insertion, 9 bp deletion in alpha-2 relative to alpha-1 (starting at base 97). [1] also reports sequences for the epsilon-1 and epsilon-2 (pseudogene) C-region genes. The authors [1] determined the physical linkage between epsilon-1 and alpha-2 and that between epsilon-2 and alpha-1. [2] also reports the alpha-1 CH2 domain and epsilon-2. This entry is part of a multigene region (region B), which includes the gamma-2, gamma-4, epsilon-1 and alpha-2 genes. See segment 1 for more comments. Complete source information: Human genomic DNA, cosmid Ig10 [1],[3]; placenta DNA [2] clone H-Ig-alpha-25; genomic DNA from TOU II-5 library clone lambda-TOU-alpha2 (for A2m(2) allele) [3].

FEATURES Location/Qualifiers

source 1..2516
/organism="Homo sapiens"
/db_xref="taxon:9606"
/map="14q32.33"
/germline

gene <1..1621
/gene="IgH"
/note="IGHA2"

intron <1..163
/gene="IgH"
/note="alpha-2 intron J-C"

CDS join(<164..469,684..1004,1227..1621)
/gene="IgH"
/note="contains constant region"
/codon_start=3
/product="immunoglobulin alpha-2 heavy chain"
/protein_id="AAB59396.1"
/db_xref="GI:184761"
/translation="SPTSPKVFPLSLDSTPQDGNVVVACLVOGFFPQEPLSVTWSESG
QNV TARNFPPSQDASGDLYTTSSQLTLTPATQCPDGKSVTCHVKHYTNPSQDVTVECPV
PPPPPCCHPRLSLHRPALEDLLGSEANLTCTLTGLRDASGATFTWTFSSGKSAVQGP
PERDLGCGYSVSSVLPGCAQPWNHGETFTCTAAHPELKTFLTANITKSGNTFRPEVHL
LPPPSEELALNELVTLTCLARGFSPKDVLRWLQGSQELPREKYLTVASRQEPSQGT
TFAVTSILRVAEDWKKGDTFSCMVGHEALPLAFTQKTIDRLAGKP THVNVSVVMAEV
DGTCY"

exon 164..469
/gene="IgH"
/note="G00-119-333"

intron 470..683
/gene="IgH"
/note="alpha-2 intron A"

exon 684..1004

FIGURE 7 F

```

intron      /gene="IgH"
            1005..1226
            /gene="IgH"
            /note="alpha-2 intron B"
exon        1227..1621
            /gene="IgH"
variation   1434
            /gene="IgH"
            /note="t in A2m(1); a in A2m(2)"
variation   1441
            /gene="IgH"
            /note="g in A2m(1); a in A2m(2)"
variation   1465
            /gene="IgH"
            /note="c in A2m(1); t in A2m(2)"
variation   1486
            /gene="IgH"
            /note="c in A2m(1); g in A2m(2)"
variation   1553
            /gene="IgH"
            /note="t in A2m(1); a in A2m(2)"
variation   1573..1574
            /gene="IgH"
            /note="tg in A2m(1); ca in A2m(2)"
variation   1602..1606
            /gene="IgH"
            /note="tggac in A2m(1); cggat in A2m(2)"
variation   2060
            /note="c in A2m(1); t in A2m(2)"
variation   2384
            /note="a in A2m(1); c in A2m(2)"
variation   2390
            /note="c in A2m(1); g in A2m(2)"
BASE COUNT 488 a    861 c    754 g    413 t
ORIGIN
1 ggtccaaccg caggcccatg gtgcaggagc tgtgtaacct atggggctgt caccaggcct
61 ctctgtgctg gggttcctcca gtgtagagga gaggcaggta cagcctgtcc tcctggggag
121 atggcatgag ggccgcgtcc tcacagcgca ttctgtgttc cagcatcccc gaccagcccc
181 aaggtcttcc cgctgagcct cgacagcacc cccaagatg ggaacgtggt cgtcgcatgc
241 ctggtccagg gcttcttccc ccaggagcca ctacagtgtga cctggagcga aagcggacag
301 aacgtgaccg ccagaaactt cccacctagc caggatgcct ccggggacct gtacaccacg
361 agcagccagc tgaccctgcc ggccacacag tgcccagacg gcaagtcctg gacatgccac
421 gtgaagcact acacgaatcc cagccaggat gtgactgtgc cctgcccagg tcagagggca
481 ggctggggag tggggcgggg ccaccccgtc ctgccctgac actgcgcctg caccctgtgt
541 cccacagggg agccgcccct tcactcacac cagagtggac cccgggccga gcccagggag
601 gtggtggtgg acaggccagg aggggagcag cgggggacag ggggaaggcg ttctgaccag
661 ctgagcccat ctctccactc cagtcccccc acctccccca tgctgccacc cccgactgtc
721 gctgcaccga ccggccctcg aggacctgct cttaggttca gaagcgaacc tcacgtgcac
781 actgaccggc ctgagagatg cctctgggtg cacttccacc tggacgcccct caagtgggaa
841 gagcgctgtt caaggaccac ctgagcgtga cctctgtggc tgctacagcg tgtccagtgt
901 cctgcctggc tgtgcccagc catggaacca tggggagacc ttcacctgca ctgctgcca
961 ccccgagttg aagaccccac taaccgcaa catcacaaa tccggtgggt ccagaccctg
1021 ctccggggccc tgctcagtgc tctggtttgc aaagcatatt cccggcctgc ctctccctc
1081 ccaatcctgg gctccagtgc tcatgccaag tacacaggga aactgaggca ggctgagggg
1141 ccaggacaca gccaggggtg cccaccagag cagaggggct ctctcatccc ctgcccagcc
1201 ccctgacctg gctctctacc ctccaggaaa cacattccgg cccgaggtcc acctgctgcc
1261 gccgccgtcg gaggagctgg ccctgaacga gctggtgacg ctgacgtgcc tggcacgtgg
1321 cttcagcccc aaggatgtgc tggttcgctg gctgcagggg tcacaggagc tgcccgcga
1381 gaagtacctg acttgggcat cccggcagga gccagccag ggcaccacca ccttcgctgt
1441 gaccagcata ctgcgcgtgg cagccgagga ctggaagaag ggggacacct tctcctgcat
1501 ggtggggccac gagggcctgc cgctggcctt cacacagaag acctcgacc gcttggcggg
1561 taaaccaccc catgtcaatg tgtctgttgt catggcggag gtggacggca cctgctactg
1621 agccgcccgc ctgtccccc cctgaataa actccatgct cccccaagca gcccacgct

```

FIGURE 7**G**

```

1681 tccatccggc gcctgtctgt ccatectcag ggtctcagca cttgggaaag ggccagggca
1741 tggacagggg agaatacccc ctgccctgag cctcgggggg cccctggcac ccccatgaga
1801 ctttccaccc tgggtgtgagt gtgagttgtg agtgtgagag tgtgtgggtgc aggaggcctc
1861 gctgggtgtga gatcttaggt ctgccaaggc aggcacagcc caggatgggt tctgagagac
1921 gcacatgccc cggacagtgc tgagttagca gtggcatggc cgtttgtccc tgagagagcc
1981 gcctctggct gtagctggga gggaataggg agggtaaaag gagcaggcta gccaagaaag
2041 gcgcaggtag tggcaggagc ggcgaggag tgaggggctg gactccaggc cccactggg
2101 aggacaagct ccaggagggc cccaccaccc tagtgggtgg gcctcaggac gtcccactga
2161 cgcattgcagg aaggggcacc tccccttaac cacactgctc tgtacggggc acgtgggcac
2221 acatgcacac tcacactcac atatacgct gagccctgca ggagtggaaac gttcacagcc
2281 cagaccaggt tccagaaaag ccaggggagt cccctcccaa gcccacaagc tcagcctgct
2341 cccccaggcc cctctggctt cctgtgttt cactgtgca cagatcaggc accaactcca
2401 cagaccctc ccaggcagcc cctgtctcct gcctggccaa gtctcccatc ccttcctaag
2461 cccaactagg acccaaagca tagacaggga ggggcgcgt ggggtggcat cagaag

```

III. HUMAN IG GAMMA-1 CHAIN C REGION - HOMO SAPIENS (HUMAN)**AMINO ACID SEQUENCE**

>sp|P01857|GC1_HUMAN IG GAMMA-1 CHAIN C REGION - Homo sapiens (Human).

```

      10      20      30      40      50      60
      |      |      |      |      |      |
ASTKGPSVFP LAPSSKSTSG GTAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS

      70      80      90     100     110     120
      |      |      |      |      |      |
GLYSLSSVVT VPSSSLGTQT YICNVNHKPS NTKVDKKVEP KSCDKTHTCP PCPAPELLGG

     130     140     150     160     170     180
      |      |      |      |      |      |
PSVFLFPPKP KDTLMIS RTP EVTCVVVDVS HEDPEVKFNW YVDGVEVHNA KTKPREEQYN

     190     200     210     220     230     240
      |      |      |      |      |      |
STYRVVSVLT VLHQDWLNGK EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSRDE

     250     260     270     280     290     300
      |      |      |      |      |      |
LTKNQVSLTC LVKGFYPSDI AVEWESNGQP ENNYKTPPV LDSDGSFFLY SKLTVDKSRW

     310     320     330
      |      |      |
QQGNVFSCSV MHEALHNHYT QKSLSLSPGK

```

CODING SEQUENCE

```

cctccaccaa gggcccatcg gtcttcccc tggcacccct ctccaagagc acctctgggg      g      -1
gcacagcggc cctgggctgc ctggtcaagg actacttccc cgaaccgggtg acggtgtogt      60
ggaactcagg cgccctgacc agcggcgctgc acaccttccc ggctgtccta cagtcctcag      120
gactctactc cctcagcagc gtggtgaccg tgccctccag cagcttgggc acccagacct      180
acatctgcaa cgtgaatcac aagcccagca acaccaaggt ggacaagaaa gttgagccca      240
aatcttgtga caaaactcac acatgcccac cgtgccagc acctgaactc ctgggggggac      300
cgtcagtcct cctcttcccc ccaaaaccca aggacaccct catgatctcc cggaccctg      360
aggtcacatg cgtgggtgggt gacgtgagcc acgaagaccc tgaggtaag ttcaactgg      420
acgtggacgg cgtggagggt cataatgcca agacaaagcc gcgggaggag cagtacaaca      480
gcacgtaccg ggtggtcagc gtcttcaccg tcctgcacca ggactggctg aatggcaagg      540
agtacaagtg caaggctctc aacaaagccc tcccagcccc catcgagaaa accatctcca      600
aagccaaagg gcagccccga gaaccacagg tgtacaccct gcccctatcc cgggatgagc      660
                                         720

```

FIGURE 7 H

```

tgaccaagaa ccaggtcagc ctgacctgcc tgggtcaaagg cttctatccc agcgacatcg      780
ccgtggagtg ggagagcaat gggcagccgg agaacaacta caagaccacg cctcccgtgc      840
tggactccga cggctccttc ttcctctaca gcaagctcac cgtggacaag agcaggtggc      900
agcaggggaa cgtcttctca tgctccgtga tgcattgaggc tctgcacaac cactacacgc      960
agaagagcct ctccctgtct ccgggtaaat ga                                  992

```

GenBank

J00228.

LOCUS HUMIGCC4 2009 bp DNA PRI 02-DEC-1998

DEFINITION Homo sapiens immunoglobulin gamma-1 heavy chain constant region (IGHG1) gene, partial cds.

ACCESSION J00228

VERSION J00228.1 GI:184739

KEYWORDS .

SOURCE human.

ORGANISM Homo sapiens
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.

REFERENCE 1 (bases 1 to 2009)

AUTHORS Takahashi,N., Ueda,S., Obata,M., Nikaido,T., Nakai,S. and Honjo,T.

TITLE Structure of human immunoglobulin gamma genes: Implications for evolution of a gene family

JOURNAL Cell 29, 671-679 (1982)

MEDLINE 83001943

COMMENT This sequence is part of a multigene region containing the immunoglobulin heavy chain gamma-3, gamma-1, pseudo-epsilon, and alpha-1 genes.

FEATURES

source Location/Qualifiers

1..2009

/organism="Homo sapiens"

/db_xref="taxon:9606"

/chromosome="14"

/map="14q32.33"

/clone="cosmid Ig13; Ig-gamma3-122"

/tissue_type="placenta; liver"

/germline

gene <1..>1803

/gene="IGHG1"

intron <1..209

/gene="IGHG1"

CDS join(<210..503,892..936,1055..1384,1481..1803)

/gene="IGHG1"

/codon_start=3

/product="immunoglobulin gamma-1 heavy chain constant region"

/protein_id="AAC82527.1"

/db_xref="GI:184747"

/translation="STKGPSVFPPLAPSSKSTSGGTAALGCLVKDYFPEPVTFSWNSGA
LTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSC
DKTHTCTPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWY
VDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTII
SKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT
PPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQSVMEALHNHYTQKSLSLSPGK"

misc_difference 563

/gene="IGHG1"

/replace=""

misc_difference 593

/gene="IGHG1"

/replace=""

misc_difference 614

/gene="IGHG1"

/replace=""

FIGURE 7

I

```

misc_difference 633
                  /gene="IGHG1"
                  /replace=""
misc_difference 643
                  /gene="IGHG1"
                  /replace=""
misc_difference 654
                  /gene="IGHG1"
                  /replace=""
misc_difference 684
                  /gene="IGHG1"
                  /replace=""
misc_difference 692
                  /gene="IGHG1"
                  /replace=""
misc_difference 765..766
                  /gene="IGHG1"
                  /replace=""
misc_difference 1475
                  /gene="IGHG1"
                  /replace=""
misc_difference 1578
                  /gene="IGHG1"
                  /replace=""
BASE COUNT      418 a      698 c      566 g      327 t
ORIGIN
1  agctttcttgg ggcaggccag gcctgacctt ggctttgggg cagggagggg gctaagggtga
61  ggcagggtggc gccagcaggt gcacacccaa tgcccatgag cccagacact ggacgctgaa
121 cctcgcggac agttaagaac ccaggggcct ctgcgcctgg gccagctct gtccacacc
181 gcggtcacat ggcaccacct ctcttgacgc ctccaccaag ggcccatcgg tcttccccct
241 ggcaccctcc tccaagagca cctctggggg cacagcggcc ctgggctgcc tgggtcaagga
301 ctacttcccc gaaccggtga cgggtgtcgtg gaactcaggc gccctgacca gcggcgtgca
361 caccttcccc gctgtcctac agtcctcagg actctactcc ctgagcagcg tggtgaccgt
421 gccctccagc agcttgggca cccagaccta catctgcaac gtgaatcaca agcccagcaa
481 caccaagggtg gacaagaaag ttggtgagag gccagcacag ggaggagggg tgtctgctgg
541 aagcaggctc agcgtcctg cctggacgca tcccggctat gcagccccag tccagggcag
601 caaggcaggc cccgtctgcc tcttcacccg gagcctctgc ccgccccact catgctcagg
661 gagagggtct tctggctttt tcccaggctc tgggcaggca caggctaggt gcccctaacc
721 caggccctgc acacaaaggg gcagggtgctg ggctcagacc tgccaagagc catatccggg
781 aggaccctgc ccctgaccta agcccacccc aaaggccaaa ctctccactc cctcagctcg
841 gacaccttct ctctcccag attccagtaa ctcccaatct tctctctgca gagcccaaat
901 cttgtgacaa aactcacaca tgcccaccgt gcccaggtaa gccagcccag gcctcgccct
961 ccagctcaag gcgggacagg tgccctagag tagcctgcat ccagggacag gcccagccg
1021 ggtgctgaca cgtccacctc catctcttcc tcagcacctg aactcctggg gggaccgtca
1081 gtcttctctt tccccccaaa acccaaggac accctcatga tctcccggac cctgagggtc
1141 acatgcgtgg tgggtggacgt gagccacgaa gacctgagg tcaagttcaa ctggtacgtg
1201 gacggcgtgg aggtgcataa tgccaagaca aagccgcggg aggagcagta caacagcacg
1261 taccgggtgg tcagcgtcct caccgtcctg caccaggact ggctgaatgg caaggagtac
1321 aagtgcgaag tctccaacaa agccctccca gcccccatcg agaaaaccat ctccaaagcc
1381 aaagggtggga cccgtggggg gcgagggcca catggacaga ggccggctcg gcccaccctc
1441 tgccctgaga gtgaccgctg taccaacctc tgtcctacag ggcagccccg agaaccacag
1501 gtgtacaccc tgcccccatc ccgggatgag ctgaccaaga accaggtcag cctgacctgc
1561 ctggtcaaa gcttctatcc cagcgacatc gccgtggagt gggagagcaa tgggcagccg
1621 gagaacaact acaagaccac gcctcccgtg ctggactcgg acggctcctt ctctctctac
1681 agcaagctca ccgtggacaa gagcagggtg cagcagggga acgtcttctc atgctccgtg
1741 atgcatgagg ctctgcacaa cacttacacg cagaagagcc tctccctgtc tccgggtaaa
1801 tgagtgcgac ggccggcaag ccccgctccc cgggctctcg cggtcgcacg aggatgcttg
1861 gcacgtaccc cctgtacata ctctccgggc gcccagcatg gaaataaagc acccagcgtc
1921 gccctgggccc cctgcgagac tgtgatggtt ctttccacgg gtcaggccga gtctgaggcc
1981 tgagtggcat gagggaggca gagcgggtc

```

FIGURE 7 J

IV. HUMAN IG GAMMA-2 CHAIN C REGION - HOMO SAPIENS (HUMAN).

AMINO ACID SEQUENCE

>sp|P01859|GC2_HUMAN IG GAMMA-2 CHAIN C REGION - Homo sapiens (Human).

```

      10      20      30      40      50      60
      |      |      |      |      |      |
ASTKGPSVFP LAPCSRSTSE STAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS

      70      80      90     100     110     120
      |      |      |      |      |      |
GLYSLSSVVT VPSSNFGTQT YTCNVDHKPS NTKVDKTVER KCCVECPPCP APPVAGPSVF

     130     140     150     160     170     180
      |      |      |      |      |      |
LFPPKPKDTL MISRTPEVTC VVVDVSHEDP EVQFNWYVDG VEVHNAKTKP REEQFNSTFR

     190     200     210     220     230     240
      |      |      |      |      |      |
VVSVLTVVHQ DWLNGKEYKC KVSNGKLPAP IEKTISKTKG QPREPQVYTL PPSREEMTKN

     250     260     270     280     290     300
      |      |      |      |      |      |
QVSLTCLVKG FYPSDIAVEW ESNQOPENNY KTTTPMLDSD GSFFLYSKLI VDKSRWQQGN

     310     320     326
      |      |      |
VFSCSVMHEA LHNHYTQKSL SLSPGK

```

CODING SEQUENCE

```

gcctccacca agggcccacc ggtcttcccc ctggcgccct gctccaggag cacctccgag      60
agcacagccg ccctgggctg cctgggtcaag gactacttcc ccgaaccggt gacgggtgtc      120
tggaactcag gcgctctgac cagcgggcgtg cacaccttcc cagctgtcct acagtcccca      180
ggactctact ccctcagcag cgtgggtgacc gtgccctcca gcaacttcgg caccagacc      240
tacacctgca acgtagatca caagcccagc aacaccaagg tggacaagac agttgagcgc      300
aaatgttgtg tcgagtgtcc accgtgtccc gcaccacctg tggcaggacc gtcagtcttc      360
ctcttcccc caaaacccaa ggacaccttc atgatctccc ggacctctga ggtcacgtgc      420
gtgggtgttg acgtgagcca cgaagacccc gaggtccagt tcaactggta cgtggacggc      480
gtggaggtgc ataatgccaa gacaaagcca cgggaggagc agttcaacag cacgttccgt      540
gtggtcagcg tcttcaccgt tgtgcaccag gactggctga acggcaagga gtacaagtgc      600
aaggctctca acaaaggcct cccagcccc atcgagaaaa ccatctccaa aaccaaggg      660
cagccccgag aaccacaggt gtacaccttg ccccatccc gggaggagat gaccaagaac      720
caggtcagcc tgacctgcct ggtcaaaggc ttctacccca gcgacatcgc cgtggagtgg      780
gagagcaatg ggcagccgga gaacaactac aagaccacac ctcccatgct ggactccgac      840
ggctccttct tctctacag caagctcacc gtggacaaga gcaggtggca gcaggggaac      900
gtcttctcat gctccgtgat gcatgaggct ctgcacaacc actacacgca gaagagcctc      960
tccctgtctc cgggtaaa

```

GenBank

J00230. Human Ig germline ...

LOCUS HUMIGCD1 2009 bp DNA PRI 11-APR-2001

DEFINITION Human Ig germline H-chain G-E-A region B: gamma-2 constant region, 3' end.

ACCESSION J00230 V00554

VERSION J00230.1 GI:184750

KEYWORDS .

SOURCE human.

ORGANISM Homo sapiens

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;

FIGURE 7 K

Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.

REFERENCE 1 (bases 1 to 2009)

AUTHORS Ellison, J. and Hood, L.

TITLE Linkage and sequence homology of two human immunoglobulin gamma heavy chain constant region genes

JOURNAL Proc. Natl. Acad. Sci. U.S.A. 79 (6), 1984-1988 (1982)

MEDLINE 82197621

PUBMED 6804948

REFERENCE 2 (bases 896 to 1256; 1749 to 1937)

AUTHORS Krawinkel, U. and Rabbitts, T.H.

TITLE Comparison of the hinge-coding segments in human immunoglobulin gamma heavy chain genes and the linkage of the gamma 2 and gamma 4 subclass genes JOURNAL EMBO J. 1 (4), 403-407 (1982)

MEDLINE 84235992 PUBMED 6329676

REFERENCE 3 (bases 475 to 1071; 1179 to 1330; 1461 to 1524)

AUTHORS Takahashi, N., Ueda, S., Obata, M., Nikaido, T., Nakai, S. and Honjo, T.

TITLE Structure of human immunoglobulin gamma genes: implications for evolution of a gene family JOURNAL Cell 29 (2), 671-679 (1982)

MEDLINE 83001943 PUBMED 6811139

COMMENT On Mar 2, 2000 this sequence version replaced gi:32759.

[2] also reports sequences for gamma-3, gamma-4, and a gamma pseudogene. Most of this sequence is 95% homologous with gamma-4. The hinge exons are only 70% homologous. The authors estimate that gamma-2 and gamma-4 diverged 6.6 million years ago. The authors in [1] speculate that intron-mediated domain transfer played an important role in the evolution of human gamma genes. They also report the hinge regions of gamma-1, gamma-3, gamma-4, and a pseudo-gamma gene. [1] estimates the divergence of the human gamma genes to be between 7.7 and 4.4 million years ago. This entry is part of a multigene region containing the gamma-2, gamma-4, epsilon-1, and alpha-2 genes. The relative locations of the four genes were determined by Flanagan and Rabbitts (Nature 300, 709-713 (1982)). They refer to this gene group as region B. The region A genes are gamma-3, gamma-1, pseudo-epsilon, alpha-1. Flanagan and Rabbitts also determined the general locations of the two regions. They place region A between the JH/mu/delta region and region B. Complete source information:

Human fetal liver DNA, library of T. Maniatis [3] and Lawn et al [2], [1]; clones p-gamma-2RPA3 [2], 5A [3], and Ig-gamma-2-15 [1].

FEATURES

Location/Qualifiers source 1..2009

/organism="Homo sapiens"

/db_xref="taxon:9606" /map="14q32.33"

/germline intron <1..215

/gene="IgH" gene <1..2009

/gene="IgH" /note="IGHG2"

exon 216..509 /gene="IgH"

/note="G00-119-338" C_region 216..508

/gene="IgH"

/note="immunoglobulin heavy chain constant region CH1"

CDS join(<216..509,902..937,1056..1382,1480..1802)

/gene="IgH" /codon_start=3

/product="immunoglobulin gamma-2 heavy chain"

/protein_id="AAB59393.1"

/db_xref="GI:184758"

/translation="STKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSNFGTQYTCNVDPKPSNTKVDKTKVERKCCVECPGPCAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVSVLTVVHQDNLNGKEYCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNYHTQKSLSLSPGK"

intron 510..901 /gene="IgH"

conflict 537 /gene="IgH"

/citation=[3] /replace=""

FIGURE 7 L

```

conflict      550..551                /gene="IgH"
               /citation=[3]          /replace=""
conflict      570                    /gene="IgH"
               /citation=[3]          /replace=""
conflict      777..778                /gene="IgH"
               /citation=[3]          /replace=""
conflict      791                    /gene="IgH"
               /citation=[3]          /replace=""
conflict      864                    /gene="IgH"
               /citation=[3]          /replace=""
C_region      901..936                /gene="IgH"
               /note="immunoglobulin heavy chain hinge"
exon          902..937                /gene="IgH"
               /note="G00-119-338"    intron      938..1055
               /gene="IgH"          C_region      1055..1381
               /gene="IgH"
               /note="immunoglobulin heavy chain constant region CH2"
exon          1056..1382              /gene="IgH"
               /note="G00-119-338"    intron      1383..1479
               /gene="IgH"          C_region      1479..1799
               /gene="IgH"
               /note="immunoglobulin heavy chain constant region CH3"
exon          1480..1802              /gene="IgH"
               /note="G00-119-338"    conflict      1493
               /gene="IgH"                /citation=[3]
               /replace=""          conflict      1802..1806
               /gene="IgH"                /citation=[2]
               /replace=""          conflict      1814..1815
               /gene="IgH"                /citation=[2]
               /replace=""          conflict      1825
               /gene="IgH"                /citation=[2]
               /replace=""          conflict      1844..1853
               /gene="IgH"                /citation=[2]
               /replace=""          conflict      1890
               /gene="IgH"                /citation=[2]
               /replace=""          polyA_signal  1903..1908
               /gene="IgH"          conflict      1909..1918
               /gene="IgH"                /citation=[2]
               /replace=""          conflict      1929..1932
               /gene="IgH"                /citation=[2]
               /replace=""          /replace=""BASE COUNT      410 a      700 c      568 g      331 t

ORIGIN
1 agctttctgg ggcgagccgg gcctgacttt ggctttgggg cagggagtgg gctaagggtga
61 ggcaggtggc gccagccagg tgcacaccca atgcccgtag gccagacac tggaccctgc
121 ctggaccctc gtggatagac aagaaccgag gggcctctgc gcctgggccc agctctgtcc
181 cacaccgagg tcacatggca ccacctctct tgcagcctcc accaagggcc catcggtctt
241 cccctggcgg ccctgctcca ggagcacctc cgagagcaca gccgcccctg gctgcctggg
301 caaggactac ttccccgaac cgggtgacgg gtctgtggaac tcaggcgctc tgaccagcgg
361 cgtgcacacc ttcccagctg tcctacagtc ctcaggactc tactccctca gcagcgtggg
421 gaccgtgccc tccagcaact tccggcaccca gacctacacc tgcaacgtag atcacaagcc
481 cagcaacacc aaggtggaca agacagttag tgagaggcca gctcaggagg ggagggtgtc
541 tgctggaagc caggctcagc cctcctgcct ggacgcaccc cggctgtgca gcccagccc
601 agggcagcaa ggcaggcccc atctgtctcc tcacccggag gcctctgccc gcccactca
661 tgctcaggga gagggctctc tggctttttc caccaggctc caggcaggca caggctgggt
721 gcccctaccc caggcccttc acacacaggg gcagggtgctt ggctcagacc tgccaaaagc
781 catatccggg aggaccctgc ccctgacctc agccgacccc aaaggccaaa ctgtccactc
841 cctcagctcg gacaccttct ctctcccag atccgagtaa ctcccaatct tctctctgca
901 gagcgcaaat gttgtgtcga gtgcccaccg tgcccaggta agccagccca ggctctgccc
961 tccagctcaa ggcgggacag gtgcccctaga gtagcctgca tccagggaca ggcccagct
1021 ggggtgctgac acgtccacct ccctctcttc ctcagcacca cctgtggcag gaccgtcagt
1081 cttctctctc cccccaaaac ccaaggacac cctcatgatc tcccggacce ctgaggtcac
1141 gtgcgtgggt gtggacgtga gccacgaaga ccccgaggtc cagttcaact ggtacgtgga
1201 cggcgtggag gtgcataatg ccaagacaaa gccacgggag gagcagttca acagcacgtt

```

FIGURE 7 M

```

1261 ccgtgtgggc agcgtectca ccgttgtgca ccaggactgg ctgaacggca aggagtacaa
1321 gtgcaaggtc tccaacaaag gcctcccagc ccccatcgag aaaaccatct ccaaaaccaa
1381 aggtgggacc cgcgggggat gagggccaca tggacagagg ccggctcggc ccaccctctg
1441 ccctgggagt gaccgctgtg ccaacctctg tccctacagg gcagccccga gaaccacagg
1501 tgtacaccct gcccccatcc cgggaggaga tgaccaagaa ccaggtcagc ctgacctgcc
1561 tgggtcaaagg cttctacccc agcgcacatcg ccgtggagtg ggagagcaat gggcagccgg
1621 agaacaacta caagaccaca cctcccatgc tggactccga cggctccttc ttcctctaca
1681 gcaagctcac cgtggacaag agcagggtggc agcaggggaa cgtcttctca tgctccgtga
1741 tgcattgaggc tctgcacaac cactacacgc agaagagcct ctccctgtct ccgggtaaat
1801 gagtggcacg gccggcaagc ccccgctccc caggctctcg gggtcgcgtg aggatgcttg
1861 gcacgtaccc cgtgtacata cttcccaggc acccagcatg gaaataaagc acccagcgt
1921 gccctgggccc cctgcgagac tgtgatgggt ctttcctgtg gtcaggccga gtctgaggcc
1981 tgagtggcat gagggaggca gagtgggtc

```

V. HUMAN IG GAMMA-3 CHAIN C REGION - HOMO SAPIENS (HUMAN).

AMINO ACID SEQUENCE

CAA27268 C gamma 3 [Homo sapiens]

```

      10      20      30      40      50      60
      |      |      |      |      |      |
ASTKGPSVFP LAPCSRSTSG GTAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS

      70      80      90     100     110     120
      |      |      |      |      |      |
GLYSLSSVVT VPSSSLGTQT YTCNVNHKPS NTKVDKRVEL KTPLGDTTHI CPRCPEPKSC

     130     140     150     160     170     180
      |      |      |      |      |      |
DTPPPCPRCP EPKSCDTPPP CPRCPEPKSC DTPPPCPRCP APELLGGPSV FLFPPKPKDT

     190     200     210     220     230     240
      |      |      |      |      |      |
LMISRTPEVT CVVVDVSHED PEVQFKWYVD GVEVHNAKTK PREEQYNSTF RVVSVLTVLH

     250     260     270     280     290     300
      |      |      |      |      |      |
QDWLNGKEYK CKVSNKALPA PIEKTISKTK GQPREPQVYT LPPSREEMTK NQVSLTCLVK

     310     320     330     340     350     360
      |      |      |      |      |      |
GFYPSDIAVE WESSGQPENN YNTTPMMLDS DGSFFLYSKL TVDKSRWQQG NIFSCSVMHE

     370     377
      |
ALHNRFTQKS LSLSPGK

```

CODING SEQUENCE

```

GCTTCCACCAAGGGCCCATCGGTCTTCCCCCTGGCGCCCTGCTCCAGGAGCACCTCTGGGGGCACAGCGGCCCTGGG
CTGCCTGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGGAAGTCAAGGCGCCCTGACCAGCGGCGTGCACA
CCTTCCCGGCTGTCTACAGTCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGC
ACCCAGACCTACACCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTTGAGCTCAAAACCCC
ACTTGGTGACACAACCTCACACATGCCACGGTGCCAGAGCCCAATCTTGTGACACACCTCCCCCGTGCCACGGT
GCCAGAGCCCAATCTTGTGACACACCTCCCCCATGCCACGGTGCCAGAGCCCAATCTTGTGACACACCTCCC
CCGTGCCCAAGGTGCCAGCACCTGAACTCCTGGGAGGACCGTCAGTCTTCTTCCCCCAAAACCCAAGGATAC
CCTTATGATTTCCCGGACCCCTGAGGTACAGTGGTGGTGGTGGACGTGAGCCACGAAGACCCGAGGTCCAGTTCA
AGTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTTCCTG

```

FIGURE 7 N

GTGGTCAGCGTCCTCACCCTCCTGCACCAGGACTGGCTGAACGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGC
 CCTCCCAGCCCCCATCGAGAAAACCATCTCCAAAACCAAAGGACAGCCCCGAGAACCACAGGTGTACACCCCTGCCCC
 CATCCCGGGAGGAGATGACCAAGAACCAGGTGACCTGACCTGGCTCAAAGGCTTCTACCCCAGCGACATCGCC
 GTGGAGTGGGAGAGCAGCGGGCAGCCGGAGAACAACACTACAACACCACGCCTCCCATGCTGGACTCCGACGGCTCCTT
 CTTCTCTACAGCAAGCTCACCCTGGACAAGAGCAGGTGGCAGCAGGGGAACATCTTCTCATGCTCCGTGATGCATG
 AGGCTCTGCACAACCGCTTCACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAATGA

GenBank

X03604 Human C gamma 3 gene for IgG G3m(b) heavy chain C-region from EZZ
 (individual II-4 of TOU) PubMed, Protein, Related Sequences, Taxonomy, OMIM,
 LinkOut

LOCUS HSIGGC3 2590 bp DNA PRI 24-NOV-1998
 DEFINITION Human C gamma 3 gene for IgG G3m(b) heavy chain C-region from EZZ
 (individual II-4 of TOU).
 ACCESSION X03604 M12958
 VERSION X03604.1 GI:33070
 KEYWORDS constant region; gamma-immunoglobulin; Ig heavy chain;
 immunoglobulin.
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.
 REFERENCE 1 (bases 1 to 2590)
 AUTHORS Huck,S., Fort,P., Crawford,D.H., Lefranc,M.P. and Lefranc,G.
 TITLE Sequence of a human immunoglobulin gamma 3 heavy chain constant
 region gene: comparison with the other human C gamma genes
 JOURNAL Nucleic Acids Res. 14 (4), 1779-1789 (1986)
 MEDLINE 86148507
 REFERENCE 2 (bases 4 to 204; 2202 to 2236)
 AUTHORS Takahashi,N., Ueda,S., Obata,M., Nikaido,T., Nakai,S. and
 Honjo,T.
 TITLE Structure of human immunoglobulin gamma genes: implications for
 evolution of a gene family
 JOURNAL Cell 29 (2), 671-679 (1982)
 MEDLINE 83001943
 FEATURES Location/Qualifiers
 source 1..2590
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 conflict 28
 /note="T is missing in [2]"
 /citation=[2]
 conflict 80..82
 /note="CGC is GCG in [2]"
 /citation=[2]
 conflict 120
 /note="C is T in [2]"
 /citation=[2]
 conflict 135
 /note="T is C in [2]"
 /citation=[2]
 CDS join(<215..509,901..951,1095..1139,1283..1327,
 1471..1515,1634..1963,2061..>2380)
 /codon_start=1
 /product="C gamma 3"
 /protein_id="CAA27268.1"

FIGURE 7

O

/db_xref="GI:577056"

/translation="ASTKGPSVFPLAPCSRSTSGGTAALGCLVKDYFPEPVTVSWNSG

ALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYTCNVNHKPSNTKVDKRVELKT

PLGDTTHTCPRCPEPKSCDTPPPCPRCPEPKSCDTPPPCPRCPEPKSCDTPPPCPRCP

APELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVQFKWYVDGVEVHNAK

TKPREEQYNSTFRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKTKGQPREP

QVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESSGQPENNYNTTTPMMLDSGGSF

FLYSKLTVDKSRWQQGNIFSCSVMEALHNRFQKSLSLSPGK"

intron 510..900

/note="intron I"

intron 952..1094

/note="intron II"

intron 1140..1282

/note="intron III"

intron 1328..1470

/note="intron IV"

intron 1516..1633

/note="intron V"

intron 1964..2060

/note="intron VI"

misc_feature 2484..2489

/note="pot. polyA signal"

BASE COUNT 541 a 925 c 703 g 421 t

ORIGIN

```

1 agcttttctgg ggcaggccag gcctgacttt ggctgggggc agggaggggg ctaagggtgac
61 gcagggtggcg ccagccaggc gcacacccaa tgcccgtag cccagacact ggaccctgcc
121 tggaccctcg tggatagaca agaaccgagg ggctctgctg ccctggggccc agctctgtcc
181 cacaccgcag tcacatggcg ccatctctct tgcagcttcc accaagggcc catcggtctt
241 ccccttgggc ccctgctcca ggagcacctc tgggggcaca gcggccctgg gctgcctggt
301 caaggactac ttccccgaac cggtagcggg gtcgtggaac tcaggcgccc tgaccagcgg
361 cgtgcacacc ttcccggctg tcctacagtc ctcaggactc tactccctca gcagcgtggg
421 gaccgtgccc tccagcagct tggggaccca gacctacacc tgcaacgtga atcacaagcc
481 cagcaacacc aagggtggaca agagagttgg tgagaggcca gcgcaggagg ggaggggtgtc
541 tgctggaagc caggctcagc cctcctgcct ggacgcattc cggctgtgca gtcccagccc
601 agggcagcaa ggcaggcccc gtctgactcc tcaccgggag cctctgcccc cccactcat
661 gctcaggagg aggggtcttct ggcttttttc accaggctcc gggcaggcac aggctggatg
721 cccctacccc aggcccttca cacacagggg cagggtgctg gctcagagct gccaaaagcc
781 atatccagga ggaccctgcc cctgacctaa gccaccccca aaggccaaac tctctactca
841 ctcagctcag acaccttctc tcttcccaga tctgagtaac tcccaatctt ctctctgcag
901 agctcaaaac cccacttggt gacacaactc acacatgccc acgggtgcca ggtaagccag
961 cccaggactc gccctccagc tcaaggcggg acaagagccc tagagtggcc tgagtccagg
1021 gacaggcccc agcagggtgc tgacgcattc acctccatcc cagatccccg taactcccaa
1081 tcttctctct gcagagccca aatcttgtga cacacctccc ccgtgcccac ggtgcccagg
1141 taagccagcc caggcctcac cctccagctc aaggcaggac aagagcccta gagtggcctg
1201 agtccaggga caggccccag cagggtgctg acgcgtccac ctccatccca gatccccgta
1261 actcccaatc ttctctctgc agagcccaaa tcttgtgaca cacctcccc atgcccacgg
1321 tgcccaggta agccagccca ggcctcgccc tccagctcaa ggcgggacaa gagccctaga
1381 gtggcctgag tccagggaca ggccccagca ggggtgctgac gcatccacct ccatcccaga
1441 tccccgtaac tcccaatctt ctctctgcag agcccaaatc ttgtgacaca cctcccccg
1501 gcccaagggtg cccaggtaag ccagcccagg cctcgccctc cagctcaagg caggacagg
1561 gccctagagt ggcctgcata caggacagg tcccagtcgg gtgctgacac atctgcctcc

```

FIGURE 7

P

```

1621 atctcttctt cagcacctga actcctggga ggaccgtcag tcttctctct ccccccaaaa
1681 cccaaggata cccttatgat ttcccggacc cctgagggtca cgtgcgtggt ggtggacgtg
1741 agccacgaag accccgaggt ccagttcaag tggtagctgg acggcgtgga ggtgcataat
1801 gccaagacaa agccgcggga ggagcagtag aacagcacgt tccgtgtggt cagcgtcctc
1861 accgtcctgc accaggactg gctgaacggc aaggagtaca agtgcaaggt ctccaacaaa
1921 gccctcccag ccccatcga gaaaaccatc tccaaaacca aagggtgggac ccgcggggta
1981 tgagggccac atggacagag gccagcttga cccaccctct gccctgggag tgaccgctgt
2041 gccaacctct gtccctacag gacagccccg agaaccacag gtgtacaccc tgcccccatc
2101 ccgggaggag atgaccaaga accagggtcag cctgacctgc ctggtcaaag gcttctaccc
2161 cagcgacatc gccgtggagt gggagagcag cgggcagccg gagaacaact acaacaccac
2221 gcctcccatg ctggactccg acggctcctt ctctctctac agcaagctca ccgtggacaa
2281 gagcaggtgg cagcagggga acatcttctc atgctccgtg atgcatgagg ctctgcacaa
2341 ccgcttcacg cagaagagcc tctccctgtc tccgggtaaa tgagtgcgac agccggcaag
2401 ccccgctccc ccgggtcttc ggggtcgcgc gaggatgctt ggcacgtacc ccgtgtacat
2461 acttcccggg caccagcat ggaaataaag caccagcgc tgccctgggc ccctgtgaga
2521 ctgtgatggt tctttccacg ggtcaggccg agtctgaggc ctgagtgaca tgagggaggg
2581 agagcgggtc

```

VI. HUMAN IG GAMMA-4 CHAIN C REGION - HOMO SAPIENS (HUMAN).

AMINO ACID SEQUENCE

>sp|P01861|GC4_HUMAN IG GAMMA-4 CHAIN C REGION - Homo sapiens (Human).

```

      10      20      30      40      50      60
      |      |      |      |      |      |
ASTKGPSVFP LAPCSRSTSE STAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS

      70      80      90     100     110     120
      |      |      |      |      |      |
GLYSLSSVVT VPSSSLGTKT YTCNVDHKPS NTKVDKRVES KYGPPCPSCP APEFLGGPSV

     130     140     150     160     170     180
      |      |      |      |      |      |
FLFPPKPKDT LMISRTPEVT CVVVDVSQED PEVQFNWYVD GVEVHNAKTK PREEQFNSTY

     190     200     210     220     230     240
      |      |      |      |      |      |
RVVSVLTVLH QDWLNGKEYK CKVSNKGLPS SIEKTISKAK GQPREPQVYT LPPSQEEMTK

     250     260     270     280     290     300
      |      |      |      |      |      |
NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPVLDS DGSFFLYSRL TVDKSRWQEG

     310     320     327
      |      |      |
NVFSCSVMHE ALHNHYTQKS LSLSLGK

```

CODING SEQUENCE

```

cttccaccaa gggcccatcc gtcttccccc tggcgccctg ctccaggagc acctccgaga      g      -1
gcacagccgc cctgggctgc ctggtcaaag actacttccc cgaaccgggtg acgggtgtcgt      60
ggaaactcagg cgccctgacc agcggcgtgc acaccttccc ggctgtccta cagtcctcag      120
gactctactc cctcagcagc gtgggtgaccg tgccctccag cagcttgggc acgaagacct      180
acacctgcaa cgtagatcac aagcccagca acaccaaggt ggacaagaga gttgagtcca      240
aatatggtcc cccatgccca tcatgccagc cacctgagtt cctgggggga ccatcagtct      300
tcctgttccc cccaaaaccc aaggacactc tcatgatctc ccggaccctt gaggtcacgt      360
                                                    420

```

FIGURE 7 Q

```

gcgtggtggt ggacgtgagc caggaagacc ccgaggtcca gttcaactgg tacgtggatg      480
gcgtggaggt gcataatgcc aagacaaagc cgcgggagga gcagttcaac agcacgtacc      540
gtgtggtcag cgtcctcacc gtcctgcacc aggactgggt gaacggcaag gagtacaagt      600
gcaaggtctc caacaaaggc ctcccgtcct ccacgcagaa aaccatctcc aaagccaaag      660
ggcagccccc agagccacag gtgtacaccc tgcccccatc ccaggaggag atgaccaaga      720
accaggtcag cctgacctgc ctgggtcaaag gcttctaccc cagcgacatc gccgtggagt      780
gggagagcaa tgggcagccg gagaacaact acaagaccac gcctcccggt ctggactccg      840
acggctcctt cttcctctac agcaggctaa ccgtggacaa gagcaggtgg caggagggga      900
atgtcttctc atgctccgtg atgcatgagg ctctgcacaa ccactacaca cagaagagcc      960
tctccctgtc tctgggtaaa tga                                           983

```

GenBank

K01316. Human Ig germline

LOCUS HUMIGCD2 2028 bp DNA PRI 11-APR-2001

DEFINITION Human Ig germline H-chain G-E-A region B: gamma-4 constant region, 3' end.

ACCESSION K01316

VERSION K01316.1 GI:184751

KEYWORDS .

SOURCE human.

ORGANISM Homo sapiens

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.

REFERENCE 1 (bases 1 to 2028)

AUTHORS Ellison,J., Buxbaum,J. and Hood,L.

TITLE Nucleotide sequence of a human immunoglobulin C gamma 4 gene

JOURNAL DNA 1 (1), 11-18 (1981)

MEDLINE 83157104

REFERENCE 2 (bases 475 to 1069; 1180 to 1331; 1432 to 1655)

AUTHORS Takahashi,N., Ueda,S., Obata,M., Nikaido,T., Nakai,S. and Honjo,T.

TITLE Structure of human immunoglobulin gamma genes: implications for evolution of a gene family

JOURNAL Cell 29 (2), 671-679 (1982)

MEDLINE 83001943

PUBMED 6811139

REFERENCE 3 (bases 894 to 1106)

AUTHORS Krawinkel,U. and Rabbitts,T.H.

TITLE Comparison of the hinge-coding segments in human immunoglobulin gamma heavy chain genes and the linkage of the gamma 2 and gamma 4 subclass genes

JOURNAL EMBO J. 1 (4), 403-407 (1982)

MEDLINE 84235992

PUBMED 6329676

REFERENCE 4 (bases 1 to 2028)

AUTHORS Ellison,J. and Hood,L.

TITLE Linkage and sequence homology of two human immunoglobulin gamma heavy chain constant region genes

JOURNAL Proc. Natl. Acad. Sci. U.S.A. 79 (6), 1984-1988 (1982)

MEDLINE 82197621

PUBMED 6804948

COMMENT [1] reports that the human C-gamma-4 gene is equally homologous to the mouse gamma-1, gamma-2a, and gamma-2b genes (about 75%). [3] also reports partial sequences for human gamma-2, gamma-3, and a gamma pseudogene. [2] presents the gamma-1, gamma-2, gamma-3, and pseudo-gamma hinge regions.

This entry is part of a multigene region (region B), which includes the gamma-2, gamma-4, epsilon-1, and alpha-2 genes. See segment 1 for more comments.

Complete source information:

Human fetal liver DNA, library of T. Maniatis [3] and Lawn et al [1],[2]; clones 24B [1], lambda-HG4.1 [3], and Ig-gamma-4-2 [2].

FEATURES Location/Qualifiers

source 1..2028

FIGURE 7

R

```

/organism="Homo sapiens"
/db_xref="taxon:9606"
/map="14q32.33"
/germline
gene      <1..2028
          /gene="IgH"
          /note="IGHG4"
intron    <1..215
          /gene="IgH"
          /note="gamma-4 intron J-C"
CDS       join(<216..509,900..935,1054..1383,1481..1803)
          /gene="IgH"
          /codon_start=3
          /product="immunoglobulin gamma-4 heavy chain"
          /protein_id="AAB59394.1"
          /db_xref="GI:184759"
          /translation="STKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGA
          LTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYTCNVDPKPSNTKVDKRVESKYG
          PPCPSCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDG
          VEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKA
          KGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPV
          LDSDGSEFFLYSRLTVDKSRWQEGNVFSCSVMEALHNHYTQKSLSLGLGK"
exon      216..509
          /gene="IgH"
          /note="G00-119-340"
intron    510..899
          /gene="IgH"
exon      900..935
          /gene="IgH"
intron    936..1053
          /gene="IgH"
exon      1054..1383
          /gene="IgH"
intron    1384..1480
          /gene="IgH"
exon      1481..1803
          /gene="IgH"
BASE COUNT      421 a      709 c      567 g      331 t
ORIGIN
    1 agcttttctgg ggcaggcccg ggcctgacttt ggctggggggc agggagggggg ctaaggtgac
   61 gcaggtggcg ccagccaggt gcacacccaa tgcccatgag ccagacact ggaccctgca
  121 tggaccatcg cggatagaca agaaccgagg ggccctctcg ccctggggcc agctctgtcc
  181 cacaccgchg tcacatggca ccacctctct tgcagcttcc accaagggcc catccgtctt
  241 ccccttggcg ccctgctcca ggagcacctc cgagagcaca gccgccctgg gctgcctggt
  301 caaggactac ttccccgaac cgggtgacggt gtcgtggaac tcaggcgccc tgaccagcgg
  361 cgtgcacacc ttcccggctg tcctacagtc ctcaggactc tactccctca gcagcgtggt
  421 gaccgtgccc tccagcagct tgggcacgaa gacctacacc tgcaacgtag atcacaagcc
  481 cagcaacacc aaggtggaca agagagttgg tgagaggcca gcacaggagg ggaggggtgtc
  541 tgctggaagc caggctcagc cctcctgcct ggacgcaccc cggctgtgca gccccagccc
  601 agggcagcaa ggcattgccc atctgtctcc tcaccgggag gcctctgacc accccactca
  661 tgctcagggg gaggggtcttc tggatttttc caccaggctc ccggcaccac aggctggatg
  721 cccctacccc aggccttgcg catacagggc aggtgctgcg ctcagacctg ccaagagcca
  781 tatccgggag gaccctgccc ctgacctaa cccaccccaa aggccaaact ctccactccc
  841 tcagctcaga cactttctct cctcccagat ctgagtaact cccaatcttc tctctgcaga
  901 gtccaaatat ggtcccccct gcccatcatg cccaggtaag ccaaccaggg cctcgccctc
  961 cagctcaagg cgggacaggt gccctagagt agcctgcac cagggacagg ccccagccgg
 1021 gtgctgacgc atccacctcc atctcttctt cagcacctga gttcctgggg ggaccatcag
 1081 tcttctgtgt cccccaaaa cccaaggaca ctctcatgat ctcccgacc cctgaggtca
 1141 cgtgcgtggt ggtggacgtg agccaggagg accccgaggt ccagttcaac tggtagtggt
 1201 atggcggtga ggtgcataat gcccaagaca agccgcggga ggagcagttc aacagcacgt
 1261 accgtgtggt cagcgtcctc accgtcctgc accaggactg gctgaacggc aaggagtaca
 1321 agtgcaaggt ctccaacaaa ggcctcccgt cctccatcga gaaaaccatc tccaaagcca
 1381 aaggtgggac ccacgggggt cgagggccac acggacagag gccagctcgg cccaccctct

```

FIGURE 7 S

```

1441 gccctgggag tgaccgctgt gccaacctct gtccctacag ggcagccccg agagccacag
1501 gtgtacaccc tgcccccatc ccaggaggag atgaccaaga accagggtcag cctgacctgc
1561 ctgggtcaaag gcttctaccc cagcgacatc gccgtggagt gggagagcaa tgggcagccg
1621 gagaacaact acaagaccac gcctcccgtg ctggactccg acggctcctt cttcctctac
1681 agcaggctaa ccgtggacaa gagcaggtgg caggagggga atgtcttctc atgctccgtg
1741 atgcatgagg ctctgcacaa ccactacaca cagaagagcc tctccctgtc tctgggtaaa
1801 tgagtgccag ggccggcaag cccccgtcc ccgggctctc ggggtcgcgc gaggatgctt
1861 ggcacgtacc ccgtctacat acttcccagg caccagcat ggaaataaag caccaccac
1921 tgccctgggc ccctgtgaga ctgtgatggt tctttccacg ggtcaggccg agtctgaggc
1981 ctgagtgaca tgaggagggc agagcgggtc ccactgtccc cacactgg

```

VII. HUMAN IG DELTA CHAIN C REGION - HOMO SAPIENS (HUMAN).**AMINO ACID SEQUENCE**

>sp|P01880|DTC_HUMAN IG DELTA CHAIN C REGION - Homo sapiens (Human).

```

      10      20      30      40      50      60
      |      |      |      |      |      |
APTKAPDVFP IISGCRHPKD NSPVVLACLI TGYHPTSVTV TWYMGTSQSQP QRTFPEIQRR

      70      80      90     100     110     120
      |      |      |      |      |      |
DSYYMTSSQL STPLQQWRQG EYKCVVQHTA SKSKKEIFRW PESPKAQASS VPTAQPOAEG

     130     140     150     160     170     180
      |      |      |      |      |      |
SLAKATTAPA TTRNTGRGGE EKKKEKEKEE QEERETKTPE CPSHTQPLGV YLLTPAVQDL

     190     200     210     220     230     240
      |      |      |      |      |      |
WLRDKATFTC FVVGSDLKDA HLTWEVAGKV PTGGVEEGLL ERHSNGSQSQ HSRLTLPRSL

     250     260     270     280     290     300
      |      |      |      |      |      |
WNAGTSVTCT LNHPSLPPQR LMALREPAAQ APVKLSLNL LASSDPPEAAS WLLCEVSGFS

     310     320     330     340     350     360
      |      |      |      |      |      |
PPNILLMWLE DQREVNTSGF APARPPPQPG STTFWAWSVL RVPAPPSPQP ATYTCVVSHE

     370     380
      |      |
DSRTLLNASR SLEVSIVTDH GPM

```

GenBank

K02876. Human germline IgD...[gi:184766] PubMed, Protein, Related Sequences, Taxonomy, OMIM, LinkOut

```

LOCUS      HUMIGCH02      300 bp      DNA      PRI      08-NOV-1994
DEFINITION Human germline IgD-chain gene, C-region, first hinge domain.
ACCESSION  K02876
VERSION    K02876.1  GI:184766
KEYWORDS   C-region; germline; hinge exon; immunoglobulin heavy chain;
           immunoglobulin-delta.
SOURCE     Homo sapiens (individual_isolate Chronic lymphocytic leukemia (CLL)
           patient) DNA.
ORGANISM   Homo sapiens
           Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;

```

FIGURE 7

T

Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.

REFERENCE 1 (bases 1 to 300)

AUTHORS White,M.B., Shen,A.L., Word,C.J., Tucker,P.W. and Blattner,F.R.

TITLE Human immunoglobulin D: genomic sequence of the delta heavy chain

JOURNAL Science 228 (4700), 733-737 (1985)

MEDLINE 85192522

COMMENT See segment 1.

FEATURES

Location/Qualifiers

source 1..300

/organism="Homo sapiens"

/isolate="Chronic lymphocytic leukemia (CLL) patient"

/db_xref="taxon:9606"

/map="14q32.33"

/cell_type="lymphocyte"

/germline

intron <1..151

/gene="IGHD"

/note="G00-120-084"

/number=1

exon 101..202

/partial

/gene="IGHD"

/note="hinge-1 domain; G00-120-084"

/number=2

intron 203..>300

/gene="IGHD"

/note="G00-120-084"

/number=2

gene join(K02875.1:1..495,1..300,K02877.1:1..300,
K02878.1:1..500,K02879.1:1..500,K02880.1:1..100,
K02881.1:1..200,K02882.1:1..52)

/gene="IGHD"

CDS join(K02875.1:101..403,101..202,K02877.1:101..172,
K02878.1:101..424,K02879.1:101..424,K02881.1:25..182,
K02882.1:44..52)

/partial

/gene="IGHD"

/note="membrane bound form"

/codon_start=3

/product="immunoglobulin delta-chain"

/protein_id="AAA52771.1"

/db_xref="GI:495872"

/db_xref="GDB:G00-120-084"

/translation="PTKAPDVFPIISGCRHPKDNSPVVLACLITGYHPTSVTVTWYMG
TQSQPQRTFPEIQRRDSYYMTSSQLSTPLQQWRQGEYKCVVQHTASKSKKEIFRWPE
PKAQASSVPTAQPOAEGSLAKATTAPATTRNTGRGGEEKKKEKEKEEQEERETKTPEC
PSHTQPLGVYLLTPAVQDLWLRDKATFTCFVVGSDLDKDAHLTWEVAGKVPTGCVVEEGL
LERHSNGSQSQHSRLTLPRSLWNAGTSVTCTLNHPSLPPQRLMALREPAAQAPVKLSL
NLLASSDPPEAASWLLCEVSGFSPPNILLMWLEDQREVNTSGFAPARPPPQPRSTTFW
AWSVLRVPAPPSPQPATYTCVVSHEDSRTLLNASRSLEVSYLAMTPLIPQSKDENSDD
YTTFDDVGSLWTTTLSTFVALFILTLTLYSGIVTFIKVK"

CDS join(K02875.1:101..403,101..202,K02877.1:101..172,
K02878.1:101..424,K02879.1:101..424,K02880.1:25..53)

/partial

/gene="IGHD"

/note="secreted form"

/codon_start=3

/product="immunoglobulin delta-chain"

/protein_id="AAA52770.1"

/db_xref="GI:495871"

/db_xref="GDB:G00-120-084"

/translation="PTKAPDVFPIISGCRHPKDNSPVVLACLITGYHPTSVTVTWYMG
TQSQPQRTFPEIQRRDSYYMTSSQLSTPLQQWRQGEYKCVVQHTASKSKKEIFRWPE

FIGURE 7

U

PKAQASSVPTAQFQAEGSLAKATTAPATTRNTGRGGEEKKKKEKEEQEERETKTPEC
 PSHTQPLGVYLLTPAVQDLWLRDKATFTCFVVGSDLKDAHLTWEVAGKVPTGGVEEGL
 LERHSNGSQSQHSRLTLPRSLWNAGTSVTCTLNHPSLPPQRLMALREPAAQAPVKLSL
 NLLASSDPPEAASWLLCEVSGFSPPNILLMWLEDQREVNTSGFAPARPPPQPRSTTFW
 AWSVLRVPAPPSPQPATYTCVVSHEDSRILLNASRSLEVSYVTDHGPMK"

BASE COUNT 59 a 133 c 52 g 56 t

ORIGIN About 300 bp after segment 1; 118 bp upstream of StuI site.

1 taggctgcct gtgccccca cctgcctgtc cacaaccag cctctggtac atccatgccc
 61 tctgccctaa gcctcacctg cacttttcct tggatttcag agtctccaaa ggcacaggcc
 121 tcctccgtgc ccactgcaca accccaagca gagggcagcc tcgccaaggc aaccacagcc
 181 ccagccacca cccgtaacac aggtgagaag ccccttcctt gcacactcca cccccacca
 241 cctgctcatt cctcagccgc ctctccagg cagcccttca taactccttg tctgagtctc

K02877. Human Ig germline ...[gi:184767] PubMed, Protein, Related Sequences, Taxonomy, OMIM, LinkOut

LOCUS HUMIGCH03 300 bp DNA PRI 08-NOV-1994
 DEFINITION Human Ig germline delta H-chain C-region gene, second hinge domain (CLL lymphocyte).
 ACCESSION K02877
 VERSION K02877.1 GI:184767
 KEYWORDS C-region; germline; hinge exon; immunoglobulin heavy chain; immunoglobulin-delta.
 SOURCE Homo sapiens (individual_isolate Chronic lymphocytic leukemia (CLL) patient) DNA.
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.
 REFERENCE 1 (bases 1 to 300)
 AUTHORS White,M.B., Shen,A.L., Word,C.J., Tucker,P.W. and Blattner,F.R.
 TITLE Human immunoglobulin D: genomic sequence of the delta heavy chain
 JOURNAL Science 228 (4700), 733-737 (1985)
 MEDLINE 85192522
 COMMENT See segment 1.
 FEATURES
 source Location/Qualifiers
 1..300
 /organism="Homo sapiens"
 /isolate="Chronic lymphocytic leukemia (CLL) patient"
 /db_xref="taxon:9606"
 /map="14q32.33"
 /cell_type="lymphocyte"
 /germline
 intron <1..100
 /gene="IGHD"
 /note="G00-120-084"
 /number=2
 exon 101..172
 /gene="IGHD"
 /note="hinge-2 domain; G00-120-084; putative"
 /number=3
 intron 173..>300
 /gene="IGHD"
 /note="G00-120-084"
 /number=3
 gene join(K02875.1:1..495,K02876.1:1..300,1..300,
 K02878.1:1..500,K02879.1:1..500,K02880.1:1..100,
 K02881.1:1..200,K02882.1:1..52)
 /gene="IGHD"
 CDS join(K02875.1:101..403,K02876.1:101..202,101..172,
 K02878.1:101..424,K02879.1:101..424,K02881.1:25..182,
 K02882.1:44..52)
 /partial
 /gene="IGHD"

FIGURE 7

V

```

/feature="membrane bound form"
/codon_start=3
/product="immunoglobulin delta-chain"
/protein_id="AAA52771.1"
/db_xref="GI:495872"
/db_xref="GDB:G00-120-084"
/translation="PTKAPDVFPPIISGCRHPKDNSPVVLACLITGYHPTSVTVTWYMG
TQSQPQRTFPEIQRRDSYYMTSSQLSTPLQQRQGEYKCVVQHTASKSKKEIFRWPE
PKAQASSVPTAQQAEGSLAKATTAPATTRNTGRGGEEKKEKEKEEQEERETKTPEC
PSHTQPLGVYLLTPAVQDLWLRDKATFTCFVVGSDLKDAHLTWEVAGKVPTGGVEEGL
LERHSNGSQSQHSRLTLPRSLWNAGTSVTCTLNHPSLPPQRLMALREPAAQAPVKLSL
NLLASSDPPEAASWLLCEVSGFSPPNILLMWLEDQREVNTSGFAPARPPPQPRSTTFW
AWSVLRVPAPPSPQATYTCVVSHEDSRTLLNASRSLEVSYLAMTPLIPQSKDENSDD
YTTFDDVGSLLWTTLSFVALFILTLTLYSGIVTFIKVK"
CDS
join(K02875.1:101..403,K02876.1:101..202,101..172,
K02878.1:101..424,K02879.1:101..424,K02880.1:25..53)
/partial
/gene="IGHD"
/feature="secreted form"
/codon_start=3
/product="immunoglobulin delta-chain"
/protein_id="AAA52770.1"
/db_xref="GI:495871"
/db_xref="GDB:G00-120-084"
/translation="PTKAPDVFPPIISGCRHPKDNSPVVLACLITGYHPTSVTVTWYMG
TQSQPQRTFPEIQRRDSYYMTSSQLSTPLQQRQGEYKCVVQHTASKSKKEIFRWPE
PKAQASSVPTAQQAEGSLAKATTAPATTRNTGRGGEEKKEKEKEEQEERETKTPEC
PSHTQPLGVYLLTPAVQDLWLRDKATFTCFVVGSDLKDAHLTWEVAGKVPTGGVEEGL
LERHSNGSQSQHSRLTLPRSLWNAGTSVTCTLNHPSLPPQRLMALREPAAQAPVKLSL
NLLASSDPPEAASWLLCEVSGFSPPNILLMWLEDQREVNTSGFAPARPPPQPRSTTFW
AWSVLRVPAPPSPQATYTCVVSHEDSRTLLNASRSLEVSYVTDHGPMK"
BASE COUNT      102 a      52 c      70 g      76 t
ORIGIN      About 1.85 kb after segment 2.
      1 gtcattagct ggatttagcc attccacaat gtacacatat ttcaaacatt gtgttgata
      61 tgataaacat gtataatttt tgtcaattaa aaatttttag gaagaggagg agaagagaag
     121 aagaaggaga aggagaaaga ggaacaagaa gagagagaga caaagacacc aggttttttc
     181 tgaccctgg gctatcaaaa cacctattgc ccaataacta gttggcgtt ggtgcctaa
     241 actattgaag cgattgctgt tatgtggatg ggccccggac acttagaaac tcgtgacccc

K02878. Human germline IgD...[gi:184768] PubMed, Protein, Related Sequences, Taxonomy,
OMIM, LinkOut

LOCUS      HUMIGCH04      500 bp      DNA      PRI      08-NOV-1994
DEFINITION Human germline IgD chain gene, C-region, C-delta-2 domain.
ACCESSION  K02878
VERSION    K02878.1  GI:184768
KEYWORDS   C-region; germline; immunoglobulin heavy chain;
            immunoglobulin-delta.
SOURCE     Homo sapiens (individual_isolate Chronic lymphocytic leukemia (CLL)
            patient) DNA.
ORGANISM   Homo sapiens
            Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
            Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 500)
AUTHORS    White,M.B., Shen,A.L., Word,C.J., Tucker,P.W. and Blattner,F.R.
TITLE      Human immunoglobulin D: genomic sequence of the delta heavy chain
JOURNAL    Science 228 (4700), 733-737 (1985)
MEDLINE    85192522
COMMENT    See segment 1.
FEATURES   Location/Qualifiers
            source      1..500
                        /organism="Homo sapiens"
                        /isolate="Chronic lymphocytic leukemia (CLL) patient"

```

FIGURE 7

W

```

/db_xref="taxon:9606"
/map="14q32.33"
/cell_type="lymphocyte"
/germline
intron    <1..100
          /gene="IGHD"
          /note="G00-120-084"
          /number=3
exon      101..424
          /gene="IGHD"
          /note="C-delta-2 domain; G00-120-084"
          /number=4
intron    425..>500
          /gene="IGHD"
          /note="G00-120-084"
          /number=4
intron    425..>500
          /gene="IGHD"
          /note="IgD-s intron D"
gene      join(K02875.1:1..495,K02876.1:1..300,K02877.1:1..300,
1..500,K02879.1:1..500,K02880.1:1..100,K02881.1:1..200,
K02882.1:1..52)
          /gene="IGHD"
CDS       join(K02875.1:101..403,K02876.1:101..202,
K02877.1:101..172,101..424,K02879.1:101..424,
K02881.1:25..182,K02882.1:44..52)
          /partial
          /gene="IGHD"
          /note="membrane bound form"
          /codon_start=3
          /product="immunoglobulin delta-chain"
          /protein_id="AAA52771.1"
          /db_xref="GI:495872"
          /db_xref="GDB:G00-120-084"
          /translation="PTKAPDVFPFIISGCRHPKDNSPVVLACLITGYHPTSVITVTWYMG
TQSQPQRTFPEIQRRDSYYMTSSQLSTPLQQRQGEYKCVVQHTASKSKKEIFRWPES
PKAQASSVPTAQPPAEGSLAKATTAPATTRNTGRGGEEKKKEKEKEEQEERETKTPEC
PSHTQPLGVYLLTPAVQDLWLRDKATFTCFVVGSDLKDAHLTWEVAGKVPTGGVEEGL
LERHSNGSQSQHSRLTLPRSLWNAGTSVTCTLNHPSLPPQRLMALREPAAQAPVKLSL
NLLASSDPPEAASWLLCEVSGFSPPNILLMWLEDQREVNTSGFAPARPPPQPRSTTFW
AWSVLRVPAPPSPQATYTCVVSHEDSRTLLNASRSLEVSYLAMTFLIPQSKDENSDD
YTTFDVGSLLWTTLSLTFVALFILTLTLYSGIVTFIKVK"
CDS       join(K02875.1:101..403,K02876.1:101..202,
K02877.1:101..172,101..424,K02879.1:101..424,
K02880.1:25..53)
          /partial
          /gene="IGHD"
          /note="secreted form"
          /codon_start=3
          /product="immunoglobulin delta-chain"
          /protein_id="AAA52770.1"
          /db_xref="GI:495871"
          /db_xref="GDB:G00-120-084"
          /translation="PTKAPDVFPFIISGCRHPKDNSPVVLACLITGYHPTSVITVTWYMG
TQSQPQRTFPEIQRRDSYYMTSSQLSTPLQQRQGEYKCVVQHTASKSKKEIFRWPES
PKAQASSVPTAQPPAEGSLAKATTAPATTRNTGRGGEEKKKEKEKEEQEERETKTPEC
PSHTQPLGVYLLTPAVQDLWLRDKATFTCFVVGSDLKDAHLTWEVAGKVPTGGVEEGL
LERHSNGSQSQHSRLTLPRSLWNAGTSVTCTLNHPSLPPQRLMALREPAAQAPVKLSL
NLLASSDPPEAASWLLCEVSGFSPPNILLMWLEDQREVNTSGFAPARPPPQPRSTTFW
AWSVLRVPAPPSPQATYTCVVSHEDSRTLLNASRSLEVSYSYVTDHGPMK"
BASE COUNT      93 a      171 c      157 g      79 t
ORIGIN          About 450 bp after segment 3; 131 bp upstream of AccI site.
1 gaagctgggg agaggagagc acagtgggta agtcagtccc tgcagcccaa ctgctcccca

```

FIGURE 7 X

```

61 aggtccggcc acagctgctc tcgtttgctc tcccctgcag agtggtccgag ccacacccag
121 cctcttggcg tctacctgct aacccttgca gtgcaggacc tgtgggtccg ggacaaagcc
181 accttcacct gcttcgtggt gggcagtgac ctgaaggatg ctcacctgac ctgggaggtg
241 gctgggaagg tccccacagg gggcgtggag gaagggctgc tggagcggca cagcaacggc
301 tcccagagcc agcacagccg tctgacctg cccaggtcct tgtggaacgc ggggacctcc
361 gtcacctgca cactgaacca tcccagcctc ccaccccaga ggttgatggc gctgagagaa
421 cccggtgagc ctggctccca ggtggggaga cgaggggtgc cacagcctgc tgacctctac
481 gccgccccca gggccatgac

```

K02879. Human Ig germline ...[gi:184769] PubMed, Protein, Related Sequences, Taxonomy, OMIM, LinkOut

```

LOCUS      HUMIGCH05      500 bp      DNA      PRI      08-NOV-1994
DEFINITION Human Ig germline delta H-chain C-region gene, C-delta-3 domain
            (CLL lymphocyte).
ACCESSION  K02879
VERSION    K02879.1  GI:184769
KEYWORDS    C-region; germline; immunoglobulin heavy chain;
            immunoglobulin-delta.
SOURCE      Homo sapiens (individual_isolate Chronic lymphocytic leukemia (CLL)
            patient) DNA.
ORGANISM    Homo sapiens
            Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
            Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE   1  (bases 1 to 500)
AUTHORS     White,M.B., Shen,A.L., Word,C.J., Tucker,P.W. and Blattner,F.R.
TITLE       Human immunoglobulin D: genomic sequence of the delta heavy chain
JOURNAL     Science 228 (4700), 733-737 (1985)
MEDLINE     85192522
COMMENT     See segment 1.
FEATURES             Location/Qualifiers
     source           1..500
                     /organism="Homo sapiens"
                     /isolate="Chronic lymphocytic leukemia (CLL) patient"
                     /db_xref="taxon:9606"
                     /map="14q32.33"
                     /cell_type="lymphocyte"
                     /germline
     intron           <1..100
                     /gene="IGHD"
                     /note="G00-120-084"
                     /number=4
     exon             101..424
                     /gene="IGHD"
                     /note="C-delta-3 domain; G00-120-084; putative"
                     /number=5
     intron           425..>500
                     /gene="IGHD"
                     /note="G00-120-084"
                     /number=5
     gene             join(K02875.1:1..495,K02876.1:1..300,K02877.1:1..300,
                     K02878.1:1..500,1..500,K02880.1:1..100,K02881.1:1..200,
                     K02882.1:1..52)
                     /gene="IGHD"
     CDS             join(K02875.1:101..403,K02876.1:101..202,
                     K02877.1:101..172,K02878.1:101..424,101..424,
                     K02881.1:25..182,K02882.1:44..52)
                     /partial
                     /gene="IGHD"
                     /note="membrane bound form"
                     /codon_start=3
                     /product="immunoglobulin delta-chain"
                     /protein_id="AAA52771.1"

```

FIGURE 7

Y

```

/db_xref="GI:495872"
/db_xref="GDB:G00-120-084"
/translation="PTKAPDVFPIISGCRHPKDNSPVVLACLITGYHPTSVTVTWYMG
TQSQPQRTFPEIQRRDSYMTSSQLSTPLQQWRQGEYKCVVQHTASKSKKEIFRWPE
PKAQASSVPTAQPAEGSLAKATTAPATTRNTGRGGEEKKKKEKEEQEERETKTPEC
PSHTQPLGVYLLTPAVQDLWLRDKATFTCFVVGSDLKDAHLTWEVAGKVPTGGVBEGL
LERHSNGSQSQHSRLTLPRSLWNAGTSVTCTLNHPSLPPQRLMALREPAAQAPVKLSL
NLLASSDPPEAASWLLCEVSGFSPPNILLMWLEDQREVNTSGFAPARPPPQPRSTTFW
AWSVLRVPAPPSPQPATYTCVVSHEDSRTLLNASRSLEVSYLAMTPLIPQSKDENSDD
YTTFDDVGSLWTTLSTFVALFILTLLYSGIVTFIKVK"
CDS
join(K02875.1:101..403,K02876.1:101..202,
K02877.1:101..172,K02878.1:101..424,101..424,
K02880.1:25..53)
/partial
/gene="IGHD"
/note="secreted form"
/codon_start=3
/product="immunoglobulin delta-chain"
/protein_id="AAA52770.1"
/db_xref="GI:495871"
/db_xref="GDB:G00-120-084"
/translation="PTKAPDVFPIISGCRHPKDNSPVVLACLITGYHPTSVTVTWYMG
TQSQPQRTFPEIQRRDSYMTSSQLSTPLQQWRQGEYKCVVQHTASKSKKEIFRWPE
PKAQASSVPTAQPAEGSLAKATTAPATTRNTGRGGEEKKKKEKEEQEERETKTPEC
PSHTQPLGVYLLTPAVQDLWLRDKATFTCFVVGSDLKDAHLTWEVAGKVPTGGVBEGL
LERHSNGSQSQHSRLTLPRSLWNAGTSVTCTLNHPSLPPQRLMALREPAAQAPVKLSL
NLLASSDPPEAASWLLCEVSGFSPPNILLMWLEDQREVNTSGFAPARPPPQPRSTTFW
AWSVLRVPAPPSPQPATYTCVVSHEDSRTLLNASRSLEVSYVTDHGPMK"
BASE COUNT      85 a      188 c      145 g      82 t
ORIGIN      About 150 bp after segment 4; 118 bp upstream of HindIII site.
      1 ccacaggaaa ggagaaggga ggcaccacac cctggccggc cccacttctc tcccagtgcc
     61 cccgtggcca gagcctgaca gccccccac ctccccgcag ctgcgaggc acccgtcaag
    121 ctttctctga acctgctggc ctgctctgac cctcccgagg cggcctcgtg gctcctgtgt
    181 gaggtgtctg gcttctcgcc ccccaacatc ctccctgatg ggctggagga ccagcgtgag
    241 gtgaacactt ctgggtttgc ccccgacgc cccctccac agcccaggag caccacgttc
    301 tgggcctgga gtgtgctgcg tgtcccagcc ccgcccagcc ctgagccagc cacctacag
    361 tgtgtggtca gccacgagga ctcccgact ctgctcaacg ccagccggag cctagaagtc
    421 agctgtgagt cccccccagg ccagggttg gacggggact ctgagggggg ccataaggag
    481 ctggaatcca tactaggcag

K01311. Human IgD germline...[gi:184716] PubMed, Protein, Taxonomy, OMIM

LOCUS      HUMIGCB9      106 bp      DNA      PRI      12-APR-2001
DEFINITION Human IgD germline chain J-delta region: C-delta CH1.
ACCESSION  K01311
VERSION    K01311.1  GI:184716
KEYWORDS   C-region; germline; immunoglobulin heavy chain;
           immunoglobulin-delta.
SOURCE     human.
  ORGANISM Homo sapiens
           Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
           Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE  1 (bases 1 to 106)
  AUTHORS  Rabbitts,T.H., Forster,A. and Milstein,C.P.
  TITLE    Human immunoglobulin heavy chain genes: evolutionary comparisons of
           C mu, C delta and C gamma genes and associated switch sequences
  JOURNAL  Nucleic Acids Res. 9 (18), 4509-4524 (1981)
  MEDLINE  82059479
  PUBMED   6795593
COMMENT    The deduced amino acid sequence is compared in [1] to the
           J/C-delta-1 junction of human ER1 protein. The delta gene occurs
           only 5 kb from the mu region. The authors [1] could not detect any
           switch-related sequences adjacent to the delta gene and state that

```

FIGURE 7 Z

this implies that the mu/delta switch cannot occur by the class switch recombination method. They speculate that the entire VH-(C-mu)-(C-delta) region is transcribed into one nuclear precursor molecule which is spliced later. This is part of a multigene region containing the J-region, switch region, C-mu-secreted, C-mu-membrane, and C-delta genes.

```

FEATURES             Location/Qualifiers
     source            1..106
                        /organism="Homo sapiens"
                        /db_xref="taxon:9606"
                        /map="14q32.33"
                        /cell_type="lymphocyte"
                        /tissue_type="placenta"
                        /tissue_type="liver"
                        /dev_stage="foetus"
                        /germline
                        /tissue_lib="of Lawn et al."
     gene              1..106
                        /gene="IGHD"
     intron            <1..26
                        /gene="IGHD"
                        /note="intron delta J-C; G00-120-084"
     CDS               <27..>106
                        /gene="IGHD"
                        /note="C-region CH1 domain"
                        /codon_start=3
                        /product="immunoglobulin delta-chain"
                        /protein_id="AAB59423.1"
                        /db_xref="GI:184735"
                        /translation="PTKAPDVFPFIISGCRHPKDNSPVVLA"
BASE COUNT           24 a      38 c      24 g      20 t
ORIGIN
      1  tgccacccca ggactctgtc ttccagcacc caccaaggct ccggatgtgt tccccatcat
     61  atcagggtgc agacacccaa aggataacag ccctgtgggc ctggca

```

K02880. Human germline IgD...[gi:184770] PubMed, Protein, Related Sequences, Taxonomy, OMIM, LinkOut

```

LOCUS       HUMIGCH06      100 bp      DNA              PRI          08-NOV-1994
DEFINITION  Human germline IgD chain gene, C-region, secreted terminus.
ACCESSION   K02880
VERSION     K02880.1  GI:184770
KEYWORDS    C-region; germline; immunoglobulin heavy chain;
            immunoglobulin-delta.
SOURCE      Homo sapiens (individual_isolate Chronic lymphocytic leukemia (CLL)
            patient) DNA.
  ORGANISM  Homo sapiens
            Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
            Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE   1  (bases 1 to 100)
  AUTHORS   White,M.B., Shen,A.L., Word,C.J., Tucker,P.W. and Blattner,F.R.
  TITLE     Human immunoglobulin D: genomic sequence of the delta heavy chain
  JOURNAL   Science 228 (4700), 733-737 (1985)
  MEDLINE   85192522
COMMENT     See segment 1.
FEATURES             Location/Qualifiers
     source            1..100
                        /organism="Homo sapiens"
                        /isolate="Chronic lymphocytic leukemia (CLL) patient"
                        /db_xref="taxon:9606"
                        /map="14q32.33"
                        /cell_type="lymphocyte"
                        /germline

```

FIGURE 7 AA

```

intron      <1..>100
            /gene="IGHD"
            /note="G00-120-084"
            /number=5
intron      <1..24
            /gene="IGHD"
            /note="G00-120-084"
            /number=5
exon        25..>53
            /gene="IGHD"
            /note="secreted terminus domain; G00-120-084"
            /number=6
gene        join(K02875.1:1..495,K02876.1:1..300,K02877.1:1..300,
K02878.1:1..500,K02879.1:1..500,1..100,K02881.1:1..200,
K02882.1:1..52)
            /gene="IGHD"
CDS         join(K02875.1:101..403,K02876.1:101..202,
K02877.1:101..172,K02878.1:101..424,K02879.1:101..424,
25..53)
            /partial
            /gene="IGHD"
            /note="secreted form"
            /codon_start=3
            /product="immunoglobulin delta-chain"
            /protein_id="AAA52770.1"
            /db_xref="GI:495871"
            /db_xref="GDB:G00-120-084"
            /translation="PTKAPDVFPPIISGCRHPKDNSPVVLACLITGYHPTSVTVTWYMG
TQSQPQRTFFPEIQRRDSYYMTSSQLSTPLQQRQGEYKCVVQHTASKSKKEIFRWPES
PKAQASSVPTAQPPAEGSLAKATTAPATTRNTGRGGEEKKEKEKEEQEERETKTPEC
PSHTQPLGVYLLTPAVQDLWLRDKATFTCFVVGSDLKDAHLTWEVAGKVPTGGVEEGL
LERHSNGSQSQHSRLTLPRSLWNAGTSVTCTLNHPSLPPQRLMALREPAAQAPVKLSL
NLLASSDPPEAASWLLCEVSGFSPPNILLMWLEDQREVNTSGFAPARPPPQPRSTTFW
AWSVLRVPAPPSPQPATYTCVVSHEDSRTLLNASRSLEVSIVTDHGPMK"
BASE COUNT      24 a      33 c      22 g      21 t
ORIGIN          About 1.8 kb after segment 5.
                1 gacacgccga ttttttggtta ttagatgtaa cagaccatgg ccccatgaaa tgatccccga
                61 ccagatccgt ccgcacccgc cactcagcag ctctggccga

```

K02881. Human germline IgD...[gi:184771] PubMed, Protein, Related Sequences, Taxonomy, OMIM, LinkOut

```

LOCUS          HUMIGCH07      200 bp      DNA          PRI          08-NOV-1994
DEFINITION     Human germline IgD-chain gene, C-region, first domain of membrane
terminus.
ACCESSION      K02881
VERSION        K02881.1  GI:184771
KEYWORDS       C-region; germline; immunoglobulin heavy chain;
immunoglobulin-delta.
SOURCE         Homo sapiens (individual_isolate Chronic lymphocytic leukemia (CLL)
patient) DNA.
ORGANISM       Homo sapiens
                Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
                Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE      1 (bases 1 to 200)
AUTHORS        White,M.B., Shen,A.L., Word,C.J., Tucker,P.W. and Blattner,F.R.
TITLE          Human immunoglobulin D: genomic sequence of the delta heavy chain
JOURNAL        Science 228 (4700), 733-737 (1985)
MEDLINE        85192522
COMMENT        See segment 1.
FEATURES       Location/Qualifiers
                source          1..200
                /organism="Homo sapiens"

```

FIGURE 7

BB

```

/isolate="Chronic lymphocytic leukemia (CLL) patient"
/db_xref="taxon:9606"
/map="14q32.33"
/cell_type="lymphocyte"
/germline
intron      <1..24
            /gene="IGHD"
            /note="G00-120-084"
            /number=5
exon        25..182
            /gene="IGHD"
            /note="first domain of membrane terminus; G00-120-084;
            putative"
            /number=6
intron      183..>200
            /gene="IGHD"
            /note="G00-120-084"
            /number=6
gene        join(K02875.1:1..495,K02876.1:1..300,K02877.1:1..300,
            K02878.1:1..500,K02879.1:1..500,K02880.1:1..100,1..200,
            K02882.1:1..52)
            /gene="IGHD"
CDS         join(K02875.1:101..403,K02876.1:101..202,
            K02877.1:101..172,K02878.1:101..424,K02879.1:101..424,
            25..182,K02882.1:44..52)
            /partial
            /gene="IGHD"
            /note="membrane bound form"
            /codon_start=3
            /product="immunoglobulin delta-chain"
            /protein_id="AAA52771.1"
            /db_xref="GI:495872"
            /db_xref="GDB:G00-120-084"
            /translation="PTKAPDVFPFIISGCRHPKDNSPVVLACLITGYHPTSVTVTWYMG
            TQSQPQRTFPEIQRRDSYYMTSSQLSTPLQQRQGEYKCVVQHTASKSKKEIFRWPES
            PKAQASSVPTAQPOAEGSLAKATTAPATTRNTGRGGEEKKEKEKEEQEERETKTPEC
            PSHTQPLGVYLLTPAVQDLWLRDKATFTCFVVGSDLKDAHLTWEVAGKVPTGGVEEGL
            LERHSNGSQSQHSRLTLPRSLWNAGTSVTCTLNHPSLPPQRLMALREPAAQAPVKLSL
            NLLASSDPPEAASWLLCEVSGFSPPNILLMWLEDQREVNTSGFAPARPPPQPRSTTFW
            AWSVLRVPAPPSPQPATYTCVVSHEDSRTLLNASRSLEVSYLAMTPLIPQSKDENSDD
            YTTFDDVGSLWTTLSTFVALFILTLISGIVTFIKVK"
BASE COUNT   37 a    72 c    49 g    42 t
ORIGIN       About 800 bp after segment 6.
            1 cgctcggccc ccgttcctcc ccagacctgg ccatgacccc cctgatccct cagagcaagg
            61 atgagaacag cgatgactac acgacctttg atgatgtggg cagcctgtgg accaccctgt
            121 ccacgtttgt ggcctctctc atcctcaccc tctctacag cggcattgtc actttcatca
            181 aggtcagggg agcggccagg

```

K02882. Human germline IgD...[gi:184772] PubMed, Protein, Related Sequences, Taxonomy, OMIM, LinkOut

LOCUS HUMIGCH08 100 bp DNA PRI 08-NOV-1994
 DEFINITION Human germline IgD-chain gene, C-region, second domain of membrane terminus.
 ACCESSION K02882
 VERSION K02882.1 GI:184772
 KEYWORDS C-region; germline; immunoglobulin heavy chain; immunoglobulin-delta.
 SOURCE Homo sapiens (individual_isolate Chronic lymphocytic leukemia (CLL) patient) DNA.
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.

FIGURE 7

CC

REFERENCE 1 (bases 1 to 100)
 AUTHORS White,M.B., Shen,A.L., Word,C.J., Tucker,P.W. and Blattner,F.R.
 TITLE Human immunoglobulin D: genomic sequence of the delta heavy chain
 JOURNAL Science 228 (4700), 733-737 (1985)
 MEDLINE 85192522
 COMMENT See segment 1.
 FEATURES
 Location/Qualifiers
 source 1..100
 /organism="Homo sapiens"
 /isolate="Chronic lymphocytic leukemia (CLL) patient"
 /db_xref="taxon:9606"
 /map="14q32.33"
 /cell_type="lymphocyte"
 /germline
 intron <1..43
 /gene="IGHD"
 /note="IgD-Mb"
 /number=6
 exon 44..>52
 /gene="IGHD"
 /note="membrane-bound form (second domain of membrane terminus); G00-120-084; putative"
 /number=7
 gene join(K02875.1:1..495,K02876.1:1..300,K02877.1:1..300,K02878.1:1..500,K02879.1:1..500,K02880.1:1..100,K02881.1:1..200,1..52)
 /gene="IGHD"
 CDS join(K02875.1:101..403,K02876.1:101..202,K02877.1:101..172,K02878.1:101..424,K02879.1:101..424,K02881.1:25..182,44..52)
 /partial
 /gene="IGHD"
 /note="membrane bound form"
 /codon_start=3
 /product="immunoglobulin delta-chain"
 /protein_id="AAA52771.1"
 /db_xref="GI:495872"
 /db_xref="GDB:G00-120-084"
 /translation="PTKAPDVFPIISGCRHPKDNSPVVLACLITGYHPTSVTVTWYMG
 TQSQPQRTFPEIQRRDSYMTSSQLSTPLQQWRQGEYKCVVQHTASKSKKEIFRWPE
 PKAQASSVPTAQFQAEGSLAKATTAPATTRNTGRGGEKKKEKEKEEQEERETKTPEC
 PSHTQPLGVYLLTPAVQDLWLRDKATFTCFVVGSDLKDAHLTWEVAGKVPTGGVEEGL
 LERHSNGSQSQHSRLTLPRSLWNAGTSVTCTLNHPSLPPQRLMALREPAAQAPVKLSL
 NLLASSDPPEAASWLLCEVSGFSPPNILLMWLEDQREVNTSGFAPARPPPQPRSTTFW
 AWSVLRVPAPPSPQPATYTCVVSHEDSRTLLNASRSLEVSYLAMTFLIPQSKDENSDD
 YTTFDVGSFLWTTLSTFVALFILTLLYSGIVTFIKVK"
 BASE COUNT 22 a 30 c 30 g 18 t
 ORIGIN About 1.3 kb after segment 7.
 1 tcaggcttct agccctgtcttgacccaggggctgtcttt caggtgaagt agcccagaa
 61 gagcaggacg ccctgtacct gcagagaagg gaagcagcct

K02875. Human germline IgD...[gi:184765] PubMed, Related Sequences, Taxonomy, OMIM, LinkOut

LOCUS HUMIGCH01 495 bp DNA PRI 08-NOV-1994
 DEFINITION Human germline IgD chain gene, C-region, C-delta-1 domain.
 ACCESSION K02875
 VERSION K02875.1 GI:184765
 KEYWORDS C-region; germline; immunoglobulin heavy chain; immunoglobulin-delta.
 SOURCE Homo sapiens (individual_isolate Chronic lymphocytic leukemia (CLL) patient) DNA.
 ORGANISM Homo sapiens

FIGURE 7 DD

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.

REFERENCE 1 (bases 1 to 495)

AUTHORS White, M.B., Shen, A.L., Word, C.J., Tucker, P.W. and Blattner, F.R.

TITLE Human immunoglobulin D: genomic sequence of the delta heavy chain

JOURNAL Science 228 (4700), 733-737 (1985)

MEDLINE 85192522

COMMENT Sequence in computer readable form and draft entry for [1] were kindly provided by M.B.White, 06-AUG-1985.
The C-delta and delta-s exon boundaries were located by comparing the translated sequences with known AA sequences [1].

FEATURES

Location/Qualifiers

source 1..495
/organism="Homo sapiens"
/isolate="Chronic lymphocytic leukemia (CLL) patient"
/db_xref="taxon:9606"
/map="14q32.33"
/cell_type="lymphocyte"
/germline

gene join(1..495, K02876.1:1..300, K02877.1:1..300, K02878.1:1..500, K02879.1:1..500, K02880.1:1..100, K02881.1:1..200, K02882.1:1..52)
/gene="IGHD"

intron <1..100
/gene="IGHD"
/note="J-C intron; G00-120-084"

CDS join(101..403, K02876.1:101..202, K02877.1:101..172, K02878.1:101..424, K02879.1:101..424, K02880.1:25..53)
/partial
/gene="IGHD"
/note="secreted form"
/codon_start=3
/product="immunoglobulin delta-chain"
/protein_id="AAA52770.1"
/db_xref="GI:495871"
/db_xref="GDB:G00-120-084"
/translation="PTKAPDVFPPIISGCRHPKDNSPVVLACLTGYHPTSVTVTWYMG
TQSQPQRTFPEIQRRDSYMTSSQLSTPLQWRQGEYKCVVQHTASKSKKEIFRWPE
PKAQASSVPTAQPAEGSLAKATTAPATTRNTGRGGEEKKKEKEKEEQEERETKTPEC
PSHTQPLGVYLLTPAVQDLWLRDKATFTCFVVGSDLKDAHLTWEVAGKVPTGGVEEGL
LERHSNGSQSQHSRLTLPRSLWNAGTSVTCTLNHPSLPPQRLMALREPAAQAPVKLSL
NLLASSDPPEAASWLLCEVSGFSPNILLMWLEDQREVNTSGFAPARPPPQPRSTTFW
AWSVLRVPAPPSPQPATYTCVVSHEDSRTLLNASRSLEVS YVTDHGPMK"

exon 101..403
/gene="IGHD"
/note="C-delta-1 domain; G00-120-084; putative"
/number=1

CDS join(101..403, K02876.1:101..202, K02877.1:101..172, K02878.1:101..424, K02879.1:101..424, K02881.1:25..182, K02882.1:44..52)
/partial
/gene="IGHD"
/note="membrane bound form"
/codon_start=3
/product="immunoglobulin delta-chain"
/protein_id="AAA52771.1"
/db_xref="GI:495872"
/db_xref="GDB:G00-120-084"
/translation="PTKAPDVFPPIISGCRHPKDNSPVVLACLTGYHPTSVTVTWYMG
TQSQPQRTFPEIQRRDSYMTSSQLSTPLQWRQGEYKCVVQHTASKSKKEIFRWPE
PKAQASSVPTAQPAEGSLAKATTAPATTRNTGRGGEEKKKEKEKEEQEERETKTPEC
PSHTQPLGVYLLTPAVQDLWLRDKATFTCFVVGSDLKDAHLTWEVAGKVPTGGVEEGL
LERHSNGSQSQHSRLTLPRSLWNAGTSVTCTLNHPSLPPQRLMALREPAAQAPVKLSL

FIGURE 7

EE

intron
 NLLASSDPPEAASWLLCEVSGFSPPNILLMWLEDQREVNTSGFAPARPPPQPRSTTFW
 AWSVLRVPAPPSPQPATYTCVVSHEDSRTLLNASRSLEVSYLAMTPLIPQSKDENSDD
 YTTFDDVGSLWTTLSTFVALFILTLLYSGIVTFIKVK"
 404..>495
 /gene="IGHD"
 /note="G00-120-084"
 /number=1
 BASE COUNT 114 a 179 c 120 g 82 t
 ORIGIN 182 bp upstream of SphI site; chromosome 14q32.3.
 1 tttccctgcc tcccgtcacc ctgccgccag ggcctctgcc ctgccctgcc ccttgcctc
 61 aggtttccag cctcagactc ccactgtgtc tgtcttccag caccaccaa ggctccgat
 121 gtgttcccca tcatatcagg gtgcagacac ccaaaggata acagccctgt ggtcctggca
 181 tgcttgataa ctgggtacca cccaacgtcc gtgactgtca cctggtacat ggggacacag
 241 agccagcccc agagaacctt ccctgagata caaagacggg acagctacta catgacaagc
 301 agccagctct ccacccccct ccagcagtgg cgccaaggcg agtacaaatg cgtggtccag
 361 cacaccgcca gcaagagtaa gaaggagatc ttccgctggc caggtaggtc gcaccggaga
 421 tcaccagaa gggcccccca ggacccccag caccttcac tcagggcctg accacaaaga
 481 cagaagcaag ggctg

VIII. HUMAN IG EPSILON CHAIN C REGION - HOMO SAPIENS (HUMAN).

AMINO ACID SEQUENCE

>sp|P01854|EPC_HUMAN IG EPSILON CHAIN C REGION - Homo sapiens (Human).

10	20	30	40	50	60
ASTQSPSVFP	LTRCCKNIPS	NATSVTLGCL	ATGYFPEPVM	VTWDTGSLNG	TTMTLPATTL
70	80	90	100	110	120
TLSGHYATIS	LLTVSGAWAK	QMFTCRVAHT	PSSTDWVDNK	TFSVCSRDFI	PPTVKILQSS
130	140	150	160	170	180
CDGGGHFPPT	IQLLCLVSGY	TPGTINITWL	EDGQVMDVDL	STASTTQEGE	LASTQSELT
190	200	210	220	230	240
SQKHWLSDRT	YTCQVTYQGH	TFEDSTKKCA	DSNPRGVSAY	LSRPSPFDLF	IRKSPTITCL
250	260	270	280	290	300
VVDLAPSKCT	VNLTWSRASG	KPVNHSTRKE	EKQRNGTLTV	TSTLPVGTRD	WIEGETYQCR
310	320	330	340	350	360
VTHPHLPRAL	MRSTTKTSGP	RAAPEVYafa	TPEWPGSRDK	RTLACLIQNF	MPEDISVQWL
370	380	390	400	410	420
HNEVQLPDAR	HSTTQPRKTK	GSGFFVFSRL	EVTRAWEQK	DEFICRAVHE	AASPSQTVQR
428					

FIGURE 7 FF

AVSVNPGK

CODING SEQUENCE

```

atggactgga cctggatcct cttcttgggtg gcagcagcca cgcgagtcca ctcccagacg      60
cagttggtgc agtctggggc tgaggtgagg aagcctgggg catcagttag ggtctcctgc      120
aaggcttctg gatacacctt catcgactcc tatatccact ggatacgaca ggcccctggg      180
cacgggcttg agtgggtggg atggatcaac cctaacagtg gtggcacaaa ctatgctccg      240
agatttcagg gcaggggtcac catgaccaga gacgcgtcct tcagtacagc ctacatggac      300
ctgagaagtc tgagatctga cgactcggcc gtgttttact gtgcgaaaag tgaccctttt      360
tggagtgatt attataaactt tgactactcg tacactttgg acgtctgggg ccaagggacc      420
acgggtcaccg tctcctcagc ctccacacag agcccatccg tcttcccctt gaccgctgc      480
tgcaaaaaca ttccctccaa tgccacctcc gtgactctgg gctgcctggc cacgggctac      540
ttcccggagc cgggtgatggg gacctgggac acaggctccc tcaacgggac aactatgacc      600
ttaccagcca ccacctcac gctctctggg cactatgcc aacacagctt gctgaccgtc      660
tcgggtgcgt gggccaagca gatgttcacc tgcctgtgtg cacacactcc atcgtccaca      720
gactgggtcg acaacaaaac cttcagcgtc tgcctcaggg acttcacccc gccacccgtg      780
aagatcttac agtcgtcctg cgacggcgcc gggcacttcc ccccgaccat ccagctcctg      840
tgctcgtctt ctgggtacac cccagggact atcaacatca cctggctgga ggacgggcag      900
gtcatggacg tggacttgte caccgcctct accacgcagg agggtgagct ggctccaca      960
caaagcgagc tcacctcag ccagaagcac tggctgtcag accgcacctt cactgccag      1020
gtcacctate aaggtcacac ctttgaggac agcaccaaga agtgtgcaga ttccaacccg      1080
agaggggtga gcgcctacct aagccggccc agcccgttcg acctgttcat ccgcaagtcg      1140
cccacgatca cctgtctggt ggtggacctg gcacccagca aggggaccgt gaacctgacc      1200
tgggtcccgg ccagtgggaa gcctgtgaac cactccacca gaaaggagga gaagcagcgc      1260
aatggcacgt taaccgtcac gtccacctg cgggtgggca ccgagactg gatcgagggg      1320
gagacctacc agtgcagggt gacccacccc cacctgcccc gggccctcat gcgggtccacg      1380
accaagacca gcggcccgcg tgctgccccg gaagtctatg cgtttgcgac gccggagtgg      1440
ccggggagcc gggacaagcg caccctcgcc tgctgatcc agaacttcat gcctgaggac      1500
atctcgggtg agtggctgca caacgagggt cagctcccg acgcccggca cagcacgacg      1560
cagccccgca agaccaaggg ctccggcttc ttctgtttca gccgcctgga ggtgaccagg      1620
gccgaatggg agcagaaaag tgagttcatc tgccgtgcag tccatgaggc agcgagcccc      1680
tcacagaccg tccagcgagc ggtgtctgta aatcccggta aatga      1725

```

GenBank

L00022. Human Ig active he...

LOCUS HUMIGHAE2 1920 bp DNA PRI 22-DEC-1994

DEFINITION Human Ig active heavy chain epsilon-1 gene, constant region.

ACCESSION L00022 J00227 V00555

VERSION L00022.1 GI:185035

KEYWORDS C-region; epsilon-immunoglobulin; immunoglobulin heavy chain; processed gene.

SOURCE Human myeloma cell line 266B1 DNA and cDNA to mRNA, clones H-Ig-epsilon-11, lambda-epsilon-1.2, pJU71, pGET2 and K85/A12 (see comment).

ORGANISM Homo sapiens

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.

REFERENCE 1 (bases 1 to 1920)

AUTHORS Flanagan,J.G. and Rabbitts,T.H.

TITLE The sequence of a human immunoglobulin epsilon heavy chain constant region gene, and evidence for three non-allelic genes

JOURNAL EMBO J. 1 (5), 655-660 (1982)

MEDLINE 84236029

REFERENCE 2 (bases 528 to 736; 1044 to 1138)

AUTHORS Nishida,Y., Miki,T., Hisajima,H. and Honjo,T.

TITLE Cloning of human immunoglobulin epsilon chain genes: evidence for multiple C epsilon genes

JOURNAL Proc. Natl. Acad. Sci. U.S.A. 79 (12), 3833-3837 (1982)

MEDLINE 82247945

REFERENCE 3 (bases 1 to 1920)

AUTHORS Kenten,J.H., Molgaard,H.V., Houghton,M., Derbyshire,R.B., Viney,J., Bell,L.O. and Gould,H.J.

TITLE Cloning and sequence determination of the gene for the human

FIGURE 7 GG

immunoglobulin epsilon chain expressed in a myeloma cell line

JOURNAL Proc. Natl. Acad. Sci. U.S.A. 79 (21), 6661-6665 (1982)

MEDLINE 83065234

REFERENCE 4 (bases 98 to 1884)

AUTHORS Seno,M., Kurokawa,T., Ono,Y., Onda,H., Sasada,R., Igarashi,K., Kikuchi,M., Sugino,Y., Nishida,Y. and Honjo,T.

TITLE Molecular cloning and nucleotide sequencing of human immunoglobulin epsilon chain cDNA

JOURNAL Nucleic Acids Res. 11 (3), 719-726 (1983)

MEDLINE 83168897

REFERENCE 5 (bases 691 to 807; 1571 to 1818; 1860 to 1885)

AUTHORS Liu,F.T., Albrandt,K.A., Bry,C.G. and Ishizaka,T.

TITLE Expression of a biologically active fragment of human IgE epsilon chain in Escherichia coli

JOURNAL Proc. Natl. Acad. Sci. U.S.A. 81 (17), 5369-5373 (1984)

MEDLINE 84298140

COMMENT [2] and [1] report the isolation of two other epsilon genes, epsilon-2 and epsilon-3. The authors in [2] claim that epsilon-3 is a pseudogene. Compared in [4] with the germline C-region sequence by Max, et al (Cell 29, 691-699 (1982)), and there are three nucleotide differences. The deduced amino acid sequence in [4] differs somewhat from the published C-region sequence. [5] produced expression of IgE in E.coli by insertion into expression vector pUC7.

Complete source information:
Human myeloma cell line 266B1 DNA [2],[1],[5] and cDNA to mRNA [3], [4], clones H-Ig-epsilon-11 [2], lambda-epsilon-1.2 [1], pJJ71 [3], pGET2 [4] and K85/A12 [5].

FEATURES

Location/Qualifiers

source 1..1920
/organism="Homo sapiens"
/db_xref="taxon:9606"
/map="14q32.33"

prim_transcript <1..1886
/note="epsilon-1 mRNA"

intron <1..97
/gene="IGHE"
/note="epsilon-1 intron J-C"

exon 98..406
/gene="IGHE"
/note="Ig heavy chain epsilon-1 (CH1 domain); G00-119-335"

intron 407..613
/gene="IGHE"
/note="epsilon-1 intron A"

exon 614..934
/gene="IGHE"
/note="Ig heavy chain epsilon-1 (CH2 domain)"

conflict 735
/gene="IGHE"
/citation=[3]
/replace=""

intron 935..1020
/gene="IGHE"
/note="epsilon-1 intron B"

exon 1021..1344
/gene="IGHE"
/note="Ig heavy chain epsilon-1 (CH3 domain)"

conflict 1124
/gene="IGHE"
/citation=[3]
/replace=""

conflict 1337
/gene="IGHE"

FIGURE 7

HH

```

/citation=[3]
/replace=""
intron 1345..1427
/gene="IGHF"
/note="epsilon-1 intron C"
exon 1428..>1759
/note="Ig heavy chain epsilon-1 (CH4 domain)"
conflict 1444..1445
/gene="IGHF"
/citation=[3]
/replace=""
conflict 1612
/gene="IGHF"
/citation=[3]
/replace=""
conflict 1785
/citation=[3]
/replace=""
gene join(L00021.1:57..495,1..1758)
/gene="IGHF"
CDS join(L00021.1:57..495,98..406,614..934,1021..1344,
1428..1759)
/partial
/gene="IGHF"
/note="Ig heavy chain epsilon-1 (V-D-J region)"
/codon_start=1
/protein_id="AAB59424.1"
/db_xref="GI:386807"
/db_xref="GDB:G00-119-335"
/translation="MDWTWILFLVAAATRVHSQTQLVQSGAEVRKPGASVRVSCKASG
YTFIDSYIHWIRQAPGHGLEWVGWINPNSGGTNYAPRFQGRVTMTRDASFSTAYMDLR
SLRSDDSAVFYCAKSDPFWSDDYNYFDYSYTLVDVWGQGTITVTVSSASTQSPSVFPLTRC
CKNIPSNATSVTLGCLATGYFPEFVMVTWDTGSLNGTMTLPATTLTSLGHYATISLL
TVSGAWAKQMFTCRVAHTPSSTDWVDNKTFSVCSRDFTPPTVKILOSSCDGGGHFPPT
IQLLCLVSGYTPGTINITWLEDGQVMDVDLSTASTTQEGELASTQSELTLSQKHWLSD
RTYTCQVTYQGHTFEDSTKKCADSNPRGVSAYLSRPSFDFIRKSPTITCLVVDLAP
SKGTVNLTWSRASGKPVNHSTRKEEKQRNGTLTVTSTLPVGTDRDWIEGETYQCRVTHP
HLPRALMRSTTKTSGPRAAPEVYAFATPEWPGSRDKRTLACLIQNFMPEDISVQWLHN
EVQLPDAHSTTQPRKTKGSGFFVFSRLEVTRAWEQKDEFICRAVHEAASPSQTVQR
AVSVNPGK"
BASE COUNT 387 a 658 c 576 g 299 t
ORIGIN About 3 kb after ; 1 bp upstream of BamHI site.
1 ggatccctgc caccggggtcc ccagctcccc catccaggcc cccagggctg atgggagctg
61 gactgagget ggcactgact aggtctctgtc ctcacagcct ccacacagag cccatccgtc
121 ttcccttga cccgctgctg caaaaacatt cctcccaatg ccacctccgt gactctgggc
181 tgcctggcca cgggctactt cccggagccg gtgatgggtga cctgggacac aggcctccctc
241 aacgggacaa ctatgacctt accagccacc accctcacgc tctctgggtca ctatgccacc
301 atcagcttgc tgaccgtctc ggggtgcgtgg gccaaagcaga tgttcacctg ccgtgtggca
361 cacactccat cgtccacaga ctgggtcgac aacaaaacct tcagcggtaa gagagggcca
421 agctcagaga ccacagttcc caggagtgc aggcgtgagg ctggcagagt gggcaggggt
481 tgaggggggt ggtgggctca aacgtgggaa caccagcat gcctggggac ccgggcccagg
541 acgtgggggc aagaggaggg cacacagagc tcagagaggc caacaacct catgaccacc
601 agctctcccc cagtctctc cagggacttc accccgcca ccgtgaagat cttacagtcg
661 tcctgcgacg gcggcgggca cttccccccg accatccagc tcctgtgcct cgtctctggg
721 tacaccccag ggactatcaa catcacctgg ctggaggacg ggcaggtcat ggacgtggac
781 ttgtccaccg cctctaccac gcaggagggt gagctggcct ccacacaaag cgagctcacc
841 ctcagccaga agcactggct gtcagaccgc acctacacct gccaggtcac ctatcaaggt
901 cacacctttg aggacagcac caagaagtgt gcaggtacgt tcccacctgc cctggtggcc
961 gccacggagg ccagagaaga ggggcgggtg ggcctcacac agccctccgg tgtaccacag
1021 attccaaccc gagaggggtg agcgccctacc taagccggcc cagcccggtc gacctgttca
1081 tccgcaagtc gccacgata acctgtctgg tgggtggacct ggcacccagc aaggggaccg
1141 tgaacctgac ctgggtcccg gccagtgga agcctgtgaa ccactccacc agaaaggagg
1201 agaagcagcg caatggcacg ttaaccgtca cgtccacct gccggtgggc acccgagact

```

FIGURE 7 II

```

1261 ggatcgaggg ggagacctac cagtgcaggg tgacccaccc ccacctgccc agggccctca
1321 tgcggtccac gaccaagacc agcgggtgagc catgggcagg ccgggggtcgt gggggaaggg
1381 agggagcgag tgagcggggc ccgggctgac cccacgtctg gccacaggcc cgcgtgctgc
1441 cccggaagtc tatgcgtttg cgacgccgga gtggccgggg agccgggaca agcgcaccct
1501 cgcctgcctg atccagaact tcatgcctga ggacatctcg gtgcagtggc tgcacaacga
1561 ggtgcagctc ccggacgccc ggcacagcac gacgcagccc cgcaagacca agggctccgg
1621 cttcttcgtc ttcagccgcc tggaggtgac cagggccgaa tgggagcaga aagatgagtt
1681 catctgccgt gcagtccatg aggcagcgag cccctcacag accgtccagc gagcgggtgc
1741 tgtaaattccc ggtaaatgac gtactcctgc ctccctccct cccagggctc catccagctg
1801 tgcagtgggg aggactggcc agaccttctg tccactgttg caatgacccc aggaagctac
1861 cccaataaaa ctgtgcctgc tcagagcccc agtacacca ttcttgggag cgggcagggc

```

IX. HUMAN IG MU CHAIN C REGION - HOMO SAPIENS (HUMAN).**AMINO ACID SEQUENCE**

>sp|P01871|MUC_HUMAN IG MU CHAIN C REGION - Homo sapiens (Human).

```

      10      20      30      40      50      60
      |      |      |      |      |      |
GSASAPTLFP LVSCENSPSD TSSVAVGCLA QDFLPDSITF SWKYKNNSDI SSTRGFPSVL
      70      80      90     100     110     120
      |      |      |      |      |      |
RGGKYAATSQ VLLPSKDVMQ GTDEHVVKV  QHPNGNKEKN VPLPVIAELP PKVSVFVPPR
     130     140     150     160     170     180
      |      |      |      |      |      |
DGFFGNPRSK SKLICQATGF SPRQIQVSWL REGKQVGSV  TTDQVQAEAK ESGPTTYKVT
     190     200     210     220     230     240
      |      |      |      |      |      |
STLTIKESDW LSQSMFTCRV DHRGLTFQQN ASSMCVPDQD TAIRVFAIPP SFASIFLTKS
     250     260     270     280     290     300
      |      |      |      |      |      |
TKLTCLVTDL TTYDSVTISW TRQNGEAVKT HTNISESHPN ATFSAVGEAS ICEDDWNSGE
     310     320     330     340     350     360
      |      |      |      |      |      |
RFTCTVIHTD LPSPLKQTIS RPKGVALHRP DVYLLPPARE QLNLRRESATI TCLVTGFSPA
     370     380     390     400     410     420
      |      |      |      |      |      |
DVFVQWMQRG QPLSPEKYVT SAPMPEPQAP GRYFAHSILT VSEEEWNTGE TYTCVVAHEA
     430     440     450
      |      |      |
LPNRVTERTV DKSTGKPTLY NVSLVMSDTA GTCY

```

CODING SEQUENCE

```

atggactgga cctggagggt cctctttgtg gtggcagcag ctacaggtgt ccagtcccag      60
gtgcagctgg tgcagtctgg ggctgagggt aagaagcctg ggctctcggt gaaggtctcc      120
tgcaaggcct ctggaggcac cttcagcagc tatgctatca gctgggtgcg acaggcccct      180
ggacaagggc ttgagtggat gggagggatc atccctatct ttggtacagc aaactacgca      240
cagaagttcc agggcagagt cagcattacc gcggacgaat ccacgagcac agcctacatg      300
gagctgagca gcctgagatc tgaggacacg gccgtgtatt actgtgcgaa aaccgggatc      360
ctggggccgt atagcagtgg ctggtacccg aactcggact actactacta cggtatggac      420
gtctggggcc aagggaccac ggtcaccgtc tcttcaggga gtgcatccgc cccaaccctt      480
ttccccctcg tctcctgtga gaattccccg tcggatacga gcagcgtggc cgttggctgc      540
ctcgcacagg acttccttcc cgactccatc actttctcct ggaaatacaa gaacaactct      600
gacatcagca gcacccgggg cttcccatca gtctgagag ggggcaagta cgcagccacc      660

```

FIGURE 7

JJ

```

tcacaggtgc tgctgccttc caaggacgtc atgcagggca cagacgaaca cgtggtgtgc      720
aaagtccagc accccaacgg caacaaagaa aagaacgtgc ctcttccagt gattgctgag      780
ctgcctccca aagtgagcgt ctctgtccca ccccgcgacg gcttcttcgg caacccccgc      840
agcaagtcca agctcatctg ccaggccacg ggtttcagtc cccggcagat tcaggtgtcc      900
tggctgcgcg aggggaagca ggtggggtct ggcgtcacca cggaccaggt gcaggctgag      960
gccaaagagt ctgggcccac gacctacaag gtgaccagca cactgaccat caaagagagc     1020
gactggctca gccagagcat gttcacctgc cgcgtggatc acaggggcct gaccttccag     1080
cagaatgcgt cctccatgtg tgtccccgat caagacacag ccatccgggt cttcgccatc     1140
ccccatcct ttgccagcat cttcctcacc aagtccacca agttgacctg cctggtcaca     1200
gacctgacca cctatgacag cgtgaccatc tcctggacct gccagaatgg cgaagctgtg     1260
aaaaccacaca ccaacatctc cgagagccac cccaatgcc ctttcagcgc cgtgggtgag     1320
gccagcatct gcgaggatga ctggaattcc ggggagaggt tcacgtgcac cgtgaccac     1380
acagacctgc cctcgccact gaagcagacc atctcccggc ccaagggggt ggccctgcac     1440
agggccgatg tctacttgct gccaccagcc cgggagcagc tgaacctgcg ggagtcggcc     1500
accatcacgt gcctggtgac gggcttctct cccgcccagc tcttcgtgca gtggatgcag     1560
agggggcagc ccttgtcccc ggagaagtat gtgaccagcg cccaatgcc tgagccccag     1620
gccccaggcc ggtacttcgc ccacagcatc ctgacctgt ccgaagagga atggaacacg     1680
ggggagacct acacctgcgt ggtggcccat gaggccctgc ccaacagggt caccgagagg     1740
accgtggaca agtccaccga gggggagggt agcgcgcgac aggagggtt tgagaacctg     1800
tgggcccacc cctccacctt catcgtcctc ttctcctga gcctcttcta cagtaccacc     1860
gtcaccttgt tcaagggtgaa atga                                           1884

```

GenBank

X17115. Human mRNA for IgM

LOCUS HSI201 2213 bp mRNA PRI 03-APR-1995

DEFINITION Human mRNA for IgM heavy chain complete sequence.

ACCESSION X17115

VERSION X17115.1 GI:33450

KEYWORDS Ig heavy chain; IgM gene; IgM heavy chain; transmembrane protein.

SOURCE human.

ORGANISM Homo sapiens

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.

REFERENCE 1 (bases 1 to 2213)

AUTHORS Friedlander, R.M.

TITLE Direct Submission

JOURNAL Submitted (03-NOV-1989) Friedlander R. M., Harvard Medical School,
Howard Hughes Medical Institute, Department of Genetics, 25
Shattuck St, Boston, MA 02115, USA

REFERENCE 2 (bases 1 to 2213)

AUTHORS Friedlander, R.M., Nussenzweig, M.C. and Leder, P.

TITLE Complete nucleotide sequence of the membrane form of the human IgM
heavy chain

JOURNAL Nucleic Acids Res. 18 (14), 4278 (1990)

MEDLINE 90332450

REFERENCE 3 (bases 1 to 2213)

AUTHORS Kristensen, T., Lopez, R. and Prydz, H.

TITLE An estimate of the sequencing error frequency in the DNA sequence
databases

JOURNAL DNA Seq. 2 (6), 343-346 (1992)

MEDLINE 93075997

REMARK Erratum: [[published erratum appears in DNA Seq 1993;3(5):337]]

COMMENT For genomic sequence see <K01306>, <X14939> and
<X14940>. The author reports various conflicts with these
sequences. Data kindly reviewed (30-MAY-1990) by Friedlander R.M.

FEATURES

Location/Qualifiers

source

1..2213

/organism="Homo sapiens"

/db_xref="taxon:9606"

/clone="201-203"

/cell_line="lymphoma 201"

/cell_type="B"

FIGURE 7

KK

```

misc_feature      /tissue_type="lymphoid"
                  1..39
                  /note="putative VECTOR sequence Bluescript SKP+"
                  /citation=[3]
CDS               73..1956
                  /note="precursor (AA -15 to 612)"
                  /codon_start=1
                  /protein_id="CAA34971.1"
                  /db_xref="GI:33451"
                  /db_xref="SWISS-PROT:P01871"
                  /db_xref="SWISS-PROT:P20769"
                  /translation="MDWTWRFLEVVAAATGVQSQVQLVQSGAEVKKPGSSSVKVSCKAS
GGTFSSYAIWVRQAPGQGLEWMGGIIPFGTANYAQKFQGRVTITADESTSTAYMEL
SSLRSEDVAVYYCAKTGILGPYSSGWYPNSDYIYYGMDVWGQGTITVTVSSGSASAPTL
FPLVSCENSPSDTSSVAVGCLAQDFLPDSITFSWKYKNSDISSTRGFPSVLRGGKYA
ATSQVLLPSKDVMOGTDEHVVCKVQHPNGNKEKNVPLFVIAELPPKVSFVFPDRDGF
GNPRSKSKLICQATGFSPRQIQVSWLREGKQVSGVTTIDQVQAEAKESGPTTYKVTST
LTIKESDWLSQSMFTCRVDHRGLTFQQNASSMCPDQDTAIRVFAIPPSFASIFLTKS
TKLTCLVTDLTITYDSVTISWTRQNGEAVKTHTNISESHENATFSAVGEASICEDDWS
GERFTCTVTHTDLPSPKQITSRPKGVALHRPDVYLLPPAREQLNLRESATITCLVTG
FSPADVVFVQWMQRGQPLSPEKYVTSAPMPEPQAPGRYFAHSILTVSEEEWNTGETYTC
VVAHEALPNRVTERTVDKSTEGEVSADDEGFENLWATASTFIVLFLLSLFYSTTVTLF
KVK"
sig_peptide       73..117
mat_peptide       118..1953
                  /product="IgM heavy chain (AA 1 to 612)"
polyA_site        2213
                  /note="polyA site"
BASE COUNT       462 a    708 c    629 g    414 t
ORIGIN
    1 gctctagaac tagtggatcc cccgggctgc aggaattctc taaagaagcc cctgggagca
   61 cagctcatca ccatggactg gacctggagg ttctctcttg tgggtggcagc agctacaggt
  121 gtccagtcce aggtgcagct ggtgcagctc ggggctgagg tgaagaagcc tgggtcctcg
  181 gtgaaggtct cctgcaaggc ttctggaggc accttcagca gctatgctat cagctgggtg
  241 cgacaggccc ctggacaagg gcttgagtgg atgggaggga tcatccctat ctttggtaca
  301 gcaaaactacg cacagaagtt ccagggcaga gtcacgatta ccgcggacga atccacgagc
  361 acagcctaca tggagctgag cagcctgaga tctgaggaca cggccgtgta ttactgtgcg
  421 aaaaccggga tcctggggcc gtatagcagt ggctgggtacc cgaactcgga ctactactac
  481 tacggatatg acgtctgggg ccaagggacc acggtcaccg tctcctcagg gagtgcctcc
  541 gccccaaacc ttttccccct cgtctcctgt gagaattccc cgtcgggatac gagcagcgtg
  601 gccgttggtg gcctcgcaca ggacttcctt cccgactcca tcactttctc ctggaaatac
  661 aagaacaact ctgacatcag cagcaccctg ggcttcccat cagtcctgag aggggggcaag
  721 tacgcagcca cctcacaggt gctgctgcct tccaaggacg tcatgcaggg cacagacgaa
  781 cacgtggtgt gcaaagtcca gcaccccaac ggcaacaaag aaaagaacgt gcctcttcca
  841 gtgattgctg agctgcctcc caaagtgagc gtcttcgtcc caccctcgga cggcttcttc
  901 ggcaaccccc gcagcaagtc caagctcatc tgccaggcca cgggtttcag tccccggcag
  961 attcagggtg cctggctgcg cgagggggaag caggtggggt ctggcgtcac cacggaccag
 1021 gtgcaggctg aggccaaaga gtctgggccc acgacctaca aggtgaccag cactactgac
 1081 atcaaagaga gcgactggct cagccagagc atgttcacct gccgcgtgga tcacaggggc
 1141 ctgaccttcc agcagaatgc gtcctccatg tgtgtccccg atcaagacac agccatccgg
 1201 gtcttcgcca tccccccatc ctttgccagc atcttctca ccaagtccac caagttgacc
 1261 tgcttggtca cagacctgac cacttatgac agcgtgacca tctcctggac ccgccagaat
 1321 ggcgaagctg tgaaaaccca caccaacatc tccgagagcc accccaatgc cactttcagc
 1381 gccgtgggtg aggccagcat ctgcgaggat gactggaatt ccggggagag gttcacgtgc
 1441 accgtgaccc acacagacct gccctcgcca ctgaagcaga ccatctcccg gccaagggg
 1501 gtggccctgc acaggcccca tgtctacttg ctgccaccag cccgggagca gctgaacctg
 1561 cgggagtcgg ccacctcac gtgcctgggt acgggcttct ctcccgcgga cgtcttcgtg
 1621 cagtggatgc agagggggca gcccttgtec ccggagaagt atgtgaccag cgcaccaatg
 1681 cctgagcccc agggcccagg ccggtacttc gccacagca tctgaccgt gtccgaagag
 1741 gaatggaaca cgggggagac ctacacctgc gtgggtggcc atgaggccct gccaacagg
 1801 gtcaccgaga ggaccgtgga caagtccacc gagggggagg tgagcgccga cgaggagggc
 1861 tttgagaacc tgtggggcac cgcctccacc ttcctcgtcc tcttctcctc gagcctcttc
 1921 tacagtacca ccgtcacctt gttcaagggt aatgatccc aacagaagaa catcggagac

```

FIGURE 7 LL

1981 cagagagagg aactcaaagg ggcgctgcct ccgggtcttg ggtcctggcc tgcgtggcct
2041 gttggcacgt gtttctcttc ccgcccggcc tccagttgtg tgetctcaca caggcttcct
2101 tctcgaccgg caggggctgg ctggcttgca ggccacgagg tgggctctac cccacactgc
2161 ttgctgtgt atacgcttgt tgccctgaaa taaatatgca cattttatcc atg

HindIII SalI ClaI

1 GAACTCGAGCAGCTGAAGCTTGCATGCCTGCAGGTCGACGGTATCGATAAGGATCCCTGAAAGCGACGTTGGATGTTAAC
81 ATCTACAAATTGCCTTTTCTTATCGACCATGTACGTAAGCGCTTACGTTTTTGGTGGACCCTTGAGGAACTGGTAGCTG
161 TTGTGGGCTGTGGTCTCAAGATGGATCATTAAATTTCCACCTTCACCTACGATGGGGGGCATCGCACCGGTGAGTAATAT
241 TGTACGGCTAAGAGCGAATTGCGCTGTAGGATCCCTGAAAGCGACGTTGGATGTTAACATCTACAAATTGCCTTTTCTT
321 ATCGACCATGTACGTAAGCGCTTACGTTTTTGGTGGACCCTTGAGGAACTGGTAGCTGTTGTGGGCTGTGGTCTCAAG
401 ATGGATCATTAAATTTCCACCTTCACCTACGATGGGGGGCATCGCACCGGTGAGTAATATTGTACGGCTAAGAGCGAATTT
481 GGCCTGTAGGATCCCTGAAAGCGACGTTGGATGTTAACATCTACAAATTGCCTTTTCTTATCGACCATGTACGTAAGCGC
561 TTACGTTTTTGGTGGACCCTTGAGGAACTGGTAGCTGTTGTGGGCTGTGGTCTCAAGATGGATCATTAAATTTCCACCT
641 TCACCTACGATGGGGGGCATCGCACCGGTGAGTAATATTGTACGGCTAAGAGCGAATTGCGCTGTAGGATCCGCGAGCT
721 GGTCAATCCCATTGCTTTTGAAGCAGCTCAACATTGATCTCTTTCTCGAGGGAGATTTTTCAAATCAGTGCGCAAGACGT
801 GACGTAAGTATCCGAGTCAGTTTTTATTTTTCTACTAATTTGGTCGTTTTATTTTCGGCGTGTAGGACATGGCAACCGGGCC
881 TGAATTTTCGCGGGTATTCTGTTTCTATTCCAACCTTTTCTTGATCCGCAGCCATTAACGACTTTTGAATAGATACGCTGA
961 CACGCCAAGCCTCGCTAGTCAAAAGTGTACCAAACAACGCTTTACAGCAAGAACGGAATGCGCGTGACGCTCGCGGTGAC
1041 GCCATTTTCGCTTTTTCAGAAATGGATAAATAGCCTTGCTTCTATTATATCTTCCCAAATTACCAATACATTACACTAGC

ClaI

1121 ATCTGAATTTTCATAACCAATCTCGATACACCAAATCGACTCTAGAGGATCTATCGATTCCCGGGTACCATGGGATCT
MetGlySer

SpeI

1198 AAACCTTTTTTGTCTCTTCTTTCATTGTCATTGCTTTTGTTTACATCTACTAGTTTG
LysProPheLeuSerLeuLeuSerLeuSerLeuLeuLeuPheThrSerThrSerLeu
1255 GCACAGACATCTGTGTCCCCCTCAAAGTCATCCTGCCCGGGGAGGCTCCGTG
AlaGlnThrSerValSerProSerLysValIleLeuProArgGlyGlySerVal
1309 CTGGTGACATGCAGCACCTCCTGTGACCAGCCCAAGTTGTTGGGCATAGAGACCCCGTTG
LeuValThrCysSerThrSerCysAspGlnProLysLeuLeuGlyIleGluThrProLeu
1369 CCTAAAAGGAGTTGCTCCTGCCTGGGAACAACCGGAAGGTGTATGAACTGAGCAATGTG
ProLysLysGluLeuLeuLeuProGlyAsnAsnArgLysValTyrGluLeuSerAsnVal
1429 CAAGAAGATAGCCAACCAATGTGCTATTCAAACCTGCCCTGATGGGCAGTCAACAGCTAAA
GlnGluAspSerGlnProMetCysTyrSerAsnCysProAspGlyGlnSerThrAlaLys
1489 ACCTTCCTCACCGTGTACTGGACTCCAGAACGGGTGGAACCTGGCACCCCTCCCCTCTTGG
ThrPheLeuThrValTyrTrpThrProGluArgValGluLeuAlaProLeuProSerTrp
1549 CAGCCAGTGGGCAAGAACCTTACCCTACGCTGCCAGGTGGAGGGTGGGGCACCCCGGGCC
GlnProValGlyLysAsnLeuThrLeuArgCysGlnValGluGlyGlyAlaProArgAla
1609 AACCTCACCGTGGTGCTGCTCCGTGGGAGAAGGAGCTGAAACGGGAGCCAGCTGTGGGG
AsnLeuThrValValLeuLeuArgGlyGluLysGluLeuLysArgGluProAlaValGly
1669 GAGCCCGCTGAGGTCACGACCACGGTGCTGGTGAGGAGAGATCACCATGGAGCCAATTTT
GluProAlaGluValThrThrThrValLeuValArgArgAspHisHisGlyAlaAsnPhe
1729 TCGTGCCGCACTGAACTGGACCTGCGGCCCCAAGGGCTGGAGCTGTTTGAGAACACCTCG
SerCysArgThrGluLeuAspLeuArgProGlnGlyLeuGluLeuPheGluAsnThrSer
1789 GCCCCCTACCAGCTCCAGACCTTTGTCTGCCAGCGACTCCCCACAACCTTGTCAGCCCC
AlaProTyrGlnLeuGlnThrPheValLeuProAlaThrProProGlnLeuValSerPro
1849 CGGGTCCTAGAGGTGGACACGCAGGGGACCGTGGTCTGTTCCCTGGACGGGCTGTTCCCA
ArgValLeuGluValAspThrGlnGlyThrValValCysSerLeuAspGlyLeuPhePro
1909 GTCTCGGAGGCCAGGTCCACCTGGCACTGGGGGACCAGAGGTGAAACCCACAGTCACC
ValSerGluAlaGlnValHisLeuAlaLeuGlyAspGlnArgLeuAsnProThrValThr
1969 TATGGCAACGACTCCTTCTCGGCCAAGGCCTCAGTCAGTGTGACCGCAGAGGACGAGGGC
TyrGlyAsnAspSerPheSerAlaLysAlaSerValSerValThrAlaGluAspGluGly
2029 ACCCAGCGGCTGACGTGTGCAGTAATACTGGGGAACCAGAGCCAGGAGACACTGCAGACA
ThrGlnArgLeuThrCysAlaValIleLeuGlyAsnGlnSerGlnGluThrLeuGlnThr
2089 GTGACCATCTACAGCTTTCCGGCGCCCAACGTGATTCTGACGAAGCCAGAGGTCTCAGAA
ValThrIleTyrSerPheProAlaProAsnValIleLeuThrLysProGluValSerGlu
2149 GGGACCGAGGTGACAGTGAAGTGTGAGGCCACCCCTAGAGCCAAGGTGACGCTGAATGGG
GlyThrGluValThrValLysCysGluAlaHisProArgAlaLysValThrLeuAsnGly
2209 GTTCCAGCCCAGCCACTGGGCCCCAGGGCCCAGCTCCTGCTGAAGGCCACCCAGAGGAC
ValProAlaGlnProLeuGlyProArgAlaGlnLeuLeuLeuLysAlaThrProGluAsp
2269 AACGGGCGCAGCTTCTCCTGCTCTGCAACCCTGGAGGTGGCCGGCCAGCTTATACACAAG
AsnGlyArgSerPheSerCysSerAlaThrLeuGluValAlaGlyGlnLeuIleHisLys

FIGURE 8 A - 1

2329 AACCAGACCCGGGAGCTTCGTGTCCTGTATGGCCCCGACTGGACGAGAGGGATTGTCCG
 AsnGlnThrArgGluLeuArgValLeuTyrGlyProArgLeuAspGluArgAspCysPro
 2389 GGAAACTGGACGTGGCCAGAAAATTCCCAGCAGACTCCAATGTGCCAGGCTTGGGGGAAC
 GlyAsnTrpThrTrpProGluAsnSerGlnGlnThrProMetCysGlnAlaTrpGlyAsn
 SacI
 2449 CCATTGCCCCGAGCTCAAGTGTCTAAAGGATGGCACTTTCCCACTGCCCATCGGGGAATCA
 ProLeuProGluLeuLysCysLeuLysAspGlyThrPheProLeuProIleGlyGluSer
 2509 GTGACTGTCACTCGAGATCTTGAGGGCACCTACCTCTGTCCGGCCAGGAGCACTCAAGGG
 ValThrValThrArgAspLeuGluGlyThrTyrLeuCysArgAlaArgSerThrGlnGly
 SpeI SacI
 2569 GAGGTCACCCGCGAGGTGACCGTGAATGTGACTAGTGGGAGCTCAGCATCCCCG
 GluValThrArgGluValThrValAsnValThrSerGlySerSerAlaSerPro
 2623 ACCAGCCCCAAGGTCTTCCCGCTGAGCCTCGACAGCACCCCCAAGATGGGAACGTGGTC
 ThrSerProLysValPheProLeuSerLeuAspSerThrProGlnAspGlyAsnValVal
 2683 GTCGCATGCCTGGTCCAGGGCTTCTTCCCCCAGGAGCCACTCAGTGTGACCTGGAGCGAA
 ValAlaCysLeuValGlnGlyPhePheProGlnGluProLeuSerValThrTrpSerGlu
 2743 AGCGGACAGAACGTGACCGCCAGAACTTCCCACCTAGCCAGGATGCCTCCGGGGACCTG
 SerGlyGlnAsnValThrAlaArgAsnPheProProSerGlnAspAlaSerGlyAspLeu
 2803 TACACCACGAGCAGCCAGCTGACCCTGCCGGCCACACAGTGCCCGAGACGGCAAGTCCGTG
 TyrThrThrSerSerGlnLeuThrLeuProAlaThrGlnCysProAspGlyLysSerVal
 2863 ACATGCCACGTGAAGCACTACACGAATTCAGCCAGGATGTGACTGTGCCCTGCCGAGTT
 ThrCysHisValLysHisTyrThrAsnSerSerGlnAspValThrValProCysArgVal
 2923 CCCCCACCTCCCCCATGCTGCCACCCCCGACTGTCTGCTGCACCGACCGGCCCTCGAGGAC
 ProProProProProCysCysHisProArgLeuSerLeuHisArgProAlaLeuGluAsp
 2983 CTGCTCTTAGGTTTCAAGCGAACCTCACGTGCACACTGACCGGCCTGAGAGATGCCTCT
 LeuLeuLeuGlySerGluAlaAsnLeuThrCysThrLeuThrGlyLeuArgAspAlaSer
 3043 GGTGCCACCTTCACCTGGACGCCCTCAAGTGGGAAGAGCGCTGTTCAAGGACCACCTGAG
 GlyAlaThrPheThrTrpThrProSerSerGlyLysSerAlaValGlnGlyProProGlu
 3103 CGTGACCTCTGTGGCTGCTACAGCGTGTCCAGAGTACTTCTTGGCTGTGCCAGCCATGG
 ArgAspLeuCysGlyCysTyrSerValSerArgValLeuProGlyCysAlaGlnProTrp
 3163 AACCATGGGGAGACCTTCACCTGCACTGCTGCCACCCCCGAGTTGAAGACCCCACTAACC
 AsnHisGlyGluThrPheThrCysThrAlaAlaHisProGluLeuLysThrProLeuThr
 3223 GCCAACATCACAAAATCCGGAAACACATTCCGGCCCCGAGGTCCACCTGCTGCCGCCGCCG
 AlaAsnIleThrLysSerGlyAsnThrPheArgProGluValHisLeuLeuProProPro
 3283 TCGGAGGAGCTGGCCCTGAACGAGCTGGTGACGCTGACGTGCCTGGCACGTGGCTTCAGC
 SerGluGluLeuAlaLeuAsnGluLeuValThrLeuThrCysLeuAlaArgGlyPheSer
 3343 CCCAAGGATGTGCTGGTTCGCTGGCTGCAGGGGTACAGGAGCTGCCCCGCGAGAAGTAC
 ProLysAspValLeuValArgTrpLeuGlnGlySerGlnGluLeuProArgGluLysTyr
 3403 CTGACTTGGGCATCCCGGAGGAGCCAGCCAGGGCACCCACCTATGCTGTGACCAGC
 LeuThrTrpAlaSerArgGlnGluProSerGlnGlyThrThrThrTyrAlaValThrSer
 3463 ATACTGCGCGTGGCAGCCGAGGACTGGAAGAAGGGGGAGACCTTCTCCTGCATGGTGGGC
 IleLeuArgValAlaAlaGluAspTrpLysLysGlyGluThrPheSerCysMetValGly
 3523 CACGAGGCCCTGCCGCTGGCCTTCACACAGAAGACCATCGACCGCTTGGCGGGTAAACCC
 HisGluAlaLeuProLeuAlaPheThrGlnLysThrIleAspArgLeuAlaGlyLysPro
 3583 ACCCATATCAATGTGTCTGTTGTATGGCGGAGGCGGACGGCACCTGCTACAGATCTGAA
 ThrHisIleAsnValSerValValMetAlaGluAlaAspGlyThrCysTyrArgSerGlu
 junEcoRI marker
 3643 AAGGATGAACCTTTAGAATTCGATATCAAGCTAATTCGCGATCGTTCAAACATTTGGCAATAAAGTTTCTTAAGAT
 LysAspGluLeu
 3718 TGAATCCTGTTGCCGGTCTTGCGATGATTATCATATAATTTCTGTTGAATTACGTTAAGCATGTAATAATTAACATGTAA
 3798 TGCATGACGTTATTTATGAGATGGGTTTTTATGATTAGAGTCCCGCAATTATACATTTAATACGCGATAGAAAACAAAAT
 3878 ATAGCGCGCAAACTAGGATAAATTATCGCGCGCGGTGTCATCTATGTTACTAGATCGGGGATCTGCCGGTCTCCCTATAG
 3958 TGAGTCGTATTAATTTTCGATAAGCCAGGTAACTTGCATTAATGAATCGGCCAACGCGCGGGGAGAGGCGGTTTGCGTAT
 4038 TGGGCGCTCTTCCGCTTCTCGCTCACTGACTCGCTGCGCTCGGTGCTTGGCTGCGGCGAGCGGTATCAGCTCACTCAA
 4118 AGGCGGTAATACGGTTATCCACAGAATCAGGGGATAACGCAGGAAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAGG
 4198 AACCCTAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACAAAATCGACGCTCAAG
 4278 TCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGCGTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTT

FIGURE 8 A - 2

4358 CGACCCTGCCGCTTACCGGATACCTGTCCGCTTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCAATGCTCACGCTGTAGG
4438 TATCTCAGTTCGGTGTAGGTCGTTTCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCCGTTCAGCCCGACCGCTGCGCCTT
4518 ATCCGGTAACTATCGTCTTGAGTCCAACCCGGTAAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTA
4598 GCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAAGGACAGTATTT
4678 GGTATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAACAAACCACCGCTGG
4758 TAGCGGTGGTTTTTTTTGTGTTGCAAGCAGCAGATTACGCGCAGAAAAAAAGGATCTCAAGAAGATCCTTTGATCTTTTCTA
4838 CGGGGTCTGACGCTCAGTGGAACGAAAACCTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGGATCTTCACCTAG
4918 ATCCTTTTAAATTAAAAATGAAGTTTAAATCAATCTAAAGTATATATGAGTAAACTTGGTCTGACAGTTACCAATGCTT
4998 AATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTTTCGTTTCATCCATAGTTGCCTGACTCCCCGTCGTGTAGATAACTA
5078 CGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGACCCACGCTCACCGGCTCCAGATTTATCA
5158 GCAATAAACCCAGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGTCCTGCAACTTTATCCGCCTCCATCCAGTCTATTAATTG
5238 TTGCCGGGAAGCTAGAGTAAGTAGTTCCGCCAGTTAATAGTTTTCGCAACGTTGTTGCCATTGCTACAGGCATCGTGGTGT
5318 CACGCTCGTCGTTTGGTATGGCTTCATTACGCTCCGTTCCCAACGATCAAGGCGAGTTACATGATCCCCCATGTTGTGC
5398 AAAAAAGCGGTTAGCTCCTTCGGTCCCTCCGATCGTTGTCAGAAGTAAGTTGGCCGAGTGTTATCACTCATGGTTATGGC
5478 AGCACTGCATAATTCTCTTACTGTCATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCT
5558 GAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATAGCAGAACTTTA
5638 AAAGTGCTCATCATTTGGAAAACGTTCTTCGGGGCGAAAACCTCTCAAGGATCTTACCGCTGTTGAGATCCAGTTCGATGTA
5718 ACCCACTCGTGACCCCACTGATCTTCAGCATCTTTTACTTTTACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAA
5798 ATGCCGCAAAAAAGGGAATAAGGGCGACACGGAAATGTTGAATACTCATACTCTTCCTTTTCAATATTATTGAAGCATT
5878 TATCAGGGTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGAAAAATAAACAAATAGGGGTTCGCGCACATT
5958 TCCCCGAAAAGTGCCACCTGACGTCTAAGAAACCATTATTATCATGACATTAACCTATAAAAAATAGGCGTATCACGAGGC
6038 CCTTTCGTCTCGCGCGTTTCGGTGATGACGGTGAAAACCTCTGACACATGCAGCTCCCGGAGACGGTCACAGCTTGTCTG
6118 TAAGCGGATGCCGGGAGCAGACAAGCCCGTCAGGGCGCGTCAGCGGGTGTGTCGGGGCTGGCTTAACTATGC
NdeI
6198 GGCATCAGAGCAGATTGTACTGAGAGTGCACCATATGGACATATTGTCGTTAGAACGCGGCTACAATTAATACATAACCT
6278 TATGTATCATAACATAACGATTTAGGTGACACTATA

FIGURE 8 A – 3

Bean Legumin Signal Peptide

MetSerLysProPheLeuSerLeuLeuSerLeuSerLeuLeuLeuPheThrSerThrCysLeuAla

FIGURE 8 B

Nucleotide and Amino Acid Sequence of Protein Coding Region of pSHuJ

1141

AGGATCTATCGATTCCCGGGTACC ATG GAG AAC CAT TTG CTT TTC TGG GGA GTC CTG GCG
met glu asn his leu leu phe trp gly val leu ala

1201/13

GTT TTT ATT AAG GCT GTT CAT GTG AAA GCC CAA GAA GAT GAA AGG ATT GTT CTT GTT GAC
val phe ile lys ala val his val lys ala gln glu asp glu arg ile val leu val asp

1261/33

AAC AAA TGT AAG TGT GCC CGG ATT ACT TCC AGG ATC ATC CGT TCT TCC GAA GAT CCT AAT
asn lys cys lys cys ala arg ile thr ser arg ile ile arg ser ser glu asp pro asn

1321/53

GAG GAC ATT GTG GAG AGA AAC ATC CGA ATT ATT GTT CCT CTG AAC AAC AGG GAG AAT ATC
glu asp ile val glu arg asn ile arg ile ile val pro leu asn asn arg glu asn ile

1381/73

TCT GAT CCC ACC TCA CCA TTG AGA ACC AGA TTT GTG TAC CAT TTG TCT GAC CTC TGT AAA
ser asp pro thr ser pro leu arg thr arg phe val tyr his leu ser asp leu cys lys

1441/93

AAA TGT GAT CCT ACA GAA GTG GAG CTG GAT AAT CAG ATA GTT ACT GCT ACC CAG AGC AAT
lys cys asp pro thr glu val glu leu asp asn gln ile val thr ala thr gln ser asn

1501/113

ATC TGT GAT GAA GAC AGT GCT ACA GAG ACC TGC TAC ACT TAT GAC AGA AAC AAG TGC TAC
ile cys asp glu asp ser ala thr glu thr cys tyr thr tyr asp arg asn lys cys tyr

1561/133

ACA GCT GTG GTC CCA CTC GTA TAT GGT GGT GAG ACC AAA ATG GTG GAA ACA GCC TTA ACC
thr ala val val pro leu val tyr gly gly glu thr lys met val glu thr ala leu thr

1621/153

CCA GAT GCC TGC TAT CCT GAC TGA ATTC
pro asp ala cys tyr pro asp

FIGURE 8 C

Nucleotide and Amino Acid Sequence of Protein Coding Region of pSHuSC

1137

GTCGATTCCCGGGTACC ATG GTG CTC TTC GTG CTC ACC TGC
met val leu phe val leu thr cys

1178/9

CTG CTG GCG GTC TTC CCA GCC ATC TCC ACG AAG AGT CCC ATA TTT GGT CCC GAG GAG GTG
leu leu ala val phe pro ala ile ser thr lys ser pro ile phe gly pro glu glu val

1238/29

AAT AGT GTG GAA GGT AAC TCA GTG TCC ATC ACG TGC TAC TAC CCA CCC ACC TCT GTC AAC
asn ser val glu gly asn ser val ser ile thr cys tyr tyr pro pro thr ser val asn

1298/49

CGG CAC ACC CGG AAG TAC TGG TGC CGG CAG GGA GCT AGA GGT GGC TGC ATA ACC CTC ATC
arg his thr arg lys tyr trp cys arg gln gly ala arg gly gly cys ile thr leu ile

1358/69

TCC TCG GAG GGC TAC GTC TCC AGC AAA TAT GCA GGC AGG GCT AAC CTC ACC AAC TTC CCG
ser ser glu gly tyr val ser ser lys tyr ala gly arg ala asn leu thr asn phe pro

1418/89

GAG AAC GGC ACA TTT GTG GTG AAC ATT GCC CAG CTG AGC CAG GAT GAC TCC GGG CGC TAC
glu asn gly thr phe val val asn ile ala gln leu ser gln asp asp ser gly arg tyr

1478/109

AAG TGT GGC CTG GGC ATC AAT AGC CGA GGC CTG TCC TTT GAT GTC AGC CTG GAG GTC AGC
lys cys gly leu gly ile asn ser arg gly leu ser phe asp val ser leu glu val ser

1538/129

CAG GGT CCT GGG CTC CTA AAT GAC ACT AAA GTC TAC ACA GTG GAC CTG GGC AGA ACG GTG
gln gly pro gly leu leu asn asp thr lys val tyr thr val asp leu gly arg thr val

1598/149

ACC ATC AAC TGC CCT TTC AAG ACT GAG AAT GCT CAA AAG AGG AAG TCC TTG TAC AAG CAG
thr ile asn cys pro phe lys thr glu asn ala gln lys arg lys ser leu tyr lys gln

1658/169

ATA GGC CTG TAC CCT GTG CTG GTC ATC GAC TCC AGT GGT TAT GTG AAT CCC AAC TAT ACA
ile gly leu tyr pro val leu val ile asp ser ser gly tyr val asn pro asn tyr thr

1718/189

GGA AGA ATA CGC CTT GAT ATT CAG GGT ACT GGC CAG TTA CTG TTC AGC GTT GTC ATC AAC
gly arg ile arg leu asp ile gln gly thr gly gln leu leu phe ser val val ile asn

1778/209

CAA CTC AGG CTC AGC GAT GCT GGG CAG TAT CTC TGC CAG GCT GGG GAT GAT TCC AAT AGT
gln leu arg leu ser asp ala gly gln tyr leu cys gln ala gly asp asp ser asn ser

1838/229

AAT AAG AAG AAT GCT GAC CTC CAA GTG CTA AAG CCC GAG CCC GAG CTG GTT TAT GAA GAC
asn lys lys asn ala asp leu gln val leu lys pro glu pro glu leu val tyr glu asp

1898/249

CTG AGG GGC TCA GTG ACC TTC CAC TGT GCC CTG GGC CCT GAG GTG GCA AAC GTG GCC AAA
leu arg gly ser val thr phe his cys ala leu gly pro glu val ala asn val ala lys

1958/269

TTT CTG TGC CGA CAG AGC AGT GGG GAA AAC TGT GAC GTG GTC GTC AAC ACC CTG GGG AAG
phe leu cys arg gln ser ser gly glu asn cys asp val val val asn thr leu gly lys

2018/289

AGG GCC CCA GCC TTT GAG GGC AGG ATC CTG CTC AAC CCC CAG GAC AAG GAT GGC TCA TTC
arg ala pro ala phe glu gly arg ile leu leu asn pro gln asp lys asp gly ser phe

2078/309

AGT GTG GTG ATC ACA GGC CTG AGG AAG GAG GAT GCA GGG CGC TAC CTG TGT GGA GCC CAT
ser val val ile thr gly leu arg lys glu asp ala gly arg tyr leu cys gly ala his

2138/329

TCG GAT GGT CAG CTG CAG GAA GGC TCG CCT ATC CAG GCC TGG CAA CTC TTC GTC AAT GAG
ser asp gly gln leu gln glu gly ser pro ile gln ala trp gln leu phe val asn glu

2198/349

GAG TCC ACG ATT CCC CGC AGC CCC ACT GTG GTG AAG GGG GTG GCA GGA AGC TCT GTG GCC

FIGURE 8 D - 1

glu ser thr ile pro arg ser pro thr val val lys gly val ala gly ser ser val ala
 2258/369
 GTG CTC TGC CCC TAC AAC CGT AAG GAA AGC AAA AGC ATC AAG TAC TGG TGT CTC TGG GAA
 val leu cys pro tyr asn arg lys glu ser lys ser ile lys tyr trp cys leu trp glu
 2318/389
 GGG GCC CAG AAT GGC CGC TGC CCC CTG CTG GTG GAC AGC GAG GGG TGG GTT AAG GCC CAG
 gly ala gln asn gly arg cys pro leu leu val asp ser glu gly trp val lys ala gln
 2378/409
 TAC GAG GGC CGC CTC TCC CTG CTG GAG GAG CCA GGC AAC GGC ACC TTC ACT GTC ATC CTC
 tyr glu gly arg leu ser leu leu glu glu pro gly asn gly thr phe thr val ile leu
 2438/429
 AAC CAG CTC ACC AGC CGG GAC GCC GGC TTC TAC TGG TGT CTG ACC AAC GGC GAT ACT CTC
 asn gln leu thr ser arg asp ala gly phe tyr trp cys leu thr asn gly asp thr leu
 2498/449
 TGG AGG ACC ACC GTG GAG ATC AAG ATT ATC GAA GGA GAA CCA AAC CTC AAG GTT CCC GGG
 trp arg thr thr val glu ile lys ile ile glu gly glu pro asn leu lys val pro gly
 2558/469
 AAT GTC ACG GCT GTG CTG GGA GAG ACT CTC AAG GTC CCC TGT CAC TTT CCA TGC AAA TTC
 asn val thr ala val leu gly glu thr leu lys val pro cys his phe pro cys lys phe
 2618/489
 TCC TCG TAC GAG AAA TAC TGG TGC AAG TGG AAT AAC ACG GGC TGC CAG GCC CTG CCC AGC
 ser ser tyr glu lys tyr trp cys lys trp asn asn thr gly cys gln ala leu pro ser
 2678/509
 CAA GAC GAA GGC CCC AGC AAG GCC TTC GTG AAC TGT GAC GAG AAC AGC CGG CTT GTC TCC
 gln asp glu gly pro ser lys ala phe val asn cys asp glu asn ser arg leu val ser
 2738/529
 CTG ACC CTG AAC CTG GTG ACC AGG GCT GAT GAG GGC TGG TAC TGG TGT GGA GTG AAG CAG
 leu thr leu asn leu val thr arg ala asp glu gly trp tyr trp cys gly val lys gln
 2798/549
 GGC CAC TTC TAT GGA GAG ACT GCA GCC GTC TAT GTG GCA GTT GAA GAG AGG AAG GCA GCG
 gly his phe tyr gly glu thr ala ala val tyr val ala val glu glu arg lys ala ala
 2858/569
 GGG TCC CGC GAT GTC AGC CTA GCG AAG GCA GAC GCT GCT CCT GAT GAG AAG GTG CTA GAC
 gly ser arg asp val ser leu ala lys ala asp ala ala pro asp glu lys val leu asp
 2918/589
 TCT GGT TTT CGG GAG ATT GAG AAC AAA GCC ATT CAG GAT CCC AGG CTT TTT GCA GAG TGA
 ser gly phe arg glu ile glu asn lys ala ile gln asp pro arg leu phe ala glu
 2978
 ATTC

1 CTGGCCGGCGCCAGATCTGGGGAACCTGTGGTTGGCATGCACATACAAATGGACGAACGGATAAACCTTTTCACGCCCTT
 81 TTAAATATCCGATTATTCTAATAAACGCTCTTTTCTCTTAGGTTTACCCGCCAATATATCCTGTCAAACACTGATAGTTT
 161 AAAGTGAAGGCGGGAAACGACAATCTGATCATGAGCGGAGAATTAAGGGAGTCACGTTATGACCCCGCCGATGACGCGG
 241 GACAAGCCGTTTTACGTTTGGAACCTGACAGAACCGCAACGATTGAAGGAGCCACTCAGCCGATCTGAATTAATCCCGAT
 321 CTAGTAACATAGATGACACCGCGCGCGATAATTTATCCTAGTTTGCGCGCTATATTTTGTCTATCGCGTATTAAATG
 401 TATAATTGCGGGACTCTAATCATAAAAACCCATCTCATAAATAACGTCATGCATTACATGTTAATTATTACATGCTTAAC
 481 GTAATTCAACAGAAATTATATGATAATCATCGCAAGACCGGCAACAGGATTCAATCTTAAGAACTTTATTGCCAAATGT
 561 TTGAACGATCGGGGAAATTCGAGCTCCACCGCGGTGGCGGCCGCTCTAGAACTAGTGGATCCCCCGGGCTGCAGGAATTC
 641 GATCAGATCTGATCAAGCTTATCGATAACCGTCGACCTCGAGGGGGGGCCCGGTACCCCTAGAGTCGATTTGGTGTATCGA
 721 GATTGGTTATGAAATTCAGATGCTAGTGTAATGTATTGGTAATTTGGGAAGATATAATAGGAAGCAAGGCTATTTATCCA
 801 TTTCTGAAAAGGCGAAATGGCGTCACCGCGAGCGTCACGCGCATTCGGTTCTTGCTGTAAAGCGTTGTTTGGTACACTTT
 881 TGACTAGCGAGGCTTGGCGTGTGAGCGTATCTATTCAAAGTCGTTAATGGCTGCGGATCAAGAAAAAGTTGGAATAGAA
 961 ACAGAATACCCGCGAAATTCAGGCCCGGTTGCCATGTCCTACACGCCGAAATAAACGACCAAATTAGTAGAAAAATAAAA
 1041 ACTGACTCGGATACTTACGTCACGTCTTGCGCACTGATTTGAAAAATCTCCCTCGATCGAGAAAGAGATCAATGTTGAGC
 1121 TGCTTCAAAGCAATGGGATTGACCAGCTCGCGGATCCTACAGGCCAAATTCGCTCTTAGCCGTACAATATTACTCACCG
 1201 GTGCGATGCCCCCATCGTAGGTGAAGGTGGAAATTAATGATCCATCTTGAGACCACAGGCCCAACAGCTACCAGTTT
 1281 CCTCAAGGGTCCACCAAAAACGTAAGCGCTTACGTACATGGTCGATAAGAAAAGGCAATTTGTAGATGTTAACATCCAAC
 1361 GTCGCTTTCAGGGATCCTACAGGCCAAATTCGCTCTTAGCCGTACAATATTACTCACCGGTGCGATGCCCCCATCGTAG

SacI NotI XmaI
 SmaI
 SalI XhoI KpnI
 BamHI
 BamHI

FIGURE 8 E - 1

1441 GTGAAGGTGGAAATTAATGATCCATCTTGAGACCACAGGCCACAAACAGCTACCAGTTTCCTCAAGGGTCCACCAAAAAC
 BamHI
 1521 GTAAGCGCTTACGTACATGGTCGATAAGAAAAGGCAATTTGTAGATGTTAACATCCAACGTCGCTTTCAGGGATCCTACA
 1601 GGCCAAATTCGCTCTTAGCCGTACAATATTACTCACCGGTGCGATGCCCCCATCGTAGGTGAAGGTGGAAATTAATGAT
 1681 CCATCTTGAGACCACAGGCCACAAACAGCTACCAGTTTCCTCAAGGGTCCACCAAAAACGTAAGCGCTTACGTACATGGT
 BamHI
 1761 CGATAAGAAAAGGCAATTTGTAGATGTTAACATCCAACGTCGCTTTCAGGGATCCGCGAGCTTATCGCGATACCGTCGAA
 1841 TATAATAATTATATTTGTAAGAATATTATTATAATAATATAAAATATATATATAAAATTATAATATATTAATTATTGTT
 1921 AATTATTAATAATATATATATTAATCATTTAGATATATAATTCTATAGCCTTAGACTCCTCATCAATAGAAGACTACGTA
 2001 TAAAAATAATCAGATAACATCTAAACATGTAGATAAATAAGTTGTTTCATATCCAACATGATGTCCAGAGCTTCACG
 2081 CTGCCGCAAGCACTCAGGGCGCAAGGGCTGCTAAAGGAAGCGGAACACGTAGAAAGCCAGTCCGCGAGAANCAGGTGCTGAC
 2161 CCCGGATGAATGTCAGCTACTGGGCTATCTGGACAAGGGAAAACGCAAGCGCANNAGAGAAAGCAGGTAGCTTGCAAGTGG
 2241 GCTTACATGGCGATAGCTAGACTGGGCGGTTTTATGGACAGCAAGCGAACCAGGAAATTGCCAGCTGGGGCGCCCTCTGGTA
 2321 AGGTTGGGAAGCCCTGCAAAGTAACTGGATGGCTTTCTTGGCCGAAGGATCTGATGGCGCAGGGGATCAAGATCATGA
 2401 GCGGAGAATTAAGGGAGTCACGTTATGACCCCGCCGATGACGCGGGACAAGCCGTTTTACGTTTGAACTGACAGAACC
 2481 GCAACGTTGAAGGAGCCACTCAGCCGCGGGTTTTCTGGAGTTTAATGAGCTAAGCACATACGTCAGAAACCATTATTGCGC
 2561 GTTCAAAGTCGCCTAAGGTCAGCTAGCAAATATTTCTTGTCAAAAATGCTCCACTGACGTTCCATAAATTCCC
 2641 CTCGGTATCCAATTAGAGTCTCATATTCATCTCAATCCAGATCTGGATCGTTTCGCATGATTGAACAAGATGGATTGCA
 MetIleGluGlnAspGlyLeuHi
 2721 CGCAGGTTCTCCGGCCGCTTGGGTGGAGAGGCTATTCCGGCTATGACTGGGCACAACAGACAATCGGCTGCTCTGATGCCG
 sAlaGlySerProAlaAlaTrpValGluArgLeuPheGlyTyrAspTrpAlaGlnGlnThrIleGlyCysSerAspAlaA
 2801 CCGTGTTCGGCTGTCAGCGCAGGGGCGCCCGGTTCTTTTTGTCAAGACCGACCTGTCCGGTGCCCTGAATGAACTGCAG
 laValPheArgLeuSerAlaGlnGlyArgProValLeuPheValLysThrAspLeuSerGlyAlaLeuAsnGluLeuGln
 2881 GACGAGGCAGCGCGGCTATCGTGGCTGGCCACGACGGGCGTTCTTGGCGAGCTGTGCTCGACGTTGTCAGTGAAGCGGG
 AspGluAlaAlaArgLeuSerTrpLeuAlaThrThrGlyValProCysAlaAlaValLeuAspValValThrGluAlaGl
 2961 AAGGGACTGGCTGCTATTGGGCGAAGTGCCGGGGCAGGATCTCCTGTATCTCACCTTGCTCCTGCCGAGAAAGTATCCA
 yArgAspTrpLeuLeuGlyGluValProGlyGlnAspLeuLeuSerSerHisLeuAlaProAlaGluLysValSerI
 3041 TCATGGCTGATGCAATGCGGCGGCTGCATACGCTTGATCCGGCTACCTGCCCATTCGACCACCAAGCGAAACATCGCATC
 leMetAlaAspAlaMetArgArgLeuHisThrLeuAspProAlaThrCysProPheAspHisGlnAlaLysHisArgIle
 3121 GAGCGAGCACGTACTCGGATGGAAGCCGGTCTTGTGATCAGGATGATCTGGACGAAGAGCATCAGGGGCTCGCGCCAGC
 GluArgAlaArgThrArgMetGluAlaGlyLeuValAspGlnAspAspLeuAspGluGluHisGlnGlyLeuAlaProAl
 3201 CGAACTGTTCCGCGAGGCTCAAGGCGCGCATGCCCGACGCGGATGATCTCGTCTGACCCATGGCGATGCCTGCTTGCCGA
 aGluLeuPheAlaArgLeuLysAlaArgMetProAspGlyAspAspLeuValValThrHisGlyAspAlaCysLeuProA
 3281 ATATCATGGTGGAAAATGGCCGCTTTTCTGGATTTCATCGACTGTGGCCGGCTGGGTGTGGCGGACCGCTATCAGGACATA
 snIleMetValGluAsnGlyArgPheSerGlyPheIleAspCysGlyArgLeuGlyValAlaAspArgTyrGlnAspIle
 3361 GCGTTGGCTACCCGTGATATTGCTGAAGAGCTTGGCGGCGAATGGGCTGACCGCTTCCTCGTGCTTTACGGTATCGCCGC
 AlaLeuAlaThrArgAspIleAlaGluGluLeuGlyGlyGluTrpAlaAspArgPheLeuValLeuTyrGlyIleAlaAl
 BamHI
 3441 TCCCGATTTCGAGCGCATCGCCTTCTATCGCCTTCTTGACGAGTTCTTCTGAGCGGGACTCTGAGGATCCCCCGATGAGC
 aProAspSerGlnArgIleAlaPheTyrArgLeuLeuAspGluPhePhe...
 3521 TAAGCTAGCTATATCATCAATTTATGTATTACACATAATATCGCACTCAGTCTTTCATCTACGGCAATGTACCAGCTGAT
 3601 ATAATCAGTTATTGAAATATTTCTGAATTTAACTTGCATCAATAAATTTATGTTTTTGCTTGGACTATAATACCTGACT
 3681 TGTTATTTTATCAATAAATATTTAACTATATTTCTTTCAAGATGGGAATTAATTCAGTGGCCGTCGTTTTACAACGTCG
 3761 TGAAGTGGGAAAACCCCTGGCGTTACCCAACCTTAATCGCCTTGCAGCACATCCCCCTTTCGCCAGCTGGCGTAATAGCGAAG
 PvuI BglI
 3841 AGGCCCCGACCGATCGCCCTTCCCAACAGTTGCGCAGCCTGAATGGCGCCCGCTCCTTTTCGCTTTCTTCCCTTCCTTTCT
 3921 CGCCACGTTCCGCGGCTTTCCCGTCAAGCTCTAAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACC
 4001 TCGACCCCAAAAACTTGATTTGGGTGATGGTTCACGTAGTGGGCCATCGCCCTGATAGACGGTTTTTCGCCCTTTGACG
 4081 TTGGAGTCCACGTTCTTTAATAGTGGACTCTTGTTCCAACTGGAACAACACTCAACCCTATCTCGGGCTATTCTTTTGA
 4161 TTTATAAGGGATTTTGCCGATTTCCGAACCACCATCAAACAGGATTTTCGCCTGCTGGGGCAAACCAGCGTGGACCGCTT
 4241 GCTGCAACTCTCTCAGGGCCAGGCGGTGAAGGGCAATCAGCTGTTGCCCGTCTCACTGGTGAAAAGAAAAACCACCCAG
 4321 TACATTAATAACGTCGCAATGTGTTATTAAGTTGTCTAAGCGTCAATTTGTTTACACCACAATATATCCTGCCACCAGC
 4401 CAGCCAACAGCTCCCCGACCGGCAGCTCGGCACAAAATCACCCTCGATACAGGCAGCCCATCAG

FIGURE 8 E - 2

FIGURE 8 F - 1

1nHisGluGluAspLeuAspAspGlnAspValLeuGlyAlaGluMetArgThrArgAlaArgGluIleArgHisLysAla
 1361 TTGGTGGTTCGAATGGGCAGGTAGCCGGATCAAGCGTATGCAGCCGCCGCATTGCATCAGCCATGATGGATACTTTCTCGG
 GlnHisAspPheProCysThrAlaProAspLeuThrHisLeuArgArgMetAlaAspAlaMetIleSerValLysGluAl
 1441 CAGGAGCAAGGTGAGATGACAGGAGATCCTGCCCCGGCACTTCGCCCCAATAGCAGCCAGTCCCTTCCCGCTTCAGTGACA
 aProAlaLeuHisSerSerLeuLeuAspGlnGlyProValGluGlyLeuLeuLeuTrpAspArgGlyAlaGluThrValV
 PstI
 1521 ACGTCGAGCACAGCTGCGCAAGGAACGCCCGTCGTGGCCAGCCACGATAGCCGCGCTGCCTCGTCCTGCAGTTCATTCAG
 alAspLeuValAlaAlaCysProValGlyThrThrAlaLeuTrpSerLeuArgAlaAlaGluAspGlnLeuGluAsnLeu
 Sfo I
 1601 GGCACCGGACAGGTCTGGTCTTGACAAAAAGAACCGGGCGCCCCTGCGCTGACAGCCGGAACACGGCGGCATCAGAGCAGC
 AlaGlySerLeuAspThrLysValPheLeuValProArgGlyGlnAlaSerLeuArgPheValAlaAlaAspSerCysGl
 1681 CGATTGTCTGTTGTGCCAGTCATAGCCGAATAGCCTCTCCACCCAAGCGGCCGAGAACCTGCGTGCAATCCATCTTGT
 yIleThrGlnGlnAlaTrpAspTyrGlyPheLeuArgGluValTrpAlaAlaProSerGlyAlaHisLeuGlyAspGlnG
 BglII
 1761 TCAATCATGCGAAACGATCCAGATCTGGATTGAGAGTGAATATGAGACTCTAATTGGATACCGAGGGGAATTTATGGAAC
 luIleMet
 1841 GTCAGTGGAGCATTTTTGGACAAGAAATATTTGCTAGCTGATAGTGACCTTAGGCGACTTTTGAACGCGCAATAATGGTTT
 SacII
 1921 CTGACGTATGTGCTTAGCTCATTAAGTCCAGAAACCCGCGGCTGAGTGGCTCCTTCAACGTTGCGGTTCTGTCAGTTCC
 2001 AAACGTAAACGGCTTGTCCCGCGTCATCGGCGGGGTCATAACGTGACTCCCTTAATTCTCCGCTCATGATCTTGATCC
 Sfo I
 2081 CCTGCGCCATCAGATCCTTGGCGGCAAGAAAGCCATCCAGTTTACTTTGCAGGGCTTCCCAACCTTACCAGAGGGCGCCC
 2161 CAGCTGGCAATTCCGGTTCGCTTGCTGTCCATAAAACCGCCAGTCTAGCTATCGCCATGTAAGCCCACTGCAAGCTACC
 2241 TGCTTCTCTTTGCGCTTGCCTTTCCCTTGCTCCAGATAGCCAGTAGCTGACATTCATCCGGGGTTCAGCACCGNTTCTG
 2321 CGGACTGGCTTTCTACGTGTTCCGCTTCCTTTAGCAGCCCTTGCGCCCTGAGTGCTTGCGGCAGCGTGAAGCTCTGGACA
 2401 TCATGTTGGATATGAAACAACCTATTATTTATCTACATGTTTTAGATGTTATCTGATTATTTTTTATACGTAGTCTTCTATT
 2481 GATGAGGAGTCTAAGGCTATAGAATTATATATCTAAATGATTAATATATATATTATTAATAATTAACAATAATTAATATA
 NruI
 2561 TTATAATTTATATATATATATATTTTATATTATTATAATAATATTCTTACAAATATAATTATTATATTCGACGGTATCGCGA
 2641 TAAGCTCGCGGATCCCTGAAAGCGACGTTGGATGTTAACATCTACAAATTGCCTTTTCTTATCGACCATGTACGTAAGCG
 2721 CTTACGTTTTTGGTGGACCCTTGAGGAACTGGTAGCTGTTGTGGGCTGTGGTCTCAAGATGGATCATTAAATTTCCACC
 2801 TTCACCTACGATGGGGGGCATCGCACCGGTGAGTAATATTGTACGGCTAAGAGCGAATTTGGCCTGTAGGATCCCTGAAA
 2881 GCGACGTTGGATGTTAACATCTACAAATTGCCTTTTCTTATCGACCATGTACGTAAGCGCTTACGTTTTTGGTGGACCCT
 2961 TGAGGAACTGGTAGCTGTTGTGGGCTGTGGTCTCAAGATGGATCATTAAATTTCCACCTTCACCTACGATGGGGGGCAT
 3041 CGCACCGGTGAGTAATATTGTACGGCTAAGAGCGAATTTGGCCTGTAGGATCCCTGAAAGCGACGTTGGATGTTAACATC
 3121 TACAAATTGCCTTTTCTTATCGACCATGTACGTAAGCGCTTACGTTTTTGGTGGACCCTTGAGGAACTGGTAGCTGTTG
 3201 TGGGCTGTGGTCTCAAGATGGATCATTAAATTTCCACCTTCACCTACGATGGGGGGCATCGCACCGGTGAGTAATATTGT
 3281 ACGGCTAAGAGCGAATTTGGCCTGTAGGATCCGCGAGCTGGTCAATCCCATTGCTTTTGAAGCAGCTCAACATTGATCTC
 3361 TTTCTCGATCGAGGGAGATTTTCAAATCAGTGCGCAAGACGTGACGTAAGTATCCGAGTCAGTTTTTATTTTCTACTA
 3441 ATTTGGTTCGTTTATTTTCGGCGTGTAGGACATGGCAACCGGGCTGAATTTTCGCGGGTATTCGTTTCTATTCCAACCTTTT
 3521 TCTTGATCCGCGAGCCATTAACGACTTTTGAATAGATACGCTGACACGCCAAGCCTCGCTAGTCAAAGTGTAACAAACAA
 3601 CGCTTTACAGCAAGAACGGAATGCGCGTGACGCTCGCGGTGACGCCATTTTCGCTTTTTCAGAAATGGATAAATAGCCTTG
 3681 CTTCTTATTATATCTTCCCAAATTACCAATACATTACACTAGCATCTGAATTTTCATAACCAATCTCGATACACCAAATCG
 KpnI
 3761 ACTCTAGGGGTACCATGGTGCTCTTCGTGCTCACCTGCCTGCTGGCGGTCTTCCCAGCCATC
 MetValLeuPheValLeuThrCysLeuLeuAlaValPheProAlaIle
 3823 TCCACGAAGAGTCCCATATTTGGTCCCAGGAGGTGAATAGTGTGGAAGGTAAGTCAAGT
 SerThrLysSerProIlePheGlyProGluGluValAsnSerValGluGlyAsnSerVal
 3883 TCCATCACGTGCTACTACCCACCCACCTCTGTCAACCGGCACACCCGGAAGTACTGGTGC
 SerIleThrCysTyrTyrProProThrSerValAsnArgHisThrArgLysTyrTrpCys
 3943 CGGCAGGGAGCTAGAGGTGGCTGCATAACCCTCATCTCCTCGGAGGGCTACGTCTCCAGC
 ArgGlnGlyAlaArgGlyGlyCysIleThrLeuIleSerSerGluGlyTyrValSerSer
 4003 AAATATGCAGGCAGGGCTAACCTCACCAACTTCCCGGAGAACGGCACATTTGTGGTGAAC
 LysTyrAlaGlyArgAlaAsnLeuThrAsnPheProGluAsnGlyThrPheValValAsn
 4063 ATTGCCAGCTGAGCCAGGATGACTCCGGGCGCTACAAGTGTGGCCTGGGCATCAATAGC
 IleAlaGlnLeuSerGlnAspAspSerGlyArgTyrLysCysGlyLeuGlyIleAsnSer
 4123 CGAGGCCTGTCCTTTGATGTCAGCCTGGAGGTCAGCCAGGGTCCTGGGCTCCTAAATGAC

FIGURE 8 F - 2

ArgGlyLeuSerPheAspValSerLeuGluValSerGlnGlyProGlyLeuLeuAsnAsp
 4183 ACTAAAGTCTACACAGTGGACCTGGGCAGAACGGTGACCATCAACTGCCCTTTCAAGACT
 ThrLysValTyrThrValAspLeuGlyArgThrValThrIleAsnCysProPheLysThr
 4243 GAGAATGCTCAAAGAGGAAGTCCTTGTACAAGCAGATAGGCCTGTACCCTGTGCTGGTC
 GluAsnAlaGlnLysArgLysSerLeuTyrLysGlnIleGlyLeuTyrProValLeuVal
 4303 ATCGACTCCAGTGGTTATGTGAATCCCAACTATACAGGAAGAATACGCCTTGATATTCAG
 IleAspSerSerGlyTyrValAsnProAsnTyrThrGlyArgIleArgLeuAspIleGln
 4363 GGTACTGGCCAGTTACTGTTTCAGCGTTGTCATCAACCAACTCAGGCTCAGCGATGCTGGG
 GlyThrGlyGlnLeuLeuPheSerValValIleAsnGlnLeuArgLeuSerAspAlaGly
 4423 CAGTATCTCTGCCAGGCTGGGGATGATTCCAATAGTAATAAGAAGAATGCTGACCTCCAA
 GlnTyrLeuCysGlnAlaGlyAspAspSerAsnSerAsnLysLysAsnAlaAspLeuGln
 4483 GTGCTAAAGCCCGAGCCCGAGCTGGTTTATGAAGACCTGAGGGGCTCAGTGACCTTCCAC
 ValLeuLysProGluProGluLeuValTyrGluAspLeuArgGlySerValThrPheHis
 4543 TGTGCCCTGGGCCCTGAGGTGGCAAACGTGGCCAAATTTCTGTGCCGACAGAGCAGTGGG
 CysAlaLeuGlyProGluValAlaAsnValAlaLysPheLeuCysArgGlnSerSerGly
 4603 GAAAACTGTGACGTGGTCGTCAACACCCTGGGGAAGAGGGCCCCAGCCTTTGAGGGCAGG
 GluAsnCysAspValValValAsnThrLeuGlyLysArgAlaProAlaPheGluGlyArg
 4663 ATCCTGCTCAACCCCCAGGACAAGGATGGCTCATTTCAGTGTGGTGATCACAGGCCTGAGG
 IleLeuLeuAsnProGlnAspLysAspGlySerPheSerValValIleThrGlyLeuArg
 4723 AAGGAGGATGCAGGGCGCTACCTGTGTGGAGCCCATTTCGGATGGTCAGCTGCAGGAAGGC
 LysGluAspAlaGlyArgTyrLeuCysGlyAlaHisSerAspGlyGlnLeuGlnGluGly
 4783 TCGCCTATCCAGGCCTGGCAACTCTTCGTCAATGAGGAGTCCACGATTCCCCGCAGCCCC
 SerProIleGlnAlaTrpGlnLeuPheValAsnGluGluSerThrIleProArgSerPro
 4843 ACTGTGGTGAAGGGGGTGGCAGGAAGCTCTGTGGCCGTGCTCTGCCCCCTACAACCGTAAG
 ThrValValLysGlyValAlaGlySerSerValAlaValLeuCysProTyrAsnArgLys
 4903 GAAAGCAAAGCATCAAGTACTGGTGTCTCTGGGAAGGGGCCAGAAATGGCCGCTGCCCC
 GluSerLysSerIleLysTyrTrpCysLeuTrpGluGlyAlaGlnAsnGlyArgCysPro
 4963 CTGCTGGTGGACAGCGAGGGGTGGGTTAAGGCCAGTACGAGGGCCGCCTCTCCCTGCTG
 LeuLeuValAspSerGluGlyTrpValLysAlaGlnTyrGluGlyArgLeuSerLeuLeu
 5023 GAGGAGCCAGGCAACGGCACCTTCACTGTCACTCAACCAGCTCACCAGCCGGGACGCC
 GluGluProGlyAsnGlyThrPheThrValIleLeuAsnGlnLeuThrSerArgAspAla
 5083 GGCTTCTACTGGTGTCTGACCAACGGCGATACTCTCTGGAGGACCACCGTGGAGATCAAG
 GlyPheTyrTrpCysLeuThrAsnGlyAspThrLeuTrpArgThrThrValGluIleLys
 5143 ATTATCGAAGGAGAACCACCTCAAGGTTCCCGGAATGTACGGCTGTGCTGGGAGAG
 IleIleGluGlyGluProAsnLeuLysValProGlyAsnValThrAlaValLeuGlyGlu
 5203 ACTCTCAAGGTCCCCTGTCACTTTCCATGCAAATTCCTCGTACGAGAAATACTGGTGC
 ThrLeuLysValProCysHisPheProCysLysPheSerSerTyrGluLysTyrTrpCys
 5263 AAGTGAATAACACGGGCTGCCAGGCCCTGCCAGCCAAGACGAAGCCCCAGCAAGGCC
 LysTrpAsnAsnThrGlyCysGlnAlaLeuProSerGlnAspGluGlyProSerLysAla
 5323 TTCGTGAACGTGACGAGAACAGCCGGCTTGTCTCCCTGACCCTGAACCTGGTGACCAGG
 PheValAsnCysAspGluAsnSerArgLeuValSerLeuThrLeuAsnLeuValThrArg
 5383 GCTGATGAGGGCTGGTACTGGTGTGGAGTGAAGCAGGGCCACTTCTATGGAGAGACTGCA
 AlaAspGluGlyTrpTyrTrpCysGlyValLysGlnGlyHisPheTyrGlyGluThrAla
 5443 GCCGTCTATGTGGCAGTTGAAGAGAGGAAGGCAGCGGGTCCCGCGATGTCAGCCTAGCG
 AlaValTyrValAlaValGluGluArgLysAlaAlaGlySerArgAspValSerLeuAla
 5503 AAGGCAGACGCTGCTCCTGATGAGAAGGTGCTAGACTCTGGTTTTTCGGGAGATTGAGAAC
 LysAlaAspAlaAlaProAspGluLysValLeuAspSerGlyPheArgGluIleGluAsn
 EcoRI
 5563 AAAGCCATTTCAGGATCCCAGGCTTTTTGTCAGAGTGAATTCCTGATCGTTCAAACATTTGGCAATAAAG
 LysAlaIleGlnAspProArgLeuPheAlaGlu
 5631 TTTCTTAAGATTGAATCCTGTTGCCGGTCTTGCGATGATTATCATATAATTTCTGTTGAATTACGTTAAGCATGTAATAA
 5711 TTAACATGTAATGCATGACGTTATTTATGAGATGGGTTTTTATGATTAGAGTCCCGCAATTATACATTTAATACGCGATA
 5791 GAAAACAAAATATAGCGCGCAAACTAGGATAAATTATCGCGCGCGGTGTCATCTATGTTACTAGATCGGGGATCCGTCGA
 ClaI
 5871 CGGTATCGATAAGGATCCCTGAAAGCGACGTTGGATGTTAACATCTACAAATTGCCTTTTCTTATCGACCATGTACGTAA
 5951 GCGCTTACGTTTTTGGTGGACCCTTGAGGAAACTGGTAGCTGTTGTGGGCTGTGGTCTCAAGATGGATCATTAATTTCC
 6031 ACCTTCACCTACGATGGGGGGCATCGCACCGGTGAGTAATATTGTACGGCTAAGAGCGAATTTGGCCTGTAGGATCCCTG

FIGURE 8 F - 3

6111 AAAGCGACGTTGGATGTTAACATCTACAAATTGCCTTTTCTTATCGACCATGTACGTAAGCGCTTACGTTTTTGGTGGAC
 6191 CCTTGAGGAACTGGTAGCTGTTGTGGGCCTGTGGTCTCAAGATGGATCATTAAATTTCCACCTTCACCTACGATGGGGGG
 6271 CATCGCACCGGTGAGTAATATTGTACGGCTAAGAGCGAATTTGGCCTGTAGGATCCCTGAAAGCGACGTTGGATGTTAAC
 6351 ATCTACAAATTGCCTTTTCTTATCGACCATGTACGTAAGCGCTTACGTTTTTGGTGGACCTTGAGGAACTGGTAGCTG
 6431 TTGTGGGCCTGTGGTCTCAAGATGGATCATTAAATTTCCACCTTCACCTACGATGGGGGGCATCGCACCGGTGAGTAATAT
 6511 TGTACGGCTAAGAGCGAATTTGGCCTGTAGGATCCGCGAGCTGGTCAATCCCATTGCTTTTGAAGCAGCTCAACATTGAT
 XhoI
 6591 CTCTTTCTCGAGGGAGATTTTTCAAATCAGTGCGCAAGACGTGACGTAAGTATCCGAGTCAGTTTTTATTTTTCTACTAA
 6671 TTTGGTCGTTTATTTTCGGCGTGTAGGACATGGCAACCGGGCCTGAATTTTCGCGGGTATTCTGTTTCTATTCCAACCTTTT
 6751 CTTGATCCGCAGCCATTAACGACTTTTGAATAGATACGCTGACACGCCAAGCCTCGCTAGTCAAAAGTGTAACCAACAAC
 6831 GCTTTACAGCAAGAACGGAATGCGCGTGACGCTCGCGGTGACGCCATTTTCGCCTTTTCAGAAATGGATAAATAGCCTTGC
 6911 TTCCTATTATATCTTCCCAAATTACCAATACATTACACTAGCATCTGAATTTTCATAACCAATCTCGATACACCAAATCGA
 XbaI
 6991 CTCTAGAGGATCTAACCATGGGATCTAAACCTTTTTTGTCTCTTCTTTTCATTGTCATTGCTT
 MetGlySerLysProPheLeuSerLeuLeuSerLeuSerLeuLeu
 SpeI
 7053 TTGTTTACATCTACTAGTTTGGCACAAGAAGATGAAAGGATTGTTCTTGTGACAACAAA
 LeuPheThrSerThrSerLeuAlaGlnGluAspGluArgIleValLeuValAspAsnLys
 7113 TGTAAGTGTGCCCCGATTACTTCCAGGATCATCCGTTCTTCCGAAGATCCTAATGAGGAC
 CysLysCysAlaArgIleThrSerArgIleIleArgSerSerGluAspProAsnGluAsp
 7173 ATTGTGGAGAGAAACATCCGAATTATTGTTCTCTGAACAACAGGGAGAATATCTCTGAT
 IleValGluArgAsnIleArgIleIleValProLeuAsnAsnArgGluAsnIleSerAsp
 7233 CCCACCTCACCATTGAGAACCAGATTTGTGTACCATTGTGTCTGACCTCTGTAAAAAATGT
 ProThrSerProLeuArgThrArgPheValTyrHisLeuSerAspLeuCysLysLysCys
 7293 GATCCTACAGAAGTGGAGCTGGATAATCAGATAGTTACTGCTACCCAGAGCAATATCTGT
 AspProThrGluValGluLeuAspAsnGlnIleValThrAlaThrGlnSerAsnIleCys
 7353 GATGAAGACAGTGCTACAGAGACCTGCTACACTTATGACAGAAACAAGTGCTACACAGCT
 AspGluAspSerAlaThrGluThrCysTyrThrTyrAspArgAsnLysCysTyrThrAla
 7413 GTGGTCCCACTCGTATATGGTGGTGAGACCAAAATGGTGGAAACAGCCTTAACCCAGAT
 ValValProLeuValTyrGlyGlyGluThrLysMetValGluThrAlaLeuThrProAsp
 SacI
 7473 GCCTGCTATCCTGACTGAGCTCGAATTTCCCGATCGTTCAAACATTTGGCAATAAAGTTTCTTAAGATTGAAT
 AlaCysTyrProAsp...
 7547 CCTGTTGCCGGTCTTGCGATGATTATCATATAATTTCTGTTGAATTACGTTAAGCATGTAATAATTAACATGTAATGCAT
 7627 GACGTTATTTATGAGATGGGTTTTTATGATTAGAGTCCCGCAATTATACATTTAATACGCGATAGAAAACAAAATATAGC
 7707 GCGCAAACTAGGATAAAATTATCGCGCGCGGTGTCATCTATGTTACTAGATCGGGAATTAATTCAGATCGGCTGAGTGGCT
 7787 CCTTCAATCGTTGCGGTTCTGTCTAGTTCCAAACGTAAACGGCTTGTCCCGCGTCATCGGCGGGGGTCATAACGTGACTC
 7867 CCTTAATTCTCCGCTCATGATCAGATTGTCGTTTCCCGCCTTCAGTTTAACTATCAGTGTTTGACAGGATATATTGGCG
 7947 GGTAACCTAAGAGAAAAGAGCGTTTATTAGAATAATCGGATATTTAAAGGGCGTGAAAAGGTTTATCCGTTCTGTCAT
 BglII Sfo I
 8027 TTGTATGTGCATGCCAACCACAGGTTCCCCAGATCTGGCGCCGGCCAG

FIGURE 8 F - 4

FIGURE 9

SEQUENCE LISTING

<110> PLANET BIOTECHNOLOGY, INCORPORATED
 Larrick, James William
 Wycoff, Keith Lynn

<120> NOVEL IMMUNOADHESION FOR THE PREVENTION
 OF RHINOVIRUS INFECTION

<130> 030905.0003.WO

<140> To be assigned

<141> 2001-04-27

<150> 60/200,298

<151> 2000-04-27

<160> 8

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1596

<212> DNA

<213> Homo Sapien

<220>

<221> CDS

<222> (1)...(1596)

<223> ICAM-1

<400> 1

atg gct ccc agc agc ccc cgg ccc gcg ctg ccc gca ctc ctg gtc ctg	48
Met Ala Pro Ser Ser Pro Arg Pro Ala Leu Pro Ala Leu Leu Val Leu	
1 5 10 15	
ctc ggg gct ctg ttc cca gga cct ggc aat gcc cag aca tct gtg tcc	96
Leu Gly Ala Leu Phe Pro Gly Pro Gly Asn Ala Gln Thr Ser Val Ser	
20 25 30	
ccc tca aaa gtc atc ctg ccc cgg gga ggc tcc gtg ctg gtg aca tgc	144
Pro Ser Lys Val Ile Leu Pro Arg Gly Gly Ser Val Leu Val Thr Cys	
35 40 45	
agc acc tcc tgt gac cag ccc aag ttg ttg ggc ata gag acc ccg ttg	192
Ser Thr Ser Cys Asp Gln Pro Lys Leu Leu Gly Ile Glu Thr Pro Leu	
50 55 60	
cct aaa aag gag ttg ctc ctg cct ggg aac aac cgg aag gtg tat gaa	240
Pro Lys Lys Glu Leu Leu Leu Pro Gly Asn Asn Arg Lys Val Tyr Glu	
65 70 75 80	
ctg agc aat gtg caa gaa gat agc caa cca atg tgc tat tca aac tgc	288
Leu Ser Asn Val Gln Glu Asp Ser Gln Pro Met Cys Tyr Ser Asn Cys	
85 90 95	

FIGURE 9 - 1

cct gat ggg cag tca aca gct aaa acc ttc ctc acc gtg tac tgg act	336
Pro Asp Gly Gln Ser Thr Ala Lys Thr Phe Leu Thr Val Tyr Trp Thr	
100 105 110	
cca gaa cgg gtg gaa ctg gca ccc ctc ccc tct tgg cag cca gtg ggc	384
Pro Glu Arg Val Glu Leu Ala Pro Leu Pro Ser Trp Gln Pro Val Gly	
115 120 125	
aag aac ctt acc cta cgc tgc cag gtg gag ggt ggg gca ccc cgg gcc	432
Lys Asn Leu Thr Leu Arg Cys Gln Val Glu Gly Gly Ala Pro Arg Ala	
130 135 140	
aac ctc acc gtg gtg ctg ctc cgt ggg gag aag gag ctg aaa cgg gag	480
Asn Leu Thr Val Val Leu Leu Arg Gly Glu Lys Glu Leu Lys Arg Glu	
145 150 155 160	
cca gct gtg ggg gag ccc gct gag gtc acg acc acg gtg ctg gtg agg	528
Pro Ala Val Gly Glu Pro Ala Glu Val Thr Thr Thr Val Leu Val Arg	
165 170 175	
aga gat cac cat gga gcc aat ttc tcg tgc cgc act gaa ctg gac ctg	576
Arg Asp His His Gly Ala Asn Phe Ser Cys Arg Thr Glu Leu Asp Leu	
180 185 190	
cgg ccc caa ggg ctg gag ctg ttt gag aac acc tcg gcc ccc tac cag	624
Arg Pro Gln Gly Leu Glu Leu Phe Glu Asn Thr Ser Ala Pro Tyr Gln	
195 200 205	
ctc cag acc ttt gtc ctg cca gcg act ccc cca caa ctt gtc agc ccc	672
Leu Gln Thr Phe Val Leu Pro Ala Thr Pro Pro Gln Leu Val Ser Pro	
210 215 220	
cgg gtc cta gag gtg gac acg cag ggg acc gtg gtc tgt tcc ctg gac	720
Arg Val Leu Glu Val Asp Thr Gln Gly Thr Val Val Cys Ser Leu Asp	
225 230 235 240	
ggg ctg ttc cca gtc tcg gag gcc cag gtc cac ctg gca ctg ggg gac	768
Gly Leu Phe Pro Val Ser Glu Ala Gln Val His Leu Ala Leu Gly Asp	
245 250 255	
cag agg ttg aac ccc aca gtc acc tat ggc aac gac tcc ttc tcg gcc	816
Gln Arg Leu Asn Pro Thr Val Thr Tyr Gly Asn Asp Ser Phe Ser Ala	
260 265 270	
aag gcc tca gtc agt gtg acc gca gag gac gag ggc acc cag cgg ctg	864
Lys Ala Ser Val Ser Val Thr Ala Glu Asp Glu Gly Thr Gln Arg Leu	
275 280 285	
acg tgt gca gta ata ctg ggg aac cag agc cag gag aca ctg cag aca	912
Thr Cys Ala Val Ile Leu Gly Asn Gln Ser Gln Glu Thr Leu Gln Thr	
290 295 300	
gtg acc atc tac agc ttt ccg gcg ccc aac gtg att ctg acg aag cca	960
Val Thr Ile Tyr Ser Phe Pro Ala Pro Asn Val Ile Leu Thr Lys Pro	
305 310 315 320	

FIGURE 9-2

gag gtc tca gaa ggg acc gag gtg aca gtg aag tgt gag gcc cac cct	1008
Glu Val Ser Glu Gly Thr Glu Val Thr Val Lys Cys Glu Ala His Pro	
325 330 335	
aga gcc aag gtg acg ctg aat ggg gtt cca gcc cag cca ctg ggc ccg	1056
Arg Ala Lys Val Thr Leu Asn Gly Val Pro Ala Gln Pro Leu Gly Pro	
340 345 350	
agg gcc cag ctc ctg ctg aag gcc acc cca gag gac aac ggg cgc agc	1104
Arg Ala Gln Leu Leu Leu Lys Ala Thr Pro Glu Asp Asn Gly Arg Ser	
355 360 365	
ttc tcc tgc tct gca acc ctg gag gtg gcc ggc cag ctt ata cac aag	1152
Phe Ser Cys Ser Ala Thr Leu Glu Val Ala Gly Gln Leu Ile His Lys	
370 375 380	
aac cag acc cgg gag ctt cgt gtc ctg tat ggc ccc cga ctg gac gag	1200
Asn Gln Thr Arg Glu Leu Arg Val Leu Tyr Gly Pro Arg Leu Asp Glu	
385 390 395 400	
agg gat tgt ccg gga aac tgg acg tgg cca gaa aat tcc cag cag act	1248
Arg Asp Cys Pro Gly Asn Trp Thr Trp Pro Glu Asn Ser Gln Gln Thr	
405 410 415	
cca atg tgc cag gct tgg ggg aac cca ttg ccc gag ctc aag tgt cta	1296
Pro Met Cys Gln Ala Trp Gly Asn Pro Leu Pro Glu Leu Lys Cys Leu	
420 425 430	
aag gat ggc act ttc cca ctg ccc atc ggg gaa tca gtg act gtc act	1344
Lys Asp Gly Thr Phe Pro Leu Pro Ile Gly Glu Ser Val Thr Val Thr	
435 440 445	
cga gat ctt gag ggc acc tac ctc tgt cgg gcc agg agc act caa ggg	1392
Arg Asp Leu Glu Gly Thr Tyr Leu Cys Arg Ala Arg Ser Thr Gln Gly	
450 455 460	
gag gtc acc cgc aag gtg acc gtg aat gtg ctc tcc ccc cgg tat gag	1440
Glu Val Thr Arg Lys Val Thr Val Asn Val Leu Ser Pro Arg Tyr Glu	
465 470 475 480	
att gtc atc atc act gtg gta gca gcc gca gtc ata atg ggc act gca	1488
Ile Val Ile Ile Thr Val Val Ala Ala Val Ile Met Gly Thr Ala	
485 490 495	
ggc ctc agc acg tac ctc tat aac cgc cag cgg aag atc aag aaa tac	1536
Gly Leu Ser Thr Tyr Leu Tyr Asn Arg Gln Arg Lys Ile Lys Lys Tyr	
500 505 510	
aga cta caa cag gcc caa aaa ggg acc ccc atg aaa ccg aac aca caa	1584
Arg Leu Gln Gln Ala Gln Lys Gly Thr Pro Met Lys Pro Asn Thr Gln	
515 520 525	
gcc acg cct ccc	1596
Ala Thr Pro Pro	
530	

FIGURE 9 – 3

<210> 2
 <211> 532
 <212> PRT
 <213> Homo Sapien

<400> 2
 Met Ala Pro Ser Ser Pro Arg Pro Ala Leu Pro Ala Leu Leu Val Leu
 1 5 10 15
 Leu Gly Ala Leu Phe Pro Gly Pro Gly Asn Ala Gln Thr Ser Val Ser
 20 25 30
 Pro Ser Lys Val Ile Leu Pro Arg Gly Gly Ser Val Leu Val Thr Cys
 35 40 45
 Ser Thr Ser Cys Asp Gln Pro Lys Leu Leu Gly Ile Glu Thr Pro Leu
 50 55 60
 Pro Lys Lys Glu Leu Leu Leu Pro Gly Asn Asn Arg Lys Val Tyr Glu
 65 70 75 80
 Leu Ser Asn Val Gln Glu Asp Ser Gln Pro Met Cys Tyr Ser Asn Cys
 85 90 95
 Pro Asp Gly Gln Ser Thr Ala Lys Thr Phe Leu Thr Val Tyr Trp Thr
 100 105 110
 Pro Glu Arg Val Glu Leu Ala Pro Leu Pro Ser Trp Gln Pro Val Gly
 115 120 125
 Lys Asn Leu Thr Leu Arg Cys Gln Val Glu Gly Gly Ala Pro Arg Ala
 130 135 140
 Asn Leu Thr Val Val Leu Leu Arg Gly Glu Lys Glu Leu Lys Arg Glu
 145 150 155 160
 Pro Ala Val Gly Glu Pro Ala Glu Val Thr Thr Thr Val Leu Val Arg
 165 170 175
 Arg Asp His His Gly Ala Asn Phe Ser Cys Arg Thr Glu Leu Asp Leu
 180 185 190
 Arg Pro Gln Gly Leu Glu Leu Phe Glu Asn Thr Ser Ala Pro Tyr Gln
 195 200 205
 Leu Gln Thr Phe Val Leu Pro Ala Thr Pro Pro Gln Leu Val Ser Pro
 210 215 220
 Arg Val Leu Glu Val Asp Thr Gln Gly Thr Val Val Cys Ser Leu Asp
 225 230 235 240
 Gly Leu Phe Pro Val Ser Glu Ala Gln Val His Leu Ala Leu Gly Asp
 245 250 255
 Gln Arg Leu Asn Pro Thr Val Thr Tyr Gly Asn Asp Ser Phe Ser Ala
 260 265 270
 Lys Ala Ser Val Ser Val Thr Ala Glu Asp Glu Gly Thr Gln Arg Leu
 275 280 285
 Thr Cys Ala Val Ile Leu Gly Asn Gln Ser Gln Glu Thr Leu Gln Thr
 290 295 300
 Val Thr Ile Tyr Ser Phe Pro Ala Pro Asn Val Ile Leu Thr Lys Pro
 305 310 315 320
 Glu Val Ser Glu Gly Thr Glu Val Thr Val Lys Cys Glu Ala His Pro
 325 330 335
 Arg Ala Lys Val Thr Leu Asn Gly Val Pro Ala Gln Pro Leu Gly Pro
 340 345 350
 Arg Ala Gln Leu Leu Leu Lys Ala Thr Pro Glu Asp Asn Gly Arg Ser
 355 360 365
 Phe Ser Cys Ser Ala Thr Leu Glu Val Ala Gly Gln Leu Ile His Lys
 370 375 380
 Asn Gln Thr Arg Glu Leu Arg Val Leu Tyr Gly Pro Arg Leu Asp Glu
 385 390 395 400
 Arg Asp Cys Pro Gly Asn Trp Thr Trp Pro Glu Asn Ser Gln Gln Thr

FIGURE 9 - 4

				405					410					415					
Pro	Met	Cys	Gln	Ala	Trp	Gly	Asn	Pro	Leu	Pro	Glu	Leu	Lys	Cys	Leu				
			420					425					430						
Lys	Asp	Gly	Thr	Phe	Pro	Leu	Pro	Ile	Gly	Glu	Ser	Val	Thr	Val	Thr				
		435					440					445							
Arg	Asp	Leu	Glu	Gly	Thr	Tyr	Leu	Cys	Arg	Ala	Arg	Ser	Thr	Gln	Gly				
	450					455					460								
Glu	Val	Thr	Arg	Lys	Val	Thr	Val	Asn	Val	Leu	Ser	Pro	Arg	Tyr	Glu				
465					470					475					480				
Ile	Val	Ile	Ile	Thr	Val	Val	Ala	Ala	Ala	Val	Ile	Met	Gly	Thr	Ala				
				485					490					495					
Gly	Leu	Ser	Thr	Tyr	Leu	Tyr	Asn	Arg	Gln	Arg	Lys	Ile	Lys	Lys	Tyr				
			500					505					510						
Arg	Leu	Gln	Gln	Ala	Gln	Lys	Gly	Thr	Pro	Met	Lys	Pro	Asn	Thr	Gln				
		515					520						525						
Ala	Thr	Pro	Pro																
		530																	

<210> 3

<211> 3003

<212> DNA

<213> Homo Sapien

<400> 3

gctataagga	tcacgcgccc	cagtcgacgc	tgagctcctc	tgctactcag	agttgcaacc	60
tcagcctcgc	tatggctccc	agcagccccc	ggcccgcgct	gcccgcactc	ctggtcctgc	120
tcggggctct	gttcccagga	cctggcaatg	cccagacatc	tgtgtccccc	tcaaaagtca	180
tcctgccccg	gggaggctcc	gtgctggtga	catgcagcac	ctcctgtgac	cagcccaagt	240
tgttgggcat	agagaccccg	ttgcctaaaa	aggagttgct	cctgcctggg	aacaaccgga	300
aggtgtatga	actgagcaat	gtgcaagaag	atagccaacc	aatgtgctat	tcaaactgcc	360
ctgatgggca	gtcaacagct	aaaaccttcc	tcaccgtgta	ctggactcca	gaacgggtgg	420
aactggcacc	cctcccctct	tggcagccag	tgggcaagaa	ccttacccta	cgtgcccagg	480
tggaggggtg	ggcaccgccg	gccaacctca	ccgtggtgct	gctccgtggg	gagaaggagc	540
tgaaacggga	gccagctgtg	ggggagcccg	ctgaggtcac	gaccacgggtg	ctggtgagga	600
gagatcacca	tggagccaat	ttctcgtgcc	gcactgaact	ggacctgcgg	ccccaaagggc	660
tggagctggt	tgagaacacc	tcggccccct	accagctcca	gacctttgtc	ctgccagcga	720
ctccccaca	acttgtcagc	ccccgggtcc	tagaggtgga	cacgcagggg	accgtggtct	780
gttccctgga	cgggctgttc	ccagtctcgg	aggcccagg	ccacctggca	ctggggggacc	840
agaggttgaa	ccccacagtc	acctatggca	acgactcctt	ctcggccaag	gcctcagtca	900
gtgtgaccgc	agaggacgag	ggcaccacgc	ggctgacgtg	tgcagtaata	ctgggggaacc	960
agagccagga	gacactgcag	acagtgacca	tctacagctt	tccggcgccc	aacgtgattc	1020
tgacgaagcc	agaggtctca	gaagggaccg	aggtgacagt	gaagtgtgag	gcccacccta	1080
gagccaaggt	gacgctgaat	gggggttccag	cccagccact	gggcccagag	gcccagctcc	1140
tgctgaaggc	caccccagag	gacaacgggc	gcagcttctc	ctgctctgca	accctggagg	1200
tggccggcca	gcttatacac	aagaaccaga	cccgggagct	tcgtgtcctg	tatggccccc	1260
gactggacga	gagggattgt	ccgggaaact	ggacgtggcc	agaaaattcc	cagcagactc	1320
caatgtgcca	ggcttggggg	aaccatttgc	ccgagctcaa	gtgtctaaag	gatggcactt	1380
tcccactgcc	catcggggaa	tcagtgactg	tcactcgaga	tcttgagggc	acctacctct	1440
gtcggggccag	gagcactcaa	ggggaggtca	cccgcagggt	gaccgtgaat	gtgctctccc	1500
cccggtatga	gattgtcatc	atcactgtgg	tagcagccgc	agtcataatg	ggcactgcag	1560
gcctcagcac	gtacctctat	aaccgccagc	ggaagatcaa	gaaatacaga	ctacaacagg	1620
cccaaaaagg	gacccccatg	aaaccgaaca	cacaagccac	gcctccctga	acctatcccg	1680
ggacagggcc	tcttcctcgg	ccttcccata	ttggtggcag	tggtgccaca	ctgaacagag	1740
tgggaagacat	atgccatgca	gctacaccta	ccggccctgg	gacgccggag	gacagggcat	1800
tgtcctcagt	cagatacaac	agcatttggg	gccatggtac	ctgcacacct	aaaacactag	1860
gccacgcac	tgatctgtag	tcacatgact	aagccaagag	gaaggagcaa	gactcaagac	1920
atgattgatg	gatgttaaag	tctagcctga	tgagagggga	agtgggtggg	gagacatagc	1980

FIGURE 9 - 5

```

cccaccatga ggacatacaa ctgggaaata ctgaaacttg ctgcctattg ggtatgctga 2040
ggccccacag acttacagaa gaagtggccc tccatagaca tgtgtagcat caaaacacaa 2100
aggccccacac ttcctgacgg atgccagctt gggcactgct gtctactgac cccaaccctt 2160
gatgatatgt atttattcat ttgttatttt accagctatt tattgagtgt cttttatgta 2220
ggctaaatga acataggtct ctggcctcac ggagctccca gtccatgtca cattcaaggt 2280
caccaggtac agttgtacag gttgtacact gcaggagagt gcctggcaaa aagatcaaatt 2340
ggggctggga cttctcattg gccaacctgc ctttcccag aaggagtgat ttttctatcg 2400
gcacaaaagc actatatgga ctggtaatgg ttcacagggt cagagattac ccagtgaggc 2460
cttattcctc ccttcccccc aaaactgaca cctttgttag ccacctcccc acccacatac 2520
atttctgcca gtgttcacaa tgacactcag cggtcatgtc tggacatgag tgcccaggga 2580
atatgcccac gctatgcctt gtcctcttgt cctgtttgca tttcactggg agcttgcact 2640
attgcagctc cagtttcttg cagtgatcag ggtcctgcaa gcagtgggga agggggccaa 2700
ggtattggag gactccctcc cagctttgga agcctcatcc gcgtgtgtgt gtgtgtgtgt 2760
atgtgtagac aagctctcgc tctgtcacc aggctggagt gcagtgggtc aatcatgggt 2820
cactgcagtc ttgacctttt gggctcaagt gatcctccca cctcagcctc ctgagtagct 2880
gggaccatag gctcacaaca ccacacctgg caaatttgat tttttttttt tttttcagag 2940
acgggggtctc gcaacattgc ccagacttcc tttgtgttag ttaataaagc tttctcaact 3000
gcc 3003

```

<210> 4
 <211> 6
 <212> PRT
 <213> Homo Sapien

<400> 4
 Ser Glu Lys Asp Glu Leu
 1 5

<210> 5
 <211> 7
 <212> PRT
 <213> Homo Sapien

<400> 5
 Arg Ser Glu Lys Asp Glu Leu
 1 5

<210> 6
 <211> 52
 <212> DNA
 <213> Homo Sapien

<400> 6
 tctgttccca ggaactagtt tggcacagac atctgtgtcc ccctcaaaag to 52

<210> 7
 <211> 38
 <212> DNA
 <213> Homo Sapien

<400> 7
 cataccgggg actagtcaca ttcacgggtca cctcgcg 38

<210> 8
 <211> 799
 <212> PRT
 <213> Homo Sapien

FIGURE 9 - 6

<220>
 <221> CDS
 <222> (1)...(448)
 <223> ICAM-1 Extracellular Domains

<220>
 <221> CDS
 <222> (453)...(799)
 <223> Human IgA2m(2)

<400> 8
 Gln Thr Ser Val Ser Pro Ser Lys Val Ile Leu Pro Arg Gly Gly Ser
 1 5 10 15
 Val Leu Val Thr Cys Ser Thr Ser Cys Asp Gln Pro Lys Leu Leu Gly
 20 25 30
 Ile Glu Thr Pro Leu Pro Lys Lys Glu Leu Leu Leu Pro Gly Asn Asn
 35 40 45
 Arg Lys Val Tyr Glu Leu Ser Asn Val Gln Glu Asp Ser Gln Pro Met
 50 55 60
 Cys Tyr Ser Asn Cys Pro Asp Gly Gln Ser Thr Ala Lys Thr Phe Leu
 65 70 75 80
 Thr Val Tyr Trp Thr Pro Glu Arg Val Glu Leu Ala Pro Leu Pro Ser
 85 90 95
 Trp Gln Pro Val Gly Lys Asn Leu Thr Leu Arg Cys Gln Val Glu Gly
 100 105 110
 Gly Ala Pro Arg Ala Asn Leu Thr Val Val Leu Leu Arg Gly Glu Lys
 115 120 125
 Glu Leu Lys Arg Glu Pro Ala Val Gly Glu Pro Ala Glu Val Thr Thr
 130 135 140
 Thr Val Leu Val Arg Arg Asp His His Gly Ala Asn Phe Ser Cys Arg
 145 150 155 160
 Thr Glu Leu Asp Leu Arg Pro Gln Gly Leu Glu Leu Phe Glu Asn Thr
 165 170 175
 Ser Ala Pro Tyr Gln Leu Gln Thr Phe Val Leu Pro Ala Thr Pro Pro
 180 185 190
 Gln Leu Val Ser Pro Arg Val Leu Glu Val Asp Thr Gln Gly Thr Val
 195 200 205
 Val Cys Ser Leu Asp Gly Leu Phe Pro Val Ser Glu Ala Gln Val His
 210 215 220
 Leu Ala Leu Gly Asp Gln Arg Leu Asn Pro Thr Val Thr Tyr Gly Asn
 225 230 235 240
 Asp Ser Phe Ser Ala Lys Ala Ser Val Ser Val Thr Ala Glu Asp Glu
 245 250 255
 Gly Thr Gln Arg Leu Thr Cys Ala Val Ile Leu Gly Asn Gln Ser Gln
 260 265 270
 Glu Thr Leu Gln Thr Val Thr Ile Tyr Ser Phe Pro Ala Pro Asn Val
 275 280 285
 Ile Leu Thr Lys Pro Glu Val Ser Glu Gly Thr Glu Val Thr Val Lys
 290 295 300
 Cys Glu Ala His Pro Arg Ala Lys Val Thr Leu Asn Gly Val Pro Ala
 305 310 315 320
 Gln Pro Leu Gly Pro Arg Ala Gln Leu Leu Leu Lys Ala Thr Pro Glu
 325 330 335
 Asp Asn Gly Arg Ser Phe Ser Cys Ser Ala Thr Leu Glu Val Ala Gly
 340 345 350
 Gln Leu Ile His Lys Asn Gln Thr Arg Glu Leu Arg Val Leu Tyr Gly
 355 360 365

FIGURE 9 - 7

Pro	Arg	Leu	Asp	Glu	Arg	Asp	Cys	Pro	Gly	Asn	Trp	Thr	Trp	Pro	Glu
370						375					380				
Asn	Ser	Gln	Gln	Thr	Pro	Met	Cys	Gln	Ala	Trp	Gly	Asn	Pro	Leu	Pro
385					390					395					400
Glu	Leu	Lys	Cys	Leu	Lys	Asp	Gly	Thr	Phe	Pro	Leu	Pro	Ile	Gly	Glu
				405					410					415	
Ser	Val	Thr	Val	Thr	Arg	Asp	Leu	Glu	Gly	Thr	Tyr	Leu	Cys	Arg	Ala
			420					425					430		
Arg	Ser	Thr	Gln	Gly	Glu	Val	Thr	Arg	Glu	Val	Thr	Val	Asn	Val	Thr
		435					440					445			
Ser	Gly	Ser	Ser	Ala	Ser	Pro	Thr	Ser	Pro	Lys	Val	Phe	Pro	Leu	Ser
	450					455					460				
Leu	Asp	Ser	Thr	Pro	Gln	Asp	Gly	Asn	Val	Val	Val	Ala	Cys	Leu	Val
465					470				475						480
Gln	Gly	Phe	Phe	Pro	Gln	Glu	Pro	Leu	Ser	Val	Thr	Trp	Ser	Glu	Ser
				485					490					495	
Gly	Gln	Asn	Val	Thr	Ala	Arg	Asn	Phe	Pro	Pro	Ser	Gln	Asp	Ala	Ser
			500					505					510		
Gly	Asp	Leu	Tyr	Thr	Thr	Ser	Ser	Gln	Leu	Thr	Leu	Pro	Ala	Thr	Gln
	515						520					525			
Cys	Pro	Asp	Gly	Lys	Ser	Val	Thr	Cys	His	Val	Lys	His	Tyr	Thr	Asn
	530					535					540				
Ser	Ser	Gln	Asp	Val	Thr	Val	Pro	Cys	Arg	Val	Pro	Pro	Pro	Pro	Pro
545					550					555					560
Cys	Cys	His	Pro	Arg	Leu	Ser	Leu	His	Arg	Pro	Ala	Leu	Glu	Asp	Leu
				565					570					575	
Leu	Leu	Gly	Ser	Glu	Ala	Asn	Leu	Thr	Cys	Thr	Leu	Thr	Gly	Leu	Arg
			580					585					590		
Asp	Ala	Ser	Gly	Ala	Thr	Phe	Thr	Trp	Thr	Pro	Ser	Ser	Gly	Lys	Ser
	595						600					605			
Ala	Val	Gln	Gly	Pro	Pro	Glu	Arg	Asp	Leu	Cys	Gly	Cys	Tyr	Ser	Val
	610					615					620				
Ser	Arg	Val	Leu	Pro	Gly	Cys	Ala	Gln	Pro	Trp	Asn	His	Gly	Glu	Thr
625					630					635					640
Phe	Thr	Cys	Thr	Ala	Ala	His	Pro	Glu	Leu	Lys	Thr	Pro	Leu	Thr	Ala
				645					650					655	
Asn	Ile	Thr	Lys	Ser	Gly	Asn	Thr	Phe	Arg	Pro	Glu	Val	His	Leu	Leu
			660					665					670		
Pro	Pro	Pro	Ser	Glu	Glu	Leu	Ala	Leu	Asn	Glu	Leu	Val	Thr	Leu	Thr
	675						680					685			
Cys	Leu	Ala	Arg	Gly	Phe	Ser	Pro	Lys	Asp	Val	Leu	Val	Arg	Trp	Leu
	690					695					700				
Gln	Gly	Ser	Gln	Glu	Leu	Pro	Arg	Glu	Lys	Tyr	Leu	Thr	Trp	Ala	Ser
705					710					715					720
Arg	Gln	Glu	Pro	Ser	Gln	Gly	Thr	Thr	Thr	Tyr	Ala	Val	Thr	Ser	Ile
				725					730					735	
Leu	Arg	Val	Ala	Ala	Glu	Asp	Trp	Lys	Lys	Gly	Glu	Thr	Phe	Ser	Cys
			740					745					750		
Met	Val	Gly	His	Glu	Ala	Leu	Pro	Leu	Ala	Phe	Thr	Gln	Lys	Thr	Ile
	755						760					765			
Asp	Arg	Leu	Ala	Gly	Lys	Pro	Thr	His	Ile	Asn	Val	Ser	Val	Val	Met
	770					775					780				
Ala	Glu	Ala	Asp	Gly	Thr	Cys	Tyr	Arg	Ser	Glu	Lys	Asp	Glu	Leu	
785					790						795				

FIGURE 9 – 8

