



US00PP13648P29

(12) **United States Plant Patent**
Uchneat

(10) **Patent No.:** **US PP13,648 P2**
(45) **Date of Patent:** **Mar. 11, 2003**

(54) **DOUBLE IMPATIENS PLANT NAMED**
'BALOLESAL'

PP11,549 P * 10/2000 Cosner et al. Plt./317
PP11,859 P2 * 5/2001 Knosher Plt./317

(75) Inventor: **Michael S. Uchneat**, St. Charles, IL
(US)

OTHER PUBLICATIONS

(73) Assignee: **Ball FloraPlant, a division of Ball Horticultural Company**, West Chicago, IL (US)

UPOV ROM GTITM computer database, GTI JOUVE retrieval software 2002/02, citations for 'Balolesal'.*

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

* cited by examiner

Primary Examiner—Bruce R. Campbell
Assistant Examiner—W C Haas

(21) Appl. No.: **10/115,052**

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(22) Filed: **Apr. 1, 2002**

(57) **ABSTRACT**

(51) **Int. Cl.**⁷ **A01H 5/00**

A new and distinct cultivar of Double Impatiens plant named 'Balolesal', characterized by its compact, upright, outwardly spreading and mounded plant habit; freely branching growth habit; dark green-colored leaves; freely flowering habit; and light red-colored flowers that are fully double and positioned above the foliage.

(52) **U.S. Cl.** **Plt./317**

(58) **Field of Search** Plt./317

(56) **References Cited**

U.S. PATENT DOCUMENTS

1 Drawing Sheet

PP9,691 P * 11/1996 Leue Plt./317

1

2

BOTANICAL CLASSIFICATION/CULTIVAR DESIGNATION

Impatiens walleriana cultivar Balolesal.

SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Balolesal'. These characteristics in combination distinguish 'Balolesal' as a new and distinct cultivar:

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct cultivar of Double Impatiens plant, botanically known as *Impatiens walleriana*, and hereinafter referred to by the name 'Balolesal'.

1. Compact, upright, outwardly spreading and mounded plant habit.
2. Freely branching growth habit.
3. Dark green-colored leaves.
4. Freely flowering habit.
5. Light red-colored flowers that are fully double and positioned above the foliage.

The new Impatiens is a product of a planned breeding program conducted by the Inventor in Elburn, Ill. The objective of the breeding program was to develop new compact Double Impatiens cultivars with fully double flowers, freely branching growth habit, and attractive flower and foliage coloration.

Plants of the new Impatiens differ primarily from plants of the parent selections in flower color.

The new Impatiens originated from a cross-pollination made by the Inventor in June, 1999, of a proprietary *Impatiens walleriana* selection identified as 3039C-1, not patented, as the female, or seed parent, with a proprietary *Impatiens walleriana* selection identified as 3111-1-9, not patented, as the male, or pollen parent. The cultivar Balolesal was discovered and selected by the Inventor as a flowering plant within the progeny of the stated cross-pollination in a controlled environment in Elburn, Ill.

The new Impatiens can be compared to the *Impatiens walleriana* cultivar Salmon Sunrise, disclosed in U.S. Plant Pat. No. 9,691. However, in side-by-side comparisons conducted in West Chicago, Ill. plants of the new Impatiens differed from plants of the cultivar Salmon Sunrise in the following characteristics:

1. Plants of the new Impatiens had longer lateral branches with longer internodes than plants of the cultivar Salmon Sunrise.
2. Plants of the new Impatiens had larger and darker green-colored leaves than plants of the cultivar Salmon Sunrise.
3. Plants of the new Impatiens were more freely flowering than plants of the cultivar Salmon Sunrise.
4. Plants of the new Impatiens had larger flowers with more petals per flower than plants of the cultivar Salmon Sunrise.

Asexual reproduction of the new cultivar by terminal cuttings taken at Elburn, Ill. since October, 1999, has shown that the unique features of this new Impatiens are stable and reproduced true to type in successive generations of asexual reproduction.

5. Flower color of plants of the new *Impatiens* was more orange than flower color of plants of the cultivar Salmon Sunrise.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Impatiens*.

The photograph comprises a side perspective view of multiple flowering plants of 'Balolesal' in a container.

DETAILED BOTANICAL DESCRIPTION

The cultivar Balolesal has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity without, however, any variance in genotype. The aforementioned photograph and following observations and measurements describe plants grown in West Chicago, Ill. under commercial practice in a polycarbonate-covered greenhouse with day temperatures ranging from 20 to 24°C., night temperatures ranging from 10 to 17° C. and light levels ranging from 4,000 to 6,000 footcandles. Rooted young plants were planted in containers and had been growing for about five to six weeks when the photographs and the description were taken.

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Impatiens walleriana* cultivar Balolesal.

Parentage:

Female parent.—Proprietary *Impatiens walleriana* selection identified as 3039C-1, not patented.

Male parent.—Proprietary *Impatiens walleriana* selection identified as 3111-1-9, not patented.

Propagation:

Type cutting.—Terminal cuttings.

Time to initiate roots.—About 7 days at 18° C.

Time to produce a rooted young plant.—About 21 days at 18° C.

Root description.—Fibrous and fine.

Rooting habit.—Freely branching.

Plant description:

General appearance.—Compact, upright, outwardly spreading and mounded plant habit. Appropriate for 10-cm and larger containers.

Growth and branching habit.—Vigorous and freely branching growth habit with about three basal branches and lateral branches potentially forming at every node; dense and bushy growth. Pinching, that is, removal of the terminal apices, is not required, but will enhance branching.

Plant height.—About 11.5 cm.

Plant diameter or spread.—About 25 cm.

Lateral branches.—Length: About 13.4 cm. Diameter: About 8 mm. Internode length: About 3.2 cm. Texture: Smooth, glabrous. Color: 146C; random dark red purple, 59A, spots and streaks at nodes.

Foliage description.—Arrangement: Opposite, simple. Length: About 4.6 cm. Width: About 3 cm. Shape: Ovate. Apex: Acuminate. Base: Attenuate. Margin: Serrate to crenate with ciliation. Texture, upper and lower surfaces: Smooth, glabrous. Venation pattern: Pinnate, arcuate. Color: Young and fully expanded foliage, upper surface: 139A. Young and fully expanded foliage, lower surface: 138B. Venation, upper and lower surfaces: 144B. Petiole. Length: About 8 mm. Diameter: About 2 mm. Texture, upper and lower surfaces: Smooth, glabrous. Color: 146D.

Flower description:

Flower type and habit.—Numerous and consistently double flowers. Flower buds open similar to a rose in fullness; flowers rounded in shape. Flowers arise from leaf axils. Freely flowering, usually about 16 flowers and flower buds per lateral branch. Flowers positioned above the foliage and typically face upright or outward. Flowers last about five to seven days under greenhouse conditions. Flowers not persistent. Flowers not fragrant.

Flowering season.—Year-round under greenhouse conditions. In the garden, flowering is continuous from spring until fall.

Flower size.—Diameter: About 4.3 cm. Depth: About 2.3 cm.

Flower buds (at stage of showing color).—Length: About 1.7 cm. Diameter: About 1.3 cm. Shape: Ovoid. Color: 48C.

Petals.—Quantity/arrangement: About 37 per flower; imbricate. Length, largest petals: About 2 cm. Width, largest petals: About 1.8 cm. Shape: Obovate. Apex: Emarginate. Base: Truncate to attenuate. Margin: Entire. Texture, upper and lower surfaces: Velvety, smooth. Color: When opening and fully opened, upper surface: 39A; becoming closer to 52A with subsequent development. When opening and fully opened, lower surface: 48C.

Sepals.—Quantity/arrangement: Five in a single whorl; one modified into an elongated spur. Calyx length: About 1 cm. Calyx diameter: About 1 cm. Shape: Elliptic. Apex: Acute. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: 144C. Spur length: About 2.9 cm. Spur color: 145A overlain with 59A.

Peduncles.—Length: About 2.1 cm. Diameter: About 1.25 mm. Angle: Acute. Strength: Strong. Color: 144B.

Reproductive organs.—None observed.

Seed/fruit.—None observed.

Disease-pest resistance: Plants of the new *Impatiens* have not been observed to be resistant to pathogens and pests common to *Impatiens*.

It is claimed:

1. A new and distinct cultivar of Double *Impatiens* plant named 'Balolesal', as illustrated and described.

* * * * *

