

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
27 May 2004 (27.05.2004)

PCT

(10) International Publication Number
WO 2004/044046 A2

- (51) International Patent Classification⁷: **C08L** William [US/US]; 18 Woodland Drive, West Paterson, NJ 08844 (US).
- (21) International Application Number: PCT/US2003/035526 (74) Agent: **KATZ, Martin, L.**; Wood, Phillips, Katz, Clark & Mortimer, 500 West Madison Street, Suite 3800, Chicago, IL 60661 (US).
- (22) International Filing Date: 7 November 2003 (07.11.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/424,928 8 November 2002 (08.11.2002) US
- (71) Applicants (for all designated States except US): **ENCY-SIVE PHARMACEUTICALS** [US/US]; 6700 West Loop South, Bellaire, TX 77401 (US). **SCHERING CORPORATION** [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **VANDERSLICE, Peter** [US/US]; 3427 Gannett, Houston, TX 77025 (US). **HOLLAND, George** [US/US]; 3331 Green Tree Park, Houston, TX 77007 (US). **SHIH, Neng-Yang** [US/US]; 1 Maple Drive, North Caldwell, NJ 07006 (US). **ASLANIAN, Robert, G.** [US/US]; 144 Philip Drive, Rockaway, NJ 07866 (US). **CHAPMAN, Richard, W.** [US/US]; 30 Ditmars Circle, Hillsboro, NJ 07866 (US). **KREUTNER,**
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMBINATION PRODUCTS WITH CARBOXYLIC ACID DERIVATIVES THAT INHIBIT THE BINDING OF INTEGRINS TO THEIR RECEPTORS AND OTHER THERAPEUTIC COMPOUNDS

(57) Abstract: A composition, method and kit comprising a compound for the inhibition of the binding of $\alpha_4\beta_1$ integrin to its receptors, for example VCAM-1 (vascular cell adhesion molecule-1) and fibronectin and other therapeutic compounds for the control or prevention of diseases states in which $\alpha_4\beta_1$ is involved.

WO 2004/044046 A2

**COMBINATION PRODUCTS WITH CARBOXYLIC ACID DERIVATIVES THAT
INHIBIT THE BINDING OF INTEGRINS TO THEIR RECEPTORS AND OTHER
THERAPEUTIC COMPOUNDS**

5 Field of the Invention

This invention is directed generally to combination products of and methods of co-administering compounds that inhibit the binding of $\alpha_4\beta_1$ integrin to its receptors, for example VCAM-1 (vascular cell adhesion molecule-1) and fibronectin and other therapeutic compounds. The invention also relates to the use of such compositions and methods for the control or prevention of disease states in which $\alpha_4\beta_1$ is involved.

Background of the Invention

When a tissue has been invaded by a microorganism or has been damaged, white blood cells, also called leukocytes, play a major role in the inflammatory response. One of the most important aspects of the inflammatory response involves the cell adhesion event.

15 Generally, white blood cells are found circulating through the bloodstream. However, when a tissue is infected or becomes damaged, the white blood cells recognize the invaded or damaged tissue, bind to the wall of the capillary and migrate through the capillary into the affected tissue. These events are mediated by a family of proteins called cell adhesion molecules.

20 There are three main types of white blood cells: granulocytes, monocytes and lymphocytes. The integrin $\alpha_4\beta_1$ (also called VLA-4 for very late antigen-4) is a heterodimeric protein expressed on the surface of monocytes, lymphocytes and two subclasses of granulocytes: eosinophils and basophils. This protein plays a key role in cell adhesion through its ability to recognize and bind VCAM-1 and fibronectin, proteins associated with

25 the endothelial cells that line the interior wall of capillaries.

Following infection or damage of tissue surrounding a capillary, endothelial cells express a series of adhesion molecules, including VCAM-1, that are critical for binding the white blood cells that are necessary for fighting infection. Prior to binding to VCAM-1 or fibronectin, the white blood cells initially bind to certain adhesion molecules to slow their

30 flow and allow the cells to "roll" along the activated endothelium. Monocytes, lymphocytes, basophils and eosinophils are then able to firmly bind to VCAM-1 or fibronectin on the blood

-2-

vessel wall *via* the $\alpha_4\beta_1$ integrin. There is evidence that such interactions are also involved in transmigration of these white blood cells into the damaged tissue as well as the initial rolling event itself.

Although white blood cell migration to the site of injury helps fight infection and destroy foreign material, in many instances this migration can become uncontrolled, with white blood cells flooding to the scene, causing widespread tissue damage. Compounds capable of blocking this process, therefore, may be beneficial as therapeutic agents. Thus, it would be useful to develop inhibitors that would prevent the binding of white blood cells to VCAM-1 and fibronectin.


Some of the diseases that might be treated by the combination of compounds for the inhibition of $\alpha_4\beta_1$ binding and other therapeutic compounds include, but are not limited to, psoriasis, atherosclerosis, rheumatoid arthritis, asthma, allergy, multiple sclerosis, Guillan-Barr Syndrome, lupus, inflammatory bowel disease, graft rejection, contact hypersensitivity, reperfusion injury and type I diabetes.

It is therefore an object of the invention to provide novel compositions comprising compounds which are inhibitors of $\alpha_4\beta_1$ binding and other therapeutic agents.

Brief Summary of the Invention

The present invention is directed to pharmaceutical compositions that comprise a compound that inhibits binding of integrins to their receptors and one or more other therapeutic agents.

More particularly, the present invention is directed to a pharmaceutical composition that comprises a compound of Formula I



Formula I

wherein Y, at each occurrence, is independently selected from the group consisting

-3-

of C(O), N, CR¹, C(R²)(R³), NR⁵, CH, O and S;

q is an integer of from 3 to 10;

A is selected from the group consisting of O, S, C(R¹⁶)(R¹⁷) and NR⁶;

E is selected from the group consisting of CH₂, O, S, and

5 NR⁷;

J is selected from the group consisting of O, S and NR⁸;

T is selected from the group consisting of C(O) and (CH₂)_b wherein b is an integer
of from 0 to 3;

M is selected from the group consisting of C(R⁹)(R¹⁰) and

10 (CH₂)_u, wherein u is an integer of from 0 to 3;

L is selected from the group consisting of O, NR¹¹, S, and

(CH₂)_n wherein n is an integer of 0 or 1;

X is selected from the group consisting of CO₂B, PO₃H₂,

SO₃H, SO₂NH₂, SO₂NHCOR¹², OPO₃H₂, C(O)NHC(O)R¹³,

15 C(O)NHSO₂R¹⁴, hydroxyl, tetrazolyl and hydrogen;

W is selected from the group consisting of C, CR¹⁵ and N;

B is selected from the group consisting of hydrogen, alkyl, alkenyl,

alkynyl, hydroxyalkyl, haloalkyl, -CF₃, cycloalkyl, cycloalkenyl,

cycloalkynyl, cycloalkylalkyl, aryl, heterocyclyl, alkylaryl, aralkenyl, aralkyl,

20 alkylheterocyclyl, heterocyclylalkyl and aryloxyalkyl; and

R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁶ and R¹⁷ at each

occurrence are independently selected from the group consisting of

hydrogen, halogen, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy,

alkynoxy, thioalkoxy, hydroxyalkyl, aliphatic acyl, -CF₃, -CO₂H, -SH,

25 -CN, -NO₂, -NH₂, -OH, alkynylamino, alkoxy carbonyl, heterocycloyl, carboxy,

-N(C₁-C₃ alkyl)-C(O)(C₁-C₃ alkyl), -NHC(O)N(C₁-C₃ alkyl)C(O)-

NH(C₁-C₃alkyl), -NHC(O)NH(C₁-C₆ alkyl), -NHSO₂(C₁-C₃ alkyl),

-NHSO₂(aryl), alkoxyalkyl, alkylamino, alkenylamino, di(C₁-C₃)amino,

-C(O)O-(C₁-C₃)alkyl, -C(O)NH-(C₁-C₃)alkyl, -C(O)N(C₁-C₃ alkyl)₂,

30 -CH=NOH, -PO₃H₂, -OPO₃H₂, haloalkyl, alkoxyalkoxy, carboxaldehyde,

carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl,

-4-


aryl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl, -SO₂-(C₁-C₃ alkyl), -SO₃-(C₁-C₃ alkyl), sulfonamido, carbamate, aryloxyalkyl and -C(O)NH(benzyl) groups;

- 5 wherein B, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁶ and R¹⁷ are unsubstituted or substituted with at least one electron donating or electron withdrawing group;
- wherein when L is NR¹¹, R⁴ and R¹¹ taken together may form a ring; and wherein when M is C(R⁹)(R¹⁰), R⁹ and R¹⁰ taken together may
- 10 form a ring;
- and wherein when A is NR⁶ and at least one Y is CR¹, R¹ and R⁶ taken together may form a ring;

or a pharmaceutically acceptable salt thereof;
and one or more other therapeutic compounds.

- 15 For Formula I, presently preferred compounds may have A as NR⁶; E as NR⁷; J as O; M as C(R⁹)(R¹⁰); q as 4 or 5; T as (CH₂)_b wherein b is 0; L as (CH₂)_n wherein n is 0; X as CO₂B; W as C or CR¹⁵; R⁴ as aryl, alkylaryl, aralkyl, heterocyclyl, alkylheterocyclyl or heterocyclylalkyl; and R⁶, R⁷, R⁹, R¹⁰ and R¹⁵ independently as hydrogen or lower alkyl.

- 20 More specifically, the compositions of this invention comprise compounds of Formula II



Formula II

wherein Y, at each occurrence, is independently selected from the group consisting of C(O), N, CR¹, C(R²)(R³), NR⁵, CH, O and S;

-5-

q is an integer of from 3 to 7;

T is selected from the group consisting of C(O) and (CH₂)_b wherein b is an integer of 0 to 3;

L is selected from the group consisting of O, NR¹¹, S, and

5 (CH₂)_n wherein n is an integer of 0 or 1;

W is selected from the group consisting of C, CR¹⁵ and N;

B is selected from the group consisting of hydrogen, alkyl, alkenyl,

alkynyl, hydroxyalkyl, haloalkyl, -CF₃, cycloalkyl, cycloalkenyl,

cycloalkynyl, cycloalkylalkyl, aryl, heterocyclyl, alkylaryl, aralkenyl, aralkyl,

10 alkylheterocyclyl, heterocyclylalkyl and aryloxyalkyl; and

R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁹, R¹⁰, R¹¹ and R¹⁵ are independently selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl,

alkoxy, alkenoxy, alkynoxy, thioalkoxy, hydroxyalkyl, aliphatic acyl, -CF₃,

-CO₂H, -SH, -CN, -NO₂, -NH₂, -OH, alkynylamino, alkoxyacetyl,

15 heterocycloyl, carboxy, -N(C₁-C₃ alkyl)-C(O)(C₁-C₃ alkyl),

-NHC(O)N(C₁-C₃ alkyl)C(O)NH(C₁-C₃alkyl), -NHC(O)NH(C₁-C₆ alkyl),

-NHSO₂(C₁-C₃ alkyl), -NHSO₂(aryl), alkoxyalkyl, alkylamino, alkenylamino,

di(C₁-C₃)amino, -C(O)O-(C₁-C₃)alkyl, -C(O)NH-(C₁-C₃)alkyl, -C(O)N(C₁-C₃ alkyl)₂, -CH=NOH, -PO₃H₂, -OPO₃H₂, haloalkyl, alkoxyalkoxy,

20 carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl,

cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino,

heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl,

sulfonyl, -SO₂-(C₁-C₃ alkyl), -SO₃-(C₁-C₃ alkyl), sulfonamido, carbamate,

aryloxyalkyl and -C(O)NH(benzyl) groups;

25 wherein B, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁹, R¹⁰, R¹¹ and R¹⁵ are unsubstituted or substituted with at least one electron donating or electron withdrawing group;

wherein when L is NR¹¹, R⁴ and R¹¹ taken together may form a ring;

and wherein R⁹ and R¹⁰ taken together may form a ring;


30 and wherein when at least one Y is CR¹, R¹ and R⁶ taken together may form a ring;

-6-

or a pharmaceutically acceptable salt thereof and one or more other therapeutic compounds.

For Formula II, presently preferred compounds may have q as 4 or 5; W as C or CR^{15} ; T as $(CH_2)_b$ wherein b is 0; L as $(CH_2)_n$ wherein n is 0; R^4 as aryl, alkylaryl, aralkyl, heterocyclyl, alkylheterocyclyl or heterocyclylalkyl; and R^6 , R^7 , R^9 , R^{10} and R^{15} as independently hydrogen or lower alkyl.

Even more specifically, the compositions of this invention comprise a compound of Formula III



10

Formula III

wherein Y , at each occurrence, is independently selected from the group consisting of $C(O)$, N , CR^1 , $C(R^2)(R^3)$, NR^5 , CH , O and S ;
 q is an integer of from 2 to 5;
 T is selected from the group consisting of $C(O)$ and $(CH_2)_b$ wherein b is an integer of 0 to 3;
 L is selected from the group consisting of O , NR^{11} , S , and $(CH_2)_n$ wherein n is an integer of 0 or 1;
 R^5 , R^6 , R^7 , R^{11} and R^{18} are each independently selected from the group consisting of alkyl, alkenyl, alkynyl, hydroxyalkyl, aliphatic acyl, alkynylamino, alkoxyalkyl, heterocyclyl, $-CH=NOH$, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl,

25

-7-

alkylheterocyclyl, heterocyclylalkyl, carbamate, aryloxyalkyl, hydrogen and –
C(O)NH(benzyl) groups;

B is selected from the group consisting of hydrogen, alkyl, alkenyl,
alkynyl, hydroxyalkyl, haloalkyl, -CF₃, cycloalkyl, cycloalkenyl,
5 cycloalkynyl, cycloalkylalkyl, aryl, heterocyclyl, alkylaryl, aralkenyl, aralkyl,
alkylheterocyclyl, heterocyclylalkyl and aryloxyalkyl; and

R¹, R², R³, R⁴, R⁹ and R¹⁰ are independently selected from the
group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl,
alkoxy, alkenoxy, alkynoxy, thioalkoxy, hydroxyalkyl, aliphatic acyl, -CF₃,
10 -CO₂H, -SH, -CN, -NO₂, -NH₂, -OH, alkynylamino, alkoxy carbonyl,
heterocycloyl, carboxy, -N(C₁-C₃ alkyl)-C(O)(C₁-C₃ alkyl),
-NHC(O)N(C₁-C₃ alkyl)C(O)NH(C₁-C₃alkyl), -NHC(O)NH(C₁-C₆ alkyl),
-NHSO₂(C₁-C₃ alkyl), -NHSO₂(aryl), alkoxyalkyl, alkylamino, alkenylamino,
di(C₁-C₃)amino, -C(O)O-(C₁-C₃)alkyl, -C(O)NH- (C₁-C₃)alkyl, -C(O)N(C₁-C₃
15 alkyl)₂, -CH=NOH, -PO₃H₂, -OPO₃H₂, haloalkyl, alkoxyalkoxy,
carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl,
cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino,
heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl,
sulfonyl, -SO₂-(C₁-C₃ alkyl), -SO₃-(C₁-C₃ alkyl), sulfonamido, carbamate,
20 aryloxyalkyl and –C(O)NH(benzyl) groups;

wherein B, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁹, R¹⁰, R¹¹ and R¹⁸ are unsubstituted or
substituted with at least one electron donating or electron withdrawing group;

wherein when L is NR¹¹, R⁴ and R¹¹ taken together may form a ring;

and wherein R⁹ and R¹⁰ taken together may form a ring;


25 and wherein when at least one Y is CR¹, R¹ and R⁶ taken
together may form a ring;

or a pharmaceutically acceptable salt thereof and one or more other therapeutic
compounds.

Presently preferred compounds of Formula III may have R¹⁸ as hydrogen, alkyl, aryl,
30 aralkyl, cycloalkyl, alkylheterocyclyl, heterocyclylalkyl or heterocyclyl; T as (CH₂)_b wherein b
is 0; L as (CH₂)_n wherein n is 0; Y as CR¹ and C(R²)(R³) and q as 2 or 3.


-8-

In Formula III, the portion of the molecule



can be

5



and pharmaceutical acceptable salts thereof

- 10 wherein R^{19} , R^{20} , R^{21} and R^{22} at each occurrence are independently selected from the group consisting of halogen, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkynoxy, thioalkoxy, hydroxyalkyl, aliphatic acyl, $-CF_3$, $-OH$, $-CO_2H$, $-SH$, $-CN$, $-NO_2$, $-NH_2$, alkynylamino, alkoxy carbonyl, heterocycloyl, carboxy, $-N(C_1-C_3 \text{ alkyl})-C(O)(C_1-C_3 \text{ alkyl})$, $-NHC(O)N(C_1-C_3 \text{ alkyl})C(O)NH(C_1-C_3 \text{ alkyl})$, $-NHC(O)NH(C_1-C_6 \text{ alkyl})$, $-NHSO_2(C_1-C_3 \text{ alkyl})$, $-NHSO_2(\text{aryl})$, alkoxyalkyl, alkylamino, alkenylamino,
- 15

di(C₁-C₃)amino, -C(O)O-(C₁-C₃)alkyl, -C(O)NH-(C₁-C₃)alkyl, -C(O)N(C₁-C₃alkyl)₂, -CH=NOH, -PO₃H₂, -OPO₃H₂, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl, -SO₂-(C₁-C₃alkyl), -SO₃-(C₁-C₃alkyl), sulfonamido, carbamate, aryloxyalkyl and -C(O)NH(benzyl) groups;

R¹⁸ is selected from the group consisting of alkyl, alkenyl, alkynyl, hydroxyalkyl, aliphatic acyl, alkynylamino, alkoxyacetyl, heterocycloyl, -CH=NOH, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, carbamate, aryloxyalkyl, hydrogen and -C(O)NH(benzyl) groups;

R²² is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkynoxy, thioalkoxy, hydroxyalkyl, aliphatic acyl, -CF₃, -CO₂H, -SH, -CN, -NO₂, -NH₂, -OH, alkynylamino, alkoxyacetyl, heterocycloyl, carboxy, -N(C₁-C₃alkyl)-C(O)(C₁-C₃alkyl), -NHC(O)N(C₁-C₃alkyl)C(O)NH(C₁-C₃alkyl), -NHC(O)NH(C₁-C₆alkyl), -NHSO₂(C₁-C₃alkyl), -NHSO₂(aryl), alkoxyalkyl, alkylamino, alkenylamino, di(C₁-C₃)amino, -C(O)O-(C₁-C₃)alkyl, -C(O)NH-(C₁-C₃)alkyl, -C(O)N(C₁-C₃alkyl)₂, -CH=NOH, -PO₃H₂, -OPO₃H₂, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl, -SO₂-(C₁-C₃alkyl), -SO₃-(C₁-C₃alkyl), sulfonamido, carbamate, aryloxyalkyl and -C(O)NH(benzyl) groups;

c is an integer of zero to two;

d is an integer of zero to three;

e is an integer of zero to four; and


-10-

i is an integer of zero to two.

In one embodiment, R^{18} is aralkyl; R^4 is aryl; T is $(CH_2)_b$ where *b* is zero; L is $(CH_2)_n$ where *n* is zero; and, B, R^6 , R^7 , R^9 and R^{10} are each independently hydrogen.

Even more specifically, the compositions of this invention comprise compounds of

5 Formula IV



Formula IV

wherein T is selected from the group consisting of C(O) and $(CH_2)_b$ wherein *b* is an integer of from 0 to 3;

10 L is selected from the group consisting of O, NR^{11} , S, and $(CH_2)_n$ wherein *n* is an integer of 0 or 1;

g is an integer of from 0 to 7;

B is selected from the group consisting of hydrogen, alkyl, alkenyl,

alkynyl, hydroxyalkyl, haloalkyl, $-CF_3$, cycloalkyl, cycloalkenyl,

15 cycloalkynyl, cycloalkylalkyl, aryl, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl and aryloxyalkyl; and

R^4 , R^9 , R^{10} and R^{23} at each occurrence are independently selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkynoxy, thioalkoxy, hydroxyalkyl, aliphatic acyl,

20 $-CF_3$, $-CO_2H$, $-SH$, $-CN$, $-NO_2$, $-NH_2$, $-OH$, alkynylamino, alkoxy carbonyl, heterocycloyl, carboxy, $-N(C_1-C_3 \text{ alkyl})-C(O)(C_1-C_3 \text{ alkyl})$,

$-NHC(O)N(C_1-C_3 \text{ alkyl})C(O)NH(C_1-C_3 \text{ alkyl})$ $-NHC(O)NH(C_1-C_6 \text{ alkyl})$,

$-NHSO_2(C_1-C_3 \text{ alkyl})$, $-NHSO_2(\text{aryl})$, alkoxyalkyl, alkylamino,

25 alkenylamino, $di(C_1-C_3)amino$, $-C(O)O-(C_1-C_3)alkyl$, $-C(O)NH-(C_1-$

-11-

C₃alkyl, -C(O)N(C₁-C₃ alkyl)₂, -CH=NOH, -PO₃H₂, -OPO₃H₂, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl, -SO₂-(C₁-C₃ alkyl), -SO₃-(C₁-C₃ alkyl), sulfonamido, carbamate, aryloxyalkyl and -C(O)NH(benzyl) groups; and


R⁶, R⁷, R¹¹ and R¹⁸ are each independently selected from the group consisting of alkyl, alkenyl, alkynyl, hydroxyalkyl, aliphatic acyl, alkynylamino, alkoxyacetyl, heterocyclyl, -CH=NOH, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, carbamate, aryloxyalkyl, hydrogen and -C(O)NH(benzyl) groups;

wherein B, R⁴, R⁶, R⁷, R⁹, R¹⁰, R¹¹, R¹⁸ and R²³ are unsubstituted or substituted with at least one electron donating or electron withdrawing group;

wherein when L is NR¹¹, R⁴ and R¹¹ taken together may form a ring; and wherein R⁹ and R¹⁰ taken together may form a ring;

or a pharmaceutically acceptable salt thereof and one or more other therapeutic compounds.

Presently preferred compositions of the present invention comprise compounds of Formula V



Formula V

25

-12-

wherein h is an integer of zero to five;

B is selected from the group consisting of hydrogen, alkyl, alkenyl,

alkynyl, hydroxyalkyl, haloalkyl, -CF₃, cycloalkyl, cycloalkenyl,

5 cycloalkynyl, cycloalkylalkyl, aryl, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl and aryloxyalkyl;

R⁹, R¹⁰, R²⁴, and R²⁵ are each independently selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkynoxy,

thioalkoxy, hydroxyalkyl, aliphatic acyl, -CF₃, -CO₂H, -SH, -CN, -NO₂,

10 -NH₂, -OH, alkynylamino, alkoxycarbonyl, heterocycloyl, carboxy,

-N(C₁-C₃ alkyl)-C(O)(C₁-C₃ alkyl),

-NHC(O)N(C₁-C₃ alkyl)C(O)NH(C₁-C₃alkyl), -NHC(O)NH(C₁-C₆ alkyl),

-NH₂SO₂(C₁-C₃ alkyl), -NH₂SO₂(aryl), alkoxyalkyl, alkylamino,

15 alkenylamino, di(C₁-C₃)amino, -C(O)O-(C₁-C₃)alkyl,

-C(O)NH-(C₁-C₃)alkyl, -C(O)N(C₁-C₃ alkyl)₂, -CH=NOH, -PO₃H₂,

-OPO₃H₂, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide,

cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl,

aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl,

20 aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl,

-SO₂-(C₁-C₃ alkyl), -SO₃-(C₁-C₃ alkyl), sulfonamido, carbamate,

aryloxyalkyl and -C(O)NH(benzyl) groups;

R²⁷, at each occurrence, is independently selected from the group consisting of

halogen, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkynoxy, thioalkoxy,

hydroxyalkyl, aliphatic acyl, -CF₃, -CO₂H, -SH, -CN, -NO₂, -NH₂, -OH,

25 alkynylamino, alkoxycarbonyl, heterocycloyl, carboxy, -N(C₁-C₃ alkyl)-

C(O)(C₁-C₃ alkyl), -NHC(O)N(C₁-C₃ alkyl)C(O)NH(C₁-C₃alkyl),

-NHC(O)NH(C₁-C₆ alkyl), -NH₂SO₂(C₁-C₃ alkyl), -NH₂SO₂(aryl), -N(C₁-C₃

alkyl)SO₂(C₁-C₃alkyl), -N(C₁-C₃alkyl)SO₂(aryl), -C

alkoxyalkyl, alkylamino, alkenylamino, di(C₁-C₃)amino, -C(O)O-(C₁-

30 C₃)alkyl, -C(O)NH-(C₁-C₃)alkyl, -C(O)N(C₁-C₃ alkyl)₂, -CH=NOH,

-PO₃H₂, -OPO₃H₂, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide,

cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl, -SO₂-(C₁-C₃ alkyl), -SO₃-(C₁-C₃ alkyl), sulfonamido, carbamate, aryloxyalkyl and -C(O)NH(benzyl) groups;

5 R⁶, R⁷ and R¹⁸ are each independently selected from the group consisting of alkyl, alkenyl, alkynyl, hydroxyalkyl, aliphatic acyl, alkynylamino, alkoxy carbonyl, heterocycloyl, -CH=NOH, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, carbamate, aryloxyalkyl, hydrogen and -C(O)NH(benzyl) groups; and,

10 R²⁶ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, hydroxyalkyl, aliphatic acyl, -CF₃, alkoxy carbonyl, heterocycloyl, carboxy, -C(O)O-(C₁-C₃)alkyl, -C(O)NH-(C₁-C₃)alkyl, -C(O)N(C₁-C₃ alkyl)₂, -PO₃H₂, haloalkyl, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, biaryl, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl, -SO₂-(C₁-C₃ alkyl), sulfonamido, aryloxyalkyl and -C(O)NH(benzyl) groups;

15 wherein B, R⁶, R⁷, R⁹, R¹⁰, R¹⁸, R²⁴, R²⁵, R²⁶ and R²⁷ are unsubstituted or substituted with at least one electron donating or electron withdrawing group;

wherein R¹⁸ and R²⁴ taken together may form a ring;

R²⁴ and R²⁵ taken together may form a ring;

R²⁵ and R²⁶ taken together may form a ring;

25 and wherein R⁹ and R¹⁰ taken together may form a ring;

or a pharmaceutically acceptable salt thereof and one or more other therapeutic compounds.


Presently preferred compounds of Formula V have B, R⁶, R⁷, R⁹, R¹⁰, R²⁴, R²⁵ and R²⁶ each independently hydrogen and R¹⁸ as substituted or unsubstituted aralkyl.

30

Other presently preferred compositions of the present invention comprise compounds

of

Formula VI



Formula VI

5

wherein Z, at each occurrence, is independently selected from the group consisting of C(O), N, CR³⁰, C(R³¹)(R³²), NR³³, CH, O and S;

z is an integer of from 3 to 6;

k is an integer of from 0 to 5;

10 T is selected from the group consisting of C(O) and (CH₂)_b wherein b is an integer of from 0 to 3;

L is selected from the group consisting of O, NR¹¹, S, and (CH₂)_n wherein n is an integer of 0 or 1;

15 R⁶, R⁷, R¹¹, R¹⁸ and R³³ are each independently selected from the group consisting of alkyl, alkenyl, alkynyl, hydroxyalkyl, aliphatic acyl, alkynylamino, alkoxy carbonyl, heterocycloyl, -CH=NOH, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, carbamate, aryloxyalkyl, hydrogen and -C(O)NH(benzyl) groups;

20

B is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, hydroxyalkyl, haloalkyl, -CF₃, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, heterocyclyl, alkylanyl, aralkenyl, aralkyl,

-15-

alkylheterocyclyl, heterocyclylalkyl and aryoxyalkyl;

R⁴, R⁹, R¹⁰, R³⁰, R³¹ and R³² at each occurrence are independently selected from
 the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, alkoxy,
 5 alkenoxy, alkynoxy, thioalkoxy, hydroxyalkyl, aliphatic acyl, -CF₃,
 -CO₂H, -SH, -CN, -NO₂, -NH₂, -OH, alkynylamino, alkoxy carbonyl,
 heterocycloyl, carboxy, -N(C₁-C₃ alkyl)-C(O)(C₁-C₃ alkyl),
 -NHC(O)N(C₁-C₃ alkyl)C(O)NH(C₁-C₃alkyl), -NHC(O)NH(C₁-C₆ alkyl),
 -NHSO₂(C₁-C₃ alkyl), -NHSO₂(aryl), alkoxyalkyl, alkylamino,
 10 alkenylamino, di(C₁-C₃)amino, -C(O)O-(C₁-C₃)alkyl,
 -C(O)NH-(C₁-C₃)alkyl, -C(O)N(C₁-C₃ alkyl)₂, -CH=NOH, -PO₃H₂,
 -OPO₃H₂, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide,
 cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl,
 aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl,
 15 aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl, -SO₂-(C₁-
 C₃ alkyl), -SO₃-(C₁-C₃ alkyl), sulfonamido, carbamate, aryloxyalkyl and
 -C(O)NH(benzyl) groups; and,

R²⁹, at each occurrence, is independently selected from the group consisting of
 halogen, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkynoxy, thioalkoxy,
 20 hydroxyalkyl, aliphatic acyl, -CF₃, -CO₂H, -SH, -CN, -NO₂, -NH₂, -OH,
 alkynylamino, alkoxy carbonyl, heterocycloyl, carboxy, -N(C₁-C₃ alkyl)-
 C(O)(C₁-C₃ alkyl), -NHC(O)N(C₁-C₃ alkyl)C(O)NH(C₁-C₃alkyl),
 -NHC(O)NH(C₁-C₆ alkyl), -NHSO₂(C₁-C₃ alkyl), -NHSO₂(aryl),
 alkoxyalkyl, alkylamino, alkenylamino, di(C₁-C₃)amino, -C(O)O-(C₁-
 25 C₃)alkyl, -C(O)NH-(C₁-C₃)alkyl, -C(O)N(C₁-C₃ alkyl)₂, -CH=NOH,
 -PO₃H₂, -OPO₃H₂, haloalkyl, alkoxyalkoxy, carboxaldehyde,
 carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl,
 aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl,
 alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl,
 30 -SO₂-(C₁-C₃ alkyl), -SO₃-(C₁-C₃ alkyl), sulfonamido, carbamate,
 aryloxyalkyl and -C(O)NH(benzyl) groups;

-16-

wherein B, R⁴, R⁶, R⁷, R⁹, R¹⁰, R¹¹, R¹⁸, R²⁹, R³⁰, R³¹, R³² and R³³ are unsubstituted or substituted with at least one electron donating or electron withdrawing group;

wherein when L is NR¹¹, R⁴ and R¹¹ taken together may form a ring; and

5 wherein R⁹ and R¹⁰ taken together may form a ring;

or a pharmaceutically acceptable salt thereof and one or more other therapeutic compounds.

Presently preferred compounds that inhibit $\alpha_4\beta_1$ binding and are useful in the compositions of the present invention include (3S)-3-[(1-(2-chlorobenzyl)-4-hydroxy-5-methyl-2-oxo-1,2-dihydropyridin-3-yl)amino]carbonylamino]-3-(4-methylphenyl)propanoic acid and pharmaceutically acceptable salts thereof; (3S)-3-[(1-(2-chlorobenzyl)-4-hydroxy-2-oxo-2,5,6,7-tetrahydro-1H-cyclopenta[b]pyridin-3-yl)amino]carbonylamino]-3-(4-methylphenyl)propanoic acid and pharmaceutically acceptable salts thereof; and (3S)-3-[(1-(2-chlorobenzyl)-4-hydroxy-5-methyl-2-oxo-1,2-dihydropyridin-3-yl)amino]carbonylamino]-3-[3-(diethylamino)phenyl]propanoic acid and pharmaceutically acceptable salts thereof.

The pharmaceutical compositions of the present invention also comprise other therapeutic compounds and physiologically acceptable diluents.

The invention also provides a method for treating disease states mediated by $\alpha_4\beta_1$ binding which comprises administration of an effective amount of a composition of the present invention to an afflicted patient.

The present invention is also directed to a method for treating disease states mediated by $\alpha_4\beta_1$ binding which comprises co-administering a therapeutically effective amount of a combination of a compound of Formulae I, II, III, IV, V or VI and a therapeutically effective amount of one or more other therapeutic compounds to an afflicted patient.

25 The present invention is further directed to a kit comprising in a single package, one container comprising a compound that inhibits binding of $\alpha_4\beta_1$ integrin to its receptors in a pharmaceutically acceptable carrier and one or more separate containers comprising other therapeutic compounds in pharmaceutically acceptable carriers, with the compound that inhibits binding of $\alpha_4\beta_1$ integrin to its receptors and the other therapeutic compounds being present in amounts such that the combination is effective to treat disease states mediated by $\alpha_4\beta_1$ integrin binding.

Detailed Description of the Invention

Definitions of Terms

The term "alkyl" as used herein, alone or in combination, refers to C₁-C₁₂ straight or
5 branched, substituted or unsubstituted saturated chain radicals derived from saturated
hydrocarbons by the removal of one hydrogen atom, unless the term alkyl is preceded by a
C_x-C_y designation. Representative examples of alkyl groups include methyl, ethyl,
n-propyl, iso-propyl, n-butyl, sec-butyl, iso-butyl, and tert-butyl among others.

The term "alkenyl" as used herein, alone or in combination, refers to a substituted or
10 unsubstituted straight-chain or substituted or unsubstituted branched-chain alkenyl radical
containing from 2 to 10 carbon atoms. Examples of such radicals include, but are not limited
to, ethenyl, E- and Z-pentenyl, decenyl and the like.

The term "alkynyl" as used herein, alone or in combination, refers to a substituted or
unsubstituted straight or substituted or unsubstituted branched chain alkynyl radical
15 containing from 2 to 10 carbon atoms. Examples of such radicals include, but are not limited
to ethynyl, propynyl, propargyl, butynyl, hexynyl, decynyl and the like.

The term "lower" modifying "alkyl", "alkenyl", "alkynyl" or "alkoxy" refers to a C₁-
C₆ unit for a particular functionality. For example lower alkyl means C₁-C₆ alkyl.

The term "aliphatic acyl" as used herein, alone or in combination, refers to radicals of
20 formula alkyl-C(O)-, alkenyl-C(O)- and alkynyl-C(O)- derived from an alkane-, alkene- or
alkynocarboxylic acid, wherein the terms "alkyl", "alkenyl" and "alkynyl" are as defined
above. Examples of such aliphatic acyl radicals include, but are not limited to, acetyl,
propionyl, butyryl, valeryl, 4-methylvaleryl, acryloyl, crotyl, propiolyl and methylpropiolyl,
among others.

The term "cycloalkyl" as used herein refers to an aliphatic ring system having 3 to 10
25 carbon atoms and 1 to 3 rings, including, but not limited to cyclopropyl, cyclopentyl,
cyclohexyl, norbornyl, and adamantyl among others. Cycloalkyl groups can be unsubstituted
or substituted with one, two or three substituents independently selected from lower alkyl,
haloalkyl, alkoxy, thioalkoxy, amino, alkylamino, dialkylamino, hydroxy, halo, mercapto,
30 nitro, carboxaldehyde, carboxy, alkoxy carbonyl and carboxamide.

"Cycloalkyl" includes cis or trans forms. Furthermore, the substituents may either be

in endo or exo positions in the bridged bicyclic systems.

The term "cycloalkenyl" as used herein alone or in combination refers to a cyclic carbocycle containing from 4 to 8 carbon atoms and one or more double bonds. Examples of such cycloalkenyl radicals include, but are not limited to, cyclopentenyl, cyclohexenyl,
5 cyclopentadienyl and the like.

The term "cycloalkylalkyl" as used herein refers to a cycloalkyl group appended to a lower alkyl radical, including, but not limited to cyclohexylmethyl.

The term "halo" or "halogen" as used herein refers to I, Br, Cl or F.

The term "haloalkyl" as used herein refers to a lower alkyl radical, to which is
10 appended at least one halogen substituent, for example chloromethyl, fluoroethyl, trifluoromethyl and pentafluoroethyl among others.

The term "alkoxy" as used herein, alone or in combination, refers to an alkyl ether radical, wherein the term "alkyl" is as defined above. Examples of suitable alkyl ether radicals include, but are not limited to, methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy,
15 iso-butoxy, sec-butoxy, tert-butoxy and the like.

The term "alkoxyalkyl" as used herein, refers to R_y-O-R_z , wherein R_y is lower alkyl as defined above, and R_z is alkylene $-(CH_2)_w-$ wherein w is an integer of from one to six. Representative examples include methoxymethyl, methoxyethyl, and ethoxyethyl among others.

The term "alkenoxy" as used herein, alone or in combination, refers to a radical of
20 formula alkenyl-O, provided that the radical is not an enol ether, wherein the term "alkenyl" is as defined above. Examples of suitable alkenoxy radicals include, but are not limited to, allyloxy, E- and Z- 3-methyl-2-propenoxy and the like.

The term "alkynoxy" as used herein, alone or in combination, refers to a radical of
25 formula alkynyl-O, provided that the radical is not an -ynol ether. Examples of suitable alkynoxy radicals include, but are not limited to, propargyloxy, 2-butynyloxy and the like.

The term "carboxy" as used herein refers to $-C(O)O-$.

The term "thioalkoxy" refers to a thioether radical of formula alkyl-S-, wherein
"alkyl" is as defined above.

30 The term "sulfonamido" as used herein refers to $-SO_2NH_2$.

The term "carboxaldehyde" as used herein refers to $-C(O)R$ wherein R is hydrogen.

-19-

The terms "carboxamide" or "amide" as used herein refer to $-C(O)NR_aR_b$ wherein R_a and R_b are each independently hydrogen, alkyl or any other suitable substituent.

The term "alkoxyalkoxy" as used herein refers to R_cO-R_dO- wherein R_c is lower alkyl as defined above and R_d is alkylene wherein alkylene is $-(CH_2)_{n'}$ wherein n' is an integer
5 from 1 to 6. Representative examples of alkoxyalkoxy groups include methoxymethoxy, ethoxymethoxy, t-butoxymethoxy among others.

The term "alkylamino" as used herein refers to R_eNH- wherein R_e is a lower alkyl group, for example, ethylamino, butylamino, among others.

The term "alkenylamino" as used herein, alone or in combination, refers to a radical of
10 formula alkenyl-NH- or $(alkenyl)_2N-$, wherein the term "alkenyl" is as defined above, provided that the radical is not an enamine. An example of such alkenylamino radical is the allylamino radical.

The term "alkynylamino" as used herein, alone or in combination, refers to a radical of formula alkynyl-NH- or $(alkynyl)_2N-$ wherein the term "alkynyl" is as defined above,
15 provided that the radical is not an amine. An example of such alkynylamino radicals is the propargyl amino radical.

The term "dialkylamino" as used herein refers to R_fR_gN- wherein R_f and R_g are independently selected from lower alkyl, for example diethylamino, and methyl propylamino, among others.

20 The term "alkoxycarbonyl" as used herein refers to an alkoxy group as previously defined appended to the parent molecular moiety through a carbonyl group. Examples of alkoxycarbonyl include methoxycarbonyl, ethoxycarbonyl, and isopropoxycarbonyl among others.

The term "aryl" or "aromatic" as used herein alone or in combination refers to a
25 substituted or unsubstituted carbocyclic aromatic group having about 6 to 12 carbon atoms such as phenyl, naphthyl, indenyl, indanyl, azulenyl, fluorenyl and anthracenyl; or a heterocyclic aromatic group containing at least one endocyclic N, O or S atom such as furyl, thienyl, pyridyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, 2-pyrazolinyl, pyrazolidinyl, isoxazolyl, isothiazolyl, 1,2,3-oxadiazolyl, 1,2,3-triazolyl, 1,3,4-thiadiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, 1,3,5-triazinyl, 1,3,5-trithianyl, indolizinyl, indolyl,
30 isoindolyl, 3H-indolyl, indolynyl, benzo[b]furanlyl, 2,3-dihydrobenzofuranlyl,

benzo[b]thiophenyl, 1H-indazolyl, benzimidazolyl, benzthiazolyl, purinyl, 4H-quinolizinyl, isoquinolinyl, cinnolinyl, phthalazinyl, quinazoliny, quinoxaliny, 1,8-naphthridinyl, pteridinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, phenoxyazinyl, pyrazolo[1,5-c]triazinyl and the like. "Aralkyl" and "alkylaryl" employ the term "alkyl" as defined above.

5 Rings may be multiply substituted.

The term "aralkyl" as used herein, alone or in combination, refers to an aryl substituted alkyl radical, wherein the terms "alkyl" and "aryl" are as defined above.

Examples of suitable aralkyl radicals include, but are not limited to, phenylmethyl, phenethyl, phenylhexyl, diphenylmethyl, pyridylmethyl, tetrazolyl methyl, furylmethyl, imidazolyl
10 methyl, indolylmethyl, thienylpropyl and the like.

The term "aralkenyl" as used herein, alone or in combination, refers to an aryl substituted alkenyl radical, wherein the terms "aryl" and "alkenyl" are as defined above.

The term "arylamino" as used herein, alone or in combination, refers to a radical of formula aryl-NH-, wherein "aryl" is as defined above. Examples of arylamino radicals
15 include, but are not limited to, phenylamino(anilido), naphthlamino, 2-, 3-, and 4-pyridylamino and the like.

The term "benzyl" as used herein refers to $C_6H_5-CH_2-$.

The term "biaryl" as used herein, alone or in combination, refers to a radical of formula aryl-aryl, wherein the term "aryl" is as defined above.

20 The term "thioaryl" as used herein, alone or in combination, refers to a radical of formula aryl-S-, wherein the term "aryl" is as defined above. An example of a thioaryl radical is the thiophenyl radical.

The term "aroyle" as used herein, alone or in combination, refers to a radical of formula aryl-CO-, wherein the term "aryl" is as defined above. Examples of suitable
25 aromatic acyl radicals include, but are not limited to, benzoyl, 4-halobenzoyl, 4-carboxybenzoyl, naphthoyl, pyridylcarbonyl and the like.

The term "heterocyclyl" as used herein, alone or in combination, refers to a non-aromatic 3- to 10- membered ring containing at least one endocyclic N, O, or S atom. The heterocycle may be optionally aryl-fused. The heterocycle may also optionally be substituted
30 with at least one substituent which is independently selected from the group consisting of hydrogen, halogen, hydroxyl, amino, nitro, trifluoromethyl, trifluoromethoxy, alkyl, aralkyl,

alkenyl, alkynyl, aryl, cyano, carboxy, carboalkoxy, carboxyalkyl, oxo, arylsulfonyl and aralkylaminocarbonyl among others.

The term "alkylheterocyclyl" as used herein refers to an alkyl group as previously defined appended to the parent molecular moiety through a heterocyclyl group, including
5 but not limited to 2-methyl-5-thiazolyl, 2-methyl-1-pyrrolyl and 5-ethyl-2-thienyl.

The term "heterocyclylalkyl" as used herein refers to a heterocyclyl group as previously defined appended to the parent molecular moiety through an alkyl group, including but not limited to 2-thienylmethyl, 2-pyridinylmethyl and 2-(1-piperidinyl) ethyl.

The term "heterocycloyl" as used herein refers to radicals of the formula heterocyclyl-
10 C(O)-, wherein the term "heterocyclyl" is as defined above.

The term "aminal" as used herein refers to a hemi-acetal of the structure $R_hC(NR_iR_j)(NR_kR_l)$ - wherein R_h , R_i , R_j , R_k and R_l are each independently hydrogen, alkyl or any other suitable substituent.

The term "ester" as used herein refers to $-C(O)R_m$, wherein R_m is hydrogen, alkyl or
15 any other suitable substituent.

The term "carbamate" as used herein refers to compounds based on carbamic acid $NH_2C(O)OH$.

The term "optical isomers" as used herein refers to compounds which differ only in the stereochemistry of at least one atom, including enantiomers, diastereomers and racemates.

20 Use of the above terms is meant to encompass substituted and unsubstituted moieties. Substitution may be by one or more groups such as alcohols, ethers, esters, amides, sulfones, sulfides, hydroxyl, nitro, cyano, carboxy, amines, heteroatoms, lower alkyl, lower alkoxy, lower alkoxy carbonyl, alkoxyalkoxy, acyloxy, halogens, trifluoromethoxy, trifluoromethyl, alkyl, aralkyl, alkenyl, alkynyl, aryl, cyano, carboxy,
25 carboalkoxy, carboxyalkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, alkylheterocyclyl, heterocyclylalkyl, oxo, arylsulfonyl and aralkylaminocarbonyl or any of the substituents of the preceding paragraphs or any of those substituents either attached directly or by suitable linkers. The linkers are typically short chains of 1-3 atoms containing any combination of $-C-$, $-C(O)-$, $-NH-$, $-S-$, $-S(O)-$, $-O-$, $-C(O)O-$ or $-S(O)O-$. Rings may be substituted
30 multiple times.

The terms "electron-withdrawing" or "electron-donating" refer to the ability of a

substituent to withdraw or donate electrons relative to that of hydrogen if hydrogen occupied the same position in the molecule. These terms are well-understood by one skilled in the art and are discussed in Advanced Organic Chemistry by J. March, 1985, pp. 16-18, incorporated herein by reference. Electron withdrawing groups include halo, nitro, carboxyl, lower
5 alkenyl, lower alkynyl, carboxaldehyde, carboxyamido, aryl, quaternary ammonium, trifluoromethyl, sulfonyl and aryl lower alkanoyl among others. Electron donating groups include such groups as hydroxy, lower alkyl, amino, lower alkylamino, di(lower alkyl)amino, aryloxy, mercapto, lower alkylthio, lower alkylmercapto, and disulfide among others. One skilled in the art will appreciate that the aforesaid substituents may have electron donating or
10 electron withdrawing properties under different chemical conditions. Moreover, the present invention contemplates any combination of substituents selected from the above-identified groups.

The most preferred electron donating or electron withdrawing substituents are halo, nitro, alkanoyl, carboxaldehyde, arylalkanoyl, aryloxy, carboxyl, carboxamide, cyano,
15 sulfonyl, sulfoxide, heterocyclyl, guanidine, quaternary ammonium, lower alkenyl, lower alkynyl, sulfonium salts, hydroxy, lower alkoxy, lower alkyl, amino, lower alkylamino, di(lower alkyl)amino, amine lower alkyl mercapto, mercaptoalkyl, alkylthio, carboxy lower alkyl, arylalkoxy, alkanoylamino, alkanoyl(lower alkyl)amino, lower alkylsulfonylamino, arylsulfonylamino, alkylsulfonyl(lower alkyl)amino, arylsulfonyl(lower alkyl)amino, lower
20 alkylcarboxamide, di(lower alkyl)carboxamide, sulfonamide, lower alkylsulfonamide, di(lower alkyl)sulfonamide, lower alkylsulfonyl, arylsulfonyl and alkylidithio.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from a combination of the specified ingredients in the specified
25 amounts.

As used herein, the term "mammals" includes humans and other animals.

The ring defined by Y in Formulae I, II and III can be a mono-cyclic heterocycle or aromatic ring, or can be a bicyclic ring.

The dotted lines used in Formulae I, II, III, IV and VI indicate that the bond at that
30 location can be either single or double. The bond between the atoms Y and W for example can be a single or double bond if Y and/or W is a substituent such as N, C or CH. Therefore,

the ring defined by Y in the Formulae can be either saturated or unsaturated, depending upon which W and/or Y is selected. In Formulae IV and VI, the dotted line indicates that the nitrogen-containing ring optionally contains double bonds at the indicated locations.

In the Formulae, certain R groups potentially substitute their associated rings a number of times. R¹⁹, R²⁰, R²¹, R²³, R²⁷, R²⁸, R²⁹ and R²⁵ may each substitute their associated rings more than once. For example for R¹⁹, when c is zero, the associated ring is unsubstituted, having hydrogens at the C-2 and C-4 positions; and for R²³, when g is zero, hydrogens are at the C-2 – C-5 positions.

Suitable substituents for the aryl, alkyl, cycloalkyl, heterocyclyl groups or the ring defined by Y and W in the formulae described above, when present, include alcohols, amines, heteroatoms, or any combination of aryl, alkoxy, alkoxyalkoxy, alkyl, cycloalkyl or heterocyclyl groups either attached directly, or *via* suitable linkers. The linkers are typically short chains of 1-3 atoms containing any combination of C, C=O, CO₂, O, N, S, S=O, SO₂, as for example ethers, amides, amines, ureas, sulfamides, sulfonamides, among others.

For example, R¹, R², R³, R⁵, R⁶, R⁷ and R⁸ in the above formulae may independently be, but are not limited to: hydrogen, alkyl, phenyl, thienylmethyl, isobutyl, n-butyl, 2-thienylmethyl, 1,3-thiazol-2-yl-methyl, benzyl, thienyl, 3-pyridinylmethyl, 3-methyl-1-benzothiophen-2-yl, allyl, 3-methoxybenzyl, propyl, 2-ethoxyethyl, cyclopropylmethyl, benzylsulfanylmethyl, benzylsulfonylmethyl, phenylsulfanylmethyl, phenethylsulfanylmethyl, 3-phenylpropylsulfanylmethyl, 4-((2-toluidinocarbonyl)amino)benzyl, 2-pyridinylethyl, 2-(1H-indol-3-yl)ethyl, 1H-benzimidazol-2-yl, 4-piperidinylmethyl, 3-hydroxy-4-methoxybenzyl, 4-hydroxyphenethyl, 4-aminobenzyl, phenylsulfonylmethyl, 4-(acetylamino)phenyl, 4-methoxyphenyl, 4-aminophenyl, 4-chlorophenyl, (4-(benzylsulfonyl)amino)phenyl, (4-(methylsulfonyl)amino)phenyl, 2-aminophenyl, 2-methylphenyl, isopropyl, 2-oxo-1-pyrrolidinyl, 3-(methylsulfanyl)propyl, (propylsulfanyl)methyl, octylsulfanylmethyl, 3-aminophenyl, 4-((2-toluidinocarbonyl)amino)phenyl, 2-((methylbenzyl)amino)benzyl, methylsulfanylethyl, hydroxy, chloro, fluoro, bromo, ureido, amino, methanesulfonylamino, acetylamino, ethylsulfanylmethyl, 2-chlorobenzyl, 2-bromobenzyl, 2-fluorobenzyl, 2-chloro-6-fluorobenzyl, 2-chloro-4-fluorobenzyl, 2,4-dichlorobenzyl, 2-chloro-6-methoxybenzyl, 2-cyanobenzyl, 2,6-difluorobenzyl, 2-chloro-

5-(trifluoromethyl)benzyl, 2-chloro-6-methylbenzyl, 2,6-dimethoxybenzyl, 2-chloro-5-(methylsulfonyl)benzyl, 2-chloro-6-cyanobenzyl, 2-chloro-6-ethoxybenzyl, 2-chloro-5-methoxybenzyl, 2-chloro-5-fluorobenzyl, 5-chloro-2-fluorobenzyl, ethyl, propyl, butyl, pentyl, cyclopropyl, tert-butylamino, propylamino, 4-methyl-1-piperazinyl, 1-azetidiny, 4-morpholino, (4-carboxyphenyl)amino, pivaloylamino, ((tert-butylamino) carbonyl)amino, trifluoromethyl, benzyloxy, 2-(2-methoxyethoxy)ethoxy, 2-(2-(2-methoxyethoxy)ethoxy)ethoxy and 2-(2-(2-(2-ethoxyethoxy)ethoxy)ethoxy)ethoxy.

The R⁴ substituent for the formulae above may be, but is not limited to 1,3-benzodioxol-5-yl, 1-naphthyl, thienyl, 4-isobutoxyphenyl, 2,6-dimethylphenyl, allyloxyphenyl, 3-bromo-4-methoxyphenyl, 4-butoxyphenyl, 1-benzofuran-2-yl, 2-thienylmethyl, phenyl, methylsulfonyl, phenylsulfonyl, phenethylsulfonyl, 4-bromo-2-thienyl, 3-methyl-2-thienyl, 4-methylphenyl, 3,5-bis(methyloxy)phenyl, 4-(methyloxy)phenyl, 4-fluorophenyl, 3-(methyloxy)phenyl, 3,4,5-tris(methyloxy)phenyl, 2,3-dihydro-1-benzofuran-5-yl, 3-fluorophenyl, 4-(trifluoromethyl)phenyl, 4-fluoro-3-(trifluoromethyl)phenyl, 4-(1,1-dimethylethyl)phenyl, 3,5-dimethylphenyl, 4-hydroxyphenyl, 3,4-dimethylphenyl, 3-methyl-4-(methyloxy)phenyl, 4-hydroxy-3-methylphenyl, 3-methylphenyl, 2,3-dihydro-inden-5-yl, 2-methylphenyl, 2,6-bis(methyloxy)phenyl, 2,6-dihydroxyphenyl, 4-chlorophenyl, 3-chlorophenyl, 3,4-dichlorophenyl, 4-((trifluoromethyl)oxy)phenyl, 4-ethylphenyl, 4-(ethyloxy)phenyl, methyl, 2-propyl, 4,5-dihydro-1,3-oxazol-2-yl, 3-(trifluoromethyl)phenyl, 4-(trifluoromethoxy)phenyl, 2,3-dihydro-1,4-benzodioxin-6-yl, 7-methoxy-1,3-benzodioxol-5-yl, 3-ethoxy-4-methoxyphenyl, 3,4-dimethoxyphenyl, 3,4-diethoxyphenyl, 3-ethoxyphenyl, 3-methoxy-4-methylphenyl, 3,5-dimethoxy-4-methylphenyl, 3-propoxyphenyl, 3-butoxyphenyl, 3-(2-methoxyethoxy)phenyl, 3,4-dipropoxyphenyl, 3-(difluoromethoxy)phenyl, 2-naphthyl, 3-isopropoxyphenyl, 1-methyl-1H-indol-5-yl, 2,3-dihydro-1-benzofuran-5-yl, 1,3-diethyl-2-oxo-2,3-dihydro-1H-benzimidazol-5-yl, 3-(trifluoromethoxy)phenyl, 1-methyl-1H-indol-6-yl, 3-(cyclopropoxy)phenyl, 3-(cyclopropylmethoxy)phenyl, 3-(difluoromethoxy)phenyl, 3-(1,1,2,2-tetrafluoroethoxy)phenyl, 1-ethyl-1H-indol-5-yl, 3-(diethylamino)phenyl, 6-methoxy-2-naphthyl, 3-[(methylsulfonyl)amino]phenyl, 3-[methyl(methylsulfonyl)amino]phenyl, 3-

[ethyl(methylsulfonyl)amino]phenyl, 1H-indol-5-yl, 3-fluoro-4-methoxyphenyl and 3-(difluoromethyl)phenyl.

Two independent R¹, R², R³ or R⁵ groups taken together may be linked to form a ring,

5 R⁴ and R¹¹ may be linked to form a ring such as 1-pyrrolidino, 1-piperidino, 4-methyl-1-piperazino, 4-acetyl-1-piperazino and 4-morpholino among others.

R⁹ and R¹⁰ may be linked to form a ring such as cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl among others.

Representative disease targets of the compositions of the present invention include
10 inflammatory diseases such as psoriasis, asthma, atherosclerosis, autoimmune diseases such as multiple sclerosis and Guillan-Barr Syndrome, rheumatoid arthritis, transplant and graft v. host disease, inflammatory bowel disease, chronic obstructive pulmonary disease and reperfusion injury.

One or more $\alpha_4\beta_1$ binding (VLA-4) inhibitors may be combined with one or more
15 other therapeutic compounds. The other therapeutic compounds include compounds for treating an inflammatory disease, i.e., inflammatory response, as, for example, IL-5 antagonists for growth, maturation and survival of eosinophils; CCR-3 antagonists for chemotaxis of eosinophils; corticosteroids for general suppression of inflammation; antihistamines for histamine early phase blockages; Leukotriene antagonists for LTD
20 bronchoconstrictor and LTB eosinophil chemotaxis; COX-I and COX-II inhibitors for prostaglandin production; mast cell stabilizers such as Chromolyn; anti-IL-5 or anti-IgE; IL-5 synthesis and release inhibitors; selectin antagonists, CD20 antagonists for suppression of B-cell mediated inflammation and syk tyrosine kinase inhibitors.

Agents known in the treatment of rheumatoid arthritis, transplant and graft v. host
25 disease, inflammatory bowel disease, chronic obstructive pulmonary disease and multiple sclerosis which can be administered in combination with the $\alpha_4\beta_1$ binding inhibitors of the present invention are as follows:

solid organ transplant rejection and graft v. host disease: immune suppressants
such as cyclosporine rapamycin and Interleukin-10 (IL-10), tacrolimus, antilymphocyte
30 globulin, OKT-3 antibody, and steroids;

-26-

inflammatory bowel disease: IL-10 (see U.S. Patent No. 5,368,854), steroids and azulfidine;

rheumatoid arthritis: methotrexate, azathioprine, cyclophosphamide, steroids and mycophenolate mofetil;

5 multiple sclerosis: interferon-beta, interferon-alpha, and steroids.

Non limitative examples of antihistamines include: astemizole, azatadine, azelastine, acrivastine, brompheniramine, cetirizine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine, descarboethoxyloratadine (also known as SCH-34117), doxylamine, dimethindene, ebastine, epinastine, efenitidine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine, mizolastine, equitazine, mianserin, noberastine, meclizine, norasteremizole, picumast, pyrilamine, promethazine, 10 terfenadine, tripeleminamine, temelastine, trimeprazine and triprolidine.

The term "leukotriene antagonist" includes any agent or compound that inhibits, restrains, retards or otherwise interacts with the action or activity of leukotrienes. Non-
15 limitative examples of leukotriene inhibitors include montelukast [R-(E)]-1[[[1-[3-[2-(7-chloro-2-quinolinyl)-ethenyl] phenyl]-3[2-(1-hydroxy-1-methylethyl)phenyl]propyl]thio]methyl]cyclo-propaneacetic acid and its sodium salt, described in EP 0 480 717; 1-(((R)-(3-(2-(6,7-difluoro-2-quinolinyl)ethenyl)phenyl)-3-(2-(2-hydroxy-2-propyl)phenyl)thio) methyl)cyclopropaneacetic acid, and its sodium salt, described in WO 97/28797 and U.S.
20 Patent 5,270,324; 1-(((1(R)-3(3-(2-(2,3-dichlorothieno[3,2-b]pyridin-5-yl)-(E)-ethenyl) phenyl)-3-(2-(1-hydroxy-1-methylethyl)phenyl)propyl)thio)methyl)cyclopropaneacetic acid, and its sodium salt, described in WO 97/28797 and U.S. Patent 5,472,964; pranlukast, N-[4-oxo-2-(H-tetrazol-5-yl)-4H-1-benzopyran-8-yl]-p-(4-phenylbutoxy) benzamide) described in WO 97/28797 and EP 173,516; zafirlukast, (cyclopentyl-3-[2-methoxy-4-[(o-tolylsulfonyl) carbamoyl]benzyl]-1-methylindole-5-carbamate) described
25 in WO 97/28797 and EP 199,543; and [2-[[2(4-*tert*-butyl-2-thiazolyl)-5-benzofuranyl]oxymethyl]phenyl]acetic acid, described in U.S. Patent 5,296,495 and Japanese patent JP08325265A.

Other therapeutic agents that may be used in conjunction with the compounds that
30 inhibit binding of $\alpha_4\beta_1$ integrin to its receptors include:

-27-

- β -agonists: albuterol, salmeterol, formoterol, levabuterol, terbutaline, pirbuterol, metaprotrenol
- muscarinic antagonists: ipratropium bromide, tiotropium bromide
- PDE 4 inhibitors: roflumilast, theophylline, rolipram, piclamilast, cilomilast,
- 5 CDP-840
- immunosuppressives: methotrexate, leflunomide, sulfasalazine, cyclosporin
- steroids: prednisolone, fluticasone, triamcinolone, beclomethasone, mometasone, budisamide, betamethasone, dexamethasone, prednisone, flunisolide, cortisone
- COX-I inhibitors: aspirin, piroxicam
- 10 COX-II inhibitors: rofecoxib, celecoxib, valdecoxib, etoricoxib
- TNF- α inhibitor: infliximab, etanercept
- Anti-IgE antibody: omalizumab
- p38 MAP kinase inhibitor: VX-745, BIRB-796
- tryptase inhibitor: BMS-262084
- 15 anticytokine/antichemokine: RO-3202947, UCB-35625
- vaccine: Allervax Cat
- cromolyn: oral cromolyn formulation from Emisphere Technologies
- Anti-CD20 antibody: rituximab
- syk tyrosine kinase inhibitor: R-112
- 20 Actual dosage levels of active ingredients in the pharmaceutical compositions of this invention can be varied so as to obtain an amount of the active compounds which are effective to achieve the desired therapeutic response for a particular patient, compositions and mode of administration. The selected dosage level will depend upon the activity of the particular compounds, the route of administration, the severity of the condition being
- 25 treated and the condition and prior medical history of the patient being treated. However, it is within the skill of the art to start doses of the compound at levels lower than required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.
- The phrase "therapeutically effective amount" of the composition of the invention
- 30 means a sufficient amount of the active compounds to treat disorders, at a reasonable benefit/risk ratio applicable to any medical treatment. It will be understood, however,

-28-

that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder;

5 activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts. For example, it is well within

10 the skill of the art to start doses of the compound at levels lower than required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

The total daily dose of the compositions of this invention administered to a human or lower animal may range from about 0.0001 to about 1000 mg/kg/day. For purposes of

15 oral administration, more preferable doses can be in the range of from about 0.001 to about 5 mg/kg/day. If desired, the effective daily dose can be divided into multiple doses for purposes of administration; consequently, single dose compositions may contain such amounts or submultiples thereof to make up the daily dose.

The pharmaceutical compositions of this invention can be administered to humans

20 and other mammals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, buccally or as an oral or nasal spray. The term "parenterally," as used herein, refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

The present invention includes one or more compounds as described above

25 formulated into compositions together with one or more non-toxic physiologically tolerable or acceptable diluents, carriers, adjuvants or vehicles that are collectively referred to herein as diluents, for parenteral injection, for intranasal delivery, for oral administration in solid or liquid form, for rectal or topical administration, among others.

The compositions can also be delivered through a catheter for local delivery at a

30 target site, *via* an intracoronary stent (a tubular device composed of a fine wire mesh), or

via a biodegradable polymer. The compounds may also be complexed to ligands, such as antibodies, for targeted delivery.

Compositions suitable for parenteral injection may comprise physiologically acceptable, sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions and sterile powders for reconstitution into sterile injectable solutions or dispersions. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (propyleneglycol, polyethyleneglycol, glycerol, and the like), vegetable oils (such as olive oil), injectable organic esters such as ethyl oleate, and suitable mixtures thereof.

These compositions can also contain adjuvants such as preserving, wetting, emulsifying, and dispensing agents. Prevention of the action of microorganisms can be ensured by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, for example sugars, sodium chloride and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin.

Suspensions, in addition to the active compounds, may contain suspending agents, as for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, or mixtures of these substances, and the like.

In some cases, in order to prolong the effect of the drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This can be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microcapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include

poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

Solid dosage forms for oral administration include capsules, tablets, pills, powders and granules. In such solid dosage forms, the active compound may be mixed with at least one inert, pharmaceutically acceptable excipient or carrier, such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol and silicic acid; b) binders such as carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose and acacia; c) humectants such as glycerol; d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates and sodium carbonate; e) solution retarding agents such as paraffin; f) absorption accelerators such as quaternary ammonium compounds; g) wetting agents such as cetyl alcohol and glycerol monostearate; h) absorbents such as kaolin and bentonite clay and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

The solid dosage forms of tablets, dragees, capsules, pills and granules can be prepared with coatings and shells such as enteric coatings and other coatings well-known in the pharmaceutical formulating art. They may optionally contain opacifying agents and may also be of a composition such that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

The active compounds can also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned excipients.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups and elixirs. In addition to the active
5 compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethyl formamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol,
10 tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan and mixtures thereof.

Besides inert diluents, the oral compositions may also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring and perfuming agents.

15 Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at room temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

20 Compounds useful in the compositions of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals which are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming
25 liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients and the like. The preferred lipids are natural and synthetic phospholipids and phosphatidyl cholines (lecithins) used separately or together.

30 Methods to form liposomes are known in the art. See, for example, Prescott, Ed., Methods in Cell Biology, Volume XIV, Academic Press, New York, N.Y. (1976), p. 33 *et seq.*

The term "pharmaceutically acceptable prodrugs" as used herein represents those prodrugs of the compounds of the present invention which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use, as well as the zwitterionic forms, where possible, of the compounds of the invention. Prodrugs of the present invention may be rapidly transformed *in vivo* to the parent compound of the above formula, for example, by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, Pro-drugs as Novel Delivery Systems, V. 14 of the A.C.S. Symposium Series, and in Edward B. Roche, ed., Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press (1987), hereby incorporated by reference.

Compounds that are useful in the compositions of the present invention that are formed by *in vivo* conversion of a different compound that was administered to a mammal are intended to be included within the scope of the present invention.

Compounds that are useful in the compositions of the present invention may exist as stereoisomers wherein asymmetric or chiral centers are present. These stereoisomers are "R" or "S" depending on the configuration of substituents around the chiral carbon atom. The present invention contemplates various stereoisomers and mixtures thereof. Stereoisomers include enantiomers and diastereomers, and mixtures of enantiomers or diastereomers. Individual stereoisomers of compounds may be prepared synthetically from commercially available starting materials which contain asymmetric or chiral centers or by preparation of racemic mixtures followed by resolution well-known to those of ordinary skill in the art. These methods of resolution are exemplified by (1) attachment of a mixture of enantiomers to a chiral auxiliary, separation of the resulting mixture of diastereomers by recrystallization or chromatography and liberation of the optically pure product from the auxiliary or (2) direct separation of the mixture of optical enantiomers on chiral chromatographic columns.

Compounds useful in the compositions of the invention can exist in unsolvated as well as solvated forms, including hydrated forms, such as hemi-hydrates. In general, the

-33-


solvated forms, with pharmaceutically acceptable solvents such as water and ethanol among others are equivalent to the unsolvated forms for the purposes of the invention.

A procedure in which a 26-amino acid peptide containing the CS 1 sequence of fibronectin with an N-terminal Cys (CDELPQLVTLPHPNLHGPEELDVPST) may be
5 coupled to maleimide activated ovalbumin was used to determine the efficacy of the compounds synthesized. Bovine serum albumin (BSA) and CS 1 conjugated ovalbumin may be coated onto 96-well polystyrene plates at 0.5 $\mu\text{g}/\text{ml}$ in TBS (50 mM TRIS, pH 7.5; 150 mM NaCl) at 4°C for 16 hours. The plates may be washed three times with TBS and blocked with TBS containing 3% BSA at room temperature for 4 hours. Blocked plates
10 may be marked three times in binding buffer (TBS; 1 mM MgCl_2 ; 1 mM CaCl_2 ; 1 mM MnCl_2) prior to assay. Ramos cells fluorescently labeled with calcein AM may be resuspended in binding buffer (10⁷ cells/ml) and diluted 1:2 with same buffer with or without compound. 100 pM of compound should be added. The cells should be added immediately to the wells (2.5 x 10⁵ cells/ well) and incubated for 30 minutes at 37°C.
15 Following three washes with binding buffer, adherent cells should be lysed and quantitated using a fluorometer. IC₅₀ is defined as the dose required to give 50% inhibition. The lower the IC₅₀ value and the greater the percentage of inhibition, the more efficient the compound is at prevention of cell adhesion.

Claims

We claim:

1. A pharmaceutical composition comprising a compound of the structure (I)



5

(I)

wherein Y, at each occurrence, is independently selected from the group consisting of C(O), N, CR¹, C(R²)(R³), NR⁵, CH, O and S;

q is an integer of from 3 to 10;

10 A is selected from the group consisting of O, S, C(R¹⁶)(R¹⁷) and NR⁶;

E is selected from the group consisting of CH₂, O, S, and NR⁷;

J is selected from the group consisting of O, S and NR⁸;

15 T is selected from the group consisting of C(O) and (CH₂)_b wherein b is an integer of from 0 to 3;

M is selected from the group consisting of C(R⁹)(R¹⁰) and (CH₂)_u, wherein u is an integer of from 0 to 3;

20 L is selected from the group consisting of O, NR¹¹, S, and (CH₂)_n wherein n is an integer of 0 or 1;

X is selected from the group consisting of CO₂B, PO₃H₂, SO₃H, SO₂NH₂, SO₂NHCOR¹², OPO₃H₂, C(O)NHC(O)R¹³, C(O)NHSO₂R¹⁴, hydroxyl, tetrazolyl and hydrogen;

W is selected from the group consisting of C, CR¹⁵ and N;

25 B is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, hydroxyalkyl, haloalkyl, -CF₃, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, heterocyclyl, alkylaryl, aralkenyl, aralkyl,

-35-


alkylheterocyclyl, heterocyclylalkyl and aryloxyalkyl; and
 $R^1, R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}$ and R^{17} at each
occurrence are independently selected from the group consisting of
hydrogen, halogen, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkynoxy,
5 thioalkoxy, hydroxyalkyl, aliphatic acyl, $-CF_3$, $-CO_2H$, $-SH$,
 $-CN$, $-NO_2$, $-NH_2$, $-OH$, alkynylamino, alkoxycarbonyl, heterocycloyl,
carboxy, $-N(C_1-C_3 \text{ alkyl})-C(O)(C_1-C_3 \text{ alkyl})$, $-NHC(O)N(C_1-C_3$
alkyl) $C(O)NH(C_1-C_3 \text{ alkyl})$, $-NHC(O)NH(C_1-C_6 \text{ alkyl})$, $-NHSO_2(C_1-C_3$
alkyl), $-NHSO_2(\text{aryl})$, alkoxyalkyl, alkylamino, alkenylamino, di(C_1-
10 C_3)amino, $-C(O)O-(C_1-C_3) \text{ alkyl}$, $-C(O)NH-(C_1-C_3) \text{ alkyl}$, $-C(O)N(C_1-C_3$
alkyl) $_2$, $-CH=NOH$, $-PO_3H_2$, $-OPO_3H_2$, haloalkyl, alkoxyalkoxy,
carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl,
cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl,
diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl,
15 heterocyclylalkyl, sulfonyl, $-SO_2-(C_1-C_3 \text{ alkyl})$, $-SO_3-(C_1-C_3 \text{ alkyl})$,
sulfonamido, carbamate, aryloxyalkyl and $-C(O)NH(\text{benzyl})$ groups;
wherein $B, R^1, R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13},$
 R^{14}, R^{15}, R^{16} and R^{17} are unsubstituted or substituted with at least
one electron donating or electron withdrawing group;
20 wherein when L is NR^{11}, R^4 and R^{11} taken together may form a ring;
and wherein when M is $C(R^9)(R^{10}), R^9$ and R^{10} taken together may
form a ring;
and wherein when A is NR^6 and at least one Y is CR^1, R^1 and R^6
taken together may form a ring;
25 or a pharmaceutically acceptable salt thereof;
one or more other therapeutically active compounds and a pharmacologically
acceptable diluent.

2. A composition of claim 1 wherein
30 A is NR^6 ;
E is NR^7 ;

-36-

- J is O;
 M is $C(R^9)(R^{10})$;
 q is 4 or 5;
 T is $(CH_2)_b$ wherein b is 0;
 5 L is $(CH_2)_n$ wherein n is 0;
 X is CO_2B ;
 W is C or CR^{15} ;
 R^4 is selected from the group consisting of aryl, alkylaryl, aralkyl,
 heterocyclyl, alkylheterocyclyl and heterocyclylalkyl; and
 10 R^6, R^7, R^9, R^{10} and R^{15} are independently selected from the
 group consisting of hydrogen and lower alkyl.

3. A pharmaceutical composition comprising a compound of the structure



- 15 wherein Y, at each occurrence, is independently selected from the group
 consisting of $C(O)$, N, CR^1 , $C(R^2)(R^3)$, NR^5 , CH, O and S;
 q is an integer of from 3 to 7;
 T is selected from the group consisting of $C(O)$ and $(CH_2)_b$ wherein b is an
 integer of 0 to 3;
 20 L is selected from the group consisting of O, NR^{11} , S, and
 $(CH_2)_n$ wherein n is an integer of 0 or 1;
 W is selected from the group consisting of C, CR^{15} and N;
 B is selected from the group consisting of hydrogen, alkyl, alkenyl,
 alkynyl, hydroxyalkyl, haloalkyl, $-CF_3$, cycloalkyl, cycloalkenyl,

-37-

cycloalkynyl, cycloalkylalkyl, aryl, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl and aryloxyalkyl; and

$R^1, R^2, R^3, R^4, R^5, R^6, R^7, R^9, R^{10}, R^{11}$ and R^{15} are independently selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkynoxy, thioalkoxy, hydroxyalkyl, aliphatic acyl, $-CF_3$, $-CO_2H$, $-SH$, $-CN$, $-NO_2$, $-NH_2$, $-OH$, alkynylamino, alkoxy-carbonyl, heterocycloyl, carboxy, $-N(C_1-C_3 \text{ alkyl})-C(O)(C_1-C_3 \text{ alkyl})$, $-NHC(O)N(C_1-C_3 \text{ alkyl})C(O)NH(C_1-C_3 \text{ alkyl})$, $-NHC(O)NH(C_1-C_6 \text{ alkyl})$, $-NHSO_2(C_1-C_3 \text{ alkyl})$, $-NHSO_2(\text{aryl})$, alkoxyalkyl, alkylamino, alkenylamino, di(C_1-C_3)amino, $-C(O)O-(C_1-C_3 \text{ alkyl})$, $-C(O)NH-(C_1-C_3 \text{ alkyl})$, $-C(O)N(C_1-C_3 \text{ alkyl})_2$, $-CH=NOH$, $-PO_3H_2$, $-OPO_3H_2$, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl, $-SO_2-(C_1-C_3 \text{ alkyl})$, $-SO_3-(C_1-C_3 \text{ alkyl})$, sulfonamido, carbamate, aryloxyalkyl and $-C(O)NH(\text{benzyl})$ groups;

wherein B, $R^1, R^2, R^3, R^4, R^5, R^6, R^7, R^9, R^{10}, R^{11}$ and R^{15} are

unsubstituted or substituted with at least one electron donating or electron withdrawing group;

wherein when L is NR^{11} , R^4 and R^{11} taken together may form a ring;

and wherein R^9 and R^{10} taken together may form a ring;

and wherein when at least one Y is CR^1 , R^1 and R^6 taken together may form a ring;

or a pharmaceutically acceptable salt thereof;

one or more other therapeutically active compounds and a pharmacologically acceptable diluent.

4. A composition of claim 3 wherein

q is 4 or 5;

W is C or CR^{15} ;

-38-


T is $(\text{CH}_2)_b$ wherein b is 0;

L is $(\text{CH}_2)_n$ wherein n is 0;

R^4 is selected from the group consisting of aryl, alkylaryl, aralkyl, heterocyclyl, alkylheterocyclyl and heterocyclylalkyl; and

5 R^6, R^7, R^9, R^{10} and R^{15} are independently selected from the group consisting of hydrogen and lower alkyl.

5. A pharmaceutical composition comprising a compound of the structure



10

wherein Y, at each occurrence, is independently selected from the group consisting of $\text{C}(\text{O})$, N , CR^1 , $\text{C}(\text{R}^2)(\text{R}^3)$, NR^5 , CH , O and S ;

q is an integer of from 2 to 5;

15 T is selected from the group consisting of $\text{C}(\text{O})$ and $(\text{CH}_2)_b$ wherein b is an integer of 0 to 3;

L is selected from the group consisting of O , NR^{11} , S , and

$(\text{CH}_2)_n$ wherein n is an integer of 0 or 1;

R^5 , R^6 , R^7 , R^{11} and R^{18} are each independently selected from the group consisting of alkyl, alkenyl, alkynyl, hydroxyalkyl, aliphatic acyl, alkynylamino, alkoxyacetyl, heterocycloyl, $-\text{CH}=\text{NOH}$, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, carbamate, aryloxyalkyl, hydrogen and $-\text{C}(\text{O})\text{NH}(\text{benzyl})$ groups;

B is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, hydroxyalkyl, haloalkyl, $-\text{CF}_3$, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl and aryloxyalkyl; and

R^1 , R^2 , R^3 , R^4 , R^9 and R^{10} are independently selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkynoxy, thioalkoxy, hydroxyalkyl, aliphatic acyl, $-\text{CF}_3$, $-\text{CO}_2\text{H}$, $-\text{SH}$, $-\text{CN}$, $-\text{NO}_2$, $-\text{NH}_2$, $-\text{OH}$, alkynylamino, alkoxyacetyl, heterocycloyl, carboxy, $-\text{N}(\text{C}_1\text{-C}_3 \text{ alkyl})-\text{C}(\text{O})(\text{C}_1\text{-C}_3 \text{ alkyl})$, $-\text{NHC}(\text{O})\text{N}(\text{C}_1\text{-C}_3 \text{ alkyl})\text{C}(\text{O})\text{NH}(\text{C}_1\text{-C}_3 \text{ alkyl})$, $-\text{NHC}(\text{O})\text{NH}(\text{C}_1\text{-C}_6 \text{ alkyl})$, $-\text{NHSO}_2(\text{C}_1\text{-C}_3 \text{ alkyl})$, $-\text{NHSO}_2(\text{aryl})$, alkoxyalkyl, alkylamino, alkenylamino, $\text{di}(\text{C}_1\text{-C}_3)\text{amino}$, $-\text{C}(\text{O})\text{O}-(\text{C}_1\text{-C}_3)\text{alkyl}$, $-\text{C}(\text{O})\text{NH}-(\text{C}_1\text{-C}_3)\text{alkyl}$, $-\text{C}(\text{O})\text{N}(\text{C}_1\text{-C}_3 \text{ alkyl})_2$, $-\text{CH}=\text{NOH}$, $-\text{PO}_3\text{H}_2$, $-\text{OPO}_3\text{H}_2$, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl, $-\text{SO}_2-(\text{C}_1\text{-C}_3 \text{ alkyl})$, $-\text{SO}_3-(\text{C}_1\text{-C}_3 \text{ alkyl})$, sulfonamido, carbamate, aryloxyalkyl and $-\text{C}(\text{O})\text{NH}(\text{benzyl})$ groups;

wherein B , R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^9 , R^{10} , R^{11} and R^{18} are

unsubstituted or substituted with at least one electron donating or electron withdrawing group;

wherein when L is NR^{11} , R^4 and R^{11} taken together may form a ring;

and wherein R^9 and R^{10} taken together may form a ring;

and wherein when at least one Y is CR¹, R¹ and R⁶ taken together may form a ring;

or a pharmaceutically acceptable salt thereof, one or more other therapeutically active compounds and a pharmacologically acceptable diluent.

5

6. A composition of claim 5 wherein R¹⁸ is selected from the group consisting of hydrogen, alkyl, aryl, aralkyl, cycloalkyl, alkylheterocyclyl, heterocyclylalkyl and heterocyclyl;

T is (CH₂)_b wherein b is 0;


10

L is (CH₂)_n wherein n is 0;

Y is selected from the group consisting of CR¹ and C(R²)(R³) and



q is 2 or 3.

7. A composition of claim 5 wherein




15

is selected from the group consisting of



,



and

;

- wherein R¹⁹, R²⁰, R²¹ and R²⁸ at each occurrence are independently selected from
- 5 the group consisting of halogen, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkynoxy, thioalkoxy, hydroxyalkyl, aliphatic acyl, -CF₃, -OH, -CO₂H, -SH, -CN, -NO₂, -NH₂, alkynylamino, alkoxy carbonyl, heterocycloyl, carboxy, -N(C₁-C₃ alkyl)-C(O)(C₁-C₃ alkyl), -NHC(O)N(C₁-C₃ alkyl)C(O)NH(C₁-C₃alkyl), -NHC(O)NH(C₁-C₆ alkyl),
- 10 -NHSO₂(C₁-C₃ alkyl), -NHSO₂(aryl), alkoxyalkyl, alkylamino, alkenylamino, di(C₁-C₃)amino, -C(O)O-(C₁-C₃)alkyl, -C(O)NH-(C₁-C₃)alkyl, -C(O)N(C₁-C₃alkyl)₂, -CH=NOH, -PO₃H₂, -OPO₃H₂, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino,
- 15 heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl, -SO₂-(C₁-C₃ alkyl), -SO₃-(C₁-C₃ alkyl), sulfonamido, carbamate, aryloxyalkyl and -C(O)NH(benzyl) groups;
- R¹⁸ is selected from the group consisting of alkyl, alkenyl, alkynyl, hydroxyalkyl, aliphatic acyl, alkynylamino, alkoxy carbonyl, heterocycloyl, -CH=NOH,
- 20 haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, carbamate, aryloxyalkyl, hydrogen and -C(O)NH(benzyl) groups;
- 25 R²² is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkynoxy, thioalkoxy, hydroxyalkyl, aliphatic acyl, -CF₃, -CO₂H, -SH, -CN, -NO₂, -NH₂, -OH, alkynylamino, alkoxy carbonyl, heterocycloyl, carboxy, -N(C₁-C₃ alkyl)-C(O)(C₁-C₃ alkyl), -NHC(O)N(C₁-C₃ alkyl)C(O)NH(C₁-C₃alkyl), -NHC(O)NH(C₁-C₆ alkyl),
- 30 -NHSO₂(C₁-C₃ alkyl), -NHSO₂(aryl), alkoxyalkyl, alkylamino, alkenylamino, di(C₁-C₃)amino, -C(O)O-(C₁-C₃)alkyl,

-42-

5
 -C(O)NH-(C₁-C₃)alkyl, -C(O)N(C₁-C₃ alkyl)₂, -CH=NOH, -PO₃H₂,
 -OPO₃H₂, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide,
 cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy,
 arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl,
 10 aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl, -SO₂-(C₁-C₃ alkyl),
 -SO₃-(C₁-C₃ alkyl), sulfonamido, carbamate, aryloxyalkyl and
 -C(O)NH(benzyl) groups;

c is an integer of zero to two;

d is an integer of zero to three;

10 e is an integer of zero to four; and

i is an integer of zero to two.

8. The composition of claim 5 wherein R¹⁸ is aralkyl;

R⁴ is aryl;


15 T is (CH₂)_b where b is zero;

L is (CH₂)_n where n is zero; and,

B, R⁶, R⁷, R⁹ and R¹⁰ are each independently hydrogen.

9. A pharmaceutical composition comprising a compound of the structure

20



wherein T is selected from the group consisting of C(O) and (CH₂)_b wherein b is an integer of from 0 to 3;

L is selected from the group consisting of O, NR¹¹, S, and

-43-

$(\text{CH}_2)_n$ wherein n is an integer of 0 or 1;

g is an integer of from 0 to 7;

B is selected from the group consisting of hydrogen, alkyl, alkenyl,

alkynyl, hydroxyalkyl, haloalkyl, $-\text{CF}_3$, cycloalkyl, cycloalkenyl,

5 cycloalkynyl, cycloalkylalkyl, aryl, heterocyclyl, alkylaryl, aralkenyl, aralkyl,

alkylheterocyclyl, heterocyclylalkyl and aryloxyalkyl;

R^4 , R^9 , R^{10} and R^{23} at each occurrence are independently selected from the group

consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy,

alkynoxy, thioalkoxy, hydroxyalkyl, aliphatic acyl,

10 $-\text{CF}_3$, $-\text{CO}_2\text{H}$, $-\text{SH}$, $-\text{CN}$, $-\text{NO}_2$, $-\text{NH}_2$, $-\text{OH}$, alkynylamino,

alkoxycarbonyl, heterocycloyl, carboxy, $-\text{N}(\text{C}_1\text{-C}_3 \text{ alkyl})\text{-C}(\text{O})(\text{C}_1\text{-C}_3$

alkyl), $-\text{NHC}(\text{O})\text{N}(\text{C}_1\text{-C}_3 \text{ alkyl})\text{C}(\text{O})\text{NH}(\text{C}_1\text{-C}_3\text{alkyl})$, $-\text{NHC}(\text{O})\text{NH}(\text{C}_1\text{-C}_6$

alkyl), $-\text{NHSO}_2(\text{C}_1\text{-C}_3 \text{ alkyl})$, $-\text{NHSO}_2(\text{aryl})$, alkoxyalkyl, alkylamino,

alkenylamino, $\text{di}(\text{C}_1\text{-C}_3)\text{amino}$, $-\text{C}(\text{O})\text{O}(\text{C}_1\text{-C}_3)\text{alkyl}$,

15 $-\text{C}(\text{O})\text{NH}(\text{C}_1\text{-C}_3)\text{alkyl}$, $-\text{C}(\text{O})\text{N}(\text{C}_1\text{-C}_3 \text{ alkyl})_2$, $-\text{CH}=\text{NOH}$, $-\text{PO}_3\text{H}_2$,

$-\text{OPO}_3\text{H}_2$, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide,

cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl,

aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl,

aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl, $-\text{SO}_2(\text{C}_1\text{-}$

20 $\text{C}_3 \text{ alkyl})$, $-\text{SO}_3(\text{C}_1\text{-C}_3 \text{ alkyl})$, sulfonamido, carbamate, aryloxyalkyl and

$-\text{C}(\text{O})\text{NH}(\text{benzyl})$ groups;

R^6 , R^7 , R^{11} and R^{18} are each independently selected from the group consisting of alkyl,

alkenyl, alkynyl, hydroxyalkyl, aliphatic acyl, alkynylamino, alkoxycarbonyl,

heterocycloyl, $-\text{CH}=\text{NOH}$, haloalkyl, alkoxyalkoxy, carboxaldehyde,

25 carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl,

aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl,

aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, carbamate,

aryloxyalkyl, hydrogen and $-\text{C}(\text{O})\text{NH}(\text{benzyl})$ groups;

wherein B, R^4 , R^6 , R^7 , R^9 , R^{10} , R^{11} , R^{18} and R^{23} are unsubstituted or substituted

30 with at least one electron donating or electron withdrawing group;


wherein when L is NR^{11} , R^4 and R^{11} taken together may form a ring;

-44-

and wherein R^9 and R^{10} taken together may form a ring;
 or a pharmaceutically acceptable salt thereof;
 one or more other therapeutically active compounds and a pharmacologically
 acceptable diluent.

5

10. A pharmaceutical composition comprising a compound of the structure



10

wherein h is an integer of zero to five;

B is selected from the group consisting of hydrogen, alkyl, alkenyl,
 alkynyl, hydroxyalkyl, haloalkyl, $-CF_3$, cycloalkyl, cycloalkenyl,
 cycloalkynyl, cycloalkylalkyl, aryl, heterocyclyl, alkylaryl, aralkenyl, aralkyl,
 alkylheterocyclyl, heterocyclylalkyl and aryloxyalkyl;

15

R^9 , R^{10} , R^{24} and R^{25} are each independently selected from the group consisting of
 hydrogen, halogen, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkynoxy,
 thioalkoxy, hydroxyalkyl, aliphatic acyl,

20

$-CF_3$, $-CO_2H$, $-SH$, $-CN$, $-NO_2$, $-NH_2$, $-OH$, alkynylamino, alkoxy carbonyl,
 heterocycloyl, carboxy, $-N(C_1-C_3 \text{ alkyl})-C(O)(C_1-C_3 \text{ alkyl})$,
 $-NHC(O)N(C_1-C_3 \text{ alkyl})C(O)NH(C_1-C_3 \text{ alkyl})$, $-NHC(O)NH(C_1-C_6 \text{ alkyl})$,
 $-NHSO_2(C_1-C_3 \text{ alkyl})$, $-NHSO_2(\text{aryl})$, alkoxyalkyl, alkylamino,
 alkenylamino, $di(C_1-C_3 \text{ amino})$, $-C(O)O-(C_1-C_3 \text{ alkyl})$, $-C(O)NH-(C_1-$

25

$C_3 \text{ alkyl})$, $-C(O)N(C_1-C_3 \text{ alkyl})_2$, $-CH=NOH$,
 $-PO_3H_2$, $-OPO_3H_2$, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide,
 cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl,
 aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl,

-45-

aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl, $-\text{SO}_2-(\text{C}_1-\text{C}_3 \text{ alkyl})$, $-\text{SO}_3-(\text{C}_1-\text{C}_3 \text{ alkyl})$, sulfonamido, carbamate, aryloxyalkyl and $-\text{C}(\text{O})\text{NH}(\text{benzyl})$ groups;

R^{27} , at each occurrence, is independently selected from the group consisting of

5 halogen, hydroxyl, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkynoxy, thioalkoxy, hydroxyalkyl, aliphatic acyl, $-\text{CF}_3$, $-\text{CO}_2\text{H}$, $-\text{SH}$, $-\text{CN}$, $-\text{NO}_2$, $-\text{NH}_2$, alkynylamino, alkoxycarbonyl, heterocycloyl, carboxy, $-\text{N}(\text{C}_1-\text{C}_3 \text{ alkyl})-\text{C}(\text{O})(\text{C}_1-\text{C}_3 \text{ alkyl})$, $-\text{NHC}(\text{O})\text{N}(\text{C}_1-\text{C}_3 \text{ alkyl})\text{C}(\text{O})\text{NH}(\text{C}_1-\text{C}_3 \text{ alkyl})$, $-\text{NHC}(\text{O})\text{NH}(\text{C}_1-\text{C}_6 \text{ alkyl})$, $-\text{NHSO}_2(\text{C}_1-\text{C}_3 \text{ alkyl})$, $-\text{NHSO}_2(\text{aryl})$, $-\text{N}(\text{C}_1-\text{C}_3 \text{ alkyl})\text{SO}_2(\text{C}_1-\text{C}_3 \text{ alkyl})$, $-\text{N}(\text{C}_1-\text{C}_3 \text{ alkyl})\text{SO}_2(\text{aryl})$, alkoxyalkyl, alkylamino, alkenylamino, di(C_1-C_3)amino, $-\text{C}(\text{O})\text{O}-(\text{C}_1-\text{C}_3)\text{alkyl}$, $-\text{C}(\text{O})\text{NH}-(\text{C}_1-\text{C}_3)\text{alkyl}$, $-\text{C}(\text{O})\text{N}(\text{C}_1-\text{C}_3 \text{ alkyl})_2$, $-\text{CH}=\text{NOH}$, $-\text{PO}_3\text{H}_2$, $-\text{OPO}_3\text{H}_2$, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide, 10 cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl, $-\text{SO}_2-(\text{C}_1-\text{C}_3 \text{ alkyl})$, $-\text{SO}_3-(\text{C}_1-\text{C}_3 \text{ alkyl})$, sulfonamido, carbamate, aryloxyalkyl and $-\text{C}(\text{O})\text{NH}(\text{benzyl})$ groups;

20 R^6 , R^7 and R^{18} are each independently selected from the group consisting of alkyl, alkenyl, alkynyl, hydroxyalkyl, aliphatic acyl, alkynylamino, alkoxycarbonyl, heterocycloyl, $-\text{CH}=\text{NOH}$, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, 25 heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, carbamate, aryloxyalkyl, hydrogen and $-\text{C}(\text{O})\text{NH}(\text{benzyl})$ groups; and,

R^{26} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, hydroxyalkyl, aliphatic acyl, $-\text{CF}_3$, alkoxycarbonyl, heterocycloyl, carboxy, $-\text{C}(\text{O})\text{O}-(\text{C}_1-\text{C}_3)\text{alkyl}$, $-\text{C}(\text{O})\text{NH}-(\text{C}_1-\text{C}_3)\text{alkyl}$, $-\text{C}(\text{O})\text{N}(\text{C}_1-\text{C}_3 \text{ alkyl})_2$, $-\text{PO}_3\text{H}_2$, haloalkyl, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, 30 cycloalkylalkyl, aryl, aroyl, biaryl, heterocyclyl, alkylaryl, aralkenyl,

-46-

aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl, $-\text{SO}_2-(\text{C}_1\text{-C}_3 \text{ alkyl})$,
 sulfonamido, aryloxyalkyl and $-\text{C}(\text{O})\text{NH}(\text{benzyl})$ groups;
 wherein B , R^6 , R^7 , R^9 , R^{10} , R^{18} , R^{24} , R^{25} , R^{26} and R^{27} are unsubstituted or
 substituted with at least one electron donating or electron withdrawing group;

5 wherein R^{18} and R^{24} taken together may form a ring;

R^{24} and R^{25} taken together may form a ring;

R^{25} and R^{26} taken together may form a ring;

and wherein R^9 and R^{10} taken together may form a ring;


or a pharmaceutically acceptable salt thereof;

10 one or more other therapeutically active compounds and a pharmacologically
 acceptable diluent.

11. The composition of claim 10 wherein B , R^6 , R^7 , R^9 , R^{10} , R^{24} , R^{25} and R^{26} are each
 independently hydrogen and R^{18} is substituted or unsubstituted aralkyl.

15

12. A pharmaceutical composition comprising a compound of the structure



20 wherein Z , at each occurrence, is independently selected from the group
 consisting of $\text{C}(\text{O})$, N , CR^{30} , $\text{C}(\text{R}^{31})(\text{R}^{32})$, NR^{33} , CH , O and S ;

z is an integer of from 3 to 6;

k is an integer of from 0 to 5;

T is selected from the group consisting of $\text{C}(\text{O})$ and $(\text{CH}_2)_b$ wherein b is an
 integer of from 0 to 3;

25 L is selected from the group consisting of O , NR^{11} , S , and

-47-

$(\text{CH}_2)_n$ wherein n is an integer of 0 or 1;

$\text{R}^6, \text{R}^7, \text{R}^{11}, \text{R}^{18}$ and R^{33} are each independently selected from the group consisting of alkyl, alkenyl, alkynyl, hydroxyalkyl, aliphatic acyl, alkynylamino, alkoxy carbonyl, heterocycloyl, $-\text{CH}=\text{NOH}$, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, carbamate, aryloxyalkyl, hydrogen and $-\text{C}(\text{O})\text{NH}(\text{benzyl})$ groups;

B is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, hydroxyalkyl, haloalkyl, $-\text{CF}_3$, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl and aryloxyalkyl;

$\text{R}^4, \text{R}^9, \text{R}^{10}, \text{R}^{30}, \text{R}^{31}$ and R^{32} at each occurrence are independently selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkynoxy, thioalkoxy, hydroxyalkyl, aliphatic acyl, $-\text{CF}_3$, $-\text{CO}_2\text{H}$, $-\text{SH}$, $-\text{OH}$, $-\text{CN}$, $-\text{NO}_2$, $-\text{NH}_2$, alkynylamino, alkoxy carbonyl, heterocycloyl, carboxy, $-\text{N}(\text{C}_1\text{-C}_3 \text{ alkyl})-\text{C}(\text{O})(\text{C}_1\text{-C}_3 \text{ alkyl})$, $-\text{NHC}(\text{O})\text{N}(\text{C}_1\text{-C}_3 \text{ alkyl})\text{C}(\text{O})\text{NH}(\text{C}_1\text{-C}_3 \text{ alkyl})$, $-\text{NHC}(\text{O})\text{NH}(\text{C}_1\text{-C}_6 \text{ alkyl})$, $-\text{NHSO}_2(\text{C}_1\text{-C}_3 \text{ alkyl})$, $-\text{NHSO}_2(\text{aryl})$, alkoxyalkyl, alkylamino, alkenylamino, $\text{di}(\text{C}_1\text{-C}_3)\text{amino}$, $-\text{C}(\text{O})\text{O}-(\text{C}_1\text{-C}_3)\text{alkyl}$, $-\text{C}(\text{O})\text{NH}-(\text{C}_1\text{-C}_3)\text{alkyl}$, $-\text{C}(\text{O})\text{N}(\text{C}_1\text{-C}_3 \text{ alkyl})_2$, $-\text{CH}=\text{NOH}$, $-\text{PO}_3\text{H}_2$, $-\text{OPO}_3\text{H}_2$, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide, cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl, aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl, aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl, $-\text{SO}_2-(\text{C}_1\text{-C}_3 \text{ alkyl})$, $-\text{SO}_3-(\text{C}_1\text{-C}_3 \text{ alkyl})$, sulfonamido, carbamate, aryloxyalkyl and $-\text{C}(\text{O})\text{NH}(\text{benzyl})$ groups; and

R^{29} , at each occurrence, is independently selected from the group consisting of halogen, alkyl, alkenyl, alkynyl, alkoxy, alkenoxy, alkynoxy, thioalkoxy, hydroxyalkyl, aliphatic acyl, $-\text{CF}_3$, $-\text{CO}_2\text{H}$, $-\text{SH}$, $-\text{CN}$, $-\text{NO}_2$, $-\text{NH}_2$, $-\text{OH}$, alkynylamino, alkoxy carbonyl, heterocycloyl, carboxy, $-\text{N}(\text{C}_1\text{-C}_3 \text{ alkyl})-$

-48-

C(O)(C₁-C₃ alkyl), -NHC(O)N(C₁-C₃ alkyl)C(O)NH(C₁-C₃alkyl),
 -NHC(O)NH(C₁-C₆ alkyl), -NHSO₂(C₁-C₃ alkyl), -NHSO₂(aryl),
 alkoxyalkyl, alkylamino, alkenylamino, di(C₁-C₃)amino, -C(O)O-(C₁-
 C₃)alkyl, -C(O)NH-(C₁-C₃)alkyl, -C(O)N(C₁-C₃ alkyl)₂, -CH=NOH,
 5 -PO₃H₂, -OPO₃H₂, haloalkyl, alkoxyalkoxy, carboxaldehyde, carboxamide,
 cycloalkyl, cycloalkenyl, cycloalkynyl, cycloalkylalkyl, aryl, aroyl,
 aryloxy, arylamino, biaryl, thioaryl, diarylamino, heterocyclyl, alkylaryl,
 aralkenyl, aralkyl, alkylheterocyclyl, heterocyclylalkyl, sulfonyl,
 -SO₂-(C₁-C₃ alkyl), -SO₃-(C₁-C₃ alkyl), sulfonamido, carbamate,
 10 aryloxyalkyl and -C(O)NH(benzyl) groups;

wherein B, R⁴, R⁵, R⁶, R⁷, R⁹, R¹⁰, R¹¹, R¹⁸, R²⁹, R³⁰, R³¹, R³² and R³³ are
 unsubstituted or substituted with at least one electron donating or electron
 withdrawing group;

wherein when L is NR¹¹, R⁴ and R¹¹ taken together may form a ring;

15 and wherein R⁹ and R¹⁰ taken together may form a ring;

or a pharmaceutically acceptable salt thereof;

one or more other therapeutically active compounds and a pharmacologically
 acceptable diluent.

20 13. The composition of claim 12 wherein z is three or four.

14. The composition of claim 1 where the compound of structure (I) is (3S)-3-[[[1-(2-
 chlorobenzyl)-4-hydroxy-5-methyl-2-oxo-1,2-dihydropyridin-3-yl]amino}carbonyl]amino]-3-
 (4-methylphenyl)propanoic acid and pharmaceutically acceptable salts thereof.

25

15. The composition of claim 1 where the compound of structure (I) is (3S)-3-[[[1-(2-
 chlorobenzyl)-4-hydroxy-2-oxo-2,5,6,7-tetrahydro-1H-cyclopenta[b]pyridin-3-
]amino}carbonyl]amino]-3-(4-methylphenyl)propanoic acid and pharmaceutically acceptable
 salts thereof.

30

16. The composition of claim 1 where the compound of structure (I) is (3S)-3-[[[1-(2-

chlorobenzyl)-4-hydroxy-5-methyl-2-oxo-1,2-dihydropyridin-3-yl]amino} carbonyl)amino]-3-[3-(diethylamino)phenyl]propanoic acid and pharmaceutically acceptable salts thereof.

17. The composition of claim 1 where the compound of structure (I) is (3S)-3-[[[1-(2-chlorobenzyl)-4-hydroxy-2-oxo-2,5,6,7-tetrahydro-1H-cyclopenta[b]pyridin-3-yl]amino} carbonyl)amino]-3-(3-isopropylphenyl)propanoic acid and pharmaceutically acceptable salts thereof.

18. The composition of claim 1 where the compound of structure (I) is (3S)-3-[[[1-(2-chlorobenzyl)-4-hydroxy-2-oxo-1,2-dihydro-1,8-naphthyridin-3-yl]amino} carbonyl)amino]-3-(4-methylphenyl)propanoic acid and pharmaceutically acceptable salts thereof.

19. The composition of claim 1 where the other therapeutically active compounds are selected from the group consisting of IL-5 antagonists, CCR-3 antagonists, corticosteroids, antihistamines, Leukotrine antagonists, COX-I and COX-II inhibitors, mast cell stabilizers, anti IL-5 and anti IgE antibodies, IL-5 synthesis and release inhibitors, TNF- α inhibitors, p38 MAP kinase inhibitors, tryptase inhibitors, anticytokine/antichemokine agents, vaccines, cromolyn, selectin antagonists, PDE 4 inhibitors, β -agonists, muscarinic antagonists and immunosuppressives, CD20 antagonists and syk tyrosine kinase inhibitors.

20. A method for treating an inflammatory disease in a mammal comprising administering to said mammal a therapeutically effective amount of a composition of claim 1.

21. The method of claim 20 wherein the inflammatory disease is selected from psoriasis, asthma, atherosclerosis, multiple sclerosis, Guillan-Barr Syndrome, rheumatoid arthritis, inflammatory bowel disease and reperfusion injury.

22. A method for treating an inflammatory disease in a mammal comprising administering to said mammal a therapeutically effective amount of a combination of a compound of structure (I) in claim 1 and an effective amount of one or more other therapeutic compounds.

23. The method of claim 22 wherein the inflammatory disease is selected from psoriasis, asthma, atherosclerosis, multiple sclerosis, Guillan-Barr Syndrome, rheumatoid arthritis, inflammatory bowel disease and reperfusion injury.

5

24. The composition of claim 19 wherein the compound of structure (I) is selected from the group consisting of (3S)-3-[(1-(2-chlorobenzyl)-4-hydroxy-5-methyl-2-oxo-1,2-dihydropyridin-3-yl)amino]carbonylamino]-3-(4-methylphenyl)propanoic acid; (3S)-3-[(1-(2-chlorobenzyl)-4-hydroxy-2-oxo-2,5,6,7-tetrahydro-1H-cyclopenta [b]pyridin-3-yl)amino]carbonylamino]-3-(4-methylphenyl)propanoic acid; (3S)-3-[(1-(2-chlorobenzyl)-4-hydroxy-5-methyl-2-oxo-1,2-dihydropyridin-3-yl)amino]carbonylamino]-3-(3-(diethylamino)phenyl)propanoic acid; (3S)-3-[(1-(2-chlorobenzyl)-4-hydroxy-2-oxo-2,5,6,7-tetrahydro-1H-cyclopenta[b]pyridin-3-yl) amino]carbonylamino]-3-(3-isopropylphenyl)propanoic acid; and (3S)-3-[(1-(2-chlorobenzyl)-4-hydroxy-2-oxo-1,2-dihydro-1,8-naphthyridin-3-yl)amino] carbonyl amino-3-(4-methylphenyl)propanoic acid and pharmaceutically acceptable salts thereof.

10
15

25. The method of claim 20 wherein the compound of structure (I) is selected from the group consisting of (3S)-3-[(1-(2-chlorobenzyl)-4-hydroxy-5-methyl-2-oxo-1,2-dihydropyridin-3-yl)amino]carbonylamino]-3-(4-methylphenyl)propanoic acid; (3S)-3-[(1-(2-chlorobenzyl)-4-hydroxy-2-oxo-2,5,6,7-tetrahydro-1H-cyclopenta [b]pyridin-3-yl)amino]carbonylamino]-3-(4-methylphenyl)propanoic acid; (3S)-3-[(1-(2-chlorobenzyl)-4-hydroxy-5-methyl-2-oxo-1,2-dihydropyridin-3-yl)amino]carbonylamino]-3-(3-(diethylamino)phenyl)propanoic acid; (3S)-3-[(1-(2-chlorobenzyl)-4-hydroxy-2-oxo-2,5,6,7-tetrahydro-1H-cyclopenta[b]pyridin-3-yl) amino]carbonylamino]-3-(3-isopropylphenyl)propanoic acid; and (3S)-3-[(1-(2-chlorobenzyl)-4-hydroxy-2-oxo-1,2-dihydro-1,8-naphthyridin-3-yl)amino] carbonyl amino-3-(4-methylphenyl)propanoic acid and pharmaceutically acceptable salts thereof.

20
25

30 26. A kit comprising in a single package, one container comprising a compound that inhibits binding of an $\alpha_4\beta_1$ integrin to its receptors as set forth in structure (I) in claim 1 in a

-51-

pharmaceutically acceptable carrier and one or more separate containers comprising other therapeutic compounds in pharmaceutically acceptable carriers, with the compound that inhibits binding of $\alpha_4\beta_1$ integrin to its receptors and the other therapeutic compounds being present in amounts such that the combination is effective to treat disease states mediated by

5 $\alpha_4\beta_1$ integrin binding.