




(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2006/09/08
(87) Date publication PCT/PCT Publication Date: 2007/03/22
(85) Entrée phase nationale/National Entry: 2008/03/13
(86) N° demande PCT/PCT Application No.: CA 2006/001477
(87) N° publication PCT/PCT Publication No.: 2007/030919
(30) Priorité/Priority: 2005/09/14 (US60/716,507)

(51) Cl.Int./Int.Cl. *C12Q 1/68* (2006.01),
C12N 15/10 (2006.01), *C12N 15/12* (2006.01),
C12N 15/18 (2006.01), *C12N 15/54* (2006.01),
C12N 15/57 (2006.01), *C12Q 1/04* (2006.01),
C40B 30/04 (2006.01), *G01N 33/53* (2006.01),
G01N 33/574 (2006.01)
(71) Demandeur/Applicant:
NATIONAL RESEARCH COUNCIL OF CANADA, CA
(72) Inventeurs/Inventors:
BELACEL, NABIL, CA;
CUPERLOVIC-CULF, MIROSLAVA, CA;
OUELLETTE, RODNEY, CA
(74) Agent: BRION, ARTURO

(54) Titre : METHODE MOLECULAIRE DE DIAGNOSTIC DU CANCER DE LA PROSTATE
(54) Title: MOLECULAR METHOD FOR DIAGNOSIS OF PROSTATE CANCER



(57) Abrégé/Abstract:

Methods for diagnosing or detecting cancerous prostate tissue in a sample using a panel of 8 specific marker genes glutathione S-transferase M4, latent transforming growth factor beta binding protein 4, D component of complement (adipsin), Nel-like 2, X-box binding protein 1, intersectin 1, folate hydrolase and hepsin The over-expression or under-expression of the 8 marker genes, compared to their expression in normal prostate tissue, is indicative of cancerous prostate tissue By using these 8 marker genes as a diagnostic tool, smaller tissue samples, such as those obtained by core needle biopsies can be used.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



PCT



(43) International Publication Date
22 March 2007 (22.03.2007)

(10) International Publication Number
WO 2007/030919 A1

(51) International Patent Classification:

C12Q 1/68 (2006.01) *C12N 15/57* (2006.01)
C12N 15/12 (2006.01) *C12Q 1/04* (2006.01)
C12N 15/10 (2006.01) *C40B 30/04* (2006.01)
C12N 15/18 (2006.01) *G01N 33/53* (2006.01)
C12N 15/54 (2006.01) *G01N 33/574* (2006.01)

[CA/CA]; 450 Leblanc Rd., Dieppe, New Brunswick E1A 7N4 (CA).

(74) Agent: CASSAN MACLEAN; Suite 401, 80 Aberdeen Street, Ottawa, Ontario K1S 5R5 (CA).

(21) International Application Number:

PCT/CA2006/001477

(22) International Filing Date:

8 September 2006 (08.09.2006)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/716,507 14 September 2005 (14.09.2005) US

(71) Applicant (for all designated States except US): NATIONAL RESEARCH COUNCIL OF CANADA [CA/CA]; 1200 Montreal Road, Ottawa, Ontario K1A 0R6 (CA).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BELACEL, Nabil [CA/CA]; 30 Algonquin Crescent, Moncton, New Brunswick E1G 2C8 (CA). CUPERLOVIC-CULF, Miroslava [CA/CA]; 355 Glencairn Drive, Moncton, New Brunswick E1G 1Y5 (CA). OUELLETTE, Rodney

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: MOLECULAR METHOD FOR DIAGNOSIS OF PROSTATE CANCER



(57) Abstract: Methods for diagnosing or detecting cancerous prostate tissue in a sample using a panel of 8 specific marker genes glutathione S- transferase M4, latent transforming growth factor beta binding protein 4, D component of complement (adipsin), Nel-like 2, X-box binding protein 1, intersectin 1, folate hydrolase and hepsin The over-expression or under-expression of the 8 marker genes, compared to their expression in normal prostate tissue, is indicative of cancerous prostate tissue By using these 8 marker genes as a diagnostic tool, smaller tissue samples, such as those obtained by core needle biopsies can be used.

WO 2007/030919 A1

MOLECULAR METHOD FOR DIAGNOSIS OF PROSTATE CANCER

Field of the Invention

[00001] The present invention relates to diagnosis methods and, more particularly, to diagnosis methods for detecting prostate cancer.

Background to the Invention

[00002] Prostate cancer is a leading cancer in men with 20,100 new cases expected in Canada in 2004 (Canadian Cancer Statistics, 2004). An even larger number of patients, following a positive PSA (prostate specific antigen) reading undergo an invasive biopsy but are not diagnosed with cancer. Unfortunately, invasive biopsy procedures require a long hospitalization with many possible side-effects. The reason for using the more invasive biopsy procedure, as opposed to the less invasive core needle biopsy is that the standard diagnostic procedure on core needle biopsy samples has not been proven to be as accurate as a more invasive biopsy and was thereby discarded as a diagnostic modality in several countries. Although the more invasive biopsies may provide a more accurate diagnosis, they are extremely traumatic for the patient. Moreover, such procedures can potentially result in long term disabilities and constitutes a significant cost to the health system.

[00003] Current diagnostic techniques for detecting prostate cancer is based on the PSA level in the serum. The final diagnosis is determined by a pathologist checking for cancer cells in the biopsy samples.

[00004] However, these present techniques are not perfect. The PSA level in the serum can be affected by factors other than cancer, including other pathologies and age. In addition, specific properties of PSA protein in serum make accurate concentration measurements very difficult. As a result, the PSA test has a large percentage of false positive as well as false negative readings. Therefore, biopsy samples are essential for more accurate diagnosis. Unfortunately, as noted above, the preferred method of biopsy, the core needle biopsy, is often inaccurate due to the very small sample size. However, genetic

testing in core needle biopsy samples will allow for the accurate diagnosis without the need for more invasive methods.

[00005] There is therefore a need for a more accurate diagnostic method that does not require an invasive biopsy to detect or diagnose prostate cancer. Ideally, such a method should be usable even with very small sample sizes and may be combined with other, pathologist-based diagnosis methods.

Summary of the Invention

[00006] The present invention provides methods for diagnosing or detecting cancerous prostate tissue. A panel of 8 specific marker genes are provided. The overexpression of some of these marker genes compared to their expression in normal prostate tissue and the underexpression of the rest of these marker genes are indicative of cancerous prostate tissue. By using these 8 marker genes as a diagnostic tool, smaller tissue samples, such as those obtained by core needle biopsies can be used.

[00007] In a first aspect, the present invention provides a method for determining if prostate cells are cancerous, the method comprising:

- a) obtaining said prostate cells;
- b) determining if at least one specific gene is overexpressed or underexpressed in said prostate cells compared to an expression of said at least one specific gene in normal prostate cells;
- c) determining that said prostate cells are cancerous based on whether said at least one gene is overexpressed or underexpressed in said prostate cells.

[00008] In another aspect, the present invention provides a use of at least one marker gene for identifying cancerous prostate tissue, an overexpression or underexpression of said at least one marker gene in prostate tissue compared to an expression of said at least one marker gene in normal prostate tissue being indicative of cancerous prostate tissue.

[00009] Yet another aspect of the invention provides a method of diagnosing prostate cancer, the method comprising:

- a) obtaining prostate tissue to be diagnosed;
- b) determining if specific marker genes are overexpressed or

underexpressed in said prostate tissue to be diagnosed compared to non-cancerous prostate tissue;

c) determining if said prostate tissue to be diagnosed is cancerous based on an underexpression or overexpression of said specific marker genes.

Brief Description of the Drawings

[00010] A better understanding of the invention will be obtained by considering the detailed description below, with reference to the following drawings in which:

Figure 1 is an expression plot for the 8 genes which is the subject of the present invention;

Figure 2 is a boxplot of expression levels for the gene GSTM obtained over several experiments on prostate tissue;

Figure 3 is a boxplot of expression levels for the gene LTPB4 obtained over several experiments on prostate tissue;

Figure 4 is a boxplot of expression levels for the gene DF (adipsin) obtained over several experiments on prostate tissue;

Figure 5 is a boxplot of expression levels for the gene NELL2 obtained over several experiments on prostate tissue;

Figure 6 is a boxplot of expression levels for the gene XBP1 obtained over several experiments on prostate tissue;

Figure 7 is a boxplot of expression levels for the gene ITSN1 obtained over several experiments on prostate tissue;

Figure 8 is a boxplot of expression levels for the gene FOLH1 obtained over several experiments on prostate tissue; and

Figure 9 is a boxplot of expression levels for the gene Hepsin obtained over several experiments on prostate tissue.

Detailed Description

[00011] The present invention relates to the use of a panel of 8 specific marker genes to diagnose or detect cancerous prostate tissue. The panel of 8 marker genes are listed in Table 1 below. Experiments have shown that this

panel of marker genes give high accuracy in prostate cancer diagnosis due to the expression levels of the marker genes in cancer tissue relative to their expression levels in normal tissue.

[00012] The panel of 8 marker genes is given in Table 1. The marker genes were determined using a method developed by the inventors from the prostate tissues (normal and cancer) gene expression dataset obtained and described by Singh, D. et al. (Singh, D. et al. Gene expression correlates of clinical prostate cancer behaviour. *Cancer Cell* 1:203 (2002)).

Table 1 - Panel of eight genes found to give high accuracy in prostate cancer diagnosis and their expression level in cancer relative to normal tissue.

GeneBank Accession Number	Gene Name	Symbol	UniGene ID	Over or Under- expressed in cancer tissue relative to normal tissue
M96233	Glutathione S-transferase M4	GSTM4	Hs.348387	Under- expressed
Y13622	Latent transforming growth factor beta binding protein 4	LTBP4	Hs.85087	Under- expressed
M84526	D component of complement (adipsin)	DF	Hs.155597	Under- expressed
D83018	Nel-like 2	NELL2	Hs.79389	Under- expressed
Z93930	X-box binding protein 1	XBP1	Hs.149923	Over- expressed

AF064243	Intersectin 1 (SH3 domain protein)	ITSN1	Hs.66392	Under-expressed
M99487	Folate hydrolase (Prostate specific membrane antigen) 1	FOLH1	Hs.1915	Over-expressed
X07732	Hepsin (transmembrane protease, serine 1)	HPN	Hs.823	Over-expressed

[00013] The genes listed above were derived using a microarray gene expression experiment, the gene expression plot being provided as Figure 1 for the 8 genes. For this expression plot, the samples are normal and tumour tissues.

[00014] Figures 2-9 are boxplots of the comparative expression of the specific marker genes in normal prostate tissue, cancerous tissue, and other types of prostate tissue. The testing was done using the prostate microarray data available at the Oncomine database (see Rhodes DR, Yu J, Shanker K, Deshpande N, Varambally R, Ghosh D, Barrette T, Pandey A, and Chinnaiyan AM. ONCOMINE: A Cancer Microarray Database and Data-Mining Platform. *Neoplasia* 2004); and Rhodes DR, Yu J, Shanker K, Deshpande N., Varambally R, Ghosh D, Barrette T, Pandey A, and Chinnaiyan AM, Large-Scale Meta-Analysis of Cancer Microarray Data Identifies Common Transcriptional Profiles of Neoplastic Transformation and Progression, *PNAS* (2004)).

[00015] The dataset used for the experiments which resulted in the expression levels shown in the boxplots of Figs 2-9 included gene expressions measured using Affymetrix technology on 52 prostate tumors and 50 non-tumor prostate samples. The pre-processing included threshold readjustment (floor 50, ceiling 16,000), filtering (max/min less than 5; max-min less than 50); logarithmic transformation and standardization (to mean of zero and variance of 1). The pre-processing resulted in the set comprising 6034 genes and 102 experiments. The

microarray experiments were performed on prostate tumour samples and adjacent prostate tissue not containing tumour cells (selected by a pathologist based on cell morphology, referred to as “normal”) collected from patients undergoing radical prostatectomy at the Brigham and Women’s Hospital, Boston, Massachusetts. Each of these samples was reviewed by pathologist determining their Gleason Score (GS) as well as serum PSA. As noted above, microarray experiments were performed using GeneChip arrays (Affymetrix Inc.). Total RNA extracted from each sample was labeled with the fluorescence dye and hybridized to the slide in a one sample per slide format. Gene expression levels were obtained by measuring fluorescence intensities for each target (spot). Data pre-processing was performed using the standard protocol for GeneChip arrays (see Welsh JB., Analysis of gene expression identifies candidate markers and pharmacological targets in prostate cancer. *Cancer Res.* **61**: 5974 (2001)) Data for each experiment was then scaled (normalized) to have overall mean of zero and standard deviation of one making data from different experiments comparable.

[00016] Referring to Figure 2, a boxplot of the expression levels for the gene GSTM4 obtained over several different experiments on prostate tissues is presented. Each box includes values for replicated experiments. The experiments included and the tissues used are summarized in Table 2:

Table 2:

Experiment	C1	C2	P-value
A1 [4]	BPH and Normal Prostate	Prostate Cancer	6.366
A2 [4]	Prostate Cancer	Metastatic Prostate Cancer	0.067
A3 [4]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8	0.0823

A4 [4]	Prostate Cancer: No PSA recur at 2 years	Prostate Cancer: PSA recur	0.8647
A5 [5]	Benign Prostatic Hyperplasia	Prostate Cancer	0.1103

[00017] As can be seen from Fig 2 in conjunction with Table 2 above, the gene GSTM4 is underexpressed in cancerous prostate tissue when compared to its expression in normal prostate tissue (see experiment A1 in Fig 2). It can also be seen that, compared to the expression of this gene in normal prostate tissue, cancerous prostate tissue has a lower expression (see C1 in experiments A2-A4). Thus, for cancerous prostate tissue, the gene GSTM4 is underexpressed when compared to its expression in normal prostate tissue.

[00018] Referring to Fig 3, a boxplot of expression levels for the gene LTPB4 obtained over several different experiments on prostate tissues is illustrated. Each box includes values for replicated experiments. The experiments included and the tissues used are summarized in Table 3 below:

Table 3 :

Experiment	C1	C2	P-value
A1 [4]	BPH and Normal Prostate	Prostate Cancer	0.0872
A2 [4]	Prostate Cancer	Metastatic Prostate Cancer	0.0069
A3 [4]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8	0.1896
A4 [4]	Prostate Cancer: No PSA recur at 2 years	Prostate Cancer: PSA recur	0.7115

A5 [6]	Normal Prostate	Prostate Cancer	0.1142
A6 [6]	Prostate Cancer	Metastatic Prostate Cancer	1.2e-5
A7 [7]	Normal Prostate	Prostate Cancer	0.158
A8 [5]	Benign Prostatic Hyperplasia	Prostate Cancer	0.0404
A9 [8]	Normal Prostate	Prostate Cancer	1.459
A10 [8]	Prostate Cancer	Metastatic Prostate Cancer	0.7855
A11 [9]	No PSA recurrence at 3 years	PSA Recurrence at 3 years	0.6524
A12 [9]	Nontumor Prostate	Prostate Cancer	3.7e-5
A13 [9]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8,9	0.3183
A14 [10]	Normal Prostate	Prostate Cancer	6.4e-6
A15 [10]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8,9	0.6032

[00019] The results illustrated in Fig 3 show that the gene LTPB4 is underexpressed in cancerous prostate tissue when compared to its expression in normal prostate tissue. The results of experiments A1, A5, A7, A9, A12, and A14 show that, compared to its expression in normal prostate tissue, the gene LTPB4 is underexpressed in cancerous prostate tissue. For prostate cancer tissue, the gene

LTPB4 is therefore underexpressed when compared to its expression in normal prostate tissue.

[00020] Referring to Fig 4, such is a boxplot of expression levels for the gene DF (adipsin) obtained over several different experiments on prostate tissues. Each box includes values for replicated experiments. The experiments and the prostate tissue included in the experiments are summarized in Table 4 below:

Table 4:

Experiment	C1	C2	P-value
A1 [4]	BPH and Normal Prostate	Prostate Cancer	0.4053
A2 [4]	Prostate Cancer	Metastatic Prostate Cancer	0.7189
A3 [4]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8	0.044
A4 [4]	Prostate Cancer: No PSA recur at 2 years	Prostate Cancer: PSA recur	0.7859
A5 [6]	Normal Prostate	Prostate Cancer	0.0006
A6 [6]	Prostate Cancer	Metastatic Prostate Cancer	0.013
A7 [7]	Normal Prostate	Prostate Cancer	0.0642
A8 [11]	Normal Prostate	Prostate Cancer	0.6993

A9 [8]	Normal Prostate	Prostate Cancer	0.0893
A10 [8]	Prostate Cancer	Metastatic Prostate Cancer	0.0031
A11 [9]	No PSA recurrence at 3 years	PSA Recurrence at 3 years	0.0518
A12 [9]	Nontumor Prostate	Prostate Cancer	6.4e-10
A13 [9]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8,9	0.2172
A14 [10]	Normal Prostate	Prostate Cancer	0.118
A15 [10]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8,9	0.9273

[00021] The results in Fig 4 show that the gene DF(adipsin) is underexpressed in cancerous prostate tissue compared to its expression in normal prostate tissue. As can be seen from the results in experiments A1, A5, A7-A9, A12, and A14, the gene DF(adipsin) has much lower expression levels in cancerous prostate tissue than in normal prostate tissue.

[00022] Referring to Figure 5, a boxplot of expression levels for gene NELL2 obtained over several different experiments on prostate tissues is illustrated. Each box includes values for replicated experiments. The experiments and the types of prostate tissue used in the experiments are summarized in Table 5 below:

Table 5:

Experiment	C1	C2	P-value
A1 [4]	BPH and Normal Prostate	Prostate Cancer	0.0039
A2 [4]	Prostate Cancer	Metastatic Prostate Cancer	0.0339
A3[4]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8	0.3201
A4 [4]	Prostate Cancer: No PSA recur at 2 years	Prostate Cancer: PSA recur	0.3781
A5 [6]	Normal Prostate	Prostate Cancer	0.0041
A6 [6]	Prostate Cancer	Metastatic Prostate Cancer	0.0018
A7 [7]	Normal Prostate	Prostate Cancer	0.9941
A8 [5]	Benign Prostatic Hyperplasia	Prostate Cancer	4.10E-06
A9 [11]	Normal Prostate	Prostate Cancer	0.3558
A10 [8]	Normal Prostate	Prostate Cancer	0.0009
A11 [8]	Prostate Cancer	Metastatic Prostate Cancer	0.832
A12 [9]	No PSA recurrence at 3 years	PSA Recurrence at 3 years	0.0291
A13 [9]	Nontumor Prostate	Prostate Cancer	1.20E-12
A14 [9]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8,9	0.8026
A15 [10]	Normal Prostate	Prostate Cancer	0.1714
A16 [10]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8,9	0.6622

[00023] The results in Fig 5 illustrate that the gene NELL2 is underexpressed in cancerous prostate tissue when compared to its expression in non-cancerous prostate tissue. The results for the experiments A1, A5, A7, A9, A10, A13, and A14 show that the gene NELL2 has a lower expression in cancerous prostate tissue than in normal prostate tissue.

[00024] Referring to Figure 6, a boxplot of expression levels for gene XBP1 obtained over several different experiments on prostate tissues is illustrated. Each box includes values for replicated experiments. The experiments and the types of prostate tissue used are summarized in Table 6 below:

Table 6 :

Experiment	C1	C2	P-value
A1 [4]	BPH and Normal Prostate	Prostate Cancer	0.0042
A2 [4]	Prostate Cancer	Metastatic Prostate Cancer	1.00E-04
A3 [4]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8	0.0456
A4 [4]	Prostate Cancer: No PSA recur at 2 years	Prostate Cancer: PSA recur	0.0581
A5 [7]	Normal Prostate	Prostate Cancer	0.5674
A6 [5]	Benign Prostatic Hyperplasia	Prostate Cancer	0.8723
A7 [11]	Normal Prostate	Prostate Cancer	0.8965
A8 [8]	Normal Prostate	Prostate Cancer	0.428
A9[8]	Prostate Cancer	Metastatic Prostate Cancer	0.0697

[00025] Figure 6 shows that the gene XBP1 is overexpressed in cancerous prostate tissue as opposed to normal prostate tissue. Experiments A1, A4, A5, A7 and A8 illustrate that the gene XBP1 has a lower expression level in normal prostate tissue than in cancerous prostate tissue.

[00026] Referring to Figure 7, a boxplot of expression levels for gene ITSN1 obtained over several different experiments on prostate tissues is illustrated. Each box includes values for replicated experiments. The experiments and the various tissues used are summarized in Table 7 below:

Table 7 :

Experiment	C1	C2	P-value
A1 [4]	BPH and Normal Prostate	Prostate Cancer	0.0024
A2 [4]	Prostate Cancer	Metastatic Prostate Cancer	0.0106
A3 [4]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8	0.2204
A4 [4]	Prostate Cancer: No PSA recur at 2 years	Prostate Cancer: PSA recur	0.6537
A5 [6]	Normal Prostate	Prostate Cancer	0.3049
A6[6]	Prostate Cancer	Metastatic Prostate Cancer	0.0051
A7 [7]	Normal Prostate	Prostate Cancer	0.2124
A8 [5]	Benign Prostatic Hyperplasia	Prostate Cancer	0.0021
A9 [9]	No PSA recurrence at 3 years	PSA Recurrence at 3 years	0.6691
A10 [9]	Nontumor Prostate	Prostate Cancer	3.50E-10
A11 [9]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8,9	0.2232
A12[10]	Normal Prostate	Prostate Cancer	0.0003
A13 [10]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8,9	0.586

[00027] Figure 7 illustrates that the gene ITSN1 has a lower expression in cancerous prostate tissue when compared to its expression in normal prostate tissue. As can be seen from the results of experiments A1, A5, A7, A10, and A12, cancerous prostate tissue has lower expression levels of ITSN1 when compared to non-cancerous prostate tissue.

[00028] Referring to Figure 8, a boxplot of expression levels for gene FOLH1 obtained over several different experiments on prostate tissues is illustrated. Each box includes values for replicated experiments. The experiments and the tissues used are summarized in Table 8 below:

Table 8:

Experiment	C1	C2	P-value
A1 [4]	BPH and Normal Prostate	Prostate Cancer	0.0011
A2 [4]	Prostate Cancer	Metastatic Prostate Cancer	0.07
A3 [4]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8	0.3748
A4 [4]	Prostate Cancer: No PSA recur at 2 years	Prostate Cancer: PSA recur	0.0801
A5 [6]	Normal Prostate	Prostate Cancer	0.0047
A6 [6]	Prostate Cancer	Metastatic Prostate Cancer	0.2598
A7 [7]	Normal Prostate	Prostate Cancer	0.0059
A8 [5]	Benign Prostatic Hyperplasia	Prostate Cancer	0.0152
A9 [11]	Normal Prostate	Prostate Cancer	0.3939
A10 [8]	Normal Prostate	Prostate Cancer	0.8983
A11 [8]	Prostate Cancer	Metastatic Prostate Cancer	0.0123
A12 [10]	Normal Prostate	Prostate Cancer	0.0039
A13 [10]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8,9	0.2576

[00029] Figure 8 illustrates that the gene FOLH1 is overexpressed in cancerous prostate tissue as compared to normal prostate tissue. The results of experiments A1, A5, A7, A9, A10, and A12 show that the gene FOLH1 has a higher expression in cancerous prostate tissue than in normal prostate tissue.

[00030] Referring to Figure 9, a boxplot of expression levels for gene Hepsin obtained over several different experiments on prostate tissues is illustrated. Each box includes values for replicated experiments. The experiments and the different types of prostate tissue used are summarized in Table 9 below:

Table 9

Experiment	C1	C2	P-value
A1 [4]	BPH and Normal Prostate	Prostate Cancer	1.60E-08
A2 [4]	Prostate Cancer	Metastatic Prostate Cancer	0.3311
A3 [4]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8	0.8772
A4 [4]	Prostate Cancer: No PSA recur at 2 years	Prostate Cancer: PSA recur	0.5244
A5 [6]	Normal Prostate	Prostate Cancer	0.0205
A6 [6]	Prostate Cancer	Metastatic Prostate Cancer	0.0351
A7 [7]	Normal Prostate	Prostate Cancer	0.0017
A8 [5]	Benign Prostatic Hyperplasia	Prostate Cancer	1.80E-07
A9 [8]	Normal Prostate	Prostate Cancer	0.0484
A10 [8]	Prostate Cancer	Metastatic Prostate Cancer	0.0662
A11 [9]	No PSA recurrence at 3 years	PSA Recurrence at 3 years	0.866
A12 [9]	Nontumor Prostate	Prostate Cancer	6.10E-25
A13 [9]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8,9	0.2081
A14 [10]	Normal Prostate	Prostate Cancer	2.10E-08
A15 [10]	Prostate Cancer: Gleason 6	Prostate Cancer: Gleason 8,9	0.7475

[00031] From Fig 9, it can be seen that the gene Hepsin is overexpressed in cancerous prostate tissue as opposed to normal prostate tissue. The results of experiments A1, A5, A7, A8, A9, A12, and A14 show that Hepsin has a higher expression in cancerous prostate tissue than in normal prostate tissue.

[00032] The qualitative agreement between the differences in expression in cancer and normal prostate tissues for the eight genes investigated here is good overall with most genes showing significant difference between cancer and normal

tissue as well as benign and malignant tumours in the majority of experiments. The results shown here prove that a subset of 8 genes is an appropriate diagnostic panel regardless of the experimental conditions. At the same time, in some cases, especially for the gene LTPB4, there are changes in the relative expression in cancer against normal prostate tissue depending on the experiment. Such variations show that it is preferable that one looks at the overall result for the whole panel and that diagnostics based on only one gene may be unreliable.

[00033] It should be noted that expression analysis can be carried out using any method for measuring gene expression. Such methods as microarrays, diagnostic panel mini-chip, PCR, real-time PCR, and other similar methods may be used. Similarly, methods for measuring protein expression may also be used.

[00034] As noted above, the cancerous prostate cells can be obtained from a patient using needle biopsy or even from prostate cancer cells present in the blood stream. Normal or non-cancerous prostate cells against which the cancerous cells can be compared can also be obtained from the patient or from other patients. Experiments have shown that the diagnosis can be possible from just a small number of cancer cells.

[00035] While it is preferable that the complete panel of 8 marker genes be used in the diagnosis of possible prostate cancer, using a subset of the 8 marker genes will also yield useful results. Using a panel of anywhere from 1 to 7 marker genes out of the 8 marker genes on suspect prostate tissue will still provide a useful indication as to whether cancerous prostate tissue may be present or whether further and more involved tests are required.

[00036] A person understanding this invention may now conceive of alternative structures and embodiments or variations of the above all of which are intended to fall within the scope of the invention as defined in the claims that follow.

We claim:

1. A method for determining if prostate cells are cancerous, the method comprising:

a) obtaining said prostate cells;

b) determining if at least one first gene is overexpressed in said prostate cells compared to an expression of said at least one first gene in normal prostate cells, said at least one first gene being selected from a group consisting of :

- Hepsin (transmembrane protease serine 1);

- X-box binding protein 1; and

- Folate hydrolase (Prostate specific membrane antigen) 1

c) determining if at least one second gene is underexpressed in said prostate cells compared to an expression of said at least one second gene in normal prostate cells, said at least one second gene being selected from a group consisting of :

- Glutathione S-transferase M4;

- Intersectin 1 (SH3 domain protein);

- Nel-like 2;

- D component of complement (adipsin); and

- Latent transforming growth factor beta binding protein 4

d) determining that said prostate cells are cancerous if :

i) said at least one first gene is overexpressed in said prostate cells compared to an expression of said at least one first gene in normal prostate cells and

ii) said at least one second gene is underexpressed in said prostate cells compared to an expression of said at least one second gene in normal prostate cells.

2. A method according to claim 1 wherein said prostate cells are obtained by a core needle biopsy.

3. Use of at least one first marker gene and at least one second marker gene for identifying cancerous prostate tissue, a combination of both :

i) an overexpression of said at least one first marker gene in prostate tissue compared to an expression of said at least one first marker gene in normal prostate tissue and

ii) an underexpression of said at least one second marker gene in prostate tissue compared to an expression of said at least one second marker gene in normal prostate tissue

being indicative of cancerous prostate tissue.

4. A use according to claim 3 wherein said at least one first marker gene is selected from a group consisting of

- Hepsin (transmembrane protease serine 1);
- X-box binding protein 1; and
- Folate hydrolase (Prostate specific membrane antigen) 1.

5. A use according to claim 3 wherein said at least one second marker gene is selected from a group consisting of:

- Glutathione S-transferase M4;
- Intersectin 1 (SH3 domain protein);
- Nel-like 2;
- D component of complement (adipsin); and
- Latent transforming growth factor beta binding protein 4.

6. A use according to claim 3 wherein said at least one first marker gene is selected from a group consisting of

- Hepsin (transmembrane protease serine 1);
- X-box binding protein 1; and
- Folate hydrolase (Prostate specific membrane antigen) 1

and wherein said at least one second marker gene is selected from a group consisting of:

- Glutathione S-transferase M4;
- Intersectin 1 (SH3 domain protein);
- Nel-like 2;
- D component of complement (adipsin); and
- Latent transforming growth factor beta binding protein 4.

7. A method of diagnosing prostate cancer, the method comprising:

- a) obtaining prostate tissue to be diagnosed;
- b) determining if at least one first marker gene is overexpressed in said prostate tissue to be diagnosed compared to non-cancerous prostate tissue;
- c) determining if at least one second marker gene is underexpressed in said prostate tissue to be diagnosed compared to non-cancerous prostate tissue;
- d) determining if said prostate tissue to be diagnosed is cancerous based on two conditions :
 - i) said at least one first marker gene is overexpressed in said prostate tissue to be diagnosed
 - ii) said at least one second marker gene is underexpressed in said prostate tissue to be diagnosed.

8. A method according to claim 7 wherein said prostate tissue is obtained by a core needle biopsy.

9. A method according to claim 7 wherein said at least one first marker gene is selected from a group consisting of:

- Hepsin (transmembrane protease serine 1);
- X-box binding protein 1; and
- Folate hydrolase (Prostate specific membrane antigen) 1.

10. A method according to claim 7 said at least one second marker gene is selected from a group consisting of:

- Glutathione S-transferase M4;
- Intersectin 1 (SH3 domain protein);
- Nel-like 2;
- D component of complement (adipsin); and
- Latent transforming growth factor beta binding protein 4.


11. A method according to claim 7 wherein said at least one first marker gene is selected from a group consisting of:

- Hepsin (transmembrane protease serine 1);
- X-box binding protein 1; and
- Folate hydrolase (Prostate specific membrane antigen) 1

and wherein said at least one second marker gene is selected from a group consisting of:

- Glutathione S-transferase M4;
- Intersectin 1 (SH3 domain protein);
- Nel-like 2;

- D component of complement (adipsin); and
- Latent transforming growth factor beta binding protein 4.






Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8






Figure 9

-3.0

1:1

3.0



"M84526"
"M96233"
"Y13622"
"AF064243"
"083018"
"Z93930"
"M99487"
"X07732"