



(86) **Date de dépôt PCT/PCT Filing Date:** 2008/05/15
(87) **Date publication PCT/PCT Publication Date:** 2008/12/04
(45) **Date de délivrance/Issue Date:** 2015/04/14
(85) **Entrée phase nationale/National Entry:** 2009/11/20
(86) **N° demande PCT/PCT Application No.:** US 2008/063630
(87) **N° publication PCT/PCT Publication No.:** 2008/147697
(30) **Priorité/Priority:** 2007/05/22 (US60/939,452)

(51) **Cl.Int./Int.Cl. C07D 235/26** (2006.01),
A61K 31/4184 (2006.01), **A61P 11/06** (2006.01),
C07D 401/06 (2006.01), **C07D 401/14** (2006.01),
C07D 403/06 (2006.01), **C07D 403/14** (2006.01),
C07D 405/12 (2006.01), **C07D 405/14** (2006.01),
C07D 409/06 (2006.01), **C07D 409/12** (2006.01),
C07D 409/14 (2006.01), ...

(72) **Inventeurs/Inventors:**
ABEYWARDANE, ASITHA, US;
COOK, BRIAN NICHOLAS, US;
DE LOMBAERT, STEPHANE, US;
EMMANUEL, MICHEL JOSE, US;
...

(54) **Titre : INHIBITEURS DE BENZIMIDAZOLONE CHYMASE**
(54) **Title: BENZIMIDAZOLONE CHYMASE INHIBITORS**

(57) **Abrégé/Abstract:**

Disclosed are small molecule inhibitors which are useful in treating various diseases and conditions involving chymase.



(51) **Cl.Int./Int.Cl. (suite/continued)** **C07D 413/06** (2006.01), **C07D 417/06** (2006.01), **C07D 417/14** (2006.01)

(72) **Inventeurs(suite)/Inventors(continued)**: GUO, XIN, US; KIM, JIN MI, US; HAO, MING-HONG, US; LO, HO YIN, US; MAN, CHUK CHUI, US; MORWICK, TINA, US; NEMOTO, PETER ALLEN, US; QIAN, KEVIN CHUNGENG, US; TAKAHASHI, HIDENORI, US; TAYLOR, STEVEN JOHN, US

(73) **Propriétaires(suite)/Owners(continued)**:BOEHRINGER INGELHEIM INTERNATIONAL GMBH, DE

(74) **Agent**: FETHERSTONHAUGH & CO.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 December 2008 (04.12.2008)

PCT

(10) International Publication Number
WO 2008/147697 A1

(51) International Patent Classification:

C07D 235/26 (2006.01) *C07D 409/12* (2006.01)
C07D 401/06 (2006.01) *C07D 409/14* (2006.01)
C07D 401/14 (2006.01) *C07D 413/06* (2006.01)
C07D 403/06 (2006.01) *C07D 417/06* (2006.01)
C07D 403/14 (2006.01) *C07D 417/14* (2006.01)
C07D 405/12 (2006.01) *A61K 31/4184* (2006.01)
C07D 405/14 (2006.01) *A61P 11/06* (2006.01)
C07D 409/06 (2006.01)

(21) International Application Number:

PCT/US2008/063630

(22) International Filing Date: 15 May 2008 (15.05.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/939,452 22 May 2007 (22.05.2007) US

(71) Applicant (*for all designated States except DE, US*):
BOEHRINGER INGELHEIM INTERNATIONAL GMBH [DE/DE]; Binger Strasse 173, 55216 Ingelheim (DE).

(71) Applicant (*for DE only*): **BOEHRINGER INGELHEIM PHARMA GMBH & CO. KG** [DE/DE]; Binger Strasse 173, 55216 Ingelheim (DE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **ABEYWARDANE, Asitha** [LK/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877-0368 (US). **COOK, Brian, Nicholas** [US/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877-0368 (US). **DELOMBAERT, Stephane** [BE/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877-0368 (US). **EMMANUEL, Michel, Jose** [US/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877-0368 (US). **GUO, Xin** [CN/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877-0368 (US). **KIM, Jin, Mi** [CA/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877-0368 (US). **HAO, Ming-Hong**

[US/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877-0368 (US). **LO, Ho, Yin** [CN/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877-0368 (US). **MAN, Chuk, Chui** [GB/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877-0368 (US). **MORWICK, Tina** [US/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877-0368 (US). **NEMOTO, Peter, Allen** [US/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877-0368 (US). **QIAN, Kevin, Chungeng** [US/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877-0368 (US). **TAKAHASHI, Hidenori** [JP/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877-0368 (US). **TAYLOR, Steven, John** [US/US]; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877-0368 (US).

(74) Agents: **MORRIS, Michael, P.** et al.; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(54) Title: BENZIMIDAZOLONE CHYMASE INHIBITORS

(57) Abstract: Disclosed are small molecule inhibitors which are useful in treating various diseases and conditions involving chymase.

WO 2008/147697 A1

DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME __1__ DE __2__

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**

THIS IS VOLUME __1__ OF __2__

NOTE: For additional volumes please contact the Canadian Patent Office.

25771-1706

Benzimidazolone Chymase Inhibitors

Field of the Invention

The invention relates to small molecule inhibitors which are useful in treating various diseases and conditions involving Chymase.

Background of the Invention

In cardiac tissue of cardiomyopathic patients, transforming growth factor- β (TGF- β), which has been demonstrated to stimulate cardiac fibrosis in animal models (Kuwahara, et al. Circulation, 2002, 106, 130), is increased (Li et al., Circulation, 1997, 96, 874). In the myocardial fibrotic area, it is known that mast cells are increased in number and may contribute to the development of fibroblast proliferation in cardiac tissues of patients with cardiomyopathy (Patella et al., Circulation, 1998, 97, 971). Chymase is a chymotrypsin-like serine protease contained in the secretory granules of mast cells. Although the precise physiological roles of Chymase have not been completely revealed, Chymase is known to transform angiotensin I to angiotensin II and may contribute to activation of TGF- β , matrix proteases, and cytokines (Taipale et al., J. Biol. Chem., 1995, 270, 4689; Takai et al., Life Sci., 1996, 58, 591; Takai et al., Circulation, 1999, 100, 654).

A potent and selective Chymase inhibitor may have potential use as a treatment of chronic heart failure, atherosclerosis, restenosis, and myocardial infarction by inhibiting local production of angiotensin II in the heart and release of TGF- β , two independent mediators of cardiac remodeling. An inhibitor may also have potential use for treatment of mast cell mediated diseases such as dermatitis, asthma, chronic obstructive pulmonary disease (COPD), and pulmonary inflammation, since Chymase is implicated in microvascular leakage, neutrophil

accumulation, the stimulation of mucus secretion, and the modulation of cytokines (He et al., Eur. J. Pharmacol., 1998, 352, 91).

Several small molecule Chymase inhibitors have been reported to be efficacious in the cardiomyopathic hamster model of heart failure (Takai et al. J. Pharmacol. Exp. Ther. 2003, 305, 17), in carotid artery injury by a balloon catheter in dogs (Takai et al. J. Pharmacol. Exp. Ther, 2003, 304, 841), and in the hamster left anterior descending coronary artery ligation model of heart failure (WO 03/018061). Additionally, a Chymase inhibitor has been demonstrated to be efficacious in a sheep asthma model (WO 2005/073214). However, there is no example of commercialization of a Chymase inhibitor as a medicament.

SUMMARY OF THE INVENTION


It is therefore an object of the invention to provide a small molecule a Chymase inhibitor as defined herein, and pharmaceutical compositions thereof.

It is also an object of the invention to provide methods of using said Chymase inhibitors to treat various diseases and conditions related thereto.

It is a further object of the invention to provide processes of preparing said Chymase inhibitors.

DETAILED DESCRIPTION OF THE INVENTION

In a first generic embodiment, there is provided a compound of the formula (I):



wherein:

A is C₁-C₈ linear or branched alkyl, C₂-C₈ linear or branched alkenyl, (CH₂)_m-aryl-(CH₂)_m, (CH₂)_m-C₃-C₈ cycloalkyl-(CH₂)_m or (CH₂)_m-C₃-C₈ cycloalkenyl-(CH₂)_m, each optionally interrupted at each possible occurrence by one or more -O-, -S-, -SO-, -SO₂- or -NRa-, and **A** is substituted with R₁ and further optionally substituted with 1 to 5 substituents independently chosen at each possible occurrence from C₁₋₅ linear or branched alkyl, C₂₋₅ linear or branched alkenyl, C₃-C₇ cycloalkyl, C₃-C₇ cycloalkenyl, C₁₋₄ alkoxy, C₁₋₄ alkoxycarbonyl, carboxyl, (C₁₋₄ alkyl)sulfonyl, hydroxyl, halogen, C₁-C₆ haloalkyl, trifluoromethylsulfonyl, trifluoromethoxy, difluoromethoxy, cyano, heterocycloalkyl, aryl, heteroaryl and carboxamido or sulfonamido wherein the nitrogen atom of the said carboxamide or sulfonamide is optionally independently mono or disubstituted with C₁₋₃ linear or branched alkyl, or the nitrogen atom of the said carboxamide or sulfonamide is part of a pyrrolidinyl, piperidinyl, morpholinyl or piperazinyl ring;

R₁ is -C(O)-R₄, -C(O)-N(R₅)-SO₂R₆, or -C(O)-NR₅R₇, wherein R₄ is chosen from hydroxyl, C₁-C₄ alkoxy and aryloxy; R₅ is chosen from hydrogen or C₁-C₃ alkyl; R₆ is chosen from aryl and heteroaryl; or R₆ is chosen from C₁-C₆ linear or branched alkyl, amino and C₃-C₆ cycloalkyl, each optionally independently substituted with 1-5 substituents chosen from C₁₋₅ alkyl, C₁₋₄ alkoxy, halogen, cyano, aryl and heteroaryl; R₇ is chosen from hydrogen, cyano, aryl and heteroaryl; or R₇ is chosen from C₁-C₆ linear or branched alkyl and C₃-C₆ cycloalkyl, each optionally independently substituted with 1-5 substituents chosen from C₁₋₅ alkyl, C₁₋₄ alkoxy, halogen, cyano, aryl and heteroaryl;

or

R₁ is -SO₂R₈ or -SO₂-NH-C(O)R₉, wherein R₈ is chosen from hydroxyl, amino, C₁-C₃ alkoxy, aryloxy, trifluoromethyl, C₁-C₃ alkylamino, C₁-C₃ dialkylamino, 1-pyrrolidino, 1-piperidino and 1-morpholino; R₉ is chosen from aryl and heteroaryl; or R₉ is chosen from C₁-C₆ linear or branched alkyl and C₃-C₆ cycloalkyl, each optionally independently substituted with 1-5 substituents chosen from C₁₋₅ alkyl, C₁₋₄ alkoxy, halogen, cyano, aryl and heteroaryl;

or

R₁ is -NH-C(O)-N(R₅)-SO₂R₁₀ wherein R₁₀ is chosen from C₁₋₅ alkyl, aryl and heteroaryl; or

R₁ is tetrazolyl or dioxothiadiazolidinone;

R₂ is aryl or heteroaryl;

R₃ is carboxyl, halogen, cyano, C₁₋₃ alkyl, -NHR₁₁, pyrrolidino, piperidino, morpholino, -C(O)-R₁₂ or -C(O)-NR₁₃R₁₄; R₁₁ is chosen from hydrogen, carboxamido, C₁-C₄ alkyl, C₁-C₄ alkoxy, carbonyl, aryl, benzyl and heteroaryl; R₁₂ is C₁-C₄ alkoxy; R₁₃ is heteroaryl, heterocyclyl or C₁-C₄ alkyl, each optionally substituted with 1-3 substituents chosen from aryl, heteroaryl, heterocyclyl, hydroxy, C₁₋₃ alkoxy, oxo, thioalkyl and dimethylamino; R₁₄ is chosen from hydrogen and C₁-C₄ alkyl optionally substituted with 1-2 hydroxyl groups ;

or R₁₃ and R₁₄ together with the nitrogen atom to which they are attached form a 5-6 membered heterocyclic ring optionally substituted with 1-3 substituents chosen from halogen, C₁₋₃ alkyl, C₁₋₃ alkoxy, hydroxyl, hydroxymethyl, carboxamido, N-acetyl-N-methyl amino and cyano;

n = 0-3

and

m = 0-6;

wherein all aryl or heteroaryl groups are optionally substituted with 1-5 substituents chosen from halogen, hydroxyl, cyano, C₁₋₆ linear or branched alkyl, C₁₋₆ linear or branched alkoxy, C₁₋₆ linear or branched alkylthio, C₁₋₆ linear or branched alkylsulfonyl, phenylsulfonyl, C₁₋₆ linear or branched acyl, C₁₋₆ linear or branched acylamino, polyhalomethyl, polyhalomethoxy, phenyl, oxo, -COORa, -CON(Rc)Ra, SO₂N(Rc)Ra, -N(Rc)Ra and phenoxy optionally substituted by one or more halogens;

Ra and Rc are each independently C1-3 alkyl or hydrogen;

with the proviso that

when **A** is C₁-C₃ linear or branched alkyl, and **R₁** is -C(O)-OH, -C(O)-NH₂, or -C(O)-OC₂CH₃ and **R₃** is lower alkyl, lower alkoxy or halogen, and **n** = 0-2,

then **R₂** is not unsubstituted naphthyl, unsubstituted thienyl, unsubstituted furanyl, or phenyl with 0-2 substituents selected from halogen, trifluoromethyl, lower alkyl or lower alkoxy;

or a pharmaceutically acceptable salt thereof.

In another embodiment, there is provided a compound as described herein above and wherein:

A is C₁-C₈ linear or branched alkyl, C₂-C₈ linear or branched alkenyl, (CH₂)_m-phenyl-(CH₂)_m, (CH₂)_m-C₃-C₆ cycloalkyl-(CH₂)_m, or (CH₂)_m-C₃-C₆ cycloalkenyl-(CH₂)_m;

and

A is substituted with R₁ and further optionally substituted with 1 to 5 substituents independently chosen at each possible occurrence from C₁₋₅ linear or branched alkyl, C₁₋₄ alkoxy, C₁₋₄ alkoxycarbonyl, carboxyl, hydroxyl, halogen, C₁-C₆ haloalkyl, cyano and carboxamido wherein the nitrogen atom of the said carboxamide is optionally independently mono or disubstituted with C₁₋₃ linear or branched alkyl; **A** is further optionally substituted with phenyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, thienyl, furanyl or tetrazolyl, each optionally independently substituted with 1-3 substituents chosen from halogen, hydroxyl, cyano, C₁₋₆ linear or branched alkyl, C₁₋₆ linear or branched alkoxy, C₁₋₆ linear or branched alkylthio, C₁₋₆ linear or branched alkylsulfonyl, phenylsulfonyl, C₁₋₆ linear or branched acyl, C₁₋₆ linear or branched acylamino, polyhalomethyl, polyhalomethoxy, phenyl, -COOR_a, -CON(R_c)R_a, SO₂N(R_c)R_a, -N(R_c)R_a and phenoxy optionally substituted by one or more halogens;

R_a and R_c are each independently C₁₋₃ alkyl or hydrogen;

R₁ is -C(O)-R₄, -C(O)-N(R₅)-SO₂R₆, or -C(O)-NR₅R₇, wherein R₄ is chosen from hydroxyl and C₁-C₄ alkoxy; R₅ is chosen from hydrogen and C₁-C₃ alkyl; R₆ is chosen from phenyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, thienyl and furanyl, each optionally independently substituted with 1-5 substituents chosen from C₁₋₅ alkyl, C₁₋₄ alkoxy, halogen, or cyano; or R₆ is chosen from C₁-C₆ linear or branched alkyl, amino and C₃-C₆ cycloalkyl; R₇ is chosen from hydrogen, cyano and benzyl;

or

R₁ is -SO₂R₈ or -SO₂-NH-C(O)R₉, wherein R₈ is chosen from hydroxyl, amino, C₁-C₃ alkoxy, phenoxy, C₁-C₃ alkylamino and C₁-C₃ dialkylamino; R₉ is chosen from phenyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, thienyl

and furanyl, each optionally independently substituted with 1-5 substituents chosen from C₁₋₅ alkyl, C₁₋₄ alkoxy, halogen and cyano; or R₉ is chosen from C₁-C₆ linear or branched alkyl and C₃-C₆ cycloalkyl;

or

R₁ is -NH-C(O)-N(R₅)-SO₂R₁₀ wherein R₁₀ is chosen from C₁₋₅ alkyl and phenyl;

or

R₁ is tetrazolyl or dioxothiadiazolidinone;

R₂ is phenyl, naphthyl or a heteroaryl ring chosen from benzothienyl, benzothiazolyl, benzoisothiazolyl, benzoxazolyl, benzisoxazolyl, indolyl, indoliziny, dihydroindolyl, pyrazolopyridinyl, benzoxazoliny and benzothiazoliny, wherein each ring is optionally independently substituted with 1-5 substituents chosen from halogen, C₁₋₃ alkyl, C₁₋₃ alkoxy, hydroxy, cyano, oxo, C₁₋₃ alkoxycarbonyl, amino, trifluormethyl, difluoromethoxy and trifluoromethoxy;

R₃ is carboxyl, halogen, cyano, C₁₋₃ alkyl, -NHR₁₁, -C(O)-R₁₂ or -C(O)-NR₁₃R₁₄; R₁₁ is chosen from hydrogen, carboxamido, C₁-C₄ alkyl, C₁-C₄ alkoxycarbonyl, phenyl, thiazolyl, oxazolyl, imidazolyl, thienyl and furanyl; R₁₂ is C₁-C₄ alkoxy; R₁₃ is 1,1-dioxotetrahydrothienyl, oxotetrahydrofuranly or C₁-C₃ alkyl, each optionally substituted with 1-3 substituents chosen from phenyl optionally mono- or di-substituted with methoxy, hydroxyl, thiomethoxy or halogen, tetrahydrofuranly, dioxolanyl, hydroxyl, C₁₋₃ alkoxy, thioalkyl and dimethylamino; R₁₄ is chosen from hydrogen and C₁-C₄ alkyl optionally substituted with 1-2 hydroxyl groups ; or R₁₃ and R₁₄ together with the nitrogen atom to which they are attached form a 5-6 membered heterocyclic ring chosen from piperidinyl and pyrrolidinyl, optionally substituted with 1-3 substituents chosen from halogen, C₁₋₃ alkyl, C₁₋₃ alkoxy, hydroxyl, hydroxymethyl, carboxamido, N-acetyl-N-methyl amino and cyano;

and

n = 0-3

m = 0-3.

In another embodiment, there is provide a compound as described hereinabove and wherein:

A is C₁-C₄ linear or branched alkyl, C₂-C₄ linear or branched alkenyl, (CH₂)_m-phenyl, (CH₂)_m - C₃-C₆ cycloalkyl-(CH₂)_m, or cyclopentenyl;

and

A is substituted with R₁ and further optionally substituted with 1 to 3 substituents independently chosen at each possible occurrence from C₁₋₅ linear or branched alkyl, C₁₋₄ alkoxy, C₁₋₄ alkoxycarbonyl, carboxyl, hydroxyl, C₁-C₆ haloalkyl, phenyl, pyridinyl and carboxamido wherein the nitrogen atom of the said carboxamide is optionally independently mono or disubstituted with C₁₋₃ linear or branched alkyl;

R₁ is -C(O)-OH, -C(O)-N(R₅)-SO₂R₆, or -C(O)-NR₅R₇; R₅ is chosen from hydrogen and methyl; R₆ is chosen from phenyl, pyridyl, imidazolyl, each optionally independently substituted with 1-3 substituents chosen from C₁₋₅ alkyl, C₁₋₄ alkoxy and halogen; or R₆ is chosen from C₁-C₆ linear or branched alkyl, amino and C₃-C₆ cycloalkyl; R₇ is chosen from hydrogen, cyano and benzyl;

or

R₁ is -SO₂R₈ or -SO₂-NH-C(O)R₉, wherein R₈ is chosen from hydroxyl, amino and C₁-C₃ alkylamino; R₉ is chosen from C₁₋₄ alkyl and phenyl;

or

R₁ is -NH-C(O)-N(R₅)-SO₂R₁₀ wherein R₁₀ is phenyl;

or

R₁ is tetrazolyl or dioxothiadiazolidinone;

R₂ is phenyl, naphthyl or a heteroaryl ring chosen from benzothienyl, benzoisothiazolyl, indolyl, indoliziny, dihydroindolyl, benzoxazoliny and benzothiazoliny, wherein each ring is optionally independently substituted with 1-4 substituents chosen from halogen, C₁₋₃ alkyl, methoxy, hydroxy, cyano, oxo and trifluoromethyl;

R₃ is halogen, cyano, methyl, -NHR₁₁, -C(O)-R₁₂ or -C(O)-NR₁₃R₁₄; R₁₁ is chosen from carboxamido, C₁-C₄ alkoxycarbonyl, phenyl and thiazolyl; R₁₂ is C₁-C₄ alkoxy; R₁₃ is 1,1-

dioxotetrahydrothienyl, oxotetrahydrofuranyl or C₁-C₃ alkyl, each optionally substituted with 1-3 substituents chosen from phenyl optionally mono- or di-substituted with methoxy, hydroxyl, thiomethoxy or halogen, tetrahydrofuranyl, dioxolanyl and hydroxy; R₁₄ is hydrogen; or R₁₃ and R₁₄ together with the nitrogen atom to which they are attached form a piperidinyl or pyrrolidinyl ring optionally substituted with hydroxyl, hydroxymethyl, carboxamido or N-acetyl-N-methyl amino;

n = 0-2

and

m = 0-2.

In another embodiment, there is provide a compound as described hereinabove and wherein:

A is C₁-C₄ linear or branched alkyl, ethenyl, phenyl, (CH₂)_m-C₃-C₆ cycloalkyl-(CH₂)_m or cyclopentenyl;

and

A is substituted with R₁ and further optionally substituted with 1 to 3 substituents independently chosen at each possible occurrence from C₁₋₅ linear or branched alkyl, C₁₋₄ alkoxy, C₁₋₄ alkoxy carbonyl, carboxyl, hydroxyl, phenyl and pyridinyl;

R₁ is -C(O)-OH, -C(O)-N(R₅)-SO₂R₆ or -C(O)-NR₅R₇; R₅ is hydrogen; R₆ is chosen from phenyl, pyridyl and imidazolyl, each optionally independently substituted with 1-3 substituents chosen from methyl, methoxy and fluoro; or R₆ is chosen from C₁-C₆ linear or branched alkyl, amino and C₃-C₆ cycloalkyl; R₇ is cyano;

or

R₁ is -SO₃H or -SO₂-NH-C(O)R₉; R₉ is chosen from methyl and phenyl;

or

R₁ is -NH-C(O)-N(R₅)-SO₂R₁₀ wherein R₁₀ is phenyl;

or

R₁ is tetrazolyl or dioxothiadiazolidinone;

R₂ is phenyl, naphthyl or a heteroaryl ring chosen from benzothienyl, benzoisothiazolyl, indolyl, indolizinyll and dihydroindolyl, wherein each ring is optionally independently substituted with 1-4 substituents chosen from halogen, C₁₋₂ alkyl, hydroxy, cyano and oxo;




R₃ is bromo, methyl or -NHR₁₁, R₁₁ is thiazolyl;







n = 0-2 and







m = 0-1.








In another embodiment, there is provided compound as described in Table I which can be made as described in the schemes and examples herein below, and by methods apparent to those of ordinary skill in the art:







Table I








	N-[(2-fluorophenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanamide
	3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)hexanamide
	N-[(4-fluorophenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanamide







	(3S)-3-{6-methyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-5-phenylpentanoic acid
	3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(pyridin-2-ylsulfonyl)hexanamide
	N-[(2-methoxyphenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide
	3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}octanoic acid
	3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-5,6-dimethyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-ethoxy-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)propanamide







	3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}heptanoic acid
	N-[(3-fluorophenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide
	3-{6-methyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-ethoxy-3-{6-methyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-{5,6-dimethyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-ethoxypropanoic acid
 Chiral	(3S)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid







	N-[(4-methoxyphenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide
	3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-ethoxy-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	N-[(3-methoxyphenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide
	(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(pyridin-3-ylsulfonyl)hexanamide








	3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	N-(cyclopropylsulfonyl)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide
	5-methoxy-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-5-oxopentanoic acid
	3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-ethoxypropanoic acid
	5-methyl-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)propanamide







	3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanedioic acid
	3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-methoxy-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(methylsulfonyl)hexanamide
	(3R)-3-{6-methyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-5-phenylpentanoic acid
	N-[(2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}ethyl)sulfonyl]benzamide







	3-ethoxy-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	(3R)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	(3R)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid
	3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid
	3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid







	2,2-dimethyl-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	5-hydroxy-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid
	3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid
	5-methyl-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid
	(2E)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}acrylic acid
	3-[3-(1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid






	3-{2-oxo-3-[(1,2,3,5-tetramethyl-1H-indol-7-yl)methyl]-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-{3-[(8-bromo-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid
	N-(isopropylsulfonyl)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanamide
	N-[(2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}ethyl)sulfonyl]acetamide






	3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(methylsulfonyl)propanamide
	3-methoxy-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	3-{3-[(5-fluoro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	4-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid
	{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}acetic acid
	2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}ethanesulfonic acid
	3-{2-oxo-3-[(1,2,3-trimethyl-1H-indol-7-yl)methyl]-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid






	1-[(4-methyl-1-benzothien-3-yl)methyl]-3-[2-(1H-tetrazol-5-yl)ethyl]-1,3-dihydro-2H-benzimidazol-2-one
	3-{2-oxo-3-[(1,2,3-trimethyl-1H-indol-7-yl)methyl]-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-(6-{[2-(3,4-dimethoxyphenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid
	3-{3-[(8-bromo-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-(6-{[2-(3-hydroxy-4-methoxyphenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid






	3-(6-{[2-(2-methoxyphenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid
	3-{3-[(4-chloro-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid
	3-({3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} methyl)benzoic acid
	3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]pentanoic acid
	3-[6-{[2-(3,4-dimethoxyphenyl)ethyl]carbamoyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	4-methyl-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]pentanoic acid






	3-[3-(2,5-dimethylbenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid
	trans-4-({3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)cyclohexanecarboxylic acid
	3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]-N-(phenylsulfonyl)propanamide
	3-[6-{[2-(3-hydroxy-4-methoxyphenyl)ethyl]carbamoyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	3-{6-(benzylcarbamoyl)-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-{6-[(2-hydroxyethyl)carbamoyl]-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid






	N-methyl-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)hexanamide
	3-(6-{[2-(4-methoxyphenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid
	(2E)-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]acrylic acid
	3-[6-{[2-(2-methoxyphenyl)ethyl]carbamoyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	3-{6-[(3-hydroxypyrrolidin-1-yl)carbonyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid







	3-(6-{[2-(3-methoxyphenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid
	3-{3-[5-chloro-2-(trifluoromethyl)benzyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(4-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-{3-(1-naphthylmethyl)-2-oxo-6-[(tetrahydrofuran-2-ylmethyl)carbamoyl]-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-{6-[(2-hydroxyethyl)carbamoyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid







	3-{[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]methyl} benzoic acid
	3-[6-{[2-(1,3-dioxolan-2-yl)ethyl]carbamoyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	3-[6-{[2-(hydroxymethyl)piperidin-1-yl]carbonyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	3-{3-[(1-methyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} propanoic acid
	3-[6-({3-[acetyl(methyl)amino]pyrrolidin-1-yl} carbonyl)-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid






	3-(6-{[2-(2-fluorophenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid
	3-[6-{[(2S)-2-carbamoylpyrrolidin-1-yl]carbonyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	3-{6-[(3-carbamoylpiperidin-1-yl)carbonyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-{3-[(1,3-dimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-(6-{[2-(2-chlorophenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid






	3-{3-[(5-bromo-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{6-[(1,1-dioxidotetrahydro-3-thienyl)carbamoyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-(6-{[2-(4-chlorophenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid
	3-[6-cyano-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}ethanesulfonamide







	N-(methylsulfonyl)-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanamide
	3-{3-[(1,2-dimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	N-benzyl-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide
	3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-6-[(2-phenylethyl)carbamoyl]-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-{3-[(5-chloro-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid







	3-{3-[(4-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-(6-{[2-(3-fluorophenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid
	3-{3-[(4-bromo-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(6-hydroxy-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-{3-(1-naphthylmethyl)-2-oxo-6-[(2-oxotetrahydrofuran-3-yl)carbamoyl]-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-(6-{[2-(3-chlorophenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid






	3-[3-(3,5-dichlorobenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid
	3-[6-({(2S)-2-hydroxy-1-(hydroxymethyl)-2-[4-(methylthio)phenyl]ethyl} carbamoyl)-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	2-({3-[2-bromo-5-(trifluoromethyl)benzyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} methyl)pentanoic acid
	3-[6-{{[(2S)-2-(hydroxymethyl)pyrrolidin-1-yl]carbonyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	3-[3-(1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	5-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]pentanoic acid







	3-{3-[(4-chloro-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
Chiral 	3-[6-{[(1R,2S)-2-hydroxy-1-(hydroxymethyl)-2-phenylethyl]carbamoyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	4-{3-[(5-chloro-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid
	3-{3-[(6-methoxy-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
Chiral 	3-[6-{[(1S,2R)-2-hydroxy-1-(hydroxymethyl)-2-phenylethyl]carbamoyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid







	3-{6-[(2-hydroxy-1-phenylethyl)carbamoyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-[6-{[1-(hydroxymethyl)-3-(methylthio)propyl]carbamoyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	3-[5-cyano-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	3-{6-[(2-methoxybenzyl)carbamoyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-[3-(1-benzothien-3-ylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid

	3-{6-[bis(2-hydroxyethyl)carbamoyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-{3-[(1,3-dimethyl-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-{3-[(4-methylpyrazolo[1,5-a]pyridin-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-[3-(5-bromo-2-methoxybenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid
	3-[3-(1,2-benzisothiazol-3-ylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	3-(3-{[6-(methoxycarbonyl)-1,2,3-trimethyl-1H-indol-7-yl]methyl}-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid







	3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanamide
	3-{3-[(5-bromo-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-[3-(1,2-benzisoxazol-3-ylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid
	3-{3-[5-chloro-2-(difluoromethoxy)benzyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	phenyl 2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}ethanesulfonate
	3-[3-(1-benzothien-7-ylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid








	3-{3-[(1-methyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-[6-{2-(dimethylamino)ethyl}carbonyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	3-[6-(tert-butoxycarbonyl)-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	3-[5-(methoxycarbonyl)-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
	3-{3-[2-bromo-5-(trifluoromethyl)benzyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid








	3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}octanoic acid
	4-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)butanamide
	2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)acetamide
	3-{5-bromo-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{5-anilino-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)hexanamide








	N-(tert-butylsulfonyl)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide
	3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(ethylsulfonyl)hexanamide
	3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-5-(1,3-thiazol-2-ylamino)-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
 Chiral	(2S)-2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
 Chiral	(2S)-3-methyl-2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid
 Chiral	(3R)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-phenylpropanoic acid







37








	2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclopent-1-ene-1-carboxylic acid
 Chiral	(1R,2R)-2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclohexanecarboxylic acid
	3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-methylbutanoic acid
	3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-[(1-methyl-1H-imidazol-4-yl)sulfonyl]hexanamide
	N-[(1,2-dimethyl-1H-imidazol-4-yl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide
	2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclopentanecarboxylic acid








	3-{3-[(5-methoxy-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
 Chiral	(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)hexanamide
	3-{3-[(1-cyano-5-methylindolizin-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-7-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
 Chiral	(3S)-N-[(1,2-dimethyl-1H-imidazol-4-yl)sulfonyl]-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide
 Chiral	(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(pyridin-2-ylsulfonyl)hexanamide
	3-[2-oxo-3-(2,3,5-trichlorobenzyl)-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid







	3-{3-[(4-fluoro-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid
	3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2,2-dimethylbutanoic acid
 Chiral	(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(methylsulfonyl)hexanamide
	1-(1-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} butyl)cyclopropanecarboxylic acid
	3-{3-[(1-ethyl-4-methyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid
 Chiral	(3S)-N-(aminosulfonyl)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanamide
 Chiral	(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-[(1-methyl-1H-imidazol-4-yl)sulfonyl]hexanamide








 Chiral	(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(isopropylsulfonyl)hexanamide
	3-{3-[(1-cyano-5-methylindolizin-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-{3-[(1-cyano-5,7-dimethylindolizin-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(5-bromo-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
 Chiral	(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(ethylsulfonyl)hexanamide
 Chiral	(3S)-N-(tert-butylsulfonyl)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide
	(1-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclopentyl)acetic acid








	(1-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclohexyl)acetic acid
	3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2,2-dimethylhexanoic acid
	3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-4-methoxybutanoic acid
 Chiral	(3S)-N-cyano-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide
	3-[3-(2,5-dichlorobenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid
	3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-4-methoxy-N-[(1-methyl-1H-imidazol-4-yl)sulfonyl]butanamide








	2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentane-1-sulfonic acid
 Chiral	(3S)-N-(cyclopropylsulfonyl)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide
	3-{3-[(2,3-dimethylindolizin-8-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}benzoic acid
	3-{3-[(4-chloro-1-methyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(6,8-dichloro-3-methylindolizin-1-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(3,8-dimethylindolizin-1-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid








	3-{3-[(4-methyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid
	2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-methoxybenzoic acid
	2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-methylbenzoic acid
	1-[2-(1,1-dioxido-4-oxo-1,2,5-thiadiazolidin-2-yl)ethyl]-3-[(4-methyl-1-benzothien-3-yl)methyl]-1,3-dihydro-2H-benzimidazol-2-one
 Chiral	(3S)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-4-phenylbutanoic acid
 Chiral	(2S)-2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid
 Chiral	(2S)-2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid








 Chiral	(2S)-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} (phenyl)acetic acid
 Chiral	(3S)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-phenylpropanoic acid
 Chiral	(3S)-3-{5-(carbamoylamino)-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid
 Chiral	(3R)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-phenylpropanoic acid
	3-{3-[(5-bromo-1-methyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid
	3-{2-oxo-3-[(1,4,6-trimethyl-1H-indol-3-yl)methyl]-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid







	3-{3-[(3-bromo-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid
	3-{3-[(6-bromo-1-methyl-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid
	3-{7-bromo-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} propanoic acid
	3-{3-[(5-bromo-1-methyl-2-oxo-2,3-dihydro-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid
	3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-4-methoxybenzoic acid
	2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2-methylpropanoic acid
 Chiral	(3S)-3-{2-oxo-3-[(1,4,6-trimethyl-1H-indol-3-yl)methyl]-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid








	3-[3-(3,5-dibromobenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid
	3-{3-[(5-chloro-1,3-dimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-4-methoxy-2,2-dimethylbutanoic acid
	3-{3-[(6-bromo-1-methyl-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	2-({3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)pentanoic acid
	2-({3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)pentanoic acid
	3-{3-[(6-bromo-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid



	3-{3-[2-methyl-5-(trifluoromethyl)benzyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-[3-(5-chloro-2-methoxybenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid
	3-{3-[(1-methyl-2-oxo-2,3-dihydro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(6-bromo-1-methyl-2-oxo-2,3-dihydro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(4,6-dimethyl-1,2-benzisothiazol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(5-chloro-1-methyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	2-({3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)butanoic acid




	2-({3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} methyl)butanoic acid
	3-{3-[(6-chloro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid
	1-({3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} methyl)cyclohexanecarboxylic acid
	1-({3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} methyl)cyclohexanecarboxylic acid
	3-[3-(2,3-dichlorobenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid
	3-[3-(2-chloro-5-methylbenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid
	3-[3-(5-bromo-2-methylbenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid





	3-[3-(5-chloro-2-methylbenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid
	3-[3-(3-hydroxy-2,5-dimethylbenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid
	3-{3-[(6-methyl-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(6-chloro-1-methyl-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2-methylpropanoic acid
	3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2-methylpropanoic acid
 Chiral	(3S)-3-{3-[(4,6-dimethyl-1,2-benzisothiazol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid





	2-({3-[(4,6-dimethyl-1,2-benzisothiazol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)pentanoic acid
	1-({3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)cyclopentanecarboxylic acid
	1-({3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)cyclopentanecarboxylic acid
	3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2-phenylpropanoic acid
 Chiral	(3R)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-pyridin-3-ylpropanoic acid
	3-{3-[(6-bromo-2-oxo-2,3-dihydro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid





	3-{3-[(5-chloro-2-oxo-2,3-dihydro-1,3-benzoxazol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[4-amino-3-(trifluoromethoxy)benzyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(6-chloro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-{3-[(6-chloro-2-oxo-2,3-dihydro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-{3-[(6-chloro-2-oxo-2,3-dihydro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
	3-{3-[(6-bromo-2-oxo-2,3-dihydro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
	3-{3-[(6-bromo-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid





 Chiral	(1R,3S)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclopentanecarboxylic acid
 Chiral	methyl (1R,3S)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclopentanecarboxylate





	3-[3-(1,4-Dimethyl-1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-6,6,6-trifluorohexanoic acid
 Chiral	(R)-3-(4-Chloro-phenyl)-3-[3-(1,4-dimethyl-1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid
 Chiral	(R)-3-(4-Cyano-phenyl)-3-[3-(1,4-dimethyl-1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid





 Chiral	(R)-3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-(4-methoxy-phenyl)-propionic acid
 Chiral	(R)-3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-(4-fluoro-phenyl)-propionic acid
 Chiral	(S)-3-[3-(5-Bromo-2-oxo-2,3-dihydro-benzothiazol-7-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid
	2-[3-(5-Chloro-1,2,3-trimethyl-1 <i>H</i> -indol-7-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid





	2-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-ylmethyl]-hexanoic acid
	3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2-phenyl-propionic acid
	3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2-hydroxymethyl-propionic acid
	1-[3-(4,6-Dimethyl-1,2-benzisothiazol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-ylmethyl]-cyclohexanecarboxylic acid





	3-[7-Bromo-3-(4-methylbenzo[b]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid
	(R)-3-[3-(4,6-Dimethyl-1,2-benzisothiazol-3-ylmethyl)-6-methyl-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid
	(S)-3-[3-(4,6-Dimethyl-1,2-benzisothiazol-3-ylmethyl)-6-methyl-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid
	(R)-3-[3-(5-Chloro-1,2,3-trimethyl-1H-indol-7-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid





	3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-(2-fluoro-phenyl)-propionic acid
	3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3- <i>o</i> -tolyl-propionic acid
	2-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-ylmethyl]-3-hydroxy-3-phenyl-propionic acid
	3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2-morpholin-4-ylmethyl-propionic acid





	3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3- <i>m</i> -tolyl-propionic acid
	2-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-ylmethyl]-3-methoxy-3-phenyl-propionic acid
	3-[7-Bromo-3-(4,6-dimethyl-1,2-benzisothiazol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid
	2-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-ylmethyl]-3-hydroxy-butyric acid





	3-[3-(1,4-Dimethyl-1<i>H</i>-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2-piperidin-1-ylmethyl-propionic acid
	3-[3-(1,4-Dimethyl-1<i>H</i>-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2-(4-methyl-piperazin-1-ylmethyl)-propionic acid
	3-[3-(1,4-Dimethyl-1<i>H</i>-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-<i>p</i>-tolyl-propionic acid
	(<i>S</i>)-3-[3-(4,6-Dimethyl-1,2-benzisothiazol-3-ylmethyl)-6-ethoxy-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid





	3-[3-(4,6-Dimethyl-1,2-benzisothiazol-3-ylmethyl)-2-oxo-7-phenyl-2,3-dihydro-benzimidazol-1-yl]-propionic acid
	3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-(4-fluoro-phenyl)-propionic acid
	2-Benzyl-3-[3-(1,4-dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid
	3-[3-(3-Isopropoxy-2,5-dimethyl-benzyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid





	(S)-3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-6-ethoxy-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid
	2-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-ylmethyl]-3-hydroxy-pentanoic acid
	2-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-ylmethyl]-3-hydroxy-hexanoic acid
	3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-(4-methoxy-phenyl)-propionic acid





	3-[3-(1,4-Dimethyl-1<i>H</i>-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-(3-methoxy-phenyl)-propionic acid
	4-[3-(1,4-Dimethyl-1<i>H</i>-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-ylmethyl]-tetrahydro-pyran-4-carboxylic acid
	3-{1-[3-(1,4-Dimethyl-1<i>H</i>-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethyl}-benzoic acid
	(<i>R</i>)-3-[2-Oxo-3-(1,4,6-trimethyl-1<i>H</i>-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid





	2-[2-Oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-ylmethyl]-pentanoic acid
	3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-(3-fluoro-phenyl)-propionic acid
	3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-(4-trifluoromethyl-phenyl)-propionic acid
	3-(4-Chloro-phenyl)-2-[3-(1,4-dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-ylmethyl]-3-hydroxy-propionic acid





	3-[7-Bromo-3-(1,4-dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid
	3-Cyclopropyl-3-[3-(1,4-dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid
	3-Cyclopropyl-3-[3-(1,4-dimethyl-1 <i>H</i> -indol-3-ylmethyl)-7-methyl-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid
	2-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-ylmethyl]-3-hydroxy-3-thiazol-2-yl-propionic acid





	(S)-3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid (2-hydroxy-ethyl)-amide
	3-Cyclopropyl-3-[3-(4,6-dimethyl-1,2-benzisothiazol-3-ylmethyl)-7-methyl-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid
	3-Cyclopropyl-3-[3-(4,6-dimethyl-1,2-benzisothiazol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid
	3-[3-(4,6-Dimethyl-1,2-benzisothiazol-3-ylmethyl)-7-methyl-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid





	3-[7-Bromo-3-(4,6-dimethyl-1,2-benzisothiazol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid
	3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-(4-hydroxy-phenyl)-propionic acid
	3-{1-[3-(4,6-Dimethyl-1,2-benzisothiazol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethyl}-benzoic acid
	3-{1-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-butyl}-benzoic acid





	3-{1-[3-(4,6-Dimethyl-1,2-benzisothiazol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-butyl}-benzoic acid
	(R)-3-[5-Cyano-3-(1,4-dimethyl-1<i>H</i>-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid
	(R)-3-[5-Cyano-3-(4,6-dimethyl-1,2-benzisothiazol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid
	(R)-3-[6-Cyano-3-(1,4-dimethyl-1<i>H</i>-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid





	(R)-3-[6-Cyano-3-(4,6-dimethyl-1,2-benzisothiazol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid
	3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2-pyridin-3-yl-propionic acid
	3-[3-(5-Methoxy-1-methyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid
	(R)-3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-6-(2-hydroxy-ethylcarbamoyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid ethyl ester





	(R)-3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-6-(2-hydroxyethylcarbamoyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid
	(R)-3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-6-(2-hydroxy-2-methylpropylcarbamoyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid
	(R)-3-[6-Carbamoyl-3-(1,4-dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid
	3-((R)-2-Carboxy-1-phenyl-ethyl)-1-(1,4-dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-1 <i>H</i> -benzimidazole-5-carboxylic acid <i>tert</i> -butyl ester





	1-(1,4-Dimethyl-1<i>H</i>-indol-3-ylmethyl)-3-((<i>R</i>)-2-ethoxycarbonyl-1-phenylethyl)-2-oxo-2,3-dihydro-1<i>H</i>-benzimidazole-5-carboxylic acid
	(<i>R</i>)-3-[3-(1,4-Dimethyl-1<i>H</i>-indol-3-ylmethyl)-6-methylcarbamoyl-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid
	(<i>R</i>)-3-[3-(1,4-Dimethyl-1<i>H</i>-indol-3-ylmethyl)-2-oxo-6-(5-oxo-tetrahydrofuran-3-ylcarbamoyl)-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid ethyl ester
	(<i>S</i>)-3-[6-Methylcarbamoyl-2-oxo-3-(1,4,6-trimethyl-1<i>H</i>-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid





	(R)-3-[6-Dimethylcarbamoyl-3-(1,4-dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid
	(R)-3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-6-(ethyl-methyl-carbamoyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid
	3-[2-Oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-2-pyridin-3-yl-propionic acid
	(R)-3-[6-Cyano-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid





	1-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-3-(6-oxo-1,6-dihydro-pyridin-3-yl)-1,3-dihydro-benzimidazol-2-one
	1-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-3-(2-oxo-1,2-dihydro-pyridin-4-ylmethyl)-1,3-dihydro-benzimidazol-2-one
	1-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-3-(6-oxo-1,6-dihydro-pyridin-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one
	(R)-3-[6-Methylcarbamoyl-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid





	(R)-3-[6-(2-Hydroxy-ethylcarbamoyl)-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid
	(R)-3-[6-(2-Hydroxy-2-methyl-propylcarbamoyl)-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid
	(R)-3-[6-Carbamoyl-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid
	(R)-3-[6-(Ethyl-methyl-carbamoyl)-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid





	(3-Naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-acetic acid
	3-((S)-1-Carboxymethyl-butyl)-2-oxo-1-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-1 <i>H</i> -benzimidazole-5-carboxylic acid <i>tert</i> -butyl ester
	1-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-ylmethyl]-cyclohexanecarboxylic acid amide
	3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2-hydroxy-propionic acid methyl ester





	3-[3-(1,4-Dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2-hydroxy-propionic acid
	3-[3-(4,6-Dimethyl-1,2-benzisothiazol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2-pyridin-3-yl-propionic acid
	(R)-3-[6-Cyano-3-(1,4-dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-(4-fluoro-phenyl)-propionic acid
	(R)-3-[6-Cyano-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-(4-fluoro-phenyl)-propionic acid





	2-Oxo-1,3-bis-(1,4,6-trimethyl-1<i>H</i>-indazol-3-ylmethyl)-2,3-dihydro-1<i>H</i>-benzimidazole-5-carbonitrile
	(<i>R</i>)-3-[6-Methanesulfonylaminocarbonyl-2-oxo-3-(1,4,6-trimethyl-1<i>H</i>-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid ethyl ester
	(<i>R</i>)-3-[6-Methanesulfonylaminocarbonyl-2-oxo-3-(1,4,6-trimethyl-1<i>H</i>-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid
	(<i>S</i>)-3-[6-Methanesulfonylaminocarbonyl-2-oxo-3-(1,4,6-trimethyl-1<i>H</i>-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester





	(S)-3-[6-Methanesulfonylamino-2-oxo-3-(1,4,6-trimethyl-1<i>H</i>-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid
	3-[6-Cyano-3-(1,4-dimethyl-1<i>H</i>-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid
	3-[6-Cyano-2-oxo-3-(1,4,6-trimethyl-1<i>H</i>-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid
	1,3-Bis-(1,4-dimethyl-1<i>H</i>-indol-3-ylmethyl)-2-oxo-2,3-dihydro-1<i>H</i>-benzimidazole-5-carbonitrile





	(R)-3-(4-Chloro-phenyl)-3-[6-cyano-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-propionic acid
	(S)-3-[6-Cyano-3-(1,4-dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid
	(S)-3-[6-Cyano-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid
	(R)-3-[6-Chloro-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid



 Chiral	(S)-3-[6-Cyano-3-(1,4-dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid
 Chiral	(S)-3-[6-Cyano-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid
	3-[6-Cyano-3-(1,4-dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-heptanoic acid
	3-[6-Cyano-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-heptanoic acid

	3-[6-Cyano-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-cyclopropyl-propionic acid
	(R)-3-(6-Methoxy-pyridin-3-yl)-3-[6-methyl-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-propionic acid
	(S)-3-[6-Cyano-2-oxo-3-(1,4,5-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid
	(S)-3-[6-Cyano-2-oxo-3-(1,5,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid

	(S)-2-[6-Cyano-3-(1,4-dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-butyric acid
	(S)-2-[6-Cyano-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-butyric acid
	(R)-3-[6-Cyano-2-oxo-3-(1,4,5-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-(4-fluorophenyl)-propionic acid
	3-[6-Cyano-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-4,4-dimethyl-pentanoic acid

Chiral 	(R)-3-[6-Cyano-2-oxo-3-(1,5,6-trimethyl-1<i>H</i>-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-(4-fluorophenyl)-propionic acid
Chiral 	(S)-[6-Cyano-3-(1,4-dimethyl-1<i>H</i>-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-cyclohexyl-acetic acid
Chiral 	(S)-[6-Cyano-2-oxo-3-(1,4,6-trimethyl-1<i>H</i>-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-cyclohexyl-acetic acid
Chiral 	(S)-2-[6-Cyano-3-(1,4-dimethyl-1<i>H</i>-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid

	(S)-2-[6-Cyano-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid
	(S)-2-[6-Cyano-3-(1,4-dimethyl-1 <i>H</i> -indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid
	(S)-2-[6-Cyano-2-oxo-3-(1,4,6-trimethyl-1 <i>H</i> -indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid
	2-[3-(5-Bromo-2-oxo-2,3-dihydro-benzothiazol-7-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid

	2-[3-(6-Bromo-2-oxo-2,3-dihydro-1 <i>H</i> -indol-4-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-ylmethyl]-pentanoic acid
	2-[3-(6-Bromo-1 <i>H</i> -indol-4-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-ylmethyl]-pentanoic acid

or a pharmaceutically acceptable salt thereof.

The following are preferred compounds of the invention:

Table II

N-[(2-fluorophenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}hexanamide

(3*R*)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}-3-phenylpropanoic acid

N-[(1,2-dimethyl-1*H*-imidazol-4-yl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}hexanamide

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}-N-(phenylsulfonyl)hexanamide

(3*S*)-N-[(1,2-dimethyl-1*H*-imidazol-4-yl)sulfonyl]-3-{3-[(1,4-dimethyl-1*H*-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}hexanamide

(3*S*)-3-{3-[(1,4-dimethyl-1*H*-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}-N-(pyridin-2-ylsulfonyl)hexanamide

2-(3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)methylpentanoic acid

3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2-phenylpropanoic acid

(2S)-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}(phenyl)acetic acid

N-[(4-fluorophenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-[(1-methyl-1H-imidazol-4-yl)sulfonyl]hexanamide

(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-[(1-methyl-1H-imidazol-4-yl)sulfonyl]hexanamide

(3R)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-phenylpropanoic acid

(3S)-3-{6-methyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-5-phenylpentanoic acid

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)hexanamide

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(pyridin-2-ylsulfonyl)hexanamide

N-[(2-methoxyphenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}octanoic acid

(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)hexanamide

(3R)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-pyridin-3-ylpropanoic acid

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}octanoic acid

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-5,6-dimethyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

3-ethoxy-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)propanamide

(3S)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-phenylpropanoic acid

(3S)-N-(tert-butylsulfonyl)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide

2-({3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)pentanoic acid

(3S)-3-{2-oxo-3-[(1,4,6-trimethyl-1H-indol-3-yl)methyl]-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}heptanoic acid

N-[(3-fluorophenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide

3-{6-methyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

3-ethoxy-3-{6-methyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

3-{5,6-dimethyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-ethoxypropanoic acid

(3S)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

3-{2-oxo-3-[(1,4,6-trimethyl-1H-indol-3-yl)methyl]-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

N-[(4-methoxyphenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

N-(tert-butylsulfonyl)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2,2-dimethylhexanoic acid

1-({3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)cyclohexanecarboxylic acid

3-ethoxy-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

N-[(3-methoxyphenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide

3-{3-[(6-bromo-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

2-({3-[(4,6-dimethyl-1,2-benzisothiazol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} methyl)pentanoic acid

(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-4-methoxy-N-[(1-methyl-1H-imidazol-4-yl)sulfonyl]butanamide

3-{3-[(6-chloro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid

(3S)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-4-phenylbutanoic acid

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(pyridin-3-ylsulfonyl)hexanamide

2-({3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} methyl)butanoic acid

2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} pentane-1-sulfonic acid

(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(isopropylsulfonyl)hexanamide

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-5-(1,3-thiazol-2-ylamino)-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid

1-({3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} methyl)cyclohexanecarboxylic acid

N-(cyclopropylsulfonyl)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanamide

5-methoxy-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-5-oxopentanoic acid

(3S)-N-cyano-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanamide

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-ethoxypropanoic acid

3-{5-bromo-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid

5-methyl-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(ethylsulfonyl)hexanamide

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)propanamide

(3S)-N-(cyclopropylsulfonyl)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanedioic acid

3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

1-(1-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butyl)cyclopropanecarboxylic acid

3-{3-[(3-bromo-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

3-{3-[(1-cyano-5,7-dimethylindolizin-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

N-[(2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}ethyl)carbamoyl]benzenesulfonamide

3-methoxy-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(methylsulfonyl)hexanamide

(1-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclohexyl)acetic acid

3-{3-[(5-chloro-1,3-dimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

(3R)-3-{6-methyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-5-phenylpentanoic acid

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(ethylsulfonyl)hexanamide

(3S)-3-{3-[(4,6-dimethyl-1,2-benzisothiazol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

N-[(2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}ethyl)sulfonyl]benzamide

(2S)-2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid

3-{3-[(3,8-dimethylindolizin-1-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

(3S)-N-(aminosulfonyl)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide

3-ethoxy-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid

(3R)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

(3R)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

2-({3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)butanoic acid

3-{3-[(4,6-dimethyl-1,2-benzisothiazol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid

1-({3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)cyclopentanecarboxylic acid

2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)acetamide

3-{3-[(1-ethyl-4-methyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

3-{3-[(5-bromo-1-methyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid

3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(methylsulfonyl)hexanamide

3-{3-[(6,8-dichloro-3-methylindolizin-1-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

4-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}heptanoic acid

(2S)-2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid

1-({3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)cyclopentanecarboxylic acid

2,2-dimethyl-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

5-hydroxy-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid

3-{3-[(6-bromo-2-oxo-2,3-dihydro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid

3-{3-[(6-bromo-1-methyl-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclopent-1-ene-1-carboxylic acid

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-4-methoxybutanoic acid

3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2-methylpropanoic acid

(1-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclopentyl)acetic acid

3-{3-[(1-cyano-5-methylindolizin-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-methoxybenzoic acid

3-{3-[(5-chloro-1-methyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

3-{7-bromo-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

5-methyl-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid

3-{3-[(6-methyl-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

- 4-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)butanamide
- (2E)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} acrylic acid
- 3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2,2-dimethylbutanoic acid
- (2S)-3-methyl-2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid
- 3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-4-methoxy-2,2-dimethylbutanoic acid
- 3-[3-(1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid
- 3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-4-methoxybenzoic acid
- 3-{2-oxo-3-[(1,2,3,5-tetramethyl-1H-indol-7-yl)methyl]-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
- 3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
- 2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}benzoic acid
- 3-{3-[(8-bromo-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
- 3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid
- N-(isopropylsulfonyl)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanamide
- N-[(2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}ethyl)sulfonyl]acetamide
- 3-{3-[(4-methyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
- (2S)-2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
- 2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-methylbenzoic acid
- 3-{3-[(6-bromo-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

- 3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(methylsulfonyl)propanamide
- 3-{3-[(4-chloro-1-methyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
- 3-{3-[(6-chloro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
- 3-methoxy-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid
- 4-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid
- 3-{3-[(5-fluoro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
- 4-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid
- 3-[3-(3-hydroxy-2,5-dimethylbenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid
- {3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}acetic acid
- 2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}ethanesulfonic acid
- 3-{2-oxo-3-[(1,2,3-trimethyl-1H-indol-7-yl)methyl]-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid
- 1-[(4-methyl-1-benzothien-3-yl)methyl]-3-[2-(1H-tetrazol-5-yl)ethyl]-1,3-dihydro-2H-benzimidazol-2-one
- 3-{3-[(6-bromo-1-methyl-2-oxo-2,3-dihydro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
- 3-[2-oxo-3-(2,3,5-trichlorobenzyl)-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid
- 3-{2-oxo-3-[(1,2,3-trimethyl-1H-indol-7-yl)methyl]-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid
- (1R,2R)-2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclohexanecarboxylic acid
- 3-(6-{[2-(3,4-dimethoxyphenyl)ethyl]carbonyl})-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-ylpropanoic acid
- 3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-methylbutanoic acid

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-7-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

3-{3-[(8-bromo-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2-methylpropanoic acid

3-(6-{[2-(3-hydroxy-4-methoxyphenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2-methylpropanoic acid

3-{3-[(6-chloro-1-methyl-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

3-(6-{[2-(2-methoxyphenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid

3-{3-[(4-chloro-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

3-({3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)benzoic acid

2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclopentanecarboxylic acid

1-[2-(1,1-dioxido-4-oxo-1,2,5-thiadiazolidin-2-yl)ethyl]-3-[(4-methyl-1-benzothien-3-yl)methyl]-1,3-dihydro-2H-benzimidazol-2-one

3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]pentanoic acid

3-[6-{[2-(3,4-dimethoxyphenyl)ethyl]carbamoyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid

4-methyl-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]pentanoic acid

3-[3-(5-bromo-2-methylbenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid

3-[3-(2,5-dimethylbenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid

3-{3-[(5-methoxy-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

trans-4-({3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} methyl)cyclohexanecarboxylic acid

3-[3-(2,5-dichlorobenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid

3-{3-[(6-chloro-2-oxo-2,3-dihydro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid

3-{3-[(5-bromo-1-methyl-2-oxo-2,3-dihydro-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid

3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]-N-(phenylsulfonyl)propanamide

3-[6-{[2-(3-hydroxy-4-methoxyphenyl)ethyl]carbamoyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid

(3S)-3-{5-(carbamoylamino)-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid

(3S)-3-{5-[(tert-butoxycarbonyl)amino]-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid

3-{6-(benzylcarbamoyl)-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} propanoic acid

3-{6-[(2-hydroxyethyl)carbamoyl]-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} propanoic acid

3-[3-(5-chloro-2-methylbenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid

N-methyl-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)hexanamide

3-(6-{[2-(4-methoxyphenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid

(2E)-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]acrylic acid

3-[6-{[2-(2-methoxyphenyl)ethyl]carbamoyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid

3-{6-[(3-hydroxypyrrolidin-1-yl)carbonyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} propanoic acid

3-{3-[(4-fluoro-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid

3-(6-{[2-(3-methoxyphenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid

3-{3-[5-chloro-2-(trifluoromethyl)benzyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid

3-{3-[(4-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} propanoic acid

3-{3-(1-naphthylmethyl)-2-oxo-6-[(tetrahydrofuran-2-ylmethyl)carbamoyl]-2,3-dihydro-1H-benzimidazol-1-yl} propanoic acid

3-{6-[(2-hydroxyethyl)carbamoyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} propanoic acid

3-{[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]methyl} benzoic acid

3-[6-{[2-(1,3-dioxolan-2-yl)ethyl]carbamoyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl] propanoic acid

3-[6-{[2-(hydroxymethyl)piperidin-1-yl]carbonyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl] propanoic acid

3-{5-anilino-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid

3-{3-[(1-methyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} propanoic acid

3-[6-({3-[acetyl(methyl)amino]pyrrolidin-1-yl} carbonyl)-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl] propanoic acid

3-(6-{[2-(2-fluorophenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl) propanoic acid

3-[6-{[(2S)-2-carbamoylpyrrolidin-1-yl]carbonyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl] propanoic acid

3-{6-[(3-carbamoylpiperidin-1-yl)carbonyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} propanoic acid

3-{3-[2-methyl-5-(trifluoromethyl)benzyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid

3-{3-[(1,3-dimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} propanoic acid

3-(6-{[2-(2-chlorophenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl) propanoic acid

3-{3-[(5-bromo-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid

3-{6-[(1,1-dioxidotetrahydro-3-thienyl)carbamoyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} propanoic acid

3-(6-{[2-(4-chlorophenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid

3-[6-cyano-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid

2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}ethanesulfonamide

N-(methylsulfonyl)-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanamide

3-{3-[(1,2-dimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

N-benzyl-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-6-[(2-phenylethyl)carbamoyl]-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

3-{3-[(1-cyano-5-methylindolizin-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

3-{3-[(5-chloro-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

3-{3-[(4-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

3-(6-{[2-(3-fluorophenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid

3-{3-[(4-bromo-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

3-{3-[(6-bromo-1-methyl-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

3-{3-[(6-hydroxy-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

3-{3-(1-naphthylmethyl)-2-oxo-6-[(2-oxotetrahydrofuran-3-yl)carbamoyl]-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

3-(6-{[2-(3-chlorophenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid

3-[3-(3,5-dibromobenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid

3-[3-(3,5-dichlorobenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid

3-[6-({(2S)-2-hydroxy-1-(hydroxymethyl)-2-[4-(methylthio)phenyl]ethyl} carbamoyl)-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid

3-{3-[(5-chloro-2-oxo-2,3-dihydro-1,3-benzoxazol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

2-({3-[2-bromo-5-(trifluoromethyl)benzyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)pentanoic acid

3-{3-[2-bromo-5-(trifluoromethyl)benzyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid

3-[6-{[(2S)-2-(hydroxymethyl)pyrrolidin-1-yl]carbonyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid

3-[3-(1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid

5-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]pentanoic acid

3-{3-[(4-chloro-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid

(R)-3-(4-Chloro-phenyl)-3-[3-(1,4-dimethyl-1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid

3-[3-(1,4-Dimethyl-1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2-hydroxymethyl-propionic acid

3-[2-Oxo-3-(1,4,6-trimethyl-1H-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-2-pyridin-3-yl-propionic acid

(R)-3-[6-Cyano-2-oxo-3-(1,4,6-trimethyl-1H-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid

(R)-3-[6-Carbamoyl-2-oxo-3-(1,4,6-trimethyl-1H-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid

(R)-3-[6-Cyano-2-oxo-3-(1,4,6-trimethyl-1H-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-(4-fluoro-phenyl)-propionic acid

(R)-3-[6-Cyano-2-oxo-3-(1,4,5-trimethyl-1H-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-(4-fluoro-phenyl)-propionic acid

or a pharmaceutically acceptable salt thereof.

In all the compounds disclosed hereinabove in this application, in the event the nomenclature is in conflict with the structure, it shall be understood that the compound is defined by the structure.

The invention includes the use of any compounds of described above containing one or more asymmetric carbon atoms may occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures, and individual diastereomers. All such isomeric forms of these compounds are expressly included in the present invention. Each stereogenic carbon may be in the R or S configuration, or a combination of configurations.

Some of the compounds of formula (I) can exist in more than one tautomeric form. The invention includes methods using all such tautomers.

All terms as used herein in this specification, unless otherwise stated, shall be understood in their ordinary meaning as known in the art. For example, C₁₋₄alkoxy includes the organic radical C₁₋₄alkyl with a terminal oxygen, such as methoxy, ethoxy, propoxy, butoxy.

All organic radicals: alkyl, alkenyl and alkynyl groups, or such groups which are incorporated in other radicals such as acyl and alkoxy, shall be understood as being branched or unbranched where structurally possible and unless otherwise specified, and may be partially or fully halogenated.

The term "lower" referred to above and hereinafter in connection with organic radicals or compounds respectively defines such as branched or unbranched with up to and including 7, preferably up to and including 4 and advantageously one or two carbon atoms.

A cyclic group shall be understood to mean carbocycle, heterocycle or heteroaryl, each may be partially or fully halogenated.

An acyl group is a radical defined as $-C(O)-R$, where R is an organic radical or a cyclic group. Acyl represents, for example, carbocyclic or heterocyclic aroyl, cycloalkylcarbonyl, (oxa or thia)-cycloalkylcarbonyl, lower alkanoyl, (lower alkoxy, hydroxy or acyloxy)-lower alkanoyl, (mono- or di- carbocyclic or heterocyclic)-(lower alkanoyl or lower alkoxy-, hydroxy- or acyloxy- substituted lower alkanoyl), or biaroyl.

Carbocycles include hydrocarbon rings containing from three to fourteen carbon atoms. These carbocycles may be either aromatic or non-aromatic ring systems. The non-aromatic ring systems may be mono- or polyunsaturated, monocyclic, bicyclic or tricyclic and may be bridged. Preferred carbocycles include but are not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cycloheptanyl, cycloheptenyl, phenyl, benzyl, indanyl, indenyl, benzocyclobutanyl, dihydronaphthyl, tetrahydronaphthyl, naphthyl, decahydronaphthyl, benzocycloheptanyl, adamantyl, norbornyl, fluorene, and benzocycloheptenyl. Certain terms for cycloalkyl such as cyclobutanyl and cyclobutyl shall be used interchangeably.

The term "heterocycle" refers to a stable nonaromatic 4-8 membered (but preferably, 5 or 6 membered) monocyclic or non-aromatic 8-11 membered bicyclic heterocycle radical which may be either saturated or unsaturated. Each heterocycle consists of carbon atoms and one or more, preferably from 1 to 4 heteroatoms chosen from nitrogen, oxygen and sulfur. The heterocycle may be attached by any atom of the cycle, which results in the creation of a stable structure. Unless otherwise stated, heterocycles include but are not limited to, for example pyrrolidinyl, pyrrolinyl, morpholinyl, thiomorpholinyl, thiomorpholinyl sulfoxide, thiomorpholinyl sulfone, dioxalanyl, piperidinyl, piperazinyl, tetrahydrofuranyl, tetrahydropyranyl, tetrahydrofuranyl, 1,3-dioxolanone, 1,3-dioxanone, 1,4-dioxanyl, piperidinonyl, tetrahydropyrimidinyl, pentamethylene sulfide, pentamethylene sulfoxide, pentamethylene sulfone, tetramethylene sulfide, tetramethylene sulfoxide and tetramethylene sulfone.

The term "heteroaryl" shall be understood to mean an aromatic 5-8 membered monocyclic or 8-11 membered bicyclic ring containing 1-4 heteroatoms such as N, O and S. Unless otherwise

stated, such heteroaryls include aziridinyl, thienyl, furanyl, isoxazolyl, oxazolyl, thiazolyl, thiadiazolyl, pyrazolyl, pyrrolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, pyranyl, quinoxalinyl, indolyl, benzimidazolyl, benzoxazolyl, benzopyrrolyl, benzothiazolyl, benzothienyl, quinolinyl, quinazolinyl, naphthyridinyl, indazolyl, triazolyl, pyrazolo[3,4-*b*]pyrimidinyl, purinyl, pyrrolo[2,3-*b*]pyridinyl, pyrazolo[3,4-*b*]pyridinyl, tubercidinyl, oxazo[4,5-*b*]pyridinyl, tetrazol-5-yl, 5-oxo-1,2,4-oxadiazol-3-yl, 5-oxo-1,2,4-thiadiazol-3-yl, 5-thioxo-1,2,4-oxadiazol-3-yl, 5-thioxo-1,2,4-thiadiazol-3-yl, 3-oxo-1,2,4-oxadiazol-5-yl, N3,5-dioxo-5-oxadiazolidino, 2-oxo-1,3,4-oxadiazol-2-yl, 3-oxo-1,2,4-triazol-5-yl, 2,dioxo-benzothiadiazin-6-yl, 1,2-dioxo-3-hydroxy-3-cyclobuten-4-yl, and imidazo[4,5-*b*]pyridinyl.

The term “heteroatom” as used herein shall be understood to mean atoms other than carbon such as oxygen, nitrogen, sulfur and phosphorous.

As used herein, “nitrogen” and “sulfur” include any oxidized form of nitrogen and sulfur and the quaternized form of any basic nitrogen. All heteroatoms in open chain or cyclic radicals include all oxidized forms.

In all alkyl groups or carbon chains one or more carbon atoms can be optionally replaced by heteroatoms: O, S or N, it shall be understood that if N is not substituted then it is NH, it shall also be understood that the heteroatoms may replace either terminal carbon atoms or internal carbon atoms within a branched or unbranched carbon chain. Such groups can be substituted as herein above described by groups such as oxo to result in definitions such as but not limited to: alkoxycarbonyl, acyl, amido and thioxo.

The term “aryl” as used herein shall be understood to mean aromatic carbocycle or heteroaryl as defined herein. Each aryl or heteroaryl unless otherwise specified includes its partially or fully hydrogenated derivative and/or is partially or fully halogenated. For example, quinolinyl may include decahydroquinolinyl and tetrahydroquinolinyl, naphthyl may include its hydrogenated derivatives such as tetrahydronaphthyl. Other partially or fully hydrogenated derivatives of the aryl and heteroaryl compounds described herein will be apparent to one of ordinary skill in the art.

The term "halogen" as used in the present specification shall be understood to mean bromine, chlorine, fluorine or iodine, preferably fluorine. The definitions "partially or fully halogenated"; partially or fully fluorinated; "substituted by one or more halogen atoms", includes for example, mono, di or tri halo derivatives on one or more carbon atoms. For alkyl, a nonlimiting example would be $-\text{CH}_2\text{CHF}_2$, $-\text{CF}_3$ etc.

The compounds of the invention are only those which are contemplated to be 'chemically stable' as will be appreciated by those skilled in the art. For example, a compound which would have a 'dangling valency', or a 'carbanion' are not compounds contemplated by the inventive methods disclosed herein.

The invention includes pharmaceutically acceptable derivatives of compounds of formula (I). A "pharmaceutically acceptable derivative" refers to any pharmaceutically acceptable salt or ester, or any other compound which, upon administration to a patient, is capable of providing (directly or indirectly) a compound useful for the invention, or a pharmacologically active metabolite or pharmacologically active residue thereof. A pharmacologically active metabolite shall be understood to mean any compound of the invention capable of being metabolized enzymatically or chemically. This includes, for example, hydroxylated or oxidized derivative compounds of the formula (I).

Pharmaceutically acceptable salts include those derived from pharmaceutically acceptable inorganic and organic acids and bases. Examples of suitable acids include hydrochloric, hydrobromic, sulfuric, nitric, perchloric, fumaric, maleic, phosphoric, glycolic, lactic, salicylic, succinic, toluene-*p*-sulfuric, tartaric, acetic, citric, methanesulfonic, formic, benzoic, malonic, naphthalene-2-sulfuric and benzenesulfonic acids. Other acids, such as oxalic acid, while not themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining the compounds and their pharmaceutically acceptable acid addition salts. Salts derived from appropriate bases include alkali metal (*e.g.*, sodium), alkaline earth metal (*e.g.*, magnesium), ammonium and $\text{N}-(\text{C}_1\text{-C}_4 \text{ alkyl})_4^+$ salts.

In addition, within the scope of the invention is use of prodrugs of compounds of the formula (I). Prodrugs include those compounds that, upon simple chemical transformation, are modified to produce compounds of the invention. Simple chemical transformations include hydrolysis, oxidation and reduction. Specifically, when a prodrug is administered to a patient, the prodrug may be transformed into a compound disclosed hereinabove, thereby imparting the desired pharmacological effect.

The compounds described herein are either commercially available or can be made by methods and any necessary intermediates well known in the art .

In order that this invention be more fully understood, the following examples are set forth. These examples are for the purpose of illustrating preferred embodiments of this invention, and are not to be construed as limiting the scope of the invention in any way.

The examples which follow are illustrative and, as recognized by one skilled in the art, particular reagents or conditions could be modified as needed for individual compounds without undue experimentation. Starting materials used in the scheme below are either commercially available or easily prepared from commercially available materials by those skilled in the art.


GENERAL SYNTHETIC METHODS

The invention also provides processes for making compounds of Formula (I). In all schemes, unless specified otherwise, R_1 , R_2 , R_3 and n in the formulas below shall have the meaning of R_1 , R_2 , R_3 and n in Formula (I) of the invention described herein above.

Optimum reaction conditions and reaction times may vary depending on the particular reactants used. Unless otherwise specified, solvents, temperatures, pressures, and other reaction conditions may be readily selected by one of ordinary skill in the art. Specific procedures are provided in the Synthetic Examples section. Typically, reaction progress may be monitored by thin layer chromatography (TLC), if desired, and intermediates and products may be purified by chromatography on silica gel and/or by recrystallization.

The examples which follow are illustrative and, as recognized by one skilled in the art, particular reagents or conditions could be modified as needed for individual compounds without undue experimentation. Starting materials and intermediates used, in the schemes below, are either commercially available or easily prepared from commercially available materials by those skilled in the art.


The compounds in this application may be synthesized according to schemes 1, 2 and 3 below:



Scheme 1


As illustrated in scheme 1, reacting an appropriately substituted benzimidazolone of formula II (wherein P is a protecting group) with an alkylating agent $\text{R}_1\text{-LG}$ (wherein LG is a leaving group such as halogen, alkylammonium salt, tosylate, an activated alcohol) provides a N-alkylated benzimidazolone of formula III. Deprotection of the compound of formula III with a suitable reagent, in a suitable solvent, provides a compound of formula IV. N-alkylation of the compound of formula IV with a suitable alkylating agent such as $\text{R}_2\text{-LG}$ (wherein LG is a leaving group such as halogen, alkylammonium salt, tosylate, an activated alcohol), in a suitable solvent provides a compound of formula (I). Alternatively, intermediate IV may be reacted with an alkene or alkyne, such as $\text{CH}_2\text{CH}_2\text{C(O)-R}_2$ or CHC-C(O)-R_2 a suitable solvent to provide a compound of formula (I)

Further modification of the initial product of formula (I) by methods known in the art and illustrated in the Examples below, may be used to prepare additional compounds of this invention

**Scheme 2**

As outlined in scheme 2, reacting an appropriately substituted starting material of formula V (wherein Hal is F, Cl, Br or I) with an amine R_1NH_2 in a suitable solvent provides a compound of formula VI. Reducing the compound of formula VI with a suitable reducing agent, in a suitable solvent, provides a diamine of formula VII. Cyclization of the diamine of formula VII with a reagent such as N,N'-carbonyldimidazole (CDI) provides a cyclized compound of formula IV. N-alkylation of the compound of formula IV with a suitable alkylating agent such as R_2-LG (wherein LG is a leaving group such as halogen, alkylammonium salt, tosylate, an activated alcohol), in a suitable solvent provides a compound of formula (I). Alternatively, intermediate IV may be reacted with an alkene or alkyne, such as $CH_2CH_2C(O)-R_2$ or $CHC-C(O)-R_2$, in a suitable solvent to provide a compound of formula (I).

Compounds of formula (I) may also be made using the method outlined in scheme 3.


**Scheme 3**

As shown in scheme 3, alkylation of a compound of formula (IV) with an alcohol R_2OH under Mitsunobu reaction conditions gives a compound of formula (I).

Further modification of the initial product of formula (I) by methods known in the art and illustrated in the Examples below, may be used to prepare additional compounds of this invention

The experimental procedures described below are included as examples.

Example 1: 3-{3-[(5-chloro-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid



1-(5-Chloro-benzo[*b*]thiophen-3-ylmethyl)-3-isopropenyl-1,3-dihydro-benzimidazol-2-one:

To a solution of 1-isopropenylbenzimidazolone (0.3 g, 1.72 mmol) in DMF (5 mL) was added 60% NaH (0.1 g, 2.6 mmol). The reaction mixture was stirred at room temperature for 15 min and added 3-(Bromomethyl)-5-chlorobenzo[*b*]thiophene (0.54 g, 2.07 mmol). The reaction mixture was heated to 65 °C and stirred at that temperature for 2 h. When the reaction was complete, the mixture was treated with water. The mixture was then diluted with ethyl acetate and washed with water (x4). The organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC to afford 0.53 g (87%) of the desired product.

1-(5-Chloro-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one:

1-(5-Chloro-benzo[*b*]thiophen-3-ylmethyl)-3-isopropenyl-1,3-dihydro-benzimidazol-2-one (0.53 g, 1.56 mmol) was treated with MeOH (5 mL) followed by water (5 mL). To this heterogeneous reaction mixture was added 37% hydrochloric acid (4 mL). This heterogeneous reaction mixture was heated to 80 °C for 30 min. A white solid was precipitated out. TLC indicated that reaction was complete (starting material R_f 0.8, product R_f 0.4, CH₂Cl₂:MeOH 95:5). The reaction mixture was diluted with CH₂Cl₂ (at this point a white precipitate was completely soluble in CH₂Cl₂) and washed with water. The organic phase was dried over Na₂SO₄ and concentrated to afford 0.44 g (89%) of the desired product as a white solid. This product was used for the next reaction without further purification. LCMS (ESMS): m/z 315.79 (M+H⁺).

3-[3-(5-Chloro-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid ethyl ester:


To a solution of 1-(5-Chloro-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one (50 mg, 0.16 mmol) in DMF (2 mL) were added ethyl acrylate (0.02 mL) and benzyltrimethyl ammonium hydroxide (20 mg, 0.01 mmol). The resulting mixture was stirred at room temperature overnight. When the reaction was complete, the reaction mixture was diluted with ethyl acetate and washed with water (x4). The organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC using 95:5 CH₂Cl₂:MeOH as an eluent to afford 50 mg of the desired product as a colorless oil. LCMS (ESMS): m/z 415.78 (M+H⁺).

3-{3-[(5-chloro-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid:


A solution of 3-[3-(5-Chloro-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid ethyl ester (50 mg, 0.12 mmol) in MeOH (3 mL) was added 2N NaOH (1 mL). The reaction mixture was stirred at room temperature for 1 h. The solution was then treated with 1M HCl solution until pH ~4 and extracted with CH₂Cl₂. The organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC using 95:5 CH₂Cl₂:MeOH as an eluent to afford 20 mg (42 %) of the desired product as a white solid. LCMS (ESMS): m/z 387.89 (M+H⁺).

Following compounds were synthesized using a similar procedure.


3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]pentanoic acid. LCMS (ESMS): m/z 375.07 ($M+H^+$).




4-methyl-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]pentanoic acid. LCMS (ESMS): m/z 389.66 ($M+H^+$).




5-methyl-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid. LCMS (ESMS): m/z 403.64 ($M+H^+$).




3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid. LCMS (ESMS): m/z 389.05 ($M+H^+$).




3-ethoxy-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid.
LCMS (ESMS): m/z 391.35 ($M+H^+$).



3-[3-(3,5-dibromobenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid. LCMS
(ESMS): m/z 497.06 ($M+H^+$).



Example 2: 3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid



3-(3-Isopropenyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid ethyl ester:

To a solution of 1-Isopropenyl-2-benzimidazolidinone (2.5 g, 14.4 mmol) in DMF (10 mL) were added ethyl acrylate (1.7 mL, 15.8 mmol) and benzyltrimethyl ammonium hydroxide in MeOH (1.4 g, 8.6 mmol). The resulting mixture was stirred at room temperature for 1.5 h. When the reaction was complete, the reaction mixture was diluted with ethyl acetate and washed with water (x4). The organic phase was dried over Na_2SO_4 and concentrated. The resulting residue was purified by silica gel prep TLC using 1:1 Hexanes: Ethyl acetate as an eluent to afford 3.0 g (76%) of the desired product along with methyl ester. Since the methyl ester was the major product, this product was used for the next reaction.

3-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid methyl ester:

3-(3-Isopropenyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid methyl ester (2.2 g, 8.3 mmol) was treated with MeOH (5 mL) followed by water (5 mL). To this heterogeneous reaction mixture was added 37% hydrochloric acid (4 mL). This reaction mixture was heated to 60 °C for 20 min. The reaction remained as a clear homogeneous solution after heating. The reaction mixture was diluted with CH_2Cl_2 and washed with water. The organic phase was dried over Na_2SO_4 and concentrated to afford 1.7 g of the desired product as a white solid. This product was used for the next reaction without further purification.

3-(4-Methyl-benzo[b]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl-propionic acid methyl ester:


To a solution of 3-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid methyl ester (0.1 g, 0.45 mmol) in DMA (2 mL) were added K_2CO_3 (0.1 g, 0.68 mmol) and trimethyl-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-ammonium; iodide (0.17 g, 0.50 mmol). The reaction mixture was heated to 150 °C for 15 min in the microwave. When the reaction was complete, the mixture was diluted with ethyl acetate and washed with water (x4). The organic phase was dried over Na_2SO_4 and concentrated. The resulting oily residue was purified by silica gel prep TLC using 98:2 CH_2Cl_2 :MeOH as an eluent to afford 0.1 g of the desired product. LCMS (ESMS): m/z 381.21 ($M+H^+$).

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid:


To a solution of 3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid methyl ester (0.1 g, 0.26 mmol) in MeOH (2 mL) was added 2 N NaOH (2 mL). The reaction mixture was stirred at room temperature for 2 h. The solution was then treated with 1M HCl until pH ~2 and extracted with CH_2Cl_2 (x2). The organic phase was dried over Na_2SO_4 and concentrated. The resulting residue was purified by silica gel prep TLC using 95:5 CH_2Cl_2 :MeOH as an eluent to afford 48 mg of the title compound as an off-white solid. LCMS (ESMS): m/z 367.20 ($M+H^+$).

The following compounds were synthesized by a similar procedure.


3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS): m/z 367.20 ($M+H^+$).




3-[3-(1-benzothien-3-ylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid. LCMS (ESMS): m/z 353.08 ($M+H^+$).




2-({3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)pentanoic acid. LCMS (ESMS): m/z 406.27 ($M+H^+$)




2-({3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)butanoic acid. LCMS (ESMS): m/z 392.02 ($M+H^+$)




3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2-methylpropanoic acid. LCMS (ESMS): m/z 378.23 ($M+H^+$)



2-({3-[(4,6-dimethyl-1,2-benzisothiazol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)pentanoic acid. LCMS (ESMS): m/z 424.18 ($M+H^+$)



Example 3: 3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid



1-Isopropenyl-3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one:
1-isopropenylbenzimidazolone (0.5 g, 2.9 mmol) in DMA (8 mL) was added and trimethyl-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-ammonium; iodide (1.0 g, 2.9 mmol) and K₂CO₃ (0.80 g, 5.74 mmol). The reaction mixture was heated to 120 °C for 15 min in a microwave and then 150 °C for another 5 min. The mixture was diluted with ethyl acetate and washed with water (x4). The organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC using 98:2 CH₂Cl₂:MeOH as an eluent to afford 0.7 g of the desired product as a white solid

1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one:

1-Isopropenyl-3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one (0.70 g, 2.1 mmol) was treated with MeOH (10 mL) followed by water (10 mL). To this heterogeneous reaction mixture was added 37% hydrochloric acid (5 mL). This reaction mixture was heated to 65 °C for 3 h. The reaction mixture maintained as a heterogeneous mixture with white precipitates. The reaction mixture was extracted with CH₂Cl₂ (x2). The combined organic phase was washed with water. The organic phase was dried over Na₂SO₄ and concentrated to afford 0.7 g of the desired product as a white scaly solid.

3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester:

To a solution of 1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one (0.2 g, 0.68 mmol) in DMA (5 mL) were added ethyl trans-2-hexanoate (0.19 g, 1.4 mmol) and benzyltrimethyl ammonium hydroxide in MeOH (0.43 g, 1.0 mmol) (Benzyltrimethyl ammonium hydroxide in MeOH can be replaced with K₂CO₃ (2eq) as a base). The reaction mixture was heated to 60 °C for 20 h. The reaction mixture was diluted with ethyl acetate and washed with water (x4). The organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was attempted to dissolve in a mixture of CH₂Cl₂ and MeOH to load to a prep TLC. However, some insoluble material was present. This insoluble solid was filtered, however, the structure of the filtered solid couldn't be characterized (no aliphatic region in NMR). The filtrate was concentrated and resulting residue was purified by silica gel prep TLC using 98:2 CH₂Cl₂:MeOH as an eluent to afford 0.2 g of the desired product (with methyl ester instead of ethyl ester) as an oil. LCMS (ESMS): *m/z* 423.46 (M+H⁺).

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid:


To a solution of the ester (0.1 g, 0.23mmol) in MeOH (5 mL) was added 2N NaOH (3 mL). The reaction mixture was stirred at room temperature for 2 h. When all the starting material was consumed, the reaction mixture was acidified with 1N HCl. The reaction mixture was extracted with CH₂Cl₂ (x2). The combined organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC using 95:5

CH_2Cl_2 :MeOH as an eluent to afford 66 mg of the title compound as an off-white powder.


LCMS (ESMS): m/z 409.18 ($\text{M}+\text{H}^+$).

Two enantiomers of 3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester were separated by chiral column chromatography (85:15 Hexanes:iPA, 8.5 mL/min, AD-H column) to afford (R) and (S) enantiomers. These (R) and (S) esters were hydrolyzed using 2N NaOH in MeOH to afford (R) and (S) acids.

(3S)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid. LCMS (ESMS): m/z 409.59($\text{M}+\text{H}^+$).




(3R)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid. LCMS (ESMS): m/z 409.24 ($\text{M}+\text{H}^+$).




The following compounds were synthesized using a similar procedure.


3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid. LCMS (ESMS): m/z 381.15 ($\text{M}+\text{H}^+$).




3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid. LCMS (ESMS): m/z 395.10 (M+H⁺).




3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}octanoic acid. LCMS (ESMS): m/z 437.33 (M+H⁺).




3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}heptanoic acid. LCMS (ESMS): m/z 423.43 (M+H⁺).




3-ethoxy-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS): m/z 411.40 (M+H⁺).




5-methyl-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid. LCMS (ESMS): m/z 423.70 (M+H⁺).




3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid. LCMS (ESMS): m/z 406.48 ($M+H^+$).




3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-ethoxypropanoic acid. LCMS (ESMS): m/z 408.49 ($M+H^+$).




3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}octanoic acid. LCMS (ESMS): m/z 437.33 ($M+H^+$).



3-{2-oxo-3-[(1,4,6-trimethyl-1H-indol-3-yl)methyl]-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid. LCMS (ESMS): m/z 420.28 ($M+H^+$).



Example 4: 3-methoxy-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid




(E)-3-(3-Naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-acrylic acid ethyl ester:
To a solution of 1-Naphthalen-1-ylmethyl-1,3-dihydro-benzimidazol-2-one (0.5 g, 1.8 mmol) in DMF (4 mL) were added ethyl propiolate (0.28 mL, 2.7 mmol) and DMAP (0.22 mL, 1.8 mmol). The reaction mixture was stirred at room temperature for 10 min. The reaction mixture turned dark brown in color. The reaction mixture was diluted with ethyl acetate and washed with water (x4). The organic phase was dried over Na_2SO_4 and concentrated to afford 0.68 g of the desired product as a dark brown foam. This crude product was used for the next reaction without further purification. LCMS (ESMS): m/z 373.33 ($\text{M}+\text{H}^+$).

3-methoxy-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid:
To a solution of (E)-3-(3-Naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-acrylic acid ethyl ester (0.1 g, 0.27 mmol) in DMF (1 mL) was added NaOCH_3 (1.1 mL, 0.5 M in MeOH). The reaction mixture was stirred at 60 °C for 2 h (due to poor solubility, the reaction


was heated). The reaction mixture was concentrated. The reaction mixture was diluted with 1N HCl and then extracted with CH₂Cl₂. The organic phase was dried over Na₂SO₄ and concentrated. Since LCMS indicated that there was a small amount of the product in methyl ester form, the resulting residue was diluted with methanol and treated with 2N NaOH for 30 min. The mixture was acidified by adding 1N HCl and extracted with CH₂Cl₂. The organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC using 95:5 CH₂Cl₂:MeOH as an eluent to afford 39 mg of the title compound as an off-white solid. LCMS (ESMS): m/z 377.31 (M+H⁺).

Example 5: (2E)-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]acrylic acid



To a solution of (E)-3-[3-(1-Naphthalen-1-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-acrylic acid ethyl ester (0.1 g, 0.30 mmol) in MeOH (10 mL) was added 2N NaOH (5 mL). The reaction mixture was stirred at room temperature for 3 h. Since the mixture was heterogeneous, the mixture was heated to 60 °C for 2 h. The reaction mixture was then treated with 1N HCl and extracted with CH₂Cl₂. The organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC using 10:1 CH₂Cl₂:MeOH as an eluent to afford 30 mg of the title compound as an off-white solid. LCMS (ESMS): m/z 345.12 (M+H⁺).

Example 6: 3-{[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]methyl}benzoic acid



3-(3-(1-Naphthalen-1-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-ylmethyl)-benzoic acid methyl ester:


To a solution of 1-(1-Naphthalen-1-ylmethyl)-1,3-dihydro-benzimidazol-2-one (0.1 g, 0.37 mmol) in DMF (3 mL) was added 60% NaH (22 mg, 0.55 mmol). The reaction mixture was stirred at room temperature for 10 min and then Methyl 3-(bromomethyl)benzoate (0.1 g, 0.44 mmol) was added to the reaction mixture. The reaction mixture was stirred at room temperature for 2 h. When the reaction was complete, the mixture was diluted with ethyl acetate and washed with water (x4). The organic phase was then dried over Na_2SO_4 and concentrated. The resulting residue was purified by silica gel prep TLC using 100% CH_2Cl_2 as an eluent to afford 0.14 g of the desired product as a white foam. LCMS (ESMS): m/z 423.84 ($\text{M}+\text{H}^+$).

3-[[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]methyl]benzoic acid:


To a solution of the ester (0.14 g, 0.34 mmol) in MeOH (3 mL), 2N NaOH (2 mL) was added. The reaction mixture was stirred at room temperature for 1 h until all the starting material was consumed. The mixture was then acidified by adding 1M HCl and extracted with CH_2Cl_2 (x2). The organic phase was dried over Na_2SO_4 and concentrated. The resulting white solid was not soluble in CH_2Cl_2 . This white solid was therefore filtered to afford 0.1 g of the title compound. LCMS (ESMS): m/z 407.70 ($\text{M}-\text{H}^+$).

The following compounds were synthesized using a similar procedure.


5-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]pentanoic acid. LCMS (ESMS): m/z 375.84 ($\text{M}+\text{H}^+$).




4-{3-[(5-chloro-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid. LCMS (ESMS): m/z 401.75 ($M+H^+$).




3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methylbenzoic acid. LCMS (ESMS): m/z 429.09 ($M+H^+$).




{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}acetic acid. LCMS (ESMS): m/z 353.28 ($M+H^+$).



4-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid. LCMS (ESMS): m/z 381.81 ($M+H^+$).



Example 7: (3S)-3-{6-methyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-5-phenylpentanoic acid



(S)-3-(5-Methyl-2-nitro-phenylamino)-5-phenyl-pentanoic acid ethyl ester:

To a solution of 3-Fluoro-4-nitrotoluene (0.7 g, 4.5 mmol) in DMF (5 mL) were added (S)-3-Amino-5-phenyl-pentanoic acid ethyl ester hydrochloride (1.2 g, 4.5 mmol) and DIPEA (1.6 g, 9.0 mmol). The reaction mixture was stirred at room temperature for 48 h. The reaction mixture was then diluted with ethyl acetate and washed with water (x4). The organic phase was then dried over Na_2SO_4 and concentrated to afford an orange solid. The resulting residue was purified by silica gel prep TLC using 4:1 Hexanes:ethyl acetate as an eluent to afford 0.9 g of the desired product.

(S)-3-(2-Amino-5-methyl-phenylamino)-5-phenyl-pentanoic acid ethyl ester:

To a solution of (S)-3-(5-Methyl-2-nitro-phenylamino)-5-phenyl-pentanoic acid ethyl ester (0.9 g, 2.5 mmol) in MeOH (20 mL) was added slurry of Pd/C (0.2 g) in MeOH. The reaction mixture was de-gassed using house vacuum. The reaction mixture was stirred under H₂ balloon for 3 h. When the reaction color changed from orange to colorless, the reaction mixture was filtered through a pad of celite. The filtrate was concentrated to afford 0.84 g the desired product. This product was used for the next reaction without further purification.

(S)-3-(6-Methyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-5-phenyl-pentanoic acid ethyl ester:

To a solution of (S)-3-(2-Amino-5-methyl-phenylamino)-5-phenyl-pentanoic acid ethyl ester (0.84 g, 2.6 mmol) in THF (15 mL) was added CDI (0.50 g, 3.1 mmol). The reaction mixture was stirred at room temperature for 16 h. When the reaction was complete, the reaction mixture was diluted with CH₂Cl₂ and washed with water. The organic phase was dried over Na₂SO₄ and concentrated to afford 0.75 g of the desired product as a dark brown solid. The resulting residue was used for the next reaction without further purification.

(S)-3-[6-Methyl-3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-5-phenyl-pentanoic acid ethyl ester:

To a solution of (S)-3-(6-Methyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-5-phenyl-pentanoic acid ethyl ester (0.2 g, 0.57 mmol), (4-Methyl-benzo[*b*]thiophen-3-yl)-methanol (0.10 g, 0.57 mmol), and triphenylphosphine (0.18 g, 0.68 mmol) was added dropwise DIAD (0.14 g, 0.68 mmol). The mixture was stirred at room temperature for 1 h. The reaction mixture was diluted with ethyl acetate and washed with water. The organic phase was dried over Na₂SO₄ and concentrated. The resulting oily residue was purified by prep TLC using 4:1 Hexane:EtOAc as an eluent to afford 0.12 g of the desired product as a brown oil.


(3S)-3-{6-methyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-5-phenylpentanoic acid:

To a solution of (S)-3-[6-Methyl-3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester (0.12 g, 0.23 mmol) in MeOH (5 mL) was added 2N NaOH (2 mL). The reaction mixture was stirred at room temperature for 2 h. When the reaction was complete, the reaction mixture was treated with 1N HCl and extracted with CH₂Cl₂


(x2). The organic phase was dried over Na_2SO_4 and concentrated. The resulting residue was purified by silica gel prep TLC using 10:1 CH_2Cl_2 :MeOH as an eluent to afford 0.1 g of the title compound as a white solid. LCMS (ESMS): m/z 485.54 ($\text{M}+\text{H}^+$).

The following compounds were synthesized following a similar procedure using appropriate starting materials.


(3R)-3-{6-methyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-5-phenylpentanoic acid. LCMS (ESMS): m/z 485.28 ($\text{M}+\text{H}^+$).




3-{6-methyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid. LCMS (ESMS): m/z 423.21 ($\text{M}+\text{H}^+$).




3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid. LCMS (ESMS): m/z 468.34 ($\text{M}+\text{H}^+$).




(3R)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-phenylpropanoic acid. LCMS (ESMS): m/z 443.27 ($M+H^+$).




(3S)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-4-phenylbutanoic acid. LCMS (ESMS): m/z 457.28 ($M+H^+$).




(3S)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-phenylpropanoic acid. LCMS (ESMS): m/z 443.23 ($M+H^+$).




1-({3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)cyclohexanecarboxylic acid. LCMS (ESMS): m/z 480.24 ($M+H^+$).



1-({3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)cyclopentanecarboxylic acid. LCMS (ESMS): m/z 466.16 ($M+H^+$).



Example 8: 3-[6-cyano-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid



3-(5-Cyano-2-nitro-phenylamino)-propionic acid ethyl ester:

To a solution of 3-Fluoro-4-nitrobenzonitrile (1.0 g, 6.0 mmol) in DMA (5 mL) were added Beta-Alanine ethyl ester HCl (1.4 g, 9.03 mmol) and DIPEA (1.6 mL, 9.03 mmol). The reaction mixture was stirred at room temperature for 16 h (the reaction was completed; the reaction mixture turned to orange from yellow in color). The reaction mixture was then diluted with ethyl acetate and washed with water (x4). The organic phase was then dried over Na₂SO₄ and concentrated to afford 1.53 g (96%) of the desired product as yellow solid.

3-(2-Amino-5-cyano-phenylamino)-propionic acid ethyl ester:

To a solution of 3-(5-Cyano-2-nitro-phenylamino)-propionic acid ethyl ester (1.53 g, 5.8 mmol) in MeOH (10 mL) was added a slurry of Pd/C (0.3 g) in MeOH. The reaction mixture was saturated with H₂ and stirred for 2 h. When the reaction was complete, the reaction mixture

was filtered and the filtrate was concentrated to afford 0.65 g (48%) of the desired product as a brown solid. The resulting residue was used for the next reaction without further purification. LCMS (ESMS): m/z 233.66 ($M+H^+$).

3-(6-Cyano-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid ethyl ester:

To a solution of 3-(2-Amino-5-cyano-phenylamino)-propionic acid ethyl ester (0.65g, 2.8 mmol) in $CHCl_3$ (10 mL) was added CDI (0.68g, 4.2 mmol). The reaction mixture was stirred at room temperature for 16 h. When the reaction was complete, the reaction mixture was diluted with CH_2Cl_2 and washed with water. The organic phase was dried over Na_2SO_4 and concentrated to afford 0.6 g (83%) of the desired product as a brown solid. The resulting residue was used for the next reaction without further purification.

3-(6-Cyano-3-naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid ethyl ester.

To a solution of 3-(6-Cyano-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid ethyl ester (0.1 g, 0.39 mmol) in DMA (3 mL) were added 1-(chloromethyl)naphthalene (0.1 g, 0.58 mmol) and K_2CO_3 (0.11g, 0.81 mmol). The reaction mixture was stirred at 65 °C for 16 h. When the reaction was complete, the reaction mixture was diluted with ethyl acetate and washed with water (x4). The organic phase was dried over Na_2SO_4 and concentrated. The resulting residue was purified by silica gel prep TLC using 98:2 CH_2Cl_2 :MeOH as an eluent to afford 20 mg (13%) of the desired product.


3-[6-cyano-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid.

To a solution of 3-(6-Cyano-3-naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid ethyl ester (20 mg, 0.05 mmol) in MeOH (5 mL) was added 2N NaOH (3 mL). The reaction mixture was stirred at room temperature for 16 h. When all the starting material was consumed, the reaction mixture was acidified with 1N HCl. The reaction mixture was extracted with CH_2Cl_2 (x2). The combined organic phase was dried over Na_2SO_4 and concentrated. The resulting residue was purified by silica gel prep TLC using 10:1 CH_2Cl_2 :MeOH as an eluent to afford 4 mg (22%) of the desired product as an off-white solid. LCMS (ESMS): m/z 372.11 ($M+H^+$).


The following compounds were synthesized by a similar procedure.

3-[5-cyano-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid.

LCMS (ESMS): m/z 372.10 ($M+H^+$).



Example 9: 3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-5,6-dimethyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid



1-(1,4-Dimethyl-1H-indol-3-ylmethyl)-3-isopropenyl-5,6-dimethyl-1,3-dihydro-benzimidazol-2-one:

To a solution of 1-Isopropenyl-5,6-dimethyl-1,3-dihydro-benzimidazol-2-one (0.25 g, 1.2 mmol) in DMF (10 mL) was added K_2CO_3 (0.51 g, 3.7 mmol). The reaction mixture was heated to 90 °C and stirred for 20 min. (1,4-Dimethyl-1H-indol-3-ylmethyl)-trimethyl-ammonium;

iodide (0.64 g, 1.8 mmol) was then added to the reaction mixture and stirred for another 3 h. The reaction mixture was treated with ethyl acetate and washed with water. The resulting residue was purified by silica gel prep TLC using 4:1 Hexane:ethyl acetate as an eluent to afford 0.35 g the desired product as an off-white solid. LCMS (ESMS): m/z 360.33 ($M+H^+$).

1-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-5,6-dimethyl-1,3-dihydro-benzimidazol-2-one:

To 1-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-3-isopropenyl-5,6-dimethyl-1,3-dihydro-benzimidazol-2-one (0.35 mmol) in MeOH (5 mL) was added water (5 mL) and conc. HCl (3 mL). The reaction mixture was heated to 60 °C for 2 h. The reaction mixture turned into pink in color and heterogeneous (milky solution). The TLC indicated that the reaction was complete and clean. The reaction mixture was therefore diluted with CH₂Cl₂ and washed with water. The organic phase remained heterogeneous (milky with fine powder). The organic phase was filtered through a fine filter to collect some of the solid which was partially soluble in DMSO-d₆. ¹H-NMR indicated this filtered solid being the desired product. Therefore, the organic phase was concentrated without drying with Na₂SO₄ to afford a pink solid. ¹H-NMR of this product in DMSO-d₆ matched well with the previous filtered solid.

3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-5,6-dimethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester:


A mixture of 1-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-5,6-dimethyl-1,3-dihydro-benzimidazol-2-one (0.1 g, 0.31 mmol), K₂CO₃ (87 mg, 0.63 mmol) and ethyl trans -2-hexenoate (0.45 g, 3.1 mmol) was heated to 130 °C for 20 min in a microwave. Since the starting material was not soluble in ethyl trans -2-hexenoate even after heating, DMA (2 mL) was added to the reaction mixture and heated again at 150 °C for 30 min. Although the un-reacted starting material was present, the reaction mixture was diluted with ethyl acetate and washed with water (x4). The organic phase was dried over Na₂SO₄ and concentrated. The resulting heterogeneous residue was purified by silica gel prep TLC using 4:1 hexanes:ethyl acetate as an eluent to afford 25 mg of the desired product as an oil. LCMS (ESMS): m/z 462.88 ($M+H^+$).

3-{3-[(1,4-dimethyl-1*H*-indol-3-yl)methyl]-5,6-dimethyl-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}hexanoic acid:


To a solution of the ester (25 mg, 0.05 mmol) in MeOH (5 mL) was added 2N NaOH (5 mL). The reaction mixture was stirred at room temperature for 3 h. When the reaction was complete, the mixture was treated with 1N HCl and extracted with CH₂Cl₂. The organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC using 10:1 CH₂Cl₂:MeOH as an eluent to afford 22 mg of the title compound as a pale gray solid. LCMS (ESMS): m/z 434.72 (M+H⁺).

The following compounds were synthesized using a similar procedure.

3-ethoxy-3-{6-methyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS): m/z 434.72 (M+H⁺).



Example 10: 3-{5,6-dimethyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-ethoxypropanoic acid



5,6-Dimethyl-1-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one:

To a slurry of 5,6-Dimethyl-2-benzimidazolinone (0.2 g, 1.2 mmol) in DMA (2 mL) (poor solubility) were added K_2CO_3 (0.10 g, 0.74 mmol) and trimethyl-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-ammonium; iodide (0.21 g, 0.62 mmol). The heterogeneous reaction mixture was heated to 130 °C for 10 min. Since there was still un-reacted starting material left, NaH (15 mg, 0.38 mmol) was added to the reaction mixture and heated to 150 °C for 10 min. The reaction was complete. The reaction mixture was then diluted with ethyl acetate (some MeOH was added to solublize the reaction mixture into the organic phase) and washed with water (x5). The organic phase was dried over Na_2SO_4 and concentrated. The resulting solid had poor solubility. To the solid, CH_2Cl_2 was added and the resulting solid was filtered. The filtered solid was identified as the desired product (with a minimal amount of excess 5,6-Dimethyl-2-benzimidazolinone). Therefore, this filtered solid was used for the next reaction. LCMS (ESMS): m/z 323.13 ($M+H^+$).

3-[5,6-Dimethyl-1-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-ethoxy-propionic acid ethyl ester:


To a mixture of 5,6-Dimethyl-1-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one (0.1 g, 0.31 mmol) and K_2CO_3 (86 mg, 0.62 mmol) in a microwave vial was added Ethyl 3-ethoxyacrylate (0.45 g, 3.1 mmol). The reaction mixture was heated to

130 °C for 20 min in a microwave. The reaction mixture was then diluted with ethyl acetate (since there was some insoluble material left, MeOH was added to the solution mixture to solublize the material). This organic phase was washed with water (x2). The organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC using 4:1 Hexane:ethyl acetate as an eluent to afford 45 mg of the desired product as an oily foam. LCMS (ESMS): m/z 467.32 (M+H⁺).

3-{5,6-dimethyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-ethoxypropanoic acid:

To a solution of the ester (45 mg, 0.1 mmol) in MeOH (5 mL) was added 2N NaOH (2 mL). The reaction mixture was stirred at room temperature for 2 h. The reaction mixture was then acidified with 1N HCl. The reaction mixture was extracted with CH₂Cl₂ and the combined organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC using 10:1 CH₂Cl₂:MeOH as an eluent to afford 37 mg of the title compound as an off-white solid. LCMS (ESMS): m/z 439.48 (M+H⁺).

Example 11: 3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid



3-(5-Methyl-2-nitro-phenylamino)-hexanoic acid ethyl ester:

To a solution of 3-Fluoro-4-nitrotoluene (1.0 g, 6.5 mmol) in DMF (5 mL) were: 3-Amino-hexanoic acid ethyl ester hydrochloride (1.3g, 6.7 mmol) and DIPEA (2.2 mL, 12.9 mmol). The reaction mixture was stirred at room temperature for 48 h. The reaction mixture was then diluted with ethyl acetate and washed with water (x4). The organic phase was then dried over Na₂SO₄ and concentrated. The resulting residue was purified by Combiflash using hexane and ethyl acetate (10% gradient) as an eluent to afford 1.4 g of the desired product as orange oil.

3-(2-Amino-5-methyl-phenylamino)-hexanoic acid ethyl ester:

To a solution of 3-(5-Methyl-2-nitro-phenylamino)-hexanoic acid ethyl ester (1.4 g, 4.8 mmol) in MeOH (10 mL) was added a slurry of Pd/C (wet) (0.25 g) in MeOH. The reaction mixture was saturated with H₂ balloon and stirred at room temperature for 2 h. When the reaction was complete, the reaction mixture was filtered through a pad of celite. The filtrate was concentrated to afford 1.4 g of the desired product as dark purplish oil. This product was used for the next reaction without further purification.

3-(6-Methyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-hexanoic acid ethyl ester:

To a solution of 3-(2-Amino-5-methyl-phenylamino)-hexanoic acid ethyl ester (1.4 g, 5.3 mmol) in THF (20 mL) was added CDI (1.0 g, 6.4 mmol). The reaction mixture was stirred at room temperature for 16 h. When the reaction was complete, the reaction mixture was diluted with ethyl acetate and washed with water. The organic phase was dried over Na₂SO₄ and concentrated to afford 1.3 g of the desired product as dark brown solid. The resulting residue was used for the next reaction without further purification. LCMS (ESMS): m/z 291.40 (M+H⁺).

3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-6-methyl-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester:

To a solution of 3-(6-Methyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-hexanoic acid ethyl ester (50 mg, 0.17 mmol) in DMF (2 mL) were added K₂CO₃ (71 mg, 0.52 mmol) and (1,4-Dimethyl-1*H*-indol-3-ylmethyl)-trimethyl-ammonium; iodide (90 mg, 0.26 mmol). The reaction mixture was heated to 80 °C for 16 h. The reaction mixture was then diluted with ethyl acetate and


washed with water (x4). The organic phase was then dried over Na_2SO_4 and concentrated. The resulting residue was purified by silica gel prep TLC using 4:1 Hexanes: ethyl acetate as an eluent to afford 35 mg of the desired product as oil. LCMS (ESMS): m/z 448.82 ($\text{M}+\text{H}^+$).

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid:


To a solution of the ester (35 mg, 0.078 mmol) in MeOH (5 mL) was added 2N NaOH (5 mL). The reaction mixture was stirred at room temperature for 3 h. When the reaction was complete, the mixture was treated with 1N HCl and extracted with CH_2Cl_2 . The organic phase was dried over Na_2SO_4 and concentrated. The resulting residue was purified by silica gel prep TLC using 10:1 CH_2Cl_2 :MeOH as an eluent to afford 27 mg of the title compound as an off-white solid. LCMS (ESMS): m/z 420.63 ($\text{M}+\text{H}^+$).

The following compounds were prepared following a similar procedure using appropriate starting materials.


(3R)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-phenylpropanoic acid. LCMS (ESMS): m/z 440.17 ($\text{M}+\text{H}^+$).




(3R)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-pyridin-3-ylpropanoic acid. LCMS (ESMS): m/z 455.51 ($\text{M}+\text{H}^+$).




3-{5-bromo-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid. LCMS (ESMS): m/z 484.38 ($M+H^+$).




(3S)-3-{5-[(tert-butoxycarbonyl)amino]-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid. LCMS (ESMS): m/z 520.62 ($M+H^+$).




(1R,2R)-2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclohexanecarboxylic acid LCMS (ESMS): m/z 418 ($M+H^+$).




1-({3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)cyclohexanecarboxylic acid. LCMS (ESMS): m/z 432.25 ($M+H^+$).



1-({3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)cyclopentanecarboxylic acid. LCMS (ESMS): m/z 418.23 ($M+H^+$).



Example 12: (2E)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}acrylic acid



(E)-3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-acrylic acid ethyl ester:

To a solution of 1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one (0.1 g, 0.34 mmol) in AcCN (2 mL) and DMF (1 mL) were added ethyl propiolate (0.04 mL, 0.34 mmol) and DMAP (0.02 mL, 0.17 mmol). The reaction mixture was stirred at room temperature for 2h. When the reaction was complete, the reaction mixture was diluted with CH₂Cl₂ and washed with water. The organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC using 95:5 CH₂Cl₂:MeOH as an eluent to afford 0.11 g of the desired product as a white solid. LCMS (ESMS): *m/z* 393.13 (M+H⁺).


(2E)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} acrylic acid:

To a solution of the ester (0.11 g, 0.28 mmol) in MeOH (10 mL) was added 2N NaOH (5 mL). The reaction mixture was stirred at 60 °C for 2 h until all the starting material was consumed. The reaction mixture was then treated with 1M HCl to pH ~2 and extracted with CH₂Cl₂ (x2).


The organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC using 10:1 CH₂Cl₂:MeOH as an eluent to afford 20 mg of the title compound as an off-white solid. LCMS (ESMS): m/z 365.03 (M+H⁺).

The by-product was obtained during hydrolysis of (E)-3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-acrylic acid ethyl ester.

3-methoxy-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid LCMS (ESMS): m/z 397.04 (M+H⁺).



Example 13: 3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid



1-(5-Chloro-1,2,3-trimethyl-1*H*-indol-7-ylmethyl)-3-isopropenyl-5-methyl-1,3-dihydro-benzimidazol-2-one:

To a solution of 1-Isopropenyl-6-methyl-1,3-dihydro-benzimidazol-2-one (50 mg, 0.27 mmol) and (5-Chloro-1,2,3-trimethyl-1*H*-indol-7-yl)-methanol (60 mg, 0.27 mmol), triphenylphosphine (84 mg, 0.32 mmol) in THF (2mL) was added dropwise DIAD (0.06 mL, 0.32 mmol). The mixture was stirred at room temperature for 1 h. The reaction mixture was diluted with ethyl acetate and washed with water (x2). The organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC using 4:1 Hexanes:ethyl acetate as an eluent to afford 60 mg of the desired product as a brownish oil. LCMS (ESMS): *m/z* 394.28 (M+H⁺).

1-(5-Chloro-1,2,3-trimethyl-1*H*-indol-7-ylmethyl)-5-methyl-1,3-dihydro-benzimidazol-2-one:

To a solution of 1-(5-Chloro-1,2,3-trimethyl-1*H*-indol-7-ylmethyl)-3-isopropenyl-5-methyl-1,3-dihydro-benzimidazol-2-one (60 mg, 0.15 mmol) in a mixture of MeOH (3 mL) and water (3 mL), was added 1N HCl (2 mL). The reaction mixture was heated to 60 °C for 2 h. When the

reaction was completed, the mixture was extracted with CH₂Cl₂ (x2). The organic phase was dried over Na₂SO₄ and concentrated to afford 40 mg of the desired product. The resulting residue was used for the next reaction without further purification.


3-[3-(5-Chloro-1,2,3-trimethyl-1*H*-indol-7-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid methyl ester:

To a solution of 1-(5-Chloro-1,2,3-trimethyl-1*H*-indol-7-yl)methyl-5-methyl-1,3-dihydro-benzimidazol-2-one (40 mg, 0.11 mmol) in DMA (2 mL) were added Methyl 2-pentenoate (26 mg, 0.23 mmol) and K₂CO₃ (31 mg, 0.23 mmol). The reaction mixture was heated to 120 °C for 20 min in a microwave. When the reaction was completed, the reaction mixture was diluted with ethyl acetate and washed with water (x4). The organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC using 4:1 Hexane:ethyl acetate to afford 40 mg of the desired product. LCMS (ESMS): *m/z* 468.36 (M+H⁺).

3-{3-[(5-chloro-1,2,3-trimethyl-1*H*-indol-7-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}pentanoic acid:

To a solution of the ester (40 mg, 0.09 mmol) in MeOH (2 mL) was added 2N NaOH (2 mL). The reaction mixture was not homogeneous at room temperature. The reaction mixture was therefore heated to 60 °C for 1 h. When the reaction was complete, the mixture was treated with 1N HCl and extracted with CH₂Cl₂ (x2). The organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC using 10:1 CH₂Cl₂:MeOH as an eluent to afford the title compound as an off-white solid. LCMS (ESMS): *m/z* 454.20 (M+H⁺). However, this product also contains approximately 50% of the other isomer, 3-{3-[(5-chloro-1,2,3-trimethyl-1*H*-indol-7-yl)methyl]-5-methyl-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}pentanoic acid.

Example 14: *N*-[3-(3-Naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionyl]-benzenesulfonamide



1-Isopropenyl-3-naphthalen-1-ylmethyl-1,3-dihydro-benzimidazol-2-one:

To a solution of 1-isopropenylbenzimidazolone (1.0 g, 5.7 mmol) in dry THF (80 mL) are added triphenylphosphine (1.8 g, 6.9 mmol) and naphthalen-1-yl-methanol (1.1 g, 6.9 mmol). Then diisopropyl azodicarboxylate (1.4 mL, 6.9 mmol) is added drop wise into the above solution at room temperature. The mixture is stirred for 16 hrs and then the solvent is removed under vacuum. The residue is purified by flash column chromatography using 10% EtOAc in hexanes to give 1.6 g (89%) of 1-isopropenyl-3-naphthalen-1-ylmethyl-1,3-dihydro-benzimidazol-2-one.

1-Naphthalen-1-ylmethyl-1,3-dihydro-benzimidazol-2-one:

1-Isopropenyl-3-naphthalen-1-ylmethyl-1,3-dihydro-benzimidazol-2-one (1.6 g, 5.1 mmol) is dissolved in MeOH (5.0 mL) and water (5.0 mL) is added into the solution. To this heterogeneous reaction mixture is added 37% hydrochloric acid (4.0 mL). The reaction mixture is heated to 60 °C for 30 min and then water (50 mL) and CH₂Cl₂ (50 mL) are added. The organic layer is separated and the aqueous layer is extracted with CH₂Cl₂ (3 x 50 mL). The organic layers are combined, washed with Sat. NaHCO₃ (100 mL), Sat. NH₄Cl (100 mL) and water (3 x 100 mL), dried over MgSO₄ and concentrated to give 1.3 g (93%) of crude product which is pure enough for the next step.

3-(3-Naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid methyl ester:

To a solution of 1-naphthalen-1-ylmethyl-1,3-dihydro-benzimidazol-2-one (200 mg, 0.73 mmol) in DMF (3.0 mL) are added methyl acrylate (0.07 mL, 0.8 mmol) and 40% benzyltrimethyl ammonium hydroxide in MeOH (0.03 mL, 0.073 mmol). The resulting mixture is stirred at room temperature for 1.5 h and then Sat. NH_4Cl (5 mL) and water (20 mL) are added. The mixture is extracted with EtOAc (3 x 50 mL) and the organic layers are combined, washed with water (3 x 75 mL), dried over MgSO_4 and concentrated to give crude product. Purification by flash column chromatography using 30% EtOAc in Hexanes affords 220 mg (84%) of 3-(3-naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid methyl ester.

3-(3-Naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid:

Lithium hydroxide monohydrate (31 mg, 0.75 mmol) is dissolved in water (3.5 mL) and this solution is added into 3-(3-naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid methyl ester (180 mg, 0.5 mmol) in methanol (3.5 mL) at room temperature. The mixture is stirred for 6 hr and 1.0 M HCl aqueous solution (1.5 mL) is added along with water (25 mL). Then the mixture is extracted with EtOAc (3 x 50 mL) and the organic layers are combined, dried over MgSO_4 and concentrated to give 170 mg of crude 3-(3-Naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid which is used in the following step without further purification.

N-[3-(3-Naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionyl]-benzenesulfonamide:


3-(3-Naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid (80 mg, 0.23 mmol) is dissolved in dry THF (3.0 mL) and 1,1'-carbonyl diimidazole (84 mg, 0.52 mmol) is added into it. The mixture is stirred for 30 min at room temperature and then it is heated at 55 °C for 1 hr. After the mixture is cooled down to room temperature, benzenesulfonamide (73 mg, 0.46 mmol) is added and after 10 min, 1,8-diazabicyclo[5,4,0]undec-7-ene (0.07 mL, 0.46 mmol) is added. The mixture is again stirred for 8 hr at room temperature. 1.0 M HCl aqueous solution (3 mL) is added followed by water (30 mL). The mixture is extracted with EtOAc (3 x 50 mL) and the organic layers are combined, dried and concentrated to give crude product. Purification by preparative TLC using 55% EtOAc in Hexanes affords 66 mg (59%) of *N*-[3-(3-

Naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionyl]-benzenesulfonamide.


LCMS (ESMS): m/z 486.05 ($M+H^+$)

The following compound is synthesized using the same procedure as Example 1 by replacing benzenesulfonamide with commercially available methane sulfonamide:

N-[3-(3-Naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionyl]-methanesulfonamide. LCMS (ESMS): m/z 424.06 ($M+H^+$)




Example 15: 3-(3-Naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionamide



To the THF (1.0 mL) suspension of 3-(3-Naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid (50 mg, 0.14 mmol) at room temperature are added ammonia 0.5 M solution in dioxane (1.4 mL, 0.72 mmol) and 1-hydroxybenzotriazole (94 mg, 0.7 mmol). Then *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide (118 mg, 0.61 mmol) is added in small portions. The reaction mixture is stirred for 20 hr and then water (5 mL) is added and the

mixture is extracted with EtOAc (3 x 10 mL). The organic layers are combined, dried over MgSO_4 and concentrated to give crude product. Purification by preparative TLC using 5% MeOH in CH_2Cl_2 affords 37 mg (74%) of 3-(3-Naphthalen-1-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionamide. LCMS (ESMS): m/z 346.12 ($\text{M}+\text{H}^+$).

Example 16: *N*-{3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionyl}-benzenesulfonamide



1-Isopropenyl-3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one:

To a solution of 1-isopropenylbenzimidazolone (500 mg, 2.8 mmol) in dry THF (40 mL) are added triphenylphosphine (902 mg, 3.4 mmol) and (4-Methyl-benzo[*b*]thiophen-3-yl)-methanol (563 mg, 3.2 mmol). Then diisopropyl azodicarboxylate (0.70 mL, 3.4 mmol) is added drop wise into the above solution after it is cooled down to 0 °C. The mixture is then stirred at room temperature for 5 hr and then the solvent is removed under vacuum and the residue is purified by flash column chromatography using gradient EtOAc/Hexane from 0% to 20% to give 630mg (66%) of 1-isopropenyl-3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one.

1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one:

1-Isopropenyl-3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one (630 mg, 1.9 mmol) is dissolved in methanol (2.0 mL), 1,4-dioxane (10 mL) and water (2.0 mL). Then 37% HCl (1.6 mL) is added and the reaction mixture is heated at 65 °C for 6 hr. A white solid is formed during the heating. Then water (80 mL) and CH₂Cl₂ (80 mL) are added. The organic layer is separated and the aqueous layer is extracted with CH₂Cl₂ (3 x 50 mL). The organic layers are combined, washed with Sat. NaHCO₃ (100 mL), Sat. NH₄Cl (100 mL) and water (3 x 100 mL), dried over MgSO₄ and concentrated to give 560 mg (99%) of 1-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one which is used for the next step without further purification.

3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid methyl ester:

To a solution of 1-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one (560 mg, 1.9 mmol) in DMF (9.0 mL) are added methyl acrylate (0.19 mL, 2.1 mmol) and 40% benzyltrimethyl ammonium hydroxide in MeOH (0.09 mL, 0.19 mmol). The resulting mixture is stirred at room temperature for 3 hr. Then Sat. NH₄Cl (5 mL) and water (50 mL) are added and the mixture is extracted with EtOAc (3 x 50 mL). The organic layers are combined, washed with water (3 x 75 mL), dried over MgSO₄ and concentrated to give crude product. Purification by flash column chromatography using 30% EtOAc in hexanes affords 600 mg (83%) of 3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid methyl ester.

3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid:

Lithium hydroxide monohydrate (99 mg, 2.4 mmol) is dissolved in water (12 mL) and this solution is added into 3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid methyl ester (600 mg, 1.6 mmol) in methanol (12 mL) at room temperature. The mixture is stirred for 2 hr and then 1.0 M HCl aqueous solution (3 mL) is added along with water (100 mL). Then the mixture is extracted with EtOAc (3 x 100 mL) and the organic layers are combined, dried over MgSO₄ and concentrated to give about 560 mg


(97%) of 3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid which is used in the next step without further purification.

N-{3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionyl}-benzenesulfonamide:


3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid (150 mg, 0.41 mmol) is dissolved in dry THF (3.0 mL) and 1,1'-carbonyl diimidazole (150 mg, 0.92 mmol) is added into it at room temperature. The mixture is stirred for 30 min and then it is heated at 55 °C for 1 hr. After the mixture is cooled down to room temperature, benzenesulfonamide (130 mg, 0.82 mmol) is added and after 10 min, 1,8-diazabicyclo[5,4,0]undec-7-ene (0.12 mL, 0.82 mmol) is added. The mixture is stirred for 5 hr at room temperature and then 1.0 M HCl (3 mL) is added followed by water (30 mL). The mixture is extracted with EtOAc (3 x 50 mL) and the organic layers are combined, dried and concentrated to give crude product. Purification by preparative TLC using 75% EtOAc in Hexanes affords 123mg (60%) of *N*-{3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionyl}-benzenesulfonamide. LCMS (ESMS): *m/z* 506.18 (*M*+*H*⁺).

The following compounds are synthesized using the same procedure.

N-{3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionyl}-methanesulfonamide. LCMS (ESMS): *m/z* 444.00 (*M*+*H*⁺).




Propane-2-sulfonic acid {3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionyl}-amide. LCMS (ESMS): *m/z* 472.07 (*M*+*H*⁺).



Propane-2-sulfonic acid amide:

Propane-2-sulfonyl chloride (0.11 mL, 0.98 mmol) is added drop wise into the mixture of 28% NH_4OH in H_2O (2.0 mL) and CHCl_3 (2.0 mL) at room temperature. The mixture is stirred for 2 hrs at that temperature and then the solvents are removed under vacuum. THF (50 mL) is added into the residue to form a white suspension. The solid is filtered and the filtrate is concentrated to give 86 mg of propane-2-sulfonic acid amide which is used in the next step without further purification.

Example 17: 2,2-Dimethyl-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid




2,2-Dimethyl-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid methyl ester:

To a stirred solution of 1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one (50 mg, 0.17 mmol), 3-Hydroxy-2,2-dimethyl-propionic acid methyl ester (0.026 mL, 0.20 mmol) and triphenylphosphine (53 mg, 0.20 mmol) in dry THF (1.5 mL) is added diisopropyl azodicarboxylate (0.041 mL, 0.20 mmol) in a drop wise manner. The mixture is stirred at room temperature for 40 hrs. Then the solvent is removed under vacuum and the residue is purified by two preparative TLC's (first one is developed with 30% EtOAc in Hexanes, second one is developed with 1% MeOH in CH₂Cl₂) to give 28 mg (41%) of 2,2-Dimethyl-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid methyl ester.

2,2-Dimethyl-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid:

Lithium hydroxide monohydrate (4.3 mg, 0.10 mmol) is dissolved in water (1.0 mL) and it is added into the solution of 2,2-Dimethyl-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid methyl ester (28 mg, 0.069 mmol) in 1,4-dioxane (2.0 mL) at room temperature. The mixture is stirred for 60 hr and LCMS then 1.0 M HCl solution (1.0 mL) is added followed by water (15 mL). Then the mixture is extracted with EtOAc (3 x 30 mL) and the organic layers are combined, dried over MgSO₄ and concentrated to give 25 mg (90%) of 2,2-Dimethyl-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid. LCMS (ESMS): *m/z* 395.17 (M+H⁺).

Example 18: *N*-{3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-benzenesulfonamide



3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester:

To a solution of 1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one (50 mg, 0.17 mmol) in dry DMF (1.0 mL) are added (E)-Hex-2-enoic acid ethyl ester (27 mg, 0.19 mmol) and 40% benzyltrimethyl ammonium hydroxide in MeOH (0.008 mL, 0.017 mmol). The resulting mixture is warmed up to 60 °C for 16 hr. Then another 27 mg of (E)-Hex-2-enoic acid ethyl ester is added the heating is continued for another 8 hr. Then saturated NH₄Cl (2 mL) and water (10 mL) are added and the mixture is extracted with EtOAc (3 x 25 mL). The organic layers are combined, dried over MgSO₄ and concentrated to give the crude product. Purification with flash column chromatography using 20% EtOAc in Hexanes affords 70 mg (95%) of 3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester.

3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid:

To a solution of 3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester (65 mg, 0.15 mmol) in MeOH (3.0 mL) is added 2.0 N NaOH (0.5 mL). The reaction mixture is stirred at room temperature for 3.5 h and then 1.0M HCl (1.5 mL) and water (50 mL) are added. The mixture is extracted with EtOAc (3 x 50 mL). The organic layers are combined, dried over MgSO₄ and concentrated to give crude product. Purification by flash column chromatography affords 56 mg (92%) of 3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid.


N-{3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-benzenesulfonamide:

3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid (56 mg, 0.14 mmol) is dissolved in dry THF (1.0 mL) and 1,1'-carbonyl diimidazole (50 mg, 0.29 mmol) is added into it at room temperature. The mixture is stirred for 30 min and then it is heated at 55 °C for 1hr. After the mixture is cooled down to room temperature, benzenesulfonamide (46 mg, 0.29 mmol) is added and after 10 min, 1,8-diazabicyclo[5,4,0]undec-7-ene (0.044 mL, 0.29 mmol) is added. The mixture is stirred for 3 hr


at room temperature. Then 1.0 M HCl (3 mL) is added followed by water (30 mL). The mixture is extracted with EtOAc (3 x 50 mL) and the organic layers are combined, dried and concentrated to give crude product. Purification by preparative TLC using 2% MeOH in CH₂Cl₂ affords 63 mg (84%) of *N*-{3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-benzenesulfonamide. LCMS (ESMS): *m/z* 548.11 (M+H⁺).

The following compounds are synthesized using the same procedure.


N-{3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-methanesulfonamide. LCMS (ESMS): *m/z* 486.29 (M+H⁺).




Cyclopropanesulfonic acid {3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-amide. LCMS (ESMS): *m/z* 512.26 (M+H⁺).




Pyridine-3-sulfonic acid {3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-amide. LCMS (ESMS): *m/z* 549.16 (M+H⁺).




4-Methoxy-*N*-{3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-benzenesulfonamide. LCMS (ESMS): m/z 578.21 ($M+H^+$).




3-Methoxy-*N*-{3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-benzenesulfonamide. LCMS (ESMS): m/z 578.19 ($M+H^+$).




Pyridine-2-sulfonic acid {3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-amide. LCMS (ESMS): m/z 549.25 ($M+H^+$).




2-Methoxy-*N*-{3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-benzenesulfonamide. LCMS (ESMS): m/z 578.35 ($M+H^+$).




2-Fluoro-*N*-{3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-benzenesulfonamide. LCMS (ESMS): m/z 566.28 ($M+H^+$).




4-Fluoro-*N*-{3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-benzenesulfonamide. LCMS (ESMS): m/z 566.17 ($M+H^+$).




3-Fluoro-*N*-(3-{3-[4-((*Z*)-1-methyl-propenyl)-5-vinyl-thiophen-3-ylmethyl]-2-oxo-2,3-dihydro-benzimidazol-1-yl}-hexanoyl)-benzenesulfonamide. LCMS (ESMS): m/z 566.17 ($M+H^+$).



3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-*N*-[(1-methyl-1H-imidazol-4-yl)sulfonyl]hexanamide. LCMS (ESMS): m/z 552.20 ($M+H^+$).




N-[(1,2-dimethyl-1H-imidazol-4-yl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide. LCMS (ESMS): m/z 566.24 ($M+H^+$).




The following compounds were prepared following the same procedure using the acids that were synthesized in Example 6.

4-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)butanamide. LCMS (ESMS): m/z 520.09 ($M+H^+$).




2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)acetamide. LCMS (ESMS): m/z 492.15 ($M+H^+$).




The following compounds were prepared following the same procedure using the acids that were synthesized in Example 11.


3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)hexanamide. LCMS (ESMS): m/z 559.31 ($M+H^+$).




N-(tert-butylsulfonyl)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide. LCMS (ESMS): m/z 539.36 ($M+H^+$).



3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(ethylsulfonyl)hexanamide. LCMS (ESMS): m/z 511.25 ($M+H^+$).



Example 19: 1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-3-[2-(1H-tetrazol-5-yl)-ethyl]-1,3-dihydro-benzimidazol-2-one




3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionitrile:

To a solution of 1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one (50 mg, 0.17 mmol) in dry DMF (1.0 mL) are added acrylonitrile (0.012 mL, 0.19 mmol) and 40% benzyltrimethyl ammonium hydroxide in MeOH (0.008 mL, 0.017 mmol). The resulting mixture is stirred at room temperature for 2hr and then Sat.NH₄Cl (2 mL) is added along with water (30 mL). Then the mixture is extracted with EtOAc (3 x 30 mL). The organic layers are combined, dried over MgSO₄ and concentrated to give crude product. Purification by flash column chromatography affords 58 mg (98%) of 3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionitrile.

1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-3-[2-(1*H*-tetrazol-5-yl)-ethyl]-1,3-dihydro-benzimidazol-2-one:

3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionitrile (58 mg, 0.17 mmol), azidotrimethylsilane (0.047 mL, 0.33 mmol) and dibutyltin oxide (4.2 mg, 0.017 mmol) are dissolved in dry DMF (2.0 mL) in microwave vial. It is then heated at 150 °C for 2 hrs in microwave reactor. Then another 0.05 mL of azidotrimethylsilane and 4.0 mg of dibutyltin oxide are added and the mixture is heated for another 2 hr at 150 °C in microwave reactor. Then water (2 mL) is added and the mixture is extracted with EtOAc (2x30 mL). The organic layers are combined and extracted with 2.0M NaOH (2 x 30 mL). The NaOH layers are combined and washed with EtOAc (30 mL). Then the NaOH layers are acidified with 1.0M HCl to pH=3 and it is then extracted with EtOAc (2 x 30 mL). All the organic layers are combined, dried (MgSO₄) and concentrated to give crude product. Purification by preparative TLC using 3% MeOH in CH₂Cl₂ affords 15 mg (65%) of 1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-3-[2-(1*H*-tetrazol-5-yl)-ethyl]-1,3-dihydro-benzimidazol-2-one. LCMS (ESMS): *m/z* 391.07 (M+H⁺).

Example 20: 5-Hydroxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid:



5-Hydroxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid methyl ester and 1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-3-(2-oxo-tetrahydro-pyran-4-yl)-1,3-dihydro-benzimidazol-2-one:

To a solution of 1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one (50 mg, 0.17 mmol) in DMF (0.7 mL) are added 5,6-Dihydro-pyran-2-one (0.059 mL, 0.68 mmol) and 40% benzyltrimethyl ammonium hydroxide in MeOH (0.008 mL, 0.017 mmol). The resulting mixture is stirred for 5 days and then saturated NH_4Cl (2 mL) and water (30 mL) are added. The mixture is extracted with EtOAc (3 x 35 mL). The organic layers are combined, dried over MgSO_4 and concentrated to give the crude product. Purification with flash column chromatography using EtOAc in Hexane afforded 51 mg of mixture of 5-Hydroxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid methyl ester and 1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-3-(2-oxo-tetrahydro-pyran-4-yl)-1,3-dihydro-benzimidazol-2-one. They are used in the next step without further separation.


5-Hydroxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid:

5-Hydroxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid methyl ester and 1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-3-(2-oxo-tetrahydro-pyran-4-yl)-1,3-dihydro-benzimidazol-2-one (51 mg) are dissolved in 1,4-dioxane (2.5 mL). Then 1.0 N NaOH solution (2.0 mL) is added and the mixture is stirred for 2 hr. Then 1.0 M HCl (3.0 mL) is added to make it acidic and the mixture is extracted with EtOAc (3 x 35 mL). The organic layers are combined, dried and concentrated to give crude product.

Purification by preparative TLC using 7.5% MeOH in CH_2Cl_2 affords 43 mg (62% for two

steps) of 5-Hydroxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid. LCMS (ESMS): m/z 411.07 ($M+H^+$).

Example 21: 3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanedioic acid monomethyl ester:



5-Hydroxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid methyl ester:


The crude mixture of 5-Hydroxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid methyl ester and 1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-3-(2-oxo-tetrahydro-pyran-4-yl)-1,3-dihydro-benzimidazol-2-one (320 mg) is dissolved in MeOH (30 mL) and then one drop of concentrated H_2SO_4 is added. The mixture is stirred at room temperature for 30 min, and sat. $NaHCO_3$ (2 mL) along with water (20 mL) are added. The mixture is then extracted with EtOAc (3 x 40 mL). The organic layers are combined, dried ($MgSO_4$) and concentrated to give crude product. Purification by flash column chromatography using EtOAc in Hexane affords 70 mg (25% for two steps) of 5-Hydroxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid methyl ester.

3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanedioic acid monomethyl ester:

5-Hydroxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid methyl ester (70 mg, 0.17 mmol) is dissolved in DMF (1.5 mL) at room temperature and pyridinium dichromate (250 mg, 0.66 mmol) is added into it. The mixture is stirred for 20 hr and water (25 mL) is added. The pH of the solution is adjusted to pH 5 by

adding acetic acids and the mixture is extracted with EtOAc (3 x 50 mL). The organic layers are combined, dried and concentrated to give crude product. Purification by preparative TLC using 80% EtOAc in Hexanes with 1% acetic acid affords 48 mg (66%) of 3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanedioic acid monomethyl ester. LCMS (ESMS): m/z 439.17 ($M+H^+$).


Example 22: 3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanedioic acid:



3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanedioic acid:

Lithium hydroxide monohydrate (7.2 mg, 0.17 mmol) is dissolved in water (1.0 mL) and it is added into the solution of 3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanedioic acid monomethyl ester (25 mg, 0.057 mmol) in methanol (1.0 mL) at room temperature. The mixture is stirred for 2 hr and then 1.0 M HCl solution (0.5 mL) is added along with water (10 mL). The mixture is extracted with EtOAc (3 x 10 mL) and the organic layers are combined, dried over $MgSO_4$ and concentrated to give crude product. Purification by preparative TLC using 5 % MeOH in CH_2Cl_2 with 1% acetic acid afforded 21 mg (87%) of 3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanedioic acid. LCMS (ESMS): m/z 425.22 ($M+H^+$).


Example 23: 3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid benzylamide:



3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid benzylamide:

To the suspension of 3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid (20 mg, 0.049 mmol) in THF (1.0 mL) at room temperature are added benzyl amine (0.027 mL, 0.245 mmol) and 1-hydroxybenzotriazole (32 mg, 0.24 mmol). Then *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide (40 mg, 0.21 mmol) is added and the reaction mixture is stirred for 16 hr. Water (5 mL) is added and the mixture is extracted with EtOAc (3 x 10 mL). The organic layers are combined, dried over MgSO_4 and concentrated to give crude product. The crude is filtered through a short path of silica gel eluting first with CH_2Cl_2 then with 50% EtOAc in hexanes to give 23 mg (95%) of 3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid benzylamide. LCMS (ESMS): m/z 498.22 ($\text{M}+\text{H}^+$).

Example 24: *N*-{3-Ethoxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionyl}-benzenesulfonamide:



3-Ethoxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid ethyl ester:

To a solution of 1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one (50 mg, 0.17 mmol) in dry DMF (0.7 mL) are added (E)-3-Ethoxy-acrylic acid ethyl ester

(0.049 mL, 0.34 mmol) and 40% benzyltrimethyl ammonium hydroxide in MeOH (0.008 mL, 0.017 mmol). The resulting mixture is warmed up to 60 °C for 16 hr and then saturated NH₄Cl solution (1 mL) is added followed by water (5 mL). The mixture is extracted with EtOAc (3 x 10 mL). The organic layers are combined, dried and concentrated to give crude product. Purification by flash column chromatography using EtOAc in hexane (from 0% to 30%) affords 50 mg (67%) of 3-Ethoxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid ethyl ester.

3-Ethoxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid:


To a solution of 3-Ethoxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid ethyl ester (50 mg, 0.11 mmol) in MeOH (3.0 mL) is added 2.0 N NaOH solution (1.0 mL). The reaction mixture is stirred at room temperature for 2 h. Then 1.0 M HCl solution (3.0 mL) and water (10 mL) are added and the reaction mixture is extracted with EtOAc (3 x 20 mL). The organic layers are combined, dried and concentrated to give 45 mg (96%) of the crude 3-Ethoxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid which is used in the next step without further purification.

N-{3-Ethoxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionyl}-benzenesulfonamide:

3-Ethoxy-3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid (45 mg, 0.11 mmol) is dissolved in dry THF (1.0 mL) and 1,1'-carbonyl diimidazole (40 mg, 0.25 mmol) is added into it at room temperature. The mixture is stirred for 30 min and then it is heated at 55 °C for 1 hr. After the mixture is cooled down to room temperature, benzenesulfonamide (34 mg, 0.22 mmol) is added and after 10 min, 1,8-diazabicyclo[5,4,0]undec-7-ene (0.033 mL, 0.22 mmol) is added. The mixture is stirred for 4 hr at room temperature and 1.0 M HCl solution (3 mL) is added followed by water (30 mL). The mixture is extracted with EtOAc (3 x 50 mL) and the organic layers are combined, dried and concentrated to give crude product. Purification by preparative TLC using 6% MeOH in CH₂Cl₂ with 1.0% concentrated aqueous NH₄OH solution (28%) affords 22 mg (36%) of *N*-{3-Ethoxy-

3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionyl}-benzenesulfonamide. LCMS (ESMS): m/z 550.15 ($M+H^+$).


Example 25: 2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethanesulfonic acid phenyl ester:



2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethanesulfonic acid phenyl ester:

To a solution of 1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one (50 mg, 0.17 mmol) in THF (1.0 mL) are added Ethenesulfonic acid phenyl ester (63 mg, 0.34 mmol) and solid NaOH (14 mg, 0.34 mmol). The resulting suspension is stirred at room temperature for 0.5 hr and it turned into clear solution in about 10 min. Then saturated NH_4Cl solution (5.0 mL) is added along with water (5 mL). The mixture is extracted with EtOAc (3 x 20 mL). The organic layers are combined, dried ($MgSO_4$) and concentrated to give crude product. Purification by flash column chromatography using EtOAc in Hexanes (from 0% to 30%) affords 61 mg (75%) of 2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethanesulfonic acid phenyl ester. LCMS (ESMS): m/z 479.11 ($M+H^+$).


Example 26: *N*-Methyl-*N*-{3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-benzenesulfonamide:



N-Methyl-*N*-{3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-benzenesulfonamide:

N-{3-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-benzenesulfonamide (33 mg, 0.061 mmol) and Iodomethane (0.015 mL, 0.24 mmol) are added into DMF (1.0 mL) at room temperature. Then cesium carbonate (80 mg, 0.24 mmol) is added and the mixture is warmed up to 65 °C for 10 min. Then it is cooled down to room temperature for 16 hrs. Saturated NH₄Cl solution (3.0 mL) and water (10 mL) are added and the mixture is extracted with EtOAc (3 x 20 mL). The organic layers are combined, dried (MgSO₄) and concentrated to give crude product. Purification by flash column chromatography using EtOAc in hexanes (from 0% to 30%) affords 24 mg (71%) of *N*-Methyl-*N*-{3-[3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-benzenesulfonamide. LCMS (ESMS): *m/z* 562.19 (M+H⁺).

Example 27: 2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethanesulfonic acid:




2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethanesulfonic acid:

2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethanesulfonic acid phenyl ester (25 mg, 0.052 mmol) is dissolved in ethanol (1.0 mL) and THF (1.0 mL), 2.0 N NaOH solution (1.0 mL) is added into it and the mixture is warmed up to 80 °C for 3hr. Then 1.0 M HCl solution (5.0 mL) is added along with water (15 mL). The mixture is extracted with EtOAc (3 x 35 mL) and the organic layers are combined, dried (MgSO₄) and concentrated to give crude product. The crude is then dissolved in 10% MeOH in CH₂Cl₂ and purified by column chromatography eluting with 10% MeOH in CH₂Cl₂ to give 12 mg (58%) of

2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethanesulfonic acid. LCMS (ESMS): m/z 403.08 ($M+H^+$).

Example 28: 2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethanesulfonic acid amide:




Ethenesulfonic acid amide:

2-Chloro-ethanesulfonyl chloride (0.32 mL, 3.1 mmol) is added into the mixture of $CHCl_3$ (1.0 mL) and 28% ammonium hydroxide aqueous solution (0.85 mL, 6.2 mmol) at room temperature. The mixture is stirred for 1.5 hr and then the solvents are removed under vacuum. THF (50 mL) is added to dissolve the organic portion of the crude. The mixture is filtered and the filtrate is concentrated to give 100 mg (31%) of ethenesulfonic acid amide.

2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethanesulfonic acid amide:

To the solution of 1-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one (50 mg, 0.17 mmol) in THF (1.0 mL) are added ethenesulfonic acid amide (40 mg, 0.34 mmol) and solid NaOH (14 mg, 0.34 mmol). The mixture is warmed up to 65 °C for 24 hr and then saturated NH_4Cl solution (5.0 mL) is added along with water (5.0 mL). The mixture is extracted with EtOAc (3 x 20 mL). The organic layers are combined, dried (Na_2SO_4) and concentrated to give crude product. Purification by preparative TLC using 4% MeOH in CH_2Cl_2 affords 31 mg (46%) of 2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethanesulfonic acid amide. LCMS (ESMS): m/z 402.10 ($M+H^+$).

Example 29: 2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethanesulfonic acid benzoylamide:





2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethanesulfonic acid benzoyl amide:

Benzoic acid (6.1 mg, 0.050 mmol) is dissolved in THF (0.50 mL) and 1,1'-carbonyl diimidazole (9.1 mg, 0.056 mmol) is added into it at room temperature. The mixture is stirred for 30 min and then it is heated at 55 °C for 1 hr. After the mixture is cooled down to room temperature, 2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethanesulfonic acid amide (10 mg, 0.025 mmol) is added and after 10 min, 1,8-diazabicyclo[5,4,0]undec-7-ene (0.007 mL, 0.050 mmol) is added. The mixture is stirred for 16 hr at room temperature. 1.0 M HCl solution (1.0 mL) is added followed by water (10 mL). The mixture is extracted with EtOAc (3 x 20 mL) and the organic layers are combined, dried and concentrated to give crude product. Purification by preparative TLC using 6% MeOH in EtOAc affords 11 mg (85%) of 2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethanesulfonic acid benzoyl amide. LCMS (ESMS): m/z 506.25 ($M+H^+$).

The following compound is synthesized using the same procedure.

2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethanesulfonic acid acetyl-amide. LCMS (ESMS): m/z 444.07 ($M+H^+$).




Example 30: 3-[3-(2,5-Dimethyl-benzyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid.


3-(2-Oxo-2,3-dihydro-benzoimidazol-1-yl)-hexanoic acid ethyl ester (40 mg, 0.145 mmol), prepared as described above, was dissolved in DMF (1 mL), and potassium carbonate (69 mg, 0.50 mmol) was added followed by 2-bromomethyl-1,4-dimethyl-benzene (56.3 mg, 0.28 mmol). The resulting mixture was heated at 80°C for 4 h, cooled, filtered, and the solution was treated directly with the Amberlyst® resin (524mg, 0.7 mmol). This reaction mixture was agitated overnight and the resin was then filtered and washed with several portions of methanol. The product was eluted by treating the resin with a 20% solution of formic acid in methanol (2x3 mL) and washing with methanol (2x3 mL). The combined eluant and final wash was evaporated to provide the title compound (25 mg, 47%). LCMS (ESMS): m/z 367 ($M+H^+$).

Using methods similar to those described in the above example, the following analogs were also synthesized:


3-[3-(5-Chloro-2-difluoromethoxy-benzyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid. LCMS (ESMS): m/z 439 ($M+H^+$).




3-[3-(5-Chloro-2-trifluoromethyl-benzyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid. LCMS (ESMS): m/z 441 ($M+H^+$).




3-{3-[2-bromo-5-(trifluoromethyl)benzyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid. LCMS (ESMS): m/z 485.7 ($M+H^+$).



3-[2-oxo-3-(2,3,5-trichlorobenzyl)-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid. LCMS (ESMS): m/z 441.0 ($M+H^+$).



Example 31: 3-[3-(5-Bromo-2-methoxy-benzyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid




3-(2-Oxo-2,3-dihydro-benzoimidazol-1-yl)-hexanoic acid ethyl ester (96.7 mg, 0.35 mmol), prepared as described above, (5-bromo-2-methoxy-phenyl)-methanol (152 mg, 0.7 mmol) and triphenylphosphine (183.6, 0.7 mmol) were combined in a reaction vial and THF (1 mL) was


added. The mixture was stirred until complete solution was attained, then cooled to 0°C, and diisopropylazodicarboxylate (0.136 mL, 0.7 mmol) was added dropwise. The reaction mixture was agitated at room temperature overnight, diluted with THF (2 mL) then treated directly with the Amberlyst® resin (3 g, 4 mmol). This reaction mixture was agitated overnight and the resin was then filtered and washed with DMF (2 x 3 mL) and several portions of methanol. The product was eluted by treating the resin with a 20% solution of formic acid in methanol (5 mL) and washing with methanol (3 mL). The combined eluant and final wash was evaporated to provide the title compound (154 mg, 99%). LCMS (ESMS): m/z 447/9 ($M+H^+$).

Using methods similar to those described in the above example, the following analogs were also synthesized:


3-[3-(3,5-Dichloro-benzyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid. LCMS (ESMS): m/z 407 ($M+H^+$).




3-[3-(5-Bromo-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid. LCMS (ESMS): m/z 473/5 ($M+H^+$).




3-[3-(1*H*-Indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid. LCMS (ESMS): m/z 378 ($M+H^+$).




-[3-(1*H*-Indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-propionic acid. LCMS (ESMS): m/z 336 ($M+H^+$).




3-{3-[(5-bromo-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}propanoic acid. LCMS (ESMS): m/z 431,433 ($M+H^+$).




3-{3-[2-methyl-5-(trifluoromethyl)benzyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}hexanoic acid. LCMS (ESMS): m/z 421.0 ($M+H^+$).




3-[3-(2,3-dichlorobenzyl)-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl]hexanoic acid. . LCMS (ESMS): m/z 407.0 ($M+H^+$).



3-[3-(2,5-dichlorobenzyl)-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl]hexanoic acid. LCMS (ESMS): m/z 407.0 ($M+H^+$).



Example 32: 3-[3-(4-Chloro-1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-propionic acid




3-(2-Oxo-2,3-dihydro-benzoimidazol-1-yl)-hexanoic acid ethyl ester (80 mg, 0.29 mmol), prepared as described above, 4-chloro-3-hydroxymethyl-indole-1-carboxylic acid *tert*-butyl ester (282 mg, 1 mmol) and triphenylphosphene (223 mg, 0.85 mmol) were combined in a reaction vial and THF (1 mL) was added. The mixture was stirred until complete solution was attained, then cooled to 0°C, and diisopropylazodicarboxylate (0.165 mL, 0.85 mmol) was added dropwise. The reaction mixture was heated in the microwave at 130°C for 20 minutes, cooled, then diluted with THF (1 mL) and treated directly with the Amberlyst® resin (2.2 g, 2.95 mmol). This reaction mixture was agitated overnight and the resin was then filtered and washed with DMF (2 x 3 mL) and several portions of methanol. The product was eluted by treating the resin with a 20% solution of formic acid in methanol (2x3 mL) and washing with methanol (3 mL). The combined eluant and final wash was evaporated and redissolved in dichloromethane (1 mL). To this solution was added TFA (0.2 mL) and the reaction mixture was agitated for 1 h. The solvent was evaporated and the residue was purified by preparative HPLC using an acetonitrile/water/formic acid gradient providing the title compound (8 mg, 7%) LCMS (ESMS): m/z 412 ($M+H^+$).

Using methods similar to those described in the above example, the following analogs were also synthesized:


3-[3-(4-Bromo-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid.

LCMS (ESMS): m/z 456.8 ($M+H^+$).




3-[3-(4-Chloro-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid.

LCMS (ESMS): m/z 370 ($M+H^+$).



Example 33: 3-[3-(1-benzothien-7-ylmethyl)-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl]propanoic acid



3-(3-Isopropenyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid methyl ester:

To a stirred solution of 1-Isopropenyl-1,3-dihydro-benzimidazol-2-one (5g, 28.7mmol) in DMF (20ml) was added methyl acrylate (2.8ml, 31.6mmol) followed by benzyltrimethylammonium hydroxide (Triton B, 40 wt% in MeOH, 3.2ml). The resulting solution was stirred at ambient temperature until the reaction was complete. After this time the reaction was poured into water

and ethyl acetate. The layers were separated and the aqueous phase was extracted twice more with ethyl acetate. The combined organic layers were dried (MgSO_4), filtered and concentrated to give the crude product which was purified *via* column chromatography (silica gel, 30% ethyl acetate/hexanes). The product-containing fractions were combined and concentrated to give 4.8g (64%) of the desired product.

3-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid methyl ester:


To a stirred solution of 3-(3-Isopropenyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid methyl ester (4.8g, 18.4mmol) in ethanol (30ml) and water (30ml) was added 37% hydrochloric acid (24ml). The resulting solution was warmed to 60°C for twenty minutes and cooled. The reaction was diluted with dichloromethane and water. The layers were separated and the organic phase was dried (Na_2SO_4), filtered and concentrated to give 3.5g (86%) of the desired product as an oil which solidified on standing. Used without further purification.

3-[3-(1-benzothien-7-ylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid:

To a solution of 3-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid methyl ester (200mg, 0.91mmol) in DMA (5ml) was added sodium hydride (60% oil dispersion, 55mg, 1.4mmol). The resulting mixture was stirred for 10 minutes at ambient temperature after which time 7-bromomethyl-benzo[*b*]thiophene (206mg, 0.91mmol) was added. The reaction was heated at 120°C in a microwave for 20 minutes. Upon cooling to ambient temperature the reaction was diluted with ethyl acetate and washed with NaHCO_3 (sat.) and brine. The aqueous phase was acidified using 1M HCl and extracted once with ethyl acetate. The organic phase was dried (MgSO_4), filtered and concentrated to give the crude product which was purified *via* column chromatography (silica gel, 5% methanol/dichloromethane). The product-containing fractions were combined and concentrated to give 20mg (6%) of the desired product. LCMS (ESMS): m/z 353.41 ($\text{M}+\text{H}^+$).


The following compounds were synthesized using a similar procedure:

3-{3-[(1-methyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS): m/z 350.38 ($\text{M}+\text{H}^+$).




3-[3-(1,2-benzisothiazol-3-ylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid.

LCMS (ESMS): m/z 354.40 ($M+H^+$).




3-{3-[(6-methoxy-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid.


LCMS (ESMS): m/z 377.41 ($M+H^+$).




3-{3-[(1-methyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS): m/z 350.38 ($M+H^+$).



3-{3-[(1,3-dimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS): m/z 364.41 ($M+H^+$).



Example 34: 3-{3-[(6-hydroxy-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid



3-{3-[6-(*tert*-Butyl-dimethyl-silanyloxy)-naphthalen-1-ylmethyl]-2-oxo-2,3-dihydro-benzimidazol-1-yl}-propionic acid methyl ester:

To a stirred solution of 3-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid methyl ester (180mg, 0.82mmol) in THF (12ml) was added [6-(*tert*-butyl-dimethyl-silanyloxy)-naphthalen-1-yl]-methanol (283mg, 0.98mmol) and triphenylphosphine (257mg, 0.98mmol). The DIAD (0.20ml, 0.98mmol) was then introduced in a dropwise manner and the resulting mixture was stirred at ambient temperature over night. Silica gel was added and the reaction mixture was concentrated to dryness. The resulting solid was purified *via* column chromatography (silica gel, 40% ethyl acetate/hexanes) to give 240mg (60%) of the desired product as an oil which solidified on standing.


3-{3-[(6-hydroxy-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid:

To a stirred solution of 3-{3-[6-(*tert*-Butyl-dimethyl-silanyloxy)-naphthalen-1-ylmethyl]-2-oxo-2,3-dihydro-benzimidazol-1-yl}-propionic acid methyl ester (200mg, 0.41mmol) in THF (7ml) and water (1ml) was added lithium hydroxide monohydrate (26mg, 0.61mmol) at ambient temperature. Upon complete consumption of the starting material the reaction was diluted with water, ethyl acetate and 6N HCl (adjusted to pH=2). The layers were separated and the aqueous phase was extracted once with ethyl acetate. The combined organics were dried (MgSO₄), filtered and concentrated. The remaining residue was diluted with acetonitrile (3ml) and allowed to stand over night. The resulting precipitated product was collected *via* filtration, washed with


acetonitrile and dried on house vacuum to give 15mg (10%) of the desired product. LCMS (ESMS): m/z 363.38 ($M+H^+$).

The following compounds were synthesized using a similar procedure:


3-{2-oxo-3-[(1,2,3-trimethyl-1H-indol-7-yl)methyl]-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS): m/z 378.44 ($M+H^+$).




3-{3-[(8-bromo-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS): m/z 427.28 ($M+2H^+$).




3-{3-[(4-methylpyrazolo[1,5-a]pyridin-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS): m/z 351.37 ($M+H^+$).




3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS): m/z 412.88 ($M+H^+$).



3-{3-[(5-methoxy-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS): m/z 408.30 ($M+H^+$).



Example 35: 3-[3-(1,2-benzisoxazol-3-ylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid



1-[2-(2-Fluoro-phenyl)-2-oxo-ethyl]-3-isopropenyl-1,3-dihydro-benzimidazol-2-one:

To a stirred solution of 1-isopropenyl-1,3-dihydro-benzimidazol-2-one (1g, 5.7mmol) in DMF (50ml) was added potassium carbonate (1.6g, 11.5mmol) followed by 2-fluorophenacyl bromide (1.5g, 6.9mmol). The resulting mixture was warmed to 80°C for 2h and cooled to ambient

temperature. The reaction was poured into water and ethyl acetate and the layers were separated. The aqueous phase was extracted with ethyl acetate (2x) and the combined organics were dried (MgSO₄). Filtration and concentration gave the crude product which was purified *via* column chromatography (silica gel, 50% EtOAc/hexanes). The product-containing fractions were combined and concentrated to give 1.6g (90%) of the desired product.

1-{2-(2-Fluoro-phenyl)-2-[hydroxyimino]-ethyl}-3-isopropenyl-1,3-dihydro-benzimidazol-2-one:

To a stirred solution of 1-[2-(2-fluoro-phenyl)-2-oxo-ethyl]-3-isopropenyl-1,3-dihydro-benzimidazol-2-one (1.3g, 4.2mmol) in ethanol (50ml) was added hydroxylamine hydrochloride (0.84g, 12.2mmol) followed by pyridine (1.7ml, 21.0mmol) at ambient temperature. The resulting mixture was warmed to 80°C for 6h, cooled to ambient temperature and concentrated. The remaining residue was diluted with water, 1M HCl and ethyl acetate. The layers were separated and the aqueous phase was extracted once more with ethyl acetate. The combined organics were dried (MgSO₄), filtered and concentrated to give the crude product which was purified *via* column chromatography (silica gel, 50% EtOAc/hexanes). The product-containing fractions were combined and concentrated to give 1.0g (73%) of the desired product as an oil which solidified on standing.

1-1,2-Benzisoxazol-3-ylmethyl-3-isopropenyl-1,3-dihydro-benzimidazol-2-one:

A 25ml pear-shaped flask was charged with sodium hydride (60% oil dispersion, 68mg, 1.7mmol) which was washed with hexanes and then diluted with DME (3ml). The 1-{2-(2-fluoro-phenyl)-2-[hydroxyimino]-ethyl}-3-isopropenyl-1,3-dihydro-benzimidazol-2-one (0.5g, 1.5mmol) was introduced as a DME (3ml) solution in a dropwise manner. The resulting mixture was warmed to 85°C for 20h and cooled to ambient temperature. The reaction was diluted with water and the product was extracted into ethyl acetate (2x). The combined organics were dried (MgSO₄), filtered and concentrated to give the crude product which was purified *via* column chromatography (silica gel, 40% EtOAc/hexanes). The product-containing fractions were combined and concentrated to give 0.26g (55%) of the desired product.

1-1,2-Benzisoxazol-3-ylmethyl-1,3-dihydro-benzimidazol-2-one:

The 1-1,2-benzisoxazol-3-ylmethyl-3-isopropenyl-1,3-dihydro-benzimidazol-2-one (0.26g, 0.85mmol) was diluted with ethanol (2ml), water (2ml) and 37% HCl (1ml). The resulting mixture was warmed to 60°C for 1h and cooled to ambient temperature. The reaction was diluted with water and the product was extracted into dichloromethane. The layers were separated and the organic phase was dried (MgSO₄). Filtration and concentration gave 0.20g (89%) of the desired product which was used without further purification.


3-(3-1,2-Benzisoxazol-3-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-hexanoic acid ethyl ester:

To a stirred solution of 1-1,2-benzisoxazol-3-ylmethyl-1,3-dihydro-benzimidazol-2-one (100mg, 0.38mmol) in DMF (5ml) was added ethyl trans-2-hexenoate (80mg, 0.57mmol) followed by benzyltrimethylammonium hydroxide (Triton B, 40 wt% in MeOH, 17.0ul). The resulting mixture was warmed to 60°C for 12h and cooled to ambient temperature. The reaction was diluted with NH₄Cl (sat.) and the product was extracted into ethyl acetate (2x). The combined organics were dried (MgSO₄), filtered and concentrated to give the crude product which was purified *via* column chromatography (silica gel, 5% EtOAc/dichloromethane). The product-containing fractions were combined and concentrated to give 70mg (46%) of the desired product.

3-[3-(1,2-benzisoxazol-3-ylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid:

To a stirred solution of 3-(3-1,2-benzisoxazol-3-ylmethyl-2-oxo-2,3-dihydro-benzimidazol-1-yl)-hexanoic acid ethyl ester (70mg, 0.17mmol) in THF (5ml) and water (0.5ml) was added lithium hydroxide monohydrate (22mg, 0.52mmol). Upon complete consumption of the starting material, the reaction was concentrated to low volume and diluted with water (20ml) and acetic acid (0.5ml). The resulting heterogeneous mixture was stirred for 1h after which time the precipitated product was collected *via* filtration. The solid was washed with water and dried on house vacuum at 40°C (18h, over P₂O₅) to give 52mg (80%) of the desired product. LCMS (ESMS): m/z 380.41 (M+H⁺).

Example 36: 3-{3-[(4-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid




3-[3-(4-Chloro-1,2,3-trimethyl-1*H*-indol-7-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid methyl ester:

To a stirred solution of 3-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid methyl ester (50mg, 0.23mmol) in DMF (15ml) was added (4-chloro-1,2,3-trimethyl-1*H*-indol-7-ylmethyl)-trimethyl-ammonium; iodide (107mg, 0.27mmol) followed by potassium carbonate (94mg, 0.68mmol). The resulting mixture was warmed at 100°C over night and cooled to ambient temperature. The reaction was diluted with water and the resulting solids were removed *via* filtration. The filtrate was extracted with ethyl acetate (3x). The combined organics were dried (MgSO₄), filtered and concentrated to give the crude product which was purified *via* column chromatography (silica gel, 80% EtOAc/hexanes). The product-containing fractions were combined and concentrated to give 40mg (41%) of the desired product.

3-{3-[(4-chloro-1,2,3-trimethyl-1*H*-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}propanoic acid:

To a stirred solution of 3-[3-(4-chloro-1,2,3-trimethyl-1*H*-indol-7-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid methyl ester (40mg, 0.09mmol) in THF (3ml) and water (1ml) was added lithium hydroxide monohydrate (10mg, 0.24mmol). The resulting mixture was stirred at ambient temperature until all of the starting material was consumed. After this time the reaction was concentrated to low volume and diluted with water (10ml) and acetic acid (0.5ml). The resulting heterogeneous mixture was stirred at ambient temperature for 18h. The solid product was collected *via* filtration and suspended in acetonitrile. The product was collected once more *via* filtration and washed with minimal acetonitrile to give 5mg (13%) of the desired product. LCMS (ESMS): *m/z* 412.88 (M+H⁺).

Example 37: 3-{3-[(4-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid




3-[3-(4-Chloro-1,2,3-trimethyl-1*H*-indol-7-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester:

To a stirred solution of 3-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-hexanoic acid ethyl ester (93mg, 0.34mmol) in DMF (10ml) was added (4-chloro-1,2,3-trimethyl-1*H*-indol-7-ylmethyl)-trimethyl-ammonium; iodide (132mg, 0.34mmol) followed by potassium carbonate (140mg, 1.01mmol). The resulting mixture was warmed at 100°C over night and cooled to ambient temperature. The reaction was diluted with water and ethyl acetate. The layers were separated and the organic phase was dried (MgSO₄), filtered and concentrated to give the crude product. Purification *via* preparative TLC (1mm silica gel, 20% EtOAc/hexanes) gave 25mg (15%) of the desired product.

3-{3-[(4-chloro-1,2,3-trimethyl-1*H*-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}hexanoic acid:

To a stirred solution of 3-[3-(4-Chloro-1,2,3-trimethyl-1*H*-indol-7-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester (25mg, 0.05mmol) in THF (5ml) and water (1ml) was added lithium hydroxide monohydrate (20mg, 0.48mmol). The resulting mixture was stirred at ambient temperature until all of the starting material was consumed. After this time the reaction was concentrated to low volume and diluted with water (5ml) and acetic acid (0.2ml). The resulting solid was collected *via* filtration and suspended in acetonitrile/water. The solid was collected once more *via* filtration and dried on house vacuum at 70°C to give 10mg (43%) of the desired product. LCMS (ESMS): *m/z* 454.96 (M+H⁺).


Example 38: 3-{3-[(1,3-dimethyl-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid



3-{3-[(1,3-dimethyl-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid:

A mixture of 3-[3-(7-Bromo-1,3-dimethyl-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid (prepared using similar conditions as described in example 2, 65mg, 0.15mmol) and 10% Pd/C (50% wet, Degussa Type, 12mg) in methanol (8ml) was degassed and placed under hydrogen atmosphere (balloon). After stirring at ambient temperature over night, LCMS showed desired product as well as the methyl ester of the desired product. The catalyst was removed *via* filtration and the filtrate was concentrated to give the crude product. Purification *via* column chromatography (silica gel, 20-100% EtOAc/hexanes) gave the methyl ester which was dissolved into THF (2.5ml) and water (0.5ml) and treated with lithium hydroxide monohydrate (8mg). Upon complete hydrolysis of the ester, the reaction was concentrated to low volume and the remaining residue was diluted with water and acetic acid. The resulting precipitate was collected *via* filtration, washed with water and dried on house vacuum to give 18mg (34%) of the desired product. LCMS (ESMS): m/z 364.41 ($M+H^+$).

Example 39: 3-[2-Oxo-3-(1,2,3-trimethyl-1H-indol-7-ylmethyl)-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid



3-(3-Isopropenyl-2-oxo-2,3-dihydro-benzoimidazol-1-yl)-hexanoic acid ethyl ester:

To a solution of 1-isopropenyl-2-benzimidazolinone (2 g, 11.5 mmol) in DMF (10 mL) were added ethyl trans-2-hexenoate (2 g, 14 mmol) and the benzyltrimethylammonium hydroxide (40% in MeOH) (0.4 mL, 2.3 mmol) at room temperature. The solution was heated to 60 °C for 12 hours. The solution was cooled down and sat. NH_4Cl sol was added. The solution was extracted with EtOAc and the combined organic layer was dried with MgSO_4 and filtered. The filtrate was concentrated and the residue was purified by CombiFlash with 20% EtOAc in Hexane to afford the desirable product 3-(3-Isopropenyl-2-oxo-2,3-dihydro-benzoimidazol-1-yl)-hexanoic acid ethyl ester (2 g, 55%) as a colorless oil.

3-(2-Oxo-2,3-dihydro-benzoimidazol-1-yl)-hexanoic acid ethyl ester:

3-(3-Isopropenyl-2-oxo-2,3-dihydro-benzoimidazol-1-yl)-hexanoic acid ethyl ester (1 g, 3.2 mmol) was dissolved in MeOH (30 mL) and HCl (30 mL) solution at room temperature. The solution was heated to 60 °C for 2 hours. The solution was cooled down and was extracted with EtOAc. The combined organic layer was dried with MgSO_4 and was filtered. The filtrate was concentrated and the residue was purified by CombiFlash with 5% MeOH in CH_2Cl_2 as the eluent to afford the desirable product 3-(2-Oxo-2,3-dihydro-benzoimidazol-1-yl)-hexanoic acid ethyl ester (700 mg, 80%) as a colorless oil.

3-[2-Oxo-3-(1,2,3-trimethyl-1*H*-indol-7-ylmethyl)-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid ethyl ester:


To a solution of 3-(2-Oxo-2,3-dihydro-benzoimidazol-1-yl)-hexanoic acid ethyl ester (100 mg, 0.36 mmol) in THF (10 mL) were added (1,2,3-Trimethyl-1*H*-indol-7-yl)-methanol (103 mg, 0.54 mmol), DIAD (diisopropyl azo-dicarboxylate) (0.14 mL, 0.7 mmol) and triphenylphosphine (95 mg, 0.36 mmol) at room temperature. The solution was stirred at the same temperature for 12 hours. The solution was filtered and the filtrate was concentrated. The residue was purified by combiFlash with 10% EtOAc in Hexane as the eluent to afford the desirable product 3-[2-Oxo-3-(1,2,3-trimethyl-1*H*-indol-7-ylmethyl)-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid ethyl ester (80 mg, 49%) as a white foam.

3-[2-Oxo-3-(1,2,3-trimethyl-1*H*-indol-7-ylmethyl)-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid:


To a solution of 3-[2-Oxo-3-(1,2,3-trimethyl-1*H*-indol-7-ylmethyl)-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid ethyl ester (50 mg, 0.11 mmol) in 1,4-dioxane (5 mL) and H₂O (2 mL) was added LiOH (5.4 mg, 0.22 mmol). The solution was stirred at room temperature for 5 hours. The solution was concentrated and the residue was purified by CombiFlash with 10% MeOH in CH₂Cl₂ as the eluent to afford the desirable product 3-[2-Oxo-3-(1,2,3-trimethyl-1*H*-indol-7-ylmethyl)-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid (14.5 mg, 31%) as a white foam. LC-MS ($M^+ + 1$): 420.21.

The following compound was prepared using a similar procedure.


3-[3-(8-Bromo-naphthalen-1-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid.
LC-MS ($M^+ + 1$): 469.04




3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid. LC-MS ($M^+ + 1$): 454.9.




3-{3-[(5-fluoro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid. LC-MS ($M^+ + 1$): 438.5.




3-{3-[(5-chloro-1,3-dimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid. LC-MS ($M^+ + 1$): 440.25




2-({3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)pentanoic acid. LC-MS ($M^+ + 1$): 454.26




3-{3-[(5-chloro-1-methyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid. LC-MS ($M^+ + 1$): 426.16




2-({3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)butanoic acid. LC-MS ($M^+ + 1$): 440.02




3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2-methylpropanoic acid. LC-MS ($M^+ + 1$): 426.16



3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2-phenylpropanoic acid. LC-MS ($M^+ + 1$): 488.17



Example 40: 7-[3-(2-Carboxy-ethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl]-1,2,3-trimethyl-1H-indole-6-carboxylic acid methyl ester



2-(3-Isopropenyl-2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl)-3-nitro-benzoic acid methyl ester:

To a solution of 1-isopropenyl-2-benzimidazolidinone (200 mg, 1.1 mmol) in DMF (10 mL) were added methyl-2-bromomethyl-3-nitrobenzoate (472 mg, 1.7 mmol) and K_2CO_3 (317 mg, 2.3 mmol) at room temperature. The solution was heated to 80 C for 2 hours. The solution was cooled down and sat. NH_4Cl solution was added. The solution was extracted with EtOAc. The combined organic layer was dried with $MgSO_4$ and was filtered. The filtrate was concentrated and the residue was purified by CombiFlash with 20% EtOAc in Hexane as the eluent to afford the desirable product 2-(3-Isopropenyl-2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl)-3-nitro-benzoic acid methyl ester (395 mg, 94%) as a pale yellow foam.

7-(3-Isopropenyl-2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl)-2,3-dimethyl-1*H*-indole-6-carboxylic acid methyl ester:

To a stirred solution of 2-(3-Isopropenyl-2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl)-3-nitro-benzoic acid methyl ester (1 g, 2.7 mmol) in THF was added 1-methyl-1-propenylmagnesium bromide solution (0.5M in THF) (16 mL, 8.6 mmol) at $-50^\circ C$ (rapid dropwise addition) under nitrogen. Upon complete addition the reaction was stirred for 45 minutes ($-50^\circ C$) and quenched (NH_4Cl , sat.) and warmed to ambient temperature. Diluted with ether and separated the layers. The organic phase was washed with NH_4Cl (Sat., 2x). The organic phase was dried ($MgSO_4$), filtered and concentrated to give the crude product which was purified *via* CombiFlash with

10% EtOAc in hexane as the eluent to afford the desirable product 7-(3-Isopropenyl-2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl)-2,3-dimethyl-1*H*-indole-6-carboxylic acid methyl ester (350 mg, 33%) as a colorless oil.

7-(3-Isopropenyl-2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl)-1,2,3-trimethyl-1*H*-indole-6-carboxylic acid methyl ester:

To a solution of 7-(3-Isopropenyl-2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl)-2,3-dimethyl-1*H*-indole-6-carboxylic acid methyl ester (107 mg, 0.3 mmol) in THF (10 mL) was added NaH (60% in mineral oil) (22 mg, 0.55 mmol) at 0 C under nitrogen atmosphere. The solution was stirred at the same temperature for 15 minutes. MeI (0.034 mL, 0.55 mmol) was added and the solution was stirred at 0 C for 2 hour. Sat. NH₄Cl solution was added and the solution was extracted with EtOAc. The combined organic layer was dried with MgSO₄ and was filtered. The filtrate was concentrated and the residue was purified by CombiFlash with 10% EtOAc in Hexane as the eluent to afford the desirable product 7-(3-Isopropenyl-2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl)-1,2,3-trimethyl-1*H*-indole-6-carboxylic acid methyl ester (100 mg, 90%) as a colorless oil.

1,2,3-Trimethyl-7-(2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl)-1*H*-indole-6-carboxylic acid methyl ester:

7-(3-Isopropenyl-2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl)-1,2,3-trimethyl-1*H*-indole-6-carboxylic acid methyl ester (100 mg, 0.25 mmol) was dissolved in MeOH (10 mL) and HCl (5 mL) solution at room temperature. The solution was heated to 60 C for 2 hours. The solution was cooled down and was extracted with EtOAc. The combined organic layer was dried with MgSO₄ and was filtered. The filtrate was concentrated and the residue was purified by CombiFlash with 10% EtOAc in hexane as the eluent to afford the desirable product 1,2,3-Trimethyl-7-(2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl)-1*H*-indole-6-carboxylic acid methyl ester (85 mg, 94%) as a colorless oil.

7-[3-(2-Ethoxycarbonyl-ethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl]-1,2,3-trimethyl-1*H*-indole-6-carboxylic acid methyl ester:


To a stirred solution of 1,2,3-Trimethyl-7-(2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl)-1*H*-indole-6-carboxylic acid methyl ester (70 mg, 0.19 mmol) in DMF (10 mL) was added the methylacrylate (0.035 mL, 0.39 mmol) followed by the benzyltrimethylammonium hydroxide (40% in MeOH) (0.13 mL, 0.29 mmol). The resulting solution was stirred at ambient temperature for 2h after which time the reaction appeared complete. The reaction was poured into water and EtOAc. The layers were separated and the aqueous phase was extracted with EtOAc (2x). The combined organics were dried with MgSO₄ and was filtered, concentrated. The residue was purified by CombiFlash with 20% EtOAc in Hexane as the eluent to afford the product 7-[3-(2-Ethoxycarbonyl-ethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl]-1,2,3-trimethyl-1*H*-indole-6-carboxylic acid methyl ester (66 mg, 76%) as a colorless oil.

7-[3-(2-Carboxy-ethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl]-1,2,3-trimethyl-1*H*-indole-6-carboxylic acid methyl ester:


To a solution of 7-[3-(2-Ethoxycarbonyl-ethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl]-1,2,3-trimethyl-1*H*-indole-6-carboxylic acid methyl ester (60 mg, 0.13 mmol) in 1,4-dioxane (5 mL) and H₂O (2 mL) was added LiOH (6.4 mg). The solution was stirred at room temperature for 5 hours. The solution was concentrated and the residue was purified by CombiFlash with 10% MeOH in CH₂Cl₂ as the eluent to afford the desirable product 7-[3-(2-Carboxy-ethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl]-1,2,3-trimethyl-1*H*-indole-6-carboxylic acid methyl ester (10 mg, 17%) as a white solid. LC-MS (M⁺): 435.36.

The following compound was prepared using a similar procedure.

3-{2-oxo-3-[(1,2,3,5-tetramethyl-1*H*-indol-7-yl)methyl]-2,3-dihydro-1*H*-benzimidazol-1-yl}propanoic acid. LCMS (M+1): 392.46.



Example 41: (S)-3-[3-(4-Methyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid



(S)-3-[Benzyl-((S)-1-phenyl-ethyl)-amino]-hexanoic acid ethyl ester:

To a solution of (S)-(-)-N-benzyl- α -methyl-benzylamine (50 mL, 239 mmol) in THF (150 mL) at 0 C was added BuLi (2.5 M in THF) (100 mL, 250 mmol) under nitrogen atmosphere. The solution was cooled to -78 C and ethyl trans-2-hexenoate (16.6 g, 117 mmol) was added. The solution was stirred for 1 hour and Sat. NH_4Cl solution was added. The solution was warmed to room temperature and was extracted with EtOAc. The combined organic layer was dried with MgSO_4 and was filtered. The filtrate was concentrated and the residue was purified by CombiFlash with 5% EtOAc in Hexane as the eluent to afford the desirable product (S)-3-[Benzyl-((S)-1-phenyl-ethyl)-amino]-hexanoic acid ethyl ester (20.8 g, 50%) as a colorless oil.

(S)-3-Amino-hexanoic acid ethyl ester:

To a solution of (S)-3-[Benzyl-((S)-1-phenyl-ethyl)-amino]-hexanoic acid ethyl ester (20.8 g, 59 mmol) in EtOH (50 mL) was added $\text{Pd}(\text{OH})_2$ (20% in carbon) (2g). The mixture was stirred under hydrogen atmosphere with 40 psi for 16 hours. The mixture was filtered and the filtrate was concentrated. The residue (8.2 g, 87%) was used in the next step of the synthesis without further purification.

(S)-3-(2-Nitro-phenylamino)-hexanoic acid ethyl ester:

To a solution of 1-fluoro-2-nitrobenzene (500 mg, 3.5 mmol) in DMF (150 mL) were added (S)-3-Amino-hexanoic acid ethyl ester (600 mg, 3.8 mmol) and diisopropyl ethylamine (0.65 mL, 3.8 mmol). The solution was heated to 90 C for 16 hours. The solution was cooled down and water (300 mL) was added. The solution was extracted with EtOAc and the combined organic layer was dried with MgSO₄. The mixture was filtered and the filtrate was concentrated. The residue was purified by CombiFlash with 5% EtOAc in Hexane as the eluent to afford the desirable product (S)-3-(2-Nitro-phenylamino)-hexanoic acid ethyl ester (795 mg, 80%) as a colorless oil.

(S)-3-(2-Amino-phenylamino)-hexanoic acid ethyl ester:

To a solution of (S)-3-(2-Nitro-phenylamino)-hexanoic acid ethyl ester (795 mg, 2.8 mmol) in EtOH (20 mL) was added Pd/C (10% wt on carbon) (300 mg). The solution was stirred at room temperature for 3 hours. The solution was filtered and the filtrate was concentrated. The residue (683 mg, 96%) was used in the next step of the synthesis without further purification.

(S)-3-(2-Oxo-2,3-dihydro-benzoimidazol-1-yl)-hexanoic acid ethyl ester:

To a solution of (S)-3-(2-Amino-phenylamino)-hexanoic acid ethyl ester (650 mg, 2.6 mmol) in THF (10 mL) was added carbonyl diimidazole [CDI] (842 mg, 5.2 mmol) at room temperature under nitrogen atmosphere. The solution was stirred at the same temperature for 12 hours. The solution was concentrated and the residue was purified by CombiFlash with 20% EtOAc in Hexane as the eluent to afford the desirable product (S)-3-(2-Oxo-2,3-dihydro-benzoimidazol-1-yl)-hexanoic acid ethyl ester (700 mg, 98%) as a colorless oil.

(S)-3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid ethyl ester:

To a solution of (S)-3-(2-Oxo-2,3-dihydro-benzoimidazol-1-yl)-hexanoic acid ethyl ester (700 mg, 2.5 mmol) in DMF (50 mL) were added (1,4-Dimethyl-1*H*-indol-3-ylmethyl)-trimethylamine iodide salt (1.3 g, 3.8 mmol) and K₂CO₃ (700 mg, 5.1 mmol) at room temperature under nitrogen atmosphere. The solution was heated to 100 C for 4 hours. The solution was cooled down and water was added. The solution was extracted with EtOAc and the combined organic


layer was dried with MgSO_4 . The solution was filtered and the filtrate was concentrated. The residue was purified by CombiFlash with 20% EtOAc in Hexane as the eluent to afford the desirable product (S)-3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid ethyl ester (737 mg, 67%) as a yellow foam.

(S)-3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid:


To a solution of (S)-3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid ethyl ester (4.2 g, 9.8 mmol) in 1,4-dioxane (100 mL) and water (10 mL) was added LiOH (234 mg, 9.8 mmol) at room temperature. The solution was stirred at the same temperature for 12 hours. The solution was concentrated and water was added to the residue. The solution was acidified by 12N HCl in an ice-bath. The solid that precipitated out from the solution was collected by filtration. The solid was dried under vacuum and was confirmed to be the desirable product (S)-3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid (3.7 g, 92%). LC-MS ($\text{M}^+ + 1$): 406.27.

The following compound was prepared following the similar procedure using appropriate starting material.

(3S)-3-{2-oxo-3-[(1,4,6-trimethyl-1*H*-indol-3-yl)methyl]-2,3-dihydro-1*H*-benzimidazol-1-yl}hexanoic acid. LCMS: m/z 420.28 ($\text{M} + \text{H}^+$).



Example 42: 3-(3-naphtalen-2-ylmethyl-2-oxo-6-[(tetrahydro-furan-2-ylmethyl)carbamoyl]-2,3-dihydro-benzoimidazol-1-yl)propionic acid



3-Fluoro-4-nitro-benzoic acid tert-butyl ester:

A solution of 3-Fluoro-4-Nitro benzoic acid (2.00g, 10.8 mmol) in 50% DMF/CH₂Cl₂ (60 ml) was put on ice for 15 minutes. To the solution EDC (2.1g, 11 mmol) was added. The solution was stirred in the cold for another 15 minutes. t-butanol (3.7g, 50 mmol) was then added, followed by DMAP (100mg). The solution was stirred at room temperature overnight. Upon completion (TLC), the reaction mixture was diluted with ethyl acetate (300ml), washed with water then with 1N NaOH solution and finally with brine. The organic phase was evaporated to dryness. The residue was washed down a column eluted with 3% methanol in CH₂Cl₂. the pooled fractions were evaporated to dryness to give 2g (77%) of the desired product.

3-(2-Ethoxycarbonyl-ethylamino)-4-nitro-benzoic acid tert-butyl ester:

3-fluoro-4-nitro-benzoic acid tert-butyl ester (1g, 4.14mmol) and Beta-Alanine ethyl ester hydrochloride 1.037g, 6.75 mmol) were dissolved in DMF (10ml). To the solution DIEA 0.872 g, 6.75 mmol) was added and the mixture stirred at room temperature overnight. The reaction mixture was diluted with water (100 ml), extracted with Ethyl acetate

3x, dried over MgSO_4 , filtered and evaporated to dryness to afford 1.2g of solid was obtained (86%).

4-Amino-3-(2-Ethoxycarbonyl-ethylamino)-benzoic acid tert-butyl ester:

3-(2-Ethoxycarbonyl-ethylamino)-4-nitro-benzoic acid tert-butyl ester (1.2g, 3.55mmol) was dissolved in ethanol (60 ml) to the solution Pd/C (10%) (300mg) was added and the reaction mixture was stirred under Hydrogen (1 atmosphere) overnight. The reaction mixture was filtered through celite and evaporated to dryness to give 1g (91%) of product.

3-(2-Ethoxycarbonyl-ethyl)-2-oxo-2,3-dihydro-1-H-benzoimidazole-5-carboxylic acid tert-butyl ester:

To a solution of 4-Amino-3-(2-Ethoxycarbonyl-ethylamino)-benzoic acid tert-butyl ester (900mg, 2.92 mmol) in THF (20ml) was added CDI (997mg, 6mmol). The solution was heated at 90 degrees for 30 minutes. The reaction mixture was evaporated to dryness, and then taken up in Ethyl Acetate. The organic phase was washed with brine, dried over MgSO_4 , filtered, evaporated to dryness. The residue was purified by flash Chromatography eluted with 5% methanol in CH_2Cl_2 . After evaporation of the desired fractions, 800mg was obtained (82%).

3-(2-Ethoxycarbonyl-ethyl)-1-naphtalen-2-ylmethyl-2-oxo-2,3-dihydro-1-H-benzoimidazole-5-carboxylic acid tert-butyl ester:

3-(2-Ethoxycarbonyl-ethyl)-2-oxo-2,3-dihydro-1-H-benzoimidazole-5-carboxylic acid tert-butyl ester (800mg, 2.4 mmol) dissolved in DMF (15 ml) and put on ice for 15 minutes. To the solution NaH (87mg, 3.6 mmol) was added portionwise. The resulting mixture was allowed to stir for another 20 minutes. The mixture was put on warm water until evolution of gas stopped. The ice bath was reintroduced and chloromethyl naphthalene dissolved in DMF (5ml) was added slowly. The reaction mixture was then stirred at room temperature overnight. The reaction mixture was diluted with water, extracted (3x) with ethyl acetate, dried over MgSO_4 , filtered, and evaporated to dryness. After purification by flash chromatography 850 mg was obtained (75%).

3-(2-Ethoxycarbonyl-ethyl)-1-naphtalen-2-ylmethyl-2-oxo-2,3-dihydro-1-H-benzoimidazole-5-carboxylic acid:

3-(2-Ethoxycarbonyl-ethyl)-1-naphtalen-2-ylmethyl-2-oxo-2,3-dihydro-1-H-benzoimidazole-5-carboxylic acid tert-butyl ester (900mg, 1.9 mmol) was dissolved in 25% TFA in CH₂Cl₂ (100ml). The solution was allowed to stand at room temperature for 4 hours. After complete disappearance of the starting material the solution was evaporated to dryness. The residue was re-dissolved in CH₂Cl₂ (100 ml) and evaporated to dryness. This was done 4 times. The residue was dried overnight under vacuum. 690mg was obtained (87%).

3-{3-Naphtalen-2-ylmethyl-2-oxo-6-[(tetrahydro-furan-2-ylmethyl)-carbamoyl]-2'3-dihydro-benzoimidazol-1-yl}-propionic acid ethyl ester:


3-(2-Ethoxycarbonyl-ethyl)-1-naphtalen-2-ylmethyl-2-oxo-2,3-dihydro-1-H-benzoimidazole-5-carboxylic acid (100mg, 0.24mmol) was dissolved in DMF (3ml). the solution was then put on ice. To the cold solution EDC (48 mg, 0.250 mmol) and HOBt (34mg, 0.250 mmol) were added. After stirring the mixture on ice for 30 minutes, C-(tetrahydro-furan-2-yl)-methyl amine (25 mg, 0.250 mmol) was added. The reaction mixture was stirred at room temperature for 4 hours. TLC showed completion of the reaction. The reaction mixture was diluted with aqueous solution of Sodium bicarbonate, extracted with Ethyl acetate, washed with brine, dried over MgSO₄, filtered, and evaporated to dryness to give 106 mg (88 %).

3-(3-naphtalen-2-ylmethyl-2-oxo-6-[(tetrahydro-furan-2-ylmethyl)carbamoyl]-2,3-dihydro-benzoimidazol-1-yl)propionic acid:


3-{3-Naphtalen-2-ylmethyl-2-oxo-6-[(tetrahydro-furan-2-ylmethyl)-carbamoyl]-2'3-dihydro-benzoimidazol-1-yl}-propionic acid ethyl ester (100mg, 0.2mmol) was dissolved in methanol (2ml). To the solution aqueous solution of sodium Hydroxide was added. The mixture was stirred at room temperature for 2 hours and evaporated to dryness. The reaction mixture was then taken up in water (10ml) acidified with 1 N HCl, extracted with Ethyl acetate, washed with brine, dried over MgSO₄, filtered, evaporated to dryness. The residue was purified by flash chromatography eluted 5% methanol in CH₂Cl₂ to give 60 mg (60%) of the desired product. LCMS: m/z 473.52 (M+H⁺).

The following compounds were synthesized using a similar procedure:


3-{6-[(2-hydroxyethyl)carbamoyl]-3-[(4-methyl-1-benzothien-3-ylmethyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS): 453.51 ($M+H^+$).




3-{6-[(Benzyl)carbamoyl]-3-[(4-methyl-1-benzothien-3-ylmethyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS): 499.6 ($M+H^+$).




3-{6-{[2-(4-methoxyphenyl) ethyl] carbamoyl}-3-[(4-methyl-1-benzothien-3-ylmethyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS): 543.54 ($M+H^+$).




3-{6-{[2-(2-methoxyphenyl) ethyl] carbamoyl}-3-[(1-naphthyl methyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS): 523.54 ($M+H^+$).




3-{6-{[3-hydroxypyrrolidin-1-yl] carbamoyl}-3-[(1-naphthyl methyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]}propanoic acid. LCMS (ESMS): 459.54 ($M+H^+$).




3-{6-{[2-(3-methoxyphenyl) ethyl] carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]}propanoic acid. LCMS (ESMS): 543.54 ($M+H^+$).




3-{6-{[2-(3-hydroxy-4-methoxyphenyl) ethyl] carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]}propanoic acid. LCMS (ESMS): 559.64 ($M+H^+$).




3-{6-{[2-(2-methoxyphenyl) ethyl] carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS):543.64 ($M+H^+$).




3-{6-{[2-(3,4-dimethoxyphenyl) ethyl] carbamoyl}-3-[(1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS): 553.64 ($M+H^+$).




3-{6-{[2-(3-hydroxy-4-methoxyphenyl) ethyl] carbamoyl}-3-[(1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid. LCMS (ESMS):539.58 ($M+H^+$).



3- {6- {[2-(3,4-dimethoxyphenyl)ethyl] carbamoyl}-3-[(4-methyl-1-benzothien-3-ylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]} propanoic acid. LCMS (ESMS):573.64 (M+H⁺).



Example 43: 3-[3-(1,2-Dimethyl-1H-indol-7-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid:



1-(1,2-Dimethyl-1H-indol-7-ylmethyl)-3-isopropenyl-1,3-dihydro-benzimidazol-2-one:

To a solution of 1-isopropenylbenzimidazolone (220 mg, 1.3 mmol) in dry THF (10 mL) are added triphenylphosphine (397 mg, 1.5 mmol) and (1,2-Dimethyl-1H-indol-7-yl)-methanol (220 mg, 1.3 mmol). Then diisopropyl azodicarboxylate (0.31 mL, 1.5 mmol) is added drop wise into the above solution at room temperature. The mixture is stirred for 16 hrs and then the solvent is removed under vacuum. The residue is purified by flash column chromatography using 10-30% EtOAc in hexanes to give 200 mg (48%) of 1-(1,2-dimethyl-1H-indol-7-ylmethyl)-3-isopropenyl-1,3-dihydro-benzimidazol-2-one.

1-(1,2-Dimethyl-1H-indol-7-ylmethyl)-1,3-dihydro-benzimidazol-2-one:

(1,2-dimethyl-1H-indol-7-ylmethyl)-3-isopropenyl-1,3-dihydro-benzimidazol-2-one (200 mg, 0.60 mmol) was suspended in ethanol (0.5 ml) and water (0.5 ml) and treated with 37% HCl (0.2 ml). The reaction was warmed to 60°C for 1.5h, cooled to room temperature and concentrated to low volume. The remaining residue was dissolved into DCM and sequentially washed with water and NaHCO₃ (sat.). The organic phase was dried (MgSO₄), filtered and concentrated to give 170 mg (97%) of 1-(1,2-dimethyl-1H-indol-7-ylmethyl)-1,3-dihydro-benzimidazol-2-one which was used without further purification.

3-[3-(1,2-Dimethyl-1H-indol-7-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid methyl ester:


To a solution of 1-(1,2-dimethyl-1H-indol-7-ylmethyl)-1,3-dihydro-benzimidazol-2-one (170 mg, 0.58 mmol) in DMF (3.0 mL) are added methyl acrylate (0.06 mL, 0.6 mmol) and 40% benzyltrimethyl ammonium hydroxide in MeOH (0.06 mL, 0.15 mmol). The resulting mixture is stirred at room temperature for 18 h after which time water (20 mL) was added. The mixture is extracted with EtOAc (3 x 50 mL) and the organic layers are combined, dried over MgSO₄ and concentrated to give crude product. Purification by flash column chromatography using 20-50% EtOAc in Hexanes affords 90 mg (41%) of 3-[3-(1,2-dimethyl-1H-indol-7-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid methyl ester.

3-[3-(1,2-Dimethyl-1H-indol-7-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid:

To a stirred solution of 3-[3-(1,2-dimethyl-1H-indol-7-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid methyl ester (90 mg, 0.24 mmol) in THF (3 ml) and water

(0.5 ml) was added lithium hydroxide monohydrate (15 mg, 0.36 mmol). The reaction was stirred for 0.5h and concentrated to low volume. The remaining residue was diluted with water and acetic acid (0.3ml). The resulting suspension was stirred at ambient temperature over night. The precipitate was collected *via* filtration, washed with water and then suspended in 1:1 acetonitrile/water. The solid precipitate was isolated *via* filtration and dried to give 65 mg (87%) of 3-[3-(1,2-dimethyl-1H-indol-7-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid. LCMS (ESMS): m/z 364.41 (M+H).

Example 44: (2S)-2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid



(S)-2-(2-Nitro-phenylamino)-propionic acid *tert*-butyl ester.

To a solution of 2-fluoronitrobenzene (1.0 g, 7.09 mmol) in DMF (5 mL) were added H-ALA-OTBU HCl (2.6 g, 14.2 mmol) and DIPEA (2.5 mL, 14.2 mmol). The reaction mixture was stirred at room temperature over the weekend. The reaction mixture was then diluted with ethyl acetate and washed with water (x4). The organic phase was then dried over Na₂SO₄ and concentrated to afford 1.8 g (95%) of the desired product as orange oil. The resulting residue was used for the next reaction without further purification.

(S)-2-(2-Amino-phenylamino)-propionic acid *tert*-butyl ester.

To a solution of (S)-2-(2-Nitro-phenylamino)-propionic acid *tert*-butyl ester (1.6 g, 6.0 mmol) in MeOH (20 mL) was added a slurry of 10% Pd/C (dry) (300 mg). The reaction mixture was degassed using house vacuum aspirator and then saturated with H₂ gas in balloon. The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was then filtered through a celite and the filtrate was concentrated to afford 1.5 g (100%) of the desired product as a dark brown oily residue. This product was used for the next reaction without purification. (S)-2-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid *tert*-butyl ester.

To a solution of (S)-2-(2-Amino-phenylamino)-propionic acid *tert*-butyl ester (1.5 g, 6.3 mmol) in THF (10 mL) was added CDI (1.03 g, 6.3 mmol). The reaction mixture was stirred at room temperature for 1 h. When the reaction was completed, the reaction mixture was diluted with ethyl acetate and washed with water. The organic phase was dried over Na₂SO₄ and concentrated to afford 1.6 g of the desired product as an oily residue. The LCMS indicated only 40% purity; however, this product was used for the next reaction without further purification. LCMS (ESMS): m/z 263.38 (M+H⁺).

(S)-2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid *tert*-butyl ester.

To a solution of (S)-2-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid *tert*-butyl ester (0.3 g, 0.46 mmol), (4-Methyl-benzo[*b*]thiophen-3-yl)-methanol (0.12 g, 0.69 mmol), and triphenylphosphine (0.14 g, 0.55 mmol) was added dropwise DIAD (0.11 mL, 0.55 mmol) at 0 °C. The mixture was slowly warmed to room temperature and stirred for 2 h. The reaction mixture was diluted with ethyl acetate and washed with water (x2). The organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC using 4:1 Hexanes:ethyl acetate as an eluent to afford 0.12 g (62%) of the desired product as a brownish oil. This product was not pure (~70% pure by ¹H-NMR), but was used for the next reaction without further purification.


(2S)-2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid:

To a solution of (S)-2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid *tert*-butyl ester (0.12 g, 0.28 mmol) in CH₂Cl₂ (5mL) was


added TFA (2 mL). The reaction mixture was stirred at room temperature for 3 h. When the reaction was complete, the reaction mixture was diluted with CH₂Cl₂ and washed with water. The organic phase was dried over Na₂SO₄ and concentrated. When a small amount of CH₂Cl₂ (~3 mL) was added to the resulting residue, a white solid came out of the solution. This solid was filtered and dried to afford 69 mg (66%) of the titled compound. LCMS (ESMS): m/z 367.23 (M+H⁺).

The following compounds were synthesized following a similar procedure.


(2S)-3-methyl-2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid. LCMS (ESMS): m/z 395.20 (M+H⁺)



(2S)-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}(phenyl)acetic acid. LCMS (ESMS): m/z 429.16 (M+H⁺)



Example 45. (2S)-2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid



(S)-2-(2-Nitro-phenylamino)-pentanoic acid *tert*-butyl ester.

To a solution of 2-fluoronitrobenzene (0.3 g, 2.1 mmol) in DMF (3 mL) were added (L)-NVA OTBU HCl (0.53 g, 2.6 mmol) and DIPEA (0.93 mL, 5.3 mmol). The reaction mixture was stirred at 70 °C overnight. The reaction mixture was then diluted with ethyl acetate and washed with water (x4). The organic phase was then dried over Na₂SO₄ and concentrated to afford 0.69 g (100%) of orange oil. The resulting residue was used for the next reaction without further purification.

(S)-2-(2-Amino-phenylamino)-pentanoic acid *tert*-butyl ester.

To a solution of (S)-2-(2-Nitro-phenylamino)-pentanoic acid *tert*-butyl ester (0.69g, 2.4 mmol) in MeOH (10 mL) was added a slurry of Pd/C (0.2 g) in MeOH (10 mL). The reaction mixture was degassed using house vacuum. The flask was saturated with H₂ and stirred under H₂ balloon for 2h. When the reaction was completed, the mixture was filtered through a pad of celite and the filtrate was concentrated to afford 0.61 g (97%) of the desired product. The resulting residue was used for the next reaction without further purification.

(S)-2-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-pentanoic acid *tert*-butyl ester.

To a solution of (S)-2-(2-Amino-phenylamino)-pentanoic acid *tert*-butyl ester (0.61 g, 2.3 mmol) in THF (8 mL) was added CDI (0.41 g, 2.5 mmol). The reaction mixture was stirred at room temperature for 3 h. When the reaction was complete, the reaction mixture was concentrated. The resulting residue was purified by silica gel prep TLC using 95:5

CH₂Cl₂:MeOH as an eluent to afford 0.51 g (76%) of the desired product as a brown solid.

LCMS (ESMS): m/z 291.31 (M+H⁺).


(S)-2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid *tert*-butyl ester.

To a solution of (S)-2-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-pentanoic acid *tert*-butyl ester (0.1 g, 0.34 mmol) in DMF (2 mL) were added K₂CO₃ (0.1 g, 0.69 mmol) and trimethyl-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-ammonium iodide (0.24 g, 0.69 mmol). The reaction mixture was heated to 80 °C for 5 h. The reaction mixture was then diluted with ethyl acetate and washed with water (x4). The organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC using 98:2 CH₂Cl₂:MeOH as an eluent to afford 0.11 g (71%) of the desired product as an oily residue. LCMS (ESMS): m/z 451.33 (M+H⁺).


To a solution of (S)-2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid *tert*-butyl ester (0.11 g, 0.24 mmol) in CH₂Cl₂ (2 mL) was added TFA (1 mL). The reaction mixture was stirred at room temperature for 2 h. When the reaction was complete, the reaction mixture was washed with water (x2). The organic phase was dried over Na₂SO₄ and concentrated. The resulting residue was purified by silica gel prep TLC using 95:5 CH₂Cl₂:MeOH as an eluent to afford 81 mg (84%) of the desired product as an off-white solid. LCMS (ESMS): m/z 395.09 (M+H⁺).

The following compound were prepared following a similar procedure.

(2S)-2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid. LCMS (ESMS): m/z 381.09 (M+H⁺).



Example 46: 3-{5-anilino-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid



3-[3-1, 4-Dimethyl-1H-indol-3-ylmethyl)-2-oxo-5-phenylamino-2,3-dihydro-benzimidazol-1-yl]hexanoic acid Ethyl ester:

In a dry pressure bottle, 3-[5-bromo-3-(1, 4-dimethyl-1H-indol-3-ylmethyl) 2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester (200mg, 0.390mmol) was dissolved in 50%Toluene/NMP (10ml). To the solution, 2-amino thiazole (59 mg, 0.585mmol), Cesium carbonate (191 mg, 0.585 mmol), 2(dicyclohexylphosphino) biphenyl (14mg, 0.040mmol), tris (dibenzylideneacetone)-dipalladium (0) 99mg, 0.010mmol) were added. The bottle was then purged for 5 minutes by passing a slow stream of argon through the mixture while stirring. As the bottle was ready to be capped, the argon flow was increased. After capping, the reaction vessel was then immersed in an oil bath heated to 100 °C for 18 hours. An aliquot was removed and checked by TLC. The cooled reaction mixture was then diluted with Ethyl Acetate (200ml) and washed with water 3 times. The organic phase was again washed with 1N HCl (200ml), followed with brine. The collected organic phase was dried over MgSO₄, filtered and evaporated to dryness to give an oil, which was purified by flash Chromatography, eluted with 30% to 50% ethyl acetate in Hexanes in 30 minutes to give 205 mg (35%) of 3-[3-1, 4-Dimethyl-1H-indol-3-ylmethyl)-2-oxo-5-phenylamino-2,3-dihydro-benzimidazol-1-yl]hexanoic acid Ethyl ester. LCMS (ESMS): m/z 524.65 (M+H).


3-[3-1, 4-Dimethyl-1H-indol-3-ylmethyl)-2-oxo-5-phenylamino-2, 3-dihydro-benzimidazol-1-yl] hexanoic acid:

To a stirred solution of 3-[3-1, 4-Dimethyl-1H-indol-3-ylmethyl)-2-oxo-5-phenylamino-2,3-dihydro-benzimidazol-1-yl]hexanoic acid ethyl ester (50mg, 0.95mmol) in Methanol (5ml) was


added a concentrated solution of NaOH in water (2ml). The mixture was stirred at RT for 2 hours. The reaction mixture was checked by TLC. The reaction mixture was then evaporated to dryness. The residue was diluted with water (10ml) and acidified with 1N HCl (15ml). The aqueous mixture was extracted with Ethyl Acetate (100ml). The separated organic layer was dried over MgSO₄, filtered and evaporated to dryness. The residue obtained was purified by flash Chromatography eluted with 0 to 10% Methanol in Dichloro methane to give 20 mg (67.6%) of 3-[3-1, 4-Dimethyl-1H-indol-3-ylmethyl]-2-oxo-5-phenylamino-2,3-dihydro-benzimidazol-1-yl]hexanoic acid. LCMS (ESMS): m/z 496.60 (M+H).

The following compound was prepared following a similar procedure using appropriate starting materials.

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-5-(1,3-thiazol-2-ylamino)-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid. LCMS (ESMS): m/z 503.62 (M+H).



Example 47: (3S)-3-{5-(carbamoylamino)-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid



3-[5-Amino-3-(1,4-Dimethyl-1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]hexanoic acid ethyl ester.

3-[5-tert-Butoxycarbonylamino-3-(1,4-Dimethyl-1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]hexanoic acid: ethyl ester (350mg, 0.638mmol) was dissolved in Dichloromethane (5ml) was put on ice for 20 minutes. To the cold solution 4N Dioxane/HCl (10ml) was added and the mixture was stirred on ice for 2 hours. The solution was evaporated to dryness. The residue obtained was neutralized with an aqueous solution of Sodium Bicarbonate (25ml), then extracted with Ethyl acetate (200ml). The organic phase was evaporated to dryness and purified by Flash Chromatography, eluted with 10% methanol in dichloromethane to give 280 mg (97.9%) of 3-[5-Amino-3-(1,4-Dimethyl-1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]hexanoic acid ethyl ester. LCMS (ESMS): m/z 448.56 (M+H).

3-[3-(1,4-Dimethyl-1H-indol-3-ylmethyl)-2-oxo-5-ureido-2,3-dihydro-benzoimidazol-1-yl]hexanoic acid ethyl ester.


To a solution of 3-[5-Amino-3-(1,4-Dimethyl-1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]hexanoic acid: ethyl ester (50mg, 0.111mmol) in THF (30ml) was added TMS isocyanate (1.73 mg, 15mmol). The solution was stirred at RT for 3 hours. The reaction was checked by LCMS. To the reaction mixture an aqueous solution of Sodium bicarbonate (60ml) was added. The resulting mixture, stirred for another 30 minutes, was then extracted with

Ethyl Acetate. The organic layer, dried over MgSO_4 , was filtered and evaporated to dryness to give a dark residue. The residue was purified by prep TLC, eluted with 10% methanol in Dichloromethane to give 10mg (18%) of 3-[3-1, 4-Dimethyl-1H-indol-3-ylmethyl)-2-oxo-5-ureido-2, 3-dihydro-benzoimidazol-1-yl] hexanoic acid ethyl ester. LCMS (ESMS): m/z 491.58 (M+H).

3-[3-1, 4-Dimethyl-1H-indol-3-ylmethyl)-2-oxo-5-ureido-2, 3-dihydro-benzoimidazol-1-yl] hexanoic acid.


To a solution of 3-[3-1, 4-Dimethyl-1H-indol-3-ylmethyl)-2-oxo-5-ureido-2, 3-dihydro-benzoimidazol-1-yl] hexanoic acid ethyl ester (10mg, 0.020mmol) in methanol (2ml) was added a concentrated solution of Sodium Hydroxide in water (3ml). The mixture was stirred at RT for 2 hours. The reaction was checked by TLC. The reaction mixture was evaporated to dryness. The residue was taken in water (3ml), neutralized with 1N HCl (4ml) and extracted with Ethyl acetate (100ml). The organic layer was dried over MgSO_4 , filtered and evaporated to dryness to give a light beige residue. The residue was purified by mass triggered HPLC to give 4mg (42.4%) of 3-[3-1, 4-Dimethyl-1H-indol-3-ylmethyl)-2-oxo-5-ureido-2, 3-dihydro-benzoimidazol-1-yl] hexanoic acid. LCMS (ESMS): m/z 463.53 (M+H).

Example 48: Sodium (3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoate



To a mixture of (S)-3-[3-(1,4-dimethyl-1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid (1.45 g, 3.6 mmol) in H_2O (25 mL) was added NaOH (1.041 M in THF) (3.1 mL, 3.2 mmol). The solution was stirred at room temperature for 2 hours. The solution was washed with EtOAc (30 mL x2) and the aqueous layer was lyophilized to give the desired sodium (S)-3-[3-(1,4-dimethyl-1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoate (1.4 g, 91%) as a white solid. LCMS (ESMS): 428.10 (M+H⁺).

Example 49: 4-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid



4-(2-Amino-phenylamino)-pentanoic acid methyl ester.

To a mixture of 1,2-phenylenediamine (3.0 g, 27.7 mmol) and levulinic acid methyl ester (2.7 mL, 21.8 mmol) in 1,2-dichloroethane (25 mL) was added sodium triacetoxyborohydride (4.6 g, 21.7 mmol) and acetic acid (1.5 mL). The mixture was stirred at room temperature for 16 hours. The reaction mixture was diluted with EtOAc (100 mL) and washed with H₂O (75 mL x3). The organic layer was dried over sodium sulfate and concentrated. The resulting residue was purified by CombiFlash with 40% EtOAc in Hexane as the eluent to afford the desired 4-(2-amino-phenylamino)-pentanoic acid methyl ester (1.05 g, 22%).

4-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-pentanoic acid methyl ester.

To a solution of 4-(2-amino-phenylamino)-pentanoic acid methyl ester (1.05 g, 4.7 mmol) in THF (15 mL) was added 1,1'-carbonyldiimidazole (0.8 g, 4.9 mmol). The solution was stirred at room temperature for 16 hours. The reaction mixture was concentrated. The resulting residue was purified by CombiFlash with 50% EtOAc in Hexane as the eluent to afford the desired 4-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-pentanoic acid methyl ester (1.1 g, 94%).

4-[3-(1,4-Dimethyl-3*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid methyl ester.


To a mixture of 4-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-pentanoic acid methyl ester (85 mg, 0.34 mmol) and (1,4-dimethyl-1*H*-indol-3-ylmethyl)-trimethylammonium iodide (175 mg, 0.51 mmol) in DMF (2.5 mL) was added K₂CO₃ (95 mg, 0.69 mmol). The mixture was stirred at 100 °C for 4 hours. The reaction mixture was allowed to cool to room temperature, diluted with EtOAc (50 mL) and washed with H₂O (50 mL x3). The organic layer was dried over sodium sulfate and concentrated. The resulting residue was purified by CombiFlash with 25% EtOAc in Hexane as the eluent to afford the desired 4-[3-(1,4-dimethyl-3*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid methyl ester (120 mg, 86%).

4-{3-[(1,4-dimethyl-1*H*-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}pentanoic acid:


To a solution of 4-[3-(1,4-dimethyl-3*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentanoic acid methyl ester (120 mg, 0.3 mmol) in dioxane (2.5 ml) was added a solution of LiOH monohydrate (25 mg, 0.6 mmol) in H₂O (2.5 mL). The solution was stirred at room temperature for 4 hours. The reaction mixture was quenched with 4M HCl in dioxane (250 μ L) and concentrated. The resulting residue was purified by CombiFlash with 10% MeOH in dichloromethane as the eluent to afford the desired 4-{3-[(1,4-dimethyl-1*H*-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}pentanoic acid (17 mg, 15%). LCMS (ESMS): 392.26 (M+H⁺).

Following compounds were synthesized using a similar procedure.


4-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-heptanoic acid. LCMS (ESMS): 420.27 (M+H⁺).



3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2,2-dimethyl-butyrlic acid. LCMS (ESMS): 406.32 ($M+H^+$).



Example 50: {1-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-cyclohexyl}-acetic acid



[1-(2-Nitro-phenylamino)-cyclohexyl]-acetic acid methyl ester:

To a mixture of 1-fluoro-2-nitrobenzene (450 μ L, 4.3 mmol) and (1-amino-cyclohexyl)-acetic acid methyl ester (880 mg, 5.2 mmol) in DMF (5 mL) was added DIPEA (1.0 mL, 5.7 mmol). The mixture was heated at 90 °C for 16 hours. The reaction mixture was allowed to cool to room temperature, diluted with EtOAc (50 mL) and washed with H₂O (50 mL x3). The organic layer was dried over sodium sulfate and concentrated. The resulting residue was purified by CombiFlash with 20% EtOAc in Hexane as the eluent to afford the desired [1-(2-nitro-phenylamino)-cyclohexyl]-acetic acid methyl ester (230 mg, 19%).

[1-(2-Amino-phenylamino)-cyclohexyl]-acetic acid methyl ester:

A mixture of [1-(2-nitro-phenylamino)-cyclohexyl]-acetic acid methyl ester (230 mg, 0.79 mmol), 10%Pd/C (80 mg) in EtOH (20 mL) under H₂ atmosphere at room temperature was stirred for 16h. The reaction mixture was filtered and the organic layer was concentrated to give the desired [1-(2-amino-phenylamino)-cyclohexyl]-acetic acid methyl ester (205 mg, 99%).

[1-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-cyclohexyl]-acetic acid methyl ester:

A mixture of [1-(2-amino-phenylamino)-cyclohexyl]-acetic acid methyl ester (205 mg, 0.79 mmol) in THF (5 mL) was added 1,1'-carbonyldiimidazole (180 mg, 1.11 mmol). The solution was stirred at room temperature for 16 hours. The reaction mixture was concentrated. The resulting residue was purified by CombiFlash with 50% EtOAc in Hexane as the eluent to afford the desired [1-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-cyclohexyl]-acetic acid methyl ester (194 mg, 86%).

{1-[3-(1,4-Dimethyl-3*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-cyclohexyl}-acetic acid methyl ester:

To a mixture of [1-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-cyclohexyl]-acetic acid methyl ester (115 mg, 0.40 mmol) and (1,4-dimethyl-1*H*-indol-3-ylmethyl)-trimethylammonium iodide (200 mg, 0.58 mmol) in DMF (2.5 mL) was added K₂CO₃ (110 mg, 0.80 mmol). The mixture was stirred at 100 °C for 4 hours. The reaction mixture was diluted with EtOAc (50 mL) and washed with H₂O (50 mL x3). The organic layer was dried over sodium sulfate and concentrated. The resulting residue was purified by CombiFlash with 30% EtOAc in Hexane as the eluent to afford the desired {1-[3-(1,4-dimethyl-3*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-cyclohexyl}-acetic acid methyl ester (131 mg, 74%).


{1-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-cyclohexyl}-acetic acid:

To a solution of {1-[3-(1,4-dimethyl-3*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-cyclohexyl}-acetic acid methyl ester (131 mg, 0.29 mmol) in dioxane (2.5 ml) was added a solution of LiOH monohydrate (25 mg, 0.60 mmol) in H₂O (2.5 mL). The solution was stirred at room temperature for 4 hours. The reaction mixture was quenched with 4M HCl in dioxane (500 μ L) and concentrated. The resulting residue was purified by CombiFlash with 3% MeOH


in dichloromethane as the eluent to afford the desired {1-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-cyclohexyl}-acetic acid (85 mg, 68%). LCMS (ESMS): 432.32 ($M+H^+$).

Following compounds were synthesized using a similar procedure.


3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-3-methylbutyric acid. LCMS (ESMS): 392.26 ($M+H^+$)



{1-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-cyclopentyl}-acetic acid. LCMS (ESMS): 418.30 ($M+H^+$)



Example 51: 3-{3-[(1,4-dimethyl-1*H*-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}-4-methoxy-2,2-dimethylbutanoic acid



4-Methoxy-2,2-dimethyl-3-oxo-butyric acid methyl ester.

To a solution of methyl 4-methoxyacetoacetate (2.0 mL, 15.5 mmol) and iodomethane (3.0 mL, 48.1 mmol) in DMSO (20 mL) was added potassium carbonate (4.7 g, 34.0 mmol). The mixture was stirred at room temperature for 16 hours. The reaction mixture was diluted with EtOAc (150 mL) and washed with H₂O (150 mL x3). The organic layer was dried over sodium sulfate and concentrated to afford the desired 4-methoxy-2,2-dimethyl-3-oxo-butyric acid methyl ester (2.7 g, 100%).

3-(2-Amino-phenylamino)-4-methoxy-2,2-dimethyl-butyl-2-oxo-2,3-dihydro-benzimidazole-1-carboxylate.

To a mixture of 1,2-phenylenediamine (900 mg, 8.3 mmol) and 4-methoxy-2,2-dimethyl-3-oxo-butyric acid methyl ester (1.7 g, 9.8 mmol) in MeOH (5 mL) was added sodium cyanoborohydride (700 mg, 11.1 mmol) and acetic acid (1.0 mL). The mixture was stirred at room temperature for 16 hours. The reaction mixture was diluted with EtOAc (100 mL) and washed with H₂O (75 mL x3). The organic layer was dried over sodium sulfate and concentrated. The resulting residue was purified by CombiFlash with 40% EtOAc in Hexane as the eluent to afford the desired 3-(2-amino-phenylamino)-4-methoxy-2,2-dimethyl-butyl-2-oxo-2,3-dihydro-benzimidazole-1-carboxylate (550 mg, 25%).

4-Methoxy-2,2-dimethyl-3-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-butyl-2-oxo-2,3-dihydro-benzimidazole-1-carboxylate.

To a solution of 3-(2-amino-phenylamino)-4-methoxy-2,2-dimethyl-butyl-2-oxo-2,3-dihydro-benzimidazole-1-carboxylate (550 mg, 2.1 mmol) in THF (25 mL) was added 1,1'-carbonyldiimidazole (500 mg, 3.1 mmol).

The solution was stirred at room temperature for 16 hours. The reaction mixture was concentrated. The resulting residue was purified by CombiFlash with 75% EtOAc in Hexane as the eluent to afford the desired 4-methoxy-2,2-dimethyl-3-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-butyric acid methyl ester (420 mg, 70%).

3-[3-(1,4-Dimethyl-3*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-4-methoxy-2,2-dimethyl-butyric acid methyl ester.


To a mixture of 4-methoxy-2,2-dimethyl-3-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-butyric acid methyl ester (137 mg, 0.47 mmol) and (1,4-dimethyl-1*H*-indol-3-ylmethyl)-trimethylammonium iodide (225 mg, 0.65 mmol) in DMF (2.5 mL) was added K₂CO₃ (200 mg, 1.45 mmol). The mixture was stirred at 100 °C for 4 hours. The reaction mixture was allowed to cool to room temperature, diluted with EtOAc (50 mL) and washed with H₂O (50 mL x3). The organic layer was dried over sodium sulfate and concentrated. The resulting residue was purified by CombiFlash with 50% EtOAc in Hexane as the eluent to afford the desired 3-[3-(1,4-dimethyl-3*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-4-methoxy-2,2-dimethyl-butyric acid methyl ester (50 mg, 24%).

3-[3-(1,4-Dimethyl-3*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-4-methoxy-2,2-dimethyl-butyric acid.


To a solution of 3-[3-(1,4-dimethyl-3*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-4-methoxy-2,2-dimethyl-butyric acid methyl ester (50 mg, 0.11 mmol) in dioxane (2.5 mL) was added a solution of LiOH monohydrate (20 mg, 0.48 mmol) in H₂O (2.5 mL). The solution was stirred at 50 °C for 16 hours. The reaction mixture was allowed to cool to rt and quenched with 4M HCl in dioxane (500 μ L) and concentrated. The resulting residue was purified by CombiFlash with 3% MeOH in dichloromethane as the eluent to afford the desired 3-[3-(1,4-dimethyl-3*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-4-methoxy-2,2-dimethyl-butyric acid (15 mg, 31%). LCMS (ESMS): 436.12 (M+H⁺).

Following compounds were synthesized using a similar procedure.


1-{1-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-butyl}-cyclopropanecarboxylic acid. LCMS (ESMS): 432.32 (M+H⁺).



3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2,2-dimethyl-hexanoic acid. LCMS (ESMS): 434.30 (M+H⁺).



Example 52: 2-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2-methyl-propionic acid



2-Methyl-2-(2-nitro-phenylamino)-propionic acid *tert*-butyl ester:

To a mixture of 1-fluoro-2-nitrobenzene (485 μ L, 4.6 mmol) and 2-amino-2-methyl-propionic acid *tert*-butyl ester hydrochloride (1.0g, 5.1 mmol) in DMF (5 mL) was added Et₃N (1.3 mL, 9.3 mmol). The mixture was heated at 85 °C for 16 hours. The reaction mixture was allowed to cool to room temperature, diluted with EtOAc (50 mL) and washed with H₂O (50 mL x3). The

organic layer was dried over sodium sulfate and concentrated. The resulting residue was purified by CombiFlash with 10% EtOAc in Hexane as the eluent to afford the desired 2-methyl-2-(2-nitro-phenylamino)-propionic acid *tert*-butyl ester (640 mg, 50%).

2-(2-Amino-phenylamino)-2-methyl-propionic acid *tert*-butyl ester:

A mixture of 2-methyl-2-(2-nitro-phenylamino)-propionic acid *tert*-butyl ester (638 mg, 2.3 mmol), 10%Pd/C (110 mg) in EtOH (25 mL) under H₂ atmosphere at room temperature was stirred for 30 min. The reaction mixture was filtered and the organic layer was concentrated to give the desired 2-(2-amino-phenylamino)-2-methyl-propionic acid *tert*-butyl ester (570 mg, 100%).

2-Methyl-2-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid *tert*-butyl ester:

A mixture of 2-(2-amino-phenylamino)-2-methyl-propionic acid *tert*-butyl ester (570 mg, 2.3 mmol) in THF (25 mL) was added 1,1'-carbonyldiimidazole (560 mg, 3.5 mmol). The solution was stirred at room temperature for 16 hours. The reaction mixture was concentrated. The resulting residue was purified by CombiFlash with 50% EtOAc in Hexane as the eluent to afford the desired 2-methyl-2-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid *tert*-butyl ester (462 mg, 72%).


2-[3-(1,4-Dimethyl-3*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2-methyl-propionic acid *tert*-butyl ester:

To a mixture of 2-methyl-2-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid *tert*-butyl ester (130 mg, 0.47 mmol) and (1,4-dimethyl-1*H*-indol-3-ylmethyl)-trimethylammonium iodide (250 mg, 0.73 mmol) in DMF (2.5 mL) was added K₂CO₃ (200 mg, 1.45 mmol). The mixture was stirred at 100 °C for 4 hours. The mixture was diluted with EtOAc (50 mL) and washed with H₂O (50 mL x3). The organic layer was dried over sodium sulfate and concentrated. The resulting residue was purified by CombiFlash with 30% EtOAc in Hexane as the eluent to afford the desired 2-[3-(1,4-dimethyl-3*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2-methyl-propionic acid *tert*-butyl ester (138 mg, 68%).

2-[3-(1,4-Dimethyl-3*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2-methyl-propionic acid:

A solution of 2-[3-(1,4-dimethyl-3*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2-methyl-propionic acid *tert*-butyl ester (138 mg, 0.32 mmol) in TFA (1.0 mL) was stirred at room temperature for 1 hour. The reaction mixture was concentrated. The resulting residue was purified by CombiFlash with 10% MeOH in dichloromethane as the eluent to afford the desired 2-[3-(1,4-dimethyl-3*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2-methyl-propionic acid (36 mg, 30%). LCMS (ESMS): 378.25 ($M+H^+$).

Example 53: N-[(2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}ethyl)carbamoyl]benzenesulfonamide




1-(2-Bromo-ethyl)-3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one:
To the stirred solution of 1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one (100 mg, 0.34 mmol), 2-Bromo-ethanol (0.03 mL, 0.41 mmol) and triphenylphosphine (107 mg, 0.41 mmol) in THF (2.0 mL) is added DIAD (0.082 mL, 0.41 mmol) in a dropwise manner. The mixture is stirred at room temperature for 20 hr and then another 0.03 mL of 2-Bromo-ethanol, 107 mg of triphenylphosphine and 0.08 mL of DIAD are added. The reaction is continued for another 24 hr and then the solvent is removed under vacuum and the residue is purified by flash column chromatography to give 75 mg (55%) of 1-(2-Bromo-ethyl)-3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one.

1-(2-Amino-ethyl)-3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one:
 1-(2-Bromo-ethyl)-3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one
 (55 mg, 0.14 mmol) is dissolved in THF (1.5 mL) and NH_4OH (1.5 mL, 27% aqueous) is added
 into it at room temperature. The mixture is then stirred for 16 hr at 65 °C and another one week
 at room temperature. Then the solvents are removed under vacuum and the residue is washed
 with THF. The solid formed is filtered and rinsed with more THF to give 28 mg (61%) of 1-(2-
 Amino-ethyl)-3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one.

N-[(2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-
 yl}ethyl)carbamoyl]benzenesulfonamide:

1-(2-Amino-ethyl)-3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one
 (28 mg, 0.083 mmol) is added into DCM (1.0 mL) at room temperature and triethylamine (0.014
 mL, 0.10 mmol) is added. Then Benzenesulfonyl isocyanate (0.014 mL, 0.10 mmol) is added
 into the above reaction mixture. After stirring for 2hr, another 0.014 mL of Benzenesulfonyl
 isocyanate is added and the mixture is further stirred for 30 min. Then 1.0 mL of 1.0M HCl is
 added along with 5 mL of water. The mixture is extracted with EtOAc (3 x 20 mL) and the
 organic layers are combined, dried (Na_2SO_4) and concentrated to give crude product.
 Purification by preparative TLC affords 33 mg (76%) of the title compound. LCMS (ESMS):
 m/z 521.17 ($\text{M}+\text{H}^+$).

**Example 54: (3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-
 benzimidazol-1-yl}-N-(phenylsulfonyl)hexanamide**




N-{(S)-3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-benzenesulfonamide:


(S)-3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid (25 mg, 0.062 mmol) is dissolved THF (0.5 mL) and CDI (22 mg, 0.14 mmol) is added into it at room temperature. The mixture is stirred for 30 min and then it is heated at 55 °C for 1 hr. After the mixture is cooled down to room temperature, Benzenesulfonamide (19 mg, 0.12 mmol) is added and after 10 min, DBU (0.018 mL, 0.12 mmol) is added. The mixture is then stirred for 16 hr at room temperature. 1 mL of 1.0 M HCl is added followed by 10 mL of water. The mixture is extracted with EtOAc (3 x 20 mL) and the organic layers are combined, dried and concentrated to give crude product. Purification by Preparative TLC affords 16 mg (47%) of *N*-{(S)-3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoyl}-benzenesulfonamide. LCMS (ESMS): *m/z* 545.32 (*M*+*H*⁺).

The following compounds are synthesized using the same procedure.


(3*S*)-*N*-[(1,2-dimethyl-1*H*-imidazol-4-yl)sulfonyl]-3-{3-[(1,4-dimethyl-1*H*-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl} hexanamide. LCMS (ESMS): *m/z* 563.35 (*M*+*H*⁺).




(3*S*)-3-{3-[(1,4-dimethyl-1*H*-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}-*N*-(pyridin-2-ylsulfonyl)hexanamide. LCMS (ESMS): *m/z* 546.30 (*M*+*H*⁺).




(3*S*)-3-{3-[(1,4-dimethyl-1*H*-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}-*N*-(methylsulfonyl)hexanamide. LCMS (ESMS): *m/z* 483.25 (*M*+*H*⁺).




(3S)-N-(aminosulfonyl)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide. LCMS (ESMS): m/z 484.23 ($M+H^+$).




(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-[(1-methyl-1H-imidazol-4-yl)sulfonyl]hexanamide LCMS (ESMS): m/z 549.47 ($M+H^+$).




(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(isopropylsulfonyl)hexanamide LCMS (ESMS): m/z 511.27 ($M+H^+$).




(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(ethylsulfonyl)hexanamide. LCMS (ESMS): m/z 497.29 ($M+H^+$).




(3S)-N-(tert-butylsulfonyl)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide. LCMS (ESMS): m/z 525.32 ($M+H^+$).




(3S)-N-cyano-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide. LCMS (ESMS): m/z 430.22 ($M+H^+$).



(3S)-N-(cyclopropylsulfonyl)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide LCMS (ESMS): m/z 509.28 ($M+H^+$).



Example 55: 3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-4-methoxybutanoic acid



3-(2-Amino-phenylamino)-4-methoxy-butyric acid methyl ester:

4-Methoxy-3-oxo-butyric acid methyl ester (0.67 mL, 5.0 mmol) and Benzene-1,2-diamine (560 mg, 5.5 mmol) are dissolved in DCM (20 mL) and HOAc (0.63 mL, 11 mmol) is added into it. The mixture is stirred for 6 hrs and then $\text{NaBH}(\text{OAc})_3$ (3.7 g, 17 mmol) is added and the mixture is stirred for another 60 hr. Then 50 mL of sat. NaHCO_3 along with 50 mL of water are added and the mixture is extracted with EtOAc (3 x 100 mL). The organic layers are combined, dried and concentrated to give crude product. Purification by flash column chromatography affords 320 mg (27%) of 3-(2-Amino-phenylamino)-4-methoxy-butyric acid methyl ester.

4-Methoxy-3-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-butyric acid methyl ester:

To a solution of 3-(2-Amino-phenylamino)-4-methoxy-butyric acid methyl ester (320 mg, 1.3 mmol) in THF (5.0 mL) is added CDI (440 mg, 2.7 mmol) at room temperature. The mixture is stirred at the same temperature for 16 hours. Then 5 mL of 1.0 M HCl solution is added along with 20 mL of water. The mixture is extracted with EtOAc (3 x 25 mL) and the organic layers are combined, dried and concentrated to give crude product. Purification by flash column chromatography affords 205 mg (58%) of 4-Methoxy-3-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-butyric acid methyl ester.


3-[3-(1,4-Dimethyl-1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-4-methoxy-butyric acid methyl ester:

To a solution of 4-Methoxy-3-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-butyric acid methyl ester (205 mg, 0.78 mmol) in DMF (10 mL) are added (1,4-Dimethyl-1*H*-indol-3-ylmethyl)-trimethyl-ammonium iodide (470 mg, 1.4 mmol) and K₂CO₃ (210 mg, 1.6 mmol) at room temperature. The solution is heated to 100 °C for 5.5 hours. The solution is cooled down and 30 mL of water is added. The mixture is extracted with EtOAc (3 x 50 mL) and the organic layers are combined, washed with water (3x100 mL), dried with MgSO₄ and concentrated to give crude product. Purification by flash column chromatography affords 200 mg (61%) of 3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-4-methoxy-butylric acid methyl ester.

3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-4-methoxy-butylric acid:

To a solution of 3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-4-methoxy-butylric acid methyl ester (200 mg, 0.48 mmol) in 1,4-dioxane (2.0 mL) is added LiOH monohydrate (25 mg, 0.59 mmol) in water (2.0 mL) at room temperature. The solution is stirred at the same temperature for 2 hours. Then 1.0 mL of 1.0 M HCl solution is added along with 10 mL of water. Then the mixture is extracted with EtOAc (3 x 25 mL) and the organic layers are combined, dried and concentrated to give 190 mg (97%) of crude product. 30 mg of this crude is purified by preparative TLC to give 25 mg of 3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-4-methoxy-butylric acid. LCMS (ESMS): *m/z* 408.25 (M+H⁺).


Example 56: 3-{3-[(1,4-dimethyl-1*H*-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}-4-methoxy-N-[(1-methyl-1*H*-imidazol-4-yl)sulfonyl]butanamide



3-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-4-methoxy-butylric acid (30 mg, 0.074 mmol) is dissolved THF (0.70 mL) and CDI (27 mg, 0.17 mmol) is

added into it at room temperature. The mixture is heated at 55 °C for 1 hr. After the mixture is cooled down to room temperature, 1-Methyl-1*H*-imidazole-4-sulfonic acid amide (24 mg, 0.15 mmol) is added and after 10 min, DBU (0.022 mL, 0.15 mmol) is added. The mixture is stirred for 16 hr at room temperature. 2.0 mL of 1.0 M HCl is added followed by 20 mL of water. The mixture is extracted with EtOAc (3 x 25 mL) and the organic layers are combined, dried and concentrated to give crude product. Purification by flash column chromatography affords 17 mg (41%) of 1-Methyl-1*H*-imidazole-4-sulfonic acid {3-[3-(1,4-dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-4-methoxy-butyl}-amide. LCMS (ESMS): m/z 551.25 ($M+H^+$).

Example 57: 2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}pentane-1-sulfonic acid



(Diethoxy-phosphoryl)-methanesulfonic acid methyl ester:

Methanesulfonic acid methyl ester (0.43 mL, 5.0 mmol) is dissolved in THF (10 mL) and it is cooled down to -78 °C. 1.6 M *n*-BuLi (3.4 mL, 5.5 mmol) in hexane is added into it dropwise and the mixture is stirred for 30 min at -78 °C. Then Phosphorochloridic acid diethyl ester (0.41 mL, 2.8 mmol) is added dropwise into the mixture and it is then gradually warmed up to -50 °C in 2 hrs. 5 mL of sat. NH_4Cl and 20 mL of water are added and it is further warmed up to room temperature. The mixture is extracted with EtOAc (3 x 40 mL) and the organic layers are combined, dried (Na_2SO_4) and concentrated to give crude product. Purification by flash column chromatography affords 350 mg (53%) of (Diethoxy-phosphoryl)-methanesulfonic acid methyl ester.


Pent-1-ene-1-sulfonic acid methyl ester:

(Diethoxy-phosphoryl)-methanesulfonic acid methyl ester (150 mg, 0.61 mmol) is dissolved in THF (5.0 mL) and it is cooled down to -78 °C. 1.6M nBuLi (0.46 mL, 0.73 mmol) in hexane is added into it dropwise. The mixture is stirred for 30 min at that temperature and Butyraldehyde (0.066 mL, 0.73 mmol) is added. Then mixture is gradually warmed up to room temperature in 3 hrs and the reaction is further stirred for 2 hr at room temperature. Then 30 mL of water is added and the mixture is extracted with EtOAc (3 x 50 mL). The organic layers are combined, dried and concentrated to give 100 mg (100%) of Pent-1-ene-1-sulfonic acid methyl ester.

2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentane-1-sulfonic acid:

To a solution of 1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one (50 mg, 0.17 mmol) in THF (1.0 mL) are added Pent-1-ene-1-sulfonic acid methyl ester (56 mg, 0.34 mmol) and solid NaOH (14 mg, 0.34 mmol) in order. The resulting suspension is stirred at room temperature for 1 hr and it turned into a clear solution. Then 5 mL of sat.NH₄Cl is added along with 5 mL of water. The mixture is extracted with EtOAc (3 x 20 mL). The organic layers are combined, dried (MgSO₄) and concentrated to give crude product. Purification by flash column chromatography affords 11 mg (15%) of 2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-pentane-1-sulfonic acid. LCMS (ESMS): *m/z* 445.13 (M+H⁺).

Example 58: 2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}benzoic acid



Nitro-phenylamino)-benzoic acid:

1-Fluoro-2-nitro-benzene (0.53 mL, 5.0 mmol), 2-Amino-benzoic acid (750 mg, 5.5 mmol) and Na_2CO_3 (420 mg, 4.0 mmol) are added together in reaction flask and the mixture is heated at 130 °C for 60 hrs. After it is cooled down to room temperature, 50 mL of water is added and HOAc is used to adjust the pH to slightly acidic. Then the mixture is extracted with EtOAc (3 x 50 mL) and the organic layers are combined, washed with water (3 x 50 mL), dried (Na_2SO_4) and concentrated to give crude product. Purification by flash column chromatography affords 560 mg (43%) of 2-(2-Nitro-phenylamino)-benzoic acid.

2-(2-Nitro-phenylamino)-benzoic acid methyl ester:

2-(2-Nitro-phenylamino)-benzoic acid (560 mg, 2.2 mmol) is dissolved in methanol (10 mL) and concentrated H_2SO_4 (0.16 mL, 98%) is added into it. The mixture is then refluxed for 60 hr. After it is cooled down to room temperature, 100 mL of water is added and the pH of this solution is adjusted to pH 7 by adding 1.0 M NaOH aqueous solution. An orange solid is formed and it is filtered and rinsed with more water. Air drying of this orange solid affords 470 mg (80%) of 2-(2-Nitro-phenylamino)-benzoic acid methyl ester.

2-(2-Amino-phenylamino)-benzoic acid methyl ester:

2-(2-Nitro-phenylamino)-benzoic acid methyl ester (200mg, 0.74 mmol) is suspended in EtOAc (5.0 mL)) and 10% Pd/C wet (78 mg, 0.073 mmol) is added into it. A H_2 balloon is attached to the reaction flask and the reaction mixture is stirred under H_2 for 20 hrs. Then the Pd/C is

filtered and the filtrate is concentrated to give 165 mg (93%) of 2-(2-Amino-phenylamino)-benzoic acid methyl ester.

2-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-benzoic acid methyl ester:

To a solution of 2-(2-Amino-phenylamino)-benzoic acid methyl ester (75 mg, 0.31 mmol) in THF (1.0 mL) is added CDI (100 mg, 0.62 mmol) at room temperature. The mixture is stirred at the same temperature for 16 hours. Then 3 mL of 1.0 M HCl solution is added along with 20 mL of water. The mixture is extracted with EtOAc (3 x 30 mL) and the organic layers are combined, dried and concentrated to give crude product. Purification by flash column chromatography affords 65 mg (78%) of 2-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-benzoic acid methyl ester.

2-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-benzoic acid methyl ester:

To a solution of 2-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-benzoic acid methyl ester (30 mg, 0.11 mmol) in DMF (1.0 mL) are added (1,4-Dimethyl-1*H*-indol-3-ylmethyl)-trimethylammonium iodide (46 mg, 0.13 mmol) and K₂CO₃ (31 mg, 0.22 mmol) at room temperature. The solution is heated to 60 °C for 16 hr and it is cooled down to room temperature for another 48 hr. Then 30 mL of water is added. The mixture is extracted with EtOAc (3 x 30 mL) and the combined organic layers are washed with water (3 x 100 mL), dried with Na₂SO₄ and concentrated to give crude product. Purification by preparative TLC affords 40 mg (86%) of 2-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-benzoic acid methyl ester.


2-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-benzoic acid:

To a solution of 2-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-benzoic acid methyl ester (45 mg, 0.11 mmol) in 1,4-dioxane (1.0 mL) is added LiOH monohydrate (5.6 mg, 0.13 mmol) in water (0.1 mL) at room temperature. The solution is stirred at the same temperature for 16 hours. Then another 1.0 mg of LiOH monohydrate is added and the reaction is continued for another 2 hr. Then 0.5 mL of 1.0 M HCl is added along with 10 mL of water. The mixture is extracted with EtOAc (3 x 15 mL) and the organic layers


are combined, dried and concentrated to give crude product. Purification by preparative TLC affords 41 mg (95%) of 2-[3-(1,4-Dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-benzoic acid. LCMS (ESMS): m/z 412.17 ($M+H^+$).

The following compounds are synthesized using the same procedure.


2-{3-[(1,4-dimethyl-1*H*-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}-3-methoxybenzoic acid. LCMS (ESMS): m/z 442.06 ($M+H^+$).




2-{3-[(1,4-dimethyl-1*H*-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}-3-methylbenzoic acid. LCMS (ESMS): m/z 426.04 ($M+H^+$).



3-{3-[(1,4-dimethyl-1*H*-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}-4-methoxybenzoic acid. LCMS (ESMS): m/z 442.18 ($M+H^+$).



Example 59: 1-[2-(1,1-dioxido-4-oxo-1,2,5-thiadiazolidin-2-yl)ethyl]-3-[(4-methyl-1-benzothien-3-yl)methyl]-1,3-dihydro-2*H*-benzimidazol-2-one



{2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethylamino}-acetic acid methyl ester:


1-(2-Bromo-ethyl)-3-(4-methyl-benzo[*b*]thiophen-3-ylmethyl)-1,3-dihydro-benzimidazol-2-one (30 mg, 0.075 mmol) is dissolved in CH₃CN (0.5 mL) and Methoxycarbonylmethyl-ammonium chloride (11 mg, 0.09 mmol) and Et₃N (0.025 mL, 0.18 mmol) are added. The mixture is first stirred at room temperature for 16 hr. And then 11 mg of Na₂CO₃ is added and the mixture is stirred at room temperature for another 48 hr. Since the reaction is not completed, another 11 mg of Methoxycarbonylmethyl-ammonium chloride, 0.025 mL of Et₃N and 11 mg of Na₂CO₃ are added. The mixture is then heated at 80 °C for 20 hrs. Then 10 mL of water is added and the mixture is extracted with EtOAc (3 x 15 mL). The organic layers are combined, dried and concentrated to give crude product. Purification by flash column chromatography affords 30 mg (98%) of {2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethylamino}-acetic acid methyl ester.

1-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-3-[2-(1,1,4-trioxo-1,6-1,2,5-thiadiazolidin-2-yl)-ethyl]-1,3-dihydro-benzimidazol-2-one:

{2-[3-(4-Methyl-benzo[*b*]thiophen-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-ethylamino}-acetic acid methyl ester (30 mg, 0.073 mmol) and sulfamide (7.0 mg, 0.073 mmol) are added into the same reaction flask and DBU (0.1 mL, 0.67 mmol) is added into them. The neat reaction mixture is heated at 130 °C for 40 min. After it is cooled down to room temperature, 1.5 mL of 1.0M HCl solution is added along with 15 mL of water. The mixture is extracted with EtOAc (3 x 15 mL) and the organic layers are combined, dried and concentrated to give crude product. Purification by preparative TLC affords 14 mg (42%) of 1-(4-Methyl-

benzo[*b*]thiophen-3-ylmethyl)-3-[2-(1,1,4-trioxo-1 $\square$ 6-1,2,5-thiadiazolidin-2-yl)-ethyl]-1,3-dihydro-benzimidazol-2-one. LCMS (ESMS): m/z 457.08 ($M+H^+$).

Example 60: 2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclopent-1-ene-1-carboxylic acid



A solution of 500 mg (4.62 mmol) of 1,2-phenylenediamine and 0.7 mL (4.62 mmol) of ethyl 2-oxocyclopentane carboxylate in 5 mL of dichloroethane was treated with 653 mg (3.08 mmol, 0.67 eq.) of sodium triacetoxyborohydride and 5 drops of acetic acid at room temperature. The resulting mixture was stirred at this temperature for h. Then the mixture was diluted with water and the organic layer was separated. The organic layer was dried over magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography to give 375 mg (33 %) of desired compound as light tan solid.


A solution of 370 mg (1.5 mmol) of 2-(2-amino-phenylamino)-cyclopent-1-enecarboxylic acid ethyl ester and 483 mg (3.0 mmol, 2.0 eq.) of CDI in 10 mL of THF was stirred room temperature for 16h. The resulting mixture was concentrated and the residue was purified by combiflash to give 120 mg (29%) of desired compound as light tan oil.

To a solution of 100mg (0.37 mmol) of 2-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-cyclopent-1-enecarboxylic acid ethyl ester in 5mL of DMF was added 101mg (0.73 mmol, 2.0 eq.) of potassium carbonate followed by 190 mg (0.55 mmol, 1.5 eq.) indole ammonium iodide at room temperature under nitrogen atmosphere. The resulting suspension was heated to 100°C for 4

hours. The solution was cooled and stirred at RT over night. The reaction was poured into water and extracted with EtOAc (3x). The combined organics were washed with water, brine and dried (magnesium sulfate). Filtration and concentration gave the crude product which was absorbed onto silica gel and purified *via* CombiFlash to give 120 mg (76 %) of the desired product as a light yellow oil.


A solution of 120 mg (0.28 mmol) of 2-[3-(1,4-dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-cyclopent-1-enecarboxylic acid ethyl ester and 23 mg (0.56 mmol, 2.0 eq.) of lithium hydroxide monohydrate in 5 mL of methanol and 1 mL of water was stirred at room temperature for 2.5 days. Then the pH of the resulting mixture was adjusted to 3 with acetic acid and diluted with water. The precipitate was collected and washed with water. The isolated light pink solid was pumped overnight to give 94 mg (84%) of 2-[3-(1,4-dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-cyclopent-1-enecarboxylic acid as a light pink solid.

Example 61: 2-{3-[(1,4-dimethyl-1*H*-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}cyclopentanecarboxylic acid



A mixture of 65 mg (0.16 mmol) of 2-{3-[(1,4-dimethyl-1*H*-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}cyclopent-1-ene-1-carboxylic acid and 30 mg of 10% Pd/C in 3 mL of methanol was stirred at room temperature under hydrogen gas atmosphere (1 atm) for 16h. The resulting mixture was filtered through Celite and the filtrate was concentrated. The residue was purified by combiflash to give 20 mg (31%) of 2-[3-(1,4-dimethyl-1*H*-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-cyclopentanecarboxylic acid as a white solid.

Example 62: 3-{3-[(2,3-dimethylindolizin-8-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}propanoic acid



2,3-Dimethyl-indolizine-8-carboxylic acid methyl ester:

A solution of methyl 2-methylnicotinate (4.9g, 32.1mmol) and 3-bromo-2-butanone (4.8g, 32.1mmol) in acetone (40ml) were warmed to reflux for 3d. Cooled to RT and partially purified the remaining residue *via* column chromatography (silica gel, 5-10% EtOAc/hexanes). The product-containing fractions were combined and concentrated to give the desired product.

(2,3-Dimethyl-indolizin-8-yl)-methanol:

A solution of the 2,3-dimethyl-indolizine-8-carboxylic acid methyl ester (265mg, 1.3mmol) in ether (10ml) was cooled to 0°C and treated with LAH (1.0M in THF, 1.6ml, 1.6mmol, dropwise addition). After 3.5h the reaction was quenched with sodium sulfate decahydrate (added in small portions until gas evolution ceased). The resulting mixture was stirred for 0.5h and the precipitated solids were removed *via* filtration and washed with EtOAc. The combined filtrates were concentrated to give the product which was used without further purification.

3-[3-(2,3-Dimethyl-indolizin-8-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid methyl ester:


To a stirred solution of the 3-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-propionic acid methyl ester (100mg, 0.45mmol) and (2,3-Dimethyl-indolizin-8-yl)-methanol (88mg, 0.50mmol) in THF (8ml) was added triphenylphosphine (143mg, 0.54mmol) followed by DIAD (0.11ml, 0.54mmol, dropwise) at ambient temperature. After five minutes the reaction became dark brown and was consequently wrapped in aluminum foil. After stirring over night the reaction was treated with silica gel and concentrated. The remaining solid was purified via column chromatography (silica

gel, 5-40% EtOAc/hexanes). The product-containing fractions were combined and concentrated to give the product which was used without further purification.

3-{3-[(2,3-dimethylindolizin-8-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid:

To a stirred solution of 3-[3-(2,3-dimethyl-indolizin-8-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid methyl ester (110mg, 0.29mmol) in THF (5ml) and water (1.5ml) was added lithium hydroxide monohydrate (25mg, 0.58mmol). After 1.5h the reaction appeared complete (LC-MS) and was subsequently concentrated. The remaining residue was diluted with water (~12ml) and acidified to pH~5 using HOAc (~5 drops). The resulting solid was dissolved into DCM and the layers were separated. The aqueous phase was extracted with DCM (1x) and the combined organics were concentrated. The remaining green residue was dissolved into a minimal volume of DCM and applied to a preparative TLC plate (1mm). Eluted with 8% MeOH/DCM. The product-band was isolated and suspended in 5% MeOH/DCM. The silica gel was removed *via* filtration and washed with DCM. The combined filtrates were concentrated to give the desired product as a light green solid. LCMS (ESMS): m/z 364.24..

Example 63: 3-{3-[(1-cyano-5-methylindolizin-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid



To a solution of 2-picoline (5.3 mL, 0.054 mol) in Et₂O (15 mL) was added chloroacetic acid (6 g, 0.064 mol) at room temperature. The solution was stirred at the same temperature for 12 hours. The solution was concentrated and the residue oil was used in the next step of the synthesis without further purification.

To a solution of crude 1-Carboxymethyl-2-methyl-pyridinium chloride (7 g, 0.037 mol) in toluene (50 mL) were added acrylonitrile (7.4 mL, 0.11 mol), MnO₂ (6.5 g, 0.075 mol) and Et₃N (2 mL) at room temperature under nitrogen atmosphere. The solution was heated to 90 °C and was stirred at the same temperature for 12 hours. The solution was cooled down and was filtered to remove the excess MnO₂. The filtrate was concentrated and the residue was purified by CombiFlash with 10% EtOAc in Hexane to afford the desirable product 5-Methyl-indolizine-1-carbonitrile, CAS: 5-Methyl-indolizine-1-carbonitrile (3g, 52%) as a white solid.

A solution of: 5-Methyl-indolizine-1-carbonitrile (200 mg, 1.28 mmol) in DMF (0.12 mL, 1.54 mmol) was cooled to ~5°C under nitrogen. Pyrophosphoryl chloride (0.2 mL, 1.54 mmol) was then slowly introduced. Upon complete addition the cooling bath was removed and the reaction was allowed to warm to ambient temperature. After a total of 45 minutes the reaction was recooled to 5°C and slowly quenched with 2N NaOH. The pH was adjusted to ~8. The solid that precipitate out from the solution was collected by filtration and was washed with water. The remaining white solid was dried in the oven and was confirmed to be the desirable product 3-Formyl-5-methyl-indolizine-1-carbonitrile, CAS: 3-Formyl-5-methyl-indolizine-1-carbonitrile (210 mg, 89%).

To a solution of 3-Formyl-5-methyl-indolizine-1-carbonitrile (200 mg, 1.09 mmol) in CH₂Cl₂ (10 mL) and dimethylamine (2M MeOH solution) (1.09 mL, 2.17 mmol) was added NaBH(OAc)₃ (460 mg, 2.17 mmol) at 0°C. Upon complete addition the cooling bath was removed and the reaction mixture was stirred at room temperature for 12h. When the reaction was complete, the mixture was diluted with CH₂Cl₂ and washed with saturated NaHCO₃. The layers were separated and the organic phase was dried over MgSO₄, filtered and concentrated. The residue was purified by CombiFlash with 20% EtOAc in Hexane as the eluent to afford the

desirable product 3-Dimethylaminomethyl-5-methyl-indolizine-1-carbonitrile, CAS: 3-Dimethylaminomethyl-5-methyl-indolizine-1-carbonitrile (150 mg, 65%) as a white solid.

To a stirred solution of 3-Dimethylaminomethyl-5-methyl-indolizine-1-carbonitrile (60 mg, 0.28 mmol) in acetonitrile (5 mL) was added iodomethane (0.02 mL, 0.34 mmol) at room temperature. The solution was stirred at the same temperature for 5 hours. The solution was concentrated and the resulting white solid was washed with small amount of acetonitrile (0.5 mL) and Et₂O was added. The white solid was collected by filtration and was washed with Et₂O. The white solid was confirmed to be the desirable product 3-trimethylaminomethyl-5-methyl-indolizine-1-carbonitrile iodide, CAS: 3-trimethylaminomethyl-5-methyl-indolizine-1-carbonitrile iodide (60 mg, 60%).

To a solution of 3-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-hexanoic acid ethyl ester (50 mg, 0.18 mmol) in DMF (5 mL) were added 3-trimethylaminomethyl-5-methyl-indolizine-1-carbonitrile iodide, CAS: 3-trimethylaminomethyl-5-methyl-indolizine-1-carbonitrile iodide (96 mg, 0.67 mmol) and K₂CO₃ (50 mg, 0.36 mmol) at room temperature under nitrogen atmosphere. The solution was heated to 100 C for 2 hours. The solution was cooled down and water was added. The solution was extracted with EtOAc and the combined organic layer was dried with MgSO₄. The solution was filtered and the filtrate was concentrated. The residue was purified by CombiFlash with 20% EtOAc in Hexane as the eluent to afford the desirable product 3-[3-(1-Cyano-5-methyl-indolizin-3-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid ethyl ester, CAS: 3-[3-(1-Cyano-5-methyl-indolizin-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester (36 mg, 45%) as yellow foam.

To a solution of 3-[3-(1-Cyano-5-methyl-indolizin-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester (30 mg, 0.067 mmol) in 1,4-dioxane (5 mL) and water (1 mL) was added LiOH (3.2 mg, 0.14 mmol) at room temperature. The solution was stirred at the same temperature for 12 hours. The solution was concentrated and water was added to the residue. The solution was acidified by 12N HCl in an ice-bath. The solid that precipitate out from the solution was collected by filtration. The solid was dried under vacuum

To a solution of 3,5-dichloropyridine (5 g, 0.034 mol) in Et₂O (10 mL) was added 2-chloropropionic acid (3.5 mL, 0.041 mol) at room temperature. The solution was stirred at the same temperature for 12 hours. The solution was concentrated and the residue oil was used in the next step of the synthesis without further purification.

To a solution of crude N-carbonyl-ethyl-3,5-Dichloro-pyridinium chloride (10 g, 0.039 mol) in toluene (100 mL) were added ethyl acrylate (12.7 mL, 0.117 mol), MnO₂ (10.2 g, 0.117 mol) and Et₃N (5 mL) at room temperature under nitrogen atmosphere. The solution was heated to 90 °C and was stirred at the same temperature for 12 hours. The solution was cooled down and was filtered to remove the excess MnO₂. The filtrate was concentrated and the residue was purified by CombiFlash with 10% EtOAc in Hexane to afford the desirable product 6,8-Dichloro-3-methyl-indolizine-1-carboxylic acid ethyl ester (150 mg, 1.4%) as a pale yellow foam.

To a solution of 6,8-Dichloro-3-methyl-indolizine-1-carboxylic acid ethyl ester (80 mg, 0.29 mmol) in Et₂O (10 mL) was added LAH (22.3 mg, 0.58 mmol) at 0 °C under nitrogen atmosphere. The solution was warmed to room temperature. Sat. NaHCO₃ was added and the solution was extracted with EtOAc. The combined organic layer was dried with MgSO₄ and was filtered. The filtrate was concentrated and the residue was subjected to ¹H-NMR analysis to confirm the presence of the desirable product (6,8-Dichloro-3-methyl-indolizine-1-yl)-methanol, CAS: (6,8-Dichloro-3-methyl-indolizine-1-yl)-methanol (60 mg, 89%).

To a solution of (6,8-Dichloro-3-methyl-indolizine-1-yl)-methanol (100 mg, 0.44 mmol) in acetone was added MnO₂ (75.5 mg, 0.87 mmol) at room temperature. The mixture was stirred at the same temperature for 3 hours. The solution was filtered and the filtrate was concentrated. The residue was purified by CombiFlash with 5% EtOAc in Hexane as the eluent to afford the desirable product 6,8-Dichloro-3-methyl-indolizine-1-carbaldehyde, CAS: 6,8-Dichloro-3-methyl-indolizine-1-carbaldehyde (50 mg, 50%) as a white solid.

To a solution of 6,8-Dichloro-3-methyl-indolizine-1-carbaldehyde (50 mg, 0.22 mmol) in CH₂Cl₂ (20 mL) and dimethylamine (2M methanol solution) (0.33 mL, 0.66 mmol) was added

NaBH(OAc)₃ (232 mg, 1.1 mmol) at 0°C. Upon complete addition the cooling bath was removed and the reaction mixture was stirred at room temperature for 48h. When the reaction was complete, the mixture was diluted with CH₂Cl₂ and washed with saturated NaHCO₃. The layers were separated and the organic phase was dried over MgSO₄, filtered and concentrated. The residue was used in the next step of the synthesis without purification.


To a stirred solution of crude (6,8-Dichloro-3-methyl-indolizin-1-ylmethyl)-dimethyl-amine (50 mg, 0.19 mmol) in acetonitrile (5 mL) was added iodomethane (0.04 mL, 0.58 mmol) at room temperature. The solution was stirred at the same temperature for 5 hours. The solution was concentrated and the resulting white solid was washed with small amount of acetonitrile (0.5 mL) and Et₂O was added. The white solid was collected by filtration and was washed with Et₂O. The white solid was confirmed to be the desirable product (6,8-Dichloro-3-methyl-indolizin-1-ylmethyl)-dimethyl-amine iodide (60 mg, 77%).

To a solution of 3-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-hexanoic acid ethyl ester (30 mg, 0.11 mmol) in DMF (10 mL) were added (6,8-Dichloro-3-methyl-indolizin-1-ylmethyl)-dimethyl-amine iodide, CAS: (6,8-Dichloro-3-methyl-indolizin-1-ylmethyl)-dimethyl-amine iodide (65 mg, 0.16 mmol), and K₂CO₃ (45 mg, 0.33 mmol) at room temperature under nitrogen atmosphere. The solution was heated to 100 C for 2 hours. The solution was cooled down and water was added. The solution was extracted with EtOAc and the combined organic layer was dried with MgSO₄. The solution was filtered and the filtrate was concentrated. The residue was purified by CombiFlash with 20% EtOAc in Hexane as the eluent to afford the desirable product 3-[3-(6,8-Dichloro-3-methyl-indolizin-1-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester (20 mg, 38%) as a yellow foam.

To a solution of 3-[3-(6,8-Dichloro-3-methyl-indolizin-1-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester (20 mg, 0.041 mmol) in 1,4-dioxane (5 mL) and water (1 mL) was added LiOH (2 mg, 0.082 mmol) at room temperature. The solution was stirred at the same temperature for 4 hours. The solution was concentrated and water was added to the residue. The solution was acidified by 12N HCl in an ice-bath. The solid that precipitate

out from the solution was collected by filtration. The solid was dried under vacuum and was confirmed to be the desirable product. LCMS (ESMS): m/z 460.17 ($M+H^+$).

Example 65: 3-{3-[(3,8-dimethylindolizin-1-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid



To a solution of 3-picoline (2 mL, 0.021 mol) in Et₂O (10 mL) was added 2-chloropropionic acid (2.1 mL, 0.025 mol) at room temperature. The solution was stirred at the same temperature for 12 hours. The solution was concentrated and the residue oil was used in the next of the synthesis without further purification.

To a solution of crude 1-(1-Carboxy-ethyl)-3-methyl-pyridinium; chloride (6g, 0.03 mol) in toluene (100 mL) were added ethyl acrylate (9.7 mL, 0.089 mol), MnO₂ (7.8 g, 0.089 mol) and Et₃N (5 mL) at room temperature under nitrogen atmosphere. The solution was heated to 90 C and was stirred at the same temperature for 12 hours. The solution was cooled down and was filtered to remove the excess MnO₂. The filtrate was concentrated and the residue was purified by CombiFlash with 10% EtOAc in Hexane to afford the desirable product 3,8-Dimethyl-indolizine-1-carboxylic acid ethyl ester (735 mg, 11.4%).

To a solution of 3,8-Dimethyl-indolizine-1-carboxylic acid ethyl ester (700 mg, 3.2 mmol) in Et₂O (30 mL) was added LAH (244.5 mg, 6.4 mmol) at 0 C under nitrogen atmosphere. The

solution was warmed to room temperature. Sat. NaHCO_3 was added and the solution was extracted with EtOAc. The combined organic layer was dried with MgSO_4 and was filtered. The filtrate was concentrated and the residue was subjected to ^1H -NMR analysis to confirm the presence of the desirable product (3,8-Dimethyl-indolizin-1-yl)-methanol, CAS: (3,8-Dimethyl-indolizin-1-yl)-methanol (450 mg, 80%).

To a solution (3,8-Dimethyl-indolizin-1-yl)-methanol (600 mg, 3.4 mmol) in acetone (20 mL) was added MnO_2 (595 mg, 6.8 mmol) at room temperature. The mixture was stirred at the same temperature for 3 hours. The solution was filtered and the filtrate was concentrated. The residue was purified by CombiFlash with 5% EtOAc in Hexane as the eluent to afford the desirable product 3,8-Dimethyl-indolizine-1-carbaldehyde (400 mg, 67%) as a grey solid.

To a solution of 3,8-Dimethyl-indolizine-1-carbaldehyde (300 mg, 1.7 mmol) in CH_2Cl_2 (60 mL) and dimethylamine (2M methanol solution)(4.3 mL, 8.7 mmol) was added $\text{NaBH}(\text{OAc})_3$ (1.8 g, 8.6 mmol) at 0°C . Upon complete addition the cooling bath was removed and the reaction mixture was stirred at room temperature for 12h. When the reaction was complete, the mixture was diluted with CH_2Cl_2 and washed with saturated NaHCO_3 . The layers were separated and the organic phase was dried over MgSO_4 , filtered and concentrated. The residue was used in the next step of the synthesis without purification.


To a stirred solution of (3,8-Dimethyl-indolizin-1-ylmethyl)-dimethyl-amine (100 mg, 0.49 mmol) in acetonitrile (5 mL) was added iodomethane (0.1 mL, 1.5 mmol) at room temperature. The solution was stirred at the same temperature for 5 hours. The solution was concentrated and the resulting white solid was washed with small amount of acetonitrile (0.5 mL) and Et_2O was added. The white solid was collected by filtration and was washed with Et_2O . The reddish brown solid was used in the next step of the synthesis without further purification.

To a solution of 3-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-hexanoic acid ethyl ester (50 mg, 0.18 mmol) in DMF (10 mL) were added (3,8-Dimethyl-indolizin-1-ylmethyl)-trimethyl-ammonium; iodide, CAS: (3,8-Dimethyl-indolizin-1-ylmethyl)-trimethyl-ammonium; iodide (93.4 mg, 0.27 mmol) and K_2CO_3 (75 mg, 0.54 mmol) at room temperature under nitrogen

atmosphere. The solution was heated to 100 C for 2 hours. The solution was cooled down and water was added. The solution was extracted with EtOAc and the combined organic layer was dried with MgSO₄. The solution was filtered and the filtrate was concentrated. The residue was purified by CombiFlash with 20% EtOAc in Hexane as the eluent to afford the desirable product 3-[3-(3,8-Dimethyl-indolizin-1-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester (20 mg, 26%) as a colorless foam.

To a solution of 3-[3-(3,8-Dimethyl-indolizin-1-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester (10 mg, 0.023 mmol) in 1,4-dioxane (5 mL) and water (1 mL) was added LiOH (1.1 mg, 0.046 mmol) at room temperature. The solution was stirred at the same temperature for 4 hours. The solution was concentrated and water was added to the residue. The solution was acidified by 12N HCl in an ice-bath. The solid that precipitate out from the solution was collected by filtration. The solid was dried under vacuum and was confirmed to be the desirable product. LCMS (ESMS): m/z 406.27 (M+H⁺).

Example 66: 3-{3-[(5-bromo-1-methyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid



A solution of the 5-bromo-indole-7-carboxylic acid methyl ester (300 mg, 1.18 mmol) in DMF (15 mL) was cooled to 0°C under nitrogen and treated with sodium hydride (60% in mineral oil) (56.6 gm, 1.4 mmol). After stirring for 15 minutes the iodomethane (0.08 mL, 1.3 mmol) was introduced and the cooling bath was removed (reaction became light purple). After 30 minutes

the reaction was quenched with water (5ml) and then concentrated to reduce the volume of DMF. The reaction was poured into water (1.6L) and EtOAc (200ml). The layers were separated and the aqueous phase was extracted with EtOAc (3x200ml). The combined organics were washed with water (3x) and then dried (MgSO₄). The solution was filtered and the filtrate was concentrated. The residue was purified by CombiFlash with 10% EtOAc in Hexane as the eluent to afford the desirable product 5-Bromo-1-methyl-1*H*-indole-7-carboxylic acid methyl ester (290 mg, 92%) as a white solid.

To a solution of 5-Bromo-1-methyl-1*H*-indole-7-carboxylic acid methyl ester (260 mg, 0.97 mmol) in Et₂O (50 mL) was added LAH (40.5 mg, 1.1 mmol) at 0 °C under nitrogen atmosphere. The solution was warmed to room temperature. Sat. NaHCO₃ was added and the solution was extracted with EtOAc. The combined organic layer was dried with MgSO₄ and was filtered. The filtrate was concentrated and the residue was purified by CombiFlash with 50% EtOAc in Hexane as the eluent to afford the desirable product (5-Bromo-1-methyl-1*H*-indol-7-yl)-methanol (200 mg, 86%) as colorless oil.

To a solution of (5-Bromo-1-methyl-1*H*-indol-7-yl)-methanol (220 mg, 0.92 mmol) in acetone (20 mL) was added MnO₂ (239 mg, 2.7 mmol) at room temperature. The mixture was stirred at the same temperature for 3 hours. The solution was filtered and the filtrate was concentrated. The residue was purified by CombiFlash with 20% EtOAc in Hexane as the eluent to afford the desirable product 5-Bromo-1-methyl-1*H*-indole-7-carbaldehyde (177 mg, 81%) as a white solid.


To a solution of 5-Bromo-1-methyl-1*H*-indole-7-carbaldehyde (175 mg, 0.74 mmol) in CH₂Cl₂ (50 mL) and dimethylamine (2M methanol solution) (0.74 mL, 1.5 mmol) was added NaBH(OAc)₃ (467 mg, 2.2 mmol) at 0°C. Upon complete addition the cooling bath was removed and the reaction mixture was stirred at room temperature for 12h. When the reaction was complete, the mixture was diluted with CH₂Cl₂ and washed with saturated NaHCO₃. The layers were separated and the organic phase was dried over MgSO₄, filtered and concentrated. The residue was used in the next step of the synthesis without purification.

To a stirred solution of crude (5-Bromo-1-methyl-1*H*-indol-7-ylmethyl)-dimethyl-amine (230 mg, 0.86 mmol) in acetonitrile (10 mL) was added iodomethane (0.06 mL, 1.0 mmol) at room temperature. The solution was stirred at the same temperature for 5 hours. The solution was concentrated and the resulting white solid was washed with small amount of acetonitrile (0.5 mL) and Et₂O was added. The white solid was collected by filtration and was washed with Et₂O. The white solid was used in the next step of the synthesis without further purification.

To a solution of 3-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-hexanoic acid ethyl ester (50 mg, 0.18 mmol) in DMF (10 mL) were added (5-Bromo-1-methyl-1*H*-indol-7-ylmethyl)-trimethyl-ammonium; iodide (111 mg, 0.27 mmol) and K₂CO₃ (75 mg, 0.54 mmol) at room temperature under nitrogen atmosphere. The solution was heated to 100 C for 2 hours. The solution was cooled down and water was added. The solution was extracted with EtOAc and the combined organic layer was dried with MgSO₄. The solution was filtered and the filtrate was concentrated. The residue was purified by CombiFlash with 20% EtOAc in Hexane as the eluent to afford the desirable product 3-[3-(5-Bromo-1-methyl-1*H*-indol-7-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester (50 mg, 55%) as a colorless foam.

To a solution of 3-[3-(5-Bromo-1-methyl-1*H*-indol-7-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester (10 mg, 0.02 mmol) in 1,4-dioxane (5 mL) and water (1 mL) was added LiOH (1 mg, 0.04 mmol) at room temperature. The solution was stirred at the same temperature for 4 hours. The solution was concentrated and water was added to the residue. The solution was acidified by 12N HCl in an ice-bath. The solid that precipitate out from the solution was collected by filtration. The solid was dried under vacuum and was confirmed to be the desirable product. LCMS (ESMS): *m/z* 471.17 (M+H⁺).


Example 67: 3-{3-[(5-bromo-1-methyl-2-oxo-2,3-dihydro-1*H*-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}hexanoic acid



To a solution of 3-[3-(5-Bromo-1-methyl-1*H*-indol-7-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid (20 mg, 0.04 mmol) in tBuOH (5 mL) was added NBS (15 mg, 0.085 mmol) at 40 °C. The solution was stirred at the same temperature for 1 hour. The solution was concentrated and the residue was purified by CombiFlash with 5% MeOH in CH₂Cl₂ as the eluent to afford the desirable product 3-[2-Oxo-3-(3,3,5-tribromo-1-methyl-2-oxo-2,3-dihydro-1*H*-indol-7-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid (20 mg, 73%) as colorless oil.

To a solution of 3-[2-Oxo-3-(3,3,5-tribromo-1-methyl-2-oxo-2,3-dihydro-1*H*-indol-7-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid (20 mg, 0.031 mmol) in acetic acid (5 mL) was added Zinc powder (10 mg, 0.16 mmol) at room temperature. The mixture was stirred at the same temperature for 30 minutes. The mixture was filtered and the filtrate was concentrated. The residue was cooled in an ice bath and water was added. The solid that precipitated out from the solution was collected by filtration and was washed with water. The white solid was dried in an oven and was confirmed to be the desirable product (8.9 mg, 59%). LCMS (ESMS): *m/z* 488.12 (M+H⁺).

Example 68: 3-{3-[(4,6-dimethyl-1,2-benzisothiazol-3-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}hexanoic acid



To a solution of AlCl_3 (0.4g, 3.6 mmol) in CH_2Cl_2 (4 mL) at -20°C under nitrogen were added 3,5-dimethylbenzenethiol (1g, 7.2 mmol) and oxalyl chloride (0.6 mL, 7.2 mmol) respectively. The solution was warmed to room temperature for 1 hour and was heated to 120°C in a microwave reactor for 15 minutes. The solution was cooled down and was pouled into crashed ice. The solution was extracted with CH_2Cl_2 and the organic layer was collected. The solution was dried with MgSO_4 and was filtered. The filtrate was concentrated and the residue was purified by Combi-Flash with 5% EtOAc in Hexane as the eluent to afford the desirable product 4,6-Dimethyl-benzo[*b*]thiophene-2,3-dione (500 mg, 35%).

To a solution of 4,6-Dimethyl-benzo[*b*]thiophene-2,3-dione (100 mg, 0.5 mmol) in NH_3 in MeOH (5 mL) was dropwisely added 30 % H_2O_2 (0.17 mL, 1.6 mmol) at room temperature. The solution was stirred at the same temperature for 2 hours. The solution was concentrated and the residue was purified by CombiFlash with 20% EtOAc in Hexane as the eluent to afford the desirable product 4,6-Dimethyl-1,2-benzisothiazole-3-carboxylic acid amide (35 mg, 33%) as a pale red solid.

To a solution of 4,6-Dimethyl-1,2-benzisothiazole-3-carboxylic acid amide (20 mg, 0.097 mmol) in EtOH (10 mL) and H_2O (2 mL) was added KOH (11 mg, 0.19 mmol). The solution was heated to 85°C for 48 hours. The solution was cooled down and was placed in an ice bath. Conc HCl was added to adjust the pH of the solution to 2. The solid the precipitated out from

the solution was collected by means of filtration. The white solid was dried and was confirmed to be the desirable product 4,6-Dimethyl-1,2-benzisothiazole-3-carboxylic acid (15 mg, 75%).

To a solution of 4,6-Dimethyl-1,2-benzisothiazole-3-carboxylic acid (10 mg, 0.048 mmol) in MeOH (5 mL) was added conc. H₂SO₄ (0.1 mL) at room temperature. The solution was heated up to 60 C for 24 hours. The solution was cooled down and was neutralized with sat. NaHCO₃. The solution was extracted with EtOAc. The combined organic layer was dried with MgSO₄ and was filtered. The filtrate was concentrated and the residue was purified by CombiFlash with 20% EtOAc in Hexane as the eluent to afford the desirable product 4,6-Dimethyl-benzo[*d*]isothiazole-3-carboxylic acid methyl ester, CAS: 4,6-Dimethyl-1,2-benzisothiazole-3-carboxylic acid methyl ester (10 mg, 94%) as colorless oil.

To a solution of 4,6-Dimethyl-1,2-benzisothiazole-3-carboxylic acid methyl ester (100 mg, 0.45 mmol) in THF (10 mL) was added LAH (34 mg, 0.9 mmol) at 0 C under nitrogen atmosphere. The solution was stirred at the same temperature for 1 hour. Sat. NaHCO₃ was added and the solution was extracted with EtOAc. The combined organic layer was dried with MgSO₄ and was filtered. The filtrate was concentrated and the residue was used in the next step of the synthesis without further purification.


To a solution of (4,6-Dimethyl-1,2-benzisothiazol-3-yl)-methanol (100 mg, 0.5 mmol) in CH₂Cl₂ (15 mL) were added PPh₃ (200 mg, 0.78 mmol) and CBr₄ (340 mg, 1 mmol) at room temperature. The solution was stirred at the same temperature for 1 hour. The solution was concentrated and the residue was purified by CombiFlash with 10% EtOAc in Hexane as the eluent to afford the desirable product 3-Bromomethyl-4,6-dimethyl-1,2-benzisothiazole (90 mg, 68%) as a white solid.

To a solution of 3-(2-Oxo-2,3-dihydro-benzimidazol-1-yl)-hexanoic acid ethyl ester (50 mg, 0.18 mmol) in DMF (10 mL) were added 3-Bromomethyl-4,6-dimethyl-benzo[*d*]isothiazole, CAS: 3-Bromomethyl-4,6-dimethyl-1,2-benzisothiazole (69.5 mg, 0.27 mmol) and K₂CO₃ (50 mg, 0.36 mmol) at room temperature under nitrogen atmosphere. The solution was heated to 100 C for 2 hours. The solution was cooled down and water was added. The solution was


extracted with EtOAc and the combined organic layer was dried with MgSO_4 . The solution was filtered and the filtrate was concentrated. The residue was purified by CombiFlash with 20% EtOAc in Hexane as the eluent to afford the desirable product 3-[3-(4,6-Dimethyl-1,2-benzisothiazol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester (45 mg, 55%) as colorless foam.

To a solution of product 3-[3-(4,6-Dimethyl-1,2-benzisothiazol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-hexanoic acid ethyl ester (23 mg, 0.051 mmol) in 1,4-dioxane (5 mL) and water (1 mL) was added LiOH (2.4 mg, 0.1 mmol) at room temperature. The solution was stirred at the same temperature for 4 hours. The solution was concentrated and water was added to the residue. The solution was acidified by 12N HCl in an ice-bath. The solid that precipitate out from the solution was collected by filtration. The solid was dried under vacuum and was confirmed to be the desirable product (10 mg, 46%). LCMS (ESMS): m/z 424.18 ($\text{M}+\text{H}^+$).

The following compound was prepared following a similar procedure using appropriate starting material.



Example 69: 3-{3-[(6-bromo-2-oxo-2,3-dihydro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid



3,3,6-Tribromo-2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid methyl ester:

6-Bromo-1H-indole-4-carboxylic acid methyl ester 1.0g (3.93 mmol, DE3525564) is dissolved in *tert*-butanol at 40°C and then pyridinium bromide perbromide (4.0g, 12.5 mmol) is added portion wise over 30 minutes. The reaction is stirred for two hours then the volatiles evaporated *in vacuo* and the residue taken up in 50 mL of ethyl acetate and 50 mL of water. The water layer is extracted with ethyl acetate (2x 25mL) and then the combined organic layers washed with brine, dried (MgSO₄) filtered and evaporated *in vacuo* to give the title compound (1.90g, 3.77 mmol 95%). LCMS (ESMS): *m/z* 428.7 (M+H⁺).

6-Bromo-2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid methyl ester:

3,3,6-Tribromo-2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid methyl ester (5.90 g, 16.9 mmol) is suspended in glacial acetic acid (153 mL) and zinc dust (10 μM Mesh, 11.5g, 169 mmol) is added over 5 minutes. The reaction is monitored by TLC and when complete (30 min) the reaction is filtered through celite and the filtrate washed with ethyl acetate. The combined filtrate and washings are evaporated *in vacuo* to yield an oil that is taken up in ethyl acetate

(500mL), washed with water (100 mL) and brine (100 mL). The organic layer is dried filtered and evaporated *in vacuo* to give the title compound that is recrystallized from ethyl acetate (2.2g, 11.5 mmol 68%). LCMS (ESMS): m/z 271.3 (M+H⁺).

6-Bromo-2-*tert*-butoxycarbonyloxy-indole-1,4-dicarboxylic acid 1-*tert*-butyl ester 4-methyl ester:

6-Bromo-2-oxo-2,3-dihydro-1*H*-indole-4-carboxylic acid methyl ester 1g 3.7 mmol and di *tert* butyldicarbonate (1.77g, 8.14 mmol) is dissolved tetrahydrofuran (75 mL) and stirred at room temperature. DMAP (5 mg, 0.37 mmol) is added and the resulting solution stirred for 4 hours. The reaction is the evaporated *in vacuo* under low heat and the resulting solid purified on silica with hexanes ethyl acetate as the eluent to give the title compound (850 mg 1.8 mmol 49 %). LCMS (ESMS): m/z 471.5(M+H⁺).

6-Bromo-2-*tert*-butoxycarbonyloxy-4-hydroxymethyl-indole-1-carboxylic acid *tert*-butyl ester:

6-Bromo-2-*tert*-butoxycarbonyloxy-indole-1,4-dicarboxylic acid 1-*tert*-butyl ester 4-methyl ester (280 mg, 0.59 mmol) is dissolved in tetrahydrofuran 3 mL and then purged with nitrogen and cooled to -78°C (dry ice IPA). DIBAL-H (1M in hexanes 3.8 mL 3.8 mmol) is added to this stirring solution at -78 over the course of 5 minutes. Upon completion (3 hours) the reaction is quenched by the slow addition of saturated Rochelle's salt (50 mL) and the resulting slurry warmed to room temperature and stirred overnight. The product is extracted with dichloromethane (3x 50 mL), dried (MgSO₄) filtered and evaporated *in vacuo*. Product purified on silica gel with dichloromethane methanol as the eluent (125 mg, 0.28 mmol 48%). LCMS (ESMS): m/z 443.7 (M+H⁺)..

6-Bromo-2-*tert*-butoxycarbonyloxy-4-[3-(1-methoxycarbonylmethyl-butyl)-2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl]-indole-1-carboxylic acid *tert*-butyl ester:

6-Bromo-2-*tert*-butoxycarbonyloxy-4-hydroxymethyl-indole-1-carboxylic acid *tert*-butyl ester (220 mg, 0.50 mmol) is and 3-(2-Oxo-2,3-dihydro-benzoimidazol-1-yl)-hexanoic acid ethyl ester (151 mg, 0.55 mmol) is taken up in 5 mL of tetrahydrofuran and then triphenyl phosphine is added (157 mg, 0.6 mmol). The reaction is stirred at 0°C and diisopropyl azodicarboxylate (116 □L, 0.6 mmol) is added drop wise to the reaction. The reaction is stirred at room

temperature for 4 hours and then the solvent evaporated *in vacuo*. Product is purified by preparative TLC 3% methanol in dichloromethane to give the title compound (200 mg 0.28 mmol 57 %). LCMS (ESMS): m/z 687.7 ($M+H^+$).

3-[3-(6-Bromo-2-oxo-2,3-dihydro-1*H*-indol-4-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid methyl ester:

6-Bromo-2-*tert*-butoxycarbonyloxy-4-[3-(1-methoxycarbonylmethyl-butyl)-2-oxo-2,3-dihydro-benzoimidazol-1-ylmethyl]-indole-1-carboxylic acid *tert*-butyl ester (200 mg, 0.28 mmol) is dissolved in dichloromethane (5 mL) and then trifluoroacetic acid is added (2 mL). The reaction is stirred for 1 hour then evaporated *in vacuo* to give an oil that is purified by preparative reverse phase LC to give the title compound (63mg 0.13 mmol 44 %). LCMS (ESMS): m/z 487.6 ($M+H^+$).

3-[3-(6-Bromo-2-oxo-2,3-dihydro-1*H*-indol-4-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid:

3-[3-(6-Bromo-2-oxo-2,3-dihydro-1*H*-indol-4-ylmethyl)-2-oxo-2,3-dihydro-benzoimidazol-1-yl]-hexanoic acid methyl ester 80 mg (0.16 mmol) is dissolved in 2.5 mL of dioxane and then lithium hydroxide (10 mg, 0.43 mmol in 1.5 mL of water) is added drop wise to the reaction mixture. The reaction is stirred at room temperature and monitored by LC/MS. Upon completion the reaction is acidified by the addition of 1N HCl, and diluted with 10 mL of water. A solid precipitated that is collected and dried to give the title compound (73 mg, 0.15 mmol 96%). LCMS (ESMS): m/z 473.1 ($M+H^+$).

Example 70: 3-{3-[(5-chloro-2-oxo-2,3-dihydro-1,3-benzoxazol-7-yl)methyl]-2-oxo-2,3-dihydro-1*H*-benzimidazol-1-yl}hexanoic acid

DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME __1__ DE __2__

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**

THIS IS VOLUME __1__ OF __2__

NOTE: For additional volumes please contact the Canadian Patent Office.

25771-1706

CLAIMS:

1. A compound:

N-[(2-fluorophenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide;

(3R)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-phenylpropanoic acid;

N-[(1,2-dimethyl-1H-imidazol-4-yl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide;

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)hexanamide;

(3S)-N-[(1,2-dimethyl-1H-imidazol-4-yl)sulfonyl]-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide;

(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(pyridin-2-ylsulfonyl)hexanamide;

2-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methylpentanoic acid;

3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2-phenylpropanoic acid;

(2S)-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}(phenyl)acetic acid;

N-[(4-fluorophenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide;

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-[(1-methyl-1H-imidazol-4-yl)sulfonyl]hexanamide;

(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-[(1-methyl-1H-imidazol-4-yl)sulfonyl]hexanamide;

(3R)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-phenylpropanoic acid;

(3S)-3-{6-methyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-5-phenylpentanoic acid;

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)hexanamide;

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(pyridin-2-ylsulfonyl)hexanamide;

N-[(2-methoxyphenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide;

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}octanoic acid;

(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)hexanamide;

25771-1706

(3R)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-pyridin-3-ylpropanoic acid;

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}octanoic acid;

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-5,6-dimethyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-ethoxy-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)propanamide;

(3S)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-phenylpropanoic acid;

(3S)-N-(tert-butylsulfonyl)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide;

2-({3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)pentanoic acid;

(3S)-3-{2-oxo-3-[(1,4,6-trimethyl-1H-indol-3-yl)methyl]-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}heptanoic acid;

N-[(3-fluorophenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide;

3-{6-methyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-ethoxy-3-{6-methyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

3-{5,6-dimethyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-ethoxypropanoic acid;

(3S)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-{2-oxo-3-[(1,4,6-trimethyl-1H-indol-3-yl)methyl]-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

N-[(4-methoxyphenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide;

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

N-(tert-butylsulfonyl)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide;

25771-1706

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2,2-dimethylhexanoic acid;

1-({3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)cyclohexanecarboxylic acid;

3-ethoxy-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

N-[(3-methoxyphenyl)sulfonyl]-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide;

3-{3-[(6-bromo-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

2-({3-[(4,6-dimethyl-1,2-benzisothiazol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)pentanoic acid;

(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-4-methoxy-N-[(1-methyl-1H-imidazol-4-yl)sulfonyl]butanamide;

3-{3-[(6-chloro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

(3S)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-4-phenylbutanoic acid;

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(pyridin-3-ylsulfonyl)hexanamide;

2-({3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)butanoic acid;

2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentane-1-sulfonic acid;

(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(isopropylsulfonyl)hexanamide;

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-5-(1,3-thiazol-2-ylamino)-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

1-({3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)cyclohexanecarboxylic acid;

25771-1706

N-(cyclopropylsulfonyl)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanamide;

5-methoxy-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-5-oxopentanoic acid;

(3S)-N-cyano-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanamide;

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-ethoxypropanoic acid;

3-{5-bromo-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid;

5-methyl-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid;

(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(ethylsulfonyl)hexanamide;

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)propanamide;

(3S)-N-(cyclopropylsulfonyl)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanamide;

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} pentanedioic acid;

3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid;

1-(1-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} butyl)cyclopropanecarboxylic acid;

3-{3-[(3-bromo-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid;

3-{3-[(1-cyano-5,7-dimethylindolizin-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} hexanoic acid;

3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} propanoic acid;

N-[(2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} ethyl)carbamoyl]benzenesulfonamide;

3-methoxy-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl} propanoic acid;

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(methylsulfonyl)hexanamide;

25771-1706

- (1-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclohexyl)acetic acid;
- 3-{3-[(5-chloro-1,3-dimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;
- (3R)-3-{6-methyl-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-5-phenylpentanoic acid;
- 3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(ethylsulfonyl)hexanamide;
- (3S)-3-{3-[(4,6-dimethyl-1,2-benzisothiazol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;
- N-[(2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}ethyl)sulfonyl]benzamide;
- (2S)-2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid;
- 3-{3-[(3,8-dimethylindolizin-1-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;
- (3S)-N-(aminosulfonyl)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide;
- 3-ethoxy-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid;
- (3R)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;
- (3R)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;
- 2-({3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)butanoic acid;
- 3-{3-[(4,6-dimethyl-1,2-benzisothiazol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;
- 3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid;
- 1-({3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)cyclopentanecarboxylic acid;
- 2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)acetamide;
- 3-{3-[(1-ethyl-4-methyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

25771-1706

3-{3-[(5-bromo-1-methyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid;

3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-6-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

(3S)-3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(methylsulfonyl)hexanamide;

3-{3-[(6,8-dichloro-3-methylindolizin-1-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

4-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}heptanoic acid;

(2S)-2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid;

1-({3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)cyclopentanecarboxylic acid;

2,2-dimethyl-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

5-hydroxy-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid;

3-{3-[(6-bromo-2-oxo-2,3-dihydro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid;

3-{3-[(6-bromo-1-methyl-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclopent-1-ene-1-carboxylic acid;

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-4-methoxybutanoic acid;

3-{3-[(5-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2-methylpropanoic acid;

(1-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclopentyl)acetic acid;

3-{3-[(1-cyano-5-methylindolizin-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

25771-1706

2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-methoxybenzoic acid;

3-{3-[(5-chloro-1-methyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-{7-bromo-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

5-methyl-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid;

3-{3-[(6-methyl-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

4-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)butanamide;

(2E)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}acrylic acid;

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2,2-dimethylbutanoic acid;

(2S)-3-methyl-2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid;

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-4-methoxy-2,2-dimethylbutanoic acid;

3-[3-(1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid;

3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-4-methoxybenzoic acid;

3-{2-oxo-3-[(1,2,3,5-tetramethyl-1H-indol-7-yl)methyl]-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}benzoic acid;

3-{3-[(8-bromo-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid;

N-(isopropylsulfonyl)-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanamide;

25771-1706

N-[(2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}ethyl)sulfonyl]acetamide;
 3-{3-[(4-methyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;
 (2S)-2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;
 2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-methylbenzoic acid;
 3-{3-[(6-bromo-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;
 3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(methylsulfonyl)propanamide;
 3-{3-[(4-chloro-1-methyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;
 3-{3-[(6-chloro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;
 3-methoxy-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid;
 4-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}pentanoic acid;
 3-{3-[(5-fluoro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;
 4-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}butanoic acid;
 3-[3-(3-hydroxy-2,5-dimethylbenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid;
 {3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}acetic acid;
 2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}ethanesulfonic acid;
 3-{2-oxo-3-[(1,2,3-trimethyl-1H-indol-7-yl)methyl]-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;
 1-[(4-methyl-1-benzothien-3-yl)methyl]-3-[2-(1H-tetrazol-5-yl)ethyl]-1,3-dihydro-2H-benzimidazol-2-one;
 3-{3-[(6-bromo-1-methyl-2-oxo-2,3-dihydro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

25771-1706

- 3-[2-oxo-3-(2,3,5-trichlorobenzyl)-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid;
- 3-{2-oxo-3-[(1,2,3-trimethyl-1H-indol-7-yl)methyl]-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;
- (1R,2R)-2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclohexanecarboxylic acid;
- 3-(6-{[2-(3,4-dimethoxyphenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid;
- 3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-3-methylbutanoic acid;
- 3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-7-methyl-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;
- 3-{3-[(8-bromo-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;
- 3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;
- 2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2-methylpropanoic acid;
- 3-(6-{[2-(3-hydroxy-4-methoxyphenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid;
- 3-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-2-methylpropanoic acid;
- 3-{3-[(6-chloro-1-methyl-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;
- 3-(6-{[2-(2-methoxyphenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid;
- 3-{3-[(4-chloro-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;
- 3-({3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)benzoic acid;
- 2-{3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}cyclopentanecarboxylic acid;
- 1-[2-(1,1-dioxido-4-oxo-1,2,5-thiadiazolidin-2-yl)ethyl]-3-[(4-methyl-1-benzothien-3-yl)methyl]-1,3-dihydro-2H-benzimidazol-2-one;
- 3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]pentanoic acid;

25771-1706

;3-[6-{[2-(3,4-dimethoxyphenyl)ethyl]carbamoyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid;

4-methyl-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]pentanoic acid;

3-[3-(5-bromo-2-methylbenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid;

3-[3-(2,5-dimethylbenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid;

3-{3-[(5-methoxy-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

trans-4-({3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)cyclohexanecarboxylic acid;

3-[3-(2,5-dichlorobenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid;

3-{3-[(6-chloro-2-oxo-2,3-dihydro-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-{3-[(5-bromo-1-methyl-2-oxo-2,3-dihydro-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]-N-(phenylsulfonyl)propanamide;

3-[6-{[2-(3-hydroxy-4-methoxyphenyl)ethyl]carbamoyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid;

(3S)-3-{5-(carbamoylamino)-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

(3S)-3-{5-[(tert-butoxycarbonyl)amino]-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-{6-(benzylcarbamoyl)-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

3-{6-[(2-hydroxyethyl)carbamoyl]-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

3-[3-(5-chloro-2-methylbenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid;

N-methyl-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}-N-(phenylsulfonyl)hexanamide;

3-(6-{[2-(4-methoxyphenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid;

25771-1706

(2E)-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]acrylic acid;

3-[6-{[2-(2-methoxyphenyl)ethyl]carbamoyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid;

3-{6-[(3-hydroxypyrrolidin-1-yl)carbonyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

3-{3-[(4-fluoro-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-(6-{[2-(3-methoxyphenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid;

3-{3-[5-chloro-2-(trifluoromethyl)benzyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-{3-[(4-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

3-{3-(1-naphthylmethyl)-2-oxo-6-[(tetrahydrofuran-2-yl)methyl]carbamoyl]-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

3-{6-[(2-hydroxyethyl)carbamoyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

3-{[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]methyl}benzoic acid;

3-[6-{[2-(1,3-dioxolan-2-yl)ethyl]carbamoyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid;

3-[6-{[2-(hydroxymethyl)piperidin-1-yl]carbonyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid;

3-{5-anilino-3-[(1,4-dimethyl-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-{3-[(1-methyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

3-[6-({3-[acetyl(methyl)amino]pyrrolidin-1-yl}carbonyl)-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid;

3-(6-{[2-(2-fluorophenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid;

3-[6-({[(2S)-2-carbamoylpyrrolidin-1-yl]carbonyl}-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid;

3-{6-[(3-carbamoylpiperidin-1-yl)carbonyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

25771-1706

3-{3-[2-methyl-5-(trifluoromethyl)benzyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-{3-[(1,3-dimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

3-(6-{[2-(2-chlorophenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid;

3-{3-[(5-bromo-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-{6-[(1,1-dioxidotetrahydro-3-thienyl)carbamoyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

3-(6-{[2-(4-chlorophenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid;

3-[6-cyano-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid;

2-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}ethanesulfonamide;

N-(methylsulfonyl)-3-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanamide;

3-{3-[(1,2-dimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

N-benzyl-3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanamide;

3-{3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-6-[(2-phenylethyl)carbamoyl]-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

3-{3-[(1-cyano-5-methylindolizin-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

3-{3-[(5-chloro-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

3-{3-[(4-chloro-1,2,3-trimethyl-1H-indol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-(6-{[2-(3-fluorophenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid;

3-{3-[(4-bromo-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-{3-[(6-bromo-1-methyl-1H-indol-4-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

25771-1706

3-{3-[(6-hydroxy-1-naphthyl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

3-{3-(1-naphthylmethyl)-2-oxo-6-[(2-oxotetrahydrofuran-3-yl)carbamoyl]-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

3-(6-{[2-(3-chlorophenyl)ethyl]carbamoyl}-3-[(4-methyl-1-benzothien-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)propanoic acid;

3-[3-(3,5-dibromobenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid;

3-[3-(3,5-dichlorobenzyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]hexanoic acid;

3-[6-((2S)-2-hydroxy-1-(hydroxymethyl)-2-[4-(methylthio)phenyl]ethyl)carbamoyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid;

3-{3-[(5-chloro-2-oxo-2,3-dihydro-1,3-benzoxazol-7-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

2-({3-[2-bromo-5-(trifluoromethyl)benzyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}methyl)pentanoic acid;

3-{3-[2-bromo-5-(trifluoromethyl)benzyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}hexanoic acid;

3-[6-((2S)-2-(hydroxymethyl)pyrrolidin-1-yl)carbonyl]-3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid;

3-[3-(1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]propanoic acid;

5-[3-(1-naphthylmethyl)-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl]pentanoic acid;

3-{3-[(4-chloro-1H-indol-3-yl)methyl]-2-oxo-2,3-dihydro-1H-benzimidazol-1-yl}propanoic acid;

(R)-3-(4-Chloro-phenyl)-3-[3-(1,4-dimethyl-1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-propionic acid;

3-[3-(1,4-Dimethyl-1H-indol-3-ylmethyl)-2-oxo-2,3-dihydro-benzimidazol-1-yl]-2-hydroxymethyl-propionic acid;

3-[2-Oxo-3-(1,4,6-trimethyl-1H-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-2-pyridin-3-yl-propionic acid;

(R)-3-[6-Cyano-2-oxo-3-(1,4,6-trimethyl-1H-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-phenyl-propionic acid;

(R)-3-[6-Carbamoyl-2-oxo-3-(1,4,6-trimethyl-1H-indol-3-ylmethyl)-2,3-dihydro-

25771-1706

benzimidazol-1-yl]-3-phenyl-propionic acid;

(R)-3-[6-Cyano-2-oxo-3-(1,4,6-trimethyl-1H-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-(4-fluoro-phenyl)-propionic acid; or

(R)-3-[6-Cyano-2-oxo-3-(1,4,5-trimethyl-1H-indol-3-ylmethyl)-2,3-dihydro-benzimidazol-1-yl]-3-(4-fluoro-phenyl)-propionic acid;

or a pharmaceutically acceptable salt of the compound.

2. A pharmaceutical composition comprising a compound as defined in claim 1, or a pharmaceutically acceptable salt thereof, and one or more pharmaceutically acceptable carriers and/or adjuvants.

10 3. Use of a pharmaceutically effective amount of a compound as defined in claim 1, or a pharmaceutically acceptable salt thereof, in the treatment of chronic heart failure (non-ischemic), post-myocardial infarction, heart failure (ischemic), acute myocardial infarction, reperfusion injury, left ventricular dysfunction, cardiac fibrosis, diastolic heart failure, hypertrophic cardiomyopathy, pulmonary hypertension, systolic hypertension,
15 resistant hypertension, coronary artery disease, peripheral arterial occlusive disease, aneurism, stable/unstable angina, restenosis, diabetic nephropathy, atrial fibrillation/ventricular arrhythmias, valvular heart disease, pericardial diseases, chronic kidney disease, end stage renal disease or stroke in a patient.

4. Use of a pharmaceutically effective amount of a compound as defined in
20 claim 1, or a pharmaceutically acceptable salt thereof, in the treatment of a disease or condition in a patient requiring a medical procedure, wherein the disease or condition is coronary artery bypass grafting, percutaneous coronary intervention or stenting.

5. The pharmaceutical composition according to claim 2, for use in the treatment of chronic heart failure (non-ischemic), post-myocardial infarction, heart failure (ischemic),
25 acute myocardial infarction, reperfusion injury, left ventricular dysfunction, cardiac fibrosis, diastolic heart failure, hypertrophic cardiomyopathy, pulmonary hypertension, systolic hypertension, resistant hypertension, coronary artery disease, peripheral arterial occlusive

25771-1706

disease, aneurism, stable/unstable angina, restenosis, diabetic nephropathy, atrial fibrillation/ventricular arrhythmias, valvular heart disease, pericardial diseases, chronic kidney disease, end stage renal disease or stroke in a patient.

6. The pharmaceutical composition according to claim 2, for use in the treatment
5 of a disease or condition in a patient requiring a medical procedure, wherein the disease or condition is coronary artery bypass grafting, percutaneous coronary intervention or stenting.