




(86) Date de dépôt PCT/PCT Filing Date: 2014/08/01
(87) Date publication PCT/PCT Publication Date: 2015/02/05
(85) Entrée phase nationale/National Entry: 2016/01/29
(86) N° demande PCT/PCT Application No.: US 2014/049459
(87) N° publication PCT/PCT Publication No.: 2015/017812
(30) Priorité/Priority: 2013/08/02 (US61/861,853)

(51) Cl.Int./Int.Cl. *A61K 31/52* (2006.01),
A61K 31/519 (2006.01), *A61P 35/00* (2006.01)
(71) Demandeur/Applicant:
PHARMACYCLICS LLC, US
(72) Inventeur/Inventor:
BALASUBRAMANIAN, SRIRAM, US
(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : METHODES PERMETTANT DE TRAITER DES TUMEURS SOLIDES
(54) Title: METHODS FOR THE TREATMENT OF SOLID TUMORS

FIG. 1



(57) **Abrégé/Abstract:**

Described herein are combinations of ACK inhibitors and HDAC inhibitors in the treatment of diseases and disorders characterized by the presence or development of solid tumors.

(51) **International Patent Classification:**
A61K 31/52 (2006.01) *A61P 35/00* (2006.01)
A61K 31/519 (2006.01)

(21) **International Application Number:**
PCT/US2014/049459

(22) **International Filing Date:**
1 August 2014 (01.08.2014)

(25) **Filing Language:** English

(26) **Publication Language:** English

(30) **Priority Data:**
61/861,853 2 August 2013 (02.08.2013) US

(71) **Applicant:** PHARMACYCLICS, INC. [US/US]; 995 East Arques Avenue, Sunnyvale, CA 94085 (US).

(72) **Inventor:** BALASUBRAMANIAN, Sriram; 1623 Howard Avenue, San Carlos, CA 94070 (US).

(74) **Agent:** HOSTETLER, Michael, J.; Wilson Sonsini Goodrich & Rosati, 650 Page Mill Road, Palo Alto, CA 94304 (US).

(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,

KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:


- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

(54) **Title:** METHODS FOR THE TREATMENT OF SOLID TUMORS

FIG. 1



(57) **Abstract:** Described herein are combinations of ACK inhibitors and HDAC inhibitors in the treatment of diseases and disorders characterized by the presence or development of solid tumors.

WO 2015/017812 A1


METHODS FOR THE TREATMENT OF SOLID TUMORS

CROSS-REFERENCE

[0001] This application claims the benefit of U.S. Application Serial No. 61/861,853, filed August 2, 2013, which is incorporated herein by reference in its entirety.

SUMMARY OF THE INVENTION

[0002] Disclosed herein, in certain embodiments, are compositions comprising a therapeutically effective amount of an ACK inhibitor compound, a therapeutically effective amount of an HDAC inhibitor compound, and a pharmaceutically acceptable excipient. In some embodiments, the ACK inhibitor compound is a BTK inhibitor. In some embodiments, the BTK inhibitor is an irreversible BTK inhibitor. In some embodiments, the ACK inhibitor is a compound of Formula (A):



Formula (A)

wherein

A is independently selected from N or CR₅;

R₁ is H, L₂-(substituted or unsubstituted alkyl), L₂-(substituted or unsubstituted cycloalkyl), L₂-(substituted or unsubstituted alkenyl), L₂-(substituted or unsubstituted cycloalkenyl), L₂-(substituted or unsubstituted heterocycle), L₂-(substituted or unsubstituted heteroaryl), or L₂-(substituted or unsubstituted aryl), where L₂ is a bond, O, S, -S(=O), -S(=O)₂, C(=O), - (substituted or unsubstituted C₁-C₆ alkyl), or -(substituted or unsubstituted C₂-C₆ alkenyl);

R₂ and R₃ are independently selected from H, lower alkyl and substituted lower alkyl;


R₄ is L₃-X-L₄-G, wherein,

L₃ is optional, and when present is a bond, optionally substituted or unsubstituted alkyl, optionally substituted or unsubstituted cycloalkyl, optionally substituted or unsubstituted alkenyl, optionally substituted or unsubstituted alkynyl;

X is optional, and when present is a bond, O, -C(=O), S, -S(=O), -S(=O)₂, -NH, -NR₉, -NHC(O), -C(O)NH, -NR₉C(O), -C(O)NR₉, -S(=O)₂NH, -NHS(=O)₂, -S(=O)₂NR₉-, -NR₉S(=O)₂-, -OC(O)NH-, -NHC(O)O-, -OC(O)NR₉-, -NR₉C(O)O-, -CH=NO-, -ON=CH-, -NR₁₀C(O)NR₁₀-, heteroaryl, aryl, -NR₁₀C(=NR₁₁)NR₁₀-, -NR₁₀C(=NR₁₁)-, -C(=NR₁₁)NR₁₀-, -OC(=NR₁₁)-, or -C(=NR₁₁)O-;

L₄ is optional, and when present is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycle;

or L₃, X and L₄ taken together form a nitrogen containing heterocyclic ring;



R₆, R₇ and R₈ are independently selected from among H, lower alkyl or substituted lower alkyl, lower heteroalkyl or substituted lower heteroalkyl, substituted or unsubstituted lower cycloalkyl, and substituted or unsubstituted lower heterocycloalkyl;

R₅ is H, halogen, -L₆-(substituted or unsubstituted C₁-C₃ alkyl), -L₆-(substituted or unsubstituted C₂-C₄ alkenyl), -L₆-(substituted or unsubstituted heteroaryl), or -L₆-(substituted or unsubstituted aryl), wherein L₆ is a bond, O, S, -S(=O), S(=O)₂, NH, C(O), -NHC(O)O, -OC(O)NH, -NHC(O), or -C(O)NH;

each R₉ is independently selected from among H, substituted or unsubstituted lower alkyl, and substituted or unsubstituted lower cycloalkyl;


each R₁₀ is independently H, substituted or unsubstituted lower alkyl, or substituted or unsubstituted lower cycloalkyl; or

two R₁₀ groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or

R₉ and R₁₀ can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or


each R₁₁ is independently selected from H, -S(=O)₂R₈, -S(=O)₂NH₂, -C(O)R₈, -CN, -NO₂, heteroaryl, or heteroalkyl.

In some embodiments, the ACK inhibitor is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib).



Ibrutinib.

[0003] In some embodiments, the ACK inhibitor is AVL-263 (Avila Therapeutics/Celgene Corporation), AVL-292 (Avila Therapeutics/Celgene Corporation), AVL-291 (Avila Therapeutics/Celgene Corporation), BMS-488516 (Bristol-Myers Squibb), BMS-509744 (Bristol-Myers Squibb), CGI-1746 (CGI Pharma/Gilead Sciences), CTA-056, GDC-0834 (Genentech), HY-11066 (also, CTK4I7891, HMS3265G21, HMS3265G22, HMS3265H21, HMS3265H22, 439574-61-5, AG-F-54930), ONO-4059 (Ono Pharmaceutical Co., Ltd.), ONO-WG37 (Ono Pharmaceutical Co., Ltd.), PLS-123 (Peking University), RN486 (Hoffmann-La Roche), or HM71224 (Hanmi Pharmaceutical Company Limited). In some embodiments, the HDAC inhibitor is a compound of Formula (B):



Formula (B)

wherein:

R^1 is hydrogen or alkyl;


X is $-O-$, $-NR^2-$, or $-S(O)_n$ where n is 0-2 and R^2 is hydrogen or alkyl;

Y is alkylene optionally substituted with cycloalkyl, optionally substituted phenyl, alkylthio, alkylsulfinyl, alkylsulfonyl, optionally substituted phenylalkylthio, optionally substituted phenylalkylsulfonyl, hydroxy, or optionally substituted phenoxy;

Ar^1 is phenylene or heteroarylene wherein said Ar^1 is optionally substituted with one or two groups independently selected from alkyl, halo, hydroxy, alkoxy, haloalkoxy, or haloalkyl;


R^3 is hydrogen, alkyl, hydroxyalkyl, or optionally substituted phenyl; and

Ar^2 is aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, heteroaralkenyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, or heterocycloalkylalkyl. In some embodiments, the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat)



Abexinostat.


[0004] In some embodiments, the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat)



Abexinostat

and,

the ACK inhibitor is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib)




Ibrutinib.

[0005] In some embodiments, the composition is in the form of a solid dosage form. In some embodiments, the composition is in the form of a capsule. In some embodiments, the composition is in the form of a solution. In some embodiments, the composition is in the form of a solution for intravenous administration. In some embodiments, the compositions comprise 140 mg of ibrutinib. In some embodiments, the composition is for use in treatment of a solid tumor. In some embodiments, the composition is for use in treatment of a sarcoma or carcinoma. In some embodiments, the composition is for use in treatment of a sarcoma. In some embodiments, the composition is for use in treatment of a carcinoma. In some embodiments, the sarcoma is

selected from alveolar rhabdomyosarcoma; alveolar soft part sarcoma; ameloblastoma; angiosarcoma; chondrosarcoma; chordoma; clear cell sarcoma of soft tissue; dedifferentiated liposarcoma; desmoid; desmoplastic small round cell tumor; embryonal rhabdomyosarcoma; epithelioid fibrosarcoma; epithelioid hemangioendothelioma; epithelioid sarcoma; esthesioneuroblastoma; Ewing sarcoma; extrarenal rhabdoid tumor; extraskeletal myxoid chondrosarcoma; extraskeletal osteosarcoma; fibrosarcoma; giant cell tumor; hemangiopericytoma; infantile fibrosarcoma; inflammatory myofibroblastic tumor; Kaposi sarcoma; leiomyosarcoma of bone; liposarcoma; liposarcoma of bone; malignant fibrous histiocytoma (MFH); malignant fibrous histiocytoma (MFH) of bone; malignant mesenchymoma; malignant peripheral nerve sheath tumor; mesenchymal chondrosarcoma; myxofibrosarcoma; myxoid liposarcoma; myxoinflammatory fibroblastic sarcoma; neoplasms with perivascular epithelioid cell differentiation; osteosarcoma; parosteal osteosarcoma; neoplasm with perivascular epithelioid cell differentiation; periosteal osteosarcoma; pleomorphic liposarcoma; pleomorphic rhabdomyosarcoma; PNET/extraskeletal Ewing tumor; rhabdomyosarcoma; round cell liposarcoma; small cell osteosarcoma; solitary fibrous tumor; synovial sarcoma; telangiectatic osteosarcoma. In some embodiments, the carcinoma is selected from an adenocarcinoma, squamous cell carcinoma, adenosquamous carcinoma, anaplastic carcinoma, large cell carcinoma, or small cell carcinoma. In some embodiments, the carcinoma is selected from anal cancer; appendix cancer; bile duct cancer (i.e., cholangiocarcinoma); bladder cancer; brain tumor; breast cancer; cervical cancer; colon cancer; cancer of Unknown Primary (CUP); esophageal cancer; eye cancer; fallopian tube cancer; kidney cancer; liver cancer; lung cancer; medulloblastoma; melanoma; oral cancer; ovarian cancer; pancreatic cancer; parathyroid disease; penile cancer; pituitary tumor; prostate cancer; rectal cancer; skin cancer; stomach cancer; testicular cancer; throat cancer; thyroid cancer; uterine cancer; vaginal cancer; or vulvar cancer. In some embodiments, the carcinoma is breast cancer. In some embodiments, the breast cancer is invasive ductal carcinoma, ductal carcinoma in situ, invasive lobular carcinoma, or lobular carcinoma in situ. In some embodiments, the carcinoma is pancreatic cancer. In some embodiments, the pancreatic cancer is adenocarcinoma, or islet cell carcinoma. In some embodiments, the carcinoma is colorectal cancer. In some embodiments, the colorectal cancer is adenocarcinoma. In some embodiments, the solid tumor is a colon polyp. In some embodiments, the colon polyp is associated with familial adenomatous polyposis. In some embodiments, the carcinoma is bladder cancer. In some embodiments, the bladder cancer is transitional cell bladder cancer, squamous cell bladder cancer, or adenocarcinoma. In some embodiments, the carcinoma is lung cancer. In some embodiments, the lung cancer is a non-

small cell lung cancer. In some embodiments, the non-small cell lung cancer is adenocarcinoma, squamous-cell lung carcinoma, or large-cell lung carcinoma. In some embodiments, the non-small cell lung cancer is large cel lung cancer. In some embodiments, the lung cancer is a small cell lung cancer. In some embodiments, the carcinoma is prostate cancer. In some embodiments, the prostate cancer is adenocarcinoma or small cell carcinoma. In some embodiments, the carcinoma is ovarian cancer. In some embodiments, the ovarian cancer is epithelial ovarian cancer. In some embodiments, the carcinoma is bile duct cancer. In some embodiments, the bile duct cancer is proximal bile duct carcinoma or distal bile duct carcinoma. In some embodiments, the combination of ibrutinib and abexinostat is 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55% or 60% more efficacious than administration of abexinostat alone. In some embodiments, the combination of the ibrutinib and abexinostat is 50% more efficacious than administration of abexinostat alone. In some embodiments, the combination of ibrutinib and abexinostat is 5%, 10%, 15%, 20%, 25%, 30%, or 35% more efficacious than administration of ibrutinib alone. In some embodiments, the combination of ibrutinib and abexinostat is 25% more efficacious than administration of ibrutinib alone.

[0006] Disclosed herein, in certain embodiments, are combinations of a therapeutically effective amount of an ACK inhibitor compound and a therapeutically effective amount of an HDAC inhibitor compound. In some embodiments, the ACK inhibitor compound is a BTK inhibitor. In some embodiments, the BTK inhibitor is an irreversible BTK inhibitor. In some embodiments, the ACK inhibitor is a compound of Formula (A):



Formula (A)

wherein

A is independently selected from N or CR₅;

R₁ is H, L₂-(substituted or unsubstituted alkyl), L₂-(substituted or unsubstituted cycloalkyl), L₂-(substituted or unsubstituted alkenyl), L₂-(substituted or unsubstituted cycloalkenyl), L₂-(substituted or unsubstituted heterocycle), L₂-(substituted or unsubstituted heteroaryl), or L₂-(substituted or unsubstituted aryl), where L₂ is a bond, O, S, -S(=O), -S(=O)₂, C(=O), - (substituted or unsubstituted C₁-C₆ alkyl), or -(substituted or unsubstituted C₂-C₆ alkenyl);

R₂ and R₃ are independently selected from H, lower alkyl and substituted lower alkyl;


R₄ is L₃-X-L₄-G, wherein,

L₃ is optional, and when present is a bond, optionally substituted or unsubstituted alkyl, optionally substituted or unsubstituted cycloalkyl, optionally substituted or unsubstituted alkenyl, optionally substituted or unsubstituted alkynyl;

X is optional, and when present is a bond, O, -C(=O), S, -S(=O), -S(=O)₂, -NH, -NR₉, -NHC(O), -C(O)NH, -NR₉C(O), -C(O)NR₉, -S(=O)₂NH, -NHS(=O)₂, -S(=O)₂NR₉-, -NR₉S(=O)₂-, -OC(O)NH-, -NHC(O)O-, -OC(O)NR₉-, -NR₉C(O)O-, -CH=NO-, -ON=CH-, -NR₁₀C(O)NR₁₀-, heteroaryl, aryl, -NR₁₀C(=NR₁₁)NR₁₀-, -NR₁₀C(=NR₁₁)-, -C(=NR₁₁)NR₁₀-, -OC(=NR₁₁)-, or -C(=NR₁₁)O-;

L₄ is optional, and when present is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycle;

or L₃, X and L₄ taken together form a nitrogen containing heterocyclic ring;



R₆, R₇ and R₈ are independently selected from among H, lower alkyl or substituted lower alkyl, lower heteroalkyl or substituted lower heteroalkyl, substituted or unsubstituted lower cycloalkyl, and substituted or unsubstituted lower heterocycloalkyl;

R₅ is H, halogen, -L₆-(substituted or unsubstituted C₁-C₃ alkyl), -L₆-(substituted or unsubstituted C₂-C₄ alkenyl), -L₆-(substituted or unsubstituted heteroaryl), or -L₆-(substituted or unsubstituted aryl), wherein L₆ is a bond, O, S, -S(=O), S(=O)₂, NH, C(O), -NHC(O)O, -OC(O)NH, -NHC(O), or -C(O)NH;

each R₉ is independently selected from among H, substituted or unsubstituted lower alkyl, and substituted or unsubstituted lower cycloalkyl;


each R₁₀ is independently H, substituted or unsubstituted lower alkyl, or substituted or unsubstituted lower cycloalkyl; or

two R₁₀ groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or

R₉ and R₁₀ can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or


each R₁₁ is independently selected from H, -S(=O)₂R₈, -S(=O)₂NH₂, -C(O)R₈, -CN, -NO₂, heteroaryl, or heteroalkyl.

In some embodiments, the ACK inhibitor is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib).



Ibrutinib.

[0007] In some embodiments, the ACK inhibitor is AVL-263 (Avila Therapeutics/Celgene Corporation), AVL-292 (Avila Therapeutics/Celgene Corporation), AVL-291 (Avila Therapeutics/Celgene Corporation), BMS-488516 (Bristol-Myers Squibb), BMS-509744 (Bristol-Myers Squibb), CGI-1746 (CGI Pharma/Gilead Sciences), CTA-056, GDC-0834 (Genentech), HY-11066 (also, CTK4I7891, HMS3265G21, HMS3265G22, HMS3265H21, HMS3265H22, 439574-61-5, AG-F-54930), ONO-4059 (Ono Pharmaceutical Co., Ltd.), ONO-WG37 (Ono Pharmaceutical Co., Ltd.), PLS-123 (Peking University), RN486 (Hoffmann-La Roche), or HM71224 (Hanmi Pharmaceutical Company Limited). In some embodiments, the HDAC inhibitor is a compound of Formula (B):



Formula (B)

wherein:

R^1 is hydrogen or alkyl;


X is $-O-$, $-NR^2-$, or $-S(O)_n$ where n is 0-2 and R^2 is hydrogen or alkyl;

Y is alkylene optionally substituted with cycloalkyl, optionally substituted phenyl, alkylthio, alkylsulfinyl, alkylsulfonyl, optionally substituted phenylalkylthio, optionally substituted phenylalkylsulfonyl, hydroxy, or optionally substituted phenoxy;

Ar^1 is phenylene or heteroarylene wherein said Ar^1 is optionally substituted with one or two groups independently selected from alkyl, halo, hydroxy, alkoxy, haloalkoxy, or haloalkyl;


R^3 is hydrogen, alkyl, hydroxyalkyl, or optionally substituted phenyl; and

Ar² is aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, heteroaralkenyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, or heterocycloalkylalkyl. In some embodiments, the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat)



Abexinostat.


[0008] In some embodiments, the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat)



Abexinostat

and,

the ACK inhibitor is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib)




Ibrutinib.

[0009] In some embodiments the combination is administered in a solid dosage form. In some embodiments the combination is administered in a capsule. In some embodiments the combination is administered in a solution. In some embodiments the combination is administered in a solution for intravenous administration. In some embodiments the combination, comprises 140 mg of ibrutinib. In some embodiments, the combination is for use in treatment of a solid tumor. In some embodiments, the combination is for use in treatment of a sarcoma or carcinoma. In some embodiments, the combination is for use in treatment of a sarcoma. In some embodiments, the combination is for use in treatment of a carcinoma. In some embodiments, the sarcoma is selected from alveolar rhabdomyosarcoma; alveolar soft part sarcoma; ameloblastoma; angiosarcoma; chondrosarcoma; chordoma; clear cell sarcoma of soft tissue; dedifferentiated liposarcoma; desmoid; desmoplastic small round cell tumor; embryonal rhabdomyosarcoma; epithelioid fibrosarcoma; epithelioid hemangioendothelioma; epithelioid sarcoma; esthesioneuroblastoma; Ewing sarcoma; extrarenal rhabdoid tumor; extraskeletal myxoid chondrosarcoma; extraskeletal osteosarcoma; fibrosarcoma; giant cell tumor; hemangiopericytoma; infantile fibrosarcoma; inflammatory myofibroblastic tumor; Kaposi sarcoma; leiomyosarcoma of bone; liposarcoma; liposarcoma of bone; malignant fibrous histiocyte (MFH); malignant fibrous histiocyte (MFH) of bone; malignant mesenchymoma; malignant peripheral nerve sheath tumor; mesenchymal chondrosarcoma; myxofibrosarcoma; myxoid liposarcoma; myxoinflammatory fibroblastic sarcoma; neoplasms with perivascular epithelioid cell differentiation; osteosarcoma; parosteal osteosarcoma; neoplasm with perivascular epithelioid cell differentiation; periosteal osteosarcoma; pleomorphic liposarcoma; pleomorphic rhabdomyosarcoma; PNET/extraskeletal Ewing tumor; rhabdomyosarcoma; round cell liposarcoma; small cell osteosarcoma; solitary fibrous tumor;

synovial sarcoma; telangiectatic osteosarcoma. In some embodiments, the carcinoma is selected from an adenocarcinoma, squamous cell carcinoma, adenosquamous carcinoma, anaplastic carcinoma, large cell carcinoma, or small cell carcinoma. In some embodiments, the carcinoma is selected from anal cancer; appendix cancer; bile duct cancer (i.e., cholangiocarcinoma); bladder cancer; brain tumor; breast cancer; cervical cancer; colon cancer; cancer of Unknown Primary (CUP); esophageal cancer; eye cancer; fallopian tube cancer; kidney cancer; liver cancer; lung cancer; medulloblastoma; melanoma; oral cancer; ovarian cancer; pancreatic cancer; parathyroid disease; penile cancer; pituitary tumor; prostate cancer; rectal cancer; skin cancer; stomach cancer; testicular cancer; throat cancer; thyroid cancer; uterine cancer; vaginal cancer; or vulvar cancer. In some embodiments, the carcinoma is breast cancer. In some embodiments, the breast cancer is invasive ductal carcinoma, ductal carcinoma in situ, invasive lobular carcinoma, or lobular carcinoma in situ. In some embodiments, the carcinoma is pancreatic cancer. In some embodiments, the pancreatic cancer is adenocarcinoma, or islet cell carcinoma. In some embodiments, the carcinoma is colorectal cancer. In some embodiments, the colorectal cancer is adenocarcinoma. In some embodiments, the solid tumor is a colon polyp. In some embodiments, the colon polyp is associated with familial adenomatous polyposis. In some embodiments, the carcinoma is bladder cancer. In some embodiments, the bladder cancer is transitional cell bladder cancer, squamous cell bladder cancer, or adenocarcinoma. In some embodiments, the carcinoma is lung cancer. In some embodiments, the lung cancer is a non-small cell lung cancer. In some embodiments, the non-small cell lung cancer is adenocarcinoma, squamous-cell lung carcinoma, or large-cell lung carcinoma. In some embodiments, the non-small cell lung cancer is large cel lung cancer. In some embodiments, the lung cancer is a small cell lung cancer. In some embodiments, the carcinoma is prostate cancer. In some embodiments, the prostate cancer is adenocarcinoma or small cell carcinoma. In some embodiments, the carcinoma is ovarian cancer. In some embodiments, the ovarian cancer is epithelial ovarian cancer. In some embodiments, the carcinoma is bile duct cancer. In some embodiments, the bile duct cancer is proximal bile duct carcinoma or distal bile duct carcinoma. In some embodiments, the combination of ibrutinib and abexinostat is 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55% or 60% more efficacious than administration of abexinostat alone. In some embodiments, the combination of the ibrutinib and abexinostat is 50% more efficacious than administration of abexinostat alone. In some embodiments, the combination of ibrutinib and abexinostat is 5%, 10%, 15%, 20%, 25%, 30%, or 35% more efficacious than administration of ibrutinib alone. In some embodiments, the combination of ibrutinib and abexinostat is 25% more efficacious than administration of ibrutinib alone. In some embodiments the combination is


administered in a unified dosage form or separate dosage forms. In some embodiments the combination is administered simultaneously or sequentially.

[0010] Disclosed herein, in certain embodiments, are methods for treating a solid tumor comprising co-administering to an individual in need thereof an ACK inhibitor compound and an HDAC inhibitor compound. In some embodiments, the ACK inhibitor is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib)



Ibrutinib.

[0011] In some embodiments, the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat)




Abexinostat.

[0012] In some embodiments, the solid tumor is a carcinoma. In some embodiments, the carcinoma is breast cancer. In some embodiments, the carcinoma is pancreatic cancer. In some embodiments, the carcinoma is colorectal cancer. In some embodiments, the carcinoma is bladder cancer. In some embodiments, the carcinoma is lung cancer. In some embodiments, the

lung cancer is a non-small cell lung cancer. In some embodiments, the lung cancer is a large cell lung cancer. In some embodiments, the carcinoma is prostate cancer. In some embodiments, the carcinoma is ovarian cancer. In some embodiments, the carcinoma is bile duct cancer. In some embodiments, co-administration of the ACK inhibitor and the HDAC inhibitor is 50% more efficacious than administration of the HDAC inhibitor alone. In some embodiments, co-administration of the ACK inhibitor and the HDAC inhibitor is 25% more efficacious than administration of the ACK inhibitor alone. In some embodiments, the ACK inhibitor and the HDAC inhibitor are administered in a unified dosage form or separate dosage forms. In some embodiments, the ACK inhibitor and the HDAC inhibitor are administered simultaneously or sequentially.

[0013] Disclosed herein, in certain embodiments, are methods for treating a solid tumor comprising co-administering to an individual in need thereof an ACK inhibitor compound and an HDAC inhibitor compound. In some embodiments, the ACK inhibitor compound is a BTK inhibitor. In some embodiments, the BTK inhibitor is an irreversible BTK inhibitor. In some embodiments, the ACK inhibitor is a compound of Formula (A):



Formula (A)

wherein

A is independently selected from N or CR₅;

R₁ is H, L₂-(substituted or unsubstituted alkyl), L₂-(substituted or unsubstituted cycloalkyl), L₂-(substituted or unsubstituted alkenyl), L₂-(substituted or unsubstituted cycloalkenyl), L₂-(substituted or unsubstituted heterocycle), L₂-(substituted or unsubstituted heteroaryl), or L₂-(substituted or unsubstituted aryl), where L₂ is a bond, O, S, -S(=O), -S(=O)₂, C(=O), - (substituted or unsubstituted C₁-C₆ alkyl), or -(substituted or unsubstituted C₂-C₆ alkenyl);

R₂ and R₃ are independently selected from H, lower alkyl and substituted lower alkyl;

R₄ is L₃-X-L₄-G, wherein,


L₃ is optional, and when present is a bond, optionally substituted or unsubstituted alkyl, optionally substituted or unsubstituted cycloalkyl, optionally substituted or unsubstituted alkenyl, optionally substituted or unsubstituted alkynyl;

X is optional, and when present is a bond, O, -C(=O), S, -S(=O), -S(=O)₂, -NH, -NR₉, -NHC(O), -C(O)NH, -NR₉C(O), -C(O)NR₉, -S(=O)₂NH, -NHS(=O)₂, -S(=O)₂NR₉-, -NR₉S(=O)₂, -

OC(O)NH-, -NHC(O)O-, -OC(O)NR₉-, -NR₉C(O)O-, -CH=NO-, -ON=CH-, -NR₁₀C(O)NR₁₀-, heteroaryl, aryl, -NR₁₀C(=NR₁₁)NR₁₀-, -NR₁₀C(=NR₁₁)-, -C(=NR₁₁)NR₁₀-, -OC(=NR₁₁)-, or -C(=NR₁₁)O-;

L₄ is optional, and when present is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycle;

or L₃, X and L₄ taken together form a nitrogen containing heterocyclic ring;



R₆, R₇ and R₈ are independently selected from among H, lower alkyl or substituted lower alkyl, lower heteroalkyl or substituted lower heteroalkyl, substituted or unsubstituted lower cycloalkyl, and substituted or unsubstituted lower heterocycloalkyl;

R₅ is H, halogen, -L₆-(substituted or unsubstituted C₁-C₃ alkyl), -L₆-(substituted or unsubstituted C₂-C₄ alkenyl), -L₆-(substituted or unsubstituted heteroaryl), or -L₆-(substituted or unsubstituted aryl), wherein L₆ is a bond, O, S, -S(=O), S(=O)₂, NH, C(O), -NHC(O)O, -OC(O)NH, -NHC(O), or -C(O)NH;

each R₉ is independently selected from among H, substituted or unsubstituted lower alkyl, and substituted or unsubstituted lower cycloalkyl;


each R₁₀ is independently H, substituted or unsubstituted lower alkyl, or substituted or unsubstituted lower cycloalkyl; or

two R₁₀ groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or

R₉ and R₁₀ can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or


each R₁₁ is independently selected from H, -S(=O)₂R₈, -S(=O)₂NH₂, -C(O)R₈, -CN, -NO₂, heteroaryl, or heteroalkyl.

[0014] In some embodiments, the ACK inhibitor is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib).



Ibrutinib.

[0015] In some embodiments, the ACK inhibitor is AVL-263 (Avila Therapeutics/Celgene Corporation), AVL-292 (Avila Therapeutics/Celgene Corporation), AVL-291 (Avila Therapeutics/Celgene Corporation), BMS-488516 (Bristol-Myers Squibb), BMS-509744 (Bristol-Myers Squibb), CGI-1746 (CGI Pharma/Gilead Sciences), CTA-056, GDC-0834 (Genentech), HY-11066 (also, CTK4I7891, HMS3265G21, HMS3265G22, HMS3265H21, HMS3265H22, 439574-61-5, AG-F-54930), ONO-4059 (Ono Pharmaceutical Co., Ltd.), ONO-WG37 (Ono Pharmaceutical Co., Ltd.), PLS-123 (Peking University), RN486 (Hoffmann-La Roche), or HM71224 (Hanmi Pharmaceutical Company Limited). In some embodiments, the HDAC inhibitor is a compound of Formula (B):



Formula (B)

wherein:

R^1 is hydrogen or alkyl;


X is $-O-$, $-NR^2-$, or $-S(O)_n$ where n is 0-2 and R^2 is hydrogen or alkyl;

Y is alkylene optionally substituted with cycloalkyl, optionally substituted phenyl, alkylthio, alkylsulfinyl, alkylsulfonyl, optionally substituted phenylalkylthio, optionally substituted phenylalkylsulfonyl, hydroxy, or optionally substituted phenoxy;

Ar^1 is phenylene or heteroarylene wherein said Ar^1 is optionally substituted with one or two groups independently selected from alkyl, halo, hydroxy, alkoxy, haloalkoxy, or haloalkyl;

R^3 is hydrogen, alkyl, hydroxyalkyl, or optionally substituted phenyl; and

Ar² is aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, heteroaralkenyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, or heterocycloalkylalkyl. In some embodiments, the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat)




Abexinostat.

[0016] In some embodiments, the solid tumor is a sarcoma or carcinoma. In some embodiments, the solid tumor is a sarcoma. In some embodiments, the solid tumor is a carcinoma. In some embodiments, the sarcoma is selected from alveolar rhabdomyosarcoma; alveolar soft part sarcoma; ameloblastoma; angiosarcoma; chondrosarcoma; chordoma; clear cell sarcoma of soft tissue; dedifferentiated liposarcoma; desmoid; desmoplastic small round cell tumor; embryonal rhabdomyosarcoma; epithelioid fibrosarcoma; epithelioid hemangioendothelioma; epithelioid sarcoma; esthesioneuroblastoma; Ewing sarcoma; extrarenal rhabdoid tumor; extraskeletal myxoid chondrosarcoma; extraskeletal osteosarcoma; fibrosarcoma; giant cell tumor; hemangiopericytoma; infantile fibrosarcoma; inflammatory myofibroblastic tumor; Kaposi sarcoma; leiomyosarcoma of bone; liposarcoma; liposarcoma of bone; malignant fibrous histiocytoma (MFH); malignant fibrous histiocytoma (MFH) of bone; malignant mesenchymoma; malignant peripheral nerve sheath tumor; mesenchymal chondrosarcoma; myxofibrosarcoma; myxoid liposarcoma; myxoinflammatory fibroblastic sarcoma; neoplasms with perivascular epithelioid cell differentiation; osteosarcoma; parosteal osteosarcoma; neoplasm with perivascular epithelioid cell differentiation; periosteal osteosarcoma; pleomorphic liposarcoma; pleomorphic rhabdomyosarcoma; PNET/extraskeletal Ewing tumor; rhabdomyosarcoma; round cell liposarcoma; small cell osteosarcoma; solitary fibrous tumor; synovial sarcoma; telangiectatic osteosarcoma. In some embodiments, the carcinoma is selected from an adenocarcinoma, squamous cell carcinoma, adenosquamous carcinoma, anaplastic carcinoma, large cell carcinoma, or small cell carcinoma. In some embodiments, the carcinoma


is selected from anal cancer; appendix cancer; bile duct cancer (i.e., cholangiocarcinoma); bladder cancer; brain tumor; breast cancer; cervical cancer; colon cancer; cancer of Unknown Primary (CUP); esophageal cancer; eye cancer; fallopian tube cancer; kidney cancer; liver cancer; lung cancer; medulloblastoma; melanoma; oral cancer; ovarian cancer; pancreatic cancer; parathyroid disease; penile cancer; pituitary tumor; prostate cancer; rectal cancer; skin cancer; stomach cancer; testicular cancer; throat cancer; thyroid cancer; uterine cancer; vaginal cancer; or vulvar cancer. In some embodiments, the carcinoma is breast cancer. In some embodiments, the breast cancer is invasive ductal carcinoma, ductal carcinoma in situ, invasive lobular carcinoma, or lobular carcinoma in situ. In some embodiments, the carcinoma is pancreatic cancer. In some embodiments, the pancreatic cancer is adenocarcinoma, or islet cell carcinoma. In some embodiments, the carcinoma is colorectal cancer. In some embodiments, the colorectal cancer is adenocarcinoma. In some embodiments, the solid tumor is a colon polyp. In some embodiments, the colon polyp is associated with familial adenomatous polyposis. In some embodiments, the carcinoma is bladder cancer. In some embodiments, the bladder cancer is transitional cell bladder cancer, squamous cell bladder cancer, or adenocarcinoma. In some embodiments, the carcinoma is lung cancer. In some embodiments, the lung cancer is a non-small cell lung cancer. In some embodiments, the non-small cell lung cancer is adenocarcinoma, squamous-cell lung carcinoma, or large-cell lung carcinoma. In some embodiments, the lung cancer is a small cell lung cancer. In some embodiments, the carcinoma is prostate cancer. In some embodiments, the prostate cancer is adenocarcinoma or small cell carcinoma. In some embodiments, the carcinoma is ovarian cancer. In some embodiments, the ovarian cancer is epithelial ovarian cancer. In some embodiments, the carcinoma is bile duct cancer. In some embodiments, the bile duct cancer is proximal bile duct carcinoma or distal bile duct carcinoma. In some embodiments, co-administration of the ACK inhibitor and the HDAC inhibitor is 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55% or 60% more efficacious than administration of the HDAC inhibitor alone. In some embodiments, co-administration of the ACK inhibitor and the HDAC inhibitor is 50% more efficacious than administration of the HDAC inhibitor alone. In some embodiments, co-administration of the ACK inhibitor and the HDAC inhibitor is 5%, 10%, 15%, 20%, 25%, 30%, or 35% more efficacious than administration of the ACK inhibitor alone. In some embodiments, co-administration of the ACK inhibitor and the HDAC inhibitor is 25% more efficacious than administration of the ACK inhibitor alone. In some embodiments, the ACK inhibitor and the HDAC inhibitor are administered in a unified dosage form or separate dosage forms. In some embodiments, the ACK inhibitor and the HDAC inhibitor are administered simultaneously or sequentially.

[0017] Disclosed herein, in certain embodiments, are methods for treating a solid tumor comprising co-administering to an individual in need thereof an HDAC inhibitor compound and (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib).



Ibrutinib.

[0018] In some embodiments, the HDAC inhibitor is a compound of Formula (B):



Formula (B)

wherein:

R¹ is hydrogen or alkyl;

X is -O-, -NR²-, or -S(O)_n where n is 0-2 and R² is hydrogen or alkyl;


Y is alkylene optionally substituted with cycloalkyl, optionally substituted phenyl, alkylthio, alkylsulfinyl, alkylsulfonyl, optionally substituted phenylalkylthio, optionally substituted phenylalkylsulfonyl, hydroxy, or optionally substituted phenoxy;

Ar¹ is phenylene or heteroarylene wherein said Ar¹ is optionally substituted with one or two groups independently selected from alkyl, halo, hydroxy, alkoxy, haloalkoxy, or haloalkyl;

R³ is hydrogen, alkyl, hydroxyalkyl, or optionally substituted phenyl; and

Ar² is aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, heteroaralkenyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, or heterocycloalkylalkyl.

In some embodiments, the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat)




Abexinostat.

[0019] In some embodiments, the solid tumor is a sarcoma or carcinoma. In some embodiments, the solid tumor is a sarcoma. In some embodiments, the solid tumor is a carcinoma. In some embodiments, the sarcoma is selected from alveolar rhabdomyosarcoma; alveolar soft part sarcoma; ameloblastoma; angiosarcoma; chondrosarcoma; chordoma; clear cell sarcoma of soft tissue; dedifferentiated liposarcoma; desmoid; desmoplastic small round cell tumor; embryonal rhabdomyosarcoma; epithelioid fibrosarcoma; epithelioid hemangioendothelioma; epithelioid sarcoma; esthesioneuroblastoma; Ewing sarcoma; extrarenal rhabdoid tumor; extraskelatal myxoid chondrosarcoma; extrasketetal osteosarcoma; fibrosarcoma; giant cell tumor; hemangiopericytoma; infantile fibrosarcoma; inflammatory myofibroblastic tumor; Kaposi sarcoma; leiomyosarcoma of bone; liposarcoma; liposarcoma of bone; malignant fibrous histiocyteoma (MFH); malignant fibrous histiocyteoma (MFH) of bone; malignant mesenchymoma; malignant peripheral nerve sheath tumor; mesenchymal chondrosarcoma; myxofibrosarcoma; myxoid liposarcoma; myxoinflammatory fibroblastic sarcoma; neoplasms with perivascular epitheiod cell differentiation; osteosarcoma; parosteal osteosarcoma; neoplasm with perivascular epitheiod cell differentiation; periosteal osteosarcoma; pleomorphic liposarcoma; pleomorphic rhabdomyosarcoma; PNET/extraskelatal Ewing tumor; rhabdomyosarcoma; round cell liposarcoma; small cell osteosarcoma; solitary fibrous tumor; synovial sarcoma; telangiectatic osteosarcoma. In some embodiments, the carcinoma is selected from an adenocarcinoma, squamous cell carcinoma, adenosquamous carcinoma, anaplastic carcinoma, large cell carcinoma, or small cell carcinoma. In some embodiments, the carcinoma is selected from anal cancer; appendix cancer; bile duct cancer (i.e., cholangiocarcinoma); bladder cancer; brain tumor; breast cancer; cervical cancer; colon cancer; cancer of Unknown Primary (CUP); esophageal cancer; eye cancer; fallopian tube cancer; kidney cancer; liver cancer; lung cancer; medulloblastoma; melanoma; oral cancer; ovarian cancer; pancreatic


cancer; parathyroid disease; penile cancer; pituitary tumor; prostate cancer; rectal cancer; skin cancer; stomach cancer; testicular cancer; throat cancer; thyroid cancer; uterine cancer; vaginal cancer; or vulvar cancer. In some embodiments, the carcinoma is breast cancer. In some embodiments, the breast cancer is invasive ductal carcinoma, ductal carcinoma in situ, invasive lobular carcinoma, or lobular carcinoma in situ. In some embodiments, the carcinoma is pancreatic cancer. In some embodiments, the pancreatic cancer is adenocarcinoma, or islet cell carcinoma. In some embodiments, the carcinoma is colorectal cancer. In some embodiments, the colorectal cancer is adenocarcinoma. In some embodiments, the solid tumor is a colon polyp. In some embodiments, the colon polyp is associated with familial adenomatous polyposis. In some embodiments, the carcinoma is bladder cancer. In some embodiments, the bladder cancer is transitional cell bladder cancer, squamous cell bladder cancer, or adenocarcinoma. In some embodiments, the carcinoma is lung cancer. In some embodiments, the lung cancer is a non-small cell lung cancer. In some embodiments, the non-small cell lung cancer is adenocarcinoma, squamous-cell lung carcinoma, or large-cell lung carcinoma. In some embodiments, the lung cancer is a small cell lung cancer. In some embodiments, the carcinoma is prostate cancer. In some embodiments, the prostate cancer is adenocarcinoma or small cell carcinoma. In some embodiments, the carcinoma is ovarian cancer. In some embodiments, the ovarian cancer is epithelial ovarian cancer. In some embodiments, the carcinoma is bile duct cancer. In some embodiments, the bile duct cancer is proximal bile duct carcinoma or distal bile duct carcinoma. In some embodiments, co-administration of ibrutinib and the HDAC inhibitor is 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55% or 60% more efficacious than administration of the HDAC inhibitor alone. In some embodiments, co-administration of ibrutinib and the HDAC inhibitor is 50% more efficacious than administration of the HDAC inhibitor alone. In some embodiments, co-administration of ibrutinib and the HDAC inhibitor is 5%, 10%, 15%, 20%, 25%, 30%, or 35% more efficacious than administration of ibrutinib alone. In some embodiments, co-administration of ibrutinib and the HDAC inhibitor is 25% more efficacious than administration of ibrutinib alone. In some embodiments, ibrutinib and the HDAC inhibitor are administered in a unified dosage form or separate dosage forms. In some embodiments, ibrutinib and the HDAC inhibitor are administered simultaneously or sequentially.

[0020] Disclosed herein, in certain embodiments, are methods for treating a solid tumor comprising co-administering to an individual in need thereof an ACK inhibitor compound and 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat)



Abexinostat.

[0021] In some embodiments, the ACK inhibitor compound is a BTK inhibitor. In some embodiments, the BTK inhibitor is an irreversible BTK inhibitor. In some embodiments, the ACK inhibitor is a compound of Formula (A):



Formula (A)

wherein

A is independently selected from N or CR₅;

R₁ is H, L₂-(substituted or unsubstituted alkyl), L₂-(substituted or unsubstituted cycloalkyl), L₂-(substituted or unsubstituted alkenyl), L₂-(substituted or unsubstituted cycloalkenyl), L₂-(substituted or unsubstituted heterocycle), L₂-(substituted or unsubstituted heteroaryl), or L₂-(substituted or unsubstituted aryl), where L₂ is a bond, O, S, -S(=O), -S(=O)₂, C(=O), -(substituted or unsubstituted C₁-C₆ alkyl), or -(substituted or unsubstituted C₂-C₆ alkenyl);

R₂ and R₃ are independently selected from H, lower alkyl and substituted lower alkyl;

R₄ is L₃-X-L₄-G, wherein,


L₃ is optional, and when present is a bond, optionally substituted or unsubstituted alkyl, optionally substituted or unsubstituted cycloalkyl, optionally substituted or unsubstituted alkenyl, optionally substituted or unsubstituted alkynyl;

X is optional, and when present is a bond, O, -C(=O), S, -S(=O), -S(=O)₂, -NH, -NR₉, -NHC(O), -C(O)NH, -NR₉C(O), -C(O)NR₉, -S(=O)₂NH, -NHS(=O)₂, -S(=O)₂NR₉-, -NR₉S(=O)₂-, -OC(O)NH-, -NHC(O)O-, -OC(O)NR₉-, -NR₉C(O)O-, -CH=NO-, -ON=CH-, -NR₁₀C(O)NR₁₀-,

heteroaryl, aryl, $-\text{NR}_{10}\text{C}(=\text{NR}_{11})\text{NR}_{10}-$, $-\text{NR}_{10}\text{C}(=\text{NR}_{11})-$, $-\text{C}(=\text{NR}_{11})\text{NR}_{10}-$, $-\text{OC}(=\text{NR}_{11})-$, or $-\text{C}(=\text{NR}_{11})\text{O}-$;

L_4 is optional, and when present is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycle;

or L_3 , X and L_4 taken together form a nitrogen containing heterocyclic ring;



R_6 , R_7 and R_8 are independently selected from among H, lower alkyl or substituted lower alkyl, lower heteroalkyl or substituted lower heteroalkyl, substituted or unsubstituted lower cycloalkyl, and substituted or unsubstituted lower heterocycloalkyl;

R_5 is H, halogen, $-\text{L}_6-(\text{substituted or unsubstituted } \text{C}_1-\text{C}_3 \text{ alkyl})$, $-\text{L}_6-(\text{substituted or unsubstituted } \text{C}_2-\text{C}_4 \text{ alkenyl})$, $-\text{L}_6-(\text{substituted or unsubstituted heteroaryl})$, or $-\text{L}_6-(\text{substituted or unsubstituted aryl})$, wherein L_6 is a bond, O, S, $-\text{S}(=\text{O})$, $\text{S}(=\text{O})_2$, NH, C(O), $-\text{NHC}(\text{O})\text{O}$, $-\text{OC}(\text{O})\text{NH}$, $-\text{NHC}(\text{O})$, or $-\text{C}(\text{O})\text{NH}$;


each R_9 is independently selected from among H, substituted or unsubstituted lower alkyl, and substituted or unsubstituted lower cycloalkyl;

each R_{10} is independently H, substituted or unsubstituted lower alkyl, or substituted or unsubstituted lower cycloalkyl; or

two R_{10} groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or

R_9 and R_{10} can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or

each R_{11} is independently selected from H, $-\text{S}(=\text{O})_2\text{R}_8$, $-\text{S}(=\text{O})_2\text{NH}_2$, $-\text{C}(\text{O})\text{R}_8$, $-\text{CN}$, $-\text{NO}_2$, heteroaryl, or heteroalkyl. In some embodiments, the ACK inhibitor is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib)




Ibrutinib.

[0022] In some embodiments, the ACK inhibitor is AVL-263 (Avila Therapeutics/Celgene Corporation), AVL-292 (Avila Therapeutics/Celgene Corporation), AVL-291 (Avila Therapeutics/Celgene Corporation), BMS-488516 (Bristol-Myers Squibb), BMS-509744 (Bristol-Myers Squibb), CGI-1746 (CGI Pharma/Gilead Sciences), CTA-056, GDC-0834 (Genentech), HY-11066 (also, CTK4I7891, HMS3265G21, HMS3265G22, HMS3265H21, HMS3265H22, 439574-61-5, AG-F-54930), ONO-4059 (Ono Pharmaceutical Co., Ltd.), ONO-WG37 (Ono Pharmaceutical Co., Ltd.), PLS-123 (Peking University), RN486 (Hoffmann-La Roche), or HM71224 (Hanmi Pharmaceutical Company Limited). In some embodiments, the solid tumor is a sarcoma or carcinoma. In some embodiments, the solid tumor is a sarcoma. In some embodiments, the solid tumor is a carcinoma. In some embodiments, the sarcoma is selected from alveolar rhabdomyosarcoma; alveolar soft part sarcoma; ameloblastoma; angiosarcoma; chondrosarcoma; chordoma; clear cell sarcoma of soft tissue; dedifferentiated liposarcoma; desmoid; desmoplastic small round cell tumor; embryonal rhabdomyosarcoma; epithelioid fibrosarcoma; epithelioid hemangioendothelioma; epithelioid sarcoma; esthesioneuroblastoma; Ewing sarcoma; extrarenal rhabdoid tumor; extraskeletal myxoid chondrosarcoma; extraskeletal osteosarcoma; fibrosarcoma; giant cell tumor; hemangiopericytoma; infantile fibrosarcoma; inflammatory myofibroblastic tumor; Kaposi sarcoma; leiomyosarcoma of bone; liposarcoma; liposarcoma of bone; malignant fibrous histiocytoma (MFH); malignant fibrous histiocytoma (MFH) of bone; malignant mesenchymoma; malignant peripheral nerve sheath tumor; mesenchymal chondrosarcoma; myxofibrosarcoma; myxoid liposarcoma; myxoinflammatory fibroblastic sarcoma; neoplasms with perivascular epithelioid cell differentiation; osteosarcoma; parosteal osteosarcoma; neoplasm with perivascular epithelioid cell differentiation; periosteal osteosarcoma; pleomorphic

liposarcoma; pleomorphic rhabdomyosarcoma; PNET/extraskeletal Ewing tumor; rhabdomyosarcoma; round cell liposarcoma; small cell osteosarcoma; solitary fibrous tumor; synovial sarcoma; telangiectatic osteosarcoma. In some embodiments, the carcinoma is selected from an adenocarcinoma, squamous cell carcinoma, adenosquamous carcinoma, anaplastic carcinoma, large cell carcinoma, or small cell carcinoma. In some embodiments, the carcinoma is selected from anal cancer; appendix cancer; bile duct cancer (i.e., cholangiocarcinoma); bladder cancer; brain tumor; breast cancer; cervical cancer; colon cancer; cancer of Unknown Primary (CUP); esophageal cancer; eye cancer; fallopian tube cancer; kidney cancer; liver cancer; lung cancer; medulloblastoma; melanoma; oral cancer; ovarian cancer; pancreatic cancer; parathyroid disease; penile cancer; pituitary tumor; prostate cancer; rectal cancer; skin cancer; stomach cancer; testicular cancer; throat cancer; thyroid cancer; uterine cancer; vaginal cancer; or vulvar cancer. In some embodiments, the carcinoma is breast cancer. In some embodiments, the breast cancer is invasive ductal carcinoma, ductal carcinoma in situ, invasive lobular carcinoma, or lobular carcinoma in situ. In some embodiments, the carcinoma is pancreatic cancer. In some embodiments, the pancreatic cancer is adenocarcinoma, or islet cell carcinoma. In some embodiments, the carcinoma is colorectal cancer. In some embodiments, the colorectal cancer is adenocarcinoma. In some embodiments, the solid tumor is a colon polyp. In some embodiments, the colon polyp is associated with familial adenomatous polyposis. In some embodiments, the carcinoma is bladder cancer. In some embodiments, the bladder cancer is transitional cell bladder cancer, squamous cell bladder cancer, or adenocarcinoma. In some embodiments, the carcinoma is lung cancer. In some embodiments, the lung cancer is a non-small cell lung cancer. In some embodiments, the non-small cell lung cancer is adenocarcinoma, squamous-cell lung carcinoma, or large-cell lung carcinoma. In some embodiments, the lung cancer is a small cell lung cancer. In some embodiments, the carcinoma is prostate cancer. In some embodiments, the prostate cancer is adenocarcinoma or small cell carcinoma. In some embodiments, the carcinoma is ovarian cancer. In some embodiments, the ovarian cancer is epithelial ovarian cancer. In some embodiments, the carcinoma is bile duct cancer. In some embodiments, the bile duct cancer is proximal bile duct carcinoma or distal bile duct carcinoma. In some embodiments, co-administration of the abexinostat and the ACK inhibitor is 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55% or 60% more efficacious than administration of abexinostat alone. In some embodiments, co-administration of the abexinostat and the ACK inhibitor is 50% more efficacious than administration of abexinostat alone. In some embodiments, co-administration of the abexinostat and the ACK inhibitor is 5%, 10%, 15%, 20%, 25%, 30%, or 35% more efficacious than administration of the ACK inhibitor alone. In


some embodiments, co-administration of the abexinostat and the ACK inhibitor is 25% more efficacious than administration of the ACK inhibitor alone. In some embodiments, the ACK inhibitor and abexinostat are administered in a unified dosage form or separate dosage forms. In some embodiments, the ACK inhibitor and abexinostat are administered simultaneously or sequentially.

[0023] Disclosed herein, in certain embodiments, are methods for treating a solid tumor comprising administering to an individual in need thereof a combination of (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib)



Ibrutinib,

and 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbonyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat)




Abexinostat.

[0024] In some embodiments, the solid tumor is a sarcoma or carcinoma. In some embodiments, the solid tumor is a sarcoma. In some embodiments, the solid tumor is a carcinoma. In some

embodiments, the sarcoma is selected from alveolar rhabdomyosarcoma; alveolar soft part sarcoma; ameloblastoma; angiosarcoma; chondrosarcoma; chordoma; clear cell sarcoma of soft tissue; dedifferentiated liposarcoma; desmoid; desmoplastic small round cell tumor; embryonal rhabdomyosarcoma; epithelioid fibrosarcoma; epithelioid hemangioendothelioma; epithelioid sarcoma; esthesioneuroblastoma; Ewing sarcoma; extrarenal rhabdoid tumor; extrasketetal myxoid chondrosarcoma; extrasketetal osteosarcoma; fibrosarcoma; giant cell tumor; hemangiopericytoma; infantile fibrosarcoma; inflammatory myofibroblastic tumor; Kaposi sarcoma; leiomyosarcoma of bone; liposarcoma; liposarcoma of bone; malignant fibrous histiocytoma (MFH); malignant fibrous histiocytoma (MFH) of bone; malignant mesenchymoma; malignant peripheral nerve sheath tumor; mesenchymal chondrosarcoma; myxofibrosarcoma; myxoid liposarcoma; myxoinflammatory fibroblastic sarcoma; neoplasms with perivascular epithelioid cell differentiation; osteosarcoma; parosteal osteosarcoma; neoplasm with perivascular epithelioid cell differentiation; periosteal osteosarcoma; pleomorphic liposarcoma; pleomorphic rhabdomyosarcoma; PNET/extrasketetal Ewing tumor; rhabdomyosarcoma; round cell liposarcoma; small cell osteosarcoma; solitary fibrous tumor; synovial sarcoma; telangiectatic osteosarcoma. In some embodiments, the carcinoma is selected from an adenocarcinoma, squamous cell carcinoma, adenosquamous carcinoma, anaplastic carcinoma, large cell carcinoma, or small cell carcinoma. In some embodiments, the carcinoma is selected from anal cancer; appendix cancer; bile duct cancer (i.e., cholangiocarcinoma); bladder cancer; brain tumor; breast cancer; cervical cancer; colon cancer; cancer of Unknown Primary (CUP); esophageal cancer; eye cancer; fallopian tube cancer; kidney cancer; liver cancer; lung cancer; medulloblastoma; melanoma; oral cancer; ovarian cancer; pancreatic cancer; parathyroid disease; penile cancer; pituitary tumor; prostate cancer; rectal cancer; skin cancer; stomach cancer; testicular cancer; throat cancer; thyroid cancer; uterine cancer; vaginal cancer; or vulvar cancer. In some embodiments, the carcinoma is breast cancer. In some embodiments, the breast cancer is invasive ductal carcinoma, ductal carcinoma in situ, invasive lobular carcinoma, or lobular carcinoma in situ. In some embodiments, the carcinoma is pancreatic cancer. In some embodiments, the pancreatic cancer is adenocarcinoma, or islet cell carcinoma. In some embodiments, the carcinoma is colorectal cancer. In some embodiments, the colorectal cancer is adenocarcinoma. In some embodiments, the solid tumor is a colon polyp. In some embodiments, the colon polyp is associated with familial adenomatous polyposis. In some embodiments, the carcinoma is bladder cancer. In some embodiments, the bladder cancer is transitional cell bladder cancer, squamous cell bladder cancer, or adenocarcinoma. In some embodiments, the carcinoma is lung cancer. In some embodiments, the lung cancer is a non-

small cell lung cancer. In some embodiments, the non-small cell lung cancer is adenocarcinoma, squamous-cell lung carcinoma, or large-cell lung carcinoma. In some embodiments, the lung cancer is a small cell lung cancer. In some embodiments, the carcinoma is prostate cancer. In some embodiments, the prostate cancer is adenocarcinoma or small cell carcinoma. In some embodiments, the carcinoma is ovarian cancer. In some embodiments, the ovarian cancer is epithelial ovarian cancer. In some embodiments, the carcinoma is bile duct cancer. In some embodiments, the bile duct cancer is proximal bile duct carcinoma or distal bile duct carcinoma. In some embodiments, co-administration of ibrutinib and abexinostat is 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55% or 60% more efficacious than administration of abexinostat alone. In some embodiments, co-administration of the ibrutinib and abexinostat is 50% more efficacious than administration of abexinostat alone. In some embodiments, co-administration of ibrutinib and abexinostat is 5%, 10%, 15%, 20%, 25%, 30%, or 35% more efficacious than administration of ibrutinib alone. In some embodiments, co-administration of ibrutinib and abexinostat is 25% more efficacious than administration of ibrutinib alone. In some embodiments, abexinostat and ibrutinib are administered in a unified dosage form or separate dosage forms. In some embodiments, abexinostat and ibrutinib are administered simultaneously or sequentially.

[0025] Disclosed herein, in certain embodiments, are compositions comprising a therapeutically effective amount of an ACK inhibitor compound, a therapeutically effective amount of an HDAC inhibitor compound, and a pharmaceutically acceptable excipient. In some embodiments, the ACK inhibitor compound is a BTK inhibitor. In some embodiments, the BTK inhibitor is an irreversible BTK inhibitor. In some embodiments, the ACK inhibitor is a compound of Formula (A):



Formula (A)

wherein

A is independently selected from N or CR₅;

R₁ is H, L₂-(substituted or unsubstituted alkyl), L₂-(substituted or unsubstituted cycloalkyl), L₂-(substituted or unsubstituted alkenyl), L₂-(substituted or unsubstituted cycloalkenyl), L₂-(substituted or unsubstituted heterocycle), L₂-(substituted or unsubstituted heteroaryl), or L₂-

(substituted or unsubstituted aryl), where L_2 is a bond, O, S, $-S(=O)$, $-S(=O)_2$, $C(=O)$, - (substituted or unsubstituted C_1-C_6 alkyl), or -(substituted or unsubstituted C_2-C_6 alkenyl); R_2 and R_3 are independently selected from H, lower alkyl and substituted lower alkyl;


R_4 is L_3-X-L_4-G , wherein,

L_3 is optional, and when present is a bond, optionally substituted or unsubstituted alkyl, optionally substituted or unsubstituted cycloalkyl, optionally substituted or unsubstituted alkenyl, optionally substituted or unsubstituted alkynyl;

X is optional, and when present is a bond, O, $-C(=O)$, S, $-S(=O)$, $-S(=O)_2$, $-NH$, $-NR_9$, $-NHC(O)$, $-C(O)NH$, $-NR_9C(O)$, $-C(O)NR_9$, $-S(=O)_2NH$, $-NHS(=O)_2$, $-S(=O)_2NR_9$, $-NR_9S(=O)_2$, $-OC(O)NH$, $-NHC(O)O$, $-OC(O)NR_9$, $-NR_9C(O)O$, $-CH=NO$, $-ON=CH$, $-NR_{10}C(O)NR_{10}$, heteroaryl, aryl, $-NR_{10}C(=NR_{11})NR_{10}$, $-NR_{10}C(=NR_{11})$, $-C(=NR_{11})NR_{10}$, $-OC(=NR_{11})$, or $-C(=NR_{11})O$;

L_4 is optional, and when present is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycle;

or L_3 , X and L_4 taken together form a nitrogen containing heterocyclic ring;



R_6 , R_7 and R_8 are independently selected from among H, lower alkyl or substituted lower alkyl, lower heteroalkyl or substituted lower heteroalkyl, substituted or unsubstituted lower cycloalkyl, and substituted or unsubstituted lower heterocycloalkyl;

R_5 is H, halogen, $-L_6$ -(substituted or unsubstituted C_1-C_3 alkyl), $-L_6$ -(substituted or unsubstituted C_2-C_4 alkenyl), $-L_6$ -(substituted or unsubstituted heteroaryl), or $-L_6$ -(substituted or unsubstituted aryl), wherein L_6 is a bond, O, S, $-S(=O)$, $S(=O)_2$, NH, $C(O)$, $-NHC(O)O$, $-OC(O)NH$, $-NHC(O)$, or $-C(O)NH$;


each R_9 is independently selected from among H, substituted or unsubstituted lower alkyl, and substituted or unsubstituted lower cycloalkyl;

each R_{10} is independently H, substituted or unsubstituted lower alkyl, or substituted or unsubstituted lower cycloalkyl; or

two R_{10} groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or


R_9 and R_{10} can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or

each R_{11} is independently selected from H, $-S(=O)_2R_8$, $-S(=O)_2NH_2$, $-C(O)R_8$, $-CN$, $-NO_2$, heteroaryl, or heteroalkyl. In some embodiments, the ACK inhibitor is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib)



Ibrutinib.

[0026] In some embodiments, the ACK inhibitor is AVL-263 (Avila Therapeutics/Celgene Corporation), AVL-292 (Avila Therapeutics/Celgene Corporation), AVL-291 (Avila Therapeutics/Celgene Corporation), BMS-488516 (Bristol-Myers Squibb), BMS-509744 (Bristol-Myers Squibb), CGI-1746 (CGI Pharma/Gilead Sciences), CTA-056, GDC-0834 (Genentech), HY-11066 (also, CTK4I7891, HMS3265G21, HMS3265G22, HMS3265H21, HMS3265H22, 439574-61-5, AG-F-54930), ONO-4059 (Ono Pharmaceutical Co., Ltd.), ONO-WG37 (Ono Pharmaceutical Co., Ltd.), PLS-123 (Peking University), RN486 (Hoffmann-La Roche), or HM71224 (Hanmi Pharmaceutical Company Limited). In some embodiments, the HDAC inhibitor is a compound of Formula (B):



Formula (B)

wherein:

R^1 is hydrogen or alkyl;


X is $-O-$, $-NR^2-$, or $-S(O)_n$ where n is 0-2 and R^2 is hydrogen or alkyl;

Y is alkylene optionally substituted with cycloalkyl, optionally substituted phenyl, alkylthio, alkylsulfinyl, alkylsulfonyl, optionally substituted phenylalkylthio, optionally substituted phenylalkylsulfonyl, hydroxy, or optionally substituted phenoxy;

Ar¹ is phenylene or heteroarylene wherein said Ar¹ is optionally substituted with one or two groups independently selected from alkyl, halo, hydroxy, alkoxy, haloalkoxy, or haloalkyl;


R³ is hydrogen, alkyl, hydroxyalkyl, or optionally substituted phenyl; and

Ar² is aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, heteroaralkenyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, or heterocycloalkylalkyl. In some embodiments, the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat)




Abexinostat.

[0027] In some embodiments, the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat)



Abexinostat

and, the ACK inhibitor is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib)



Ibrutinib.

[0028] In some embodiments, the composition is in the form of a solid dosage form. In some embodiments, the composition is in the form of a capsule. In some embodiments, the composition is in the form of a solution. In some embodiments, the composition is in the form of a solution for intravenous administration. In some embodiments, the composition comprises 140 mg of ibrutinib.

[0029] Other objects, features and advantages of the methods and compositions described herein will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating specific embodiments, are given by way of illustration only. The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described.

BRIEF DESCRIPTION OF THE FIGURES

[0030] **Figure 1:** Is an illustrative example of the average number of tumors on the lung surface for control and drug-treated (abexinostat, ibrutinib, or abexinostat + ibrutinib) Grg1 transgenic mice. The bars at the left show mice treated for 4 weeks beginning at 2 months (3 month samples) and the bars in the middle show mice treated for 4 weeks beginning at 5 months (6 month samples). The average number of tumors over 1 mm for 6 month samples is also shown.

[0031] **Figure 2:** Is an illustrative example of the number of surface tumors per Grg1 transgenic mice at 3 months after 4 weeks of treatment with abexinostat, ibrutinib, or the combination of abexinostat and ibrutinib.

[0032] **Figure 3:** Is an illustrative example of tumor sections from Grg1 transgenic mice at 3 months. Tumor sections from control mice (A, B), Abexinostat-treated mice (C, D), and

Ibrutinib-treated mice (E, F). Arrows in A, C, E point to tumors. Original magnification of images in the left column was 50X and in the right column was 400X.

[0033] Figure 4: Is an illustrative example of the number of surface tumors per Grg1 transgenic mice at 6 months after 4 weeks of treatment with abexinostat, ibrutinib, or the combination of abexinostat and ibrutinib.

[0034] Figure 5: Is an illustrative example of tumor diameter in tissue sections at 6 months for Grg1 transgenic mice after 4 weeks of treatment with abexinostat, ibrutinib, or the combination of abexinostat and ibrutinib. Each column represents one mouse sample, and the size of each tumor present in the sectioned slides for that mouse sample is given in the column. Each bar in the graph represents the diameter of one tumor and each group of bars represents one mouse. Four mice were analyzed from the control group, and two mice were analyzed for each of the drug-treated groups.

[0035] Figure 6: Is an illustrative example of the number of tumors over 1 mm in 6 month samples from Grg1 transgenic mice after 4 weeks of treatment with abexinostat, ibrutinib, or the combination of abexinostat and ibrutinib. Each number in the table and bar in the graph represents the number of tumors >1 mm for one mouse. The average number of large tumors per mouse for each treatment group is shown in the bottom row of the table and the right-most bar for each group.

[0036] Figure 7: Is an illustrative example of whole left lung lobes from 6 month Grg1 transgenic mice after 4 weeks of treatment with abexinostat, ibrutinib, or the combination of abexinostat and ibrutinib. Three large tumors and one small tumor are visible in the control sample (A); medium and small tumors are visible in Abexinostat-treated (B), Ibrutinib-treated (C) and Abexinostat + Ibrutinib-treated (D) samples.

[0037] Figure 8: Is an illustrative example of whole lung lobes from 3 month Grg1 transgenic mice after 4 weeks of treatment with abexinostat, ibrutinib, or the combination of abexinostat and ibrutinib. Four large tumors are visible in the control sample (A); one small tumor is visible in Abexinostat-treated (B) and Ibrutinib-treated (C) samples, and no visible tumors are seen in the Abexinostat + Ibrutinib-treated (D) samples.

[0038] Figure 9: Is a table exemplifying the effects on tumor diameter in 6 month sections of the combination of abexinostat and ibrutinib. The first bar is the control. The second bar shows the results following treatment with abexinostat alone. The third bar shows the results following treatment with ibrutinib alone. The fourth bar shows the results following treatment with abexinostat and ibrutinib. There were many large tumors in controls compared to treated groups. Interestingly, there were no small tumors in Ib-treated sections, just one large tumor.

[0039] **Figure 10:** Is an illustrative example of lung tissue sections from 6 month Grg1 transgenic mice after 4 weeks of treatment with abexinostat, ibrutinib, or the combination of abexinostat and ibrutinib. In the tissue sections, tumors in control mice exhibit lymphocyte infiltration (L), necrosis (N) and a high number of apoptotic macrophages (Figure 8A and B). Tumors from Abexinostat-treated mice have lymphocyte infiltration, necrosis and large areas with apoptotic cells (Figure 8C and 8D). Tumors from Ibrutinib-treated mice have lymphocyte infiltration but not the apoptotic macrophages seen in the control and in Abexinostat-treated mice (Figure 8E and 8F). Tumors from Abexinostat+Ibrutinib-treated mice have lymphocyte infiltration and macrophage present but not a high number of apoptotic macrophage (Figure 8G and 8H).

[0040] **Figure 11:** Is a table exemplifying the effects of the combination of abexinostat and ibrutinib is MDA-MB-453 cells (breast carcinoma), DLD-1 cells (colorectal adenocarcinoma), NCI-H460 cells (large cell lung cancer), A549 cells (lung carcinoma) and LNCap cells (prostate carcinoma). The combination showed synergy for MDA-MB-453 cells, weak synergy for DLD-1 cells, additivity for NCI-H460 cells, and no additional effect for A549 and LNCaP cells.

[0041] **Figure 12:** Is an illustrative example of the the effects of the combination of abexinostat and ibrutinib in MDA-MB-453 cells (breast carcinoma). 24781 is abexinostat and 32765 is ibrutinib. CI value < 1 indicates synergy, CI value = 1 indicates additive, and CI value > 1 indicate antagonist.

[0042] **Figure 13:** Is an illustrative example of the the effects of the combination of abexinostat and ibrutinib in DLD-1 cells (colorectal adenocarcinoma). 24781 is abexinostat and 32765 is ibrutinib. CI value < 1 indicates synergy, CI value = 1 indicates additive, and CI value > 1 indicate antagonist.

[0043] **Figure 14:** Is an illustrative example of the the effects of the combination of abexinostat and ibrutinib in NCI-H460 cells (large cell lung cancer). 24781 is abexinostat and 32765 is ibrutinib. CI value < 1 indicates synergy, CI value = 1 indicates additive, and CI value > 1 indicate antagonist.


[0044] **Figure 15:** Is an illustrative example of the the effects of the combination of abexinostat and ibrutinib in A549 cells (lung carcinoma). 24781 is abexinostat and 32765 is ibrutinib.

[0045] **Figure 16:** Is an illustrative example of the the effects of the combination of abexinostat and ibrutinib in LNCaP cells (prostate carcinoma). 24781 is abexinostat and 32765 is ibrutinib.

DETAILED DESCRIPTION OF THE INVENTION


[0046] Described herein methods of treating solid tumors comprising co-administering an ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor,

such as for example ibrutinib) and an HDAC inhibitor compound. In some embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) is a compound of Formula (A). In some embodiments, the ACK inhibitor compound is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib).



Ibrutinib

In some embodiments, the ACK inhibitor is AVL-263 (Avila Therapeutics/Celgene Corporation), AVL-292 (Avila Therapeutics/Celgene Corporation), AVL-291 (Avila Therapeutics/Celgene Corporation), BMS-488516 (Bristol-Myers Squibb), BMS-509744 (Bristol-Myers Squibb), CGI-1746 (CGI Pharma/Gilead Sciences), CTA-056, GDC-0834 (Genentech), HY-11066 (also, CTK4I7891, HMS3265G21, HMS3265G22, HMS3265H21, HMS3265H22, 439574-61-5, AG-F-54930), ONO-4059 (Ono Pharmaceutical Co., Ltd.), ONO-WG37 (Ono Pharmaceutical Co., Ltd.), PLS-123 (Peking University), RN486 (Hoffmann-La Roche), or HM71224 (Hanmi Pharmaceutical Company Limited). In some embodiments, the HDAC inhibitor is a compound of Formula (B). In some embodiments, the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat).



Abexinostat

Certain Terminology

[0047] It is to be understood that the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of any subject matter claimed. In this application, the use of the singular includes the plural unless specifically stated otherwise. It must be noted that, as used in the specification and the appended claims, the singular forms “a,” “an” and “the” include plural referents unless the context clearly dictates otherwise. In this application, the use of “or” means “and/or” unless stated otherwise. Furthermore, use of the term “including” as well as other forms, such as “include”, “includes,” and “included,” is not limiting.

[0048] As used herein, “ACK” and “Accessible Cysteine Kinase” are synonyms. They mean a kinase with an accessible cysteine residue. ACKs include, but are not limited to, BTK, ITK, Bmx/ETK, TEC, EFGR, HER4, HER4, LCK, BLK, C-src, FGR, Fyn, HCK, Lyn, YES, ABL, Brk, CSK, FER, JAK3, SYK. In some embodiments, the ACK is HER4.

[0049] As used herein, “amelioration” refers to any lessening of severity, delay in onset, slowing of growth, slowing of metastasis, or shortening of duration of a solid tumor, whether permanent or temporary, lasting or transient that can be attributed to or associated with administration of the compound or composition.

[0050] The term “Bruton’s tyrosine kinase,” as used herein, refers to Bruton’s tyrosine kinase from *Homo sapiens*, as disclosed in, e.g., U.S. Patent No. 6,326,469 (GenBank Accession No. NP_000052).

[0051] The term “Bruton’s tyrosine kinase homolog,” as used herein, refers to orthologs of Bruton’s tyrosine kinase, e.g., the orthologs from mouse (GenBank Accession No. AAB47246), dog (GenBank Accession No. XP_549139.), rat (GenBank Accession No. NP_001007799), chicken (GenBank Accession No. NP_989564), or zebra fish (GenBank Accession No.

XP_698117), and fusion proteins of any of the foregoing that exhibit kinase activity towards one or more substrates of Bruton's tyrosine kinase (e.g. a peptide substrate having the amino acid sequence "AVLESEEELYSSARQ").

[0052] The term "HER4", also known as ERBB4, also known as "V-erb-a erythroblastic leukemia viral oncogene homolog 4" means either (a) the nucleic acid sequence encoding a receptor tyrosine kinase that is a member of the epidermal growth factor receptor subfamily, or (b) the protein thereof. For the nucleic acid sequence that comprises the human HER4 gene *see* GenBank Accession No. NM_001042599. For the amino acid sequence that comprises the human HER4 protein *see* GenBank Accession No. NP_001036064.

[0053] The terms "co-administration" or the like, as used herein, encompass administration of an ACK inhibitor compound and an HDAC inhibitor compound to a single patient, and are intended to include treatment regimens in which the agents are administered by the same or different route of administration, in the same or a different dosage form, and at the same or different time.

[0054] The term "homologous cysteine," as used herein refers to a cysteine residue found within a sequence position that is homologous to that of cysteine 481 of Bruton's tyrosine kinase, as defined herein. For example, cysteine 482 is the homologous cysteine of the rat ortholog of Bruton's tyrosine kinase; cysteine 479 is the homologous cysteine of the chicken ortholog; and cysteine 481 is the homologous cysteine in the zebra fish ortholog. In another example, the homologous cysteine of TXK, a Tec kinase family member related to Bruton's tyrosine, is Cys 350.

[0055] The term "irreversible Btk inhibitor," as used herein, refers to an inhibitor of Btk that can form a covalent bond with an amino acid residue of Btk. In one embodiment, the irreversible inhibitor of Btk can form a covalent bond with a Cys residue of Btk; in particular embodiments, the irreversible inhibitor can form a covalent bond with a Cys 481 residue (or a homolog thereof) of Btk or a cysteine residue in the homologous corresponding position of another tyrosine kinase, as shown in Fig. 7.

[0056] As used herein, the term "pERK" refers to phosphorylated ERK1 and ERK2 at Thr202/Tyr 204 as detected by commercially available phospho-specific antibodies (e.g. Cell Signaling Technologies #4377).

[0057] The terms "individual", "patient" and "subject" are used interchangeable. They refer to a mammal (e.g., a human) which is the object of treatment, or observation. The term is not to be construed as requiring the supervision of a medical practitioner (e.g., a physician, physician's assistant, nurse, orderly, hospice care worker).

[0058] The terms “treat,” “treating” or “treatment”, as used herein, include lessening of severity of a solid tumor, delay in onset of a solid tumor, slowing the growth of a solid tumor, slowing metastasis of cells of a solid tumor, shortening of duration of a solid tumor, arresting the development of o a solid tumor, causing regression of a solid tumor, relieving a condition caused by of a solid tumor, or stopping symptoms which result from a solid tumor. The terms “treat,” “treating” or “treatment”, include, but are not limited to, prophylactic and/or therapeutic treatments.

Therapeutic Methods

[0059] Described herein methods of treating solid tumors comprising co-administering an ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and an HDAC inhibitor compound. In some embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) is a compound of Formula (A). In some embodiments, the ACK inhibitor compound is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib). In some embodiments, the ACK inhibitor is AVL-263 (Avila Therapeutics/Celgene Corporation), AVL-292 (Avila Therapeutics/Celgene Corporation), AVL-291 (Avila Therapeutics/Celgene Corporation), BMS-488516 (Bristol-Myers Squibb), BMS-509744 (Bristol-Myers Squibb), CGI-1746 (CGI Pharma/Gilead Sciences), CTA-056, GDC-0834 (Genentech), HY-11066 (also, CTK4I7891, HMS3265G21, HMS3265G22, HMS3265H21, HMS3265H22, 439574-61-5, AG-F-54930), ONO-4059 (Ono Pharmaceutical Co., Ltd.), ONO-WG37 (Ono Pharmaceutical Co., Ltd.), PLS-123 (Peking University), RN486 (Hoffmann-La Roche), or HM71224 (Hanmi Pharmaceutical Company Limited). In some embodiments, the HDAC inhibitor is a compound of Formula (B). In some embodiments, the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2- carboxamide (i.e. PCI-24781/abexinostat).

Solid Tumors

[0060] As used herein, a “solid tumor” is an abnormal mass of tissue resulting from the abnormal growth or division of cells (i.e., neoplasia). Solid tumors are characterized by an absence of liquid areas. In some embodiments, the solid tumor is benign. In some embodiments, the solid tumor is malignant (i.e., a cancer).

[0061] In some embodiments, the solid tumor is a sarcoma, or carcinoma.

[0062] In some embodiments, the solid tumor is a sarcoma. Sarcomas are cancers of the bone, cartilage, fat, muscle, blood vessels, or other connective or supportive tissue. Sarcomas include,

but are not limited to, alveolar rhabdomyosarcoma; alveolar soft part sarcoma; ameloblastoma; angiosarcoma; chondrosarcoma; chordoma; clear cell sarcoma of soft tissue; dedifferentiated liposarcoma; desmoid; desmoplastic small round cell tumor; embryonal rhabdomyosarcoma; epithelioid fibrosarcoma; epithelioid hemangioendothelioma; epithelioid sarcoma; esthesioneuroblastoma; Ewing sarcoma; extrarenal rhabdoid tumor; extraskeletal myxoid chondrosarcoma; extraskeletal osteosarcoma; fibrosarcoma; giant cell tumor; hemangiopericytoma; infantile fibrosarcoma; inflammatory myofibroblastic tumor; Kaposi sarcoma; leiomyosarcoma of bone; liposarcoma; liposarcoma of bone; malignant fibrous histiocytoma (MFH); malignant fibrous histiocytoma (MFH) of bone; malignant mesenchymoma; malignant peripheral nerve sheath tumor; mesenchymal chondrosarcoma; myxofibrosarcoma; myxoid liposarcoma; myxoinflammatory fibroblastic sarcoma; neoplasms with perivascular epithelioid cell differentiation; osteosarcoma; parosteal osteosarcoma; neoplasm with perivascular epithelioid cell differentiation; periosteal osteosarcoma; pleomorphic liposarcoma; pleomorphic rhabdomyosarcoma; PNET/extraskeletal Ewing tumor; rhabdomyosarcoma; round cell liposarcoma; small cell osteosarcoma; solitary fibrous tumor; synovial sarcoma; telangiectatic osteosarcoma.

[0063] In some embodiments, the solid tumor is a carcinoma. Carcinomas are cancers that begin in the epithelial cells. Carcinomas are classified as adenocarcinoma, squamous cell carcinoma, adenosquamous carcinoma, anaplastic carcinoma, large cell carcinoma, small cell carcinoma. By way of non-limiting example, carcinomas include, anal cancer; appendix cancer; bile duct cancer (i.e., cholangiocarcinoma); bladder cancer; brain tumor; breast cancer; cervical cancer; colon cancer; cancer of Unknown Primary (CUP); esophageal cancer; eye cancer; fallopian tube cancer; kidney cancer; liver cancer; lung cancer; medulloblastoma; melanoma; oral cancer; ovarian cancer; pancreatic cancer; parathyroid disease; penile cancer; pituitary tumor; prostate cancer; rectal cancer; skin cancer (e.g., basal cell carcinoma, squamous cell carcinoma, malignant melanoma); stomach cancer; testicular cancer; throat cancer; thyroid cancer; uterine cancer; vaginal cancer; or vulvar cancer.

[0064] In some embodiments, the solid tumor is breast cancer. In some embodiments, the breast cancer is invasive ductal carcinoma (e.g., triple-negative breast cancer), ductal carcinoma in situ, invasive lobular carcinoma, or lobular carcinoma in situ.

[0065] In some embodiments, the solid tumor is pancreatic cancer. In some embodiments, the pancreatic cancer is adenocarcinoma, or islet cell carcinoma.

[0066] In some embodiments, the solid tumor is colorectal cancer. In some embodiments, the colorectal cancer is adenocarcinoma. In some embodiments, the solid tumor is a colon polyp, for example associated with familial adenomatous polyposis.

[0067] In some embodiments, the solid tumor is bladder cancer. In some embodiments, the bladder cancer is transitional cell bladder cancer, squamous cell bladder cancer, or adenocarcinoma.

[0068] In some embodiments, the solid tumor is lung cancer. In some embodiments, the lung cancer is a non-small cell lung cancer. In some embodiments, the lung cancer is a small cell lung cancer. In some embodiments, the non-small cell lung cancer is adenocarcinoma, squamous-cell lung carcinoma, or large-cell lung carcinoma. In some embodiments, the lung cancer is large cell lung cancer.

[0069] In some embodiments, the solid tumor is prostate cancer. In some embodiments, the prostate cancer is adenocarcinoma or small cell carcinoma.

[0070] In some embodiments, the solid tumor is ovarian cancer (e.g., epithelial ovarian cancer).

[0071] In some embodiments, the solid tumor is bile duct cancer. In some embodiments, the bile duct cancer is proximal bile duct carcinoma or distal bile duct carcinoma.

Synergistic Effects

[0072] Described herein methods of treating solid tumors comprising co-administering an ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and an HDAC inhibitor compound. In some embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) is a compound of Formula (A). In some embodiments, the ACK inhibitor compound is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib). In some embodiments, the ACK inhibitor is AVL-263 (Avila Therapeutics/Celgene Corporation), AVL-292 (Avila Therapeutics/Celgene Corporation), AVL-291 (Avila Therapeutics/Celgene Corporation), BMS-488516 (Bristol-Myers Squibb), BMS-509744 (Bristol-Myers Squibb), CGI-1746 (CGI Pharma/Gilead Sciences), CTA-056, GDC-0834 (Genentech), HY-11066 (also, CTK4I7891, HMS3265G21, HMS3265G22, HMS3265H21, HMS3265H22, 439574-61-5, AG-F-54930), ONO-4059 (Ono Pharmaceutical Co., Ltd.), ONO-WG37 (Ono Pharmaceutical Co., Ltd.), PLS-123 (Peking University), RN486 (Hoffmann-La Roche), or HM71224 (Hanmi Pharmaceutical Company Limited). In some embodiments, the HDAC inhibitor is a compound of Formula (B). In some embodiments, the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-

[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2- carboxamide (i.e. PCI-24781/abexinostat).

[0073] In some embodiments, the co-administration of and ACK inhibitor (e.g., ibrutinib) and an HDAC inhibitor (e.g., abexinostat) results in a synergistic effect for the treatment of a solid tumor (e.g., reduction in the number of tumors, growth/size of tumors, metastasis of tumors)

[0074] In some embodiments, co-administration of an ACK inhibitor (e.g., ibrutinib) and an HDAC inhibitor (e.g., abexinostat) is 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55% or 60% more efficacious than administration of an HDAC inhibitor (e.g., abexinostat) alone. In some embodiments, co-administration of an ACK inhibitor (e.g., ibrutinib) and an HDAC inhibitor (e.g., abexinostat) is 50% more efficacious than administration of an HDAC inhibitor (e.g., abexinostat) alone. In some embodiments, co-administration of an ACK inhibitor (e.g., ibrutinib) and an HDAC inhibitor (e.g., abexinostat) is 1.1x, 1.2x, 1.3x, 1.4x, 1.5x, or 1.6x more efficacious than administration of an HDAC inhibitor (e.g., abexinostat) alone. In some embodiments, co-administration of an ACK inhibitor (e.g., ibrutinib) and an HDAC inhibitor (e.g., abexinostat) is 1.5x more efficacious than administration of an HDAC inhibitor (e.g., abexinostat) alone.

[0075] In some embodiments, co-administration of an ACK inhibitor (e.g., ibrutinib) and an HDAC inhibitor (e.g., abexinostat) is 5%, 10%, 15%, 20%, 25%, 30%, or 35% more efficacious than administration of an ACK inhibitor (e.g., ibrutinib) alone. In some embodiments, co-administration of an ACK inhibitor (e.g., ibrutinib) and an HDAC inhibitor (e.g., abexinostat) is 25% more efficacious than administration of an ACK inhibitor (e.g., ibrutinib) alone. In some embodiments, co-administration of an ACK inhibitor (e.g., ibrutinib) and an HDAC inhibitor (e.g., abexinostat) is 1.1x, 1.15x, 1.2x, 1.25x, or 1.3x more efficacious than administration of an ACK inhibitor (e.g., ibrutinib) alone. In some embodiments, co-administration of an ACK inhibitor (e.g., ibrutinib) and an HDAC inhibitor (e.g., abexinostat) is 1.25x more efficacious than administration of an ACK inhibitor (e.g., ibrutinib) alone.

[0076] In some embodiments, co-administration of ibrutinib and abexinostat is 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55% or 60% more efficacious than administration of abexinostat alone. In some embodiments, co-administration of ibrutinib and abexinostat is 50% more efficacious than administration of abexinostat alone. In some embodiments, co-administration of ibrutinib and abexinostat is 1.1x, 1.2x, 1.3x, 1.4x, 1.5x, or 1.6x more efficacious than administration of abexinostat alone. In some embodiments, co-administration of ibrutinib and abexinostat is 1.5x more efficacious than administration of abexinostat alone.

[0077] In some embodiments, co-administration of ibrutinib and abexinostat is 5%, 10%, 15%, 20%, 25%, 30%, or 35% more efficacious than administration of ibrutinib alone. In some embodiments, co-administration of ibrutinib and abexinostat is 25% more efficacious than administration of ibrutinib alone. In some embodiments, co-administration of ibrutinib and abexinostat is 1.1x, 1.15x, 1.2x, 1.25x, or 1.3x more efficacious than administration of ibrutinib alone. In some embodiments, co-administration of ibrutinib and abexinostat is 1.25x more efficacious than administration of ibrutinib alone.

Administration

[0078] Described herein methods of treating solid tumors comprising co-administering an ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and an HDAC inhibitor compound. In some embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) is a compound of Formula (A). In some embodiments, the ACK inhibitor compound is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib). In some embodiments, the ACK inhibitor is AVL-263 (Avila Therapeutics/Celgene Corporation), AVL-292 (Avila Therapeutics/Celgene Corporation), AVL-291 (Avila Therapeutics/Celgene Corporation), BMS-488516 (Bristol-Myers Squibb), BMS-509744 (Bristol-Myers Squibb), CGI-1746 (CGI Pharma/Gilead Sciences), CTA-056, GDC-0834 (Genentech), HY-11066 (also, CTK4I7891, HMS3265G21, HMS3265G22, HMS3265H21, HMS3265H22, 439574-61-5, AG-F-54930), ONO-4059 (Ono Pharmaceutical Co., Ltd.), ONO-WG37 (Ono Pharmaceutical Co., Ltd.), PLS-123 (Peking University), RN486 (Hoffmann-La Roche), or HM71224 (Hanmi Pharmaceutical Company Limited). In some embodiments, the HDAC inhibitor is a compound of Formula (B). In some embodiments, the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat).

[0079] In some embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example, an irreversible BTK inhibitor, such as for example, ibrutinib) and the HDAC inhibitor are administered in the same composition. In some embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and the HDAC inhibitor are not administered in the same composition. In some embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and the HDAC inhibitor are administered by different routes. In some embodiments, the ACK inhibitor compound (e.g., a

BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and the HDAC inhibitor are administered simultaneously or sequentially.

[0080] If simultaneously, the multiple therapeutic agents are optionally provided in a single, unified form, or in multiple forms (by way of example only, either as a single pill or as two separate pills). In some embodiments, one of the therapeutic agents is given in multiple doses, or both are given as multiple doses. If not simultaneous, the timing between the multiple doses is from about more than zero weeks to less than about four weeks.

[0081] In some embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and the HDAC inhibitor are administered in a combined dosage form, or in separate dosage forms intended for substantially simultaneous administration. In some embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and the HDAC inhibitor are administered sequentially, with either therapeutic compound being administered by a regimen calling for two-step administration. In some embodiments, the two-step administration regimen calls for sequential administration of the active agents or spaced-apart administration of the separate active agents. The time period between the multiple administration steps ranges from a few minutes to several hours, depending upon the properties of each pharmaceutical agent, such as potency, solubility, bioavailability, plasma half-life and kinetic profile of the pharmaceutical agent. In some embodiments, circadian variation of the target molecule concentration determines the optimal dose interval.

[0082] The ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and the HDAC inhibitor compounds described herein are administered before, during or after the development of a solid tumor. In some embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and the HDAC inhibitor compounds are used as a prophylactic and are administered continuously to subjects with a propensity to develop a solid tumor. In some embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and the HDAC inhibitor compounds are administered to an individual during or as soon as possible after the development of a solid tumor. In some embodiments, the administration of the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and the HDAC inhibitor compounds is initiated within the first 48 hours of the onset of the symptoms, within the first 6 hours of the onset of the symptoms, or

within 3 hours of the onset of the symptoms. In some embodiments, the initial administration of the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and the HDAC inhibitor compounds is via any route practical, such as, for example, an intravenous injection, a bolus injection, infusion over 5 minutes to about 5 hours, a pill, a capsule, transdermal patch, buccal delivery, and the like, or combination thereof. The ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and the HDAC inhibitor compounds should be administered as soon as is practicable after the onset of a disorder is detected or suspected, and for a length of time necessary for the treatment of the disease, such as, for example, from about 1 month to about 3 months. The length of treatment can vary for each subject, and the length can be determined using the known criteria. In some embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and the HDAC inhibitor compounds are administered for at least 2 weeks, between about 1 month to about 5 years, or from about 1 month to about 3 years.

[0083] Therapeutically effective amounts will depend on the severity and course of the disorder, previous therapy, the patient's health status, weight, and response to the drugs, and the judgment of the treating physician. Prophylactically effective amounts depend on the patient's state of health, weight, the severity and course of the disease, previous therapy, response to the drugs, and the judgment of the treating physician.

[0084] In some embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and HDAC inhibitor compounds are administered to the patient on a regular basis, e.g., three times a day, two times a day, once a day, every other day or every 3 days. In other embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and HDAC inhibitor compounds are administered to the patient on an intermittent basis, e.g., twice a day followed by once a day followed by three times a day; or the first two days of every week; or the first, second and third day of a week. In some embodiments, intermittent dosing is as effective as regular dosing. In further or alternative embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and HDAC inhibitor compounds are administered only when the patient exhibits a particular symptom, e.g., the onset of pain, or the onset of a fever, or the onset of an inflammation, or the onset of a skin disorder.

Dosing schedules of each compound may depend on the other or may be independent of the other.

[0085] In the case wherein the patient's condition does not improve, upon the doctor's discretion the administration of the compounds may be administered chronically, that is, for an extended period of time, including throughout the duration of the patient's life in order to ameliorate or otherwise control or limit the symptoms of the patient's disorder.

[0086] In the case wherein the patient's status does improve, upon the doctor's discretion the administration of the compounds may be given continuously; alternatively, the dose of drug being administered may be temporarily reduced or temporarily suspended for a certain length of time (i.e., a "drug holiday"). The length of the drug holiday can vary between 2 days and 1 year, including by way of example only, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 10 days, 12 days, 15 days, 20 days, 28 days, 35 days, 50 days, 70 days, 100 days, 120 days, 150 days, 180 days, 200 days, 250 days, 280 days, 300 days, 320 days, 350 days, or 365 days. The dose reduction during a drug holiday may be from 10%-100%, including, by way of example only, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100%.

[0087] Once improvement of the patient's conditions has occurred, a maintenance regimen is administered if necessary. Subsequently, the dosage or the frequency of administration, or both, of one or both of the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and HDAC inhibitor compounds can be reduced, as a function of the symptoms, to a level at which the individual's improved condition is retained. Individuals can, however, require intermittent treatment on a long-term basis upon any recurrence of symptoms.

[0088] The amount of the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and HDAC inhibitor compounds will vary depending upon factors such as the particular compound, disorder and its severity, the identity (e.g., weight) of the subject or host in need of treatment, and is determined according to the particular circumstances surrounding the case, including, e.g., the specific agents being administered, the routes of administration, the solid tumor being treated, and the subject or host being treated. In general, however, doses employed for adult human treatment will typically be in the range of 0.02-5000 mg per day, or from about 1-1500 mg per day of each compound. The desired dose of each compound may be presented in a single dose or as divided doses administered simultaneously (or over a short period of time) or at appropriate intervals, for example as two, three, four or more sub-doses per day.

[0089] In some embodiments, the amount of the ACK inhibitor (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) is from 300 mg/day up to, and including, 1000 mg/day. In some embodiments, the amount of the irreversible Btk inhibitor is from 420 mg/day up to, and including, 840 mg/day. In some embodiments, the amount of the irreversible Btk inhibitor is about 420 mg/day, about 560 mg/day, or about 840 mg/day. In some embodiments, the amount of the irreversible Btk inhibitor is about 420 mg/day.

[0090] In some embodiments, the HDAC inhibitor (e.g., abexinostat) is administered for 4 consecutive days, followed by three consecutive days without administration of the HDAC inhibitor (e.g., abexinostat).

[0091] The ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and HDAC inhibitor compounds described herein may be individually or combined into unit dosage forms suitable for single administration of precise dosages. In unit dosage form, the formulation is divided into unit doses containing appropriate quantities of one or both compounds. The unit dosage may be in the form of a package containing discrete quantities of the formulation. Non-limiting examples are packaged tablets or capsules, and powders in vials or ampoules. Aqueous suspension compositions can be packaged in single-dose non-reclosable containers. Alternatively, multiple-dose reclosable containers can be used, in which case it is typical to include a preservative in the composition. By way of example only, formulations for parenteral injection may be presented in unit dosage form, which include, but are not limited to ampoules, or in multi-dose containers, with an added preservative.

[0092] It is understood that a medical professional will determine the dosage regimen in accordance with a variety of factors. These factors include the solid tumor from which the subject suffers, the degree of metastasis, as well as the age, weight, sex, diet, and medical condition of the subject.

Compounds

[0093] Described herein methods of treating solid tumors comprising co-administering an ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and an HDAC inhibitor compound.

[0094] Definition of standard chemistry terms are found in reference works, including Carey and Sundberg "ADVANCED ORGANIC CHEMISTRY 4TH ED." Vols. A (2000) and B (2001), Plenum Press, New York. Unless otherwise indicated, conventional methods of mass spectroscopy, NMR, HPLC, protein chemistry, biochemistry, recombinant DNA techniques and pharmacology, within the skill of the art are employed. Unless specific definitions are provided,

the nomenclature employed in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those known in the art. Standard techniques are optionally used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients. Standard techniques are optionally used for recombinant DNA, oligonucleotide synthesis, and tissue culture and transformation (e.g., electroporation, lipofection). Reactions and purification techniques are performed using documented methodologies or as described herein.

[0095] It is to be understood that the methods and compositions described herein are not limited to the particular methodology, protocols, cell lines, constructs, and reagents described herein and as such optionally vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the methods and compositions described herein, which will be limited only by the appended claims.

[0096] Unless stated otherwise, the terms used for complex moieties (i.e., multiple chains of moieties) are to be read equivalently either from left to right or right to left. For example, the group alkylencycloalkylene refers both to an alkylene group followed by a cycloalkylene group or as a cycloalkylene group followed by an alkylene group.

[0097] The suffix “ene” appended to a group indicates that such a group is a diradical. By way of example only, a methylene is a diradical of a methyl group, that is, it is a $-\text{CH}_2-$ group; and an ethylene is a diradical of an ethyl group, i.e., $-\text{CH}_2\text{CH}_2-$.

[0098] An “alkyl” group refers to an aliphatic hydrocarbon group. The alkyl moiety includes a “saturated alkyl” group, which means that it does not contain any alkene or alkyne moieties. The alkyl moiety also includes an “unsaturated alkyl” moiety, which means that it contains at least one alkene or alkyne moiety. An “alkene” moiety refers to a group that has at least one carbon-carbon double bond, and an “alkyne” moiety refers to a group that has at least one carbon-carbon triple bond. The alkyl moiety, whether saturated or unsaturated, includes branched, straight chain, or cyclic moieties. Depending on the structure, an alkyl group includes a monoradical or a diradical (i.e., an alkylene group), and if a “lower alkyl” having 1 to 6 carbon atoms.

[0099] As used herein, $\text{C}_1\text{-C}_x$ includes $\text{C}_1\text{-C}_2$, $\text{C}_1\text{-C}_3$. . . $\text{C}_1\text{-C}_x$.

[00100] The “alkyl” moiety optionally has 1 to 10 carbon atoms (whenever it appears herein, a numerical range such as “1 to 10” refers to each integer in the given range; e.g., “1 to 10 carbon atoms” means that the alkyl group is selected from a moiety having 1 carbon atom, 2 carbon

atoms, 3 carbon atoms, *etc.*, up to and including 10 carbon atoms, although the present definition also covers the occurrence of the term “alkyl” where no numerical range is designated). The alkyl group of the compounds described herein may be designated as “C₁-C₄ alkyl” or similar designations. By way of example only, “C₁-C₄ alkyl” indicates that there are one to four carbon atoms in the alkyl chain, *i.e.*, the alkyl chain is selected from among methyl, ethyl, propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, and t-butyl. Thus C₁-C₄ alkyl includes C₁-C₂ alkyl and C₁-C₃ alkyl. Alkyl groups are optionally substituted or unsubstituted. Typical alkyl groups include, but are in no way limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tertiary butyl, pentyl, hexyl, ethenyl, propenyl, butenyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and the like.

[00101] The term “alkenyl” refers to a type of alkyl group in which the first two atoms of the alkyl group form a double bond that is not part of an aromatic group. That is, an alkenyl group begins with the atoms —C(R)=C(R)—R , wherein R refers to the remaining portions of the alkenyl group, which are either the same or different. The alkenyl moiety is optionally branched, straight chain, or cyclic (in which case, it is also known as a “cycloalkenyl” group). Depending on the structure, an alkenyl group includes a monoradical or a diradical (*i.e.*, an alkenylene group). Alkenyl groups are optionally substituted. Non-limiting examples of an alkenyl group include —CH=CH_2 , $\text{—C(CH}_3\text{)=CH}_2$, —CH=CHCH_3 , $\text{—C(CH}_3\text{)=CHCH}_3$. Alkenylene groups include, but are not limited to, —CH=CH— , $\text{—C(CH}_3\text{)=CH—}$, $\text{—CH=CHCH}_2\text{—}$, $\text{—CH=CHCH}_2\text{CH}_2\text{—}$ and $\text{—C(CH}_3\text{)=CHCH}_2\text{—}$. Alkenyl groups optionally have 2 to 10 carbons, and if a “lower alkenyl” having 2 to 6 carbon atoms.

[00102] The term “alkynyl” refers to a type of alkyl group in which the first two atoms of the alkyl group form a triple bond. That is, an alkynyl group begins with the atoms $\text{—C}\equiv\text{C—R}$, wherein R refers to the remaining portions of the alkynyl group, which is either the same or different. The “R” portion of the alkynyl moiety may be branched, straight chain, or cyclic. Depending on the structure, an alkynyl group includes a monoradical or a diradical (*i.e.*, an alkynylene group). Alkynyl groups are optionally substituted. Non-limiting examples of an alkynyl group include, but are not limited to, $\text{—C}\equiv\text{CH}$, $\text{—C}\equiv\text{CCH}_3$, $\text{—C}\equiv\text{CCH}_2\text{CH}_3$, $\text{—C}\equiv\text{C—}$, and $\text{—C}\equiv\text{CCH}_2\text{—}$. Alkynyl groups optionally have 2 to 10 carbons, and if a “lower alkynyl” having 2 to 6 carbon atoms.

[00103] An “alkoxy” group refers to a (alkyl)O- group, where alkyl is as defined herein.

[00104] “Hydroxyalkyl” refers to an alkyl radical, as defined herein, substituted with at least one hydroxy group. Non-limiting examples of a hydroxyalkyl include, but are not limited to, hydroxymethyl, 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, 1-(hydroxymethyl)-2-methylpropyl, 2-hydroxybutyl, 3-hydroxybutyl, 4-hydroxybutyl, 2,3-dihydroxypropyl,

1-(hydroxymethyl)-2-hydroxyethyl, 2,3-dihydroxybutyl, 3,4-dihydroxybutyl and 2-(hydroxymethyl)-3-hydroxypropyl.

[00105] “Alkoxyalkyl” refers to an alkyl radical, as defined herein, substituted with an alkoxy group, as defined herein.

[00106] The term “alkylamine” refers to the $-N(\text{alkyl})_x\text{H}_y$ group, where x and y are selected from among x=1, y=1 and x=2, y=0. When x=2, the alkyl groups, taken together with the N atom to which they are attached, optionally form a cyclic ring system.

[00107] “Alkylaminoalkyl” refers to an alkyl radical, as defined herein, substituted with an alkylamine, as defined herein.

[00108] “Hydroxyalkylaminoalkyl” refers to an alkyl radical, as defined herein, substituted with an alkylamine, and alkylhydroxy, as defined herein.

[00109] “Alkoxyalkylaminoalkyl” refers to an alkyl radical, as defined herein, substituted with an alkylamine and substituted with an alkylalkoxy, as defined herein.

[00110] An “amide” is a chemical moiety with the formula $-C(O)NHR$ or $-NHC(O)R$, where R is selected from among alkyl, cycloalkyl, aryl, heteroaryl (bonded through a ring carbon) and heteroalicyclic (bonded through a ring carbon). In some embodiments, an amide moiety forms a linkage between an amino acid or a peptide molecule and a compound described herein, thereby forming a prodrug. Any amine, or carboxyl side chain on the compounds described herein can be amidified. The procedures and specific groups to make such amides are found in sources such as Greene and Wuts, *Protective Groups in Organic Synthesis*, 3rd Ed., John Wiley & Sons, New York, NY, 1999, which is incorporated herein by reference for this disclosure.

[00111] The term “ester” refers to a chemical moiety with formula $-COOR$, where R is selected from among alkyl, cycloalkyl, aryl, heteroaryl (bonded through a ring carbon) and heteroalicyclic (bonded through a ring carbon). Any hydroxy, or carboxyl side chain on the compounds described herein can be esterified. The procedures and specific groups to make such esters are found in sources such as Greene and Wuts, *Protective Groups in Organic Synthesis*, 3rd Ed., John Wiley & Sons, New York, NY, 1999, which is incorporated herein by reference for this disclosure.

[00112] As used herein, the term “ring” refers to any covalently closed structure. Rings include, for example, carbocycles (e.g., aryls and cycloalkyls), heterocycles (e.g., heteroaryl and non-aromatic heterocycles), aromatics (e.g., aryls and heteroaryl), and non-aromatics (e.g., cycloalkyls and non-aromatic heterocycles). Rings can be optionally substituted. Rings can be monocyclic or polycyclic.

[00113] As used herein, the term “ring system” refers to one, or more than one ring.

[00114] The term “membered ring” can embrace any cyclic structure. The term “membered” is meant to denote the number of skeletal atoms that constitute the ring. Thus, for example, cyclohexyl, pyridine, pyran and thiopyran are 6-membered rings and cyclopentyl, pyrrole, furan, and thiophene are 5-membered rings.

[00115] The term “fused” refers to structures in which two or more rings share one or more bonds.

[00116] The term “carbocyclic” or “carbocycle” refers to a ring wherein each of the atoms forming the ring is a carbon atom. Carbocycle includes aryl and cycloalkyl. The term thus distinguishes carbocycle from heterocycle (“heterocyclic”) in which the ring backbone contains at least one atom which is different from carbon (i.e. a heteroatom). Heterocycle includes heteroaryl and heterocycloalkyl. Carbocycles and heterocycles can be optionally substituted.


[00117] The term “aromatic” refers to a planar ring having a delocalized π -electron system containing $4n+2$ π electrons, where n is an integer. Aromatic rings can be formed from five, six, seven, eight, nine, or more than nine atoms. Aromatics can be optionally substituted. The term “aromatic” includes both carbocyclic aryl (e.g., phenyl) and heterocyclic aryl (or “heteroaryl” or “heteroaromatic”) groups (e.g., pyridine). The term includes monocyclic or fused-ring polycyclic (i.e., rings which share adjacent pairs of carbon atoms) groups.

[00118] As used herein, the term “aryl” refers to an aromatic ring wherein each of the atoms forming the ring is a carbon atom. Aryl rings can be formed by five, six, seven, eight, nine, or more than nine carbon atoms. Aryl groups can be optionally substituted. Examples of aryl groups include, but are not limited to phenyl, naphthalenyl, phenanthrenyl, anthracenyl, fluorenyl, and indenyl. Depending on the structure, an aryl group can be a monoradical or a diradical (i.e., an arylene group).

[00119] An “aryloxy” group refers to an (aryl)O- group, where aryl is as defined herein.

[00120] The term “carbonyl” as used herein refers to a group containing a moiety selected from the group consisting of -C(O)-, -S(O)-, -S(O)₂-, and -C(S)-, including, but not limited to, groups containing at least one ketone group, and/or at least one aldehyde group, and/or at least one ester group, and/or at least one carboxylic acid group, and/or at least one thioester group. Such carbonyl groups include ketones, aldehydes, carboxylic acids, esters, and thioesters. In some embodiments, such groups are a part of linear, branched, or cyclic molecules.

[00121] The term “cycloalkyl” refers to a monocyclic or polycyclic radical that contains only carbon and hydrogen, and is optionally saturated, partially unsaturated, or fully unsaturated. Cycloalkyl groups include groups having from 3 to 10 ring atoms. Illustrative examples of cycloalkyl groups include the following moieties:




, and the like. Depending on the structure, a cycloalkyl group is either a monoradical or a diradical (e.g., an cycloalkylene group), and if a “lower cycloalkyl” having 3 to 8 carbon atoms.

[00122] “Cycloalkylalkyl” means an alkyl radical, as defined herein, substituted with a cycloalkyl group. Non-limiting cycloalkylalkyl groups include cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, and the like.

[00123] The term “heterocycle” refers to heteroaromatic and heteroalicyclic groups containing one to four heteroatoms each selected from O, S and N, wherein each heterocyclic group has from 4 to 10 atoms in its ring system, and with the proviso that the ring of said group does not contain two adjacent O or S atoms. Herein, whenever the number of carbon atoms in a heterocycle is indicated (e.g., C₁-C₆ heterocycle), at least one other atom (the heteroatom) must be present in the ring. Designations such as “C₁-C₆ heterocycle” refer only to the number of carbon atoms in the ring and do not refer to the total number of atoms in the ring. It is understood that the heterocyclic ring can have additional heteroatoms in the ring. Designations such as “4-6 membered heterocycle” refer to the total number of atoms that are contained in the ring (i.e., a four, five, or six membered ring, in which at least one atom is a carbon atom, at least one atom is a heteroatom and the remaining two to four atoms are either carbon atoms or heteroatoms). In heterocycles that have two or more heteroatoms, those two or more heteroatoms can be the same or different from one another. Heterocycles can be optionally substituted. Binding to a heterocycle can be at a heteroatom or via a carbon atom. Non-aromatic heterocyclic groups include groups having only 4 atoms in their ring system, but aromatic heterocyclic groups must have at least 5 atoms in their ring system. The heterocyclic groups include benzo-fused ring systems. An example of a 4-membered heterocyclic group is azetidiny (derived from azetidine). An example of a 5-membered heterocyclic group is thiazolyl. An example of a 6-membered heterocyclic group is pyridyl, and an example of a 10-membered heterocyclic group is quinolinyl. Examples of non-aromatic heterocyclic groups are pyrrolidinyl,


tetrahydrofuranyl, dihydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, dihydropyranyl, tetrahydrothiopyranyl, piperidino, morpholino, thiomorpholino, thioxanyl, piperazinyl, azetidiny, oxetanyl, thietanyl, homopiperidiny, oxepanyl, thiepanyl, oxazepiny, diazepiny, thiazepiny, 1,2,3,6-tetrahydropyridiny, 2-pyrroliny, 3-pyrroliny, indoliny, 2H-pyranyl, 4H-pyranyl, dioxanyl, 1,3-dioxolanyl, pyrazoliny, dithianyl, dithiolanyl, dihydropyranyl, dihydrothienyl, dihydrofuranyl, pyrazolidiny, imidazoliny, imidazolidiny, 3-azabicyclo[3.1.0]hexanyl, 3-azabicyclo[4.1.0]heptanyl, 3H-indolyl and quinoliziny. Examples of aromatic heterocyclic groups are pyridiny, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyraziny, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrroly, quinoliny, isoquinoliny, indolyl, benzimidazolyl, benzofuranyl, cinnoliny, indazolyl, indoliziny, phthalaziny, pyridaziny, triaziny, isoindolyl, pteridinyl, puriny, oxadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxaliny, naphthyridiny, and furopyridiny. The foregoing groups, as derived from the groups listed above, are optionally C-attached or *N*-attached where such is possible. For instance, a group derived from pyrrole includes pyrrol-1-yl (*N*-attached) or pyrrol-3-yl (C-attached). Further, a group derived from imidazole includes imidazol-1-yl or imidazol-3-yl (both *N*-attached) or imidazol-2-yl, imidazol-4-yl or imidazol-5-yl (all C-attached). The heterocyclic groups include benzo-fused ring systems and ring systems substituted with one or two oxo (=O) moieties such as pyrrolidin-2-one. Depending on the structure, a heterocycle group can be a monoradical or a diradical (i.e., a heterocyclene group).

[00124] The terms “heteroaryl” or, alternatively, “heteroaromatic” refers to an aromatic group that includes one or more ring heteroatoms selected from nitrogen, oxygen and sulfur. An *N*-containing “heteroaromatic” or “heteroaryl” moiety refers to an aromatic group in which at least one of the skeletal atoms of the ring is a nitrogen atom. Illustrative examples of heteroaryl groups include the following moieties:



and the like. Depending on the structure, a heteroaryl group can be a monoradical or a diradical (i.e., a heteroarylene group).

[00125] As used herein, the term “non-aromatic heterocycle”, “heterocycloalkyl” or “heteroalicyclic” refers to a non-aromatic ring wherein one or more atoms forming the ring is a heteroatom. A “non-aromatic heterocycle” or “heterocycloalkyl” group refers to a cycloalkyl group that includes at least one heteroatom selected from nitrogen, oxygen and sulfur. In some embodiments, the radicals are fused with an aryl or heteroaryl. Heterocycloalkyl rings can be formed by three, four, five, six, seven, eight, nine, or more than nine atoms. Heterocycloalkyl rings can be optionally substituted. In certain embodiments, non-aromatic heterocycles contain one or more carbonyl or thiocarbonyl groups such as, for example, oxo- and thio-containing groups. Examples of heterocycloalkyls include, but are not limited to, lactams, lactones, cyclic imides, cyclic thioimides, cyclic carbamates, tetrahydrothiopyran, 4H-pyran, tetrahydropyran, piperidine, 1,3-dioxin, 1,3-dioxane, 1,4-dioxin, 1,4-dioxane, piperazine, 1,3-oxathiane, 1,4-oxathiin, 1,4-oxathiane, tetrahydro-1,4-thiazine, 2H-1,2-oxazine, maleimide, succinimide, barbituric acid, thiobarbituric acid, dioxopiperazine, hydantoin, dihydrouracil, morpholine, trioxane, hexahydro-1,3,5-triazine, tetrahydrothiophene, tetrahydrofuran, pyrroline, pyrrolidine, pyrrolidone, pyrazoline, pyrazolidine, imidazoline, imidazolidine, 1,3-dioxole, 1,3-dioxolane, 1,3-dithiole, 1,3-dithiolane, isoxazoline, isoxazolidine, oxazoline, oxazolidine, oxazolidinone, thiazoline, thiazolidine, and 1,3-oxathiolane. Illustrative examples of heterocycloalkyl groups, also referred to as non-aromatic heterocycles, include:



the like. The term heteroalicyclic also includes all ring forms of the carbohydrates, including but not limited to the monosaccharides, the disaccharides and the oligosaccharides. Depending on the structure, a heterocycloalkyl group can be a monoradical or a diradical (i.e., a heterocycloalkylene group).

[00126] The term “halo” or, alternatively, “halogen” or “halide” means fluoro, chloro, bromo and iodo.

[00127] The term “haloalkyl,” refers to alkyl structures in which at least one hydrogen is replaced with a halogen atom. In certain embodiments in which two or more hydrogen atoms are

replaced with halogen atoms, the halogen atoms are all the same as one another. In other embodiments in which two or more hydrogen atoms are replaced with halogen atoms, the halogen atoms are not all the same as one another.

[00128] The term “fluoroalkyl,” as used herein, refers to alkyl group in which at least one hydrogen is replaced with a fluorine atom. Examples of fluoroalkyl groups include, but are not limited to, $-\text{CF}_3$, $-\text{CH}_2\text{CF}_3$, $-\text{CF}_2\text{CF}_3$, $-\text{CH}_2\text{CH}_2\text{CF}_3$ and the like.

[00129] As used herein, the term “heteroalkyl” refers to optionally substituted alkyl radicals in which one or more skeletal chain atoms is a heteroatom, *e.g.*, oxygen, nitrogen, sulfur, silicon, phosphorus or combinations thereof. The heteroatom(s) are placed at any interior position of the heteroalkyl group or at the position at which the heteroalkyl group is attached to the remainder of the molecule. Examples include, but are not limited to, $-\text{CH}_2-\text{O}-\text{CH}_3$, $-\text{CH}_2-\text{CH}_2-\text{O}-\text{CH}_3$, $-\text{CH}_2-\text{NH}-\text{CH}_3$, $-\text{CH}_2-\text{CH}_2-\text{NH}-\text{CH}_3$, $-\text{CH}_2-\text{N}(\text{CH}_3)-\text{CH}_3$, $-\text{CH}_2-\text{CH}_2-\text{NH}-\text{CH}_3$, $-\text{CH}_2-\text{CH}_2-\text{N}(\text{CH}_3)-\text{CH}_3$, $-\text{CH}_2-\text{S}-\text{CH}_2-\text{CH}_3$, $-\text{CH}_2-\text{CH}_2-\text{S}(\text{O})-\text{CH}_3$, $-\text{CH}_2-\text{CH}_2-\text{S}(\text{O})_2-\text{CH}_3$, $-\text{CH}=\text{CH}-\text{O}-\text{CH}_3$, $-\text{Si}(\text{CH}_3)_3$, $-\text{CH}_2-\text{CH}=\text{N}-\text{OCH}_3$, and $-\text{CH}=\text{CH}-\text{N}(\text{CH}_3)-\text{CH}_3$. In addition, in some embodiments, up to two heteroatoms are consecutive, such as, by way of example, $-\text{CH}_2-\text{NH}-\text{OCH}_3$ and $-\text{CH}_2-\text{O}-\text{Si}(\text{CH}_3)_3$.

[00130] The term “heteroatom” refers to an atom other than carbon or hydrogen. Heteroatoms are typically independently selected from among oxygen, sulfur, nitrogen, silicon and phosphorus, but are not limited to these atoms. In embodiments in which two or more heteroatoms are present, the two or more heteroatoms can all be the same as one another, or some or all of the two or more heteroatoms can each be different from the others.

[00131] The term “bond” or “single bond” refers to a chemical bond between two atoms, or two moieties when the atoms joined by the bond are considered to be part of larger substructure.

[00132] The term “moiety” refers to a specific segment or functional group of a molecule. Chemical moieties are often recognized chemical entities embedded in or appended to a molecule.

[00133] A “thioalkoxy” or “alkylthio” group refers to a $-\text{S}$ -alkyl group.


[00134] A “SH” group is also referred to either as a thiol group or a sulfhydryl group.

[00135] The term “optionally substituted” or “substituted” means that the referenced group may be substituted with one or more additional group(s) individually and independently selected from alkyl, cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, alkoxy, aryloxy, alkylthio, arylthio, alkylsulfoxide, arylsulfoxide, alkylsulfone, arylsulfone, cyano, halo, acyl, nitro, haloalkyl, fluoroalkyl, amino, including mono- and di-substituted amino groups, and the protected derivatives thereof. By way of example an optional substituents may be L_sR_s , wherein

each L_s is independently selected from a bond, -O-, -C(=O)-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -NHC(O)-, -C(O)NH-, S(=O)₂NH-, -NHS(=O)₂-, -OC(O)NH-, -NHC(O)O-, -(substituted or unsubstituted C₁-C₆ alkyl), or -(substituted or unsubstituted C₂-C₆ alkenyl); and each R_s is independently selected from H, (substituted or unsubstituted C₁-C₄alkyl), (substituted or unsubstituted C₃-C₆cycloalkyl), heteroaryl, or heteroalkyl. The protecting groups that form the protective derivatives of the above substituents include those found in sources such as Greene and Wuts, above.

ACK Inhibitor Compounds

[00136] Described herein methods of treating solid tumors comprising co-administering an ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and an HDAC inhibitor compound. In some embodiments, the HDAC inhibitor is a compound of Formula (B). In some embodiments, the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat).




Abexinostat

[00137] The ACK inhibitor compounds described herein are selective for kinases having an accessible cysteine that is able to form a covalent bond with a Michael acceptor moiety on the inhibitor compound. In some embodiments, the cysteine residue is accessible or becomes accessible when the binding site moiety of the irreversible inhibitor binds to the kinase. That is, the binding site moiety of the irreversible inhibitor binds to an active site of the ACK and the Michael acceptor moiety of irreversible inhibitor gains access (in one embodiment the step of binding leads to a conformational change in the ACK, thus exposing the cysteine) or is otherwise exposed to the cysteine residue of the ACK; as a result a covalent bond is formed between the “S” of the cysteine residue and the Michael acceptor of the irreversible inhibitor.

Consequently, the binding site moiety of the irreversible inhibitor remains bound or otherwise blocks the active site of the ACK.

[00138] In one embodiment, the ACK is Btk, a homolog of Btk or a tyrosine kinase having a cysteine residue in an amino acid sequence position that is homologous to the amino acid sequence position of cysteine 481 in Btk. In some embodiments, the ACK is HER4. Inhibitor compounds described herein include a Michael acceptor moiety, a binding site moiety and a linker that links the binding site moiety and the Michael acceptor moiety (and in some embodiments, the structure of the linker provides a conformation, or otherwise directs the Michael acceptor moiety, so as to improve the selectivity of the irreversible inhibitor for a particular ACK).

[00139] In some embodiments, the ACK inhibitor is a compound of Formula (A):



Formula (A)

wherein

A is independently selected from N or CR₅;

R₁ is H, L₂-(substituted or unsubstituted alkyl), L₂-(substituted or unsubstituted cycloalkyl), L₂-(substituted or unsubstituted alkenyl), L₂-(substituted or unsubstituted cycloalkenyl), L₂-(substituted or unsubstituted heterocycle), L₂-(substituted or unsubstituted heteroaryl), or L₂-(substituted or unsubstituted aryl), where L₂ is a bond, O, S, -S(=O), -S(=O)₂, C(=O), -(substituted or unsubstituted C₁-C₆ alkyl), or -(substituted or unsubstituted C₂-C₆ alkenyl);

R₂ and R₃ are independently selected from H, lower alkyl and substituted lower alkyl;


R₄ is L₃-X-L₄-G, wherein,

L₃ is optional, and when present is a bond, optionally substituted or unsubstituted alkyl, optionally substituted or unsubstituted cycloalkyl, optionally substituted or unsubstituted alkenyl, optionally substituted or unsubstituted alkynyl;

X is optional, and when present is a bond, O, -C(=O), S, -S(=O), -S(=O)₂, -NH, -NR₉, -NHC(O), -C(O)NH, -NR₉C(O), -C(O)NR₉, -S(=O)₂NH, -NHS(=O)₂, -S(=O)₂NR₉, -NR₉S(=O)₂, -OC(O)NH-, -NHC(O)O-, -OC(O)NR₉-, -NR₉C(O)O-, -CH=NO-, -ON=CH-, -NR₁₀C(O)NR₁₀-, heteroaryl, aryl, -NR₁₀C(=NR₁₁)NR₁₀-, -NR₁₀C(=NR₁₁)-, -C(=NR₁₁)NR₁₀-, -OC(=NR₁₁)-, or -C(=NR₁₁)O-;

L₄ is optional, and when present is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycle;

or L₃, X and L₄ taken together form a nitrogen containing heterocyclic ring;



R₆, R₇ and R₈ are independently selected from among H, lower alkyl or substituted lower alkyl, lower heteroalkyl or substituted lower heteroalkyl, substituted or unsubstituted lower cycloalkyl, and substituted or unsubstituted lower heterocycloalkyl;

R₅ is H, halogen, -L₆-(substituted or unsubstituted C₁-C₃ alkyl), -L₆-(substituted or unsubstituted C₂-C₄ alkenyl), -L₆-(substituted or unsubstituted heteroaryl), or -L₆-(substituted or unsubstituted aryl), wherein L₆ is a bond, O, S, -S(=O), S(=O)₂, NH, C(O), -NHC(O)O, -OC(O)NH, -NHC(O), or -C(O)NH;

each R₉ is independently selected from among H, substituted or unsubstituted lower alkyl, and substituted or unsubstituted lower cycloalkyl;

each R₁₀ is independently H, substituted or unsubstituted lower alkyl, or substituted or unsubstituted lower cycloalkyl; or


two R₁₀ groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or

R₉ and R₁₀ can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or

each R₁₁ is independently selected from H, -S(=O)₂R₈, -S(=O)₂NH₂, -C(O)R₈, -CN, -NO₂, heteroaryl, or heteroalkyl; and

pharmaceutically active metabolites, pharmaceutically acceptable solvates, pharmaceutically acceptable salts, or pharmaceutically acceptable prodrugs thereof.

[00140] In some embodiments, the ACK inhibitor is R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib)
















Ibrutinib.


[00141] In some embodiments, the ACK inhibitor is AVL-263 (Avila Therapeutics/Celgene Corporation), AVL-292 (Avila Therapeutics/Celgene Corporation), AVL-291 (Avila Therapeutics/Celgene Corporation), BMS-488516 (Bristol-Myers Squibb), BMS-509744 (Bristol-Myers Squibb), CGI-1746 (CGI Pharma/Gilead Sciences), CTA-056, GDC-0834 (Genentech), HY-11066 (also, CTK4I7891, HMS3265G21, HMS3265G22, HMS3265H21, HMS3265H22, 439574-61-5, AG-F-54930), ONO-4059 (Ono Pharmaceutical Co., Ltd.), ONO-WG37 (Ono Pharmaceutical Co., Ltd.), PLS-123 (Peking University), RN486 (Hoffmann-La Roche), or HM71224 (Hanmi Pharmaceutical Company Limited).

[00142] In some embodiments, the ACK inhibitor is 4-(tert-butyl)-N-(2-methyl-3-(4-methyl-6-((4-(morpholine-4-carbonyl)phenyl)amino)-5-oxo-4,5-dihydropyrazin-2-yl)phenyl)benzamide (CGI-1746); 7-benzyl-1-(3-(piperidin-1-yl)propyl)-2-(4-(pyridin-4-yl)phenyl)-1H-imidazo[4,5-g]quinoxalin-6(5H)-one (CTA-056); (R)-N-(3-(6-(4-(1,4-dimethyl-3-oxopiperazin-2-yl)phenylamino)-4-methyl-5-oxo-4,5-dihydropyrazin-2-yl)-2-methylphenyl)-4,5,6,7-tetrahydrobenzo[b]thiophene-2-carboxamide (GDC-0834); 6-cyclopropyl-8-fluoro-2-(2-hydroxymethyl-3-{1-methyl-5-[5-(4-methyl-piperazin-1-yl)-pyridin-2-ylamino]-6-oxo-1,6-dihydro-pyridin-3-yl}-phenyl)-2H-isoquinolin-1-one (RN-486); N-[5-[5-(4-acetylpiperazine-1-carbonyl)-4-methoxy-2-methylphenyl]sulfanyl-1,3-thiazol-2-yl]-4-[(3,3-dimethylbutan-2-ylamino)methyl]benzamide (BMS-509744, HY-11092); or N-(5-((5-(4-Acetylpiperazine-1-carbonyl)-4-methoxy-2-methylphenyl)thio)thiazol-2-yl)-4-(((3-methylbutan-2-yl)amino)methyl)benzamide (HY11066).

[00143] In some embodiments, the ACK inhibitor is:








HDAC Inhibitor Compounds

[00144] Described herein methods of treating solid tumors comprising co-administering an ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and an HDAC inhibitor compound. In some embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) is a compound of Formula (A). In some embodiments, the ACK inhibitor compound is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib).




Ibrutinib

In some embodiments, the ACK inhibitor is AVL-263 (Avila Therapeutics/Celgene Corporation), AVL-292 (Avila Therapeutics/Celgene Corporation), AVL-291 (Avila

Therapeutics/Celgene Corporation), BMS-488516 (Bristol-Myers Squibb), BMS-509744 (Bristol-Myers Squibb), CGI-1746 (CGI Pharma/Gilead Sciences), CTA-056, GDC-0834 (Genentech), HY-11066 (also, CTK4I7891, HMS3265G21, HMS3265G22, HMS3265H21, HMS3265H22, 439574-61-5, AG-F-54930), ONO-4059 (Ono Pharmaceutical Co., Ltd.), ONO-WG37 (Ono Pharmaceutical Co., Ltd.), PLS-123 (Peking University), RN486 (Hoffmann-La Roche), or HM71224 (Hanmi Pharmaceutical Company Limited).

[00145] In some embodiments, the HDAC inhibitor has the structure of Formula (B):



Formula (B)

wherein:

R¹ is hydrogen or alkyl;

X is -O-, -NR²-, or -S(O)_n where n is 0-2 and R² is hydrogen or alkyl;

Y is alkylene optionally substituted with cycloalkyl, optionally substituted phenyl, alkylthio, alkylsulfinyl, alkylsulfonyl, optionally substituted phenylalkylthio, optionally substituted phenylalkylsulfonyl, hydroxy, or optionally substituted phenoxy;


Ar¹ is phenylene or heteroarylene wherein said Ar¹ is optionally substituted with one or two groups independently selected from alkyl, halo, hydroxy, alkoxy, haloalkoxy, or haloalkyl;

R³ is hydrogen, alkyl, hydroxyalkyl, or optionally substituted phenyl; and

Ar² is aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, heteroaralkenyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, or heterocycloalkylalkyl;

and individual stereoisomers, individual geometric isomers, or mixtures thereof; or a pharmaceutically acceptable salt thereof.

[00146] In some embodiments, the histone deacetylase inhibitor is 3-((dimethylamino)methyl)-N-(2-(4-(hydroxycarbamoyl)phenoxy)ethyl)benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat).



Abexinostat

Pharmaceutical Compositions/Formulations

[00147] Disclosed herein, in certain embodiments, are compositions comprising a therapeutically effective amount of an ACK inhibitor compound and/or a therapeutically effective amount of an HDAC inhibitor compound, and a pharmaceutically acceptable excipient. Further disclosed herein, in certain embodiments, are compositions comprising a therapeutically effective amount of an ACK inhibitor compound, a therapeutically effective amount of an HDAC inhibitor compound, and a pharmaceutically acceptable excipient. In some embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) is a compound of Formula (A). In some embodiments, the ACK inhibitor compound is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib). In some embodiments, the ACK inhibitor is AVL-263 (Avila Therapeutics/Celgene Corporation), AVL-292 (Avila Therapeutics/Celgene Corporation), AVL-291 (Avila Therapeutics/Celgene Corporation), BMS-488516 (Bristol-Myers Squibb), BMS-509744 (Bristol-Myers Squibb), CGI-1746 (CGI Pharma/Gilead Sciences), CTA-056, GDC-0834 (Genentech), HY-11066 (also, CTK4I7891, HMS3265G21, HMS3265G22, HMS3265H21, HMS3265H22, 439574-61-5, AG-F-54930), ONO-4059 (Ono Pharmaceutical Co., Ltd.), ONO-WG37 (Ono Pharmaceutical Co., Ltd.), PLS-123 (Peking University), RN486 (Hoffmann-La Roche), or HM71224 (Hanmi Pharmaceutical Company Limited). In some embodiments, the HDAC inhibitor is a compound of Formula (B). In some embodiments, the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbonyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat).

[00148] Pharmaceutical compositions of ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and/or HDAC

inhibitor compound are formulated in a conventional manner using one or more physiologically acceptable carriers including excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen. A summary of pharmaceutical compositions described herein is found, for example, in Remington: The Science and Practice of Pharmacy, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., Pharmaceutical Dosage Forms, Marcel Decker, New York, N.Y., 1980; and Pharmaceutical Dosage Forms and Drug Delivery Systems, Seventh Ed. (Lippincott Williams & Wilkins 1999).

[00149] A pharmaceutical composition, as used herein, refers to a mixture of an ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and/or HDAC inhibitor compound with other chemical components, such as carriers, stabilizers, diluents, dispersing agents, suspending agents, thickening agents, and/or excipients.

[00150] Pharmaceutical compositions are optionally manufactured in a conventional manner, such as, by way of example only, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or compression processes.

[00151] The pharmaceutical formulations described herein are administered by any suitable administration route, including but not limited to, oral, parenteral (e.g., intravenous, subcutaneous, intramuscular), intranasal, buccal, topical, rectal, or transdermal administration routes.

[00152] The pharmaceutical compositions described herein are formulated into any suitable dosage form, including but not limited to, aqueous oral dispersions, liquids, gels, syrups, elixirs, slurries, suspensions and the like, for oral ingestion by an individual to be treated, solid oral dosage forms, aerosols, controlled release formulations, fast melt formulations, effervescent formulations, lyophilized formulations, tablets, powders, pills, dragees, capsules, delayed release formulations, extended release formulations, pulsatile release formulations, multiparticulate formulations, and mixed immediate release and controlled release formulations. In some embodiments, the compositions are formulated into capsules. In some embodiments, the compositions are formulated into solutions (for example, for IV administration).

[00153] The pharmaceutical solid dosage forms described herein optionally include a compound described herein and one or more pharmaceutically acceptable additives such as a compatible carrier, binder, filling agent, suspending agent, flavoring agent, sweetening agent, disintegrating

agent, dispersing agent, surfactant, lubricant, colorant, diluent, solubilizer, moistening agent, plasticizer, stabilizer, penetration enhancer, wetting agent, anti-foaming agent, antioxidant, preservative, or one or more combination thereof.

[00154] In still other aspects, using standard coating procedures, such as those described in Remington's Pharmaceutical Sciences, 20th Edition (2000), a film coating is provided around the compositions. In some embodiments, the compositions are formulated into particles (for example for administration by capsule) and some or all of the particles are coated. In some embodiments, the compositions are formulated into particles (for example for administration by capsule) and some or all of the particles are microencapsulated. In some embodiments, the compositions are formulated into particles (for example for administration by capsule) and some or all of the particles are not microencapsulated and are uncoated.

[00155] In some embodiments, the pharmaceutical compositions are formulated such that the amount of the ACK inhibitor (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) in each unit dosage form is about 140 mg per.

Kits/Articles of Manufacture

[00156] Described herein methods of treating solid tumors comprising co-administering an ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) and an HDAC inhibitor compound. In some embodiments, the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor, such as for example ibrutinib) is a compound of Formula (A). In some embodiments, the ACK inhibitor compound is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib). In some embodiments, the ACK inhibitor is AVL-263 (Avila Therapeutics/Celgene Corporation), AVL-292 (Avila Therapeutics/Celgene Corporation), AVL-291 (Avila Therapeutics/Celgene Corporation), BMS-488516 (Bristol-Myers Squibb), BMS-509744 (Bristol-Myers Squibb), CGI-1746 (CGI Pharma/Gilead Sciences), CTA-056, GDC-0834 (Genentech), HY-11066 (also, CTK4I7891, HMS3265G21, HMS3265G22, HMS3265H21, HMS3265H22, 439574-61-5, AG-F-54930), ONO-4059 (Ono Pharmaceutical Co., Ltd.), ONO-WG37 (Ono Pharmaceutical Co., Ltd.), PLS-123 (Peking University), RN486 (Hoffmann-La Roche), or HM71224 (Hanmi Pharmaceutical Company Limited). In some embodiments, the HDAC inhibitor is a compound of Formula (B). In some embodiments, the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat).

[00157] For use in the therapeutic applications described herein, kits and articles of manufacture are also described herein. In some embodiments, such kits include a carrier, package, or container that is compartmentalized to receive one or more containers such as vials, tubes, and the like, each of the container(s) including one of the separate elements to be used in a method described herein. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers can be formed from a variety of materials such as glass or plastic.

[00158] The articles of manufacture provided herein contain packaging materials. Examples of pharmaceutical packaging materials include, but are not limited to, blister packs, bottles, tubes, inhalers, pumps, bags, vials, containers, syringes, bottles, and any packaging material suitable for a selected formulation and intended mode of administration and treatment. A wide array of formulations of the compounds and compositions provided herein are contemplated as are a variety of treatments for any disorder that benefit by inhibition of Btk, or in which Btk is a mediator or contributor to the symptoms or cause.

[00159] For example, a container may include one or both of the irreversible BTK inhibitor and HDAC inhibitor compounds. The container(s) optionally have a sterile access port (for example the container is an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). Such kits optionally comprising a compound with an identifying description or label or instructions relating to its use in the methods described herein.

[00160] A kit will typically include one or more additional containers, each with one or more of various materials (such as reagents, optionally in concentrated form, and/or devices) desirable from a commercial and user standpoint for use of a compound described herein. Non-limiting examples of such materials include, but not limited to, buffers, diluents, filters, needles, syringes; carrier, package, container, vial and/or tube labels listing contents and/or instructions for use, and package inserts with instructions for use. A set of instructions will also typically be included.

[00161] In some embodiments, a label is on or associated with the container. A label can be on a container when letters, numbers or other characters forming the label are attached, molded or etched into the container itself; a label can be associated with a container when it is present within a receptacle or carrier that also holds the container, e.g., as a package insert. A label can be used to indicate that the contents are to be used for a specific therapeutic application. The label can also indicate directions for use of the contents, such as in the methods described herein.

[00162] In certain embodiments, a pharmaceutical composition comprising one or both the ACK inhibitor compound (e.g., a BTK inhibitor, such as for example an irreversible BTK inhibitor,

such as for example ibrutinib) and the HDAC inhibitor compounds is presented in a pack or dispenser device which can contain one or more unit dosage forms. The pack can for example contain metal or plastic foil, such as a blister pack. The pack or dispenser device can be accompanied by instructions for administration. The pack or dispenser can also be accompanied with a notice associated with the container in form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the drug for human or veterinary administration. Such notice, for example, can be the labeling approved by the U.S. Food and Drug Administration for prescription drugs, or the approved product insert. Compositions containing a compound provided herein formulated in a compatible pharmaceutical carrier can also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

EXAMPLES

[00163] The following specific and non-limiting examples are to be construed as merely illustrative, and do not limit the present disclosure in any way whatsoever.

Example 1

[00164] The aim of the experiment is to evaluate the effect of two drugs on tumor development in a mouse model of non-small cell lung cancer.

[00165] The two drugs tested were Abexinostat (PCI-24781), and Ibrutinib (PCI-32765).

[00166] Grg1 transgenic mice were used to evaluate the drugs. Grg1 transgenic mice overexpress the Groucho-related gene 1 (Grg1). The mice develop lung tumors that resemble human non-small cell lung cancer. Tumors initiate at 1 month of age and progress to invasive adenocarcinoma by 8 months of age. The drugs were administered to 2 month-old mice or to 5 month-old mice, and mice were treated for 4 weeks. The former group is referred to as the 3 month samples and the latter as the 6 month samples.

Summary

[00167] Drug treatment of 2 to 3 month old mice reduced tumor number compared to the control group. The combination of abexinostat and ibrutinib reduced tumor number by 75%.

[00168] With the combination of abexinostat and ibrutinib treatment for 5 to 6 months, there were no large tumors.

Methods

[00169] a. Mice

[00170] Mice were housed and treated following the applicable Standard Operating Procedures and according to the Canadian Animal Care Committee standards.

[00171] Mice were bred to generate double transgenic (Grg1/Cre recombinase) mice on a CD1 background. Four groups of female double transgenic mice were established:

[00172] Group 1 – Control (no treatment, injection of PBS, or water with carrier and no drug)

[00173] Group 2 – Abexinostat-treated

[00174] Group 3 – Ibrutinib-treated

[00175] Group 4 – Abexinostat + Ibrutinib-treated

[00176] At least 8 mice in each group were treated at 2 months old and 8 mice in each group were treated at 5 months old. Treatments lasted 4 weeks, and mice were sacrificed at 3 months and 6 months. Therefore samples are referred to as 3 month samples or 6 month samples.

[00177] b. Drug preparation and dosage

[00178] Abexinostat was administered by intraperitoneal (i.p.) injection, twice per day for five days each week. Mice were weighed and injected with 6 µl per gram body weight of a 1 mg/ml solution. The dosage was 12 mg/kg BID.

[00179] Ibrutinib was administered by supplying it in the drinking water, for a dosage of ~ 22 mg/kg/day.

[00180] Mice were treated for 4 weeks. After treatment, mice were euthanized and lung tissue was examined.

[00181] Drugs were prepared and administered as follows:

[00182] *Abexinostat*

[00183] Formulation: Add DMSO to powder stock in vial to make a 200mg/ml solution. Aliquot out into 100 µl. Every two weeks, as required, thaw a 100 µl aliquot and add 900 µl sterile water. Aliquot into 20 µl aliquots of 20 mg/ml. Store this stock at 4 C.

[00184] Dosage

[00185] 20 µl of 20mg/ml stock in the fridge; inject at 1 mg/ml concentration in sterile water. On the day of injection, add 380 µl sterile water or PBS and pipet well to mix thoroughly. Inject 6 µl per g body weight TWICE DAILY.

[00186] *Ibrutinib*

[00187] Formulation: 100 ml of the 10x concentrate of Ibrutinib. Dilute 1 part of concentrate with 9 parts water. Both the 10x and 1x dilutions can be stored at room temperature. PCI-32765 is >99% stable in this formulation after 6 weeks at room temperature (22°C).

[00188] Dosage

[00189] Administered by drinking water; Average consumption is ~4 mL/day/mouse, corresponding to a final dose of ~22 mg/kg/day. Thus 4ml x 4 mice = 16ml/day x 7 days = 140ml per week

[00190] c. Quantitation

[00191] For quantitation of tumors, the total visible tumors on the lung surface were counted using a dissecting microscope. In addition, the left lung lobe was fixed in 4% PFA, paraffin-embedded and 5 µm sections were taken for 100 µm at two levels. The first level began 30 µm below the lung surface and the second level began 130 µm below the lung surface. Alternate slides were H&E stained, and tumors were detected and counted using a dissecting microscope. Tumor diameter was measured using an inverted microscope and reticule.

[00192] The right lung lobes were snap-frozen and stored for DNA or protein analysis.

[00193] d. Blood Samples

[00194] For 6 month-old mice, blood samples were collected in EDTA tubes, and peripheral blood mononuclear cells (PBMC) and plasma were prepared, frozen and stored as follows:

[00195] 1. A minimum of 500 µL of blood is drawn into the EDTA tube

[00196] 2. Transfer the blood into 1.5 mL eppendorf tubes and centrifuge at 350 x g for 5 minutes to separate the plasma and the cells.

[00197] 3. Save the supernatant/plasma.

[00198] 4. Add 1 ml of 1X RBC Lysis Buffer (Sigma #R7757) to the pellet

[00199] 5. Gently vortex each tube immediately after adding the lysing solution.

[00200] 6. Incubate at room temperature, protected from light, for 5 minutes.

[00201] 7. Centrifuge 350 x g for 2 minutes.

[00202] 8. Aspirate supernatant without disturbing pellet.

[00203] 9. Wash the pellet once with 1 mL of PBS to the sample and centrifuge 350 x g for 2 minutes

[00204] 10. Aspirate supernatant without disturbing the pellet.

[00205] 11. Freeze the PBMC pellet at -80C

Results

[00206] a. Three month old

[00207] The total number of surface tumors visible at 3 months, after 4 weeks of treatment, is shown in a table and corresponding bar graph in Figure 2. Each number in the table represents the number of tumors in one mouse. The average number of tumors per mouse for each treatment is shown in the bottom row of the table and right-most bar for each group. For the control group, the average number of tumors was 5.9 (n=10). For the Abexinostat-treated mice,

the average tumor number was 3.0 (n=10), for Ibrutinib it was 2.6 (n=8), and for treatment with both drugs 1.4 (n=11).

[00208] Tumors in the 3 month samples were further analyzed for four mice in each treatment group by sectioning the left lung lobe, and H&E staining alternate slides. The tumor histology is presented in Figure 3 and all images are available on disk.

[00209] The tumors are carcinoma that are less than 400 μm . The tumor histology appears similar in treated and control mice.

[00210] b. Six month old

[00211] The total number of surface tumors visible at 6 months, after 4 weeks of treatment, is shown in a table and corresponding bar graph in Figure 4. Each number in the table represents the number of tumors in one mouse. The average number of tumors per mouse for each treatment is shown in the bottom row and the right-most bar for each group. For the control group, the average number of tumors was 7.8 (n=8). For the Abexinostat-treated mice, the average tumor number was 4.6 (n=8), for Ibrutinib it was 7.6 (n=5), and for treatment with both drugs 8.4 (n=5).

[00212] The number of visible surface tumors in the 6 month-old mice was quite variable and did not show a trend between treatment groups.

[00213] The tumors were further analyzed for two to four mice in each group by sectioning the left lung lobe, and H&E staining alternate slides. To measure and compare the tumor load in the mice in the 6 month samples, the tumor diameter in the tissue sections was measured. The number of tumors in the sections and the size of each tumor is recorded in the table and corresponding bar graph in Figure 5. Each column in the table and group of bars in the graph shows the tumor sizes for one mouse.

[00214] The treated mice had fewer large tumors than the control mice. The number of tumors over 1 mm in diameter for each mouse is shown in Figure 6. Control mice had an average of 1.5 large tumors per mouse, Abexinostat- and Ibrutinib-treated mice had an average of 0.5 large tumors per mouse and Abexinostat+Ibrutinib-treated mice had 0 large tumors per mouse.

[00215] Images of the whole left lung lobe from control and treated mice are shown in Figure 7 and the tissue sections showing the tumor histology are shown in Figure 8.

Conclusion

[00216] The data analysis reveals that both Abexinostat and Ibrutinib greatly reduce tumor number in mice treated from 2 to 3 months and the drugs appear to act synergistically. The reduction was 50-60% for each drug alone and 75% when the drugs were used together.

Although the drugs reduce tumor number, the tumors appear morphologically similar between treated and control samples at 3 months.

[00217] At 6 months, the tumor number is variable, but in treated mice the number of large tumors (over 1 mm) is much lower. The drugs also appear to act synergistically at this stage to reduce the number of large tumors. The tumors for control and treated mice have lymphocyte infiltration. However tumors in mice receiving Abexinostat have a high number of apoptotic cells. Tumors in mice receiving Ibrutinib lack the high number of apoptotic macrophages seen in the control and Abexinostat-treated mice, and have less tumor necrosis than the control samples.

Example 2: Clinical Trial for Pancreatic Cancer

[00218] Study Type: Interventional

[00219] Study Design: Allocation: Non-Randomized

[00220] Endpoint Classification: Safety/Efficacy Study

[00221] Intervention Model: Single Group Assignment

[00222] Masking: Open Label

[00223] Primary Purpose: Treatment

[00224] Primary Outcome Measures: To assess the response rate associated with ibrutinib and abexinostat in patients with advanced pancreatic cancer

[00225] Secondary Outcome Measures: To assess the side effects of ibrutinib and abexinostat in patients with advanced pancreatic cancer

[00226] Detailed Description

[00227] Patients will be treated three times daily ibrutinib (140 mg/dose) and once daily with abexinostat for 4 consecutive days, followed by 3 days off of abexinostat (a cycle is 7 days). At day 1 of each cycle a physical exam and blood work will be performed. Reassessment of tumor size will be conducted at 6 weeks, 12 weeks and then every 9 weeks thereafter. Patients will remain on treatment until one of the following occur: disease progression, illness that prevents further treatment or unacceptable adverse events.

Eligibility

[00228] Ages Eligible for Study: 18 Years and older

[00229] Genders Eligible for Study: Both

[00230] Accepts Healthy Volunteers: No

[00231] Inclusion Criteria:

[00232] Metastatic pancreatic carcinoma (excluding pancreatic endocrine tumors)

[00233] Only patients with measurable disease

[00234] ECOG performance status < or equal to 1

[00235] Life expectancy >12 weeks

[00236] Signed informed consent

[00237] Failed or intolerance to first-line therapy for metastatic disease with a gemcitabine-containing regimen. Patients may have received adjuvant therapy in addition to one prior regimen for metastatic disease.

[00238] >4 weeks must have elapsed from the completion of previous chemotherapy and patients must have recovered from any related toxicities

[00239] >4 weeks must have elapsed from the participation in any investigational drug study

[00240] Laboratory values: ANC > 1500/mm³; Hemoglobin > 9.0 gm/dl; Platelets > 100,000/mm³; SGOT <2.5 X upper limit of normal; or <5 X upper limit of normal if evidence of liver metastases; Alkaline phosphatase < 2.5 X upper limit of normal; or < 5 X upper limit of normal if evidence of liver metastases; Total bilirubin < 1.5 X upper limit of normal; Creatinine clearance > 50 cc/min (by Cockcroft - Gault or as determined from a 24-hour urine collection).

[00241] Exclusion Criteria:

[00242] More than one prior chemotherapy treatment regimen for metastatic disease

[00243] Clinically apparent central nervous system (CNS) metastases or carcinomatous meningitis

[00244] Clinically significant cardiac disease (e.g. congestive heart failure, symptomatic coronary artery disease and cardiac arrhythmias not well controlled with medication or heart attack within the last 12 months).

[00245] Major surgery within 4 weeks of the start of study treatment, without complete recovery.

[00246] Evidence of CNS metastases (unless CNS metastases have been stable for > 3 months) or history of uncontrolled seizures, central nervous system disorders

[00247] Uncontrolled serious medical or psychiatric illness

[00248] Women must not be pregnant or lactating

[00249] Concurrent radiation therapy

[00250] Other active malignancy

[00251] Inability to swallow capsules

[00252] Patients lacking physical integrity of the upper gastrointestinal tract or who have malabsorption syndrome

Example 3: Colon Cancer

[00253] Study Type: Interventional

[00254] Study Design: Allocation: Randomized

[00255] Endpoint Classification: Efficacy Study

[00256] Intervention Model: Parallel Assignment

[00257] Masking: Double Blind (Subject, Investigator)

[00258] Primary Outcome Measures: Disease-free Survival following therapy with ibrutinib and abexinostat

[00259] Secondary Outcome Measures: Survival as Assessed by Death From Any Cause

Detailed Description

[00260] Patients will be treated three times daily ibrutinib (140 mg/dose) and once daily with abexinostat for 4 consecutive days, followed by 3 days off of abexinostat (a cycle is 7 days). At day 1 of each cycle a physical exam and blood work will be performed. Reassessment of tumor size will be conducted at 6 weeks, 12 weeks and then every 9 weeks thereafter. Patients will remain on treatment until one of the following occur: disease progression, illness that prevents further treatment or unacceptable adverse events.

Eligibility

[00261] Ages Eligible for Study: 18 Years and older

[00262] Genders Eligible for Study: Both

[00263] Accepts Healthy Volunteers: No

[00264] Inclusion Criteria:

[00265] Patients must consent to be in the study and must have signed and dated an IRB approved consent form conforming to federal and institutional guidelines

[00266] Randomization must occur during the three-week interval beginning on postoperative day 29 and ending on postoperative day 50

[00267] The distal extent of the tumor must be ≥ 12 cm from the anal verge on endoscopy; if the patient is not a candidate for endoscopy, then the distal extent of the tumor must be ≥ 12 cm from the anal verge as determined by surgical examination

[00268] The patient must have had an en bloc complete gross resection of tumor (curative resection) by open laparotomy or laparoscopically-assisted colectomy; patients who have had a two-stage surgical procedure, to first provide a decompressive colostomy and then in a later procedure to have the definitive surgical resection, are eligible

[00269] Patients must have histologically confirmed adenocarcinoma of the colon that meets one of the following criteria: Stage II carcinoma (T3,4 N0 M0); the tumor invades through the muscularis propria into the subserosa or into non-peritonealized pericolic or perirectal tissues (T3); or directly invades other organs or structures, and/or perforates visceral peritoneum (T4);

Stage III carcinoma (any T N1,2 M0); the tumor has invaded to any depth, with involvement of regional lymph nodes

[00270] Patients with T4 tumors that have involved an adjacent structure (e.g., bladder, small intestine, ovary, etc.) by direct extension from the primary tumor are eligible if all of the following conditions are met: All or a portion of the adjacent structure was removed en bloc with the primary tumor; In the opinion of the surgeon, all grossly visible tumor was completely resected ("curative resection"); Histologic evaluation by the pathologist confirms the margins of the resected specimen are not involved by malignant cells; and Local radiation therapy will not be utilized

[00271] Patients with more than one synchronous primary colon tumor are eligible; (staging classification will be based on the more advanced primary tumor)

[00272] Patients must have an ECOG performance status of 0 or 1

[00273] At the time of randomization, postoperative absolute granulocyte count (AGC) must be $\geq 1500/\text{mm}^3$ (or $< 1500/\text{mm}^3$, if in the opinion of the investigator, this represents an ethnic or racial variation of normal)

[00274] At the time of randomization, the postoperative platelet count must be $\geq 100,000/\text{mm}^3$

[00275] Bilirubin must be $\leq$ ULN for the lab unless the patient has a chronic grade 1 bilirubin elevation due to Gilbert's disease or similar syndrome due to slow conjugation of bilirubin

[00276] Alkaline phosphatase must be $< 2.5 \times$ ULN for the lab

[00277] AST must be $< 1.5 \times$ ULN for the lab. If AST is $> \text{ULN}$, serologic testing for hepatitis B and C must be performed and results must be negative

[00278] Serum creatinine $\leq 1.5 \times$ ULN for the lab

[00279] Urine protein/creatinine (UPC) ratio of < 1.0 ; patients with a UPC ratio ≥ 1.0 must undergo a 24-hour urine collection, which must be an adequate collection and must demonstrate < 1 gm of protein in the 24-hour urine collection in order to participate in the study

[00280] Patients with prior malignancies, including colorectal cancers, are eligible if they have been disease-free for ≥ 5 years and are deemed by their physician to be at low risk for recurrence; patients with squamous or basal cell carcinoma of the skin, melanoma in situ, carcinoma in situ of the cervix, or carcinoma in situ of the colon or rectum that have been effectively treated are eligible, even if these conditions were diagnosed within 5 years prior to randomization

[00281] Exclusion Criteria:

[00282] Patients < 18 years old

- [00283] Colon tumor other than adenocarcinoma
- [00284] Rectal tumors, i.e. a tumor located < 12 cm from the anal verge on endoscopy, or by surgical exam if the patient is not a candidate for endoscopy
- [00285] Isolated, distant, or non-contiguous intra-abdominal metastases, even if resected
- [00286] Any systemic or radiation therapy initiated for this malignancy
- [00287] Any significant bleeding that is not related to the primary colon tumor within 6 months before study entry
- [00288] Serious or non-healing wound, skin ulcers, or bone fracture
- [00289] Gastroduodenal ulcer(s) determined by endoscopy to be active
- [00290] Invasive procedures defined as follows: Major surgical procedure, open biopsy, or significant traumatic injury within 28 days prior to randomization; Anticipation of need for major surgical procedures during the course of the study; Core biopsy or other minor procedure, excluding placement of a vascular access device, within 7 days prior to randomization
- [00291] Uncontrolled blood pressure defined as > 150/90 mmHg
- [00292] History of TIA or CVA
- [00293] History of arterial thrombotic event within 12 months before study entry
- [00294] Symptomatic peripheral vascular disease
- [00295] PT/INR > 1.5, unless the patient is on therapeutic doses of warfarin. If so, the following criteria must be met for enrollment: The subject must have an in-range INR (usually between 2 and 3) on a stable dose of warfarin; The subject must not have active bleeding or a pathological condition that is associated with a high-risk of bleeding
- [00296] Concomitant halogenated antiviral agents
- [00297] Clinically significant peripheral neuropathy at the time of randomization (defined in the NCI Common Terminology Criteria for Adverse Events Version 3.0 [CTCAE v3.0] as grade 2 or greater neurosensory or neuromotor toxicity)
- [00298] Non-malignant systemic disease (cardiovascular, renal, hepatic, etc.) that would preclude any of the study therapy drugs; specifically excluded are the following cardiac conditions: New York Heart Association Class III or IV cardiac disease; History of myocardial infarction within 12 months before study entry; Unstable angina within 12 months before study entry; and Symptomatic arrhythmia
- [00299] History of chronic or persistent viral hepatitis or other chronic liver disease
- [00300] Psychiatric or addictive disorders or other conditions that, in the opinion of the investigator, would preclude the patient from meeting the study requirements

Example 4: Breast and Ovarian Cancer

[00301] Study Type: Interventional

[00302] Study Design: Allocation: Non-Randomized

[00303] Endpoint Classification: Safety Study

[00304] Intervention Model: Single Group Assignment

[00305] Masking: Open Label

[00306] Primary Purpose: Treatment

[00307] Primary Outcome Measures: Determine the safety and toxicity of the combination of ibrutinib and abexinostat in BRCA 1/2-associated recurrent breast and ovarian cancer patients

[00308] Secondary Outcome Measures: Assess clinical activity of the combination;

Detailed Description

[00309] Patients will be treated three times daily ibrutinib (140 mg/dose) and once daily with abexinostat for 4 consecutive days, followed by 3 days off of abexinostat (a cycle is 7 days). At day 1 of each cycle a physical exam and blood work will be performed. Reassessment of tumor size will be conducted at 6 weeks, 12 weeks and then every 9 weeks thereafter. Patients will remain on treatment until one of the following occur: disease progression, illness that prevents further treatment or unacceptable adverse events.

Eligibility

[00310] Ages Eligible for Study: 18 Years and older

[00311] Genders Eligible for Study: Both

[00312] Accepts Healthy Volunteers: No

[00313] Inclusion Criteria

[00314] Patients must have breast and/or epithelial ovarian cancer, primary peritoneal cancer, and/or fallopian tube cancer histologically or cytologically confirmed at the NCI that is metastatic or unresectable and for which standard curative measures do not exist or are no longer effective.

[00315] All patients in cohort 1 must have measurable and/or evaluable disease.

[00316] Patients in the expansion cohort 2 must have safely biopsible disease as determine by an interventional radiologist and must agree to the first mandatory biopsy (the other two biopsies optional).

[00317] Breast cancer patients with locally advanced, unresectable disease must have been previously treated with standard therapy.

[00318] There is no limit on number of prior therapy.

[00319] ECOG performance status less than or equal to 2 (Karnofsky greater than or equal to 60%).

[00320] Life expectancy greater than 3 months.

[00321] Patients must have normal organ and marrow function defined as follows: hemoglobin greater than or equal to 10g/dL; leukocytes greater than or equal to 3,000/mcL; absolute neutrophil count greater than or equal to 1,500/mcL; platelets greater than or equal to 100,000/mcL; total bilirubin less than or equal to upper limit of normal (ULN) in the absence of Gilbert's syndrome; AST(SGOT)/ALT(SGPT) less than or equal to 2.5 X ULN; creatinine clearance greater than or equal to 60 mL/min by 24-hour urine; corrected or Ionized Calcium less than or equal to ULN; potassium within normal limits

[00322] A documented deleterious BRCA 1/2 germline mutation or BRCAPRO score of greater than or equal to 30% for patients enrolling in Group A.

[00323] For patients enrolling in the sporadic serous epithelial ovarian cancer group, Group B, a negative family history (BRCAPRO score less than or equal to 20% or negative BRCA1/2 mutation test).

[00324] For patients enrolling in the triple negative breast cancer (ER-/PR- /Her2-) group, Group B, a negative family history and /or BRCAPRO score less than or equal to 10% or negative BRCA1/2 mutation test).

[00325] Exclusion Criteria

[00326] Patients who have had chemotherapy, biological therapy, hormonal therapy (with the exception of raloxifene or others approved for bone health) or radiotherapy within 4 weeks (6 weeks for nitrosoureas or mitomycin C) prior to entering the study.

[00327] Patients may not be receiving any other investigational agents or had them in the previous 28 days.

[00328] Patients with known brain metastases diagnosed within 1 year should be excluded from this clinical trial because of their poor prognosis and because they often develop progressive neurological dysfunction that would confound the evaluation of neurologic and other adverse events.

[00329] Clinically significant bleeding.

[00330] Inability to swallow capsules.

[00331] Uncontrolled intercurrent illness including, but not limited to, ongoing or active infection, symptomatic congestive heart failure, unstable angina pectoris, cardiac arrhythmia, or psychiatric illness/social situations that would limit compliance with study requirements.

[00332] Pregnant and breast-feeding women.

[00333] HIV-positive patients on combination antiretroviral therapy are ineligible because of the potential for pharmacokinetic interactions with AZD2281. In addition, these patients are at increased risk of lethal infections when treated with marrow-suppressive therapy such as carboplatin.

[00334] Major surgery within the past 28 days.

[00335] Patients with locally advanced breast tumors presenting for their initial therapy, or patients with local (only in breast or chest wall) recurrence only will not be eligible for this trial

[00336] For subjects in the dose-expansion cohorts, history of prior invasive malignancies within the past 5 years (with the exception of non-melanomatous skin cancers, non-invasive bladder cancer, stage I endometrial cancer or cervical cancer cured by surgical resection).

Example 5: Bladder Cancer

[00337] Study Type: Interventional

[00338] Study Design: Allocation: Randomized

[00339] Primary Purpose: Treatment with combination of ibrutinib and abexinostat

Objectives

[00340] I. Determine whether treatment with ibrutinib and abexinostat is effective in preventing recurrence of tumor after transurethral resection in patients with low grade, superficial transitional cell carcinoma of the bladder.

[00341] II. Determine the incidence and severity of toxicities associated with the long-term use of this drug in this patient population.

Eligibility

[00342] Ages Eligible for Study: 18 Years and older

[00343] Genders Eligible for Study: Both

[00344] Accepts Healthy Volunteers: No

Criteria

[00345] Disease Characteristics

[00346] Histologically confirmed low grade (grade 1 or 2), superficial (stage Ta or T1) transitional cell carcinoma (TCC) of the bladder

[00347] Newly diagnosed or recurrent

[00348] All visible tumor must have been resected within the past 12 weeks

[00349] Standard clinical management determined to be expectant observation without further surgery, intravesical therapy, or systemic therapy

[00350] No prior upper tract TCC

[00351] No history of grade 3 TCC, carcinoma in situ including severe dysplasia, non-TCC histology, or TCC greater than or equal to T2

[00352] No involvement of upper urinary tract prior to or at the time of initial tumor resection

[00353] Abdominal CT scan, IVP, or retrograde pyelogram within the past 3 months to rule out upper urinary tract tumor

[00354] Patient Characteristics

[00355] Not pregnant or nursing

[00356] Negative pregnancy test

[00357] Fertile patients must use effective contraception

[00358] No prior malignancy within the past 5 years and no concurrent malignancy except nonmelanomatous skin cancer or carcinoma in situ of the cervix

[00359] No clinically significant hearing loss (i.e., hearing loss effects everyday life and/or wears a hearing aide)

[00360] No other significant medical or psychiatric condition

Example 6: Lung Cancer

[00361] Study Type: Interventional

[00362] Study Design: Allocation: Randomized

[00363] Endpoint Classification: Safety/Efficacy Study

[00364] Masking: Double Blind (Subject, Investigator)

[00365] Primary Purpose: Treatment with ibrutinib and abexinostat

[00366] Primary Outcome Measures: Immune-related Progression-free Survival (irPFS) in Participants With Nonsmall-cell Lung Cancer (NSCLC) Per Immune-related Response Criteria (irRC) [Time Frame: Tumor assessed at screening, every 6 weeks on treatment to Week 24, and every 12 weeks on maintenance until immune-related Progressive Disease (irPD) or death (of censored, maximum reached: 16.5 months)]

[00367] irPFS is defined as the time between the randomization date and date of immune-related Progressive Disease (irPD) (at least 25% increase percentage change in total tumor burden, including new lesions) or death, whichever occurs first. For patients with no recorded postbaseline tumor assessments, irPFS is censored at randomization. Participant who die without reported irPD are considered to have progressed on the date of death. For those who remain alive and have no irPD, irPFS is censored on the date of last evaluable tumor assessment.

Independent review committee performed tumor assessment

Detailed Description

[00368] Patients will be treated three times daily ibrutinib (140 mg/dose) and once daily with abexinostat for 4 consecutive days, followed by 3 days off of abexinostat (a cycle is 7 days). At day 1 of each cycle a physical exam and blood work will be performed. Reassessment of tumor size will be conducted at 6 weeks, 12 weeks and then every 9 weeks thereafter. Patients will remain on treatment until one of the following occur: disease progression, illness that prevents further treatment or unacceptable adverse events.

[00369] Inclusion Criteria

[00370] Histologically or cytologically confirmed lung cancer (Stage IIIb/IV nonsmall-cell lung cancer or extensive stage small-cell lung cancer [SCLC])

[00371] Measurable tumor lesion (as long as it is not located in a previously irradiated area) as defined by modified World Health Organization criteria

[00372] Eastern Cooperative Oncology Group performance status of ≤ 1 at study entry

[00373] Accessible for treatment and follow-up

[00374] Exclusion Criteria:

[00375] Brain metastases

[00376] Malignant pleural effusion

[00377] Autoimmune disease

[00378] Motor neuropathy of autoimmune origin

[00379] SCLC-related paraneoplastic syndromes

[00380] Any concurrent malignancy other than nonmelanoma skin cancer; carcinoma in situ of the cervix or breast; or prostate cancer treated with systemic therapy (participants with a previous malignancy but without evidence of disease for 5 years were allowed to enter the study)

[00381] Prior systemic therapy for lung cancer. Prior radiation therapy or locoregional surgeries performed later than at least 3 weeks prior to randomization date were allowed.

[00382] Known HIV or hepatitis B or C infection

[00383] Chronic use of immunosuppressants and/or systemic corticosteroids (used in the management of cancer or noncancer-related illnesses). However, use of corticosteroids was allowed if used as premedication for paclitaxel infusion or for treating immune-related adverse events or adrenal insufficiencies.

[00384] Inadequate hematologic function defined by an absolute neutrophil count $< 1,500/\text{mm}^3$, a platelet count $< 100,000/\text{mm}^3$, or hemoglobin level $< 9 \text{ g/dL}$.

[00385] Inadequate hepatic function defined by a total bilirubin level >2.0 times the upper limit of normal (ULN), or ≥ 2.5 times the ULN if liver metastases are present, aspartate aminotransferase and alanine aminotransferase levels ≥ 2.5 times the ULN or ≥ 5 times the ULN if liver metastases are present.

[00386] Inadequate renal function defined by a serum creatinine level ≥ 2.5 times the ULN

[00387] Inadequate creatinine clearance defined as less than 50 mL/min.

Example 7: Prostate Cancer

[00388] Study Type: Interventional

[00389] Study Design: Endpoint Classification: Safety/Efficacy Study

[00390] Intervention Model: Single Group Assignment

[00391] Masking: Open Label

[00392] Primary Purpose: Treatment

[00393] Primary Outcome Measures: PSA response rate with abexinostat and ibrutinib therapy in androgen independent non-metastatic prostate cancer [Time Frame: An average every 6 weeks for up to 3 months]

[00394] Secondary Outcome Measures: Overall survival of androgen independent non-metastatic prostate cancer patients treated with abexinostat and ibrutinib [Time Frame: Every 3 months]

Detailed Description

[00395] Patients will be treated three times daily ibrutinib (140 mg/dose) and once daily with abexinostat for 4 consecutive days, followed by 3 days off of abexinostat (a cycle is 7 days). At day 1 of each cycle a physical exam and blood work will be performed. Reassessment of tumor size will be conducted at 6 weeks, 12 weeks and then every 9 weeks thereafter. Patients will remain on treatment until one of the following occur: disease progression, illness that prevents further treatment or unacceptable adverse events.

Eligibility

[00396] Ages Eligible for Study: 18 Years and older

[00397] Genders Eligible for Study: Both

[00398] Accepts Healthy Volunteers: No

Inclusion Criteria:

[00399] A histologic diagnosis of prostate adenocarcinoma.

[00400] No evidence of bone/visceral metastases as visualized on standard imaging such as bone scan, chest X-ray, CT scan or MRI of abdomen and pelvis.

[00401] PSA-only progression despite androgen deprivation therapy. PSA progression is defined as 3 rising levels, with a minimum interval of 2 weeks between each determination. The last determination must have a minimum value of 1ng/ml and be determined within two weeks prior to registration. If the second or third confirmatory value is less than the previous value, the patient will still be eligible if a repeat value (No. 4) is found to be greater than all the prior values.

[00402] If patient has been on antiandrogen in the past 28 days, then PSA progression after withdrawal period (28 days for flutamide and 42 days for bicalutamide or nilutamide) is required.

[00403] ECOG performance status of 0-1.

[00404] No investigational or commercial agents or therapies (except LHRH agonists) may be administered concurrently with the intent to treat the patient's malignancy. Patients on LHRH agonists must continue the use of LHRH agonist therapy. Bisphosphonates can be administered per treating physician discretion.

[00405] At least 4 weeks must have elapsed since prior systemic therapy, except for LHRH analogue therapy and steroids. If steroids are being used for therapy of prostate cancer, these should be discontinued prior to starting avastin therapy.

[00406] Life expectancy of at least 6 months.

[00407] Ability to understand and the willingness to sign a written informed consent that is approved by the Institutional Human Investigation Committee.

[00408] Use of effective means of contraception in subjects.

Exclusion Criteria:

[00409] Inability to comply with study and/or follow-up procedures.

[00410] Inadequately controlled hypertension (defined as systolic blood pressure >150 and/or diastolic blood pressure > 100 mmHg on antihypertensive medications).

[00411] Any prior history of hypertensive crisis or hypertensive encephalopathy.

[00412] New York Heart Association (NYHA) Grade II or greater congestive heart failure

[00413] History of myocardial infarction or unstable angina within last 12 months prior to study enrollment.

[00414] History of stroke or transient ischemic attack within 6 months prior to study enrollment.

[00415] Known CNS disease.

[00416] Significant vascular disease (e.g., aortic aneurysm, aortic dissection).

[00417] Symptomatic peripheral vascular disease.

[00418] Evidence of bleeding diathesis or coagulopathy.

[00419] Patients on anticoagulants are allowed if patient has been on therapy for at least 4 weeks and patient has no acute thromboembolic activity.

[00420] Major surgical procedure, open biopsy, or significant traumatic injury within 28 days prior to study enrollment or anticipation of need for major surgical procedure during the course of the study.

[00421] Core biopsy or other minor surgical procedure, excluding placement of a vascular access device, within 7 days prior to study enrollment.

[00422] History of abdominal fistula, gastrointestinal perforation, or intra-abdominal abscess within 6 months prior to study enrollment.

[00423] Serious, non-healing wound, ulcer, or bone fracture.

[00424] Proteinuria at screening as demonstrated by: Urine protein:creatinine (UPC) ratio ≥ 1.0 at screening

[00425] Refusal to use effective means of contraception.

[00426] Patients with known brain metastases should be excluded from this clinical trial because of their poor prognosis and because they often develop progressive neurologic dysfunction that would confound the evaluation of neurologic and other adverse events.

[00427] History of allergic reactions attributed to compounds of similar chemical or biologic composition to avastin.

[00428] Uncontrolled intercurrent illness including, but not limited to, ongoing or active infection, symptomatic congestive heart failure, unstable angina pectoris, cardiac arrhythmia, or psychiatric illness/social situations that would limit compliance with study requirements.

[00429] Patients with immune deficiency such as HIV-positive patients or those receiving combination anti-retroviral therapy are excluded from the study because of lack of safety data for avastin in these patients.

Example 8: Bile Duct Cancer

[00430] Study Type: Interventional

[00431] Study Design: Endpoint Classification: Safety/Efficacy Study

[00432] Intervention Model: Single Group Assignment

[00433] Masking: Open Label

[00434] Primary Purpose: Treatment

[00435] Primary Outcome Measures: Overall survival with abexinostat and ibrutinib therapy [Time Frame: 3 years]

[00436] Secondary Outcome Measures: Response rate defined as proportion of patients with complete response (CR), partial response (PR), stable disease (SD), or progressive disease (PD), using based on the RECIST version 1.1

Detailed Description

[00437] Patients will be treated three times daily ibrutinib (140 mg/dose) and once daily with abexinostat for 4 consecutive days, followed by 3 days off of abexinostat (a cycle is 7 days). At day 1 of each cycle a physical exam and blood work will be performed. Reassessment of tumor size will be conducted at 6 weeks, 12 weeks and then every 9 weeks thereafter. Patients will remain on treatment until one of the following occur: disease progression, illness that prevents further treatment or unacceptable adverse events.

Eligibility

[00438] Ages Eligible for Study: 18 Years and older

[00439] Genders Eligible for Study: Both

[00440] Accepts Healthy Volunteers: No

Inclusion Criteria:

[00441] Histologically and cytologically proven cholangiocarcinoma of any type (including intrahepatic cholangiocarcinoma, extrahepatic primary cholangiocarcinoma, hilar cholangiocarcinomas, cholangiocarcinomas located in the gall bladder or hepatic capsule effraction, and carcinoma of the Ampulla of Vater, etc.) that is not amenable to surgery, radiation, or combined modality therapy with curative intent, and has failed or is not eligible for available chemotherapies such as gemcitabine with or without platinum

[00442] Local, locally-advanced, or metastatic disease documented as having shown progression on a scan (e.g., computed tomography [CT], magnetic resonance imaging [MRI])

[00443] Measurable tumor according to Response Evaluation Criteria in Solid Tumors (RECIST) 1.1 criteria with at least one unidimensionally measurable target lesion

[00444] No evidence of biliary duct obstruction, unless obstruction is controlled by local treatment or, in whom the biliary tree can be decompressed by endoscopic or percutaneous stenting with subsequent reduction in bilirubin to below 1.5 x upper level of normal (ULN)

[00445] Eastern Cooperative Oncology Group (ECOG) performance status being 0-3

[00446] Expected survival > 3 months

[00447] Women of child-bearing potential (i.e., women who are pre-menopausal or not surgically sterile) must use accepted contraceptive methods (abstinence, intrauterine device [IUD], oral contraceptive or double barrier device) during the study, and must have a negative serum or urine pregnancy test within 1 week prior to treatment initiation

[00448] Fertile men must practice effective contraceptive methods during the study, unless documentation of infertility exists

[00449] Granulocyte count $\geq 1500/\text{mm}^3$

[00450] White blood cell (WBC) $\geq 3500 \text{ cells}/\text{mm}^3$ or $\geq 3.5 \text{ bil}/\text{L}$

[00451] Platelet count $\geq 150,000 \text{ cells}/\text{mm}^3$ or $\geq 150 \text{ bil}/\text{L}$

[00452] Absolute neutrophil count (ANC) $\geq 1500 \text{ cells}/\text{mm}^3$ or $\geq 1.5 \text{ bil}/\text{L}$

[00453] Hemoglobin $\geq 9 \text{ g}/\text{dL}$ or $\geq 90 \text{ g}/\text{L}$

[00454] Aspartate aminotransferase (AST/serum glutamic oxaloacetic transaminase [SGOT]) $\leq 3 \times$ upper normal limit (UNL), alanine aminotransferase (ALT/serum glutamate pyruvate transaminase [SGPT]) $\leq 3 \times$ UNL ($\leq 5 \times$ UNL if liver metastases present)

[00455] Bilirubin $\leq 1.5 \times$ UNL

[00456] Serum creatinine $\leq 2.0 \text{ mg}/\text{dL}$ or $177 \mu\text{mol}/\text{L}$

[00457] International normalized ratio or INR must be ≤ 1.5

[00458] No evidence of active infection and no serious infection within the past month

[00459] Mentally competent, ability to understand and willingness to sign the informed consent form

Exclusion Criteria:

[00460] Patients receiving any other standard or investigational treatment for their cancer

[00461] Serious medical illness that would potentially increase patients' risk for toxicity

[00462] Any active uncontrolled bleeding, and any patients with a bleeding diathesis (e.g., active peptic ulcer disease)

[00463] Pregnant women, or women of child-bearing potential not using reliable means of contraception

[00464] Lactating females

[00465] Fertile men unwilling to practice contraceptive methods during the study period

[00466] Life expectancy less than 3 months

[00467] Unwilling or unable to follow protocol requirements

[00468] Dyspnea with moderate exertion; patients with pleural or pericardial effusions

[00469] Active heart disease including but not limited to symptomatic congestive heart failure, symptomatic coronary artery disease, symptomatic angina pectoris, symptomatic myocardial infarction, arrhythmias requiring medication, or symptomatic congestive heart failure; also patients with a history of myocardial infarction that is < 1 year prior to registration, or patients with previous congestive heart failure (< 1 year prior to registration) requiring pharmacologic support or with left ventricular ejection fraction $< 50\%$

[00470] A marked baseline prolongation of QT/corrected QT (QTc) interval (e.g., repeated exhibition of a QTc interval > 470 ms)

[00471] A history of additional risk factors for torsade de pointes (e.g., heart failure, hypokalemia, family history of long QT syndrome)

[00472] Evidence of active infection, or serious infection within the past month

[00473] Patients with known human immunodeficiency virus (HIV) infection

[00474] Requirement for immediate palliative treatment of any kind including surgery

[00475] Patients that have received a chemotherapy regimen with stem cell support in the previous 6 months

[00476] Prior illicit drug addiction

[00477] Any condition or abnormality which may, in the opinion of the investigator, compromise the safety of the patient

Example 9: Large Cell Lung Cancer

[00478] Study Type: Interventional

[00479] Study Design: Endpoint Classification: Safety/Efficacy Study

[00480] Intervention Model: Single Group Assignment

[00481] Masking: Open Label

[00482] Primary Purpose: Treatment

[00483] Primary Outcome Measures: The proportion of subjects progression-free at Month 3. Time Frame [Time Frame: 3 months]

[00484] Secondary Outcome Measures: the proportion of subjects progression-free at Month 6 [Time Frame: 6 months]; overall response rate (ORR) is the proportion of patients with a best overall response of Complete Response (CR) or Partial Response (PR) at Month 3 [Time Frame: 3 months]; disease control rate (DCR) is the proportion of patients with a best overall response of CR or PR or Stable Disease (SD) at Month 3 (C4D21) [Time Frame: 3 months]; progression Free Survival [Time Frame: approximately 3-6 months]; overall Survival [Time Frame: estimated 12 months]

Detailed Description

[00485] Patients will be treated three times daily ibrutinib (140 mg/dose) and once daily with abexinostat for 4 consecutive days, followed by 3 days off of abexinostat (a cycle is 7 days). At day 1 of each cycle a physical exam and blood work will be performed. Reassessment of tumor size will be conducted at 6 weeks, 12 weeks and then every 9 weeks thereafter. Patients will remain on treatment until one of the following occur: disease progression, illness that prevents further treatment or unacceptable adverse events.

Eligibility

[00486] Ages Eligible for Study: 18 Years and older

[00487] Genders Eligible for Study: Both

[00488] Accepts Healthy Volunteers: No

Inclusion Criteria:

[00489] Written informed consent obtained according to local guidelines

[00490] Histologically confirmed diagnosis of stage IV lung cancer of LC-NEC type according to WHO classification:

- Histological analysis of newly diagnosed disease must not be older than 8 weeks from signed consent
- Relapse must be confirmed by histology
- Neuroendocrine differentiation

[00491] 3. World Health Organisation (WHO) performance status grade ≤ 1

[00492] 4. measurable disease

[00493] 5. Adequate bone marrow function

[00494] 6. Adequate liver function

[00495] 7. Adequate renal function

Exclusion Criteria:

[00496] Clinical evidence of central nervous system (CNS) metastases.

[00497] Presence of SCLC cells

[00498] Patients who have a history of another primary malignancy ≤ 3 years, with the exception of inactive basal or squamous cell carcinoma of the skin or cervical cancer in situ, early stages of breast cancer (LCIS and DCIS) and prostate cancer (stage T1a)


[00499] Prior chemotherapy for the treatment of advanced lung cancer and/or not having recovered from the side effects of any other therapy (adjuvant treatment for earlier stages I-III is allowed if finished at least one year before study entry)

[00500] Treatment with any investigational drug ≤ 28 days before starting study treatment or who have not recovered from side effects of such therapy

[00501] Women who are pregnant or lactating

WHAT IS CLAIMED IS:

1. A composition comprising a therapeutically effective amount of an ACK inhibitor compound, a therapeutically effective amount of an HDAC inhibitor compound, and a pharmaceutically acceptable excipient.
2. The composition of claim 1, wherein the ACK inhibitor compound is a BTK inhibitor.
3. The composition of claim 2, wherein the BTK inhibitor is an irreversible BTK inhibitor.
4. The composition of claim 1, wherein the ACK inhibitor is a compound of Formula (A):



Formula (A)

wherein

A is independently selected from N or CR₅;

R₁ is H, L₂-(substituted or unsubstituted alkyl), L₂-(substituted or unsubstituted cycloalkyl), L₂-(substituted or unsubstituted alkenyl), L₂-(substituted or unsubstituted cycloalkenyl), L₂-(substituted or unsubstituted heterocycle), L₂-(substituted or unsubstituted heteroaryl), or L₂-(substituted or unsubstituted aryl), where L₂ is a bond, O, S, -S(=O), -S(=O)₂, C(=O), -(substituted or unsubstituted C₁-C₆ alkyl), or -(substituted or unsubstituted C₂-C₆ alkenyl);

R₂ and R₃ are independently selected from H, lower alkyl and substituted lower alkyl;

R₄ is L₃-X-L₄-G, wherein,


L₃ is optional, and when present is a bond, optionally substituted or unsubstituted alkyl, optionally substituted or unsubstituted cycloalkyl, optionally substituted or unsubstituted alkenyl, optionally substituted or unsubstituted alkynyl;

X is optional, and when present is a bond, O, -C(=O), S, -S(=O), -S(=O)₂, -NH, -NR₉, -NHC(O), -C(O)NH, -NR₉C(O), -C(O)NR₉, -S(=O)₂NH, -NHS(=O)₂, -S(=O)₂NR₉-, -NR₉S(=O)₂-, -OC(O)NH-, -NHC(O)O-, -OC(O)NR₉-, -NR₉C(O)O-, -CH=NO-, -ON=CH-, -NR₁₀C(O)NR₁₀-, heteroaryl, aryl, -NR₁₀C(=NR₁₁)NR₁₀-, -NR₁₀C(=NR₁₁)-, -C(=NR₁₁)NR₁₀-, -OC(=NR₁₁)-, or -C(=NR₁₁)O-;

L₄ is optional, and when present is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted

heteroaryl, substituted or unsubstituted heterocycle;

or L₃, X and L₄ taken together form a nitrogen containing heterocyclic ring;



R₆, R₇ and R₈ are independently selected from among H, lower alkyl or substituted lower alkyl, lower heteroalkyl or substituted lower heteroalkyl, substituted or unsubstituted lower cycloalkyl, and substituted or unsubstituted lower heterocycloalkyl;

R₅ is H, halogen, -L₆-(substituted or unsubstituted C₁-C₃ alkyl), -L₆-(substituted or unsubstituted C₂-C₄ alkenyl), -L₆-(substituted or unsubstituted heteroaryl), or -L₆-(substituted or unsubstituted aryl), wherein L₆ is a bond, O, S, -S(=O), S(=O)₂, NH, C(O), -NHC(O)O, -OC(O)NH, -NHC(O), or -C(O)NH;

each R₉ is independently selected from among H, substituted or unsubstituted lower alkyl, and substituted or unsubstituted lower cycloalkyl;


each R₁₀ is independently H, substituted or unsubstituted lower alkyl, or substituted or unsubstituted lower cycloalkyl; or

two R₁₀ groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or

R₉ and R₁₀ can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or

each R₁₁ is independently selected from H, -S(=O)₂R₈, -S(=O)₂NH₂, -C(O)R₈, -CN, -NO₂, heteroaryl, or heteroalkyl.


5. The composition of claim 1, wherein the ACK inhibitor is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib).



Ibrutinib.

6. The composition of claim 1, wherein the ACK inhibitor is AVL-263 (Avila Therapeutics/Celgene Corporation), AVL-292 (Avila Therapeutics/Celgene Corporation), AVL-291 (Avila Therapeutics/Celgene Corporation), BMS-488516 (Bristol-Myers Squibb), BMS-509744 (Bristol-Myers Squibb), CGI-1746 (CGI Pharma/Gilead Sciences), CTA-056, GDC-0834 (Genentech), HY-11066 (also, CTK4I7891, HMS3265G21, HMS3265G22, HMS3265H21, HMS3265H22, 439574-61-5, AG-F-54930), ONO-4059 (Ono Pharmaceutical Co., Ltd.), ONO-WG37 (Ono Pharmaceutical Co., Ltd.), PLS-123 (Peking University), RN486 (Hoffmann-La Roche), or HM71224 (Hanmi Pharmaceutical Company Limited).

7. The composition of claim 1, wherein the HDAC inhibitor is a compound of Formula (B):



Formula (B)

wherein:

R¹ is hydrogen or alkyl;

X is -O-, -NR²-, or -S(O)_n where n is 0-2 and R² is hydrogen or alkyl;


Y is alkylene optionally substituted with cycloalkyl, optionally substituted phenyl, alkylthio, alkylsulfinyl, alkylsulfonyl, optionally substituted phenylalkylthio, optionally substituted phenylalkylsulfonyl, hydroxy, or optionally substituted phenoxy;

Ar¹ is phenylene or heteroarylene wherein said Ar¹ is optionally substituted with one or two groups independently selected from alkyl, halo, hydroxy, alkoxy, haloalkoxy, or haloalkyl;

R³ is hydrogen, alkyl, hydroxyalkyl, or optionally substituted phenyl; and


Ar² is aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, heteroaralkenyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, or heterocycloalkylalkyl.

8. The composition of claim 1, wherein the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat)



Abexinostat.


9. The composition of claim 1, wherein the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2- carboxamide (i.e. PCI-24781/abexinostat)



Abexinostat

and,


the ACK inhibitor is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib)



Ibrutinib.

10. The composition of claim 1, in the form of a solid dosage form.
11. The composition of claim 10, in the form of a capsule.
12. The composition of claim 1, in the form of a solution.
13. The composition of claim 12, in the form of a solution for intravenous administration.
14. The composition of claim 1, comprising 140 mg of ibrutinib.
15. The composition of claim 1 for use in treatment of a solid tumor.
16. The composition of claim 1 for use in treatment of a carcinoma.
17. The composition of claim 16, wherein the carcinoma is breast cancer.
18. The composition of claim 17, wherein the breast cancer is invasive ductal carcinoma, ductal carcinoma in situ, invasive lobular carcinoma, or lobular carcinoma in situ.
19. The composition of claim 16, wherein the carcinoma is pancreatic cancer.
20. The composition of claim 19, wherein the pancreatic cancer is adenocarcinoma, or islet cell carcinoma.
21. The composition of claim 16, wherein the carcinoma is colorectal cancer.
22. The composition of claim 21, wherein the colorectal cancer is adenocarcinoma.
23. The composition of claim 15, wherein the solid tumor is a colon polyp.
24. The composition of claim 23, wherein the colon polyp is associated with familial adenomatous polyposis.
25. The composition of claim 16, wherein the carcinoma is bladder cancer.
26. The composition of claim 25, wherein the bladder cancer is transitional cell bladder cancer, squamous cell bladder cancer, or adenocarcinoma.
27. The composition of claim 16, wherein the carcinoma is lung cancer.
28. The composition of claim 27, wherein the lung cancer is a non-small cell lung cancer.

29. The composition of claim 28, wherein the non-small cell lung cancer is adenocarcinoma, squamous-cell lung carcinoma, or large-cell lung carcinoma.
30. The composition of claim 28, wherein the non-small cell lung cancer is large cel lung cancer.
31. The composition of claim 16, wherein the carcinoma is prostate cancer.
32. The composition of claim 31, wherein the prostate cancer is adenocarcinoma or small cell carcinoma.
33. The composition of claim 16, wherein the carcinoma is ovarian cancer.
34. The composition of claim 33, wherein the ovarian cancer is epithelial ovarian cancer.
35. The composition of claim 16, wherein the carcinoma is bile duct cancer.
36. The composition of claim 35, wherein the bile duct cancer is proximal bile duct carcinoma or distal bile duct carcinoma.
37. The composition of claim 1, wherein combination of ibrutinib and abexinostat is 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55% or 60% more efficacious than administration of abexinostat alone.
38. The composition of claim 37, wherein combination of the ibrutinib and abexinostat is 50% more efficacious than administration of abexinostat alone.
39. The composition of claim 37, wherein combination of ibrutinib and abexinostat is 5%, 10%, 15%, 20%, 25%, 30%, or 35% more efficacious than administration of ibrutinib alone.
40. The composition of claim 37, wherein combination of ibrutinib and abexinostat is 25% more efficacious than administration of ibrutinib alone.
41. Combination of a therapeutically effective amount of an ACK inhibitor compound and a therapeutically effective amount of an HDAC inhibitor compound.
42. The combination of claim 41, wherein the ACK inhibitor compound is a BTK inhibitor.
43. The combination of claim 42, wherein the BTK inhibitor is an irreversible BTK inhibitor.
44. The combination of claim 41, wherein the ACK inhibitor is a compound of Formula (A):



Formula (A)

wherein

A is independently selected from N or CR₅;

R_1 is H, L_2 -(substituted or unsubstituted alkyl), L_2 -(substituted or unsubstituted cycloalkyl), L_2 -(substituted or unsubstituted alkenyl), L_2 -(substituted or unsubstituted cycloalkenyl), L_2 -(substituted or unsubstituted heterocycle), L_2 -(substituted or unsubstituted heteroaryl), or L_2 -(substituted or unsubstituted aryl), where L_2 is a bond, O, S, -S(=O), -S(=O)₂, C(=O), -(substituted or unsubstituted C₁-C₆ alkyl), or -(substituted or unsubstituted C₂-C₆ alkenyl);

R_2 and R_3 are independently selected from H, lower alkyl and substituted lower alkyl;


R_4 is L_3 -X- L_4 -G, wherein,

L_3 is optional, and when present is a bond, optionally substituted or unsubstituted alkyl, optionally substituted or unsubstituted cycloalkyl, optionally substituted or unsubstituted alkenyl, optionally substituted or unsubstituted alkynyl;

X is optional, and when present is a bond, O, -C(=O), S, -S(=O), -S(=O)₂, -NH, -NR₉, -NHC(O), -C(O)NH, -NR₉C(O), -C(O)NR₉, -S(=O)₂NH, -NHS(=O)₂, -S(=O)₂NR₉-, -NR₉S(=O)₂, -OC(O)NH-, -NHC(O)O-, -OC(O)NR₉-, -NR₉C(O)O-, -CH=NO-, -ON=CH-, -NR₁₀C(O)NR₁₀-, heteroaryl, aryl, -NR₁₀C(=NR₁₁)NR₁₀-, -NR₁₀C(=NR₁₁)-, -C(=NR₁₁)NR₁₀-, -OC(=NR₁₁)-, or -C(=NR₁₁)O-;

L_4 is optional, and when present is a bond, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycle;

or L_3 , X and L_4 taken together form a nitrogen containing heterocyclic ring;




R_6 , R_7 and R_8 are independently selected from among H, lower alkyl or substituted lower alkyl, lower heteroalkyl or substituted lower heteroalkyl, substituted or unsubstituted lower cycloalkyl, and substituted or unsubstituted lower heterocycloalkyl;

R_5 is H, halogen, - L_6 -(substituted or unsubstituted C₁-C₃ alkyl), - L_6 -(substituted or unsubstituted C₂-C₄ alkenyl), - L_6 -(substituted or unsubstituted heteroaryl), or - L_6 -(substituted or unsubstituted aryl), wherein L_6 is a bond, O, S, -S(=O), S(=O)₂, NH, C(O), -NHC(O)O, -OC(O)NH, -NHC(O), or -C(O)NH;

each R_9 is independently selected from among H, substituted or unsubstituted lower alkyl, and substituted or unsubstituted lower cycloalkyl;

each R_{10} is independently H, substituted or unsubstituted lower alkyl, or substituted or unsubstituted lower cycloalkyl; or
 two R_{10} groups can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or
 R_9 and R_{10} can together form a 5-, 6-, 7-, or 8-membered heterocyclic ring; or
 each R_{11} is independently selected from H, $-S(=O)_2R_8$, $-S(=O)_2NH_2$, $-C(O)R_8$, $-CN$, $-NO_2$, heteroaryl, or heteroalkyl.


45. The combination of claim 41, wherein the ACK inhibitor is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib).



Ibrutinib.

46. The combination of claim 41, wherein the ACK inhibitor is AVL-263 (Avila Therapeutics/Celgene Corporation), AVL-292 (Avila Therapeutics/Celgene Corporation), AVL-291 (Avila Therapeutics/Celgene Corporation), BMS-488516 (Bristol-Myers Squibb), BMS-509744 (Bristol-Myers Squibb), CGI-1746 (CGI Pharma/Gilead Sciences), CTA-056, GDC-0834 (Genentech), HY-11066 (also, CTK4I7891, HMS3265G21, HMS3265G22, HMS3265H21, HMS3265H22, 439574-61-5, AG-F-54930), ONO-4059 (Ono Pharmaceutical Co., Ltd.), ONO-WG37 (Ono Pharmaceutical Co., Ltd.), PLS-123 (Peking University), RN486 (Hoffmann-La Roche), or HM71224 (Hanmi Pharmaceutical Company Limited).

47. The combination of claim 41, wherein the HDAC inhibitor is a compound of Formula (B):



Formula (B)

wherein:

R^1 is hydrogen or alkyl;

X is -O-, -NR²-, or -S(O)_n where n is 0-2 and R² is hydrogen or alkyl;


Y is alkylene optionally substituted with cycloalkyl, optionally substituted phenyl, alkylthio, alkylsulfinyl, alkylsulfonyl, optionally substituted phenylalkylthio, optionally substituted phenylalkylsulfonyl, hydroxy, or optionally substituted phenoxy;

Ar¹ is phenylene or heteroarylene wherein said Ar¹ is optionally substituted with one or two groups independently selected from alkyl, halo, hydroxy, alkoxy, haloalkoxy, or haloalkyl;

R³ is hydrogen, alkyl, hydroxyalkyl, or optionally substituted phenyl; and


Ar² is aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, heteroaralkenyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, or heterocycloalkylalkyl.

48. The combination of claim 41, wherein the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat)



Abexinostat.


49. The combination of claim 41, wherein the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat)



Abexinostat

and,

the ACK inhibitor is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib)




Ibrutinib.

50. The combination of claim 41 administered in a solid dosage form.
51. The combination of claim 41 administered in a capsule.
52. The combination of claim 41 administered in a solution.
53. The combination of claim 41 administered in a solution for intravenous administration.
54. The combination of claim 41, comprising 140 mg of ibrutinib.
55. The combination of claim 41 for use in treatment of a solid tumor.
56. The combination of claim 41 for use in treatment of a sarcoma or carcinoma.
57. The combination of claim 41 for use in treatment of a carcinoma.
58. The combination of claim 57, wherein the carcinoma is breast cancer.


59. The combination of claim 58, wherein the breast cancer is invasive ductal carcinoma, ductal carcinoma in situ, invasive lobular carcinoma, or lobular carcinoma in situ.
60. The combination of claim 57, wherein the carcinoma is pancreatic cancer.
61. The combination of claim 60, wherein the pancreatic cancer is adenocarcinoma, or islet cell carcinoma.
62. The combination of claim 57, wherein the carcinoma is colorectal cancer.
63. The combination of claim 62, wherein the colorectal cancer is adenocarcinoma.
64. The combination of claim 55, wherein the solid tumor is a colon polyp.
65. The combination of claim 64, wherein the colon polyp is associated with familial adenomatous polyposis.
66. The combination of claim 57, wherein the carcinoma is bladder cancer.
67. The combination of claim 66, wherein the bladder cancer is transitional cell bladder cancer, squamous cell bladder cancer, or adenocarcinoma.
68. The combination of claim 57, wherein the carcinoma is lung cancer.
69. The combination of claim 68, wherein the lung cancer is a non-small cell lung cancer.
70. The combination of claim 69, wherein the non-small cell lung cancer is adenocarcinoma, squamous-cell lung carcinoma, or large-cell lung carcinoma.
71. The combination of claim 69, wherein the non-small cell lung cancer is large cell lung cancer.
72. The combination of claim 57, wherein the carcinoma is prostate cancer.
73. The combination of claim 72, wherein the prostate cancer is adenocarcinoma or small cell carcinoma.
74. The combination of claim 57, wherein the carcinoma is ovarian cancer.
75. The combination of claim 74, wherein the ovarian cancer is epithelial ovarian cancer.
76. The combination of claim 57, wherein the carcinoma is bile duct cancer.
77. The combination of claim 76, wherein the bile duct cancer is proximal bile duct carcinoma or distal bile duct carcinoma.
78. The combination of claim 41, wherein combination of ibrutinib and abexinostat is 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55% or 60% more efficacious than administration of abexinostat alone.
79. The combination of claim 78, wherein combination of the ibrutinib and abexinostat is 50% more efficacious than administration of abexinostat alone.
80. The combination of claim 78, wherein combination of ibrutinib and abexinostat is 5%, 10%, 15%, 20%, 25%, 30%, or 35% more efficacious than administration of ibrutinib alone.

81. The combination of claim 78, wherein combination of ibrutinib and abexinostat is 25% more efficacious than administration of ibrutinib alone.
82. The combination of claim 41, administered in a unified dosage form or separate dosage forms.
83. The combination of claim 41 administered simultaneously or sequentially.
84. A method for treating a solid tumor comprising co-administering to an individual in need thereof an ACK inhibitor compound and an HDAC inhibitor compound.
85. The method of claim 84, wherein the ACK inhibitor is (R)-1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)piperidin-1-yl)prop-2-en-1-one (i.e. PCI-32765/ibrutinib)



Ibrutinib.

86. The method of claim 84, wherein the HDAC inhibitor is 3-[(dimethylamino)methyl]-N-{2-[4-(hydroxycarbamoyl)phenoxy]ethyl}-1-benzofuran-2-carboxamide (i.e. PCI-24781/abexinostat)



Abexinostat.

87. The method of claim 84, wherein the solid tumor is a carcinoma.
88. The method of claim 87, wherein the carcinoma is breast cancer.
89. The method of claim 87, wherein the carcinoma is pancreatic cancer.
90. The method of claim 87, wherein the carcinoma is colorectal cancer.
91. The method of claim 87, wherein the carcinoma is bladder cancer.
92. The method of claim 87, wherein the carcinoma is lung cancer.
93. The method of claim 92, wherein the lung cancer is a non-small cell lung cancer.
94. The method of claim 92, wherein the lung cancer is a large cell lung cancer.
95. The method of claim 87, wherein the carcinoma is prostate cancer.
96. The method of claim 87, wherein the carcinoma is ovarian cancer.
97. The method of claim 87, wherein the carcinoma is bile duct cancer.
98. The method of claim 84, wherein co-administration of the ACK inhibitor and the HDAC inhibitor is 50% more efficacious than administration of the HDAC inhibitor alone.
99. The method of claim 84, wherein co-administration of the ACK inhibitor and the HDAC inhibitor is 25% more efficacious than administration of the ACK inhibitor alone.
100. The method of claim 84, wherein the ACK inhibitor and the HDAC inhibitor are administered in a unified dosage form or separate dosage forms.
101. The method of claim 84, wherein the ACK inhibitor and the HDAC inhibitor are administered simultaneously or sequentially.

1/16

FIG. 1






FIG. 2




3/16

FIG. 3




4/16

FIG. 4




5/16

FIG. 5




6/16

FIG. 6



7/16


FIG. 7



8/16

FIG. 8


A



Control


4 tumors in photo, 8 tumors total

B




HDACi

1 small tumor, 2 tumors total



BTKi

1 small tumor, 1 tumor total




BTKi + HDACi

no visible tumors

C


D

9/16

FIG. 9

10/16


FIG. 10



11/16
FIG. 11


Cell Line Name	Cancer Type	Combination Effects of ibrutinib and abiraterone
MDA-MB-453	Breast carcinoma	Synergy
DLD-1	colorectal adenocarcinoma	Weak Synergy
NCI-H460	lung	Additive
A549	lung	no additional effect when combined
LNCaP	prostate carcinoma	no additional effect when combined

FIG. 12



ibrutinib/abexinostat			ibrutinib/abexinostat			ibrutinib/abexinostat		
μM	μM	CI	μM	μM	CI	μM	μM	CI
0.04	0.004	0.64	0.33	0.004	0.197	1	0.004	0.311
0.04	0.008	0.84	0.33	0.008	0.435	1	0.008	0.264
0.04	0.016	0.52	0.33	0.016	0.266	1	0.016	0.292
0.04	0.031	0.64	0.33	0.031	0.222	1	0.031	0.264
0.04	0.063	0.38	0.33	0.063	0.199	1	0.063	0.206

FIG. 14



ibrutinib μ M	abexinostat μ M	CI	ibrutinib μ M	abexinostat μ M	CI	ibrutinib μ M	abexinostat μ M	CI
0.04	0.125	1.15	0.11	0.125	1.107	1	0.125	0.927
0.04	0.25	1.12	0.11	0.25	1.227	1	0.25	1.152
0.04	0.5	1.02	0.11	0.5	0.966	1	0.5	1.049
0.04	1	0.99	0.11	1	0.972	1	1	1.002
0.04	2	1.17	0.11	2	1.174	1	2	1.122

FIG. 1

