



US00RE39614E

(19) **United States**
 (12) **Reissued Patent**
Sim et al.

(10) **Patent Number:** **US RE39,614 E**
 (45) **Date of Reissued Patent:** **May 8, 2007**

- (54) **CANINE IL-4 IMMUNOREGULATORY PROTEINS AND USES THEREOF**
 (75) Inventors: **Gek-Kee Sim**, Denver, CO (US);
Matthew J. Dreitz, Firestone, CO (US)
 (73) Assignee: **Heska Corporation**, Loveland, CO (US)
 (21) Appl. No.: **10/916,286**
 (22) Filed: **Aug. 11, 2004**

Related U.S. Patent Documents

Reissue of:

- (64) Patent No.: **6,471,957**
 Issued: **Oct. 29, 2002**
 Appl. No.: **09/322,409**
 Filed: **May 28, 1999**

U.S. Applications:

- (60) Provisional application No. 60/087,306, filed on May 29, 1998.
 (51) **Int. Cl.**
A61K 45/00 (2006.01)
C07K 1/00 (2006.01)
C12P 21/06 (2006.01)
 C12N 5/00; C12N 15/00; C12N 15/63; C07H 21/04
 (52) **U.S. Cl.** **424/85.2**; 424/85.1; 435/69.1;
 435/320.1; 435/325; 435/455; 530/350;
 530/351; 536/23.1; 536/23.5
 (58) **Field of Classification Search** 536/23.1;
 530/350
 See application file for complete search history.

(56) References Cited**U.S. PATENT DOCUMENTS**

5,831,023 A 11/1998 Capon et al. 530/351

FOREIGN PATENT DOCUMENTS

EP	0 186 098 A1	7/1986
EP	0 322 870 A2	7/1989
EP	0 414 355 A1	2/1991
EP	0 759 468 A1	2/1997
EP	0 875 251 A1	11/1998
WO	WO 94/04680	3/1994
WO	WO 99/61618	12/1999

OTHER PUBLICATIONS

Armitage et al., *Seminars in Immunology*, vol. 5, 1993, pp. 401–412.
 Armitage et al., *Nature*, vol. 357, 1992, pp. 80–82.
 Azuma et al., *Nucleic Acids Research*, vol. 14, No. 22, 1986, pp. 9149–9158.
 Brown et al., *Journal of Immunology*, vol. 142, No. 2, 1989, pp. 679–687.
 Cantrell et al., *Proc. Natl. Acad. Sci. USA*, vol. 82, 1985, pp. 6250–6254.
 Daugherty et al., *Journal of Interferon Research*, vol. 4, 1984, pp. 635–643.
 Dion et al., *Biochemical and Biophysical Research Communications*, vol. 138, No. 2, 1986, pp. 826–834.
 Drexler, *Leukemia*, vol. 10, 1996, pp. 588–599.

Feng et al., *J. Mol. Evol.*, vol. 21, 1985, pp. 112–125.
 Gauchat et al., *Res. Immunol.*, vol. 145(3), Mar.–Apr. 1994, pp. 240–249.
 Gauchat et al., *FEBS 11964*, vol. 315, No. 3, 1993, pp. 259–266.
 Goeddel et al., *Nature*, vol. 290, 1981, pp. 20–26.
 van der Kaaij et al., Molecular cloning and sequencing of the cDNA for dog interleukin-4, 1999, *Immunogenetics*, vol. 49, pp. 142–143.
 McKenzie, Regulation of T helper type 2 cell immunity by interleukin-4 and interleukin-13, 2000, *Pharmacology & Therapeutics*, vol. 88, pp. 143–151.
 Gough et al., *The EMBO Journal*, vol. 4, No. 3, 1985, pp. 645–653.
 Graf et al., *Eur. J. Immunol.*, vol. 22, 1992, pp. 3191–3194.
 Grimaldi et al., *Journal of Immunology*, vol. 149, No. 12, 1992, pp. 3921–3926.
 Hannum et al., *Nature*, vol. 368, 1994, pp. 643–648.
 Heussler et al., *Gene*, vol. 114, 1992, pp. 273–278.
 Himmler et al., *Journal of Interferon Research*, vol. 7, 1987, pp. 173–183.
 Hirano et al., *Immunology*, vol. 90, 1997, pp. 294–300.
 Hollenbaugh et al., *The EMBO Journal*, vol. 11, No. 12, 1992, pp. 4313–4321.
 Imumaru et al., *Immunology and Cell Biology*, vol. 73, 1995, pp. 474–476.
 Johnson et al., *J. Mol. Biol.*, vol. 233, 1993, pp. 716–738.
 Kelley et al., *Nucleic Acids Research*, vol. 13, No. 3, 1985, pp. 805–823.
 Lakkis et al., *Biochemical and Biophysical Research Communications*, vol. 197, No. 2, 1993, pp. 612–618.
 Leong et al., *Veterinary Immunology and Immunopathology*, vol. 21, 1989, pp. 261–278.
 Lerner et al., GenBank Accession No. U39634, submitted Oct. 27, 1995.

(Continued)

Primary Examiner—Sumesh Kaushal

(57) ABSTRACT

The present invention relates to canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, and/or feline GM-CSF proteins; to canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, and/or feline GM-CSF nucleic acid molecules, including those that encode canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, and/or feline GM-CSF proteins, respectively; to antibodies raised against such proteins; and to inhibitory compounds that regulate such proteins. The present invention also includes methods to identify and obtain such proteins, nucleic acid molecules, antibodies, and inhibitory compounds. Also included in the present invention are therapeutic compositions comprising such proteins, nucleic acid molecules, antibodies and/or inhibitory compounds as well as the use of such therapeutic compositions to regulate an immune response in an animal.

10 Claims, No Drawings

OTHER PUBLICATIONS

- Lerner et al., GenBank Accession No. AAB42052 (U39634.1), submitted Oct. 27, 1995.
- Lyman et al., *Blood*, vol. 83, No. 10, 1994, pp. 2795–2801.
- Lyman et al., *Oncogene*, vol. 10, 1995, pp. 149–157.
- Lyman et al., *Cell*, vol. 75, 1993, pp. 1157–1167.
- Lyman et al., *Oncogene*, vol. 11, 1995, pp. 1165–1172.
- McClanahan et al., *Blood*, vol. 88, No. 9, 1996, pp. 3371–3382.
- McInnes et al., *Gene*, vol. 105, 1991, pp. 275–279.
- McKenzie et al., *Proc. Natl. Acad. Sci. USA*, vol. 90, 1993, pp. 3735–3739.
- Mertens et al., *Immunogenetics*, vol. 42, 1995, pp. 430–431.
- Minty et al., *Nature*, vol. 362, 1993, pp. 248–250.
- Nagata et al., *Nature*, vol. 287, 1980, pp. 401–408.
- Nakamura et al., *Biosci. Biotech. Biochem.*, vol. 56, No. 2, 1992, pp. 211–214.
- Nash et al., *Blood*, vol. 78, No. 4, 1991, pp. 930–937.
- Navarro et al., *J. Gen. Virol.* vol. 70, 1989, pp. 1381–1389.
- O'Brien et al., *Immunology and Cell Biology*, vol. 69, 1991, pp. 51–55.
- Osorio et al., *Vaccine*, vol. 17, 1999, pp. 1109–1116.
- Padrid et al., *AJVR*, vol. 59, No. 10, 1998, pp. 1263–1269.
- Patterson et al., *Journal of Clinical Investigation*, vol. 44, No. 1, 1965, pp. 140–148.
- Seow et al., *Gene*, vol. 124, 1993, pp. 291–293.
- Sideras et al., *Adv. Exp. Med. Biol.*, vol. 213, 1987, pp. 227–236.
- Stamenkovic et al., *The EMBO Journal*, vol. 8, No. 5, 1989, pp. 1403–1410.
- Torres et al., *Journal of Immunology*, vol. 148, No. 2, 1992, pp. 620–626.
- Wong et al., *Science*, vol. 228, 1985, pp. 810–815.
- Yokota et al., *Proc. Natl. Acad. Sci. USA*, vol. 83, 1986, pp. 5894–5898.
- Zhou et al., GenBank Accession No. L12991.
- Van Der Kaaij, et al., 1999, “Immunogenetics,” vol. 49, pp. 142–143.
- Ueda, et al., 1993, *Journal of Veterinary Medical Science*, vol. 55, No. 2, pp. 251–258, XP001001462.
- Venta et al., GenBank Accession No. L77382.
- Venta et al., GenBank Accession No. L77383.
- Khatlani, et al., *J. Vet. Med. Sci.*, vol. 61(8), 1999, pp. 967–969.
- McKenzie, *Pharmacology & Therapeutics*, vol. 88, 2000, pp. 143–151.
- Santra, et al., *Cancer Immunology, Immunotherapy*, vol. 44, Issue 5, 1997, pp. 291–300, abstract.
- Wondimu, et al., *Cytokine*, vol. 16, No. 3, Nov. 7, 2001, pp. 88–92.

CANINE IL-4 IMMUNOREGULATORY PROTEINS AND USES THEREOF

Matter enclosed in heavy brackets [] appears in the original patent but forms no part of this reissue specification; matter printed in *italics* indicates the additions made by reissue.

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority to prior pending U.S. Provisional Patent Application Serial No. 60/087,306, filed May 29, 1998, entitled "CANINE INTERLEUKIN-4 AND FLT-3 LIGAND PROTEINS, NUCLEIC ACID MOLECULES, AND USES THEREOF", which is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

The present invention relates to canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF nucleic acid molecules, proteins encoded by such nucleic acid molecules, antibodies raised against such proteins and/or inhibitors of such proteins or nucleic acid molecules. The present invention also includes therapeutic compositions comprising such nucleic acid molecules, proteins, antibodies and/or inhibitors, as well as their use to regulate an immune response in an animal.

BACKGROUND OF THE INVENTION

Regulating immune responses in animals is important in disease management. Immune responses can be regulated by modifying the activity of immunoregulatory molecules and immune cells.

Several immunoregulatory molecules have been found in humans and other mammal species. Interleukin-4, produced by activated type 2 helper cells (T_H2 cells), has a number of functions. These functions include promotion of naive T cells and B cells to differentiate and proliferate. IL-4 promotes T_H2 differentiation and inhibits T_H1 development. FMS-like tyrosine kinase 3, (Flt-3 ligand) stimulates the expansion and mobilization of hematopoietic precursor cell stimulating activity. CD40 is a type I transmembrane protein expressed on antigen presenting cells, such as B lymphocytes, and other types of cells such as endothelial cells, epithelial cells, and fibroblasts. CD40 ligand (also known as CD154) is a type II transmembrane protein that is preferentially expressed on activated T lymphocytes. The CD40-CD154 interaction regulates diverse pathways of the immune system, including B cell proliferation, immunoglobulin production and class switching by B cells, activation and clonal expansion of T cells, activity of antigen presenting cells, growth and differentiation of epithelial cells, and regulation of inflammatory responses at mucosal and cutaneous sites. Interleukin-5 is produced by activated type 2 helper cells (T_H2), mast cells, and eosinophils. Its main functions include promotion of growth and differentiation of eosinophils and generation of cytotoxic T cells from thymocytes. Interleukin-13 is produced by T_H1 and T_H2 cells, and promotes growth and differentiation of B cells, up-regulation of MHC class II and CD23 expression on monocytes/macrophages and B cells; and inhibition of production of inflammatory cytokines such as IL-1 α , IL-1 β , IL-6, IL-8, IL-10, IL-12, among others. Interferon alpha is an antiviral protein that has three major functions: it inhibits

viral replication by activating cellular genes that destroy mRNA and inhibit protein translation, it induces MHC class I expression in non virally-infected cells, increasing resistance to NK cells, and can activate NK cells. GM-CSF, (granulocyte-macrophage colony-stimulating factor) stimulates the production of granulocytes and macrophages.

Prior investigators have disclosed sequences encoding feline IL-4 (Lemer et al., Genbank Accession No. U39634); porcine IL-4 (Zhou et al., Genbank Accession No. L12991); bovine IL-4 (Heussler, V. T., et al., Gene, vol. 114, pp. 273-278, 1992); ovine IL-4 (Seow, H.-F., et al., Gene, vol. 124, pp. 291-293, 1993); human IL-4 (Yokota, T., et al. Proc. Natl. Acad. Sci. U.S.A., vol. 83(16), pp. 5894-5898, 1986); and murine IL-4 (Sideras, P., et al., Adv. Exp. Med. Biol., vol. 213, pp. 227-236, 1987). Prior investigators have disclosed sequences encoding murine Flt-3 ligand (McClanahan et al., Genbank Accession No. U44024); and human Flt-3 ligand (Lyman et al., Blood, vol. 83, pp. 2795-2801, 1994). Prior investigators have disclosed sequences encoding human CD40 (Stamenkovic et al., EMBO J., vol. 8:1403-1410, 1989, GenBank Accession No. X60592), bovine CD40 (Hirano et al., Immunology, vol. 90, pp. 294-300, 1997, GenBank Accession No. U57745), and murine CD40 (Grimaldi et al., J. Immunol., vol. 143, pp. 3921-3926, 1992; Torres and Clark, J. Immunol., vol. 148, pp. 620-626, 1992, GenBank Accession No. M83312). Prior investigators have disclosed sequences encoding human CD154 (Graf et al., Eur. J. Immunol., vol. 22, pp. 3191-3194, 1992; Hollenbaugh, et al., EMBO J., vol. 11:4313-4321, 1992; Gauchat et al., FEBS lett., vol., 315, pp. 259-266, 1993; GenBank Accession Nos L07414, X68550, Z15017, X67878, respectively); bovine CD154 (Mertens et al., Immunogenetics, vol. 42, pp. 430-431, GenBank Accession No. Z48468); and murine CD154 (Armitage et al., Nature, vol. 357, pp. 80-82, 1992, GenBank Accession No. X65453). Prior investigators have disclosed sequences encoding feline interleukin-5 (Padrid et al., Am. J. Vet. Res., vol. 59, pp. 1263-1269, 1988, GenBank Accession No. AF025436) and human interleukin-5 (Azuma et al., Nucleic Acids Res., vol. 14, pp. 9149-9158, 1986, GenBank Accession No. X04688). Prior investigators have disclosed sequences encoding human interleukin-13 (McKenzie et al., Proc. Natl Acad. Sci. USA, vol. 90, pp. 3735-3739, 1993; Minty et al., Nature, vol. 362, pp. 248-250, 1993, GenBank Accession Nos L06801 and X69079, respectively); murine interleukin-13 (Brown et al., J. Immunol., vol. 142, pp. 679-687; 1989, GenBank Accession No. M23504); and rat interleukin-13 (Lakkis et al., Biochem. Biophys. Res. Commun., Vol. 197, pp. 612-618, 1993, GenBank Accession No. L26913). Prior investigators have disclosed sequences encoding feline interferon (Nakamura, N., Sudo, T., Matsuda, S., Yanai, A., Biosci. Biotechnol. Biochem. (1992)Vol: 56 pp 211-214, GenBank accession #E02521). Prior investigators have also disclosed sequences encoding feline GM-CSF (direct submission to GenBank, Accession No. AF053007)

There remains a need for compounds and methods to regulate an immune response by manipulation of the function of canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF.

SUMMARY OF THE INVENTION

The present invention relates to canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13,

feline interferon alpha, or feline GM-CSF nucleic acid molecules, proteins encoded by such nucleic acid molecules, antibodies raised against such proteins and/or inhibitors of such proteins or nucleic acid molecules. Identification of the nucleic acid molecules of the present invention is unexpected because initial attempts to obtain nucleic acid molecules using PCR were unsuccessful. After numerous attempts, the inventors discovered specific primers that were useful for isolating such nucleic acid molecules.

One embodiment of the present invention is an isolated nucleic acid molecule selected from the group consisting of: (a) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:19, and/or SEQ ID NO:21; or a homolog thereof, wherein said homolog has an at least about 50 contiguous nucleotide region identical in sequence to a 50 contiguous nucleotide region of a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:19, and/or SEQ ID NO:21; (b) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:36, and/or SEQ ID NO:37 or a homolog thereof, wherein said homolog has an at least 40 contiguous nucleotide region identical in sequence to a 40 contiguous nucleotide region of a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:36, and/or SEQ ID NO:37; (c) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, and/or SEQ ID NO:50, and/or a homolog thereof, wherein said homolog has an at least 30 contiguous nucleotide region identical in sequence to a 30 contiguous nucleotide region of a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, and/or SEQ ID NO:50; (d) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, and/or SEQ ID NO:59, and/or a homolog thereof, wherein said homolog has an at least 40 contiguous nucleotide region identical in sequence to a 40 contiguous nucleotide region of a nucleic acid sequence selected from the group consisting of SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, and/or SEQ ID NO:59; (e) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:60 and/or SEQ ID NO:62 and/or a homolog thereof, wherein said homolog has an at least 30 contiguous nucleotide region identical in sequence to a 30 contiguous nucleotide region of a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:60 and/or SEQ ID NO:62; (f) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:66, SEQ ID

NO:67, SEQ ID NO:68, SEQ ID NO:69 and/or SEQ ID NO:71, and/or a homolog thereof, wherein said homolog has an at least 45 contiguous nucleotide region identical in sequence to a 45 nucleotide region of a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69 and/or SEQ ID NO:71; (g) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, and/or SEQ ID NO:79, and/or a homolog thereof, wherein said homolog has an at least 35 contiguous nucleotide region identical in sequence to a 35 contiguous nucleotide region of a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, and/or SEQ ID NO:79; (h) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, and/or SEQ ID NO:87, and/or a homolog thereof, wherein said homolog has an at least 45 contiguous nucleotide region identical in sequence to a 45 contiguous nucleotide region of a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, and/or SEQ ID NO:87; (i) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, and/or SEQ ID NO:106, and/or a homolog thereof, wherein said homolog has an at least 15 contiguous nucleotide region identical to a 15 contiguous nucleotide region of a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, and/or SEQ ID NO:106 (j) an isolated nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:118; and/or (k) an isolated nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:122, SEQ ID NO:123, SEQ ID NO:124, SEQ ID NO:126.

Another embodiment of the present invention is an isolated nucleic acid molecule selected from the group consisting of: (a) a nucleic acid molecule having a nucleic acid sequence that is at least about 92 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:19, and/or SEQ ID NO:21; (b) a nucleic acid molecule having a nucleic acid sequence that is at least about 75 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:36, and/or SEQ ID NO:37; (c) a nucleic acid molecule having a nucleic acid sequence that is at least about 75 percent identical to a nucleic acid

5

sequence selected from the group consisting of SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, and/or SEQ ID NO:50; (d) a nucleic acid molecule having a nucleic acid sequence that is at least about 70 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, and/or SEQ ID NO:59; (e) a nucleic acid molecule having a nucleic acid sequence that is at least about 70 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:60 and/or SEQ ID NO:62; (f) a nucleic acid molecule having a nucleic acid sequence that is at least about 85 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69 and/or SEQ ID NO:71; (g) a nucleic acid molecule having a nucleic acid sequence that is at least about 91 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, and/or SEQ ID NO:79; (h) a nucleic acid molecule having a nucleic acid sequence that is at least about 90 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, and/or SEQ ID NO:87; (i) a nucleic acid molecule having a nucleic acid sequence that is at least about 65 percent identical to a nucleic acid sequence selected from the group consisting of SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, and/or SEQ ID NO:106; (j) a nucleic acid molecule having a nucleic acid sequence that is selected from the group consisting of SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:122, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:116, and/or SEQ ID NO:118; and/or (k) a nucleic acid molecule having a nucleic acid sequence that is selected from the group consisting of SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:122, SEQ ID NO:123, SEQ ID NO:124, and/or SEQ ID NO:126.

Yet another embodiment of the present invention is an isolated nucleic acid molecule selected from the group consisting of: (a) a nucleic acid molecule having a nucleic acid sequence encoding an IL-4 protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 85 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:2 and/or SEQ ID NO:20 and/or (ii) a protein comprising a fragment of at least 20 amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:2 and/or SEQ ID NO:20; (b) a nucleic acid molecule having a nucleic acid sequence encoding a Flt-3 ligand protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 75 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:7, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:31, and/or SEQ ID NO:34 and/or (ii) a protein comprising a fragment of at least 25 amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:7, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:31, and/or SEQ ID NO:34; (c) a nucleic acid molecule having a nucleic acid sequence encoding a Flt-3 ligand protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 75 percent identical to an amino acid sequence selected from

6

the group consisting of SEQ ID NO:44 and/or SEQ ID NO:49 and/or (ii) a protein comprising a fragment of at least 25 amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:44 and/or SEQ ID NO:49; (d) a nucleic acid molecule having a nucleic acid sequence encoding a CD40 protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 70 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:53 and/or SEQ ID NO:58 and/or (ii) a protein comprising a fragment of at least 30 amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:53 and/or SEQ ID NO:58; (e) a nucleic acid molecule having a nucleic acid sequence encoding a CD40 protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 60 percent identical to an amino acid sequence comprising SEQ ID NO:61 and/or (ii) a protein comprising a fragment of at least 20 amino acids of an amino acid sequence comprising SEQ ID NO:61; (f) a nucleic acid molecule having a nucleic acid sequence encoding a CD154 protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 80 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:65 and/or SEQ ID NO:70, and/or (ii) a protein comprising a fragment of at least 35 amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:65 and/or SEQ ID NO:70; (g) a nucleic acid molecule having a nucleic acid sequence encoding a CD154 protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 85 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:73 and/or SEQ ID NO:78, and/or (ii) a protein comprising a fragment of at least 50 amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:73 and/or SEQ ID NO:78; (h) a nucleic acid molecule having a nucleic acid sequence encoding an IL-5 protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 85 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:81 and/or SEQ ID NO:86 and/or (ii) a protein comprising a fragment of at least 20 amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:81 and/or SEQ ID NO:86; (i) a nucleic acid molecule having a nucleic acid sequence encoding an IL-13 protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 70 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:92, SEQ ID NO:97, SEQ ID NO:100, and/or SEQ ID NO:105 and/or (ii) a protein comprising a fragment of at least 15 amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:92, SEQ ID NO:97, SEQ ID NO:100, and/or SEQ ID NO:105; (j) a nucleic acid molecule having a nucleic acid sequence encoding an interferon alpha protein having an amino acid sequence that is selected from the group consisting of amino acid sequence SEQ ID NO:108, SEQ ID NO:111, SEQ ID NO:114, and/or SEQ ID NO:117; (k) a nucleic acid molecule having a nucleic acid sequence encoding a GM-CSF protein having an amino acid sequence that is selected from the group consisting of amino acid sequence SEQ ID NO:120, SEQ ID NO:126, and/or (l) a nucleic acid molecule comprising any of said nucleic acid molecules as set forth in (a), (b), (c), (d), (e), (f), (g), (h), (i), (j), and/or (k), wherein said IL-4 protein elicits an immune response against an IL-4 protein selected from the group consisting of SEQ ID NO:2 and/or SEQ ID NO:20

and/or is a protein with interleukin-4 activity said Flt-3 ligand protein elicits an immune response against a Flt-3 ligand protein selected from the group consisting of SEQ ID NO:7, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:44, and/or SEQ ID NO:49 and/or is a protein with Flt-3 ligand activity, said CD40 protein elicits an immune response against a CD40 protein selected from the group consisting of SEQ ID NO:53, SEQ ID NO:58, and/or SEQ ID NO:61 and/or is a protein with CD40 activity, said CD154 protein elicits an immune response against a CD154 protein selected from the group consisting of SEQ ID NO:65, SEQ ID NO:70, SEQ ID NO:73, and/or SEQ ID NO:78 and/or is a protein with CD154 activity, said IL-5 protein elicits an immune response against a IL-5 protein selected from the group consisting of SEQ ID NO:81 and/or SEQ ID NO:86 and/or is a protein with IL-5 activity, said IL-13 protein elicits an immune response against an IL-13 protein selected from the group consisting of SEQ ID NO:92, SEQ ID NO:97, SEQ ID NO:100, and/or SEQ ID NO:105 and/or is a protein with IL-13 activity, said interferon alpha protein elicits an immune response against an interferon alpha protein selected from the group consisting of SEQ ID NO:108, SEQ ID NO:111, SEQ ID NO:114, and/or SEQ ID NO:117 and/or is a protein with interferon alpha activity, and/or said GM-CSF protein elicits an immune response against a GM-CSF protein selected from the group consisting of SEQ ID NO:120 and/or SEQ ID NO:125 and/or is a protein with GM-CSF activity.

The present invention also includes methods to produce any of the proteins of the present invention using nucleic acid molecules of the present invention and recombinantly using such nucleic acid molecules.

The present invention also includes an isolated protein selected from the group consisting of: (a) (i) an isolated protein of at least about 20 amino acids in length, wherein said protein is encoded by a nucleic acid molecule, wherein said nucleic acid molecule has an at least 60 contiguous nucleotide region identical in sequence to a 60 contiguous nucleotide region of a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:4, and/or SEQ ID NO:19; and/or (ii) an isolated protein of at least about 20 amino acids in length, wherein said protein has an at least 20 contiguous amino acid region identical in sequence to a 20 contiguous amino acid region selected from the group consisting of SEQ ID NO:2 and/or SEQ ID NO:20, wherein said isolated protein elicits an immune response against a canine IL-4 protein and/or has IL-4 activity; (b) (i) an isolated protein of at least about 20 amino acids in length, wherein said protein is encoded by a nucleic acid molecule, wherein said nucleic acid molecule has an at least contiguous nucleotide region identical in sequence to a 60 contiguous nucleotide region of a nucleic acid sequence selected from the group consisting of SEQ ID NO:6, SEQ ID NO:9, SEQ ID NO:22, SEQ ID NO:25, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:33, and/or SEQ ID NO:36; and/or (ii) an isolated protein of at least about 20 amino acids in length, wherein said protein has an at least 20 contiguous amino acid region identical in sequence to a 20 contiguous amino acid region selected from the group consisting of SEQ ID NO:7, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:31, and/or SEQ ID NO:34, wherein said isolated protein is capable of eliciting an immune response against a canine Flt-3 ligand protein and/or has Flt-3 activity; (c) (i) an isolated protein of at least about 20 amino acids in length, wherein said protein is encoded by a nucleic acid molecule, wherein said nucleic acid molecule has an at

least 60 contiguous nucleotide region identical in sequence to a 60 contiguous nucleotide region of a nucleic acid sequence selected from the group consisting of SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:46, and/or SEQ ID NO:48; and/or (ii) an isolated protein of at least about 20 amino acids in length, wherein said protein has an at least 20 contiguous amino acid region identical in sequence to a 20 contiguous amino acid region selected from the group consisting of SEQ ID NO:44 and/or SEQ ID NO:49, wherein said isolated protein is capable of eliciting an immune response against a feline Flt-3 ligand protein and/or has Flt-3 activity; (d) (i) an isolated protein of at least about 30 amino acids in length, wherein said protein is encoded by a nucleic acid molecule, wherein said nucleic acid molecule has an at least 90 contiguous nucleotide region identical in sequence to a 90 contiguous nucleotide region of a nucleic acid sequence selected from the group consisting of SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:55, and/or SEQ ID NO:57 and/or (ii) an isolated protein of at least about 30 amino acids in length, wherein said protein has an at least 30 contiguous amino acid region identical in sequence to a 30 contiguous amino acid region selected from the group consisting of SEQ ID NO:53, SEQ ID NO:58, wherein said isolated protein is capable of eliciting an immune response against a canine CD40 protein and/or has CD40 activity; (e) (i) an isolated protein of at least about 20 amino acids in length, wherein said protein is encoded by a nucleic acid molecule, wherein said nucleic acid molecule has an at least 60 contiguous nucleotide region identical in sequence to a 60 contiguous nucleotide region of a nucleic acid sequence comprising Seq id no:60; and/or (ii) an isolated protein of at least about 20 amino acids in length, wherein said protein has an at least 20 contiguous amino acid region identical in sequence to a 20 contiguous amino acid region comprising the amino acid sequence SEQ ID NO:61, wherein said isolated protein is capable of eliciting an immune response against a feline CD40 protein and/or has CD40 activity; (f) (i) an isolated protein of at least about 35 amino acids in length, wherein said protein is encoded by a nucleic acid molecule, wherein said nucleic acid molecule has an at least 5 contiguous nucleotide region identical in sequence to a 105 contiguous nucleotide region of a nucleic acid sequence selected from the group consisting of SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:67, and/or SEQ ID NO:69; and/or (ii) an isolated protein of at least about 35 amino acids in length, wherein said protein has an at least 35 contiguous amino acid region identical in sequence to a 35 contiguous amino acid region selected from the group consisting of SEQ ID NO:65 and/or SEQ ID NO:70, wherein said isolated protein is capable of eliciting an immune response against a canine CD154 protein and/or has CD154 activity; (g) (i) an isolated protein of at least about 50 amino acids in length, wherein said protein is encoded by a nucleic acid molecule, wherein said nucleic acid molecule has an at least 150 contiguous nucleotide region identical in sequence to a 150 contiguous nucleotide region of a nucleic acid sequence selected from the group consisting of SEQ ID NO:72, SEQ ID NO:75, and/or SEQ ID NO:77; and/or (ii) an isolated protein of at least about 50 amino acids in length, wherein said protein has an at least 50 contiguous amino acid region identical in sequence to a 50 contiguous amino acid region selected from the group consisting of SEQ ID NO:73 and/or SEQ ID NO:78, wherein said isolated protein is capable of eliciting an immune response against a feline CD154 protein and/or has CD154 activity; (h) (i) an isolated protein of at least about 20 amino acids in length, wherein said protein is encoded by a nucleic acid molecule, wherein said nucleic

acid molecule has an at least contiguous nucleotide region identical in sequence to a 60 contiguous nucleotide region of a nucleic acid sequence selected from the group consisting of SEQ ID NO:80, SEQ ID NO:83, and/or SEQ ID NO:85; and/or (ii) an isolated protein of at least about 20 amino acids in length, wherein said protein has an at least 20 contiguous amino acid region identical in sequence to a 20 contiguous amino acid region selected from the group consisting of SEQ ID NO:81 and/or SEQ ID NO:86, wherein said isolated protein is capable of eliciting an immune response against a canine IL-5 protein and/or has IL-5 activity; (i) (i) an isolated protein of at least about 15 amino acids in length, wherein said protein is encoded by a nucleic acid molecule, wherein said nucleic acid molecule has an at least 45 contiguous nucleotide region identical in sequence to a 45 contiguous nucleotide region of a nucleic acid sequence selected from the group consisting of SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:99, SEQ ID NO:102, and/or SEQ ID NO:104; and/or (ii) an isolated protein of at least about 15 amino acids in length, wherein said protein has an at least 15 contiguous amino acid region identical in sequence to a 15 contiguous amino acid region selected from the group consisting of SEQ ID NO:92, SEQ ID NO:97, SEQ ID NO:100, and/or SEQ ID NO:105, wherein said isolated protein is capable of eliciting an immune response against a canine IL-13 protein and/or has IL-13 activity; (j) (i) an isolated protein encoded by a nucleic acid molecule selected from the group consisting of SEQ ID NO:107, SEQ ID NO:110, SEQ ID NO:113, and/or SEQ ID NO:116, and/or (ii) an isolated protein selected from the group consisting of SEQ ID NO:108, SEQ ID NO:11, SEQ ID NO:114, and/or SEQ ID NO:117, wherein said isolated protein is capable of eliciting an immune response against a feline interferon alpha protein and/or has interferon alpha activity; (k) (i) an isolated protein encoded by a nucleic acid molecule selected from the group consisting of SEQ ID NO:119, SEQ ID NO:122, and/or SEQ ID NO:124, and/or (ii) an isolated protein selected from the group consisting of SEQ ID NO:120 and/or SEQ ID NO:125, wherein said isolated protein is capable of eliciting an immune response against a feline GM-CSF and/or has GM-CSF activity.

The present invention also includes an isolated protein selected from the group consisting of: (a) a protein having an amino acid sequence that is at least about 85 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:2 and/or SEQ ID NO:20; (b) a protein having an amino acid sequence that is at least about 75 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:7, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:31, and/or SEQ ID NO:34; (c) a protein having an amino acid sequence that is at least about 75 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:44 and/or SEQ ID NO:49; and/or (d) a protein having an amino acid sequence that is at least about 70 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:53 and/or SEQ ID NO:58; (e) a protein having an amino acid sequence that is at least about 60 percent identical to an amino acid sequence comprising SEQ ID NO:61; (f) a protein having an amino acid sequence that is at least about 80 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:65 and/or SEQ ID NO:70; (g) a protein having an amino acid sequence that is at least about 85 percent identical to the amino acid sequence SEQ ID NO:73 and/or SEQ ID NO:78; (h) a protein having an amino acid sequence that is at least about 85 percent

identical to an amino acid sequence selected from the group consisting of SEQ ID NO:81 and/or SEQ ID NO:86; (i) a protein having an amino acid sequence that is at least about 70 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:92, SEQ ID NO:97, SEQ ID NO:100, and/or SEQ ID NO:105; (j) a protein having an amino acid sequence selected from the group consisting of SEQ ID NO:108, SEQ ID NO:111, SEQ ID NO:114, and/or SEQ ID NO:117; and/or (k) a protein having an amino acid sequence selected from the group consisting of SEQ ID NO:120, and/or SEQ ID NO:125.

The present invention also includes isolated antibodies that selectively bind to a protein of the present invention.

One aspect of the present invention is a therapeutic composition that, when administered to an animal, regulates an immune response in said animal, said therapeutic composition comprising a therapeutic compound selected from the group consisting of: an immunoregulatory protein of the present invention; a mimotope of any of said immunoregulatory proteins; and a multimeric form of any of said immunoregulatory proteins; an isolated nucleic acid molecule of the present invention; an antibody that selectively binds to any of said immunoregulatory protein; and/or an inhibitor of a immunoregulatory protein activity identified by its ability to inhibit the activity of any of said immunoregulatory proteins. Yet another aspect of the present invention is a method to regulate an immune response in an animal comprising administering to the animal a therapeutic composition of the present invention.

The present invention also includes a method to produce an immunoregulatory protein, said method comprising culturing a cell capable of expressing said protein, said protein being encoded by a nucleic acid molecule of the present invention.

One embodiment of the present invention is a method to identify a compound capable of regulating an immune response in an animal, said method comprising: (a) contacting an isolated canine IL-4 protein of the present invention with a putative inhibitory compound under conditions in which, in the absence of said compound, said protein has T cell proliferation stimulating activity; and determining if said putative inhibitory compound inhibits said activity; (b) contacting an isolated canine Flt-3 ligand protein of the present invention with a putative inhibitory compound under conditions in which, in the absence of said compound, said protein has dendritic precursor cell proliferation stimulating activity; and determining if said putative inhibitory compound inhibits said activity; (c) contacting an isolated feline Flt-3 ligand protein of the present invention with a putative inhibitory compound under conditions in which, in the absence of said compound, said protein has dendritic precursor cell proliferation stimulating activity; and determining if said putative inhibitory compound inhibits said activity; (d) contacting an isolated canine CD40 protein of the present invention with a putative inhibitory compound under conditions in which, in the absence of said compound, said protein has CD40 ligand binding activity; and determining if said putative inhibitory compound inhibits said activity; (e) contacting an isolated feline CD40 protein of the present invention with a putative inhibitory compound under conditions in which, in the absence of said compound, said protein has CD40 ligand binding activity; and determining if said putative inhibitory compound inhibits said activity; (f) contacting an isolated canine CD154 protein of the present invention with a putative inhibitory compound under conditions in which, in the absence of said compound, said protein has B cell proliferation activity; and determining if

said putative inhibitory compound inhibits said activity; (g) contacting an isolated feline CD154 protein of the present invention with a putative inhibitory compound under conditions in which, in the absence of said compound, said protein has B cell proliferation activity; and determining if said putative inhibitory compound inhibits said activity; (h) contacting an isolated canine IL-5 protein of the present invention with a putative inhibitory compound under conditions in which, in the absence of said compound, said protein has TF-1 cell proliferation activity; and determining if said putative inhibitory compound inhibits said activity; (i) contacting an isolated canine IL-13 protein of the present invention with a putative inhibitory compound under conditions in which, in the absence of said compound, said protein has TF-1 cell proliferation activity; and determining if said putative inhibitory compound inhibits said activity; (j) contacting an isolated feline IFN α protein of the present invention with a putative inhibitory compound under conditions in which, in the absence of said compound, said protein has inhibition of proliferation of GM-CSF stimulated TF-1 cell activity; and determining if said putative inhibitory compound inhibits said activity; or (k) contacting an isolated feline GM-CSF protein of the present invention with a putative inhibitory compound under conditions in which, in the absence of said compound, said protein has TF-1 cell proliferation activity; and determining if said putative inhibitory compound inhibits said activity.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides for isolated canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF proteins, isolated canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF nucleic acid molecules, antibodies directed against canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF proteins, and compounds derived therefrom that regulate the immune response of an animal (e.g. inhibitors, antibodies and peptides).

Canine IL-4 protein can refer to a canine IL-4 protein, including homologs thereof. Canine Flt-3 ligand protein can refer to a canine Flt-3 ligand, including homologs thereof, and feline Flt-3 ligand can refer to feline Flt-3 ligand, including homologs thereof. Canine CD40 can refer to a canine CD40, including homologs thereof; feline CD40 can refer to a feline CD40, including homologs thereof. Canine CD154 can refer to a canine CD154, including homologs thereof; feline CD154 can refer to a feline CD154, including homologs thereof. Canine IL-5 can refer to canine IL-5, including homologs thereof; canine IL-13 can refer to canine IL-13, including homologs thereof. Feline IFN α can refer to a feline IFN α , including homologs thereof, and feline GM-CSF can refer to a feline GM-CSF, including homologs thereof. As used herein, the phrase "regulate an immune response" refers to modulating the activity of cells or molecules involved in an immune response. The term "regulate" can refer to increasing or decreasing an immune response. Regulation of an immune response can be determined using methods known in the art as well as methods disclosed herein. The term, "immunoregulatory protein" refers to a protein that can modulate the activity of cells or of molecules involved in an immune response. An immu-

noregulatory protein of the present invention refers to a canine IL-4, a canine and/or feline CD40, a canine and/or feline Flt3 ligand, a canine and/or feline CD154, a canine IL-5, a canine IL-13, a feline IFN α and/or a feline GM-CSF protein as described herein. As used herein, the terms isolated canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF proteins and/or isolated canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF nucleic acid molecules refer to canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF proteins and/or canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF nucleic acid molecules derived from mammals and, as such, can be obtained from their natural source, or can be produced using, for example, recombinant nucleic acid technology or chemical synthesis. Also included in the present invention is the use of these proteins, nucleic acid molecules, antibodies, and/or compounds derived therefrom as therapeutic compositions to regulate the immune response of an animal as well as in other applications, such as those disclosed below.

One embodiment of the present invention is an isolated protein that includes a canine IL-4 protein, a canine and/or feline Flt-3 ligand protein, a canine and/or feline CD40 protein, a canine and/or feline CD154 protein, a canine interleukin-5 protein, a canine interleukin-13 protein, a feline interferon alpha protein, and/or a feline GM-CSF protein. It is to be noted that the term "a" or "an" entity refers to one or more of that entity: for example, a protein refers to one or more proteins or at least one protein. As such, the terms "a" (or "an"), "one or more" and "at least one" can be used interchangeably herein. It is also to be noted that the terms "comprising", "including", and "having" can be used interchangeably. According to the present invention, an isolated, or biologically pure, protein, is a protein that has been removed from its natural milieu. As such, "isolated" and/or "biologically pure" do not necessarily reflect the extent to which the protein has been purified. An isolated protein of the present invention can be obtained from its natural source, can be produced using recombinant DNA technology, or can be produced by chemical synthesis. Nucleic acid molecules of the present invention of known length isolated from *Canis familiaris* are denoted as follows: IL-4 is denoted as nCaIL-4_x, for example, nCaIL-4₅₄₉, wherein "#" refers to the number of nucleotides in that molecule; and in a similar fashion, Flt-3 ligand nucleic acid molecules are referred to as nCaFlt3L_x; CD40, nCaCD40_x; CD154, nCaCD154_x; IL-5, nCaIL-5_x; and IL-13, nCaIL-13_x. In a similar fashion, Flt-3 ligand nucleic acid molecules of the present invention of known length isolated from *Felis catus* are denoted as nFeFlt3L_x, CD40, nFeCD40_x; CD154, nFeCD154_x; IFN α , nFeIFN α _x; and GM-CSF (also denoted GMCSF), nFeGM-CSF_x. Similarly, proteins of the present invention of known length isolated from *Felis catus* are denoted as PFeFlt3I_x, PFeCD40_x, PFeCD154_x, PFeIFN α _x, and/or PFeGM-CSF_x; and proteins of the present invention of known length isolated from *Canis familiaris* are denoted PCaIL-4_x, PCaFlt3L_x, PCaCD40_x, PCaCD154_x, PCaIL-5_x, and/or PCaIL-13_x.

As used herein, an isolated canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline

CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, and/or feline GM-CSF ligand protein of the present invention (i.e., an canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF protein, respectively) can be a full-length protein or any homolog of such a protein. An isolated IL-4 protein of the present invention, including a homolog, can be identified in a straight-forward manner by the protein's ability to elicit an immune response against, (or to) an IL4 protein, bind to an IL-4 receptor, stimulate B cell differentiation or activation or stimulate production of immunoglobulin by a B cell. An isolated Flt-3 ligand protein of the present invention, including a homolog, can be identified in a straight-forward manner by the protein's ability to elicit an immune response against a Flt-3 ligand protein, bind to Flt-3 receptor or stimulate Flt-3 receptor-bearing hematopoietic stem cells, early hematopoietic progenitor cells or immature lymphocytes. An isolated CD40 protein of the present invention, including a homolog, can be identified in a straight-forward manner by the protein's ability to elicit an immune response against a CD40 protein, bind to CD154 or stimulate CD154-bearing B cells, T cells, and/or epithelial cells. An isolated CD154 protein of the present invention, including a homolog, can be identified in a straight-forward manner by the protein's ability to elicit an immune response to a CD154 protein, bind to CD40 or stimulate CD40-bearing B cells, T cells, and/or epithelial cells. An isolated IL-5 protein of the present invention, including a homolog, can be identified in a straight-forward manner by the protein's ability to elicit an immune response to an IL-5 protein, bind to an IL-5 receptor, and/or stimulate eosinophils and/or cause thymocytes to product cytotoxic T cells. An isolated IL-13 protein of the present invention, including a homolog, can be identified in a straight-forward manner by the protein's ability to elicit an immune response to an IL-13 protein, bind to an IL-13 receptor, and/or stimulate B cells, up-regulate expression of MHC class II and/or CD23 on monocytes, macrophages and/or B cells; and/or inhibition of proinflammatory cytokines. An isolated interferon alpha protein of the present invention, including a homolog, can be identified in a straight-forward manner by the protein's ability to elicit an immune response to an interferon alpha protein, bind to an interferon-alpha receptor, and/or activate NK cells and/or inhibit viral replication. An isolated GM-CSF protein of the present invention, including a homolog, can be identified in a straight-forward manner by the protein's ability to elicit an immune response to a GM-CSF protein, bind to a GM-CSF receptor, and/or active granulocytes and/or macrophages. Examples of protein homologs of the present invention include immunoregulatory proteins of the present invention in which amino acids have been deleted (e.g., a truncated version of the protein, such as a peptide), inserted, inverted, substituted and/or derivatized (e.g., by glycosylation, phosphorylation, acetylation, myristoylation, prenylation, palmitoylation, amidation and/or addition of glycerophosphatidyl inositol) such that the protein homolog includes at least one epitope capable of eliciting an immune response against the parent protein, of binding to an antibody directed against the parent protein and/or of binding to the parent's receptor, where the term parent refers to the longer and/or full-length protein that the homolog is derived from. That is, when the homolog is administered to an animal as an immunogen, using techniques known to those skilled in the art, the animal will produce an immune response against at least one epitope of an immunoregulatory protein of the

present invention, depending upon which protein is administered to an animal. The ability of a protein to effect an immune response can be measured using techniques known to those skilled in the art. As used herein, the term "epitope" refers to the smallest portion of a protein capable of selectively binding to the antigen binding site of an antibody. It is well accepted by those skilled in the art that the minimal size of a protein epitope capable of selectively binding to the antigen binding site of an antibody is about five or six to seven amino acids.

Homologs of immunoregulatory proteins of the present invention can be the result of natural allelic variation, including natural mutation. Protein homologs of the present invention can also be produced using techniques known in the art including, but not limited to, direct modifications to the protein and/or modifications to the gene encoding the protein using, for example, classic or recombinant DNA techniques to effect random or targeted mutagenesis.

Immunoregulatory proteins of the present invention include variants of a full-length protein of a protein of the present invention. Such variants include proteins that are less than full-length. As used herein, variants of the present invention refer to nucleic acid molecules that are naturally-occurring as defined below, and may result from alternative RNA splicing, alternative termination of an amino acid sequence or DNA recombination. Examples of variants include allelic variants as defined below. It is to be noted that a variant is an example of a homolog of the present invention.

Immunoregulatory proteins of the present invention are encoded by nucleic acid molecules of the present invention. As used herein, an IL-4 nucleic acid molecule includes nucleic acid sequences related to a natural canine IL-4 gene. As used herein, a Flt-3 ligand nucleic acid molecule includes nucleic acid sequences related to a natural canine Flt-3 ligand gene. As used herein, a CD40 nucleic acid molecule includes nucleic acid sequences related to a natural CD40 gene. As used herein, a CD154 nucleic acid molecule includes nucleic acid sequences related to a natural CD154 gene. As used herein, an IL-5 nucleic acid molecule includes nucleic acid sequences related to a natural IL-5 gene. As used herein, an IL-13 nucleic acid molecule includes nucleic acid sequences related to a natural IL-13 gene. As used herein, an IFN α nucleic acid molecule includes nucleic acid sequences related to a natural IFN α gene. As used herein, a GM-CSF nucleic acid molecule includes nucleic acid sequences related to a natural GM-CSF gene. As used herein, a canine IL-4, a canine and/or feline CD40, a canine and/or feline Flt3 ligand, a canine and/or feline CD154, a canine IL-5, a canine IL-13, a feline IFN α , and/or a feline GM-CSF gene refers to the natural genomic elements that encode an canine IL-4, a canine and/or feline CD40, a canine and/or feline Flt3 ligand, a canine and/or feline CD154, a canine IL-5, a canine IL-13, a feline IFN α , and/or a feline GM-CSF protein, respectively, and includes all regions such as regulatory regions that control production of the protein encoded by the gene (such as, but not limited to, transcription, translation or post-translation control regions) as well as the coding region itself, and any introns or non-translated coding regions. As used herein, a gene that "includes" or "comprises" a sequence may include that sequence in one contiguous array, or may include the sequence as fragmented exons. As used herein, the term "coding region" refers to a continuous linear array of nucleotides that translates into a protein. A full-length coding region is that region that is translated into a full-length, i.e., a complete, protein as would be initially translated in its natural milieu, prior to any post-translational modifications.

In one embodiment, an IL-4 gene of the present invention includes the nucleic acid sequence SEQ ID NO:1, as well as the complement of SEQ ID NO:1. Nucleic acid sequence SEQ ID NO:1 represents the deduced sequence of the coding strand of a cDNA (complementary DNA) denoted herein as nucleic acid molecule nCaIL-4₅₄₉, the production of which is disclosed in the Examples. Nucleic acid molecule nCaIL-4₅₄₉ comprises an apparently full-length coding region of canine IL-4. The complement of SEQ ID NO:1 (represented herein by SEQ ID NO:3) refers to the nucleic acid sequence of the strand fully complementary to the strand having SEQ ID NO:1, which can easily be determined by those skilled in the art. Likewise, a nucleic acid sequence complement of any nucleic acid sequence of the present invention refers to the nucleic acid sequence of the nucleic acid strand that is fully complementary to (i.e., can form a double helix with) the strand for which the sequence is cited. It should be noted that since nucleic acid sequencing technology is not entirely error-free, SEQ ID NO:1 (as well as other nucleic acid and protein sequences presented herein) represents an apparatus nucleic acid sequence of the nucleic acid molecule encoding an immunoregulatory protein of the present invention.

In another embodiment, a Flt-3 ligand gene of the present invention includes the nucleic acid sequence SEQ ID NO:6, as well as the complement represented by SEQ ID NO:8. Nucleic acid sequence SEQ ID NO:6 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nCaFlt3L₁₀₁₃, the production of which is disclosed in the Examples. Nucleic acid molecule nCaFlt3L₁₀₁₃ comprises an apparently full-length coding region of canine Flt-3 ligand.

In another embodiment, a Flt-3 ligand gene of the present invention includes the nucleic acid sequence SEQ ID NO:43, as well as the complement represented by SEQ ID NO:45. Nucleic acid sequence SEQ ID NO:43 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFeFlt3L₉₄₂, the production of which is disclosed in the Examples. Nucleic acid molecule nFeFlt3L₉₄₂ comprises an apparently full-length coding region of feline Flt-3 ligand.

In another embodiment, a CD40 gene of the present invention includes the nucleic acid sequence SEQ ID NO:52, as well as the complement represented by SEQ ID NO:54. Nucleic acid sequence SEQ ID NO:52 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nCaCD40₁₄₂₅, the production of which is disclosed in the Examples. Nucleic acid molecule nCaCD40₁₄₂₅ comprises an apparently full-length coding region of canine CD40.

In another embodiment, a CD40 gene of the present invention includes the nucleic acid sequence SEQ ID NO:60, as well as the complement represented by SEQ ID NO:62. Nucleic acid sequence SEQ ID NO:60 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFeCD40₃₃₆, the production of which is disclosed in the Examples. Nucleic acid molecule nFeCD40₃₃₆ comprises an apparent portion of the coding region of feline CD40.

In another embodiment, a CD154 gene of the present invention includes the nucleic acid sequence SEQ ID NO:64, as well as the complement represented by SEQ ID NO:66. Nucleic acid sequence SEQ ID NO:64 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nCaCD154₁₈₇₈, the production of which is disclosed in the Examples. Nucleic acid

molecule nCaCD154₁₈₇₈ comprises an apparently full-length coding region of canine CD154.

In another embodiment, a CD154 gene of the present invention includes the nucleic acid sequence SEQ ID NO:72, as well as the complement represented by SEQ ID NO:74. Nucleic acid sequence SEQ ID NO:72 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFeCD154₈₈₅, the production of which is disclosed in the Examples. Nucleic acid molecule nFeCD154₈₈₅ comprises an apparently full-length coding region of feline CD154.

In another embodiment, an IL-5 gene of the present invention includes the nucleic acid sequence SEQ ID NO:80, as well as the complement represented by SEQ ID NO:82. Nucleic acid sequence SEQ ID NO:80 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nCaIL-5₆₁₀, the production of which is disclosed in the Examples. Nucleic acid molecule nCaIL-5₆₁₀ comprises an apparently full-length coding region of canine IL-5.

In another embodiment, an IL-13 gene of the present invention includes the nucleic acid sequence SEQ ID NO:91, as well as the complement represented by SEQ ID NO:93. Nucleic acid sequence SEQ ID NO:91 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nCaIL-13₁₃₀₂, the production of which is disclosed in the Examples. Nucleic acid molecule nCaIL-13₁₃₀₂ comprises an apparently full-length coding region of canine IL-13.

In another embodiment, an IFN α gene of the present invention includes the nucleic acid sequence SEQ ID NO:107, as well as the complement represented by SEQ ID NO:109. Nucleic acid sequence SEQ ID NO:107 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFeIFN α _{567a}, the production of which is disclosed in the Examples. Nucleic acid molecule nFeIFN α _{567a} comprises an apparently full-length coding region of feline IFN α .

In another embodiment, a GM-CSF gene of the present invention includes the nucleic acid sequence SEQ ID NO:119, as well as the complement represented by SEQ ID NO:121. Nucleic acid sequence SEQ ID NO:119 represents the deduced sequence of the coding strand of a cDNA denoted herein as nucleic acid molecule nFeGM-CSF₄₄₄, the production of which is disclosed in the Examples. Nucleic acid molecule nFeGM-CSF₄₄₄ comprises an apparently full-length coding region of feline GM-CSF.

Additional immunoregulatory nucleic acid molecules and proteins of the present invention having specific sequence identifiers are described in Table 1.

TABLE 1

Sequence identification numbers (SEQ ID NOs) and their corresponding nucleic acid molecules or proteins.

SEQ ID NO	DESCRIPTION
1	nCaIL-4 ₅₄₀ coding strand
2	PCaIL-4 ₁₃₂
3	nCaIL-4 ₅₄₉ complementary strand
4	nCaIL-4 ₃₉₆ coding strand
5	nCaIL-4 ₃₉₆ complementary strand
6	nCaFlt3L ₁₀₁₃ coding strand
7	PCaFlt3L ₂₉₄
8	nCaFlt3L ₁₀₁₃ complementary strand
9	nCaFlt3L ₈₈₂ coding strand

TABLE 1-continued

Sequence identification numbers (SEQ ID NOs) and their corresponding nucleic acid molecules or proteins.

SEQ ID NO	DESCRIPTION
10	nCaFlt3L ₈₈₂ complementary strand
19	nCaIL-4 ₃₂₄ coding strand
20	PCaIL-4 ₁₀₈
21	nCaIL-4 ₃₂₄ complementary strand
22	nCaFlt3L ₈₀₄ coding strand
23	PCaFlt3L ₂₆₈
24	nCaFlt3L ₈₀₄ complementary strand
25	nCaFlt3L ₉₈₅ coding strand
26	PCaFlt3L ₂₇₆
27	nCaFlt3L ₉₈₅ complementary strand
28	nCaFlt3L ₈₂₈ coding strand
29	nCaFlt3L ₈₂₈ complementary strand
30	nCaFlt3L ₇₅₀ coding strand
31	PCaFlt3L ₂₅₀
32	nCaFlt3L ₇₅₀ complementary strand
33	nCaFlt3L ₁₀₁₉ coding strand
34	PCaFlt3L ₃₁
35	nCaFlt3L ₁₀₁₉ complementary strand
36	nCaFlt3L ₉₃ coding strand
37	nCaFlt3L ₉₃ complementary strand
41	nFeFlt3L ₃₉₅ coding strand
42	nFeFlt3L ₇₉₃ coding strand
43	nFeFlt3L ₉₄₂ coding strand
44	PFeFlt3L ₂₉₁
45	nFeFlt3L ₉₄₂ complementary strand
46	nFeFlt3L ₈₇₃ coding strand
47	nFeFlt3L ₈₇₃ complementary strand
48	nFeFlt3L ₇₉₅ coding strand
49	PFeFlt3L ₂₆₅
50	nFeFlt3L ₇₉₅ complementary strand
51	nCaCD40 ₃₂₄ coding strand
52	nCaCD40 ₁₄₂₅ coding strand
53	PCaCD40 ₂₇₄
54	nCaCD40 ₁₄₂₅ complementary strand
55	nCaCD40 ₈₂₂ coding strand
56	nCaCD40 ₈₂₂ complementary strand
57	nCaCD40 ₇₆₅ coding strand
58	PCaCD40 ₂₅₅
59	nCaCD40 ₇₆₅ complementary strand
60	nFeCD40 ₃₃₆ coding strand
61	PFeCD40 ₁₁₂
62	nFeCD40 ₃₃₆ complementary strand
63	nCaCD154 ₃₉₀ coding strand
64	nCaCD154 ₁₈₇₈ coding strand
65	PCaCD154 ₂₆₀
66	nCaCD154 ₁₈₇₈ complementary strand
67	nCaCD154 ₇₈₀ coding strand
68	nCaCD154 ₇₈₀ complementary strand
69	nCaCD154 ₆₃₃ coding strand
70	PCaCD154 ₂₁₁
71	nCaCD154 ₆₅₃ complementary strand
72	nFeCD154 ₈₈₅ coding strand
73	PFeCD154 ₂₆₀
74	nFeCD154 ₈₈₅ complementary strand
75	nFeCD154 ₇₅₀ coding strand
76	nFeCD154 ₇₅₀ complementary strand
77	nFeCD154 ₈₃₃ coding strand
78	PFeCD154 ₄
79	nFeCD154 ₈₃₃ complementary strand
80	nCaIL-5 ₆₁₀ coding strand
81	PCaIL-5 ₁₃₄
82	nCaIL-5 ₆₁₀ complementary strand
83	nCaIL-5 ₄₀₂ coding strand
84	nIL-5 ₄₀₂ complementary strand
85	nCaIL-5 ₃₄₉ coding strand
86	PCaIL-5 ₁₁₅
87	nCaIL-5 ₃₄₅ complementary strand
88	nCaIL-13 ₁₆₆ coding strand
89	nCaIL-13 ₂₇₂ coding strand
90	nCaIL-13 ₂₇₈ coding strand
91	nCaIL-13 ₁₃₀₂ coding strand
92	PCaIL-13 ₁₃₁
93	nCaIL-13 ₁₃₀₂ complementary strand

TABLE 1-continued

Sequence identification numbers (SEQ ID NOs) and their corresponding nucleic acid molecules or proteins.

SEQ ID NO	DESCRIPTION
94	nCaIL-13 ₃₉₃ coding strand
95	nCaIL-13 ₃₉₃ complementary strand
96	nCaIL-13 ₃₃₃ coding strand
97	PaIL-13 ₁₁₁
98	nCaIL-13 ₃₃₃ complementary strand
99	nCaIL-13 ₁₂₆₉ coding strand
100	PCaIL-13 ₁₁₀
101	nCaIL-13 ₁₂₆₉ complementary strand
102	nCaIL-13 ₃₉₀ coding strand
103	nCaIL-13 ₃₉₀ complementary strand
104	nCaIL-13 ₃₃₀ coding strand
105	PCaIL-13 ₁₁₀
106	nCaIL-13 ₃₃₀ complementary strand
107	nFeIFN _{567a} coding strand
108	PFeIFN _{189a}
109	nFeIFN _{567a} complementary strand
110	nFeIFN _{567b} coding strand
111	PFeIFN _{180b}
112	nFeIFN _{567b} complementary strand
113	nFeIFN _{498a} coding strand
114	PFeIFN _{166a}
115	nFeIFN _{498a} complementary strand
116	nFeFeIFN _{498b} coding strand
117	PFeIFN _{166b}
118	nFeIFN _{498b} complementary strand
119	nFeGMCSF ₄₄₄ coding strand
120	PFeGMCSF ₁₆₄
121	nFeGMCSF ₄₄₄ complementary strand
122	nFeGMCSF ₄₃₂ coding strand
123	nFeGMCSF ₄₃₂ complementary strand
124	nFeGMCSF ₃₈₁ coding strand
125	PFeGMCSF ₁₂₇
126	nFeGMCSF ₃₈₁ complementary strand

In another embodiment, an IL-4 gene or nucleic acid molecule can be an allelic variant that includes a similar but not identical sequence to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:19, SEQ ID NO:21, and/or any other IL-4 nucleic acid sequence cited herein. In another embodiment, a Flt-3 ligand gene or nucleic acid molecule can be an allelic variant that includes a similar but not identical sequence to SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:50 and/or any other Flt-3 ligand nucleic acid sequence cited herein. In another embodiment, a CD40 gene or nucleic acid molecule can be an allelic variant that includes a similar but not identical sequence to SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:62 and/or any other CD40 nucleic acid sequence cited herein. In another embodiment, a CD154 gene or nucleic acid molecule can be an allelic variant that includes a similar but not identical sequence to SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:79 and/or any other CD154 nucleic acid sequence cited herein. In another embodiment, an IL-5 gene or nucleic acid molecule can be an allelic variant that includes a similar but not identical sequence to SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84,

SEQ ID NO:85, SEQ ID NO:87 and/or any other IL-5 nucleic acid sequence cited herein. In another embodiment, an IL-13 gene or nucleic acid molecule can be an allelic variant that includes a similar but not identical sequence to SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:106 and/or any other IL-13 nucleic acid sequence cited herein. In another embodiment, an IFN α gene or nucleic acid molecule can be an allelic variant that includes a similar but not identical sequence to SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:116, and/or SEQ ID NO:118 and/or any other IFN α nucleic acid sequence cited herein. In another embodiment, a GM-CSF gene or nucleic acid molecule can be an allelic variant that includes a similar but not identical sequence to SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:122, SEQ ID NO:123, SEQ ID NO:124, and/or SEQ ID NO:126 and/or any other GM-CSF nucleic acid sequence cited herein. An allelic variant of a canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF gene, including the particular SEQ ID NO's cited herein, is a gene that occurs at essentially the same locus (or loci) in the genome as the gene including the particular SEQ ID NO's cited herein, but which, due to natural variations caused by, for example, mutation or recombination, has a similar but not identical sequence. Also included in the term allelic variant are allelic variants of cDNAs derived from such genes. Because natural selection typically selects against alterations that affect function, allelic variants usually encode proteins having similar activity to that of the protein encoded by the gene to which they are being compared. Allelic variants of genes or nucleic acid molecules can also comprise alterations in the 5' or 3' untranslated regions of the gene (e.g., in regulatory control regions), or can involve alternative splicing of a nascent transcript, thereby brining alternative exons into juxtaposition. Allelic variants are well known to those skilled in the art and would be expected to be found within a given animal, since the respective genomes are diploid, and sexual reproduction will result in the reassortment of alleles.

The minimal size of an canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF protein homolog of the present invention is a size sufficient to be encoded by a nucleic acid molecule capable of forming a stable hybrid (i.e., hybridize under stringent hybridization conditions) with the complementary sequence of a nucleic acid molecule encoding the corresponding natural protein. Stringent hybridization conditions are determined based on defined physical properties of the gene to which the nucleic acid molecule is being hybridized, and can be defined mathematically. Stringent hybridization conditions are those experimental parameters that allow an individual skilled in the art to identify significant similarities between heterologous nucleic acid molecules. These conditions are well known to those skilled in the art. See, for example, Sambrook, et al., 1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Labs Press, and Meinkoth, et al., 1984, Anal. Biochem. 138, 267-284, each of which is incorporated herein by this reference. As explained in detail in the cited references, the determination of hybridization conditions

involves the manipulation of a set of variables including the ionic strength (M. in moles/liter), the hybridization temperature ($^{\circ}$ C.), the concentration of nucleic acid helix destabilizing agents, such as formamide, the average length of the shortest hybrid duplex (n), and the percent G+C composition of the fragment to which an unknown nucleic acid molecule is being hybridized. For nucleic acid molecules of at least about 150 nucleotides, these variables are inserted into a standard mathematical formula to calculate the melting temperature, or T_m , of a given nucleic acid molecule. As defined in the formula below, T_m is the temperature at which two complementary nucleic acid molecule strands will disassociate, assuming 100% complementarity between the two strands:

$$T_m = 81.5^{\circ} \text{ C.} + 16.6 \log M + 0.41(\%G+C) - 500/n - 0.61(\% \text{ formamide}).$$

For nucleic acid molecules smaller than about 50 nucleotides, hybrid stability is defined by the dissociation temperature (T_d), which is defined as the temperature at which 50% of the duplexes dissociate. For these smaller molecules, the stability at a standard ionic strength is defined by the following equation:

$$T_d = 4(G+C) + 2(A+T).$$

A temperature of 5° C. below T_d is used to detect hybridization between perfectly matched molecules.

Also well known to those skilled in the art is how base pair mismatch, i.e. differences between two nucleic acid molecules being compared, including non-complementarity of bases at a given location, and gaps due to insertion or deletion of one or more bases at a given location on either of the nucleic acid molecules being compared, will affect T_m or T_d for nucleic acid molecules of different sizes. For example, T_m decreases about 1° C. for each 1% of mismatched base pairs for hybrids greater than about 150 bp, and T_d decreases about 5° C. for each mismatched base pair for hybrids below about 50 bp. Conditions for hybrids between about 50 and about 150 base pairs can be determined empirically and without undue experimentation using standard laboratory procedures well known to those skilled in the art. These simple procedures allow one skilled in the art to set the hybridization conditions, by altering, for example, the salt concentration, the formamide concentration, or the temperature, so that only nucleic acid hybrids with greater than a specified % base pair mismatch will hybridize. Stringent hybridization conditions are commonly understood by those skilled in the art to be those experimental conditions that will allow about 30% base pair mismatch, i.e., about 70% identity. Because one skilled in the art can easily determine whether a given nucleic acid molecule to be tested is less than or greater than about 50 nucleotides, and can therefore choose the appropriate formula for determining hybridization conditions, he or she can determine whether the nucleic acid molecule will hybridize with a given gene or specified nucleic acid molecule under stringent hybridization conditions and similarly whether the nucleic acid molecule will hybridize under conditions designed to allow a desired amount of base pair mismatch.

Hybridization reactions are often carried out by attaching the nucleic acid molecule to be hybridized to a solid support such as a membrane, and then hybridizing with a labeled nucleic acid molecule, typically referred to as a probe, suspended in a hybridization solution. Examples of common hybridization reaction techniques include, but are not limited to, the well-known Southern and Northern blotting procedures. Typically, the actual hybridization reaction is done

under non-stringent conditions, i.e., at a lower temperature and/or a higher salt concentration, and then high stringency is achieved by washing the membrane in a solution with a higher temperature and/or lower salt concentration in order to achieve the desired stringency.

Preferred portions, or fragments, of a canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF, protein of the present invention include at least 15 amino acids, at least 20 amino acids, at least 25 amino acids, at least 30 amino acids, at least 35 amino acids, at least 40 amino acids, at least 45 amino acids, at least 50 amino acids, at least 60 amino acids, at least 75 amino acids or at least 100 amino acids. An IL-4, IL-5, and/or IL-13 protein of the present invention can include at least a portion of an IL-4, IL-5, and/or IL-13 protein that is capable of binding to an IL-4, IL-5, and/or IL-13 receptor, respectively. IL-4, IL-5, and IL-13 receptors are known to those of skill in the art, and are described in Janeway et al., in *Immunobiology, the Immune System in Health and Disease*, Garland Publishing, Inc., NY, 1996 (which is incorporated herein by this reference in its entirety). The IL-4, IL-5, and/or IL-13 receptor-binding portion of an IL-4, IL-5, and/or IL-13 protein, respectively, can be determined by incubating the protein with an isolated IL-4, IL-5, and/or IL-13 receptor, as appropriate, or a cell having an IL-4, IL-5, and/or IL-13 receptor on its surface, as appropriate, IL-4, IL-5, and/or IL-13 protein binding to purified IL-4, IL-5, and/or IL-13 receptor, respectively, can be determined using methods known in the art including Biacore® screening, confocal immunofluorescent microscopy, immunoprecipitation, gel chromatography, determination of inhibition of binding of antibodies that bind specifically to the IL-4, IL-5, and/or IL-13 binding domain of an IL-4, IL-5, and/or IL-13 receptor. ELISA using an IL-4, IL-5, and/or IL-13 receptor, respectively, labeled with a detectable tag such as an enzyme or chemiluminescent tag or yeast-2 hybrid technology. A Flt-3 ligand protein of the present invention can include at least a portion of a Flt-3 ligand protein that is capable of binding to Flt-3 receptor or stimulating Flt-3 receptor-bearing hematopoietic stem cells, early hematopoietic progenitor cells or immature lymphocytes. Flt-3 receptors are known to those of skill in the art, and are described in Drexler, *Leukemia*, vol. 10, pp. 588-599, 1996 (which is incorporated herein in its entirety by this reference). The Flt-3 receptor-binding portion of a Flt-3 ligand protein can be determined by incubating the protein with isolated Flt-3 receptor or a cell having a Flt-3 receptor on its surface. Flt-3 ligand protein binding to purified Flt-3 receptor can be determined using methods known in the art including Biacore® screening, confocal immunofluorescent microscopy, immunoprecipitation, gel chromatography, determination of inhibition of binding of antibodies that bind specifically to the Flt-3 ligand binding domain of a Flt-3 receptor. ELISA using a Flt-3 receptor labeled with a detectable tag such as an enzyme or chemiluminescent tag or yeast-2 hybrid technology. A CD40 and/or CD154 protein of the present invention can include at least a portion of a CD40 and/or CD154 protein that is capable of binding to a CD40 and/or CD154 receptor, respectively, or stimulating CD40 and/or CD154 receptor-bearing hematopoietic stem cells, early hematopoietic progenitor cells or immature lymphocytes. The CD40 and/or CD154 receptor-binding portion of a CD40 and/or CD154 protein can be determined by incubating the protein with isolated CD40 and/or CD154 receptor, as appropriate, or a cell having a CD40 and/or CD154 receptor on its surface, as

appropriate. CD40 and/or CD154 protein binding to CD154 and/or CD40, respectively, can be determined using methods known in the art including Biacore® screening, confocal immunofluorescent microscopy, immunoprecipitation, gel chromatography, determination of inhibition of binding of antibodies that bind specifically to the CD40 and/or CD154 binding domain of CD40 and/or CD154, as appropriate, ELISA using a CD40 and/or CD154 labeled with a detectable tag such as an enzyme or chemiluminescent tag or yeast-2 hybrid technology.

The present invention also includes mimetopes of canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF proteins of the present invention. As used herein, a mimetope of an immunoregulatory protein of the present invention refers to any compound that is able to mimic the activity of such a canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF protein, respectively, often because the mimetope has a structure that mimics the particular protein. Mimetopes can be, but are not limited to: peptides that have been modified to decrease their susceptibility to degradation such as all-D retro peptides; anti-idiotypic and/or catalytic antibodies, or fragments thereof; non-proteinaceous immunogenic portions of an isolated protein (e.g., carbohydrate structures); and/or synthetic or natural organic molecules, including nucleic acids. Such mimetopes can be designed using computer-generated structures of proteins of the present invention. Mimetopes can also be obtained by generating random samples of molecules, such as oligonucleotides, peptides or other organic molecules, and screening such samples by affinity chromatography techniques using the corresponding binding partner.

One embodiment of an immunoregulatory protein of the present invention is a fusion protein that includes either a canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF protein-containing domain, each attached to one or more fusion segments. Suitable fusion segments for use with the present invention include, but are not limited to, segments that can: link two or more immunoregulatory proteins of the present invention, to form multimeric forms of an immunoregulatory protein of the present invention; enhance a protein's stability; act as an immunopotentiator to enhance an immune response against a canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF protein; and/or assist in purification of a canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF protein (e.g., by affinity chromatography). A suitable fusion segment can be a domain of any size that has the desired function (e.g., imparts increased stability, imparts increased immunogenicity to a protein, and/or simplifies purification of a protein). Fusion segments can be joined to amino and/or carboxyl termini of the IL-4-containing domain, or the Flt-3 ligand-containing domain, or the CD40-containing domain, or the CD154-containing domain, or the IL-5-containing domain, or the IL-13-containing domain, or the IFN α -containing domain, or GM-CSF-containing domain, of a protein and can be susceptible to cleavage in order to enable straightforward recovery of either canine interleukin-4, canine or

feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF protein, respectively. Fusion proteins are preferably produced by culturing a recombinant cell transformed with a fusion nucleic acid molecule that encodes a protein including the fusion segment attached to either the carboxyl and/or amino terminal end of an canine interleukin-4-, canine or feline Flt-3 ligand-, canine or feline CD40-, canine or feline CD154-, canine interleukin-5-, canine interleukin-13-, feline interferon alpha-, or feline GM-CSF-containing domain. Preferred fusion segments include a metal binding domain (e.g., a poly-histidine segment); an immunoglobulin binding domain (e.g., Protein A; Protein G; T cell; B cell; Fc receptor or complement protein antibody-binding domain); a sugar binding domain (e.g., a maltose binding domain); and/or a "tag" domain (e.g., at least a portion of -galactosidase, a strep tag peptide, a T7 tag peptide, a FlagTM peptide, or other domains that can be purified using compounds that bind to the domain, such as monoclonal antibodies). More preferred fusion segments include metal binding domains, such as a poly-histidine segment; a maltose binding domain; a strep tag peptide, such as that available from Biometra in Tampa, Fla.; and an S10 peptide.

A suitable fusion segment that links one IL-4 protein to another IL-4 protein, or one Flt-3 ligand protein to another Flt-3 ligand protein, or one CD40 protein to another CD40 protein, or one CD154 protein to another CD154 protein, or one IL-5 protein to another IL-5 protein to another IL-5 protein, or one IL-13 protein to another IL-13 protein, or one IFN α protein to another IFN α protein, or one GM-CSF protein to another GM-CSF protein, includes any amino acid sequence that enables such proteins to be linked while maintaining the biological function of either the canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF proteins, respectively. Selection of a suitable linker is dependent upon how many proteins are to be linked to form one multimeric molecule and from where on either the canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF molecule the linker extends. Preferably, a linker fusion segment of the present invention comprises a peptide of from about 6 amino acid residues to about 40 residues, more preferably from about 6 residues to about 30 residues in length.

In another embodiment, an canine interleukin-4, canine or feline, Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF protein of the present invention also includes at least one additional protein segment that is capable of targeting either canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF protein, respectively, to a desired cell or receptive molecule. Such a multivalent targeting protein can be produced by culturing a cell transformed with a nucleic acid molecule comprising two or more nucleic acid domains joined together in such a manner that the resulting nucleic acid molecule is expressed as a multivalent targeting protein containing a canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF protein or portion thereof and/or at least one targeting

compound capable of delivering the canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF protein, respectively, to a desired site in an animal.

Examples of multivalent targeting proteins include, but are not limited to, a canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF protein of the present invention attached to one or more compounds that can bind to a receptive molecule on the surface of a cell located in an area of an animal where regulation of an immune response is desired. One of skill in the art can select appropriate targeting fusion segments depending upon the cell or receptive molecule being targeted.

Another example of a multivalent protein of the present invention includes, but is not limited to, a canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF protein of the present invention attached to one or more proteins that are potentially antigenic in mammals. Thus, immunogenicity of the potentially antigenic protein could be enhanced by administering to a mammal together with an immunoregulatory protein of the present invention.

A naturally-occurring variant of a canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF protein of the present invention is preferably isolated from (including isolation of the natural protein or production of the protein by recombinant or synthetic techniques) from mammals, including but not limited to dogs (i.e., canids), cats (i.e., felids), horses (i.e., equids), humans, cattle, chinchillas, ferrets, goats, mice, minks, rabbits, raccoons, rats, sheep, squirrels, swine, chickens, ostriches, quail and/or turkeys as well as other furry animals, pets, zoo animals, work animals and/or food animals. Particularly preferred animals from which to isolate canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF proteins are dogs, cats, horses, and/or humans.

A preferred isolated protein of the present invention is a protein encoded by at least one of the following nucleic acid molecules: nCaIL-4₅₄₉, nCaIL-4₃₉₆, nCaIL-4₃₂₄, nCaFlt3L₁₀₁₃, nCaFlt3L₈₈₂, nCaFlt3L₈₀₄, nCaFlt3L₈₂₈, nCaFlt3L₉₈₅, nCaFlt3L₁₀₁₉, nCaFlt3L₉₃, nCaFlt3L₇₅₀, nFeFlt3L₃₉₅, nFeFlt3L₇₉₃, nFeFlt3L₉₄₂, nFeFlt3L₈₇₃, nFeFlt3L₇₉₅, nCaCD40₃₂₁, nCaCD40₁₄₂₅, nCaCD40₈₂₂, nCaCD40₇₆₅, nFeCD40₃₃₆, nCaCD154₃₉₀, nCaCD154₈₇₈, nCaCD154₇₈₀, nCaCD154₆₃₃, nFeCD154₈₈₅, nFeCD154₇₈₀, nFeCD154₆₃₃, nCaIL-5₆₁₀, nCaIL-5₄₀₂, nCaIL-5₃₄₅, nCaIL-13₁₆₆, nCaIL-13₂₇₂, nCaIL-13₂₇₈, nCaIL-13₁₃₀₂, nCaIL-13₃₉₃, nCaIL-13₃₃₃, nCaIL-13₁₂₆₉, nCaIL-13₃₉₀, nCaIL-13₃₃₀, nFeIFN α _{567a}, nFeIFN α _{567b}, nFeIFN α _{498a}, nFeIFN α _{498b}, nFeGMCSF₄₄₄, nFeGMCSF₄₃₂, nFeGMCSF₃₈₁ and/or allelic variants of any of these nucleic acid molecules. Also preferred is an isolated protein that is encoded by a nucleic acid molecule the having nucleic acid sequence SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:19, SEQ ID NO:6, SEQ ID NO:9, SEQ ID NO:22, SEQ ID NO:25, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:46, SEQ ID NO:48, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:55, SEQ ID NO:57, SEQ ID

NO:60, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:72, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:80, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:99, SEQ ID NO:102, SEQ ID NO:104, SEQ ID NO:107, SEQ ID NO:110, SEQ ID NO:113, SEQ ID NO:116, SEQ ID NO:119, SEQ ID NO:122, SEQ ID NO:124; and/or an allelic variant of such a nucleic acid molecule.

Translation of SEQ ID NO:1, the coding strand of nCaIL-4₅₄₉, yields a protein of about 132 amino acids, denoted herein as PCaIL-4₁₃₂, the amino acid sequence of which is presented in SEQ ID NO:2, assuming an open reading frame having an initiation codon spanning from nucleotide 43 through nucleotide 45 of SEQ ID NO:1 and a stop codon spanning from nucleotide 439 through nucleotide 441 of SEQ ID NO:1.

Translation of SEQ ID NO:6, the coding strand of nCaFlt3L₁₀₁₃, yields a protein of about 294 amino acids, denoted herein as PCaFlt3L₂₉₄, the amino acid sequence of which is presented in SEQ ID NO:7, assuming an open reading frame having an initiation codon spanning from nucleotide 35 through nucleotide 37 of SEQ ID NO:6 and a stop codon spanning from nucleotide 917 through nucleotide 919 of SEQ ID NO:6.

Translation of SEQ ID NO:43, the coding strand for nFeFlt3L₉₄₂, yields a protein of about 291 amino acids, denoted herein as PFeFlt3L₂₉₁, the amino acid sequence of which is presented in SEQ ID NO:44, assuming an open reading frame having an initiation codon spanning from nucleotide 31 through nucleotide 33 of SEQ ID NO:43 and a stop codon spanning from nucleotide 904 through nucleotide 906 of SEQ ID NO:43.

Translation of SEQ ID NO:52, the coding strand for nCaCD40₁₄₂₅, yields a protein of about 274 amino acids, denoted herein as PCaCD40₂₇₄, the amino acid sequence of which is presented in SEQ ID NO:53, assuming an open reading frame having an initiation codon spanning from nucleotide 196 through nucleotide 198 of SEQ ID NO:52 and a stop codon spanning from about nucleotide 1018 through nucleotide 1020 of SEQ ID NO:52.

Translation of SEQ ID NO:60, the coding strand for nFeCD40₃₃₆, yields a protein of about 112 amino acids, denoted herein as PFeCD40₁₁₂, the amino acid sequence of which is presented in SEQ ID NO:61, assuming an open reading frame having an initiation codon spanning from nucleotide 1 through nucleotide 3 of SEQ ID NO:60.

Translation of SEQ ID NO:64, the coding strand for nCaCD154₁₈₇₈, yields a protein of about 260 amino acids, denoted herein as PCaCD154₂₆₀, the amino acid sequence of which is presented in SEQ ID NO:65, assuming an open reading frame having an initiation codon spanning from nucleotide 284 through nucleotide 286 of SEQ ID NO:64 and a stop codon spanning from nucleotide 1064 through nucleotide 1066 of SEQ ID NO:64.

Translation of SEQ ID NO:72, the coding strand for nFeCD154₈₈₅, yields a protein of about 260 amino acids, denoted herein as PFeCD154₂₆₀, the amino acid sequence of which is presented in SEQ ID NO:73, assuming an open reading frame having an initiation codon spanning from nucleotide 29 through nucleotide 31 of SEQ ID NO:72, and a stop codon spanning from nucleotide 809 through nucleotide 811 of SEQ ID NO:72.

Translation of SEQ ID NO:80, the coding strand for nCaIL-5₆₁₀, yields a protein of about 134 amino acids, denoted herein as PCaIL-5₁₃₄, the amino acid sequence of which is presented in SEQ ID NO:81, assuming an open

reading frame having an initiation codon spanning from nucleotide 29 through nucleotide 31 of SEQ ID NO:80, and a stop codon spanning from nucleotide 431 through nucleotide 433 of SEQ ID NO:80.

Translation of SEQ ID NO:91, the coding strand for nCaIL-13₁₃₀₂, yields a protein of about 131 amino acids, denoted herein as PCaIL-13₁₃₁, the amino acid sequence of which is presented in SEQ ID NO:92, assuming an open reading frame having an initiation codon spanning from nucleotide 52 through nucleotide 54 of SEQ ID NO:91 and a stop codon spanning from nucleotide 445 through nucleotide 447 of SEQ ID NO:91.

Translation of SEQ ID NO:107, the coding strand for nFeIFN α _{567a}, yields a protein of about 189 amino acids, denoted herein as PFeIFN α _{189a}, the amino acid sequence of which is presented in SEQ ID NO:108, assuming an open reading frame having an initiation codon spanning from nucleotide 1 through nucleotide 3 and a last codon prior to a stop codon spanning from nucleotide 565 through nucleotide 567 of SEQ ID NO:107.

Translation of SEQ ID NO:119, the coding strand for nFeGMCSF₄₄₄, yields a protein of about 144 amino acids, denoted herein as PFeGMCSF₁₄₄, the amino acid sequence of which is presented in SEQ ID NO:120, assuming an open reading frame having an initiation codon spanning from nucleotide 10 through nucleotide 12 of SEQ ID NO:119 and a stop codon spanning from nucleotide 442 through nucleotide 444 of SEQ ID NO:119.

Preferred IL-4 proteins of the present invention include proteins that are at least about 85%, preferably at least about 90%, and even more preferably at least about 95% identical to PCaIL-4₁₃₂, PCaIL-4₁₀₈, or fragments thereof. Preferred Flt-3 ligand proteins of the present invention include proteins that are at least about 75%, even more preferably at least about 80%, even more preferably at least about 85%, even more preferably at least about 90%, and even more preferably at least about 95% identical to PCaFlt3L₂₉₄, PCaFlt3L₂₆₈, PCaFlt3L₂₇₆, PCaFlt3L₂₅₀, PCaFlt3L₃₁, and/or fragments thereof. Additional preferred Flt-3 ligand proteins of the present invention includes proteins that are at least about 75%, even more preferably at least about 80%, even more preferably at least about 85%, even more preferably at least about 90%, and even more preferably at least about 95% identical to PFeFlt3L₂₉₁, PFeFlt3L₂₆₅ and/or fragments thereof. Preferred CD40 proteins of the present invention includes proteins that are at least about 70%, preferably at least about 75%, even more preferably at least about 80%, even more preferably at least about 85%, even more preferably at least about 90%, and even more preferably at least about 95% identical to PCaCD40₂₇₄, PCaCD40₂₅₅ and/or fragments thereof. Additional preferred CD40 proteins of the present invention includes proteins that are at least about 60%, at least about 65%, preferably at least about 70%, preferably at least about 75%, even more preferably at least about 80%, even more preferably at least about 85%, even more preferably at least about 90%, and even more preferably at least about 95% identical to PFeCD40₁₁₂ and/or fragments thereof. Preferred CD154 proteins of the present invention includes proteins that are at least about 80% identical, preferably at least about 85% identical, even more preferably at least about 90%, and even more preferably at least about 95% identical to PCaCD154₂₆₀, PCaCD154₂₁₁ and/or fragments thereof. Additional preferred CD154 proteins of the present invention includes proteins that are at least about 85% identical, even more preferably at least about 90%, and even more preferably at least about 95% identical to PFeCD154₂₆₀,

PFeCD154₂₁₁ and/or fragments thereof. Preferred IL-5 proteins of the present invention includes proteins that are at least about 85% identical, even more preferably at least about 90%, and even more preferably at least about 95% identical to PCaIL-5₁₃₄, PCaIL-5₁₁₅ and/or fragments thereof. Preferred IL-13 proteins of the present invention includes proteins that are at least about 70% identical, preferably at least about 75% identical, more preferably at least about 80% identical, more preferably at least about 85% identical, even more preferably at least about 90%, and even more preferably at least about 95% identical to PCaIL-13₁₃₁, PCaIL-13₁₁₁, PCaIL-13₁₃₀, PCaIL-13110, and/or fragments thereof. Preferred IFN α proteins of the present invention include PFeIFN α _{189a}, PFeIFN α _{189b}, PFeIFN α _{166a}, and/or PFeIFN α _{166b}. Preferred GM-CSF proteins of the present invention include PFeGMCSF₁₄₄, and/or PFeGMCSF₁₂₇.

More preferred are IL-4 proteins comprising PCaIL-4₁₃₂, PCaIL-4₁₀₈, and/or proteins encoded by allelic variants of a nucleic acid molecule encoding proteins PCaIL-4₁₃₂ and/or PCaIL-4₁₀₈. More preferred are Flt-3 ligand proteins comprising PCaFlt3L₂₉₄, PCaFlt3L₂₆₈, PCaFlt3L₂₇₆, PCaFlt3L₂₅₀, PCaFlt3L₃₁, PFeFlt3L₂₉₁, PFeFlt3L₂₆₅ and/or proteins encoded by allelic variants of a nucleic acid molecule encoding proteins PCaFlt3L₂₉₄, PCaFlt3L₂₆₈, PCaFlt3L₂₇₆, PCaFlt3L₂₅₀, PCaFlt3L₃₁, PFeFlt3L₂₉₁, and/or PFeFlt3L₂₆₅. More preferred are CD40 proteins comprising PCaCD40₂₇₄, PCaCD40₂₅₅, and/or PFeCD40₁₁₂ and/or proteins encoded by allelic variants of a nucleic acid molecule encoding proteins PCaCD40₂₇₄, PCaCD40₂₅₅, and/or PFeCD40₁₁₂. More preferred are CD154 proteins comprising PCaCD154₂₆₀, PCaCD154₂₁₁, PFeCD154₂₆₀, PFeCD154₂₁₁ and/or proteins encoded by allelic variants of a nucleic acid molecule encoding one of proteins PCaCD154₂₆₀, PCaCD154₂₁₁, PFeCD154₂₆₀, PFeCD154₂₁₁. More preferred are IL-5 proteins comprising PCaIL-5₁₃₄, PCaIL-5₁₁₅ and/or proteins encoded by allelic variants of a nucleic acid molecule encoding one of the proteins PCaIL-5₁₃₄ and/or PCaIL-5₁₁₅. More preferred are IL-13 proteins comprising PCaIL-13₁₃₁, PCaIL-13₁₁₁, PCaIL-13₁₃₀, PCaIL-13₁₁₀, and/or proteins encoded by allelic variants of a nucleic acid molecule encoding one of the proteins PCaIL-13₁₃₁, PCaIL-13₁₁₁, PCaIL-13₁₃₀, PCaIL-13₁₁₀.

Also preferred are IL-4 proteins of the present invention having amino acid sequences that are at least about 85%, preferably at least about 90%, and even more preferably at least about 95% identical to SEQ ID NO:2, SEQ ID NO:20 and/or fragments thereof. Also preferred are Flt-3 ligand proteins of the present invention having amino acids sequences that are at least about 75%, even more preferably at least about 80%, even more preferably at least about 85%, even more preferably at least about 90%, and even more preferably at least about 95% identical to SEQ ID NO:7, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:31, and/or SEQ ID NO:34 and/or fragments thereof. Additional preferred are Flt-3 ligand proteins of the present invention includes proteins that are at least about 75%, even more preferably at least about 80%, even more preferably at least about 85%, even more preferably at least about 90%, and/or even more preferably at least about 95% identical to

SEQ ID NO:53, SEQ ID NO:58 and/or fragments thereof. Additional preferred CD40 proteins of the present invention includes proteins that are at least about 60%, at least about 65%, preferably at least about 70%, preferably at least about 75%, even more preferably at least about 80%, even more preferably at least about 85%, even more preferably at least about 90%, and even more preferably at least about 95% identical to SEQ ID NO:61 and/or fragments thereof. Preferred CD154 proteins of the present invention includes proteins that are at least about 80% identical, preferably at least about 85% identical, even more preferably at least about 90%, and even more preferably at least about 95% identical to SEQ ID NO:65, SEQ ID NO:70 and/or fragments thereof. Additional preferred CD154 proteins of the present invention includes proteins that are at least about 85% identical, even more preferably at least about 90%, and even more preferably at least about 95% identical to SEQ ID NO:73, SEQ ID NO:78 and/or fragments thereof. Preferred IL-5 proteins of the present invention includes proteins that are at least about 85% identical, even more preferably at least about 90%, and even more preferably at least about 95% identical to SEQ ID NO:81, SEQ ID NO:86 and/or fragments thereof. Preferred IL-13 proteins of the present invention includes proteins that are at least about 70% identical, preferably at least about 75% identical, more preferably at least about 80% identical, more preferably at least about 85% identical, even more preferably at least about 90%, and even more preferably at least about 95% identical to SEQ ID NO:92, SEQ ID NO:97, SEQ ID NO:100, SEQ ID NO:105, and/or fragments thereof. Preferred IFN α proteins of the present invention include SEQ ID NO:108, SEQ ID NO:11, SEQ ID NO:114, SEQ ID NO:117. Preferred GM-CSF proteins of the present invention include SEQ ID NO:120, SEQ ID NO:125.

More preferred are IL-4 proteins comprising the amino acid sequence SEQ ID NO:2, SEQ ID NO:20; and/or IL-4 proteins encoded by allelic variants of nucleic acid molecules encoding IL-4 proteins having the amino acid sequence SEQ ID NO:2, SEQ ID NO:20. More preferred are Flt-3 ligand proteins comprising SEQ ID NO:7, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:31 and/or SEQ ID NO:34, SEQ ID NO:44, SEQ ID NO:49 and/or proteins encoded by allelic variants of a nucleic acid molecule encoding proteins SEQ ID NO:7, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:44, and/or SEQ ID NO:49. More preferred are CD40 proteins comprising SEQ ID NO:53, SEQ ID NO:58, SEQ ID NO:61 and/or proteins encoded by allelic variants of a nucleic acid molecule encoding proteins SEQ ID NO:53, SEQ ID NO:58 and/or SEQ ID NO:61. More preferred are CD154 proteins comprising SEQ ID NO:65, SEQ ID NO:70, SEQ ID NO:73, SEQ ID NO:78 and/or proteins encoded by allelic variants of a nucleic acid molecule encoding one of proteins SEQ ID NO:65, SEQ ID NO:70, SEQ ID NO:73, and/or SEQ ID NO:78. More preferred are IL-5 proteins comprising SEQ ID NO:81, SEQ ID NO:86 and/or proteins encoded by allelic variants of a nucleic acid molecule encoding one of the proteins SEQ ID NO:81, and/or SEQ ID NO:86. More preferred are IL-13 proteins comprising SEQ ID NO:92, SEQ ID NO:97, SEQ ID NO:100, SEQ ID NO:105, and/or proteins encoded by allelic variants of a nucleic acid molecule encoding one of the proteins SEQ ID NO:92, SEQ ID NO:97, SEQ ID NO:100, and/or SEQ ID NO:105.

Percent identities between amino acid or nucleic acid sequences can be determined using standard methods known to those of skill in the art. It is known in the art that methods to determine the percentage identity and the number of gaps

are substantially similar when different methods for determining sequence similarity are used and when the degree of similarity is greater than 30% amino acid identity, as described by Johnson et al., J. Mol. Biol., vol. 233, pages 716-738, 1993, and Feng et al., J. Mol. Evol., vol. 21, pages 112-125, 1985, which are incorporated by reference herein in their entirety. Preferred methods to determine percentage identities between amino acid sequences and between nucleic acid sequences include comparisons using various computer programs such as GCG™ program (available from Genetics Computer Group, Madison, Wis.), DNAsis™ program (available from Hitachi Software, San Bruno, Calif.) or the MacVector™ program (available from the Eastman Kodak Company, New Haven, Conn.). Preferred settings for sequence comparisons using the DNAsis™ computer program or the GAP GCG™ program are disclosed herein in the Examples section.

Additional preferred IL-4 proteins of the present invention include proteins encoded by nucleic acid molecules comprising at least a portion of nCaIL-4₅₄₉, nCaIL-4₃₉₆, and/or nCaIL-4₃₂₄, as well as IL-4 proteins encoded by allelic variants of such nucleic acid molecules. Additional preferred Flt-3 ligand proteins of the present invention include proteins encoded by nucleic acid molecules comprising at least a portion of nCaFlt3L₁₀₁₃, nCaFlt3L₈₈₂, nCaFlt3L₈₀₄, nCaFlt3L₈₂₈, nCaFlt3L₉₈₅, nCaFlt3L₁₀₁₉, nCaFlt3L₉₃, nCaFlt3L₇₅₀, nFeFlt3L₃₉₅, nFeFlt3L₇₉₃, nFeFlt3L₉₄₂, nCaFlt3L₈₇₃, and/or nFeFlt3L₇₉₅ as well as Flt-3 ligand proteins encoded by allelic variants of such nucleic acid molecules. Additional preferred CD40 proteins of the present invention include proteins encoded by nucleic acid molecules encoding at least a portion of nCaCD40₃₂₁, nCaCD40₁₄₂₅, nCaCD40₈₂₂, nCaCD40₇₆₅, and/or nFeCD40₃₃₆ as well as CD40 proteins encoded by allelic variants of such nucleic acid molecules. Additional preferred CD154 proteins of the present invention include proteins encoded by nucleic acid molecules encoding at least a portion of nCaCD154₃₉₀, nCaCD154₈₇₉, nCaCD154₇₈₀, nCaCD154₆₃₃, nFeCD154₈₈₅, nFeCD154₇₈₀, and/or nFeCD154₆₃₃ as well as CD154 proteins encoded by allelic variants of such nucleic acid molecules. Additional preferred IL-5 proteins of the present invention include proteins encoded by nucleic acid molecules encoding at least a portion of nCaIL-5₆₁₀, nCaIL-5₄₀₂, and/or nCaIL-5₃₄₅ as well as IL-5 proteins encoded by allelic variants of such nucleic acid molecules. Additional preferred IL-13 proteins of the present invention include proteins encoded by nucleic acid molecules encoding at least a portion of nCaIL-5₆₁₀, nCaIL-5₄₀₂, and/or nCaIL-5₃₄₅ as well as IL-13 proteins encoded by allelic variants of such nucleic acid molecules.

Also preferred are IL-4 proteins encoded by nucleic acid molecules having nucleic acid sequences comprising at least a portion of SEQ ID NO:1, SEQ ID NO:4, and/or SEQ ID NO:19, as well as allelic variants of these nucleic acid molecules. Also preferred are Flt-3 ligand proteins encoded by nucleic acid molecules having nucleic acid sequences comprising at least a portion of SEQ ID NO:6, SEQ ID NO:9, SEQ ID NO:22, SEQ ID NO:25, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:33, SEQ ID NO:36, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:46, and/or SEQ ID NO:48, as well as allelic variants of these nucleic acid molecules. Also preferred are CD40 proteins encoded by nucleic acid molecules having nucleic acid sequences comprising at least a portion of SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:55, SEQ ID NO:57, and/or SEQ ID NO:60, as well as allelic variants of these nucleic acid molecules. Also preferred are CD154 proteins encoded

by nucleic acid molecules having nucleic acid sequences comprising at least a portion of SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:67, SEQ ID NO:69, SEQ ID NO:72, SEQ ID NO:75, and/or SEQ ID NO:77, as well as allelic variants of these nucleic acid molecules. Also preferred are IL-5 proteins encoded by nucleic acid molecules having nucleic acid sequences comprising at least a portion of SEQ ID NO:80, SEQ ID NO:83, and/or SEQ ID NO:85, as well as allelic variants of these nucleic acid molecules. Also preferred are IL-13 proteins encoded by nucleic acid molecules having nucleic acid sequences comprising at least a portion of SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:99, SEQ ID NO:102, and/or SEQ ID NO:104, as well as allelic variants of these nucleic acid molecules.

Another embodiment of the present invention is a canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF nucleic acid molecule that includes one or more regulatory regions, full-length or partial coding regions, or combinations thereof. The minimal size of a nucleic acid molecule of the present invention is a size sufficient to allow the formation of stable hybrid (i.e., hybridization under stringent hybridization conditions) with the complementary sequence of another nucleic acid molecule. As such, the minimal size of a canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF nucleic acid molecule of the present invention is from about 12 to about 18 nucleotides in length.

In accordance with the present invention, an isolated nucleic acid molecule is a nucleic acid molecule that has been removed from its natural milieu (i.e., that has been subjected to human manipulation) and can include DNA, RNA, or derivatives of either DNA or RNA. As such, "isolated" does not reflect the extent to which the nucleic acid molecule has been purified. An isolated canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF nucleic acid molecule of the present invention can be isolated from its natural source or produced using recombinant DNA technology (e.g., polymerase chain reaction (PCR) amplification or cloning) or chemical synthesis. Isolated canine, interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, and/or feline GM-CSF, nucleic acid molecules can include, for example, natural allelic variants and/or nucleic acid molecules modified by nucleotide insertions, deletions, substitutions, and/or inversions in a manner such that the modifications do not substantially interfere with the nucleic acid molecule's ability to encode an canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, and/or feline GM-CSF protein of the present invention.

A canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, and/or feline GM-CSF ligand nucleic acid molecule homolog can be produced using a number of methods known to those skilled in the art, see, for example, Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Labs Press: Sambrook et al., *ibid.*, is incorporated by reference herein in its entirety. For

example, nucleic acid molecules can be modified using a variety of techniques including, but not limited to, classic mutagenesis and recombinant DNA techniques such as site-directed mutagenesis, chemical treatment, restriction enzyme cleavage, ligation of nucleic acid fragments, PCR amplification, synthesis of oligonucleotide mixtures and ligation of mixture groups to "build" a mixture of nucleic acid molecules, and combinations thereof. Nucleic acid molecule homologs can be selected by hybridization with either a canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF nucleic acid molecule or by screening the function of a protein encoded by the nucleic acid molecule (e.g., ability to elicit an immune response against at least one epitope of a canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF protein, respectively).

An isolated nucleic acid molecule of the present invention can include a nucleic acid sequence that encodes at least one canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF protein of the present invention, examples of such proteins being disclosed herein. Although the phrase "nucleic acid molecule" primarily refers to the physical nucleic acid molecule and the phrase "nucleic acid sequence" primarily refers to the sequence of nucleotides on the nucleic acid molecule, the two phrases can be used interchangeably, especially with respect to a nucleic acid molecule, or a nucleic acid sequence, being capable of encoding a canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF ligand protein.

A preferred nucleic acid molecule of the present invention, when administered to an animal, is capable of regulating an immune response in an animal. As will be disclosed in more detail below, such a nucleic acid molecule can be, or encode, an antisense RNA, a molecule capable of triple helix formation, a ribozyme, or other nucleic acid-based drug compound. In additional embodiments, a nucleic acid molecule of the present invention can encode an immunoregulatory protein (e.g., a cell-bound or soluble protein of the present invention), the nucleic acid molecule being delivered to the animal, for example, by direct injection (i.e. as a genetic vaccine) or in a vehicle such as a recombinant virus vaccine or a recombinant cell vaccine.

One embodiment of the present invention is an IL-4 nucleic acid molecule comprising all or part (i.e., a fragment of the IL-4 nucleic acid molecule) of nucleic acid molecules nCaIL-4₅₄₉, nCaIL-4₃₉₆, and/or nCaIL-4₃₂₄, or allelic variants of these nucleic acid molecules. One embodiment of the present invention is a Flt-3 ligand nucleic acid molecule comprising all or part (i.e., a fragment of the Flt-3 ligand nucleic acid molecule) of nucleic acid molecules nCaFlt3L₁₀₁₃, nCaFlt3L₈₈₂, nCaFlt3L₈₀₄, nCaFlt3L₈₂₈, nCaFlt3L₉₈₅, nCaFlt3L₁₀₁₉, nCaFlt3L₉₃, nCaFlt3L₇₅₀, nFeFlt3L₃₉₅, nFeFlt3L₇₉₃, nFeFlt3L₉₄₂, nFeFlt3L₈₇₃, and/or nFeFlt3L₇₉₅ and/or allelic variants of these nucleic acid molecules. One embodiment of the present invention is a CD40 nucleic acid molecule comprising all or part (i.e., a fragment of the CD40 nucleic acid molecule) of nucleic acid molecules nCaCD40₃₂₁, nCaCD40₁₄₂₅, nCaCD40₈₂₂, nCaCD40₇₆₅, and/or nFeCD40₃₃₆ and/or allelic variants of these nucleic acid molecules. One embodiment of the

present invention is a CD154 nucleic acid molecule comprising all or part of nucleic acid molecules nCaCD154₃₉₀, nCaCD154₁₈₇₈, nCaCD154₇₈₀, nCaCD154₆₃₃, nFeCD154₈₈₅, nFeCD154₇₈₀, and/or nFeCD154₆₃₃, and/or allelic variants of these nucleic acid molecules. One embodiment of the present invention is an IL-5 nucleic acid molecule comprising all or part of nucleic acid molecules nCaIL-5₆₁₀, nCaIL-5₄₀₂, and/or nCaIL-5₃₄₅, and/or allelic variants of these nucleic acid molecules. One embodiment of the present invention is an IL-13 nucleic acid molecule comprising all or part of nucleic acid molecules nCaIL-13₁₆₆, nCaIL-13₂₇₂, nCaIL-13₂₇₈, nCaIL-13₁₃₀₂, nCaIL-13₃₉₃, nCaIL-13₃₃₃, nCaIL-13₁₂₆₉, nCaIL-13₃₉₀, and/or nCaIL-13₃₃₀, and/or allelic variants of these nucleic acid molecules. Another preferred nucleic acid molecule of the present invention includes at least a portion of (i.e., a fragment of the nucleic acid molecule) nucleic acid sequence SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:118, SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:122, SEQ ID NO:123, SEQ ID NO:124; and/or SEQ ID NO:126, as well as allelic variants of nucleic acid molecules having these nucleic acid sequences. Such nucleic acid molecules can include nucleotides in addition to those included in the SEQ ID NOs, such as, but not limited to, a full-length gene, a full-length coding region, a nucleic acid molecule encoding a fusion protein, and/or a nucleic acid molecule encoding a multivalent therapeutic compound.

One embodiment of an isolated nucleic acid molecule of the present invention is a nucleic acid molecule that can be any of the following: (a) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:19, and/or SEQ ID NO:21 and/or a homolog thereof, wherein said homolog has an at least 50 contiguous nucleotide region identical in sequence to a 50 contiguous nucleotide region of a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:19, and/or SEQ ID NO:21; (b) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:33,

SEQ ID NO:35, SEQ ID NO:36, and/or SEQ ID NO:37, and/or a homolog thereof, wherein said homolog has an at least 40 contiguous nucleotide region identical in sequence to a 40 contiguous nucleotide region of a nucleic acid sequence having a nucleic acid sequence selected from the group consisting of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:36, and/or SEQ ID NO:37; (c) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, and/or SEQ ID NO:50, and/or a homolog thereof, wherein said homolog has an at least 30 contiguous nucleotide region identical in sequence to a 30 contiguous nucleotide region of a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, and/or SEQ ID NO:50; (d) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, and/or SEQ ID NO:59, and/or a homolog thereof, wherein said homolog has an at least 40 contiguous nucleotide region identical in sequence to a 40 contiguous nucleotide region of a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, and/or SEQ ID NO:59; (e) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:60 and/or SEQ ID NO:62 and/or a homolog thereof, wherein said homolog has an at least 30 contiguous nucleotide region identical in sequence to a 30 contiguous nucleotide region of a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:60 and/or SEQ ID NO:62; (f) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69 and/or SEQ ID NO:71, and/or a homolog thereof, wherein said homolog has an at least 45 contiguous nucleotide region identical in sequence to a 45 contiguous nucleotide region of a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, and/or SEQ ID NO:71; (g) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, and/or SEQ ID NO:79, and/or a homolog thereof, wherein said homolog has an at least 35 contiguous nucleotide region identical in sequence to a 35 contiguous nucleotide region of a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, and/or SEQ ID NO:79; (h) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, and/or SEQ ID NO:87, and/or a homolog thereof, wherein said homolog has an at least 45 contiguous nucleotide region identical in sequence to a 45 contiguous nucleotide region of a nucleic acid molecule having a nucleic acid sequence selected from

the group consisting of SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85 and/or SEQ ID NO:87; (i) an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, and/or SEQ ID NO:106, and/or a homolog thereof, wherein said homolog has an at least 15 contiguous nucleotide region identical in sequence to a 15 contiguous nucleotide region of a nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, and/or SEQ ID NO:106; (j) an isolated nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:116, and/or SEQ ID NO:118, and/or (k) an isolated nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:122, SEQ ID NO:123, SEQ ID NO:124, and/or SEQ ID NO:126. The phrase, a homolog having an at least "x" contiguous nucleotide region identical in sequence to an "x" contiguous nucleotide region of a nucleic acid molecule selected from the group consisting of SEQ ID NO:"y", refers to an "x"-nucleotide in length nucleic acid molecule that is identical in sequence to an "x"-nucleotide portion of SEQ ID NO:"y", as well as to nucleic acid molecules that are longer in length than "x". The additional length may be in the form of nucleotides that extend from either the 5' or the 3' end(s) of the contiguous identical "x"-nucleotide portion. The 5' and/or 3' extensions can include one or more extensions that have no identity to an immunoregulatory molecule of the present invention, as well as extensions that show similarity or identity to cited nucleic acids sequences or portions thereof.

In another embodiment, an isolated nucleic acid molecule of the present invention can be any of the following: (a) a nucleic acid molecule having a nucleic acid sequence encoding an IL-4 protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 85 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:2 and/or SEQ ID NO:20 and/or (ii) a protein comprising a fragment of at least 20 amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:2 and/or SEQ ID NO:20; (b) a nucleic acid molecule having a nucleic acid sequence encoding a Flt-3 ligand protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 75 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:7, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:31, and/or SEQ ID NO:34, and/or (ii) a protein comprising a fragment of at least 25 amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:7, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:31, and/or SEQ ID NO:34; (c) a nucleic acid molecule having a nucleic acid sequence encoding a Flt-3 ligand protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 75 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:44 and/or SEQ ID NO:49 and/or

(ii) a protein comprising a fragment of at least 25 amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:44 and/or SEQ ID NO:49; (d) a nucleic acid molecule having a nucleic acid sequence encoding a CD40 protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 70 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:53 and/or SEQ ID NO:58 and/or (ii) a protein comprising a fragment of at least 30 amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:53 and/or SEQ ID NO:58; (e) a nucleic acid molecule having a nucleic acid sequence encoding a CD40 protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 60 percent identical to an amino acid sequence comprising SEQ ID NO:61 and/or (ii) a protein comprising a fragment of at least 20 amino acids of an amino acid sequence comprising SEQ ID NO:61; (f) a nucleic acid molecule having a nucleic acid sequence encoding a CD154 protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 80 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:65 and/or SEQ ID NO:70; and/or (ii) a protein comprising a fragment of at least 35 amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:65 and/or SEQ ID NO:70; (g) a nucleic acid molecule having a nucleic acid sequence encoding a CD154 protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 85 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:73 and/or SEQ ID NO:78; and/or (ii) a protein comprising a fragment of at least 50 amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:73 and/or SEQ ID NO:78; (h) a nucleic acid molecule having a nucleic acid sequence encoding an IL-5 protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 85 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:81 and/or SEQ ID NO:86 and/or (ii) a protein comprising a fragment of at least 20 amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:81 and/or SEQ ID NO:86; (i) a nucleic acid molecule having a nucleic acid sequence encoding an IL-13 protein selected from the group consisting of (i) a protein having an amino acid sequence that is at least about 70 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:92, SEQ ID NO:97, SEQ ID NO:100, and/or SEQ ID NO:105 and/or (ii) a protein comprising a fragment of at least 15 amino acids of an amino acid sequence selected from the group consisting of SEQ ID NO:92, SEQ ID NO:97, SEQ ID NO:100, and/or SEQ ID NO:105; (j) a nucleic acid molecule having a nucleic acid sequence encoding an interferon alpha protein having an amino acid sequence that is selected from the group consisting of amino acid sequence SEQ ID NO:108, SEQ ID NO:111, SEQ ID NO:114, and/or SEQ ID NO:117; (k) a nucleic acid molecule having a nucleic acid sequence encoding a GM-CSF protein having an amino acid sequence that is selected from the group consisting of amino acid sequence SEQ ID NO:120, SEQ ID NO:126 and/or (l) a nucleic acid molecule comprising a complement of any of the before-mentioned nucleic acid sequences, wherein said IL-4 protein elicits an immune response against an IL-4 protein selected from the group consisting of SEQ ID NO:2 and/or SEQ ID NO:20 and/or is a protein with interleukin-4 activity, said Flt-3 ligand protein elicits an immune response against a

Flt-3 ligand protein selected from the group consisting of SEQ ID NO:7, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:44, and/or SEQ ID NO:49 and/or is a protein with Flt-3 ligand activity, said CD40 protein elicits an immune response against a CD40 protein selected from the group consisting of SEQ ID NO:53, SEQ ID NO:58, and/or SEQ ID NO:61 and/or is a protein with CD40 activity, said CD154 protein elicits an immune response against a CD154 protein selected from the group consisting of SEQ ID NO:65, SEQ ID NO:70, SEQ ID NO:73, and/or SEQ ID NO:78 and/or is a protein with CD154 activity, said IL-5 protein elicits an immune response against a IL-5 protein selected from the group consisting of SEQ ID NO:81 and/or SEQ ID NO:86 and/or is a protein with IL-5 activity, said IL-13 protein elicits an immune response against an IL 13 protein selected from the group consisting of SEQ ID NO:92, SEQ ID NO:97, SEQ ID NO:100, and/or SEQ ID NO:105 and/or is a protein with IL-13 activity, said interferon alpha protein elicits an immune response against an interferon alpha protein selected from the group consisting of SEQ ID NO:108, SEQ ID NO:111, SEQ ID NO:114, and/or SEQ ID NO:117 and/or is a protein with interferon alpha activity, and said GM-CSF protein elicits an immune response against a GM-CSF protein selected from the group consisting of SEQ ID NO:120 and/or SEQ ID NO:125 and/or is a protein with GM-CSF activity.

In one embodiment, an IL-4 nucleic acid molecule of the present invention encodes a protein that is at least about 85%, preferably at least about 90%, preferably at least about 92%, and even more preferably at least about 95% identical to PCaIL-4₁₃₂ and/or PCaIL-4₁₀₈. In one embodiment, a Flt-3 ligand nucleic acid molecule of the present invention encodes a protein that is at least about 75%, even more preferably at least about 80%, even more preferably at least about 85%, even more preferably at least about 90%, and even more preferably at least about 95% identical to PCaFlt3L₂₉₄, PCaFlt3L₂₆₈, PCaFlt3L₂₇₆, PCaFlt3L₂₅₀, and/or PCaFlt3L₃₁. In one embodiment, a Flt-3 ligand nucleic acid molecule of the present invention encodes a protein that is at least about 75%, more preferably at least about 80%, even more preferably at least about 85%, even more preferably at least about 90%, and even more preferably at least about 95% identical to PFeFlt3L₂₉₁, and/or PFeFlt3L₂₆₅. In one embodiment, a CD40 nucleic acid molecule of the present invention encodes a protein that is at least about 60%, preferably at least about 65%, preferably at least about 70%, preferably at least about 75%, even more preferably at least about 80%, even more preferably at least about 85%, even more preferably at least about 90%, and even more preferably at least about 95% identical to PFeCD40₁₁₂. In one embodiment, a CD154 nucleic acid molecule of the present invention encodes a protein that is at least about 80%, at least about 85%, more preferably at least about 90%, and even more preferably at least about 95% identical to PCaCD154₂₆₀, and/or PCaCD154₂₁₁. In one embodiment, a CD154 nucleic acid molecule of the present invention encodes a protein that is at least about 85%, more preferably at least about 90%, and even more preferably at least about 95% identical to PFeCD154₂₆₀, PFeCD154₂₁₁. In one embodiment, an IL-5 nucleic acid molecule of the present invention encodes a protein that is at least about 85%, more preferably at least about 90%, and even more preferably at least about 95% identical to PCaIL-5₁₃₄, and/or PCaIL-5₁₁₅. In one

embodiment, an IL-13 nucleic acid molecule of the present invention encodes a protein that is at least about 70%, at least about 75%, at least about 80%, preferably at least about 85%, more preferably at least about 90%, and even more preferably at least about 95% identical to PCaIL-13₁₃₁, PCaIL-13₁₁₁, PCaIL-13₁₃₀, PCaIL-13₁₁₀. Even more preferred is a nucleic acid molecule encoding PCaIL-4₁₃₂, PCaIL-4₁₀₈, PCaFlt3L₂₉₄, PCaFlt3L₂₆₈, PCaFlt3L₂₇₆, PCaFlt3L₂₅₀, PCaFlt3L₃₁, PFeFlt3L₂₉₁, PFeFlt3L₂₆₅, PCaCD40₂₇₄, PCaCD40₂₅₅, PFeCD40₁₁₂, PCaCD154₂₆₀, PCaCD154₂₁₁, PFeCD154₂₆₀, PFeCD154₂₁₁, PCaIL-5₁₃₄, PCaIL-5₁₁₅, PCaIL-13₁₃₁, PCaIL-13₁₁₁, PCaIL-13₁₃₀, PCaIL-13₁₁₀ and/or an allelic variant of such a nucleic acid molecule.

In another embodiment, an IL-4 nucleic acid molecule of the present invention encodes a protein having an amino acid sequence that is at least about 85%, preferably at least about 90%, and even more preferably at least about 95% identical to SEQ ID NO:2, SEQ ID NO:20. The present invention also includes an IL-4 nucleic acid molecule encoding a protein having at least a portion of SEQ ID NO:2, and/or SEQ ID NO:20, as well as allelic variants of an IL-4 nucleic acid molecule encoding a protein having these sequences, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In another embodiment, a Flt-3 ligand nucleic acid molecule of the present invention encodes a protein having an amino acid sequence that is at least about 75%, even more preferably at least about 80%, even more preferably at least about 85%, even more preferably at least about 90%, and even more preferably at least about 95% identical to SEQ ID NO:7, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:31, and/or SEQ ID NO:34. The present invention also includes a Flt-3 ligand nucleic acid molecule encoding a protein having at least a portion of SEQ ID NO:7, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:31, and/or SEQ ID NO:34, as well as allelic variants of a Flt-3 ligand nucleic acid molecule encoding a protein having these sequences, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In another embodiment, a Flt-3 ligand nucleic acid molecule of the present invention encodes a protein having an amino acid sequence that is at least about 75%, more preferably at least about 80%, even more preferably at least about 85%, even more preferably at least about 90%, and even more preferably at least about 95% identical to SEQ ID NO:44, and/or SEQ ID NO:49. The present invention also includes a Flt-3 ligand nucleic acid molecule encoding a protein having at least a portion of SEQ ID NO:44, and/or SEQ ID NO:49, as well as allelic variants of a Flt-3 ligand nucleic acid molecule encoding a protein having these sequences, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In another embodiment, a CD40 nucleic acid molecule of the present invention encodes a protein having an amino acid sequence that is at least about 70%, preferably at least about 75%, even more preferably at least about 80%, even more preferably at least about 85%, even more preferably at least about 90%, and even more preferably at least about 95% identical to SEQ ID NO:53 and/or SEQ ID NO:58. The present invention also includes a CD40 nucleic acid molecule encoding a protein having at least a portion of SEQ ID

NO:53 and/or SEQ ID NO:58, as well as allelic variants of a CD40 nucleic acid molecule encoding a protein having these sequences, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In another embodiment, a CD40 nucleic acid molecule of the present invention encodes a protein having an amino acid sequence that is at least about 60%, preferably at least about 65%, preferably at least about 70%, preferably at least about 75%, even more preferably at least about 80%, even more preferably at least about 85%, even more preferably at least about 90%, and even more preferably at least about 95% identical to SEQ ID NO:60. The present invention also includes a CD40 nucleic acid molecule encoding a protein having at least a portion of SEQ ID NO:60, as well as allelic variants of a CD40 nucleic acid molecule encoding a protein having these sequences, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In another embodiment, a CD154 nucleic acid molecule of the present invention encodes a protein having an amino acid sequence that is at least about at least about 80%, at least about 85%, more preferably at least about 90%, and even more preferably at least about 95% identical to SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:67, and/or SEQ ID NO:69. The present invention also includes a CD154 nucleic acid molecule encoding a protein having at least a portion of SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:67, and/or SEQ ID NO:69, as well as allelic variants of a CD154 nucleic acid molecule encoding a protein having these sequences, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In another embodiment, a CD154 nucleic acid molecule of the present invention encodes a protein having an amino acid sequence that is at least about at least about 85%, more preferably at least about 90%, and even more preferably at least about 95% identical to SEQ ID NO:72, SEQ ID NO:75, and/or SEQ ID NO:77. The present invention also includes a CD154 nucleic acid molecule encoding a protein having at least a portion of SEQ ID NO:72, SEQ ID NO:75, and/or SEQ ID NO:77, as well as allelic variants of a CD154 nucleic acid molecule encoding a protein having these sequences, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In another embodiment, an IL-5 nucleic acid molecule of the present invention encodes a protein having an amino acid sequence that is at least about at least about 85%, at least about 85%, more preferably at least about 90%, and even more preferably at least about 95% identical to SEQ ID NO:80, SEQ ID NO:83, and/or SEQ ID NO:85. The present invention also includes an IL-5 nucleic acid molecule encoding a protein having at least a portion of SEQ ID NO:80, SEQ ID NO:83, and/or SEQ ID NO:85, as well as allelic variants of an IL-5 nucleic acid molecule encoding a protein having these sequences, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In another embodiment, an IL-13 nucleic acid molecule of the present invention encodes a protein having an amino acid sequence that is at least about at least about 70%, at

least about 75%, at least about 80%, preferably at least about 85%, more preferably at least about 90%, and even more preferably at least about 95% identical to SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:99, SEQ ID NO:102, and/or SEQ ID NO:104. The present invention also includes an IL-13 nucleic acid molecule encoding a protein having at least a portion of SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:94, SEQ ID NO:96, SEQ ID NO:99, SEQ ID NO:102, and/or SEQ ID NO:104, as well as allelic variants of an IL-13 nucleic acid molecule encoding a protein having these sequences, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In one embodiment, an IL-4 nucleic acid molecule of the present invention is at least about 90%, and preferably at least about 95% identical to nCaIL-4₅₄₉. Even more preferred is a nucleic acid molecule comprising nCaIL4₅₄₉, nCaIL-4₃₉₆, nCaIL-4₃₂₄, and/or an allelic variant of such a nucleic acid molecule. In another embodiment, a Flt-3 ligand nucleic acid molecule of the present invention is at least about 75%, more preferably at least about 80%, more preferably at least about 85%, more preferably at least about 90% and even more preferably at least about 95% identical to nCaFlt3L₁₀₁₃. Even more preferred is a nucleic acid molecule comprising nCaFlt3L₁₀₁₃, nCaFlt3L₈₈₂, nCaFlt3L₈₀₄, nCaFlt3L₈₂₈, nCaFlt3L₉₈₅, nCaFlt3L₁₀₁₉, nCaFlt3L₉₃, and/or nCaFlt3L₇₅₀, and/or an allelic variant of such a nucleic acid molecule. In one embodiment, a Flt-3 ligand nucleic acid molecule of the present invention is at least about 75%, more preferably at least about 80%, more preferably at least about 85%, more preferably at least about 90% and even more preferably at least about 95% identical to nFeFlt3L₉₄₂. Even more preferred is a nucleic acid molecule comprising nFeFlt3L₃₉₅, nFeFlt3L₇₉₃, nFeFlt3L₉₄₂, nFeFlt3L₈₇₃, and/or nFeFlt3L₇₉₅, and/or an allelic variant of such a nucleic acid molecule. In one embodiment, a CD40 nucleic acid molecule of the present invention is at least about 70%, at least about 75%, more preferably at least about 80%, more preferably at least about 85%, more preferably at least about 90% and even more preferably at least about 95% identical to nCaCD40₃₂₁, nCaCD40₁₄₂₅, nCaCD40₈₂₂, and/or nCaCD40₇₆₅, and/or an allelic variant of such a nucleic acid molecule. In one embodiment, a CD40 nucleic acid molecule of the present invention is at least about 70%, at least about 75%, more preferably at least about 80%, more preferably at least about 85%, more preferably at least about 90% and even more preferably at least about 95% identical to nFeCD40₃₃₆, and/or an allelic variant of such a nucleic acid molecule. In one embodiment, a CD154 nucleic acid molecule of the present invention is at least about 85%, preferably at least about 85%, more preferably at least about 90% and even more preferably at least about 95% identical to nCaCD154₃₉₀, nCaCD154₁₈₇₈, nCaCD154₇₈₀, CaCD154₆₃₃, and/or an allelic variant of such a nucleic acid molecule. In one embodiment, a CD154 nucleic acid molecule of the present invention is at least about 91%, and preferably about 95% identical to nFeCD154₈₈₅, nFeCD154₇₈₀, and/or nFeCD154₆₃₃, and/or an allelic variant of such a nucleic acid molecule. In one embodiment, an IL-5 molecule of the present invention is at least about 90% and preferably at least about 95% identical to nCaIL-5₆₁₀, nCaIL-5₄₀₂, and/or nCaIL-5₃₄₅, and/or an allelic variant of such a nucleic acid molecule. In another embodiment, an IL-13 molecule of the present invention is at least about

65%, at least about 70%, preferably at least about 75%, more preferably at least about 80%, more preferably at least about 85%, more preferably at least about 90% and even more preferably at least about 95% identical to nCaIL-13₁₆₆, nCaIL-13₂₇₂, nCaIL-13₂₇₈, nCaIL-13₁₃₀₂, nCaIL13₃₉₃, nCaIL-13₃₃₃, nCaIL13₁₂₆₉, nCaIL-13₃₉₀, and/or nCaIL13₃₃₀, and/or an allelic variant of such a nucleic acid molecule.

In another embodiment, an IL-4 nucleic acid molecule of the present invention comprises a nucleic acid sequence that is at least about 90%, and preferably at least about 95% identical to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:19, and/or SEQ ID NO:21. The present invention also includes an IL-4 nucleic acid molecule comprising at least a portion of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:19, and/or SEQ ID NO:21, as well as allelic variants of such IL4 nucleic acid molecules, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In another embodiment, a Flt-3 ligand nucleic acid molecule of the present invention comprises a nucleic acid sequence that is at least about 75%, preferably at least about 80%, more preferably at least about 85%, more preferably at least about 90% and even more preferably at least about 95% identical to SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:36, and/or SEQ ID NO:37. The present invention also includes a Flt-3 ligand- nucleic acid molecule comprising at least a portion of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:35, SEQ ID NO:36, and/or SEQ ID NO:37, as well as allelic variants of such Flt-3 ligand nucleic acid molecules, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In one embodiment, a Flt-3 ligand nucleic acid molecule of the present invention comprises a nucleic acid sequence that is at least about 75%, more preferably at least about 80%, more preferably at least about 85%, more preferably at least about 90% and even more preferably at least about 95% identical to SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, and/or SEQ ID NO:50. The present invention also includes a Flt-3 ligand- nucleic acid molecule comprising at least a portion of SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, and/or SEQ ID NO:50, as well as allelic variants of such Flt-3 ligand nucleic acid molecules, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In one embodiment, a CD40 nucleic acid molecule of the present invention comprises a nucleic acid sequence that is at least about 70%, at least about 75%, more preferably at least about 80%, more preferably at least about 85%, more preferably at least about 90% and even more preferably at least about 95% identical to SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, and/or SEQ ID NO:59. The present invention also includes a CD40 nucleic acid molecule comprising at least

a portion of SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, and/or SEQ ID NO:59, as well as allelic variants of such CD40 nucleic acid molecules, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In one embodiment, a CD40 nucleic acid molecule of the present invention comprises a nucleic acid sequence that is at least about 70%, at least about 75%, more preferably at least about 80%, more preferably at least about 85%, more preferably at least about 90% and even more preferably at least about 95% identical to SEQ ID NO:60 and/or SEQ ID NO:62. The present invention also includes a CD40 nucleic acid molecule comprising at least a portion of SEQ ID NO:60 and/or SEQ ID NO:62, as well as allelic variants of such CD40 nucleic acid molecules, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In one embodiment, a CD154 nucleic acid molecule of the present invention comprises a nucleic acid sequence that is at least about 85%, preferably at least about 85%, more preferably at least about 90% and even more preferably at least about 95% identical to SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, and/or SEQ ID NO:71. The present invention also includes a CD154 nucleic acid molecule comprising at least a portion of SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, and/or SEQ ID NO:71, as well as allelic variants of such CD154 nucleic acid molecules, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In one embodiment, a CD154 nucleic acid molecule of the present invention comprises a nucleic acid sequence that is at least about 91%, and preferably about 95% identical to SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, and/or SEQ ID NO:79. The present invention also includes a CD154 nucleic acid molecule comprising at least a portion of SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, and/or SEQ ID NO:79, as well as allelic variants of such CD154 nucleic acid molecules, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In one embodiment, an IL-5 nucleic acid molecule of the present invention comprises a nucleic acid sequence that is at least about 90% and preferably at least about 95% identical to SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, and/or SEQ ID NO:87. The present invention also includes an IL-5 nucleic acid molecule comprising at least a portion of SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, and/or SEQ ID NO:87, as well as allelic variants of such IL-5 nucleic acid molecules, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In one embodiment, an IL-13 nucleic acid molecule of the present invention comprises a nucleic acid sequence that is at least about 65%, at least about 70%, preferably at least about 75%, more preferably at least about 80%, more preferably at least about 85%, more preferably at least about 90% and even more preferably at least about 95% identical

to SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, and/or SEQ ID NO:106. The present invention also includes an IL-13 nucleic acid molecule comprising at least a portion of SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, and/or SEQ ID NO:106, as well as allelic variants of such IL-13 nucleic acid molecules, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In one embodiment, an IFN α nucleic acid molecule of the present invention is identical to SEQ ID NO:107, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:116, and/or SEQ ID NO:118.

In another embodiment, a GM-CSF nucleic acid molecule of the present invention is identical to SEQ ID NO:119, SEQ ID NO:121, SEQ ID NO:122, SEQ ID NO:123, SEQ ID NO:124, and/or SEQ ID NO:126.

Knowing the nucleic acid sequences of certain immunoregulatory nucleic acid molecules of the present invention allows one skilled in the art to, for example, (a) make copies of those nucleic acid molecules, (b) obtain nucleic acid molecules including at least a portion of such nucleic acid molecules (e.g., nucleic acid molecules including full-length genes, full-length coding regions, regulatory control sequences, truncated coding regions), and/or (c) obtain other immunoregulatory nucleic acid molecules. Such nucleic acid molecules can be obtained in a variety of ways including screening appropriate expression libraries with antibodies of the present invention; traditional cloning techniques using oligonucleotide probes of the present invention to screen appropriate libraries; and PCR amplification of appropriate libraries or DNA using oligonucleotide primers of the present invention. Preferred libraries to screen or from which to amplify nucleic acid molecules include mammalian cDNA libraries as well as genomic DNA libraries. Similarly, preferred DNA sources from which to amplify nucleic acid molecules include mammalian cDNA and genomic DNA. Techniques to clone and amplify genes are disclosed, for example, in Sambrook et al., *ibid*.

The present invention also includes nucleic acid molecules that are oligonucleotides capable of hybridizing, under stringent hybridization conditions, with complementary regions of other, preferably longer, nucleic acid molecules of the present invention such as those comprising canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF nucleic acid molecules. Oligonucleotides of the present invention can be RNA, DNA, or derivatives of either. The minimum size of such oligonucleotides in the size required for formation of a stable hybrid between an oligonucleotide and a complementary sequence on a nucleic acid molecule of the present invention. A preferred oligonucleotide of the present invention has a maximum size of about 100 nucleotides. The present invention includes oligonucleotides that can be used as, for example, probes to identify nucleic acid molecules, primers to produce nucleic acid molecules, or therapeutic reagents to inhibit canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine

interleukin-13, feline interferon alpha, or feline GM-CSF protein production or activity (e.g., as antisense-, triplex formation-, ribozyme- and/or RNA drug-based reagents). The present invention also includes the use of such oligonucleotides to protect animals from disease using one or more of such technologies. Appropriate oligonucleotide-containing therapeutic compositions can be administered to an animal using techniques known to those skilled in the art.

One embodiment of the present invention includes a recombinant vector, which includes at least one isolated nucleic acid molecule of the present invention, inserted into any vector capable of delivering the nucleic acid molecule into a host cell. Such a vector contains heterologous nucleic acid sequences, that is nucleic acid sequences that are not naturally found adjacent to nucleic acid molecules of the present invention and that preferably are derived from a species other than the species from which the nucleic acid molecule(s) are derived. The vector can be either RNA or DNA, either prokaryotic or eukaryotic, and typically is a virus or a plasmid. Recombinant vectors can be used in the cloning, sequencing, and/or otherwise manipulating immunoregulatory nucleic acid molecules of the present invention.

One type of recombinant vector, referred to herein as a recombinant molecule, comprises a nucleic acid molecule of the present invention operatively linked to an expression vector. The phrase operatively linked refers to insertion of a nucleic acid molecule into an expression vector in a manner such that the molecule is able to be expressed when transformed into a host cell. As used herein, an expression vector is a DNA or RNA vector that is capable of transforming a host cell and of effecting expression of a specific nucleic acid molecule. Preferably, the expression vector is also capable of replicating within the host cell. Expression vectors can be either prokaryotic or eukaryotic, and are typically viruses or plasmids. Expression vectors of the present invention include any vectors that function (i.e., direct gene expression) in recombinant cells of the present invention, including in bacterial, fungal, parasite, insect, other animal, and plant cells. Preferred expression vectors of the present invention can direct gene expression in bacterial, yeast, insect and mammalian cells, and more preferably in the cell types disclosed herein, more preferably in vivo.

In particular, expression vectors of the present invention contain regulatory sequences such as transcription control sequences, translation control sequences, origins of replication, and other regulatory sequences that are compatible with the recombinant cell and that control the expression of nucleic acid molecules of the present invention. In particular, recombinant molecules of the present invention include transcription control sequences. Transcription control sequences are sequences which control the initiation, elongation, and termination of transcription. Particularly important transcription control sequences are those which control transcription initiation, such as promoter, enhancer, operator and repressor sequences. Suitable transcription control sequences include any transcription control sequence that can function in at least one of the recombinant cells of the present invention. A variety of such transcription control sequences are known to those skilled in the art. Preferred transcription control sequences include those which function in bacterial, yeast, helminth and/or other endoparasite, insect and mammalian cells, such as, but not limited to, *tac*, *lac*, *trp*, *trc*, *oxy-pro*, *omp/lpp*, *rrnB*, bacteriophage lambda (such as lambda *p_L* and lambda *p_R* and fusions that include such promoters), bacteriophage T7, T7lac, bacteriophage T3, bacteriophage SP6, bacteriophage

SPO1, metallothionein, alpha-mating factor, Pichia alcohol oxidase, alphavirus subgenomic promoter, antibiotic resistance gene, baculovirus. Heliothis zea insect virus, vaccinia virus, herpesvirus, raccoon poxvirus, other poxvirus, adenovirus, cytomegalovirus (such as immediate early promoter), simian virus 40, retrovirus, actin, retroviral long terminal repeat. Rous sarcoma virus, heat shock, phosphate and nitrate transcription control sequences as well as other sequences capable of controlling gene expression in prokaryotic or eukaryotic cells. Additional suitable transcription control sequences include tissue-specific promoters and enhancers as well as lymphokine-inducible promoters (e.g., promoters inducible by interferons or interleukins). Transcription control sequences of the present invention can also include naturally occurring transcription control sequences naturally associated with mammals, such as dog, cat, horse or human transcription control sequences.

Suitable and preferred nucleic acid molecules to include in recombinant vectors of the present invention are as disclosed herein. Preferred nucleic acid molecules to include in recombinant vectors, and particularly in recombinant molecules, include nCaIL-4₅₄₉, nCaIL-4₃₉₆, nCaIL-4₃₂₄, nCaFlt3L₁₀₁₃, nCaFlt3L₈₈₂, nCaFlt3L₈₀₄, nCaFlt3L₈₂₈, nCaFlt3L₉₈₅, nCaFlt3L₁₀₁₉, nCaFlt3L₉₃, nCaFlt3L₇₅₀, nFeFlt3L₃₉₅, nFeFlt3L₇₉₃, nFeFlt3L₉₄₂, nFeFlt3L₈₇₃, nFeFlt3L₇₉₅, nCaCD40₃₂₁, nCaCD40₁₄₂₅, nCaCD40₈₂₂, nCaCD40₇₆₅, nFeCD40₃₃₆, nCaCD154₃₉₀, nCaCD154₈₇₈, nCaCD154₇₈₀, nCaCD154₆₃₃, nFeCD154₈₈₅, nFeCD154₇₈₀, nFeCD154₆₃₃, nCaIL-5₆₁₀, nCaIL-5₄₀₂, nCaIL-5₃₄₅, nCaIL-13₁₆₆, nCaIL-13₂₇₂, nCaIL-13₂₇₈, nCaIL-13₁₃₀₂, nCaIL-13₃₉₃, nCaIL-13₃₃₃, nCaIL-13₁₂₆₉, nCaIL-13₃₉₀, nCaIL-13₃₃₀, nFeIFN α _{567a}, nFeIFN α _{567b}, nFeIFN α _{498a}, nFeIFN α _{498b}, nFeGMCSF₄₄₄, nFeGMCSF₄₃₂, and/or nFeGMCSF₃₈₁.

Recombinant molecules of the present invention may also (a) contain secretory signals (i.e., signal segment nucleic acid sequences) to enable an expressed parasitic helminth protein of the present invention to be secreted from the cell that produces the protein and/or (b) contain fusion sequences which lead to the expression of nucleic acid molecules of the present invention as fusion proteins. Examples of suitable signal segments include any signal segment capable of directing the secretion of a protein of the present invention. Preferred signal segments include, but are not limited to, tissue plasminogen activator (t-PA), interferon, interleukin, growth hormone, histocompatibility and viral envelope glycoprotein signal segments. Suitable fusion segments encoded by fusion segment nucleic acids are disclosed herein. In addition, a nucleic acid molecule of the present invention can be joined to a fusion segment that directs the encoded protein to the proteosome, such as a ubiquitin fusion segment. Eukaryotic recombinant molecules may also include intervening and/or untranslated sequences surrounding and/or within the nucleic acid sequences of nucleic acid molecules of the present invention.

Another embodiment of the present invention includes a recombinant cell comprising a host cell transformed with one or more recombinant molecules of the present invention. Transformation of a nucleic acid molecule into a cell can be accomplished by any method by which a nucleic acid molecule can be inserted into the cell. Transformation techniques include, but are not limited to, transfection, electroporation, microinjection, lipofection, adsorption, and protoplast fusion. A recombinant cell may remain unicellular or may grow into a tissue, organ or a multicellular organism. Transformed nucleic acid molecules of the present invention can remain extrachromosomal or can integrate into one or

more sites within a chromosome of the transformed (i.e., recombinant) cell in such a manner that their ability to be expressed is retained. Preferred nucleic acid molecules with which to transform a cell include immunoregulatory nucleic acid molecules of the present invention disclosed herein. Particularly preferred nucleic acid molecules with which to transform a cell include nCaIL-4₅₄₉, nCaIL-4₃₉₆, nCaIL-4₃₂₄, nCaFlt3L₁₀₁₃, nCaFlt3L₈₈₂, nCaFlt3L₈₀₄, nCaFlt3L₈₂₈, nCaFlt3L₉₈₅, nCaFlt3L₁₀₁₉, nCaFlt3L₉₃, nCaFlt3L₇₅₀, nFeFlt3L₃₉₅, nFeFlt3L₇₉₃, nFeFlt3L₉₄₂, nFeFlt3L₈₇₃, nFeFlt3L₇₉₅, nCaCD40₃₂₁, nCaCD40₁₄₂₅, nCaCD40₈₂₂, nCaCD40₇₆₅, nFeCD40₃₃₆, nCaCD154₃₉₀, nCaCD154₁₈₇₈, nCaCD154₇₈₀, nCaCD154₆₃₃, nFeCD154₈₈₅, nFeCD154₇₈₀, nFeCD154₆₃₃, nCaIL-5₆₁₀, nCaIL-5₄₀₂, nCaIL-5₃₄₅, nCaIL-13₁₆₆, nCaIL-13₂₇₂, nCaIL-13₂₇₈, nCaIL13₁₃₀₂, nCaIL-13₃₉₃, nCaIL-13₃₃₃, nCaIL13₁₂₆₉, nCaIL-13₃₉₀, nCaIL-13₃₃₀, nFeIFN α _{567a}, nFeIFN α _{567b}, nFeIFN α _{498a}, nFeIFN α _{498b}, nFeGMCSF₄₄₄, nFeGMCSF₄₃₂, and/or nFeGMCSF₃₈₁.

Suitable host cells to transform include any cell that can be transformed with a nucleic acid molecule of the present invention. Host cells can be either untransformed cells or cells that are already transformed with at least one nucleic acid molecule (e.g., nucleic acid molecules encoding one or more proteins of the present invention and/or other proteins useful in the production of multivalent vaccines). Host cells of the present invention either can be endogenously (i.e., naturally) capable of producing immunoregulatory proteins of the present invention or can be capable of producing such proteins after being transformed with at least one nucleic acid molecule of the present invention. Host cells of the present invention can be any cell capable of producing at least one protein of the present invention, and include bacterial, fungal (including yeast), parasite (including helminth, protozoa and ectoparasite), other insect, other animal and plant cells. Preferred host cells include bacterial, mycobacterial, yeast, helminth, insect and mammalian cells. More preferred host cells include *Salmonella*, *Escherichia*, *Bacillus*, *Listeria*, *Saccharomyces*, *Spodoptera*, *Mycobacteria*, *Trichoplusia*, BHK (baby hamster kidney) cells, MDCK cells (Madin-Darby canine kidney cell line), CRFK cells (Crandell feline kidney cell line), CV-1 cells (African monkey kidney cell line used, for example, to culture racoon poxvirus), COS (e.g., COS-7) cells, chinese hamster ovary (CHO) cells, Ltk cells and Vero cells. Particularly preferred host cells are *Escherichia coli*, including *E. coli* K-12 derivatives; *Salmonella typhi*; *Salmonella typhimurium*, including attenuated strains such as UK-1₀₃₉₈₇ and SR-11₀₄₀₇₂; *Spodoptera frugiperda*; *Trichoplusia ni*; BHK cells; MDCK cells; CRFK cells; CV-1 cells; COS cells; Vero cells; and non-tumorigenic mouse myoblast G8 cells (e.g., ATCC CRL 1246). Additional appropriate mammalian cell hosts include other kidney cell lines, other fibroblast cell lines (e.g., human, murine or chicken embryo fibroblast cell lines), myeloma cell lines. Chinese hamster ovary cells, mouse NIH/3T3 cells, LMTK³¹ cells and/or HeLa cells. In one embodiment, the proteins may be expressed as heterologous proteins in myeloma cell lines employing immunoglobulin promoters.

A recombinant cell is preferably produced by transforming a host cell with one or more recombinant molecules, each comprising one or more nucleic acid molecules of the present invention operatively linked to an expression vector containing one or more transcription control sequences, examples of which are disclosed herein.

A recombinant cell of the present invention includes any cell transformed with at least one of any nucleic acid

molecule of the present invention. Suitable and preferred nucleic acid molecules as well as suitable and preferred recombinant molecules with which to transfer cells are disclosed herein.

Recombinant cells of the present invention can also be co-transformed with one or more recombinant molecules including any of canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF nucleic acid molecule encoding one or more proteins of the present invention and/or one or more other nucleic acid molecules encoding other therapeutic compounds, as disclosed herein (e.g., to produce multivalent vaccines).

Recombinant DNA technologies can be used to improve expression of transformed nucleic acid molecules by manipulating, for example, the number of copies of the nucleic acid molecules within a host cell, the efficiency with which those nucleic acid molecules are transcribed, the efficiency with which the resultant transcripts are translated, and the efficiency of post-translational modifications. Recombinant techniques useful for increasing the expression of nucleic acid molecules of the present invention include, but are not limited to, operatively linking nucleic acid molecules to high-copy number plasmids, integration of the nucleic acid molecules into one or more host cell chromosomes, addition of vector stability sequences to plasmids, substitutions or modifications of transcription control signals (e.g., promoters, operators, enhancers), substitutions or modifications of translational control signals (e.g., ribosome binding sites, Shine-Dalgarno sequences), modification of nucleic acid molecules of the present invention to correspond to the codon usage of the host cell, deletion of sequences that destabilize transcripts, and use of control signals that temporally separate recombinant cell growth from recombinant enzyme production during fermentation. The activity of an expressed recombinant protein of the present invention may be improved by fragmenting, modifying, or derivatizing nucleic acid molecules encoding such a protein.

Isolated immunoregulatory proteins of the present invention can be produced in a variety of ways, including production and/or recovery of natural proteins, production and/or recovery of recombinant proteins, and/or chemical synthesis of the proteins. In one embodiment, an isolated protein of the present invention is produced by culturing a cell capable of expressing the protein under conditions effective to produce the protein, and recovering the protein. A preferred cell to culture is a recombinant cell of the present invention. Effective culture conditions include, but are not limited to, effective media, bioreactor, temperature, pH and oxygen conditions that permit protein production. An effective medium refers to any medium in which a cell is cultured to produce an immunoregulatory protein of the present invention. Such medium typically comprises an aqueous medium having assimilable carbon, nitrogen and phosphate sources, and appropriate salts, minerals, metals and other nutrients, such as vitamins. Cells of the present invention can be cultured in conventional fermentation bioreactors, shake flasks, test tubes, microtiter dishes, and petri plates. Culturing can be carried out at a temperature, pH and oxygen content appropriate for a recombinant cell. Such culturing conditions are within the expertise of one of ordinary skill in the art.

Depending on the vector and host system used for production, resultant proteins of the present invention may either remain within the recombinant cell; be secreted into

the fermentation medium; be secreted into a space between two cellular membranes, such as the periplasmic space in *E. coli*; or be retained on the outer surface of a cell or viral membrane.

The phrase "recovering the protein", as well as similar phrases, refers to collecting the whole fermentation medium containing the protein and need not imply additional steps of separation or purification. Proteins of the present invention can be purified using a variety of standard protein purification techniques, such as, but not limited to, affinity chromatography, ion exchange chromatography, filtration, electrophoresis, hydrophobic interaction chromatography, gel filtration chromatography, reverse phase chromatography, concanavalin A chromatography, chromatofocusing and/or differential solubilization. Proteins of the present invention are preferably retrieved in "substantially pure" form. As used herein, "substantially pure" refers to a purity that allows for the effective use of the protein as a therapeutic composition or diagnostic. A therapeutic composition for animals, for example, should exhibit no substantial toxicity and preferably should be capable of stimulating the production of antibodies in a treated animal.

The present invention also includes isolated (i.e., removed from their natural milieu) antibodies that selectively bind to an immunoregulatory protein of the present invention and/or a mimotope thereof (e.g., anti-IL-4 antibodies, anti-Flt-3 ligand antibodies, anti-CD40 antibodies, anti-CD154 antibodies, anti-IL-5 antibodies, anti-IL-13 antibodies, anti-IFN α antibodies, and/or anti-GM-CSF antibodies). As used herein, the term "selectively binds to" an immunoregulatory protein of the present invention, refers to the ability of antibodies of the present invention to preferentially bind to specified proteins and/or mimetopes thereof of the present invention. Binding can be measured using a variety of methods standard in the art including enzyme immunoassays (e.g., ELISA), immunoblot assays, etc.; see, for example, Sambrook et al., *ibid.*, and Harlow, et al., 1988, *Antibodies*, a Laboratory Manual, Cold Spring Harbor Labs Press; Harlow et al., *ibid.*, is incorporated by this reference herein in its entirety. An anti-IL-4 antibody of the present invention preferably selectively binds to an IL-4 protein in such a way as to inhibit the function of that protein. An anti-Flt-3 ligand antibody of the present invention preferably selectively binds to a Flt-3 ligand- protein in such a way as to inhibit the function of that protein. An anti-CD40 antibody of the present invention preferably selectively binds to a CD40 protein in such a way as to inhibit the function of that protein. An anti-CD154 antibody of the present invention preferably selectively binds to a CD154 protein in such a way as to inhibit the function of that protein. An anti-IL-5 antibody of the present invention preferably selectively binds to an IL-5 protein in such a way as to inhibit the function of that protein. An anti-IL-13 antibody of the present invention preferably selectively binds to an IL-13 protein in such a way as to inhibit the function of that protein. An anti-IFN α antibody of the present invention preferably selectively binds to an IFN α protein in such a way as to inhibit the function of that protein. An anti-GM-CSF antibody of the present invention preferably selectively binds to a GM-CSF protein in such a way as to inhibit the function of that protein.

Isolated antibodies of the present invention can include antibodies in serum, or antibodies that have been purified to varying degrees. Antibodies of the present invention can be polyclonal or monoclonal, or can be functional equivalents such as antibody fragments and/or genetically-engineered antibodies, including single chain antibodies or chimeric antibodies that can bind to one or more epitopes.

A preferred method to produce antibodies of the present invention includes (a) administering to an animal an effective amount of a protein, peptide and/or mimotope thereof of the present invention to produce the antibodies and (b) recovering the antibodies. In another method, antibodies of the present invention are produced recombinantly using techniques as heretofore disclosed to produce any of the immunoregulatory proteins of the present invention. Antibodies raised against defined proteins or mimetopes can be advantageous because such antibodies are not substantially contaminated with antibodies against other substances that might otherwise cause interference in a diagnostic assay or side effects if used in a therapeutic composition.

Antibodies of the present invention have a variety of potential uses that are within the scope of the present invention. For example, such antibodies can be used (a) as reagents in assays to detect an immunoregulatory protein of the present invention, (b) as reagents in assays to modulate cellular activity through an immunoregulatory protein of the present invention (e.g., mimicking ligand binding to a canine interleukin-4, canine or feline Flt-3 ligand, canine or feline CD40, canine or feline CD154, canine interleukin-5, canine interleukin-13, feline interferon alpha, or feline GM-CSF protein, as appropriate), and/or (c) as tools to screen expression libraries and/or to recover desired proteins of the present invention from a mixture of proteins and other contaminants. Furthermore, antibodies of the present invention can be used to target compounds (e.g., nucleic acid molecules, drugs or proteins) to antigen presenting cells. Targeting can be accomplished by conjugating (i.e., stably joining) such antibodies to the compounds using techniques known to those skilled in the art. Suitable compounds are known to those skilled in the art.

One embodiment of the present invention is a therapeutic composition that, when administered to an animal in an effective manner, is capable of regulating an immune response in an animal. Therapeutic compositions of the present invention can include at least one of the following therapeutic compounds: an isolated IL-4, Flt-3 ligand, CD40, CD154, IL-5, IL-13, IFN α , and/or GM-CSF protein of the present invention and/or a mimotope thereof; an isolated IL-4, Flt-3 ligand, CD40, CD154, IL-5, IL-13, IFN α , and/or GM-CSF nucleic acid molecule of the present invention; an isolated antibody that selectively binds to an IL-4, Flt-3 ligand, CD40, CD154, IL-5, IL-13, IFN α , and/or GM-CSF protein of the present invention; an inhibitor of canine IL-4, Flt-3 ligand, CD40, CD154, IL-5, IL-13, IFN α , and/or GM-CSF function identified by its ability to bind to an IL-4, Flt-3 ligand, CD40, CD154, IL-5, IL-13, IFN α , and/or GM-CSF protein, respectively, of the present invention; such an inhibitor can inhibit binding of the respective immunoregulatory protein with its respective receptor, or inhibit the activity the respective protein. Methods to perform such assays to measure binding and/or activity of an immunoregulatory protein of the present invention are known to those of skill in the art, and are described, for example, in Janeway et al., *ibid.* As used herein, a therapeutic compound refers to a compound that, when administered to an animal in an effective manner, is able to treat, ameliorate, and/or prevent a disease. Examples of proteins, nucleic acid molecules, antibodies and/or inhibitors of the present invention are disclosed herein.

The present invention also includes a therapeutic composition comprising at least one IL-4-, Flt-3 ligand-, CD40-, CD154-, IL-5-, IL-13-, IFN α -, and/or GM-CSF-based compound of the present invention in combination with at least one additional therapeutic compound. Examples of such compounds are disclosed herein.

Therapeutic compositions of the present invention can be administered to any animal susceptible to such therapy, preferably to mammals, and more preferably to dogs, cats, humans, ferrets, horses, cattle, sheep and/or other pets, economic food animals and/or zoo animals. Preferred animals include dogs, cats, horses and/or humans.

A therapeutic composition of the present invention is administered to an animal in an effective manner such that the composition is capable of regulating an immune response in that animal. Therapeutic compositions of the present invention can be administered to animals prior to onset of a disease (i.e., as a preventative vaccine) and/or can be administered to animals after onset of a disease in order to treat the disease (i.e., as a therapeutic vaccine). Preferred diseases to prevent and/or treat include autoimmune diseases, allergic reactions, infectious diseases, tumor development, inflammatory diseases and/or graft rejection. In one embodiment, a therapeutic composition of the present invention is administered with an antigen to enhance an immune response against that antigen.

Therapeutic compositions of the present invention can be formulated in an excipient that the animal to be treated can tolerate. Examples of such excipients include water, saline, Ringer's solution, dextrose solution, Hank's solution, and/or other aqueous physiologically balanced salt solutions. Non-aqueous vehicles, such as fixed oils, sesame oil, ethyl oleate, or triglycerides may also be used. Other useful formulations include suspensions containing viscosity enhancing agents, such as sodium carboxymethylcellulose, sorbitol, or dextran. Excipients can also contain minor amounts of additives, such as substances that enhance isotonicity and chemical stability. Examples of buffers include phosphate buffer, bicarbonate buffer and/or Tris buffer, while examples of preservatives include thimerosal, o-cresol, formalin and/or benzyl alcohol. Standard formulations can either be liquid injectables or solids which can be taken up in a suitable liquid as a suspension or solution for injection. Thus, in a non-liquid formulation, the excipient can comprise dextrose, human serum albumin, preservatives, etc., to which sterile water or saline can be added prior to administration.

In one embodiment of the present invention, a therapeutic composition can include an adjuvant. Adjuvants are agents that are capable of enhancing the immune response of an animal to a specific antigen. Suitable adjuvants include, but are not limited to, cytokines, chemokines, and/or compounds that induce the production of cytokines and/or chemokines (e.g., granulocyte macrophage colony stimulating factor (GM-CSF), granulocyte colony stimulating factor (G-CSF), macrophage colony stimulating factor (M-CSF), colony stimulating factor (CSF), erythropoietin (EPO), interleukin 2 (IL-2), interleukin-3 (IL-3), interleukin 5 (IL-5), interleukin 6 (IL-6), interleukin 7 (IL-7), interleukin 8 (IL-8), interleukin 10 (IL-10), interleukin 12 (IL-12), interferon gamma, interferon gamma inducing factor I (IGIF), transforming growth factor beta, RANTES (regulated upon activation, normal T cell expressed and presumably secreted), macrophage inflammatory proteins (e.g., MIP-1 alpha and MIP-1 beta), and Leishmania elongation initiating factor (LEIF)); bacterial components (e.g., endotoxins, in particular superantigens, exotoxins and cell wall components); aluminum-based salts; calcium-based salts; silica; polynucleotides; toxoids; serum proteins, viral coat proteins; block copolymer adjuvants (e.g., Hunter's Titermax™ adjuvant (Vaxcel™, Inc. Norcross, Ga.), Ribi adjuvants (Ribi ImmunoChem Research, Inc., Hamilton, Mont.); and saponins and their derivatives (e.g., Quil A (Superfos Biosector A/S, Denmark). Protein adjuvants of the present

invention can be delivered in the form of the protein themselves or of nucleic acid molecules encoding such proteins using the methods described herein.

In one embodiment of the present invention, a therapeutic composition can include a carrier. Carriers include compounds that increase the half-life of a therapeutic composition in the treated animal. Suitable carriers include, but are not limited to, polymeric controlled release vehicles, biodegradable implants, liposomes, bacteria, viruses, other cells, oils, esters, and glycols.

One embodiment of the present invention is a controlled release formulation that is capable of slowly releasing a composition of the present invention into an animal. As used herein, a controlled release formulation comprises a composition of the present invention in a controlled release vehicle. Suitable controlled release vehicles include, but are not limited to, biocompatible polymers, other polymeric matrices, capsules, microcapsules, microparticles, bolus preparations, osmotic pumps, diffusion devices, liposomes, lipospheres, and transdermal delivery systems. Other controlled release formulations of the present invention include liquids that, upon administration to an animal, form a solid or a gel in situ. Preferred controlled release formulations are biodegradable (i.e., bioerodible).

A preferred controlled release formulation of the present invention is capable of releasing a composition of the present invention into the blood of the treated animal at a constant rate sufficient to attain therapeutic dose levels of the composition to regulate an immune response in an animal. The therapeutic composition is preferably released over a period of time ranging from about 1 to about 12 months. A controlled release formulation of the present invention is capable of effecting a treatment preferably for at least about 1 month, more preferably for at least about 3 months, even more preferably for at least about 6 months, even more preferably for at least about 9 months, and even more preferably for at least about 12 months.

Therapeutic compositions of the present invention can be administered to animals prior to and/or after onset of disease. Acceptable protocols to administer therapeutic compositions in an effective manner include individual dose size, number of doses, frequency of dose administration, and/or mode of administration. Determination of such protocols can be accomplished by those skilled in the art. A suitable single dose is a dose that is capable of regulating the immune response in an animal when administered one or more times over a suitable time period. For example, a preferred single dose of a protein, mimotope or antibody therapeutic composition is from about 1 microgram (μg) to about 10 milligrams (mg) of the therapeutic composition per kilogram body weight of the animal. Booster vaccinations can be administered from about 2 weeks to several years after the original administration. Booster administrations preferably are administered when the immune response of the animal becomes insufficient to protect the animal from disease. A preferred administration schedule is one in which from about 10 μg to about 1 mg of the therapeutic composition per kg body weight of the animal is administered from about one to about two times over a time period of from about 2 weeks to about 12 months. Modes of administration can include, but are not limited to, subcutaneous, intradermal, intravenous, intranasal, intraocular, oral, transdermal and/or intramuscular routes.

According to one embodiment, a nucleic acid molecule of the present invention can be administered to an animal in a fashion to enable expression of that nucleic acid molecule into a therapeutic protein or therapeutic RNA (e.g., antisense

RNA, ribozyme, triple helix forms or RNA drug) in the animal. Nucleic acid molecules can be delivered to an animal in a variety of methods including, but not limited to, (a) administering a naked (i.e., not packaged in a viral coat or cellular membrane) nucleic acid as a genetic vaccine (e.g., as naked DNA or RNA molecules, such as is taught, for example in Wolff et al., 1990, Science 247, 1465-1468) or (b) administering a nucleic acid molecule packaged as a recombinant virus vaccine or as a recombinant cell vaccine (i.e., the nucleic acid molecule is delivered by a viral or cellular vehicle).

A genetic (i.e., naked nucleic acid) vaccine of the present invention includes a nucleic acid molecule of the present invention and preferably includes a recombinant molecule of the present invention that preferably is replication, or otherwise amplification, competent. A genetic vaccine of the present invention can comprise one or more nucleic acid molecules of the present invention in the form of, for example, a dicistronic recombinant molecule. Preferred genetic vaccines include at least a portion of a viral genome (i.e., a viral vector). Preferred viral vectors include those based on alphaviruses, poxviruses, adenoviruses, herpesviruses, picomaviruses, and/or retroviruses, with those based on alphaviruses (such as Sindbis or Semliki forest virus), species-specific herpesviruses and/or poxviruses being particularly preferred. Any suitable transcription control sequence can be used, including those disclosed as suitable for protein production. Particularly preferred transcription control sequences include cytomegalovirus immediate early (preferably in conjunction with Intron-A), Rous sarcoma virus long terminal repeat, and tissue-specific transcription control sequences, as well as transcription control sequences endogenous to viral vectors if viral vectors are used. The incorporation of a "strong" polyadenylation signal is also preferred.

Genetic vaccines of the present invention can be administered in a variety of ways, with intramuscular, subcutaneous, intradermal, transdermal, intranasal and/or oral routes of administration being preferred. A preferred single dose of a genetic vaccine ranges from about 1 nanogram (ng) to about 600 µg, depending on the route of administration and/or method of delivery, as can be determined by those skilled in the art. Suitable delivery methods include, for example, by injection, as drops, aerosolized and/or topically. Genetic vaccines of the present invention can be contained in an aqueous excipient (e.g., phosphate buffered saline) alone or in a carrier (e.g., lipid-based vehicles).

A recombinant virus vaccine of the present invention includes a recombinant molecule of the present invention that is packaged in a viral coat and that can be expressed in an animal after administration. Preferably, the recombinant molecule is packaging- or replication-deficient and/or encodes an attenuated virus. A number of recombinant viruses can be used, including, but not limited to, those based on alphaviruses, poxviruses, adenoviruses, herpesviruses, picomaviruses, and/or retroviruses. Preferred recombinant virus vaccines are those based on alphaviruses (such as Sindbis virus), raccoon poxviruses, species-specific herpesviruses and/or species-specific poxviruses. An example of methods to produce and use alphavirus recombinant virus vaccines are disclosed in U.S. Pat. No. 5,766,602 by Xiong et al., issued Jun. 16, 1998, which is incorporated by this reference herein in its entirety.

When administered to an animal, a recombinant virus vaccine of the present invention infects cells within the immunized animal and directs the production of a therapeutic

protein or RNA nucleic acid molecule that is capable of protecting the animal from disease caused by a parasitic helminth as disclosed herein. For example, a recombinant virus vaccine comprising an immunoregulatory nucleic acid molecule of the present invention is administered according to a protocol that results in the regulation of an immune response in an animal. A preferred single dose of a recombinant virus vaccine of the present invention is from about 1×10^4 to about 1×10^8 virus plaque forming units (pfu) per kilogram body weight of the animal. Administration protocols are similar to those described herein for protein-based vaccines, with subcutaneous, intramuscular, intranasal, intraocular and/or oral administration routes being preferred.

A recombinant cell vaccine of the present invention includes recombinant cells of the present invention that express at least one protein of the present invention. Preferred recombinant cells for this embodiment include *Salmonella*, *E. coli*, *Listeria*, *Mycobacterium*, *S. frugiperda*, yeast, (including *Saccharomyces cerevisiae* and *Pichia pastoris*), BHK, CV-1, myoblast G8, COS (e.g., COS-7), Vero, MDCK and CRFK recombinant cells. Recombinant cell vaccines of the present invention can be administered in a variety of ways but have the advantage that they can be administered orally, preferably at doses ranging from about 10^8 to about 10^{12} cells per kilogram body weight. Administration protocols are similar to those described herein for protein-based vaccines. Recombinant cell vaccines can comprise whole cells, cells stripped of cell walls or cell lysates.

The efficacy of a therapeutic composition of the present invention to regulate the immune response in an animal can be tested in a variety of ways including, but not limited to, detection of cellular immunity within the treated animal, determining lymphocyte or dendritic cell activity, detection of immunoglobulin levels, determining hematopoietic stem cell or hematopoietic early progenitor cell development, determining dendritic cell development or challenge of the treated animal with an infectious agent to determine whether the treated animal is resistant to disease. In one embodiment, therapeutic compositions can be tested in animal models such as mice. Such techniques are known to those skilled in the art.

One embodiment of the present invention is an inhibitory compound. Preferably, such an inhibitory compound is derived from an IL-4, Flt-3 ligand, CD40, CD154, IL-5, IL-13, IFN α , and/or GM-CSF protein of the present invention. Examples of inhibitory compounds include an antibody of the present invention, that is administered to an animal in an effective manner (i.e., is administered in an amount so as to be present in the animal at a titer that is sufficient, upon interaction of that antibody with a native IL-4, Flt-3 ligand, CD40, CD154, IL-5, IL-13, IFN α , and/or GM-CSF protein, to decrease the activity of such proteins in an animal, at least temporarily). Oligonucleotide nucleic acid molecules of the present invention can also be administered in an effective manner, thereby reducing expression of either an IL-4, Flt-3 ligand, CD40, CD154, IL-5, IL-13, IFN α , and/or GM-CSF protein, in order to interfere with the protein activity targeted in accordance with the present invention. Peptides of an IL-4, Flt-3 ligand, CD40, CD154, IL-5, IL-13, IFN α , and/or GM-CSF protein of the present invention can also be administered in an effective manner, thereby reducing binding of IL-4, Flt-3 ligand, CD40, CD154, IL-5, IL-13, IFN α , and/or GM-CSF proteins to the appropriate receptor, in order to interfere with the protein activity targeted in accordance with the present invention. An inhibitory compound of an IL-4, Flt-3 ligand, CD40, CD154, IL-5, IL-13, IFN α , and/or

GM-CSF function can be identified using IL-4, Flt-3 ligand, CD40, CD154, IL-5, IL-13, IFN α , and/or GM-CSF proteins of the present invention, respectively.

One embodiment of the present invention is a method to identify a compound capable of inhibiting IL-4 function. Such a method includes the steps of: (a) contacting (e.g., combining, mixing) an isolated IL-4 protein of the present invention, with a putative inhibitory compound under conditions in which, in the absence of the compound, the IL-4 protein binds to IL-4 receptor or stimulates T cells in a T cell proliferation assay, and (b) determining if the putative inhibitory compound inhibits the binding of IL-4 protein to IL-4 receptor or the stimulation of T cells in a T cell proliferation assay. Another embodiment of the present invention is a method to identify a compound capable of inhibiting Flt-3 ligand function. Such a method includes the steps of: (a) contacting an isolated Flt-3 ligand protein of the present invention, with a putative inhibitory compound under conditions in which, in the absence of the compound, the Flt-3 ligand protein binds to Flt-3 receptor or stimulates dendritic precursor cells in a proliferation assay, and (b) determining if the putative inhibitory compound inhibits the binding of Flt-3 ligand protein to Flt-3 receptor or the stimulation of dendritic precursor cells in a proliferation assay. Another embodiment of the present invention is a method to identify a compound capable of inhibiting CD40 function. Such a method includes the steps of (a) contacting an isolated CD40 protein of the present invention, with a putative inhibitory compound under conditions in which, in the absence of the compound, the CD40 protein binds to a CD40 binding partner (e.g., CD154) and (b) determining if the putative inhibitory compound inhibits the binding of CD40 protein to the CD40 binding partner. A CD40 binding partner is a molecule that selectively binds to CD40 protein. Likewise, a binding partner for any other immunoregulatory protein of the present invention includes molecules that selectively bind to that particular immunoregulatory protein. Another embodiment of the present invention is a method to identify a compound capable of inhibiting CD154 function. Such a method includes the steps of (a) contacting an isolated CD154 protein of the present invention, with a putative inhibitory compound under conditions in which, in the absence of the compound, the CD154 protein binds to a CD154 binding partner (e.g., CD40) and (b) determining if the putative inhibitory compound inhibits the binding of CD154 protein to the CD154 binding partner. Yet another embodiment of the present invention is a method to identify a compound capable of inhibiting IL-5 function. Such a method includes the steps of: (a) contacting an isolated IL-5 protein of the present invention, with a putative inhibitory compound under conditions in which, in the absence of the compound, the IL-5 protein binds to IL-5 receptor or stimulates T cells in a T cell proliferation assay, and (b) determining if the putative inhibitory compound inhibits the binding of IL-5 protein to IL-5 receptor or the stimulation of T cells in a T cell proliferation assay. Another embodiment of the present invention is a method to identify a compound capable of inhibiting IL-13 function. Such a method includes the steps of: (a) contacting an isolated IL-13 protein of the present invention, with a putative inhibitory compound under conditions in which, in the absence of the compound, the IL-13 protein binds to IL-13 receptor or stimulates T cells in a T cell proliferation assay, and (b) determining if the putative inhibitory compound inhibits the binding of IL-13 protein to IL-13 receptor or the stimulation of T cells in a T cell proliferation assay. Another embodiment of the present invention is a method to identify a

compound capable of inhibiting IFN α function. Such a method includes the steps of: (a) contacting an isolated IFN α protein of the present invention, with a putative inhibitory compound under conditions in which, in the absence of the compound, the IFN α protein binds to IFN α receptor or inhibits proliferation of GM-CSF stimulated TF-1 cells, and (b) determining if the putative inhibitory compound inhibits the binding of IFN α protein to IFN α receptor or inhibits proliferation of GM-CSF stimulated TF-1 cells. Another embodiment of the present invention is a method to identify a compound capable of inhibiting GM-CSF function. Such a method includes the steps of: (a) contacting an isolated GM-CSF protein of the present invention, with a putative inhibitory compound under conditions in which, in the absence of said compound, the GM-CSF protein bind to GM-CSF receptor or stimulates T cells in a T cell proliferation assay, and (b) determining if the putative inhibitory compound inhibits the binding of GM-CSF protein to GM-CSF receptor or the stimulation of T cells in a T cell proliferation assay.

Putative inhibitory compounds to screen include small organic molecules, antibodies (including mimetopes thereof), and/or ligand analogs. Such compounds are also screened to identify those that are substantially not toxic in host animals.

Preferred IL-4, Flt-3 ligand, CD40, CD154, IL-5, IL-13, IFN α , and/or GM-CSF, proteins to inhibit are those produced by dogs, cats, horses or humans, even more preferred IL-4, Flt-3 ligand, CD40, CD154, IL-5, IL-13, IFN α , and/or GM-CSF proteins to inhibit are those produced by domestic dogs or cats. A particularly preferred inhibitor of the present invention is capable of regulating an immune response in an animal. It is also within the scope of the present invention to use inhibitors of the present invention to target diseases involving undesired immune activity in animals. Compositions comprising inhibitors of IL-4, Flt-3 ligand, CD40, CD154, IL-5, IL-13, IFN α , and/or GM-CSF function can be administered to animals in an effective manner to regulate the immune response in the animals, and preferably to prevent autoimmune disease, allergy, infectious disease, inflammation or prevent graft rejection in animals, or to treat animals with such diseases. Effective amounts and/or dosing regimens can be determined using techniques known to those skilled in the art.

It is also within the scope of the present invention to use isolated proteins, mimetopes, nucleic acid molecules and/or antibodies of the present invention as diagnostic reagents. Methods to use such diagnostic reagents are well known to those skilled in the art, see, for example, Janeway, et al., *ibid.*, and/or PCT Publication No. WO 98/23964, published Jun. 4, 1998.

The following examples are provided for the purposes of illustration and are not intended to limit the scope of the present invention.

EXAMPLES

It is to be noted that the examples include a number of molecular biology, microbiology, immunology and biochemistry techniques considered to be familiar to those skilled in the art. Disclosure of such techniques can be found, for example, in Sambrook et al., *ibid.*, and Ausubel, et al., 1993, *Current Protocols in Molecular Biology*, Greene/Wiley Interscience, New York, N.Y., and related references. Ausubel, et al. *ibid.* is incorporated by reference herein in its entirety.

Example 1

This example describes the isolation and sequencing of canine interleukin-4 (IL-4) nucleic acid molecules of the

present invention. This example also describes expression of recombinant canine IL-4 in *E. coli* and mammalian cells; development of monoclonal and polyclonal antibodies to *E. coli*-expressed canine IL-4; and bioactivity of mammalian-expressed canine IL-4.

A. Isolation and Sequencing of a Canine IL-4 Nucleic Acid Molecule.

Initial attempts to isolate a canine IL-4 nucleic acid molecule using various primers corresponding to putative conserved regions of IL-4 nucleic acid molecules failed. Forward and reverse primers were then designed using a sequence tag site (IL-4sts) described by Venta et al. in GenBank. The forward primer was designated as IL-4stsA, having the nucleic acid sequence 5' CTATTAATGG GTCT-CACCTC CCAA CT 3', designated herein as SEQ ID NO:11. The reverse primer was designated as prIL-4stsB, having the nucleic acid sequence 5' TCAACTCGGT GCA-CAGAGTC TTGG 3', designated herein as SEQ ID NO:12. The primers were used to amplify PCR products from a *C. familiaris* mitogen activated PBMC cDNA library that was constructed in the Uni-ZAP® XR vector (available from Stratagene Cloning Systems, La Jolla, Calif.), using Stratagene's ZAP-cDNA® Synthesis Kit and the manufacturer's protocol. The mRNA was isolated from *C. familiaris* peripheral blood mononuclear cells about 4 hours after they were activated by a polyclonal activating agent in culture. Four PCR products were produced that had the expected size range. The PCR products were cloned and sequenced using standard techniques. A portion of one of the four products was found to be somewhat homologous with an IL-4 nucleic acid sequence reported in GenBank.

To identify a cDNA encoding a full-length canine IL-4 protein, the PCR product showing some homology with the IL-4 sequence reported in GenBank was used to generate an about 549 base pair DNA fragment as follows. The PCR product was labeled with ³²P and used as a probe to screen the canine PBMC cDNA library. Hybridization was done at about 6×SSC, 5×Denhardt's solution, 0.5% SDS, 100 µg/ml of ssDNA and 100 µg/ml of tRNA, at about 68° C., for about 36 hr. (the compositions of SSC and Denhardt's are described in Sambrook et al., *ibid.*). The filters were washed 3 times, for about 30 minutes per wash, at about 55° C. in about 2×SSC, 0.2% SDS, followed by a final wash of about 30 minutes in the same buffer except using about 1×SSC. Positive clones were isolated and the cDNA inserts were sequenced for both strands using vector flanking primers and gene-specific internal primers. Sequence analysis was performed using the GAP program of GCG (available from the University of Wisconsin) using the alignment settings of: gap weight set at 50, length weight set at 3, and average match set at 10 for nucleic acid sequence comparisons; and gap weight set at 12, length weight set at 4, and average match set at 2.912 for amino acid sequence comparisons.

A cDNA nucleic acid molecule was isolated, referred to herein as nCaIL-4₅₄₉, the coding strand of which was shown to have a nucleic acid sequence denoted herein as SEQ ID NO:1. The complement of SEQ ID NO:1 is represented herein by SEQ ID NO:3. Translation of SEQ ID NO:1 suggests that nucleic acid molecule nCaIL-4₅₄₉ encodes a full-length IL-4 protein of about 132 amino acids, denoted herein as PCaIL-4₁₃₂, the amino acid sequence of which is presented in SEQ ID NO:2, assuming an open reading frame having an initiation codon spanning from nucleotide 43 through nucleotide 45 of SEQ ID NO:1 and a stop codon spanning from nucleotide 439 through nucleotide 441 of SEQ ID NO:1. The coding region encoding PCaIL-4₁₃₂ is

presented herein as nCaIL-4₃₉₆, which has the nucleotide sequence SEQ ID NO:4 (the coding strand) and SEQ ID NO:5 (the complementary strand). A putative signal sequence coding region extends from nucleotide 43 through nucleotide 114 of SEQ ID NO:1. The proposed mature protein (i.e., canine IL-4 protein from which the signal sequence has been cleaved), denoted herein as PCaIL-4₁₀₈, contains about 108 amino acids, extending from residue 25 through residue 132 of SEQ ID NO:2; PCaIL-4₁₀₈ amino acid sequence is represented herein as SEQ ID NO:20. The nucleic acid molecule encoding PCaIL-4₁₀₈ is denoted herein as nCaIL-4₃₂₄, extending from nucleotide 115 through nucleotide 438 of SEQ ID NO:1, nCaIL-4₃₂₄ has a coding sequence denoted SEQ ID NO:19 and a complementary sequence denoted SEQ ID NO:21.

Comparison of nucleic acid sequence SEQ ID NO:1 with nucleic acid sequences reported in GenBank indicates that SEQ ID NO:1 showed the most homology, i.e., about 89.3% identity, with a feline IL-4 gene. Comparison of amino acid sequence SEQ ID NO:2 with amino acid sequences reported in GenBank indicates that SEQ ID NO:2 showed the most homology, i.e., about 82.6% identity, with a feline IL-4 protein. Sequence analysis was performed using the GCG GAP program as described above.

B. Expression of Recombinant Canine IL-4 in *E. coli* and Mammalian Cells

i. *E. coli* Expression

A recombinant molecule capable of expressing the mature form of canine IL-4, denoted herein as pGEX-nCaIL-4₃₂₇, was produced as follows. A 340-nucleotide fragment was PCR amplified from nucleic acid molecule nCaIL-4₅₄₉ (having coding strand SEQ ID NO:1) using the following primer sequences: positive strand 5' TGAATTCGGA CAT-AACTTCA ATATTAC 3' (SEQ ID NO:38) (EcoRI site in bold) and 5' TCTCGAGATT CAGCTTCATG CCTGTA 3' (SEQ ID NO:39) (XhoI site in bold). The resulting 340-base pair fragment was digested with EcoRI and XhoI restriction enzymes (available from New England Biolabs, Beverly, Mass.), according to the manufacturer's directions, and gel-purified using standard techniques. The digested 340-base pair fragment, now 327 base pairs, was then ligated into pGEX-6P-1 (available from Amersham Pharmacia, Piscataway, N.J.), which had been previously digested with EcoRI and XhoI and gel purified, to produce recombinant molecule pGEX-nCaIL-4₃₂₇. Recombinant molecules of pGEX produce the protein of interest as a glutathione s-transferase (GST) fusion protein. The recombinant molecule pGEX-nCaIL-4₃₂₇ was transformed into DH5α cells (available from Life Technologies, Gaithersburg, Md.), a recombination deficient strain of *E. coli*, to produce recombinant cell DH5-pGEX-nCaIL-4₃₂₇. The recombinant cells were screened for presence of insert by PCR and confirmed by enzyme restriction analysis and nucleic acid sequencing, using standard techniques. Several clonal recombinant molecules were transformed into BL21 cells (available from Amersham Pharmacia, Piscataway, N.J.), a protease deficient strain of *E. coli*, to produce recombinant cell BL21-pGEX-nCaIL-4₃₂₇. These recombinant cells were screened, and the clone with the highest level of protein yield was selected for scaling up for larger-scale protein production. The resultant recombinant protein is referred to herein as *E. coli*PCaIL-4₁₀₉.

To produce and purify *E. coli*PCaIL-4₁₀₉, bacterial cultures of recombinant cell BL21:pGEX-nCaIL-4₃₂₇ were grown in shake flasks at 37° C. and induced with 0.1 mM IPTG (isopropyl-β-D-thiogalactopyranoside), (available

from Sigma Chemical Company, St. Louis, Mo.) when OD_{600nm} reached about 0.8 units. Growth was allowed to continue for about 4 hours; then bacteria were harvested by centrifugation at 4000×g (times gravity) for 20 minutes. The bacterial pellet was washed and resuspended in phosphate buffered saline (PBS) (for recipe, see Sambrook et al. *ibid.*), then lysed by exposure to gaseous nitrogen pressure in a Parr pressure vessel (available from Parr Instrument Co., Moline, Ill.), according to the manufacturer's instructions. Cell debris was removed by centrifugation at 10,000×g for 20 minutes. The IL-4-GST fusion protein E. coliPCaIL-4₁₀₉ was purified from the supernatant by allowing incubation with glutathione-conjugated resin, removing unbound proteins and then removing the GST tag with PRESCISSON™ protease; all reagents were available from Amersham Pharmacia and all were used according to the manufacturer's directions.

Concentration and purity of E. coliPCaIL-4₁₀₉ were estimated by BCA Protein Assay kit (available from Pierce, Rockford, Ill.) and SDS-PAGE followed by Coomassie staining, respectively. The purified material exhibited a single band of approximately 14 kilodaltons (kD) by Coomassie stained SDS-PAGE.

ii. CHO Cell Expression

A recombinant molecule denoted herein as pCMV-nCaIL-4₃₉₀, capable of expressing a full length form of canine IL-4 (including signal sequence) was produced as follows. A 422-nucleotide fragment was PCR amplified from nucleic acid molecule nCaIL-4₁₄₉. Using the following primers: 5' CCCAAGCTTA TGGGTCTCACC TCCCAAC (HindIII site in bold), denoted SEQ ID NO:40, and 3' CCTCGAGATT CAGCTTTCAA TGCCTGTA (XhoI site in bold), denoted SEQ ID NO:127. The 422-base pair PCR product was digested with the restriction endonucleases HindIII and xhoI, both available from New England Biolabs. The resulting 399-base pair product encoding full-length canine IL-4 was gel purified using standard techniques and ligated into the cytomegalovirus (CMV) immediate-early transcription control region of the pCMV-Int A plasmid vector that had been digested with HindIII and XhoI (available from New England Biolabs), and gel purified, to produce the recombinant molecule pCMV-nCaIL-4₃₉₉. The pCMV-Int A plasmid vector was generated as referenced by J. E. Osorio et al., 1999, Vaccine 17, 1109–1116. Briefly, vector pRc/RSV, (available from Invitrogen Corp., San Diego, Calif.) was cleaved with restriction enzyme PvuII (available from New England Biolabs), and the 2963-base pair PvuII fragment was gel purified. The fragment was self-ligated to form the vector pRc/RSV(Pvu), which contains a Rous Sarcoma Virus (RSV) long terminal repeat, a multiple cloning site, a bovine growth hormone polyadenylation sequence, a bacterial origin of replication, and an ampicillin resistance gene. Vector pRc/RSV(Pvu) was restriction enzyme digested using HindIII and NruI. A HindIII/SspI fragment containing the HCMV intermediate early promoter and first intron (i.e. intron A) was ligated into the digested pRc/RSV(Pvu) vector to produce the vector pCMV-Int A.

Stable expression of CaIL-4 in mammalian cells was carried out by transfecting the recombinant molecule pCMV-nCaIL-4₃₉₉ into Chinese Hamster Ovary cells, (CHO, available from ATCC) as follows. Six-well polystyrene tissue culture plates (available from Corning Costar, Acton, Mass.) were seeded with approximately 5×10⁵ cells/well in 2 milliliter (ml) cell culture media, consisting of Minimal Essential Media (MEM) supplemented with 100 mM L-glutamine, 100 mM gentamicin, and 10% fetal

bovine serum (FBS), (all available from Life Technologies). Cells were grown to about 80% confluence (for about 18 hours) before transfection. The recombinant molecules to be transfected were purified using the Plasmid Midi Kit (available from Qiagen, Valencia, Calif.) and used according to the manufacturer's instructions. The recombinant molecule pCMV-nCaIL-4₃₉₉ was linearized using the restriction enzyme PvuI (available from New England Biolabs). The plasmid pcDNA3, (available from Invitrogen), which contains the neomycin resistance gene, was linearized using the restriction enzyme EcoRI. Approximately 2 µg of pCMV-nCaIL-4₃₉₉ was mixed with about 2 ng of linearized pcDNA3 in about 100 µl OPTIMEM™ media, available from Life Technologies. About 10 µl Lipofectamine, (available from Life Technologies) was mixed with 100 µl OPTIMEM. The nucleic acid molecule-containing mixture was then added to the Lipofectamine mixture and incubated at room temperature for about 45 minutes. After incubation, about 0.8 ml OPTIMEM was added, and the mixture was overlaid onto the CHO cells which had been previously rinsed with OPTIMEM. Cells were incubated for about 5 hours at 37° C. 5% CO₂, 95% relative humidity. Approximately 1 ml of cell culture media as described previously, with 20% FBS, was added and the cells were incubated overnight. The media was changed at 24 hours, and at 72 hours post transfection, the cells were split 1:4 and put into fresh cell culture media containing about 500 µg/ml geneticin (G418, available from Life Technologies). The media was changed every 3–5 days. After several weeks, G418 resistant colonies were trypsinized using sterile filter papers, 5–6 mm in diameter that were soaked in trypsin, which were then placed over individual wells of 24 well plates that contained separated widely spaced colonies of CHO cells. After 3 days, the papers were removed. The resulting recombinant cells are referred to herein as CHO-pCMV-nCaIL-4₃₉₉. The recombinant cells were then expanded and tested for the presence of nIL-4399 RNA by RT-PCR and tested for the presence of PCaIL-4₁₃₃ protein by Western blot analysis. Westerns were developed with rabbit anti-E. coliPCaIL-4₁₀₉ serum and 607.1 monoclonal antibody, a monoclonal antibody that selectively binds to E. coliPCaIL-4₁₀₉ protein. See Example 1C for a description of how these antibodies were produced.

C. Monoclonal and Polyclonal Antibodies to Recombinant canine IL-4 (i.e., Anti-canine IL-4 Antibodies)

The following describes the development of monoclonal and polyclonal antibodies that selectively bind to E. coliPCaIL-4₁₀₉.

Female Balb/C mice, 6–8 weeks old, were injected subcutaneously, at about 4 sites, with a total of 25 µg E. coliPCaIL-4₁₀₉ (produced as described in Example 1Bi) in Freund's Complete Adjuvant (day 0). Fourteen days later, the mice received an intraperitoneal boost of 25 µg E. coliPCaIL-4₁₀₉ in Freund's Incomplete Adjuvant (day 14). Fourteen days later, serum was tested for antibody titer to E. coliPCaIL-4₁₀₉ by ELISA (day 28). Three days prior to fusion, mice were boosted intravenously with 20 µg E. coliPCaIL-4₁₀₉ in PBS (day 35). Splenocytes were harvested from mice demonstrating the highest serum titer by ELISA and depleted of CD4+ and CD8+ cells. This depletion was achieved by incubation of the splenocytes with biotinylated rat anti-mouse CD4 and anti-mouse CD8 monoclonal antibodies, available from PharMingen, San Diego, Calif. Antibody-labeled cells were then removed by incubation with M-280 streptavidin coated magnetic beads,

available from Dynal, Oslo, Norway. Depleted splenocytes were fused to SP2/0 cells (available from ATCC) using 50% polyethylene glycol in unsupplemented Iscove's Modified Dulbecco's Media (IMDM), following established protocols; see, for example, Harlow E., and Lane D., eds., 1995, *Antibodies, A Laboratory Manual*, Monoclonal Antibodies, Cold Spring Harbor Laboratories; Harlow et al. *ibid.*, is incorporated by reference herein in its entirety. Fused cells were plated in 96-well plates using IMDM cell culture media, (available from Life Technologies, Inc., Rockville, Md.), which was supplemented with 10% fetal bovine serum, 2 mM L-glutamine, 1 mM sodium pyruvate, 1xnonessential amino acids, 1xMEM amino acids, 0.05 mg/ml gentamicin, and 0.5 mM β -mercaptoethanol (all reagents available from Life Technologies). Additionally, 100 μ M hypoxanthine, 0.4 μ M aminopterin, and 16 μ M thymidine, all available from Sigma Chemical Corporation, St Louis, Mo., were added.

After about 7 days, wells positive for hybridoma growth were screened by ELISA to E. coliPCaIL-4₁₀₉. Immulon II 96-well plates (available from VWR, Denver, Colo.) were coated, overnight, with 100 ng/ml E. coliPCaIL-4₁₀₉ in 0.1 M carbonate/bicarbonate buffer, Ph 9.6. After blocking the wells with 20% FBS in Tris buffered saline (TBS), culture supernatants were allowed to bind. Presence of anti-E. coliPCaIL-4₁₀₉ mouse antibody was detected with polyclonal goat anti-mouse IgG conjugated to horseradish peroxidase, (available from KPL, Gaithersburg, Md.), and color developed with 3,3',5,5'-tetramethylbenzidine dihydrochloride (TMB), available from Pierce, Rockford, Ill. Specificity of the ELISA reactivity was verified by Western blot analysis to E. coliPCaIL-4₁₀₉, developed with polyclonal goat anti-mouse IgG conjugated to alkaline phosphatase and nitro-blue tetrazolium/5-bromo-4-chloro-3'-indolylphosphate p-toluidine salt substrate (NBT/BCIP, available from Sigma). Western blots exhibited a single band of approximately 14 kD. Immunoglobulin isotype of the monoclonal antibodies was determined using IsoStrips, available from Boehringer Mannheim, Indianapolis, Ind. Twenty-three monoclonal antibodies were generated to E. coliPCaIL-4₁₀₉, 22 of which were of the IgM isotype and one of which was IgG1, and is referred to herein as 607.1.

Polyclonal rabbit serum was produced by repeated immunization (over a 10 month period) of a male, New Zealand White rabbit 12-16 months old. Initial immunization was 50 μ g E. coliPCaIL-4₁₀₉ (prepared as described in Example 1bi) in Freund's Complete Adjuvant, at several sites subcutaneously and intradermally. One month later, and at one month intervals thereafter, the rabbit was boosted intradermally with 50 μ g E. coliPCaIL-4₁₀₉ in Freund's Incomplete Adjuvant. Serum was collected bi-weekly and titers monitored by ELISA and Western blot to E. coliPCaIL-4₁₀₉. Serum that selectively bound to E. coliPCaIL-4₁₀₉ protein is referred to as anti-E. coliPCaIL-4₁₀₉ serum.

D. Bioactivity of Mammalian-expressed Canine IL-4

4

The following describes a bioassay to detect the expression of canine IL-4 protein expressed in the supernatants from CHO-pCMV-nCaIL-4₃₉₉ recombinant cells by screening for production of CD23.

About 100 μ l Ramos cells, available from ATCC, at a concentration of about 3.5×10^3 cells/ml were seeded into 96-well flat bottom plates, available from Becton Dickinson, Franklin Lakes, N.J.). These cells were grown in RPMI media supplemented with 100 mM L-glutamine,

gentamicin, and 10% FBS (called TCM). The Ramos cells were then treated in 5% CO₂ for 37° C. for approximately 48 h. with one of the following:

Group Treatment

1 TCM

2 CHO-pCMV (a transfectant cell line containing the empty pCMV vector) supernatant (1:4 final dilution in TCM)

3 CHO-pCMV-nCaIL-4₃₉₉ supernatant (1:20 final dilution in TCM)

Triplicate samples for each treatment group were pooled for staining to look for increased expression of CD23 (one of the reported effects of IL-4). Briefly, 1×10^5 cells from each treatment group were incubated in phosphate buffered saline (PBS) containing 30% FBS for 15-30 min on ice. The cells were collected and incubated with the following:

Condition	Primary Incubation	Secondary Location
A	PBS	Goat anti mouse PE
B	Mouse anti human CD23	Goat anti mouse PE

Condition Primary Incubation Secondary Incubation
Mouse anti-human CD23 monoclonal antibody, available from Pharmingen, was used at about 10 μ g/ml. Goat (Fab'2) anti mouse IgG PE, available from Southern Biotechnologies was used at about 2.5 μ g/ml. These reagents were diluted in PBS with 5% FBS. Primary incubations were performed for 30-60 minutes on ice, and secondary incubations were performed for 20-30 min on ice. Three washes of PBS/5% FBS were performed in between each incubation. Cells were then analyzed on a flow cytometer (e.g., MoFlow Desk Top System, available from Cytomation, Ft. Collins, Colo.) with the fluorescein gate set at 10¹. The results are shown below in Table 2.

TABLE 2

Induction of CD23 on Ramos cells post-treatment with supernatants from CHO-pCMV-nCaIL-4 ₃₉₉		
Treatment	Condition	% positive
1	A	0
	B	1
2	A	8
	B	1
3	A	3
	B	99

Table 2 shows that the canine IL-4 expressed by the CHO transfectant CHO-pCMV-nCaIL-4₃₉₉ is biologically active, demonstrated by its ability to induce expression of CD23 in Ramos cells.

Example 2

This example describes the isolation and sequencing of certain canine Flt-3 ligand and feline Flt-3 nucleic acid molecules and proteins of the present invention. The example also describes expression of a canine Flt-3 ligand protein of the present invention in CHO cells, as well as detection of the expressed canine Flt-3 ligand protein.

A. Canine Flt-3 Ligand Nucleic Acid Molecules and Proteins.

i. This example describes the isolation and sequencing of certain canine Flt-3 ligand nucleic acid molecules and proteins of the present invention.

61

A canine Flt-3 ligand nucleic acid molecule was produced as follows. A pair of primers was initially used to amplify DNA from the *C. familiaris* mitogen activated PBMC cDNA library described above in Example 1. A forward primer referred to as FLT3F1, having the nucleic acid sequence 5' CTGGCGCCAG CCTGGAGCCC 3', designated herein as SEQ ID NO:13 was used in combination with a reverse primer referred to herein as FLT3B1, having the nucleic acid sequence 5' GGGAGATGTT GGTCTGGACG 3', referred to herein as SEQ ID NO:14 to amplify Flt-3 ligand DNA from the cDNA library by polymerase chain reaction (PCR). The primers were designed using conserved regions of IL-4 cDNA sequences from other species in the public databases corresponding to the positions shown below:

Database	Accession number	Nucleotides	Animal
gb	U04806	102-121	human
gb	L23636	41-60	mouse
gb	U04806	77-458	human
gb	L23636	419-400	mouse

A 360-base pair (bp) PCR product was generated in the above reaction that was purified, radiolabeled and used as a probe to screen the cDNA library. Hybridization was performed in 6×SSC, 5×Denhardt's solution, 0.5% SDS, 100 µg/ml ssDNA and 100 µg/ml of tRNA, at 68° C., for about 36 hr. The filters were washed 3 times, for about 30 minutes per wash, at 55° C. in 2×SSC, 0.1% SDS, followed by a final wash in 0.25×SSC, for about 30 minutes, at 55° C. Several positive phage clones were identified and shown to produce PCR products when used as templates in combination with the FLT3F1 and FLT3B1 primers. The DNA inserts in the phage clones were sequenced using standard techniques and failed to yield any clones containing DNA inserts having a portion homologous to published Flt-3 ligand sequences. The 360-bp PCR fragment generated above was then cloned into the vector pcDNA 2.1 (available from Invitrogen Corp., San Diego, Calif.). Several independent colonies were generated and the sequences of their inserts determined. One clone was identified that which contained insert sequence having a portion, that was somewhat homologous to published human or murine Flt-3 ligand sequence.

Two canine Flt-3 ligand-specific primers were then designed using the nucleic acid sequence obtained using the 360-bp PCR product described above.

Primer	Sequence	SEQ ID NO
DFLB1	5'GACCAGGCGCCAGAACGC 3'	SEQ ID NO:15
DFLF1	5'CGGTCACCATCCGCAAGC 3'	SEQ ID NO:16

The 5' region of a Flt-3 ligand nucleic acid molecule was PCR amplified from the cDNA library using the DFLB1 primer in combination with the 5' T3 vector primer from the Uni-ZAP® XR vector (available from Stratagene). The 3' region of a Flt-3 ligand nucleic acid molecule was PCR amplified from the cDNA library using the DFLF1 in combination with the 3' T7 primer from the Uni-ZAP® XR vector (available from Stratagene). A 855-bp PCR product was obtained representing the 5' region of a Flt-3 ligand nucleic acid molecule. A 265-bp PCR product was obtained representing the 3' region of a Flt-3 ligand nucleic acid molecule. Both the 855-bp PCR product and 265-bp PCR

62

product were cloned and sequenced using standard methods. Additional canine Flt-3 ligand-specific primers were designed using the nucleic acid sequence obtained from the sequence of the 855-bp PCR product and 265-bp PCR products.

Primer	Sequence	SEQ ID NO
DFLB2	5'TGGCAAGGCAGTGGCCTC 3'	SEQ ID NO:17
DFLF2	5'GCCGAGATGATAGTGCTGGC 3'	SEQ ID NO:18

A 546-bp PCR product was generated using the primer DFLF2 in combination with the primer DFLB2 to amplify a PCR product from the cDNA library. The 546-bp PCR product was then purified, radiolabeled and used as a probe to screen the cDNA library. Hybridization was performed in 6×SSC, 5×Denhardt's solution, 0.5% SDS, 100 µg/ml of ssDNA and 100 µg/ml of tRNA, at 68° C. for about 36 hr. The filters were washed in 1.25×SSC, for about 30 minutes, at 55° C. Four cDNA clones encoding full-length canine Flt-3 ligand were isolated. Nucleic acid sequence was obtained using standard techniques.

A Flt-3 ligand clone was isolated, referred to herein as nCaFlt3L₁₀₁₃, the coding strand of which was shown to have a nucleic acid sequence denoted herein as SEQ ID NO:6. The complement of SEQ ID NO:6 is represented herein by SEQ ID NO:8. Translation of SEQ ID NO:6 suggests that nucleic acid molecule nCaFlt3L₁₀₁₃ encodes a full-length Flt-3 ligand protein of about 294 amino acids, denoted herein as PCaFlt3L₂₉₄, the amino acid sequence of which is presented in SEQ ID NO:7, assuming an open reading frame having an initiation codon spanning from nucleotide 35 through nucleotide 37 of SEQ ID NO:6 and a stop codon spanning from nucleotide 917 through nucleotide 919 of SEQ ID NO:6. The coding region encoding PCaFlt3L₂₉₄ is presented herein as nCaFlt3L₈₈₂, which has the nucleotide sequence SEQ ID NO:9 (the coding strand) and SEQ ID NO:10 (the complementary strand). A putative signal sequence coding region extends from nucleotide 35 through nucleotide 112 of SEQ ID NO:6. The proposed mature protein (i.e., canine Flt-3 ligand protein from which the signal sequence has been cleaved), denoted herein as PCaFlt3L₂₆₈ (SEQ ID NO:23), contains about 268 amino acids, extending from residue 27 through residue 294 of SEQ ID NO:7. The nucleic acid molecule encoding PCaFlt3L₂₆₈ is denoted herein as nCaFlt3L₈₀₄, extending from nucleotide 113 through nucleotide 916 of SEQ ID NO:6, nCaFlt3L₈₀₄ has a coding sequence denoted SEQ ID NO:22 and a complementary sequence denoted SEQ ID NO:24.

Below is a description of the identification of alternatively spliced Canis Flt3 ligand transcripts. Besides cDNA clones such as nucleic acid molecule nCaFlt3L₁₀₁₃ encoding the full-length canine FLT3 ligand protein, two splice variants of canine Flt3 ligand cDNA clones were also isolated, using the same hybridization conditions as mentioned previously in this Example. One such variant (Clone 1), denoted herein as nCaFlt3L₉₈₅, has a coding strand the nucleic acid sequence of which is represented as SEQ ID NO:25. The complement of SEQ ID NO:25 is represented herein by SEQ ID NO:27. Translation of SEQ ID NO:25 suggests that nucleic acid molecule nCaFlt3L₉₈₅ encodes a Flt-3 ligand protein of 276 amino acids, denoted herein as PCaFlt3L₂₇₆, the amino acid sequence of which is represented by SEQ ID NO:26, assuming an open reading frame having an initiation

codon spanning from nucleotide 74 through nucleotide 76 of SEQ ID NO:25 and a stop codon spanning from nucleotide 902 through nucleotide 904 of SEQ ID NO:25. The coding region encoding PCaFlt3L₂₇₆ is represented herein as nCaFlt3L₈₂₈, which has the nucleotide sequence SEQ ID NO:28 (the coding strand) and SEQ ID NO:29 (the complementary strand). Alignment of nucleic acid molecules nCaFlt3L₈₁₂ and nCaFlt3L₈₂₈ indicates that the nucleic acid molecules are identical except for a deletion in nCaFlt3L₈₂₈, which spans from nucleotide 343 through nucleotide 396 of nCaFlt3L₈₈₂. Accordingly, nCaFlt3L₈₂₈ encodes 18 fewer amino acids than nCaFlt3L₈₈₂. The deletion in PCaFlt3L₂₇₆, which spans from residue 115 through residue 132 of PCaFlt3L₂₉₄, occurs between helix III and helix IV of the canine Flt3 ligand protein inferred from alignment with the human and mouse Flt3 ligand protein (Lyman et al., Cell, vol. 75, pp. 1157–1167, 1993; Hannum et al., Nature, vol. 368, pp. 643–648, 1994; Lyamn et al., Blood, vol. 83, pp. 2795–2801, 1994). In addition, the alignment shows that there are 39 more nucleotides in the 5' untranslated region of nucleic acid molecule nCaFlt3L₉₈₅ (nucleotides 1 to 39) than nucleic acid molecule nCaFlt3L₁₀₁₃ and there are 2 more nucleotides in the 3' untranslated region of nucleic acid molecule nCaFlt3L₉₈₅ (nucleotides 922 to 923) than nucleic acid molecule nCaFlt3L₁₀₁₃. The remaining sequences between nCaFlt3L₉₈₅ and nCaFlt3L₁₀₁₃ are identical. A putative mature form of nCaFlt3L₉₈₅ (without the signal sequence) is predicted. The putative signal sequence coding region extends from nucleotide 74 to nucleotide 151 of SEQ ID NO:25. The proposed mature protein, denoted herein as PCaFlt3L₂₅₀, represented by SEQ ID NO:31, contains about 250 amino acids, extending from residue 27 through residue 276 of SEQ ID NO:26. The nucleic acid molecule encoding PCaFlt3L₂₅₀, extending from nucleotide 152 through nucleotide 901 of SEQ ID NO:6, denoted herein as nCaFlt3L₇₅₀, is represented by SEQ ID NO:30 (the coding strand) and SEQ ID NO:32 (the complement strand).

A second variant (Clone 19) is represented by nucleic acid molecule nCaFlt3L₁₀₁₉, the coding strand of which is denoted herein as SEQ ID NO:33. The complement of SEQ ID NO:33 is denoted herein as SEQ ID NO:35. Translation of SEQ ID NO:33 suggests that nCaFlt3L₁₀₁₉ encodes a Flt-3 ligand protein of 31 amino acids, PCaFlt3L₃₁, denoted SEQ ID NO:34, assuming an initiation codon spanning from nucleotide 74 through nucleotide 76 and a stop codon spanning nucleotide 167 through nucleotide 169 of SEQ ID NO:33. The coding region encoding PCaFlt3L₃₁ is represented herein as nCaFlt3L₉₃, which has the nucleotide sequence SEQ ID NO:36 (the coding strand) and SEQ ID NO:37 (the complementary strand). Alignment of nucleic acid molecules nCaFlt3L₉₈₅ and nCaFlt3L₁₀₁₉ indicates the presence of an insertion of 91 nucleotides in nCaFlt3L₁₀₁₉. The insertion spans nucleotide 107 through nucleotide 198 of nCaFlt3L₁₀₁₉. A stop codon is found in this insertion in frame with the predicted initiation codon, which span nucleotide 74 through nucleotide 76 of SEQ ID NO:6. Since this insertion (with an in-frame stop codon) occurs in or close to the signal peptide, it is likely that nucleic acid molecule nCaFlt3L₁₀₁₉ encodes a nonfunctional Flt-3 ligand protein.

Comparison of nucleic acid sequence SEQ ID NO:6 with nucleic acid sequences reported in GenBank indicates that SEQ ID NO:6 showed the most homology, i.e., about 69.8% identity, with a human Flt-3 ligand gene. Comparison of amino acid sequence SEQ ID NO:7 with amino acid sequences reported in GenBank indicates that SEQ ID NO:7 showed the most homology, i.e. about 71% identity, with a human Flt-3 ligand protein. Sequence analysis was per-

formed with DNAsis™ using the alignment settings of: gap penalty set at 5; number of top diagonals set at 5; fixed gap penalty set at 10; K-tuple set at 2; window size set at 5 and floating gap penalty set at 10.

ii. This example describes the production of a recombinant molecule encoding a full length canine Flt-3 ligand protein and expression of that protein by a recombinant cell of the present invention.

A recombinant molecule, denoted herein as pCMV-nCaFlt3L₈₈₂ and capable of expressing a full length form of Flt-3 ligand, was produced as follows. Nucleic acid molecule nCaFlt3L₈₈₂ was digested with the restriction endonucleases EcoRI and XbaI, gel purified and ligated into pCMV-Int A (prepared by methods described in Example 1) to produce recombinant molecule pCMV-nCaFlt3L₈₁₂. Insert size and identity were confirmed by restriction digestion, PCR, and sequencing analyses.

Stable transfectants expressing the recombinant molecule pCMV-nCaFlt3L₈₈₂ were established in Chinese Hamster Ovary cells (CHO, available from ATCC) as follows. Briefly, six-well polystyrene tissue culture plates were seeded with approximately 4×10^5 cells per well in 2 ml of MEM (available from Life Technologies, Gaithersburg, Md.) supplemented with 100 mM L-glutamine, gentamicin, and 10% FBS (TCM). Cells were grown to about 80% confluence (about 18 hr). The recombinant molecule to be transfected was prepared using the Qiagen Endotoxin-Free Plasmid Maxi Kit as per the manufacturer's instructions. The recombinant molecule was linearized with the restriction enzyme PvuI, extracted with phenol, and precipitated with isopropanol. The plasmid pcDNA 3, available from Invitrogen, which contains the neomycin resistance gene, was linearized with the restriction enzyme EcoRI. Approximately 1 μ g of recombinant plasmid DNA and 100 ng of pcDNA3 were mixed with about 100 μ l OptiMEM medium, available from Life Technologies. About 10 μ l Lipofectamine (available from Life Technologies) was mixed with 100 μ l OptiMEM. The DNA-containing mixture was then added to the Lipofectamine mixture and incubated at room temperature for about 30 min. After incubation, about 800 μ l of OptiMEM was added, and the entire mixture was overlaid onto the CHO cells that had been rinsed with OptiMEM. Cells were incubated for 6 hours at 37° C., 5% CO₂, 95% relative humidity. Approximately 1 ml of TCM with 20% FBS was added, and the cells were incubated overnight. The media was changed after about 24 hr. About 72 hr post transfection, the cells were split 1:4 and put into selection TCM containing 500 μ g/ml Geneticin (G418), available from Life Technologies. The medium was changed every 3–5 days. After several weeks, G418-resistant colonies were trypsinized, and the cells were plated into 24 well plates. The resulting recombinant cells are referred to herein as CHO-pCMV-nCaFlt3L₈₈₂. The recombinant cells were then expanded for testing.

iii. The following describes the detection of expression of a canine Flt-3 ligand protein of the present invention by CHO-pCMV-nCaFlt3L₈₈₂, a recombinant cell of the present invention.

Recombinant cells CHO-pCMV-nCaFlt3L₈₈₂, produced as described in Example 2, part (B)(ii) above, were tested for surface expression of canine Flt-3 ligand using a cross-reactive goat anti-human Flt-3 ligand polyclonal antibody as follows. Briefly, 1×10^5 CHO-pCMV-nCaFlt3L₈₈₂ cells or CHO-pCMV cells (i.e., cells transfected with an empty vector as described in Example 1) were incubated in phosphate buffered saline (PBS) containing 30% fetal bovine

65

serum (FBS) for about 30 min on ice. The cells were then spun down and treated with the following:

Condition	Primary Incubation	Secondary Incubation
1	PBS	Rabbit (Fab' 2) anti sheep (H + L) FITC
2	Goat anti-human Flt3 ligand	Rabbit (Fab' 2) anti sheep (H + L) FITC

Goat anti-human Flt3 ligand, available from R and D Systems, Minneapolis, Minn. was used at about 20µg/ml. Rabbit (Fab' 2) anti sheep (H+L) FITC, available from Southern Biotechnology Associates, Inc., was used at about 10 µg/ml. These reagents were diluted in PBS/5%FBS. All incubations were in 50 µl for about 1 hr on ice with 2 washes of PBS/5%FBS in between each incubation. Cells were then analyzed on a flow cytometer (e.g., MoFlow Desk Top System, available from Cytomation, Ft. Collins, Colo.) with the fluorescein gate set at 10¹. The results are shown below in Table 3.

TABLE 3

Expression of canine Flt3 ligand on CHO transfectants.		
Cells	% positive	
	Condition 1	Condition 2
CHO-pCMV	1	1
CHO-pCMV nCaFlt3L ₈₈₂	2	48
CHO-pCMV nCaFlt3L ₈₈₂	1	20

Table 3 shows that canine Flt3 ligand is expressed on the surface of the CHO transfectants.

B. Feline Flt-3 Ligand Nucleic Acid Molecules and Proteins.

This example describes the production of certain feline Flt-3 ligand nucleic acid molecules and proteins of the present invention.

A nucleic acid molecule encoding a feline Flt-3 ligand was isolated from a feline PBMC cDNA library as follows. A *Felis catus* mitogen activated PBMC cDNA library was constructed in the Uni-Zap-R XRTM vector, available from Stratagene, La Jolla, Calif., using Stratagene's Zap cDNA-RTM Synthesis Kit and the manufacturer's protocol using mRNA isolated from *F. catus* peripheral blood mononuclear cells about 4 hours after they were activated by a polyclonal activating agent in culture. PCR amplification to isolate a feline Flt 3 ligand nucleic acid molecule was conducted according to the following set of steps: one initial denaturation step at 95° C. for 3 minutes; then 35 cycles of the following; 94° C. for 30 seconds, 53.8° C. for 30 seconds, and 72° C. for 105 seconds; then one final extension step at 72° C. for 8 minutes. A 395 nucleotide cDNA fragment containing the 5' end of feline Flt3 ligand coding region, denoted nFeFlt3L₃₉₅, was amplified from the feline PMBC cDNA library using the following primers: vector primer T3 having nucleic acid sequence 5' AATTAACCT CAC-TAAAGGG 3' (SEQ ID NO:142) (available from Stratagene) and the antisense primer having SEQ ID NO:14, described in Example 2A. The nucleic acid sequence of the coding strand of nFeFlt3L₃₉₅ is denoted SEQ ID NO:41. A 793-nucleotide cDNA fragment containing the 3' end of feline Flt3 ligand coding region, denoted nFeFlt3L₇₉₃, was

66

isolated using sense primer 2 having the nucleic acid sequence 5' CACAGYCCCA TCTCTCC 3' (where Y was either T or C) denoted herein as SEQ ID NO:151, in conjunction with vector primer T7 having the nucleic acid sequence 5' GTAATACGAC TCACTATAGG GC 3' (SEQ ID NO:152). The nucleic acid sequence of the coding strand of nFeFlt3L₇₉₃ is denoted SEQ ID NO:42. Nucleic acid molecules nFeFlt3L₃₉₃ and nFeFlt3L₇₉₃ overlap by 246 nucleotides and form a composite sequence encoding a Flt3 ligand protein that is similar in length to that of PCaFlt3L₂₉₄. This composite feline Flt3 ligand cDNA is referred to herein as nFeFlt3L₉₄₂, the coding strand of which was shown to have nucleic acid sequence SEQ ID NO:43. The reverse complement of SEQ ID NO:43 is referred to herein as SEQ ID NO:45. Translation of SEQ ID NO:43 suggests that nucleic acid molecule nFeFlt3L₉₄₂ encodes a Flt3 ligand protein of 291 amino acids, denoted herein as PFeFlt3L₂₉₁, the amino acid sequence of which is presented in SEQ ID NO:44, assuming an open reading frame having an initiation codon spanning from nucleotide 31 through nucleotide 33 of SEQ ID NO:43 and a stop codon spanning from nucleotide 904 through nucleotide 906 of SEQ ID NO:43. The coding region encoding PFeFlt3L₂₉₁, not including the termination codon, is presented herein as nFeFlt3L₈₇₃, which has the nucleotide sequence SEQ ID NO:46 (the coding strand) and SEQ ID NO:47 (the complementary strand). A putative signal sequence coding region extends from nucleotide 31 to nucleotide 108 of SEQ ID NO:43. The proposed mature protein, denoted herein as PFeFlt3L₂₆₅, denoted SEQ ID NO:49, contains about 265 amino acids, extending from residue 27 through residue 291 of SEQ ID NO:44. The nucleic acid molecule encoding PFeFlt3L₂₆₅, is denoted herein as nFeFlt3L₇₉₅, (SEQ ID NO:48) extending from nucleotide 109 through nucleotide 903 of SEQ ID NO:43. SEQ ID NO:48 has a complementary strand denoted SEQ ID NO:50.

Sequence alignment indicates that nucleic acid sequence SEQ ID NO:43 shares the highest (67.8%) identity with the nucleic acid sequence of human Flt-3 ligand (GenBank accession numbers U04806 and U03858). Amino acid sequence SEQ ID NO:44 shares the highest (70.2%) identity with human Flt-3 ligand protein (GenBank accession numbers U04806 and U03858).

Example 3.

This example describes the isolation and sequencing of certain canine CD40 and feline CD40 nucleic acid molecules and proteins of the present invention.

A. Canine CD40 Nucleic Acid Molecules and Proteins

This example describes the production of certain canine CD40 nucleic acid molecules and proteins of the present invention.

A canine CD40 nucleic acid molecule of the present invention was produced by PCR amplification as follows. A 321-nucleotide canine CD40 nucleic acid molecule, denoted nCaCD40₃₂₁, was amplified from a canine PBMC cDNA library, prepared as described in Example 1, using two degenerate oligonucleotide primers designed in accordance with conserved regions of human, bovine, rabbit, and mouse CD40 gene sequences available in GenBank; sense primer, 5' TGCCCRSTCG GCTTCTTCTC C 3', denoted herein as SEQ ID NO:128; and antisense primer, 5' CGACTCTCTT TRCCRTCCTC CTG 3', denoted herein as SEQ ID NO:129, where R was either A or G and S was either G or C. PCR

67

conditions were as follows: one initial denaturation step at 95° C. for 3 minutes; then 35 cycles of the following: 94° C. for 30 seconds, then 53° C. for 30 seconds, then 72° C. for 105 seconds; followed by one final extension at 72° C. for 5 minutes. The resulting PCR product, i.e., nCaCD40₃₂₁, with a coding strand represented by SEQ ID NO:51, was radiolabeled using standard techniques and used to screen the canine PBMC cDNA library, under the following hybridization conditions: hybridized in 6×SSC, 5×Denhardt's solution, 0.5% SDS, 100 µg/ml single stranded DNA, 100 µg/ml tRNA for 36 hours at 68° C., followed by a wash of 0.1% SDS, 1×SSC at 55° C. for 60 minutes. A clone (Clone 18B) containing a 1425-nucleotide canine CD40 nucleic acid molecule, denoted nCaCD40₁₄₂₅, was obtained. The nucleic acid sequence of the coding strand of nCaCD40₁₄₂₅ is represented as SEQ ID NO:52. The reverse complement of SEQ ID NO:52 is referred to herein as SEQ ID NO:54. Translation of SEQ ID NO:52 suggests that nucleic acid molecule nCaCD40₁₄₂₅ encodes a canine CD40 protein of 274 amino acids, denoted herein as PCaCD40₂₇₄, the amino acid sequence of which is presented in SEQ ID NO:53, assuming an open reading frame having an initiation codon spanning from nucleotide 196 through nucleotide 198 of SEQ ID NO:52 and a stop codon spanning from nucleotide 1018 through nucleotide 1020 of SEQ ID NO:52. The coding region encoding PCaCD40₂₇₄, not including the termination codon, is presented herein as nCaCD40₈₂₂, which has the nucleotide sequence SEQ ID NO:55 (the coding strand) and SEQ ID NO:56 (the complementary strand).

A putative signal sequence coding region extends from nucleotide 196 through nucleotide 252 of SEQ ID NO:52. The proposed mature protein, denoted herein as PCaCD40₂₅₅, represented by SEQ ID NO:58, contains about 255 amino acids, extending from residue 20 through residue 274 of SEQ ID NO:53. The nucleotide sequence encoding PCaCD40₂₅₅, which extends from nucleotide 253 through nucleotide 1017 of SEQ ID NO:52, is denoted herein as nucleic acid molecule nCaCD40₇₆₅, represented by SEQ ID NO:57 (the coding strand) and SEQ ID NO:59 (the complement strand).

Sequence analysis was performed with DNAsis™ using the alignment settings of: gap penalty set at 5; number of top diagonals set at 5; fixed gap penalty set at 10; k-tuple set at 2; window size set at 5 and floating gap penalty set at 10. At the amino acid level, PCaCD40₂₇₄ shares 65.3%, 50.1%, and 42.3% identity with the CD40 proteins of human, bovine, and mouse, respectively (Stamenkovic et al., EMBO J., vol. 8:1403–1410, 1989; Hirano et al., Immunology, vol. 90, pp. 294–300, 1997; Grimaldi et al., J. Immunol., vol. 143, pp. 3921–3926; Torres and Clark, J. Immunol., vol. 148, pp. 620–626). At the nucleotide level, nCaCD40₁₄₂₅ shares 69.3%, 69.4%, and 40.4% identity with the cDNA sequences of human, bovine, and mouse CD40, respectively.

B. Feline CD40 Nucleic Acid Molecules and Proteins

This example describes the isolation and sequencing of certain nucleic acid molecules of the present invention that encode certain feline CD40 proteins of the present invention.

A 336-nucleotide feline CD40 nucleic acid molecule, denoted nFeCD40₃₃₆, was amplified from a feline PBMC cDNA library, prepared as described in Example 2, using PCR conditions and primers as described in Example 3A, i.e., a sense primer having SEQ ID NO:128; and an antisense

68

primer having SEQ ID NO:129. The resulting PCR product, i.e., nFeCD40₃₃₆, was shown to have a coding strand the nucleic acid sequence of which is represented as SEQ ID NO:60. The reverse complement of SEQ ID NO:60 is referred to herein as SEQ ID NO:62. Translation of SEQ ID NO:60 suggests that nucleic acid molecule nFeCD40₃₃₆ encodes a partial CD40 protein of 112 amino acids, denoted herein as PFeCD40₁₁₂, the amino acid sequence of which is presented in SEQ ID NO:61, assuming an open reading frame spanning from nucleotide 1 through nucleotide 336 of SEQ ID NO:60.

Comparison of nucleic acid sequence SEQ ID NO:60 with nucleic acid molecules reported in GenBank indicates that SEQ ID NO:60 showed the most homology, i.e. 67.2% identity, with a human CD40 gene. Comparison of amino acid sequence SEQ ID NO:61 with amino acid sequences reported in GenBank indicates that SEQ ID NO:61 showed the most homology, i.e. about 54.4% identity, with a human CD40 protein. Sequence analysis was performed using the GCG GAP program as described above.

Example 4

This example describes the isolation and sequencing of certain canine CD154 (canine CD40 ligand) and feline CD154 (feline CD40 ligand) nucleic acid molecules and proteins of the present invention.

A. Canine CD154 (CD40 Ligand) Nucleic Acid Molecules and Proteins

The following describes the isolation and sequencing of certain cDNA nucleic acid molecules encoding certain canine CD154 (CD40 ligand) proteins of the present invention.

A canine CD154 nucleic acid molecule of the present invention was produced by PCR amplification as follows. A 390-nucleotide canine CD40 nucleic acid molecule, denoted nCaCD154₃₉₀, was amplified from a canine PBMC cDNA library, prepared as described in Example 1, using two degenerate oligonucleotide primers designed in accordance with human CD154 gene sequences available in GenBank; sense primer, 5' CCTCAAATTG CGGCACATGT C 3', denoted herein as SEQ ID NO:130; and antisense primer, 5' CTGTTCAGAG TTTGAGTAAG CC 3', denoted herein as SEQ ID NO:131. PCR conditions used for canine CD154 cDNA amplification were standard conditions for PCR amplification (Sambrook, et al., *ibid.*). The resulting PCR product, i.e., nCaCD154₃₉₀, with a coding strand represented by SEQ ID NO:63, was radiolabeled using standard techniques and used to screen the canine PBMC cDNA library, under the hybridization conditions described in Example 3. A clone containing a 1878-nucleotide canine CD154 nucleic acid molecule, denoted nCaCD154₁₈₇₈, was obtained. The nucleic acid sequence of the coding strand of nCaCD154₁₈₇₈ is represented as SEQ ID NO:64. The reverse complement of SEQ ID NO:64 is referred to herein as SEQ ID NO:66. Translation of SEQ ID NO:64 suggests that nucleic acid molecule of nCaCD154₁₈₇₈ encodes a CD154 protein of 260 amino acids, denoted herein as PCaCD154₂₆₀, the amino acid sequence of which is presented in SEQ ID NO:65, assuming an open reading frame having an initiation codon spanning from nucleotide 284 through nucleotide 286 of SEQ ID NO:64 and a stop codon spanning from nucleotide 1064 through nucleotide 1066 of SEQ ID NO:64. The coding region encoding PCaCD154₂₆₀, not including the termination codon, is presented herein as nCaCD154₇₈₀, which has the nucleotide sequence SEQ ID

NO:67 (the coding strand) and SEQ ID NO:68 (the complementary strand).

A putative signal/membrane anchor sequence coding region extends from nucleotide 284 through nucleotide 430 of SEQ ID NO:64. The proposed soluble CD154 protein, denoted herein as PCaCD154₂₁₁, represented by SEQ ID NO:70, contains about 211 amino acids, extending from residue 50 through residue 260 of SEQ ID NO:65. The nucleotide sequence encoding PCaCD154₂₁₁, which extends from nucleotide 431 through nucleotide 1063 of SEQ ID NO:64, is denoted herein as nucleic acid molecule nCaCD154₆₃₃, represented by SEQ ID NO:69 (the coding strand) and SEQ ID NO:71 (the complement strand).

Sequence analysis was performed with DNAsis™ using the alignment settings of: gap penalty set at 5; number of top diagonals set at 5; fixed gap penalty set at 10; k-tuple set at 2; window size set at 5 and floating gap penalty set at 10. At the amino acid level, PCaCD154₂₆₀ shares 78.0%, 77.6%, and 67.6% identity with the CD154 proteins of human, bovine, and mouse, respectively (Graf et al., Eur. J. Immunol., vol. 22, pp. 3191–3194, 1992; Hollenbaugh, et al., EMBO J., vol. 11:4313–4321, 1992; Gauchat et al., FEBS lett., vol. 315, pp. 259–266, 1993; Mertens et al., Immunogenetics, vol. 42, pp. 430–431; Armitage et al., Nature, vol. 357, pp. 80–82; 1992). At the nucleotide level, nCaCD154₁₈₇₈ shares 81.1%, 81.5%, and 74.4% identity with the sequences of human, bovine, and mouse CD154 cDNAs, respectively.

B. Feline CD154 (CD40 ligand) Nucleic Acid Molecules and Proteins

This example describes the isolation and sequencing of certain nucleic acid molecules encoding certain feline CD154 (CD40 ligand) proteins of the present invention.

A feline CD154 nucleic acid molecule was isolated by PCR amplification from a feline PBMC cDNA library, prepared as described in Example 2, using Amplitaq DNA polymerase (available from PE Applied Biosystems Inc, Foster City, Calif.) under the following PCR protocol; one initial denaturation step at 95° C. for 5 minutes; then 40 cycles of the following: 94° C. for 45 seconds, then 48° C. for 45 seconds, then 72° C. for 120 seconds; followed by a final extension at 72° C. for 7 minutes. The forward and reverse primers used were based on human CD154 cDNA sequences outside the open reading frame in the 5' and 3' untranslated regions, respectively, so that the open reading frame in the PCR product contained only feline sequences. The sequence of the forward primer was 5' GAAGATACCA TTTCAACTTT AACACAGC 3' SEQ ID NO:132, and that of the reverse primer was 5' TGCTGTATTG TGAA-GACTCC CAGC 3' SEQ ID NO:133. PCR products were cloned into the TA cloning vector (available from Invitrogen Corporation, Carlsbad, Calif.), and the resulting clones were sequenced using an ABI Prism™ Model 377 Automatic DNA Sequencer (available from PE Applied Biosystems Inc.). DNA sequencing reactions were performed using Prism™ dRhodamine Terminator Cycle Sequencing Ready Reaction kits (available from PE Applied Biosystems Inc.).

The PCR product was sequenced and found to contain 885 nucleotides, and is denoted as nFeCD154₈₈₅. The nucleotide sequence of the coding strand of nFeCD154₈₈₅ is represented herein as SEQ ID NO:72, and its complement is denoted SEQ ID NO:74. Translation of the open reading frame in SEQ ID NO:72 suggests that nFeCD154₈₈₅ encodes a protein containing 260 amino acids, referred to herein as PFeCD154₂₆₀, the amino acid sequence of which

is presented as SEQ ID NO:73, assuming an open reading frame in which the first codon spans from nucleotide 29 through nucleotide 31 of SEQ ID NO:72, and the stop codon spans from nucleotide 809 through nucleotide 811 of SEQ ID NO:72. The encoded protein has a predicted molecular weight of 28.6 kDa for the precursor protein and 27.2 kDa for the mature protein. The coding region encoding PFeCD154₂₆₀, not including the termination codon, is presented herein as nFeCD154₇₈₀, which has the nucleotide sequence SEQ ID NO:75 (the coding strand) and SEQ ID NO:76 (the complementary strand).

A putative signal/membrane anchor sequence coding region extends from nucleotide 29 through nucleotide 175 of SEQ ID NO:72. The proposed soluble feline CD154 protein, denoted herein as PFeCD154₂₁₁, represented by SEQ ID NO:78, contains about 211 amino acids, extending from residue 50 through residue 260 of SEQ ID NO:73. The nucleotide sequence encoding PFeCD154₂₁₁, denoted herein as nFeCD154₆₃₃ which extends from nucleotide 176 through nucleotide 808 of SEQ ID NO:72, is represented herein by SEQ ID NO:77 (the coding strand) and SEQ ID NO:79 (the complementary strand).

Comparison of feline CD154 nucleotide and amino acid sequences with those of other species published in GenBank reveals that the feline CD154 nucleotide sequence SEQ ID NO:75 is 86%, 88% and 75% identical to the human, bovine and murine CD154 gene sequences, respectively (Genbank accession number L07414, Z48469 and X56453 respectively). At the amino acid sequence level, SEQ ID NO:73 is 81%, 82%, and 67% identical to the human, bovine and murine CD154 amino acid sequences, respectively. Hydrophobicity analysis of feline CD154 proteins results in a pattern similar to those of human, bovine and murine CD154 proteins. A putative N-glycosylation site was identified at position 239 in PFeCD154₂₆₀ that is conserved in the human, bovine and murine amino acid sequences. Five cysteine residues are present in the feline CD154 protein sequence SEQ ID NO:73. Four of the five residues, located at positions 72, 84, 177 and 217 of PFeCD154₂₆₀, are conserved in all four species and are likely involved in disulfide bond formation. The cysteine residue located at position 193 of PFeCD154₂₆₀ is present in all but the murine sequence.

Example 5

This example describes the isolation and sequencing of certain canine IL-5 nucleic acid molecules and proteins of the present invention. This example also describes expression of canine IL-5 in a Pichia expression system.

A. Isolation and Sequencing of Canine IL-5 Nucleic Acid Molecules and Proteins

A canine IL-5 cDNA nucleic acid molecule encoding a canine IL-5 protein was isolated by PCR amplification from a canine PBMC cDNA library (prepared as described in Example 1) using PCR conditions as described in Example 4B and the following primers. Degenerate oligonucleotide primers were designed in accordance with conserved regions of human and cat IL-5 gene sequences available in GenBank; sense primer, 5' ATGCACTTTC TTGCC 3', denoted herein as SEQ ID NO:134; antisense primer 1, 5' CTGGAG-GAAA AKACTTCRAT GATTCTGATA TCTGAAATAT AT 3', denoted herein as SEQ ID NO:135; and antisense primer 2, 5' CTGACYCTTK STTGGSCCTC ATTCTCA 3', denoted herein as SEQ ID NO:136, where K was G or T, R was either A or G, S was either G or C, and Y was either T or C.

An about 610-nucleotide canine IL-5 nucleic acid molecule, denoted nCaIL-5₆₁₀, was obtained using primers having SEQ ID NO:134 and SEQ ID NO:135, respectively. The sequence of the coding strand of nCaIL-5₆₁₀ is represented herein as SEQ ID NO:80. The reverse complement of SEQ ID NO:80 is referred to herein as SEQ ID NO:82. Translation of SEQ ID NO:80 suggests that nucleic acid molecule nCaIL-5₆₁₀ encodes an IL-5 protein of 134 amino acids, denoted herein as PCaIL-5₁₃₄, the amino acid sequence of which is presented in SEQ ID NO:81, assuming an open reading frame having an initiation codon spanning from nucleotide 29 through nucleotide 31 of SEQ ID NO:80 and a stop codon spanning from nucleotide 431 through nucleotide 433 of SEQ ID NO:80. The coding region encoding PCaIL-13₁₃₄, not including the termination codon, is presented herein as nCaIL-5₄₀₂, which has the nucleotide sequence SEQ ID NO:83 (the coding strand) and SEQ ID NO:84 (the complementary strand).

An about 488-nucleotide fragment, denoted herein as nCaIL-5₄₈₈, isolated by PCR with primers having SEQ ID NO:134 and SEQ ID NO:136, respectively, corresponds to nucleotide 1 through nucleotide 488 of nCaIL-5₆₁₀.

A putative signal sequence coding region extends from nucleotide 29 through nucleotide 85 of SEQ ID NO:80. The proposed mature protein, denoted herein as PCaIL-5₁₁₅, represented by SEQ ID NO:86, contains about 115 amino acids, extending from residue 20 through residue 134 of SEQ ID NO:81. The nucleotide sequence encoding PCaIL-5₁₁₅, which extends from nucleotide 86 through nucleotide 430 of SEQ ID NO:80, is denoted herein as nucleic acid molecule nCaIL-5₃₄₅, represented by SEQ ID NO:85 (coding strand) and SEQ ID NO:87 (the complement strand).

Sequence analysis was performed with DNAsis™ using the alignment settings of: gap penalty set at 5; number of top diagonals set at 5; fixed gap penalty set at 10; k-tuple set at 2; window size set at 5 and floating gap penalty set at 10. At the amino acid level, PCaIL-5₁₃₄ shared 82.8% and 57.4% identity with feline and human IL-5 proteins, respectively (Padrid et al., *Am. J. Vet. Res.*, vol. 59, pp. 1263-1269, 1998; Azuma et al., *Nucleic Acids Res.*, vol. 14, pp. 9149-9158, 1986). At the nucleotide level, nCaIL-5₆₁₀ shared 81.7% and 75% identity with the cDNA sequences of the feline and human IL-5, respectively.

B. Expression of Canine IL-5 in Pichia

This example describes the expression in Pichia of a canine IL-5 cDNA fragment, namely a canine IL-5 nucleic acid molecule denoted nCaIL-5₃₄₈, the coding strand of which consists of nucleotides 86-433 of SEQ ID NO:80, and as such, encodes a predicted mature canine IL-5 protein having SEQ ID NO:86. Nucleic acid molecule nCaIL-5₃₄₈, was PCR amplified from nCaIL-5₆₁₀ using sense primer 5' GGGCTCGAGA AAAGATTTGC TGTAAGAAAT CCCATG 3' denoted herein as SEQ ID NO:137, with nucleotides 16-36 corresponding to nucleotides 86-106 of SEQ ID NO:80; and antisense primer 5' CCCGCGGCCG CTCAACTTTC CGGTGTCCAC TC 3', denoted herein as SEQ ID NO:138, with nucleotides 12-32 corresponding to the reverse complement of nucleotides 413-433 of SEQ ID NO:80. To facilitate cloning, an XhoI site (shown in bold) was added to the sense primer and a NotI site (shown in bold) was added to the antisense primer. The PCR-amplified fragment was digested with restriction endonucleases XhoI and NotI, gel purified and ligated into pPICZαA plasmid vector, available from Invitrogen, that has been digested by Aho I and Not I and gel purified, to produce recombinant

molecule pPICZαA-nCaIL-5₃₄₈. The insert in the recombinant molecule was verified by DNA sequencing. The recombinant molecule was used to transform Pichia pastoris strain X-33 by electroporation to produce recombinant cell Pichia-pPICZαA-nCaIL-5₃₄₈. Recombinant cell Pichia-pPICZαA-nCaIL-5₃₄₈ was cultured using techniques known to those skilled in the art and IL-5 expression was induced with methanol. The supernatant was recovered and submitted to SDS-PAGE. Silver staining of the resultant gel indicated a band of about 18 kDa only seen in the supernatant of Pichia transformed with recombinant molecule pPICZαA-nCaIL-5₃₄₈.

Example 6

This example describes the isolation and sequencing of certain canine IL-13 nucleic acid molecules and proteins of the present invention. This example also describes expression of canine IL-13 in E. coli.

A. Isolation and Sequencing of Canine IL-13 Nucleic Acid Molecules and Proteins

A canine IL-13 cDNA nucleic acid molecule encoding a canine IL-13 protein was isolated by PCR amplification from a canine PBMC cDNA library (prepared as described in Example 1) using the following primers and PCR conditions: Degenerate oligonucleotide primers were designed in accordance with conserved regions of human and cat IL-5 gene sequences available in GenBank; sense primer, 5' GTCMTKGCTC TYRCTTGCCT TGG 3', denoted herein as SEQ ID NO:139; antisense primer 1, 5' AAAGTGGGCY ACYTCGATTT TGG 3', denoted herein as SEQ ID NO:140; antisense primer 2, 5' GTGATGTTGM YCAGCTCCTC 3', denoted herein as SEQ ID NO:141, where M was either A or C, K was G or T, R was either A or G, S was either G or C, and Y was either T or C. PCR conditions used were as follows: One initial denaturation step at 95° C. for 3 minutes; then 38 cycles of the following: 94° C. for 30 seconds, 51.8° C. for 45 seconds, then 72° C. for 105 seconds; then a final extension at 72° C. for 5 minutes.

An about 272-nucleotide canine IL-13 nucleic acid molecule, denoted nCaIL-13₂₇₂ and having a coding strand represented by SEQ ID NO:89, was PCR amplified using primers having nucleic acid sequences of SEQ ID NO:139 and SEQ ID NO:140, respectively. An about 166-nucleotide canine IL-13 nucleic acid molecule, denoted nCaIL-13₁₆₆ and having a coding strand represented by SEQ ID NO:88, was isolated using primers having nucleic acid sequences of SEQ ID NO:142 (see Example 2B) and SEQ ID NO:141, respectively. Nucleic acid molecules nCaIL-13₂₇₂ and nCaIL-13₁₆₆ form an overlapping composite fragment of 383 nucleotides, denoted nCaIL-13₃₈₃. Two canine IL-13 specific primers (i.e., sense primer, 5' ATGGCGCTCT GGT-TGACTGT 3', denoted herein as SEQ ID NO:143; and antisense primer, 5' GGCTTTTGAG AGCACAGTGC 3', denoted herein as SEQ ID NO:144) were derived from nCaIL-13₃₈₃ and were used to isolate a 278-nucleotide fragment, denoted nCaIL-13₂₇₈ and having a coding strand represented by SEQ ID NO:90. Nucleic acid molecule nCaIL-13₂₇₈ was radiolabeled and used to screen the canine PBMC cDNA library under the following hybridization conditions: hybridization took place in 6×SSC, 5×Denhardt's solution, 0.5% SDS, 100 µg/ml single stranded DNA, 100 µg/ml tRNA, for 36 hours at 60° C.; the final wash solution was 0.1% SDS, 0.125×SSC at 60° C. for 30 minutes. Two clones were selected, namely clone 80 and clone 78.

Sequence analysis of Clone 80 indicated that the clone includes an about 1302-nucleotide canine IL-13 nucleic acid molecule referred to herein as nCaIL-13₁₃₀₂, the coding strand of which was shown to have nucleic acid sequence SEQ ID NO:91. The reverse complement of SEQ ID NO:91 is referred to herein as SEQ ID NO:93. Translation of SEQ ID NO:91 suggests that nucleic acid molecule nCaIL-13₁₃₀₂ encodes an IL-13 protein of 131 amino acids, denoted herein as PCaIL-13₁₃₁, the amino acid sequence of which is presented in SEQ ID NO:92, assuming an open reading frame having an initiation codon spanning from nucleotide 52 through nucleotide 54 of SEQ ID NO:91 and a stop codon spanning from nucleotide 445 through nucleotide 447 of SEQ ID NO:91. The coding region encoding PCaIL-13₁₃₁, not including the termination codon, is presented herein as nCaIL-13₃₉₃, which has the nucleotide sequence SEQ ID NO:94 (the coding strand) and SEQ ID NO:95 (the complementary strand).

A putative signal sequence coding region extends from nucleotide 52 to nucleotide 111 of SEQ ID NO:91. The proposed mature protein, denoted herein as PCaIL-13₁₁₁, represented by SEQ ID NO:97, contains 111 amino acids, extending from residue 21 through residue 131 of SEQ ID NO:91. The nucleotide sequence encoding PCaIL-13₁₁₁, which extends from nucleotide 112 through nucleotide 444 of SEQ ID NO:91, is denoted herein as nucleic acid molecule nCaIL-13₃₃₃, represented by SEQ ID NO:96 (coding strand) and SEQ ID NO:98 (the complement strand).

Sequence analysis of Clone 78 indicated that the clone includes an about 1269-nucleotide canine IL-13 nucleic acid molecule referred to herein as nCaIL-13₁₂₆₉, the coding strand of which was shown to have nucleic acid sequence SEQ ID NO:99. The reverse complement of SEQ ID NO:99 is referred to herein as SEQ ID NO:101. Translation of SEQ ID NO:99 suggests that nucleic acid molecule nCaIL-13₁₂₆₉ encodes an IL-13 protein of 130 amino acids, denoted herein as PCaIL-13₁₃₀, the amino acid sequence of which is presented in SEQ ID NO:100, assuming an open reading frame having an initiation codon spanning from nucleotide 57 through nucleotide 59 of SEQ ID NO:99 and a stop codon spanning from nucleotide 447 through nucleotide 449 of SEQ ID NO:99. The coding region encoding PCaIL-13₁₃₀, not including the termination codon, is represented herein as nCaIL-13₃₉₀, which has the nucleotide sequence SEQ ID NO:102 (the coding strand) and SEQ ID NO:103 (the complementary strand). PCaIL-13₁₃₀ is missing one amino acid compared to PCaIL-13₁₃₃, namely amino acid position Q97 of PCaIL-13₁₃₃.

A putative signal sequence coding region extends from nucleotide 57 to nucleotide 116 of SEQ ID NO:99. The proposed mature protein, denoted herein as PCaIL-13₁₁₀, represented by SEQ ID NO:105, contains 110 amino acids, extending from residue 21 through residue 130 of SEQ ID NO:100. The nucleotide sequence encoding PCaIL-13₁₁₀, which extends from nucleotide 117 through nucleotide 446 of SEQ ID NO:99, is denoted herein as nucleic acid molecule nCaIL-13₃₃₀, represented by SEQ ID NO:104 (coding strand) and SEQ ID NO:106 (the complement strand).

Sequence analysis was performed with DNAsis™ using the alignment settings of: gap penalty set at 5; number of top diagonals set at 5; fixed gap penalty set at 10; k-tuple set at 2; window size set at 5 and floating gap penalty set at 10. At the amino acid level, PCaIL-13₁₃₁ shared 61.7%, 39.6%, 36.6% identity with the IL-13 proteins of human, mouse, and rat (Brown et al., J. Immunol., vol. 142, pp 679–687, 1989; Lakkis et al., Biochem. Biophys. Res. Commun., Vol. 197, pp. 612–618, 1993; McKenzie et al., Proc. Natl Acad.

Sci. USA, vol. 90, pp. 3735–3739, 1993; Minty et al., Nature, vol. 362, pp. 248–250, 1993), respectively. At the nucleotide level, nCaIL-13₁₃₀₂ shared 56.0%, 57.1%, and 45.9% identity with the sequences of human, rat, and mouse IL-13 cDNAs, respectively.

B. Expression of Canine IL-13 in E. coli

This example describes the expression in E. coli of a canine IL-13 cDNA fragment, namely a canine IL-13 nucleic acid molecule denoted nCaIL-13₃₃₆, the coding strand of which consists of nucleotides 112–447 of SEQ ID NO:91, and as such, encodes a predicted mature canine IL-13 protein having SEQ ID NO:97. Nucleic acid molecule nCaIL-13₃₃₆ was PCR amplified from nCaIL-13₁₃₀₂ using sense primer 5' CCCCATATGAGCCCTGTGAC TCCCTC-CCC 3' denoted herein as SEQ ID: 145, with nucleotides 10–29 corresponding to nucleotides 112–1131 of SEQ ID NO:91; and antisense primer 5' GGGAATTCT CATCTGAAAT TTCCATGGCG 3', denoted herein as SEQ ID NO:146, with nucleotides 10–30 corresponding to the reverse complement of nucleotides 427–447 of SEQ ID NO:91. To facilitate cloning, an NdeI site (shown in bold) was added to the sense primer and an EcoRI site (shown in bold) was added to the antisense primer. The resulting PCR fragment was digested with restriction endonucleases NdeI and EcoRI, gel purified and ligated into λcro plasmid vector, the production of which is described in U.S. Pat. No. 5,569,603 by Tripp et al., issued Oct. 29, 1996, that had been digested by NdeI and EcoRI and gel purified to produce recombinant molecule pλcro-nCaIL-13₃₃₆. The insert in the recombinant molecule was verified by DNA sequencing. Recombinant molecule pλcro-nCaIL-13₃₃₆ was used to transform E. coli strain HCE101 (BL21), thereby producing BL21-pλcro-nCaIL-13₃₃₆. PCaIL-13₁₁₁ was produced under conditions as described in U.S. Pat. No. 5,569,603, *ibid.*, protein expression being induced by switching the fermentation temperature from 32° C. to 42° C. SDS-PAGE and Commassie blue staining analysis indicated that a band of about 11 kD was only produced by induced BL21-pλcro-nCaIL-13₃₃₆ recombinant cells. The 11-kD band showed a positive reaction with a rabbit polyclonal antibody against human IL-13 (available from PeproTech Inc, Rocky Hill, N.J.), indicating expression of canine IL-13 in E. coli.

Example 7

This example describes the isolation and sequencing of feline interferon alpha nucleic acid molecules and proteins of the present invention.

Feline IFN-alpha nucleic acid molecules were PCR amplified from a feline cDNA library as follows. Total RNA was isolated from cat (kitten) mesenteric lymph node cells stimulated with PMA (phorbol myristate acetate) for 48 hours using Tri Reagent™ (available from Molecular Research Center, Cincinnati, Ohio). cDNA was made from the RNA using the cDNA synthesis kit containing Ready to Go -You Prime First-Strand Beads™ (available from Amersham Pharmacia Biotech, Piscataway, N.J.). An aliquot of this cDNA was used as a template to isolate a feline IFN-alpha nucleic acid molecule by PCR amplification using Amplitaq DNA polymerase™ (available from PE Applied Biosystems Inc, Foster City, Calif.) and the following primers and conditions. The sequence of the forward primer was 5' ATGGCGCTGC CCTCTTCCTT CTTG 3' (SEQ ID NO:143), and that of the reverse primer was 5' TCATTCTCG CTCCTTAATC TTTTCTGC 3' (SEQ ID NO:148). The following PCR protocol was used: one initial

denaturation step at 95° C. for 5 minutes; then 43 cycles of the following: 94° C. for 45 seconds, then 47° C. for 45 seconds, then 72° C. for 120 seconds; followed by a final extension at 72° C. for 7 minutes. PCR products were cloned into the TA cloning vector (available from Invitrogen Corporation) and the clones were sequenced using an ABI Prism™ Model 377 Automatic DNA Sequencer (available from PE Applied Biosystems Inc.). DNA sequencing reactions were performed using Prism™ dRhodamine Terminator Cycle Sequencing Ready Reaction kits (available from PE Applied Biosystems Inc.).

Two PCR products were generated and sequenced. Both contained 570 nucleotides (including the termination codons), and are distinguished as Clone #2 and Clone #3 as there were differences in the sequences of the clones.

Clone #2 includes a feline IFN-alpha nucleic acid molecule that is represented herein as nFeIFN α_{567a} , the coding strand of which was shown to have a nucleic acid sequence denoted herein as SEQ ID NO:107. The complement of SEQ ID NO:107 is represented herein by SEQ ID NO:109. Translation of SEQ ID NO:107 suggests that nFeIFN α_{567a} encodes a protein containing 189 amino acids, referred to herein as PFeIFN α_{189a} , with an amino acid sequence denoted SEQ ID NO:108. The open reading frame of SEQ ID NO:107 is assumed to be the following: the first codon spans from nucleotide 1 through nucleotide 3 and the last codon before the stop codon spans from nucleotide 565 to nucleotide 567 of SEQ ID NO:107. The encoded protein has a predicted molecular weight of 21 kDa. The putative signal peptide cleavage site occurs between amino acid positions 23 and 24, based on homology with the human and canine interferon-alpha proteins. The proposed mature protein (i.e. feline IFN α protein from which the signal sequence has been cleaved), denoted herein as PFeIFN α_{166a} , contains about 166 amino acids, extending from residue 24 to residue 166 of SEQ ID NO:108; the amino acid sequence is denoted herein as SEQ ID NO:114. The nucleic acid molecule encoding PFeIFN α_{166a} is denoted herein as nFeIFN α_{498a} , which is represented by SEQ ID NO:113, with a complementary sequence represented by SEQ ID NO:115. A putative N-glycosylation site and an interferon alpha-beta-delta family signature motif are present at amino acid positions 102 and 145, respectively, of PFeIFN α_{189a} .

Clone #3 includes a feline IFN-alpha nucleic acid molecule that is represented herein as nFeIFN α_{567b} , the coding strand of which was shown to have a nucleic acid sequence denoted herein as SEQ ID NO:110. The complement of SEQ ID NO:110 is represented herein by SEQ ID NO:112. Translation of SEQ ID NO:110 suggests that nFeIFN α_{567b} encodes a protein containing 189 amino acids, referred to herein as PFeIFN α_{189b} , with an amino acid sequence denoted SEQ ID NO:111. The open reading frame of SEQ ID NO:110 is assumed to be the following: the first codon spans from nucleotide 1 through nucleotide 3 and the last codon before the stop codon spans from nucleotide 565 through nucleotide 567 of SEQ ID NO:110. The encoded protein has a predicted molecular weight of 21 kDa. The putative signal peptide cleavage site occurs between amino acid positions 23 and 24, based on homology with the human and canine interferon-alpha proteins. The proposed mature protein (i.e. feline IFN α protein from which the signal sequence has been cleaved), denoted herein as PFeIFN α_{166b} , contains about 166 amino acids, extending from residue 24 to residue 166 of SEQ ID NO:111, the amino acid sequence is denoted herein as SEQ ID NO:117. The nucleic acid molecule encoding PFeIFN α_{166b} is denoted herein as nFeIFN α_{498b} , which is represented by

SEQ ID NO:116, with a complementary sequence represented by SEQ ID NO:118. A putative N-glycosylation site and an interferon alpha-beta-delta family signature motif are present at amino acid positions 102 and 145, respectively, of PFeIFN α_{189b} .

The differences between the proteins encoded by SEQ ID NO:107 and SEQ ID NO:110 are detailed as follows: The amino acid residues at position 56 in SEQ ID NO:108 (i.e., the protein encoded by SEQ ID NO:107) and at position 56 in SEQ ID NO:111 (i.e., the protein encoded by SEQ ID NO:110) are both arginines, although the corresponding codons in SEQ ID NO:107 and SEQ ID NO:110 are AGA and AGG, respectively. The amino acid residues at position 74 in SEQ ID NO:108 and at position 74 in SEQ ID NO:111 are both alanines, although the corresponding codons in SEQ ID NO:107 and SEQ ID NO:110 are GCC and GCT, respectively. The amino acid residue at position 86 in SEQ ID NO:108 is lysine, encoded by AAG in SEQ ID NO:107, whereas the amino acid residue at position 86 in SEQ ID NO:111 is glutamic acid, encoded by GAG in SEQ ID NO:110. The amino acid residue at position 125 in SEQ ID NO:108 is methionine, encoded by CTG in SEQ ID NO:107, whereas the amino acid residue at position 125 in SEQ ID NO:111 is valine, encoded by GTG in SEQ ID NO:110. The amino acid residue at position 141 in SEQ ID NO:108 is isovaline, encoded by ATC in SEQ ID NO:107, whereas the amino acid residue at position 141 in SEQ ID NO:111 is leucine, encoded by CTC in SEQ ID NO:110.

Feline IFN-alpha proteins of the present invention PFeIFN α_{189a} and PFeIFN α_{189b} are five amino acids shorter than the GenBank entry for feline IFN-alpha, accession #E02521. In addition, there are 3 non-conservative and 2 conservative changes at the amino acid level between this GenBank entry and SEQ ID NO:108 (PFeIFN α_{189a}) as well as 4 non-conservative and 3 conservative changes at the amino acid level between this GenBank entry and SEQ ID NO:111 (PFeIFN α_{189b}). The lengths of SEQ ID NO:108 and SEQ ID NO:111, when compared with those of IFN-alpha proteins of other species, are two amino acids longer than published canine interferon-alpha subtype 1, 2 and 3 sequences, two amino acids longer than published human interferon-alpha type I,B,D, F, and J, sequences, three amino acids longer than the published human interferon-alpha sequence type A sequence and two amino acids longer than published murine interferon-alpha type B, 8, 7, 11, and 19 sequences.

Example 8

This example describes the isolation and sequencing of feline granulocyte-macrophage colony-stimulating factor (GMCSF) nucleic acid molecules and proteins of the present invention. This example also describes expression of a feline GMCSF protein of the present invention.

Nucleic acid molecules encoding feline GMCSF were isolated as follows. A cDNA library was prepared from feline PBMCs stimulated with Con A for 12 hours, as previously described in Example 2. An aliquot of this library was used as a template to amplify feline GMCSF nucleic acid molecules by PCR using Amplitaq DNA polymerase™ (PE Applied Biosystems Inc, Foster City, Calif.) and the following primers and conditions. The sequence of the forward primer was 5' CAGGATCCA CCATGTGGCT GCA-GAACCTG CTTTTC 3' (SEQ ID NO:149), and that of the reverse primer was 5' TTACTTCTGG TCTGGTCCCC AGCAGTCAAA GGGGTGTTA AACAGAAAAT 3' (SEQ ID NO:150). The following PCR protocol was used:

one initial denaturation step at 95° C. for 5 minutes; then 35 cycles of the following: 94° C. for 30 seconds, then 50° C. for 30 seconds, then 72° C. for 90 seconds; followed by a final extension at 72° C. for 7 minutes. PCR products were cloned into the CMV-Intron A vector and the clones were sequenced as described in Example 7.

A PCR product was isolated, referred to herein as nFeGMCSF₄₄₄, the coding strand of which is represented herein as SEQ ID NO:119, and its complement is denoted SEQ ID NO:121. Translation of the open reading frame in SEQ ID NO:119 suggests that nucleic acid molecule nFeGMCSF₄₄₄ encodes a protein containing 144 amino acids, referred to herein as PFeGMCSF₁₄₄, with an amino acid sequence denoted SEQ ID NO:120, assuming an open reading frame in which the first codon spans from nucleotide 10 through nucleotide 12 of SEQ ID NO:119, and the stop codon spans from nucleotide 442 through nucleotide 444 of SEQ ID NO:121. The encoded protein has a predicted molecular weight of 16 kDa. The coding region encoding PFeGMCSF₁₄₄ is presented herein as nFeGMCSF₄₃₂ which has the nucleotide sequence SEQ ID NO:122 (the coding strand) and SEQ ID NO:123 (the complementary strand). A putative signal peptide cleavage site is between amino acid positions 17 and 18, based on homology with human, mouse and ovine GMCSF proteins. The nucleic acid molecule encoding the proposed mature protein is denoted as nFeGMCSF₃₈₁ and has a nucleotide sequence represented herein as SEQ ID NO:124 and a complementary sequence represented herein as SEQ ID NO:126. The amino acid sequence

of the putative mature protein, referred to herein as PFeGMCSF₁₂₇, has an amino acid sequence represented herein as SEQ ID NO:125. The number of amino acids in the feline GMCSF protein is the same compared to human, porcine, ovine and canine GMCSF proteins. The feline GMCSF protein is one amino acid longer than bovine GMCSF and three amino acid longer than murine GMCSF.

The deduced amino acid sequence of the full-length feline GMCSF protein of the present invention has four non-conservative changes and one conservative change compared to a GenBank entry for feline GMCSF (accession #AF053007). Amino acids asparagine, methionine, threonine, and lysine at positions 10, 36, 56 and 126 of the GenBank entry have been changed to glycine, isoleucine, alanine and asparagine, respectively, in PFeGMCSF₁₄₄, PFeGMCSF₁₄₄, containing the above-noted amino acid substitutions, appears to have GMCSF activity, as demonstrated by an experiment in which supernatant collected from Chinese Hamster Ovary (CHO) cells that were transiently transfected with a recombinant molecule encoding a feline GMCSF protein of the present invention was able to induce proliferation of TF-1 cells.

While various embodiments of the present invention have been described in detail, it is apparent that modifications and adaptations of those embodiments will occur to those skilled in the art. It is to be expressly understood, however, that such modifications and adaptations are within the scope of the present invention, as set forth in the following claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 154

<210> SEQ ID NO 1
<211> LENGTH: 549
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (43)..(438)

<400> SEQUENCE: 1

```

ggcagcaggt ctgctattgt cactgcaaat agagatctat ta atg ggt ctc acc      54
                                     Met Gly Leu Thr
                                     1

tcc caa ctg att cca act ctg gtc tgc tta cta gca ctc acc agc acc      102
Ser Gln Leu Ile Pro Thr Leu Val Cys Leu Leu Ala Leu Thr Ser Thr
  5              10              15              20

ttt gtc cac gga cat aac ttc aat att act att aaa gag atc atc aaa      150
Phe Val His Gly His Asn Phe Asn Ile Thr Ile Lys Glu Ile Ile Lys
  25              30              35

atg ttg aac atc ctc aca gcg aga aac gac tcg tgc atg gag ctg act      198
Met Leu Asn Ile Leu Thr Ala Arg Asn Asp Ser Cys Met Glu Leu Thr
  40              45              50

gtc aag gac gtc ttc act gct cca aag aac aca agc gat aag gaa atc      246
Val Lys Asp Val Phe Thr Ala Pro Lys Asn Thr Ser Asp Lys Glu Ile
  55              60              65

ttc tgc aga gct gct act gta ctg cgg cag atc tat aca cac aac tgc      294
Phe Cys Arg Ala Ala Thr Val Leu Arg Gln Ile Tyr Thr His Asn Cys
  70              75              80

tcc aac aga tat ctc aga gga ctc tac agg aac ctc agc agc atg gca      342
Ser Asn Arg Tyr Leu Arg Gly Leu Tyr Arg Asn Leu Ser Ser Met Ala

```

-continued

85	90	95	100	
aac aag acc tgt tct atg aat gaa atc aag aag agt aca ctg aaa gac				390
Asn Lys Thr Cys Ser Met Asn Glu Ile Lys Lys Ser Thr Leu Lys Asp				
	105	110	115	
ttc ttg gaa agg cta aaa gtg atc atg cag aag aaa tac tac agg cat				438
Phe Leu Glu Arg Leu Lys Val Ile Met Gln Lys Lys Tyr Tyr Arg His				
	120	125	130	
tgaagctgaa tattttaatt tatgagtttt taaatagctt tattttaaaa atatttatat				498
atttataaca taataaaata aaatatatat agaaaaaaaa aaaaaaaaaa a				549

<210> SEQ ID NO 2
 <211> LENGTH: 132
 <212> TYPE: PRT
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 2

Met Gly Leu Thr Ser Gln Leu Ile Pro Thr Leu Val Cys Leu Leu Ala			
1	5	10	15
Leu Thr Ser Thr Phe Val His Gly His Asn Phe Asn Ile Thr Ile Lys			
	20	25	30
Glu Ile Ile Lys Met Leu Asn Ile Leu Thr Ala Arg Asn Asp Ser Cys			
	35	40	45
Met Glu Leu Thr Val Lys Asp Val Phe Thr Ala Pro Lys Asn Thr Ser			
	50	55	60
Asp Lys Glu Ile Phe Cys Arg Ala Ala Thr Val Leu Arg Gln Ile Tyr			
	65	70	75
Thr His Asn Cys Ser Asn Arg Tyr Leu Arg Gly Leu Tyr Arg Asn Leu			
	85	90	95
Ser Ser Met Ala Asn Lys Thr Cys Ser Met Asn Glu Ile Lys Lys Ser			
	100	105	110
Thr Leu Lys Asp Phe Leu Glu Arg Leu Lys Val Ile Met Gln Lys Lys			
	115	120	125
Tyr Tyr Arg His			
	130		

<210> SEQ ID NO 3
 <211> LENGTH: 549
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 3

tttttttttt tttttttttt tatatatatt ttatttttatt atgttataaa tatataaata	60
ttttttaaatt aaagctattt aaaaactcat aaattaaaat attcagcttc aatgcctgta	120
gtatttccttc tgcattgatca cttttagcct ttccaagaag tctttcagtg tactcttctt	180
gatttcattc atagaacagg tcttggttgc catgctgctg aggttcctgt agagtcctct	240
gagatatctg ttggagcagt tgtgtgtata gatctgccgc agtacagtag cagctctgca	300
gaagatttcc ttatcgcttg tgttcttttg agcagtgaag acgtccttga cagtcagctc	360
catgcacgag tcgtttctcg ctgtgaggat gttcaacatt ttgatgatct cttaaatagt	420
aatattgaag ttatgtccgt ggacaaaggt gctggtgagt gctagtaagc agaccagagt	480
tggaatcagt tgggaggtga gaccattaa tagatctcta tttgcagtga caatagcaga	540
cctcgtgcc	549

<210> SEQ ID NO 4

-continued

<211> LENGTH: 396

<212> TYPE: DNA

<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 4

```

atgggtctca cctcccaact gattccaact ctggtctgct tactagcact caccagcacc      60
tttgtccacg gacataactt caatattact attaaagaga tcataaaaat gttgaacatc     120
ctcacagcga gaaacgactc gtgcatggag ctgactgtca aggacgtctt cactgctcca     180
aagaacacaa gcgataagga aatcttctgc agagctgcta ctgtactgcg gcagatctat     240
acacacaact gctccaacag atatctcaga ggactctaca ggaacctcag cagcatggca     300
aacaagacct gttctatgaa tgaaatcaag aagagtacac tgaaagactt cttggaaagg     360
ctaaaagtga tcatgcagaa gaaatactac aggcatt                                396

```

<210> SEQ ID NO 5

<211> LENGTH: 396

<212> TYPE: DNA

<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 5

```

atgcctgtag tatttcttct gcatgatcac ttttagcctt tccaagaagt ctttcagtgt      60
actcttcttg atttcattca tagaacaggt cttgtttgcc atgctgctga ggttcctgta     120
gagtcctctg agatatctgt tggagcagtt gtgtgtatag atctgccgca gtacagtagc     180
agctctgcag aagatttctt tatcgcttgt gttcttttga gcagtgaaga cgtccttgac     240
agtcagctcc atgcacgagt cgtttctcgc tgtgaggatg ttcaacattt tgatgatctc     300
tttaatagta atattgaagt tatgtccgtg gacaaagggt ctggtgagtg ctagtaagca     360
gaccagagtt ggaatcagtt gggaggtgag acccat                                396

```

<210> SEQ ID NO 6

<211> LENGTH: 1013

<212> TYPE: DNA

<213> ORGANISM: Canis familiaris

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (35)..(916)

<400> SEQUENCE: 6

```

atctgaccaa aggcattgagg ggcctccggc cgag atg ata gtg ctg gcg cca gcc      55
                               Met Ile Val Leu Ala Pro Ala
                               1           5

tgg agc cca act gcc tcc ctg ttg ctg ctg ctg ctg ctc agc ccc ggc      103
Trp Ser Pro Thr Ala Ser Leu Leu Leu Leu Leu Ser Pro Gly
      10           15           20

ctc cgc ggg acc ccc gac tgc tcc ttc agc cac agc ccc atc tcc tcc      151
Leu Arg Gly Thr Pro Asp Cys Ser Phe Ser His Ser Pro Ile Ser Ser
      25           30           35

acc ttc gcg gtc acc atc cgc aag ctg tct gat tac ctg ctt cag gac      199
Thr Phe Ala Val Thr Ile Arg Lys Leu Ser Asp Tyr Leu Leu Gln Asp
      40           45           50           55

tat cca gtc act gtc gcc tcc aac ctg cag gac gac gag ctc tgc ggg      247
Tyr Pro Val Thr Val Ala Ser Asn Leu Gln Asp Asp Glu Leu Cys Gly
      60           65           70

gcg ttc tgg cgc ctg gtc ctg gcc cag cgc tgg atg gtg cgg ctc cag      295
Ala Phe Trp Arg Leu Val Leu Ala Gln Arg Trp Met Val Arg Leu Gln
      75           80           85

gct gtg gct gga tcc caa atg caa atc ctg ctg gag gct gtc aac acg      343
Ala Val Ala Gly Ser Gln Met Gln Ile Leu Leu Glu Ala Val Asn Thr

```

-continued

90	95	100		
gag ata cac ttt gtc acc ttc tgt gcc ttc cag ccc ctc ccc agc tgt			391	
Glu Ile His Phe Val Thr Phe Cys Ala Phe Gln Pro Leu Pro Ser Cys				
105	110	115		
ctt cgc ttc gtc cag acc aac atc tcc cac ctc ctg cag gac acc tcc			439	
Leu Arg Phe Val Gln Thr Asn Ile Ser His Leu Leu Gln Asp Thr Ser				
120	125	130	135	
cag cag ctg gcc gcc ctg aag ccc tgg atc acc cgc agg aat ttc tcc			487	
Gln Gln Leu Ala Ala Leu Lys Pro Trp Ile Thr Arg Arg Asn Phe Ser				
	140	145	150	
ggg tgc ctg gag ctg cag tgt cag ccc gac tcc tct aca ttg gtg ccc			535	
Gly Cys Leu Glu Leu Gln Cys Gln Pro Asp Ser Ser Thr Leu Val Pro				
	155	160	165	
cca agg agc ccc ggg gcc ctg gag gcc act gcc ttg cca gcc cct cag			583	
Pro Arg Ser Pro Gly Ala Leu Glu Ala Thr Ala Leu Pro Ala Pro Gln				
	170	175	180	
gca cct cgg ctg ctc ctc ctg ctg ctg ctg ccc gtg gct ctc ctg ctg			631	
Ala Pro Arg Leu Leu Leu Leu Leu Leu Leu Pro Val Ala Leu Leu Leu				
	185	190	195	
atg tcc act gcc tgg tgc ctg cat tgg cga agg agg cgg cgg cgg agg			679	
Met Ser Thr Ala Trp Cys Leu His Trp Arg Arg Arg Arg Arg Arg				
	200	205	210	215
tca ccc tac cct ggg gag cag agg aca ctg agg ccc agc gag cgg agc			727	
Ser Pro Tyr Pro Gly Glu Gln Arg Thr Leu Arg Pro Ser Glu Arg Ser				
	220	225	230	
cat ctg ccc gag gac aca gag ctg gga cct gga ggg agt cag cta gag			775	
His Leu Pro Glu Asp Thr Glu Leu Gly Pro Gly Gly Ser Gln Leu Glu				
	235	240	245	
act ggt ccc ttc ctc gac cac gca gcc cgg ctc gct ccc tcc cca gga			823	
Thr Gly Pro Phe Leu Asp His Ala Ala Pro Leu Ala Pro Ser Pro Gly				
	250	255	260	
tca agg caa cgc cgg ccc cca acg ccc cca aag cca gcc cca gcc cca			871	
Ser Arg Gln Arg Pro Pro Pro Thr Pro Pro Lys Pro Ala Pro Ala Pro				
	265	270	275	
cct ctc ccc ctc tgt aca aag tcc ttg ccc cca aga aat tgt ata			916	
Pro Leu Pro Leu Cys Thr Lys Ser Leu Pro Pro Arg Asn Cys Ile				
	280	285	290	
taaatcatcc ttttctacca aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa			976	
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaa			1013	
<210> SEQ ID NO 7				
<211> LENGTH: 294				
<212> TYPE: PRT				
<213> ORGANISM: Canis familiaris				
<400> SEQUENCE: 7				
Met Ile Val Leu Ala Pro Ala Trp Ser Pro Thr Ala Ser Leu Leu Leu				
1	5	10	15	
Leu Leu Leu Leu Ser Pro Gly Leu Arg Gly Thr Pro Asp Cys Ser Phe				
	20	25	30	
Ser His Ser Pro Ile Ser Ser Thr Phe Ala Val Thr Ile Arg Lys Leu				
	35	40	45	
Ser Asp Tyr Leu Leu Gln Asp Tyr Pro Val Thr Val Ala Ser Asn Leu				
	50	55	60	
Gln Asp Asp Glu Leu Cys Gly Ala Phe Trp Arg Leu Val Leu Ala Gln				
	65	70	75	80
Arg Trp Met Val Arg Leu Gln Ala Val Ala Gly Ser Gln Met Gln Ile				
	85	90	95	

-continued

Leu Leu Glu Ala Val Asn Thr Glu Ile His Phe Val Thr Phe Cys Ala
 100 105 110
 Phe Gln Pro Leu Pro Ser Cys Leu Arg Phe Val Gln Thr Asn Ile Ser
 115 120 125
 His Leu Leu Gln Asp Thr Ser Gln Gln Leu Ala Ala Leu Lys Pro Trp
 130 135 140
 Ile Thr Arg Arg Asn Phe Ser Gly Cys Leu Glu Leu Gln Cys Gln Pro
 145 150 155 160
 Asp Ser Ser Thr Leu Val Pro Pro Arg Ser Pro Gly Ala Leu Glu Ala
 165 170 175
 Thr Ala Leu Pro Ala Pro Gln Ala Pro Arg Leu Leu Leu Leu Leu
 180 185 190
 Leu Pro Val Ala Leu Leu Leu Met Ser Thr Ala Trp Cys Leu His Trp
 195 200 205
 Arg Arg Arg Arg Arg Arg Arg Ser Pro Tyr Pro Gly Glu Gln Arg Thr
 210 215 220
 Leu Arg Pro Ser Glu Arg Ser His Leu Pro Glu Asp Thr Glu Leu Gly
 225 230 235 240
 Pro Gly Gly Ser Gln Leu Glu Thr Gly Pro Phe Leu Asp His Ala Ala
 245 250 255
 Pro Leu Ala Pro Ser Pro Gly Ser Arg Gln Arg Pro Pro Pro Thr Pro
 260 265 270
 Pro Lys Pro Ala Pro Ala Pro Pro Leu Pro Leu Cys Thr Lys Ser Leu
 275 280 285
 Pro Pro Arg Asn Cys Ile
 290

<210> SEQ ID NO 8
 <211> LENGTH: 1013
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 8

tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt 60
 tttttttttt ttttttttgg tagaaaagga tgatttatat acaatttcctt gggggcaagg 120
 actttgtaca gagggggaga ggtggggctg gggctggctt tgggggcgtt gggggcgggc 180
 gttgccttga tcctggggag ggagcgagcg gggctgcgtg gtcgaggaag ggaccagtct 240
 ctagctgact ccctccaggt ccagctctg tgtcctcggg cagatggctc cgctcgtggt 300
 gcctcagtgt cctctgctcc ccagggtagg gtgacctcgc ccgccgcctc ctcgccaat 360
 gcaggcacca ggcagtggac atcagcagga gagccacggg cagcagcagc aggaggagca 420
 gccgaggtgc ctgaggggct ggcaaggcag tggcctccag ggccccgggg ctccctgggg 480
 gcaccaatgt agaggagtgc ggctgacact gcagctccag gcacccggag aaattcctgc 540
 ggggtgatcca gggcttcagg gcggccagct gctgggaggt gtcctgcagg aggtgggaga 600
 tgttggtctg gacgaagcga agacagctgg ggaggggctg gaaggcacag aaggtgacaa 660
 agtgtatctc cgtgttgaca gcctccagca ggatttgcatt ttgggatcca gccacagcct 720
 ggagccgcac catccagcgc tgggccagga ccaggcgcca gaacgccccg cagagctcgt 780
 cgtcctgcag gttggaggcg acagtgactg gatagtcctg aagcaggtaa tcagacagct 840
 tgcggatggt gaccgcgaag gtggaggaga tggggctgtg gctgaaggag cagtcggggg 900
 tcccgcggag gccggggctg agcagcagca gcagcaacag ggaggcagtt gggctccagg 960

-continued

ctggcgccag cactatcctc tcggcggag gccctcatg cctatggtca gat 1013

<210> SEQ ID NO 9

<211> LENGTH: 882

<212> TYPE: DNA

<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 9

atgatagtgc	tgggcgccag	ctggagccca	actgcctccc	tgttgctgct	gctgctgctc	60
agccccggcc	tccgcgggac	ccccgactgc	tccttcagcc	acagccccat	ctcctccacc	120
ttcgcggtca	ccatccgcaa	gctgtctgat	tacctgcttc	aggactatcc	agtcactgtc	180
gcctccaacc	tgagagacga	cgagctctgc	ggggcggtct	ggcgctggt	cctggccag	240
cgctggatgg	tgcggtccca	ggctgtggct	ggatcccaaa	tgcaaatcct	gctggaggct	300
gtcaacacgg	agatacactt	tgtcaccttc	tgtgccttcc	agccccctcc	cagctgtctt	360
cgcttcgtcc	agaccaacat	ctcccacctc	ctgcaggaca	cctcccagca	gctggccgcc	420
ctgaagccct	ggatcaccgg	caggaatttc	tccgggtgcc	tggagctgca	gtgtcagccc	480
gactcctcta	cattgggtgc	cccaaggagc	cccggggccc	tggaggccac	tgcttgcca	540
gccccctcag	caactcggct	gctcctcctg	ctgctgctgc	ccgtggctct	cctgctgatg	600
tccactgcct	ggtgcctgca	ttggcgaaag	aggcgggcgc	ggaggtcacc	ctaccctggg	660
gagcagagga	cactgaggcc	cagcgagcgg	agccatctgc	ccgaggacac	agagctggga	720
cctggaggga	gtcagctaga	gactgggtccc	ttcctcgacc	acgcagcccc	gctcgtcccc	780
tccccaggat	caaggcaacg	cccgccccca	acgcccccaa	agccagcccc	agccccacct	840
ctccccctct	gtacaaagtc	cttgccccca	agaaattgta	ta		882

<210> SEQ ID NO 10

<211> LENGTH: 882

<212> TYPE: DNA

<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 10

tatacaatth	cttgggggca	aggactttgt	acagaggggg	agaggtgggg	ctggggctgg	60
ctttgggggc	gttgggggag	ggcgttgctt	tgatcctggg	gagggagcga	gcggggctgc	120
gtggtcgagg	aaggggaccg	tctctagctg	actccctcca	ggtcccagct	ctgtgtcttc	180
gggcagatgg	ctccgctcgc	tgggcctcag	tgtcctctgc	tccccagggt	agggtgacct	240
ccgcgcggcg	ctccttcgcc	aatgcaggca	ccaggcagtg	gacatcagca	ggagagccac	300
gggcagcagc	agcaggagga	gcagccgagg	tgcctgaggg	gctggcaagg	cagtggcctc	360
cagggccccc	gggctccttg	ggggcaccaa	tgtagaggag	tcgggctgac	actgcagctc	420
caggcaccgc	gagaaattcc	tgcggtgat	ccagggtctc	agggcgccca	gctgctggga	480
ggtgtcctgc	aggaggtggg	agatgttggt	ctggacgaag	cgaagacagc	tggggagggg	540
ctggaaggca	cagaaggtga	caaagtgtat	ctccgtgttg	acagcctcca	gcaggatttg	600
catttgggat	ccagccacag	cctggagccg	caccatccag	cgctgggcca	ggaccaggcg	660
ccagaacgcc	ccgcagagct	cgctcgtcctg	caggttgagg	gcgacagtga	ctggatagtc	720
ctgaagcagg	taatcagaca	gcttgccgat	ggtgaccgcg	aaggtggagg	agatggggct	780
gtggctgaag	gagcagtcgg	gggtcccgcg	gagggcgggg	ctgagcagca	gcagcagcaa	840
cagggaggca	gttgggctcc	aggctggcgc	cagcactatc	at		882

-continued

<210> SEQ ID NO 11
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 11

ctattaatgg gtctcacctc ccaact 26

<210> SEQ ID NO 12
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 12

tcaactcggg gcacagagtc ttgg 24

<210> SEQ ID NO 13
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 13

ctggcgccag cctggagccc 20

<210> SEQ ID NO 14
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 14

gggagatggt ggtctggacg 20

<210> SEQ ID NO 15
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 15

gaccaggcgc cagaacgc 18

<210> SEQ ID NO 16
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 16

cggtcaccat ccgcaagc 18

-continued

```

<210> SEQ ID NO 17
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Primer

<400> SEQUENCE: 17

tggcaaggca gtggcctc                                     18

<210> SEQ ID NO 18
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Primer

<400> SEQUENCE: 18

gccgagatga tagtgctggc                                     20

<210> SEQ ID NO 19
<211> LENGTH: 324
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(324)

<400> SEQUENCE: 19

cat aac ttc aat att act att aaa gag atc atc aaa atg ttg aac atc      48
His Asn Phe Asn Ile Thr Ile Lys Glu Ile Ile Lys Met Leu Asn Ile
  1             5             10             15

ctc aca gcg aga aac gac tcg tgc atg gag ctg act gtc aag gac gtc      96
Leu Thr Ala Arg Asn Asp Ser Cys Met Glu Leu Thr Val Lys Asp Val
             20             25             30

ttc act gct cca aag aac aca agc gat aag gaa atc ttc tgc aga gct     144
Phe Thr Ala Pro Lys Asn Thr Ser Asp Lys Glu Ile Phe Cys Arg Ala
             35             40             45

gct act gta ctg cgg cag atc tat aca cac aac tgc tcc aac aga tat     192
Ala Thr Val Leu Arg Gln Ile Tyr Thr His Asn Cys Ser Asn Arg Tyr
  50             55             60

ctc aga gga ctc tac agg aac ctc agc agc atg gca aac aag acc tgt     240
Leu Arg Gly Leu Tyr Arg Asn Leu Ser Ser Met Ala Asn Lys Thr Cys
  65             70             75             80

tct atg aat gaa atc aag aag agt aca ctg aaa gac ttc ttg gaa agg     288
Ser Met Asn Glu Ile Lys Lys Ser Thr Leu Lys Asp Phe Leu Glu Arg
             85             90             95

cta aaa gtg atc atg cag aag aaa tac tac agg cat                     324
Leu Lys Val Ile Met Gln Lys Lys Tyr Tyr Arg His
  100             105

<210> SEQ ID NO 20
<211> LENGTH: 108
<212> TYPE: PRT
<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 20

His Asn Phe Asn Ile Thr Ile Lys Glu Ile Ile Lys Met Leu Asn Ile
  1             5             10             15

Leu Thr Ala Arg Asn Asp Ser Cys Met Glu Leu Thr Val Lys Asp Val
             20             25             30

Phe Thr Ala Pro Lys Asn Thr Ser Asp Lys Glu Ile Phe Cys Arg Ala

```

-continued

35	40	45	
Ala Thr Val Leu Arg Gln Ile Tyr Thr His Asn Cys Ser Asn Arg Tyr			
50	55	60	
Leu Arg Gly Leu Tyr Arg Asn Leu Ser Ser Met Ala Asn Lys Thr Cys			
65	70	75	80
Ser Met Asn Glu Ile Lys Lys Ser Thr Leu Lys Asp Phe Leu Glu Arg			
	85	90	95
Leu Lys Val Ile Met Gln Lys Lys Tyr Tyr Arg His			
100	105		
 <210> SEQ ID NO 21			
<211> LENGTH: 324			
<212> TYPE: DNA			
<213> ORGANISM: Canis familiaris			
 <400> SEQUENCE: 21			
atgcctgtag tatttcttct gcatgatcac ttttagcctt tccaagaagt ctttcagtgt			60
actcttcttg atttcattca tagaacaggt cttgtttgcc atgctgctga ggttcctgta			120
gagtcctctg agatatctgt tggagcagtt gtgtgtatag atctgccgca gtacagtagc			180
agctctgcag aagatttctt tatcgcttgt gttctttgga gcagtgaaga cgtccttgac			240
agtcagctcc atgcacgagt cgtttctcgc tgtgaggatg ttcaacattt tgatgatctc			300
tttaatagta atattgaagt tatg			324
 <210> SEQ ID NO 22			
<211> LENGTH: 804			
<212> TYPE: DNA			
<213> ORGANISM: Canis familiaris			
<220> FEATURE:			
<221> NAME/KEY: CDS			
<222> LOCATION: (1)..(804)			
 <400> SEQUENCE: 22			
acc ccc gac tgc tcc ttc agc cac agc ccc atc tcc tcc acc ttc gcg			48
Thr Pro Asp Cys Ser Phe Ser His Ser Pro Ile Ser Ser Thr Phe Ala			
1	5	10	15
gtc acc atc cgc aag ctg tct gat tac ctg ctt cag gac tat cca gtc			96
Val Thr Ile Arg Lys Leu Ser Asp Tyr Leu Leu Gln Asp Tyr Pro Val			
	20	25	30
act gtc gcc tcc aac ctg cag gac gac gag ctc tgc ggg gcg ttc tgg			144
Thr Val Ala Ser Asn Leu Gln Asp Asp Glu Leu Cys Gly Ala Phe Trp			
35	40	45	
cgc ctg gtc ctg gcc cag cgc tgg atg gtg cgg ctc cag gct gtg gct			192
Arg Leu Val Leu Ala Gln Arg Trp Met Val Arg Leu Gln Ala Val Ala			
50	55	60	
gga tcc caa atg caa atc ctg ctg gag gct gtc aac acg gag ata cac			240
Gly Ser Gln Met Gln Ile Leu Leu Glu Ala Val Asn Thr Glu Ile His			
65	70	75	80
ttt gtc acc ttc tgt gcc ttc cag ccc ctc ccc agc tgt ctt cgc ttc			288
Phe Val Thr Phe Cys Ala Phe Gln Pro Leu Pro Ser Cys Leu Arg Phe			
	85	90	95
gtc cag acc aac atc tcc cac ctc ctg cag gac acc tcc cag cag ctg			336
Val Gln Thr Asn Ile Ser His Leu Leu Gln Asp Thr Ser Gln Gln Leu			
100	105	110	
gcc gcc ctg aag ccc tgg atc acc cgc agg aat ttc tcc ggg tgc ctg			384
Ala Ala Leu Lys Pro Trp Ile Thr Arg Arg Asn Phe Ser Gly Cys Leu			
115	120	125	
gag ctg cag tgt cag ccc gac tcc tct aca ttg gtg ccc cca agg agc			432
Glu Leu Gln Cys Gln Pro Asp Ser Ser Thr Leu Val Pro Pro Arg Ser			

-continued

130	135	140	
ccc ggg gcc ctg gag gcc act gcc ttg cca gcc cct cag gca cct cgg			480
Pro Gly Ala Leu Glu Ala Thr Ala Leu Pro Ala Pro Gln Ala Pro Arg			
145	150	155	160
ctg ctc ctc ctg ctg ctg ctg ccc gtg gct ctc ctg ctg atg tcc act			528
Leu Leu Leu Leu Leu Leu Leu Pro Val Ala Leu Leu Leu Met Ser Thr			
	165	170	175
gcc tgg tgc ctg cat tgg cga agg agg cgg cgg cgg agg tca ccc tac			576
Ala Trp Cys Leu His Trp Arg Arg Arg Arg Arg Arg Ser Pro Tyr			
	180	185	190
cct ggg gag cag agg aca ctg agg ccc agc gag cgg agc cat ctg ccc			624
Pro Gly Glu Gln Arg Thr Leu Arg Pro Ser Glu Arg Ser His Leu Pro			
	195	200	205
gag gac aca gag ctg gga cct gga ggg agt cag cta gag act ggt ccc			672
Glu Asp Thr Glu Leu Gly Pro Gly Gly Ser Gln Leu Glu Thr Gly Pro			
	210	215	220
ttc ctc gac cac gca gcc ccg ctc gct ccc tcc cca gga tca agg caa			720
Phe Leu Asp His Ala Ala Pro Leu Ala Pro Ser Pro Gly Ser Arg Gln			
	225	230	235
cgc ccg ccc cca acg ccc cca aag cca gcc cca gcc cca cct ctc ccc			768
Arg Pro Pro Pro Thr Pro Pro Lys Pro Ala Pro Ala Pro Pro Leu Pro			
	245	250	255
ctc tgt aca aag tcc ttg ccc cca aga aat tgt ata			804
Leu Cys Thr Lys Ser Leu Pro Pro Arg Asn Cys Ile			
	260	265	
<210> SEQ ID NO 23			
<211> LENGTH: 268			
<212> TYPE: PRT			
<213> ORGANISM: Canis familiaris			
<400> SEQUENCE: 23			
Thr Pro Asp Cys Ser Phe Ser His Ser Pro Ile Ser Ser Thr Phe Ala			
1	5	10	15
Val Thr Ile Arg Lys Leu Ser Asp Tyr Leu Leu Gln Asp Tyr Pro Val			
	20	25	30
Thr Val Ala Ser Asn Leu Gln Asp Asp Glu Leu Cys Gly Ala Phe Trp			
	35	40	45
Arg Leu Val Leu Ala Gln Arg Trp Met Val Arg Leu Gln Ala Val Ala			
	50	55	60
Gly Ser Gln Met Gln Ile Leu Leu Glu Ala Val Asn Thr Glu Ile His			
	65	70	75
Phe Val Thr Phe Cys Ala Phe Gln Pro Leu Pro Ser Cys Leu Arg Phe			
	85	90	95
Val Gln Thr Asn Ile Ser His Leu Leu Gln Asp Thr Ser Gln Gln Leu			
	100	105	110
Ala Ala Leu Lys Pro Trp Ile Thr Arg Arg Asn Phe Ser Gly Cys Leu			
	115	120	125
Glu Leu Gln Cys Gln Pro Asp Ser Ser Thr Leu Val Pro Pro Arg Ser			
	130	135	140
Pro Gly Ala Leu Glu Ala Thr Ala Leu Pro Ala Pro Gln Ala Pro Arg			
	145	150	155
Leu Leu Leu Leu Leu Leu Leu Pro Val Ala Leu Leu Leu Met Ser Thr			
	165	170	175
Ala Trp Cys Leu His Trp Arg Arg Arg Arg Arg Arg Ser Pro Tyr			
	180	185	190
Pro Gly Glu Gln Arg Thr Leu Arg Pro Ser Glu Arg Ser His Leu Pro			

-continued

195	200	205	
Glu Asp Thr Glu Leu Gly Pro Gly Gly Ser Gln Leu Glu Thr Gly Pro			
210	215	220	
Phe Leu Asp His Ala Ala Pro Leu Ala Pro Ser Pro Gly Ser Arg Gln			
225	230	235	240
Arg Pro Pro Pro Thr Pro Pro Lys Pro Ala Pro Ala Pro Pro Leu Pro			
	245	250	255
Leu Cys Thr Lys Ser Leu Pro Pro Arg Asn Cys Ile			
	260	265	
<210> SEQ ID NO 24			
<211> LENGTH: 804			
<212> TYPE: DNA			
<213> ORGANISM: Canis familiaris			
<400> SEQUENCE: 24			
tatacaattt cttgggggca aggactttgt acagaggggg agaggtgggg ctggggctgg			60
ctttgggggc gttgggggag ggcgttgccct tgatcctggg gagggagcga gcggggctgc			120
gtggtcgagg aagggaccag tctctagctg actccctcca ggtcccagct ctgtgtcctc			180
gggcagatgg ctccgcctgc tgggcctcag tgcctctgc tccccagggt agggtgacct			240
ccgcgcgccg ctcccttcgcc aatgcaggca ccaggcagtg gacatcagca ggagagccac			300
gggcagcagc agcaggagga gcagccgagg tgcctgaggg gctggcaagg cagtggcctc			360
cagggccccc gggctccttg ggggcaccaa tgtagaggag tcgggctgac actgcagctc			420
cagggcaccg gagaaattcc tgcgggtgat ccagggcttc agggcgccca gctgctggga			480
ggtgtcctgc aggaggtggg agatgttggt ctggacgaag cgaagacagc tggggagggg			540
ctggaaggca cagaagtgta caaagtgtat ctccgtgttg acagcctcca gcaggatttg			600
catttgggat ccagccacag cctggagccg caccatccag cgctgggcca ggaccaggcg			660
ccagaacgcc ccgcagagct cgtcgtcctg caggttgag ggcacagtga ctggatagtc			720
ctgaagcagg taatcagaca gcttgccgat ggtgaccgcg aaggtggagg agatggggct			780
gtggctgaag gagcagtcgg ggg			804
<210> SEQ ID NO 25			
<211> LENGTH: 985			
<212> TYPE: DNA			
<213> ORGANISM: Canis familiaris			
<220> FEATURE:			
<221> NAME/KEY: CDS			
<222> LOCATION: (74)..(901)			
<400> SEQUENCE: 25			
ccggcctggc cccttcacag ccagctggg gcaagcctga tctgaccata ggcatgaggg			60
gcctccggcc gag atg ata gtg ctg gcg cca gcc tgg agc cca act gcc			109
Met Ile Val Leu Ala Pro Ala Trp Ser Pro Thr Ala			
1 5 10			
tcc ctg ttg ctg ctg ctg ctg ctc agc ccc ggc ctc cgc ggg acc ccc			157
Ser Leu Leu Leu Leu Leu Ser Pro Gly Leu Arg Gly Thr Pro			
15 20 25			
gac tgc tcc ttc agc cac agc ccc atc tcc tcc acc ttc gcg gtc acc			205
Asp Cys Ser Phe Ser His Ser Pro Ile Ser Ser Thr Phe Ala Val Thr			
30 35 40			
atc cgc aag ctg tct gat tac ctg ctt cag gac tat cca gtc act gtc			253
Ile Arg Lys Leu Ser Asp Tyr Leu Leu Gln Asp Tyr Pro Val Thr Val			
45 50 55 60			

-continued

gcc tcc aac ctg cag gac gac gag ctc tgc ggg gcg ttc tgg cgc ctg Ala Ser Asn Leu Gln Asp Asp Glu Leu Cys Gly Ala Phe Trp Arg Leu	301
65 70 75	
gtc ctg gcc cag cgc tgg atg gtg cgg ctc cag gct gtg gct gga tcc Val Leu Ala Gln Arg Trp Met Val Arg Leu Gln Ala Val Ala Gly Ser	349
80 85 90	
caa atg caa atc ctg ctg gag gct gtc aac acg gag ata cac ttt gtc Gln Met Gln Ile Leu Leu Glu Ala Val Asn Thr Glu Ile His Phe Val	397
95 100 105	
acc ttc tgt gcc ttc cag gac acc tcc cag cag ctg gcc gcc ctg aag Thr Phe Cys Ala Phe Gln Asp Thr Ser Gln Gln Leu Ala Ala Leu Lys	445
110 115 120	
ccc tgg atc acc cgc agg aat ttc tcc ggg tgc ctg gag ctg cag tgt Pro Trp Ile Thr Arg Arg Asn Phe Ser Gly Cys Leu Glu Leu Gln Cys	493
125 130 135 140	
cag ccc gac tcc tct aca ttg gtg ccc cca agg agc ccc ggg gcc ctg Gln Pro Asp Ser Ser Thr Leu Val Pro Pro Arg Ser Pro Gly Ala Leu	541
145 150 155	
gag gcc act gcc ttg cca gcc cct cag gca cct cgg ctg ctc ctc ctg Glu Ala Thr Ala Leu Pro Ala Pro Gln Ala Pro Arg Leu Leu Leu Leu	589
160 165 170	
ctg ctg ctg ccc gtg gct ctc ctg ctg atg tcc act gcc tgg tgc ctg Leu Leu Leu Pro Val Ala Leu Leu Met Ser Thr Ala Trp Cys Leu	637
175 180 185	
cat tgg cga agg agg cgg cgg cgg agg tca ccc tac cct ggg gag cag His Trp Arg Arg Arg Arg Arg Arg Arg Ser Pro Tyr Pro Gly Glu Gln	685
190 195 200	
agg aca ctg agg ccc agc gag cgg agc cat ctg ccc gag gac aca gag Arg Thr Leu Arg Pro Ser Glu Arg Ser His Leu Pro Glu Asp Thr Glu	733
205 210 215 220	
ctg gga cct gga ggg agt cag cta gag act ggt ccc ttc ctc gac cac Leu Gly Pro Gly Gly Ser Gln Leu Glu Thr Gly Pro Phe Leu Asp His	781
225 230 235	
gca gcc ccg ctc gct ccc tcc cca gga tca agg caa cgc ccg ccc cca Ala Ala Pro Leu Ala Pro Ser Pro Gly Ser Arg Gln Arg Pro Pro Pro	829
240 245 250	
acg ccc cca aag cca gcc cca gcc cca cct ctc ccc ctc tgt aca aag Thr Pro Pro Lys Pro Ala Pro Ala Pro Pro Leu Pro Leu Cys Thr Lys	877
255 260 265	
tcc ttg ccc cca aga aat tgt ata taaatcatcc ttttctacca gcaaaaaaaaaa Ser Leu Pro Pro Arg Asn Cys Ile	931
270 275	
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa	985

<210> SEQ ID NO 26

<211> LENGTH: 276

<212> TYPE: PRT

<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 26

Met Ile Val Leu Ala Pro Ala Trp Ser Pro Thr Ala Ser Leu Leu Leu	1 5 10 15
Leu Leu Leu Leu Ser Pro Gly Leu Arg Gly Thr Pro Asp Cys Ser Phe	20 25 30
Ser His Ser Pro Ile Ser Ser Thr Phe Ala Val Thr Ile Arg Lys Leu	35 40 45
Ser Asp Tyr Leu Leu Gln Asp Tyr Pro Val Thr Val Ala Ser Asn Leu	50 55 60
Gln Asp Asp Glu Leu Cys Gly Ala Phe Trp Arg Leu Val Leu Ala Gln	

-continued

65	70	75	80
Arg Trp Met Val	Arg Leu Gln Ala Val	Ala Gly Ser Gln Met	Gln Ile
	85	90	95
Leu Leu Glu Ala Val	Asn Thr Glu Ile His Phe Val	Thr Phe Cys Ala	
	100	105	110
Phe Gln Asp Thr Ser	Gln Gln Leu Ala Ala Leu Lys	Pro Trp Ile Thr	
	115	120	125
Arg Arg Asn Phe Ser	Gly Cys Leu Glu Leu Gln Cys	Gln Pro Asp Ser	
	130	135	140
Ser Thr Leu Val Pro	Pro Arg Ser Pro Gly Ala Leu Glu Ala Thr	Ala	
	145	150	155
Leu Pro Ala Pro Gln	Ala Pro Arg Leu Leu Leu Leu Leu Leu	Pro	
	165	170	175
Val Ala Leu Leu Leu	Met Ser Thr Ala Trp Cys Leu His	Trp Arg Arg	
	180	185	190
Arg Arg Arg Arg Arg	Ser Pro Tyr Pro Gly Glu Gln Arg	Thr Leu Arg	
	195	200	205
Pro Ser Glu Arg Ser	His Leu Pro Glu Asp Thr Glu Leu Gly	Pro Gly	
	210	215	220
Gly Ser Gln Leu Glu	Thr Gly Pro Phe Leu Asp His Ala Ala	Pro Leu	
	225	230	235
Ala Pro Ser Pro Gly	Ser Arg Gln Arg Pro Pro Pro Thr Pro	Pro Lys	
	245	250	255
Pro Ala Pro Ala Pro	Pro Leu Pro Leu Cys Thr Lys Ser Leu Pro	Pro	
	260	265	270
Arg Asn Cys Ile			
	275		

<210> SEQ ID NO 27
 <211> LENGTH: 985
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 27

tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	60
ttgtcgttag aaaaggatga tttatataca atttcttggg ggcaaggact ttgtacagag	120
ggggagaggt ggggctgggg ctggccttgg gggcgttggg ggcgggcgtt gccttgatcc	180
tggggagggg gcgagcgggg ctgcgtggtc gaggaaggga ccagtctcta gctgactccc	240
tccaggtccc agctctgtgt cctcgggcag atggctccgc tcgctgggccc tcagtgtcct	300
ctgctcccca gggtaggggtg acctccgccg ccgcctcctt cgccaatgca ggcaccaggc	360
agtggacatc agcaggagag ccacgggcag cagcagcagg aggagcagcc gaggtgcctg	420
aggggctggc aaggcagtgg cctccagggc cccggggctc cttgggggca ccaatgtaga	480
ggagtcgggc tgacactgca gctccaggca cccggagaaa ttcctgcggg tgatccaggg	540
cttcagggag gccagctgct gggaggtgtc ctggaaggca cagaaggtga caaagtgtat	600
ctcgtgtttg acagcctcca gcaggatttg catttgggat ccagccacag cctggagccg	660
caccatccag cgctggggcca ggaccaggcg ccagaacgcc ccgagagct cgctgtcctg	720
caggttgagg gcgacagtga ctggatagtc ctgaagcagg taatcagaca gcttgccgat	780
ggtgaccgcg aaggtggagg agatggggct gtggctgaag gagcagtcgg gggccccgcg	840
gaggccgggg ctgagcagca gcagcagcaa cagggaggca gttgggctcc aggctggcgc	900

-continued

cagcactatc atctcggccg gaggccctc atgcctatgg tcagatcagg cttgccccag	960
ctgggcgtgg aaggggccag gccgg	985

<210> SEQ ID NO 28
 <211> LENGTH: 828
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 28

atgatagtgc tggcgccagc ctggagccca actgcctccc tgttgctgct gctgctgctc	60
agccccggcc tccgcgggac ccccgactgc tccttcagcc acagcccat ctctccacc	120
ttcgcggtca ccatccgcaa gctgtctgat tacctgcttc aggactatcc agtcactgtc	180
gcctccaacc tgcaggacga cgagctctgc ggggcgttct ggcgcctggt cctggccccag	240
cgctggatgg tgcggctcca ggctgtggct ggatcccaa tgcaaactct gctggaggct	300
gtcaacacgg agatacactt tgtcaccttc tgtgccttcc aggacacctc ccagcagctg	360
gccgccctga agccctggat caccgcaggg aatttctccg ggtgcctgga gctgcagtgt	420
cagccccgact cctctacatt ggtgccccca aggagccccg gggccctgga ggccactgcc	480
ttgccagccc ctccaggacc tcggctgctc ctctgctgc tgcctgccgt ggctctcctg	540
ctgatgtcca ctgcctgggt cctgcattgg cgaaggaggc ggcggcggag gtcaccctac	600
cctggggagc agaggacact gagggccagc gagcggagcc atctgcccga ggacacagag	660
ctgggacctg gagggagtca gctagagact ggtcccttcc tcgaccacgc agccccgctc	720
gtccctctcc caggatcaag gcaacgccc cccccaacgc ccccaaagcc agccccagcc	780
ccacctctcc ccctctgtac aaagtccctg cccccaagaa attgtata	828

<210> SEQ ID NO 29
 <211> LENGTH: 828
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 29

tatacaattt cttgggggca aggactttgt acagaggggg agaggtgggg ctggggctgg	60
ctttgggggc gttgggggag ggcgttgctt tgatcctggg gagggagcga gcggggctgc	120
gtggtcgagg aagggaccag tctctagctg actccctcca ggtcccagct ctgtgtcttc	180
gggcagatgg ctccgctcgc tgggcctcag tgtcctctgc tccccagggt aggggtgacct	240
ccgcgcgcgc ctccctcgcc aatgcaggca ccaggcagtg gacatcagca ggagagccac	300
gggcagcagc agcaggagga gcagccgagg tgcctgaggg gctggcaagg cagtggcctc	360
cagggccccc gggctccttg ggggcaccaa tgtagaggag tcgggctgac actgcagctc	420
caggcacccg gagaaattcc tgcgggtgat ccagggcctc agggcgccca gctgctggga	480
ggtgtcctgg aaggcacaga aggtgacaaa gtgtatctcc gtgttgacag cctccagcag	540
gatttgctt tgggatccag ccacagcctg gagccgcacc atccagcgt gggccaggac	600
caggcgccag aacgccccgc agagctcgtc gtcctgcagg ttggaggcga cagtgactgg	660
atagtcctga agcaggtaat cagacagctt gcggatggtg accgcgaagg tggaggagat	720
ggggctgtgg ctgaaggagc agtcgggggt ccgcggagg ccggggctga gcagcagcag	780
cagcaacagg gaggcagttg ggctccagcc tggcgccagc actatcat	828

<210> SEQ ID NO 30
 <211> LENGTH: 750

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Canis familiaris
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(750)

<400> SEQUENCE: 30

acc ccc gac tgc tcc ttc agc cac agc ccc atc tcc tcc acc ttc gcg      48
Thr Pro Asp Cys Ser Phe Ser His Ser Pro Ile Ser Ser Thr Phe Ala
   1             5             10             15

gtc acc atc cgc aag ctg tct gat tac ctg ctt cag gac tat cca gtc      96
Val Thr Ile Arg Lys Leu Ser Asp Tyr Leu Leu Gln Asp Tyr Pro Val
           20           25           30

act gtc gcc tcc aac ctg cag gac gac gag ctc tgc ggg gcg ttc tgg     144
Thr Val Ala Ser Asn Leu Gln Asp Asp Glu Leu Cys Gly Ala Phe Trp
           35           40           45

cgc ctg gtc ctg gcc cag cgc tgg atg gtg cgg ctc cag gct gtg gct     192
Arg Leu Val Leu Ala Gln Arg Trp Met Val Arg Leu Gln Ala Val Ala
           50           55           60

gga tcc caa atg caa atc ctg ctg gag gct gtc aac acg gag ata cac     240
Gly Ser Gln Met Gln Ile Leu Leu Glu Ala Val Asn Thr Glu Ile His
           65           70           75           80

ttt gtc acc ttc tgt gcc ttc cag gac acc tcc cag cag ctg gcc gcc     288
Phe Val Thr Phe Cys Ala Phe Gln Asp Thr Ser Gln Gln Leu Ala Ala
           85           90           95

ctg aag ccc tgg atc acc cgc agg aat ttc tcc ggg tgc ctg gag ctg     336
Leu Lys Pro Trp Ile Thr Arg Arg Asn Phe Ser Gly Cys Leu Glu Leu
          100          105          110

cag tgt cag ccc gac tcc tct aca ttg gtg ccc cca agg agc ccc ggg     384
Gln Cys Gln Pro Asp Ser Ser Thr Leu Val Pro Pro Arg Ser Pro Gly
          115          120          125

gcc ctg gag gcc act gcc ttg cca gcc cct cag gca cct cgg ctg ctc     432
Ala Leu Glu Ala Thr Ala Leu Pro Ala Pro Gln Ala Pro Arg Leu Leu
          130          135          140

ctc ctg ctg ctg ctg ccc gtg gct ctc ctg ctg atg tcc act gcc tgg     480
Leu Leu Leu Leu Leu Pro Val Ala Leu Leu Leu Met Ser Thr Ala Trp
          145          150          155          160

tgc ctg cat tgg cga agg agg cgg cgg cgg agg tca ccc tac cct ggg     528
Cys Leu His Trp Arg Arg Arg Arg Arg Arg Ser Pro Tyr Pro Gly
          165          170          175

gag cag agg aca ctg agg ccc agc gag cgg agc cat ctg ccc gag gac     576
Glu Gln Arg Thr Leu Arg Pro Ser Glu Arg Ser His Leu Pro Glu Asp
          180          185          190

aca gag ctg gga cct gga ggg agt cag cta gag act ggt ccc ttc ctc     624
Thr Glu Leu Gly Pro Gly Gly Ser Gln Leu Glu Thr Gly Pro Phe Leu
          195          200          205

gac cac gca gcc ccg ctc gct ccc tcc cca gga tca agg caa cgc ccg     672
Asp His Ala Ala Pro Leu Ala Pro Ser Pro Gly Ser Arg Gln Arg Pro
          210          215          220

ccc cca acg ccc cca aag cca gcc cca gcc cca cct ctc ccc ctc tgt     720
Pro Pro Thr Pro Pro Lys Pro Ala Pro Ala Pro Pro Leu Pro Leu Cys
          225          230          235          240

aca aag tcc ttg ccc cca aga aat tgt ata                               750
Thr Lys Ser Leu Pro Pro Arg Asn Cys Ile
          245          250

<210> SEQ ID NO 31
<211> LENGTH: 250
<212> TYPE: PRT
<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 31

```

-continued

Thr Pro Asp Cys Ser Phe Ser His Ser Pro Ile Ser Ser Thr Phe Ala
 1 5 10 15
 Val Thr Ile Arg Lys Leu Ser Asp Tyr Leu Leu Gln Asp Tyr Pro Val
 20 25 30
 Thr Val Ala Ser Asn Leu Gln Asp Asp Glu Leu Cys Gly Ala Phe Trp
 35 40 45
 Arg Leu Val Leu Ala Gln Arg Trp Met Val Arg Leu Gln Ala Val Ala
 50 55 60
 Gly Ser Gln Met Gln Ile Leu Leu Glu Ala Val Asn Thr Glu Ile His
 65 70 75 80
 Phe Val Thr Phe Cys Ala Phe Gln Asp Thr Ser Gln Gln Leu Ala Ala
 85 90 95
 Leu Lys Pro Trp Ile Thr Arg Arg Asn Phe Ser Gly Cys Leu Glu Leu
 100 105 110
 Gln Cys Gln Pro Asp Ser Ser Thr Leu Val Pro Pro Arg Ser Pro Gly
 115 120 125
 Ala Leu Glu Ala Thr Ala Leu Pro Ala Pro Gln Ala Pro Arg Leu Leu
 130 135 140
 Leu Leu Leu Leu Leu Pro Val Ala Leu Leu Leu Met Ser Thr Ala Trp
 145 150 155 160
 Cys Leu His Trp Arg Arg Arg Arg Arg Arg Ser Pro Tyr Pro Gly
 165 170 175
 Glu Gln Arg Thr Leu Arg Pro Ser Glu Arg Ser His Leu Pro Glu Asp
 180 185 190
 Thr Glu Leu Gly Pro Gly Gly Ser Gln Leu Glu Thr Gly Pro Phe Leu
 195 200 205
 Asp His Ala Ala Pro Leu Ala Pro Ser Pro Gly Ser Arg Gln Arg Pro
 210 215 220
 Pro Pro Thr Pro Pro Lys Pro Ala Pro Ala Pro Pro Leu Pro Leu Cys
 225 230 235 240
 Thr Lys Ser Leu Pro Pro Arg Asn Cys Ile
 245 250

<210> SEQ ID NO 32

<211> LENGTH: 750

<212> TYPE: DNA

<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 32

tatacaattt cttgggggca aggactttgt acagaggggg agaggtgggg ctggggctgg 60
 ctttgggggc gttgggggcg ggcgttgccct tgatcctggg gagggagcga gcggggctgc 120
 gtggtcgagg aagggaccag tctctagctg actccctcca ggtcccagct ctgtgtcctc 180
 gggcagatgg ctccgctcgc tgggcctcag tgtcctctgc tccccagggt aggggtgacct 240
 ccgccgccgc ctccctcgcc aatgcaggca ccaggcagtg gacatcagca ggagagccac 300
 gggcagcagc agcaggagga gcagccgagg tgccctgagg gctggcaagg cagtggcctc 360
 cagggccccc gggctccttg ggggcaccaa tgtagaggag tcgggctgac actgcagctc 420
 caggcacccg gagaaattcc tgcgggtgat ccagggcttc agggcggccca gctgctggga 480
 ggtgtcctgg aaggcacaga aggtgacaaa gtgtatctcc gtgttgacag cctccagcag 540
 gatttgcat tgggtaccag ccacagcctg gagccgcacc atccagcgt gggccaggac 600
 caggcgccag aacgccccgc agagctcgtc gtcctgcagg ttggaggcga cagtgactgg 660

-continued

```

atagtctga agcaggtaat cagacagctt gcggatggtg accgcgaagg tggaggagat 720
ggggctgtgg ctgaaggagc agtcgggggt 750

```

```

<210> SEQ ID NO 33
<211> LENGTH: 1019
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (74)..(166)

```

```

<400> SEQUENCE: 33

```

```

ccggcctggc cccttccacg cccagctggg gcaagcctga tctgaccata ggcattgagg 60
gcctccggcc gag atg ata gtg ctg gcg cca gcc tgg agc cca act gtg 109
      Met Ile Val Leu Ala Pro Ala Trp Ser Pro Thr Val
      1             5             10

cgt ata ccc ggg gga caa ggc ggg gga cag gca gag cgc tac cga gct 157
Arg Ile Pro Gly Gly Gln Gly Gly Gly Gln Ala Glu Arg Tyr Arg Ala
      15             20             25

ggg cag agc tgagagagca gacggacaga ggccctccctg ttgctgctgc 206
Gly Gln Ser
      30

tgctgctcag ccccggcctc cgcgggaccc ccgactgctc cttcagccac agcccatct 266
cctccacctt cgcggtcacc atccgcaagc tgtctgatta cctgcttcag gactatccag 326
tcaactgtgc ctccaacctg caggacgacg agctctgcgg ggcgttcttg cgcctgggtcc 386
tggcccagcg ctggatggtg cggctccagg ctgtggctgg atcccaaatg caaatcctgc 446
tggaggctgt caacacggag atacactttg tcaccttctg tgccttcag gacacctccc 506
agcagctggc cgcctgaag ccctggatca cccgcaggaa tttctccggg tgcctggagc 566
tgcaagtgtca gcccactcc tctacattgg tgccccaag gagccccggg gccctggagg 626
ccactgcctt gccagccct caggcacctc ggctgctcct cctgctgctg ctgcccgtgg 686
ctctcctgct gatgtccact gcctgggtgcc tgcattggcg aaggaggcgg cggcggaggt 746
caccctaccc tggggagcag aggacactga ggcccagcga gcggagccat ctgcccagag 806
acacagagct gggacctgga gggagtcagc tagagactgg tcccttcctc gaccacgcag 866
ccccgtcgc tccctcccca ggatcaaggc aacgcccgcc cccaacgccc ccaaagccag 926
ccccagcccc acctctcccc ctctgtacaa agtccttgcc ccaagaaat tgtatataaa 986
tcataccttt ctacaaaaa aaaaaaaaaa aaa 1019

```

```

<210> SEQ ID NO 34
<211> LENGTH: 31
<212> TYPE: PRT
<213> ORGANISM: Canis familiaris

```

```

<400> SEQUENCE: 34

```

```

Met Ile Val Leu Ala Pro Ala Trp Ser Pro Thr Val Arg Ile Pro Gly
  1             5             10             15

Gly Gln Gly Gly Gly Gln Ala Glu Arg Tyr Arg Ala Gly Gln Ser
      20             25             30

```

```

<210> SEQ ID NO 35
<211> LENGTH: 1019
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris

```

```

<400> SEQUENCE: 35

```

-continued

```

tttttttttt ttttttttgg tagaaaagga tgatttatat acaatttctt gggggcaagg      60
actttgtaca gagggggaga ggtggggctg gggctggctt tgggggcgtt gggggcgggc      120
gttgcttga tcctggggag ggagcgagcg gggctgcgtg gtcgaggaag ggaccagtct      180
ctagctgact ccctccaggt cccagctctg tgcctcggg cagatggctc cgctcgctgg      240
gcctcagtg cctctgctcc ccagggtagg gtgacctccg ccgccgcctc ctctcgccaat      300
gcaggcacca ggagtgagc atcagcagga gagccacggg cagcagcagc aggaggagca      360
gccgaggtgc ctgaggggct ggcaaggcag tggcctccag gggcccgggg ctctctgggg      420
gcaccaatgt agaggagtcg ggctgacct gcagctccag gcacccggag aaattcctgc      480
gggtgatcca gggcttcagg gcggccagct gctgggaggt gtcctggaag gcacagaagg      540
tgacaaagtg tatctccgtg ttgacagcct ccagcaggat ttgcatttgg gatccagcca      600
cagcctggag ccgcaccatc cagcgtctgg ccaggaccag gcgccagaac gcccgcaga      660
gctcgtcgtc ctgcaggttg gaggcgacag tgactggata gtcctgaagc aggtaatcag      720
acagcttgcg gatggtgacc gcgaagggtg aggagatggg gctgtggctg aaggagcagt      780
cgggggtccc gcggaggccg gggctgagca gcagcagcag caacagggag gcctctgtcc      840
gtctgtcttc tcagctctgc ccagctcgtt agcgtctctg ctgtcccccg ccttgtcccc      900
cgggtatacg cacagtgtgg ctccaggctg gcgccagcac tatcatctcg gccggaggcc      960
cctcatgcct atggtcagat caggcttgcc ccagctgggc gtggaagggg ccaggccgg      1019

```

```

<210> SEQ ID NO 36
<211> LENGTH: 93
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris

```

```

<400> SEQUENCE: 36

```

```

atgatagtgc tggcgccagc ctggagccca actgtgcgta taccggggg acaaggcggg      60
ggacaggcag agcgtaccg agctgggcag agc                                93

```

```

<210> SEQ ID NO 37
<211> LENGTH: 93
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris

```

```

<400> SEQUENCE: 37

```

```

gctctgcccc gctcggtagc gctctgctg tccccgcct tgtcccccg gtatacgcac      60
agttgggctc caggctggcg ccagcactat cat                                93

```

```

<210> SEQ ID NO 38
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

```

```

<400> SEQUENCE: 38

```

```

tgaattcgga cataacttca atattac                                27

```

```

<210> SEQ ID NO 39
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

```

-continued

<400> SEQUENCE: 39

tctcagagatt cagcttcaat gcttgta

27

<210> SEQ ID NO 40

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 40

cccaagctta tgggtctcac ctccaac

28

<210> SEQ ID NO 41

<211> LENGTH: 395

<212> TYPE: DNA

<213> ORGANISM: Felis catus

<400> SEQUENCE: 41

ggccataggc atgaagggcc tccggccgag atgatagtgc tggcgccagc ctggagccca 60

actacctccc tgctgctgct gctactgctc agccctggcc tccgcggtc ccccgactgt 120

tccttcagcc acagcccat ctcctccacc ttcaaggcca ccacccgaaa gctgtctgat 180

tacctgcttc aggattaccc agtcaccgct gcctccaacc tacaggacga cgagctctgt 240

gggccattct ggcacctggt cctggcccag cgctggatgg gtcggctcaa ggctgtggct 300

gggtcccaga tgcaaacct gctggaggcg gtcaaacacc agatacattt tgtcaccttg 360

tgtgccttcc agccctccc cagctgtctt cgatt 395

<210> SEQ ID NO 42

<211> LENGTH: 793

<212> TYPE: DNA

<213> ORGANISM: Felis catus

<400> SEQUENCE: 42

cttcaaggct accatccgaa agctgtctga ttacctgctt caggattacc cagtcaccgt 60

cgctccaac ctacaggacg acgagctctg tgggccattc tggcacctgg tcctggccca 120

gcgctggatg ggtcggtcca aggtgtggc tgggtcccag atgcaaagcc tgcaggaggc 180

ggtaaacacc gagatacatt ttgtcacctt gtgtgccttc cagccctcc ccagctgtct 240

tcgattcgtc cagaccaaca tctcccacct cctgcaggac acctccgagc agctggcggc 300

cttgaagccc tggatcaccg gcaggaattt ctcggggtgc ctggagctac agtgtcagcc 360

cgactcctcc accccactgc ccccaaggag cccaggggcc ttggaggcca cagccctgcc 420

agccctcag gcccctctgc tgctcctcct gctgctgttg cctgtggctc tcttgctgat 480

gtccgcccgc tgggtcctgc actggcgaag aaggagatgg agaacgcctt accccaggga 540

gcagagggaag aactgaggc ccagagagag gaatcacctg cccaggaca cagagccggg 600

actcgggaaa agtcagctag agactgggtc cttcctcgac cacgctgcc cgctcactct 660

ccccccggga tggaggcaac gccagcccc aacgccagcc ccagaccac ctatccccct 720

ctgtacaaa tgctgtctct caggaaattg tatataaatc atccttttct accaaaaaaa 780

aaaaaaaaaaa aaa 793

<210> SEQ ID NO 43

<211> LENGTH: 942

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Felis catus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (31)..(903)

<400> SEQUENCE: 43

ggccataggc atgaagggcc tccggccgag atg ata gtg ctg gcg cca gcc tgg      54
                               Met Ile Val Leu Ala Pro Ala Trp
                               1           5

agc cca act acc tcc ctg ctg ctg ctg cta ctg ctc agc cct ggc ctc      102
Ser Pro Thr Thr Ser Leu Leu Leu Leu Leu Leu Ser Pro Gly Leu
10           15           20

cgc ggg tcc ccc gac tgt tcc ttc agc cac agc ccc atc tcc tcc acc      150
Arg Gly Ser Pro Asp Cys Ser Phe Ser His Ser Pro Ile Ser Ser Thr
25           30           35           40

ttc aag gtc acc atc cga aag ctg tct gat tac ctg ctt cag gat tac      198
Phe Lys Val Thr Ile Arg Lys Leu Ser Asp Tyr Leu Leu Gln Asp Tyr
45           50           55

cca gtc acc gtc gcc tcc aac cta cag gac gac gag ctc tgt ggg cca      246
Pro Val Thr Val Ala Ser Asn Leu Gln Asp Asp Glu Leu Cys Gly Pro
60           65           70

ttc tgg cac ctg gtc ctg gcc cag cgc tgg atg ggt cgg ctc aag gct      294
Phe Trp His Leu Val Leu Ala Gln Arg Trp Met Gly Arg Leu Lys Ala
75           80           85

gtg gct ggg tcc cag atg caa agc ctg ctg gag gcg gtc aac acc gag      342
Val Ala Gly Ser Gln Met Gln Ser Leu Leu Glu Ala Val Asn Thr Glu
90           95           100

ata cat ttt gtc acc ttg tgt gcc ttc cag ccc ctc ccc agc tgt ctt      390
Ile His Phe Val Thr Leu Cys Ala Phe Gln Pro Leu Pro Ser Cys Leu
105          110          115          120

cga ttc gtc cag acc aac atc tcc cac ctc ctg cag gac acc tcc gag      438
Arg Phe Val Gln Thr Asn Ile Ser His Leu Leu Gln Asp Thr Ser Glu
125          130          135

cag ctg gcg gcc ttg aag ccc tgg atc acc cgc agg aat ttc tcg ggg      486
Gln Leu Ala Ala Leu Lys Pro Trp Ile Thr Arg Arg Asn Phe Ser Gly
140          145          150

tgc ctg gag cta cag tgt cag ccc gac tcc tcc acc cca ctg ccc cca      534
Cys Leu Glu Leu Gln Cys Gln Pro Asp Ser Ser Thr Pro Leu Pro Pro
155          160          165

agg agc ccc agg gcc ttg gag gcc aca gcc ctg cca gcc cct cag gcc      582
Arg Ser Pro Arg Ala Leu Glu Ala Thr Ala Leu Pro Ala Pro Gln Ala
170          175          180

cct ctg ctg ctc ctc ctg ctg ctg ttg cct gtg gct ctc ttg ctg atg      630
Pro Leu Leu Leu Leu Leu Leu Leu Leu Pro Val Ala Leu Leu Leu Met
185          190          195          200

tcc gcc gcc tgg tgc ctg cac tgg cga aga agg aga tgg aga acg ccc      678
Ser Ala Ala Trp Cys Leu His Trp Arg Arg Arg Arg Trp Arg Thr Pro
205          210          215

tac ccc agg gag cag agg aag aca ctg agg ccc aga gag agg aat cac      726
Tyr Pro Arg Glu Gln Arg Lys Thr Leu Arg Pro Arg Glu Arg Asn His
220          225          230

ctg ccc gag gac aca gag ccg gga ctc gga gaa agt cag cta gag act      774
Leu Pro Glu Asp Thr Glu Pro Gly Leu Gly Glu Ser Gln Leu Glu Thr
235          240          245

ggg tcc ttc ctc gac cac gct gcc ccg ctc act ctc ccc ccg gga tgg      822
Gly Ser Phe Leu Asp His Ala Ala Pro Leu Thr Leu Pro Pro Gly Trp
250          255          260

agg caa cgc cag ccc cca acg cca gcc cca gac cca cct atc ccc ctc      870
Arg Gln Arg Gln Pro Pro Thr Pro Ala Pro Asp Pro Pro Ile Pro Leu
265          270          275          280

```


-continued

tgt aca aag tcc ttg tcc tca gga aat tgt ata taaatcatcc ttttctacca 923
 Cys Thr Lys Ser Leu Ser Ser Gly Asn Cys Ile
 285 290

aaaaaaaaa aaaaaaaaaa 942

<210> SEQ ID NO 44
 <211> LENGTH: 291
 <212> TYPE: PRT
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 44

Met Ile Val Leu Ala Pro Ala Trp Ser Pro Thr Thr Ser Leu Leu Leu
 1 5 10 15
 Leu Leu Leu Leu Ser Pro Gly Leu Arg Gly Ser Pro Asp Cys Ser Phe
 20 25 30
 Ser His Ser Pro Ile Ser Ser Thr Phe Lys Val Thr Ile Arg Lys Leu
 35 40 45
 Ser Asp Tyr Leu Leu Gln Asp Tyr Pro Val Thr Val Ala Ser Asn Leu
 50 55 60
 Gln Asp Asp Glu Leu Cys Gly Pro Phe Trp His Leu Val Leu Ala Gln
 65 70 75 80
 Arg Trp Met Gly Arg Leu Lys Ala Val Ala Gly Ser Gln Met Gln Ser
 85 90 95
 Leu Leu Glu Ala Val Asn Thr Glu Ile His Phe Val Thr Leu Cys Ala
 100 105 110
 Phe Gln Pro Leu Pro Ser Cys Leu Arg Phe Val Gln Thr Asn Ile Ser
 115 120 125
 His Leu Leu Gln Asp Thr Ser Glu Gln Leu Ala Ala Leu Lys Pro Trp
 130 135 140
 Ile Thr Arg Arg Asn Phe Ser Gly Cys Leu Glu Leu Gln Cys Gln Pro
 145 150 155 160
 Asp Ser Ser Thr Pro Leu Pro Pro Arg Ser Pro Arg Ala Leu Glu Ala
 165 170 175
 Thr Ala Leu Pro Ala Pro Gln Ala Pro Leu Leu Leu Leu Leu Leu
 180 185 190
 Leu Pro Val Ala Leu Leu Leu Met Ser Ala Ala Trp Cys Leu His Trp
 195 200 205
 Arg Arg Arg Arg Trp Arg Thr Pro Tyr Pro Arg Glu Gln Arg Lys Thr
 210 215 220
 Leu Arg Pro Arg Glu Arg Asn His Leu Pro Glu Asp Thr Glu Pro Gly
 225 230 235 240
 Leu Gly Glu Ser Gln Leu Glu Thr Gly Ser Phe Leu Asp His Ala Ala
 245 250 255
 Pro Leu Thr Leu Pro Pro Gly Trp Arg Gln Arg Gln Pro Pro Thr Pro
 260 265 270
 Ala Pro Asp Pro Pro Ile Pro Leu Cys Thr Lys Ser Leu Ser Ser Gly
 275 280 285
 Asn Cys Ile
 290

<210> SEQ ID NO 45
 <211> LENGTH: 942
 <212> TYPE: DNA
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 45

-continued

```

tttttttttt tttttttttt ggtagaaaag gatgatttat atacaatttc ctgaggacaa    60
ggacttttga cagaggggga taggtgggtc tggggctggc gttgggggct ggcgttgct    120
ccatcccggg gggagagtga gcggggcagc gtggtcgagg aaggaaccag tctctagctg    180
actttctccg agtcccggct ctgtgtcctc gggcaggtga ttcctctctc tgggcctcag    240
tgtcttcctc tgctccctgg ggtagggcgt tctccatctc cttcttcgcc agtgcaggca    300
ccaggcggcg gacatcagca agagagccac aggcaacagc agcaggagga gcagcagagg    360
ggcctgaggg gctggcaggg ctgtggcctc caaggccctg gggctccttg ggggcagtgg    420
ggtggaggag tcgggctgac actgtagctc caggcaccac gagaaattcc tgcgggtgat    480
ccagggtctc aaggccgcca gctgctcgga ggtgtcctgc aggaggtggg agatgttggg    540
ctggacgaat cgaagacagc tggggagggg ctggaaggca cacaaggtga caaaatgtat    600
ctcgtgtgtt accgcctcca gcaggctttg catctgggac ccagccacag ccttgagccg    660
acccatccag cgctgggcca ggaccaggtg ccagaatggc ccacagagct cgtcgtcctg    720
taggttgagg gcgacggtga ctgggtaatc ctgaagcagg taatcagaca gctttcggat    780
ggtgaccttg aaggtggagg agatggggct gtggctgaag gaacagtcgg gggacccgcg    840
gaggccaggg ctgagcagta gcagcagcag cagggaggta gttgggctcc aggctggcgc    900
cagcactatc atctcggccg gagggccttc atgcctatgg cc                      942

```

<210> SEQ ID NO 46

<211> LENGTH: 873

<212> TYPE: DNA

<213> ORGANISM: Felis catus

<400> SEQUENCE: 46

```

atgatatgtc tggcgccagc ctggagccca actacctccc tgctgctgct gctactgctc    60
agccctggcc tccgcgggtc ccccgactgt tccttcagcc acagcccat ctcctccacc    120
ttcaaggta ccatccgaaa gctgtctgat tacctgcttc aggattaccc agtcaccgtc    180
gcctccaacc tacaggacga cgagctctgt gggccattct ggcacctggt cctggccag    240
cgctggatgg gtcggctcaa ggctgtggct gggteccaga tgcaaagcct gctggaggcg    300
gtcaacaccg agatacattt tgtcaccttg tgtgccttec agccctccc cagctgtctt    360
cgattcgtcc agaccaacat ctcccacctc ctgcaggaca cctccgagca gctggcggcc    420
ttgaagccct ggtaccccg caggaatttc tcgggggtgcc tggagctaca gtgtcagccc    480
gactcctcca cccactgcc cccaaggagc ccagggcctt tggaggccac agccctgcca    540
gcccctcagg cccctctgct gctcctcctg ctgctgttgc ctgtggctct cttgctgatg    600
tcggccgctt ggtgcctgca ctggcggaaga aggagatgga gaacgcccta cccagggag    660
cagaggaaga cactgaggcc cagagagagg aatcacctgc ccgaggacac agagccggga    720
ctcgagaaaa gtcagctaga gactggttcc ttcctcgacc acgctgcccc gctcactctc    780
ccccgggat ggaggcaacg ccagcccca acgccagccc cagaccacc tatcccctc    840
tgtacaaagt ccttgtcctc aggaattgt ata                      873

```

<210> SEQ ID NO 47

<211> LENGTH: 873

<212> TYPE: DNA

<213> ORGANISM: Felis catus

<400> SEQUENCE: 47

-continued

tatacaattt cctgaggaca aggactttgt acagaggggg ataggtgggt ctggggctgg	60
cggtgggggc tggcgttgcc tccatcccgg ggggagagtg agcggggcag cgtggctgag	120
gaaggaacca gtctctagct gactttctcc gagtcccggc tctgtgtcct cgggcaggtg	180
attcctctct ctggggcctca gtgtcttcct ctgctccctg gggtagggcg ttctccatct	240
ccttcttcgc cagtgcaggc accaggcggc ggacatcagc aagagagcca caggcaacag	300
cagcaggagg agcagcagag gggcctgagg ggctggcagg gctgtggcct ccaaggccct	360
ggggctcctt gggggcagtg ggggtggagga gtcgggctga cactgtagct ccaggcaccc	420
cgagaaattc ctgcggtgga tccagggtt caaggccgcc agctgctcgg aggtgtcctg	480
caggaggtgg gagatgttg tctggacgaa tcgaagacag ctggggaggg gctggaaggc	540
acacaaggtg aaaaaatgta tctcgggtgtt gaccgcctcc agcaggcttt gcatctggga	600
cccagccaca gccttgagcc gacccatcca gcgctgggcc aggaccagggt gccagaatgg	660
cccacagagc tcgtcgtcct gtaggttgga ggcgacggtg actgggtaat cctgaagcag	720
gtaatcagac agctttcggga tggtagcctt gaaggtggag gagatggggc tgtggctgaa	780
ggaacagtcg ggggacccgc ggaggccagg gctgagcagt agcagcagca gcagggaggt	840
agttgggctc caggctggcg ccagcactat cat	873

<210> SEQ ID NO 48
 <211> LENGTH: 795
 <212> TYPE: DNA
 <213> ORGANISM: Felis catus
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(795)

<400> SEQUENCE: 48

tcc ccc gac tgt tcc ttc agc cac agc ccc atc tcc tcc acc ttc aag	48
Ser Pro Asp Cys Ser Phe Ser His Ser Pro Ile Ser Ser Thr Phe Lys	
1 5 10 15	
gtc acc atc cga aag ctg tct gat tac ctg ctt cag gat tac cca gtc	96
Val Thr Ile Arg Lys Leu Ser Asp Tyr Leu Leu Gln Asp Tyr Pro Val	
20 25 30	
acc gtc gcc tcc aac cta cag gac gac gag ctc tgt ggg cca ttc tgg	144
Thr Val Ala Ser Asn Leu Gln Asp Asp Glu Leu Cys Gly Pro Phe Trp	
35 40 45	
cac ctg gtc ctg gcc cag cgc tgg atg ggt cgg ctc aag gct gtg gct	192
His Leu Val Leu Ala Gln Arg Trp Met Gly Arg Leu Lys Ala Val Ala	
50 55 60	
ggg tcc cag atg caa agc ctg ctg gag gcg gtc aac acc gag ata cat	240
Gly Ser Gln Met Gln Ser Leu Leu Glu Ala Val Asn Thr Glu Ile His	
65 70 75 80	
ttt gtc acc ttg tgt gcc ttc cag ccc ctc ccc agc tgt ctt cga ttc	288
Phe Val Thr Leu Cys Ala Phe Gln Pro Leu Pro Ser Cys Leu Arg Phe	
85 90 95	
gtc cag acc aac atc tcc cac ctc ctg cag gac acc tcc gag cag ctg	336
Val Gln Thr Asn Ile Ser His Leu Leu Gln Asp Thr Ser Glu Gln Leu	
100 105 110	
gcg gcc ttg aag ccc tgg atc acc cgc agg aat ttc tcg ggg tgc ctg	384
Ala Ala Leu Lys Pro Trp Ile Thr Arg Arg Asn Phe Ser Gly Cys Leu	
115 120 125	
gag cta cag tgt cag ccc gac tcc tcc acc cca ctg ccc cca agg agc	432
Glu Leu Gln Cys Gln Pro Asp Ser Ser Thr Pro Leu Pro Pro Arg Ser	
130 135 140	
ccc agg gcc ttg gag gcc aca gcc ctg cca gcc cct cag gcc cct ctg	480
Pro Arg Ala Leu Glu Ala Thr Ala Leu Pro Ala Pro Gln Ala Pro Leu	

-continued

145	150	155	160	
ctg ctc ctc ctg ctg ctg ttg cct gtg gct ctc ttg ctg atg tcc gcc				528
Leu Leu Leu Leu Leu Leu Leu Pro Val Ala Leu Leu Leu Met Ser Ala	165	170	175	
gcc tgg tgc ctg cac tgg cga aga agg aga tgg aga acg ccc tac ccc				576
Ala Trp Cys Leu His Trp Arg Arg Arg Trp Arg Thr Pro Tyr Pro	180	185	190	
agg gag cag agg aag aca ctg agg ccc aga gag agg aat cac ctg ccc				624
Arg Glu Gln Arg Lys Thr Leu Arg Pro Arg Glu Arg Asn His Leu Pro	195	200	205	
gag gac aca gag ccg gga ctc gga gaa agt cag cta gag act ggt tcc				672
Glu Asp Thr Glu Pro Gly Leu Gly Glu Ser Gln Leu Glu Thr Gly Ser	210	215	220	
ttc ctc gac cac gct gcc ccg ctc act ctc ccc ccg gga tgg agg caa				720
Phe Leu Asp His Ala Ala Pro Leu Thr Leu Pro Pro Gly Trp Arg Gln	225	230	235	240
cgc cag ccc cca acg cca gcc cca gac cca cct atc ccc ctc tgt aca				768
Arg Gln Pro Pro Thr Pro Ala Pro Asp Pro Pro Ile Pro Leu Cys Thr	245	250	255	
aag tcc ttg tcc tca gga aat tgt ata				795
Lys Ser Leu Ser Ser Gly Asn Cys Ile	260	265		

<210> SEQ ID NO 49

<211> LENGTH: 265

<212> TYPE: PRT

<213> ORGANISM: Felis catus

<400> SEQUENCE: 49

Ser Pro Asp Cys Ser Phe Ser His Ser Pro Ile Ser Ser Thr Phe Lys	1	5	10	15
Val Thr Ile Arg Lys Leu Ser Asp Tyr Leu Leu Gln Asp Tyr Pro Val	20	25	30	
Thr Val Ala Ser Asn Leu Gln Asp Asp Glu Leu Cys Gly Pro Phe Trp	35	40	45	
His Leu Val Leu Ala Gln Arg Trp Met Gly Arg Leu Lys Ala Val Ala	50	55	60	
Gly Ser Gln Met Gln Ser Leu Leu Glu Ala Val Asn Thr Glu Ile His	65	70	75	80
Phe Val Thr Leu Cys Ala Phe Gln Pro Leu Pro Ser Cys Leu Arg Phe	85	90	95	
Val Gln Thr Asn Ile Ser His Leu Leu Gln Asp Thr Ser Glu Gln Leu	100	105	110	
Ala Ala Leu Lys Pro Trp Ile Thr Arg Arg Asn Phe Ser Gly Cys Leu	115	120	125	
Glu Leu Gln Cys Gln Pro Asp Ser Ser Thr Pro Leu Pro Pro Arg Ser	130	135	140	
Pro Arg Ala Leu Glu Ala Thr Ala Leu Pro Ala Pro Gln Ala Pro Leu	145	150	155	160
Leu Leu Leu Leu Leu Leu Leu Pro Val Ala Leu Leu Leu Met Ser Ala	165	170	175	
Ala Trp Cys Leu His Trp Arg Arg Arg Arg Trp Arg Thr Pro Tyr Pro	180	185	190	
Arg Glu Gln Arg Lys Thr Leu Arg Pro Arg Glu Arg Asn His Leu Pro	195	200	205	
Glu Asp Thr Glu Pro Gly Leu Gly Glu Ser Gln Leu Glu Thr Gly Ser	210	215	220	

-continued

Phe Leu Asp His Ala Ala Pro Leu Thr Leu Pro Pro Gly Trp Arg Gln
225 230 235 240

Arg Gln Pro Pro Thr Pro Ala Pro Asp Pro Pro Ile Pro Leu Cys Thr
245 250 255

Lys Ser Leu Ser Ser Gly Asn Cys Ile
260 265

<210> SEQ ID NO 50
<211> LENGTH: 795
<212> TYPE: DNA
<213> ORGANISM: Felis catus

<400> SEQUENCE: 50

```
tatacaattt cctgaggaca aggactttgt acagaggggg ataggtgggt ctggggcttg      60
cgttgggggc tggcgttgcc tccatcccgg ggggagagtg agcggggcag cgtggtcgag     120
gaaggaacca gtctctagtct gactttctcc gagtcccggc tctgtgtcct cgggcaggtg     180
attcctctct ctggggcctca gtgtcttctc ctgctccctg gggtagggcg ttctccatct     240
ccttcttcgc cagtgcaggc accaggcggc ggacatcagc aagagagcca caggcaacag     300
cagcaggagg agcagcagag gggcctgagg ggctggcagg gctgtggcct ccaaggccct     360
ggggctcctt gggggcagtg ggggtggagga gtcgggctga cactgtagct ccaggcaccc     420
cgagaaattc ctgcgggtga tccagggtt caaggccgcc agctgctcgg aggtgtcctg     480
caggaggttg gagatgttg tctggacgaa tcgaagacag ctggggaggg gctggaaggc     540
acacaaggtg aaaaaatgta tctcgggtgt gaccgcctcc agcaggcttt gcatctggga     600
cccagccaca gccttgagcc gacccatcca gcgctgggcc aggaccaggt gccagaatgg     660
cccacagagc tcgtctcctc gtaggttgga ggcgacggtg actgggtaat cctgaagcag     720
gtaatcagac agctttcgga tgggtgacct gaaggtggag gagatggggc tgtggctgaa     780
ggaacagtcg gggga                                           795
```

<210> SEQ ID NO 51
<211> LENGTH: 321
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 51

```
aatgtgtctt ctgctttgga aaagtgtcac ccttgacaa gctgtgaaac caaaggcctg      60
gtgaagggtc agcggggaac taacaagact gatgttatct gtgttcccca gcctcgggta     120
agagccctag tgggtgtccc catcattatg gggatcctgc ttgttgcctt gttggtgtct     180
gcctgcatcc gaaaggtggt caagaagcca gagaataagg ttatgtatca ggaccctgtg     240
gaggacttgg aggaatttcc tatgcccccg cactccattg ctccggtgca agagacctta     300
catgggtgcc agcccgtcac c                                           321
```

<210> SEQ ID NO 52
<211> LENGTH: 1425
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (196)..(1017)

<400> SEQUENCE: 52

```
tagactcccc ggaatattca ggggaactcc cggcgctaag ggtctccagg agtccgcccc      60
```

-continued

tgcccaacga agcgcggccac gattggtccc cgaagacccc gcccatctcc tggcgggggc	120
ggcggggggc aagggtctggg gagttactaa agacatcccc gcgcccctac tccgctgcct	180
gctattcacc tcgcc atg gtt ctc ctg cct ctg cgc tgt ctc ttc tgg ggc	231
Met Val Leu Leu Pro Leu Arg Cys Leu Phe Trp Gly	
1 5 10	
tcc ttg ttg acc acc gtc tac cca gaa cca cgc act gca tgc aga gaa	279
Ser Leu Leu Thr Thr Val Tyr Pro Glu Pro Arg Thr Ala Cys Arg Glu	
15 20 25	
aag caa tac cta gta gac agt cag tgc tgt aat atg tgc cca cca gga	327
Lys Gln Tyr Leu Val Asp Ser Gln Cys Cys Asn Met Cys Pro Pro Gly	
30 35 40	
gag aaa ctg gtg aat gac tgc cta cat acc att gac acg gaa tgc act	375
Glu Lys Leu Val Asn Asp Cys Leu His Thr Ile Asp Thr Glu Cys Thr	
45 50 55 60	
cgt tgc caa aca ggc gaa ttc cta gac act tgg aac gca gag aga cac	423
Arg Cys Gln Thr Gly Glu Phe Leu Asp Thr Trp Asn Ala Glu Arg His	
65 70 75	
tgt cac cag cac aaa tac tgc gac ccc aac cta ggg ctc cat gtc gag	471
Cys His Gln His Lys Tyr Cys Asp Pro Asn Leu Gly Leu His Val Glu	
80 85 90	
aag gag ggc acg tca gaa aca gac acc act tgc aca tgc gat gaa ggt	519
Lys Glu Gly Thr Ser Glu Thr Asp Thr Thr Cys Thr Cys Asp Glu Gly	
95 100 105	
ctg cat tgt acc aac gct gcc tgt gag agc tgc acc atg cac agc ctg	567
Leu His Cys Thr Asn Ala Ala Cys Glu Ser Cys Thr Met His Ser Leu	
110 115 120	
tgc ccc cct ggc ctg gga gtc aaa cag atc gct aca ggg att tct gat	615
Cys Pro Pro Gly Leu Gly Val Lys Gln Ile Ala Thr Gly Ile Ser Asp	
125 130 135 140	
acc atc tgc gat ccc tgc ccc atc ggc ttc ttc tcc aat gtg tct tct	663
Thr Ile Cys Asp Pro Cys Pro Ile Gly Phe Phe Ser Asn Val Ser Ser	
145 150 155	
gct ttg gaa aag tgt cac cct tgg aca agc tgt gaa acc aaa ggc ctg	711
Ala Leu Glu Lys Cys His Pro Trp Thr Ser Cys Glu Thr Lys Gly Leu	
160 165 170	
gtg aag gtt cag gcg gga act aac aag act gat gtt atc tgt ggt ccc	759
Val Lys Val Gln Ala Gly Thr Asn Lys Thr Asp Val Ile Cys Gly Pro	
175 180 185	
cag cct cgg tta aga gcc cta gtg gtg gtc ccc atc att atg ggg atc	807
Gln Pro Arg Leu Arg Ala Leu Val Val Val Pro Ile Ile Met Gly Ile	
190 195 200	
ctg ctt gtt gtc ctg ttg gtg tct gcc tgc atc cga aag gtg gtc aag	855
Leu Leu Val Val Leu Leu Val Ser Ala Cys Ile Arg Lys Val Val Lys	
205 210 215 220	
aag cca gag aat aag gtt atg tat cag gac cct gtg gag gac ttg gag	903
Lys Pro Glu Asn Lys Val Met Tyr Gln Asp Pro Val Glu Asp Leu Glu	
225 230 235	
gaa ttt cct atg ccc ccg cac tcc att gct ccg gtg caa gag acc tta	951
Glu Phe Pro Met Pro Pro His Ser Ile Ala Pro Val Gln Glu Thr Leu	
240 245 250	
cat ggg tgc cag ccc gtc acc cag gag gac ggc aaa gag agc cgc atc	999
His Gly Cys Gln Pro Val Thr Gln Glu Asp Gly Lys Glu Ser Arg Ile	
255 260 265	
tcc gtg cag gag aga gtg tgaggcagcg tgtgcccagg agtgtgacag	1047
Ser Val Gln Glu Arg Val	
270	
cgtgggagag tgggcgcgtg gctggagagc ctggagctgc tggaggggca tgaaggggcg	1107
gtgtccctt gcctgcaccc ctgtgtgtgca gaaacagaga accttccacc ccaccccttg	1167

-continued

```

agcccatcc acctcccaac ttgcttttaa agatggagat gaaacttttg gggggccaga 1227
tagtaatatc caccaaccca gcatttcagg gccctgaggt gtatatcacg gtggtttcta 1287
cgagcccagg aagaccacag aagagccatt gtggcattgt ttgtgacagt ggacaactgg 1347
aggccactta gctgttcagc agcaggggac tggctaaata aaatttgtaa tatattata 1407
aaaaaaaaa aaaaaaaaaa 1425

```

```

<210> SEQ ID NO 53
<211> LENGTH: 274
<212> TYPE: PRT
<213> ORGANISM: Canis familiaris

```

```

<400> SEQUENCE: 53

```

```

Met Val Leu Leu Pro Leu Arg Cys Leu Phe Trp Gly Ser Leu Leu Thr
  1             5             10            15
Thr Val Tyr Pro Glu Pro Arg Thr Ala Cys Arg Glu Lys Gln Tyr Leu
      20            25            30
Val Asp Ser Gln Cys Cys Asn Met Cys Pro Pro Gly Glu Lys Leu Val
      35            40            45
Asn Asp Cys Leu His Thr Ile Asp Thr Glu Cys Thr Arg Cys Gln Thr
      50            55            60
Gly Glu Phe Leu Asp Thr Trp Asn Ala Glu Arg His Cys His Gln His
      65            70            75            80
Lys Tyr Cys Asp Pro Asn Leu Gly Leu His Val Glu Lys Glu Gly Thr
      85            90            95
Ser Glu Thr Asp Thr Thr Cys Thr Cys Asp Glu Gly Leu His Cys Thr
      100           105           110
Asn Ala Ala Cys Glu Ser Cys Thr Met His Ser Leu Cys Pro Pro Gly
      115           120           125
Leu Gly Val Lys Gln Ile Ala Thr Gly Ile Ser Asp Thr Ile Cys Asp
      130           135           140
Pro Cys Pro Ile Gly Phe Phe Ser Asn Val Ser Ser Ala Leu Glu Lys
      145           150           155           160
Cys His Pro Trp Thr Ser Cys Glu Thr Lys Gly Leu Val Lys Val Gln
      165           170           175
Ala Gly Thr Asn Lys Thr Asp Val Ile Cys Gly Pro Gln Pro Arg Leu
      180           185           190
Arg Ala Leu Val Val Val Pro Ile Ile Met Gly Ile Leu Leu Val Val
      195           200           205
Leu Leu Val Ser Ala Cys Ile Arg Lys Val Val Lys Lys Pro Glu Asn
      210           215           220
Lys Val Met Tyr Gln Asp Pro Val Glu Asp Leu Glu Glu Phe Pro Met
      225           230           235           240
Pro Pro His Ser Ile Ala Pro Val Gln Glu Thr Leu His Gly Cys Gln
      245           250           255
Pro Val Thr Gln Glu Asp Gly Lys Glu Ser Arg Ile Ser Val Gln Glu
      260           265           270
Arg Val

```

```

<210> SEQ ID NO 54
<211> LENGTH: 1425
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris

```

```

<400> SEQUENCE: 54

```

-continued

```

tttttttttt ttttttttta taaatatatt acaaatttta tttagccagt cccctgctgc      60
tgaacagcta agtggcctcc agttgtccac tgtcacaac aatgccacaa tggctcttcg      120
tgggtcttcc tgggctcgta gaaaccaccg tgatatacac ctgagggcc tgaatgctg      180
ggttggtgga tattactatc tggcccccca aaagtttcat ctccatcttt aaaagcaagt      240
tgggaggtgg aatgggctcc aggggtgggg tgggaaggttc tctgtttctg cagcacaggg      300
gtgcaggcag gggagcaccg ccccttcatg cccctccagc agctccaggc tctccagcca      360
cgcgccact ctcccacgct gtcacactcc tgggcacacg ctgcctcaca ctctctcctg      420
cacggagatg cggctctctt tgccgtcttc ctgggtgacg ggctggcacc catgtaaggt      480
ctcttgacc ggagcaatgg agtgcggggg cataggaaat tcctccaagt cctccacagg      540
gtcctgatac ataaccttat tctctggctt cttgaccacc tttcggatgc aggagacac      600
caacaggaca acaagcagga tccccataat gatggggacc accactaggg ctcttaaccg      660
aggctgggga ccacagataa catcagtctt gttagtctcc gcctgaacct tcaccaggcc      720
tttggtttca cagcttgtcc aagggtgaca cttttccaaa gcagaagaca cattggagaa      780
gaagccgatg gggcagggat cgcagatggt atcagaaatc cctgtagcga tctgtttgac      840
tcccaggcca ggggggcaca ggctgtgcat ggtgcagctc tcacaggcag cgttggtaaca      900
atgcagacct tcatcgcatg tgcaagtggg gtctgtttct gacgtgccct ccttctcgac      960
atggagccct aggttggggt cgcagtattt gtgctggtga cagtgtctct ctgcgttcca     1020
agtgtctagg aattcgctg tttggcaacg agtgcattcc gtgtcaatgg tatgtaggca     1080
gtcattcacc agtttctctc ctgggtggga catattacag cactgactgt ctactaggta     1140
ttgtctttct ctgcctgcat tgccgtggtc tgggttagac gtggtcaaca aggagcccca     1200
gaagagacag cgcagaggga ggagaacctt ggcgaggtga atagcaggca gcggagtagg     1260
ggcgcgggga tgtctttagt aactccccag cccttgcccc cgcccgcccc gccagggaga     1320
tgggcggggg cttcggggac caatcgtggc cggtctcgtt gggcagggcg gagctcctgg     1380
agacccttag cgccgggagt tcccctgaat attcccggga gtcta                        1425

```

```

<210> SEQ ID NO 55
<211> LENGTH: 822
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris

```

```

<400> SEQUENCE: 55
atggttctcc tgccctgctg ctgtctcttc tggggctcct tgttgaccac cgtctaccca      60
gaaccacgca ctgcctgcat agaaaagcaa tacctagtag acagtcatg ctgtaatatg      120
tgccaccag gagagaaact ggtgaatgac tgccacata ccattgacac ggaatgcact      180
cgttgccaaa caggcgaatt cctagacact tggaaacgag agagacactg tcaccagcac      240
aaatactgct accccaacct agggctccat gtcgagaagg agggcacgtc agaaacagac      300
accacttgca catgcatgta aggtctgcat tgtaccaacg ctgcctgtga gagctgcacc      360
atgcacagcc tgtgcccccc tggcctggga gtcaaacaga tcgctacagg gatctctgat      420
accatctgct atccctgccc catcggtctc ttctccaatg tgtcttctgc tttggaaaag      480
tgtcacccct ggacaagctg tgaaacaaaa ggctgtgtga aggttcaggc gggaaactaac      540
aagactgatg ttatctgtgg tccccagcct cggttaagag ccctagtggg ggtcccatc      600
attatgggga tcctgcttgt tgtcctgttg gtgtctgcct gcacccgaaa ggtggtaag      660

```


-continued

```

aagccagaga ataaggttat gtatcaggac cctgtggagg acttggagga atttcctatg 720
ccccgcact ccattgctcc ggtgcaagag accttacatg ggtgccagcc cgtcaccag 780
gaggacggca aagagagccg catctccgtg caggagagag tg 822

```

```

<210> SEQ ID NO 56
<211> LENGTH: 822
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris

```

```

<400> SEQUENCE: 56

```

```

cactctctcc tgcacggaga tgcggtcttc tttgccgtcc tcctgggtga cgggctggca 60
cccatgtaag gtctcttgca cgggagcaat ggagtgcggg ggcataggaa attcctccaa 120
gtcctccaca gggctctgat acataacctt attctctggc ttcttgacca ctttctcgat 180
gcaggcagac accaacagga caacaagcag gatccccata atgatgggga ccaccactag 240
ggctcttaac cgaggctggg gaccacagat aacatcagtc ttgttagttc ccgcctgaac 300
cttcaccagg cttttggttt cacagcttgt ccaaggggtga cacttttcca aagcagaaga 360
cacattggag aagaagccga tggggcaggg atcgccagatg gtatcagaaa tccctgtagc 420
gatctgtttg actcccaggc caggggggca caggctgtgc atggtgcagc tctcacaggc 480
agcgttggtg caatgcagac cttcatcgca tgtgcaagtg gtgtctgttt ctgacgtgcc 540
ctccttctcg acatggagcc ctagggtggg gtcgcagtat ttgtgctggt gacagtgtct 600
ctctgcgttc caagtgtcta ggaattcgcc tgtttggcaa cgagtgcatt ccgtgtcaat 660
ggtagttagg cagtcattca ccagtttctc tcctgggtggg cacatattac agcactgact 720
gtctactagg tattgctttt ctctgcatgc agtgcggtggt tctgggtaga cggtggtcaa 780
caaggagccc cagaagagac agcgagagg caggagaacc at 822

```

```

<210> SEQ ID NO 57
<211> LENGTH: 765
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(765)

```

```

<400> SEQUENCE: 57

```

```

cca gaa cca cgc act gca tgc aga gaa aag caa tac cta gta gac agt 48
Pro Glu Pro Arg Thr Ala Cys Arg Glu Lys Gln Tyr Leu Val Asp Ser
  1             5             10            15

cag tgc tgt aat atg tgc cca cca gga gag aaa ctg gtg aat gac tgc 96
Gln Cys Cys Asn Met Cys Pro Pro Gly Glu Lys Leu Val Asn Asp Cys
  20            25            30

cta cat acc att gac acg gaa tgc act cgt tgc caa aca ggc gaa ttc 144
Leu His Thr Ile Asp Thr Glu Cys Thr Arg Cys Gln Thr Gly Glu Phe
  35            40            45

cta gac act tgg aac gca gag aga cac tgt cac cag cac aaa tac tgc 192
Leu Asp Thr Trp Asn Ala Glu Arg His Cys His Gln His Lys Tyr Cys
  50            55            60

gac ccc aac cta ggg ctc cat gtc gag aag gag ggc acg tca gaa aca 240
Asp Pro Asn Leu Gly Leu His Val Glu Lys Glu Gly Thr Ser Glu Thr
  65            70            75            80

gac acc act tgc aca tgc gat gaa ggt ctg cat tgt acc aac gct gcc 288
Asp Thr Thr Cys Thr Cys Asp Glu Gly Leu His Cys Thr Asn Ala Ala
  85            90            95

tgt gag agc tgc acc atg cac agc ctg tgc ccc cct ggc ctg gga gtc 336
Cys Glu Ser Cys Thr Met His Ser Leu Cys Pro Pro Gly Leu Gly Val

```

-continued

100	105	110	
aaa cag atc gct aca ggg att tct gat acc atc tgc gat ccc tgc ccc			384
Lys Gln Ile Ala Thr Gly Ile Ser Asp Thr Ile Cys Asp Pro Cys Pro			
115	120	125	
atc ggc ttc ttc tcc aat gtg tct tct gct ttg gaa aag tgt cac cct			432
Ile Gly Phe Phe Ser Asn Val Ser Ser Ala Leu Glu Lys Cys His Pro			
130	135	140	
tgg aca agc tgt gaa acc aaa ggc ctg gtg aag gtt cag gcg gga act			480
Trp Thr Ser Cys Glu Thr Lys Gly Leu Val Lys Val Gln Ala Gly Thr			
145	150	155	160
aac aag act gat gtt atc tgt ggt ccc cag cct cgg tta aga gcc cta			528
Asn Lys Thr Asp Val Ile Cys Gly Pro Gln Pro Arg Leu Arg Ala Leu			
165	170	175	
gtg gtg gtc ccc atc att atg ggg atc ctg ctt gtt gtc ctg ttg gtg			576
Val Val Val Pro Ile Ile Met Gly Ile Leu Leu Val Val Leu Leu Val			
180	185	190	
tct gcc tgc atc cga aag gtg gtc aag aag cca gag aat aag gtt atg			624
Ser Ala Cys Ile Arg Lys Val Val Lys Lys Pro Glu Asn Lys Val Met			
195	200	205	
tat cag gac cct gtg gag gac ttg gag gaa ttt cct atg ccc ccg cac			672
Tyr Gln Asp Pro Val Glu Asp Leu Glu Glu Phe Pro Met Pro Pro His			
210	215	220	
tcc att gct ccg gtg caa gag acc tta cat ggg tgc cag ccc gtc acc			720
Ser Ile Ala Pro Val Gln Glu Thr Leu His Gly Cys Gln Pro Val Thr			
225	230	235	240
cag gag gac ggc aaa gag agc cgc atc tcc gtg cag gag aga gtg			765
Gln Glu Asp Gly Lys Glu Ser Arg Ile Ser Val Gln Glu Arg Val			
245	250	255	

<210> SEQ ID NO 58

<211> LENGTH: 255

<212> TYPE: PRT

<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 58

Pro Glu Pro Arg Thr Ala Cys Arg Glu Lys Gln Tyr Leu Val Asp Ser	
1 5 10 15	
Gln Cys Cys Asn Met Cys Pro Pro Gly Glu Lys Leu Val Asn Asp Cys	
20 25 30	
Leu His Thr Ile Asp Thr Glu Cys Thr Arg Cys Gln Thr Gly Glu Phe	
35 40 45	
Leu Asp Thr Trp Asn Ala Glu Arg His Cys His Gln His Lys Tyr Cys	
50 55 60	
Asp Pro Asn Leu Gly Leu His Val Glu Lys Glu Gly Thr Ser Glu Thr	
65 70 75 80	
Asp Thr Thr Cys Thr Cys Asp Glu Gly Leu His Cys Thr Asn Ala Ala	
85 90 95	
Cys Glu Ser Cys Thr Met His Ser Leu Cys Pro Pro Gly Leu Gly Val	
100 105 110	
Lys Gln Ile Ala Thr Gly Ile Ser Asp Thr Ile Cys Asp Pro Cys Pro	
115 120 125	
Ile Gly Phe Phe Ser Asn Val Ser Ser Ala Leu Glu Lys Cys His Pro	
130 135 140	
Trp Thr Ser Cys Glu Thr Lys Gly Leu Val Lys Val Gln Ala Gly Thr	
145 150 155 160	
Asn Lys Thr Asp Val Ile Cys Gly Pro Gln Pro Arg Leu Arg Ala Leu	
165 170 175	

-continued

Val	Val	Val	Pro	Ile	Ile	Met	Gly	Ile	Leu	Leu	Val	Val	Leu	Leu	Val
			180					185					190		
Ser	Ala	Cys	Ile	Arg	Lys	Val	Val	Lys	Lys	Pro	Glu	Asn	Lys	Val	Met
		195					200					205			
Tyr	Gln	Asp	Pro	Val	Glu	Asp	Leu	Glu	Glu	Phe	Pro	Met	Pro	Pro	His
	210					215					220				
Ser	Ile	Ala	Pro	Val	Gln	Glu	Thr	Leu	His	Gly	Cys	Gln	Pro	Val	Thr
	225				230					235					240
Gln	Glu	Asp	Gly	Lys	Glu	Ser	Arg	Ile	Ser	Val	Gln	Glu	Arg	Val	
				245					250					255	

<210> SEQ ID NO 59

<211> LENGTH: 765

<212> TYPE: DNA

<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 59

cactctctcc tgcacggaga tgcggctctc tttgccgtcc tcctgggtga cgggctggca	60
cccatgtaag gtctcttgca cggagcaat ggagtgcggg ggcataggaa attcctccaa	120
gtcctccaca gggctctgat acataacctt attctctggc ttcttgacca cctttcggat	180
gcaggcagac accaacagga caacaagcag gatccccata atgatgggga ccaccactag	240
ggctcttaac cgaggctggg gaccacagat aacatcagtc ttgttagttc ccgcctgaac	300
cttcaccagg cctttggttt cacagcttgt ccaagggtga cacttttcca aagcagaaga	360
cacattggag aagaagccga tggggcaggg atcgcagatg gtatcagaaa tccctgtagc	420
gatctgtttg actcccaggc caggggggca caggctgtgc atggtgcagc tctcacaggc	480
agcgttggtg caatgcagac cttcatcgca tgtgcaagtg gtgtctgttt ctgacgtgcc	540
ctccttctcg acatggagcc ctagggtggg gtcgcagtat ttgtgctggt gacagtgtct	600
ctctgcgttc caagtgtcta ggaattcgcc tgtttggcaa cgagtgcatt ccgtgtcaat	660
ggtagttagg cagtcattca ccagtttctc tcctgggtggg cacatattac agcactgact	720
gtctactagg tattgctttt ctctgcatgc agtgcggtgt tctgg	765

<210> SEQ ID NO 60

<211> LENGTH: 336

<212> TYPE: DNA

<213> ORGANISM: Felis catus

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (1)..(336)

<400> SEQUENCE: 60

aat	gtg	tca	tct	gct	tcg	gaa	aag	tgt	cac	cct	tg	acg	agg	tgt	gag
Asn	Val	Ser	Ser	Ala	Ser	Glu	Lys	Cys	His	Pro	Trp	Thr	Arg	Cys	Glu
1				5					10					15	
acc	aaa	ggc	ctg	gtg	gag	ctt	cag	gcg	ggg	acc	aac	aag	acg	gat	gcc
Thr	Lys	Gly	Leu	Val	Glu	Leu	Gln	Ala	Gly	Thr	Asn	Lys	Thr	Asp	Ala
			20					25					30		
gtc	tcg	ggg	ttc	cag	gat	cg	ata	aga	gcc	ctg	gtg	gtg	atc	ccc	atc
Val	Cys	Gly	Phe	Gln	Asp	Arg	Ile	Arg	Ala	Leu	Val	Val	Ile	Pro	Ile
		35					40					45			
acg	atg	gtg	gtc	ctg	ctt	gct	gtc	ttg	ttg	gtg	tct	gcg	tat	atc	aga
Thr	Met	Val	Val	Leu	Leu	Ala	Val	Leu	Leu	Val	Ser	Ala	Tyr	Ile	Arg
		50				55				60					
aag	gtg	acc	aag	aag	cca	gag	aat	aag	gtc	ctc	cag	cct	aag	gct	gtg
Lys	Val	Thr	Lys	Lys	Pro	Glu	Asn	Lys	Val	Leu	Gln	Pro	Lys	Ala	Val
		65			70				75					80	

-continued

tcg cag gac cct gtg gag gac ttg gag gtc ctt cct gtc ccc ctc cac 288
 Ser Gln Asp Pro Val Glu Asp Leu Glu Val Leu Pro Val Pro Leu His
 85 90 95

ccc att gct ccg gtg cag gag acc tta cac ggg tgc cag ccg gtc acc 336
 Pro Ile Ala Pro Val Gln Glu Thr Leu His Gly Cys Gln Pro Val Thr
 100 105 110

<210> SEQ ID NO 61
 <211> LENGTH: 112
 <212> TYPE: PRT
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 61

Asn Val Ser Ser Ala Ser Glu Lys Cys His Pro Trp Thr Arg Cys Glu
 1 5 10 15
 Thr Lys Gly Leu Val Glu Leu Gln Ala Gly Thr Asn Lys Thr Asp Ala
 20 25 30
 Val Cys Gly Phe Gln Asp Arg Ile Arg Ala Leu Val Val Ile Pro Ile
 35 40 45
 Thr Met Val Val Leu Leu Ala Val Leu Leu Val Ser Ala Tyr Ile Arg
 50 55 60
 Lys Val Thr Lys Lys Pro Glu Asn Lys Val Leu Gln Pro Lys Ala Val
 65 70 75 80
 Ser Gln Asp Pro Val Glu Asp Leu Glu Val Leu Pro Val Pro Leu His
 85 90 95
 Pro Ile Ala Pro Val Gln Glu Thr Leu His Gly Cys Gln Pro Val Thr
 100 105 110

<210> SEQ ID NO 62
 <211> LENGTH: 336
 <212> TYPE: DNA
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 62

ggtgaccggc tggcaccggt gtaaggtctc ctgcaccgga gcaatggggt ggagggggac 60
 aggaaggacc tccaagtctc ccacagggtc ctgcgacaca gccttaggct ggaggacctt 120
 attctctggc ttcttgggtca cttttctgat atacgcagac accaacaaga cagcaagcag 180
 gaccaccatc gtgatgggga tcaccaccag ggctcttctc cgatcctgga aaccgcagac 240
 ggcatccgtc ttgttgggtc ccgcctgaag ctccaccagg cctttgggtc caccctcgt 300
 ccaagggtga cacttttccg aagcagatga cacatt 336

<210> SEQ ID NO 63
 <211> LENGTH: 390
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 63

ataagtgagg ctagttagtaa ccagcgtcc gttctgcggt gggcgccaaa aggggtactac 60
 accataagca gcaacctggt gagcctcgag aatgggaaac agttggccgt gaaaagacaa 120
 ggactctatt acgtctatgc ccaagtcacc ttctgctcca atcgggcagc ttcgagtcaa 180
 gctccgttcg tcgccagcct atgcctccat tccccgagtg gaacggagag agtcttactc 240
 cgcgccgcga gctcccgcgg ctctgccaaa ccttgcggcc aacagtccat ccacttgggga 300
 ggagtatttg aattgcatcc aggtgcttcg gtgttcgtca acgtgactga tccaagccaa 360
 gtgagccacg ggaccggctt cactctttt 390

-continued

```

<210> SEQ ID NO 64
<211> LENGTH: 1878
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (284)..(1063)

<400> SEQUENCE: 64

aatgtatgga agaagaaact tgtttcttct ttactaacia aagggaaagc ctggaagtga      60
atgatatggg tataattaaa aaaaaaaaaa aaaaaaaaaa aaaaccttta cgtaactttt      120
tttgctggga gagaagacta cgaagcacat tttccaggaa gtgtgggctg caacgattgt      180
gcgctcttaa ctaatcctga gtaaggtggc cactttgaca gtgttttcat gctgcctctg      240
ccaccttctc ggtctgaaga tatcatttca actctaacac agc atg atc gaa aca      295
                                         Met Ile Glu Thr
                                         1

tat agc caa act gct ccc cga tct gtg gcc act gga cca ccc gtc agt      343
Tyr Ser Gln Thr Ala Pro Arg Ser Val Ala Thr Gly Pro Pro Val Ser
   5              10              15              20

atg aaa att ttt atg tat ttg ctt act gtt ttt ctc atc acc cag atg      391
Met Lys Ile Phe Met Tyr Leu Leu Thr Val Phe Leu Ile Thr Gln Met
              25              30              35

att gga tcg gca ctc ttt gct gta tat ctt cac aga aga ttg gac aag      439
Ile Gly Ser Ala Leu Phe Ala Val Tyr Leu His Arg Arg Leu Asp Lys
              40              45              50

ata gaa gat gaa agg aat ctt tat gaa gat ttt gtg ttc atg aaa acg      487
Ile Glu Asp Glu Arg Asn Leu Tyr Glu Asp Phe Val Phe Met Lys Thr
              55              60              65

tta cag aaa tgc aac aaa ggg gag ggg tcc ttg tcc tta ctg aac tgt      535
Leu Gln Lys Cys Asn Lys Gly Glu Gly Ser Leu Ser Leu Leu Asn Cys
              70              75              80

gag gaa att aaa agc caa ttt gaa gcc ttt ctc aag gag ata atg cta      583
Glu Glu Ile Lys Ser Gln Phe Glu Ala Phe Leu Lys Glu Ile Met Leu
              85              90              95              100

aac aac gaa atg aag aaa gaa gaa aac att gca atg caa aaa ggt gat      631
Asn Asn Glu Met Lys Lys Glu Glu Asn Ile Ala Met Gln Lys Gly Asp
              105              110              115

cag gat cct cga att gca gcc cat gtc ata agt gag gct agt agt aac      679
Gln Asp Pro Arg Ile Ala Ala His Val Ile Ser Glu Ala Ser Ser Asn
              120              125              130

cca gcg tcc gtt ctg cgg tgg gcg cca aaa ggg tac tac acc ata agc      727
Pro Ala Ser Val Leu Arg Trp Ala Pro Lys Gly Tyr Tyr Thr Ile Ser
              135              140              145

agc aac ctg gtg agc ctc gag aat ggg aaa cag ttg gcc gtg aaa aga      775
Ser Asn Leu Val Ser Leu Glu Asn Gly Lys Gln Leu Ala Val Lys Arg
              150              155              160

caa gga ctc tat tac gtc tat gcc caa gtc acc ttc tgc tcc aat cgg      823
Gln Gly Leu Tyr Tyr Val Tyr Ala Gln Val Thr Phe Cys Ser Asn Arg
              165              170              175              180

gca gct tcg agt caa gct ccg ttc gtc gcc agc cta tgc ctc cat tcc      871
Ala Ala Ser Ser Gln Ala Pro Phe Val Ala Ser Leu Cys Leu His Ser
              185              190              195

ccg agt gga acg gag aga gtc tta ctc cgc gcc gcg agc tcc cgc ggc      919
Pro Ser Gly Thr Glu Arg Val Leu Leu Arg Ala Ala Ser Ser Arg Gly
              200              205              210

tcg tcc aaa cct tgc ggc caa cag tcc atc cac ttg gga gga gta ttt      967
Ser Ser Lys Pro Cys Gly Gln Gln Ser Ile His Leu Gly Gly Val Phe
              215              220              225

```

-continued

```

gaa ttg cat cca ggt gct tcg gtg ttc gtc aac gtg act gat cca agc 1015
Glu Leu His Pro Gly Ala Ser Val Phe Val Asn Val Thr Asp Pro Ser
    230                235                240

caa gtg agc cac ggg acc ggc ttc acg tct ttt ggc tta ctc aaa ctc 1063
Gln Val Ser His Gly Thr Gly Phe Thr Ser Phe Gly Leu Leu Lys Leu
    245                250                255                260

tgagtgtctgg caccctcacag gctgcagctc agctcctgtt ggtggtcttc gtaatacggc 1123

cgagcagtta agaccaccac ccctgttgaa ctgcctatctt ataaccctag gatcctcctc 1183

gtggagaact atttattata cccccccagg cgtggagggc tgcaagaagg gaatgacagg 1243

gcgggggcag cgccaacagg ccccggtcgg taagagttga tattctggaa gcagccgccc 1303

cactgatgca gacatccaga gagtcccatg aaaaagacga gactattatg cacagattga 1363

atcctcagta aacggcagat aattagtcca gtttcgtttt gtttctttgc atgcagtgtc 1423

tttctactga gaatgtactc gatttccccg cgaagatgct gaagggcaac agggagcctc 1483

agctcacagt cagttacggt tgaccggggg tccccggggc cccgatggag gggacaggct 1543

ccagaaaagtc tgatggcgcg gagaactgga aaaccctgcc cccaccagcc accctgacac 1603

tcattctctc cctcctccgc cccccctccc ccacagtcag gctgttgcta atcggttatc 1663

ttatttcaac cctgttgctt ctccaccagt gtaggcggga ggagagagca gaggctgccc 1723

actcctcctc ctgaaatgac tgtattttaa ggaaatctct cctacctacc tgcagtctcc 1783

attgtttcca gagtgaactt gtgattatct tgttatttat ttttgaata ataaagcgcc 1843

cttaacgtta aaaaaaaaaa aaaaaaaaaa aaaaaa 1878

```

```

<210> SEQ ID NO 65
<211> LENGTH: 260
<212> TYPE: PRT
<213> ORGANISM: Canis familiaris

```

```

<400> SEQUENCE: 65

```

```

Met Ile Glu Thr Tyr Ser Gln Thr Ala Pro Arg Ser Val Ala Thr Gly
  1             5             10            15

Pro Pro Val Ser Met Lys Ile Phe Met Tyr Leu Leu Thr Val Phe Leu
    20            25            30

Ile Thr Gln Met Ile Gly Ser Ala Leu Phe Ala Val Tyr Leu His Arg
    35            40            45

Arg Leu Asp Lys Ile Glu Asp Glu Arg Asn Leu Tyr Glu Asp Phe Val
    50            55            60

Phe Met Lys Thr Leu Gln Lys Cys Asn Lys Gly Glu Gly Ser Leu Ser
    65            70            75            80

Leu Leu Asn Cys Glu Glu Ile Lys Ser Gln Phe Glu Ala Phe Leu Lys
    85            90            95

Glu Ile Met Leu Asn Asn Glu Met Lys Lys Glu Glu Asn Ile Ala Met
    100           105           110

Gln Lys Gly Asp Gln Asp Pro Arg Ile Ala Ala His Val Ile Ser Glu
    115           120           125

Ala Ser Ser Asn Pro Ala Ser Val Leu Arg Trp Ala Pro Lys Gly Tyr
    130           135           140

Tyr Thr Ile Ser Ser Asn Leu Val Ser Leu Glu Asn Gly Lys Gln Leu
    145           150           155           160

Ala Val Lys Arg Gln Gly Leu Tyr Tyr Val Tyr Ala Gln Val Thr Phe
    165           170           175

Cys Ser Asn Arg Ala Ala Ser Ser Gln Ala Pro Phe Val Ala Ser Leu

```

-continued

180	185	190	
Cys Leu His Ser Pro Ser Gly Thr Glu Arg Val Leu Leu Arg Ala Ala			
195	200	205	
Ser Ser Arg Gly Ser Ser Lys Pro Cys Gly Gln Gln Ser Ile His Leu			
210	215	220	
Gly Gly Val Phe Glu Leu His Pro Gly Ala Ser Val Phe Val Asn Val			
225	230	235	240
Thr Asp Pro Ser Gln Val Ser His Gly Thr Gly Phe Thr Ser Phe Gly			
245	250	255	
Leu Leu Lys Leu			
260			
<210> SEQ ID NO 66			
<211> LENGTH: 1878			
<212> TYPE: DNA			
<213> ORGANISM: Canis familiaris			
<400> SEQUENCE: 66			
tttttttttt tttttttttt ttttttaacg ttaagggcgc tttattattc aaaaaataaa	60		
taacaagata atcacaagtt cactctggaa acaatggaga ctgcaggtag gtaggagaga	120		
tttcctttaa atacagtcac ttcaggagga ggagtgggca gcctctgctc tctcctcccg	180		
cctacactgg tggagaggca acaggggtga aataagataa ccgattagca acagcctgac	240		
tgtgggggga ggggggcgga ggagggagag aatgagtgtc aggggtggctg gtgggggcag	300		
ggttttccag ttctccgcgc catcagactt totggagcct gtcccctcca tcggggcccc	360		
ggggaccctg ggtcaaccgt aactgactgt gagctgagga tccctgttgc ccttcagcat	420		
cttcgcgggg aaatcgagta cattctccag tgaaagacac tgcattgaaa gaaacaaaac	480		
gaaactggac taattatctg ccgtttactg aggattcaat ctgtgcataa tagtctcgtc	540		
tttttcatgg gactctctgg atgtctgcat cagtggggcg gctgcttcca gaatatcaac	600		
tcttaccgac cggggcctgt tggcgctgcc ccgcctctgt cattcccttc ttgcagccct	660		
ccacgcctgg ggggtgataa taaatagttc tccacgagga ggatcctagg gttataaata	720		
ggcagttcaa caggggtggt ggtcttaact gctcggccgt attacgaaga ccaccaacag	780		
gagctgagct gcagcctgtg aggtgccagc actcagagtt tgagtaagcc aaaagacgtg	840		
aagccgggtcc cgtggctcac ttggcttgga tcagtcacgt tgacgaacac cgaagcacct	900		
ggatgcaatt caaatactcc tcccaagtgg atggactgtt ggcgcgaagg tttggacgag	960		
ccgcggggagc tcgcggcgcg gagtaagact ctctccgttc cactcgggga atggaggcat	1020		
aggctggcga cgaacggagc ttgactcgaa gctgcccgat tggagcagaa ggtgacttgg	1080		
gcatagacgt aatagagtcc ttgtcttttc acggccaact gtttccatt ctcgaggctc	1140		
accaggttgc tgcttatggt gtagtaccct tttggcgccc accgcagaac ggaagctggg	1200		
ttactactag cctcacttat gacatgggct gcaattcgag gatcctgatc acctttttgc	1260		
attgcaatgt tttcttcttt cttcatttcg ttgttttagca ttatctcctt gagaaaggct	1320		
tcaaattggc ttttaatttc ctcacagttc agtaaggaca aggacccctc ccctttgttg	1380		
catttctgta acgttttcat gaacacaaaa tottcataaa gattcctttc atcttctatc	1440		
ttgtccaatc ttctgtgaag atatacagca aagagtgcg atccaatcat ctgggtgatg	1500		
agaaaaacag taagcaaata cataaaaatt ttcatactga cgggtgtgccc agtggccaca	1560		
gatcggggag cagtttggct atatgtttcg atcatgctgt gttagagttg aaatgatatc	1620		

-continued

ttcagaccga gaaggtggca gaggcagcat gaaaacactg tcaaagtggc caccttactc	1680
aggattagtt aagagcgcac aatcgttgca gccacactt cctggaaaat gtgcttcgta	1740
gtcttctctc ccagcaaaaa aagttacgta aaggtttttt tttttttttt tttttttttt	1800
taattatacc catatcattc acttccaggc tttccctttt gttagtaaag aagaaacaag	1860
tttcttcttc catacatt	1878

<210> SEQ ID NO 67
 <211> LENGTH: 780
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 67

atgatcgaaa catatagcca aactgctccc cgatctgtgg ccaactggacc acccgtcagt	60
atgaaaattt ttatgtattt gcttactgtt tttctcatca ccagatgat tggatcggca	120
ctctttgctg tatatcttca cagaagattg gacaagatag aagatgaaag gaatctttat	180
gaagattttg tgttcataaa aacgttacag aaatgcaaca aaggggaggg gtccttgctc	240
ttactgaact gtgaggaaat taaaagccaa tttgaagcct tttcaagga gataatgcta	300
aaacacgaaa tgaagaaaga agaaaacatt gcaatgcaaa aaggtgatca ggatcctcga	360
attgcagccc atgtcataag tgaggctagt agtaaccag cgtccgttct gcggtggcg	420
ccaaaagggg actacaccat aagcagcaac ctggtgagcc tcgagaatgg gaaacagttg	480
gccgtgaaaa gacaaggact ctattacgtc tatgcccaag tcaccttctg ctccaatcgg	540
gcagcttcga gtcaagctcc gttcgtcgcc agcctatgcc tccattcccc gagtggaacg	600
gagagagtct tactccgcgc cgcgagctcc cgcggctcgt ccaaaccctg cggccaacag	660
tccatccact tgggaggagt atttgaattg catccagtg cttcggtgtt cgtcaacgtg	720
actgatccaa gccaaagtga ccacgggacc ggcttcacgt cttttggctt actcaaaactc	780

<210> SEQ ID NO 68
 <211> LENGTH: 780
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 68

gagtttgagt aagccaaaag acgtgaagcc ggtcccgtgg ctacttggc ttggatcagt	60
cacgttgacg aacaccgaag cacctggatg caattcaaat actcctccca agtggatgga	120
ctgttggccg caaggttttg acgagccgcg ggagctcgcg gcgcggagta agactctctc	180
cgttccactc ggggaatgga ggcataggct ggcgacgaac ggagcttgac tcgaagctgc	240
ccgattggag cagaagtgta cttgggcata gacgtaatag agtccctgtc ttttcacggc	300
caactgtttc ccattctcga ggctcaccag gttgctgctt atggtgtagt acccttttgg	360
cgccaccgcg agaacggacg ctgggttact actagcctca cttatgacat gggctgcaat	420
tcgaggatcc tgatcacctt ttgcatgtc aatgttttct tctttcttca ttctgtgtt	480
tagcattatc tccttgagaa aggcttcaaa ttggctttta atttctcac agttcagtaa	540
ggacaaggac ccctccctt ttgttcattt ctgtaacgtt ttcataaaca caaaatcttc	600
ataaagattc ctttcatctt ctatctgtgc caatcttctg tgaagatata cagcaaaagag	660
tgccgatcca atcatctggg tgatgagaaa aacagtaagc aaatacataa aaattttcat	720
actgacgggt ggtccagtgg ccacagatcg gggagcagtt tggctatatg ttccgatcat	780

-continued

```

<210> SEQ ID NO 69
<211> LENGTH: 633
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(633)

<400> SEQUENCE: 69

ttg gac aag ata gaa gat gaa agg aat ctt tat gaa gat ttt gtg ttc      48
Leu Asp Lys Ile Glu Asp Glu Arg Asn Leu Tyr Glu Asp Phe Val Phe
   1             5             10             15

atg aaa acg tta cag aaa tgc aac aaa ggg gag ggg tcc ttg tcc tta      96
Met Lys Thr Leu Gln Lys Cys Asn Lys Gly Glu Gly Ser Leu Ser Leu
   20             25             30

ctg aac tgt gag gaa att aaa agc caa ttt gaa gcc ttt ctc aag gag     144
Leu Asn Cys Glu Glu Ile Lys Ser Gln Phe Glu Ala Phe Leu Lys Glu
   35             40             45

ata atg cta aac aac gaa atg aag aaa gaa gaa aac att gca atg caa     192
Ile Met Leu Asn Asn Glu Met Lys Lys Glu Glu Asn Ile Ala Met Gln
   50             55             60

aaa ggt gat cag gat cct cga att gca gcc cat gtc ata agt gag gct     240
Lys Gly Asp Gln Asp Pro Arg Ile Ala Ala His Val Ile Ser Glu Ala
   65             70             75             80

agt agt aac cca gcg tcc gtt ctg cgg tgg gcg cca aaa ggg tac tac     288
Ser Ser Asn Pro Ala Ser Val Leu Arg Trp Ala Pro Lys Gly Tyr Tyr
   85             90             95

acc ata agc agc aac ctg gtg agc ctc gag aat ggg aaa cag ttg gcc     336
Thr Ile Ser Ser Asn Leu Val Ser Leu Glu Asn Gly Lys Gln Leu Ala
   100            105            110

gtg aaa aga caa gga ctc tat tac gtc tat gcc caa gtc acc ttc tgc     384
Val Lys Arg Gln Gly Leu Tyr Tyr Val Tyr Ala Gln Val Thr Phe Cys
   115            120            125

tcc aat cgg gca gct tcg agt caa gct ccg ttc gtc gcc agc cta tgc     432
Ser Asn Arg Ala Ala Ser Ser Gln Ala Pro Phe Val Ala Ser Leu Cys
   130            135            140

ctc cat tcc ccg agt gga acg gag aga gtc tta ctc cgc gcc gcg agc     480
Leu His Ser Pro Ser Gly Thr Glu Arg Val Leu Leu Arg Ala Ala Ser
   145            150            155            160

tcc cgc ggc tcg tcc aaa cct tgc ggc caa cag tcc atc cac ttg gga     528
Ser Arg Gly Ser Ser Lys Pro Cys Gly Gln Gln Ser Ile His Leu Gly
   165            170            175

gga gta ttt gaa ttg cat cca ggt gct tcg gtg ttc gtc aac gtg act     576
Gly Val Phe Glu Leu His Pro Gly Ala Ser Val Phe Val Asn Val Thr
   180            185            190

gat cca agc caa gtg agc cac ggg acc ggc ttc acg tct ttt ggc tta     624
Asp Pro Ser Ser Gln Val Ser His Gly Thr Gly Phe Thr Ser Phe Gly Leu
   195            200            205

ctc aaa ctc
Leu Lys Leu
   210

```

```

<210> SEQ ID NO 70
<211> LENGTH: 211
<212> TYPE: PRT
<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 70

Leu Asp Lys Ile Glu Asp Glu Arg Asn Leu Tyr Glu Asp Phe Val Phe
   1             5             10             15

Met Lys Thr Leu Gln Lys Cys Asn Lys Gly Glu Gly Ser Leu Ser Leu
   20             25             30

```

-continued

Leu Asn Cys Glu Glu Ile Lys Ser Gln Phe Glu Ala Phe Leu Lys Glu
 35 40 45
 Ile Met Leu Asn Asn Glu Met Lys Lys Glu Glu Asn Ile Ala Met Gln
 50 55 60
 Lys Gly Asp Gln Asp Pro Arg Ile Ala Ala His Val Ile Ser Glu Ala
 65 70 75 80
 Ser Ser Asn Pro Ala Ser Val Leu Arg Trp Ala Pro Lys Gly Tyr Tyr
 85 90 95
 Thr Ile Ser Ser Asn Leu Val Ser Leu Glu Asn Gly Lys Gln Leu Ala
 100 105 110
 Val Lys Arg Gln Gly Leu Tyr Tyr Val Tyr Ala Gln Val Thr Phe Cys
 115 120 125
 Ser Asn Arg Ala Ala Ser Ser Gln Ala Pro Phe Val Ala Ser Leu Cys
 130 135 140
 Leu His Ser Pro Ser Gly Thr Glu Arg Val Leu Leu Arg Ala Ala Ser
 145 150 155 160
 Ser Arg Gly Ser Ser Lys Pro Cys Gly Gln Gln Ser Ile His Leu Gly
 165 170 175
 Gly Val Phe Glu Leu His Pro Gly Ala Ser Val Phe Val Asn Val Thr
 180 185 190
 Asp Pro Ser Gln Val Ser His Gly Thr Gly Phe Thr Ser Phe Gly Leu
 195 200 205
 Leu Lys Leu
 210

<210> SEQ ID NO 71
 <211> LENGTH: 633
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 71

gagtttgagt aagccaaaag acgtgaagcc ggtcccgtgg ctcaacttggc ttggatcagt 60
 cacgttgacg aacaccgaag cacctggatg caattcaaact actcctccca agtggatgga 120
 ctgttgccg caaggttttg acgagccgcg ggagctcgcg gcgcggagta agactctctc 180
 cgttccactc ggggaatgga ggcataaggct ggcgacgaac ggagcttgac tcgaagctgc 240
 ccgattggag cagaaggtga cttgggcata gacgtaatag agtccttgtc ttttcacggc 300
 caactgtttc ccattctcga ggctcaccag gttgctgctt atggtgtagt acccttttgg 360
 cgcccaccgc agaacggacg ctgggttact actagcctca cttatgacat gggtgcgaat 420
 tcgaggatcc tgatcacctt tttgcattgc aatgttttct tctttcttca tttcgttggt 480
 tagcattatc tcottgagaa aggcttcaaa ttggctttta atttcctcac agttcagtaa 540
 ggacaaggac ccctcccctt tgttgcatth ctgtaacgtt ttcatgaaca caaaatcttc 600
 ataaagattc ctttcatctt ctatcttgctc caa 633

<210> SEQ ID NO 72
 <211> LENGTH: 885
 <212> TYPE: DNA
 <213> ORGANISM: Felis catus
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (29)..(808)

<400> SEQUENCE: 72

gaagataacca tttaactttt aacacagc atg atc gaa aca tat agc caa act 52

-continued

Met Ile Glu Thr Tyr Ser Gln Thr																100	
1								5									
gct	ccc	cgc	tcc	gtg	gcc	cct	gga	cca	ccc	gtc	agt	atg	aaa	att	ttt		
Ala	Pro	Arg	Ser	Val	Ala	Pro	Gly	Pro	Pro	Val	Ser	Met	Lys	Ile	Phe		
10						15				20							
atg	tat	tta	ctt	act	gtg	ttt	ctc	atc	acc	cag	atg	att	ggg	tca	gca		
Met	Tyr	Leu	Leu	Thr	Val	Phe	Leu	Ile	Thr	Gln	Met	Ile	Gly	Ser	Ala		
25				30						35				40			
ctc	ttt	gct	gtg	tat	ctt	cac	aga	aga	ctg	gac	aag	ata	gaa	gat	gaa		
Leu	Phe	Ala	Val	Tyr	Leu	His	Arg	Arg	Leu	Asp	Lys	Ile	Glu	Asp	Glu		
				45				50						55			
agg	aat	ctt	tat	gaa	gat	ttt	gtg	ttc	atg	aaa	aca	tta	cag	aaa	tgc		
Arg	Asn	Leu	Tyr	Glu	Asp	Phe	Val	Phe	Met	Lys	Thr	Leu	Gln	Lys	Cys		
		60						65				70					
aac	aaa	gga	gag	ggg	gcc	tta	tcc	tta	ctg	aac	tgt	gag	gaa	att	aaa		
Asn	Lys	Gly	Glu	Gly	Ala	Leu	Ser	Leu	Leu	Asn	Cys	Glu	Glu	Ile	Lys		
		75				80						85					
agc	cgg	ttt	gaa	gcc	ttt	ctc	aag	gag	ata	atg	cta	aac	aaa	gaa	acg		
Ser	Arg	Phe	Glu	Ala	Phe	Leu	Lys	Glu	Ile	Met	Leu	Asn	Lys	Glu	Thr		
90						95				100							
aag	aaa	gaa	aaa	aat	gtt	gca	atg	caa	aaa	ggc	gac	cag	gat	cct	cga		
Lys	Lys	Glu	Lys	Asn	Val	Ala	Met	Gln	Lys	Gly	Asp	Gln	Asp	Pro	Arg		
105				110						115				120			
gtt	gca	gca	cat	gtc	ata	agt	gag	gcc	agc	agt	agc	aca	gcg	tct	gtt		
Val	Ala	Ala	His	Val	Ile	Ser	Glu	Ala	Ser	Ser	Ser	Thr	Ala	Ser	Val		
				125				130						135			
ctc	cag	tgg	gcc	ccc	aaa	ggc	tac	tac	acc	ata	agc	agc	aac	ttg	gtg		
Leu	Gln	Trp	Ala	Pro	Lys	Gly	Tyr	Tyr	Thr	Ile	Ser	Ser	Asn	Leu	Val		
		140				145						150					
acc	ctc	gag	aac	ggg	aag	cag	ctg	gcc	gtt	aaa	aga	caa	gga	ctc	tat		
Thr	Leu	Glu	Asn	Gly	Lys	Gln	Leu	Ala	Val	Lys	Arg	Gln	Gly	Leu	Tyr		
		155				160						165					
tat	atc	tac	gcc	caa	gtc	acc	ttc	tgt	tcc	aat	cgg	gaa	gct	tcg	agt		
Tyr	Ile	Tyr	Ala	Gln	Val	Thr	Phe	Cys	Ser	Asn	Arg	Glu	Ala	Ser	Ser		
170						175				180							
caa	gct	ccg	ttc	ata	gcc	agc	ctc	tgc	ctg	cat	tcc	ccg	agt	gga	tcc		
Gln	Ala	Pro	Phe	Ile	Ala	Ser	Leu	Cys	Leu	His	Ser	Pro	Ser	Gly	Ser		
185				190						195				200			
gag	aga	gtc	tta	ctc	aga	gct	gca	aat	gcc	cgc	agt	tcc	tcc	aaa	ccc		
Glu	Arg	Val	Leu	Leu	Arg	Ala	Ala	Asn	Ala	Arg	Ser	Ser	Ser	Lys	Pro		
		205				210						215					
tgt	ggg	cag	caa	tcc	att	cac	ttg	gga	gga	gtc	ttc	gaa	ctg	cat	cca		
Cys	Gly	Gln	Gln	Ser	Ile	His	Leu	Gly	Gly	Val	Phe	Glu	Leu	His	Pro		
		220				225				230							
ggt	gct	tcg	gtg	ttc	gtg	aac	gtg	act	gat	ccg	agc	caa	gtg	agc	cac		
Gly	Ala	Ser	Val	Phe	Val	Asn	Val	Thr	Asp	Pro	Ser	Gln	Val	Ser	His		
235						240				245							
ggg	acg	ggc	ttc	acg	tct	ttt	ggc	ttg	ctc	aaa	ctc	tgaacactgg					
Gly	Thr	Gly	Phe	Thr	Ser	Phe	Gly	Leu	Leu	Lys	Leu						
250						255				260							
cacctgcgag gccgcgaggc ctgcaggccg cggtgagct cacgctggga gtcttcacaa																	878
tacagca																	885

<210> SEQ ID NO 73

<211> LENGTH: 260

<212> TYPE: PRT

<213> ORGANISM: Felis catus

<400> SEQUENCE: 73

-continued

tcctcacagt tcagtaagga taaggccccc tctcctttgt tgcattttctg taatgttttc	660
atgaacacaa aatcttcata aagatttcctt tcattcttcta tcttgtccag tcttctgtga	720
agatacacag caaagagtgc tgacccaatc atctgggtga tgagaaacac agtaagtaaa	780
tacataaaaa ttttcatact gacgggtggg ccagggggcca cggagcgggg agcagtttgg	840
ctatatgttt cgatcatgct gtgttaaagt tgaaatggta tcttc	885

<210> SEQ ID NO 75
 <211> LENGTH: 780
 <212> TYPE: DNA
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 75

atgatcgaaa catatagcca aactgctccc cgctccgtgg cccctggacc acccgtcagt	60
atgaaaaattt ttatgtattt acttactgtg tttctcatca ccagatgat tgggtcagca	120
ctctttgctg tgtatcttca cagaagactg gacaagatag aagatgaaag gaatctttat	180
gaagattttg tgttcataaa aacattacag aaatgcaaca aggagagagg ggccttatcc	240
ttactgaaat gtgaggaaat taaaagccgg tttgaagcct ttctcaagga gataatgcta	300
aacaaagaaa cgaagaaaga aaaaaatgtt gcaatgcaaa aaggcgacca ggatcctcga	360
gttgacgac atgtcataag tgaggccagc agtagcacag cgtctgttct ccagtgggcc	420
cccaaaggct actacaccat aagcagcaac ttggtgaccc tcgagaacgg gaagcagctg	480
gccgttaaaa gacaaggact ctattatatac tacgcccaag tcaccttctg ttccaatcgg	540
gaagcttcga gtcaagctcc gttcatagcc agcctctgcc tgcattcccc gagtggatcc	600
gagagagtct tactcagagc tgcaaatgcc cgcagttcct ccaaaccctg tgggcagcaa	660
tccattcact tgggaggagt cttcgaactg catccagggtg cttcggtgtt cgtgaacgtg	720
actgatccga gccaaagtga ccacgggacg ggcttcacgt cttttggctt gctcaaaactc	780

<210> SEQ ID NO 76
 <211> LENGTH: 780
 <212> TYPE: DNA
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 76

gagtttgagc aagccaaaag acgtgaagcc cgtcccgtgg ctcacttggc tcggatcagt	60
cacgttcacg aacaccgaag cacctggatg cagttcgaag actcctccca agtgaatgga	120
ttgtgcccc caggggtttg aggaactgcg ggcatattgca gctctgagta agactctctc	180
ggatccactc ggggaatgca ggagaggct ggctatgaac ggagcttgac tcgaagcttc	240
ccgattggaa cagaagtgta cttggggcta gatataatag agtccttgtc ttttaacggc	300
cagctgcttc ccgttctcga gggtcaccaa gttgctgctt atggtgtagt agcctttggg	360
ggcccaactg agaacagacg ctgtgctact gctggcctca cttatgacat gtgctgcaac	420
tcgaggatcc tggtcgcctt ttgtcattgc aacatttttt tctttctctg tttctttgtt	480
tagcattatc tccttgagaa aggcttcaaa ccggctttta atttctctac agttcagtaa	540
ggataaggcc ccctctcctt tgttgcatct ctgtaatgtt ttcataaaca caaaatcttc	600
ataaagattc ctttcattct ctatcttgct cagtcctctg tgaagatata cagcaaaagag	660
tgctgaccca atcatctggg tgatgagaaa cacagtaagt aaatacataa aaattttcat	720
actgacgggt ggtccagggg ccacggagcg gggagcagtt tggctatatg tttcgatcat	780

-continued

```

<210> SEQ ID NO 77
<211> LENGTH: 633
<212> TYPE: DNA
<213> ORGANISM: Felis catus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(633)

<400> SEQUENCE: 77

ctg gac aag ata gaa gat gaa agg aat ctt tat gaa gat ttt gtg ttc      48
Leu Asp Lys Ile Glu Asp Glu Arg Asn Leu Tyr Glu Asp Phe Val Phe
   1             5             10             15

atg aaa aca tta cag aaa tgc aac aaa gga gag ggg gcc tta tcc tta      96
Met Lys Thr Leu Gln Lys Cys Asn Lys Gly Glu Gly Ala Leu Ser Leu
   20             25             30

ctg aac tgt gag gaa att aaa agc cgg ttt gaa gcc ttt ctc aag gag      144
Leu Asn Cys Glu Glu Ile Lys Ser Arg Phe Glu Ala Phe Leu Lys Glu
   35             40             45

ata atg cta aac aaa gaa acg aag aaa gaa aaa aat gtt gca atg caa      192
Ile Met Leu Asn Lys Glu Thr Lys Lys Glu Lys Asn Val Ala Met Gln
   50             55             60

aaa ggc gac cag gat cct cga gtt gca gca cat gtc ata agt gag gcc      240
Lys Gly Asp Gln Asp Pro Arg Val Ala Ala His Val Ile Ser Glu Ala
   65             70             75             80

agc agt agc aca gcg tct gtt ctc cag tgg gcc ccc aaa ggc tac tac      288
Ser Ser Ser Thr Ala Ser Val Leu Gln Trp Ala Pro Lys Gly Tyr Tyr
   85             90             95

acc ata agc agc aac ttg gtg acc ctc gag aac ggg aag cag ctg gcc      336
Thr Ile Ser Ser Asn Leu Val Thr Leu Glu Asn Gly Lys Gln Leu Ala
   100            105            110

gtt aaa aga caa gga ctc tat tat atc tac gcc caa gtc acc ttc tgt      384
Val Lys Arg Gln Gly Leu Tyr Tyr Ile Tyr Ala Gln Val Thr Phe Cys
   115            120            125

tcc aat cgg gaa gct tcg agt caa gct ccg ttc ata gcc agc ctc tgc      432
Ser Asn Arg Glu Ala Ser Ser Gln Ala Pro Phe Ile Ala Ser Leu Cys
   130            135            140

ctg cat tcc ccg agt gga tcc gag aga gtc tta ctc aga gct gca aat      480
Leu His Ser Pro Ser Gly Ser Glu Arg Val Leu Leu Arg Ala Ala Asn
   145            150            155            160

gcc cgc agt tcc tcc aaa ccc tgt ggg cag caa tcc att cac ttg gga      528
Ala Arg Ser Ser Ser Lys Pro Cys Gly Gln Gln Ser Ile His Leu Gly
   165            170            175

gga gtc ttc gaa ctg cat cca ggt gct tcg gtg ttc gtg aac gtg act      576
Gly Val Phe Glu Leu His Pro Gly Ala Ser Val Phe Val Asn Val Thr
   180            185            190

gat ccg agc caa gtg agc cac ggg acg ggc ttc acg tct ttt ggc ttg      624
Asp Pro Ser Ser Gln Val Ser His Gly Thr Gly Phe Thr Ser Phe Gly Leu
   195            200            205

ctc aaa ctc
Leu Lys Leu
   210

```

```

<210> SEQ ID NO 78
<211> LENGTH: 211
<212> TYPE: PRT
<213> ORGANISM: Felis catus

<400> SEQUENCE: 78

Leu Asp Lys Ile Glu Asp Glu Arg Asn Leu Tyr Glu Asp Phe Val Phe
   1             5             10             15

Met Lys Thr Leu Gln Lys Cys Asn Lys Gly Glu Gly Ala Leu Ser Leu
   20             25             30

```

-continued

Leu Asn Cys Glu Glu Ile Lys Ser Arg Phe Glu Ala Phe Leu Lys Glu
 35 40 45
 Ile Met Leu Asn Lys Glu Thr Lys Lys Glu Lys Asn Val Ala Met Gln
 50 55 60
 Lys Gly Asp Gln Asp Pro Arg Val Ala Ala His Val Ile Ser Glu Ala
 65 70 75 80
 Ser Ser Ser Thr Ala Ser Val Leu Gln Trp Ala Pro Lys Gly Tyr Tyr
 85 90 95
 Thr Ile Ser Ser Asn Leu Val Thr Leu Glu Asn Gly Lys Gln Leu Ala
 100 105 110
 Val Lys Arg Gln Gly Leu Tyr Tyr Ile Tyr Ala Gln Val Thr Phe Cys
 115 120 125
 Ser Asn Arg Glu Ala Ser Ser Gln Ala Pro Phe Ile Ala Ser Leu Cys
 130 135 140
 Leu His Ser Pro Ser Gly Ser Glu Arg Val Leu Leu Arg Ala Ala Asn
 145 150 155 160
 Ala Arg Ser Ser Ser Lys Pro Cys Gly Gln Gln Ser Ile His Leu Gly
 165 170 175
 Gly Val Phe Glu Leu His Pro Gly Ala Ser Val Phe Val Asn Val Thr
 180 185 190
 Asp Pro Ser Gln Val Ser His Gly Thr Gly Phe Thr Ser Phe Gly Leu
 195 200 205
 Leu Lys Leu
 210

<210> SEQ ID NO 79
 <211> LENGTH: 633
 <212> TYPE: DNA
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 79

gagtttgagc aagccaaaag acgtgaagcc cgtcccgtgg ctcaactggc tcggatcagt 60
 cacgttcacg aacaccgaag cacctggatg cagttcgaag actcctccca agtgaatgga 120
 ttgctgcccc cagggttttg aggaactgcg ggcatttgca gctctgagta agactctctc 180
 ggatccactc ggggaatgca ggcagaggct ggctatgaac ggagcttgac tcgaagcttc 240
 ccgattggaa cagaagggtga cttgggcgta gatataatag agtccttgtc tttaaaggc 300
 cagctgcttc ccgttctcga gggtcaccaa gttgctgctt atggtgtagt agcctttggg 360
 ggcccactgg agaacagacg ctgtgtact gctggcctca cttatgacat gtgctgcaac 420
 tcgaggatcc tggtcgcctt tttgcattgc aacatttttt tctttcttcg tttctttggt 480
 tagcattatc tccttgagaa aggcttcaaa ccggctttta atttcctcac agttcagtaa 540
 ggataaggcc ccctctcctt tgttgcatth ctgtaatggt ttcatagaaca caaaatcttc 600
 ataaagattc ctttcatctt ctatcttgct cag 633

<210> SEQ ID NO 80
 <211> LENGTH: 610
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (29)..(430)

<400> SEQUENCE: 80

caaggcaaac actgaacatt tcagagct atg aga atg ctt ctg aat ttg agt 52

-continued

	Met	Arg	Met	Leu	Leu	Asn	Leu	Ser	
	1			5					
ttg cta gct ctt ggg gct gcc tat gtt tct gcc ttt gct gta gaa aat									100
Leu Leu Ala Leu Gly Ala Ala Tyr Val Ser Ala Phe Ala Val Glu Asn									
10 15 20									
ccc atg aat aga ctg gtg gca gag acc ttg aca ctg ctc tcc act cat									148
Pro Met Asn Arg Leu Val Ala Glu Thr Leu Thr Leu Leu Ser Thr His									
25 30 35 40									
cga act tgg ctg ata ggc gat ggg aac ctg atg att cct act cct gaa									196
Arg Thr Trp Leu Ile Gly Asp Gly Asn Leu Met Ile Pro Thr Pro Glu									
45 50 55									
aat aaa aat cac caa ctg tgc att aaa gaa gtt ttt cag ggt ata gac									244
Asn Lys Asn His Gln Leu Cys Ile Lys Glu Val Phe Gln Gly Ile Asp									
60 65 70									
aca ttg aag aac caa act gcc cac ggg gag gct gtg gat aaa cta ttc									292
Thr Leu Lys Asn Gln Thr Ala His Gly Glu Ala Val Asp Lys Leu Phe									
75 80 85									
caa aac ttg tct tta ata aaa gaa cac ata gag cgc caa aaa aaa agg									340
Gln Asn Leu Ser Leu Ile Lys Glu His Ile Glu Arg Gln Lys Lys Arg									
90 95 100									
tgt gca gga gaa aga tgg aga gtg aca aag ttc cta gac tac ctg caa									388
Cys Ala Gly Glu Arg Trp Arg Val Thr Lys Phe Leu Asp Tyr Leu Gln									
105 110 115 120									
gta ttt ctt ggt gta ata aac acc gag tgg aca ccg gaa agt									430
Val Phe Leu Gly Val Ile Asn Thr Glu Trp Thr Pro Glu Ser									
125 130									
tgagaacaaa ccggcttatt gtagtggaag attttggaaga agaatgggttt tttggcgatg									490
agaatgaggg ccaaccaaca gtagggactt aatggccagt ataactaagc ttcagagaca									550
aagtaaatat ttcaggcatc ctactacttt atcacttcac acagatgaaa tatatttgag									610

<210> SEQ ID NO 81
 <211> LENGTH: 134
 <212> TYPE: PRT
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 81

Met	Arg	Met	Leu	Leu	Asn	Leu	Ser	Leu	Leu	Ala	Leu	Gly	Ala	Ala	Tyr
1				5					10				15		
Val	Ser	Ala	Phe	Ala	Val	Glu	Asn	Pro	Met	Asn	Arg	Leu	Val	Ala	Glu
		20					25					30			
Thr	Leu	Thr	Leu	Leu	Ser	Thr	His	Arg	Thr	Trp	Leu	Ile	Gly	Asp	Gly
	35					40					45				
Asn	Leu	Met	Ile	Pro	Thr	Pro	Glu	Asn	Lys	Asn	His	Gln	Leu	Cys	Ile
	50				55					60					
Lys	Glu	Val	Phe	Gln	Gly	Ile	Asp	Thr	Leu	Lys	Asn	Gln	Thr	Ala	His
	65			70					75					80	
Gly	Glu	Ala	Val	Asp	Lys	Leu	Phe	Gln	Asn	Leu	Ser	Leu	Ile	Lys	Glu
		85					90					95			
His	Ile	Glu	Arg	Gln	Lys	Lys	Arg	Cys	Ala	Gly	Glu	Arg	Trp	Arg	Val
		100					105					110			
Thr	Lys	Phe	Leu	Asp	Tyr	Leu	Gln	Val	Phe	Leu	Gly	Val	Ile	Asn	Thr
		115				120					125				
Glu	Trp	Thr	Pro	Glu	Ser										
	130														

<210> SEQ ID NO 82
 <211> LENGTH: 610

-continued

<212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 82

```
ctcaaatata tttcatctgt gtgaagtgat aaagtagtag gatgcctgaa atatttactt    60
tgtctctgaa gcttagttat actggccatt aagtccttac tgttggttgg ccctcattct    120
catcgccaaa aaaccattct tctccaaaat ctccactac aataagccgg ttgtttctca    180
actttccggt gtccactcgg tgtttattac accaagaaat acttgcaggt agtctaggaa    240
ctttgtcact ctccatcttt ctccctgcaca cctttttttt tggcgctcta tgtgttcttt    300
tattaaagac aagttttgga atagttttatc cacagcctcc ccgtgggcag ttggttcttt    360
caatgtgtct ataccctgaa aaacttcttt aatgcacagt tggtgatttt tattttcagg    420
agtaggaatc atcaggttcc catcgccctat cagccaagtt cgatgagtgg agagcagtgt    480
caaggtctct gccaccagtc tattcatggg attttctaca gcaaaggcag aaacataggc    540
agccccaaga gctagcaaac tcaaattcag aagcattctc atagctctga aatgttcagt    600
gtttgccttg                                     610
```

<210> SEQ ID NO 83
 <211> LENGTH: 402
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 83

```
atgagaatgc ttctgaattt gagtttgcta gctcttgggg ctgcctatgt ttctgccttt    60
gctgtagaaa atcccatgaa tagactggtg gcagagacct tgacactgct ctccactcat    120
cgaacttggc tgataggcga tgggaacctg atgattccta ctctgaaaa taaaaatcac    180
caactgtgca ttaaagaagt ttttcaggg atagacacat tgaagaacca aactgcccac    240
ggggaggctg tggataaact attccaaaac ttgtctttaa taaagaaca catagagcgc    300
caaaaaaaaa ggtgtgcagg agaaagatgg agagtgacaa agttcctaga ctacctgcaa    360
gtatttcttg gtgtaataaa caccgagtg acaccggaaa gt                                     402
```

<210> SEQ ID NO 84
 <211> LENGTH: 402
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 84

```
actttccggt gtccactcgg tgtttattac accaagaaat acttgcaggt agtctaggaa    60
ctttgtcact ctccatcttt ctccctgcaca cctttttttt tggcgctcta tgtgttcttt    120
tattaaagac aagttttgga atagttttatc cacagcctcc ccgtgggcag ttggttcttt    180
caatgtgtct ataccctgaa aaacttcttt aatgcacagt tggtgatttt tattttcagg    240
agtaggaatc atcaggttcc catcgccctat cagccaagtt cgatgagtgg agagcagtgt    300
caaggtctct gccaccagtc tattcatggg attttctaca gcaaaggcag aaacataggc    360
agccccaaga gctagcaaac tcaaattcag aagcattctc at                                     402
```

<210> SEQ ID NO 85
 <211> LENGTH: 345
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(345)

-continued

<400> SEQUENCE: 85

```

ttt gct gta gaa aat ccc atg aat aga ctg gtg gca gag acc ttg aca      48
Phe Ala Val Glu Asn Pro Met Asn Arg Leu Val Ala Glu Thr Leu Thr
  1             5             10             15

ctg ctc tcc act cat cga act tgg ctg ata ggc gat ggg aac ctg atg      96
Leu Leu Ser Thr His Arg Thr Trp Leu Ile Gly Asp Gly Asn Leu Met
             20             25             30

att cct act cct gaa aat aaa aat cac caa ctg tgc att aaa gaa gtt     144
Ile Pro Thr Pro Glu Asn Lys Asn His Gln Leu Cys Ile Lys Glu Val
             35             40             45

ttt cag ggt ata gac aca ttg aag aac caa act gcc cac ggg gag gct     192
Phe Gln Gly Ile Asp Thr Leu Lys Asn Gln Thr Ala His Gly Glu Ala
             50             55             60

gtg gat aaa cta ttc caa aac ttg tct tta ata aaa gaa cac ata gag     240
Val Asp Lys Leu Phe Gln Asn Leu Ser Leu Ile Lys Glu His Ile Glu
             65             70             75             80

cgc caa aaa aaa agg tgt gca gga gaa aga tgg aga gtg aca aag ttc     288
Arg Gln Lys Lys Arg Cys Ala Gly Glu Arg Trp Arg Val Thr Lys Phe
             85             90             95

cta gac tac ctg caa gta ttt ctt ggt gta ata aac acc gag tgg aca     336
Leu Asp Tyr Leu Gln Val Phe Leu Gly Val Ile Asn Thr Glu Trp Thr
             100             105             110

ccg gaa agt
Pro Glu Ser
             115

```

<210> SEQ ID NO 86

<211> LENGTH: 115

<212> TYPE: PRT

<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 86

```

Phe Ala Val Glu Asn Pro Met Asn Arg Leu Val Ala Glu Thr Leu Thr
  1             5             10             15

Leu Leu Ser Thr His Arg Thr Trp Leu Ile Gly Asp Gly Asn Leu Met
             20             25             30

Ile Pro Thr Pro Glu Asn Lys Asn His Gln Leu Cys Ile Lys Glu Val
             35             40             45

Phe Gln Gly Ile Asp Thr Leu Lys Asn Gln Thr Ala His Gly Glu Ala
             50             55             60

Val Asp Lys Leu Phe Gln Asn Leu Ser Leu Ile Lys Glu His Ile Glu
             65             70             75             80

Arg Gln Lys Lys Arg Cys Ala Gly Glu Arg Trp Arg Val Thr Lys Phe
             85             90             95

Leu Asp Tyr Leu Gln Val Phe Leu Gly Val Ile Asn Thr Glu Trp Thr
             100             105             110

Pro Glu Ser
             115

```

<210> SEQ ID NO 87

<211> LENGTH: 345

<212> TYPE: DNA

<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 87

```

actttccggt gtccactcgg tgtttattac accaagaaat acttgcaggt agtctaggaa      60
ctttgtcact ctccatcttt ctccctgcaca cctttttttt tggcgctcta tgtgttcttt     120

```

-continued

tatttaaagac aagttttgga atagttttatc cacagcctcc ccgtgggcag tttggttctt	180
caatgtgtct ataccctgaa aaacttcttt aatgcacagt tggatgtttt ttttttcagg	240
agtaggaatc atcaggttcc catcgcttat cagccaagtt cgatgagtgg agagcagtgt	300
caaggctctct gccaccagtc tattcatggg attttctaca gcaaa	345

<210> SEQ ID NO 88
 <211> LENGTH: 166
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 88

ctcagcttag gccagcctac gacctgcctg ctcttccctc gtcctcctg cattggctct	60
gggtcccatg gcgctctggt tgactgtggt cattgctctc acctgcctcg gtggccttgc	120
ctccccgagc cctgtgactc cctccccaac cctcaaggag ctcat	166

<210> SEQ ID NO 89
 <211> LENGTH: 272
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 89

tggccttgcc tccccgagcc ctgtgactcc ctccccaacc ctcaaggagc tcattgagga	60
gctggtcaac atcaccaga atcaggcatc cctctgcaac ggcagcatgg tgtggagcgt	120
caacctgacc gccggcatgt actgcgcagc tctagaatct ctgatcaatg tctccgactg	180
cagcgccatc caaaggaccc agaggatgct gaaagcactg tgctctcaaa agcccgcggc	240
agggcagatt tccagtgaac gcagccgaga ca	272

<210> SEQ ID NO 90
 <211> LENGTH: 278
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 90

atggcgctct ggttgactgt ggtcattgct ctcacctgcc tcggtggcct tgcctccccg	60
agccctgtga ctccctcccc aacctcaag gagctcattg aggagctggt caacatcacc	120
cagaatcagg catccctctg caacggcagc atggtgtgga gcgtcaacct gaccgcggc	180
atgtactgcg cagctctaga atctctgac aatgtctcgg actgcagcgc catccaaagg	240
acccagagga tgctgaaagc actgtgctct caaaagcc	278

<210> SEQ ID NO 91
 <211> LENGTH: 1302
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (52)..(444)

<400> SEQUENCE: 91

ctacgacctg cctgctcttc cctcgctcct cctgcattgg ctctgggctc c atg gcg	57
	Met Ala
	1
ctc tgg ttg act gtg gtc att gct ctc acc tgc ctc ggt ggc ctt gcc	105
Leu Trp Leu Thr Val Val Ile Ala Leu Thr Cys Leu Gly Gly Leu Ala	
5 10 15	
tcc ccg agc cct gtg act ccc tcc cca acc ctc aag gag ctc att gag	153
Ser Pro Ser Pro Val Thr Pro Ser Pro Thr Leu Lys Glu Leu Ile Glu	

-continued

20	25	30	
gag ctg gtc aac atc acc cag aat cag gca tcc ctc tgc aac ggc agc			201
Glu Leu Val Asn Ile Thr Gln Asn Gln Ala Ser Leu Cys Asn Gly Ser			
35 40 45 50			
atg gtg tgg agc gtc aac ctg acc gcc ggc atg tac tgc gca gct cta			249
Met Val Trp Ser Val Asn Leu Thr Ala Gly Met Tyr Cys Ala Ala Leu			
55 60 65			
gaa tct ctg atc aat gtc tcc gac tgc agc gcc atc caa agg acc cag			297
Glu Ser Leu Ile Asn Val Ser Asp Cys Ser Ala Ile Gln Arg Thr Gln			
70 75 80			
agg atg ctg aaa gca ctg tgc tct caa aag ccc gcg gca ggg cag att			345
Arg Met Leu Lys Ala Leu Cys Ser Gln Lys Pro Ala Ala Gly Gln Ile			
85 90 95			
tcc agt gaa cgc agc cga gac acc aaa att gaa gtg atc cag ttg gtg			393
Ser Ser Glu Arg Ser Arg Asp Thr Lys Ile Glu Val Ile Gln Leu Val			
100 105 110			
aaa aac ctg ctc acc tat gta agg gga gtt tat cgc cat gga aat ttc			441
Lys Asn Leu Leu Thr Tyr Val Arg Gly Val Tyr Arg His Gly Asn Phe			
115 120 125 130			
aga tgaagcatga aaacttagca tccttatctg tagaccacaga cctgaccact			494
Arg			
taagttccag attcattttt ctttccgacg tcacaaattt cttagggagg tggggggggg			554
ggagaacccat ttccctcagct gggacccctcag cctgcaccgc ctgcctccat ggagctgagc			614
ccagccacccc ctgccttggt gcatggggcc cagccgggtg gccctcctcc gtctgcactt			674
catcaacgct gagggaaagc actgcatccc atgactgtcc cctcctcaga gcaaagtgca			734
gcattacagt ggaggcagat atgtgtggga gggggtcttg ctgtacctgg gagtggcaca			794
gacatgtttc ttcttagcct tatttattat tgtgtgttat ttaaacaagt gtctttgttt			854
gtgtgtggga cagggagtgg cttggagctg ggggccctg gactcgggtt tagagagtcc			914
ctgggaataa gcaactgtgtg taaaattctg ctacctcact gggatcctgg ggcgcacaca			974
ggggacagga gaaaggggtca gagatgtgc tcttgtctgc cactcagcag ctggccctca			1034
gccaaagcagt aatttattgt ttttccttgt atttaaagtt aagaaataaa atatgttatc			1094
aaagagttaa taatatatag aagagtagcc taaaaggctg catttggtgt gtgtggccag			1154
gccggggcgg gtggggggga ggggtgtgtc actgaatgtg ctctttcact gactttgtca			1214
aactggaagc cagaaataaa gatggtgaca agagaaaaaa aaaaaaaaaa aaaaaaaaaa			1274
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa			1302

<210> SEQ ID NO 92

<211> LENGTH: 131

<212> TYPE: PRT

<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 92

Met Ala Leu Trp Leu Thr Val Val Ile Ala Leu Thr Cys Leu Gly Gly
1 5 10 15

Leu Ala Ser Pro Ser Pro Val Thr Pro Ser Pro Thr Leu Lys Glu Leu
20 25 30

Ile Glu Glu Leu Val Asn Ile Thr Gln Asn Gln Ala Ser Leu Cys Asn
35 40 45

Gly Ser Met Val Trp Ser Val Asn Leu Thr Ala Gly Met Tyr Cys Ala
50 55 60

Ala Leu Glu Ser Leu Ile Asn Val Ser Asp Cys Ser Ala Ile Gln Arg
65 70 75 80

-continued

Thr Gln Arg Met Leu Lys Ala Leu Cys Ser Gln Lys Pro Ala Ala Gly
 85 90 95
 Gln Ile Ser Ser Glu Arg Ser Arg Asp Thr Lys Ile Glu Val Ile Gln
 100 105 110
 Leu Val Lys Asn Leu Leu Thr Tyr Val Arg Gly Val Tyr Arg His Gly
 115 120 125
 Asn Phe Arg
 130

<210> SEQ ID NO 93
 <211> LENGTH: 1302
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 93

```

tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt ttttctcttg      60
tcaccatctt tatttctggc ttccagtttg acaaagtcag tgaaagagca cattcagtga      120
caacaccctc cccccccacc gccccggcct ggccacacac accaaatgca gccttttagg      180
ctactcttct atatatattt aactctttga taacatatatt tatttcttaa ctttaaatac      240
aaggaaaaac aataaattac tgcttggtcg agggccagct gctgagtggtc agacaagagc      300
agcatctctg accctttctc ctgtccccctg tgtcggcccc aggatcccag tgaggtagca      360
gaattttaca cacagtgcct attcccaggg actctctaaa ccgagtcac tgggccccca      420
gctccaagcc actccctgtc ccagcacaca acaaagacac ttgtttaaat aacacacaat      480
aataaataag gctaagaaga aacatgtctg tgccactccc aggtacagca agacccctc      540
ccacacatat ctgcctccac tgtaatgctg cactttgctc tgaggagggtg acagtcatgg      600
gatgcagtg cttccctcag cgttgatgaa gtgcagacgg aggaggggcca cccggctggg      660
ccccatgcac caaggcaggg gtggctgggc tcagctccat ggaggcaggc ggtgcaggct      720
gagggtcccg ctgaggaaat ggttctcccc cccccccacc tccctaagaa atttgtgacg      780
tcggaagaaa aaatgaatct ggaacttaag tggtcaggtc tgggtctaca gataaggatg      840
ctaagttttc atgcttcctc tgaaatttcc atggcgataa actcccctta cataggtagg      900
cagggttttc accaactgga tcacttcaat ttgtgtgtct cggctgcgtt cactggaaat      960
ctgccctgcc gcgggctttt gagagcacag tgctttcagc atcctctggg tcctttggat      1020
ggcgtgcag tcggagacat tgatcagaga ttctagagct gcgcagtaca tgccggcggt      1080
caggttgacg ctccacacca tgctgccgtt gcagagggat gcctgattct ggtgatggtt      1140
gaccagctcc tcaatgagct ccttgagggt tggggaggga gtcacagggc tcggggaggc      1200
aaggccaccg aggcagggtg gagcaatgac cacagtcaac cagagcgcca tggagcccag      1260
agccaatgca ggaggagcga gggaagagca ggcaggctcg ag                                1302

```

<210> SEQ ID NO 94
 <211> LENGTH: 393
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 94

```

atggcgctct ggttgactgt ggtcattgct ctcacctgcc tcggtggcct tgctccccg      60
agcctgtgta ctccctcccc aacctcaag gagctcattg aggagctggt caacatcacc      120
cagaatcagg catccctctg caacggcagc atggtgtgga gcgtcaacct gaccgccggc      180

```

-continued

```

atgtactgcg cagctctaga atctctgac aatgtctccg actgcagcgc catccaaagg 240
acctcagagga tgctgaaagc actgtgctct caaaagcccg cggcagggca gatttccagt 300
gaacgcagcc gagacaccaa aattgaagtg atccagttgg tgaaaaacct gctcacctat 360
gtaaggggag tttatcgcca tggaaatttc aga 393

```

```

<210> SEQ ID NO 95
<211> LENGTH: 393
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris

```

```

<400> SEQUENCE: 95

```

```

tctgaaatct ccatggcgat aaactcccct tacataggtg agcaggtttt tcaccaactg 60
gatcacttca attttggtgt ctcggtgcg ttcactggaa atctgccctg ccgcgggctt 120
ttgagagcac agtgctttca gcatcctctg ggtcctttgg atggcgctgc agtcggagac 180
attgatcaga gattctagag ctgcgcagta catgccggcg gtcagggtga cgctccacac 240
catgctgccg ttgcagaggg atgctgatt ctgggtgatg ttgaccagct cctcaatgag 300
ctccttgagg gttggggagg gagtcacagg gctcggggag gcaaggccac cgaggcaggt 360
gagagcaatg accacagtca accagagcgc cat 393

```

```

<210> SEQ ID NO 96
<211> LENGTH: 333
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(333)

```

```

<400> SEQUENCE: 96

```

```

agc cct gtg act ccc tcc cca acc ctc aag gag ctc att gag gag ctg 48
Ser Pro Val Thr Pro Ser Pro Thr Leu Lys Glu Leu Ile Glu Glu Leu
1 5 10 15
gtc aac atc acc cag aat cag gca tcc ctc tgc aac ggc agc atg gtg 96
Val Asn Ile Thr Gln Asn Gln Ala Ser Leu Cys Asn Gly Ser Met Val
20 25 30
tgg agc gtc aac ctg acc gcc ggc atg tac tgc gca gct cta gaa tct 144
Trp Ser Val Asn Leu Thr Ala Gly Met Tyr Cys Ala Ala Leu Glu Ser
35 40 45
ctg atc aat gtc tcc gac tgc agc gcc atc caa agg acc cag agg atg 192
Leu Ile Asn Val Ser Asp Cys Ser Ala Ile Gln Arg Thr Gln Arg Met
50 55 60
ctg aaa gca ctg tgc tct caa aag ccc gcg gca ggg cag att tcc agt 240
Leu Lys Ala Leu Cys Ser Gln Lys Pro Ala Ala Gly Gln Ile Ser Ser
65 70 75 80
gaa cgc agc cga gac acc aaa att gaa gtg atc cag ttg gtg aaa aac 288
Glu Arg Ser Arg Asp Thr Lys Ile Glu Val Ile Gln Leu Val Lys Asn
85 90 95
ctg ctc acc tat gta agg gga gtt tat cgc cat gga aat ttc aga 333
Leu Leu Thr Tyr Val Arg Gly Val Tyr Arg His Gly Asn Phe Arg
100 105 110

```

```

<210> SEQ ID NO 97
<211> LENGTH: 111
<212> TYPE: PRT
<213> ORGANISM: Canis familiaris

```

```

<400> SEQUENCE: 97

```

```

Ser Pro Val Thr Pro Ser Pro Thr Leu Lys Glu Leu Ile Glu Glu Leu
1 5 10 15

```


Ser	Ser	Glu	Arg	Ser	Arg	Asp	Thr	Lys	Ile	Glu	Val	Ile	Gln	Leu	Val	
100							105			110						
aaa	aac	ctg	ctc	acc	tat	gta	agg	gga	gtt	tat	cgc	cat	gga	aat	ttc	443
Lys	Asn	Leu	Leu	Thr	Tyr	Val	Arg	Gly	Val	Tyr	Arg	His	Gly	Asn	Phe	
115							120			125						
aga	tga	agcatga	aaacttagca	tccttatctg	tagacc	caga	ctgacc	act								496
Arg																
130																
taagt	tccag	attcatt	ttt	ctt	ccgacg	tcacaa	attt	cttagg	gagg	tg	gggggggg					556
ggaga	aaccat	ttcctcag	ct	gggac	ctcag	cctgcac	cgc	ctgcct	ccat	ggag	ctgag					616
ccagc	acacc	ctgcct	tgg	gc	atggggc	cagccg	ggtg	gccctc	ctcc	gtctg	caact					676
catca	acgct	gaggg	aaagc	actgc	atccc	atgact	gtcc	cctcct	caga	gcaa	agtgca					736
gcatt	acagt	ggagg	cagat	atgtg	tggga	ggggg	tcttg	ctgtac	ctgg	gagt	ggcaca					796
gacat	gtttc	ttcttag	ctct	tattt	attat	tgtgt	gttat	ttaaa	caagt	gtctt	tgttt					856
gtgct	gggga	caggg	agtg	cttgg	agctg	ggggc	ccagt	gactc	gggt	taga	gagtc					916
ctgga	aataa	gcactg	tgtg	taaa	aattctg	ctacct	cact	gggat	cttg	ggccg	acaca					976
gggg	acagga	gaaagg	gtca	gagat	gtcgc	tctt	gtctgc	cactc	agcag	ctgg	ccctca					1036
gcca	agcagt	aattt	attgt	tttct	cttgt	attt	aaagtt	aagaa	aataaa	atat	gtttatc					1096
aaag	agttaa	taatat	atag	aagag	tagcc	taaa	aggctg	cattt	ggtgt	gtgtg	gccag					1156
gccg	gggcg	gtggg	ggga	gggt	gttgtc	actga	atgtg	ctctt	tact	gactt	gttca					1216
aact	ggaagc	cagaa	aataaa	gatg	gtgaca	agag	aaaaaa	aaaa	aaaaaa	aaa						1269

<400> SEQUENCE: 100

[illegible]

<400> SEQUENCE: 101

-continued

tttttttttt tttttttttt tcttgtcacc atcttttattt ctggcttcca gtttgacaaa	60
gtcagtgaaa gagcacattc agtgacaaca cctccccccc caccgcgccc ggctgggcca	120
cacacaccaa atgcagcctt ttaggtactt cttctatata ttattaactc tttgataaca	180
tattttattt ctaacttta aatacaagga aaaacaataa attactgctt ggctgagggc	240
cagctgctga gtggcagaca agagcagcat ctctgaccct ttctcctgtc cctgtgtctg	300
gccccaggat cccagtgagg tagcagaatt ttacacacag tgcttattcc cagggactct	360
ctaaacccga gtcactgggc ccccagctcc aagccactcc ctgtcccag cacaaacaaa	420
gacacttggt taaataacac acaataataa ataaggctaa gaagaaacat gtctgtgcca	480
ctcccaggta cagcaagacc ccctcccaca catactgtgc tccactgtaa tgctgcactt	540
tgctctgagg aggggacagt catgggatgc agtgctttcc ctcagcgttg atgaagtga	600
gacggaggag ggccaccggy ctgggcccga tgcaccaagg caggggtggc tgggctcagc	660
tccatggagg cagggcgtgc aggctgaggt cccagctgag gaaatgggtc tcccccccc	720
ccacctccct aagaaatttg tgacgtcgga aagaaaaatg aatctggaac ttaagtggtc	780
aggctctggg ctacagataa ggatgctaag ttttcatgct tcactctgaaa tttccatggc	840
gataaactcc ccttacatag gtgagcaggt ttttcaccaa ctggatcact tcaatttttg	900
tgctctggct gcgttactg gaaatccctg ccgcgggctt ttgagagcac agtgctttca	960
gcatcctctg ggtccttttg atggcgtgc agtcggagac attgatcaga gattctagag	1020
ctgcgcagta catgccggcg gtcaggttga cgctccacac catgctgccg ttgcagaggg	1080
atgctgatt ctgggtgatg ttgaccagct cctcaatgag ctccttgagg gttggggagg	1140
gagtcacagg gctcggggag gcaaggccac cgaggcaggt gagagcaatg accacagtca	1200
accagagcgc catggagccc agagccaatg caggaggagc gagggaagag caggcaggtc	1260
gtaggctgg	1269

<210> SEQ ID NO 102
 <211> LENGTH: 390
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 102

atggcgctct ggttgactgt ggtcattgct ctcacctgcc tcggtggcct tgcctccccg	60
agccctgtga ctccctcccc aacctcaag gagctcattg aggagctggg caacatcacc	120
cagaatcagg catccctctg caacggcagc atggtgtgga gcgtcaacct gaccgccggc	180
atgtactgcy cagctctaga atctctgac aatgtctccg actgcagcgc catccaaagg	240
acccagagga tgctgaaagc actgtgctct caaaagcccg cggcagggat ttccagtga	300
cgcagccgag acaccaaagt tgaagtgatc cagttggtga aaaacctgct cacctatgta	360
aggggagttt atcgccatgg aaatttcaga	390

<210> SEQ ID NO 103
 <211> LENGTH: 390
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 103

tctgaaattt ccatggcgat aaactccctt tacatagggt agcaggtttt tcaccaactg	60
gatcacttca attttggtgt ctcggctgcy ttcactggaa atccctgccg cgggcttttg	120

-continued

```

agagcacagt gctttcagca tcctctgggt cctttggatg gcgctgcagt cggagacatt 180
gatcagagat tctagagctg cgcagtacat gccggcggtc aggttgacgc tccacaccat 240
gtgcccgttg cagaggggatg cctgattctg ggtgatgttg accagctcct caatgagctc 300
cttgagggtt ggggaggag tcacagggt cggggaggca aggccaccga ggcaggtgag 360
agcaatgacc acagtcaacc agagcgccat 390

```

```

<210> SEQ ID NO 104
<211> LENGTH: 330
<212> TYPE: DNA
<213> ORGANISM: Canis familiaris
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(330)

```

```

<400> SEQUENCE: 104

```

```

agc cct gtg act ccc tcc cca acc ctc aag gag ctc att gag gag ctg 48
Ser Pro Val Thr Pro Ser Pro Thr Leu Lys Glu Leu Ile Glu Glu Leu
1 5 10 15

gtc aac atc acc cag aat cag gca tcc ctc tgc aac ggc agc atg gtg 96
Val Asn Ile Thr Gln Asn Gln Ala Ser Leu Cys Asn Gly Ser Met Val
20 25 30

tgg agc gtc aac ctg acc gcc ggc atg tac tgc gca gct cta gaa tct 144
Trp Ser Val Asn Leu Thr Ala Gly Met Tyr Cys Ala Ala Leu Glu Ser
35 40 45

ctg atc aat gtc tcc gac tgc agc gcc atc caa agg acc cag agg atg 192
Leu Ile Asn Val Ser Asp Cys Ser Ala Ile Gln Arg Thr Gln Arg Met
50 55 60

ctg aaa gca ctg tgc tct caa aag ccc gcg gca ggg att tcc agt gaa 240
Leu Lys Ala Leu Cys Ser Gln Lys Pro Ala Ala Gly Ile Ser Ser Glu
65 70 75 80

cgc agc cga gac acc aaa att gaa gtg atc cag ttg gtg aaa aac ctg 288
Arg Ser Arg Asp Thr Lys Ile Glu Val Ile Gln Leu Val Lys Asn Leu
85 90 95

ctc acc tat gta agg gga gtt tat cgc cat gga aat ttc aga 330
Leu Thr Tyr Val Arg Gly Val Tyr Arg His Gly Asn Phe Arg
100 105 110

```

```

<210> SEQ ID NO 105
<211> LENGTH: 110
<212> TYPE: PRT
<213> ORGANISM: Canis familiaris

```

```

<400> SEQUENCE: 105

```

```

Ser Pro Val Thr Pro Ser Pro Thr Leu Lys Glu Leu Ile Glu Glu Leu
1 5 10 15

Val Asn Ile Thr Gln Asn Gln Ala Ser Leu Cys Asn Gly Ser Met Val
20 25 30

Trp Ser Val Asn Leu Thr Ala Gly Met Tyr Cys Ala Ala Leu Glu Ser
35 40 45

Leu Ile Asn Val Ser Asp Cys Ser Ala Ile Gln Arg Thr Gln Arg Met
50 55 60

Leu Lys Ala Leu Cys Ser Gln Lys Pro Ala Ala Gly Ile Ser Ser Glu
65 70 75 80

Arg Ser Arg Asp Thr Lys Ile Glu Val Ile Gln Leu Val Lys Asn Leu
85 90 95

Leu Thr Tyr Val Arg Gly Val Tyr Arg His Gly Asn Phe Arg
100 105 110

```

-continued

<210> SEQ ID NO 106
 <211> LENGTH: 330
 <212> TYPE: DNA
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 106

tctgaaattt ccatggcgat aaactcccct tacataggtg agcagggttt tcaccaactg	60
gatacattca attttggtgt ctgcggtgcg ttacttgga atccctgccg cgggcttttg	120
agagcacagt gctttcagca tcctctgggt cctttggatg gcgctgcagt cggagacatt	180
gatcagagat tctagagctg cgcagtacat gccggcggtc aggttgacgc tccacacat	240
gctgccgttg cagagggatg cctgattctg ggtgatgttg accagctcct caatgagctc	300
cttgagggtt ggggaggag tcacagggt	330

<210> SEQ ID NO 107
 <211> LENGTH: 567
 <212> TYPE: DNA
 <213> ORGANISM: Felis catus
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(567)

<400> SEQUENCE: 107

atg gcg ctg ccc tct tcc ttc ttg gtg gcc ctg gtg gcg ctg ggc tgc	48
Met Ala Leu Pro Ser Ser Phe Leu Val Ala Leu Val Ala Leu Gly Cys	
1 5 10 15	
aac tcc gtc tgc tct ctg ggc tgt gac ctg cct cag acc cac ggc ctg	96
Asn Ser Val Cys Ser Leu Gly Cys Asp Leu Pro Gln Thr His Gly Leu	
20 25 30	
ctg aac agg agg gcc ttg acg ctc ctg gga caa atg agg aga ctc cct	144
Leu Asn Arg Arg Ala Leu Thr Leu Leu Gly Gln Met Arg Arg Leu Pro	
35 40 45	
gcc agc tcc tgt cag aag gac aga aat gac ttc gcc ttc ccc cag gac	192
Ala Ser Ser Cys Gln Lys Asp Arg Asn Asp Phe Ala Phe Pro Gln Asp	
50 55 60	
gtg ttt ggt gga gac cag tcc cac aag gcc caa gcc ctc tcg gtg gtg	240
Val Phe Gly Gly Asp Gln Ser His Lys Ala Gln Ala Leu Ser Val Val	
65 70 75 80	
cac gtg acg aac cag aag atc ttc cac ttc ttc tgc aca gag gcg tcc	288
His Val Thr Asn Gln Lys Ile Phe His Phe Phe Cys Thr Glu Ala Ser	
85 90 95	
tcg tct gct gct tgg aac acc acc ctc ctg gag gaa ttc tgc acg gga	336
Ser Ser Ala Ala Trp Asn Thr Thr Leu Leu Glu Glu Phe Cys Thr Gly	
100 105 110	
ctt gat tgg cag ctg acc cgc ctg gaa gcc tgt gtc atg cag gag gtg	384
Leu Asp Trp Gln Leu Thr Arg Leu Glu Ala Cys Val Met Gln Glu Val	
115 120 125	
ggg gag gga gag gct ccc ctc acg aac gag gac tcc atc ctg agg aac	432
Gly Glu Gly Glu Ala Pro Leu Thr Asn Glu Asp Ser Ile Leu Arg Asn	
130 135 140	
tac ttc caa aga ctc tcc ctc tac ctg caa gag aag aaa tac agc cct	480
Tyr Phe Gln Arg Leu Ser Leu Tyr Leu Gln Glu Lys Lys Tyr Ser Pro	
145 150 155 160	
tgt gcc tgg gag atc gtc aga gca gaa atc atg aga tcc ttg tat tat	528
Cys Ala Trp Glu Ile Val Arg Ala Glu Ile Met Arg Ser Leu Tyr Tyr	
165 170 175	
tca tca aca gcc ttg cag aaa aga tta agg agc gag aaa	567
Ser Ser Thr Ala Leu Gln Lys Arg Leu Arg Ser Glu Lys	
180 185	

-continued

<210> SEQ ID NO 108
 <211> LENGTH: 189
 <212> TYPE: PRT
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 108

```

Met Ala Leu Pro Ser Ser Phe Leu Val Ala Leu Val Ala Leu Gly Cys
 1             5             10             15
Asn Ser Val Cys Ser Leu Gly Cys Asp Leu Pro Gln Thr His Gly Leu
      20             25             30
Leu Asn Arg Arg Ala Leu Thr Leu Leu Gly Gln Met Arg Arg Leu Pro
      35             40             45
Ala Ser Ser Cys Gln Lys Asp Arg Asn Asp Phe Ala Phe Pro Gln Asp
      50             55             60
Val Phe Gly Gly Asp Gln Ser His Lys Ala Gln Ala Leu Ser Val Val
      65             70             75             80
His Val Thr Asn Gln Lys Ile Phe His Phe Phe Cys Thr Glu Ala Ser
      85             90             95
Ser Ser Ala Ala Trp Asn Thr Thr Leu Leu Glu Glu Phe Cys Thr Gly
      100            105            110
Leu Asp Trp Gln Leu Thr Arg Leu Glu Ala Cys Val Met Gln Glu Val
      115            120            125
Gly Glu Gly Glu Ala Pro Leu Thr Asn Glu Asp Ser Ile Leu Arg Asn
      130            135            140
Tyr Phe Gln Arg Leu Ser Leu Tyr Leu Gln Glu Lys Lys Tyr Ser Pro
      145            150            155            160
Cys Ala Trp Glu Ile Val Arg Ala Glu Ile Met Arg Ser Leu Tyr Tyr
      165            170            175
Ser Ser Thr Ala Leu Gln Lys Arg Leu Arg Ser Glu Lys
      180            185

```

<210> SEQ ID NO 109
 <211> LENGTH: 567
 <212> TYPE: DNA
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 109

```

tttctcgctc cttaatcttt tctgcaaggc tgttgatgaa taatacaagg atctcatgat      60
ttctgctctg acgatctccc aggcacaagg gctgtatttc ttctcttgca ggtagaggga      120
gagtcttttg aagtagttcc tcaggatgga gtcctcgttc gtgaggggag cctctccttc      180
ccccacctcc tgcgatgacac aggcctccag gcgggtcagc tgccaatcaa gtcccgtgca      240
gaattcctcc aggagggtgg tgttccaagc agcagacgag gacgcctctg tgcagaagaa      300
gtggaagatc ttctggttcg tcacgtgcac caccgagagg gcttgggcct tgtgggactg      360
gtctccacca aacacgtcct gggggaaggc gaagtcattt ctgtccttct gacaggagct      420
ggcagggagt ctctcattt gtcccaggag cgtcaaggcc ctctgttca gcaggccgtg      480
ggtctgaggc aggtcacagc ccagagagca gacggagttg cagcccagcg ccaccagggc      540
caccaagaag gaagagggca gcgccat                                     567

```

<210> SEQ ID NO 110
 <211> LENGTH: 567
 <212> TYPE: DNA
 <213> ORGANISM: Felis catus
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(567)

-continued

<400> SEQUENCE: 110

```

atg gcg ctg ccc tct tcc ttc ttg gtg gcc ctg gtg gcg ctg ggc tgc      48
Met Ala Leu Pro Ser Ser Phe Leu Val Ala Leu Val Ala Leu Gly Cys
  1             5             10             15

aac tcc gtc tgc tct ctg ggc tgt gac ctg cct cag acc cac ggc ctg      96
Asn Ser Val Cys Ser Leu Gly Cys Asp Leu Pro Gln Thr His Gly Leu
          20             25             30

ctg aac agg agg gcc ttg acg ctc ctg gga caa atg agg aga ctc cct     144
Leu Asn Arg Arg Ala Leu Thr Leu Leu Gly Gln Met Arg Arg Leu Pro
          35             40             45

gcc agc tcc tgt cag aag gac agg aat gac ttc gcc ttc ccc cag gac     192
Ala Ser Ser Cys Gln Lys Asp Arg Asn Asp Phe Ala Phe Pro Gln Asp
          50             55             60

gtg ttc ggt gga gac cag tcc cac aag gct caa gcc ctc tcg gtg gtg     240
Val Phe Gly Gly Asp Gln Ser His Lys Ala Gln Ala Leu Ser Val Val
          65             70             75             80

cac gtg acg aac cag gag atc ttc cac ttc ttc tgc aca gag gcg tcc     288
His Val Thr Asn Gln Glu Ile Phe His Phe Phe Cys Thr Glu Ala Ser
          85             90             95

tcg tct gct gct tgg aac acc acc ctc ctg gag gaa ttc tgc acg gga     336
Ser Ser Ala Ala Trp Asn Thr Thr Leu Leu Glu Glu Phe Cys Thr Gly
          100             105             110

ctt gat cgg cag ctg acc cgc ctg gaa gcc tgt gtc gtg cag gag gtg     384
Leu Asp Arg Gln Leu Thr Arg Leu Glu Ala Cys Val Val Gln Glu Val
          115             120             125

ggg gag gga gag gct ccc ctc acg aac gag gac tcc ctc ctg agg aac     432
Gly Glu Gly Glu Ala Pro Leu Thr Asn Glu Asp Ser Leu Leu Arg Asn
          130             135             140

tac ttc caa aga ctc tcc ctc tac ctg caa gag aag aaa tac agc cct     480
Tyr Phe Gln Arg Leu Ser Leu Tyr Leu Gln Glu Lys Lys Tyr Ser Pro
          145             150             155             160

tgt gcc tgg gag atc gtc aga gca gaa atc atg aga tcc ttg tat tat     528
Cys Ala Trp Glu Ile Val Arg Ala Glu Ile Met Arg Ser Leu Tyr Tyr
          165             170             175

tca tca aca gcc ttg caa aaa aga tta agg agc gag aaa                567
Ser Ser Thr Ala Leu Gln Lys Arg Leu Arg Ser Glu Lys
          180             185

```

<210> SEQ ID NO 111

<211> LENGTH: 189

<212> TYPE: PRT

<213> ORGANISM: Felis catus

<400> SEQUENCE: 111

```

Met Ala Leu Pro Ser Ser Phe Leu Val Ala Leu Val Ala Leu Gly Cys
  1             5             10             15

Asn Ser Val Cys Ser Leu Gly Cys Asp Leu Pro Gln Thr His Gly Leu
          20             25             30

Leu Asn Arg Arg Ala Leu Thr Leu Leu Gly Gln Met Arg Arg Leu Pro
          35             40             45

Ala Ser Ser Cys Gln Lys Asp Arg Asn Asp Phe Ala Phe Pro Gln Asp
          50             55             60

Val Phe Gly Gly Asp Gln Ser His Lys Ala Gln Ala Leu Ser Val Val
          65             70             75             80

His Val Thr Asn Gln Glu Ile Phe His Phe Phe Cys Thr Glu Ala Ser
          85             90             95

Ser Ser Ala Ala Trp Asn Thr Thr Leu Leu Glu Glu Phe Cys Thr Gly
          100             105             110

```

-continued

Leu Asp Arg Gln Leu Thr Arg Leu Glu Ala Cys Val Val Gln Glu Val
 115 120 125
 Gly Glu Gly Glu Ala Pro Leu Thr Asn Glu Asp Ser Leu Leu Arg Asn
 130 135 140
 Tyr Phe Gln Arg Leu Ser Leu Tyr Leu Gln Glu Lys Lys Tyr Ser Pro
 145 150 155 160
 Cys Ala Trp Glu Ile Val Arg Ala Glu Ile Met Arg Ser Leu Tyr Tyr
 165 170 175
 Ser Ser Thr Ala Leu Gln Lys Arg Leu Arg Ser Glu Lys
 180 185

<210> SEQ ID NO 112
 <211> LENGTH: 567
 <212> TYPE: DNA
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 112

```

ttctcgcctc cttaatcttt ttgcaaggc tgttgatgaa taatacaagg atctcatgat      60
ttctgctctg acgatctccc aggcacaagg gctgtatttc ttctcttgca ggtagaggga    120
gagtccttgg aagtagttcc tcaggaggga gtcctcgttc gtgaggggag cctctccctc    180
ccccacctcc tgcacgacac aggcctccag gcgggtcagc tgccgatcaa gtcccgtgca    240
gaattcctcc aggagggtgg tgttccaagc agcagacgag gacgcctctg tgcagaagaa    300
gtggaagatc tcctggttcg tcacgtgcac caccgagagg gcttgagcct tgtgggactg    360
gtctccaccg aacacgtcct gggggaaggc gaagtcattc ctgtccttct gacaggagct    420
ggcagggagt ctctcattt gtcccaggag cgtcaaggcc ctctgttca gcaggccgtg    480
ggtctgaggc aggtcacagc ccagagagca gacggagttg cagcccagcg ccaccagggc    540
caccaagaag gaagagggca gcgccat                                     567

```

<210> SEQ ID NO 113
 <211> LENGTH: 498
 <212> TYPE: DNA
 <213> ORGANISM: Felis catus
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(498)

<400> SEQUENCE: 113

```

tgt gac ctg cct cag acc cac ggc ctg ctg aac agg agg gcc ttg acg      48
Cys Asp Leu Pro Gln Thr His Gly Leu Leu Asn Arg Arg Ala Leu Thr
  1 5 10 15
ctc ctg gga caa atg agg aga ctc cct gcc agc tcc tgt cag aag gac      96
Leu Leu Gly Gln Met Arg Arg Leu Pro Ala Ser Ser Cys Gln Lys Asp
  20 25 30
aga aat gac ttc gcc ttc ccc cag gac gtg ttt ggt gga gac cag tcc    144
Arg Asn Asp Phe Ala Phe Pro Gln Asp Val Phe Gly Gly Asp Gln Ser
  35 40 45
cac aag gcc caa gcc ctc tcg gtg gtg cac gtg acg aac cag aag atc    192
His Lys Ala Gln Ala Leu Ser Val Val His Val Thr Asn Gln Lys Ile
  50 55 60
ttc cac ttc ttc tgc aca gag gcg tcc tcg tct gct gct tgg aac acc    240
Phe His Phe Phe Cys Thr Glu Ala Ser Ser Ser Ala Ala Trp Asn Thr
  65 70 75 80
acc ctc ctg gag gaa ttc tgc acg gga ctt gat tgg cag ctg acc cgc    288
Thr Leu Leu Glu Glu Phe Cys Thr Gly Leu Asp Trp Gln Leu Thr Arg
  85 90 95

```

-continued

ctg gaa gcc tgt gtc atg cag gag gtg ggg gag gga gag gct ccc ctc	336
Leu Glu Ala Cys Val Met Gln Glu Val Gly Glu Gly Glu Ala Pro Leu	
100 105 110	
acg aac gag gac tcc atc ctg agg aac tac ttc caa aga ctc tcc ctc	384
Thr Asn Glu Asp Ser Ile Leu Arg Asn Tyr Phe Gln Arg Leu Ser Leu	
115 120 125	
tac ctg caa gag aag aaa tac agc cct tgt gcc tgg gag atc gtc aga	432
Tyr Leu Gln Glu Lys Lys Tyr Ser Pro Cys Ala Trp Glu Ile Val Arg	
130 135 140	
gca gaa atc atg aga tcc ttg tat tat tca tca aca gcc ttg cag aaa	480
Ala Glu Ile Met Arg Ser Leu Tyr Tyr Ser Ser Thr Ala Leu Gln Lys	
145 150 155 160	
aga tta agg agc gag aaa	498
Arg Leu Arg Ser Glu Lys	
165	

<210> SEQ ID NO 114
 <211> LENGTH: 166
 <212> TYPE: PRT
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 114

Cys Asp Leu Pro Gln Thr His Gly Leu Leu Asn Arg Arg Ala Leu Thr	
1 5 10 15	
Leu Leu Gly Gln Met Arg Arg Leu Pro Ala Ser Ser Cys Gln Lys Asp	
20 25 30	
Arg Asn Asp Phe Ala Phe Pro Gln Asp Val Phe Gly Gly Asp Gln Ser	
35 40 45	
His Lys Ala Gln Ala Leu Ser Val Val His Val Thr Asn Gln Lys Ile	
50 55 60	
Phe His Phe Phe Cys Thr Glu Ala Ser Ser Ser Ala Ala Trp Asn Thr	
65 70 75 80	
Thr Leu Leu Glu Glu Phe Cys Thr Gly Leu Asp Trp Gln Leu Thr Arg	
85 90 95	
Leu Glu Ala Cys Val Met Gln Glu Val Gly Glu Gly Glu Ala Pro Leu	
100 105 110	
Thr Asn Glu Asp Ser Ile Leu Arg Asn Tyr Phe Gln Arg Leu Ser Leu	
115 120 125	
Tyr Leu Gln Glu Lys Lys Tyr Ser Pro Cys Ala Trp Glu Ile Val Arg	
130 135 140	
Ala Glu Ile Met Arg Ser Leu Tyr Tyr Ser Ser Thr Ala Leu Gln Lys	
145 150 155 160	
Arg Leu Arg Ser Glu Lys	
165	

<210> SEQ ID NO 115
 <211> LENGTH: 498
 <212> TYPE: DNA
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 115

tttctcgctc cttaatcttt tctgcaaggc tgttgatgaa taatacaagg atctcatgat	60
ttctgctctg acgatctccc aggcacaagg gctgtatttc ttctcttgca ggtagaggga	120
gagtctttgg aagtagttcc tcaggatgga gtcctcgttc gtgaggggag cctctccctc	180
ccccacctcc tgcattgacac aggcttcag gcgggtcagc tgccaatcaa gtcccgtgca	240
gaattctctc aggaggggtg tgttccaagc agcagacgag gacgcctctg tgcagaagaa	300

-continued

```

gtggaagatc ttctggttcg tcacgtgcac caccgagagg gcttgggcct tgtgggactg 360
gtctccacca aacacgtcct gggggaaggc gaagtcattt ctgtccttct gacaggagct 420
ggcagggagt ctctcattt gtcccaggag cgtcaaggcc ctctgttca gcaggccgtg 480
ggtctgaggc aggtcaca 498

```

```

<210> SEQ ID NO 116
<211> LENGTH: 498
<212> TYPE: DNA
<213> ORGANISM: Felis catus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(498)

```

```

<400> SEQUENCE: 116

```

```

tgt gac ctg cct cag acc cac ggc ctg ctg aac agg agg gcc ttg acg 48
Cys Asp Leu Pro Gln Thr His Gly Leu Leu Asn Arg Arg Ala Leu Thr
  1             5             10             15

ctc ctg gga caa atg agg aga ctc cct gcc agc tcc tgt cag aag gac 96
Leu Leu Gly Gln Met Arg Arg Leu Pro Ala Ser Ser Cys Gln Lys Asp
             20             25             30

agg aat gac ttc gcc ttc ccc cag gac gtg ttc ggt gga gac cag tcc 144
Arg Asn Asp Phe Ala Phe Pro Gln Asp Val Phe Gly Gly Asp Gln Ser
             35             40             45

cac aag gct caa gcc ctc tcg gtg gtg cac gtg acg aac cag gag atc 192
His Lys Ala Gln Ala Leu Ser Val Val His Val Thr Asn Gln Glu Ile
             50             55             60

ttc cac ttc ttc tgc aca gag gcg tcc tcg tct gct gct tgg aac acc 240
Phe His Phe Phe Cys Thr Glu Ala Ser Ser Ser Ala Ala Trp Asn Thr
             65             70             75             80

acc ctc ctg gag gaa ttc tgc acg gga ctt gat cgg cag ctg acc cgc 288
Thr Leu Leu Glu Glu Phe Cys Thr Gly Leu Asp Arg Gln Leu Thr Arg
             85             90             95

ctg gaa gcc tgt gtc gtg cag gag gtg ggg gag gga gag gct ccc ctc 336
Leu Glu Ala Cys Val Val Gln Glu Val Gly Glu Gly Glu Ala Pro Leu
             100            105            110

acg aac gag gac tcc ctc ctg agg aac tac ttc caa aga ctc tcc ctc 384
Thr Asn Glu Asp Ser Leu Leu Arg Asn Tyr Phe Gln Arg Leu Ser Leu
             115            120            125

tac ctg caa gag aag aaa tac agc cct tgt gcc tgg gag atc gtc aga 432
Tyr Leu Gln Glu Lys Lys Tyr Ser Pro Cys Ala Trp Glu Ile Val Arg
             130            135            140

gca gaa atc atg aga tcc ttg tat tat tca tca aca gcc ttg caa aaa 480
Ala Glu Ile Met Arg Ser Leu Tyr Tyr Ser Ser Thr Ala Leu Gln Lys
             145            150            155            160

aga tta agg agc gag aaa 498
Arg Leu Arg Ser Glu Lys
             165

```

```

<210> SEQ ID NO 117
<211> LENGTH: 166
<212> TYPE: PRT
<213> ORGANISM: Felis catus

```

```

<400> SEQUENCE: 117

```

```

Cys Asp Leu Pro Gln Thr His Gly Leu Leu Asn Arg Arg Ala Leu Thr
  1             5             10             15

Leu Leu Gly Gln Met Arg Arg Leu Pro Ala Ser Ser Cys Gln Lys Asp
             20             25             30

Arg Asn Asp Phe Ala Phe Pro Gln Asp Val Phe Gly Gly Asp Gln Ser
             35             40             45

```


-continued

His Lys Ala Gln Ala Leu Ser Val Val His Val Thr Asn Gln Glu Ile
50 55 60

Phe His Phe Phe Cys Thr Glu Ala Ser Ser Ser Ala Ala Trp Asn Thr
65 70 75 80

Thr Leu Leu Glu Glu Phe Cys Thr Gly Leu Asp Arg Gln Leu Thr Arg
85 90 95

Leu Glu Ala Cys Val Val Gln Glu Val Gly Glu Gly Glu Ala Pro Leu
100 105 110

Thr Asn Glu Asp Ser Leu Leu Arg Asn Tyr Phe Gln Arg Leu Ser Leu
115 120 125

Tyr Leu Gln Glu Lys Lys Tyr Ser Pro Cys Ala Trp Glu Ile Val Arg
130 135 140

Ala Glu Ile Met Arg Ser Leu Tyr Tyr Ser Ser Thr Ala Leu Gln Lys
145 150 155 160

Arg Leu Arg Ser Glu Lys
165

<210> SEQ ID NO 118
<211> LENGTH: 498
<212> TYPE: DNA
<213> ORGANISM: Felis catus

<400> SEQUENCE: 118

```

tttctcgctc cttaatcttt ttgcaaggc tgttgatgaa taatacaagg atctcatgat      60
ttctgctctg acgatctccc aggcacaagg gctgtatttc ttctcttgca ggtagaggga    120
gagtcctttg aagtagttcc tcaggaggga gtcctcgttc gtgaggggag cctctccctc    180
ccccacctcc tgcacgacac aggcctccag gcgggtcagc tgccgatcaa gtcccgtgca    240
gaattcctcc aggagggtgg tgttccaagc agcagacgag gacgcctctg tgcagaagaa    300
gtggaagatc tcctggttcg tcacgtgcac caccgagagg gcttgagcct tgtgggactg    360
gtctccaccg aacacgtcct gggggaaggc gaagtcattc ctgtccttct gacaggagct    420
ggcagggagt ctctcattt gtcccaggag cgtcaaggcc ctctgttca gcaggccgtg    480
ggtctgaggc aggtcaca                                     498

```

<210> SEQ ID NO 119
<211> LENGTH: 444
<212> TYPE: DNA
<213> ORGANISM: Felis catus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (10)..(441)

<400> SEQUENCE: 119

```

ggatccacc atg tgg ctg cag aac ctg ctt ttc ctg ggc act gtg gtc tgc      51
      Met Trp Leu Gln Asn Leu Leu Phe Leu Gly Thr Val Val Cys
          1          5          10

agc atc tct gca ccc acc agt tca ccc agc tct gtc act cgg ccc tgg      99
Ser Ile Ser Ala Pro Thr Ser Ser Pro Ser Ser Val Thr Arg Pro Trp
15          20          25          30

caa cac gtg gat gcc atc aag gag gcc ctg agc ctt ctg aac aac agt      147
Gln His Val Asp Ala Ile Lys Glu Ala Leu Ser Leu Leu Asn Asn Ser
35          40          45

agt gaa ata act gct gtg atg aat gaa gca gta gaa gtc gtc tct gaa      195
Ser Glu Ile Thr Ala Val Met Asn Glu Ala Val Glu Val Val Ser Glu
50          55          60

atg ttt gac cct gag gag ccg aaa tgc ctg cag act cac cta aag ctg      243

```

[illegible]

<400> SEQUENCE: 120

Met	Trp	Leu	Gln	Asn	Leu	Leu	Phe	Leu	Gly	Thr	Val	Val	Cys	Ser	Ile
1				5					10					15	
Ser	Ala	Pro	Thr	Ser	Ser	Pro	Ser	Ser	Val	Thr	Arg	Pro	Trp	Gln	His
			20					25					30		
Val	Asp	Ala	Ile	Lys	Glu	Ala	Leu	Ser	Leu	Leu	Asn	Asn	Ser	Ser	Glu
		35					40					45			
Ile	Thr	Ala	Val	Met	Asn	Glu	Ala	Val	Glu	Val	Val	Ser	Glu	Met	Phe
	50					55					60				
Asp	Pro	Glu	Glu	Pro	Lys	Cys	Leu	Gln	Thr	His	Leu	Lys	Leu	Tyr	Glu
65					70					75					80
Gln	Gly	Leu	Arg	Gly	Ser	Leu	Ile	Ser	Leu	Lys	Glu	Pro	Leu	Arg	Met
				85					90					95	
Met	Ala	Asn	His	Tyr	Lys	Gln	His	Cys	Pro	Leu	Thr	Pro	Glu	Thr	Pro
			100					105					110		
Cys	Glu	Thr	Gln	Thr	Ile	Thr	Phe	Lys	Asn	Phe	Lys	Glu	Asn	Leu	Lys
		115					120					125			
Asp	Phe	Leu	Phe	Asn	Asn	Pro	Phe	Asp	Cys	Trp	Gly	Pro	Asp	Gln	Lys
	130					135					140				

<400> SEQUENCE: 121

ttactttctgg tctggtccccc agcagtc meta ggggttgcta aacagaaaat ccttcagatt	60
ctctttgaaa tttttgaagg tgatagtcctg ggtttcacag ggcgtttccg gagtaagggg	120
gcagtgctgc ttgtaattgt tggccatcat tctcagaggc tccttgaggc tgatgaggct	180
gccccgtagg ccctgctcgt acagctttag gtgagtcctgc aggcatttcg gtccttcagg	240
gtcaaacatt tcagagacga cttctactgc ttcattcatc acagcagtta tttcactact	300
gttgttcaga aggctcaggg cctccttgat ggcattccacg tgttgccagg gccgagtgac	360
aqagctgggt gaactggttg gtgcagagat gctgcagacc acagtgccta gaaaaagcaq	420

-continued

```

gtttctgcagc cacatggtgg atcc 444

<210> SEQ ID NO 122
<211> LENGTH: 432
<212> TYPE: DNA
<213> ORGANISM: Felis catus

<400> SEQUENCE: 122
atgttgctgc agaacctgct tttcctgggc actgtggtct gcagcatctc tgcacccacc 60
agttcaccca gctctgtcac tcggccctgg caacacgtgg atgccatcaa ggaggccctg 120
agccttctga acaacagtag tgaaataact gctgtgatga atgaagcagt agaagtcgtc 180
tctgaaatgt ttgacctga ggagccgaaa tgcctgcaga ctcacctaaa gctgtacgag 240
cagggcctac ggggcagcct catcagcctc aaggagcctc tgagaatgat ggccaaccat 300
tacaagcagc actgccccct tactccggaa acgccctgtg aaaccagac tatcaccttc 360
aaaaatttca aagagaatct gaaggatttt ctgtttaaca acccctttga ctgctgggga 420
ccagaccaga ag 432

<210> SEQ ID NO 123
<211> LENGTH: 432
<212> TYPE: DNA
<213> ORGANISM: Felis catus

<400> SEQUENCE: 123
cttctggtct ggtccccagc agtcaaaggg gttgttaaac agaaaatcct tcagattctc 60
tttgaaattt ttgaaggatga tagtctgggt ttcacagggc gtttccggag taagggggca 120
gtgctgcttg taatggttg ccattcattct cagaggctcc ttgaggctga tgaggctgcc 180
ccgtaggccc tgctctgaca gctttagggt agtctgcagg catttcggct cctcagggtc 240
aaacatttca gagacgactt ctactgttc attcatcaca gcagttattt cactactgtt 300
gttcagaagg ctccaggcct ccttgatggc atccacgtgt tgccagggcc gagtgacaga 360
gtggtgtgaa ctggtgggtg cagagatgct gcagaccaca gtgccagga aaagcagggt 420
ctgcagccac at 432

<210> SEQ ID NO 124
<211> LENGTH: 381
<212> TYPE: DNA
<213> ORGANISM: Felis catus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(381)

<400> SEQUENCE: 124
gca ccc acc agt tca ccc agc tct gtc act cgg ccc tgg caa cac gtg 48
Ala Pro Thr Ser Ser Pro Ser Ser Val Thr Arg Pro Trp Gln His Val
1 5 10 15
gat gcc atc aag gag gcc ctg agc ctt ctg aac aac agt agt gaa ata 96
Asp Ala Ile Lys Glu Ala Leu Ser Leu Leu Asn Asn Ser Ser Glu Ile
20 25 30
act gct gtg atg aat gaa gca gta gaa gtc gtc tct gaa atg ttt gac 144
Thr Ala Val Met Asn Glu Ala Val Glu Val Val Ser Glu Met Phe Asp
35 40 45
cct gag gag ccg aaa tgc ctg cag act cac cta aag ctg tac gag cag 192
Pro Glu Glu Pro Lys Cys Leu Gln Thr His Leu Lys Leu Tyr Glu Gln
50 55 60
ggc cta cgg gcc agc ctc atc agc ctc aag gag cct ctg aga atg atg 240
Gly Leu Arg Gly Ser Leu Ile Ser Leu Lys Glu Pro Leu Arg Met Met

```

-continued

65	70	75	80	
gcc aac cat tac aag	cag cac tgc ccc	ctt act ccg gaa acg ccc	tgt	288
Ala Asn His Tyr Lys	Gln His Cys Pro	Leu Thr Pro Glu Thr	Pro Cys	
	85	90	95	
gaa acc cag act atc acc ttc aaa aat ttc aaa gag aat ctg aag gat				336
Glu Thr Gln Thr Ile Thr Phe Lys	Asn Phe Lys Glu Asn Leu Lys	Asp		
	100	105	110	
ttt ctg ttt aac aac ccc ttt gac tgc tgg gga cca gac cag aag				381
Phe Leu Phe Asn Asn Pro Phe Asp Cys Trp Gly	Pro Asp Gln Lys			
	115	120	125	

<210> SEQ ID NO 125
 <211> LENGTH: 127
 <212> TYPE: PRT
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 125

Ala Pro Thr Ser Ser Pro Ser Ser Val Thr Arg Pro Trp Gln His Val	
1 5 10 15	
Asp Ala Ile Lys Glu Ala Leu Ser Leu Leu Asn Asn Ser Ser Glu Ile	
20 25 30	
Thr Ala Val Met Asn Glu Ala Val Glu Val Val Ser Glu Met Phe Asp	
35 40 45	
Pro Glu Glu Pro Lys Cys Leu Gln Thr His Leu Lys Leu Tyr Glu Gln	
50 55 60	
Gly Leu Arg Gly Ser Leu Ile Ser Leu Lys Glu Pro Leu Arg Met Met	
65 70 75 80	
Ala Asn His Tyr Lys Gln His Cys Pro Leu Thr Pro Glu Thr Pro Cys	
85 90 95	
Glu Thr Gln Thr Ile Thr Phe Lys Asn Phe Lys Glu Asn Leu Lys Asp	
100 105 110	
Phe Leu Phe Asn Asn Pro Phe Asp Cys Trp Gly Pro Asp Gln Lys	
115 120 125	

<210> SEQ ID NO 126
 <211> LENGTH: 381
 <212> TYPE: DNA
 <213> ORGANISM: Felis catus

<400> SEQUENCE: 126

cttctggtct ggtccccagc agtcaaagg gttgttaaac agaaaatcct tcagattctc	60
tttgaaatth ttgaagggtga tagtctgggt ttcacagggc gtttccggag taagggggca	120
gtgctgcttg taatggttg ccatcattct cagaggctcc ttgaggctga tgaggctgcc	180
ccgtaggccc tgctcgtaca gctttaggtg agtctgcagg catttcggct cctcagggtc	240
aaacatttca gagacgactt ctactgcttc attcatcaca gcagttatth cactactggt	300
gttcagaagg ctcaggccct ccttgatggc atccacgtgt tgccagggcc gagtgacaga	360
gctgggtgaa ctggtgggtg c	381

<210> SEQ ID NO 127
 <211> LENGTH: 28
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Primer

<400> SEQUENCE: 127

-continued

cctcgagatt cagctttcaa tgctgtga 28

<210> SEQ ID NO 128
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 128

tgcccrstcg gcttcttctc c 21

<210> SEQ ID NO 129
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 129

cgactctctt tccrctctc ctg 23

<210> SEQ ID NO 130
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 130

cctcaaattg cggcacatgt c 21

<210> SEQ ID NO 131
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 131

ctgttcagag tttagtaag cc 22

<210> SEQ ID NO 132
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 132

gaagatacca tttaacttt aacacagc 28

<210> SEQ ID NO 133
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 133

tgctgtattg tgaagactcc cagc 24

-continued

<210> SEQ ID NO 134
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 134
atgcactttc ttgccc 16

<210> SEQ ID NO 135
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 135
ctggaggaaa akacttcrat gattctgata tctgaaatat at 42

<210> SEQ ID NO 136
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 136
ctgacycttk sttgscctc attctca 27

<210> SEQ ID NO 137
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 137
gggctcgaga aaagatttgc ttagaaaaat cccatg 36

<210> SEQ ID NO 138
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 138
cccgcggcgc ctcaactttc cgggtgccac tc 32

<210> SEQ ID NO 139
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 139
gtcmtggctc tyrcttgccct tgg 23

-continued

<210> SEQ ID NO 140
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 140

aaastgggcy acytcgattt tgg 23

<210> SEQ ID NO 141
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 141

gtgatgttgm ycagctcctc 20

<210> SEQ ID NO 142
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 142

aattaaccct cactaaaggg 20

<210> SEQ ID NO 143
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 143

atggcgctct ggttgactgt 20

<210> SEQ ID NO 144
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 144

ggcttttgag agcacagtgc 20

<210> SEQ ID NO 145
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 145

ccccatatga gccctgtgac tccctcccc 29

-continued

<210> SEQ ID NO 146
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 146
ggggaattct catctgaaat ttccatggcg 30

<210> SEQ ID NO 147
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 147
atggcgctgc cctcttcctt ctg 24

<210> SEQ ID NO 148
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 148
tcatttctcg ctccttaac ttttctgc 28

<210> SEQ ID NO 149
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 149
cagggatcca ccatgtggct gcagaacctg cttttcc 37

<210> SEQ ID NO 150
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 150
ttacttctgg tctggtcccc agcagtc aaa ggggttgta aacagaaaat 50

<210> SEQ ID NO 151
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Primer

<400> SEQUENCE: 151
cacagyccca tctcctcc 18

<210> SEQ ID NO 152

-continued

```

<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Primer

```

```

<400> SEQUENCE: 152

```

```

gtaatacgac tcactatagg gc

```

22

```

<210> SEQ ID NO 153
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Primer

```

```

<400> SEQUENCE: 153

```

```

acggaattcg agatgatagt gctggc

```

26

```

<210> SEQ ID NO 154
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Primer

```

```

<400> SEQUENCE: 154

```

```

gtgtctagat ttggtagaaa aggatgat

```

28

What is claimed is:

1. An isolated protein selected from the group consisting of:

(a) an isolated protein of at least about 20 amino acids in length, wherein said 20 amino acids are encoded by a nucleic acid molecule that has at least 60 contiguous nucleotide region identical in sequence to a 60 contiguous nucleotide region of a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:4, and SEQ ID NO:19; and

(b) an isolated protein of at least about 20 amino acids in length, wherein said protein has an at least 20 contiguous amino acid region identical in sequence to a 20 contiguous amino acid region selected from the group consisting of SEQ ID NO:2 and SEQ ID NO:20,

wherein said isolated protein of (a) or (b) elicits an immune response against a canine IL-4 protein or has IL-4 activity.

2. The isolated protein of claim 1, wherein said protein has an amino acid sequence selected from the group consisting of SEQ ID NO:2 and SEQ ID NO:20.

3. The isolated protein of claim 2, wherein the protein has the amino acid sequence of SEQ ID NO:2.

35 4. The isolated protein of claim 2, wherein the protein has the amino acid sequence of SEQ ID NO:20.

5. An isolated protein having an amino acid sequence that is at least about 85 percent identical to an amino acid sequence selected from the group consisting of SEQ ID NO:2 and SEQ ID NO:20, wherein said isolated protein elicits and immune response against a canine IL-4 protein or has IL-4 activity.

6. A therapeutic composition comprising the isolated protein of claim 1.

7. The composition of claim 6, wherein said composition further comprises a component selected from the group consisting of an excipient, an adjuvant and a carrier.

8. A method to regulate an immune response in an animal comprising administering to the animal the therapeutic composition of claim 6.

9. The method of claim 8, wherein said animal is a canid.

10. The method of claim 8, wherein said composition further comprises a component selected from the group consisting of an excipient, an adjuvant and a carrier.

* * * * *