

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2017271665 B2**

(54) Title
Treatment of primary ciliary dyskinesia with synthetic messenger RNA

(51) International Patent Classification(s)
A61K 48/00 (2006.01) **A61K 31/713** (2006.01)
A61K 31/711 (2006.01)

(21) Application No: **2017271665** (22) Date of Filing: **2017.05.26**

(87) WIPO No: **WO17/205767**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/342,784	2016.05.27	US

(43) Publication Date: **2017.11.30**

(44) Accepted Journal Date: **2023.11.23**

(71) Applicant(s)
Transcriptx, Inc.

(72) Inventor(s)
Lockhart, David J.;Wustman, Brandon;Hennig, Mirko;Ishimaru, Daniella

(74) Agent / Attorney
Pizzeys Patent and Trade Mark Attorneys Pty Ltd, PO Box 291, WODEN, ACT, 2606, AU

(56) Related Art
BRIGITTE CHHIN ET AL: "Ciliary Beating Recovery in Deficient Human Airway Epithelial Cells after Lentivirus Ex Vivo Gene Therapy", PLOS GENETICS, vol. 5, no. 3, 20 March 2009 (2009-03-20), pages e1000422, DOI: 10.1371/journal.pgen.1000422



(51) International Patent Classification:

A61K 48/00 (2006.01) A61K 31/713 (2006.01)
A61K 31/711 (2006.01)

(21) International Application Number:

PCT/US2017/034723

(22) International Filing Date:

26 May 2017 (26.05.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/342,784 27 May 2016 (27.05.2016) US

(71) Applicant: TRANSCRIPTX, INC. [US/US]; 675 Al-
manor Avenue, Sunnyvale, CA 94085 (US).

(72) Inventors: LOCKHART, David, J.; 1005 Lakeview Way,
Emerald Hills, CA 94062 (US). WUSTMAN, Brandon;
2020 Fort Stockton Drive, San Diego, CA 92103 (US).
HENNIG, Mirko; 865 Carlisle Way #122, Sunnyvale,
CA 94087 (US). ISHIMARU, Daniella; 865 Carlisle Way
#122, Sunnyvale, CA 94087 (US).

(74) Agent: COLE, Adam, J. et al.; Wilson Sonsini Goodrich
& Rosati, 650 Page Mill Road, Palo Alto, CA 94304-1050
(US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR,
KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

(54) Title: TREATMENT OF PRIMARY CILIARY DYSKINESIA WITH SYNTHETIC MESSENGER RNA




FIGURE 3

(57) Abstract: Polynucleotides encoding peptides, proteins, enzymes, and functional fragments thereof are disclosed. The polynucleotides of the disclosure can be effectively delivered to an organ, such as the lung, and expressed within cells of the organ. The polyribonucleotides of the disclosure can be used to treat a disease or condition associated with cilia maintenance and function, impaired function of the axoneme, such as DNAI1 or DNAH5.



Published:

— *with international search report (Art. 21(3))*

TREATMENT OF PRIMARY CILIARY DYSKINESIA WITH SYNTHETIC MESSENGER RNA

CROSS-REFERENCE

[0001] This application is a continuation of U.S. Provisional Patent Application Serial No. 62/342,784, filed on May 27, 2016, which is incorporated herein by reference in its entirety and to which application we claim priority under 35 U.S.C. § 120.

BACKGROUND

[0002] Messenger RNAs (mRNA) are polymers containing a number of linked nucleotides, each composed of a sugar, a phosphate, and a base. Each mRNA polymer stores genetic information along the nucleotide chain. Messenger RNA polymers carry the genetic information from the DNA in the nucleus of the cell to the cytoplasm where proteins are made. Each triplet of nucleotides in the mRNA is called a codon, and each codon specifies the identity of an amino acid in the translated protein.

[0003] A cell can also take up and translate an exogenous RNA, but many factors influence efficient uptake and translation. For instance, the immune system recognizes many exogenous RNAs as foreign and triggers a response that is aimed at inactivating the RNAs.

SUMMARY

[0004] The present disclosure provides polyribonucleotides, and compositions comprising the same, that can encode a protein of choice. In some cases, the disclosure provides a method for treating a subject having or at risk of having primary ciliary dyskinesia, the method comprising administering to the subject a composition that comprises a nucleic acid construct that encodes dynein axonemal intermediate chain 1 protein or a variant thereof, which nucleic acid construct includes codons that provide for heterologous or enhanced expression of the dynein axonemal intermediate chain 1 protein or a variant thereof within cells of the subject, thereby treating the subject having or at risk of having primary ciliary dyskinesia. The nucleic acid construct can be, for example a complementary deoxyribonucleic acid DNA template. The nucleic acid construct may encode dynein axonemal intermediate chain 1 protein or a variant thereof at a level that is increased by a factor of at least about 1.5, at least about 5, or another suitable amount as compared to levels within cells exposed to a composition comprising a nucleic acid construct that does not include the codons encoding dynein axonemal intermediate chain 1 protein or a variant thereof. In some instances, the codons of the construct are at least 70% homologous to a mammalian, such as a human, dynein axonemal intermediate chain 1 mRNA.

[0005] In some cases, the construct comprises a 5' and/or 3' untranslated region (UTR) flanking the codon sequence which encodes the dynein axonemal intermediate chain 1, wherein the untranslated region(s) enhance(s) the expression of the protein within cells of the subject. The 3' noncoding region may comprise a 3'-cap independent translation enhancer (3'-CITEs). In some instances, the 3' noncoding region may also comprise at least one intermediate sequence region between the codon sequence and either the 3' noncoding region or the 5' noncoding region or a 3'-stem loop region derived from the nucleotide sequence of a histone protein. In some cases, the codon sequence comprises an open reading frame (ORF). The 3' noncoding region flanking the codon sequence (e.g., ORF) may comprise a poly adenosine tail, wherein the number of adenosines in the poly adenosine tail improves the translation efficiency and increases the half-life of the dynein axonemal intermediate chain 1 mRNA. In some cases, the length of the poly adenosine tail is at most 200 adenosines. The poly adenosine tail may comprise a percentage of chemically modified nucleotides. In some instances, fewer than 20% of the nucleotides in the poly adenosine tail are chemically modified. In some instances, fewer than 30% of the nucleotides encoding dynein axonemal intermediate chain 1 in a construct are chemically modified nucleotides. In the instances where the nucleotides comprise a chemically modified nucleotide, the chemically modified nucleotide can be selected from the group consisting of pseudouridine, 1-methylpseudouridine, 2-thiouridine, 5-iodouridine, 5-methyluridine, 5-methylcytidine, and 5-iodocytidine. In some cases, the chemically modified nucleotide is 1-methylpseudouridine. In some cases, the modified nucleotide is pseudouridine. In other cases, the modified nucleotides are a combination of 1-methylpseudouridine and pseudouridine. In addition to the composition comprising a polyribonucleotide for treating a subject having or at risk of having primary ciliary dyskinesia, in some cases, the present disclosure further provides a composition comprising at least one additional nucleic acid construct that encodes a protein selected from the group consisting of: armadillo repeat containing 4 (ARMC4), chromosome 21 open reading frame 59 (C21orf59), coiled-coil domain containing 103 (CCDC103), coiled-coil domain containing 114 (CCDC114), coiled-coil domain containing 39 (CCDC39), coiled-coil domain containing 40 (CCDC40), coiled-coil domain containing 65 (CCDC65), cyclin O (CCNO), dynein (axonemal) assembly factor 1 (DNAAF1), dynein (axonemal) assembly factor 2 (DNAAF2), dynein (axonemal) assembly factor 3 (DNAAF3), dynein (axonemal) assembly factor 5 (DNAAF5), dynein axonemal heavy chain 11 (DNAH11), dynein axonemal heavy chain 5 (DNAH5), dynein axonemal heavy chain 6 (DNAH6), dynein axonemal heavy chain 8 (DNAH8), dynein axonemal intermediate chain 2 (DNAI2), dynein axonemal light chain 1 (DNAL1), dynein regulatory complex subunit 1 (DRC1), dyslexia susceptibility 1 candidate 1

(DYX1C1), growth arrest specific 8 (GAS8), axonemal central pair apparatus protein (HYDIN), leucine rich repeat containing 6 (LRR6), NME/NM23 family member 8 (NME8), oral-facial-digital syndrome 1 (OFD1), retinitis pigmentosa GTPase regulator (RPGR), radial spoke head 1 homolog (Chlamydomonas) (RSPH1), radial spoke head 4 homolog A (Chlamydomonas) (RSPH4A), radial spoke head 9 homolog (Chlamydomonas) (RSPH9), sperm associated antigen 1 (SPAG1), and zinc finger MYND-type containing 10 (ZMYND10).

[0006] The disclosure provides a composition comprising a nucleic acid construct encoding dynein axonemal intermediate chain 1, which nucleic acid construct includes codons that provide for heterologous or enhanced expression of the dynein axonemal intermediate chain 1 protein or a variant thereof within cells of a subject having or at risk of having primary ciliary dyskinesia. The compositions described herein may comprise a ratio of moles of amine groups of cationic polymers to moles of phosphate groups of the modified polyribonucleotide of at least about 4. In some cases, the composition is formulated in a nanoparticle or nanocapsule. In other cases, the composition is formulated in a cationic lipid, cationic polymer, or nanoemulsion. The composition may be formulated for administration to a subject. The nucleic acid constructs in the composition may include codons that provide for heterologous or enhanced expression of the dynein axonemal intermediate chain 1 protein or a variant thereof within cells of a subject having or at risk of having primary ciliary dyskinesia. In some cases, fewer than 30% of the ribonucleotides encoding dynein axonemal intermediate chain 1 are chemically modified nucleotides. In some instances, the codons of the construct are at least 70% homologous to a mammalian, such as a human, dynein axonemal intermediate chain 1 mRNA. In some cases, the construct comprises a 5' or 3' noncoding region flanking the codon sequence which encodes the dynein axonemal intermediate chain 1, wherein the noncoding region enhances the expression of the protein within cells the subject. In other cases, the construct comprises a 3' noncoding region flanking the codon sequence which encodes the dynein axonemal intermediate chain 1, wherein the 3' noncoding region comprises a 3'-cap independent translation enhancer (3'-CITEs). The 3' noncoding region may comprise a 3'-stem loop region derived from the nucleotide sequence of a histone protein. The 3' noncoding region may comprise a 3'-triple helical structure derived from the nucleotide sequence of metastasis-associated lung adenocarcinoma transcript 1 (MALAT1). The 3' noncoding region flanking the codon sequence may comprise a poly adenosine tail, wherein the number of adenosines in the poly adenosine tail improves the translation efficiency of the dynein axonemal intermediate chain 1 protein. In some cases, the number of adenosines in the poly adenosine tail improves the half-life of the dynein axonemal intermediate chain 1 protein. In some cases, the length of the poly adenosine tail is at most 200 adenosines. In some

instances, a percentage of the poly adenosine tail comprises modified nucleotides. In some instances, fewer than 20% of the adenosines in the poly(A)tail are modified. In some cases, the construct comprises a percentage of chemically modified nucleotides. In some instances, fewer than 30% of the nucleotides encoding dynein axonemal intermediate chain 1 are chemically modified. When chemically modified nucleotides are present, they may be selected from the group consisting of pseudouridine, 1-methylpseudouridine, 5-methoxyuridine, 2-thiouridine, 5-iodouridine, 5-methyluridine, 5-methylcytidine, 2'-amino-2'-deoxycytidine, 2'-fluoro-2'-deoxycytidine, and 5-iodocytidine. In some cases, the chemically modified nucleotide is pseudouridine or 1-methyl pseudouridine. In some instances, the composition further comprises at least one additional nucleic acid construct. The at least one additional nucleic acid construct encodes a protein selected from the group consisting of: armadillo repeat containing 4 (ARMC4), chromosome 21 open reading frame 59 (C21orf59), coiled-coil domain containing 103 (CCDC103), coiled-coil domain containing 114 (CCDC114), coiled-coil domain containing 39 (CCDC39), coiled-coil domain containing 40 (CCDC40), coiled-coil domain containing 65 (CCDC65), cyclin O (CCNO), dynein (axonemal) assembly factor 1 (DNAAF1), dynein (axonemal) assembly factor 2 (DNAAF2), dynein (axonemal) assembly factor 3 (DNAAF3), dynein (axonemal) assembly factor 5 (DNAAF5), dynein axonemal heavy chain 11 (DNAH11), dynein axonemal heavy chain 5 (DNAH5), dynein axonemal heavy chain 6 (DNAH6), dynein axonemal heavy chain 8 (DNAH8), dynein axonemal intermediate chain 2 (DNAI2), dynein axonemal light chain 1 (DNAL1), dynein regulatory complex subunit 1 (DRC1), dyslexia susceptibility 1 candidate 1 (DYX1C1), growth arrest specific 8 (GAS8), axonemal central pair apparatus protein (HYDIN), leucine rich repeat containing 6 (LRRC6), NME/NM23 family member 8 (NME8), oral-facial-digital syndrome 1 (OFD1), retinitis pigmentosa GTPase regulator (RPGR), radial spoke head 1 homolog (Chlamydomonas) (RSPH1), radial spoke head 4 homolog A (Chlamydomonas) (RSPH4A), radial spoke head 9 homolog (Chlamydomonas) (RSPH9), sperm associated antigen 1 (SPAG1), and zinc finger MYND-type containing 10 (ZMYND10).

[0007] The present disclosure also provides a nucleic acid construct, a vector, or an isolated nucleic acid that is/are formulated for administration to a subject. In some cases, the formulation includes a therapeutically effective amount of the nucleic acid construct encoding dynein axonemal intermediate chain 1. The nucleic acid construct can be a cDNA construct that encodes dynein axonemal intermediate chain 1 protein or a variant thereof, or any one of the aforementioned additional nucleic acid constructs. In some cases, the present disclosure provides a composition comprising a nucleic acid construct encoding dynein axonemal

intermediate chain 1, wherein the nucleic acid construct comprises any one of SEQ ID NOs 14-16. In some cases, the present disclosure provides a composition comprising a nucleic acid construct encoding dynein axonemal heavy chain 5, wherein the nucleic acid construct comprises any one of SEQ ID NOs 17-18.

[0008] Additional aspects and advantages of the present disclosure will become readily apparent to those skilled in this art from the following detailed description, wherein only illustrative embodiments of the present disclosure are shown and described. As will be realized, the present disclosure is capable of other and different embodiments, and its several details are capable of modifications in various obvious respects, all without departing from the disclosure. Accordingly, the drawings and description are to be regarded as illustrative in nature, and not as restrictive.

INCORPORATION BY REFERENCE

[0009] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.

BRIEF DESCRIPTION OF THE DRAWINGS

[0010] The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings of which:

[0011] FIGURE 1 is an agarose gel illustrating the production of capped and uncapped DNAI1 RNA.

[0012] FIGURE 2 is a western blot illustrating the translations of DNAI1 mRNA in HEK-293 cells at 6 hours, 24 hours, and 48 hours post-transfection.

[0013] FIGURE 3 illustrates fragment analyzer data of a posttranscriptionally poly-adenylated RNA transcript encoding dynein axonemal intermediate chain 1 (DNAI1).

[0014] FIGURE 4 illustrates fragment analyzer data of a posttranscriptionally poly-adenylated RNA transcript encoding dynein axonemal intermediate chain 1 (DNAI1).

[0015] FIGURE 5 illustrates PAGE data of the size of poly adenylated tail of the plasmid encoding dynein axonemal intermediate chain 1 (DNAI1).

[0016] FIGURE 6 illustrates the fragment analyzer data of an *in vitro* transcribed DNAI1 mRNA comprising the unmodified nucleotides.

[0017] **FIGURE 7** illustrates fragment analyzer data of an *in vitro* transcribed DNAI1 mRNA comprising 50% pseudouridine (Ψ).

[0018] **FIGURE 8** illustrates fragment analyzer data of an *in vitro* transcribed DNAI1 mRNA comprising 100% pseudouridine (Ψ).

[0019] **FIGURE 9** illustrates fragment analyzer data of an *in vitro* transcribed DNAI1 mRNA comprising 100% 1-methylpseudouridine that was post-transcriptionally poly adenylated.

[0020] **FIGURE 10** illustrates double-stranded RNA content as detected by dot-blot.

[0021] **FIGURE 11** illustrates the HPLC-based nucleotide composition analysis of an *in vitro* transcribed nucleic acid construct that encodes dynein axonemal intermediate chain 1, transcribed with unmodified nucleotides.

[0022] **FIGURE 12** illustrates the HPLC-based nucleotide composition analysis of an *in vitro* transcribed nucleic acid construct that encodes dynein axonemal intermediate chain 1, transcribed with 50% Ψ .

[0023] **FIGURE 13** illustrates the HPLC-based nucleotide composition analysis of an *in vitro* transcribed nucleic acid construct that encodes dynein axonemal intermediate chain 1, transcribed with 100% Ψ .

[0024] **FIGURE 14** is a graph illustrating the relative expression levels of DNAI1 protein in HEK-293, A549, and MLE-15 cells.

[0025] **FIGURE 15** illustrates the induction of IL-6 in A549 cells transfected with the DNAI1 mRNA variants.

[0026] **FIGURE 16** illustrates the induction of IL-6 in A549 cells transfected with the DNAI1 mRNA variants.

[0027] **FIGURE 17** is a graph illustrating the relative expression of DNAI1 protein in HEK-293, A549, and MLE-15 cells.

[0028] **FIGURE 18** illustrates induction of IL-6 in A549 cells by DNAI1 transcripts.

[0029] **FIGURE 19** illustrates induction of IL-6 in A549 cells by DNAI1 transcripts.

[0030] **FIGURE 20** illustrates cell viability of A549 cells after transfection with various amounts of each DNAI1 mRNA measured using the CellTiter-Glo assay.

[0031] **FIGURE 21** illustrates cell viability of A549 cells after transfection with various amounts of each DNAI1 mRNA measured using the CellTiter-Glo assay.

[0032] **FIGURE 22** illustrates induction of IP-10 in HepG2 cells by DNAI1 transcripts.

[0033] **FIGURE 23** illustrates cell viability of HepG2 cells after transfection with various amounts of each DNAI1 mRNA measured using the CellTiter-Glo assay.

[0034] **FIGURE 24** illustrates the peak expression of dynein axonemal intermediate chain 1 (DNAI1) protein or other controls in HEK-293 cells.

[0035] **FIGURE 25** expression of DNAI1 in fully differentiated human airway epithelial cells.

[0036] **FIGURE 26** illustrates an overall quality improvement in DNAI1 expressing a polyribonucleotide of SEQ ID NO 15 (**B**) as compared to a polyribonucleotide of SEQ ID NO 14 (**A**).

[0037] **FIGURE 27** illustrates an overall improvement in translation efficiency in A549 cells of a polyribonucleotide of SEQ ID NO 15 (**B**) as compared to a polyribonucleotide of SEQ ID NO 14 (**A**).

[0038] **FIGURE 28** illustrates an analysis of double-stranded RNA content of a polyribonucleotide of SEQ ID NO 15 as compared to known concentrations of known concentrations of poly-IC.

[0039] **FIGURE 29** illustrates HPLC-purification of unmodified and 100% m¹Ψ-containing DNAI1 mRNA.

[0040] **FIGURE 30** illustrates example translation activity and immunogenicity for fractions enriched in full-length, unmodified mRNA transcripts in A549 cells using HPLC-purification.

DETAILED DESCRIPTION

[0041] While various embodiments of the invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions may occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed.

[0042] The term “subject,” as used herein generally refers to a human. In some instances, a subject can also be an animal, such as a mouse, a rat, a guinea pig, a dog, a cat, a horse, a rabbit, and various other animals. A subject can be of any age, for example, a subject can be an infant, a toddler, a child, a pre-adolescent, an adolescent, an adult, or an elderly individual.

[0043] The term “disease,” as used herein, generally refers to an abnormal physiological condition that affects part or all of a subject, such as an illness (e.g., primary ciliary dyskinesia) or another abnormality that causes defects in the action of cilia in, for example, the lining the respiratory tract (lower and upper, sinuses, Eustachian tube, middle ear), in a variety of lung cells, in the fallopian tube, or flagella of sperm cells.

[0044] The term “polynucleotide” or “nucleic acid” as used herein refers to a polymeric form of nucleotides of any length, either ribonucleotides or deoxyribonucleotides, that comprise purine and pyrimidine bases, purine and pyrimidine analogues, chemically or biochemically modified,

natural or non-natural, or derivatized nucleotide bases. Polynucleotides include sequences of deoxyribonucleic acid (DNA), ribonucleic acid (RNA), or DNA copies of ribonucleic acid (cDNA), all of which can be recombinantly produced, artificially synthesized, or isolated and purified from natural sources. The polynucleotides and nucleic acids may exist as single-stranded or double-stranded. The backbone of the polynucleotide can comprise sugars and phosphate groups, as may typically be found in RNA or DNA, or analogues or substituted sugar or phosphate groups. A polynucleotide may comprise naturally occurring or non-naturally occurring nucleotides, such as methylated nucleotides and nucleotide analogues (or analogs).

[0045] The term “polyribonucleotide,” as used herein, generally refers to polynucleotide polymers that comprise ribonucleic acids. The term also refers to polynucleotide polymers that comprise chemically modified ribonucleotides. A polyribonucleotide can be formed of D-ribose sugars, which can be found in nature.

[0046] The term “polypeptides,” as used herein, generally refers to polymer chains comprised of amino acid residue monomers which are joined together through amide bonds (peptide bonds). A polypeptide can be a chain of at least three amino acids, a protein, a recombinant protein, an antigen, an epitope, an enzyme, a receptor, or a structure analogue or combinations thereof. As used herein, the abbreviations for the L-enantiomeric amino acids that form a polypeptide are as follows: alanine (A, Ala); arginine (R, Arg); asparagine (N, Asn); aspartic acid (D, Asp); cysteine (C, Cys); glutamic acid (E, Glu); glutamine (Q, Gln); glycine (G, Gly); histidine (H, His); isoleucine (I, Ile); leucine (L, Leu); lysine (K, Lys); methionine (M, Met); phenylalanine (F, Phe); proline (P, Pro); serine (S, Ser); threonine (T, Thr); tryptophan (W, Trp); tyrosine (Y, Tyr); valine (V, Val). X or Xaa can indicate any amino acid.

[0047] The term “engineered,” as used herein, generally refers to polynucleotides, vectors, and nucleic acid constructs that have been genetically designed and manipulated to provide a polynucleotide intracellularly. An engineered polynucleotide can be partially or fully synthesized *in vitro*. An engineered polynucleotide can also be cloned. An engineered polyribonucleotide can contain one or more base or sugar analogues, such as ribonucleotides not naturally-found in messenger RNAs. An engineered polyribonucleotide can contain nucleotide analogues that exist in transfer RNAs (tRNAs), ribosomal RNAs (rRNAs), guide RNAs (gRNAs), small nuclear RNA (snRNA), small nucleolar RNA (snoRNA), SmY RNA, spliced leader RNA (SL RNA), CRISPR RNA, long noncoding RNA (lncRNA), microRNA (miRNA), or another suitable RNA.

Overview

[0048] The present disclosure provides compositions and methods for the treatment of conditions associated with cilia maintenance and function, with nucleic acids encoding a protein or protein fragment(s). Numerous eukaryotic cells carry appendages, which are often referred to as cilia or flagella, whose inner core comprises a cytoskeletal structure called the axoneme. The axoneme can function as the skeleton of cellular cytoskeletal structures, both giving support to the structure and, in some instances, causing it to bend. Usually, the internal structure of the axoneme is common to both cilia and flagella. Cilia are often found in the linings of the airway, the reproductive system, and other organs and tissues. Flagella are tail-like structures that, similarly to cilia, can propel cells forward, such as sperm cells.

[0049] Without properly functioning cilia in the airway, bacteria can remain in the respiratory tract and cause infection. In the respiratory tract, cilia move back and forth in a coordinated way to move mucus towards the throat. This movement of mucus helps to eliminate fluid, bacteria, and particles from the lungs. Many infants afflicted with cilia and flagella malfunction experience breathing problems at birth, which suggests that cilia play an important role in clearing fetal fluid from the lungs. Beginning in early childhood, subjects afflicted with cilia malfunction can develop frequent respiratory tract infections.

[0050] Primary ciliary dyskinesia is a condition characterized by chronic respiratory tract infections, abnormally positioned internal organs, and the inability to have children (infertility). The signs and symptoms of this condition are caused by abnormal cilia and flagella. Subjects afflicted with primary ciliary dyskinesia often have year-round nasal congestion and a chronic cough. Chronic respiratory tract infections can result in a condition called bronchiectasis, which damages the passages, called bronchi, leading from the windpipe to the lungs and can cause life-threatening breathing problems.

[0051] In some instances, a nucleic acid construct, vector, or composition of the disclosure comprises one or more nucleotide sequences that encode dynein axonemal intermediate chain 1 protein or a variant thereof, and the sequences provide for heterologous or enhanced expression of the dynein axonemal intermediate chain 1 protein or a variant thereof within cells of a subject. In some instances, the nucleic acid construct, vector, or composition also comprises the genetic code of 5' untranslated regions (UTRs) and 3' UTRs of SEQ ID NOs 1-9, as shown below.

TABLE 1	
UTR	DNA sequence (from 5' to 3')
α-globin 5' UTR (HBA1)	GGGAGACATAAACCCCTGGCGCGCTCGCGGCCCGGCACTCTTCTGGTCCCC ACAGACTCAGAGAGAAGCCACC (SEQ ID NO: 1)
α-globin 5' UTR (HBA2)	GGGAGACATAAACCCCTGGCGCGCTCGCGGGCCCGGCACTCTTCTGGTCCCC ACAGACTCAGAGAGAAGCCACC (SEQ ID NO: 2)
α-globin 5' UTR	GGGAGACTCTTCTGGTCCCCACAGACTCAGAGAGAACGCCACC (SEQ ID NO: 3)
IRES of EMCV 5'-UTR	GTTATTTTCCACCATATTGCCGTCTTTTGGCAATGTGAGGGCCCGGAAACC TGGCCCTGTCTTCTTGACGAGCATTCTAGGGGTCTTTCCCTCTCGCCAA AGGAATGCAAGGTCTGTTGAATGTCGTGAAGGAAGCAGTTCCTCTGGAAG CTTCTTGAAGACAAACAACGTCTGTAGCGACCCTTTGCAGGCAGCGGAAC CCCCACCTGGCGACAGGTGCCTCTGCGGCCAAAAGCCACGTGTATAAGA TACACCTGCAAAGGCGGCACAACCCAGTGCCACGTTGTGAGTTGGATAG TTGTGGAAAGAGTCAAATGGCTCTCCTCAAGCGTATTCAACAAGGGGCTG AAGGATGCCCAGAAGGTACCCCATTTGTATGGGATCTGATCTGGGGCCTCG GTGCACATGCTTTACGTGTGTTTAGTCGAGGTTAAAAAACGTCTAGGCC CCCGAACCACGGGGACGTGGTTTTTCTTTGAAAAACACGATGATAATATG GCCACAACC (SEQ ID NO: 4)
IRES of TEV 5'-UTR	AAATAACAAATCTCAACACAACATATACAAAACAAACGAATCTCAAGCA ATCAAGCATTCTACTTCTATTGCAGCAATTTAAATCATTCTTTTAAAGCA AAAGCAATTTTCTGAAAATTTTACCATTACGAACGATAGCA (SEQ ID NO: 5)
ssRNA1 5'UTR	GGGAGACAAGAGAGAAAAGAAGAGCAAGAAGAAATATAAGAGCCACC (SEQ ID NO: 6)
ssRNA2 5'UTR	GGGAGACCCAAGCTGGCTAGCGTTTAAACTTAAGCTTGGCAATCCGGTAC TGTTGGTAAAGCCACC (SEQ ID NO: 7)
ssRNA 3 + native 5' UTR	GGGAGACCCAAGCTGGCTAGCGTTTAAACTTAAGCTTTCCTTTCCGGGCC GGCTGGGCGCGCCGAAGCGCCTGCGCCTTGGCTGCTGGTCCGGTTGCTGGG TAACCGCGTCAGGGAGTTGGATTCTATCCTGCAAGGGCACGGGGACCCAC AACGACGGCTGTCCCTAAAGAACCGTTGCGACTGGTAACTGAAGTGGAA GAGAGTCCAGATTTCTTGTGTGGTCAAGGAGACGGACAAACTTTTTGT CTTCAGACGAGGGAGCGTTTTGTAGGCTCTCCAGGGGTTGAG (SEQ ID NO: 8)
TMV 3'-UTR	GGATTGTGTCCGTAATCACACGTGGTGCCTACGATAACGCATAGTGTTTT TCCCTCCACTTAAATCGAAGGGTGTGTCTTGGATCGCGCGGGTCAAATG TATATGGTTCATATACATCCGCAGGCACGTAATAAAGCGAGGGGTTTCGAA TCCCCCGTTACCCCGGTAGGGGCCATTGTCTTC (SEQ ID NO: 9)
MALAT1 3'-UTR	TCAGTAGGGTCATGAAGGTTTTTCTTTTCTGAGAAAACAACACGTATTGT TTTCTCAGGTTTTGCTTTTTGGCCTTTTTCTAGCTTAAAAAAAAAAAAAAGC AAAATTGTCTTC (SEQ ID NO: 10)

NEAT2 3'-UTR	TCAGTAGGGTTGTAAAGGTTTTTCTTTTCCTGAGAAAACAACCTTTTGTTT TCTCAGGTTTTGCTTTTTGGCCTTTCCCTAGCTTTAAAAAAAAAAAAAAGCAA AATTGTCTTC (SEQ ID NO: 11)
histone cluster 2, H3c 3'-UTR	GAAGTGGCGGTTTCGGCCGGAGGTTCCATCGTATCCAAAAGGCTCTTTTCA GAGCCACCCATTGTCTTC (SEQ ID NO: 12)
Native 3' UTR	GGGGCTGGCCTCAGTCTCTGTCCCATCGCTTGAATACAGTACTCCTAGGG CTTGACCCTGGTACCCAGCCCAGCCTTAGCACCCAGCATGTGACCCCACT CCTGATCAGGTCCCAGCATCTTCCCTTCTTGTTCTGTTCCCTTAAGGTCCCA GCACCTTACCCAGGACTTGGTCTTCAACCACCATTACCCCTCTAACTTTG CACAAATAAACCTGTGTAGAAACCCACCCCAAAAAAA (SEQ ID NO: 13)

Primary Ciliary Dyskinesia, Related Conditions and Treatments Thereof

[0052] The methods, constructs, and compositions of this disclosure provide a method to treat primary ciliary dyskinesia (PCD), also known as immotile ciliary syndrome or Kartagener syndrome. PCD is typically considered to be a rare, ciliopathic, autosomal recessive genetic disorder that often causes defects in the action of cilia lining the respiratory tract (lower and upper, sinuses, Eustachian tube, middle ear) and fallopian tube, as well as in the flagella of sperm cells.

[0053] Some individuals with primary ciliary dyskinesia have abnormally placed organs within their chest and abdomen. These abnormalities arise early in embryonic development when the differences between the left and right sides of the body are established. About 50 percent of people with primary ciliary dyskinesia have a mirror-image reversal of their internal organs (*situs inversus totalis*). For example, in these individuals the heart is on the right side of the body instead of on the left. When someone afflicted with primary ciliary dyskinesia has *situs inversus totalis*, they are often said to have Kartagener syndrome.

[0054] Approximately 12 percent of people with primary ciliary dyskinesia have a condition known as heterotaxy syndrome or *situs ambiguus*, which is characterized by abnormalities of the heart, liver, intestines, or spleen. These organs may be structurally abnormal or improperly positioned. In addition, affected individuals may lack a spleen (asplenia) or have multiple spleens (polysplenia). Heterotaxy syndrome results from problems establishing the left and right sides of the body during embryonic development. The severity of heterotaxy varies widely among affected individuals.

[0055] Primary ciliary dyskinesia can also lead to infertility. Vigorous movements of the flagella are can be needed to propel the sperm cells forward to the female egg cell. Because the sperm of subjects afflicted with primary ciliary dyskinesia does not move properly, males with primary

ciliary dyskinesia are usually unable to father children. Infertility occurs in some affected females and it is usually associated with abnormal cilia in the fallopian tubes.

[0056] Another feature of primary ciliary dyskinesia is recurrent ear infections (otitis media), especially in young children. Otitis media can lead to permanent hearing loss if untreated. The ear infections are likely related to abnormal cilia within the inner ear.

[0057] Rarely, individuals with primary ciliary dyskinesia have an accumulation of fluid in the brain (hydrocephalus), likely due to abnormal cilia in the brain.

[0058] The polyribonucleotides of the disclosure can be used, for example, to treat a subject having or at risk of having primary ciliary dyskinesia or any other condition associated with a defect or malfunction of a gene whose function is linked to cilia maintenance and function. Non limiting examples of genes that have been associated with primary ciliary dyskinesia include: armadillo repeat containing 4 (ARMC4), chromosome 21 open reading frame 59 (C21orf59), coiled-coil domain containing 103 (CCDC103), coiled-coil domain containing 114 (CCDC114), coiled-coil domain containing 39 (CCDC39), coiled-coil domain containing 40 (CCDC40), coiled-coil domain containing 65 (CCDC65), cyclin O (CCNO), dynein (axonemal) assembly factor 1 (DNAAF1), dynein (axonemal) assembly factor 2 (DNAAF2), dynein (axonemal) assembly factor 3 (DNAAF3), dynein (axonemal) assembly factor 5 (DNAAF5), dynein axonemal heavy chain 11 (DNAH11), dynein axonemal heavy chain 5 (DNAH5), dynein axonemal heavy chain 6 (DNAH6), dynein axonemal heavy chain 8 (DNAH8), dynein axonemal intermediate chain 2 (DNAI2), dynein axonemal light chain 1 (DNAL1), dynein regulatory complex subunit 1 (DRC1), dyslexia susceptibility 1 candidate 1 (DYX1C1), growth arrest specific 8 (GAS8), axonemal central pair apparatus protein (HYDIN), leucine rich repeat containing 6 (LRRC6), NME/NM23 family member 8 (NME8), oral-facial-digital syndrome 1 (OFD1), retinitis pigmentosa GTPase regulator (RPGR), radial spoke head 1 homolog (Chlamydomonas) (RSPH1), radial spoke head 4 homolog A (Chlamydomonas) (RSPH4A), radial spoke head 9 homolog (Chlamydomonas) (RSPH9), sperm associated antigen 1 (SPAG1), and zinc finger MYND-type containing 10 (ZMYND10).

[0059] In some cases, the composition comprises a nucleic acid construct encoding dynein axonemal intermediate chain 1 (DNAI1), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having primary ciliary dyskinesia. The DNAI1 gene can provide instructions for making a protein that is part of a group (complex) of proteins called dynein. This complex functions within the cilia. Coordinated back and forth movement of cilia can move the cell or the fluid surrounding the cell and dynein produces the force needed for cilia to move. Within the core of cilia (the axoneme), dynein

complexes are part of structures known as inner dynein arms (IDAs) and outer dynein arms (ODAs) depending on their location. Coordinated movement of the dynein arms causes the entire axoneme to bend back and forth. IDAs and ODAs have different combinations of protein components (subunits) that are classified by weight as heavy, intermediate, or light chains. The DNAI1 gene provides instructions for making intermediate chain 1, which is found in ODAs. Other subunits can be produced from different genes administered to the subject in the same or in a separate composition. Alternatively, other subunits can be produced by a single nucleic acid construct that encodes a functional component of an inner dynein arm or an outer dynein arm.

[0060] At least 21 mutations in the DNAI1 gene have been found to cause primary ciliary dyskinesia, which is a condition characterized by respiratory tract infections, abnormal organ placement, and an inability to have children (infertility). DNAI1 gene mutations result in an absent or abnormal intermediate chain 1. Without a normal version of this subunit, the ODAs cannot form properly and may be shortened or absent. As a result, cilia cannot produce the force needed to bend back and forth. Defective cilia lead to the features of primary ciliary dyskinesia. In some cases, the disclosure provides a nucleic acid that is engineered to replace or to supplement the function of the endogenous DNAI1 protein comprising the IVS1+2_3insT (219+3insT) mutation. In some cases, the disclosure provides a nucleic acid that is engineered to replace or to supplement the function of the endogenous DNAI1 protein comprising the A538T mutation, the second most common.

[0061] In some cases, the composition comprises a nucleic acid construct encoding dynein axonemal intermediate chain 2 (DNAI2), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having primary ciliary dyskinesia. The DNAI2 gene is part of the dynein complex of respiratory cilia and sperm flagella. Mutations in this gene are associated with primary ciliary dyskinesia type 9, a disorder characterized by abnormalities of motile cilia, respiratory infections leading to chronic inflammation and bronchiectasis, and abnormalities in sperm tails.

[0062] In some cases, the composition comprises a nucleic acid construct encoding armadillo repeat containing 4 (ARMC4), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the ARMC4 gene comprises ten Armadillo repeat motifs (ARMs) and one HEAT repeat, and has been shown to localize to the ciliary axonemes and at the ciliary base of respiratory cells. Mutations in the ARMC4 gene can cause partial outer dynein arm (ODA) defects in respiratory cilia.

[0063] In some cases, the composition comprises a nucleic acid construct encoding chromosome 21 open reading frame 59 (C21orf59), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the C21orf59 gene can play a critical role in dynein arm assembly and motile cilia function. Mutations in this gene can result in primary ciliary dyskinesia.

[0064] In some cases, the composition comprises a nucleic acid construct encoding coiled-coil domain containing 103 (CCDC103), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the CCDC103 gene can function as a dynein-attachment factor required for cilia motility.

[0065] In some cases, the composition comprises a nucleic acid construct encoding coiled-coil domain containing 114 (CCDC114), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the CCDC114 gene can function as a component of the outer dynein arm docking complex in cilia cells. Mutations in this gene can cause primary ciliary dyskinesia 20.

[0066] In some cases, the composition comprises a nucleic acid construct encoding coiled-coil domain containing 39 (CCDC39), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the CCDC39 gene can function as the assembly of dynein regulatory and inner dynein arm complexes, which regulate ciliary beat. Defects in this gene are a cause of primary ciliary dyskinesia type 14 (CCDC39).

[0067] In some cases, the composition comprises a nucleic acid construct encoding coiled-coil domain containing 40 (CCDC40), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the CCDC40 gene can function together with CCDC39 to form a molecular ruler that determines the 96 nanometer (nm) repeat length and arrangements of components in cilia and flagella (by similarity). CCDC40 may not be required for outer dynein arm complexes assembly, but it may be required for axonemal recruitment of CCDC39. In some cases, CCD40 and CCD39 can be produced from different genes administered to the subject in the same or in a separate composition. Alternatively, CCD40 and CCD39 can be produced by a single nucleic acid construct that encodes a functional component of an inner dynein arm or an

outer dynein arm. Defects in the CCD40 gene are a cause of primary ciliary dyskinesia type 14 (CILD14).

[0068] In some cases, the composition comprises a nucleic acid construct encoding coiled-coil domain containing 65 (CCDC65), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the CCDC65 gene can function as a sperm cell protein. CCDC65 has been shown to be highly expressed in adult testis, spermatocytes and spermatids. The protein plays a critical role in the assembly of the nexin-dynein regulatory complex. Mutations in this gene have been associated with primary ciliary dyskinesia type 27.

[0069] In some cases, the composition comprises a nucleic acid construct encoding cyclin O (CCNO), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia.

[0070] In some cases, the composition comprises a nucleic acid construct encoding dynein (axonemal) assembly factor 1 (DNAAF1), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the DNAAF1 gene is thought to be cilium-specific and it can be required for the stability of the ciliary architecture. Mutations in this gene have been associated with primary ciliary dyskinesia type 13.

[0071] In some cases, the composition comprises a nucleic acid construct encoding dynein (axonemal) assembly factor 2 (DNAAF2), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the DNAAF2 gene can be involved in the preassembly of dynein arm complexes which power cilia. Mutations in this gene have been associated with primary ciliary dyskinesia type 10 (CILD10).

[0072] In some cases, the composition comprises a nucleic acid construct encoding dynein (axonemal) assembly factor 3 (DNAAF3), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the DNAAF3 gene can be required for the assembly of axonemal inner and outer dynein arms and it can play a role in assembling dynein complexes for transport into cilia. Mutations in this gene have been associated with primary ciliary dyskinesia type 2 (CILD2).

[0073] In some cases, the composition comprises a nucleic acid construct encoding dynein (axonemal) assembly factor 5 (DNAAF5), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary

dyskinesia. The protein encoded by the DNAAF5 gene is thought to be required for the preassembly or stability of axonemal dynein arms, and is found only in organisms with motile cilia and flagella. Mutations in this gene have been associated with primary ciliary dyskinesia-18.

[0074] In some cases, the composition comprises a nucleic acid construct encoding dynein axonemal heavy chain 11 (DNAH11), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the DNAH11 gene can produce a ciliary outer dynein arm protein. DNAH11 is thought to be a microtubule-dependent motor ATPase involved in the movement of respiratory cilia. Mutations in this gene have been associated with primary ciliary dyskinesia type 7 (CILD7) and heterotaxy syndrome.

[0075] In some cases, the composition comprises a nucleic acid construct encoding dynein axonemal heavy chain 5 (DNAH5), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having primary ciliary dyskinesia. The DNAH5 gene can provide instructions for making a protein that is part of a group (complex) of proteins called dynein. Coordinated back and forth movement of cilia can move the cell or the fluid surrounding the cell. Dynein can produce the force needed for cilia to move. More than 80 mutations of the DNAH5 have been associated with primary ciliary dyskinesia. Mutations in this gene have been associated with primary ciliary dyskinesia and heterotaxy syndrome.

[0076] In some cases, the composition comprises a nucleic acid construct encoding dynein axonemal heavy chain 6 (DNAH6), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having primary ciliary dyskinesia.

[0077] In some cases, the composition comprises a nucleic acid construct encoding dynein axonemal heavy chain 8 (DNAH8), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the DNAH8 gene can function as a force generating protein of respiratory cilia. DNAH8 can produce force towards the minus ends of microtubules. Dynein has ATPase activity; the force-producing power stroke is thought to occur on release of ADP. DNAH8 can be involved in sperm motility and in sperm flagellar assembly. DNAH8 is also known as ATPase and hdhc9.

[0078] In some cases, the composition comprises a nucleic acid construct encoding dynein axonemal light chain 1 (DNAL1), and upon translation within the cells of a subject the construct

yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the DNAL1 gene can function as a force generating protein of respiratory cilia. DNAL1 can function as a component of the outer dynein arms complex. This complex acts as the molecular motor that provides the force to move cilia in an ATP-dependent manner. Mutations in this gene have been associated with primary ciliary dyskinesia type 16 (CILD16).

[0079] In some cases, the composition comprises a nucleic acid construct encoding dynein regulatory complex subunit 1 (DRC1), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the DRC1 gene can function as a force generating protein of respiratory cilia. DRC1 can encode a central component of the nexin-dynein complex (N-DRC), which regulates the assembly of ciliary dynein. Mutations in this gene have been associated with primary ciliary dyskinesia type 21 (CILD21).

[0080] In some cases, the composition comprises a nucleic acid construct encoding dyslexia susceptibility 1 candidate 1 (DYX1C1), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the DYX1C1 gene can function as a force generating protein of respiratory cilia. DYX1C1 can encode a tetratricopeptide repeat domain-containing protein. The encoded protein can interact with estrogen receptors and the heat shock proteins, Hsp70 and Hsp90. Mutations in this gene are also associated with deficits in reading and spelling, and a chromosomal translocation involving this gene is associated with a susceptibility to developmental dyslexia.

[0081] In some cases, the composition comprises a nucleic acid construct encoding growth arrest specific 8 (GAS8), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia.

[0082] In some cases, the composition comprises a nucleic acid construct encoding axonemal central pair apparatus protein (HYDIN), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the HYDIN gene can function in cilia motility. Mutations in this gene have been associated with primary ciliary dyskinesia type 5 (CILD5).

[0083] In some cases, the composition comprises a nucleic acid construct encoding leucine rich repeat containing 6 (LRRC6), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the LRRC6 gene contains several leucine-rich repeat domains and

appears to be involved in the motility of cilia. Mutations in this gene have been associated with primary ciliary dyskinesia type 19 (CILD19).

[0084] In some cases, the composition comprises a nucleic acid construct encoding NME/NM23 family member 8 (NME8), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the NME8 gene can function as a force generating protein of respiratory cilia. The NME8 protein comprises an N-terminal thioredoxin domain and three C-terminal nucleoside diphosphate kinase (NDK) domains. Mutations in this gene have been associated with primary ciliary dyskinesia type 6 (CILD6).

[0085] In some cases, the composition comprises a nucleic acid construct encoding oral-facial-digital syndrome 1 (OFD1), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The function of the protein produced by the OFD1 gene is not well understood, but it may play a role play a critical role in the early development of many parts of the body, including the brain, face, limbs, and kidneys. About 100 mutations in the OFD1 gene have been found in people with oral-facial-digital syndrome type I, which is the most common form of the disorder. Mutations in this gene have been associated with primary ciliary dyskinesia and Joubert syndrome.

[0086] In some cases, the composition comprises a nucleic acid construct encoding retinitis pigmentosa GTPase regulator (RPGR), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the RPGR gene can be important for normal vision and for the function of the cilia. Mutations in this gene have been associated with primary ciliary dyskinesia, X-linked retinitis pigmentosa, progressive vision loss, chronic respiratory and sinus infections, recurrent ear infections (otitis media), and hearing loss.

[0087] In some cases, the composition comprises a nucleic acid construct encoding radial spoke head 1 homolog (RSPH1), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the RSPH1 gene may play an important role in male meiosis and in the building of the axonemal central pair and radial spokes. Mutations in this gene have been associated with primary ciliary dyskinesia type 24 (CILD24).

[0088] In some cases, the composition comprises a nucleic acid construct encoding radial spoke head 4 homolog A (RSPH4A), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia.

The protein encoded by the RSPH4A gene may be a component the radial spoke head. Mutations in this gene have been associated with primary ciliary dyskinesia type 11 (CILD11).

[0089] In some cases, the composition comprises a nucleic acid construct encoding radial spoke head 9 homolog (RSPH9), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the RSPH9 gene may be a component the radial spoke head in motile cilia and flagella. Mutations in this gene have been associated with primary ciliary dyskinesia type 12 (CILD12).

[0090] In some cases, the composition comprises a nucleic acid construct encoding sperm associated antigen 1 (SPAG1), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the SPAG1 gene may play a role in the cytoplasmic assembly of the ciliary dynein arms. Mutations in this gene have been associated with primary ciliary dyskinesia type 28 (CILD28).

[0091] In some cases, the composition comprises a nucleic acid construct encoding zinc finger MYND-type containing 10 (ZMYND10), and upon translation within the cells of a subject the construct yields a polypeptide that treats a subject having or at risk of having of primary ciliary dyskinesia. The protein encoded by the ZMYND10 can function in axonemal assembly of inner and outer dynein arms (IDA and ODA, respectively) for proper axoneme building for cilia motility. Mutations in this gene have been associated with primary ciliary dyskinesia type 22 (CILD22).

[0092] The treatment may comprise treating a subject (*e.g.*, a patient with a disease and/or a lab animal with a condition). In some cases, the condition is primary ciliary dyskinesia (PCD) or Kartagener syndrome. In some cases, the condition is broadly associated with defects in one or more proteins that function within cell structures known as cilia. In some cases, the subject is a human. Treatment may be provided to the subject before clinical onset of disease. Treatment may be provided to the subject after clinical onset of disease. Treatment may be provided to the subject on or after 1 minute, 5 minutes, 10 minutes, 30 minutes, 1 hour, 2 hours, 3 hours, 4 hours, 5 hours, 6 hours, 12 hours, 1 day, 1 week, 6 months, 12 months, or 2 years after clinical onset of the disease. Treatment may be provided to the subject for a time period that is greater than or equal to 1 minute, 10 minutes, 30 minutes, 1 hour, 2 hours, 3 hours, 4 hours, 5 hours, 6 hours, 12 hours, 1 day, 1 week, 1 month, 6 months, 12 months, 2 years or more after clinical onset of the disease. Treatment may be provided to the subject for a time period that is less than or equal to 2 years, 12 months, 6 months, 1 month, 1 week, 1 day, 12 hours, 6 hours, 5 hours, 4

hours, 3 hours, 2 hours, 1 hour, 30 minutes, 10 minutes, or 1 minute after clinical onset of the disease. Treatment may also include treating a human in a clinical trial.

[0093] Compositions containing the engineered polyribonucleotides described herein can be administered for prophylactic and/or therapeutic treatments. In therapeutic applications, the nucleic acid constructs or vectors can be administered to a subject already suffering from a disease, such as a primary ciliary dyskinesia, in the amount sufficient to provide the amount of the encoded polypeptide that cures or at least improves the symptoms of the disease. Nucleic acid constructs, vectors, engineered polyribonucleotides, or compositions can also be administered to lessen a likelihood of developing, contracting, or worsening a disease. Amounts effective for this use can vary based on the severity and course of the disease or condition, the efficiency of transfection of a nucleic acid construct(s), vector(s), engineered polyribonucleotide(s), or composition(s), the affinity of an encoded polypeptide to a target molecule, previous therapy, the subject's health status, weight, response to the drugs, and the judgment of the treating physician.

[0094] In some cases, a polynucleotide of the disclosure can encode a polypeptide that is at least 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% homologous to a protein associated with primary ciliary dyskinesia, such as armadillo repeat containing 4 (ARMC4), chromosome 21 open reading frame 59 (C21orf59), coiled-coil domain containing 103 (CCDC103), coiled-coil domain containing 114 (CCDC114), coiled-coil domain containing 39 (CCDC39), coiled-coil domain containing 40 (CCDC40), coiled-coil domain containing 65 (CCDC65), cyclin O (CCNO), dynein (axonemal) assembly factor 1 (DNAAF1), dynein (axonemal) assembly factor 2 (DNAAF2), dynein (axonemal) assembly factor 3 (DNAAF3), dynein (axonemal) assembly factor 5 (DNAAF5), dynein axonemal heavy chain 11 (DNAH11), dynein axonemal heavy chain 5 (DNAH5), dynein axonemal heavy chain 6 (DNAH6), dynein axonemal heavy chain 8 (DNAH8), dynein axonemal intermediate chain 2 (DNAI2), dynein axonemal light chain 1 (DNAL1), dynein regulatory complex subunit 1 (DRC1), dyslexia susceptibility 1 candidate 1 (DYX1C1), growth arrest specific 8 (GAS8), axonemal central pair apparatus protein (HYDIN), leucine rich repeat containing 6 (LRRC6), NME/NM23 family member 8 (NME8), oral-facial-digital syndrome 1 (OFD1), retinitis pigmentosa GTPase regulator (RPGR), radial spoke head 1 homolog (Chlamydomonas) (RSPH1), radial spoke head 4 homolog A (Chlamydomonas) (RSPH4A), radial spoke head 9 homolog (Chlamydomonas) (RSPH9), sperm associated antigen 1 (SPAG1), and zinc finger MYND-type containing 10 (ZMYND10).

[0095] Multiple nucleic acid constructs, vectors, engineered polyribonucleotides, or compositions can be administered in any order or simultaneously. The nucleic acid constructs, vectors, engineered polyribonucleotides, or compositions can be packed together or separately, in a single package comprising polyribonucleotides that target the same target molecule or in a plurality of packages. One or all of the nucleic acid constructs, vectors, engineered polyribonucleotides, or compositions can be given in multiple doses. If not simultaneous, the timing between the multiple doses may vary.

[0096] The nucleic acid constructs, vectors, engineered polyribonucleotides, or compositions can be administered to a subject as soon as possible after the onset of the symptoms. A nucleic acid construct(s), a vector(s), engineered polyribonucleotide(s), or compositions can be administered as soon as is practical after the onset of a disease or condition is detected or suspected, and for a length of time necessary for the treatment of the disease, such as, for example, for about 1 month, for about 6 months, for about 12 months, for about 18 months, for about 24 months, or any appropriate length of time. The length of treatment can vary for each subject.

Altered nucleotide usage in coding regions to increase mRNA stability for transcript therapy

[0097] Hydrolysis of oligonucleotides suggests that the reactivity of the phosphodiester bond linking two ribonucleotides in single-stranded (ss)RNA depends on the nature of those nucleotides. At pH 8.5, dinucleotide cleavage susceptibility when embedded in ssRNA dodecamers may vary by an order of magnitude. Under near physiological conditions, hydrolysis of RNA usually involves an S_N2-type attack by the 2'-oxygen nucleophile on the adjacent phosphorus target center on the opposing side of the 5'-oxyanion leaving group, yielding two RNA fragments with 2',3'-cyclic phosphate and 5'-hydroxyl termini. More reactive scissile phosphodiester bonds may include 5'-UpA-3' (R₁ = U, R₂ = A) and 5'-CpA-3' (R₁ = C, R₂ = A) because the backbone at these steps can most easily adopt the "in-line" conformation that is required for S_N2-type nucleophilic attack by the 2'-OH on the adjacent phosphodiester linkage. In addition, interferon-regulated dsRNA-activated antiviral pathways produce 2'-5' oligoadenylates which bind to ankyrin repeats leading to activation of RNase L endoribonuclease. RNase L cleaves ssRNA efficiently at UA and UU dinucleotides. Lastly, U-rich sequences are potent activators of RNA sensors including Toll-like receptor 7 and 8 and RIG-I making global uridine content reduction a potentially attractive approach to reduce immunogenicity of therapeutic mRNAs.

[0098] Altered nucleotide usage schemes aiming to reduce the number of more reactive 5'-U(U/A)-3' dinucleotides within codons as well as across codons of modified mRNAs partially

alleviate limitations imposed by the inherent chemical instability of RNA. At the same time, lowering the U-content in RNA transcripts renders them less immunogenic. The present disclosure relates to RNA transcripts comprising altered open reading frames (ORF). In particular, a method comprising a substantial reduction of 5'-U(U/A)-3' dinucleotides within protein coding regions leading to stabilized therapeutic mRNAs is proposed.

TABLE 2

Construct	DNA sequence (from 5' to 3')
DNAII GeneScript Codon	ATGATCCCAGCAAGCGCCAAGGCACCACACAAGCAGCCCCACAAGCAGA GCATCTCCATCGGCAGGGGCACAAGGAAGAGGGACGAGGATAGCGGAAC CGAAGTGGGAGAGGGAAACAGACGAGTGGGCACAGTCCAAGGCAACCGTG CGCCACCTGACCAGCTGGAGCTGACAGATGCCGAGCTGAAGGAGGAGT TCACCAGGATCCTGACAGCCAACAATCCACACGCCCCCCAGAACATCGTG CGTACTCTTTCAAGGAGGGCACATATAAGCCAATCGGCTTTGTGAACCA GCTGGCCGTGCACTATACCCAAGTGGGCAATCTGATCCCCAAGGACTCCG ATGAGGGGCCGGAGACAGCACTACAGGGACGAGCTGGTGGCAGGATCCCA GGAGTCTGTGAAAGTGATCTCTGAGACCGGCAATCTGGAGGAGGACGAG GAGCCAAAGGAGCTGGAGACCGAGCCAGGAAGCCAGACAGATGTGCCTG CAGCAGGAGCAGCAGAGAAGGTGACCGAGGAGGAGCTGATGACACCTAA GCAGCCAAAGGAGCGGAAGCTGACCAACCAGTTCAATTTTTCCGAGAGA GCCTCTCAGACATAACAATCCAGTGCGGGACAGAGAGTGCCAGACCG AGCCACCCCCTAGAACCAACTTTTTCCGCCACAGCCAATCAGTGGGAGATC TACGATGCCTATGTGGAGGAGCTGGAGAAGCAGGAGAAGACCAAGGAGA AGGAGAAGGCCAAGACACCCGTGGCCAAGAAGTCCGGCAAGATGGCCAT GCGGAAGCTGACCAGCATGGAGTCCCAGACAGACGATCTGATCAAGCTG TCTCAGGCCGCCAAGATCATGGAGAGAATGGTGAACCAGAATACCTATG ACGATATCGCCCAGGACTTCAAGTACTATGACGATGCAGCAGACGAGTAC AGGGATCAAGTGGGCACACTGCTGCCTCTGTGGAAGTTTCAGAACGATAA GGCCAAGAGGCTGAGCGTGACCGCCCTGTGCTGGAATCCAAAGTACAGG GACCTGTTTCGACAGTGGGATACGGATCTTATGACTTCATGAAGCAGAGCAG AGGCATGCTGCTGCTGTATTCCCTGAAGAACCCCTCTTTCCCTGAGTACAT GTTTAGCTCCAATTCCGGCGTGATGTGCCTGGACATCCACGTGGATCACC CCTACCTGGTGGCCGTGGGCCACTATGACGGCAACGTGGCCATCTACAAT CTGAAGAAGCCTCACTCTCAGCCCAGCTTCTGTTCTAGCGCCAAGAGCGG CAAGCACTCCGATCCCGTGTGGCAGGTGAAGTGGCAGAAGGACGATATG GACCAGAACCTGAATTTCTTTTCCGTGTCCTCTGATGGCAGGATCGTGTCT TGGACCCTGGTGAAGCGCAAGCTGGTGCACATCGACGTGATCAAGCTGA AGGTGGAGGGCAGCACCACAGAGGTGCCAGAGGGACTGCAGCTGCACCC AGTGGGATGCGGCACAGCCTTCGACTTTCACAAGGAGATCGATTATATGT TCCTGGTGGGCACCGAGGAGGGCAAGATCTACAAGTGTCTAAGAGCTAT AGCTCCAGTTTCTGGACACATATGATGCCACAACATGAGCGTGGATAC CGTGTCTGGAATCCTTACCACACAAAGGTGTTTCATGAGCTGCTCTAGCG ACTGGACCGTGAAGATCTGGGATCACACCATCAAGACACCTATGTTTATC TATGACCTGAACTCCGCCGTGGGCGATGTGGCATGGGCACCATACTCCTC TACAGTGTTTCGACAGTACCACAGACGGCAAGGCACACATCTTTGATC TGGCCATCAACAAGTACGAGGCCATCTGTAATCAGCCCGTGGCCGCCAAG AAGAACAGGCTGACCCACGTGCAGTTCAATCTGATCCACCCTATCATCAT CGTGGGCGACGATCGGGGCCACATCATCTCTCTGAAGCTGAGCCCCAACC TGAGAAAGATGCCTAAGGAGAAGAAGGGACAGGAGGTGCAGAAGGGAC

	CAGCAGTGGAGATCGCAAAGCTGGACAAGCTGCTGAATCTGGTGCGCGA GGTGAAGATCAAGACCTGA (SEQ ID NO: 14)
DNAI1 Altered Nucleotide Usage 1	ATGATCCCAGCAAGCGCCAAGGCACCACACAAGCAGCCCCACAAGCAGA GCATCAGCATCGGCAGGGGCACAAGGAAGAGGGACGAGGACAGCGGAA CCGAAGTGGGAGAGGGAACAGACGAGTGGGCACAGAGCAAGGCAACCG TGCGCCACCCGACCAGCTGGAGCTGACAGACGCCGAGCTGAAGGAGGA GTTACCAGGATCCTGACAGCCAACAACCCACACGCCCCCCAGAACATCG TGCGCTACAGCTTCAAGGAGGGCACATACAAGCCAATCGGCTTCGTGAAC CAGCTGGCCGTGCACTACACCCAAGTGGGCAACCTGATCCCCAAGGACA GCGACGAGGGCCGGAGACAGCACTACAGGGACGAGCTGGTGGCAGGAA GCCAGGAGAGCGTGAAAGTGATCAGCGAGACCGGCAACCTGGAGGAGGA CGAGGAGCCAAAGGAGCTGGAGACCGAGCCAGGAAGCCAGACAGACGT GCCCGCAGCAGGAGCAGCAGAGAAGGTGACCGAGGAGGAGCTGATGAC ACCAAGCAGCCAAAGGAGCGGAAGCTGACCAACCAGTTCAACTTCAGC GAGAGAGCCAGCCAGACATAACAACCCAGTGCGGGACAGAGAGTGCC AGACCGAGCCACCCCCAGAACCAACTTCAGCGCCACAGCCAACCAAGTG GGAGATCTACGACGCCTACGTGGAGGAGCTGGAGAAGCAGGAGAAGACC AAGGAGAAGGAGAAGGCCAAGACACCCGTGGCCAAGAAGAGCGGCAAG ATGGCCATGCGGAAGCTGACCAGCATGGAGAGCCAGACAGACGACCTGA TCAAGCTGAGCCAGGCCGCAAGATCATGGAGAGAATGGTGAACCAGAA CACCTACGACGACATCGCCAGGACTTCAAGTACTACGACGACGCAGCA GACGAGTACAGGGACCAAGTGGGCACACTGCTGCCCTGTGGAAGTTCC AGAACGACAAGGCCAAGAGGCTGAGCGTGACCGCCCTGTGCTGGAACCC AAAGTACAGGGACCTGTTTCGAGTGGGATACGGAAGCTACGACTTCATG AAGCAGAGCAGAGGCATGCTGCTGCTGTACAGCCTGAAGAACCCAGCT TCCCCGAGTACATGTTACAGCAGCAACAGCGGCGTGATGTGCCTGGACATC CACGTGGACCACCCCTACCTGGTGGCCGTGGGCCACTACGACGGCAACGT GGCCATCTACAACCTGAAGAAGCCCCACAGCCAGCCAGCTTCTGCAGCA GCGCCAAGAGCGGCAAGCACAGCGACCCCGTGTGGCAGGTGAAGTGGCA GAAGGACGACATGGACCAGAACCTGAACTTCTTCAGCGTGAGCAGCGAC GGCAGGATCGTGAGCTGGACCCTGGTGAAGCGCAAGCTGGTGCACATCG ACGTGATCAAGCTGAAGGTGGAGGGCAGCACACAGAGGTGCCAGAGGG ACTGCAGCTGCACCCAGTGGGATGCGGCACAGCCTTCGACTTCCACAAGG AGATCGACTACATGTTCTTGGTGGGCACCGAGGAGGGCAAGATCTACAA GTGCAGCAAGAGCTACAGCAGCCAGTTCCTGGACACATACGACGCCAC AACATGAGCGTGGACACCGTGAGCTGGAACCCCTACCACACAAGGTGT TCATGAGCTGCAGCAGCGACTGGACCGTGAAGATCTGGGACCACACCATC AAGACACCCATGTTTCATCTACGACCTGAACAGCGCCGTGGGCGACGTGGC ATGGGCACCATACAGCAGCACAGTGTTCGACAGCAGTGACCACAGACGGC AAGGCACACATCTTCGACCTGGCCATCAACAAGTACGAGGCCATCTGCAA CCAGCCCGTGGCCGCAAGAAGAACAGGCTGACCCACGTGCAGTTCAAC CTGATCCACCCCATCATCATCGTGGGCGACGACCGGGGCCACATCATCAG CCTGAAGCTGAGCCCCAACCTGAGAAAGATGCCCAAGGAGAAGAAGGGA CAGGAGGTGCAGAAGGGACCAGCAGTGGAGATCGCAAAGCTGGACAAGC TGCTGAACCTGGTGCGCGAGGTGAAGATCAAGACCTGA (SEQ ID NO: 15)
DNAI1 Altered Nucleotide Usage 2	ATGATCCCAGCAAGCGCCAAGGCACCACACAAGCAGCCCCACAAGCAGA GCATCTCCATCGGCAGGGGCACAAGGAAGAGGGACGAGGACAGCGGAAC CGAAGTGGGAGAGGGAACAGACGAGTGGGCACAGTCCAAGGCAACCGTG CGCCACCTGACCAGCTGGAGCTGACAGATGCCGAGCTGAAGGAGGAGT TCACCAGGATCCTGACAGCCAACAATCCACACGCCCCCCAGAACATCGTG CGCTACAGCTTCAAGGAGGGCACATACAAGCCAATCGGCTTCGTGAACCA

	<p>GCTGGCCGTGCACTACACCCAAGTGGGCAATCTGATCCCCAAGGACTCCG ATGAGGGCCGGAGACAGCACTACAGGGACGAGCTGGTGGCAGGATCCCA GGAGTCTGTGAAAGTGATCTCTGAGACCGGCAATCTGGAGGAGGACGAG GAGCCAAAGGAGCTGGAGACCGAGCCAGGAAGCCAGACAGATGTGCCTG CAGCAGGAGCAGCAGAGAAGGTGACCGAGGAGGAGCTGATGACACCCA AGCAGCCAAAGGAGCGGAAGCTGACCAACCAGTTCAACTTCTCCGAGAG AGCCTCTCAGACATAACAATCCAGTGCGGGACAGAGAGTGCCAGACC GAGCCACCCCCAGAACCAACTTCTCCGCCACAGCCAATCAGTGGGAGAT CTACGATGCCTACGTGGAGGAGCTGGAGAAGCAGGAGAAGACCAAGGAG AAGGAGAAGGCCAAGACACCCGTGGCCAAGAAGTCCGGCAAGATGGCCA TGCGGAAGCTGACCAGCATGGAGTCCAGACAGACGATCTGATCAAGCT GTCTCAGGCCGCCAAGATCATGGAGAGAATGGTGAACCAGAACACCTAC GACGACATCGCCCAGGACTTCAAGTACTACGACGATGCAGCAGACGAGT ACAGGGATCAAGTGGGCACACTGCTGCCTCTGTGGAAGTTCAGAACGAC AAGGCCAAGAGGCTGAGCGTGACCGCCCTGTGCTGGAATCCAAAGTACA GGGACCTGTTCGCAGTGGGATACGGAAGCTACGACTTCATGAAGCAGAG CAGAGGCATGCTGCTGCTGTACTCCCTGAAGAACCCAGCTTCCCTGAGT ACATGTTCAGCTCCAACCTCCGGCGTGATGTGCCTGGACATCCACGTGGAT CACCCCTACCTGGTGGCCGTGGGCCACTACGACGGCAACGTGGCCATCTA CAATCTGAAGAAGCCTCACTCTCAGCCCAGCTTCTGCAGCAGCGCCAAGA GCGGCAAGCACTCCGATCCCGTGTGGCAGGTGAAGTGGCAGAAGGACGA CATGGACCAGAACCTGAACTTCTTCTCCGTGTCTCTGATGGCAGGATCG TGAGCTGGACCCTGGTGAAGCGCAAGCTGGTGCACATCGACGTGATCAA GCTGAAGGTGGAGGGCAGCACACAGAGGTGCCAGAGGGACTGCAGCTG CACCCAGTGGGATGCGGCACAGCCTTCGACTTCCACAAGGAGATCGACTA CATGTTCCCTGGTGGGCACCGAGGAGGGCAAGATCTACAAGTGCAGCAAG AGCTACAGCTCCCAGTTCCTGGACACATACGATGCCACAACATGAGCGT GGACACCGTGTCTGGAATCCCTACCACACAAAGGTGTTTCATGAGCTGCA GCAGCGACTGGACCGTGAAGATCTGGGATCACACCATCAAGACACCCAT GTTTCATCTACGACCTGAACTCCGCCGTGGGCGATGTGGCATGGGCACCAT ACTCCAGCACAGTGTTTCGCAGCAGTGACCACAGACGGCAAGGCACACAT CTTCGATCTGGCCATCAACAAGTACGAGGCCATCTGCAATCAGCCCGTGG CCGCCAAGAAGAACAGGCTGACCCACGTGCAGTTCAATCTGATCCACCCC ATCATCATCGTGGGCGACGATCGGGGCCACATCATCTCTCTGAAGCTGAG CCCCAACCTGAGAAAGATGCCCAAGGAGAAGAAGGGACAGGAGGTGCA GAAGGGACCAGCAGTGGAGATCGCAAAGCTGGACAAGCTGCTGAATCTG GTGCGCGAGGTGAAGATCAAGACCTGA (SEQ ID NO: 16)</p>
<p>DNAH5 Altered Nucleotide Usage 1</p>	<p>ATGTTCAGAATCGGCAGACGGCAGCTGTGGAAGCACAGCGTGACCAGAG TGCTGACCCAGCGGCTGAAGGGCGAGAAAGAGGCCAAGAGAGCCCTGCT GGACGCCCGGCACAAcTACCTGTTTCGCCATCGTGGCCAGCTGCCTGGACC TGAACAAGACCGAGGTGGAAGACGCCATCCTGGAAGGCAACCAGATCGA GCGGATCGACCAGCTGTTTCGCCGTGGGCGGACTGCGGCACCTGATGTTCT ACTACCAAGACGTGGAAGAGGCCGAGACAGGCCAGCTGGGAAGCCTGGG CGGAGTGAACCTGGTGGAGCGGCAAGATCAAGAAACCCAAGGTGTTTCGTG ACCGAGGGCAACGACGTGGCCCTGACAGGCGTGTGCGTGTTCATCAG AACCGACCCCAGCAAGGCCATCACCCCGACAACATCCACCAGGAAGTG AGCTTCAACATGCTGGACGCCGCCGACGGCGGCCTGCTGAACAGCGTGCG GAGACTGCTGAGCGACATTTTCATCCCCGCCCTGAGAGCCACAAGCCACG GCTGGGGAGAGCTGGAAGGACTGCAGGACGCCGCCAACATCCGGCAGGA ATTCTGAGCAGCCTGGAAGGATTCGTGAACGTGCTGAGCGGCGCCCAGG AAAGCCTGAAAGAAAAGTGAACCTGCGGAAGTGCGACATCCTGGAAC</p>

GAAAACCCTGAAAGAGCCCACCGACTACCTGACCCTGGCCAACAACCCC
GAGACACTGGGCAAGATCGAGGACTGCATGAAAGTGTGGATCAAGCAGA
CCGAACAGGTGCTGGCCGAGAACAACCAGCTGCTGAAAGAAGCCGACGA
CGTGGGCCCAAGAGCCGAGCTGGAACACTGGAAGAAGCGGCTGAGCAAG
TTCAACTACCTGCTGGAACAGCTGAAGAGCCCCGACGTGAAGGCCGTGCT
GGCCGTGCTGGCAGCCGCCAAGAGCAAACCTGCTGAAAACCTGGCGCGAG
ATGGACATCCGGATCACCGACGCCACCAACGAGGCCAAGGACAACGTGA
AGTACCTGTACACCCTGGAAAAGTGTGCTGCGACCCCCTGTACAGCAGCGAC
CCCCTGAGCATGATGGACGCCATCCCCACCCTGATCAACGCCATCAAGAT
GATCTACAGCATCAGCCACTACTACAACACCAGCGAGAAGATCACCGAC
CTGTTTCGTGAAAGTGACCAACCAGATCATCAGCGCCTGCAAGGCCTACAT
CACCAACAACGGCACCCGCCAGCATCTGGAACCAGCCCCAGGACGTGGTG
GAAGAGAAGATCCTGAGCGCCATCAAGCTGAAGCAGGAATACCAGCTGT
GCTTCCACAAGACCAAGCAGAAGCTGAAACAGAACCCCAACGCCAAGCA
GTTTCGACTTCAGCGAGATGTACATCTTCGGCAAGTTCGAGACATTCCACC
GGCGGCTGGCCAAGATCATCGACATCTTACCACCCTGAAAACATACAGC
GTGCTGCAGGACAGCACCATCGAGGGCCTGGAAGACATGGCCACCAAGT
ACCAGGGCATCGTGGCCACCATCAAGAAGAAAGAGTACAACCTCCTGGA
CCAGCGCAAGATGGACTTCGACCAGGACTACGAGGAATTCTGCAAGCAG
ACAAACGACCTGCACAACGAGCTGCGCAAGTTCATGGACGTGACCTTCGC
CAAGATCCAGAACACCAACCAGGCCCTGCGGATGCTGAAGAAGTTCGAG
AGACTGAACATCCCCAACCTGGGCATCGACGACAAGTACCAGCTGATCCT
GGAAAACCTACGGCGCCGACATCGACATGATCAGCAAGCTGTACACAAAG
CAGAAGTACGACCCCCCTGGCCCGGAACCAGCCCCCATCGCCGGCAA
AATCCTGTGGGCCAGACAGCTGTTCCACCGGATCCAGCAGCCCATGCAGC
TGTTCCAGCAGCACCCCGCCGTGCTGAGCACAGCCGAGGCCAAACCCATC
ATCCGGAGCTACAACCGGATGGCCAAGGTGCTGCTGGAATTCGAGGTGCT
GTTCCACCGGGCCTGGCTGCGGCAGATCGAAGAGATCCACGTGGGACTG
GAAGCCAGCCTGCTCGTGAAGGCCCCCGGAACCGGCGAGCTGTTTCGTGA
ACTTCGACCCCCAGATCCTGATCCTGTTCCGGGAAACCGAGTGCATGGCC
CAGATGGGGCTGGAAGTGAGCCCCCTGGCCACCAGCCTGTTCCAGAAGC
GGGACCGGTACAAGCGGAACTTCAGCAACATGAAGATGATGCTGGCCGA
GTACCAGCGCGTGAAGAGCAAGATCCCCGCCGCCATCGAGCAGCTGATC
GTGCCCCACCTGGCCAAAGTGGACGAGGCCCTGCAGCCAGGACTGGCCG
CCCTGACATGGACCAGCCTGAACATCGAGGCCTACCTGGAAAACACATTC
GCCAAAATCAAGGACCTGGAACCTGCTGCTGGACCGCGTGAACGACCTGA
TCGAGTTCGGATCGACGCCATCCTGGAAGAGATGAGCAGCACCCCCCTG
TGCCAGCTGCCCCAGGAAGAACCCTGACCTGCGAAGAGTTCCTGCAGAT
GACCAAGGACCTGTGCGTGAACGGCGCCCAGATCCTGCACTTCAAGAGC
AGCCTGGTGAAGAAGCCGTGAACGAGCTCGTGAACATGCTGCTGGACG
TGGAAGTGCTGAGCGAGGAAGAGAGCGAGAAGATCAGCAACGAGAACA
GCGTGAACCTACAAGAACGAGAGCAGCGCCAAGCGGGAAGAGGGCAACTT
CGACACCCTGACCAGCAGCATCAACGCCAGAGCCAACGCCCTGCTGCTGA
CCACCGTGACCCGGAAGAAAAAAGAAACCGAGATGCTGGGCGAAGAGGC
CAGAGAGCTGCTGAGCCACTTCAACCACCAGAACATGGACGCCCTGCTGA
AAGTGACACGGAACACCCTGGAAGCCATCCGGAAGCGGATCCACAGCAG
CCACACCATCAACTTCGGGACAGCAACAGCGCCAGCAACATGAAGCAG
AACAGCCTGCCATCTTCCGGGCCAGCGTGACACTGGCCATCCCCAACAT
CGTGATGGCCCCCGCCCTGGAAGACGTGCAGCAGACACTGAACAAGGCC
GTGGAATGCATCATCAGCGTGCCCAAGGGCGTGCGGCAGTGGAGCAGCG
AACTGCTGAGCAAGAAGAAGATCCAGGAACGGAAAATGGCCGCCCTGCA

GAGCAACGAGGACAGCGACAGCGACGTGGAAATGGGCGAGAACGAGCT
GCAGGACACACTGGAAATCGCCAGCGTGAACCTGCCCATCCCCGTGCAG
ACCAAGAATACTACAAGAACGTGAGCGAAAACAAAGAAATCGTGAAGC
TGGTGAGCGTGCTGAGCACCATCATCAACAGCACCAAGAAAGAAGTGAT
CACCAGCATGGACTGCTTCAAGCGGTACAACCACATCTGGCAGAAGGGC
AAAGAAGAGGGCCATCAAGACCTTCATCACCCAGAGCCCCCTGCTGAGCG
AGTTCGAGAGCCAGATCCTGTACTTCCAGAACCTGGAACAGGAAATCAAC
GCCGAGCCCGAGTACGTGTGCGTGGGCAGCATCGCCCTGTACACCGCCGA
CCTGAAGTTCGCCCTGACCGCCGAGACAAAGGCCTGGATGGTTCGTGATCG
GCCGGCACTGCAACAAAAAGTACAGAAGCGAGATGGAAAACATCTTCAT
GCTGATCGAGGAATTCAACAAGAACTGAACCGGCCATCAAGGACCTG
GACGACATCAGAATCGCCATGGCCGCACTGAAAGAGATCAGAGAGGAAAC
AGATCAGCATCGACTTCCAAGTGGGCCCCATCGAGGAAAGCTACGCCCTG
CTGAACAGATACGGACTGCTGATCGCCCGGGAAGAGATCGACAAGGTGG
ACACCTTGCACTACGCCTGGGAGAAGCTGCTGGCCAGAGCCGGCGAGGT
GCAGAACAACTGGTGAGCCTGCAGCCAGCTTCAAGAAAGAACTGATC
AGCGCCGTGGAAGTGTTCTGCAGGACTGCCACCAGTTCTACCTGGACTA
CGACCTGAACGGCCCCATGGCCAGCGGCCTGAAACCCCAAGGAAGCCAGC
GACCGGCTGATCATGTTCCAGAACCAGTTCGACAACATCTACCGGAAGTA
CATCACCTACACAGGCGGCGAGGAACTGTTCCGGCCTGCCCGCCACACAGT
ACCCCCAGCTGCTGGAAATCAAGAAGCAGCTGAACCTGCTGCAGAAGAT
CTACACCCTGTACAACAGCGTGATCGAGACAGTGAACAGCTACTACGACA
TCCTGTGGAGCGAAGTGAACATCGAGAAGATCAACAACGAACTGCTGGA
ATTCCAGAACCGGTGCCGGAAGCTGCCAGAGCACTGAAGGACTGGCAG
GCCTTCTGGACCTGAAGAAAATCATCGACGACTTCAGCGAGTGCTGCC
CCTGCTGGAGTACATGGCCAGCAAGGCCATGATGGAACGGCACTGGGAG
AGAATCACCACTGACCGGCCACAGCCTGGACGTGGGCAACGAGAGCT
TCAAGCTGCGGAACATCATGGAAGCCCCACTGCTGAAGTACAAAGAGGA
AATCGAGGACATCTGCATCAGCGCCGTGAAAGAGCGGGACATCGAGCAG
AACTGAAACAAGTGATCAACGAGTGGGACAACAAGACCTTCACCTTCG
GCAGCTTCAAGACCAGAGGGCGAGCTGCTGCTGCGGGGCGACAGCACCAG
CGAGATCATCGCCAACATGGAAGACAGCCTGATGCTGCTGGGCAGCCTGC
TGAGCAACCGGTACAACATGCCCTTCAAGGCCAGATCCAGAAATGGGT
GCAGTACCTGAGCAACAGCACCGACATCATCGAGAGCTGGATGACCGTG
CAGAACCTGTGGATCTACCTGGAAGCCGTGTTCTGTTGGGCGGCGACATCGC
CAAGCAGCTGCCCAAAGAGGGCCAAGCGGTTACGCAACATCGACAAGAGC
TGGGTCAAGATCATGACCAGAGCCCACGAGGTGCCAGCGTGGTGCAGT
GCTGCGTGGGCGACGAAACACTGGGACAGCTGCTGCCCCACCTGCTGGAC
CAGCTGGAAATCTGCCAGAAGAGCCTGACCGGCTACCTGGAAAAGAAAC
GGCTGTGCTTCCCCCGGTTCTTCTTCGTGAGCGACCCCGCCCTGCTGGAAA
TCCTGGGCCAGGCCAGCGACAGCCACACAATCCAGGCCACCTGCTGAAC
GTGTTTCGACAACATCAAGAGCGTGAAGTTCCACGAGAAAATCTACGACC
GGATCCTGAGCATCAGCAGCCAGGAAGGCGAGACAATCGAGCTGGACAA
GCCCCGTGATGGCCGAGGGAAACGTGGAAGTGTGGCTGAACAGCCTGCTG
GAAGAGAGCCAGAGCAGCCTGCACCTCGTGATCAGACAGGCCGCCGCCA
ACATCCAGGAAACCGGCTTCCAGCTGACCGAGTTCCTGAGCAGCTTCCA
GCACAAGTGGGACTGCTGGGCATCCAGATGATCTGGACCAGAGACAGCG
AAGAGGCCCTGAGAAACGCCAAGTTCGACAAGAAAATCATGCAGAAAAC
AAACCAGGCATTCTGGAAGTGTGTAACACCTGATCGACGTGACCACCC
GGGACCTGAGCAGCACCGAGAGAGTGAAGTACGAGACACTGATCACCAT
CCACGTGCACCAGCGGGACATCTTCGACGACCTGTGCCACATGCACATCA

AGAGCCCCATGGACTTCGAGTGGCTGAAGCAGTGCAGGTTCTACTTCAAC
GAGGACAGCGACAAGATGATGATCCACATCACCGACGTGGCCTTCATCTA
CCAGAACGAGTTCCTGGGCTGCACCGACCGCCTCGTGATCACCCCTGA
CCGACCGGTGCTACATCACTGGCCCAGGCACTGGGCATGAGCATGGG
AGGCGCACCCAGCAGGACCCGCCGGCACAGGCAAGACCGAAACCACCAAG
GACATGGGACGCTGCCTGGGCAAATACGTGGTGGTGTTCAACTGCAGCGA
CCAGATGGACTTCCGGGGCCTGGGCCGGATCTTCAAGGGCCTGGCACAGA
GCGGAAGCTGGGGCTGCTTCGACGAGTTCAACAGAATCGACCTGCCCGTG
CTGAGCGTGGCCGCACAGCAGATCAGCATCATCCTGACATGCAAAAAAG
AGCACAAGAAGAGCTTCATCTTCACCGACGGCGACAACGTGACCATGAA
CCCCGAGTTCGGCCTGTTCTGACAATGAACCCCGGCTACGCCGGACGGC
AGGAACTGCCCGAGAACCTGAAGATCAACTTCGGGAGCGTGGCCATGAT
GGTGCCCGACCGGCAGATCATCATCAGAGTGAAACTGGCCAGCTGCGGCT
TCATCGACAACGTGGTGTGGCCCCGGAAGTTCTTCACACTGTACAAGCTG
TGCGAAGAACAGCTGAGCAAACAGGTGCACTACGACTTCGGCCTGAGGA
ACATCCTGAGCGTGCTGAGAACCCTGGGAGCCGCCAAGCGGGCCAACCC
CATGGACACCGAGAGCACAATCGTGATGCGGGTGCTGCGGGACATGAAC
CTGAGCAAGCTGATCGACGAGGACGAGCCCCTGTTCTGAGCCTGATCGA
GGACCTGTTCCCAACATCCTGCTGGACAAGGCCGGCTACCCCGAACTGG
AAGCCGCCATCAGCAGACAGGTGGAAGAGGCCGGCCTGATCAACCACCC
CCCCTGGAAACTGAAAGTGATCCAGCTGTTTCGAGACACAGCGCGTGCGGC
ACGGCATGATGACACTGGGACCCAGCGGAGCCGGCAAGACCACCTGCAT
CCACACACTGATGCGGGCCATGACCGACTGCGGCAAGCCCCACCGCGAG
ATGCGGATGAAC
CCCAAGGCCATCACCGCCCCCAGATGTTTCGGCAGACTGGACGTGGCCAC
CAACGACTGGACCGACGGCATCTTCAGCACCCCTGTGGCGCAAGACCCTGC
GGGCCAAGAAGGGCGAGCACATCTGGATCATCCTGGACGGCCCCGTGGA
CGCCATCTGGATCGAGAACCTGAACAGCGTGCTGGACGACAACAAGACA
CTGACCCTGGCCAACGGCGACCGGATCCCCATGGCCCCCAACTGCAAGAT
CATCTTCGAGCCCCACAACATCGACAACGCCAGCCCCGCCACCGTGAGCA
GAAACGGCATGGTGTTCATGAGCAGCAGCATCCTGGACTGGAGCCCCATC
CTGGAAGGCTTCCTGAAGAAGCGGAGCCCCCAGGAAGCCGAGATCCTGA
GACAGCTGTACACCGAGAGCTTCCCCGACCTGTACCGGTTCTGCATCCAG
AACCTGGAGTACAAGATGGAAGTGCTGGAAGCCTTCGTGATCACCCAGA
GCATCAACATGCTGCAGGGCCTGATCCCCCTGAAAGAACAGGGGCGGAGA
AGTGAGCCAGGCCACCTGGGCAGACTGTTTCGTGTTTCGCCCTGCTGTGGA
GCGCCGGCGCCGCCCTGGAACCTGGACGGAAGGCCGGAGACTGGAACCTGTG
GCTGCGGAGCAGACCCACCGGCACCCCTGGAACCTGCCCCACCAGCCGGA
CCCGGCGACACCGCCTTCGACTACTACGTGGCCCCCGACGGCACCTGGAC
CCACTGGAACACCCGGACCCAGGAATACTGTACCCAGCGACACCACCC
CCGAGTACGGCAGCATCCTGGTGCCCAACGTGGACAACGTGCGGACCGA
CTTCCTGATCCAGACAATCGCCAAGCAGGGAAAGGCCGTGCTGCTGATCG
GCGAGCAGGGCACAGCCAAGACCGTGATCATCAAGGGCTTCATGAGCAA
GTACGACCCCGAGTGCCACATGATCAAGAGCCTGAACTTCAGCAGCGCCA
CCACCCACTGATGTTCCAGCGGACCATCGAGAGCTACGTGGACAAGCGG
ATGGGCACCACCTACGGCCCCCAGCCGGCAAGAAAATGACCGTGTTTCAT
CGACGACGTGAACATGCCATCATCAACGAGTGGGGCGACCAAGTGACC
AACGAGATCGTGCGGCAGCTGATGGAACAGAACGGCTTCTACAACCTGG
AAAAGCCCGGCGAGTTCACCAGCATCGTGGACATCCAGTTCCTGGCCGCC
ATGATCCACCCCGGCGGGCGGAAGAAACGACATCCCCAGCGGCTGAAGC
GGCAGTTCAGCATCTTCAACTGCACCCTGCCAGCGAGGCCAGCGTGGAC

AAGATCTTCGGCGTGATCGGGCGTGGGCCACTACTGCACCCAGAGAGGCTT
CAGCGAGGAAGTGCGGGACAGCGTGACCAAGCTGGTGCCCCTGACAAGA
CGGCTGTGGCAGATGACCAAGATCAAGATGCTGCCACCCCCGCCAAGTT
CCACTACGTGTTCAACCTGCGGGACCTGAGCAGAGTGTGGCAGGGAATGC
TGAACACCACCAGCGAAGTGATCAAAGAGCCCAACGACCTGCTGAAGCT
GTGGAAGCACGAGTGCAAGAGAGTGATCGCCGACCGGTTACCCGTGAGC
AGCGACGTGACATGGTTCGACAAGGCCCTGGTGAGCCTGGTGGAAGAGG
AATTCGGCGAAGAGAAGAACTGCTGGTGGACTGCGGCATCGACACCTA
CTTCGTGGACTTCCTGCGCGACGCCCCGAAGCCGCCGGCGAGACAAGCG
AAGAGGCCGACGCCGAGACACCCAAGATCTACGAGCCCATCGAGAGCTT
CAGCCACCTGAAAGAAAGGCTGAACATGTTCTGCAGCTGTACAACGAG
AGCATCCGGGGAGCCGGCATGGACATGGTGTTCCTTCGCCGACGCCATGGT
GCACCTCGTGAAGATCAGCAGAGTGATCCGGACCCCCAGGGCAACGCC
CTGCTCGTGGGAGTGGGAGGCAGCGGCAAGCAGAGCCTGACCAGACTGG
CCAGCTTCATCGCCGGCTACGTGAGCTTCCAGATCACCTGACCCGGAGC
TACAACACCAGCAACCTGATGGAAGACCTGAAGGTGCTGTACCCGGACAG
CCGGCCAGCAGGGGAAGGGCATCACCTTCATCTTCACCGACAACGAGATC
AAGGACGAGAGCTTCTGGAGTACATGAACAACGTGCTGAGCAGCGGCG
AGGTGAGCAACCTGTTCGCCCGGGACGAGATCGACGAGATCAACAGCGA
CCTGGCCAGCGTGATGAAGAAAGAATTCCCCCGGTGCCTGCCACAAACG
AGAACCTGCACGACTACTTCATGAGCAGAGTGCGGCAGAACCTGCACATC
GTGCTGTGCTTCAGCCCCGTGGGCGAGAAGTTCAGAAACCGGGCCCTGAA
GTTCCCCGCCCTGATCAGCGGCTGCACCATCGACTGGTTCAGCCGGTGGC
CCAAGGACGCCCTGGTGGCCGTGAGCGAGCACTTCTGACCAGCTACGAC
ATCGACTGCAGCCTGGAAATCAAGAAAGAGGTGGTGCAGTGCATGGGCA
GTTCCAGGACGGCGTGGCCGAGAAATGCGTGGACTACTTCCAGCGGTTT
CGGCGGAGCACCCACGTGACCCCAAGAGCTACCTGAGCTTCATCCAGGG
CTACAAGTTCATCTACGGCGAGAAGCACGTGGAAGTGCGCACACTGGCC
AACCGGATGAACACCGGCCTGGAAAAACTGAAAGAGGCCAGCGAGAGCG
TGGCCGCCCTGAGCAAAGAAGTGGAAAGCCAAAGAAAAAGAACTGCAGGT
GGCCAACGACAAGGCCGACATGGTGTGAAAGAAGTGACCATGAAGGCC
CAGGCCGCCGAGAAAGTGAAAGCCGAGGTGCAGAAAGTGAAAGGCCG
GCCAGGCCATCGTGGACAGCATCAGCAAGGACAAGGCCATCGCCGAGG
AAAAGCTGGAAGCAGCCAAGCCC GCCCTGGAAGAGGCAGAAGCCGCCCT
GCAGACCATCCGGCCCAGCGACATCGCCACAGTGCAGGACCCTGGGAAGG
CCCCCCCACCTGATCATGCGGATCATGGACTGCGTGCTGCTGCTGTTCCA
GAGAAAGGTGAGCGCCGTGAAGATCGACCTGGAAAAAAGCTGCACCATG
CCCAGCTGGCAGGAAAGCCTGAAGCTGATGACCGCCGGCAACTTCTGCA
GAACCTGCAGCAGTTCCCCAAGGACACCATCAACGAGGAAGTGATCGAG
TTCTGAGCCCCTACTTCGAGATGCCCGACTACAACATCGAAACCGCCAA
ACGCGTGTGCGGCAACGTGGCCGGACTGTGCAGCTGGACCAAGGCCATG
GCCAGCTTCTTCAGCATCAACAAAGAGGTGCTGCCCTGAAGGCCAACCT
GGTGGTGCAGGAAAACCGGCACCTGCTGGCCATGCAGGACCTGCAGAAA
GCCAGGCCGAGCTGGACGACAAGCAGGCCGAGCTGGACGTGGTGCAGG
CCGAGTACGAGCAGGCCATGACCGAGAAGCAGACCCTGCTGGAAGACGC
AGAGCGGTGCAGACACAAGATGCAGACCGCCAGCACCCCTGATCAGCGGA
CTGGCCGGCGAAAAAGAGCGGTGGACCGAGCAGAGCCAGGAATTCGCCG
CCCAGACCAAGCGGCTCGTGGGAGACGTGCTGCTGGCCACCGCCTTCTG
AGCTACAGCGGCCCTTCAACCAGGAATTCAGGGACCTGCTGCTGAACGA
CTGGCGGAAAGAGATGAAGGCCAGAAAGATCCCCTTCGGCAAGAACCTG
AACCTGAGCGAGATGCTGATCGACGCCCCACCATCAGCGAGTGGAACT

GCAGGGACTGCCCAACGACGACCTGAGCATCCAGAACGGAATCATCGTG
ACCAAAGCCAGCAGATACCCCTGCTGATCGACCCCCAGACACAGGGCA
AGATCTGGATCAAGAACAAGAGAGCCGGAACGAGCTGCAGATCACCAG
CCTGAACCACAAGTACTTCCGGAACCACCTGGAAGACAGCCTGAGCCTGG
GCAGGCCACTGCTGATCGAGGACGTGGGCGAGGAACTGGACCCAGCCCT
GGACAACGTGCTGGAACGGAACCTTCATCAAGACCGGCAGCACCTTCAA
GTGAAAGTGGGCGACAAAGAAGTGGACGTGCTGGACGGCTTCCGGCTGT
ACATCACCACCAAGCTGCCCAACCCCGCCTACACCCCCGAGATCAGCGCC
CGGACCAGCATCATCGACTTCACCGTGACAATGAAGGGACTGGAAGACC
AGCTGCTGGGACGCGTGATCCTGACAGAGAAGCAGGAACTGGAAAAAGA
ACGGACCCACCTGATGGAAGACGTGACCGCCAACAAGCGGCGGATGAAG
GAACTGGAAGACAACCTGCTGTACAGGCTGACCAGCACCCAGGGCAGCC
TGGTGAAGACGAGAGCCTGATCGTGGTGTGAGCAACACCAAGCGGAC
CGCAGAGGAAGTGACCCAGAAGCTGGAAATCAGCGCCGAGACAGAGGTG
CAGATCAACAGCGCCAGAGAAGAGTACCGGCCCGTGGCCACCCGGGGAA
GCATCCTGTACTTCTGATCACCGAGATGCGGCTCGTGAACGAGATGTAC
CAGACCAGCCTGCGGCAGTTCCTGGGCCTGTTTCGACCTGAGCCTGGCCAG
AAGCGTGAAGAGCCCCATCACCGCAAGAGAATCGCCAACATCATCGAG
CACATGACCTACGAGGTGTACAAATACGCCGCCAGAGGCCTGTACGAGG
AACACAAGTTCCTGTTCACACTGCTGCTGACCCTGAAGATCGACATCCAG
CGGAACAGAGTGAAGCACGAAGAGTTCCTGACACTGATCAAGGGGGGAG
CCAGCCTGGACCTGAAGGCCTGCCCCCCCAAGCCAGCAAGTGGATCCTG
GACATCACCTGGCTGAACCTGGTGAAGCTGAGCAAGCTGAGACAGTTCA
GCGACGTGCTGGACCAGATCAGCCGCAACGAGAAGATGTGGAAGATCTG
GTTTCGACAAAGAGAACCCCGAGGAAGAACCCTGCCAACGCCTACGAC
AAGAGCCTGGACTGCTTCCGGCGGCTGCTGCTGATCAGAAGCTGGTGGCC
CGACCGGACAATCGCCAGGCCCGCAAGTACATCGTGGACAGCATGGGA
GAGAAGTACGCCGAGGGCGTGATCCTGGACCTGGAAAAGACCTGGGAGG
AAAGCGACCCAGAACCCCTGATCTGCCTGCTGAGCATGGGCAGCGAC
CCCACCGACAGCATCATCGCCCTGGGCAAGAGACTGAAGATCGAGACAA
GATACGTGAGCATGGGCCAGGGCCAGGAAGTGCACGCCAGAAAGCTGCT
GCAGCAGACCATGGCCAACGGCGGCTGGGCCCTGCTGCAGAACTGCCAC
CTGGGGCTGGACTTCATGGACGAACTGATGGACATCATCATCGAGACAGA
GCTGGTGCACGACGCCTTCAGACTGTGGATGACCACCGAGGCCACAAGC
AGTTCCCATCACCTGCTGCAGATGAGCATCAAGTTCGCCAACGACCCC
CCCAGGGACTGAGAGCCGGCCTGAAGAGAACCTACAGCGGCGTGAGCC
AGGACCTGCTGGACGTGAGCAGCGGCAGCCAGTGGAAAGCCATGCTGTA
CGCCGTGGCATTCTGCACAGCACCGTGCAGGAACGGCGGAAGTTCGGC
GCCCTGGGATGGAACATCCCCTACGAGTTCAACCAGGCCGACTTCAACGC
CACCGTGCAGTTCATCCAGAACCACCTGGACGACATGGACGTGAAGAAA
GGGGTGAGCTGGACAACCATCCGGTACATGATCGGAGAGATCCAGTACG
GCGGCAGAGTGACCGACGACTACGACAAGAGGCTGCTGAACACCTTCGC
CAAAGTGTGGTTCAGCGAGAACATGTTTCGGCCCCGACTTCAGCTTCTACC
AGGGCTACAACATCCCCAAGTGCAGCACCGTGGACAACCTACCTGCAGTAC
ATCCAGAGCCTGCCCGCCTACGACAGCCCCGAGGTGTTCCGACTGCACCC
CAACGCCGACATCACCTACCAGAGCAAACCTGGCCAAGGACGTGCTGGAC
ACCATCCTGGGCATCCAGCCCAAGGACACCAGCGGCGGAGGCGACGAAA
CCCGGGAAGCAGTGGTGGCCAGACTGGCCGACGACATGCTGGAAAAGCT
GCCCCCGACTACGTGCCCTTCGAAGTGAAAGAACGCCTGCAGAAGATG
GGCCCCCTTCCAGCCATGAACATCTTCTGAGGCAGGAAATCGACCGGAT
GCAGCGGGTGTGAGCCTCGTGCGGAGCACACTGACCGAGCTGAAACTG

	<p>GCCATCGACGGCACCATCATCATGAGCGAGAACCTGCGGGACGCACTGG ACTGCATGTTTCGACGCCAGAATCCCCGCATGGTGGAAAAAGGCCAGCTG GATCAGCAGCACCCCTGGGCTTCTGGTTCACCGAACTGATCGAGAGAAACA GCCAGTTCACCAGCTGGGTGTTCAACGGCAGACCCCACTGCTTCTGGATG ACCGGCTTCTTCAACCCACAAGGCTTCTGACAGCAATGCGCCAGGAAAT CACCAGAGCCAACAAGGGCTGGGCCCTGGACAACATGGTGTGTGCAAC GAAGTGACCAAGTGGATGAAGGACGACATCAGCGCCCCCCCCACAGAGG GCGTGTACGTGTACGGCCTGTACCTGGAAGGCGCCGGATGGGACAAGAG AAACATGAAGCTGATCGAGAGCAAGCCAAGGTGCTGTTTCGAGCTGATG CCCGTGATCAGGATCTACGCCGAGAACAACACCCTGAGGGACCCCGGTT CTACAGCTGCCCCATCTACAAGAAACCCGTGCGCACCGACCTGAACTACA TCGCCGCCGTGGACCTGAGGACAGCCAGACACCCGAGCACTGGGTGCT GAGAGGCGTGGCACTGCTGTGCGACGTGAAGTGA (SEQ ID NO: 17)</p>
<p>DNAH5 Altered Nucleotide Usage 2</p>	<p>ATGTTTCAGAATCGGCAGACGGCAGCTGTGGAAGCACAGCGTGACCAGAG TGCTGACCCAGCGGCTGAAGGGCGAGAAAGAGGCCAAGAGAGCCCTGCT GGACGCCCGGCACAATTACCTGTTTGCCATCGTGGCCAGCTGCCTGGACC TGAACAAGACCGAGGTGGAAGATGCCATCCTGGAAGGCAACCAGATCGA GCGGATCGACCAGCTGTTTGCCGTGGGCGGACTGCGGCACCTGATGTTCT ATTATCAAGACGTGGAAGAGGCCGAGACAGGCCAGCTGGGATCTCTGGG CGGAGTGAATCTGGTGTCCGGCAAGATCAAGAAACCAAGGTGTTTCGTG ACCGAGGGCAACGACGTGGCCCTGACAGGCGTGTGCGTGTTCATCAG AACCGACCCAGCAAGGCCATCACCCCGACAACATCCACCAGGAAGTG TCCTTCAACATGCTGGATGCCGCCGATGGCGGCCTGCTGAATTCTGTGCG GAGACTGCTGAGCGACATCTTCATCCCCGCCCTGAGAGCCACATCTCACG GCTGGGGAGAGCTGGAAGGACTGCAGGACGCCGCCAATATCCGGCAGGA ATTTCTGAGCAGCCTGGAAGGATTCGTGAACGTGCTGTCTGGCGCCAGG AAAGCCTGAAAGAAAAAGTGAACCTGCGGAAGTGCGATATCCTGGAAC GAAAACCCTGAAAGAGCCCACCGACTACCTGACCCTGGCCAACAACCCT GAGACACTGGGCAAGATCGAGGACTGCATGAAAGTGTGGATCAAGCAGA CCGAACAGGTGCTGGCCGAGAACAACCAGCTGCTGAAAGAAGCCGACGA CGTGGGCCCAAGAGCCGAGCTGGAACACTGGAAGAAGCGGCTGAGCAAG TTCAACTACCTGCTGGAACAGCTGAAGTCCCCCGACGTGAAGGCCGTGCT GGCTGTGCTGGCAGCCGCCAAGAGCAAACCTGCTGAAAACCTGGCGCGAG ATGGACATCCGGATCACCGACGCCACCAACGAGGCCAAGGACAACGTGA AGTACCTGTACACCCTGGAAAAGTGTGCTGCGACCCCTGTACAGCAGCGAC CCTCTGAGCATGATGGACGCCATCCCTACCCTGATCAACGCCATCAAGAT GATCTACAGCATCAGCCACTACTACAACACCAGCGAGAAGATCACCAGC CTGTTTCGTGAAAGTGACCAATCAGATCATCAGCGCCTGCAAGGCCTACAT CACCACAACGGCACCCGCCAGCATCTGGAACCAGCCCCAGGATGTGGTG GAAGAGAAGATCCTGTCTGCCATCAAGCTGAAGCAGGAATACCAGCTGT GTTTTCAAGACCAAGCAGAAGCTGAAACAGAACCCCAACGCCAAGCA GTTTCGACTTCAGCGAGATGTATATCTTCGGCAAGTTCGAGACATTCCACC GGCGGCTGGCCAAGATCATCGACATCTTACCACCCTGAAAACATACAGC GTGCTGCAGGACAGCACCATCGAGGGCCTGGAAGATATGGCCACCAAGT ACCAGGGCATTGTGGCCACCATCAAGAAGAAAGAGTACAACCTCCTGGA CCAGCGCAAGATGGACTTCGACCAGGACTACGAGGAATTCTGCAAGCAG ACAAACGACCTGCACAACGAGCTGCGCAAGTTTATGGACGTGACCTTCGC CAAGATCCAGAACACCAACCAGGCCCTGCGGATGCTGAAGAAGTTTGAG AGACTGAACATCCCCAACCTGGGCATCGACGATAAGTACCAGCTGATCCT GGAAAACCTACGGCGCCGACATCGACATGATCAGCAAGCTGTACACAAAG CAGAAGTACGACCCCCCCTGGCCCAGGAATCAGCCTCCTATCGCCGGCAA</p>

AATCCTGTGGGCTAGACAGCTGTTTCACCGGATCCAGCAGCCCATGCAGC
 TGTTCCAGCAGCACCCCTGCCGTGCTGAGCACAGCCGAGGCCAAACCCATC
 ATCCGGTCTTACAACCGGATGGCCAAGGTGCTGCTGGAATTCGAGGTGCT
 GTTCCACCGGGCCTGGCTGCGGCAGATCGAAGAGATTCACGTGGGACTGG
 AAGCCAGCCTGCTCGTGAAGGCTCCTGGAACCGGCCGAGCTGTTTGTGAAC
 TTCGACCCCCAGATCCTGATCCTGTTCCGGGAAACCGAGTGCATGGCCCA
 GATGGGGCTGGAAGTGTCTCCTCTGGCCACCTCCCTGTTCCAGAAGCGGG
 ACCGGTACAAGCGGAACTTCAGCAACATGAAGATGATGCTGGCTGAGTA
 CCAGCGCGTGAAGTCCAAGATCCCCGCTGCCATCGAGCAGCTGATCGTGC
 CTCACCTGGCCAAAGTGGACGAGGCCCTGCAGCCAGGACTGGCCGCTCTG
 ACATGGACCAGCCTGAACATCGAGGGCCTATCTGGAAAACACATTCGCCAA
 AATCAAGGATCTGGAAGTGTGCTGGACCGCGTGAACGACCTGATCGAGT
 TCCGGATCGACGCCATTCTGGAAGAGATGTCCAGCACCCCCCTGTGTGTCAG
 CTGCCCCAGGAAGAACCCCTGACCTGCGAAGAGTTCCTGCAGATGACCAA
 GGACCTGTGCGTGAACGGCGCCAGATTCTGCACTTCAAGTCCAGCCTGG
 TGAAGAAGCCGTGAACGAGCTCGTGAATATGCTGCTGGATGTGGAAGT
 GCTGAGCGAGGAAGAGTCCGAGAAGATCTCCAACGAGAACAGCGTGAAC
 TACAAGAACGAGTCCAGCGCCAAGCGGGAAGAGGGCAACTTCGACACCC
 TGACCAGCTCCATCAATGCCAGAGCCAACGCCCTGCTGCTGACCACCGTG
 ACCCGGAAGAAAAAAGAAACCGAGATGCTGGGCGAAGAGGCTAGAGAG
 CTGCTGTCCACTTCAACCACCAGAACATGGATGCCCTGCTGAAAGTGAC
 ACGGAATACCCTGGAAGCCATCCGGAAGCGGATCCACAGCAGCCACACC
 ATCAACTTCCGGGACAGCAACAGCGCCAGCAATATGAAGCAGAACAGCC
 TGCCCATCTTCCGGGCCTCCGTGACACTGGCCATCCCCAATATCGTGATG
 GCCCCTGCTCTGGAAGATGTGCAGCAGACACTGAACAAGGCCGTGGAAT
 GCATCATCTCCGTGCCAAGGGCGTGCGGCAGTGGTCTAGCGAACTGCTG
 TCCAAGAAGAAGATCCAGGAACGGAAAATGGCCGCCCTGCAGTCTAACG
 AGGACAGCGACTCCGACGTGGAAATGGGCGAGAATGAGCTGCAGGATAC
 ACTGGAATCGCCTCTGTGAATCTGCCCATCCCCGTGCAGACCAAGAACT
 ACTATAAGAACGTGTCCGAAAACAAAGAAATCGTGAAGCTGGTGTCTGT
 GCTGTCCACCATCATCAACAGCACCAAGAAAGAAGTGATCACCTCCATGG
 ACTGTTCAAGCGGTACAACCACATCTGGCAGAAGGGCAAAGAAGAGGC
 CATTAAAGACCTTCATCACCCAGAGCCCCCTGCTGTCCGAGTTCGAGTCTC
 AGATCCTGTACTTCCAGAACCTGGAACAGGAAATCAACGCCGAGCCCGA
 GTACGTGTGTGTGGGCTCTATCGCCCTGTATACCGCCGACCTGAAGTTCG
 CCCTGACCGCCGAGACAAAGGCCCTGGATGGTCTGATCGGCCGGCACTGC
 AACAAAAAGTACAGATCCGAGATGGAAAACATCTTTATGCTGATTGAGG
 AATTCACAAGAAACTGAACCGGCCCATTAAGGACCTGGACGACATCAG
 AATCGCCATGGCCGCACTGAAAGAGATCAGAGAGGAACAGATCAGCATC
 GACTTCCAAGTGGGCCCCATCGAGGAAAGCTACGCTCTGCTGAACAGATA
 CGGACTGCTGATCGCCCGGAAGAGATCGACAAGGTGGACACCCTGCAC
 TACGCCCTGGGAGAAGCTGCTGGCTAGAGCCGGCGAGGTGCAGAACAAC
 TGGTGTCTCTGCAGCCCAGCTTTAAGAAAGAAGTATCTCCGCCGTGGAA
 GTGTTTCTGCAGGACTGCCACCAGTTCTACCTGGACTACGACCTGAACGG
 CCCCATGGCCTCTGGCCTGAAACCTCAGGAAGCCTCCGACCGGCTGATTA
 TGTTTCAGAACCAGTTCGACAATATCTACCGGAAGTACATCACCTACACA
 GGCGGCGAGGAACTGTTCCGGCCTGCCTGCCACACAGTACCCCCAGCTGCT
 GGAAATCAAGAAGCAGCTGAACCTGCTGCAGAAGATCTACACCCTGTAC
 AACTCCGTGATCGAGACAGTGAACAGCTACTACGACATCCTGTGGAGCGA
 AGTGAACATTGAGAAGATTAACAATGAACTGCTGGAATTCAGAACCGGT
 GCCGGAAGCTGCCAGAGCACTGAAGGATTGGCAGGCCTTTCTGGATCTG

AAGAAAATCATCGACGACTTCTCCGAGTGCTGCCCTCTGCTGGAGTACAT
 GGCCTCCAAGGCCATGATGGAACGGCACTGGGAGAGAATCACCACACTG
 ACCGGCCACAGCCTGGACGTGGGCAACGAGAGCTTCAAGCTGCGGAACA
 TCATGGAAGCCCCACTGCTGAAGTACAAAGAGGAAATCGAGGACATCTG
 TATCAGCGCCGTGAAAGAGCGGGATATCGAGCAGAAACTGAAACAAGTG
 ATCAACGAGTGGGACAACAAGACCTTTACCTTCGGCAGCTTCAAGACCAG
 AGGCGAGCTGCTGCTGCGGGGCGATAGCACCTCTGAGATCATTGCCAACA
 TGGAAGATAGCCTGATGCTGCTGGGCTCCCTGCTGAGCAACCGGTATAAC
 ATGCCCTTCAAGGCTCAGATTCAGAAATGGGTGCAGTACCTGAGCAACTC
 CACCGACATCATCGAGTCCTGGATGACCGTGCAGAACCTGTGGATCTACC
 TGGAAGCCGTGTTCTGCGGGCGGACATTGCCAAGCAGCTGCCCAAAGA
 GGCTAAGCGGTTCTCCAACATCGACAAGAGCTGGGTCAAGATCATGACCA
 GAGCCCACGAGGTGCCCAGCGTGGTGCAGTGCTGTGTGGGCGACGAAAC
 ACTGGGACAGCTGCTGCCTCATCTGCTGGACCAGCTGGAAATCTGCCAGA
 AGTCCCTGACCGGCTACCTGGAAAAGAAACGGCTGTGTTTCCCCCGGTTT
 TTCTTCGTGTCCGACCCCGCCCTGCTGGAAATCTGGGCCAGGCCAGCGA
 CTCACACACAATTCAGGCCATCTGCTGAATGTGTTTCGATAACATCAAGA
 GCGTGAAGTTCCACGAGAAAATCTACGACCGGATCCTGAGCATCAGCTCC
 CAGGAAGGCGAGACAATCGAGCTGGACAAGCCTGTGATGGCCGAGGGAA
 ACGTGGAAAGTGTGGCTGAACAGCCTGCTGGAAGAGTCCCAGAGCAGCCT
 GCACCTCGTGATCAGACAGGCCGCTGCCAACATCCAGGAAACCGGCTTTC
 AGCTGACCGAGTTCCTGTCCAGCTTCCCAGCACAAGTGGGACTGCTGGGC
 ATCCAGATGATTTGGACCAGAGACTCCGAAGAGGCCCTGAGAAACGCCA
 AGTTCGATAAGAAAATTATGCAGAAAACAAATCAGGCATTTCTGGAACTG
 CTGAACACCCTGATCGACGTGACCACCCGGGACCTGAGCAGCACCAGAGA
 GAGTGAAGTACGAGACACTGATCACCATCCACGTGCACCAGCGGGACAT
 CTTTCGACGACCTGTGCCACATGCACATCAAGTCTCCCATGGATTTTCGAGT
 GGCTGAAGCAGTGCAGGTTCTACTTCAACGAGGACTCCGACAAGATGATG
 ATCCACATCACCAGTGTGGCCTTTATCTATCAGAATGAGTTCCTGGGCTGT
 ACCGATCGCCTCGTGATTACCCCCCTGACCGACCGGTGTTACATCACACT
 GGCCAGGCACTGGGCATGTCTATGGGAGGCGCACCCAGCAGGACCTGCC
 GGCACAGGCAAGACCGAAACCACCAAGGACATGGGACGCTGCCTGGGCA
 AATACGTGGTGGTGTTCAACTGCAGCGACCAGATGGATTTCCGGGGCCTG
 GGCCGGATCTTTAAGGGCCTGGCACAGAGCGGAAGCTGGGGCTGCTTCG
 ACGAGTTCAACAGAATCGACCTGCCCGTGCTGTCCGTGGCCGCACAGCAG
 ATCTCCATCATCCTGACATGCAAAAAGAGCACAAGAAGTCCTTCATCTT
 CACCGACGGCGACAATGTGACCATGAACCCCGAGTTTGGCCTGTTCTGTA
 CAATGAACCCTGGCTACGCCGGACGGCAGGAACTGCCCGAGAACCTGAA
 GATCAACTTTCGGAGTGTGGCTATGATGGTGCCCCGACCGGCAGATCATT
 TCAGAGTGAAACTGGCCTCCTGCGGCTTCATCGACAACGTGGTGTGGCT
 CGGAAGTTCTTCACACTGTACAAGCTGTGCGAAGAACAGCTGAGTAAACA
 GGTGCACTACGACTTCGGCCTGAGGAACATCCTGAGCGTGCTGAGAACTC
 TGGGAGCCGCTAAGCGGGCCAACCCCATGGATACCGAGAGCACAATCGT
 GATGCGGGTGCTGCGGGACATGAACCTGTCCAAGCTGATCGATGAGGAC
 GAGCCCCTGTTTCTGTCTCTGATCGAGGATCTGTTTCCCAACATTCTGCTG
 GATAAGGCCGGCTACCCCGAACTGGAAGCTGCTATCAGCAGACAGGTGG
 AAGAGGCTGGCCTGATCAACCACCCCCCTGGAAACTGAAAGTGATCCA
 GCTGTTTCGAGACACAGCGCGTGCGGCACGGCATGATGACACTGGGACCT
 AGCGGAGCCGGCAAGACCACCTGTATCCACACACTGATGCGGGCCATGA
 CCGATTGCGGCAAGCCCCACCGCGAGATGCGGGATGAAC
 CCCAAGGCCATTACCGCCCCTCAGATGTTTCGGCAGACTGGACGTGGCCAC

CAACGACTGGACCGACGGCATCTTCAGCACCCCTGTGGCGCAAGACCCTGC
 GGGCCAAGAAGGGCGAGCACATCTGGATCATCCTGGACGGCCCCGTGGA
 CGCCATCTGGATTGAGAACCTGAACAGCGTGCTGGACGACAACAAGACA
 CTGACCCTGGCCAACGGCGACCGGATCCCCATGGCCCCCAACTGCAAGAT
 CATCTTCGAGCCCCACAACATCGACAACGCCAGCCCTGCCACCGTGTCCA
 GAAACGGCATGGTGTTCATGAGCAGCAGCATCCTGGATTGGAGCCCTATC
 CTGGAAGGCTTCTGAAGAAGCGGAGCCCCAGGAAGCCGAGATCCTGA
 GACAGCTGTACACCGAGAGCTTCCCCGACCTGTACCGGTTCTGCATCCAG
 AATCTGGAGTACAAGATGGAAGTGCTGGAAGCCTTTGTGATCACCCAGAG
 CATCAACATGCTGCAGGGCCTGATCCCCCTGAAAGAACAGGGCGGAGAA
 GTGTCCCAGGCCACCTGGGCAGACTGTTCGTGTTTGCCCTGCTGTGGAG
 CGCTGGCGCCGCTCTGGAAGTGGATGGAAGGCGGAGACTGGAAGTGTGG
 CTGCGGAGCAGACCTACCGGCACCCTGGAAGTGCCTCCACCAGCTGGACC
 TGGCGACACCGCCTTCGATTACTACGTGGCCCCTGACGGCACCTGGACCC
 ACTGGAATACCCGGACCCAGGAATACCTGTACCCCAGCGACACCACCCCC
 GAGTACGGCTCTATCCTGGTGCCCAACGTGGACAACGTGCGGACCGACTT
 CCTGATCCAGACAATCGCCAAGCAGGGAAAGGCCGTGCTGCTGATCGGC
 GAGCAGGGCACAGCCAAGACCGTGATCATCAAGGGCTTTATGTCTAAGTA
 CGACCCCGAGTGCCACATGATCAAGAGCCTGAACTTCAGCTCCGCCACCA
 CCCCCTGATGTTCCAGCGGACCATCGAGAGCTATGTGGACAAGCGGATG
 GGCACCACCTACGGCCCTCCAGCCGGCAAGAAAATGACCGTGTTTCATCGA
 CGACGTGAACATGCCCATCATCAACGAGTGGGGCGACCAAGTGACCAAC
 GAGATCGTGCGGCAGCTGATGGAACAGAACGGCTTCTACAACCTGGAAA
 AGCCCGGCGAGTTCACCTCTATCGTGGACATCCAGTTTCTGGCCGCCATG
 ATCCACCCTGGCGGCGGAAGAAACGACATCCCCAGCGGCTGAAGCGGC
 AGTTCAGCATCTTCAACTGCACCCTGCCAGCGAGGCCAGCGTGGACAAG
 ATCTTTGGCGTGATCGGCGTGGGCCACTACTGCACCCAGAGAGGCTTCAG
 CGAGGAAGTGCGGGACAGCGTGACCAAGCTGGTGCCTCTGACAAGACGG
 CTGTGGCAGATGACCAAGATCAAGATGCTGCCACCCCCGCCAAGTTCCA
 CTACGTGTTCAACCTGCGGGACCTGAGCAGAGTGTGGCAGGGAATGCTGA
 ACACCACCAGCGAAGTGATCAAAGAGCCCAACGACCTGCTGAAGCTGTG
 GAAGCACGAGTGCAAGAGAGTGATCGCCGACCGGTTACCGTGTCTAGC
 GACGTGACATGGTTCGACAAGGCCCTGGTGTCCCTGGTGGAAAGAGGAATT
 CGGCGAAGAGAAGAACTGCTGGTGGACTGCGGCATCGATACTACTTC
 GTGGACTTCTGCGCGACGCCCTGAAGCCGCTGGCGAGACAAGTGAAG
 AGGCCGACGCCGAGACACCCAAGATCTACGAGCCCATCGAGTCCCTCAGC
 CATCTGAAAGAAAGGCTGAATATGTTCTGACAGCTGTATAACGAGTCCAT
 CCGGGGAGCCGGCATGGATATGGTGTCTTTGCCGACGCCATGGTGCACC
 TCGTGAAGATCAGCAGAGTGATCCGGACCCCCAGGGCAACGCTCTGCTC
 GTGGGAGTGGGAGGCTCTGGCAAGCAGAGCCTGACCAGACTGGCCAGCT
 TTATCGCCGGCTACGTGTCCTTCCAGATCACCTGACCCGGTCTTACAACA
 CCAGCAACCTGATGGAAGATCTGAAGGTGCTGTACCGGACAGCCGGCCA
 GCAGGGGAAGGGCATCACCTTCATCTTACCGACAATGAGATCAAGGAC
 GAGTCTTTCCTGGAGTATATGAACAATGTGCTGAGCAGCGGCGAGGTGTC
 CAACCTGTTCCGCCGGGACGAGATCGACGAGATTAACAGCGACCTGGCCT
 CCGTGATGAAGAAAGAATTCCCCCGGTGCCTGCCACAAACGAGAACCT
 GCACGACTACTTCATGTCCAGAGTGCGGCAGAATCTGCACATCGTGCTGT
 GCTTCAGCCCCGTGGGCGAGAAGTTCAGAAACCGGGCCCTGAAGTTCCCC
 GCCCTGATCAGCGGCTGCACCATCGACTGGTTCAGCCGGTGGCCTAAGGA
 TGCCCTGGTGGCCGTGTCCGAGCACTTCTGACCAGCTACGACATCGACT
 GCAGCCTGGAAATCAAGAAAGAGGTGGTGCAGTGCATGGGCAGCTTCCA

GGACGGCGTGGCCGAGAAATGCGTGGACTACTTCCAGCGGTTCCGGCGG
AGCACCCACGTGACCCCTAAGAGCTACCTGAGCTTCATCCAGGGCTACAA
GTTTCATCTACGGCGAGAAGCACGTGGAAGTGCGCACACTGGCCAACCGG
ATGAACACCGGCCTGGAAAACTGAAAGAGGCCTCCGAGAGCGTGGCCG
CCCTGAGCAAAGAAGCTGGAAGCCAAAGAAAAAGAAGTGCAGGTGGCCAA
CGATAAGGCCGACATGGTGCTGAAAGAAGTGACCATGAAGGCCCAGGCC
GCCGAGAAAGTGAAAGCCGAGGTGCAGAAAGTGAAGGACCGGGCCAG
GCCATCGTGGACTCCATCAGCAAGGACAAGGCCATTGCCGAGGAAAAGC
TGGAAGCAGCCAAGCCCGCCCTGGAAGAGGCAGAAGCTGCTCTGCAGAC
CATCCGGCCCTCCGATATTGCCACAGTGCAGGACCCTGGGAAGGCCCCCTC
ACCTGATCATGCGGATCATGGACTGTGTGCTGCTGCTGTTCCAGAGAAAG
GTGTCCGCCGTGAAGATCGACCTGGAAAAATCCTGCACCATGCCTAGCTG
GCAGGAATCCCTGAAGCTGATGACCGCCGGCAACTTCCTGCAGAACCTGC
AGCAGTTCCCAAGGACACCATCAATGAGGAAGTGATCGAGTTCCTGAGC
CCCTACTTCGAGATGCCCGACTACAATATCGAAACCGCCAAACGCGTGTG
CGGCAACGTGGCCGACTGTGCTCTTGACCAAGGCTATGGCTAGCTTCT
TTAGCATTAAACAAAGAGGTGCTGCCTCTGAAGGCCAACCTGGTGGTGCAG
GAAAACCGGCATCTGCTGGCCATGCAGGACCTGCAGAAAGCCAGGCCG
AGCTGGACGATAAGCAGGCTGAGCTGGATGTGGTGCAGGCCGAGTACGA
GCAGGCCATGACCGAGAAGCAGACCCTGCTGGAAGATGCAGAGCGGTGC
AGACACAAGATGCAGACCGCCAGCACCCCTGATCTCTGGACTGGCCGGCG
AAAAAGAGCGGTGGACCGAGCAGTCCCAGGAATTCGCCGCCAGACCAA
GCGGCTCGTGGGAGATGTGCTGCTGGCCACCGCCTTTCTGAGCTACAGCG
GCCCCTTCAATCAGGAATTCAGGGACCTGCTGCTGAACGACTGGCGGAAA
GAGATGAAGGCCAGAAAGATCCCCTTCGGCAAGAATCTGAACCTGAGCG
AGATGCTGATCGACGCCCCACCATCTCCGAGTGGAAATCTGCAGGGACTG
CCCAACGATGACCTGTCCATCCAGAACGGAATCATCGTGACCAAAGCCTC
CAGATACCCCCTGCTGATTGACCCCCAGACACAGGGCAAGATTTGGATCA
AGAACAAGAGAGGCCGGAACGAGCTGCAGATCACCAGCCTGAACCACAA
GTACTTCCGGAACCACCTGGAAGATAGCCTGAGCCTGGGCAGGCCACTGC
TGATCGAGGATGTGGGCGAGGAACTGGACCCAGCCCTGGATAACGTGCT
GGAACGGAAC TTCATCAAGACCGGCTCCACCTTCAAAGTGAAAGTGGGC
GACAAAGAAGTGGACGTGCTGGATGGCTTCCGGCTGTACATCACCACCAA
GCTGCCTAACCCCGCCTACACCCCTGAGATCAGCGCCCGGACCAGCATCA
TCGACTTCACCGTGACAATGAAGGGACTGGAAGATCAGCTGCTGGGACG
CGTGATCCTGACAGAGAAGCAGGAACTGGAAAAAGAACGGACCCATCTG
ATGGAAGATGTGACCGCCAACAAGCGGCGGATGAAGGAACTGGAAGATA
ACCTGCTGTACAGGCTGACCAGCACCCAGGGCAGTCTGGTGGAAAGATGA
GAGCCTGATCGTGGTGTGCTGTCCAACACCAAGCGGACCGCAGAGGAAGTG
ACCCAGAAGCTGGAAATCAGCGCCGAGACAGAGGTGCAGATCAACAGCG
CCAGAGAAGAGTACCGGCCTGTGGCCACCCGGGGATCCATCCTGTACTTT
CTGATCACCGAGATGCGGCTCGTGAACGAGATGTACCAGACCAGCCTGCG
GCAGTTCCTGGGCCTGTTTCGATCTGTCCCTGGCCAGAAGCGTGAAGTCCC
CCATCACAGCAAGAGAATCGCCAACATCATCGAGCACATGACCTACGA
GGTGTACAAATACGCCGCCAGAGGCCTGTACGAGGAACACAAGTTTCTGT
TCACACTGCTGCTGACCCCTGAAGATCGATATCCAGCGGAACAGAGTGAAG
CACGAAGAGTTTCTGACACTGATCAAGGGGGGAGCCTCCCTGGACCTGAA
GGCCTGTCTCCCAAGCCCAGCAAGTGGATCCTGGACATCACCTGGCTGA
ATCTGGTGGAACTGAGCAAGCTGAGACAGTTCTCCGATGTGCTGGACCAG
ATCAGCCGCAACGAGAAGATGTGGAAGATTTGGTTTGACAAAGAGAACC
CCGAGGAAGAACCCTGCCTAACGCTACGATAAGAGCCTGGACTGCTTC

CGGCGGCTGCTGCTGATTAGAAGCTGGTGTCCCACCGGACAATCGCCCA
 GGCCCGCAAGTACATCGTGGATAGCATGGGAGAGAAGTACGCCGAGGGC
 GTGATCCTGGACCTGGAAAAGACCTGGGAGGAAAGCGACCCCAGAACCC
 CCCTGATCTGCCTGCTGAGCATGGGCTCCGACCCACCGACAGCATTATC
 GCCCTGGGCAAGAGACTGAAGATTGAGACAAGATACGTGTCCATGGGCC
 AGGGCCAGGAAGTGCACGCTAGAAAGCTGCTGCAGCAGACTATGGCCAA
 TGGCGGCTGGGCCCTGCTGCAGAATTGTCACCTGGGGCTGGACTTCATGG
 ACGAACTGATGGACATCATCATTGAGACAGAGCTGGTGCACGACGCCTTC
 AGACTGTGGATGACCACCGAGGCCATAAAGCAGTTTCCCATTACCCTGCT
 GCAGATGAGCATCAAGTTCGCCAACGACCCCCCTCAGGGACTGAGAGCC
 GGCCTGAAGAGAACCTACTCCGGCGTGTACAGGATCTGCTGGACGTGTC
 CTCTGGCAGCCAGTGGAAAGCCTATGCTGTACGCCGTGGCATTCTGCACA
 GCACCGTGCAGGAACGGCGGAAGTTTGGCGCCCTGGGATGGAACATCCC
 CTACGAGTTTAACCAGGCCGACTTCAACGCCACTGTGCAGTTTATCCAGA
 ACCATCTGGACGACATGGACGTGAAGAAAGGGGTGTCTTGGACAACCAT
 CCGGTACATGATCGGAGAGATCCAGTACGGCGGCAGAGTGACCGACGAC
 TACGACAAGAGGCTGCTGAATACCTTCGCCAAAGTGTGGTTCTCCGAGAA
 CATGTTTGGCCCCGACTTCAGCTTTTACCAGGGCTATAACATCCCCAAGTG
 CTCCACCGTGGATAACTACCTGCAGTACATCCAGAGCCTGCCCGCCTACG
 ACAGCCCTGAGGTGTTTCGGACTGCACCCCAACGCCGATATCACCTACCAG
 AGCAAACCTGGCCAAGGATGTGCTGGATAACCATCCTGGGCATCCAGCCAA
 GGATAACAGTGGCGGAGGCGACGAAACCCGGGAAGCAGTGGTGGCTAGA
 CTGGCCGACGACATGCTGGAAAAGCTGCCCCCCGACTACGTGCCCTTTGA
 AGTGAAAGAACGCCTGCAGAAGATGGGCCCCTTCCAGCCTATGAACATCT
 TCCTGAGGCAGGAAATCGACCGGATGCAGCGGGTGTGTCTCTCGTGC GG
 AGCACACTGACCGAGCTGAAACTGGCTATCGACGGCACCATCATCATGAG
 CGAGAATCTGCGGGATGCACTGGACTGCATGTTCGACGCCAGAATCCCCG
 CATGGTGGAAAAGGCCAGCTGGATCAGCTCTACCCTGGGCTTCTGGTTC
 ACCGAACTGATCGAGAGAAACAGCCAGTTTACCAGCTGGGTGTTCAACG
 GCAGACCTCACTGCTTCTGGATGACCGGCTTCTTCAATCCACAAGGCTTTC
 TGACAGCAATGCGCCAGGAAATCACCAGAGCCAACAAGGGCTGGGCTCT
 GGACAATATGGTGCTGTGTAACGAAGTGAAGTAAAGTGGATGAAGGACGAC
 ATCAGCGCCCCTCCCACAGAGGGCGTGTACGTGTACGGCCTGTACCTGGA
 AGGCGCCGGATGGGACAAGAGAAACATGAAGCTGATCGAGAGCAAGCCC
 AAGGTGCTGTTTCGAGCTGATGCCCGTGTACAGGATCTATGCCGAGAACAA
 CACCCTGAGGGACCCCCGTTCTACAGCTGCCCATCTACAAGAAACCCG
 TGCGCACCGACCTGAACTATATCGCCGCCGTGGACCTGAGGACAGCCCAG
 ACACCTGAGCATTGGGTGCTGAGAGGCGTGGCACTGCTGTGCGACGTGAA
 GTGA (SEQ ID NO: 18)

Nucleic Acid Constructs, Vectors, and Engineered Polyribonucleotides

[0099] The present disclosure provides nucleic acid molecules, such as polynucleotides, which encode one or more polypeptides of interest. The term nucleic acid includes any compound and/or substance that comprise a polymer of nucleotides. Nucleotide polymers that contain greater than 50% of ribose bases or ribonucleotide analogues are referred to as polyribonucleotides. Nucleotide polymers may use altered nucleotide usage that encode a protein or functional fragment thereof, such as DNAI1 or DNAH5. The sequence of the engineered

polynucleotides can be derived from, for example, DNA, RNA, mRNA transcripts, genomic DNA, mitochondrial DNA, mitochondrial RNA, or another suitable nucleic acid that comprises the genetic information of a gene of interest. The nucleic acid constructs, vectors, engineered polyribonucleotides, or compositions can be derived from nucleic acids carrying mutated genes and polymorphisms.

[00100] In addition to the four canonical ribonucleotides, namely, adenosine, guanosine, cytidine and uridine, several cellular RNAs also contain a number of structurally diverse ribonucleotides. About a hundred structurally different nucleotides or nucleotide analogues have been identified in transfer RNAs (tRNAs), ribosomal RNAs (rRNAs), messenger RNAs (mRNAs) and small nuclear RNAs (snRNAs). In tRNAs, some nucleotides can be important determinants of the specificity and efficiency of aminoacylation and codon recognition. Such structurally diverse ribonucleotides can be a modified ribonucleotide or a nucleotide analogue. In some cases, a polynucleotide of the disclosure is engineered to comprise a ribonucleotide analogue.

[00101] In some cases, a nucleic acid construct, a vector, or a polynucleotide is engineered to contain the four classical ribonucleotides and can be modified post-transcriptionally, after being administered to a subject. For instance, in some cases the disclosure provides a composition, vector, or a nucleic acid construct comprising a nucleic acid construct encoding dynein axonemal intermediate chain 1, wherein fewer than 30% of the nucleic acids encoding dynein axonemal intermediate chain 1 are nucleotide analogues. In other cases, fewer than 27.5%, fewer than 25%, fewer than 22.5%, fewer than 20%, fewer than 17.5%, fewer than 15%, fewer than 12.5%, fewer than 10%, fewer than 7.5%, fewer than 5%, or fewer than 2.5% of the nucleotides encoding dynein axonemal intermediate chain 1 are nucleotide analogues.

[00102] Exemplary nucleic acids that can form a polynucleotide of the disclosure include, but are not limited to, ribonucleic acids (RNAs), deoxyribonucleic acids (DNAs), or hybrids thereof. Exemplary modified nucleotides that can form at least a fraction of a polynucleotide of the disclosure include, but are not limited to, pseudouridine (Ψ) and 1-methylpseudouridine ($m^1\Psi$).

[00103] A chemical modification can be located on one or more nucleoside(s) or the backbone of the nucleic acid molecule. They can be located on both a nucleoside and a backbone linkage. A modification can be engineered into a polynucleotide *in vitro*. Modified ribonucleotides and nucleic acid analogues can also be introduced post-transcriptionally by covalent modification of the classical ribonucleotides.

[00104] A nucleic acid construct, a vector, or an engineered polyribonucleotide of the disclosure can comprise purine and pyrimidine analogues. In some cases, a polyribonucleotide of the disclosure comprises a modified pyrimidine, such as a modified uridine. In some cases a uridine analogue is selected from pseudouridine (Ψ), 1-methylpseudouridine ($m^1\Psi$), 2-thiouridine (s^2U), 5-methyluridine (m^5U), 5-methoxyuridine (mo^5U), 4-thiouridine (s^4U), 5-bromouridine (Br^5U), 2'-O-methyluridine ($U2'm$), 2'-amino-2'-deoxyuridine ($U2'NH_2$), 2'-azido-2'-deoxyuridine ($U2'N_3$), and 2'-fluoro-2'-deoxyuridine ($U2'F$).

[00105] In some instances, the nucleic acid construct(s), vector(s), engineered polyribonucleotide(s), or composition(s) encodes dynein axonemal intermediate chain 1 protein or a variant thereof at a level that is increased by a factor of at least about 1.5 as compared to levels within cells exposed to a composition comprising a nucleic acid construct that does not include the codons encoding dynein axonemal intermediate chain 1 protein or a variant thereof. In some cases, the factor is at least about 1.1, at least about 1.2, at least about 1.3, at least about 1.4, at least about 1.5, at least about 2, at least about 3, at least about 4, at least about 5, at least about 10, at least about 20, at least about 30, at least about 40, at least about 50, at least about 60, at least about 70, at least about 80, at least about 90, or at least about 100.

[00106] A polyribonucleotide can have the same or a mixture of different nucleotide analogues or modified nucleotides. The nucleotide analogues or modified nucleotides can have structural changes that are naturally or not naturally occurring in messenger RNA. A mixture of various analogues or modified nucleotides can be used. For example one or more analogues within a polynucleotide can have natural modifications, while another part has modifications that are not naturally found in mRNA. Additionally, some analogues or modified ribonucleotides can have a base modification, while other modified ribonucleotides have a sugar modification. In the same way, it is possible that all modifications are base modifications or all modifications are sugar modifications or any suitable mixture thereof.

[00107] A nucleotide analogue or modified nucleotide can be selected from the group comprising pyridin-4-one ribonucleoside, 5-aza-uridine, 2-thio-5-aza-uridine, 2-thiouridine, 4-thio-pseudouridine, 2-thio-pseudouridine, 5-hydroxyuridine, 3-methyluridine, 5-carboxymethyluridine, 1-carboxymethyl-pseudouridine, 5-propynyl-uridine, 1-propynyl-pseudouridine, 5-taurinomethyluridine, 1-taurinomethyl-pseudouridine, 5-taurinomethyl-2-thio-uridine, 1-taurinomethyl-4-thio-uridine, 5-methyl-uridine, 1-methyl-pseudouridine, 4-thio-1-methyl-pseudouridine, 2-thio-1-methyl-pseudouridine, 1-methyl-1-deaza-pseudouridine, 2-thio-1-methyl-1-deaza-pseudouridine, dihydrouridine, dihydropseudouridine, 2-thio-dihydrouridine, 2-thio-dihydropseudouridine, 2-methoxyuridine, 2-methoxy-4-thio-uridine, 4-methoxy-pseudouridine,

4-methoxy-2-thio-pseudouridine, 5-aza-cytidine, pseudoisocytidine, 3-methyl-cytidine, N4-acetylcytidine, 5-formylcytidine, N4-methylcytidine, 5-hydroxymethylcytidine, 1-methyl-pseudoisocytidine, pyrrolo-cytidine, pyrrolo-pseudoisocytidine, 2-thio-cytidine, 2-thio-5-methyl-cytidine, 4-thio-pseudoisocytidine, 4-thio-1-methyl-pseudoisocytidine, 4-thio-1-methyl-1-deaza-pseudoisocytidine, 1-methyl-1-deaza-pseudoisocytidine, zebularine, 5-aza-zebularine, 5-methyl-zebularine, 5-aza-2-thio-zebularine, 2-thio-zebularine, 2-methoxy-cytidine, 2-methoxy-5-methyl-cytidine, 4-methoxy-pseudoisocytidine, 4-methoxy-1-methyl-pseudoisocytidine, 2-aminopurine, 2, 6-diaminopurine, 7-deaza-adenine, 7-deaza-8-aza-adenine, 7-deaza-2-aminopurine, 7-deaza-8-aza-2-aminopurine, 7-deaza-2,6-diaminopurine, 7-deaza-8-aza-2,6-diaminopurine, 1-methyladenosine, N6-methyladenosine, N6-isopentenyladenosine, N6-(cis-hydroxyisopentenyl)adenosine, 2-methylthio-N6-(cis-hydroxyisopentenyl) adenosine, N6-glycinylicarbamoyladenosine, N6-threonylicarbamoyladenosine, 2-methylthio-N6-threonyl carbamoyladenosine, N6,N6-dimethyladenosine, 7-methyladenine, 2-methylthio-adenine, 2-methoxy-adenine, inosine, 1-methyl-inosine, wyosine, wybutosine, 7-deaza-guanosine, 7-deaza-8-aza-guanosine, 6-thio-guanosine, 6-thio-7-deaza-guanosine, 6-thio-7-deaza-8-aza-guanosine, 7-methyl-guanosine, 6-thio-7-methyl-guanosine, 7-methylinosine, 6-methoxy-guanosine, 1-methyl-guanosine, N2-methyl-guanosine, N2,N2-dimethyl-guanosine, 8-oxo-guanosine, 7-methyl-8-oxo-guanosine, 1-methyl-6-thio-guanosine, N2-methyl-6-thio-guanosine, and N2,N2-dimethyl-6-thio-guanosine.

[00108] In some cases, at least about 5 % of the nucleic acid construct(s), a vector(s), engineered polyribonucleotide(s), or compositions includes non-naturally occurring (e.g., modified, analogues, or engineered) uridine, adenosine, guanine, or cytosine, such as the nucleotides described herein. In some cases, 100% of the modified nucleotides in the composition are either 1-methylpseudouridine or pseudouridine. In some cases, at least about 10 %, 15 %, 20 %, 25 %, 30 %, 35 %, 40 %, 45 %, 50 %, 55 %, 60 %, 65 %, 70 %, 75 %, 80 %, 85 %, 90 %, 95 % of the nucleic acid construct(s), a vector(s), engineered polyribonucleotide(s), or compositions includes non-naturally occurring uracil, adenine, guanine, or cytosine. In some cases, at most about 99 %, 95 %, 90 %, 85 %, 80 %, 75 %, 70 %, 65 %, 60 %, 55 %, 50 %, 45 %, 40 %, 35 %, 30 %, 25 %, 20 %, 15 %, 10 %, 5 %, 1 %, of the nucleic acid construct(s), a vector(s), engineered polyribonucleotide(s), or compositions includes non-naturally occurring uracil, adenine, guanine, or cytosine.

[00109] A nucleic acid construct(s), a vector(s), or an engineered polyribonucleotide(s) of the disclosure can comprise one or more promoter sequences and any associated regulatory sequences. A promoter sequence and/or an associated regulatory sequence can comprise any

number of modified or unmodified nucleotides, and any number of nucleic acid analogues. Promoter sequences and/or any associated regulatory sequences can comprise, for example, at least 2 bases or base pairs, 3 bases or base pairs, 4 bases or base pairs, 5 bases or base pairs, 6 bases or base pairs, 7 bases or base pairs, 8 bases or base pairs, 9 bases or base pairs, 10 bases or base pairs, 11 bases or base pairs, 12 bases or base pairs, 13 bases or base pairs, 14 bases or base pairs, 15 bases or base pairs, 16 bases or base pairs, 17 bases or base pairs, 18 bases or base pairs, 19 bases or base pairs, 20 bases or base pairs, 21 bases or base pairs, 22 bases or base pairs, 23 bases or base pairs, 24 bases or base pairs, 25 bases or base pairs, 26 bases or base pairs, 27 bases or base pairs, 28 bases or base pairs, 29 bases or base pairs, 30 bases or base pairs, 35 bases or base pairs, 40 bases or base pairs, 50 bases or base pairs, 75 bases or base pairs, 100 bases or base pairs, 150 bases or base pairs, 200 bases or base pairs, 300 bases or base pairs, 400 bases or base pairs, 500 bases or base pairs, 600 bases or base pairs, 700 bases or base pairs, 800 bases or base pairs, 900 bases or base pairs, 1000 bases or base pairs, 2000 bases or base pairs, 3000 bases or base pairs, 4000 bases or base pairs, 5000 bases or base pairs, at least 10000 bases or base pairs or more. A promoter sequence and/or an associated regulatory sequence can comprise any number of modified or unmodified nucleotides, for example, at most 10000 bases or base pairs, 5000 bases or base pairs, 4000 bases or base pairs, 3000 bases or base pairs, 2000 bases or base pairs, 1000 bases or base pairs, 900 bases or base pairs, 800 bases or base pairs, 700 bases or base pairs, 600 bases or base pairs, 500 bases or base pairs, 400 bases or base pairs, 300 bases or base pairs, 200 bases or base pairs, 100 bases or base pairs, 75 bases or base pairs, 50 bases or base pairs, 40 bases or base pairs, 35 bases or base pairs, 30 bases or base pairs, 29 bases or base pairs, 28 bases or base pairs, 27 bases or base pairs, 26 bases or base pairs, 25 bases or base pairs, 24 bases or base pairs, 23 bases or base pairs, 22 bases or base pairs, 21 bases or base pairs, 20 bases or base pairs, 19 bases or base pairs, 18 bases or base pairs, 17 bases or base pairs, 16 bases or base pairs, 15 bases or base pairs, 14 bases or base pairs, 13 bases or base pairs, 12 bases or base pairs, 11 bases or base pairs, 10 bases or base pairs, 9 bases or base pairs, 8 bases or base pairs, 7 bases or base pairs, 6 bases or base pairs, 5 bases or base pairs, 4 bases or base pairs, 3 bases or base pairs or 2 bases or base pairs.

[00110] In some cases, less than all of the nucleotides in the promoter sequence or associated regulatory region are nucleotide analogues or modified nucleotides. For instance, in some cases, less than or equal to 99%, 95%, 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, or 5% of the nucleotides in a promoter or associated regulatory region. In some cases, all of the nucleotides in a promoter or associated regulatory region are nucleic acid analogues or modified nucleotides.

[00111] A nucleic acid construct(s), a vector(s), an engineered polyribonucleotide(s), or compositions of the disclosure can comprise an engineered 5' cap structure, or a 5'-cap can be added to a polyribonucleotide intracellularly. The 5' cap structure of an mRNA can be involved in binding to the mRNA Cap Binding Protein (CBP), which is responsible for mRNA stability in the cell and translation competency through the association of CBP with poly(A) binding protein to form the mature pseudo-circular mRNA species. The 5' cap structure can also be involved in nuclear export, increases in mRNA stability, and in assisting the removal of 5' proximal introns during mRNA splicing.

[00112] A nucleic acid construct(s), a vector(s), or an engineered polyribonucleotide(s) can be 5'-end capped generating a 5'-GpppN-3'-triphosphate linkage between a terminal guanosine cap residue and the 5'-terminal transcribed sense nucleotide of the mRNA molecule. The cap-structure can comprise a modified or unmodified 7-methylguanosine linked to the first nucleotide via a 5'-5' triphosphate bridge. This 5'-guanylate cap can then be methylated to generate an N7-methyl-guanylate residue (Cap-0 structure). The ribose sugars of the terminal and/or antiterminal transcribed nucleotides of the 5' end of the mRNA may optionally also be 2'-O-methylated (Cap-1 structure). 5'-decapping through hydrolysis and cleavage of the guanylate cap structure may target a nucleic acid molecule, such as an mRNA molecule, for degradation.

[00113] In some cases, a cap can comprise further modifications, including the methylation of the 2' hydroxy-groups of the first 2 ribose sugars of the 5' end of the mRNA. For instance, an eukaryotic cap-1 has a methylated 2'-hydroxy group on the first ribose sugar, while a cap-2 has methylated 2'-hydroxy groups on the first two ribose sugars. The 5' cap can be chemically similar to the 3' end of an RNA molecule (the 5' carbon of the cap ribose is bonded, and the free 3'-hydroxyls on both 5'- and 3'- ends of the capped transcripts. Such double modification can provide significant resistance to 5' exonucleases. Non-limiting examples of 5' cap structures that can be used with an engineered polyribonucleotide include, but are not limited to, m⁷G(5')ppp(5')N (Cap-0), m⁷G(5')ppp(5')N1mpNp (Cap-1), and m⁷G(5')ppp(5')N1mpN2mp (Cap-2).

[00114] Modifications to the modified mRNA of the present disclosure may generate a non-hydrolyzable cap structure preventing decapping and thus increasing mRNA half-life while facilitating efficient translation. Because cap structure hydrolysis requires cleavage of 5'-ppp-5' triphosphate linkages, modified nucleotides may be used during the capping reaction. For example, a Vaccinia Capping Enzyme from New England Biolabs (Ipswich, MA) may be used with guanosine α -thiophosphate nucleotides according to the manufacturer's instructions to create a phosphorothioate linkage in the 5'-ppp-5' cap. Additional modified guanosine

nucleotides may be used such as α -methyl-phosphonate and seleno-phosphate nucleotides. Additional modifications include, but are not limited to, 2'-O-methylation of the ribose sugars of 5'-terminal and/or 5'-antiterminal nucleotides of the mRNA on the 2'-hydroxyl group of the sugar ring. Multiple distinct 5'-cap structures can be used to generate the 5'-cap of a polyribonucleotide.

[00115] The modified mRNA may be capped post-transcriptionally. According to the present disclosure, 5' terminal caps may include endogenous caps or cap analogues. According to the present disclosure, a 5' terminal cap may comprise a guanine analogue. Useful guanine analogues include, but are not limited to, inosine, N1-methyl-guanosine, 2' fluoro-guanosine, 7-deaza-guanosine, 8-oxo-guanosine, 2-amino-guanosine, LNA-guanosine, and 2-azido-guanosine.

[00116] Further, a nucleic acid construct(s), a vector(s), or an engineered polyribonucleotide(s) can contain one or more internal ribosome entry site(s) (IRES). IRES sequences can initiate protein synthesis in absence of the 5' cap structure. An IRES sequence can also be the sole ribosome binding site, or it can serve as one of multiple ribosome binding sites of an mRNA. Engineered polyribonucleotides containing more than one functional ribosome binding site can encode several peptides or polypeptides that are translated by the ribosomes ("polycistronic or multicistronic polynucleotides"). An engineered polynucleotide described here can comprise at least 1 IRES sequence, two IRES sequences, three IRES sequences, four IRES sequences, five IRES sequences, six IRES sequences, seven IRES sequences, eight IRES sequences, nine IRES sequences, ten IRES sequences, or another suitable number are present in an engineered polyribonucleotide. Examples of IRES sequences that can be used according to the present disclosure include without limitation, those from tobacco etch virus (TEV), picornaviruses (e.g., FMDV), pest viruses (CFFV), polio viruses (PV), encephalomyocarditis viruses (EMCV), foot-and-mouth disease viruses (FMDV), hepatitis C viruses (HCV), classical swine fever viruses (CSFV), murine leukemia virus (MLV), simian immune deficiency viruses (SIV) or cricket paralysis viruses (CrPV). An IRES sequence can be derived, for example, from commercially available vectors such as the IRES sequences available from ClontechTM, GeneCopoeiaTM, or Sigma-AldrichTM. IRES sequences can be, for example, at least 150 bases or base pairs, 200 bases or base pairs, 300 bases or base pairs, 400 bases or base pairs, 500 bases or base pairs, 600 bases or base pairs, 700 bases or base pairs, 800 bases or base pairs, 900 bases or base pairs, 1000 bases or base pairs, 2000 bases or base pairs, 3000 bases or base pairs, 4000 bases or base pairs, 5000 bases or base pairs, or 10000 bases or base pairs. IRES sequences can at most 10000 bases or base pairs, 5000 bases or base pairs, 4000 bases or base pairs, 3000 bases or base pairs, 2000 bases or base pairs, 1000 bases or base pairs, 900 bases or base pairs, 800

bases or base pairs, 700 bases or base pairs, 600 bases or base pairs, 500 bases or base pairs, 400 bases or base pairs, 300 bases or base pairs, 200 bases or base pairs, 100 bases or base pairs, 50 bases or base pairs, or 10 bases or base pairs.

[00117] A nucleic acid construct(s), a vector(s), or an engineered polyribonucleotide(s) of the disclosure can comprise one or more untranslated regions. An untranslated region can comprise any number of modified or unmodified nucleotides. Untranslated regions (UTRs) of a gene are transcribed but not translated into a polypeptide. In some cases, an untranslated sequence can increase the stability of the nucleic acid molecule and the efficiency of translation. The regulatory features of a UTR can be incorporated into the modified mRNA molecules of the present disclosure, for instance, to increase the stability of the molecule. The specific features can also be incorporated to ensure controlled down-regulation of the transcript in case they are misdirected to undesired organs sites. Some 5' UTRs play roles in translation initiation. A 5' UTR can comprise a Kozak sequence which is involved in the process by which the ribosome initiates translation of many genes. Kozak sequences can have the consensus GCC(R)CCAUGG, where R is a purine (adenine or guanine) that is located three bases upstream of the start codon (AUG). 5' UTRs may form secondary structures which are involved in binding of translation elongation factor. In some cases, one can increase the stability and protein production of the engineered polynucleotide molecules of the disclosure, by engineering the features typically found in abundantly expressed genes of specific target organs. For example, introduction of 5' UTR of liver-expressed mRNA, such as albumin, serum amyloid A, Apolipoprotein A/B/E, transferrin, alpha fetoprotein, erythropoietin, or Factor VIII, can be used to increase expression of an engineered polynucleotide in a liver. Likewise, use of 5' UTR from muscle proteins (MyoD, Myosin, Myoglobin, Myogenin, Herculin), for endothelial cells (Tie-1, CD36), for myeloid cells (C/EBP, AML1, G-CSF, GM-CSF, CD11b, MSR, Fr-1, i-NOS), for leukocytes (CD45, CD18), for adipose tissue (CD36, GLUT4, ACRP30, adiponectin) and for lung epithelial cells (SP-A/B/C/D) can be used to increase expression of an engineered polynucleotide in a desired cell or tissue.

[00118] Other non-UTR sequences can be incorporated into the 5' (or 3' UTR) UTRs of the polyribonucleotides of the present disclosure. The 5' and/or 3' UTRs can provide stability and/or translation efficiency of polyribonucleotides. For example, introns or portions of intron sequences can be incorporated into the flanking regions of an engineered polyribonucleotide. Incorporation of intronic sequences can also increase the rate of translation of the polyribonucleotide.

[00119] 3' UTRs may have stretches of Adenosines and Uridines embedded therein. These AU rich signatures are particularly prevalent in genes with high rates of turnover. Based on their sequence features and functional properties, the AU rich elements (AREs) can be separated into classes: Class I AREs contain several dispersed copies of an AUUUA motif within U-rich regions. C-Myc and MyoD contain class I AREs. Class II AREs possess two or more overlapping UUAUUUA(U/A)(U/A) nonamers. Molecules containing this type of AREs include GM-CSF and TNF- α . Class III AREs are less well defined. These U rich regions do not contain an AUUUA motif c-Jun and Myogenin are two well-studied examples of this class. Proteins binding to the AREs may destabilize the messenger, whereas members of the ELAV family, such as HuR, may increase the stability of mRNA. HuR may bind to AREs of all the three classes. Engineering the HuR specific binding sites into the 3' UTR of nucleic acid molecules can lead to HuR binding and thus, stabilization of the message *in vivo*.

[00120] Engineering of 3' UTR AU rich elements (AREs) can be used to modulate the stability of an engineered polyribonucleotide. One or more copies of an ARE can be engineered into a polyribonucleotide to modulate the stability of a polyribonucleotide. AREs can be identified, removed or mutated to increase the intracellular stability and thus increase translation and production of the resultant protein. Transfection experiments can be conducted in relevant cell lines, using engineered polyribonucleotides and protein production can be assayed at various time points post-transfection. For example, cells can be transfected with different ARE-engineering molecules and by using an ELISA kit to the relevant protein and assaying protein produced at 6 hours, 12 hours, 24 hours, 48 hours, and 7 days post-transfection.

[00121] An untranslated region can comprise any number of nucleotides. An untranslated region can comprise a length of about 1 to about 10 bases or base pairs, about 10 to about 20 bases or base pairs, about 20 to about 50 bases or base pairs, about 50 to about 100 bases or base pairs, about 100 to about 500 bases or base pairs, about 500 to about 1000 bases or base pairs, about 1000 to about 2000 bases or base pairs, about 2000 to about 3000 bases or base pairs, about 3000 to about 4000 bases or base pairs, about 4000 to about 5000 bases or base pairs, about 5000 to about 6000 bases or base pairs, about 6000 to about 7000 bases or base pairs, about 7000 to about 8000 bases or base pairs, about 8000 to about 9000 bases or base pairs, or about 9000 to about 10000 bases or base pairs in length. An untranslated region can comprise a length of for example, at least 1 base or base pair, 2 bases or base pairs, 3 bases or base pairs, 4 bases or base pairs, 5 bases or base pairs, 6 bases or base pairs, 7 bases or base pairs, 8 bases or base pairs, 9 bases or base pairs, 10 bases or base pairs, 20 bases or base pairs, 30 bases or base pairs, 40 bases or base pairs, 50 bases or base pairs, 60 bases or base pairs, 70 bases or base

pairs, 80 bases or base pairs, 90 bases or base pairs, 100 bases or base pairs, 200 bases or base pairs, 300 bases or base pairs, 400 bases or base pairs, 500 bases or base pairs, 600 bases or base pairs, 700 bases or base pairs, 800 bases or base pairs, 900 bases or base pairs, 1000 bases or base pairs, 2000 bases or base pairs, 3000 bases or base pairs, 4000 bases or base pairs, 5000 bases or base pairs, 6000 bases or base pairs, 7000 bases or base pairs, 8000 bases or base pairs, 9000 bases or base pairs, or 10000 bases or base pairs in length.

[00122] An engineered polyribonucleotide of the disclosure can comprise one or more introns. An intron can comprise any number of modified or unmodified nucleotides. An intron can comprise, for example, at least 1 base or base pair, 50 bases or base pairs, 100 bases or base pairs, 150 bases or base pairs, 200 bases or base pairs, 300 bases or base pairs, 400 bases or base pairs, 500 bases or base pairs, 600 bases or base pairs, 700 bases or base pairs, 800 bases or base pairs, 900 bases or base pairs, 1000 bases or base pairs, 2000 bases or base pairs, 3000 bases or base pairs, 4000 bases or base pairs, or 5000 bases or base pairs. In some cases, an intron can comprise, for example, at most 10000 bases or base pairs, 5000 bases or base pairs, 4000 bases or base pairs, 3000 bases or base pairs, 2000 bases or base pairs, 1000 bases or base pairs, 900 bases or base pairs, 800 bases or base pairs, 700 bases or base pairs, 600 bases or base pairs, 500 bases or base pairs, 400 bases or base pairs, 300 bases or base pairs, 200 bases or base pairs, or 100 bases or base pairs.

[00123] In some cases, a percentage of the nucleotides in an intron are modified. For instance, in some cases, fewer than 99%, 95%, 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, 5% or 1% of the nucleotides in an intron are modified. In some cases, all of the nucleotides in an intron are modified.

[00124] An engineered polyribonucleotide of the disclosure can comprise a polyA sequence. A polyA sequence (e.g., polyA tail) can comprise any number of nucleotides. A polyA sequence can comprise a length of about 1 to about 10 bases or base pairs, about 10 to about 20 bases or base pairs, about 20 to about 50 bases or base pairs, about 50 to about 100 bases or base pairs, about 100 to about 500 bases or base pairs, about 500 to about 1000 bases or base pairs, about 1000 to about 2000 bases or base pairs, about 2000 to about 3000 bases or base pairs, about 3000 to about 4000 bases or base pairs, about 4000 to about 5000 bases or base pairs, about 5000 to about 6000 bases or base pairs, about 6000 to about 7000 bases or base pairs, about 7000 to about 8000 bases or base pairs, about 8000 to about 9000 bases or base pairs, or about 9000 to about 10000 bases or base pairs in length. In some examples, a polyA sequence is at least about 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, or 200 nucleotides in length. A polyA sequence can comprise a length of for example, at least 1 base or base pair, 2 bases or

base pairs, 3 bases or base pairs, 4 bases or base pairs, 5 bases or base pairs, 6 bases or base pairs, 7 bases or base pairs, 8 bases or base pairs, 9 bases or base pairs, 10 bases or base pairs, 20 bases or base pairs, 30 bases or base pairs, 40 bases or base pairs, 50 bases or base pairs, 60 bases or base pairs, 70 bases or base pairs, 80 bases or base pairs, 90 bases or base pairs, 100 bases or base pairs, 200 bases or base pairs, 300 bases or base pairs, 400 bases or base pairs, 500 bases or base pairs, 600 bases or base pairs, 700 bases or base pairs, 800 bases or base pairs, 900 bases or base pairs, 1000 bases or base pairs, 2000 bases or base pairs, 3000 bases or base pairs, 4000 bases or base pairs, 5000 bases or base pairs, 6000 bases or base pairs, 7000 bases or base pairs, 8000 bases or base pairs, 9000 bases or base pairs, or 10000 bases or base pairs in length. A polyA sequence can comprise a length of at most 100 bases or base pairs, 90 bases or base pairs, 80 bases or base pairs, 70 bases or base pairs, 60 bases or base pairs, 50 bases or base pairs, 40 bases or base pairs, 30 bases or base pairs, 20 bases or base pairs, 10 bases or base pairs, or 5 bases or base pairs.

[00125] In some cases, a percentage of the nucleotides in a poly-A sequence are modified. For instance, in some cases, fewer than 99%, 95%, 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, 5% or 1% of the nucleotides in a poly-A sequence are modified. In some cases, all of the nucleotides in a poly-A are modified.

[00126] A linker sequence can comprise any number of nucleotides. A linker can be attached to the modified nucleobase at an N-3 or C-5 position. The linker attached to the nucleobase can be diethylene glycol, dipropylene glycol, triethylene glycol, tripropylene glycol, tetraethylene glycol, tetraethylene glycol, divalent alkyl, alkenyl, alkynyl moiety, ester, amide, or an ether moiety. A linker sequence can comprise a length of about 1 to about 10 bases or base pairs, about 10 to about 20 bases or base pairs, about 20 to about 50 bases or base pairs, about 50 to about 100 bases or base pairs, about 100 to about 500 bases or base pairs, about 500 to about 1000 bases or base pairs, about 1000 to about 2000 bases or base pairs, about 2000 to about 3000 bases or base pairs, about 3000 to about 4000 bases or base pairs, about 4000 to about 5000 bases or base pairs, about 5000 to about 6000 bases or base pairs, about 6000 to about 7000 bases or base pairs, about 7000 to about 8000 bases or base pairs, about 8000 to about 9000 bases or base pairs, or about 9000 to about 10000 bases or base pairs in length. A linker sequence can comprise a length of for example, at least 1 base or base pair, 2 bases or base pairs, 3 bases or base pairs, 4 bases or base pairs, 5 bases or base pairs, 6 bases or base pairs, 7 bases or base pairs, 8 bases or base pairs, 9 bases or base pairs, 10 bases or base pairs, 20 bases or base pairs, 30 bases or base pairs, 40 bases or base pairs, 50 bases or base pairs, 60 bases or base pairs, 70 bases or base pairs, 80 bases or base pairs, 90 bases or base pairs, 100 bases or base

pairs, 200 bases or base pairs, 300 bases or base pairs, 400 bases or base pairs, 500 bases or base pairs, 600 bases or base pairs, 700 bases or base pairs, 800 bases or base pairs, 900 bases or base pairs, 1000 bases or base pairs, 2000 bases or base pairs, 3000 bases or base pairs, 4000 bases or base pairs, 5000 bases or base pairs, 6000 bases or base pairs, 7000 bases or base pairs, 8000 bases or base pairs, 9000 bases or base pairs, or at least 10000 bases or base pairs in length. A linker at most 10000 bases or base pairs, 5000 bases or base pairs, 4000 bases or base pairs, 3000 bases or base pairs, 2000 bases or base pairs, 1000 bases or base pairs, 900 bases or base pairs, 800 bases or base pairs, 700 bases or base pairs, 600 bases or base pairs, 500 bases or base pairs, 400 bases or base pairs, 300 bases or base pairs, 200 bases or base pairs, or 100 bases or base pairs in length.

[00127] In some cases, a percentage of the nucleotides in a linker sequence are modified. For instance, in some cases, fewer than 99%, 95%, 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, 5% or 1% of the nucleotides in a linker sequence are modified. In some cases, all of the nucleotides in a linker sequence are modified.

[00128] In some cases, a nucleic acid construct(s), a vector(s), or an engineered polyribonucleotide(s) can include at least one stop codon before the 3' untranslated region (UTR). In some cases, a nucleic acid construct(s), a vector(s), or an engineered polyribonucleotide(s) includes multiple stop codons. The stop codon can be selected from TGA, TAA and TAG. The stop codon may be modified or unmodified. In some cases, the nucleic acid construct(s), vector(s), or engineered polyribonucleotide(s) includes the stop codon TGA and one additional stop codon. In some cases, the nucleic acid construct(s), vector(s), or engineered polyribonucleotide(s) includes the addition of the TAA stop codon.

Encoded Polypeptides

[00129] In some cases, the disclosure provides a method for treating a subject having or at risk of having primary ciliary dyskinesia, the method comprising administering to the subject a composition that comprises a nucleic acid construct that encodes dynein axonemal intermediate chain 1 protein (DNAI1), armadillo repeat containing 4 (ARMC4), chromosome 21 open reading frame 59 (C21orf59), coiled-coil domain containing 103 (CCDC103), coiled-coil domain containing 114 (CCDC114), coiled-coil domain containing 39 (CCDC39), coiled-coil domain containing 40 (CCDC40), coiled-coil domain containing 65 (CCDC65), dynein (axonemal) assembly factor 1 (DNAAF1), dynein (axonemal) assembly factor 2 (DNAAF2), dynein (axonemal) assembly factor 3 (DNAAF3), dynein (axonemal) assembly factor 5 (DNAAF5),

dynein axonemal heavy chain 11 (DNAH11), dynein axonemal heavy chain 5 (DNAH5), dynein axonemal heavy chain 8 (DNAH8), dynein axonemal intermediate chain 2 (DNAI2), dynein axonemal light chain 1 (DNAL1), dynein regulatory complex subunit 1 (DRC1), dyslexia susceptibility 1 candidate 1 (DYX1C1), axonemal central pair apparatus protein (HYDIN), leucine rich repeat containing 6 (LRRC6), NME/NM23 family member 8 (NME8), oral-facial-digital syndrome 1 (OFD1), retinitis pigmentosa GTPase regulator (RPGR), radial spoke head 1 homolog (Chlamydomonas) (RSPH1), radial spoke head 4 homolog A (Chlamydomonas) (RSPH4A), radial spoke head 9 homolog (Chlamydomonas) (RSPH9), sperm associated antigen 1 (SPAG1), and zinc finger MYND-type containing 10 (ZMYND10), or a variant of any of the aforementioned, which nucleic acid construct includes codons that provide for heterologous or enhanced expression of said protein(s) or a variant thereof within cells of the subject, thereby treating the subject having or at risk of having primary ciliary dyskinesia.

[00130] The encoded polypeptides are polymer chains comprised of amino acid residue monomers which are joined together through amide bonds (peptide bonds). The amino acids may be of the L-optical isomer, the D-optical isomer or a combination thereof. A polypeptide can be a chain of at least three amino acids, peptide-mimetics, a protein, a recombinant protein, an antibody (monoclonal or polyclonal), an antigen, an epitope, an enzyme, a receptor, a vitamin, or a structure analogue or combinations thereof. A polyribonucleotide that is translated within a subject's body can generate an ample supply of specific peptides or proteins within a cell, a tissue, or across many cells and tissues of a subject. In some cases, a polyribonucleotide can be translated *in vivo* within the cytosol of a specific target cell(s) type or target tissue. In some cases, a polyribonucleotide can be translated *in vivo* to provide a protein whose gene has been associated with primary ciliary dyskinesia, a functional fragment thereof, or a protein that is at least 70% homologous to a human DNAI1 or a human DNAH5 protein. In some cases, a polyribonucleotide can be translated *in vivo* in various non-target cell types or target tissue(s). Non-limiting examples of cells that be target or non-target cells include: a) skin cells, e.g.: keratinocytes, melanocytes, urothelial cells; b) neural cells, e.g.: neurons, Schwann cells, oligodendrocytes, astrocytes; c) liver cells, e.g.: hepatocytes; d) intestinal cells, e.g.: goblet cell, enterocytes; e) blood cells; e.g.: lymphoid or myeloid cells; and f) germ cells; e.g.: sperm and eggs. Non-limiting examples of tissues include connective tissue, muscle tissue, nervous tissue, or epithelial tissue. In some cases, a target cell or a target tissue is a cancerous cell, tissue, or organ.

[00131] A polynucleotide sequence can be derived from one or more species. For example, a polynucleotide sequence can be derived from a human (*Homo sapiens*), a mouse

(e.g., *Mus musculus*), a rat (e.g., *Rattus norvegicus* or *Rattus rattus*), a microorganism (e.g., Chlamydomonas genus), or any other suitable creature. A polynucleotide sequence can be a chimeric combination of the sequence of one or more species.

[00132] In some cases, the endogenous translational machinery can add a post-translational modification to the encoded peptide. A post-translational modification can involve the addition of hydrophobic groups that can target the polypeptide for membrane localization, the addition of cofactors for increased enzymatic activity, or the addition of smaller chemical groups. The encoded polypeptide can also be post-translationally modified to receive the addition of other peptides or protein moieties. For instance, ubiquitination can lead to the covalent linkage of ubiquitin to the encoded polypeptide, SUMOylation can lead to the covalent linkage of SUMO (Small Ubiquitin-related MOdifier) to the encoded polypeptide, ISGylation can lead to the covalent linkage of ISG15 (Interferon-Stimulate Gene 15).

[00133] In some cases, the encoded polypeptide can be post-translationally modified to undergo other types of structural changes. For instance, the encoded polypeptide can be proteolytically cleaved, and one or more proteolytic fragments can modulate the activity of an intracellular pathway. The encoded polypeptide can be folded intracellularly. In some cases, the encoded polypeptide is folded in the presence of co-factors and molecular chaperones. A folded polypeptide can have a secondary structure and a tertiary structure. A folded polypeptide can associate with other folded peptides to form a quaternary structure. A folded-peptide can form a functional multi-subunit complex, such as an antibody molecule, which has a tetrameric quaternary structure. Various polypeptides that form classes or isotypes of antibodies can be expressed from a polyribonucleotide.

[00134] The encoded polypeptide can be post-translationally modified to change the chemical nature of the encoded amino acids. For instance, the encoded polypeptide can undergo post-translational citrullination or deimination, the conversion of arginine to citrulline. The encoded polypeptide can undergo post-translation deamidation; the conversion of glutamine to glutamic acid or asparagine to aspartic acid. The encoded polypeptide can undergo elimination, the conversion of an alkene by beta-elimination of phosphothreonine and phosphoserine, or dehydration of threonine and serine, as well as by decarboxylation of cysteine. The encoded peptide can also undergo carbamylation, the conversion of lysine to homocitrulline. An encoded peptide can also undergo racemization, for example, racemization of proline by prolyl isomerase or racemization of serine by protein-serine epimerase. In some cases, an encoded peptide can undergo serine, threonine, and tyrosine phosphorylation.

[00135] The activity of a plurality of biomolecules can be modulated by a molecule encoded by a polyribonucleotide. Non-limiting examples of molecules whose activities can be modulated by an encoded polynucleotide include: amino acids, peptides, peptide-mimetics, proteins, recombinant proteins antibodies (monoclonal or polyclonal), antibody fragments, antigens, epitopes, carbohydrates, lipids, fatty acids, enzymes, natural products, nucleic acids (including DNA, RNA, nucleosides, nucleotides, structure analogues or combinations thereof), nutrients, receptors, and vitamins.

[00136] Non-limiting examples of nucleotide sequences that can be a part of a polynucleotide of the disclosure are disclosed in **TABLE 3**.

[00137]

TABLE 3	
Name	Sequence
dynein axonemal intermediate chain 1 (DNAI1)	SEQ ID NOs: 14-16
dynein axonemal heavy chain 5 (DNAH5)	SEQ ID NOs: 17-18

[00138] A polypeptide sequence can be engineered to have a desired altered codon usage, such as the altered codon usage of SEQ ID NOs 15-16 or the altered codon usage of SEQ ID NOs 17-18. Computer software can be used, for example, to generate the codon usage of SEQ ID NO 14. A polypeptide sequence can share a % homology to an amino acid sequence of an endogenous polypeptide. A polypeptide sequence can share at most 10% homology, at most 20% homology, at most 30% homology, at most 40% homology, at most 50% homology, at most 60% homology, at most 70% homology, at most 80% homology, at most 90% homology, or at most 99% homology with an amino acid sequence of an endogenous polypeptide. Various methods and software programs can be used to determine the homology between two or more peptides, such as NCBI BLAST, Clustal W, MAFFT, Clustal Omega, AlignMe, Praline, or another suitable method or algorithm.

Immunogenicity

[00139] Many pharmaceutical agents, including compositions comprising molecules of various sizes (polynucleotides, proteins, or enzymes) can trigger an immune response when administered to a subject. In many cases, the immune system recognizes the composition as a foreign body and neutralizes its pharmaceutical action. A polyribonucleotide and a composition of the present disclosure can have low immunogenicity or be non-immunogenic, thereby triggering a small response by the immune system, or not triggering any immune response at all.

[00140] The immunogenicity can also be determined by measurement of, for example, the TNF- α and IL-8 levels and the binding capacity to TLR-3, TLR-7, TLR-8 and helicase RIG-1. In order thereby to establish whether a polyribonucleotide has a desired low immunogenicity, the quantity of one or more of the factors can be measured after administration of the polyribonucleotide to a subject. The immunogenicity of a polypeptide can be determined in relation to an increase in the number of white blood cells upon administration of the polypeptide to the subject. In some cases, upon administration of the composition to the subject, the subject exhibits an increase in the number of white blood cells that is less than 90%, less than 80%, less than 70%, less than 60%, less than 50%, less than 40%, less than 30%, less than 20%, or less than 10%. A polyribonucleotide of the disclosure can trigger minimum or insignificant inflammatory or immunological reactions.

[00141] For the determination of the immunogenicity of a polyribonucleotide, various methods can be used. A very suitable method is the determination of inflammatory markers in cells or a simple white cell blood count, as a reaction to the administration of the polyribonucleotide. Such a method is described in the examples. Cytokines which are associated with inflammation, such as, for example TNF- α , IFN- α , IFN- β , IP-10, IL-8, IL-6, and/or IL-12, can be measured. The expression of dendritic cell activation markers can also be used for the estimation of immunogenicity. A further indication of an immunological reaction can be the detection of binding to the Toll-like receptors TLR-3, TLR-7 and TLR-8 and to helicase RIG-1.

[00142] The immunogenicity of a polyribonucleotide can be determined as an overall increase in the level of inflammatory marker or white blood cell count as compared to a level prior to the administration of the polyribonucleotide. For instance, an engineered polyribonucleotide that is unmodified or modified can be administered to cells, or to a subject, and the secretion of inflammatory markers in a defined time interval as a reaction to the administration of the polyribonucleotide can be measured.

Compositions

[00143] In some cases, the disclosure provides a composition comprising a nucleic acid construct encoding dynein axonemal intermediate chain 1, wherein the nucleic acid construct comprises a complementary deoxyribonucleic acid encoding dynein axonemal intermediate chain 1, which composition is formulated for administration to a subject. In some cases, the disclosure provides a composition comprising a nucleic acid construct encoding dynein axonemal intermediate chain 1, which nucleic acid construct includes codons that provide for heterologous or enhanced expression of the dynein axonemal intermediate chain 1 protein or a

variant thereof within cells of a subject having or at risk of having primary ciliary dyskinesia. In some cases, the disclosure provides a composition comprising a nucleic acid construct encoding dynein axonemal intermediate chain 1, wherein fewer than 30% of the nucleic acids encoding dynein axonemal intermediate chain 1 are nucleic acid analogues, such as pseudouridine or 1-methyl pseudouridine. In some cases, the coding sequence of these constructs is engineered to have an altered nucleotide usage in the protein coding regions to increase its stability.

[00144] In some cases, the codons of the construct are at least 70% homologous to a mammalian or to a human dynein axonemal intermediate chain 1 protein. The construct may also comprise a 3' or 5' noncoding region flanking the codon sequence which encodes a protein of interest, such as dynein axonemal intermediate chain 1, wherein the noncoding region enhances the expression of the protein within cells the subject. The 3' noncoding region flanking the codon can comprise a 3'-cap independent translation enhancer (3'-CITEs) or a 3'-stem loop region derived from the nucleotide sequence of a histone protein or a 3'-triple helical structure derived from the nucleotide sequence of metastasis-associated lung adenocarcinoma transcript 1 (MALAT1). The 3' noncoding region flanking the codon can comprise a poly adenosine tail, wherein the number of adenosines in the poly adenosine tail improves the translation efficiency or the half-life of the protein of interest, such as dynein axonemal intermediate chain 1 protein. In some cases, the length of the poly adenosine tail is at most 200 adenosines. In some cases, a percentage of the poly adenosine tail comprises nucleic acid analogues. Fewer than 50%, 40%, 30%, 20%, 10%, or 5% of the nucleic acids in the poly adenosine tail can be nucleic acid analogues.

[00145] When the composition comprises a percentage of nucleotide analogues the nucleotide analogues can be selected from the group consisting of pseudouridine, 1-methylpseudouridine, 2-thiouridine, 5-methyluridine, 5-methoxyuridine, 5-methylcytidine, 2'-amino-2'-deoxycytidine, 2'-fluoro-2'-deoxycytidine, and. In some cases, the nucleic acid analogue is pseudouridine or 1-methylpseudouridine. In some cases, the nucleic acid analogue is 5-methoxyuridine.

[00146] In some cases, the composition comprises a nucleic acid encoding dynein axonemal intermediate chain 1 and/or nucleic acid analogues. Optionally, the composition can further comprise at least one additional nucleic acid construct. The at least one additional nucleic acid construct may encode a protein selected from the group consisting of: armadillo repeat containing 4 (ARMC4), chromosome 21 open reading frame 59 (C21orf59), coiled-coil domain containing 103 (CCDC103), coiled-coil domain containing 114 (CCDC114), coiled-coil domain containing 39 (CCDC39), coiled-coil domain containing 40 (CCDC40), coiled-coil domain

containing 65 (CCDC65), dynein (axonemal) assembly factor 1 (DNAAF1), dynein (axonemal) assembly factor 2 (DNAAF2), dynein (axonemal) assembly factor 3 (DNAAF3), dynein (axonemal) assembly factor 5 (DNAAF5), dynein axonemal heavy chain 11 (DNAH11), dynein axonemal heavy chain 5 (DNAH5), dynein axonemal heavy chain 8 (DNAH8), dynein axonemal intermediate chain 2 (DNAI2), dynein axonemal light chain 1 (DNAL1), dynein regulatory complex subunit 1 (DRC1), dyslexia susceptibility 1 candidate 1 (DYX1C1), axonemal central pair apparatus protein (HYDIN), leucine rich repeat containing 6 (LRRC6), NME/NM23 family member 8 (NME8), oral-facial-digital syndrome 1 (OFD1), retinitis pigmentosa GTPase regulator (RPGR), radial spoke head 1 homolog (Chlamydomonas) (RSPH1), radial spoke head 4 homolog A (Chlamydomonas) (RSPH4A), radial spoke head 9 homolog (Chlamydomonas) (RSPH9), sperm associated antigen 1 (SPAG1), and zinc finger MYND-type containing 10 (ZMYND10).

[00147] The compositions may comprise engineered polyribonucleotides, vectors, or nucleic acid constructs. “Naked” polynucleotide compositions can be successfully administered to a subject, and uptaken by a subject’s cell, without the aid of carriers, stabilizers, diluents, dispersing agents, suspending agents, thickening agents, and/or excipients (Wolff et al. 1990, Science, 247, 1465-1468). However, in many instances, encapsulation of polynucleotides with formulations that can increase the endocytotic uptake can increase the effectiveness of a composition of the disclosure. To overcome this challenge, in some cases, the composition comprises a nucleic acid construct, a vector, or an isolated nucleic acid encoding dynein axonemal intermediate chain 1, wherein the nucleic acid construct comprises a complementary deoxyribonucleic acid encoding dynein axonemal intermediate chain 1, which composition is formulated for administration to a subject.

[00148] Another technical challenge underlying the delivery of polyribonucleotides to multicellular organisms is to identify a composition that provides a high efficiency delivery of polyribonucleotides that are translated within a cell or a tissue of a subject. It has been recognized that administration of naked nucleic acids may be highly inefficient and may not provide a suitable approach for administration of a polynucleotide to a multicellular organism.

[00149] To solve this challenge, a composition comprising an engineered polyribonucleotide can be encapsulated or formulated with a pharmaceutical carrier. The formulation may be, but is not limited to, nanoparticles, poly(lactic-co-glycolic acid) (PLGA) microspheres, lipidoids, lipoplex, liposome, polymers, carbohydrates (including simple sugars), cationic lipids, fibrin gel, fibrin hydrogel, fibrin glue, fibrin sealant, fibrinogen, thrombin, rapidly eliminated lipid nanoparticles (reLNPs) and combinations thereof. A composition comprising an

engineered polyribonucleotide disclosed herein can comprise from about 1% to about 99% weight by volume of a carrier system. The amount of carrier present in a carrier system is based upon several different factors or choices made by the formulator, for example, the final concentration of the polyribonucleotide and the amount of solubilizing agent. Various carriers have been shown useful in delivery of different classes of therapeutic agents. Among these carriers, biodegradable nanoparticles formulated from biocompatible polymers poly(D,L-lactide-co-glycolide) (PLGA) and polylactide (PLA) have shown the potential for sustained intracellular delivery of different therapeutic agents.

[00150] The loading weight percent of the engineered polynucleotide in a composition may be at least 0.05%, 0.1%, 0.2%, 0.3%, 0.4%, 0.5%, 1%, 2%, 4 %, 5 %, 6 %, 7 %, 8 %, 9 %, or 10 %. The encapsulation efficiency of the modified mRNA in the PLGA microsphere may be at least 50%, at least 70%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%.

[00151] The present disclosure describes nanoparticles, oligomers, polymers or lipidoids comprising oligo(alkylene amines) containing alternating, non-identical alkylene amine units which are useful for delivering a polynucleotide, in some cases an engineered polyribonucleotides, into a cell or into a tissue. A composition disclosed herein can be stable for at least about 1 minute, 5 minutes, 10 minutes, 30 minutes, 1 hour, 2 hours, 3 hours, 4 hours, 5 hours, 6 hours, 12 hours, 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days, 2 weeks, 4 weeks, 6 weeks, 8 weeks, 10 weeks, 12 weeks, 3 months, 4 months, 5 months, 6 months, 7 months, 8 months, 9 months, 10 months, 11 months, or one year. A formulation disclosed herein can be stable, for example, at a temperature of at least about 0°C, 5°C, 10°C, 15°C, 20°C, 25°C, 30°C, 35°C, 40°C, 45°C, 50°C, 60°C, 70°C, or 80°C. A composition of the disclosure can have a desired density. The density of a composition can improve a property of the composition, such as the rheology of the composition.

Nanoparticles

[00152] The present disclosure also provides nanoparticle based formulations of nucleic acid constructs, engineered polyribonucleotides, or vectors that are able to translocate following administration to a subject. In some instances, the administration is pulmonary and the engineered polyribonucleotides can move intact either actively or passively from the site of administration to the systemic blood supply and subsequently to be deposited in different cells or tissues, such as, e.g., the breast. This translocation of the nanoparticle comprising an engineered polyribonucleotide encoding a therapeutic protein, such as, e.g., dynein axonemal intermediate

chain 1 (DNAI1), armadillo repeat containing 4 (ARMC4), chromosome 21 open reading frame 59 (C21orf59), coiled-coil domain containing 103 (CCDC103), coiled-coil domain containing 114 (CCDC114), coiled-coil domain containing 39 (CCDC39), coiled-coil domain containing 40 (CCDC40), coiled-coil domain containing 65 (CCDC65), cyclin O (CCNO), dynein (axonemal) assembly factor 1 (DNAAF1), dynein (axonemal) assembly factor 2 (DNAAF2), dynein (axonemal) assembly factor 3 (DNAAF3), dynein (axonemal) assembly factor 5 (DNAAF5), dynein axonemal heavy chain 11 (DNAH11), dynein axonemal heavy chain 5 (DNAH5), dynein axonemal heavy chain 6 (DNAH6), dynein axonemal heavy chain 8 (DNAH8), dynein axonemal intermediate chain 2 (DNAI2), dynein axonemal light chain 1 (DNAL1), dynein regulatory complex subunit 1 (DRC1), dyslexia susceptibility 1 candidate 1 (DYX1C1), growth arrest specific 8 (GAS8), axonemal central pair apparatus protein (HYDIN), leucine rich repeat containing 6 (LRRC6), NME/NM23 family member 8 (NME8), oral-facial-digital syndrome 1 (OFD1), retinitis pigmentosa GTPase regulator (RPGR), radial spoke head 1 homolog (Chlamydomonas) (RSPH1), radial spoke head 4 homolog A (Chlamydomonas) (RSPH4A), radial spoke head 9 homolog (Chlamydomonas) (RSPH9), sperm associated antigen 1 (SPAG1), and zinc finger MYND-type containing 10 (ZMYND10) or a functional fragment thereof, constitutes non-invasive systemic delivery of an active pharmaceutical ingredient beyond the lung to result in the production of a functional protein to systemically accessible non-lung cells or tissues.

[00153] A nanoparticle can be a particle of particle size from about 10 nanometers (nm) to 5000 nm, 10 nm to 1000 nm, or 60 nm to 500 nm, or 70 nm to 300 nm. In some examples, a nanoparticle has a particle size from about 60 nm to 225 nm. The nanoparticle can include an encapsulating agent (e.g., coating) that encapsulates one or more polyribonucleotides, which may be engineered polyribonucleotides. The nanoparticle can include engineered and/or naturally occurring polyribonucleotides. The encapsulating agent can be a polymeric material, such as PEI or PEG.

[00154] A lipidoid or lipid nanoparticle which may be used as a delivery agent may include a lipid which may be selected from the group consisting of C12-200, MD1, 98N12-5, DLin-DMA, DLin-K-DMA, DLin-KC2-DMA, DLin-MC3-DMA, PLGA, PEG, PEG-DMG, PEGylated lipids and analogues thereof. A suitable nanoparticle can comprise one or more lipids in various ratios. For example, a composition of the disclosure can comprise a 40:30:25:5 ratio of C12-200:DOPE:Cholesterol:DMG-PEG2000 or a 40:20:35:5 ratio of HGT5001:DOPE:Cholesterol:DMG-PEG2000. A nanoparticle can include at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 lipids or another suitable number of lipids. A nanoparticle can be formed of any

suitable ratio of lipids selected from the group consisting of C12-200, MD1, 98N12-5, DLin-DMA, DLin-K-DMA, DLin-KC2-DMA, DLin-MC3-DMA, PLGA, PEG, PEG-DMG.


[00155] The mean size of the nanoparticle formulation may comprise the modified mRNA between 60 nanometers (nm) and 225 nm. The polydispersity index PDI of the nanoparticle formulation comprising the modified mRNA can be between 0.03 and 0.15. The zeta potential of the nanoparticle formulation may be from -10 to +10 at a pH of 7.4. The formulations of modified mRNA may comprise a fusogenic lipid, cholesterol and a PEG lipid. The formulation may have a molar ratio 50:10:38.5:1.5-3.0 (cationic lipid: fusogenic lipid: cholesterol: polyethylene glycol (PEG) lipid). The PEG lipid may be selected from, but is not limited to PEG-c-DOMG, PEG-DMG. The fusogenic lipid may be DSPC. A lipid nanoparticle of the present disclosure can be formulated in a sealant such as, but not limited to, a fibrin sealant.

Oligo(alkylene amine groups)

[00156] It has also been recognized that although encapsulation of polynucleotides with some formulations can increase the endocytotic uptake of a composition, the polynucleotide that is taken up by a cell may not be effectively translated within the cell. Some formulations may be effectively used for plasmid DNA and/or siRNA delivery, while not practical for use in the delivery of polyribonucleotides. The present disclosure provides formulations that can be employed for effective delivery and translation of polyribonucleotide compositions to a subject.

[00157] A composition of the disclosure can be designed to provide a polyribonucleotide that is effectively translated within a cell. A composition of the disclosure can comprise an arrangement of alkylene amine units of alternating length in groups of three or more units and containing an ethyleneamine unit in compositions for transfecting a cell with any polynucleotide, such as an engineered polyribonucleotide. A composition of the disclosure can provide a more efficacious delivery of a polyribonucleotide to a cell than analogous arrangements of alkylene amine units of non-alternating length.

[00158] Oligomers, polymers or lipidoids can be provided which share a common structural entity which is illustrated in formula (I):




[00159] A composition of the disclosure can comprise an oligo(alkylene amine) that is selected from:

m is 1 or 2; n is 0 or 1 and $m + n \geq 2$; and

R^2 to R^5 are, independently of each other, selected from hydrogen; a group $-\text{CH}_2-\text{CH}(\text{OH})-\text{R}^7$, $-\text{CH}(\text{R}^7)-\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{CH}_2-(\text{C}=\text{O})-\text{O}-\text{R}^7$, $-\text{CH}_2-\text{CH}_2-(\text{C}=\text{O})-\text{NH}-\text{R}^7$, $-\text{CH}_2-\text{R}^7$ or $-\text{CH}_2-$ wherein R^7 is selected from C3–C18 alkyl or C3–C18 alkenyl having one C–C double bond; a protecting group for an amino group; and a poly(ethylene glycol) chain;

and wherein one or more of the nitrogen atoms indicated in formula (III) may be protonated to provide a cationic group of formula (III).

[00162] c) a lipidoid having the structure of formula (IV):



wherein the variables a , b , p , m , n , and R^2 to R^6 are defined as follows:

a is 1 and b is an integer of 2 to 4; or a is an integer of 2 to 4 and b is 1,

p is 1 or 2,

m is 1 or 2; n is 0 or 1 and $m + n \geq 2$; and


R^2 to R^6 are, independently of each other, selected from hydrogen; a group $-\text{CH}_2-\text{CH}(\text{OH})-\text{R}^7$, $-\text{CH}(\text{R}^7)-\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{CH}_2-(\text{C}=\text{O})-\text{O}-\text{R}^7$, $-\text{CH}_2-\text{CH}_2-(\text{C}=\text{O})-\text{NH}-\text{R}^7$, or $-\text{CH}_2-\text{R}^7$ wherein R^7 is selected from C3–C18 alkyl or C3–C18 alkenyl having one C–C double bond; a protecting group for an amino group; and a poly(ethylene glycol) chain; and a receptor ligand; provided that at least two residues among R^1 to R^6 are a group $-\text{CH}_2-\text{CH}(\text{OH})-\text{R}^7$, $-\text{CH}(\text{R}^7)-\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{CH}_2-(\text{C}=\text{O})-\text{O}-\text{R}^7$, $-\text{CH}_2-\text{CH}_2-(\text{C}=\text{O})-\text{NH}-\text{R}^7$, or $-\text{CH}_2-\text{R}^7$ wherein R^7 is selected from C3–C18 alkyl or C3–C18 alkenyl having one C–C double bond;

and wherein one or more of the nitrogen atoms indicated in formula (IV) may be protonated to provide a cationic group of formula (IV).

[00163] Non-limiting examples of alkenyl and alkenylene groups include straight, branched, and cyclic alkenyl groups. The olefin or olefins of an alkenyl group can be, for example, *E*, *Z*, *cis*, *trans*, terminal, or *exo*-methylene. An alkenylene group can be, for example, a C_2 , C_3 , C_4 , C_5 , C_6 , C_7 , C_8 , C_9 , C_{10} , C_{11} , C_{12} , C_{13} , C_{14} , C_{15} , C_{16} , C_{17} , C_{18} , C_{19} , C_{20} , C_{21} , C_{22} , C_{23} , C_{24} , C_{25} , C_{26} , C_{27} , C_{28} , C_{29} , C_{30} , C_{31} , C_{32} , C_{33} , C_{34} , C_{35} , C_{36} , C_{37} , C_{38} , C_{39} , C_{40} , C_{41} , C_{42} , C_{43} , C_{44} , C_{45} , C_{46} , C_{47} , C_{48} , C_{49} , or C_{50} group that is substituted or unsubstituted.

[00164] The oligo(alkylene amine) structures of formulae (II), (III) and (IV) are characterized in that they can combine shorter (also referred to for illustration as “S”) ethylene amine units (i.e., a or b is 1) with longer (also referred to for illustration as “L”) alkylene amine units (i.e., the other one of a or b is an integer of 2 to 4) in an alternating manner. Such an arrangement of the protonatable units can provide advantages in terms of the suitability of the resulting group to provide a vehicle for delivering polyribonucleotides into a cell.

[00165] A composition of the disclosure can comprise a plurality of oligo(alkylene amine) groups of formula (II) as a side chain or as a terminal group:



wherein the variables a, b, p, m, n, and R² to R⁶ are defined as follows, independently for each group of formula (II) in a plurality of such groups:

a is 1 and b is an integer of 2 to 4; or a is an integer of 2 to 4 and b is 1,

p is 1 or 2,

m is 1 or 2; n is 0 or 1 and m + n is ≥ 2; and

R² to R⁵ are, independently of each other, selected from hydrogen; a group -CH₂-CH(OH)-R⁷, -CH(R⁷)-CH₂-OH, -CH₂-CH₂-(C=O)-O-R⁷, -CH₂-CH₂-(C=O)-NH-R⁷, or -CH₂-R⁷ wherein R⁷ is selected from C3-C18 alkyl or C3-C18 alkenyl having one C-C double bond; a protecting group for an amino group; -C(NH)-NH₂-; and a poly(ethylene glycol) chain;

R⁶ is selected from hydrogen; a group -CH₂-CH(OH)-R⁷, -CH(R⁷)-CH₂-OH, -CH₂-CH₂-(C=O)-O-R⁷, -CH₂-CH₂-(C=O)-NH-R⁷, or -CH₂-R⁷ wherein R⁷ is selected from C3-C16 alkyl or C3-C16 alkenyl having one C-C double bond; a protecting group for an amino group; -C(NH)-NH; a poly(ethylene glycol) chain; and a receptor ligand.

[00167] In some cases, R² to R⁵ are hydrogen and R⁶ is selected from hydrogen, a protecting group for an amino group; -C(NH)-NH₂ and a poly(ethylene glycol) chain. In some cases, R² to R⁶ are hydrogen. In some cases, R⁷ is selected from C8-C18 alkyl or C8-C18 alkenyl having one C-C double bond, or from C8-C12 alkyl or C8-C12 alkenyl having one C-C double bond, or from C10-C12 alkyl or C10-C12 alkenyl having one C-C double bond. A composition of the disclosure can comprise one, or multiple alkylene groups of formulas (II)-(IV).

[00168] In some cases, the oligomers or polymers which can be used in the compositions in accordance with the present disclosure comprise a plurality of oligo (alkylene amine) groups of formula (III) as repeating units:

[00169] $\text{NR}^2\{\text{CH}_2-(\text{CH}_2)_a-\text{NR}^3-[\text{CH}_2-(\text{CH}_2)_b-\text{NR}^4]_p\}_m-[\text{CH}_2-(\text{CH}_2)_a-\text{NR}^5]_n-$ (III)

wherein the variables a, b, p, m, n, and R^2 to R^5 are defined as follows, independently for each group of formula (III) in a plurality of such groups:

a is 1 and b is an integer of 2 to 4; or a is an integer of 2 to 4 and b is 1,

p is 1 or 2,

m is 1 or 2; n is 0 or 1 and $m + n \geq 2$; and


R^2 to R^5 are, independently of each other, selected from hydrogen; a group $-\text{CH}_2-\text{CH}(\text{OH})-\text{R}^7$, $-\text{CH}(\text{R}^7)-\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{CH}_2-(\text{C}=\text{O})-\text{O}-\text{R}^7$, $-\text{CH}_2-\text{CH}_2-(\text{C}=\text{O})-\text{NH}-\text{R}^7$, $-\text{CH}_2-\text{R}^7$ or $-\text{CH}_2-$ wherein R^7 is selected from C3–C18 alkyl or C3–C18 alkenyl having one C–C double bond; a protecting group for an amino group; $-\text{C}(\text{NH})-\text{NH}_2$; a poly(ethylene glycol) chain; and endosomal escape effector and a receptor ligand. In some cases, R^2 to R^5 are hydrogen. In some cases, R^7 is selected from C8–C18 alkyl or C8–C18 alkenyl having one C–C. R^7 may be selected from C8–C12 alkyl or C8–C12 alkenyl having one C–C. As an alternative, R^7 may be selected from C10–C12 alkyl or C10–C12 alkenyl having one C–C.

[00170] One or more of the nitrogen atoms indicated in formula (III) may be protonated to provide a cationic group of formula (III).


[00171] Optionally, the oligomers or polymers which comprise a plurality of groups of formula (III) as repeating units can comprise, in addition, one or more oligo(alkylene amine) group(s) of formula (II) as a side chain and/or as a terminal group.

[00172] In a plurality of groups of formula (III) as repeating units, two, three or more of the groups of formula (III) can be contained in the oligomers or polymers. Generally, substances comprising 2 to 9 repeating units are referred to herein as oligomers, those comprising 10 and more repeating units as polymers. Thus, in the polymers containing a plurality of groups of formula (III) as repeating units, 10 or more groups of formula (III) may be present. It will be understood that the groups of formula (III) can have the same structure within a polymer or oligomer, or can have two or more different structures within the scope of formula (III). In some cases, the oligomers or polymers containing a plurality of groups of formula (III) as repeating units can be provided in the form of a library of sequence defined polymers which are prepared from different groups of formula (III) in a controlled, stepwise polymerization.

[00173] In line with formulae (II) and (III) above, an alkylene amine unit may be repeated once in an alternating chain such that oligo(alkylene amine) moieties of the type –S–L–L–S– or –L–S–S–L– may result, wherein S represents a shorter ethylene amine unit, and L represents a longer alkylene amine unit. In some cases, groups of formula (II) and (III) are those wherein no repetition occurs, i.e., wherein p is 1, such that the shorter or longer units do not appear in pairs. The group of formula (II) can be an oligo(alkylene amine) group of formula (IIa) and the group of formula (III) can be an oligo(alkylene amine) group of (IIIa):




wherein a, b, m, n, and R² to R⁶ are defined as in formula (II), and wherein one or more of the nitrogen atoms indicated in formula (IIa) may be protonated to provide a cationic oligomer or polymer structure;




wherein a, b, m, n, and R² to R⁵ are defined as in formula (III), and wherein one or more of the nitrogen atoms indicated in formula (IIa) can be protonated to provide a cationic oligomer or polymer structure.

[00176] Moreover, in some cases, the oligo(alkylene amine) group of formulae (II) and (III) can have an n of 1. In some cases, m is 1 and n is 1. In some cases, the group of formula (II) is an oligo(alkylene amine) group of formula (IIb), and the group of formula (III) is an oligo(alkylene amine) group of formula (IIIb):




wherein a, b, and R² to R⁶ are defined as in formula (II), and wherein one or more of the nitrogen atoms indicated in formula (IIb) can be protonated to provide a cationic oligomer or polymer structure;




wherein a, b, and R² to R⁵ are defined as in formula (III) and wherein one or more of the nitrogen atoms indicated in formula (IIIb) can be protonated to provide a cationic oligomer or polymer structure.

[00179] With respect to the length of the alkylene amine units in the oligo(alkylene amine) groups of formula (II), (IIa), (IIb) and (III), (IIIa), (IIIb), one of the alternating units can be an ethylene amine unit (i.e., either a or b is 1). The other alternating unit can be a propylene amine

unit, a butylene amine unit or a pentylene amine unit (i.e., the other one of a or b can be an integer from 2 to 4. In some cases, the other of a or b can be 2 or 3, and in some cases, a is 1 and b is 2, or a is 2 and b is 1. In some cases, an oligo(alkylene amine) group of formula (IIc) is employed instead of or in addition to group (II), and/or an oligo(alkylene amine) group of formula (IIIc) is employed instead of or in addition to group (III). The formulae of group (IIc) and group (IIIc) are as follows:



[00181] wherein R^2 to R^6 are as defined in formula (II), and wherein R^2 to R^6 are hydrogen, and wherein one or more of the nitrogen atoms indicated in formula (IIc) can be protonated to provide a cationic oligomer or polymer structure;



[00183] wherein R^2 to R^5 are as defined in formula (III), and wherein one or more of the nitrogen atoms indicated in formula (IIIc) can be protonated to provide a cationic oligomer or polymer structure.

[00184] In some cases, the groups R^2 to R^6 in formula (II), (IIa), (IIb) and (IIc) or the groups R^2 to R^5 in formula (III), (IIIa), (IIIb) and (IIIc) can be protecting group for an amino group. Non-limiting examples of protecting groups include t-butoxycarbonyl (Boc), 9-fluorenylmethoxycarbonyl (Fmoc), or carbobenzyloxy (Cbz).

[00185] In some cases, the groups R^1 to R^6 in formula (II), (IIa), (IIb) and (IIIc) or the groups R^2 to R^5 in formula (III), (IIIa), (IIIb) and (IIIc) are a receptor ligand, such as the receptor ligands described in Philipp and Wagner in "Gene and Cell Therapy - Therapeutic Mechanisms and Strategy", 3rd Edition, Chapter 15, CRC Press, Taylor & Francis Group LLC, Boca Raton 2009. Examples of receptor ligands that target the lung tissue are described in Pfeifer et al. 2010, Ther. Deliv. 1 (1): 133-48. Receptor ligands can include synthetic cyclic or linear peptides such as derived from screening peptide libraries for binding to a particular cell surface structure or particular cell type, cyclic or linear RGD peptides, synthetic or natural carbohydrates such as sialic acid, galactose or mannose or synthetic ligands derived from reacting a carbohydrate for example with a peptide, antibodies specifically recognizing cell surface structures, folic acid, epidermal growth factor and peptides derived thereof, transferrin, anti-transferrin receptor antibodies, nanobodies and antibody fragments, approved drugs that may bind to cell surface molecules (e.g., cell surface receptors), etc.

[00186] As far as any of the groups R^1 to R^6 in formula (II), (IIa), (IIb) and (IIc) or the groups R^2 to R^5 in formula (III), (IIIa), (IIIb) and (IIIc) are a poly(ethylene glycol) chain, the

molecular weight of the poly(ethylene glycol) chain can be from about 100 g/mol to 20,000 g/mol, from about 1,000 g/mol to 10,000 g/mol or from about 1,000 g/mol to 5,000 g/mol.

[00187] In some cases, group (II) can be an oligo(alkylene amine) group of formula (II):

[00188] $-\text{NH}-\text{CH}_2-\text{CH}_2-\text{NH}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{NH}-\text{CH}_2-\text{CH}_2-\text{NH}-\text{H}$ (II),

[00189] wherein one or more of the nitrogen atoms indicated in formula (II) may be protonated to provide a cationic polymer or dendrimer structure. In some cases, group (III) is an oligo(alkylene amine) group of formula (III):

[00190] $-\text{NH}-\text{CH}_2-\text{CH}_2-\text{NH}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{NH}-\text{CH}_2-\text{CH}_2-\text{NH}-$ (III)

[00191] wherein one or more of the nitrogen atoms indicated in formula (III) may be protonated to provide a cationic polymer or dendrimer structure.

Lipidoids

[00192] An engineered polyribonucleotide can be encapsulated in a lipidoid formulation. A lipidoid formulation can be any material that has characteristics of a lipid, such as fats, waxes, sterols, fat-soluble vitamins (such as vitamins A, D, E, and K), monoglycerides, diglycerides, triglycerides, phospholipids, and others. For example, a lipid or lipidoid formulation can include lipids such as cholesterol, DOPE, DOPC or DSPC which are referred to as helper lipids in the scientific literature, and/or PEGylated lipids or any other lipid useful for preparing lipoplexes. The formulation comprising the engineered polyribonucleotide may be a nanoparticle which may comprise at least one lipid. A lipidoid formulation can be a lipid nanoparticle. The lipid may be selected from, but is not limited to, DOPE, DOPC, DSPC, cholesterol, DLin-DMA, DLin-K-DMA, 98N12-5, C12-200, DLin-MC3-DMA, DLin-KC2-DMA, DODMA, PLGA, PEG, PEG-DMG and PEGylated lipids. In another aspect, the lipid may be a cationic lipid such as, but not limited to, DLin-DMA, DLin-D-DMA, DLin-MC3-DMA, DLin-KC2-DMA and DODMA.

[00193] The composition containing a lipidoid may be about 40-60% lipidoid, about 40-60 % cholesterol, and about 5-20% PEG-lipid (in percent by weight, based on the total weight of the composition). The composition containing a lipidoid may be about 50-60% lipidoid, about 40- 50 % cholesterol, and about 5-10% PEG-lipid. The composition containing a lipidoid may be about 50-75% lipidoid, about 20-40% cholesterol, and about 1-10% PEG-lipid. The composition containing a lipidoid may be about 60-70% lipidoid, about 25-35% cholesterol, and about 5-10% PEG-lipid. The composition may be provided with techniques described in, for example, Akinc et al, 2007, Nat Biotech, 26, 561-569; Akinc et al, 2009, Mol Ther, 17, 872-9; Love et al, 2010, PNAS, 107, 1864-9; US 8,450,298, O2006/138380). RNA/lipidoid complexes may form

particles that are useful in the delivery of RNA, such as single-stranded RNAs or mRNAs, into cells.

[00194] A composition of the disclosure can be an engineered polyribonucleotide encapsulated by a lipidoid of formula (IV)

[00195] $R^1-NR^2\{CH_2-(CH_2)_a-NR^3-[CH_2-(CH_2)_b-NR^4]_p\}_m-[CH_2-(CH_2)_a-NR^5]_n-R^6$
(IV),

[00196] wherein the variables a, b, p, m, n and R¹ to R⁶ are defined as follows:

a is 1 and b is an integer of 2 to 4; or a is an integer of 2 to 4 and b is 1,

p is 1 or 2,

m is 1 or 2; n is 0 or 1 and m + n is ≥ 2 ; and

R¹ to R⁶ are independently of each other selected from hydrogen; a group $-CH_2-CH(OH)-R^7$, $-CH(R^7)-CH_2-OH$, $-CH_2-CH_2-(C=O)-O-R^7$, $-CH_2-CH_2-(C=O)-NH-R^7$ or $-CH_2-R^7$ wherein R⁷ is selected from C3-C18 alkyl or C3-C18 alkenyl having one C-C double bond; a protecting group for an amino group; $-C(NH)-NH_2$; a poly(ethylene glycol) chain; and a receptor ligand; provided that at least two residues among R¹ to R⁶ are a group $-CH_2-CH(OH)-R^7$, $-CH(R^7)-CH_2-OH$, $-CH_2-CH_2-(C=O)-O-R^7$, $-CH_2-CH_2-(C=O)-NH-R^7$ or $-CH_2-R^7$ wherein R⁷ is selected from C3-C18 alkyl or C3-C18 alkenyl having one C-C double bond.

[00197] In some cases, R¹ to R⁶ are independently selected from hydrogen; a group $-CH_2-C(OH)H-R^7$ or $-CH(R^7)-CH_2-OH$, wherein R⁷ is selected from C3-C18 alkyl or C3-C18 alkenyl having one C-C double bond; a protecting group for an amino group; and a poly(ethylene glycol) chain; provided that at least two residues among R¹ to R⁶ are a group $-CH_2-C(OH)H-R^7$ or $-CH(R^7)-CH_2-OH$, wherein R⁷ is selected from C3-C18 alkyl or C3-C18 alkenyl having one C-C double bond. In some cases, R¹ to R⁶ are independently selected from hydrogen; and a group $-CH_2-CH(OH)-R^7$ or $-CH(R^7)-CH_2-OH$ wherein R⁷ is selected from C3-C16 alkyl or C3-C16 alkenyl having one C-C double bond; provided that at least two residues among R¹ to R⁶ are a group $-CH_2-CH(OH)-R^7$ or $-CH(R^7)-CH_2-OH$, wherein R⁷ is selected from C3-C18 alkyl or C3-C18 alkenyl having one C-C double bond. In some cases, R¹ and R⁶ are independently selected from hydrogen; and a group $-CH_2-CH(OH)-R^7$ or $-CH(R^7)-CH_2-OH$ wherein R⁷ is selected from C3-C18 alkyl or C3-C18 alkenyl having one C-C double bond; and R² to R⁵ are all a group $-CH_2-CH(OH)-R^7$ or $-CH(R^7)-CH_2-OH$ wherein R⁷ is selected from C3-C18 alkyl or C3-C18 alkenyl having one C-C double bond. In some cases, R⁷ is selected from C8-C16 alkyl or C8-C18 alkenyl having

one C–C double bond, or from C8–C12 alkyl or C8–C12 alkenyl having one C–C double bond, or from C10–C12 alkyl or C10–C12 alkenyl having one C–C double bond.

[00198] One or more of the nitrogen atoms indicated in formula (IV) may be protonated to provide a cationic lipidoid of formula (IV).

[00199] In line with formula (IV) above, an alkylene amine unit may be repeated once in an alternating chain such that oligo(alkylene amine) moieties of the type –S–L–L–S– or –L–S–S–L– may result, wherein S represents a shorter ethylene amine unit, and L represents a longer alkylene amine unit. In some cases, a lipidoid of formula (IV) is one wherein no repetition occurs, i.e., wherein p is 1, such that the shorter or longer units do not appear in pairs. The lipidoid of formula (IV) can be a lipidoid of (IVa):

[00200] $R^1-NR^2\{CH_2-(CH_2)_a-NR^3-CH_2-(CH_2)_b-NR\}_m-[CH_2-(CH_2)_a-NR^5]_n-R^6$
(IVa),

[00201] wherein a, b, m, n, and R^1 to R^6 are defined as in formula (IV) and wherein one or more of the nitrogen atoms indicated in formula (IVa) may be protonated to provide a cationic lipidoid;

[00202] In some cases, the lipidoid is a lipidoid of formula (IV). In some cases ‘n’ is 1 in a lipidoid of formula (IV). In some cases, ‘m’ is 1 and n is 1 in a lipidoid of formula (IV). In some cases, the lipidoid of formula (IV) is a lipidoid of formula (IVb):

[00203] $R-NR^2-CH_2-(CH_2)_a-NR^3-CH_2-(CH_2)_b-NR^4-CH_2-(CH_2)_a-NR^5-R^6$ (IVb),

[00204] wherein a, b, and R^1 to R^6 are defined as in formula (IV) wherein one or more of the nitrogen atoms indicated in formula (IVb) may be protonated to provide a cationic lipidoid.

[00205] As regards the length of the alkylene amine units in the lipidoid of formula (IV), (IVa) and (IVb), it will be understood that one of the alternating units needs to be an ethylene amine unit (i.e., either a or b is 1). The other alternating unit can be a propylene amine unit, a butylene amine unit, a pentylene amine unit, or another suitable unit (i.e., the other one of a or b is an integer of 2 to 4). In some cases, a lipidoid of formula (IV) is a lipidoid of formula (IVc):

[00206] $R^1-NR^2-CH_2-CH_2-NR^3-CH_2-CH_2-CH_2-NR^4-CH_2-CH_2-NR^5-R^6$ (IVc)

[00207] wherein R^1 to R^6 are as defined in formula (IV) and wherein one or more of the nitrogen atoms indicated in formula (IVc) can be protonated to provide a cationic lipidoid;

[00208] In some cases, the groups R^1 to R^6 in formula (IV), (IVa), (IVb) and (IVc) are a protecting group for an amino group. Non-limiting examples of protecting groups include t-butoxycarbonyl (Boc), 9-fluorenylmethoxycarbonyl (Fmoc), or carbobenzyloxy (Cbz).

[00209] As far as the groups R^1 to R^6 in formula (IV), (IVa), (IVb) and (IVc) are a receptor ligand, such as the receptor ligands described in Philipp and Wagner in “Gene and Cell

Therapy - Therapeutic Mechanisms and Strategy”, 3rd Edition, Chapter 15, CRC Press, Taylor & Francis Group LLC, Boca Raton 2009. Examples of receptor ligands that target the lung tissue are described in Pfeifer et al. 2010, Ther. Deliv. 1 (1): 133-48. Receptor ligands can include synthetic cyclic or linear peptides such as derived from screening peptide libraries for binding to a particular cell surface structure or particular cell type, cyclic or linear RGD peptides, synthetic or natural carbohydrates such as sialic acid, galactose or mannose or synthetic ligands derived from reacting a carbohydrate for example with a peptide, antibodies specifically recognizing cell surface structures, folic acid, epidermal growth factor and peptides derived thereof, transferrin, anti-transferrin receptor antibodies, nanobodies and antibody fragments, approved drugs that may bind to cell surface molecules (e.g., cell surface receptors), etc.

[00210] As far as the groups R^1 to R^6 in formula (IV), (IVa), (IVb) and (IVc) are a poly(ethylene glycol) chain, the molecular weight of the poly(ethylene glycol) chain can be from about 100 g/mol to 20,000 g/mol, from about 1,000 g/mol to 10,000 g/mol or from about 1,000 g/mol to 5,000 g/mol. In some cases, a molecular weight of the PEG chain can provide a composition with a desired density.

[00211] Multiple lipidoid molecules can be associated with an engineered polyribonucleotide. For example, a composition can comprise 1 engineered polyribonucleotide to 100 lipidoid molecules, 1 engineered polyribonucleotide to 1,000 lipidoid molecules, 10 engineered polyribonucleotide to 1,000 lipidoid molecules, or 100 engineered polyribonucleotide to 10,000 lipidoid molecules. The complex of engineered polyribonucleotide and lipidoid can form a particle. The diameter of the particles may range, e.g., from 10 nanometers to 1,200 nanometers. In some cases the diameter of the particles ranges from 10 nanometers to 500 nanometers. In some cases, the diameters of the particles are from 20 nanometers to 150 nanometers.

Administration to a subject

[00212] Further described herein are methods for the administration of a polynucleotide (e.g., polyribonucleotide, nucleic acid construct, or vector) to a subject. The polyribonucleotide can be provided to the subject via a delivery agent, such as a particle or capsule with an encapsulating agent that encapsulates the polyribonucleotide. The delivery agent can be a therapeutic agent. The subject can be a human, such as a human afflicted with a disease or condition (e.g., primary ciliary dyskinesia (PCD), Kartagener Syndrome or cancer). The delivery agent can be administered to the subject (e.g., self-administration or administration by a third party, such as a healthcare provider) at a given dosage, and the dosage can be increased with

time, decreased with time, or kept constant. The dosage can be changed based on a progression or regression of a disease in the subject, such as a rare disease or a cancer.

[00213] A polyribonucleotide of the disclosure can be formulated with one or more pharmaceutically acceptable carrier(s) to be administered to a subject. In some cases, the polyribonucleotide can be formulated for targeted delivery to a target cell or cell population. In some cases, the polyribonucleotide can be formulated for untargeted delivery to a cell or cell population. The encoded polypeptide product of the polyribonucleotide is then transcribed and it accumulates within the recipient cell.

[00214] A composition can be a combination of any engineered polyribonucleotide described herein with other chemical components, such as carriers, stabilizers, diluents, dispersing agents, suspending agents, thickening agents, and/or excipients. The composition facilitates administration of the compound to an organism. Pharmaceutical compositions can be administered in therapeutically-effective amounts as pharmaceutical compositions by various forms and routes including, for example, intravenous, subcutaneous, intramuscular, oral, rectal, aerosol, parenteral, ophthalmic, pulmonary, transdermal, vaginal, otic, nasal, and topical administration.

[00215] A composition can be administered in a local or systemic manner, for example, via injection of the compound directly into an organ, optionally in a depot or sustained release formulation. Pharmaceutical compositions can be provided in the form of a rapid release formulation, in the form of an extended release formulation, or in the form of an intermediate release formulation. A rapid release form can provide an immediate release. An extended release formulation can provide a controlled release or a sustained delayed release.

[00216] For administration by inhalation, the active compounds can be in a form as an aerosol, a mist, a vapor, a spray, or a powder. Pharmaceutical compositions are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, for example, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, for example, gelatin for use in an inhaler or insufflator can be formulated containing a powder mix of the compounds and a suitable powder base such as lactose or starch.

[00217] The eye comprises several structurally and functionally distinct vascular beds that supply ocular components critical to the maintenance of vision. These beds include the retinal

and choroidal vasculatures, which supply the inner and outer portions of the retina, respectively, and the limbal vasculature located at the periphery of the cornea.

[00218] A pharmaceutical composition comprising an engineered polyribonucleotide can be administered to the eye via any suitable form or route including, for example, topical, oral, systemic, intravitreal, intracameral, subconjunctival, subtenon, retrobulbar, intraocular, posterior juxtасcleral, periocular, subretinal, and suprachoroidal administration. The compositions can be administered by injecting the formulation in any part of the eye including anterior chamber, posterior chamber, vitreous chamber (intravitreal), retina proper, and/or subretinal space. The compositions can also be delivered via a non-invasive method. Non-invasive modes of administering the formulation can include using a needleless injection device. Multiple administration routes can be employed for efficient delivery of the pharmaceutical compositions.

[00219] An engineered polynucleotide of the disclosure can be delivered to any suitable ocular cell including for example, endothelial cells such as vascular endothelial cells, cells of the retina such as retinal pigment epithelium (RPE), corneal cells, fibroblasts, astrocytes, glial cells, pericytes, iris epithelial cells, cells of neural origin, ciliary epithelial cells, mueller cells, muscle cells surrounding and attached to the eye such as cells of the lateral rectus muscle, orbital fat cells, cells of the sclera and episclera, cells of the trabecular meshwork, and connective tissue cells.

[00220] A composition that is disclosed herein, upon administration to a subject, can have a transfection efficiency of at least about 80%, 90%, or 95% by the cell of the subject. In some cases, the transfection efficiency of an encapsulated composition, upon administration to a subject, is at least about 50%, 60%, 70%, 80%, 90%, 95%, 100%, 110%, 120%, 130%, 140%, 150%, 160%, 170%, 180%, 190%, 200%, 225%, 250%, 275%, 300%, 325%, 350%, 375%, 400%, 450%, or 500% relative to an unencapsulated polyribonucleotide. In some situations, transfection efficiency of a composition comprising a modified polyribonucleotide (in some cases also comprising an unmodified polyribonucleotide), upon administration to a subject, is at least about 50%, 60%, 70%, 80%, 90%, 95%, 100%, 110%, 120%, 130%, 140%, 150%, 160%, 170%, 180%, 190%, 200%, 225%, 250%, 275%, 300%, 325%, 350%, 375%, 400%, 450%, or 500% relative to composition solely containing an unmodified polyribonucleotide. The transfection efficiency of a composition can be increased by addition of a carrier, such as a cell penetrating peptide or a cationic coating to the outer layer of the composition. The transfection efficiency of a composition can be modulated by the density of a composition.

[00221] Methods for the preparation of compositions comprising the engineered polyribonucleotides described herein include formulating the compounds with one or more inert,

pharmaceutically-acceptable excipients or carriers to form a solid, semi-solid, or liquid composition. Solid compositions include, for example, powders, tablets, dispersible granules, capsules, cachets, and suppositories. Liquid compositions include, for example, solutions in which a compound is dissolved, emulsions comprising a compound, or a solution containing liposomes, micelles, or nanoparticles comprising a compound as disclosed herein. Semi-solid compositions include, for example, gels, suspensions and creams. The compositions can be in liquid solutions or suspensions, solid forms suitable for solution or suspension in a liquid prior to use, or as emulsions. These compositions can also contain minor amounts of nontoxic, auxiliary substances, such as wetting or emulsifying agents, pH buffering agents, and other pharmaceutically-acceptable additives.

[00222] Non-limiting examples of pharmaceutically-acceptable excipients can be found, for example, in *Remington: The Science and Practice of Pharmacy*, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., *Pharmaceutical Dosage Forms*, Marcel Decker, New York, N.Y., 1980; and *Pharmaceutical Dosage Forms and Drug Delivery Systems*, Seventh Ed. (Lippincott Williams & Wilkins 1999), each of which is incorporated by reference in its entirety.

[00223] A composition comprising a polynucleotide (e.g., polyribonucleotide) can be provided in various dosages. A dose of a polynucleotide, or a polyribonucleotide, can be from about 1 μg to about 1000 μg , about 1 μg to about 500 μg , about 1 μg to about 1000 μg , about 10 μg to about 500 μg , about 20 μg to about 500 μg , about 25 μg to about 500 μg , about 30 μg to about 500 μg , about 40 μg to about 500 μg , about 50 μg to about 500 μg , about 10 μg to about 250 μg , about 20 μg to about 250 μg , about 30 μg to about 250 μg , about 40 μg to about 250 μg , about 50 μg to about 250 μg , about 1 μg to about 200 μg , about 10 μg to about 200 μg , about 20 μg to about 200 μg , about 30 μg to about 200 μg , about 40 μg to about 200 μg , about 50 μg to about 200 μg , about 25 μg to about 50 μg , about 25 μg to about 100 μg , about 25 μg to about 150 μg , about 25 μg to about 200 μg , about 25 μg to about 250 μg , about 25 μg to about 300 μg , about 25 μg to about 350 μg , about 25 μg to about 400 μg , about 25 μg to about 450 μg , about 25 μg to about 500 μg , about 50 μg to about 750 μg , or about 25 μg to about 1000 μg of the engineered polyribonucleotide. In some cases, a dose of a polynucleotide is about 1 mg to about 100 mg, about 1 mg to about 50 mg, about 10 mg to about 50 mg, about 20 mg to about 50 mg, about 25 mg to about 50 mg, about 30 mg to about 50 mg, about 40 mg to about 50 mg, about 50 mg to about 100 mg, about 1 mg to about 25 mg, about 2 mg to about 25 mg, about 3 mg to about 25 mg, about 4 mg to about 25 mg, about 5 mg to about 25 mg, about 1 mg to about 20

mg, about 1 mg to about 20 mg, about 2 mg to about 20 mg, about 3 mg to about 20 mg, about 4 mg to about 20 mg, or about 5 mg to about 20 mg of an engineered polyribonucleotide.

[00224] The percentage of a polyribonucleotide in a formulation (e.g., within an encapsulated agent) can be greater than or equal to 0.25 % polyribonucleotide, 0.5 % polyribonucleotide, 0.75 % polyribonucleotide, 1 % polyribonucleotide, 1.25 % polyribonucleotide, 1.5 % polyribonucleotide, 1.75 % polyribonucleotide, 2 % polyribonucleotide, 2.25 % polyribonucleotide, 2.5 % polyribonucleotide, 2.75 % polyribonucleotide, 3 % polyribonucleotide, 3.25 % polyribonucleotide, 3.5 % polyribonucleotide, 3.75 % polyribonucleotide, 4 % polyribonucleotide, 4.25 % polyribonucleotide, 4.5 % polyribonucleotide, 4.75 % polyribonucleotide, 5 % polyribonucleotide, 5.25 % polyribonucleotide, 5.5 % polyribonucleotide, 5.75 % polyribonucleotide, 6 % polyribonucleotide, 6.25 % polyribonucleotide, 6.5 % polyribonucleotide, 6.75 % polyribonucleotide, 7 % polyribonucleotide, 7.25 % polyribonucleotide, 7.5 % polyribonucleotide, 7.75 % polyribonucleotide, 8 % polyribonucleotide, 8.25 % polyribonucleotide, 8.5 % polyribonucleotide, 8.75 % polyribonucleotide, 9 % polyribonucleotide, 9.25 % polyribonucleotide, 9.5 % polyribonucleotide, 9.75 % polyribonucleotide, 10 % polyribonucleotide, 10.25 % polyribonucleotide, 10.5 % polyribonucleotide, 10.75 % polyribonucleotide, 11 % polyribonucleotide, 11.25 % polyribonucleotide, 11.5 % polyribonucleotide, 11.75 % polyribonucleotide, 12 % polyribonucleotide, 12.25 % polyribonucleotide, 12.5 % polyribonucleotide, 12.75 % polyribonucleotide, 13 % polyribonucleotide, 13.25 % polyribonucleotide, 13.5 % polyribonucleotide, 13.75 % polyribonucleotide, 14 % polyribonucleotide, 14.25 % polyribonucleotide, 14.5 % polyribonucleotide, 14.75 % polyribonucleotide, 15 % polyribonucleotide, 15.25 % polyribonucleotide, 15.5 % polyribonucleotide, 15.75 % polyribonucleotide, 16 % polyribonucleotide, 16.25 % polyribonucleotide, 16.5 % polyribonucleotide, 16.75 % polyribonucleotide, 17 % polyribonucleotide, 17.25 % polyribonucleotide, 17.5 % polyribonucleotide, 17.75 % polyribonucleotide, 18 % polyribonucleotide, 18.25 % polyribonucleotide, 18.5 % polyribonucleotide, 18.75 % polyribonucleotide, 19 % polyribonucleotide, 19.25 % polyribonucleotide, 19.5 % polyribonucleotide, 19.75 % polyribonucleotide, 20 % polyribonucleotide, 20.5 % polyribonucleotide, 21 % polyribonucleotide, 21.5 % polyribonucleotide, 22 % polyribonucleotide, 22.5 % polyribonucleotide, 23 % polyribonucleotide, 23.5 % polyribonucleotide, 24 % polyribonucleotide, 24.5 % polyribonucleotide, or 25% polyribonucleotide by weight. Alternatively, the percentage of the

polyribonucleotide in the formulation (e.g., within an encapsulated agent) can be less than about 25 % polyribonucleotide, 24.5 % polyribonucleotide, 24 % polyribonucleotide, 23.5 % polyribonucleotide, 23 % polyribonucleotide, 22.5 % polyribonucleotide, 22 % polyribonucleotide, 21.5 % polyribonucleotide, 21% polyribonucleotide, 20.5 % polyribonucleotide, 20 % polyribonucleotide, 19.5 % polyribonucleotide, 19% polyribonucleotide, 18.5 % polyribonucleotide, 18 % polyribonucleotide, 17.5 % polyribonucleotide, 17 % polyribonucleotide, 16.5 % polyribonucleotide, 16 % polyribonucleotide, 15.5 % polyribonucleotide, 15% polyribonucleotide, 14.5 % polyribonucleotide, 14 % polyribonucleotide, 13.5 % polyribonucleotide, 13 % polyribonucleotide, 12.5 % polyribonucleotide, 12 % polyribonucleotide, 11.5 % polyribonucleotide, 11 % polyribonucleotide, 10.5 % polyribonucleotide, 10 % polyribonucleotide, 9.5 % polyribonucleotide, 9 % polyribonucleotide, 8.5 % polyribonucleotide, 8 % polyribonucleotide, 7.5 % polyribonucleotide, 7 % polyribonucleotide, 6.5 % polyribonucleotide, 6 % polyribonucleotide, 5.5 % polyribonucleotide, 5 % polyribonucleotide, 4.5 % polyribonucleotide, 4% polyribonucleotide, 3.5 % polyribonucleotide, 3% polyribonucleotide, 2.5 % polyribonucleotide, 2 % polyribonucleotide, 1.5 % polyribonucleotide, 1% polyribonucleotide, 0.5% polyribonucleotide, or 0.1 % polyribonucleotide.

[00225] In some cases, an encapsulated composition of the disclosure can produce a plasma, serum or blood concentration of the polyribonucleotide, pharmaceutical carrier, encapsulating agent, or polymeric material (e.g.: polyethylene glycol or polyethylenimine) in a subject within about 1 second to about 30 minutes, about 1 second to 20 minutes, about 1 second to 10 minutes, about 1 second to 5 minutes, about 1 second to 2 minutes, about 1 second to 1 minute, about 1 second to about 30 seconds, about 30 seconds to 30 minutes, about 30 seconds to 20 minutes, about 30 seconds to 10 minutes, about 30 seconds to 5 minutes, about 30 seconds to 2 minutes, about 30 seconds to about 1 minute, about 1 minute to about 30 minutes, about 1 minute to about 25 minutes, about 1 minute to about 20 minutes, about 1 minute to about 15 minutes, about 1 minute to about 10 minutes, about 5 minutes to about 30 minutes, about 5 minutes to about 25 minutes, about 5 minutes to about 20 minutes, about 5 minutes to about 15 minutes, about 5 minutes to about 10 minutes, about 10 minutes to about 30 minutes, about 10 minutes to about 25 minutes, about 10 minutes to about 20 minutes, or about 10 minutes to about 15 minutes of use of the device. The plasma, serum or blood concentration of the polyribonucleotide, pharmaceutical carrier, encapsulating agent, or polymeric material (e.g.: polyethylene glycol or polyethylenimine) concentration can be a peak concentration or an average concentration.

EXAMPLES

EXAMPLE 1: Production of DNAI1 RNA comprising

[00226] This experiment demonstrates the production of a DNAI1 complementary deoxyribonucleic acid construct.

[00227] Methods: DNAI1 was synthesized at GenScript. pUC57/DNAI1 was digested with HindIII and EcoRI HF restriction enzymes. Moreover, a digested pVAX120 vector and DNAI1 cDNA were gel purified and ligated (the ORF for DNAI1 is codon optimized). Standard *in vitro* translation procedure was used for RNA production utilizing unmodified nucleotides. Capping reaction was carried out using Vaccinia Virus capping system and cap 2'-O-methyl transferase. **FIGURE 1** is an agarose gel illustrating the production of capped and uncapped DNAI1 RNA. Note that in this experiment, the DNAI1 cDNA was ligated into pVAX120 to provide a construct that comprises a poly(A) tail.

EXAMPLE 2: Expression of DNAI Ribonucleic Acid in Mammalian Cells

[00228] This experiment demonstrates the expression (translation) of DNAI1 in HEK-293 cells. **FIGURE 2** is a western blot illustrating the translations of DNAI1 mRNA in 293 cells at 6 hours, 24 hours, and 48 hours post-transfection. For this experiment, 5×10^5 293 cells/well in a 6 well plate were transfected with 2.5 μ g of DNAI1 RNA using 3.75 μ l messenger max transfection reagent. 6, 24, and 48 hours post transfection, cells were scraped from the wells, pelleted, and the pellet was lysed in RIPA buffer. The blot was probed with anti-DNAI1 ab166912 from Abcam. A C-terminal FLAG tagged DNAI1 plasmid DNA was transfected as a control, and the difference in MW between the plasmid and mRNA is likely due to the FLAG tag in the pENTRY vector.

EXAMPLE 3: Formulation of a Composition Comprising an Engineered Polyribonucleotide for the Treatment of Human Subjects Afflicted with Primary Ciliary Dyskinesia.

[00229] Compositions are formulated as follows:

[00230] A nucleic acid construct encoding the DNAI1 gene sequence, NCBI Reference Sequence: NM_012144, is prepared as described in Example 1. Branched polyethylenimine is purchased from Sigma Aldrich™. Linear *in vivo*-jetPEI® (polyethylenimine) is purchased from Polyplus transfection® (Illkirch, France) and used without further purification. Following the manufacturer protocol, jetPEI is diluted in 5% glucose (final concentration) using the sterile 10%

glucose solution provided by the manufacturer and HPLC-grade water purchased from Sigma-Aldrich (St. Louis, MO). After diluting the nucleic acid construct in 5% glucose (final concentration), the RNA and jetPEI solutions are combined/mixed at a ratio of 1:1 with a final N/P ratio of 8. The mRNA is then administered by intranasal instillation. Alternatively, the nucleic acid construct could also be formulated for administration by nebulizing or sniffing with a lipoplex formulation.

EXAMPLE 4: Effects of posttranscriptional polyadenylation reaction times on RNA Quality

[00231] The effect of the post-transcriptional polyadenylation reaction times on RNA quality was tested. Post *in vitro* transcription (IVT) poly-adenylation reaction times are typically 60-90 minutes long and usually provide polyA lengths that are at most about ~200 As. Because mRNA is susceptible to hydrolysis, it often degrades over time during the posttranscriptional poly-adenylation reaction. To maintain an optimal length for the DNAI1 poly A tail and to maximize RNA quality, a nucleic acid construct that encodes dynein axonemal intermediate chain 1 protein or a variant thereof with a poly A tail already included in the template was constructed.

[00232] A summary of the nucleic acid constructs encoding the DNAI1 gene sequence both with and without a poly-A sequence that were used to generate DNAI1 mRNAs are shown below:

TABLE 4					
Nucleic acid constructs encoding the DNAI1 gene sequence	Vector	Enzyme for Linearization	Codon-optimized	Nucleotide composition	Poly(A) post-IVT
DNAI1	pCMV6Entry	Pme I	No	unmodified	yes
DNAI1 (SEQ ID NO: 14)	pVAX	NotI	Yes, GenScript	unmodified	No, Poly(A) sequence in template

[00233] **FIGURE 3** illustrates fragment analyzer data to determine the length of DNAI1 mRNAs produced from the DNAI1-pCMV6Entry plasmid that were post-transcriptionally polyadenylated with reaction times from 0 to 60 min. This demonstrates increasing transcripts lengths with longer polyadenylation reaction times. **FIGURE 4** illustrates fragment analyzer

data to examine the quality of these DNAI1 mRNAs that were post-transcriptionally polyadenylated with reaction times from 0 to 60 min. These results indicate that the RNA undergoes degradation as the poly-adenylation reaction proceeds as demonstrated by the reduction in % peak and increase in pre-peak smear % with longer reaction times. **FIGURE 5** illustrates the length of the poly- A sequence in the DNAI1-pVAX plasmid template as determined by 8% PAGE.

EXAMPLE 5: RNA Production and Quality Control *in vitro*

[00234] The following experiment was conducted to compare the effect of incorporating specific chemically-modified nucleotides, in varying ratios, on translation efficiency in different cell types and immunogenicity.

[00235] The experiments involved: 1) *in vitro* transcription of nucleic acid constructs; 2) *in vitro* capping of the nucleic acid constructs; 3) analysis of the integrity of the transcribed RNAs; 4) immuno-Dot-Blot Assay for dsRNAs; and 5) analysis of the nucleotide composition of the transcribed RNAs. General protocols for *in vitro* transcription (IVT) and capping of the RNAs were followed with a few modifications. IVT reactions for nucleic acid constructs encoding the DNAI1 gene were performed at 37°C lasted for 6 h in the presence of 20 mM MgCl₂ and 7.5 mM of each ribonucleotide.

[00236] **TABLE 5** illustrates various specific chemically-modified nucleotides that were transcribed *in vitro* from a nucleic acid construct that encodes dynein axonemal intermediate chain 1.

TABLE 5						
	Sample	Modified nucleotide composition of <i>in vitro</i> transcription reaction	5' Cap	Vector backbone	5' UTR	3' UTR
2	DNAI1-RNA-002.2a (SEQ ID NO: 14)	Unmodified	Yes	pVAX	vector	vector
3	DNAI1-RNA-003.2a (SEQ ID NO: 14)	50% Ψ	Yes	pVAX	vector	vector
4	DNAI1-RNA-004.2a (SEQ ID NO: 14)	100% Ψ	Yes	pVAX	vector	vector

5	DNAI1-RNA-037.1a (SEQ ID NO: 14)	100% m1Ψ	Yes	pVAX	vector	vector
---	----------------------------------	----------	-----	------	--------	--------

[00237] Results:

[00238] UV measurements

	Sample	Modified nucleotide composition of <i>in vitro</i> transcription reaction	mg/mL	260/280
2	DNAI1-RNA-002.2a	unmodified	0.944	2.22
3	DNAI1-RNA-003.2a	50% Ψ	0.971	2.16
4	DNAI1-RNA-004.2a	100% Ψ	0.940	2.13
5	DNAI1-RNA-037.1a	100% m1Ψ	1.092	1.94

[00239] Template Poly(A) Length - Fragment Analyzer

[00240] Analysis of poly(A) tail length of the DNAI1 nucleic acid construct (SEQ ID NO: 5) used as a template for *in vitro* transcription indicated that the number of A residues was maintained in comparison to the original cloning vector (pVAX-A120). The initial vector contained 120 adenosine nucleotides, while a band between 100 and 150 bp was detected on this nucleic acid construct (**FIGURE 5**). Templates were digested with Eco RI and Not I to remove the poly(A) fragment: 12 non-poly(A) nucleotides are expected to be part of the fragment. G*AATTCtgcag – poly(A) – GC*GGCCGC = 12 nt plus the poly(A) in the EcoRI/NotI generated fragment.

[00241] RNA Smear Analysis - Fragment Analyzer

[00242] For all transcripts generated for DNAI1 a peak around 2,000 nt was detected. Evaluation of the *in vitro* generated transcript on a Fragment Analyzer indicated that capped transcripts maintained good integrity: limited detection of smear content (an indicator of RNA degradation and/or hydrolysis) was observed in repeated experiments. Briefly, 2 μL of 200 ng/μL samples were analyzed on a Fragment Analyzer (DNF-471 Standard Sensitivity RNA Analysis Kit (15nt Lower Marker). Data analysis was conducted using PROSize 2.0 software. The sizing accuracy is approximately within ± 5%; the sizing precision is approximately within 5% CV, the quantification accuracy is approximately within ± 20%; and the quantification precision is approximately 10% CV. **FIGURE 6** illustrates the fragment analyzer data for the *in*

in vitro reaction comprising the canonical nucleotides only, namely: adenosine 5'-triphosphate, guanosine 5'-triphosphate, cytidine 5'-triphosphate, and uridine 5'-triphosphate. **FIGURE 7** illustrates the fragment analyzer data for the *in vitro* reaction comprising 50%/50% mixtures of pseudouridine and uridine 5'-triphosphate. **FIGURE 8** illustrates the fragment analyzer data for the *in vitro* reaction comprising 100% pseudouridine 5'-triphosphate. **FIGURE 9** illustrates the fragment analyzer data for the *in vitro* reaction comprising 100% 1-methyl-pseudouridine 5'-triphosphate. **TABLE 7** summarizes the results of the RNA smear analysis.

Sample	% smear pre-peak	% smear post-peak	% full-length	Length
DNAI1-RNA-002.2a	24.9	3.9	71.2	2366
DNAI1-RNA-003.2a	18.5	2.5	79.0	2338
DNAI1-RNA-004.2a	16.4	3.0	80.6	2298
DNAI1-RNA-037.1a	17.7	0.4	81.9	2338

[00243] Double-stranded RNA content as detected by dot-blot showed reactivity with J2 antibody. **FIGURE 10** illustrates double-stranded RNA content, as detected by dot-blot analysis, of *in vitro* transcribed RNAs from a nucleic acid construct that encodes dynein axonemal intermediate chain 1 as well as the double-stranded RNA content of DNAI1 constructs transcribed with the various specific modifications shown on **TABLE 5**.

[00244] Nucleoside Composition Analysis

[00245] **TABLES 8-10** illustrate the nucleotide composition analysis of *in vitro* transcribed RNAs from a nucleic acid construct that encodes dynein axonemal intermediate chain 1. The various specific nucleotide modifications are shown on **TABLE 5**. **FIGURES 11-13** illustrate the corresponding HPLC chromatograms of individual ribonucleotides obtained after nuclease digestion of the transcripts and subsequent dephosphorylation.

[00246] **TABLE 8** illustrates the nucleotide composition analysis of *in vitro* transcribed RNA from a nucleic acid construct that encodes dynein axonemal intermediate chain 1, transcribed with unmodified nucleotides. **FIGURE 11** illustrates the corresponding HPLC chromatogram.

Nucleotide composition (unmodified)	% abundance exp. [theoretical]
A	28.9 [27.3]
C	26.9 [26.5]

G	26.6 [28.9]
U	17.7 [17.3]

[00247] **TABLE 9** illustrates the nucleotide composition analysis of *in vitro* transcribed RNA from a nucleic acid construct that encodes dynein axonemal intermediate chain 1, transcribed with 50% pseudouridine. **FIGURE 12** illustrates the corresponding HPLC chromatogram. The retention time for the hydrophobic 1-methyl-pseudouridine using identical reverse-phase HPLC conditions averages ca. 9.5 minutes and is well separated from all other ribonucleotides investigated (data not shown).

TABLE 9	
Nucleotide composition	% abundance exp.
A	28.2
C	27.0
G	26.9
U	8.1
Ψ	9.7 (ϵ_{262}) *

* concentration estimated using empirical absorption coefficient ratio of $\epsilon_{262}/\epsilon_{260}=1.001$ (Ψ)

[00248] **TABLE 10** illustrates the nucleotide composition analysis of *in vitro* transcribed RNA from a nucleic acid construct that encodes dynein axonemal intermediate chain 1, transcribed with 100% pseudouridine. **FIGURE 13** illustrates the corresponding HPLC chromatogram.

TABLE 10	
Nucleotide composition	% abundance exp.
A	28.8
C	26.5
G	26.9
Ψ	17.8 (ϵ_{262}) *

* concentration estimated using empirical absorption coefficient ratio of $\epsilon_{262}/\epsilon_{260}=1.001$ (Ψ)

EXAMPLE 6: Translation Efficiency

[00249] The translation efficiency of the aforementioned DNAI1 transcripts was assessed in three cell lines: 1) HEK-293 human embryonic kidney cells; 2) A549 adenocarcinomic human

alveolar basal epithelial cells; and 3) MLE-15 murine lung epithelial cells. Each cell line was transfected in triplicate with each DNAI1 transcript and the resulting cell extracts were analyzed for DNAI1 protein expression with western blotting. Briefly, either 1×10^6 (HEK-293, MLE-15) or 2×10^6 (A549) cells per well were plated 18 hours before transfection in 6-well plates. Cells were transfected with about 100 ng of each RNA using MessengerMax transfection reagent at a RNA: MessengerMax ratio of 1:37.5. Cells were harvested at 6 hours post-transfection and whole cell extracts were prepared in RIPA buffer (50 mM Tris-HCl pH 8, 150 mM NaCl, 1 % Triton X-100, 0.1 % SDS, 0.5 % Sodium taurocholate). 3.5 μ g of total protein from each extract was prepared in 1x LDS sample buffer containing 2.5 % beta-mercaptoethanol and loaded on a 4-12% Bis-Tris SDS-PAGE gel. The gel was then run for 30 min. at 30 V constant voltage followed by 1 hour at 150 V. The proteins were transferred to PVDF membrane for 1 hour at 25 V in 1x NuPAGE transfer buffer containing 10% methanol. Following the transfer, DNAI1 protein was detected by western blot using an anti-DNAI1 antibody and developed using an alkaline phosphatase chemiluminescent substrate. The western blot values were normalized using Sypro Ruby total protein staining as a loading control and are expressed as relative expression to unmodified RNA. Each data point is the mean \pm standard deviation of three biological (transfection) replicates.

[00250] FIGURES 14, 15, and 16 illustrate the expression of DNAI1 protein in HEK-293, A549, and MLE-15 cells respectively. DNAI1 was expressed as a 699 amino acid, 79.3 kDa protein. The pseudouridine (Ψ) containing transcripts express well in all three cell types, with expression levels at or above the unmodified RNA. Similarly, the 1-methylpseudouridine ($m^1\Psi$)-containing transcript produced expression levels at or above the unmodified RNA in A549 and MLE15 cells (Expression in HEK-293 cells was not tested for this transcript). Importantly, the expression levels of each transcript and their relative rankings were similar in each cell line, indicating that there were no cell-type specific effects on DNAI1 translation. FIGURE 17 is a graph illustrating the relative expression of DNAI1 protein in HEK-293, A549, and MLE-15 cells. Western blot signal values were normalized using total protein staining and are plotted as the mean expression \pm std. dev. relative to the unmodified DNAI1 mRNA.

[00251] TABLE 11 is a summary of the relative expression of DNAI1 protein in each of the aforementioned cell lines.

TABLE 11				
Sample	Modified Nucleotide	HEK-293 Cells Relative Expression (% \pm)	A549 Cells Relative Expression (% \pm)	MLE15 Cells Relative Expression (% \pm)

	Composition	Std. Dev).	Std. Dev).	Std. Dev).
DNAI1-RNA-002.2a	Unmodified	100 ± 9.57	100 ± 11.09	100 ± 5.35
DNAI1-RNA-003.2a	50% Ψ	152.21 ± 31.21	120.36 ± 12.40	125.18 ± 35.84
DNAI1-RNA-004.2a	100% Ψ	97.52 ± 23.12	127.57 ± 12.90	100.26 ± 15.71
DNAI1-RNA-037.3a	100% m1Ψ	n.d.	126 ± 3	164 ± 17

[00252] **EXAMPLE 7: Immunogenicity of Nucleic Acid Constructs Encoding Human DNAI1 *in vitro***

[00253] The immunogenicity of the aforementioned transcripts was tested in two cell lines namely, A549 adenocarcinomic human alveolar basal epithelial cells and HepG2 human liver carcinoma cells, by measuring cytokine production. Production of IL-6 in response to the transcripts was measured in A549 cells, while production of IP-10 was measured in HepG2 cells. Each cell line was transfected in triplicate with a titration of each RNA. Briefly, either 20,000 (A549) or 40,000 (HepG2) cells per well were plated 24 hours prior to transfection in 96 well plates. The cells were then transfected with a titration of each transcript: From 250 ng to 7 ng per well for unmodified, 50% Ψ, and 100% Ψ transcripts; and from 1000 ng to 32 ng per well for the 100% m1Ψ mRNA using MessengerMax reagent at a RNA: MessengerMax ratio of 1:1.5.

[00254] Culture supernatants were harvested at 18 hours post-transfection. Cell viability was measured immediately following supernatant removal using the CellTiter-Glo assay kit (Promega) which measures ATP levels as an indication of metabolically active cells. For IL-6 detection, A549 cell culture supernatants were diluted 1:20 in assay buffer and IL-6 levels were measured using the IL-6 High Sensitivity Human ELISA kit (Abcam ab46042). IP-10 was detected in undiluted HepG2 cell culture supernatants using the Human IP-10 ELISA Kit SimpleStep (Abcam ab173194).

[00255] **FIGURES 18 and 19** illustrate the induction of IL-6 in A549 cells treated with the DNAI1 transcripts. For the assay shown in **FIGURE 18**, cells were exposed to the RNA-MessengerMax complexes for 18 hrs, while for the assay shown in **FIGURE 19** the RNA-MessengerMax complexes were removed at 2 hrs post-transfection. In both cases the cell culture supernatants were harvested at 18 hrs for detection of IL-6 by ELISA. **FIGURES 20 and 21** illustrate cell viability for the assay shown in **FIGURES 18 and 19** as measured using the CellTiter-Glo assay. **FIGURE 22** illustrates induction of IP-10 in HepG2 cells by DNAI1 transcripts. For this assay, cells were exposed to the RNA-MessengerMax complexes for 18 hrs. IP-10 expression induced by various amounts of each DNAI1 mRNA was then measured by ELISA. **FIGURE 23** illustrates cell viability for the assay shown in **FIGURE 22** as measured using the CellTiter-Glo assay.

[00256] **EXAMPLE 8: Translation of DNAI1 mRNA in HEK293 Cells**

[00257] **FIGURE 24** illustrates the peak expression of a nucleic acid encoding dynein axonemal intermediate chain 1 (DNAI1), or nucleic acid controls, in HEK293 cells. As shown in **FIGURE 24**, in HEK293 cells, translation of DNAI1 nucleic acid construct in HEK293 cells peaks at 6 hours but is still present at 48 hours.

[00258] **EXAMPLE 9: Expression of DNAI1 Protein in Undifferentiated and Fully-Differentiated Human Airway Epithelial Cells (HAECs) and Mouse Tracheal Epithelial Cells (MTECs) Following Administration of Lipoplex-Formulated 100% m1Ψ-containing DNAI1 mRNA.**

[00259] Expression of DNAI1 protein in primary human airway epithelial cells and mouse tracheal epithelial cells following treatment with lipoplex-formulated DNAI1-HA mRNA was assessed by western blot. The 100% m1Ψ-containing transcript used for this experiment was produced from a DNAI1 alternate codon usage template (SEQ ID NO 15) that contains an HA epitope tag. Primary human epithelial cells were maintained in submerged liquid culture for undifferentiated cultures or maintained at an air-liquid interface and allowed to differentiate for ~ 3 weeks into fully-differentiated ciliated epithelia. Next 12 or 24 μg of lipoplex-formulated DNAI1-HA mRNA was applied to the apical side of the fully-differentiated cultures or directly to the undifferentiated liquid cultures. Cells were treated either once or once per day for two consecutive days. Cells were then harvested at 24 or 48 hrs after the final treatment and whole cell extracts were prepared in RIPA buffer (50 mM Tris-HCl pH 8, 150 mM NaCl, 1 % Triton X-100, 0.1 % SDS, 0.5 % Sodium taurocholate). Total protein from each extract was separated on a 4-12% Bis-Tris SDS-PAGE gel and transferred to PVDF membrane. DNAI1-HA protein

was detected by western blot using an anti-HA antibody and developed using an enhanced chemiluminescent substrate. As shown in **FIGURE 25**, DNAI1-HA protein was expressed at high levels in both the undifferentiated and fully-differentiated, ciliated human airway epithelial cells and in mouse tracheal epithelial cells.

[00260] **EXAMPLE 10:** Altered Nucleotide Usage in Coding Regions Increases mRNA Stability for Transcript Therapy.

[00261] Altered nucleotide usage schemes aiming to reduce the number of more reactive dinucleotides within codons as well as across codons of modified mRNAs partially alleviate limitations imposed by the inherent chemical instability of RNA. At the same time, lowering the U-content in RNA transcripts renders them less immunogenic. Traditional codon optimization (CO) can be performed prior to (+) removal of reactive dinucleotide and (+) U-reduction in general yielding ORFs that are termed CO⁺⁺.

[00262] **FIGURE 26** illustrates an overall quality improvement in DNAI1 expressing a polyribonucleotide of SEQ ID NO 15 (**B**) as compared to a polyribonucleotide of SEQ ID NO 14 (**A**). The overall quality improvement is judged by the increasing main RNA peak % of the fragment analyzer traces in the polyribonucleotide engineered with the altered codon usage strategy. Furthermore, DNAI1 mRNA featuring the CO⁺⁺-optimized open reading frame, i.e., altered codon usage, show an improvement in translation efficiency in transfected A549 cells when compared with transcripts that have been traditionally optimized (CO) (**see FIGURE 27**). Here, 1.25×10^6 cells per well were plated 18 hours before transfection in 6-well plates. Cells were transfected with about 100 ng of each RNA using MessengerMax transfection reagent at a RNA: MessengerMax ratio of 1:12 and harvested 6 h post transfection. Western blotting using an anti-DNAI1 antibody revealed DNAI1 protein expression as a 699 amino acid, 79.3 kDa protein. Relative translation efficiencies are indicated as the mean of three biological (transfection) replicates.

[00263] Altered reactivity with J2 antibody sensing double-stranded RNA content was observed as shown in **FIGURE 28**. Based on comparison with known concentrations of poly-IC, the dsRNA content of DNAI1-coding mRNA featuring the CO⁺⁺ ORF averaged 39 ng while RNAs with a CO ORF were estimated to contain 68 ng of dsRNA contaminants when 200 ng of *in vitro* transcribed mRNA were dotted.

[00264] **EXAMPLE 11:** HPLC-purification of unmodified and 100% m¹Ψ-containing DNAI1 mRNA.

[00265] Reverse phase high-performance liquid chromatography (HPLC) of DNAI1 mRNA was employed to purify full-length RNA and remove contaminants such as long dsRNA generated during the *in vitro* transcription using T7 RNA polymerase. Fractionation and purification results obtained using a non-porous RNASep C18 semi prep (100 mm x 21.2 mm, column volume (CV) ca. 2.4 mL) column together with mobile phases containing triethylammonium acetate (TEAA) as an ion-pairing agent and increasing Acetonitrile content are shown in **FIGURE 29**. Judged by fragment analyzer evaluation of purified fractions, an overall quality improvement and full-length RNA enrichment using semi-prep RNASep column was observed. This quality improvement was achieved for both, unmodified (**A, B**) and 100% m¹Ψ-containing DNAI1 mRNA species (**C, D**).

[00266] As shown in **FIGURE 30**, a moderate improvement of translation activity was observed for fractions enriched in full-length, unmodified mRNA transcripts in A549 cells (**A** and **B**). Here, 1x10⁶ A549 cells per well were plated 18 hours before transfection in 6-well plates. Cells were transfected with about 100 ng of each RNA using MessengerMax transfection reagent at a RNA:MessengerMax ratio of 1:12 and harvested 6 h post transfection. Western blotting using a 1:2000 rabbit-anti-DNAI1 (AbCam ab166912, rabbit monoclonal to recombinant DNAI1 fragment anti-DNAI1) antibody revealed DNAI1 protein expression as a 699 amino acid, 79.3 kDa protein (**C**). Relative translation efficiencies are indicated as the mean ± standard deviation of three biological (transfection) replicates.

[00267] Importantly, HPLC readily removes late-eluting dot-blot reactive species at semi-prep scale. Detectable double-stranded RNA content reacting with J2 antibody was observed exclusively within the late-eluting fraction F7 and the unpurified control transcript while undetectable in all other HPLC-purified DNAI1 mRNA fractions F1 through F6 (**D**). The immunogenicity of unmodified, HPLC-purified transcripts was further tested in A549 cells by monitoring production of IL-6 in response to the transfected mRNA. Each cell line was transfected in triplicate with a titration of each RNA. Briefly, 20,000 cells per well were plated 18 hours prior to transfection in 96 well plates. The cells were then transfected with a titration of each transcript, from 250 ng to 7 ng per well, using MessengerMax reagent at a RNA:MessengerMax ratio of 1:1.5. A reduction of IL-6 response was observed for the HPLC-purified fraction F3 producing the highest relative DNAI1 protein level (unpurified reference DNAI1 mRNA: (727±109 pg/mL) > F3 (73±30 pg/mL) for cells transfected with 125 ng RNA) (**E**).

[00268] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided

by way of example only. It is not intended that the invention be limited by the specific examples provided within the specification. While the invention has been described with reference to the aforementioned specification, the descriptions and illustrations of the embodiments herein are not meant to be construed in a limiting sense. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. Furthermore, it shall be understood that all aspects of the invention are not limited to the specific depictions, configurations or relative proportions set forth herein which depend upon a variety of conditions and variables. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is therefore contemplated that the invention shall also cover any such alternatives, modifications, variations or equivalents. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

CLAIMSWHAT IS CLAIMED IS:

1. A method for treating a subject having or suspected of having primary ciliary dyskinesia, the method comprising administering to said subject in need thereof a polynucleotide comprising nucleotides 1-1,000 of SEQ ID NO: 15, which polynucleotide includes codons that provide for heterologous expression of a dynein axonemal intermediate chain 1 (DNAI1) protein within ciliated cells of said subject.
2. The method of claim 1, wherein said subject is a human.
3. The method of claim 1 or 2, wherein said ciliated cells are in a lung of said subject.
4. The method of any one of claims 1-3, wherein said ciliated cells are ciliated epithelial cells.
5. The method of claim 4, wherein said ciliated epithelial cells are ciliated airway epithelial cells.
6. The method of claim 4 or 5, wherein said ciliated epithelial cells are undifferentiated.
7. The method of claim 4 or 5, wherein said ciliated epithelial cells are differentiated.
8. The method of any one of claims 1-7, wherein said polynucleotide is a messenger ribonucleic acid (mRNA).
9. The method of any one of claims 1-8, wherein fewer than 15% of nucleotides within said polynucleotide are nucleotide analogues.
10. The method of any one of claims 1-9, wherein substantially all nucleotides replacing uridine within said polynucleotide are nucleotide analogues.

11. The method of any one of claims 1-10, wherein said polynucleotide comprises 1-methylpseudouridine.
12. The method of any one of claims 1-11, wherein said polynucleotide is formulated for administration to said subject in a formulation comprising a cationic lipid, a fusogenic lipid, a cholesterol, a polyethylene glycol (PEG) lipid, or a combination thereof.
13. The method of any one of claims 1-12, wherein said polynucleotide is formulated for administration to said subject in a formulation using a cationic lipid or a cationic polymer.
14. The method of any one of claims 1-13, wherein said polynucleotide is formulated for administration to said subject in a formulation using a nanoparticle or a nanocapsule.
15. The method of any one of claims 1-14, wherein said polynucleotide further comprises a 3' or 5' noncoding region, wherein said 3' or 5' noncoding region enhances the expression of said dynein axonemal intermediate chain 1 polypeptide within cells of said subject.
16. The method of claim 15, wherein said 3' noncoding region comprises a poly adenosine tail.
17. The method of claim 16, wherein said poly adenosine tail improves the half-life of said polynucleotide.
18. The method of claim 16 or 17, wherein the length of said poly adenosine tail is at most 200 adenosines.
19. The method of any one of claims 1-18, wherein said polynucleotide further comprises a 5' cap structure.
20. A pharmaceutical formulation comprising a polynucleotide comprising nucleotides 1-1,000 of SEQ ID NO: 15.
21. The pharmaceutical formulation of claim 20, wherein substantially all nucleotides replacing uridine within said polynucleotide are nucleotide analogues.

22. The pharmaceutical formulation of claim 20 or 21, wherein said polynucleotide comprises 1-methylpseudouridine.
23. The pharmaceutical formulation of any one of claims 20-22, wherein fewer than 15% of nucleotides within said polynucleotide are nucleotide analogues.
24. The pharmaceutical formulation of any one of claims 20-23, wherein said pharmaceutical formulation is formulated for administration to a subject.
25. The pharmaceutical formulation of any one of claims 20-24, wherein said polynucleotide is a messenger ribonucleic acid (mRNA).
26. The pharmaceutical formulation of any one of claims 20-25, wherein said pharmaceutical formulation further comprises one or more members selected from the group consisting of a cationic lipid, a fusogenic lipid, a cholesterol, a polyethylene glycol (PEG) lipid, or a combination thereof.
27. The pharmaceutical formulation of any one of claims 20-26, wherein said pharmaceutical formulation comprises a cationic lipid or a cationic polymer.
28. The pharmaceutical formulation of any one of claims 20-27, wherein said pharmaceutical formulation is formulated in a nanoparticle or nanocapsule.
29. The pharmaceutical formulation of any one of claims 20-28, wherein said polynucleotide further comprises a 3' or 5' noncoding region, wherein said 3' or 5' noncoding region enhances expression of said polynucleotide within a cell of said subject.
30. The pharmaceutical formulation of claim 29, wherein said 3' noncoding region comprises a poly adenosine tail.
31. The pharmaceutical formulation of claim 30, wherein said poly adenosine tail enhances a half-life of said polynucleotide.

32. The pharmaceutical formulation of claim 30 or 31, wherein a length of said poly adenosine tail is at most 200 adenosines.

33. The pharmaceutical formulation of any one of claims 20-32, wherein said polynucleotide further comprises a 5' cap structure.




FIGURE 1




FIGURE 2





FIGURE 3



	% smear pre-peak	% smear post-peak	% peak (estimated)	Length (nt)
0 min	24.2	3.2	72.6	2312
10 min	30.0	3.1	66.9	2325
20 min	32.9	1.6	65.5	2364
30 min	38.1	2.9	59.0	2390
60 min	43.4	1.5	55.1	2416

FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8




FIGURE 9




FIGURE 10




FIGURE 11




FIGURE 12




FIGURE 13

HEK-293




FIGURE 14




FIGURE 15

MLE-15




FIGURE 16




FIGURE 17




FIGURE 18




FIGURE 19




FIGURE 20




FIGURE 21




FIGURE 22




FIGURE 23




FIGURE 24




FIGURE 25




FIGURE 26




FIGURE 27




FIGURE 28




FIGURE 29




FIGURE 30

Sequence Listing

2017271665 01 Nov 2023

1	Sequence Listing Information	
1-1	File Name	58530-708.681_Sequence Listing.xml
1-2	DTD Version	V1_3
1-3	Software Name	WIPO Sequence
1-4	Software Version	2.3.0
1-5	Production Date	2023-10-05
1-6	Original free text language code	
1-7	Non English free text language code	
2	General Information	
2-1	Current application: IP Office	
2-2	Current application: Application number	
2-3	Current application: Filing date	
2-4	Current application: Applicant file reference	58530-708.681
2-5	Earliest priority application: IP Office	US
2-6	Earliest priority application: Application number	62/342,784
2-7	Earliest priority application: Filing date	2016-05-27
2-8en	Applicant name	Transcriptx, Inc.
2-8	Applicant name: Name Latin	
2-9en	Inventor name	LOCKHART, David, J.
2-9	Inventor name: Name Latin	
2-10en	Invention title	TREATMENT OF PRIMARY CILIARY DYSKINESIA WITH SYNTHETIC MESSENGER RNA
2-11	Sequence Total Quantity	20

3-1	Sequences	
3-1-1	Sequence Number [ID]	1
3-1-2	Molecule Type	DNA
3-1-3	Length	72
3-1-4	Features	misc_feature 1..72
	Location/Qualifiers	note=Description of Artificial Sequence: Synthetic oligonucleotide source 1..72 mol_type=other DNA organism=synthetic construct
	NonEnglishQualifier Value	
3-1-5	Residues	gggagacata aaccctggcg cgctcgcggc cgggactct tctggcccc acagactcag 60 agagaagcca cc 72
3-2	Sequences	
3-2-1	Sequence Number [ID]	2
3-2-2	Molecule Type	DNA
3-2-3	Length	72
3-2-4	Features	misc_feature 1..72
	Location/Qualifiers	note=Description of Artificial Sequence: Synthetic oligonucleotide source 1..72 mol_type=other DNA organism=synthetic construct
	NonEnglishQualifier Value	
3-2-5	Residues	gggagacata aaccctggcg cgctcgcggg cgggactct tctggcccc acagactcag 60 agagaagcca cc 72
3-3	Sequences	
3-3-1	Sequence Number [ID]	3
3-3-2	Molecule Type	DNA
3-3-3	Length	43
3-3-4	Features	misc_feature 1..43
	Location/Qualifiers	note=Description of Artificial Sequence: Synthetic oligonucleotide source 1..43 mol_type=other DNA organism=synthetic construct
	NonEnglishQualifier Value	
3-3-5	Residues	gggagactct tctggcccc acagactcag agagaacgcc acc 43
3-4	Sequences	
3-4-1	Sequence Number [ID]	4
3-4-2	Molecule Type	DNA
3-4-3	Length	511
3-4-4	Features	misc_feature 1..511
	Location/Qualifiers	note=Description of Artificial Sequence: Synthetic polynucleotide source 1..511 mol_type=other DNA organism=synthetic construct
	NonEnglishQualifier Value	
3-4-5	Residues	gttattttcc accatattgc cgtcttttgg caatgtgagg gcccgaaac ctggcctgt 60 cttcttgacg agcattccta ggggtcttcc cctctcgcgc aaaggaatgc aaggtctgtt 120 gaatgtcgtg aaggaagcag ttcctctgga agcttcttga agacaaacaa cgtctgtagc 180 gaccctttgc aggcagcggg acccccacc tggcgacagg tgcctctgcg gccaaaagcc 240 acgtgtataa gatacacctg caaaggcggc acaaccaccg tgccacgttg tgagttggat 300 agttgtggaa agagtcaaat ggctctcctc aagcgtattc aacaaggggc tgaaggatgc 360 ccagaaggta cccattgta tgggatctga tctggggcct cgggtgcacat gctttacgtg 420 tgtttagtcg aggttaaaaa acgtctaggc cccccgaacc acggggacgt ggttttccct 480 tgaaaaacac gatgataata tggccacaac c 511
3-5	Sequences	
3-5-1	Sequence Number [ID]	5
3-5-2	Molecule Type	DNA
3-5-3	Length	143
3-5-4	Features	misc_feature 1..143
	Location/Qualifiers	note=Description of Artificial Sequence: Synthetic polynucleotide source 1..143 mol_type=other DNA organism=synthetic construct
	NonEnglishQualifier Value	
3-5-5	Residues	aaataacaaa tctcaacaca acatatacaa aacaaacgaa tctcaagcaa tcaagcattc 60 tacttctatt gcagcaattt aaatcatttc ttttaaagca aaagcaattt tctgaaaatt 120 ttcaccattt acgaacgata gca 143
3-6	Sequences	
3-6-1	Sequence Number [ID]	6
3-6-2	Molecule Type	DNA

3-6-3	Length	47
3-6-4	Features Location/Qualifiers	misc_feature 1..47 note=Description of Artificial Sequence: Synthetic oligonucleotide source 1..47 mol_type=other DNA organism=synthetic construct
3-6-5	NonEnglishQualifier Value Residues	gggagacaag agagaaaaga agagcaagaa gaaatataag agccacc 47
3-7	Sequences	
3-7-1	Sequence Number [ID]	7
3-7-2	Molecule Type	DNA
3-7-3	Length	66
3-7-4	Features Location/Qualifiers	misc_feature 1..66 note=Description of Artificial Sequence: Synthetic oligonucleotide source 1..66 mol_type=other DNA organism=synthetic construct
3-7-5	NonEnglishQualifier Value Residues	gggagaccca agctggctag cgtttaaact taagcttggc aatccggtac tgttggttaa 60 gccacc 66
3-8	Sequences	
3-8-1	Sequence Number [ID]	8
3-8-2	Molecule Type	DNA
3-8-3	Length	291
3-8-4	Features Location/Qualifiers	misc_feature 1..291 note=Description of Artificial Sequence: Synthetic polynucleotide source 1..291 mol_type=other DNA organism=synthetic construct
3-8-5	NonEnglishQualifier Value Residues	gggagaccca agctggctag cgtttaaact taagctttcc tttccgggcc ggctgggccc 60 gccgaagcgc ctgcgccttg gctgctggtc ggttgctggg taaccgcgctc agggagttag 120 attctatcct gcaagggcac ggggaccac aacgacggct gtcctaaag aaccgttgcg 180 actggtaact gaagtgggag agagtccaga tttcttgtgt gtggccaagg agacggacaa 240 actttttgtc ttcagacgag ggagcgtttt gtaggctctc caggggttga g 291
3-9	Sequences	
3-9-1	Sequence Number [ID]	9
3-9-2	Molecule Type	DNA
3-9-3	Length	186
3-9-4	Features Location/Qualifiers	misc_feature 1..186 note=Description of Artificial Sequence: Synthetic polynucleotide source 1..186 mol_type=other DNA organism=synthetic construct
3-9-5	NonEnglishQualifier Value Residues	ggattgtgtc cgtaatcaca cgtgggtgctg acgataacgc atagtgtttt tccctccact 60 taaatcgaag ggttgtgtct tggatcgcgc gggtaaatg tatatggttc atatacatcc 120 gcaggcacgt aataaagcga ggggttcgaa tccccccggt acccccggtg ggggccatt 180 gtcttc 186
3-10	Sequences	
3-10-1	Sequence Number [ID]	10
3-10-2	Molecule Type	DNA
3-10-3	Length	114
3-10-4	Features Location/Qualifiers	misc_feature 1..114 note=Description of Artificial Sequence: Synthetic polynucleotide source 1..114 mol_type=other DNA organism=synthetic construct
3-10-5	NonEnglishQualifier Value Residues	tcagtaggt catgaaggtt tttcttttcc tgagaaaaca acacgtattg tttctcagg 60 tttgctttt tggcctttt ctgacttaa aaaaaaaaa gcaaaattgt ctcc 114
3-11	Sequences	
3-11-1	Sequence Number [ID]	11
3-11-2	Molecule Type	DNA
3-11-3	Length	112
3-11-4	Features Location/Qualifiers	misc_feature 1..112 note=Description of Artificial Sequence: Synthetic polynucleotide source 1..112 mol_type=other DNA

3-11-5	NonEnglishQualifier Value Residues	organism=synthetic construct tcagtagggt tgtaaagggt tttcttttcc tgagaaaaca accttttggt ttctcaggtt 60 ttgctttttg gcctttccct agctttaaaa aaaaaaaagc aaaattgtct tc 112
3-12	Sequences	
3-12-1	Sequence Number [ID]	12
3-12-2	Molecule Type	DNA
3-12-3	Length	68
3-12-4	Features	misc_feature 1..68
	Location/Qualifiers	note=Description of Artificial Sequence: Synthetic oligonucleotide source 1..68 mol_type=other DNA organism=synthetic construct
3-12-5	NonEnglishQualifier Value Residues	gaagtggcgg ttcgcccgga ggttccatcg tatccaaaag gctcttttca gagccacca 60 ttgtcttc 68
3-13	Sequences	
3-13-1	Sequence Number [ID]	13
3-13-2	Molecule Type	DNA
3-13-3	Length	239
3-13-4	Features	misc_feature 1..239
	Location/Qualifiers	note=Description of Artificial Sequence: Synthetic polynucleotide source 1..239 mol_type=other DNA organism=synthetic construct
3-13-5	NonEnglishQualifier Value Residues	ggggctggcc tcagtctctg tcccctcgt tgaatacagt actcctaggg cttgaccctg 60 gtaccagcc cagccttagc acccagcatg tgacccact cctgatcagg tcccagcctc 120 ttcccttctt gttctgttcc ttaaggctcc agcaccttac cccaggactt ggtcttcaac 180 caccattacc cctctaactt tgcacaaaata aacctgtgta gaaaccacc ccaaaaaa 239
3-14	Sequences	
3-14-1	Sequence Number [ID]	14
3-14-2	Molecule Type	DNA
3-14-3	Length	2100
3-14-4	Features	misc_feature 1..2100
	Location/Qualifiers	note=Description of Artificial Sequence: Synthetic polynucleotide source 1..2100 mol_type=other DNA organism=synthetic construct
3-14-5	NonEnglishQualifier Value Residues	atgatcccag caagcgcca ggcaccacac aagcagccc acaagcagag catctccatc 60 ggcaggggca caaggaagag ggcagaggat agcggaaacc aagtgggaga gggaaacagac 120 gagtgggac agtccaaggc aaccgtgcgc ccacctgacc agctggagct gacagatgcc 180 gagctgaagg aggagttcac caggatcctg acagccaaca atccacacgc ccccagaac 240 atcgtgcgct actctttcaa ggagggcaca tataagccaa tcggctttgt gaaccagctg 300 gccgtgcact ataoccaaagt gggcaatctg atccccaagg actccgatga gggccggaga 360 cagcactaca gggacgagct ggtggcagga tcccaggagt ctgtgaaagt gatctctgag 420 accggcaatc tggaggagga cgaggagcca aaggagctgg agaccgagcc aggaagccag 480 acagatgtgc ctgcagcagg agcagcagag aaggtgaccg agggagagct gatgacacct 540 aagcagccaa aggagcggaa gctgaccaac cagttcaatt tttccgagag agcctctcag 600 acatacaaca atccagtgcg ggacagagag tgccagaccg agccaccccc tagaaccaac 660 ttttccgcca cagccaatca gtgggagatc tacgatgctt atgtggagga gctggagaag 720 caggagaaga ccaaggagaa ggagaaggcc aagacacccc tggccaagaa gtcccgcaag 780 atggccatgc ggaagctgac cagcatggag tcccagacag acgatctgat caagctgtct 840 caggccgcca agatcatgga gagaatggtg aaccagaata cctatgacga tatcgcccag 900 gacttcaagt actatgacga tgcagcagac gagtacaggg atcaagtggg cacactgctg 960 cctctgtgga agtttcagaa cgataaggcc aagaggctga gcgtgaccgc cctgtgctgg 1020 aatccaaagt acagggacct gttcgcagtg ggatacggat cttatgactt catgaagcag 1080 agcagaggca tgctgctgct gtattccctg aagaaccctt ctttccctga gtacatgttt 1140 agctccaatt ccggcgtgat gtgcctggac atccactgg atcaccoccta cctggtggcc 1200 gtgggccact atgacggcaa cgtggccatc tacaatctga agaagcctca ctctcagccc 1260 agcttctgtt ctagcgccaa gagcggcaag cactccgatc ccgtgtggca ggtgaagtgg 1320 cagaaggacg atatggacca gaacctgaat ttcttttccg tctcctctga tggcaggatc 1380 gtgtcttgga cctggtgaa gcgcaagctg gtgcacatcg agctgatcaa gctgaagggtg 1440 gagggcagca ccacagaggt gccagaggga ctgcagctgc acccagtggtg atgcccagca 1500 gccttcgact ttcacaagga gatcgattat atgttctctg tgggcaccga ggagggcaag 1560 atctacaagt gttctaagag ctatagctcc cagtttctgg acacatatga tgcccacaac 1620 atgagcgtgg ataccgtgtc ctggaatcct taccacacaa aggtgttcat gagctgctct 1680 agcagactgga ccgtgaagat ctgggatcac accatcaaga cacctatggt tatctatgac 1740 ctgaactccg ccgtgggcca tgtggcatgg gcaccatact cctctacagt gtcgcagca 1800 gtgaccacag acggcaaggc acacatcttt gatctggcca tcaacaagta ctaggcacatc 1860 tgtaatcagc ccgtggcgc caagaagaac aggtgaccac acgtgcagtt caatctgatc 1920

		caccctatca tcatcgtggg cgacgatcgg ggccacatca tctctctgaa gctgagcccc 1980 aacctgagaa agatgcctaa ggagaagaag ggacaggagg tgcagaaggg accagcagtg 2040 gagatcgcaa agctggacaa gctgctgaat ctggtgctgg aggtgaagat caagacctga 2100
3-15	Sequences	
3-15-1	Sequence Number [ID]	15
3-15-2	Molecule Type	DNA
3-15-3	Length	2100
3-15-4	Features	misc_feature 1..2100
	Location/Qualifiers	note=Description of Artificial Sequence: Synthetic polynucleotide source 1..2100 mol_type=other DNA organism=synthetic construct
	NonEnglishQualifier Value	
3-15-5	Residues	atgatcccag caagcgccaa ggcaccacac aagcagcccc acaagcagag catcagcatc 60 ggcaggggca caaggaagag ggacgaggac agcggaaaccg aagtgggaga gggaaacagac 120 gagtggggcac agagcaaggc aaccgctgctg ccaccctgacc agctggagct gacagatgcc 180 gagctgaagg aggagttcac caggatcctg acagccaaca atccacacgc cccccagaa 240 atcgtgctgct acagcttcaa ggagggcaca tacaagccaa tccgctctctg gaaccagctg 300 gccgtgact acaccctgctg gggcaactctg atcccccaagg actccgatga gggccggaga 360 cagcactaca gggacgagct ggtggcagga tcccaggagt ctgtgaaagt gatctctgag 420 accggcaatc tggaggagga cgaggagcca aaggagctgg agaccgagcc aggaagccag 480 acagactgctg ccgagcagag agcagcagag aaggtgaccg aggaggagct gatgacaccc 540 aagcagccaa aggagcggaa gctgaccaac cagttcaact tctccgagag agcctctcag 600 acatacaaca atccagctgct ggacagagag tgccagacct agccaccccc cagaaccaac 660 ttctccgcca cagccaatca gtggggagatc tacgatgctt acgtggagga gctggagaag 720 caggagaaga ccaaggagaa ggagaaggcc aagacacccg tggccaagaa gtccggcaag 780 atggccatgc ggaagctgac cagcatggag tcccagacag acgatctgat caagctgtct 840 caggccgcca agatcatgga gagaatggtg aaccagaaca cctacgacga catcgccag 900 gacttcaagt actacgacga cgcagcagag gactacaggg accaagtggg cacactgctg 960 ccctgtgga agttccagaa cgacaaggcc aagaggctga gcgtgaccgc cctgtgctgg 1020 aaccctgctg acagggacct gttcgcagtg ggatacggaa gctacgactt catgaagcag 1080 agcagaggca tgcgtgctgct gtacagcctg aagaaccccc gcttccccga gtacatgttc 1140 agcagcaaca gcggcgtgat gtgcctggac atcccactgg accaccctta cctgtggtggc 1200 gtgggcaact acgacggcaa cgtggccatc tacaacctga agaagcccca cagccagccc 1260 agcttctgca gcagcgccaa gagcggcaag cacagcgacc ccggtgagga ggtgaaagtgg 1320 cagaaggcag acatggacca gaacctgaac ttcttcagcg tgagcagcga cggcaggatc 1380 gtgagctgga ccctggtgaa gcgcaagctg gtgcacatcg acgtgatcaa gctgaaggtg 1440 gagggcagca ccacagaggt gccagaggga ctgcagctgc acccagtggt atgcccagca 1500 gccttcgact tcacaagga gatcgactac atgttctctg tgggcaccga ggaggcaag 1560 atctacaagt gcagcaagag ctacagcagc cagttctctg acacatacga cccccacaac 1620 atgagcgtgg acaccgtgag ctggaacccc taccacaca aggtgttcat gagctgcagc 1680 agcagctgga ccgtgaagat ctgggaccac accatcaaga caccctggtt catctacgac 1740 ctgaacagcg ccgtgggca cgtggcagtg gcaccataca gcagcagagt gttcgcagca 1800 gtgaccacag acggcaaggc acacatcttc gacctggcca tcaacaagta cgaggccatc 1860 tgcaaccagc ccgtggccgc caagaagaac aggctgacct acgtgcagtt caacctgatc 1920 caccctatca tcatcgtggg cgacgaccgg ggccacatca tcagcctgaa gctgagcccc 1980 aacctgagaa agatgcccga ggagaagaag ggacaggagg tgcagaaggg accagcagtg 2040 gagatcgcaa agctggacaa gctgctgaac ctggtgctgg aggtgaagat caagacctga 2100
3-16	Sequences	
3-16-1	Sequence Number [ID]	16
3-16-2	Molecule Type	DNA
3-16-3	Length	2100
3-16-4	Features	misc_feature 1..2100
	Location/Qualifiers	note=Description of Artificial Sequence: Synthetic polynucleotide source 1..2100 mol_type=other DNA organism=synthetic construct
	NonEnglishQualifier Value	
3-16-5	Residues	atgatcccag caagcgccaa ggcaccacac aagcagcccc acaagcagag catctccatc 60 ggcaggggca caaggaagag ggacgaggac agcggaaaccg aagtgggaga gggaaacagac 120 gagtggggcac agtccaaggc aaccgctgctg ccaccctgacc agctggagct gacagatgcc 180 gagctgaagg aggagttcac caggatcctg acagccaaca atccacacgc cccccagaa 240 atcgtgctgct acagcttcaa ggagggcaca tacaagccaa tccgctctctg gaaccagctg 300 gccgtgact acaccctgctg gggcaactctg atcccccaagg actccgatga gggccggaga 360 cagcactaca gggacgagct ggtggcagga tcccaggagt ctgtgaaagt gatctctgag 420 accggcaatc tggaggagga cgaggagcca aaggagctgg agaccgagcc aggaagccag 480 acagatgtgc ctgcagcagag agcagcagag aaggtgaccg aggaggagct gatgacaccc 540 aagcagccaa aggagcggaa gctgaccaac cagttcaact tctccgagag agcctctcag 600 acatacaaca atccagctgct ggacagagag tgccagacct agccaccccc cagaaccaac 660 ttctccgcca cagccaatca gtggggagatc tacgatgctt acgtggagga gctggagaag 720 caggagaaga ccaaggagaa ggagaaggcc aagacacccg tggccaagaa gtccggcaag 780 atggccatgc ggaagctgac cagcatggag tcccagacag acgatctgat caagctgtct 840 caggccgcca agatcatgga gagaatggtg aaccagaaca cctacgacga catcgccag 900

		<p>gacttcaagt actacgacga tgcagcagac gagtacaggg atcaagtggg cacactgctg 960 cctctgtgga agttccagaa cgacaaggcc aagaggctga gcgtgaccgc cctgtgctgg 1020 aatccaaagt acagggacct gttcgcagtg ggatacggaa gctacgactt catgaagcag 1080 agcagaggca tgctgctgct gtactccctg aagaacccca gcttccctga gtacatgttc 1140 agctccaact ccggcgtgat gtgcctggac atcccactgg atcaccctca cctgggtggc 1200 gtggggcact acgacggcaa cgtggccatc tacaatctga agaagcctca ctctcagccc 1260 agcttctgca gcagcgcaa gagcggcaag cactccgatc ccgtgtggca ggtgaagtgg 1320 cagaaggacg acatggacca gaacctgaac ttcttctccg tgtcctctga tggcaggatc 1380 gtgagctgga ccctggtgaa gcgcaagctg gtgcacatcg acgtgatcaa gctgaaggtg 1440 gagggcagca ccacagaggt gccagaggga ctgcagctgc acccagtggt atgcccagca 1500 gccttcgact tcacaagga gatcgactac atgttctcgg tgggcaccga ggagggcaag 1560 atctacaagt gcagcaagag ctacagctcc cagttctcgg acacatacga tgcccacaac 1620 atgagcgtgg acaccgtgtc ctggaatccc taccacacaa aggtgttcat gagctgcagc 1680 agcgactgga cgtgaagat ctgggatcac accatcaaga caccatggt catctacgac 1740 ctgaactccg ccgtgggcca tgtggcatgg gcaccatact ccagcacagt gttcgagca 1800 gtgaccacag acggcaaggc acacatcttc gatctggcca tcaacaagta cgaggccatc 1860 tgcaatcagc ccgtggccgc caagaagaac aggttgacc acgtgcagtt caatctgatc 1920 caccocatca tcatcgtggg cgacgatcgg ggccacatca tctctctgaa gctgagcccc 1980 aacctgagaa agatgcccga ggagaagaag ggacaggagg tgcagaaggg accagcagtg 2040 gagatcgcaa agctggacaa gctgctgaa ctggtgcggg aggtgaagat caagacctga 2100</p>
3-17	Sequences	
3-17-1	Sequence Number [ID]	17
3-17-2	Molecule Type	DNA
3-17-3	Length	13875
3-17-4	Features	misc_feature 1..13875
	Location/Qualifiers	note=Description of Artificial Sequence: Synthetic polynucleotide source 1..13875 mol_type=other DNA organism=synthetic construct
3-17-5	NonEnglishQualifier Value Residues	<p>atgttcagaa tggcagacg gcagctgtgg aagcacagcg tgaccagagt gctgaccag 60 cggctgaagg gcgagaaaga gccaagaga gccctgctgg acgcccggca caactacctg 120 ttcgccatcg tggccagctg cctggacctg aacaagaccg aggtggaaga cgccatcctg 180 gaaggcaacc agatcgagcg gatcgaccag ctgttcgccc tggggcagct gcggcacctg 240 atgttctact accaagacgt ggaagagggc gagacagggc agctgggaaag cctggggcga 300 gtgaacctgg tgagcggcaa gatcaagaaa cccaaggtgt tcgtgaccga gggcaacgac 360 gtggccctga caggcgtgtg cgtgttcttc atcagaaccg accccagcaa ggccatcacc 420 cccgacaaca tccaccagga agtgagcttc aacatgctgg acgcccgcga cggcggcctg 480 ctgaacagcg tgcggagact gctgagcgac atcttcatcc ccgcccctgag agccacaagc 540 cacggctggg gagagctgga aggactgcag gacgcccga acatccggca ggaattcctg 600 agcagcctgg aaggattcgt gaacgtgctg agcggcgccc aggaaaagct gaaagaaaaa 660 gtgaacctgc ggaagtgcga catcctggaa ctgaaaaccg tgaagagacc cactgactac 720 ctgacctggg ccaacaaccg cgagacactg ggcaagatcg aggactgcat gaaagtgtgg 780 atcaagcaga ccgaacaggt gctggccgag aacaaccagc tgctgaaaga agccgacgac 840 gtgggcccga gagccgagct ggaacactgg aagaagcggc tgagcaagtt caactacctg 900 ctggaacagc tgaagagccc cgacgtgaag gccgtgctgg ccgtgctggc agccgccaag 960 agcaaactgc tgaaacctg gcgagagatg gacatccgga tcaccgacgc caccaacgag 1020 gccaaaggca acgtgaagta cctgtacacc ctggaaaagt gctgcgaccc cctgtacagc 1080 agcgaccccc tgagcatgat ggaagccatc cccaccctga tcaaacgcat caagatgatc 1140 tacagcatca gccactacta caacaccagc gagaagatca ccagcctggt cgtgaaagtg 1200 accaaccaga tcatcagcgc ctgcaaggcc tacatcacca acaacggcac cgccagcatc 1260 tggaaaccag cccaggacgt ggtggaagag aagatcctga gcgccatcaa gctgaagcag 1320 gaataaccagc tgtgcttcca caagaccaag cagaagctga aacagaaccc caacgccaag 1380 cagttcgact tcagcgagat gtacatcttc ggcaagttcg agacattcca ccggcggctg 1440 gccaaagatc tcgacatctt caccaccctg aaaacatata gcgtgctgca ggacagcacc 1500 atcgagggcc tggaaagacat gccaccgaag taccagggca tcgtggccac catcaagaag 1560 aaagagtaca acttctctgga ccagcgcaag atggacttcg accaggacta cgaggaattc 1620 tgcaagcaga caaacgacct gcacaacgag ctgcgcaagt tcatggacgt gaccttcgcc 1680 aagatccaga acaccaacca gccctgcgg atgctgaaga agttcgagag actgaacatc 1740 cccaacctgg gcatcgacga caagtaccag ctgatcctgg aaaactacgg cgccgacatc 1800 gacatgatca gcaagctgta cacaagcag aagtacgacc cccccctggc ccggaaccag 1860 ccccccatcg ccggcaaaat cctgtgggac agacagctgt tccaccggtt ccagcagccc 1920 atcgagctgt tcagcagca ccccgccgtg ctgagcaacg ccgagggcaa acccatcatc 1980 cggagctaca accggatggc caaggtgctg ctggaattcg aggtgctggt ccaccgggccc 2040 tggctgcggc agatcgaaga gatccacgtg ggactggaag ccagcctgct cgtgaaggcc 2100 cccggaacgg gcgagctgtt cgtgaacttc gaccccaga tctgtatcct gttccgggaa 2160 accgagtgca tggcccagat ggggctggaa gtgagcccc tggccaccag cctgttccag 2220 aagcgggacc ggtacaagcg gaacttcagc aacatgaaga tgatgctggc cgagtaccag 2280 cgcgtgaaga gcaagatccc cgccgcatc gagcagctga tcgtgcccc cctggccaaa 2340 gtggacgagc cctgcagcc aggactggcc gccctgacat ggaccagcct gaacatcgag 2400 gcctacctgg aaaacacatt cgccaaaatc aaggacctgg aactgctgct ggaccgctg 2460 aacgacctga tcgagttccg gatcgacgcc atcctggaag agatgagcag cccccctg 2520 tgccagctgc ccaggaaga acccctgacc tgcgaagagt tctgcagat gaccaaggac 2580 ctgtgcgtga acggcgcca gatcctgcac tcaagagca gcctggtgga agaagcctg 2640 aacgagctcg tgaacatgct gctggacgtg gaagtgtcga gcgaggaaga gagcgagaag 2700</p>

atcagcaacg	agaacagcgt	gaactacaag	aacgagagca	gcgccaagcg	ggaagagggc	2760
aacttcgaca	cctgaccag	cagcatcaac	gccagagcca	acgcccctgct	gctgaccacc	2820
gtgacccgga	agaaaaaaga	aaccgagatg	ctgggccaag	aggccagaga	gctgctgagc	2880
cacttcaacc	accagaacat	ggacgcccctg	ctgaaagtga	cacggaacac	cctggaagcc	2940
atccggaagc	ggatccacag	cagccacacc	atcaacttcc	gggacagcaa	cagcgcaccg	3000
aacatgaagc	agaacagcct	gcccattctc	cgggccagcg	tgacactggc	catccccaac	3060
atcgtgatgg	ccccgcctt	ggaagacgtg	cagcagacac	tgaacaaggc	ctggaatgc	3120
atcatcagc	tgcccagggg	cgtgcgcag	tggagcagc	aactgcctgag	caagaagaag	3180
atccaggaac	ggaaaatggc	gcgccctcag	agcaacgagg	acagcgcagc	cgacgtggaa	3240
atggggcaga	acgagctgca	ggacacactg	gaaatcgcca	gcgtgaaact	gccccatccc	3300
gtgcagacca	agaactacta	caagaaactg	agcgaaaaca	aagaaatcgt	gaagctgggtg	3360
agcgtgctga	gcaccatcat	caacagcacc	aagaaaagaag	tgatcaccag	catggactgc	3420
ttcaagcggg	acaaccacat	ctggcagaag	ggcaaaagaag	aggccatcaa	gaccttcac	3480
accagagcc	cctgctgag	cgagttcgag	agccagatcc	tgtacttcca	gaacctggaa	3540
caggaaatca	acgccagccc	cgagtaactg	tgcgtgggca	gcactgccct	gtcacaccgc	3600
gacctgaagt	tcgccctgac	gcgcgagaca	aaggcctgga	tggtcgtgat	cggccggcac	3660
tgcaacaaaa	agtacagaag	cgagatggaa	aacatcttca	tgctgatcga	ggaattcaac	3720
aagaaactga	accggcccat	caaggacctg	gacgacaatc	gaatcgccat	ggccgcaactg	3780
aaagagatca	gagaggaaca	gatcagcatc	gacttccaag	tgggccccc	cgaggaaagc	3840
tacgccctgc	tgaacagata	cggactgctg	atcgcccggg	aagagatcga	caaggtggac	3900
acctgcaact	acgctggga	gaagctgctg	gccagagccc	gcgaggtgca	gaacaaactg	3960
gtgagcctgc	agcccagctt	caagaaagaa	ctgatcagc	ccgtgggaag	gttccctgag	4020
gactgccacc	agttctacct	ggactacgac	ctgaacggcc	ccatggccag	cggcctgaaa	4080
ccccaggaag	ccagcgaccg	gctgatcatg	ttccagaacc	agttcgacaa	catctaccgg	4140
aagtacatca	cctacacagc	cggcagggaa	ctggttcggc	tgcccgccac	acagtacccc	4200
cagctgctgg	aaatcaagaa	gcagctgaac	ctgctgcaga	agatctacac	cctgtacaac	4260
agcgtgatcg	agacagtga	cagctactac	gacatcctgt	ggagcgaagt	gaacatcgag	4320
aagatcaaca	acgaactgct	ggaattccag	aaccggctgc	ggaagctgcc	cgagcactg	4380
aaggactggc	aggccttccc	ggactgaag	aaaatcatcg	acgacttcag	cgagtgctgc	4440
ccctgctgg	agtacatggc	cagcaaggcc	atgatggaa	ggcactggga	gagaatcacc	4500
acactgaccg	gccacagcct	ggacgtgggc	aacgagagct	tcaagctgcg	gaacatcatg	4560
gaagcccac	tgctgaagta	caaagaggaa	atcgaggaca	tctgcatcag	cgccctgaaa	4620
gagcgggaca	tcgagcagaa	actgaaacaa	gtgatcaacg	agtgggacaa	caagaccttc	4680
accttcggca	gcttcaagac	cagaggcgag	ctgctgctgc	ggggcgacag	caccagcgag	4740
atcatcgcca	acatggaaga	cagcctgatg	ctgctgggca	gcctgctgag	caaccggtac	4800
aacatgcctc	tcaaggccca	gatccagaaa	tgggtgcagt	acctgagcaa	cagcaccgac	4860
atcatcgaga	gctggatgac	cgtgcagaac	ctgtggatct	acctggaagc	cgtgttcgtg	4920
ggcggcgaca	tcgccaagca	gctgcccaaa	gaggccaagc	ggttcagcaa	catcgacaag	4980
agctgggtca	agatcatgac	cagagcccac	gaggtgcccc	gcgtggtgca	gtgctgctg	5040
ggcgacgaaa	cactgggaca	gctgctgccc	cacctgctgg	accagctgga	aatctgccag	5100
aagagcctga	ccggctacct	ggaaaagaaa	cggctgtgct	tcccccggtt	cttcttcgct	5160
agcgaccccg	cctgctgga	aatcctgggc	caggccagcg	acagccacac	aatccagggc	5220
caactgctga	acgtgttcga	caacatcaag	agcgtgaagt	tccacgagaa	aatctacgac	5280
cggatcctga	gcatcagcag	ccaggaaggc	gagacaatcg	agctggacaa	gcccgtgatg	5340
gccgagggaa	acgtggaagt	gtggctgaac	agcctgctgg	aagagagcca	gagcagcctg	5400
cacctcgtga	tcagacagc	gcgcgccaac	atccaggaaa	ccggcttcca	gctgaccgag	5460
ttcctgagca	gcttcccagc	acaagtggga	ctgctgggca	tccagatgat	ctggaccaga	5520
gacagcgaag	aggccttgag	aaacgccaag	ttcgacaaga	aatcatgca	gaaaacaaa	5580
caggcattcc	tggaactgct	gaacaccctg	atcgactgga	ccaccgggga	aatccagcagc	5640
accgagagag	tgaagtacga	gacactgatc	accatccacg	tgccaccagc	ggacatcttc	5700
gacgacctgt	gccacatgca	catcaagagc	cccattggact	tcgagtggct	gaagcagtgc	5760
aggttctact	tcaacgagga	cagcgacaag	atgatgatcc	acatcaccga	cgtggccttc	5820
atctaccaga	acgagttcct	gggctgcacc	gaccgcctcg	tgatcaccac	cctgaccgac	5880
cggtgctaca	tcacactggc	ccaggcactg	ggcatgagca	tgaggaggcg	accagcagga	5940
cccgcggca	caggcaagac	cgaaaccacc	aaggacatgg	gacgctgctc	gggcaaaata	6000
gtggtggtgt	tcaactgacg	cgaccagatg	gacttccggg	gcctgggccc	gatcttcaag	6060
ggcctggcac	agagcggaa	ctggggctgc	ttcgacgagt	tcaacagaat	gacctgccc	6120
gtgctgagcg	tgcccgaca	gcagatcagc	atcatcctga	catgcaaaaa	agagcacaag	6180
aagagcttca	tcttaccgca	cggcgacaac	gtgacctga	accccagatt	cggcctgttc	6240
ctgacaatga	accccggcta	gcgcggacgg	caggaactgc	ccgagaacct	gaagatcaac	6300
ttccggagcg	tgcccatgat	ggtgcccgac	cggcagatca	tcatcagagt	gaaactggcc	6360
agctgcccgt	tcactgacaa	cgtggtgctg	gcccgggaag	tcttccacct	gtacaagctg	6420
tgcaagaaac	agctgagcaa	acaggtgcac	tacgacttcg	gcctgaggaa	catcctgagc	6480
gtgctgagaa	ccctgggagc	gcaccaagcg	gccaacccca	tggacaccga	gagcacaatc	6540
gtgatgcggg	tgctgcccga	catgaacctg	agcaagctga	tcgacgagga	cgagcccctg	6600
ttcctgagcc	tgatcgagga	cctgttcccc	aacatcctgc	tggacaaggc	cggctacccc	6660
gaactggaag	ccgccatcag	cagacaggtg	gaagaggccc	gcctgatcaa	ccaccccccc	6720
tggaaactga	aagtgatcca	gctgttcgag	acacagcgcg	tgccggcacg	catgatgaca	6780
ctgggacca	gcggagccgg	caagaccacc	tgcatccaca	cactgatcgg	ggcctcagc	6840
gactgcggca	agccccaccg	cgagatcgcg	atgaaccaca	aggccatcac	gcccccacc	6900
atgttcggca	gactggacgt	ggccaccaac	gactggaccg	acggcatctt	cagcaccctg	6960
tggcgcaaga	ccctgcccgg	caagaagggc	gagcacatct	ggatcatcct	ggacggcccc	7020
gtggacgcca	tctggatcga	gaacctgaac	agcgtgctgg	acgacaacaa	gacactgacc	7080
ctggccaacg	gcgaccggat	ccccatggcc	cccaactgca	agatcatctt	cgagccccac	7140
aacatcgaca	acgccagccc	gcgccaccgtg	agcagaaacg	gcattggtgt	catgagcagc	7200
agcatcctgg	actggagccc	catcctggaa	ggcttctgca	agaagcggag	ccccaggaa	7260

gccgagatcc	tgagacagct	gtacaccgag	agcttccccg	acctgtaccg	gttctgcatc	7320
cagaacctgg	agtacaagat	ggaagtgctg	gaagccttcg	tgatcaccca	gagcatcaac	7380
atgctgcagg	gcctgatccc	cctgaaagaa	cagggcggag	aagtgagcca	ggccccactg	7440
ggcagactgt	tcgtgttcgc	cctgctgtgg	agcgcggcgg	ccgcccctgga	actggacgga	7500
aggcggagac	tggaactgtg	gctgcccagc	agaccaaccg	gcacctgga	actgccccca	7560
ccagccggac	ccggcgacac	cgcttctgac	tactactgtg	cccccgacgg	cacctggacc	7620
actggaaca	ccggaccaca	ggaatacctg	taccccagcg	acaccacccc	cgagtacggc	7680
agcatcctgg	tgcccaactg	ggacaactgt	cggaccgact	tcctgatcca	gacaatcgcc	7740
aagcagggaa	aggccgtgct	gctgatcggc	gagcagggca	cagccaagac	cgtgatcatc	7800
aagggcttca	tgagcaagta	cgaccccag	tgccacatga	tcaagagcct	gaacttcagc	7860
agcgcaccca	cccactgat	gttccagcgg	accatcgaga	gctacgtgga	caagcggatg	7920
ggcaccacct	acggcccccc	agccggcaag	aaaatgaccg	tgttcatcga	cgacgtgaac	7980
atgcccata	tcaacgagtg	gggcgaccac	gtgaccaacg	agatcgtgcg	gcagctgatg	8040
gaacagaacg	gcttctacaa	cctggaaaag	ccggcgagtg	tcaccagcat	cgtggacatc	8100
cagtctctgg	ccgccatgat	ccaccocggc	ggcggaaaga	acgacatccc	gcagcggctg	8160
aagcggcagt	tcagcatctt	caactgcacc	ctgcccagcg	aggccagcgt	ggacaagatc	8220
ttcggcgtga	tcggcgtggg	ccactactgc	accagagag	gcttcagcga	ggaagtgcgg	8280
gacagcgtga	ccaagctggt	gcccctgaca	agacggctgt	ggcagatgac	caagatcaag	8340
atgctgcccc	cccccgcaa	gttccactac	gtgttcaacc	tgccggacct	gagcagagtg	8400
tggcagggaa	tgctgaacac	caccagcgaa	gtgatcaaac	agccccacga	cctgctgaag	8460
ctgtggaagc	acgagtgcaa	gagagtgatc	gccgaccggt	tcaccgtgag	cagcgcagtg	8520
acatggttgc	acaaggccct	gtgagcctg	gtggaagag	aattcggcga	agagaagaaa	8580
ctgctggtgg	actgcggcat	cgacacctac	ttcgtggact	tcctgcgcga	cgcccccgaa	8640
gccgcggcgg	agacaagcga	agaggccgac	gcccagacac	ccaagatcta	cgagcccac	8700
gagagcttca	gcacactgaa	agaaaggctg	aacatgttcc	tgacagctga	caacgagagc	8760
atcgggggag	ccggcatgga	catggtgttc	ttcgcggcag	ccatggtgca	cctcgtgaag	8820
atcagcagag	tgatccggac	cccccagggc	aacgcctcgc	tcgtgggagt	gggagggcgc	8880
ggcaagcaga	gcctgaccag	actggccagc	ttcatcgccg	gctacgtgag	cttccagatc	8940
accctgacct	ggagctacaa	caccagcaac	ctgatggaag	acctgaaggt	gctgtaccgg	9000
acagccggcc	agcaggggaa	gggcatcacc	ttcatcttca	ccgacaacga	gatcaaggac	9060
gagagcttcc	tggagtacat	gaacaactgt	ctgagcagcg	gagaggtgag	caactctgtc	9120
gcccgggcag	agatcgacga	gatcaacagc	gacctggcca	gcgtgatgaa	gaaagaattc	9180
ccccggtgcc	tgcccacaaa	cgagaacctg	cacgactact	tcattgagcag	agtgcggcag	9240
aacctgcaca	tcgtgctgtg	cttcagcccc	gtgggcgaga	agttcagaaa	ccggcccctg	9300
aagttccccg	ccctgatcag	cggtgcacc	atcgactggt	tcagccgggt	gccccaggac	9360
gcctggtgg	ccgtgagcga	gcacttctcg	accagctacg	acatcgactg	cagcctggaa	9420
atcaagaaag	aggtggtgca	gtgcatgggc	agcttccagg	acggcgtggc	cgagaaatgc	9480
gtggactact	tcagcgggtt	ccggcggagc	accacgtgta	cccccaagag	ctacctgagc	9540
ttcatccag	gctacaagtt	catctacggc	gagaagcacg	tggaagtgcg	cacctggccc	9600
aaccggatga	acaccggcct	ggaaaaactg	aaagaggcca	gcgagagcgt	ggccgcccctg	9660
agcaaaagac	tggaagccaa	agaaaaagaa	ctgcaggtgg	ccaacgacaa	ggccgacatg	9720
gtgctgaaa	aagtgaccat	gaaggccag	gcccggcaga	aagtgaagac	cgaggtgcag	9780
aaagtgaag	accgggcccc	ggccatcgtg	gacagcatca	gcaaggacaa	ggccatcgcc	9840
gaggaaaaag	tggaagcagc	caagcccggc	ctggaagagg	cagaagccgc	cctgcagacc	9900
atccggcccc	gcgacatcgc	cacagtgcgg	accctgggaa	ggcccccca	cctgatcatg	9960
cggatcatgg	actgcgtgct	gctgctgttc	cagagaaagg	tgagcggcgt	gaagatcgac	10020
ctggaaaaaa	gctgcaccat	gcccagctgg	caggaagacc	tgaagctgat	gaccgcccgc	10080
aacttctctg	agaacctgca	gcagttcccc	aaggacacca	tcaacgagga	agtgatcgag	10140
ttcctgagcc	cctacttcca	gatgcccagc	tacaacatcg	aaaccgcaaa	acgctgtctg	10200
ggcaactggt	ccggactgtg	cagctggacc	aaggccatgg	ccagcttctt	cagcatcaac	10260
aaagaggtgc	tgcccctgaa	ggccaacctg	gtggtgcagg	aaaaccggca	cctgctggcc	10320
atgcaggacc	tgcaaaaagc	ccaggccgag	ctggacgaca	agcaggccga	gctggacgtg	10380
gtgcaggccg	agtacgagca	ggccatgacc	gagaagcaga	ccctgctgga	agacgcagag	10440
cggtgcagac	acaagatgca	gaccgccagc	accctgatca	gcggactggc	cgggcaaaaa	10500
gagcgggtga	ccgagcagag	ccaggaattc	gcccggccag	ccaagcggct	cggtggagac	10560
gtgctgctgg	ccaccgcctt	cctgagctac	agcggcccct	tcaaccagga	attcagggag	10620
ctgctgctga	acgactggcg	gaaagagatg	aaggccagaa	agatcccctt	cggaagaagc	10680
ctgaacctga	gcgagatgct	gatcgacgcc	cccaccatca	gcgagtggaa	cctgcagggg	10740
ctgcccacag	acgacctgag	catccagaac	ggaatcatcg	tgaccaaacg	cagcagatca	10800
cccctgctga	tcgaccccc	gacacagggc	aagatctgga	tcaagaacaa	agagagccgg	10860
aacgagctgc	agatcaccag	cctgaaccac	aagtacttcc	ggaaccacct	ggaagacagc	10920
ctgagcctgg	gcaggccact	gctgatcgag	gacgtggggc	aggaactgga	cccagccctg	10980
gacaactgct	tggaacggaa	cttcatcaag	accggcagca	ccttcaaggt	gaaagtgggc	11040
gacaaagaag	tggactgctg	ggacggcttc	cggtgttaca	tcaccaccaa	gctgcccac	11100
cccgcctaca	ccccgagat	cagcgcggcg	accagcatca	tcgacttcac	cgtgacaatg	11160
aagggactgg	aagaccagct	gctgggacgc	gtgatcctga	cagagaagca	ggaactggaa	11220
aaagaacgga	ccacactgat	ggaagacgtg	accgccaaca	agcggcggat	gaaggaactg	11280
gaagacaacc	tgctgtacag	gctgaccagc	accaggggca	gcctggtgga	agacgagagc	11340
ctgatcgtgg	tgctgagcaa	caccaagcgg	accgcagagg	aagtgaccac	gaagctggaa	11400
atcagccggc	agacagaggt	gcagatcaac	agcggccagag	aagagtaccg	gcccgtggcc	11460
accgggggaa	gcatcctgta	cttctctgat	accgagatgc	ggctcgtgaa	cgagatgtac	11520
cagaccagcc	tgccgcagtt	cctgggctcg	ttcgacctga	gcctggccag	aagcgtgaag	11580
agccccatca	ccagcaagag	aatcgccaac	atcatcgagc	acatgacctc	cgaggtgtac	11640
aaatacgccg	ccagaggcct	gtacgaggaa	cacaagtctc	tgttcacact	gctgctgacc	11700
ctgaagatcg	acatccagcg	gaacagagtg	aagcacgaag	agttcctgac	actgatcaag	11760
gggggagcca	gcctggacct	gaaggcctgc	cccccaagc	ccagcaagtg	gatcctggac	11820

		<p>atcacctggc tgaacctggt ggaactgagc aagctgagac agttcagcga cgtgctggac 11880 cagatcagcc gcaacgagaa gatgtggaag atctggttcc acaaagagaa ccccaggaa 11940 gaaccctgc ccaacgcta cgacaagagc ctggactgct tccggcggct gctgctgac 12000 agaagctggt gccccgaccg gacaatcgcc caggcccgca agtacatcgt ggacagcatg 12060 ggagagaagt acgccgaggg cgtgatcctg gacctggaaa agacctggga ggaaagcgac 12120 cccagaacct cctgatctg cctgctgagc atgggacagc accccaccga cagcatcatc 12180 gcctgggca agagactgaa gatcgagaca agatactgga gcatgggcca gggccaggaa 12240 gtgcagttca gaaagctgct gcagcagacc atggccaacg gcggctgggc cctgctgca 12300 aactgccacc tgggctgga ctctcatggac gaactgatgg acatcatcat cgagacagag 12360 ctggtgcacg acgccttcag actgtggatg accaccgagg cccacaagca gttccccatc 12420 acctgctgc agatgagcat caagttcgcc aacgaccccc cccagggact gagagccggc 12480 ctgaagagaa cctacagcgg cgtgagccag gacctgctgg acgtgagcag cggcagccag 12540 tggaaagcca tgcgttacgc cgtggcattc ctgcacagca ccgtgcagga acggcggaag 12600 ttcggcgccc tgggatggaa catcccctac gagttcaacc aggcgactt caacgccacc 12660 gtgcagttca tccagaacca cctggacgac atggacgtga agaaaagggt gagctggaca 12720 accatccggt acatgatcgg agagatccag tacggcggca gagtgaccga cgactacgac 12780 aagaggctgc tgaacacctt cgccaaagtg tggttcagcg agaactggt cggccccgac 12840 ttcagcttct accagggcta caacatcccc aagtgcagca ccgtggacaa ctacctgca 12900 tacctccaga gcctgccgc ctacgacagc cccgaggtgt tcggactgca ccccaacgcc 12960 gacatcacct accagagcaa actggccaag gacgtgctgg acaccatcct gggcatccag 13020 cccaggaca ccagcggcgg aggcgacgaa acccggaag cagtggctgg ccagctggcc 13080 gacgacatgc tggaaaagct gcccccgcac tacgtgctct tcgaagtgaa agaacgctg 13140 cagaagatgg gcccttcca gcccatgaac atcttctga ggcaggaaat cgaccggatg 13200 cagcgggtgc tgagcctcgt gcggagcaca ctgaccgagc tgaactggc catcgacggc 13260 accatcatca tgagcgagaa cctgcccggac gcaactggact gcatgttcca cgccagaatc 13320 cccgatggt ggaaaaaggc cagctggatc agcagcacc tgggcttctg gttaccggaa 13380 ctgatcgaga gaaacagcca gttcaccagc tgggtgttca acggcagacc ccactgcttc 13440 tggatgaccg gcttcttcaa cccacaaggc ttctgacag caatgcgcca ggaactcacc 13500 agagccaaca agggctgggc cctggacaac atgggtgctgt gcaacgaagt gaccaagtg 13560 atgaaggagc acatcagcgc cccccccaca gagggcgtgt acgtgtacgg cctgtacctg 13620 gaaggcgccg gatgggacaa gagaaacatg aagctgatcg agagcaagcc caaggtgctg 13680 ttcgagctga tgcccgtgat caggatctac gccgagaaca acacctgag ggacccccgg 13740 ttctacagct gccccatcta caagaaacct gtgcgcaccg acctgaaacta catcgccgcc 13800 gtggacctga ggacagccca gacaccggag cactgggtgc tgagaggcgt ggcactgctg 13860 tgcgacgtga agtga 13875</p>
3-18	Sequences	
3-18-1	Sequence Number [ID]	18
3-18-2	Molecule Type	DNA
3-18-3	Length	13875
3-18-4	Features	misc_feature 1..13875
	Location/Qualifiers	note=Description of Artificial Sequence: Synthetic polynucleotide source 1..13875 mol_type=other DNA organism=synthetic construct
3-18-5	NonEnglishQualifier Value Residues	<p>atgttcagaa tcggcagacg gcagctgtgg aagcacagcg tgaccagagt gctgaccag 60 cggctgaagg gcgagaaaga ggccaagaga gccctgctgg acgcccggca caattacctg 120 tttgccatcg tggccagctg cctggacctg aacaagacc aggtggaaga tgccatctg 180 gaaggcaacc agatcgagcg gatcgaccag ctgtttgccc tggggaggact gcggcaacctg 240 atgttctatt atcaagacgt ggaagaggcc gagacaggcc agctgggatc tctgggcgga 300 gtgaatctgg tgtccggcaa gatcaagaaa cccaaggtgt tcgtgaccga gggcaacgac 360 gtggocctga caggcgtgtg cgtgttcttc atcagaaccc accccagcaa ggccatcacc 420 cccgacaaca tccaccagga agtgtccttc aacatgctgg atgcccgcca tggcggcctg 480 ctgaattctg tcgggagact gctgagcagc atcttcatcc ccgcccctgag agccacatct 540 cacggctggg gagagctgga aggactgagc gacgcccgca atatccggca ggaatttctg 600 agcagcctgg aaggattcgt gaacgtgctg tctggcgccc aggaaaacct gaaagaaaaa 660 gtgaacctgc ggaagtgcga tctcctggaa ctgaaaacc tgaaagagcc caccgactac 720 ctgacctgga ccaacaacct tgagacactg ggcaagatcg aggactgcat gaaagtgtgg 780 atcaagcaga ccgaacaggt gctggccgag aacaaccagc tgctgaaaga agccgacgac 840 gtgggcccga gagccgagct ggaacactgg aagaagcggc tgagcaagtt caactacctg 900 ctggaacagc tgaagtcccc gcagctgaag gccgtgctgg ctgtgctggc agccgccaag 960 agcaaacctg tgaaacctg gcgagagatg gacatccgga tcaccgagc cccaacagag 1020 gccaaggaca acgtgaagta cctgtacacc ctggaaaagt gctgagcacc cctgtacagc 1080 agcagacctc tgagcatgat ggacgccatc cctacctga tcaacgcat caagatgatc 1140 tacagcatca gccactacta caaccaccg gagaaagatc ccagcctggt cgtgaaagtg 1200 accaatcaga tcatcagcgc ctgcaaggcc tacatcacca acaacggcac cgccagcatc 1260 tggaaaccagc ccaggatgt ggtggaagag aagatcctgt ctgccatcaa gctgaaagcag 1320 gaataaccagc tgtgttttca caagaccaag cagaagctga aacagaacct caacgccaag 1380 cagttcgact tcagcgagat gtatatcttc ggcaagttcg agacattcca ccggcggctg 1440 gccaagatca tcgacatctt taccacctg aaaacataca gcgtgctgca gcgacagcacc 1500 atcgagggcc tggaaagatat ggccaccaag taccagggca ttgtggccac catcaagaag 1560 aaagagtaca acttcttggc ccagcgcaag atggacttgc accaggacta cgaggaaattc 1620 tgcaagcaga caaacgacct gcacaacgag ctgcccgaagt ttatggacgt gacctcgcc 1680 aagatccaga acaccaacca gcccctcggc atgctgaaga agtttgagag actgaaatc 1740 cccaacctgg gcatcgagca taagtaccag ctgatcctgg aaaactacgg cgccgacatc 1800</p>

gacatgatca	gcaagctgta	cacaaagcag	aagtacgacc	ccccctggc	cgggaatcag	1860
cctcctatcg	ccggcaaaat	cctgtgggct	agacagctgt	ttcaccggat	ccagcagccc	1920
atgcagctgt	tccagcagca	cctgcccgtg	ctgagcacag	ccgaggccaa	acctcatc	1980
cggtcctaca	accgatggc	caaggtgctg	ctggaatctg	agggtgctgt	ccaccgggccc	2040
tggctgcgcc	agatcgaaga	gattcacgtg	ggactggaag	ccagcctgct	cgtgaaggct	2100
cctggaaccg	gcgagctgtt	tgtgaacttc	gacccccaga	tctctgatcct	gttccgggaa	2160
accgagtgca	tggcccagat	ggggctggaa	gtgtctctct	tggccacctc	cctgttcccag	2220
aagcgggacc	ggtacaagcg	gaacttcagc	aacatgaaga	tgatgtggc	tgtagtcag	2280
cgctggaagt	ccaagatccc	cgctgccatc	gagcagctga	tctgtcctca	cctggccaaa	2340
gtggacgagg	ccctgcagcc	aggactggcc	gctctgacat	ggaccagcct	gaacatcgag	2400
gcctatctgg	aaaacacatt	cgccaaaatc	aaggatctgg	aactgtgct	ggaccgctg	2460
aacgacctga	tcgagttccg	gatcgacgcc	attctggaag	agatgtccag	cacccccctg	2520
tgtcagctgc	ccaggaaga	accctgacc	tgcgaagagt	tctctcagat	gaccaaggac	2580
ctgtgctga	acggcgcca	gattctgcac	ttcaagtcca	gcctgggga	agaagccgtg	2640
aacgagctcg	tgaatctgct	gctggatgtg	gaagtgtgta	gcgaggaaga	gtccgagaag	2700
atctccaacg	agaacagcgt	gaactacaag	aacgagtcca	gcgccaagcg	ggaagagggc	2760
aacttcgaca	ccctgaccag	ctccatcaat	gccagagcca	acgccctgct	gctgaccacc	2820
gtgacctcga	agaaaaaaga	aaccgagatg	ctgggccaag	aggctagaga	gctgtgtcc	2880
cacttcaacc	accagaacat	ggatgcctcg	ctgaaagtga	cacggaatac	cctggaagcc	2940
atccggaagc	ggatccacag	cagccacacc	atcaacttcc	gggacagcaa	cagcggcacc	3000
aatatgaagc	agaacagcct	gccatcttc	cgggcctccg	tgacctggc	catccccatc	3060
atcgtgatgg	ccctgctct	gtaagatgtg	cagcagacac	tgaaccaaggc	cgtggaatgc	3120
atcatctccg	tgcccaaggg	cgctggcgag	tggctctagc	aactgtgtgc	caagaagaag	3180
atccaggaac	ggaaaatggc	cgccctgcag	tctaacgagg	acagcgactc	cgactgtgaa	3240
atggcgagca	atgagctgca	ggatacactg	gaaatcgct	ctgtgaaatc	gccccatccc	3300
gtgcagacca	agaactacta	taagaactgt	tccgaaaaca	aagaaatcgt	gaagtgggtg	3360
tctgtgctgt	ccaccatcat	caacagcacc	aagaaagaag	tgatcacctc	catggactgc	3420
ttcaagcggg	acaaccacat	ctggcagaag	ggcaaagaag	aggccattaa	gaacttcatc	3480
accagagcc	ccctgctgtc	cgagttcgag	tctcagatcc	tgtacttcca	gaacctggaa	3540
caggaaatca	acgccgagcc	cgagtacgtg	tgtgtgggct	ctatcgccct	gtataaccgc	3600
gacctgaagt	tgcacctgac	cgccgagaca	aaggcctgga	tggctgtgat	cggccggcac	3660
tgcaacaaaa	agtacagatc	cgagatggaa	aacatcttta	tgctgattga	ggaattcaac	3720
aagaaactga	accggcccat	taaggacctg	gacgacatca	gaatcgccat	ggccgcaactg	3780
aaagagatca	gagaggaaca	gatcagcatc	gacttccaag	tgggccccat	cgaggaaagc	3840
tacgctctgc	tgaacagata	cggactgctg	atcgcccggg	aagagatcga	caagttggag	3900
accctgcact	acgcctggga	gaagctgctg	gctagagccg	gcgaggtgca	gaacaaactg	3960
gtgtctctgc	agcccagctt	taagaaagaa	ctgatctccg	ccgtggaagt	gtttctgcag	4020
gactgccacc	agttctacct	ggactacgac	ctgaacggcc	ccatggctc	tggcctgaaa	4080
cctcaggaag	cctccgaccg	gctgattatg	tttcagaacc	agttcgacaa	tatctaccgg	4140
aagtacatca	cctacacagg	cgccgagga	ctggtccggc	tgcctgccac	acagtacccc	4200
cagctgctgg	aaatcaagaa	gcagctgaac	ctgctgcaga	agatctacac	cctgtacaac	4260
tccgtgatcg	agacagtga	cagctactac	gacatcctgt	ggagcgaagt	gaacattgag	4320
aagattaaca	atgaactgct	ggaatttcag	aaccggtgcc	ggaagctgcc	cagagcactg	4380
aaggattggc	aggcctttct	ggatctgaag	aaaatcatcg	acgacttctc	cgagtgtctg	4440
cctctgctgg	agtacatggc	ctccaaggcc	atgatggaac	ggcactggga	gagaatcacc	4500
acactgaccg	gccacagcct	ggacgtgggc	aacgagagct	tcaagctgcg	gaacatcatg	4560
gaagccccac	tgtgaagta	caaagaggaa	atcgaggaca	tctgtatcag	cgccgtgaaa	4620
gagcgggata	tgcagcagaa	actgaaaca	gtgatcaacg	agtgggacaa	caagaccttt	4680
accttcggca	gcttcaagac	cagaggcgag	ctgctgctgc	ggggcgatag	cacctctgag	4740
atcattgcca	acatggaaga	tagcctgatg	ctgctgggct	ccctgtgag	caaccggtat	4800
aacatgccct	tcaaggetca	gattcagaaa	tgggtgcagt	acctgagcaa	ctccaccgac	4860
atcatcgagt	cctggatgac	cgtgcagaac	ctgtggatct	acctggaagc	cgtgttcgtg	4920
ggcggcgaca	tgccaagca	gctgccaaa	gaggctaagc	ggttctccaa	catcgacaag	4980
agctgggtca	agatcatgac	cagagcccac	gagggtccca	gcgtgggtgca	gtgctgtgtg	5040
ggcgacgaaa	cactgggaca	gctgctgct	catctgctgg	accagctgga	aatctgccag	5100
aagtccctga	cggctacct	ggaaaagaaa	cggctgtgtt	tccccgggtt	cttctctgtg	5160
tccgaccccg	ccctgctgga	aattctgggc	caggccaagc	actcacacac	aattcaggcc	5220
catctgctga	atgtgttcga	taacatcaag	agcgtgaagt	tccacgagaa	aatctacgac	5280
cggatcctga	gcatcagctc	ccaggaaggc	gagacaatcg	agctggacaa	gcctgtgatg	5340
gccgagggaa	acgtggaagt	gtggctgaac	agcctgctgg	aagagtccca	gagcagcctg	5400
cacctcgtga	tcagacaggg	cgctgccaac	atccagga	ccggctttca	gctgaccgag	5460
ttcctgtcca	gcttcccagc	acaagtgga	ctgctgggca	tccagatgat	ttgaccaga	5520
gactccgaag	aggccctgag	aaacgccaag	tccgataaga	aaattatgca	gaaaacaaat	5580
caggcatttc	tggaactgct	gaacacctg	atcgactgta	ccaccgggga	cctgagcagc	5640
accgagagag	tgaagtacga	gacactgatc	accatccacg	tgcaccagcg	ggacatcttc	5700
gacgacctgt	gcoacatgca	catcaagtct	cccatggatt	tccagtggtc	gaagcagctg	5760
aggttctact	tcaacgagga	ctccgacaag	atgatgatcc	acatcaccga	tgtggccttt	5820
atctatcaga	atgagttcct	gggctgtacc	gatcgctcgc	tgattacccc	cctgaccgac	5880
cggtgttaca	tcacactggc	ccaggcactg	ggcatgtcta	tgggagggcc	accagcagga	5940
cctgcggca	caggcaagac	cgaaaccacc	aaggacatgg	gacgctgcct	gggcaaatc	6000
gtgggtgtgt	tcaactgcag	cgaccagatg	gatttccggg	gcctggggccg	gacttttaag	6060
ggcctggcac	agagcgggaag	ctggggctgc	ttcgacgagt	tcaacagaat	cgacctgccc	6120
gtgctgtccg	tggccgcaca	gcagatctcc	atcatctcta	catgcaaaaa	agagcacaag	6180
aagtccctca	tcttcaccga	cggcgacaat	gtgacctgta	accccgagtt	tggcctgttc	6240
ctgacaatga	accctggcta	cgccggacgg	caggaactgc	ccgagaacct	gaagatcaac	6300
tttcggagtg	tggctatgat	ggtgcccgac	cggcagatca	ttatcagagt	gaaactggcc	6360

tctcgcgct	tcacgacaa	cggtgtgctg	gctcgggaag	tcttcacact	gtacaagctg	6420
tgcaagaac	agctgagtaa	acaggtgcac	tacgacttcg	gcctgaggaa	catcctgagc	6480
gtgctgagaa	ctctgggagc	cgtaagcgg	gccaacccca	tggataccga	gagcacaatc	6540
gtgatgcggg	tgctgcggga	catgaacctg	tccaagctga	tcgatgagga	cgagcccctg	6600
tttctgtctc	tgatcgagga	tctgtttccc	aacattctgc	tggataaggg	cggtaccccc	6660
gaactggaag	ctgctatcag	cagacaggtg	gaagaggtcg	gcctgatcaa	ccaccccccc	6720
tggaaactga	aagtgatcca	gctgttctgag	acacagcgg	tgcggcagcg	catgatgaca	6780
ctgggacctg	gcggagccgg	caagaccacc	tgtatccaca	cactgatgcg	ggcccagacc	6840
gattgcgcca	agccccaccg	cgagatgctg	atgaacccca	aggccattac	cgcccctcag	6900
atgttcggca	gactggacgt	ggccaccaac	gactggaccg	acggcatctt	cagcacctcg	6960
tggcgcaaga	ccctgcgggc	caagaagggc	gagcacatct	ggatcatcct	ggacggcccc	7020
gtggacgcca	tctggattga	gaacctgaac	agcgtgctgg	acgacaacaa	gacactgacc	7080
ctggccaacg	gcgaccggat	ccccatggcc	cccaactgca	agatcatctt	cgagccccac	7140
aacatcgaca	acgccagccc	tgccaccgtg	tccagaaacg	gcatggtggt	catgagcagc	7200
agctgacctg	attggagccc	tatcctggaa	ggcttccctga	agaagcggag	ggcccaggaa	7260
gccgagatcc	tgagacagct	gtacaccgag	agcttccccg	acctgtaccg	gttctgcatc	7320
cagaatctgg	agtacaagat	ggaagtgtcg	gaagcctttg	tgatcaccca	gagcatcaac	7380
atgctgcagg	gcctgatccc	cctgaaagaa	cagggcggag	aagtgtccca	ggcccacctg	7440
ggcagactgt	tcgtgtttgc	cctgctgtgg	agcgtcggcg	ccgctctgga	actggatgga	7500
aggcggagac	tggaaactgtg	gctgcgggagc	agacctaccg	gcacctgga	actgcctcca	7560
ccagctggac	ctggcgacac	cgcttctgat	tactacgtgg	cccctgacgg	cactggacc	7620
cactggaata	ccgggaccca	ggaatacctg	taccaccagc	acaccacccc	cgagtacggc	7680
tctatcctgg	tgcccaactg	ggacaactgt	cggaccgact	tctgatcca	gacaatcgcc	7740
aagcagggaa	aggcctgtct	gctgatcggc	gagcagggca	cagccaagac	cgtgatcatc	7800
aagggcttta	tgtctaagta	cgaccocgag	tgccacatga	tcaagagcct	gaacttcagc	7860
tccgccacca	cccactgat	gttccagcgg	accatcgaga	gctatgtgga	caagcggatg	7920
ggcaccacct	acggccctcc	agccggcaag	aaaatgaccg	tgttcatcga	cgactggaac	7980
atgcccatac	tcaacgagtg	ggcgaccaa	gtgaccaacg	agatcgtgcg	gcactgatg	8040
gaacagaacg	gcttctacaa	cctggaaaag	ccggcgagtg	tcacctctat	ctgggacatc	8100
cagtttctgg	ccgccatgat	ccaccctggc	ggcggaaaga	acgacatccc	ccagcggctg	8160
aagcggcagt	tcagcatctt	caactgcacc	ctgccacgcg	aggccagcgt	ggacaagatc	8220
tttggcgtga	tcggcgtggg	ccactactgc	accagagag	gcttccagca	ggaagtgcgg	8280
gacagcgtga	ccaagctggt	gcctctgaca	agacggctgt	ggcagatgac	caagatcaag	8340
atgctgcca	ccccgccea	gttccactac	gtgttcaaac	tgcgggacct	gagcagatg	8400
tggcagggaa	tgctgaacac	caccagcgaa	tgatcaaacg	agcccaacga	ctcgtgaag	8460
ctgtggaagc	acgagtgcaa	gagagtgatc	gccgaccggt	tcacctgttc	tagcagcgtg	8520
acatggttcg	acaaggccct	ggtgtccctg	gtggaagagg	aattcggcga	agagaagaaa	8580
ctgctggtgg	actgcggcat	cgatacctac	ttcgtggact	tctgcgcgca	cgcccctgaa	8640
gccgctggcg	agacaagtga	agagggcgac	gccgagacac	ccaagatcta	cgagcccatc	8700
gagtccttca	gccatctgaa	agaaaggctg	aatatgttcc	tgcagctgta	taacgagtcc	8760
atccggggag	ccggcatgga	tatggtgttc	tttgccgagc	ccatggtgca	cctcgtgaag	8820
atcacgagag	tgatccggac	cccccagggc	aacgctctgc	tcgtgggagt	gggagcctc	8880
ggcaagcaga	gcctgaccag	actggccagc	tttatcgccg	gctacgtgtc	cttccagatc	8940
accctgaccc	ggtcctacaa	caccagcaac	ctgatggaag	atctgaaggt	gctgtaccgg	9000
acagccggcc	agcaggggaa	gggcatcacc	ttcatcttca	ccgacaatga	gatcaaggac	9060
gagtctttcc	tggagtatat	gaacaatgtg	ctgagcagcg	gcgaggtgtc	caacctgttc	9120
gcccgggagc	agatcgacga	gattaacagc	gacctggcct	ccgtgatgaa	gaaagaattc	9180
ccccgggtgc	tgcccacaaa	cgagaacctg	cacgactact	tcatgtccag	agtgcggcag	9240
aatctgcaca	tcgtgctgtg	cttcagcccc	gtgggcgaga	agttcagaaa	ccgggcccctg	9300
aagttccccg	ccctgatcag	cggtgtcacc	atcgactggt	tcagccggtg	gcctaaggat	9360
gccctggtgg	ccgtgtccga	gcactttctg	accagctacg	acatcgactg	cagcctggaa	9420
atcaagaaa	aggtggtgca	gtgcatgggc	agottccagg	acggcgtggc	cgagaatgac	9480
gtggactact	tcagcgggtt	ccggcggagc	acccactgta	cccctaagag	ctacctgagc	9540
ttcatccagg	gtacaagtt	catctacggc	gagaagcacg	tggaagtgcg	cacctggcc	9600
aaccggatga	acaccggcct	ggaaaaactg	aaagaggcct	ccgagagcgt	ggcccgcctc	9660
agcaaaaga	tcgaaagcaa	agaaaaagaa	ctgcaggtg	ccaacgataa	ggccgacatg	9720
gtgctgaaa	aagtgacct	gaaggcccag	gccgcccaga	aagtgaagc	cgaggtgcag	9780
aaagtgaagg	accgggccc	ggccatctg	gactccatca	gcaaggacaa	ggccattgcc	9840
gaggaaaagc	tggaagcagc	caagcccggc	ctggaagagg	cagaagctgc	tctgcagacc	9900
atccggccct	ccgatattgc	cacagtgcgg	accctgggaa	ggccccctca	cctgatcatg	9960
cggatcatgg	actgtgtgct	gctgctgttc	cagagaaagg	tgtccgctg	gaagatcgac	10020
ctggaaaaat	cctgcaccat	gcctagctg	caggaatccc	tgaagctgat	gaccgcccgg	10080
aaacttccctg	agaacctgca	gcagttcccc	aaggacacca	tcaatgagga	agtgatcgag	10140
ttcctgagcc	cctactctga	gatgcccggc	tacaatatcg	aaaccgcaa	acgcgtgtgc	10200
ggcaacgtgg	ccggactgtg	ctcttgagcc	aaggctatgg	ctagcttctt	tagcattaac	10260
aaagaggtgc	tgctctgaa	ggccaacctg	gtggtgcagg	aaaaccggca	tctgctggcc	10320
atgcaggacc	tgagaaaagc	ccaggccgag	ctggacgata	agcaggtgta	gctggatgtg	10380
gtgcaggccg	agtacgagca	ggccatgacc	gagaagcaga	ccctgctgga	agatgcagag	10440
cggtgcagac	acaagatgca	gaccgcccag	acocctgatc	ctggactggc	cggcgaaaaa	10500
gagcgggtgga	ccgagcagtc	ccaggaattc	gccgcccaga	ccaagcggct	cgtgggagat	10560
gtgctgctgg	ccaccgctt	tctgagctac	agcggcccc	tcaatcagga	attcagggac	10620
ctgctgctga	acgactggcg	gaaagagatg	aaggccagaa	agatccccct	cggcaagaat	10680
ctgaacctga	gcgagatgct	gatcgacgcc	cccaccatct	ccgagtggaa	tctgcaggg	10740
ctgcccacag	atgacctgtc	catccagaac	ggaatcatcg	tgaccaaaagc	ctccagatac	10800
ccctgctgga	ttgaccccca	gacacagggc	aagatttgg	tcaagaacaa	agagagccgg	10860
aacgagctgc	agatcaccag	cctgaaccac	aagtacttcc	ggaaccacct	ggaagatagc	10920

		<p>ctgagcctgg gcagggcact gctgatcgag gatgtgggag aggaactgga cccagccctg 10980 gataacgtgc tggaacggaa cttcatcaag accggctcca ccttcaaagt gaaagtgggc 11040 gacaaagaag tggacgtgct ggatggcttc cggtgtaca tcaccaccaa gctgcctaac 11100 cccgcctaca ccctgagat cagcgcccg accagcatca tcgacttcac cgtgacaatg 11160 aagggactgg aagatcagct gctgggacgc gtgatcctga cagagaagca ggaactggaa 11220 aaagaacgga cccatctgat ggaagatgtg accgccaaca agcggcggat gaaggactg 11280 gaagataacc tgctgtacag gctgaccag acccagggga gtctgggtga agatgagagc 11340 ctgatcgtgg tgctgtccaa caccacgcgg accgcagagg aagtgaacca gaagctggaa 11400 atcagcgccg agacagaggt gcagatcaac agcggccagag aagagtaccg gcctgtggcc 11460 accgggggat ccatcctgta ctttctgatc accgagatgc ggctcgtgaa cgagatgtac 11520 cagaccagcc tgcggcagtt cctgggctctg ttogatctgt ccctggccag aagcgtgaa 11580 tcccccatca ccagcaagag aatcgccaac atcatcgagc acatgacctg cgaggtgtac 11640 aaatacgccg ccagaggcct gtacgaggaa cacaagtttc tgttcacact gctgtgacc 11700 ctgaagatcg atatccagcg gaacagagtg aagcacgaag agtttctgac actgatcaag 11760 gggggagcct ccctggacaa gaaggcctgt cctccaagc ccagcaagtg gatcctggac 11820 atcacctggc tgaatctggt ggaactgagc aagctgagac agttctccga tgtgtggac 11880 cagatcagcc gcaacgagaa gatgtggaag atttggtttg acaaagagaa ccccaggaa 11940 gaaccctgc ctaacgccta cgataagagc ctggactgct tccggcggct gctgtgatt 12000 agaagctggt gtcccagccg gacaatcgcc caggcccgcga agtacatcgt ggatagcatg 12060 ggagagaagt acgcccaggg cgtgatcctg gacctggaaa agacctggga ggaagcgac 12120 cccagaacc cctgatctg cctgctgagc atgggctcgc accccaccga cagcattatc 12180 gcctgggca agagactgaa gattgagaca agatacgtgt ccattgggcca gggccaggaa 12240 gtgcacgcta gaaagctgct gcagcagact atggccaatg gcggctgggc cctgtgagc 12300 aattgtcacc tggggctgga cttcatggac gaactgatgg acatcatcat tgagacagag 12360 ctggtgcagc acgcttcag actgtggatg accaccgagg cccataagca gtttccatt 12420 accctgctgc agatgagcat caagttcgcc aacgaccccc ctcagggact gagagccggc 12480 ctgaagagaa cctactccgg cgtgtcacag gatctgctgg acgtgtcctc tggcagccag 12540 tggaaagccta tgctgtacgc cgtggcattc ctgacacaga ccgtgcagga acggcggaag 12600 tttggcgccc tgggatggaa catcccctac gagtttaacc aggcgactt caacgccact 12660 gtgcagttta tccagaacca tctggacgac atggagctga agaaaggggt gtccctggaca 12720 accatccggt acatgatcgg agagatccag taoggcgcca gactgaccga cgactacgac 12780 aagaggctgc tgaatacctt cgccaaagtg tggttctccg agaacatggt tggccccgac 12840 ttcagctttt accagggcta taacatcccc aagtgtccca ccgtggataa ctacctgagc 12900 tacctccaga gcctgcccgc ctacgacagc cctgaggtgt tcggactgca ccccaaccg 12960 gatatacct accagagcaa actggccaag gatgtgctgg ataccatcct gggcatccag 13020 cccaaggata ccagtggcgg aggcgacgaa acccggaag cagtgtggc tagactggcc 13080 gacgacatgc tggaaaagct gccccccgac tacgtgccc tgaagtga agaacgctg 13140 cagaagatgg gcccttcca gcctatgaac atcttctga ggcaggaaat cgaccggatg 13200 cagcgggtgc tgtctctcgt gcggagcaca ctgaccgagc tgaactggc tatcgacggc 13260 accatcatca tgagcgagaa tctgcccggat gcaactggact gcatgttcca cgccagaatc 13320 cccgatggt ggaaaaaggc cagctggatc agctctacc tgggcttctg gttaccgga 13380 ctgatcgaga gaaacagcca gttaccagc tgggtgttca acggcagacc tcaactgttc 13440 tggatgaccg gcttcttcaa tccacaaggc tttctgacag caatgcgcca ggaatcacc 13500 agagccaaca agggctgggc tctggacaat atgggtgctgt gtaacgaagt gactaagtgg 13560 atgaaggacg acatcagcgc cctcccaca gagggcgtgt acgtgtacgg cctgtacctg 13620 gaaggcgcgg gatgggacaa gagaacatg aagctgatcg agagcaagcc caaggtgctg 13680 ttcgagctga tgcctgatg caggatctat gccgagaaca acacctgag ggacccccg 13740 ttctacagct gcccctcta caagaaacc gtgcgcaacc acctgaacta tatcgccgc 13800 gtggacctga ggacagcca gacacctgag cattgggtgc tgagaggcgt ggcactgctg 13860 tgcgacgtga agtga 13875</p>
3-19	Sequences	
3-19-1	Sequence Number [ID]	19
3-19-2	Molecule Type	RNA
3-19-3	Length	10
3-19-4	Features	misc_feature 1..10
	Location/Qualifiers	note=Description of Artificial Sequence: Synthetic oligonucleotide source 1..10 mol_type=other RNA organism=synthetic construct
3-19-5	NonEnglishQualifier Value Residues	gcccrcatgg 10
3-20	Sequences	
3-20-1	Sequence Number [ID]	20
3-20-2	Molecule Type	DNA
3-20-3	Length	11
3-20-4	Features	misc_feature 1..11
	Location/Qualifiers	note=Description of Artificial Sequence: Synthetic oligonucleotide source 1..11 mol_type=other DNA organism=synthetic construct
3-20-5	NonEnglishQualifier Value Residues	gaattctgca g 11