

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
3 February 2005 (03.02.2005)

PCT

(10) International Publication Number
WO 2005/010142 A2

- (51) International Patent Classification⁷: **C12N**
- (21) International Application Number:
PCT/US2003/020381
- (22) International Filing Date: 27 June 2003 (27.06.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (71) Applicant: **CHROMATIN, INC.** [US/US]; 2255 West Harrison St., Chicago, IL 60612 (US).
- (72) Inventors: **MACH, Jennifer**; 2958 North Kilbourn Avenue, Chicago, IL 60641 (US). **ZIELER, Helge**; 910 West VanBuren Street, #194, Chicago, IL 60607 (US). **JIN, RongGuan**; 837 East 56th Street, Chicago, IL 60637 (US). **KEITH, Kevin**; 1320 South Federal Street, Unit F, Chicago, IL 60605 (US). **COPENHAVER, Gregory**; 112 Windover Drive, Chapel Hill, NC 27514 (US). **PREUSS, Daphne**; 5807 South Dorchester, Apt. 2E, Chicago, IL 60637 (US).
- (74) Agent: **BARRETT, Robert**; Bell, Boyd & Lloyd LLC, P.O. Box 1135, Chicago, IL 60690 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PLANT CENTROMERE COMPOSITIONS

(57) Abstract: The present invention provides for the nucleic acid sequences of plant centromeres. This will permit construction of stably inherited recombinant DNA constructs and minichromosomes which can serve as vectors for the construction of transgenic plant and animal cells.

WO 2005/010142 A2

SPECIFICATION

TITLE OF THE INVENTION

“Plant Centromere Compositions”

5 This application is a continuation-in-part of U.S. Patent Application Serial No. 09/553,231, filed March 13, 2002, which is a continuation of U.S. Patent Application Serial No. 09/090,051, filed June 3, 1998, now U.S. Patent No. 6,156,953 which claims the priority of U.S. Provisional Patent Application Serial No. 60/048,451, filed June 3, 1997; and U.S. Provisional Patent Application Serial No. 60/073,741, filed
10 February 5, 1998, both of the disclosures of which are specifically incorporated herein by reference in their entirety. This application is also a continuation-in-part of U.S. Patent Application Serial No. 09/531,120, filed March 17, 2000, which claims the priority of U.S. Provisional Application Ser. No. 60/125,219, filed March 18, 1999; U.S. Provisional Application Ser. No. 60/127,409, filed April 1, 1999; U.S.
15 Provisional Application Ser. No. 60/134,770, filed May 18, 1999; U.S. Provisional Application Ser. No. 60/153,584, filed September 13, 1999, U.S. Provisional Application Ser. No. 60/154,603, filed September 17, 1999 and U.S. Provisional Application Ser. No. 60/172,493, filed December 16, 1999, each of which disclosures is specifically incorporated herein by reference in its entirety.

20

 The government owns rights in the present invention pursuant to U.S. Department of Agriculture Grant No. 96-35304-3491 and Grant No. DE-FC05-920R22072 from the Consortium for Plant Biotechnology Research, National Science Foundation Grant No. 9872641, and Department of Energy Small Business Innovation
25 Research Grants DE-FG02-01ER83163, DE-FG02-01ER83165, and DE-FG02-01ER83166.

BACKGROUND OF THE INVENTION

30 Two general approaches are used for introduction of new genetic information (“transformation”) into cells. One approach is to introduce the new genetic information as part of another DNA molecule, referred to as an “episomal vector,” or

“minichromosome”, which can be maintained as an independent unit (an episome) apart from the host chromosomal DNA molecule(s). Episomal vectors contain all the necessary DNA sequence elements required for DNA replication and maintenance of the vector within the cell. Many episomal vectors are available for use in bacterial
5 cells (for example, see Maniatis *et al.*, 1982). However, only a few episomal vectors that function in higher eukaryotic cells have been developed. Higher eukaryotic episomal vectors were primarily based on naturally occurring viruses. In higher plant systems gemini viruses are double-stranded DNA viruses that replicate through a double-stranded intermediate upon which an episomal vector could be based, although
10 the gemini virus is limited to an approximately 800 bp insert. Although an episomal plant vector based on the Cauliflower Mosaic Virus has been developed, its capacity to carry new genetic information also is limited (Brisson *et al.*, 1984).

The other general method of genetic transformation involves integration of
15 introduced DNA sequences into the recipient cell's chromosomes, permitting the new information to be replicated and partitioned to the cell's progeny as a part of the natural chromosomes. The introduced DNA usually is broken and joined together in various combinations before it is integrated at random sites into the cell's chromosome (see, for example Wigler *et al.*, 1977). Common problems with this procedure are the
20 rearrangement of introduced DNA sequences and unpredictable levels of expression due to the location of the transgene in the genome or so called “position effect variation” (Shingo *et al.*, 1986). Further, unlike episomal DNA, integrated DNA cannot normally be precisely removed. A more refined form of integrative transformation can be achieved by exploiting naturally occurring viruses that integrate
25 into the host's chromosomes as part of their life cycle, such as retroviruses (see Cepko *et al.*, 1984). In mouse, homologous integration has recently become common, although it is significantly more difficult to use in plants (Lam *et al.* 1996).

The most common genetic transformation method used in higher plants is
30 based on the transfer of bacterial DNA into plant chromosomes that occurs during infection by the phytopathogenic soil bacterium *Agrobacterium* (see Nester *et al.*, 1984). By substituting genes of interest for the naturally transferred

bacterial sequences (called T-DNA), investigators have been able to introduce new DNA into plant cells. However, even this more "refined" integrative transformation system is limited in three major ways. First, DNA sequences introduced into plant cells using the *Agrobacterium* T-DNA system are frequently rearranged (see Jones *et al.*, 1987). Second, the expression of the introduced DNA sequences varies between individual transformants (see Jones *et al.*, 1985). This variability is presumably caused by rearranged sequences and the influence of surrounding sequences in the plant chromosome (*i.e.*, position effects), as well as methylation of the transgene. A third drawback of the *Agrobacterium* T-DNA system is the reliance on a "gene addition" mechanism: the new genetic information is added to the genome (*i.e.*, all the genetic information a cell possesses) but does not replace information already present in the genome.

One attractive alternative to commonly used methods of transformation is the use of an artificial chromosome. Artificial chromosomes are man-made linear or circular DNA molecules constructed from cis-acting DNA sequence elements that provide replication and partitioning of the constructed chromosomes (see Murray *et al.*, 1983). Desired elements include: (1) Autonomous Replication Sequences (ARS) (these have properties of replication origins, which are the sites for initiation of DNA replication), (2) Centromeres (site of kinetochore assembly and responsible for proper distribution of replicated chromosomes at mitosis or meiosis), and (3) if the chromosome is linear, telomeres (specialized DNA structures at the ends of linear chromosomes that function to stabilize the ends and facilitate the complete replication of the extreme termini of the DNA molecule).

25

The essential chromosomal elements for construction of artificial chromosomes have been precisely characterized in lower eukaryotic species, and more recently in mouse and human. ARSs have been isolated from unicellular fungi, including *Saccharomyces cerevisiae* (brewer's yeast) and *Schizosaccharomyces pombe* (see Stinchcomb *et al.*, 1979 and Hsiao *et al.*, 1979). An ARS behaves like a replication origin allowing DNA molecules that contain the ARS to be replicated as an episome after introduction into the cell nuclei of these fungi. DNA molecules containing these

sequences replicate, but in the absence of a centromere they are partitioned randomly into daughter cells.

Artificial chromosomes have been constructed in yeast using the three cloned
5 essential chromosomal elements. Murray *et al.*, 1983, disclose a cloning system based on the in vitro construction of linear DNA molecules that can be transformed into yeast, where they are maintained as artificial chromosomes. These yeast artificial chromosomes (YACs) contain cloned genes, origins of replication, centromeres and telomeres and are segregated in daughter cells with high fidelity when the YAC is at
10 least 100 kB in length. Smaller CEN-containing vectors may be stably segregated, however, when in circular form.

None of the essential components identified in unicellular organisms, however, function in higher eukaryotic systems. For example, a yeast CEN sequence will not
15 confer stable inheritance upon vectors transformed into higher eukaryotes. While such DNA fragments can be readily introduced, they do not stably exist as episomes in the host cell. This has seriously hampered efforts to produce artificial chromosomes in higher organisms.

20 In one case, a plant artificial chromosome was discussed (Richards *et al.*, U.S. Patent No. 5,270,201). However, this vector was based on plant telomeres, as a functional plant centromere was not disclosed. While telomeres are important in maintaining the stability of chromosomal termini, they do not encode the information needed to ensure stable inheritance of an artificial chromosome. It is well documented
25 that centromere function is crucial for stable chromosomal inheritance in almost all eukaryotic organisms (reviewed in Nicklas 1988). For example, broken chromosomes that lack a centromere (acentric chromosomes) are rapidly lost from cell lines, while fragments that have a centromere are faithfully segregated. The centromere accomplishes this by attaching, via centromere binding proteins, to the spindle fibers
30 during mitosis and meiosis, thus ensuring proper gene segregation during cell divisions.

In contrast to the detailed studies done in *S. cerevisiae* and *S. pombe*, less is known about the molecular structure of functional centromeric DNA of higher eukaryotes. Ultrastructural studies indicate that higher eukaryotic kinetochores, which are specialized complexes of proteins that form on the centromere during late
5 prophase, are large structures (mammalian kinetochore plates are approximately 0.3 μm in diameter) which possess multiple microtubule attachment sites (reviewed in Rieder, 1982). It is therefore possible that the centromeric DNA regions of these organisms will be correspondingly large, although the minimal amount of DNA necessary for centromere function may be much smaller.

10

The above studies have been useful in elucidating the structure and function of centromeres-. The extensive literature indicating both the necessity of centromeres for stable inheritance of chromosomes, and the non-functionality of yeast centromeres in higher organisms, demonstrate that cloning of a functional centromere from a higher
15 eukaryote is a necessary first step in the production of artificial chromosomes suitable for use in higher plants and animals. The production of artificial chromosomes with centromeres which function in higher eukaryotes would overcome many of the problems associated with the prior art and represent a significant breakthrough in biotechnology research.

20

SUMMARY OF THE INVENTION

The present invention allows the isolation and identification of plant centromere DNA sequences from the total genomic DNA of an organism or fractions
25 thereof. With centromere DNA sequences, it is possible to construct chromosomes having functional centromeres and carrying large number of genes. Genes for producing a vast set of products have been identified, but technologies used within the industry severely limit the delivery of these genes to plant cells. One or at most a few genes are typically inserted into random locations in the host chromosomes, which can
30 irreversibly disrupt host gene functions while causing variable and uncontrolled expression of the introduced genes. The present invention makes it possible to

overcome the technical limitations associated with gene delivery in crop species, thereby allowing for the ability to shorten the time required for crop development.

In one aspect, the invention provides a method to obtain a centromere DNA
5 sequence from a selected organism, the method comprising the steps of preparing a sample of genomic DNA from a selected organism, obtaining a plurality of nucleic acid segments from the genomic DNA and screening the nucleic acid segments to identify one or more centromere nucleic acid sequences. In an embodiment, the method of obtaining the plurality of nucleic acid segments comprises contacting said
10 genomic DNA with a restriction endonuclease and selecting nucleic acid segments containing repetitive DNA to obtain said plurality of nucleic acid segments. In another embodiment, the method of obtaining the plurality of nucleic acid segments comprises contacting said genomic DNA with a methylation sensitive restriction endonuclease and selecting nucleic acid segments exhibiting resistance to cleavage with said
15 methylation sensitive restriction endonuclease to obtain said plurality of nucleic acid segments. In yet another embodiment, the method of obtaining the plurality of nucleic acid segments comprises contacting said genomic DNA with a restriction endonuclease or physically shearing said genomic DNA and selecting nucleic acid segments that anneal rapidly after denaturation to obtain said plurality of nucleic acid
20 segments.

In another aspect, the invention provides a method for identifying a centromere nucleic acid sequence from a dataset of the genomic sequences of an organism. The method comprises the steps of (1) providing a first dataset consisting of the genomic
25 sequences, or a representative fraction of genomic sequence, of the organism; (2) identifying and eliminating known non-centromeric repeat sequences from the first dataset by using the BLAST sequence comparison algorithm to create a second dataset; (3) comparing each sequence in the second dataset to itself by using the BLAST sequence comparison algorithm, obtaining a BLAST score for each pair of
30 sequence compared, and collecting high score pairs to create a third dataset; (4) examining the BLAST score of each high score pair in the third dataset and eliminating the pairs having a score greater than 10^{-20} to create a fourth dataset; (5)

eliminating the high score pairs in the fourth dataset having less than 80 bp or more than 250 bp to create a fifth dataset; (6) examining the nucleotide position of each high score pair in the fifth dataset and eliminating pairs having 100% identity as well as identical nucleotide positions to create a sixth dataset; (7) examining the nucleotide position of each high score pair in the sixth dataset and eliminating pairs having opposite orientation of the nucleotides to create a seventh dataset; (8) examining the nucleotide position of both sequences for each high score pair in the seventh dataset and eliminating sequences that are overlapping to create an eighth dataset; and (9) examining the nucleotide position of each sequence in the eighth dataset and eliminating sequences not having at least one neighboring sequence within 250 bp to create a ninth dataset; and (10) comparing each sequence in the ninth dataset to all other sequences in the ninth dataset by using the BLAST sequence comparison algorithm and selecting the most common sequence as a centromere sequence of the organism. In one embodiment, the known non-centromeric repeat sequence in the second step is a ribosomal DNA.

In another aspect, the invention provides a *Brassica oleracea* centromere comprising *Brassica oleracea* centromere DNA. In one embodiment, the *Brassica oleracea* centromere is defined as comprising n copies of a repeated nucleotide sequence, wherein n is at least 2. Potentially any number of repeat copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100, 150, 200, 300, 400, 500, 750, 1,000, 1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 and about 100,000, including all ranges in-between such copy numbers. In one embodiment, the repeated nucleotide sequence is isolated from *Brassica oleracea* given by SEQ ID NO:1, 2, 3, or 4.

In yet another aspect, the invention provides a *Glycine max* centromere comprising *glycine max* centromere DNA. In an embodiment, the *Glycine max* centromere is defined as comprising n copies of a repeated nucleotide sequence, wherein n is at least 2. Potentially any number of repeat copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100, 150, 200, 300, 400, 500, 750, 1,000,

1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 and about 100,000, including all ranges in-between such copy numbers. In one embodiment, the repeated nucleotide sequence is isolated from *Glycine max* given by SEQ ID NO:5, 6, 7, or 8.

5

In yet another aspect, the invention provides a *Lycopersicon esculentum* centromere comprising *Lycopersicon esculentum* centromere DNA. In an embodiment, the *Lycopersicon esculentum* centromere is defined as comprising n copies of a repeated nucleotide sequence, wherein n is at least 2. Potentially any number of repeat
10 copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100, 150, 200, 300, 400, 500, 750, 1,000, 1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 and about 100,000, including all ranges in-between such copy numbers. In one embodiment, the repeated nucleotide
15 sequence is isolated from *Lycopersicon esculentum* given by SEQ ID NO:9 or 10.

In yet another aspect, the invention provides a *Zea mays* centromere comprising *Zea mays* centromere DNA. In an embodiment, the centromere is defined as comprising n copies of a repeated nucleotide sequence, wherein n is at least 2.
20 Potentially any number of repeat copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100, 150, 200, 300, 400, 500, 750, 1,000, 1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 and about 100,000, including all ranges in-between such copy numbers. In one
25 embodiment, the repeated nucleotide sequence is isolated from *Zea mays* given by SEQ ID NO:11, 12 or 13.

In yet another aspect, the invention provides a recombinant DNA construct comprising a plant centromere sequence of the present invention. The recombinant
30 DNA construct may additionally comprise any other desired sequences, for example, a telomere. Still further, one may wish to include a structural gene on the construct, or multiple genes. Examples of structural genes one may wish to use include a selectable

or screenable marker gene, an antibiotic resistance gene, a ligand gene, an enzyme gene, a herbicide resistance gene, a nitrogen fixation gene, a plant pathogen defense gene, a plant stress-induced gene, a toxin gene, a receptor gene, a gene encoding an enzyme, a gene encoding an antibody, a gene encoding an antigen for a vaccine, a
5 transcription factor, a cytoskeletal protein, a DNA-binding protein, a protease, an endonuclease, a lipid, a seed storage gene, an interleukin gene, a clotting factor gene, a cytokine gene, a growth factor gene and a biosynthetic gene for producing pharmaceutically active proteins, small molecules with medicinal properties, chemicals with industrial utility, nutraceuticals, carbohydrates, RNAs, lipids, fuels,
10 dyes, pigments, vitamins, scents, flavors, vaccines, antibodies, and hormones. In one embodiment of the invention, the construct is capable of expressing the structural gene, for example, in a prokaryote or eukaryote, including a lower eukaryote, or a higher eukaryote such as a plant. Moreover, the recombinant construct could contain other useful non-coding sequences, including promoters, terminators, boundary
15 elements that regulate gene expression, sequences that alter maintenance, inheritance, or stability of the construct, and sequences that allow subsequent modification of the composition of the construct.

In still yet another aspect, the invention provides a recombinant DNA construct
20 comprising a plant centromere sequence of the present invention and which is capable of being maintained as a chromosome, wherein the chromosome is transmitted in dividing cells. The plant centromere may be from any plant or may be from any other source of DNA or may be partially or entirely synthetic in origin.

25 In yet another aspect, the invention provides a recombinant DNA construct comprising a plant centromere sequence of the present invention and which is a plasmid. The plasmid may contain any desired sequences, such as an origin of replication. The plasmid may also comprise a selection marker.

30 In still yet another aspect, the invention provides a minichromosome comprising a plant centromere sequence of the present invention and may also contain a telomere sequence. Any additional desired sequences may be added to the

minichromosome, such as an autonomous replicating sequence and a structural gene such as those described above. The minichromosome may comprise any of the centromere compositions disclosed herein.

5 The minichromosome also may contain “negative” selectable markers which confer susceptibility to an antibiotic, herbicide or other agent, thereby allowing for selection against plants, plant cells or cells of any other organism of interest containing a minichromosome. The minichromosome also may include genes or other sequences which control the copy number of the minichromosome within a cell. One or more
10 structural genes also may be included in the minichromosome. Specifically contemplated as being useful will be as many structural genes as may be inserted into the minichromosome.

 In still yet another aspect, the invention provides a cell transformed with a
15 recombinant DNA construct comprising a plant centromere sequence of the present invention. The cell may be of any type, including a prokaryotic cell or eukaryotic cell. Where the cell is a eukaryotic cell, the cell may be, for example, a yeast cell or a higher eukaryotic cell, such as plant cell. The plant cell may be from a dicotyledonous plant, such as tobacco, tomato, potato, soybean, canola, sunflower, alfalfa, cotton and
20 *Arabidopsis*, or may be a monocotyledonous plant cell, such as wheat, maize, rye, rice, turfgrass, oat, barley, sorghum, millet, and sugarcane. In one embodiment of the invention, the plant centromere is a centromere chosen from the group consisting of *Brassica oleracea*, *Glycine max*, *Lycopersicon esculentum*, and *Zea mays* and the cell may be a cell chosen from one of the above species or any other species. The
25 recombinant DNA construct may comprise additional sequences, such as a telomere, an autonomous replicating sequence (ARS), a structural gene or genes, or a selectable or screenable marker gene or genes, including as many of such sequences as may physically be placed on said recombinant DNA construct. In one embodiment of the invention, the cell is further defined as capable of expressing said structural gene. In
30 another embodiment of the invention, a plant is provided comprising the aforementioned cells.

In still yet another aspect, the invention provides a method for preparing a transgenic plant cell. The method comprises the steps of contacting a starting plant cell with a recombinant DNA construct comprising a plant centromere sequence of the present invention, whereby the starting plant cell is transformed with the recombinant
5 DNA construct.

In still yet another aspect, the invention provides a transgenic crop comprising a minichromosome, wherein the minichromosome comprises a plant centromere sequence of the present invention. The minichromosome may further comprise a
10 telomere sequence, an autonomous replicating sequence or a structural gene, such as a selectable or screenable marker gene, an antibiotic resistance gene, a ligand gene, an enzyme gene, a herbicide resistance gene, a nitrogen fixation gene, a plant pathogen defense gene, a plant stress-induced gene, a toxin gene, a receptor gene, a gene encoding an enzyme, a gene encoding an antibody, a gene encoding an antigen for a
15 vaccine, a transcription factor, a cytoskeletal protein, a DNA-binding protein, a protease, an endonuclease, a lipid, a seed storage gene, an interleukin gene, a clotting factor gene, a cytokine gene, a growth factor gene and a biosynthetic gene for producing pharmaceutically active proteins, small molecules with medicinal properties, chemicals with industrial utility, nutraceuticals, carbohydrates, RNAs,
20 lipids, fuels, dyes, pigments, vitamins, scents, flavors, vaccines, antibodies, and hormones. The transgenic crop may be any type of crop, such as a dicotyledonous plant, for example, tobacco, tomato, potato, pea, carrot, cauliflower, broccoli, soybean, canola, sunflower, alfalfa, cotton and *Arabidopsis*, or may be a monocotyledonous plant, such as wheat, maize, rye, rice, turfgrass, oat, barley, sorghum, millet, and
25 sugarcane.

In still yet another aspect, the invention provides a method for preparing a transgenic crop tissue. The method comprises the steps of contacting a starting crop tissue with a recombinant DNA construct comprising a plant centromere sequence of
30 the present invention, whereby the starting crop tissue is transformed with the recombinant DNA construct.

In still yet another aspect, the invention provides a method for preparing a transgenic crop seed. The method comprises the steps of contacting a starting crop, crop tissue, or crop cell, with a recombinant DNA construct comprising a plant centromere sequence of the present invention, whereby the starting crop, crop tissue, or crop cell is transformed with the recombinant DNA construct. These transformed crops, crop tissues, or crop cells are allowed to develop into mature crops, using standard agricultural techniques. Transgenic seed is then collected from these crops.

In still yet another aspect, the invention provides a method for preparing an extract of a transgenic crop, crop tissue, crop seed, or crop cell. The method comprises the steps of contacting a starting crop, crop tissue, or crop cell with a recombinant DNA construct comprising a plant centromere sequence of the present invention, whereby the starting crop cell is transformed with the recombinant DNA construct. The resulting transgenic crop, crop tissue, crop seed, or crop cell is then extracted and processed to yield the desirable product. One preferred desirable product is a food product. Another preferred desirable product is a pharmaceutical product. Yet another preferred desirable product is a chemical product.

Additional features and advantages of the present invention are described in, and will be apparent from, the following Detailed Description of the Invention and the figures.

BRIEF DESCRIPTION OF THE FIGURES

The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

FIG. 1A-F Consensus sequences of repeats from *Brassica oleracea*. FIG. 1A is the consensus sequence of ChrBo1. This consensus was assembled from 33 sequences

collected by the inventors. The length of this repeat is 180 ± 0.86 base pairs and A and T compose 60% of the consensus. FIG. 1B is the consensus sequence of ChrBo2. This consensus was assembled from 7 sequences collected by the inventors. The length of this repeat is 180 ± 0.45 base pairs and A and T compose 63% of the consensus. FIG. 1C is a comparison of the consensus sequences of ChrBo1 and ChrBo2. The two repeats (ChrBo1 and ChrBo2) were aligned to each other using the ClustalX program (ClustalX is a free multiple sequence alignment program for Windows. Those sites with significant differences between the two sequences (Chi-squared, $P < 0.05$) are highlighted. FIG. 1D is a revised consensus sequence of ChrBo1. This consensus was assembled from 33 DNA sequences collected by the inventors and 18 sequences from Genbank, identified by the accession numbers:

M30962	M30963	M31436	M31435
M31438	M31434	M31439	M31437
X68786	X12736	X07519	X16589
X15291	X68783	X68784	X61583
AJ228348	Z22947		

FIG. 1E is a revised consensus sequence of ChrBo2. This consensus was assembled from 7 DNA sequences collected by the inventors and 5 sequences from Genbank, identified by the accession numbers AJ228347, M30962, X12736, X61583, and X68785. FIG. 1F is a comparison of the revised consensus sequences of ChrBo1 and ChrBo2, aligned as for FIG. 1C.

FIG. 2A-F Consensus sequences of repeats from *Glycine max*. FIG. 2A is a consensus sequence of ChrGm1. This consensus was assembled from 32 sequences collected by the inventors. The length of this repeat is 92 ± 0.79 base pairs and A and T compose 63% of the consensus. FIG. 2B is a consensus sequence of ChrGm2. This consensus was assembled from 21 sequences collected by the inventors. The length of this repeat is 91 ± 0.48 base pairs and A and T compose 62% of the consensus. FIG. 2C is a comparison of the consensus sequences of ChrGm1 and ChrGm2. The two repeats (ChrGm1 and ChrGm2) were aligned to each other using the ClustalX program (ClustalX is a free multiple sequence alignment program for Windows. Those sites with significant differences between the two sequences (Chi-squared, $P < 0.05$) are highlighted. FIG. 2D is a revised consensus sequence of ChrGm1. This consensus

was assembled from 32 DNA sequences collected by the inventors and 1 sequence from Genbank, identified by the accession number Z26334. FIG. 2E is a revised consensus sequence of ChrGm2. This consensus was assembled from 21 DNA sequences collected by the inventors and 13 sequences from Genbank, identified by the accession numbers AF297983, AF297984, and AF297985. FIG. 2F is a comparison of the revised consensus sequences of ChrGm1 and ChrGm2, aligned as for FIG. 2C.

FIG. 3A-B Consensus sequences of repeats from *Lycopersicon esculentum*. FIG. 3A is a consensus sequence of ChrLe1. This consensus was assembled from 42 sequences collected by the inventors. The length of this repeat is 181 0.61 base pairs and A and T compose 50% of the consensus. FIG. 3B is a revised consensus sequence of ChrLe1. This consensus was assembled from 32 sequences collected by the inventors and 2 Genbank sequences identified by the accession numbers X87233 and AY007367.

FIG. 4A-C Consensus sequences of repeats from *Zea mays*. FIG. 4A is a consensus sequence of ChrZm1. This consensus was assembled from 38 sequences collected by the inventors. The length of this repeat is 180 1.15 base pairs and A and T compose 56% of the consensus. FIG. 4B is a revised consensus sequence of ChrZm1. This consensus was assembled from 38 sequences collected by the inventors and 26 sequences from Genbank, identified by the accession numbers:

M32521	M32522	M32523	M32524	M32525	M32526
M32527	M32528	M32529	M32530	M32531	M32532
M32533	M32534	M325375	M32536	M32537	M32538
M35408	AF030934	AF030935	AF030936	AF030937	AF030938
AF030939	AF030940				

FIG. 4C is a consensus sequence of ChrZm2. This consensus was assembled from 6 sequences collected from Genbank identified by the accession numbers:

AF078918	AF078919	AF078920
AF0789121	AF078922	AF078923

The length of this repeat is 158 1.6 base pairs and A and T compose 53% of

the consensus.

FIG. 5 Minichromosome containing centromere sequences as well as minichromosome vector sequences

5 **FIG. 6** Minichromosome construct formed by minichromosome vector tailing method.

FIG. 7A-7N. Exemplary Minichromosome vectors: The vectors shown in **FIG. 7A, FIG. 7B, FIG. 7E, FIG. 7F, FIG. 7I** and **FIG. 7J** have an *E. coli* origin of replication which can be high copy number, low copy number or single copy. In **FIGS. 7A-7N**, the vectors include a multiple cloning site which can contain recognition sequences for conventional restriction endonucleases with 4-8 bp specificity as well as recognition sequences for very rare cutting enzymes such as, for example, I-Ppo I, I-Cue I, PI-Tli, PI-Psp I, Not I, and PI Sce I. In **FIG. 7A-7N**, the centromere is flanked by Lox sites which can act as targets for the site specific recombinase Cre. **FIG. 7A.** Shows an *E. coli* plant circular shuttle vector with a plant ARS. **FIG. 7B.** Shows a plant circular vector without a plant ARS. The vector relies on a plant origin of replication function found in other DNA sequences such as selectable or screenable markers. **FIG. 7C.** Shows a yeast-plant circular shuttle vector with a plant ARS. The yeast ARS is included twice, once on either side of multiple cloning site to ensure that large inserts are stable. **FIG. 7D.** Shows a yeast-plant circular shuttle vector without a plant ARS. The vector relies on a plant origin of replication function found in other plant DNA sequences such as selectable markers. The yeast ARS is included twice, once on either side of the multiple cloning site to ensure that large inserts are stable. **FIG. 7E.** Shows an *E. coli-Agrobacterium*-plant circular shuttle vector with a plant ARS. Vir functions for T-DNA transfer would be provided in trans by a using the appropriate *Agrobacterium* strain. **FIG. 7F.** Shows an *E. coli-Agrobacterium*-plant circular shuttle vector without a plant ARS. The vector relies on a plant origin of replication function found in other plant DNA sequences such as selectable markers. Vir functions for T-DNA transfer would be provided in trans by a using the appropriate *Agrobacterium* strain. **FIG. 7G.** Shows a linear plant vector with a plant ARS. The linear vector could be assembled

in vitro and then transferred into the plant by, for example, mechanical means such as micro projectile bombardment, electroporation, or PEG-mediated transformation. **FIG. 7H.** Shows a linear plant vector without a plant ARS. The linear vector could be assembled *in vitro* and then transferred into the plant by, for example, mechanical means such as micro projectile bombardment, electroporation, or PEG-mediated transformation. **FIGs. 7I-7N.** The figures are identical to **FIGs. 7A-7F**, respectively, with the exception that they do not contain plant telomeres. These vectors will remain circular once delivered into the plant cell and therefore do not require telomeres to stabilize their ends.

10

FIG. 8. Sequence features at Arabidopsis *CEN2* (A) and *CEN4* (B). Central bars depict annotated genomic sequence of indicated BAC clones; black, genetically-defined centromeres; white, regions flanking the centromeres. Sequences corresponding to genes and repetitive features, filled boxes (above and below the bars, respectively), are defined as in FIG. 11A-T; predicted nonmobile genes, red; genes carried by mobile elements, black; nonmobile pseudogenes, pink; pseudogenes carried by mobile elements, gray; retroelements, yellow; transposons, green; previously defined centromeric repeats, dark blue; 180 bp repeats, pale blue. Chromosome-specific centromere features include a large mitochondrial DNA insertion (orange; *CEN2*), and a novel array of tandem repeats (purple; *CEN4*). Gaps in the physical maps (/), unannotated regions (hatched boxes), and expressed genes (filled circles) are shown.

FIG. 9. Method for converting a BAC clone (or any other bacterial clone) into a minichromosome. A portion of the conversion vector will integrate into the BAC clone (or other bacterial clone of interest) either through non-homologous recombination (transposable element mediated) or by the action of a site specific recombinase system, such as Cre-Lox or FLP-FRT.

FIG. 10A-G. Method for converting a BAC clone (or any other bacterial clone) into a minichromosome. The necessary selectable markers and origins of replication for propagation of genetic material in *E. coli*, *Agrobacterium* and

Arabidopsis as well as the necessary genetic loci for *Agrobacterium* mediated transformation into *Arabidopsis* are cloned into a conversion vector. Using Cre/loxP recombination, the conversion vectors are recombined into BACs containing centromere fragments to form minichromosomes.

5

FIG. 11A-T. Properties of centromeric regions on chromosomes II and IV of *Arabidopsis*. (Top) Drawing of genetically-defined centromeres (gray shading, *CEN2*, left; *CEN4*, right), adjacent pericentromeric DNA, and a distal segment of each chromosome, scaled in Mb as determined by DNA sequencing (gaps in the grey shading correspond to gaps in the physical maps). Positions in cM on the RI map (http://nasc.nott.ac.uk/new_ri_map.html) and physical distances in Mb, beginning at the northern telomere and at the centromeric gap, are shown. (Bottom) The density of each feature (**FIGs. 11A-11T**) is plotted relative to the position on the chromosome in Mb. (**FIG. 11A, 11K**) cM positions for markers on the RI map (solid squares) and a curve representing the genomic average of 1 cM/221 kb (dashed line). A single crossover within *CEN4* in the RI mapping population (http://nasc.nott.ac.uk/new_ri_map.html; Somerville and Somerville, 1999) may reflect a difference between male meiotic recombination monitored here and recombination in female meiosis. (**FIGs. 11B-11E and FIGs. 11L-11O**) The % of DNA occupied by repetitive elements was calculated for a 100 kb window with a sliding interval of 10 kb. (**FIGs. 11B, 11L**) 180 bp repeats; (**FIGs. 12C, 12M**) sequences with similarity to retroelements, including *del*, *Ta1*, *Ta11*, *copia*, *Athila*, *LINE*, *Ty3*, *TSCL*, 106B (*Athila*-like), *Tat1*, LTRs and *Cinful*; (**FIGs. 11D, 11N**) sequences with similarity to transposons, including *Tag1*, *En/Spm*, *Ac/Ds*, *Tam1* *MuDR*, *Limpet*, *MITEs* and *Mariner*; (**FIGs. 11E, 11O**) previously described centromeric repeats including 163A, 164A, 164B, 278A, 11B7RE, *mi167*, *pAT27*, 160-, 180- and 500-bp repeats, and telomeric sequences. (**FIGs. 11F, 11P**) % adenosine + thymidine was calculated for a 50 kb window with a sliding interval of 25 kb (**FIGs. 11G-11J, 11Q-11T**). The number of predicted genes or pseudogenes was plotted over a window of 100 kb with a sliding interval of 10 kb. (**FIGs. 11G, 11I, 11Q, 11S**) predicted genes (**FIGs. 11G, 11Q**) and pseudogenes (**FIGs. 11I, 11S**) typically not found on mobile DNA elements; (**FIGs. 11H, 11J, 11R, 11T**) predicted genes (**FIGs. 11H, 11R**) and pseudogenes

(FIGs. 11J, 11T) often carried on mobile DNA, including reverse transcriptase, transposase, and retroviral polypeptides. Dashed lines indicate regions in which sequencing or annotation is in progress, annotation was obtained from GenBank records (<http://www.ncbi.nlm.nih.gov/Entrez/nucleotide.html>), from the AGAD database (<http://www.tigr.org/tdb/at/agad/>), and by BLAST comparisons to the database of repetitive *Arabidopsis* sequences (<http://nucleus.cshl.org/protarab/AtRepBase.htm>); though updates to annotation records may change individual entries, the overall structure of the region will not be significantly altered.

10

FIG. 12. Methods for converting a BAC clone containing centromere DNA into a minichromosome for introduction into plant cells. The specific elements described are provided for exemplary purposes and are not limiting. A) diagram of the BAC clone, noting the position of the centromere DNA, a site-specific recombination site (for example, lox P), and the F origin of replication. B) Conversion vector containing selectable and color markers (for example, 35S-Bar, nptII, LAT52-GUS, Scarecrow-GFP), telomeres, a site-specific recombination site (for example, lox P), antibiotic resistance markers (for example, amp or spc/str), *Agrobacterium* T-DNA borders (Agro Left and Right) and origin of replication (RiA4). C) The product of site specific recombination with the Cre recombinase at the lox P sites yields a circular product with centromeric DNA and markers flanked by telomeres. D) Minichromosome immediately after transformation into plants; subsequently, the left and right borders will likely be removed by the plant cell and additional telomeric sequence added by the plant telomerase.

25

FIG. 13A-B. Conservation of *Arabidopsis* centromere DNA. BAC clones (bars) used to sequence *CEN2* (FIG. 13A) and *CEN4* (FIG. 13B) are indicated; arrows denote the boundaries of the genetically-defined centromeres. PCR primer pairs yielding products from only Columbia (filled circles) or from both Landsberg and Columbia (open circles); BACs encoding DNA with homology to the mitochondrial genome (gray bars); 180 bp repeats (gray boxes); unsequenced DNA (dashed lines); and gaps in the physical map (double slashes) are shown.

30

FIG. 14A-B. Primers used to analyze conservation of centromere sequences in the *A. thaliana* Columbia and Landsberg ecotypes. **FIG. 14A:** Primers used for amplification of chromosome 2 sequences. **FIG. 14B:** Primers used for amplification of chromosome 4 sequences.

FIG. 15. Sequences common to *Arabidopsis CEN2* and *CEN4*. Genetically-defined centromeres (bold lines), sequenced (thin lines), and unannotated (dashed lines) BAC clones are displayed as in FIG. 14A, B. Repeats AtCCS1 (*A. thaliana* centromere conserved sequence) and AtCCS2 (closed and open circles, respectively), AtCCS3 (triangles), and AtCCS4-7 (4-7, respectively) are indicated (GenBank Accession numbers AF204874 to AF204880), and were identified using BLAST 2.0 (<http://blast.wustl.edu>).

DETAILED DESCRIPTION OF THE INVENTION

The inventors have overcome the deficiencies in the prior art by providing the nucleic acid sequences of plant centromeres. The significance of this achievement relative to the prior art is exemplified by the general lack of detailed information in the art regarding the centromeres of multicellular organisms in general. To date, the most extensive and reliable characterization of centromere sequences has come from studies of lower eukaryotes such as *S. cerevisiae* and *S. pombe*, where the ability to analyze centromere functions has provided a clear picture of the desired DNA sequences. The *S. cerevisiae* centromere consists of three essential regions, CDEI, CDEII, and CDEIII, totaling only 125 bp, or approximately 0.006 to 0.06% of each yeast chromosome (Carbon *et al.*, 1990; Bloom 1993). *S. pombe* centromeres are between 40 and 100 kB in length and consist of repetitive elements that comprise 1 to 3% of each chromosome (Baum *et al.*, 1994). Subsequent studies, using tetrad analysis to follow the segregation of artificial chromosomes, demonstrated that less than 1/5 of the naturally occurring *S. pombe* centromere is sufficient for centromere function (Baum *et al.*, 1994).

In contrast, the centromeres of mammals and other higher eukaryotes are less understood. Although DNA fragments that hybridize to centromeric regions in higher eukaryotes have been identified, in many cases, little is known regarding the functionality of these sequences (see Tyler-Smith *et al.*, 1993). Centromere repeats
5 often correlate with centromere location, with probes to the repeats mapping both cytologically and genetically to centromere regions. Many of these sequences are tandemly-repeated satellite elements and dispersed repeated sequences in arrays ranging from 300 kB to 5000 kB in length (Willard 1990). To date, only one of these repeats, a 171 bp element known as the alphoid satellite, has been shown by in situ
10 hybridization to be present at each human centromere (Tyler-Smith *et al.*, 1993). Whether repeats themselves represent functional centromeres remains controversial, as other genomic DNA can be required to confer efficient inheritance upon a region of DNA (Willard, 1997). Alternatively, the positions of some higher eukaryotic centromeres have been estimated by analyzing the segregation of chromosome
15 fragments. This approach is imprecise, however, because a limited set of fragments can be obtained, and because normal centromere function is influenced by surrounding chromosomal sequences (for example, see Koornneef, 1983; FIG. 2).

A more precise method for mapping centromeres that can be used in intact
20 chromosomes is tetrad analysis (Mortimer *et al.*, 1981), which provides a functional definition of a centromere in its native chromosomal context. Centromeres that have been mapped in this manner include those from the yeasts *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, and *Kluyveromyces lactis* (Carbon *et al.*, 1990; Hegemann *et al.*, 1993). In many of these systems, accurate mapping of the
25 centromeres made it possible to clone centromeric DNA, using a chromosome walking strategy (Clarke *et al.*, 1980). Subsequently, artificial chromosome assays were used to define more precisely the centromere sequences (Hegemann *et al.*, 1993; Baum *et al.*, 1994).

30 Attempts to develop a reliable centromeric assay in mammals have yielded ambiguous results. For example, Hadlaczký *et al.*, (1991) identified a 14 kB human fragment that can, at low frequency, result in *de novo* centromere formation in a mouse

cell line. *In situ* hybridization studies, however, have shown that this fragment is absent from naturally occurring centromeres, calling into question the reliability of this approach for testing centromere function (Tyler-Smith *et al.*, 1993). Similarly, transfection of alphoid satellites into cell lines results in the formation of new
5 chromosomes, yet some of these chromosomes also required host sequences that could contribute centromere activity (Haaf *et al.*, 1992; Willard, 1997). Further, the novel chromosomes can have alphoid DNA spread throughout their length yet have only a single centromeric constriction, indicating that a block of alphoid DNA alone may be insufficient for centromere function (Tyler-Smith *et al.*, 1993).

10

Although plant centromeres can be visualized easily in condensed chromosomes, they have not been characterized as extensively as centromeres from yeast or mammals. Genetic characterization has relied on segregation analysis of chromosome fragments, and in particular on analysis of trisomic strains that carry a
15 genetically marked, telocentric fragment (for example, see Koornneef 1983). In addition, repetitive elements have been identified that are either genetically (Richards *et al.*, 1991) or physically (Alfenito *et al.*, 1993; Maluszynska *et al.*, 1991) linked to a centromere. In no case, however, has the functional significance of these sequences been tested.

20

Cytology in *Arabidopsis thaliana* has served to correlate centromere structure with repeat sequences. A fluorescent dye, DAPI, allows visualization of centromeric chromatin domains in metaphase chromosomes. A fluorescence *in situ* hybridization (FISH) probe based on 180 bp pAL1 repeat sequences colocalized with the DAPI
25 signature near the centromeres of all five *Arabidopsis* chromosomes (Maluszynska *et al.*, 1991; Martinez-Zapater *et al.*, 1986). Although a functional role for pAL1 has been proposed, more recent studies have failed to detect this sequence near the centromeres in species closely related to *Arabidopsis thaliana* (Maluszynska *et al.*, 1993). These results are particularly troubling because one of the
30 species tested, *A. pumila*, is thought to be an amphidiploid, derived from a cross between *A. thaliana* and another close relative (Maluszynska *et al.*, 1991; Price *et al.*, 1995). Another repetitive sequence, pAtT12, has been genetically mapped

to within 5 cM of the centromere on chromosome 1 and to the central region of chromosome 5 (Richards *et al.*, 1991), although its presence on other chromosomes has not been established. Like pAL1, a role for pAtT12 in centromere function remains to be demonstrated.

5

Due to the fact that kinetochores constitute a necessary link between centromeric DNA and the spindle apparatus, the proteins that are associated with these structures recently have been the focus of intense investigation (Bloom 1993; Earnshaw 1991). Human autoantibodies that bind specifically in the vicinity of the centromere have facilitated the cloning of centromere-associated proteins (CENPs, Rattner 1991), and at least one of these proteins belongs to the kinesin superfamily of microtubule-based motors (Yen 1991). Yeast centromere-binding proteins also have been identified, both through genetic and biochemical studies (Bloom 1993; Lechner *et al.*, 1991).

15

The centromeres of *Arabidopsis thaliana* have been mapped using trisomic strains, where the segregation of chromosome fragments (Koornneef 1983) or whole chromosomes (Sears *et al.*, 1970) was used to localize four of the centromeres to within 5, 12, 17 and 38 cM, respectively. These positions have not been refined by more recent studies because the method is limited the difficulty of obtaining viable trisomic strains (Koornneef 1983). These factors introduce significant error into the calculated position of the centromere, and in *Arabidopsis*, where 1 cM corresponds roughly to 200 kB (Koornneef 1987; Hwang *et al.*, 1991), this method did not map any of the centromeres with sufficient precision to make chromosome walking strategies practical. Mapping of the *Arabidopsis* genome was also discussed by (Hauge *et al.*, 1991).

25

I. Isolation of centromere clones

The present invention relates to methods of isolating and identifying centromere DNA sequences from total genomic DNA of an organism without genetic mapping of the organism. Centromere DNA can be purified from total genomic DNA using several methods which include: 1) digesting genomic DNA with restriction

30

enzymes and separating the fragments on agarose gels, to reveal major classes of repetitive DNA; 2) digesting genomic DNA with restriction enzymes sensitive to DNA methylation and separating the fragments on agarose gels to reveal the heavily methylated fraction of the genome; and 3) collecting the rapidly annealing fraction of denatured genomic DNA. These three methods isolate centromere DNA; therefore, these methods are expected to independently isolate the same sequences, thus validating the sequences' centromere origin. It is anticipated that each of these methods can be applied to genomic DNA from any organism, including some lower organisms such as yeasts, as well as higher organisms such as plants and animals. Each of these methods is described in detail below.

1. Isolation of repetitive DNA

Centromere regions often contain many copies of the same DNA sequence (repetitive DNA); such repeats can range in size from a few nucleotides long to hundreds or thousands of bases. Such repetitive DNA can be identified following digestion of genomic DNA with restriction endonucleases. Digestion of non-repetitive genomic DNA with a particular restriction enzyme produces a distribution of size fragments; in contrast, digestion of repeats with a restriction enzyme that cuts within each repeat produces a fragment of a typical size. Thus, genomic DNA that has been cut with a restriction enzyme can be size fractionated by agarose gel electrophoresis to reveal repetitive DNA elements; after staining the gel to reveal the DNA, the repetitive fragment can be excised and purified using conventional techniques or commercial kits. Such repeats can be introduced into cloning vectors and characterized as described below. By using this method with a variety of restriction enzymes, different repetitive elements can be purified from genomic DNA.

2. Purification of methylated DNA

This method is disclosed in detail in co-pending U.S. Patent Application Serial No.09/888,220, filed June 22, 2001, the disclosure of which is incorporated herein by reference in its entirety and made a part hereof. Plant centromere DNA is often extensively modified by methylation; the presence of this methylation can be used to purify centromere fragments. Digestion of genomic DNA with a methylation-sensitive

restriction endonuclease (for example Sau3A or HpaII) yields a range of fragment sizes; endonuclease sites that are methylated are protected from digestion. Heavily methylated DNA molecules, such as centromere DNA, yield large fragments after digestion and can therefore be separated from the lightly or non-methylated fraction by virtue of their size. For example agarose gel electrophoresis, acrylamide gel electrophoresis, sucrose gradient fractionation, or other size fractionation techniques can be used to separate these fragments into pools of "large" (7-12kb) and "smaller" fragments (3-7 kb and 0-3kb).

10 3. Isolation of rapidly annealing DNA.

The rapidity with which denatured single stranded DNA can reanneal with another single stranded DNA molecule of complementary sequence upon renaturation is dependant upon its abundance. Therefore when genomic DNA is denatured and allowed to renature, the repetitive fraction of the genome, including centromere DNA, will renature before the unique and low copy fractions of the genome. Thus by fragmenting purified genomic DNA, denaturing it, collecting fractions at specific time points (such as 2, 4, 6, 8, and 10 minutes) during renaturation and treating those fractions to remove unannealed DNA it is possible to purify repetitive DNA from total genomic DNA. Several methods can be used to remove unannealed from annealed DNA including treatment of the sample with an enzyme, such as S1 nuclease, that degrades single-stranded DNA or exposure to an agent that binds single-stranded DNA such as hydroxylapatite. By varying the time at which fractions are collected during renaturation it is possible to separate DNA fragments into highly repetitive, moderately repetitive, and non-repetitive fractions.

25 II. Cloning and sequencing small fragments of centromere DNA

Repetitive or methylated DNA fragments isolated using the methods described above can be ligated (using T4 DNA ligase, for example) to a plasmid vector and cloned by transformation into *E. coli*. These clones can then be propagated, sequenced, used to assemble minichromosomes, or used to identify larger centromere clones, generate molecular markers that facilitate genetic mapping of centromeres, or create

probes for chromosome mapping experiments such as fluorescent in situ hybridization (FISH).

III. Identifying centromere clones in genomic libraries

5 A genomic library can be screened for clones carrying centromere DNA by arraying the clones onto solid supports, such as membrane filters, and probing with labeled fragments of purified centromere DNA, including cloned repetitive or methylated DNA fragments described above, or alternatively, the entire set of rapidly annealing genomic DNA or highly methylated genomic DNA fragments. Probes can
10 be used singly or in combination. Typically these probes are labeled by incorporation of radionucleotides, fluorescent nucleotides, or other chemical or enzymatic ligands that enable easy detection. The labeled probe DNA is denatured and hybridized to the arrayed library using standard molecular biology techniques. Hybridization is performed at a temperature that will discourage non-specific DNA annealing while
15 promoting the hybridization of the labeled probe to complementary sequences. After incubation, the arrayed library is washed to remove unannealed probe, and a detection method appropriate to the label incorporated in the probe is used. For example, if the probe is radiolabeled, the labeled filter is exposed to X-ray film.

20 To identify centromere clones, the results of several hybridization experiments are quantitated and compared. In some cases, centromere clones may hybridize to only one probe; in other cases, the clones will hybridize to multiple probes. The hybridization intensity of each clone to each probe can be measured and stored in a database. A preferred method for this analysis is to use software that digitizes the
25 hybridization signals, assigns each signal to its corresponding clone address, ensures that duplicate copies of the clones successfully hybridized, and enters the resulting information into a relational database (MySQL for example). Another possible method for this analysis is to examine the hybridization results visually, estimate the hybridization intensity, and tabulate the resulting information.

30

The results of each hybridization experiment can be classified by grouping

clones that show hybridization to each probe above a threshold value. For example, a computerized relational database can be queried for clones giving hybridization signals above a certain threshold for individual probes or for multiple probes. Based on these hybridization patterns, clones can be grouped into categories, and representative
5 members of each category can be tested in minichromosomes.

IV. Identifying Centromere Sequences of an Organism from Genomic Sequence

Datasets

It is possible to devise computational algorithms to search databases of
10 genomic sequences and select centromere sequences by identifying those with the characteristics of centromeres. For example, by selecting the most abundant tandem repeat of a particular size will yield centromere sequences. Other sets of characteristics could also be useful. The following is an example of a computational algorithm designed to extract centromere sequences from genomic sequence datasets.
15 It is important to note that this algorithm examines primary sequence data and does not rely on prior annotation of the sequence. The algorithm consists of steps 1 through 10. However, not all the steps must occur in the listed order without altering the output. Other rearrangements are easily recognizable by one skilled in the art. The following terms are used in describing the algorithm. BLAST is Basic Local Alignment Search
20 Tool, a family of freely available algorithms for sequence database searches. BLAST aligns two sequences and yields an estimate of the probability that this alignment is significant, i.e. that it did not occur by chance. The two sequences compared by BLAST are called the 'query', usually a single sequence of interest, and the 'subject', often part of a large database of sequences that are compared to the query. The query
25 sequence (query) can also be part of a database of sequences. The outputs of BLAST are High Scoring Pairs (HSPs) that are alignments of subject and query sequences. Nucleotide position describes the position of a given nucleotide within the sequence, relative to the first nucleotide of the sequence. BLAST score (e value) is the likelihood that a given sequence alignment is significant (the lower the value the
30 higher the significance). The algorithm is as follows:

- (1) provide a first dataset consisting of the genomic sequences, or a representative fraction of genomic sequence, of the organism of interest;
- (2) identify and eliminate known non-centromeric repeat sequences from the first dataset by using the BLAST sequence comparison algorithm to create a second
5 dataset;
- (3) compare each sequence in the second dataset to itself by using the BLAST sequence comparison algorithm, obtain a BLAST score for each pair of sequence compared, and collect high score pairs to create a third dataset;
- (4) examine the BLAST score of each high score pair in the third dataset and
10 eliminate the pairs having a score greater than 10^{-20} to create a fourth dataset;
- (5) eliminate the high score pairs in the fourth dataset having less than 80 bp or more than 250 bp to create a fifth dataset;
- (6) examine the nucleotide position of each high score pair in the fifth dataset and eliminate pairs having 100% identity and identical nucleotide positions (i.e. self
15 matches) to create a sixth dataset;
- (7) examine the nucleotide position of each high score pair in the sixth dataset and eliminate pairs having opposite orientation of the nucleotides to create a seventh dataset;
- (8) examine the nucleotide position of both sequences for each high score pair
20 in the seventh dataset and eliminate sequences that are overlapping to create an eighth dataset; and
- (9) examine the nucleotide position of each sequence in the eighth dataset and eliminate sequences not having at least one neighboring sequence within 250 bp to create a ninth dataset; and
- 25 (10) compare each sequence in the ninth dataset to all other sequences in the ninth dataset by using the BLAST sequence comparison algorithm and select the most common sequence as a centromere sequence of the organism.

Optimally, the dataset used in step (1) in the above algorithm would be the
30 whole genome dataset such as the *Arabidopsis* genome which was derived by methodical sequencing of mapped clones or the rice genome dataset which was derived by shotgun sequencing. Alternatively, the algorithm would also work well on

representative genome datasets. By the term "representative genome datasets", it is meant that the genomic sequences in the dataset is a subset of the sequences of the whole genome collected from the whole genome without bias, such as bias toward coding sequences. These sequences would be representative of the genome as a whole. For example, the use of a 0.5X or even a 0.1X library of *Arabidopsis* with representative genome datasets would return a true positive result. On the contrary, the use of a subset of genomic sequences of the whole genome which are not representative of the whole genome and biased toward certain sequences, such as the coding sequence, would return false positive results.

10

V. Centromere Compositions

The present invention concerns nucleic acid segments, isolatable from various plant cells, that are enriched relative to total genomic DNA, or isolated from other sources or chemically synthesized with a novel sequence, or other nucleic acids that are capable of conferring centromere activity to a recombinant molecule when incorporated into the host cell. As used herein, the term "nucleic acid segment" refers to a nucleic acid molecule that has been purified from total genomic nucleic acids of a particular species. Therefore, a nucleic acid segment conferring centromere function refers to a nucleic acid segment that contains centromere sequences yet is isolated away from, or purified free from, total genomic nucleic acids. Included within the term "nucleic acid segment", are nucleic acid segments and smaller fragments of such segments, and also recombinant vectors, including, for example, minichromosomes, artificial chromosomes, BACs, YACs, plasmids, cosmids, phage, viruses, and the like.

25

Similarly, a nucleic acid segment comprising an isolated or purified centromeric sequence refers to a nucleic acid segment including centromere sequences and, in certain aspects, regulatory sequences, isolated substantially away from other naturally occurring sequences, or other nucleic acid sequences. In this respect, the term "gene" is used for simplicity to refer to a protein, polypeptide- or peptide-encoding unit. As will be understood by those in the art, this functional term includes

30

both genomic sequences, cDNA sequences and smaller engineered gene segments that may express, or may be adapted to express, proteins, polypeptides or peptides.

"Isolated substantially away from other sequences" means that the sequences
5 of interest, in this case centromere sequences, are included within the genomic nucleic acid clones provided herein. Of course, this refers to the nucleic acid segment as originally isolated, and does not exclude all genes or coding regions.

In particular embodiments, the invention concerns isolated nucleic acid
10 segments and recombinant vectors incorporating nucleic acid sequences that encode a centromere functional sequence that includes a contiguous sequence from the centromeres of the current invention. Again, nucleic acid segments that exhibit centromere function activity will be most preferred.

15 In still yet another aspect, the invention provides a plant centromere which is further defined as an *Arabidopsis thaliana* centromere. In yet another embodiment of the invention, the plant centromere comprises an *Arabidopsis thaliana* chromosome 2 centromere. The chromosome 2 centromere may comprise, for example, from about 100 to about 611,000, about 500 to about 611,000, about 1,000 to about 611,000, about
20 10,000 to about 611,000, about 20,000 to about 611,000, about 40,000 to about 611,000, about 80,000 to about 611,000, about 150,000 to about 611,000, or about 300,000 to about 611,000 contiguous nucleotides of a first nucleic acid sequence flanking a first series of 180 bp repeats in centromere 2 of *A. thaliana*. The centromere may also be defined as comprising from about 100 to about 50,959, about
25 500 to about 50,959, about 1,000 to about 50,959, about 5,000 to about 50,959, about 10,000 to about 50,959, 20,000 to about 50,959, about 30,000 to about 50,959, or about 40,000 to about 50,959 contiguous nucleotides of a second nucleic acid sequence flanking a second series of 180 bp repeats in centromere 2 of *A. thaliana*. The centromere may comprise sequences from both of the third and the fourth
30 sequences, including the aforementioned fragments, or the entirety of these sequences. In particular embodiments, the inventors contemplate a 3' fragment of the first

sequence can be fused to a 5' fragment of the second sequence, optionally including one or more 180 bp repeat sequence disposed therebetween.

In still yet another aspect, the invention provides an *Arabidopsis thaliana* chromosome 4 centromere. In certain embodiments of the invention, the centromere may comprise from about 100 to about 1,082,000, about 500 to about 1,082,000, about 1,000 to about 1,082,000, about 5,000 to about 1,082,000, about 10,000 to about 1,082,000, about 50,000 to about 1,082,000, about 100,000 to about 1,082,000, about 200,000 to about 1,082,000, about 400,000 to about 1,082,000, or about 800,000 to about 1,082,000 contiguous nucleotides of a third nucleic acid sequence flanking a third series of repeated sequences, including comprising the nucleic acid sequence of the third sequence. The centromere may also be defined as comprising from about 100 to about 163,317, about 500 to about 163,317, about 1,000 to about 163,317, about 5,000 to about 163,317, about 10,000 to about 163,317, about 30,000 to about 163,317, about 50,000 to about 163,317, about 80,000 to about 163,317, or about 120,000 to about 163,317 contiguous nucleotides of the nucleic acid sequence of a fourth sequence flanking a fourth series of repeated sequences, and may be defined as comprising the nucleic acid sequence of the fourth sequence. The centromere may comprise sequences from both the third and the fourth sequences, including the aforementioned fragments, or the entirety of the third and the fourth sequences. In particular embodiments, the inventors contemplate a 3' fragment of the third sequence can be fused to a 5' fragment of the fourth sequence, optionally including one or more 180 bp repeat sequence disposed therebetween.

In yet another embodiment, there is provided a *Arabidopsis thaliana* chromosome 1, 3 or 5 centromere selected from the nucleic acid sequence given by one of the repeated sequences in these chromosomes, or fragments thereof. The length of the repeat used may vary, but will preferably range from about 20 bp to about 250 bp, from about 50 bp to about 225 bp, from about 75 bp to about 210 bp, from about 100 bp to about 205 bp, from about 125 bp to about 200 bp, from about 150 bp to about 195 bp, from about 160 bp to about 190 and from about 170 bp to about 185 bp including about 180 bp. In one embodiment, the construct comprises at least 100 base

pairs, up to an including the full length, of one of the preceding sequences. In addition, the construct may include 1 or more 180 base pair repeats.

In one embodiment, the centromere n copies of a repeated nucleotide sequence
5 obtained by the method disclosed herein, wherein n is at least 2. Potentially any number of repeat copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100, 150, 200, 300, 400, 500, 750, 1,000, 1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 and about 100,000,
10 including all ranges in-between such copy numbers. Moreover, the copies, while largely identical, can vary from each other. Such repeat variation is commonly observed in naturally occurring centromeres.

In another embodiment, the centromere is a *Brassica oleracea* centromere comprising *Brassica oleracea* centromere DNA. In one embodiment, the *Brassica*
15 *oleracea* centromere is defined as comprising n copies of a repeated nucleotide sequence, wherein n is at least 2. Potentially any number of repeat copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100, 150, 200, 300, 400, 500, 750, 1,000, 1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000,
20 60,000, 70,000, 80,000, 90,000 and about 100,000, including all ranges in-between such copy numbers. In one embodiment, the repeated nucleotide sequence is isolated from *Brassica oleracea* given by SEQ ID NO:1, 2, 3, or 4.

In yet another embodiment, the centromere is a *Glycine max* centromere
25 comprising *glycine max* centromere DNA. In an embodiment, the *Glycine max* centromere is defined as comprising n copies of a repeated nucleotide sequence, wherein n is at least 2. Potentially any number of repeat copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100, 150, 200, 300, 400, 500, 750, 1,000,
30 1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 and about 100,000, including all ranges in-between such copy

numbers. In one embodiment, the repeated nucleotide sequence is isolated from *Glycine max* given by SEQ ID NO:5, 6, 7, or 8.

In yet another embodiment, the centromere is a *Lycopersicon esculentum* centromere comprising *Lycopersicon esculentum* centromere DNA. In an embodiment, the *Lycopersicon esculentum* centromere is defined as comprising n copies of a repeated nucleotide sequence, wherein n is at least 2. Potentially any number of repeat copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100, 150, 200, 300, 400, 500, 750, 1,000, 1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 and about 100,000, including all ranges in-between such copy numbers. In one embodiment, the repeated nucleotide sequence is isolated from *Lycopersicon esculentum* given by SEQ ID NO:9 or 10.

In yet another embodiment, the centromere is a *Zea mays* centromere comprising *Zea mays* centromere DNA. In an embodiment, the centromere is defined as comprising n copies of a repeated nucleotide sequence, wherein n is at least 2. Potentially any number of repeat copies capable of physically being placed on the recombinant construct could be included on the construct, including about 5, 10, 15, 20, 30, 50, 75, 100, 150, 200, 300, 400, 500, 750, 1,000, 1,500, 2,000, 3,000, 5,000, 7,500, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 and about 100,000, including all ranges in-between such copy numbers. In one embodiment, the repeated nucleotide sequence is isolated from *Zea mays* given by SEQ ID NO:11, 12 or 13.

25

The centromere can additionally be defined as the region of the chromosome where the sister chromatids pair during cell division. The centromere is also the chromosomal region where the kinetochore (the chromosomal attachment structure for the spindle) and the spindle (the cellular machinery that provides the motive force for chromosome segregation) attach to the chromosome during mitosis and meiosis. The centromere is also defined as the region of the primary constriction in a condensed

chromosome. The DNA of the centromere is characteristically heavily methylated, repetitive, and condensed (heterochromatic).

VI. Minichromosome construction

5 Minichromosomes are constructed by combining fragments of centromere DNA with other DNA sequences useful for propagation of the resultant recombinant DNA molecule in *E. coli*, other bacteria, yeast or plants. Recombinant plasmids containing large fragments of centromere DNA are referred to as centromere clones. Centromere sequences removed from centromere clones, or centromere sequences
10 derived directly from genomic DNA, are referred to as centromere fragments. Recombinant constructs containing DNA sequences necessary for the propagation, delivery, selection, and detection of minichromosomes will be referred to as minichromosome vector sequences or minichromosome vectors; these sequences can include but are not limited to selectable marker genes, visible marker genes, origins of
15 replication, restriction endonuclease recognition sites, homing endonuclease recognition sites, sequences recognized by site specific recombinase enzymes, telomere sequences, and sequences required for delivery of minichromosomes into bacteria, yeast or plant cells. Recombinant constructs containing both large centromere fragments as well as minichromosome vector sequences are referred to as
20 minichromosomes. The process of assembling minichromosomes from centromere clones/fragments and minichromosome vector sequences can be done in several ways, and involves techniques that are common practice among those trained in molecular biology:

1) Joining centromere fragments to minichromosome vector sequences:

25 Centromere DNA fragments and minichromosome vector DNA fragments are generated and purified using conventional techniques, some of which include restriction enzyme digestion, agarose gel electrophoresis, gel purification of specific fragments, anion-exchange purification and ethanol precipitation. The resulting purified centromere and vector fragments are enzymatically joined *in vitro*, using for
30 example T4 DNA ligase. The ends of the fragments can be cohesive, as the result of digestion with compatible restriction endonucleases or from the addition of compatible oligonucleotide linkers; alternatively the ends of the fragments can be blunt and can be

directly joined. Following ligation, the resulting minichromosomes are introduced into *E. coli*, other bacteria, yeast, or plant cells using chemical or physical transformation methods. The structure of the resulting minichromosomes can be determined by recovering them from the host organism and assessing DNA fragment size and composition.

2) Transfer of minichromosome vector sequences into centromere clones by site-specific recombination:

The minichromosome vector sequences can be constructed to include site-specific recombination sequences (for example those recognized by the bacteriophage P1 Cre recombinase, or the bacteriophage lambda integrase, or similar recombination enzymes). A compatible recombination site, or a pair of such sites, can also be included in the centromere clones. Incubation of the minichromosome vector and the centromere clone in the presence of the recombinase enzyme causes strand exchange to occur between the recombination sites in the two plasmids; the resulting minichromosomes contain centromere sequences as well as minichromosome vector sequences (FIG. 5). Introducing the DNA molecules formed in such recombination reactions into *E. coli*, other bacteria, yeast or plant cells can be followed by selection for marker genes present on both parental plasmids, allowing the isolation of minichromosomes.

3) *Minichromosome vector tailing method for minichromosome construction:*

Centromere DNA fragments isolated from genomic DNA or from centromere clones can be modified on their ends by treatment with restriction endonucleases, or by ligation with DNA molecules including, but not limited to, oligonucleotide linkers, or by the addition of nucleotides, to produce a desired cohesive or blunt end. These fragments are size-fractionated by, agarose gel electrophoresis or other methods, and the centromere fragments purified using conventional techniques. Minichromosome vector fragments are generated and purified in a similar manner, resulting in linear minichromosome vector sequences with DNA ends compatible with those on the centromere fragments. Compatible ends in this case are defined by ends that can be joined in vitro by the action of a ligase enzyme. As shown in FIG. 6, the two fragments are then mixed so that the minichromosome vector molecules are present in at least two-fold molar excess over the centromere fragments. The fragments are joined by the

addition of a ligase enzyme (for example bacteriophage T4 DNA ligase), resulting in the formation of DNA molecules in which minichromosome vector molecules have been joined to both ends of the same centromere fragment. Digestion of the ligation mixture with a rare-cutting restriction or homing endonuclease (for example
5 endonucleases with recognition sequences of 8 or more bases) results in linear minichromosome precursors consisting of a fragment of the original minichromosome vector attached to each end of the centromere fragment. The ends of this hybrid molecule are compatible because they were created by the same restriction enzyme. This linear minichromosome precursor is purified, for example, by agarose gel
10 electrophoresis followed by gel purification of the DNA fragments of the expected length. The purified DNA molecules are circularized by joining the ends, for example by treatment with a DNA ligase enzyme. The resulting minichromosome molecules can be introduced into *E. coli*, other bacteria, yeast or plant cells, followed by purification and characterization using conventional methods.

15

VII. Use of minichromosomes for plant transformation

1) Delivery of minichromosomes into plant cells:

Minichromosomes are purified and delivered into plant cells, either
20 individually or as a mixture. The minichromosomes can be either circular or linear or mixtures thereof. The plant cells used for minichromosome delivery can be either intact seedlings, immature or mature plants, parts of seedlings or plants, specific plant tissues (for example leaves, stems, roots, flowers, fruits), differentiated tissues cultured in vitro (for example roots), or undifferentiated cells (for example callus) cultured in
25 vitro. The minichromosome DNA can be delivered into plant cells by a variety of methods including but not limited to the following: electroporation; *Agrobacterium*-mediated DNA delivery; virus-mediated DNA delivery; delivery mediated by salts or lipids that facilitate the cellular uptake of DNA; microinjection of DNA; manipulation into a cell of DNA-coated or DNA-containing particles, droplets, micelles,
30 microspheres, or chemical complexes using a variety of techniques, including biolistic particle bombardment, optical tweezers, particle beams, and electrospray apparatus; manipulation of DNA-coated magnetic particles into the cells by magnetic fields;

DNA delivery into cells by cell wounding using micro-needles (for example silicon carbide needles); sonication or other acoustic treatment of the cells to facilitate DNA uptake; fusion of plant cells with other cell types carrying a minichromosome, including bacterial, yeast, or other plant cells; any other electrical, chemical, physical, or biological mechanism that results in the introduction of minichromosome DNA into the plant cell

2) Isolating plant cells containing minichromosomes:

Following minichromosome delivery, plant cells, plant tissues, or complete plants carrying the minichromosome can be isolated by a variety of selection methods. Selection involves subjecting the plant cells, tissues or plants to chemical, environmental, or mechanical treatments that enrich for those cells, tissue or plants that contain a minichromosome. The selection methods include but are not limited to: fluorescence-activated cell sorting of cells, cell clumps, or cell protoplasts based on expression of a marker protein encoded by the minichromosome (for example, a fluorescent protein such as DsRed); affinity purification of cells, cell clumps, or protoplasts based on expression of a cell wall protein, membrane protein, or membrane-associated protein encoded by the minichromosome; any cell fractionation method capable of separating cells based on their density, size or shape to enrich for cells with a property that differs from that of the starting population and is conferred by the minichromosome; selection of cells for resistance to an antibiotic conferred by the minichromosome; selection of cells for resistance to an herbicide conferred by the minichromosome; selection of cells for resistance to a toxic metal, salt, mineral or other substance conferred by the minichromosome; selection of cells for resistance to abiotic stress (for example heat, cold, acid, base, osmotic stress) conferred by the minichromosome; selection of cells capable of utilizing a carbon source or other nutrient source not normally utilized by plant cells, this utilization function being conferred by the minichromosome. As a result of the treatment, a population of plant cells can be obtained that contain minichromosomes. Individual clones or sub-populations of these cells can be expanded in culture for further characterization.

Alternatively, plant cells, plant tissues, or complete plants that carry minichromosomes can be identified by direct screening. Such methods involve subjecting each cell, plant, or tissue to diagnostic tests indicative of the presence of the minichromosome. These tests can include direct assays for the presence of minichromosome DNA, or indirect assays for properties conferred by the minichromosome. Direct assays for the presence of the minichromosome DNA include but are not limited to: staining of cells with DNA-binding molecules to allow detection of an additional chromosome; *in situ* hybridization with labeled DNA probes corresponding to sequences present on the minichromosome; southern blots or dot blots of DNA extracted from the cells, plant or tissue and probed with labeled DNA sequences corresponding to sequences present on the minichromosome; electrophoresis of genomic DNA extracted from the cells, plant or tissue under conditions that allow identification of the minichromosome; amplification of specific sequences present on the minichromosome from genomic DNA extracted from the cells, plant or tissue using the polymerase chain reaction. Indirect assays for properties conferred by the minichromosome include but are not limited to: detection of the expression of a fluorescent marker encoded by the minichromosome by fluorescence microscopy, flow cytometry or fluorimetry; detection of the expression of a protein encoded by the minichromosome by use of specific antibodies, or any other reagent capable of specifically binding to the protein; use of cell fractionation methods capable of detecting a specific density, size or shape of the cells or tissues, that is conferred by the minichromosome; growth of cells, seedlings, plants or tissues on an antibiotic-containing medium to determine the presence of an antibiotic-resistance gene encoded by the minichromosome; growth of cells, seedlings, plants or tissues on an herbicide-containing medium to determine the presence of an herbicide-resistance gene encoded by the minichromosome; growth of cells, seedlings, plants or tissues on a medium containing a toxic metal, salt, mineral or other substance to determine the presence of an gene conferring resistance to this substance encoded by the minichromosome; growth of cells, tissues or plants under conditions of abiotic stress (for example heat, cold, acid, base, osmotic stress) to determine the presence of a gene conferring resistance to this stress encoded by the minichromosome; growth of cells on a medium

containing a carbon source or other nutrient source normally not utilized by plant cells, to determine the presence of a utilization function conferred by the minichromosome.

3) Characterization of plant cell clones containing minichromosomes

5 Plant cells, tissues, or entire plants containing minichromosomes can be further characterized to determine whether the minichromosome is an autonomous DNA molecule, or whether it is associated with one of the plant cell's chromosomes by integration. The methods used for this analysis include, but are not limited to, the following:

10 1) Detection of marker protein expression by microscopy, flow cytometry, fluorimetry, enzymatic assays, cell staining or any other technique that allows the detection of a marker protein having a specific enzymatic activity, or conferring a specific color, or fluorescence property onto the cells. For example, if a cell line has been selected for containing a minichromosome by selecting for the function of a resistance gene
15 encoded by the minichromosome, and if a marker protein is also encoded by the minichromosome, then expression of this marker protein in the selected cells is an indication of the presence of the entire minichromosome, and could indicate autonomy of this minichromosome from the cell's other chromosomes.

2) Use of gel electrophoresis to detect a minichromosome in genomic DNA isolated
20 from the plant cells, tissue or entire plants. For example, genomic DNA isolated from the cells, tissues or plants can be fractionated by gel electrophoresis, either intact or following digestion with restriction endonucleases or homing endonucleases, allowing the detection of a minichromosome or a fragment of a minichromosome.

3) Use of southern blots or dot blots of DNA extracted from the cells, tissue or plants
25 to detect the presence of specific sequences contained on the minichromosome. For example, digestion of genomic DNA extracted from the cells, tissues or plants can be fractionated by agarose gel electrophoresis, blotted onto a DNA-binding membrane, and probed with labeled DNA sequences corresponding to sequences present on the minichromosome to detect specific fragments of minichromosome DNA, and thus
30 allowing the determination of the autonomous, or integrated structure of the minichromosome.

4) Cytological techniques for directly visualizing the minichromosome in the transformed cells, such as staining of cells with DNA-binding dyes or in situ hybridization with labeled DNA probes corresponding to sequences present on the minichromosome.

- 5) Genetic analysis of marker segregation by scoring marker inheritance in progeny of a plant containing a minichromosome. For example, markers present on an autonomous minichromosome will segregate independently from markers on the arms of the host chromosomes in a population of F2 progeny generated from a cross between a line carrying a minichromosome and a second marked line that doesn't carry the minichromosome. Markers include but are not limited to: visible markers conferring a visible characteristic to the plant; selectable markers, conferring resistance to an antibiotic, herbicide, or other toxic compound; enzymatic markers, conferring an enzymatic activity that can be assayed in the plant or in extracts made from the plant; protein markers, allowing the specific detection of a protein expressed in the plant; molecular markers, such as restriction fragment length polymorphisms, amplified fragment length polymorphisms, short sequence repeat (microsatellite) markers, presence of certain sequences in the DNA of the plant as detected by the polymerase chain reaction, single nucleotide polymorphisms or cleavable amplified polymorphic sites.

4) *Plant regeneration from transformed cell clones:*

Plant cells or tissues that harbor minichromosomes can be used to regenerate entire plants. This will be accomplished with standard techniques of plant regeneration from differentiated tissues or undifferentiated cells. Typically, transformed tissues or callus are subjected to a series of treatments with media containing various mixtures of plant hormones and growth regulators that promote the formation of a plant embryo, specific plant tissues or organs, or a complete plant (roots and shoot) from the starting cells or tissues. Following plant regeneration, the plant can be grown either in sterile media or in soil.

VIII. Testing minichromosome inheritance in plant cells

The inheritance of minichromosomes can be measured through one or more cell divisions. After isolating cells, tissues, or entire plants that contain the minichromosome, the population of cells is allowed to grow (either with or without selection), and the presence of the minichromosome is monitored as the cells divide.

5 Minichromosomes can be detected in cells by a variety of methods, including but not limited to: detection of fluorescence or any other visual characteristic arising from a marker protein gene present on the minichromosome; resistance to an antibiotic, herbicide, toxic metal, salt, mineral or other substance, or abiotic stress as outlined above (*Isolating plant cells containing minichromosomes*); staining of cells with

10 DNA-binding molecules to allow detection of an additional chromosome; in situ hybridization with labeled DNA probes corresponding to sequences present on the minichromosome; southern blots or dot blots of DNA extracted from the cell population and probed with labeled DNA sequences corresponding to sequences present on the minichromosome; expression of a marker enzyme encoded by a gene

15 present on the minichromosome (i.e. luciferase, alkaline phosphatase, beta-galactosidase, etc.) that can be assayed in the cells or in an extract made from the cells.

The percentage of cells containing the chromosome is determined at regular intervals during this growth phase. The change in the fraction of cells harboring the minichromosome, divided by the number of cell divisions, represents the average

20 minichromosome loss rate. Minichromosomes with the lowest loss rates have the highest level of inheritance.

IX. Recovery of minichromosomes from plant cells

Recovery of minichromosomes from plant cells can be achieved by a variety of

25 techniques, including, but not limited to, the following:

- 1) Extracting the genomic DNA of transformed plant cells and introducing that DNA into *E. coli*, other bacteria or yeast and selecting for the antibiotic resistance genes present on the minichromosome.
 - 2) Isolation of chromosomes from cells, tissues or plants containing
- 30 minichromosomes, and sorting these by flow cytometry to allow the separation of chromosomes of different size;

- 3) Isolation of individual chromosomes from a cell harboring minichromosomes by micro-manipulation involving mechanical devices such as needles made of glass, metal or other suitable substances, or other techniques such as optical tweezers, or micro-suction devices.
- 5 4) Combinations of the above, for example chromosome isolation by flow cytometry or micromanipulation followed by introduction into *E. coli*, other bacteria, yeast or plant cells.

The resulting minichromosomes "rescued" in this fashion may differ from their
10 parental molecules in total size, size of the centromere, presence or absence of additional sequences, and overall arrangement of the sequences. These procedures allow the isolation of DNA molecules capable of replicating and segregating in plant cells without having to test minichromosomes individually. For example, after delivery of pools of minichromosomes, or pools of centromere clones into plant cells, tissues or
15 whole plants, and recovering them by the methods listed above, facilitates the selection of specific minichromosomes or centromere clones that remain autonomous in plant cells. Whereas plant transformation with minichromosomes relies on the sequences contributed by minichromosome vectors, the recovery methods do not necessarily require minichromosome vector sequences; as a result, pools of centromere clones can
20 be delivered into plant cells followed by recovery of the ones that replicated and persist.

X. Exogenous Genes for Expression in Plants

One particularly important advance of the present invention is that it provides
25 methods and compositions for expression of exogenous genes in plant cells. One advance of the constructs of the current invention is that they enable the introduction of multiple genes (often referred to as gene "stacking"), potentially representing an entire biochemical pathway, or any combination of genes encoding different biochemical processes or pathways. Significantly, the current invention allows for the
30 transformation of plant cells with a minichromosome comprising a number of structural genes. Another advantage is that more than one minichromosome could be introduced, allowing combinations of genes to be moved and shuffled. Moreover, the

ability to eliminate a minichromosome from a plant would provide additional flexibility, making it possible to alter the set of genes contained within a plant. Further, by using site-specific recombinases, it should be possible to add genes to an existing minichromosome once it is in a plant.

5

Added genes often will be genes that direct the expression of a particular protein or polypeptide product, but they also may be non-expressible DNA segments, *e.g.*, transposons such as Ds that do not direct their own transposition. As used herein, an "expressible gene" is any gene that is capable of being transcribed into RNA (*e.g.*,
10 mRNA, antisense RNA, *etc.*) or translated into a protein, expressed as a trait of interest, or the like, *etc.*, and is not limited to selectable, screenable or non-selectable marker genes. The inventors also contemplate that, where both an expressible gene that is not necessarily a marker gene is employed in combination with a marker gene, one may employ the separate genes on either the same or different DNA segments for
15 transformation. In the latter case, the different vectors may be delivered concurrently to recipient cells to maximize cotransformation or may be delivered sequentially.

The choice of the particular DNA segments to be delivered to the recipient cells often will depend on the purpose of the transformation. One of the major
20 purposes of transformation of crop plants is to add some commercially desirable, agronomically important traits to the plant. Such traits include, but are not limited to, herbicide resistance or tolerance; insect resistance or tolerance; disease resistance or tolerance (viral, bacterial, fungal, nematode); stress tolerance and/or resistance, as exemplified by resistance or tolerance to drought, heat, chilling, freezing, excessive
25 moisture, salt stress; oxidative stress; increased yields; food content and makeup; physical appearance; male sterility; drydown; standability; prolificacy; starch quantity and quality; oil quantity and quality; protein quality and quantity; amino acid composition; the production of a pharmaceutically active protein; the production of a small molecule with medicinal properties; the production of a chemical including those
30 with industrial utility; the production of nutraceuticals, carbohydrates, RNAs, lipids, fuels, dyes, pigments, vitamins, scents, flavors, vaccines, antibodies, hormones, and the like. Additionally one could create a library of an entire genome from any

organism or organelle including mammals, plants, microbes, fungi, bacteria, represented on minichromosomes. Furthermore one could incorporate a desired genomic segment such as one that includes a quantitative trait onto a minichromosome. One may desire to incorporate one or more genes conferring any
5 such desirable trait or traits, such as, for example, a gene or genes encoding herbicide resistance.

In certain embodiments, the present invention contemplates the transformation of a recipient cell with minichromosomes comprising more than one exogenous gene.
10 An "exogenous gene," can be a gene not normally found in the host genome in an identical context, or alternatively, the minichromosome could be used to introduce extra copies of host genes into a cell. The gene may be isolated from a different species than that of the host genome, or alternatively, isolated from the host genome but operably linked to one or more regulatory regions which differ from those found in
15 the unaltered, native gene. Two or more exogenous genes also can be supplied in a single transformation event using either distinct transgene-encoding vectors, or using a single vector incorporating two or more gene coding sequences. For example, plasmids bearing the *bar* and *aroA* expression units in either convergent, divergent, or colinear orientation, are considered to be particularly useful. Further preferred
20 combinations are those of an insect resistance gene, such as a Bt gene, along with a protease inhibitor gene such as *pinII*, or the use of *bar* in combination with either of the above genes. Of course, any two or more transgenes of any description, such as those conferring herbicide, insect, disease (viral, bacterial, fungal, nematode) or drought resistance, male sterility, drydown, standability, prolificacy, starch properties,
25 oil quantity and quality, modified chemical production, pharmaceutical or nutraceutical properties, bioremediation properties, increased biomass, altered growth rate, altered fitness, altered salinity tolerance, altered thermal tolerance, altered growth form, altered composition, altered metabolism, altered biodegradability, altered CO₂ fixation, altered stress tolerance, presence of bioindicator activity, altered digestibility
30 by humans or animals, altered allergenicity, altered mating characteristics, altered pollen dispersal, altered appearance, improved environmental impact, nitrogen fixation capability, or those increasing yield or nutritional quality may be employed as desired.

(i) *Herbicide Resistance*

The genes encoding phosphinothricin acetyltransferase (*bar* and *pat*), glyphosate tolerant EPSP synthase genes, the glyphosate degradative enzyme gene *gox* encoding glyphosate oxidoreductase, *deh* (encoding a dehalogenase enzyme that inactivates dalapon), herbicide resistant (*e.g.*, sulfonylurea and imidazolinone) acetolactate synthase, and *bxn* genes (encoding a nitrilase enzyme that degrades bromoxynil) are good examples of herbicide resistant genes for use in transformation. The *bar* and *pat* genes code for an enzyme, phosphinothricin acetyltransferase (PAT), which inactivates the herbicide phosphinothricin and prevents this compound from inhibiting glutamine synthetase enzymes. The enzyme 5-enolpyruvylshikimate 3-phosphate synthase (EPSP Synthase), is normally inhibited by the herbicide N-(phosphonomethyl)glycine (glyphosate). However, genes are known that encode glyphosate-resistant EPSP synthase enzymes. These genes are particularly contemplated for use in plant transformation. The *deh* gene encodes the enzyme dalapon dehalogenase and confers resistance to the herbicide dalapon. The *bxn* gene codes for a specific nitrilase enzyme that converts bromoxynil to a non-herbicidal degradation product.

20 (ii) *Insect Resistance*

Potential insect resistance genes that can be introduced include *Bacillus thuringiensis* crystal toxin genes or Bt genes (Watrud *et al.*, 1985). Bt genes may provide resistance to lepidopteran or coleopteran pests such as European Corn Borer (ECB). Preferred Bt toxin genes for use in such embodiments include the *CryIA(b)* and *CryIA(c)* genes. Endotoxin genes from other species of *B. thuringiensis* which affect insect growth or development also may be employed in this regard.

It is contemplated that preferred Bt genes for use in the transformation protocols disclosed herein will be those in which the coding sequence has been modified to effect increased expression in plants, and more particularly, in monocot plants. Means for preparing synthetic genes are well known in the art and are disclosed in, for example, U.S. Patent No. 5,500,365 and U.S. Patent Number No.

5,689,052, each of the disclosures of which are specifically incorporated herein by reference in their entirety. Examples of such modified Bt toxin genes include a synthetic Bt *CryIA(b)* gene (Perlak *et al.*, 1991), and the synthetic *CryIA(c)* gene termed 1800b (PCT Application WO 95/06128). Some examples of other Bt toxin

5 genes known to those of skill in the art are given in Table 1 below.

Table 1: *Bacillus thuringiensis* δ -Endotoxin Genes^a

New Nomenclature	Old Nomenclature	GenBank Accession
Cry1Aa	CryIA(a)	M11250
Cry1Ab	CryIA(b)	M13898
Cry1Ac	CryIA(c)	M11068
Cry1Ad	CryIA(d)	M73250
Cry1Ae	CryIA(e)	M65252
Cry1Ba	CryIB	X06711
Cry1Bb	ET5	L32020
Cry1Bc	PEG5	Z46442
Cry1Bd	CryE1	U70726
Cry1Ca	CryIC	X07518
Cry1Cb	CryIC(b)	M97880
Cry1Da	CryID	X54160
Cry1Db	PrtB	Z22511
Cry1Ea	CryIE	X53985
Cry1Eb	CryIE(b)	M73253
Cry1Fa	CryIF	M63897
Cry1Fb	PrtD	Z22512
Cry1Ga	PrtA	Z22510
Cry1Gb	CryH2	U70725
Cry1Ha	PrtC	Z22513
Cry1Hb		U35780
Cry1Ia	CryV	X62821
Cry1Ib	CryV	U07642

New Nomenclature	Old Nomenclature	GenBank Accession
Cry1Ja	ET4	L32019
Cry1Jb	ET1	U31527
Cry1K		U28801
Cry2Aa	CryIIA	M31738
Cry2Ab	CryIIB	M23724
Cry2Ac	CryIIC	X57252
Cry3A	CryIIIA	M22472
Cry3Ba	CryIIIB	X17123
Cry3Bb	CryIIIB2	M89794
Cry3C	CryIIID	X59797
Cry4A	CryIVA	Y00423
Cry4B	CryIVB	X07423
Cry5Aa	CryVA(a)	L07025
Cry5Ab	CryVA(b)	L07026
Cry6A	CryVIA	L07022
Cry6B	CryVIB	L07024
Cry7Aa	CryIIIC	M64478
Cry7Ab	CryIIICb	U04367
Cry8A	CryIIIE	U04364
Cry8B	CryIIIG	U04365
Cry8C	CryIIIF	U04366
Cry9A	CryIG	X58120
Cry9B	CryIX	X75019
Cry9C	CryIH	Z37527
Cry10A	CryIVC	M12662
Cry11A	CryIVD	M31737
Cry11B	Jeg80	X86902
Cry12A	CryVB	L07027
Cry13A	CryVC	L07023
Cry14A	CryVD	U13955

New Nomenclature	Old Nomenclature	GenBank Accession
Cry15A	34kDa	M76442
Cry16A	cbm71	X94146
Cry17A	cbm71	X99478
Cry18A	CryBP1	X99049
Cry19A	Jeg65	Y08920
Cyt1Aa	CytA	X03182
Cyt1Ab	CytM	X98793
Cyt2A	CytB	Z14147
Cyt2B	CytB	U52043

^aAdapted from: http://epunix.biols.susx.ac.uk/Home/Neil_Crickmore/Bt/index.html

Protease inhibitors also may provide insect resistance (Johnson *et al.*, 1989), and will thus have utility in plant transformation. The use of a protease inhibitor II gene, *pinII*, from tomato or potato is envisioned to be particularly useful. Even more advantageous is the use of a *pinII* gene in combination with a Bt toxin gene, the combined effect of which has been discovered to produce synergistic insecticidal activity. Other genes which encode inhibitors of the insect's digestive system, or those that encode enzymes or co-factors that facilitate the production of inhibitors, also may be useful. This group may be exemplified by oryzacystatin and amylase inhibitors such as those from wheat and barley.

Also, genes encoding lectins may confer additional or alternative insecticide properties. Lectins (originally termed phytohemagglutinins) are multivalent carbohydrate-binding proteins which have the ability to agglutinate red blood cells from a range of species. Lectins have been identified recently as insecticidal agents with activity against weevils, ECB and rootworm (Murdock *et al.*, 1990; Czapla & Lang, 1990). Lectin genes contemplated to be useful include, for example, barley and wheat germ agglutinin (WGA) and rice lectins (Gatehouse *et al.*, 1984), with WGA being preferred.

Genes controlling the production of large or small polypeptides active against insects when introduced into the insect pests, such as, *e.g.*, lytic peptides, peptide hormones and toxins and venoms, form another aspect of the invention. For example, it is contemplated that the expression of juvenile hormone esterase, directed towards
5 specific insect pests, also may result in insecticidal activity, or perhaps cause cessation of metamorphosis (Hammock *et al.*, 1990).

Transgenic plants expressing genes which encode enzymes that affect the integrity of the insect cuticle form yet another aspect of the invention. Such genes
10 include those encoding, *e.g.*, chitinase, proteases, lipases and also genes for the production of nikkomycin, a compound that inhibits chitin synthesis, the introduction of any of which is contemplated to produce insect resistant plants. Genes that code for activities that affect insect molting, such as those affecting the production of ecdysteroid UDP-glucosyl transferase, also fall within the scope of the useful
15 transgenes of the present invention.

Genes that code for enzymes that facilitate the production of compounds that reduce the nutritional quality of the host plant to insect pests also are encompassed by the present invention. It may be possible, for instance, to confer insecticidal activity
20 on a plant by altering its sterol composition. Sterols are obtained by insects from their diet and are used for hormone synthesis and membrane stability. Therefore alterations in plant sterol composition by expression of novel genes, *e.g.*, those that directly promote the production of undesirable sterols or those that convert desirable sterols into undesirable forms, could have a negative effect on insect growth and/or
25 development and hence endow the plant with insecticidal activity. Lipxygenases are naturally occurring plant enzymes that have been shown to exhibit anti-nutritional effects on insects and to reduce the nutritional quality of their diet. Therefore, further embodiments of the invention concern transgenic plants with enhanced lipxygenase activity which may be resistant to insect feeding.

30

Tripsacum dactyloides is a species of grass that is resistant to certain insects, including corn root worm. It is anticipated that genes encoding proteins that are toxic

to insects or are involved in the biosynthesis of compounds toxic to insects will be isolated from *Tripsacum* and that these novel genes will be useful in conferring resistance to insects. It is known that the basis of insect resistance in *Tripsacum* is genetic, because said resistance has been transferred to *Zea mays* via sexual crosses (Branson and Guss, 1972). It is further anticipated that other cereal, monocot or dicot plant species may have genes encoding proteins that are toxic to insects which would be useful for producing insect resistant plants.

Further genes encoding proteins characterized as having potential insecticidal activity also may be used as transgenes in accordance herewith. Such genes include, for example, the cowpea trypsin inhibitor (CpTI; Hilder *et al.*, 1987) which may be used as a rootworm deterrent; genes encoding avermectin (*Avermectin and Abamectin.*, Campbell, W.C., Ed., 1989; Ikeda *et al.*, 1987) which may prove particularly useful as a corn rootworm deterrent; ribosome inactivating protein genes; and even genes that regulate plant structures. Transgenic plants including anti-insect antibody genes and genes that code for enzymes that can convert a non-toxic insecticide (pro-insecticide) applied to the outside of the plant into an insecticide inside the plant also are contemplated.

(iii) *Environment or Stress Resistance*

Improvement of a plants ability to tolerate various environmental stresses such as, but not limited to, drought, excess moisture, chilling, freezing, high temperature, salt, and oxidative stress, also can be effected through expression of novel genes. It is proposed that benefits may be realized in terms of increased resistance to freezing temperatures through the introduction of an "antifreeze" protein such as that of the Winter Flounder (Cutler *et al.*, 1989) or synthetic gene derivatives thereof. Improved chilling tolerance also may be conferred through increased expression of glycerol-3-phosphate acetyltransferase in chloroplasts (Wolter *et al.*, 1992). Resistance to oxidative stress (often exacerbated by conditions such as chilling temperatures in combination with high light intensities) can be conferred by expression of superoxide dismutase (Gupta *et al.*, 1993), and may be improved by glutathione reductase (Bowler *et al.*, 1992). Such strategies may allow for tolerance to freezing in

newly emerged fields as well as extending later maturity higher yielding varieties to earlier relative maturity zones.

It is contemplated that the expression of novel genes that favorably effect plant water content, total water potential, osmotic potential, and turgor will enhance the ability of the plant to tolerate drought. As used herein, the terms "drought resistance" and "drought tolerance" are used to refer to a plants increased resistance or tolerance to stress induced by a reduction in water availability, as compared to normal circumstances, and the ability of the plant to function and survive in lower-water environments. In this aspect of the invention it is proposed, for example, that the expression of genes encoding for the biosynthesis of osmotically-active solutes, such as polyol compounds, may impart protection against drought. Within this class are genes encoding for mannitol-L-phosphate dehydrogenase (Lee and Saier, 1982) and trehalose-6-phosphate synthase (Kaasen *et al.*, 1992). Through the subsequent action of native phosphatases in the cell or by the introduction and coexpression of a specific phosphatase, these introduced genes will result in the accumulation of either mannitol or trehalose, respectively, both of which have been well documented as protective compounds able to mitigate the effects of stress. Mannitol accumulation in transgenic tobacco has been verified and preliminary results indicate that plants expressing high levels of this metabolite are able to tolerate an applied osmotic stress (Tarczynski *et al.*, 1992, 1993).

Similarly, the efficacy of other metabolites in protecting either enzyme function (*e.g.*, alanopine or propionic acid) or membrane integrity (*e.g.*, alanopine) has been documented (Loomis *et al.*, 1989), and therefore expression of genes encoding for the biosynthesis of these compounds might confer drought resistance in a manner similar to or complimentary to mannitol. Other examples of naturally occurring metabolites that are osmotically active and/or provide some direct protective effect during drought and/or desiccation include fructose, erythritol (Coxson *et al.*, 1992), sorbitol, dulcitol (Karsten *et al.*, 1992), glucosylglycerol (Reed *et al.*, 1984; ErdMann *et al.*, 1992), sucrose, stachyose (Koster and Leopold, 1988; Blackman *et al.*, 1992), raffinose (Bernal-Lugo and Leopold, 1992), proline

(Rensburg *et al.*, 1993), glycine betaine, ononitol and pinitol (Vernon and Bohnert, 1992). Continued canopy growth and increased reproductive fitness during times of stress will be augmented by introduction and expression of genes such as those controlling the osmotically active compounds discussed above and other such
5 compounds. Currently preferred genes which promote the synthesis of an osmotically active polyol compound are genes which encode the enzymes mannitol-1-phosphate dehydrogenase, trehalose-6-phosphate synthase and myoinositol 0-methyltransferase.

It is contemplated that the expression of specific proteins also may increase
10 drought tolerance. Three classes of Late Embryogenic Proteins have been assigned based on structural similarities (see Dure *et al.*, 1989). All three classes of LEAs have been demonstrated in maturing (*i.e.* desiccating) seeds. Within these 3 types of LEA proteins, the Type-II (dehydrin-type) have generally been implicated in drought and/or desiccation tolerance in vegetative plant parts (*i.e.* Mundy and Chua, 1988;
15 Piatkowski *et al.*, 1990; Yamaguchi-Shinozaki *et al.*, 1992). Recently, expression of a Type-III LEA (HVA-1) in tobacco was found to influence plant height, maturity and drought tolerance (Fitzpatrick, 1993). In rice, expression of the HVA-1 gene influenced tolerance to water deficit and salinity (Xu *et al.*, 1996). Expression of structural genes from all three LEA groups may therefore confer drought tolerance.
20 Other types of proteins induced during water stress include thiol proteases, aldolases and transmembrane transporters (Guerrero *et al.*, 1990), which may confer various protective and/or repair-type functions during drought stress. It also is contemplated that genes that effect lipid biosynthesis and hence membrane composition might also be useful in conferring drought resistance on the plant.

25

Many of these genes for improving drought resistance have complementary modes of action. Thus, it is envisaged that combinations of these genes might have additive and/or synergistic effects in improving drought resistance in plants. Many of these genes also improve freezing tolerance (or resistance); the physical stresses
30 incurred during freezing and drought are similar in nature and may be mitigated in similar fashion. Benefit may be conferred via constitutive expression of these genes, but the preferred means of expressing these novel genes may be through the use of a

turgor-induced promoter (such as the promoters for the turgor-induced genes described in Guerrero *et al.*, 1990 and Shagan *et al.*, 1993 which are incorporated herein by reference). Spatial and temporal expression patterns of these genes may enable plants to better withstand stress.

5

It is proposed that expression of genes that are involved with specific morphological traits that allow for increased water extractions from drying soil would be of benefit. For example, introduction and expression of genes that alter root characteristics may enhance water uptake. It also is contemplated that expression of genes that enhance reproductive fitness during times of stress would be of significant value. For example, expression of genes that improve the synchrony of pollen shed and receptiveness of the female flower parts, *i.e.*, silks, would be of benefit. In addition it is proposed that expression of genes that minimize kernel abortion during times of stress would increase the amount of grain to be harvested and hence be of value.

15

Given the overall role of water in determining yield, it is contemplated that enabling plants to utilize water more efficiently, through the introduction and expression of novel genes, will improve overall performance even when soil water availability is not limiting. By introducing genes that improve the ability of plants to maximize water usage across a full range of stresses relating to water availability, yield stability or consistency of yield performance may be realized.

20

(iv) *Disease Resistance*

It is proposed that increased resistance to diseases may be realized through introduction of genes into plants, for example, into monocotyledonous plants such as maize. It is possible to produce resistance to diseases caused by viruses, bacteria, fungi and nematodes. It also is contemplated that control of mycotoxin producing organisms may be realized through expression of introduced genes.

25

30

Resistance to viruses may be produced through expression of novel genes. For example, it has been demonstrated that expression of a viral coat protein in a

transgenic plant can impart resistance to infection of the plant by that virus and perhaps other closely related viruses (Cuozzo *et al.*, 1988, Hemenway *et al.*, 1988, Abel *et al.*, 1986). It is contemplated that expression of antisense genes targeted at essential viral functions may also impart resistance to viruses. For example, an antisense gene targeted at the gene responsible for replication of viral nucleic acid may inhibit replication and lead to resistance to the virus. It is believed that interference with other viral functions through the use of antisense genes also may increase resistance to viruses. Further, it is proposed that it may be possible to achieve resistance to viruses through other approaches, including, but not limited to the use of satellite viruses.

It is proposed that increased resistance to diseases caused by bacteria and fungi may be realized through introduction of novel genes. It is contemplated that genes encoding so-called "peptide antibiotics," pathogenesis related (PR) proteins, toxin resistance, and proteins affecting host-pathogen interactions such as morphological characteristics will be useful. Peptide antibiotics are polypeptide sequences which are inhibitory to growth of bacteria and other microorganisms. For example, the classes of peptides referred to as cecropins and magainins inhibit growth of many species of bacteria and fungi. It is proposed that expression of PR proteins in monocotyledonous plants such as maize may be useful in conferring resistance to bacterial disease. These genes are induced following pathogen attack on a host plant and have been divided into at least five classes of proteins (Bol, Linthorst, and Cornelissen, 1990). Included amongst the PR proteins are β -1, 3-glucanases, chitinases, and osmotin and other proteins that are believed to function in plant resistance to disease organisms. Other genes have been identified that have antifungal properties, *e.g.*, UDA (stinging nettle lectin) and hevein (Brookaert *et al.*, 1989; Barkai-Golan *et al.*, 1978). It is known that certain plant diseases are caused by the production of phytotoxins. It is proposed that resistance to these diseases would be achieved through expression of a novel gene that encodes an enzyme capable of degrading or otherwise inactivating the phytotoxin. It also is contemplated that expression of novel genes that alter the interactions between the host plant and pathogen may be useful in reducing the ability of the disease

organism to invade the tissues of the host plant, e.g., an increase in the waxiness of the leaf cuticle or other morphological characteristics.

(v) *Plant Agronomic Characteristics*

5 Two of the factors determining where crop plants can be grown are the average daily temperature during the growing season and the length of time between frosts. Within the areas where it is possible to grow a particular crop, there are varying limitations on the maximal time it is allowed to grow to maturity and be harvested. For example, a variety to be grown in a particular area is selected for its ability to
10 mature and dry down to harvestable moisture content within the required period of time with maximum possible yield. Therefore, crops of varying maturities is developed for different growing locations. Apart from the need to dry down sufficiently to permit harvest, it is desirable to have maximal drying take place in the field to minimize the amount of energy required for additional drying post-harvest.
15 Also, the more readily a product such as grain can dry down, the more time there is available for growth and kernel fill. It is considered that genes that influence maturity and/or dry down can be identified and introduced into plant lines using transformation techniques to create new varieties adapted to different growing locations or the same growing location, but having improved yield to moisture ratio at harvest. Expression
20 of genes that are involved in regulation of plant development may be especially useful.

It is contemplated that genes may be introduced into plants that would improve standability and other plant growth characteristics. Expression of novel genes in plants which confer stronger stalks, improved root systems, or prevent or reduce ear
25 droppage would be of great value to the farmer. It is proposed that introduction and expression of genes that increase the total amount of photoassimilate available by, for example, increasing light distribution and/or interception would be advantageous. In addition, the expression of genes that increase the efficiency of photosynthesis and/or the leaf canopy would further increase gains in productivity. It is contemplated that
30 expression of a phytochrome gene in crop plants may be advantageous. Expression of such a gene may reduce apical dominance, confer semidwarfism on a plant, and

increase shade tolerance (U.S. Patent No. 5,268,526). Such approaches would allow for increased plant populations in the field.

(vi) *Nutrient Utilization*

5 The ability to utilize available nutrients may be a limiting factor in growth of crop plants. It is proposed that it would be possible to alter nutrient uptake, tolerate pH extremes, mobilization through the plant, storage pools, and availability for metabolic activities by the introduction of novel genes. These modifications would allow a plant such as maize to more efficiently utilize available nutrients. It is
10 contemplated that an increase in the activity of, for example, an enzyme that is normally present in the plant and involved in nutrient utilization would increase the availability of a nutrient. An example of such an enzyme would be phytase. It is further contemplated that enhanced nitrogen utilization by a plant is desirable. Expression of a glutamate dehydrogenase gene in plants, *e.g.*, *E. coli gdhA* genes, may
15 lead to increased fixation of nitrogen in organic compounds. Furthermore, expression of *gdhA* in plants may lead to enhanced resistance to the herbicide glufosinate by incorporation of excess ammonia into glutamate, thereby detoxifying the ammonia. It also is contemplated that expression of a novel gene may make a nutrient source available that was previously not accessible, *e.g.*, an enzyme that releases a component
20 of nutrient value from a more complex molecule, perhaps a macromolecule.

(vii) *Male Sterility*

Male sterility is useful in the production of hybrid seed. It is proposed that male sterility may be produced through expression of novel genes. For example, it has
25 been shown that expression of genes that encode proteins that interfere with development of the male inflorescence and/or gametophyte result in male sterility. Chimeric ribonuclease genes that express in the anthers of transgenic tobacco and oilseed rape have been demonstrated to lead to male sterility (Mariani *et al.*, 1990).

30 A number of mutations were discovered in maize that confer cytoplasmic male sterility. One mutation in particular, referred to as T cytoplasm, also correlates with sensitivity to Southern corn leaf blight. A DNA sequence, designated TURF-13

(Levings, 1990), was identified that correlates with T cytoplasm. It is proposed that it would be possible through the introduction of TURF-13 via transformation, to separate male sterility from disease sensitivity. As it is necessary to be able to restore male fertility for breeding purposes and for grain production, it is proposed that genes
5 encoding restoration of male fertility also may be introduced.

(viii) Improved Nutritional Content

Genes may be introduced into plants to improve the nutrient quality or content of a particular crop. Introduction of genes that alter the nutrient composition of a crop
10 may greatly enhance the feed or food value. For example, the protein of many grains is suboptimal for feed and food purposes, especially when fed to pigs, poultry, and humans. The protein is deficient in several amino acids that are essential in the diet of these species, requiring the addition of supplements to the grain. Limiting essential amino acids may include lysine, methionine, tryptophan, threonine, valine, arginine,
15 and histidine. Some amino acids become limiting only after corn is supplemented with other inputs for feed formulations. The levels of these essential amino acids in seeds and grain may be elevated by mechanisms which include, but are not limited to, the introduction of genes to increase the biosynthesis of the amino acids, decrease the degradation of the amino acids, increase the storage of the amino acids in proteins, or
20 increase transport of the amino acids to the seeds or grain.

The protein composition of a crop may be altered to improve the balance of amino acids in a variety of ways including elevating expression of native proteins, decreasing expression of those with poor composition, changing the composition of
25 native proteins, or introducing genes encoding entirely new proteins possessing superior composition.

The introduction of genes that alter the oil content of a crop plant may also be of value. Increases in oil content may result in increases in metabolizable-energy-
30 content and density of the seeds for use in feed and food. The introduced genes may encode enzymes that remove or reduce rate-limitations or regulated steps in fatty acid or lipid biosynthesis. Such genes may include, but are not limited to, those that encode

acetyl-CoA carboxylase, ACP-acyltransferase, β -ketoacyl-ACP synthase, plus other well known fatty acid biosynthetic activities. Other possibilities are genes that encode proteins that do not possess enzymatic activity such as acyl carrier protein. Genes may be introduced that alter the balance of fatty acids present in the oil providing a more
5 healthful or nutritive feedstuff. The introduced DNA also may encode sequences that block expression of enzymes involved in fatty acid biosynthesis, altering the proportions of fatty acids present in crops.

Genes may be introduced that enhance the nutritive value of the starch
10 component of crops, for example by increasing the degree of branching, resulting in improved utilization of the starch in livestock by delaying its metabolism. Additionally, other major constituents of a crop may be altered, including genes that affect a variety of other nutritive, processing, or other quality aspects. For example, pigmentation may be increased or decreased.

15

Feed or food crops may also possess sub-optimal quantities of vitamins, antioxidants or other nutraceuticals, requiring supplementation to provide adequate nutritive value and ideal health value. Introduction of genes that enhance vitamin biosynthesis may be envisioned including, for example, vitamins A, E, B₁₂, choline,
20 and the like. Mineral content may also be sub-optimal. Thus genes that affect the accumulation or availability of compounds containing phosphorus, sulfur, calcium, manganese, zinc, and iron among others would be valuable.

Numerous other examples of improvements of crops may be used with the
25 invention. The improvements may not necessarily involve grain, but may, for example, improve the value of a crop for silage. Introduction of DNA to accomplish this might include sequences that alter lignin production such as those that result in the "brown midrib" phenotype associated with superior feed value for cattle.

30 In addition to direct improvements in feed or food value, genes also may be introduced which improve the processing of crops and improve the value of the products resulting from the processing. One use of crops is via wetmilling. Thus

novel genes that increase the efficiency and reduce the cost of such processing, for example by decreasing steeping time, may also find use. Improving the value of wetmilling products may include altering the quantity or quality of starch, oil, corn gluten meal, or the components of gluten feed. Elevation of starch may be achieved
5 through the identification and elimination of rate limiting steps in starch biosynthesis or by decreasing levels of the other components of crops resulting in proportional increases in starch.

Oil is another product of wetmilling, the value of which may be improved by
10 introduction and expression of genes. Oil properties may be altered to improve its performance in the production and use of cooking oil, shortenings, lubricants or other oil-derived products or improvement of its health attributes when used in the food-related applications. Novel fatty acids also may be synthesized which upon extraction can serve as starting materials for chemical syntheses. The changes in oil properties
15 may be achieved by altering the type, level, or lipid arrangement of the fatty acids present in the oil. This in turn may be accomplished by the addition of genes that encode enzymes that catalyze the synthesis of novel fatty acids and the lipids possessing them or by increasing levels of native fatty acids while possibly reducing levels of precursors. Alternatively, DNA sequences may be introduced which slow or
20 block steps in fatty acid biosynthesis resulting in the increase in precursor fatty acid intermediates. Genes that might be added include desaturases, epoxidases, hydratases, dehydratases, and other enzymes that catalyze reactions involving fatty acid intermediates. Representative examples of catalytic steps that might be blocked include the desaturations from stearic to oleic acid and oleic to linolenic acid resulting
25 in the respective accumulations of stearic and oleic acids. Another example is the blockage of elongation steps resulting in the accumulation of C₈ to C₁₂ saturated fatty acids.

(ix) *Production or Assimilation of Chemicals or Biologicals*

It may further be considered that a transgenic plant prepared in accordance with the invention may be used for the production or manufacturing of useful biological compounds that were either not produced at all, or not produced at the same
5 level, in the corn plant previously. Alternatively, plants produced in accordance with the invention may be made to metabolize or absorb and concentrate certain compounds, such as hazardous wastes, thereby allowing bioremediation of these compounds.

10 The novel plants producing these compounds are made possible by the introduction and expression of one or potentially many genes with the constructs provided by the invention. The vast array of possibilities include but are not limited to any biological compound which is presently produced by any organism such as proteins, nucleic acids, primary and intermediary metabolites, carbohydrate polymers,
15 enzymes for uses in bioremediation, enzymes for modifying pathways that produce secondary plant metabolites such as flavonoids or vitamins, enzymes that could produce pharmaceuticals, and for introducing enzymes that could produce compounds of interest to the manufacturing industry such as specialty chemicals and plastics. The compounds may be produced by the plant, extracted upon harvest and/or processing,
20 and used for any presently recognized useful purpose such as pharmaceuticals, fragrances, and industrial enzymes to name a few.

(x) *Non-Protein-Expressing Sequences*

DNA may be introduced into plants for the purpose of expressing RNA
25 transcripts that function to affect plant phenotype yet are not translated into protein. Two examples are antisense RNA and RNA with ribozyme activity. Both may serve possible functions in reducing or eliminating expression of native or introduced plant genes. However, as detailed below, DNA need not be expressed to effect the phenotype of a plant.

30

1. Antisense RNA

Genes may be constructed or isolated, which when transcribed, produce antisense RNA that is complementary to all or part(s) of a targeted messenger RNA(s). The antisense RNA reduces production of the polypeptide product of the messenger RNA. Genes may also be constructed to produce double-stranded RNA molecules complementary to all or part of the targeted messenger RNA(s). Genes designed in this manner will be referred to as RNAi constructs; the double-stranded RNA or RNAi constructs can trigger the sequence-specific degradation of the target messenger RNA. The polypeptide product of the target messenger RNA may be any protein. The
10 aforementioned genes will be referred to as antisense genes and RNAi constructs, respectively. An antisense gene or RNAi construct may thus be introduced into a plant by transformation methods to produce a novel transgenic plant with reduced expression of a selected protein of interest. For example, the protein may be an enzyme that catalyzes a reaction in the plant. Reduction of the enzyme activity may
15 reduce or eliminate products of the reaction which include any enzymatically synthesized compound in the plant such as fatty acids, amino acids, carbohydrates, nucleic acids and the like. Alternatively, the protein may be a storage protein, such as a zein, or a structural protein, the decreased expression of which may lead to changes in seed amino acid composition or plant morphological changes respectively. The
20 possibilities cited above are provided only by way of example and do not represent the full range of applications.

2. Ribozymes

Genes also may be constructed or isolated, which when transcribed, produce
25 RNA enzymes (ribozymes) which can act as endoribonucleases and catalyze the cleavage of RNA molecules with selected sequences. The cleavage of selected messenger RNAs can result in the reduced production of their encoded polypeptide products. These genes may be used to prepare novel transgenic plants which possess them. The transgenic plants may possess reduced levels of polypeptides including, but
30 not limited to, the polypeptides cited above.

Ribozymes are RNA-protein complexes that cleave nucleic acids in a site-specific fashion. Ribozymes have specific catalytic domains that possess endonuclease activity (Kim and Cech, 1987; Gerlach *et al.*, 1987; Forster and Symons, 1987). For example, a large number of ribozymes accelerate phosphoester transfer
5 reactions with a high degree of specificity, often cleaving only one of several phosphoesters in an oligonucleotide substrate (Cech *et al.*, 1981; Michel and Westhof, 1990; Reinhold-Hurek and Shub, 1992). This specificity has been attributed to the requirement that the substrate bind *via* specific base-pairing interactions to the internal guide sequence ("IGS") of the ribozyme prior to chemical reaction.

10

Ribozyme catalysis has primarily been observed as part of sequence-specific cleavage/ligation reactions involving nucleic acids (Joyce, 1989; Cech *et al.*, 1981). For example, U. S. Patent 5,354,855 reports that certain ribozymes can act as endonucleases with a sequence specificity greater than that of known ribonucleases
15 and approaching that of the DNA restriction enzymes.

Several different ribozyme motifs have been described with RNA cleavage activity (Symons, 1992). Examples include sequences from the Group I self splicing introns including Tobacco Ringspot Virus (Prody *et al.*, 1986), Avocado Sunblotch
20 Viroid (Palukaitis *et al.*, 1979; Symons, 1981), and Lucerne Transient Streak Virus (Forster and Symons, 1987). Sequences from these and related viruses are referred to as hammerhead ribozyme based on a predicted folded secondary structure.

Other suitable ribozymes include sequences from RNase P with RNA cleavage
25 activity (Yuan *et al.*, 1992, Yuan and Altman, 1994, U. S. Patents 5,168,053 and 5,624,824), hairpin ribozyme structures (Berzal-Herranz *et al.*, 1992; Chowrira *et al.*, 1993) and Hepatitis Delta virus based ribozymes (U. S. Patent 5,625,047). The general design and optimization of ribozyme directed RNA cleavage activity has been discussed in detail (Haseloff and Gerlach, 1988, Symons, 1992,
30 Chowrira *et al.*, 1994; Thompson *et al.*, 1995).

The other variable on ribozyme design is the selection of a cleavage site on a given target RNA. Ribozymes are targeted to a given sequence by virtue of annealing to a site by complimentary base pair interactions. Two stretches of homology are required for this targeting. These stretches of homologous sequences flank the catalytic ribozyme structure defined above. Each stretch of homologous sequence can vary in length from 7 to 15 nucleotides. The only requirement for defining the homologous sequences is that, on the target RNA, they are separated by a specific sequence which is the cleavage site. For hammerhead ribozyme, the cleavage site is a dinucleotide sequence on the target RNA is a uracil (U) followed by either an adenine, cytosine or uracil (A,C or U) (Perriman *et al.*, 1992; Thompson *et al.*, 1995). The frequency of this dinucleotide occurring in any given RNA is statistically 3 out of 16. Therefore, for a given target messenger RNA of 1,000 bases, 187 dinucleotide cleavage sites are statistically possible.

Designing and testing ribozymes for efficient cleavage of a target RNA is a process well known to those skilled in the art. Examples of scientific methods for designing and testing ribozymes are described by Chowrira *et al.*, (1994) and Lieber and Strauss (1995), each incorporated by reference. The identification of operative and preferred sequences for use in down regulating a given gene is simply a matter of preparing and testing a given sequence, and is a routinely practiced "screening" method known to those of skill in the art.

3. Induction of Gene Silencing

It also is possible that genes may be introduced to produce novel transgenic plants which have reduced expression of a native gene product by the mechanism of co-suppression. It has been demonstrated in tobacco, tomato, and petunia (Goring *et al.*, 1991; Smith *et al.*, 1990; Napoli *et al.*, 1990; van der Krol *et al.*, 1990) that expression of the sense transcript of a native gene will reduce or eliminate expression of the native gene in a manner similar to that observed for antisense genes. The introduced gene may encode all or part of the targeted native protein but its translation may not be required for reduction of levels of that native protein.

4. Non-RNA-Expressing Sequences

DNA elements including those of transposable elements such as Ds, Ac, or Mu, may be inserted into a gene to cause mutations. These DNA elements may be inserted in order to inactivate (or activate) a gene and thereby "tag" a particular trait. In this instance the transposable element does not cause instability of the tagged mutation, because the utility of the element does not depend on its ability to move in the genome. Once a desired trait is tagged, the introduced DNA sequence may be used to clone the corresponding gene, *e.g.*, using the introduced DNA sequence as a PCR primer together with PCR gene cloning techniques (Shapiro, 1983; Dellaporta *et al.*, 1988). Once identified, the entire gene(s) for the particular trait, including control or regulatory regions where desired, may be isolated, cloned and manipulated as desired. The utility of DNA elements introduced into an organism for purposes of gene tagging is independent of the DNA sequence and does not depend on any biological activity of the DNA sequence, *i.e.*, transcription into RNA or translation into protein. The sole function of the DNA element is to disrupt the DNA sequence of a gene.

It is contemplated that unexpressed DNA sequences, including novel synthetic sequences, could be introduced into cells as proprietary "labels" of those cells and plants and seeds thereof. It would not be necessary for a label DNA element to disrupt the function of a gene endogenous to the host organism, as the sole function of this DNA would be to identify the origin of the organism. For example, one could introduce a unique DNA sequence into a plant and this DNA element would identify all cells, plants, and progeny of these cells as having arisen from that labeled source. It is proposed that inclusion of label DNAs would enable one to distinguish proprietary germplasm or germplasm derived from such, from unlabelled germplasm.

Another possible element which may be introduced is a matrix attachment region element (MAR), such as the chicken lysozyme A element (Stief, 1989), which can be positioned around an expressible gene of interest to effect an increase in overall expression of the gene and diminish position dependent effects upon incorporation into the plant genome (Stief *et al.*, 1989; Phi-Van *et al.*, 1990).

5. Other

Other examples of non-protein expressing sequences specifically envisioned for use with the invention include tRNA sequences, for example, to alter codon usage, and rRNA variants, for example, which may confer resistance to various agents such as antibiotics.

XI. Biological Functional Equivalents

Modification and changes may be made in the centromeric DNA segments of the current invention and still obtain a functional molecule with desirable characteristics. The following is a discussion based upon changing the nucleic acids of a centromere to create an equivalent, or even an improved, second-generation molecule.

In particular embodiments of the invention, mutated centromeric sequences are contemplated to be useful for increasing the utility of the centromere. It is specifically contemplated that the function of the centromeres of the current invention may be based upon the secondary structure of the DNA sequences of the centromere, modification of the DNA with methyl groups or other adducts, and / or the proteins which interact with the centromere. By changing the DNA sequence of the centromere, one may alter the affinity of one or more centromere-associated protein(s) for the centromere and / or the secondary structure or modification of the centromeric sequences, thereby changing the activity of the centromere. Alternatively, changes may be made in the centromeres of the invention which do not affect the activity of the centromere. Changes in the centromeric sequences which reduce the size of the DNA segment needed to confer centromere activity are contemplated to be particularly useful in the current invention, as would changes which increased the fidelity with which the centromere was transmitted during mitosis and meiosis.

XII. Plants

The term "plant," as used herein, refers to any type of plant. The inventors have provided below an exemplary description of some plants that may be used with

the invention. However, the list is not in any way limiting, as other types of plants will be known to those of skill in the art and could be used with the invention.

A common class of plants exploited in agriculture are vegetable crops, including artichokes, kohlrabi, arugula, leeks, asparagus, lettuce (*e.g.*, head, leaf, romaine), bok choy, malanga, broccoli, melons (*e.g.*, muskmelon, watermelon, crenshaw, honeydew, cantaloupe), brussels sprouts, cabbage, cardoni, carrots, napa, cauliflower, okra, onions, celery, parsley, chick peas, parsnips, chicory, chinese cabbage, peppers, collards, potatoes, cucumber plants (marrows, cucumbers), pumpkins, cucurbits, radishes, dry bulb onions, rutabaga, eggplant, salsify, escarole, shallots, endive, garlic, spinach, green onions, squash, greens, beet (sugar beet and fodder beet), sweet potatoes, swiss chard, horseradish, tomatoes, kale, turnips, and spices.

Other types of plants frequently finding commercial use include fruit and vine crops such as apples, apricots, cherries, nectarines, peaches, pears, plums, prunes, quince almonds, chestnuts, filberts, pecans, pistachios, walnuts, citrus, blueberries, boysenberries, cranberries, currants, loganberries, raspberries, strawberries, blackberries, grapes, avocados, bananas, kiwi, persimmons, pomegranate, pineapple, tropical fruits, pomes, melon, mango, papaya, and lychee.

Many of the most widely grown plants are field crop plants such as evening primrose, meadow foam, corn (field, sweet, popcorn), hops, jojoba, peanuts, rice, safflower, small grains (barley, oats, rye, wheat, etc.), sorghum, tobacco, kapok, leguminous plants (beans, lentils, peas, soybeans), oil plants (rape, mustard, poppy, olives, sunflowers, coconut, castor oil plants, cocoa beans, groundnuts), fibre plants (cotton, flax, hemp, jute), lauraceae (cinnamon, camphor), or plants such as coffee, sugarcane, tea, and natural rubber plants.

Still other examples of plants include bedding plants such as flowers, cactus, succulents and ornamental plants, as well as trees such as forest (broad-leaved trees

and evergreens, such as conifers), fruit, ornamental, and nut-bearing trees, as well as shrubs and other nursery stock.

XIII. Definitions

5 As used herein, the terms “autonomous replicating sequence” or “ARS” or “origin of replication ” refer to an origin of DNA replication recognized by proteins that initiate DNA replication.

10 As used herein, the terms “binary BAC” or “binary bacterial artificial chromosome” refer to a bacterial vector that contains the T-DNA border sequences necessary for *Agrobacterium* mediated transformation (see, for example, Hamilton *et al.*, 1996; Hamilton, 1997; and Liu *et al.*, 1999).

15 As used herein, the term "candidate centromere sequence" refers to a nucleic acid sequence which one wishes to assay for potential centromere function.

20 As used herein, a “centromere” is any DNA sequence that confers an ability to segregate to daughter cells through cell division. In one context, this sequence may produce a segregation efficiency to daughter cells ranging from about 1% to about 100%, including to about 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or about 95% of daughter cells. Variations in such a segregation efficiency may find important applications within the scope of the invention; for example, minichromosomes carrying centromeres that confer 100% stability could be maintained in all daughter cells without selection, while those that confer 1% stability
25 could be temporarily introduced into a transgenic organism, but be eliminated when desired. In particular embodiments of the invention, the centromere may confer stable segregation of a nucleic acid sequence, including a recombinant construct comprising the centromere, through mitotic or meiotic divisions, including through both meiotic and mitotic divisions. A plant centromere is not necessarily derived from plants, but
30 has the ability to promote DNA segregation in plant cells.

As used herein, the term "centromere-associated protein" refers to a protein encoded by a sequence of the centromere or a protein which is encoded by host DNA and binds with relatively high affinity to the centromere.

5 As used herein, the term "circular permutations" refer to variants of a sequence that begin at base n within the sequence, proceed to the end of the sequence, resume with base number one of the sequence, and proceed to base $n - 1$. For this analysis, n may be any number less than or equal to the length of the sequence. For example, circular permutations of the sequence ABCD are: ABCD, BCDA, CDAB, and DABC.

10

As used herein, the term "crop" includes any plant or portion of a plant grown or harvested for commercial or beneficial purposes.

15 As used herein, "eukaryote" refers to living organisms whose cells contain nuclei. A eukaryote may be distinguished from a "prokaryote" which is an organism which lacks nuclei. Prokaryotes and eukaryotes differ fundamentally in the way their genetic information is organized, as well as their patterns of RNA and protein synthesis.

20 As used herein, the term "expression" refers to the process by which a structural gene produces an RNA molecule, typically termed messenger RNA (mRNA). The mRNA is typically, but not always, translated into polypeptide(s).

25 As used herein, the term "genome" refers to all of the genes and DNA sequences that comprise the genetic information within a given cell of an organism. Usually, this is taken to mean the information contained within the nucleus, but also includes the organelles.

30 As used herein, the term "higher eukaryote" means a multicellular eukaryote, typically characterized by its greater complex physiological mechanisms and relatively large size. Generally, complex organisms such as plants and animals are included in this category. Preferred higher eukaryotes to be transformed by the present invention

include, for example, monocot and dicot angiosperm species, gymnosperm species, fern species, plant tissue culture cells of these species, animal cells and algal cells. It will of course be understood that prokaryotes and eukaryotes alike may be transformed by the methods of this invention.

5

As used herein, the term "host" refers to any organism that contains a plasmid, expression vector, or integrated construct comprising a plant centromere. Preferred examples of host cells for cloning, useful in the present invention, are bacteria such as *Escherichia coli*, *Bacillus subtilis*, *Pseudomonas*, *Streptomyces*, *Salmonella*, and yeast cells such as *S. cerevisiae*. Host cells which can be targeted for expression of a minichromosome may be plant cells of any source and specifically include *Arabidopsis*, maize, rice, sugarcane, sorghum, barley, soybeans, tobacco, wheat, tomato, potato, citrus, or any other agronomically or scientifically important species.

15 As used herein, the term "hybridization" refers to the pairing of complementary RNA and DNA strands to produce an RNA-DNA hybrid, or alternatively, the pairing of two DNA single strands from genetically different or the same sources to produce a double stranded DNA molecule.

20 As used herein, the term "linker" refers to a DNA molecule, generally up to 50 or 60 nucleotides long and synthesized chemically, or cloned from other vectors. In a preferred embodiment, this fragment contains one, or preferably more than one, restriction enzyme site for a blunt-cutting enzyme and a staggered-cutting enzyme, such as *Bam*HI. One end of the linker fragment is adapted to be ligatable to one end of the linear molecule and the other end is adapted to be ligatable to the other end of the linear molecule.

25 As used herein, a "library" is a pool of random DNA fragments which are cloned. In principle, any gene can be isolated by screening the library with a specific hybridization probe (see, for example, Young *et al.*, 1977). Each library may contain the DNA of a given organism inserted as discrete restriction enzyme-generated fragments or as randomly sheered fragments into many thousands of plasmid vectors.

For purposes of the present invention, *E. coli*, yeast, and *Salmonella* plasmids are particularly useful when the genome inserts come from other organisms.

As used herein, the term "lower eukaryote" refers to a eukaryote characterized by a comparatively simple physiology and composition, and most often unicellularity. Examples of lower eukaryotes include flagellates, ciliates, and yeast.

As used herein, a "minichromosome" is a recombinant DNA construct including a centromere and capable of transmission to daughter cells. Minichromosome may remain separate from the host genome (as episomes) or may integrate into host chromosomes. The stability of this construct through cell division could range between from about 1% to about 100%, including about 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% and about 95%. The minichromosome construct may be a circular or linear molecule. It may include elements such as one or more telomeres, ARS sequences, and genes. The number of such sequences included is only limited by the physical size limitations of the construct itself. It could contain DNA derived from a natural centromere, although it may be preferable to limit the amount of DNA to the minimal amount required to obtain a segregation efficiency in the range of 1-100%. The minichromosome could also contain a synthetic centromere composed of tandem arrays of repeats of any sequence, either derived from a natural centromere, or of synthetic DNA. The minichromosome could also contain DNA derived from multiple natural centromeres. The minichromosome may be inherited through mitosis or meiosis, or through both meiosis and mitosis. As used herein, the term minichromosome specifically encompasses and includes the terms "plant artificial chromosome" or "PLAC," or engineered chromosomes or microchromosomes and all teachings relevant to a PLAC or plant artificial chromosome specifically apply to constructs within the meaning of the term minichromosome.

As used herein, by "minichromosome-encoded protein" it is meant a polypeptide which is encoded by a sequence of a minichromosome of the current invention. This includes sequences such as selectable markers, telomeres, *etc.*, as well

as those proteins encoded by any other selected functional genes on the minichromosome.

As used herein, the term "plant" includes plant cells, plant protoplasts, plant calli, and the like, as well as whole plants regenerated therefrom.

As used herein, the term "plasmid" or "cloning vector" refers to a closed covalently circular extrachromosomal DNA or linear DNA which is able to replicate in a host cell and which is normally nonessential to the survival of the cell. A wide variety of plasmids and other vectors are known and commonly used in the art (see, for example, Cohen *et al.*, U.S. Patent No. 4,468,464, which discloses examples of DNA plasmids, and which is specifically incorporated herein by reference).

As used herein, a "probe" is any biochemical reagent (usually tagged in some way for ease of identification), used to identify or isolate a gene, a gene product, a DNA segment or a protein.

As used herein, the term "recombination" refers to any genetic exchange that involves breaking and rejoining of DNA strands.

As used herein the term "regulatory sequence" refers to any DNA sequence that influences the efficiency of transcription or translation of any gene. The term includes, but is not limited to, sequences comprising promoters, enhancers and terminators.

As used herein, a "selectable marker" is a gene whose presence results in a clear phenotype, and most often a growth advantage for cells that contain the marker. This growth advantage may be present under standard conditions, altered conditions such as elevated temperature, or in the presence of certain chemicals such as herbicides or antibiotics. Use of selectable markers is described, for example, in Broach *et al.* (1979). Examples of selectable markers include the thymidine kinase gene, the cellular adenine-phosphoribosyltransferase gene and the dihydrolfolate

reductase gene, hygromycin phosphotransferase genes, the bar gene and neomycin phosphotransferase genes, among others. Preferred selectable markers in the present invention include genes whose expression confer antibiotic or herbicide resistance to the host cell, sufficient to enable the maintenance of a vector within the host cell, and
5 which facilitate the manipulation of the plasmid into new host cells. Of particular interest in the present invention are proteins conferring cellular resistance to ampicillin, chloramphenicol, tetracycline, G-418, bialaphos, and glyphosate for example.

10 As used herein, a "screenable marker" is a gene whose presence results in an identifiable phenotype. This phenotype may be observable under standard conditions, altered conditions such as elevated temperature, or in the presence of certain chemicals used to detect the phenotype.

15 As used herein, the term "site-specific recombination" refers to any genetic exchange that involves breaking and rejoining of DNA strands at a specific DNA sequence.

As used herein, a "structural gene" is a sequence which codes for a polypeptide
20 or RNA and includes 5' and 3' ends. The structural gene may be from the host into which the structural gene is transformed or from another species. A structural gene will preferably, but not necessarily, include one or more regulatory sequences which modulate the expression of the structural gene, such as a promoter, terminator or enhancer. A structural gene will preferably, but not necessarily, confer some useful
25 phenotype upon an organism comprising the structural gene, for example, herbicide resistance. In one embodiment of the invention, a structural gene may encode an RNA sequence which is not translated into a protein, for example a tRNA or rRNA gene.

As used herein, the term "telomere" refers to a sequence capable of capping the
30 ends of a chromosome, thereby preventing degradation of the chromosome end, ensuring replication and preventing fusion to other chromosome sequences.

Telomeres can include naturally occurring telomere sequences or synthetic sequences. Telomeres from one species may confer telomere activity in another species.

As used herein, the terms "transformation" or "transfection" refer to the acquisition in cells of new DNA sequences through the chromosomal or extra-chromosomal addition of DNA. This is the process by which naked DNA, DNA coated with protein, or whole minichromosomes are introduced into a cell, resulting in a potentially heritable change.

As used herein the term "consensus" refers to a nucleic acid sequence derived by comparing two or more related sequences. A consensus sequence defines both the conserved and variable sites between the sequences being compared. Any one of the sequences used to derive the consensus or any permutation defined by the consensus may be useful in construction minichromosomes.

As used herein the term "repeated nucleotide sequence" refers to any nucleic acid sequence of at least 25 bp present in a genome or a recombinant molecule that occurs at least two or more times and that are preferably at least 80% identical either in head to tail or head to head orientation either with or without intervening sequence between repeat units.

XIV. Examples

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skilled the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventors to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the

same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

5

EXAMPLE 1

Isolation of Genomic DNA

Tissue from various plants are harvested for DNA extraction. For DNA extraction, leaf tissue is cooled in liquid nitrogen, ground to a fine powder and
10 transferred to an organic solvent-resistant test tube or beaker. Warm CTAB extraction solution (2% (w/v) CTAB, 100 mM Tris-Cl, pH 9.5, 20 mM EDTA, pH 8.0, 1.4 M NaCl, 1% polyethylene glycol) is added in a ratio of 20 ml per gram of tissue and mixed thoroughly. For each 20 ml extraction buffer, 50 microliters of β -mercaptoethanol and 30 microliters of 30 mg/ml RNase A are added and the mixture
15 is incubated for 10-60 min. at 65°C with occasional mixing. The homogenate is extracted with an equal volume of chloroform, and is then centrifuged 5 min at 7500 x g (8000 rpm in JA20; 10,000 rpm in a microcentrifuge, for smaller samples), 4°C. The top (aqueous) phase is recovered and nucleic acids are precipitated by adding 1 volume isopropanol. After mixing, the precipitate is pelleted at 15 min at 7500 x g, 4°C. The
20 pellet is washed with 70% ethanol, dried and resuspended in a minimal volume of TE (10 mM Tris-Cl, pH 8.0, 0.1 mM EDTA, pH 8.0).

EXAMPLE 2

Brassica oleracea centromere repeat sequences

25

We purified repetitive sequences from *Brassica oleracea* (*Brassica oleracea* fast plants, obtained from the Wisconsin Crucifer Cooperative). We set forth herein two centromere repeats, termed ChrBo1 and ChrBo2. We determined the consensus of each repeat as described in Example 6.

30

The consensus sequence of ChrBo1 is shown in FIG. 1A (SEQ ID NO:1). This consensus was assembled from DNA sequences collected by the inventors. Twenty-

four of these sequences completely spanned the repeat, and nine others partially covered the repeat. The length of this repeat is 180 ± 0.86 base pairs, and A and T comprise of 60% of the consensus.

5 The consensus sequence of ChrBo2 is shown in FIG. 1B (SEQ ID NO:2). This consensus was assembled from DNA sequences collected by the inventors. Five of these sequences completely spanned the repeat, and two others partially covered the repeat. The length of this repeat is 180 ± 0.45 base pairs, and A and T comprise 63% of the consensus.

10

 The two repeats (ChrBo1 and ChrBo2) were aligned to each other using the ClustalX program (ClustalX is a free multiple sequence alignment program for Windows. Thompson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F. and Higgins, D.G. (1997) The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. Nucleic Acids Research, 24:4876-4882.).
15 The two consensus sequences differ significantly from each other at several bases. Those sites with significant differences (chi-squared, $P < 0.05$) are highlighted as shown in FIG. 1C.

20 The GenBank nt database (<http://ftp.ncbi.nlm.nih.gov/blast/db/>, March 29 version, downloaded on 04/07/2002) and the plant satellite DNA database (<http://w3lamc.umbr.cas.cz/PlantSat/>, downloaded on 4/14/2002) were compared to the inventors' consensus sequences using the blastn program and an Expect value threshold score of -3. Consensus sequences were assembled using all inventors' and
25 GenBank sequences that matched with an Expect (E) value of less than -45.

 The revised consensus sequence of ChrBo1 is shown in FIG. 1D (SEQ ID NO:3). This consensus was assembled from thirty-three DNA sequences collected by the inventors and eighteen GenBank sequences (Table 10). Thirty of these sequences
30 completely spanned the repeat, and twenty-one others partially covered the repeat. The length of this repeat is 180 ± 0.81 base pairs, and A and T comprise of 59% of the consensus.

Table 1. GenBank sequences (accession numbers) that match inventors' ChrBo1 consensus

M30962	M30963	M31436	M31435
M31438	M31434	M31439	M31437
X68786	X12736	X07519	X16589
X15291	X68783	X68784	X61583
AJ228348	Z22947		

5

The revised consensus sequence of ChrBo2 is shown in FIG. 1E (SEQ ID NO:4). This consensus was assembled from seven DNA sequences collected by the inventors and five GenBank sequences (Table 2). Seven of these sequences completely spanned the repeat, and five others partially covered the repeat. The length of this repeat is 180 ± 0.44 base pairs, and A and T comprise of 63% of the consensus.

10

Table 2. GenBank sequences (accession numbers) that match inventors' ChrBo2 consensus

15

AJ228347	M30962	X12736
X61583	X68785	

The two revised consensus sequences (ChrBo1 and ChrBo2) were aligned to each other using the ClustalX program. The two consensus sequences differ significantly (chi-squared, $P < 0.05$) from each other at several bases (highlighted as shown in FIG. 1F).

20

A total of 20 GenBank entries match the *Brassica oleracea* centromere sequences defined by the inventors. These are annotated as follows:

- Xle7-2EB gene
- 25 Xle4-7B gene
- Xle6-14H gene
- Satellite tandem repeat monomer
- HindIII satellite repeat

Satellite DNA inverted direct repeat

Tandem repeated DNA

Highly repetitive DNA

They are not annotated as centromere repeats in GenBank. A completed list of these
5 sequences are shown in Table 3.

**Table 3. GenBank entries match the *Brassica oleracea* centromere sequences
defined by the inventors**

GenBank Accession No.	Annotation	Repeat	Position	No. of base pairs	% Identity
X68786	B. juncea Xle7- 2EB gene	Complete	472-651	180	97
X68786	B. juncea Xle7- 2EB gene	Complete	763-942	180	94
X68786	B. juncea Xle7- 2EB gene	Partial	648-761	115	96
X12736	B. campestris DNA for satellite tandem repeat monomer (consensus sequence)	Complete	181-2	180	97
X07519	Wild cabbage satellite DNA	Complete	179-1	179	97
X61583	B. napus Canrep highly repetitive DNA	Complete	2-173	176	98
X68783	B. juncea repetitive DNA sequence canrep subfamily A	Partial	2-173	172	97
X68784	B. juncea Xle4-7B gene	Complete	983- 1162	180	95
X68784	B. juncea Xle4-7B gene	Partial	815-986	172	94
AJ228348	B. carinata DNA, HindIII satellite repeat (clone pBcar3)	Partial	2-173	172	96
M31438	B. oleracea satellite DNA inverted direct repeat	Partial	176-1	176	94
X16589	B. nigra tandem repeat DNA (clone	Partial	177-1	177	94

	BN1G 9, BN1G 23, BG1G 14)				
M31434	B. oleracea satellite DNA inverted direct repeat	Partial	176-8	169	95
M31437	B. oleracea satellite DNA inverted direct repeat	Partial	175-1	175	94
M30963	B. juncea tandemly repeated DNA	Complete	181-2	180	93
M31435	B. oleracea satellite DNA inverted direct repeat	Partial	174-8	169	94
M31436	B. oleracea satellite DNA inverted direct repeat	Partial	175-1	177	94
X15291	B. juncea satellite DNA	Partial	1-161	161	95
M31439	B. oleracea satellite DNA inverted direct repeat	Partial	176-1	177	90
Z22947	B. campestris satellite DNA	Partial	181-347	170	90
Z22947	B. campestris satellite DNA	Partial	2-179	178	89
M30962	B. campestris tandemly repeated DNA	Complete	181-2	180	87
X68785	B. juncea Xle6-14H gene	Complete	580-758	180	92
X68785	B. juncea Xle6-14H gene	Partial	404-568	165	90
AJ228347	B. carinata DNA, HindIII satellite repeat (clone pBcar5)	Partial	177-2	176	90

EXAMPLE 3

5

Glycine max centromere repeat sequences

We purified repetitive sequences from soybean (*Glycine max*, variety Williams 82), and set forth herein two centromere repeats, termed ChrGm1 and ChrGm2. We determined the consensus of each repeat as shown in Example 6.

5 The consensus sequence for ChrGm1 is shown in FIG. 2A (SEQ ID NO:5). This consensus was assembled from DNA sequences collected by the inventors. Seven of these sequences completely spanned the repeat, and twenty-five others partially covered the repeat. It is 92 ± 0.79 base pairs in length, and A and T comprise of 63% of the consensus.

10

 The consensus sequence for ChrGm2 is shown in FIG. 2B (SEQ ID NO:6). This consensus was assembled from DNA sequences collected by the inventors. Ten of these sequences completely spanned the repeat, and eleven others partially covered the repeat. It is 91 ± 0.48 base pairs in length, and A and T comprise of 62% of the
15 consensus.

 The two repeats (ChrGm1 and ChrGm2) were aligned to each other using the ClustalX program. Those sites which differ significantly from each other (chi-squared, $P < 0.05$) are highlighted in FIG. 2C.

20

 The GenBank nt database (<http://ftp.ncbi.nlm.nih.gov/blast/db/>, March 29 version, downloaded on 04/07/2002) and the plant satellite DNA database (<http://w31amc.umbr.cas.cz/PlantSat/> downloaded on 4/14/2002) were compared to the inventors' consensus sequences using the blastn program and an Expect value
25 threshold of -3. Consensus sequences were built using all inventors' and GenBank sequences that matched with an Expect (E) value of less than -25.

 The revised consensus sequence for ChrGm1 is shown in FIG. 2D (SEQ ID NO:7). This consensus was assembled from thirty-two DNA sequences collected by
30 the inventors and one matching sequence from GenBank (accession number Z26334). Eight of these sequences completely spanned the repeat, and twenty-five others partially covered the repeat. It is 92 ± 0.74 base pairs in length, and A and T

comprise of 56% of the consensus.

The revised consensus sequence for ChrGm2 is shown in FIG. 2E (SEQ ID NO:8). This consensus was assembled from twenty-one DNA sequences collected by the inventors and three matching sequences from GenBank (accession numbers AF297983, AF297984, AF297985). Ten of these sequences completely spanned the repeat, and fourteen others partially covered the repeat. It is 91 ± 0.53 base pairs in length, and A and T comprise of 61% of the consensus.

The two repeats (ChrGm1 and ChrGm2) were aligned to each other using the ClustalX program. Those sites with significant differences (chi-squared, $P < 0.05$) are highlighted in FIG. 2F.

A total of 4 GenBank entries match the *Glycine max* centromere sequences defined by the inventors. These are annotated as follows:

Satellite DNA

Tospovirus resistance protein C (Sw5-c), tospovirus resistance protein D (Sw5-d), and tospovirus resistance protein E (Sw5-e) genes

They are not annotated as centromere repeats in GenBank. A complete list of these sequences is shown in Table 4:

Table 4: GenBank entries match the *Glycine max* centromere sequences defined by the inventors

GenBank Accession No.	Annotation	Repeat	Position	No. of base pairs	% Identity
Z26334	G. max satellite DNA	Complete	92-1	92	95
AF297985	G. max clone TRS3 tandem repetitive repeat region	Partial	259-173	87	93
AF297985	G. max clone TRS3 tandem repetitive repeat region	Partial	78-3	76	94
AF297985	G. max clone	Partial	168-83	86	90

	TRS3 tandem repetitive repeat region				
AF297984	G. max clone TRS2 tandem repetitive repeat region	Partial	170-84	87	91
AF297984	G. max clone TRS2 tandem repetitive repeat region	Partial	260-175	86	88
AF297984	G. max clone TRS2 tandem repetitive repeat region	Partial	79-3	77	89
AF297983	G. max clone TRS1 tandem repetitive repeat region	Partial	77-3	75	94

EXAMPLE 4

5 *Lycopersicon esculentum* centromere repeat sequences

We purified repetitive sequences from tomato (*Lycopersicon esculentum*, variety Microtom) and set forth herein one centromere repeat. We determined the consensus of this repeat as shown in Example 6.

10

The consensus sequence of ChrLe1 is shown in FIG. 3A (SEQ ID NO:9). This consensus was assembled from forty-two DNA sequences collected by the inventors. Eighteen of these sequences completely spanned the repeat, and twenty-four others partially covered the repeat. The repeat is 181 ± 0.61 base pairs in length, and A and

15

T comprise of 50% of the consensus.

The GenBank nt database (<http://ftp.ncbi.nlm.nih.gov/blast/db/>, March 29 version, downloaded on 04/07/2002) and the plant satellite DNA database (<http://w3lamc.umbr.cas.cz/PlantSat/>, downloaded on 4/14/2002) were compared to

20

the inventors' consensus sequences using the blastn program and an Expect value

threshold value of -3. Consensus sequences were built using all inventors' and GenBank sequences matched with an Expect (E) value of less than -40.

We determined the consensus of this repeat. The repeat is 181 ± 0.61 base
5 pairs in length, and A and T comprise of 50% of the consensus.

The revised consensus sequence of ChrLe1 is shown in FIG. 3B (SEQ ID NO:10). This consensus was assembled from forty-two sequences collected by the inventors and two GenBank sequence (nt database at
10 [ftp://ftp.ncbi.nlm.nih.gov/blast/db/](http://ftp.ncbi.nlm.nih.gov/blast/db/). March 29 version, downloaded on 04/07/2002). Eighteen of these sequences completely spanned the repeat, and twenty-six others partially covered the repeat. The GenBank sequences are accession numbers X87233 and AY007367.

15 Neither of the 2 GenBank entries that match the *Lycopersicon esculentum* centromere sequences defined by the inventors are complete repeats; they match only a portion of the sequence identified by the company. These are annotated as follows:

Satellite DNA

Tandem repetitive repeat region

20 They are not annotated as centromere repeats in GenBank. A complete list of these sequences is shown in Table 5.

Table 5: GenBank entries match the *Lycopersicon esculentum* centromere sequences defined by the inventors

GenBank Accession No.	Annotation	Repeat	Position	No. of base pairs	% Identity
X87233	L. esculentum satellite DNA	Partial	163-1	161	93
AY007367	L. esculentum tospovirus resistance protein C (Sw5-c), tospovirus resistance protein D (Sw5-d), and	Partial	12003-12156	154	93

	tospovirus resistance protein E (Sw5-e) genes				
AY007367	L. esculentum tospovirus resistance protein C (Sw5-c), tospovirus resistance protein D (Sw5-d), and tospovirus resistance protein E (Sw5-e) genes	Partial	12184- 12344	161	90
AY007367	L. esculentum tospovirus resistance protein C (Sw5-c), tospovirus resistance protein D (Sw5-d), and tospovirus resistance protein E (Sw5-e) genes	Partial	12546- 12700	155	90
AY007367	L. esculentum tospovirus resistance protein C (Sw5-c), tospovirus resistance protein D (Sw5-d), and tospovirus resistance protein E (Sw5-e) genes	Partial	12365- 12526	157	89

5

EXAMPLE 5***Zea mays* centromere repeat sequences**

We purified repetitive sequences from corn (*Zea mays*, variety B73), and set forth herein one centromere repeat, termed ChrZm1. We determined the consensus of the repeat as shown in Example 5. The repeat is 180 ± 1.15 base pairs in length, and A and T comprise of 56% of the consensus.

The consensus sequence of ChrZm1 is shown in FIG. 4A (SEQ ID NO:11). This consensus was assembled from thirty-eight DNA sequences collected by the inventors. Three of these sequences completely spanned the repeat, and thirty-five
 5 others partially covered the repeat.

The GenBank nt database (<http://ftp.ncbi.nlm.nih.gov/blast/db/>, March 29 version, downloaded on 04/07/2002) and the plant satellite DNA database (<http://w3lamc.umbr.cas.cz/PlanSat/>, downloaded on 4/14/2002) were compared to the
 10 inventors' consensus sequences using the blastn program and an Expect value threshold score of -3. Consensus sequences were built using all inventors' and GenBank sequences matched with an Expect (E) value of -50.

The revised consensus sequence of ChrZm1 is shown in FIG. 4B (SEQ ID
 15 NO:12). This consensus was assembled from thirty-eight DNA sequences collected by the inventors and twenty-six matching GenBank sequences (Table 6). Twenty of these sequences completely spanned the repeat, and forty-four others partially covered the repeat. The length of the repeat is 180 ± 0.51 base pairs, and A and T comprise the consensus.

20

Table 6. GenBank sequences that match the inventors' ChrZm1 consensus

M32521	M32522	M32523	M32524	M32525	M32526
M32527	M32528	M32529	M32530	M32531	M32532
M32533	M32534	M325375	M32536	M32537	M32538
M35408	AF030934	AF030935	AF030936	AF030937	AF030938
AF030939	AF030940				

A total of 26 GenBank entries match the *Zea mays* centromere sequences
 25 defined by the inventors. These are annotated as follows:

180-bp knob-specific repeat region

heterochromatin repetitive DNA

They are not annotated as centromere repeats in GenBank. A complete list of these
 sequences is shown in Table 7.

30

Table 7: GenBank entries match the *Lycopersicon esculentum* centromere sequences defined by the inventors

GenBank Accession No.	Annotation	Repeat	Position	No. of base pairs	% Identity
M32522	Maize 180-bp knob-specific repeat region	Complete	1-180	180	96
M32521	Maize 180-bp knob-specific repeat region	Complete	1-180	180	96
M32533	Z. mays subsp. mexicana 180-bp knob-specific repeat region	Complete	1-180	180	96
M32525	Maize 180-bp knob-specific repeat region	Complete	1-180	180	96
M32524	Maize 180-bp knob-specific repeat region	Complete	1-180	180	96
M32523	Maize 180-bp knob-specific repeat region	Complete	1-180	180	96
M35408	Corn heterochromatin repetitive DNA	Complete	1-180	180	96
M32526	Maize 180-bp knob-specific repeat region	Complete	1-180	180	95
AF030939	Z. mays 180-bp knob-associated tandem repeat 15-T3-2	Complete	1-180	180	95
M32528	Maize 180-bp knob-specific repeat region	Complete	1-180	180	95
M32534	Z. mays subsp. mexicana 180-bp knob-specific repeat region	Complete	1-180	180	94
M32527	Maize 180-bp knob-specific repeat region	Partial	8-179	172	95
M32538	T. dactyloides (Tripsacum dactyloides, gama grass) 180-bp knob-specific	Complete	1-179	179	94

	repeat region				
M32529	Maize 180-bp knob-specific repeat region	Complete	1-180	180	93
AF030938	Z. mays 180-bp knob-associated tandem repeat 15-T3-1	Partial	4-180	177	93
M32532	Maize 180-bp knob-specific repeat region	Complete	1-180	180	93
AF030937	Z. mays 180-bp knob-associated tandem repeat 1-T7-2	Complete	1-180	180	92
AF030940	Z. mays 180-bp knob-associated tandem repeat 15-T7-1	Complete	1-180	180	92
AF030936	Z. mays 180-bp knob-associated tandem repeat 1-T7-1	Partial	10-180	172	93
M32537	T. dactyloides 180-bp knob-specific repeat region	Complete	1-180	180	92
M32530	Maize 180-bp knob-specific repeat region	Complete	1-180	180	92
M32531	Maize 180-bp knob-specific repeat region	Complete	1-179, introduced one gap	180	91
AF030935	Z. mays 180-bp knob-associated tandem repeat 1-T3-2	Partial	1-175	175	90
AF030934	Z. mays 180-bp knob-associated tandem repeat 1-T3-1	Partial	47-201	155	92
M32536	T. dactyloides 180-bp knob-specific repeat region	Complete	1-180	180	94
M32535	T. dactyloides 180-bp knob-specific repeat region	Complete	1-177, 2% gaps	177	91

Six GenBank sequences of *Zea mays* centromeric repeat CentC were collected (Table 13) and assigned the identifier ChrZm2. The consensus of the repeat was determined as shown in Example 6. The repeat is 158 ± 1.6 base pairs in length. A and T comprises of 53% of the bases. All 6 sequences are of unit length.

5

The consensus sequence of ChrZm2 (SEQ ID NO:13) is shown in FIG. 4C.

Table 8. GenBank sequences of *Zea mays* centromeric repeat ChrZm2

AF078918	AF078919	AF078920
AF0789121	AF078922	AF078923

10

EXAMPLE 6

15

Determining consensus sequences

Sequences were first aligned and edited in Vector NTI suite7 (InforMax, 7600 Wisconsin Ave., Suite 1100, Bethesda, MD 20814) and exported as a fasta file. A perl program, consensus.pl, was written and used to determine the consensus for each position within the repeats based on the following rules:

20

The most common base is designated as the consensus if it occurs three times more frequently than the second most common base.

25

If the occurrence of the most common base is not three times more frequent than the second most common base, but the combined frequency of the two most common bases is three times that of the third most common base, and the frequency of the second most common base is greater than the frequency of the third most common base, then the second and first bases are together considered as a consensus polymorphism, and designated using the IUPAC codes (M=A or C, R= A or G, W= A or T, S=C or G, Y= C or T, k= G or T, V= A or C or G, H=A or C or T, D=A or G or T, B=C or G or T, N= G or T or C or A).

30

If the combined frequency of the two most common bases is not three times greater than that of the third most common base, but the combined frequency of the three most common bases is three times that of the fourth most common base, and the third most common base is more common than the fourth most common base, and the
5 frequency of occurrence of the fourth base is less than or equal to 22%, the consensus is assigned according to the IUPAC ambiguity codes for the three most common bases. If the four bases occur approximately equally (23-27%), the consensus is assigned as N.

10

EXAMPLE 7

Constructing BAC Vectors for Testing Centromere Function

15 A BAC clone may be retrofitted with one or more plant telomeres and selectable markers together with the DNA elements necessary for *Agrobacterium* transformation (FIG. 9). This method will provide a means to deliver any BAC clone into plant cells and to test it for centromere function.

20 The method works in the following way. The conversion vector contains a retrofitting cassette. The retrofitting cassette is flanked by Tn10, Tn5, Tn7, Mu or other transposable elements and contains an origin of replication and a selectable marker for *Agrobacterium*, a plant telomere array followed by T-DNA right and left borders followed by a second plant telomere array and a plant selectable marker (FIG.
25 9). The conversion vector is transformed into an *E. coli* strain carrying the target BAC. The transposable elements flanking the retrofitting cassette then mediate transposition of the cassette randomly into the BAC clone. The retrofitted BAC clone can now be transformed into an appropriate strain of *Agrobacterium* and then into plant cells where it can be tested for high fidelity meiotic and mitotic transmission
30 which would indicate that the clone contained a complete functional plant centromere.

EXAMPLE 8

Sequence Analysis of *Arabidopsis* Centromeres

A. Abundance of genes in the centromeric regions

5 Expressed genes are located within 1 kb of essential centromere sequences in *S. cerevisiae*, and multiple copies of tRNA genes reside within an 80 kb fragment necessary for centromere function in *S. pombe* (Kuhn *et al.*, 1991). In contrast, genes are thought to be relatively rare in the centromeres of higher eukaryotes, though there are notable exceptions. The *Drosophila light*, *concertina*, *responder*, and *rolled* loci
10 all map to the centromeric region of chromosome 2, and translocations that remove *light* from its native heterochromatic context inhibit gene expression. In contrast, many *Drosophila* and human genes that normally reside in euchromatin become inactive when they are inserted near a centromere. Thus, genes that reside near centromeres likely have special control elements that allow expression (Karpen, 1994;
15 Lohe and Hilliker, 1995). The sequences of *Arabidopsis CEN2* and *CEN4*, provided herein, provide a powerful resource for understanding how gene density and expression correlate with centromere position and associated chromatin.

Annotation of chromosome II and IV ([http://](http://www.ncbi.nlm.nih.gov/Entrez/nucleotide.html)
20 www.ncbi.nlm.nih.gov/Entrez/nucleotide.html) identified many genes within and adjacent to *CEN2* and *CEN4* (FIG. 8, FIGs. 11A-11T). The density of predicted genes on *Arabidopsis* chromosome arms averages 25 per 100 kb, and in the repeat-rich regions flanking *CEN2* and *CEN4* this decreases to 9 and 7 genes per 100 kb, respectively (Bevan *et al.*, 1999). Many predicted genes also reside within the
25 recombination-deficient, genetically-defined centromeres. Within *CEN2*, there were 5 predicted genes per 100 kb; while *CEN4* was strikingly different, with 12 genes per 100 kb.

There was strong evidence that several of the predicted centromeric genes are
30 transcribed. The phosphoenolpyruvate gene (*CUE1*) defines one *CEN5* border; mutations in this gene cause defects in light-regulated gene expression (Li *et al.*, 1995). Within the sequenced portions of *CEN2* and *CEN4*, 17% (27/160) of

the predicted genes shared >95% identity with cloned cDNAs (ESTs), with three-fold more matches in *CEN4* than in *CEN2* (<http://www.tigr.org/tdb/at/agad/>). Twenty-four of these genes have multiple exons, and four correspond to single-copy genes with known functions. A list of the predicted genes identified is given in Table 9, below. A list of additional genes encoded within the boundaries of *CEN4* are listed in Table 10. The identification of these genes is significant in that the genes may themselves contain unique regulatory elements or may reside in genomic locations flanking unique control or regulatory elements involved in centromere function or gene expression. In particular, the current inventors contemplate use of these genes, or DNA sequences 0 to 5 kb upstream or downstream of these sequences, for insertion into a gene of choice in a minichromosome. It is expected that such elements could potentially yield beneficial regulatory controls of the expression of these genes, even when in the unique environment of a centromere.

To investigate whether the remaining 23 genes were uniquely encoded at the centromere, a search was made in the database of annotated genomic *Arabidopsis* sequences. With the exception of two genes, no homologs with >95% identity were found elsewhere in the 80% of the genome that has been sequenced. The number of independent cDNA clones that correspond to a single-copy gene provides an estimate of the level of gene expression. On chromosome II, predicted genes with high quality matches to the cDNA database (> 95% identity) match an average of four independent cDNA clones (range 1-78). Within *CEN2* and *CEN4*, 11/27 genes exceed this average (Table 9). Finally, genes encoded at *CEN2* and *CEN4* are not members of a single gene family, nor do they correspond to genes predicted to play a role in centromere functions, but instead have diverse roles.

Many genes in the *Arabidopsis* centromeric regions are nonfunctional due to early stop codons or disrupted open reading frames, but few pseudogenes were found on the chromosome arms. Though a large fraction of these pseudogenes have homology to mobile elements, many correspond to genes that are typically not mobile (FIGs. 11I-J and FIGs. 11S-T). Within the genetically-defined centromeres there were 1.0 (*CEN2*) and 0.7 (*CEN4*) of these nonmobile pseudogenes per 100 kb; the

repeat-rich regions bordering the centromeres have 1.5 and 0.9 per 100 kb respectively. The distributions of pseudogenes and transposable elements are overlapping, indicating that DNA insertions in these regions contributed to gene disruptions.

5

Table 9: Predicted genes within *CEN2* and *CEN4* that correspond to the cDNA database.

Putative function	GenBank protein accession	# of EST matches*
<i>CEN2</i>		
Unknown	AAC69124	1
SH3 domain protein	AAD15528	5
Unknown	AAD15529	1
unknown†	AAD37022	1
RNA helicase‡	AAC26676	2
40S ribosomal protein S16	AAD22696	9
<i>CEN4</i>		
Unknown	AAD36948	1
Unknown	AAD36947	4
leucyl tRNA synthetase	AAD36946	4
aspartic protease	AAD29758	6
Peroxisomal membrane protein (<i>PPM2</i>) §	AAD29759	5
5'-adenylylsulfate reductase §	AAD29775	14
symbiosis-related protein	AAD29776	3
ATP synthase gamma chain 1 (<i>APC1</i>) §	AAD48955	3

protein kinase and EF hand	AAD03453	3
ABC transporter	AAD03441	1
Transcriptional regulator	AAD03444	14
Unknown	AAD03446	12
human PCF11p homolog	AAD03447	6
NSF protein	AAD17345	2
1,3-beta-glucan synthase	AAD48971	2
pyridine nucleotide-disulphide oxidoreductase	AAD48975	4
Polyubiquitin (<i>UBQ11</i>) §	AAD48980	72
wound induced protein	AAD48981	6
short chain dehydrogenase/reductase	AAD48959	7
SL15†	AAD48939	2
WD40-repeat protein	AAD48948	2

* Independent cDNAs with >95% identity, † related gene present in non-centromeric DNA, ‡ potentially associated with a mobile DNA element, § characterized gene (B. Tugal, 1999; J.F. Gutierrez-Marcos, 1996; N. Inohara, 1991; J. Callis, 1995).

5 Table 10: List of additional genes encoded within the boundaries of CEN4.

Putative Function	GenBank accession	Nucleotide Position
3'(2'),5'-Bisphosphate Nucleotidase	AC012392	71298 -73681
Transcriptional regulator	AC012392	80611 -81844
Equilibrative nucleoside transporter 1	AC012392	88570 -90739
Equilibrative nucleoside transporter 1	AC012392	94940 -96878

Equilibrative nucleoside transporter 1	AC012392	98929 -101019
Equilibrative nucleoside transporter 1	AC012392	113069 -115262
unknown	AC012392	122486 -124729
4-coumarate--CoA ligase	AC012392	126505 -128601
ethylene responsive protein	AC012392	130044 -131421
Oxygen-evolving enhancer protein precursor	AC012392	134147 -135224
Kinesin	AC012392	137630 -141536
receptor-like protein kinase	AC012392	141847 -144363
LpxD-like protein	AC012392	144921 -146953
hypersensitivity induced protein	AC012392	147158 -147838
ubiquitin	AC012392	149057 -149677
unknown	AC012392	150254 -151072
ubiquitin-like protein	AC012392	153514 -154470
ubiquitin-like protein	AC012392	155734 -156513
ubiquitin-like protein	AC012392	156993 -157382
unknown	AC012392	159635 -165559
unknown	AC012392	166279 -166920
unknown	AC012392	167724 -170212
ubiquitin-like protein	AC012392	176819 -178066
polyubiquitin (<i>UBQ10</i>) §	AC012392	180613 -182007
phosphatidylinositol-3,4,5-triphosphate binding protein	AC012477	89384 -91291
Mitochondrial ATPase	AC012477	94302 -94677
RING-H2 finger protein	AC012477	95522 -96142

unknown	AC012477	104747 -105196
Mitochondrial ATPase	AC012477	105758 -106595
ferredoxin--NADP+ reductase	AC012477	107451 -109095
unknown	AC012477	109868 -110620
U3 snoRNP-associated protein	AC012477	111841 -114133
UV-damaged DNA binding factor	AC012477	114900 -121275
Glucan endo-1,3-Beta-Glucosidase precursor	AC012477	122194 -122895
D123 -like protein	AC012477	125886 -126887
Adrenodoxin Precursor	AC012477	127660 -129246
N7 like-protein	AC012477	129718 -131012
N7 like-protein	AC012477	131868 -133963
N7 like-protein	AC012477	134215 -136569
N7 like-protein	AC012477	139656 -140864

§ characterized gene (J. Callis, 1995).

B. Conservation of centromeric DNA

To investigate the conservation of *CEN2* and *CEN4* sequences, PCR primer
5 pairs were designed that correspond to unique regions in the Columbia sequence and
used to survey the centromeric regions of Landsberg and Columbia at ~20 kb intervals
(FIGs. 13A, B). The primers used for the analysis are listed in FIGs. 14A, B.
Amplification products of the appropriate length were obtained in both ecotypes for
most primer pairs (85%), indicating that the amplified regions were highly similar. In
10 the remaining cases, primer pairs amplified Columbia, but not Landsberg DNA, even
at very low stringencies. In these regions, additional primers were designed to
determine the extent of nonhomology. In addition to a large insertion of mitochondrial
DNA in *CEN2*, two other non-conserved regions were identified (FIGs. 13A, B).
Because this DNA is absent from Landsberg centromeres, it is unlikely to be required
15 for centromere function; consequently, the relevant portion of the centromeric

sequence is reduced to 577 kb (*CEN2*) and 1250 kb (*CEN4*). The high degree of sequence conservation between Landsberg and Columbia centromeres indicated that the inhibition of recombination frequencies was not due to large regions of nonhomology, but instead was a property of the centromeres themselves.

5

C. Sequence similarity between *CEN2* and *CEN4*

In order to discern centromere function, a search was conducted for novel sequence motifs shared between *CEN2* and *CEN4*, excluding from the comparison retroelements, transposons, characterized centromeric repeats, and coding sequences resembling mobile genes. After masking simple repetitive sequences, including homopolymer tracts and microsatellites, contigs of unique sequence measuring 417 kb and 851 kb for *CEN2* and *CEN4*, respectively, were compared with BLAST (<http://blast.wustl.edu>).

15 The comparison showed that the complex DNA within the centromere regions was not homologous over the entire sequence length. However, 16 DNA segments in *CEN2* matched 11 regions in *CEN4* with >60% identity (FIG. 15). The sequences were grouped into families of related sequences, and were designated AtCCS1-7 (*Arabidopsis thaliana* centromere conserved sequences 1-7). These sequences were not previously known to be repeated in the *Arabidopsis* genome. The sequences comprised a total of 17 kb (4%) of *CEN2* DNA, had an average length of 1017 bp, and had an A + T content of 65%. Based on similarity, the matching sequences were sorted into groups, including two families containing 8 sequences each, 3 sequences from a small family encoding a putative open reading frame, and 4 sequences found once within the centromeres, one of which corresponds to predicted *CEN2* and *CEN4* proteins with similarity throughout their exons and introns (FIG. 15).

Searches of the *Arabidopsis* genomic sequence database demonstrated that AtCCS1 - AtCCS5 were moderately repeated sequences that appear in centromeric and pericentromeric regions. The remaining sequences were present only in the genetically-defined centromeres. Similar comparisons of all 16 *S. cerevisiae* centromeres defined a consensus consisting of a conserved 8 bp CDEI motif, an

AT-rich 85 bp CDEII element, and a 26 bp CDEII region with 7 highly conserved nucleotides (Fleig *et al.*, 1995). In contrast, surveys of the three *S. pombe* centromeres revealed conservation of overall centromere structure, but no universally conserved motifs (Clark, 1998).

5

EXAMPLE 9

Construction of Plant Minichromosomes

Minichromosomes are constructed by combining the previously isolated
10 essential chromosomal elements. Exemplary minichromosome vectors include those designed to be "shuttle vectors"; *i.e.*, they can be maintained in a convenient host (such as *E. coli*, *Agrobacterium* or yeast) as well as plant cells.

A. General Techniques for Minichromosome Construction

15 A minichromosome can be maintained in *E. coli* or other bacterial cells as a circular molecule by placing a removable stuffer fragment between the telomeric sequence blocks. The stuffer fragment is a dispensable DNA sequence, bordered by unique restriction sites, which can be removed by restriction digestion of the circular DNAs to create linear molecules with telomeric ends. The linear minichromosome can
20 then be isolated by, for example, gel electrophoresis. In addition to the stuffer fragment and the plant telomeres, the minichromosome contains a replication origin and selectable marker that can function in plants to allow the circular molecules to be maintained in bacterial cells. The minichromosomes also include a plant selectable marker, a plant centromere, and a plant ARS to allow replication and maintenance of
25 the DNA molecules in plant cells. Finally, the minichromosome includes several unique restriction sites where additional DNA sequence inserts can be cloned. The most expeditious method of physically constructing such a minichromosome, *i.e.*, ligating the various essential elements together for example, will be apparent to those of ordinary skill in this art.

30

A number of minichromosome vectors have been designed by the current inventors and are disclosed herein for the purpose of illustration (FIGs. 7A-7H). These

vectors are not limiting however, as it will be apparent to those of skill in the art that many changes and alterations may be made and still obtain a functional vector.

B. Modified Technique for Minichromosome Construction

5 A two step method was developed for construction of minichromosomes, which allows adding essential elements to BAC clones containing centromeric DNA. These procedures can take place *in vivo*, eliminating problems of chromosome breakage that often happen in the test tube. The details and advantages of the techniques are as follows:

10

1.) One plasmid can be created that contains markers, origins and border sequences for *Agrobacterium* transfer, markers for selection and screening in plants, plant telomeres, and a loxP site or other site useful for site-specific recombination *in vivo* or *in vitro*. The second plasmid can be an existing BAC
15 clone, isolated from the available genomic libraries (FIG. 10A).

20

2.) The two plasmids are mixed, either within a single *E. coli* cell, or in a test tube, and the site-specific recombinase *cre* is introduced. This will cause the two plasmids to fuse at the loxP sites (FIG. 10B).

25

3.) If deemed necessary, useful restriction sites (AseI/PacI or Not I) are included to remove excess material. (for example other selectable markers or replication origins)

4.) Variations include vectors with or without a Kan^R gene (FIGs. 10B, 10C), with or without a LAT52 GUS gene, with a LAT52 GFP gene, and with a GUS gene under the control of other plant promoters. (FIGs. 10C, 10D and 10E).

C. Method for Preparation of Stable Non-Integrated Minichromosomes

A technique has been developed to ensure that minichromosomes do not integrate into the host genome (FIG. 10F). In particular, minichromosomes must be maintained as distinct elements separate from the host chromosomes. In one method for ensuring that the introduced minichromosome does not integrate, the inventors envision a variety that would encode a lethal plant gene (such as diphtheria toxin or any other gene product that, when expressed, causes lethality in plants). This gene could be located between the right *Agrobacterium* border and the telomere.

Minichromosomes that enter a plant nucleus and integrate into a host chromosome would result in lethality. However, if the minichromosome remains separate, and further, if the ends of this construct are degraded up to the telomeres, then the lethal gene would be removed and the cells would survive.

It should be understood that various changes and modifications to the presently preferred embodiments described herein will be apparent to those skilled in the art. Such changes and modifications can be made without departing from the spirit and scope of the present invention and without diminishing its intended advantages. It is therefore intended that such changes and modifications be covered by the appended claims.

REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated
5 herein by reference.

- Abdullah *et al.*, *Biotechnology*, 4:1087, 1986.
- Abel *et al.*, *Science*, 232:738-743, 1986.
- Alfenito *et al.*, "Molecular characterization of a maize B chromosome centric
10 sequence," *Genetics*, 135:589-597, 1993.
- Alfenito and Birchler, "Molecular characterization of a maize B chromosome centric
sequence," *Genetics* 135: 589-597, 1993.
- Ananiev *et al.*, "A knob-associated tandem repeat in maize capable of forming fold-
back DNA segments: are chromosome knobs megatransposons?" *Proc. Natl.*
15 *Acad. Sci. U.S.A.* 95 (18), 10785-10790, 1998.
- Ananiev *et al.*, "Chromosome-specific molecular organization of maize (*Zea mays* L.)
centromeric regions," *Proc. Natl. Acad. Sci. U.S.A.* 95 (22), 13073-13078,
1998.
- Ananiev *et al.*, "Complex structure of knob DNA on maize chromosome 9.
20 Retrotransposon invasion into heterochromatin," *Genetics* 149 (4), 2025-2037,
1998
- Araki *et al.*, "Site-specific recombinase, R, encoded by yeast plasmid pSR1," *J. Mol.*
Biol. 225:25-37, 1992.
- Armstrong *et al.*, "Physical mapping of DNA repetitive sequences to mitotic and
25 meiotic chromosomes of *Brassica oleracea* var. *alboglabra* by fluorescence in
situ hybridization," *Heredity* 81: 666-673, 1998.
- Barkai-Golan *et al.*, *Arch. Microbiol.*, 116:119-124, 1978.
- Baum *et al.*, "The centromeric K-type repeat and the central core are together
sufficient to establish a functional *Schizosaccharomyces pombe* centromere,"
30 *Mol. Bio. Cell.*, 5:747-761, 1994.
- Bell *et al.*, "Assignment of 30 microsatellite loci to the linkage map of *Arabidopsis*,"
Genomics, 19:137-144, 1994.

- Bernal-Lugo and Leopold, *Plant Physiol.*, 98:1207-1210, 1992.
- Berzal-Herranz *et al.*, *Genes and Devel.*, 6:129-134, 1992.
- Bevan *et al.*, *Nucleic Acids Research*, 11(2):369-385, 1983.
- Bevan *et al.*, *BioEssays* 21:110, 1999.
- 5 Blackman *et al.*, *Plant Physiol.*, 100:225-230, 1992.
- Bloom, "The centromere frontier: Kinetochore components, microtubule-based motility, and the CEN-value paradox," *Cell*, 73:621-624, 1993.
- Bol *et al.*, *Annu. Rev. Phytopath.*, 28:113-138, 1990.
- Bowler *et al.*, *Ann Rev. Plant Physiol.*, 43:83-116, 1992.
- 10 Brandes *et al.*, *Chrom. Res.*, 5:238, 1997.
- Branson and Guss, *Proceedings North Central Branch Entomological Society of America*, 27:91-95, 1972.
- Brisson *et al.*, *Nature*, 310:511, 1984.
- Broach *et al.*, *Gene*, 8:121-133, 1979.
- 15 Broekaert *et al.*, *Science*, 245:1100-1102, 1989.
- Burke *et al.*, *Science*, 236:806-812, 1987.
- Bytebier *et al.*, *Proc. Natl Acad. Sci. USA*, 84:5345, 1987.
- Callis *et al.*, *Genes and Development*, 1:1183, 1987.
- Cambareri *et al.*, *Mol. Cell. Biol.*, 18:5465, 1998.
- 20 Campbell (ed.), *In: Avermectin and Abamectin*, 1989.
- Campbell, "Monoclonal Antibody Technology, Laboratory Techniques in Biochemistry and Molecular Biology," Vol. 13, Burden and Von Knippenberg, Eds. pp. 75-83, Elsevier, Amsterdam, 1984.
- Capecchi, "High efficiency transformation by direct microinjection of DNA into cultured mammalian cells," *Cell* 22(2):479-488, 1980.
- 25 Carbon *et al.*, *In: Recombinant Molecules: Impact on Science and Society* (Raven Press), 335-378, 1977.
- Carbon *et al.*, "Centromere structure and function in budding and fission yeasts," *New Biologist*, 2:10-19, 1990.
- 30 Carpenter *et al.*, "The control of the distribution of meiotic exchange in *Drosophilla melanogaster*," *Genetics*, 101:81-90, 1982.

- Cech *et al.*, "In vitro splicing of the ribosomal RNA precursor of Tetrahymena: involvement of a guanosine nucleotide in the excision of the intervening sequence," *Cell*, 27:487-496, 1981.
- Chandler *et al.*, *The Plant Cell*, 1:1175-1183, 1989.
- 5 Chang *et al.*, "Restriction fragment length polymorphism linkage map for *Arabidopsis thaliana*," *Proc. Natl Acad. Sci., USA*, 85:6856-6860, 1988.
- Charlesworth *et al.*, *Nature*, 371:215, 1994.
- Charlesworth, C.H. Langley, W. Stephan, 112:947, 1986
- Chepko, *Cell*, 37:1053, 1984.
- 10 Choi *et al.*, *Plant Mol Biol Rep*, 13:124-29, 1995.
- Choo, K.H.A. *Genome Res.* 8:81, 1998.
- Chowrira *et al.*, "In vitro and in vivo comparison of hammerhead, hairpin, and hepatitis delta virus self-processing ribozyme cassettes," *J. Biol. Chem.*, 269:25856-25864, 1994.
- 15 Chu *et al.*, "Separation of large DNA molecules by contour-clamped homogeneous electric fields" *Science*, 234, 1582-1585, 1986.
- Chye *et al.*, *Plant Mol. Biol.*, 35:893, 1997.
- Clapp, "Somatic gene therapy into hematopoietic cells. Current status and future implications," *Clin. Perinatol.* 20(1):155-168, 1993.
- 20 Clark, L. *Curr. Op. Gen. & Dev.*, 8:212, 1998
- Clarke *et al.*, "Isolation of a yeast centromere and construction of functional small circular chromosomes," *Nature*, 287:504-509, 1980.
- Cohen *et al.*, *Proc. Nat'l Acad. Sci. USA*, 70:3240, 1973.
- Conkling *et al.*, *Plant Physiol.*, 93:1203-1211, 1990.
- 25 Copenhaver and Pikaard, "RFLP and physical mapping with an rDNA-specific endonuclease reveals that nucleolus organizer regions of *Arabidopsis thaliana* adjoin the telomeres on chromosomes 2 and 4," *Plant J.*, 9:259-276, 1996.
- Copenhaver *et al.*, "Use of RFLPs larger than 100 kbp to map position and internal organization of the nucleolus organizer region on chromosome 2 in
- 30 *Arabidopsis thaliana*," *Plant J.* 7, 273-286, 1995.
- Copenhaver *et al.*, *Proc. Natl. Acad. Sci.* 95:247, 1998.
- Copenhaver *et al.*, *Science*. 286:2468-2474, 1999.

- Copenhaver and Preuss, *Plant Biology*, 2:104-108, 1999.
- Coxson *et al.*, *Biotropica*, 24:121-133, 1992.
- Creusot *et al.*, *Plant Journal*, 8:763-70, 1995
- Cristou *et al.*, *Plant Physiol*, 87:671-674, 1988.
- 5 Cuozzo *et al.*, *Bio/Technology*, 6:549-553, 1988.
- Curiel *et al.*, "Adenovirus enhancement of transferrin-polylysine-mediated gene delivery," *Proc. Natl Acad. Sci. USA* 88(19):8850-8854, 1991.
- Curiel *et al.*, high-efficiency gene transfer mediated by adenovirus coupled to DNA-polylysine complexes," *Hum. Gen. Ther.* 3(2):147-154, 1992.
- 10 Cutler *et al.*, *J. Plant Physiol.*, 135:351-354, 1989.
- Czapla and Lang, *J. Econ. Entomol.*, 83:2480-2485, 1990.
- Davies *et al.*, *Plant Physiol.*, 93:588-595, 1990.
- Dellaporta *et al.*, In: *Chromosome Structure and Function: Impact of New Concepts, 18th Stadler Genetics Symposium*, 11:263-282, 1988.
- 15 Dennis and Peacock, "Knob heterochromatin homology in maize and its relatives," *J. Mol. Evol.* 20, 341-350, 1984.
- Depicker *et al.*, *Plant Cell Reports*, 7:63-66, 1988.
- DiLaurenzio *et al.*, *Cell*, 86:423-33, 1996
- Dillon *et al.*, *Recombinant DNA Methodology*, 1985.
- 20 Donahue *et al.*, "The nucleotide sequence of the HIS4 region of yeast," *Gene* Apr;18(1):47-59, 1982.
- Dure *et al.*, *Plant Molecular Biology*, 12:475-486, 1989.
- Earnshaw *et al.*, "Proteins of the inner and outer centromere of mitotic chromosomes," *Genome*, 31:541-552, 1989.
- 25 Earnshaw, "When is a centromere not a kinetochore?," *J. Cell Sci.*, 99:1-4, 1991.
- Ebert *et al.*, 84:5745-5749, *Proc. Nat'l Acad. Sci. USA*, 1987
- Ecker, JR, *Genomics*, 19:137-144
- Ecker, *Methods*, 1:186-94, 1990.
- Eglitis *et al.*, "Retroviral vectors for introduction of genes into mammalian cells,"
- 30 *Biotechniques* 6(7):608-614, 1988.
- Eglitis *et al.*, "Retroviral-mediated gene transfer into hemopoietic cells," *Adv. Exp. Med. Biol.* 241:19-27, 1988.

- Enomoto *et al.*, "Mapping of the pin locus coding for a site-specific recombinase that causes flagellar-phase variation in *Escherichia coli* K-12," *J. Bacteriol.*, 156:663-668, 1983.
- Erdmann *et al.*, *J. Gen. Microbiology*, 138:363-368, 1992.
- 5 Ferrin *et al.*, "Selective cleavage of human DNA: RecA-Assited Restriction Endonuclease (RARE) cleavage," *Science*, 254:1494-1497, 1991.
- Fitzpatrick, *Gen. Engineering News*, 22:7, 1993.
- Fleig, U. *et al.*, "Functional selection for the centromere DNA from yeast chromosome VIII," *Nuc. Acids. Res.* 23:922-924, 1995.
- 10 Forster and Symons, "Self-cleavage of plus and minus RNAs of a virusoid and a structural model for the active sites," *Cell*, 49:211-220, 1987.
- Fraley *et al.*, *Biotechnology*, 3:629, 1985.
- Franz *et al.*, *Plant J.*, 13:867, 1998.
- Fromm *et al.*, *Nature*, 312:791-793, 1986.
- 15 Fromm *et al.*, "Expression of genes transferred into monocot and dicot plant cells by electroporation," *Proc. Natl Acad. Sci. USA* 82(17):5824-5828, 1985.
- Fujimura *et al.*, *Plant Tissue Culture Letters*, 2:74, 1985.
- Fynan *et al.*, "DNA vaccines: protective immunizations by parenteral, mucosal, and gene gun inoculations," *Proc. Nat'l Acad. Sci. USA* 90(24):11478-11482, 1993.
- 20 Gatehouse *et al.*, *J. Sci. Food. Agric.*, 35:373-380, 1984.
- Gefter *et al.*, *Somatic Cell Genet.* 3:231-236, 1977.
- Gerlach *et al.*, "Construction of a plant disease resistance gene from the satellite RNA of tobacco rinspot virus," *Nature (London)*, 328:802-805, 1987.
- Goding, "Monoclonal Antibodies: Principles and Practice," pp. 60-74. 2nd Edition,
- 25 Academic Press, Orlando, FL, 1986.
- Golic and Lindquist, "The FLP recombinase of yeast catalyses site-specific recombination in the *Drosophila* genome," *Cell*, 59:499-509, 1989.
- Goring *et al.*, *Proc. Natl. Acad. Sci. USA*, 88:1770-1774, 1991.
- Graham *et al.*, "Transformation of rat cells by DNA of human adenovirus 5," *Virology*
- 30 54(2):536-539, 1973.
- Grellet *et al.*, "Organization and evolution of a higher plant alphoid-like satellite DNA sequence," *J. Mol. Biol.* 187:495-507, 1986.

- Grill and Somerville, *Mol Gen Genet*, 226:484-90, 1991
- Guerrero *et al.*, *Plant Molecular Biology*, 15:11-26, 1990.
- Gupta *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:1629-1633, 1993.
- Gutierrez-Marcos *et al.*, *Proc. Natl. Acad. Sci.*, USA, 93:13377, 1996.
- 5 Haaf *et al.*, "Integration of human α -satellite DNA into simian chromosomes: centromere protein binding and disruption of normal chromosome segregation," *Cell*, 70:681-696, 1992.
- Hadlaczky *et al.*, "Centromere formation in mouse cells cotransformed with human DNA and a dominant marker gene," *Proc. Natl Acad. Sci. USA*, 88:8106-8110,
- 10 1991.
- Hamilton *et al.*, "Stable transfer of intact high molecular weight DNA into plant chromosomes," *Proc Natl Acad Sci U S A* 93(18):9975-9, 1996
- Hamilton, "A binary-BAC system for plant transformation with high-molecular-weight DNA," *Gene*, 4;200(1-2):107-16, 1997.
- 15 Hammock *et al.*, *Nature*, 344:458-461, 1990.
- Harrison and Heslop-Harrison, "Centromeric repetitive DNA sequences in the genus Brassica," *Theor. Appl. Genet.* 90:157-165, 1995.
- Haseloff *et al.*, *Proc. Nat'l Acad. Sci. USA* 94(6):2122-2127, 1997.
- Hauge *et al.*, *Symp Soc Exp Biol*, 45:45-56, 1991
- 20 Hegemann *et al.*, "The centromere of budding yeast," *Bioassays*, 15(7):451-460, 1993.
- Hemenway *et al.*, *The EMBO J.*, 7:1273-1280, 1988.
- Heslop-Harrison *et al.*, *Plant Cell*, 11:31, 1999.
- Hilder *et al.*, *Nature*, 330:160-163, 1987.
- Hinchee *et al.*, *Bio/technol.*, 6:915-922, 1988.
- 25 Hoess *et al.*, *Proc Natl Acad Sci*, 79:3398-402, 1982
- Hsiao *et al.*, *J. Proc. Nat'l Acad. Sci. USA*, 76:3829-3833, 1979.
- Hudspeth and Grula, *Plant Mol. Biol.*, 12:579-589, 1989.
- Hwang *et al.*, "Identification and map position of YAC clones comprising one-third of the *Arabidopsis* genome," *The Plant Journal*, 1:367-374, 1991.
- 30 Ikeda *et al.*, *J. Bacteriol.*, 169:5615-5621, 1987.
- Ikuta *et al.*, *Bio/technol.*, 8:241-242, 1990.
- Inohara *et al.*, *J. Biol. Chem.*, 266, 7333, 1991.

- Johnston *et al.*, "Gene gun transfection of animal cells and genetic immunization," *Methods Cell. Biol.* 43(A):353-365, 1994.
- Jones, *Embo J.*, 4:2411-2418, 1985.
- Jones, *Mol Gen. Genet.*, 207:478, 1987.
- 5 Jorgensen *et al.*, *Mol. Gen. Genet.*, 207:471, 1987.
- Jouanin *et al.*, *Mol Gene Genet*, 201:370-4, 1985
- Joyce, "RNA evolution and the origins of life," *Nature*, 338:217-244, 1989.
- Kaasen *et al.*, *J. Bacteriology*, 174:889-898, 1992.
- Karpen, *Curr. Op. Gen. & Dev.*, 4:281, 1994.
- 10 Karsten *et al.*, *Botanica Marina*, 35:11-19, 1992.
- Katz *et al.*, *J. Gen. Microbiol.*, 129:2703-2714, 1983.
- Kim and Cech, "Three dimensional model of the active site of the self-splicing rRNA precursor of Tetrahymena," *Proc. Natl. Acad. Sci. USA*, 84:8788-8792, 1987.
- Klee *et al.*, *Bio/Technology* 3:637-642, 1985.
- 15 Klein *et al.*, *Nature*, 327:70-73, 1987.
- Klein *et al.*, *Proc. Nat'l Acad. Sci. USA*, 85:8502-8505, 1988.
- Kohler *et al.*, *Eur. J. Immunol.* 6:511-519, 1976.
- Kohler *et al.*, *Nature* 256:495-497, 1975.
- Kolchinski and Gresshoff, "A major satellite DNA of soybean is a 92-base pairs
20 tandem repeat," *Theor. Appl. Genet.* 90(5): 621-626, 1995.
- Konieczny *et al.*, "A procedure for mapping *Arabidopsis* mutations using codominant ecotype-specific PCR-based markers," *The Plant Journal*, 4:403-410, 1993.
- Konieczny *et al.*, *Genetics*, 127:801, 1991.
- Koorneef *et al.*, *Genetica*, 61:41-46, 1983.
- 25 Koorneef, "Linkage map of *Arabidopsis thaliana* (2n=10)," In SJ O'Brien, ed, *Genetic Maps 1987: A compilation of linkage and restriction maps of genetically studied organisms*, 724-745, 1987.
- Koorneef, "The use of telotrisomics for centromere mapping in *Arabidopsis thaliana* (L.) Heynh, *Genetica*, 62:33-40, 1983.
- 30 Koster and Leopold, *Plant Physiol.*, 88:829-832, 1988.
- Kuby, J., *Immunology 2nd Edition*, W. H. Freeman & Company, NY, 1994
- Kuhn *et al.*, *Proc. Natl. Acad. Sci.*, 88:1306, 1991.

- Kyte *et al.*, A simple method for displaying the hydropathic character of a protein," *J. Mol. Biol.* 157(1):105-132, 1982.
- Lawton *et al.*, *Plant Mol. Biol.* 9:315-324, 1987.
- Lakshmikumaran and Ranade, "Isolation and characterization of a highly repetitive
5 DNA of *Brassica campestris*," *Plant Mol. Biol.* 14:447-448, 1990.
- Lechner *et al.*, "A 240 kd multisubunit protein complex, CBF3 is a major component of the budding yeast centromere," *Cell*, 64:717-725, 1991.
- Lee and Saier, *J. of Bacteriol.*, 153-685, 1983.
- Levings, *Science*, 250:942-947, 1990.
- 10 Lewin, *Genes II*, John Wiley & Sons, Publishers, N.Y., 1985.
- Li *et al.*, *Plant Cell*, 7:1599, 1995.
- Li *et al.*, *Proc. Natl. Acad. Sci.*, 87:4580-4584, 1990.
- Lieber and Strauss, "Selection of efficient cleavage sites in target RNAs by using a ribozyme expression library." *Mol. Cell. Biol.*, 15: 540-551, 1995.
- 15 Lin, S., Kaul, S. Rounsley, T.P. Shea, M-I. Benito, C.D. Town, C.Y. Fujii, T. Mason, C. L. Bowman, M. Barnstead, T. Feldblyum, C.R. Buell, K.A. Ketchum, C.M. Ronning, H. Koo, K. Moffat, L. Cronin, M. Shen, G. Pai, S. Van Aken, L., Umayam, L. Tallon, J. Gill, M.D. Adams, A.J. Carrera, T.H. Creasy, H.M. Goodman, C.R. Somerville, G.P. Copenhaver, D. Preuss, W.C. Nierman, O.
20 White, J.A. Eisen, S. Salzberg, C.M. Fraser, and J.C. Venter, "Sequence and Analysis of Chromosome 2 of *Arabidopsis thaliana*," *Nature* 402: 761-768, 1999.
- Liu, YG., Shirano, Y., Fukaki, H., Yanai, Y., Tasaka, M., Tabata, S., Shibata, D, *Proc. Natl Acad Sci USA* 96: 6535-40, 1999.
- 25 Lohe and Hilliker, *Curr. Op. Gen. & Dev.*, 5:746, 1995.
- Loomis *et al.*, *J. Expt. Zoology*, 252:9-15, 1989.
- Lorz *et al.*, *Mol. Gen. Genet.*, 199:178, 1985.
- Louis, EJ, "Corrected sequence for the right telomere of *Saccharomyces cerevisiae* chromosome III," *Yeast*, 10(2):271-4, 1994.
- 30 Lu *et al.*, "High efficiency retroviral mediated gene transduction into single isolated immature and replatable CD34(3+) hematopoietic stem/progenitor cells from human umbilical cord blood," *J. Exp. Med.* 178(6):2089-2096, 1993.

- Maeser and Kahmann, "The GIN recombinase of phage Mu can catalyze site-specific recombination in plant protoplasts," *Mol. Gen. Genet.*, 230:170-176, 1991.
- Mahtani, M.M. and Willard, H.F. *Genome Res.* 8:100, 1998.
- Maloy, S. R., "Experimental Techniques in Bacterial Genetics" Jones and Bartlett
- 5 Prokop, A., and Bajpai, R. K. "Recombinant DNA Technology I" *Ann. N. Y. Acad. Sci.* vol. 646, 1991.
- Maluszynaska *et al.*, "Molecular cytogenetics of the genus *Arabidopsis*: *In situ* localization of rDNA sites, chromosome numbers and diversity in centromeric heterochromatin," *Annals Botany*, 71:479-484, 1993.
- 10 Maluszynska *et al.*, "Localization of tandemly repeated DNA sequences in *Arabidopsis thaliana*," *Plant Jour.*, 1(2):159-166, 1991.
- Maniatis *et al.*, "Molecular Cloning: a Laboratory Manual," Cold Spring Harbor Laboratory, Cold Spring Harbor, NY., 1982.
- Marcotte *et al.*, *Nature*, 335:454, 1988.
- 15 Mariani *et al.*, *Nature*, 347:737-741, 1990.
- Marra *et al.*, *Nature Genet.* 22:265, 1999.
- Martinez-Zapater *et al.*, *Mol. Gen. Genet.*, 204:417-423, 1986.
- Matsuura *et al.*, *Journal of Bacteriology*, 178:3374-6. 1996
- McCabe *et al.*, *Biotechnology*, 6:923, 1988.
- 20 Michel and Westhof, "Modeling of the three-dimensional architecture of group I catalytic introns based on comparative sequence analysis," *J. Mol. Biol.*, 216:585-610, 1990.
- Mortimer *et al.*, "Genetic mapping in *Saccharomyces cerevisiae*," *Life Cycle and Inheritance, In: The Molecular Biology of the Yeast Saccharomyces*, 11-26,
- 25 1981.
- Mozo *et al.*, *Mol Gen Genet*, 258:562-70, 1998.
- Mozo *et al.*, *Nature Genet.* 22:271, 1999.
- Mundy and Chua, *The EMBO J.*, 7:2279-2286, 1988.
- Murakami *et al.*, *Mol. Gen. Genet.*, 205:42-50, 1986.
- 30 Murata *et al.*, *Plant J.*, 12:31, 1997.
- Murdock *et al.*, *Phytochemistry*, 29:85-89, 1990.
- Murray *et al.*, *Nature*, 305:189-193, 1983.

- Mysore *et al.*, "An *arabidopsis* histone H2A mutant is deficient in *agrobacterium* T-DNA integration," *Proc Natl Acad Sci U S A* 18;97(2):948-53, 2000a.
- Mysore *et al.*, "*Arabidopsis* ecotypes and mutants that are recalcitrant to *Agrobacterium* root transformation are susceptible to germ-line transformation,"
5 *Plant J* 21(1):9-16, 2000b.
- Napoli, Lemieux, Jorgensen, "Introduction of a chimeric chalcone synthase gene into petunia results in reversible co-suppression of homologous genes *in trans*,"
Plant Cell, 2:279-289, 1990.
- Negrutiu, I., Hinnisdaels, S., Cammaerts, D., Cherdshewasart, W., Gharti-Chhetri, G.,
10 and Jacobs, M. "Plant protoplasts as genetic tool: selectable markers for developmental studies," *Int. J. Dev. Biol.* 36: 73-84, 1992.
- Nester, *Ann. Rev. Plant Phys.*, 35:387-413, 1984.
- Nicklas, "The forces that move chromosomes in mitosis," *Annu. Rev. Biophys. Biophys. Chem.*, 17:431-39, 1988.
- 15 Nussbaum *et al.*, *Proc. Nat'l Acad. Sci USA*, 73:1068, 1976.
- Odell *et al.*, *Nature*, 313:810-812, 1985.
- Ohmori and Tomizawa, "Nucleotide sequence of the region required for maintenance of colicin E1 plasmid," *Mol Gen Genet*, Oct 3;176(2):161-70, 1979.
- Omirulleh *et al.*, *Plant Molecular Biology*, 21:415-428, 1993.
- 20 Ow *et al.*, *Science*, 234:856-859, 1986.
- Page *et al.*, "Characterization of a maize chromosome 4 centromeric sequence: evidence for an evolutionary relationship with the B chromosome centromere,"
Genetics 159: 291-302, 2001.
- Palukaitis *et al.*, "Characterization of a viroid associated with avacado sunblotch
25 disease," *Virology*, 99:145-151, 1979.
- Peacock *et al.*, "Highly repeated DNA sequence limited to knob heterochromatin in maize," *Proc. Natl. Acad. Sci. U.S.A.* 78, 4490-4494, 1981.
- Pelissier *et al.*, *Genetica*, 97:141, 1996.
- Pelissier *et al.*, *Plant Mol. Biol.*, 26:441, 1995.
- 30 Perkins, "The detection of linkage in tetrad analysis," *Genetics*, 38, 187-197, 1953.
- Perlak *et al.*, *Proc. Natl. Acad. Sci. USA*, 88:3324-3328, 1991.

- Perriman *et al.*, "Extended target-site specificity for a hammerhead ribozyme," *Gene*, 113:157-163, 1992.
- Peterson *et al.*, "Production of transgenic mice with yeast artificial chromosomes," *Trends Genet.* 13: 61-66, 1997.
- 5 Phi-Van *et al.*, *Mol. Cell. Biol.*, 10:2302-2307, 1990.
- Piatkowski *et al.*, *Plant Physiol.*, 94:1682-1688, 1990.
- Potrykus *et al.*, *Mol. Gen. Genet.*, 199:183-188, 1985.
- Prasher *et al.*, *Biochem. Biophys. Res. Commun.*, 126(3):1259-1268, 1985.
- Preuss *et al.*, "Tetrad analysis possible in *Arabidopsis* with mutation of the QUARTET
10 (QRT) genes," *Science*, 264:1458, 1994.
- Price *et al.*, "Systematic relationships of *Arabidopsis*: a molecular and morphological perspective", in: Somerville, C. and Meyerowitz, E. (eds.) *Arabidopsis*, Cold Spring Harbor Press, NY, 1995.
- Prody *et al.*, "Autolytic processing of dimeric plant virus satellite RNA." *Science*,
15 231:1577-1580, 1986.
- Prokop *et al.*, *Ann. N. Y. Acad. Sci.* 646, 1991
- Puechberty, J. *Genomics* 56:247, 1999
- Rathore *et al.*, *Plant Mol Biol*, 21:871-84, 1993
- Rattner, "The structure of the mammalian centromere," *Bioassays*, 13(2):51-56, 1991.
- 20 Ravatn *et al.*, *Journal of Bacteriology*, 180:5505-14, 1998.
- Reed *et al.*, *J. Gen. Microbiology*, 130:1-4, 1984.
- Reichel *et al.*, *Proc. Nat'l Acad. Sci. USA*, 93 (12) p. 5888-5893, 1996
- Reinhold-Hurek and Shub, "Self-splicing introns in tRNA genes of widely divergent bacteria," *Nature*, 357:173-176, 1992.
- 25 Rensburg *et al.*, *J. Plant Physiol.*, 141:188-194, 1993.
- Richards and Ausubel, "Isolation of a higher eukaryotic telomere from *Arabidopsis thaliana*," *Cell*, 8:53(1):127-36, 1988.
- Richards *et al.*, "The centromere region of *Arabidopsis thaliana* chromosome 1 contains telomere-similar sequences," *Nucleic Acids Research*,
30 19(12):3351-3357, 1991.
- Rieder, "The formation, structure and composition of the mammalian kinetochore and kinetochore fiber," *Int. Rev. Cytol*, 79:1-58, 1982.

- Rogers *et al.*, *Meth. in Enzymol.*, 153:253-277, 1987.
- Rosenberg *et al.*, "RFLP subtraction: A method for making libraries of polymorphic markers," *Proc. Natl Acad. Sci. USA*, 91:6113-6117, 1994.
- Round *et al.*, *Genome Res*, 7, 1053, 1997.
- 5 Sauer, "Functional expression of the cre-lox site-specific recombination system in the yeast *Saccharomyces cerevisiae*," *Mol. and Cell. Biol.*, 7: 2087-2096, 1987.
- Schmidt *et al.*, *Plant Journal*, 5:735-44, 1994
- Schwartz *et al.*, Cold Spring Harbor Symp. *Quant. Biol.*, 47, 195-198, 1982.
- Schweizer *et al.*, "Species specific sequences for the identification of somatic hybrids
10 between *Lycopersicon esculentum* and *Solanum acaule*," *Theor. Appl. Genet.* 75, 679-684, 1998
- Sears *et al.*, "Cytogenetic studies in *Arabidopsis thaliana*," *Can. J. Genet. Cytol.*, 12:217-233, 1970.
- Segal, "Biochemical Calculations" 2nd Edition. John Wiley & Sons, New York, 1976.
- 15 Setlow *et al.*, *Genetic Engineering: Principles and Methods*, 1979.
- Shagan and Bar-Zvi, *Plant Physiol.*, 101:1397-1398, 1993.
- Shapiro, In: *Mobile Genetic Elements*, 1983.
- Sheen *et al.*, *Plant Journal*, 8(5):777-784, 1995.
- Shingo *et al.*, *Mol. Cell. Biol.*, 6:1787, 1986.
- 20 Simoens *et al.*, *Nuc. Acids Res.*, 16:6753, 1988.
- Smith, Watson, Bird, Ray, Schuch, Grierson, "Expression of a truncated tomato polygalacturonase gene inhibits expression of the endogenous gene in transgenic plants," *Mol. Gen. Genet.*, 224:447-481, 1990.
- Smithies *et al.*, *Nature*, 317:230-234, 1985.
- 25 Smythe, "Pollen clusters," *Current Biology*, 4:851-853, 1994.
- Somerville, C. and Somerville, S., *Science* 285:380, 1999.
- Spielmann *et al.*, *Mol. Gen. Genet.*, 205:34, 1986.
- Stalker *et al.*, *Science*, 242:419-422, 1988.
- Stiefel *et al.*, *Nature*, 341:343, 1989.
- 30 Stinchcomb *et al.*, *Nature*, 282:39-43, 1979.
- Stougaard, *The Plant Journal*, 3:755-761, 1993.
- Sullivan, Christensen, Quail, *Mol. Gen. Genet.*, 215(3):431-440, 1989.

- Sun *et al.*, *Cell*, 91:1007, 1997.
- Sutcliffe, *Proc. Nat'l Acad. Sci. USA*, 75:3737-3741, 1978.
- Symington *et al.*, *Cell*, 52:237-240, 1988.
- Symons, "Avacado sunblotch viroid: primary sequence and proposed secondary
5 structure." *Nucl. Acids Res.*, 9:6527-6537, 1981.
- Symons, "Small catalytic RNAs." *Annu. Rev. Biochem.*, 61:641-671, 1992.
- Tarczyński *et al.*, "Expression of a bacterial *mtlD* gene in transgenic tobacco leads to
production and accumulation of mannitol," *Proc. Natl. Acad. Sci. USA*, 89:1-5,
1992.
- 10 Tarczyński *et al.*, "Stress Protection of Transgenic Tobacco by Production of the
Osmolyte Mannitol," *Science*, 259:508-510, 1993.
- Thillet *et al.*, *J. Biol. Chem.*, 263:12500-12508, 1988.
- Thomas *et al.*, *Cell*, 44:419-428, 1986.
- Thomas *et al.*, *Proc. Natl Acad. Sci. USA*, 71:4579, 1974.
- 15 Thompson *et al.*, "Decreased expression of BRCA1 accelerates growth and is often
present during sporadic breast cancer progression," *Nature Genet.*, 9:444-450,
1995.
- Thompson *et al.*, *Nuc. Acids Res.*, 24:3017, 1996.
- Tian, Sequin, Charest, *Plant Cell Rep.*, 16:267-271, 1997.
- 20 Tominaga, *Microbiology*, 143:2057-63, 1997
- Toriyama *et al.*, *Theor Appl. Genet.*, 73:16, 1986.
- Tsay *et al.*, *Science*, 260:342, 1993.
- Tugal *et al.*, *Plant Physiol.*, 120:309, 1999
- Twell *et al.*, *Genes Dev* 5:496-507, 1991
- 25 Twell *et al.*, *Plant Physiol* 91:1270-1274, 1989.
- Tyler-Smith *et al.*, "Mammalian chromosome structure," *Current Biology*, 3:390-397,
1993.
- Uchimiya *et al.*, *Mol. Gen. Genet.*, 204:204, 1986.
- Vahedian *et al.*, "Genomic organization and evolution of the soybean SB92 satellite
30 sequence," *Plant Mol. Biol.* 29: 857-862, 1997.

- Van der Krol, Mur, Beld, Mol, Stuitje, "Flavonoid genes in petunia: addition of a limiting number of copies may lead to a suppression of gene expression," *Plant Cell*, 2:291-99, 1990.
- Van't Hof, Kuniyuki, Bjerken, "The size and number of replicon families of chromosomal DNA of *Arabidopsis thaliana*," *Chromosoma*, 68: 269-285, 1978.
- Vasil *et al.*, "Herbicide-resistant fertile transgenic wheat plants obtained by microprojectile bombardment of regenerable embryogenic callus," *Biotechnology*, 10:667-674, 1992.
- 10 Vasil, *Biotechnology*, 6:397, 1988.
- Vernon and Bohnert, *The EMBO J.*, 11:2077-2085, 1992.
- Voytas and Ausubel, *Nature*, 336:242, 1988.
- Wagner *et al.*, "Coupling of adenovirus to transferrin-polylysine/DNA complexes greatly enhances receptor-mediated gene delivery and expression of transfected genes," *Proc. Nat'l Acad. Sci. USA* 89 (13):6099-6103, 1992.
- 15 Walker *et al.*, *Proc. Nat'l Acad. Sci. USA*, 84:6624-6628, 1987.
- Wang *et al.*, *Molecular and Cellular Biology*, 12(8):3399-3406, 1992.
- Watrud *et al.*, In: *Engineered Organisms and the Environment*, 1985.
- Watson *et al.*, *Recombinant DNA: A Short Course*, 1983.
- 20 Weide *et al.*, "Paracentromeric sequences on tomato chromosome 6 show homology to human satellite III and to the mammalian CENP-B binding box," *Mol. Gen. Genet.* 259 (2): 190-197, 1998.
- Weinsink *et al.*, *Cell*, 3:315-325, 1974.
- Wevrick *et al.*, "Partial deletion of alpha satellite DNA association with reduced amounts of the centromere protein CENP-B in a mitotically stable human chromosome rearrangement," *Mol Cell Biol.*, 10:6374-6380, 1990.
- 25 Whitehouse, *Nature*, No. 4205: 893, 1950.
- Wigler *et al.*, *Cell*, 11:223, 1977.
- Willard, H., *Nature Genetics* 15:345-354, 1997
- 30 Willard, H., "Centromeres of mammalian chromosomes" *Trends Genet.*, 6:410-416, 1990.
- Wolter *et al.*, *The EMBO J.*, 4685-4692, 1992.

- Wong *et al.*, "Electric field mediated gene transfer," *Biochim. Biophys. Res. Commun.* 107(2):584-587, 1982.
- Wright *et al.*, *Genetics*, 142:569, 1996.
- Xia, X. *et al.*, "Structure and evolution of a highly repetitive DNA sequence from
5 *Brassica napus*," *Plant Mol. Biol.* 21:213-224, 1993.
- Xia, X. *et al.*, "Genomic organization of the canrep repetitive DNA in *Brassica juncea*," *Plant Mol. Biol.* 26:817-832, 1994.
- Xiang and Guerra, *Plant Physiol.*, 102:287-293, 1993.
- Xu *et al.*, *Plant Physiol.*, 110:249-257, 1996.
- 10 Yamada *et al.*, *Plant Cell Rep.*, 4:85, 1986.
- Yamaguchi-Shinozaki *et al.*, *Plant Cell Physiol.*, 33:217-224, 1992.
- Yang and Russell, *Proc. Nat'l Acad. Sci. USA*, 87:4144-4148, 1990.
- Yen, *Embo J.* 10(5), 1245-1254, 1991.
- Young *et al.*, In: *Eukaryotic Genetic Systems ICN-UCLA Symposia on Molecular and
15 Cellular Biology, VII*, 315-331, 1977.
- Yuan and Altman, "Selection of guide sequences that direct efficient cleavage of mRNA by human ribonuclease P," *Science*, 263:1269-1273, 1994.
- Yuan *et al.*, "Targeted cleavage of mRNA by human RNase P," *Proc. Natl. Acad. Sci. USA*, 89:8006-8010, 1992.
- 20 Zatloukal *et al.*, "Transferrinfection: a highly efficient way to express gene constructs in eukaryotic cells," *Ann. N.Y. Acad. Sci.*, 660:136-153, 1992.
- Zhang *et al.*, *Gene*, 202:139-46, 1997
- Zhang *et al.*, Zea mays B chromosome centromere repeat sequence
Zea_mays_MBSC216
25 pMBSC216 unpublished
- Zukowsky *et al.*, *Proc. Nat'l Acad. Sci. USA*, 80:1101-1105, 1983.

CLAIMS

The invention is claimed as follows:

1. A method for obtaining a centromere nucleic acid sequence from a selected organism, the method comprising the steps of:
 - 5 preparing a sample of genomic DNA from a selected organism;
 - obtaining a plurality of nucleic acid segments from the genomic DNA; and
 - screening the nucleic acid segments to identify one or more centromere nucleic acid sequences.
- 10 2. The method of claim 1, wherein the step of obtaining comprises contacting said genomic DNA with a restriction endonuclease and selecting nucleic acid segments containing repetitive DNA to obtain said plurality of nucleic acid segments.
- 15 3. The method of claim 1, wherein said the step of obtaining comprises contacting said genomic DNA with a methylation sensitive restriction endonuclease and selecting nucleic acid segments exhibiting resistance to cleavage with said methylation sensitive restriction endonuclease to obtain said plurality of nucleic acid segments.
- 20 4. The method of claim 1, wherein said obtaining comprises fragmenting said genomic DNA and selecting nucleic acid segments that anneal rapidly after denaturation to obtain said plurality of nucleic acid segments.
5. The method of claim 1, wherein the selected organism is a plant.
6. The method of claim 5, wherein the plant is a dicotyledonous plant.
- 25 7. The method of claim 6, wherein the dicotyledonous plant is selected from the group consisting of tobacco, tomato, potato, sugar beet, pea, carrot, cauliflower, broccoli, soybean, canola, sunflower, alfalfa, and cotton.
8. The method of claim 6, wherein the plant is a monocotyledonous plant.
- 30 9. The method of claim 8, wherein the monocotyledonous plant is selected from the group consisting of wheat, maize, rye, rice, turfgrass, oat, barley, sorghum, millet, and sugarcane.

10. A method for identifying a centromere nucleic acid sequence from an organism, the method comprising the steps of:
- providing a first dataset consisting of genomic sequences of the organism wherein the dataset is selected from the group consisting of total genome datasets, and representative genome datasets;
 - identifying and eliminating known non-centromeric repeat sequences from the first dataset by using a BLAST sequence comparison algorithm to create a second dataset;
 - comparing each sequence in the second dataset to itself by using the BLAST sequence comparison algorithm, obtaining a BLAST score for each pair of sequence compared, and collecting high score pairs to create a third dataset;
 - examining the BLAST score of each high score pair in the third dataset and eliminating the pairs having a score greater than 10^{-20} to create a fourth dataset;
 - eliminating the high score pairs in the fourth dataset having less than 80 bp or more than 250 bp to create a fifth dataset;
 - examining the nucleotide position of each high score pair in the fifth dataset and eliminating pairs having 100% identity and identical nucleotide positions to create a sixth dataset;
 - examining the nucleotide position of each high score pair in the sixth dataset and eliminating pairs having opposite orientation of the nucleotides to create a seventh dataset;
 - examining the nucleotide position of both sequences for each high score pair in the seventh dataset and eliminating sequences that are overlapping to create an eighth dataset; and
 - examining the nucleotide position of each sequence in the eighth dataset and eliminating sequences not having at least one neighboring sequence within 250 bp to create a ninth dataset; and
 - comparing each sequence in the ninth dataset to all other sequences in the ninth dataset by using the BLAST sequence

comparison algorithm and selecting the most common sequence as a centromere sequence of the organism.

11. The method of claim 10, wherein the known non-centromeric repeat sequence in the second step is a ribosomal DNA.
- 5 12. The method of claim 10, wherein the steps occur in the listed order.
13. A method for identifying centromere sequences comprising the steps of using a BLAST algorithm to identify centromere sequences from a genomic sequence database.
14. An isolated nucleic acid comprising a plant centromere having n copies of a repeated nucleotide sequence isolated from total genomic DNA of the plant, wherein n is at least 2.
- 10 15. The isolated nucleic acid of claim 14, wherein the repeated nucleotide sequence is provided by any circular permutation of the repeated nucleotide sequence.
- 15 16. The isolated nucleic acid of claim 14, wherein n is from about 5 to about 100,000.
17. The isolated nucleic acid of claim 14, wherein n is from about 10 to about 80,000.
18. The isolated nucleic acid of claim 14, wherein n is from about 25 to about 60,000.
- 20 19. The isolated nucleic acid of claim 14, wherein n is from about 100 to about 50,000.
20. The isolated nucleic acid of claim 14, wherein n is from about 200 to about 40,000.
- 25 21. The isolated nucleic acid of claim 14, wherein n is from about 400 to about 30,000.
22. The isolated nucleic acid of claim 14, wherein n is from about 1,000 to about 30,000.
23. The isolated nucleic acid of claim 14, wherein n is from about 5,000 to about 20,000.
- 30 24. The isolated nucleic acid of claim 14, wherein n is from about 10,000 to about 15,000.

25. An isolated nucleic acid comprising a *Brassica oleracea* centromere, wherein the centromere comprises n copies of a repeated nucleotide sequence isolated from *Brassica oleracea*, and wherein n is at least 2.
- 5 26. The isolated nucleic acid of claim 25, wherein the repeated nucleotide sequence is given by SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3 or SEQ ID NO:4.
27. The isolated nucleic acid of claim 25, wherein the repeated nucleotide sequence is provided by any circular permutation of the repeated nucleotide sequence.
- 10 28. The isolated nucleic acid of claim 25, wherein n is from about 5 to about 100,000.
29. The isolated nucleic acid of claim 25, wherein n is from about 10 to about 80,000.
- 15 30. The isolated nucleic acid of claim 25, wherein n is from about 25 to about 60,000.
31. The isolated nucleic acid of claim 25, wherein n is from about 100 to about 50,000.
32. The isolated nucleic acid of claim 25, wherein n is from about 200 to about 40,000.
- 20 33. The isolated nucleic acid of claim 25, wherein n is from about 400 to about 30,000.
34. The isolated nucleic acid of claim 25, wherein n is from about 1,000 to about 30,000.
- 25 35. The isolated nucleic acid of claim 25, wherein n is from about 5,000 to about 20,000.
36. The isolated nucleic acid of claim 25, wherein n is from about 10,000 to about 15,000.
- 30 37. An isolated nucleic acid comprising a *Glycine max* centromere, wherein the centromere comprises n copies of a repeated nucleotide sequence isolated from *Glycine max*, and wherein n is at least 2.

38. The isolated nucleic acid of claim 33, wherein the repeated nucleotide sequence is given by SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7 or SEQ ID NO:8.
- 5 39. The isolated nucleic acid of claim 37, wherein the repeated nucleotide sequence is provided by any circular permutation of the repeated nucleotide sequence.
40. The isolated nucleic acid of claim 37, wherein n is from about 5 to about 100,000.
- 10 41. The isolated nucleic acid of claim 37, wherein n is from about 10 to about 80,000.
42. The isolated nucleic acid of claim 37, wherein n is from about 25 to about 60,000.
43. The isolated nucleic acid of claim 37, wherein n is from about 100 to about 50,000.
- 15 44. The isolated nucleic acid of claim 37, wherein n is from about 200 to about 40,000.
45. The isolated nucleic acid of claim 37, wherein n is from about 400 to about 30,000.
- 20 46. The isolated nucleic acid of claim 37, wherein n is from about 1,000 to about 30,000.
47. The isolated nucleic acid of claim 37, wherein n is from about 5,000 to about 20,000.
48. The isolated nucleic acid of claim 37, wherein n is from about 10,000 to about 15,000.
- 25 49. An isolated nucleic acid comprising a *Lycopersicon esculentum* centromere, wherein the centromere comprises n copies of a repeated nucleotide sequence isolated from *Lycopersicon esculentum*, and wherein n is at least 2.
- 30 50. The isolated nucleic acid of claim 49, wherein the repeated nucleotide sequence is given by SEQ ID NO:9 or SEQ ID:10.

51. The isolated nucleic acid of claim 49, wherein the repeated nucleotide sequence is provided by any circular permutation of the repeated nucleotide sequence.
52. The isolated nucleic acid of claim 49, wherein n is from about 5 to about 100,000.
53. The isolated nucleic acid of claim 49, wherein n is from about 10 to about 80,000.
54. The isolated nucleic acid of claim 49, wherein n is from about 25 to about 60,000.
55. The isolated nucleic acid of claim 49, wherein n is from about 100 to about 50,000.
56. The isolated nucleic acid of claim 49, wherein n is from about 200 to about 40,000.
57. The isolated nucleic acid of claim 49, wherein n is from about 400 to about 30,000.
58. The isolated nucleic acid of claim 47, wherein n is from about 1,000 to about 30,000.
59. The isolated nucleic acid of claim 47, wherein n is from about 5,000 to about 20,000.
60. The isolated nucleic acid of claim 47, wherein n is from about 10,000 to about 15,000.
61. An isolated nucleic acid sequence comprising a *Zea mays* centromere, wherein the centromere comprises n copies of a repeated nucleotide sequence isolated from *Zea mays*, and wherein n is at least 2.
62. The isolated nucleic acid sequence of claim 61, wherein the repeated nucleotide sequence is give by SEQ ID NO:11, SEQ ID NO: 12 or SEQ ID NO:13.
63. The isolated nucleic acid sequence of claim 61, wherein the repeated nucleotide sequence is provided by any circular permutation of the repeated nucleotide sequence.
64. The isolated nucleic acid of claim 61, wherein n is from about 5 to about 100,000.

65. The isolated nucleic acid of claim 61, wherein n is from about 10 to about 80,000.
66. The isolated nucleic acid of claim 61, wherein n is from about 25 to about 60,000. The isolated nucleic acid of claim 61, wherein n is from about 100 to about 50,000.
67. The isolated nucleic acid of claim 61, wherein n is from about 200 to about 40,000.
68. The isolated nucleic acid of claim 61, wherein n is from about 400 to about 30,000.
69. The isolated nucleic acid of claim 61, wherein n is from about 1,000 to about 30,000.
70. The isolated nucleic acid of claim 61, wherein n is from about 5,000 to about 20,000.
71. The isolated nucleic acid of claim 61, wherein n is from about 10,000 to about 15,000.
72. A recombinant DNA construct comprising a plant centromere comprising n copies of a repeated nucleotide sequence obtained from total genomic DNA of the plant, and wherein n is at least 2.
73. The recombinant DNA construct of claim 72, which additionally comprises a telomere.
74. The recombinant DNA construct of claim 72, which additionally comprises an autonomous replicating sequence (ARS).
75. The recombinant DNA construct of claim 72, which additionally comprises a selectable marker gene.
76. The recombinant DNA construct of claim 72, which additionally comprises at least a structural gene.
77. The recombinant DNA construct of claim 76, wherein the structural gene is selected from the group consisting of a selectable or screenable marker gene, an antibiotic resistance gene, a ligand gene, an enzyme gene, a herbicide resistance gene, a nitrogen fixation gene, a plant pathogen defense gene, a plant stress-induced gene, a toxin gene, a receptor gene, a gene encoding an enzyme, a gene encoding an

- antibody, a gene encoding an antigen for a vaccine, a transcription factor, a cytoskeletal protein, a DNA-binding protein, a protease, an endonuclease, a lipid, a seed storage gene, an interleukin gene, a clotting factor gene, a cytokine gene, a growth factor gene and a biosynthetic gene for producing pharmaceutically active proteins, small molecules with medicinal properties, chemicals with industrial utility, nutraceuticals, carbohydrates, RNAs, lipids, fuels, dyes, pigments, vitamins, scents, flavors, vaccines, antibodies, and hormones.
- 5
78. The recombinant DNA construct of claim 76, wherein the construct is capable of expressing the desired structural gene.
- 10
79. The recombinant DNA construct of claim 72, which is capable of being maintained as a chromosome, wherein the chromosome is stably transmitted in dividing cells.
80. A recombinant DNA construct comprising a plant centromere, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2, and wherein the repeated nucleotide sequence is given by SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, or SEQ ID NO:13.
- 15
- 20
81. A plasmid comprising a recombinant DNA construct, wherein the recombinant DNA comprises a plant centromere, the centromere comprises n copies of a repeated nucleotide sequence obtained from total genomic DNA of the plant, and wherein n is at least 2.
- 25
82. The plasmid of claim 81, which additionally comprises a telomere.
83. The plasmid of claim 81, which additionally comprises an autonomous replicating sequence (ARS).
84. The plasmid of claim 81, which additionally comprises a selectable marker gene.
- 30
85. The plasmid of claim 81, which additionally comprises at least a structural gene.

86. The plasmid of claim 85, wherein the structural gene is selected from the group consisting of a selectable or screenable marker gene, an antibiotic resistance gene, a ligand gene, an enzyme gene, a herbicide resistance gene, a nitrogen fixation gene, a plant pathogen defense gene, a plant stress-induced gene, a toxin gene, a receptor gene, a gene encoding an enzyme, a gene encoding an antibody, a gene encoding an antigen for a vaccine, a transcription factor, a cytoskeletal protein, a DNA-binding protein, a protease, an endonuclease, a lipid, a seed storage gene, an interleukin gene, a clotting factor gene, a cytokine gene, a growth factor gene and a biosynthetic gene for producing pharmaceutically active proteins, small molecules with medicinal properties, chemicals with industrial utility, nutraceuticals, carbohydrates, RNAs, lipids, fuels, dyes, pigments, vitamins, scents, flavors, vaccines, antibodies, and hormones.
87. The plasmid of claim 85, wherein the construct is capable of expressing the desired structural gene.
88. A recombinant DNA construct, wherein the construct comprises a plant centromere, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2, and wherein the repeated nucleotide sequence is given by SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, or SEQ ID NO:13.
89. A host transformed with a recombinant DNA construct comprising a plant centromere, the centromere comprises n copies of a repeated nucleotide sequence obtained from total genomic DNA of the plant, and wherein n is at least 2.
90. The host of claim 89, wherein the cell is a eukaryotic cell.
91. The host of claim 89, wherein the cell is a yeast cell.
92. The host of claim 89, wherein the cell is a higher eukaryotic cell.
93. The host of claim 92, wherein the cell is a crop cell.

- 5 94. A host transformed with a recombinant DNA construct comprising a plant centromere, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2, and wherein the repeated nucleotide sequence is given by SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, or SEQ ID NO:13.
- 10 95. A method of preparing a transgenic plant cell comprising contacting a starting plant cell with a recombinant DNA construct comprising a plant centromere, the centromere comprises n copies of a repeated nucleotide sequence obtained from total genomic DNA of the plant, wherein n is at least 2, and whereby the starting plant cell is transformed with the recombinant DNA construct.
- 15 96. A method of preparing a transgenic plant cell comprising contacting a starting plant cell with a recombinant DNA construct comprising a plant centromere, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2 and the repeated nucleotide sequence is given by SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, or SEQ ID NO:13, and whereby the starting plant cell is transformed with the recombinant DNA construct.
- 20 97. A minichromosome comprising a plant centromere, wherein the plant centromere comprises n copies of a repeated nucleotide sequence obtained from total genomic DNA of the plant, and wherein n is at least 2.
- 25 98. The minichromosome of claim 97, further comprising a telomere.
99. The minichromosome of claim 97, further comprising an autonomous replicating sequence.
- 30 100. The minichromosome of claim 97, further comprising a structural gene.
101. The minichromosome of claim 100, wherein the structural gene is selected from the group consisting of a selectable or screenable marker

- gene, an antibiotic resistance gene, a ligand gene, an enzyme gene, a herbicide resistance gene, a nitrogen fixation gene, a plant pathogen defense gene, a plant stress-induced gene, a toxin gene, a receptor gene, a gene encoding an enzyme, a gene encoding an antibody, a gene encoding an antigen for a vaccine, a transcription factor, a cytoskeletal protein, a DNA-binding protein, a protease, an endonuclease, a lipid, a seed storage gene, an interleukin gene, a clotting factor gene, a cytokine gene, a growth factor gene and a biosynthetic gene for producing pharmaceutically active proteins, small molecules with medicinal properties, chemicals with industrial utility, nutraceuticals, carbohydrates, RNAs, lipids, fuels, dyes, pigments, vitamins, scents, flavors, vaccines, antibodies, and hormones.
102. The minichromosome of claim 100, wherein the minichromosome is capable of expressing the desired structural gene.
103. A minichromosome comprising a plant centromere, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2, and wherein the repeated nucleotide sequence is given by SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, or SEQ ID NO:13.
104. A transgenic crop comprising a crop cell transformed with a minichromosome that comprises a plant centromere, the centromere comprises n copies of a repeated nucleotide sequence, and wherein n is at least 2.
105. The transgenic crop of claim 104, wherein the minichromosome further comprises at least one structural gene.
106. The transgenic crop of claim 105, wherein the structural gene is selected from the group consisting of a selectable or screenable marker gene, an antibiotic resistance gene, a ligand gene, an enzyme gene, a herbicide resistance gene, a nitrogen fixation gene, a plant pathogen defense gene, a plant stress-induced gene, a toxin gene, a receptor gene, a gene encoding an enzyme, a gene encoding an antibody, a gene

- 5 encoding an antigen for a vaccine, a transcription factor, a cytoskeletal protein, a DNA-binding protein, a protease, an endonuclease, a lipid, a seed storage gene, an interleukin gene, a clotting factor gene, a cytokine gene, a growth factor gene and a biosynthetic gene for producing pharmaceutically active proteins, small molecules with medicinal properties, chemicals with industrial utility, nutraceuticals, carbohydrates, RNAs, lipids, fuels, dyes, pigments, vitamins, scents, flavors, vaccines, antibodies, and hormones.
- 10 107. The transgenic crop of claim 104, wherein the crop is a dicotyledonous plant.
108. The transgenic crop of claim 105, wherein the dicotyledonous plant is selected from the group consisting of tobacco, tomato, potato, sugar beet, pea, carrot, cauliflower, broccoli, soybean, canola, sunflower, alfalfa, and cotton.
- 15 109. The transgenic crop of claim 104, wherein the crop is a monocotyledonous plant.
110. The transgenic crop of claim 109, wherein the monocotyledonous plant is selected from the group consisting of wheat, maize, rye, rice, turfgrass, oat, barley, sorghum, millet, and sugarcane.
- 20 111. A transgenic crop comprising a cell transformed with a minichromosome, wherein the minichromosome comprises a plant centromere, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2, and the repeated nucleotide sequence is given by SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, or SEQ ID NO:13.
- 25 112. A method for preparing a transgenic crop tissue, the method comprising the steps of contacting a starting crop tissue with a recombinant DNA construct comprising a plant centromere, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2,
- 30

whereby the starting crop tissue is transformed with the recombinant DNA construct.

- 5 113. A method for shortening the time required to develop a new crop, the method comprising the steps of providing a plant minichromosome comprising a plant centromere, transforming a plant cell with the minichromosome, and generating a transgenic plant from the plant cell, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2, and the minichromosome comprises at least a structural gene.
- 10 114. The method of claim 113, wherein the new crop is a vegetable crop.
115. The method of claim 113, wherein the new crop is a field crop plant.
116. The method of claim 113, wherein the new crop is a fruit and vine crop.
117. The method of claim 113, wherein the new crop has at least one improved agricultural trait as compared to existing crops of the same species.
- 15 118. A method of providing a vegetable crop comprising the steps of providing a plant minichromosome comprising a plant centromere, providing a vegetable crop cell of the vegetable crop, transforming the vegetable crop cell with the plant minichromosome, and generating the vegetable crop from the transformed vegetable crop cell, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2, and the minichromosome comprises at least a structural gene.
- 20 119. A method of providing a fruit and vine crop comprising the steps of providing a plant minichromosome comprising a plant centromere, providing a fruit and vine crop cell of the fruit and vine crop, transforming the fruit and vine crop cell with the plant minichromosome, and generating the fruit and vine crop from the transformed fruit and vine crop cell, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2, and the minichromosome comprises at least a structural gene.
- 25 120. A method of providing a field crop comprising the steps of providing a plant minichromosome comprising a plant centromere, providing a field
- 30

5 crop cell of the field crop, transforming the field crop cell with the plant minichromosome, and generating the field crop from the transformed field crop cell, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2, and the minichromosome comprises at least a structural gene.

10 121. A method of providing a vegetable crop seed comprising the steps of providing a plant minichromosome comprising a plant centromere, providing a vegetable crop cell of the vegetable crop, transforming the vegetable crop cell with the plant minichromosome, generating the vegetable crop from the transformed vegetable crop cell, fertilizing the vegetable crop, and collecting the seeds from the fertilized vegetable crop, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2, and the minichromosome comprises at least a structural gene.

15 122. A method of providing a fruit and vine crop seed comprising the steps of providing a plant minichromosome comprising a plant centromere, providing a fruit and vine crop cell of the fruit and vine crop, transforming the fruit and vine crop cell with the plant minichromosome, generating the fruit and vine crop from the transformed fruit and vine crop cell, fertilizing the fruit and vine crop, and collecting the seeds from the fertilized fruit and vine crop, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2, and the minichromosome comprises at least a structural gene.

20 25 123. A method of providing a field crop seed comprising the steps of providing a plant minichromosome comprising a plant centromere, providing a field crop cell of the field crop, transforming the field crop cell with the plant minichromosome, generating the field crop from the transformed field crop cell, fertilizing the field crop, and collecting the seeds from the fertilized field crop, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2, and the minichromosome comprises at least a structural gene.

124. A method of providing a crop seed comprising the steps of providing a plant minichromosome comprising a plant centromere, providing a cell of the crop, transforming the cell with the plant minichromosome, generating the crop from the transformed crop cell, fertilizing the crop, and collecting the seeds from the fertilized crop, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2, and the minichromosome comprises at least a structural gene.
125. A method for providing a food product, the method comprising the steps of transforming a crop with a minichromosome comprising a plant centromere and a structural gene, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2, and the structural gene is capable of expressing the food product, and extracting the food product from the transgenic crop.
126. A method for providing a pharmaceutical product, the method comprising the steps of transforming a crop with a minichromosome comprising a plant centromere and a structural gene, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2, and the structural gene is capable of expressing the pharmaceutical product, and extracting the pharmaceutical product from the transgenic crop.
127. A method for providing a chemical product, the method comprising the steps of transforming a crop with a minichromosome comprising a plant centromere and a structural gene, the centromere comprises n copies of a repeated nucleotide sequence, wherein n is at least 2, and the structural gene is capable of expressing the chemical product, and extracting the chemical product from the transgenic crop.

1/47

AGCTTGATTGGATACATAAAGTGGTGGAGAAATCACCAGGAAGT
TGAATAAATCTCATAGGAGTTGGCATGAAGAAGTTATCCCACTT
TCAAATCAGGTGATTCCAGTTTCCCAGTTTGGGAATAGCACAGC
TTCTTCGTCGTTCCAATCAAAACCAGGATGAATCWCCTTTGIRARA
AGCT

FIG. 1A

AGCTTGATTTTGATACATAAAGTAGTGGAGAAATCA YTWGGAAGT
GGAAATAAATCTCATAGGAGTTAGGATGAAGAAGCTATCMCACTT
TCAAATCAGGTGATCCCARTTTCCCTGTTTGGGAATATGACAAAC
TTMTTGTCA TTC TAATCAAAACCAGGAWGAATCGBATGTAARA
AGCT

FIG. 1B

<u>Chrbo1</u>	AGCTTGATTGGATACATAAAGTGGTGGAGAAATCA	CCA	GGAAGTTGAATAAATCTCATAG	
<u>ChrBo2</u>	AGCTTGATTGGATACATAAAGTAGTGGAGAAATCA	YTW	GGAAGTGAATAAATCTCATAG	
<u>position</u>	1.....10.....20.....30.....40.....50.....60			
<u>Chrbo1</u>	GAGTTGGCATGAAGAAGTATCC	CACATTTCAAATCAGGTGATTCCAGTTT	CCCA	GTTTGG
<u>ChrBo2</u>	GAGTTAGGATGAAGAAGCTATCM	CACATTTCAAATCAGGTGATCCCAR	TTTCCCT	GTTTGG
<u>position</u>	61.....70.....80.....90.....100.....110.....120			
<u>Chrbo1</u>	GAATAGCACAGCTTCTTCGTCTTC	CAATCAAACCQGGQTGAATCMCT	TTGTR	AAAGCT
<u>ChrBo2</u>	GAATAATCACAAC	TTCTTGTCAATCAAACCQGGQWGAATCGCB	TTGTAA	RAAGCT
<u>position</u>	121....130.....140.....150.....160.....170.....180			

FIG. 1C

3/47

AGCTTGATTGGATACATAAAGTGGTGGAGAAATCACCAGGAAGT
TGAATAAATCTCATAGGAGTTGGSATGAAGAAGTTATCCCACTT
TCAAATCAGGTGATTCAGTTTCCAGTTTGGGAATAGCACAGC
TTCTTCGTCGTTCCAATCAAAACCAGGATGAATCACTTTGTRAGA
AGCT

FIG. 1D

AGCTTGATTTTGATACATAAAGTARTGGAGAAATCAYYAGGAAGT
KGAATAAATCTCATAGGAGTTAGGATGAAGAAGCTATCCCACTT
TCAAATCAGGTGATCCCARTTTTCCTGTTTGGGAATAKGACARC
TTCTTTGTCATTCTAATCAAAACCAGGAWGAATCGKATGTAAARA
AGCT

FIG. 1E

4/47

Chrbo1
ChrBo2
position
AGCTTGATTGGATACATAAAGTGGTGGAGAATCA¹CCAGGAAGTTGAATAAACTCATAG
AGCTTGATTGGATACATAAAGTARITGGAGAATCA²YAGGAAGTKGAATAAACTCATAG
1.....10.....20.....30.....40.....50.....60

Chrbo1
ChrBo2
position
GAGTTGGSATGAAGAAGTTATCCCACTTTCAAATCAGGTGATTTCCA¹GTTTCCAGTTTGG
GAGTTAGGATGAAGAAGCTATCCCACTTTCAAATCAGGTGATCCCA²RITTTCCCTGTTTGG
61.....70.....80.....90.....100.....110.....120

Chrbo1
ChrBo2
position
GAATAGCACAGCTTCTTCTCGTCGTTCC¹AATCAAACCAGGA²TGAATCACTTTGT³RAGAAGCT
GAATAKGACARCTTCTTTGTCA¹TTCT²AATCAAACCAGGA³WGAATCGCKATGT⁴AARAAGCT
121....130....140....150....160....170....180

FIG. 1F

5/47

AAATTCAAATGGTCATAACTTTTMACWC GGAKGTCCGATTCAGG
CGCATAATATATCGAGACGCTCGAAATTGAACAA YGGAAGCTCT
CGAG

FIG. 2A

AAATTCAAACGACAATAACTTTTACTCGGATGTCYGATTGAGT
CCCGTAATATATCGAGACGCTCGAAATTGAATRYTGAAAGCTCTG
AGC

FIG. 2B

<u>Chrbo1</u>	AAATTCAAAT	G	GTC	ATAACTTTTT	M	ACWCGGAK	GTCCGATT	CAGGC	CC	CC	ATAATATATCGAG
<u>ChrBo2</u>	AAATTCAAAC	CGACA	ATAACTTTTT	T	ACTCGGAT	GTCYGATT	GAGT	CCCC	GTAAATATATCGAG		
<u>position</u>	1	10	20	30	40	50	60				
<u>Chrbo1</u>	ACGCTCGAAATTGAA	CAAY	GGAAGCTCTCGAG	92							
<u>ChrBo2</u>	ACGCTCGAAATTGAA	TRYT	GGAAGCTCT-GAGC	91							
<u>position</u>	61	70	80	90							

FIG. 2C

AAATTCAAATGGTCATAAACTTTTMACWCGGAKGTCCGATTTCAGG
CGCATAATATATCGAGACGCTCGAAATTGAACAAYGGAAGCTCT
CGAG

FIG. 2D

AAATTCAAACGACAAATAACTTTTACTCGGATGTCYGATTGAGT
CCCGTAATATATCGAGACGCTCGAAATTGAATRYTGAAAGCTCTG
AGC

FIG. 2E

<u>Chrbo1</u>	AAATTCAAATGGTCATAACTTTT	MACWCGGAKGTC	CGATT	CAGCGCCA	TAATATATCGAG
<u>ChrBo2</u>	AAATTCAAACGACAATAACTTTT	TACTCGGAT	GTCYGATT	GAGTCCCG	TAATATATCGAG
<u>position</u>	1.....10.....20.....30.....40.....50.....60				

<u>Chrbo1</u>	ACGCTCGAAATTGAA	CAAY	GGAAGCTCTCGAG	92
<u>ChrBo2</u>	ACGCTCGAAATTGAA	TRYT	GGAAGCTCT-GAGC	91
<u>position</u>	61.....70.....80.....90			

FIG. 2F

8/47

CCATCACGGGTTTCTGGGCCRTTGGAAAGGTCAAACGAGCCCC
GGAGCGAGCATACGCCCTCATTTTGACGATTTTCGTGCTATTG
CACACCATTTTGGGTGATCGGGATTCGACGTCAAAAATGCC
AAATTGTTCGTGGACGTCCGTCAAGACGTTGTCTATGCATACG
GTTCG

FIG. 3A

CCATCACGGGTTTCTGGGCCRTTGGAAAGGTCAAACGAGCCCC
GRAGCGAGCATACGCCCTCATTTTGACGATTTTCGTGCTATTG
CACACCATTTTGGGTGATCGGGATTCGACGTCAAAAATGCC
AAATTGTTCGTGGACGTCCGTCAAGACGTTGTCTATGCATACG
GTTCG

FIG. 3B

9/47

GGCCACACAACCCCCATTTTTGTCTCGAAAAATAGCCATGAACGACC
ATTTTCAATAATACYRAAGGCTAACACCTACGGATTTTTRACCA
AGAAATGGTCTCCACCAGAAATCCAAGAATGTGATCTATGGCAA
GGAAACATATGTGGGTGAGGTGTAYGAGCCTCTGCTCGAYGAT
CAAT

FIG. 4A

GGCCACACAACCCCCATTTTTGTCTCGAAAAATAGCCATGAAYCGACC
ATTTTCAATAATACCGAAGGCTAACACCTACGGATTTTIGACCA
AGAAATGGTCTCCACCAGAAATCCAAGAATGTGATCTATGGCAA
GGAAACATATGTGGGTGAGGTGTAYGAGCCTCTGCTCGAYGAT
CAAT

FIG. 4B

10/47


GGTTCCGGTGGCAAAACTCGTAGCTTTGTATGCACCCMGACAAC
CCGTTTTCGGAATGGGTGACGTGYGACAACAGAAATTGCCMGAA
ACCACCCCAAACATGAGTTTGGKACCCTAAAGTAGTGGAATTGGGC
ATGTTTCGTTGTGYGAAACGAAGAAAT

FIG. 4C

11/47

FIG. 5

Mini-chromosome construction:
BAC retrofitting by Cre-Lox recombination



12/47

Mini-chromosome construction: Subcloning of BAC inserts by vector tailing





FIG. 6


13/47

FIG. 7A




14/47

FIG. 7B




15/47

FIG. 7C




16/47

FIG. 7D




17/47

FIG. 7E



18/47

FIG. 7F



19/47

FIG. 7G






FIG. 7H




20/47

FIG. 7I




21/47

FIG. 7J




22/47

FIG. 7K




23/47

FIG. 7L




24/47

FIG. 7M



25/47

FIG. 7N




26/47

FIG. 8A



FIG. 8B



28/47

FIG. 9




FIG. 10A




FIG. 10B




FIG. 10C




FIG. 10D




FIG. 10E




FIG. 10F







FIG. 10G



36/47






FIG. 12A




FIG. 12B

FIG. 13A





FIG. 13B



39/47

FIG. 14A/1

Chromosome 2

Sequenced clone	marker name	marker or primer pair	Marker location	marker properties, position	Forward Primer	Reverse Primer
T13E11	T13E11.01	primer	1755-2385	Lan & Col	AGCGCTGGGATGGGT TGGTTG	TTAATGCGGCAATGGCTGA ACAAG
T13E11	T13E11.30	primer	30628-31270	Lan & Col	CAGGTTGCGGTTACTA CATGGGTTTCAG	TATGCATGCGAGTTGGTGG AGGTAAAG
T13E11	T13E11.48	primer	48187-48969	Lan & Col	AACCCCGTAAATTA AACCACAC	CGATACGGGCATGACTCCA G
T13E11	T13E11.63	primer	63886-64530	Lan & Col	ACGCGCGTTGAGAGG AGAAGC	CCCCAAACGCAGCAAGAC AATC
T13E11	T13E11.78	primer	78190-78878	Lan & Col	AACAAAACAAATGCC AGGTCAGG	CTCCGGTCGCAAAGTTACA TACAG
T13E11	T13E11.93	primer	93907-94579	Lan & Col	GTTACCCCGGTCCTGA GATTGAG	TTGGGGAGCAGGATTTGAT GTG
F27C21	F27C21.18	primer	18383-19057	Lan & Col	CAGGCGATTGTCTTTT ATAGGCTGTAAG	TTTTGCTGGAACGGAGGGA GTAC
F27C21	F27C21.02	primer	2570-3293	Lan & Col	ACAAAAGCCGAATC GTGGAAG	TGCCCTGGTTGATTATTGCT GAAAG
F9A16	F9A16.71	primer	71978-72592	Lan & Col	AATGCTTTTGC GACTC TTTTGAC	TTGTTATTTTGGGTTTGGG TTGG
F9A16	F9A16.53	primer	53253-53921	Dom	CGCAGGCGGCTACTT GTTTG	GATGAATTGATCCGTTGTT TTATGTCT
F9A16	F9A16.38	primer	37116-37733	Dom	CAAGGCGGGAAAACA ACTC	AATTATTTTCAACGGCTCT TTACC
F9A16	F9A16.22	primer	22166-22889	Dom	GTATTTAGCATTATGT TAGTCTGTTAGTGG	GCTCCTTGCGTATTCTTCAC C
F9A16	F9A16.03		3308-4091	Lan & Col	GATCCAGCAACCTTA GCCTCCTTC	GATCCCTCAGTTCGAAATC AATCTTC
T5M2	mito border	marker	17685	Dom	AAGGCATCAACGTTT GTGTG	CTACCAGGTAGGTGAAACG
T5E7	T5E7.23	primer	109092-109688	Lan & Col	ATTATTGGCTGCTGCA CTTCTGTCAC	AGGGCCGAGCTCGTATGG A
T5E7	T5E7.73	primer	73460-74120	Lan & Col	TCTCGGAGTAGGGG CTTGTCTCTG	CGGTCGCCCTCGTTCTGTA TCTG
T5E7	T5E7.58	primer	57942-58583	Lan & Col	AGGGGGCTTACAAGA ATGAAC	TCGCCAATGAAAGAGGGT AG
T5E7	T5E7.40	primer	40913-41537	Lan & Col	GTAGCCGGCTCAGTCT CATAACATC	CTAAATCCCGAAAACCCAA ACCAC
T5E7	mito border	marker	13507	Dom	CGAATTCCTTCAGATG ATGC	TTGCCTTACCTATACCCGAC
T5E7	T5E7.02	primer	2919-3585	Lan & Col	AAATGGCAGAAGCAG AAGCAGGAATAG	TAATGCAAGGGTCTCGTAA TGGAATG
T12J2	T12J2.01	primer	1373-1998	Lan & Col	GTGTGGCCTCGTGTGA CCTGAC	AGTTTGCTTCGTTGCTTGCT TATTATG

40/47

FIG. 14A/2

T12J2	T12J2.19	primer	19369-20038	Lan & Col	CTGGCCCATCCCTTAT CGGTTTAC	AGCGCAATCAAGCTATCCC TACATA
T12J2	T12J2.37	primer	37750-38359	Lan & Col	ACCTCCCCACACTTA AACGACACTG	CTCAGAATCCCAAAACAGA GCCACAC
T12J2	ATEDA.59	marker	50592	Lan & Col	TGAATGCTATGAAAG ATGGATGAAAC	AGACGGCTAGTGATTGGT GG
T12J2	T12J2.56	primer	56455-57533	Lan & Col	AATCGGGCTCGGTTGT GTAGAAAC	ATGGCGCAATCAAAAGCA ATCC
T12J2	T12J2.73	primer	73911-74556	Lan & Col	ACTTGTAGGCCCTTTG ATGTTCTG	TGCTTTGTGTTGCTTTGATT ATTCTATTAG
T14C8	T14C8.1	primer	8862-9544	Lan & Col	ACGAACCCGACGACC ACTG	ACGCCTTTGATTCCATTCT TACC
T14C8	T14C8.6	primer	10837-11485	Lan & Col	GACGGGTTGAAAGAA AGCACAG	AGAAGATGATGGCAAGTT ACGAAGAG
T14C8	T14C8.7	primer	45334-46016	Lan & Col	CGACCATTACGACCC ATAC	AAGCCCATTCAAAAGAGTT AGGAGAG
T14C8	T14C8.3	primer	46672-47283	Lan & Col	ATAGCGTCAGCCCTCA TTTCAG	ACCCTTTTGCTTGTATTTTC GTG
T14C8	T14C8.8	primer	48833-49538	Lan & Col	CAAATGGGCGGAGGG GG	TGTGGCAAGTCATGGGTAA GGAG
T14C8	T14C8.9	primer	71115-71878	Lan & Col	GGTGCGGGAGAACGA TGAC	CCGGTTTCTGCGATATTG GTTAG
T14C8	T14C8.5	primer	73841-74456	Lan & Col	TTCCGCGCCCAAAAG GTG	ACGCCTTTGATTCCATTCT TACC
T14C8	T14C8.4	primer	75442-76122	Lan & Col	AAGAAGGCTGGAAAT TGGTTGAG	GAGCGGAAGTAGATGCAG AATGTC
F7B19	F7B19.1	primer	108-757	Lan & Col	AAGGTCCGGCGGTGG TGAG	GGGTCGAGTGATGTGATTG AGTG
F7B19	F7B19.12	primer	13831	Lan & Col	TAACGTCATCAGCGGT AGGAAAC	TTACAAGCGAGAAAAGAT GAGAAGC
F7B19	F7B19.27	primer	27033	Lan & Col	CCCCGCTGAACTGACT GACTACGAG	TCCGCCACCGATAAGATA CGAC
F7B19	F7B19.2	primer	30189-30791	Lan & Col	GCTCGTTGCGGTTGCT GTTC	CCGCGGTGGCTGCTTTTAG
F7B19	F7B19.43	primer	43142	Lan & Col	CAGGGAAAGTGGTTG GATTGATG	TGCTCTCCGGAAGTGGT G
F7B19	F7B19.3	primer	55446-56209	Lan & Col	ATTGTGTTGCCATCGT CCTTC	TTATCAATGTATTTCCCTG TGTATC
F7B19	F7B19.59	primer	59771	Lan & Col	TTGAAGAATGCGCAT AGCCGTAG	TCTGGGATGAAGAGAAAG AGAACTGTC
F7B19	F7B19.4	primer	70859-71492	Lan & Col	AAAAACCGTGAGACC CATAAATG	TCCAAATCGCGAAAGTGAC AG
F7B19	F7B19.77	primer	77633	Lan & Col	ATAACCGAAGAAGCC GAGAAATC	ATCCGGAGACGAAAATGA ACTTAG
F7B19	F7B19.95	primer	95351	Lan & Col	TCGATTGCCAGCAGA GTCAGAAC	TGGGGGCTTGTAAGGAGG AGTAAC

41/47

FIG. 14A/3

F7B19	F7B19.5	primer	98977-99658	Dom?	AGATGGGGTGCTATTC TTGTATG	GCGGTCGAGTGATTGCTG TAG
F7B19	F7B19.6	primer	112337-113039	Lan & Col	AGGGCGAAACTTTGA GAGCAC	TATCGGGTTTTGAAGAGGG AAGG
T15D9	T15D9.3	primer	2985	Lan & Col	AGCGTCGGCGGTGTG GAG	TCCTGGCAAATTGTCTTCT CGTTG
T15D9	T15D9.1	primer	12299-12914	Lan & Col	GCTCCGCCATCTCCTC GTC	GAAGTCCCATGCCTATCCC TG
T15D9	T15D9.19	MARKER	18991	LAST RECOM- BINANT Col	GAGCCCTTCTATGAGC CTACCTGTTT	AGAGATCCCCTGTTACTAA AGCCTATTCTG
T15D9	T15D9.2	primer	37103-37728	Lan & Col	ATGGGGTAATCGAAT AGTGTGGTC	CCCTAGGGCATCCGTTTTT ATCTC
T15D9	T15D9.3a	primer	52189-52811	Col	CGGAGAAAGTTGGGG GTTAGTTG	GAGAGGTTTGGGTTGGGCT TGTAAG
T15D9	T15D9.55	primer	55134	Lan & Col	GCTGCGAACCACAC TTTGCTC	ATGTTATCGTCGCCCGGTT TTATG
T15D9	T15D9.73	primer	72993	Lan & Col	AACCGGTTGATAGTA GACGAGATG	TCCGGGGTTGCGATAGAG
T15D9	T15D9.4	primer	73930-74552	Lan & Col	GTAAGACGGAGCCCC TGAAG	AACATGTTAAAGCCAATAC CCTCTC
T15D9	T15D9.5	primer	86724-87494	Lan & Col	TCGGAAGGCTAGAG ATGGGTAATG	ATTGGACTATATGGGCCTC GTGAC
T15D9	T15D9.93	primer	93763	Lan & Col	TTTGCGATATTCTAA AGGTGATG	TACTATTGCGCTGCTGTTG AGG
F7K9	F7K9.3	primer	21647-22276	Lan & Col	GGATGCAATGCCCGTI ATGATG	TCGAGGGAGGATGCTGAGT ATG
F7K9	F7K9.2	primer	12216-12843	Lan & Col	CAAAGCGGCCATCTC CTTC	GCAATTCATACCGCCACAT CTG
F7K9	F7K9.1	primer	3590-4226	Lan & Col	ACTATGCGTGGGTGG CTTTGTG	CAGGGGCATGCGGAATCTC
F12P23	F12P23.3	primer	61772-62430	Lan & Col	AGCGAGGTTATCTATC AGGGTTG	GATTAGGTCGCTTCTTCC AGTTAG
F12P23	F12P23.5	primer	44870-45511	Lan & Col	CTTCATTGTCATCATC GTTATTAG	GGTGTGAAGTCTGAGGCTC CC
F12P23	F12P23.4	primer	40880-41507	Lan & Col	TACCCATGCCTTGACT GCTG	TTCTGAACGTGTGTGTCT ATTG
F12P23	F12P23.2	primer	22431-23107	Lan & Col	TCGTGGAATAATTGG TGGGAAC	TCGGGAAGAGTGCCTAAG AG
F12P23	F12P23.1	primer	3352-4026	Lan & Col	ACAATGGCAACAATG GGCTGATAG	TTGGGTCGTTGTTCTAA AG
T4D8	T4D8.5	primer	81647-82250	Lan & Col	CTCGGTCTGGTAATGT GAAGTGGT	ATACGTCGCGGGAGTTGAG
T4D8	T4D8.3	primer	47146-47883	Lan & Col	GCCCGTCTGCCATCTC TATC	CGCCTCCTTCACAGCCACA A

42/47

FIG. 14A/4

T4D8	T4D8.2	primer	21848-22453	Lan & Col	AAACTCGCCGCCTCGT GTAAC	AGGATAAACCCATAGCTTG ACCAG
T4D8	T4D8.1	primer	18915-19589	Lan & Col	CTCGTCTCATCCAAAT CCGTCC	CAATATAACCCCGTCCCGT GAAG

FIG. 14B/1

Chromosome 4

Sequenced clone	marker name	marker or primer pair	Marker location	marker properties, position	Forward Primer	Reverse Primer
T5H22	T5H22.00	primer	248-2654	Lan & Col SSLP	TTTGTTACCCCTTTGGC TCGGACTGG	AAGGGGACACGCAC AAAAACGCTCTC
T5H22	T5H22.21	primer	21508-22868	Col Dom	GTCGCCCTTGGTCTAG TAAATGG	CTGTTTCGTCGCCTTC TGCTG
T5H22	T5H22.41.3	primer	35072-35719	Lan & Col	TTGCGAGAAACTTGCG AGGAACATC	TTAGAAAAGCATCG GGCACCAAAC
T5H22	T5H22.41.4	primer	64404-65591	Col Dom	CTCCCTCGCATATTTTG TGACTG	GTTGCCAAAGTTCTC TACGATTC
T5H22	T5H22.65	primer	65036-66470	Lan & Col	TCAACCTAAGGCAAAT TTTCTAAG	TTTAATGAAGGCC AACACC
T7M24	T7M24.04	primer	4816-8214	Lan & Col	GTGCATGGCCTAAACA ACAG	GTTCTCATAACGGGT CAGTCC
T7M24	T7M24.46	primer	46240-47868	Col Dom	ATGTTATGTTTACGTC GGGGTTGTGTTG	TCTCGGCTCCGGATG CTATTGTATTTTC
T25H8	T25H8.01	primer	1889-2953	Lan & Col	TGACGAAGAAGGGGG AAAAGTTG	TGACGTGGTGAAAG TAGGCTGTGAAG
T25H8	T25H8.17	primer	16846-17990	Col Dom	ACTAAAGCCCCAACTG AAGAGGAAG	AAACCGCCACTACC GCCATAA
T25H8	T25H8.22.9	primer	22482-25074	Ler & Col	AATCGATCCGTCTTTC ACCAAC	CTTCCTGCAGCCGTT CTTC
T24M8	T24M8.65	primer	65402-66309	Lan & Col	CGGCATGACCAAACCC TAAACTC	AGGGGAAAGATGAA AGATGAAATAAG
T24M8	T24M8.54	primer	53851-54655	Col Dom	TAATAAACCGCTCAGC CACCACCTAAG	GGGCTGCTCCAATCT CGCTACAC
T24M8	T24M8.43	primer	42439-43274	Lan & Col	CTTAAATTGCCCGTGA TGATGGTTG	GATGGAGTCGGCAA AAGATAGGATG
T24M8	T24M8.22	primer	22640-23386	Lan & Col	ACGAGAAGCGAAAAC CGAAGATAG	CGAACCTAAACCAA ACCTAAACTGAATC
T24M8	T24M8.09	primer	5961-8374	Lan & Col	GAGTTCTGGGGTAATT TCCTCTCG	ATTCTTGOGTGTCCC TGGTGTAAC
T24H24	T24H24.82	primer	82814-82890	Ler & Col	AGACAGCCGGAAGCA ATGGTGG	TCTCGTGCTGGACA TACTCACTCAC
T24H24	T24H24.66	primer	66082-66765	Ler & Col	GTTGAAGGACCGGAGT TGTTAGAC	TGTGGATCGGTTATT GGAGGG

43/47

FIG. 14B/2

T24H24	T24H24.48	primer	47836-48636	Ler & Col	CCCCCAGCCCATTGAG TGAGTAG	AGCGGCGGCCTTGA GAGTATC
T24H24	T24H24.11	primer	11212-11867	Ler & Col	GGCGGCGTAGTTATGT TGATTGAG	TACCACGGCCCCGA GATACTAAC
T27D20	T27D20.77	primer	77681-78420	Lan & Col	TCGCGCAAATGGGACA CG	CGGGGAGGCTCGGG AATC
T27D20	T27D20.64	primer	64198-66686	Ler & Col	ACCTGCGATAGAGTTG TGAGTTC	CTGCCITTTGCCGATA ATAGTC
T27D20	T27D20.51	primer	51084-51783	Lan & Col	CCGCCGGCTTATGCTG AG	GAAGAGAAATGCCC TGTGAGTCC
T27D20	T27D20.41	primer	41203-42770	Lan & Col	GTGATTCCCAGGACAT TGAGTG	TACATTTTTGCAGCC ATTTTGIG
T27D20	T27D20.06	primer	6107-6887	Col Dom	TCGAATGGCTGAAAGA AAAGAATAAGAG	AAAACGGGTGGCGG AGAATG
T19B17	T19B17.96	primer	96402-97060	Lan & Col	CGTCTCCCGTGAGGTG GC	ATTTTCATAATTATT TGGCGTGTGC
T19B17	T19B17.77	primer	77318-78093	Lan & Col	CGAACCCCATCCGAAC TAAC	TGCCACAACAACT CCACTATG
T19B17	T19B17.59	primer	59092-59808	Lan & Col	AGCGGTCAATGTTCTT CAATGTCGTAG	TATCGCGGCGGAGT CAGGAG
T19B17	T19B17.44	primer	44057-44788	Col Dom	CTGCCCCGACCACCTT TCAAC	TTGCGGATTGCTTAT GCTGTTCTC
T19B17	T19B17.30	primer	30680-31352	Lan & Col	ATCGCCGCCGTCTTCTT CAG	CAGGTTTACCCCGTT CAACTATAATC
T19B17	T19B17.11	primer	11260-12044	Lan & Col	TCATTGCGTCTAGAG GTGGAGTGC	GGGGTAGAAAGAAG CGAGAGGGATAG
T26N6	T26N6.12	primer	12724-13462	Lan & Col	CACGGCATCATTATC AAACGAG	GTAGGATCCGGTG AATAGTGGTGG
T26N6	T26N6.27	primer	27839-28536	Lan & Col	TCTTCCGATGACGACA ACGACAC	ATTCTGCTGCTGCTGCTG ATTCTG
T26N6	T26N6.42	primer	43996-44639	Col Only	GACGGCCTTTTCATTCT CACACAG	TTTCATATTTGCTCA TCTAACCCCTTC
T26N6	T26N6.59	primer	59333-59938	Col Only	GCCTCGAACCTACAC CTCCAC	AGTCGCCGTAGCAA ATGAAACC
T26N6	T26N6.74	primer	74460-75083	Lan & Col	ATGGGGCCCTTTGACT ACTGAG	TCCGGAGACGATTTT GATGAC
T26N6	T26N6.93	primer	93352-93986	Lan & Col	TTCCCGCATGCATTAG TTCTTGIG	TTGCCATCATCTTTC TGTGTTTGCTATC
F4H6	F4H6.44	primer	42567-43173	Lan & Col	GCAGACGCGAGGACA CAGACAG	CAGCCTAAGCCCAT TTGTTTTGAAG
F4H6	F4H6.60	primer	60209-60835	Lan & Col	GTCCAACGCTAGCAA GGTCTG	AGGGGCCAACATGC ACTACAAG
F4H6	F4H6.82	primer	82859-83642	Lan & Col	ACAAATCAGAGGCCCA AAGTCAATG	TGGGCCGAATAACA GCAAGTCC
F4H6	F4H6.100	primer	100331-101001	Lan & Col	ATCCAAACGCCCAAAT GTCAAC	TTAAGTGCGGTGCG GTTCAAATAC

44/47

FIG. 14B/3

T19J18	T19J18.12	primer	12781-13435	Lan & Col	ATGCCCATAAAGAAAG CCCAGTC	CGCCTATCTTCGGTG TCTCGTC
T19J18	T19J18.27	primer	28093-29954	Lan & Col	CAGCGCTGTACAGTGG TCAAATG	CGTGGGTCAGGTGG GTCAGG
T4B21	T19J18.71	primer	6380-7009	Lan & Col	CATTACTTACCCGCTTC CGTCTTTATC	AATGTTAGTGCGAG TTTATGGTTGTGTC
T4B21	T19J18.42	primer	20045-20648	Lan & Col	TGTCGCCTTACTCCATT CGTTC AAC	CGGCCGCCTTCATGT ATCTATCTC
T4B21	T4B21.20	primer	21757-22522	SSLP polymorphic	AATAGGCTTTTCCGGT GCTTCTC	AATTGATTTGGGGT TTCTCTGTTCTC
T4B21	T4B21.35	primer	373446-38074	Lan & Col	GTGAAAGGAGCAGCA GGAACAGTG	ATTATAGGCCAAT GACCCAATCG
T4B21	T19J18.57	primer	38498-39157	Lan & Col	CTATCAAACGCAGTCA AAGAAAGG	AGAAGGTGAGCCAA AGAGATTAGTG
T4B21	T4B21.52	primer	54320-55077	Lan & Col	ATAGACAAAATTGGCA ACACATAACC	CACGCCACTCTTCAT CTCCTTTC
T4B21	T4B21-68	primer	69927-70543	Lan & Col	TTGTCATTGGCGCTGC TCTATC	GCTTTCCCCACCAAT ATCCTTTC
T4B21	T4B21.83	primer	85772-86299	Lan & Col	AAGCCC GCGATTG GT TC	CGCTACGCATGGGT CTATTTG
T1J1	T1J1.08	primer	8862-9483	Lan & Col	TAGAGCGGTAACCTAA CGAATGTGC	ATGTGGGGCCAAAT AAATCAAAAC
T1J1	T1J1.23	primer	23155-23843	Lan & Col	TGGAGGGCTTG CATGT GAGAGTG	CAGAGCCGATGAG AAAACAGAGC
T1J1	T15D16	marker	38027	CAPS	AATCAATTGGTTTCTA CTTTT TAG	AACTCCGACTGAAG GTATAGC
T1J1	T1J1.39	primer	39177-40174	Lan & Col	ACCGGCTCATTGGCTA AAAAGTTC	TTAAGGGTTGGGGT TCATCTGTAC
T1J1	T1J1.50	primer	50248-50937	Lan & Col	AAGTCTGGAAGAGG ATGAGAAACCC	ATAAAGTACGCCGC CCATCAATAG
T32N4	T32N4.09	primer	10175-11108	Lan & Col	GGCAGATACGGCGGGT CCATAC	TCTGAATCGCATCTC CTCGTGTAAG
T32N4	T32N4.24	primer	24917-25724	Lan & Col	CGTGGGAGCTGCCGTA GAAG	GCCGTTGATGATGA AAATAGGGTG
T32N4	T32N4.45	primer	45840-46451	Lan & Col	CGCCCCCTTCAGGTTAG TCC	GTTTGCTCCCCTCCC AGTG
T32N4	T32N4.46	primer	46637-47558	Col Dom	CTGGCGTACGAGAGTG CTTG TG	ATGACCCTGTGCTTT TGCTCCTC
T32N4	T32N4.60	primer	60777-61645	Lan & Col	CTCTCGGCGTTGCTTCT GG	GCCCGGCTGGTGCT ATTCTC
T32N4	T32N4.66	primer	66497-67374	Col Dom	AAAGAAGCGAAACAA CATAACCATAG	GGAGACAAAGAAAT CGGCAGAGTAG
T1J24	T1J24.114	primer	114825-115648	Lan & Col	CATGCCCCGAATTACGA CACCTC	GCGCCAAATCTCTA ACAACACTC
T1J24	T1J24.90	primer	90665-91646	Lan & Col	AATGAATGGGACGAA AACGAAACT	GCATCCCCGGTACT GGTGAG

45/47
FIG. 14B/4

T1J24	T1J24.81	primer	80921-81638	Col Dom	AATCGCGACTTTGCCT TCC	TAAACTACTATCCCA CCACCACTACC
T1J24	T1J24.79	primer	79569-80351	Lan & Col	GTGTATCGGGGGCCAT CTCAG	GCTCAACATCGCCG CAATCT
T1J24	T1J24.61	primer	60440-61245	Lan & Col	CCCAAAGTATAAGCGC CCACCTA	TAAGCGCCTCACTTC ACCATTG
T1J24	T1J24.51	primer	51061-51798	Lan & Col	TCCGGAAGGAGCCACA TAAG	TCCCCAGACCTCTCG TTGAC
T1J24	T1J24.27	primer	27855-28895	Lan & Col	GGCCGGGAGTTGGTCA TAAGG	TCAATTTCAATCCCC GCTGGTC
T1J24	T1J24.23	primer	23943-24800	Lan & Col	TGGTCGGGCATATTGT TTTTCTTGTC	CGGCGCTGTCCCTG GTTCC
T1J24	T1J24.01	primer	683-1663	Lan & Col	TTCCCCAAAAATCGTT CAGC	ACATCGCCTCTTCAA CCCACTC
F6H8	F6H8.70	primer	unknown-unique seq	Lan & Col	ACCCGAGAAGCCGATG ACC	AAATTTGGGGGAGT TGATAAGTGTG
F6H8	F6H8.51	primer	unknown	Lan & Col	GCTAAGCCATCCAAGT TCTGAG	GTTTGAGTCTTTGGC TTTGTATGTTC
F6H8	F6H8.94	primer	unknown	Lan & Col	CGTGCAGGGGAGTGTC GTG	CAATTTCAATCCCCG CTGGTC
F6H8	F6H8.114	primer	unknown	Lan & Col	CGCGGCTGCCTTCATG TATCTATC	GCCCATTTGTCGCCT TATTCTATTC
F2112	F2112.82	primer	82463-83233	Lan & Col	TTTTTG GG GATAGGGA TTGAGTGTG	TAAGCGGAAGGAGA GGTTTGAAGTTG
F2112	F2112.70	primer	70415-71220	Lan & Col	TGCTGGCCTTTGTCATC TATTTGTC	CCGCGGGGACTGCC TACTC
F2112	F2112.68	primer	68874-69938	Lan & Col	CCAGAGCCGGGGAAA GCAATAC	TAGCCGGGGTGGTC TCGTGC
F2112	F2112.50	primer	50288-50891	Lan & Col	TGACTATAGGGGCGGT TGTGGTAAAG	TTGGCTTGAGTTTG CGTCGTC
F2112	F2112.48	primer	48960-50345	Col	ACCTTTCTTCCTCAACG CACCTCACC	AACCCCTTGGCATAT AACTCCGACTC
F2112	F2112.29	primer	29895-30702	Lan & Col	GTGGGGTCGAGTGGTG TGGTAG	GGATCCCCCTGTTACT TAAGCCTATTC
F2112	F2112.02	primer	2313-3098	Lan & Col	AAAATCCTCCCGCGTC AACATC	CATCATCCCAATCCC AAATACAAGTC
F14G16	F14G16.100	primer	3496-4174	Lan & Col	AAACTTTGCGCACTCT CCTCTATTATG	ATTTGCGTAAGGCG TTGATGACTC
F14G16	F14G16.81	primer	22905-23604	Lan & Col	CGTCTTCATCGGCTTC GTTTCAAG	TGGGGAGCGGAGGA TTCTTG
F14G16	F14G16.66	primer	37689-38299	Lan & Col	AGCGATTGTACCCCCA CCATTTC	GCTCCGGCAATCTTC TTCCTCTC
F14G16	F14G16.49	primer	54150-54777	Lan & Col	ACTTTGGGCAATGAAG CGTATG	AACCCCTTAGGATT ATTCGTAGTGTTC
F28D6	F14G16.32	primer	8172-8825	Lan & Col	TCTCGCAGTTGCAGAG ATGGTG	TCCGCGAAGAGAAG AGTGATGG

46/47

FIG. 14B/5

F28D6	F14G16.66	primer	9445-10055	Lan & Col	AGCGATTGTACCCCCA CCATTC	GCTCCGGCAATCTTC TTCCTCTC
F28D6	F14G16.16	primer	24251-24873	Lan & Col	TGGTGTATTTTGTCTT GTTTCTCAGG	GTTGTCCGCTATGG GGCTAAGG
F28D6	F14G16.01	primer	39801-40577	Lan & Col	GTGCGGAAATGTCTGG GCTC	AATCACTCAACCGC GAAACTCTATC
F28D6	F28D6.42	primer	42565-43225	Lan & Col	ATCAACCCCCAAATCA CCAGAAAC	AATCGCGGTTAGCC ACTTCATC
F28D6	F28D6.50K	MARKER	50323	Col Dom	CGGCTGGCTTTATTAT CTGAGTTG	TTCGGGAAGCCTGT GGAAG
F28D6	F28D6.58	primer	58994-59869	Col Dom	ACCCCGAGCTCAACTT CTTAGG	GGACGGGAGATGGG ATTACC
F28D6	F28D6.76	primer	75571-77289	Lan & Col	AGAATAGGAGCTGGG AGGTCAAAC	ATACTTAGATGCAA TGGGTGTGGTG
F28D6	F28D6.93	primer	93823-94512	Lan & Col	CCCCATCCTGCCGACA TAAAG	TACTCCGCATCATCT TCCATCTCTTC
F28D6	F28D6.120	primer	7985-8702	Lan & Col	GAGGGGCGAGTAGTTG AATCTGC	CCTAAGCCCCGAAAC CAAGTGAG




FIG. 15

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Pruess, Daphne
Copenhaver, Gregory
Keith, Kevin
- (ii) TITLE OF INVENTION: PLANT CENTROMERE COMPOSITIONS
- (iii) NUMBER OF SEQUENCES: 13
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Bell, Boyd & Lloyd
 - (B) STREET: 70 West Madison
 - (C) CITY: Chicago
 - (D) STATE: Illinois
 - (E) COUNTRY: USA
 - (F) ZIP: 60602
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC Compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn 3.1 Version 3.1.15
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: US Unknown
 - (B) FILING DATE: Concurrently Herewith
 - (C) CLASSIFICATION: Unknown
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 09/553,231
 - (B) FILING DATE: 2002-03-13
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 09/090,051
 - (B) FILING DATE: 1998-06-03
- (vii) PRIOR APPLICATION DATA:
 - A) APPLICATION NUMBER: 60/048,451
 - (B) FILING DATE: 1997-06-03
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 60/073,741
 - (B) FILING DATE: 1998-02-05
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 09/531,120
 - (B) FILING DATE: 2000-03-17

(vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 60/125,219
(B) FILING DATE: 1999-03-18

(vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 60/127,409
(B) FILING DATE: 1999-04-01

(vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 60/134,770
(B) FILING DATE: 1999-05-18

(vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 60/153,584
(B) FILING DATE: 1999-09-13

(vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 60/154,603
(B) FILING DATE: 1999-09-17

(vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 60/172,493
(B) FILING DATE: 1999-12-16

(viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Barrett, Robert M.
(B) REGISTRATION NUMBER: 30,142
(C) REFERENCE/DOCKET NUMBER: 0112283-042

(ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: 312-807-4204
(B) TELEFAX: 312-827-1270

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 180
(B) TYPE: DNA
(C) ORGANISM: Brassica oleracea
(D) SEQUENCE DESCRIPTION:

```
agcttgattt ggatacataa agtggtggag aatcaccagg aagttgaata aatctcatag 60
gagttggcat gaagaagtta tcccactttc aaatcaggtg attccagttt cccagtttgg 120
gaatagcaca gcttcttcgt cgttccaatc aaaccaggat gaatcwcttt gtraraagct 180
```

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 180

(B) TYPE: DNA

(C) ORGANISM: Brassica oleracea

(D) SEQUENCE DESCRIPTION:

agcttgattt tgatacataa agtagtggag aatcaytwgg aagtggaata aatctcatag 60
gagttaggat gaagaagcta tcmactttc aaatcagggtg atcccarttt tcctgtttgg 120
gaatagcaca acttmtttgt cattctaate aaaccaggaw gaatcgcbat gtaaraagct 180

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 180

(B) TYPE: DNA

(C) ORGANISM: Brassica oleracea

(D) SEQUENCE DESCRIPTION:

agcttgattt ggatacataa agtggtggag aatcaccagg aagttgaata aatctcatag 60
gagttggsat gaagaagtta tcccactttc aaatcagggtg attccagttt cccagtttgg 120
gaatagcaca gcttcttcgt cgttccaatc aaaccaggat gaatcacttt gtragaagct 180

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 180

(B) TYPE: DNA

(C) ORGANISM: Brassica oleracea

(D) SEQUENCE DESCRIPTION:

agcttgattt tgatacataa agtartggag aatcayyagg aagtkgaata aatctcatag 60
gagttaggat gaagaagcta tcccactttc aaatcagggtg atcccarttt tcctgtttgg 120
gaatakacaca rcttctttgt cattctaate aaaccaggaw gaatcgckat gtaaraagct 180

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 92

(B) TYPE: DNA

(C) ORGANISM: Glycine max

(D) SEQUENCE DESCRIPTION:

aaattcaaat ggtcataact tttmacwgg akgtccgatt caggcgcata atatatcgag 60
acgctcgaaa ttgaacaayg gaagctctcg ag 92

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 91

(B) TYPE: DNA

(C) ORGANISM: Glycine max

(D) SEQUENCE DESCRIPTION:

aaattcaaac gacaataact ttttactcgg atgtcygatt gagtcccgtata atatatcgag 60
acgctcgaaa ttgaatrytg aagctctgag c 91

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 92
- (B) TYPE: DNA
- (C) ORGANISM: Glycine max
- (D) SEQUENCE DESCRIPTION:

aaattcaaat ggtcataact tttmacwgg akgtccgatt caggcgcata atatatcgag 60
acgctcgaaa ttgaacaayg gaagctctcg ag 92

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 91
- (B) TYPE: DNA
- (C) ORGANISM: Glycine max
- (D) SEQUENCE DESCRIPTION:

aaattcaaac gacaataact ttttactcgg atgtcygatt gagtcccgtg atatatcgag 60
acgctcgaaa ttgaatrytg aagctctgag c 91

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 181
- (B) TYPE: DNA
- (C) ORGANISM: Lycopersicon esculentum
- (D) SEQUENCE DESCRIPTION:

ccatcacggg ttttctgggc crtttggaag gtcaaacgag ccccgagcgc agcatacgcc 60
tcattttgac gatttttcgtg tgctattgca caccatTTTT tgggtgatcg ggattccgac 120
gtcaaaaatg ccaaatttgt tcgtggacgt ccgtaagac gttgtctatg catacggttg 180
g 181

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 181
- (B) TYPE: DNA
- (C) ORGANISM: Lycopersicon esculentum
- (D) SEQUENCE DESCRIPTION:

ccatcacggg ttttctgggc crtttggaag gtcaaacgag ccccgagcgc agcatacgcc 60
tcattttgac gatttttcgtg tgctattgca caccatTTTT tgggtgatcg ggattccgac 120
gtcaaaaatg ccaaatttgt tcgtggacgt ccgtaagac gttgtctatg catacggttg 180
g 181

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 180
- (B) TYPE: DNA
- (C) ORGANISM: Zea mays
- (D) SEQUENCE DESCRIPTION:

ggccacacaa cccccatttt tgtcgaaaat agccatgaac gaccattttc aataatacyr 60
aaggctaaca cctacggatt tttraccaag aaatgggtctc caccagaaat ccaagaatgt 120
gatctatggc aaggaaacat atgtgggggtg aggtgtayga gcctctggtc gaygatcaat 180

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 180

(B) TYPE: DNA

(C) ORGANISM: Zea mays

(D) SEQUENCE DESCRIPTION:

ggccacacaa cccccatttt tgtcgaaaat agccatgaay gaccattttc aataataaccg 60
aaggctaaca cctacggatt ttgaccaag aaatgggtctc caccagaaat ccaagaatgt 120
gatctatggc aaggaaacat atgtgggggtg aggtgtayga gcctctggtc gatgatcaat 180

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 158

(B) TYPE: DNA

(C) ORGANISM: Zea mays

(D) SEQUENCE DESCRIPTION:

ggttccggtg gcaaaaactc gtagctttgt atgcacccmg acaccggtt tcggaatggg 60
tgacgtgyga caacagaaat tgcgmgaaac caccctaaac atgagttttg kacctaaagt 120
agtggattgg gcatgttcgt tgygaaaaac gaagaaat 158