

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
12 February 2004 (12.02.2004)

PCT

(10) International Publication Number
WO 2004/013072 A2

(51) International Patent Classification⁷: **C07C**

(21) International Application Number:
PCT/US2003/023673

(22) International Filing Date: 29 July 2003 (29.07.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/400,662 2 August 2002 (02.08.2002) US

(71) Applicant: **RHODIA PHARMA SOLUTIONS INC.**
[US/US]; 56 Roland Street, Boston, MA 02129 (US).

(72) Inventor: **HICKS, Frederick**; 18 Granite Street,
Somerville, MA 02143 (US).

(74) Agent: **MC VEIGH, Kevin, E.**; Rhodia Inc., 259 Prospect
Plains Road, Cranbury, NJ 08512 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COPPER CATALYZED ARYLATION

(57) Abstract: The present invention is directed to an arylation of a nucleophile by reacting the nucleophile with a substrate aromatic compound ArX in the presence of a copper catalyst, a base and water, wherein Ar is aryl, heteroaryl or alkenyl, X is halo, sulfonate or phosphonate, the base comprises an alkaline earth carbonate, bicarbonate, hydroxide or phosphate, and the copper catalyst comprises a copper atom or ion and a ligand.



WO 2004/013072 A2

- 1 -

COPPER CATALYZED ARYLATION

Field of the Invention

5

This invention relates to aromatic bond formation, more specifically to copper-catalyzed formation of aryl and heteroaryl carbon-nitrogen bonds.

Background of the Invention

10

Palladium catalyzed arylation of indoles is known, see, for example, Stephen L. Buchwald *Org. Lett.*, 2, 1403-1406. More recently, copper catalyzed arylation of indoles and amides has been reported, see, for example, Klapars, A., Antilla, J.C., Huang, X., and Buchwald, S.L., *J. Am. Chem. Soc.* 2001, 123, 7727-7729 and Buchwald, S. L. *J. Am. Chem. Soc.* 2002, 124, 7421. Buchwald et. al. disclose copper-catalyzed indole arylation with catalytic amounts, that is, from 0.1 to 10 mol%, copper under mild temperatures in solvents such as toluene and 1,4-dioxane and inorganic carbonate or phosphate bases such as K_2CO_3 , Cs_2CO_3 , or K_3PO_4 . The key to this reactivity under mild conditions even in noncoordinating solvents are the use of ligands, for example, diamines, such as 1,2-diaminocyclohexane or N,N'-dimethylethylenediamine.

15
20

Current methods for copper-catalyzed indole arylation employing aryl bromides suffers from slow reaction rates, a variable induction period, use of large amounts of inorganic bases such as K_3PO_4 , which lead to reactor agitation issues at minimal solvent volumes or without organic solvents and reactor capacity issues as well.

25

- 2 -

Summary of the Invention


The present invention is directed to cross-coupling reaction method for
5 arylating a nucleophile comprising reacting the nucleophile with a substrate
aromatic compound ArX in the presence of a copper catalyst, a base and
water, wherein Ar is aryl, heteroaryl or alkenyl, X is halo, sulfonate or
phosphonate, the base comprises an alkaline earth carbonate, bicarbonate,
hydroxide or phosphate, and the copper catalyst comprises a copper atom or
10 ion and a ligand.

The method of the present invention allows the amount of base to be
reduced compared to prior methods, thus minimizing reactor agitation and
capacity issues.

15

In one embodiment, the present invention is directed to a method for
arylation of a HN-containing heterocycle, comprising reacting the HN-containing
heterocycle with a substrate aromatic compound ArX according to the
reaction scheme:

20




wherein Ar is aryl, heteroaryl or alkenyl,
25 X is halo, sulfonate or phosphonate,
the base comprises an alkaline earth carbonate, bicarbonate, hydroxide or
phosphate, and


- 3 -

the copper catalyst comprises a copper atom or ion and a ligand.

In a second embodiment, the present invention is directed to a method for arylating a HN-containing compound according to the formula $\text{HN}(\text{R}^1)\text{R}^2$,
 5 comprising reacting the HN-containing compound with a substrate aromatic compound, ArX , according to the reaction scheme:




10 wherein Ar is aryl, heteroaryl or alkenyl,
 X is halo, sulfonate or phosphonate
 R^1 is H, alkyl or aryl
 R^2 is according to the formula:



15

wherein R^3 is H, alkyl, aryl, heteroaryl, alkenyl, $-\text{OR}^5$ or $-\text{NR}^6_2$, and R^5 and R^6 are each independently alkyl, aryl, or



20

- 4 -

wherein R^7 is alkyl or aryl,
the base comprises an alkaline earth carbonate, bicarbonate, hydroxide or
phosphate, and
the copper catalyst comprises a copper atom or ion and a ligand.

5

Detailed Description of Invention and Preferred Embodiments

As used herein, the term "substituted" denotes the conceptual
replacement of a hydrogen atom of a given organic moiety with a substituent
10 group other than a hydrogen atom and includes all permissible substituent
groups, including acyclic hydrocarbon groups, alicyclic hydrocarbon groups,
monocyclic aromatic hydrocarbon groups, polycyclic aromatic hydrocarbon
groups, heteroacyclic groups, heterocyclic groups, fused ring systems and
bridged ring systems, of which the substituents specifically described below
15 are illustrative examples.

"Alkyl" refers to a linear, branched or cyclic saturated hydrocarbon
group, preferably a (C_1 - C_{30}) linear, branched or cyclic saturated hydrocarbon
group that may, optionally, contain one or more heteroatoms, such as, for
20 example, methyl, ethyl, propyl, n-butyl, isobutyl, t-butyl, neopentyl,
cyclopentyl, hexyl, cyclohexyl, decyl, stearyl, eicosyl, methoxy, triacontyl,
2,5,7-trioxanonanyl, 2,5,8-triazadecenyl, and that may, optionally, be
substituted at one or more positions with other moieties, such as, for
example, alkyl, alkenyl, alkynyl, aryl, heterocyclyl, halo, hydroxy, sulfhydryl,
25 hydroperoxy, carbonyl-containing groups (including, for example, carboxy,
ketone, ester and aldehyde groups), alkyloxy, alkylodioxy, amino, amido,
imino, hydrazino, nitro, cyanato, thiocyanato, mercapto, thiocarbonyl-
containing groups (including, for example, thioketone groups, thioester
groups and thioaldehyde groups), sulfonyl-containing groups (including, for

- 5 -

example, sulfate, sulfonate and sulfamoyl groups), silyl, siloxy and phosphorus-containing substituent groups (including, for example, phosphoranyl, phosphinyl, phosphinothioyl, phosphinimyl). Such substituent groups may themselves be further substituted with, for example, any of the groups described above as suitable substituents for alkyl groups, to form compound substituent groups, such as, for example, aralkyl, aminoalkyl, haloalkyl, heterocyclalkyl.

As used herein the term "heteroatom" means an element other than carbon, such as for example, oxygen, nitrogen and sulfur.

As used herein, "halo" means fluoro, chloro, bromo or iodo, "hydroxy" means -OH, "sulfhydryl" means -SH, "hydroperoxy" means -OOH, "carbonyl" means -C(O)-, "carboxy" means -COOH, a ketone group is a group containing a carbonyl moiety that is attached to two carbon atoms, an ester group is a group containing a -C(O)OR moiety, an aldehyde group is a group containing a -CHO moiety, "alkyloxy" means -OR', "alkyldioxy" means -OOR', "amino" is conceptually a derivative of NH₃ in which one or more hydrogen atoms are replaced by nonacyl organic groups and includes primary, secondary and tertiary amines, "amido" includes, for example, -C(O)NR₂, "imino" means =NH, "hydrazino" includes, for example, -HNNR₂, "nitro" is -NO₂, "cyanato" is -OCN, "thiocyanato" is -SCN, "mercapto" is -SH, "thiocarbonyl" is -C(S)-, a "thioketone group" is a group containing a thiocarbonyl moiety that is attached to two carbon atoms, a "thioester group" is -C(S)OR, a "thioaldehyde" group is a group containing a -CHS moiety, "sulfonyl" is -SO₂-, "sulfate" includes, for example, -OSO₂OR", "sulfonate" includes, for example, -O₂SOR", "sulfamoyl" includes, for example, -O₂SNR₂, "silyl" is -SiR₃, "siloxy" is -OSiR₃, "phosphino" includes -PR₂, "phosphoranyl" includes, for example, -PR₄, "phosphinyl"

- 6 -

includes, for example, $-P(O)R''_2$, "phosphinothioyl" includes, for example, $-P(S)R''_2$, "phosphinimyl" includes, for example, $-P(NH)R''_2$, "aralkyl" means an alkyl group substituted with an aryl group, such as, for example, benzyl, "aminoalkyl" means an alkyl group substituted with an amino group, such as, for example, dimethylaminoethyl, "haloalkyl" means an alkyl group substituted with a halogen atom, such as, for example, chloromethyl, "heterocyclalkyl" means an alkyl group substituted with a heterocycl group, such as, for example, pyrrolidinyethyl, wherein, as used above in this paragraph, R is an organic group, R' is alkyl and R'' is H, alkyl or aryl.

10

"Alkenyl" refers to a linear, branched or cyclic hydrocarbon group, preferably a (C_2-C_{20}) linear, branched or cyclic hydrocarbon group, that contains one or more carbon-carbon double bonds per group and that may, optionally, contain one or more heteroatoms, such as, for example, ethenyl, propenyl, allyl, isopropenyl, ethenylidenyl, cyclopentyl, cyclohexadienyl, azanonenyl, and that may, optionally, be substituted at one or more positions with other moieties, such as, for example, any of the possible substituents described above in respect to alkyl groups.

15

20

"Alkynyl" refers to a linear, branched or cyclic unsaturated hydrocarbon group, preferably a (C_2-C_{20}) unsaturated hydrocarbon group, that contains one or more carbon-carbon triple bonds per group and that may, optionally, contain one or more heteroatoms, such as, for example, ethynyl, propynyl, thianonyl, and that may, optionally, be substituted at one or more positions with other moieties, such as, for example, any of the possible substituents described above in respect to alkyl groups.

25

"Aryl" refers to an unsaturated hydrocarbon group that contains one or more six membered rings in each of which the unsaturation may be

- 7 -

represented by three conjugated carbon-carbon double bonds, including monocyclic and polycyclic ring systems, such as, for example, phenyl, naphthyl, anthryl, phenanthryl, indenyl, fluorenyl, which may, optionally, be substituted at one or more positions with other moieties, such as, for
5 example, any of the possible substituents described above in respect to alkyl groups.

"Heterocyclyl" and "heterocycle" refer to a saturated or unsaturated organic group or compound that contains one or more rings in which one or
10 more ring members is a heteroatom, preferably a nitrogen, sulfur or oxygen heteroatom, such as, for example, thiacyclopentadienyl, thiaindenyl, thianthrenyl, oxacyclopentadienyl, oxaindenyl, isobenzylfuranyl, pyranal, azacyclopentadienyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolinyl, quinolinyl, isoquinolinyl, phthalazinyl, cinnolinyl, azafluorenyl,
15 phenanthrolinyl, phenazinyl, phenothiazinyl, phenarsazinyl, isothiazolyl, isoxazolyl, phenoxazinyl, pyrrolidinyl, pyrimadinyl, imidazolidinyl, piperidinyl, piperizinyl, oxathiaanthracenyl, isoxazolyl, oxaazaanthracenyl, isothiazolyl, morpholinyl, and which may, optionally, be substituted at one or more positions with other moieties, such as, for example, any of the possible
20 substituents described above in respect to alkyl groups.

As used herein, the term "nucleophile" refers to a chemical moiety having a reactive pair of electrons and the terms "electrophile" and "electrophilic" refer to a chemical moiety that can accept a pair of electrons from a
25 nucleophile.

Compounds suitable as the substrate aromatic compound ArX component of the method of the present invention are those compounds that


- 8 -

contain an electrophilic atom bonded to leaving group X that is susceptible to the above reaction with a nucleophile.

5 In one embodiment, Ar comprises a phenyl ring, which may in addition to the X substituent, be further substituted on one or more carbons of the ring with, for example, any of the groups described above as suitable substituents for alkyl groups. In a preferred embodiment, Ar comprises a phenyl ring, which is further substituted, in addition to the X substituent, on one or more carbons of the ring with one or more substituent groups each independently
10 selected from alkyl, alkoxy, alkenyl, alkynyl, aryl, heteroaryl, cyano, carbonyl, amino, amido or sulfonyl.

In one embodiment, X is a halo, sulfonate or phosphonate group,
more preferably halo.
15

Suitable sulfonate groups include, for example, those according to the general formula:




20

wherein R⁸ is alkyl, aryl, fluoroalkyl, preferably trifluoromethyl, perfluoroalkyl.

- 9 -

Suitable phosphonate groups include, for example, those according to the general formula:



5

wherein each R⁹ is independently alkyl or aryl.

Compounds suitable as the substrate aromatic compound ArX include, for example, 4-bromobenzonitrile, 4-N,N'-dimethyl-bromoaniline, 2-bromothiophene, 3-bromoquinoline, 1-nitro-2-iodobenzene, 4-chlorotoluene, 4-bromofluorobenzene, 2-bromoanisole, 4-iodoaniline, 3-bromoacetophenone, and 4-bromothioanisole. In a highly preferred embodiment, the substrate aromatic compound is 4-bromofluorobenzene.

15 Suitable substrate aromatic compounds are made by known synthetic methods. The substituents on the substrate aromatic compound are selected based on structure of the desired product.


20 Compounds suitable as the nucleophile component of the method of the present invention include HN-containing heterocycles and HN-containing compound according to the formula HN(R¹)R², wherein R¹ and R² are as described above.

25 Compounds suitable as the HN-containing heterocycle component of one of the preferred embodiments of the method of the present invention

- 10 -

include, for example, substituted or unsubstituted triazoles, pyrroles, pyrazoles, imidazoles, indoles, azaindoles, benzotriazoles, benzimidazoles, indazoles, and carbazoles, such as, for example, 3-methylpyrazole, 2-phenylindole, 5-methoxyindole, 5-aminoindole, 5-nitroindole, 3-carbomethoxyindole, benzimidazolylacetonitrile, pyrrole, 7-azaindole, 1,2,4-triazole, and carbazole.

In one embodiment, the HN-containing heterocycle comprises a monocyclic system according to the formula:



10

wherein n is 0 or an integer of from 1 to 3 and R^{10} is substituted alkyl, substituted N, or O, such as, for example, 2-pyrimidinone, phthalazinone, 2-azetidinone, 2-pyrrolidinone, 2-oxazolidinone, or imidazolidinone.

15

In a preferred embodiment, the N-containing heterocycle is 5-chloroindole or 2-pyrrolidinone.

Compounds suitable as the HN-containing compound according to the formula $HN(R^1)R^2$ include amides, carbamates, ureas, and sulfonamides.

20

In one embodiment, the HN-containing compound according to the formula $HN(R^1)R^2$ is an amide, such as for example, benzamide, 4-aminobenzamide, cyclohexylamide, trans-cinnamamide, N-phenylacetamide, N-methylformamide, N-benzylformamide, or N-cyclohexylformamide. In a

25

- 11 -

more highly preferred embodiment, the $\text{HN}(\text{R}^1)\text{R}^2$ compound is selected from benzamide and N-methylformamide.

5 In another embodiment, the HN-containing compound according to the formula $\text{HN}(\text{R}^1)\text{R}^2$ is a carbamate, urea or sulfonamide, such as, for example, N-phenyl-*tert*-butyl carbamate, N-methylimidazolidinone, or p-toluenesulfonamide.

10 The copper atom or ion component of the method of the present invention may be derived from any copper-containing material. In a preferred embodiment, the copper atom or ion is derived from copper metal, Cu_2O or a copper salt, such as CuCl , CuBr , CuBr_2 or CuI . In a preferred embodiment, the copper catalyst is CuI .

15 In a preferred embodiment, the copper catalyst is present in the reaction mixture as a metal-ligand complex wherein the copper catalyst is bound to a supporting ligand.

20 Compounds suitable as the ligand component of the method of the present invention are those compounds that are capable of solubilizing the copper species in the reaction mixture. In a preferred embodiment, the ligand is a 1,2-diamine compound, such as, for example, 1,2-di(aminomethyl)cyclohexane, N,N'-dimethylethylenediamine, 1-propyl-1,2-N,N'-dimethylethylenediamine. In a more highly preferred embodiment, the
25 ligand is 1,2-di(aminomethyl)cyclohexane.

The copper atom or ion component and ligand component may be added to the reaction mixture as separate compounds. Alternatively, a copper-ligand catalyst complex may be formed prior to addition to the

- 12 -

reaction mixture and then added to the reaction mixture as the copper-ligand complex.

5 In general, the coupling reaction is run in the presence of a catalytic amount of the copper catalyst. As used herein, a "catalytic amount" of catalyst refers to an amount of catalyst that provides an increase in the rate of the reaction of the method of the present invention, compared to the rate of the same reaction conducted under analogous conditions, but absent the catalyst. Typically, the amount of copper catalyst ranges from about 0.01 to
10 about 10 mole%, more preferably from about 0.5 to about 5 mole%, based on the amount of limiting reactant.

In a preferred embodiment, the coupling reaction is run in the presence of from about 0.8 to 3 equivalents, more preferably from about 1.0
15 to about 2.0 equivalents of ArX, based on the amount of nucleophile.

Compounds suitable as the base component of the method of the present invention include, for example, magnesium bicarbonate, potassium carbonate, cesium carbonate, potassium phosphate, sodium hydroxide and
20 potassium hydroxide. In a preferred embodiment, the base comprises one or more of sodium hydroxide and potassium hydroxide. In a preferred embodiment, the reaction mixture includes from about 1 to about 5 equivalents, more preferably from about 1.2 to about 4 equivalents, of base, based on the amount of nucleophile.

25

The base is generally added to the reaction mixture as solid or as an aqueous solution. The full amount of base to be used in the reaction may be added to the reaction mixture at one time or may be added to the reaction

- 13 -

mixture over time. In any case, it is preferred that the base be added after formation of the copper-ligand catalyst complex.

5 In a preferred embodiment, the reaction is conducted in the presence of from about 1 to about 80 percent by volume ("vol%") water, more preferably from about 10 to about 50 vol% water, based on the total volume of reaction mixture.

10 In one embodiment, the reaction mixture contains no organic solvent and in a preferred embodiment, the reaction mixture consists essentially of the reactants, copper catalyst, base and water.


15 In an alternative embodiment, the reaction mixture further comprises a solvent selected from aliphatic or aromatic hydrocarbon solvents such as pentane, hexane, benzene, xylene and toluene, ethers such as diethyl ether and t-butyl methyl ether, tetrahydrofuran, 1,4-dioxane and 1,2-dimethoxyethane.

20 In general, the coupling reaction is run under mild conditions that will not adversely affect the reactants, catalyst or product. In a preferred embodiment, the coupling reaction is run at a temperature of from about 25°C to about 300°C, more preferably from about 25°C to about 150°C.

25 In a preferred embodiment, the coupling reaction is run in an inert atmosphere, such as, for example, under an argon or nitrogen atmosphere.

- 14 -

Examples 1 and 2 and Comparative Examples C1 and C2: Synthesis of 5-chloro-1-(4-fluorophenyl)indole



5

In the reaction of Example 1, a reaction vessel composed of a 250 mL 3-necked round bottom flask equipped with a reflux condenser, overhead mechanical stirrer and Ar inlet was flushed with Ar for 30 mins. The reaction vessel was then charged with 4-bromofluorobenzene **3** (6.6 mL, 60 mmol) and H₂O (5 mL, purged with bubbling Ar for 1 h) and the overhead stirrer was engaged. 1,2-di(aminomethyl)cyclohexane **1** (426 mg, 3 mmol), CuI (114 mg, 0.60 mmol), and 5-chloroindole **2** (4.5g, 30 mmol) were added and reaction mixture was stirred for 5 minutes. KOH (5.05g, 90 mmol) was charged to the reaction vessel and the reaction flask was placed in a 120 °C oil bath. After 4 hours, 95% conversion was indicated by GC analysis of aliquots diluted with EtOAc. The oil bath was removed and hot reaction mixture was charged with H₂O (4 volumes), which led to a rapid precipitation of the organic product. Once reaction mixture had cooled to ambient

10

15

- 15 -

temperature, the solid product was collected by filtration and dried overnight on house vacuum. Yield: 7.03g (95%). GC Purity: 97.5%.

The reactions of Example 2 and Comparative Examples C1 and C2 were conducted under conditions analogous to those of Example 1, except
5 as indicated in TABLE I:

TABLE I

Example #	Water	Base
Ex. 1	5 mL	5.05 g KOH
Ex. 2	5 mL	14.88 g K ₃ PO ₄
C. Ex. C1	none	5.05 g KOH
C. Ex. C2	none	14.88 g K ₃ PO ₄

The percent conversion over time is given below in TABLE II for each of the
10 examples.

TABLE II


Example #	Conversion (%)		
	1 hour	2 hours	4 hours
Ex. 1	56	86	96
Ex. 2	42	66	90
C.Ex. C1	2	3.7	6
C. Ex. C2	54	65	83

The use of water/KOH offered comparable kinetics to using 2 eq ArBr
15 as solvent with 2 eq K₃PO₄, either with or without water, and provided the advantages of 1) a cheaper and more volume efficient base and 2) easier reaction agitation which should help scale up.


- 16 -

Claims:

1. A method for arylating a nucleophile, comprising reacting the nucleophile with a substrate aromatic compound, ArX, in the presence of a copper catalyst, a base and water, wherein Ar is aryl, heteroaryl or alkenyl, X is halo, sulfonate or phosphonate, the base comprises an alkaline earth carbonate, bicarbonate, hydroxide or phosphate, and the copper catalyst comprises a copper atom or ion and a ligand.
2. The method of claim 1, wherein the nucleophile comprises a HN-containing heterocycle or a HN-containing compound according to the formula $\text{HN}(\text{R}^1)\text{R}^2$, wherein R^1 is H, alkyl or aryl, and R^2 is a substituent according to the formula:



wherein R^3 is H, alkyl, aryl, heteroaryl, alkenyl, $-\text{OR}^5$ or $-\text{NR}^6_2$, and R^5 and R^6 are each independently alkyl, aryl, or a substituent according to the formula:




wherein R^7 is alkyl or aryl.

- 17 -

3. The method of claim 2, wherein the nucleophile comprises a HN-containing heterocycle selected from triazoles, pyrroles, pyrazoles, imidazoles, indoles, azaindoles, benzotriazoles, benzimidazoles, indazoles,
5 and carbazoles, each of which may be substituted or unsubstituted.

4. The method of claim 2, wherein the nucleophile comprises a HN-containing heterocycle comprises a monocyclic system according to the formula:
10



wherein n is 0 or an integer of from 1 to 3 and R¹⁰ is substituted alkyl, substituted N, or O.
15

5. The method of claim 4, wherein nucleophile comprises an amide, a carbamate, a urea, or a sulfonamide.

6. The method of claim 5, wherein nucleophile comprises benzamide, 4-aminobenzamide, cyclohexylamide, trans-cinnamamide, N-phenylacetamide, N-methylformamide, N-benzylformamide, N-cyclohexylformamide, N-phenyl-*tert*-butyl carbamate, N-methylimidazolidinone, or p-toluenesulfonamide.

7. The method of claim 1, wherein Ar comprises a phenyl ring which is, other than the X substituent, unsubstituted or is further substituted, in addition to the X substituent, on one or more carbons of the ring with one or
25


- 18 -

more substituent groups each independently selected from alkyl, alkoxy, alkenyl, alkynyl, aryl, heteroaryl, cyano, carbonyl, amino, amido or sulfonyl.

8. The method of claim 1, wherein X is a halo.

5


9. The method of claim 1, wherein X is a sulfonate group according to the formula:



10

wherein R⁸ is alkyl, aryl, fluoroalkyl, preferably trifluoromethyl, perfluoroalkyl.

10. The method of claim 1, wherein X is a phosphonate group according to the formula:



15

wherein each R⁹ is independently alkyl or aryl.

11. The method of claim 1, wherein ArX comprises 4-bromobenzonitrile, 4-N,N'-dimethyl-bromoaniline, 2-bromothiophene, 3-bromoquinoline, 1-nitro-2-iodobenzene, 4-chlorotoluene, 4-bromofluorobenzene, 2-bromoanisole, 4-iodoaniline, 3-bromoacetophenone, or 4-bromothioanisole.


20

- 19 -

12. The method of claim 1, wherein the copper atom or ion is derived from copper metal, from Cu_2O or from a copper salt selected from CuCl , CuBr , CuBr_2 and CuI .
- 5 13. The method of claim 1, wherein the ligand comprises a 1,2-diamine compound.
14. The method of claim 1, wherein the ligand comprises 1,2-di(aminomethyl)cyclohexane, $\text{N,N}'$ -dimethylethylenediamine, or 1-propyl-1,2-
10 $\text{N,N}'$ - dimethylethylenediamine.
15. The method of claim 1, wherein the copper catalyst is present in the reaction mixture as complex comprising the copper atom or ion and the ligand.
- 15 16. The method of claim 1, wherein the copper catalyst is present in an amount of from about 0.01 to about 10 mole%, based on the amount of limiting reactant.
- 20 17. The method of claim 1, wherein the reaction is conducted in the presence of from about 0.8 to 3 equivalents of ArX , based on the amount of nucleophile.
- 25 18. The method of claim 1, wherein the reaction is conducted in the presence of from about 1 to about 80 percent by volume water, based on the total volume of reaction mixture.


- 20 -

19. A method for arylating a HN-containing heterocycle, comprising reacting the HN-containing heterocycle with a substrate aromatic compound, ArX, according to the reaction scheme:




wherein Ar is aryl, heteroaryl or alkenyl,
 X is halo, sulfonate or phosphonate,
 the base comprises an alkaline earth carbonate, bicarbonate, hydroxide or
 10 phosphate, and
 the copper catalyst comprises a copper atom or ion and a ligand.

20. A method for arylating a HN-containing compound according to the formula $\text{HN}(\text{R}^1)\text{R}^2$, comprising reacting the HN-containing compound with a
 15 substrate aromatic compound, ArX, according to the reaction scheme:




wherein Ar is aryl, heteroaryl or alkenyl,
 X is halo, sulfonate or phosphonate
 20 R^1 is H, alkyl or aryl
 R^2 is according to the formula:



- 21 -

wherein R^3 is H, alkyl, aryl, heteroaryl, alkenyl, $-OR^5$ or $-NR^6_2$, and R^5 and R^6 are each independently alkyl, aryl, or



5

wherein R^7 is alkyl or aryl,

the base comprises an alkaline earth carbonate, bicarbonate, hydroxide or phosphate, and

10 the copper catalyst comprises a copper atom or ion and a ligand.