

(19)



(11)

EP 1 882 731 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

30.01.2008 Bulletin 2008/05

(51) Int Cl.:

C11D 3/06 (2006.01)

C11D 3/12 (2006.01)

C11D 3/386 (2006.01)

(21) Application number: **06115574.3**

(22) Date of filing: **16.06.2006**

(84) Designated Contracting States:

**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI
SK TR**

Designated Extension States:

AL BA HR MK YU

(71) Applicant: **The Procter and Gamble Company**
Cincinnati, Ohio 45202 (US)

(72) Inventors:

- **Lant, Neil**
Newcastle upon Tyne, NE3 5RP (GB)
- **Patterson, Steven**
Tyne & Wear, NE37 1RZ (GB)

(74) Representative: **Morelle, Evelyne Charlotte**
Isabelle

N.V. Procter & Gamble Services Company S.A.
Temselaan 100
1853 Strombeek-Bever (BE)

(54) **Detergent compositions**

(57) Detergent compositions containing an alkaline bacterial enzyme exhibiting endo-beta-1,4-glucanase activity (E.C. 3.2.1.4) and specific detergent formulations comprising less than 10 wt% zeolite and phosphate builder are described. Preferred formulations comprise sur-

factants selected from alkyl benzene sulphonates in combination with alkyl ethoxylated sulfates or MES or non-ionic surfactants.

EP 1 882 731 A1

Description

Technical field

5 [0001] The present invention relates to laundry detergent compositions and in particular to detergents comprising an alkaline bacterial enzyme exhibiting endo-beta-1,4-glucanase activity (E.C. 3.2.1.4)

Background of the Invention and Prior Art

10 [0002] Cellulase enzymes have been used in detergent compositions for many years now for their known benefits of depilling, softness and colour care. However, the use of most of cellulases has been limited because of the negative impact that cellulase may have on the tensile strength of the fabrics' fibers by hydrolysing crystalline cellulose. Recently, cellulases with a high specificity towards amorphous cellulose have been developed to exploit the cleaning potential of cellulases while avoiding the negative tensile strength loss. Especially alkaline endo-glucanases have been developed
15 to suit better the use in alkaline detergent conditions.

[0003] For example, Novozymes in WO02/099091 discloses a novel enzyme exhibiting endo-beta-glucanase activity (EC 3.2.1.4) endogenous to the strain *Bacillus sp.*, DSM 12648; for use in detergent and textile applications. Novozymes further describes in WO04/053039 detergent compositions comprising an anti-redeposition endo-glucanase and its combination with certain cellulases having increased stability towards anionic surfactant and/or further specific enzymes.
20 Kao's EP 265 832 describes novel alkaline cellulase K, CMCase I and CMCase II obtained by isolation from a culture product of *Bacillus sp* KSM-635. Kao further describes in EP 1 350 843, alkaline cellulase which acts favourably in an alkaline environment and can be mass produced readily because of having high secretion capacity or having enhanced specific activity.

[0004] The problem facing the present inventors was how to maximise performance from this new generation of
25 cellulases. The present inventors found that whilst a small benefit could be achieved formulating such enzymes according to present day detergent formulations by simply replacing existing cellulase enzymes with the new generation of enzymes, a considerable improvement in performance was found by formulating the detergent compositions in a different way and even reducing the levels of some conventional detergent ingredients. Indeed, it has been surprisingly found that the use of very low levels of builder, or even complete absence of inorganic builder, enhances the cleaning performance of the
30 bacterial alkaline cellulase. Without wishing to be bound by theory, it is believed that (i) inorganic builders such as STPP, zeolites and silicates interact with hardness to form insoluble materials that deposit on fabrics and potentially interfere with the catalytic mechanism of the cellulase and that (ii) hardness ions such as Ca²⁺ and Mg²⁺ stabilise the enzyme in solution, and promote deposition of enzyme onto fabric surfaces. Therefore, the removal or reduction in builder is thus expected to increase free hardness levels leading to increased enzyme stability and surface deposition while
35 increasing its activity on fabrics through reduced levels of encrustation.

Definition of the Invention

[0005] In accordance with the present invention there is provided a detergent composition comprising an alkaline
40 bacterial enzyme exhibiting endo-beta-1,4-glucanase activity (E.C. 3.2.1.4) and comprising less than 10 wt% aluminosilicate (anhydrous basis) builder and less than 10 wt% phosphate builder, the composition having a reserve alkalinity of greater than 4.

[0006] In a preferred aspect of the invention, the detergent compositions of the invention comprise less than 10 wt% builders selected from aluminosilicate (zeolite) builder and/or phosphate builder. In a further preferred aspect of the
45 invention, the compositions comprise less than 8wt% zeolite, or even less than 4wt% and less than 8 wt% phosphate builder or even less than 4wt%.

Sequence listings

50 [0007] SEQ ID NO: 1 shows the amino acid sequence of an endoglucanase from *Bacillus sp.* AA349
[0008] SEQ ID NO: 2 shows the amino acid sequence of an endoglucanase from *Bacillus sp* KSM-S237

Detailed Description of the Invention

55 SUITABLE ENDOGLUCANASE

[0009] The endoglucanase to be incorporated into the detergent composition of the present invention is one or more bacterial alkaline enzyme(s) exhibiting endo-beta-1,4-glucanase activity (E.C. 3.2.1.4) and is typically comprised at a

level of from 0.00005% to 0.15%, from 0.0002% to 0.02%, or even from 0.0005% to 0.01% by weight of pure enzyme of one or more endoglucanase.

[0010] Preferably, the endoglucanase is a bacterial polypeptide endogenous to a member of the genus *Bacillus*.

[0011] More preferably, the alkaline enzyme exhibiting endo-beta-1,4-glucanase activity (E.C. 3.2.1.4), is a polypeptide containing (i) at least one family 17 carbohydrate binding module (Family 17 CBM) and/or (ii) at least one family 28 carbohydrate binding module (Family 28 CBM). Please refer for example to: Current Opinion in Structural Biology, 2001, 593-600 by Y. Bourne and B. Henrissat in their article entitled: "Glycoside hydrolases and glycosyltransferases: families and functional modules" for the definition and classification of CBMs. Please refer further to Biochemical Journal, 2002, v361, 35-40 by A.B. Boraston et al in their article entitled: "Identification and glucan-binding properties of a new carbohydrate-binding module family" for the properties of the family 17 and 28 CBM's.

[0012] In a more preferred embodiment, said enzyme comprises a polypeptide (or variant thereof) endogenous to one of the following *Bacillus species*:

Bacillus sp.	As described in:
AA349 (DSM 12648)	WO 2002/099091A (Novozymes) p2, line 25 WO 2004/053039A (Novozymes) p3, line19
KSM S237	EP 1350843A (Kao) p3, line 18
1139	EP 1350843A (Kao) p3, line 22
KSM 64	EP 1350843A (Kao) p3, line 24
KSM N131	EP 1350843A (Kao) p3, line 25
KSM 635, FERM BP 1485	EP 265 832A (Kao) p7, line 45
KSM 534, FERM BP 1508	EP 0271044 A (Kao) p9, line 21
KSM 539, FERM BP 1509	EP 0271044 A (Kao) p9, line 22
KSM 577, FERM BP 1510	EP 0271044 A (Kao) p9, line 22
KSM 521, FERM BP 1507	EP 0271044 A (Kao) p9, line 19
KSM 580, FERM BP 1511	EP 0271044 A (Kao) p9, line 20
KSM 588, FERM BP 1513	EP 0271044 A (Kao) p9, line 23
KSM 597, FERM BP 1514	EP 0271044 A (Kao) p9, line 24
KSM 522, FERM BP 1512	EP 0271044 A (Kao) p9, line 20
KSM 3445, FERM BP 1506	EP 0271044 A (Kao) p10, line 3
KSM 425. FERM BP 1505	EP 0271044 A (Kao) p10, line 3

[0013] Suitable endoglucanases for the compositions of the present invention are:

1) An enzyme exhibiting endo-beta-1,4-glucanase activity (E.C. 3.2.1.4), which has a sequence of at least 90%, preferably 94%, more preferably 97% and even more preferably 99%, 100% identity to the amino acid sequence of position 1 to position 773 of SEQ ID NO:1 (Corresponding to SEQ ID NO:2 in WO02/099091); or a fragment thereof has endo-beta-1,4-glucanase activity, when identity is determined by GAP provided in the GCG program using a GAP creation penalty of 3.0 and GAP extension penalty of 0.1. The enzyme and the corresponding method of production is described extensively in the patent application WO02/099091 published by Novozymes A/S on December 12, 2002. Please refer to the detailed description pages 4 to 17 and to the examples page 20 to page 26. One of such enzyme is commercially available under the tradename Celluclean™ by Novozymes A/S. GCG refers to the sequence analysis software package provided by Accelrys, San Diego, CA, USA. This incorporates a program called GAP which uses the algorithm of Needleman and Wunsch to find the alignment of two complete sequences that maximises the number of matches and minimises the number of gaps.

2) Also suitable are the alkaline endoglucanase enzymes described in EP 1 350 843 published by Kao corporation on October 8, 2003. Please refer to the detailed description [0011] to [0039] and examples 1 to 4 [0067] to [0077] for a detailed description of the enzymes and its production. The alkaline cellulase variants are obtained by substituting the amino acid residue of a cellulase having an amino acid sequence exhibiting at least 90%, preferably 95%, more preferably 98% and even 100% identity with the amino acid sequence represented by SEQ. ID NO:2 (Corresponding to SEQ. ID NO:1 in EP 1 350 843 on pages 11-13) at (a) position 10, (b) position 16, (c) position 22, (d) position 33, (e) position 39,

(f) position 76, (g) position 109, (h) position 242, (i) position 263, (j) position 308, (k) position 462, (l) position 466, (m) position 468, (n) position 552, (o) position 564, or (p) position 608 in SEQ ID NO:2 or at a position corresponding thereto with another amino acid residue

Examples of the "alkaline cellulase having the amino acid sequence represented by SEQ. ID NO:2" include Egl-237 [derived from *Bacillus* sp. strain KSM-S237 (FERM BP-7875), Hakamada, et al., Biosci. Biotechnol. Biochem., 64, 2281-2289, 2000]. Examples of the "alkaline cellulase having an amino acid sequence exhibiting at least 90% homology with the amino acid sequence represented by SEQ. ID NO:2" include alkaline cellulases having an amino acid sequence exhibiting preferably at least 95% homology, more preferably at least 98% homology, with the amino acid sequence represented by SEQ. ID NO:2. Specific examples include alkaline cellulase derived from *Bacillus* sp. strain 1139 (Egl-1139) (Fukumori, et al., J. Gen. Microbiol., 132, 2329-2335) (91.4% homology), alkaline cellulases derived from *Bacillus* sp. strain KSM-64 (Egl-64) (Sumitomo, et al., Biosci. Biotechnol. Biochem., 56, 872-877, 1992) (homology: 91.9%), and cellulase derived from *Bacillus* sp. strain KSM-N131 (Egl-N131b) (Japanese Patent Application No. 2000-47237) (homology: 95.0%).

The amino acid is preferably substituted by: glutamine, alanine, proline or methionine, especially glutamine is preferred at position (a), asparagine or arginine, especially asparagine is preferred at position (b), proline is preferred at position (c), histidine is preferred at position (d), alanine, threonine or tyrosine, especially alanine is preferred at position (e), histidine, methionine, valine, threonine or alanine, especially histidine is preferred at position (f), isoleucine, leucine, serine or valine, especially isoleucine is preferred at position (g), alanine, phenylalanine, valine, serine, aspartic acid, glutamic acid, leucine, isoleucine, tyrosine, threonine, methionine or glycine, especially alanine, phenylalanine or serine is preferred at position (h), isoleucine, leucine, proline or valine, especially isoleucine is preferred at position (i), alanine, serine, glycine or valine, especially alanine is preferred at position (j), threonine, leucine, phenylalanine or arginine, especially threonine is preferred at position (k), leucine, alanine or serine, especially leucine is preferred at position (l), alanine, aspartic acid, glycine or lysine, especially alanine is preferred at position (m), methionine is preferred at position (n), valine, threonine or leucine, especially valine is preferred at position (o) and isoleucine or arginine, especially isoleucine is preferred at position (p).

The "amino acid residue at a position corresponding thereto" can be identified by comparing amino acid sequences by using known algorithm, for example, that of Lipman-Pearson's method, and giving a maximum similarity score to the multiple regions of similarity in the amino acid sequence of each alkaline cellulase. The position of the homologous amino acid residue in the sequence of each cellulase can be determined, irrespective of insertion or depletion existing in the amino acid sequence, by aligning the amino acid sequence of the cellulase in such manner (Fig. 1 of EP 1 350 843). It is presumed that the homologous position exists at the three-dimensionally same position and it brings about similar effects with regard to a specific function of the target cellulase.

With regard to another alkaline cellulase having an amino acid sequence exhibiting at least 90% homology with SEQ. ID NO:1, specific examples of the positions corresponding to (a) position 10, (b), position 16, (c) position 22, (d) position 33, (e) position 39, (f) position 76, (g) position 109, (h) position 242, (i) position 263, (j) position 308, (k) position 462, (l) position 466, (m) position 468, (n) position 552, (o) position 564 and (p) position 608 of the alkaline cellulase (Egl-237) represented by SEQ. ID NO: 2 and amino acid residues at these positions will be shown below:

	Egl-237	Egl-1139	Egl-64	Egl-N131b
(a)	10Leu	10Leu	10Leu	10Leu
(b)	16Ile	16Ile	16Ile	Nothing corresponding thereto
(c)	22Ser	22Ser	22Ser	Nothing corresponding thereto
(d)	33Asn	33Asn	33Asn	19Asn
(e)	39Phe	39Phe	39Phe	25Phe
(f)	76Ile	76Ile	76Ile	62Ile
(g)	109Met	109Met	109Met	95Met
(h)	242Gln	242Gln	242Gln	228Gln
(i)	263Phe	263Phe	263Phe	249Phe
(j)	308Thr	308Thr	308Thr	294Thr
(k)	462Asn	461Asn	461Asn	448Asn
(l)	466Lys	465Lys	465Lys	452Lys
(m)	468Val	467Val	467Val	454Val

EP 1 882 731 A1

(continued)

	Egl-237	Egl-1139	Egl-64	Egl-N131b
(n)	552Ile	550Ile	550Ile	538Ile
(o)	564Ile	562Ile	562Ile	550Ile
(p)	608Ser	606Ser	606Ser	594Ser

3) Also suitable are the alkaline endoglucanase enzymes described in EP 265 832 published by Kao on May 4, 1988. Please refer to the description page 4, line 35 to page 12, line 22 and examples 1 and 2 on page 19 for a detailed description of the enzymes and its production: The alkaline cellulase K has the following physical and chemical properties:

- (1) Activity: Having a C_x enzymatic activity of acting on carboxymethyl cellulose along with a weak C₁ enzymatic activity and a weak beta-glucosidase activity;
- (2) Specificity on Substrates: Acting on carboxymethyl cellulose(CMC), crystalline cellulose, Avicell, cellobiose, and p-nitrophenyl cellobioside(PNPC);
- (3) Having a working pH in the range of 4 to 12 and an optimum pH in the range of 9 to 10;
- (4) Having stable pH values of 4.5 to 10.5 and 6.8 to 10 when allowed to stand at 40°C for 10 minutes and 30 minutes, respectively;
- (5) Working in a wide temperature range of from 10 to 65°C with an optimum temperature being recognized at about 40°C;
- (6) Influences of chelating agents: The activity not impeded with ethylenediamine tetraacetic acid (EDTA), ethyleneglycol-bis-(β-aminoethylether) N,N,N',N''-tetraacetic acid (EGTA), N,N-bis(carboxymethyl)glycine (nitrilotriacetic acid) (NTA), sodium tripolyphosphate (STPP) and zeolite;
- (7) Influences of surface active agents: Undergoing little inhibition of activity by means of surface active agents such as sodium linear alkylbenzenesulfonates (LAS), sodium alkylsulfates (AS), sodium polyoxyethylene alkylsulfates (ES), sodium alphaolefinsulfonates (AOS), sodium alpha-sulfonated aliphatic acid esters (alpha-SFE), sodium alkyl-sulfonates (SAS), polyoxyethylene secondary alkyl ethers, fatty acid salts (sodium salts), and dimethyldialkylammonium chloride;
- (8) Having a strong resistance to proteinases; and
- (9) Molecular weight (determined by gel chromatography): Having a maximum peak at 180,000 ± 10,000.

Preferably such enzyme is obtained by isolation from a culture product of Bacillus sp KSM-635.

One or more of such enzyme is commercially available by the Kao Corporation: e.g. the cellulase preparation Eg-X known as KAC® being a mixture of E-H and E-L both from Bacillus sp. KSM-635 bacterium. Cellulases E-H and E-L have been described in S. Ito, Extremophiles, 1997, v1, 61-66 and in S. Ito et al, Agric Biol Chem, 1989, v53, 1275-1278.

4) Finally the alkaline bacterial endoglucanases described in EP 271 004 published by Kao on June 15, 1988 are also suitable for the purpose of the present invention. Please refer to the detailed description page 9, line 15 to page 23, line 17 and page 31, line 1 to page 33, line 17 for a detailed description of the enzymes and its production. Those are:

Alkaline Cellulase K-534 from KSM 534, FERM BP 1508,
Alkaline Cellulase K-539 from KSM 539, FERM BP 1509,
Alkaline Cellulase K-577 from KSM 577, FERM BP 1510,
Alkaline Cellulase K-521 from KSM 521, FERM BP 1507,
Alkaline Cellulase K-580 from KSM 580, FERM BP 1511,
Alkaline Cellulase K-588 from KSM 588, FERM BP 1513,
Alkaline Cellulase K-597 from KSM 597, FERM BP 1514,
Alkaline Cellulase K-522 from KSM 522, FERM BP 1512,
Alkaline Cellulase E-II from KSM 522, FERM BP 1512,

Alkaline Cellulase E-III from KSM 522, FERM BP 1512.
 Alkaline Cellulase K-344 from KSM 344, FERM BP 1506, and
 Alkaline Cellulase K-425 from KSM 425, FERM BP 1505.

BUILDERS

[0014] Commercially available laundry detergents comprise strong inorganic builder, with either phosphate builder typically sodium tripolyphosphate (STPP), or zeolite typically sodium aluminosilicate builder, being used as the predominant strong builder. Generally such strong builders are present at relatively high levels such as 15 to 20 wt% or even higher, for example even up to 40 wt%. In accordance with the present invention, the amount of strong builder selected from phosphate and/or zeolite builder is no greater than 10 wt% based on the total weight of the detergent composition, preferably below 8 wt%, or even below 5 or 4 or 3 or 2 or 1 wt%.

[0015] Thus, the compositions of the invention may comprise from 0 wt% to 10wt% zeolite builder, and 0 wt% to 10 wt% phosphate builder, the total amount of phosphate and/or zeolite not exceeding 10 wt%, and preferably being below 10 wt% as described above. Preferably the compositions of the invention comprise from 0 wt% to 8 wt%, or from 0 wt% to 5 or 4 wt%, or from 0 wt% to 3 or even less than 2 wt% zeolite builder. It may even be preferred for the composition to be essentially free from zeolite builder. By essentially free from zeolite builder it is typically meant that the composition comprises no deliberately added zeolite builder. This is especially preferred if it is desirable for the composition to be very highly soluble, to minimise the amount of water-insoluble residues (for example, which may deposit on fabric surfaces), and also when it is highly desirable to have transparent wash liquor. Zeolite builders include zeolite A, zeolite X, zeolite P and zeolite MAP.

[0016] The compositions of the invention may comprise from 0 wt% to 10 wt% phosphate builder. The composition preferably comprises from 0 wt% to 8 wt%, or from 0 wt% to 5 or 4 wt%, or from 0 wt% to 3 or even 2 wt% phosphate builder. It may even be preferred for the composition to be essentially free from phosphate builder. By essentially free from phosphate builder it is typically meant that the composition comprises no deliberately added phosphate builder. This is especially preferred if it is desirable for the composition to have a very good environmental profile. Phosphate builders include sodium tripolyphosphate.

[0017] In a further preferred aspect of the invention, the total level of weak builders selected from layered silicate (SKS-6), citric acid, citrate salts and nitrilo triacetic acid or salt thereof is below 15 wt%, more preferably below 8 wt%, more preferably below 4 wt% or even below 3 or 2 wt% based on the total weight of the detergent composition. Typically the level of each of layered silicate, citric acid, citrate salts and nitrilo triacetic acid or salt thereof will be below 10 wt% or even below 5 wt% or wt% based on the total weight of the composition.

[0018] Although builders bring several benefits to the formulator, their main role is to sequester divalent metal ions (such as calcium and magnesium ions) from the wash solution that would otherwise interact negatively with the surfactant system. Builders are also effective at removing metal ions and inorganic soils from the fabric surface too, leading to improved removal of particulate and beverage stains. It would therefore be expected that reduction of their levels would negatively impact on cleaning performance and therefore, preparation of detergent compositions that are effective with the claimed reduced levels of phosphate and zeolite builders is surprising.

Reserve Alkalinity

[0019] As used herein, the term "reserve alkalinity" is a measure of the buffering capacity of the detergent composition (g/NaOH/100g detergent composition) determined by titrating a 1% (w/v) solution of detergent composition with hydrochloric acid to pH 7.5 i.e in order to calculate Reserve Alkalinity as defined herein:

Reserve Alkalinity (to pH 7.5) as % alkali in g NaOH/100 g product =

$$\frac{T \times M \times 40 \times \text{Vol}}{10 \times \text{Wt} \times \text{Aliquot}}$$

T = titre (ml) to pH 7.5

M = Molarity of HCl = 0.2

40 = Molecular weight of NaOH

Vol = Total volume (ie. 1000 ml)

Wt = Weight of product (10 g)

Aliquot = (100 ml)

[0020] Obtain a 10g sample accurately weighed to two decimal places, of fully formulated detergent composition. The sample should be obtained using a Pascall sampler in a dust cabinet. Add the 10g sample to a plastic beaker and add 200 ml of carbon dioxide-free deionised water. Agitate using a magnetic stirrer on a stirring plate at 150 rpm until fully dissolved and for at least 15 minutes. Transfer the contents of the beaker to a 1 litre volumetric flask and make up to 1 litre with deionised water. Mix well and take a 100 mls $\pm$ 1 ml aliquot using a 100 mls pipette immediately. Measure and record the pH and temperature of the sample using a pH meter capable of reading to +0.01pH units, with stirring, ensuring temperature is 21°C $\pm$ 2°C. Titrate whilst stirring with 0.2M hydrochloric acid until pH measures exactly 7.5. Note the millilitres of hydrochloric acid used. Take the average titre of three identical repeats. Carry out the calculation described above to calculate RA to pH 7.5.

The RA will be greater than 4 and preferably greater than 6 and most preferably greater than 7.5 or even greater than 8 or 8.5 or higher.

[0021] It has been found that a robust alkalinity system is beneficial in the detergent compositions of the invention. Adequate reserve alkalinity may be provided, for example, by one or more of alkali metal silicates (excluding crystalline layered silicate), typically amorphous silicate salts, generally 1.2 to 2.2 ratio sodium salts, alkali metal typically sodium carbonate, bicarbonate and/or sesquicarbonates. STPP and persalts such as perborates and percarbonates also contribute to alkalinity. Buffering is necessary to maintain an alkaline pH during the wash process counteracting the acidity of soils.

[0022] The detergent composition preferably comprises from 0 wt% to 50 wt% silicate salt, more usually 5 to 30 wt% silicate salt, or 7 to 20 wt% silicate salt, usually sodium silicate.

[0023] In order to provide the desired reserve alkalinity the detergent compositions of the invention may comprise a carbonate salt, typically from 1 wt% to 70 wt%, or from 5 wt% to 50 wt% or from 10 wt% to 30 wt% carbonate salt. Preferred carbonate salts are sodium carbonate and/or sodium bicarbonate and/or sodium sesquicarbonate. The carbonate salt may be incorporated into the detergent composition wholly or partially via a mixed salt such as Burkeite. A highly preferred carbonate salt is sodium carbonate. Preferably, the composition may comprise from 5 wt% to 50 wt% sodium carbonate, or from 10 to 40 wt% or even 15 to 35 wt% sodium carbonate. It may also be desired for the composition to comprise from 1wt% to 20 wt% sodium bicarbonate, or even 2 to 10 or 8 wt%.

[0024] If zeolite is present, it may be desired for the weight ratio of sodium carbonate and/or sodium silicate to zeolite builder to be at least 5:1, preferably at least 10:1, or at least 15:1, or at least 20:1 or even at least 25:1


[0025] The carbonate salt, or at least part thereof, is typically in particulate form, typically having a weight average particle size in the range of from 200 to 500 micrometers. However, it may be preferred for the carbonate salt, or at least part thereof, to be in micronised particulate form, typically having a weight average particle size in the range of from 4 to 40 micrometers; this is especially preferred when the carbonate salt, or at least part thereof, is in the form of a co-particulate admixture with a deterative surfactant, such as an alkoxyated anionic deterative surfactant.

[0026] In order to provide the required reserve alkalinity, preferably the levels of carbonate and/or silicate salts, typically sodium carbonate and sodium silicate will be from 10 to 70 wt%, or from 10 or even 15 to 50 wt% based on the total weight of the composition.

FURTHER INGREDIENTS

[0027] The compositions of the present invention may comprise further ingredients as described below. Preferred are the chelants and especially hydroxyethane-dimethylene-phosphonic acid (HEDP), 2-phosphonobutane-1,2,4-tricarboxylic acid (PBTC) and 4,5-dihydroxy-m-benzenedisulfonic acid, disodium salt (Tiron®). Indeed it is believed that the combination of the endoglucanase within the low builder system of the present invention with these chelants provide even better stain removal, cleaning and whiteness.

[0028] Another preferred ingredient is a fluorescent whitening agent, especially the following:




wherein R1 and R2, together with the nitrogen atom linking them, form an unsubstituted or C1-C4 alkyl-substituted morpholino, piperidine or pyrrolidine ring. Indeed it is believed that the combination of the endoglucanase within the low builder system of the present invention with these fluorescent whitening agents provide even better cleaning and whiteness.

Surfactant


[0029] A highly preferred adjunct component of the compositions of the invention is a surfactant. Preferably, the detergent composition comprises one or more surfactants. Typically, the detergent composition comprises (by weight of the composition) from 0% to 50%, preferably from 5% and more preferably from 10 or even 15 wt% to 40%, or to 30%, or to 20% one or more surfactants. Preferred surfactants are anionic surfactants, non-ionic surfactants, cationic surfactants, zwitterionic surfactants, amphoteric surfactants, cationic surfactants and mixtures thereof.

Anionic surfactants

[0030] Suitable anionic surfactants typically comprise one or more moieties selected from the group consisting of carbonate, phosphate, phosphonate, sulphate, sulphonate, carboxylate and mixtures thereof. The anionic surfactant may be one or mixtures of more than one of C₈₋₁₈ alkyl sulphates and C₈₋₁₈ alkyl sulphonates. Suitable anionic surfactants incorporated alone or in mixtures in the compositions of the invention are also the C₈₋₁₈ alkyl sulphates and/or C₈₋₁₈ alkyl sulphonates optionally condensed with from 1 to 9 moles of C₁₋₄ alkylene oxide per mole of C₈₋₁₈ alkyl sulphate and/or C₈₋₁₈ alkyl sulphonate. The alkyl chain of the C₈₋₁₈ alkyl sulphates and/or C₈₋₁₈ alkyl sulphonates may be linear or branched, preferred branched alkyl chains comprise one or more branched moieties that are C₁₋₆ alkyl groups. More particularly, suitable anionic surfactants include the C₁₀₋₂₀ primary, branched-chain, linear-chain and random-chain alkyl sulphates (AS), typically having the following formula:



wherein, M is hydrogen or a cation which provides charge neutrality, preferred cations are sodium and ammonium cations, wherein x is an integer of at least 7, preferably at least 9; C₁₀₋₁₈ secondary (2,3) alkyl sulphates, typically having the following formulae:



wherein, M is hydrogen or a cation which provides charge neutrality, preferred cations include sodium and ammonium cations, wherein x is an integer of at least 7, preferably at least 9, y is an integer of at least 8, preferably at least 9; C₁₀₋₁₈ alkyl alkoxy carboxylates; mid-chain branched alkyl sulphates as described in more detail in US 6,020,303 and US 6,060,443; modified alkylbenzene sulphonate (MLAS) as described in more detail in WO 99/05243, WO 99/05242, WO 99/05244, WO 99/05082, WO 99/05084, WO 99/05241, WO 99/07656, WO 00/23549, WO 00/23548 and mixtures thereof.

[0031] Preferred anionic surfactants are C₈₋₁₈ alkyl benzene sulphates and/or C₈₋₁₈ alkyl benzene sulphonates. The alkyl chain of the C₈₋₁₈ alkyl benzene sulphates and/or C₈₋₁₈ alkyl benzene sulphonates may be linear or branched, preferred branched alkyl chains comprise one or more branched moieties that are C₁₋₆ alkyl groups.

[0032] Other preferred anionic surfactants are selected from the group consisting of: C₈₋₁₈ alkenyl sulphates, C₈₋₁₈ alkenyl sulphonates, C₈₋₁₈ alkenyl benzene sulphates, C₈₋₁₈ alkenyl benzene sulphonates, C₈₋₁₈ alkyl di-methyl benzene sulphate, C₈₋₁₈ alkyl di-methyl benzene sulphonate, fatty acid ester sulphonates, di-alkyl sulphosuccinates, and combinations thereof. Other useful anionic surfactants herein include the esters of alpha-sulfonated fatty acids, typically containing from 6 to 20 carbon atoms in the fatty acid group and from 1 to 10 carbon atoms in the ester group; 2-acyloxy-alkane-1-sulfonic acid and salts thereof, typically containing from about 2 to 9 carbon atoms in the acyl group and from about 9 to 23 carbon atoms in the alkane moiety; alpha-olefin sulfonates (AOS), typically containing from about 12 to 24 carbon atoms; and beta-alkoxy alkane sulfonates, typically containing from about 1 to 3 carbon atoms in the alkyl group and from about 8 to 20 carbon atoms in the alkane moiety. Also useful are the sulphonation products of fatty acid esters containing an alkyl group typically with from 10 to 20 carbon atoms. Preferred are C₁₋₄, most preferably methyl ester sulphonates. Preferred are C₁₆₋₁₈ methyl ester sulphonates (MES).

[0033] The anionic surfactants may be present in the salt form. For example, the anionic surfactant(s) may be an alkali metal salt of any of the above. Preferred alkali metals are sodium, potassium and mixtures thereof.

[0034] Preferred anionic deterative surfactants are selected from the group consisting of: linear or branched, substituted or unsubstituted, C₁₂₋₁₈ alkyl sulphates; linear or branched, substituted or unsubstituted, C₁₀₋₁₃ alkylbenzene sulphonates, preferably linear C₁₀₋₁₃ alkylbenzene sulphonates; and mixtures thereof. Highly preferred are linear C₁₀₋₁₃ alkylbenzene sulphonates. Highly preferred are linear C₁₀₋₁₃ alkylbenzene sulphonates that are obtainable, preferably obtained, by sulphonating commercially available linear alkyl benzenes (LAB); suitable LAB include low 2-phenyl LAB, such as those supplied by Sasol under the tradename Isochem® or those supplied by Petresa under the tradename Petrelab®, other suitable LAB include high 2-phenyl LAB, such as those supplied by Sasol under the tradename Hyblene®.

[0035] It may be preferred for the anionic deterative surfactant to be structurally modified in such a manner as to cause the anionic deterative surfactant to be more calcium tolerant and less likely to precipitate out of the wash liquor in the presence of free calcium ions. This structural modification could be the introduction of a methyl or ethyl moiety in the vicinity of the head group of the anionic deterative surfactant, as this can lead to a more calcium tolerant anionic deterative surfactant due to steric hindrance of the head group, which may reduce the affinity of the anionic deterative surfactant for complexing with free calcium cations in such a manner as to cause precipitation out of solution. Other structural modifications include the introduction of functional moieties, such as an amine moiety, in the alkyl chain of the anionic deterative surfactant; this can lead to a more calcium tolerant anionic deterative surfactant because the presence of a functional group in the alkyl chain of an anionic deterative surfactant may minimise the undesirable physicochemical property of the anionic deterative surfactant to form a smooth crystal structure in the presence of free calcium ions in the wash liquor. This may reduce the tendency of the anionic deterative surfactant to precipitate out of solution.

Alkoxylated anionic surfactants

[0036] The composition may comprise an alkoxylated anionic surfactant. Where present such a surfactant will generally be present in amounts from 0.1 wt% to 40 wt%, generally 0.1 to 10 wt% based on the detergent composition as a whole. It may be preferred for the composition to comprise from 3wt% to 5wt% alkoxylated anionic deterative surfactant, or it may be preferred for the composition to comprise from 1wt% to 3wt% alkoxylated anionic deterative surfactant.

[0037] Preferably, the alkoxylated anionic deterative surfactant is a linear or branched, substituted or unsubstituted C₁₂₋₁₈ alkyl alkoxylated sulphate having an average degree of alkoxylation of from 1 to 30, preferably from 1 to 10. Preferably, the alkoxylated anionic deterative surfactant is a linear or branched, substituted or unsubstituted C₁₂₋₁₈ alkyl ethoxylated sulphate having an average degree of ethoxylation of from 1 to 10. Most preferably, the alkoxylated anionic deterative surfactant is a linear unsubstituted C₁₂₋₁₈ alkyl ethoxylated sulphate having an average degree of ethoxylation of from 3 to 7.

[0038] The alkoxylated anionic deterative surfactant may also increase the non-alkoxylated anionic deterative surfactant activity by making the non-alkoxylated anionic deterative surfactant less likely to precipitate out of solution in the presence of free calcium cations. Preferably, the weight ratio of non-alkoxylated anionic deterative surfactant to alkoxylated anionic deterative surfactant is less than 5:1, or less than 3:1, or less than 1.7:1, or even less than 1.5:1. This ratio gives optimal whiteness maintenance performance combined with a good hardness tolerency profile and a good sudsing profile. However, it may be preferred that the weight ratio of non-alkoxylated anionic deterative surfactant to alkoxylated anionic deterative surfactant is greater than 5:1, or greater than 6:1, or greater than 7:1, or even greater than 10:1. This ratio gives optimal greasy soil cleaning performance combined with a good hardness tolerency profile, and a good sudsing profile.

Suitable alkoxylated anionic deterative surfactants are: Texapan LEST™ by Cognis; Cosmacol AEST™ by Sasol; BES151™ by Stephan; Empicol ESC70/U™; and mixtures thereof.

Non-ionic deterative surfactant

[0039] The compositions of the invention may comprise non-ionic surfactant. Where present it is generally present in amounts of from 0.5wt% to 20, more typically 0.5 to 10 wt% based on the total weight of the composition. The composition may comprise from 1wt% to 7wt% or from 2wt% to 4wt% non-ionic deterative surfactant. The inclusion of non-ionic deterative surfactant in the composition helps to provide a good overall cleaning profile, especially when laundering at high temperatures such as 60°C or higher.

[0040] The non-ionic deterative surfactant can be selected from the group consisting of: C_{12-C18} alkyl ethoxylates, such as, NEODOL® non-ionic surfactants from Shell; C_{6-C12} alkyl phenol alkoxylates wherein the alkoxylate units are ethyleneoxy units, propyleneoxy units or a mixture thereof; C_{12-C18} alcohol and C_{6-C12} alkyl phenol condensates with ethylene oxide/propylene oxide block polymers such as Pluronic® from BASF; C_{14-C22} mid-chain branched alcohols, BA, as described in more detail in US 6,150,322; C_{14-C22} mid-chain branched alkyl alkoxylates, BAE_x, wherein x = from 1 to 30, as described in more detail in US 6,153,577, US 6,020,303 and US 6,093,856; alkylpolysaccharides as described


in more detail in US 4,565,647, specifically alkylpolyglycosides as described in more detail in US 4,483,780 and US 4,483,779; polyhydroxy fatty acid amides as described in more detail in US 5,332,528, WO 92/06162, WO 93/19146, WO 93/19038, and WO 94/09099; ether capped poly(oxyalkylated) alcohol surfactants as described in more detail in US 6,482,994 and WO 01/42408; and mixtures thereof.

[0041] The non-ionic deterative surfactant could be an alkyl polyglucoside and/or an alkyl alkoxyated alcohol. Preferably the non-ionic deterative surfactant is a linear or branched, substituted or unsubstituted C₈₋₁₈ alkyl ethoxylated alcohol having an average degree of ethoxylation of from 1 to 50, more preferably from 3 to 40. Non-ionic surfactants having a degree of ethoxylation from 3 to 9 may be especially useful either. Nonionic surfactants having an HLB value of from 13 to 25, such as C₈₋₁₈ alkyl ethoxylated alcohols having an average degree of ethoxylation from 15 to 50, or even from 20 to 50 may also be preferred non-ionic surfactants in the compositions of the invention. Examples of these latter non-ionic surfactants are Lutensol AO30 and similar materials disclosed in WO04/041982. These may be beneficial as they have good lime soap dispersant properties.

[0042] The non-ionic deterative surfactant not only provides additional soil cleaning performance but may also increase the anionic deterative surfactant activity by making the anionic deterative surfactant less likely to precipitate out of solution in the presence of free calcium cations. Preferably, the weight ratio of non-alkoxyated anionic deterative surfactant to non-ionic deterative surfactant is in the range of less than 8:1, or less than 7:1, or less than 6:1 or less than 5:1, preferably from 1:1 to 5:1, or from 2:1 to 5:1, or even from 3:1 to 4:1.

Cationic deterative surfactant

[0043] In one aspect of the invention, the detergent compositions are free of cationic surfactant. However, the composition optionally may comprise from 0.1 wt% to 10 or 5wt% cationic deterative surfactant. When present however, preferably the composition comprises from 0.5wt% to 3wt%, or from 1% to 3wt%, or even from 1wt% to 2wt% cationic deterative surfactant. This is the optimal level of cationic deterative surfactant to provide good cleaning. Suitable cationic deterative surfactants are alkyl pyridinium compounds, alkyl quaternary ammonium compounds, alkyl quaternary phosphonium compounds, and alkyl ternary sulphonium compounds. The cationic deterative surfactant can be selected from the group consisting of: alkoxyate quaternary ammonium (AQA) surfactants as described in more detail in US 6,136,769; dimethyl hydroxyethyl quaternary ammonium as described in more detail in US 6,004,922; polyamine cationic surfactants as described in more detail in WO 98/35002, WO 98/35003, WO 98/35004, WO 98/35005, and WO 98/35006; cationic ester surfactants as described in more detail in US 4,228,042, US 4,239,660, US 4,260,529 and US 6,022,844; amino surfactants as described in more detail in US 6,221,825 and WO 00/47708, specifically amido propyldimethyl amine; and mixtures thereof. Preferred cationic deterative surfactants are quaternary ammonium compounds having the general formula:



wherein, R is a linear or branched, substituted or unsubstituted C₆₋₁₈ alkyl or alkenyl moiety, R¹ and R² are independently selected from methyl or ethyl moieties, R³ is a hydroxyl, hydroxymethyl or a hydroxyethyl moiety, X is an anion which provides charge neutrality, preferred anions include halides (such as chloride), sulphate and sulphonate. Preferred cationic deterative surfactants are mono-C₆₋₁₈ alkyl mono-hydroxyethyl di-methyl quaternary ammonium chlorides. Highly preferred cationic deterative surfactants are mono-C₈₋₁₀ alkyl mono-hydroxyethyl di-methyl quaternary ammonium chloride, mono-C₁₀₋₁₂ alkyl mono-hydroxyethyl di-methyl quaternary ammonium chloride and mono-C₁₀ alkyl mono-hydroxyethyl di-methyl quaternary ammonium chloride. Cationic surfactants such as Praepagen HY (tradename Clariant) may be useful and may also be useful as a suds booster.

[0044] The cationic deterative surfactant provides additional greasy soil cleaning performance. However, the cationic deterative surfactant may increase the tendency of any non-alkoxyated anionic deterative surfactant to precipitate out of solution. Preferably, the cationic deterative surfactant and any non-alkoxyated anionic deterative surfactant are separated in the detergent composition of the invention, for example if cationic surfactant is present, preferably the cationic and any anionic surfactant, particularly non-alkoxyated anionic surfactant will be present in the composition in separate particles. This minimises any effect that any cationic deterative surfactant may have on the undesirable precipitation of the anionic deterative surfactant, and also ensures that upon contact with water, the resultant wash liquor is not cloudy. If cationic surfactant is present, preferably the weight ratio of non-alkoxyated anionic deterative surfactant to cationic deterative surfactant is in the range of from 5:1 to 25:1, more preferably from 5:1 to 20:1 or from 6:1 to 15:1, or from 7:1 to 10:1, or even from 8:1 to 9:1.

[0045] Typically, the detergent composition comprises from 1 to 50 wt% anionic surfactant, more typically from 2 to 40 wt%. Alkyl benzene sulphonates are preferred anionic surfactants.

[0046] Preferred compositions of the present invention comprise at least two different surfactants in combination comprising at least one selected from a first group, the first group comprising alkyl benzene sulphonate and MES

surfactant; and at least one selected from a second group, the second group comprising alkoxylated anionic surfactant, MES and alkoxylated non-ionic surfactant and alpha olefin sulfonates (AOS). A particularly preferred combination comprises alkyl benzene sulphonate, preferably LAS in combination with MES. A further particularly preferred combination comprises alkyl benzene sulphonate, preferably LAS with an alkoxylated anionic surfactant, preferably C₈₋₁₈ alkyl alkoxylated sulphate having an average degree of alkoxylation of from 1 to 10. A third particularly preferred combination comprises alkyl benzene sulphonate, preferably LAS in combination with an alkoxylated non-ionic surfactant, preferably C₈₋₁₈ alkyl ethoxylated alcohol having a degree of alkoxylation of from 15 to 50, preferably from 20 to 40.

[0047] The weight ratio of the surfactant from the first group to the weight ratio of the surfactant from the second group is typically 1:5 to 100:1, preferably 1:2 to 100:1 or 1:1 to 50:1 or even to 20:1 or 10:1. The levels of the surfactants are as described above under the specific classes of surfactants. Presence of AE3S and/or MES in the system is preferred on account of their exceptional hardness-tolerance and ability to disperse lime soaps which are formed during the wash by lipase.

[0048] In a further embodiment, the surfactant in the detergent compositions of the invention comprises at least three surfactants, at least one from each of the first and second groups defined above and in addition a third surfactant, preferably also from the first or second groups defined above.


[0049] The detergent compositions of the invention may surprisingly contain relatively low levels of surfactant and yet still perform good cleaning, on account of the soil removal functionality delivered by the lipase, so that the overall level of surfactant may be below 12 wt%, or 10 wt% or 8 wt% based on total weight of the composition

Polymeric polycarboxylate

[0050] It may be desired for the compositions of the invention to comprise at least 0.1 wt%, or at least 0.5 wt%, or at least 2 or 3 wt%, or even at least 5 wt% polymeric polycarboxylates up to levels of 30 wt% or 20 wt% or 10 wt%. Preferred polymeric polycarboxylates include: polyacrylates, preferably having a weight average molecular weight of from 1,000Da to 20,000Da; co-polymers of maleic acid and acrylic acid, preferably having a molar ratio of maleic acid monomers to acrylic acid monomers of from 1:1 to 1:10 and a weight average molecular weight of from 10,000Da to 200,000Da, or preferably having a molar ratio of maleic acid monomers to acrylic acid monomers of from 0.3:1 to 3:1 and a weight average molecular weight of from 1,000Da to 50,000Da. Suitable polycarboxylates are the Sokalan CP, PA and HP ranges (BASF) such as Sokalan CP5, PA40 and HP22, and the Alcosperse range of polymers (Alco) such as Alcosperse 725, 747, 408, 412 and 420.

Soil dispersant

[0051] It may also be preferred for the composition to comprise a soil dispersant having the formula:



wherein, n = from 20 to 30, and x = from 3 to 8. Other suitable soil dispersants are sulphonated or sulphated soil dispersants having the formula: sulphonated or sulphated bis((C₂H₅O)(C₂H₄O)_n)(CH₃)-N⁺-C_xH_{2x}-N⁺-(CH₃)-bis((C₂H₅O)(C₂H₄O)_n)

wherein, n = from 20 to 30, and x = from 3 to 8. Preferably, the composition comprises at least 1wt%, or at least 2wt%, or at least 3wt% soil dispersants.

[0052] In a preferred embodiment of the invention, the detergent composition also comprises a suds booster, typically in amounts from 0.01 to 10 wt%, preferably in amounts from 0.02 to 5 wt% based on the total weight of the composition. Suitable suds boosters include fatty acid amides, fatty acid alkalonamides, betaines, sulfobetaines and amine oxides. Particularly preferred materials are cocamidopropyl betaine, cocomonoethanolamide and amine oxide. A suitable amine oxide is Admox 12, supplied by Albemarle.

Lime Soap Dispersants

[0053] It may also be preferred for the composition to comprise, especially when a lipase is present, anti-redeposition polymers such as the polymeric polycarboxylates described above. In addition, or alternatively, cellulose ethers such as carboxymethyl cellulose (CMC) will be useful. A suitable CMC is Tylose CR1500 G2, sold by Clariant. Suitable polymers are also sold by Andercol, Colombia under the Textilán brand name.

[0054] It is especially preferred to include additives with lime soap dispersancy functionality such as the aforementioned MES, AES, highly ethoxylated nonionic surfactant or polymers showing excellent lime soap dispersancy such as Acusol 460N (Rohm & Haas). Lists of suitable lime soap dispersants are given in the following references and documents cited therein.

[0055] WO9407974 (P&G), WO9407984 (P&G), WO9407985 (P&G), WO9504806 (P&G), WO9703379 (P&G), US6770610 (Clariant), EP0324568 (Rohm & Haas), EP0768370 (Rohm & Haas), M.K. Nagarajan and W.F. Masler, *Cosmetics and Toiletries*, 1989, 104, pp71-73, W. M. Linfield, *Tenside Surf. Det.*, 1990, 27, pp159-161, R.G. Bistline et al, *J. Am. Oil Chem. Soc.*, 1972, 49, pp63-69

[0056] Presence of a soil release polymer has been found to be especially beneficial in further strengthening the stain removal and cleaning benefits of the development, especially on synthetic fibres. Modified cellulose ethers such as methyl hydroxyethyl cellulose (MHEC), for example as sold by Clariant as Tylose MH50 G4 and Tylose MH300 G4, are preferred. Polyester-based Soil Release Polymers are especially preferred as they can also be effective as lime soap dispersants. Examples of suitable materials are Repel-o-Tex PF (supplied by Rhodia), Texcare SRA100 (supplied by Clariant) and Sokalan SR100 (BASF)

[0057] The detergent compositions of the invention may be in any convenient form such as solids such as powdered or granular or tablet solids, bars. Any of these forms may be partially or completely encapsulated. However, the present invention particularly relates to solid detergent compositions, especially granular compositions. Where the detergent compositions of the invention are solid, conventionally, surfactants are incorporated into agglomerates, extrudates or spray dried particles along with solid materials, usually builders, and these may be admixed to produce a fully formulated detergent composition according to the invention. When present in the granular form the detergent compositions of the present invention are preferably those having an overall bulk density of from 350 to 1200 g/l, more preferably 450 to 1000g/l or even 500 to 900g/l. Preferably, the detergent particles of the detergent composition in a granular form have a size average particle size of from 200 μ m to 2000 μ m, preferably from 350 μ m to 600 μ m.

[0058] Generally the detergent compositions of the invention will comprise a mixture of detergent particles including combinations of agglomerates, spray-dried powders and/or dry added materials such as bleaching agents, enzymes etc.

[0059] In one aspect of the invention the detergent compositions of the invention comprise an anionic surfactant from the list above which is a non-alkoxylated anionic deterative surfactant and this is preferably incorporated into the detergent composition in particulate form, such as via an agglomerate, a spray-dried powder, an extrudate, a bead, a noodle, a needle or a flake. Spray-dried particles are preferred. If via an agglomerate, the agglomerate preferably comprises at least 20%, by weight of the agglomerate, of a non-alkoxylated anionic deterative surfactant, more preferably from 25wt% to 65wt%, by weight of the agglomerate, of a non-alkoxylated anionic deterative surfactant. It may be preferred for part of the non-alkoxylated anionic deterative surfactant to be in the form of a spray-dried powder (e.g. a blown powder), and for part of the non-alkoxylated anionic deterative surfactant to be in the form of a non-spray-dried powder (e.g. an agglomerate, or an extrudate, or a flake such as a linear alkyl benzene sulphonate flake; suitable linear alkyl benzene sulphonate flakes are supplied by Pilot Chemical under the tradename F90[®], or by Stepan under the tradename Nacconol 90G[®]). This is especially preferred when it is desirable to incorporate high levels of non-alkoxylated anionic deterative surfactant in the composition.

[0060] Any alkoxylated anionic deterative surfactant may be incorporated into the detergent compositions of the invention via a spray-dried particle of a non-spray-dried powder such as an extrudate, agglomerate, preferably an agglomerate. Non-spray dried particles are preferred when it is desirable to incorporate high levels of alkoxylated anionic deterative surfactant in the composition

[0061] Any non-ionic deterative surfactant, or at least part thereof, can be incorporated into the composition in the form of a liquid spray-on, wherein the non-ionic deterative surfactant, or at least part thereof, in liquid form (e.g. in the form of a hot-melt) is sprayed onto the remainder of the composition. The non-ionic deterative surfactant, or at least part thereof, may be included into a particulate for incorporation into the detergent composition of the invention and the non-ionic deterative surfactant, or at least part thereof, may be dry-added to the remainder of the composition. The non-ionic surfactant, or at least part thereof, may be in the form of a co-particulate admixture with a solid carrier material such as carbonate salt, sulphate salt, borate, silica or any mixture thereof.

[0062] Any non-ionic deterative surfactant, or at least part thereof, may be in a co-particulate admixture with either an alkoxylated anionic deterative surfactant, a non-alkoxylated anionic deterative surfactant or a cationic deterative surfactant. The non-ionic deterative surfactant, or at least part thereof, may be agglomerated or extruded with either an alkoxylated anionic deterative surfactant, a non-alkoxylated anionic deterative surfactant or a cationic deterative surfactant.

[0063] The cationic deterative surfactant if present may be incorporated into the composition by incorporation in a particulate, such as a spray-dried powder, an agglomerate, an extrudate, a flake, a noodle, a needle, or any combination thereof. Preferably, the cationic deterative surfactant, or at least part thereof, is in the form of a spray-dried powder or an agglomerate.

First, second and third surfactant components

[0064] In a further aspect of the invention a detergent composition is provided comprising granular components, and comprising at least two separate surfactant components or even at least three separate surfactant components: a first, a second and an optional third surfactant component. These separate surfactant components may be present in separate

particulates so that at least two surfactant components are separate from one another in the detergent composition.

[0065] The composition preferably comprises at least two separate surfactant components, each in particulate form. It may be preferred for the composition to comprise at least three separate surfactant components, each in particulate form.

[0066] The first surfactant component predominantly comprises an alkoxyated deterative surfactant. By predominantly comprises, it is meant that the first surfactant component comprises greater than 50%, by weight of the first surfactant component, of an alkoxyated anionic deterative surfactant, preferably greater than 60%, or greater than 70%, or greater than 80%, or greater than 90% or even essentially 100%, by weight of the first surfactant component, of an alkoxyated anionic deterative surfactant. Preferably, the first surfactant component comprises less than 10%, by weight of the first surfactant component, of a non-alkoxyated anionic deterative surfactant, preferably less than 5%, or less than 2%, or even 0%, by weight of the first surfactant component, of a non-alkoxyated anionic deterative surfactant. Preferably, the first surfactant component is essentially free from non-alkoxyated anionic deterative surfactant. By essentially free from non-alkoxyated anionic deterative surfactant it is typically meant that the first surfactant component comprises no deliberately added non-alkoxyated anionic deterative surfactant. This is especially preferred in order to ensure that the composition has good dispensing and dissolution profiles, and also to ensure that the composition provides a clear wash liquor upon dissolution in water.

[0067] If cationic deterative surfactant is present in the composition, then preferably the first surfactant component comprises less than 10%, by weight of the first surfactant component, of a cationic deterative surfactant, preferably less than 5%, or less than 2%, or even 0%, by weight of the first surfactant component, of a cationic deterative surfactant. Preferably, the first surfactant component is essentially free from cationic deterative surfactant. By essentially free from cationic deterative surfactant it is typically meant that the first surfactant component comprises no deliberately added cationic deterative surfactant. This is especially preferred in order to reduce the degree of surfactant gelling in the wash liquor.

[0068] The first surfactant component is preferably in the form of a spray-dried powder, an agglomerate, an extrudate or a flake. If the first surfactant component is in the form of an agglomerate particle or an extrudate particle, then preferably the particle comprises from 20% to 65%, by weight of the particle, of an alkoxyated anionic deterative surfactant. If the first surfactant component is in spray-dried particle form, then preferably the particle comprises from 10wt% to 30wt%, by weight of the particle, of an alkoxyated anionic deterative surfactant. The first surfactant component may be in the form of a co-particulate admixture with a solid carrier material. The solid carrier material can be a sulphate salt and/or a carbonate salt, preferably sodium sulphate and/or sodium carbonate.

[0069] The second surfactant component predominantly comprises a non-alkoxyated deterative surfactant. By predominantly comprises, it is meant the second surfactant component comprises greater than 50%, by weight of the second surfactant component, of a non-alkoxyated anionic deterative surfactant, preferably greater than 60%, or greater than 70%, or greater than 80%, or greater than 90% or even essentially 100%, by weight of the second surfactant component, of a non-alkoxyated anionic deterative surfactant. Preferably, the second surfactant component comprises less than 10%, by weight of the second surfactant component, of an alkoxyated anionic deterative surfactant, preferably less than 5%, or less than 2%, or even 0%, by weight of the second surfactant component, of an alkoxyated anionic deterative surfactant. If cationic deterative surfactant is present in the composition, then preferably the second surfactant component comprises less than 10%, by weight of the second surfactant component, of a cationic deterative surfactant, preferably less than 5%, or less than 2%, or even 0%, by weight of the second surfactant component, of a cationic deterative surfactant. Preferably, the second surfactant component is essentially free from alkoxyated anionic deterative surfactant. By essentially free from alkoxyated anionic deterative surfactant it is typically meant that the second surfactant component comprises no deliberately added alkoxyated anionic deterative surfactant. Preferably, the second surfactant component is essentially free from cationic deterative surfactant. By essentially free from cationic deterative surfactant it is typically meant that the second surfactant component comprises no deliberately added cationic deterative surfactant. This is especially preferred in order to ensure that the composition has good dispensing and dissolution profiles, and also to ensure that the composition provides a clear wash liquor upon dissolution in water.

[0070] The second surfactant component may be in the form of a spray-dried powder, a flash-dried powder, an agglomerate or an extrudate. If the second surfactant component is in the form of an agglomerate particle, then preferably the particle from 5% to 50%, by weight of the particle, of a non-alkoxyated anionic deterative surfactant, or from 5wt% to 25wt% non-alkoxyated anionic deterative surfactant. The second surfactant component may be in form of a co-particulate admixture with a solid carrier material. The solid carrier material can be a sulphate salt and/or a carbonate salt, preferably sodium sulphate and/or sodium carbonate.

[0071] Although the detergent compositions of the invention may be substantially free of cationic surfactant, if present, the cationic surfactant may be present in a third surfactant component or may be incorporated into a spray-dried particle with at least some anionic surfactant. If present in a third component, it may be beneficial to have the third surfactant component predominantly comprising a cationic deterative surfactant. By predominantly comprises, it is meant the third surfactant component comprises greater than 50%, by weight of the third surfactant component, of a cationic deterative surfactant, preferably greater than 60%, or greater than 70%, or greater than 80%, or greater than 90% or even essentially

100%, by weight of the third surfactant component, of a cationic deterative surfactant. Preferably, the third surfactant component comprises less than 10%, by weight of the third surfactant component, of an alkoxyated anionic deterative surfactant, preferably less than 5%, or less than 2%, or even essentially 0%, by weight of the third surfactant component, of an alkoxyated anionic deterative surfactant. Preferably the third surfactant component comprises less than 10%, by weight of the third surfactant component, of a non-alkoxyated anionic deterative surfactant, preferably less than 5%, or less than 2%, or even 0%, by weight of the third surfactant component, of a non-alkoxyated anionic deterative surfactant. Preferably, the third surfactant component is essentially free from alkoxyated anionic deterative surfactant. By essentially free from alkoxyated anionic deterative surfactant it is typically meant that the third surfactant component comprises no deliberately added alkoxyated anionic deterative surfactant. Preferably, the third surfactant component is essentially free from non-alkoxyated anionic deterative surfactant. By essentially free from non-alkoxyated anionic deterative surfactant it is typically meant that the third surfactant component comprises no deliberately added non-alkoxyated anionic deterative surfactant. This is especially preferred in order to ensure that the composition has good dispensing and dissolution profiles, and also to ensure that the composition provides a clear wash liquor upon dissolution in water.

[0072] The third surfactant component is preferably in the form of a spray-dried powder, a flash-dried powder, an agglomerate or an extrudate. If the third surfactant component is in the form of an agglomerate particle, then preferably the particle comprises from 5% to 50%, by weight of the particle, of cationic deterative surfactant, or from 5wt% to 25wt% cationic deterative surfactant. The third surfactant component may be in form of a co-particulate admixture with a solid carrier material. The solid carrier material can be a sulphate salt and/or a carbonate salt, preferably sodium sulphate and/or sodium carbonate.

Optional Deterative Adjuncts

[0073] Optionally, the detergent ingredients can include one or more other deterative adjuncts or other materials for assisting or enhancing cleaning performance, treatment of the substrate to be cleaned, or to modify the aesthetics of the detergent composition. Usual deterative adjuncts of detergent compositions include the ingredients set forth in U.S. Pat. No. 3,936,537, Baskerville et al. and in Great Britain Patent Application No. 9705617.0, Trinh et al., published September 24, 1997. Such adjuncts are included in detergent compositions at their conventional art-established levels of use, generally from 0 wt% to about 80 wt% of the detergent ingredients, preferably from about 0.5 wt% to about 20wt % and can include color speckles, suds boosters, suds suppressors, anti-tarnish and/or anticorrosion agents, soil-suspending agents, soil release agents, dyes, fillers, optical brighteners, germicides, alkalinity sources, hydrotropes, antioxidants, enzymes, enzyme stabilizing agents, solvents, solubilizing agents, chelating agents, clay soil removal/anti-redeposition agents, polymeric dispersing agents, processing aids, fabric softening components, static control agents, bleaching agents, bleaching activators, bleach stabilizers, dye-transfer inhibitors, flocculants, fabric softeners, suds suppressors, fabric integrity agents, perfumes, whitening agents, alkali metal sulphate salts, sulphamic acid, sodium sulphate and sulphamic acid complexes, etc and combinations thereof. The precise nature of these additional components, and levels of incorporation thereof will depend on the physical form of the composition or component, and the precise nature of the washing operation for which it is to be used.

[0074] Preferred zwitterionic surfactants comprise one or more quaternized nitrogen atoms and one or more moieties selected from the group consisting of: carbonate, phosphate, sulphate, sulphonate, and combinations thereof. Preferred zwitterionic surfactants are alkyl betaines. Other preferred zwitterionic surfactants are alkyl amine oxides. Catanionic surfactants which are complexes comprising a cationic surfactant and an anionic surfactant may also be included. Typically, the molar ratio of the cationic surfactant to anionic surfactant in the complex is greater than 1:1, so that the complex has a net positive charge.

[0075] A preferred adjunct component is a bleaching agent. Preferably, the detergent composition comprises one or more bleaching agents. Typically, the composition comprises (by weight of the composition) from 1% to 50% of one or more bleaching agent. Preferred bleaching agents are selected from the group consisting of sources of peroxide, sources of peracid, bleach boosters, bleach catalysts, photo-bleaches, and combinations thereof. Preferred sources of peroxide are selected from the group consisting of: perborate monohydrate, perborate tetra-hydrate, percarbonate, salts thereof, and combinations thereof. Preferred sources of peracid are selected from the group consisting of: bleach activator typically with a peroxide source such as perborate or percarbonate, preformed peracids, and combinations thereof. Preferred bleach activators are selected from the group consisting of: oxy-benzene-sulphonate bleach activators, lactam bleach activators, imide bleach activators, and combinations thereof. A preferred source of peracid is tetra-acetyl ethylene diamine (TAED) and peroxide source such as percarbonate. Preferred oxy-benzene-sulphonate bleach activators are selected from the group consisting of: nonanoyl-oxy-benzene-sulphonate, 6-nonamido-caproyl-oxy-benzene-sulphonate, salts thereof, and combinations thereof. Preferred lactam bleach activators are acyl- caprolactams and/or acyl-valerolactams. A preferred imide bleach activator is N-nonanoyl-N-methyl-acetamide.

[0076] Preferred preformed peracids are selected from the group consisting of N,N-pthaloyl-amino-peroxycaproic acid, nonyl-amido-peroxyadipic acid, salts thereof, and combinations thereof. Preferably, the STW-composition com-

prises one or more sources of peroxide and one or more sources of peracid. Preferred bleach catalysts comprise one or more transition metal ions. Other preferred bleaching agents are di-acyl peroxides. Preferred bleach boosters are selected from the group consisting of zwitterionic imines, anionic imine polyions, quaternary oxaziridinium salts, and combinations thereof. Highly preferred bleach boosters are selected from the group consisting of: aryliminium zwitterions, aryliminium polyions, and combinations thereof. Suitable bleach boosters are described in US360568, US5360569 and US5370826.

[0077] A preferred adjunct component is an anti-redeposition agent. Preferably, the detergent composition comprises one or more anti-redeposition agents. Preferred anti-redeposition agents are cellulosic polymeric components, most preferably carboxymethyl celluloses.

[0078] A preferred adjunct component is a chelant. Preferably, the detergent composition comprises one or more chelants. Preferably, the detergent composition comprises (by weight of the composition) from 0.01% to 10% chelant, or 0.01 to 5 wt% or 4 wt% or 2 wt%. Preferred chelants are selected from the group consisting of: hydroxyethane-dimethylene-phosphonic acid (HEDP), 2-phosphonobutane-1,2,4-tricarboxylic acid (PBTC), ethylene diamine tetra (methylene phosphonic) acid, diethylene triamine pentacetate, ethylene diamine tetraacetate, diethylene triamine penta (methyl phosphonic) acid, ethylene diamine disuccinic acid, and combinations thereof. A further preferred chelant is an anionically modified catechol. An anionically modified catechol, as used herein, means 1,2-benzenediol having one or two anionic substitutions on the benzene ring. The anionic substitutions may be selected from sulfonate, sulfate, carbonate, phosphonate, phosphate, fluoride, and mixtures thereof. One embodiment of an anionically modified catechol having two sulfate moieties having a sodium cation on the benzene ring is 4,5-dihydroxy-m-benzenedisulfonic acid, disodium salt (Tiron®). Preferably, the anionically modified catechol is essentially free (less than 3%) of catechol (1,2-benzenediol), to avoid skin irritation when present.

[0079] A preferred adjunct component is a dye transfer inhibitor. Preferably, the detergent composition comprises one or more dye transfer inhibitors. Typically, dye transfer inhibitors are polymeric components that trap dye molecules and retain the dye molecules by suspending them in the wash liquor. Preferred dye transfer inhibitors are selected from the group consisting of polyvinylpyrrolidones, polyvinylpyridine N-oxides, polyvinylpyrrolidone-polyvinylimidazole copolymers, and combinations thereof.


[0080] Preferred adjunct components include other enzymes. Preferred enzymes are selected from the group consisting of: amylases, arabinosidases, carbohydrases, cellulases, chondroitinases, cutinases, dextranases, esterases, β -glucanases, glucoamylases, hyaluronidases, keratanases, laccases, ligninases, lipoxygenases, malanases, mannanases, oxidases, pectinases, pentosanases, peroxidases, phenoloxidases, phospholipases, proteases, pullulanases, reductases, tannases, transferases, xylanases, xyloglucanases, and combinations thereof. Preferred additional enzymes are selected from the group consisting of: lipases, amylases, carbohydrases, cellulases, proteases, and combinations thereof, more preferably a lipase, for further improved cleaning and whitening performance.

[0081] Preferred adjunct components include fluorescent whitening agents. Any fluorescent whitening agent suitable for use in a laundry detergent composition may be used in the composition of the present invention. The most commonly used fluorescent whitening agents are those belonging to the classes of diaminostilbene-sulphonic acid derivatives, diarylpyrazoline derivatives and bisphenyl-distyryl derivatives. Examples of the diaminostilbene-sulphonic acid derivative type of fluorescent whitening agents include the sodium salts of:

4,4'-bis-(2-diethanolamino-4-anilino-s-triazin-6-ylamino) stilbene-2,2'-disulphonate,
 4,4'-bis-(2,4-dianilino-s-triazin-6-ylamino) stilbene-2,2'-disulphonate,
 4,4'-bis-(2-anilino-4(N-methyl-N-2-hydroxy-ethylamino)-s-triazin-6-ylamino) stilbene-2,2'-disulphonate,
 4,4'-bis-(4-phenyl-2,1,3-triazol-2-yl)stilbene-2,2'-disulphonate,
 4,4'-bis-(2-anilino-4(1-methyl-2-hydroxy-ethylamino)-s-triazin-6-ylamino) stilbene-2,2'-disulphonate and,
 2-(stilbyl-4"-naptho-1.,2':4,5)-1,2,3-trizole-2"-sulphonate.

Preferred fluorescent whitening agents are Tinopal® DMS and Tinopal® CBS available from Ciba-Geigy AG, Basel, Switzerland. Tinopal® DMS is the disodium salt of 4,4'-bis-(2-morpholino-4 anilino-s-triazin-6-ylamino) stilbene disulphonate. Tinopal® CBS is the disodium salt of 2,2'-bis-(phenyl-styryl) disulphonate.

[0082] Also preferred are fluorescent whitening agents of the structure:



wherein R1 and R2, together with the nitrogen atom linking them, form an unsubstituted or C1-C4 alkyl-substituted morpholino, piperidine or pyrrolidine ring, preferably a morpholino ring (commercially available as Parawhite KX, supplied by Paramount Minerals and Chemicals, Mumbai, India). Other fluorescent whitening agents suitable for use in the invention include the 1-3-diaryl pyrazolines and the 7-alkylaminocoumarins. Typical levels of fluorescent whitening agent in the composition are from 0.03-0.5%, preferably 0.05 % to 0.3 % by weight.

[0083] A preferred adjunct component is a fabric integrity agent. Preferably, the detergent composition comprises one or more fabric integrity agents. Typically, fabric integrity agents are polymeric components that deposit on the fabric surface and prevent fabric damage during the laundering process. Preferred fabric integrity agents are hydrophobically modified celluloses. These hydrophobically modified celluloses reduce fabric abrasion, enhance fibre-fibre interactions and reduce dye loss from the fabric. A preferred hydrophobically modified cellulose is described in WO99/14245. Other preferred fabric integrity agents are polymeric components and/or oligomeric components that are obtainable, preferably obtained, by a process comprising the step of condensing imidazole and epichlorohydrin.

[0084] A preferred adjunct component is a salt. Preferably, the detergent composition comprises one or more salts. The salts can act as alkalinity agents, buffers, builders, co-builders, encrustation inhibitors, fillers, pH regulators, stability agents, and combinations thereof. Typically, the detergent composition comprises (by weight of the composition) from 5% to 60% salt. Preferred salts are alkali metal salts of aluminate, carbonate, chloride, bicarbonate, nitrate, phosphate, silicate, sulphate, and combinations thereof. Other preferred salts are alkaline earth metal salts of aluminate, carbonate, chloride, bicarbonate, nitrate, phosphate, silicate, sulphate, and combinations thereof. Especially preferred salts are sodium sulphate, sodium carbonate, sodium bicarbonate, sodium silicate, sodium sulphate, and combinations thereof. Optionally, the alkali metal salts and/or alkaline earth metal salts may be anhydrous.

[0085] A preferred adjunct component is a soil release agent. Preferably, the detergent composition comprises one or more soil release agents. Typically, soil release agents are polymeric compounds that modify the fabric surface and prevent the redeposition of soil on the fabric. Preferred soil release agents are copolymers, preferably block copolymers, comprising one or more terephthalate unit. Preferred soil release agents are copolymers that are synthesised from dimethylterephthalate, 1,2-propyl glycol and methyl capped polyethyleneglycol. Other preferred soil release agents are anionically end capped polyesters.

Softening system

[0086] The detergent compositions of the invention may comprise softening agents for softening through the wash such as clay optionally also with flocculant and enzymes. Further more specific description of suitable detergent components can be found in WO97/11151.

Washing Method

[0087] The invention also includes methods of washing textiles comprising contacting textiles with an aqueous solution comprising the detergent composition of the invention. The invention may be particularly beneficial at low water temperatures such as below 30°C or below 25 or 20°C. Typically the aqueous wash liquor will comprise at least 100 ppm, or at least 500ppm of the detergent composition.

Examples

[0088] The following are examples of the invention.

Examples 1-6

[0089] Granular laundry detergent compositions designed for handwashing or top-loading washing machines.

EP 1 882 731 A1

	1 (wt%)	2 (wt%)	3 (wt%)	4 (wt%)	5 (wt%)	6 (wt%)
5	Linear alkylbenzenesulfonate	20	22	20	15	20
	C ₁₂₋₁₄ Dimethylhydroxyethyl ammonium chloride	0.7	0.2	1	0.6	0
10	AE3S	0.9	1	0.9	0.0	0.9
	AE7	0.0	0.0	0.0	1	3
	Sodium tripolyphosphate	5	0.0	4	3	2
15	Zeolite A	0.0	1	0.0	1	4
	1.6R Silicate (SiO ₂ : Na ₂ O at ratio 1.6:1)	4	5	2	3	5
	Sodium Carbonate	25	20	25	17	25
20	Polyacrylate MW 4500	1	0.6	1	1	1.5
	Carboxy Methyl Cellulose	1	0.3	1	1	1
25	Celluclean® (15.6mg/g)	0.1	0.2	0.1	0.2	0.1
	Savinase® 32.89mg/g	0.1	0.1	0.1	0.1	0.1
	Natalase® 8.65mg/g	0.1	0.0	0.1	0.0	0.1
30	Lipex® 18mg/g	0.03	0.07	0.3	0.1	0.07
	Fluorescent Brightener 1	0.06	0.0	0.06	0.18	0.06
	Fluorescent Brightener 2	0.1	0.06	0.1	0.0	0.1
35	Diethylenetriamine pentaacetic acid or Ethylene diamine tetraacetic acid	0.6	0	0.6	0.25	0.6
40	MgSO ₄	1	1	1	0.5	1
	Sodium Percarbonate	0.0	0	0.1	0.0	0.0
	Sodium Perborate Monohydrat	4.4	0.0	3.85	2.09	0.78
45	NOBS	1.9	0.0	1.66	0.0	0.33
	TAED	0.58	0	0.51	0.0	0.015
	Perfume spray-on	0.4	0.4	0.6	1	0.3
50	Starch encapsulated perfume	0.3	0.2	0.3	0.2	0.3
	Sulfate/Moisture	Balance to 100%	Balance to 100%	Balance to 100%	Balance to 100%	Balance to 100%

Examples 7-12

[0090] Granular laundry detergent compositions designed for front-loading automatic washing machines.

EP 1 882 731 A1

	7 (wt%)	8 (wt%)	9 (wt%)	10 (wt%)	11 (wt%)	12 (wt%)
5						
	Linear alkylbenzenesulfonate	8	7.1	7	6.5	7.5
	AE3S	0	4.8	0	5.2	4
	Alkylsulfate	1	0	1	0	0
10	AE7	2.2	0	3.2	0	0
	C ₁₀₋₁₂ Dimethyl hydroxyethylammonium chloride	0.75	0.94	0.98	0.98	0
15	Crystalline layered silicate (δ -Na ₂ Si ₂ O ₅)	2.0	0	2.0	0	0
	Zeolite A	7	0	7	0	2
	Citric Acid	3	5	3	4	2.5
20	Sodium Carbonate	15	20	14	20	23
	Silicate 2R (SiO ₂ :Na ₂ O at ratio 2:1)	0.08	0	0.11	0	0
	Soil release agent	0.75	0.72	0.71	0.72	0
25	Acrylic Acid/Maleic Acid Copolymer	1.1	3.7	1.0	3.7	2.6
	Carboxymethylcellulose	0.15	1.4	0.2	1.4	1
	Protease (84mg active/g)	0.2	0.2	0.3	0.15	0.12
30	Celluclean® (15.6mg active/g)	0.2	0.15	0.2	0.3	0.15
	Lipex®(18.00mg active/g)	0.05	0.15	0.1	0	0
35	Termamyl® (25mg active/g)	0.1	0.1	0.1	0.12	0.1
	Natalase® (8.65mg active/g)	0.1	0.2	0	0	0.15
40	Termamyl® (25 mg active/g)	0.2	0.1	0.2	0	0.1
	TAED	3.6	4.0	3.6	4.0	2.2
45	Percarbonate	13	13.2	13	13.2	16
	Na salt of Ethylenediamine-N,N'-disuccinic acid, (S,S) isomer (EDDS)	0.2	0.2	0.2	0.2	0.2
50	Hydroxyethane di phosphonate (HEDP)	0.2	0.2	0.2	0.2	0.2
	MgSO ₄	0.42	0.42	0.42	0.42	0.4
55	Perfume	0.5	0.6	0.5	0.6	0.6
	Starch Encapsulated Perfume	0.2	0.5	0.3	0.4	0.3

EP 1 882 731 A1

(continued)

	7 (wt%)	8 (wt%)	9 (wt%)	10 (wt%)	11 (wt%)	12 (wt%)
Suds suppressor agglomerate	0.05	0.1	0.05	0.1	0.06	0.05
Soap	0.45	0.45	0.45	0.45	0	0
Sulfate/ Water & Miscellaneous	Balance to 100%	Balance to 100%	Balance to 100%	Balance to 100%	Balance to 100%	Balance to 100%

Any of the above compositions is used to launder fabrics at a concentration of 7000 to 10000 ppm in water, 20-90 °C, and a 5:1 water:cloth ratio. The typical pH is about 10.

Raw Materials and Notes For Composition Examples 1-12

[0091] Linear alkylbenzenesulfonate having an average aliphatic carbon chain length C_{11} - C_{12} supplied by Stepan, Northfield, Illinois, USA

C_{12-14} Dimethylhydroxyethyl ammonium chloride, supplied by Clariant GmbH, Sulzbach, Germany

AE3S is C_{12-15} alkyl ethoxy (3) sulfate supplied by Stepan, Northfield, Illinois, USA

AE7 is C_{12-15} alcohol ethoxylate, with an average degree of ethoxylation of 7, supplied by Huntsman, Salt Lake City, Utah, USA

Sodium tripolyphosphate is supplied by Rhodia, Paris, France

Zeolite A was supplied by Industrial Zeolite (UK) Ltd, Grays, Essex, UK 1.6R Silicate was supplied by Koma, Nestemica, Czech Republic

Sodium Carbonate was supplied by Solvay, Houston, Texas, USA

Polyacrylate MW 4500 is supplied by BASF, Ludwigshafen, Germany

Carboxy Methyl Cellulose is Finnfix® BDA supplied by CPKelco, Arnhem, Netherlands

Savinase®, Natalase®, Lipex®, Termamyl®, Mannaway®, Celluclean® supplied by Novozymes, Bagsvaerd, Denmark

Protease (examples 7-12) described in patent application US 6312936B1 was supplied by Genencor International, Palo Alto, California, USA

Fluorescent Brightener 1 is Tinopal® AMS, Fluorescent Brightener 2 is Tinopal® CBS-X. Sulphonated zinc phthalocyanine supplied by Ciba Specialty Chemicals, Basel, Switzerland

Diethylenetriamine pentacetic acid was supplied by Dow Chemical, Midland, Michigan, USA

Sodium percarbonate supplied by Solvay, Houston, Texas, USA

Sodium perborate was supplied by Degussa, Hanau, Germany

NOBS is sodium nonanoyloxybenzenesulfonate, supplied by Eastman, Batesville, Arkansas, USA

TAED is tetraacetylenediamine, supplied under the Peractive® brand name by Clariant GmbH, Sulzbach, Germany

S-ACMC is carboxymethylcellulose conjugated with C.I. Reactive Blue 19, sold by Megazyme, Wicklow, Ireland under the product name AZO-CM-CELLULOSE, product code S-ACMC.

Soil release agent is Repel-o-tex® PF, supplied by Rhodia, Paris, France

Acrylic Acid/Maleic Acid Copolymer is molecular weight 70,000 and acrylate:maleate ratio 70:30, supplied by BASF, Ludwigshafen, Germany

Na salt of Ethylenediamine-N,N'-disuccinic acid, (S,S) isomer (EDDS) was supplied by Octel, Ellesmere Port, UK

Hydroxyethane di phosphonate (HEDP) was supplied by Dow Chemical, Midland, Michigan, USA

Suds suppressor agglomerate was supplied by Dow Coming, Midland, Michigan, USA

SEQUENCE LISTING

5 <110> THE PROCTER & GAMBLE COMPANY
 <120> Detergent compositions
 <130> CM3098F
 <160> 2
 10 <170> PatentIn version 3.3
 <210> 1
 <211> 773
 <212> PRT
 15 <213> Bacillus sp.
 <400> 1
 Ala Glu Gly Asn Thr Arg Glu Asp Asn Phe Lys His Leu Leu Gly Asn
 1 5 10 15
 20 Asp Asn Val Lys Arg Pro Ser Glu Ala Gly Ala Leu Gln Leu Gln Glu
 20 25 30
 Val Asp Gly Gln Met Thr Leu Val Asp Gln His Gly Glu Lys Ile Gln
 35 40 45
 25 Leu Arg Gly Met Ser Thr His Gly Leu Gln Trp Phe Pro Glu Ile Leu
 50 55 60
 30 Asn Asp Asn Ala Tyr Lys Ala Leu Ala Asn Asp Trp Glu Ser Asn Met
 65 70 75 80
 Ile Arg Leu Ala Met Tyr Val Gly Glu Asn Gly Tyr Ala Ser Asn Pro
 85 90 95
 35 Glu Leu Ile Lys Ser Arg Val Ile Lys Gly Ile Asp Leu Ala Ile Glu
 100 105 110
 40 Asn Asp Met Tyr Val Ile Val Asp Trp His Val His Ala Pro Gly Asp
 115 120 125
 Pro Arg Asp Pro Val Tyr Ala Gly Ala Glu Asp Phe Phe Arg Asp Ile
 130 135 140
 45 Ala Ala Leu Tyr Pro Asn Asn Pro His Ile Ile Tyr Glu Leu Ala Asn
 145 150 155 160
 Glu Pro Ser Ser Asn Asn Asn Gly Gly Ala Gly Ile Pro Asn Asn Glu
 165 170 175
 50 Glu Gly Trp Asn Ala Val Lys Glu Tyr Ala Asp Pro Ile Val Glu Met
 180 185 190
 55 Leu Arg Asp Ser Gly Asn Ala Asp Asp Asn Ile Ile Ile Val Gly Ser

EP 1 882 731 A1

5	Pro	Asn 210	Trp	Ser	Gln	Arg	Pro 215	Asp	Leu	Ala	Ala	Asp 220	Asn	Pro	Ile	Asn
10	Asp 225	His	His	Thr	Met	Tyr 230	Thr	Val	His	Phe	Tyr 235	Thr	Gly	Ser	His	Ala 240
15	Ala	Ser	Thr	Glu	Ser 245	Tyr	Pro	Pro	Glu	Thr 250	Pro	Asn	Ser	Glu	Arg 255	Gly
20	Asn	Val	Met	Ser 260	Asn	Thr	Arg	Tyr	Ala 265	Leu	Glu	Asn	Gly	Val 270	Ala	Val
25	Phe	Ala	Thr 275	Glu	Trp	Gly	Thr	Ser 280	Gln	Ala	Asn	Gly	Asp 285	Gly	Gly	Pro
30	Tyr	Phe 290	Asp	Glu	Ala	Asp	Val 295	Trp	Ile	Glu	Phe	Leu 300	Asn	Glu	Asn	Asn
35	Ile 305	Ser	Trp	Ala	Asn	Trp 310	Ser	Leu	Thr	Asn	Lys 315	Asn	Glu	Val	Ser	Gly 320
40	Ala	Phe	Thr	Pro	Phe 325	Glu	Leu	Gly	Lys	Ser 330	Asn	Ala	Thr	Asn	Leu 335	Asp
45	Pro	Gly	Pro	Asp 340	His	Val	Trp	Ala	Pro 345	Glu	Glu	Leu	Ser	Leu 350	Ser	Gly
50	Glu	Tyr	Val 355	Arg	Ala	Arg	Ile	Lys 360	Gly	Val	Asn	Tyr	Glu 365	Pro	Ile	Asp
55	Arg	Thr 370	Lys	Tyr	Thr	Lys	Val 375	Leu	Trp	Asp	Phe	Asn 380	Asp	Gly	Thr	Lys
60	Gln	Gly	Phe	Gly	Val	Asn 390	Ser	Asp	Ser	Pro	Asn 395	Lys	Glu	Leu	Ile	Ala 400
65	Val	Asp	Asn	Glu	Asn 405	Asn	Thr	Leu	Lys	Val 410	Ser	Gly	Leu	Asp	Val 415	Ser
70	Asn	Asp	Val	Ser 420	Asp	Gly	Asn	Phe	Trp 425	Ala	Asn	Ala	Arg	Leu 430	Ser	Ala
75	Asp	Gly	Trp 435	Gly	Lys	Ser	Val	Asp 440	Ile	Leu	Gly	Ala	Glu 445	Lys	Leu	Thr
80	Met	Asp 450	Val	Ile	Val	Asp	Glu 455	Pro	Thr	Thr	Val	Ala 460	Ile	Ala	Ala	Ile
85	Pro	Gln	Ser	Ser	Lys	Ser	Gly	Trp	Ala	Asn	Pro	Glu	Arg	Ala	Val	Arg

EP 1 882 731 A1

	465					470										475													480
5	Val	Asn	Ala	Glu	Asp 485	Phe	Val	Gln	Gln	Thr 490	Asp	Gly	Lys	Tyr	Lys 495	Ala													
	Gly	Leu	Thr	Ile 500	Thr	Gly	Glu	Asp	Ala 505	Pro	Asn	Leu	Lys	Asn 510	Ile	Ala													
10																													
	Phe	His	Glu 515	Glu	Asp	Asn	Asn	Met 520	Asn	Asn	Ile	Ile	Leu 525	Phe	Val	Gly													
	Thr	Asp 530	Ala	Ala	Asp	Val	Ile 535	Tyr	Leu	Asp	Asn	Ile 540	Lys	Val	Ile	Gly													
15																													
	Thr	Glu	Val	Glu	Ile	Pro 550	Val	Val	His	Asp	Pro 555	Lys	Gly	Glu	Ala	Val 560													
20																													
	Leu	Pro	Ser	Val	Phe 565	Glu	Asp	Gly	Thr	Arg 570	Gln	Gly	Trp	Asp	Trp 575	Ala													
	Gly	Glu	Ser	Gly 580	Val	Lys	Thr	Ala	Leu 585	Thr	Ile	Glu	Glu	Ala 590	Asn	Gly													
25																													
	Ser	Asn	Ala 595	Leu	Ser	Trp	Glu	Phe 600	Gly	Tyr	Pro	Glu	Val 605	Lys	Pro	Ser													
30																													
	Asp	Asn 610	Trp	Ala	Thr	Ala	Pro 615	Arg	Leu	Asp	Phe	Trp 620	Lys	Ser	Asp	Leu													
	Val	Arg	Gly	Glu	Asn	Asp 630	Tyr	Val	Ala	Phe	Asp 635	Phe	Tyr	Leu	Asp	Pro 640													
35																													
	Val	Arg	Ala	Thr	Glu 645	Gly	Ala	Met	Asn	Ile 650	Asn	Leu	Val	Phe	Gln 655	Pro													
40																													
	Pro	Thr	Asn	Gly 660	Tyr	Trp	Val	Gln	Ala 665	Pro	Lys	Thr	Tyr	Thr 670	Ile	Asn													
	Phe	Asp	Glu 675	Leu	Glu	Glu	Ala	Asn 680	Gln	Val	Asn	Gly	Leu 685	Tyr	His	Tyr													
45																													
	Glu	Val 690	Lys	Ile	Asn	Val	Arg 695	Asp	Ile	Thr	Asn	Ile 700	Gln	Asp	Asp	Thr													
50																													
	Leu	Leu	Arg	Asn	Met	Met 710	Ile	Ile	Phe	Ala	Asp 715	Val	Glu	Ser	Asp	Phe 720													
	Ala	Gly	Arg	Val	Phe 725	Val	Asp	Asn	Val	Arg 730	Phe	Glu	Gly	Ala	Ala														

EP 1 882 731 A1

	740	745	750
5	Pro Val Asp Glu Lys Glu Ala Lys Lys Glu Gln Lys Glu Ala Glu Lys		
	755	760	765
	Glu Glu Lys Glu Glu		
10	770		
	<210> 2		
	<211> 824		
	<212> PRT		
	<213> Bacillus sp.		
15	<220>		
	<223> Bacillus sp. KSM-S237		
	<400> 2		
20	Met Met Leu Arg Lys Lys Thr Lys Gln Leu Ile Ser Ser Ile Leu Ile		
	1	5	10
	Leu Val Leu Leu Leu Ser Leu Phe Pro Ala Ala Leu Ala Ala Glu Gly		
	20	25	30
25	Asn Thr Arg Glu Asp Asn Phe Lys His Leu Leu Gly Asn Asp Asn Val		
	35	40	45
30	Lys Arg Pro Ser Glu Ala Gly Ala Leu Gln Leu Gln Glu Val Asp Gly		
	50	55	60
	Gln Met Thr Leu Val Asp Gln His Gly Glu Lys Ile Gln Leu Arg Gly		
	65	70	75
35	Met Ser Thr His Gly Leu Gln Trp Phe Pro Glu Ile Leu Asn Asp Asn		
	85	90	95
40	Ala Tyr Lys Ala Leu Ser Asn Asp Trp Asp Ser Asn Met Ile Arg Leu		
	100	105	110
	Ala Met Tyr Val Gly Glu Asn Gly Tyr Ala Thr Asn Pro Glu Leu Ile		
	115	120	125
45	Lys Gln Arg Val Ile Asp Gly Ile Glu Leu Ala Ile Glu Asn Asp Met		
	130	135	140
50	Tyr Val Ile Val Asp Trp His Val His Ala Pro Gly Asp Pro Arg Asp		
	145	150	155
	Pro Val Tyr Ala Gly Ala Lys Asp Phe Phe Arg Glu Ile Ala Ala Leu		
	165	170	175
55	Tyr Pro Asn Asn Pro His Ile Ile Tyr Glu Leu Ala Asn Glu Pro Ser		
	180	185	190

EP 1 882 731 A1

Ser Asn Asn Asn Gly Gly Ala Gly Ile Pro Asn Asn Glu Glu Gly Trp
 195 200 205
 5
 Lys Ala Val Lys Glu Tyr Ala Asp Pro Ile Val Glu Met Leu Arg Lys
 210 215 220
 10
 Ser Gly Asn Ala Asp Asp Asn Ile Ile Ile Val Gly Ser Pro Asn Trp
 225 230 235 240
 Ser Gln Arg Pro Asp Leu Ala Ala Asp Asn Pro Ile Asp Asp His His
 245 250 255
 15
 Thr Met Tyr Thr Val His Phe Tyr Thr Gly Ser His Ala Ala Ser Thr
 260 265 270
 Glu Ser Tyr Pro Ser Glu Thr Pro Asn Ser Glu Arg Gly Asn Val Met
 275 280 285
 20
 Ser Asn Thr Arg Tyr Ala Leu Glu Asn Gly Val Ala Val Phe Ala Thr
 290 295 300
 25
 Glu Trp Gly Thr Ser Gln Ala Ser Gly Asp Gly Gly Pro Tyr Phe Asp
 305 310 315 320
 Glu Ala Asp Val Trp Ile Glu Phe Leu Asn Glu Asn Asn Ile Ser Trp
 325 330 335
 30
 Ala Asn Trp Ser Leu Thr Asn Lys Asn Glu Val Ser Gly Ala Phe Thr
 340 345 350
 35
 Pro Phe Glu Leu Gly Lys Ser Asn Ala Thr Asn Leu Asp Pro Gly Pro
 355 360 365
 Asp His Val Trp Ala Pro Glu Glu Leu Ser Leu Ser Gly Glu Tyr Val
 370 375 380
 40
 Arg Ala Arg Ile Lys Gly Val Asn Tyr Glu Pro Ile Asp Arg Thr Lys
 385 390 395 400
 45
 Tyr Thr Lys Val Leu Trp Asp Phe Asn Asp Gly Thr Lys Gln Gly Phe
 405 410 415
 Gly Val Asn Ser Asp Ser Pro Asn Lys Glu Leu Ile Ala Val Asp Asn
 420 425 430
 50
 Glu Asn Asn Thr Leu Lys Val Ser Gly Leu Asp Val Ser Asn Asp Val
 435 440 445
 55
 Ser Asp Gly Asn Phe Trp Ala Asn Ala Arg Leu Ser Ala Asn Gly Trp
 450 455 460

EP 1 882 731 A1

Gly Lys Ser Val Asp Ile Leu Gly Ala Glu Lys Leu Thr Met Asp Val
 465 470 475 480
 5
 Ile Val Asp Glu Pro Thr Thr Val Ala Ile Ala Ala Ile Pro Gln Ser
 485 490 495
 10
 Ser Lys Ser Gly Trp Ala Asn Pro Glu Arg Ala Val Arg Val Asn Ala
 500 505 510
 Glu Asp Phe Val Gln Gln Thr Asp Gly Lys Tyr Lys Ala Gly Leu Thr
 515 520 525
 15
 Ile Thr Gly Glu Asp Ala Pro Asn Leu Lys Asn Ile Ala Phe His Glu
 530 535 540
 Glu Asp Asn Asn Met Asn Asn Ile Ile Leu Phe Val Gly Thr Asp Ala
 545 550 555 560
 20
 Ala Asp Val Ile Tyr Leu Asp Asn Ile Lys Val Ile Gly Thr Glu Val
 565 570 575
 25
 Glu Ile Pro Val Val His Asp Pro Lys Gly Glu Ala Val Leu Pro Ser
 580 585 590
 Val Phe Glu Asp Gly Thr Arg Gln Gly Trp Asp Trp Ala Gly Glu Ser
 595 600 605
 30
 Gly Val Lys Thr Ala Leu Thr Ile Glu Glu Ala Asn Gly Ser Asn Ala
 610 615 620
 35
 Leu Ser Trp Glu Phe Gly Tyr Pro Glu Val Lys Pro Ser Asp Asn Trp
 625 630 635 640
 Ala Thr Ala Pro Arg Leu Asp Phe Trp Lys Ser Asp Leu Val Arg Gly
 645 650 655
 40
 Glu Asn Asp Tyr Val Ala Phe Asp Phe Tyr Leu Asp Pro Val Arg Ala
 660 665 670
 45
 Thr Glu Gly Ala Met Asn Ile Asn Leu Val Phe Gln Pro Pro Thr Asn
 675 680 685
 Gly Tyr Trp Val Gln Ala Pro Lys Thr Tyr Thr Ile Asn Phe Asp Glu
 690 695 700
 50
 Leu Glu Glu Ala Asn Gln Val Asn Gly Leu Tyr His Tyr Glu Val Lys
 705 710 715 720
 55
 Ile Asn Val Arg Asp Ile Thr Asn Ile Gln Asp Asp Thr Leu Leu Arg
 725 730 735

EP 1 882 731 A1

5 Asn Met Met Ile Ile Phe Ala Asp Val Glu Ser Asp Phe Ala Gly Arg
 740 745 750
 Val Phe Val Asp Asn Val Arg Phe Glu Gly Ala Ala Thr Thr Glu Pro
 755 760 765
 10 Val Glu Pro Glu Pro Val Asp Pro Gly Glu Glu Thr Pro Pro Val Asp
 770 775 780
 15 Glu Lys Glu Ala Lys Lys Glu Gln Lys Glu Ala Glu Lys Glu Glu Lys
 785 790 795 800
 Glu Ala Val Lys Glu Glu Lys Lys Glu Ala Lys Glu Glu Lys Lys Ala
 805 810 815
 20 Val Lys Asn Glu Ala Lys Lys Lys
 820
 25
 30
 35
 40
 45
 50
 55

<First Sequence;protein/1;Bacillus sp.>

AEGNTREDNFKHLLGNDNVKRPSEAGALQLQEVDGQMTLVDQHGEKIQLR
 GMSTHGLQWFPEILNDNAYKALANDWESNMIRLAMYVGENGYSNPPELIK
 SRVIKIDLAENDMYVIVDWHVHAPGDPRDPVYAGAEDFFRDIAALYPN
 NPHIYELANGPSSNNNGGAGIPNNEEGWNAVKEYADPIVEMLRDSGNAD
 DNIIIVGSPNWSQRPDLAADNPINDHHTMYTVHFYTGSHAASTESYPPE
 PNSEKGNVMSNTRYALENGVAVFATEWGTSGANGDGGPYFDEADVWIEFL
 NENNISWANWSLTNKNEVSGAFTPFELGKSNAATNLDPGPDHVWAPPELSL
 SGEYVRARIKGVNYEPIDRTKYTKVLWDFNDGTKQGFGVNSDSPNKELIA
 VDNENNTLKVSGLDVSDNDVSDGNFWANARLSADGWGKSVDILGAEKLTMD
 VIVDEPTTVAIAAIPQSSKSGWANPERAVRVNAEDFVQQTGDGKYKAGLTI
 TGEDAPNLKNIAFHEEDNNMNNIILFVGTDAAADVILYLDNIKVGITEVEIP
 VVHDPKGEAVLPSVFEDGTRQGWWDWAGESGVKTALTIEEANGSNALSWEF
 GYPEVKPSDNWATAPRLDFWKSDLVRGENDYVAFDFYLDPVPRATEGAMNI
 NLVFQPPTNGYWVQAPKTYTINFDELEEEANQVNGLYHYEVKINVRDITNI
 QDDTLLRNMMIIFADVESDFAGRVFVDNVRFEGAATTEPVEPEPVDPGEE
 TPPVDEKEAKKEQKEAEKEEKEE

<Second Sequence;protein/1;Bacillus sp. KSM-S237>

MMLRKKTKQLISSILILVLLSLFPAALAAEGNTREDNFKHLLGNDNVKR
 PSEAGALQLQEVDGQMTLVDQHGEKIQLRGMSTHGLQWFPEILNDNAYKA
 LSNDWDSNMIRLAMYVGENGYSNPPELIKQRVIDGIELAIENDMYVIVDW
 HVHAPGDPRDPVYAGAKDFFREIAALYPNNPHIYELANEPSSNNNGGAG
 IPNNEEGWKAVKEYADPIVEMLRKSGNADDNIIIVGSPNWSQRPDLAADN
 PIDDHHTMYTVHFYTGSHAASTESYPSETPNSEKGNVMSNTRYALENGVA
 VFATEWGTSGASGDGGPYFDEADVWIEFLNENNISWANWSLTNKNEVSGA
 FTPFELGKSNAATNLDPGPDHVWAPPELSLSGEYVRARIKGVNYEPIDRTK
 YTKVLWDFNDGTKQGFGVNSDSPNKELIAVDNENNTLKVSGLDVSDNDVSD
 GNFWANARLSANGWGKSVDILGAEKLTMDVIVDEPTTVAIAAIPQSSKSG
 WANPERAVRVNAEDFVQQTGDGKYKAGLTTITGEDAPNLKNIAFHEEDNNM
 NIILFVGTDAAADVILYLDNIKVGITEVEIPVVHDPKGEAVLPSVFEDGTRQ
 GWDWAGESGVKTALTIEEANGSNALSWEFGYPEVKPSDNWATAPRLDFWK
 SDLVRGENDYVAFDFYLDPVPRATEGAMNINLVFQPPTNGYWVQAPKTYTI
 NFDELEEEANQVNGLYHYEVKINVRDITNIQDDTLLRNMMIIFADVESDFA
 GRVFVDNVRFEGAATTEPVEPEPVDPGEETPPVDEKEAKKEQKEAEKEEK
 EAVKEEKKEAKKEKKAVKNEAKKK

Claims

1. A detergent composition comprising an alkaline bacterial enzyme exhibiting endo-beta-1,4-glucanase activity (E.C. 3.2.1.4), up to 10 wt% aluminosilicate (anhydrous basis) and/or phosphate builder, the composition having a reserve alkalinity of greater than 4.
2. A composition according to claim 1 wherein enzyme is a bacterial polypeptide endogenous to a member of the genus *Bacillus*
3. A composition according to claims 1-2 wherein the enzyme is a polypeptide containing (i) at least one family 17 carbohydrate binding module and/or (ii) at least one family 28 carbohydrate binding module.
4. A composition according to claims 1-3 wherein the enzyme comprises a polypeptide endogenous to one of the following *Bacillus* species selected from the group consisting of: AA349 (DSM 12648), KSM S237, 1139, KSM 64, KSM N131, KSM 635 (FERM BP 1485), KSM 534 (FERM BP 1508), KSM 53 (FERM BP 1509), KSM 577 (FERM

BP 1510), KSM 521 (FERM BP 1507), KSM 580 (FERM BP 1511), KSM 588 (FERM BP 1513), KSM 597 (FERM BP 1514), KSM 522 (FERM BP 1512), KSM 3445 (FERM BP 1506) or KSM 425 (FERM BP 1505).

5. A composition according to claims 1-4 wherein the enzyme is selected from:

- (i) the endoglucanase having the amino acid sequence of positions 1 to position 773 of SEQ ID NO:1;
- (ii) an endoglucanase having a sequence of at least 90%, preferably 94%, more preferably 97% and even more preferably 99%, 100% identity to the amino acid sequence of position 1 to position 773 of SEQ ID NO:1; or a fragment thereof has endo-beta-1,4-glucanase activity, when identity is determined by GAP provided in the GCG program using a GAP creation penalty of 3.0 and GAP extension penalty of 0.1.

6. A composition according to claims 1-4 wherein the enzyme is an alkaline endoglucanase variant obtained by substituting the amino acid residue of a cellulase having an amino acid sequence exhibiting at least 90%, preferably 95%, more preferably 98%, 100% identity with the amino acid sequence represented by SEQ. ID NO:2 at (a) position 10, (b) position 16, (c) position 22, (d) position 33, (e) position 39, (f) position 76, (g) position 109, (h) position 242, (i) position 263, (j) position 308, (k) position 462, (l) position 466, (m) position 468, (n) position 552, (o) position 564, or (p) position 608 in SEQ ID NO:1 or at a position corresponding thereto with another amino acid residue.

7. A composition according to claim 5 wherein the enzyme is **characterised by** the following substitutions:

- (a) at position 10: glutamine, alanine, proline or methionine, preferably glutamine;
- (b) at position 16: asparagine or arginine, preferably asparagine;
- (c) at position 22: proline;
- (d) at position 33: histidine;
- (e) at position 39: alanine, threonine or tyrosine, preferably alanine;
- (f) at position 76: histidine, methionine, valine, threonine or alanine, preferably histidine;
- (g) at position 109: isoleucine, leucine, serine or valine, preferably isoleucine;
- (h) at position 242: alanine, phenylalanine, valine, serine, aspartic acid, glutamic acid, leucine, isoleucine, tyrosine, threonine, methionine or glycine, preferably alanine, phenylalanine or serine;
- (i) at position 263: isoleucine, leucine, proline or valine, preferably isoleucine;
- (j) at position 308: alanine, serine, glycine or valine, preferably alanine;
- (k) at position 462: threonine, leucine, phenylalanine or arginine, preferably threonine;
- (l) at position 466: leucine, alanine or serine, preferably leucine;
- (m) at position 468: alanine, aspartic acid, glycine or lysine, preferably alanine;
- (n) at position 552: methionine;
- (o) at position 564: valine, threonine or leucine, preferably valine; and
- (p) at position 608: isoleucine or arginine, preferably isoleucine.

8. A composition according to claims 6 and 7 wherein the enzyme is selected from the group consisting of the following endoglucanase variants: Egl-237, Egl-1139, Egl-64, Egl-N131b and mixtures thereof.

9. A composition according to claims 1-4 wherein the enzyme is an alkaline cellulase K having the following physical and chemical properties:

- (1) Activity: Having a C_x enzymatic activity of acting on carboxymethyl cellulose along with a weak C₁ enzymatic activity and a weak beta-glucosidase activity;
- (2) Specificity on Substrates: Acting on carboxymethyl cellulose(CMC), crystalline cellulose, Avicell, cellobiose, and p-nitrophenyl cellobioside(PNPC);
- (3) Having a working pH in the range of 4 to 12 and an optimum pH in the range of 9 to 10;
- (4) Having stable pH values of 4.5 to 10.5 and 6.8 to 10 when allowed to stand at 40°C for 10 minutes and 30 minutes, respectively;
- (5) Working in a wide temperature range of from 10 to 65°C with an optimum temperature being recognized at about 40°C;
- (6) Influences of chelating agents: The activity not impeded with ethylenediamine tetraacetic acid (EDTA), ethyleneglycol-bis-(β-aminoethylether) N,N,N',N"-tetraacetic acid (EGTA), N,N-bis(carboxymethyl)glycine (nitrilotriacetic acid) (NTA), sodium tripolyphosphate (STPP) and zeolite;
- (7) Influences of surface active agents: Undergoing little inhibition of activity by means of surface active agents such as sodium linear alkylbenzenesulfonates (LAS), sodium alkylsulfates (AS), sodium polyoxyethylene alkyl-

sulfates (ES), sodium alphaolefinsulfonates (AOS), sodium alpha-sulfonated aliphatic acid esters (alpha-SFE), sodium alkylsulfonates (SAS), polyoxyethylene secondary alkyl ethers, fatty acid salts (sodium salts), and dimethyldialkylammonium chloride;

(8) Having a strong resistance to proteinases; and

(9) Molecular weight (determined by gel chromatography): Having a maximum peak at $180,000 \pm 10,000$.

10. A composition according to claim 8 wherein the alkaline cellulase K is obtained by isolation from a culture product of *Bacillus* sp KSM-635.

11. A composition according to claims 1-4 wherein the enzyme is selected from the group consisting of:

Alkaline Cellulase K-534 from KSM 534, FERM BP 1508,
Alkaline Cellulase K-539 from KSM 539, FERM BP 1509,
Alkaline Cellulase K-577 from KSM 577, FERM BP 1510,
Alkaline Cellulase K-521 from KSM 521, FERM BP 1507,
Alkaline Cellulase K-580 from KSM 580, FERM BP 1511,
Alkaline Cellulase K-588 from KSM 588, FERM BP 1513,
Alkaline Cellulase K-597 from KSM 597, FERM BP 1514,
Alkaline Cellulase K-522 from KSM 522, FERM BP 1512,
Alkaline Cellulase E-II from KSM 522, FERM BP 1512,
Alkaline Cellulase E-III from KSM 522, FERM BP 1512.
Alkaline Cellulase K-344 from KSM 344, FERM BP 1506,
Alkaline Cellulase K-425 from KSM 425, FERM BP 1505, and mixtures thereof.

12. A composition according to any of the preceding claims wherein the bacterial alkaline enzyme exhibiting endo-beta-1,4-glucanase activity is comprised at a level of from 0.00005% to 0.15%, preferably from 0.0002% to 0.02%, or more preferably from 0.0005% to 0.01% by weight of pure enzyme.

13. A detergent composition according to any of the preceding claims comprising less than 8 wt% aluminosilicate (anhydrous basis) and/or phosphate builder.

14. A detergent composition according to any of the preceding claims comprising less than 5 wt% aluminosilicate (anhydrous basis) and/or phosphate builder.

15. A detergent composition according to any preceding claim having a reserve alkalinity greater than 7.5.


16. A detergent composition according to any preceding claim comprising from 0.1 to 40 wt% alkoxyated alkyl sulphate surfactant and/or from 0.1 to 40 wt% C₁₋₄ alkyl ester sulphonate, preferably methyl ester sulphonate (MES).

17. A detergent composition according to any preceding claim comprising a suds booster in an amount from 0.05 to 2 wt%, preferably selected from fatty acid amides, fatty acid alkanolamides, betaines, sulfobetaines and amine oxides or mixtures thereof.

18. A detergent composition according to any preceding claim comprising from 0.05 to 5, preferably from 0.1 to 1 wt% soil release polymer, preferably selected from modified cellulose ethers such as methyl hydroxyethyl cellulose (MHEC) or polyester-based soil release polymers or mixtures thereof.

19. A detergent composition according to any of the preceding claims comprising from 0.01wt% to 10wt% of a chelant, preferably selected from the group consisting of 4,5-dihydroxy-m-benzenedisulfonic acid, disodium salt (Tiron®); hydroxyethane-dimethylene-phosphonic acid (HEDP); 2-phosphonobutane-1,2,4-tricarboxylic acid (PBTC) and mixtures thereof.

20. A detergent composition according to any of the preceding claims comprising an optical brightener of the following structure, wherein R1 and R2, together with the nitrogen atom linking them, form an unsubstituted or C1-C4 alkyl-substituted morpholino, piperidine or pyrrolidine ring :



21. A detergent composition according to any preceding claim further comprising a lipase enzyme (E.C. 3.1.1.3).
22. A detergent composition according to any preceding claim which is a solid detergent composition, preferably granular.
23. A washing process comprising laundering textile articles in an aqueous solution comprising the detergent composition according to any preceding claim.
24. A washing process according to claim 10 in which the aqueous solution is at a temperature below 30°C.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 06 11 5574

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
D,Y	WO 2004/053039 A (NOVOZYMES AS [DK]; GIBSON KEITH [DK]; HANSEN LONE [DK]) 24 June 2004 (2004-06-24) * claims 1-27 *	1-25	INV. C11D3/06 C11D3/12 C11D3/386
D,Y	WO 02/099091 A (NOVOZYMES AS [DK]; OUTTRUP HELLE [DK]; SCHUELEIN MARTIN [DK]; ESKELUND) 12 December 2002 (2002-12-12) * claims 1,27 *	1-25	
A	WO 99/09133 A (PROCTER & GAMBLE [US]; BETTIOL JEAN LUC PHILIPPE [BE]; THOEN CHRISTIAA) 25 February 1999 (1999-02-25) * page 5, line 28 - line 32; examples 1,2,5 *	1-25	
			TECHNICAL FIELDS SEARCHED (IPC)
			C11D
The present search report has been drawn up for all claims			
Place of search The Hague		Date of completion of the search 10 December 2007	Examiner Richards, Michael
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

2

EPO FORM 1503 03.82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 06 11 5574

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

10-12-2007

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 2004053039	A	24-06-2004	AU 2003302905 A1	30-06-2004
			BR 0317124 A	25-10-2005
			CN 1738899 A	22-02-2006
			EP 1572996 A2	14-09-2005
			JP 2006509850 T	23-03-2006
			MX PA05006071 A	30-09-2005
			US 2006035800 A1	16-02-2006

WO 02099091	A	12-12-2002	CN 1633496 A	29-06-2005
			EP 1399543 A2	24-03-2004
			JP 2004536593 T	09-12-2004
			MX PA03011194 A	26-02-2004

WO 9909133	A	25-02-1999	AT 363527 T	15-06-2007
			AT 230013 T	15-01-2003
			AT 276344 T	15-10-2004
			AT 332958 T	15-08-2006
			AU 7832798 A	08-03-1999
			AU 7833398 A	08-03-1999
			AU 7833498 A	08-03-1999
			AU 7958198 A	08-03-1999
			AU 8064198 A	08-03-1999
			AU 8064298 A	08-03-1999
			AU 8065198 A	08-03-1999
			AU 8065398 A	08-03-1999
			BR 9811186 A	25-07-2000
			BR 9811187 A	25-07-2000
			BR 9811189 A	25-07-2000
			BR 9811190 A	18-07-2000
			BR 9811191 A	18-07-2000
			BR 9811192 A	18-07-2000
			BR 9811195 A	25-07-2000
			BR 9811196 A	25-07-2000
			CA 2299410 A1	25-02-1999
			CA 2300696 A1	25-02-1999
			CA 2301156 A1	25-02-1999
			CA 2301167 A1	25-02-1999
			CA 2301168 A1	25-02-1999
			CA 2301200 A1	25-02-1999
			CA 2301205 A1	25-02-1999
			CA 2301404 A1	25-02-1999
			CN 1306566 A	01-08-2001
			CN 1336953 A	20-02-2002
			CN 1469919 A	21-01-2004
			CN 1301294 A	27-06-2001
			CN 1276005 A	06-12-2000

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 06 11 5574

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

10-12-2007

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9909133 A		CN 1276824 A	13-12-2000
		CN 1276825 A	13-12-2000
		CN 1276826 A	13-12-2000
		DE 69810309 D1	30-01-2003
		DE 69810309 T2	16-10-2003
		DE 69826294 D1	21-10-2004
		DE 69826294 T2	17-11-2005
		DE 69835214 T2	21-06-2007
		DK 1009795 T3	24-01-2005
		EP 0896998 A1	17-02-1999
		ES 2185172 T3	16-04-2003
		ES 2227845 T3	01-04-2005
		ES 2268780 T3	16-03-2007
		HU 0003670 A2	28-05-2001
		ID 23442 A	20-04-2000
		JP 2001515126 T	18-09-2001

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 02099091 A [0003] [0013] [0013]
- WO 04053039 A [0003]
- EP 265832 A [0003] [0012] [0013]
- EP 1350843 A [0003] [0012] [0012] [0012] [0012] [0013] [0013] [0013]
- WO 2002099091 A [0012]
- WO 2004053039 A [0012]
- EP 0271044 A [0012] [0012] [0012] [0012] [0012] [0012] [0012] [0012] [0012]
- JP 2000047237 A [0013]
- EP 271004 A, Kao [0013]
- US 6020303 A [0030] [0040]
- US 6060443 A [0030]
- WO 9905243 A [0030]
- WO 9905242 A [0030]
- WO 9905244 A [0030]
- WO 9905082 A [0030]
- WO 9905084 A [0030]
- WO 9905241 A [0030]
- WO 9907656 A [0030]
- WO 0023549 A [0030]
- WO 0023548 A [0030]
- US 6150322 A [0040]
- US 6153577 A [0040]
- US 6093856 A [0040]
- US 4565647 A [0040]
- US 4483780 A [0040]
- US 4483779 A [0040]
- US 5332528 A [0040]
- WO 9206162 A [0040]
- WO 9319146 A [0040]
- WO 9319038 A [0040]
- WO 9409099 A [0040]
- US 6482994 B [0040]
- WO 0142408 A [0040]
- WO 04041982 A [0041]
- US 6136769 A [0043]
- US 6004922 A [0043]
- WO 9835002 A [0043]
- WO 9835003 A [0043]
- WO 9835004 A [0043]
- WO 9835005 A [0043]
- WO 9835006 A [0043]
- US 4228042 A [0043]
- US 4239660 A [0043]
- US 4260529 A [0043]
- US 6022844 A [0043]
- US 6221825 B [0043]
- WO 0047708 A [0043]
- WO 9407974 P [0055]
- WO 9407984 P [0055]
- WO 9407985 P [0055]
- WO 9504806 P [0055]
- WO 9703379 P [0055]
- US 6770610 B, Clariant [0055]
- EP 0324568 A [0055]
- EP 0768370 A [0055]
- US 3936537 A, Baskerville [0073]
- GB 9705617 A, Trinh [0073]
- US 360568 A [0076]
- US 5360569 A [0076]
- US 5370826 A [0076]
- WO 9914245 A [0083]
- WO 9711151 A [0086]
- US 6312936 B1 [0091]

Non-patent literature cited in the description

- **Y. BOURNE ; B. HENRISSAT.** Glycoside hydrolases and glycosyltransferases: families and functional modules. *Current Opinion in Structural Biology*, 2001, 593-600 [0011]
- **A.B. BORASTON et al.** Identification and glucan-binding properties of a new carbohydrate-binding module family. *Biochemical Journal*, 2002, vol. 361, 35-40 [0011]
- **HAKAMADA et al.** Bacillus sp. strain KSM-S237 (FERM BP-7875. *Biosci. Biotechnol. Biochem.*, 2000, vol. 64, 2281-2289 [0013]
- **FUKUMORI et al.** *J. Gen. Microbiol.*, vol. 132, 2329-2335 [0013]
- **SUMITOMO et al.** *Biosci. Biotechnol. Biochem.*, 1992, vol. 56, 872-877 [0013]
- **S. ITO.** *Extremophiles*, 1997, vol. 1, 61-66 [0013]
- **S. ITO et al.** *Agric Biol Chem*, 1989, vol. 53, 1275-1278 [0013]
- **M.K. NAGARAJAN ; W.F. MASLER.** *Cosmetics and Toiletries*, 1989, vol. 104, 71-73 [0055]
- **W. M. LINFIELD.** *Tenside Surf. Det.*, 1990, vol. 27, 159-161 [0055]

- **R.G. BISTLINE et al.** *J. Am. Oil Chem. Soc.*, 1972, vol. 49, 63-69 **[0055]**