

FASCICULE DE BREVET D'INVENTION

21 Numéro de dépôt : 1201600164
PCT/US2014/064412

22 Date de dépôt : 06/11/2014

30 Priorité(s) :

US n° 61/902,544 du 11/11/2013

24 Délivré le : 06/01/2021

45 Publié le : 26.02.2021

73 Titulaire(s) :

GILEAD SCIENCES, INC.,
333 Lakeside Drive,
FOSTER CITY, CA 94404 (US)

72 Inventeur(s) :

SIEGEL Dustin (US);
DOERFFLER Edward (US);
MACKMAN Richard L. (US);
CLARKE Michael O'neil Hanrahan (US)


74 Mandataire :

54 Titre : Pyrrolo[1,2-f][1,2,4]Triazines useful for treating respiratory syncytial virus infections.

57 Abrégé :

Provided herein are formulations, methods and substituted tetrahydrofuranyl-pyrrolo[1,2-f][1,2,4]-triazine-4-amine compounds of formula (I) for treating Pneumovirinae virus infections, including respiratory syncytial virus infections, as well as methods and intermediates for synthesis of tetrahydrofuranyl-pyrrolo[1,2-f][1,2,4]triazine-4-amine compounds.

Formula (I)



PYRROLO[1,2-*f*][1,2,4]TRIAZINES USEFUL FOR TREATING
RESPIRATORY SYNCYTIAL VIRUS INFECTIONS

FIELD

5 Provided herein are substituted tetrahydrofuranyl-pyrrolo[1,2-*f*][1,2,4]triazine-4-amine compounds, methods and pharmaceutical formulations for treating *Pneumovirinae* virus infections, particularly including respiratory syncytial virus infections, as well as methods and intermediates useful for preparing the compounds.

BACKGROUND

10 *Pneumovirinae* viruses are negative-sense, single-stranded, RNA viruses that are responsible for many prevalent human and animal diseases. The *Pneumovirinae* sub-family of viruses is a part of the family *Paramyxoviridae* and includes human respiratory syncytial virus (HRSV). Almost all children will have had an HRSV infection by their second birthday. HRSV is the major cause of lower respiratory tract infections in infancy and childhood with
15 0.5% to 2% of those infected requiring hospitalization. The elderly and adults with chronic heart, lung disease or those that are immunosuppressed also have a high risk for developing severe HRSV disease (<http://www.cdc.gov/rsv/index.html>). No vaccine to prevent HRSV infection is currently available. The monoclonal antibody palivizumab is available for immunoprophylaxis, but its use is restricted to infants at high risk, e.g., premature infants or
20 those with either congenital heart or lung disease, and the cost for general use is often prohibitive. In addition, nucleoside analog ribavirin has been approved as the only antiviral agent to treat HRSV infections but has limited efficacy. Therefore, there is a need for anti-*Pneumovirinae* therapeutics.

 Examples of pyrrolo[2,3-*d*]pyrimidine compounds useful for treating viral infections
25 are described in U.S. 2012/0009147 A1 (Cho et al.), U.S. 2012/0020921 A1 (Cho et al.), WO 2008/089105 A2 (Babu et al.), WO 2008/141079 A1 (Babu et al.), WO 2009/132135 A1 (Butler et al.), WO 2010/002877 A2 (Francom), WO 2011/035231 A1 (Cho et al.), WO 2011/035250 A1 (Butler et al.), WO 2011/150288 A1 (Cho et al.), WO 2012/012465 (Cho et al.), WO 2012/012776 A1 (Mackman et al.), WO 2012/037038 (Clarke et al.), WO
30 2012/087596 A1 (Delaney et al.), and WO 2012/142075 A1 (Girijavallabhan et al.).


There remains a need for new antiviral agents useful in treating *Paramyxoviridae* viral infections, including *Pneumovirinae* viral infections, such as HRSV infections, that are effective and have acceptable toxicity profiles.

5

SUMMARY

Provided are compounds, methods, and pharmaceutical formulations for the treatment of infections caused by the *Pneumovirinae* virus family, including treatment of infections caused by human respiratory syncytial virus.

10 Provided is a compound of the Formula (I), or a pharmaceutically acceptable salt thereof:



wherein:

R^1 is H or F;



R^2 is H or F;

15 R^3 is OH or F;

R^4 is CN, C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_3 - C_4 cycloalkyl, azido, halogen, or C_1 - C_2 haloalkyl;

R^6 is OH;

R^5 is selected from the group of H and:




wherein:

n' is selected from 1, 2, 3, and 4;

5 R^8 is selected from C_1 - C_8 alkyl, - O - C_1 - C_8 alkyl, benzyl, - O -benzyl, - CH_2 - C_3 - C_6 cycloalkyl, - O - CH_2 - C_3 - C_6 cycloalkyl, and CF_3 ;

R^9 is selected from phenyl, 1-naphthyl, 2-naphthyl,



R^{10} is selected from H and CH_3 ;

10 R^{11} is selected from H or C_1 - C_6 alkyl;

R^{12} is selected from H, C_1 - C_8 alkyl, benzyl, C_3 - C_6 cycloalkyl, and - CH_2 - C_3 - C_6 cycloalkyl.

DETAILED DESCRIPTION

An embodiment herein comprises a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^1 is H, and all other variables, including R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , and n' are as defined above for Formula (I).

5 Another embodiment herein comprises a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein R^2 is H, and all other variables, including R^1 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , and n' are as defined above for Formula (I).


A further embodiment herein comprises a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein both R^1 and R^2 are H, and all other
10 variables, including R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , and n' are as defined above for Formula (I).

Still another embodiment herein comprises a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein both R^1 , R^2 , and R^5 are H, and all other variables, including R^3 , R^4 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , and n' are as defined above for
15 Formula (I).

Another separate embodiment herein comprises a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein both R^1 and R^2 are H, R^3 is OH, and all other variables, including R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , and n' are as defined above for Formula (I).

20 Another separate embodiment herein comprises a compound of Formula (I), or a pharmaceutically acceptable salt thereof, wherein both R^1 and R^2 are H, R^3 is F, and all other variables, including R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , and n' are as defined above for Formula (I).

Another embodiment provided herein comprises a compound of Formula (II), or a
25 pharmaceutically acceptable salt thereof:





wherein:

R^3 is OH or F;

R^4 is CN, C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_3 - C_4 cycloalkyl, azido, halogen,
5 or C_1 - C_2 haloalkyl;

R^5 is selected from the group of H and:



wherein:

10 n' is selected from 1, 2, 3, and 4;

R^8 is selected from C_1 - C_8 alkyl, -O- C_1 - C_8 alkyl, benzyl, -O-benzyl, -CH₂- C_3 - C_6 cycloalkyl, -O-CH₂- C_3 - C_6 cycloalkyl, and CF₃;

R⁹ is phenyl;

R¹⁰ is selected from H and CH₃;

R¹¹ is selected from H or C₁-C₆ alkyl;

R¹² is selected from H, C₁-C₈ alkyl, benzyl, C₃-C₆ cycloalkyl, and -CH₂-C₃-C₆
5 cycloalkyl.

A further embodiment comprises a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein:

R³ is OH or F;

R⁴ is CN, methyl, ethyl, ethenyl, ethynyl, azido, F, Cl, -CH₂Cl, -CH₂F, -CHF₂, or -CF₃;

10 and R⁵ and all other groups are as defined for Formula (II).

Also provided is an embodiment comprising a compound of Formula (II) or a pharmaceutically acceptable salt thereof, as described above, wherein R³ is F.

Also provided is an embodiment comprising a compound of Formula (II) or a pharmaceutically acceptable salt thereof, as described above, wherein R³ is OH.

15 Also provided is an embodiment comprising a compound of Formula (II) or a pharmaceutically acceptable salt thereof, as described above, wherein R³ is F and R⁴ is CN.

Also provided is an embodiment comprising a compound of Formula (II) or a pharmaceutically acceptable salt thereof, as described above, wherein R³ is OH and R⁴ is CN.

20 Also provided is an embodiment comprising a compound of Formula (II) or a pharmaceutically acceptable salt thereof, as described above, wherein both R¹ and R² are H, R³ is F, and R⁴ is methyl, ethyl, vinyl, or ethynyl.

Also provided is an embodiment comprising a compound of Formula (II) or a pharmaceutically acceptable salt thereof, as described above, wherein R³ is OH, and R⁴ is
25 methyl, ethyl, vinyl, or ethynyl.

Also provided is an embodiment comprising a compound of Formula (II) or a pharmaceutically acceptable salt thereof, as described above, wherein R^3 is F and R^4 is halomethyl.

Also provided is an embodiment comprising a compound of Formula (II) or a pharmaceutically acceptable salt thereof, as described above, wherein R^3 is OH and R^4 is halomethyl.


Also provided is an embodiment comprising a compound of Formula (II) or a pharmaceutically acceptable salt thereof, as described above, wherein R^5 is H.


Also provided is an embodiment comprising a compound of Formula (II) or a pharmaceutically acceptable salt thereof, as described above, wherein each of R^5 is H, R^3 is OH, and R^4 is methyl, ethyl, vinyl, or ethynyl.

Also provided is an embodiment comprising a compound of Formula (II) or a pharmaceutically acceptable salt thereof, as described above, wherein R^5 is H, R^3 is F, and R^4 is halomethyl.

Also provided is an embodiment comprising a compound of Formula (II) or a pharmaceutically acceptable salt thereof, as described above, wherein R^5 are H, R^3 is OH, and R^4 is halomethyl.

Within each of the embodiments described above comprising a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein R^5 may be other than H, there is a further embodiment wherein all other variables are as described for the embodiment and R^5 is selected from the group of:





wherein:

R^8 is selected from C_1 - C_8 alkyl, $-O$ - C_1 - C_8 alkyl, benzyl, and $-CH_2$ - C_3 - C_6 cycloalkyl;

and

5 R^{12} is selected from C_1 - C_8 alkyl, benzyl, C_3 - C_6 cycloalkyl, and $-CH_2$ - C_3 - C_6 cycloalkyl.

Within each of the embodiments described immediately above there is a further embodiment comprising a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein all other variables are as described immediately above, except that R^8 and R^9 are each selected from C_1 - C_8 alkyl. Within each of the embodiments described in the last sentence there is a further embodiment comprising a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein all other variables are as described immediately above, except that R^8 and R^9 are each selected from C_1 - C_6 alkyl. Within each of the embodiments described in the last sentence there is a further embodiment comprising a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein all other variables are as described immediately above, except that R^8 and R^9 are each selected from C_1 - C_5 alkyl. Within each of the embodiments described in the last sentence there is a further embodiment comprising a compound of Formula (II), or a pharmaceutically acceptable salt thereof, wherein all other variables are as described immediately above, except that R^8 and R^9 are each selected from C_1 - C_4 alkyl.

20 Within each of the embodiments described herein comprising a compound of Formula (I) or of Formula (II) there is a further embodiment wherein all variables are as defined for the particular embodiment and further comprising the proviso that when R^3 is F, R^4 is not methyl.

Definitions

The terms halo and halogen refer to halogen atoms selected from F, Cl, Br, and I .

“Azido” refers to an azide group, i.e. the group $-N_3$. The term “n” as used herein refers to an integer, such as an integer selected from 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, and 20, i.e. 2 to 20 or 2-20. In some instances, “n” refers to groups of integers such as 1 to 3, 1 to 4, 1 to 6, 1 to 8, 2 to 4, 2 to 6, 2 to 8, etc.

The term “haloalkyl” as used herein refers to an alkyl as defined herein, wherein one or more hydrogen atoms are each replaced by a halo substituent. For example, a (C_1-C_6) haloalkyl is a (C_1-C_6) alkyl wherein one or more of the hydrogen atoms have been replaced by a halo substituent. Such a range includes one halo substituent on the alkyl group t to complete halogenation of the alkyl group.

The term “ (C_{1-n}) haloalkyl” as used herein, wherein n is an integer, either alone or in combination with another radical, is intended to mean an alkyl radical having 1 to n carbon atoms as defined above wherein one or more hydrogen atoms are each replaced by a halo substituent. Examples of (C_{1-n}) haloalkyl, wherein n is 2 include, but are not limited to, chloromethyl, chloroethyl, dichloroethyl, bromomethyl, bromoethyl, dibromoethyl, fluoromethyl, difluoromethyl, trifluoromethyl, fluoroethyl and difluoroethyl.

The term “ (C_{1-n}) alkyl” as used herein, wherein n is an integer, either alone or in combination with another radical, is intended to mean acyclic, straight or branched chain alkyl radicals containing from 1 to n carbon atoms. “ (C_{1-4}) alkyl” includes, but is not limited to, methyl, ethyl, propyl (*n*-propyl), butyl (*n*-butyl), 1-methylethyl (*iso*-propyl), 1-methylpropyl (*sec*-butyl), 2-methylpropyl (*iso*-butyl), and 1,1-dimethylethyl (*tert*-butyl or *t*-butyl). The abbreviation Me denotes a methyl group; Et denotes an ethyl group, Pr denotes a propyl group, iPr denotes a 1-methylethyl group, Bu denotes a butyl group and tBu denotes a 1,1-dimethylethyl group.

The term “alkyl” refers to a hydrocarbon containing normal, secondary, or tertiary atoms. For example, an alkyl group can have 1 to 4 carbon atoms (*i.e.*, (C_1-C_4) alkyl), 1 to 3 carbon atoms (*i.e.*, (C_1-C_3) alkyl), or 1 or 2 carbon atoms (*i.e.*, (C_1-C_2) alkyl). Examples of suitable alkyl groups include, but are not limited to, methyl (Me, $-CH_3$), ethyl (Et, $-CH_2CH_3$), 1-propyl ($\underline{n}$ -Pr, $\underline{n}$ -propyl, $-CH_2CH_2CH_3$), 2-propyl ($\underline{i}$ -Pr, $\underline{i}$ -propyl, $-CH(CH_3)_2$), 1-butyl ($\underline{n}$ -Bu, $\underline{n}$ -butyl, $-CH_2CH_2CH_2CH_3$), 2-methyl-1-propyl ($\underline{i}$ -Bu, $\underline{i}$ -butyl, $-CH_2CH(CH_3)_2$), 2-butyl ($\underline{s}$ -Bu, $\underline{s}$ -butyl, $-CH(CH_3)CH_2CH_3$), and 2-methyl-2-propyl ($\underline{t}$ -Bu, $\underline{t}$ -butyl, $-C(CH_3)_3$). “Alkyl” also refers

to a saturated, branched or straight chain hydrocarbon radical having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkane. Typical alkyl radicals include, but are not limited to, methylene (-CH₂-), 1,1-ethyl (-CH(CH₃)-), 1,2-ethyl (-CH₂CH₂-), 1,1-propyl (-CH(CH₂CH₃)-), 1,2-propyl (-CH₂CH(CH₃)-), 1,3-propyl (-CH₂CH₂CH₂-), 1,4-butyl (-CH₂CH₂CH₂CH₂-), and the like.

"Alkenyl" is a straight or branched hydrocarbon containing normal, secondary or tertiary carbon atoms with at least one site of unsaturation, *i.e.* a carbon-carbon, *sp*² double bond. As examples, an alkenyl group can have 2 to 4 carbon atoms (*i.e.*, C₂-C₄ alkenyl), or 2 to 3 carbon atoms (*i.e.*, C₂-C₃ alkenyl). Examples of suitable alkenyl groups include, but are not limited to, ethylene or vinyl (-CH=CH₂) and allyl (-CH₂CH=CH₂).

The term "(C_{2-n})alkenyl", as used herein, wherein n is an integer, either alone or in combination with another radical, is intended to mean an unsaturated, acyclic straight or branched chain radical containing two to n carbon atoms, at least two of which are bonded to each other by a double bond. Examples of such radicals include, but are not limited to, ethenyl (vinyl), 1-propenyl, 2-propenyl, and 1-butenyl. Unless specified otherwise, the term "(C_{2-n})alkenyl" is understood to encompass individual stereoisomers where possible, including but not limited to (*E*) and (*Z*) isomers, and mixtures thereof. When a (C_{2-n})alkenyl group is substituted, it is understood to be substituted on any carbon atom thereof which would otherwise bear a hydrogen atom, unless specified otherwise, such that the substitution would give rise to a chemically stable compound, such as are recognized by those skilled in the art.

"Alkynyl" is a straight or branched hydrocarbon containing normal, secondary or tertiary carbon atoms with at least one site of unsaturation, *i.e.* a carbon-carbon, *sp* triple bond. For example, an alkynyl group can have 2 to 4 carbon atoms (*i.e.*, C₂-C₄ alkynyl) or 2 to 3 carbon atoms (*i.e.*, C₂-C₃ alkyne,). Examples of suitable alkynyl groups include, but are not limited to, acetylenic (-C≡CH), propargyl (-CH₂C≡CH), and the like.

The term "(C_{2-n})alkynyl", as used herein, wherein n is an integer, either alone or in combination with another radical, is intended to mean an unsaturated, acyclic straight or branched chain radical containing two to n carbon atoms, at least two of which are bonded to each other by a triple bond. Examples of such radicals in which n is 4 include, but are not limited to, ethynyl, 1-propynyl, 2-propynyl, and 1-butynyl. When a (C_{2-n})alkynyl group is substituted, it is understood to be substituted on any carbon atom thereof which would otherwise bear a hydrogen atom, unless specified otherwise, such that the substitution

would give rise to a chemically stable compound, such as are recognized by those skilled in the art.

The term cycloalkyl refers to a cyclic aliphatic group. The cycloalkyl groups herein may be referenced by the number of carbon atoms in their ring, such as "C₃-C₄ cycloalkyl" referring to a cycloalkyl ring with 3 or 4 carbon ring atoms or "C₃-C₆ cycloalkyl" indicating a cycloalkyl ring with 3, 4, 5, or 6 carbon ring atoms, i.e. a cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl ring.

The term "carbocycle" or "carbocyclyl" refers to a saturated (i.e., cycloalkyl) or partially unsaturated (e.g., cycloalkenyl, cycloalkadienyl, etc.) ring having the number of carbon atoms specified, such as 3 to 4 carbon atoms or 3 to 6 carbon atoms as a monocyclic ring system. In one embodiment the carbocycle is a monocycle comprising 3-6 ring carbons (i.e. (C₃-C₆)carbocycle). Non-limiting examples of monocyclic carbocycles include cyclopropyl, cyclobutyl, cyclopentyl, 1-cyclopent-1-enyl, 1-cyclopent-2-enyl, 1-cyclopent-3-enyl, cyclohexyl, 1-cyclohex-1-enyl, 1-cyclohex-2-enyl, 1-cyclohex-3-enyl, and cyclohexa-1,3-dienyl rings.

Each carbocyclyl group may be substituted by 0, 1, 2, or 3 substituents independently selected from halogen, -OH, -CN, -NO₂, -NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, C₁-C₆ alkyl, C₁-C₆ alkoxy, and -CF₃.

Pharmaceutical Formulations

Also provided herein is a pharmaceutical formulation comprising a pharmaceutically effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and a pharmaceutically acceptable carrier or excipient. Also provided are separate pharmaceutical formulations, each comprising a pharmaceutically effective amount of a compound of Formula (II) or one of the specific compounds of the examples herein, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and a pharmaceutically acceptable carrier or excipient.

The compounds herein are formulated with conventional carriers and excipients, which will be selected in accord with ordinary practice. Tablets will contain excipients, glidants, fillers, binders and the like. Aqueous formulations are prepared in sterile form, and when intended for delivery by other than oral administration generally will be isotonic. All formulations will optionally contain excipients such as those set forth in the "Handbook of

Pharmaceutical Excipients" (1986). Excipients include ascorbic acid and other antioxidants, chelating agents such as EDTA, carbohydrates such as dextran, hydroxyalkylcellulose, hydroxyalkylmethylcellulose, stearic acid and the like. The pH of the formulations ranges from about 3 to about 11, but is ordinarily about 7 to 10.

5 While it is possible for the active ingredients to be administered alone it may be preferable to present them as pharmaceutical formulations. The formulations, both for veterinary and for human use, comprise at least one active ingredient, as above defined, together with one or more acceptable carriers and optionally other therapeutic ingredients, particularly those additional therapeutic ingredients as discussed herein. The carrier(s) must
10 be "acceptable" in the sense of being compatible with the other ingredients of the formulation and physiologically innocuous to the recipient thereof.

 The formulations include those suitable for the foregoing administration routes. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. Techniques and formulations
15 generally are found in Remington's Pharmaceutical Sciences (Mack Publishing Co., Easton, PA). Such methods include the step of bringing into association the active ingredient with the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the
20 product.

 Formulations suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The
25 active ingredient may also be administered as a bolus, electuary or paste.

 A tablet is made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, surface active or dispersing agent.
30 Molded tablets may be made by molding in a suitable machine a mixture of the powdered active ingredient moistened with an inert liquid diluent. The tablets may optionally be coated or scored and optionally are formulated so as to provide slow or controlled release of the

active ingredient therefrom.

For infections of the eye or other external tissues e.g. mouth and skin, the formulations are preferably applied as a topical ointment or cream containing the active ingredient(s) in an amount of, for example, 0.075 to 20% w/w (including active ingredient(s) in a range between 0.1% and 20% in increments of 0.1% w/w such as 0.6% w/w, 0.7% w/w, etc.), preferably 0.2 to 15% w/w and most preferably 0.5 to 10% w/w. When formulated in an ointment, the active ingredients may be employed with either a paraffinic or a water-miscible ointment base. Alternatively, the active ingredients may be formulated in a cream with an oil-in-water cream base.

If desired, the aqueous phase of the cream base may include, for example, at least 30% w/w of a polyhydric alcohol, i.e. an alcohol having two or more hydroxyl groups such as propylene glycol, butane 1,3-diol, mannitol, sorbitol, glycerol and polyethylene glycol (including PEG 400) and mixtures thereof. The topical formulations may desirably include a compound which enhances absorption or penetration of the active ingredient through the skin or other affected areas. Examples of such dermal penetration enhancers include dimethyl sulphoxide and related analogs.

The oily phase of the emulsions may be constituted from known ingredients in a known manner. While the phase may comprise merely an emulsifier (otherwise known as an emulgent), it desirably comprises a mixture of at least one emulsifier with a fat or an oil or with both a fat and an oil. Preferably, a hydrophilic emulsifier is included together with a lipophilic emulsifier which acts as a stabilizer. It is also preferred to include both an oil and a fat. Together, the emulsifier(s) with or without stabilizer(s) make up the so-called emulsifying wax, and the wax together with the oil and fat make up the so-called emulsifying ointment base which forms the oily dispersed phase of the cream formulations.

Emulgents and emulsion stabilizers suitable for use in the formulation include Tween[®] 60, Span[®] 80, cetostearyl alcohol, benzyl alcohol, myristyl alcohol, glyceryl monostearate and sodium lauryl sulfate.

The choice of suitable oils or fats for the formulation is based on achieving the desired cosmetic properties. The cream should preferably be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or other containers. Straight or branched chain, mono- or dibasic alkyl esters such as di-isoadipate,

isocetyl stearate, propylene glycol diester of coconut fatty acids, isopropyl myristate, decyl oleate, isopropyl palmitate, butyl stearate, 2-ethylhexyl palmitate or a blend of branched chain esters known as Crodamol CAP may be used, the last three being preferred esters. These may be used alone or in combination depending on the properties required.

- 5 Alternatively, high melting point lipids such as white soft paraffin and/or liquid paraffin or other mineral oils are used.

Pharmaceutical formulations herein comprise a combination together with one or more pharmaceutically acceptable carriers or excipients and optionally other therapeutic agents. Pharmaceutical formulations containing the active ingredient may be in any form
10 suitable for the intended method of administration. When used for oral use for example, tablets, troches, lozenges, aqueous or oil suspensions, dispersible powders or granules, emulsions, hard or soft capsules, solutions, syrups or elixirs may be prepared. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain
15 one or more agents including sweetening agents, flavoring agents, coloring agents and preserving agents, in order to provide a palatable preparation. Tablets containing the active ingredient in admixture with non-toxic pharmaceutically acceptable excipient which are suitable for manufacture of tablets are acceptable. These excipients may be, for example, inert diluents, such as calcium or sodium carbonate, lactose, calcium or sodium phosphate;
20 granulating and disintegrating agents, such as maize starch, or alginic acid; binding agents, such as starch, gelatin or acacia; and lubricating agents, such as magnesium stearate, stearic acid or talc. Tablets may be uncoated or may be coated by known techniques including microencapsulation to delay disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay
25 material such as glyceryl monostearate or glyceryl distearate alone or with a wax may be employed.

Formulations for oral use may be also presented as hard gelatin capsules where the active ingredient is mixed with an inert solid diluent, for example calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil
30 medium, such as peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients include a suspending

agent, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropyl methylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia, and dispersing or wetting agents such as a naturally-occurring phosphatide (e.g., lecithin), a condensation product of an alkylene oxide with a fatty acid (e.g., polyoxyethylene stearate),
5 a condensation product of ethylene oxide with a long chain aliphatic alcohol (e.g., heptadecaethyleneoxycetanol), a condensation product of ethylene oxide with a partial ester derived from a fatty acid and a hexitol anhydride (e.g., polyoxyethylene sorbitan monooleate). The aqueous suspension may also contain one or more preservatives such as ethyl or n-propyl p-hydroxy-benzoate, one or more coloring agents, one or more flavoring
10 agents and one or more sweetening agents, such as sucrose or saccharin.

Oil suspensions may be formulated by suspending the active ingredient in a vegetable oil, such as arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oral suspensions may contain a thickening agent, such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set forth above, and
15 flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an antioxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, a suspending agent, and one or more preservatives. Suitable dispersing or
20 wetting agents and suspending agents are exemplified by those disclosed above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

The pharmaceutical compositions may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil, a mineral oil, such as liquid paraffin, or a mixture of these. Suitable emulsifying agents include naturally-occurring
25 gums, such as gum acacia and gum tragacanth, naturally-occurring phosphatides, such as soybean lecithin, esters or partial esters derived from fatty acids and hexitol anhydrides, such as sorbitan monooleate, and condensation products of these partial esters with ethylene oxide, such as polyoxyethylene sorbitan monooleate. The emulsion may also contain sweetening and flavoring agents. Syrups and elixirs may be formulated with
30 sweetening agents, such as glycerol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative, a flavoring or a coloring agent.

The pharmaceutical compositions may be in the form of a sterile injectable or intravenous preparations, such as a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above.

5 The sterile injectable or intravenous preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, such as a solution in 1,3-butane-diol or prepared as a lyophilized powder. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile fixed oils may conventionally be employed as a solvent or
10 suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid may likewise be used in the preparation of injectables.

The amount of active ingredient that may be combined with the carrier material to produce a single dosage form will vary depending upon the host treated and the particular
15 mode of administration. For example, a time-release formulation intended for oral administration to humans may contain approximately 1 to 1000 mg of active material compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 95% of the total compositions (weight:weight). The pharmaceutical composition can be prepared to provide easily measurable amounts for administration. For
20 example, an aqueous solution intended for intravenous infusion may contain from about 3 to 500 μ g of the active ingredient per milliliter of solution in order that infusion of a suitable volume at a rate of about 30 mL/hr can occur.

Formulations suitable for topical administration to the eye also include eye drops wherein the active ingredient is dissolved or suspended in a suitable carrier, especially an
25 aqueous solvent for the active ingredient. The active ingredient is preferably present in such formulations in a concentration of 0.5 to 20%, advantageously 0.5 to 10%, and particularly about 1.5% w/w.

Formulations suitable for topical administration in the mouth include lozenges comprising the active ingredient in a flavored basis, usually sucrose and acacia or
30 tragacanth; pastilles comprising the active ingredient in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

Formulations for rectal administration may be presented as a suppository with a suitable base comprising for example cocoa butter or a salicylate.

Formulations suitable for intrapulmonary or nasal administration have a particle size for example in the range of 0.1 to 500 microns, such as 0.5, 1, 30, 35 etc., which is administered by rapid inhalation through the nasal passage or by inhalation through the mouth so as to reach the alveolar sacs. Suitable formulations include aqueous or oily solutions of the active ingredient. Formulations suitable for aerosol or dry powder administration may be prepared according to conventional methods and may be delivered with other therapeutic agents such as compounds heretofore used in the treatment or prophylaxis of *Pneumovirinae* infections as described below.

Another embodiment provides a novel, efficacious, safe, nonirritating and physiologically compatible inhalable composition comprising a compound of Formula (I) or Formula (II), or a pharmaceutically acceptable salt thereof, suitable for treating *Pneumovirinae* infections and potentially associated bronchiolitis. Preferred pharmaceutically acceptable salts are inorganic acid salts including hydrochloride, hydrobromide, sulfate or phosphate salts as they may cause less pulmonary irritation. Preferably, the inhalable formulation is delivered to the endobronchial space in an aerosol comprising particles with a mass median aerodynamic diameter (MMAD) between about 1 and about 5 μm . Preferably, the compound of Formula (I) or Formula (II) is formulated for aerosol delivery using a nebulizer, pressurized metered dose inhaler (pMDI), or dry powder inhaler (DPI).

Non-limiting examples of nebulizers include atomizing, jet, ultrasonic, pressurized, vibrating porous plate, or equivalent nebulizers including those nebulizers utilizing adaptive aerosol delivery technology (Denyer, *J. Aerosol medicine Pulmonary Drug Delivery* 2010, 23 Supp 1, S1-S10). A jet nebulizer utilizes air pressure to break a liquid solution into aerosol droplets. An ultrasonic nebulizer works by a piezoelectric crystal that shears a liquid into small aerosol droplets. A pressurized nebulization system forces solution under pressure through small pores to generate aerosol droplets. A vibrating porous plate device utilizes rapid vibration to shear a stream of liquid into appropriate droplet sizes.

In a preferred embodiment, the formulation for nebulization is delivered to the endobronchial space in an aerosol comprising particles with a MMAD predominantly between about 1 μm and about 5 μm using a nebulizer able to aerosolize the formulation of

the compound of Formula (I) or Formula (II) into particles of the required MMAD. To be optimally therapeutically effective and to avoid upper respiratory and systemic side effects, the majority of aerosolized particles should not have a MMAD greater than about 5 μm . If an aerosol contains a large number of particles with a MMAD larger than 5 μm , the particles are deposited in the upper airways decreasing the amount of drug delivered to the site of inflammation and bronchoconstriction in the lower respiratory tract. If the MMAD of the aerosol is smaller than about 1 μm , then the particles have a tendency to remain suspended in the inhaled air and are subsequently exhaled during expiration.

When formulated and delivered according to the method herein, the aerosol formulation for nebulization delivers a therapeutically efficacious dose of the compound of Formula (I) or Formula (II) to the site of *Pneumovirinae* infection sufficient to treat the *Pneumovirinae* infection. The amount of drug administered must be adjusted to reflect the efficiency of the delivery of a therapeutically efficacious dose of the compound of Formula (I) or Formula (II). In a preferred embodiment, a combination of the aqueous aerosol formulation with the atomizing, jet, pressurized, vibrating porous plate, or ultrasonic nebulizer permits, depending on the nebulizer, about, at least, 20, to about 90%, typically about 70% delivery of the administered dose of the compound of Formula (I) or Formula (II) into the airways. In a preferred embodiment, at least about 30 to about 50% of the active compound is delivered. More preferably, about 70 to about 90% of the active compound is delivered.

In another embodiment, a compound of Formula (I) or Formula (II) or a pharmaceutically acceptable salt thereof, is delivered as a dry inhalable powder. The compounds are administered endobronchially as a dry powder formulation to efficacious deliver fine particles of compound into the endobronchial space using dry powder or metered dose inhalers. For delivery by DPI, the compound of Formula (I) or Formula (II) is processed into particles with, predominantly, MMAD between about 1 μm and about 5 μm by milling spray drying, critical fluid processing, or precipitation from solution. Media milling, jet milling and spray-drying devices and procedures capable of producing the particle sizes with a MMAD between about 1 μm and about 5 μm are well known in the art. In one embodiment, excipients are added to the compound of Formula (I) or Formula (II) before processing into particles of the required sizes. In another embodiment, excipients are blended with the particles of the required size to aid in dispersion of the drug particles, for example by using lactose as an excipient.

Particle size determinations are made using devices well known in the art. For example a multi-stage Anderson cascade impactor or other suitable method such as those specifically cited within the US Pharmacopoeia Chapter 601 as characterizing devices for aerosols within metered-dose and dry powder inhalers.

5 In another preferred embodiment, a compound of Formula (I) or Formula (II) is delivered as a dry powder using a device such as a dry powder inhaler or other dry powder dispersion devices. Non-limiting examples of dry powder inhalers and devices include those disclosed in US5,458,135; US5,740,794; US5775320; US5,785,049; US3,906,950; US4,013,075; US4,069,819; US4,995,385; US5,522,385; US4,668,218; US4,667,668;
10 US4,805,811 and US5,388,572. There are two major designs of dry powder inhalers. One design is a metering device in which a reservoir for the drug is place within the device and the patient adds a dose of the drug into the inhalation chamber. The second design is a factory-metered device in which each individual dose has been manufactured in a separate container. Both systems depend on the formulation of the drug into small particles of MMAD
15 from 1 μm and about 5 μm and often involve co-formulation with larger excipient particles such as, but not limited to, lactose. Drug powder is placed in the inhalation chamber (either by device metering or by breakage of a factory-metered dosage) and the inspiratory flow of the patient accelerates the powder out of the device and into the oral cavity. Non-laminar flow characteristics of the powder path cause the excipient-drug aggregates to decompose,
20 and the mass of the large excipient particles causes their impaction at the back of the throat, while the smaller drug particles are deposited deep in the lungs. In preferred embodiments, a compound of Formula (I) or Formula (II), or a pharmaceutically acceptable salt thereof, is delivered as a dry powder using either type of dry powder inhaler as described herein, wherein the MMAD of the dry powder, exclusive of any excipients, is predominantly in the
25 range of 1 μm to about 5 μm .

 In another embodiment, a compound of Formula (I) or Formula (II) is delivered as a dry powder using a metered dose inhaler. Non-limiting examples of metered dose inhalers and devices include those disclosed in US5,261,538; US5,544,647; US5,622,163; US4,955,371; US3,565,070; US3,361306 and US6,116,234. In preferred embodiments, a
30 compound of Formula (I) or Formula (II), or a pharmaceutically acceptable salt thereof, is delivered as a dry powder using a metered dose inhaler wherein the MMAD of the dry powder, exclusive of any excipients, is predominantly in the range of about 1-5 μm .

Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

5 Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents.

10 The formulations are presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injection, immediately prior to use. Extemporaneous injection solutions and suspensions are prepared from sterile powders, granules and tablets of the kind previously described. Preferred unit dosage formulations are those containing a daily dose or unit daily sub-dose, as herein
15 above recited, or an appropriate fraction thereof, of the active ingredient.

It should be understood that in addition to the ingredients particularly mentioned above the formulations may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavoring agents.

20 Further provided are veterinary compositions comprising at least one active ingredient as above defined together with a veterinary carrier therefor.

Veterinary carriers are materials useful for the purpose of administering the composition and may be solid, liquid or gaseous materials which are otherwise inert or acceptable in the veterinary art and are compatible with the active ingredient. These
25 veterinary compositions may be administered orally, parenterally or by any other desired route.

Compounds herein are used to provide controlled release pharmaceutical formulations containing as active ingredient one or more of the compounds ("controlled release formulations") in which the release of the active ingredient are controlled and
30 regulated to allow less frequency dosing or to improve the pharmacokinetic or toxicity profile of a given active ingredient.

Effective dose of active ingredient depends at least on the nature of the condition being treated, toxicity, whether the compound is being used prophylactically (lower doses) or against an active viral infection, the method of delivery, and the pharmaceutical formulation, and will be determined by the clinician using conventional dose escalation studies. It can be expected to be from about 0.0001 to about 100 mg/kg body weight per day; typically, from about 0.01 to about 10 mg/kg body weight per day; more typically, from about .01 to about 5 mg/kg body weight per day; most typically, from about .05 to about 0.5 mg/kg body weight per day. For example, the daily candidate dose for an adult human of approximately 70 kg body weight will range from 1 mg to 1000 mg, preferably between 5 mg and 500 mg, and may take the form of single or multiple doses.

Routes of Administration

One or more of the compounds (herein referred to as the active ingredients) are administered by any route appropriate to the condition to be treated. Suitable routes include oral, rectal, nasal, pulmonary, topical (including buccal and sublingual), vaginal and parenteral (including subcutaneous, intramuscular, intravenous, intradermal, intrathecal and epidural), and the like. It will be appreciated that the preferred route may vary with for example the condition of the recipient. An advantage of the compounds herein is that they are orally bioavailable and can be dosed orally.

Combination Therapy

Compositions are also used in combination with other active ingredients. For the treatment of *Pneumovirinae* virus infections, preferably, the other active therapeutic agent is active against *Pneumovirinae* virus infections, particularly respiratory syncytial virus infections. Non-limiting examples of these other active therapeutic agents are ribavirin, palivizumab, motavizumab, RSV-IGIV (RespiGam®), MEDI-557, A-60444 (also known as RSV604), MDT-637, BMS-433771, ALN-RSV0, ALX-0171 and mixtures thereof.

Many of the infections of the *Pneumovirinae* viruses are respiratory infections. Therefore, additional active therapeutics used to treat respiratory symptoms and sequelae of infection may be used in combination with the compounds of Formula (I) or Formula (II). The additional agents are preferably administered orally or by direct inhalation. For example, other preferred additional therapeutic agents in combination with the compounds of

Formula (I) or Formula (II) for the treatment of viral respiratory infections include, but are not limited to, bronchodilators and corticosteroids.

Glucocorticoids, which were first introduced as an asthma therapy in 1950 (Carrier, Journal of Allergy, 21, 282-287, 1950), remain the most potent and consistently effective therapy for this disease, although their mechanism of action is not yet fully understood (Morris, J. Allergy Clin. Immunol., 75 (1 Pt) 1-13, 1985). Unfortunately, oral glucocorticoid therapies are associated with profound undesirable side effects such as truncal obesity, hypertension, glaucoma, glucose intolerance, acceleration of cataract formation, bone mineral loss, and psychological effects, all of which limit their use as long-term therapeutic agents (Goodman and Gilman, 10th edition, 2001). A solution to systemic side effects is to deliver steroid drugs directly to the site of inflammation. Inhaled corticosteroids (ICS) have been developed to mitigate the severe adverse effects of oral steroids. Non-limiting examples of corticosteroids that may be used in combinations with the compounds of Formula (I) or Formula (II) are dexamethasone, dexamethasone sodium phosphate, fluorometholone, fluorometholone acetate, loteprednol, loteprednol etabonate, hydrocortisone, prednisolone, fludrocortisone, triamcinolone, triamcinolone acetonide, betamethasone, beclomethasone dipropionate, methylprednisolone, fluocinolone, fluocinolone acetonide, flunisolide, flucortin-21-butyrate, flumethasone, flumetasone pivalate, budesonide, halobetasol propionate, mometasone furoate, fluticasone propionate, ciclesonide; or a pharmaceutically acceptable salts thereof.

Other anti-inflammatory agents working through anti-inflammatory cascade mechanisms are also useful as additional therapeutic agents in combination with the compounds of Formula (I) or Formula (II) for the treatment of viral respiratory infections. Applying "anti-inflammatory signal transduction modulators" (referred to in this text as AISTM), like phosphodiesterase inhibitors (e.g. PDE-4, PDE-5, or PDE-7 specific), transcription factor inhibitors (e.g. blocking NFκB through IKK inhibition), or kinase inhibitors (e.g. blocking P38 MAP, JNK, PI3K, EGFR or Syk) is a logical approach to switching off inflammation as these small molecules target a limited number of common intracellular pathways - those signal transduction pathways that are critical points for the anti-inflammatory therapeutic intervention (see review by P.J. Barnes, 2006). These non-limiting additional therapeutic agents include: 5-(2,4-Difluoro-phenoxy)-1-isobutyl-1H-indazole-6-carboxylic acid (2-dimethylamino-ethyl)-amide (P38 Map kinase inhibitor ARRY-797); 3-Cyclopropylmethoxy-N-(3,5-dichloro-pyridin-4-yl)-4-difluoromethoxy-benzamide (PDE-4 inhibitor Roflumilast); 4-

[2-(3-cyclopentyloxy-4-methoxyphenyl)-2-phenyl-ethyl]-pyridine (PDE-4 inhibitor CDP-840); N-(3,5-dichloro-4-pyridinyl)-4-(difluoromethoxy)-8-[(methylsulfonyl)amino]-1-dibenzofurancarboxamide (PDE-4 inhibitor Oglemilast); N-(3,5-Dichloro-pyridin-4-yl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxo-acetamide (PDE-4 inhibitor AWD 12-281); 8-Methoxy-2-trifluoromethyl-quinoline-5-carboxylic acid (3,5-dichloro-1-oxy-pyridin-4-yl)-amide (PDE-4 inhibitor Sch 351591); 4-[5-(4-Fluorophenyl)-2-(4-methanesulfinyl-phenyl)-1H-imidazol-4-yl]-pyridine (P38 inhibitor SB-203850); 4-[4-(4-Fluoro-phenyl)-1-(3-phenyl-propyl)-5-pyridin-4-yl-1H-imidazol-2-yl]-but-3-yn-1-ol (P38 inhibitor RWJ-67657); 4-Cyano-4-(3-cyclopentyloxy-4-methoxy-phenyl)-cyclohexanecarboxylic acid 2-diethylamino-ethyl ester (2-diethyl-ethyl ester prodrug of Cilomilast, PDE-4 inhibitor); (3-Chloro-4-fluorophenyl)-[7-methoxy-6-(3-morpholin-4-yl-propoxy)-quinazolin-4-yl]-amine (Gefitinib, EGFR inhibitor); and 4-(4-Methyl-piperazin-1-ylmethyl)-N-[4-methyl-3-(4-pyridin-3-yl-pyrimidin-2-ylamino)-phenyl]-benzamide (Imatinib, EGFR inhibitor).

Combinations comprising inhaled β 2-adrenoreceptor agonist bronchodilators such as formoterol, albuterol or salmeterol with the compounds of Formula (I) or Formula (II) are also suitable, but non-limiting, combinations useful for the treatment of respiratory viral infections.

Combinations of inhaled β 2-adrenoreceptor agonist bronchodilators such as formoterol or salmeterol with ICS's are also used to treat both the bronchoconstriction and the inflammation (Symbicort® and Advair®, respectively). The combinations comprising these ICS and β 2-adrenoreceptor agonist combinations along with the compounds of Formula (I) or Formula (II) are also suitable, but non-limiting, combinations useful for the treatment of respiratory viral infections.

For the treatment or prophylaxis of pulmonary broncho-constriction, anticholinergics are of potential use and, therefore, useful as an additional therapeutic agents in combination with the compounds of Formula (I) or Formula (II) for the treatment of viral respiratory infections. These anticholinergics include, but are not limited to, antagonists of the muscarinic receptor (particularly of the M3 subtype) which have shown therapeutic efficacy in man for the control of cholinergic tone in COPD (Witek, 1999); 1-{4-Hydroxy-1-[3,3,3-tris-(4-fluoro-phenyl)-propionyl]-pyrrolidine-2-carbonyl}-pyrrolidine-2-carboxylic acid (1-methyl-piperidin-4-ylmethyl)-amide; 3-[3-(2-Diethylamino-acetoxy)-2-phenyl-propionyloxy]-8-isopropyl-8-methyl-8-azonia-bicyclo[3.2.1]octane (Ipratropium-N,N-diethylglycinate); 1-Cyclohexyl-3,4-dihydro-1H-isoquinoline-2-carboxylic acid 1-aza-bicyclo[2.2.2]oct-3-yl ester

(Solifenacin); 2-Hydroxymethyl-4-methanesulfinyl-2-phenyl-butyric acid 1-azabicyclo[2.2.2]oct-3-yl ester (Revatropate); 2-{1-[2-(2,3-Dihydro-benzofuran-5-yl)-ethyl]-pyrrolidin-3-yl}-2,2-diphenyl-acetamide (Darifenacin); 4-Azepan-1-yl-2,2-diphenyl-butyramide (Buzepide); 7-[3-(2-Diethylamino-acetoxy)-2-phenyl-propionyloxy]-9-ethyl-9-methyl-3-oxa-9-azonia-tricyclo[3.3.1.0^{2,4}]nonane (Oxitropium-N,N-diethylglycinate); 7-[2-(2-Diethylamino-acetoxy)-2,2-di-thiophen-2-yl-acetoxy]-9,9-dimethyl-3-oxa-9-azonia-tricyclo[3.3.1.0^{2,4}]nonane (Tiotropium-N,N-diethylglycinate); Dimethylamino-acetic acid 2-(3-diisopropylamino-1-phenyl-propyl)-4-methyl-phenyl ester (Tolterodine-N,N-dimethylglycinate); 3-[4,4-Bis-(4-fluoro-phenyl)-2-oxo-imidazolidin-1-yl]-1-methyl-1-(2-oxo-2-pyridin-2-yl-ethyl)-pyrrolidinium; 1-[1-(3-Fluoro-benzyl)-piperidin-4-yl]-4,4-bis-(4-fluoro-phenyl)-imidazolidin-2-one; 1-Cyclooctyl-3-(3-methoxy-1-aza-bicyclo[2.2.2]oct-3-yl)-1-phenyl-prop-2-yn-1-ol; 3-[2-(2-Diethylamino-acetoxy)-2,2-di-thiophen-2-yl-acetoxy]-1-(3-phenoxy-propyl)-1-azonia-bicyclo[2.2.2]octane (Acidinium-N,N-diethylglycinate); or (2-Diethylamino-acetoxy)-di-thiophen-2-yl-acetic acid 1-methyl-1-(2-phenoxy-ethyl)-piperidin-4-yl ester.

The compounds of Formula (I) or Formula (II) may also be combined with mucolytic agents to treat both the infection and symptoms of respiratory infections. A non-limiting example of a mucolytic agent is ambroxol. Similarly, the compounds of Formula (I) or Formula (II) may be combined with expectorants to treat both the infection and symptoms of respiratory infections. A non-limiting example of an expectorant is guaifenesin.

Nebulized hypertonic saline is used to improve immediate and long-term clearance of small airways in patients with lung diseases (Kuzik, *J. Pediatrics* 2007, 266). The compounds of Formula (I) or Formula (II) may also be combined with nebulized hypertonic saline particularly when the *Pneumovirinae* virus infection is complicated with bronchiolitis. The combination of the compounds of Formula (I) or Formula (II) with hypertonic saline may also comprise any of the additional agents discussed above. In one embodiment, nebulized about 3% hypertonic saline is used.

It is also possible to combine any compound with one or more additional active therapeutic agents in a unitary dosage form for simultaneous or sequential administration to a patient. The combination therapy may be administered as a simultaneous or sequential regimen. When administered sequentially, the combination may be administered in two or more administrations.

Co-administration of a compound herein with one or more other active therapeutic agents generally refers to simultaneous or sequential administration of a compound and one or more other active therapeutic agents, such that therapeutically effective amounts of the compound and one or more other active therapeutic agents are both present in the body of the patient.

Co-administration includes administration of unit dosages of the compounds before or after administration of unit dosages of one or more other active therapeutic agents, for example, administration of the compounds within seconds, minutes, or hours of the administration of one or more other active therapeutic agents. For example, a unit dose of a compound can be administered first, followed within seconds or minutes by administration of a unit dose of one or more other active therapeutic agents. Alternatively, a unit dose of one or more other therapeutic agents can be administered first, followed by administration of a unit dose of a compound within seconds or minutes. In some cases, it may be desirable to administer a unit dose of a compound first, followed, after a period of hours (e.g., 1-12 hours), by administration of a unit dose of one or more other active therapeutic agents. In other cases, it may be desirable to administer a unit dose of one or more other active therapeutic agents first, followed, after a period of hours (e.g., 1-12 hours), by administration of a unit dose of a compound herein.

The combination therapy may provide "synergy" and "synergistic", i.e. the effect achieved when the active ingredients used together is greater than the sum of the effects that results from using the compounds separately. A synergistic effect may be attained when the active ingredients are: (1) co-formulated and administered or delivered simultaneously in a combined formulation; (2) delivered by alternation or in parallel as separate formulations; or (3) by some other regimen. When delivered in alternation therapy, a synergistic effect may be attained when the compounds are administered or delivered sequentially, e.g. in separate tablets, pills or capsules, or by different injections in separate syringes. In general, during alternation therapy, an effective dosage of each active ingredient is administered sequentially, i.e. serially, whereas in combination therapy, effective dosages of two or more active ingredients are administered together. A synergistic anti-viral effect denotes an antiviral effect which is greater than the predicted purely additive effects of the individual compounds of the combination.

In still yet another embodiment, the present application provides a method of treating

Pneumovirinae virus infection in a human, the method comprising administering to the human a therapeutically effective amount of a compound of Formula (I), or a pharmaceutically acceptable salt, solvate, and/or ester thereof. Also provided are separate methods of treating *Pneumovirinae* virus infection in a human, each comprising

- 5 administering to the human a therapeutically effective a pharmaceutically effective amount of a compound of Formula (II) or one of the specific compounds of the examples herein, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and a pharmaceutically acceptable carrier or excipient.

- 10 In another embodiment, provided is a method of treating a *Pneumovirinae* infection in a human by administering to the human a therapeutically effective amount of a racemate, enantiomer, diastereomer, tautomer, polymorph, pseudopolymorph, amorphous form, hydrate or solvate of a compound of a compound of Formula (I), or a pharmaceutically acceptable salt or ester thereof.

- 15 Further provided are separate methods of treating a *Pneumovirinae* infection in a human in need thereof, each method comprising administering to the human a therapeutically effective amount of a racemate, enantiomer, diastereomer, tautomer, polymorph, pseudopolymorph, amorphous form, hydrate or solvate of a compound of Formula (II) or one of the specific compounds of the examples herein, or a pharmaceutically acceptable salt, solvate, and/or ester thereof.

- 20 In still yet another embodiment, the present application provides for a method of treating human respiratory syncytial virus infection in a human, the method comprising administering to the human a therapeutically effective amount of a compound of Formula (I), or a pharmaceutically acceptable salt, solvate, and/or ester thereof.

- 25 In still yet another embodiment, the present application provides for a method of treating human respiratory syncytial virus infection in a human, the method comprising administering to the human a therapeutically effective amount of a compound of Formula (I), or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and at least one additional active therapeutic agent.

- 30 Further provided are separate methods of treating a human respiratory syncytial virus infection in a human in need thereof, each method comprising administering to the human a therapeutically effective amount of a compound of Formula (II) or one of the specific

compounds of the examples herein, or a pharmaceutically acceptable salt, solvate, and/or ester thereof.

Also provided are separate methods of treating a human respiratory syncytial virus infection in a human in need thereof, each method comprising administering to the human a therapeutically effective amount of a compound of Formula (II) or one of the specific compounds of the examples herein, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and at least one additional active therapeutic agent.

Also provided are separate methods of treating a human respiratory syncytial virus infection in a human in need thereof, wherein the human is also experiencing bronchiolitis, each method comprising administering to the human a therapeutically effective amount of a compound of Formula (I), Formula (II), or one of the specific compounds of the examples herein, or a pharmaceutically acceptable salt, solvate, and/or ester thereof.

Also provided are separate methods of treating a human respiratory syncytial virus infection in a human in need thereof, wherein the human is also experiencing pneumonia, each method comprising administering to the human a therapeutically effective amount of a compound of Formula (I), Formula (II), or one of the specific compounds of the examples herein, or a pharmaceutically acceptable salt, solvate, and/or ester thereof.

Also provided are separate methods of improving respiratory symptoms in a human experiencing a human respiratory syncytial virus infection, each method comprising administering to the human a therapeutically effective amount of a compound of Formula (I), Formula (II), or one of the specific compounds of the examples herein, or a pharmaceutically acceptable salt, solvate, and/or ester thereof.

Respiratory symptoms in a human experiencing a respiratory syncytial virus infection may include congested or runny nose, coughing, wheezing, sneezing, rapid breathing or difficulty breathing, apnea, bronchiolitis, and pneumonia.

Also provided is an embodiment comprising the use of a compound of Formula (I), or a pharmaceutically acceptable salt, solvate, and/or ester thereof, for the manufacture of a medicament for the treatment of a *Pneumovirinae* virus infection or a respiratory syncytial virus infection.

Also provided is an embodiment comprising the use of a compound of Formula (II) or one of the specific compounds of the examples herein, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, for the manufacture of a medicament for the treatment of a *Pneumovirinae* virus infection or a respiratory syncytial virus infection.

5 Also provided is a pharmaceutical formulation comprising a pharmaceutically effective amount of a compound of Formula (I), or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and a pharmaceutically acceptable carrier or excipient. Further provided is a pharmaceutical formulation comprising a pharmaceutically effective amount of a compound of Formula (II) or one of the specific compounds of the examples herein, or a
10 pharmaceutically acceptable salt, solvate, and/or ester thereof, and a pharmaceutically acceptable carrier or excipient.

 Also provided is a pharmaceutical formulation comprising a pharmaceutically effective amount of a compound of Formula (I), or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and a pharmaceutically acceptable carrier or excipient and a
15 pharmaceutically effective amount of at least one additional active therapeutic agent. Further provided is a pharmaceutical formulation comprising a pharmaceutically effective amount of a compound of Formula (II) or one of the specific compounds of the examples herein, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and a
20 pharmaceutically acceptable carrier or excipient and a pharmaceutically effective amount of at least one additional active therapeutic agent.

 Also provided are separate embodiments comprising a compound of Formula (I), Formula (II), or one of the specific compounds of the examples herein, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, for use in the treatment of a *Pneumovirinae* virus infection or a respiratory syncytial virus infection in a human.

25 Also provided are separate embodiments comprising a compound of Formula (I), Formula (II) or one of the specific compounds of the examples herein, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, for use as a medicament.

 Also provided are separate embodiments comprising a method for manufacturing a medicament intended for treatment of a *Pneumovirinae* virus infection or a respiratory
30 syncytial virus infection in a human, characterised in that a compound of Formula (I),

Formula (II), or one of the specific compounds of the examples herein, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, is used.

Also provided is a compound of Formula (I), or a pharmaceutically acceptable salt, solvate, and/or ester thereof, for the treatment of a *Pneumovirinae* virus infection or a respiratory syncytial virus infection in a human.

Also provided are separate embodiments comprising that a compound of Formula (II) or one of the specific compounds of the examples herein, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, for the treatment of a *Pneumovirinae* virus infection or a respiratory syncytial virus infection in a human.

Further provided is a compound as described in this specification. Also provided is a pharmaceutical composition as described in this specification. Also provided is a method of using a compound of Formula (I), as described in this specification. Further provided is a method of making a compound of Formula (I), as described in this specification.

Metabolites of the Compounds

Also falling within the scope herein are the *in vivo* metabolic products of the compounds described herein, to the extent such products are novel and unobvious over the prior art. Such products may result for example from the oxidation, reduction, hydrolysis, amidation, esterification and the like of the administered compound, primarily due to enzymatic processes. Accordingly, included are novel and unobvious compounds produced by a process comprising contacting a compound with a mammal for a period of time sufficient to yield a metabolic product thereof. Such products typically are identified by preparing a radiolabelled (e.g. ^{14}C or ^3H) compound, administering it parenterally in a detectable dose (e.g. greater than about 0.5 mg/kg) to an animal such as rat, mouse, guinea pig, monkey, or to man, allowing sufficient time for metabolism to occur (typically about 30 seconds to 30 hours) and isolating its conversion products from the urine, blood or other biological samples. These products are easily isolated since they are labeled (others are isolated by the use of antibodies capable of binding epitopes surviving in the metabolite). The metabolite structures are determined in conventional fashion, e.g. by MS or NMR analysis. In general, analysis of metabolites is done in the same way as conventional drug metabolism studies well-known to those skilled in the art. The conversion products, so long as they are not otherwise found *in vivo*, are useful in diagnostic assays for therapeutic

dosing of the compounds even if they possess no HSV antiviral activity of their own.

Recipes and methods for determining stability of compounds in surrogate gastrointestinal secretions are known. Compounds are defined herein as stable in the gastrointestinal tract where less than about 50 mole percent of the protected groups are deprotected in surrogate intestinal or gastric juice upon incubation for 1 hour at 37°C. Simply because the compounds are stable to the gastrointestinal tract does not mean that they cannot be hydrolyzed *in vivo*. The prodrugs typically will be stable in the digestive system but may be substantially hydrolyzed to the parental drug in the digestive lumen, liver, lung or other metabolic organ, or within cells in general. As used herein, a prodrug is understood to be a compound that is chemically designed to efficiently liberate the parent drug after overcoming biological barriers to oral delivery.

Abbreviations

Certain abbreviations and acronyms are used in describing the experimental details. Although most of these would be understood by one skilled in the art, Table 1 contains a list of many of these abbreviations and acronyms.


Table 1. List of abbreviations and acronyms.

Abbreviation	Meaning
Ac	acetate
ACN	acetonitrile
AIBN	azobisisobutyronitrile
Bn	benzyl
Bu	butyl
Bz	benzoyl
BzCl	benzoyl chloride
CDI	1,1'-carbonyldiimidazole

DAST	diethylaminosulfur trifluoride
DCE	1,2-dichloroethane
DCM	dichloromethane
DMAP	4-dimethylamiopyridine
DMDO	dimethyldioxirane
DMSO	dimethylsulfoxide
DMF	dimethylformamide
DMTrCl	4,4'-dimethoxytritylchloride
DMTr	4,4'-dimethoxytrityl
EDCI	<i>N</i> -(3-dimethylaminopropyl)- <i>N'</i> -ethylcarbodiimide hydrochloride
Et	ethyl
Imid	imidazole
KOtBu	potassium tert-butoxide
LC	liquid chromatography
MCPBA	meta-chloroperbenzoic acid
Me	methyl
m/z	mass to charge ratio
MS or ms	mass spectrum
NIS	<i>N</i> -iodosuccinimide
NMP	<i>N</i> -methyl-2-pyrrolidone
Ph	phenyl


Ph ₃ P	triphenylphosphine
PMB	<i>para</i> -methoxybenzyl
PMBCl	<i>para</i> -methoxybenzyl chloride
PhOC(S)Cl	phenylchlorothionoformate
(PhO) ₃ PMeI	methyltriphenoxyphosphonium iodide
Pyr	pyridine
RT	room temperature
TBAF	tetrabutylammonium fluoride
TBS	<i>tert</i> -butyldimethylsilyl
TBSCl	<i>tert</i> -Butyldimethylsilyl chloride
TMSN ₃	trimethylsilyl azide
TEA	triethylamine
TES	triethylsilane
TFA	trifluoroacetic acid
THF	tetrahydrofuran
TMS	trimethylsilyl
TMSCl	trimethylsilyl chloride
Ts	4-toluenesulfonyl
TsOH	tosylic acid
δ	parts per million referenced to residual non-deuterated solvent peak

General Schemes




Scheme 1.

Scheme 1 shows a general synthesis of intermediates beginning with an iodination reaction (e.g. NIS) to generate nucleobase **S1b**.




Scheme 2.

Scheme 2 shows a general synthesis of intermediates beginning with a fluorination reaction (e.g. HBF_4 , NaNO_2) in a manner similar to that described in WO2012037038A1 to afford intermediate **S2b**. Intermediate **S2b** can then be iodinated (e.g. NIS) to generate nucleobase **S2c**.




Scheme 3.

Scheme 3 shows a general synthesis of compounds beginning with a lithium-halogen exchange (e.g. *n*-BuLi, [-CH₂SiMe₂Cl]₂) reaction with an appropriate nucleobase **S3b**, followed by addition to the lactone **S3a**. Reduction of the pendant 1' hydroxyl group under Lewis acidic conditions (e.g. BF₃·Et₂O, Et₃SiH) generates intermediate **S3c**. Intermediate **S3c** can first be protected (e.g. BzCl, Pyr; NH₄OH) at the nitrogen functionality, and the benzyl groups can then be removed under reducing conditions (e.g. HCO₂H, Pd/C, BCl₃, BBr₃) to afford intermediate **S3d**. A sequence involving first selective 5' hydroxyl protection (e.g. DMTrCl), then 2' and 3' hydroxyl protection (e.g. TBSCl), followed by selective removal of the 5' hydroxyl protecting group under acidic conditions (e.g. TsOH) furnishes intermediate **S3e**. The 5' hydroxyl group can then be converted to the corresponding iodide (e.g. (PhO)₃PMel), which is then exposed to basic conditions (i.e. KO^tBu) to effect an elimination reaction generating intermediate **S3f**. Oxidation of the olefin **S3f** (e.g. DMDO) followed by treatment with an appropriate nucleophile (e.g. TMSCN) under Lewis acidic conditions (e.g. InBr₃) and removal of the hydroxyl protecting groups (e.g. CsF) affords intermediate **S3g**. Removal of the nitrogen protecting group (e.g. NH₂Me) yields the final compounds of the type **S3h**.




Scheme 4.

Scheme 4 shows a general synthesis of compounds beginning with oxidation of the olefin **S3f** (e.g. DMDO) followed by treatment with an appropriate nucleophile (e.g. TMSN₃) under Lewis acidic conditions (e.g. InBr₃) in a manner similar to that described in *J. Med. Chem.* **2007**, *50*, 5463-5470. Removal of the hydroxyl protecting groups (e.g. CsF) then affords intermediate **S4a**. Removal of the nitrogen protecting group (e.g. NH₂Me) yields the final compounds of the type **S4b**.




Scheme 5.

Scheme 5 shows a general synthesis of compounds beginning with oxidation of the olefin **S3f** (e.g. DMDO) in the presence of the appropriate alcohol HOR^a followed by removal of the hydroxyl protecting groups (e.g. CsF) to afford intermediate **S5a**. Removal of the nitrogen protecting group (e.g. NH₂Me) yields the final compounds of the type **S5b**.




Scheme 6.

Scheme 6 shows a general synthesis of compounds beginning with oxidation of the olefin **S3f** (e.g. DMDO) followed by treatment with an appropriate nucleophile (e.g. (HC≡C)₃Al) in a manner similar to that described in *Nucleosides, Nucleotides, and Nucleic Acids* **2005**, 24, 343-347. Removal of the hydroxyl protecting groups (e.g. CsF) then affords intermediate **S6a**. Removal of the nitrogen protecting group (e.g. NH₂Me) yields the final compounds of the type **S6b**. Elaboration of the final compound through hydrogenation conditions (e.g. H₂, Pd/C or Lindlar's conditions) can selectively afford the final compounds of the type **S6c** and **S6d** respectively. Elaboration of the final compound **S6d** through cyclopropanation conditions (e.g. CH₂N₂) can yield the final compounds of the type **S6e**.



Scheme 7.


Scheme 7 shows a general synthesis of compounds beginning with a synthetic sequence to protect the nitrogen (e.g. TMSCl, Pyr; BzCl, Pyr; NH₄OH) of intermediate **S7a**, synthesized in a similar manner as described in WO2012037038A1. Selective protection of the 5' hydroxyl group (e.g. DMTrCl) then generates intermediate **S7b**. Protection of the 2' hydroxyl group (e.g. TBSCl) followed by removal of the 5' hydroxyl group under acidic conditions (e.g. TsOH) furnishes intermediate **S7e**. Conversion of the 5' hydroxyl group to the aldehyde under oxidative conditions (e.g. EDCI·HCl, Pyr, TFA, DMSO) followed by condensation of the corresponding enolate with formaldehyde and reduction (e.g. NaBH₄) yields intermediate **S7f**. Sequential selective protection of the hydroxyl moieties with orthogonal protecting groups (e.g. DMTrCl and TBSCl) followed by removal of the more labile protecting group under acidic conditions (e.g. TsOH) then affords intermediate **S7g**. Conversion of the hydroxyl group to the aldehyde under oxidative conditions (e.g. EDCI·HCl, Pyr, TFA, DMSO) generates intermediate **S7h**. Elaboration of the aldehyde **S7h** to the halo-olefin intermediate **S7i** can be effected under Wittig olefination conditions (e.g. [Ph₃PCH₂Br]⁺Br⁻, KOtBu). An elimination reaction under basic conditions (e.g. KOtBu) generates the alkyne, and removal of the hydroxyl protecting groups (e.g. TBAF) and nitrogen protecting group (e.g. NH₄OH) yields the final compounds of the type **S7j**.



Scheme 8.


Scheme 8 shows a general synthesis of compounds beginning with oxime formation (e.g. $\text{NH}_2\text{OH} \cdot \text{HCl}$), followed by conversion of the oxime to a nitrile group (e.g. CDI).

- 5 Removal of the nitrogen protecting group (e.g. NH_4OH), and hydroxyl protecting groups (e.g. $\text{HF} \cdot \text{Pyr}$) then yields the final compounds of the type **S8a**.




Scheme 9.

- 10 Scheme 9 shows a general synthesis of compounds beginning with elaboration of the aldehyde **S7h** to the olefin with Wittig olefination conditions (e.g. $[\text{Ph}_3\text{PCH}_3]^+\text{Br}^-$, KOtBu). Removal of the hydroxyl protecting groups (e.g. TBAF), and nitrogen protecting group (e.g. NH_4OH) yields the final compounds of the type **S9a**. Reducing conditions (e.g. H_2 , Pd/C) then can generate the final compounds of the type **S9b**, and cyclopropanation conditions (e.g. CH_2N_2) can generate the final compounds of the type **S9c**.




Scheme 10.

Scheme 10 shows a general synthesis of compounds beginning with an Appel reaction (e.g. Ph_3P , CCl_4) to convert the hydroxyl group into a chloride. Removal of the hydroxyl protecting groups (e.g. TBAF), and nitrogen protecting group (e.g. NH_4OH) yields the final compounds of the type **S10a**.



Scheme 11.


Scheme 11 shows a general synthesis of compounds beginning with protection of the free hydroxyl group of intermediate **S7g** with a labile protecting group (e.g. PMBCl, K_2CO_3). Selective removal of the 2' and 5' silyloxy protecting groups (e.g. TBAF), followed by re-protection with robust protecting groups (e.g. BnBr , NaH), and removal of the labile hydroxyl protecting group under acidic conditions (e.g. AcOH) affords intermediate **S11a**. Conversion of the hydroxyl group to the fluorine (e.g. DAST) followed by removal of the hydroxyl protecting groups (e.g. BBr_3), and nitrogen protecting group (e.g. NH_4OH) yields the final compounds of the type **S11b**.



Scheme 12.


Scheme 12 shows a general synthesis of compounds beginning with conversion of the 5' hydroxyl group to the corresponding iodide (e.g. $(\text{PhO})_3\text{PMel}$), which is then treated with basic conditions (i.e. KOtBu) to effect an elimination reaction generating intermediate **S12a**. Oxidation of the olefin **S12a** (e.g. DMDO) followed by treatment with an appropriate nucleophile (e.g. TMSN_3) under Lewis acidic conditions (e.g. SnCl_4) in a manner similar to that described in *J. Med. Chem.* **2007**, 50, 5463-5470, and removal of the hydroxyl

protecting groups (e.g. CsF) and nitrogen protecting group (e.g. NH_4OH) yields the final compounds of the type **S12b**.




Scheme 13.

5 Scheme 13 shows a general synthesis of compounds beginning with oxidation of the olefin **S12a** (e.g. DMDO) in the presence of the appropriate alcohol HOR^a and removal of the hydroxyl protecting groups (e.g. CsF). Removal of the nitrogen protecting group (e.g. NH₂Me) yields the final compounds of the type **S13a**.



10 **Scheme 14.**


Scheme 14 shows a general synthesis of compounds beginning with xanthate formation (e.g. PhOC(S)Cl, DMAP) followed by a Barton–McCombie deoxygenation reaction (e.g. (TMS)₃SiH, AIBN) to generate intermediate **S14a**. Removal of the hydroxyl protecting groups (e.g. TBAF), and nitrogen protecting group (e.g. NH₄OH) yields the final compounds of the type **S14b**.



Scheme 15.


Scheme 15 shows a general synthesis of compounds beginning with intermediate **S15a** prepared in a manner similar to that described in *Biosci. Biotech. Biochem.* **1993**, 57,

1433-1438. Removal of the acetate protecting groups using hydrolytic conditions (e.g. K_2CO_3 , MeOH), followed by chemoselective oxidation conditions (e.g. NIS, Bu_4NI), and protection of the 2' hydroxyl group (e.g. $BnBr$, Ag_2O) generates intermediate **S15b**. Lithium-halogen exchange (e.g. $n-BuLi$, $[-CH_2SiMe_2Cl]_2$) with an appropriate nucleobase **S3b** and addition to the lactone **S15b**, followed by reduction of the 1' hydroxyl group under Lewis acidic conditions (e.g. $BF_3 \cdot Et_2O$, Et_3SiH), and deprotection (e.g. H_2 , Pd black) yields the final compounds of the type **S15c**.




Scheme 16.

Scheme 16 shows a general synthesis of compounds beginning with conversion of the 5' hydroxyl group to the aldehyde under oxidative conditions (e.g. EDCI·HCl, Pyr, TFA, DMSO) followed by condensation of the corresponding enolate with formaldehyde and reduction (e.g. $NaBH_4$) to yields intermediate **S16a**. Sequential selective protection of the hydroxyl moieties with orthogonal protecting groups (e.g. DMTrCl and TBSCl) followed by removal of the more labile protecting group under acidic conditions (e.g. TsOH) then affords intermediate **S16b**. An Appel reaction (e.g. Ph_3P , CCl_4) then can convert the hydroxyl group into a chloride, and removal of the hydroxyl protecting groups (e.g. TBAF), and nitrogen protecting group (e.g. NH_4OH) yields the final compounds of the type **S16c**.




Scheme 17.

Scheme 17 shows a general synthesis of compounds beginning with protection of the free hydroxyl group of intermediate **S16b** with a labile protecting group (e.g. PMBCl, K_2CO_3). Selective removal of the 2', 3', and 5' silyloxy protecting groups (e.g. TBAF), followed by reprotection of with robust protecting groups (e.g. BnBr, NaH), and removal of the labile hydroxyl protecting group under acidic conditions (e.g. AcOH) affords intermediate **S17a**. Conversion of the hydroxyl group to the fluorine (e.g. DAST) followed by removal of the hydroxyl protecting groups (e.g. BBr_3), and nitrogen protecting group (e.g. NH_4OH) yields the final compounds of the type **S17b**.



Scheme 18.


Scheme 18 shows a general synthesis of compounds through appropriate electrophilic halogenation reactions of intermediate **S18a** to afford the final compounds of the type **S18b** (e.g. NCS), **S18c** (e.g. NIS), and **S18d** (e.g. Selectfluor).



Scheme 19.


Scheme 19 shows a general synthesis of compounds beginning with a cross-coupling reaction (e.g. $Zn(CN)_2$, $Pd(PtBu)_3$) to yield the final compounds of the type **S19a**.

Compound **S19a** can then be elaborated through a hydrolysis reaction of the nitrile (e.g. H_2O_2 , NH_4OH , H_2O) to afford compounds of the type **S19b**.




Scheme 20.

5 Scheme 20 shows a general synthesis of compounds beginning with a Sonogashira reaction (e.g. CuI , $\text{PdCl}_2(\text{PPh}_3)_2$) to yield the final compounds of the type **S20a**.




Scheme 21.

10 Scheme 21 shows a general synthesis of compounds beginning with a cross-coupling reaction (e.g. $\text{Pd}(\text{dppf})\text{Cl}_2$, Cs_2CO_3) to yield the final compounds of the type **S21a**.




Scheme 22.

Scheme 22 shows a general synthesis of compounds involving synthesis of phosphorylated analogs of the type **S22b**.



Scheme 23.


Scheme 23 shows a general synthesis of compounds involving synthesis of phosphorylated analogs of the type **S23b**.



Scheme 24.

Scheme 24 shows a general synthesis of compounds involving synthesis of phosphorylated analogs of the type **S24b**.

Experimentals



Intermediate 1b.

To a solution of intermediate **1a** (50 mg, 373 μ mol) in DMF (1 mL) was charged *N*-iodosuccinimide (84 mg, 373 μ mol) as a solid at RT. After 1.5 h, the reaction mixture was diluted 1M NaOH solution (10 mL), and the resulting slurry was stirred at RT. After 1 h, the solids were collected by vacuum filtration and dried under reduced pressure to afford intermediate **1b**.


^1H NMR (400 MHz, DMSO- d_6) δ 7.89 (s, 1H), 7.78 (br-s, 1H), 6.98 (d, J = 4.4 Hz, 1H), 6.82 (d, J = 4.4 Hz, 1H), 3.30 (br-s, 1H).

LC/MS: $t_R = 1.21$ min, MS $m/z = 261.02$ $[M+1]$; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm

Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-2.0 min 2-100% ACN, 2.0 min-3.05 min 100% ACN, 3.05 min-3.2 min 100%-2% ACN, 3.2 min-3.5 min 2% ACN at 2 μ l/min.

HPLC: $t_R = 1.536$ min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.



Intermediate 1d - (2S,3R,4R,5R)-2-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-3,4-bis(benzyloxy)-5-(benzyloxymethyl)tetrahydrofuran-2-ol.

n-Butyllithium (2.5M in hexanes, 34.4 mL, 86.0 mmol) was added rapidly to a suspension of 7-iodopyrrolo[1,2-f][1,2,4]triazin-4-amine **1b** (6.84 g, 26.3 mmol) and 1,2-bis(chlorodimethylsilyl)ethane (5.66 g, 26.3 mmol) in THF (200 mL) at -78 °C under an argon atmosphere. Over the course of the addition the internal temperature of the reaction mixture rose to -40.5 °C, and the reaction mixture became a clear brown solution. After 15 min, a solution of (3R,4R,5R)-3,4-bis(benzyloxy)-5-(benzyloxymethyl)dihydrofuran-2(3H)-one (**1c**, Purchased from Carbosynth, 10 g, 23.9 mmol) in tetrahydrofuran (40 mL) precooled to -78 °C, was added rapidly via cannula. After 1 h, the reaction mixture was quenched with acetic acid (15 mL), and the resulting mixture was allowed to warm to RT. The resulting mixture was diluted with ethyl acetate (800 mL) and was washed with saturated aqueous sodium bicarbonate solution (500 mL) and brine (500 mL). The organic layer was dried over anhydrous sodium sulfate, and was concentrated under reduced pressure. The crude residue was purified via SiO₂ column chromatography (220 g SiO₂ Combiflash HP Gold Column, 0–100% ethyl acetate/hexanes) to afford intermediate **1d**.


LC/MS: t_R = 1.50 min, MS m/z = 553.34 $[M+1]$; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm

Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-1.5 min 2-100% ACN, 1.5 min-2.2 min 100% ACN, 2.2 min-2.4 min 100%-2% ACN, 2.4 min-2.5 min 2% ACN.

HPLC: t_R = 3.442 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.

TLC: eluent: ethyl acetate, R_f = 0.5 (UV)



Intermediate 1e - 7-((2S,3S,4R,5R)-3,4-bis(benzyloxy)-5-(benzyloxymethyl)tetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-amine.

To a solution of intermediate **1d** (4.74 g, 8.58 mmol) and triethylsilane (3.56 mL, 22.3 mmol), in DCM (43 mL) was added boron trifluoride diethyl etherate (1.59 mL, 12.9 mmol) slowly via syringe at 0 °C under an argon atmosphere. After 2 h, the reaction mixture was slowly diluted with saturated aqueous sodium bicarbonate solution (100 mL), and the resulting mixture was extracted with ethyl acetate (2 x 150 mL), was dried over anhydrous sodium sulfate, and was concentrated under reduced pressure. The crude residue was purified via SiO₂ column chromatography (24 g SiO₂ Combiflash HP Gold Column, 0–100% ethyl acetate/hexanes) to afford intermediate **1e**.


¹H NMR (400 MHz, CDCl₃) δ 7.88 (s, 1H), 7.37 – 7.22 (m, 15H), 6.73 (d, J = 4.6 Hz, 1H), 6.71 (d, J = 4.6 Hz, 1H), 5.66 (d, J = 4.2 Hz, 1H), 4.71 (s, 2H), 4.60 (d, J = 12.0 Hz, 1H), 4.54 (s, 2H), 4.45 (d, J = 11.9 Hz, 1H), 4.39 (dt, J = 7.1, 3.6 Hz, 1H), 4.25 (t, J = 4.6 Hz, 1H), 4.14 – 4.10 (m, 1H), 3.78 (dd, J = 10.7, 3.4 Hz, 1H), 3.65 (dd, J = 10.7, 4.0 Hz, 1H).

LC/MS: t_R = 2.01 min, MS m/z = 537.41 $[M+1]$; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm. Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-1.5 min 2-100% ACN, 1.5 min-2.2 min 100% ACN, 2.2 min-2.4 min 100%-2% ACN, 2.4 min-2.5 min 2% ACN at 2 μ l/min.

- 5 HPLC: t_R = 3.596 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.

TLC: eluent: ethyl acetate, R_f = 0.3 (UV)



- 10 **Intermediate 1f - N-(7-((2S,3S,4R,5R)-3,4-bis(benzyloxy)-5-(benzyloxymethyl)tetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.**


- To a solution of intermediate **1e** (3.94 g, 7.34 mmol) in pyridine (36.7 mL) was added benzoyl chloride (1.69 mL, 14.68 mmol) slowly at RT under an argon atmosphere. After 1 h, additional benzoyl chloride (1.69 mL, 14.68 mmol) was added slowly. After 19 h, water (20 mL) was added slowly and the reaction mixture became slightly cloudy. Ammonium Hydroxide (~10 mL) was then added slowly until the reaction mixture was basic at pH=10. After 1 h, water (150 mL) was added dropwise via addition funnel and white solids slowly began to precipitate from the reaction mixture over the course of the addition. The resulting mixture was stirred for 24 h and the white solids were collected by vacuum filtration and were dried azeotropically from toluene to afford intermediate **1f**.

^1H NMR (400 MHz, CDCl_3) δ 8.23 (br s, 1H), 7.62 (t, J = 7.8 Hz, 1H), 7.53 (t, J = 7.6 Hz, 2H), 7.38 – 7.21 (m, 18H), 7.17 (d, J = 7.6 Hz, 1H), 5.69 (d, J = 4.1 Hz, 1H), 4.71 (s, 2H), 4.63 – 4.44 (m, 4H), 4.43 – 4.39 (m, 1H), 4.22 (t, J = 4.5 Hz, 1H), 4.15 – 4.10 (m, 1H), 3.79 (dd, J = 10.8, 3.2 Hz, 1H), 3.65 (dd, J = 10.7, 3.7 Hz, 1H).

- 25 LC/MS: t_R = 1.91 min, MS m/z = 641.18 [$M+1$]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-1.5 min 2-100% ACN, 1.5 min-2.2 min 100% ACN, 2.2 min-2.4 min 100%-2% ACN, 2.4 min-2.5 min 2% ACN at 2 μ l/min.

5 TLC: eluent: 50% ethyl acetate in hexanes, R_f = 0.6 (UV)



Intermediate 1g - N-(7-((2S,3R,4S,5R)-3,4-dihydroxy-5-(hydroxymethyl)tetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

10 Ethanol (68.5 mL) and formic acid (51.7 mL, 1.37 mol) were added sequentially to a mixture of intermediate **1f** (4.39 g, 6.85 mmol) and palladium on carbon (10% by wt, 2.2 g) at RT under an argon atmosphere. After 3 d, the reaction mixture was filtered through a pad of celite, and the filtrate was concentrated under reduced pressure. The crude residue was azeotroped with toluene (3 x 20 mL) to afford intermediate **1g**, which was used directly in the next step without further purification.

15 ^1H NMR (400 MHz, CD_3OD) δ 8.15 (s, 1H), 7.67 – 7.40 (m, 5H), 7.23 (d, J = 4.7 Hz, 1H), 7.00 (d, J = 4.7 Hz, 1H), 5.40 (d, J = 6.0 Hz, 1H), 4.44 (t, J = 5.7 Hz, 1H), 4.17 (t, J = 5.1 Hz, 1H), 4.03 (q, J = 4.3 Hz, 1H), 3.81 (dd, J = 12.1, 3.5 Hz, 1H), 3.71 (dd, J = 12.0, 4.5 Hz, 1H).


LC/MS: t_R = 1.04 min, MS m/z = 371.15 [$M+1$]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm

20 Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-1.5 min 2-100% ACN, 1.5 min-2.2 min 100% ACN, 2.2 min-2.4 min 100%-2% ACN, 2.4 min-2.5 min 2% ACN at 2 μ l/min. ;

HPLC: t_R = 2.055 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.

25



Intermediate 1h - N-(7-((2S,3R,4S,5R)-5-((bis(4-methoxyphenyl)(phenyl)methoxy)methyl)-3,4-dihydroxytetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

5 4,4'-Dimethoxytrityl chloride (2.23 g, 6.59 mmol) was added as a solid in one portion to a solution of intermediate **1g** (2.44 g, 6.59 mmol) in pyridine (32.5 mL) at RT. After 5.5 h, the reaction mixture was diluted with ethyl acetate (300 mL) and the resulting mixture was washed with brine (3 x 200 mL). The organic layer was concentrated under reduced pressure, and the crude residue was purified via SiO₂ column chromatography (80 g SiO₂ Combiflash HP Gold Column, 0–100% ethyl acetate/hexanes) to afford intermediate **1h**.
10


LC/MS: t_R = 1.68 min, MS m/z = 673.22 [M+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm;

Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-1.5 min 2-100% ACN, 1.5 min-2.2 min 100% ACN, 2.2 min-2.4 min 100%-2% ACN, 2.4 min-2.5 min 2% ACN at 2 μ L/min.
15

HPLC: t_R = 4.270 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.

TLC: eluent: 50% ethyl acetate in hexanes, R_f = 0.15 (UV)



20

Intermediate 1i - N-(7-((2S,3S,4R,5R)-5-((bis(4-methoxyphenyl)(phenyl)methoxy)methyl)-3,4-bis(tert-butylidimethylsilyloxy)tetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

tert-Butyldimethylsilyl chloride (2.47 g, 16.4 mmol) was added to a solution of intermediate **1h** (1.84 g, 2.74 mmol) and imidazole (2.23 g, 32.8 mmol) in *N,N*-dimethylformamide (28.2 mL) at RT. After 17 h, saturated aqueous sodium bicarbonate solution (500 mL) was added slowly to the reaction mixture. The resulting mixture was extracted with ethyl acetate (500 mL), and the organic layer was washed with brine (2 x 400 mL), was dried over anhydrous sodium sulfate, and was concentrated under reduced pressure. The crude residue was purified via SiO₂ column chromatography (80 g SiO₂ Combiflash HP Gold Column, 0–100% ethyl acetate/hexanes) to afford intermediate **1i**.


LC/MS: t_R = 3.43 min, MS m/z = 901.37 [M+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm

Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-2.0 min 2-100% ACN, 2.0 min-3.55 min 100% ACN, 3.55 min-4.2 min 100%-2% ACN at 2 μ L/min

HPLC: t_R = 5.724 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.

TLC: eluent: 50% ethyl acetate in hexanes, R_f = 0.75 (UV)



Intermediate 1j - N-(7-((2S,3S,4R,5R)-3,4-bis(tert-butylidimethylsilyloxy)-5-(hydroxymethyl)tetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

A solution of *p*-toluenesulfonic acid monohydrate (509 mg, 2.67 mmol) in methanol (3.7 mL) was slowly added to a solution of intermediate **1i** (2.41 g, 2.67 mmol) in

dichloromethane (22.3 mL) at 0 °C. After 1.5 h, the reaction mixture was diluted with saturated aqueous bicarbonate solution (100 mL), and the resulting mixture was extracted with dichloromethane (2 x 100 mL). The combined organic extracts were dried over anhydrous sodium sulfate, and were concentrated under reduced pressure. The crude residue was purified via SiO₂ column chromatography (120 g SiO₂ Combiflash HP Gold Column, 0–100% ethyl acetate/hexanes) to afford intermediate **1j**.

¹H NMR (400 MHz, CDCl₃) δ 8.72 (br-s, 1H), 8.16 (br-t, *J* = 7.1 Hz, 2H), 8.07 (br-t, *J* = 7.7 Hz, 3H), 7.49 – 7.43 (m, 1H), 5.75 (d, *J* = 8.2 Hz, 1H), 5.28 (dd, *J* = 8.1, 4.7 Hz, 1H), 4.81 (d, *J* = 5.0 Hz, 1H), 4.70 – 4.63 (m, 1H), 4.44 (d, *J* = 12.3 Hz, 1H), 4.24 (d, *J* = 12.4 Hz, 1H), 1.48 (s, 9H), 1.30 (s, 9H), 0.65 (s, 3H), 0.64 (s, 3H), 0.41 (s, 3H), 0.00 (s, 3H).


LC/MS: *t*_R = 2.66 min, MS *m/z* = 599.19 [*M*+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ XB-C18 100A, 50 x 4.6 mm

Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min–2.0 min 2–100% ACN, 2.0 min–3.05 min 100% ACN, 3.05 min–3.2 min 100%–2% ACN, 3.2 min–3.5 min 2% ACN at 2μl/min.

HPLC: *t*_R = 5.622 min; HPLC system: Agilent 1100 series.; Column: Gemini 5μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min–5.0 min 2–98% ACN, 5.0 min–6.0 min 98% ACN at 2 mL/min.

TLC: eluent: 50% ethyl acetate in hexanes, *R*_f = 0.55 (UV)



Intermediate 1k - N-(7-((2*S*,3*S*,4*R*,5*S*)-3,4-bis(tert-butyldimethylsilyloxy)-5-(iodomethyl)tetrahydrofuran-2-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4-yl)benzamide.

Intermediate **1j** (1.19 g, 1.99 mmol) was added to a solution of methyltriphenoxyphosphonium iodide (0.99 g, 2.19 mmol) in DMF (9.9 mL) at RT. After 3 h, an additional portion of methyltriphenoxyphosphonium iodide (0.99 g, 2.19 mmol) was

added. After 1 h, the reaction mixture was diluted with ethyl acetate (200 mL) and was washed with brine (3 x 100 mL). The organic layer was dried over anhydrous sodium sulfate, and was concentrated under reduced pressure. The crude residue was purified via SiO₂ column chromatography (80 g SiO₂ Combiflash HP Gold Column, 0–100% ethyl acetate/hexanes) to afford intermediate **1k**.


¹H NMR (400 MHz, CDCl₃) δ 8.21 (br s, 1H), 7.61 (br t, *J* = 7.2 Hz, 1H), 7.53 (br t, *J* = 7.5 Hz, 3H), 7.05 (br s, 1H), 5.44 (d, *J* = 4.5 Hz, 1H), 4.52 (t, *J* = 4.3 Hz, 1H), 4.08 – 3.99 (m, 2H), 3.55 (dd, *J* = 10.7, 5.2 Hz, 1H), 3.38 (dd, *J* = 10.7, 5.0 Hz, 1H), 0.93 (s, 9H), 0.85 (s, 9H), 0.16 (s, 3H), 0.11 (s, 3H), -0.01 (s, 3H), -0.11 (s, 1H).

LC/MS: *t_R* = 3.06 min, MS *m/z* = 709.16 [M+1]; LC system: Thermo Accela 1250 UHPLC
MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ XB-C18 100A, 50 x 4.6 mm

Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-2.0 min 2-100% ACN, 2.0 min-3.05 min 100% ACN, 3.05 min-3.2 min 100%-2% ACN, 3.2 min-3.5 min 2% ACN at 2μl/min.

HPLC: *t_R* = 5.837 min; HPLC system: Agilent 1100 series.; Column: Gemini 5μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.

TLC: eluent: 20% ethyl acetate in hexanes, *R_f* = 0.45 (UV)



Intermediate 1l - N-(7-((2S,3S,4S)-3,4-bis(tert-butyldimethylsilyloxy)-5-methylenetetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

Potassium *t*-butoxide (700 mg, 6.24 mmol) was added to a solution of intermediate **1k** (1.77 g, 2.5 mmol) in pyridine (25 mL) at RT. After 2 h, the reaction mixture was diluted with saturated aqueous sodium bicarbonate solution (25 mL) and brine (200 mL). The resulting mixture was extracted with ethyl acetate (300 mL). The organic layer was then

washed with brine (200 mL), was dried over anhydrous sodium sulfate, and was concentrated under reduced pressure. The crude residue was purified via SiO₂ column chromatography (40 g SiO₂ Combiflash HP Gold Column, 0–100% ethyl acetate/hexanes) to afford intermediate **1l**.

5 LC/MS: t_R = 2.87 min, MS m/z = 581.37 [M+1]; LC system: Thermo Accela 1250 UHPLC


MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm

Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid

Gradient: 0 min–2.0 min 2–100% ACN, 2.0 min–3.05 min 100% ACN, 3.05 min–3.2 min 100%–2% ACN, 3.2 min–3.5 min 2% ACN at 2 μ L/min.

10 HPLC: t_R = 5.750 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min–5.0 min 2–98% ACN, 5.0 min–6.0 min 98% ACN at 2 mL/min.

TLC: eluent: 50% ethyl acetate in hexanes, R_f = 0.20 (UV)



15 **Intermediate 1m - N-(7-((2S,3S,4S)-3,4-bis(tert-butyldimethylsilyloxy)-5-hydroxy-5-(hydroxymethyl)tetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.**


DMDO (0.07M solution in acetone, 13.8 mL, 0.964 mmol) was added to a solution of intermediate **1l** (560 mg, 0.964 mmol) in acetone (4.82 mL) at 0 °C. After 10 min, the reaction mixture was concentrated under reduced pressure was dried azeotropically with toluene (2 x 1 mL) to afford **1m** that was used immediately in the next step without further purification.

20 LC/MS: t_R = 2.57 min, MS m/z = 615.14 [M+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm

Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid

Gradient: 0 min-2.0 min 2-100% ACN, 2.0 min-3.05 min 100% ACN, 3.05 min-3.2 min 100%-2% ACN, 3.2 min-3.5 min 2% ACN at 2 μ l/min.




5 Intermediate 1n - N-(7-((2S,3S,4S)-3,4-bis(tert-butylidimethylsilyloxy)-5-cyano-5-((trimethylsilyloxy)methyl)tetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

To a solution of crude intermediate **1m** (~592 mg, ~0.964 mmol) and trimethylsilyl cyanide (640 μ l, 4.80 mmol) in dichloromethane (19.2 mL) was added indium (III) bromide (681 mg, 1.92 mmol) at 0 °C under an argon atmosphere. After 4.5 h, the reaction mixture was quenched with saturated aqueous sodium bicarbonate solution (6 mL) and was allowed to warm to RT. The resulting mixture was partitioned between dichloromethane (20 mL) and saturated aqueous sodium bicarbonate solution (20 mL). The phases were split and the aqueous layer was extracted with dichloromethane (20 mL). The combined organic layers were dried over anhydrous sodium sulfate and were concentrated under reduced pressure to afford intermediate **1n** (1:1 diastereomeric mixture) (710 mg) that was used directly in the next step without further purification.

LC/MS: first eluting isomer t_r = 2.91 min, MS m/z = 696.28 [M+1], second eluting isomer t_R = 3.02 min, MS m/z = 696.19 [M+1],; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm

Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-2.0 min 2-100% ACN, 2.0 min-3.05 min 100% ACN, 3.05 min-3.2 min 100%-2% ACN, 3.2 min-3.5 min 2% ACN at 2 μ l/min.




Intermediate 1o - N-(7-((2S,3R,4S)-5-cyano-3,4-dihydroxy-5-(hydroxymethyl)tetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

To a solution of crude intermediate **1n** (668.23 mg, 0.96 mmol) in DMF (9.6 mL) was added cesium fluoride (729 mg, 4.8 mmol) at RT. After 5 h, the reaction mixture was diluted with brine (100 mL), and the resulting mixture was extracted with dichloromethane (3 x 100 mL). The combined organic layers were dried over anhydrous sodium sulfate and were concentrated under reduced pressure to afford intermediate **1o** (1:1 diastereomeric mixture) that was used directly in the next step without further purification.

LC/MS: first eluting isomer $t_R = 1.31$ min, MS $m/z = 396.19$ [M+1], second eluting isomer $t_R = 1.32$ min, MS $m/z = 396.19$ [M+1],; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm

Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-2.0 min 2-100% ACN, 2.0 min-3.05 min 100% ACN, 3.05 min-3.2 min 100%-2% ACN, 3.2 min-3.5 min 2% ACN at 2 μ L/min.



Example 1 - (2R,3S,4R,5S)-5-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-3,4-dihydroxy-2-(hydroxymethyl)tetrahydrofuran-2-carbonitrile.

Methylamine (40% in water, 0.3 mL) was added to a solution of crude intermediate **1o** in methanol (1 mL) at RT. After 2.5 h, the reaction mixture was concentrated under

reduced pressure, and was directly purified by preparatory HPLC (Phenominex Luna 5u C18 100Å 100 x 30 mm column, 5-15% acetonitrile/water gradient, 25 min). The fractions containing the desired product and the 4' anomer were combined and were concentrated under reduced pressure. The 4' anomers were then separated by preparatory HPLC (Phenominex Luna 5u C18 100Å 100 x 30 mm column, 5-15% acetonitrile/water gradient, 25 min). The fractions containing the desired product were combined and were lyophilized to afford Example 1.


¹H NMR (400 MHz, CD₃OD) δ 7.79 (s, 1H), 6.85 (d, *J* = 4.5 Hz, 1H), 6.76 (d, *J* = 4.5 Hz, 1H), 5.45 (d, *J* = 5.9 Hz, 1H), 4.59 (t, *J* = 5.7 Hz, 1H), 4.40 (d, *J* = 5.6 Hz, 1H), 3.88 (d, *J* = 12.0 Hz, 1H), 3.80 (d, *J* = 12.0 Hz, 1H).

LC/MS: *t_R* = 0.29 min, MS *m/z* = 292.16 [M+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ XB-C18 100A, 50 x 4.6 mm

Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-1.5 min 2-100% ACN, 1.5 min-2.2 min 100% ACN, 2.2 min-2.4 min 100%-2% ACN, 2.4 min-2.5 min 2% ACN at 2μl/min.

HPLC: *t_R* = 0.377 min; HPLC system: Agilent 1100 series.; Column: Gemini 5μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.; HPLC: *t_R* = 6.643 min; HPLC system: Agilent 1100 series.; Column: Luna 5μ C18(2) 110A, 250 x 4.6 mm; Solvents: Acetonitrile, Water; Gradient: 5-15% ACN over 10 min at 2 mL/min



Intermediate 2b - N-(7-((2S,3R,4R,5R)-3-fluoro-4-hydroxy-5-(hydroxymethyl)tetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

To an N₂ purged flask was added intermediate **2a**, (2R,3R,4R,5S)-5-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-4-fluoro-2-(hydroxymethyl)tetrahydrofuran-3-ol (prepared according to WO2012037038A1, 1.20 g, 4.01 mmol) (dried by coevaporation with

pyridine 3 times), which was then dissolved in pyridine (18 mL). Chlorotrimethylsilane (1.54 mL, 13.13 mmol) was added in one portion at 0 °C and the resulting mixture was stirred under an N₂ atmosphere for 1 h. Benzoyl chloride (675 μL, 5.82 mmol) was added dropwise and the reaction mixture was stirred for 1h. An additional portion of benzoyl chloride (100 μL) was added to consume the remaining starting material. A mixture of mono- and bis-Bz protected products was observed. The reaction was quenched with H₂O (5 mL), and the resulting mixture was stirred for 5 min. Then concentrated NH₄OH_(aq) (8 mL) was added in one portion and allowed to stir for 15 min at which point the bis-Bz product was converted to the desired product. The solvents were removed under reduced pressure, and then the residue was coevaporated with CH₃OH. Intermediate **2b** was isolated after purification by silica gel chromatography, using an eluent ramp of 50%-100% EtOAc in hexanes.

¹H NMR (400 MHz, DMSO-*d*₆) δ 8.26 – 7.91 (m, 2H), 7.88 – 7.79 (m, 1H), 7.61 (t, *J* = 7.5 Hz, 1H), 7.56 – 7.36 (m, 2H), 7.37 – 7.24 (m, 1H), 7.10 (d, *J* = 4.6 Hz, 1H), 7.01 (s, 1H), 5.53 (d, *J* = 23.4 Hz, 1H), 5.45 (d, *J* = 6.5 Hz, 1H), 5.16 – 4.91 (m, 1H), 4.86 (t, *J* = 5.6 Hz, 1H), 4.21 – 4.04 (m, 1H), 3.82 (dd, *J* = 8.0, 3.9 Hz, 1H), 3.71 (ddd, *J* = 12.3, 5.6, 2.6 Hz, 1H), 3.52 (ddd, *J* = 12.2, 5.7, 4.5 Hz, 1H).


¹⁹F NMR (376 MHz, DMSO-*d*₆) δ -196.33 (dt, *J* = 55.1, 22.5 Hz).

LC/MS: *t*_R = 0.77 min, MS *m/z* = 373.14 [*M*+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN.



Intermediate 2c - N-(7-((2S,3R,4R,5R)-5-((bis(4-methoxyphenyl)(phenyl)methoxy)methyl)-3-fluoro-4-hydroxytetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

Intermediate **2b** (1.3 g, 3.49 mmol) was dried by coevaporation with pyridine. The dried material was then dissolved in pyridine (15 mL) under an N₂ atmosphere. 4,4'-Dimethoxytrityl chloride (1.71 g, 5.0 mmol) was added in one portion at room temperature and allowed to stir for 2 h. Ethanol (2 mL) was added and the resulting solution was stirred for 5 min. Solvents were removed under reduced pressure. The residue was purified by silica gel chromatography, using an eluent ramp of 0%-100% EtOAc in hexanes, to afford intermediate **2c**.

¹H NMR (400 MHz, DMSO-*d*₆) δ 8.28 – 8.02 (m, 2H), 7.61 (t, *J* = 7.5 Hz, 1H), 7.51 (t, *J* = 7.6 Hz, 2H), 7.36 (ddt, *J* = 6.0, 4.7, 2.0 Hz, 2H), 7.31 – 7.04 (m, 7H), 6.90 – 6.73 (m, 4H), 5.62 (d, *J* = 24.4 Hz, 1H), 5.49 (d, *J* = 7.0 Hz, 1H), 5.27 – 5.01 (m, 1H), 4.32 – 4.13 (m, 1H), 4.06 – 3.95 (m, 1H), 3.69 (s, 4H), 3.28 (s, 3H), 3.12 (dd, *J* = 10.4, 5.2 Hz, 1H).


¹⁹F NMR (376 MHz, DMSO-*d*₆) δ -195.58 (dt, *J* = 52.1, 24.7 Hz).

LC/MS: *t*_R = 1.37 min, MS *m/z* = 675.29 [M+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN.



Intermediate 2d - N-(7-((2S,3S,4R,5R)-4-(tert-butyldimethylsilyloxy)-3-fluoro-5-(hydroxymethyl)tetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

To a solution of intermediate **2c** (1.47 g, 2.18 mmol) in DMF (8 mL), prepared under an N₂ atmosphere, was added imidazole (251 mg, 3.70 mmol), followed by *tert*-

butylchlorodimethylsilane (492 mg, 3.27 mmol). The solution was allowed to stir at room temperature for 16 h. The solution was diluted with H₂O (5 mL) and then the solvents were removed under reduced pressure. The residue was partitioned between EtOAc and H₂O. The layers were separated and the organic phase was then washed with brine. The organics were dried over Na₂SO₄, which was removed by filtration, and the filtrate was concentrated under reduced pressure. The crude material was used as is in the next step.

The crude material was dissolved in CHCl₃ (15 mL) and cooled to 0 °C. *p*-Toluenesulfonic acid hydrate (414 mg, 2.18 mmol) dissolved in CH₃OH (6 mL) was added dropwise to the mixture and was allowed to stir for 15 min. The reaction was quenched with sat. NaHCO_{3(aq)}. The organics were washed with brine and dried over Na₂SO₄, filtered and the solvent was removed under reduced pressure. The residue was purified by silica gel chromatography, using an eluent ramp of 0%-50% EtOAc in hexanes, to afford intermediate **2d**.

¹H NMR (400 MHz, DMSO-*d*₆) δ 8.21 – 7.98 (m, 2H), 7.51 (dt, *J* = 39.9, 7.5 Hz, 3H), 7.12 – 6.86 (m, 2H), 5.48 (d, *J* = 21.9 Hz, 1H), 5.07 (dt, *J* = 54.6, 3.8 Hz, 1H), 4.86 (t, *J* = 5.5 Hz, 1H), 4.28 (ddd, *J* = 17.8, 7.1, 4.4 Hz, 1H), 3.83 – 3.72 (m, 1H), 3.63 (ddd, *J* = 12.1, 5.2, 3.0 Hz, 1H), 3.43 (ddd, *J* = 12.1, 6.0, 4.2 Hz, 1H), 0.80 (s, 9H), 0.01 (s, 3H), 0.00 (s, 3H).


¹⁹F NMR (376 MHz, DMSO-*d*₆) δ -198.05 (dt, *J* = 54.2, 19.8 Hz).

LC/MS: *t*_R = 1.35 min, MS *m/z* = 487.24 [*M*+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN.



Intermediate 2e - N-(7-((2S,3S,4R)-4-(tert-butyldimethylsilyloxy)-3-fluoro-5,5-bis(hydroxymethyl)tetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

To solution of intermediate **2d** (856 mg, 1.75 mmol) in toluene (4 mL), and DMSO (6 mL), prepared under an N₂ atmosphere, was added 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride (EDCI) (504 mg, 2.63 mmol). To this mixture was added pyridine (150 µL), and TFA (70 µL) and the mixture was stirred at RT for 15 min. Additional EDCI (100 mg) and pyridine (100 µL) was added and the mixture was stirred for an additional 45 min. The reaction was quenched with H₂O (10 mL) and CH₂Cl₂ (10 mL). The organics were washed with brine and dried over Na₂SO₄, filtered, and the solvent was removed under reduced pressure to afford the crude aldehyde, which was used as is for the next step.

The crude aldehyde was dissolved in dioxane (5 mL) and 37% formaldehyde_(aq) (925 µL) was added followed by 2N NaOH_(aq) (925 µL). After stirring at room temperature for 3 h, the reaction was quenched with AcOH, diluted with EtOAc, and washed with H₂O. The organics were dried over Na₂SO₄, filtered, and the solvent was removed under reduced pressure to afford the crude aldol product, which was carried forward as is to the next step.

The crude aldol product was dissolved in EtOH (9 mL) under an N₂ atmosphere and cooled to 0 °C. NaBH₄ (80 mg, 2.1 mmol) was added in one portion and the reaction was stirred for 10 min. The reaction was quenched with AcOH, diluted with CH₂Cl₂ and washed with a 1:1 solution of water and sat. NaHCO_{3(aq)}. The organic phase was dried over Na₂SO₄, filtered, and the solvent was removed under reduced pressure. The residue was purified by silica gel chromatography, using an eluent ramp from 0%-100% EtOAc in hexanes, to afford intermediate **2e**.

¹H NMR (400 MHz, DMSO-*d*₆) δ 8.20 – 7.99 (m, 2H), 7.51 (dt, *J* = 39.5, 7.5 Hz, 3H), 7.15 – 6.93 (m, 2H), 5.50 (d, *J* = 14.1 Hz, 1H), 5.24 (dt, *J* = 54.2, 5.4 Hz, 1H), 4.78 (t, *J* = 5.6 Hz, 1H), 4.48 (dd, *J* = 10.7, 4.8 Hz, 1H), 4.34 (dd, *J* = 6.7, 4.9 Hz, 1H), 3.62 (dd, *J* = 11.9, 4.9 Hz, 1H), 3.55 – 3.35 (m, 3H), 0.82 (s, 9H), 0.07 (s, 3H), -0.09 (s, 3H).

¹⁹F NMR (376 MHz, DMSO-*d*₆) δ -200.37 (d, *J* = 51.1 Hz).

LC/MS: *t*_R = 1.25 min, MS *m/z* = 517.21 [*M*+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6µ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN.



5 Intermediate 2f - N-(7-((2S,3S,4R,5R)-5-((bis(4-methoxyphenyl)(phenyl)methoxy)methyl)-4-(tert-butyldimethylsilyloxy)-5-((tert-butyldimethylsilyloxy)methyl)-3-fluorotetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

Intermediate **2e** (370 mg, 0.717 mmol) was dissolved in CH₂Cl₂ (10 mL) and TEA (200 μL) under an N₂ atmosphere, and was then cooled to 0 °C. 4,4'-Dimethoxytrityl chloride 4,4'-Dimethoxytrityl chloride (0.364g, 1.07mmol) was added in one portion and the reaction mixture was allowed to stir for 30 min. CH₃OH (2 mL) was added and the solution was diluted with CH₂Cl₂ and sat. NaHCO_{3(aq)}. The organic layer was washed with brine, dried over Na₂SO₄, filtered, and the solvents were removed under reduced pressure. The residue was purified by silica gel chromatography, using an eluent ramp of 5%-100% EtOAc in hexanes, to afford the crude product as a mixture of bis-DMTr and 4'β products. This mixture was carried forward without further purification.


To the crude product (574 mg, mixture) in DMF (3 mL), under an N₂ atmosphere, was added imidazole (143 mg, 2.10 mmol), followed by *tert*-butylchlorodimethylsilane (158 mg, 1.05 mmol). The solution was allowed to stir at room temperature for 2 h. The solution was diluted with CH₃OH (1 mL) and EtOAc. The organics were washed with H₂O and then with brine. The organic layer was dried over Na₂SO₄, filtered, and the solvent was removed under reduced pressure. The residue was purified by silica gel chromatography, using an eluent ramp of 0%-50% EtOAc in hexanes, to afford intermediate **2f** that contained some bis-DMTr material.

LC/MS: *t_R* = 2.25 min, MS *m/z* = 933.52 [M+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-1.5 min 2-100% ACN, 1.5 min-2.8 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3 min 2% ACN.



5

Intermediate 2g - N-(7-((2S,3S,4R,5R)-4-(tert-butyldimethylsilyloxy)-5-((tert-butyldimethylsilyloxy)methyl)-3-fluoro-5-(hydroxymethyl)tetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

Intermediate **2f** was dissolved in CHCl₃ (5 mL) and was cooled to 0 °C. *p*-Toluenesulfonic acid hydrate (90 mg, 0.474 mmol), dissolved in CH₃OH (4 mL) was added dropwise to the mixture and the reaction mixture was allowed to stir for 5 min. The reaction mixture was quenched with sat. NaHCO_{3(aq)}. The organics were washed with brine, dried over Na₂SO₄, filtered, and the solvent was removed under reduced pressure. The residue was purified by silica gel chromatography, using an eluent ramp of 0%-40% EtOAc in hexanes, to afford intermediate **2g**.

15

¹H NMR (400 MHz, CDCl₃) δ 8.09 – 7.88 (m, 2H), 7.48 (dt, *J* = 35.4, 7.4 Hz, 3H), 7.30 (d, *J* = 4.6 Hz, 1H), 6.88 (s, 1H), 5.69 (dd, *J* = 18.8, 4.2 Hz, 1H), 5.05 (dt, *J* = 54.6, 4.7 Hz, 1H), 4.62 (dd, *J* = 14.9, 5.1 Hz, 1H), 3.91 – 3.64 (m, 3H), 0.92 – 0.70 (m, 18H), 0.13 – 0.04 (m, 6H), 0.01 (m, 6H).

¹⁹F NMR (376 MHz, CDCl₃) δ -196 (m)


20

LC/MS: *t*_R = 2.61 min, MS *m/z* = 631.43 [M+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-1.5 min 2-100% ACN, 1.5 min-2.8 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3 min 2% ACN.



Intermediate 2h - N-(7-((2S,3S,4R,5R)-5-((E)-2-bromovinyl)-4-(tert-

5 butyldimethylsilyloxy)-5-((tert-butyldimethylsilyloxy)methyl)-3-fluorotetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

To solution of intermediate **2g** (228 mg, 0.361 mmol) in toluene (0.75 mL), and DMSO (0.15 mL), was added 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDCI) (208 mg, 1.08 mmol) under an N₂ atmosphere. To this mixture was added pyridine (30 μ L), and TFA (15 μ L) and the mixture was stirred at RT for 30 min. The reaction was diluted with EtOAc and washed with H₂O followed by brine. The organics were dried over Na₂SO₄, filtered, and solvent was removed under reduced pressure to afford the crude aldehyde. This material was used as is for the next step.

To a suspension of bromomethyltriphenylphosphonium bromide (314 mg, 0.72 mmol) in THF (4 mL) at -40°C was added KOtBu (1.0M in THF, 1.08 mL, 1.08 mmol) and the reaction mixture was stirred for 2 h under an N₂ atmosphere. The crude aldehyde was dissolved in THF (4 mL) and added dropwise. The reaction mixture was removed from the cold bath and allowed to warm to 10 °C over 1 h. The reaction mixture was re-cooled to -40 °C and the reaction mixture was quenched with sat NH₄Cl_(aq). The layers were separated and the organics were dried over Na₂SO₄, filtered, and solvent was removed under reduced pressure. The residue was purified by silica gel chromatography, using an eluent of 0%-50% EtOAc in hexanes, to afford intermediate **2h**.


¹H NMR (400 MHz, CDCl₃) δ 8.08 – 7.85 (m, 2H), 7.62 – 7.36 (m, 2H), 7.34 – 7.01 (m, 3H), 6.92 (s, 1H), 6.60 – 6.45 (m, 1H), 6.35 (d, *J* = 8.2 Hz, 1H), 5.61 (d, *J* = 25.1 Hz, 1H), 4.82 (dd, *J* = 56.3, 4.9 Hz, 1H), 4.69 – 4.46 (m, 1H), 3.97 (d, *J* = 11.4 Hz, 1H), 3.53 (d, *J* = 11.4 Hz, 1H), 0.84 (d, *J* = 3.9 Hz, 18H), 0.13 – -0.10 (m, 12H).

^{19}F NMR (376 MHz, CDCl_3) δ -190.60 (m).

LC/MS: t_R = 2.10 min, MS m/z = 705.54/707.29 [$M+1$]; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

- 5 Gradient: 0 min-1.5 min 2-100% ACN, 1.5 min-2.8 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3 min 2% ACN.



- 10 **Intermediate 2i** - N-(7-((2S,3S,4R,5R)-4-(tert-butyldimethylsilyloxy)-5-((tert-butyldimethylsilyloxy)methyl)-5-ethynyl-3-fluorotetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

- Intermediate **2h** (204 mg, 0.289 mmol) was dissolved in THF (8 mL) under an N_2 atmosphere and cooled to -40°C . KOtBu (1.0M in THF, 1.08 mL, 1.08 mmol) was slowly added. The reaction was allowed to stir for 20 min as was quenched with sat. $\text{NH}_4\text{Cl}_{(\text{aq})}$. The solution was diluted with EtOAc and washed with brine. The organics were dried over
15 Na_2SO_4 , filtered, and solvent was removed under reduced pressure. The residue was purified by silica gel chromatography, using an eluent ramp of 0%-50% EtOAc in hexanes, to afford intermediate **2i**.

- ^1H NMR (400 MHz, CDCl_3) δ 8.12 – 7.88 (m, 2H), 7.51 (dt, J = 36.5, 7.5 Hz, 2H), 7.32 (d, J = 4.6 Hz, 1H), 6.88 (s, 1H), 5.79 (d, J = 22.1 Hz, 1H), 5.02 (ddd, J = 55.3, 5.1, 3.2 Hz, 1H),
20 4.56 (dd, J = 18.1, 5.1 Hz, 1H), 3.91 (d, J = 11.3 Hz, 1H), 3.83 – 3.62 (m, 1H), 2.55 (m, 1H), 0.90 (dd, J = 25.3, 1.6 Hz, 18H), 0.20 – -0.08 (m, 12H).


^{19}F NMR (376 MHz, CDCl_3) δ -193.10 (broad-s).

LC/MS: t_R = 1.88 min, MS m/z = 625.24 [$M+1$]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-1.5 min 2-100% ACN, 1.5 min-2.8 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3 min 2% ACN.



5 Intermediate 2j - N-(7-((2S,3R,4R,5R)-5-ethynyl-3-fluoro-4-hydroxy-5-(hydroxymethyl)tetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

To a solution of intermediate **2i** (152 mg, 0.243 mmol) in THF (3.5 mL) under an N₂ atmosphere was added TBAF (1.0M in THF, 700 µL, 0.700 mmol) at room temperature and the mixture was allowed to stir for 30 min. Solvents were removed under reduced pressure.


10 The residue was purified by silica gel chromatography, using an eluent ramp of 40%-100% EtOAc in hexanes, to afford intermediate **2j**.

LC/MS: t_R = 0.88 min, MS m/z = 397.16 [M+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6µ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

15 Gradient: 0 min-1.5 min 2-100% ACN, 1.5 min-2.8 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3 min 2% ACN.



Example 2 - (2R,3R,4R,5S)-5-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-2-ethynyl-4-fluoro-2-(hydroxymethyl)tetrahydrofuran-3-ol.

To a solution of intermediate **2j** (71 mg, 0.179 mmol) in CH₃OH (2 mL) was added con. NH₄OH_(aq) (0.7 mL) and the resulting solution was stirred at room temperature for 16 h. Solvents were removed under reduced pressure. The residue was purified by silica gel chromatography, using an eluent ramp of 0%-20% CH₃OH in CH₂Cl₂, followed by reverse phase HPLC, using an eluent ramp of 0%-20% ACN in H₂O, to yield example **2**.

¹H NMR (400 MHz, CD₃OD) δ 7.77 (s, 1H), 6.90 – 6.70 (m, 2H), 5.62 (dd, *J* = 25.5, 2.6 Hz, 1H), 5.18 (ddd, *J* = 56.0, 5.4, 2.7 Hz, 1H), 4.57 (dd, *J* = 20.5, 5.4 Hz, 1H), 3.93 – 3.59 (m, 2H), 3.02 (d, *J* = 0.7 Hz, 1H).

¹⁹F NMR (376 MHz, CD₃OD) δ -193.76 (ddd, *J* = 56.0, 25.5, 20.4 Hz).

LC/MS: *t_R* = 0.45 min, MS *m/z* = 293.13 [M+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid


Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN.

HPLC: *t_R* = 3.112 min; HPLC system: Agilent 1100 series.

Column: Phenomenex Kinetex C18 2.6 μm 100A, 4.6 x 100 mm

Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA

Gradient: 0 min-8.0 min 2-98% ACN at 1.5 mL/min.



Intermediate 3a - N-(7-((2S,3S,4R,5R)-4-(tert-butyldimethylsilyloxy)-5-((tert-butyldimethylsilyloxy)methyl)-3-fluoro-5-formyltetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

To a solution of intermediate **2g** (1.13 g, 1.79 mmol) in DMSO (1 mL) and toluene (10 mL), prepared under an N₂ atmosphere, was added EDCI·HCl (1.02 g, 5.36 mmol) and

pyridine (149 μ L, 1.92 mmol). TFA (74 μ L, 0.97 mmol) was added drop-wise. After 1 hr the reaction was checked by LC/MS. A single peak was observed, with a retention time similar to the starting material's, but with an M+1 peak equal to that expected for the product.


Another 50 μ L of pyridine was added and the reaction was stirred for another 15 min. No change by LC/MS. The reaction was diluted with EtOAc and quenched with a 1:1 mixture of sat. $\text{NaHCO}_3(\text{aq.})$ and H_2O . The mixture was partitioned between EtOAc and more H_2O . The organic layer was separated and washed with H_2O , brine, and then dried over Na_2SO_4 . The drying agent was removed by vacuum filtration and the filtrate was concentrated. The residue was taken up in CH_2Cl_2 , concentrated, and the resulting material was placed under high vacuum for 1 h. The product, intermediate **3a** was used as is in the next reaction.

LC/MS: $t_R = 1.90$ min, MS $m/z = 629.46$ [M+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-1.5 min 2-100% ACN, 1.5 min-2.8 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3 min 2% ACN.



Intermediate 3b - N-(7-((2S,3S,4R,5R)-4-(tert-butyldimethylsilyloxy)-5-((tert-butyldimethylsilyloxy)methyl)-3-fluoro-5-((E)-(hydroxyimino)methyl)tetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

To a solution of intermediate **3a** (crude material from previous step, assumed 1.79 mmol) in pyridine (11 mL), prepared under an N_2 atmosphere, was added $\text{HONH}_2 \cdot \text{HCl}$ in one portion at room temperature. The reaction was checked by LC/MS 5 minutes later; starting material was consumed. The reaction was checked again 25 minutes later. There was no change from the first time point. The reaction was concentrated and the residue was partitioned between EtOAc and H_2O . The organic layer was separated, washed with brine, dried over Na_2SO_4 and filtered. The filtrate was concentrated, taken up in CH_2Cl_2 ,


concentrated again, and the residue was placed under high vacuum. The product intermediate **3b** was used as is in the next reaction.

LC/MS: $t_R = 1.83$ min, MS $m/z = 644.55$ $[M+1]$; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ C18 100A, 50 x 3.00 mm

5 Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-1.5 min 2-100% ACN, 1.5 min-2.8 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3 min 2% ACN.



10 **Intermediate 3c - N-(7-((2S,3S,4R,5R)-4-(tert-butyldimethylsilyloxy)-5-((tert-butyldimethylsilyloxy)methyl)-5-cyano-3-fluorotetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.**

To a solution of intermediate **3b** (crude material from the previous step, assumed 1.79 mmol) in ACN (16 mL) was added CDI (436 mg, 2.69 mmol) in a single portion. The reaction was run under an N₂ atmosphere. The reaction was checked by LC/MS after 20 minutes. Peaks with the mass of starting material and product are barely time resolved. The reaction was checked 1.5 h later. The UV peak corresponding to the starting material was nearly gone and the intensity of the mass peak was diminished. The reaction was diluted with CH₂Cl₂ and quenched with a 1:1 mixture of sat. NaHCO_{3(aq.)} and H₂O. The layers were separated, the aqueous was back extracted with CH₂Cl₂ and the combined organics layers were extracted with a 1:1 mixture of brine and H₂O, dried over Na₂SO₄ and filtered. The filtrate was concentrated and intermediate **3c** was isolated by silica gel column chromatography using the following solvent ramp: 0% EtOAc in hexanes ramping to 20% EtOAc in hexanes, pause at 20% EtOAc in hexanes and then ramping to 40% EtOAc in hexanes.

25 ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.31 (s, 1H), 8.07 (s, 1H), 7.65 (t, $J = 8$ Hz, 1H), 7.54 (t, $J = 8$ Hz, 1H), 7.15 (d, 4 Hz, 1H), 7.02 (s, 1H), 5.82 (d, $J = 24$ Hz, 1H), 5.51 (ddd, $J = 52, 4.8,$

2.8 Hz, 1H), 4.70 (dd, $J = 18.4, 4.4$ Hz, 1H), 3.94 (dd, $J = 53.2, 11.2$ Hz, 2H), 0.93 (s, 9H,), 0.84 (s, 9H), 0.17 (s, 3H), 0.16 (s, 3H), 0.05 (s, 3H), 0.01 (s, 3H).


^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ -194.658 (dt, $J = 53, 21.4$ Hz).

LC/MS: $t_R = 1.84$ min, MS $m/z = 626.60$ $[\text{M}+1]$; LC system: Thermo Accela 1250 UHPLC

5 MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ C18 100A, 50×3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-1.5 min 2-100% ACN, 1.5 min-2.8 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3 min 2% ACN.



10 **Intermediate 3d - (2R,3R,4S,5S)-5-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-3-(tert-butylidimethylsilyloxy)-2-((tert-butylidimethylsilyloxy)methyl)-4-fluorotetrahydrofuran-2-carbonitrile.**

To a solution of intermediate **3c** (770 mg, 1.23 mmol) in MeOH (11.2 mL), cooled in an ice water bath, was added concentrated $\text{NH}_4\text{OH}_{(\text{aq})}$ (3.74 mL). The cold bath was removed and the resulting heterogeneous solution was stirred overnight. The next day the reaction was incomplete as determined by LC/MS. Additional concentrated $\text{NH}_4\text{OH}_{(\text{aq})}$ (4 mL) and 2-MeTHF (12 mL) was added. The reaction became homogenous but after 20 minutes there was no additional reaction progress. The reaction was concentrated and the residue was dissolved in THF (15 mL). To this mixture was added concentrated $\text{NH}_4\text{OH}_{(\text{aq})}$ (5 mL) and enough MeOH (1.9 mL) to turn the solution homogenous and monophasic. The reaction was stirred at room temperature for 22 h. The reaction was nearly complete (about 5% of the starting material remains). The reaction was diluted with CH_2Cl_2 and H_2O . The layers were separated, and the aqueous layer was diluted with saturated $\text{NaHCO}_{3(\text{aq})}$ and extracted with CH_2Cl_2 . The aqueous layer was neutralized with 2N HCl and then extracted with CH_2Cl_2 . The combined organic layers were dried over Na_2SO_4 , which was removed by filtration. The filtrate was concentrated and intermediate **3d** was isolated from the residue by silica gel


column chromatography using the following solvent ramp: 0% EtOAc in hexanes ramping to 50% EtOAc in hexanes, pause at 50% EtOAc in hexanes and then ramping to 100% EtOAc in hexanes.

LC/MS: t_R = 1.59 min, MS m/z = 522.47 [M+1]; LC system: Thermo Accela 1250 UHPLC

5 MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.8 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN.



10 **Example 3 - (2R,3R,4R,5S)-5-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-4-fluoro-3-hydroxy-2-(hydroxymethyl)tetrahydrofuran-2-carbonitrile.**

To a solution of intermediate **3d** (100 mg, 0.191 mmol) in THF (2 mL), in a polypropylene tube, was added 70% HF-Pyridine in pyridine (60 μ L, 0.478 mmol) at 0 $^{\circ}$ C, under an N_2 atmosphere. The reaction was checked after 20 minutes; there was no reaction, so additional 70% HF-Pyridine in pyridine (150 μ L) was added and the cold bath was removed. After 1 hour and 50 minutes additional 70% HF-Pyridine in pyridine (150 μ L) was added. After another 2 hours additional 70% HF-Pyridine in pyridine (300 μ L) was added. After another 2 hours and 15 minutes additional 70% HF-Pyridine in pyridine (1 mL) was added. The reaction turns clear and homogenous upon this final addition of 70% HF-Pyridine in pyridine. The reaction was stirred overnight. The reaction was complete the next day. The reaction was cooled in an ice bath and quenched with H_2O and a small amount of saturated $NaHCO_{3(aq)}$. The mixture was concentrated and the residue was taken up in DMSO. Remaining insoluble material was removed by filtration and the filtrate was semipurified by HPLC. The isolated material was dissolved in DMF and purified by HPLC. Example **3** was isolated, with 0.5% as a TFA salt.

15

20

25

¹H NMR (400 MHz, DMF-*d*₇) δ 7.92 (s, 1H), 6.99 (d, *J* = 4.4 Hz, 1H), 6.89 (d, *J* = 4.9 Hz, 1H), 6.64 (d, *J* = 6 Hz, 1H), 5.92 (t, *J* = 6.4 Hz, 1H), 5.83 (dd, *J* = 25.2, 2 Hz, 1H), 5.40 (ddd, *J* = 54.8, 4.8, 2.4 Hz, 1H), 4.75 (dt, *J* = 22, 5.2 Hz, 1H), 4.02 (dd, *J* = 12, 6.4 Hz, 1H), 3.87 (dd, *J* = 12, 6.4 Hz, 1H).

5 ¹⁹F NMR (376 MHz, DMF-*d*₇) δ -74.92 (s), -193.726 (dt, *J* = 54.5, 23.3 Hz).

LC/MS: $t_R = 0.56$ min, MS $m/z = 294.10$ [M+1]; LC system: Thermo Accela 1250 UHPLC


MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

10 Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.8 min 100% ACN, 1.8 min-1.85 min 100%-
2% ACN, 1.85 min-2 min 2% ACN.

HPLC: t_R = 3.220 min; HPLC system: Agilent 1100 series.; Column: Phenomenex Kinetex C18 2.6 μ m 100A, 4.6 x 100 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA


Gradient: 0 min-8.0 min 2-98% ACN at 1.5 mL/min.



Intermediate 4b - (3aR,5R,6aS)-5-((R)-2,2-dimethyl-1,3-dioxolan-4-yl)-2,2-dimethyldihydrofuro[3,2-d][1,3]dioxol-6(3aH)-one.


Into a 10-L 4-necked round-bottom flask, was placed a solution of intermediate **4a**, (3aR,5S,6S,6aR)-5-((R)-2,2-dimethyl-1,3-dioxolan-4-yl)-2,2-dimethyltetrahydrofuro[3,2-d][1,3]dioxol-6-ol (500 g, 1.90 mol) in dichloromethane/water (2.7 L / 2.3 L) at room temperature. To this was added sodium carbonate (290 g, 3.42 mol). Addition of potassium carbonate (451 g, 3.24 mol) was next. This was followed by the addition of 2,2,6,6-tetramethylpiperidinoxy (TEMPO, 15.2 g, 96.31 mmol). To the mixture was added tetrabutylammonium bromide (31 g, 95.20 mmol). To the above was added *N*-bromosuccinimide (514 g, 2.86 mol), in portions at 35 °C. The resulting solution was allowed to react, with stirring, for 2 h at room temperature. The resulting solution was extracted with

2 x 1 L of dichloromethane and the organic layers combined. The resulting mixture was washed with 1 x 1.5 L of water. The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in (crude) intermediate **4b**.



5 Intermediate 4c - (3aR,5S,6R,6aR)-5-((R)-2,2-dimethyl-1,3-dioxolan-4-yl)-2,2-dimethyltetrahydrofuro[3,2-d][1,3]dioxol-6-ol.


Into a 2-L 4-necked round-bottom flask, was placed a solution of intermediate **4b** (370 g, 1.29 mol) in methanol (1300 mL). To the above was added sodium borohydride (26.4 g, 706.38 mmol), in portions at room temperature. The resulting solution was allowed to react, with stirring, for 2 h at room temperature. The resulting mixture was concentrated under vacuum. The reaction was then quenched by the addition of 1000 mL of 5% aqueous ammonium chloride solution. The resulting solution was extracted with 3 x 500 mL of dichloromethane and the organic layers combined. The resulting solution was washed with 2 x 300 mL of water. The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. The crude product was purified by re-crystallization from petroleum ether. This resulted in intermediate **4c**.



Intermediate 4d.


Into a 5000-mL 4-necked round-bottom flask, was placed a solution of intermediate **4c** (350 g, 1.34 mol) in dichloromethane (700 mL). To this was added tetrabutylammonium bromide (476.8 g, 1.48 mol). To the mixture was added 50% sodium hydroxide/water (700 g). To the above was added 2-(bromomethyl) naphthalene (340 g, 1.54 mol) in several batches. The resulting solution was allowed to react, with stirring, for 4 h at room temperature. The reaction was then quenched by the addition of 1800 mL of dichloromethane/water (1:1). The resulting solution was extracted with 2 x 1 L of dichloromethane and the organic layers combined. The resulting mixture was washed with 1

x 1000 mL of water. The residue was dissolved in 1000/1000 mL of petroleum ether/water. The crude product was purified by re-crystallization from petroleum ether. This resulted in intermediate **4d**.



5 Intermediate 4e - (R)-1-((3aR,5R,6R,6aR)-2,2-dimethyl-6-(naphthalen-2-ylmethoxy)tetrahydrofuro[3,2-d][1,3]dioxol-5-yl)ethane-1,2-diol.


Into a 5-L 4-necked round-bottom flask, was placed intermediate **4d** (500 g, 1.25 mol). To this was added acetic acid (1.8 L). To the mixture was added water (600 mL). The resulting solution was allowed to react, with stirring, overnight at room temperature. The solids were filtered out. The resulting solution was extracted with 3 x 1 L of petroleum ether and the aqueous layers combined. The resulting solution was diluted with 2 L of ethyl acetate. The resulting mixture was washed with 2 x 2 L of sodium chloride_(aq). The pH value of the solution was adjusted to 8 with sodium carbonate (50 %). The resulting solution was extracted with 2 x 1 L of ethyl acetate and the organic layers combined and concentrated under vacuum. This resulted in intermediate **4e**.



Intermediate 4f - (3aR,5S,6R,6aR)-2,2-dimethyl-6-(naphthalen-2-ylmethoxy)tetrahydrofuro[3,2-d][1,3]dioxole-5-carbaldehyde.


Into a 10 L 4-necked roundbottom flask, was placed a solution of intermediate **4e** (300 g, 833.33 mmol) in 1,4-dioxane (2100 mL) at room temperature. This was followed by the addition of a solution of sodium periodate (250 g) in water (4000 mL) at room temperature in 0.5 h. The resulting solution was allowed to react, with stirring, for 0.5 hr at room temperature. The solids were filtered out. The resulting solution was extracted with 3 x 1000 mL of ethyl acetate and the organic layers combined. The resulting mixture was

washed with 2 x 1000 mL of sodium chloride (aq). The resulting mixture was concentrated under vacuum. This resulted in intermediate **4f**.



Intermediate 4g - ((3aR,6S,6aR)-2,2-dimethyl-6-(naphthalen-2-ylmethoxy)tetrahydrofuro[3,2-d][1,3]dioxole-5,5-diyl)dimethanol.


Into a 10-L 4-necked round-bottom flask, was placed a solution of intermediate **4f** (250 g, 761.36 mmol) in water/tetrahydrofuran (1250/1250 mL) at room temperature. To the above was added 2N sodium hydroxide (aq) (1500 mL) dropwise with stirring at 0-15 °C. To the mixture was added formaldehyde (620 mL). The resulting solution was allowed to react, with stirring, overnight at room temperature. The resulting solution was extracted with 2 x 2000 mL of ethyl acetate. The resulting mixture was washed with 2 x 2000 mL of sodium chloride (aq). The organic layers combined and dried over anhydrous sodium sulfate. The residue was applied onto a silica gel column with petroleum ether: ethyl acetate (2/1). The crude product was re-crystallized from ethyl acetate:ethanol in the ratio of 1g/(1 mL:1 mL). This resulted in intermediate **4g**.



Intermediate 4h - ((3aR,5R,6S,6aR)-5-((tert-butyldiphenylsilyloxy)methyl)-2,2-dimethyl-6-(naphthalen-2-ylmethoxy)tetrahydrofuro[3,2-d][1,3]dioxol-5-yl)methanol.


Into a 5-L 4-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed a solution of intermediate **4g** (125 g, 346.84 mmol) in dichloromethane (2500 mL) at room temperature. To the mixture was added triethylamine (157.5 mL) at room temperature. To the above was added *tert*-butyldiphenylsilyl chloride (157.5 mL) dropwise with stirring at 0-10 °C. The resulting solution was allowed to react, with stirring, overnight at room temperature. The reaction was then quenched by the addition of

37.5 mL of methanol. The resulting mixture was washed with 2 x 500 mL of 5% hydrogen chloride_(aq) and 2 x 500 mL of sodium bicarbonate_(aq). The resulting mixture was washed with 2 x 500 mL of 1N sodium hydroxide_(aq). The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. The crude product was purified by re-crystallization from dichloromethane/hexane. This resulted in intermediate **4h**.




Intermediate 4i - tert-butyl(((3aR,5R,6S,6aR)-5-(iodomethyl)-2,2-dimethyl-6-(naphthalen-2-ylmethoxy)tetrahydrofuro[3,2-d][1,3]dioxol-5-yl)methoxy)diphenylsilane.

Into a 1000-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed a solution of intermediate **4h** (20 g, 31.73 mmol) in toluene (320 mL), triphenylphosphine (35 g, 132.11 mmol), imidazole (8.96 g, 132.26 mmol). This was followed by the addition of iodine (16.95 g, 66.8 mmol) in several batches at 60 °C. The resulting solution was stirred overnight at 80 °C in an oil bath. The reaction mixture was cooled to room temperature with a water/ice bath. The resulting solution was diluted with 1000 mL of ethyl acetate. The resulting mixture was washed with 2 x 300 mL of sodium thiosulfate_(aq). The resulting mixture was washed with 1 x 300 mL of sodium chloride_(aq). The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:10). This resulted in intermediate **4i**.




Intermediate 4j - tert-butylidiphenyl(((3aR,5R,6S,6aR)-2,2,5-trimethyl-6-(naphthalen-2-ylmethoxy)tetrahydrofuro[3,2-d][1,3]dioxol-5-yl)methoxy)silane.

Into a 2000-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed a solution of intermediate **4i** (66 g, 88.47 mmol) in ethanol/ethyl acetate (600/600 mL), triethylamine (20.7 g, 202.52 mmol), palladium on carbon (10% wt, 24.8 g, 23.30 mmol). The resulting solution was stirred for 3 h at 40 °C. The solids were
 5 filtered out. The resulting mixture was concentrated under vacuum. The resulting solution was diluted with 1500 mL of ethyl acetate. The resulting mixture was washed with 1 x 500 mL of sodium chloride_(aq). The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in intermediate **4j**.



10 **Intermediate 4k - (3aR,5R,6S,6aR)-5-((tert-butyldiphenylsilyloxy)methyl)-2,2,5-trimethyltetrahydrofuro[3,2-d][1,3]dioxol-6-ol.**

Into a 500-mL round-bottom flask, was placed a solution of intermediate **4j** (1.0 g, 1.63 mmol) in dichloromethane (15 mL), water (1.25 mL), and 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ, 780 mg, 3.40 mmol). The resulting solution was stirred for 1 h at room
 15 temperature. The resulting solution was diluted with 50 mL of dichloromethane. The resulting mixture was washed with 1 x 30 mL of water and 2 x 30 mL of sodium bicarbonate_(aq). The resulting mixture was washed with 1 x 30 mL of sodium chloride_(aq). The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:20~1:10). This resulted in
 20 intermediate **4k**.




Intermediate 4l - (3aR,5R,6S,6aR)-5-(hydroxymethyl)-2,2,5-trimethyltetrahydrofuro[3,2-d][1,3]dioxol-6-ol.

Into a 50-mL round-bottom flask, was placed a solution of intermediate **4k** (520 mg, 1.12 mmol) in tetrahydrofuran (9 mL), Tetrabutylammonium fluoride (369 mg, 1.40 mmol).
 25


The resulting solution was stirred for 2 h at room temperature. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with dichloromethane/methanol (100:1). This resulted in intermediate **4l**.

¹H NMR (300 MHz, DMSO-*d*₆): δ 5.64 (d, *J* = 3.9 Hz, 1H), 4.96 (d, *J* = 6.6 Hz, 1H), 4.67 (m, 1H), 4.55 (m, 1H), 4.05 (m, 1H), 3.24 – 3.30 (m, 1H), 3.11 – 3.18 (m, 1H), 1.50 (s, 3H), 1.27 (s, 3H), 1.16 (s, 1H).



Intermediate 4m - (3aR,5R,6S,6aR)-6-(benzyloxy)-5-(benzyloxymethyl)-2,2,5-trimethyltetrahydrofuro[3,2-d][1,3]dioxole.

Into a 50-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed a solution of intermediate **4l** (180 mg, 0.84 mmol) in tetrahydrofuran (4 mL). This was followed by the addition of sodium hydride (60% wt., 140 mg, 3.50 mmol), in portions at 0 °C. The resulting solution was stirred for 30 min at 0 °C. The resulting solution was allowed to react, with stirring, for an additional 30 min at room temperature. To this was added benzyl bromide (452 mg, 2.62 mmol) dropwise with stirring at 0 °C. The resulting solution was allowed to react, with stirring, for an additional 3 h at room temperature. The reaction was then quenched by the addition of 30 mL of ammonium chloride_(aq). The resulting solution was extracted with 50 mL of dichloromethane and the organic layers were combined and dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:30-1:20). This resulted in intermediate **4m**.




Intermediate 4n - (2R,3R,4S,5R)-4-(benzyloxy)-5-(benzyloxymethyl)-5-methyltetrahydrofuran-2,3-diyl diacetate.

Into a 1000-mL round-bottom flask, was placed intermediate **4m** (Also prepared according to *Biosci. Biotech. Biochem.* **1993**, 57, 1433-1438, 45 g, 111.19 mmol), acetic acid (270 mL), acetic anhydride (90 mL), sulfuric acid (45 d). The resulting solution was stirred for 30 min at room temperature. The reaction was then quenched by the addition of 1000 mL of water/ice. The resulting solution was diluted with 3000 mL of ethyl acetate. The resulting mixture was washed with 2 x 1000 mL of water and 4 x 1000 mL of sodium bicarbonate_(aq). The resulting mixture was washed with 2 x 1000 mL of sodium chloride_(aq). The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:30-1:20). This resulted in intermediate **4n**.

¹H NMR (300 MHz, CDCl₃): δ 7.28 – 7.38 (m, 10H), 6.13 (s, 1H), 5.37 (d, *J* = 4.8 Hz, 1H), 4.44 – 4.68 (m, 4H), 4.33 (d, *J* = 5.1 Hz, 1H), 3.33 – 3.45 (m, 2H), 2.15 (s, 3H), 1.88 (s, 3H), 1.35 (s, 3H).

MS *m/z* = 451[M+Na]



Intermediate 4o - (3R,4S,5R)-4-(benzyloxy)-5-(benzyloxymethyl)-3-hydroxy-5-methyldihydrofuran-2(3H)-one.


Intermediate **4n** (1.3 g, 3 mmol) was dissolved in anhydrous MeOH (15 mL). Potassium carbonate powder (456 mg, 3.3 mmol) was added, and the reaction mixture was stirred for 1 h. The reaction mixture was then concentrated under reduced pressure. Added acetonitrile and stirred for 5 minutes. Filtered off insoluble and washed with acetonitrile. Concentrated filtrate under reduced pressure. Dissolved resulting material in anhydrous DCM (20 mL). Added tetrabutylammonium iodine (1.66 g, 4.5 mmol) and *N*-iodo-succinimide (NIS, 1.69 g, 2.5 mmol). Stirred the reaction mixture in the dark for 16 h. Added more NIS (0.85 g, 1.25 mmol) and stirred for 4 hours. Added more NIS (0.85 g, 1.25 mmol) and stirred for 2 days in the dark. Diluted reaction with EtOAc and washed with aqueous sodium thiosulfate solution twice and then with saturated aqueous sodium chloride solution. Dried organic portion over anhydrous sodium sulfate and concentrated under reduced pressure. Purified with silica gel column (0-30% EtOAc in hexanes) to afford intermediate **4o**.

^1H NMR (400 MHz, CDCl_3): δ 7.35 – 7.22 (m, 10H), 4.82 (bs, 1H), 4.75 – 4.66 (m, 2H), 4.55 – 4.44 (m, 2H), 4.13 (d, J = 8Hz, 1H), 3.70-3.45 (m, 2H), 1.38 (s, 3H).

LC/MS: t_R = 2.58 min, MS m/z = 342.9 $[\text{M}+1]$, 360.0 $[\text{M}+\text{H}_2\text{O}]$; LC/MS system: Thermo LCQ Advantage; Phenomenex Gemini, C_{18} , 5u, 110A, 30 x 4.6 mm; Buffer A: 0.1% Acetic acid in Water; Buffer B: 0.1% Acetic acid in Acetonitrile

5-100% Buffer B in 2.5 mins then 100% for 0.9min @ 2 mL/min.

HPLC: t_R = 3.78 min; HPLC system: Agilent 1100; Phenomenex Gemini, C_{18} , 5u, 110A, 50 x 4.6 mm; Buffer A: 0.05% TFA in Water; Buffer B: 0.05% TFA in Acetonitrile; 2-98% Buffer B in 5 minutes @ 2mL/min.



Intermediate 4p - (3R,4S,5R)-3,4-bis(benzyloxy)-5-(benzyloxymethyl)-5-methyldihydrofuran-2(3H)-one.


Dissolved intermediate **4o** (955 mg, 2.79 mmol) in EtOAc (10 mL). Added benzyl bromide (400 μL , 3.35 mmol) and silver(I) oxide (712 mg, 3.07 mmol). Stirred at 60 $^\circ\text{C}$ under $\text{N}_2(\text{g})$ in the dark for 3 h. Added more benzyl bromide (400 μL , 3.35 mmol) and stirred at 60 $^\circ\text{C}$ under $\text{N}_2(\text{g})$ in the dark for 16 h. Added more silver(I) oxide (350 mg, 1.5 mmol) and stirred at 60 $^\circ\text{C}$ under $\text{N}_2(\text{g})$ in the dark for 8 h. Cooled to room temperature. Filtered off solids and washed with EtOAc. Concentrated filtrate under reduced pressure to give an oil. Added hexanes and stirred for 2 h to give solid. Collected solid and washed with hexanes. Dried solid under high vacuum to afford intermediate **4p**.

^1H NMR (400 MHz, CDCl_3): δ 7.35 – 7.16 (m, 15H), 5.03 (d, J = 12Hz, 1H), 4.79 – 4.71 (m, 2H), 4.52 – 4.40 (m, 4H), 4.06 (d, J = 6Hz, 1H), 3.49 – 3.39 (m, 2H), 1.38 (s, 3H).

LC/MS: t_R = 2.91 min, MS m/z = 433.1 $[\text{M}+1]$, 450.1 $[\text{M}+\text{H}_2\text{O}]$; LC/MS system: Thermo LCQ Advantage; Phenomenex Gemini, C_{18} , 5u, 110A, 30 x 4.6 mm; Buffer A: 0.1% Acetic acid in Water; Buffer B: 0.1% Acetic acid in Acetonitrile; 5-100% Buffer B in 2.5 mins then 100% for 0.9min @ 2 mL/min.

HPLC: t_R = 4.54 min; HPLC system: Agilent 1100; Phenomenex Gemini, C₁₈, 5u, 110A, 50 x 4.6 mm; Buffer A: 0.05% TFA in Water; Buffer B: 0.05% TFA in Acetonitrile

2-98% Buffer B in 5 minutes @ 2mL/min.




5 Intermediate 4q - (3R,4S,5R)-2-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-3,4-bis(benzyloxy)-5-(benzyloxymethyl)-5-methyltetrahydrofuran-2-ol.

Intermediate **1b** (148 mg, 0.570 mmol) and 1,2-bis(chlorodimethylsilyl)ethane (123 mg, 0.570 mmol) were dissolved in anhydrous THF (20 mL) and stirred under Ar(g) at -70 °C. Added *n*-butyllithium (2.5M solution in hexanes, 684 μ L, 1.71 mmol) dropwise to the
 10 reaction mixture while maintaining internal temperature below -65 °C. Allowed the reaction to warm to -40 °C and kept for 15 min. A solution of intermediate **4p** (224 mg, 0.518 mmol) in THF (10 mL) precooled to -70 °C was then added to the reaction mixture under Ar(g). The resulting solution was stirred for 2 h at -40 °C. The reaction mixture was then poured into a stirring mixture of EtOAc and citric acid_(aq). Stirred for 5 min. Collected organic layer and
 15 washed with saturated NaCl_(aq). Dried organic layer over anhydrous Na₂SO₄ and concentrated under reduced pressure. Purified with prep HPLC to afford intermediate **4q**.

LC/MS: t_R = 2.60 min, MS m/z = 567.3 [M+1], 565.1 [M-1]; LC/MS system: Thermo LCQ Advantage; Phenomenex Gemini, C₁₈, 5u, 110A, 30 x 4.6 mm; Buffer A: 0.1% Acetic acid in Water; Buffer B: 0.1% Acetic acid in Acetonitrile; 5-100% Buffer B in 2.5 mins then 100% for
 20 0.9min @ 2 mL/min.

HPLC: t_R = 3.22 min; HPLC system: Agilent 1100; Phenomenex Gemini, C₁₈, 5u, 110A, 50 x 4.6 mm; Buffer A: 0.05% TFA in Water; Buffer B: 0.05% TFA in Acetonitrile; 2-98% Buffer B in 5 minutes @ 2mL/min.



Intermediate 4r - 7-((2S,3S,4S,5R)-3,4-bis(benzyloxy)-5-(benzyloxymethyl)-5-methyltetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-amine.


Intermediate **4q** (81 mg, 0.143 mmol) was dissolved in anhydrous DCM (15 mL) and stirred under $\text{N}_2(\text{g})$ in an ice bath. Added triethylsilane (114 μL , 0.715 mmol) in one portion. Added boron trifluoride diethyl etherate (27 μL , 0.215 mmol) dropwise. Stirred for 15 min and then removed ice bath. Stirred for 60 min. Added triethylamine (100 μL , 0.715 mmol) and concentrated under reduced pressure. Dissolved in EtOAc and washed with saturated $\text{NaHCO}_{3(\text{aq})}$ (2x) and then saturated with $\text{NaCl}_{(\text{aq})}$. Dried organic over anhydrous Na_2SO_4 and concentrated under reduced pressure. Purified with silica gel column (0-80% EtOAc in hexanes) to afford intermediate **4r**.

^1H NMR (400 MHz, CDCl_3): δ 7.71 (s, 1H), 7.35 – 7.10 (m, 16H), 6.82-6.78 (m, 1H), 5.57 (d, $J=4.4\text{Hz}$, 1H), 4.70 – 4.45 (m, 6H), 4.25 – 4.15 (m, 2H), 3.55 – 3.40 (m, 2H), 1.42 (s, 3H).

LC/MS: $t_R = 2.75\text{min}$, MS $m/z = 551.4$ $[\text{M}+1]$; LC/MS system: Thermo LCQ Advantage

Phenomenex Gemini, C_{18} , 5u, 110A, 30 x 4.6 mm; Buffer A: 0.1% Acetic acid in Water; Buffer B: 0.1% Acetic acid in Acetonitrile; 5-100% Buffer B in 2.5 mins then 100% for 0.9min @ 2 mL/min.

HPLC: $t_R = 3.57\text{min}$; HPLC system: Agilent 1100; Phenomenex Gemini, C_{18} , 5u, 110A, 50 x 4.6 mm; Buffer A: 0.05% TFA in Water; Buffer B: 0.05% TFA in Acetonitrile; 2-98% Buffer B in 5 minutes @ 2mL/min.




f[1,2,4]triazin-4-yl)benzamide, (~110 mg, ~0.18 mmol) and azidotrimethylsilane (242 μ l, 1.84 mmol) in dichloromethane (1.5 mL) was added indium (III) bromide (130 mg, 0.369 mmol) at RT under an argon atmosphere. After 1 h, the reaction mixture was quenched with saturated aqueous sodium bicarbonate solution (1 mL). The resulting mixture was partitioned between
 5 dichloromethane (20 mL) and saturated aqueous sodium bicarbonate solution (20 mL). The phases were split and the aqueous layer was extracted with dichloromethane (20 mL). The combined organic layers were dried over anhydrous sodium sulfate and were concentrated under reduced pressure to afford intermediate **5a** that was used directly in the next step without further purification.

10 LC/MS: t_R = 3.52 min, MS m/z = 712.16 [M+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm

Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-2.0 min 2-100% ACN, 2.0 min-3.55 min 100% ACN, 3.55 min-4.2 min 100%-2% ACN at 2 μ l/min.




Intermediate 5b - N-(7-((2S,3R,4S,5R)-5-azido-3,4-dihydroxy-5-(hydroxymethyl)tetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide.

To a solution of crude intermediate **5a** (~120 mg, ~0.168 mmol) in DMF (5 mL) was added cesium fluoride (256 mg, 1.68 mmol) at RT. After 25 h, the reaction mixture was
 20 diluted with brine (100 mL), and the resulting mixture was extracted with ethyl acetate (3 x 100 mL). The combined organic layers were dried over anhydrous sodium sulfate and were concentrated under reduced pressure to afford intermediate **5b** that was used directly in the next step without further purification.

LC/MS: t_R = 1.40 min, MS m/z = 412.17 [M+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm. Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-2.0 min 2-100% ACN, 2.0 min-3.05 min 100% ACN, 3.05 min-3.2 min 100%-2% ACN, 3.2 min-3.5 min 2% ACN at 2 μ l/min.

5 HPLC: t_R = 2.46 min' HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.



Example 5 - (2R,3S,4R,5S)-5-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-2-azido-2-(hydroxymethyl)tetrahydrofuran-3,4-diol.

Concentrated ammonium hydroxide (1 mL) was added to a solution of crude intermediate **5b** in methanol (1 mL) at RT. After 2 d, the reaction mixture was concentrated under reduced pressure, and was directly purified by preparatory HPLC (Phenomenex Synergi 4u Hydro-RR 80Å 150 x 30 mm column, 5-100% acetonitrile/water gradient). The fractions containing the desired were combined and were concentrated under reduced pressure. The residue repurified via SiO₂ column chromatography (4 g SiO₂ Combiflash HP Gold Column, 0–20% methanol/dichloromethane) to afford example **5**.


¹H NMR (400 MHz, CD₃OD) δ 7.79 (s, 1H), 6.86 (d, J = 4.5 Hz, 1H), 6.77 (d, J = 4.5 Hz, 1H), 5.51 (d, J = 6.0 Hz, 1H), 4.63 (t, J = 5.8 Hz, 1H), 4.37 (d, J = 5.7 Hz, 1H), 3.69 (d, J = 12.0 Hz, 1H), 3.59 (d, J = 12.0 Hz, 1H).

LC/MS: t_R = 0.76 min, MS m/z = 308.08 [M+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-1.5 min 2-100% ACN, 1.5 min-2.2 min 100% ACN, 2.2 min-2.4 min 100%-2% ACN, 2.4 min-2.5 min 2% ACN at 2 μ l/min.

HPLC: t_R = 1.287 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.;

TLC: eluent: 20% methanol in dichloromethane, R_f = 0.4 (UV)



Example 6 (also referred to as TP-1) - ((2R,3R,4R,5S)-5-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-2-cyano-4-fluoro-3-hydroxytetrahydrofuran-2-yl)methyl tetrahydrogen triphosphate.


Example 3 (15.0 mg, 0.05 mmol) was dried in a flask under vacuum overnight.

Trimethylphosphate (0.5 mL) and 1,8-bis(dimethylamino)naphthalene (25 mg, 0.12 mmol) were added to the flask and the solution was allowed to stir under N_2 cooled by an ice/water bath. Distilled phosphorus oxychloride (10 μ L, 0.11 mmol) was added and the reaction was allowed to stir for 4 h with cooling. Tributylamine (0.1 mL, 0.42 mmol) and tributylammonium pyrophosphate (0.8 mL of a 0.5 M solution in DMF, 0.4 mmol) were added and the reaction was allowed to stir for an additional 45 min with cooling. The reaction was quenched with triethylammonium bicarbonate (0.5 M, 5 mL). The solvents were removed by rotary evaporation and the remaining crude mixture was dissolved in 2 mL of water. The product was purified using a Sephadex DEAE A-25 column with a linear gradient of 0–1 M triethylammonium bicarbonate. The product containing fractions were pooled and concentrated to give example 6 (TP1), which was then dissolved in 1 mL of water to give a 10 mM solution.

MS m/z = 532.0 [M-1]

Ion Exchange HPLC Retention time: 12.015 min; Column: DNAPac PA-100 4 x 250 mm SN

Solvent A: milliQ water; Solvent B: 0.5 M tetraethylammonium bromide; Solvent gradient program: equilibrate using 100%A for 10 min, then ramp 0-80% B over 14min.; Flow: 1 mL/min.




5 **Example 7 (also TP2) - ((2R,3R,4R,5S)-5-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-2-ethynyl-4-fluoro-3-hydroxytetrahydrofuran-2-yl)methyl tetrahydrogen triphosphate.**

Example 2 (16.0 mg, 0.055 mmol) was dried in a flask under vacuum overnight. Trimethylphosphate (0.5 mL) and 1,8-bis(dimethylamino)naphthalene (28 mg, 0.13 mmol) were added to the flask and the solution was allowed to stir under N₂ cooled by an ice/water bath. Distilled phosphorus oxychloride (11 µL, 0.12 mmol) was added and the reaction was
 10 allowed to stir for 4 h with cooling. Tributylamine (0.11 mL, 0.42 mmol) and tributylammonium pyrophosphate (0.9 mL of a 0.5 M solution in DMF, 0.45 mmol) were added and the reaction was allowed to stir for an additional 45 min with cooling. The reaction was quenched with triethylammonium bicarbonate (0.5 M, 5 mL). The solvents were
 15 removed by rotary evaporation and the remaining crude mixture was dissolved in 2 mL of water. The product was purified using a Sephadex DEAE A-25 column with a linear gradient of 0–1 M triethylammonium bicarbonate. The product containing fractions were pooled and concentrated to give example 7 (TP2), which was then dissolved in 1.4 mL of water to give a 10 mM solution.

20 MS *m/z* = 531.0 [M-1]

Ion Exchange HPLC Retention time: 19.829 min; Column: DNAPac PA-100 4 x 250 mm SN

Solvent A: milliQ water; Solvent B: 0.5 M tetraethylammonium bromide; Solvent gradient program: equilibrate using 100%A for 10 min, then ramp 0-80% B over 14 min.; Flow: 1 mL/min




Example 8 (TP3) - ((2R,3S,4R,5S)-5-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-2-cyano-3,4-dihydroxytetrahydrofuran-2-yl)methyl tetrahydrogen triphosphate.

To a solution of example 1 (5.0 mg, 0.017 mmol) in PO(OMe)₃ (0.6 mL) at 0 °C was added POCl₃ (45 mg, 0.29 mmol). The reaction mixture was stirred at 0 °C for 10 h, at which point ion-exchange HPLC showed approximately 50% conversion. A solution of pyrophosphate tributylamine salts (250 mg) in ACN (0.6 mL) was added, followed by tributylamine (110 mg, 0.59 mmol). The reaction mixture was stirred at 0 °C for 0.5 h, and ion-exchange HPLC showed the reaction was completed. The reaction was quenched with triethylammonium bicarbonate buffer (1 M, 5 mL). The reaction mixture was stirred at RT for 0.5 h, then concentrated and co-evaporated with water twice. The residue was dissolved in H₂O (5 mL) and loaded to a ion-exchange column, eluted with H₂O, then 5-35% triethylammonium bicarbonate buffer (1M)-H₂O. The product fractions were combined, concentrated and co-evaporated with H₂O. The residue was purified by ion-exchange column again to give crude material. ³¹P NMR showed this material contained impurities, so the material was repurified with C-18 column, eluted with 0-15% ACN-H₂O containing 0.05% TEA, and the fractions containing product were combined and concentrated to give 3.6 mg material, which contained only 1.5 equiv of TEA as indicated by ¹H NMR analysis. The material was dissolved in H₂O (1 mL) and triethylammonium bicarbonate buffer (1 M, 0.1 mL) was added. The resulting mixture was concentrated under reduced pressure and co-evaporated with H₂O twice under reduced pressure to give example 8 (TP3), as a tetra-TEA salt.

¹H NMR (400 MHz, D₂O): δ 7.78 (s, 1H), 6.85 (d, *J* = 2.4 Hz, 1H), 6.82 (d, *J* = 2.4 Hz, 1H), 5.51 (d, *J* = 3.0 Hz, 1H), 4.65 – 4.55 (m, 2H), 4.20 – 4.08 (m, 2H), 3.15 – 3.00 (m, 24H), 1.18 – 1.08 (m, 36H).

³¹P NMR (162 MHz, D₂O): δ -6.25 (d, *J* = 52 Hz), -12.21 (d, *J* = 52 Hz), -22.32 (t, *J* = 52 Hz).

MS m/z = 530.2 [M-1], 532.1 [M+1]




Example 9 (TP4) - ((2R,3S,4R,5S)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-azido-3,4-dihydroxytetrahydrofuran-2-yl)methyl tetrahydrogen triphosphate.

5 To a solution of example 5 (6.0 mg, 0.019 mmol) in PO(OMe)₃ (0.6 mL) at 0 °C was added POCl₃ (45 mg, 0.29 mmol). The reaction mixture was stirred at 0 °C for 10 h, at which point ion-exchange HPLC showed approximately 50% conversion. A solution of pyrophosphate tributylamine salts (250 mg) in ACN (0.6 mL) was added, followed by tributylamine (110 mg, 0.59 mmol). The reaction mixture was stirred at 0 °C for 6 h. The reaction was quenched with triethylammonium bicarbonate buffer (1 M, 5 mL). The reaction mixture was stirred at RT for 0.5 h, then concentrated and co-evaporated with water twice. The residue was dissolved in H₂O (5 mL) and loaded to a ion-exchange column, eluted with H₂O, then 5-35% triethylammonium bicarbonate buffer (1M)-H₂O. The product fractions were combined, concentrated and co-evaporated with H₂O. The residue was purified by ion-exchange column again to give crude material. ³¹P NMR showed this material contained impurities, so the material was repurified with ion-exchange column again to give crude material. The material was treated with NaHCO₃ (10 mg) and the mixture was concentrated under reduced pressure. The solid residue was dissolved in 0.5 mL of H₂O and 40 μL of NaOH (1N) was added. The resulting mixture was purified with C-18 column, eluted with H₂O, and the fractions containing product were combined and concentrated under reduced pressure to afford Example 9 (TP4) as the tetra-sodium salt.

¹H NMR (400 MHz, D₂O): δ 7.76 (s, 1H), 6.88 (d, J = 4.3 Hz, 1H), 6.81 (d, J = 4.6 Hz, 1H), 5.59 (d, J = 5.5 Hz, 1H), 4.60 (t, J = 5.6 Hz, 1H), 4.55 (d, J = 5.8 Hz, 1H), 3.99 (qd, J = 11.2, 5.5 Hz, 3H).

25 ³¹P NMR (162 MHz, D₂O): δ -8.13 (d, J = 19.8 Hz), -14.04 (d, J = 18.9 Hz), -24.00 (t, J = 19.3 Hz).


MS m/z = 546.1 [M-1], 547.9 [M+1]



Intermediate PD1a - S-2-hydroxyethyl 2,2-dimethylpropanethioate.

5 To a solution of 2-thioethanol (3.50 mL, 50.0 mmol) and triethylamine (7.02 mL, 50.0 mmol) in CH_2Cl_2 , that has been cooled to -78°C was added pivalyl chloride (6.15 mL, 50.0 mmol) dropwise over 30 min. The reaction was allowed to warm slowly to room temperature and progression was monitored by TLC. After 30 min, the reaction was determined to be complete and was quenched with water. The layers were separated and the aqueous was
10 washed with CH_2Cl_2 . The organics were combined and dried over sodium sulfate. The solids were filtered and the solvent was removed under reduced pressure. The crude was purified by silica gel chromatography 0- 50% EtOAc/hexanes to afford intermediate **PD1a**.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 4.89 (t, J = 5.5 Hz, 1H), 3.49 – 3.36 (m, 2H), 2.86 (t, J = 6.7 Hz, 2H), 1.14 (s, 9H).




15 Take up phosphorous oxychloride (281 μL , 3.08 mmol) in CH_2Cl_2 (5 mL) and cool the solution to -78°C . Take up the thioester **PD1a** (1.00 g, 6.17 mmol) in CH_2Cl_2 (5 mL) and add slowly to the POCl_3 solution. Next add TEA (891 μL , 6.16 mmol) dropwise and allow to stir cold for 30 min. Then warm to room temperature and allow to stir for 2 h. Add the *p*-
20 nitrophenol (428 mg, 3.08 mmol) in one portion followed by a slow addition of TEA (449 μL , 3.08 mmol). Stir at room temperature for 30 min. TLC (70:30 Hexanes/EtOAc) showed only one spot, but LC/MS had two peaks (product and bis-*p*-nitrophenolate). The solution was diluted with ether and the solids were removed by filtration and discarded. The mother liquor was concentrated and purified by silica gel chromatography to give a mixture of product and

bis-*p*-nitrophenolate. The mixture was then repurified by HPLC to afford intermediate **PD1b**, *S,S'*-2,2'-((4-nitrophenoxy)phosphoryl)bis(oxy)bis(ethane-2,1-diyl) bis(2,2-dimethylpropanethioate).

¹H NMR (400 MHz, CDCl₃) δ 8.29 – 8.21 (m, 2H), 7.46 – 7.36 (m, 2H), 4.23 (br q, *J* = 7.7 Hz, 4H), 3.16 (br t, *J* = 6.7 Hz, 4H), 1.23 (s, 18H).

³¹P NMR (162 MHz, CDCl₃) δ -7.72 (s).



Example 10 (also referred to as PD1) - *S,S'*-2,2'-((((2*R*,3*S*,4*R*,5*S*)-5-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-2-cyano-3,4-dihydroxytetrahydrofuran-2-yl)methoxy)phosphoryl)bis (oxy)bis(ethane-2,1-diyl) bis(2,2-dimethylpropanethioate).

Example 1 (6.0 mg, 0.02 mmol) was dissolved in NMP (0.1 mL), and THF (0.2 mL) was added. *tert*-Butyl magnesium chloride (1.0 M solution in THF, 0.031 mL, 0.031 mmol) was then added at RT under an argon atmosphere. After 10 min, a solution of intermediate **PD1b** (15.7 mg, 0.031 mmol) in THF (0.1 mL) was added and the resulting mixture was warmed to 50 °C. After 5 h, the resulting residue was purified by preparatory HPLC (Phenominex Synergi 4u Hydro-RR 80Å 150 x 30 mm column, 40-100% acetonitrile/water gradient). The fractions containing the desired product were combined and were lyophilized to afford **example 10 (PD1)**.

¹H NMR (400 MHz, CDCl₃) δ 7.82 (s, 1H), 6.69 (d, *J* = 4.5 Hz, 1H), 6.64 (d, *J* = 4.5 Hz, 1H), 5.56 (d, *J* = 3.4 Hz, 1H), 4.61 (br s, 2H), 4.45 – 4.32 (m, 2H), 4.22 – 4.06 (m, 4H), 3.13 (dt, *J* = 11.7, 6.7 Hz, 4H), 1.23 (s, 9H), 1.21 (s, 9H).

³¹P NMR (162 MHz, CDCl₃) δ -2.34 (s).


LC/MS: *t*_R = 1.70 min, MS *m/z* = 660.02 [*M*+1],; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ XB-C18 100A, 50 x 4.6 mm

Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-2.0 min 2-100% ACN, 2.0 min-3.05 min 100% ACN, 3.05 min-3.2 min 100%-2% ACN, 3.2 min-3.5 min 2% ACN at 2 μ l/min.

HPLC: t_R = 3.204 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA

Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.



Example 11 (PD2) - S,S'-2,2'-(((2R,3R,4R,5S)-5-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-2-cyano-4-fluoro-3-hydroxytetrahydrofuran-2-yl)methoxy)phosphoryl)bis(oxy)bis(ethane-2,1-diyl) bis(2,2-dimethylpropanethioate).

Example 3 (10.5 mg, 0.036 mmol) was dissolved in NMP (0.1 mL), and THF (0.1 mL) was added. *tert*-Butyl magnesium chloride (1.0 M solution in THF, 0.054 mL, 0.054 mmol) was then added at RT under an argon atmosphere. After 10 min, a solution of intermediate **PD1b** (27.3 mg, 0.054 mmol) in THF (0.1 mL) was added and the resulting mixture was warmed to 50 °C. After 24 h, the resulting residue was purified by preparatory HPLC (Phenominex Synergi 4u Hydro-RR 80Å 150 x 30 mm column, 40-100% acetonitrile/water gradient). The fractions containing the desired product were combined and were lyophilized to afford example 11 (**PD2**).

^1H NMR (400 MHz, CDCl_3) δ 7.94 (s, 1H), 6.75 (d, J = 4.5 Hz, 1H), 6.67 (d, J = 4.5 Hz, 1H), 5.77 (dd, J = 27.8, 1.4 Hz, 1H), 5.43 (ddd, J = 55.2, 4.9, 1.3 Hz, 1H), 4.93 (dd, J = 21.2, 4.9 Hz, 1H), 4.49 (dd, J = 11.3, 7.8 Hz, 1H), 4.40 (dd, J = 11.3, 7.8 Hz, 1H), 4.10 (ddt, J = 15.9, 8.0, 6.7 Hz, 4H), 3.16 – 3.04 (m, 4H), 1.23 (s, 9H), 1.21 (s, 9H).


^{31}P NMR (162 MHz, CDCl_3) δ -2.10 (s).

^{19}F NMR (376 MHz, CDCl_3) δ -191.64 (ddd, J = 55.0, 27.8, 21.3 Hz).

LC/MS: t_R = 1.85 min, MS m/z = 662.03 [$M+1$]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-2.0 min 2-100% ACN, 2.0 min-3.05 min 100% ACN, 3.05 min-3.2 min 100%-2% ACN, 3.2 min-3.5 min 2% ACN at 2 μ L/min.

- 5 HPLC: t_R = 3.385 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.



Intermediate PD3c - (2S)-2-ethylbutyl 2-((4-nitrophenoxy)(phenoxy)phosphorylamino)propanoate.


- 10 Dissolved phenyl dichlorophosphate PD3a (1.5 mL, 10 mmol) in 30 mL anhydrous DCM and stirred under N₂(g) in an ice bath. Added amino ester HCl salt PD3b, (S)-2-ethylbutyl 2-aminopropanoate hydrochloride, (prepared according to *Eur. J. Med. Chem.* **2009**,44, 3765-3770, 2.1 g, 10 mmol) in one portion. Added TEA (3 mL, 22 mmol) dropwise.
- 15 Stirred for 1 h at 0 °C. Added p-nitrophenol (1.4 g, 10 mmol) in one portion and TEA (1.5 mL, 11 mmol). The reaction mixture was then stirred at room temperature for 16 h. Diluted with DCM and washed with saturated NaHCO_{3(aq)}. Dried organic over anhydrous Na₂SO₄ and concentrated under reduced pressure. Purified with silica gel column (0-15% EtOAc in hexanes) to afford intermediate PD3c.

- 20 ¹H NMR (400 MHz, CDCl₃) δ 8.23 (d, J = 8.8 Hz, 2H), 7.41 – 7.30 (m, 4H), 7.25 – 7.19 (m, 3H), 4.10 – 4.00 (m, 3H), 3.90-3.83 (m, 1H), 1.55 – 1.45 (m, 1H), 1.42 – 1.31 (m, 7H), 0.87 (t, J = 7.2Hz, 6H).

³¹P NMR (162 MHz, CDCl₃) δ -3.04 (s), -3.10 (s).

- LC/MS: t_R = 2.87min, MS m/z = 451.1 [M+1], 449.0 [M-1]; LC/MS system: Thermo LCQ Advantage; Phenomenex Gemini, C₁₈, 5 μ , 110A, 30 x 4.6 mm; Buffer A: 0.1% Acetic acid in Water; Buffer B: 0.1% Acetic acid in Acetonitrile; 5-100% Buffer B in 2.5 mins then 100% for 0.9min @ 2 mL/min.

HPLC: t_R = 4.40min; HPLC system: Agilent 1100; Phenomenex Gemini, C₁₈, 5u, 110A, 50 x 4.6 mm; Buffer A: 0.05% TFA in Water; Buffer B: 0.05% TFA in Acetonitrile; 2-98% Buffer B in 5 minutes @ 2mL/min.



5 **Example 12 (PD3) - (2S)-2-ethylbutyl 2-(((2R,3R,4R,5S)-5-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-2-cyano-4-fluoro-3-hydroxytetrahydrofuran-2-yl)methoxy)(phenoxy)phosphorylamino)propanoate.**

Dissolved example 3 (15 mg, 0.051 mmol) in anhydrous DMF (1 mL) and stirred under N₂(g). Dissolved *p*-nitrophenylphosphoamidate **PD3c** (35 mg, 0.077 mmol) in anhydrous DMF (0.5 mL) and added to the reaction mixture in one portion. Added *t*BuMgCl in THF (1M in THF, 77 μ L, 0.077 mmol) dropwise. Stirred for 2 h. Added more *p*-nitrophenylphosphoamidate (35 mg in 0.5 mL anhydrous DMF) and more *t*BuMgCl (1M in THF, 50 μ L, 0.050 mmol). Stirred for 2 h. Added more *p*-nitrophenylphosphoamidate (35 mg in 0.5 mL anhydrous DMF) and more *t*BuMgCl solution (1M in THF, 50 μ L, 0.050 mmol). Stirred for 16 hours. Diluted with EtOAc and washed with saturated NaHCO_{3(aq)} (3x). Washed with saturated NaCl_(aq) and dried organic over anhydrous Na₂SO₄. Concentrated under reduced pressure. Purified with silica gel column (0-5% MeOH in DCM). Combined fractions and concentrated under reduced pressure. Purified with preparatory HPLC with TFA as modifier to afford example 12 (**PD3**).

20 Prep HPLC system: Gilson 215 Liquid Handler; Phenomenex Gemini, C₁₈ 4u, 100 x 30.0 mm

Buffer A: 0.1% TFA in Water ; Buffer B: 0.1% TFA in Acetonitrile; 5-100% Buffer B in 13 minutes @ 20 mL/min.


¹H NMR (400 MHz, CDCl₃) δ 7.89 (s, 1H), 7.31 – 7.13 (m, 6H), 6.80-6.75 (m, 1H), 5.80 – 5.70 (m, 1H), 5.35 – 5.20 (m, 1H), 4.80 – 4.62 (m, 1H), 4.60 – 4.45 (m, 2H), 4.35 – 4.10 (m, 1H), 4.06 – 3.96 (m, 3H), 1.49 – 1.28 (m, 8H), 0.90 – 0.82 (m, 6H).

³¹P NMR (162 MHz, CDCl₃) δ 2.36 (s), 2.22 (s).

HPLC: t_R = 3.00min; HPLC system: Agilent 1100; Phenomenex Gemini, C₁₈, 5u, 110A, 50 x 4.6 mm; Buffer A: 0.05% TFA in Water; Buffer B: 0.05% TFA in Acetonitrile; 2-98% Buffer B in 5 minutes @ 2mL/min.

LC/MS: t_R = 2.39min, MS m/z = 605.1 [M+1], 603.0 [M-1]; LC/MS system: Thermo LCQ

5 Advantage; Phenomenex Gemini, C₁₈, 5u, 110A, 30 x 4.6 mm; Buffer A: 0.1% Acetic acid in Water; Buffer B: 0.1% Acetic acid in Acetonitrile; 5-100% Buffer B in 2.5 mins then 100% for 0.9min @ 2 mL/min.



10 **Intermediate 6a - N-(7-((2S,3S,4R,5R)-4-(tert-butyldimethylsilyloxy)-5-((tert-butyldimethylsilyloxy)methyl)-3-fluoro-5-vinyltetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide**

Intermediate **2g**, N-(7-((2S,3S,4R,5R)-4-(tert-butyldimethylsilyloxy)-5-((tert-butyldimethylsilyloxy)methyl)-3-fluoro-5-(hydroxymethyl)tetrahydrofuran-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4-yl)benzamide, (220 mg, 0.35 mmol) was dissolved in 5 mL of anhydrous DMSO and stirred under N₂(g). Added EDCI (100 mg, 0.52 mmol) and then TFA-Pyridine (34 mg, 0.18 mmol). Stirred for 1 h. Added more EDCI (100 mg, 0.52 mmol) and stirred for 1 h. Monitoring by LC/MS showed starting material alcohol remained. Added more EDCI (100 mg, 0.52 mmol) and stirred for 1 h. Monitoring by LC/MS showed that the reaction reached full conversion. Diluted with ethyl acetate and washed with saturated NaHCO_{3(aq)} (2x) and then saturated NaCl_(aq). Dried organic layer over anhydrous Na₂SO₄ and concentrated under reduced pressure. Purified with silica gel column (0-20% EtOAc in hexanes). Combined fractions and concentrated under reduced pressure to give aldehyde as solid. Suspended Methyl triphenylphosphonium bromide (500 mg, 1.40 mmol) in 10mL anhydrous THF and stirred at -78 °C under Ar(g). Added 2.5 M *n*-butyllithium solution in hexane (560 µL, 1.40 mmol) dropwise. Stirred reaction mixture in an ice bath for 1 h to give a yellow mixture. Dissolved above prepared aldehyde in 5 mL of anhydrous THF and added to reaction dropwise. Removed ice bath and let reaction warmed to RT. Stirred for 3 h at RT. Added saturated aqueous NH₄Cl solution and extracted with ethyl acetate. Washed organic extract


15
20
25

with saturated $\text{NaHCO}_{3(\text{aq})}$ and then saturated $\text{NaCl}_{(\text{aq})}$. Dried organic over anhydrous Na_2SO_4 and concentrated under reduced pressure. Purified with silica gel column (0-20% EtOAc in hexanes) to afford intermediate **6a**.

¹H NMR (400 MHz, DMSO-*d*₆) δ 8.22 (br s, 1H), 8.03 (br s, 2H), 7.58 (dt, *J* = 40.4, 7.4 Hz, 3H), 7.12 (d, *J* = 4.7 Hz, 1H), 6.97 (s, 1H), 6.01 (dd, *J* = 17.5, 10.9 Hz, 1H), 5.58 (d, *J* = 22.8 Hz, 1H), 5.46 (dd, *J* = 17.5, 2.1 Hz, 1H), 5.25 (dd, *J* = 11.0, 2.0 Hz, 1H), 5.14 (ddd, *J* = 55.4, 4.9, 2.7 Hz, 1H), 4.61 (dd, *J* = 20.8, 4.8 Hz, 1H), 3.63 – 3.40 (m, 2H), 0.89 (s, 9H), 0.84 (s, 9H), 0.09 (d, *J* = 8.4 Hz, 6H), 0.00 (d, *J* = 14.1 Hz, 6H).

¹⁹F NMR (376 MHz, DMSO-*d*₆) δ -191.86 (d, *J* = 56.8 Hz).

10 MS m/z = 627.3 [M+1].




Example 13 - (2R,3R,4R,5S)-5-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-4-fluoro-2-(hydroxymethyl)-2-vinyltetrahydrofuran-3-ol

Intermediate **6a** (146 mg, 0.23 mmol) was dissolved into THF (10 mL) and the resulting solution was stirred in an ice bath. Added 1M TBAF solution in THF (700 μ L, 0.70 mmol) and stirred for 2 h. Diluted with EtOAc and washed with saturated NaCl_(aq) (5x). Dried organic layer over anhydrous Na₂SO₄ and concentrated under reduced pressure. Dissolved in 7 M ammonia in MeOH (7 mL) and stirred for 18 h. Concentrated reaction under reduced pressure. Purified with C₁₈ preparatory HPLC with TFA as modifier. Combined fractions and concentrated under reduced pressure. Dissolved in NaHCO_{3(aq)} and repurified with preparatory HPLC under neutral condition. Combined fractions and freeze-dried to afford example **13**.

25 ¹H NMR (400 MHz, D₂O) δ 7.54 (s, 1H), 6.62 – 6.49 (m, 2H), 5.98 – 5.79 (m, 1H), 5.55 – 5.36 (m, 2H), 5.31 (d, *J* = 11.1 Hz, 1H), 5.11 (ddd, *J* = 54.8, 5.2, 2.9 Hz, 1H), 4.42 (dd, *J* = 20.6, 4.8 Hz, 1H), 3.62 – 3.43 (m, 2H).

^{19}F NMR (376 MHz, D_2O) δ -193.23 (dd, J = 54.7, 44.2 Hz).

MS m/z = 295.2 $[\text{M}+1]$




5 Example 14 - (2R,3R,4R,5S)-5-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-2-ethyl-4-fluoro-2-(hydroxymethyl)tetrahydrofuran-3-ol

Example 13 (5 mg, 0.017 mmol) was dissolved into methanol (2 mL). 10% Pd/C Degussa Catalyst (2 mg) was then added and the resulting mixture was stirred under atmosphere of hydrogen gas. After 40 min, the resulting mixture was filtered to remove Pd/C and the filtrate was concentrated under reduced pressure. The residue was dissolved in water and freeze-dried to afford example 14.

^1H NMR (400 MHz, D_2O) δ 7.67 (s, 1H), 6.79 – 6.55 (m, 2H), 5.54 – 5.12 (m, 2H), 4.46 (dd, J = 15.1, 5.5 Hz, 1H), 3.65 – 3.44 (m, 2H), 1.89 – 1.44 (m, 2H), 0.84 (t, J = 7.6 Hz, 3H).

^{19}F NMR (376 MHz, D_2O) δ -197.62 (ddd, J = 54.5, 20.6, 15.0 Hz).

15 MS m/z = 297.3 $[\text{M}+1]$.




Example 15 (PD4) - S,S'-2,2'-(((2R,3R,4R,5S)-5-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-4-fluoro-3-hydroxy-2-vinyltetrahydrofuran-2-yl)methoxyphosphoryl)bis(oxy)bis(ethane-2,1-diyl) bis(2,2-dimethylpropanethioate)

Example 13 (5 mg, 0.017 mmol) was dissolved into anhydrous DMF (0.5 mL). Added *p*-nitro-phenonate (13 mg, 0.026 mmol) in one portion. Added 1M *t*-butylmagnesium chloride in THF (25 μ L, 0.026 mmol) dropwise. Stirred for 1 h. Warmed to 50 $^{\circ}$ C and stirred for 2 h. Added more *p*-nitro-phenonate (13 mg, 0.026 mmol) and stirred for 2 h. Added more 1M *t*-butylmagnesium chloride in THF (25 μ L, 0.026 mmol) and stirred for 16 h at 50 $^{\circ}$ C. Cooled to RT. The resulting mixture was purified directly by preparatory HPLC column and eluted with linear gradient 0-100% ACN in water to afford example 15 (PD4).

^1H NMR (400 MHz, CD_3OD) δ 7.84 (s, 1H), 6.92 (d, J = 4.5 Hz, 1H), 6.80 (d, J = 4.5 Hz, 1H), 6.10 (dd, J = 17.4, 10.9 Hz, 1H), 5.67 (dd, J = 5.8, 1.9 Hz, 1H), 5.61 (s, 1H), 5.45 – 5.35 (m, 1H), 5.15 (ddd, J = 55.6, 5.0, 2.2 Hz, 1H), 4.65 (dd, J = 22.5, 5.1 Hz, 1H), 4.13 (dd, J = 11.1, 5.2 Hz, 1H), 4.08 – 3.95 (m, 5H), 3.06 (dd, J = 7.0, 6.1 Hz, 4H), 1.21 (s, 9H), 1.18 (s, 9H).

^{19}F NMR (376 MHz, CD_3OD) δ 192.99 (td, J = 55.7, 23.6 Hz).

MS m/z = 663.0 $[\text{M}+1]$.



Example 16 (PD5) - (2S)-2-ethylbutyl 2-((((2R,3S,4R,5S)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-azido-3,4-dihydroxytetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate.

Example 5 (5 mg, 0.016 mmol) was dissolved in anhydrous *N*-methyl-2-pyrrolidone (0.2 mL) and THF (0.1 mL) was added under an argon atmosphere. *tert*-Butyl magnesium chloride (1M in THF, 24 μ L, 0.024 mmol) was then added at RT, and white solids precipitated. After 5 min, a solution of *p*-nitrophenylphosphoramidate PD3c (15 mg, 0.032 mmol) in THF (0.1 mL) was added to the reaction mixture in one portion, and the resulting mixture was heated to 50 $^{\circ}$ C. After 3.5 h, the reaction mixture was allowed to cool to RT and was stirred for 18 h. *p*-Nitrophenylphosphoramidate PD3c (50 mg, 0.111 mmol) and *tert*-butyl magnesium chloride (1M in THF, 24 μ L, 0.024 mmol) were then added and the reaction mixture was stirred for an additional 5 d. The resulting residue was then purified directly by preparatory HPLC (Phenomenex Synergi 4u Hydro-RR 80 $\text{\AA}$ 150 x 30 mm column, 40-100%

acetonitrile/water gradient). The fractions containing the desired product were combined and were lyophilized to afford example **16 (PD5)** (2:1 diastereomeric mixture).


^1H NMR (400 MHz, CDCl_3) δ 7.88 (br s, 1H), 7.33 – 7.22 (br m, 2H), 7.22 – 7.10 (br m, 3H), 6.69 (br d, J = 4.4 Hz, 1H), 6.61 (br d, J = 4.5 Hz, 1H), 5.64 – 5.56 (m, 1H), 4.54 (d, J = 6.3 Hz, 1H), 4.50 – 4.20 (m, 3H), 4.11 – 3.94 (m, 3H), 3.90 – 3.76 (m, 1H), 1.49 (s, J = 6.2 Hz, 1H), 1.40 – 1.24 (m, 7H), 0.86 (t, J = 7.4 Hz, 6H).

^{31}P NMR (162 MHz, CDCl_3) δ 2.68 (s), 2.56 (s).

LC/MS: t_R = 1.70 min, MS m/z = 619.09 $[\text{M}+1]^+$; LC system: Thermo Accela 1250 UHPLC.

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-2.0 min 2-100% ACN, 2.0 min-3.05 min 100% ACN, 3.05 min-3.2 min 100%-2% ACN, 3.2 min-3.5 min 2% ACN at 2 μ L/min.

HPLC: t_R = 3.010 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.



Example 17 (TP5) - ((2R,3R,4R,5S)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-4-fluoro-3-hydroxy-2-vinyltetrahydrofuran-2-yl)methyl tetrahydrogen triphosphate.

To a solution of example **13** (6.0 mg, 0.020 mmol) in $\text{PO}(\text{OMe})_3$ (0.6 mL) at 0 °C was added POCl_3 (50 mg, 0.32 mmol). The reaction mixture was stirred at 0 °C for 6 h, at which point ion-exchange HPLC showed approximately 90% conversion. A solution of pyrophosphate tributylamine salts (250 mg) in ACN (0.6 mL) was added, followed by tributylamine (110 mg, 0.59 mmol). The reaction mixture was stirred at 0 °C for 1 h. The reaction was quenched with triethylammonium bicarbonate buffer (1 M, 5 mL). The reaction mixture was stirred at RT for 0.5 h, then concentrated and co-evaporated with water twice.


The residue was dissolved in H₂O (5 mL) and loaded to a ion-exchange column, eluted with H₂O, then 5-35% triethylammonium bicarbonate buffer (1M)-H₂O. The product fractions were combined, concentrated and co-evaporated with H₂O. The solid residue was dissolved in 3 mL of H₂O and 100 μ L of NaOH (1N) was added. The resulting mixture was purified with C-18 column, eluted with H₂O, and the fractions containing product were combined and concentrated under reduced pressure to afford Example 17 (TP5) as the tetra-sodium salt.

¹H NMR (400 MHz, D₂O): δ 7.74 (s, 1H), 6.89 (d, J = 4.4 Hz, 1H), 6.81 (d, J = 4.6 Hz, 1H), 6.00 (dd, J = 17.4, 11.1 Hz, 1H), 5.72 (d, J = 23.3 Hz, 1H), 5.49 (d, J = 16.9 Hz, 1H), 5.32 (d, J = 11.1 Hz, 1H), 5.14 (dd, J = 54.0, 4.6 Hz, 1H), 4.72 (dd, J = 23.7, 4.5 Hz, 1H), 4.09 (dd, J = 11.3, 5.8 Hz, 1H), 3.79 (dd, J = 11.6, 3.8 Hz, 1H).

³¹P NMR (162 MHz, D₂O): δ -8.38 (d, J = 20.5 Hz), -13.67 (d, J = 19.3 Hz), -24.20 (t, J = 19.9 Hz).

¹⁹F NMR (376 MHz, CDCl₃) δ -194.58 (dt, J = 55.0, 23.8 Hz).

MS m/z = 533.0[M-1], 535.0 [M+1].



Example 18 (TP6) - ((2R,3R,4R,5S)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-ethyl-4-fluoro-3-hydroxytetrahydrofuran-2-yl)methyl tetrahydrogen triphosphate.

To a solution of example 14 (5.0 mg, 0.017 mmol) in PO(OMe)₃ (0.6 mL) at 0 °C was added POCl₃ (45 mg, 0.30 mmol). The reaction mixture was stirred at 0 °C for 6 h, at which point ion-exchange HPLC showed approximately 90% conversion. A solution of pyrophosphate tributylamine salts (250 mg) in ACN (0.6 mL) was added, followed by tributylamine (110 mg, 0.59 mmol). The reaction mixture was stirred at 0 °C for 1 h. The reaction was quenched with triethylammonium bicarbonate buffer (1 M, 5 mL). The reaction mixture was stirred at RT for 0.5 h, then concentrated and co-evaporated with water twice.

The residue was dissolved in H₂O (5 mL) and loaded to a ion-exchange column, eluted with


H₂O, then 5-35% triethylammonium bicarbonate buffer (1M)-H₂O. The product fractions were combined, concentrated and co-evaporated with H₂O. The solid residue was dissolved in 3 mL of H₂O and 100 μ L of NaOH (1N) was added. The resulting mixture was purified with C-18 column, eluted with H₂O, and the fractions containing product were combined and concentrated under reduced pressure to afford **18 (TP6)** as the tetra-sodium salt.

¹H NMR (400 MHz, D₂O): δ 7.73 (s, 1H), 6.86 (d, J = 4.6 Hz, 1H), 6.80 (d, J = 4.6 Hz, 1H), 5.60 (dd, J = 21.9, 3.5 Hz, 1H), 5.23 (dt, J = 55.2, 4.2 Hz, 1H), 4.65 (dd, J = 20.6, 5.3 Hz, 1H), 4.08 – 3.84 (m, 3H), 1.83 (dq, J = 14.4, 7.4, 6.9 Hz, 1H), 1.62 (dq, J = 15.0, 7.5 Hz, 1H), 0.87 (t, J = 7.5 Hz, 3H).

³¹P NMR (162 MHz, D₂O): -5.72 (d, J = 20.2 Hz), -10.81 (d, J = 19.3 Hz), -21.60 (t, J = 19.8 Hz).

¹⁹F NMR (376 MHz, CDCl₃) δ -194.77 (dt, J = 55.2, 21.2 Hz).

MS m/z = 535.1[M-1], 536.9.0 [M+1].



Intermediate 8a - N-(7-((2S,3R,4R,5S)-3-fluoro-4-hydroxy-5-(iodomethyl)tetrahydrofuran-2-yl)pyrrolo[2,1-f][1,2,4]triazin-4-yl)benzamide.

To an argon purged flask was added **2b** (68 mg, 0.183 mmol) in DMF (2 mL.) followed by methyltriphenoxyphosphonium iodide (124 mg, 0.274 mmol). The reaction was allowed to stir at room temperature for 5 min where complete conversion to product was observed by LCMS. The reaction was quenched with methanol and solvents were removed under reduced pressure. The crude material was partitioned between EtOAc and H₂O. The organics were separated and washed with brine. The resulting material was dried over Na₂SO₄, filtered and solvent removed under reduced pressure. The crude material was purified by silica gel chromatography (20 – 100 % EtOAc/Hexanes) to afford intermediate **8a**.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.17 (m, 3H), 7.63 (t, $J = 7.4$ Hz, 1H), 7.53 (t, $J = 7.6$ Hz, 2H), 7.17 – 6.96 (m, 2H), 5.74 (s, 1H), 5.60 (d, $J = 24.9$ Hz, 1H), 5.19 (ddd, $J = 54.6, 4.5, 2.1$ Hz, 1H), 4.09 – 3.92 (m, 1H), 3.72 (t, $J = 6.4$ Hz, 1H), 3.63 (dd, $J = 11.0, 3.4$ Hz, 1H), 3.44 (dd, $J = 11.0, 5.9$ Hz, 1H).

5 ^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ -194.23 (m).


LC/MS: $t_R = 1.13$ min, MS $m/z = 483.23$ [M+1]

LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

10 Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN.



Intermediate 8b - (3R,4R,5S)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-4-fluoro-2-methylenetetrahydrofuran-3-ol.

15 Intermediate **8a** (80 mg, 0.166 mmol) was dissolved into THF. DBU (0.074 mL, 0.498 mmol) was added in one portion. The reaction was then heated to 60 °C in an oil bath for 16 h. The reaction was cooled to room temperature and the solvent was removed under reduced pressure. The crude material was purified by silica gel chromatography (0-70% EtOAc/Hex) to afford intermediate **8b**.

20 ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.34-8.05 (m, 3H), 7.63 (t, $J = 7.4$ Hz, 1H), 7.53 (t, $J = 7.6$ Hz, 2H), 7.13 (d, $J = 4.7$ Hz, 1H), 6.85 (d, $J = 4.4$ Hz, 1H), 5.97 – 5.82 (m, 2H), 5.39 – 5.13 (m, 1H), 4.89 – 4.69 (m, 1H), 4.38 (d, $J = 2.1$ Hz, 1H), 4.16 (t, $J = 1.8$ Hz, 1H).

^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ -198.14 (ddd, $J = 53.9, 24.7, 20.9$ Hz).

LC/MS: $t_R = 1.05$ min, MS $m/z = 355.15$ [M+1]


LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min

5 100%-2% ACN, 1.85 min-2 min 2% ACN.



Intermediate 8c - (2S,3R,4R,5S)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-azido-4-fluoro-2-(iodomethyl)tetrahydrofuran-3-ol.

10 Benzyltrimethylammonium chloride (55 mg, 0.296 mmol) and sodium azide (19.3 mg, 0.296 mmol) were dissolved into ACN (1 mL). The resulting mixture was stirred at RT overnight and was then filtered and added via syringe to a solution of intermediate **8b** (50 mg, 0.141 mmol) in THF (1 mL). *N*-methylmorpholine (0.078 mL, 0.706 mmol) was then added followed by the dropwise addition of a solution of iodine (65 mg, 0.25 mmol) in THF (1 mL). After 15 min, *N*-acetyl cysteine was added until the evolution of gas was no longer

15 observable. Then saturated aqueous sodium thiosulfate was added until the solution was light yellow. The crude mixture was partitioned between EtOAc and H₂O. The phases were split and the organic layer was dried over sodium sulfate, filtered, and concentrated under reduced pressure. The crude material was purified by silica gel chromatography (0-60% EtOAc/Hex) to afford intermediate **8c**.

20 ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.31-8.05 (m, 3H), 7.63 (t, *J* = 7.5 Hz, 1H), 7.53 (t, *J* = 7.6 Hz, 2H), 7.14 (d, *J* = 4.6 Hz, 1H), 7.04 (s, 1H), 6.34 (d, *J* = 6.9 Hz, 1H), 5.80 (d, *J* = 23.7 Hz, 1H), 5.55 – 5.31 (m, 1H), 4.62 (dt, *J* = 21.9, 5.9 Hz, 1H), 3.78 – 3.56 (m, 2H).


¹⁹F NMR (376 MHz, DMSO-*d*₆) δ -194.44 (dt, *J* = 54.7, 22.8 Hz).

LC/MS: *t*_R = 1.19 min, MS *m/z* = 524.09 [*M*+1]; LC system: Thermo Accela 1250 UHPLC

25 MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN.



Intermediate 8d - (2S,3R,4S,5S)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-azido-4-fluoro-2-(iodomethyl)tetrahydrofuran-3-yl acetate.

To a solution of intermediate **8c** (40 mg, 0.076 mmol) in THF (1mL) was added acetic anhydride (0.009 mL, 0.092 mmol) followed by DMAP (10 mg, 0.082 mmol) at RT. After 15 min, the reaction mixture was quenched with methanol, and the resulting mixture concentrated under reduced pressure. The crude was purified by silica gel chromatography (0-50% EtOAc/Hex) to afford intermediate **8d**.

¹H NMR (400 MHz, CDCl₃) δ 8.15-8.02 (m, 3H), 7.62 (t, *J* = 7.3 Hz, 1H), 7.53 (t, *J* = 7.6 Hz, 2H), 7.44 (d, *J* = 4.6 Hz, 1H), 7.00 (d, *J* = 4.6 Hz, 1H), 5.95 – 5.80 (m, 1H), 5.70 – 5.43 (m, 2H), 3.71 (d, *J* = 11.3 Hz, 1H), 3.60 (d, *J* = 11.3 Hz, 1H), 2.25 (s, 3H).


¹⁹F NMR (376 MHz, CDCl₃) δ -192.78 (ddd, *J* = 55.7, 24.6, 18.5 Hz).

LC/MS: *t_R* = 1.35 min, MS *m/z* = 566.14 [*M*+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN.



Intermediate 8e - ((2R,3R,4S,5S)-3-acetoxy-2-azido-5-(4-benzamidopyrrolo[2,1-f][1,2,4]triazin-7-yl)-4-fluorotetrahydrofuran-2-yl)methyl benzoate.

To a solution of intermediate **8d** (30 mgs, 0.053 mmol) in DMF (2 mL) was added 15-Crown-5 (0.105 mL, 0.531 mmol) and sodium benzoate (77 mg, 0.531 mmol) at RT. The reaction was then heated to 105 °C. After 30 h, the reaction mixture was allowed to RT and was partitioned between 5% LiCl_(aq) and EtOAc. The phases were split and the aqueous phase was washed with EtOAc (2 x). The combined organic extracts were dried over anhydrous sodium sulfate, and were concentrated under reduced pressure. The crude residue was purified by silica gel chromatography (0 - 60% EtOAc/Hex) to afford intermediate **8e**.

¹H NMR (400 MHz, CDCl₃) δ 8.25 – 7.97 (m, 4H), 7.69 – 7.40 (m, 6H), 7.36 (d, *J* = 4.7 Hz, 1H), 6.95 – 6.80 (m, 1H), 5.90 (d, *J* = 25.0 Hz, 1H), 5.65 (d, *J* = 1.9 Hz, 1H), 5.62 – 5.48 (m, 1H), 4.69 (dd, *J* = 79.3, 12.0 Hz, 2H), 2.20 (s, 3H).


¹⁹F NMR (376 MHz, CDCl₃) δ -192.57 (ddd, *J* = 53.9, 25.1, 22.0 Hz).

LC/MS: *t_R* = 1.45 min, MS *m/z* = 560.14 [M+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN.



Example 19 - (2R,3R,4R,5S)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-azido-4-fluoro-2-(hydroxymethyl)tetrahydrofuran-3-ol.

To intermediate **8e** (24 mg, 0.043 mmol) was added 7N NH₃ in CH₃OH (2 mL) at RT. After 16 h, the resulting mixture was concentrated under reduced pressure. The crude residue was purified by reverse phase HPLC without acid modifier to afford example **19**.

¹H NMR (400 MHz, Methanol-*d*₄) δ 7.81 (s, 1H), 6.85 (d, *J* = 4.5 Hz, 1H), 6.80 (d, *J* = 4.5 Hz, 1H), 5.80 (dd, *J* = 24.7, 1.9 Hz, 1H), 5.22 (ddd, *J* = 55.6, 5.1, 1.9 Hz, 1H), 4.63 (dd, *J* = 22.9, 5.1 Hz, 1H), 3.81 (d, *J* = 12.1 Hz, 1H), 3.70 (d, *J* = 12.2 Hz, 1H).


¹⁹F NMR (376 MHz, Methanol-*d*₄) δ -195.30 (ddd, *J* = 55.5, 24.6, 22.9 Hz).

LC/MS: *t*_R = 0.61 min, MS *m/z* = 310.02 [M+1]; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ C18 100A, 50 x 3.00 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN.



Intermediate 9b - ((3aR,5R,6S,6aR)-6-(benzyloxy)-5-((benzyloxy)methyl)-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-5-yl)methanol

((3aR,6S,6aR)-6-(benzyloxy)-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxole-5,5-diyl)dimethanol (**9a**, Purchased from Carbosynth, 10.0 g, 32.2 mmol) was added to a solution of sodium hydride (60% by wt, 1.55 g, 38.7 mmol) in THF (100 mL) at 0 °C under an argon atmosphere. After 10 min, benzyl bromide (4.54 mL, 38.6 mmol) was added and the reaction mixture was allowed to warm to RT. After 2 h, the reaction was quenched with saturated aqueous ammonium chloride solution (500 mL). The resulting mixture was extracted with ethyl acetate (500 mL). The organic phase was then washed with brine (400 mL), was dried over anhydrous sodium sulfate, and was concentrated under reduced


pressure to afford a colorless oil. The crude residue was purified via SiO₂ column chromatography (220 g SiO₂ Combiflash HP Gold Column, 0–100% ethyl acetate/hexanes) to afford intermediate **9a** (9.49 g, 73%) as a colorless oil.

¹H NMR (400 MHz, DMSO-*d*₆) δ 7.38 – 7.19 (m, 10H), 5.68 (app t, *J* = 3.6 Hz, 1H), 4.73 (q, *J* = 4.4 Hz, 1H), 4.63 (d, *J* = 12.1 Hz, 1H), 4.49 – 4.36 (m, 3H), 4.24 (br s, 1H), 4.20 – 4.13 (m, 1H), 3.81 (d, *J* = 11.9 Hz, 1H), 3.56 (d, *J* = 11.9 Hz, 1H), 3.46 (q, *J* = 10.3 Hz, 2H), 1.47 (s, 3H), 1.25 (s, 3H).

LC/MS: *t*_R = 1.88 min, MS *m/z* = 423.31 [M+Na]; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ XB-C18 100A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min–2.0 min 2–100% ACN, 2.0 min–3.05 min 100% ACN, 3.05 min–3.2 min 100%–2% ACN, 3.2 min–3.5 min 2% ACN at 2μl/min.

HPLC: *t*_R = 3.79 min; HPLC system: Agilent 1100 series.; Column: Gemini 5μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min–5.0 min 2–98% ACN, 5.0 min–6.0 min 98% ACN at 2 mL/min.

TLC: eluent: 40% ethyl acetate in hexanes, *R*_f = 0.4 (UV)




Hz, 1H), 4.46 (d, J = 12.0 Hz, 1H), 4.37 (d, J = 4.4 Hz, 1H), 3.68 (d, J = 11.0 Hz, 1H), 3.61 (d, J = 11.0 Hz, 1H), 1.60 (s, 3H), 1.35 (s, 3H).

LC/MS: t_R = 1.99 min, MS m/z = 421.25 $[M+Na]$; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-2.0 min 2-100% ACN, 2.0 min-3.05 min 100% ACN, 3.05 min-3.2 min 100%-2% ACN, 3.2 min-3.5 min 2% ACN at 2 μ L/min.

HPLC: t_R = 4.09 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.

TLC: eluent: 40% ethyl acetate in hexanes, R_f = 0.6 (UV)



Intermediate 9d - (3aR,5R,6S,6aR)-6-(benzyloxy)-5-((benzyloxy)methyl)-2,2-dimethyl-5-vinyltetrahydrofuro[2,3-d][1,3]dioxole


2.5M n-butyllithium (6.02 mL) was added to a solution of methyltriphenylphosphonium bromide (5.38 g, 15.1 mmol) in tetrahydrofuran (20 mL) at -78 °C. The reaction was allowed to warm to 0 °C, and a solution of intermediate **9c** (2.00 g, 5.02 mmol) in tetrahydrofuran (5 mL) was added slowly via syringe. The reaction mixture was allowed to warm to RT, and was stirred for 4 h. The reaction mixture was then quenched with saturated aqueous ammonium chloride solution (10 mL) and was partitioned between water (200 mL) and ethyl acetate (200 mL). The layers were split and the organic layer was washed with brine (200 mL), dried over anhydrous sodium sulfate, and was concentrated under reduced pressure. The crude residue was purified via SiO₂ column chromatography (120 g SiO₂ Combiflash HP Gold Column, 0–50% ethyl acetate/hexanes) to afford intermediate **9d** (1.01 g, 51%) as a colorless oil.

^1H NMR (400 MHz, CDCl_3) δ 7.42 – 7.17 (m, 10H), 6.19 (dd, J = 17.6, 11.0 Hz, 1H), 5.76 (d, J = 3.9 Hz, 1H), 5.52 (dd, J = 17.5, 1.9 Hz, 1H), 5.25 (dd, J = 11.1, 1.8 Hz, 1H), 4.76 (d, J = 12.3 Hz, 1H), 4.62 – 4.55 (m, 2H), 4.52 (d, J = 12.1 Hz, 1H), 4.41 (d, J = 12.1 Hz, 1H), 4.25 (d, J = 4.9 Hz, 1H), 3.32 (d, J = 1.5 Hz, 2H), 1.52 (s, 3H), 1.29 (s, 3H)

5 LC/MS: t_R = 2.13 min, MS m/z = 419.24 $[\text{M}+\text{Na}]$; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-2.0 min 2-100% ACN, 2.0 min-3.05 min 100% ACN, 3.05 min-3.2 min 100%-2% ACN, 3.2 min-3.5 min 2% ACN at 2 μ L/min.

10 HPLC: t_R = 4.37 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.

TLC: eluent: 50% ethyl acetate in hexanes, R_f = 0.55 (UV)



15 **Intermediate 9e - (3R,4S,5R)-4-(benzyloxy)-5-((benzyloxy)methyl)-2-methoxy-5-vinyltetrahydrofuran-3-ol**


4M HCl in dioxane (320 μ L) was added to a solution of intermediate **9d** (1.01 g, 2.55 mmol) in methanol (12.5 mL) at RT. After 1.25 h, the reaction mixture was partitioned
 20 between ethyl acetate (100 mL) and saturated aqueous sodium bicarbonate solution (100 mL). The phases were split and the organic phase was washed with brine (100 mL), was dried over anhydrous sodium sulfate, and was concentrated under reduced pressure to afford crude intermediate **9e** (1.05 g, ~2.5 : 1 mixture of 1' anomers) as a colorless oil.

LC/MS: major anomer t_R = 2.00 min, MS m/z = 393.22 $[\text{M}+\text{Na}]$, minor anomer t_R = 1.98 min,
 25 MS m/z = 393.22 $[\text{M}+\text{Na}]$; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm; Solvents: Acetonitrile with

0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-2.0 min 2-100% ACN, 2.0 min-3.05 min 100% ACN, 3.05 min-3.2 min 100%-2% ACN, 3.2 min-3.5 min 2% ACN at 2 μ L/min.

HPLC: major anomer t_R = 4.01 min, minor anomer t_R = 3.955 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.

TLC: eluent: 25% ethyl acetate in hexanes, Major anomer R_f = 0.30 (UV), Minor anomer R_f = 0.25 (UV)




Intermediate 9f - (2R,3S,4R)-3,4-bis(benzyloxy)-2-((benzyloxy)methyl)-5-methoxy-2-vinyltetrahydrofuran

NaH (60% wt, 130 mg, 3.2 mmol) was added as a solid to a solution of intermediate **9e** (1.0 g, 2.7 mmol) in THF (13.5 mL) at RT under an argon atmosphere. After 15 min, benzyl bromide (0.38 mL, 3.2 mmol) was added and the reaction mixture was stirred for 4 h. The reaction mixture was quenched with saturated aqueous ammonium chloride solution (5 mL) and was partitioned between ethyl acetate (100 mL) and brine (100 mL). The phases were split and the organic layer was dried over anhydrous sodium sulfate, and was concentrated under reduced pressure to afford crude intermediate **9f** (1.57 g, ~2:1 mixture of 1' anomers) as a colorless oil.

LC/MS: major anomer t_R = 1.88 min, MS m/z = 483.36 [M+Na], minor anomer t_R = 1.83 min, MS m/z = 483.36 [M+Na]; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-1.5 min 2-100% ACN, 1.5 min-2.2 min 100% ACN, 2.2 min-2.4 min 100%-2% ACN, 2.4 min-2.5 min 2% ACN at 2 μ L/min.

HPLC: major anomer t_R = 4.83 min, minor anomer t_R = 4.62 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.




5

Intermediate 9g - (3R,4S,5R)-3,4-bis(benzyloxy)-5-((benzyloxy)methyl)-5-vinyltetrahydrofuran-2-ol

A solution of TFA (16 mL) and water (1.6 mL) was added to intermediate **9f** (1.5 g, 3.2 mmol) at 0 °C, and the reaction mixture was allowed to warm to RT. After 9 h, water (1 mL) was added and the reaction mixture was allowed to stir an additional 10 h. The reaction mixture was then concentrated under reduced pressure. The crude residue was dissolved into ethyl acetate (200 mL) and was washed with saturated aqueous sodium bicarbonate solution (2 x 150 mL) and brine (150 mL). The organic layer was dried over anhydrous sodium sulfate and was concentrated under reduced pressure. The residue was purified via SiO₂ column chromatography (24 g SiO₂ Combiflash HP Gold Column, 0–100% ethyl acetate/hexanes). The fractions containing the desired product were combined to afford intermediate **9g** (580 mg) as a colorless oil that was a mixture with other impurities. The mixture was used directly in the following step.

LC/MS: t_R = 3.13 min, MS m/z = 463.88 [M+OH]; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-2.0 min 2-100% ACN, 2.0 min-3.05 min 100% ACN, 3.05 min-3.2 min 100%-2% ACN, 3.2 min-3.5 min 2% ACN at 2 μ L/min.

HPLC: t_R = 4.34 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.



Intermediate 9h - (3R,4S,5R)-3,4-bis(benzyloxy)-5-((benzyloxy)methyl)-5-vinyldihydrofuran-2(3H)-one


Tetrapropylammonium perruthenate (45.7 mg, 130 μmol) and 4-methylmorpholine *N*-oxide (457 mg, 3.89 mmol) were added to a solution of intermediate **9g** (580 mg, 1.30 mmol) and 4Å MS (100 mg) in DCM (6.45 mL) at RT. After 1 h, silica gel (~500 mg) was added to the reaction mixture and the resulting slurry was filtered through a plug of silica gel (~1 g). The filtrate was concentrated under reduced pressure. The crude residue was purified via SiO₂ column chromatography (12 g SiO₂ Combiflash HP Gold Column, 0–100% ethyl acetate/hexanes) to afford intermediate **9h** (254 mg, 18% over two steps) as a colorless oil.

¹H NMR (400 MHz, CDCl₃) δ 7.38 – 7.23 (m, 13H), 7.20 – 7.13 (m, 2H), 5.91 (dd, *J* = 17.5, 11.2 Hz, 1H), 5.49 (dd, *J* = 17.5, 0.9 Hz, 1H), 5.33 (dd, *J* = 11.2, 0.9 Hz, 1H), 4.96 (d, *J* = 12.0 Hz, 1H), 4.74 – 4.68 (m, 2H), 4.55 – 4.47 (m, 3H), 4.39 (d, *J* = 11.9 Hz, 1H), 4.20 (d, *J* = 6.0 Hz, 1H), 3.55 (d, *J* = 10.8 Hz, 1H), 3.46 (d, *J* = 10.8 Hz, 1H)

LC/MS: *t_R* = 2.19 min, MS *m/z* = 444.78 [M+H]⁺; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min–2.0 min 2–100% ACN, 2.0 min–3.05 min 100% ACN, 3.05 min–3.2 min 100%–2% ACN, 3.2 min–3.5 min 2% ACN at 2 μ L/min.

HPLC: *t_R* = 4.53 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min–5.0 min 2–98% ACN, 5.0 min–6.0 min 98% ACN at 2 mL/min.

TLC: eluent: 25% ethyl acetate in hexanes, *R_f* = 0.45 (UV)




Intermediate 9i - (3R,4S,5R)-2-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-3,4-bis(benzyloxy)-5-((benzyloxy)methyl)-5-vinyltetrahydrofuran-2-ol

n-Butyllithium (2.5M in hexanes, 1.0 mL, 2.5 mmol) was added rapidly to a suspension of intermediate **1b** (0.21 g, 0.81 mmol) and 1,2-bis(chlorodimethylsilyl)ethane (0.17 g, 0.81 mmol) in THF (4 mL) at -78 °C under an argon atmosphere. The resulting mixture was then transferred via cannula to a solution of **9h** (0.18 g, 0.41 mmol) in THF (1 mL) at -78 °C under an argon atmosphere. After 20 min, the reaction mixture was allowed to warm to 0 °C and was stirred for 15 min. The reaction mixture was quenched with saturated aqueous ammonium chloride solution (1 mL). The resulting mixture was diluted with ethyl acetate (100 mL) and was washed with saturated aqueous sodium bicarbonate solution (100 mL) and brine (100 mL). The organic layer was dried over anhydrous sodium sulfate, and was concentrated under reduced pressure. The crude residue was purified via SiO₂ column chromatography (12 g SiO₂ Combiflash HP Gold Column, 0–100% ethyl acetate/hexanes) to afford intermediate **9i** (11.1 mg, 5%, mixture of isomers) as a colorless oil.

LC/MS: t_R = 1.97 min, MS m/z = 579.27 [M+H]⁺; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ XB-C18 100A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-2.0 min 2-100% ACN, 2.0 min-3.05 min 100% ACN, 3.05 min-3.2 min 100%-2% ACN, 3.2 min-3.5 min 2% ACN at 2μl/min.

HPLC: t_R = 3.37 min; HPLC system: Agilent 1100 series.; Column: Gemini 5μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.

TLC: eluent: ethyl acetate, R_f = 0.3 (UV)



Intermediate 9j - 7-((2S,3S,4S,5R)-3,4-bis(benzyloxy)-5-((benzyloxy)methyl)-5-vinyltetrahydrofuran-2-yl)pyrrolo[2,1-f][1,2,4]triazin-4-amine


To a solution of intermediate **9i** (11.0 mg, 19.0 μmol) and triethylsilane (0.5 mL) in DCM (1 mL) was added boron trifluoride diethyl etherate (0.1 mL) slowly at 0 °C under an argon atmosphere. After 1 h, the reaction mixture was slowly diluted with saturated aqueous sodium bicarbonate solution (10 mL), and the resulting mixture was extracted with ethyl acetate (2 x 10 mL), was dried over anhydrous sodium sulfate, and was concentrated under reduced pressure. The crude residue was purified via SiO_2 column chromatography (4 g SiO_2 Combiflash HP Gold Column, 0–100% ethyl acetate/hexanes) to afford intermediate **9j** (7.7 mg, 72%) as a colorless film.

^1H NMR (400 MHz, CDCl_3) δ 7.87 (s, 1H), 7.37 – 7.17 (m, 15H), 6.73 (d, J = 4.5 Hz, 1H), 6.51 (d, J = 4.5 Hz, 1H), 6.23 (dd, J = 17.5, 10.9 Hz, 1H), 5.70 (d, J = 3.9 Hz, 1H), 5.59 (dd, J = 17.5, 1.8 Hz, 1H), 5.32 (dd, J = 10.9, 1.7 Hz, 1H), 4.72 – 4.56 (m, 4H), 4.49 (d, J = 11.9 Hz, 2H), 4.43 (d, J = 5.6 Hz, 1H), 4.25 (dd, J = 5.6, 4.0 Hz, 1H), 3.56 (s, 2H).

LC/MS: t_R = 2.32 min, MS m/z = 563.33 $[\text{M}+\text{H}]^+$; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min–2.0 min 2–100% ACN, 2.0 min–3.05 min 100% ACN, 3.05 min–3.2 min 100%–2% ACN, 3.2 min–3.5 min 2% ACN at 2 $\mu\text{L}/\text{min}$.

HPLC: t_R = 3.51 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min–5.0 min 2–98% ACN, 5.0 min–6.0 min 98% ACN at 2 mL/min.

TLC: eluent: ethyl acetate, R_f = 0.40 (UV)




Example 20 - (2R,3S,4R,5S)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-(hydroxymethyl)-2-vinyltetrahydrofuran-3,4-diol

Boron tribromide (1M, 0.06 mL, 60 μ mol) was added dropwise to a solution of intermediate **9j** (7.7 mg, 13.7 μ mol) in dichloromethane (1 mL) at -78 $^{\circ}$ C under an argon atmosphere. After 1 h, the reaction mixture was allowed to warm to 0 $^{\circ}$ C, and was stirred an additional 1.5 h. The reaction was cooled to -78 $^{\circ}$ C and was quenched with a 2:1 methanol/pyridine solution (1.5 mL). The resulting mixture was allowed to warm to RT, and was concentrated under reduced pressure. The crude residue was purified by preparatory HPLC (Phenominex Synergi 4u Hydro-RR 80 $\text{\AA}$ 150 x 30 mm column, 0-100% acetonitrile/water gradient) to afford example **20** (0.5 mg, 13%) as a white solid.

¹H NMR (400 MHz, CD₃OD) δ 7.78 (s, 1H), 6.88 (d, *J* = 4.5 Hz, 1H), 6.76 (d, *J* = 4.5 Hz, 1H), 6.02 (dd, *J* = 17.4, 11.0 Hz, 1H), 5.47 (dd, *J* = 17.4, 2.0 Hz, 1H), 5.23 (dd, *J* = 10.9, 2.1 Hz, 1H), 5.15 (d, *J* = 8.3 Hz, 1H), 4.72 (dd, *J* = 8.2, 5.7 Hz, 1H), 4.34 (d, *J* = 5.7 Hz, 1H), 3.60 (d, *J* = 11.8 Hz, 1H), 3.49 (d, *J* = 11.8 Hz, 1H)

LC/MS: t_R = 0.84 min, MS m/z = 293.19 [M+H]⁺; LC system: Thermo Accela 1250 UHPLC;
MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm; Solvents:
Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-1.5 min 2-
100% ACN, 1.5 min-2.2 min 100% ACN, 2.2 min-2.4 min 100%-2% ACN, 2.4 min-2.5 min
20 2% ACN at 2 μ l/min

HPLC: t_R = 2.181 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.




Example 21 - (2R,3R,4R,5S)-5-(4-amino-5-fluoropyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-azido-4-fluoro-2-(hydroxymethyl)tetrahydrofuran-3-ol

Example **19** (237 mg, 0.766 mmol) and selectfluor (407 mg, 1.15 mmol) were suspended in acetonitrile (5 mL), and AcOH (0.2 mL) was added. The resulting mixture was stirred at room temperature for 30 min, and was then neutralized with sodium bicarbonate solution and filtered to remove solids. Upon concentration *in vacuo*, the residue was purified by preparative HPLC (acetonitrile 0 to 30% in water) to give example **21** (27 mg, 11%) as an off-white solid.

^1H NMR (400 MHz, CD_3OD) δ 7.73 (s, 1H), 6.63 (s, 1H), 5.80 (dd, $J = 23.9, 1.7$ Hz, 1H), 5.16 (ddd, $J = 55.3, 5.0, 1.7$ Hz, 1H), 4.56 (dd, $J = 23.5, 5.0$ Hz, 1H), 3.94 – 3.60 (m, 2H)

^{19}F NMR (376 MHz, CD_3OD) δ -161.76 (s), -195.42 (d, $J = 55.4$ Hz)

MS $m/z = 328$ $[\text{M}+\text{H}]$. MS system: Thermo LCQ Fleet



Intermediate 11b - 2-fluoropyrrolo[2,1-f][1,2,4]triazin-4-amine.


Intermediate **11a** (2.0 g, 13.4 mmol) was charged to a polyTube vessel. The reaction vessel was then placed in an ice bath and both 70% HF/Pyr (18 mL) and pyridine (9 mL) were added sequentially. Immediately after the addition of pyridine, $t\text{BuNO}_2$ (2.07 mL, 17.43 mmol) was then added slowly over 20 min. The solution goes from tan to black with an exotherm and outgassing. The reaction was then stirred for an additional 20 min, and the reaction mixture was then diluted with water, and was concentrated under reduced pressure. The crude residue was partitioned between ethyl acetate and water. The organics were

separated and the aqueous was washed with ethyl acetate three times. The organics were combined and washed with brine. The crude was dried over Na₂SO₄, filtered, and the solvent was removed under reduced pressure. The crude residue was purified by silica gel chromatography (50-100% EtOAc/ Hex) to afford intermediate **11b** (1.68 g, 82%) as a tan solid.

¹H NMR (400 MHz, DMSO-*d*₆) δ 8.49 – 8.09 (m, 2H), 7.58 (t, *J* = 2.0 Hz, 1H), 6.95 (d, *J* = 4.5, 1H), 6.59 (d, *J* = 4.5, 1H).

¹⁹F NMR (376 MHz, DMSO-*d*₆) δ -73.42 (s).

LC/MS: *t*_R = 1.03 min, MS *m/z* = 153.08 [M+H]⁺; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ XB-C18 100A, 50 x 3.00 mm; Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN at 1.8mL/min.




Intermediate 11c - 7-bromo-2-fluoropyrrolo[2,1-f][1,2,4]triazin-4-amine.

A solution of **11b** (3.36 g, 22.1 mmol) in DMF (50 mL) was cooled to 0 °C in an ice bath. A solution of 1,3-dibromo-5,5-dimethylhydantoin (3.16 g, 11.0 mmol) in DMF (50 mL) was added dropwise by addition funnel over 40 min. After 1 h, the reaction was quenched with sat. Na₂S₂O_{3(aq)} and the crude was partitioned between EtOAc and 5% LiCl_(aq). The organics were extracted with 5% LiCl_(aq) (4 x) followed by brine. The organics were dried over Na₂SO₄, the solids were removed by filtration, and the filtrate was concentrated under reduced pressure. The crude residue was sonicated with CH₂Cl₂, and the solids were collected by filtration, and were dried under high vacuum to afford intermediate **11c** (3.93 g, 77%) as a yellow solid.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.49-8.44 (m, 2H), 7.08 (d, $J = 4.6$ Hz, 1H), 6.76 (d, $J = 4.6$ Hz, 1H).

^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ -71.45 (s).

LC/MS: $t_R = 1.22$ min, MS $m/z = 232.98$ $[\text{M}+\text{H}]^+$; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 3.00 mm; Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN at 1.8mL/min.




Intermediate 11e - (3R,4R,5R)-2-(4-amino-2-fluoropyrrolo[2,1-f][1,2,4]triazin-7-yl)-4-(benzyloxy)-5-((benzyloxy)methyl)-3-fluorotetrahydrofuran-2-ol.

To a solution of **11c** (2.09 g, 9.08 mmol) in THF (30 mL) was added 1,2-bis(chlorodimethylsilyl)ethane (STABASE, 1.96 g, 9.08 mmol) in one portion, and the resulting mixture was allowed to stir at ambient temperature for 1 h. The reaction was then cooled to $-78\text{ }^\circ\text{C}$ using a dry ice bath with methanol. $n\text{BuLi}$ (2.5M in hexanes, 10.9 mL, 27.2 mmol) was added in a manner to maintain an internal temperature of $-65\text{ }^\circ\text{C}$. A solution of intermediate **11d** (Prepared according to WO2012012776, 2.5 g, 7.5 mmol) in THF (25 mL) was then added to the reaction mixture over 1 min. After 5 min, the reaction mixture was quenched with acetic acid and was allowed to come to ambient temperature. The solvents were removed under reduced pressure and the residue was taken up into ethyl acetate. The organics were washed with water followed by brine. The layers were separated and the organics were dried over Na_2SO_4 , filtered, and concentrated under reduced pressure to afford crude **11e** as a mixture of isomers, which was used as is for the next step.

LC/MS: $t_R = 1.32$ and 1.40 min, MS $m/z = 483.15$ $[\text{M}+\text{H}]^+$; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 3.00 mm; Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0


min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN at 1.8mL/min.



Intermediate 11f - 7-((2S,3S,4R,5R)-4-(benzyloxy)-5-((benzyloxy)methyl)-3-fluorotetrahydrofuran-2-yl)-2-fluoropyrrolo[2,1-f][1,2,4]triazin-4-amine.

Intermediate **11e** (1.99 g, 3.72 mmol) was dissolved into CH_2Cl_2 (80 mL) and TES (4.75 mL, 29.7 mmol) was added to the mixture. The reaction mixture was cooled to 0 °C and $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (1.07 mL, 4.09 mmol) was slowly added. After 15 min, the reaction mixture was quenched with sat. $\text{NaHCO}_3(\text{aq})$ and the layers were separated. The aqueous layer was washed with CH_2Cl_2 . The organic layers were combined and were washed with sat. $\text{NaHCO}_3(\text{aq})$. The organics were dried over Na_2SO_4 , filtered, and were concentrated under reduced pressure. The crude was purified by silica gel chromatography (0-60% EtOAc/Hex) to afford intermediate **11f** (1.12 g, 64%, 2:1 mixture of 1' anomers).

LC/MS: $t_R = 1.55$ min, MS $m/z = 467.47$ $[\text{M}+\text{H}]^+$; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 3.00 mm; Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN at 1.8mL/min.




Intermediate 11g - (2R,3R,4R,5S)-5-(4-amino-2-fluoropyrrolo[2,1-f][1,2,4]triazin-7-yl)-4-fluoro-2-(hydroxymethyl)tetrahydrofuran-3-ol

Intermediate 11f (0.82 g, 1.76 mmol) was dissolved in acetic acid (25 mL). The reaction vessel was purged with Argon and 10% Pd/C (468 mg, 0.439 mmol) was added.

5 The vessel was evacuated and backfilled with $H_{2(g)}$ (3 x). After 1 h, the reaction vessel was purged with nitrogen. The resulting mixture was filtered through a pad of celite and the filter cake was washed with CH_3OH . The filtrate was concentrated under reduced pressure and then coevaporated with ethyl acetate followed by hexanes to afford intermediate 11g (503 mg, 98%, 2:1 mixture of 1' anomers).

10 LC/MS: t_R = 0.81 min, MS m/z = 286.97 [M+H]⁺; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 3.00 mm; Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN at 1.8mL/min.

15



Intermediate 11h - (2S,3R,4R,5S)-5-(4-amino-2-fluoropyrrolo[2,1-f][1,2,4]triazin-7-yl)-4-fluoro-2-(iodomethyl)tetrahydrofuran-3-ol.

To an argon purged flask was added a solution of 11g (283 mg, 0.989 mmol) in DMF (10 mL) followed by a solution of methyl triphenoxyposphonium iodide (0.536 g, 1.19 mmol) in 4mL DMF. The reaction mixture was allowed to stir at 0 °C for 10 min and was then warmed to ambient temperature. After 30 min, the reaction was quenched with sat.

Na₂S₂O_{3(aq)}. The crude material was partitioned between EtOAc and 5% LiCl_(aq). The organics were separated and washed with brine. The organics were dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The crude residue was taken up in ACN and purified by HPLC without acid modifier to afford intermediate 11h (201 mg, 52%, 2:1


20

25

mixture of 1' anomers) as a white solid.

LC/MS: t_R = 1.08 min, MS m/z = 397.12 $[M+H]^+$; LC system: Thermo Accela 1250 UHPLC;
MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 3.00 mm;

- 5 Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN at 1.8 mL/min.




- 10 **Intermediate 11i - (3R,4R,5S)-5-(4-amino-2-fluoropyrrolo[2,1-f][1,2,4]triazin-7-yl)-4-fluoro-2-methylenetetrahydrofuran-3-ol.**

To a solution of **11h** (356 mg, 0.899 mmol) in THF (8 mL) was added DBU (0.403 mL, 2.70 mmol) and the resulting mixture was heated to 60 °C. After 3 h, the reaction mixture was concentrated under reduced pressure. The crude residue was purified by silica gel chromatography (40-100% EtOAc/Hex) to afford intermediate **11i** (201 mg, 83%, 2:1 mixture of 1' anomers) as a white solid.

LC/MS: t_R = 1.04 min, MS m/z = 269.14 $[M+1]^+$; LC system: Thermo Accela 1250 UHPLC;
MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 3.00 mm;

- 20 Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN at 1.8 mL/min.




Intermediate 11j - (2S,3R,4R,5S)-5-(4-amino-2-fluoropyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-azido-4-fluoro-2-(iodomethyl)tetrahydrofuran-3-ol

5 Benzyltrimethylammonium chloride (292 mg, 1.57 mmol) and sodium azide (102 mg, 1.57 mmol) were dissolved in ACN (4 mL), and the resulting mixture was stirred at ambient temperature for 4 h. The mixture was filtered and the filtrate was added to a solution of **11i** (0.201 g, 0.749 mmol) in THF (4 mL). NMM (0.412 mL, 3.75 mmol) was added followed by the dropwise addition of a solution of iodine (0.342 g, 1.35 mmol) in THF (4 mL). After 15 min, N-acetyl cysteine was added portion wise until no outgassing was observed. Sat. $\text{Na}_2\text{S}_2\text{O}_3(\text{aq})$ was added until the solution was a light yellow. The resulting mixture was partitioned between water and ethyl acetate. The layers were separated and the organic layer was dried over Na_2SO_4 , filtered, and concentrated under reduced pressure. The crude residue was purified by silica gel chromatography (20-100% EtOAc/Hex) to afford intermediate **11j** (183 mg, 56%) as a single isomer.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.44 (d, J = 28.6 Hz, 2H), 6.98 (d, J = 4.6 Hz, 1H), 6.79 (d, J = 4.6 Hz, 1H), 6.32 (d, J = 6.9 Hz, 1H), 5.60 (dd, J = 23.8, 2.5 Hz, 1H), 5.36 (ddd, J = 54.9, 5.0, 2.6 Hz, 1H), 4.60 (ddd, J = 21.5, 6.9, 5.0 Hz, 1H), 3.63 (ABq, $\Delta\delta$ = 0.09 ppm, J = 8 Hz, 2H).

^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ -71.74 (s), -194.57 (ddd, J = 54.9, 24.0, 21.7 Hz)

LC/MS: t_R = 1.71 min, MS m/z = 437.93 [$M+1$]; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 3.00 mm; Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-2.4 min 2-100% ACN, 2.4 min-2.80 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3.0 min 2% ACN at 1.8 mL/min.



Intermediate 11k - (2S,3R,4S,5S)-5-(4-amino-2-fluoropyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-azido-4-fluoro-2-(iodomethyl)tetrahydrofuran-3-yl isobutyrate.

To a solution of **11j** (0.183 g, 0.419 mmol) in THF (10 mL) was added isobutyric anhydride (0.083 mL, 0.502 mmol), TEA (0.118 mL, 0.837 mmol), and DMAP (10 mg, 0.084 mmol). The reaction was allowed to stir at ambient temperature for 15 min, and the reaction was quenched with CH₃OH. The reaction mixture was concentrated under reduced pressure and the crude residue was purified by silica gel chromatography (0-50% EtOAc/Hex) to afford intermediate **11k** (0.198 g, 93%) as a white solid.


¹H NMR (400 MHz, DMSO-*d*₆) δ 8.48 (d, *J* = 30.7 Hz, 2H), 6.99 (d, *J* = 4.6 Hz, 1H), 6.84 (d, *J* = 4.6 Hz, 1H), 5.77 – 5.47 (m, 3H), 3.69 (ABq, Δδ = 0.05ppm, *J* = 12 Hz, 2H), 2.70 (p, *J* = 7.0 Hz, 1H), 1.24 – 1.05 (d, *J* = 7.0 Hz, 6H).

¹⁹F NMR (376 MHz, DMSO-*d*₆) δ -71.58 (s), -194.89 (ddd, *J* = 55.0, 24.3, 16.8 Hz).

LC/MS: *t*_R = 1.56 min, MS *m/z* = 508.13 [M+1]; LC system: Thermo Accela 1250 UHPLC;

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ XB-C18 100A, 50 x 3.00 mm;

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-2.4 min 2-100% ACN, 2.4 min-2.80 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3.0 min 2% ACN at 1.8mL/min.




Intermediate 11I - ((2R,3R,4S,5S)-5-(4-amino-2-fluoropyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-azido-4-fluoro-3-(isobutyryloxy)tetrahydrofuran-2-yl)methyl 3-chlorobenzoate.

Intermediate **11k** (0.153 g, 0.302 mmol) was dissolved in CH_2Cl_2 (10 mL) and H_2O (6 mL). Potassium phosphate dibasic (0.138 g, 0.603 mmol), tetrabutylammonium bisulfate (0.210 g, 0.618 mmol), and 3-chlorobenzoic acid (0.097 g, 0.618 mmol) were added sequentially. The resulting mixture was cooled to 0 °C and MCPBA (0.203 g, 0.905 mmol) was added. The reaction mixture was allowed to warm to ambient temperature and was stirred for 16 h. The reaction mixture was then quenched with sat $\text{Na}_2\text{S}_2\text{O}_3(\text{aq})$, and was concentrated under reduced pressure. The crude aqueous residue was diluted with ACN and purified by preparative HPLC without acid modifier to afford intermediate **11I** (20 mg, 13%) as a white solid.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.46 (d, J = 26.4 Hz, 2H), 7.96 – 7.81 (m, 2H), 7.81 – 7.66 (m, 1H), 7.62 – 7.46 (m, 1H), 6.94 (d, J = 4.5 Hz, 1H), 6.80 (d, J = 4.5 Hz, 1H), 5.73 (s, 3H), 4.60 (ABq, $\Delta\delta$ = 0.08 ppm, J = 12 Hz, 2H), 2.66 (p, J = 7.0 Hz, 1H), 1.20 – 1.01 (m, 6H).

^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ = -71.45 (s), -193.41 (ddd, J = 54.4, 25.4, 21.2 Hz.).

LC/MS: t_R = 2.22 min, MS m/z = 536.17 [$M+1$]; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 3.00 mm; Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-2.4 min 2-100% ACN, 2.4 min-2.80 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3.0 min 2% ACN at 1.8 mL/min.



Example 22 - (2R,3R,4R,5S)-5-(4-amino-2-fluoropyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-azido-4-fluoro-2-(hydroxymethyl)tetrahydrofuran-3-ol


To a solution of intermediate **11I** (22 mg, 0.041 mmol) in CH_3OH (1 mL) was added

conc. NH_4OH (1 mL) at RT. After 30 min, the reaction mixture was concentrated under reduced pressure. The crude residue was diluted with minimal H_2O and was purified by preparative HPLC without modifier to afford example **22** (10 mg, 77%) as a white solid.

- 5 ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.40 (d, $J = 30.9$ Hz, 2H), 6.95 (d, $J = 4.5$ Hz, 1H), 6.78 (d, $J = 4.5$ Hz, 1H), 5.89 (d, $J = 7.5$ Hz, 1H), 5.61 (dd, $J = 23.8, 2.1$ Hz, 1H), 5.44 (t, $J = 6.1$ Hz, 1H), 5.18 (ddd, $J = 55.3, 5.1, 2.2$ Hz, 1H), 4.44 (ddd, $J = 23.7, 7.5, 5.0$ Hz, 1H), 3.59 (ddd, $J = 48.5, 12.0, 6.1$ Hz, 2H).

^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ -71.18 (s), -193.48 (dt, $J = 55.3, 23.8$ Hz).

- 10 LC/MS: $t_R = 1.13$ min, MS $m/z = 327.86$ [$\text{M}+1$]; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex $2.6\mu\text{XB-C18}$ 100A, 50×3.00 mm; Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-2.4 min 2-100% ACN, 2.4 min-2.80 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3.0 min 2% ACN at 1.8mL/min.



Intermediate 12a - (2R,3R,4S,5S)-5-(4-amino-5-fluoropyrrolo[2,1-f][1,2,4]triazin-7-yl)-3-(((tert-butyldimethylsilyl)oxy)-2-(((tert-butyldimethylsilyl)oxy)methyl)-4-fluorotetrahydrofuran-2-carbonitrile


- 20 To a solution of intermediate **3d** (57 mg, 0.109 mmol) in ACN (3 mL), was added Selectfluor II in one portion (52 mg, 0.164 mmol). After 1.5 h, the reaction was quenched by the addition of saturated $\text{NaHCO}_{3(\text{aq})}$. Ethyl acetate (4mL) was added and the biphasic mixture was stirred vigorously for 5 min. The reaction was further diluted with EtOAc and saturated $\text{NaHCO}_{3(\text{aq})}$. The layers were separated and the organic phase was extracted with
- 25 water and then brine. The organic layer was dried over Na_2SO_4 . The drying agent was removed by vacuum filtration and the filtrate was concentrated under reduced pressure. Intermediate **12a** (11 mg, 18.7 %) was isolated from the concentrated crude material by silica gel column chromatography using the following solvent ramp: 0% EtOAc in hexanes

ramping to 70% EtOAc in hexanes, quickly ramping to 100% EtOAc once the starting material elutes off the column.

^1H NMR (400 MHz, CD_3OD) δ 7.72 (s, 1H), 7.55 (s, 1H), 5.65 (dd, J = 24.8, 2.4 Hz, 1H), 5.38 (dq, J = 54.4, 2 Hz, 1H), 4.88 (dd, J = 19.2, 4.4 Hz, 1H), 3.96 (ABq, $\Delta\delta_{\text{AB}}$ = 0.141ppm, J = 11 Hz, 2H), 0.99 (s, 9H), 0.86 (s, 9H), 0.21 (s, 6H), 0.07 (s, 3H), -0.02 (s, 3H).

^{19}F NMR (376 MHz, CD_3OD) δ -161.795 (s), -194.806 (ddd, J = 54.5, 19.2, 18.8 Hz).

LC/MS: R_T = 2.06 min, MS m/z = 540.64 [$M+1$]; LC system: Thermo Accela 1250 UHPLC; MS: Thermo LCQ Fleet; Column: Kinetex 2.6 μ C18 100A, 50 x 3.00 mm; Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-2.4 min 2-100% ACN, 2.4 min-2.8 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3 min 2% ACN.



Example 23 - (2R,3R,4R,5S)-5-(4-amino-5-fluoropyrrolo[2,1-f][1,2,4]triazin-7-yl)-4-fluoro-3-hydroxy-2-(hydroxymethyl)tetrahydrofuran-2-carbonitrile

To a solution of intermediate **12a** (29 mg, 0.054 mmol) in THF (2 mL) in a polypropylene tube was added 70% HF-Pyridine in pyridine (51 μL , 1.97 mmol) at 0 $^{\circ}\text{C}$, under an N_2 atmosphere. After 1.5 h, the reaction was removed from the ice bath. Additional 70% HF-Pyridine in pyridine was added after 3 h (150 μL), 5 h 45 min (200 μL), and 21 h 15 min (0.7 mL). The reaction mixture was then stirred for another 24 h at which point the reaction mixture was cooled in an ice bath and then quenched with water and saturated $\text{NaHCO}_{3(\text{aq})}$. The mixture was concentrated under reduced pressure and the residue was taken up in DMF. The resulting solution/suspension was filtered through a syringe filter (Whatman 0.45 μm PTFE w/GMF). The filtrate was injected on an HPLC and the semi-purified product was further purified by silica gel column chromatography using the following

solvent ramp: 0% MeOH in DCM ramping to 20% MeOH in DCM. Product containing fractions were concentrated and the residue was lyophilized to afford example **23** (5 mg, 30%) as a white powder.


¹H NMR (400 MHz, DMF-*d*₇) δ 7.74 (s, 1H), 6.61 (s, 1H), 5.75 (dd, *J* = 25.2, 1.6 Hz, 1H), 5.23 (ddd, *J* = 54.8, 4.8, 1.6 Hz, 1H), 4.64 (dd, *J* = 22, 4.4 Hz, 1H), 3.90 (ABq, Δδ_{AB} = 0.151 ppm, *J* = 12 Hz, 2H).

¹⁹F NMR (376 MHz, DMF-*d*₇) δ -161.727 (s), -193.726 (ddd, *J* = 54.5, 22.9, 21.8 Hz).

10

LC/MS: R_T = 0.81 min, MS *m/z* = 312.13 [M+1]; LC: Thermo Accela 1250 UHPLC; MS: Thermo LCQ Fleet; Column: Kinetex 2.6μ C18 100A, 50 x 3.00 mm; Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-2.4 min 2-100% ACN, 2.4 min-2.8 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3 min 2% ACN.

15




Intermediate 13a - N-(7-((2S,3S,4R,5R)-4-((tert-butyldimethylsilyl)oxy)-5-(((tert-butyldimethylsilyl)oxy)methyl)-3-fluoro-5-(iodomethyl)tetrahydrofuran-2-yl)pyrrolo[2,1-f][1,2,4]triazin-4-yl)benzamide.

To a solution of triphenylphosphine (973 mg, 3.71 mmol) and imidazole (252 mg, 3.71 mmol) in THF (5 mL) was added iodine (253 mg, 1.86 mmol) at room temperature. Upon complete dissolution of iodine, a solution of compound **2g** (650 mg, 0.93 mmol) in THF (5 mL) was added dropwise slowly. The resulting mixture was stirred at 80 °C for 3 days and was then concentrated *in vacuo*. The residue was purified by silica gel column chromatography (0 to 50% EtOAc in hexanes) to give intermediate **13a** (230 mg, 33%) as an oil.

25


MS *m/z* = 742 [M+H]. MS system: Thermo LCQ Fleet.



Intermediate 13b - N-(7-((2S,3S,4R,5R)-4-((tert-butyldimethylsilyl)oxy)-5-(((tert-butyldimethylsilyl)oxy)methyl)-3-fluoro-5-methyltetrahydrofuran-2-yl)pyrrolo[2,1-f][1,2,4]triazin-4-yl)benzamide.

Intermediate **13a** (200 mg, 0.243 mmol) was dissolved in methanol (10 mL) and under nitrogen atmosphere, 10% Pd/C (100 mg, 0.094 mmol) and TEA (0.035 mL, 0.243 mmol) were added. The resulting mixture was then stirred under H₂ atmosphere (balloon) at room temperature for 40 min. The resulting mixture was filtered, and concentrated *in vacuo* and the residue was purified by silica gel column chromatography (0 to 40% EtOAc in hexanes) to give intermediate **13b** (145 mg, 72%) as a white solid with 75% purity.

MS *m/z* = 616 [M+H]. MS system: Thermo LCQ Fleet



Example 24 - (2R,3R,4R,5S)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-4-fluoro-2-(hydroxymethyl)-2-methyltetrahydrofuran-3-ol.


Intermediate **13b** (145 mg, 75% purity, 0.177 mmol) was dissolved in THF (10 mL) and TBAF (1M in THF, 0.53 mL, 0.531 mmol) was added. The resulting mixture was stirred at room temperature for 2 h and then methanolic ammonia (7N, 10 mL) was added. The resulting mixture was stirred for 24 h and was concentrated under reduced pressure. The

crude residue was purified by preparative HPLC (0 to 35% acetonitrile in water in 20 min) to afford example **24** (30 mg, 60%) as a white solid.

^1H NMR (400 MHz, CD_3OD) δ 7.78 (s, 1H), 6.84 (d, J = 4.5 Hz, 1H), 6.76 (d, J = 4.5 Hz, 1H), 5.53 (dd, J = 21.5, 4.0 Hz, 1H), 5.25 (ddd, J = 55.5, 5.3, 4.1 Hz, 1H), 4.44 (dd, J = 17.2, 5.2 Hz, 1H), 3.65 – 3.43 (m, 2H), 1.27 (s, 3H)

^{19}F NMR (376 MHz, CD_3OD) δ -197.08 (ddd, J = 55.4, 21.5, 17.1 Hz)

MS m/z = 282 $[\text{M}+\text{H}]$. MS system: Thermo LCQ Fleet.



Intermediate 14a - (2S,3R,4S,5R)-2-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-5-(hydroxymethyl)tetrahydrofuran-3,4-diol.

Intermediate **1e** (2.64 g, 4.91 mmol) was dissolved in acetic acid (50 mL). The flask was purged with Argon and 10% Pd/C (1.05 g, 0.982 mmol) was added. The flask was evacuated and backfilled with $\text{H}_{2(g)}$ three times. The reaction mixture was stirred under an atmosphere of $\text{H}_{2(g)}$. After 1h, the flask was purged with nitrogen and the reaction mixture was filtered through a celite pad with CH_3OH washings. The filtrate was concentrated under reduced pressure and then coevaporated with EtOAc followed by hexanes. The residue was placed under high vacuum to afford intermediate **14a** (1.31 g, 99%) as a white solid.


^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.80 (s, 1H), 7.66 (s, 2H), 6.82 (d, J = 4.4 Hz, 1H), 6.66 (d, J = 4.4 Hz, 1H), 5.09 (d, J = 6.5 Hz, 1H), 5.06-4.56 (m, 3H), 4.21 (t, J = 5.9 Hz, 1H), 3.93 (t, J = 4.9 Hz, 1H), 3.77 (q, J = 4.5 Hz, 1H), 3.48 (ddd, J = 38.9, 11.8, 4.4 Hz, 2H).

LC/MS: t_R = 0.47 min, MS m/z = 267.13 $[\text{M}+\text{H}]$; LC system: Thermo Accela 1250 UHPLC

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 3.00 mm;

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-

2.4 min 2-100% ACN, 2.4 min-2.80 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3.0 min 2% ACN at 1.8mL/min.




Intermediate 14b - ((3aR,4R,6S,6aS)-6-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-2,2-dimethyltetrahydrofuro[3,4-d][1,3]dioxol-4-yl)methanol.

Intermediate **14a** (3.13 g, 11.7 mmol) was dissolved in acetone (80 mL) and TsOH (6.00 g, 31.5 mmol) was added. Triethylorthoformate (6.0 mL, 36.1 mmol) was slowly added over 10 min. The resulting mixture was allowed to stir at ambient temperature overnight. Saturated aqueous sodium carbonate solution was added until the reaction mixture was pH=8. The solids were removed by filtration and the filtrate was concentrated under reduced pressure. The crude residue was partitioned between EtOAc and brine. The phases were split and the organics were dried over Na₂SO₄, were filtered, and were concentrated under reduced pressure. The crude was purified by silica gel chromatography (60-100% EtOAc/Hex - 20% MeOH/EtOAc) to afford intermediate **14b** (2.55 g, 71%) as a white solid.

¹H NMR (400 MHz, DMSO-*d*₆) δ 7.83 (s, 1H), 7.71 (s, 2H), 6.83 (d, *J* = 4.4 Hz, 1H), 6.73 (d, *J* = 4.5 Hz, 1H), 5.21 (d, *J* = 4.9 Hz, 1H), 5.01 (dd, *J* = 6.6, 4.9 Hz, 1H), 4.84 (t, *J* = 5.7 Hz, 1H), 4.71 (dd, *J* = 6.7, 3.7 Hz, 1H), 3.99 – 3.85 (m, 1H), 3.46 (t, *J* = 5.5 Hz, 2H), 1.48 (s, 3H), 1.29 (s, 3H).

LC/MS: *t*_R = 0.87 min, MS *m/z* = 307.21 [M+H]⁺; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ XB-C18 100A, 50 x 3.00 mm; Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN at 1.8mL/min.




Intermediate 14c - 7-((3a*S*,4*S*,6*R*,6a*R*)-6-(((tert-butyl dimethylsilyl)oxy)methyl)-2,2-dimethyltetrahydrofuro[3,4-*d*][1,3]dioxol-4-yl)pyrrolo[2,1-*f*][1,2,4]triazin-4-amine.

Intermediate **14b** (2.55 g, 8.32 mmol) was dissolved in DCM (50 mL) and the mixture
 5 was cooled to 0 °C. Imidazole (1.70 g, 24.9 mmol) was added followed by TBSCl (1.88 g, 12.5 mmol). After 16 h, the reaction was quenched with methanol. The resulting mixture was concentrated under reduced pressure and the crude residue was partitioned between water and EtOAc. The organics were dried over Na₂SO₄, were filtered, and were concentrated under reduced pressure. The crude residue was purified by silica gel
 10 chromatography (50-100% EtOAc/Hex) to afford intermediate **14c** (2.60 g, 74%) as a white solid.

¹H NMR (400 MHz, DMSO-*d*₆) δ 7.83 (s, 1H), 7.74 (s, 2H), 6.82 (d, *J* = 4.4 Hz, 1H), 6.68 (d, *J* = 4.4 Hz, 1H), 5.26 (d, *J* = 4.4 Hz, 1H), 5.00 (dd, *J* = 6.5, 4.5 Hz, 1H), 4.71 (dd, *J* = 6.5, 3.7
 15 Hz, 1H), 3.97 (td, *J* = 5.1, 3.6 Hz, 1H), 3.64 (d, *J* = 5.2 Hz, 2H), 1.48 (s, 3H), 1.28 (s, 3H), 0.83 (s, 9H), -0.02 (s, 6H).

LC/MS: *t*_R = 1.91 min, MS *m/z* = 421.60 [M+H]⁺; LC system: Thermo Accela 1250 UHPLC
 MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ XB-C18 100A, 50 x 3.00 mm;
 20 Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-2.4 min 2-100% ACN, 2.4 min-2.80 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3.0 min 2% ACN at 1.8mL/min.



Intermediate 14d - tert-butyl (7-((3aS,4S,6R,6aR)-6-(((tert-butyl)dimethylsilyl)oxy)methyl)-2,2-dimethyltetrahydrofuro[3,4-d][1,3]dioxol-4-yl)pyrrolo[2,1-f][1,2,4]triazin-4-yl)carbamate.


5 Intermediate **14c** (2.59 g, 6.16 mmol) was dissolved in THF (60 mL) and the resulting solution was cooled to 0 °C. Boc_2O (2.69 g, 12.3 mmol) and DMAP (0.3 g, 2.46 mmol) were then added. TEA (2.56 mL, 18.3 mmol) was slowly added and the reaction mixture was allowed to warm to RT. After 3 h, the reaction mixture was cooled to 0 °C and MeOH (10 mL) was added followed conc. $\text{NH}_4\text{OH}_{(\text{aq})}$ (50 mL). The resulting mixture was allowed to
 10 warm to RT and was stirred overnight. The reaction mixture was concentrated under reduced pressure and the crude residue was partitioned between EtOAc and water. The layers were separated and the organic layer was dried over Na_2SO_4 , was filtered, and was concentrated under reduced pressure. The crude residue was purified by silica gel chromatography (0-100% EtOAc/Hex) to afford intermediate **14d** (2.82 g, 88%) as a white
 15 solid.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.46 (s, 1H), 8.20 (s, 1H), 7.19 (d, $J = 4.6$ Hz, 1H), 6.90 (d, $J = 4.6$ Hz, 1H), 5.33 (d, $J = 4.2$ Hz, 1H), 5.02 (dd, $J = 6.5, 4.3$ Hz, 1H), 4.72 (dd, $J = 6.5, 3.6$ Hz, 1H), 4.01 (q, $J = 5.0$ Hz, 1H), 3.64 (d, $J = 5.1$ Hz, 2H), 1.49 (s, 9H), 1.32 (d, $J = 22.7$ Hz, 6H), 0.82 (s, 9H), -0.03 (s, 6H).
 20

LC/MS: $t_R = 1.89$ min, MS $m/z = 521.27$ $[\text{M}+\text{H}]^+$; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 3.00 mm; Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-2.4 min 2-100% ACN, 2.4 min-2.80 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3.0 min 2% ACN at 1.8mL/min.
 25

Intermediate **14d** (2.8 g, 5.4 mmol) was dissolved in THF (50 mL), and TBAF (1.0M in THF, 5.92 mL, 5.92 mmol) was added. After 30 min, additional TBAF (1.0M in THF, 5.92 mL, 5.92 mmol) was added. After another 30 min, the reaction mixture was quenched with water and the resulting mixture was extracted with EtOAc (2 x). The combine the organic layers were washed with brine, were dried over Na₂SO₄, were filtered, and were concentrated under reduced pressure. The crude residue was purified by silica gel chromatography (10-100% EtOAc/Hex) to afford intermediate **14e** (2.19 g, 86%) as a white solid.

LC/MS: t_R = 1.52 min, MS m/z = 407.05 [M+H]⁺; LC system: Thermo Accela 1250 UHPLC;
MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 3.00 mm;
Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-
2.4 min 2-100% ACN, 2.4 min-2.80 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85
min-3.0 min 2% ACN at 1.8mL/min.



Intermediate 14f - tert-butyl (7-((3aS,4S,6aS)-6,6-bis(hydroxymethyl)-2,2-dimethyltetrahydrofuro[3,4-d][1,3]dioxol-4-yl)pyrrolo[2,1-f][1,2,4]triazin-4-yl)carbamate.

5 Intermediate **14e** (1.78 g, 4.38 mmol) was dissolved in DMSO (20 mL) and toluene (15mL). Pyridine (0.35 mL, 4.38 mmol) and EDCI (1.26 g, 6.56 mmol) were added followed by TFA (0.178 mL, 2.39 mmol). After 90 min, additional pyridine (0.35 mL, 4.38 mmol) and EDCI (1.26 g, 6.56 mmol) were added and the reaction mixture was stirred for an additional 30 min. The reaction was quenched with water and the resulting mixture was extracted with
10 CH_2Cl_2 . The aqueous was back extracted with CH_2Cl_2 . The organic layers were combined and were washed with brine, were dried over Na_2SO_4 , were filtered, and were concentrated under reduced pressure. The crude was put under high vacuum for 15 min then was used as is for the next step.

The crude residue was dissolved in dioxane (15 mL) and formaldehyde (37% in
15 water, 5.0 mL, 37.2 mmol) and 2N NaOH (5.34 mL, 10.7 mmol) were added sequentially. After 10 min, the reaction was quenched with AcOH and the resulting mixture was partitioned between sat. $\text{NaHCO}_3(\text{aq})$ and CH_2Cl_2 . The aqueous layer was back extracted with CH_2Cl_2 . The organic layers were combined and were wash with brine, were dried over Na_2SO_4 , were filtered, and were concentrated under reduced pressure. The crude
20 was placed under high vacuum for 15 min then were taken directly into the next reaction.


The crude residue was dissolved in EtOH (50 mL), and NaBH_4 (0.324 g, 8.76 mmol) was added in small portions. After 20 min, the reaction mixture was quenched with AcOH and was concentrated under reduced pressure. The crude residue was partitioned between with EtOAc and sat. $\text{NaHCO}_3(\text{aq})$. The organic layer was split, was dried over Na_2SO_4 , was
25 filtered, and was concentrated under reduced pressure. The crude residue was purified by silica gel chromatography (50-100% EtOAc/Hex) to afford intermediate **14f** (1.91 g, 68%) as a white solid.

¹H NMR (400 MHz, DMSO-*d*₆) δ 10.45 (s, 1H), 8.20 (s, 1H), 7.19 (d, *J* = 4.3 Hz, 1H), 6.95 (d, *J* = 4.7 Hz, 1H), 5.35 (d, *J* = 5.2 Hz, 1H), 5.06 (t, *J* = 5.7 Hz, 1H), 4.79-4.74 (m, 2H), 4.45 (t, *J* = 5.8 Hz, 1H), 3.73 – 3.46 (m, 3H), 3.40 – 3.30 (m, 1H), 1.50 (s, 12H), 1.27 (s, 3H).

5

LC/MS: *t*_R = 1.45 min, MS *m/z* = 437.09 [M+H]⁺; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ XB-C18 100A, 50 x 3.00 mm; Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-2.4 min 2-100% ACN, 2.4 min-2.80 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3.0 min 2% ACN at 1.8mL/min.

10



Intermediate 14g - tert-butyl (7-((3a*S*,4*S*,6*S*,6a*S*)-6-((bis(4-methoxyphenyl)(phenyl)methoxy)methyl)-6-(hydroxymethyl)-2,2-dimethyltetrahydrofuro[3,4-*d*][1,3]dioxol-4-yl)pyrrolo[2,1-*f*][1,2,4]triazin-4-yl)carbamate.

15

Intermediate **14f** (1.15 g, 2.63 mmol) was dissolved in CH₂Cl₂ (50 mL) and TEA (0.73 mL, 5.27 mmol) was added. The resulting solution was cooled to 0 °C and DMTrCl (1.35 g, 3.95 mmol) was added. After 10 min, the reaction mixture was quenched with CH₃OH as was then diluted with CH₂Cl₂. The resulting mixture was washed with sat NaHCO_{3(aq)} and brine. The organic layer was dried over Na₂SO₄, was filtered, and was concentrated under reduced pressure. The crude residue was purified by silica gel chromatography (0-100% EtOAc/Hex) to afford **14g** (1.95 g, 79%) as an off-white solid.

20


25

¹H NMR (400 MHz, DMSO-*d*₆) δ 10.46 (s, 1H), 8.23 (s, 1H), 7.56 – 7.07 (m, 10H), 7.07 –

6.70 (m, 5H), 5.24 (d, $J = 5.2$ Hz, 1H), 5.04 (t, $J = 5.9$ Hz, 1H), 4.93 – 4.71 (m, 2H), 3.80 – 3.59 (m, 7H), 3.52 (dd, $J = 10.9, 4.8$ Hz, 1H), 3.25 (d, $J = 9.9$ Hz, 1H), 3.09 (d, $J = 9.9$ Hz, 1H), 1.50 (s, 9H), 1.25 (s, 3H), 1.21 (s, 3H).

- 5 LC/MS: $t_R = 2.54$ min, MS $m/z = 739.28$ [M+H]⁺; LC system: Thermo Accela 1250 UHPLC MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 3.00 mm; Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-2.4 min 2-100% ACN, 2.4 min-2.80 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3.0 min 2% ACN at 1.8mL/min.

10



Intermediate 14h - tert-butyl (7-((3aS,4S,6R,6aS)-6-((bis(4-methoxyphenyl)(phenyl)methoxy)methyl)-6-(((tert-butyldimethylsilyl)oxy)methyl)-2,2-dimethyltetrahydrofuro[3,4-d][1,3]dioxol-4-yl)pyrrolo[2,1-f][1,2,4]triazin-4-yl)carbamate.

15

Intermediate **14g** (1.53 g, 2.08 mmol) was dissolved in DMF (10 mL) and imidazole (0.42 g, 6.23 mmol) was added followed by TBSCl (0.47 g, 3.11 mmol). After 1 h, the reaction was quenched with methanol and partitioned between EtOAc and 5% LiCl_(aq). The phases were split and the organic layer was washed with brine, was dried over Na₂SO₄, was filtered, and was concentrated under reduced pressure. The crude residue was purified by silica gel chromatography (0-50% EtOAc/Hex) to afford **14h** (1.77 g, 78%) as an off-white solid.


20

¹H NMR (400 MHz, DMSO-*d*₆) δ 10.47 (s, 1H), 8.23 (s, 1H), 7.56 – 6.66 (m, 15H), 5.31 (d, $J = 4.9$ Hz, 1H), 5.14 (dd, $J = 6.5, 4.9$ Hz, 1H), 4.73 (d, $J = 6.5$ Hz, 1H), 3.87 (d, $J = 9.7$ Hz, 1H), 3.72 (s, 6H), 3.53 (d, $J = 9.7$ Hz, 1H), 3.31 (m, 1H), 3.08 (d, $J = 9.8$ Hz, 1H), 1.50 (s,

25

MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 3.00 mm;
 Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-
 1.0 min 2-100% ACN, 1.0 min-2.80 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85
 min-3.0 min 2% ACN at 1.8mL/min.

5



Intermediate 14j - tert-butyl (7-((3aS,4S,6R,6aS)-6-(((tert-butylidimethylsilyl)oxy)methyl)-6-cyano-2,2-dimethyltetrahydrofuro[3,4-d][1,3]dioxol-4-yl)pyrrolo[2,1-f][1,2,4]triazin-4-yl)carbamate.

10

Intermediate **14i** (0.838 g, 1.52 mmol) was dissolved in DMSO (5 mL) and toluene (3 mL). Pyridine (0.14 mL, 1.67 mmol) and EDCI (0.438 g, 2.28 mmol) were added followed by TFA (0.057 mL, 0.761 mmol). After 30 min, additional pyridine (0.14 mL, 1.67 mmol) and EDCI (0.438 g, 2.28 mmol) were added. After 1 h, additional pyridine (0.14 mL, 1.67 mmol) and EDCI (0.438 g, 2.28 mmol) were added. After 2 h, the reaction mixture was quenched with $\frac{1}{2}$ sat. NaHCO_{3(aq)} and was partitioned between EtOAc and $\frac{1}{2}$ sat. NaHCO_{3(aq)}. The layers were separated and the organic layer was dried over Na₂SO₄, was filtered, and was concentrated under reduced pressure. The residue was dissolved in CH₂Cl₂ was concentrated under high vacuum for 1 h to afford a residue that was used directly in the next step.

15

20


The residue was dissolved in pyridine (8 mL) and hydroxylamine hydrochloride (0.159 g, 2.28 mmol) was added in one portion. After 15 min, the reaction mixture was concentrated under reduced pressure and was partitioned between EtOAc and water. The organic layer was dried over Na₂SO₄, was filtered, and was concentrated under reduced pressure. The crude residue was placed under high vacuum for 30 min and used as is for the third step.

25

The crude residue was dissolved in ACN (8 mL). CDI (0.37 g, 2.28 mmol) was added in one portion. After 45 min, additional CDI (0.37 g, 2.28 mmol) was added. After 1 h, the reaction was quenched with $\frac{1}{2}$ sat. $\text{NaHCO}_{3(\text{aq})}$. The crude was partitioned between EtOAc and $\frac{1}{2}$ sat. $\text{NaHCO}_{3(\text{aq})}$. The layers were separated and the organic layer was dried over Na_2SO_4 , was filtered, and was concentrated under reduced pressure. The crude residue was purified by silica gel chromatography (0-50% EtOAc/Hex) to afford intermediate **14j** (0.72 g, 87%) as a white solid.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.53 (s, 1H), 8.25 (s, 1H), 7.21 (s, 1H), 7.00 (d, $J = 4.6$ Hz, 1H), 5.62 (d, $J = 3.6$ Hz, 1H), 5.28 (dd, $J = 6.6, 3.7$ Hz, 1H), 4.93 (d, $J = 6.6$ Hz, 1H), 3.83 (s, 2H), 1.62 (s, 3H), 1.50 (s, 9H), 1.33 (s, 3H), 0.83 (s, 9H), 0.00 (s, 6H).

LC/MS: $t_R = 2.50$ min, MS $m/z = 546.15$ $[\text{M}+\text{H}]$; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 3.00 mm; Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-2.4 min 2-100% ACN, 2.4 min-2.80 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3.0 min 2% ACN at 1.8mL/min.




Intermediate 14k - (3aS,4R,6S,6aS)-6-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-4-(((tert-butyldimethylsilyl)oxy)methyl)-2,2-dimethyltetrahydrofuro[3,4-d][1,3]dioxole-4-carbonitrile.

Intermediate **14j** (0.688 g, 1.26 mmol) was dissolved in CH_2Cl_2 (15 mL). Zinc bromide (0.567 g, 2.52 mmol) was added in one portion and the reaction mixture was stirred at ambient temperature. After 3 h, the reaction mixture was added to a silica load cartridge and was purified by silica gel chromatography (40-100% EtOAc/Hex) to afford intermediate **14k** (0.56 g, 99%) as a white solid.

10


138



Intermediate **14l** (80 mg, 0.164 mmol) was dissolved in DMF (2 mL) and NBS (29 mg, 0.164 mmol) was added in one portion. After 45 min, the reaction was diluted with methanol. The solvent was removed under reduced pressure. The crude residue was purified by silica gel chromatography (0-50% EtOAc/Hex) to afford intermediate **14m** (50 mg, 54%) as an off-white solid.

¹H NMR (400 MHz, DMSO-*d*₆) δ 10.13 (s, 1H), 8.42 (s, 1H), 7.29 (s, 1H), 5.65 (d, *J* = 3.2 Hz, 1H), 5.29 (dd, *J* = 6.6, 3.2 Hz, 1H), 4.91 (d, *J* = 6.5 Hz, 1H), 3.84 (d, *J* = 1.6 Hz, 2H), 2.27 (s, 3H), 1.62 (s, 3H), 1.33 (s, 3H), 0.83 (s, 9H), 0.02 (s, 3H), 0.00 (s, 3H).

LC/MS: t_R = 1.79 min, MS m/z = 566.40 [M+1]; LC system: Thermo Accela 1250 UHPLC;
MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 3.00 mm;
Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-
1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85
min-2 min 2% ACN at 1.8mL/min.




Intermediate 14n - N-(7-((3a*S*,4*S*,6*R*,6a*S*)-6-(((*tert*-butyldimethylsilyl)oxy)methyl)-6-cyano-2,2-dimethyltetrahydrofuro[3,4-*d*][1,3]dioxol-4-yl)-5-fluoropyrrolo[2,1-*f*][1,2,4]triazin-4-yl)acetamide.

5 Intermediate **14m** (50 mg, 0.088 mmol) was dissolved in THF (2 mL) and the solution was cooled to -78 °C. *n*BuLi (2.5M in hexanes, 0.071 mL, 0.18 mmol) was added. After 5 min, *N*-fluorobenzenesulfonimide (NFSI, 33.4 mg, 0.106 mmol) was added and the reaction mixture was stirred for 5 min. The reaction was then quenched with AcOH. The solvents were removed under reduced pressure. The crude residue was purified by reverse phase HPLC to afford intermediate **14n** (10 mg, 22%) as a white solid.

¹H NMR (400 MHz, Methanol-*d*₄) δ 8.13 (s, 1H), 6.80 (s, 1H), 5.65 (d, *J* = 3.5 Hz, 1H), 5.23 (dd, *J* = 6.7, 3.6 Hz, 1H), 4.97 (d, *J* = 6.7 Hz, 1H), 3.92 (d, *J* = 1.7 Hz, 2H), 2.37 (s, 3H), 1.70 (s, 3H), 1.38 (s, 3H), 0.90 (s, 9H), 0.08 (s, 6H).

¹⁹F NMR (376 MHz, Methanol-*d*₄) δ -156.43 (s).

LC/MS: *t*_R = 1.65 min, MS *m/z* = 506.18 [M+H]⁺; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6μ XB-C18 100A, 50 x 3.00 mm; Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-2.4 min 2-100% ACN, 2.4 min-2.80 min 100% ACN, 2.8 min-2.85 min 100%-2% ACN, 2.85 min-3.0 min 2% ACN at 1.8mL/min.




Example 25 - (2R,3S,4R,5S)-5-(4-amino-5-fluoropyrrolo[2,1-f][1,2,4]triazin-7-yl)-3,4-dihydroxy-2-(hydroxymethyl)tetrahydrofuran-2-carbonitrile.

Intermediate **14n** (11 mg, 0.022 mmol) was taken up into a solution of 50% TFA in water at ambient temperature. After 2 h, the reaction mixture was quenched with solid Na_2CO_3 to achieve a pH = 8. The solvent was removed under reduced pressure and the crude residue was purified by reverse phase HPLC. The fractions containing example **25** were combined and set aside and the fractions containing the N6-Acyl were combined and concentrated under reduced pressure. The N6-Acyl intermediate residue was taken up in conc. $\text{NH}_4\text{OH}_{(\text{aq})}$ (1 mL) and the mixture was allowed to stir at ambient temperature. After 30 min, the resulting mixture was concentrated under reduced pressure and the crude residue was purified by HPLC. The fractions containing example **25** were combined with the previously set aside fractions containing example **25** to afford example **25** (4 mg, 53%) as a white solid.

^1H NMR (400 MHz, Methanol- d_6) δ 7.71 (s, 1H), 6.56 (s, 1H), 5.44 (d, J = 5.6 Hz, 1H), 4.48 (t, J = 5.6 Hz, 1H), 4.36 (d, J = 5.5 Hz, 1H), 3.83 ((ABq, $\Delta\delta$ = 0.05ppm, J = 12 Hz, 2H).

^{19}F NMR (376 MHz, Methanol- d_4) δ -161.81 (s).


LC/MS: t_R = 0.47 min, MS m/z = 310.13 $[\text{M}+\text{H}]^+$; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 3.00 mm; Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid; Gradient: 0 min-1.4 min 2-100% ACN, 1.4 min-1.80 min 100% ACN, 1.8 min-1.85 min 100%-2% ACN, 1.85 min-2 min 2% ACN at 1.8mL/min.



Intermediate 15a - tert-butyl (7-((3aS,4S,6R,6aS)-6-(((tert-butylidimethylsilyl)oxy)methyl)-6-(chloromethyl)-2,2-dimethyltetrahydrofuro[3,4-d][1,3]dioxol-4-yl)pyrrolo[2,1-f][1,2,4]triazin-4-yl)carbamate.

Intermediate **14i** (100 mg, 0.18 mmol) was dissolved in anhydrous pyridine (5 mL). Trifluoromethanesulfonyl chloride (23 μ L, 0.22 mmol) was added in one portion and the reaction mixture was stirred for 45 min at RT. Additional trifluoromethanesulfonyl chloride (100 μ L) was then added. After 30 min, additional trifluoromethanesulfonyl chloride (100 μ L) was added. After an additional 30 min, more trifluoromethanesulfonyl chloride (100 μ L) was added and the reaction was stirred for 30 min, at which point the reaction mixture was concentrated under reduced pressure. The crude residue was dissolved in anhydrous DMF (5 mL) and lithium chloride (153 mg, 3.6 mmol) was then added in one portion. The resulting mixture was stirred at RT for 16 h. The reaction mixture was diluted with ethyl acetate (50 mL) and was washed with saturated aqueous sodium chloride solution (3 x 20 mL). The organic layer was dried over anhydrous sodium sulfate and was concentrated under reduced pressure. The crude residue was purified with silica gel chromatography (0-20% ethyl acetate in hexanes) to afford intermediate **15a**.

MS m/z = 569.0 [M+H]. MS system: Thermo LCQ Advantage




Example 26 - (2R,3S,4R,5S)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-(chloromethyl)-2-(hydroxymethyl)tetrahydrofuran-3,4-diol.

Intermediate **15a** was dissolved in a solution of TFA and water (1:1, 5 mL) and the resulting mixture was stirred for 16 h. The reaction mixture was then concentrated under reduced pressure. The crude residue was dissolved in aqueous sodium bicarbonate solution and acetonitrile and was purified with prep HPLC to afford example **26** (19 mg, 34%) as white powder.

^1H NMR (400 MHz, D_2O) δ 7.61 (s, 1H), 6.72 – 6.64 (m, 2H), 5.19 (d, J = 9.1 Hz, 1H), 4.73 – 4.66 (m, 1H), 4.28 (d, J = 5.2 Hz, 1H), 3.78 (s, 2H), 3.72 – 3.57 (m, 2H).

MS m/z = 315.3 $[\text{M}+\text{H}]$. MS system: Thermo LCQ Advantage




Example 27 (also TP7) - ((2R,3R,4R,5S)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-azido-4-fluoro-3-hydroxytetrahydrofuran-2-yl)methyl tetrahydrogen triphosphate.

Example **27** was prepared as the tetra-sodium salt in a manner similar to that described for example **TP4** starting with example **19**.

^1H NMR (400 MHz, D_2O) δ 7.76 (s, 1H), 6.83 (d, J = 4.4 Hz, 1H), 6.80 (d, J = 4.8 Hz, 1H), 5.94 (d, J = 25.2 Hz, 1H), 5.24 (dd, J = 55.2, 5.2 Hz, 1H), 4.78 (dd, J = 26.8, 5.2 Hz, 1H), 4.08 – 4.18 (m, 2H).

^{19}F NMR (376 MHz, D_2O) δ -193.74 – -194.02 (m).

^{31}P NMR (162 MHz, D_2O) δ -4.60 (d, J = 53.2 Hz, 1P), -10.25 (d, J = 48.4 Hz, 1P), -20.28 (t, J = 48.4 Hz, 1P).



Example 28 - ((2R,3R,4R,5S)-5-(4-amino-5-fluoropyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-azido-4-fluoro-3-hydroxytetrahydrofuran-2-yl)methyl tetrahydrogen triphosphate.


Example **28** was prepared as the tetra-sodium salt in a manner similar to that described for example **TP4** starting with example **21**.

^1H NMR (400 MHz, D_2O) δ 7.64 (s, 1H), 6.60 (s, 1H), 5.90 (d, J = 24.4 Hz, 1H), 5.20 (dd, J = 54.8, 4.8 Hz, 1H), 4.72 (dd, J = 27.2, 4.8 Hz, 1H), 4.05 – 4.18 (m, 2H).

^{19}F NMR (376 MHz, D_2O) δ -161.00 (s), -196.39 – -196.69 (m).

^{31}P NMR (162 MHz, D_2O) δ -8.24 (d, J = 50.4 Hz), -14.20 (d, J = 46.0 Hz), -24.08 (t, J = 48.4 Hz).

MS m/z = 567.87 [$\text{M}+1$]. MS system: Thermo LCQ Advantage



Example 29 - ((2R,3R,4R,5S)-5-(4-amino-2-fluoropyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-azido-4-fluoro-3-hydroxytetrahydrofuran-2-yl)methyl tetrahydrogen triphosphate.


Example **29** was prepared as the tetra-sodium salt in a manner similar to that described for example **TP4** starting with example **22**.

^1H NMR (400 MHz, D_2O) δ 6.81 (d, J = 4.4 Hz, 1H), 6.75 (d, J = 4.8 Hz, 1H), 5.81 (d, J = 24.4 Hz, 1H), 5.16 (dd, J = 54.4, 4.8 Hz, 1H), 4.70 (dd, J = 26.8, 4.4 Hz, 1H), 4.02-4.12 (m, 2H).

^{19}F NMR (376 MHz, D_2O) δ -75.95 (s), -196.51 – -196.80 (m).

^{31}P NMR (162 MHz, D_2O) δ -8.29 (d, J = 53.2 Hz), -14.22 (d, J = 48.4 Hz), -24.09 (t, J = 48.4 Hz).

MS m/z = 567.59 $[\text{M}+1]$. MS system: Thermo LCQ Advantage



Example 30 - ((2R,3R,4R,5S)-5-(4-amino-5-fluoropyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-cyano-4-fluoro-3-hydroxytetrahydrofuran-2-yl)methyl tetrahydrogen triphosphate.


Example **30** was prepared as the tetra-sodium salt in a manner similar to that described for example **TP4** starting with example **23**.

^1H NMR (400 MHz, D_2O) δ 7.64 (s, 1H), 6.57 (s, 1H), 5.87 (d, J = 24.8 Hz, 1H), 5.26 (dd, J = 53.6, 4.0 Hz, 1H), 4.82 (dd, J = 25.2, 4.4 Hz, 1H), 4.26-4.35 (m, 2H).

^{19}F NMR (376 MHz, D_2O) δ -161.05 (s), -194.92 – -195.19 (m).

^{31}P NMR (162 MHz, D_2O) δ -8.22 (d, J = 50.8 Hz), -14.48 (d, J = 48.4 Hz), -24.01 (t, J = 48.4 Hz).

MS m/z = 551.91 $[\text{M}+1]$. MS system: Thermo LCQ Advantage




Example 31 (also TP11) - ((2R,3R,4R,5S)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-4-fluoro-3-hydroxy-2-methyltetrahydrofuran-2-yl)methyl tetrahydrogen triphosphate.

Example 31 was prepared as the tetra-sodium salt in a manner similar to that described for example TP4 starting with example 24

^1H NMR (400 MHz, D_2O) δ 7.66 (s, 1H), 6.78 (d, J = 4.8 Hz, 1H), 6.72 (d, J = 4.4 Hz, 1H), 5.58 (dd, J = 23.6, 2.4 Hz, 1H), 5.16 (ddd, J = 55.2, 5.2, 2.8 Hz, 1H), 4.51 (dd, J = 23.2, 5.2 Hz, 1H), 3.88 (dd, J = 11.6, 6.0 Hz, 1H), 3.78 (dd, J = 10.8, 4.0 Hz, 1H), 1.2 (s, 3H).

^{19}F NMR (376 MHz, D_2O) δ -195.74 – -196.01 (m).

^{31}P NMR (162 MHz, D_2O) δ -8.24 (d, J = 50.4 Hz), -13.54 (d, J = 45.6 Hz), -24.11 (t, J = 48.0 Hz).



Example 32 (also TP12) - ((2R,3S,4R,5S)-5-(4-amino-5-fluoropyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-cyano-3,4-dihydroxytetrahydrofuran-2-yl)methyl tetrahydrogen triphosphate.


Example 32 was prepared as the tetra-sodium salt in a manner similar to that described for example TP4 starting with example 25.

^1H NMR (400 MHz, D_2O) δ 7.59 (s, 1H), 6.57 (s, 1H), 5.44 (d, J = 6.0 Hz, 1H), 4.56 (d, J = 5.2 Hz, 1H), 4.48 (dd, J = 5.6 Hz, 1H), 4.16 (dd, J = 11.6, 6.0 Hz, 1H), 4.08 (dd, J = 11.2, 5.2 Hz, 1H).

^{19}F NMR (376 MHz, D_2O) δ -161.25 (s).

5 ^{31}P NMR (162 MHz, D_2O) δ -8.29 (d, J = 48.4 Hz), -14.49 (d, J = 53.2 Hz), -24.15 (t, J = 48.4 Hz).

MS m/z = 549.90 $[\text{M}+1]$. MS system: Thermo LCQ Advantage




10 **Example 33 (also TP13) - ((2R,3S,4R,5S)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-(chloromethyl)-3,4-dihydroxytetrahydrofuran-2-yl)methyl tetrahydrogen triphosphate.**

Example 33 was prepared as the tetra-triethylamine salt in a manner similar to that described for example TP3 starting with example 26.

15 ^1H NMR (400 MHz, D_2O) δ 7.76 (s, 1H), 6.92 (br s, 1H), 6.85 (br s, 1H), 5.32 (d, J = 9.6 Hz, 1H), 4.78 (dd, J = 8, 6.4 Hz, 1H), 4.53 (d, J = 5.6 Hz, 1H), 4.08 (dd, J = 10.0, 4.0 Hz, 1H), 3.83 – 3.95 (m, 3H), 3.07 (q, J = 7.6 Hz, 24 H), 1.16 (t, J = 7.6 Hz, 36 H).

^{31}P NMR (162 MHz, D_2O) δ -9.44 (d, J = 45.6 Hz), -11.51 (d, J = 48.8 Hz), -22.95 (t, J = 48.4 Hz).

MS m/z = 555.06 $[\text{M}+1]$. MS system: Thermo LCQ Advantage



Example 34 (also PD6) - (2S)-2-ethylbutyl 2-((((2R,3R,4R,5S)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-cyano-4-fluoro-3-hydroxytetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate.


Example 1 (3.8 mg, 0.013 mmol) was dissolved in anhydrous *N*-methyl-2-pyrrolidone (0.2 mL) and THF (0.1 mL) was added under an argon atmosphere. *tert*-Butyl magnesium chloride (1M in THF, 20 μ L, 0.024 mmol) was then added at RT, and white solids precipitated. After 5 min, a solution of *p*-nitrophenylphosphoramidate **PD3c** (12 mg, 0.026 mmol) in THF (0.1 mL) was added to the reaction mixture in one portion, and the resulting mixture was heated to 50 $^{\circ}$ C. After 20 h, the reaction mixture was allowed to cool to RT and was then purified directly by preparatory HPLC (Phenominex Synergi 4u Hydro-RR 80 $\text{\AA}$ 150 x 30 mm column, 40-100% acetonitrile/water gradient). The fractions containing the desired product were combined and were lyophilized to afford example **34** (2.9 mg, 37%, 3:2 diastereomeric mixture) as a white solid.

^1H NMR (400 MHz, CD_3OD) δ 7.80 (s, 0.3H), 7.78 (s, 0.6H), 7.38 – 7.10 (m, 5H), 6.85 (br dd, J = 4.7, 2.2 Hz, 1H), 6.75 – 6.71 (m, 1H), 5.54 – 5.46 (m, 1H), 4.65 – 4.58 (m, 1H), 4.53 – 4.31 (m, 3H), 4.07 – 3.84 (m, 3H), 1.54 – 1.39 (m, 1H), 1.38 – 1.19 (m, 7H), 0.92 – 0.81 (m, 6H) 29H

^{31}P NMR (162 MHz, CD_3OD) δ 3.25 (br s).

LC/MS: t_R = 1.55 min, MS m/z = 603.19 $[\text{M}+\text{H}]^+$; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-2.0 min 2-100% ACN, 2.0 min-3.05 min 100% ACN, 3.05 min-3.2 min 100%-2% ACN, 3.2 min-3.5 min 2% ACN at 2 μ L/min.

HPLC: t_R = 2.98 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.



5

Example 35 (also PD7) - (2S)-ethyl 2-((((2R,3S,4R,5S)-5-(4-aminopyrrolo[2,1-f][1,2,4]triazin-7-yl)-2-cyano-3,4-dihydroxytetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino) propanoate.

Example 1 (19 mg, 65.3 μ mol) was dissolved in NMP (0.2 mL). THF (0.1 mL) was added followed by tert-butyl magnesium chloride (1.0M solution in tetrahydrofuran, 0.098 mL) at RT under an argon atmosphere. After 5 min, a solution of intermediate PD7a (Prepared according to US20120009147A1, 51.4 mg, 130 μ mol) in THF (0.1 mL) was added, and the resulting mixture was warmed to 50 $^{\circ}$ C. After 1 h, the reaction mixture was allowed to cool to RT and was purified directly by preparatory HPLC (Phenominex Synergi 4u Hydro-RR 80 $\text{\AA}$ 150 x 30 mm column, 5-100% acetonitrile/water gradient). The fractions containing product were combined, and concentrated and the resulting residue was repurified by preparatory HPLC (Phenominex Luna 5u C18 100 x 30 mm column, 5-100% acetonitrile/water gradient) to afford example 35 (12 mg, 34%, 3:2 mixture of diastereomers) as a white solid.

^1H NMR (400 MHz, CD_3OD) δ 7.80 (d, J = 2.3 Hz, 0.4H), 7.78 (d, J = 2.3 Hz, 0.6H), 7.36 – 7.12 (m, 5H), 6.88 – 6.81 (m, 1H), 6.76 – 6.70 (m, 1H), 5.53 – 5.46 (m, 1H), 4.66 – 4.60 (m, 1H), 4.55 – 4.30 (m, 3H), 4.15 – 3.98 (m, 2H), 3.93 – 3.79 (m, 1H), 1.30 – 1.12 (m, 6H).


^{31}P NMR (162 MHz, CD_3OD) δ 3.27 (br s).

LC/MS: major diastereomer t_R = 1.28 min, MS m/z = 547.14 [M+H], minor diastereomer t_R = 1.30 min, MS m/z = 547.04 [M+H]; LC system: Thermo Accela 1250 UHPLC; MS system: Thermo LCQ Fleet; Column: Kinetex 2.6 μ XB-C18 100A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% acetic acid, Water with 0.1% acetic acid; Gradient: 0 min-2.0 min 2-100% ACN,

2.0 min-3.05 min 100% ACN, 3.05 min-3.2 min 100%-2% ACN, 3.2 min-3.5 min 2% ACN at 2 μ l/min.

HPLC: major diastereomer t_R = 2.44 min, minor diastereomer t_R = 2.46 min; HPLC system: Agilent 1100 series.; Column: Gemini 5 μ C18 110A, 50 x 4.6 mm; Solvents: Acetonitrile with 0.1% TFA, Water with 0.1% TFA; Gradient: 0 min-5.0 min 2-98% ACN, 5.0 min-6.0 min 98% ACN at 2 mL/min.

Also provided are separate embodiments comprising, respectively, compounds of Formula (A), Formula (B), Formula (C), Formula (D), and Formula (E):



10 wherein, in each instance, X^1 represents an oxygen protecting group and X^2 represents an amine protecting group.

Useful oxygen protecting groups include a silyl ether protecting group or a benzyl-type protecting group, including methoxybenzyl groups.

Useful silyl ether protecting groups include Trimethylsilyl (TMS), Triethylsilyl (TES),
 15 Dimethylisopropylsilyl (IPDMS), Diethylisopropylsilyl (DEIPS), Dimethylthexylsilyl (TDS), *t*-Butyldimethylsilyl (TBS or TBDMS), *t*-Butyldiphenylsilyl (TBDPS), Tribenzylsilyl, Tri-*p*-xylylsilyl, Triisopropylsilyl (TIPS), Diphenylmethylsilyl (DPMS), Di-*t*-butylmethylsilyl (DTBMS), Triphenylsilyl (TPS), Methylphenylsilyl (MDPS), *t*-butylmethoxyphenylsilyl, Tris(trimethylsilyl)silyl (sisyl), (2-Hydroxystyryl)dimethylsilyl (HSDMS), (2-
 20 Hydroxystyryl)diisopropylsilyl (HSDIS). *t*-Butylmethoxyphenylsilyl (TBMPS), and *t*-Butoxydiphenylsilyl (DPTBOS) protecting groups.

Useful benzyl-type protecting groups include benzyl, halogenated benzyl, *p*-methoxybenzyl, benzyloxymethyl, 2,4-dimethoxybenzyl, 3,4-dimethoxybenzyl, 2,6-dimethoxybenzyl, *p*-CF₃-benzyl, *p*-methylbenzyl, *p*-methoxybenzyl, 3,5-dimethylbenzyl, *p*-tert-butylbenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, *p*-halobenzyl, including *p*-Br-benzyl, 2,6-dichlorobenzyl, *p*-cyanobenzyl, *p*-phenylbenzyl, 2,6-difluorobenzyl, *p*-acylaminobenzyl (PAB), *p*-azidobenzyl (Azb), 4-azido-3-chlorobenzyl, 2-trifluoromethylbenzyl, *p*-(methylsulfinyl)benzyl, 2-picolyl, 4-picolyl, 3-methyl-2-picolyl N-oxido, 2-quinolinylmethyl, diphenylmethyl (DPM), *p,p'*-dinitrobenzhydryl, triphenylmethyl, alpha-naphthylmethyl, *p*-methoxyphenyldiphenylmethyl, di(*p*-methoxyphenyl)phenylmethyl, tri(*p*-methoxyphenyl)methyl, 4,4',4''-tris(benzoyloxyphenyl)methyl, and 2-naphthylmethyl protecting groups.

Useful amine protecting groups include *p*-methoxybenzyl carbonyl (Moz or MeOZ), acetyl (Ac), benzoyl (Bz), *p*-methoxybenzyl (PMB), 3,4-dimethoxybenzyl (DMPM), *p*-methoxyphenyl (PMP), tosyl (Ts or Tos), trifluoroacetamide, and trityl protecting groups. Useful amine protecting groups also include carbamate and amide protecting groups. Examples of carbamate protecting groups include methyl and ethyl carbamates such as 9-fluorenylmethyloxycarbonyl (Fmoc), 9-(2-sulfo)fluorenylmethyl, 9-(2,7-dibromo)fluorenylmethyl, 17-tetrabenz[a,c,g,i]fluorenylmethyl (Tbfmoc), 2-chloro-3-indenylmethyl (Climoc), benz[*f*]inden-3-ylmethyl (Bimoc), 2,7-di-*t*-butyl[9-(10,10-dioxo-10,10,10,10-tetrahydrothioxanyl)]methyl (DBD-Tmoc), [2-(1,3-dithianyl)methyl (Dmoc), and 1,1-dioxobenzo[*b*]thiophene-2-ylmethyl (Bsmoc) carbamates.

Examples of useful substituted ethyl carbamates include 1,1-dimethyl-2-cyanoethyl, 2-phosphonioethyl (Peoc), 2-methylthioethyl, 2-(*p*-toluenesulfonyl)ethyl, 2,2,2-trichloroethyl (Troc), 2-(trimethylsilyl)ethyl (Teoc), 2-phenylethyl (hZ), 1-(1-adamantyl)-1-methylethyl (Adpoc), 1,1-dimethyl-2-bromoethyl, 1,1-dimethyl-2-chloroethyl, 1,1-dimethyl-2,2-dibromoethyl (DB-*t*-BOC), 1,1-dimethyl-2,2,2-trichloroethyl (TCBOC), 1-methyl-1-(4-biphenyl)ethyl (Bpoc), 1-(3,5-di-*t*-butylphenyl)-1-methylethyl (*t*-Bumeoc), 2-(2'-pyridyl)ethyl, 2-(4'-pyridyl)ethyl, 2,2-bis(4'-nitrophenyl)ethyl (Bnpeoc), *N*-(2-pivaloylamino)-1,1-dimethylethyl, 2-[(2-nitrophenyl)dithio]-1-phenylethyl (NpSSPeoc), 2-(*N,N*-dicyclohexylcarboxamido)ethyl, *t*-butyl (Boc or BOC), 1-adamantyl (1-Adoc), 2-adamantyl (2-Adoc), vinyl (Voc), allyl (Aloc or alloc), 1-isopropylallyl (Ipaoc), cinnamyl (Coc), 4-nitrocinnamyl (Noc), 3-(3'-pyridyl)prop-2-enyl (Paloc), 8-quinolyl, and *N*-hydroxypiperidinyl,

carbamates, as well as alkylidithio carbamates, including methyldithio, ethyldithio, isopropyldithio, t-butyldithio, and phenyldithio carbamates.

Also useful are aryl-containing and substituted aryl-containing carbamates such as benzyl, *p*-methoxybenzyl, *p*-nitrobenzyl, *p*-bromobenzyl, *p*-chlorobenzyl, 2,4-dichlorobenzyl, 4-methylsulfinylbenzyl (MsZ), 9-anthrylmethyl, 4-methylthiophenyl (Mtpc), 1-methyl-1-(triphenylphosphonio)ethyl (2-triphenylphosphonioisopropyl) (Ppoc), 2-dansylethyl (Dnseoc), 2-(4-nitrophenyl)ethyl (Npeoc), 4-phenylacetoxybenzyl (PhAcOZ), 4-azidobenzyl (ACBZ), 4-azidomethoxybenzyl, *m*-chloro-*p*-acyloxybenzyl, *p*-(dihydroxyboryl)benzyl, carbobenzyloxy (Cbz), 4-benzisoxazolymethyl (Bic), 2-(trifluoromethyl)-6-chromonylmethyl (Tcroc), phenyl, and diphenylmethyl carbamates. Additional carbamates include butynyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, 1-methylcyclobutyl, 1-methylcyclohexyl, 1,1-dimethylpropynyl, and 1-methyl-1-cyclopropylmethyl carbamates.

Useful amide protecting groups for amines include *N*-formyl, *N*-acetyl, *N*-chloroacetyl, *N*-trichloroacetyl, *N*-trifluoroacetyl (TFA), *N*-phenylacetyl, *N*-3-phenylpropionyl, *N*-4-pentenoyl, *N*-picolinoyl, *N*-3-pyridylcarboxamido, *N*-benzoylphenylalanyl, *N*-benzoyl, and *N*-*p*-phenylbenzoyl amides.

Antiviral Activity

Another embodiment relates to methods of inhibiting viral infections, comprising the step of treating a sample or subject suspected of needing such inhibition with a composition herein.

Considered herein are samples suspected of containing a virus include natural or man-made materials such as living organisms; tissue or cell cultures; biological samples such as biological material samples (blood, serum, urine, cerebrospinal fluid, tears, sputum, saliva, tissue samples, and the like); laboratory samples; food, water, or air samples; bioproduct samples such as extracts of cells, particularly recombinant cells synthesizing a desired glycoprotein; and the like. Typically the sample will be suspected of containing an organism which induces a viral infection, frequently a pathogenic organism such as a tumor virus. Samples can be contained in any medium including water and organic solvent/water mixtures. Samples include living organisms such as humans, and manmade materials such as cell cultures.

If desired, the anti-virus activity of a compound after application of the composition

can be observed by any method including direct and indirect methods of detecting such activity. Quantitative, qualitative, and semiquantitative methods of determining such activity are all contemplated. Typically one of the screening methods described above are applied, however, any other method such as observation of the physiological properties of a living organism are also applicable.

The antiviral activity of a compound can be measured using standard screening protocols that are known. For example, the antiviral activity of a compound can be measured using the following general protocols.

Respiratory syncytial virus (RSV) antiviral activity and cytotoxicity assays

Anti-RSV activity

Antiviral activity against RSV is determined using an infectious cytopathic cell protection assay in HEp-2 cells. In this assay, compounds inhibiting viral infection and/or replication produce a cytoprotective effect against the virus-induced cell killing that can be quantified using a cell viability reagent. The techniques used here are novel adaptations of methods described in published literature (Chapman et al., *Antimicrob Agents Chemother.* 2007, 51(9):3346-53.)

HEp-2 cells are obtained from ATCC (Manassas, VI) and maintained in MEM media supplemented with 10% fetal bovine serum and penicillin/streptomycin. Cells are passaged twice a week and kept at subconfluent stage. Commercial stock of RSV strain A2 (Advanced Biotechnologies, Columbia, MD) is titered before compound testing to determine the appropriate dilution of the virus stock that generates desirable cytopathic effect in HEp-2 cells.

For antiviral tests, HEp-2 cells are grown in large cell culture flasks to near confluency but not fully so. The compounds to be tested are prediluted in DMSO in 384-well compound dilution plates, either in an 8 or 40 sample per plate standardized dose response format. 3-fold serial dilution increments of each test compound are prepared in the plates and test samples are transferred via acoustic transfer apparatus (Echo, Labcyte) at 100 nl per well into cell culture assay 384-well plates. Each compound dilution is transferred in single or quadruplicate samples into dry assay plates, which are stored until assay is ready to go. The positive and negative controls are laid out in opposite on ends of the plate in vertical blocks (1 column).

Subsequently, an infectious mixture is prepared using an appropriate dilution of virus stock previously determined by titration with cells at a density of 50,000/ml and 20 μ L/well is added to test plates w/compounds via automation (uFlow, Biotek). Each plate includes negative and positive controls (16 replicates each) to create 0% and 100% virus inhibition standards, respectively. Following the infection with RSV, testing plates are incubated for 4 days in a 37 °C cell culture incubator. After the incubation, a cell viability reagent, Cell TiterGlo (Promega, Madison, WI) is added to the assay plates, which are incubated briefly, and a luminescent readout is measured (Envision, Perkin Elmer) in all the assay plates. The RSV-induced cytopathic effect, percentage inhibition, is determined from the levels of remaining cell viability. These numbers are calculated for each tested concentration relative to the 0% and 100% inhibition controls, and the EC₅₀ value for each compound is determined by non-linear regression as a concentration inhibiting the RSV-induced cytopathic effect by 50%. Various potent anti-RSV tool compounds are used as positive controls for antiviral activity.

Cytotoxicity Assay in HEp-2 Cells

Cytotoxicity of tested compounds is determined in uninfected HEp-2 cells in parallel with the antiviral activity using the cell viability reagent in a similar fashion as described before for other cell types (Cihlar et al., *Antimicrob Agents Chemother.* **2008**,52(2):655-65.). The same protocol as for the determination of antiviral activity is used for the measurement of compound cytotoxicity except that the cells are not infected with RSV. Instead, an uninfected cell mixture at the same density is added at 20 μ L/well to plates containing prediluted compounds, also at 100 nL/sample. Assay plates are then incubated for 4 days followed by a cell viability test using the same CellTiter Glo reagent addition and measurement of luminescent readouts. Untreated cell and cells treated at 2 μ M puromycin (Sigma, St. Louis, MO) serve as 100% and 0% cell viability control, respectively. The percent of cell viability is calculated for each tested compound concentration relative to the 0% and 100% controls and the CC₅₀ value is determined by non-linear regression as a compound concentration reducing the cell viability by 50%.

Cytotoxicity Assay in MT-4 Cells

The MT-4 cell line was obtained from the NIH AIDS Research and Reference Reagent Program (Germantown, MD) and cultured in RPMI-1640 medium (Irvine Scientific, Santa Ana, CA, Cat # 9160) supplemented with 10% FBS, 100 units/mL penicillin, 100

units/mL streptomycin, and 2 mM L-Glutamine. The MT-4 cells were passaged twice per week to maintain cell densities below 0.6×10^6 cells/mL. Complete RPMI-1640 media containing 100x concentrations of 3-fold serially diluted compound, ranging from 26 nM to 530 μ M, were stamped in quadruplicate into black 384-well plates. After compound addition,

5 2×10^3 MT-4 cells were added to each well using a MicroFlo liquid dispenser (BioTek, Winooski, VT) and the cells were cultured for 5 days at 37 °C in a 5% CO₂ incubator. Following the incubation the cells were allowed to equilibrate to 25 °C and cell viability was determined by adding 25 μ L of Cell-Titer Glo viability reagent. The mixture was incubated for 10 minutes at 25 °C, and the luminescence signal was quantified on a Victor

10 Luminescence plate reader. The CC₅₀ value is defined as the concentration of compound that reduces cell viability by 50% as determined by the Cell-Titer Glo signal. The data were analyzed using Pipeline Pilot Plate Data Analytics Collection software (Version 7.0, Accelrys, San Diego, CA). CC₅₀ values were calculated from a non-linear regression analysis using a 4-parameter sigmoidal dose-response equation: $Y = \text{Bottom} + (\text{Top} - \text{Bottom}) / (1 + 10^{[(\text{LogCC}_{50} - X) * \text{HillSlope}]})$ where the Top and Bottom were fixed at 100% and 0% cell viability, respectively. CC₅₀ values were calculated as the average $\pm$ standard deviation of 3 independent experiments.


15

Example	EC ₅₀ / μ M	HEp-2 CC ₅₀ / μ M	MT-4 CC ₅₀ / μ M
1	7.3	>50	>53
2	9.6	>100	>106
3	2.0	>100	>106
4	30	>50	>53
5	2.1	>50	>53
10 (PD1)	1.0	39	7.3
11 (PD2)	1.2	42	19.7
12 (PD3)	3.0	>50	27
13	10.5	>100	>106


14	40	>100	>106	
15 (PD4)	1.6	35	7.9	
16 (PD5)	0.3	18	25	
19	0.40	>44	16	
20	>50	>50		
21	50	36	15	
22	0.57	>100	32	
23	23.5	72	21	
24	1.0	>100	11	
25	9.2	>100	>93	
26	>50	42	>57	
34 (PD6)	0.21	>50	>50	
35 (PD7)	0.34	>50	47	

Another benefit relates to advantage that compounds bearing R⁴ substitution provides in comparison to compounds lacking R⁴ substitution (i.e. those in which R⁴ = H) with regard to MT-4 cytotoxicity. For example, compound (2*S*,3*R*,4*S*,5*R*)-2-(4-aminopyrrolo[1,2-

- 5 *f*][1,2,4]triazin-7-yl)-5-(hydroxymethyl)tetrahydrofuran-3,4-diol (Patil, S. A.; Otter, R. A.; Klein, R. S. *Tetrahedron Lett.* **1994**, 35, 5339-5342), having the structure:



exhibits a MT-4 CC_{50} = 0.007 μ M; whereas Examples **1**, **4**, **5**, **20**, and **26** all exhibit a MT-4 CC_{50} > 53 μ M. Furthermore, compound (2*R*,3*R*,4*R*,5*S*)-5-(4-aminopyrrolo[1,2-
f][1,2,4]triazin-7-yl)-4-fluoro-2-(hydroxymethyl)tetrahydrofuran-3-ol (WO2012037038A1),
5 having the structure:



exhibits a MT-4 CC_{50} = 30 μ M; whereas Examples **2**, **3**, **13**, and **14** all exhibit a MT-4 CC_{50} > 106 μ M.

Another benefit relates to advantage that R^3 = F compounds bearing R^4 substitution provides
10 in comparison to compounds lacking R^4 substitution (i.e. those in which R^4 = H) with regard
to HEp-2 anti-RSV activity. For example the compound (2*R*,3*R*,4*R*,5*S*)-5-(4-
aminopyrrolo[1,2-*f*][1,2,4]triazin-7-yl)-4-fluoro-2-(hydroxymethyl)tetrahydrofuran-3-ol
(WO2012037038A1) having the structure above has a HEp-2 EC_{50} = >100 μ M; whereas
examples **2**, **3**, **13**, **14**, **19**, **21**, **22**, **23**, and **24** all exhibit EC_{50} = <100 μ M.

RSV RNP Preparation

RSV ribonucleoprotein (RNP) complexes were prepared from a method modified from Mason *et al* (1). HEp-2 cells were plated at a density of 7.1×10^4 cells/cm² in MEM + 10% fetal bovine serum (FBS) and allowed to attach overnight at 37°C (5% CO₂). Following attachment, the cells were infected with RSV A2 (MOI=5) in 35 mL MEM + 2% FBS. At 20 hours post-infection, the media was replaced with MEM + 2% FBS supplemented with 2 µg/mL actinomycin D and returned to 37°C for one hour. The cells were then washed once with PBS and treated with 35 mL of PBS + 250 µg/mL lyso-lecithin for one minute, after which all liquid was aspirated. The cells were harvested by scrapping them into 1.2 mL of buffer A [50 mM TRIS acetate (pH 8.0), 100 mM potassium acetate, 1 mM DTT and 2 µg/mL actinomycin D] and lysed by repeated passage through an 18 gauge needle (10 times). The cell lysate was placed in ice for 10 minutes and then centrifuged at 2400g for 10 minutes at 4°C. The supernatant (S1) was removed and the pellet (P1) was disrupted in 600 µL of Buffer B [10 mM TRIS acetate (pH 8.0), 10mM potassium acetate and 1.5 mM MgCl₂] supplemented with 1% Triton X-100 by repeated passage through an 18 gauge needle (10 times). The resuspended pellet was placed in ice for 10 minutes and then centrifuged at 2400g for 10 minutes at 4°C. The supernatant (S2) was removed and the pellet (P2) was disrupted in 600 µL of Buffer B supplemented with 0.5% deoxycholate and 0.1% Tween 40. The resuspended pellet was placed in ice for 10 minutes and then centrifuged at 2400g for 10 minutes at 4°C. The supernatant (S3) fraction, containing the enriched RSV RNP^o complexes, was collected and the protein concentration determined by UV absorbance at 280 nm. Aliquoted RSV RNP S3 fractions were stored at -80°C.

RSV RNP Assay

Transcription reactions contained 25µg of crude RSV RNP complexes in 30 µL of reaction buffer [50 mM TRIS-acetate (pH 8.0), 120 mM potassium acetate, 5% glycerol, 4.5 mM MgCl₂, 3 mM DTT, 2 mM ethyleneglycol-bis(2-aminoethylether)-tetraacetic acid (EGTA), 50 µg/mL BSA, 2.5 U RNasin (Promega), ATP, GTP, UTP, CTP and 1.5 uCi [α -³²P] NTP (3000 Ci/mmol)]. The radiolabeled nucleotide used in the transcription assay was selected to match the nucleotide analog being evaluated for inhibition of RSV RNP transcription. Cold, competitive NTP was added at a final concentration of one-half its K_m (ATP= 20 µM, GTP= 12.5 µM, UTP= 6 µM and CTP= 2 µM). The three remaining nucleotides were added at a final concentration of 100 µM.

To determine whether nucleotide analogs inhibited RSV RNP transcription, compounds were added using a 6 step serial dilution in 5-fold increments. Following a 90 minute incubation at 30°C, the RNP reactions were stopped with 350 µL of Qiagen RLT lysis buffer and the RNA was purified using a Qiagen RNeasy 96 kit. Purified RNA was


5 denatured in RNA sample loading buffer (Sigma) at 65°C for 10 minutes and run on a 1.2% agarose/MOPS gel containing 2M formaldehyde. The agarose gel was dried and exposed to a Storm phosphorimager screen and developed using a Storm phosphorimager (GE Healthcare). The concentration of compound that reduced total radiolabeled transcripts by 50% (IC₅₀) was calculated by non-linear regression analysis of two replicates.

- 10 **Reference** Mason, S., Lawetz, C., Gaudette, Y., Do, F., Scouten, E., Lagace, L., Simoneau, B. and Liuzzi, M. (2004) Polyadenylation-dependent screening assay for respiratory syncytial virus RNA transcriptase activity and identification of an inhibitor. *Nucleic Acids Research*, **32**, 4758-4767.

Example	IC ₅₀ / µM
6 (TP1)	0.086
7 (TP2)	1
8 (TP3)	0.025
9 (TP4)	0.12
17 (TP5)	0.56
18 (TP6)	0.42
27 (TP7)	0.097
28 (TP8)	0.086
29 (TP9)	0.080
30 (TP10)	
31 (TP11)	
32 (TP12)	0.022
33 (TP13)	


A further consideration relates to advantage that compounds exemplified exhibit potent inhibition of the RSV RNP transcription in comparison to compounds with 2'CMe substitution. For example, ((2*R*,3*R*,4*R*,5*S*)-5-(4-aminopyrrolo[2,1-*f*][1,2,4]triazin-7-yl)-3,4-dihydroxy-4-methyltetrahydrofuran-2-yl)methyl tetrahydrogen triphosphate

5 (WO2008089105A2, and WO2010002877A2), having the structure:



exhibits an IC_{50} = 8.5 μ M; whereas Example **TP3** exhibits an IC_{50} = 0.025 μ M. Furthermore, compound ((2*R*,3*R*,4*R*,5*S*)-5-(4-aminopyrrolo[2,1-*f*][1,2,4]triazin-7-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methyl tetrahydrogen triphosphate (WO 2011035231 A1), having


10 the structure:



exhibits an IC_{50} = >100 μ M; whereas Examples **TP1** exhibits an IC_{50} = 0.086 μ M. and **TP2** exhibits an IC_{50} = 1 μ M.

What is Claimed:

1. A compound of Formula (I), or a pharmaceutically acceptable salt thereof:



wherein:

R^1 is H or F;


R^2 is H or F;


R^3 is OH or F;

R^4 is CN, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_2\text{-C}_4$ alkenyl, $\text{C}_2\text{-C}_4$ alkynyl, $\text{C}_3\text{-C}_4$ cycloalkyl, azido, halogen, or $\text{C}_1\text{-C}_2$ haloalkyl;

R^6 is OH;

R^5 is selected from the group of H and:






wherein:

n' is selected from 1, 2, 3, and 4;

R^8 is selected from C_1 - C_8 alkyl, - O - C_1 - C_8 alkyl, benzyl, - O -benzyl, - CH_2 - C_3 - C_6 cycloalkyl, - O - CH_2 - C_3 - C_6 cycloalkyl, and CF_3 ;

R^9 is selected from phenyl, 1-naphthyl, 2-naphthyl,



R^{10} is selected from H and CH_3 ;

R^{11} is selected from H or C_1 - C_6 alkyl;


R^{12} is selected from H, C_1 - C_8 alkyl, benzyl, C_3 - C_6 cycloalkyl, and - CH_2 - C_3 - C_6 cycloalkyl.

2. A compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein R^1 is H.

3. A compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein R^2 is H.

4. A compound according to any of Claims 1, 2, or 3 wherein both R^1 and R^2 are H.

5. A compound according to any of Claims 1, 2, 3, or 4, or a pharmaceutically acceptable salt thereof, wherein R^1 , R^2 , and R^5 are each H.
6. A compound according to any of Claims 1, 2, 3, 4, or 5, or a pharmaceutically acceptable salt thereof, wherein both R^1 and R^2 are H, and R^3 is OH.
7. A compound according to any of Claims 1, 2, 3, 4, or 5, or a pharmaceutically acceptable salt thereof, wherein both R^1 and R^2 are H, R^3 is F.
8. A compound according to any of Claims 1, 2, 3, or 4, or a pharmaceutically acceptable salt thereof, of Formula (II):





wherein:

R^3 is OH or F;

R^4 is CN, C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_3 - C_4 cycloalkyl, azido, halogen, or C_1 - C_2 haloalkyl;

R^5 is selected from the group of H and:





wherein:

n^1 is selected from 1, 2, 3, and 4;

R^8 is selected from C_1 - C_8 alkyl, - O - C_1 - C_8 alkyl, benzyl, - O -benzyl, - CH_2 - C_3 - C_6 cycloalkyl, - O - CH_2 - C_3 - C_6 cycloalkyl, and CF_3 ;

R^9 is phenyl;

R^{10} is selected from H and CH_3 ;

R^{11} is selected from H or C_1 - C_6 alkyl;


R^{12} is selected from H, C_1 - C_8 alkyl, benzyl, C_3 - C_6 cycloalkyl, and - CH_2 - C_3 - C_6 cycloalkyl.

9. A compound according to any of Claims 1, 2, 3, 4, 5, or 8, or a pharmaceutically acceptable salt thereof, wherein R^4 is CN, methyl, ethyl, ethenyl, ethynyl, azido, F, Cl, - CH_2 Cl, - CH_2 F, - CHF_2 , or - CF_3 .

10. A compound according to any of Claims 1, 2, 3, 4, 5, 7, 8, or 9, or a pharmaceutically acceptable salt thereof, wherein R^3 is F.

11. A compound according to any of Claims 1, 2, 3, 4, 5, 6, 8, or 9, or a pharmaceutically acceptable salt thereof, wherein R^3 is OH.

12. A compound according to any of Claims 1, 2, 3, 4, 5, 7, 8, 9, or 10, or a pharmaceutically acceptable salt thereof, wherein R^3 is F and R^4 is CN.
13. A compound according to any of Claims 1, 2, 3, 4, 5, 7, 8, 9, or 11, or a pharmaceutically acceptable salt thereof, wherein R^3 is OH and R^4 is CN.
14. A compound according to any of Claims 1, 2, 3, 4, 5, 7, 8, 9, or 10, or a pharmaceutically acceptable salt thereof, wherein both R^1 and R^2 are H, R^3 is F, and R^4 is methyl, ethyl, vinyl, or ethynyl.
15. A compound according to any of Claims 1, 2, 3, 4, 5, 7, 8, 9, or 11, or a pharmaceutically acceptable salt thereof, wherein R^3 is OH, and R^4 is methyl, ethyl, vinyl, or ethynyl.
16. A compound according to any of Claims 1, 2, 3, 4, 5, 7, 8, 9, or 10, or a pharmaceutically acceptable salt thereof, wherein R^3 is F and R^4 is halomethyl.
17. A compound according to any of Claims 1, 2, 3, 4, 5, 7, 8, 9, or 11, or a pharmaceutically acceptable salt thereof, wherein R^3 is OH and R^4 is halomethyl.
18. A compound according to any of Claims 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, or 17, or a pharmaceutically acceptable salt thereof, wherein R^5 is H.
19. A compound according to any of Claims 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, or 17, or a pharmaceutically acceptable salt thereof, wherein R^5 is selected from the group of:



wherein:

R⁸ is selected from C₁-C₈ alkyl, -O-C₁-C₈ alkyl, benzyl, and -CH₂-C₃-C₆ cycloalkyl;
and

R¹² is selected from C₁-C₈ alkyl, benzyl, C₃-C₆ cycloalkyl, and -CH₂-C₃-C₆ cycloalkyl.

20. A compound according to any of Claims 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or 19, or a pharmaceutically acceptable salt thereof, wherein R⁸ and R⁹ are each C₁-C₈ alkyl.


21. A compound according to any of Claims 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or 19, or a pharmaceutically acceptable salt thereof, wherein R⁸ and R⁹ are each selected from C₁-C₆ alkyl.

22. A compound according to any of Claims 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, or 21, or a pharmaceutically acceptable salt thereof, wherein R⁸ and R⁹ are each selected from C₁-C₅ alkyl.


23. A compound according to any of Claims 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, 21, or 22, or a pharmaceutically acceptable salt thereof, wherein R^8 and R^9 are each selected from C_1 - C_4 alkyl.

24. A compound according to any of Claims 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, or 23, or a pharmaceutically acceptable salt thereof, further comprising the proviso that when R^3 is F, R^4 is not methyl.


25. A compound according to any of Claims 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24, or a pharmaceutically acceptable salt thereof, wherein R^5 is selected from the group of H and:




26. A compound according to any of Claims 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, 21, 22, 23, 24, or 25, or a pharmaceutically acceptable salt thereof, wherein R^5 is:




27. A compound according to any of Claims 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, 21, 22, 23, 24, or 25, or a pharmaceutically acceptable salt thereof, wherein R^5 is:




28. A compound according to any of Claims 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, 21, 22, 23, 24, or 25, or a pharmaceutically acceptable salt thereof, wherein R^5 is:




29. A compound according to any of Claims 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, or 28, or a pharmaceutically acceptable salt thereof, selected from the group of:




30. A compound according to any of Claims 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, or 28, or a pharmaceutically acceptable salt thereof, selected from the group of:




31. A compound according to any of Claims 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, or 28, or a pharmaceutically acceptable salt thereof, selected from the group of:




32. A compound according to any of Claims 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, or 28, or a pharmaceutically acceptable salt thereof, selected from the group of:



33. A compound according to any of Claims 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, or 28, or a pharmaceutically acceptable salt thereof, selected from the group of:



34. A compound according to any of Claims 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, or 28, or a pharmaceutically acceptable salt thereof, selected from the group of:



35. A compound of any of Claims 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, or 34, or a pharmaceutically acceptable salt thereof for use in the treatment of a *Pneumovirinae* virus infection in a human.


36. A compound of any of Claims 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, or 34, or a pharmaceutically acceptable salt thereof for use in the treatment of a respiratory syncytial virus infection in a human.

37. A pharmaceutical formulation comprising a pharmaceutically effective amount of a compound of any of Claims 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, or 34, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or excipient.

38. Use of a compound of any of Claims 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, or 34, or a pharmaceutically


acceptable salt thereof in the manufacture of a medicament for treating a *Pneumovirinae* virus infection or a respiratory syncytial virus infection in a human.

39. A compound selected from Formula (A), Formula (B), Formula (C), Formula (D), and Formula (E):



wherein X^1 is an oxygen protecting group and X^2 is an amine protecting group.

40. A compound of Claim 39 selected from:



ABSTRACT OF THE DISCLOSURE

Provided herein are formulations, methods and substituted tetrahydrofuranyl-pyrrolo[1,2-*f*][1,2,4]triazine-4-amine compounds of Formula (I) for treating *Pneumovirinae* virus infections, including respiratory syncytial virus infections, as well as methods and intermediates for synthesis of tetrahydrofuranyl-pyrrolo[1,2-*f*][1,2,4]triazine-4-amine compounds.

