

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2003295784 B2**

(54) Title
Composition and method for treating lupus nephritis

(51) International Patent Classification(s)
C12N 1/19 (2006.01) **C12N 15/85** (2006.01)
C07H 21/04 (2006.01) **C12N 15/86** (2006.01)
C07K 14/705 (2006.01) **A61K 38/00** (2006.01)
C12N 1/21 (2006.01)

(21) Application No: **2003295784** (22) Date of Filing: **2003.11.21**

(87) WIPO No: **WO04/047740**

(30) Priority Data

(31) Number (32) Date (33) Country
60/428,065 **2002.11.21** **US**

(43) Publication Date: **2004.06.18**

(43) Publication Journal Date: **2004.07.22**

(44) Accepted Journal Date: **2009.06.11**

(71) Applicant(s)
Wyeth

(72) Inventor(s)
Shojaee, Negin; O'Toole, Margot; Mounts, William M.

(74) Agent / Attorney
Phillips Ormonde Fitzpatrick, Level 22 367 Collins Street, Melbourne, VIC, 3000

(56) Related Art
Uniprot Accession: NoQ9VFS6
WO 2002081731
WO 2004030615
WO 2002092759
WO 2002059260
US 2002/0137077

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
10 June 2004 (10.06.2004)

PCT

(10) International Publication Number
WO 2004/047740 A3

(51) International Patent Classification⁷: C12N 1/19, 1/21,
15/11, 15/85, 15/86, C07K 5/00, 14/00, C07H 21/04

(74) Agent: NIXON PEABODY LLP; SUITE 900 401, 9TH
STREET, N.W., Washington, DC 20004-2128 (US).

(21) International Application Number:
PCT/US2003/037317

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,
MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT,
RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date:
21 November 2003 (21.11.2003)

(25) Filing Language: English

(26) Publication Language: English

(84) Designated States (*regional*): ARIPO patent (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(30) Priority Data:
60/428,065 21 November 2002 (21.11.2002) US

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:
US 60/428,065 (CIP)
Filed on 21 November 2002 (21.11.2002)

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(71) Applicant (*for all designated States except US*): WYETH
[US/US]; 87 Cambridge Park Drive, Cambridge, MA
02140 (US).

(88) Date of publication of the international search report:
17 February 2005

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): O'TOOLE, Margot
[US/US]; 353 Albermarle Road, Newton, MA 02460 (US).
MOUNTS, William, M. [US/US]; 6 Island Way, Andover,
MA 01810 (US). SHOJAEI, Negin [US/US]; 487 James
Road #201, Palo Alto, CA 94306 (US).

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: COMPOSITION AND METHOD FOR TREATING LUPUS NEPHRITIS

(57) Abstract: The present invention provides novel isolated BFLP1698 polynucleotides and polypeptides encoded by the BFLP1698 polynucleotides. Also provided are the antibodies that immunospecifically bind to a BFLP1698 polypeptide or any derivative (including fusion derivative), variant, mutant, or fragment of the BFLP1698 polypeptide, polynucleotide or antibody. The invention additionally provides methods in which the BFLP1698 polypeptide, polynucleotide and antibody are utilized in the detection and treatment of a broad range of pathological states, as well as to other uses.



WO 2004/047740 A3

COMPOSITION AND METHOD FOR TREATING LUPUS NEPHRITIS

5

FIELD OF THE INVENTION

The invention relates generally to nucleic acids and polypeptides and more specifically to nucleic acids and polypeptides encoding polypeptides useful for detecting and treating lupus nephritis, as well as for identifying therapeutic agents for treating the same.

10

BACKGROUND OF THE INVENTION

Lupus nephritis is an example of a "classical" auto-immune disease in which the patient's immune system attacks his/her own organs. It has been estimated that 45-75% of lupus patients eventually suffer from some form or other of kidney damage. Lupus varies greatly in severity from mild cases requiring minimal intervention to those in which significant damage occurs to vital organs such as lungs, kidneys, heart and brain, and which ultimately can be fatal. Lupus is predominantly a female disease, with an approximate female to male ratio being 9:1. In North America, it is estimated to affect 1 in 500 females mainly between the age of 20 to 40 years.

There is no known cure for lupus. Treatment is typically directed at controlling the symptoms with the hope of putting the disease into remission. Recently, the antibiotic rapamycin has been demonstrated to be an effective therapy in treating lupus nephritis in a murine model of the disease.

The discussion of documents, acts, materials, devices, articles and the like is included in this specification solely for the purpose of providing a context for the present invention. It is not suggested or represented that any or all of these matters formed part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed before the priority date of each claim of this application.

Throughout the description and claims of the specification, the word "comprise" and variations of the word, such as "comprising" and "comprises", is not intended to exclude other additives, components, integers or steps. Throughout the description and claims of the specification, the word "comprise" and variations of the word, such as "comprising" and "comprises", is not intended to exclude other additives, components, integers or steps.

SUMMARY OF THE INVENTION

The invention is based, in part, upon the discovery of a gene, named BFLP1698, whose expression is increased in kidney tissue in mice with lupus nephritis; however, the expression level of the gene does not decrease markedly in response to treatment with rapamycin. This expression profile indicates that the product of the BFLP1698 gene interacts with rapamycin when this antibiotic is administered to ameliorate the symptoms of lupus nephritis. In the absence of rapamycin, the gene product is free to bring about the diseased state, and its effects can include the activation of genes required to bring about the diseased state. In the presence of rapamycin, the BFLP1698 gene product is inactive and the disease state diminishes. Accordingly, the BFLP1698 protein is useful as a target for identifying agents that, like rapamycin, are useful in treating symptoms of lupus nephritis.

In one aspect, the invention provides an isolated nucleic acid molecule that includes the sequence of a nucleotide sequence encoding a BFLP1698 gene product. In a preferred embodiment, the nucleotide sequence includes the sequence of SEQ ID NO:1, or a fragment, homolog, analog or derivative thereof. The nucleic acid can include, *e.g.*, a nucleic acid sequence encoding a polypeptide at least 70%, *e.g.*, 80%, 85%, 90%, 95%, 98%, or even 99% or more identical to a polypeptide that includes the amino acid sequences of SEQ ID NO:2. The nucleic acid can be, *e.g.*, a genomic DNA fragment, or a cDNA molecule.

Also included in the invention is a vector containing one or more of the nucleic acids described herein, and a cell containing the vectors or nucleic acids described herein.

The invention is also directed to host cells transformed with a vector comprising any of the nucleic acid molecules described above.

In another aspect, the invention includes a pharmaceutical composition that includes a BFLP1698 nucleic acid and a pharmaceutically acceptable carrier or diluent.

In a further aspect, the invention includes a substantially purified BFLP1698 polypeptide, *e.g.*, any of the BFLP1698 polypeptides encoded by a BFLP1698 nucleic acid, and fragments, homologs, analogs, and derivatives thereof.

In another aspect, the present invention provides a substantially purified polypeptide comprising an amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID NO:2.

In yet another aspect the present invention provides an isolated polypeptide comprising a rapamycin-binding domain of the amino acid sequence of SEQ ID NO:2.

Even further, the present invention provides an isolated polypeptide at least 993 amino acids in length comprising at least five contiguous amino acids of SEQ ID NO:2, wherein said polypeptide comprises a rapamycin-binding domain, provided that said polypeptide comprises an amino acid sequence other than SEQ ID NO:21.

Still further, the present invention provides a polypeptide comprising at least five contiguous amino acids of SEQ ID NO:15.

2003295784 25 May 2009

The present invention further provides a fusion polypeptide comprising a rapamycin-binding domain of a polypeptide as described herein.

The invention also includes a pharmaceutical composition that includes a polypeptide as described herein and a pharmaceutically acceptable carrier or diluent.

5 In a still further aspect, the invention provides an antibody that binds specifically to a polypeptide as described herein. The antibody can be, *e.g.*, a monoclonal or polyclonal antibody, and fragments, homologs, analogs, and derivatives thereof. The invention also includes a pharmaceutical composition including antibody to a polypeptide as described herein and a pharmaceutically acceptable carrier or diluent. The invention is also directed to isolated
10 antibodies that bind to an epitope on a polypeptide encoded by any of the nucleic acid molecules described above.

The invention also includes kits comprising in one or more containers one or more of a compound that is a BFLP1698 nucleic acid, a BFLP1698 polypeptide and/or an antibody to a BFLP1698 polypeptide. The kit is preferably provided with instructions for use. If desired, the
15 compounds in the kits are provided along with a pharmaceutically acceptable carrier.

The invention further provides a method for producing a polypeptide as described herein by providing a cell containing nucleic acid as described herein, *e.g.*, a vector that includes a BFLP1698 nucleic acid, and culturing the cell under conditions sufficient to express the polypeptide encoded by the nucleic acid. The expressed polypeptide is then recovered from
20 the cell. Preferably, the cell produces little or no endogenous BFLP1698 polypeptide. The cell can be, *e.g.*, a prokaryotic cell or eukaryotic cell.

The invention is also directed to methods of identifying a polypeptide or nucleic acid as described herein in a sample by contacting the sample with a compound that specifically binds to the polypeptide or nucleic acid, and detecting complex formation, if present.

25 Thus in another aspect, the present invention provides a method of detecting the presence of a nucleic acid molecule as described herein in a biological sample, the method comprising:

contacting the sample with a nucleic acid probe that binds specifically to a said nucleic acid; and

30 identifying the bound probe, if present,
thereby detecting the presence of said nucleic acid molecule in said sample.

In a further aspect, the present invention provides a method of detecting the presence of a polypeptide as described herein in a sample, the method comprising:

contacting the sample with a compound that selectively binds to said polypeptide under
35 conditions allowing for formation of a complex between said polypeptide and said compound;
and

detecting said complex, if present, thereby identifying said polypeptide in said sample.

2003295784 25 May 2009

The invention further provides methods of identifying a compound that modulates the activity of a BFLP1698 polypeptide by contacting a BFLP1698 polypeptide with a compound and determining whether the BFLP1698 polypeptide activity is modified.

5 The invention is also directed to compounds that modulate BFLP1698 polypeptide activity is identified by contacting a BFLP1698 polypeptide with the compound and determining whether the compound modifies activity of the BFLP1698 polypeptide, binds to the BFLP1698 polypeptide, or binds to a nucleic acid molecule encoding a BFLP1698 polypeptide.

10 In another aspect, the invention provides a method of determining the presence of or predisposition of a BFLP1698 -associated disorder in a subject. The method includes providing a sample from the subject and measuring the amount of a polypeptide as described herein in the subject sample. The amount of said polypeptide in the subject sample is then compared to the amount of said polypeptide in a control sample. An alteration in the amount of said polypeptide in the subject protein sample relative to the amount of said polypeptide in the control protein sample indicates the subject has a tissue proliferation-associated condition. A control sample is
15 preferably taken from a matched individual, *i.e.*, an individual of similar age, sex or other general condition but who is not suspected of having a tissue proliferation-associated condition. Alternatively, the control sample may be taken from the subject at a time when the subject is not suspected of having a tissue proliferation-associated disorder. In some embodiments, the BFLP1698 is detecting using an antibody to a polypeptide as described herein.

20 Thus in one aspect, the present invention provides a method for determining the presence of or predisposition to lupus nephritis in a subject, the method comprising:

- a) measuring the amount of a polypeptide as described herein in a sample from said subject; and
- b) comparing the amount of said polypeptide to the amount of nucleic acid present
25 in a control sample from a subject without lupus nephritis,

wherein an increase in the level of said polypeptide as compared to the level of the polypeptide in the control sample indicates the presence or predisposition to lupus nephritis in said subject.

30 In a further aspect, the invention provides a method of determining the presence of or predisposition of a BFLP1698 -associated disorder in a subject. The method includes providing a nucleic acid sample, *e.g.*, RNA or DNA, or both, from the subject and measuring the amount of a nucleic acid as described herein in the subject nucleic acid sample. The amount of said nucleic acid sample in the subject nucleic acid sample is then compared to the amount of a said nucleic acid in a control sample. An alteration in the amount of said nucleic acid in the sample
35 relative to the amount of said in the control sample indicates the subject has a tissue proliferation-associated disorder.

2003295784 25 May 2009

Thus in one aspect, the present invention provides a method for determining the presence of or predisposition to lupus nephritis in a subject, the method comprising:

a) measuring the amount of a nucleic acid molecule as described herein in a sample from said subject; and

b) comparing the amount of said nucleic acid in step to the amount of the nucleic acid present in a control sample from a subject without lupus nephritis,

wherein an increase in the level of said nucleic acid in step (a) as compared to the level of the nucleic acid in the control sample indicates the presence of or predisposition to lupus nephritis in said subject.

In a still further aspect, the invention provides a method of treating or preventing or delaying a BFLP1698-associated disorder. The method includes administering to a subject in which such treatment or prevention or delay is desired a nucleic acid as described herein, a polypeptide as described herein, or an antibody to a polypeptide as described herein in an amount sufficient to treat, prevent, or delay a tissue proliferation-associated disorder in the subject. Examples of such disorders include rheumatoid arthritis and multiple sclerosis.

Thus in another aspect, the present invention provides a method of treating lupus nephritis in a subject, the method comprising administering to said subject a therapeutically effective amount of an agent that inhibits activity of a polypeptide as described herein in said subject.

The present invention also provides a method for screening for a therapeutic agent for treating an autoimmune disorder, the method comprising:

contacting a test compound with a polypeptide as described herein; and
determining if said test compound binds to said polypeptide as described

herein,

wherein binding of said test compound to said polypeptide indicates the test compound is a therapeutic agent for an autoimmune disorder.

The present invention further provides a pharmaceutical composition comprising an agent that inhibits activity of a polypeptide as described herein in a subject and a pharmaceutically acceptable carrier.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be

used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In the case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

Other features and advantages of the invention will be apparent from the following detailed description and claims.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a histogram showing relative levels of gene expression in the mouse ortholog of the human BFLP1698 gene in NZB x NZWF1 kidneys before, during, and after rapamycin treatment, as well as in various control mouse strains and conditions.

DETAILED DESCRIPTION OF THE INVENTION

The BFLP1698 nucleic acid sequences disclosed herein were identified based on changes in expression of the gene in kidneys of a lupus nephritis model mouse as compared to expression of the gene in kidneys from non-diseased mice. More particularly, the gene is expressed at relatively low levels in young mice and mice that do not show symptoms of lupus nephritis. Gene expression is elevated in mice with lupus nephritis, and is lower in mice that have been successfully treated with rapamycin or anti-B7 antibodies. The observation that expression levels return to normal when kidney function is normal indicates that elevated levels are related to, and diagnostic of, disease progression. Blocking the function of these genes may inhibit or retard disease progression. Expression levels can also be used to assess and compare effectiveness of various therapeutic interventions.

Accordingly, the BFLP1698 nucleic acid sequences are useful for detecting the presence of lupus nephritis in a subject. Elevated levels of BFLP1698 transcripts or polypeptides relative to levels in control samples indicate the presence of lupus nephritis in the subject. BFLP1698 nucleic acid sequences can also be used to monitor the effectiveness of treatments for lupus

nephritis: a decrease in expression of BFLP1698 genes relative to levels in diseased treatments demonstrates that the treatment is effective.

The BFLP1698 sequences can additionally be used to identify therapeutic agents for treating or preventing lupus nephritis in a subject. For example, a BFLP1698 polypeptide can be contacted with a test agent. Binding of the BFLP1698 polypeptide to the test agent reveals that the test agent modulates BFLP1698 activity. The BFLP1698-binding agent can be further tested to determine if it acts to promote or inhibit lupus symptoms in a test organism (e.g., a NZB X NZW mouse). Inhibition of lupus symptoms reveals that the agent is useful for treating or preventing lupus nephritis, or symptoms associated with lupus nephritis. Additional utilities are disclosed herein.

A 3652 nucleotide sequence that includes a human BFLP1698 nucleic acid is shown in Table 1 (SEQ ID NO:1). The human sequence was identified as the human ortholog of a murine gene whose expression is increased in a NZB X NZW mouse with lupus nephritis-like symptoms.

Nucleotides 1-3486 of the sequence shown in Table 1 encode a polypeptide of 1162 amino acids, whose sequence is shown in Table 2 (SEQ ID NO:2).

Table 1

ATGGCCCTTGTGCCAGGGAGAAGCAAGGAGGATGGGCTTTGGACTAGAAATAGCCCAGGCTCCTCCCAGCATCCAGAA
 AGTCCCAGGCTGCCCAACCTCTCTGGGACAGAGGAAAAATTGGCAAGGTTGAAGGTCACCAGCACATTCAGGATTTT
 TCTCAAAAGTCCCATCTGCCGTCTATTGTGGTGGAATCCAGTGAGGTGAATGAAGAGAGTGGGGATCTCCATTTGCCC
 CATGAGGAGCTGCTGCTGCTCACTGATGGTGAGGAAGAGGATGCTGAGGCCTTCTTCCAAGACCAAAGTGAAGAGCCA
 GGGGCGGCACGTCCCACATCAGGCTCGGCAAGTGGAGCATTCGACGCAGCGCGGCCATCTGGAGATTCCGGGAGCTG
 AAGAAGAAGCTGTTCAAACGCCGGCGGGTGTGAATCGGGAGCGGCGTCTGAGGCACCGGGTGGTCCGGGCTGTGATA
 GACCAAGGGCTGATCAGCGGCACCACTCAAGAAGCGGGCTGCTCAGGAGCTGTCCCAGGAAATCAAGGCTTTTCTG
 ACTGGCGTAGACCCCATTTCTGGGCCACCAACTCTCAGCCCGGAACATGCTCGCTGTGGTCTTCTCCTGCTCCGTTCT
 TTGCCACCTGCTCGGGCTGCTGTGCTTGACCACTTGAGAGGTGTCTTTGATGAGAGTGTCCGGGCCACCTGGCTGCC
 CTGGATGAAACCCCTGTGGCTGGTCCACCTCACCTCCGTCCACCTCCACCTCTCATGTCCCTGCTGGTGGACCTGGT
 CTAGAGGATGTGGTTAGGAAGTGCAGCAGGTGCTGTCTGAGTTTATCCGGGCCAACCCAAAGGCCTGGGCACCTGTG
 ATTAGTGATGGTCCATTGACCTCATGGGGCAACTGAGCAGCAGTACTCAGGCCAGCACCAGCGTGTCCCCACGCT
 ACTGGCGCTCTTAATGAAGTGTACAGCTGGATGGGTTGTAGGGCCACGCGTACATTAATGGACATCTATGTGCAG
 TGCTCTCGGCTCTCATTTGGTAGCTGCCAGATGCGTGTGTGGATGCCTTGCTGGATACCTCTGTTTACGATTTCTCCA
 CACTTTGACTGGGTTGTGGCACATATTGGCTCCTCTTTTCTGGCACCATCATTTCCCGGGTTCTCTCTGTGGCCTT
 AAGGACTTTTGTGTCCATGGTGGGGCTGGAGGTGGAGCTGGCAGTAGTGGTGGAAAGCTCTTCTCAGACCCCTCTACA
 GACCCCTTCCCTGGATCTCCTGCCATTCTTGCGGAGAAACGGGTGCCCAAGATTGCCTCAGTTGTAGGCATCCTAGGT
 CACCTGGCCTCCGCCACGGAGATAGCATCCGACGGGAGCTCCTGCGAATGTTCCATGATAGCTGGCAGGGGGATCT
 GGAGGCCGAGTGGGGACCCCTCCCTTCAGGCCACGGTTCCGTTCCTACTGCAGCTGGCAGTCATGTACACAGCTTTG
 CTGGGCACTGTCTCTGGAGAGCTTGTGGATTGCCTCAAGCCCCAGCTGTGCTGAGCCAGCTGCAGCAACACCTTCAA
 GGATTTCCCCCGAGAGGAGCTGGACAACATGTTGAACCTGGCTGTGCACCTGGTGGAGCCAGGCCTCTGGGGCAGGTGCC
 TACCGCTTGCTGCAGTTCTTGGTGGACACAGCTATGCCTGCTTCGGTCATTACCACCCAGGGCCTGGCTGTGCCAGAC
 ACCGTGCGTGAGGCTTGTGACCGGCTAATCCAGCTGCTGCTGCTGCACCTGCAAAAAGTGGTTCATCACCAGGGGAGGG
 TCTCCTGGGGAAGGGTGCTAGGCCCGCCCCACCTCCCCGCTTGGTGCCCTTTTATAGATGCGCTCAAAAACCATGTT

5 GGAGAGCTGTGTGGAGAGACGTTACGATTGGAACGGAAGCGCTTCCTCTGGCAGCACCAGCTCTTGGGCCTGCTGTCT
 GTCTATACCCGGCCTAGCTGTGGACCTGAGGCCCTTGGGCCATCTGCTGAGCCGAGCCGAAGCCCTGAAGAGTTGAGT
 TTGGCCACCCAGTTATATGACAGGGCTAGTGGTCAGCCTCTCTGGCCCTCTGCCCCCTGGCTTTCCGAAGCTGTCTGGCT
 CGGGTGCATGCAGGGACATTACAGCCTCCCTTCACGGCCCGGTTCTGCGCAACTTGGCACTGCTAGTAGGGTGGGAA
 CAGCAGGGTGGCGAGGGCCCTGCAGCCCTAGGGGCGCACTTTGGGGAATCTGCCTCAGCCCATCTGTCTGACCTGGCT
 CCTCTCCTGCTACATCCTGAGGAGGAAGTAGCTGAAGCTGCTGCCTCTCTCCTGGCCATTTGTCCCTTTCTCTTGAA
 GCCTTATCCCCCTCCCAGCTCCTGGGACTGGTAAGGGCTGGGGTGCACCGCTTCTTTGCCTCTCTGAGGCTGCATGGA
 CCCCCAGGTGTGGCCTCAGCCTGTGAGCTTCTACCCGCCTGTCTCAGACATCCCAGCTGGGCTCAAGGCTGTCTG
 10 CAGCTGCTGGTTGAAGGAGCCTTACATCGAGGCAACACAGAAGTGTGGTGGGCAAGTAGATGGGGACAATGAGACT
 CTCTCAGTTGTTTTAGCTTCTTTGGCTTCTGCCTCCCTGTTGGACACTAACCGGAGGCACACTGCAGCTGTGCCAGGT
 CCTGGAGGGATTTGGTCAGTTTTCATGCTGGAGTCATCGGCCCTGGCTTAAAGCCACCAAGTTTGTCCAGTCACGA
 AATCAGCAGGAAGTGATCTATAACACCCAGAGCCTCCTCAGCCTCCTGGTTCACTGCTGCAGTGCCCCAGGGGGCACT
 GAATGTGGGGAATGCTGGGGGACCCATCTTGTAGTCCAGAGGCAGCCAAAGCAGTGGCAGTGACCTTGGTGGAGAGT
 GTGTGTCCCGATGCAGCTGGTGCAGAGCTGGCCTGGCCCCCGAGGAACACGCCCGGGCCACCGTGGAGCGGGATCTC
 15 CGCATTTGGCCGGCGCTTCCGCGAACAGCCCCGCTCTTTGAGCTGTTAAAGCTGGTAGCAGCTGCACCCCCAGCCCTG
 TGCTACTGTTCCGTGCTGCTTCGGGGGCTGCTGGCCGCCCTCTTGGGCCATTGGGAAGCCTCTCGCCACCCTGACACG
 ACCCACTCCCCCTGGCACCTGGAGGCATCCTGCACCTTAGTGGCTGTCATGGCTGAGGGAAGCCTCCTGCCTCCGGCC
 CTGGGTAATATGCATGAAGTATTTAGCCAACCTGGCACCTTTTCAGGTGCGTCTGCTGCTCAGTGTCTGGGGTTTT
 CTCGGGAGCATGGGCCCTTGCTCAGAAGTTCATCTTCCAATCAGAGCGGGGTGCGTTTCATTCGGGACTTCTCCAGG
 20 GAGGGTGGAGGTGAGGGTGGACCCCATCTGGCTGTGCTGCACAGTGTCTCCACCGCAACATCGACCGCCTAGGTCTT
 TTCTCTGGCCGTTTCCAGGCACCTTCACCGTCCACTCTCCTTCGACAGGGGACGTAGCCTTTTCTTGCTCTGGAAGCC
 CAGGGAGGTTGAGCAGTGAGAGAGGGAAGGGACTAACGTGCTCCGGAAGGGTGGAGGTTTCTCTTCTAAGTCTTGGT
 CTAAGAGCGCTGTCACTTTTTTCTCTCCACTTTTTTTTTTCTAAATAAAATTTGCCAACTTG (SEQ ID NO:1)

25 **Table 2**

MALVPGRSKEDGLWTRNSPGSSQHPESPRLPNPLWDRGKIGKVEGHQHIQDFSQKSHLPSIVVESSEVNEESGDLHLP
 HEELLLLTGDEEEDAEAFQDQSEEPGAARPHHQARQVEHSTQRGHLEIRELKKKLFKRRRVLNRERRLRHRVVGAVI
 DQGLITRHHLLKKRAAQELSQEIKAFLTGVDPILGHQLSAREHARCGLLLRLSLPPARAVALDHLRGVFDSEVRAHLAA
 30 LDETPVAGPPHLRPPPPSHVPAGGPGLEDVQEVQVQLSEFIRANPKAWAPVISAWSIDLMOQLSSTYSQHQVRVPHA
 TGALNELLQLWMGCRATRTLMDIYVQCLSALIGSCPDACVDALLDTSVQHSPhFDWVVAHIGSSFPGTIIISVLSGCL
 KDFCVHGGAGGGAGSSGSSQTPSTDPFPGSPAIPAERKVPKIASVVGILGHLASRHGDSIRRELLRMFHDLAGGS
 GGRSGDPSLQATVPFLLQLAVMSPALLGTVSGELVDCLKPPAVLSQLQOHLQGFPREELDNMLNLAVHLVSQASGAGA
 YRLLQFLVDTAMPASVITTOGLAVPDTVREACDRLIQLLLLHLQKLVHHRGGSPGEGVLGPPPPPRLPVFLDALKNHV
 35 GELCGETLRLERKRFLWQHLLGLLSVYTRPSCGPEALGHLLSRARSPEELSLATQLYAGLVVSLGSLPLAFRSCLA
 RVHAGTLQPPFTARFLRNALLVGWEQQGGEPAALGAHFGESASAHLSDLAPLLLHPPEEEVAAEAASLLATCFPSE
 ALSPSOLLGLVRA'GVHRFFASRLRHGPPGVASACQLLRLSQTSPAGLKAVLQLLVEGALHRGNTEFGGQVDGNET
 LSVVSASLASASLLDTNRRHTAAVPGPGGIWSVFHAGVIGRGLKPPKFVQSRNQEVITYNTQSLLSLLVHCCSAPGGT
 ECGECWGAPILSPEAAKAVAVTLVESVCPDAAGAEALAWPPEEHARATVERDLRIGRRFREQLLFELLKLVAAPPAL
 40 CYCSVLLRGLLAALLGHWEASRHPDTHSPWHLEASCTLVAVMAEGSLLPPALGNMHEVFSQLAPFEVRLLLSVWGF
 LREHGPLPQKFIFQSERGRFIRDFSREGGEGGPHLAVLHLSVLHRNIDRLGLFSGRFQAPSPSTLLRQGT (SEQ ID
 NO:2)

BFLP1698-like nucleic acids and polypeptides of the invention (including those shown in Table 1) are referred to herein as "BFLP1698" nucleic acids and polypeptides.

45 A BFLP1698 nucleic acid, and the encoded polypeptide, according to the invention are useful in a variety of applications and contexts.

BFLP1698 shows homology to other proteins as shown in the BLAST results described in Table 3. KIAA0169, IMAGE: 3461492, and 3598686, and cDNA: FLJ21639 are all proteins

encoded from partial reading frames (expressed sequence tags (ESTs)) found in genomic DNA. Because BFLP1698 has homology to these proteins, it is also encoded from either an entire open reading frame, or part of a larger open reading frame (EST).

Table 3: Blast Results for KIAA1698

Gene Index/ Identifier	Protein/ Organism	Length (aa)	Identity (%)	Positives (%)	Expect
gi 20480454 ref XP_167747.1 (XM_167747)	similar to KIAA1698 protein [Homo sapiens]	1019	823/993 (82%)	824/993 (82%)	0.0
gi 12697941 dbj BAB21789.1 (AB051485)	KIAA1698 protein [Homo sapiens]	908	770/902 (85%)	770/902 (85%)	0.0
gi 20380408 gb AAH28025.1 (BC028025)	Similar to KIAA1698 protein [Homo sapiens]	833	705/829 (85%)	705/829 (85%)	0.0
gi 20849521 ref XP_144111.1 (XM_144111)	similar to KIAA1698 protein [Homo sapiens] [Mus musculus]	963	169/229 (73%)	175/229 (75%)	3e-87
gi 21296297 gb EAA08442.1 (AAAB01008879)	agCP2919 [Anopheles gambiae str. PEST]	900	80/292 (27%)	121/292 (41%)	1e-16

5 Table 4 shows a ClustalW alignment of BFLP1698 (SEQ ID NO:2) against the proteins described above in Table 3.

Table 4. ClustalW Analysis of SEQ ID NO:2

1) SEQ ID NO:2	
2) gi 20480454 ref XP_167747.1 (XM_167747)	(SEQ ID NO:16)
3) gi 12697941 dbj BAB21789.1 (AB051485)	(SEQ ID NO:17)
4) gi 20380408 gb AAH28025.1 (BC028025)	(SEQ ID NO:18)
5) gi 20849521 ref XP_144111.1 (XM_144111)	(SEQ ID NO:19)
6) gi 21296297 gb EAA08442.1 (AAAB01008879)	(SEQ ID NO:20)
10	
15	
20	
25	
30	
35	

```

      10      20      30      40      50      60
MALVPGRSKEDGLMTRNSPGSSQHPESPRLPNELWDRGKIGKVEGHHQIDFSQKSHLPS 60
gi|20480454|ref -----
gi|12697941|dbj -----
gi|20380408|gb| -----
gi|20849521|ref -----MILMITLFTTATFLVLGVSVVVLKEILTVMVPPPIQVRVFMHLYFFQ 50
gi|21296297|gb| -----
      70      80      90      100     110     120
IVVESSEVNEESGDLHLPHEELLLLTDGEEEDAEAFFDQDSEEPGAARPHHQARQVEHST 120
gi|20480454|ref -----MSALCDPEGAPGPPGPAPATHG-- 22
gi|12697941|dbj -----
gi|20380408|gb| -----
gi|20849521|ref LTIALGNVLERMKICPMRFFCFIQDLLVSKNNFGVLVKNMHFGTIPVRLFQPKATSSGP 110
gi|21296297|gb| -----
      130     140     150     160     170     180
QRGHLEIRELKKKLFKARRVLNRERRLRHVVGAVIDQGLITRHHLLKRAAQELSQEIK 180
SEQ ID NO:2 -----

```

5	gi 20480454 ref	-----PAP--LS-----AQELSQEIKA	37
	gi 12697941 dbj	-----	1
	gi 20380408 gb	-----	1
	gi 20849521 ref	RKGLIFYHGGGVFGSLDSYHNTCSYLAHET-----DSVVMVAVGYRK	152
	gi 21296297 gb	-----	1
10	SEQ ID NO:2190.....200.....210.....220.....230.....240.....	
	gi 20480454 ref	FLTGVDPI LGHQLSAREHARCGLLLRLSLPPARA AVL DHLRGV FDES VRAHLAALDETPV	240
	gi 12697941 dbj	FLTGVDPI LGHQLSAREHARCGLLLRLSLPPARA AVL DHLRGV FDES VRAHLAALDETPV	97
	gi 20380408 gb	-----	1
	gi 20849521 ref	LPDHHPTAYHDCLNATVHFLKELKTYGVDPARVVVSGESIGAGAAAIQAQVVLARKDLP	212
15	gi 21296297 gb	-----	1
20	SEQ ID NO:2250.....260.....270.....280.....290.....300.....	
	gi 20480454 ref	AGPPHLRPPPPSHVPAGGPGL EDVVQEVQVLS EETIRANPKAWAPVISAWSIDLMG----	296
	gi 12697941 dbj	AGPPHLRPPPPSHVPAGGPGL EDVVQEVQVLS EETIRANPKAWAPVISAWSIDLMG----	153
	gi 20380408 gb	-----PAGGPGL EDVVQEVQVLS EETIRANPKAWAPVISAWSIDLMG----	42
	gi 20849521 ref	QFRAQVLINPVQGVNFQLPSPYQYSDVPLSRRLMTCAKYLALDQSWKDAMLKGTFFI	272
25	gi 21296297 gb	-----KNLPDPSPVDPAVQETHEALERLVTVGETAWCCEVTSWCLKILG----	44
30	SEQ ID NO:2310.....320.....330.....340.....350.....360.....	
	gi 20480454 ref	--QLSSTVSGQHQR--VPHATGALNELLLQWMGCRATRTLMDIYVQCLSALIGSCPDACV	352
	gi 12697941 dbj	--QLSSTVSGQHQR--VPHATGALNELLLQWMGCRATRTLMDIYVQCLSALIGSCPDACV	209
	gi 20380408 gb	--QLSSTVSGQHQR--VPHATGALNELLLQWMGCRATRTLMDIYVQCLSALIGSCPDACV	98
	gi 20849521 ref	PPDHWKKAKWLSSDNTEQRFKSGQROPEFPGPFNESAYLETNHIFSLETSPLLADKIT	332
35	gi 21296297 gb	--EVCKKHCRRRPP-----DIRG---ACNLWLGC SATRYLLSTLSALCFERLDQREMBEIT	94
40	SEQ ID NO:2370.....380.....390.....400.....410.....420.....	
	gi 20480454 ref	DALLDT--SVQHS PHFDWVVAHIGSSFPGTIIISRVLS CGLKDFCVHGGAGGGAGSSGGS	409
	gi 12697941 dbj	DALLDT--SVQHS PHFDWVVAHIGSSFPGTIIISRVLS CGLKDFCVHGGAGGGAGSSGGS	266
	gi 20380408 gb	DALLDT--SVQHS PHFDWVVAHIGSSFPGTIIISRVLS CGLKDFCVHGGAGGGAGSSGGS	155
	gi 20849521 ref	DALLDT--SVQHS PHFDWVVAHIGSSFPGTIIISRVLS CGLKDFCVHGGAGGGAGSSGGS	80
45	gi 21296297 gb	AQLPETFLVSS EYDVL RD TLLYKKRL EEQGVPTWLVVGLPDVRVVP-LSQCPRAPGPP	391
		NEMVIYIG--THTEFDWVVARIGGCFFLRVMSMLSMGVARFTGDFDQPSSEVEVLS	151
50	SEQ ID NO:2430.....440.....450.....460.....470.....480.....	
	gi 20480454 ref	SSQTPSTDFFPGSPAIPA EKRVPKIASVVGILGH--LASRHGDSIRRELLRMFHDSL AGGS	468
	gi 12697941 dbj	SSQTPSTDFFPGSPAIPA EKRVPKIASVVGILGH--LASRHGDSIRRELLRMFHDSL AGGS	325
	gi 20380408 gb	SSQTPSTDFFPGSPAIPA EKRVPKIASVVGILGH--LASRHGDSIRRELLRMFHDSL AGGS	214
	gi 20849521 ref	SSQTPSTDFFPGSPAIPA EKRVPKIASVVGILGH--LASRHGDSIRRELLRMFHDSL AGGS	139
55	gi 21296297 gb	GPAPATHGEVPLSAQELSQEIKAFLTGVDPI LGHQLSAREHAQCGLLLLRLSLPPAQAVAL	451
		YLG LAHESDLRKALKSTL EHVASYKQPIPYLL--MLAKASETTSQALVAVFL ELHD---	205
60	SEQ ID NO:2490.....500.....510.....520.....530.....540.....	
	gi 20480454 ref	GGRSG--DPSLQATVPFLLQLAVMSPALLGTVSGELVDCLKP--PAVLSQLQOHLQGFPR	524
	gi 12697941 dbj	GGRSG--DPSLQATVPFLLQLAVMSPALLGTVSGELVDCLKP--PAVLSQLQOHLQGFPR	381
	gi 20380408 gb	GGRSG--DPSLQATVPFLLQLAVMSPALLGTVSGELVDCLKP--PAVLSQLQOHLQGFPR	270
	gi 20849521 ref	GGRSG--DPSLQATVPFLLQLAVMSPALLGTVSGELVDCLKP--PAVLSQLQOHLQGFPR	195
65	gi 21296297 gb	DHLRGV FDES VRAHLAALDETPVAGP PHLRPPSPSHVPTGGEGLEDVVHEVQCVLCETIR	511
		EN-----RLPTLTVL EKNWPANIGL EYVLTVAQLLLKMKKHAI RVTLLILAKMSTCHSWC	260
70	SEQ ID NO:2550.....560.....570.....580.....590.....600.....	
	gi 20480454 ref	E-----ELDNMLNLAVHLVSQASGAG-----	545
	gi 12697941 dbj	E-----ELDNMLNLAVHLVSQASGAG-----	402
	gi 20380408 gb	E-----ELDNMLNLAVHLVSQASGAG-----	291
	gi 20849521 ref	E-----ELDNMLNLAVHLVSQASGAG-----	216
75	gi 21296297 gb	ANPKVWAPVISAWSIDLMGQLSSTYSQGHQRVPHATGSRNELLLQWMSCRDTRTLMDIYV	571
		Q-----ELLEMMFTELETIVLDKHTA-----	281
80	SEQ ID NO:2610.....620.....630.....640.....650.....660.....	
		-----AYRLI-----	550

10

5	gi 20480454 ref	VTLVESVCPDAAGAEIANPPEEHARATVERDLRIGRRFREQPLLFECLKLVAAAPPALCY	873
	gi 12697941 dbj	VTLVESVCPDAAGAEIANPPEEHARATVERDLRIGRRFREQPLLFECLKLVAAAPPALCY	762
	gi 20380408 gb	VTLVESVCPDAAGAEIANPPEEHARATVERDLRIGRRFREQPLLFECLKLVAAAPPALCY	687
	gi 20849521 ref	VTLVESVCPDAAGAEIANPPEEHARATVERDLRIGRRFREQPLLFECLKLVAAAPPALCY	963
	gi 21296297 gb	VTLVESVCPDAAGAEIANPPEEHARATVERDLRIGRRFREQPLLFECLKLVAAAPPALCY	743
10	SEQ ID NO:2	CSVLLRGLLAALLGHWEASRHPDIT--HSEWHLEASCTLVAVMAEGSLLPPALGNMHEVFS	1075
	gi 20480454 ref	CSVLLRGLLAALLGHWEASRHPDIT--HSEWHLEASCTLVAVMAEGSLLPPALGNMHEVFS	932
	gi 12697941 dbj	CSVLLRGLLAALLGHWEASRHPDIT--HSEWHLEASCTLVAVMAEGSLLPPALGNMHEVFS	821
	gi 20380408 gb	CSVLLRGLLAALLGHWEASRHPDIT--HSEWHLEASCTLVAVMAEGSLLPPALGNMHEVFS	746
	gi 20849521 ref	CSVLLRGLLAALLGHWEASRHPDIT--HSEWHLEASCTLVAVMAEGSLLPPALGNMHEVFS	963
15	gi 21296297 gb	CSVLLRGLLAALLGHWEASRHPDIT--HSEWHLEASCTLVAVMAEGSLLPPALGNMHEVFS	803
20	SEQ ID NO:2	QLAPFEVF--LLLLSVNGFLREHGFLPQKEIFQSERGRFIRDFS--REGGGEGGPHLAVLHS	1133
	gi 20480454 ref	QLAPFEVF--LLLLSVNGFLREHGFLPQKEIFQSERGRFIRDFS--REGGGEGGPHLAVLHS	990
	gi 12697941 dbj	QLAPFEVF--LLLLSVNGFLREHGFLPQKEIFQSERGRFIRDFS--REGGGEGGPHLAVLHS	879
	gi 20380408 gb	QLAPFEVF--LLLLSVNGFLREHGFLPQKEIFQSERGRFIRDFS--REGGGEGGPHLAVLHS	804
	gi 20849521 ref	QLAPFEVF--LLLLSVNGFLREHGFLPQKEIFQSERGRFIRDFS--REGGGEGGPHLAVLHS	963
25	gi 21296297 gb	YFDGPE--DAYF--KECV--VYKDI--VPSV--VCDPTGFHW--RDPLTS--PPPLQYTN--LRNT--MOK	863
30	SEQ ID NO:2	VLHRN--IDRLGLFSGRFOAPSPSTLLRQGT-----	1162
	gi 20480454 ref	VLHRN--IDRLGLFSGRFOAPSPSTLLRQGT-----	1019
	gi 12697941 dbj	VLHRN--IDRLGLFSGRFOAPSPSTLLRQGT-----	908
	gi 20380408 gb	VLHRN--IDRLGLFSGRFOAPSPSTLLRQGT-----	833
	gi 20849521 ref	VLHRN--IDRLGLFSGRFOAPSPSTLLRQGT-----	963
35	gi 21296297 gb	K--TKVGHLYHQMFVGP--EL--NS--NSG--PTQ--PLVQ--	900

Residues 1-170 of SEQ ID NO:2 are referred to herein as SEQ ID NO:15. The fragment of
 35 SEQ ID NO:16 that includes amino acids 1-27 is referred to herein as SEQ ID NO:21.

BFLP1698 Nucleic Acids

The nucleic acids of the invention include those that encode a BFLP1698 polypeptide or protein. As used herein, the terms polypeptide and protein are interchangeable.

In some embodiments, a BFLP1698 nucleic acid encodes a mature BFLP1698
 40 polypeptide. As used herein, a "mature" form of a polypeptide or protein described herein relates to the product of a naturally occurring polypeptide or precursor form or proprotein. The naturally occurring polypeptide, precursor or proprotein includes, by way of nonlimiting example, the full length gene product, encoded by the corresponding gene. Alternatively, it may be defined as the polypeptide, precursor or proprotein encoded by an open reading frame described herein. The
 45 product "mature" form arises, again by way of nonlimiting example, as a result of one or more naturally occurring processing steps that may take place within the cell in which the gene product arises. Examples of such processing steps leading to a "mature" form of a polypeptide or protein

include the cleavage of the N-terminal methionine residue encoded by the initiation codon of an open reading frame, or the proteolytic cleavage of a signal peptide or leader sequence. Thus a mature form arising from a precursor polypeptide or protein that has residues 1 to N, where residue 1 is the N-terminal methionine, would have residues 2 through N remaining after removal of the N-terminal methionine. Alternatively, a mature form arising from a precursor polypeptide or protein having residues 1 to N, in which an N-terminal signal sequence from residue 1 to residue M is cleaved, would have the residues from residue M+1 to residue N remaining. Further as used herein, a "mature" form of a polypeptide or protein may arise from a step of post-translational modification other than a proteolytic cleavage event. Such additional processes include, by way of non-limiting example, glycosylation, myristoylation or phosphorylation. In general, a mature polypeptide or protein may result from the operation of only one of these processes, or a combination of any of them.

The invention includes mutant or variant nucleic acids of SEQ ID NO:1, or a fragment thereof, any of whose bases may be changed from the corresponding bases shown in SEQ ID NO:1, while still encoding a protein that maintains at least one of its BFLP1698 -like activities and physiological functions (*i.e.*, modulating angiogenesis, neuronal development). The invention further includes the complement of the nucleic acid sequence of SEQ ID NO:1, including fragments, derivatives, analogs and homologs thereof. The invention additionally includes nucleic acids or nucleic acid fragments, or complements thereto, whose structures include chemical modifications.

One aspect of the invention pertains to isolated nucleic acid molecules that encode BFLP1698 proteins or biologically active portions thereof. Also included are nucleic acid fragments sufficient for use as hybridization probes to identify BFLP1698 -encoding nucleic acids (*e.g.*, BFLP1698 mRNA) and fragments for use as polymerase chain reaction (PCR) primers for the amplification or mutation of BFLP1698 nucleic acid molecules. As used herein, the term "nucleic acid molecule" is intended to include DNA molecules (*e.g.*, cDNA or genomic DNA), RNA molecules (*e.g.*, mRNA), analogs of the DNA or RNA generated using nucleotide analogs, and derivatives, fragments and homologs thereof. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA.

“Probes” refer to nucleic acid sequences of variable length, preferably between at least about 10 nucleotides (nt), 100 nt, or as many as about, *e.g.*, 6,000 nt, depending on use. Probes are used in the detection of identical, similar, or complementary nucleic acid sequences. Longer length probes are usually obtained from a natural or recombinant source, are highly specific and much slower to hybridize than oligomers. Probes may be single- or double-stranded and designed to have specificity in PCR, membrane-based hybridization technologies, or ELISA-like technologies.

An “isolated” nucleic acid molecule is one that is separated from other nucleic acid molecules that are present in the natural source of the nucleic acid. Examples of isolated nucleic acid molecules include, but are not limited to, recombinant DNA molecules contained in a vector, recombinant DNA molecules maintained in a heterologous host cell, partially or substantially purified nucleic acid molecules, and synthetic DNA or RNA molecules. Preferably, an “isolated” nucleic acid is free of sequences which naturally flank the nucleic acid (*i.e.*, sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated BFLP1698 nucleic acid molecule can contain less than about 50 kb, 25 kb, 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. Moreover, an “isolated” nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material or culture medium when produced by recombinant techniques, or of chemical precursors or other chemicals when chemically synthesized.

A nucleic acid molecule of the present invention, *e.g.*, a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:1, or a complement thereof, can be isolated using standard molecular biology techniques and the sequence information provided herein. Using all or a portion of the nucleic acid sequence of SEQ ID NO:1 as a hybridization probe, BFLP1698 nucleic acid sequences can be isolated using standard hybridization and cloning techniques (*e.g.*, as described in Sambrook *et al.*, eds., MOLECULAR CLONING: A LABORATORY MANUAL 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989; and Ausubel, *et al.*, eds., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, NY, 1993.)

A nucleic acid of the invention can be amplified using cDNA, mRNA or alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. Furthermore, oligonucleotides corresponding to
5 BFLP1698 nucleotide sequences can be prepared by standard synthetic techniques, *e.g.*, using an automated DNA synthesizer.

As used herein, the term "oligonucleotide" refers to a series of linked nucleotide residues, which oligonucleotide has a sufficient number of nucleotide bases to be used in a PCR reaction. A short oligonucleotide sequence may be based on, or designed from, a genomic or cDNA
10 sequence and is used to amplify, confirm, or reveal the presence of an identical, similar or complementary DNA or RNA in a particular cell or tissue. Oligonucleotides comprise portions of a nucleic acid sequence having about 10 nt, 50 nt, or 100 nt in length, preferably about 15 nt to 30 nt in length. In one embodiment, an oligonucleotide comprising a nucleic acid molecule less than 100 nt in length would further comprise at least 6 contiguous nucleotides of SEQ ID NO:1,
15 or a complement thereof. Oligonucleotides may be chemically synthesized and may be used as probes.

In another embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule that is a complement of the nucleotide sequence shown in SEQ ID NO:1, or a portion of this nucleotide sequence. A nucleic acid molecule that is complementary to the
20 nucleotide sequence shown in SEQ ID NO:1 is one that is sufficiently complementary to the nucleotide sequence shown in SEQ ID NO:1 that it can hydrogen bond with little or no mismatches to the nucleotide sequence shown in SEQ ID NO:1, thereby forming a stable duplex.

As used herein, the term "complementary" refers to Watson-Crick or Hoogsteen base pairing between nucleotide units of a nucleic acid molecule, and the term "binding" means the
25 physical or chemical interaction between two polypeptides or compounds or associated polypeptides or compounds or combinations thereof. Binding includes ionic, non-ionic, Van der Waals, hydrophobic interactions, etc. A physical interaction can be either direct or indirect. Indirect interactions may be through or due to the effects of another polypeptide or compound. Direct binding refers to interactions that do not take place through, or due to, the effect of another
30 polypeptide or compound, but instead are without other substantial chemical intermediates.

Moreover, the nucleic acid molecule of the invention can comprise only a portion of the nucleic acid sequence of SEQ ID NO:1, *e.g.*, a fragment that can be used as a probe or primer, or a fragment encoding a biologically active portion of BFLP1698. Fragments provided herein are defined as sequences of at least 6 (contiguous) nucleic acids or at least 4 (contiguous) amino acids, a length sufficient to allow for specific hybridization in the case of nucleic acids or for specific recognition of an epitope in the case of amino acids, respectively, and are at most some portion less than a full length sequence. Fragments may be derived from any contiguous portion of a nucleic acid or amino acid sequence of choice. Fragments can include as many as 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 amino acids. Derivatives are nucleic acid sequences or amino acid sequences formed from the native compounds either directly or by modification or partial substitution. Analogs are nucleic acid sequences or amino acid sequences that have a structure similar to, but not identical to, the native compound but differs from it in respect to certain components or side chains. Analogs may be synthetic or from a different evolutionary origin and may have a similar or opposite metabolic activity compared to wild type.

Derivatives and analogs may be full length or other than full length, if the derivative or analog contains a modified nucleic acid or amino acid, as described below. Derivatives or analogs of the nucleic acids or proteins of the invention include, but are not limited to, molecules comprising regions that are substantially homologous to the nucleic acids or proteins of the invention, in various embodiments, by at least about 70%, 80%, 85%, 90%, 95%, 98%, or even 99% identity (with a preferred identity of 80-99%) over a nucleic acid or amino acid sequence of identical size or when compared to an aligned sequence in which the alignment is done by a computer homology program known in the art, or whose encoding nucleic acid is capable of hybridizing to the complement of a sequence encoding the aforementioned proteins under stringent, moderately stringent, or low stringent conditions. An exemplary program is the Gap program (Wisconsin Sequence Analysis Package, Version 8 for UNIX, Genetics Computer Group, University Research Park, Madison, WI) using the default settings, which uses the algorithm of Smith and Waterman.

A "homologous nucleic acid sequence" or "homologous amino acid sequence," or variations thereof, refer to sequences characterized by a homology at the nucleotide level or amino acid level as discussed above. Homologous nucleotide sequences encode those sequences

coding for isoforms of a BFLP1698 polypeptide. Isoforms can be expressed in different tissues of the same organism as a result of, for example, alternative splicing of RNA. Alternatively, isoforms can be encoded by different genes. In the present invention, homologous nucleotide sequences include nucleotide sequences encoding for a BFLP1698 polypeptide of species other
5 than humans, including, but not limited to, mammals, and thus can include, *e.g.*, mouse, rat, rabbit, dog, cat, cow, horse, and other organisms. Homologous nucleotide sequences also include, but are not limited to, naturally occurring allelic variations and mutations of the nucleotide sequences set forth herein. A homologous nucleotide sequence does not, however, include the nucleotide sequence encoding human BFLP1698 protein. Homologous nucleic acid
10 sequences include those nucleic acid sequences that encode conservative amino acid substitutions (see below) in SEQ ID NO:2, as well as a polypeptide having BFLP1698 activity. Biological activities of the BFLP1698 proteins are described below. A homologous amino acid sequence does not encode the amino acid sequence of a human BFLP1698 polypeptide.

The nucleotide sequence determined from the cloning of the human BFLP1698 gene
15 allows for the generation of probes and primers designed for use in identifying and/or cloning BFLP1698 homologues in other cell types, *e.g.*, from other tissues, as well as BFLP1698 homologues from other mammals. The probe/primer typically comprises a substantially purified oligonucleotide. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, 25, 50, 100, 150, 200, 250, 300, 350 or
20 400 or more consecutive sense strand nucleotide sequence of SEQ ID NO:1; or an anti-sense strand nucleotide sequence of SEQ ID NO:1; or of a naturally occurring mutant of SEQ ID NO:1.

Probes based on the human BFLP1698 nucleotide sequence can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. In various embodiments, the probe further comprises a label group attached thereto, *e.g.*, the label group can
25 be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells or tissue which misexpress a BFLP1698 protein, such as by measuring a level of a BFLP1698-encoding nucleic acid in a sample of cells from a subject *e.g.*, detecting BFLP1698 mRNA levels or determining whether a genomic BFLP1698 gene has been mutated or deleted.

A "polypeptide having a biologically active portion of BFLP1698 " refers to polypeptides exhibiting activity similar, but not necessarily identical to, an activity of a polypeptide of the present invention, including mature forms, as measured in a particular biological assay, with or without dose dependency. A nucleic acid fragment encoding a "biologically active portion of BFLP1698 " can be prepared by isolating a portion of SEQ ID NO:1 that encodes a polypeptide having a BFLP1698 biological activity (biological activities of the BFLP1698 proteins are described below), expressing the encoded portion of BFLP1698 protein (*e.g.*, by recombinant expression *in vitro*) and assessing the activity of the encoded portion of BFLP1698.

The invention also provides polymorphic forms of BFLP1698 nucleic acid sequences as well as methods of detecting polymorphic sequences in BFLP1698 sequences. The polymorphic forms include genomic sequences corresponding to exons and/or introns associated with BFLP1698.

Individuals carrying polymorphic alleles of the invention may be detected at either the DNA, the RNA, or the protein level using a variety of techniques that are well known in the art. The present methods usually employ pre-characterized polymorphisms. That is, the genotyping location and nature of polymorphic forms present at a site have already been determined. The availability of this information allows sets of probes to be designed for specific identification of the known polymorphic forms.

The genomic DNA used for the diagnosis may be obtained from any nucleated cells of the body, such as those present in peripheral blood, urine, saliva, buccal samples, surgical specimen, and autopsy specimens. The DNA may be used directly or may be amplified enzymatically in vitro through use of PCR or other in vitro amplification methods such as the ligase chain reaction (LCR), strand displacement amplification (SDA), self-sustained sequence replication (3SR), prior to mutation analysis.

The detection of polymorphisms in specific DNA sequences, can be accomplished by a variety of methods including, but not limited to, restriction-fragment-length-polymorphism detection based on allele-specific restriction-endonuclease cleavage, hybridization with allele-specific oligonucleotide probes, including immobilized oligonucleotides or oligonucleotide arrays, allele-specific PCR, mismatch-repair detection (MRD), binding of MutS protein,

denaturing-gradient gel electrophoresis (DGGE), single-strand-conformation-polymorphism detection, RNAase cleavage at mismatched base-pairs, chemical or enzymatic cleavage of heteroduplex DNA, methods based on allele specific primer extension, genetic bit analysis (GBA), the oligonucleotide-ligation assay (OLA), the allele-specific ligation chain reaction (LCR), gap-LCR, radioactive and/or fluorescent DNA sequencing using standard procedures well known in the art, and peptide nucleic acid (PNA) assays.

BFLP1698 Variants

The invention further encompasses nucleic acid molecules that differ from the nucleotide sequences shown in SEQ ID NO:1 due to the degeneracy of the genetic code. These nucleic acids thus encode the same BFLP1698 protein as that encoded by the nucleotide sequence shown in SEQ ID NO:1, *e.g.*, the polypeptide of SEQ ID NO:2. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence shown in SEQ ID NO:2.

In addition to the human BFLP1698 nucleotide sequence shown in SEQ ID NO:1, it will be appreciated by those skilled in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequences of BFLP1698 may exist within a population (*e.g.*, the human population). Such genetic polymorphism in the BFLP1698 gene may exist among individuals within a population due to natural allelic variation. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame encoding a BFLP1698 protein, preferably a mammalian BFLP1698 protein. Such natural allelic variations can typically result in 1-5% variance in the nucleotide sequence of the BFLP1698 gene. Any and all such nucleotide variations and resulting amino acid polymorphisms in BFLP1698 that are the result of natural allelic variation and that do not alter the functional activity of BFLP1698 are intended to be within the scope of the invention.

Moreover, nucleic acid molecules encoding BFLP1698 proteins from other species, and thus that have a nucleotide sequence that differs from the human sequence of SEQ ID NO:1 are intended to be within the scope of the invention. Nucleic acid molecules corresponding to natural allelic variants and homologues of the BFLP1698 cDNAs of the invention can be isolated based on their homology to the human BFLP1698 nucleic acids disclosed herein using the human

cDNAs, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions. For example, a soluble human BFLP1698 cDNA can be isolated based on its homology to human membrane-bound BFLP1698. Likewise, a membrane-bound human BFLP1698 cDNA can be isolated based on its homology to soluble human BFLP1698.

Accordingly, in another embodiment, an isolated nucleic acid molecule of the invention is at least 6 nucleotides in length and hybridizes under stringent conditions to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1. In another embodiment, the nucleic acid is at least 10, 25, 50, 100, 250, 500 or 750 nucleotides in length. In another embodiment, an isolated nucleic acid molecule of the invention hybridizes to the coding region. As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% homologous to each other typically remain hybridized to each other.

Homologs (*i.e.*, nucleic acids encoding BFLP1698 proteins derived from species other than human) or other related sequences (*e.g.*, paralogs) can be obtained by low, moderate or high stringency hybridization with all or a portion of the particular human sequence as a probe using methods well known in the art for nucleic acid hybridization and cloning.

Thus, the present invention also includes polynucleotides capable of hybridizing under reduced stringency conditions, more preferably stringent conditions, and most preferably highly stringent conditions, to polynucleotides described herein. Examples of stringency conditions are shown in the table below: highly stringent conditions are those that are at least as stringent as, for example, conditions A-F; stringent conditions are at least as stringent as, for example, conditions G-L; and reduced stringency conditions are at least as stringent as, for example, conditions M-R.

Table 5. Stringency Conditions

Stringency Condition	Polynucleotide Hybrid	Hybrid Length (bp) ¹	Hybridization Temperature and Buffer ²	Wash Temperature and Buffer ²
----------------------	-----------------------	---------------------------------	---	--

Stringency Condition	Polynucleotide Hybrid	Hybrid Length (bp) ¹	Hybridization Temperature and Buffer ^H	Wash Temperature and Buffer ^H
A	DNA:DNA	≥50	65°C; 1xSSC -or- 42°C; 1xSSC, 50% formamide	65°C; 0.3xSSC
B	DNA:DNA	<50	T _B *; 1xSSC	T _B *; 1xSSC 67°C; 0.3xSSC
C	DNA:RNA	≥ 50	67°C; 1xSSC -or- 45°C; 1xSSC, 50% formamide	
D	DNA:RNA	<50	T _D *; 1xSSC	T _D *; 1xSSC 70°C; 0.3xSSC
E	RNA:RNA	≥50	70°C; 1xSSC -or- 50°C; 1xSSC, 50% formamide	
F	RNA:RNA	<50	T _F *; 1xSSC	T _F *; 1xSSC 65°C; 1xSSC
G	DNA:DNA	≥50	65°C; 4xSSC -or- 42°C; 4xSSC, 50% formamide	
H	DNA:DNA	<50	T _H *; 4xSSC	T _H *; 4xSSC 67°C; 1xSSC
I	DNA:RNA	≥ 50	67°C; 4xSSC -or- 45°C; 4xSSC, 50% formamide	
J	DNA:RNA	<50	T _J *; 4xSSC	T _J *; 4xSSC 67°C; 1xSSC
K	RNA:RNA	≥ 50	70°C; 4xSSC -or- 50°C; 4xSSC, 50% formamide	
L	RNA:RNA	<50	T _L *; 2xSSC	T _L *; 2xSSC 50°C; 2xSSC
M	DNA:DNA	> 50	50°C; 4xSSC -or- 40°C; 6xSSC, 50% formamide	

Stringency Condition	Polynucleotide Hybrid	Hybrid Length (bp) ¹	Hybridization Temperature and Buffer ^H	Wash Temperature and Buffer ^H
N	DNA:DNA	<50	T _N *; 6xSSC	T _N *; 6xSSC 55°C; 2xSSC
O	DNA:RNA	> 50	55°C; 4xSSC -or- 42°C; 6xSSC, 50% formamide	
P	DNA:RNA	<50	T _P *; 6xSSC	T _P *; 6xSSC 60°C; 2xSSC
Q	RNA:RNA	> 50	60°C; 4xSSC -or- 45°C; 6xSSC, 50% formamide	
R	RNA:RNA	<50	T _R *; 4xSSC	T _R *; 4xSSC

1: The hybrid length is that anticipated for the hybridized region(s) of the hybridizing polynucleotides. When hybridizing a polynucleotide to a target polynucleotide of unknown sequence, the hybrid length is assumed to be that of the hybridizing polynucleotide. When polynucleotides of known sequence are hybridized, the hybrid length can be determined by aligning the sequences of the polynucleotides and identifying the region or regions of optimal sequence complementarity.

^H: SSPE (1xSSPE is 0.15M NaCl, 10mM NaH₂PO₄, and 1.25mM EDTA, pH 7.4) can be substituted for SSC (1xSSC is 0.15M NaCl and 15mM sodium citrate) in the hybridization and wash buffers; washes are performed for 15 minutes after hybridization is complete.

T_B* - T_R*: The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length should be 5-10°C less than the melting temperature (T_m) of the hybrid, where T_m is determined according to the following equations. For hybrids less than 18 base pairs in length, T_m(°C) = 2(# of A + T bases) + 4(# of G + C bases). For hybrids between 18 and 49 base pairs in length, T_m(°C) = 81.5 + 16.6(log₁₀Na⁺) + 0.41(%G+C) - (600/N), where N is the number of bases in the hybrid, and Na⁺ is the concentration of sodium ions in the hybridization buffer (Na⁺ for 1xSSC = 0.165 M).

Preferably, each such hybridizing polynucleotide has a length that is at least 25% (more preferably at least 50%, and most preferably at least 75%) of the length of the polynucleotide of the present invention to which it hybridizes, and has at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with the polynucleotide of the present invention to which it hybridizes, where sequence identity is determined by comparing the sequences of the hybridizing polynucleotides when aligned so as to maximize overlap and identity while minimizing sequence gaps.

A non-limiting example of stringent hybridization conditions is hybridization in a high salt buffer comprising 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 mg/ml denatured salmon sperm DNA at 65°C. This hybridization is followed by one or more washes in 0.2X SSC, 0.01% BSA at 50°C. An isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence of SEQ ID NO:1 corresponds to a naturally occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature (*e.g.*, encodes a natural protein).

In a second embodiment, a nucleic acid sequence that is hybridizable to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, or fragments, analogs or derivatives thereof, under conditions of moderate stringency is provided. A non-limiting example of moderate stringency hybridization conditions are hybridization in 6X SSC, 5X Denhardt's solution, 0.5% SDS and 100 mg/ml denatured salmon sperm DNA at 55°C, followed by one or more washes in 1X SSC, 0.1% SDS at 37°C.

In a third embodiment, a nucleic acid that is hybridizable to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, or fragments, analogs or derivatives thereof, under conditions of low stringency, is provided. A non-limiting example of low stringency hybridization conditions are hybridization in 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 mg/ml denatured salmon sperm DNA, 10% (wt/vol) dextran sulfate at 40°C, followed by one or more washes in 2X SSC, 25 mM Tris-HCl (pH 7.4), 5 mM EDTA, and 0.1% SDS at 50°C.

Conservative mutations

In addition to naturally-occurring allelic variants of the BFLP1698 sequence that may exist in the population, the skilled artisan will further appreciate that changes can be introduced by mutation into the nucleotide sequence of SEQ ID NO:1, thereby leading to changes in the amino acid sequence of the encoded BFLP1698 protein, without altering the functional ability of the BFLP1698 protein. For example, nucleotide substitutions leading to amino acid substitutions at "non-essential" amino acid residues can be made in the sequence of SEQ ID NO:1. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence of BFLP1698 without altering the biological activity, whereas an "essential" amino acid residue is required for biological activity. For example, altering amino acid residues that are conserved among the BFLP1698 proteins of the present invention, is likely to result in loss of activity of the BFLP1698 protein.

Another aspect of the invention pertains to nucleic acid molecules encoding BFLP1698 proteins that contain changes in amino acid residues that are not essential for activity. Such BFLP1698 proteins differ in amino acid sequence from SEQ ID NO:2, yet retain biological activity. In one embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence encoding a protein, wherein the protein comprises an amino acid sequence at least about 75% homologous to the amino acid sequence of SEQ ID NO:2. Preferably, the protein encoded by the nucleic acid is at least about 80% homologous to SEQ ID NO:2, more preferably at least about 90%, 95%, 98%, and most preferably at least about 99% homologous to SEQ ID NO:2.

An isolated nucleic acid molecule encoding a BFLP1698 protein homologous to the protein of SEQ ID NO:2 can be created by introducing one or more nucleotide substitutions, additions or deletions into the nucleotide sequence of SEQ ID NO:1, such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein.

Mutations can be introduced into the nucleotide sequence of SEQ ID NO:1 by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic

acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in BFLP1698 is replaced with another amino acid residue from the same side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of a BFLP1698 coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for BFLP1698 biological activity to identify mutants that retain activity. Following mutagenesis of SEQ ID NO:1 the encoded protein can be expressed by any recombinant technology known in the art and the activity of the protein can be determined.

In one embodiment, a mutant BFLP1698 protein can be assayed for (1) the ability to form protein:protein interactions with other BFLP1698 proteins, other cell-surface proteins, or biologically active portions thereof, (2) complex formation between a mutant BFLP1698 protein and a BFLP1698 receptor; (3) the ability of a mutant BFLP1698 protein to bind to an intracellular target protein or biologically active portion thereof; (*e.g.*, avidin proteins); (4) the ability to bind BFLP1698 protein; or (5) the ability to specifically bind an anti-BFLP1698 protein antibody.

Antisense BFLP1698 Nucleic Acids

Another aspect of the invention pertains to isolated antisense nucleic acid molecules that are hybridizable to or complementary to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, or fragments, analogs or derivatives thereof. An "antisense" nucleic acid comprises a nucleotide sequence that is complementary to a "sense" nucleic acid encoding a protein, *e.g.*, complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. In specific aspects, antisense nucleic acid molecules are provided that comprise a sequence complementary to at least about 10, 25, 50, 100, 250 or 500 nucleotides or an entire BFLP1698 coding strand, or to only a portion thereof. Nucleic acid molecules encoding fragments, homologs, derivatives and analogs of a BFLP1698 protein of

SEQ ID NO:2, or antisense nucleic acids complementary to a BFLP1698 nucleic acid sequence of SEQ ID NO:1 are additionally provided.

In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence encoding BFLP1698. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues (*e.g.*, the protein coding region of human BFLP1698 corresponds to SEQ ID NO:2). In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence encoding BFLP1698. The term "noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (*i.e.*, also referred to as 5' and 3' untranslated regions).

Given the coding strand sequences encoding BFLP1698 disclosed herein (*e.g.*, SEQ ID NO:1), antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick or Hoogsteen base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of BFLP1698 mRNA, but more preferably is an oligonucleotide that is antisense to only a portion of the coding or noncoding region of BFLP1698 mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of BFLP1698 mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic acid of the invention can be constructed using chemical synthesis or enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (*e.g.*, an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, *e.g.*, phosphorothioate derivatives and acridine substituted nucleotides can be used.

Examples of modified nucleotides that can be used to generate the antisense nucleic acid include: 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine,

7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (*i.e.*, RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a BFLP1698 protein to thereby inhibit expression of the protein, *e.g.*, by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule that binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention includes direct injection at a tissue site. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, *e.g.*, by linking the antisense nucleic acid molecules to peptides or antibodies that bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the

strands run parallel to each other. The antisense nucleic acid molecule can also comprise a 2'-o-methylribonucleotide.

Such modifications include, by way of nonlimiting example, modified bases, and nucleic acids whose sugar phosphate backbones are modified or derivatized. These modifications are carried out at least in part to enhance the chemical stability of the modified nucleic acid, such that they may be used, for example, as antisense binding nucleic acids in therapeutic applications in a subject.

BFLP1698 Ribozymes and PNA moieties

In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity that are capable of cleaving a single-stranded nucleic acid, such as a mRNA, to which they have a complementary region. Thus, ribozymes (*e.g.*, hammerhead ribozymes) can be used to catalytically cleave BFLP1698 mRNA transcripts to thereby inhibit translation of BFLP1698 mRNA. A ribozyme having specificity for a BFLP1698-encoding nucleic acid can be designed based upon the nucleotide sequence of a BFLP1698 DNA disclosed herein (*i.e.*, SEQ ID NO:1). For example, a derivative of a Tetrahymena L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in a BFLP1698-encoding mRNA. Alternatively, BFLP1698 mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules.

Alternatively, BFLP1698 gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of the BFLP1698 (*e.g.*, the BFLP1698 promoter and/or enhancers) to form triple helical structures that prevent transcription of the BFLP1698 gene in target cells.

In various embodiments, the nucleic acids of BFLP1698 can be modified at the base moiety, sugar moiety or phosphate backbone to improve, *e.g.*, the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acids can be modified to generate peptide nucleic acids. As used herein, the terms "peptide nucleic acids" or "PNAs" refer to nucleic acid mimics, *e.g.*, DNA mimics, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four

natural nucleobases are retained. The neutral backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using standard solid phase peptide synthesis protocols.

PNAs of BFLP1698 can be used in therapeutic and diagnostic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, *e.g.*, inducing transcription or translation arrest or inhibiting replication. PNAs of BFLP1698 can also be used, *e.g.*, in the analysis of single base pair mutations in a gene by, *e.g.*, PNA directed PCR clamping; as artificial restriction enzymes when used in combination with other enzymes, *e.g.*, S1 nucleases; or as probes or primers for DNA sequence and hybridization.

In another embodiment, PNAs of BFLP1698 can be modified, *e.g.*, to enhance their stability or cellular uptake, by attaching lipophilic or other helper groups to PNA, by the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery known in the art. For example, PNA-DNA chimeras of BFLP1698 can be generated that may combine the advantageous properties of PNA and DNA.

The oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane or the blood-brain barrier. In addition, oligonucleotides can be modified with hybridization triggered cleavage agents or intercalating agents. To this end, the oligonucleotide may be conjugated to another molecule, *e.g.*, a peptide, a hybridization triggered cross-linking agent, a transport agent, a hybridization-triggered cleavage agent, etc.

BFLP1698 Interfering Nucleic Acids

Also provided by the invention is an isolated double-stranded nucleic acid (DNA or RNA) that is capable of mediating specific inhibition of BFLP1698 gene expression. In preferred embodiments, one or both strands of the double-stranded molecule is an RNA molecule. Preferably, each RNA strand has a length from 19-25, particularly from 19-23 nucleotides, more particularly from 20-22 nucleotides, and is capable of mediating BFLP1698 target-specific nucleic acid modifications, particularly RNA interference and/or DNA methylation. The double-stranded BFLP1698 molecule may be double stranded or have an overhang at one or both the 5' and/or 3' terminus. For example, the molecule may have a 3'

overhang. The length of the 3'-overhang can be, e.g., 1-6 nucleotides, 2-5 nucleotides, 3-4 nucleotides, or 2 nucleotides. The length of the overhang may be the same or different for each strand. In one embodiment, dsRNAs are composed of two 21 nucleotide strands that are paired such that 1, 2, or 3 nucleotide overhangs are present on both ends of the double-stranded RNA.

5 The RNA strands preferably have 3'-hydroxyl groups. The 5'-terminus preferably includes a phosphate, diphosphate, triphosphate or hydroxyl group. If desired, the 3'-overhangs may be stabilized against degradation. For example, they may be selected such that they consist of purine nucleotides, particularly adenosine or guanosine nucleotides. Alternatively, pyrimidine nucleotides may be replaced with modified analogues, e.g. substitution of uridine -2 nucleotide
10 3' overhangs by 2'-deoxythymidine is tolerated, and does not affect the efficiency of RNA interference. The RNA molecule may contain at least one modified nucleotide analogue. The nucleotide analogues may be located at positions where the target-specific activity, e.g. the RNAi mediating activity is not substantially affected. The modified nucleotide is preferably present in a region at the 5'-end and/or the 3'-end of the double-stranded RNA molecule. In some
15 embodiments, overhangs are stabilized by incorporating modified nucleotide analogues.

Nucleotide analogues can include sugar- or backbone-modified ribonucleotides. Other suitable nucleotides include a non-naturally occurring nucleobase instead of a naturally occurring nucleobases. For example, analogues can include uridines or cytidines modified at the 5-position, e.g. 5-(2-amino)propyl uridine, 5-bromo uridine; adenosines and guanosines modified
20 at the 8-position, e.g. 8-bromo guanosine; deaza nucleotides, e.g. 7-deaza-adenosine; O- and N-alkylated nucleotides, e.g. N6-methyl adenosine are suitable. In preferred sugar-modified ribonucleotides the 2' OH-group is replaced by a group selected from H, OR, R, halo, SH, SR, NH₂, NHR, NR₂ or CN, wherein R is C₁-C₆ alkyl, alkenyl or alkynyl and halo is F, Cl, Br or I. In a preferred embodiment, where backbone-modified ribonucleotides are used as the phosphoester
25 group connecting to adjacent ribonucleotides, they are replaced by a modified group, e.g. a phosphothioate group. It should be noted that the above modifications may be combined.

The BFLP1698 interfering RNA molecule can be a naturally isolated RNA molecule or can be a synthetic RNA molecule. Preferably, the BFLP1698 interfering RNA molecule is substantially free from contaminants occurring in cell extracts, e.g. from *Drosophila* embryos.

Further, the BFLP1698 interfering RNA molecule is preferably substantially free from any non-target-specific contaminants, particularly non-target-specific RNA molecules e.g. from contaminants occurring in cell extracts.

Isolated double-stranded BFLP1698 interfering molecules can be used for mediating BFLP1698 target-specific nucleic acid modifications, particularly RNAi, in mammalian cells, particularly in human cells.

The sequence of the double-stranded BFLP1698 interfering molecule of the present invention is of sufficient identity to a nucleic acid BFLP1698 target molecule in order to effect target-specific interference of BFLP1698 gene expression and/or DNA methylation. Preferably, the sequence has an identity of at least 50%, particularly of at least 70% to the desired target molecule in the double-stranded portion of the RNA molecule. More preferably, the identity is at least 85% and most preferably 100% in the double-stranded portion of the RNA molecule. The identity of a BFLP1698 double-stranded interfering RNA molecule to a predetermined nucleic acid target molecule, e.g. an BFLP1698 mRNA target molecule with the sequence shown in SEQ ID NO:1, may be determined using the equation: $I = (n/L) \times 100$, wherein I is the identity in percent, n is the number of identical nucleotides in the double-stranded portion of the ds RNA and the target and L is the length of the sequence overlap of the double-stranded portion of the dsRNA and the target.

Alternatively, the identity of the double-stranded RNA molecule relative to the target sequence may also be defined including the 3' overhang, particularly an overhang having a length from 1-3 nucleotides. In this case the sequence identity is preferably at least 50%, more preferably at least 70% and most preferably at least 85% to the target sequence. For example, the nucleotides from the 3' overhang and up to 2 nucleotides from the 5' and/or 3' terminus of the double strand may be modified without significant loss of activity.

A double-stranded BFLP1698 RNA molecule may be prepared by a method that includes synthesizing two RNA strands each having a length from 19-25, e.g. from 19-23 nucleotides, wherein said RNA strands are capable of forming a double-stranded RNA molecule, wherein preferably at least one strand has a 3'-overhang from 1-5 nucleotides, and (b) combining the synthesized RNA strands under conditions, wherein a double-stranded RNA molecule is

formed. The double-stranded RNA molecule is capable of mediating target-specific nucleic acid modifications, particularly RNA interference and/or DNA methylation.

Methods of synthesizing RNA molecules are known in the art. The single-stranded RNAs can also be prepared by enzymatic transcription from synthetic DNA templates or from DNA
5 plasmids isolated from recombinant bacteria. Typically, phage RNA polymerases are used such as T7, T3 or SP6 RNA polymerase.

A further aspect of the present invention relates to a method of mediating BFLP1698-specific nucleic acid modifications, particularly RNA interference and/or DNA methylation in a cell or an organism by contacting the cell or organism with the double-stranded RNA molecule
10 of the invention under conditions wherein target-specific nucleic acid modifications may occur and mediating a target-specific nucleic acid modification effected by the double-stranded RNA towards a BFLP1698 target nucleic acid.

15 **BFLP1698 Polypeptides**

A BFLP1698 polypeptide of the invention includes the BFLP1698-like protein whose sequence is provided in SEQ ID NO:2. The invention also includes a mutant or variant form of the disclosed BFLP1698 polypeptide, or of any of the fragments of the herein disclosed BFLP1698 polypeptide sequences.

20 Thus, a BFLP1698 polypeptide includes one in which any residues may be changed from the corresponding residue shown in SEQ ID NO:2 while still encoding a protein that maintains its BFLP1698-like activities and physiological functions, or a functional fragment thereof. In some embodiments, up to 20% or more of the residues may be so changed in the mutant or variant protein. In some embodiments, the BFLP1698 polypeptide according to the invention is a
25 mature polypeptide.

Rapamycin Binding Domains

To identify regions of a BFLP1698 polypeptide sequence (e.g., a polypeptide including all or a portion of SEQ ID NO:2) containing rapamycin binding domains, the entire coding

sequence, or a fragment of a BFLP1698 polypeptide sequence, is tested for its ability to bind rapamycin. Any technique known in the art for determining binding of a polypeptide to a small molecule can be used. For example, rapamycin can be labeled (*i.e.*, with a non-radioactive label or with a radiolabel (e.g., ^{14}C , ^{32}P , ^3H , or ^{125}I), and mixed with a polypeptide containing some or all of a BFLP1698 polypeptide sequence. The polypeptide optionally includes a moiety that facilitates detection, e.g., the polypeptide can be a fusion polypeptide that includes a BFLP1698 sequence and a non-BFLP1698 polypeptide sequence.

A reagent specific for the polypeptide containing the BFLP1698 polypeptide sequence (e.g., an antibody specific for BFLP1698 or a probe specific for the non-BFLP1698 polypeptide in the case of a fusion polypeptide) is added to the mixture. Complexes that bind to the reagent are isolated, and the presence of label, which reveals the presence of rapamycin, is determined.

In general, a BFLP1698-like variant that preserves BFLP1698-like function includes any variant in which residues at a particular position in the sequence have been substituted by other amino acids, and further include the possibility of inserting an additional residue or residues between two residues of the parent protein as well as the possibility of deleting one or more residues from the parent sequence. Any amino acid substitution, insertion, or deletion is encompassed by the invention. In favorable circumstances, the substitution is a conservative substitution as defined above.

One aspect of the invention pertains to isolated BFLP1698 proteins, and biologically active portions thereof, or derivatives, fragments, analogs or homologs thereof. Fragments can comprise contiguous stretches of SEQ ID NO:2, or interspersed segments of SEQ ID NO:2. Also provided are polypeptide fragments suitable for use as immunogens to raise anti-BFLP1698 antibodies. In one embodiment, native BFLP1698 proteins can be isolated from cells or tissue sources by an appropriate purification scheme using standard protein purification techniques. In another embodiment, BFLP1698 proteins are produced by recombinant DNA techniques. Alternative to recombinant expression, a BFLP1698 protein or polypeptide can be synthesized chemically using standard peptide synthesis techniques.

A "purified" protein or biologically active portion thereof is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the BFLP1698

protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of BFLP1698 protein in which the protein is separated from cellular components of the cells from which it is isolated or recombinantly produced. In one embodiment, the language

5 "substantially free of cellular material" includes preparations of BFLP1698 protein having less than about 30% (by dry weight) of non-BFLP1698 protein (also referred to herein as a "contaminating protein"), more preferably less than about 20% of non-BFLP1698 protein, still more preferably less than about 10% of non-BFLP1698 protein, and most preferably less than about 5% non-BFLP1698 protein. When the BFLP1698 protein or biologically active portion

10 thereof is recombinantly produced, it is also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes preparations of BFLP1698 protein in which the protein is separated from chemical precursors or

15 other chemicals that are involved in the synthesis of the protein. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of BFLP1698 protein having less than about 30% (by dry weight) of chemical precursors or non-BFLP1698 chemicals, more preferably less than about 20% chemical precursors or non-BFLP1698 chemicals, still more preferably less than about 10% chemical precursors or

20 non-BFLP1698 chemicals, and most preferably less than about 5% chemical precursors or non-BFLP1698 chemicals.

Biologically active portions of a BFLP1698 protein include peptides comprising amino acid sequences sufficiently homologous to or derived from the amino acid sequence of the BFLP1698 protein, *e.g.*, the amino acid sequence shown in SEQ ID NO:2 that include fewer

25 amino acids than the full length BFLP1698 proteins, and exhibit at least one activity of a BFLP1698 protein. Typically, biologically active portions comprise a domain or motif with at least one activity of the BFLP1698 protein. A biologically active portion of a BFLP1698 protein can be a polypeptide which is, for example, 10, 25, 50, 100 or more amino acids in length.

A biologically active portion of a BFLP1698 protein of the present invention may contain

30 at least one of the above-identified domains conserved between the BFLP1698 proteins.

Moreover, other biologically active portions, in which other regions of the protein are deleted, can be prepared by recombinant techniques and evaluated for one or more of the functional activities of a native BFLP1698 protein.

In an embodiment, the BFLP1698 protein has an amino acid sequence shown in SEQ ID NO:2. In other embodiments, the BFLP1698 protein is substantially homologous to SEQ ID NO:2 and retains the functional activity of the protein of SEQ ID NO:2, yet differs in amino acid sequence due to natural allelic variation or mutagenesis, as described in detail below. Accordingly, in another embodiment, the BFLP1698 protein is a protein that comprises an amino acid sequence at least about 45% homologous to the amino acid sequence of SEQ ID NO:2 and retains the functional activity of the BFLP1698 proteins of SEQ ID NO:2.

Determining homology between two or more sequences

To determine the percent homology of two amino acid sequences or of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in either of the sequences being compared for optimal alignment between the sequences). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are homologous at that position (*i.e.*, as used herein amino acid or nucleic acid "homology" is equivalent to amino acid or nucleic acid "identity").

The nucleic acid sequence homology may be determined as the degree of identity between two sequences. The homology may be determined using computer programs known in the art, such as GAP software provided in the GCG program package. Using GCG GAP software with the following settings for nucleic acid sequence comparison: GAP creation penalty of 5.0 and GAP extension penalty of 0.3, the coding region of the analogous nucleic acid sequences referred to above exhibits a degree of identity preferably of at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, or 99%, with the CDS (encoding) part of the DNA sequence shown in SEQ ID NO:1.

The term "sequence identity" refers to the degree to which two polynucleotide or polypeptide sequences are identical on a residue-by-residue basis over a particular region of comparison. The term "percentage of sequence identity" is calculated by comparing two

optimally aligned sequences over that region of comparison, determining the number of positions at which the identical nucleic acid base (*e.g.*, A, T, C, G, U, or I, in the case of nucleic acids) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the region of comparison (*i.e.*, the window size), and multiplying the result by 100 to yield the percentage of sequence identity. The term "substantial identity" as used herein denotes a characteristic of a polynucleotide sequence, wherein the polynucleotide comprises a sequence that has at least 80 percent sequence identity, preferably at least 85 percent identity and often 90 to 95 percent sequence identity, more usually at least 99 percent sequence identity as compared to a reference sequence over a comparison region. The term "percentage of positive residues" is calculated by comparing two optimally aligned sequences over that region of comparison, determining the number of positions at which the identical and conservative amino acid substitutions, as defined above, occur in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the region of comparison (*i.e.*, the window size), and multiplying the result by 100 to yield the percentage of positive residues.

Chimeric and fusion proteins

The invention also provides BFLP1698 chimeric or fusion proteins. As used herein, a BFLP1698 "chimeric protein" or "fusion protein" comprises a BFLP1698 polypeptide operatively linked to a non-BFLP1698 polypeptide. A "BFLP1698 polypeptide" refers to a polypeptide having an amino acid sequence corresponding to BFLP1698, whereas a "non-BFLP1698 polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein that is not substantially homologous to the BFLP1698 protein, *e.g.*, a protein that is different from the BFLP1698 protein and that is derived from the same or a different organism. Within a BFLP1698 fusion protein the BFLP1698 polypeptide can correspond to all or a portion of a BFLP1698 protein. An example of a BFLP1698 fusion polypeptide is one that includes amino acids 21-230 of SEQ ID NO:2 (*e.g.*, a polypeptide that includes amino acids 1-246 or amino acids 21-246 of SEQ ID NO:2). In one embodiment, a BFLP1698 fusion protein comprises at least one biologically active portion of a BFLP1698 protein. In another embodiment, a BFLP1698 fusion protein comprises at least two biologically active portions of a BFLP1698

protein. Within the fusion protein, the term "operatively linked" is intended to indicate that the BFLP1698 polypeptide and the non-BFLP1698 polypeptide are fused in-frame to each other. The non-BFLP1698 polypeptide can be fused to the N-terminus or C-terminus of the BFLP1698 polypeptide.

5 For example, in one embodiment a BFLP1698 fusion protein comprises a BFLP1698 polypeptide operably linked to either an extracellular domain of a second protein, *i.e.*, non-BFLP1698 protein, or to the transmembrane and intracellular domain of a second protein, *i.e.*, non-BFLP1698 protein. Such fusion proteins can be further utilized in screening assays for compounds that modulate BFLP1698 activity (such assays are described in detail below).

10 In another embodiment, the fusion protein is a GST-BFLP1698 fusion protein in which the BFLP1698 sequences are fused to the C-terminus of the GST (*i.e.*, glutathione S-transferase) sequences. Such fusion proteins can facilitate the purification of recombinant BFLP1698.

 In another embodiment, the fusion protein is a BFLP1698-immunoglobulin fusion protein in which the BFLP1698 sequences comprising one or more domains are fused to sequences
15 derived from a member of the immunoglobulin protein family.

 Inhibition of the BFLP1698 ligand/BFLP1698 interaction can be used therapeutically for both the treatment of proliferative and differentiative disorders, *e.g.*, cancer, modulating (*e.g.*, promoting or inhibiting) cell survival as well as immunomodulatory disorders, autoimmunity, transplantation, and inflammation by alteration of cytokine and chemokine cascade
20 mechanisms. Moreover, the BFLP1698-immunoglobulin fusion proteins of the invention can be used as immunogens to produce anti-BFLP1698 antibodies in a subject, to purify BFLP1698 ligands, and in screening assays to identify molecules that inhibit the interaction of BFLP1698 with a BFLP1698 ligand.

 A BFLP1698 chimeric or fusion protein of the invention can be produced by standard
25 recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, *e.g.*, by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another
30 embodiment, the fusion gene can be synthesized by conventional techniques including automated

DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers that give rise to complementary overhangs between two consecutive gene fragments that can subsequently be annealed and reamplified to generate a chimeric gene sequence. Moreover, many expression vectors are commercially available that already encode a fusion moiety (*e.g.*, a GST polypeptide). A BFLP1698-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the BFLP1698 protein.

If desired, libraries of fragments of the BFLP1698 protein coding sequence can be used to generate a variegated population of BFLP1698 fragments for screening and subsequent selection of variants of a BFLP1698 protein.

BFLP1698 Antibodies

Also included in the invention are antibodies to BFLP1698 proteins, or fragments of BFLP1698 proteins. The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin (Ig) molecules, *i.e.*, molecules that contain an antigen binding site that specifically binds (immunoreacts with) an antigen. Such antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, F_{ab} , F_{ab}' and $F_{(ab)2}$ fragments, and an F_{ab} expression library. In general, an antibody molecule obtained from humans relates to any of the classes IgG, IgM, IgA, IgE and IgD, which differ from one another by the nature of the heavy chain present in the molecule. Certain classes have subclasses as well, such as IgG₁, IgG₂, and others. Furthermore, in humans, the light chain may be a kappa chain or a lambda chain. Reference herein to antibodies includes a reference to all such classes, subclasses and types of human antibody species.


An isolated BFLP1698-related protein of the invention may be intended to serve as an antigen, or a portion or fragment thereof, and additionally can be used as an immunogen to generate antibodies that immunospecifically bind the antigen, using standard techniques for polyclonal and monoclonal antibody preparation. The full-length protein can be used or, alternatively, the invention provides antigenic peptide fragments of the antigen for use as immunogens. An antigenic peptide fragment comprises at least 6 amino acid residues of the amino acid sequence of the full length protein, such as an amino acid sequence shown in SEQ ID

NO:2, and encompasses an epitope thereof such that an antibody raised against the peptide forms a specific immune complex with the full length protein or with any fragment that contains the epitope. Preferably, the antigenic peptide comprises at least 10 amino acid residues, or at least 15 amino acid residues, or at least 20 amino acid residues, or at least 30 amino acid residues.

- 5 Preferred epitopes encompassed by the antigenic peptide are regions of the protein that are located on its surface; commonly these are hydrophilic regions.

In certain embodiments of the invention, at least one epitope encompassed by the antigenic peptide is a region of BFLP1698-related protein that is located on the surface of the protein, *e.g.*, a hydrophilic region. A hydrophobicity analysis of the human BFLP1698-related
 10 protein sequence will indicate which regions of a BFLP1698-related protein are particularly hydrophilic and, therefore, are likely to encode surface residues useful for targeting antibody production. As a means for targeting antibody production, hydropathy plots showing regions of hydrophilicity and hydrophobicity may be generated by any method well known in the art, including, for example, the Kyte Doolittle or the Hopp Woods methods, either with or without
 15 Fourier transformation. A Kyte & Doolittle plot was generated for the BFLP1698 protein, and is shown in Table 5 below.

Table 6. Kyte & Doolittle Plot for BFLP1698



The novel nucleic acid encoding the BFLP1698 protein of the invention, or fragments
 20 thereof, may further be useful in diagnostic applications, wherein the presence or amount of the

nucleic acid or the protein are to be assessed. These materials are further useful in the generation of antibodies that bind immunospecifically to the novel substances of the invention for use in therapeutic or diagnostic methods. The disclosed BFLP1698 protein has multiple hydrophilic regions, each of which can be used as an immunogen. In one embodiment, a contemplated
5 BFLP1698 epitope is from about amino acids 1 to 60. In another embodiment, a BFLP1698 epitope is from about amino acids 70 to 80. In additional embodiments, BFLP1698 epitopes are from about amino acids 85 to 170, from about amino acids 180 to 190, from about amino acids 210 to 220, from about amino acids 230 to 260, from about amino acids 290 to 310, from about amino acids 350 to 360, from about amino acids 370 to 380, from about amino acids 400 to 430,
10 from about amino acids 450 to 480, from about amino acids 520 to 540, from about amino acids 600 to 620, from about amino acids 630 to 640, from about amino acids 680 to 690, from about amino acids 730 to 740, from about amino acids 850 to 860, from about amino acids 870 to 890, from about amino acids 970 to 1010, from about amino acids 1030 to 1050, from about amino acids 1080 to 1130, from about amino acids 1150 to 1160, and from about amino acids 1180 to
15 1190.

Antibodies that are specific for one or more domains within an antigenic protein, or derivatives, fragments, analogs or homologs thereof, are also provided herein.

A protein of the invention, or a derivative, fragment, analog, homolog or ortholog thereof, may be utilized as an immunogen in the generation of antibodies that immunospecifically bind
20 these protein components.

Various procedures known within the art may be used for the production of polyclonal or monoclonal antibodies directed against a protein of the invention, or against derivatives, fragments, analogs homologs or orthologs thereof. The term "monoclonal antibody" (MAb) or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules
25 that contain only one molecular species of antibody molecule consisting of a unique light chain gene product and a unique heavy chain gene product.

The antibodies directed against the protein antigens of the invention can further comprise humanized antibodies or human antibodies. The humanized forms of antibodies include chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab',

F(ab')₂ or other antigen-binding subsequences of antibodies) that are principally comprised of the sequence of a human immunoglobulin, and contain minimal sequence derived from a non-human immunoglobulin.

5 The antibodies can also be human antibodies, e.g., antibody molecules in which essentially the entire sequences of both the light chain and the heavy chain, including the CDRs, arise from human genes. Human monoclonal antibodies can be prepared by the trioma technique; the human B-cell hybridoma technique and the EBV hybridoma technique.

10 Human antibodies can also be produced using phage display libraries, or by introducing human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Human antibodies may additionally be produced using transgenic nonhuman animals that are modified so as to produce fully human antibodies rather than the animal's endogenous antibodies in response to challenge by an antigen.

15 The invention also provides single-chain antibodies specific to an antigenic protein of the invention. In addition, methods can be adapted for the construction of F_{ab} expression libraries to allow rapid and effective identification of monoclonal F_{ab} fragments with the desired specificity for a protein or derivatives, fragments, analogs or homologs thereof. Antibody fragments that contain the idiotypes to a protein antigen may be produced by techniques known in the art including, but not limited to: (i) an F_{(ab')₂} fragment produced by pepsin digestion of an antibody
20 molecule; (ii) an F_{ab} fragment generated by reducing the disulfide bridges of an F_{(ab')₂} fragment; (iii) an F_{ab} fragment generated by the treatment of the antibody molecule with papain and a reducing agent and (iv) F_v fragments.

Also provided by the invention are bispecific antibodies. Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at
25 least two different antigens. One of the binding specificities is for an antigenic protein of the invention. The second binding target is any other antigen, and advantageously is a cell-surface protein or receptor or receptor subunit.

If desired, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The

fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions.

5 Bispecific antibodies can be provided as full length antibodies or antibody fragments (e.g. F(ab')₂ bispecific antibodies).

Also within the invention are antibodies with more than two valencies (such as trispecific antibodies).

Exemplary bispecific antibodies bind to two different epitopes, at least one of which
10 originates in the protein antigen of the invention.

The invention also includes heteroconjugate antibodies, which include two covalently joined antibodies.

The antibody of the invention can be modified to alter (e.g., enhance or diminish) its function. For example, cysteine residue(s) can be introduced into the Fc region, thereby allowing
15 interchain disulfide bond formation in this region. The invention also includes immunoconjugates that include an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate).

Enzymatically active toxins and fragments thereof that can be used include diphtheria A
20 chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, *sapaonaria officinalis* inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are
25 available for the production of radioconjugated antibodies. Examples include ²¹²Bi, ¹³¹I, ¹³¹In, ⁹⁰Y, and ¹⁸⁶Re.

The antibody can be conjugated to a "receptor" (such streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by

removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) that is in turn conjugated to a cytotoxic agent.

BFLP1698 Recombinant Expression Vectors and Host Cells

5 Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding a BFLP1698 protein, or derivatives, fragments, analogs or homologs thereof. As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be
10 ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are
15 replicated along with the host genomic sequence into which they have integrated. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively-linked. Such vectors are referred to herein as "expression vectors". "Plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors
20 (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

 Within a recombinant expression vector, "operably-linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner that allows for expression of the nucleotide sequence (e.g., in an *in vitro* transcription/translation system or in a
25 host cell when the vector is introduced into the host cell). The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (e.g., polyadenylation signals). Regulatory sequences include those that direct constitutive expression of a nucleotide sequence in many types of host cell and those that direct expression of the nucleotide sequence only in certain host cells (e.g., tissue-specific regulatory sequences). The

expression vectors of the invention can be introduced into host cells to thereby produce proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein (e.g., BFLP1698 proteins, mutant forms of BFLP1698 proteins, fusion proteins, etc.).

5 The recombinant expression vectors of the invention can be designed for expression of BFLP1698 proteins in prokaryotic or eukaryotic cells. For example, BFLP1698 proteins can be expressed in bacterial cells such as *Escherichia coli*, insect cells (using baculovirus expression vectors) yeast cells or mammalian cells. Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

10 In yet another embodiment, a nucleic acid of the invention is expressed in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 and pMT2PC. When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, adenovirus 2, cytomegalovirus, and simian virus 40. For other suitable
15 expression systems for both prokaryotic and eukaryotic cells.

In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (e.g., tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the
20 albumin promoter (liver-specific), lymphoid-specific promoters, in particular promoters of T cell receptors and immunoglobulins, neuron-specific promoters (e.g., the neurofilament promoter), pancreas-specific promoters, and mammary gland-specific promoters (e.g., milk whey promoter). Developmentally-regulated promoters are also encompassed, e.g., the murine hox promoters and the α -fetoprotein promoter.

25 The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an antisense orientation. That is, the DNA molecule is operatively-linked to a regulatory sequence in a manner that allows for expression (by transcription of the DNA molecule) of an RNA molecule that is antisense to BFLP1698 mRNA. Regulatory sequences operatively linked to a nucleic acid cloned in the

antisense orientation can be chosen that direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be chosen that direct constitutive, tissue specific or cell type specific expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense nucleic acids are produced under the control of a high efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced.

Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but also to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

A host cell can be any prokaryotic or eukaryotic cell. For example, BFLP1698 protein can be expressed in bacterial cells such as *E. coli*, insect cells, yeast or mammalian cells (such as human, Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

A gene that encodes a selectable marker (*e.g.*, resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Various selectable markers include those that confer resistance to drugs, such as G418, hygromycin and methotrexate. A nucleic acid encoding a selectable marker can be introduced into a host cell on the same vector as that encoding BFLP1698 or can be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (*e.g.*, cells that have incorporated the selectable marker gene will survive, while the other cells die).

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce (*i.e.*, express) BFLP1698 protein. Accordingly, the invention further provides methods for producing BFLP1698 protein using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention (into which a

recombinant expression vector encoding BFLP1698 protein has been introduced) in a suitable medium such that BFLP1698 protein is produced. In another embodiment, the method further comprises isolating BFLP1698 protein from the medium or the host cell.

Transgenic BFLP1698 Animals

5 The host cells of the invention can also be used to produce non-human transgenic animals. For example, in one embodiment, a host cell of the invention is a fertilized oocyte or an embryonic stem cell into which BFLP1698 protein-coding sequences have been introduced. Such host cells can then be used to create non-human transgenic animals in which exogenous BFLP1698 sequences have been introduced into their genome or homologous recombinant
10 animals in which endogenous BFLP1698 sequences have been altered. Such animals are useful for studying the function and/or activity of BFLP1698 protein and for identifying and/or evaluating modulators of BFLP1698 protein activity. As used herein, a "transgenic animal" is a non-human animal, preferably a mammal, more preferably a rodent such as a rat or mouse, in which one or more of the cells of the animal includes a transgene. Other examples of transgenic
15 animals include non-human primates, sheep, dogs, cows, goats, chickens, amphibians, etc. A transgene is exogenous DNA that is integrated into the genome of a cell from which a transgenic animal develops and that remains in the genome of the mature animal, thereby directing the expression of an encoded gene product in one or more cell types or tissues of the transgenic animal. As used herein, a "homologous recombinant animal" is a non-human animal, preferably
20 a mammal, more preferably a mouse, in which an endogenous BFLP1698 gene has been altered by homologous recombination between the endogenous gene and an exogenous DNA molecule introduced into a cell of the animal, *e.g.*, an embryonic cell of the animal, prior to development of the animal.

 A transgenic animal of the invention can be created by introducing BFLP1698-encoding
25 nucleic acid into the male pronuclei of a fertilized oocyte (*e.g.*, by microinjection, retroviral infection) and allowing the oocyte to develop in a pseudopregnant female foster animal. Sequences including SEQ ID NO:1 can be introduced as a transgene into the genome of a non-human animal. Alternatively, a non-human homologue of the human BFLP1698 gene, such as a mouse BFLP1698 gene, can be isolated based on hybridization to the human BFLP1698

cDNA (described further *supra*) and used as a transgene. Intronic sequences and polyadenylation signals can also be included in the transgene to increase the efficiency of expression of the transgene. A tissue-specific regulatory sequence(s) can be operably-linked to the BFLP1698 transgene to direct expression of BFLP1698 protein to particular cells. Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art. Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the BFLP1698 transgene in its genome and/or expression of BFLP1698 mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene-encoding BFLP1698 protein can further be bred to other transgenic animals carrying other transgenes.

To create a homologous recombinant animal, a vector is prepared which contains at least a portion of a BFLP1698 gene into which a deletion, addition or substitution has been introduced to thereby alter, *e.g.*, functionally disrupt, the BFLP1698 gene. The BFLP1698 gene can be a human gene (*e.g.*, the DNA of SEQ ID NO:1), but more preferably, is a non-human homologue of a human BFLP1698 gene. For example, a mouse homologue of human BFLP1698 gene of SEQ ID NO:1 can be used to construct a homologous recombination vector suitable for altering an endogenous BFLP1698 gene in the mouse genome. In one embodiment, the vector is designed such that, upon homologous recombination, the endogenous BFLP1698 gene is functionally disrupted (*i.e.*, no longer encodes a functional protein; also referred to as a "knock out" vector).

Alternatively, the vector can be designed such that, upon homologous recombination, the endogenous BFLP1698 gene is mutated or otherwise altered but still encodes functional protein (*e.g.*, the upstream regulatory region can be altered to thereby alter the expression of the endogenous BFLP1698 protein). In the homologous recombination vector, the altered portion of the BFLP1698 gene is flanked at its 5'- and 3'-termini by additional nucleic acid of the BFLP1698 gene to allow for homologous recombination to occur between the exogenous BFLP1698 gene carried by the vector and an endogenous BFLP1698 gene in an embryonic stem cell. The additional flanking BFLP1698 nucleic acid is of sufficient length for successful

homologous recombination with the endogenous gene. Typically, several kilobases of flanking DNA (both at the 5'- and 3'-termini) are included in the vector. The vector is then introduced into an embryonic stem cell line (*e.g.*, by electroporation) and cells in which the introduced BFLP1698 gene has homologously-recombined with the endogenous BFLP1698 gene are
5 selected.

The selected cells are then injected into a blastocyst of an animal (*e.g.*, a mouse) to form aggregation chimeras. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term. Progeny harboring the homologously-recombined DNA in their germ cells can be used to breed animals in which all cells of the animal
10 contain the homologously-recombined DNA by germline transmission of the transgene.

In another embodiment, transgenic non-humans animals can be produced that contain selected systems that allow for regulated expression of the transgene. One example of such a system is the cre/loxP recombinase system of bacteriophage P1. For a description of the cre/loxP recombinase system. Another example of a recombinase system is the FLP recombinase system
15 of *Saccharomyces cerevisiae*. If a cre/loxP recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the Cre recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, *e.g.*, by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase.

20 Clones of the non-human transgenic animals described herein can also be produced according to the methods described in the art. In brief, a cell (*e.g.*, a somatic cell) from the transgenic animal can be isolated and induced to exit the growth cycle and enter G₀ phase. The quiescent cell can then be fused, *e.g.*, through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed
25 oocyte is then cultured such that it develops to morula or blastocyte and then transferred to pseudopregnant female foster animal. The offspring borne of this female foster animal will be a clone of the animal from which the cell (*e.g.*, the somatic cell) is isolated.

Methods of Detecting BFLP1698 Nucleic Acids and Diagnosing Lupus Nephritis

Reagents that detect BFLP1698 nucleic acids and/or polypeptides can be used to detect levels of BFLP1698 RNA and/or proteins sequences in a sample. Because elevated levels of BFLP1698 RNA are found in animals with lupus nephritis, detection of enhanced levels of BFLP1698 RNA and/or BFLP1698 polypeptides indicates the presence or predisposition to lupus in the subject. In addition, lowered levels of BFLP1698 RNA in treated lupus subjects as compared to untreated lupus indicates a return to a non-lupus state. Thus, the efficacy of lupus treatment can be monitored by comparing BFLP1698 RNA or protein levels in a sample from a treated population to samples in a diseased but untreated sample, (or a sample from an individual that has been treated for a shorter period of time).

Levels of BFLP1698 RNA can be assessed by comparing levels in a test cell population, from a subject whose lupus status is unknown, to levels in a reference cell population whose lupus status is known. Thus, the test cell population will typically include at least one cell that is capable of expressing a BFLP1698 gene. By "capable of expressing" is meant that the gene is present in an intact form in the cell and can be expressed. Expression of the BFLP1698 sequence is then detected, if present, and, preferably, measured using methods known in the art. For example, the BFLP1698 sequences disclosed herein can be used to construct probes for detecting BFLP1698 RNA sequences in, *e.g.*, northern blot hybridization analyses or methods which specifically, and, preferably, quantitatively amplify BFLP1698 specific nucleic acid sequences. Alternatively, the sequences can be used to construct primers for specifically amplifying the BFLP1698 sequences in, *e.g.*, amplification-based detection methods such as reverse-transcription based polymerase chain reaction.

BFLP1698 expression can be also measured at the protein level, *i.e.*, by measuring the levels of BFLP1698 polypeptides. Such methods are well known in the art and include, *e.g.*, immunoassays based on antibodies to proteins encoded by the genes.

Expression of sequences in test and control populations of cells can be compared using any art-recognized method for comparing expression of nucleic acid sequences. Whether or not comparison of the gene expression profile in the test cell population to the reference cell population reveals the presence, or degree, of the measured parameter depends on the composition of the reference cell population. For example, if the reference cell population is composed of cells from a lupus free subject, a similar gene expression level in the test cell

population and a reference cell population indicates the test cell population is from a lupus free subject.. Conversely, if the reference cell population is made up of cells from a diseased subject, a similar gene expression profile between the test cell population and the reference cell population indicates the test cell population is from a subject with lupus.

5 In various embodiments, a BFLP1698 sequence in a test cell population is considered comparable in expression level to the expression level of the ADIPO sequence in the reference cell population if its expression level varies within a factor of 2.0, 1.5, or 1.0 fold to the level of the BFLP1698 transcript in the reference cell population. In various embodiments, a BFLP1698 sequence in a test cell population can be considered altered in levels of expression if its
10 expression level varies from the reference cell population by more than 1.0, 1.5, 2.0 or more fold from the expression level of the corresponding BFLP1698 sequence in the reference cell population.

 If desired, comparison of differentially expressed sequences between a test cell population and a reference cell population can be done with respect to a control nucleic acid
15 whose expression is independent of the parameter or condition being measured. Expression levels of the control nucleic acid in the test and reference nucleic acid can be used to normalize signal levels in the compared populations. Suitable control nucleic acids can readily be determined by one of ordinary skill in the art.

 In some embodiments, the test cell population is compared to multiple reference cell
20 populations. Each of the multiple reference populations may differ in the known parameter. Thus, a test cell population may be compared to a first reference cell population from a subject known to have lupus, as well as a second reference population known to not have lupus.

 The test cell population that is exposed can be any number of cells, *i.e.*, one or more cells, and can be provided *in vitro*, *in vivo*, or *ex vivo*.

25 Preferably, cells in the reference cell population are derived from a tissue type as similar as possible to test cell, *e.g.*, renal tissue. In some embodiments, the control cell is derived from the same subject as the test cell. In other embodiments, the reference cell population is derived

from a plurality of cells from multiple subjects. For example, the reference cell population can be a database of expression patterns from previously tested cells.

The subject is preferably a mammal. The mammal can be, *e.g.*, a human, non-human primate, mouse, rat, dog, cat, horse, or cow.

5 **Pharmaceutical Compositions**

The BFLP1698 nucleic acid molecules, BFLP1698 proteins, and anti-BFLP1698 antibodies (also referred to herein as "active compounds") of the invention, and derivatives, fragments, analogs and homologs thereof, can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule,
10 protein, or antibody and a pharmaceutically acceptable carrier. As used herein, "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Suitable carriers are described in the most recent edition of Remington's Pharmaceutical Sciences, a standard reference text in the
15 field, which is incorporated herein by reference. Preferred examples of such carriers or diluents include, but are not limited to, water, saline, Ringer's solutions, dextrose solution, and 5% human serum albumin. Liposomes and non-aqueous vehicles such as fixed oils may also be used. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active
20 compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (*i.e.*, topical),
25 transmucosal, and rectal administration.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion.

Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated
5 by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

Sustained-release preparations can be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the
10 antibody, which matrices are in the form of shaped articles, e.g., films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides, copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOTTM (injectable microspheres composed of lactic
15 acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

20 **Screening and Detection Methods**

The isolated nucleic acid molecules of the invention can be used to express BFLP1698 protein (*e.g.*, via a recombinant expression vector in a host cell in gene therapy applications), to detect BFLP1698 mRNA (*e.g.*, in a biological sample) or a genetic lesion in a BFLP1698 gene, and to modulate BFLP1698 activity, as described further, below. In addition, the BFLP1698
25 proteins can be used to screen drugs or compounds that modulate the BFLP1698 protein activity or expression as well as to treat disorders characterized by insufficient or excessive production of BFLP1698 protein or production of BFLP1698 protein forms that have decreased or aberrant activity compared to BFLP1698 wild-type protein. In addition, the anti-BFLP1698 antibodies of

the invention can be used to detect and isolate BFLP1698 proteins and modulate BFLP1698 activity. For example, BFLP1698 activity includes T-cell or NK cell growth and differentiation, antibody production, and tumor growth.

The invention further pertains to novel agents identified by the screening assays described
5 herein and uses thereof for treatments as described, *supra*.

Screening Assays

The invention provides a method (also referred to herein as a "screening assay") for identifying modulators, *i.e.*, candidate or test compounds or agents (*e.g.*, peptides, peptidomimetics, small molecules or other drugs) that bind to BFLP1698 proteins or have a
10 stimulatory or inhibitory effect on, *e.g.*, BFLP1698 protein expression or BFLP1698 protein activity. The invention also includes compounds identified in the screening assays described herein.

In one embodiment, the screening assays are used to identify therapeutic agents for treating autoimmune diseases. The autoimmune disease can be, *e.g.*, lupus, including lupus
15 nephritis.

In one embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of the membrane-bound form of a BFLP1698 protein or polypeptide or biologically-active portion thereof. The test compounds of the invention can be obtained using any of the numerous approaches in combinatorial library
20 methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the "one-bead one-compound" library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule
25 libraries of compounds.

A "small molecule" as used herein, is meant to refer to a composition that has a molecular weight of less than about 5 kD and most preferably less than about 4 kD. Small molecules can be, *e.g.*, rapamycin, nucleic acids, peptides, polypeptides, peptidomimetics, carbohydrates, lipids

or other organic or inorganic molecules. Libraries of chemical and/or biological mixtures, such as fungal, bacterial, or algal extracts, are known in the art and can be screened with any of the assays of the invention. The libraries of compounds may be presented in solution, or on beads, on chips, bacteria, spores, plasmids or on phage

- 5 In one embodiment, an assay is a cell-based assay in which a cell which expresses a membrane-bound form of BFLP1698 protein, or a biologically-active portion thereof, on the cell surface is contacted with a test compound and the ability of the test compound to bind to a BFLP1698 protein determined. The cell, for example, can be of mammalian origin or a yeast cell. Determining the ability of the test compound to bind to the BFLP1698 protein can be
- 10 accomplished, for example, by coupling the test compound with a radioisotope or enzymatic label such that binding of the test compound to the BFLP1698 protein or biologically-active portion thereof can be determined by detecting the labeled compound in a complex. For example, test compounds can be labeled with ^{125}I , ^{35}S , ^{14}C , or ^3H , either directly or indirectly, and the radioisotope detected by direct counting of radioemission or by scintillation counting.
- 15 Alternatively, test compounds can be enzymatically-labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product. In one embodiment, the assay comprises contacting a cell which expresses a membrane-bound form of BFLP1698 protein, or a biologically-active portion thereof, on the cell surface with a known compound
- 20 which binds BFLP1698 to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with a BFLP1698 protein, wherein determining the ability of the test compound to interact with a BFLP1698 protein comprises determining the ability of the test compound to preferentially bind to BFLP1698 protein or a biologically-active portion thereof as compared to the known compound.
- 25 In another embodiment, an assay is a cell-based assay comprising contacting a cell expressing a membrane-bound form of BFLP1698 protein, or a biologically-active portion thereof, on the cell surface with a test compound and determining the ability of the test compound to modulate (*e.g.*, stimulate or inhibit) the activity of the BFLP1698 protein or biologically-active portion thereof. Determining the ability of the test compound to modulate the

activity of BFLP1698 or a biologically-active portion thereof can be accomplished, for example, by determining the ability of the BFLP1698 protein to bind to or interact with a BFLP1698 target molecule. As used herein, a "target molecule" is a molecule with which a BFLP1698 protein binds or interacts in nature, for example, a molecule on the surface of a cell which expresses a BFLP1698 interacting protein, a molecule on the surface of a second cell, a molecule in the extracellular milieu, a molecule associated with the internal surface of a cell membrane or a cytoplasmic molecule. A BFLP1698 target molecule can be a non-BFLP1698 molecule or a BFLP1698 protein or polypeptide of the invention. In one embodiment, a BFLP1698 target molecule is a component of a signal transduction pathway that facilitates transduction of an extracellular signal (*e.g.* a signal generated by binding of a compound to a membrane-bound BFLP1698 molecule) through the cell membrane and into the cell. The target, for example, can be a second intercellular protein that has catalytic activity or a protein that facilitates the association of downstream signaling molecules with BFLP1698.

Determining the ability of the BFLP1698 protein to bind to or interact with a BFLP1698 target molecule can be accomplished by one of the methods described above for determining direct binding. In one embodiment, determining the ability of the BFLP1698 protein to bind to or interact with a BFLP1698 target molecule can be accomplished by determining the activity of the target molecule. For example, the activity of the target molecule can be determined by detecting induction of a cellular second messenger of the target (*i.e.* intracellular Ca^{2+} , diacylglycerol, IP_3 , etc.), detecting catalytic/enzymatic activity of the target an appropriate substrate, detecting the induction of a reporter gene (comprising a BFLP1698-responsive regulatory element operatively linked to a nucleic acid encoding a detectable marker, *e.g.*, luciferase), or detecting a cellular response, for example, cell survival, cellular differentiation, or cell proliferation.

In yet another embodiment, an assay of the invention is a cell-free assay comprising contacting a BFLP1698 protein or biologically-active portion thereof with a test compound and determining the ability of the test compound to bind to the BFLP1698 protein or biologically-active portion thereof. Binding of the test compound to the BFLP1698 protein can be determined either directly or indirectly as described above. In one such embodiment, the assay comprises contacting the BFLP1698 protein or biologically-active portion thereof with a known compound

which binds BFLP1698 to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with a BFLP1698 protein, wherein determining the ability of the test compound to interact with a BFLP1698 protein comprises determining the ability of the test compound to preferentially bind to
5 BFLP1698 or a biologically-active portion thereof as compared to the known compound.

In still another embodiment, an assay is a cell-free assay comprising contacting BFLP1698 protein or a biologically-active portion thereof with a test compound and determining the ability of the test compound to modulate (*e.g.* stimulate or inhibit) the activity of the BFLP1698 protein or a biologically-active portion thereof. Determining the ability of the test
10 compound to modulate the activity of BFLP1698 can be accomplished, for example, by determining the ability of the BFLP1698 protein to bind to a BFLP1698 target molecule by one of the methods described above for determining direct binding. In an alternative embodiment, determining the ability of the test compound to modulate the activity of BFLP1698 protein can be accomplished by determining the ability of the BFLP1698 protein further modulate a
15 BFLP1698 target molecule. For example, the catalytic/enzymatic activity of the target molecule on an appropriate substrate can be determined as described above.

In yet another embodiment, the cell-free assay comprises contacting the BFLP1698 protein or a biologically-active portion thereof with a known compound which binds BFLP1698 protein to form an assay mixture, contacting the assay mixture with a test compound, and
20 determining the ability of the test compound to interact with a BFLP1698 protein, wherein determining the ability of the test compound to interact with a BFLP1698 protein comprises determining the ability of the BFLP1698 protein to preferentially bind to or modulate the activity of a BFLP1698 target molecule.

The cell-free assays of the invention are amenable for use with both the soluble form or
25 the membrane-bound form of BFLP1698 protein. In the case of cell-free assays comprising the membrane-bound form of BFLP1698 protein, it may be desirable to utilize a solubilizing agent such that the membrane-bound form of BFLP1698 protein is maintained in solution. Examples of such solubilizing agents include non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide,

decanoyl-N-methylglucamide, Triton[®] X-100, Triton[®] X-114, Thesit[®], Isotridecypoly(ethylene glycol ether)_n, N-dodecyl--N,N-dimethyl-3-ammonio-1-propane sulfonate, 3-(3-cholamidopropyl) dimethylamminiol-1-propane sulfonate (CHAPS), or 3-(3-cholamidopropyl)dimethylamminiol-2-hydroxy-1-propane sulfonate (CHAPSO).

5 In more than one embodiment of the above assay methods of the invention, it may be desirable to immobilize either BFLP1698 protein or its target molecule to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to BFLP1698 protein, or interaction of BFLP1698 protein with a target molecule in the presence and absence of a candidate compound,
10 can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be provided that adds a domain that allows one or both of the proteins to be bound to a matrix. For example, GST-BFLP1698 fusion proteins or GST-target fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or
15 glutathione derivatized microtiter plates, that are then combined with the test compound or the test compound and either the non-adsorbed target protein or BFLP1698 protein, and the mixture is incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components, the matrix immobilized in the case of beads, complex determined
20 either directly or indirectly, for example, as described, *supra*. Alternatively, the complexes can be dissociated from the matrix, and the level of BFLP1698 protein binding or activity determined using standard techniques.

 Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either the BFLP1698 protein or its target molecule can be
25 immobilized utilizing conjugation of biotin and streptavidin. Biotinylated BFLP1698 protein or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well-known within the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, Ill.), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with BFLP1698 protein or target molecules, but which do not interfere with

binding of the BFLP1698 protein to its target molecule, can be derivatized to the wells of the plate, and unbound target or BFLP1698 protein trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive
5 with the BFLP1698 protein or target molecule, as well as enzyme-linked assays that rely on detecting an enzymatic activity associated with the BFLP1698 protein or target molecule.

In another embodiment, modulators of BFLP1698 protein expression are identified in a method wherein a cell is contacted with a candidate compound and the expression of BFLP1698 mRNA or protein in the cell is determined. The level of expression of BFLP1698 mRNA or
10 protein in the presence of the candidate compound is compared to the level of expression of BFLP1698 mRNA or protein in the absence of the candidate compound. The candidate compound can then be identified as a modulator of BFLP1698 mRNA or protein expression based upon this comparison. For example, when expression of BFLP1698 mRNA or protein is greater (*i.e.*, statistically significantly greater) in the presence of the candidate compound than in
15 its absence, the candidate compound is identified as a stimulator of BFLP1698 mRNA or protein expression. Alternatively, when expression of BFLP1698 mRNA or protein is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of BFLP1698 mRNA or protein expression. The level of BFLP1698 mRNA or protein expression in the cells can be determined by methods described
20 herein for detecting BFLP1698 mRNA or protein.

In yet another aspect of the invention, the BFLP1698 proteins can be used as "bait proteins" in a two-hybrid assay or three hybrid assay, to identify other proteins that bind to or interact with BFLP1698 ("BFLP1698-binding proteins" or "BFLP1698-bp") and modulate BFLP1698 activity. Such BFLP1698 -binding proteins are also likely to be involved in the
25 propagation of signals by the BFLP1698 proteins as, for example, upstream or downstream elements of the BFLP1698 pathway.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for BFLP1698 is fused to a gene

encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a
5 BFLP1698-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (*e.g.*, LacZ) that is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene that encodes
10 the protein which interacts with BFLP1698.

The invention further pertains to novel agents identified by the aforementioned screening assays and uses thereof for treatments as described herein.

The invention will be illustrated in the following non-limiting examples.

**Example 1. Expression patterns of murine BFLP1698 sequence in disease-free, lupus
15 nephritis simulated disease, and rapamycin-treated diseased mice**

The expression of murine BFLP1698 sequences were examined in mice that developed lupus nephritis-like symptoms in the NZB X NZW murine model. Expression in diseased mice was compared to expression of the sequences in non-diseased mice of varying ages, and in mice whose lupus nephritis-like symptoms diminished following treatment with rapamycin or anti-B7
20 antibodies.

Mice were obtained from Jackson Laboratories at 6 to 8 weeks of age and aged on site. Data were obtained from kidneys of mice and harvested at the indicated time point: C57BL/6 female mice at 8, and 32 weeks, F1(NZBxNZW) female mice 12, 25, and 42 weeks, mice treated with rapamycin at 42 and 55 weeks, mice treated with antibodies to B7.1 and B7.2 at 52 weeks.
25 Each group contained three mice.

Rapamycin treated mice received 5 mg/kg rapamycin subcutaneous injection 3 times per week for 8 weeks starting at 29 weeks of age. Control mice received injections of vehicle (methyl

cellulose) on the same schedule. Effectiveness of therapy was determined by normalization of proteinuria and kidney histology (data not shown). Gene expression analysis was performed on mice sacrificed at the end of the treatment course (36 weeks of age, data not shown), and at 42 weeks (6 weeks after treatment) and 55 weeks (20 weeks after treatment).

- 5 Mice treated with anti-B7 received 200µg of anti-B7.1 (1G10F9 monoclonal) and 200µg of anti-B7.2 (GL1 monoclonal) by intra-peritoneal injections 3 times per week for two weeks starting at 29 weeks of age. Gene expression analysis was performed 21 weeks after treatment.

RNA isolation and hybridization to oligonucleotide arrays

- 10 Kidneys from both male and female mice were collected and snap frozen for RNA isolation. One half each kidney was used. A longitudinal section of the left kidney and a cross section of the right kidney was used in for each individual animal.

- Snap frozen mouse kidney tissue was homogenized using homogenizer suspended in RLT buffer plus 2ME for 30 to 45 seconds. Total RNA was prepared using the Qiagen Midi Kit following the manufacturer's protocol. RNA was suspended in DEPC treated H₂O and
15 quantified by OD 280.

cDNA was synthesized from 5µg of total RNA using the Superscript Kit (BRL). cDNA was purified using phenol:chloroform:isoamyl alcohol (25:24:1) with a Phage lock gel tube following the Phage lock protocol. Supernatant was collected and cleaned up using EtOH. Sample was resuspended in DEPC treated H₂O.

- 20 In vitro T7 polymerase driven transcription reactions for synthesis and biotin labeling of antisense cRNA. Qiagen RNeasy spin column purification used used to purify the cRNA. GeneChip hybridization mixtures contained 15µg fragmented cRNA, 0.5mg/ml acetylated BSA, 0.1mg/ml herring sperm DNA, in 1X MES buffer in a total volume of 200ul as per manufactures instructions. Reaction mixtures were hybridized for 16hr at 45 °C to Affymetrix Mu11KsubA
25 and Mu11KsubB oligonucleotide arrays. The hybridization mixtures were removed and the arrays were washed and stained with Streptavidin R-phycoerthrin (Molecular Probes) using GeneChip Fluidics Station 400 and scanned with a Hewlett Packard GeneArray Scanner

following manufactures instructions. Fluorescent data was collected and converted to gene specific difference average using MicroArray Suite software.

Analysis of Oligonucleotide Array Data

An eleven member standard curve, comprised of gene fragments derived from cloned
5 bacterial and bacteriophage sequences were spiked into each hybridization mixture at
concentrations ranging from 0.5pM to 150pM representing RNA frequencies of approximately
3.3 to 1000 parts per million (ppm). The biotinylated standard curve fragments were synthesized
by T7-polymerase driven IVT reactions from plasmid-based templates. The spiked biotinylated
RNA fragments serve both as an internal standard to assess chip sensitivity and as standard curve
10 to convert measured fluorescent difference averages from individual genes into RNA frequencies
in ppm as described by Hill et al.

Gene expression frequencies from each individual mouse kidney were measured and the
expression data subjected to statistical analysis. Frequency values determined from individual
measurements for a given group of mice were averaged. Genes whose frequencies differed
15 significantly between C57Bl6 kidneys at 12 and 32 weeks of age were classified as changing as a
result of the normal aging process, and not due to a disease process.

Expression frequencies in young (disease-free), old (diseased), and effectively treated old
(disease-free) F1(NZBxNZW) mice and C57BL6 control mice of oligonucleotide sequence
identified on the Affymetrix Murine 11K chip by the qualifier aa002653_s_at are shown. This
20 sequence represents an unknown mouse gene.

The results are shown in FIG. 1. Shown is a histogram showing gene expression levels in
kidneys from the indicated mice. Expression levels of BFLP1698 do not vary significantly
between C57BL/6 kidneys at 12 weeks of age and kidney at 32 weeks of age, indicating that
expression levels do not increase with age in kidneys of non-diseased mice. In (NXBxNZW)F1
25 kidneys, the gene is expressed at normal levels prior to disease onset (12 weeks of age). As the
mice age and disease progresses, increasing expression levels are observed at 25 weeks, 36
weeks (data not shown for 36 weeks), and 42 weeks. By 55 weeks of age, the mice have died
due to kidney failure. Mice treated with rapamycin for 8 weeks with treatment starting at 29
weeks of age, remain healthy past 55 weeks of age. Kidneys of mice that have received effective

therapy (either rapamycin therapy or anti-B7 therapy) express normal levels of BFLP1698, and these normal levels persist in asymptomatic kidney 20 weeks after cessation of rapamycin therapy and 15 weeks after cessation of anti-B7 therapy. The observation that expression levels return to normal when kidney function is normal indicates that elevated levels are related to, and diagnostic of, disease progression. Blocking the function of these genes may inhibit or retard disease progression. Expression levels may also be used to assess and compare effectiveness of various therapeutic interventions.

Example 2. A variant of the human BFLP1698 polypeptide sequence shown in Table 2

A polypeptide sequence varying by one amino acid from the BFLP1698 amino acid sequence presented in Table 2 is shown below. For the sequence shown, the F at position 97 of the BFLP1698 sequence shown in Table 2 has been replaced by an L, which is shown in bold font.

MALVPGRSKEDGLWTRNSPGSSQHPESPRLPNPLWDRGKIGKVEGHQHIQDFSQKSHLPSTIVVESSEVNEESGDLHLP
HEELLLLTDGEEEDAEAFQDQSEEPGAARPHHQARQVEHSTQRGHLEIRELKKKLFKRRRVLNRRRLRHRVVGAVI
DQGLITRHHLLKKRAAQELSQEIKAFLTGVDPILGHQLSAREHARCGLLLLRSLPPARAAVLDHLRGVFEDESVRHLAA
LDETPVAGPPHRLRPPPPSHVPAGGPGLEDVVQEVQQLSEFIRANPKAWAPVISAWSIDLGMQLSSTYSGQHQRVPHA
TGALNELQLQWMGCRATRTLMDIYVQCLSALIGSCP DACVDALLDTSVQHSPPHFDWVVAHIGSSFPGTIISRVLSCGL
KDFCVHGGAGGGAGSSGSSSQTPSTDPFPGSPAIPAEKRVPKIASVVGILGHLASRHGDSIRRELLRMFHDLSLAGGS
GGRSGDPSLQATVPFLLQLAVMSPALLGTVSGELVDCPKPPAVLSQLQQLQGFPREELDNMLNLAVHLVQSASGAGA
YRLLQFLVDTAMPASVITQGLAVPDTVREACDRLIQLLLLHLQKLVHHRGGSPGEGVLGPPPPPRLVFFLDALKNHV
GELCGETRLRLERKRLWQHLLGLLSVYTRPSCGPALGHLLSRARSPEELSLATQLYAGLVVSLSGLLPLAFRSCLA
RVHAGTLQPPFTARFLRNALLLVGWEQQGGEGPAALGAHFGESASAHLSDLAPLLHP EEEVAEAAASLLAICFPFSE
ALSPSOLLGLVRAGVHRFFASRLRHGPPGVASACQLLTRLSQTSAPGLKAVLQLLVEGALHARGNTELFGGQVDGNET
LSVVSASLASASLLDTRNRHTAAVPGPGGIWSVFHAGVIGRGLKPPKPVQSRNQEQVIYNTQSLLSLLVHCCSAPGGT
ECGECWGAFILSFEAAKAVAVTLVESVCPDAAGAE LAWPPEEHARATVERDLRIGRRFREQLLFELLKLVAAPPAL
CYCSVLLRGLLAALLGHWEASRHPDTHSPWHLEASCTLVAVMAEGSLLPPALGNMHEVFSQLAPFEVRLLLLSVWGF
LREHGFLPQKFIFQSERGRFIRDFSREGGGEGGPHLAVLHSLVLRNIDRLGLFSGRFGQAPSPSTLLRQGT (SEQ ID
NO: 3)

Example 3. A variant of the human BFLP1698 polypeptide sequence shown in Table 2

A polypeptide sequence varying by one amino acid from the BFLP1698 amino acid sequence presented in Table 2 is shown below. For the sequence shown, the Q at position 192 of the BFLP1698 sequence shown in Table 2 has been replaced by a N, which is shown in bold font.

MALVPGRSKEDGLWTRNSPGSSQHPESPRLPNPLWDRGKIGKVEGHQHIQDFSQKSHLPSTIVVESSEVNEESGDLHLP
HEELLLLTDGEEEDAEAFQDQSEEPGAARPHHQARQVEHSTQRGHLEIRELKKKLFKRRRVLNRRRLRHRVVGAVI
DQGLITRHHLLKKRAAQELSQEIKAFLTGVDPILGHNLSAREHARCGLLLLRSLPPARAAVLDHLRGVFEDESVRHLAA
LDETPVAGPPHRLRPPPPSHVPAGGPGLEDVVQEVQQLSEFIRANPKAWAPVISAWSIDLGMQLSSTYSGQHQRVPHA
TGALNELQLQWMGCRATRTLMDIYVQCLSALIGSCP DACVDALLDTSVQHSPPHFDWVVAHIGSSFPGTIISRVLSCGL

5 KDFCVHGGAGGGAGSSGGSSSQTPSTDPFPGSPAIPAEKRVPKIASVVGILGHLASRHGDSIRRELLRMFHDSLAGGS
 GGRSGDPSLQATVPFLLQLAVMSPALLGTVSGELVDCLKPPAVLSQLQQHLQGFPREELDNMLNLAVHLVSQASGAGA
 YRLQLFLVDTAMPASVITTOGLAVPDTVREACDRLIQLLLLHLQKLVHHRGGSPGEGVLGPPPPRLVPFLDALKNHV
 10 GELCGETLRLEKRFRLWQHLLGLLSVYTRPSCGPEALGHLLSRARSPEELSLATQLYAGLVVSLSGLLPLAFRSCLA
 RVHAGTLQPPFTARFLRNLALLVGWEQQGGEGPAALGAHFGESASAHLSDLAPLLHPEEEVAAEAAASLLAICFPFSE
 ALSPSQLLGLVRAGVHRFFASLRLHGPVGASACQLLRLSQTSPAGLKAVLQLLVEGALHRGNTLFGGQVGDGNET
 LSVVSASLASASLLDTNRRHTAAVPGPGGIWSVFHAGVIGRGLKPPKFVQSRNQEQVIYNTQSLLSLLVHCCSAPGGT
 ECCECWGAPILSPEAAKAVAVTLVESVCPDAAGAEALWPPEEHARATVERDLRIGRRFREQLLFELLKLVAAPPAL
 15 CYCSVLLRGLLAALLGHWEASRHPDTHSPWHLEASCTLVAVMAEGSLLPPALGNMHEVFSQLAPFEVRLLLSVWGF
 LREHGPLPQKFIFQSERGRFIRDFSREGGEGGPHLAVLHSLVLRNIDRLGLFSGRFQAPSPSTLLRQGT (SEQ ID
 NO: 4)

Example 4. A variant of the human BFLP1698 polypeptide sequence shown in Table 2

15 A polypeptide sequence varying by one amino acid from the BFLP1698 amino acid
 sequence presented in Table 2 is shown below. For the sequence shown, the S at position 288 of
 the BFLP1698 sequence shown in Table 2 has been replaced by an G, which is shown in bold
 font.

20 MALVPGRSKEDGLWTRNSPGSSQHPESPRLPNPLWDRGKIGKVEGHQHIQDFSQKSHLPSIVVESSEVNEESGDLHLP
 HEELLLLTGDGEEEDAEAFQDQSEEPGAARPHHQARQVEHSTQRGHLEIRELKKKLFKRRRVNLNRERRLRHRVVGAVI
 DOGLITRHHLLKKRAAQELSQEIKAFLTGVDPILGHQLSAREHARCGLLLRLSLPPARAVALDHLRGVFDESVAHLAA
 LDETPVAGPPHRLRPPPPSHVPAGGPGLEDVVQEVQVQLSEFIRANPKAWAPVIGAWSIDLGMQLSSTYSGQHQRVPHA
 25 TGALNELQLWMGCRATRTLMDIYVQCLSALIGSCPDACVDALLDTSVQHSPhFDWVVAHIGSSFPGTIIISRVLSGGL
 KDFCVHGGAGGGAGSSGGSSSQTPSTDPFPGSPAIPAEKRVPKIASVVGILGHLASRHGDSIRRELLRMFHDSLAGGS
 GGRSGDPSLQATVPFLLQLAVMSPALLGTVSGELVDCLKPPAVLSQLQQHLQGFPREELDNMLNLAVHLVSQASGAGA
 YRLQLFLVDTAMPASVITTOGLAVPDTVREACDRLIQLLLLHLQKLVHHRGGSPGEGVLGPPPPRLVPFLDALKNHV
 30 GELCGETLRLEKRFRLWQHLLGLLSVYTRPSCGPEALGHLLSRARSPEELSLATQLYAGLVVSLSGLLPLAFRSCLA
 RVHAGTLQPPFTARFLRNLALLVGWEQQGGEGPAALGAHFGESASAHLSDLAPLLHPEEEVAAEAAASLLAICFPFSE
 ALSPSQLLGLVRAGVHRFFASLRLHGPVGASACQLLRLSQTSPAGLKAVLQLLVEGALHRGNTLFGGQVGDGNET
 LSVVSASLASASLLDTNRRHTAAVPGPGGIWSVFHAGVIGRGLKPPKFVQSRNQEQVIYNTQSLLSLLVHCCSAPGGT
 ECCECWGAPILSPEAAKAVAVTLVESVCPDAAGAEALWPPEEHARATVERDLRIGRRFREQLLFELLKLVAAPPAL
 35 CYCSVLLRGLLAALLGHWEASRHPDTHSPWHLEASCTLVAVMAEGSLLPPALGNMHEVFSQLAPFEVRLLLSVWGF
 LREHGPLPQKFIFQSERGRFIRDFSREGGEGGPHLAVLHSLVLRNIDRLGLFSGRFQAPSPSTLLRQGT (SEQ ID
 NO: 5)

Example 5. A variant of the human BFLP1698 polypeptide sequence shown in Table 2

40 A polypeptide sequence varying by one amino acid from the BFLP1698 amino acid
 sequence presented in Table 2 is shown below. For the sequence shown, the H at position 365 of
 the BFLP1698 sequence shown in Table 2 has been replaced by an R, which is shown in bold
 font.

MALVPGRSKEDGLWTRNSPGSSQHPESPRLPNPLWDRGKIGKVEGHQHIQDFSQKSHLPSIVVESSEVNEESGDLHLP
 HEELLLLTGDGEEEDAEAFQDQSEEPGAARPHHQARQVEHSTQRGHLEIRELKKKLFKRRRVNLNRERRLRHRVVGAVI
 DOGLITRHHLLKKRAAQELSQEIKAFLTGVDPILGHQLSAREHARCGLLLRLSLPPARAVALDHLRGVFDESVAHLAA
 LDETPVAGPPHRLRPPPPSHVPAGGPGLEDVVQEVQVQLSEFIRANPKAWAPVISAWSIDLGMQLSSTYSGQHQRVPHA

5 TGALNELQLWMGCRATRTLMIDIYVQCLSGALIGSCPDACVDALLDTSVQHSRFDWVVAHIGSSFPGTIIISRVLSGCL
 KDFCVHGGAGGGAGSSGSSSQTPTDPPFGSPAIPAEKRVPKIASVVGILGHLASRHGDSIRRELLRMFHDSLGGG
 GGRSGDPSLQATVPFLLQAVMSPALLGTVSSELVCLKPPAVLSQLQQLHQLQGFPREELDNMLNLAVHLVSQASGAGA
 YRLQLFLVDTAMPASVITTQGLAVPDTVREACDRLIQLLLLHLQKLVHHRGSSPGEGVLGPPPPRPLVPFLDALKNHV
 10 GELCGETLRLERKRFLWQHQLLGLLSVYTRPSCGPEALGHLSSRARSPEELSLATQLYAGLVVSLSGLLPLAFRSCLA
 RVHAGTLQPPFTARFLRNALLVGWEQQGGEGPAALGAHFGESASAHLSDLAPLLHPEEEVAAASLLAICPFPSE
 ALSPSOLLGLVRAGVHRFFASLRLHGPVGASACQLLRLSQTSPAGLKAVLQLLVEGALHRGNTLFGGQVQDGNET
 LSVVSASLASASLLDNRHTAAVPGPGGIWSVFHAGVIGRGLKPPKVVQSRNQEVYINTQSLLSLLVHCCSAPGGT
 ECGECWGAIPILSPEAAKAVAVTLVESVCPDAAGAEALWPPEEHARATVERDLRIGRRFREQLLFELLKLVAAPPAL
 CYCSVLLRGLLAALLGHWEASRHPDTHSPWHLEASCTLVAVMAEGSLLPPALGNMHEVFSQLAPFEVRLLLSVWGF
 LREHGFLPQKFIFQSERGRFIRDFSREGGEGGPHLAVLHSLVLRNIDRLGLFSGRFQAPSPSTLLRQGT (SEQ ID
 NO: 6)

Example 6. A variant of the human BFLP1698 polypeptide sequence shown in Table 2

15 A polypeptide sequence varying by one amino acid from the BFLP1698 amino acid
 sequence presented in Table 2 is shown below. For the sequence shown, the V at position 481 of
 the BFLP1698 sequence shown in Table 2 has been replaced by an M, which is shown in bold
 font.

20 MALVPGRSKEDGLWTRNSPGSSQHPESPRLPNPLWDRGKIGKVEGHQHIQDFSQKSHLPSIVVESSEVNEESGDLHLP
 HEELLLLTDGEEEDAEAFFQDQSEEPGAARPHHQARQVEHSTQRGHLEIRELKKKLFKRRRVNRRRLRHRVVGAVI
 DQGLITRHLKRAAQELSQEIKAFITGVDPILGHQLSAREHARCGLLLRLSLPPARAVALDHLRGVDFESVRAHLAA
 LDETPVAGPPHLRPPPPSHVPAGGPGLEDVVQEVQVLSSEFIRANPKAWAPVISAWSIDLMGQLSSTYSQGHQVRVPHA
 25 TGALNELQLWMGCRATRTLMIDIYVQCLSGALIGSCPDACVDALLDTSVQHSRFDWVVAHIGSSFPGTIIISRVLSGCL
 KDFCVHGGAGGGAGSSGSSSQTPTDPPFGSPAIPAEKRVPKIASVVGILGHLASRHGDSIRRELLRMFHDSLGGG
 GGRSGDPSLQATMPFLLQAVMSPALLGTVSSELVCLKPPAVLSQLQQLHQLQGFPREELDNMLNLAVHLVSQASGAGA
 YRLQLFLVDTAMPASVITTQGLAVPDTVREACDRLIQLLLLHLQKLVHHRGSSPGEGVLGPPPPRPLVPFLDALKNHV
 GELCGETLRLERKRFLWQHQLLGLLSVYTRPSCGPEALGHLSSRARSPEELSLATQLYAGLVVSLSGLLPLAFRSCLA
 RVHAGTLQPPFTARFLRNALLVGWEQQGGEGPAALGAHFGESASAHLSDLAPLLHPEEEVAAASLLAICPFPSE
 30 ALSPSOLLGLVRAGVHRFFASLRLHGPVGASACQLLRLSQTSPAGLKAVLQLLVEGALHRGNTLFGGQVQDGNET
 LSVVSASLASASLLDNRHTAAVPGPGGIWSVFHAGVIGRGLKPPKVVQSRNQEVYINTQSLLSLLVHCCSAPGGT
 ECGECWGAIPILSPEAAKAVAVTLVESVCPDAAGAEALWPPEEHARATVERDLRIGRRFREQLLFELLKLVAAPPAL
 CYCSVLLRGLLAALLGHWEASRHPDTHSPWHLEASCTLVAVMAEGSLLPPALGNMHEVFSQLAPFEVRLLLSVWGF
 LREHGFLPQKFIFQSERGRFIRDFSREGGEGGPHLAVLHSLVLRNIDRLGLFSGRFQAPSPSTLLRQGT (SEQ ID
 NO: 7)

35

Example 7. A variant of the human BFLP1698 polypeptide sequence shown in Table 2

A polypeptide sequence varying by one amino acid from the BFLP1698 amino acid
 sequence presented in Table 2 is shown below. For the sequence shown, the T at position 556 of
 the BFLP1698 sequence shown in Table 2 has been replaced by a Y, which is shown in bold font.

40 MALVPGRSKEDGLWTRNSPGSSQHPESPRLPNPLWDRGKIGKVEGHQHIQDFSQKSHLPSIVVESSEVNEESGDLHLP
 HEELLLLTDGEEEDAEAFFQDQSEEPGAARPHHQARQVEHSTQRGHLEIRELKKKLFKRRRVNRRRLRHRVVGAVI
 DQGLITRHLKRAAQELSQEIKAFITGVDPILGHQLSAREHARCGLLLRLSLPPARAVALDHLRGVDFESVRAHLAA
 LDETPVAGPPHLRPPPPSHVPAGGPGLEDVVQEVQVLSSEFIRANPKAWAPVISAWSIDLMGQLSSTYSQGHQVRVPHA
 45 TGALNELQLWMGCRATRTLMIDIYVQCLSGALIGSCPDACVDALLDTSVQHSRFDWVVAHIGSSFPGTIIISRVLSGCL
 KDFCVHGGAGGGAGSSGSSSQTPTDPPFGSPAIPAEKRVPKIASVVGILGHLASRHGDSIRRELLRMFHDSLGGG

5 GGRSGDPSLQATVPFLLQLAVMSPALLGTVSGELVDCLKPPAVLSQLQQHLQGFPREELDNMLNLAVHLVSQASGAGA
YRLQLFLVDYAMPASVITTTQGLAVPDTVREACDRLIQLLLLHLQKLVHHRGGSPGEGVLGPPPPPRVLPFLDALKNHV
GELCGETRLRLERKRFWQHQLLGLLSVYTRPSCGPEALGHLLSRARSPEELSLATQLYAGLVVSLSGLLPLAFRSCLA
RVHAGTLQPPFTARFLRNLALLVGWEQQGGEGPAALGAHFGESASAHLSDLAPLLLHPPEEVAEAAAASLLAICPPFSE
10 ALSPSQQLLGLVRAGVHRFFASRLRHGPPGVASACQLLRLSQTSPAGLKAVLQLLVEGALHRGNTLFGGQVGDGNET
LSVVSASLASASLLDTNRRHTAAVPGPGGIWSVFHAGVIGRGLKPPKFVQSRNQEQEVIYNTQSLLSLVHCCSAPGGT
ECGECWGAPILSPEAAKAVAVTLVESVCPDAAGAEALWPPEEHARATVERDLRIGRRFREQLLFFELLKLVAAPAL
CYCSVLLRGLLAALLGHWEASRHPDTHSPWHLEASCTLVAVMAEGSLLPPALGNMHEVFSQLAPFEVRLLLSVWGF
LREHGFLPQKFIQSERGRFIRDFSREGGEGGPHLAVLHSLVLRNIDRLGLFSGRFQAPSPSTLLRQGT (SEQ ID
NO: 8)

Example 8. A variant of the human BFLP1698 polypeptide sequence shown in Table 2

A polypeptide sequence varying by one amino acid from the BFLP1698 amino acid
sequence presented in Table 2 is shown below. For the sequence shown, the G at position 663 of
15 the BFLP1698 sequence shown in Table 2 has been replaced by a P, which is shown in bold font.

20 MALVPGRSKEDGLWTRNSPGSSQHPESPRLPNPLWDRGKIGKVEGHQHIQDFSQKSHLPSIVVESSEVNEESGDLHLPL
HEELLLLTGEEEDAEAFFQDQSEEPGAARPHHQARQVEHSTQQRGHLEIRELKKKLFKRRRVNLNRERRLRHRVVGAVI
DQGLITRHHLLKKRAAQELSQEIKAFLTGVDPILGHQLSAREHARCGLLLRLSLPPARAALVDHLRGVFDSEVRAHLAA
LDETPVAGPPHLRPPPPSHVPAGGPGLEDVVQEVQVQLSEFIRANPKAWAPVISAWSIDLGMQLSSTYSGQHQRVPHA
TGALNELQLQWMGCRATRLMDIYVQCLSALIGSCPDACVDALLDTSVQHSPPHFDWVVAHIGSSFPGTIISRVLSGCL
KDFCVHGGAGGGGAGSSGSSSQTPSTDPFPGSPAIPAERKVPKIASVVGILGHLSRHDGSIIRRELLRMFHDLAGGS
25 GGRSGDPSLQATVPFLLQLAVMSPALLGTVSGELVDCLKPPAVLSQLQQHLQGFPREELDNMLNLAVHLVSQASGAGA
YRLQLFLVDYAMPASVITTTQGLAVPDTVREACDRLIQLLLLHLQKLVHHRGGSPGEGVLGPPPPPRVLPFLDALKNHV
GELCGETRLRLERKRFWQHQLLGLLSVYTRPSCGPEALPHLLSRARSPEELSLATQLYAGLVVSLSGLLPLAFRSCLA
RVHAGTLQPPFTARFLRNLALLVGWEQQGGEGPAALGAHFGESASAHLSDLAPLLLHPPEEVAEAAAASLLAICPPFSE
ALSPSQQLLGLVRAGVHRFFASRLRHGPPGVASACQLLRLSQTSPAGLKAVLQLLVEGALHRGNTLFGGQVGDGNET
LSVVSASLASASLLDTNRRHTAAVPGPGGIWSVFHAGVIGRGLKPPKFVQSRNQEQEVIYNTQSLLSLVHCCSAPGGT
ECGECWGAPILSPEAAKAVAVTLVESVCPDAAGAEALWPPEEHARATVERDLRIGRRFREQLLFFELLKLVAAPAL
30 CYCSVLLRGLLAALLGHWEASRHPDTHSPWHLEASCTLVAVMAEGSLLPPALGNMHEVFSQLAPFEVRLLLSVWGF
LREHGFLPQKFIQSERGRFIRDFSREGGEGGPHLAVLHSLVLRNIDRLGLFSGRFQAPSPSTLLRQGT (SEQ ID
NO: 9)

Example 9. A variant of the human BFLP1698 polypeptide sequence shown in Table 2

A polypeptide sequence varying by one amino acid from the BFLP1698 amino acid
35 sequence presented in Table 2 is shown below. For the sequence shown, the E at position 733 of
the BFLP1698 sequence shown in Table 2 has been replaced by a D, which is shown in bold font.

40 MALVPGRSKEDGLWTRNSPGSSQHPESPRLPNPLWDRGKIGKVEGHQHIQDFSQKSHLPSIVVESSEVNEESGDLHLPL
HEELLLLTGEEEDAEAFFQDQSEEPGAARPHHQARQVEHSTQQRGHLEIRELKKKLFKRRRVNLNRERRLRHRVVGAVI
DQGLITRHHLLKKRAAQELSQEIKAFLTGVDPILGHQLSAREHARCGLLLRLSLPPARAALVDHLRGVFDSEVRAHLAA
LDETPVAGPPHLRPPPPSHVPAGGPGLEDVVQEVQVQLSEFIRANPKAWAPVISAWSIDLGMQLSSTYSGQHQRVPHA
TGALNELQLQWMGCRATRLMDIYVQCLSALIGSCPDACVDALLDTSVQHSPPHFDWVVAHIGSSFPGTIISRVLSGCL
KDFCVHGGAGGGGAGSSGSSSQTPSTDPFPGSPAIPAERKVPKIASVVGILGHLSRHDGSIIRRELLRMFHDLAGGS
45 GGRSGDPSLQATVPFLLQLAVMSPALLGTVSGELVDCLKPPAVLSQLQQHLQGFPREELDNMLNLAVHLVSQASGAGA
YRLQLFLVDYAMPASVITTTQGLAVPDTVREACDRLIQLLLLHLQKLVHHRGGSPGEGVLGPPPPPRVLPFLDALKNHV
GELCGETRLRLERKRFWQHQLLGLLSVYTRPSCGPEALGHLLSRARSPEELSLATQLYAGLVVSLSGLLPLAFRSCLA
RVHAGTLQPPFTARFLRNLALLVGWEQQGGGPAALGAHFGESASAHLSDLAPLLLHPPEEVAEAAAASLLAICPPFSE

5 ALSPSQLLGLVRAGVHRFFASLRHLHGPPGVASACQLLTRLSQTSAPGLKAVLQLLVEGALHRGNTELFGGQVDGNET
LSVVSASLASASLLDTNRRHTAAVPGPGGIWSVFHAGVIGRGLKPPKEVQSRNQVEVIYNTQSLLSLVHCCSAPGGT
ECGECWGAPILSPEAAKAVAVTLVESVCPDAAGAEALWPPEEHARATVERDLRIGRRFREQPLLFECLKLVAAAPPAL
CYCSVLLRGLLAALLGHWEASRHPDTHSPWHEASCTLVAVMAEGSLLPPALGNMHEVFSQAPFEVRLLLSVWGF
LREHGFLPQKFIFQSERGRFIRDFSRGGGEGGPHLAVLHSLVLRNIDRLGLFSGRFQAPSPSTLLRQGT (SEQ ID
NO:10)

Example 10. A variant of the human BFLP1698 polypeptide sequence shown in Table 2

10 A polypeptide sequence varying by one amino acid from the BFLP1698 amino acid
sequence presented in Table 2 is shown below. For the sequence shown, the T at position 858 of
the BFLP1698 sequence shown in Table 2 has been replaced by a A, which is shown in bold font.

15 MALVPGRSKEDGLWTRNSPGSSQHPESPRLPNPLWDRGKIGKVEGHQHIQDFSQKSHLPSIVVESSEVNEESGDLHLP
HEELLLLDGEEEDAEAFFQDQSEEPGAARPHHQARQVEHSTQRGHLEIRELKKKLFKRRRVLNREERRLRHRVVGAVI
DQGLITRHHLLKKRAAQELSQEIKAFITGVDPILGHQLSAREHARCGLLLRLSLPPARAAVLDHLRGVFDESVAHLAA
LDETPVAGPPHLRPPPPSHVPAGGPGLEDVQEVQVQLSEFIRANPKAWAPVISAWSIDLMGQLSSTYSGQHQRVPHA
TGALNELLQLWMGCRATRTLMDIYVQCLSAIGSCPDACVDALLDTSVQHSPhFDWVVAHIGSSFFGTIISRVLSGGL
KDFCVHGGAGGGAGSSGGSSSQTPTDPPGSPAIPAEKRVPKIASVVGILGHLSRHDGDSIRRELLRMFHDLSLAGGS
GGRSGDPSLQATVPFLLQLAVMSPALLGTVSGELVDCLKPPAVLSQLQOHLQGFPREELDNMLNLAVHLVSQASGAGA
YRLLQFLVDTPAMPASVITTOGLAVPDTVREACDRLIQLLLLHLQKLVHHRGGSPGEGVLGPPPPPRLPVFLDALKNHV
20 GELCGETLRLEKRFRLWQHQLLGLLSVYTRPSCGPEALGHLLSRARSPEELSLATQLYAGLVVSLSGLLPLAFRSCLA
RVHAGTLQPPFTARFLRNALLVGWEQOGGEGPAALGAHFGESASAHLSDLAPLLHPEEEVAEAAASLLAICFPFSE
ALSPSQLLGLVRAGVHRFFASLRHLHGPPGVASACQLLTRLSQTSAPGLKAVLQLLVEGALHRGNTELFGGQVDGNEA
LSVVSASLASASLLDTNRRHTAAVPGPGGIWSVFHAGVIGRGLKPPKEVQSRNQVEVIYNTQSLLSLVHCCSAPGGT
25 ECGECWGAPILSPEAAKAVAVTLVESVCPDAAGAEALWPPEEHARATVERDLRIGRRFREQPLLFECLKLVAAAPPAL
CYCSVLLRGLLAALLGHWEASRHPDTHSPWHEASCTLVAVMAEGSLLPPALGNMHEVFSQAPFEVRLLLSVWGF
LREHGFLPQKFIFQSERGRFIRDFSRGGGEGGPHLAVLHSLVLRNIDRLGLFSGRFQAPSPSTLLRQGT (SEQ ID
NO:11)

Example 11. A variant of the human BFLP1698 polypeptide sequence shown in Table 2

30 A polypeptide sequence varying by one amino acid from the BFLP1698 amino acid
sequence presented in Table 2 is shown below. For the sequence shown, the W at position 974
of the BFLP1698 sequence shown in Table 2 has been replaced by a H, which is shown in bold
font.

35 MALVPGRSKEDGLWTRNSPGSSQHPESPRLPNPLWDRGKIGKVEGHQHIQDFSQKSHLPSIVVESSEVNEESGDLHLP
HEELLLLDGEEEDAEAFFQDQSEEPGAARPHHQARQVEHSTQRGHLEIRELKKKLFKRRRVLNREERRLRHRVVGAVI
DQGLITRHHLLKKRAAQELSQEIKAFITGVDPILGHQLSAREHARCGLLLRLSLPPARAAVLDHLRGVFDESVAHLAA
LDETPVAGPPHLRPPPPSHVPAGGPGLEDVQEVQVQLSEFIRANPKAWAPVISAWSIDLMGQLSSTYSGQHQRVPHA
TGALNELLQLWMGCRATRTLMDIYVQCLSAIGSCPDACVDALLDTSVQHSPhFDWVVAHIGSSFFGTIISRVLSGGL
40 KDFCVHGGAGGGAGSSGGSSSQTPTDPPGSPAIPAEKRVPKIASVVGILGHLSRHDGDSIRRELLRMFHDLSLAGGS
GGRSGDPSLQATVPFLLQLAVMSPALLGTVSGELVDCLKPPAVLSQLQOHLQGFPREELDNMLNLAVHLVSQASGAGA
YRLLQFLVDTPAMPASVITTOGLAVPDTVREACDRLIQLLLLHLQKLVHHRGGSPGEGVLGPPPPPRLPVFLDALKNHV
GELCGETLRLEKRFRLWQHQLLGLLSVYTRPSCGPEALGHLLSRARSPEELSLATQLYAGLVVSLSGLLPLAFRSCLA
RVHAGTLQPPFTARFLRNALLVGWEQOGGEGPAALGAHFGESASAHLSDLAPLLHPEEEVAEAAASLLAICFPFSE
45 -ALSPSQLLGLVRAGVHRFFASLRHLHGPPGVASACQLLTRLSQTSAPGLKAVLQLLVEGALHRGNTELFGGQVDGNET
LSVVSASLASASLLDTNRRHTAAVPGPGGIWSVFHAGVIGRGLKPPKEVQSRNQVEVIYNTQSLLSLVHCCSAPGGT
ECGECWGAPILSPEAAKAVAVTLVESVCPDAAGAEALHPPEEHARATVERDLRIGRRFREQPLLFECLKLVAAAPPAL

CYCSVLLRGLLAALLGHWEASRHPDTHSPWHLEASCTLVAVMAEGSLLPPALGNMHEVFSQLAPFEVRLLLLSVWGF
LREHGFLPQKFIFQSERGRFIRDFSREGGEGGPHLAVLHSLVLRNIDRLGLFSGRFQAPSPSTLLRQGT (SEQ ID
NO: 12)

5 **Example 12. A variant of the human BFLP1698 polypeptide sequence shown in Table 2**

A polypeptide sequence varying by one amino acid from the BFLP1698 amino acid
sequence presented in Table 2 is shown below. For the sequence shown, the P at position 1038
of the BFLP1698 sequence shown in Table 2 has been replaced by a T, which is shown in bold
font.

10 MALVPGRSKEDGLWTRNSPGSSQHPESPRLPNPLWDRGKIGKVEGHQHIQDFSQKSHLPISIVVESSEVNEESGDLHLP
HEELLLLTGEEEEDAEAFQDQSEEPGAARPHHQARQVEHSTQRGHLEIRELKKKLFKRRRVNLRERRLRHRVVGAVI
DQGLITRHHLLKKRAAQELSQEIKAFITGVDPILGHQLSAREHARCGLLLRLSLPPARAALVDHLRGVDFESVRAHLAA
15 LDETPVAGPPHLRPPPPSHVPAGGPGLEDVVQEVQVQLSEFIRANPKAWAPVISAWSIDLMGQLSSTYSQGQHRVPHA
TGALNELLQLWMGCRATRTLMDIYVQCLSAIGSCPDACVDALLDTSVQHSPhFDWVVAHIGSSFPGTIISRVLSCGL
KDFCVHGGAGGGAGSSGSSSQTPSTDPPFGSPAIPAERKVPKIASVVGILGHLSRHDGSIIRRELLRMFHDLSAGGS
GGRSGDPSLQATVPFLLQLAVMSPALLGTVSGELVDCLKPPAVLSQLQQLHQLGFPREELDNMLNLAVHLVLSQASGAGA
YRLLQFLVDLTAMPASVITTTQGLAVPDTVREACDRLIQLLLHLQKLVHHRGGSPGEGVLGPPPPPRLPVFLDALKNHV
20 GELCGETLRERKRFLWQHQLLGLLSVYTRPSCGPEALGHLLSRARSPEELSLATQLYAGLVVLSGLLPLAFRSCIA
RVHAGTLQPPFTARFLRNALLVGWEQQGEGPAALGAHFGESASAHLSDLAPLLHPEEEVAEAAAALLAICFPFSE
ALSPSOLLGLVRAGVHRFFASLRLHGPPGVASACQLLRLSQTSPAGLKAVLQLLVEGALHRGNTLEFGGQVDGNET
LSVVSASLASASLLDNRNHTAAVPGPGGIWSVFHAGVIGRGLKPPKQVSRNQEVYINTQSLLSLVHCCSAPGGT
25 ECGECWGAPILSPEAAKAVAVTLVESVCPDAAGAEALWPPEEHARATVERDLRIGRRFREQLLFELLKLVAAPAL
CYCSVLLRGLLAALLGHWEASRHTDTHSPWHLEASCTLVAVMAEGSLLPPALGNMHEVFSQLAPFEVRLLLLSVWGF
LREHGFLPQKFIFQSERGRFIRDFSREGGEGGPHLAVLHSLVLRNIDRLGLFSGRFQAPSPSTLLRQGT (SEQ ID
NO: 13)

Example 13. A variant of the human BFLP1698 polypeptide sequence shown in Table 2

A polypeptide sequence varying by one amino acid from the BFLP1698 amino acid
sequence presented in Table 2 is shown below. For the sequence shown, the I at position 1139 of
the BFLP1698 sequence shown in Table 2 has been replaced by a L, which is shown in bold font.

30 MALVPGRSKEDGLWTRNSPGSSQHPESPRLPNPLWDRGKIGKVEGHQHIQDFSQKSHLPISIVVESSEVNEESGDLHLP
HEELLLLTGEEEEDAEAFQDQSEEPGAARPHHQARQVEHSTQRGHLEIRELKKKLFKRRRVNLRERRLRHRVVGAVI
DQGLITRHHLLKKRAAQELSQEIKAFITGVDPILGHQLSAREHARCGLLLRLSLPPARAALVDHLRGVDFESVRAHLAA
35 LDETPVAGPPHLRPPPPSHVPAGGPGLEDVVQEVQVQLSEFIRANPKAWAPVISAWSIDLMGQLSSTYSQGQHRVPHA
TGALNELLQLWMGCRATRTLMDIYVQCLSAIGSCPDACVDALLDTSVQHSPhFDWVVAHIGSSFPGTIISRVLSCGL
KDFCVHGGAGGGAGSSGSSSQTPSTDPPFGSPAIPAERKVPKIASVVGILGHLSRHDGSIIRRELLRMFHDLSAGGS
GGRSGDPSLQATVPFLLQLAVMSPALLGTVSGELVDCLKPPAVLSQLQQLHQLGFPREELDNMLNLAVHLVLSQASGAGA
YRLLQFLVDLTAMPASVITTTQGLAVPDTVREACDRLIQLLLHLQKLVHHRGGSPGEGVLGPPPPPRLPVFLDALKNHV
40 GELCGETLRERKRFLWQHQLLGLLSVYTRPSCGPEALGHLLSRARSPEELSLATQLYAGLVVLSGLLPLAFRSCIA
RVHAGTLQPPFTARFLRNALLVGWEQQGEGPAALGAHFGESASAHLSDLAPLLHPEEEVAEAAAALLAICFPFSE
ALSPSOLLGLVRAGVHRFFASLRLHGPPGVASACQLLRLSQTSPAGLKAVLQLLVEGALHRGNTLEFGGQVDGNET
LSVVSASLASASLLDNRNHTAAVPGPGGIWSVFHAGVIGRGLKPPKQVSRNQEVYINTQSLLSLVHCCSAPGGT
ECGECWGAPILSPEAAKAVAVTLVESVCPDAAGAEALWPPEEHARATVERDLRIGRRFREQLLFELLKLVAAPAL
CYCSVLLRGLLAALLGHWEASRHTDTHSPWHLEASCTLVAVMAEGSLLPPALGNMHEVFSQLAPFEVRLLLLSVWGF

2003295784 19 Aug 2005

WO 2004/047740

PCT/US2003/037317

LREHGPLPQKFIFQSERGRFIRDFSRGGGEGGPHLAVLHSLVLRNLDRLGLFSGRFQAPSPSTLLRQGT (SEQ ID NO:14)

OTHER EMBODIMENTS

While the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and
5 modifications are within the scope of the following claims.

2003295784 25 May 2009

The Claims Defining The Invention Are As Follows:

1. An isolated nucleic acid molecule encoding a polypeptide comprising an amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID NO:2.
2. The isolated nucleic acid molecule of claim 1, wherein said nucleic acid molecule encodes a polypeptide that binds rapamycin.
3. The isolated nucleic acid molecule of claim 1, wherein said nucleic acid molecule encodes a polypeptide at least 98% identical to the amino acid sequence of SEQ ID NO:2.
4. The nucleic acid molecule of claim 1, wherein said molecule hybridizes under stringent conditions to a nucleic acid sequence complementary to a nucleic acid molecule comprising nucleotides 1-3486 of SEQ ID NO:1.
5. The isolated nucleic acid molecule of claim 1, wherein said nucleic acid molecule encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2.
6. The isolated nucleic acid molecule of claim 1, wherein said nucleic acid molecule comprises nucleotides 1-3486 of SEQ ID NO:1.
7. A vector comprising the nucleic acid molecule of any one of claims 1 to 6.
8. A cell including the vector of claim 7.
9. A substantially purified polypeptide comprising an amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID NO:2.
10. The polypeptide of claim 9, wherein said polypeptide binds rapamycin.
11. The polypeptide of claim 9, wherein the amino acid sequence of said polypeptide is at least 98% identical to the amino acid sequence of SEQ ID NO:2.
12. The polypeptide of claim 9, wherein the amino acid sequence of said polypeptide is at least 99% identical to the amino acid sequence of SEQ ID NO:2.

W:\Mark Wickham\700000-799999\743901\743901_clean_speci_220509.doc

2003295784 25 May 2009

13. The polypeptide of claim 9, wherein the amino acid sequence of said polypeptide comprises the amino acid sequence of SEQ ID NO:2.

5 14. The polypeptide of claim 9, wherein the amino acid sequence of said polypeptide consists of the amino acid sequence of SEQ ID NO:2.

15. A fusion polypeptide comprising the polypeptide of claim 9 operably linked to a non-BFLP1698 polypeptide.

10 16. The fusion polypeptide of claim 15, wherein said non-BFLP1698 polypeptide comprises at least one member selected from the group consisting of an Fc region of an immunoglobulin molecules or a FLAG epitope, a HIS tag, and a MYC tag.

15 17. A fusion polypeptide comprising a rapamycin-binding domain of a polypeptide of claim 9.

18. The fusion polypeptide of claim 17, wherein the rapamycin-binding domain is operably linked to a non-BFLP1698 polypeptide, wherein said non-BFLP1698 polypeptide comprises at least one member selected from the group consisting of an Fc region of an immunoglobulin molecules or a FLAG epitope, a HIS tag, and a MYC tag.

20 19. A pharmaceutical composition comprising the fusion polypeptide of claim 17 and a pharmaceutically acceptable carrier.

25 20. An antibody that binds selectively to the polypeptide of any one of claims 9 to 14.

21. The antibody of claim 20, wherein said antibody inhibits binding of a polypeptide of claim 9 to rapamycin.

30 22. The antibody of claim 20, wherein said antibody is a polyclonal antibody.

23. The antibody of claim 20, wherein said antibody is a monoclonal antibody.

35 24. The monoclonal antibody of claim 23, wherein said monoclonal antibody is selected from the group consisting of a murine monoclonal antibody, and a humanized monoclonal antibody.

W:\Mark Wickham\700000-789999\743901\clean spec\ 220509.doc

2003295784 25 May 2009

25. A method of producing a polypeptide of claim 9, said method comprising culturing a cell including the nucleic acid molecule of any one of claims 1 to 6 under conditions allowing for expression of said polypeptide encoded by said nucleic acid molecule.

5

26. A method of detecting the presence of a nucleic acid molecule of claim 1 in a biological sample, the method comprising:

contacting the sample with a nucleic acid probe that binds specifically to said nucleic acid; and

10

identifying the bound probe, if present, thereby detecting the presence of said nucleic acid molecule in said sample.

27. A method of detecting the presence of a polypeptide of claim 9 in a sample, the method comprising:

15

contacting the sample with a compound that selectively binds to said polypeptide under conditions allowing for formation of a complex between said polypeptide and said compound; and detecting said complex, if present, thereby identifying said polypeptide in said sample.

28. The method of claim 27, wherein said compound is rapamycin.

20

29. The method of claim 27, wherein said compound is an antibody to the polypeptide of claim 9.

30. A method for determining the presence of or predisposition to lupus nephritis in a subject, the method comprising:

25

a) measuring the amount of a nucleic acid molecule of claim 1 in a sample from said subject; and

b) comparing the amount of said nucleic acid in step to the amount of the nucleic acid present in a control sample from a subject without lupus nephritis,

30

wherein an increase in the level of said nucleic acid in step (a) as compared to the level of the nucleic acid in the control sample indicates the presence of or predisposition to lupus nephritis in said subject.

31. The method of claim 30, wherein said subject is a human.

25 May 2009
2003295784

32. A method for determining the presence of or predisposition to lupus nephritis in a subject, the method comprising:

a) measuring the amount of a polypeptide of claim 9 in a sample from said subject; and

b) comparing the amount of said polypeptide to the amount of the nucleic acid present in a control sample from a subject without lupus nephritis,

wherein an increase in the level of said polypeptide as compared to the level of the polypeptide in the control sample indicates the presence of or predisposition to lupus nephritis in said subject.

33. The method of claim 32, wherein said subject is a human.

34. A method for screening for a therapeutic agent for treating an autoimmune disorder, the method comprising:

contacting a test compound with a polypeptide of claim 9; and

determining if said test compound binds to said polypeptide,

wherein binding of said test compound to said polypeptide indicates the test compound is a therapeutic agent for an autoimmune disorder.

35. The method of claim 34, wherein said immune disorder is an autoimmune disorder.

36. The method of claim 35, wherein said autoimmune disorder is lupus.

37. The method of claim 35, wherein said autoimmune disorder is lupus nephritis.

38. The method of claim 34, wherein said polypeptide is provided in a cell-free extract.

39. The method of claim 34, wherein said polypeptide is provided in a cell.

40. A method of treating lupus nephritis in a subject, the method comprising administering to said subject a therapeutically effective amount of an agent that inhibits activity of a polypeptide of claim 9 in said subject.

41. The method of claim 40, wherein said subject is a human.

W:\Mark\Wickham\700060-790000\745901\745901_clean_speci_220509.doc

2003295784 25 May 2009

42. The method of claim 40 or claim 41, wherein said agent is an antibody to the polypeptide of claim 9.

5 43. The pharmaceutical composition comprising an agent that inhibits activity of a polypeptide of claim 9 in a subject and a pharmaceutically acceptable carrier, wherein said agent is an antibody to the polypeptide of claim 9.

10 44. A polypeptide prepared by the method of claim 25.

45. An isolated nucleic acid according to claim 1 substantially as hereinbefore described with reference to the Examples and/or Figure and/or Sequence Listing.

15 46. A polypeptide according to claim 9 substantially as hereinbefore described with reference to the Examples and/or Figure and/or Sequence Listing.

47. A fusion polypeptide according to claim 17 substantially as hereinbefore described with reference to the Examples and/or Figure and/or Sequence Listing.

20 48. A method according to any one of claims 26, 27, 30, 32, 34 or 40, substantially as hereinbefore described with reference to the Examples and/or Figure and/or Sequence Listing.

GENE EXPRESSION LEVELS IN (NZB x NZW)F1 KIDNEYS OF MOUSE ORTHOLOG OF HUMAN GENE BFLP1698 AND THE EFFECT OF THERAPY ON GENE EXPRESSION LEVELS




Fig. 1

SEQUENCE LISTING

<110> Wyeth
O'Toole, Margot
Mounts, William M
Shojaee, Negin

<120> COMPOSITION AND METHOD FOR TREATING LUPUS NEPHRITIS

<130> 031896-091000

<140> US 10/718,834

<141> 2003-11-21

<150> US 60/428,065

<151> 2002-11-21

<160> 21

<170> PatentIn version 3.3

<210> 1

<211> 3652

<212> DNA

<213> Homo sapiens

<400> 1

```

atggcccttg tgccagggag aagcaaggag gatgggcttt ggactagaaa tagcccaggc      60
tcctcccagc atccagaaag tcccaggctg cccaaccctc tctgggacag aggaaaaatt      120
ggcaagggttg aaggtcacca gcacattcag gatttctctc aaaagtccca tctgccgtct      180
attgtggttg aatccagtga ggtgaatgaa gagagtgggg atctccattt gccccatgag      240
gagctgctgc tgctcactga tggtagaggaa gaggatgctg aggccttctt ccaagaccaa      300
agtgaagagc caggggcggc acgtcccat catcaggctc ggcaagtgga gcattcgacg      360
cagcgcggcc atctggagat tcgggagctg aagaagaagc tgttcaaacg ccggcggggtg      420
ttgaatcggg agcggcgtct gaggcaccgg gtggtcgggg ctgtgataga ccaagggtg      480
atcacgcggc accacctcaa gaagcgggct gctcaggagc tgtcccagga aatcaaggct      540
tttctgactg gcgtagacct cattctgggc caccaactct cagcccggga acatgctcgc      600
tgtggtcttc tcctgctccg ttctttgcca cctgctcggg ctgctgtgct tgaccacttg      660
agagggtgtc ttgatgagag tgtccgggcc cacctggctg ccctggatga aaccctgtg      720
gctggtccac ctcacctccg tccacctcca ccctctcatg tcctgctgg tggacctggt      780
ctagaggatg tggttcagga agtgcagcag gtgctgtctg agtttatccg ggccaaccca      840
aaggcctggg cacctgtgat tagtgcattg tccattgacc tcatggggca actgagcagc      900
acgtactcag gccagcacca gcgtgttccc cacgctactg gcgctcttaa tgaactgcta      960
cagctgtgga tgggtttagg ggccacgcgt acattaatgg acatctatgt gcagtgcctc     1020
tcggctctca ttggtagctg cccagatgcg tgtgtggatg ccttgctgga tacctctggt     1080

```

2003295784 19 Aug 2005

2003295784 19 Aug 2005

cagcattctc	cacactttga	ctgggttggtg	gcacatattg	gctcctcttt	tcctggcacc	1140
atcattttccc	gggttctctc	ctgtggccctt	aaggactttt	gtgtccatgg	tggggctgga	1200
ggtggagctg	gcagtagtgg	tggaagctct	tctcagaccc	cctctacaga	ccccttcctt	1260
ggatctcctg	ccattcctgc	ggagaaacgg	gtgccaaga	ttgcctcagt	tgtaggcatc	1320
ctaggtcacc	tggcctcccc	ccacggagat	agcatccgac	gggagctcct	gcgaatgttc	1380
catgatagcc	tggcaggggg	atctggaggc	cgcagtgggg	accctccctt	tcaggccacg	1440
gttccgttcc	tactgcagct	ggcagtcatg	tcaccagctt	tgctgggcac	tgtctctgga	1500
gagcttgtgg	attgcctcaa	gccccagct	gtgctgagcc	agctgcagca	acaccttcaa	1560
ggattcccc	gagaggagct	ggacaacatg	ttgaacctgg	ctgtgcacct	ggtgagccag	1620
gcctctgggg	caggtgccta	ccgcttgctg	cagttcctgg	tggacacagc	tatgcctgct	1680
tcggtcatta	ccaccaggg	cctggctgtg	ccagacaccg	tgcgtgaggc	ttgtgaccgg	1740
ctaateccagc	tgtgtctgct	gcacctgcaa	aaactggttc	atcaccgggg	agggctctct	1800
ggggaagggg	tgctaggccc	gccccacct	ccccgcttgg	tgcccttttt	agatgcgctc	1860
aaaaaccatg	ttggagagct	gtgtggagag	acgttacgat	tggaacggaa	gcgcttcctc	1920
tggcagcacc	agctcttggg	cctgctgtct	gtctataccc	ggcctagctg	tggacctgag	1980
gccttggggc	atctgctgag	ccgagcccga	agccctgaag	agttgagttt	ggccaccag	2040
ttatatgcag	ggctagtgg	cagcctctct	ggcctcctgc	ccctggcttt	ccgaagctgt	2100
ctggctcggg	tgcattgcagg	gacattacag	cctcccttca	cggcccgggt	cctgcgcaac	2160
ttggcactgc	tagtaggggtg	ggaacagcag	ggtggcgagg	gccctgcagc	cctagggggc	2220
cactttgggg	aatctgcctc	agcccatctg	tctgacctgg	ctcctctcct	gctacatcct	2280
gaggaggaag	tagctgaagc	tgctgcctct	ctcctggcca	tttgtccctt	tccttctgaa	2340
gccttatccc	cctcccagct	cctgggactg	gtaagggctg	gggtgcaccg	cttctttgcc	2400
tctctgaggc	tgcattggacc	cccaggtgtg	gcctcagcct	gtcagcttct	caccgcctg	2460
tctcagacat	ccccagctgg	gctcaaggct	gtcctgcagc	tgctggttga	aggagcctta	2520
catcgaggca	acacagaact	gtttgggtgg	caagtagatg	gggacaatga	gactctctca	2580
gttggtttcag	cttcttttgg	ttctgcctcc	ctgttgagca	ctaaccggag	gcacactgca	2640
gctgtgccag	gtcctggagg	gatttggtca	gttttccatg	ctggagtcac	cggccgtggc	2700
ttaaagccac	ccaagtttgt	ccagtcacga	aatcagcagg	aagtgatcta	taacaccag	2760
agcctcctca	gcctcctggt	tactgctgc	agtgcctcag	ggggcactga	atgtggggaa	2820
tgctgggggg	cacccatctt	gagtccagag	gcagccaaag	cagtggcagt	gaccttgggtg	2880
gagagtgtgt	gtcccgatgc	agctgggtgca	gagctggcct	ggccccccga	ggaacacgcc	2940

2003295784 19 Aug 2005

```

cggggccaccg tggagcggga tctccgcatt ggccggcgct tccgcgaaca gcccctgctc 3000
tttgagctgt taaagctggt agcagctgca cccccagccc tgtgctactg ttccgtgctg 3060
cttcggggggc tgctggccgc cctcttgggc cattgggaag cctctcgcca ccctgacacg 3120
accactccc cctggcacct ggaggcatcc tgcaccttag tggctgtcat ggctgaggga 3180
agcctcctgc ctccggccct gggtaatatg catgaagtat ttagccaact ggcacctttc 3240
gagggtgcgtc tgctgtgtct cagtgtcttg ggttttctcc gggagcatgg gcccttgctt 3300
cagaagttca tcttccaatc agagcggggg cgcttcattc gggacttctc cagggagggt 3360
ggagggtgagg gtggacccca tctggctgtg ctgcacagtg tcctccaccg caacatcgac 3420
cgcctagggtc ttttctctgg ccgtttccag gcaccttcac cgtccactct ccttcgacag 3480
gggacgtagc cttttcttgc tctggaagcc cagggagggt gagcagtgag agaggggaagg 3540
gactaacgtg ctccggaagg gtggagggtt ctcttctaag tccttggtct aaagagcgct 3600
gtcacttttt tctctccac tttttttttt ctaaataaaa tttgccaact tg 3652

```

```

<210> 2
<211> 1162
<212> PRT
<213> Homo sapiens
<400> 2

```

```

Met Ala Leu Val Pro Gly Arg Ser Lys Glu Asp Gly Leu Trp Thr Arg
1           5           10          15

Asn Ser Pro Gly Ser Ser Gln His Pro Glu Ser Pro Arg Leu Pro Asn
          20          25          30

Pro Leu Trp Asp Arg Gly Lys Ile Gly Lys Val Glu Gly His Gln His
          35          40          45

Ile Gln Asp Phe Ser Gln Lys Ser His Leu Pro Ser Ile Val Val Glu
          50          55          60

Ser Ser Glu Val Asn Glu Glu Ser Gly Asp Leu His Leu Pro His Glu
65          70          75          80

Glu Leu Leu Leu Leu Thr Asp Gly Glu Glu Glu Asp Ala Glu Ala Phe
          85          90          95

Phe Gln Asp Gln Ser Glu Glu Pro Gly Ala Ala Arg Pro His His Gln
          100         105         110

Ala Arg Gln Val Glu His Ser Thr Gln Arg Gly His Leu Glu Ile Arg
          115         120         125

Glu Leu Lys Lys Lys Leu Phe Lys Arg Arg Arg Val Leu Asn Arg Glu
          130         135         140

Arg Arg Leu Arg His Arg Val Val Gly Ala Val Ile Asp Gln Gly Leu
          145         150         155         160

```


2003295784 19 Aug 2005

Ile	Thr	Arg	His	His	Leu	Lys	Lys	Arg	Ala	Ala	Gln	Glu	Leu	Ser	Gln	
				165					170					175		
Glu	Ile	Lys	Ala	Phe	Leu	Thr	Gly	Val	Asp	Pro	Ile	Leu	Gly	His	Gln	
			180					185					190			
Leu	Ser	Ala	Arg	Glu	His	Ala	Arg	Cys	Gly	Leu	Leu	Leu	Leu	Arg	Ser	
		195					200					205				
Leu	Pro	Pro	Ala	Arg	Ala	Ala	Val	Leu	Asp	His	Leu	Arg	Gly	Val	Phe	
	210					215					220					
Asp	Glu	Ser	Val	Arg	Ala	His	Leu	Ala	Ala	Leu	Asp	Glu	Thr	Pro	Val	
225					230					235					240	
Ala	Gly	Pro	Pro	His	Leu	Arg	Pro	Pro	Pro	Pro	Ser	His	Val	Pro	Ala	
				245					250					255		
Gly	Gly	Pro	Gly	Leu	Glu	Asp	Val	Val	Gln	Glu	Val	Gln	Gln	Val	Leu	
			260					265					270			
Ser	Glu	Phe	Ile	Arg	Ala	Asn	Pro	Lys	Ala	Trp	Ala	Pro	Val	Ile	Ser	
		275					280					285				
Ala	Trp	Ser	Ile	Asp	Leu	Met	Gly	Gln	Leu	Ser	Ser	Thr	Tyr	Ser	Gly	
	290					295					300					
Gln	His	Gln	Arg	Val	Pro	His	Ala	Thr	Gly	Ala	Leu	Asn	Glu	Leu	Leu	
305					310					315					320	
Gln	Leu	Trp	Met	Gly	Cys	Arg	Ala	Thr	Arg	Thr	Leu	Met	Asp	Ile	Tyr	
			325						330					335		
Val	Gln	Cys	Leu	Ser	Ala	Leu	Ile	Gly	Ser	Cys	Pro	Asp	Ala	Cys	Val	
			340					345					350			
Asp	Ala	Leu	Leu	Asp	Thr	Ser	Val	Gln	His	Ser	Pro	His	Phe	Asp	Trp	
	355						360					365				
Val	Val	Ala	His	Ile	Gly	Ser	Ser	Phe	Pro	Gly	Thr	Ile	Ile	Ser	Arg	
	370					375					380					
Val	Leu	Ser	Cys	Gly	Leu	Lys	Asp	Phe	Cys	Val	His	Gly	Gly	Ala	Gly	
385					390					395					400	
Gly	Gly	Ala	Gly	Ser	Ser	Gly	Gly	Ser	Ser	Ser	Gln	Thr	Pro	Ser	Thr	
				405					410					415		
Asp	Pro	Phe	Pro	Gly	Ser	Pro	Ala	Ile	Pro	Ala	Glu	Lys	Arg	Val	Pro	
			420					425					430			
Lys	Ile	Ala	Ser	Val	Val	Gly	Ile	Leu	Gly	His	Leu	Ala	Ser	Arg	His	
		435					440					445				
Gly	Asp	Ser	Ile	Arg	Arg	Glu	Leu	Leu	Arg	Met	Phe	His	Asp	Ser	Leu	
	450					455					460					
Ala	Gly	Gly	Ser	Gly	Gly	Arg	Ser	Gly	Asp	Pro	Ser	Leu	Gln	Ala	Thr	
465					470					475					480	
Val	Pro	Phe	Leu	Leu	Gln	Leu	Ala	Val	Met	Ser	Pro	Ala	Leu	Leu	Gly	

2003295784 19 Aug 2005

485										490					495				
Thr	Val	Ser	Gly	Glu	Leu	Val	Asp	Cys	Leu	Lys	Pro	Pro	Ala	Val	Leu				
			500					505						510					
Ser	Gln	Leu	Gln	Gln	His	Leu	Gln	Gly	Phe	Pro	Arg	Glu	Glu	Leu	Asp				
		515					520					525							
Asn	Met	Leu	Asn	Leu	Ala	Val	His	Leu	Val	Ser	Gln	Ala	Ser	Gly	Ala				
		530				535					540								
Gly	Ala	Tyr	Arg	Leu	Leu	Gln	Phe	Leu	Val	Asp	Thr	Ala	Met	Pro	Ala				
545					550					555					560				
Ser	Val	Ile	Thr	Thr	Gln	Gly	Leu	Ala	Val	Pro	Asp	Thr	Val	Arg	Glu				
				565					570					575					
Ala	Cys	Asp	Arg	Leu	Ile	Gln	Leu	Leu	Leu	His	Leu	Gln	Lys	Leu					
			580				585					590							
Val	His	His	Arg	Gly	Gly	Ser	Pro	Gly	Glu	Gly	Val	Leu	Gly	Pro	Pro				
		595					600					605							
Pro	Pro	Pro	Arg	Leu	Val	Pro	Phe	Leu	Asp	Ala	Leu	Lys	Asn	His	Val				
		610				615					620								
Gly	Glu	Leu	Cys	Gly	Glu	Thr	Leu	Arg	Leu	Glu	Arg	Lys	Arg	Phe	Leu				
625					630					635					640				
Trp	Gln	His	Gln	Leu	Leu	Gly	Leu	Leu	Ser	Val	Tyr	Thr	Arg	Pro	Ser				
				645					650					655					
Cys	Gly	Pro	Glu	Ala	Leu	Gly	His	Leu	Leu	Ser	Arg	Ala	Arg	Ser	Pro				
			660					665					670						
Glu	Glu	Leu	Ser	Leu	Ala	Thr	Gln	Leu	Tyr	Ala	Gly	Leu	Val	Val	Ser				
		675					680					685							
Leu	Ser	Gly	Leu	Leu	Pro	Leu	Ala	Phe	Arg	Ser	Cys	Leu	Ala	Arg	Val				
		690				695					700								
His	Ala	Gly	Thr	Leu	Gln	Pro	Pro	Phe	Thr	Ala	Arg	Phe	Leu	Arg	Asn				
705					710					715					720				
Leu	Ala	Leu	Leu	Val	Gly	Trp	Glu	Gln	Gln	Gly	Gly	Glu	Gly	Pro	Ala				
				725					730					735					
Ala	Leu	Gly	Ala	His	Phe	Gly	Glu	Ser	Ala	Ser	Ala	His	Leu	Ser	Asp				
			740					745					750						
Leu	Ala	Pro	Leu	Leu	Leu	His	Pro	Glu	Glu	Glu	Val	Ala	Glu	Ala	Ala				
		755					760					765							
Ala	Ser	Leu	Leu	Ala	Ile	Cys	Pro	Phe	Pro	Ser	Glu	Ala	Leu	Ser	Pro				
		770				775					780								
Ser	Gln	Leu	Leu	Gly	Leu	Val	Arg	Ala	Gly	Val	His	Arg	Phe	Phe	Ala				
785					790						795				800				
Ser	Leu	Arg	Leu	His	Gly	Pro	Pro	Gly	Val	Ala	Ser	Ala	Cys	Gln	Leu				
				805					810					815					

2003295784 19 Aug 2005

Leu Thr Arg	Leu Ser Gln Thr Ser	Pro Ala Gly	Leu Lys	Ala Val Leu
820		825		830
Gln Leu Leu Val Glu Gly	Ala Leu His Arg	Gly Asn Thr	Glu Leu Phe	
835	840		845	
Gly Gly Gln Val Asp Gly	Asp Asn Glu Thr	Leu Ser Val	Val Ser Ala	
850	855		860	
Ser Leu Ala Ser Ala Ser	Leu Leu Asp Thr	Asn Arg Arg	His Thr Ala	
865	870	875	880	
Ala Val Pro Gly Pro Gly Gly	Ile Trp Ser Val	Phe His Ala	Gly Val	
	885	890	895	
Ile Gly Arg Gly Leu Lys Pro	Pro Lys Phe Val	Gln Ser Arg	Asn Gln	
900	905		910	
Gln Glu Val Ile Tyr Asn Thr	Gln Ser Leu Leu Ser	Leu Leu Val	His	
915	920	925		
Cys Cys Ser Ala Pro Gly Gly	Thr Glu Cys Gly	Glu Cys Trp	Gly Ala	
930	935	940		
Pro Ile Leu Ser Pro Glu Ala	Ala Lys Ala Val	Ala Val Thr	Leu Val	
945	950	955	960	
Glu Ser Val Cys Pro Asp Ala	Ala Gly Ala Glu	Leu Ala Trp	Pro Pro	
	965	970	975	
Glu Glu His Ala Arg Ala Thr	Val Glu Arg Asp	Leu Arg Ile	Gly Arg	
980	985	990		
Arg Phe Arg Glu Gln Pro Leu	Leu Phe Glu Leu Leu	Lys Leu Val	Ala	
995	1000	1005		
Ala Ala Pro Pro Ala Leu Cys	Tyr Cys Ser Val	Leu Leu Arg	Gly	
1010	1015	1020		
Leu Leu Ala Ala Leu Leu Gly	His Trp Glu Ala	Ser Arg His	Pro	
1025	1030	1035		
Asp Thr Thr His Ser Pro Trp	His Leu Glu Ala	Ser Cys Thr	Leu	
1040	1045	1050		
Val Ala Val Met Ala Glu Gly	Ser Leu Leu Pro	Pro Ala Leu	Gly	
1055	1060	1065		
Asn Met His Glu Val Phe Ser	Gln Leu Ala Pro	Phe Glu Val	Arg	
1070	1075	1080		
Leu Leu Leu Leu Ser Val Trp	Gly Phe Leu Arg	Glu His Gly	Pro	
1085	1090	1095		
Leu Pro Gln Lys Phe Ile Phe	Gln Ser Glu Arg	Gly Arg Phe	Ile	
1100	1105	1110		
Arg Asp Phe Ser Arg Glu Gly	Gly Gly Glu Gly	Gly Pro His	Leu	
1115	1120	1125		
Ala Val Leu His Ser Val Leu	His Arg Asn Ile	Asp Arg Leu	Gly	
1130	1135	1140		

2003295784 19 Aug 2005

Leu Phe Ser Gly Arg Phe Gln Ala Pro Ser Pro Ser Thr Leu Leu
 1145 1150 1155
 Arg Gln Gly Thr
 1160
 <210> 3
 <211> 1162
 <212> PRT
 <213> Artificial
 <220>
 <223> A variant of the human BFLP1698 polypeptide
 <400> 3
 Met Ala Leu Val Pro Gly Arg Ser Lys Glu Asp Gly Leu Trp Thr Arg
 1 5 10 15
 Asn Ser Pro Gly Ser Ser Gln His Pro Glu Ser Pro Arg Leu Pro Asn
 20 25 30
 Pro Leu Trp Asp Arg Gly Lys Ile Gly Lys Val Glu Gly His Gln His
 35 40 45
 Ile Gln Asp Phe Ser Gln Lys Ser His Leu Pro Ser Ile Val Val Glu
 50 55 60
 Ser Ser Glu Val Asn Glu Glu Ser Gly Asp Leu His Leu Pro His Glu
 65 70 75 80
 Glu Leu Leu Leu Leu Thr Asp Gly Glu Glu Glu Asp Ala Glu Ala Phe
 85 90 95
 Leu Gln Asp Gln Ser Glu Glu Pro Gly Ala Ala Arg Pro His His Gln
 100 105 110
 Ala Arg Gln Val Glu His Ser Thr Gln Arg Gly His Leu Glu Ile Arg
 115 120 125
 Glu Leu Lys Lys Lys Leu Phe Lys Arg Arg Arg Val Leu Asn Arg Glu
 130 135 140
 Arg Arg Leu Arg His Arg Val Val Gly Ala Val Ile Asp Gln Gly Leu
 145 150 155 160
 Ile Thr Arg His His Leu Lys Lys Arg Ala Ala Gln Glu Leu Ser Gln
 165 170 175
 Glu Ile Lys Ala Phe Leu Thr Gly Val Asp Pro Ile Leu Gly His Gln
 180 185 190
 Leu Ser Ala Arg Glu His Ala Arg Cys Gly Leu Leu Leu Leu Arg Ser
 195 200 205
 Leu Pro Pro Ala Arg Ala Ala Val Leu Asp His Leu Arg Gly Val Phe
 210 215 220
 Asp Glu Ser Val Arg Ala His Leu Ala Ala Leu Asp Glu Thr Pro Val
 225 230 235 240

2003295784 19 Aug 2005

Ala	Gly	Pro	Pro	His	Leu	Arg	Pro	Pro	Pro	Pro	Ser	His	Val	Pro	Ala	245	250	255	
Gly	Gly	Pro	Gly	Leu	Glu	Asp	Val	Val	Gln	Glu	Val	Gln	Gln	Val	Leu	260	265	270	
Ser	Glu	Phe	Ile	Arg	Ala	Asn	Pro	Lys	Ala	Trp	Ala	Pro	Val	Ile	Ser	275	280	285	
Ala	Trp	Ser	Ile	Asp	Leu	Met	Gly	Gln	Leu	Ser	Ser	Thr	Tyr	Ser	Gly	290	295	300	
Gln	His	Gln	Arg	Val	Pro	His	Ala	Thr	Gly	Ala	Leu	Asn	Glu	Leu	Leu	305	310	315	320
Gln	Leu	Trp	Met	Gly	Cys	Arg	Ala	Thr	Arg	Thr	Leu	Met	Asp	Ile	Tyr	325	330	335	
Val	Gln	Cys	Leu	Ser	Ala	Leu	Ile	Gly	Ser	Cys	Pro	Asp	Ala	Cys	Val	340	345	350	
Asp	Ala	Leu	Leu	Asp	Thr	Ser	Val	Gln	His	Ser	Pro	His	Phe	Asp	Trp	355	360	365	
Val	Val	Ala	His	Ile	Gly	Ser	Ser	Phe	Pro	Gly	Thr	Ile	Ile	Ser	Arg	370	375	380	
Val	Leu	Ser	Cys	Gly	Leu	Lys	Asp	Phe	Cys	Val	His	Gly	Gly	Ala	Gly	385	390	395	400
Gly	Gly	Ala	Gly	Ser	Ser	Gly	Gly	Ser	Ser	Ser	Gln	Thr	Pro	Ser	Thr	405	410	415	
Asp	Pro	Phe	Pro	Gly	Ser	Pro	Ala	Ile	Pro	Ala	Glu	Lys	Arg	Val	Pro	420	425	430	
Lys	Ile	Ala	Ser	Val	Val	Gly	Ile	Leu	Gly	His	Leu	Ala	Ser	Arg	His	435	440	445	
Gly	Asp	Ser	Ile	Arg	Arg	Glu	Leu	Leu	Arg	Met	Phe	His	Asp	Ser	Leu	450	455	460	
Ala	Gly	Gly	Ser	Gly	Gly	Arg	Ser	Gly	Asp	Pro	Ser	Leu	Gln	Ala	Thr	465	470	475	480
Val	Pro	Phe	Leu	Leu	Gln	Leu	Ala	Val	Met	Ser	Pro	Ala	Leu	Leu	Gly	485	490	495	
Thr	Val	Ser	Gly	Glu	Leu	Val	Asp	Cys	Leu	Lys	Pro	Pro	Ala	Val	Leu	500	505	510	
Ser	Gln	Leu	Gln	Gln	His	Leu	Gln	Gly	Phe	Pro	Arg	Glu	Glu	Leu	Asp	515	520	525	
Asn	Met	Leu	Asn	Leu	Ala	Val	His	Leu	Val	Ser	Gln	Ala	Ser	Gly	Ala	530	535	540	
Gly	Ala	Tyr	Arg	Leu	Leu	Gln	Phe	Leu	Val	Asp	Thr	Ala	Met	Pro	Ala	545	550	555	560
Ser	Val	Ile	Thr	Thr	Gln	Gly	Leu	Ala	Val	Pro	Asp	Thr	Val	Arg	Glu				

2003295784 19 Aug 2005

565					570					575				
Ala	Cys	Asp	Arg	Leu	Ile	Gln	Leu	Leu	Leu	His	Leu	Gln	Lys	Leu
580					585					590				
Val	His	His	Arg	Gly	Gly	Ser	Pro	Gly	Glu	Gly	Val	Leu	Gly	Pro
595					600					605				
Pro	Pro	Pro	Arg	Leu	Val	Pro	Phe	Leu	Asp	Ala	Leu	Lys	Asn	His
610					615					620				
Gly	Glu	Leu	Cys	Gly	Glu	Thr	Leu	Arg	Leu	Glu	Arg	Lys	Arg	Phe
625					630					635				
Trp	Gln	His	Gln	Leu	Leu	Gly	Leu	Leu	Ser	Val	Tyr	Thr	Arg	Pro
645					650					655				
Cys	Gly	Pro	Glu	Ala	Leu	Gly	His	Leu	Leu	Ser	Arg	Ala	Arg	Ser
660					665					670				
Glu	Glu	Leu	Ser	Leu	Ala	Thr	Gln	Leu	Tyr	Ala	Gly	Leu	Val	Val
675					680					685				
Leu	Ser	Gly	Leu	Leu	Pro	Leu	Ala	Phe	Arg	Ser	Cys	Leu	Ala	Arg
690					695					700				
His	Ala	Gly	Thr	Leu	Gln	Pro	Pro	Phe	Thr	Ala	Arg	Phe	Leu	Arg
705					710					715				
Leu	Ala	Leu	Leu	Val	Gly	Trp	Glu	Gln	Gln	Gly	Gly	Glu	Gly	Pro
725					730					735				
Ala	Leu	Gly	Ala	His	Phe	Gly	Glu	Ser	Ala	Ser	Ala	His	Leu	Ser
740					745					750				
Leu	Ala	Pro	Leu	Leu	Leu	His	Pro	Glu	Glu	Glu	Val	Ala	Glu	Ala
755					760					765				
Ala	Ser	Leu	Leu	Ala	Ile	Cys	Pro	Phe	Pro	Ser	Glu	Ala	Leu	Ser
770					775					780				
Ser	Gln	Leu	Leu	Gly	Leu	Val	Arg	Ala	Gly	Val	His	Arg	Phe	Phe
785					790					795				
Ser	Leu	Arg	Leu	His	Gly	Pro	Pro	Gly	Val	Ala	Ser	Ala	Cys	Gln
805					810					815				
Leu	Thr	Arg	Leu	Ser	Gln	Thr	Ser	Pro	Ala	Gly	Leu	Lys	Ala	Val
820					825					830				
Gln	Leu	Leu	Val	Glu	Gly	Ala	Leu	His	Arg	Gly	Asn	Thr	Glu	Leu
835					840					845				
Gly	Gly	Gln	Val	Asp	Gly	Asp	Asn	Glu	Thr	Leu	Ser	Val	Val	Ser
850					855					860				
Ser	Leu	Ala	Ser	Ala	Ser	Leu	Leu	Asp	Thr	Asn	Arg	Arg	His	Thr
865					870					875				
Ala	Val	Pro	Gly	Pro	Gly	Gly	Ile	Trp	Ser	Val	Phe	His	Ala	Gly
885					890					895				

2003295784 19 Aug 2005

```

Ile Gly Arg Gly Leu Lys Pro Pro Lys Phe Val Gln Ser Arg Asn Gln
    900                                905                                910

Gln Glu Val Ile Tyr Asn Thr Gln Ser Leu Leu Ser Leu Leu Val His
    915                                920                                925

Cys Cys Ser Ala Pro Gly Gly Thr Glu Cys Gly Glu Cys Trp Gly Ala
    930                                935                                940

Pro Ile Leu Ser Pro Glu Ala Ala Lys Ala Val Ala Val Thr Leu Val
    945                                950                                955                                960

Glu Ser Val Cys Pro Asp Ala Ala Gly Ala Glu Leu Ala Trp Pro Pro
    965                                970                                975

Glu Glu His Ala Arg Ala Thr Val Glu Arg Asp Leu Arg Ile Gly Arg
    980                                985                                990

Arg Phe Arg Glu Gln Pro Leu Leu Phe Glu Leu Leu Lys Leu Val Ala
    995                                1000                                1005

Ala Ala Pro Pro Ala Leu Cys Tyr Cys Ser Val Leu Leu Arg Gly
    1010                                1015                                1020

Leu Leu Ala Ala Leu Leu Gly His Trp Glu Ala Ser Arg His Pro
    1025                                1030                                1035

Asp Thr Thr His Ser Pro Trp His Leu Glu Ala Ser Cys Thr Leu
    1040                                1045                                1050

Val Ala Val Met Ala Glu Gly Ser Leu Leu Pro Pro Ala Leu Gly
    1055                                1060                                1065

Asn Met His Glu Val Phe Ser Gln Leu Ala Pro Phe Glu Val Arg
    1070                                1075                                1080

Leu Leu Leu Leu Ser Val Trp Gly Phe Leu Arg Glu His Gly Pro
    1085                                1090                                1095

Leu Pro Gln Lys Phe Ile Phe Gln Ser Glu Arg Gly Arg Phe Ile
    1100                                1105                                1110

Arg Asp Phe Ser Arg Glu Gly Gly Gly Glu Gly Gly Pro His Leu
    1115                                1120                                1125

Ala Val Leu His Ser Val Leu His Arg Asn Ile Asp Arg Leu Gly
    1130                                1135                                1140

Leu Phe Ser Gly Arg Phe Gln Ala Pro Ser Pro Ser Thr Leu Leu
    1145                                1150                                1155

Arg Gln Gly Thr
    1160

```

<210> 4

<211> 1162

<212> PRT

<213> Artificial

<220>

<223> A variant of the human BFLP1698 polypeptide

2003295784 19 Aug 2005

<400> 4

Met Ala Leu Val Pro Gly Arg Ser Lys Glu Asp Gly Leu Trp Thr Arg
 1 5 10 15
 Asn Ser Pro Gly Ser Ser Gln His Pro Glu Ser Pro Arg Leu Pro Asn
 20 25 30
 Pro Leu Trp Asp Arg Gly Lys Ile Gly Lys Val Glu Gly His Gln His
 35 40 45
 Ile Gln Asp Phe Ser Gln Lys Ser His Leu Pro Ser Ile Val Val Glu
 50 55 60
 Ser Ser Glu Val Asn Glu Glu Ser Gly Asp Leu His Leu Pro His Glu
 65 70 75 80
 Glu Leu Leu Leu Leu Thr Asp Gly Glu Glu Glu Asp Ala Glu Ala Phe
 85 90 95
 Phe Gln Asp Gln Ser Glu Glu Pro Gly Ala Ala Arg Pro His His Gln
 100 105 110
 Ala Arg Gln Val Glu His Ser Thr Gln Arg Gly His Leu Glu Ile Arg
 115 120 125
 Glu Leu Lys Lys Lys Leu Phe Lys Arg Arg Arg Val Leu Asn Arg Glu
 130 135 140
 Arg Arg Leu Arg His Arg Val Val Gly Ala Val Ile Asp Gln Gly Leu
 145 150 155 160
 Ile Thr Arg His His Leu Lys Lys Arg Ala Ala Gln Glu Leu Ser Gln
 165 170 175
 Glu Ile Lys Ala Phe Leu Thr Gly Val Asp Pro Ile Leu Gly His Asn
 180 185 190
 Leu Ser Ala Arg Glu His Ala Arg Cys Gly Leu Leu Leu Leu Arg Ser
 195 200 205
 Leu Pro Pro Ala Arg Ala Ala Val Leu Asp His Leu Arg Gly Val Phe
 210 215 220
 Asp Glu Ser Val Arg Ala His Leu Ala Ala Leu Asp Glu Thr Pro Val
 225 230 235 240
 Ala Gly Pro Pro His Leu Arg Pro Pro Pro Pro Ser His Val Pro Ala
 245 250 255
 Gly Gly Pro Gly Leu Glu Asp Val Val Gln Glu Val Gln Gln Val Leu
 260 265 270
 Ser Glu Phe Ile Arg Ala Asn Pro Lys Ala Trp Ala Pro Val Ile Ser
 275 280 285
 Ala Trp Ser Ile Asp Leu Met Gly Gln Leu Ser Ser Thr Tyr Ser Gly
 290 295 300
 Gln His Gln Arg Val Pro His Ala Thr Gly Ala Leu Asn Glu Leu Leu
 305 310 315 320

2003295784 19 Aug 2005

Gln	Leu	Trp	Met	Gly	Cys	Arg	Ala	Thr	Arg	Thr	Leu	Met	Asp	Ile	Tyr	325	330	335
Val	Gln	Cys	Leu	Ser	Ala	Leu	Ile	Gly	Ser	Cys	Pro	Asp	Ala	Cys	Val	340	345	350
Asp	Ala	Leu	Leu	Asp	Thr	Ser	Val	Gln	His	Ser	Pro	His	Phe	Asp	Trp	355	360	365
Val	Val	Ala	His	Ile	Gly	Ser	Ser	Phe	Pro	Gly	Thr	Ile	Ile	Ser	Arg	370	375	380
Val	Leu	Ser	Cys	Gly	Leu	Lys	Asp	Phe	Cys	Val	His	Gly	Gly	Ala	Gly	385	390	395
Gly	Gly	Ala	Gly	Ser	Ser	Gly	Gly	Ser	Ser	Ser	Gln	Thr	Pro	Ser	Thr	405	410	415
Asp	Pro	Phe	Pro	Gly	Ser	Pro	Ala	Ile	Pro	Ala	Glu	Lys	Arg	Val	Pro	420	425	430
Lys	Ile	Ala	Ser	Val	Val	Gly	Ile	Leu	Gly	His	Leu	Ala	Ser	Arg	His	435	440	445
Gly	Asp	Ser	Ile	Arg	Arg	Glu	Leu	Leu	Arg	Met	Phe	His	Asp	Ser	Leu	450	455	460
Ala	Gly	Gly	Ser	Gly	Gly	Arg	Ser	Gly	Asp	Pro	Ser	Leu	Gln	Ala	Thr	465	470	475
Val	Pro	Phe	Leu	Leu	Gln	Leu	Ala	Val	Met	Ser	Pro	Ala	Leu	Leu	Gly	485	490	495
Thr	Val	Ser	Gly	Glu	Leu	Val	Asp	Cys	Leu	Lys	Pro	Pro	Ala	Val	Leu	500	505	510
Ser	Gln	Leu	Gln	Gln	His	Leu	Gln	Gly	Phe	Pro	Arg	Glu	Glu	Leu	Asp	515	520	525
Asn	Met	Leu	Asn	Leu	Ala	Val	His	Leu	Val	Ser	Gln	Ala	Ser	Gly	Ala	530	535	540
Gly	Ala	Tyr	Arg	Leu	Leu	Gln	Phe	Leu	Val	Asp	Thr	Ala	Met	Pro	Ala	545	550	555
Ser	Val	Ile	Thr	Thr	Gln	Gly	Leu	Ala	Val	Pro	Asp	Thr	Val	Arg	Glu	565	570	575
Ala	Cys	Asp	Arg	Leu	Ile	Gln	Leu	Leu	Leu	Leu	His	Leu	Gln	Lys	Leu	580	585	590
Val	His	His	Arg	Gly	Gly	Ser	Pro	Gly	Glu	Gly	Val	Leu	Gly	Pro	Pro	595	600	605
Pro	Pro	Pro	Arg	Leu	Val	Pro	Phe	Leu	Asp	Ala	Leu	Lys	Asn	His	Val	610	615	620
Gly	Glu	Leu	Cys	Gly	Glu	Thr	Leu	Arg	Leu	Glu	Arg	Lys	Arg	Phe	Leu	625	630	635
Trp	Gln	His	Gln	Leu	Leu	Gly	Leu	Leu	Ser	Val	Tyr	Thr	Arg	Pro	Ser			

2003295784 19 Aug 2005

				645				650				655			
Cys	Gly	Pro	Glu	Ala	Leu	Gly	His	Leu	Leu	Ser	Arg	Ala	Arg	Ser	Pro
660				665				670							
Glu	Glu	Leu	Ser	Leu	Ala	Thr	Gln	Leu	Tyr	Ala	Gly	Leu	Val	Val	Ser
675				680				685							
Leu	Ser	Gly	Leu	Leu	Pro	Leu	Ala	Phe	Arg	Ser	Cys	Leu	Ala	Arg	Val
690				695				700							
His	Ala	Gly	Thr	Leu	Gln	Pro	Pro	Phe	Thr	Ala	Arg	Phe	Leu	Arg	Asn
705				710				715				720			
Leu	Ala	Leu	Leu	Val	Gly	Trp	Glu	Gln	Gln	Gly	Gly	Glu	Gly	Pro	Ala
725				730				735							
Ala	Leu	Gly	Ala	His	Phe	Gly	Glu	Ser	Ala	Ser	Ala	His	Leu	Ser	Asp
740				745				750							
Leu	Ala	Pro	Leu	Leu	Leu	His	Pro	Glu	Glu	Glu	Val	Ala	Glu	Ala	Ala
755				760				765							
Ala	Ser	Leu	Leu	Ala	Ile	Cys	Pro	Phe	Pro	Ser	Glu	Ala	Leu	Ser	Pro
770				775				780							
Ser	Gln	Leu	Leu	Gly	Leu	Val	Arg	Ala	Gly	Val	His	Arg	Phe	Phe	Ala
785				790				795				800			
Ser	Leu	Arg	Leu	His	Gly	Pro	Pro	Gly	Val	Ala	Ser	Ala	Cys	Gln	Leu
805				810				815							
Leu	Thr	Arg	Leu	Ser	Gln	Thr	Ser	Pro	Ala	Gly	Leu	Lys	Ala	Val	Leu
820				825				830							
Gln	Leu	Leu	Val	Glu	Gly	Ala	Leu	His	Arg	Gly	Asn	Thr	Glu	Leu	Phe
835				840				845							
Gly	Gly	Gln	Val	Asp	Gly	Asp	Asn	Glu	Thr	Leu	Ser	Val	Val	Ser	Ala
850				855				860							
Ser	Leu	Ala	Ser	Ala	Ser	Leu	Leu	Asp	Thr	Asn	Arg	Arg	His	Thr	Ala
865				870				875				880			
Ala	Val	Pro	Gly	Pro	Gly	Gly	Ile	Trp	Ser	Val	Phe	His	Ala	Gly	Val
885				890				895							
Ile	Gly	Arg	Gly	Leu	Lys	Pro	Pro	Lys	Phe	Val	Gln	Ser	Arg	Asn	Gln
900				905				910							
Gln	Glu	Val	Ile	Tyr	Asn	Thr	Gln	Ser	Leu	Leu	Ser	Leu	Leu	Val	His
915				920				925							
Cys	Cys	Ser	Ala	Pro	Gly	Gly	Thr	Glu	Cys	Gly	Glu	Cys	Trp	Gly	Ala
930				935				940							
Pro	Ile	Leu	Ser	Pro	Glu	Ala	Ala	Lys	Ala	Val	Ala	Val	Thr	Leu	Val
945				950				955				960			
Glu	Ser	Val	Cys	Pro	Asp	Ala	Ala	Gly	Ala	Glu	Leu	Ala	Trp	Pro	Pro
965				970				975							

2003295784 19 Aug 2005

Glu Glu His Ala Arg Ala Thr Val Glu Arg Asp Leu Arg Ile Gly Arg
 980 985 990
 Arg Phe Arg Glu Gln Pro Leu Leu Phe Glu Leu Leu Lys Leu Val Ala
 995 1000 1005
 Ala Ala Pro Pro Ala Leu Cys Tyr Cys Ser Val Leu Leu Arg Gly
 1010 1015 1020
 Leu Leu Ala Ala Leu Leu Gly His Trp Glu Ala Ser Arg His Pro
 1025 1030 1035
 Asp Thr Thr His Ser Pro Trp His Leu Glu Ala Ser Cys Thr Leu
 1040 1045 1050
 Val Ala Val Met Ala Glu Gly Ser Leu Leu Pro Pro Ala Leu Gly
 1055 1060 1065
 Asn Met His Glu Val Phe Ser Gln Leu Ala Pro Phe Glu Val Arg
 1070 1075 1080
 Leu Leu Leu Leu Ser Val Trp Gly Phe Leu Arg Glu His Gly Pro
 1085 1090 1095
 Leu Pro Gln Lys Phe Ile Phe Gln Ser Glu Arg Gly Arg Phe Ile
 1100 1105 1110
 Arg Asp Phe Ser Arg Glu Gly Gly Gly Glu Gly Gly Pro His Leu
 1115 1120 1125
 Ala Val Leu His Ser Val Leu His Arg Asn Ile Asp Arg Leu Gly
 1130 1135 1140
 Leu Phe Ser Gly Arg Phe Gln Ala Pro Ser Pro Ser Thr Leu Leu
 1145 1150 1155
 Arg Gln Gly Thr
 1160

<210> 5
 <211> 1162
 <212> PRT
 <213> Artificial

<220>
 <223> A variant of the human BFLP1698 polypeptide

<400> 5

Met Ala Leu Val Pro Gly Arg Ser Lys Glu Asp Gly Leu Trp Thr Arg
 1 5 10 15
 Asn Ser Pro Gly Ser Ser Gln His Pro Glu Ser Pro Arg Leu Pro Asn
 20 25 30
 Pro Leu Trp Asp Arg Gly Lys Ile Gly Lys Val Glu Gly His Gln His
 35 40 45
 Ile Gln Asp Phe Ser Gln Lys Ser His Leu Pro Ser Ile Val Val Glu
 50 55 60

2003295784 19 Aug 2005

Ser	Ser	Glu	Val	Asn	Glu	Glu	Ser	Gly	Asp	Leu	His	Leu	Pro	His	Glu	65	70	75	80
Glu	Leu	Leu	Leu	Leu	Thr	Asp	Gly	Glu	Glu	Glu	Asp	Ala	Glu	Ala	Phe	85	90	95	
Phe	Gln	Asp	Gln	Ser	Glu	Glu	Pro	Gly	Ala	Ala	Arg	Pro	His	His	Gln	100	105	110	
Ala	Arg	Gln	Val	Glu	His	Ser	Thr	Gln	Arg	Gly	His	Leu	Glu	Ile	Arg	115	120	125	
Glu	Leu	Lys	Lys	Lys	Leu	Phe	Lys	Arg	Arg	Arg	Val	Leu	Asn	Arg	Glu	130	135	140	
Arg	Arg	Leu	Arg	His	Arg	Val	Val	Gly	Ala	Val	Ile	Asp	Gln	Gly	Leu	145	150	155	160
Ile	Thr	Arg	His	His	Leu	Lys	Lys	Arg	Ala	Ala	Gln	Glu	Leu	Ser	Gln	165	170	175	
Glu	Ile	Lys	Ala	Phe	Leu	Thr	Gly	Val	Asp	Pro	Ile	Leu	Gly	His	Gln	180	185	190	
Leu	Ser	Ala	Arg	Glu	His	Ala	Arg	Cys	Gly	Leu	Leu	Leu	Leu	Arg	Ser	195	200	205	
Leu	Pro	Pro	Ala	Arg	Ala	Ala	Val	Leu	Asp	His	Leu	Arg	Gly	Val	Phe	210	215	220	
Asp	Glu	Ser	Val	Arg	Ala	His	Leu	Ala	Ala	Leu	Asp	Glu	Thr	Pro	Val	225	230	235	240
Ala	Gly	Pro	Pro	His	Leu	Arg	Pro	Pro	Pro	Pro	Ser	His	Val	Pro	Ala	245	250	255	
Gly	Gly	Pro	Gly	Leu	Glu	Asp	Val	Val	Gln	Glu	Val	Gln	Gln	Val	Leu	260	265	270	
Ser	Glu	Phe	Ile	Arg	Ala	Asn	Pro	Lys	Ala	Trp	Ala	Pro	Val	Ile	Gly	275	280	285	
Ala	Trp	Ser	Ile	Asp	Leu	Met	Gly	Gln	Leu	Ser	Ser	Thr	Tyr	Ser	Gly	290	295	300	
Gln	His	Gln	Arg	Val	Pro	His	Ala	Thr	Gly	Ala	Leu	Asn	Glu	Leu	Leu	305	310	315	320
Gln	Leu	Trp	Met	Gly	Cys	Arg	Ala	Thr	Arg	Thr	Leu	Met	Asp	Ile	Tyr	325	330	335	
Val	Gln	Cys	Leu	Ser	Ala	Leu	Ile	Gly	Ser	Cys	Pro	Asp	Ala	Cys	Val	340	345	350	
Asp	Ala	Leu	Leu	Asp	Thr	Ser	Val	Gln	His	Ser	Pro	His	Phe	Asp	Trp	355	360	365	
Val	Val	Ala	His	Ile	Gly	Ser	Ser	Phe	Pro	Gly	Thr	Ile	Ile	Ser	Arg	370	375	380	
Val	Leu	Ser	Cys	Gly	Leu	Lys	Asp	Phe	Cys	Val	His	Gly	Gly	Ala	Gly	385	390	395	400

2003295784 19 Aug 2005

Gly	Gly	Ala	Gly	Ser	Ser	Gly	Gly	Ser	Ser	Ser	Gln	Thr	Pro	Ser	Thr	405	410	415
Asp	Pro	Phe	Pro	Gly	Ser	Pro	Ala	Ile	Pro	Ala	Glu	Lys	Arg	Val	Pro	420	425	430
Lys	Ile	Ala	Ser	Val	Val	Gly	Ile	Leu	Gly	His	Leu	Ala	Ser	Arg	His	435	440	445
Gly	Asp	Ser	Ile	Arg	Arg	Glu	Leu	Leu	Arg	Met	Phe	His	Asp	Ser	Leu	450	455	460
Ala	Gly	Gly	Ser	Gly	Gly	Arg	Ser	Gly	Asp	Pro	Ser	Leu	Gln	Ala	Thr	465	470	475
Val	Pro	Phe	Leu	Leu	Gln	Leu	Ala	Val	Met	Ser	Pro	Ala	Leu	Leu	Gly	485	490	495
Thr	Val	Ser	Gly	Glu	Leu	Val	Asp	Cys	Leu	Lys	Pro	Pro	Ala	Val	Leu	500	505	510
Ser	Gln	Leu	Gln	Gln	His	Leu	Gln	Gly	Phe	Pro	Arg	Glu	Glu	Leu	Asp	515	520	525
Asn	Met	Leu	Asn	Leu	Ala	Val	His	Leu	Val	Ser	Gln	Ala	Ser	Gly	Ala	530	535	540
Gly	Ala	Tyr	Arg	Leu	Leu	Gln	Phe	Leu	Val	Asp	Thr	Ala	Met	Pro	Ala	545	550	555
Ser	Val	Ile	Thr	Thr	Gln	Gly	Leu	Ala	Val	Pro	Asp	Thr	Val	Arg	Glu	565	570	575
Ala	Cys	Asp	Arg	Leu	Ile	Gln	Leu	Leu	Leu	His	Leu	Gln	Lys	Leu		580	585	590
Val	His	His	Arg	Gly	Gly	Ser	Pro	Gly	Glu	Gly	Val	Leu	Gly	Pro	Pro	595	600	605
Pro	Pro	Pro	Arg	Leu	Val	Pro	Phe	Leu	Asp	Ala	Leu	Lys	Asn	His	Val	610	615	620
Gly	Glu	Leu	Cys	Gly	Glu	Thr	Leu	Arg	Leu	Glu	Arg	Lys	Arg	Phe	Leu	625	630	635
Trp	Gln	His	Gln	Leu	Leu	Gly	Leu	Leu	Ser	Val	Tyr	Thr	Arg	Pro	Ser	645	650	655
Cys	Gly	Pro	Glu	Ala	Leu	Gly	His	Leu	Leu	Ser	Arg	Ala	Arg	Ser	Pro	660	665	670
Glu	Glu	Leu	Ser	Leu	Ala	Thr	Gln	Leu	Tyr	Ala	Gly	Leu	Val	Val	Ser	675	680	685
Leu	Ser	Gly	Leu	Leu	Pro	Leu	Ala	Phe	Arg	Ser	Cys	Leu	Ala	Arg	Val	690	695	700
His	Ala	Gly	Thr	Leu	Gln	Pro	Pro	Phe	Thr	Ala	Arg	Phe	Leu	Arg	Asn	705	710	715
Leu	Ala	Leu	Leu	Val	Gly	Trp	Glu	Gln	Gln	Gly	Gly	Glu	Gly	Pro	Ala			

2003295784 19 Aug 2005

725										730					735				
Ala	Leu	Gly	Ala	His	Phe	Gly	Glu	Ser	Ala	Ser	Ala	His	Leu	Ser	Asp				
			740						745				750						
Leu	Ala	Pro	Leu	Leu	Leu	His	Pro	Glu	Glu	Glu	Val	Ala	Glu	Ala	Ala				
		755					760					765							
Ala	Ser	Leu	Leu	Ala	Ile	Cys	Pro	Phe	Pro	Ser	Glu	Ala	Leu	Ser	Pro				
		770				775					780								
Ser	Gln	Leu	Leu	Gly	Leu	Val	Arg	Ala	Gly	Val	His	Arg	Phe	Phe	Ala				
		785			790						795				800				
Ser	Leu	Arg	Leu	His	Gly	Pro	Pro	Gly	Val	Ala	Ser	Ala	Cys	Gln	Leu				
				805					810					815					
Leu	Thr	Arg	Leu	Ser	Gln	Thr	Ser	Pro	Ala	Gly	Leu	Lys	Ala	Val	Leu				
			820					825					830						
Gln	Leu	Leu	Val	Glu	Gly	Ala	Leu	His	Arg	Gly	Asn	Thr	Glu	Leu	Phe				
		835					840					845							
Gly	Gly	Gln	Val	Asp	Gly	Asp	Asn	Glu	Thr	Leu	Ser	Val	Val	Ser	Ala				
		850				855					860								
Ser	Leu	Ala	Ser	Ala	Ser	Leu	Leu	Asp	Thr	Asn	Arg	Arg	His	Thr	Ala				
					870						875				880				
Ala	Val	Pro	Gly	Pro	Gly	Gly	Ile	Trp	Ser	Val	Phe	His	Ala	Gly	Val				
				885					890					895					
Ile	Gly	Arg	Gly	Leu	Lys	Pro	Pro	Lys	Phe	Val	Gln	Ser	Arg	Asn	Gln				
			900					905					910						
Gln	Glu	Val	Ile	Tyr	Asn	Thr	Gln	Ser	Leu	Leu	Ser	Leu	Leu	Val	His				
		915					920					925							
Cys	Cys	Ser	Ala	Pro	Gly	Gly	Thr	Glu	Cys	Gly	Glu	Cys	Trp	Gly	Ala				
		930				935					940								
Pro	Ile	Leu	Ser	Pro	Glu	Ala	Ala	Lys	Ala	Val	Ala	Val	Thr	Leu	Val				
					950						955				960				
Glu	Ser	Val	Cys	Pro	Asp	Ala	Ala	Gly	Ala	Glu	Leu	Ala	Trp	Pro	Pro				
				965					970					975					
Glu	Glu	His	Ala	Arg	Ala	Thr	Val	Glu	Arg	Asp	Leu	Arg	Ile	Gly	Arg				
			980					985					990						
Arg	Phe	Arg	Glu	Gln	Pro	Leu	Leu	Phe	Glu	Leu	Leu	Lys	Leu	Val	Ala				
			995				1000						1005						
Ala	Ala	Pro	Pro	Ala	Leu	Cys	Tyr	Cys	Ser	Val	Leu	Leu	Arg	Gly					
						1015					1020								
Leu	Leu	Ala	Ala	Leu	Leu	Gly	His	Trp	Glu	Ala	Ser	Arg	His	Pro					
						1030					1035								
Asp	Thr	Thr	His	Ser	Pro	Trp	His	Leu	Glu	Ala	Ser	Cys	Thr	Leu					
						1045					1050								

2003295784 19 Aug 2005

Val Ala Val Met Ala Glu Gly Ser Leu Leu Pro Pro Ala Leu Gly
 1055 1060 1065
 Asn Met His Glu Val Phe Ser Gln Leu Ala Pro Phe Glu Val Arg
 1070 1075 1080
 Leu Leu Leu Leu Ser Val Trp Gly Phe Leu Arg Glu His Gly Pro
 1085 1090 1095
 Leu Pro Gln Lys Phe Ile Phe Gln Ser Glu Arg Gly Arg Phe Ile
 1100 1105 1110
 Arg Asp Phe Ser Arg Glu Gly Gly Gly Glu Gly Gly Pro His Leu
 1115 1120 1125
 Ala Val Leu His Ser Val Leu His Arg Asn Ile Asp Arg Leu Gly
 1130 1135 1140
 Leu Phe Ser Gly Arg Phe Gln Ala Pro Ser Pro Ser Thr Leu Leu
 1145 1150 1155
 Arg Gln Gly Thr
 1160

<210> 6
 <211> 1162
 <212> PRT
 <213> Artificial

<220>
 <223> A variant of the human BFLP1698 polypeptide

<400> 6

Met Ala Leu Val Pro Gly Arg Ser Lys Glu Asp Gly Leu Trp Thr Arg
 1 5 10 15
 Asn Ser Pro Gly Ser Ser Gln His Pro Glu Ser Pro Arg Leu Pro Asn
 20 25 30
 Pro Leu Trp Asp Arg Gly Lys Ile Gly Lys Val Glu Gly His Gln His
 35 40 45
 Ile Gln Asp Phe Ser Gln Lys Ser His Leu Pro Ser Ile Val Val Glu
 50 55 60
 Ser Ser Glu Val Asn Glu Glu Ser Gly Asp Leu His Leu Pro His Glu
 65 70 75 80
 Glu Leu Leu Leu Leu Thr Asp Gly Glu Glu Glu Asp Ala Glu Ala Phe
 85 90 95
 Phe Gln Asp Gln Ser Glu Glu Pro Gly Ala Ala Arg Pro His His Gln
 100 105 110
 Ala Arg Gln Val Glu His Ser Thr Gln Arg Gly His Leu Glu Ile Arg
 115 120 125
 Glu Leu Lys Lys Lys Leu Phe Lys Arg Arg Arg Val Leu Asn Arg Glu
 130 135 140

2003295784 19 Aug 2005

Arg	Arg	Leu	Arg	His	Arg	Val	Val	Gly	Ala	Val	Ile	Asp	Gln	Gly	Leu	
145					150					155					160	
Ile	Thr	Arg	His	His	Leu	Lys	Lys	Arg	Ala	Ala	Gln	Glu	Leu	Ser	Gln	
				165					170					175		
Glu	Ile	Lys	Ala	Phe	Leu	Thr	Gly	Val	Asp	Pro	Ile	Leu	Gly	His	Gln	
			180					185					190			
Leu	Ser	Ala	Arg	Glu	His	Ala	Arg	Cys	Gly	Leu	Leu	Leu	Leu	Arg	Ser	
		195					200					205				
Leu	Pro	Pro	Ala	Arg	Ala	Ala	Val	Leu	Asp	His	Leu	Arg	Gly	Val	Phe	
	210					215					220					
Asp	Glu	Ser	Val	Arg	Ala	His	Leu	Ala	Ala	Leu	Asp	Glu	Thr	Pro	Val	
225					230					235					240	
Ala	Gly	Pro	Pro	His	Leu	Arg	Pro	Pro	Pro	Pro	Ser	His	Val	Pro	Ala	
				245					250					255		
Gly	Gly	Pro	Gly	Leu	Glu	Asp	Val	Val	Gln	Glu	Val	Gln	Gln	Val	Leu	
			260					265					270			
Ser	Glu	Phe	Ile	Arg	Ala	Asn	Pro	Lys	Ala	Trp	Ala	Pro	Val	Ile	Ser	
		275					280					285				
Ala	Trp	Ser	Ile	Asp	Leu	Met	Gly	Gln	Leu	Ser	Ser	Thr	Tyr	Ser	Gly	
	290					295					300					
Gln	His	Gln	Arg	Val	Pro	His	Ala	Thr	Gly	Ala	Leu	Asn	Glu	Leu	Leu	
305					310					315					320	
Gln	Leu	Trp	Met	Gly	Cys	Arg	Ala	Thr	Arg	Thr	Leu	Met	Asp	Ile	Tyr	
			325						330					335		
Val	Gln	Cys	Leu	Ser	Ala	Leu	Ile	Gly	Ser	Cys	Pro	Asp	Ala	Cys	Val	
			340					345					350			
Asp	Ala	Leu	Leu	Asp	Thr	Ser	Val	Gln	His	Ser	Pro	Arg	Phe	Asp	Trp	
		355					360					365				
Val	Val	Ala	His	Ile	Gly	Ser	Ser	Phe	Pro	Gly	Thr	Ile	Ile	Ser	Arg	
	370					375					380					
Val	Leu	Ser	Cys	Gly	Leu	Lys	Asp	Phe	Cys	Val	His	Gly	Gly	Ala	Gly	
385					390					395					400	
Gly	Gly	Ala	Gly	Ser	Ser	Gly	Gly	Ser	Ser	Ser	Gln	Thr	Pro	Ser	Thr	
				405					410					415		
Asp	Pro	Phe	Pro	Gly	Ser	Pro	Ala	Ile	Pro	Ala	Glu	Lys	Arg	Val	Pro	
			420					425					430			
Lys	Ile	Ala	Ser	Val	Val	Gly	Ile	Leu	Gly	His	Leu	Ala	Ser	Arg	His	
		435					440					445				
Gly	Asp	Ser	Ile	Arg	Arg	Glu	Leu	Leu	Arg	Met	Phe	His	Asp	Ser	Leu	
	450					455					460					
Ala	Gly	Gly	Ser	Gly	Gly	Arg	Ser	Gly	Asp	Pro	Ser	Leu	Gln	Ala	Thr	
465					470					475					480	

2003295784 19 Aug 2005

Val	Pro	Phe	Leu	Leu	Gln	Leu	Ala	Val	Met	Ser	Pro	Ala	Leu	Leu	Gly	485	490	495
Thr	Val	Ser	Gly	Glu	Leu	Val	Asp	Cys	Leu	Lys	Pro	Pro	Ala	Val	Leu	500	505	510
Ser	Gln	Leu	Gln	Gln	His	Leu	Gln	Gly	Phe	Pro	Arg	Glu	Glu	Leu	Asp	515	520	525
Asn	Met	Leu	Asn	Leu	Ala	Val	His	Leu	Val	Ser	Gln	Ala	Ser	Gly	Ala	530	535	540
Gly	Ala	Tyr	Arg	Leu	Leu	Gln	Phe	Leu	Val	Asp	Thr	Ala	Met	Pro	Ala	545	550	555
Ser	Val	Ile	Thr	Thr	Gln	Gly	Leu	Ala	Val	Pro	Asp	Thr	Val	Arg	Glu	565	570	575
Ala	Cys	Asp	Arg	Leu	Ile	Gln	Leu	Leu	Leu	His	Leu	Gln	Lys	Leu		580	585	590
Val	His	His	Arg	Gly	Gly	Ser	Pro	Gly	Glu	Gly	Val	Leu	Gly	Pro	Pro	595	600	605
Pro	Pro	Pro	Arg	Leu	Val	Pro	Phe	Leu	Asp	Ala	Leu	Lys	Asn	His	Val	610	615	620
Gly	Glu	Leu	Cys	Gly	Glu	Thr	Leu	Arg	Leu	Glu	Arg	Lys	Arg	Phe	Leu	625	630	635
Trp	Gln	His	Gln	Leu	Leu	Gly	Leu	Leu	Ser	Val	Tyr	Thr	Arg	Pro	Ser	645	650	655
Cys	Gly	Pro	Glu	Ala	Leu	Gly	His	Leu	Leu	Ser	Arg	Ala	Arg	Ser	Pro	660	665	670
Glu	Glu	Leu	Ser	Leu	Ala	Thr	Gln	Leu	Tyr	Ala	Gly	Leu	Val	Val	Ser	675	680	685
Leu	Ser	Gly	Leu	Leu	Pro	Leu	Ala	Phe	Arg	Ser	Cys	Leu	Ala	Arg	Val	690	695	700
His	Ala	Gly	Thr	Leu	Gln	Pro	Pro	Phe	Thr	Ala	Arg	Phe	Leu	Arg	Asn	705	710	715
Leu	Ala	Leu	Leu	Val	Gly	Trp	Glu	Gln	Gln	Gly	Gly	Glu	Gly	Pro	Ala	725	730	735
Ala	Leu	Gly	Ala	His	Phe	Gly	Glu	Ser	Ala	Ser	Ala	His	Leu	Ser	Asp	740	745	750
Leu	Ala	Pro	Leu	Leu	Leu	His	Pro	Glu	Glu	Glu	Val	Ala	Glu	Ala	Ala	755	760	765
Ala	Ser	Leu	Leu	Ala	Ile	Cys	Pro	Phe	Pro	Ser	Glu	Ala	Leu	Ser	Pro	770	775	780
Ser	Gln	Leu	Leu	Gly	Leu	Val	Arg	Ala	Gly	Val	His	Arg	Phe	Phe	Ala	785	790	795
Ser	Leu	Arg	Leu	His	Gly	Pro	Pro	Gly	Val	Ala	Ser	Ala	Cys	Gln	Leu			

2003295784 19 Aug 2005

805					810					815						
Leu	Thr	Arg	Leu	Ser	Gln	Thr	Ser	Pro	Ala	Gly	Leu	Lys	Ala	Val	Leu	
820					825					830						
Gln	Leu	Leu	Val	Glu	Gly	Ala	Leu	His	Arg	Gly	Asn	Thr	Glu	Leu	Phe	
835					840					845						
Gly	Gly	Gln	Val	Asp	Gly	Asp	Asn	Glu	Thr	Leu	Ser	Val	Val	Ser	Ala	
850					855					860						
Ser	Leu	Ala	Ser	Ala	Ser	Leu	Leu	Asp	Thr	Asn	Arg	Arg	His	Thr	Ala	
865					870					875					880	
Ala	Val	Pro	Gly	Pro	Gly	Gly	Ile	Trp	Ser	Val	Phe	His	Ala	Gly	Val	
885					890					895						
Ile	Gly	Arg	Gly	Leu	Lys	Pro	Pro	Lys	Phe	Val	Gln	Ser	Arg	Asn	Gln	
900					905					910						
Gln	Glu	Val	Ile	Tyr	Asn	Thr	Gln	Ser	Leu	Leu	Ser	Leu	Leu	Val	His	
915					920					925						
Cys	Cys	Ser	Ala	Pro	Gly	Gly	Thr	Glu	Cys	Gly	Glu	Cys	Trp	Gly	Ala	
930					935					940						
Pro	Ile	Leu	Ser	Pro	Glu	Ala	Ala	Lys	Ala	Val	Ala	Val	Thr	Leu	Val	
945					950					955					960	
Glu	Ser	Val	Cys	Pro	Asp	Ala	Ala	Gly	Ala	Glu	Leu	Ala	Trp	Pro	Pro	
965					970					975						
Glu	Glu	His	Ala	Arg	Ala	Thr	Val	Glu	Arg	Asp	Leu	Arg	Ile	Gly	Arg	
980					985					990						
Arg	Phe	Arg	Glu	Gln	Pro	Leu	Leu	Phe	Glu	Leu	Leu	Lys	Leu	Val	Ala	
995					1000					1005						
Ala	Ala	Pro	Pro	Ala	Leu	Cys	Tyr	Cys	Ser	Val	Leu	Leu	Arg	Gly		
1010					1015					1020						
Leu	Leu	Ala	Ala	Leu	Leu	Gly	His	Trp	Glu	Ala	Ser	Arg	His	Pro		
1025					1030					1035						
Asp	Thr	Thr	His	Ser	Pro	Trp	His	Leu	Glu	Ala	Ser	Cys	Thr	Leu		
1040					1045					1050						
Val	Ala	Val	Met	Ala	Glu	Gly	Ser	Leu	Leu	Pro	Pro	Ala	Leu	Gly		
1055					1060					1065						
Asn	Met	His	Glu	Val	Phe	Ser	Gln	Leu	Ala	Pro	Phe	Glu	Val	Arg		
1070					1075					1080						
Leu	Leu	Leu	Leu	Ser	Val	Trp	Gly	Phe	Leu	Arg	Glu	His	Gly	Pro		
1085					1090					1095						
Leu	Pro	Gln	Lys	Phe	Ile	Phe	Gln	Ser	Glu	Arg	Gly	Arg	Phe	Ile		
1100					1105					1110						
Arg	Asp	Phe	Ser	Arg	Glu	Gly	Gly	Gly	Glu	Gly	Gly	Pro	His	Leu		
1115					1120					1125						

2003295784 19 Aug 2005

Ala Val Leu His Ser Val Leu His Arg Asn Ile Asp Arg Leu Gly
1130 1135 1140

Leu Phe Ser Gly Arg Phe Gln Ala Pro Ser Pro Ser Thr Leu Leu
1145 1150 1155

Arg Gln Gly Thr
1160

<210> 7

<211> 1162

<212> PRT

<213> Artificial

<220>

<223> A variant of the human BFLP1698 polypeptide

<400> 7

Met Ala Leu Val Pro Gly Arg Ser Lys Glu Asp Gly Leu Trp Thr Arg
1 5 10 15

Asn Ser Pro Gly Ser Ser Gln His Pro Glu Ser Pro Arg Leu Pro Asn
20 25 30

Pro Leu Trp Asp Arg Gly Lys Ile Gly Lys Val Glu Gly His Gln His
35 40 45

Ile Gln Asp Phe Ser Gln Lys Ser His Leu Pro Ser Ile Val Val Glu
50 55 60

Ser Ser Glu Val Asn Glu Glu Ser Gly Asp Leu His Leu Pro His Glu
65 70 75 80

Glu Leu Leu Leu Leu Thr Asp Gly Glu Glu Glu Asp Ala Glu Ala Phe
85 90 95

Phe Gln Asp Gln Ser Glu Glu Pro Gly Ala Ala Arg Pro His His Gln
100 105 110

Ala Arg Gln Val Glu His Ser Thr Gln Arg Gly His Leu Glu Ile Arg
115 120 125

Glu Leu Lys Lys Lys Leu Phe Lys Arg Arg Arg Val Leu Asn Arg Glu
130 135 140

Arg Arg Leu Arg His Arg Val Val Gly Ala Val Ile Asp Gln Gly Leu
145 150 155 160

Ile Thr Arg His His Leu Lys Lys Arg Ala Ala Gln Glu Leu Ser Gln
165 170 175

Glu Ile Lys Ala Phe Leu Thr Gly Val Asp Pro Ile Leu Gly His Gln
180 185 190

Leu Ser Ala Arg Glu His Ala Arg Cys Gly Leu Leu Leu Leu Arg Ser
195 200 205

Leu Pro Pro Ala Arg Ala Ala Val Leu Asp His Leu Arg Gly Val Phe
210 215 220

2003295784 19 Aug 2005

Asp	Glu	Ser	Val	Arg	Ala	His	Leu	Ala	Ala	Leu	Asp	Glu	Thr	Pro	Val	225	230	235	240
Ala	Gly	Pro	Pro	His	Leu	Arg	Pro	Pro	Pro	Pro	Ser	His	Val	Pro	Ala	245	250		255
Gly	Gly	Pro	Gly	Leu	Glu	Asp	Val	Val	Gln	Glu	Val	Gln	Gln	Val	Leu	260	265		270
Ser	Glu	Phe	Ile	Arg	Ala	Asn	Pro	Lys	Ala	Trp	Ala	Pro	Val	Ile	Ser	275	280		285
Ala	Trp	Ser	Ile	Asp	Leu	Met	Gly	Gln	Leu	Ser	Ser	Thr	Tyr	Ser	Gly	290	295		300
Gln	His	Gln	Arg	Val	Pro	His	Ala	Thr	Gly	Ala	Leu	Asn	Glu	Leu	Leu	305	310	315	320
Gln	Leu	Trp	Met	Gly	Cys	Arg	Ala	Thr	Arg	Thr	Leu	Met	Asp	Ile	Tyr	325	330		335
Val	Gln	Cys	Leu	Ser	Ala	Leu	Ile	Gly	Ser	Cys	Pro	Asp	Ala	Cys	Val	340	345		350
Asp	Ala	Leu	Leu	Asp	Thr	Ser	Val	Gln	His	Ser	Pro	His	Phe	Asp	Trp	355	360		365
Val	Val	Ala	His	Ile	Gly	Ser	Ser	Phe	Pro	Gly	Thr	Ile	Ile	Ser	Arg	370	375		380
Val	Leu	Ser	Cys	Gly	Leu	Lys	Asp	Phe	Cys	Val	His	Gly	Gly	Ala	Gly	385	390	395	400
Gly	Gly	Ala	Gly	Ser	Ser	Gly	Gly	Ser	Ser	Ser	Gln	Thr	Pro	Ser	Thr	405	410		415
Asp	Pro	Phe	Pro	Gly	Ser	Pro	Ala	Ile	Pro	Ala	Glu	Lys	Arg	Val	Pro	420	425		430
Lys	Ile	Ala	Ser	Val	Val	Gly	Ile	Leu	Gly	His	Leu	Ala	Ser	Arg	His	435	440		445
Gly	Asp	Ser	Ile	Arg	Arg	Glu	Leu	Leu	Arg	Met	Phe	His	Asp	Ser	Leu	450	455	460	
Ala	Gly	Gly	Ser	Gly	Gly	Arg	Ser	Gly	Asp	Pro	Ser	Leu	Gln	Ala	Thr	465	470	475	480
Met	Pro	Phe	Leu	Leu	Gln	Leu	Ala	Val	Met	Ser	Pro	Ala	Leu	Leu	Gly	485	490		495
Thr	Val	Ser	Gly	Glu	Leu	Val	Asp	Cys	Leu	Lys	Pro	Pro	Ala	Val	Leu	500	505		510
Ser	Gln	Leu	Gln	Gln	His	Leu	Gln	Gly	Phe	Pro	Arg	Glu	Glu	Leu	Asp	515	520	525	
Asn	Met	Leu	Asn	Leu	Ala	Val	His	Leu	Val	Ser	Gln	Ala	Ser	Gly	Ala	530	535	540	
Gly	Ala	Tyr	Arg	Leu	Leu	Gln	Phe	Leu	Val	Asp	Thr	Ala	Met	Pro	Ala	545	550	555	560

2003295784 19 Aug 2005

Ser Val Ile Thr Thr Gln Gly Leu Ala Val Pro Asp Thr Val Arg Glu
 565 570 575
 Ala Cys Asp Arg Leu Ile Gln Leu Leu Leu His Leu Gln Lys Leu
 580 585 590
 Val His His Arg Gly Gly Ser Pro Gly Glu Gly Val Leu Gly Pro Pro
 595 600 605
 Pro Pro Pro Arg Leu Val Pro Phe Leu Asp Ala Leu Lys Asn His Val
 610 615 620
 Gly Glu Leu Cys Gly Glu Thr Leu Arg Leu Glu Arg Lys Arg Phe Leu
 625 630 635 640
 Trp Gln His Gln Leu Leu Gly Leu Leu Ser Val Tyr Thr Arg Pro Ser
 645 650 655
 Cys Gly Pro Glu Ala Leu Gly His Leu Leu Ser Arg Ala Arg Ser Pro
 660 665 670
 Glu Glu Leu Ser Leu Ala Thr Gln Leu Tyr Ala Gly Leu Val Val Ser
 675 680 685
 Leu Ser Gly Leu Leu Pro Leu Ala Phe Arg Ser Cys Leu Ala Arg Val
 690 695 700
 His Ala Gly Thr Leu Gln Pro Pro Phe Thr Ala Arg Phe Leu Arg Asn
 705 710 715 720
 Leu Ala Leu Leu Val Gly Trp Glu Gln Gln Gly Gly Glu Gly Pro Ala
 725 730 735
 Ala Leu Gly Ala His Phe Gly Glu Ser Ala Ser Ala His Leu Ser Asp
 740 745 750
 Leu Ala Pro Leu Leu Leu His Pro Glu Glu Glu Val Ala Glu Ala Ala
 755 760 765
 Ala Ser Leu Leu Ala Ile Cys Pro Phe Pro Ser Glu Ala Leu Ser Pro
 770 775 780
 Ser Gln Leu Leu Gly Leu Val Arg Ala Gly Val His Arg Phe Phe Ala
 785 790 795 800
 Ser Leu Arg Leu His Gly Pro Pro Gly Val Ala Ser Ala Cys Gln Leu
 805 810 815
 Leu Thr Arg Leu Ser Gln Thr Ser Pro Ala Gly Leu Lys Ala Val Leu
 820 825 830
 Gln Leu Leu Val Glu Gly Ala Leu His Arg Gly Asn Thr Glu Leu Phe
 835 840 845
 Gly Gly Gln Val Asp Gly Asp Asn Glu Thr Leu Ser Val Val Ser Ala
 850 855 860
 Ser Leu Ala Ser Ala Ser Leu Leu Asp Thr Asn Arg Arg His Thr Ala
 865 870 875 880
 Ala Val Pro Gly Pro Gly Gly Ile Trp Ser Val Phe His Ala Gly Val

2003295784 19 Aug 2005

	885	890	895
Ile Gly Arg Gly Leu Lys Pro Pro Lys Phe Val Gln Ser Arg Asn Gln	900	905	910
Gln Glu Val Ile Tyr Asn Thr Gln Ser Leu Leu Ser Leu Leu Val His	915	920	925
Cys Cys Ser Ala Pro Gly Gly Thr Glu Cys Gly Glu Cys Trp Gly Ala	930	935	940
Pro Ile Leu Ser Pro Glu Ala Ala Lys Ala Val Ala Val Thr Leu Val	945	950	955
Glu Ser Val Cys Pro Asp Ala Ala Gly Ala Glu Leu Ala Trp Pro Pro	965	970	975
Glu Glu His Ala Arg Ala Thr Val Glu Arg Asp Leu Arg Ile Gly Arg	980	985	990
Arg Phe Arg Glu Gln Pro Leu Leu Phe Glu Leu Leu Lys Leu Val Ala	995	1000	1005
Ala Ala Pro Pro Ala Leu Cys Tyr Cys Ser Val Leu Leu Arg Gly	1010	1015	1020
Leu Leu Ala Ala Leu Leu Gly His Trp Glu Ala Ser Arg His Pro	1025	1030	1035
Asp Thr Thr His Ser Pro Trp His Leu Glu Ala Ser Cys Thr Leu	1040	1045	1050
Val Ala Val Met Ala Glu Gly Ser Leu Leu Pro Pro Ala Leu Gly	1055	1060	1065
Asn Met His Glu Val Phe Ser Gln Leu Ala Pro Phe Glu Val Arg	1070	1075	1080
Leu Leu Leu Leu Ser Val Trp Gly Phe Leu Arg Glu His Gly Pro	1085	1090	1095
Leu Pro Gln Lys Phe Ile Phe Gln Ser Glu Arg Gly Arg Phe Ile	1100	1105	1110
Arg Asp Phe Ser Arg Glu Gly Gly Gly Glu Gly Gly Pro His Leu	1115	1120	1125
Ala Val Leu His Ser Val Leu His Arg Asn Ile Asp Arg Leu Gly	1130	1135	1140
Leu Phe Ser Gly Arg Phe Gln Ala Pro Ser Pro Ser Thr Leu Leu	1145	1150	1155
Arg Gln Gly Thr	1160		

<210> 8
 <211> 1162
 <212> PRT
 <213> Artificial

2003295784 19 Aug 2005

<220>

<223> A variant of the human BFLP1698 polypeptide

<400> 8

```

Met Ala Leu Val Pro Gly Arg Ser Lys Glu Asp Gly Leu Trp Thr Arg
1          5          10          15

Asn Ser Pro Gly Ser Ser Gln His Pro Glu Ser Pro Arg Leu Pro Asn
          20          25          30

Pro Leu Trp Asp Arg Gly Lys Ile Gly Lys Val Glu Gly His Gln His
          35          40          45

Ile Gln Asp Phe Ser Gln Lys Ser His Leu Pro Ser Ile Val Val Glu
          50          55          60

Ser Ser Glu Val Asn Glu Glu Ser Gly Asp Leu His Leu Pro His Glu
65          70          75          80

Glu Leu Leu Leu Leu Thr Asp Gly Glu Glu Glu Asp Ala Glu Ala Phe
          85          90          95

Phe Gln Asp Gln Ser Glu Glu Pro Gly Ala Ala Arg Pro His His Gln
          100          105          110

Ala Arg Gln Val Glu His Ser Thr Gln Arg Gly His Leu Glu Ile Arg
          115          120          125

Glu Leu Lys Lys Lys Leu Phe Lys Arg Arg Arg Val Leu Asn Arg Glu
          130          135          140

Arg Arg Leu Arg His Arg Val Val Gly Ala Val Ile Asp Gln Gly Leu
          145          150          155          160

Ile Thr Arg His His Leu Lys Lys Arg Ala Ala Gln Glu Leu Ser Gln
          165          170          175

Glu Ile Lys Ala Phe Leu Thr Gly Val Asp Pro Ile Leu Gly His Gln
          180          185          190

Leu Ser Ala Arg Glu His Ala Arg Cys Gly Leu Leu Leu Leu Arg Ser
          195          200          205

Leu Pro Pro Ala Arg Ala Ala Val Leu Asp His Leu Arg Gly Val Phe
          210          215          220

Asp Glu Ser Val Arg Ala His Leu Ala Ala Leu Asp Glu Thr Pro Val
          225          230          235          240

Ala Gly Pro Pro His Leu Arg Pro Pro Pro Pro Ser His Val Pro Ala
          245          250          255

Gly Gly Pro Gly Leu Glu Asp Val Val Gln Glu Val Gln Gln Val Leu
          260          265          270

Ser Glu Phe Ile Arg Ala Asn Pro Lys Ala Trp Ala Pro Val Ile Ser
          275          280          285

Ala Trp Ser Ile Asp Leu Met Gly Gln Leu Ser Ser Thr Tyr Ser Gly
          290          295          300

```

2003295784 19 Aug 2005

Gln	His	Gln	Arg	Val	Pro	His	Ala	Thr	Gly	Ala	Leu	Asn	Glu	Leu	Leu	305	310	315	320
Gln	Leu	Trp	Met	Gly	Cys	Arg	Ala	Thr	Arg	Thr	Leu	Met	Asp	Ile	Tyr	325	330	335	
Val	Gln	Cys	Leu	Ser	Ala	Leu	Ile	Gly	Ser	Cys	Pro	Asp	Ala	Cys	Val	340	345	350	
Asp	Ala	Leu	Leu	Asp	Thr	Ser	Val	Gln	His	Ser	Pro	His	Phe	Asp	Trp	355	360	365	
Val	Val	Ala	His	Ile	Gly	Ser	Ser	Phe	Pro	Gly	Thr	Ile	Ile	Ser	Arg	370	375	380	
Val	Leu	Ser	Cys	Gly	Leu	Lys	Asp	Phe	Cys	Val	His	Gly	Gly	Ala	Gly	385	390	395	400
Gly	Gly	Ala	Gly	Ser	Ser	Gly	Gly	Ser	Ser	Ser	Gln	Thr	Pro	Ser	Thr	405	410	415	
Asp	Pro	Phe	Pro	Gly	Ser	Pro	Ala	Ile	Pro	Ala	Glu	Lys	Arg	Val	Pro	420	425	430	
Lys	Ile	Ala	Ser	Val	Val	Gly	Ile	Leu	Gly	His	Leu	Ala	Ser	Arg	His	435	440	445	
Gly	Asp	Ser	Ile	Arg	Arg	Glu	Leu	Leu	Arg	Met	Phe	His	Asp	Ser	Leu	450	455	460	
Ala	Gly	Gly	Ser	Gly	Gly	Arg	Ser	Gly	Asp	Pro	Ser	Leu	Gln	Ala	Thr	465	470	475	480
Val	Pro	Phe	Leu	Leu	Gln	Leu	Ala	Val	Met	Ser	Pro	Ala	Leu	Leu	Gly	485	490	495	
Thr	Val	Ser	Gly	Glu	Leu	Val	Asp	Cys	Leu	Lys	Pro	Pro	Ala	Val	Leu	500	505	510	
Ser	Gln	Leu	Gln	Gln	His	Leu	Gln	Gly	Phe	Pro	Arg	Glu	Glu	Leu	Asp	515	520	525	
Asn	Met	Leu	Asn	Leu	Ala	Val	His	Leu	Val	Ser	Gln	Ala	Ser	Gly	Ala	530	535	540	
Gly	Ala	Tyr	Arg	Leu	Leu	Gln	Phe	Leu	Val	Asp	Tyr	Ala	Met	Pro	Ala	545	550	555	560
Ser	Val	Ile	Thr	Thr	Gln	Gly	Leu	Ala	Val	Pro	Asp	Thr	Val	Arg	Glu	565	570	575	
Ala	Cys	Asp	Arg	Leu	Ile	Gln	Leu	Leu	Leu	His	Leu	Gln	Lys	Leu		580	585	590	
Val	His	His	Arg	Gly	Gly	Ser	Pro	Gly	Glu	Gly	Val	Leu	Gly	Pro	Pro	595	600	605	
Pro	Pro	Pro	Arg	Leu	Val	Pro	Phe	Leu	Asp	Ala	Leu	Lys	Asn	His	Val	610	615	620	
Gly	Glu	Leu	Cys	Gly	Glu	Thr	Leu	Arg	Leu	Glu	Arg	Lys	Arg	Phe	Leu	625	630	635	640

2003295784 19 Aug 2005

Trp	Gln	His	Gln	Leu	Leu	Gly	Leu	Leu	Ser	Val	Tyr	Thr	Arg	Pro	Ser	
				645					650						655	
Cys	Gly	Pro	Glu	Ala	Leu	Gly	His	Leu	Leu	Ser	Arg	Ala	Arg	Ser	Pro	
			660					665						670		
Glu	Glu	Leu	Ser	Leu	Ala	Thr	Gln	Leu	Tyr	Ala	Gly	Leu	Val	Val	Ser	
		675					680					685				
Leu	Ser	Gly	Leu	Leu	Pro	Leu	Ala	Phe	Arg	Ser	Cys	Leu	Ala	Arg	Val	
	690					695					700					
His	Ala	Gly	Thr	Leu	Gln	Pro	Pro	Phe	Thr	Ala	Arg	Phe	Leu	Arg	Asn	
705					710					715					720	
Leu	Ala	Leu	Leu	Val	Gly	Trp	Glu	Gln	Gln	Gly	Gly	Glu	Gly	Pro	Ala	
				725				730						735		
Ala	Leu	Gly	Ala	His	Phe	Gly	Glu	Ser	Ala	Ser	Ala	His	Leu	Ser	Asp	
			740					745						750		
Leu	Ala	Pro	Leu	Leu	Leu	His	Pro	Glu	Glu	Glu	Val	Ala	Glu	Ala	Ala	
		755					760					765				
Ala	Ser	Leu	Leu	Ala	Ile	Cys	Pro	Phe	Pro	Ser	Glu	Ala	Leu	Ser	Pro	
	770					775					780					
Ser	Gln	Leu	Leu	Gly	Leu	Val	Arg	Ala	Gly	Val	His	Arg	Phe	Phe	Ala	
785				790						795					800	
Ser	Leu	Arg	Leu	His	Gly	Pro	Pro	Gly	Val	Ala	Ser	Ala	Cys	Gln	Leu	
			805						810					815		
Leu	Thr	Arg	Leu	Ser	Gln	Thr	Ser	Pro	Ala	Gly	Leu	Lys	Ala	Val	Leu	
		820						825					830			
Gln	Leu	Leu	Val	Glu	Gly	Ala	Leu	His	Arg	Gly	Asn	Thr	Glu	Leu	Phe	
		835					840					845				
Gly	Gly	Gln	Val	Asp	Gly	Asp	Asn	Glu	Thr	Leu	Ser	Val	Val	Ser	Ala	
	850					855					860					
Ser	Leu	Ala	Ser	Ala	Ser	Leu	Leu	Asp	Thr	Asn	Arg	Arg	His	Thr	Ala	
865					870					875					880	
Ala	Val	Pro	Gly	Pro	Gly	Gly	Ile	Trp	Ser	Val	Phe	His	Ala	Gly	Val	
			885					890						895		
Ile	Gly	Arg	Gly	Leu	Lys	Pro	Pro	Lys	Phe	Val	Gln	Ser	Arg	Asn	Gln	
		900						905					910			
Gln	Glu	Val	Ile	Tyr	Asn	Thr	Gln	Ser	Leu	Leu	Ser	Leu	Leu	Val	His	
		915					920					925				
Cys	Cys	Ser	Ala	Pro	Gly	Gly	Thr	Glu	Cys	Gly	Glu	Cys	Trp	Gly	Ala	
	930					935					940					
Pro	Ile	Leu	Ser	Pro	Glu	Ala	Ala	Lys	Ala	Val	Ala	Val	Thr	Leu	Val	
945					950					955					960	
Glu	Ser	Val	Cys	Pro	Asp	Ala	Ala	Gly	Ala	Glu	Leu	Ala	Trp	Pro	Pro	

2003295784 19 Aug 2005

	965	970	975
Glu Glu His	Ala Arg Ala Thr Val	Glu Arg Asp Leu Arg	Ile Gly Arg
	980	985	990
Arg Phe Arg	Glu Gln Pro Leu Leu	Phe Glu Leu Leu Lys	Leu Val Ala
	995	1000	1005
Ala Ala	Pro Pro Ala Leu Cys	Tyr Cys Ser Val	Leu Leu Arg Gly
	1010	1015	1020
Leu Leu	Ala Ala Leu Leu Gly	His Trp Glu Ala Ser	Arg His Pro
	1025	1030	1035
Asp Thr	Thr His Ser Pro Trp	His Leu Glu Ala Ser	Cys Thr Leu
	1040	1045	1050
Val Ala	Val Met Ala Glu Gly	Ser Leu Leu Pro Pro	Ala Leu Gly
	1055	1060	1065
Asn Met	His Glu Val Phe Ser	Gln Leu Ala Pro Phe	Glu Val Arg
	1070	1075	1080
Leu Leu	Leu Leu Ser Val Trp	Gly Phe Leu Arg Glu	His Gly Pro
	1085	1090	1095
Leu Pro	Gln Lys Phe Ile Phe	Gln Ser Glu Arg Gly	Arg Phe Ile
	1100	1105	1110
Arg Asp	Phe Ser Arg Glu Gly	Gly Gly Glu Gly Gly	Pro His Leu
	1115	1120	1125
Ala Val	Leu His Ser Val Leu	His Arg Asn Ile Asp	Arg Leu Gly
	1130	1135	1140
Leu Phe	Ser Gly Arg Phe Gln	Ala Pro Ser Pro Ser	Thr Leu Leu
	1145	1150	1155
Arg Gln	Gly Thr		
	1160		

<210> 9

<211> 1162

<212> PRT

<213> Artificial

<220>

<223> A variant of the human BFLP1698 polypeptide

<400> 9

Met Ala Leu Val	Pro Gly Arg Ser Lys	Glu Asp Gly Leu Trp Thr Arg
1	5	10 15
Asn Ser Pro Gly	Ser Ser Gln His Pro	Glu Ser Pro Arg Leu Pro Asn
	20	25 30
Pro Leu Trp Asp	Arg Gly Lys Ile Gly Lys Val	Glu Gly His Gln His
	35	40 45
Ile Gln Asp Phe	Ser Gln Lys Ser His Leu Pro	Ser Ile Val Val Glu

2003295784 19 Aug 2005

50	55	60
Ser 65	Ser 70	Glu 75
Glu 85	Leu 85	Thr 90
Phe 100	Gln 100	Asp 105
Ala 115	Arg 115	Gln 120
Glu 130	Leu 130	Lys 135
Arg 145	Arg 145	Leu 150
Ile 165	Thr 165	Arg 170
Glu 180	Ile 180	Lys 185
Leu 195	Ser 195	Ala 200
Leu 210	Pro 210	Ala 215
Asp 225	Glu 225	Ser 230
Ala 245	Gly 245	Pro 250
Gly 260	Gly 260	Pro 265
Ser 275	Glu 275	Phe 280
Ala 290	Trp 290	Ser 295
Gln 305	His 305	Gln 310
Gln 325	Leu 325	Trp 330
Val 340	Gln 340	Cys 345
Asp 355	Ala 355	Leu 360
Val 370	Val 370	Ala 375

2003295784 19 Aug 2005

Val	Leu	Ser	Cys	Gly	Leu	Lys	Asp	Phe	Cys	Val	His	Gly	Gly	Ala	Gly		
385					390					395					400		
Gly	Gly	Ala	Gly	Ser	Ser	Gly	Gly	Ser	Ser	Ser	Gln	Thr	Pro	Ser	Thr		
				405					410						415		
Asp	Pro	Phe	Pro	Gly	Ser	Pro	Ala	Ile	Pro	Ala	Glu	Lys	Arg	Val	Pro		
			420					425					430				
Lys	Ile	Ala	Ser	Val	Val	Gly	Ile	Leu	Gly	His	Leu	Ala	Ser	Arg	His		
		435					440					445					
Gly	Asp	Ser	Ile	Arg	Arg	Glu	Leu	Leu	Arg	Met	Phe	His	Asp	Ser	Leu		
	450					455					460						
Ala	Gly	Gly	Ser	Gly	Gly	Arg	Ser	Gly	Asp	Pro	Ser	Leu	Gln	Ala	Thr		
465					470					475					480		
Val	Pro	Phe	Leu	Leu	Gln	Leu	Ala	Val	Met	Ser	Pro	Ala	Leu	Leu	Gly		
			485					490						495			
Thr	Val	Ser	Gly	Glu	Leu	Val	Asp	Cys	Leu	Lys	Pro	Pro	Ala	Val	Leu		
			500					505					510				
Ser	Gln	Leu	Gln	Gln	His	Leu	Gln	Gly	Phe	Pro	Arg	Glu	Glu	Leu	Asp		
		515					520					525					
Asn	Met	Leu	Asn	Leu	Ala	Val	His	Leu	Val	Ser	Gln	Ala	Ser	Gly	Ala		
	530					535					540						
Gly	Ala	Tyr	Arg	Leu	Leu	Gln	Phe	Leu	Val	Asp	Thr	Ala	Met	Pro	Ala		
545					550					555					560		
Ser	Val	Ile	Thr	Thr	Gln	Gly	Leu	Ala	Val	Pro	Asp	Thr	Val	Arg	Glu		
				565					570					575			
Ala	Cys	Asp	Arg	Leu	Ile	Gln	Leu	Leu	Leu	His	Leu	Gln	Lys	Leu			
			580				585					590					
Val	His	His	Arg	Gly	Gly	Ser	Pro	Gly	Glu	Gly	Val	Leu	Gly	Pro	Pro		
		595					600					605					
Pro	Pro	Pro	Arg	Leu	Val	Pro	Phe	Leu	Asp	Ala	Leu	Lys	Asn	His	Val		
	610					615					620						
Gly	Glu	Leu	Cys	Gly	Glu	Thr	Leu	Arg	Leu	Glu	Arg	Lys	Arg	Phe	Leu		
625					630					635					640		
Trp	Gln	His	Gln	Leu	Leu	Gly	Leu	Leu	Ser	Val	Tyr	Thr	Arg	Pro	Ser		
				645					650					655			
Cys	Gly	Pro	Glu	Ala	Leu	Pro	His	Leu	Leu	Ser	Arg	Ala	Arg	Ser	Pro		
			660					665					670				
Glu	Glu	Leu	Ser	Leu	Ala	Thr	Gln	Leu	Tyr	Ala	Gly	Leu	Val	Val	Ser		
		675					680					685					
Leu	Ser	Gly	Leu	Leu	Pro	Leu	Ala	Phe	Arg	Ser	Cys	Leu	Ala	Arg	Val		
	690					695					700						
His	Ala	Gly	Thr	Leu	Gln	Pro	Pro	Phe	Thr	Ala	Arg	Phe	Leu	Arg	Asn		
705					710					715					720		

2003295784 19 Aug 2005

Leu Ala Leu Leu Val Gly Trp Glu Gln Gln Gly Gly Glu Gly Pro Ala
 725 730 735
 Ala Leu Gly Ala His Phe Gly Glu Ser Ala Ser Ala His Leu Ser Asp
 740 745 750
 Leu Ala Pro Leu Leu Leu His Pro Glu Glu Glu Val Ala Glu Ala Ala
 755 760 765
 Ala Ser Leu Leu Ala Ile Cys Pro Phe Pro Ser Glu Ala Leu Ser Pro
 770 775 780
 Ser Gln Leu Leu Gly Leu Val Arg Ala Gly Val His Arg Phe Phe Ala
 785 790 795 800
 Ser Leu Arg Leu His Gly Pro Pro Gly Val Ala Ser Ala Cys Gln Leu
 805 810 815
 Leu Thr Arg Leu Ser Gln Thr Ser Pro Ala Gly Leu Lys Ala Val Leu
 820 825 830
 Gln Leu Leu Val Glu Gly Ala Leu His Arg Gly Asn Thr Glu Leu Phe
 835 840 845
 Gly Gly Gln Val Asp Gly Asp Asn Glu Thr Leu Ser Val Val Ser Ala
 850 855 860
 Ser Leu Ala Ser Ala Ser Leu Leu Asp Thr Asn Arg Arg His Thr Ala
 865 870 875 880
 Ala Val Pro Gly Pro Gly Gly Ile Trp Ser Val Phe His Ala Gly Val
 885 890 895
 Ile Gly Arg Gly Leu Lys Pro Pro Lys Phe Val Gln Ser Arg Asn Gln
 900 905 910
 Gln Glu Val Ile Tyr Asn Thr Gln Ser Leu Leu Ser Leu Leu Val His
 915 920 925
 Cys Cys Ser Ala Pro Gly Gly Thr Glu Cys Gly Glu Cys Trp Gly Ala
 930 935 940
 Pro Ile Leu Ser Pro Glu Ala Ala Lys Ala Val Ala Val Thr Leu Val
 945 950 955 960
 Glu Ser Val Cys Pro Asp Ala Ala Gly Ala Glu Leu Ala Trp Pro Pro
 965 970 975
 Glu Glu His Ala Arg Ala Thr Val Glu Arg Asp Leu Arg Ile Gly Arg
 980 985 990
 Arg Phe Arg Glu Gln Pro Leu Leu Phe Glu Leu Leu Lys Leu Val Ala
 995 1000 1005
 Ala Ala Pro Pro Ala Leu Cys Tyr Cys Ser Val Leu Leu Arg Gly
 1010 1015 1020
 Leu Leu Ala Ala Leu Leu Gly His Trp Glu Ala Ser Arg His Pro
 1025 1030 1035
 Asp Thr Thr His Ser Pro Trp His Leu Glu Ala Ser Cys Thr Leu

2003295784 19 Aug 2005

1040	1045	1050
Val Ala Val Met Ala Glu Gly Ser Leu Leu Pro Pro Ala Leu Gly		
1055	1060	1065
Asn Met His Glu Val Phe Ser Gln Leu Ala Pro Phe Glu Val Arg		
1070	1075	1080
Leu Leu Leu Leu Ser Val Trp Gly Phe Leu Arg Glu His Gly Pro		
1085	1090	1095
Leu Pro Gln Lys Phe Ile Phe Gln Ser Glu Arg Gly Arg Phe Ile		
1100	1105	1110
Arg Asp Phe Ser Arg Glu Gly Gly Gly Glu Gly Gly Pro His Leu		
1115	1120	1125
Ala Val Leu His Ser Val Leu His Arg Asn Ile Asp Arg Leu Gly		
1130	1135	1140
Leu Phe Ser Gly Arg Phe Gln Ala Pro Ser Pro Ser Thr Leu Leu		
1145	1150	1155
Arg Gln Gly Thr		
1160		

<210> 10
 <211> 1162
 <212> PRT
 <213> Artificial

<220>
 <223> A variant of the human BFLP1698 polypeptide

<400> 10

Met Ala Leu Val Pro Gly Arg Ser Lys Glu Asp Gly Leu Trp Thr Arg		
1	5	10 15
Asn Ser Pro Gly Ser Ser Gln His Pro Glu Ser Pro Arg Leu Pro Asn		
20	25	30
Pro Leu Trp Asp Arg Gly Lys Ile Gly Lys Val Glu Gly His Gln His		
35	40	45
Ile Gln Asp Phe Ser Gln Lys Ser His Leu Pro Ser Ile Val Val Glu		
50	55	60
Ser Ser Glu Val Asn Glu Glu Ser Gly Asp Leu His Leu Pro His Glu		
65	70	75 80
Glu Leu Leu Leu Leu Thr Asp Gly Glu Glu Glu Asp Ala Glu Ala Phe		
85	90	95
Phe Gln Asp Gln Ser Glu Glu Pro Gly Ala Ala Arg Pro His His Gln		
100	105	110
Ala Arg Gln Val Glu His Ser Thr Gln Arg Gly His Leu Glu Ile Arg		
115	120	125
Glu Leu Lys Lys Lys Leu Phe Lys Arg Arg Arg Val Leu Asn Arg Glu		

2003295784 19 Aug 2005

130	135	140
Arg Arg Leu Arg His Arg Val Val Gly Ala Val Ile Asp Gln Gly Leu		
145	150	155
Ile Thr Arg His His Leu Lys Lys Arg Ala Ala Gln Glu Leu Ser Gln		
	165	170
Glu Ile Lys Ala Phe Leu Thr Gly Val Asp Pro Ile Leu Gly His Gln		
	180	185
Leu Ser Ala Arg Glu His Ala Arg Cys Gly Leu Leu Leu Leu Arg Ser		
	195	200
Leu Pro Pro Ala Arg Ala Ala Val Leu Asp His Leu Arg Gly Val Phe		
	210	215
Asp Glu Ser Val Arg Ala His Leu Ala Ala Leu Asp Glu Thr Pro Val		
	225	230
Ala Gly Pro Pro His Leu Arg Pro Pro Pro Pro Ser His Val Pro Ala		
	245	250
Gly Gly Pro Gly Leu Glu Asp Val Val Gln Glu Val Gln Gln Val Leu		
	260	265
Ser Glu Phe Ile Arg Ala Asn Pro Lys Ala Trp Ala Pro Val Ile Ser		
	275	280
Ala Trp Ser Ile Asp Leu Met Gly Gln Leu Ser Ser Thr Tyr Ser Gly		
	290	295
Gln His Gln Arg Val Pro His Ala Thr Gly Ala Leu Asn Glu Leu Leu		
	305	310
Gln Leu Trp Met Gly Cys Arg Ala Thr Arg Thr Leu Met Asp Ile Tyr		
	325	330
Val Gln Cys Leu Ser Ala Leu Ile Gly Ser Cys Pro Asp Ala Cys Val		
	340	345
Asp Ala Leu Leu Asp Thr Ser Val Gln His Ser Pro His Phe Asp Trp		
	355	360
Val Val Ala His Ile Gly Ser Ser Phe Pro Gly Thr Ile Ile Ser Arg		
	370	375
Val Leu Ser Cys Gly Leu Lys Asp Phe Cys Val His Gly Gly Ala Gly		
	385	390
Gly Gly Ala Gly Ser Ser Gly Gly Ser Ser Ser Gln Thr Pro Ser Thr		
	405	410
Asp Pro Phe Pro Gly Ser Pro Ala Ile Pro Ala Glu Lys Arg Val Pro		
	420	425
Lys Ile Ala Ser Val Val Gly Ile Leu Gly His Leu Ala Ser Arg His		
	435	440
Gly Asp Ser Ile Arg Arg Glu Leu Leu Arg Met Phe His Asp Ser Leu		
	450	455
		460

2003295784 19 Aug 2005

Ala	Gly	Gly	Ser	Gly	Gly	Arg	Ser	Gly	Asp	Pro	Ser	Leu	Gln	Ala	Thr	465	470	475	480
Val	Pro	Phe	Leu	Leu	Gln	Leu	Ala	Val	Met	Ser	Pro	Ala	Leu	Leu	Gly	485	490		495
Thr	Val	Ser	Gly	Glu	Leu	Val	Asp	Cys	Leu	Lys	Pro	Pro	Ala	Val	Leu	500	505		510
Ser	Gln	Leu	Gln	Gln	His	Leu	Gln	Gly	Phe	Pro	Arg	Glu	Glu	Leu	Asp	515	520		525
Asn	Met	Leu	Asn	Leu	Ala	Val	His	Leu	Val	Ser	Gln	Ala	Ser	Gly	Ala	530	535		540
Gly	Ala	Tyr	Arg	Leu	Leu	Gln	Phe	Leu	Val	Asp	Thr	Ala	Met	Pro	Ala	545	550	555	560
Ser	Val	Ile	Thr	Thr	Gln	Gly	Leu	Ala	Val	Pro	Asp	Thr	Val	Arg	Glu	565	570		575
Ala	Cys	Asp	Arg	Leu	Ile	Gln	Leu	Leu	Leu	His	Leu	Gln	Lys	Leu		580	585		590
Val	His	His	Arg	Gly	Gly	Ser	Pro	Gly	Glu	Gly	Val	Leu	Gly	Pro	Pro	595	600		605
Pro	Pro	Pro	Arg	Leu	Val	Pro	Phe	Leu	Asp	Ala	Leu	Lys	Asn	His	Val	610	615		620
Gly	Glu	Leu	Cys	Gly	Glu	Thr	Leu	Arg	Leu	Glu	Arg	Lys	Arg	Phe	Leu	625	630	635	640
Trp	Gln	His	Gln	Leu	Leu	Gly	Leu	Leu	Ser	Val	Tyr	Thr	Arg	Pro	Ser	645	650		655
Cys	Gly	Pro	Glu	Ala	Leu	Gly	His	Leu	Leu	Ser	Arg	Ala	Arg	Ser	Pro	660	665		670
Glu	Glu	Leu	Ser	Leu	Ala	Thr	Gln	Leu	Tyr	Ala	Gly	Leu	Val	Val	Ser	675	680	685	
Leu	Ser	Gly	Leu	Leu	Pro	Leu	Ala	Phe	Arg	Ser	Cys	Leu	Ala	Arg	Val	690	695	700	
His	Ala	Gly	Thr	Leu	Gln	Pro	Pro	Phe	Thr	Ala	Arg	Phe	Leu	Arg	Asn	705	710	715	720
Leu	Ala	Leu	Leu	Val	Gly	Trp	Glu	Gln	Gln	Gly	Gly	Asp	Gly	Pro	Ala	725	730		735
Ala	Leu	Gly	Ala	His	Phe	Gly	Glu	Ser	Ala	Ser	Ala	His	Leu	Ser	Asp	740	745		750
Leu	Ala	Pro	Leu	Leu	Leu	His	Pro	Glu	Glu	Glu	Val	Ala	Glu	Ala	Ala	755	760	765	
Ala	Ser	Leu	Leu	Ala	Ile	Cys	Pro	Phe	Pro	Ser	Glu	Ala	Leu	Ser	Pro	770	775	780	
Ser	Gln	Leu	Leu	Gly	Leu	Val	Arg	Ala	Gly	Val	His	Arg	Phe	Phe	Ala	785	790	795	800

2003295784 19 Aug 2005

Ser Leu Arg Leu His Gly Pro Pro Gly Val Ala Ser Ala Cys Gln Leu
 805 810 815
 Leu Thr Arg Leu Ser Gln Thr Ser Pro Ala Gly Leu Lys Ala Val Leu
 820 825 830
 Gln Leu Leu Val Glu Gly Ala Leu His Arg Gly Asn Thr Glu Leu Phe
 835 840 845
 Gly Gly Gln Val Asp Gly Asp Asn Glu Thr Leu Ser Val Val Ser Ala
 850 855 860
 Ser Leu Ala Ser Ala Ser Leu Leu Asp Thr Asn Arg Arg His Thr Ala
 865 870 875 880
 Ala Val Pro Gly Pro Gly Gly Ile Trp Ser Val Phe His Ala Gly Val
 885 890 895
 Ile Gly Arg Gly Leu Lys Pro Pro Lys Phe Val Gln Ser Arg Asn Gln
 900 905 910
 Gln Glu Val Ile Tyr Asn Thr Gln Ser Leu Leu Ser Leu Leu Val His
 915 920 925
 Cys Cys Ser Ala Pro Gly Gly Thr Glu Cys Gly Glu Cys Trp Gly Ala
 930 935 940
 Pro Ile Leu Ser Pro Glu Ala Ala Lys Ala Val Ala Val Thr Leu Val
 945 950 955 960
 Glu Ser Val Cys Pro Asp Ala Ala Gly Ala Glu Leu Ala Trp Pro Pro
 965 970 975
 Glu Glu His Ala Arg Ala Thr Val Glu Arg Asp Leu Arg Ile Gly Arg
 980 985 990
 Arg Phe Arg Glu Gln Pro Leu Leu Phe Glu Leu Leu Lys Leu Val Ala
 995 1000 1005
 Ala Ala Pro Pro Ala Leu Cys Tyr Cys Ser Val Leu Leu Arg Gly
 1010 1015 1020
 Leu Leu Ala Ala Leu Leu Gly His Trp Glu Ala Ser Arg His Pro
 1025 1030 1035
 Asp Thr Thr His Ser Pro Trp His Leu Glu Ala Ser Cys Thr Leu
 1040 1045 1050
 Val Ala Val Met Ala Glu Gly Ser Leu Leu Pro Pro Ala Leu Gly
 1055 1060 1065
 Asn Met His Glu Val Phe Ser Gln Leu Ala Pro Phe Glu Val Arg
 1070 1075 1080
 Leu Leu Leu Leu Ser Val Trp Gly Phe Leu Arg Glu His Gly Pro
 1085 1090 1095
 Leu Pro Gln Lys Phe Ile Phe Gln Ser Glu Arg Gly Arg Phe Ile
 1100 1105 1110
 Arg Asp Phe Ser Arg Glu Gly Gly Gly Glu Gly Gly Pro His Leu

2003295784 19 Aug 2005

1115	1120	1125
Ala Val Leu His Ser Val	Leu His Arg Asn Ile Asp	Arg Leu Gly
1130	1135	1140
Leu Phe Ser Gly Arg Phe	Gln Ala Pro Ser Pro Ser	Thr Leu Leu
1145	1150	1155
Arg Gln Gly Thr		
1160		

<210> 11
 <211> 1162
 <212> PRT
 <213> Artificial

 <220>
 <223> A variant of the human BFLP1698 polypeptide

 <400> 11

Met	Ala	Leu	Val	Pro	Gly	Arg	Ser	Lys	Glu	Asp	Gly	Leu	Trp	Thr	Arg
1				5					10					15	
Asn	Ser	Pro	Gly	Ser	Ser	Gln	His	Pro	Glu	Ser	Pro	Arg	Leu	Pro	Asn
			20					25					30		
Pro	Leu	Trp	Asp	Arg	Gly	Lys	Ile	Gly	Lys	Val	Glu	Gly	His	Gln	His
		35					40					45			
Ile	Gln	Asp	Phe	Ser	Gln	Lys	Ser	His	Leu	Pro	Ser	Ile	Val	Val	Glu
	50					55					60				
Ser	Ser	Glu	Val	Asn	Glu	Glu	Ser	Gly	Asp	Leu	His	Leu	Pro	His	Glu
65					70				75					80	
Glu	Leu	Leu	Leu	Leu	Thr	Asp	Gly	Glu	Glu	Glu	Asp	Ala	Glu	Ala	Phe
				85				90						95	
Phe	Gln	Asp	Gln	Ser	Glu	Glu	Pro	Gly	Ala	Ala	Arg	Pro	His	His	Gln
			100					105					110		
Ala	Arg	Gln	Val	Glu	His	Ser	Thr	Gln	Arg	Gly	His	Leu	Glu	Ile	Arg
			115					120					125		
Glu	Leu	Lys	Lys	Lys	Leu	Phe	Lys	Arg	Arg	Arg	Val	Leu	Asn	Arg	Glu
	130					135					140				
Arg	Arg	Leu	Arg	His	Arg	Val	Val	Gly	Ala	Val	Ile	Asp	Gln	Gly	Leu
145					150					155					160
Ile	Thr	Arg	His	His	Leu	Lys	Lys	Arg	Ala	Ala	Gln	Glu	Leu	Ser	Gln
				165					170					175	
Glu	Ile	Lys	Ala	Phe	Leu	Thr	Gly	Val	Asp	Pro	Ile	Leu	Gly	His	Gln
			180					185					190		
Leu	Ser	Ala	Arg	Glu	His	Ala	Arg	Cys	Gly	Leu	Leu	Leu	Leu	Arg	Ser
		195					200					205			
Leu	Pro	Pro	Ala	Arg	Ala	Ala	Val	Leu	Asp	His	Leu	Arg	Gly	Val	Phe

2003295784 19 Aug 2005

210	215	220
Asp Glu Ser Val Arg	Ala His Leu Ala Ala	Leu Asp Glu Thr Pro Val
225	230	235 240
Ala Gly Pro Pro His	Leu Arg Pro Pro Pro	Ser His Val Pro Ala
	245	250 255
Gly Gly Pro Gly Leu	Glu Asp Val Val Gln	Glu Val Gln Gln Val Leu
	260	265 270
Ser Glu Phe Ile Arg	Ala Asn Pro Lys Ala Trp	Ala Pro Val Ile Ser
	275	280 285
Ala Trp Ser Ile Asp	Leu Met Gly Gln Leu Ser	Ser Thr Tyr Ser Gly
	290	295 300
Gln His Gln Arg Val	Pro His Ala Thr Gly	Ala Leu Asn Glu Leu Leu
	305	310 315 320
Gln Leu Trp Met Gly	Cys Arg Ala Thr Arg	Thr Leu Met Asp Ile Tyr
	325	330 335
Val Gln Cys Leu Ser	Ala Leu Ile Gly Ser	Cys Pro Asp Ala Cys Val
	340	345 350
Asp Ala Leu Leu Asp	Thr Ser Val Gln His Ser	Pro His Phe Asp Trp
	355	360 365
Val Val Ala His Ile	Gly Ser Ser Phe Pro Gly	Thr Ile Ile Ser Arg
	370	375 380
Val Leu Ser Cys Gly	Leu Lys Asp Phe Cys	Val His Gly Gly Ala Gly
	385	390 395 400
Gly Gly Ala Gly Ser	Ser Gly Gly Ser Ser	Ser Gln Thr Pro Ser Thr
	405	410 415
Asp Pro Phe Pro Gly	Ser Pro Ala Ile Pro	Ala Glu Lys Arg Val Pro
	420	425 430
Lys Ile Ala Ser Val	Val Gly Ile Leu Gly	His Leu Ala Ser Arg His
	435	440 445
Gly Asp Ser Ile Arg	Arg Glu Leu Leu Arg	Met Phe His Asp Ser Leu
	450	455 460
Ala Gly Gly Ser Gly	Gly Arg Ser Gly Asp	Pro Ser Leu Gln Ala Thr
	465	470 475 480
Val Pro Phe Leu Leu	Gln Leu Ala Val Met	Ser Pro Ala Leu Leu Gly
	485	490 495
Thr Val Ser Gly Glu	Leu Val Asp Cys Leu	Lys Pro Pro Ala Val Leu
	500	505 510
Ser Gln Leu Gln Gln	His Leu Gln Gly Phe	Pro Arg Glu Glu Leu Asp
	515	520 525
Asn Met Leu Asn Leu	Ala Val His Leu Val	Ser Gln Ala Ser Gly Ala
	530	535 540

2003295784 19 Aug 2005

Gly	Ala	Tyr	Arg	Leu	Leu	Gln	Phe	Leu	Val	Asp	Thr	Ala	Met	Pro	Ala	545	550	555	560
Ser	Val	Ile	Thr	Thr	Gln	Gly	Leu	Ala	Val	Pro	Asp	Thr	Val	Arg	Glu	565	570	575	
Ala	Cys	Asp	Arg	Leu	Ile	Gln	Leu	Leu	Leu	His	Leu	Gln	Lys	Leu		580	585	590	
Val	His	His	Arg	Gly	Gly	Ser	Pro	Gly	Glu	Gly	Val	Leu	Gly	Pro	Pro	595	600	605	
Pro	Pro	Pro	Arg	Leu	Val	Pro	Phe	Leu	Asp	Ala	Leu	Lys	Asn	His	Val	610	615	620	
Gly	Glu	Leu	Cys	Gly	Glu	Thr	Leu	Arg	Leu	Glu	Arg	Lys	Arg	Phe	Leu	625	630	635	640
Trp	Gln	His	Gln	Leu	Leu	Gly	Leu	Leu	Ser	Val	Tyr	Thr	Arg	Pro	Ser	645	650	655	
Cys	Gly	Pro	Glu	Ala	Leu	Gly	His	Leu	Leu	Ser	Arg	Ala	Arg	Ser	Pro	660	665	670	
Glu	Glu	Leu	Ser	Leu	Ala	Thr	Gln	Leu	Tyr	Ala	Gly	Leu	Val	Val	Ser	675	680	685	
Leu	Ser	Gly	Leu	Leu	Pro	Leu	Ala	Phe	Arg	Ser	Cys	Leu	Ala	Arg	Val	690	695	700	
His	Ala	Gly	Thr	Leu	Gln	Pro	Pro	Phe	Thr	Ala	Arg	Phe	Leu	Arg	Asn	705	710	715	720
Leu	Ala	Leu	Leu	Val	Gly	Trp	Glu	Gln	Gln	Gly	Gly	Glu	Gly	Pro	Ala	725	730	735	
Ala	Leu	Gly	Ala	His	Phe	Gly	Glu	Ser	Ala	Ser	Ala	His	Leu	Ser	Asp	740	745	750	
Leu	Ala	Pro	Leu	Leu	Leu	His	Pro	Glu	Glu	Glu	Val	Ala	Glu	Ala	Ala	755	760	765	
Ala	Ser	Leu	Leu	Ala	Ile	Cys	Pro	Phe	Pro	Ser	Glu	Ala	Leu	Ser	Pro	770	775	780	
Ser	Gln	Leu	Leu	Gly	Leu	Val	Arg	Ala	Gly	Val	His	Arg	Phe	Phe	Ala	785	790	795	800
Ser	Leu	Arg	Leu	His	Gly	Pro	Pro	Gly	Val	Ala	Ser	Ala	Cys	Gln	Leu	805	810	815	
Leu	Thr	Arg	Leu	Ser	Gln	Thr	Ser	Pro	Ala	Gly	Leu	Lys	Ala	Val	Leu	820	825	830	
Gln	Leu	Leu	Val	Glu	Gly	Ala	Leu	His	Arg	Gly	Asn	Thr	Glu	Leu	Phe	835	840	845	
Gly	Gly	Gln	Val	Asp	Gly	Asp	Asn	Glu	Ala	Leu	Ser	Val	Val	Ser	Ala	850	855	860	
Ser	Leu	Ala	Ser	Ala	Ser	Leu	Leu	Asp	Thr	Asn	Arg	Arg	His	Thr	Ala	865	870	875	880

2003295784 19 Aug 2005

Ala Val Pro Gly Pro Gly Gly Ile Trp Ser Val Phe His Ala Gly Val
885 890 895

Ile Gly Arg Gly Leu Lys Pro Pro Lys Phe Val Gln Ser Arg Asn Gln
900 905 910

Gln Glu Val Ile Tyr Asn Thr Gln Ser Leu Leu Ser Leu Leu Val His
915 920 925

Cys Cys Ser Ala Pro Gly Gly Thr Glu Cys Gly Glu Cys Trp Gly Ala
930 935 940

Pro Ile Leu Ser Pro Glu Ala Ala Lys Ala Val Ala Val Thr Leu Val
945 950 955 960

Glu Ser Val Cys Pro Asp Ala Ala Gly Ala Glu Leu Ala Trp Pro Pro
965 970 975

Glu Glu His Ala Arg Ala Thr Val Glu Arg Asp Leu Arg Ile Gly Arg
980 985 990

Arg Phe Arg Glu Gln Pro Leu Leu Phe Glu Leu Leu Lys Leu Val Ala
995 1000 1005

Ala Ala Pro Pro Ala Leu Cys Tyr Cys Ser Val Leu Leu Arg Gly
1010 1015 1020

Leu Leu Ala Ala Leu Leu Gly His Trp Glu Ala Ser Arg His Pro
1025 1030 1035

Asp Thr Thr His Ser Pro Trp His Leu Glu Ala Ser Cys Thr Leu
1040 1045 1050

Val Ala Val Met Ala Glu Gly Ser Leu Leu Pro Pro Ala Leu Gly
1055 1060 1065

Asn Met His Glu Val Phe Ser Gln Leu Ala Pro Phe Glu Val Arg
1070 1075 1080

Leu Leu Leu Leu Ser Val Trp Gly Phe Leu Arg Glu His Gly Pro
1085 1090 1095

Leu Pro Gln Lys Phe Ile Phe Gln Ser Glu Arg Gly Arg Phe Ile
1100 1105 1110

Arg Asp Phe Ser Arg Glu Gly Gly Gly Glu Gly Gly Pro His Leu
1115 1120 1125

Ala Val Leu His Ser Val Leu His Arg Asn Ile Asp Arg Leu Gly
1130 1135 1140

Leu Phe Ser Gly Arg Phe Gln Ala Pro Ser Pro Ser Thr Leu Leu
1145 1150 1155

Arg Gln Gly Thr
1160

<210> 12

<211> 1162

<212> PRT

2003295784 19 Aug 2005

<213> Artificial

<220>

<223> A variant of the human BFLP1698 polypeptide

<400> 12

Met Ala Leu Val Pro Gly Arg Ser Lys Glu Asp Gly Leu Trp Thr Arg
1 5 10 15

Asn Ser Pro Gly Ser Ser Gln His Pro Glu Ser Pro Arg Leu Pro Asn
20 25 30

Pro Leu Trp Asp Arg Gly Lys Ile Gly Lys Val Glu Gly His Gln His
35 40 45

Ile Gln Asp Phe Ser Gln Lys Ser His Leu Pro Ser Ile Val Val Glu
50 55 60

Ser Ser Glu Val Asn Glu Glu Ser Gly Asp Leu His Leu Pro His Glu
65 70 75 80

Glu Leu Leu Leu Leu Thr Asp Gly Glu Glu Glu Asp Ala Glu Ala Phe
85 90 95

Phe Gln Asp Gln Ser Glu Glu Pro Gly Ala Ala Arg Pro His His Gln
100 105 110

Ala Arg Gln Val Glu His Ser Thr Gln Arg Gly His Leu Glu Ile Arg
115 120 125

Glu Leu Lys Lys Lys Leu Phe Lys Arg Arg Arg Val Leu Asn Arg Glu
130 135 140

Arg Arg Leu Arg His Arg Val Val Gly Ala Val Ile Asp Gln Gly Leu
145 150 155 160

Ile Thr Arg His His Leu Lys Lys Arg Ala Ala Gln Glu Leu Ser Gln
165 170 175

Glu Ile Lys Ala Phe Leu Thr Gly Val Asp Pro Ile Leu Gly His Gln
180 185 190

Leu Ser Ala Arg Glu His Ala Arg Cys Gly Leu Leu Leu Leu Arg Ser
195 200 205

Leu Pro Pro Ala Arg Ala Ala Val Leu Asp His Leu Arg Gly Val Phe
210 215 220

Asp Glu Ser Val Arg Ala His Leu Ala Ala Leu Asp Glu Thr Pro Val
225 230 235 240

Ala Gly Pro Pro His Leu Arg Pro Pro Pro Pro Ser His Val Pro Ala
245 250 255

Gly Gly Pro Gly Leu Glu Asp Val Val Gln Glu Val Gln Gln Val Leu
260 265 270

Ser Glu Phe Ile Arg Ala Asn Pro Lys Ala Trp Ala Pro Val Ile Ser
275 280 285

Ala Trp Ser Ile Asp Leu Met Gly Gln Leu Ser Ser Thr Tyr Ser Gly

2003295784 19 Aug 2005

290	295	300
Gln His Gln Arg Val 305	Pro His Ala Thr Gly 310	Ala Leu Asn Glu Leu Leu 315 320
Gln Leu Trp Met Gly 325	Cys Arg Ala Thr Arg 330	Thr Leu Met Asp Ile Tyr 335
Val Gln Cys Leu Ser 340	Ala Leu Ile Gly 345	Ser Cys Pro Asp Ala Cys Val 350
Asp Ala Leu Leu Asp 355	Thr Ser Val Gln His 360	Ser Pro His Phe Asp Trp 365
Val Val Ala His Ile 370	Gly Ser Ser Phe Pro 375	Gly Thr Ile Ile Ser Arg 380
Val Leu Ser Cys Gly 385	Leu Lys Asp Phe Cys 390	Val His Gly Gly Ala Gly 395 400
Gly Gly Ala Gly Ser 405	Ser Gly Gly Ser 410	Ser Gln Thr Pro Ser Thr 415
Asp Pro Phe Pro Gly 420	Ser Pro Ala Ile 425	Pro Ala Glu Lys Arg Val Pro 430
Lys Ile Ala Ser Val 435	Val Gly Ile Leu 440	Gly His Leu Ala Ser Arg His 445
Gly Asp Ser Ile Arg 450	Arg Glu Leu Leu Arg 455	Met Phe His Asp Ser Leu 460
Ala Gly Gly Ser Gly 465	Gly Arg Ser Gly Asp 470	Pro Ser Leu Gln Ala Thr 475 480
Val Pro Phe Leu Leu 485	Gln Leu Ala Val 490	Met Ser Pro Ala Leu Leu Gly 495
Thr Val Ser Gly Glu 500	Leu Val Asp Cys 505	Leu Lys Pro Pro Ala Val Leu 510
Ser Gln Leu Gln Gln 515	His Leu Gln Gly 520	Phe Pro Arg Glu Glu Leu Asp 525
Asn Met Leu Asn Leu 530	Ala Val His Leu 535	Val Ser Gln Ala Ser Gly Ala 540
Gly Ala Tyr Arg Leu 545	Leu Gln Phe Leu 550	Val Asp Thr Ala Met Pro Ala 555 560
Ser Val Ile Thr Thr 565	Gln Gly Leu Ala 570	Val Pro Asp Thr Val Arg Glu 575
Ala Cys Asp Arg Leu 580	Ile Gln Leu Leu 585	Leu Leu His Leu Gln Lys Leu 590
Val His His Arg Gly 595	Gly Ser Pro Gly 600	Glu Gly Val Leu Gly Pro Pro 605
Pro Pro Pro Arg Leu 610	Val Pro Phe Leu 615	Asp Ala Leu Lys Asn His Val 620

2003295784 19 Aug 2005

Gly 625	Glu	Leu	Cys	Gly 630	Glu	Thr	Leu	Arg	Leu	Glu 635	Arg	Lys	Arg	Phe	Leu 640
Trp	Gln	His	Gln 645	Leu	Leu	Gly	Leu	Leu	Ser 650	Val	Tyr	Thr	Arg	Pro	Ser 655
Cys	Gly	Pro	Glu 660	Ala	Leu	Gly	His	Leu	Leu	Ser	Arg	Ala	Arg	Ser	Pro 670
Glu	Glu	Leu	Ser 675	Leu	Ala	Thr	Gln	Leu	Tyr	Ala	Gly	Leu	Val	Val	Ser 685
Leu	Ser	Gly	Leu	Leu	Pro	Leu	Ala	Phe	Arg	Ser	Cys	Leu	Ala	Arg	Val 700
His 705	Ala	Gly	Thr	Leu	Gln 710	Pro	Pro	Phe	Thr	Ala	Arg	Phe	Leu	Arg	Asn 720
Leu	Ala	Leu	Leu	Val 725	Gly	Trp	Glu	Gln	Gln	Gly	Gly	Glu	Gly	Pro	Ala 735
Ala	Leu	Gly	Ala	His	Phe	Gly	Glu	Ser	Ala	Ser	Ala	His	Leu	Ser	Asp 750
Leu	Ala	Pro	Leu	Leu	Leu	His	Pro	Glu	Glu	Glu	Val	Ala	Glu	Ala	Ala 765
Ala	Ser	Leu	Leu	Ala	Ile	Cys	Pro	Phe	Pro	Ser	Glu	Ala	Leu	Ser	Pro 780
Ser 785	Gln	Leu	Leu	Gly	Leu	Val	Arg	Ala	Gly	Val	His	Arg	Phe	Phe	Ala 800
Ser	Leu	Arg	Leu	His	Gly	Pro	Pro	Gly	Val	Ala	Ser	Ala	Cys	Gln	Leu 815
Leu	Thr	Arg	Leu	Ser	Gln	Thr	Ser	Pro	Ala	Gly	Leu	Lys	Ala	Val	Leu 830
Gln	Leu	Leu	Val	Glu	Gly	Ala	Leu	His	Arg	Gly	Asn	Thr	Glu	Leu	Phe 845
Gly 850	Gly	Gln	Val	Asp	Gly	Asp	Asn	Glu	Thr	Leu	Ser	Val	Val	Ser	Ala 860
Ser 865	Leu	Ala	Ser	Ala	Ser	Leu	Leu	Asp	Thr	Asn	Arg	Arg	His	Thr	Ala 880
Ala	Val	Pro	Gly	Pro	Gly	Gly	Ile	Trp	Ser	Val	Phe	His	Ala	Gly	Val 895
Ile	Gly	Arg	Gly	Leu	Lys	Pro	Pro	Lys	Phe	Val	Gln	Ser	Arg	Asn	Gln 910
Gln	Glu	Val	Ile	Tyr	Asn	Thr	Gln	Ser	Leu	Leu	Ser	Leu	Leu	Val	His 925
Cys	Cys	Ser	Ala	Pro	Gly	Gly	Thr	Glu	Cys	Gly	Glu	Cys	Trp	Gly	Ala 940
Pro 945	Ile	Leu	Ser	Pro	Glu	Ala	Ala	Lys	Ala	Val	Ala	Val	Thr	Leu	Val 960

2003295784 19 Aug 2005

Glu Ser Val Cys Pro Asp Ala Ala Gly Ala Glu Leu Ala His Pro Pro
 965 970 975
 Glu Glu His Ala Arg Ala Thr Val Glu Arg Asp Leu Arg Ile Gly Arg
 980 985 990
 Arg Phe Arg Glu Gln Pro Leu Leu Phe Glu Leu Leu Lys Leu Val Ala
 995 1000 1005
 Ala Ala Pro Pro Ala Leu Cys Tyr Cys Ser Val Leu Leu Arg Gly
 1010 1015 1020
 Leu Leu Ala Ala Leu Leu Gly His Trp Glu Ala Ser Arg His Pro
 1025 1030 1035
 Asp Thr Thr His Ser Pro Trp His Leu Glu Ala Ser Cys Thr Leu
 1040 1045 1050
 Val Ala Val Met Ala Glu Gly Ser Leu Leu Pro Pro Ala Leu Gly
 1055 1060 1065
 Asn Met His Glu Val Phe Ser Gln Leu Ala Pro Phe Glu Val Arg
 1070 1075 1080
 Leu Leu Leu Leu Ser Val Trp Gly Phe Leu Arg Glu His Gly Pro
 1085 1090 1095
 Leu Pro Gln Lys Phe Ile Phe Gln Ser Glu Arg Gly Arg Phe Ile
 1100 1105 1110
 Arg Asp Phe Ser Arg Glu Gly Gly Gly Glu Gly Gly Pro His Leu
 1115 1120 1125
 Ala Val Leu His Ser Val Leu His Arg Asn Ile Asp Arg Leu Gly
 1130 1135 1140
 Leu Phe Ser Gly Arg Phe Gln Ala Pro Ser Pro Ser Thr Leu Leu
 1145 1150 1155
 Arg Gln Gly Thr
 1160

<210> 13
 <211> 1162
 <212> PRT
 <213> Artificial

<220>

<223> A variant of the human BFLP1698 polypeptide

<400> 13

Met Ala Leu Val Pro Gly Arg Ser Lys Glu Asp Gly Leu Trp Thr Arg
 1 5 10 15
 Asn Ser Pro Gly Ser Ser Gln His Pro Glu Ser Pro Arg Leu Pro Asn
 20 25 30
 Pro Leu Trp Asp Arg Gly Lys Ile Gly Lys Val Glu Gly His Gln His
 35 40 45

2003295784 19 Aug 2005

Ile	Gln	Asp	Phe	Ser	Gln	Lys	Ser	His	Leu	Pro	Ser	Ile	Val	Val	Glu	
50						55					60					
Ser	Ser	Glu	Val	Asn	Glu	Glu	Ser	Gly	Asp	Leu	His	Leu	Pro	His	Glu	
65					70					75					80	
Glu	Leu	Leu	Leu	Leu	Thr	Asp	Gly	Glu	Glu	Glu	Asp	Ala	Glu	Ala	Phe	
				85					90					95		
Phe	Gln	Asp	Gln	Ser	Glu	Glu	Pro	Gly	Ala	Ala	Arg	Pro	His	His	Gln	
			100					105					110			
Ala	Arg	Gln	Val	Glu	His	Ser	Thr	Gln	Arg	Gly	His	Leu	Glu	Ile	Arg	
		115					120					125				
Glu	Leu	Lys	Lys	Lys	Leu	Phe	Lys	Arg	Arg	Arg	Val	Leu	Asn	Arg	Glu	
130						135					140					
Arg	Arg	Leu	Arg	His	Arg	Val	Val	Gly	Ala	Val	Ile	Asp	Gln	Gly	Leu	
145					150					155					160	
Ile	Thr	Arg	His	His	Leu	Lys	Lys	Arg	Ala	Ala	Gln	Glu	Leu	Ser	Gln	
				165					170					175		
Glu	Ile	Lys	Ala	Phe	Leu	Thr	Gly	Val	Asp	Pro	Ile	Leu	Gly	His	Gln	
			180					185					190			
Leu	Ser	Ala	Arg	Glu	His	Ala	Arg	Cys	Gly	Leu	Leu	Leu	Leu	Arg	Ser	
		195					200					205				
Leu	Pro	Pro	Ala	Arg	Ala	Ala	Val	Leu	Asp	His	Leu	Arg	Gly	Val	Phe	
	210					215					220					
Asp	Glu	Ser	Val	Arg	Ala	His	Leu	Ala	Ala	Leu	Asp	Glu	Thr	Pro	Val	
225					230					235					240	
Ala	Gly	Pro	Pro	His	Leu	Arg	Pro	Pro	Pro	Pro	Ser	His	Val	Pro	Ala	
				245					250					255		
Gly	Gly	Pro	Gly	Leu	Glu	Asp	Val	Val	Gln	Glu	Val	Gln	Gln	Val	Leu	
			260					265					270			
Ser	Glu	Phe	Ile	Arg	Ala	Asn	Pro	Lys	Ala	Trp	Ala	Pro	Val	Ile	Ser	
		275					280					285				
Ala	Trp	Ser	Ile	Asp	Leu	Met	Gly	Gln	Leu	Ser	Ser	Thr	Tyr	Ser	Gly	
	290					295					300					
Gln	His	Gln	Arg	Val	Pro	His	Ala	Thr	Gly	Ala	Leu	Asn	Glu	Leu	Leu	
305					310					315					320	
Gln	Leu	Trp	Met	Gly	Cys	Arg	Ala	Thr	Arg	Thr	Leu	Met	Asp	Ile	Tyr	
			325						330					335		
Val	Gln	Cys	Leu	Ser	Ala	Leu	Ile	Gly	Ser	Cys	Pro	Asp	Ala	Cys	Val	
			340					345					350			
Asp	Ala	Leu	Leu	Asp	Thr	Ser	Val	Gln	His	Ser	Pro	His	Phe	Asp	Trp	
	355						360					365				
Val	Val	Ala	His	Ile	Gly	Ser	Ser	Phe	Pro	Gly	Thr	Ile	Ile	Ser	Arg	

2003295784 19 Aug 2005

370	375	380
Val Leu Ser Cys Gly	Leu Lys Asp Phe Cys	Val His Gly Gly Ala Gly
385	390	395
Gly Gly Ala Gly Ser	Ser Gly Gly Ser Ser	Ser Gln Thr Pro Ser Thr
405	410	415
Asp Pro Phe Pro Gly	Ser Pro Ala Ile Pro Ala	Glu Lys Arg Val Pro
420	425	430
Lys Ile Ala Ser Val Val	Gly Ile Leu Gly His Leu	Ala Ser Arg His
435	440	445
Gly Asp Ser Ile Arg Arg	Glu Leu Leu Arg Met Phe	His Asp Ser Leu
450	455	460
Ala Gly Gly Ser Gly Gly	Arg Ser Gly Asp Pro Ser	Leu Gln Ala Thr
465	470	475
Val Pro Phe Leu Leu Gln	Leu Ala Val Met Ser Pro	Ala Leu Leu Gly
485	490	495
Thr Val Ser Gly Glu Leu	Val Asp Cys Leu Lys Pro	Pro Ala Val Leu
500	505	510
Ser Gln Leu Gln Gln His	Leu Gln Gly Phe Pro Arg	Glu Glu Leu Asp
515	520	525
Asn Met Leu Asn Leu Ala	Val His Leu Val Ser Gln	Ala Ser Gly Ala
530	535	540
Gly Ala Tyr Arg Leu Leu	Gln Phe Leu Val Asp Thr	Ala Met Pro Ala
545	550	555
Ser Val Ile Thr Thr Gln	Gly Leu Ala Val Pro Asp	Thr Val Arg Glu
565	570	575
Ala Cys Asp Arg Leu Ile	Gln Leu Leu Leu His Leu	Gln Lys Leu
580	585	590
Val His His Arg Gly Gly	Ser Pro Gly Glu Gly Val	Leu Gly Pro Pro
595	600	605
Pro Pro Pro Arg Leu Val	Pro Phe Leu Asp Ala Leu	Lys Asn His Val
610	615	620
Gly Glu Leu Cys Gly Glu	Thr Leu Arg Leu Glu Arg	Lys Arg Phe Leu
625	630	635
Trp Gln His Gln Leu Leu	Gly Leu Leu Ser Val Tyr	Thr Arg Pro Ser
645	650	655
Cys Gly Pro Glu Ala Leu	Gly His Leu Leu Ser Arg	Ala Arg Ser Pro
660	665	670
Glu Glu Leu Ser Leu Ala	Thr Gln Leu Tyr Ala Gly	Leu Val Val Ser
675	680	685
Leu Ser Gly Leu Leu Pro	Leu Ala Phe Arg Ser Cys	Leu Ala Arg Val
690	695	700

His 705	Ala	Gly	Thr	Leu	Gln	Pro	Pro	Phe	Thr	Ala	Arg	Phe	Leu	Arg	Asn 720
Leu	Ala	Leu	Leu	Val	Gly	Trp	Glu	Gln	Gln	Gly	Gly	Glu	Gly	Pro	Ala
				725					730					735	
Ala	Leu	Gly	Ala	His	Phe	Gly	Glu	Ser	Ala	Ser	Ala	His	Leu	Ser	Asp
			740					745					750		
Leu	Ala	Pro	Leu	Leu	Leu	His	Pro	Glu	Glu	Glu	Val	Ala	Glu	Ala	Ala
		755					760					765			
Ala	Ser	Leu	Leu	Ala	Ile	Cys	Pro	Phe	Pro	Ser	Glu	Ala	Leu	Ser	Pro
	770					775					780				
Ser	Gln	Leu	Leu	Gly	Leu	Val	Arg	Ala	Gly	Val	His	Arg	Phe	Phe	Ala
785					790					795					800
Ser	Leu	Arg	Leu	His	Gly	Pro	Pro	Gly	Val	Ala	Ser	Ala	Cys	Gln	Leu
				805					810					815	
Leu	Thr	Arg	Leu	Ser	Gln	Thr	Ser	Pro	Ala	Gly	Leu	Lys	Ala	Val	Leu
			820					825					830		
Gln	Leu	Leu	Val	Glu	Gly	Ala	Leu	His	Arg	Gly	Asn	Thr	Glu	Leu	Phe
			835				840					845			
Gly	Gly	Gln	Val	Asp	Gly	Asp	Asn	Glu	Thr	Leu	Ser	Val	Val	Ser	Ala
	850					855					860				
Ser	Leu	Ala	Ser	Ala	Ser	Leu	Leu	Asp	Thr	Asn	Arg	Arg	His	Thr	Ala
865					870					875					880
Ala	Val	Pro	Gly	Pro	Gly	Gly	Ile	Trp	Ser	Val	Phe	His	Ala	Gly	Val
				885					890					895	
Ile	Gly	Arg	Gly	Leu	Lys	Pro	Pro	Lys	Phe	Val	Gln	Ser	Arg	Asn	Gln
			900					905					910		
Gln	Glu	Val	Ile	Tyr	Asn	Thr	Gln	Ser	Leu	Leu	Ser	Leu	Leu	Val	His
		915					920					925			
Cys	Cys	Ser	Ala	Pro	Gly	Gly	Thr	Glu	Cys	Gly	Glu	Cys	Trp	Gly	Ala
	930					935					940				
Pro	Ile	Leu	Ser	Pro	Glu	Ala	Ala	Lys	Ala	Val	Ala	Val	Thr	Leu	Val
945					950					955					960
Glu	Ser	Val	Cys	Pro	Asp	Ala	Ala	Gly	Ala	Glu	Leu	Ala	Trp	Pro	Pro
				965					970					975	
Glu	Glu	His	Ala	Arg	Ala	Thr	Val	Glu	Arg	Asp	Leu	Arg	Ile	Gly	Arg
			980					985					990		
Arg	Phe	Arg	Glu	Gln	Pro	Leu	Leu	Phe	Glu	Leu	Leu	Lys	Leu	Val	Ala
			995				1000					1005			
Ala	Ala	Pro	Pro	Ala	Leu	Cys	Tyr	Cys	Ser	Val	Leu	Leu	Arg	Gly	
	1010					1015					1020				
Leu	Leu	Ala	Ala	Leu	Leu	Gly	His	Trp	Glu	Ala	Ser	Arg	His	Thr	
	1025					1030					1035				

2003295784 19 Aug 2005

Asp Thr Thr His Ser Pro Trp His Leu Glu Ala Ser Cys Thr Leu
 1040 1045 1050
 Val Ala Val Met Ala Glu Gly Ser Leu Leu Pro Pro Ala Leu Gly
 1055 1060 1065
 Asn Met His Glu Val Phe Ser Gln Leu Ala Pro Phe Glu Val Arg
 1070 1075 1080
 Leu Leu Leu Leu Ser Val Trp Gly Phe Leu Arg Glu His Gly Pro
 1085 1090 1095
 Leu Pro Gln Lys Phe Ile Phe Gln Ser Glu Arg Gly Arg Phe Ile
 1100 1105 1110
 Arg Asp Phe Ser Arg Glu Gly Gly Gly Glu Gly Gly Pro His Leu
 1115 1120 1125
 Ala Val Leu His Ser Val Leu His Arg Asn Ile Asp Arg Leu Gly
 1130 1135 1140
 Leu Phe Ser Gly Arg Phe Gln Ala Pro Ser Pro Ser Thr Leu Leu
 1145 1150 1155
 Arg Gln Gly Thr
 1160

<210> 14
 <211> 1162
 <212> PRT
 <213> Artificial

<220>
 <223> A variant of the human BFLP1698 polypeptide

<400> 14

Met Ala Leu Val Pro Gly Arg Ser Lys Glu Asp Gly Leu Trp Thr Arg
 1 5 10 15
 Asn Ser Pro Gly Ser Ser Gln His Pro Glu Ser Pro Arg Leu Pro Asn
 20 25 30
 Pro Leu Trp Asp Arg Gly Lys Ile Gly Lys Val Glu Gly His Gln His
 35 40 45
 Ile Gln Asp Phe Ser Gln Lys Ser His Leu Pro Ser Ile Val Val Glu
 50 55 60
 Ser Ser Glu Val Asn Glu Glu Ser Gly Asp Leu His Leu Pro His Glu
 65 70 75 80
 Glu Leu Leu Leu Leu Thr Asp Gly Glu Glu Glu Asp Ala Glu Ala Phe
 85 90 95
 Phe Gln Asp Gln Ser Glu Glu Pro Gly Ala Ala Arg Pro His His Gln
 100 105 110
 Ala Arg Gln Val Glu His Ser Thr Gln Arg Gly His Leu Glu Ile Arg
 115 120 125

2003295784 19 Aug 2005

Glu	Leu	Lys	Lys	Lys	Leu	Phe	Lys	Arg	Arg	Arg	Val	Leu	Asn	Arg	Glu	130	135	140
Arg	Arg	Leu	Arg	His	Arg	Val	Val	Gly	Ala	Val	Ile	Asp	Gln	Gly	Leu	145	150	155
Ile	Thr	Arg	His	His	Leu	Lys	Lys	Arg	Ala	Ala	Gln	Glu	Leu	Ser	Gln	165	170	175
Glu	Ile	Lys	Ala	Phe	Leu	Thr	Gly	Val	Asp	Pro	Ile	Leu	Gly	His	Gln	180	185	190
Leu	Ser	Ala	Arg	Glu	His	Ala	Arg	Cys	Gly	Leu	Leu	Leu	Leu	Arg	Ser	195	200	205
Leu	Pro	Pro	Ala	Arg	Ala	Ala	Val	Leu	Asp	His	Leu	Arg	Gly	Val	Phe	210	215	220
Asp	Glu	Ser	Val	Arg	Ala	His	Leu	Ala	Ala	Leu	Asp	Glu	Thr	Pro	Val	225	230	235
Ala	Gly	Pro	Pro	His	Leu	Arg	Pro	Pro	Pro	Pro	Ser	His	Val	Pro	Ala	245	250	255
Gly	Gly	Pro	Gly	Leu	Glu	Asp	Val	Val	Gln	Glu	Val	Gln	Gln	Val	Leu	260	265	270
Ser	Glu	Phe	Ile	Arg	Ala	Asn	Pro	Lys	Ala	Trp	Ala	Pro	Val	Ile	Ser	275	280	285
Ala	Trp	Ser	Ile	Asp	Leu	Met	Gly	Gln	Leu	Ser	Ser	Thr	Tyr	Ser	Gly	290	295	300
Gln	His	Gln	Arg	Val	Pro	His	Ala	Thr	Gly	Ala	Leu	Asn	Glu	Leu	Leu	305	310	315
Gln	Leu	Trp	Met	Gly	Cys	Arg	Ala	Thr	Arg	Thr	Leu	Met	Asp	Ile	Tyr	325	330	335
Val	Gln	Cys	Leu	Ser	Ala	Leu	Ile	Gly	Ser	Cys	Pro	Asp	Ala	Cys	Val	340	345	350
Asp	Ala	Leu	Leu	Asp	Thr	Ser	Val	Gln	His	Ser	Pro	His	Phe	Asp	Trp	355	360	365
Val	Val	Ala	His	Ile	Gly	Ser	Ser	Phe	Pro	Gly	Thr	Ile	Ile	Ser	Arg	370	375	380
Val	Leu	Ser	Cys	Gly	Leu	Lys	Asp	Phe	Cys	Val	His	Gly	Gly	Ala	Gly	385	390	395
Gly	Gly	Ala	Gly	Ser	Ser	Gly	Gly	Ser	Ser	Ser	Gln	Thr	Pro	Ser	Thr	405	410	415
Asp	Pro	Phe	Pro	Gly	Ser	Pro	Ala	Ile	Pro	Ala	Glu	Lys	Arg	Val	Pro	420	425	430
Lys	Ile	Ala	Ser	Val	Val	Gly	Ile	Leu	Gly	His	Leu	Ala	Ser	Arg	His	435	440	445
Gly	Asp	Ser	Ile	Arg	Arg	Glu	Leu	Leu	Arg	Met	Phe	His	Asp	Ser	Leu			

2003295784 19 Aug 2005

450	455	460
Ala Gly Gly Ser Gly Gly Arg Ser Gly Asp Pro Ser Leu Gln Ala Thr		
465	470	475
Val Pro Phe Leu Leu Gln Leu Ala Val Met Ser Pro Ala Leu Leu Gly		
	485	490
Thr Val Ser Gly Glu Leu Val Asp Cys Leu Lys Pro Pro Ala Val Leu		
	500	505
Ser Gln Leu Gln Gln His Leu Gln Gly Phe Pro Arg Glu Glu Leu Asp		
	515	520
Asn Met Leu Asn Leu Ala Val His Leu Val Ser Gln Ala Ser Gly Ala		
	530	535
Gly Ala Tyr Arg Leu Leu Gln Phe Leu Val Asp Thr Ala Met Pro Ala		
	545	550
Ser Val Ile Thr Thr Gln Gly Leu Ala Val Pro Asp Thr Val Arg Glu		
	565	570
Ala Cys Asp Arg Leu Ile Gln Leu Leu Leu Leu His Leu Gln Lys Leu		
	580	585
Val His His Arg Gly Gly Ser Pro Gly Glu Gly Val Leu Gly Pro Pro		
	595	600
Pro Pro Pro Arg Leu Val Pro Phe Leu Asp Ala Leu Lys Asn His Val		
	610	615
Gly Glu Leu Cys Gly Glu Thr Leu Arg Leu Glu Arg Lys Arg Phe Leu		
	625	630
Trp Gln His Gln Leu Leu Gly Leu Leu Ser Val Tyr Thr Arg Pro Ser		
	645	650
Cys Gly Pro Glu Ala Leu Gly His Leu Leu Ser Arg Ala Arg Ser Pro		
	660	665
Glu Glu Leu Ser Leu Ala Thr Gln Leu Tyr Ala Gly Leu Val Val Ser		
	675	680
Leu Ser Gly Leu Leu Pro Leu Ala Phe Arg Ser Cys Leu Ala Arg Val		
	690	695
His Ala Gly Thr Leu Gln Pro Pro Phe Thr Ala Arg Phe Leu Arg Asn		
	705	710
Leu Ala Leu Leu Val Gly Trp Glu Gln Gln Gly Gly Glu Gly Pro Ala		
	725	730
Ala Leu Gly Ala His Phe Gly Glu Ser Ala Ser Ala His Leu Ser Asp		
	740	745
Leu Ala Pro Leu Leu Leu His Pro Glu Glu Glu Val Ala Glu Ala Ala		
	755	760
Ala Ser Leu Leu Ala Ile Cys Pro Phe Pro Ser Glu Ala Leu Ser Pro		
	770	775
		780

2003295784 19 Aug 2005

Ser Gln Leu Leu Gly Leu Val Arg Ala Gly Val His Arg Phe Phe Ala
 785 790 795 800
 Ser Leu Arg Leu His Gly Pro Pro Gly Val Ala Ser Ala Cys Gln Leu
 805 810 815
 Leu Thr Arg Leu Ser Gln Thr Ser Pro Ala Gly Leu Lys Ala Val Leu
 820 825 830
 Gln Leu Leu Val Glu Gly Ala Leu His Arg Gly Asn Thr Glu Leu Phe
 835 840 845
 Gly Gly Gln Val Asp Gly Asp Asn Glu Thr Leu Ser Val Val Ser Ala
 850 855 860
 Ser Leu Ala Ser Ala Ser Leu Leu Asp Thr Asn Arg Arg His Thr Ala
 865 870 875 880
 Ala Val Pro Gly Pro Gly Gly Ile Trp Ser Val Phe His Ala Gly Val
 885 890 895
 Ile Gly Arg Gly Leu Lys Pro Pro Lys Phe Val Gln Ser Arg Asn Gln
 900 905 910
 Gln Glu Val Ile Tyr Asn Thr Gln Ser Leu Leu Ser Leu Leu Val His
 915 920 925
 Cys Cys Ser Ala Pro Gly Gly Thr Glu Cys Gly Glu Cys Trp Gly Ala
 930 935 940
 Pro Ile Leu Ser Pro Glu Ala Ala Lys Ala Val Ala Val Thr Leu Val
 945 950 955 960
 Glu Ser Val Cys Pro Asp Ala Ala Gly Ala Glu Leu Ala Trp Pro Pro
 965 970 975
 Glu Glu His Ala Arg Ala Thr Val Glu Arg Asp Leu Arg Ile Gly Arg
 980 985 990
 Arg Phe Arg Glu Gln Pro Leu Leu Phe Glu Leu Leu Lys Leu Val Ala
 995 1000 1005
 Ala Ala Pro Pro Ala Leu Cys Tyr Cys Ser Val Leu Leu Arg Gly
 1010 1015 1020
 Leu Leu Ala Ala Leu Leu Gly His Trp Glu Ala Ser Arg His Thr
 1025 1030 1035
 Asp Thr Thr His Ser Pro Trp His Leu Glu Ala Ser Cys Thr Leu
 1040 1045 1050
 Val Ala Val Met Ala Glu Gly Ser Leu Leu Pro Pro Ala Leu Gly
 1055 1060 1065
 Asn Met His Glu Val Phe Ser Gln Leu Ala Pro Phe Glu Val Arg
 1070 1075 1080
 Leu Leu Leu Leu Ser Val Trp Gly Phe Leu Arg Glu His Gly Pro
 1085 1090 1095
 Leu Pro Gln Lys Phe Ile Phe Gln Ser Glu Arg Gly Arg Phe Ile
 1100 1105 1110

2003295784 19 Aug 2005

Arg Asp Phe Ser Arg Glu Gly Gly Gly Glu Gly Gly Pro His Leu
1115 1120 1125

Ala Val Leu His Ser Val Leu His Arg Asn Leu Asp Arg Leu Gly
1130 1135 1140

Leu Phe Ser Gly Arg Phe Gln Ala Pro Ser Pro Ser Thr Leu Leu
1145 1150 1155

Arg Gln Gly Thr
1160

<210> 15
<211> 170
<212> PRT
<213> Homo sapiens

<400> 15

Met Ala Leu Val Pro Gly Arg Ser Lys Glu Asp Gly Leu Trp Thr Arg
1 5 10 15

Asn Ser Pro Gly Ser Ser Gln His Pro Glu Ser Pro Arg Leu Pro Asn
20 25 30

Pro Leu Trp Asp Arg Gly Lys Ile Gly Lys Val Glu Gly His Gln His
35 40 45

Ile Gln Asp Phe Ser Gln Lys Ser His Leu Pro Ser Ile Val Val Glu
50 55 60

Ser Ser Glu Val Asn Glu Glu Ser Gly Asp Leu His Leu Pro His Glu
65 70 75 80

Glu Leu Leu Leu Leu Thr Asp Gly Glu Glu Glu Asp Ala Glu Ala Phe
85 90 95

Phe Gln Asp Gln Ser Glu Glu Pro Gly Ala Ala Arg Pro His His Gln
100 105 110

Ala Arg Gln Val Glu His Ser Thr Gln Arg Gly His Leu Glu Ile Arg
115 120 125

Glu Leu Lys Lys Lys Leu Phe Lys Arg Arg Arg Val Leu Asn Arg Glu
130 135 140

Arg Arg Leu Arg His Arg Val Val Gly Ala Val Ile Asp Gln Gly Leu
145 150 155 160

Ile Thr Arg His His Leu Lys Lys Arg Ala
165 170

<210> 16
<211> 1019
<212> PRT
<213> Homo sapiens

<400> 16

2003295784 19 Aug 2005

Met	Ser	Ala	Leu	Cys	Asp	Pro	Pro	Gly	Ala	Pro	Gly	Pro	Pro	Gly	Pro		
1				5					10					15			
Ala	Pro	Ala	Thr	His	Gly	Pro	Ala	Pro	Leu	Ser	Ala	Gln	Glu	Leu	Ser		
			20					25					30				
Gln	Glu	Ile	Lys	Ala	Phe	Leu	Thr	Gly	Val	Asp	Pro	Ile	Leu	Gly	His		
		35					40					45					
Gln	Leu	Ser	Ala	Arg	Glu	His	Ala	Arg	Cys	Gly	Leu	Leu	Leu	Leu	Arg		
	50					55					60						
Ser	Leu	Pro	Pro	Ala	Arg	Ala	Ala	Val	Leu	Asp	His	Leu	Arg	Gly	Val		
65					70					75					80		
Phe	Asp	Glu	Ser	Val	Arg	Ala	His	Leu	Ala	Ala	Leu	Asp	Glu	Thr	Pro		
				85					90					95			
Val	Ala	Gly	Pro	Pro	His	Leu	Arg	Pro	Pro	Pro	Pro	Ser	His	Val	Pro		
			100					105					110				
Ala	Gly	Gly	Pro	Gly	Leu	Glu	Asp	Val	Val	Gln	Glu	Val	Gln	Gln	Val		
		115					120					125					
Leu	Ser	Glu	Phe	Ile	Arg	Ala	Asn	Pro	Lys	Ala	Trp	Ala	Pro	Val	Ile		
	130					135					140						
Ser	Ala	Trp	Ser	Ile	Asp	Leu	Met	Gly	Gln	Leu	Ser	Ser	Thr	Tyr	Ser		
145					150					155					160		
Gly	Gln	His	Gln	Arg	Val	Pro	His	Ala	Thr	Gly	Ala	Leu	Asn	Glu	Leu		
				165					170					175			
Leu	Gln	Leu	Trp	Met	Gly	Cys	Arg	Ala	Thr	Arg	Thr	Leu	Met	Asp	Ile		
			180					185					190				
Tyr	Val	Gln	Cys	Leu	Ser	Ala	Leu	Ile	Gly	Ser	Cys	Pro	Asp	Ala	Cys		
		195					200					205					
Val	Asp	Ala	Leu	Leu	Asp	Thr	Ser	Val	Gln	His	Ser	Pro	His	Phe	Asp		
	210					215					220						
Trp	Val	Val	Ala	His	Ile	Gly	Ser	Ser	Phe	Pro	Gly	Thr	Ile	Ile	Ser		
225					230					235					240		
Arg	Val	Leu	Ser	Cys	Gly	Leu	Lys	Asp	Phe	Cys	Val	His	Gly	Gly	Ala		
				245					250					255			
Gly	Gly	Gly	Ala	Gly	Ser	Ser	Gly	Gly	Ser	Ser	Ser	Gln	Thr	Pro	Ser		
			260					265					270				
Thr	Asp	Pro	Phe	Pro	Gly	Ser	Pro	Ala	Ile	Pro	Ala	Glu	Lys	Arg	Val		
		275					280					285					
Pro	Lys	Ile	Ala	Ser	Val	Val	Gly	Ile	Leu	Gly	His	Leu	Ala	Ser	Arg		
	290					295					300						
His	Gly	Asp	Ser	Ile	Arg	Arg	Glu	Leu	Leu	Arg	Met	Phe	His	Asp	Ser		
305					310					315					320		
Leu	Ala	Gly	Gly	Ser	Gly	Gly	Arg	Ser	Gly	Asp	Pro	Ser	Leu	Gln	Ala		
				325					330					335			

2003295784 19 Aug 2005

Thr Val Pro Phe Leu Leu Gln Leu Ala Val Met Ser Pro Ala Leu Leu
 340 345 350
 Gly Thr Val Ser Gly Glu Leu Val Asp Cys Leu Lys Pro Pro Ala Val
 355 360 365
 Leu Ser Gln Leu Gln Gln His Leu Gln Gly Phe Pro Arg Glu Glu Leu
 370 375 380
 Asp Asn Met Leu Asn Leu Ala Val His Leu Val Ser Gln Ala Ser Gly
 385 390 395 400
 Ala Gly Ala Tyr Arg Leu Leu Gln Phe Leu Val Asp Thr Ala Met Pro
 405 410 415
 Ala Ser Val Ile Thr Thr Gln Gly Leu Ala Val Pro Asp Thr Val Arg
 420 425 430
 Glu Ala Cys Asp Arg Leu Ile Gln Leu Leu Leu Leu His Leu Gln Lys
 435 440 445
 Leu Val His His Arg Gly Gly Ser Pro Gly Glu Gly Val Leu Gly Pro
 450 455 460
 Pro Pro Pro Pro Arg Leu Val Pro Phe Leu Asp Ala Leu Lys Asn His
 465 470 475 480
 Val Gly Glu Leu Cys Gly Glu Thr Leu Arg Leu Glu Arg Lys Arg Phe
 485 490 495
 Leu Trp Gln His Gln Leu Leu Gly Leu Leu Ser Val Tyr Thr Arg Pro
 500 505 510
 Ser Cys Gly Pro Glu Ala Leu Gly His Leu Leu Ser Arg Ala Arg Ser
 515 520 525
 Pro Glu Glu Leu Ser Leu Ala Thr Gln Leu Tyr Ala Gly Leu Val Val
 530 535 540
 Ser Leu Ser Gly Leu Leu Pro Leu Ala Phe Arg Ser Cys Leu Ala Arg
 545 550 555 560
 Val His Ala Gly Thr Leu Gln Pro Pro Phe Thr Ala Arg Phe Leu Arg
 565 570 575
 Asn Leu Ala Leu Leu Val Gly Trp Glu Gln Gln Gly Gly Glu Gly Pro
 580 585 590
 Ala Ala Leu Gly Ala His Phe Gly Glu Ser Ala Ser Ala His Leu Ser
 595 600 605
 Asp Leu Ala Pro Leu Leu Leu His Pro Glu Glu Glu Val Ala Glu Ala
 610 615 620
 Ala Ala Ser Leu Leu Ala Ile Cys Pro Phe Pro Ser Glu Ala Leu Ser
 625 630 635 640
 Pro Ser Gln Leu Leu Gly Leu Val Arg Ala Gly Val His Arg Phe Phe
 645 650 655
 Ala Ser Leu Arg Leu His Gly Pro Pro Gly Val Ala Ser Ala Cys Gln

2003295784 19 Aug 2005

660					665					670						
Leu	Leu	Thr	Arg	Leu	Ser	Gln	Thr	Ser	Pro	Ala	Gly	Leu	Lys	Ala	Val	
675					680					685						
Leu	Gln	Leu	Leu	Val	Glu	Gly	Ala	Leu	His	Arg	Gly	Asn	Thr	Glu	Leu	
690					695					700						
Phe	Gly	Gly	Gln	Val	Asp	Gly	Asp	Asn	Glu	Thr	Leu	Ser	Val	Val	Ser	
705					710					715					720	
Ala	Ser	Leu	Ala	Ser	Ala	Ser	Leu	Leu	Asp	Thr	Asn	Arg	Arg	His	Thr	
725					730					735						
Ala	Ala	Val	Pro	Gly	Pro	Gly	Gly	Ile	Trp	Ser	Val	Phe	His	Ala	Gly	
740					745					750						
Val	Ile	Gly	Arg	Gly	Leu	Lys	Pro	Pro	Lys	Phe	Val	Gln	Ser	Arg	Asn	
755					760					765						
Gln	Gln	Glu	Val	Ile	Tyr	Asn	Thr	Gln	Ser	Leu	Leu	Ser	Leu	Leu	Val	
770					775					780						
His	Cys	Cys	Ser	Ala	Pro	Gly	Gly	Thr	Glu	Cys	Gly	Glu	Cys	Trp	Gly	
785					790					795					800	
Ala	Pro	Ile	Leu	Ser	Pro	Glu	Ala	Ala	Lys	Ala	Val	Ala	Val	Thr	Leu	
805					810					815						
Val	Glu	Ser	Val	Cys	Pro	Asp	Ala	Ala	Gly	Ala	Glu	Leu	Ala	Trp	Pro	
820					825					830						
Pro	Glu	Glu	His	Ala	Arg	Ala	Thr	Val	Glu	Arg	Asp	Leu	Arg	Ile	Gly	
835					840					845						
Arg	Arg	Phe	Arg	Glu	Gln	Pro	Leu	Leu	Phe	Glu	Leu	Leu	Lys	Leu	Val	
850					855					860						
Ala	Ala	Ala	Pro	Pro	Ala	Leu	Cys	Tyr	Cys	Ser	Val	Leu	Leu	Arg	Gly	
865					870					875					880	
Leu	Leu	Ala	Ala	Leu	Leu	Gly	His	Trp	Glu	Ala	Ser	Arg	His	Pro	Asp	
885					890					895						
Thr	Thr	His	Ser	Pro	Trp	His	Leu	Glu	Ala	Ser	Cys	Thr	Leu	Val	Ala	
900					905					910						
Val	Met	Ala	Glu	Gly	Ser	Leu	Leu	Pro	Pro	Ala	Leu	Gly	Asn	Met	His	
915					920					925						
Glu	Val	Phe	Ser	Gln	Leu	Ala	Pro	Phe	Glu	Val	Arg	Leu	Leu	Leu	Leu	
930					935					940						
Ser	Val	Trp	Gly	Phe	Leu	Arg	Glu	His	Gly	Pro	Leu	Pro	Gln	Lys	Phe	
945					950					955					960	
Ile	Phe	Gln	Ser	Glu	Arg	Gly	Arg	Phe	Ile	Arg	Asp	Phe	Ser	Arg	Glu	
965					970					975						
Gly	Gly	Gly	Glu	Gly	Gly	Pro	His	Leu	Ala	Val	Leu	His	Ser	Val	Leu	
980					985					990						

2003295784 19 Aug 2005

His Arg Asn Ile Asp Arg Leu Gly Leu Phe Ser Gly Arg Phe Gln Ala
 995 1000 1005

Pro Ser Pro Ser Thr Leu Leu Arg Gln Gly Thr
 1010 1015

<210> 17
 <211> 908
 <212> PRT
 <213> Homo sapiens

<400> 17

Pro Ala Gly Gly Pro Gly Leu Glu Asp Val Val Gln Glu Val Gln Gln
 1 5 10 15

Val Leu Ser Glu Phe Ile Arg Ala Asn Pro Lys Ala Trp Ala Pro Val
 20 25 30

Ile Ser Ala Trp Ser Ile Asp Leu Met Gly Gln Leu Ser Ser Thr Tyr
 35 40 45

Ser Gly Gln His Gln Arg Val Pro His Ala Thr Gly Ala Leu Asn Glu
 50 55 60

Leu Leu Gln Leu Trp Met Gly Cys Arg Ala Thr Arg Thr Leu Met Asp
 65 70 75 80

Ile Tyr Val Gln Cys Leu Ser Ala Leu Ile Gly Ser Cys Pro Asp Ala
 85 90 95

Cys Val Asp Ala Leu Leu Asp Thr Ser Val Gln His Ser Pro His Phe
 100 105 110

Asp Trp Val Val Ala His Ile Gly Ser Ser Phe Pro Gly Thr Ile Ile
 115 120 125

Ser Arg Val Leu Ser Cys Gly Leu Lys Asp Phe Cys Val His Gly Gly
 130 135 140

Ala Gly Gly Gly Ala Gly Ser Ser Gly Gly Ser Ser Ser Gln Thr Pro
 145 150 155 160

Ser Thr Asp Pro Phe Pro Gly Ser Pro Ala Ile Pro Ala Glu Lys Arg
 165 170 175

Val Pro Lys Ile Ala Ser Val Val Gly Ile Leu Gly His Leu Ala Ser
 180 185 190

Arg His Gly Asp Ser Ile Arg Arg Glu Leu Leu Arg Met Phe His Asp
 195 200 205

Ser Leu Ala Gly Gly Ser Gly Gly Arg Ser Gly Asp Pro Ser Leu Gln
 210 215 220

Ala Thr Val Pro Phe Leu Leu Gln Leu Ala Val Met Ser Pro Ala Leu
 225 230 235 240

Leu Gly Thr Val Ser Gly Glu Leu Val Asp Cys Leu Lys Pro Pro Ala
 245 250 255

2003295784 19 Aug 2005

Val	Leu	Ser	Gln	Leu	Gln	Gln	His	Leu	Gln	Gly	Phe	Pro	Arg	Glu	Glu	
			260					265						270		
Leu	Asp	Asn	Met	Leu	Asn	Leu	Ala	Val	His	Leu	Val	Ser	Gln	Ala	Ser	
		275					280					285				
Gly	Ala	Gly	Ala	Tyr	Arg	Leu	Leu	Gln	Phe	Leu	Val	Asp	Thr	Ala	Met	
	290					295					300					
Pro	Ala	Ser	Val	Ile	Thr	Thr	Gln	Gly	Leu	Ala	Val	Pro	Asp	Thr	Val	
305					310					315					320	
Arg	Glu	Ala	Cys	Asp	Arg	Leu	Ile	Gln	Leu	Leu	Leu	Leu	His	Leu	Gln	
			325					330						335		
Lys	Leu	Val	His	His	Arg	Gly	Gly	Ser	Pro	Gly	Glu	Gly	Val	Leu	Gly	
		340					345						350			
Pro	Pro	Pro	Pro	Pro	Arg	Leu	Val	Pro	Phe	Leu	Asp	Ala	Leu	Lys	Asn	
		355				360						365				
His	Val	Gly	Glu	Leu	Cys	Gly	Glu	Thr	Leu	Arg	Leu	Glu	Arg	Lys	Arg	
	370				375					380						
Phe	Leu	Trp	Gln	His	Gln	Leu	Leu	Gly	Leu	Leu	Ser	Val	Tyr	Thr	Arg	
385					390				395						400	
Pro	Ser	Cys	Gly	Pro	Glu	Ala	Leu	Gly	His	Leu	Leu	Ser	Arg	Ala	Arg	
			405					410						415		
Ser	Pro	Glu	Glu	Leu	Ser	Leu	Ala	Thr	Gln	Leu	Tyr	Ala	Gly	Leu	Val	
		420						425					430			
Val	Ser	Leu	Ser	Gly	Leu	Leu	Pro	Leu	Ala	Phe	Arg	Ser	Cys	Leu	Ala	
		435					440					445				
Arg	Val	His	Ala	Gly	Thr	Leu	Gln	Pro	Pro	Phe	Thr	Ala	Arg	Phe	Leu	
	450					455					460					
Arg	Asn	Leu	Ala	Leu	Leu	Val	Gly	Trp	Glu	Gln	Gln	Gly	Gly	Glu	Gly	
465				470					475					480		
Pro	Ala	Ala	Leu	Gly	Ala	His	Phe	Gly	Glu	Ser	Ala	Ser	Ala	His	Leu	
			485					490						495		
Ser	Asp	Leu	Ala	Pro	Leu	Leu	Leu	His	Pro	Glu	Glu	Glu	Val	Ala	Glu	
		500						505					510			
Ala	Ala	Ala	Ser	Leu	Leu	Ala	Ile	Cys	Pro	Phe	Pro	Ser	Glu	Ala	Leu	
		515					520					525				
Ser	Pro	Ser	Gln	Leu	Leu	Gly	Leu	Val	Arg	Ala	Gly	Val	His	Arg	Phe	
	530					535					540					
Phe	Ala	Ser	Leu	Arg	Leu	His	Gly	Pro	Pro	Gly	Val	Ala	Ser	Ala	Cys	
545					550					555					560	
Gln	Leu	Leu	Thr	Arg	Leu	Ser	Gln	Thr	Ser	Pro	Ala	Gly	Leu	Lys	Ala	
			565						570					575		
Val	Leu	Gln	Leu	Leu	Val	Glu	Gly	Ala	Leu	His	Arg	Gly	Asn	Thr	Glu	
			580					585					590			

2003295784 19 Aug 2005

Leu Phe Gly Gly Gln Val Asp Gly Asp Asn Glu Thr Leu Ser Val Val
 595 600 605
 Ser Ala Ser Leu Ala Ser Ala Ser Leu Leu Asp Thr Asn Arg Arg His
 610 615 620
 Thr Ala Ala Val Pro Gly Pro Gly Gly Ile Trp Ser Val Phe His Ala
 625 630 635 640
 Gly Val Ile Gly Arg Gly Leu Lys Pro Pro Lys Phe Val Gln Ser Arg
 645 650 655
 Asn Gln Gln Glu Val Ile Tyr Asn Thr Gln Ser Leu Leu Ser Leu Leu
 660 665 670
 Val His Cys Cys Ser Ala Pro Gly Gly Thr Glu Cys Gly Glu Cys Trp
 675 680 685
 Gly Ala Pro Ile Leu Ser Pro Glu Ala Ala Lys Ala Val Ala Val Thr
 690 695 700
 Leu Val Glu Ser Val Cys Pro Asp Ala Ala Gly Ala Glu Leu Ala Trp
 705 710 715 720
 Pro Pro Glu Glu His Ala Arg Ala Thr Val Glu Arg Asp Leu Arg Ile
 725 730 735
 Gly Arg Arg Phe Arg Glu Gln Pro Leu Leu Phe Glu Leu Leu Lys Leu
 740 745 750
 Val Ala Ala Ala Pro Pro Ala Leu Cys Tyr Cys Ser Val Leu Leu Arg
 755 760 765
 Gly Leu Leu Ala Ala Leu Leu Gly His Trp Glu Ala Ser Arg His Pro
 770 775 780
 Asp Thr Thr His Ser Pro Trp His Leu Glu Ala Ser Cys Thr Leu Val
 785 790 795 800
 Ala Val Met Ala Glu Gly Ser Leu Leu Pro Pro Ala Leu Gly Asn Met
 805 810 815
 His Glu Val Phe Ser Gln Leu Ala Pro Phe Glu Val Arg Leu Leu Leu
 820 825 830
 Leu Ser Val Trp Gly Phe Leu Arg Glu His Gly Pro Leu Pro Gln Lys
 835 840 845
 Phe Ile Phe Gln Ser Glu Arg Gly Arg Phe Ile Arg Asp Phe Ser Arg
 850 855 860
 Glu Gly Gly Gly Glu Gly Gly Pro His Leu Ala Val Leu His Ser Val
 865 870 875 880
 Leu His Arg Asn Ile Asp Arg Leu Gly Leu Phe Ser Gly Arg Phe Gln
 885 890 895
 Ala Pro Ser Pro Ser Thr Leu Leu Arg Gln Gly Thr
 900 905

2003295784 19 Aug 2005

<210> 18
 <211> 833
 <212> PRT
 <213> Homo sapiens

<400> 18

Pro Arg Val Arg Asp Ile Tyr Val Gln Cys Leu Ser Ala Leu Ile Gly
 1 5 10 15

Ser Cys Pro Asp Ala Cys Val Asp Ala Leu Leu Asp Thr Ser Val Gln
 20 25 30

His Ser Pro His Phe Asp Trp Val Val Ala His Ile Gly Ser Ser Phe
 35 40 45

Pro Gly Thr Ile Ile Ser Arg Val Leu Ser Cys Gly Leu Lys Asp Phe
 50 55 60

Cys Val His Gly Gly Ala Gly Gly Gly Ala Gly Ser Ser Gly Gly Ser
 65 70 75 80

Ser Ser Gln Thr Pro Ser Thr Asp Pro Phe Pro Gly Ser Pro Ala Ile
 85 90 95

Pro Ala Glu Lys Arg Val Pro Lys Ile Ala Ser Val Val Gly Ile Leu
 100 105 110

Gly His Leu Ala Ser Arg His Gly Asp Ser Ile Arg Arg Glu Leu Leu
 115 120 125

Arg Met Phe His Asp Ser Leu Ala Gly Gly Ser Gly Gly Arg Ser Gly
 130 135 140

Asp Pro Ser Leu Gln Ala Thr Val Pro Phe Leu Leu Gln Leu Ala Val
 145 150 155 160

Met Ser Pro Ala Leu Leu Gly Thr Val Ser Gly Glu Leu Val Asp Cys
 165 170 175

Leu Lys Pro Pro Ala Val Leu Ser Gln Leu Gln Gln His Leu Gln Gly
 180 185 190

Phe Pro Arg Glu Glu Leu Asp Asn Met Leu Asn Leu Ala Val His Leu
 195 200 205

Val Ser Gln Ala Ser Gly Ala Gly Ala Tyr Arg Leu Leu Gln Phe Leu
 210 215 220

Val Asp Thr Ala Met Pro Ala Ser Val Ile Thr Thr Gln Gly Leu Ala
 225 230 235 240

Val Pro Asp Thr Val Arg Glu Ala Cys Asp Arg Leu Ile Gln Leu Leu
 245 250 255

Leu Leu His Leu Gln Lys Leu Val His His Arg Gly Gly Ser Pro Gly
 260 265 270

Glu Gly Val Leu Gly Pro Pro Pro Pro Pro Arg Leu Val Pro Phe Leu
 275 280 285

Asp Ala Leu Lys Asn His Val Gly Glu Leu Cys Gly Glu Thr Leu Arg

2003295784 19 Aug 2005

290	295	300
Leu Glu Arg Lys Arg Phe 305 310	Leu Trp Gln His 315	Gln Leu Leu Gly Leu Leu 320
Ser Val Tyr Thr Arg Pro Ser Cys Gly Pro Glu Ala Leu Gly His Leu 325 330 335		
Leu Ser Arg Ala Arg Ser Pro Glu Glu Leu Ser Leu Ala Thr Gln Leu 340 345 350		
Tyr Ala Gly Leu Val Val Ser Leu Ser Gly Leu Leu Pro Leu Ala Phe 355 360 365		
Arg Ser Cys Leu Ala Arg Val His Ala Gly Thr Leu Gln Pro Pro Phe 370 375 380		
Thr Ala Arg Phe Leu Arg Asn Leu Ala Leu Leu Val Gly Trp Glu Gln 385 390 395 400		
Gln Gly Gly Glu Gly Pro Ala Ala Leu Gly Ala His Phe Gly Glu Ser 405 410 415		
Ala Ser Ala His Leu Ser Asp Leu Ala Pro Leu Leu Leu His Pro Glu 420 425 430		
Glu Glu Val Ala Glu Ala Ala Ala Ser Leu Leu Ala Ile Cys Pro Phe 435 440 445		
Pro Ser Glu Ala Leu Ser Pro Ser Gln Leu Leu Gly Leu Val Arg Ala 450 455 460		
Gly Val His Arg Phe Phe Ala Ser Leu Arg Leu His Gly Pro Pro Gly 465 470 475 480		
Val Ala Ser Ala Cys Gln Leu Leu Thr Arg Leu Ser Gln Thr Ser Pro 485 490 495		
Ala Gly Leu Lys Ala Val Leu Gln Leu Leu Val Glu Gly Ala Leu His 500 505 510		
Arg Gly Asn Thr Glu Leu Phe Gly Gly Gln Val Asp Gly Asp Asn Glu 515 520 525		
Thr Leu Ser Val Val Ser Ala Ser Leu Ala Ser Ala Ser Leu Leu Asp 530 535 540		
Thr Asn Arg Arg His Thr Ala Ala Val Pro Gly Pro Gly Gly Ile Trp 545 550 555 560		
Ser Val Phe His Ala Gly Val Ile Gly Arg Gly Leu Lys Pro Pro Lys 565 570 575		
Phe Val Gln Ser Arg Asn Gln Gln Glu Val Ile Tyr Asn Thr Gln Ser 580 585 590		
Leu Leu Ser Leu Leu Val His Cys Cys Ser Ala Pro Gly Gly Thr Glu 595 600 605		
Cys Gly Glu Cys Trp Gly Ala Pro Ile Leu Ser Pro Glu Ala Ala Lys 610 615 620		

2003295784 19 Aug 2005

Ala Val Ala Val Thr Leu Val Glu Ser Val Cys Pro Asp Ala Ala Gly
625 630 635 640

Ala Glu Leu Ala Trp Pro Pro Glu Glu His Ala Arg Ala Thr Val Glu
645 650 655

Arg Asp Leu Arg Ile Gly Arg Arg Phe Arg Glu Gln Pro Leu Leu Phe
660 665 670

Glu Leu Leu Lys Leu Val Ala Ala Ala Pro Pro Ala Leu Cys Tyr Cys
675 680 685

Ser Val Leu Leu Arg Gly Leu Leu Ala Ala Leu Leu Gly His Trp Glu
690 695 700

Ala Ser Arg His Pro Asp Thr Thr His Ser Pro Trp His Leu Glu Ala
705 710 715 720

Ser Cys Thr Leu Val Ala Val Met Ala Glu Gly Ser Leu Leu Pro Pro
725 730 735

Ala Leu Gly Asn Met His Glu Val Phe Ser Gln Leu Ala Pro Phe Glu
740 745 750

Val Arg Leu Leu Leu Leu Ser Val Trp Gly Phe Leu Arg Glu His Gly
755 760 765

Pro Leu Pro Gln Lys Phe Ile Phe Gln Ser Glu Arg Gly Arg Phe Ile
770 775 780

Arg Asp Phe Ser Arg Glu Gly Gly Gly Glu Gly Gly Pro His Leu Ala
785 790 795 800

Val Leu His Ser Val Leu His Arg Asn Ile Asp Arg Leu Gly Leu Phe
805 810 815

Ser Gly Arg Phe Gln Ala Pro Ser Pro Ser Thr Leu Leu Arg Gln Gly
820 825 830

Thr

<210> 19
 <211> 963
 <212> PRT
 <213> Mus musculus

<400> 19

Met Ile Leu Met Ile Thr Leu Phe Thr Thr Ala Thr Phe Leu Val Leu
1 5 10 15

Gly Val Ser Val Trp Val Leu Ile Lys Glu Ile Leu Thr Val His Val
20 25 30

Pro Pro Pro Ile Pro Gln Arg Val Lys Phe His Met Leu His Tyr Phe
35 40 45

Phe Gln Leu Thr Ile Ala Leu Gly Asn Val Leu Glu Lys Met Lys Ile
50 55 60

2003295784 19 Aug 2005

Cys 65	Pro	Met	Pro	Arg	Phe 70	Phe	Cys	Phe	Ile	Gln 75	Asp	Leu	Leu	Val	Ser 80
Lys	Asn	Asn	Phe	Gly 85	Val	Leu	Val	Lys	Asn 90	Met	His	Phe	Gly	Thr 95	Ile
Pro	Val	Arg	Leu 100	Phe	Gln	Pro	Lys	Ala 105	Thr	Ser	Ser	Gly	Pro	Arg	Lys
Gly	Ile	Ile	Phe	Tyr	His	Gly	Gly	Gly	Gly	Val	Phe	Gly	Ser	Leu	Asp
		115						120					125		
Ser	Tyr	His	Asn	Thr	Cys	Ser	Tyr	Leu	Ala	His	Glu	Thr	Asp	Ser	Val
	130					135					140				
Val	Met	Ala	Val	Gly	Tyr	Arg	Lys	Leu	Pro	Asp	His	His	His	Pro	Thr
145					150					155					160
Ala	Tyr	His	Asp	Cys	Leu	Asn	Ala	Thr	Val	His	Phe	Leu	Lys	Glu	Leu
				165					170					175	
Lys	Thr	Tyr	Gly	Val	Asp	Pro	Ala	Arg	Val	Val	Val	Ser	Gly	Glu	Ser
			180					185					190		
Ile	Gly	Ala	Gly	Ala	Ala	Ala	Ile	Ile	Ala	Gln	Val	Val	Leu	Ala	Arg
		195					200						205		
Lys	Asp	Leu	Pro	Gln	Phe	Arg	Ala	Gln	Val	Leu	Ile	Asn	Pro	Val	Val
	210					215						220			
Gln	Gly	Val	Asn	Phe	Gln	Leu	Pro	Ser	Tyr	Gln	Gln	Tyr	Ser	Asp	Val
225					230					235					240
Pro	Phe	Leu	Ser	Arg	Lys	Phe	Leu	Met	Thr	Cys	Ala	Cys	Lys	Tyr	Leu
				245					250					255	
Ala	Ile	Asp	Gln	Ser	Trp	Lys	Asp	Ala	Met	Leu	Lys	Gly	Thr	Phe	Ile
			260					265					270		
Pro	Pro	Asp	His	Trp	Lys	Lys	Tyr	Ala	Lys	Trp	Leu	Ser	Ser	Asp	Asn
		275					280					285			
Ile	Pro	Gln	Arg	Phe	Lys	Ser	Gln	Gly	Arg	Gln	Pro	Glu	Phe	Pro	Gly
	290					295					300				
Pro	Phe	Asn	Glu	Ser	Ala	Tyr	Leu	Glu	Thr	Asn	His	Ile	Phe	Ser	Leu
305					310					315					320
Glu	Thr	Ser	Pro	Leu	Leu	Ala	Asp	Asp	Lys	Ile	Ile	Ala	Gln	Leu	Pro
				325					330					335	
Glu	Thr	Phe	Leu	Val	Ser	Ser	Glu	Tyr	Asp	Val	Leu	Arg	Asp	Asp	Thr
			340					345					350		
Leu	Leu	Tyr	Lys	Lys	Arg	Leu	Glu	Glu	Gln	Gly	Val	Pro	Val	Thr	Trp
		355					360					365			
Leu	Trp	Val	Gly	Leu	Pro	Asp	Val	Arg	Val	Val	Pro	Leu	Ser	Gln	Gly
	370					375					380				
Pro	Arg	Ala	Pro	Gly	Pro	Pro	Gly	Pro	Ala	Pro	Ala	Thr	His	Gly	Pro
385					390					395					400

2003295784 19 Aug 2005

Val	Pro	Leu	Ser	Ala	Gln	Glu	Leu	Ser	Gln	Glu	Ile	Lys	Ala	Phe	Leu	405	410	415	
Thr	Gly	Val	Asp	Pro	Ile	Leu	Gly	His	Gln	Leu	Ser	Ala	Arg	Glu	His	420	425	430	
Ala	Gln	Cys	Gly	Leu	Leu	Leu	Leu	Arg	Ser	Leu	Pro	Pro	Ala	Gln	Ala	435	440	445	
Ala	Val	Leu	Asp	His	Leu	Arg	Gly	Val	Phe	Asp	Glu	Ser	Val	Gln	Ala	450	455	460	
His	Leu	Ala	Ala	Leu	Glu	Glu	Ser	Pro	Val	Ala	Gly	Pro	Pro	His	Leu	465	470	475	480
Arg	Pro	Pro	Ser	Pro	Ser	His	Val	Pro	Thr	Gly	Gly	Pro	Gly	Leu	Glu	485	490	495	
Asp	Val	Val	His	Glu	Val	Gln	Gln	Val	Leu	Cys	Glu	Phe	Ile	Arg	Ala	500	505	510	
Asn	Pro	Lys	Val	Trp	Ala	Pro	Val	Ile	Ser	Ala	Trp	Ser	Ile	Asp	Leu	515	520	525	
Met	Gly	Gln	Leu	Ser	Ser	Thr	Tyr	Ser	Gly	Gln	His	Gln	Arg	Val	Pro	530	535	540	
His	Ala	Thr	Gly	Ser	Arg	Asn	Glu	Leu	Leu	Gln	Leu	Trp	Met	Ser	Cys	545	550	555	560
Arg	Asp	Thr	Arg	Thr	Leu	Met	Asp	Ile	Tyr	Val	Gln	Cys	Leu	Ser	Ala	565	570	575	
Leu	Ile	Gly	Ser	Cys	Pro	Asp	Ala	Tyr	Ser	Phe	Pro	Gly	Phe	Pro	Ala	580	585	590	
Ile	Pro	Gly	Glu	Lys	Arg	Val	Pro	Lys	Ile	Ala	Ser	Ala	Val	Gly	Ile	595	600	605	
Gln	Val	Thr	Trp	Leu	Ser	Ala	Met	Glu	Thr	Ala	Ser	Asp	Gly	Asn	Cys	610	615	620	
Cys	Ala	Cys	Phe	Met	Ile	Val	Trp	Gln	Arg	Phe	Leu	Val	Asp	Thr	Ala	625	630	635	640
Met	Pro	Ala	Ala	Val	Ile	Thr	Thr	Gln	Gly	Leu	Ala	Val	Pro	Asp	Thr	645	650	655	
Met	Arg	Glu	Ala	Tyr	Asp	Arg	Leu	Ile	Gln	Leu	Leu	Leu	Leu	His	Leu	660	665	670	
Gln	Lys	Leu	Val	His	His	Arg	Gly	Gly	Ala	Pro	Gly	Glu	Gly	Val	Leu	675	680	685	
Gly	Pro	Pro	Ser	Pro	Pro	Leu	Pro	Val	Pro	Phe	Leu	Asp	Ala	Leu	Arg	690	695	700	
Asn	His	Val	Gly	Glu	Leu	Cys	Gly	Lys	Thr	Leu	Arg	Leu	Glu	Arg	Lys	705	710	715	720
Arg	Phe	Leu	Trp	Gln	His	Gln	Leu	Leu	Ala	Tyr	Ser	Trp	Phe	Leu	Arg				

2003295784 19 Aug 2005

	725	730	735
Lys Leu Ala Leu Leu Val Gly Trp Glu Gln Gln Gly Asp Glu Gly Pro	740	745	750
Ser Ala Leu Gly Ala Arg Phe Gly Glu Ser Ala Ser Ala His Leu Ser	755	760	765
Asp Leu Ala Pro Leu Leu Leu His Pro Glu Glu Glu Val Ala Glu Ala	770	775	780
Ala Ala Ser Leu Leu Ala Val Cys Pro Phe Pro Ser Glu Ala Leu Ser	785	790	800
Pro Ser Gln Leu Leu Gly Leu Val Arg Ala Gly Val His His Phe Phe	805	810	815
Asn Ser Leu Arg Leu His Gly Pro Pro Gly Val Ala Ser Ala Ser Gln	820	825	830
Leu Leu Thr Arg Leu Ser Gln Thr Ser Pro Ala Gly Leu Lys Ala Val	835	840	845
Leu Gln Leu Leu Val Glu Val Ala Leu His Arg Gly Asn Thr Glu Leu	850	855	860
Phe Gly Glu Glu Met Val Gly Asp Asn Glu Thr Leu Ser Ile Val Ser	865	870	875
Thr Pro Leu Ala Ser Ala Ser Leu Leu Asp Ile Asn Arg Arg His Thr	885	890	895
Ala Ala Val Pro Gly Pro Gly Gly Ile Trp Ser Val Phe His Ala Gly	900	905	910
Val Ile Gly Arg Gly Leu Lys Ser Pro Lys Ile Val Gln Ser Arg Asn	915	920	925
His Gln Glu Val Ile Tyr Asn Thr Gln Ser Leu Ile Ser Leu Leu Val	930	935	940
His Cys Cys Ser Ala Ser Gly Ser Ser Glu His Lys Gly Tyr Trp Gly	945	950	955
			960

Ala Pro Thr

<210> 20
 <211> 900
 <212> PRT
 <213> Anopheles gambiae

<400> 20

Lys Asn Leu Pro Asp Pro Ser Val Asp Asp Glu Ala Val Gln Glu Ile	1	5	10	15
---	---	---	----	----

His Glu Ala Leu Glu Arg Leu Val Thr Val Gly Pro Thr Ala Trp Cys	20	25	30
---	----	----	----

Pro Val Ile Ser Ser Trp Cys Leu Lys Leu Leu Gly Glu Val Cys Lys

	35					40					45				
Lys	His	Cys	Arg	Arg	Arg	Pro	Pro	Asp	Ile	Arg	Gly	Ala	Cys	Asn	Leu
	50					55					60				
Trp	Leu	Gly	Cys	Ser	Ala	Ile	Arg	Tyr	Leu	Leu	Ser	Leu	Ser	Ala	Leu
65					70					75					80
Cys	Phe	Glu	Lys	Leu	Asp	Gln	Arg	Glu	Met	Asp	Glu	Cys	Ile	Asn	Glu
				85					90					95	
Met	Leu	Val	Ile	Tyr	Gly	Thr	His	Thr	Pro	Phe	Phe	Asp	Trp	Val	Val
			100					105					110		
Ala	Arg	Leu	Gly	Gly	Cys	Phe	Pro	Leu	Arg	Val	Met	Ser	Ser	Met	Leu
		115					120					125			
Ser	Met	Gly	Val	Ala	Arg	Phe	Thr	Gly	Asp	Phe	Asp	Gln	Pro	Ser	Glu
	130					135					140				
Ser	Glu	Val	Glu	Val	Leu	Ser	Tyr	Leu	Gly	Leu	Ala	His	Glu	Ser	Asp
145					150					155					160
Leu	Arg	Lys	Ala	Leu	Lys	Ser	Thr	Leu	Glu	His	Val	Ala	Ser	Tyr	Lys
				165					170					175	
Gln	Pro	Ile	Pro	Tyr	Leu	Leu	Met	Leu	Ala	Lys	Ala	Ser	Glu	Thr	Ile
			180					185					190		
Ser	Gln	Ala	Leu	Val	Ala	Val	Phe	Leu	Glu	Leu	His	Asp	Glu	Asn	Arg
		195					200					205			
Leu	Pro	Thr	Leu	Thr	Val	Leu	Pro	Lys	Asn	Trp	Pro	Ala	Asn	Ile	Gly
	210					215					220				
Leu	Pro	Tyr	Val	Leu	His	Thr	Val	Ala	Gly	Leu	Leu	Leu	Lys	Met	Lys
225					230					235					240
Lys	His	Ala	Ile	Arg	Val	Thr	Leu	Ile	Leu	Ala	Lys	Met	Ser	Thr	Gln
				245					250					255	
His	Ser	Trp	Cys	Gln	Glu	Leu	Leu	Glu	Met	Met	Phe	Ile	Glu	Leu	Glu
			260					265					270		
Thr	Leu	Val	Leu	Asp	Lys	His	Thr	Ala	Ala	Leu	Leu	Glu	Asp	Ile	Ile
		275					280					285			
Arg	Asp	Gly	Met	Arg	Glu	Met	Leu	Trp	Asn	Ser	Cys	Thr	Ser	Asp	Val
	290					295					300				
Pro	Tyr	Leu	Gln	Gln	Val	Ala	Val	Arg	Leu	Ile	Leu	Leu	Ala	Ser	Phe
305					310					315					320
Lys	Ser	Asn	Ser	Val	Phe	His	Gln	Thr	Ile	Val	Tyr	Leu	Leu	Ser	Val
				325					330					335	
Ser	Glu	Pro	Ala	Leu	Ala	Val	Ser	Thr	Lys	Pro	His	Leu	Asn	Ala	Leu
			340					345					350		
Val	Arg	Val	Leu	Gly	Gly	Pro	His	Gly	Thr	Val	Asp	Val	Pro	Lys	Val
		355					360					365			

2003295784 19 Aug 2005

Lys	Pro	Ala	Phe	Glu	Thr	Ala	Phe	Glu	Lys	Ile	Leu	Ile	Ser	Pro	Cys	
370						375					380					
Lys	Arg	Val	Glu	Cys	Trp	Asn	Ile	Leu	His	Asn	Leu	Val	Glu	Leu	Leu	
385					390					395					400	
Lys	Leu	Glu	Arg	Thr	Ala	Ile	Leu	Gly	Ser	Thr	Leu	Arg	Lys	Val	Asn	
				405					410					415		
Cys	Thr	Gly	Met	Met	His	Glu	Leu	Leu	Asp	Arg	Val	Leu	Lys	Ile	Trp	
			420					425					430			
Glu	Asn	Phe	Met	Ser	Arg	Glu	Arg	Gln	Asp	Asp	Thr	Gln	Gly	Gly	Gly	
		435						440					445			
Cys	Thr	Val	Arg	Ala	Thr	Gln	Glu	Val	Gln	Glu	Asn	Ala	Asn	Glu	Pro	
	450					455					460					
Gly	Lys	Arg	Val	Lys	Asn	Glu	Arg	Pro	Glu	Pro	Met	Glu	Thr	Asp	Glu	
465					470					475					480	
Gln	His	Arg	Leu	Gly	Thr	Gly	Ser	Arg	Thr	Val	Thr	Tyr	Lys	Asp	Leu	
			485					490						495		
Ile	His	Glu	Thr	Val	Arg	Leu	Ile	Glu	Cys	Met	Asp	Leu	Gly	Lys	Ser	
			500					505					510			
Val	Thr	Ile	Gly	Thr	Ala	Gln	Thr	Leu	Lys	Leu	Ser	Gln	Leu	Leu	Val	
		515					520						525			
Lys	Tyr	Phe	Phe	Tyr	Cys	Leu	Lys	Leu	Ser	Thr	Ala	Gly	Thr	Ser	Val	
	530					535					540					
Pro	Ser	Gly	Thr	Val	Pro	Glu	Ser	Leu	Asp	Glu	Ser	Leu	Asn	Arg	Val	
545					550					555					560	
Tyr	Ser	Leu	Leu	Ser	Lys	His	Cys	Gly	His	Arg	Lys	Ala	Ala	Arg	Thr	
				565					570					575		
Ala	Ala	Leu	Arg	Glu	Leu	Leu	Glu	Gly	Ala	Leu	Phe	Leu	Tyr	Gly	Asp	
			580					585					590			
Leu	Phe	Gly	Ser	Gln	Ala	Glu	Ser	Gln	Ala	Tyr	Ser	Phe	Asp	Lys	Pro	
		595						600					605			
Asp	Asp	Leu	Leu	Ile	Arg	Leu	Asn	Gln	Lys	Gln	Gly	Ile	Ala	Leu	Asn	
	610					615					620					
Ala	Ser	Arg	Ala	Thr	Val	Leu	His	Ala	Gly	Ile	Ile	Gly	Gln	Gly	Pro	
625					630					635					640	
Lys	Ile	Pro	Ser	Lys	Lys	Ala	Val	Gly	Pro	Ala	Ser	Glu	Met	Gln	Asn	
				645					650					655		
His	Leu	Leu	Asn	Ala	Ile	Val	Ala	Cys	Cys	Gln	Asp	Val	Asn	Asp	His	
			660					665					670			
Gln	Ala	Thr	Ile	Asp	Gly	Phe	Ser	Tyr	Val	Ser	Leu	Leu	Leu	Val	Glu	
		675					680					685				
Met	Ile	Ser	Pro	Asp	Val	Met	Tyr	Asn	Gly	Leu	Pro	Trp	Pro	Glu	Glu	
						695					700					

2003295784 19 Aug 2005

```

Asp Phe Ile Arg Val Thr Met Glu Arg Asp Leu Gln Ile Gln Arg Thr
705                               710                               715                               720

Phe Arg His Ser Pro Ile Leu Trp Ser Ile Leu Gly Leu Val Ala Cys
              . 725                               730                               735

Tyr Arg Pro Ser Leu Cys Tyr Cys Ser Val Leu Leu Arg Ala Leu Cys
              740                               745                               750

Ala Ser Ala Leu His Gln Trp Arg Ser Lys Thr Ala Glu Thr Leu Asn
              755                               760                               765

Gly Gln Lys Thr Asp Leu Leu Tyr Met Thr Thr Lys Leu Leu Glu Leu
              770                               775                               780

Met Ala Leu Ala Gln Leu Leu Pro Pro Pro Leu Ser Tyr Leu His Ile
785                               790                               795                               800

Val Leu Glu Tyr Phe Asp Gly Pro Glu Ile Ala Tyr Val Leu Lys Glu
              805                               810                               815

Cys Val Trp Asn Tyr Met Lys Asp His Val Pro Ser Pro Val Leu Phe
              820                               825                               830

Val Cys Asp Pro Thr Gly Phe His Trp Arg Asp Pro Leu Thr Ser Arg
              835                               840                               845

Pro Pro Leu Gln Tyr Thr Asn Pro Leu Arg Asn Thr Met Gln Lys Lys
              850                               855                               860

Leu Thr Lys Val Gly His Leu Tyr His Gln Met Phe Val Gly Pro Glu
865                               870                               875                               880

Leu Arg Asn Pro Ser Ala Ser Asn Ser Gly Gln Pro Thr Gln Gln Pro
              885                               890                               895

Leu Val Gln Gly
              900

```

```

<210> 21
<211> 27
<212> PRT
<213> Homo sapiens

```

```

<400> 21

```

```

Met Ser Ala Leu Cys Asp Pro Pro Gly Ala Pro Gly Pro Pro Gly Pro
1              5              10              15

Ala Pro Ala Thr His Gly Pro Ala Pro Leu Ser
              20              25

```