

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 221 918 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:

16.03.2005 Bulletin 2005/11

(21) Application number: 00968673.4

(22) Date of filing: 04.10.2000

(51) Int Cl.7: A61F 9/00

(86) International application number:
PCT/US2000/027367

(87) International publication number:
WO 2001/028473 (26.04.2001 Gazette 2001/17)

(54) SUB-TENON DRUG DELIVERY

MEDIKAMENTENVERSORGUNG DER SUB-TENON

ADMINISTRATION SOUS-TENDINEUSE DE MEDICAMENTS

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

(30) Priority: 21.10.1999 US 161660 P

(43) Date of publication of application:
17.07.2002 Bulletin 2002/29

(60) Divisional application:
05100176.6

(73) Proprietor: Alcon Inc.
6331 Hünenberg (CH)

(72) Inventors:

- CLARK, Abbot, F.
Arlington, TX 76017 (US)
- DAHLIN, David, C.
Arlington, TX 76016 (US)

- MARSH, David Allen
Fort Worth, TX 76132 (US)
- STRUBLE, Craig, B.
Arlington, TX 76006 (US)
- YAACOBI, Yoseph
Forth Worth, TX 76132 (US)
- YORK, Billie M.
Conroe, TX 77304 (US)

(74) Representative: Hanna, Peter William Derek et al
Hanna, Moore & Curley,
11 Mespil Road,
Dublin 4 (IE)

(56) References cited:

WO-A-93/20784	US-A- 4 759 746
US-A- 5 167 618	US-A- 5 665 069
US-A- 5 817 075	US-A- 5 824 073

EP 1 221 918 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description**Field of the Invention**

[0001] The present invention generally pertains to the delivery of ophthalmically acceptable pharmaceutically active agents to the back of the eye. More particularly, but not by way of limitation, the present invention pertains to apparatus for sub-Tenon delivery of a drug depot to the posterior segment of a human eye proximate the macula.

Description of the Related Art

[0002] Several diseases and conditions of the posterior segment of the eye threaten vision. Age related macular degeneration (ARMD), choroidal neovascularization (CNV), retinopathies (e.g., diabetic retinopathy, vitreoretinopathy), retinitis (e.g., cytomegalovirus (CMV) retinitis), uveitis, macular edema, and glaucoma are several examples.

[0003] Age related macular degeneration (ARMD) is the leading cause of blindness in the elderly. ARMD attacks the center of vision and blurs it, making reading, driving, and other detailed tasks difficult or impossible. About 200,000 new cases of ARMD occur each year in the United States alone. Current estimates reveal that approximately forty percent of the population over age 75, and approximately twenty percent of the population over age 60, suffer from some degree of macular degeneration. "Wet" ARMD is the type of ARMD that most often causes blindness. In wet ARMD, newly formed choroidal blood vessels (choroidal neovascularization (CNV)) leak fluid and cause progressive damage to the retina.

[0004] In the particular case of CNV in ARMD, two main methods of treatment are currently being developed, (a) photocoagulation and (b) the use of angiogenesis inhibitors. However, photocoagulation can be harmful to the retina and is impractical when the CNV is in proximity of the fovea. Furthermore, photocoagulation often results in recurrent CNV over time. Oral administration of anti-angiogenic compounds is also being tested as a systemic treatment for ARMD. However, due to drug-specific metabolic restrictions, systemic administration usually provides sub-therapeutic drug levels to the eye. Therefore, to achieve effective intraocular drug concentrations, either an unacceptably high dose or repetitive conventional doses are required. Various implants have also been developed for delivery of anti-angiogenic compounds locally to the eye. Examples of such implants are disclosed in U.S. Patent Nos. 5,824,072 to Wong, U.S. Patent No. 5,476,511 to Gwon et al., and U.S. Patent No. 5,773,019 to Ashton et al.

[0005] In addition, it is known to use a straight, 5/8 inch long, 25 gauge needle to perform sub-Tenon injection of corticosteroids for the treatment of posterior uveitis or macular edema associated with uveitis or anterior

segment surgery. See Uveitis: A Clinical Approach to Diagnosis and Management (Second Edition), Ronald E. Smith and Robert A. Nozik, 1989, pp. 63-68; "Echographic Localization of Corticosteroids After Periocular Injection", William R. Freeman, Ronald L. Green, and Ronald E. Smith, American Journal of Ophthalmology 103:281-288, March 1987. In such methods, a physician attempts to dispose the tip of the needle near the macula but without penetrating the posterior ciliary arteries or

the optic nerve. However, because the physician cannot see the tip, as well as movement of the eyeball within the orbit due to contact with the straight needle, it is very difficult to precisely place the tip at the desired location near the macula. For the same reasons, it is also very difficult to determine whether the tip is correctly positioned below the Tenon's capsule. Such methods do not insure a consistent delivery of a specific quantity of drug to a region over the macula. In fact, the literature reports that only about 57 percent of injections using this method result in drug being placed in the sub-Tenon space overlying the macular area. "Echographic Localization of Corticosteroids After Periocular Injection", pp. 283-285. In addition, moving a straight needle along the curved surface of the sclera causes "tenting" or stretching of the overlying Tenon's capsule. Such movement may cause penetration of the Tenon's capsule, allowing drug to be injected into surrounding tissues. Furthermore, such movement may also cause inadvertent penetration of the sclera, resulting in injection of drug into the vitreous cavity. More importantly, penetration of the sclera may result in significant damage to the eye or even a loss of sight. Documented complications of such penetrations include orbital hemorrhage, central retinal vein occlusion, and central retinal artery occlusion.

[0006] Referring to FIG. 6, it is also known to use a blunt 19 gage cannula 200 having a hub 201, a straight proximal portion 202, and an angled distal portion 204 to perform sub-Tenon injection of anesthesia for cataract and vitreoretinal surgery. See "Local Anesthesia for Vitreoretinal Surgery", Calvin E. Mein and Michael G. Woodcock, Retina 10: 47-49, 1990; "Ocular Anesthesia for Cataract Surgery: A Direct Sub-Tenon's Approach", Ophthalmic Surgery 21:696-699, 1990; "Single Quadrant Sub-Tenon's Block: Evaluation of a New Local Anaesthetic Technique for Eye Surgery", Anaesthesia and Intensive Care 24: 241-244, April 1996. However, such cannulae also suffer from the above-described "tenting" and penetration problems if used to deliver drugs into the sub-Tenon's space above the macula.

[0007] It is also known to use a gently curved cannula 210 as shown in FIG. 7 to perform sub-Tenon injection of anesthesia for cataract surgery. See "Curved, Sub-Tenon Cannula for Local Anesthesia", Julian D. Stevens, Ophthalmic Surgery 24:121-122, February 1993. However, this cannula also suffers from the above-described "tenting" and penetration problems if used to deliver drugs into the sub-Tenon's space above the macula.

[0008] It is also known to use a 24 gauge cannula that has a straight proximal portion and a curved distal portion that is disposed at a 90 degree angle to the straight portion to inject a local anesthetic solution below the Tenon's capsule. The straight portion has a length of 5 mm. The curved portion has a radius of curvature of 14 mm and an arc length of 27 mm. See "A Modified Sub-Tenon's Cannula for Local Anesthesia", P. Muthusamy and Richard F. Hommersom, Asia-Pacific Journal of Ophthalmology, Volume 8, No. 3 (July 1996). However, because of its geometry, this cannula is not suitable for the delivery of drugs in the form of suspensions, emulsions, ointments, or gels, or drugs in such forms including bioerodible polymers or non-bioerodible polymers.

[0009] Document US-A-5 817 075 discloses a cannula comprising a straight proximal portion which curves distally into a curved arc with an aperture at the tip of the cannula.

[0010] US 4,759,746 is another example of a type of needle that can be used in ocular surgery and is specifically directed towards an instrument and method for administering retro-bulbar (or peri-bulbar) anaesthetics. It comprises a curved needle portion that terminates in a straight needle portion.

[0011] Therefore, a need exists in the field of ophthalmology for improved apparatus and methods for sub-Tenon delivery of a drug depot to the posterior segment of a human eye proximate the macula that do not suffer from the above-described limitations. The improved apparatus and methods should be safe for the patient, easy for the physician to use, capable of delivering a wide spectrum of formulations, and capable of being performed in an outpatient setting.

Summary of the Invention

[0012] The present invention provides a cannula as defined in claim 1.

[0013] One aspect of the present invention is that the cannula includes a distal portion having a radius of curvature substantially equal to a radius of curvature of a globe of the human eye, a proximal portion, and a bend separating the distal portion and the proximal portion. A tangent of the distal portion at the bend is disposed at an angle of no more than about 56 degrees with respect to the proximal portion.

[0014] The present invention allows the delivery of a drug to the human eye. A cannula is inserted below the Tenon's capsule and above the sclera of the human eye at a point posterior to a limbus of the eye. The cannula includes a distal portion having a radius of curvature substantially equal to a radius of curvature of the globe of the human eye. A drug is injected through the cannula to form a drug depot on an outer surface of the sclera. The drug comprises a pharmaceutically active agent selected from the group consisting of 4,9(11)-Pregnadien-17 α ,21-diol-3,20-dione and 4,9(11)-Pregnadien-17 α ,21-diol-3,20-dione-21-acetate.

[0015] In another preferred aspect, the present invention comprises a cannula including a hub for removably coupling to a syringe, a proximal portion, and a distal portion. The distal portion has a tip, an orifice proximate the tip, and a radius of curvature substantially equal to a radius of curvature of the globe of the human eye. The part of the distal portion proximate the tip comprises a plastic.

10 Brief Description of the Drawings

[0016] For a more complete understanding of the present invention, and for further objects and advantages thereof, reference is made to the following description taken in conjunction with the accompanying drawings in which:

20 FIG. 1 is a side sectional view schematically illustrating the human eye and a drug depot according to a preferred embodiment of the present invention; FIG. 2 is a side, partially sectional view schematically illustrating a cannula for creating the drug depot of FIG. 1 according to a preferred embodiment of the present invention;

25 FIGS. 3A through 3E are side views schematically illustrating additional variations of the cannula of FIG. 2;

30 FIG. 4 is a side view schematically illustrating a variation of the cannula of FIG. 2 having dual lumens; FIG. 5 is a side view schematically illustrating a variation of the cannula of FIG. 2 having dual lumens; FIG. 6 is a side view schematically illustrating a first conventional cannula; and

35 FIG. 7 is a side view schematically illustrating a second conventional cannula.

Detailed Description of the Preferred Embodiments

[0017] The preferred embodiments of the present invention and their advantages are best understood by referring to FIGS. 1 through 5 of the drawings, like numerals being used for like and corresponding parts of the various drawings.

[0018] FIG. 1 schematically illustrates a human eye 10. Eye 10 has a cornea 12, a lens 14, a sclera 16, a choroid 18, a retina 20, and an optic nerve 22. An anterior segment 24 of eye 10 generally includes the portions of eye 10 anterior of line 25. A posterior segment 26 of eye 10 generally includes the portions of eye 10 posterior of line 25. Retina 20 is physically attached to choroid 18 in a circumferential manner proximate pars plana 28. Retina 20 has a macula 30 located slightly lateral to optic nerve 22. As is well known in the ophthalmic art, macula 30 is comprised primarily of retinal cones and is the region of maximum visual acuity in retina 20. A Tenon's capsule or Tenon's membrane 34 is disposed on sclera 16. A conjunctiva 36 covers a short area of the globe of eye 10 posterior to limbus 32 (the bulbar conjunctiva)

and folds up (the upper cul-de-sac) or down (the lower cul-de-sac) to cover the inner areas of upper eyelid 35 and lower eyelid 37, respectively. Conjunctiva 36 is disposed on top of Tenon's capsule 34.

[0019] Sclera 16 and Tenon's capsule 34 define the exterior surface of the globe of eye 10. For treatment of ARMD, CNV, retinopathies, retinitis, uveitis, cystoid macular edema (CME), glaucoma, and other diseases or conditions of posterior segment 26, it is preferable to dispose a depot 38 of a specific quantity of an ophthalmically acceptable pharmaceutically active agent directly on the outer surface of sclera 16 and below Tenon's capsule 34. In addition, in cases of ARMD and CME it is most preferable to dispose depot 38 directly on the outer surface of sclera 16, below Tenon's capsule 34, and generally above macula 30. In a study using New Zealand White rabbits, a drug depot of 4,9(11)-Pregnadien-17 α ,21-diol-,20-dione-21-acetate, an angiostatic steroid available from Steraloids, Inc. of Wilton, New Hampshire, was disposed directly on the outer surface of the sclera, below the Tenon's capsule, and slightly posterior of the equator of the rabbit eyes. Such a drug depot resulted in a concentration of the angiostatic steroid, averaged over the entire retina and measured the day after the injection, about ten times greater than a similar concentration delivered by a depot located below the conjunctiva but above the Tenon's capsule of the rabbit eyes. Given the fact that the Tenon's capsule of a New Zealand White rabbit is very thin, these beneficial results are highly unexpected. It is important to note that Tenon's capsule 34 of human eye 10 is also very thin. 4,9(11)-Pregnadien-17 α ,21-diol-3,20-dione-21-acetate, and the related compound 4,9(11)-Pregnadien-17 α ,21-diol-3,20-dione, are more fully described in U.S. Patent Nos. 5,770,592 and 5,679,666.

[0020] Referring now to FIG. 2, a cannula 50 for creating drug depot 38 according to a preferred embodiment of the present invention is schematically illustrated. Cannula 50 generally includes a distal portion 52, a proximal portion 54, and a hub 56. A bend 57 separates distal portion 52 and proximal portion 54. A hollow bore 58 runs axially within distal portion 52 and proximal portion 54 and is fluidly coupled with a hollow bore 60 within hub 56.

[0021] Distal portion 52 preferably has a blunt tip 62 to prevent damage to blood vessels in the periocular tissues and to pass smoothly over sclera 16. An orifice 64 is located proximate tip 62 for delivery of a drug formulation. Orifice 64 is preferably located about 1 mm from tip 62 on the lower or interior side of distal portion 52 to minimize the possibility of connective tissue blockage. Orifice 64 may alternatively be located on the end, on the upper or exterior side, or on other portions of distal portion 52. In addition, distal portion 52 may have multiple orifices, if desired. Orifice 64 is preferably circular and preferably has a 0.025 inch diameter that insures a smooth, controlled delivery of drug. Alternatively, other shapes and sizes of orifice 64 may be used.

[0022] Distal portion 52 and proximal portion 54 are preferably formed out of 19 gauge needle stock. However, other sizes of tubing may be utilized depending on the viscosity and/or volume of material to be injected.

5 Distal portion 52 and proximal portion 54 are preferably made of surgical stainless steel. Other conventional materials such as Teflon, other metals, metal alloys, polyethylene, polypropylene, other conventional plastics, or combinations of the foregoing may also be used. For 10 example, distal portion 52 may be made from a plastic. As another example, a part of distal portion 52 proximate tip 62 may be made from plastic, and the remainder of distal portion 52 and proximal portion 54 may be made from metal. The plastic preferably has sufficient softness and/or flexibility to minimize the possibility of penetration of sclera 16 or Tenon's capsule 34 when cannula 50 is inserted into eye 10, as described hereinbelow. In addition, the length of the plastic portion of distal portion 52, as well as the specific plastic, are preferably 15 selected so that distal portion 52 maintains its radius of curvature B when cannula 50 is inserted into eye 10.

[0023] Hub 56 is for removably coupling to a conventional syringe (not shown). Hub 56 preferably complies with Luer Taper Specification 70.1 of the American 20 Standards Association. Hub 56 preferably includes a locator protuberance 66 that is coplanar with distal portion 52 and proximal portion 54. Protuberance 66 allows a physician to know the orientation of distal portion 52 even when it is inserted below Tenon's capsule 34. Hub 25 56 is preferably made of conventional plastics.

[0024] Referring to both FIGS. 1 and 2, distal portion 52 preferably has an arc length A and a radius of curvature B that closely approximate the curvature of sclera 16 of an adult human eye 10 from insertion points 70a or 70b, each of which is about 5 mm to about 10 mm 30 posterior of limbus 32. Arc length A and radius of curvature B insure that drug depot 38, and more specifically, a specific quantity of pharmaceutically active agent, is deposited on the outer surface of sclera 16 generally above macula 30.

[0025] Arc length A and bend 57 also limit the depth of insertion of cannula 50 along sclera 16, preventing tip 62 from accidentally contacting and damaging posterior ciliary arteries 40 or optic nerve 22. For an adult human 35 eye 10 and for superior temporal or inferior temporal sub-Tenon insertion of cannula 50, arc length A is preferably about 15 mm to about 18 mm. Arc length A may be varied for patient's with smaller or larger than average adult eyes, for pediatric patient's with smaller eyes, or for different insertion points into Tenon's capsule 34. A tangent 72 of distal portion 52 at bend 57 is preferably 40 formed at an angle C with respect to proximal portion 54. In addition to making bend 57 a physical limit to the insertion of cannula 50, angle C also raises the angle of hub 56 so that the face, bridge of the nose, and eyebrows of a patient to not interfere with the attached syringe. Angle C is also important to the successful delivery of drugs in the form of suspensions, emulsions, oint- 45

ments, or gels, or drugs in such forms including bioerodible polymers or non-bioerodible polymers. Angle C is no more than about 56 degrees. Angle C is most preferably about 56 degrees. Proximal portion 54 preferably has a length D of about 15 mm. Other angles and lengths may be used for angle C and length D for specific applications of cannula 50.

[0026] Radius of curvature B insures that distal portion 52 does not drag or put pressure on sclera 16 as cannula 50 is advanced to the proper position, minimizing the risk of scleral penetration. In addition, radius of curvature B eliminates the "tenting" or pulling away of Tenon's capsule 34 from sclera 16, minimizing the risk of penetration into the periocular tissues. For an adult human eye 10 and for superior temporal or inferior temporal sub-Tenon insertion of cannula 50, radius of curvature B is preferably about 11.5 mm to about 14 mm, and most preferably about 12.5 mm. Radius of curvature B may be varied for patients with smaller or larger than average adult eyes, for pediatric patients with smaller eyes, or for different insertion points into Tenon's capsule 34.

[0027] Cannula 50 may be used to inject a wide variety of drug formulations using the following preferred techniques. In a first preferred technique, a physician first anesthetizes eye 10 using conventional topical anesthetic drops. The patient is then instructed to look down and toward his or her nose. Next, the physician uses a 25 gage, 5/8 inch needle to penetrate both conjunctiva 36 and Tenon's capsule 34 at a point about 4 mm posterior to limbus 32 in the superior temporal quadrant of eye 10. The needle is then advanced along the outer surface of sclera 16 to a point about 8 mm to about 9 mm posterior of limbus 32. The physician then makes a small bleb of anesthesia, preferably about 1 mm to about 2 mm long, at this point. The physician then grasps the tissue raised by the bleb with a forceps, and then punctures a hole through conjunctiva 36 and Tenon's capsule 34 using an introducer needle. The introducer needle preferably has an outer diameter with the same gage as cannula 50 or one gage larger than cannula 50. Next, the physician draws a drug formulation into a conventional syringe using a conventional straight needle. The needle is removed and cannula 50 is attached to the syringe. All air is removed from the syringe and cannula 50 so that the drug formulation is at tip 62. The physician then introduces cannula 50 through the hole made by the introducer needle, with orifice 64 facing sclera 16. With distal portion 52 in close contact with the outer surface of sclera 16, cannula 50 is advanced toward the back of the eye until bend 57 is at the site of the hole made by the introducer needle. At this point, tip 62 is preferably located about 5 mm to about 6 mm from the center of optic nerve 22, and about 2 mm to about 3 mm from macula 30. The physician then injects the drug formulation by actuating the syringe plunger, creating drug depot 38 on the outer surface of sclera 16 generally above macula 30. Alternatively, the above-de-

scribed technique may be performed in the inferior temporal quadrant of eye 10, in which case the patient is instructed to look up and toward his or her nose.

[0028] In a second preferred technique, a physician first anesthetizes eye 10 using conventional topical anesthetic drops. Next, the patient is instructed to look down and toward his or her nose. Next, the physician creates a small incision in conjunctiva 36 and Tenon's capsule 34 at a point about 8 mm to about 9 mm posterior to limbus 32 in the superior temporal quadrant of eye 10 using fine scissors. The physician then draws a drug formulation into a conventional syringe, and then attaches cannula 50 to the syringe, as described above. Cannula 50 is then inserted through the incision with orifice 64 facing sclera 16. With distal portion 52 in close contact with the outer surface of sclera 16, cannula 50 is advanced toward the back of the eye until bend 57 is at the site of the incision. At this point, tip 62 is preferably located about 5 mm to about 6 mm from the center of optic nerve 22, and about 2 mm to about 3 mm from macula 30. The physician then injects the drug formulation by actuating the syringe plunger, creating drug depot 38 on the outer surface of sclera 16 generally above macula 30. If necessary, the incision may be sealed around cannula 50 using a purse suture to prevent reflux of the injected drug formulation. Alternatively, the above-described technique may be performed in the inferior temporal quadrant of eye 10, in which case the patient is instructed to look up and toward his or her nose.

[0029] Depending on the physicochemical properties of the drug formulation, drug depot 38 preferably provides controlled release of a pharmaceutically active agent to macula 30 and retina 20 via sclera 16 and choroid 18 for a period of weeks or months. As the use of cannula 50 causes no "tenting" or substantial stretching of Tenon's capsule 34, cannula 50 should result in significantly less trauma to eye 10 than conventional cannulae when repeated injections are required.

[0030] Cannula 50 can be used to deliver a wide variety of drug formulations to treat a wide variety of diseases of posterior segment 26. The drug formulation used to form drug depot 38 may be a solution, a suspension, an emulsion, an ointment, a gel forming solution, a gel, a bioerodible polymer, or a non-bioerodible polymer. The drug formulation used to form drug depot 38 may include one or more ophthalmically acceptable pharmaceutically active agents, and may also include conventional non-active incipients. Examples of pharmaceutically active agents suitable for this drug formulation are anti-infectives, including, without limitation, antibiotics, antivirals, and antifungals; antiallergenic agents and mast cell stabilizers; steroid and non-steroidal anti-inflammatory agents; cyclooxygenase inhibitors, including, without limitation, Cox I and Cox II inhibitors; combinations of anti-infective and anti-inflammatory agents; decongestants; anti-glaucoma agents, including, without limitation, adrenergics, β -adrenergic

blocking agents, α -adrenergic agonists, parasympathomimetic agents, cholinesterase inhibitors, carbonic anhydrase inhibitors, and prostaglandins; combinations of anti-glaucoma agents; antioxidants; nutritional supplements; drugs for the treatment of cystoid macular edema including, without limitation, non-steroidal anti-inflammatory agents; drugs for the treatment of ARMD, including, without limitation, angiogenesis inhibitors and nutritional supplements; drugs for the treatment of herpetic infections and CMV ocular infections; drugs for the treatment of proliferative vitreoretinopathy including, without limitation, antimetabolites and fibrinolytics; wound modulating agents, including, without limitation, growth factors; antimetabolites; neuroprotective drugs, including, without limitation, eliprodil; and angiostatic steroids for the treatment of diseases or conditions of posterior segment 26, including, without limitation, ARMD, CNV, retinopathies, retinitis, uveitis, macular edema, and glaucoma. Such angiostatic steroids are more fully disclosed in U.S. Patent Nos. 5,679,666 and 5,770,592. Preferred ones of such angiostatic steroids include 4,9(11)-Pregnadien-17 α ,21-diol-3,20-dione and 4,9(11)-Pregnadien-17 α ,21-diol-3,20-dione-21-acetate. These preferred angiostatic steroids are preferably formulated as a suspension. A preferred non-steroidal anti-inflammatory for the treatment of cystoid macular edema is nepatenac. The conventional non-active excipients may include, but are not limited to, ingredients to enhance the stability, solubility, penetrability, or other properties of the pharmaceutically active agent or drug depot 38.

[0031] FIGS. 3A through 3E show additional preferred embodiments of cannulae for the creation of drug depot 38. Each of these cannulae have a proximal portion 54, hub 56, bend 57, and hollow bore 58 substantially identical to that described above for cannula 50 of FIG. 2. However, each of these cannulae has a unique distal portion. Hub 56 of each of these cannulae is removably coupled to a conventional syringe 80. Each of these cannulae can be used to create drug depot 38 directly on the outer surface of sclera 16 generally above macula 30 in a manner substantially similar to the techniques described above for cannula 50.

[0032] Cannula 82 of FIG. 3A has a distal portion 52a with a geometry substantially identical to distal portion 52 of cannula 50, except that an orifice 64a is located on the end of tip 62 of distal portion 52a. Similarly, cannula 84 of FIG. 3B has a distal portion 52b with a geometry substantially identical to distal portion 52 of cannula 50, except that tip 62 has two orifices, 64 and 64b. Orifice 64b is located on the upper or exterior side of distal portion 52b. Alternatively, although not shown in FIG. 3B, orifices 64 and 64b may be located laterally, on opposite sides of distal portion 52b.

[0033] Cannula 86 of FIG. 3C has a distal portion 52c with a geometry substantially identical to distal portion 52 of cannula 50, except that a plurality of orifices 64c, 64d, 64e, and 64f are disposed on distal portion 52c

proximate tip 62. Orifices 64c through 64f are preferably disposed on distal portion 52c in the alternating pattern shown in FIG. 3C. Cannula 86 is useful when it is desirable to create a larger drug depot 38. Distal portion 52c 5 may be formed with more or less than the four orifices shown in FIG. 3C, or with a different pattern of orifices than shown in FIG. 3C, if desired.

[0034] Cannula 88 of FIG. 3D has a distal portion 52d with a geometry substantially identical to distal portion 10 52a of cannula 82 of FIG. 3A, except that tip 62d has a globular or olive shape. Tip 62 thus serves as a scleral depressor, allowing a physician to view tip 62d through an ophthalmoscope as he or she guides cannula 88 along the outer surface of sclera 16. Tip 62 is preferably 15 sized to create as small a pathway as possible between Tenon's capsule 34 and sclera 16, but still function as a scleral depressor. A small pathway minimizes the possibility of drug formulation flowing anteriorly from drug depot 38.

[0035] Cannula 90 of FIG. 3E has a distal portion 52e with a geometry substantially identical to distal portion 52 of cannula 50 of FIG. 2, except that tip 62e is also 20 equipped with a fiber optic light source 92, allowing a physician to view tip 62e through an ophthalmoscope 25 as he or she guides cannula 90 along the outer surface of sclera 16. A conventional power source 94 is electrically coupled to fiber optic light source 92 via conventional electrical wiring 96 that is preferably at least partially disposed within the wall of distal portion 52e and proximal portion 54. Fiber optic light source 92, power 30 source 94, and wiring 96 may be incorporated into any of the cannulae disclosed in this application, if desired.

[0036] FIG. 4 schematically illustrates a cannula 100 for the creation of drug depot 38 which is not part of the 35 present invention. Cannula 100 includes a lumen 102 having a geometry substantially identical to cannula 82 of FIG. 3A. Cannula 100 also includes a second, separate lumen 104 disposed adjacent to lumen 102 and having a geometry substantially identical to cannula 82 40 of FIG. 3A. Lumen 102 has a hub 56 removably coupled to a lumen 106a of a dual lumen syringe 106. Lumen 104 has a hub 56 removably coupled to a lumen 106b of dual lumen syringe 106.

[0037] Cannula 100 can be used to create drug depot 45 38 directly on the outer surface of sclera 16 generally above macula 30 in a manner substantially similar to the techniques described above for cannula 50. However, cannula 100 allows the delivery of two separate drug formulations while creating drug depot 38. Alternatively, lumen 104 of cannula 100 can be used to aspirate a non-bioerodable drug depot 38 that has dispensed all of its pharmaceutically active agent, and lumen 102 of cannula 100 can be used to inject a new drug depot 38.

[0038] FIG. 5 schematically illustrates a cannula 120 for the creation of drug depot 38 which is not part of the 50 present invention. Cannula 120 has a geometry substantially identical to that of cannula 100, except that lumen 102 and lumen 104 join at a point 122 proximate a

single orifice 64f near the distal portions of the cannula. [0039] Cannula 120 can be used to create drug depot 38 directly on the outer surface of sclera 16 generally above macula 30 in a manner substantially similar to the techniques described above for cannula 50. However, cannula 120 allows the delivery of two separate formulations that require mixing just prior to injection out of orifice 64f.

From the above, it may be appreciated that the present invention provides an improved apparatus for sub-Tenon delivery of a drug depot to the posterior segment of a human eye proximate the macula. The apparatus and methods of the present invention increase patient safety, are easy for the physician to use, are capable of delivering a wide spectrum of formulations, and are capable of being performed in an outpatient setting. The apparatus and methods of the present invention are especially useful for localized delivery of pharmaceutically active agents to the posterior segment of the eye to combat ARMD, CNV, retinopathies, retinitis, uveitis, macular edema, glaucoma, and other posterior segment diseases. The apparatus of the present invention are also particularly useful for the sub-Tenon delivery of drugs in the form of suspensions, emulsions, ointments, or gels, or drugs in such forms including bioerodable polymers or non-bioerodable polymers.

[0040] The present invention is illustrated herein by example, and various modifications may be made by a person of ordinary skill in the art. For example, although the cannulae of the present invention have been described above in connection with the preferred sub-Tenon drug delivery generally above the macula, the cannulae can be used to deliver drugs directly on the outer surface of the sclera, below the Tenon's capsule, and generally above portions of the retina other than the macula. As another example, the arc length and/or radius of curvature of the distal portions of the cannulae may be modified to deliver drugs within the Tenon's capsule or the sclera, generally above the macula or other portions of the retina, if desired.

[0041] It is believed that the operation and construction of the present invention will be apparent from the foregoing description. While the apparatus shown or described above have been characterized as being preferred, various changes and modifications may be made therein without departing from the scope of the invention as defined in the following claims.

Claims

1. A cannula (50, 82, 84, 86, 88, 90, 100, 120), comprising:

a distal portion (52, 52a, 52b, 52c, 52d, 52e)
a straight proximal portion (54); and
a bend (57) separating said distal portion and said proximal portion,

wherein the distal portion is defined by an arc having an arc length and a radius of curvature over said length, the distal portion extending from the bend to the tip of the cannula, and a tangent of said distal portion at said bend is disposed at an angle of no more than about 56 degrees with respect to said proximal portion, and wherein at least one orifice (64, 64a, 64b, 64c, 64d, 64e, 64f) is provided proximate said tip.

5 2. The cannula of claim 1, wherein said distal portion comprises an arc length of about 15 mm to about 18 mm.

10 3. The cannula of claim 1, further comprising:
a hub (56) coupled to said proximal portion for removably coupling to a syringe (80); and
20 a hollow bore (58) disposed axially within said distal portion and said proximal portion.

4. The cannula of claim 1, wherein said distal portion comprises an interior side and the at least one orifice (64, 64a, 64b, 64c, 64d, 64e, 64f) is provided on said interior side.

25 5. The cannula of claim 1, wherein said radius of curvature of said distal portion (52) is about 11.5 mm to about 14 mm.

30 6. The cannula of claim 1, wherein said distal portion (52) comprises a plastic.

35 7. The cannula of claim 1, wherein a part of said distal portion proximate said tip comprises a plastic, and a remainder of said distal portion comprises a metal.

40 8. The cannula of claim 1, wherein said angle is about 56 degrees.

45 9. The cannula of claim 3, wherein said distal portion (52c) comprises a plurality of orifices (64c, 64d, 64e, 64f), each of said orifices communicating with said hollow bore (58).

Patentansprüche

50 1. Kanüle (50, 82, 84, 86, 88, 90, 100, 120), die folgendes umfaßt:
55 einen distalen Abschnitt (52, 52a, 52b, 52c, 52d, 52e),
einen geraden proximalen Abschnitt (54) und eine Biegung (57), die den distalen Abschnitt und den proximalen Abschnitt trennt, wobei der distale Abschnitt durch einen Bogen mit einer

Bogenlänge und einem Krümmungsradius über die Länge festgelegt ist, der distale Abschnitt sich von der Biegung zu der Spitze der Kanüle erstreckt und eine Tangente des distalen Abschnitts an der Biegung in einem Winkel von nicht mehr als 56° gegenüber dem proximalen Abschnitt verläuft und wobei wenigstens eine Öffnung (64, 64a, 64b, 64c, 64d, 64e, 64f) in der Nähe der Spitze vorgesehen ist.

2. Kanüle nach Anspruch 1, bei welcher der distale Abschnitt eine Bogenlänge von etwa 15 mm bis etwa 18 mm umfaßt.

3. Kanüle nach Anspruch 1, die weiterhin folgendes umfaßt:

eine Nabe (56), die mit dem proximalen Abschnitt verbunden ist, zum lösbar Verbinden mit einer Spritze (80) und
eine Hohlbohrung (58), die axial in dem distalen Abschnitt und dem proximalen Abschnitt angeordnet ist.

4. Kanüle nach Anspruch 1, bei welcher der distale Abschnitt eine Innenseite aufweist und die eine Öffnung (64, 64a, 64b, 64c, 64d, 64e, 64f) auf der Innenseite vorgesehen ist.

5. Kanüle nach Anspruch 1, bei welcher der Krümmungsradius des distalen Abschnitts (52) etwa 11,5 mm bis etwa 14 mm beträgt.

6. Kanüle nach Anspruch 1, bei welcher der distale Abschnitt (52) einen Kunststoff umfaßt.

7. Kanüle nach Anspruch 1, bei welcher ein Teil des distalen Abschnitts proximal der Spitze einen Kunststoff umfaßt und ein Rest des distalen Abschnitts ein Metall umfaßt.

8. Kanüle nach Anspruch 1, bei welcher der Winkel etwa 56° beträgt.

9. Kanüle nach Anspruch 3, bei welcher der distale Abschnitt (52c) mehrere Öffnungen (64c, 64d, 64e, 64f) umfaßt, wobei jede der Öffnungen mit der Hohlbohrung (58) in Verbindung steht.

ladite partie proximale,

dans laquelle la partie distale est définie par un arc ayant une longueur d'arc et un rayon de courbure sur ladite longueur, la partie distale s'étendant du coude jusqu'à la pointe de la canule, et une tangente de ladite partie distale audit coude est disposée selon un angle ne dépassant pas environ 56 degrés par rapport à ladite partie proximale, et dans laquelle au moins un orifice (64, 64a, 64b, 64c, 64d, 64e, 64f) est fourni à proximité de ladite pointe.

2. Canule selon la revendication 1, dans laquelle ladite partie distale comprend une longueur d'arc d'environ 15 mm à environ 18 mm.

3. Canule selon la revendication 1 comprenant en outre :

- un raccord (56) couplé à ladite partie proximale pour coupler de manière amovible à une seringue (80) ; et
- un alésage creux (58) disposé de manière axiale à l'intérieur de ladite partie distale et ladite partie proximale.

4. Canule selon la revendication 1, dans laquelle ladite portion distale comprend une face intérieure et au moins un orifice est fourni sur ladite face intérieure.

5. Canule selon la revendication 1, dans laquelle ledit rayon de courbure de ladite partie distale (52) est d'environ 11,5 mm à environ 14 mm.

6. Canule selon la revendication 1, dans laquelle ladite partie distale (52) comprend un plastique.

7. Canule selon la revendication 1, dans laquelle une portion de ladite partie distale située à proximité de ladite pointe comprend un plastique et le reste de ladite partie distale comprend un métal.

8. Canule selon la revendication 1, dans laquelle ledit angle est d'environ 56 degrés.

9. Canule selon la revendication 3, dans laquelle ladite partie distale (52c) comprend une pluralité d'orifices (64c, 64d, 64e, 64f), chacun desdits orifices communiquant avec ledit alésage creux (58).

Revendications

1. Canule (50, 82, 84, 86, 88, 90, 100, 120) comprenant :

- une partie distale (52, 52a, 52b, 52c, 52d, 52e)
- une partie proximale droite (54) ; et
- un coude (57) séparant ladite partie distale et




FIG. 1

FIG. 2




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D





FIG. 3E






FIG. 4




FIG. 5





FIG. 7