



(86) Date de dépôt PCT/PCT Filing Date: 2012/08/09
(87) Date publication PCT/PCT Publication Date: 2014/02/13
(45) Date de délivrance/Issue Date: 2018/07/24
(85) Entrée phase nationale/National Entry: 2015/02/06
(86) N° demande PCT/PCT Application No.: CN 2012/079897
(87) N° publication PCT/PCT Publication No.: 2014/023007


(51) Cl.Int./Int.Cl. *C07K 14/435*(2006.01),
A61K 38/10(2006.01), *A61K 38/17*(2006.01),
A61P 19/00(2006.01), *A61P 21/00*(2006.01),
C07K 7/08(2006.01), *C12N 5/074*(2010.01)

(72) Inventeurs/Inventors:
TSAO, YEOU-PING, CN;
HO, TSUNG-CHUAN, CN

(73) Propriétaire/Owner:
MACKAY MEMORIAL HOSPITAL, TW

(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : UTILISATION DE POLYPEPTIDES DERIVES DE PEDF POUR FAVORISER LA REGENERATION DE MUSCLES OU DE TENDONS OU L'ARTERIOGENESE
(54) Title: USE OF PEDF-DERIVED POLYPEPTIDES FOR PROMOTING MUSCLE OR TENDON REGENERATION OR ARTERIOGENESIS



(57) Abrégé/Abstract:

Provided is a synthetic peptide derived from PEDF, which has 20-39 amino acid residues. Also provided are pharmaceutical compositions containing the synthetic peptide and the use thereof. The peptide is useful in promoting muscle or tendon regeneration, and/or arteriogenesis in a subject.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(10) International Publication Number

WO 2014/023007 A1

(43) International Publication Date
13 February 2014 (13.02.2014)

WIPO | PCT

(51) International Patent Classification:

C07K 14/435 (2006.01) *A61K 38/10* (2006.01)
C07K 7/08 (2006.01) *A61P 19/00* (2006.01)
C12N 5/074 (2010.01) *A61P 21/00* (2006.01)
A61K 38/17 (2006.01)

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:

PCT/CN2012/079897

(22) International Filing Date:

9 August 2012 (09.08.2012)

(25) Filing Language: English
 (26) Publication Language: English
 (72) Inventors; and
 (71) Applicants : **TSAO, Yeou-Ping** [CN/CN]; No.92, Sec. 2, Zhongshan N Rd., Zhongshan District, Taipei City 104, Taiwan (CN). **HO, Tsung-Chuan** [CN/CN]; No.92, Sec. 2, Zhongshan N Rd., Zhongshan District, Taipei City 104, Taiwan (CN).
 (74) Agent: **SHANGHAI PATENT & TRADEMARK LAW OFFICE, LLC**; 435 Guiping Road, Shanghai 200233 (CN).
 (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))



WO 2014/023007 A1

(54) Title: USE OF PEDF-DERIVED POLYPEPTIDES FOR PROMOTING MUSCLE OR TENDON REGENERATION OR ARTERIOGENESIS

(57) Abstract: Provided is a synthetic peptide derived from PEDF, which has 20-39 amino acid residues. Also provided are pharmaceutical compositions containing the synthetic peptide and the use thereof. The peptide is useful in promoting muscle or tendon regeneration, and/or arteriogenesis in a subject.

USE OF PEDF-DERIVED POLYPEPTIDES FOR PROMOTING MUSCLE OR TENDON REGENERATION OR ARTERIOGENESIS

BACKGROUND OF THE INVENTION

5 [0001] 1. FIELD OF THE INVENTION

[0002] The present disclosure relates to the treatment of tissue damages. In particular, the disclosed invention relates to the use of PEDF-derived polypeptides for promoting muscle or tendon regeneration or arteriogenesis in the treatment of tissue damages.

10 [0003] 2. DESCRIPTION OF RELATED ART

[0004] Muscle tissues are classified as skeletal, cardiac or smooth muscles. Muscle is capable of repairing its damage. After injury, skeletal muscle is repaired by a spontaneous process to remove damaged myofibers and synthesizing new muscle fibers. However, such spontaneous tissue repair mechanism is absent in some tissue damage or inadequate to effect a full recovery of the tissue. For example, some pathologic conditions (such as severe injury, advanced age, muscle disuse, cancer, and tissue ischemia) or genetic defects (such as muscular dystrophy) may lead to impaired healing. Failure of repair may lead to permanent loss of muscle mass, disease progression, and functional deficiency.

20 [0005] A tendon is a tough band of fibrous connective tissue that usually connects muscle to bone. Tendon injuries generally result in inflammation and degeneration or weakening of the tendons, which may eventually lead to tendon rupture. Tendon healing is a long and intricate process that typically takes months,

and over a time period of about one year, the tissue will gradually turn from fibrous to scar-like. Such scar tissue may result in reduced elasticity and mobility of the tendon and increased propensity for recurrence of injury. Tendon-derived stem cells (TSCs) and bone marrow-derived mesenchymal stem cells (BM-MSCs) offer limited 5 autologous healing of tendonitis lesions.

[0006] Episodes of ischemia are another cause of considerable tissue damage. Ischemic episodes leading to tissue damage result in myocardial infarctions, stroke, and other disorders. Short episodes of ischemia cause mild damage from which a cell can recover, while longer periods of ischemia cause irreversible cell damage, leading to 10 cell death. In the latter case, even if blood circulation is reestablished, total functional recovery of the damaged cell is impossible. Furthermore, loss of function always precedes cell death.

[0007] No present treatment for these conditions offers a cure or facilitates regeneration of the damaged, nonfunctional tissue. Thus, there exists a need in the 15 art for means that promotes regeneration of tissue. In particular, it would be desirable to provide a composition and method for promoting arteriogenesis so as to promote blood flow in or adjacent to the damaged tissue region and to permit quasi-normal function to the tissue.

20

SUMMARY

[0008] The following presents a simplified summary of the disclosure in order to provide a basic understanding to the reader. This summary is not an extensive overview of the disclosure and it does not identify key/critical elements of the present

invention or delineate the scope of the present invention. Its sole purpose is to present some concepts disclosed herein in a simplified form as a prelude to the more detailed description that is presented later.

[0009] The present disclosure is based, at least, on the finding that synthetic peptides derived from pigment epithelium-derived factor (PEDF) may promote the muscle regeneration or tendon regeneration as well as arteriogenesis in a subject. The PEDF-derived synthetic peptides of this invention are, therefore, useful as an agent or a medicament for treating tissue damages (in particular, those associated with ischemia).

[0010] Accordingly, in one aspect, the present disclosure is directed to a synthetic peptide for promoting muscle or tendon regeneration in a subject.

[0011] According to embodiments of the present disclosure, the synthetic peptide is 20–39 amino acid residues in length, and has an amino acid sequence that is at least 80% identical to SEQ ID NO: 1. Also, the amino acid sequence comprises at least 20 consecutive residues, which is at least 90% identical to residues 11–30 of SEQ ID NO: 1, such that the synthetic peptide is useful in promoting the muscle or tendon regeneration in a subject.

[0012] According to optional embodiments of the present disclosure, at least 4 consecutive residues of the synthetic peptide are identical to residues 11–14 of SEQ ID NO: 1. Non-limiting examples of such synthetic peptides include those respectively having an amino acid sequence of SEQ ID NO: 1 (39-mer), SEQ ID NO: 2 (34-mer), SEQ ID NO: 3 (29-mer), SEQ ID NO: 5 (24-mer), SEQ ID NO: 6 (20-mer), SEQ ID NO: 8 (MO 29-mer), and SEQ ID NO: 9 (MO 20-mer). In some embodiments of the present

disclosure, the amino acid sequence of the synthetic peptide is any of SEQ ID NO: 3 (29-mer), SEQ ID NO: 5 (24-mer), or SEQ ID NO: 6 (20-mer).

[0013] In another aspect, the present disclosure is directed to a pharmaceutical composition for promoting muscle or tendon regeneration in a subject. The subject 5 may be any animal classified as a mammal, including human.

[0014] According to one embodiment of the present disclosure, the pharmaceutical composition comprises a synthetic peptide according to any of the above-mentioned aspect/embodiments, and the synthetic peptide is present in an effective amount sufficient to promote muscle or tendon regeneration in the subject.

10 The pharmaceutical composition also comprises a pharmaceutically acceptable carrier for the synthetic peptide.

[0015] According to optional embodiments of the present disclosure, the pharmaceutically acceptable carrier is a polymeric material, which may be any of alginate, gelatin, collagen, or poly(lactide-co-glycolide).

15 [0016] According to optional embodiments of the present disclosure, the synthetic peptide is present in the pharmaceutical composition in an amount of about 1–100 μ M, and preferably, about 10 μ M.

[0017] In yet another aspect, the present invention is directed to a method for promoting muscle or tendon regeneration in or adjacent to a damaged region of a 20 subject. The subject may be any animal classified as a mammal, including human.

[0018] In one embodiment, the method comprises administering, to a treatment region of the subject, a therapeutically effective amount of the synthetic peptide according to the above-mentioned aspect/embodiments of the present

disclosure, wherein the treatment region is adjacent to the damaged region so as to promote muscle or tendon regeneration in or adjacent to the damaged region of the subject.

[0019] According to optional embodiments, the synthetic peptide is formulated 5 into a pharmaceutical composition according to the above-mentioned aspect/embodiments of the present disclosure. In practice, the pharmaceutical composition may be administered via intramuscular injection.

[0020] According to some embodiments, the subject may be suffering from muscle injury, muscle disuse, muscular dystrophy, amyotrophic lateral sclerosis, 10 tendon injury, tissue ischemia, cerebral ischemia, peripheral arterial diseases, or myocardial infarction, which causes the muscle or tendon damage in the damaged region.

[0021] Also, in another aspect, the present disclosure is directed to a synthetic peptide for promoting arteriogenesis in a subject. The subject may be any animal 15 classified as a mammal, including human.

[0022] According to embodiments of the present disclosure, the synthetic peptide is 20–39 amino acid residues in length, and has an amino acid sequence that is at least 80% identical to SEQ ID NO: 1. Also, the amino acid sequence comprises at least 20 consecutive residues, which is at least 90% identical to residues 11–30 of SEQ 20 ID NO: 1, such that the synthetic peptide is useful in promoting the arteriogenesis in a subject.

[0023] According to optional embodiments of the present disclosure, at least 4 consecutive residues of the synthetic peptide are identical to residues 11–14 of SEQ ID

NO: 1. Non-limiting examples of such synthetic peptides include those respectively having an amino acid sequence of SEQ ID NO: 1 (39-mer), SEQ ID NO: 2 (34-mer), SEQ ID NO: 3 (29-mer), SEQ ID NO: 5 (24-mer), SEQ ID NO: 6 (20-mer), SEQ ID NO: 8 (MO 29-mer), and SEQ ID NO: 9 (MO 20-mer). In some embodiments of the present disclosure, the amino acid sequence of the synthetic peptide is any of SEQ ID NO: 3 (29-mer), SEQ ID NO: 5 (24-mer), or SEQ ID NO: 6 (20-mer).

[0024] In another aspect, the present disclosure is directed to a pharmaceutical composition for promoting arteriogenesis in a subject. The subject may be any animal classified as a mammal, including human.

10 [0025] According to one embodiment of the present disclosure, the pharmaceutical composition comprises a synthetic peptide according to any of the above-mentioned aspect/embodiments, and the synthetic peptide is present in an effective amount sufficient to promote arteriogenesis in the subject. The pharmaceutical composition also comprises a pharmaceutically acceptable carrier for 15 the synthetic peptide.

[0026] According to optional embodiments of the present disclosure, the pharmaceutically acceptable carrier is a polymeric material, which may be any of alginate, gelatin, collagen, or poly(lactide-co-glycolide).

20 [0027] According to optional embodiments of the present disclosure, the synthetic peptide is present in the pharmaceutical composition in an amount of about 1–100 μ M, and preferably, about 10 μ M.

[0028] In yet another aspect, the present invention is directed to a method for promoting arteriogenesis in or adjacent to an ischemic region of a subject. The subject may be any animal classified as a mammal, including human.

[0029] In one embodiment, the method comprises administering, to a 5 treatment region of the subject, a therapeutically effective amount of the synthetic peptide according to the above-mentioned aspect/embodiments of the present disclosure, wherein the treatment region is adjacent to the ischemic region so as to promote arteriogenesis in or adjacent to the ischemic region of the subject.

[0030] According to optional embodiments, the synthetic peptide is formulated 10 into a pharmaceutical composition according to the above-mentioned aspect/embodiments of the present disclosure. In practice, the pharmaceutical composition may be administered via intramuscular injection.

[0031] According to some embodiments, the subject may be suffered from 15 muscle injury, muscle disuse, muscular dystrophy, amyotrophic lateral sclerosis, tendon injury, tissue ischemia, cerebral ischemia, peripheral arterial diseases, or myocardial infarction, which causes the blood flow at the ischemic region to be hindered or blocked.

[0032] Many of the attendant features and advantages of the present 20 disclosure will becomes better understood with reference to the following detailed description considered in connection with the accompanying drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

[0033] The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawings will be provided by the Office upon request and payment of the necessary fee.

5 [0034] The present description will be better understood from the following detailed description read in light of the accompanying drawings.

[0035] Figure 1 illustrates the cumulative *in vitro* release of PEDF peptides from alginate gel in PBS at 37°C. The results are presented as the means \pm standard deviation for three separate experiments.

10 [0036] Figure 2 provides representative LDPI images illustrating the blood perfusion of ischemic hindlimbs over a time period of 4 weeks.

[0037] Figure 3 illustrates the blood perfusion analysis of mice hindlimbs treated with blank alginate gel, sustained-release formulation containing 29-mer, 24-mer, 20-mer, or 18-mer, and bolus formulation containing 29-mer. The results are 15 presented as the means \pm standard deviation for three separate experiments; $n \geq 6$.

* $P < 0.05$ versus blank control.

[0038] Figure 4A provides representative photographs from tibialis muscle specimens stained by Masson trichrome (original magnification, $\times 40$), and Figure 4B provides representative photographs from the same specimens at higher magnification 20 to highlight the extent of necrosis after surgical induction of hindlimb ischemia for 2 and 7 weeks (original magnification, $\times 200$).

[0039] Figure 5 provides representative immunostained images of arterioles in adductor magnus muscle after 2 weeks of ischemia. Arterioles were labeled with anti- α -SMA (brown) and nuclei were labeled with hematoxylin.

[0040] Figure 6 provides representative photographs of aortic ring explants 5 cultured for 4 days in either basal MCDB131 medium (untreated control) or medium supplemented with known angiogenic factors (FGF2 or VEGF), the control PEDF peptides (25-mer or 18-mer), or the PEDF peptides according to embodiments of the present disclosure (29-mer, 24-mer, 20-mer, Mo 29-mer, or Mo 20-mer).

[0041] Figure 7 provides representative dual-immunostained images 10 illustrating vascular smooth muscle cells (vSMCs) outgrowth from aortic rings cultured in medium supplemented with PEDF peptide (29-mer, 20-mer and 18-mer), in which endothelial cells were detected by Alexa Fluor 594-labeled isolectin B4 (IB4; red; left panel) and vSMCs were labeled with anti- α -SMA (green; middle panel). Merged images are located on the right (yellow). Nuclei were visualized with Hoechst 33258 15 staining. Original magnification, $\times 400$. Images are representative of four independent experiments.

[0042] Figure 8 provides representative photographs from soleus muscle specimens stained by H&E at day 14 following bupivacaine injection.

[0043] Figure 9 is a diagram illustrating muscle fiber size distributions of 20 muscles from animals in various experimental conditions.

[0044] Figure 10 provides representative photographs illustrating regenerating tissue (↑) at the inner part of tendon at week 3 post-injury. Original magnification, $\times 100$.

[0045] Figure 11 provides representative photographs of H&E-stained sections of Achillis tendon at 3 weeks after injury. Original magnification, $\times 400$; scale bar = 50 μM . Images are representative of three independent experiments.

[0046] Figure 12 provides representative photographs of tissue sections 5 stained by Masson's trichrome to highlight the collagen fibers at 3 weeks post-injury. Stars (*) represent the uninjured area in tendon. Original magnification, $\times 400$; scale bar = 50 μM . Images are representative of three independent experiments.

[0047] Figure 13 provides representative immunostained photographs of newly 10 formed type 1 collagen (brown color) in regenerating tendon at 3 weeks after surgery. Nuclei labeled with hematoxylin. Boxed regions are shown at higher magnification below. Scale bar = 50 μM . Images are representative of three independent experiments.

[0048] Figure 14 is a representative gel electrophoresis image illustrating 15 enhanced expression level of tenomodulin (TNMD) gene by the present PEDF peptides (29-mer and 20-mer) according to one working example of the present disclosure. Expression of tenomodulin (TNMD) gene is indicative of BM-MSC differentiation into tenocyte. The image is representative of three independent experiments.

DESCRIPTION

20 [0049] The detailed description provided below in connection with the appended drawings is intended as a description of the present examples and is not intended to represent the only forms in which the present example may be constructed or utilized. The description sets forth the functions of the example and

the sequence of steps for constructing and operating the example. However, the same or equivalent functions and sequences may be accomplished by different examples.

[0050] For convenience, certain terms employed in the entire application 5 (including the specification, examples, and appended claims) are collected here. Unless otherwise defined herein, scientific and technical terminologies employed in the present disclosure shall have the meanings that are commonly understood and used by one of ordinary skill in the related art. Unless otherwise required by context, it will be understood that singular terms shall include plural forms of the same and 10 plural terms shall include the singular. Specifically, as used herein and in the claims, the singular forms “a” and “an” include the plural reference unless the context clearly indicates otherwise.

[0051] Notwithstanding that the numerical ranges and parameters setting forth the broad scope of the invention are approximations, the numerical values set 15 forth in the specific examples are reported as precisely as possible. Any numerical value, however, inherently contains certain errors necessarily resulting from the standard deviation found in the respective testing measurements. Also, as used herein, the term “about” generally means within 10%, 5%, 1%, or 0.5% of a given value or range. Alternatively, the term “about” means within an acceptable standard error 20 of the mean when considered by one of ordinary skill in the art. Other than in the operating/working examples, or unless otherwise expressly specified, all of the numerical ranges, amounts, values and percentages such as those for quantities of materials, durations of times, temperatures, operating conditions, ratios of amounts,

and the likes thereof disclosed herein should be understood as modified in all instances by the term “about.” Accordingly, unless indicated to the contrary, the numerical parameters set forth in the present disclosure and attached claims are approximations that can vary as desired. At the very least, each numerical parameter 5 should at least be construed in light of the number of reported significant digits and by applying ordinary rounding techniques.

[0052] As used herein, the term “peptide” denotes a polymer of amino acid residues. By the term “synthetic peptide” as used herein, it is meant a peptide which does not comprise an entire naturally occurring protein molecule. The peptide is 10 “synthetic” in that it may be produced by human intervention using such techniques as chemical synthesis, recombinant genetic techniques, or fragmentation of the whole protein or the like. Throughout the present disclosure, the positions of any specified amino acid residues within a peptide are numbered starting from the N terminus of the peptide.

15 [0053] The term “stem cell” as used herein, refers to a cell that retains the capacity, under certain circumstances, to proliferate without substantially differentiating; as well as the capacity or potential, under particular circumstances, to differentiate to a more specialized or differentiated phenotype.

[0054] As used herein, “proliferating” and “proliferation” refers to an increase 20 in the number of cells in a population by means of cell division.

[0055] As used herein, the term “muscle cell” refers to any cell which contributes to muscle tissue, and encompasses myoblasts, satellite cells, myotubes, and myofibril tissues. “Muscle regeneration” as used herein refers to the process by

which new muscle fibers form from muscle progenitor cells. The regeneration of muscle in or adjacent to the damaged region may be evidenced by the increase in the number, diameter (size), wet weight, and/or the protein content of the muscle fibers in or adjacent to the damaged region. Also, the muscle regeneration may be

5 monitored by the proliferative activity of muscle cells and/or satellite cells in or adjacent to the damaged region.

[0056] As used herein, the term “tendon” refers to a fibrous tissue composed of parallel arrays of closely packed collagen fibers that connects muscle to bone. The healing of damaged tendon is a slow process and usually associated with scar

10 formation which may result in a defective tendon that cannot resume normal or original tendon function. As used herein, the term “tendon regeneration” refers to a tendon healing process in which type I collagen is formed, and the newly formed collagen fibers align parallel to the direction of load application, whereby resulting in minimal scar formation. The regeneration of tendon in or adjacent to the damaged

15 region may be evidenced by the increase in the number of the collagen fibrils with an organized orientation in or adjacent to the damaged region. Also, the tendon regeneration may be monitored by the proliferative activity of tendon stem cells in or adjacent to the damaged region.

[0057] As used herein, the term “arteriogenesis” is to be distinguished from

20 “angiogenesis.” Angiogenesis is a process by which new capillary blood vessels sprout from a pre-existing blood vessel. It is important to recognize that these newly formed capillary tubes lack vascular smooth muscle cells. Accordingly, they are fragile and prone to rupture. These capillary tubes would not go through vasculature

remodeling process, and hence are unable to sustain and/or restore proper circulation in and/or adjacent to the damaged region. In contrast to the capillary sprouting, arteriogenesis refers to the *in situ* recruitment and expansion of arteries or collateral arteries by proliferation of endothelial and smooth muscle cells from pre-existing arteriolar connections. These newly formed arteries or collateral arteries would develop into a functional network of arteries (or collateral arteries) which constitute natural bypasses capable of supplying sufficient blood to the damaged or ischemic tissue or site of inflammation.

[0058] The term “ischemia” as used herein relates to a condition that may occur in any tissue and/or organ that suffers from a lack of oxygen supply and/or from abnormal accumulation of metabolites, which occurs when there is an imbalance between oxygen supply and demand, due to inadequate perfusion, e.g., caused by atherosclerosis, restenotic lesions, anemia, stroke or clogged arteries just to name a few, that leads to insufficient oxygen to tissues such as, for example, the muscle, heart or brain. However, ischemia is not limited to the aforementioned organs or tissues, since it may occur in any organ/tissue.

[0059] The term “promote” or “promoting” is meant to refer to a positive alteration; in particular a statistically significant positive alteration. The positive alteration means an increase of at least 10% as compared to a reference level.

[0060] “Percentage (%) amino acid sequence identity” with respect to the synthetic polypeptide sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the specific polypeptide sequence, after aligning the sequences and

introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity.

Alignment for purposes of determining percentage sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available

5 computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software.

Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full

length of the sequences being compared. For purposes herein, sequence comparison

between two amino acid sequences was carried out by computer program Blastp

10 (protein-protein BLAST) provided online by Nation Center for Biotechnology

Information (NCBI). The percentage amino acid sequence identity of a given amino

acid sequence A to a given amino acid sequence B (which can alternatively be phrased

as a given amino acid sequence A that has a certain % amino acid sequence identity to

a given amino acid sequence B) is calculated by the formula as follows:

15
$$\frac{X}{Y} \times 100\%$$

where X is the number of amino acid residues scored as identical matches by the

sequence alignment program BLAST in that program's alignment of A and B, and where

Y is the total number of amino acid residues in A or B, whichever is shorter.

[0061] The phrase "pharmaceutically acceptable carrier" as used herein means

20 a pharmaceutically acceptable material, composition or vehicle, such as a liquid or

solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or

transporting the subject agents from one organ, or a portion of the body, to another

organ, or another portion of the body. Each carrier must be "acceptable" in the

sense of being compatible with the other ingredients of the formulation. The carrier can be in the form of a solid, semi-solid, or liquid diluent, cream or a capsule.

[0062] The terms “treatment” and “treating” are used herein to generally mean obtaining a desired pharmaceutical and/or physiological effect. Preferably, the 5 effect is therapeutic in terms of partially or completely curing the muscle damage, tendon damage, or ischemia. The term “treating” as used herein refers to application or administration of the synthetic peptide or pharmaceutical composition of the present disclosure to a subject, who has a medical condition, a symptom of the condition, a disease or disorder secondary to the condition, or a predisposition toward 10 the condition, with the purpose to partially or completely alleviate, ameliorate, relieve, delay onset of, inhibit progression of, reduce severity of, and/or reduce incidence of one or more symptoms or features of a particular disease, disorder, and/or condition. Treatment may be administered to a subject who does not exhibit signs of a disease, disorder, and/or condition and/or to a subject who exhibits only early signs of a 15 disease, disorder, and/or condition for the purpose of decreasing the risk of developing pathology associated with the disease, disorder, and/or condition. Treatment is generally “effective” if one or more symptoms or clinical markers are reduced as that term is defined herein. Alternatively, a treatment is “effective” if the progression of a disease is reduced or halted. That is, “treatment” includes not just 20 the improvement of symptoms or decrease of markers of the disease, but also a cessation or slowing of progress or worsening of a symptom that would be expected in absence of treatment. Beneficial or desired clinical results include, but are not limited to, alleviation of one or more symptom(s), diminishment of extent of disease,

stabilized (i.e., not worsening) state of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, and remission (whether partial or total), whether detectable or undetectable.

[0063] The term “effective amount” as used herein refers to the quantity of a component which is sufficient to yield a desired response. The term “therapeutically effective amount” as used herein refers to the amount of therapeutically agent of pharmaceutical composition to result in a desired “effective treatment” as defined hereinabove. The specific therapeutically effective amount will vary with such factors as the particular condition being treated, the physical condition of the patient (e.g., the patient's body mass, age, or gender), the type of mammal or animal being treated, the duration of the treatment, the nature of concurrent therapy (if any), and the specific formulations employed. A therapeutically effective amount is also one in which any toxic or detrimental effects of the compound or composition are outweighed by the therapeutically beneficial effects.

[0064] The term “subject” refers to a mammal including the human species that is treatable with the synthetic peptides, compositions, and/or methods of the present invention. The term “subject” is intended to refer to both the male and female gender unless one gender is specifically indicated.

[0065] Pigment epithelium-derived factor (PEDF) is a multifunctional secreted protein that has anti-angiogenic, anti-tumorigenic, and neurotrophic functions. Human PEDF protein (SEQ ID No: 14) is a secreted protein of roughly 50 kDa size and 418 amino acids in length. A 34-mer fragment (residues 44–77) and a 44-mer

fragment (residues 78–121) of PEDF have been identified to have anti-angiogenic and neurotrophic properties, respectively.

[0066] The present disclosure is based, at least, on the finding that synthetic peptides derived from PEDF may promote the regeneration of muscle or tendon tissue and arteriogenesis in a subject. In particular, the present disclosure is the first to identify a link between the local delivery of PEDF-derived peptides and the healing of muscle or tendon tissues suffering from damage and/or ischemia or the formation of (collateral) arteries in or adjacent to the ischemic region. Another inventive feature of the present invention lies in that the synthetic peptides are much shorter (39 amino acid residues at most) than the full-length PEDF and thus overcomes the limitations associated with the clinical use of conventional protein drugs, including high manufacturing cost, low bioavailability, and poor pharmacokinetics. Accordingly, the present synthetic peptides are useful for treating muscle or tendon damages as well as tissues or organs suffering from ischemia.

[0067] Thus, in one aspect, the present disclosure is directed to a synthetic peptide for promoting muscle or tendon regeneration in a subject. In another aspect, the present disclosure is directed to a synthetic peptide for promoting arteriogenesis in a subject. Embodiments applicable to either or both of these two aspects are discussed below.

[0068] According to embodiments of the present disclosure, the synthetic peptide has 20–39 amino acid residues in length, and has at least 80% amino acid sequence identity with the amino acid sequence of LSVATALSALSLGAEQRTEIIHRALYYDLISSPDIHGT (SEQ ID NO: 1). For example, the

synthetic peptide may have an amino acid sequence identity of about 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100 percent with SEQ ID NO: 1. Also, the synthetic peptide comprises at least 20 consecutive residues that are at least 90% identical to residues 11–30 of SEQ ID NO: 1. Specifically, the 20 consecutive amino acid residues may have about 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100 percent amino acid sequence identity with residues 11–30 of SEQ ID NO: 1.

[0069] In one embodiment, the synthetic peptide has the sequence of SEQ ID NO: 1, which has 39 amino acids in length. This synthetic peptide is referred to as 10 39-mer in the description hereinbelow. This 39-mer peptide corresponds to residues 83–121 of human PEDF and hence is a short variant derived from the known PEDF 44-mer (corresponding to residues 78–121 of PEDF).

[0070] Prior experiments conducted by the present inventors, such as those disclosed in US Patent No. 9,051,547

and experiments provided hereinbelow, reveal that 15 several short, synthetic PEDF peptides derived from the 39-mer, are capable of promoting muscle or tendon regeneration and/or arteriogenesis in a subject.

[0071] For example, based on experiments disclosed in both the prior application and the present application, a 34-mer synthetic peptide having the sequence of ALSALSLGAEQRTESIIHRALYYDLISSPDIHGT (SEQ ID NO: 2) is effective in 20 promoting muscle or tendon regeneration and/or arteriogenesis in a subject. This 34-mer peptide corresponds to residues 88–121 of human PEDF. According to the process for estimating percentage of sequence identity between any two given sequences provided above, the 34-mer has a 100% amino acid sequence identity to

the 39-mer, and the 6th–25th amino acid residues of the 34-mer has a 100% amino acid sequence identity to the amino acid residues 11–30 of the 39-mer.

[0072] Additionally, according to various examples hereinbelow, a 29-mer synthetic peptide having the sequence of SLGAEQRTEESIIHRALYYDLISSPDIHGT (SEQ ID NO: 3) has been confirmed to be effective in promoting muscle or tendon regeneration as well as arteriogenesis in a subject. This 29-mer peptide corresponds to residues 93–121 of human PEDF with a 100% amino acid sequence identity to the 39-mer. Also, the 1st –20th amino acid residues of the 29-mer has a 100% amino acid sequence identity to the amino acid residues 11–30 of the 39-mer.

[0073] In some examples, a 24-mer has been confirmed to be effective in promoting tendon regeneration and arteriogenesis in a subject. The 24-mer has the sequence of SLGAEQRTEESIIHRALYYDLISSP (SEQ ID NO: 5), which corresponds to residues 93–116 of human PEDF. This 24-mer peptide has a 100% amino acid sequence identity to the 39-mer in which the first twenty amino acid residues thereof has a 100% amino acid sequence identity to the amino acid residues 11–30 of the 39-mer.

[0074] In other examples, it has been established that a 20-mer may promote muscle or tendon regeneration as well as arteriogenesis in a subject. The 20-mer has the sequence of SLGAEQRTEESIIHRALYYDL (SEQ ID NO: 6), which corresponds to residues 93–112 of human PEDF. This 20-mer peptide is completely identical to the amino acid residues 11–30 of the 39-mer (100% amino acid sequence identity), and has a 100% amino acid sequence identity to the 39-mer.

[0075] Two synthetic peptides derived from mouse PEDF may also promote muscle or tendon regeneration and/or arteriogenesis in a subject based on experiments disclosed in both the prior application and the present application. The first mouse-derived peptide is referred to as "Mo 29-mer" in the present disclosure.

5 The Mo 29-mer has a sequence of SLGAEHRTESVIHRALYYDLITNPDIHST (SEQ ID NO: 8), which has a 83% amino acid sequence identity to 39-mer, and the first 20 amino acid residues thereof has a 90% amino acid sequence identity to the 11–30 amino acid residues of the 39-mer. Another mouse-derived peptide, Mo 20-mer has a sequence of SLGAEHRTESVIHRALYYDL (SEQ ID NO: 9). The Mo 20-mer has a 90% amino acid sequence identity to either the 39-mer or the 11–30 amino acid residues of the 10 39-mer.

[0076] Optionally, the synthetic peptide comprises 4 consecutive residues identical to residues 11–14 of SEQ ID NO: 1. It is believed that residues 11–14 (i.e., SLGA) of SEQ ID NO: 1 play an important role in maintaining the biological function of 15 the short PEDF peptides. For example, according to various examples provided below, a 18-mer peptide (EQRTESIIHRALYYDLIS; SEQ ID NO: 7) without the SLGA residues fail to elicit any arteriogenesis in a subject. Also, based on experiments disclosed in both the prior application and the present application, it is suggested that a 25-mer peptide (EQRTESIIHRALYYDLISSPDIHGT; SEQ ID NO: 4) is ineffective in promoting muscle or 20 tendon regeneration and/or arteriogenesis in a subject.

[0077] The synthetic Peptides of the invention can be synthesized by commonly used methods such as t-BOC or Fmoc protection of alpha-amino groups. Both methods involve stepwise syntheses whereby a single amino acid is added at

each step starting from the C terminus of the peptide. Peptides of the present invention can also be synthesized by the well-known solid phase peptide synthesis methods.

[0078] Other synthetic peptides with conservative variation with respect to the 5 39-mer are also contemplated. The term “conservative variation” as used herein denotes the replacement of an amino acid residue by another, biologically similar residue. Examples of conservative variations include the substitution of one hydrophobic residue such as isoleucine, valine, leucine or methionine for one another, or the substitution of one polar residue for another, such as the substitution of 10 arginine for lysine, glutamic for aspartic acids, or glutamine for asparagine, and the like. The term “conservative variation” also includes the use of a substituted amino acid in place of an unsubstituted parent amino acid provided that antibodies raised to the substituted polypeptide also immunoreact with the unsubstituted polypeptide.

[0079] The synthetic peptides according to above-mentioned embodiments 15 may be formulated into pharmaceutical compositions for promoting muscle or tendon regeneration and/or arteriogenesis in a subject, which falls within other aspects of the present disclosure.

[0080] According to one embodiment of the present disclosure, the pharmaceutical composition comprises a synthetic peptide according to any of the 20 above-mentioned aspects/embodiments, and the synthetic peptide is present in an effective amount sufficient to promote the muscle or tendon regeneration and/or arteriogenesis in the subject. The pharmaceutical composition also comprises a pharmaceutically acceptable carrier for the synthetic peptide.

[0081] The choice of a pharmaceutically acceptable carrier to be used in conjunction with a synthetic peptide is basically determined by the way the pharmaceutical composition is to be administered. According to one optional embodiment of the present disclosure, the pharmaceutical composition may be 5 administered locally via intramuscular injection. In this case, the synthetic peptide may be formulated with a pharmaceutically acceptable carrier such as a sterile aqueous solution, which is preferably isotonic with the blood of the recipient. Such formulations may be prepared by dissolving or suspending the solid active ingredient in water containing physiologically compatible substances such as sodium chloride, 10 glycine, and the like, and having a buffered pH compatible with physiological conditions to produce an aqueous solution, and rendering said solution sterile.

[0082] Still optionally, the synthetic peptide may be formulated in a sustained-release dosage form so as to ensure a more prolonged therapeutic action of the treatment. There are several polymeric materials suitable for prolonging drug 15 release, examples of which include, but are not limited to, alginate, gelatin, collagen, and poly(lactide-co-glycolide).

[0083] According to some working examples of the present disclosure, the present synthetic peptides are embedded in a matrix of cross-linked alginate gel, and the final concentration of the synthetic peptides is about 1–100 μM , and preferably, 20 about 10 μM . For example, the concentration of the synthetic peptides may be about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 μM .

[0084] Pharmaceutical compositions of the invention can also comprise various additives known to those skilled in the art. For example, solvents, including relatively small amounts of alcohol, may be used to solubilize certain drug substances. Other optional pharmaceutically acceptable additives include opacifiers, antioxidants, 5 fragrance, colorant, gelling agents, thickening agents, stabilizers, surfactants, and the like. Other agents may also be added, such as antimicrobial agents, to prevent spoilage upon storage, i.e., to inhibit growth of microbes such as yeasts and molds. Permeation enhancers and/or irritation-mitigating additives may also be included in the composition of the present invention.

10 [0085] In yet another aspect, the present invention is directed to a method for promoting muscle or tendon regeneration in or adjacent to a damaged region of a subject; and in still another aspect, the present invention is directed to a method for promoting arteriogenesis in or adjacent to an ischemic region of a subject. In either embodiment, the subject may be any animal classified as a mammal, including human. 15 Embodiments applicable to either or both of these two aspects are discussed below.

[0086] In one embodiment, the method for promoting muscle or tendon regeneration in or adjacent to a damaged region of a subject comprises administering, to a treatment region of the subject, a therapeutically effective amount of the synthetic peptide of the present disclosure, wherein the treatment region is adjacent 20 to the damaged region so as to promote the muscle or tendon to regenerate in or adjacent to the damaged region of the subject to regenerate.

[0087] In another embodiment, the method for promoting arteriogenesis in or adjacent to an ischemic region of a subject comprises administering, to a treatment

region of the subject, a therapeutically effective amount of the synthetic peptide of the present disclosure, wherein the treatment region is adjacent to the ischemic region, so as to promote arteriogenesis in or adjacent to the ischemic region of the subject.

5 [0088] According to optional embodiments, the synthetic peptide is formulated in a pharmaceutical composition according to the above-mentioned aspect/embodiments of the present disclosure. In practice, the pharmaceutical composition may be administered via intramuscular injection.

10 [0089] According to some embodiments, the subject may be suffering from muscle injury, muscle disuse, muscular dystrophy, amyotrophic lateral sclerosis, tendon injury, tissue ischemia, cerebral ischemia, peripheral arterial diseases, or myocardial infarction, which causes the blood flow at the ischemic region to be hindered or blocked.

15 [0090] The following Examples are provided to elucidate certain aspects of the present invention and to aid those of skilled in the art in practicing this invention. These Examples are in no way to be considered to limit the scope of the invention in any manner. Without further elaboration, it is believed that one skilled in the art can, based on the description herein, utilize the present invention to its fullest extent.

20

EXAMPLES

[0091] **Materials and Methods**

[0092] *Materials*

[0093] Dulbecco's modified Eagle's medium (DMEM), fetal bovine serum (FBS), 0.25% trypsin, anti-BrdU antibody, MCDB131 medium, TRIzol, and Dynabeads were purchased from Invitrogen (Carlsbad, CA). Ultrapure alginate (6000 Da), dimethyl sulfoxide (DMSO), bovine serum albumin (BSA), 5-bromo-2'-deoxyuridine (BrdU), 5 Hoechst 33258 dye, and Masson's Trichrome were all from Sigma-Aldrich (St. Louis, MO). Collagenase type I and dispase II were obtained from Roche (Indianapolis, IN). All the fluorescent dye-conjugated secondary antibodies were purchased from BioLegend (San Diego, CA). Hematoxylin and eosin (H&E) dyes were purchased from Merck (Rayway, NJ, USA). Anti-collagen 1A1 antibody was obtained from Santa Cruz Biotechnology (Santa Cruz, CA). Matrigel was purchased from BD Biosciences (Bedford, MA). Anti-alpha-smooth muscle actin (anti- α -SMA) antibody (ab5694) and anti- nucleostemin antibody were from Abcam (Cambridge, MA). Anti-Pax7 antibody (GTX62311) was from GeneTex (Taipei, Taiwan). Isolectin B4 (IB4)-Alexa Fluor 568 was from Molecular Probes (Eugene, OR).

15 [0094] Short synthetic PEDF peptides, including 29-mer (SEQ ID No: 3), 25-mer (SEQ ID No: 4), 24-mer (SEQ ID No: 5), 20-mer (SEQ ID No: 6), 18-mer (SEQ ID No: 7), MO 29-mer (SEQ ID No: 8), and MO 20-mer (SEQ ID No: 9) were synthesized and modified with acetylated at the NH₂ termini and amidated at the COOH termini for stability and characterized by mass spectrometry (>95% purity) to order at GenScript 20 (Piscataway, NJ).

[0095] All animals used in embodiments of the present disclosure were housed in an animal room under temperature control (24-25°C) and 12:12 light-dark cycle. Standard laboratory chow and tap water were available *ad libitum*. The experiments

procedures were approved by the Mackay Memorial Hospital Review Board (New Taipei City, Taiwan, R.O.C.) and were performed in compliance with national animal welfare regulations.

[0096] PEDF peptide/alginate gel formulation and bolus formulation

5 [0097] Each PEDF-derived short synthetic peptide (the 29-mer, 25-mer, 24-mer, 20-mer, 18-mer, MO 29-mer, or MO 20-mer; hereinbelow, PEDF peptide) was reconstituted in DMSO as stock (5 mM). Then, ultrapure alginate was mixed with the stock to obtain a 2% wt/vol alginate solution with PEDF peptide at a final concentration of 10 μ M. The alginate solution was then filtered by membrane filter 10 (pore size, 0.22 μ m) and mixed with filtered calcium sulfate (0.21 g CaSO₄/mL of dH₂O) at a ratio of 25:1 (40 μ L of CaSO₄ per 1 mL of the filtered alginate solution). The mixture was let standing at RT for about 1 hour to allow for the cross-linking of the alginate. The resultant sustained-release formulation was then used in the treatment of muscle or tendon damage and ischemia.

15 [0098] For bolus delivery, a final PEDF concentration of 10 μ M was used by performing serial dilutions from the 5 mM stock solution.

[0099] Histology, Immunohistochemistry and Quantification

20 [0100] The gracilis, adductor magnus, soleus, and tibialis muscles were fixed in 4% paraformaldehyde, dehydrated with graded ethanol series, and paraffinized. Fixed samples were de-paraffinized in xylene and rehydrated in a graded series of ethanol. Tissues were sliced into 5- μ m sections. General histology was performed using H&E dye.

[0101] De-paraffinized tissue sections were blocked with 10% goat serum for 1 hour. Staining was done using primary antibodies against BrdU (1:50 dilution; GTX42641) or type I collagen 1A1 (1:50 dilution) overnight at 4°C, followed by incubation with the appropriate peroxidase-labeled donkey immunoglobulin for 30 5 min and then with chromogen substrate (3,3'-diaminobenzidine) for 2 min before counterstaining with hematoxylin. Quantification was estimated based on high quality images (1208 X 960 pixels) captured using a Nikon Eclipse 80i light microscope.

[0102] The muscle fiber size was determined on H&E-stained muscle cross section and quantified using the minimum distance of parallel tangents at opposing 10 particle borders (minimal “Feret’s diameter”). Pictures were captured using a Nikon Eclipse 80i light microscope, and the minimal Feret’s diameter was measured using the Image-Pro Plus 4.5.1 software (Media Cybernetics). Normalization of the number of fibers in each fiber Feret class of 5 µm was based on the total number of muscle fibers in each picture.

15 [0103] To ascertain the number of centrally nucleated muscle fibers, sections were stained with H&E and then photographed as described above. At least 100 stained fibers were randomly chosen from each photo. Muscle fibers were judged centrally nucleated if one or more nuclei were not located at the periphery of the fiber. The data were expressed as a % of the total number of muscle fibers counted. 20 Results were evaluated from 6 sections per muscle section, and 10 mice at each group.

[0104] De-paraffinized tendon tissue sections were stained using Masson’s Trichrome according to the manufacturer’s instructions. For semi-quantitative analysis of collagen area, 10 fields from each slide were randomly selected under a

light microscope, and the repairing area per intact tendon area of the cross section (mm^2/mm^2) was measured using the Image-Pro Plus 4.5.1 system.

[0105] *Isolation and culture of tendon stem cells*

[0106] New Zealand White rabbits (6-8 months old, 3.0-4.0 kg) were used in 5 this study. Achilles tendons were removed from the rabbits by cutting through their bony attachments. The tendon sheath was stripped away and the core portion of the tendons was minced into small fragments. Each 100 mg of fragment was then digested in a solution containing 3 mg/mL of type I collagenase and 4 mg/mL of dispase in 1 ml Dulbecco's Modified Eagle Medium (DMEM-high glucose) at 37°C for 2 10 hours. The resultant cell suspension was centrifuged at 1,000 rpm for 15 minutes to obtain a cell pellet which was then resuspended in a growth medium consisting of DMEM supplemented with 10% heat inactivated fetal bovine serum (FBS), 100 μM 2-mercaptoethanol, and 100 U/ml penicillin and 100 $\mu\text{g}/\text{ml}$ streptomycin. For 15 passage, near-confluent cells were harvested with 0.25% trypsin and then 1×10^5 subcultured cells were further cultured in medium.

[0107] *TSCs proliferation assay*

[0108] The TSCs at passage 4 were seeded at gelatin-coated slide in a 6-well plate at a density of 2×10^5 cells per well and cultured in growth medium (DMEM + 10% FBS) for 24 hours before being replaced by a basal growth medium with 5% FBS 20 only (control group) or with 5% FBS plus an additional 50 nM of PEDF-derived peptide (i.e., 29-mer, 24-mer, 20-mer, 18-mer, Mo 29-mer, or Mo 20-mer) for 24 hours. For BrdU labeling assay, BrdU (final concentration, 10 μM) was added to the culture for 4 hours. After fixing with 4% paraformaldehyde, cells were exposed to cold methanol

for 2 minutes, and then treated with 1 N HCl at RT for 1 hour before performing immunofluorescence. The phenotype of passage 4 TSC was determined by immunocytochemistry of nucleostemin and type I collagen. Almost all of expanded TSCs were nucleostemin and type I collagen-double positive cells.

5 [0109] *In vivo detection of DNA synthesis*

[0110] For the detection of cell expansion, BrdU was reconstituted in DMSO as stock (80 mM). 10 µl of BrdU mixed with 90 µl of PBS was intraperitoneally injected into the mouse 16 hours prior to euthanasia. Also, 150 µl of BrdU mixed with 350 µl of PBS was intraperitoneally injected into the rat 16 hours prior to euthanasia. DNA synthesis was assessed by BrdU labeling with anti-BrdU antibodies.

10 [0111] *Immunofluorescence Analysis*

[0112] De-paraffinized tissue sections or 4% paraformaldehyde fixed rabbit tendon stem cells (TSCs) were blocked with 10% goat serum and 5% BSA for 1 hour. Double staining was done using primary antibodies against α-SMA (1:100 dilution), IB4 (5 µg/ml), Pax7 (1:100 dilution), nucleostemin (1:100 dilution) and type I collagen 1A1 (1:50) at 37°C for 2 hours, followed by incubation with the appropriate rhodamine- or FITC-conjugated donkey IgG for 1 hour at RT. Nuclei were located by counterstaining with Hoechst 33258 for 7 minutes. Images were captured using a Zeiss epifluorescence microscope with a CCD camera.

15 [0113] The small artery densities (α-SMA positive cells surrounding the whole circumference of the vessel) were measured and images were taken from 10 randomly-selected areas of adductor magnus muscle (200x magnification) in each sample, and blinded quantification was performed in triplicate by manually counting

within each section; values from five sections were then averaged and expressed as arteriole density per mm².

[0114] Bone-marrow-derived Mesenchymal Stem Cells (BM-MSCs) Isolation, Cell Culture, and Treatments

[0115] Primary rat BM-MSCs were isolated from femur of male Sprague-Dawley rats (300–450 g). Femora were aseptically removed and dissected free of adhering tissues, and then the marrow cavities were flushed by injection of DMEM medium. Collected bone marrow cells were incubated in a 100 × 15-mm Petri dish in DMEM medium supplemented with 10% FBS, 100 U/ml penicillin, and 100 µg/ml streptomycin for 2 weeks in 5% CO₂ at 37°C. The medium was replaced every 2 to 3 days. For passage, near-confluent cells were detached by 0.25% trypsin and then 2 × 10⁵ subcultured cells were seeded in a well of 6-well plate and further cultured in the 10% FBS-DMEM. Before treatment, cells were starved for 12 hours in DMEM supplemented with 1% FBS followed by treatment with 50 nM PEDF-derived peptide (29-mer or 20-mer) in fresh 1% FBS-DMEM for either 24 or 48 hours.

[0116] RNA Extraction and Reverse Transcription–Polymerase Chain Reaction

[0117] The total RNA was extracted from cells using TRIzol and treated with RNase-free DNase I (Qiagen, Santa Clarita, CA) to remove genomic DNA and then purified with an RNA purification kit (Dynabeads). 1 µg of total RNA retrieved from BM-MSCs was reverse-transcribed into cDNA by 200 units of expand Reverse-Transcriptase (Roche, Mannheim, Germany) in 20 µl of reaction buffer containing 0.25 µg of random primers and 0.8 mM dNTPs at 42°C for 1 hour. 2 µl of the cDNA was used as templates in subsequent PCR reaction.

[0118] PCR was performed using a reaction volume of 30 μ l containing 15 μ l of EconoTaq[®] PLUS GREEN 2 \times Master Mix (Lucigen[®] Corp.), 1 μ M of each primer and 2 μ l of template DNA. cDNA was synthesized in an 18-22 cycle amplification reaction (denaturation, 20s, 94°C; annealing, 30s, 57°C; and polymerization, 40s, 72°C). Cycle 5 number for each primer set was established to be in the linear range of amplification. The primer set for the amplification of rat Tenomodulin gene (TNMD; accession number: NM_022290) included a forward primer of AGAATGAGCAATGGGTGGTC (SEQ ID No: 10) and a reverse primer of CTCGACCTCCTTGGTAGCAG (SEQ ID No: 11), and PCR products of about 240 bp were observed. Analysis of rat glyceraldehyde 3-phosphate 10 dehydrogenase (GAPDH; accession number: X02231.1) gene was used as a housekeeping gene for the normalization of the expression level. For the amplification of GAPDH gene, the primer set including a forward primer of AGACAGCCGCATCTTCTTGT (SEQ ID No: 12) and a reverse primer of CTTGCCGTGGGTAGAGTCAT (SEQ ID No: 13) was used, and PCR products of about 207 15 bp were observed.

[0119] The PCR products were electrophoresed in a 2% agarose gel containing ethidium bromide and visualized by UV illumination. The intensities of the PCR products were quantified densitometrically using a FUJI LAS-3000 system and Multi Gauge Ver. 1.01 software (Fujifilm, Tokyo, Japan).

20 [0120] *Statistics*

[0121] Results were expressed as the mean \pm standard error of the mean (SEM). One-way ANOVA was used for statistical comparisons. $P < 0.05$ was considered significant, unless otherwise specified.

[0122] Example 1**[0123] Sustained Release of PEDF Peptides from Alginate Gel**

[0124] To determine the release kinetics of 29-mer and 20mer, 100 μ g of 5 FITC-conjugated PEDF peptide was mixed with 100 μ L alginate solution, and then hydrogels were prepared as set forth in the “Materials and Methods” section. Thereafter, 100 mg hydrogel was incubated in 1.5 ml of PBS (pH 7.4) in microcentrifuge tube and placed in an orbital shaking incubator over a 6-day period at 37°C. The tube was centrifuged at each predetermined time point and then 200 μ L of supernatant was 10 removed and stored at -80°C for further analysis, and 200 μ L of fresh PBS was added to the tube to replace the supernatant withdrawn. The concentration of FITC-conjugated PEDF peptide present in the collected supernatants was determined using a fluorimeter in 96-well format. A known non-encapsulated FITC-peptide was used to generate a standard curve. Triplicate data were used for analysis.

[0125] The results of the assay, as summarized in Figure 1, revealed that the 15 embedded PEDF peptides were released in a sustained manner over a 6-day period. Specifically, approximately 48% of 29-mer and 35% of 20-mer peptide remained in the alginate gel matrix after 24 hours. Most of the 29-mer peptides (90%) were released within the first 4 days, after which time the release rate decreased significantly 20 thereby resulting in a plateau of the cumulative release curve. The 20-mer peptides were released in a slightly faster rate in which about 90% of the loaded 20-mer was released in the first 3 days.

[0126] Example 2**[0127] Sustained Release of PEDF Peptides Reduces Ischemic Damages**

[0128] Ischemic muscle injury typically leads to necrosis and loss of tissue and function. Hence, ischemic animal model was employed in the present examples to 5 investigate the possibility that the local delivery of the PEDF peptide/alginate gel formulation (herein “the sustained-release formulation”) may promote the recovery of tissue or organ functions in the case of tissue or organ damages. Various conditions associated with ischemic damages, such as, limb perfusion, tissue necrosis, arteriogenesis, and neovessel sprouting, were analyzed in the examples as follows.

10 [0129] 6-week-old female C57BL/6 wild-type mice were anesthetized by an intraperitoneal injection of a mixture of zoletil (6 mg/kg) and xylazine (3 mg/kg). Hair was removed from the hindquarter with a depilating cream. To establish hindlimb ischemia, unilateral external iliac and femoral arteries and veins were ligated, cut, and excised. After surgery, the mice were randomly assigned to several experimental 15 groups (n = 6, each group) and treated as follows. In the blank control group, the mice were treated with 50 μ l of blank alginate gel, whereas in the bolus control group, the mice received the bolus formulation containing 29-mer. In the PEDF peptide/alginate gel treatment groups, the mice received 50 μ l of the sustained-release formulation, which comprised either 29-mer, 24-mer, or 20-mer. 20 Additionally, in a PEDF 18-mer control group, mice were treated with a sustained-release formulation containing a PEDF 18-mer peptide. Treatments were applied by way of a single intramuscular injection to the gracilis muscle immediately

after femoral artery and vein excision operation. The incision was closed after the wound was irrigated with sterile saline.

[0130] Example 2.1

[0131] Sustained Release of PEDF Peptides Enhances Limb Perfusion

5 [0132] A laser Doppler perfusion imaging (LDPI) analyzer (Moor Instruments, USA) was used to quantify hindlimb blood perfusion before surgery (pre OP), immediately after surgery (post OP), and over time after surgery. To minimize vasoconstriction by anesthetic heat loss, animals were kept on a heating plate at 37°C for 5 min before measurement. Representative LDPI images illustrating the blood 10 perfusion of ischemic hindlimbs over a time period of 4 weeks were provided in Figure 2 in which dark blue color represents low blood flow. Blood perfusion is expressed as LDPI index representing the ratio of operated (ischemic) versus non-operated (non-ischemic) limb blood flow of the same mouse, and the results were summarized in Figure 3 and Table 1 ($n \geq 6$). Blood flow was displayed as changes in the laser 15 frequency, represented by different color pixels.

Table 1

Treatment	Ischemic/Non-ischemic Perfusion Ratio (%)					
	Pre OP	Post OP	7 days	14 days	21 days	28 days
Blank	99.4 \pm 1.5	8.1 \pm 0.87	30.6 \pm 1.9	28.4 \pm 3.9	46.3 \pm 3.8	50.0 \pm 6.5
Bolus	112.9 \pm 6.2	9.0 \pm 0.80	22.9 \pm 4.6	31.6 \pm 2.1	44.1 \pm 8.4	55.3 \pm 2.8
18-mer	104.5 \pm 2.5	7.5 \pm 0.67	23.8 \pm 4.5	30.3 \pm 0.94	46.8 \pm 4.3	52.8 \pm 7.4
29-mer	108.2 \pm 8.8	7.0 \pm 3.1	44.8 \pm 2.0*	77.6 \pm 6.8*	101 \pm 7.0*	105 \pm 4.8*
24-mer	91.5 \pm 5.6	8.0 \pm 0.03	53.1 \pm 0.37*	67.2 \pm 5.8*	86.1 \pm 5.9*	91.8 \pm 5.5*
20-mer	98.0 \pm 7.8	4.6 \pm 0.64	43.3 \pm 7.2*	60.0 \pm 9.0*	75.5 \pm 5.2*	92.7 \pm 3.0*

* $P < 0.05$ versus blank control.

[0133] As illustrated in Figure 3, after the surgery, the regional blood flow (post OP) was immediately reduced to about 8% of the non-ischemic limb of the same animal in all groups, as expected. Blank (alginate gel-only) control led to a slow increase in reperfusion over time. It should be noted that results from the bolus 5 delivery was similar to that of the blank control, indicating that sustained release of PEDF peptides is essential for exerting its protective effect. Also, mice treated with sustained-release formulation containing the control 18-mer peptide did not exhibit improved blood perfusion as compared with that of the blank control or the bolus control; suggesting that the 18-mer peptide is ineffective in treating ischemia. In 10 contrast, the present PEDF treatments significantly improved blood perfusion over that of the blank, bolus, and PEDF 18-mer control groups. In particular, animals treated with sustained formulations containing 29-mer, 24-mer, or 20-mer exhibited a marked increase in blood flow (at least about 60% of normal limbs) starting around the second weeks after the surgery. By four weeks after the surgery, the perfusion in animals 15 treated with 29-mer, 24-mer, and 20-mer delivered with the sustained-release formulations lead to a final recovery of, respectively, 105%, 92%, and 93% of normal limbs, as compared with 50% in the blank control and 55% in the bolus control.

[0134] Example 2.2

[0135] **Sustained Release of PEDF Peptides Prevents Ischemia-induced Tissue 20 Necrosis**

[0136] In most hindlimb ischemia models, tissue necrosis generally occurs in the muscles below the knee. For example, tibialis anterior muscle, which is distant to the gracilis muscle where the treatment was administered, often undergoes extensive

necrosis with regeneration after femoral artery excision. The intensity of Masson's trichrome blue color staining depended on the content of collagen fibers in the investigated tissue, and fibrosis is the result of necrosis. Hence, two weeks and seven weeks after the surgery and the treatment, samples from the tibialis anterior muscle 5 were analyzed by Masson's trichrome staining to assess the degree of fibrosis and hence necrosis. Results from representative samples are illustrated in Figures 4A and 4B.

[0137] As depicted in Figure 4A, at the second week after the surgery, muscle tissue from the blank control groups exhibited extensive fibrosis (indicated by the blue 10 stain), whereas muscle tissue treated with the present sustained-release formulation exhibited a relatively smaller fibrosis region. Note also in Figure 4A, at week 7 post-surgery, the treatment with the present sustained-release formulation effectively reduced the areas of necrosis and fibrosis thereby achieved a complete recovery of muscle tissue.

[0138] After ischemic injury, muscle fiber regeneration is achieved by 15 proliferation of satellite cells. The newly formed muscle fiber is marked by centrally located nuclei. Also, the necrotic area is evidenced by necrotic myofibers exhibiting a pale eosinophilic cytoplasm with oedema and a loss of peripheral nuclei. As revealed in Figure 4B, two weeks after the surgery, the regeneration of myofibers with centrally 20 located nuclei was more significant in mice treated with the present sustained-release formulation than that in mice treated with the blank control. Still referring to the upper panels of Figure 4B, the large pale red area in the sample from the blank control group, as compared with the sample from the 20-mer treatment group, also suggested

that the present sustained-release formulation was effective in preventing necrosis.

At week 7 post-surgery, small bundles of muscle fiber with intersperse fat droplet remained in 15% of muscle area in tibialis anterior muscle in groups treated with blank control (Figure 4B; lower left panel).

5 [0139] Statistical analyses regarding injured area (necrotic area + fibrotic area) and numbers of centrally nucleated fibers were also performed at two weeks after the surgery, and the results were summarized in Table 2. The injured area is expressed as the percent of total stained area (%), and the centrally nucleated fibers is expressed as the total number of muscle fibers counted (%).

10 [0140] The data summarized in Table 2 revealed that the injection of the present sustained-release formulation may substantially reduce tissue injury as compared with that of blank or bolus controls. Specifically, the injured areas of the PEDF treatment groups were reduced to about 45–48% of those of the blank or bolus control group. Also, these data suggested that the treatment with 29-mer, 24-mer, 15 or 20-mer formulation resulted in an increase (about 3–3.7-fold) in the number of centrally nucleated fibers in tibialis anterior muscle, as compared with that of blank or bolus control.

Table 2

Treatment	Injured Area (%)	Centrally nucleated fibers (%)
Blank	81.0±3.3	19.5±2.2
Bolus	80.1±4.1	20.2.±3.2
18-mer	78.5±5.1	20.5±3.3
29-mer	39.5±4.2*	72.5±5.2†
24-mer	36.8±5.5*	67.8±5.3†
20-mer	38.0±5.2*	61.2±5.8†

*P < 0.001 versus blank control.

20 †P < 0.02 versus blank control.

[0141] In sum, results in Example 2.2 suggested that the treatment with the present sustained formulation containing either 29-mer, 24-mer, or 20-mer may prevent necrosis and fibrosis induced by ischemia, and thereby may improve the 5 recovery of muscle tissue. Also, the increase of recovery of tibialis muscle in mice treated with the present sustained-release formulation provides additional evidence to support its effect on the promotion of blood perfusion in ischemic limb (Example 2.1 above).

[0142] Example 2.3

10 **[0143] Sustained Release of PEDF Peptides Stimulates Arteriogenesis That Supplements Ischemic Tissue with Collateral Circulation**

[0144] In the case of acute occlusion of a major artery (such as coronary artery or femoral artery), pre-existing arteriolar connections can be recruited to bypass the site of occlusion. This process is termed arteriogenesis which differs in many aspects 15 from angiogenesis. From the anatomical aspect, these pre-existing collateral arteries, unlike capillaries formed during angiogenesis, are microvascular, thin-walled conduits that are composed of an endothelial lining, an internal elastic lamina, and one or two layers of smooth muscle cells. Under normal conditions, these endogenous pre-existing thin-walled arterioles may not be utilized to provide perfusion. However, 20 following occlusion of a major artery, these vessels can dramatically increase their lumen by growth, bypassing the site of occlusion so as to provide enhanced perfusion to the jeopardized ischemic regions. During chronic or acute occlusion of a major artery, collateral arteries may ameliorate the ensuing detrimental effects in many

regions of the body (hindlimb, heart, brain, kidney, etc.). It is important to recognize that arteriogenesis is not a simple process of passive dilatation of pre-existing collateral arteries; rather, it is associated with active proliferation and remodeling by growth of pre-existing arteriolar connections into true collateral arteries. It is 5 established that vessel radius is the dominant influence on blood flow, and hence, the collateral arteries, after adaptive growth, are capable of conducting relatively large blood volumes per unit of time. Therefore, stimulation of arteriogenesis is probably the more efficient mechanism for the survival of ischemic limbs or internal organs such as heart and brain, in comparison to angiogenesis. In contrast, angiogenesis is the 10 formation of capillaries composed of endothelial cells from the pre-existing vessels; these capillaries are fruitless in proving higher profusion to the damaged ischemic region. Thus, the increase in blood flow to the potentially ischemic tissue, as caused by the development of two or three large collateral arteries, cannot be equaled by newly formed capillaries, however numerous.

15 [0145] To investigate the arteriogenic effect of the present sustained-release formulation, adductor magnus muscles (located at the same level as femoral artery excision and in which arteriogenesis responsible for establishing collateral circulation is expected to be found) were harvested from animals in each experimental condition, two weeks after the surgery. Arterioles in muscle cross sections were identified by 20 immunohistological staining for vascular smooth muscle cells (α -SMA; brown), and nuclei were labeled with hematoxylin; representative photographs were provided in Figure 5. Quantitative analysis was also performed and the results were summarized

in Table 3, and the data were expressed as α -SMA-positive arterioles per mm^2 in the peri-injury region.

Table 3

Treatment	Arteriole Density per mm^2
Blank	3.3 \pm 0.88
Bolus	3.7 \pm 1.2
18-mer	4.0 \pm 0.58
29-mer	11.7 \pm 1.5*
24-mer	10.0 \pm 0.58*
20-mer	10.7 \pm 1.2*

* $P < 0.001$ versus blank control.

5 [0146] These data revealed that the administration of the present sustained-release formulation increased arteriole density in adductor magnus muscle adjacent to the femoral artery excision, as compared with that of the blank and bolus control groups. Therefore, the sustained release of PEDF peptides provides arteriogenic activity to establish collateral circulation after the acute disruption of
10 blood supply. The dramatically increase the lumen of these vessels by growth provides enhanced perfusion to the jeopardized ischemic regions. This well-developed collateral network leads to the recovery from ischemic events.

[0147] Example 2.4

[0148] PEDF Peptide Stimulate *Ex Vivo* Neovessel Sprouting

15 [0149] To further confirm the neovessel development promoted by PEDF peptides, rat aortic ring sprouting assay was performed. Thoracic aortas were removed from euthanized rats and gently stripped of peri-aortic fibroadipose tissue. Aortas were sectioned into about 2-mm length rings, which were then embedded in a growth factor-reduced Matrigel. Gels containing the aortic rings were polymerized in

12-well plates incubated at 37°C for 30 minutes. 1 ml of MCDB131 medium supplemented with 100 units/ml penicillin and 100 ng/ml streptomycin, 1% FBS, and a supplemental agent (50 ng/ml VEGF-A, 20 ng/ml FGF-2, or 50 ng/ml 29-mer, 24-mer, 20-mer, Mo 29-mer, Mo 20-mer, 25-mer, or 18-mer) were added to the 5 Matrigel-containing explants. The cultures were propagated at 37 °C in a humidified incubator for up to 4 days, with media changes every other day. Neovessel sprouting was assessed until day 4 using an inverted microscope platform (Leica) with bright-field optics; representative photographs were provided in Figure 6. Quantification of neovessel sprouting was assessed using Image- Pro Plus 6.0 software 10 (Dendrites program). Results were expressed as a fold of untreated aortic ring, as summarized in Table 4. The experiment was repeated in triplicate.

[0150] In the untreated control (UT) in which no supplemental factor was administered, minimal neovessel sprouting was observed at day 4. It is also noted that the control PEDF peptides (i.e., 25-mer and 18-mer) did not substantially enhance 15 the neovessel sprouting, compared with the untreated control.

Table 4

Treatment	Neovessel Sprouting Fold
Untreated	1
VEGF	3.4*
FGF2	3.5*
18-mer	0.97
25-mer	0.89
29-mer	5.6*
24-mer	6.2*
20-mer	6.5*
Mo 29-mer	6.1*
Mo 20-mer	6.4*

*P < 0.02 versus untreated control.

[0151] As expected, the well-known angiogenic factors, VEGF and FGF2, induced substantial neovessel sprouting. The neovessel sprouting in samples treated with VEGF and FGF-2 increased for about 3.4-fold and 3.5-fold, respectively; as compared to that of the UT control.

5 [0152] The data in Table 4 also indicated that the present PEDF peptides (including 29-mer, 24-mer, 20-mer, Mo 29-mer, and Mo 20-mer) stimulated more neovessel sprouting than either VEGF or FGF2. These neovessels were examined by dual-staining immunofluorescence assay for α -smooth muscle actin (a marker of arteriole wall smooth muscle cell (SMC)) and isolectin B4 (IB4, a marker of endothelial 10 cells), and representative photographs were provided in Figure 7. As could be seen in Figure 7, samples treated with the present PEDF peptide (29-mer or 20-mer) displayed arteriole phenotype with a SMC coating. In contrast, the formation of endothelial tube and SMC proliferation was barely detected upon treatment with PEDF 18-mer. This result indicated that PEDF peptides according to embodiments of the present 15 disclosure can stimulate neovessel formation beyond the angiogenesis of capillary which only contains endothelial cells in culture. It thus supports the notion that the present PEDF peptides stimulate arteriogenesis *in vivo*.

[0153] In conclusion, data presented in Example 2 (including Examples 2.1 to 2.4) demonstrated that the present PEDF peptides were effective in enhancing limb 20 perfusion, reducing tissue necrosis and fibrosis, and promoting arteriogenesis and neovessel sprouting, and hence, the administration of the PEDF peptides (in particular, the sustained-release formulation containing either of the PEDF peptides) would reduce ischemic damages and facilitate the structural and functional recoveries of the

tissue or organ. It should be noted that it has been established that a 34-mer fragment of PEDF (residues 44–77) has anti-angiogenic properties, and a 44-mer fragment of PEDF (residues 78–121) has neurotrophic properties. However, the present disclosure is the first to confirm that short PEDF fragments (at least the 29-mer, 5 24-mer, 20-mer, Mo 29-mer, and Mo 20-mer) exhibit an arteriogenic activity.

[0154] Example 3

[0155] Sustained Release of PEDF Peptides Promotes Muscle Regeneration

[0156] To investigate the effects of the present PEDF peptides on muscle regeneration, a rat myonecrosis model of a single injection of bupivacaine into the soleus muscle was employed. Adult 10-week-old male Sprague-Dawley rats (initial body weight = 312 ± 11 g) were anesthetized by an intraperitoneal injection of a xylazine (10 mg/kg). Then, the soleus muscle was injured by unilaterally injecting 0.5 ml bupivacaine (AstraZeneca) with a disposable syringe with a 26-gauge needle. 15 Briefly, the needle was inserted into the distal portion of the soleus muscle and then receded longitudinally to the proximal portion accompanying evenly bupivacaine solution injection. The solution was then injected throughout the entire length of the muscle as the needle was slowly withdrawn.

[0157] After bupivacaine injection, rats were divided equally ($n = 10$ /group) 20 into four experimental groups and treated as follows. In the blank control group, the mice were treated with 50 μ l of blank alginate gel. In the treatments groups, the mice received 50 μ l of the sustained-release formulation (29-mer or 20-mer). The mice in the bolus control group received the bolus formulation (29-mer). Treatments

were applied by way of a single intramuscular injection to the soleus muscle immediately after the bupivacaine perfusion.

[0158] At day 4 after bupivacaine injection, histology of soleus muscle cross section consisted of general necrosis with disintegrating myofibers and abundant 5 infiltrating inflammatory cells occupying a great major part of the soleus muscle (photographs not shown). Only in the peripheral were some muscle fibers with relative normal structure remained. The degrees of muscle fiber necrosis were the same in blank control group and peptide treatment group. This result indicated that the necrosis levels induced by bupivacaine in different groups were substantially the 10 same.

[0159] Example 3.1

[0160] Sustained Release of PEDF Peptides Promotes Cell Proliferation

[0161] Muscle regeneration involves proliferation of muscle fibers, or muscle cells. The muscle fiber proliferation was assayed by the incorporation of BrdU in the 15 proliferating nuclei. Satellite cells proliferation is the key step of muscle regeneration. Hence, the soleus muscle specimens were also stained for satellite cell marker, Pax7, so as to investigate the muscle regeneration activity. Detailed assay procedures are as described in "Materials and Methods." The level of BrdU-positive cells was expressed as labeling index (%), which was computed as the number of labeled cells 20 divided by the total number of cells. The labeling index (%) of Pax7-positive cells was computed as the number of labeled cells divided by the total number of cells with nuclei. Quantitative results were evaluated from 6 sections per muscle section and 10 mice at each group, and were summarized in Table 5.

Table 5

Treatment	BrdU Labeling Index (%)	Pax7 Labeling Index (%)
Blank	4.2±0.9	2.0±0.71
Bolus	4.4±1.4	2.6±0.68
29-mer	15.4±1.7	16.2±2.0
20-mer	13.6±3.0	14.6±1.9

[0162] These results revealed that the numbers of BrdU-positive cells in wounds treated with the 29-mer- or 20-mer-containing sustained-release formulation 5 were significantly increased, as compared with wounds treated with the blank or bolus control. Regarding the proliferative activity of satellite cells, the data revealed that the present sustained-release formulations lead to higher percentages of Pax7-positive cells as compared with blank and bolus controls. Together, these data suggested that the administration of the present sustained-release formulations enhances the 10 proliferative activities of muscle fibers and/or satellite cells, which in turn may promote the muscle to regenerate.

[0163] Example 3.2

[0164] Sustained Release of PEDF Peptides Promotes Muscle Fiber Regeneration

[0165] In the process of muscle regeneration, newly generated muscle fibers 15 typically contain central located nuclei. Hence, the percentage of such centrally nucleated muscle fibers is also an indicator of the regenerative activity of muscle. Statistical analysis regarding the percentage of centrally nucleated fibers was performed at day 7 after the bupivacaine injection, and the results were summarized 20 in Table 6.

Table 6

Treatment	Centrally nucleated fibers (%)
Blank	11.0±1.7
Bolus	13.8±2.2
29-mer	75.2±7.0
20-mer	63.3±7.4

[0166] As could be seen in Table 6, there were higher percentages of muscle fibers containing centrally located nuclei in animals treated with the 29-mer- or 5 20-mer-containing sustained-release formulation, as compared with that of blank or bolus control groups. These results indicated that the administration of the present sustained-release formulations was effective in promoting muscle regeneration.

[0167] Also, at 14 days after the bupivacaine injection, necrotic myofibers were replaced by newly formed myotubes in soleus muscle in all experimental groups. 10 However, a number of centrally nucleated fibers remained in regenerating muscles treated with blank or bolus control (Figure 8), suggesting an incomplete muscle regeneration. In contrast, muscle sections from animals treated with the sustained-release formulation containing 29-mer or 20-mer exhibited much less centrally nucleated fibers. Together, these data indicated that the muscle fiber 15 regeneration was facilitated by the sustained release of the PEDF peptides.

[0168] Example 3.3

[0169] Sustained Release of PEDF Peptides Promotes Maturation of Regenerated Muscle Fiber

[0170] In the later stage of muscle regeneration, newly generated muscle 20 fibers start to gain size. Muscle specimens were collected at 14 days after injury, and

respective fiber diameters were measured in accordance with the procedure set forth in the section of "Materials and Methods." Results were summarized in Figure 9.

[0171] On average, diameters of muscle fibers from animals treated with the sustained-release formulation containing 29-mer or 20-mer were larger than those 5 from animals in the blank or bolus control group. In addition, the size distribution of the 20-mer-treated muscles was in close resemblance to that of uninjured, intact muscles. Specifically, about 56.6% of muscle fibers from animals treated with 20-mer and about 53.2% of intact muscle fibers had a minimal Feret's diameter between 15–25 μm , whereas about 59.6% and about 56.2% of the regenerated fibers from the 10 blank and bolus control groups had minimal Feret's diameters between about 10–20 μm . These data indicated that the administration of the present sustained-release formulation was effective in increasing the mass of the regenerated muscles.

[0172] In conclusion, data presented in Example 3 (including Examples 3.1 to 3.3) demonstrated that the present PEDF peptides are effective in promoting the 15 proliferations of muscle fibers and satellite cells, regeneration of muscle fibers, and the maturation of regenerated muscle fibers, and hence, the administration of the PEDF peptides (in particular, the sustained-release formulation containing either of the PEDF peptides) would promote the muscle regeneration process and facilitate the structural 20 and functional recoveries of the muscle tissue. The present disclosure is the first to discover that short PEDF fragments (at least the 29-mer and 20-mer) are capable of promoting muscle regeneration.

[0173] **Example 4**

[0174] Sustained Release of PEDF Peptides Promotes Tendon Regeneration

[0175] To investigate the effects of the present PEDF peptides on tendon regeneration, a rat model with tendon injury was established as follows. Adult 10-week-old male Sprague-Dawley rats (total n = 50; initial body weight = 312 ± 11 g) were anesthetized by an intraperitoneal injection of a xylazine (10 mg/kg). Then, the left tendo Achilles injury was created by full-thickness insertion of an 18-gauge needle through tendo Achilles 1 cm proximal to its insertion into the calcaneum. This created a horizontal (transaction) wound which was flanked by intact tendon tissue on both sides to prevent the retraction of severed ends.

[0176] The rats were randomly assigned (n = 10/group) to five experimental groups and treated as follows. In the blank control group, the mice were treated with 150 µl of blank alginate gel. For the bolus control group, 150 µl of the bolus formulation (29-mer) was administered. In the treatments groups, the mice received 150 µl of the sustained-release formulation (29-mer, 24-mer, or 20-mer). Treatments were injected subcutaneously near tendon lesion immediately after the injury, and the incision was closed after the wound was irrigated with sterile saline.

[0177] Example 4.1**[0178] Sustained Release of PEDF Peptides Promotes Tendon Healing**

[0179] Three weeks after tendon injury, histologic analysis was performed to observe the healing of the tendon. Representative photographs were provided in Figure 10. As could be seen in the upper panel of Figure 10, in animals treated with blank control, a broad band of disorganized fibrous scar was formed between the two cut ends, and the pink stained collagen bundle was minimal in the scar tissue. In

contrast, the cut ends of tendons treated with sustained-release formulation containing 29-mer were healed with much lesser scar tissue as compared with that of the blank control specimen, and the pinkish stain of mature collagen bundle extended into the damaged region of the tendon; also, tendon fibers from the cut ends seemed 5 to joint together in some area (Figure 10; lower panel). Also, the fibrous tissue in the scar was more organized and in parallel direction in the 29-mer treatment group.

[0180] Figure 11 provides representative photographs of histologic analysis at higher magnification. Normal tendon had a relative scarcity of cells among the collagen fibers, and the nuclei were mostly elongated. In the blank and bolus control 10 groups, after healing for three weeks, the more abundant presence of fibroblasts (characterized by the presence of round- or spindle-shaped fibroblast-like nuclei) was observed in the tendon, and the newly formed collagen fibers were structurally disorganized (the undamaged tissue was indicated by *). These morphological changes suggested poor healing of the tendon wounds in the blank and bolus control 15 groups.

[0181] In contrast, still referring to Figure 11, in tendons treated with the present sustained-release formulations, the healing regions had thin, elongated nuclei which were morphologically similar to nuclei of mature tenocyte, and the collagen fibers were well organized and parallel to the native tendon (deep pink); suggesting a 20 better healing of the tendon wounds. These results indicated that the tendon wound healing process may benefit from the present sustained-release formulation.

[0182] Further, Masson trichrome staining was performed to evaluate the structure and organization of collagen fibers, and representative photographs of the

specimens were provided in Figure 12. In the uninjured tendon, collagens fibers were substantially parallel to one another (Figure 12; left panel). By contrast, the injured tendon treated with blank control had disorganized collagen fibers in the healed region (Figure 12; middle panel; wound margin marked by *). The injured tendon treated 5 with the present sustained-release formulation, however, had well-organized collagen fibers that were aligned in substantially the same orientation as the uninjured tendon tissue beyond the wound margin (Figure 12; right panel; wound margin marked by *). These highly oriented and organized collagen fibers suggested a better tendon wound healing effect in animals treated with the present sustained-release formulation.

10 [0183] Quantitative analysis was also performed to assess the percentage of collagen (%) in the regenerated area, and results were summarized in Table 7.

Table 7

Treatment	Collagen in Regenerated Area (%)
Blank	56.0 \pm 6.87
Bolus	58.25 \pm 8.84
29-mer	88.75 \pm 1.89
24-mer	85.75 \pm 2.56
20-mer	82.01 \pm 6.55

15 [0184] As could be seen in Table 7, animals in the PEDF treatment groups (29-mer, 24-mer or 20-mer) had higher collagen contents in the regenerated area, as compared with those in the blank and bolus control groups. These data suggested that collagen synthesis in the wound region may be promoted by the administration of the present sustained-release formulation.

20 [0185] The specimens were also subjected to immunostaining of type I collagen, and nuclei were labeled with hematoxylin. Representative photographs

were provided in Figure 13, in which lower panels are photographs of the magnified regions respectively enclosed by dash lines in the upper panels. As could be appreciated, the collagen fibrils in the uninjured tendon were well cross-linked with one another, and hence, they were unlikely to be recognized by the anti-collagen 1A1 antibody. Therefore, only minimal amount of type I collagen (brown stain) was observed in the left panel of Figure 13. By comparing the photographs of the blank control group (Figure 13; middle panel) and the PEDF treatment group (Figure 13; right panel), it was certain that type I collagen (brown) was more abundant in the PEDF treatment groups than in the blank control group.

[0186] Collectively, these results suggested that the administration of sustained-release formulation containing the present PEDF peptide would stimulate type 1 collagen synthesis in cells in injured tendon tissues, facilitate collagen deposition in healed tissues, and promote a more organized alignment of collagen fibers, and thereby promote tendon regeneration.

[0187] **Example 4.2**

[0188] **PEDF Peptides Induces *In Vitro* TSC Proliferation**

[0189] It has been reported that during the tendon healing process, tendon stem cells (TSCs) would proliferate and differentiate into tenocytes. To investigate whether present PEDF peptides would induce the proliferation of TSCs *in vitro*, tendons stem cells were isolated and cultured as described in the "Materials and Methods" section. The purity of TSCs was confirmed by a TSC marker, nucleostemin, as well as the expression of type I collagen by TSCs; together, these analyses indicated a TSC purity of near 100% (data not shown). The proliferation of TSCs was confirmed

by BrdU pulse-labeling for 2 hours. Quantitative analysis of the level of BrdU-positive cells was performed as described above, and the results were summarized in Table 8.

Table 8

Treatment	BrdU Labeling Index (%)
Control	9.6 ± 2.1
29-mer	31.8 ± 3.6*
24-mer	29.0 ± 4.6*
20-mer	33.2 ± 6.6*
Mo 29-mer	33.2 ± 6.6*
Mo 20-mer	31.8 ± 3.1*

*P < 0.002 versus untreated control cells.

5

[0190] These data revealed that, as compared with TSCs cultured in control medium, TSCs cultured in media containing the present PEDF peptide (29-mer, 24-mer, 20-mer, Mo 29-mer, or Mo 20-mer) were more proliferative. Also, it should be noted that Mo 29-mer and Mo 20-mer are derived from the mouse PEDF peptide, and does not have 100% amino acid sequence identities to the 11–30 amino acid residues of the 10 39-mer. However, they respectively exhibited similar mitogenic activity to the short PEDF peptides (e.g., 29-mer, 24-mer, and 20-mer) derived from human PEDF.

[0191] Example 4.3

[0192] Sustained Release of PEDF Peptides Promotes *In Vivo* TSC Proliferation

15 After Tendon Injury

[0193] Specimens obtained from animals in different experimental groups of Example 4.1 were stained for nucleostemin (green) to investigate whether *in vivo* TSC proliferation would be promoted by the present sustained-release formulation during the tendon wound healing process. In quantitative analysis, ten randomly selected

microscopic fields in each experimental group were photographed, and the percentage of nucleostemin-positive cells per total cells (counterstained by Hoechst 33258; blue) was calculated. Quantitative results were summarized in Table 9.

Table 9

Treatment	Nucleostemin Labeling Index (%)
Blank	5.8 ± 1.8
Bolus	5.6 ± 1.4
29-mer	16.4 ± 2.9*
24-mer	16.8 ± 4.2*
20-mer	15.0 ± 3.9*

5

*P < 0.001 versus blank control.

[0194] These data revealed that levels of nucleostemin-positive TSC cells in animals treated with 29-mer, 24-mer, or 20-mer were elevated, as compared with those in the blank and bolus control groups. Taken together, results from Examples 10 4.1 and 4.3 suggested that the *in vivo* expansion of TSCs promoted by the administration of the present sustained-release formulation was coincident with the more prominent tendon healing effect, as compared with the native healing process.

[0195] Example 4.4

[0196] **PEDF Peptide Induces Tenocyte-like Cell Generation from Bone 15 Marrow-Derived Mesenchymal Stem Cells (BM-MSCs)**

[0197] Recently, it is established that adult mesenchymal stem cells (MSCs) could be used to regenerate functional tendons. In this example, BM-MSCs were cultured in a control medium or a medium containing either PEDF 29-mer or 20-mer to investigate the ability of the present PEDF peptides in promoting BM-MSC 20 differentiation into tenocytes. Tenomodulin gene (TNMD) is a gene predominantly

expressed in tendons, and is considered as the most reliable phenotypic marker of the tenocytic lineage. Hence, tenocyte differentiation was evaluated based on the expression of TNMD. Representative image from RT-PCR analysis was provided in Figure 14.

5 [0198] The results revealed that the present PEDF peptide (29-mer or 20-mer) is an effective inducer of tenocyte-like cell differentiation in cultured BM-MSCs. Since the mobilization and differentiation of BM-MSCs is a proposed mechanism of tendon repair *in vivo*, this observation suggested that the present PEDF peptide may repair tendon damage by promoting the differentiation of BM-MSCs into tenocytes. It also
10 indicated the potential of the present PEDF peptide to facilitate the synthesis of artificial tendons from scaffold matrix culture of BM-MSCs.

[0199] In conclusion, data presented in Example 4 (including Examples 4.1 to 4.4) demonstrated that the present PEDF peptides were effective in promoting the synthesis of well-organized collagen (in particular, type one collagen) fibrils and
15 proliferation of tendon stem cells, and hence, the administration of the present PEDF peptides (in particular, the sustained-release formulation containing either of the present PEDF peptides) would promote the tendon regeneration process and facilitate the structural and functional recoveries of the tendon tissue. The present disclosure is the first to discover that short PEDF fragments (at least the 29-mer and 20-mer) are
20 capable of promoting tendon regeneration and BM-MSCs differentiation into tenocytes.

[0200] Collectively, results from the preceding examples established that the present synthetic PEDF peptides (such as the 29-mer, 24-mer, 20-mer, Mo 29-mer, and

Mo 20-mer) may promote arteriogenesis in or adjacent to the ischemic region, muscle and tendon regeneration in or adjacent to the injured region. Accordingly, the present synthetic PEDF peptides are suitable for use as a therapeutic agent to promote muscle and tendon wound-healing and reduce ischemic damages.

5 [0201] It will be understood that the above description of embodiments is given by way of example only and that various modifications may be made by those with ordinary skill in the art. The above specification, examples, and data provide a complete description of the structure and use of exemplary embodiments of the invention. Although various embodiments of the invention have been described
10 above with a certain degree of particularity, or with reference to one or more individual embodiments, those with ordinary skill in the art could make numerous alterations to the disclosed embodiments without departing from the scope of this invention.

CLAIMS:

1. Use of a synthetic peptide for the manufacture of a medicament for promoting muscle regeneration, tendon regeneration, or arteriogenesis in a subject, wherein the synthetic peptide consists of an amino acid sequence having 20-39 amino acid residues in length and at least 80% amino acid sequence identity to SEQ ID NO: 1, wherein the amino acid sequence comprises at least 20 consecutive residues having at least 90% amino acid sequence identity to residues 11-30 of SEQ ID NO: 1, and wherein the amino acid sequence comprises 4 consecutive residues that are identical to residues 11-14 of SEQ ID NO: 1.
2. The use according to claim 1, wherein the synthetic peptide has the amino acid sequence of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 8, or SEQ ID NO: 9.
3. The use according to claim 1, wherein the synthetic peptide has the amino acid sequence of SEQ ID NO: 3, SEQ ID NO: 5, or SEQ ID NO: 6.
4. The use according to any one of claims 1-3, wherein the medicament comprises a pharmaceutically acceptable carrier.
5. The use according to claim 4, wherein the pharmaceutically acceptable carrier is a polymeric material selected from the group consisting of alginate, gelatin, collagen, and poly(lactide-co-glycolide).
6. The use according to claim 5, wherein the polymeric material is alginate.

7. The use according to any one of claims 4-6, wherein the synthetic peptide is present in the medicament at a concentration of 1-100 μ M.
8. The use according to any one of claims 4-6, wherein the synthetic peptide is present in the medicament at a concentration of 10 μ M.
9. The use according to any one of claims 4-8, wherein the medicament is formulated in a sustained-release form.
10. The use according to any one of claims 4-8, wherein the pharmaceutical composition is formulated for intramuscular injection.
11. A synthetic peptide for promoting muscle regeneration, tendon regeneration, or arteriogenesis in a subject, consisting of an amino acid sequence having 20-39 amino acid residues in length and at least 80% amino acid sequence identity to SEQ ID NO: 1, wherein the amino acid sequence comprises at least 20 consecutive residues having at least 90% amino acid sequence identity to residues 11-30 of SEQ ID NO: 1, and wherein the amino acid sequence comprises 4 consecutive residues that are identical to residues 11-14 of SEQ ID NO: 1.
12. The synthetic peptide for promoting muscle regeneration, tendon regeneration, or arteriogenesis in a subject according to claim 11, wherein the synthetic peptide has the amino acid sequence of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 8, or SEQ ID NO: 9.
13. The synthetic peptide for promoting muscle regeneration, tendon regeneration, or arteriogenesis in a subject according to claim 11, wherein the synthetic peptide has the amino acid sequence of SEQ ID NO: 3, SEQ ID NO: 5, or SEQ ID NO: 6.

14. A pharmaceutical composition for promoting muscle regeneration, tendon regeneration, or arteriogenesis in a subject, comprising,
the synthetic peptide according to any one of claims 11-13; and
a pharmaceutically acceptable carrier.
15. The pharmaceutical composition of claim 14, wherein the pharmaceutically acceptable carrier is a polymeric material selected from the group consisting of alginate, gelatin, collagen, and poly(lactide-co-glycolide).
16. The pharmaceutical composition of claim 15, wherein the polymeric material is alginate.
17. The pharmaceutical composition according to any one of claims 14-16, wherein the synthetic peptide is present at a concentration of 1-100 μ M.
18. The pharmaceutical composition according to any one of claims 14-16, wherein the synthetic peptide is present at a concentration of 10 μ M.
19. The pharmaceutical composition according to any one of claims 14-18, wherein the pharmaceutical composition is formulated in a sustained-release form.
20. The pharmaceutical composition according to any one of claims 14-18, wherein the pharmaceutical composition is formulated for intramuscular injection.
21. Use of a synthetic peptide for promoting muscle regeneration, tendon regeneration, or arteriogenesis in a subject, wherein the synthetic peptide consists of an amino acid sequence having 20-39 amino acid residues in length and at least 80% amino

acid sequence identity to SEQ ID NO: 1, wherein the amino acid sequence comprises at least 20 consecutive residues having at least 90% amino acid sequence identity to residues 11-30 of SEQ ID NO: 1, and wherein the amino acid sequence comprises 4 consecutive residues that are identical to residues 11-14 of SEQ ID NO: 1.

22. The use according to claim 21, wherein the synthetic peptide has the amino acid sequence of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 8, or SEQ ID NO: 9.

23. The use according to claim 21, wherein the synthetic peptide has the amino acid sequence of SEQ ID NO: 3, SEQ ID NO: 5, or SEQ ID NO: 6.

24. The use according to any one of claims 21-23, wherein the synthetic peptide is formulated with a pharmaceutically acceptable carrier.

25. The use according to claim 24, wherein the pharmaceutically acceptable carrier is a polymeric material selected from the group consisting of alginate, gelatin, collagen, and poly(lactide-co-glycolide).

26. The use according to claim 25, wherein the polymeric material is alginate.

27. The use according to any one of claims 24-26, wherein the synthetic peptide is present at a concentration of 1-100 μ M.

28. The use according to any one of claims 24-26, wherein the synthetic peptide is present at a concentration of 10 μ M.

29. The use according to any one of claims 24-28, wherein the synthetic peptide is formulated in a sustained-release form.
30. The use according to any one of claims 24-28, wherein the synthetic peptide is formulated for intramuscular injection.
31. The use according to any one of claims 1-10, wherein the subject suffers from a muscle injury, muscle disuse, muscular dystrophy, amyotrophic lateral sclerosis, tendon injury, tissue ischemia, cerebral ischemia, peripheral arterial diseases, or myocardial infarction.
32. The synthetic peptide according to any one of claims 11-13, wherein the subject suffers from a muscle injury, muscle disuse, muscular dystrophy, amyotrophic lateral sclerosis, tendon injury, tissue ischemia, cerebral ischemia, peripheral arterial diseases, or myocardial infarction.
33. The pharmaceutical composition according to any one of claims 14-20, wherein the subject suffers from a muscle injury, muscle disuse, muscular dystrophy, amyotrophic lateral sclerosis, tendon injury, tissue ischemia, cerebral ischemia, peripheral arterial diseases, or myocardial infarction.
34. The use according to any one of claims 21-30, wherein the subject suffers from a muscle injury, muscle disuse, muscular dystrophy, amyotrophic lateral sclerosis, tendon injury, tissue ischemia, cerebral ischemia, peripheral arterial diseases, or myocardial infarction.

1/10




Fig. 1




Fig. 2

2/10




Fig. 3

3/10




Fig. 4A




Fig. 4B

4/10




Fig. 5

5/10




Fig. 6




Fig. 7




Fig. 8

8/10




Fig. 9




Fig. 10

9/10




Fig. 11




Fig. 12

10/10




Fig. 13






Fig. 14


Normal




Blank



Bolus



29-mer



Bar: 50 μ M